

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2017/058859 A1

(43) International Publication Date

6 April 2017 (06.04.2017)

(51) International Patent Classification: (71) Applicant: CELGENE CORPORATION [US/US]; 86  
*A61K 39/395* (2006.01) *C07K 16/28* (2006.01)  
*C07K 16/18* (2006.01) Morris Avenue, Summit, NJ 07901 (US).

(21) International Application Number:

PCT/US2016/054089

(72) Inventors: BENNETT, Brydon; 10835 Ashlar Place, San Diego, CA 92131 (US). CHAMBERLAIN, Philip, Paul; 2829 Brant Street, San Diego, CA 92130 (US). HARI-HARAN, Kandasamy; 6575 Little McGonigle Ranch Road, San Diego, CA 92130 (US). JACKSON, Pilgrim; 4421 Collwood Lane, San Diego, CA 92115 (US).

(22) International Filing Date:

28 September 2016 (28.09.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/234,535 29 September 2015 (29.09.2015) US

KEHRY, Marilyn; 10440 Rock Creek Drive, San Diego, CA 92131 (US). KHATTRI, Roli; 9990 Scripps Vista Way, Unit 88, San Diego, CA 92131 (US). LEUNG, Monica; 2755 Havasupai Avenue, San Diego, CA 92117 (US). MORRISON, Lisa; 4918 Coach Horse Court, San Diego, CA 92130 (US). SUN, Jeonghoon; 4992 35th Street, San Diego, CA 92116 (US). TORRES, Sanaa; 2526 Corbel

[Continued on next page]

(54) Title: PD-1 BINDING PROTEINS AND METHODS OF USE THEREOF

(57) Abstract: Provided herein are compositions, methods and uses involving antibodies that specifically bind to Programmed Death-1 (PD-1) and modulate the expression and/or activity of PD-1.

- MDX 4H1
- PD1AB-1
- ▲ PD1AB-2
- ▲ PD1AB-6
- △ Isotype control
- ★ 10nM PD-L1-DyL650




FIG. 1A



Way, San Marcos, CA 92078 (US). **CHAN, Henry, H.**; 18787 Heritage Drive, Poway, CA 92064 (US).

(74) **Agents:** WEISSER, Tamera, M. et al.; JONES DAY, 250 Vesey Street, New York, NY 10281-1047 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

## PD-1 BINDING PROTEINS AND METHODS OF USE THEREOF

### CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority to U.S. Provisional Patent Application No. 62/234,535, filed September 29, 2015, the disclosure of which is incorporated by reference herein in its entirety.

#### 1. FIELD

**[0001]** Provided herein are compositions, methods, and uses involving antibodies that specifically bind to human Programmed Death-1 (PD-1) and modulate the expression and/or activity of PD-1.

#### 2. SUMMARY

**[0002]** The present disclosure provides proteins that bind to PD-1 (e.g., human PD-1, SEQ ID NO:43), including binding proteins such as antibodies that bind to PD-1. Such binding proteins, including antibodies, can bind to a PD-1 polypeptide, a PD-1 fragment, and/or a PD-1 epitope. Such binding proteins, including antibodies, can be agonists (e.g., induce PD-1 ligand-like signaling). In some embodiments, the binding proteins do not compete with PD-1 ligand (e.g., PD-L1 and PD-L2) for the interaction with PD-1 (e.g., a non-blocking antibody).

**[0003]** The present disclosure also provides, in certain embodiments, binding proteins, including antibodies or fragments thereof, that (i) bind to human PD-1, (ii) induce PD-1 ligand-like signaling, and (iii) do not compete with PD-L1 and/or PD-L2 for the interaction with PD-1.

**[0004]** Also provided herein are polynucleotides and vectors comprising sequences encoding such antibodies, cells (e.g., host cells) comprising such polynucleotides or vectors, and compositions, reagents, and kits comprising such antibodies. In another aspect, provided herein are methods for modulating PD-1 activity (e.g., activating PD-1 signaling) or PD-1 expression levels, diagnostic methods and uses of such anti-PD-1 antibodies.

**[0005]** In some embodiments, a binding protein (e.g., an anti-PD-1 antibody) comprises six complementarity determining regions (CDRs) or fewer than six CDRs. In other embodiments, a binding protein (e.g., an anti-PD-1 antibody) comprises one, two, three, four, five, or six CDRs selected from heavy chain variable region (VH) CDR1, VH CDR2, VH CDR3, light chain

variable region (VL) CDR1, VL CDR2, and/or VL CDR3. In certain embodiments, a binding protein (e.g., an anti-PD-1 antibody) comprises one, two, three, four, five, or six CDRs selected from VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of a monoclonal antibody designated as PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as described herein, or a humanized variant thereof. In some embodiments, a binding protein (e.g., an anti-PD-1 antibody) further comprises a scaffold region or framework region (FR), including a VH FR1, VH FR2, VH FR3, VH FR4, VL FR1, VL FR2, VL FR3, and/or VL FR4 of a human immunoglobulin amino acid sequence or a variant thereof.

**[0006]** In some embodiments, the antibody or antigen-binding fragment thereof binds to an epitope of human PD-1 recognized by an antibody comprising a light chain variable region having an amino acid sequence of SEQ ID NO:8 and a heavy chain variable region having an amino acid sequence of SEQ ID NO:13.

**[0007]** In other embodiments, the antibody or antibody fragment thereof competes for the binding to human PD-1 with an antibody comprising a light chain variable region having an amino acid sequence of SEQ ID NO:8 and a heavy chain variable region having an amino acid sequence of SEQ ID NO:13.

**[0008]** In some embodiments, the antibody or antigen-binding fragment thereof comprises a VL comprising VL CDR1, VL CDR2, and VL CDR3 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in **Table 1**.

**[0009]** In other embodiments, the antibody or antigen-binding fragment thereof comprises a VH comprising VH CDR1, VH CDR2, and VH CDR3 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in **Table 2**.

**[0010]** In other embodiments, the antibody or antigen-binding fragment thereof comprises:

- (a) a VL comprising VL FR1, VL FR2, VL FR3, and VL FR4 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in **Table 3**; and
- (b) a VH comprising VH FR1, VH FR2, VH FR3, and VH FR4 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in **Table 4**.

**[0011]** In certain embodiments, the VL CDR1, VL CDR2, and VL CDR3 of the antibody or antigen-binding fragment thereof comprise amino acid sequences of SEQ ID NOS:1, 2, and 3,

respectively, and the VH CDR1, VH CDR2, and VH CDR3 of the antibody or antigen-binding fragment thereof comprise amino acid sequences of SEQ ID NOS:4, 5, and 6, respectively.

**[0012]** In yet another embodiment, the VL CDR1, VL CDR2, and VL CDR3 of the antibody or antigen-binding fragment thereof comprise amino acid sequences of SEQ ID NOS:7, 2, and 3, respectively, and the VH CDR1, VH CDR2, and VH CDR3 of the antibody or antigen-binding fragment thereof comprise amino acid sequences of SEQ ID NOS:4, 5, and 6, respectively.

**[0013]** In another embodiment, the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:8. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0014]** In certain embodiments, the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:9. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0015]** In some embodiments, the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:10. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0016]** In certain embodiments, the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:11. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0017]** In other embodiments, the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:12. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0018]** In another embodiment, the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:13. In some embodiments, the amino acid sequence comprises one or more conservative modifications thereof.

**[0019]** In certain embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and (b) a VH comprising an amino acid sequence of SEQ ID NO:11.

**[0020]** In some embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and (b) a VH comprising an amino acid sequence of SEQ ID NO:11.

**[0021]** In other embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and (b) a VH comprising an amino acid sequence of SEQ ID NO:11.

**[0022]** In one embodiment, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and (b) a VH comprising an amino acid sequence of SEQ ID NO:12.

**[0023]** In another embodiment, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and (b) a VH comprising an amino acid sequence of SEQ ID NO:12.

**[0024]** In certain embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and (b) a VH comprising an amino acid sequence of SEQ ID NO:12.

**[0025]** In some embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

**[0026]** In other embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

**[0027]** In certain embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

**[0028]** In some embodiments, the amino acid sequence of the VL comprises one or more conservative modifications thereof. In some embodiments, the amino acid sequence of the VH comprises one or more conservative modifications thereof. In some embodiments, the amino acid sequence of the VL and the VH comprises one or more conservative modifications thereof.

**[0029]** In some embodiments, the antibody comprises a human IgG1 Fc region. In other embodiments, the antibody comprises a variant human IgG1 Fc region.

**[0030]** In one embodiment, the antibody comprises a human IgG1-K322A Fc region.

**[0031]** In some embodiments, the antibody comprises a human IgG4 Fc region. In other embodiments, the antibody comprises a variant human IgG4 Fc region.

**[0032]** In another embodiment, the antibody comprises a human IgG4P Fc region.

[0033] In still another embodiment, the antibody comprises a human IgG4PE Fc region.

[0034] In some embodiments, the antibody or antigen-binding fragment thereof further comprises a light chain constant region comprising an amino acid sequence of SEQ ID NO:41.

[0035] In other embodiments, the antibody or antigen-binding fragment thereof further comprises a heavy chain Fc region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:36-40.

[0036] In yet another embodiment, the antibody or antigen-binding fragment thereof further comprises a light chain constant region comprising an amino acid sequence of SEQ ID NO:41; and a heavy chain Fc region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:36-40.

[0037] In some embodiments, the antibody or antigen-binding fragment thereof comprises a light chain comprising an amino acid sequence of SEQ ID NO:31.

[0038] In another embodiment, the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:32.

[0039] In other embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:32.

[0040] In certain embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:33.

[0041] In other embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:33.

[0042] In one embodiment, the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:34.

[0043] In yet another embodiment, the antibody or antigen-binding fragment thereof comprises: (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:34.

[0044] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:35.

**[0045]** In other embodiments, the antibody or antigen-binding fragment thereof comprises: (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:35.

**[0046]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one of residues 100-109 within an amino acid sequence of SEQ ID NO:42.

**[0047]** In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one of residues 100-105 within an amino acid sequence of SEQ ID NO:42.

**[0048]** In particular embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one residue selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0049]** In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to two or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0050]** In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to three or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0051]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to four or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0052]** In one embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to five or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0053]** In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to six or more residues selected from the group consisting of N33, T51,

S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0054]** In yet another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to seven or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0055]** In still another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to eight or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0056]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to nine or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0057]** In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to all ten residues from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0058]** In one embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to N33 within an amino acid sequence of SEQ ID NO:42.

**[0059]** In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to T51 within an amino acid sequence of SEQ ID NO:42.

**[0060]** In a particular embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to S57 within an amino acid sequence of SEQ ID NO:42.

**[0061]** In one specific embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to L100 within an amino acid sequence of SEQ ID NO:42.

**[0062]** In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to N102 within an amino acid sequence of SEQ ID NO:42.

**[0063]** In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to G103 within an amino acid sequence of SEQ ID NO:42.

**[0064]** In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to R104 within an amino acid sequence of SEQ ID NO:42.

**[0065]** In yet another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to G103 and R104 within an amino acid sequence of SEQ ID NO:42.

**[0066]** In still another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to D105 within an amino acid sequence of SEQ ID NO:42.

**[0067]** In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to H107 within an amino acid sequence of SEQ ID NO:42.

**[0068]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to S109 within an amino acid sequence of SEQ ID NO:42.

**[0069]** In one embodiment, the epitope of human PD-1 is distinct from the PD-L1 binding site. In another embodiment, the epitope of human PD-1 is distinct from the PD-L2 binding site. In a specific embodiment, the epitope of human PD-1 is distinct from both the PD-L1 binding site and the PD-L2 binding site.

**[0070]** In an embodiment, the antibody or antigen-binding fragment thereof specifically binds to human PD-1 and/or monkey PD-1 (for example, cynomolgus monkey), but not rodent PD-1.

**[0071]** In certain embodiments, the antibody or antigen-binding fragment thereof has attenuated antibody dependent cellular cytotoxicity (ADCC) activity. In other embodiments, the antibody or antigen-binding fragment thereof has attenuated complement dependent cytotoxicity (CDC) activity. In some embodiments, the antibody or antigen-binding fragment thereof has attenuated ADCC and/or attenuated CDC activity.

**[0072]** In one aspect, provided herein is an antibody or antigen-binding fragment thereof that binds to an epitope of human PD-1, wherein the antibody or antigen-binding fragment thereof: (a) attenuates T cell activity; and/or (b) downregulates PD-1 expression on the surface of T cells.

**[0073]** In one embodiment, the antibody attenuates T cell activity. In another embodiment, the antibody downregulates PD-1 expression on the surface of T cells.

**[0074]** In some embodiments, the attenuation of T cell activity is measured by a T cell effector function.

**[0075]** In certain embodiments, the attenuation of T cell activity by the antibody or antigen-binding fragment thereof occurs in human PBMC or whole blood samples.

**[0076]** In some embodiments, the attenuation of T cell activity is measured by inhibition of cytokine production.

**[0077]** In other embodiments, the cytokine that is inhibited by the antibody or antigen-binding fragment thereof comprises IL-2, IL-17, and/or IFN- $\gamma$ . In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In some embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-6. In another embodiment, the cytokine is IL-12. In some other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IL-22. In still other embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is IFN- $\gamma$ . In yet other embodiments, the cytokine is TNF- $\alpha$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0078]** In certain embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof. In another embodiment, the downregulation occurs as early as 6 hours after the contact. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact. In still another embodiment, the downregulation occurs as early as 10 hours after the contact. In one embodiment, the downregulation occurs as early as 12 hours after the contact. In another embodiment, the downregulation occurs as early as 14 hours after the contact. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact. In still another embodiment, the downregulation occurs as early as 18 hours after the contact. In one embodiment, the downregulation occurs as early as 20 hours after the contact. In another embodiment, the downregulation occurs as early as 22 hours after the contact. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact. In some embodiments, the contact is with the antibody. In other embodiments, the contact is with an antigen-binding fragment thereof.

**[0079]** In some embodiments, the downregulation of PD-1 expression on the surface of T cells precedes cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the

downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition.

**[0080]** In other embodiments, the downregulation of PD-1 expression on the surface of T cells is concurrent with cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another

embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition.

**[0081]** In yet other embodiments, the downregulation of PD-1 expression on the surface of T cells is after cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In another embodiment, the downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In another embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In

another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition.

**[0082]** In one embodiment, the  $K_D$  of the antibody or antigen-binding fragment thereof for binding to purified human PD-1 is from about 1 nM to about 100 nM. In another embodiment, the  $K_D$  of the antibody or antigen-binding fragment thereof for binding to human PD-1 expressed on a cell surface is from about 100 pM to about 10 nM. In another embodiment, the  $K_D$  of the antibody or antigen-binding fragment thereof for binding to monkey PD-1 expressed on a cell surface is from about 100 pM to about 10 nM.

**[0083]** In some embodiments, the  $EC_{50}$  of the antibody or antigen-binding fragment thereof for attenuating T cell activity is from about 1 pM to about 10 pM, from about 10 pM to about 100 pM, from about 100 pM to about 1 nM, from about 1 nM to about 10 nM, or from about 10 nM to about 100 nM.

**[0084]** In other embodiments, the maximal percent attenuation of T cell activity by the antibody or antigen-binding fragment thereof is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

**[0085]** In another embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

**[0086]** In certain embodiments, the antibody is a monoclonal antibody. In some embodiments, the antibody is a humanized, human, or chimeric antibody. In another embodiment, the humanized antibody is a deimmunized antibody or a composite human antibody. In certain embodiments, the antibody is a humanized antibody. In specific embodiments, the antibody is a humanized antibody that specifically binds human PD-1.

**[0087]** In certain embodiments, the antibody or antigen-binding fragment thereof is a Fab, a Fab', a  $F(ab')_2$ , a Fv, a scFv, a dsFv, a diabody, a triabody, or a tetrabody. In some embodiments, the antibody or antigen-binding fragment thereof is a multispecific antibody formed from antibody fragments. In other embodiments, the antibody or antigen-binding fragment thereof is a bispecific antibody.

**[0088]** In some embodiments, the antibody or antigen-binding fragment thereof is conjugated to an agent. In one embodiment, the agent is a radioisotope, a metal chelator, an enzyme, a fluorescent compound, a bioluminescent compound, or a chemiluminescent compound.

**[0089]** Also provided herein is a composition comprising an antibody or antigen-binding fragment thereof provided herein. In certain embodiments, the composition further comprises a pharmaceutically acceptable carrier.

**[0090]** Also provided herein is a polynucleotide comprising nucleotide sequences encoding a VH, a VL, or both a VH and a VL of an antibody provided herein.

**[0091]** The present disclosure also provides, in certain embodiments, a polynucleotide comprising nucleotide sequences encoding a heavy chain, a light chain, or both a heavy chain and a light chain of an antibody provided herein.

**[0092]** In certain embodiments, the polynucleotide is operably linked to a promoter.

**[0093]** Also provided herein is a population of polynucleotides comprising: (a) a first polynucleotide comprising nucleotide sequences encoding a VH or a heavy chain of an antibody provided herein, and (b) a second polynucleotide comprising nucleotide sequences encoding a VL or a light chain of an antibody provided herein. In certain embodiments, the first polynucleotide is operably linked to a first promoter, and the second polynucleotide is operably linked to a second promoter.

**[0094]** Also provided herein is a vector comprising a polynucleotide provided herein.

**[0095]** The present disclosure also provides, in certain embodiments, a population of vectors comprising: (a) a first vector comprising nucleotide sequences encoding a VH or a heavy chain of an antibody provided herein, and (b) a second vector comprising nucleotide sequences encoding a VL or a light chain of an antibody provided herein.

**[0096]** The present disclosure further provides, in certain embodiments, a population of vectors comprising: (a) a first vector comprising nucleotide sequences encoding a VH or a heavy chain of an antibody provided herein operably linked to a first promoter, and (b) a second vector comprising nucleotide sequences encoding a VL or a light chain of an antibody provided herein operably linked to a second promoter.

**[0097]** The present disclosure provides, in certain embodiments, a cell comprising a polynucleotide or a population of polynucleotides provided herein.

**[0098]** The present disclosure also provides, in certain embodiments, a cell comprising a vector or a population of vectors provided herein.

**[0099]** The present disclosure further provides, in certain embodiments, an isolated cell producing an antibody or antigen-binding fragment thereof provided herein.

**[0100]** Also provided herein is a population of cells comprising: (a) a first host cell comprising a polynucleotide comprising nucleotide sequences encoding a VH or a heavy chain of an antibody provided herein, and (b) a second host cell comprising a polynucleotide comprising nucleotide sequences encoding a VL or a light chain of an antibody provided herein.

**[0101]** Further provided herein is a population of cells comprising: (a) a first host cell comprising a polynucleotide comprising nucleotide sequences encoding a VH or a heavy chain of an antibody provided herein operably linked to a first promoter, and (b) a second host cell comprising a polynucleotide comprising nucleotide sequences encoding a VL or a light chain of an antibody provided herein operably linked to a second promoter.

**[0102]** Also provided herein is a kit comprising an antibody or antigen-binding fragment thereof provided herein.

**[0103]** Also provided herein is a method of making an antibody or antigen-binding fragment thereof that specifically binds to an epitope of human PD-1. In certain embodiments, the method comprises culturing a cell provided herein to express the antibody or antigen-binding fragment thereof. In other embodiments, the method comprises expressing a polynucleotide provided herein. In one embodiment, the epitope of human PD-1 is distinct from the PD-L1 binding site. In another embodiment, the epitope of human PD-1 is distinct from the PD-L2 binding site. In a specific embodiment, the epitope of human PD-1 is distinct from both the PD-L1 binding site and the PD-L2 binding site.

**[0104]** In another aspect, provided herein is a method of attenuating activity of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen-binding fragment thereof provided herein. In one embodiment, the cell is contacted with an antibody provided herein. In another embodiment, the cell is contacted with an antigen-binding fragment of an antibody provided herein. In one embodiment, the maximal percent attenuation of T cell activity is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%. In some embodiments, the attenuation of T cell activity is measured by inhibition of cytokine production. In some embodiments, the cytokine is selected from the

group consisting of IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In some embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-6. In another embodiment, the cytokine is IL-12. In some other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IL-22. In still other embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is IFN- $\gamma$ . In yet other embodiments, the cytokine is TNF- $\alpha$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated. In certain embodiments, the inhibition of cytokine production is preceded by downregulation of PD-1 expression on the surface of the T cell. In some embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, 14 hours, 16 hours, 18 hours, 20 hours, 22 hours, or 24 hours after the contact with the antibody or antigen-binding fragment thereof. In one embodiment, the downregulation occurs as early as 4 hours after the contact, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 6 hours after the contact, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 8 hours after the contact, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 10 hours after the contact, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 12 hours after the contact, and precedes cytokine inhibition. In an embodiment, the downregulation occurs as early as 14 hours after the contact, and precedes cytokine inhibition. In other embodiments, the downregulation occurs as early as 16 hours after the contact, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 18 hours after the contact, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 20 hours after the contact, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 22 hours after the contact, and precedes cytokine inhibition. In some embodiments, the downregulation occurs as early as 24 hours after the contact, and precedes cytokine inhibition. In some embodiments, the inhibition of cytokine production is concurrent with downregulation of PD-1 expression on the surface of the

T cell. In one embodiment, the downregulation occurs as early as 4 hours after the contact, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 6 hours after the contact, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 8 hours after the contact, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 10 hours after the contact, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 12 hours after the contact, and is concurrent with cytokine inhibition. In an embodiment, the downregulation occurs as early as 14 hours after the contact, and is concurrent with cytokine inhibition. In other embodiments, the downregulation occurs as early as 16 hours after the contact, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 18 hours after the contact, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 20 hours after the contact, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 22 hours after the contact, and is concurrent with cytokine inhibition. In some embodiments, the downregulation occurs as early as 24 hours after the contact, and is concurrent with cytokine inhibition. In some embodiments, the inhibition of cytokine production precedes downregulation of PD-1 expression on the surface of the T cell. In one embodiment, the downregulation occurs as early as 4 hours after the contact, and is preceded by cytokine inhibition. In one embodiment, the downregulation occurs as early as 6 hours after the contact, and is preceded by cytokine inhibition. In one embodiment, the downregulation occurs as early as 8 hours after the contact, and is preceded by cytokine inhibition. In one embodiment, the downregulation occurs as early as 10 hours after the contact, and is preceded by cytokine inhibition. In another embodiment, the downregulation occurs as early as 12 hours after the contact, and is preceded by cytokine inhibition. In an embodiment, the downregulation occurs as early as 14 hours after the contact, and is preceded by cytokine inhibition. In other embodiments, the downregulation occurs as early as 16 hours after the contact, and is preceded by cytokine inhibition. In one embodiment, the downregulation occurs as early as 18 hours after the contact, and is preceded by cytokine inhibition. In another embodiment, the downregulation occurs as early as 20 hours after the contact, and is preceded by cytokine inhibition. In one embodiment, the downregulation occurs as early as 22 hours after the contact, and is preceded by cytokine inhibition.

cytokine inhibition. In some embodiments, the downregulation occurs as early as 24 hours after the contact, and is preceded by cytokine inhibition.

**[0105]** In yet another aspect, provided herein is a method of downregulating PD-1 expression on the surface of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen-binding fragment thereof provided herein. In one embodiment, the cell is contacted with an antibody provided herein. In another embodiment, the cell is contacted with an antigen-binding fragment of an antibody provided herein. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%. In another embodiment, the downregulation of PD-1 expression on the surface of the T cell occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof. In one embodiment, the downregulation of PD-1 expression on the surface of the T cell precedes cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of the T cell is concurrent with cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of the T cell is preceded by cytokine inhibition. In certain embodiments, the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In some embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-6. In another embodiment, the cytokine is IL-12. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IL-22. In still other embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is IFN- $\gamma$ . In yet other embodiments, the cytokine is TNF- $\alpha$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0106]** In another aspect, provided herein is a method of inhibiting cytokine production by a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen-binding fragment thereof described herein. In one embodiment, the cell is contacted with an antibody provided herein. In another embodiment, the cell is contacted with an antigen

binding fragment of an antibody provided herein. In some embodiments, the cytokine is selected from the group consisting of IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In some embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-6. In another embodiment, the cytokine is IL-12. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IL-22. In still other embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is IFN- $\gamma$ . In yet other embodiments, the cytokine is TNF- $\alpha$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

### 3. BRIEF DESCRIPTION OF THE FIGURES

[0107] FIGS. 1A-1B show that the T cell attenuating anti-PD-1 antibodies (PD1AB) do not compete with PD-L1 (PD-L1-DyL650 denotes PD-L1 conjugated with the dye DyL650) binding to PD-1: (A) PD1AB-1, PD1AB-2, and PD1AB-6; (B) PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, and PD1AB-5. MDX 4H1, an antagonist antibody, blocks PD-L1 binding to PD-1.

[0108] FIG. 2 depicts that the PD-1:PD1AB-6 Fab interaction site is at a distal side of PD-1 relative to the PD-1:PD-L1 interaction site.

[0109] FIG. 3 depicts that PD1AB-6 Fab binds against a PD-1  $\beta$  sheet, with substantial interactions formed with a PD-1 loop composed of residues 100-105.

[0110] FIG. 4 shows the amino acid sequences of heavy chain (HC) and light chain (LC) of PD1AB-6-IgG1 and HC of its variants PD1AB-6-K3 and PD1AB-6-4P.

[0111] FIGS. 5A-5B depict the PD1AB-6-IgG1 affinity for cynomolgus (A) or human (B) PD-1 expressed on CHO cells.

[0112] FIG. 6 depicts the binding of PD1AB-6-IgG1, isotype control, and human PD-L1 Fc fusion protein (hPD-L1 Fc) to activated human PBMC gated on CD4+ T cells.

[0113] FIG. 7 depicts the binding of PD1AB-6-IgG1, isotype control, and human PD-L1 Fc fusion protein (hPD-L1 Fc) to activated cynomolgus PBMC gated on CD4+ T cells.

[0114] FIGS. 8A-8D show the PD1AB-6 variants binding to Fc $\gamma$ RI (A), Fc $\gamma$ RIIIa (V158) (B), or Fc $\gamma$ RIIb (C) expressed on HEK293 cells using Cisbio Tag-lite<sup>TM</sup> detection, and (D) the EC<sub>50</sub> values of the PD1AB-6 variants binding to Fc $\gamma$ RI, Fc $\gamma$ RIIIa (V158), or Fc $\gamma$ RIIb.

[0115] FIGS. 9A-9C depict the PD1AB-6 variants binding to Fc $\gamma$ RIIIa (V158) (A) or Fc $\gamma$ RI (B) expressed on CHO cells using FACS, and (C) the EC<sub>50</sub> values of the PD-1 antibody variants binding to Fc $\gamma$ RI or Fc $\gamma$ RIIIa.

[0116] FIGS. 10A-10B depict the ADCC activity of the PD1AB-6 variants and a control human IgG1 Fc among two representatives of four individual healthy donors: (A) Donor 7 and (B) Donor 8.

[0117] FIG. 11 depicts the CDC activity of the PD1AB-6 variants. Data are representative of 3 independent experiments: (i) CDC activity of PD1AB-6-IgG1 and anti-human CD20 IgG1; (ii) CDC activity of PD1AB-6-IgG1 and PD1AB-6-K3; (iii) CDC activity of PD1AB-6-4P and commercial human IgG4 isotype control antibody and human IgG1 Fc protein.

[0118] FIG. 12 depicts the potent attenuating activity of PD1AB-6 variants in human PBMC assay, measured by IL-2 levels in culture supernatants at 24 hours post-stimulation.

[0119] FIG. 13 depicts the activity of PD1AB-6-K3 in human whole blood assay. The graph shows a representative curve from donor 4 used to calculate EC<sub>50</sub> of IFN- $\gamma$  inhibition. The table shows EC<sub>50</sub> values of IFN- $\gamma$  inhibition for 4 healthy donors with PD1AB-6 variants and CTLA4Ig.

[0120] FIGS. 14A-14C depict downregulation of PD-1 expression by PD1AB-6-IgG1 as determined by (A) isotype vs. PD-1 staining on CD3+ T cells in human PBMC activated with anti-CD3 + anti-CD28 for 48 hours, (B) PD-1 expression in isotype IgG1 vs. PD1AB-6-IgG1 treated PBMC (the detection anti-PD-1 antibody is not blocked by PD1AB-6), and (C) PD-1 expression on CD3+ T cells in human PBMC from 3 different donors, activated with anti-CD3 + anti-CD28 and three different concentrations of either isotype IgG1 or PD1AB-6-IgG1.

[0121] FIGS. 15A-15C show (A) PD1AB-6-IgG1, (B) PD1AB-6-4P, and (C) PD1AB-6-K3 binding to PD-1 antigen on Biacore<sup>®</sup> T200.

#### 4. DETAILED DESCRIPTION

[0122] Binding proteins, such as antibodies that bind to PD-1, including human and/or cynomolgus PD-1, are provided herein. In some embodiments, the binding proteins provided herein, such as antibodies that bind to human and/or cynomolgus PD-1, do not bind to rodent

PD-1. In certain embodiments, the PD-1 binding proteins, including antibodies disclosed herein, are agonists (e.g., can mimic the effect of PD-1 ligand and induce PD-1 signaling). In some embodiments, the binding proteins such as antibodies to PD-1 provided herein (i) bind to human and/or cynomolgus PD-1, (ii) do not compete for binding with PD-1 ligand (e.g., PD-L1 and/or PD-L2), and/or (iii) induce PD-1 signaling. In one embodiment, the PD-1 antibodies bind to human PD-1. In one embodiment, the PD-1 antibodies bind to cynomolgus PD-1. In one embodiment, the PD-1 antibodies bind to both human PD-1 and cynomolgus PD-1. In some embodiments, the PD-1 antibodies do not compete with PD-L1 for binding to PD-1. In other embodiments, the PD-1 antibodies do not compete with PD-L2 for binding to PD-1. In yet other embodiments, the PD-1 antibodies do not compete with either PD-L1 or PD-L2 for binding to PD-1. In other embodiments, the PD-1 antibodies induce PD-1 signaling. In specific embodiments, the PD-1 antibodies provided herein bind to both human PD-1 and cynomolgus PD-1, do not compete for binding to PD-1 with either PD-L1 or PD-L2, and induce PD-1 signaling. In some embodiments, the binding, competition, and/or signaling is assayed *in vitro*, e.g., in a cell-based assay. In other embodiments, the binding, competition, and/or signaling is assayed *ex vivo*, e.g., in a T cell function assay. In other embodiments, the binding, competition, and/or signaling is assayed using a sample from a subject (e.g., a human subject). In certain embodiments, assays include (1) a human or cynomolgus PBMC assay (see, e.g., Examples 5.2.1 and 5.2.2); (2) a human whole blood sample assay (see, e.g., Example 5.2.1). In certain embodiments, binding proteins, such as anti-PD-1 antibodies, as described herein, exhibit activities that are consistent with the natural biological function of PD-L1 and/or PD-L2. In some embodiments, the activities are exhibited *in vitro*. In other embodiments, the activities are exhibited *ex vivo*.

**[0123]** In specific embodiments, the binding proteins, such as antibodies that bind to PD-1, provided herein share the common feature of competing with each other for the binding of PD-1. This competitive inhibition can indicate that each antibody binds to the same region of PD-1 (e.g., the same epitope), thereby asserting similar effects. In certain embodiments, anti-PD-1 antibodies provided herein include humanized anti-PD-1 antibodies, such as those derived from or based on antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, and/or PD1AB-6. In other embodiments, anti-PD-1 antibodies provided herein compete for binding with an antibody derived from or based on PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5,

and/or PD1AB-6. In some embodiments, the anti-PD-1 antibodies have CDR sequences as described in **Tables 1-2**. In certain embodiments, the anti-PD-1 antibodies bind to a specific domain or epitope of human PD-1 (e.g., residues 100-105; *see Example 5.1.4*). Moreover, such binding can be largely attributed to particular amino acid residues within the region (e.g., G103 and R104; *see Example 5.1.4*), which comprise the epitope recognized by the anti-PD-1 antibodies provided herein. Taken together, the results described herein demonstrate that the effects observed for an anti-PD-1 antibody that is derived from or based on PD1AB-6, including an antibody having one or more CDRs described in **Tables 1-2**, can be extrapolated to other anti-PD-1 antibodies provided herein having the same or similar epitope specificity (e.g., the same or similar CDRs). For example, the activities of antibodies as shown in Examples 5.1.2-3, 5.1.7-10, 5.2.1-3, and 5.3.1, for an exemplary humanized anti-PD-1 antibody, are representative of the activities and effects of the anti-PD-1 antibodies provided herein.

**[0124]** In some embodiments of the present disclosure, the binding proteins such as anti-PD-1 antibodies may comprise immunoglobulin variable regions which comprise one or more CDRs as described in **Tables 1-2**. In such binding proteins (e.g., anti-PD-1 antibodies), the CDRs may be joined with one or more scaffold regions or framework regions (FRs), which orient(s) the CDR(s) such that the proper antigen binding properties of the CDR(s) is achieved. Such binding proteins, including anti-PD-1 antibodies as described herein, can induce PD-1 signaling.

#### **4.1 General Techniques**

**[0125]** Techniques and procedures described or referenced herein include those that are generally well understood and/or commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized methodologies described in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (3d ed. 2001); Current Protocols in Molecular Biology (Ausubel *et al.* eds., 2003); Therapeutic Monoclonal Antibodies: From Bench to Clinic (An ed. 2009); Monoclonal Antibodies: Methods and Protocols (Albitar ed. 2010); and Antibody Engineering Vols 1 and 2 (Kontermann and Dübel eds., 2d ed. 2010).

#### **4.2 Terminology**

**[0126]** Unless described otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art. For purposes of interpreting this specification, the following description of terms will apply and whenever

appropriate, terms used in the singular will also include the plural and vice versa. All patents, applications, published applications, and other publications are incorporated by reference in their entirety. In the event that any description of terms set forth conflicts with any document incorporated herein by reference, the description of term set forth below shall control.

**[0127]** The terms “Programmed Death 1,” “Programmed Cell Death 1,” “Protein PD-1,” “PD-1,” “PD-1 polypeptide,” or “PD1” encompasses a polypeptide (“polypeptide” and “protein” are used interchangeably herein), including any native polypeptide, from any vertebrate source, including mammals such as primates (*e.g.*, humans and cynomolgus monkeys (*cynomolgus*)), dogs, and rodents (*e.g.*, mice and rats), unless otherwise indicated. In certain embodiments, the terms include “related PD-1 polypeptides,” including SNP variants thereof. The term “PD-1” also encompasses “full-length,” unprocessed PD-1 as well as any form of PD-1 that results from processing in the cell. In some embodiments, the PD1 has an amino acid sequence of SEQ ID NO:43. GenBank™ accession number U64863 provides another exemplary human PD-1 nucleic acid sequence.

**[0128]** “Related PD-1 polypeptides” include allelic variants (*e.g.*, SNP variants); splice variants; fragments; derivatives; substitution, deletion, and insertion variants; fusion polypeptides; and interspecies homologs, which can retain PD-1 activity. As those skilled in the art will appreciate, an anti-PD-1 antibody provided herein can bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 antigen, and/or a PD-1 epitope. An “epitope” may be part of a larger PD-1 antigen, which may be part of a larger PD-1 polypeptide fragment, which, in turn, may be part of a larger PD-1 polypeptide. PD-1 may exist in a native or denatured form. PD-1 polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. Orthologs to the PD-1 polypeptide are also well known in the art.

**[0129]** A PD-1 polypeptide “extracellular domain” or “ECD” refers to a form of the PD-1 polypeptide that is essentially free of the transmembrane and cytoplasmic domains. For example, a PD-1 polypeptide ECD may have less than 1% of such transmembrane and/or cytoplasmic domains and can have less than 0.5% of such domains.

**[0130]** The terms “PD1AB-6-IgG1,” “PD1AB-6 IgG1,” “PD1AB-6\_IgG1,” “IgG1\_PD1AB-6,” and “IgG1-PD1AB-6” are used interchangeably, and refer to the antibody PD1AB-6 having an IgG1 Fc region. In certain embodiments, the antibody PD1AB-6 comprises

a light chain amino acid sequence of LC\_PD1AB-6-IgG1 (SEQ ID NO:31) and a heavy chain amino acid sequence of HC\_PD1AB-6-IgG1 (SEQ ID NO:32), *e.g.*, as shown in **FIG. 4**.

**[0131]** The terms “PD1AB-6-K3,” “PD1AB-6-IgG1-K322A,” “PD1AB-6-K322A,” “IgG1\_PD1AB-6\_K322A,” “IgG1\_PD1AB-6\_K3,” “IgG1-PD1AB-6-K322A,” and “IgG1-PD1AB-6-K3” are used interchangeably and refer to the PD1AB-6 variant having a K322A substitution in the IgG1 Fc region. In certain embodiments, the PD1AB-6 variant has a heavy chain amino acid sequence of HC\_PD1AB-6-IgG1-K322A (SEQ ID NO:33), *e.g.*, as shown in **FIG. 4**.

**[0132]** The terms “PD1AB-6-4P,” “IgG4P\_PD1AB-6,” “IgG4P-PD1AB-6,” “PD1AB-6\_IgG4P,” and “PD1AB-6-IgG4P” are used interchangeably and refer to the PD1AB-6 variant having an IgG4P Fc region. In certain embodiments, the PD-1 antibody variant has a heavy chain amino acid sequence of HC\_PD1AB-6-IgG4P (SEQ ID NO:34), *e.g.*, as shown in **FIG. 4**.

**[0133]** The terms “PD1AB-6-4PE,” “IgG4PE\_PD1AB-6,” “IgG4PE-PD1AB-6,” and “PD1AB-6\_IgG4PE,” and “PD1AB-6-IgG4PE” are used interchangeably and refer to the PD1AB-6 variant having an IgG4PE heavy chain amino acid sequence as HC\_PD1AB-6-IgG4PE (SEQ ID NO:35).

**[0134]** The term “PD-1 ligand” refers to a molecule that binds to PD-1, *e.g.*, *in vivo* or *in vitro*. Non-limiting examples of PD-1 ligand include naturally occurring ligands, *e.g.*, PD-1 ligand 1 (PD-L1, also known as B7-H1 or CD274) and PD-1 ligand 2 (PD-L2, also known as B7-DC or CD273), and artificially generated ligands.

**[0135]** The terms “PD-L1” and “PDL-1” are used interchangeably herein and refer to PD-1 ligand 1 (also known as B7-H1 or CD274).

**[0136]** The terms “PD-1 activity,” “PD-1 signaling,” and “PD-1 ligand-like signaling” when applied to a binding protein such as an antibody that binds to PD-1 of the present disclosure, means that the binding protein (*e.g.*, antibody) mimics or modulates a biological effect induced by the binding of PD-1 ligand, and induces a biological response that otherwise would result from PD-1 ligand binding to PD-1, *e.g.*, *in vivo* or *in vitro*. In assessing the binding specificity of anti-PD-1 antibody, for example, an antibody or fragment thereof that binds to PD-1 (*e.g.*, human PD-1), the antibody is deemed to induce a biological response when the response is equal to or greater than 5%, such as equal to or greater than 10%, 15%, 20%, 25%, 30%, 35%, 40%,

45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100%, 125%, 150%, 175%, or 200% of the activity of a wild type PD-1 ligand standard. In one embodiment, the anti-PD-1 antibody or the PD-1 ligand is immobilized (for example, on a plastic surface or bead). In certain embodiments, the antibody has the following properties: exhibits an efficacy level of equal to or more than 5% of a PD-1 ligand standard, with an EC<sub>50</sub> of equal to or less than 100 nM, *e.g.*, 90 nM, 80 nM, 70 nM, 60 nM, 50 nM, 40 nM, 30 nM, 20 nM, 10 nM, 5 nM, 2 nM, 1 nM, 0.5 nM, 0.2 nM, or 0.1 nM in (1) human or cynomolgus PBMC assay (*see, e.g.*, Examples 4.2.1 and 4.2.2); or (2) human whole blood sample assay (*see, e.g.*, Example 4.2.1).

**[0137]** The term “binding protein” refers to a protein comprising a portion (*e.g.*, one or more binding regions such as CDRs) that binds to PD-1, including human and/or cynomolgus PD-1 and, optionally, a scaffold or framework portion (*e.g.*, one or more scaffold or framework regions) that allows the binding portion to adopt a conformation that promotes binding of the binding protein to a PD-1 polypeptide, fragment, or epitope. Examples of such binding proteins include antibodies, such as a human antibody, a humanized antibody, a chimeric antibody, a recombinant antibody, a single chain antibody, a diabody, a triabody, a tetrabody, a Fab fragment, a F(ab')<sub>2</sub> fragment, an IgD antibody, an IgE antibody, an IgM antibody, an IgG1 antibody, an IgG2 antibody, an IgG3 antibody, or an IgG4 antibody, and fragments thereof. The binding protein can comprise, for example, an alternative protein scaffold or artificial scaffold with grafted CDRs or CDR derivatives. Such scaffolds include, but are not limited to, antibody-derived scaffolds comprising mutations introduced to, for example, stabilize the three-dimensional structure of the binding protein as well as wholly synthetic scaffolds comprising, for example, a biocompatible polymer. *See, e.g.*, Korndorfer *et al.*, 2003, Proteins: Structure, Function, and Bioinformatics 53(1):121-29; and Roque *et al.*, 2004, Biotechnol. Prog. 20:639-54. In addition, peptide antibody mimetics (“PAMs”) can be used, as well as scaffolds based on antibody mimetics utilizing fibronectin components as a scaffold. In the context of the present disclosure, a binding protein is said to specifically bind or selectively bind to PD-1, for example, when the dissociation constant (K<sub>D</sub>) is  $\leq 10^{-7}$  M. In some embodiments, the binding proteins (*e.g.*, antibodies) may specifically bind to PD-1 with a K<sub>D</sub> of from about 10<sup>-7</sup> M to about 10<sup>-12</sup> M. In certain embodiments, the binding protein (*e.g.*, antibody) may specifically bind to PD-1 with high affinity when the K<sub>D</sub> is  $\leq 10^{-8}$  M or K<sub>D</sub> is  $\leq 10^{-9}$  M. In one embodiment, the binding proteins (*e.g.*, antibodies) may specifically bind to purified human PD-1 with a K<sub>D</sub> of

from  $1 \times 10^{-9}$  M to  $10 \times 10^{-9}$  M as measured by Biacore<sup>®</sup>. In another embodiment, the binding proteins (e.g., antibodies) may specifically bind to purified human PD-1 with a  $K_D$  of from  $0.1 \times 10^{-9}$  M to  $1 \times 10^{-9}$  M as measured by KinExA<sup>TM</sup> (Sapidyne, Boise, ID). In yet another embodiment, the binding proteins (e.g., antibodies) specifically bind to human PD-1 expressed on cells with a  $K_D$  of from  $0.1 \times 10^{-9}$  M to  $10 \times 10^{-9}$  M. In certain embodiments, the binding proteins (e.g., antibodies) specifically bind to human PD-1 expressed on cells with a  $K_D$  of from  $0.1 \times 10^{-9}$  M to  $1 \times 10^{-9}$  M. In some embodiments, the binding proteins (e.g., antibodies) specifically bind to human PD-1 expressed on cells with a  $K_D$  of  $1 \times 10^{-9}$  M to  $10 \times 10^{-9}$  M. In certain embodiments, the binding proteins (e.g., antibodies) specifically bind to human PD-1 expressed on cells with a  $K_D$  of about  $0.1 \times 10^{-9}$  M, about  $0.5 \times 10^{-9}$  M, about  $1 \times 10^{-9}$  M, about  $5 \times 10^{-9}$  M, about  $10 \times 10^{-9}$  M, or any range or interval thereof. In still another embodiment, the binding proteins (e.g., antibodies) may specifically bind to cynomolgus PD-1 expressed on cells with a  $K_D$  of  $0.1 \times 10^{-9}$  M to  $10 \times 10^{-9}$  M. In certain embodiments, the binding proteins (e.g., antibodies) specifically bind to cynomolgus PD-1 expressed on cells with a  $K_D$  of from  $0.1 \times 10^{-9}$  M to  $1 \times 10^{-9}$  M. In some embodiments, the binding proteins (e.g., antibodies) specifically bind to cynomolgus PD-1 expressed on cells with a  $K_D$  of  $1 \times 10^{-9}$  M to  $10 \times 10^{-9}$  M. In certain embodiments, the binding proteins (e.g., antibodies) specifically bind to cynomolgus PD-1 expressed on cells with a  $K_D$  of about  $0.1 \times 10^{-9}$  M, about  $0.5 \times 10^{-9}$  M, about  $1 \times 10^{-9}$  M, about  $5 \times 10^{-9}$  M, about  $10 \times 10^{-9}$  M, or any range or interval thereof.

**[0138]** The term “antibody,” “immunoglobulin,” or “Ig” is used interchangeably herein, and is used in the broadest sense and specifically encompasses, for example, individual anti-PD-1 monoclonal antibodies (including agonist, antagonist, neutralizing antibodies, full length or intact monoclonal antibodies), anti-PD-1 antibody compositions with polyepitopic or monoepitopic specificity, polyclonal or monovalent antibodies, multivalent antibodies, multispecific antibodies (e.g., bispecific antibodies so long as they exhibit the desired biological activity), formed from at least two intact antibodies, single chain anti-PD-1 antibodies, and fragments of anti-PD-1 antibodies, as described below. An antibody can be human, humanized, chimeric and/or affinity matured, as well as an antibody from other species, for example, mouse and rabbit, *etc.* The term “antibody” is intended to include a polypeptide product of B cells within the immunoglobulin class of polypeptides that is able to bind to a specific molecular antigen and is composed of two identical pairs of polypeptide chains, wherein each pair has one

heavy chain (about 50-70 kDa) and one light chain (about 25 kDa), each amino-terminal portion of each chain includes a variable region of about 100 to about 130 or more amino acids, and each carboxy-terminal portion of each chain includes a constant region. *See, e.g., Antibody Engineering (Borrebaeck ed., 2d ed. 1995); and Kuby, Immunology (3d ed. 1997).* In specific embodiments, the specific molecular antigen can be bound by an antibody provided herein, including a PD-1 polypeptide, a PD-1 fragment, or a PD-1 epitope. Antibodies also include, but are not limited to, synthetic antibodies, recombinantly produced antibodies, camelized antibodies, intrabodies, anti-idiotypic (anti-Id) antibodies, and functional fragments (*e.g.*, antigen-binding fragments such as PD-1-binding fragments) of any of the above, which refers to a portion of an antibody heavy or light chain polypeptide that retains some or all of the binding activity of the antibody from which the fragment was derived. Non-limiting examples of functional fragments (*e.g.*, antigen-binding fragments such as PD-1-binding fragments) include single-chain Fvs (scFv) (*e.g.*, including monospecific, bispecific, *etc.*), Fab fragments, F(ab') fragments, F(ab)<sub>2</sub> fragments, F(ab')<sub>2</sub> fragments, disulfide-linked Fvs (dsFv), Fd fragments, Fv fragments, diabody, triabody, tetrabody, and minibody. In particular, antibodies provided herein include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, for example, antigen-binding domains or molecules that contain an antigen-binding site that binds to a PD-1 antigen (*e.g.*, one or more CDRs of an anti-PD-1 antibody). Such antibody fragments can be found in, for example, Harlow and Lane, Antibodies: A Laboratory Manual (1989); Mol. Biology and Biotechnology: A Comprehensive Desk Reference (Myers ed., 1995); Huston *et al.*, 1993, Cell Biophysics 22:189-224; Plückthun and Skerra, 1989, Meth. Enzymol. 178:497-515; and Day, Advanced Immunochemistry (2d ed. 1990). The antibodies provided herein can be of any class (*e.g.*, IgG, IgE, IgM, IgD, and IgA) or any subclass (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2) of immunoglobulin molecule. Anti-PD-1 antibodies may be agonistic antibodies or antagonistic antibodies. Provided herein are agonistic antibodies to PD-1, including antibodies that induce PD-1 signaling. In specific embodiments, agonistic antibodies to PD-1 do not compete for the binding of PD-L1 and/or PD-L2 to PD-1.

**[0139]** The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *e.g.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts, and each monoclonal antibody will typically recognize a single epitope on the

antigen. In specific embodiments, a “monoclonal antibody,” as used herein, is an antibody produced by a single hybridoma or other cell, wherein the antibody binds to only a PD-1 epitope as determined, for example, by ELISA or other antigen-binding or competitive binding assay known in the art. The term “monoclonal” is not limited to any particular method for making the antibody. For example, the monoclonal antibodies useful in the present disclosure may be prepared by the hybridoma methodology first described by Kohler *et al.*, 1975, *Nature* 256:495, or may be made using recombinant DNA methods in bacterial or eukaryotic animal or plant cells (see, e.g., U.S. Pat. No. 4,816,567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, 1991, *Nature* 352:624-28 and Marks *et al.*, 1991, *J. Mol. Biol.* 222:581-97, for example. Other methods for the preparation of clonal cell lines and of monoclonal antibodies expressed thereby are well known in the art. *See, e.g., Short Protocols in Molecular Biology* (Ausubel *et al.* eds., 5th ed. 2002). Exemplary methods of producing monoclonal antibodies are provided in the Examples herein.

**[0140]** “Polyclonal antibodies” as used herein refer to an antibody population generated in an immunogenic response to a protein having many epitopes and thus includes a variety of different antibodies directed to the same or different epitopes within the protein. Methods for producing polyclonal antibodies are known in the art (*See, e.g., Short Protocols in Molecular Biology* (Ausubel *et al.* eds., 5th ed. 2002)).

**[0141]** In the context of a peptide or polypeptide, the term “fragment” as used herein refers to a peptide or polypeptide that comprises less than the full length amino acid sequence. Such a fragment may arise, for example, from a truncation at the amino terminus, a truncation at the carboxy terminus, and/or an internal deletion of a residue(s) from the amino acid sequence. Fragments may, for example, result from alternative RNA splicing or from *in vivo* protease activity. In certain embodiments, PD-1 fragments or anti-PD-1 antibody fragments include polypeptides comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 30 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino residues, at least 70 contiguous amino acid residues, at least 80 contiguous amino acid residues, at least 90 contiguous amino

acid residues, at least contiguous 100 amino acid residues, at least 125 contiguous amino acid residues, at least 150 contiguous amino acid residues, at least 175 contiguous amino acid residues, at least 200 contiguous amino acid residues, at least 250, at least 300, at least 350, at least 400, at least 450, at least 500, at least 550, at least 600, at least 650, at least 700, at least 750, at least 800, at least 850, at least 900, or at least 950 contiguous amino acid residues of the amino acid sequence of a PD-1 polypeptide or an anti-PD-1 antibody. In a specific embodiment, a fragment of a PD-1 polypeptide or an anti-PD-1 antibody retains at least 1, at least 2, at least 3, or more functions of the polypeptide or antibody.

**[0142]** An “antigen” is a predetermined antigen to which an antibody can selectively bind. A target antigen may be a polypeptide, carbohydrate, nucleic acid, lipid, hapten, or other naturally occurring or synthetic compound. In some embodiments, the target antigen is a polypeptide.

**[0143]** The terms “antigen-binding fragment,” “antigen-binding domain,” “antigen-binding region,” and similar terms refer to that portion of an antibody, which comprises the amino acid residues that interact with an antigen and confer on the binding agent its specificity and affinity for the antigen (*e.g.*, the CDRs).

**[0144]** An “epitope” is the site on the surface of an antigen molecule to which a single antibody molecule binds, such as a localized region on the surface of an antigen, such as a PD-1 polypeptide or a PD-1 polypeptide fragment, that is capable of being bound to one or more antigen binding regions of an antibody, and that has antigenic or immunogenic activity in an animal, such as a mammal (*e.g.*, a human), that is capable of eliciting an immune response. An epitope having immunogenic activity is a portion of a polypeptide that elicits an antibody response in an animal. An epitope having antigenic activity is a portion of a polypeptide to which an antibody binds as determined by any method well known in the art, including, for example, by an immunoassay. Antigenic epitopes need not necessarily be immunogenic. Epitopes often consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and have specific three dimensional structural characteristics as well as specific charge characteristics. Antibody epitopes may be linear epitopes or conformational epitopes. Linear epitopes are formed by a continuous sequence of amino acids in a protein. Conformational epitopes are formed of amino acids that are discontinuous in the protein sequence, but which are brought together upon folding of the protein into its three-dimensional structure. Induced epitopes are formed when the three dimensional structure of the protein is in

an altered conformation, such as following activation or binding of another protein or ligand. In certain embodiments, a PD-1 epitope is a three-dimensional surface feature of a PD-1 polypeptide. In other embodiments, a PD-1 epitope is linear feature of a PD-1 polypeptide. Generally an antigen has several or many different epitopes and may react with many different antibodies.

**[0145]** An antibody binds “an epitope,” “essentially the same epitope,” or “the same epitope” as a reference antibody, when the two antibodies recognize identical, overlapping, or adjacent epitopes in a three-dimensional space. The most widely used and rapid methods for determining whether two antibodies bind to identical, overlapping, or adjacent epitopes in a three-dimensional space are competition assays, which can be configured in a number of different formats, for example, using either labeled antigen or labeled antibody. In some assays, the antigen is immobilized on a 96-well plate, or expressed on a cell surface, and the ability of unlabeled antibodies to block the binding of labeled antibodies is measured using radioactive, fluorescent, or enzyme labels.

**[0146]** “Epitope mapping” is the process of identifying the binding sites, or epitopes, of antibodies on their target antigens. “Epitope binning” is the process of grouping antibodies based on the epitopes they recognize. More particularly, epitope binning comprises methods and systems for discriminating the epitope recognition properties of different antibodies, using competition assays combined with computational processes for clustering antibodies based on their epitope recognition properties and identifying antibodies having distinct binding specificities.

**[0147]** The terms “binds” or “binding” refer to an interaction between molecules including, for example, to form a complex. Interactions can be, for example, non-covalent interactions including hydrogen bonds, ionic bonds, hydrophobic interactions, and/or van der Waals interactions. A complex can also include the binding of two or more molecules held together by covalent or non-covalent bonds, interactions, or forces. The strength of the total non-covalent interactions between a single antigen-binding site on an antibody and a single epitope of a target molecule, such as PD-1, is the affinity of the antibody or functional fragment for that epitope. The ratio of dissociation rate ( $k_{off}$ ) to association rate ( $k_{on}$ ) of an antibody to a monovalent antigen ( $k_{off}/k_{on}$ ) is the dissociation constant  $K_D$ , which is inversely related to affinity. The lower the  $K_D$  value, the higher the affinity of the antibody. The value of  $K_D$  varies for different

complexes of antibody and antigen and depends on both  $k_{on}$  and  $k_{off}$ . The dissociation constant  $K_D$  for an antibody provided herein can be determined using any method provided herein or any other method well known to those skilled in the art. The affinity at one binding site does not always reflect the true strength of the interaction between an antibody and an antigen. When complex antigens containing multiple, repeating antigenic determinants, such as a polyvalent PD-1, come in contact with antibodies containing multiple binding sites, the interaction of antibody with antigen at one site will increase the probability of a reaction at a second site. The strength of such multiple interactions between a multivalent antibody and antigen is called the avidity. The avidity of an antibody can be a better measure of its binding capacity than is the affinity of its individual binding sites. For example, high avidity can compensate for low affinity as is sometimes found for pentameric IgM antibodies, which can have a lower affinity than IgG, but the high avidity of IgM, resulting from its multivalence, enables it to bind antigen effectively.

**[0148]** The terms “antibodies that specifically bind to PD-1,” “antibodies that specifically bind to a PD-1 epitope,” and analogous terms are also used interchangeably herein and refer to antibodies that specifically bind to a PD-1 polypeptide, such as a PD-1 antigen, or fragment, or epitope (e.g., human PD-1 such as a human PD-1 polypeptide, antigen, or epitope). An antibody that specifically binds to PD-1 (e.g., human PD-1) may bind to the extracellular domain or a peptide derived from the extracellular domain of PD-1. An antibody that specifically binds to a PD-1 antigen (e.g., human PD-1) may be cross-reactive with related antigens (e.g., cynomolgus PD-1). In certain embodiments, an antibody that specifically binds to a PD-1 antigen does not cross-react with other antigens. An antibody that specifically binds to a PD-1 antigen can be identified, for example, by immunoassays, Biacore<sup>®</sup>, or other techniques known to those of skill in the art. An antibody binds specifically to a PD-1 antigen when it binds to a PD-1 antigen with higher affinity than to any cross-reactive antigen as determined using experimental techniques, such as radioimmunoassays (RIA) and enzyme linked immunosorbent assays (ELISAs).

Typically a specific or selective reaction will be at least twice background signal or noise and may be more than 10 times background. See, e.g., Fundamental Immunology 332-36 (Paul ed., 2d ed. 1989) for a discussion regarding antibody specificity. An antibody which “binds an antigen of interest” (e.g., a target antigen such as PD-1) is one that binds the antigen with sufficient affinity such that the antibody is useful as a therapeutic agent in targeting a cell or tissue expressing the antigen, and does not significantly cross-react with other proteins. In such

embodiments, the extent of binding of the antibody to a “non-target” protein will be less than about 10% of the binding of the antibody to its particular target protein, for example, as determined by fluorescence activated cell sorting (FACS) analysis or RIA. With regard to the binding of an antibody to a target molecule, the term “specific binding,” “specifically binds to,” or “is specific for” a particular polypeptide or an epitope on a particular polypeptide target means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target, for example, an excess of non-labeled target. In this case, specific binding is indicated if the binding of the labeled target to a probe is competitively inhibited by excess unlabeled target. The term “anti-PD-1 antibody” or “an antibody that binds to PD-1” includes an antibody that is capable of binding PD-1 with sufficient affinity such that the antibody is useful, for example, as a diagnostic agent in targeting PD-1. The term “specific binding,” “specifically binds to,” or “is specific for” a particular polypeptide or an epitope on a particular polypeptide target as used herein refers to binding where a molecule binds to a particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope. In certain embodiments, an antibody that binds to PD-1 has a dissociation constant ( $K_D$ ) of less than or equal to 10 nM, 5 nM, 4 nM, 3 nM, 2 nM, 1 nM, 0.9 nM, 0.8 nM, 0.7 nM, 0.6 nM, 0.5 nM, 0.4 nM, 0.3 nM, 0.2 nM, or 0.1 nM. In certain embodiments, anti-PD-1 antibody binds to an epitope of PD-1 that is conserved among PD-1 from different species (*e.g.*, between human and cynomolgus PD-1).

**[0149]** The term “compete” when used in the context of anti-PD-1 antibodies (*e.g.*, agonistic antibodies and binding proteins that bind to PD-1 and compete for the same epitope or binding site on a target) means competition as determined by an assay in which the antibody (or binding fragment) thereof under study prevents or inhibits the specific binding of a reference molecule (*e.g.*, a reference ligand or reference antigen-binding protein, such as a reference antibody) to a common antigen (*e.g.*, PD-1 or a fragment thereof). Numerous types of competitive binding assays can be used to determine if a test antibody competes with a reference antibody for binding to PD-1 (*e.g.*, human PD-1). Examples of assays that can be employed include solid phase direct or indirect RIA, solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition

assay (see, e.g., Stahli *et al.*, 1983, Methods in Enzymology 9:242-53), solid phase direct biotin-avidin EIA (see, e.g., Kirkland *et al.*, 1986, J. Immunol. 137:3614-19), solid phase direct labeled assay, solid phase direct labeled sandwich assay (see, e.g., Harlow and Lane, Antibodies, A Laboratory Manual (1988)), solid phase direct label RIA using I-125 label (see, e.g., Morel *et al.*, 1988, Mol. Immunol. 25:7-15), and direct labeled RIA (Moldenhauer *et al.*, 1990, Scand. J. Immunol. 32:77-82). Typically, such an assay involves the use of a purified antigen (e.g., PD-1 such as human PD-1) bound to a solid surface, or cells bearing either of an unlabelled test antigen-binding protein (e.g., test anti-PD-1 antibody) or a labeled reference antigen-binding protein (e.g., reference anti-PD-1 antibody). Competitive inhibition may be measured by determining the amount of label bound to the solid surface or cells in the presence of the test antigen-binding protein. Usually the test antigen-binding protein is present in excess.

Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and/or antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference for antibodies steric hindrance to occur. Additional details regarding methods for determining competitive binding are described herein. Usually, when a competing antibody protein is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 30%, for example 40%, 45%, 50%, 55%, 60%, 65%, 70%, or 75%. In some instance, binding is inhibited by at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or more.

**[0150]** An “isolated” antibody is substantially free of cellular material or other contaminating proteins from the cell or tissue source and/or other contaminant components from which the antibody is derived, or substantially free of chemical precursors or other chemicals when chemically synthesized. The language “substantially free of cellular material” includes preparations of an antibody in which the antibody is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, an antibody that is substantially free of cellular material includes preparations of antibody having less than about 30%, 25%, 20%, 15%, 10%, 5%, or 1% (by dry weight) of heterologous protein (also referred to herein as a “contaminating protein”). In certain embodiments, when the antibody is recombinantly produced, it is substantially free of culture medium, e.g., culture medium represents less than about 20%, 15%, 10%, 5%, or 1% of the volume of the protein preparation. In certain embodiments, when the antibody is produced by chemical synthesis, it is substantially free of

chemical precursors or other chemicals, for example, it is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. Accordingly such preparations of the antibody have less than about 30%, 25%, 20%, 15%, 10%, 5%, or 1% (by dry weight) of chemical precursors or compounds other than the antibody of interest. Contaminant components can also include, but are not limited to, materials that would interfere with therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In certain embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method (Lowry *et al.*, 1951, *J. Bio. Chem.* 193: 265-75), such as 96%, 97%, 98%, or 99%, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step. In specific embodiments, antibodies provided herein are isolated.

**[0151]** A 4-chain antibody unit is a heterotetrameric glycoprotein composed of two identical light (L) chains and two identical heavy (H) chains. In the case of IgGs, the 4-chain unit is generally about 150,000 daltons. Each L chain is linked to an H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. Each H and L chain also has regularly spaced intrachain disulfide bridges. Each H chain has at the N-terminus, a variable domain (VH) followed by three constant domains (CH) for each of the  $\alpha$  and  $\gamma$  chains and four CH domains for  $\mu$  and  $\epsilon$  isotypes. Each L chain has at the N-terminus, a variable domain (VL) followed by a constant domain (CL) at its other end. The VL is aligned with the VH, and the CL is aligned with the first constant domain of the heavy chain (CH1). Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains. The pairing of a VH and VL together forms a single antigen-binding site. For the structure and properties of the different classes of antibodies, see, for example, Basic and Clinical Immunology 71 (Stites *et al.* eds., 8th ed. 1994).

**[0152]** The term "heavy chain" when used in reference to an antibody refers to a polypeptide chain of about 50-70 kDa, wherein the amino-terminal portion includes a variable region of about 120 to 130 or more amino acids, and a carboxy-terminal portion includes a constant

region. The constant region can be one of five distinct types, (e.g., isotypes) referred to as alpha ( $\alpha$ ), delta ( $\delta$ ), epsilon ( $\epsilon$ ), gamma ( $\gamma$ ), and mu ( $\mu$ ), based on the amino acid sequence of the heavy chain constant region. The distinct heavy chains differ in size:  $\alpha$ ,  $\delta$ , and  $\gamma$  contain approximately 450 amino acids, while  $\mu$  and  $\epsilon$  contain approximately 550 amino acids. When combined with a light chain, these distinct types of heavy chains give rise to five well known classes (e.g., isotypes) of antibodies, IgA, IgD, IgE, IgG, and IgM, respectively, including four subclasses of IgG, namely IgG1, IgG2, IgG3, and IgG4. A heavy chain can be a human heavy chain.

**[0153]** The term “light chain” when used in reference to an antibody refers to a polypeptide chain of about 25 kDa, wherein the amino-terminal portion includes a variable region of about 100 to about 110 or more amino acids, and a carboxy-terminal portion includes a constant region. The approximate length of a light chain is 211 to 217 amino acids. There are two distinct types, referred to as kappa ( $\kappa$ ) or lambda ( $\lambda$ ) based on the amino acid sequence of the constant domains. Light chain amino acid sequences are well known in the art. A light chain can be a human light chain.

**[0154]** The term “variable region,” “variable domain,” “V region,” or “V domain” refers to a portion of the light or heavy chains of an antibody that is generally located at the amino-terminal of the light or heavy chain and has a length of about 120 to 130 amino acids in the heavy chain and about 100 to 110 amino acids in the light chain, and are used in the binding and specificity of each particular antibody for its particular antigen. The variable region of the heavy chain may be referred to as “VH.” The variable region of the light chain may be referred to as “VL.” The term “variable” refers to the fact that certain segments of the variable regions differ extensively in sequence among antibodies. The V region mediates antigen binding and defines specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the 110-amino acid span of the variable regions. Instead, the V regions consist of less variable (e.g., relatively invariant) stretches called framework regions (FRs) of about 15-30 amino acids separated by shorter regions of greater variability (e.g., extreme variability) called “hypervariable regions” that are each about 9-12 amino acids long. The variable regions of heavy and light chains each comprise four FRs, largely adopting a  $\beta$  sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases form part of, the  $\beta$  sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the

formation of the antigen-binding site of antibodies (see, e.g., Kabat *et al.*, Sequences of Proteins of Immunological Interest (5th ed. 1991)). The constant regions are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC) and complement dependent cytotoxicity (CDC). The variable regions differ extensively in sequence between different antibodies. In specific embodiments, the variable region is a human variable region.

**[0155]** The term “variable region residue numbering as in Kabat” or “amino acid position numbering as in Kabat”, and variations thereof, refer to the numbering system used for heavy chain variable regions or light chain variable regions of the compilation of antibodies in Kabat *et al.*, *supra*. Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, an FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 and three inserted residues (e.g., residues 82a, 82b, and 82c, *etc.* according to Kabat) after residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a “standard” Kabat numbered sequence. The Kabat numbering system is generally used when referring to a residue in the variable domain (approximately residues 1-107 of the light chain and residues 1-113 of the heavy chain) (e.g., Kabat *et al.*, *supra*). The “EU numbering system” or “EU index” is generally used when referring to a residue in an immunoglobulin heavy chain constant region (e.g., the EU index reported in Kabat *et al.*, *supra*). The “EU index as in Kabat” refers to the residue numbering of the human IgG 1 EU antibody. Other numbering systems have been described, for example, by AbM, Chothia, Contact, IMGT, and AHon.

**[0156]** A “CDR” refers to one of three hypervariable regions (H1, H2 or H3) within the non-framework region of the immunoglobulin (Ig or antibody) VH  $\beta$ -sheet framework, or one of three hypervariable regions (L1, L2 or L3) within the non-framework region of the antibody VL  $\beta$ -sheet framework. Accordingly, CDRs are variable region sequences interspersed within the framework region sequences. CDR regions are well known to those skilled in the art and have been defined by, for example, Kabat as the regions of most hypervariability within the antibody variable (V) domains (Kabat *et al.*, 1997, J. Biol. Chem. 252:6609-16; Kabat, 1978, Adv. Prot. Chem. 32:1-75). CDR region sequences also have been defined structurally by Chothia as those

residues that are not part of the conserved  $\beta$ -sheet framework, and thus are able to adapt different conformations (Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-17). Both terminologies are well recognized in the art. CDR region sequences have also been defined by AbM, Contact, and IMGT. The positions of CDRs within a canonical antibody variable region have been determined by comparison of numerous structures (Al-Lazikani *et al.*, 1997, *J. Mol. Biol.* 273:927-48; Morea *et al.*, 2000, *Methods* 20:267-79). Because the number of residues within a hypervariable region varies in different antibodies, additional residues relative to the canonical positions are conventionally numbered with a, b, c and so forth next to the residue number in the canonical variable region numbering scheme (Al-Lazikani *et al.*, *supra*). Such nomenclature is similarly well known to those skilled in the art.

**[0157]** The term “hypervariable region,” “HVR,” or “HV,” when used herein refers to the regions of an antibody variable region that are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six hypervariable regions, three in the VH (H1, H2, H3) and three in the VL (L1, L2, L3). A number of hypervariable region delineations are in use and are encompassed herein. The Kabat Complementarity Determining Regions (CDRs) are based on sequence variability and are the most commonly used (*see, e.g.*, Kabat *et al.*, *supra*). Chothia refers instead to the location of the structural loops (*see, e.g.*, Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-17). The end of the Chothia CDR-H1 loop when numbered using the Kabat numbering convention varies between H32 and H34 depending on the length of the loop (this is because the Kabat numbering scheme places the insertions at H35A and H35B; if neither 35A nor 35B is present, the loop ends at 32; if only 35A is present, the loop ends at 33; if both 35A and 35B are present, the loop ends at 34). The AbM hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular’s AbM antibody modeling software (*see, e.g.*, Antibody Engineering Vol. 2 (Kontermann and Dübel eds., 2d ed. 2010)). The “contact” hypervariable regions are based on an analysis of the available complex crystal structures. The residues from each of these hypervariable regions or CDRs are noted below.

**[0158]** Recently, a universal numbering system has been developed and widely adopted, ImMunoGeneTics (IMGT) Information System<sup>®</sup> (Lafranc *et al.*, 2003, *Dev. Comp. Immunol.* 27(1):55-77). IMGT is an integrated information system specializing in immunoglobulins (IG), T cell receptors (TCR), and major histocompatibility complex (MHC) of human and other

vertebrates. Herein, the CDRs are referred to in terms of both the amino acid sequence and the location within the light or heavy chain. As the “location” of the CDRs within the structure of the immunoglobulin variable domain is conserved between species and present in structures called loops, by using numbering systems that align variable domain sequences according to structural features, CDR and framework residues are readily identified. This information can be used in grafting and replacement of CDR residues from immunoglobulins of one species into an acceptor framework from, typically, a human antibody. An additional numbering system (AHon) has been developed by Honegger and Plückthun, 2001, *J. Mol. Biol.* 309: 657-70. Correspondence between the numbering system, including, for example, the Kabat numbering and the IMGT unique numbering system, is well known to one skilled in the art (*see, e.g.*, Kabat, *supra*; Chothia and Lesk, *supra*; Martin, *supra*; Lefranc *et al.*, *supra*). In some embodiments, the CDRs are as defined by the IMGT numbering system. In other embodiments, the CDRs are as defined by the Kabat numbering system. In certain embodiments, the CDRs are as defined by the AbM numbering system. In other embodiments, the CDRs are as defined by the Chothia system. In yet other embodiments, the CDRs are as defined by the Contact numbering system.

	<u>IMGT</u>	<u>Kabat</u>	<u>AbM</u>	<u>Chothia</u>	<u>Contact</u>
V <sub>H</sub> CDR1	27-38	31-35	26-35	26-32	30-35
V <sub>H</sub> CDR2	56-65	50-65	50-58	53-55	47-58
V <sub>H</sub> CDR3	105-117	95-102	95-102	96-101	93-101
V <sub>L</sub> CDR1	27-38	24-34	24-34	26-32	30-36
V <sub>L</sub> CDR2	56-65	50-56	50-56	50-52	46-55
V <sub>L</sub> CDR3	105-117	89-97	89-97	91-96	89-96

**[0159]** Hypervariable regions may comprise “extended hypervariable regions” as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2), and 89-97 or 89-96 (L3) in the VL, and 26-35 or 26-35A (H1), 50-65 or 49-65 (H2), and 93-102, 94-102, or 95-102 (H3) in the VH. As used herein, the terms “HVR” and “CDR” are used interchangeably.

**[0160]** The term “constant region” or “constant domain” refers to a carboxy terminal portion of the light and heavy chain which is not directly involved in binding of the antibody to antigen but exhibits various effector function, such as interaction with the Fc receptor. The term refers to

the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable region, which contains the antigen binding site. The constant region may contain the CH1, CH2, and CH3 regions of the heavy chain and the CL region of the light chain.

**[0161]** The term “framework” or “FR” refers to those variable region residues flanking the CDRs. FR residues are present, for example, in chimeric, humanized, human, domain antibodies, diabodies, linear antibodies, and bispecific antibodies. FR residues are those variable domain residues other than the hypervariable region residues or CDR residues.

**[0162]** The term “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including, for example, native sequence Fc regions, recombinant Fc regions, and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is often defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue.

**[0163]** A “functional Fc region” possesses an “effector function” of a native sequence Fc region. Exemplary “effector functions” include C1q binding; CDC; Fc receptor binding; ADCC; phagocytosis; downregulation of cell surface receptors (*e.g.*, B cell receptor), *etc.* Such effector functions generally require the Fc region to be combined with a binding region or binding domain (*e.g.*, an antibody variable region or domain) and can be assessed using various assays as disclosed.

**[0164]** A “native sequence Fc region” comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature, and not manipulated, modified, and/or changed (*e.g.*, isolated, purified, selected, including or combining with other sequences such as variable region sequences) by a human. Native sequence human IgG1 Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region

as well as naturally occurring variants thereof. For example, a native human IgG1 Fc region amino acid sequence is provided below:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPAPELLGGPSVFLFPP  
KPKDTLMISRTPEVTCVVVDVSQEDPEVFKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL  
HQDWLNGKEYKCVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTQNQVSLTCLVKGFY  
PSDIAVEWESNGQPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT  
QKSLSLSPGK (SEQ ID NO:36, K322 emphasized).

An exemplary native human IgG4 Fc region sequence is provided below:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKTYTCNVDHKPSNTKVDKRVESKYGPPCPSCPAPEFLGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQD  
WLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:38, S228 and L235 emphasized).

**[0165]** A “variant Fc region” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification (e.g., substituting, addition, or deletion). In certain embodiments, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, for example, from about one to about ten amino acid substitutions, or from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of a parent polypeptide. The variant Fc region herein can possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, or at least about 90% homology therewith, for example, at least about 95% homology therewith. For example, a variant with one amino acid K change to A at 322 position in the human IgG1 Fc amino acid sequence, IgG1-K322A Fc region, is provided below:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPAPELLGGPSVFLFPP  
KPKDTLMISRTPEVTCVVVDVSQEDPEVFKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL  
HQDWLNGKEYKAVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTQNQVSLTCLVKGFY

PSDIAVEWESNGQPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:37, K322A substitution emphasized).

An exemplary variant with one amino acid S change to P at 228 position in the human IgG4 Fc amino acid sequence, IgG4P Fc region, is provided below:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKYTCNVVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:39, S228P substitution emphasized).

An exemplary variant with two amino acid changes at 228 and 235 positions in the human IgG4 Fc amino acid sequence, IgG4PE Fc region, is provided below:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKYTCNVVDHKPSNTKVDKRVESKYGPPCPPCPAPEEGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:40, S228P and L235E substitutions emphasized).

**[0166]** The term “variant” when used in relation to PD-1 or to an anti-PD-1 antibody may refer to a peptide or polypeptide comprising one or more (such as, for example, about 1 to about 25, about 1 to about 20, about 1 to about 15, about 1 to about 10, or about 1 to about 5) amino acid sequence substitutions, deletions, and/or additions as compared to a native or unmodified sequence. For example, a PD-1 variant may result from one or more (such as, for example, about 1 to about 25, about 1 to about 20, about 1 to about 15, about 1 to about 10, or about 1 to about 5) changes to an amino acid sequence of a native PD-1. Also by way of example, a variant of an anti-PD-1 antibody may result from one or more (such as, for example, about 1 to about 25, about 1 to about 20, about 1 to about 15, about 1 to about 10, or about 1 to about 5) changes to an amino acid sequence of a native or previously unmodified anti-PD-1 antibody. Variants may be naturally occurring, such as allelic or splice variants, or may be artificially constructed.

Polypeptide variants may be prepared from the corresponding nucleic acid molecules encoding

the variants. In specific embodiments, the PD-1 variant or anti-PD-1 antibody variant at least retains PD-1 or anti-PD-1 antibody functional activity, respectively. In specific embodiments, an anti-PD-1 antibody variant binds PD-1 and/or is antagonistic to PD-1 activity. In specific embodiments, an anti-PD-1 antibody variant binds PD-1 and/or is agonistic to PD-1 activity. In certain embodiments, the variant is encoded by a single nucleotide polymorphism (SNP) variant of a nucleic acid molecule that encodes PD-1 or anti-PD-1 antibody VH or VL regions or subregions, such as one or more CDRs.

**[0167]** An “intact” antibody is one comprising an antigen-binding site as well as a CL and at least heavy chain constant regions, CH1, CH2 and CH3. The constant regions may include human constant regions or amino acid sequence variants thereof. In certain embodiments, an intact antibody has one or more effector functions.

**[0168]** “Antibody fragments” comprise a portion of an intact antibody, such as the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include, without limitation, Fab, Fab’, F(ab’)2, and Fv fragments; diabodies and di-diabodies (see, e.g., Holliger *et al.*, 1993, Proc. Natl. Acad. Sci. 90:6444-48; Lu *et al.*, 2005, J. Biol. Chem. 280:19665-72; Hudson *et al.*, 2003, Nat. Med. 9:129-34; WO 93/11161; and U.S. Pat. Nos. 5,837,242 and 6,492,123); single-chain antibody molecules (see, e.g., U.S. Pat. Nos. 4,946,778; 5,260,203; 5,482,858; and 5,476,786); dual variable domain antibodies (see, e.g., U.S. Pat. No. 7,612,181); single variable domain antibodies (sdAbs) (see, e.g., Woolven *et al.*, 1999, Immunogenetics 50: 98-101; and Streltsov *et al.*, 2004, Proc Natl Acad Sci USA. 101:12444-49); and multispecific antibodies formed from antibody fragments.

**[0169]** A “functional fragment,” “binding fragment,” or “antigen-binding fragment” of a therapeutic antibody will exhibit at least one if not some or all of the biological functions attributed to the intact antibody, the function comprising at least binding to the target antigen (e.g., a PD-1 binding fragment or fragment that binds to PD-1).

**[0170]** The term “fusion protein” as used herein refers to a polypeptide that comprises an amino acid sequence of an antibody and an amino acid sequence of a heterologous polypeptide or protein (e.g., a polypeptide or protein not normally a part of the antibody (e.g., a non-anti-PD-1 antigen-binding antibody)). The term “fusion” when used in relation to PD-1 or to an anti-PD-1 antibody refers to the joining of a peptide or polypeptide, or fragment, variant, and/or derivative thereof, with a heterologous peptide or polypeptide. In certain embodiments,

the fusion protein retains the biological activity of the PD-1 or anti-PD-1 antibody. In certain embodiments, the fusion protein comprises a PD-1 antibody VH region, VL region, VH CDR (one, two, or three VH CDRs), and/or VL CDR (one, two, or three VL CDRs), wherein the fusion protein binds to a PD-1 epitope, a PD-1 fragment, and/or a PD-1 polypeptide.

**[0171]** The term “native” when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to those which are found in nature and not manipulated, modified, and/or changed (e.g., isolated, purified, selected) by a human being.

**[0172]** The antibodies provided herein can include “chimeric” antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see U.S. Pat. No. 4,816,567; and Morrison *et al.*, 1984, Proc. Natl. Acad. Sci. USA 81:6851-55).

**[0173]** “Humanized” forms of nonhuman (e.g., murine) antibodies are chimeric antibodies that include human immunoglobulins (e.g., recipient antibody) in which the native CDR residues are replaced by residues from the corresponding CDR of a nonhuman species (e.g., donor antibody) such as mouse, rat, rabbit, or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, one or more FR region residues of the human immunoglobulin are replaced by corresponding nonhuman residues. Furthermore, humanized antibodies can comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. A humanized antibody heavy or light chain can comprise substantially all of at least one or more variable regions, in which all or substantially all of the CDRs correspond to those of a nonhuman immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. In certain embodiments, the humanized antibody will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see, Jones *et al.*, 1986, Nature 321:522-25; Riechmann *et al.*, 1988, Nature 332:323-29;

Presta, 1992, *Curr. Op. Struct. Biol.* 2:593-96; Carter *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:4285-89; U.S. Pat. Nos: 6,800,738; 6,719,971; 6,639,055; 6,407,213; and 6,054,297.

**[0174]** A “human antibody” is one that possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries (Hoogenboom and Winter, 1991, *J. Mol. Biol.* 227:381; Marks *et al.*, 1991, *J. Mol. Biol.* 222:581) and yeast display libraries (Chao *et al.*, 2006, *Nature Protocols* 1: 755-68). Also available for the preparation of human monoclonal antibodies are methods described in Cole *et al.*, Monoclonal Antibodies and Cancer Therapy 77 (1985); Boerner *et al.*, 1991, *J. Immunol.* 147(1):86-95; and van Dijk and van de Winkel, 2001, *Curr. Opin. Pharmacol.* 5: 368-74. Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, mice (*see, e.g.*, Jakobovits, 1995, *Curr. Opin. Biotechnol.* 6(5):561-66; Brüggemann and Taussing, 1997, *Curr. Opin. Biotechnol.* 8(4):455-58; and U.S. Pat. Nos. 6,075,181 and 6,150,584 regarding XENOMOUSE<sup>TM</sup> technology). See also, for example, Li *et al.*, 2006, *Proc. Natl. Acad. Sci. USA* 103:3557-62 regarding human antibodies generated via a human B-cell hybridoma technology.

**[0175]** An “affinity matured” antibody is one with one or more alterations (*e.g.*, amino acid sequence variations, including changes, additions, and/or deletions) in one or more HVRs thereof which result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Affinity matured antibodies can have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. For review, see Hudson and Souriau, 2003, *Nature Medicine* 9:129-34; Hoogenboom, 2005, *Nature Biotechnol.* 23:1105-16; Quiroz and Sinclair, 2010, *Revista Ingeneria Biomedia* 4:39-51.

**[0176]** A “blocking” antibody or an “antagonist” antibody is one which inhibits or reduces biological activity of the antigen it binds. For example, blocking antibodies or antagonist antibodies may substantially or completely inhibit the biological activity of the antigen.

**[0177]** An “agonist” antibody is an antibody that triggers a response, *e.g.*, one that mimics at least one of the functional activities of a polypeptide of interest (*e.g.*, PD-L1). An agonist antibody includes an antibody that is a ligand mimetic, for example, wherein a ligand binds to a cell surface receptor and the binding induces cell signaling or activities via an intercellular cell signaling pathway and wherein the antibody induces a similar cell signaling or activation. An “agonist” of PD-1 refers to a molecule that is capable of activating or otherwise increasing one or more of the biological activities of PD-1, such as in a cell expressing PD-1. In some embodiments, an agonist of PD-1 (*e.g.*, an agonistic antibody as described herein) may, for example, act by activating or otherwise increasing the activation and/or cell signaling pathways of a cell expressing a PD-1 protein, thereby increasing a PD-1-mediated biological activity of the cell relative to the PD-1-mediated biological activity in the absence of agonist. In some embodiments the antibodies provided herein are agonistic anti-PD-1 antibodies, including antibodies that induce PD-1 signaling.

**[0178]** “Binding affinity” generally refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (*e.g.*, a binding protein such as an antibody) and its binding partner (*e.g.*, an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (*e.g.*, antibody and antigen). The affinity of a binding molecule X for its binding partner Y can generally be represented by the dissociation constant ( $K_D$ ). Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies generally bind antigen faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present disclosure. Specific illustrative embodiments include the following. In one embodiment, the “ $K_D$ ” or “ $K_D$  value” may be measured by assays known in the art, for example by a binding assay. The  $K_D$  may be measured in a RIA, for example, performed with the Fab version of an antibody of interest and its antigen (Chen *et al.*, 1999, *J. Mol Biol* 293:865-81). The  $K_D$  or  $K_D$  value may also be measured by using surface plasmon resonance assays by Biacore<sup>®</sup>, using, for example, a Biacore<sup>®</sup>TM-2000 or a Biacore<sup>®</sup>TM-3000, or by biolayer interferometry using, for example, the Octet<sup>®</sup>QK384 system. An “on-rate” or “rate of association” or “association rate” or “ $k_{on}$ ” may also be determined with the same surface

plasmon resonance or biolayer interferometry techniques described above using, for example, a Biacore®TM-2000 or a Biacore®TM-3000, or the Octet®QK384 system.

**[0179]** The term “inhibition” or “inhibit,” when used herein, refers to partial (such as, 1%, 2%, 5%, 10%, 20%, 25%, 50%, 75%, 90%, 95%, 99%) or complete (*i.e.*, 100%) inhibition.

**[0180]** The term “attenuate,” “attenuation,” or “attenuated,” when used herein, refers to partial (such as, 1%, 2%, 5%, 10%, 20%, 25%, 50%, 75%, 90%, 95%, 99%) or complete (*i.e.*, 100%) reduction in a property, activity, effect, or value.

**[0181]** “Antibody effector functions” refer to the biological activities attributable to the Fc region (*e.g.*, a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype. Examples of antibody effector functions include but are not limited to: C1q binding; CDC; Fc receptor binding; ADCC; phagocytosis; downregulation of cell surface receptors (*e.g.*, B cell receptor); and B cell activation.

**[0182]** “T cell effector functions” refer to the biological activities attributable to various types of T cells, including but not limited to cytotoxic T cells, T helper cells, and memory T cells. Examples of T cell effector functions include: increasing T cell proliferation, secreting cytokines, releasing cytotoxins, expressing membrane-associated molecules, killing target cells, activating macrophages, and activating B cells.

**[0183]** “Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which secreted immunoglobulin bound onto Fc receptors (FcRs) present on certain cytotoxic cells (*e.g.*, Natural Killer (NK) cells, neutrophils, and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The antibodies “arm” the cytotoxic cells and are absolutely required for such killing. NK cells, the primary cells for mediating ADCC, express Fc $\gamma$ RIII only, whereas monocytes express Fc $\gamma$ RI, Fc $\gamma$ RII, and Fc $\gamma$ RIII. FcR expression on hematopoietic cells is known (*see, e.g.*, Ravetch and Kinet, 1991, *Annu. Rev. Immunol.* 9:457-92). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay (*see, e.g.*, US Pat. Nos. 5,500,362 and 5,821,337) can be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, for example, in an animal model (*see, e.g.*, Clynes *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95:652-56). Antibodies with little or no ADCC activity may be selected for use.

**[0184]** “Antibody-dependent cellular phagocytosis” or “ADCP” refers to the destruction of target cells via monocyte or macrophage-mediated phagocytosis when immunoglobulin bound onto Fc receptors (FcRs) present on certain phagocytotic cells (e.g., neutrophils, monocytes, and macrophages) enable these phagocytotic cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell. To assess ADCP activity of a molecule of interest, an *in vitro* ADCP assay (see, e.g., Bracher *et al.*, 2007, *J. Immunol. Methods* 323:160-71) can be performed. Useful phagocytotic cells for such assays include peripheral blood mononuclear cells (PBMC), purified monocytes from PBMC, or U937 cells differentiated to the mononuclear type. Alternatively or additionally, ADCP activity of the molecule of interest may be assessed *in vivo*, for example, in an animal model (see, e.g., Wallace *et al.*, 2001, *J. Immunol. Methods* 248:167-82). Antibodies with little or no ADCP activity may be selected for use.

**[0185]** “Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. An exemplary FcR is a native sequence human FcR. Moreover, an exemplary FcR is one that binds an IgG antibody (e.g., a gamma receptor) and includes receptors of the Fc $\gamma$ RI, Fc $\gamma$ RII, and Fc $\gamma$ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc $\gamma$ RII receptors include Fc $\gamma$ RIIA (an “activating receptor”) and Fc $\gamma$ RIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof (see, e.g., Daëron, 1997, *Annu. Rev. Immunol.* 15:203-34). Various FcRs are known (see, e.g., Ravetch and Kinet, 1991, *Annu. Rev. Immunol.* 9:457-92; Capel *et al.*, 1994, *Immunomethods* 4:25-34; and de Haas *et al.*, 1995, *J. Lab. Clin. Med.* 126:330-41). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (see, e.g., Guyer *et al.*, 1976, *J. Immunol.* 117:587-93; and Kim *et al.*, 1994, *Eur. J. Immunol.* 24:2429-34). Antibody variants with improved or diminished binding to FcRs have been described (see, e.g., WO 2000/42072; U.S. Pat. Nos. 7,183,387; 7,332,581; and 7,335,742; Shields *et al.* 2001, *J. Biol. Chem.* 9(2):6591-604).

**[0186]** “Complement dependent cytotoxicity” or “CDC” refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass) which are bound to their cognate antigen. To assess complement activation, a CDC assay (see, e.g., Gazzano-Santoro *et al.*, 1996, *J. Immunol. Methods* 202:163) may be performed.

Polypeptide variants with altered Fc region amino acid sequences (polypeptides with a variant Fc region) and increased or decreased C1q binding capability have been described (see, e.g., US Pat. No. 6,194,551; WO 1999/51642; Idusogie *et al.*, 2000, *J. Immunol.* 164: 4178-84).

Antibodies with little or no CDC activity may be selected for use.

**[0187]** The term “identity” refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by aligning and comparing the sequences. “Percent (%) amino acid sequence identity” with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, or MEGALIGN (DNAStar, Inc.) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

**[0188]** A “modification” of an amino acid residue/position refers to a change of a primary amino acid sequence as compared to a starting amino acid sequence, wherein the change results from a sequence alteration involving said amino acid residue/position. For example, typical modifications include substitution of the residue with another amino acid (e.g., a conservative or non-conservative substitution), insertion of one or more (e.g., generally fewer than 5, 4, or 3) amino acids adjacent to said residue/position, and/or deletion of said residue/position.

**[0189]** In the context of a polypeptide, the term “analog” as used herein refers to a polypeptide that possesses a similar or identical function as a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody but does not necessarily comprise a similar or identical amino acid sequence of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody, or possess a similar or identical structure of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody. A polypeptide that has a similar amino acid sequence refers to a polypeptide that satisfies at least one of the followings: (a) a polypeptide having an amino acid sequence that is at least 30%, at least 35%, at least 40%, at least 45%, at

least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody provided herein; (b) a polypeptide encoded by a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence encoding a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody (or VH or VL region thereof) described herein at least 5 amino acid residues, at least 10 amino acid residues, at least 15 amino acid residues, at least 20 amino acid residues, at least 25 amino acid residues, at least 30 amino acid residues, at least 40 amino acid residues, at least 50 amino acid residues, at least 60 amino residues, at least 70 amino acid residues, at least 80 amino acid residues, at least 90 amino acid residues, at least 100 amino acid residues, at least 125 amino acid residues, or at least 150 amino acid residues (*see, e.g.*, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (2001); and Maniatis *et al.*, Molecular Cloning: A Laboratory Manual (1982)); or (c) a polypeptide encoded by a nucleotide sequence that is at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the nucleotide sequence encoding a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody (or VH or VL region thereof) described herein. A polypeptide with similar structure to a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody provided herein refers to a polypeptide that has a similar secondary, tertiary, or quaternary structure of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody provided herein. The structure of a polypeptide can be determined by methods known to those skilled in the art, including but not limited to, X-ray crystallography, nuclear magnetic resonance, and crystallographic electron microscopy.

**[0190]** In the context of a polypeptide, the term “derivative” as used herein refers to a polypeptide that comprises an amino acid sequence of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an antibody that binds to a PD-1 polypeptide which has been altered by the introduction of amino acid residue substitutions, deletions, or additions. The term “derivative” as used herein also refers to a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an antibody that binds to a PD-1 polypeptide which has been chemically modified, *e.g.*, by the covalent attachment of any type of molecule to the polypeptide. For example, but not by way of limitation, a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody may

be chemically modified, *e.g.*, by increase or decrease of glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, chemical cleavage, linkage to a cellular ligand or other protein, *etc.* The derivatives are modified in a manner that is different from naturally occurring or starting peptide or polypeptides, either in the type or location of the molecules attached. Derivatives further include deletion of one or more chemical groups which are naturally present on the peptide or polypeptide. Further, a derivative of a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody may contain one or more non-classical amino acids. A polypeptide derivative possesses a similar or identical function as a PD-1 polypeptide, a fragment of a PD-1 polypeptide, or an anti-PD-1 antibody provided herein.

**[0191]** The term “host” as used herein refers to an animal, such as a mammal (*e.g.*, a human).

**[0192]** The term “host cell” as used herein refers to a particular subject cell that may be transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

**[0193]** The term “vector” refers to a substance that is used to carry or include a nucleic acid sequence, including for example, a nucleic acid sequence encoding an anti-PD-1 antibody as described herein, in order to introduce a nucleic acid sequence into a host cell. Vectors applicable for use include, for example, expression vectors, plasmids, phage vectors, viral vectors, episomes, and artificial chromosomes, which can include selection sequences or markers operable for stable integration into a host cell’s chromosome. Additionally, the vectors can include one or more selectable marker genes and appropriate expression control sequences. Selectable marker genes that can be included, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like, which are well known in the art. When two or more nucleic acid molecules are to be co-expressed (*e.g.*, both an antibody heavy and light chain or an antibody VH and VL), both nucleic acid molecules can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector expression, the encoding nucleic acids can be operationally linked to one common expression control sequence

or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The introduction of nucleic acid molecules into a host cell can be confirmed using methods well known in the art. Such methods include, for example, nucleic acid analysis such as Northern blots or polymerase chain reaction (PCR) amplification of mRNA, immunoblotting for expression of gene products, or other suitable analytical methods to test the expression of an introduced nucleic acid sequence or its corresponding gene product. It is understood by those skilled in the art that the nucleic acid molecules are expressed in a sufficient amount to produce a desired product (e.g., an anti-PD-1 antibody as described herein), and it is further understood that expression levels can be optimized to obtain sufficient expression using methods well known in the art.

**[0194]** An “isolated nucleic acid” is a nucleic acid, for example, an RNA, DNA, or a mixed nucleic acids, which is substantially separated from other genome DNA sequences as well as proteins or complexes such as ribosomes and polymerases, which naturally accompany a native sequence. An “isolated” nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Moreover, an “isolated” nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. In a specific embodiment, one or more nucleic acid molecules encoding an antibody as described herein are isolated or purified. The term embraces nucleic acid sequences that have been removed from their naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogues or analogues biologically synthesized by heterologous systems. A substantially pure molecule may include isolated forms of the molecule.

**[0195]** “Polynucleotide” or “nucleic acid,” as used interchangeably herein, refers to polymers of nucleotides of any length and includes DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase or by a synthetic reaction. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and their analogs. “Oligonucleotide,” as used herein, refers to short, generally single-stranded, synthetic polynucleotides that are generally, but not necessarily, fewer than about 200 nucleotides in length. The terms “oligonucleotide” and “polynucleotide” are not

mutually exclusive. The description above for polynucleotides is equally and fully applicable to oligonucleotides. A cell that produces an anti-PD-1 antibody of the present disclosure may include a parent hybridoma cell, as well as bacterial and eukaryotic host cells into which nucleic acids encoding the antibodies have been introduced. Suitable host cells are disclosed below.

**[0196]** Unless specified otherwise, the left-hand end of any single-stranded polynucleotide sequence disclosed herein is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA transcript that are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA transcript that are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences."

**[0197]** The term "encoding nucleic acid" or grammatical equivalents thereof as it is used in reference to nucleic acid molecule refers to a nucleic acid molecule in its native state or when manipulated by methods well known to those skilled in the art that can be transcribed to produce mRNA, which is then translated into a polypeptide and/or a fragment thereof. The antisense strand is the complement of such a nucleic acid molecule, and the encoding sequence can be deduced therefrom.

**[0198]** The term "recombinant antibody" refers to an antibody that is prepared, expressed, created, or isolated by recombinant means. Recombinant antibodies can be antibodies expressed using a recombinant expression vector transfected into a host cell, antibodies isolated from a recombinant, combinatorial antibody library, antibodies isolated from an animal (e.g., a mouse or cow) that is transgenic and/or transchromosomal for human immunoglobulin genes (see, e.g., Taylor *et al.*, 1992, Nucl. Acids Res. 20:6287-95), or antibodies prepared, expressed, created, or isolated by any other means that involves splicing of immunoglobulin gene sequences to other DNA sequences. Such recombinant antibodies can have variable and constant regions, including those derived from human germline immunoglobulin sequences (See Kabat *et al.*, *supra*). In certain embodiments, however, such recombinant antibodies may be subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis), thus the amino acid sequences of the VH and VL regions of the recombinant

antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

**[0199]** The term “detectable probe” refers to a composition that provides a detectable signal. The term includes, without limitation, any fluorophore, chromophore, radiolabel, enzyme, antibody or antibody fragment, and the like, that provide a detectable signal via its activity.

**[0200]** The term “detectable agent” refers to a substance that can be used to ascertain the existence or presence of a desired molecule, such as an anti-PD-1 antibody as described herein, in a sample or subject. A detectable agent can be a substance that is capable of being visualized or a substance that is otherwise able to be determined and/or measured (e.g., by quantitation).

**[0201]** The term “diagnostic agent” refers to a substance administered to a subject that aids in the diagnosis of a disease, disorder, or condition. Such substances can be used to reveal, pinpoint, and/or define the localization of a disease causing process. In certain embodiments, a diagnostic agent includes a substance that is conjugated to an anti-PD-1 antibody as described herein, that when administered to a subject or contacted with a sample from a subject aids in the diagnosis of a PD-1-mediated disease.

**[0202]** The term “composition” is intended to encompass a product containing the specified ingredients (e.g., an antibody provided herein) in, optionally, the specified amounts.

**[0203]** “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers, such as phosphate, citrate, and other organic acids; antioxidants, including ascorbic acid; low molecular weight (e.g., fewer than about 10 amino acid residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers, such as polyvinylpyrrolidone; amino acids, such as glycine, glutamine, asparagine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates, including glucose, mannose, or dextrins; chelating agents, such as EDTA; sugar alcohols, such as mannitol or sorbitol; salt-forming counterions, such as sodium; and/or nonionic surfactants, such as TWEEN<sup>TM</sup>, polyethylene glycol (PEG), and PLURONICS<sup>TM</sup>. The term “carrier” can also refer to a diluent, adjuvant (e.g., Freund’s adjuvant (complete or incomplete)), excipient, or vehicle. Such carriers, including pharmaceutical carriers, can be sterile liquids, such as water and oils, including those of petroleum, animal,

vegetable, or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water is an exemplary carrier when a composition (e.g., a pharmaceutical composition) is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable excipients (e.g., pharmaceutical excipients) include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. Compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations, and the like. Oral compositions, including formulations, can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in Remington and Gennaro, Remington's Pharmaceutical Sciences (18th ed. 1990). Compositions, including pharmaceutical compounds, may contain an anti-PD-1 antibody, for example, in isolated or purified form, together with a suitable amount of carriers.

**[0204]** The term "pharmaceutically acceptable" as used herein means being approved by a regulatory agency of the Federal or a state government, or listed in United States Pharmacopeia, European Pharmacopeia, or other generally recognized Pharmacopeia for use in animals, and more particularly in humans.

**[0205]** The term "excipient" refers to an inert substance which is commonly used as a diluent, vehicle, preservative, binder, or stabilizing agent, and includes, but is not limited to, proteins (e.g., serum albumin, etc.), amino acids (e.g., aspartic acid, glutamic acid, lysine, arginine, glycine, histidine, etc.), fatty acids and phospholipids (e.g., alkyl sulfonates, caprylate, etc.), surfactants (e.g., SDS, polysorbate, nonionic surfactant, etc.), saccharides (e.g., sucrose, maltose, trehalose, etc.), and polyols (e.g., mannitol, sorbitol, etc.). See, also, Remington and Gennaro, Remington's Pharmaceutical Sciences (18th ed. 1990), which is hereby incorporated by reference in its entirety.

**[0206]** The terms "subject" and "patient" may be used interchangeably. As used herein, in certain embodiments, a subject is a mammal, such as a non-primate (e.g., cow, pig, horse, cat,

dog, rat, *etc.*) or a primate (*e.g.*, monkey and human). In specific embodiments, the subject is a human.

**[0207]** “Administer” or “administration” refers to the act of injecting or otherwise physically delivering a substance as it exists outside the body (*e.g.*, an anti-PD-1 antibody as described herein) into a patient, such as by mucosal, intradermal, intravenous, intramuscular delivery, and/or any other method of physical delivery described herein or known in the art.

**[0208]** The term “effective amount” as used herein refers to the amount of an antibody or pharmaceutical composition provided herein which is sufficient to result in the desired outcome.

**[0209]** The terms “about” and “approximately” mean within 20%, within 15%, within 10%, within 9%, within 8%, within 7%, within 6%, within 5%, within 4%, within 3%, within 2%, within 1%, or less of a given value or range.

**[0210]** “Substantially all” refers to at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or about 100%.

**[0211]** The phrase “substantially similar” or “substantially the same” denotes a sufficiently high degree of similarity between two numeric values (*e.g.*, one associated with an antibody of the present disclosure and the other associated with a reference antibody) such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by the values (*e.g.*,  $K_D$  values). For example, the difference between the two values may be less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 10%, or less than about 5%, as a function of the value for the reference antibody.

**[0212]** The phrase “substantially increased,” “substantially reduced,” or “substantially different,” as used herein, denotes a sufficiently high degree of difference between two numeric values (*e.g.*, one associated with an antibody of the present disclosure and the other associated with a reference antibody) such that one of skill in the art would consider the difference between the two values to be of statistical significance within the context of the biological characteristic measured by the values. For example, the difference between said two values can be greater than about 10%, greater than about 20%, greater than about 30%, greater than about 40%, or greater than about 50%, as a function of the value for the reference antibody.

#### 4.3 Compositions and Methods of Making the Same

[0213] Provided herein are antibodies that bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope.

[0214] In certain embodiments, the antibodies provided herein bind to human and/or cynomolgus PD-1. In one embodiment, the PD-1 antibodies bind to human PD-1. In one embodiment, the PD-1 antibodies bind to cynomolgus PD-1. In one embodiment, the PD-1 antibodies bind to both human PD-1 and cynomolgus PD-1. In other embodiments, the antibodies provided herein do not bind to rodent PD-1.

[0215] In some embodiments, the anti-PD-1 antibodies bind to the extracellular domain (ECD) of PD-1. In certain embodiments, the anti-PD-1 antibodies bind to an epitope in the ECD of PD-1, which is distinct from the PD-L1 binding site. In certain embodiments, the anti-PD-1 antibodies bind to an epitope in the ECD of PD-1, which is distinct from the PD-L2 binding site. In certain embodiments, the anti-PD-1 antibodies bind to an epitope in the ECD of PD-1, which is distinct from both the PD-L1 and PD-L2-binding site.

[0216] Also provided are antibodies that competitively block an anti-PD-1 antibody provided herein from binding to a PD-1 polypeptide.

[0217] Also provided are antibodies that compete for binding to a PD-1 polypeptide with an anti-PD-1 antibody provided herein.

[0218] In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L1 to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L2 to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L1 or PD-L2 to a PD-1 polypeptide.

[0219] In some embodiments, the anti-PD-1 antibodies do not compete with PD-L1 for binding to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not compete with PD-L2 for binding to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not compete with PD-L1 or PD-L2 for binding to a PD-1 polypeptide.

[0220] In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0221]** The anti-PD-1 antibodies provided herein can also be conjugated or recombinantly fused, *e.g.*, to a diagnostic agent or detectable agent. Further provided are compositions comprising an anti-PD-1 antibody.

**[0222]** Also provided herein are isolated nucleic acid molecules encoding an immunoglobulin heavy chain, light chain, VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of anti-PD-1 antibodies that bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope.

**[0223]** Further provided are vectors and host cells comprising nucleic acid molecules encoding anti-PD-1 antibodies that bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope. Also provided are methods of making antibodies that bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope.

#### 4.3.1 Anti-PD-1 antibodies

**[0224]** In one embodiment, the present disclosure provides anti-PD-1 antibodies that may find use herein as diagnostic agents. Exemplary antibodies include polyclonal, monoclonal, humanized, human, bispecific, and heteroconjugate antibodies, as well as variants thereof having improved affinity or other properties.

**[0225]** In some embodiments, provided herein are antibodies that bind to PD-1, including a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope. In certain embodiments, the antibodies provided herein bind to human and/or cynomolgus PD-1. In other embodiments, the antibodies provided herein do not bind to rodent PD-1 (*e.g.*, a mouse PD-1). In one embodiment, an antibody provided herein binds to human PD-1. In another embodiment, an antibody provided herein binds to cynomolgus PD-1. In another embodiment, an antibody provided herein binds to human PD-1 and cynomolgus PD-1. In some embodiments, an antibody provided herein binds to human PD-1 and does not bind to a rodent PD-1 (*e.g.*, a mouse PD-1). In some embodiments, an antibody provided herein binds to cynomolgus PD-1 and does not bind to a rodent PD-1 (*e.g.*, a mouse PD-1). In some embodiments, an antibody provided herein binds to human PD-1, binds to a cynomolgus PD-1, and does not bind to a rodent PD-1 (*e.g.*, a mouse PD-1). In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L1 to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L2 to a PD-1 polypeptide. In some embodiments, the anti-PD-1 antibodies do not block the binding of PD-L1 or PD-L2 to a PD-1 polypeptide. In other embodiments, the

anti-PD-1 antibodies are humanized antibodies (e.g., comprising human constant regions) that bind PD-1, including a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope.

**[0226]** In certain embodiments, the anti-PD-1 antibody comprises a VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of any one of the murine monoclonal antibodies provided herein, such as an amino acid sequence depicted in **Tables 1-6**. Accordingly, in some embodiments, the isolated antibody or functional fragment thereof provided herein comprises one, two, and/or three heavy chain CDRs and/or one, two, and/or three light chain CDRs from: (a) the antibody PD1AB-1, (b) the antibody PD1AB-2, (c) the antibody PD1AB-3, (d) the antibody PD1AB-4, (e) the antibody PD1AB-5, or (f) the antibody PD1AB-6, as shown in **Tables 1-2**.

**Table 1. VL CDR Amino Acid Sequences**

Antibody	VL CDR1 (SEQ ID NO:)	VL CDR2 (SEQ ID NO:)	VL CDR3 (SEQ ID NO:)
PD1AB-1	KSGQSVLYSSNQKNFLA (SEQ ID NO:1)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)
PD1AB-2	KSSQSVLYSSNNKNYLA (SEQ ID NO:7)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)
PD1AB-3	KSGQSVLYSSNQKNFLA (SEQ ID NO:1)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)
PD1AB-4	KSSQSVLYSSNNKNYLA (SEQ ID NO:7)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)
PD1AB-5	KSSQSVLYSSNNKNYLA (SEQ ID NO:7)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)
PD1AB-6	KSGQSVLYSSNQKNFLA (SEQ ID NO:1)	WASTRES (SEQ ID NO:2)	HQYLYSWT (SEQ ID NO:3)

**Table 2. VH CDR Amino Acid Sequences**

Antibody	VH CDR1 (SEQ ID NO:)	VH CDR2 (SEQ ID NO:)	VH CDR3 (SEQ ID NO:)
PD1AB-1	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)
PD1AB-2	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)
PD1AB-3	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)
PD1AB-4	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)
PD1AB-5	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)
PD1AB-6	GFNIKDTYMH (SEQ ID NO:4)	RIDPANGDRK (SEQ ID NO:5)	SGPVYYYGSSYVMDY (SEQ ID NO:6)

**[0227]** In some embodiments, an antibody provided herein comprises or consists of six CDRs, for example, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 identified in **Tables 1-2**. In some embodiments, an antibody provided herein can comprise fewer than six CDRs. In some embodiments, the antibody comprises or consists of one, two, three, four, or five CDRs selected from the group consisting of VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 identified in **Tables 1-2**. In some embodiments, the antibody comprises or consists of one, two, three, four, or five CDRs selected from the group consisting of VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of the monoclonal antibody selected from the group consisting of: (a) the antibody PD1AB-1, (b) the antibody PD1AB-2, (c) the antibody PD1AB-3, (d) the antibody PD1AB-4, (e) the antibody PD1AB-5, and (f) the antibody PD1AB-6, described herein. Accordingly, in some embodiments, the antibody comprises or consists of one, two, three, four, or five CDRs of anyone of the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 identified in **Tables 1-2**.

**[0228]** In some embodiments, the antibodies provided herein comprise one or more (*e.g.*, one, two, or three) VH CDRs listed in **Table 2**. In other embodiments, the antibodies provided herein comprise one or more (*e.g.*, one, two, or three) VL CDRs listed in **Table 1**. In yet other embodiments, the antibodies provided herein comprise one or more (*e.g.*, one, two, or three) VH CDRs listed in **Table 2** and one or more VL CDRs listed in **Table 1**. Accordingly, in some embodiments, the antibodies comprise a VH CDR1 having an amino acid sequence of SEQ ID NO:4. In some embodiments, the antibodies comprise a VH CDR2 having an amino acid sequence of SEQ ID NO:5. In some embodiments, the antibodies comprise a VH CDR3 having an amino acid sequence of SEQ ID NO:6. In some embodiments, the antibodies comprise a VH CDR1 and/or a VH CDR2 and/or a VH CDR3 independently selected from any one of the VH CDR1, VH CDR2, VH CDR3 amino acid sequence(s) as depicted in **Table 2**. In some embodiments, the antibodies comprise a VL CDR1 having an amino acid sequence of any one of SEQ ID NOS:1 and 7. In another embodiment, the antibodies comprise a VL CDR2 having an amino acid sequence of SEQ ID NO:2. In some embodiments, the antibodies comprise a VL CDR3 having an amino acid sequence of SEQ ID NO:3. In some embodiments, the antibodies comprise a VL CDR1 and/or a VL CDR2 and/or a VL CDR3 independently selected from any one of the VL CDR1, VL CDR2, VL CDR3 amino acid sequences as depicted in **Table 1**.

**[0229]** In certain embodiments, the antibodies provided herein comprise a VH region comprising: (1) a VH CDR1 having an amino acid sequence of SEQ ID NO:4; (2) a VH CDR2 having an amino acid sequence of SEQ ID NO:5; and (3) a VH CDR3 having an amino acid sequence of SEQ ID NO:6; and a VL region comprising: (1) a VL CDR1 having an amino acid sequence of SEQ ID NO:1; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3.

**[0230]** In certain embodiments, the antibodies provided herein comprise a VH region comprising: (1) a VH CDR1 having an amino acid sequence of SEQ ID NO:4; (2) a VH CDR2 having an amino acid sequence of SEQ ID NO:5; and (3) a VH CDR3 having an amino acid sequence of SEQ ID NO:6; and a VL region comprising: (1) a VL CDR1 having an amino acid sequence of SEQ ID NO:7; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3.

**[0231]** In some embodiments, the antibodies provided herein comprise a VH region comprising: (1) a VH CDR1 having an amino acid sequence of SEQ ID NO:4; (2) a VH CDR2 having an amino acid sequence of SEQ ID NO:5; and (3) a VH CDR3 having an amino acid sequence of SEQ ID NO:6.

**[0232]** In other embodiments, the antibodies provided herein comprise a VL region comprising: (1) a VL CDR1 having an amino acid sequence of SEQ ID NO:1; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3.

**[0233]** In some embodiments, the antibodies provided herein comprise a VL region comprising: (1) a VL CDR1 having an amino acid sequence of SEQ ID NO:7; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3.

**[0234]** Also provided herein are antibodies comprising one or more (e.g., one, two, or three) VH CDRs and one or more (e.g., one, two, or three) VL CDRs listed in **Tables 1-2**. In particular, provided herein is an antibody comprising a VH CDR1 (SEQ ID NO:4) and a VL CDR1 (SEQ ID NO:1 or 7). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4) and a VL CDR2 (SEQ ID NO:2). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4) and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5) and a VL CDR1 (SEQ ID NO:1 or 7). In some

embodiments, the antibody comprises a VH CDR2 (SEQ ID NO:5) and a VL CDR2 (SEQ ID NO:2). In one embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5) and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR3 (SEQ ID NO:6) and a VL CDR1 (SEQ ID NOS:1 or 7). In other embodiments, the antibody comprises a VH CDR3 (SEQ ID NO:6) and a VL CDR2 (SEQ ID NO:2). In some embodiments, the antibody comprises a VH CDR3 (SEQ ID NO:6) and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), and a VL CDR1 (SEQ ID NOS:1 or 7). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), and a VL CDR2 (SEQ ID NO:2). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), and a VL CDR3 (SEQ ID NOS:3). In another embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR1 (SEQ ID NOS:1 or 7). In some embodiments, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR2 (SEQ ID NO:2). In one embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), and a VL CDR1 (SEQ ID NOS:1 or 7). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), and a VL CDR2 (SEQ ID NO:2). In some embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In some embodiments, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In other embodiments, the antibody comprises a VH CDR3 (SEQ ID NO:6), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3).

NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In some embodiments, the antibody comprises a VH CDR3 (SEQ ID NO:6), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR1 (SEQ ID NOS:1 or 7). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR2 (SEQ ID NO:2). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In some embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In some embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In one embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In other embodiments, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR2 (SEQ ID NO:2). In some embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), and a VL CDR3 (SEQ ID NO:3). In one embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3).

CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR2 (SEQ ID NO:5), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In other embodiments, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In some embodiments, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises a VH CDR1 (SEQ ID NO:4), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In one embodiment, the antibody comprises a VH CDR2 (SEQ ID NO:5), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In other embodiments, the antibody comprises a VH CDR3 (SEQ ID NO:6), a VL CDR1 (SEQ ID NOS:1 or 7), a VL CDR2 (SEQ ID NO:2), and a VL CDR3 (SEQ ID NO:3). In another embodiment, the antibody comprises any combination thereof of the VH CDRs and VL CDRs listed in **Tables 1-2**.

**[0235]** In yet another aspect, the CDRs disclosed herein include consensus sequences derived from groups of related antibodies (*see, e.g.*, **Tables 1-2**). As described herein, a “consensus sequence” refers to amino acid sequences having conserved amino acids common among a number of sequences and variable amino acids that vary within a given amino acid sequences.

**[0236]** In some embodiments, the isolated antibody or functional fragment thereof provided herein further comprises one, two, three, and/or four heavy chain FRs and/or one, two, three, and/or four light chain FRs from: (a) the antibody PD1AB-1, (b) the antibody PD1AB-2, (c) the antibody PD1AB-3, (d) the antibody PD1AB-4, (e) the antibody PD1AB-5, or (f) the antibody PD1AB-6, as shown in **Tables 3-4**.

**Table 3. VL FR Amino Acid Sequences**

Antibody	VL FR1 (SEQ ID NO:)	VL FR2 (SEQ ID NO:)	VL FR3 (SEQ ID NO:)	VL FR4 (SEQ ID NO:)
PD1AB-1	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDF LTISLQAEDVAVYYC (SEQ ID NO:16)	FGQGTKLEIKR (SEQ ID NO:17)
PD1AB-2	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDF LTISLQAEDVAVYYC (SEQ ID NO:16)	FGQGTKLEIKR (SEQ ID NO:17)

PD1AB-3	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDFT LTISNLQAEDVAVYYC (SEQ ID NO:18)	FGQGTKLEIKR (SEQ ID NO:17)
PD1AB-4	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDFT LTISSLQAEDVAVYYC (SEQ ID NO:16)	FGQGTKLEIKR (SEQ ID NO:17)
PD1AB-5	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDFT LTISSLQAEDVAVYYC (SEQ ID NO:16)	FGQGTKLEIKR (SEQ ID NO:17)
PD1AB-6	DIVMTQSPDSLAVS LGERATINC (SEQ ID NO:14)	WYQQKPGQPPKLLIY (SEQ ID NO:15)	GVPDRFSGSGSGTDFT LTISSLQAEDVAVYYC (SEQ ID NO:16)	FGQGTKLEIKR (SEQ ID NO:17)

**Table 4. VH FR Amino Acid Sequences**

Antibody	VH FR1 (SEQ ID NO:)	VH FR2 (SEQ ID NO:)	VH FR3 (SEQ ID NO:)	VH FR4 (SEQ ID NO:)
PD1AB-1	EVQLVQSGAEVKKP GATVKISCKVS (SEQ ID NO:19)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TDTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:21)	WGQGTTVTVSS (SEQ ID NO:22)
PD1AB-2	EVQLVQSGAEVKKP GATVKISCKVS (SEQ ID NO:19)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TDTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:21)	WGQGTTVTVSS (SEQ ID NO:22)
PD1AB-3	EVQLVQSGAEVKKP GATVKISCKVS (SEQ ID NO:19)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TNTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:23)	WGQGTTVTVSS (SEQ ID NO:22)
PD1AB-4	EVQLVQSGAEVKKP GATVKISCKVS (SEQ ID NO:19)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TNTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:23)	WGQGTTVTVSS (SEQ ID NO:22)
PD1AB-5	EVQLVQSGAEVKKP GATVKISCKAS (SEQ ID NO:24)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TDTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:21)	WGQGTTVTVSS (SEQ ID NO:22)
PD1AB-6	EVQLVQSGAEVKKP GATVKISCKAS (SEQ ID NO:24)	WVQQAPGKGLEWMG (SEQ ID NO:20)	YDPKFQGRVTITADTS TDTAYMELSSLRSEDT AVYYCAR (SEQ ID NO:21)	WGQGTTVTVSS (SEQ ID NO:22)

**[0237]** In certain embodiments, the isolated antibody or functional fragment thereof provided herein further comprises one, two, three, and/or four heavy chain FRs from: (a) the antibody PD1AB-1, (b) the antibody PD1AB-2, (c) the antibody PD1AB-3, (d) the antibody PD1AB-4, (e)

the antibody PD1AB-5, or (f) the antibody PD1AB-6, as shown in **Table 4**. In some embodiments, the antibody heavy chain FR(s) is from the antibody PD1AB-1. In some embodiments, the antibody heavy chain FR(s) is from the antibody PD1AB-2. In other embodiments, the antibody heavy chain FR(s) is from the antibody PD1AB-3. In certain embodiments, the antibody heavy chain FR(s) is from the antibody PD1AB-4. In other embodiments, the antibody heavy chain FR(s) is from the antibody PD1AB-5. In another embodiment, the antibody heavy chain FR(s) is from the antibody PD1AB-6.

**[0238]** In some embodiments, the isolated antibody or functional fragment thereof provided herein further comprises one, two, three, and/or four light chain FRs from: (a) the antibody PD1AB-1, (b) the antibody PD1AB-2, (c) the antibody PD1AB-3, (d) the antibody PD1AB-4, (e) the antibody PD1AB-5, or (f) the antibody PD1AB-6, as shown in **Table 3**. In some embodiments, the antibody light chain FR(s) is from the antibody PD1AB-1. In some embodiments, the antibody light chain FR(s) is from the antibody PD1AB-2. In other embodiments, the antibody light chain FR(s) is from the antibody PD1AB-3. In certain embodiments, the antibody light chain FR(s) is from the antibody PD1AB-4. In other embodiments, the antibody light chain FR(s) is from the antibody PD1AB-5. In another embodiment, the antibody light chain FR(s) is from the antibody PD1AB-6.

**[0239]** In certain embodiments, an antibody of fragment thereof described herein comprises a VH region that comprises: (1) a VH FR1 having an amino acid sequence selected from the group consisting of SEQ ID NOS:19 and 24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:21 and 23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22. In certain embodiments, an antibody of fragment thereof described herein comprises a VH region that comprises: (1) a VH FR1 having an amino acid of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22. In certain embodiments, an antibody of fragment thereof described herein comprises a VH region that comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO: 23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22. In certain embodiments, an antibody of fragment thereof described herein comprises a VH region

that comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO: 24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22. In certain embodiments, an antibody of fragment thereof described herein comprises a VH region that comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO: 23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22. In specific embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4.

**[0240]** Accordingly, in some embodiments, the humanized antibody comprises a VH region that includes a VH FR1 having an amino acid sequence selected from the group consisting of SEQ ID NOS:19 and 24. In one embodiment, the humanized antibody comprises a VH region that includes a VH FR1 having an amino acid sequence of SEQ ID NO:19. In one embodiment, the humanized antibody comprises a VH region that includes a VH FR1 having an amino acid sequence of SEQ ID NO:24. In some embodiments, the humanized antibody comprises a VH region that includes a VH FR2 having an amino acid sequence of SEQ ID NO: 20. In some embodiments, the humanized antibody comprises a VH region that includes a VH FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:21 and 23. In one embodiment, the humanized antibody comprises a VH region that includes a VH FR3 having an amino acid sequence of SEQ ID NO:21. In one embodiment, the humanized antibody comprises a VH region that includes a VH FR3 having an amino acid sequence of SEQ ID NO:23. In other embodiments, the humanized antibody comprises a VH region that includes a VH FR4 having an amino acid sequence of SEQ ID NO:22.

**[0241]** In certain embodiments, an antibody of fragment thereof described herein comprises a VL region that comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:16 and 18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NOS:16; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In

other embodiments, the VL region that comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO: 18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17.

**[0242]** Accordingly, in some embodiments, the humanized antibody comprises a VL region that includes a VL FR1 having an amino acid sequence of SEQ ID NO:14. In certain embodiments, the humanized antibody comprises a VL region that includes a VL FR2 having an amino acid sequence of SEQ ID NO:15. In other embodiments, the humanized antibody comprises a VL region that includes a VL FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:16 and 18. In one embodiment, the humanized antibody comprises a VL region that includes a VL FR3 having an amino acid sequence of SEQ ID NOS:16. In other embodiments, the humanized antibody comprises a VL region that includes a VL FR3 having an amino acid sequence of SEQ ID NO: 18. In yet other embodiments, the humanized antibody comprises a VL region that includes a VL FR4 having an amino acid sequence of SEQ ID NO:17.

**[0243]** In certain embodiments, an antibody of fragment thereof described herein comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence selected from the group consisting of SEQ ID NOS:19 and 24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:21 and 23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence selected from the group consisting of SEQ ID NOS:16 and 18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0244]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:16; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0245]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0246]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3

having an amino acid sequence of SEQ ID NO:23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:16; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0247]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:19; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0248]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ

ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:16; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0249]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:21; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3 and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3 and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0250]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:16; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3,

and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0251]** In some embodiments, an antibody of fragment thereof comprises a VH region and a VL region, wherein the VH region comprises: (1) a VH FR1 having an amino acid sequence of SEQ ID NO:24; (2) a VH FR2 having an amino acid sequence of SEQ ID NO:20; (3) a VH FR3 having an amino acid sequence of SEQ ID NO:23; and/or (4) a VH FR4 having an amino acid sequence of SEQ ID NO:22; and wherein the VL region comprises: (1) a VL FR1 having an amino acid sequence of SEQ ID NO:14; (2) a VL FR2 having an amino acid sequence of SEQ ID NO:15; (3) a VL FR3 having an amino acid sequence of SEQ ID NO:18; and/or (4) a VL FR4 having an amino acid sequence of SEQ ID NO:17. In some embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4. In other embodiments, the antibody comprises a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4. In yet other embodiments, the antibody comprises a VH region comprising all four of the above-referenced VH FR1, VH FR2, VH FR3, and VH FR4, and a VL region comprising all four of the above-referenced VL FR1, VL FR2, VL FR3, and VL FR4.

**[0252]** Also provided herein are antibodies comprising one or more (e.g., one, two, three, or four) VH FRs and one or more (e.g., one, two, three, or four) VL FRs listed in **Tables 3-4**. In particular, provided herein is an antibody comprising a VH FR1 (SEQ ID NOS:19 or 24) and a VL FR1 (SEQ ID NO:14). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24) and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24) and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24) and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20) and a VL FR1 (SEQ ID NO:14). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20) and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20) and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20) and a VL FR4 (SEQ ID NO:17). In one embodiment, the

antibody comprises a VH FR3 (SEQ ID NO:21) and a VL FR1 (SEQ ID NO:14). In other embodiments, the antibody comprises a VH FR3 (SEQ ID NO:21) and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NO:21) and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR3 (SEQ ID NO:21) and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22) and a VL FR1 (SEQ ID NO:14). In another embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22) and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22) and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22) and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), and a VL FR1 (SEQ ID NO:14). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), and a VL FR1 (SEQ ID NO:14). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), and a VL FR3 (SEQ ID NOS:16 or 18). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR2 (SEQ ID NO:15) and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR2 (SEQ ID NO:15) and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NO:19 or 24), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID

NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15) and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15) and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15) and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15) and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In other embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15) and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15) and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), and a VL FR1 (SEQ ID NO:14). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), and a VL

FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), and a VL FR1 (SEQ ID NO:14). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR1 (SEQ ID NO:14). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR1 (SEQ ID NO:14). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody

comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS: 16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NO:21), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID

NO:14), and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NO:21), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In other embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17).

NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises

a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR1 (SEQ ID NO:14). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody

comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another

embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17).

NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH

FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR2 (SEQ ID NO:15). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a

VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR3 (SEQ ID NOS:16 or 18). In some embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID NO:14), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In other embodiments, the antibody comprises a VH FR2 (SEQ ID NO:20), a VH FR3 (SEQ ID NOS:21 or 23), a VH FR4 (SEQ ID NO:22), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In one embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In another embodiment, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR2 (SEQ ID NO:20), a VL FR1 (SEQ ID NO:14), a VL FR2 (SEQ ID NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises a VH FR1 (SEQ ID NOS:19 or 24), a VH FR4 (SEQ ID NO:22), a VL FR1 (SEQ ID ID



NO:15), a VL FR3 (SEQ ID NOS:16 or 18), and a VL FR4 (SEQ ID NO:17). In some embodiments, the antibody comprises any combination thereof of the VH FRs (SEQ ID NOS:19-24) and the VL FRs (SEQ ID NOS:14-18) listed in **Tables 3-4**.

**[0253]** In some embodiments, the antibodies provided herein comprise a VH region or VH domain. In other embodiments, the antibodies provided herein comprise a VL region or VL domain. In certain embodiments, the antibodies provided herein have a combination of (i) a VH domain or VH region; and/or (ii) a VL domain or VL region. In yet other embodiments, the antibodies provided herein have a combination of (i) a VH domain or VH region; and/or (ii) a VL domain or VL region selected from the group consisting of SEQ ID NOS: 8-13 as set forth in **Tables 5-6**. In still other embodiments, the antibodies provided herein have a combination of (i) a VH domain or VH region; and/or (ii) a VL domain or VL region of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6, as set forth in **Tables 5-6**.

**[0254]** In certain embodiments, the antibodies provided herein comprise a VH region comprising: (1) a VH CDR1 having an amino acid sequence of SEQ ID NO:4; (2) a VH CDR2 having an amino acid sequence of SEQ ID NO:5; and (3) a VH CDR3 having an amino acid sequence of SEQ ID NO:6; and a VL region selected from the group consisting of SEQ ID NOS:8-10 as set forth in **Table 5**. In some embodiments, the VL region has an amino acid sequence of SEQ ID NO:8. In other embodiments, the VL region has an amino acid sequence of SEQ ID NO:9. In some embodiments, the VL region has an amino acid sequence of SEQ ID NO:10.

**[0255]** In other embodiments, the antibodies provided herein comprise a VH region selected from the group consisting of SEQ ID NOS:11-13 as set forth in **Table 6**; and a VL region comprising: (1) a VL CDR1 having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 7; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3. In yet some embodiments, the antibodies provided herein comprise a VH region selected from the group consisting of SEQ ID NOS:11-13 as set forth in **Table 6**; and a VL region comprising: (1) a VL CDR1 having an amino acid sequence of SEQ ID NO:1; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3. In still other embodiments, the antibodies provided herein comprise a VH region selected from the group consisting of SEQ ID NOS:11-13 as set forth in **Table 6**; and a VL region comprising: (1) a VL

CDR1 having an amino acid sequence of SEQ ID NO:7; (2) a VL CDR2 having an amino acid sequence of SEQ ID NO:2; and (3) a VL CDR3 having an amino acid sequence of SEQ ID NO:3. In some embodiments, the VH region has an amino acid sequence of SEQ ID NO:11. In some embodiments, the VH region has an amino acid sequence of SEQ ID NO:12. In some embodiments, the VH region has an amino acid sequence of SEQ ID NO:13.

**Table 5. VL Domain Amino Acid Sequences**

Antibody	VL (SEQ ID NO:)
PD1AB-1	DIVMTQSPDSLAVSLGERATINCKSGQSVLYSSNQKNFLAWYQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 8)
PD1AB-2	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWSQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 9)
PD1AB-3	DIVMTQSPDSLAVSLGERATINCKSGQSVLYSSNQKNFLAWYQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 10)
PD1AB-4	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWSQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 9)
PD1AB-5	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWSQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 9)
PD1AB-6	DIVMTQSPDSLAVSLGERATINCKSGQSVLYSSNQKNFLAWYQQKPGQPPKLLIYWASTRESGVPDFRSGSGSGTDFTLTISLQAEDVAVYYCHQYLYSFTFGQGTKLEIKR (SEQ ID NO: 8)

**Table 6. VH Domain Amino Acid Sequences**

Antibody	VH (SEQ ID NO:)
PD1AB-1	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 11)
PD1AB-2	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 11)
PD1AB-3	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 12)
PD1AB-4	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 12)
PD1AB-5	EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 13)
PD1AB-6	EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHWWQQAPGKGLEWMGRIDPANGDRKYDPKFQGRVTITADTSTDATYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSS (SEQ ID NO: 13)

**[0256]** Also provided herein are isolated nucleic acid molecules encoding an immunoglobulin heavy chain, light chain, VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of anti-PD-1 antibodies that bind to a PD-1 polypeptide, a PD-1 polypeptide fragment, a PD-1 peptide, or a PD-1 epitope. The exemplary nucleic acid sequences for the VL region and the VH region of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, and PD1AB-6 are shown in **Tables 7-8**.

**Table 7. VL Nucleic Acid Sequences**

Antibody	Nucleotide sequences
PD1AB-1	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCACCACCAA CTGCAAGTCCGGTCAAAGTGTGTTATACAGTTCAAATCAGAAGAACTCTTGGCCTGGTACCGC AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCCACTAGGGAACTCTGGGTC GACCGATTCAAGTGGCAGCAGGGTCTGGGACAGATTCACTCTCACCACGAGCCTGCAAGCTGA AGATGTGGCAGTTATTACTGTATCAATAACCTCTACTCGTGGACGTTGGCCAGGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:25)
PD1AB-2	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCACCACCAA CTGCAAGTCCAGCCAGAGTGTGTTATACAGCTCAAACAATAAGAAACTACTTAGCTTGGTACCGC AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCTACCCGGGAATCCGGGTC GACCGATTCAAGTGGCAGCAGGGTCTGGGACAGATTCACTCTCACCACGAGCAGCCTGCAAGCTGA AGATGTGGCAGTTATTACTGTATCAATAACCTCTACTCGTGGACGTTGGCCAGGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:26)
PD1AB-3	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCACCACCAA CTGCAAGTCCGGTCAAAGTGTGTTATACAGTTCAAATCAGAAGAACTCTTGGCCTGGTACCGC AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCCACTAGGGAACTCTGGGTC GACCGATTCAAGTGGCAGCAGGGTCTGGGACAGATTCACTCTCACCACGAGCAGCCTGCAAGCTGA AGATGTGGCAGTTATTACTGTATCAATAACCTCTACTCGTGGACGTTGGCCAGGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:27)
PD1AB-4	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCACCACCAA CTGCAAGTCCAGCCAGAGTGTGTTATACAGCTCAAACAATAAGAAACTACTTAGCTTGGTACCGC AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCTACCCGGGAATCCGGGTC GACCGATTCAAGTGGCAGCAGGGTCTGGGACAGATTCACTCTCACCACGAGCAGCCTGCAAGCTGA AGATGTGGCAGTTATTACTGTATCAATAACCTCTACTCGTGGACGTTGGCCAGGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:26)
PD1AB-5	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCACCACCAA CTGCAAGTCCAGCCAGAGTGTGTTATACAGCTCAAACAATAAGAAACTACTTAGCTTGGTACCGC AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCTACCCGGGAATCCGGGTC GACCGATTCAAGTGGCAGCAGGGTCTGGGACAGATTCACTCTCACCACGAGCAGCCTGCAAGCTGA AGATGTGGCAGTTATTACTGTATCAATAACCTCTACTCGTGGACGTTGGCCAGGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:26)

Antibody	Nucleotide sequences
PD1AB-6	GACATCGTGTGACCCAGTCTCCAGACTCCCTGGCTGTCTCTGGCGAGAGGGCCACCATCAA CTGCAAGTCCGGTCAAAGTGTCTTATACAGTTCAAATCAGAAGAACTCTTGGCTGGTACCA AGAAACCAGGACAGCCTCTAAGCTGCTCATTACTGGGCATCCACTAGGAAATCTGGGTCC GACCGATTCACTGGCAGCGGGCTGGACAGATTCACTCTACCGTCAAGCTGA AGATGTGGCAGTTATTACTGTCAATACCTACTCGTGGACGTTGCCAGGGACCAAGC TGGAGATCAAACGGAC (SEQ ID NO:25)

**Table 8. VH Nucleic Acid Sequences**

Antibody	Nucleotide sequences
PD1AB-1	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAGACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:28)
PD1AB-2	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAAACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:28)
PD1AB-3	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAAACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:29)
PD1AB-4	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAGACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:29)
PD1AB-5	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAGACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:30)
PD1AB-6	GAGGTCCAGCTGGTACAGTCTGGGCTGAGGTGAAGAAGCCTGGGCTACAGTGAAAATCTCCTG CAAGGTTCTGGATTCAACATTAAAGACACGTATATGCACTGGGTGCAACAGGCCCTGGAAAAG GGCTTGAGTGGATGGGAAGGATTGATCCTGCGAATGGTGTAGGAAATATGACCCGAAGTCCAG GGCAGAGTCACCATAACCGCGGACACGTCTACAGACACAGCCTACATGGAGCTGAGCAGCCTGAG ATCTGAGGACACGGCCGTGTATTACTGTGCTAGATCAGGCCCTGTTATTACTACGGTAGTAGCT ACGTTATGGACTACTGGGTCAAGGAACCACAGTCACCGTCTCCTCA (SEQ ID NO:30)

**[0257]** In some embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-1. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:11, and a VL amino acid sequence of SEQ ID NO:8.

**[0258]** In other embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-2. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:11, and a VL amino acid sequence of SEQ ID NO:9.

**[0259]** In some embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-3. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:12, and a VL amino acid sequence of SEQ ID NO:10.

**[0260]** In other embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-4. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:12, and a VL amino acid sequence of SEQ ID NO:9.

**[0261]** In some embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-5. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:13, and a VL amino acid sequence of SEQ ID NO:9.

**[0262]** In other embodiments, an antibody provided herein has a VH and a VL amino acid sequence of PD1AB-6. In some embodiments, an antibody comprises a VH amino acid sequence of SEQ ID NO:13, and a VL amino acid sequence of SEQ ID NO:8.

**[0263]** In certain embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (*e.g.*, an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the light chain comprises a constant region having an amino acid sequence of:

TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDST  
YSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:41).

**[0264]** In other embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (*e.g.*, an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises a human IgG1 Fc region having an amino acid sequence of:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP

KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:36, K322 emphasized).

In some embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain does not comprise a human IgG1 Fc region having an amino acid sequence of SEQ ID NO:36.

**[0265]** In certain embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises a human IgG1-K322A Fc region having an amino acid sequence of:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP  
KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL  
HQDWLNGKEYKCAVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY  
PSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT  
QKSLSLSPGK (SEQ ID NO:37, K322A substitution emphasized).

**[0266]** In some embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises a human IgG4 Fc region having an amino acid sequence of:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPSSCPAPEFLGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQD  
WLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:38, S228 and L235 emphasized).

**[0267]** In another embodiment, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises a

human IgG4P Fc region having an amino acid sequence of:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTPREEQFNSTYRVVSVLTVLHQD  
WLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQOPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:39, S228P substitution emphasized).

**[0268]** In yet another embodiment, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises a human IgG4PE Fc region having an amino acid sequence of:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPAVLQSSGLYS  
LSSVVTVPSSSLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFEGGPSVFLFPPKPK  
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTPREEQFNSTYRVVSVLTVLHQD  
WLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQOPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKS  
LSLSPGK (SEQ ID NO:40, S228P and L235E substitutions emphasized).

In some embodiments, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain does not comprise a human IgG4PE Fc region having an amino acid sequence of SEQ ID NO:40.

**[0269]** In still another embodiment, an antibody or antigen-binding fragment thereof described herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the light chain comprises a constant region having an amino acid sequence of SEQ ID NO:41; and the heavy chain comprises an Fc region having an amino acid sequence selected from the group consisting of SEQ ID NOS:36-40.

**[0270]** In certain embodiments, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the light chain comprises an amino acid sequence as follows:

DIVMTQSPDSLAVSLGERATINCKSGQSVLYSSNQKNFLAWYQQKPGQPPKLLIYWASTRESGV  
PDRFSGSGSGTDFTLTISSLQAEDVAVYYCHQYLYSWTFQGKLEIKRTVAAPSVFIFPPSDE  
QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTTLSKADYE  
KHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:31, LC\_PD1AB-6-IgG1).

**[0271]** In some embodiments, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises an amino acid sequence as follows:

EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHVQQAPGKGLEWMGRIDPANGDRKYDPKF  
QGRVTITADTSTDAYMELSSLRSEDTAVYYCARSGPVYYGSSYVMDYWGQGTTVTVSSASTK  
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAPLQSSGLYSLSSV  
VTVPSSSLGTQTYICNVNHPKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKD  
TLMISRTPEVTCVVVDVSCHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW  
LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI  
AVEWESNGQPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVVMHEALHNHYTQKSL  
SLSPGK (SEQ ID NO:32, HC\_PD1AB-6-IgG1, K322 emphasized).

**[0272]** In other embodiments, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises an amino acid sequence as follows:

EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHVQQAPGKGLEWMGRIDPANGDRKYDPKF  
QGRVTITADTSTDAYMELSSLRSEDTAVYYCARSGPVYYGSSYVMDYWGQGTTVTVSSASTK  
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAPLQSSGLYSLSSV  
VTVPSSSLGTQTYICNVNHPKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKD  
TLMISRTPEVTCVVVDVSCHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW  
LNGKEYKCAVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI  
AVEWESNGQPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVVMHEALHNHYTQKSL  
SLSPGK (SEQ ID NO:33, HC\_PD1AB-6-IgG1-K322A, K322A substitution emphasized).

**[0273]** In another embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises an amino acid sequence as follows:

EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHVQQAPGKGLEWMGRIDPANGDRKYDPKF  
QGRVTITADTSTDAYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSSASTK  
GPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV  
VTVPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLM  
ISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVSVLTVLHQDWLNG  
KEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVE  
WESNGQPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSL

PGK (SEQ ID NO:34, HC PD1AB-6-IgG4P, IgG4P Fc backbone italicized and underlined).

**[0274]** In yet another embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein the heavy chain comprises an amino acid sequence as follows:

EVQLVQSGAEVKKPGATVKISCKASGFNIKDTYMHVQQAPGKGLEWMGRIDPANGDRKYDPKF  
QGRVTITADTSTDAYMELSSLRSEDTAVYYCARSGPVYYYGSSYVMDYWGQGTTVTVSSASTK  
GPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV  
VTVPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFEGGPSVFLFPPKPKDTLM  
ISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVSVLTVLHQDWLNG  
KEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVE  
WESNGQPENNYKTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSL  
PGK (SEQ ID NO:35, HC PD1AB-6-IgG4PE, IgG4PE Fc backbone italicized and underlined).

**[0275]** In one particular embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein (i) the light chain comprises an amino acid sequence of SEQ ID NO:31; and (ii) the heavy chain comprises an amino acid sequence of SEQ ID NO:32.

**[0276]** In another particular embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein (i) the light chain comprises an amino acid sequence of SEQ ID NO:31; and (ii) the heavy chain comprises an amino acid sequence of SEQ ID NO:33.

**[0277]** In yet another particular embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (e.g., an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein (i) the light chain comprises an amino acid sequence of SEQ ID NO:31; and (ii) the heavy chain comprises an amino acid sequence of SEQ ID NO:34.

**[0278]** In still another particular embodiment, an antibody provided herein, which specifically binds to a PD-1 polypeptide (*e.g.*, an ECD of PD-1, for example human PD-1), comprises a light chain and a heavy chain, wherein (i) the light chain comprises an amino acid sequence of SEQ ID NO:31; and (ii) the heavy chain comprises an amino acid sequence of SEQ ID NO:35.

**[0279]** In yet another aspect, antibodies are provided that compete with one of the exemplified antibodies or functional fragments for binding to PD-1. Such antibodies may also bind to the same epitope as one of the herein exemplified antibodies, or an overlapping epitope. Antibodies and fragments that compete with or bind to the same epitope as the exemplified antibodies are expected to show similar functional properties. The exemplified antigen-binding proteins and fragments include those with the VH and VL regions, and CDRs provided herein, including those in **Tables 1-6**. Thus, as a specific example, the antibodies that are provided include those that compete with an antibody comprising: (a) 1, 2, 3, 4, 5, or all 6 of the CDRs listed for an antibody listed in **Tables 1-2**; (b) a VH and a VL selected from the VH and the VL regions listed for an antibody listed in **Tables 5-6**; or (c) two light chains and two heavy chains comprising a VH and a VL as specified for an antibody listed in **Tables 5-6**. In some embodiments, the antibody is PD1AB-1. In some embodiments, the antibody is PD1AB-2. In some embodiments, the antibody is PD1AB-3. In some embodiments, the antibody is PD1AB-4. In some embodiments, the antibody is PD1AB-5. In some embodiments, the antibody is PD1AB-6.

**[0280]** In another aspect, antibodies or antigen-binding fragments thereof provided herein bind to a region, including an epitope, of human PD-1 or cynomolgus PD-1. For example, in some embodiments, an antibody provided herein binds to a region of human PD-1 (SEQ ID NO:42) comprising amino acid residues 33 to 109 of human PD-1. In still another aspect, antibodies provided herein bind to a specific epitope of human PD-1.

**[0281]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one of residues 100-109 (SEQ ID NO:43) within an amino acid sequence of SEQ ID NO:42. In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one of residues 100-105 (SEQ ID NO:44) within an amino acid sequence of SEQ ID NO:42.

**[0282]** In particular embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one residue selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42. In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to at least one residue selected from the group consisting of L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0283]** In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to two or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0284]** In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to three or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0285]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to four or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0286]** In one embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to five or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0287]** In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to six or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0288]** In yet another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to seven or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0289]** In still another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to eight or more residues selected from the group consisting of N33, T51,

S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0290]** In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to nine or more residues selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0291]** In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to all ten residues from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.

**[0292]** In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to N33 within an amino acid sequence of SEQ ID NO:42. In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to T51 within an amino acid sequence of SEQ ID NO:42. In a particular embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to S57 within an amino acid sequence of SEQ ID NO:42. In one specific embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to L100 within an amino acid sequence of SEQ ID NO:42. In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to N102 within an amino acid sequence of SEQ ID NO:42. In other embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to G103 within an amino acid sequence of SEQ ID NO:42. In another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to R104 within an amino acid sequence of SEQ ID NO:42. In yet another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to G103 and R104 within an amino acid sequence of SEQ ID NO:42. In still another embodiment, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to D105 within an amino acid sequence of SEQ ID NO:42. In some embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to H107 within an amino acid sequence of SEQ ID NO:42. In certain embodiments, the antibody or antigen-binding fragment thereof, when bound to PD-1, binds to S109 within an amino acid sequence of SEQ ID NO:42. Any combination of two, three, four, five, six, seven, eight, nine, ten or more of the above-referenced amino acid PD-1 binding sites is also contemplated.

**[0293]** In one aspect, provided herein are antibodies that specifically bind to PD-1 and can modulate PD-1 activity and/or expression (*e.g.*, activate PD-1 signaling and/or inhibit PD-1 expression). In certain embodiments, a PD-1 agonist is provided herein that is an antibody provided herein that specifically binds to an ECD of human PD-1, and activates (*e.g.*, partially activates) at least one PD-1 activity (*e.g.*, inhibiting cytokine production). In certain embodiments, a PD-1 agonist provided herein is an antibody provided herein that specifically binds to an ECD of human PD-1, and downregulates PD-1 expression. In certain embodiments, provided herein are antibodies that specifically bind to PD-1 and that (a) attenuate T cell activity, *e.g.*, as determined by inhibition of cytokine production; and/or (b) downregulate PD-1 expression in a cell. In certain embodiments, provided herein are antibodies that specifically bind to PD-1 and that (a) attenuate T cell activity, *e.g.*, as determined by inhibition of cytokine production; (b) downregulate PD-1 expression in a cell; and/or (c) do not inhibit PD-L1 and/or PD-L2 binding to PD-1. In certain embodiments, the antibodies that specifically bind to PD-1 bind to an ECD of human PD-1, or an epitope of an ECD of human PD-1 thereof. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0294]** PD-1 activity can relate to any activity of PD-1 such as those known or described in the art. PD-1 activity and PD-1 signaling are used interchangeably herein. In certain aspects, PD-1 activity is induced by PD-1 ligand (*e.g.*, PD-L1) binding to PD-1. Expression levels of PD-1 can be assessed by methods described herein or known to one of skill in the art (*e.g.*, Western blotting, ELISA, immunohistochemistry, or flow cytometry). In certain embodiments, provided herein are antibodies that specifically bind to PD-1 and decrease PD-1 expression. In certain embodiments, provided herein are antibodies that specifically bind to PD-1 and attenuate T cell activity. In certain embodiments, provided herein are antibodies that specifically bind to PD-1 and inhibit cytokine production. In certain embodiments, provided herein are antibodies

that specifically bind to PD-1 and activate (e.g., partially activate) PD-1 signaling. In certain embodiments, the antibodies that specifically bind to PD-1 bind to an ECD of human PD-1, or an epitope of an ECD of human PD-1 thereof. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibodies specifically bind to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0295]** In certain embodiments, an anti-PD-1 antibody provided herein attenuates (e.g., partially attenuates) T cell activity. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 10%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 15%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 20%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 25%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 30%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 35%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 40%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 45%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 50%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 55%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 60%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 65%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 70%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 75%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 80%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about

85%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 90%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 95%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 98%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 99%. In some embodiments, an anti-PD-1 antibody provided herein attenuates T cell activity by at least about 100%. In certain embodiments, an anti-PD-1 antibody provided herein can attenuate (e.g., partially attenuate) T cell activity by at least about 25% to about 65%. In specific embodiments, the T cell activity attenuation is assessed by methods described herein. In some embodiments, the T cell activity attenuation is assessed by methods known to one of skill in the art. In certain embodiments, the T cell activity attenuation is relative to T cell activity in the presence of stimulation without any anti-PD-1 antibody. In certain embodiments, the T cell activity attenuation is relative to T cell activity in the presence of stimulation with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1). A non-limiting example of T cell activity is secretion of a cytokine. In some embodiments, the cytokine is selected from the group consisting of IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In some embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-6. In another embodiment, the cytokine is IL-12. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IL-22. In still other embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is IFN- $\gamma$ . In yet other embodiments, the cytokine is TNF- $\alpha$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ .

**[0296]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-2 secretion (e.g., from a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 5%. In some

embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-2 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-2 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-2 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-2 secretion is assessed by

methods known to one of skill in the art (e.g., MesoScale<sup>TM</sup> Discovery (MSD) multiplex assay). In a specific embodiment, the IL-2 secretion is inhibited relative to IL-2 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-2 secretion is inhibited relative to IL-2 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0297]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-2 secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at

least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-2 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0298]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-17 secretion (e.g., from a cell, for example, T cells). In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 5%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 10%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 15%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 20%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 25%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 30%. In some

embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 35%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 40%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 45%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 50%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 55%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 65%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 70%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 80%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 85%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 95%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 98%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-17 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-17 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-17 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-17 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-17 secretion is inhibited relative to IL-17 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-17 secretion is inhibited relative to IL-17 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0299]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-

binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-17 secretion. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 50 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 40 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 20 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 1 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 50 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at

least about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-17 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0300]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IFN- $\gamma$  secretion (e.g., from a cell, for example, T cells). In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 5%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 10%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 15%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 25%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 30%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 35%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 40%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 45%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 50%. In one embodiment, an antibody provided herein specifically

binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 55%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 60%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 65%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 70%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 75%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 80%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 85%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 90%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 95%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 98%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IFN- $\gamma$  secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IFN- $\gamma$  secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IFN- $\gamma$  secretion is assessed by methods described herein. In other embodiments, the inhibition of IFN- $\gamma$  secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion in the absence of anti-PD-1 antibody. In other embodiments, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0301]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IFN- $\gamma$  secretion. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 50 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 40 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 30 nM. In

another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 20 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 10 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.75 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.05 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.005 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at most about 0.001 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 50 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 40 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 20 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 10 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 1 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.75 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.1 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.05 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.01 nM. In some embodiments, an

anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.005 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IFN- $\gamma$  secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0302]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-1 secretion (e.g., from a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion

by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-1 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-1 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-1 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-1 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-1 secretion is inhibited relative to IL-1 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-1 secretion is inhibited relative to IL-1 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0303]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-1 secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided

herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-1 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0304]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding

fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-6 secretion (*e.g.*, from a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-6 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and

inhibits IL-6 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-6 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-6 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-6 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-6 secretion is inhibited relative to IL-6 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-6 secretion is inhibited relative to IL-6 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0305]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-6 secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein

inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-6 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0306]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-12 secretion (e.g., from a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12

secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-12 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-12 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-12 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-12 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-12 secretion is inhibited relative to IL-12 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-12 secretion is

inhibited relative to IL-12 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0307]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-12 secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at

least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-12 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0308]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-22 secretion (e.g., from a cell, for example, a T cell). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 35%. In one embodiment, an antibody provided herein

specifically binds to PD-1 and inhibits IL-22 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-22 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-22 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-22 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-22 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-22 secretion is inhibited relative to IL-22 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-22 secretion is inhibited relative to IL-22 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0309]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-22 secretion.

In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.1 nM.

antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-22 secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0310]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit IL-23 secretion (e.g., from a cell, for example, a T cell). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23

secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits IL-23 secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit IL-23 secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-23 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-23 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-23 secretion is inhibited relative to IL-23 secretion in the absence of anti-PD-1 antibody. In other embodiments, the IL-23 secretion is inhibited relative to IL-23 secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0311]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits IL-23 secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23

secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits IL-23 secretion with an EC<sub>50</sub> of at least about 0.001 nM.

at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0312]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit GM-CSF secretion (e.g., from a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and

inhibits GM-CSF secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits GM-CSF secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit GM-CSF secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of GM-CSF secretion is assessed by methods described herein. In other embodiments, the inhibition of GM-CSF secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the GM-CSF secretion is inhibited relative to GM-CSF secretion in the absence of anti-PD-1 antibody. In other embodiments, the GM-CSF secretion is inhibited relative to GM-CSF secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0313]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits GM-CSF secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.75 nM. In

another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits GM-CSF secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0314]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and inhibit TNF- $\alpha$  secretion (e.g., from a cell, for example, a T cell). In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 5%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 15%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 20%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 30%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 35%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 45%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 50%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 60%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 75%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 80%. In other embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 90%. In one embodiment, an antibody provided herein specifically binds to PD-1 and

inhibits TNF- $\alpha$  secretion by at least about 95%. In some embodiments, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and inhibits TNF- $\alpha$  secretion by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and inhibit TNF- $\alpha$  secretion by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of TNF- $\alpha$  secretion is assessed by methods described herein. In other embodiments, the inhibition of TNF- $\alpha$  secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion in the absence of anti-PD-1 antibody. In other embodiments, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0315]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) inhibits TNF- $\alpha$  secretion. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about

0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein inhibits TNF- $\alpha$  secretion with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the EC<sub>50</sub> is assessed by MSD multiplex assay.

**[0316]** In specific embodiments, antibodies provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) specifically bind to PD-1 and downregulate PD-1 expression (e.g., in a cell, for example, T cells). In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at

least about 5%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 10%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 15%. In some embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 20%. In other embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 25%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 30%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 35%. In some embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 40%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 45%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 50%. In other embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 55%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 60%. In some embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 65%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 70%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 75%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 80%. In some embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 85%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 90%. In other embodiments, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 95%. In one embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 98%. In another embodiment, an antibody provided herein specifically binds to PD-1 and downregulates PD-1 expression by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and

downregulates PD-1 expression by at least about 25% or 35%, optionally to about 75%. In some embodiments, the downregulation of PD-1 expression is assessed by methods described herein. In other embodiments, the downregulation of PD-1 expression is assessed by methods known to one of skill in the art (e.g., flow cytometry, Western blotting, Northern blotting, or RT-PCR). In a specific embodiment, the downregulation of PD-1 expression is assessed by flow cytometry. In another embodiment, the downregulation of PD-1 expression is assessed by Western blotting. In yet another embodiment, the downregulation of PD-1 expression is assessed by Northern blotting. In still another embodiment, the downregulation of PD-1 expression is assessed by RT-PCR. In a specific embodiment, the PD-1 expression is downregulated relative to PD-1 expression downregulation in the absence of anti-PD-1 antibody. In other embodiments, the PD-1 expression is downregulated relative to PD-1 expression downregulation in the presence of an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0317]** In certain embodiments, an anti-PD-1 antibody provided herein (e.g., any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6) downregulates PD-1 expression. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with

an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, an anti-PD-1 antibody provided herein downregulates PD-1 expression with an EC<sub>50</sub> of at least about 0.001 nM. In specific embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by other methods known to one of skill in the art (*e.g.*, flow cytometry, Western blotting, Northern blotting, or RT-PCR). In a specific embodiment, the EC<sub>50</sub> is assessed by flow cytometry. In another embodiment, the EC<sub>50</sub> is assessed by Western blotting. In yet another embodiment, the EC<sub>50</sub> is assessed by Northern blotting. In still another embodiment, the EC<sub>50</sub> is assessed by RT-PCR.

**[0318]** In certain embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof. In another embodiment, the downregulation occurs as early as 6 hours after the contact. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact. In still another embodiment, the downregulation occurs as early as 10 hours after the contact. In one embodiment, the downregulation occurs as early as 12 hours after the contact. In another embodiment, the downregulation occurs as early as 14 hours after the contact. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact. In still another embodiment, the downregulation occurs as early as 18 hours after the contact. In one embodiment, the downregulation occurs as early as 20 hours after the contact. In another embodiment, the downregulation occurs as early as 22 hours after the contact. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact. In some embodiments, the contact is with the antibody. In other embodiments, the contact is with an antigen-binding fragment thereof.

**[0319]** In some embodiments, the downregulation of PD-1 expression on the surface of T cells precedes cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine

inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and precedes cytokine inhibition.

**[0320]** In other embodiments, the downregulation of PD-1 expression on the surface of T cells is concurrent with cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and is concurrent with cytokine inhibition.

**[0321]** In yet other embodiments, the downregulation of PD-1 expression on the surface of T cells is after cytokine inhibition. In one embodiment, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In another embodiment, the downregulation occurs as early as 6 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 8 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In still another embodiment, the downregulation occurs as early as 10 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In one embodiment, the downregulation occurs as early as 12 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In another embodiment, the downregulation occurs as early as 14 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 16 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In still another embodiment, the downregulation occurs as early as 18 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In one embodiment, the downregulation occurs as early as 20 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In another embodiment, the downregulation occurs as early as 22 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition. In yet another embodiment, the downregulation occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof, and is after cytokine inhibition.

#### **4.3.1.1 Polyclonal Antibodies**

**[0322]** The antibodies of the present disclosure may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include a PD-1 polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized

or to immunize the mammal with the protein and one or more adjuvants. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Ribi, CpG, Poly 1C, Freund's complete adjuvant, and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation. The mammal can then be bled, and the serum assayed for PD-1 antibody titer. If desired, the mammal can be boosted until the antibody titer increases or plateaus. Additionally or alternatively, lymphocytes may be obtained from the immunized animal for fusion and preparation of monoclonal antibodies from hybridoma as described below.

#### **4.3.1.2 Monoclonal Antibodies**

**[0323]** The antibodies of the present disclosure may alternatively be monoclonal antibodies. Monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, 1975, *Nature* 256:495-97, or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567).

**[0324]** In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as described above to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. After immunization, lymphocytes are isolated and then fused with a myeloma cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (*Goding, Monoclonal Antibodies: Principles and Practice* 59-103 (1986)).

**[0325]** The hybridoma cells thus prepared are seeded and grown in a suitable culture medium which, in certain embodiments, contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells (also referred to as fusion partner). For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the selective culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which prevent the growth of HGPRT-deficient cells.

**[0326]** Exemplary fusion partner myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a

selective medium that selects against the unfused parental cells. Exemplary myeloma cell lines are murine myeloma lines, such as SP-2 and derivatives, for example, X63-Ag8-653 cells available from the American Type Culture Collection (Manassas, VA), and those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center (San Diego, CA). Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, 1984, *Immunol. 133:3001-05*; and Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications 51-63 (1987)).

**[0327]** Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. The binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as RIA or ELISA. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis described in Munson *et al.*, 1980, *Anal. Biochem. 107:220-39*.

**[0328]** Once hybridoma cells that produce antibodies of the desired specificity, affinity, and/or activity are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *supra*). Suitable culture media for this purpose include, for example, DMEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal, for example, by i.p. injection of the cells into mice.

**[0329]** The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional antibody purification procedures such as, for example, affinity chromatography (e.g., using protein A or protein G-Sepharose) or ion-exchange chromatography, hydroxylapatite chromatography, gel electrophoresis, dialysis, *etc.*

**[0330]** DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells can serve as a source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells, such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce antibody protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et*

*al.*, 1993, Curr. Opinion in Immunol. 5:256-62 and Plückthun, 1992, Immunol. Revs. 130:151-88.

**[0331]** In some embodiments, an antibody that binds a PD-1 epitope comprises an amino acid sequence of a VH domain and/or an amino acid sequence of a VL domain encoded by a nucleotide sequence that hybridizes to (1) the complement of a nucleotide sequence encoding any one of the VH and/or VL domain described herein under stringent conditions (e.g., hybridization to filter-bound DNA in 6X sodium chloride/sodium citrate (SSC) at about 45 °C followed by one or more washes in 0.2X SSC/0.1% SDS at about 50-65 °C), under highly stringent conditions (e.g., hybridization to filter-bound nucleic acid in 6X SSC at about 45 °C followed by one or more washes in 0.1X SSC/0.2% SDS at about 68 °C), or under other stringent hybridization conditions which are known to those of skill in the art. *See, e.g., Current Protocols in Molecular Biology* Vol. I, 6.3.1-6.3.6 and 2.10.3 (Ausubel *et al.* eds., 1989).

**[0332]** In some embodiments, an antibody that binds a PD-1 epitope comprises an amino acid sequence of a VH CDR or an amino acid sequence of a VL CDR encoded by a nucleotide sequence that hybridizes to the complement of a nucleotide sequence encoding any one of the VH CDRs and/or VL CDRs depicted in **Tables 1-2** under stringent conditions (e.g., hybridization to filter-bound DNA in 6X SSC at about 45 °C followed by one or more washes in 0.2X SSC/0.1% SDS at about 50-65 °C), under highly stringent conditions (e.g., hybridization to filter-bound nucleic acid in 6X SSC at about 45 °C followed by one or more washes in 0.1X SSC/0.2% SDS at about 68 °C), or under other stringent hybridization conditions which are known to those of skill in the art (*see, e.g., Ausubel et al., supra*).

**[0333]** In a further embodiment, monoclonal antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in, for example, Antibody Phage Display: Methods and Protocols (O'Brien and Aitken eds., 2002). In principle, synthetic antibody clones are selected by screening phage libraries containing phages that display various fragments of antibody variable region (Fv) fused to phage coat protein. Such phage libraries are screened against the desired antigen. Clones expressing Fv fragments capable of binding to the desired antigen are adsorbed to the antigen and thus separated from the non-binding clones in the library. The binding clones are then eluted from the antigen and can be further enriched by additional cycles of antigen adsorption/elution.

**[0334]** Variable domains can be displayed functionally on phage, either as single-chain Fv (scFv) fragments, in which VH and VL are covalently linked through a short, flexible peptide, or as Fab fragments, in which they are each fused to a constant domain and interact non-covalently, as described, for example, in Winter *et al.*, 1994, Ann. Rev. Immunol. 12:433-55.

**[0335]** Repertoires of VH and VL genes can be separately cloned by PCR and recombined randomly in phage libraries, which can then be searched for antigen-binding clones as described in Winter *et al.*, *supra*. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned to provide a single source of human antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths *et al.*, 1993, EMBO J 12:725-34. Finally, naive libraries can also be made synthetically by cloning the unarranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement *in vitro* as described, for example, by Hoogenboom and Winter, 1992, J. Mol. Biol. 227:381-88.

**[0336]** Screening of the libraries can be accomplished by various techniques known in the art. For example, PD-1 (*e.g.*, a PD-1 polypeptide, fragment, or epitope) can be used to coat the wells of adsorption plates, expressed on host cells affixed to adsorption plates or used in cell sorting, conjugated to biotin for capture with streptavidin-coated beads, or used in any other method for panning display libraries. The selection of antibodies with slow dissociation kinetics (*e.g.*, good binding affinities) can be promoted by use of long washes and monovalent phage display as described in Bass *et al.*, 1990, Proteins 8:309-14 and WO 92/09690, and by use of a low coating density of antigen as described in Marks *et al.*, 1992, Biotechnol. 10:779-83.

**[0337]** Anti-PD-1 antibodies can be obtained by designing a suitable antigen screening procedure to select for the phage clone of interest followed by construction of a full length anti-PD-1 antibody clone using VH and/or VL sequences (*e.g.*, the Fv sequences), or various CDR sequences from VH and VL sequences, from the phage clone of interest and suitable constant region (*e.g.*, Fc) sequences described in Kabat *et al.*, *supra*.

**[0338]** In another embodiment, anti-PD-1 antibody is generated by using methods as described in Bowers *et al.*, 2011, Proc Natl Acad Sci USA. 108:20455-60, *e.g.*, the SHM-XHL<sup>TM</sup> platform (AnaptysBio, San Diego, CA). Briefly, in this approach, a fully human library of IgGs

is constructed in a mammalian cell line (*e.g.*, HEK293) as a starting library. Mammalian cells displaying immunoglobulin that binds to a target peptide or epitope are selected (*e.g.*, by FACS sorting), then activation-induced cytidine deaminase (AID)-triggered somatic hypermutation is reproduced *in vitro* to expand diversity of the initially selected pool of antibodies. After several rounds of affinity maturation by coupling mammalian cell surface display with *in vitro* somatic hypermutation, high affinity, high specificity anti-PD-1 antibodies are generated. Further methods that can be used to generate antibody libraries and/or antibody affinity maturation are disclosed, *e.g.*, in U.S. Patent Nos. 8,685,897 and 8,603,930, and U.S. Publ. Nos. 2014/0170705, 2014/0094392, 2012/0028301, 2011/0183855, and 2009/0075378, each of which are incorporated herein by reference.

#### **4.3.1.3 Antibody Fragments**

**[0339]** The present disclosure provides antibodies and antibody fragments that bind to PD-1. In certain circumstances there are advantages of using antibody fragments, rather than whole antibodies. The smaller size of the fragments allows for rapid clearance, and may lead to improved access to cells, tissues, or organs. For a review of certain antibody fragments, see Hudson *et al.*, 2003, *Nature Med.* 9:129-34.

**[0340]** Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (*see, e.g.*, Morimoto *et al.*, 1992, *J. Biochem. Biophys. Methods* 24:107-17; and Brennan *et al.*, 1985, *Science* 229:81-83). However, these fragments can now be produced directly by recombinant host cells. Fab, Fv, and scFv antibody fragments can all be expressed in and secreted from *E. coli* or yeast cells, thus allowing the facile production of large amounts of these fragments. Antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')<sub>2</sub> fragments (Carter *et al.*, 1992, *Bio/Technology* 10:163-67). According to another approach, F(ab')<sub>2</sub> fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')<sub>2</sub> fragment with increased *in vivo* half-life comprising salvage receptor binding epitope residues are described in, for example, U.S. Pat. No. 5,869,046. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In certain embodiments, an antibody is a single chain Fv fragment (scFv) (*see, e.g.*, WO 93/16185; U.S. Pat. Nos. 5,571,894 and 5,587,458). Fv and scFv have intact combining sites that are devoid of

constant regions; thus, they may be suitable for reduced nonspecific binding during *in vivo* use. scFv fusion proteins may be constructed to yield fusion of an effector protein at either the amino or the carboxy terminus of an scFv (See, *e.g.*, Borrebaeck ed., *supra*). The antibody fragment may also be a “linear antibody,” for example, as described in the references cited above. Such linear antibodies may be monospecific or multi-specific, such as bispecific.

**[0341]** Smaller antibody-derived binding structures are the separate variable domains (V domains) also termed single variable domain antibodies (sdAbs). Certain types of organisms, the camelids and cartilaginous fish, possess high affinity single V-like domains mounted on an Fc equivalent domain structure as part of their immune system. (Woolven *et al.*, 1999, Immunogenetics 50: 98-101; and Streltsov *et al.*, 2004, Proc Natl Acad Sci USA. 101:12444-49). The V-like domains (called VhH in camelids and V-NAR in sharks) typically display long surface loops, which allow penetration of cavities of target antigens. They also stabilize isolated VH domains by masking hydrophobic surface patches.

**[0342]** These VhH and V-NAR domains have been used to engineer sdAbs. Human V domain variants have been designed using selection from phage libraries and other approaches that have resulted in stable, high binding VL- and VH-derived domains.

**[0343]** Antibodies provided herein include, but are not limited to, immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, for example, molecules that contain an antigen binding site that bind to a PD-1 epitope. The immunoglobulin molecules provided herein can be of any class (*e.g.*, IgG, IgE, IgM, IgD, and IgA) or any subclass (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2) of immunoglobulin molecule.

**[0344]** Variants and derivatives of antibodies include antibody functional fragments that retain the ability to bind to a PD-1 epitope. Exemplary functional fragments include Fab fragments (*e.g.*, an antibody fragment that contains the antigen-binding domain and comprises a light chain and part of a heavy chain bridged by a disulfide bond); Fab' (*e.g.*, an antibody fragment containing a single antigen-binding domain comprising an Fab and an additional portion of the heavy chain through the hinge region); F(ab')<sub>2</sub> (*e.g.*, two Fab' molecules joined by interchain disulfide bonds in the hinge regions of the heavy chains; the Fab' molecules may be directed toward the same or different epitopes); a bispecific Fab (*e.g.*, a Fab molecule having two antigen binding domains, each of which may be directed to a different epitope); a single chain comprising a variable region, also known as, scFv (*e.g.*, the variable, antigen-binding

determinative region of a single light and heavy chain of an antibody linked together by a chain of 10-25 amino acids); a disulfide-linked Fv, or dsFv (e.g., the variable, antigen-binding determinative region of a single light and heavy chain of an antibody linked together by a disulfide bond); a camelized VH (e.g., the variable, antigen-binding determinative region of a single heavy chain of an antibody in which some amino acids at the VH interface are those found in the heavy chain of naturally occurring camel antibodies); a bispecific scFv (e.g., an scFv or a dsFv molecule having two antigen-binding domains, each of which may be directed to a different epitope); a diabody (e.g., a dimerized scFv formed when the VH domain of a first scFv assembles with the VL domain of a second scFv and the VL domain of the first scFv assembles with the VH domain of the second scFv; the two antigen-binding regions of the diabody may be directed towards the same or different epitopes); and a triabody (e.g., a trimerized scFv, formed in a manner similar to a diabody, but in which three antigen-binding domains are created in a single complex; the three antigen-binding domains may be directed towards the same or different epitopes).

#### 4.3.1.4 Humanized Antibodies

**[0345]** In some embodiments, antibodies provided herein can be humanized antibodies that bind PD-1, including human and/or cynomolgus PD-1. For example, humanized antibodies of the present disclosure may comprise one or more CDRs as shown in **Tables 1-2**. Various methods for humanizing non-human antibodies are known in the art. For example, a humanized antibody can have one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as “import” residues, which are typically taken from an “import” variable domain. Humanization may be performed, for example, following the method of Jones *et al.*, 1986, *Nature* 321:522-25; Riechmann *et al.*, 1988, *Nature* 332:323-27; and Verhoeyen *et al.*, 1988, *Science* 239:1534-36), by substituting hypervariable region sequences for the corresponding sequences of a human antibody.

**[0346]** In some cases, the humanized antibodies are constructed by CDR grafting, in which the amino acid sequences of the six CDRs of the parent non-human antibody (e.g., rodent) are grafted onto a human antibody framework. For example, Padlan *et al.* determined that only about one third of the residues in the CDRs actually contact the antigen, and termed these the “specificity determining residues,” or SDRs (Padlan *et al.*, 1995, *FASEB J.* 9:133-39). In the

technique of SDR grafting, only the SDR residues are grafted onto the human antibody framework (see, e.g., Kashmiri *et al.*, 2005, Methods 36:25-34).

**[0347]** The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies can be important to reduce antigenicity. For example, according to the so-called “best-fit” method, the sequence of the variable domain of a non-human (e.g., rodent) antibody is screened against the entire library of known human variable-domain sequences. The human sequence that is closest to that of the rodent may be selected as the human framework for the humanized antibody (Sims *et al.*, 1993, J. Immunol. 151:2296-308; and Chothia *et al.*, 1987, J. Mol. Biol. 196:901-17). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:4285-89; and Presta *et al.*, 1993, J. Immunol. 151:2623-32). In some cases, the framework is derived from the consensus sequences of the most abundant human subclasses, V<sub>L</sub>6 subgroup I (V<sub>L</sub>6I) and V<sub>H</sub> subgroup III (V<sub>H</sub>III). In another method, human germline genes are used as the source of the framework regions.

**[0348]** In an alternative paradigm based on comparison of CDRs, called superhumanization, FR homology is irrelevant. The method consists of comparison of the non-human sequence with the functional human germline gene repertoire. Those genes encoding the same or closely related canonical structures to the murine sequences are then selected. Next, within the genes sharing the canonical structures with the non-human antibody, those with highest homology within the CDRs are chosen as FR donors. Finally, the non-human CDRs are grafted onto these FRs (see, e.g., Tan *et al.*, 2002, J. Immunol. 169:1119-25).

**[0349]** It is further generally desirable that antibodies be humanized with retention of their affinity for the antigen and other favorable biological properties. To achieve this goal, according to one method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. These include, for example, WAM (Whitelegg and Rees, 2000, Protein Eng. 13:819-24), Modeller (Sali and Blundell, 1993, J. Mol. Biol. 234:779-815), and

Swiss PDB Viewer (Guex and Peitsch, 1997, Electrophoresis 18:2714-23). Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *e.g.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

**[0350]** Another method for antibody humanization is based on a metric of antibody humanness termed Human String Content (HSC). This method compares the mouse sequence with the repertoire of human germline genes, and the differences are scored as HSC. The target sequence is then humanized by maximizing its HSC rather than using a global identity measure to generate multiple diverse humanized variants (Lazar *et al.*, 2007, Mol. Immunol. 44:1986-98).

**[0351]** In addition to the methods described above, empirical methods may be used to generate and select humanized antibodies. These methods include those that are based upon the generation of large libraries of humanized variants and selection of the best clones using enrichment technologies or high throughput screening techniques. Antibody variants may be isolated from phage, ribosome, and yeast display libraries as well as by bacterial colony screening (*see, e.g.*, Hoogenboom, 2005, Nat. Biotechnol. 23:1105-16; Dufner *et al.*, 2006, Trends Biotechnol. 24:523-29; Feldhaus *et al.*, 2003, Nat. Biotechnol. 21:163-70; and Schlapschy *et al.*, 2004, Protein Eng. Des. Sel. 17:847-60).

**[0352]** In the FR library approach, a collection of residue variants are introduced at specific positions in the FR followed by screening of the library to select the FR that best supports the grafted CDR. The residues to be substituted may include some or all of the “Vernier” residues identified as potentially contributing to CDR structure (*see, e.g.*, Foote and Winter, 1992, J. Mol. Biol. 224:487-99), or from the more limited set of target residues identified by Baca *et al.* (1997, J. Biol. Chem. 272:10678-84).

**[0353]** In FR shuffling, whole FRs are combined with the non-human CDRs instead of creating combinatorial libraries of selected residue variants (*see, e.g.*, Dall’Acqua *et al.*, 2005, Methods 36:43-60). The libraries may be screened for binding in a two-step process, first humanizing VL, followed by VH. Alternatively, a one-step FR shuffling process may be used. Such a process has been shown to be more efficient than the two-step screening, as the resulting

antibodies exhibited improved biochemical and physicochemical properties including enhanced expression, increased affinity, and thermal stability (see, e.g., Damschroder *et al.*, 2007, *Mol. Immunol.* 44:3049-60).

**[0354]** The “humaneering” method is based on experimental identification of essential minimum specificity determinants (MSDs) and is based on sequential replacement of non-human fragments into libraries of human FRs and assessment of binding. It begins with regions of the CDR3 of non-human VH and VL chains and progressively replaces other regions of the non-human antibody into the human FRs, including the CDR1 and CDR2 of both VH and VL. This methodology typically results in epitope retention and identification of antibodies from multiple subclasses with distinct human V-segment CDRs. Humaneering allows for isolation of antibodies that are 91-96% homologous to human germline gene antibodies (see, e.g., Alfenito, Cambridge Healthtech Institute’s Third Annual PEGS, The Protein Engineering Summit, 2007).

**[0355]** The “human engineering” method involves altering a non-human antibody or antibody fragment, such as a mouse or chimeric antibody or antibody fragment, by making specific changes to the amino acid sequence of the antibody so as to produce a modified antibody with reduced immunogenicity in a human that nonetheless retains the desirable binding properties of the original non-human antibodies. Generally, the technique involves classifying amino acid residues of a non-human (e.g., mouse) antibody as “low risk,” “moderate risk,” or “high risk” residues. The classification is performed using a global risk/reward calculation that evaluates the predicted benefits of making particular substitution (e.g., for immunogenicity in humans) against the risk that the substitution will affect the resulting antibody’s folding. The particular human amino acid residue to be substituted at a given position (e.g., low or moderate risk) of a non-human (e.g., mouse) antibody sequence can be selected by aligning an amino acid sequence from the non-human antibody’s variable regions with the corresponding region of a specific or consensus human antibody sequence. The amino acid residues at low or moderate risk positions in the non-human sequence can be substituted for the corresponding residues in the human antibody sequence according to the alignment. Techniques for making human engineered proteins are described in greater detail in Studnicka *et al.*, 1994, *Protein Engineering* 7:805-14; U.S. Pat. Nos. 5,766,886; 5,770,196; 5,821,123; and 5,869,619; and PCT Publication WO 93/11794.

#### 4.3.1.5 Human Antibodies

**[0356]** Human anti-PD-1 antibodies can be constructed by combining Fv clone variable domain sequence(s) selected from human-derived phage display libraries with known human constant domain sequences(s). Alternatively, human monoclonal anti-PD-1 antibodies of the present disclosure can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described, for example, by Kozbor, 1984, *J. Immunol.* 133:3001-05; Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications 51-63 (1987); and Boerner *et al.*, 1991, *J. Immunol.* 147:86-95.

**[0357]** It is also possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. Transgenic mice that express human antibody repertoires have been used to generate high-affinity human sequence monoclonal antibodies against a wide variety of potential drug targets (*see, e.g.*, Jakobovits, A., 1995, *Curr. Opin. Biotechnol.* 6(5):561-66; Brüggemann and Taussing, 1997, *Curr. Opin. Biotechnol.* 8(4):455-58; U.S. Pat. Nos. 6,075,181 and 6,150,584; and Lonberg *et al.*, 2005, *Nature Biotechnol.* 23:1117-25).

**[0358]** Alternatively, the human antibody may be prepared via immortalization of human B lymphocytes producing an antibody directed against a target antigen (*e.g.*, such B lymphocytes may be recovered from an individual or may have been immunized *in vitro*) (*see, e.g.*, Cole *et al.*, Monoclonal Antibodies and Cancer Therapy (1985); Boerner *et al.*, 1991, *J. Immunol.* 147(1):86-95; and U.S. Pat. No. 5,750,373).

**[0359]** Gene shuffling can also be used to derive human antibodies from non-human, for example, rodent, antibodies, where the human antibody has similar affinities and specificities to the starting non-human antibody. According to this method, which is also called “epitope imprinting” or “guided selection,” either the heavy or light chain variable region of a non-human antibody fragment obtained by phage display techniques as described herein is replaced with a repertoire of human V domain genes, creating a population of non-human chain/human chain scFv or Fab chimeras. Selection with antigen results in isolation of a non-human chain/human chain chimeric scFv or Fab wherein the human chain restores the antigen binding site destroyed upon removal of the corresponding non-human chain in the primary phage display clone (*e.g.*, the epitope guides (imprints) the choice of the human chain partner). When the process is

repeated in order to replace the remaining non-human chain, a human antibody is obtained (*see, e.g.*, PCT WO 93/06213; and Osbourn *et al.*, 2005, Methods 36:61-68). Unlike traditional humanization of non-human antibodies by CDR grafting, this technique provides completely human antibodies, which have no FR or CDR residues of non-human origin. Examples of guided selection to humanize mouse antibodies towards cell surface antigens include the folate-binding protein present on ovarian cancer cells (*see, e.g.*, Figini *et al.*, 1998, Cancer Res. 58:991-96) and CD147, which is highly expressed on hepatocellular carcinoma (*see, e.g.*, Bao *et al.*, 2005, Cancer Biol. Ther. 4:1374-80).

**[0360]** A potential disadvantage of the guided selection approach is that shuffling of one antibody chain while keeping the other constant could result in epitope drift. In order to maintain the epitope recognized by the non-human antibody, CDR retention can be applied (*see, e.g.*, Klimka *et al.*, 2000, Br. J. Cancer. 83:252-60; and Beiboeer *et al.*, 2000, J. Mol. Biol. 296:833-49). In this method, the non-human VH CDR3 is commonly retained, as this CDR may be at the center of the antigen-binding site and may be the most important region of the antibody for antigen recognition. In some instances, however, VH CDR3 and VL CDR3, as well as VH CDR2, VL CDR2, and VL CDR1 of the non-human antibody may be retained.

#### **4.3.1.6 Bispecific Antibodies**

**[0361]** Bispecific antibodies are monoclonal antibodies that have binding specificities for at least two different antigens. In certain embodiments, bispecific antibodies are human or humanized antibodies. In certain embodiments, one of the binding specificities is for PD-1 and the other is for any other antigen. In some embodiments, one of the binding specificities is for PD-1, and the other is for another surface antigen expressed on cells expressing PD-1. In certain embodiments, bispecific antibodies may bind to two different epitopes of PD-1. Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.*, F(ab')<sub>2</sub>) bispecific antibodies).

**[0362]** Methods for making bispecific antibodies are known in the art, such as, by co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (*see, e.g.*, Milstein and Cuello, 1983, Nature 305:537-40). For further details of generating bispecific antibodies, see, for example, Bispecific Antibodies (Kontermann ed., 2011).

#### **4.3.1.7 Multivalent Antibodies**

**[0363]** A multivalent antibody may be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present disclosure can be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (*e.g.*, tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or more antigen binding sites. In certain embodiments, the dimerization domain comprises (or consists of) an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. In certain embodiments, a multivalent antibody comprises (or consists of) three to about eight antigen binding sites. In one such embodiment, a multivalent antibody comprises (or consists of) four antigen binding sites. The multivalent antibody comprises at least one polypeptide chain (*e.g.*, two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise VD1-(X1)<sub>n</sub>-VD2-(X2)<sub>n</sub>-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. For instance, the polypeptide chain(s) may comprise: VH-CH1-flexible linker-VH-CH1-Fc region chain; or VH-CH1-VH-CH1-Fc region chain. The multivalent antibody herein may further comprise at least two (*e.g.*, four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

#### **4.3.1.8 Fc Engineering**

**[0364]** It may be desirable to modify an anti-PD-1 antibody provided herein by Fc engineering. In certain embodiments, the modification to the Fc region of the antibody results in the decrease or elimination of an effector function of the antibody. In certain embodiments, the effector function is ADCC, ADCP, and/or CDC. In some embodiments, the effector function is ADCC. In other embodiments, the effector function is ADCP. In other embodiments, the effector function is CDC. In one embodiment, the effector function is ADCC and ADCP. In one embodiment, the effector function is ADCC and CDC. In one embodiment, the effector function

is ADCP and CDC. In one embodiment, the effector function is ADCC, ADCP and CDC. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. For example, substitutions into human IgG1 using IgG2 residues at positions 233-236 and IgG4 residues at positions 327, 330, and 331 were shown to greatly reduce ADCC and CDC (see, e.g., Armour *et al.*, 1999, Eur. J. Immunol. 29(8):2613-24; and Shields *et al.*, 2001, J. Biol. Chem. 276(9): 6591-604). Other Fc variants are provided elsewhere herein.

**[0365]** To increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment), for example, as described in U.S. Pat. No. 5,739,277. Term “salvage receptor binding epitope” refers to an epitope of the Fc region of an IgG molecule (e.g., IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

#### **4.3.1.9** Alternative Binding Agents

**[0366]** The present disclosure encompasses non-immunoglobulin binding agents that specifically bind to the same epitope as an anti-PD-1 antibody disclosed herein. In some embodiments, a non-immunoglobulin binding agent is identified as an agent that displaces or is displaced by an anti-PD-1 antibody of the present disclosure in a competitive binding assay. These alternative binding agents may include, for example, any of the engineered protein scaffolds known in the art. Such scaffolds may comprise one or more CDRs as shown in **Tables 1-2**. Such scaffolds include, for example, anticalins, which are based upon the lipocalin scaffold, a protein structure characterized by a rigid beta-barrel that supports four hypervariable loops which form the ligand binding site. Novel binding specificities may be engineered by targeted random mutagenesis in the loop regions, in combination with functional display and guided selection (see, e.g., Skerra, 2008, FEBS J. 275:2677-83). Other suitable scaffolds may include, for example, adnectins, or monobodies, based on the tenth extracellular domain of human fibronectin III (see, e.g., Koide and Koide, 2007, Methods Mol. Biol. 352: 95-109); affibodies, based on the Z domain of staphylococcal protein A (see, e.g., Nygren *et al.*, 2008, FEBS J. 275:2668-76); DARPinS, based on ankyrin repeat proteins (see, e.g., Stumpp *et al.*, 2008, Drug. Discov. Today 13:695-701); fynomers, based on the SH3 domain of the human Fyn protein kinase (see, e.g., Grabulovski *et al.*, 2007, J. Biol. Chem. 282:3196-204); affitins, based on Sac7d from *Sulfolobus acidolarius* (see, e.g., Krehenbrink *et al.*, 2008, J. Mol. Biol. 383:1058-68); affilins, based on human  $\gamma$ -B-crystallin (see, e.g., Ebersbach *et al.*, 2007, J. Mol.

Biol. 372:172-85); avimers, based on the A domain of membrane receptor proteins (*see, e.g.*, Silverman *et al.*, 2005, Biotechnol. 23:1556-61); cysteine-rich knottin peptides (*see, e.g.*, Kolmar, 2008, FEBS J. 275:2684-90); and engineered Kunitz-type inhibitors (*see, e.g.*, Nixon and Wood, 2006, Curr. Opin. Drug. Discov. Dev. 9:261-68). For a review, see, for example, Gebauer and Skerra, 2009, Curr. Opin. Chem. Biol. 13:245-55.

#### 4.3.2 Antibody variants

**[0367]** In some embodiments, amino acid sequence modification(s) of the antibodies that bind to PD-1 or described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody, including but not limited to specificity, thermostability, expression level, effector functions, glycosylation, reduced immunogenicity, or solubility. Thus, in addition to the anti-PD-1 antibodies provided herein, it is contemplated that anti-PD-1 antibody variants can be prepared. For example, anti-PD-1 antibody variants can be prepared by introducing appropriate nucleotide changes into the encoding DNA, and/or by synthesis of the desired antibody or polypeptide. Those skilled in the art who appreciate that amino acid changes may alter post-translational processes of the anti-PD-1 antibody, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

**[0368]** In some embodiments, antibodies provided herein are chemically modified, for example, by the covalent attachment of any type of molecule to the antibody. The antibody derivatives may include antibodies that have been chemically modified, for example, by increase or decrease of glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, chemical cleavage, proteolytic cleavage, linkage to a cellular ligand or other protein, *etc.* Additionally, the antibody may contain one or more non-classical amino acids.

**[0369]** Variations may be a substitution, deletion, or insertion of one or more codons encoding the antibody or polypeptide that results in a change in the amino acid sequence as compared with the native sequence antibody or polypeptide. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *e.g.*, conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. In certain embodiments, the substitution, deletion, or insertion includes fewer than 25

amino acid substitutions, fewer than 20 amino acid substitutions, fewer than 15 amino acid substitutions, fewer than 10 amino acid substitutions, fewer than 5 amino acid substitutions, fewer than 4 amino acid substitutions, fewer than 3 amino acid substitutions, or fewer than 2 amino acid substitutions relative to the original molecule. In a specific embodiment, the substitution is a conservative amino acid substitution made at one or more predicted non-essential amino acid residues. The variation allowed may be determined by systematically making insertions, deletions, or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

**[0370]** Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.*, for antibody-directed enzyme prodrug therapy) or a polypeptide which increases the serum half-life of the antibody.

**[0371]** Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Alternatively, conservative (*e.g.*, within an amino acid group with similar properties and/or side chains) substitutions may be made, so as to maintain or not significantly change the properties. Amino acids may be grouped according to similarities in the properties of their side chains (*see, e.g.*, Lehninger, Biochemistry 73-75 (2d ed. 1975)):

- (1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M);
- (2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q);
- (3) acidic: Asp (D), Glu (E); and
- (4) basic: Lys (K), Arg (R), His (H).

**[0372]** Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties: (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: Asp, Glu; (4) basic: His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6) aromatic: Trp, Tyr, Phe.

**[0373]** Non-conservative substitutions entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, into the remaining (non-conserved) sites. Accordingly, in one embodiment, an antibody or fragment thereof that binds to a PD-1 epitope comprises an amino acid sequence that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of a murine monoclonal antibody provided herein. In one embodiment, an antibody or fragment thereof that binds to a PD-1 epitope comprises an amino acid sequence that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to an amino acid sequence depicted in **Tables 1-6**. In yet another embodiment, an antibody or fragment thereof that binds to a PD-1 epitope comprises a VH CDR and/or a VL CDR amino acid sequence that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to a VH CDR amino acid sequence depicted in **Table 2** and/or a VL CDR amino acid sequence depicted in **Table 1**. The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (see, e.g., Carter, 1986, *Biochem J.* 237:1-7; and Zoller *et al.*, 1982, *Nucl. Acids Res.* 10:6487-500), cassette mutagenesis (see, e.g., Wells *et al.*, 1985, *Gene* 34:315-23), or other known techniques can be performed on the cloned DNA to produce the anti-PD-1 antibody variant DNA.

**[0374]** Any cysteine residue not involved in maintaining the proper conformation of the anti-PD-1 antibody also may be substituted, for example, with another amino acid, such as alanine or serine, to improve the oxidative stability of the molecule and to prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the anti-PD-1 antibody to improve its stability (e.g., where the antibody is an antibody fragment such as an Fv fragment).

**[0375]** In some embodiments, an anti-PD-1 antibody molecule of the present disclosure is a “de-immunized” antibody. A “de-immunized” anti-PD-1 antibody is an antibody derived from a humanized or chimeric anti-PD-1 antibody, which has one or more alterations in its amino acid sequence resulting in a reduction of immunogenicity of the antibody, compared to the respective

original non-de-immunized antibody. One of the procedures for generating such antibody mutants involves the identification and removal of T cell epitopes of the antibody molecule. In a first step, the immunogenicity of the antibody molecule can be determined by several methods, for example, by *in vitro* determination of T cell epitopes or *in silico* prediction of such epitopes, as known in the art. Once the critical residues for T cell epitope function have been identified, mutations can be made to remove immunogenicity and retain antibody activity. For review, see, for example, Jones *et al.*, 2009, Methods in Molecular Biology 525:405-23.

#### **4.3.2.1 In vitro Affinity Maturation**

**[0376]** In some embodiments, antibody variants having an improved property such as affinity, stability, or expression level as compared to a parent antibody may be prepared by *in vitro* affinity maturation. Like the natural prototype, *in vitro* affinity maturation is based on the principles of mutation and selection. Libraries of antibodies are displayed as Fab, scFv, or V domain fragments either on the surface of an organism (e.g., phage, bacteria, yeast, or mammalian cell) or in association (e.g., covalently or non-covalently) with their encoding mRNA or DNA. Affinity selection of the displayed antibodies allows isolation of organisms or complexes carrying the genetic information encoding the antibodies. Two or three rounds of mutation and selection using display methods such as phage display usually results in antibody fragments with affinities in the low nanomolar range. Affinity matured antibodies can have nanomolar or even picomolar affinities for the target antigen.

**[0377]** Phage display is a widespread method for display and selection of antibodies. The antibodies are displayed on the surface of Fd or M13 bacteriophages as fusions to the bacteriophage coat protein. Selection involves exposure to antigen to allow phage-displayed antibodies to bind their targets, a process referred to as “panning.” Phage bound to antigen are recovered and used to infect bacteria to produce phage for further rounds of selection. For review, see, for example, Hoogenboom, 2002, Methods. Mol. Biol. 178:1-37; and Bradbury and Marks, 2004, J. Immunol. Methods 290:29-49.

**[0378]** In a yeast display system (see, e.g., Boder *et al.*, 1997, Nat. Biotech. 15:553-57; and Chao *et al.*, 2006, Nat. Protocols 1:755-68), the antibody may be displayed as single-chain variable fusions (scFv) in which the heavy and light chains are connected by a flexible linker. The scFv is fused to the adhesion subunit of the yeast agglutinin protein Aga2p, which attaches to the yeast cell wall through disulfide bonds to Aga1p. Display of a protein via Aga2p projects

the protein away from the cell surface, minimizing potential interactions with other molecules on the yeast cell wall. Magnetic separation and flow cytometry are used to screen the library to select for antibodies with improved affinity or stability. Binding to a soluble antigen of interest is determined by labeling of yeast with biotinylated antigen and a secondary reagent such as streptavidin conjugated to a fluorophore. Variations in surface expression of the antibody can be measured through immunofluorescence labeling of either the hemagglutinin or c-Myc epitope tag flanking the scFv. Expression has been shown to correlate with the stability of the displayed protein, and thus antibodies can be selected for improved stability as well as affinity (see, e.g., Shusta *et al.*, 1999, *J. Mol. Biol.* 292:949-56). An additional advantage of yeast display is that displayed proteins are folded in the endoplasmic reticulum of the eukaryotic yeast cells, taking advantage of endoplasmic reticulum chaperones and quality-control machinery. Once maturation is complete, antibody affinity can be conveniently “titrated” while displayed on the surface of the yeast, eliminating the need for expression and purification of each clone. A theoretical limitation of yeast surface display is the potentially smaller functional library size than that of other display methods; however, a recent approach uses the yeast cells’ mating system to create combinatorial diversity estimated to be  $10^{14}$  in size (see, e.g., U.S. Pat. Publication 2003/0186374; and Blaise *et al.*, 2004, *Gene* 342:211–18).

**[0379]** In ribosome display, antibody-ribosome-mRNA (ARM) complexes are generated for selection in a cell-free system. The DNA library coding for a particular library of antibodies is genetically fused to a spacer sequence lacking a stop codon. This spacer sequence, when translated, is still attached to the peptidyl tRNA and occupies the ribosomal tunnel, and thus allows the protein of interest to protrude out of the ribosome and fold. The resulting complex of mRNA, ribosome, and protein can bind to surface-bound ligand, allowing simultaneous isolation of the antibody and its encoding mRNA through affinity capture with the ligand. The ribosome-bound mRNA is then reverse transcribed back into cDNA, which can then undergo mutagenesis and be used in the next round of selection (see, e.g., Fukuda *et al.*, 2006, *Nucleic Acids Res.* 34:e127). In mRNA display, a covalent bond between antibody and mRNA is established using puromycin as an adaptor molecule (Wilson *et al.*, 2001, *Proc. Natl. Acad. Sci. USA* 98:3750-55).

**[0380]** As these methods are performed entirely *in vitro*, they provide two main advantages over other selection technologies. First, the diversity of the library is not limited by the transformation efficiency of bacterial cells, but only by the number of ribosomes and different

mRNA molecules present in the test tube. Second, random mutations can be introduced easily after each selection round, for example, by non-proofreading polymerases, as no library must be transformed after any diversification step.

**[0381]** In a mammalian cell display system (see, e.g., Bowers *et al.*, 2011, Proc Natl Acad Sci USA. 108:20455-60), a fully human library of IgGs is constructed based on germline sequence V-gene segments joined to prerecombined D(J) regions. Full-length V regions for heavy chain and light chain are assembled with human heavy chain and light chain constant regions and transfected into a mammalian cell line (e.g., HEK293). The transfected library is expanded and subjected to several rounds of negative selection against streptavidin (SA)-coupled magnetic beads, followed by a round of positive selection against SA-coupled magnetic beads coated with biotinylated target protein, peptide fragment, or epitope. Positively selected cells are expanded, and then sorted by rounds of FACS to isolate single cell clones displaying antibodies that specifically bind to the target protein, peptide fragment, or epitope. Heavy and light chain pairs from these single cell clones are retransfected with AID for further maturation. Several rounds of mammalian cell display, coupled with AID-triggered somatic hypermutation, generate high specificity, high affinity antibodies.

**[0382]** Diversity may also be introduced into the CDRs or the whole V genes of the antibody libraries in a targeted manner or via random introduction. The former approach includes sequentially targeting all the CDRs of an antibody via a high or low level of mutagenesis or targeting isolated hot spots of somatic hypermutations (see, e.g., Ho *et al.*, 2005, J. Biol. Chem. 280:607-17) or residues suspected of affecting affinity on experimental basis or structural reasons. In a specific embodiment, somatic hypermutation is performed by AID-triggered somatic hypermutation, e.g., using the SHM-XEL<sup>TM</sup> platform (AnaptysBio, San Diego, CA). Random mutations can be introduced throughout the whole V gene using *E. coli* mutator strains, error-prone replication with DNA polymerases (see, e.g., Hawkins *et al.*, 1992, J. Mol. Biol. 226:889-96), or RNA replicases. Diversity may also be introduced by replacement of regions that are naturally diverse via DNA shuffling or similar techniques (see, e.g., Lu *et al.*, 2003, J. Biol. Chem. 278:43496-507; U.S. Pat. Nos. 5,565,332 and 6,989,250). Alternative techniques target hypervariable loops extending into framework-region residues (see, e.g., Bond *et al.*, 2005, J. Mol. Biol. 348:699-709) employ loop deletions and insertions in CDRs or use hybridization-based diversification (see, e.g., U.S. Pat. Publication No. 2004/0005709). Additional methods of

generating diversity in CDRs are disclosed, for example, in U.S. Pat. No. 7,985,840. Further methods that can be used to generate antibody libraries and/or antibody affinity maturation are disclosed, *e.g.*, in U.S. Patent Nos. 8,685,897 and 8,603,930, and U.S. Publ. Nos. 2014/0170705, 2014/0094392, 2012/0028301, 2011/0183855, and 2009/0075378, each of which are incorporated herein by reference.

**[0383]** Screening of the libraries can be accomplished by various techniques known in the art. For example, PD-1 can be immobilized onto solid supports, columns, pins, or cellulose/poly(vinylidene fluoride) membranes/other filters, expressed on host cells affixed to adsorption plates or used in cell sorting, or conjugated to biotin for capture with streptavidin-coated beads or used in any other method for panning display libraries.

**[0384]** For review of *in vitro* affinity maturation methods, see, *e.g.*, Hoogenboom, 2005, Nature Biotechnology 23:1105-16; Quiroz and Sinclair, 2010, Revista Ingeneria Biomedia 4:39-51; and references therein.

#### **4.3.2.2 Modifications of Anti-PD-1 Antibodies**

**[0385]** Covalent modifications of anti-PD-1 antibodies are included within the scope of the present disclosure. Covalent modifications include reacting targeted amino acid residues of an anti-PD-1 antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the anti-PD-1 antibody. Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains (*see, e.g.*, Creighton, Proteins: Structure and Molecular Properties 79-86 (1983)), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

**[0386]** Other types of covalent modification of the anti-PD-1 antibody included within the scope of this present disclosure include altering the native glycosylation pattern of the antibody or polypeptide (*see, e.g.*, Beck *et al.*, 2008, Curr. Pharm. Biotechnol. 9:482-501; and Walsh, 2010, Drug Discov. Today 15:773-80), and linking the antibody to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth, for example, in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192; or 4,179,337.

**[0387]** An anti-PD-1 antibody of the present disclosure may also be modified to form chimeric molecules comprising an anti-PD-1 antibody fused to another, heterologous polypeptide or amino acid sequence, for example, an epitope tag (*see, e.g.*, Terpe, 2003, *Appl. Microbiol. Biotechnol.* 60:523-33) or the Fc region of an IgG molecule (*see, e.g.*, Aruffo, Antibody Fusion Proteins 221-42 (Chamow and Ashkenazi eds., 1999)).

**[0388]** Also provided herein are fusion proteins comprising an antibody provided herein that binds to a PD-1 antigen and a heterologous polypeptide. In some embodiments, the heterologous polypeptide to which the antibody is fused is useful for targeting the antibody to cells having cell surface-expressed PD-1.

**[0389]** Also provided herein are panels of antibodies that bind to a PD-1 antigen. In specific embodiments, the panels of antibodies have different association rates, different dissociation rates, different affinities for a PD-1 antigen, and/or different specificities for a PD-1 antigen. In some embodiments, the panels comprise or consist of about 10, about 25, about 50, about 75, about 100, about 125, about 150, about 175, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 antibodies or more. Panels of antibodies can be used, for example, in 96-well or 384-well plates, for assays such as ELISAs.

#### 4.3.3 Preparation of anti-PD-1 antibodies

**[0390]** Anti-PD-1 antibodies may be produced by culturing cells transformed or transfected with a vector containing anti-PD-1 antibody-encoding nucleic acids. Polynucleotide sequences encoding polypeptide components of the antibody of the present disclosure can be obtained using standard recombinant techniques. Desired polynucleotide sequences may be isolated and sequenced from antibody producing cells such as hybridomas cells. Alternatively, polynucleotides can be synthesized using nucleotide synthesizer or PCR techniques. Once obtained, sequences encoding the polypeptides are inserted into a recombinant vector capable of replicating and expressing heterologous polynucleotides in host cells. Many vectors that are available and known in the art can be used for the purpose of the present disclosure. Selection of an appropriate vector will depend mainly on the size of the nucleic acids to be inserted into the vector and the particular host cell to be transformed with the vector. Host cells suitable for expressing antibodies of the present disclosure include prokaryotes such as Archaeabacteria and Eubacteria, including Gram-negative or Gram-positive organisms, eukaryotic microbes such as

filamentous fungi or yeast, invertebrate cells such as insect or plant cells, and vertebrate cells such as mammalian host cell lines. Host cells are transformed with the above-described expression vectors and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Antibodies produced by the host cells are purified using standard protein purification methods as known in the art.

**[0391]** Methods for antibody production including vector construction, expression, and purification are further described in Plückthun *et al.*, Antibody Engineering: Producing antibodies in Escherichia coli: From PCR to fermentation 203-52 (McCafferty *et al.* eds., 1996); Kwong and Rader, *E. coli Expression and Purification of Fab Antibody Fragments*, in Current Protocols in Protein Science (2009); Tachibana and Takekoshi, *Production of Antibody Fab Fragments in Escherichia coli*, in Antibody Expression and Production (Al-Rubeai ed., 2011); and Therapeutic Monoclonal Antibodies: From Bench to Clinic (An ed., 2009).

**[0392]** It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare anti-PD-1 antibodies. For instance, the appropriate amino acid sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques (see, e.g., Stewart *et al.*, Solid-Phase Peptide Synthesis (1969); and Merrifield, 1963, J. Am. Chem. Soc. 85:2149-54). *In vitro* protein synthesis may be performed using manual techniques or by automation. Various portions of the anti-PD-1 antibody may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the desired anti-PD-1 antibody. Alternatively, antibodies may be purified from cells or bodily fluids, such as milk, of a transgenic animal engineered to express the antibody, as disclosed, for example, in U.S. Pat. Nos. 5,545,807 and 5,827,690.

#### 4.3.4 Immunoconjugates

**[0393]** The present disclosure also provides conjugates comprising any one of the anti-PD-1 antibodies of the present disclosure covalently bound by a synthetic linker to one or more non-antibody agents.

**[0394]** In some embodiments, antibodies provided herein are conjugated or recombinantly fused, e.g., to a diagnostic or detectable molecule. The conjugated or recombinantly fused antibodies can be useful, for example, for monitoring or prognosing the onset, development, progression, and/or severity of a PD-1-mediated disease.

**[0395]** Such diagnosis and detection can be accomplished, for example, by coupling the antibody to detectable substances including, but not limited to, various enzymes, such as, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as, but not limited to, streptavidin/biotin or avidin/biotin; fluorescent materials, such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride, or phycoerythrin; luminescent materials, such as, but not limited to, luminol; bioluminescent materials, such as, but not limited to, luciferase, luciferin, or aequorin; chemiluminescent material, such as, but not limited to, an acridinium based compound or a HALOTAG; radioactive materials, such as, but not limited to, iodine ( $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ , and  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{115}\text{In}$ ,  $^{113}\text{In}$ ,  $^{112}\text{In}$ , and  $^{111}\text{In}$ ), technetium ( $^{99}\text{Tc}$ ), thallium ( $^{201}\text{Tl}$ ), gallium ( $^{68}\text{Ga}$  and  $^{67}\text{Ga}$ ), palladium ( $^{103}\text{Pd}$ ), molybdenum ( $^{99}\text{Mo}$ ), xenon ( $^{133}\text{Xe}$ ), fluorine ( $^{18}\text{F}$ ),  $^{153}\text{Sm}$ ,  $^{177}\text{Lu}$ ,  $^{159}\text{Gd}$ ,  $^{149}\text{Pm}$ ,  $^{140}\text{La}$ ,  $^{175}\text{Yb}$ ,  $^{166}\text{Ho}$ ,  $^{90}\text{Y}$ ,  $^{47}\text{Sc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{142}\text{Pr}$ ,  $^{105}\text{Rh}$ ,  $^{97}\text{Ru}$ ,  $^{68}\text{Ge}$ ,  $^{57}\text{Co}$ ,  $^{65}\text{Zn}$ ,  $^{85}\text{Sr}$ ,  $^{32}\text{P}$ ,  $^{153}\text{Gd}$ ,  $^{169}\text{Yb}$ ,  $^{51}\text{Cr}$ ,  $^{54}\text{Mn}$ ,  $^{75}\text{Se}$ ,  $^{113}\text{Sn}$ , or  $^{117}\text{Sn}$ ; positron emitting metals using various positron emission tomographies; and non-radioactive paramagnetic metal ions.

**[0396]** Also provided herein are antibodies that are recombinantly fused or chemically conjugated (covalent or non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, for example, to a polypeptide of about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, or about 100 amino acids) to generate fusion proteins, as well as uses thereof. In particular, provided herein are fusion proteins comprising an antigen-binding fragment of an antibody provided herein (e.g., a Fab fragment, Fc fragment, Fv fragment, F(ab)<sub>2</sub> fragment, a VH domain, a VH CDR, a VL domain, or a VL CDR) and a heterologous protein, polypeptide, or peptide. In one embodiment, the heterologous protein, polypeptide, or peptide that the antibody is fused to is useful for targeting the antibody to a particular cell type, such as a cell that expresses PD-1. For example, an antibody that binds to a cell surface receptor expressed by a particular cell type may be fused or conjugated to a modified antibody provided herein.

**[0397]** Moreover, antibodies provided herein can be fused to marker or “tag” sequences, such as a peptide, to facilitate purification. In specific embodiments, the marker or tag amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (see, e.g., QIAGEN, Inc.), among others, many of which are commercially available. For example, as

described in Gentz *et al.*, 1989, Proc. Natl. Acad. Sci. USA 86:821-24, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin (“HA”) tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, 1984, Cell 37:767-78), and the “FLAG” tag.

**[0398]** Methods for fusing or conjugating moieties (including polypeptides) to antibodies are known (see, e.g., Arnon *et al.*, *Monoclonal Antibodies for Immunotargeting of Drugs in Cancer Therapy*, in Monoclonal Antibodies and Cancer Therapy 243-56 (Reisfeld *et al.* eds., 1985); Hellstrom *et al.*, *Antibodies for Drug Delivery*, in Controlled Drug Delivery 623-53 (Robinson *et al.* eds., 2d ed. 1987); Thorpe, *Antibody Carriers of Cytotoxic Agents in Cancer Therapy: A Review*, in Monoclonal Antibodies: Biological and Clinical Applications 475-506 (Pinchera *et al.* eds., 1985); *Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy*, in Monoclonal Antibodies for Cancer Detection and Therapy 303-16 (Baldwin *et al.* eds., 1985); Thorpe *et al.*, 1982, Immunol. Rev. 62:119-58; U.S. Pat. Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,723,125; 5,783,181; 5,908,626; 5,844,095; and 5,112,946; EP 307,434; EP 367,166; EP 394,827; PCT publications WO 91/06570, WO 96/04388, WO 96/22024, WO 97/34631, and WO 99/04813; Ashkenazi *et al.*, 1991, Proc. Natl. Acad. Sci. USA, 88: 10535-39; Traunecker *et al.*, 1988, Nature, 331:84-86; Zheng *et al.*, 1995, J. Immunol. 154:5590-600; and Vil *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:11337-41).

**[0399]** Fusion proteins may be generated, for example, through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as “DNA shuffling”). DNA shuffling may be employed to alter the activities of anti-PD-1 antibodies as provided herein, including, for example, antibodies with higher affinities and lower dissociation rates (see, e.g., U.S. Pat. Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458; Patten *et al.*, 1997, Curr. Opinion Biotechnol. 8:724-33; Harayama, 1998, Trends Biotechnol. 16(2):76-82; Hansson *et al.*, 1999, J. Mol. Biol. 287:265-76; and Lorenzo and Blasco, 1998, Biotechniques 24(2):308-13). Antibodies, or the encoded antibodies, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion, or other methods prior to recombination. A polynucleotide encoding an antibody

provided herein may be recombined with one or more components, motifs, sections, parts, domains, fragments, *etc.* of one or more heterologous molecules.

**[0400]** An antibody provided herein can also be conjugated to a second antibody to form an antibody heteroconjugate as described, for example, in U.S. Pat. No. 4,676,980.

**[0401]** Antibodies that bind to PD-1 as provided herein may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride, or polypropylene.

**[0402]** The linker may be a “cleavable linker” facilitating release of the conjugated agent in the cell, but non-cleavable linkers are also contemplated herein. Linkers for use in the conjugates of the present disclosure include, without limitation, acid labile linkers (*e.g.*, hydrazone linkers), disulfide-containing linkers, peptidase-sensitive linkers (*e.g.*, peptide linkers comprising amino acids, for example, valine and/or citrulline such as citrulline-valine or phenylalanine-lysine), photolabile linkers, dimethyl linkers (*see, e.g.*, Chari *et al.*, 1992, Cancer Res. 52:127-31; and U.S. Pat. No. 5,208,020), thioether linkers, or hydrophilic linkers designed to evade multidrug transporter-mediated resistance (*see, e.g.*, Kovtun *et al.*, 2010, Cancer Res. 70:2528-37).

**[0403]** Conjugates of the antibody and agent may be made using a variety of bifunctional protein coupling agents such as BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate). The present disclosure further contemplates that conjugates of antibodies and agents may be prepared using any suitable methods as disclosed in the art (*see, e.g.*, Bioconjugate Techniques (Hermanson ed., 2d ed. 2008)).

**[0404]** Conventional conjugation strategies for antibodies and agents have been based on random conjugation chemistries involving the  $\epsilon$ -amino group of Lys residues or the thiol group of Cys residues, which results in heterogenous conjugates. Recently developed techniques allow site-specific conjugation to antibodies, resulting in homogeneous loading and avoiding conjugate subpopulations with altered antigen-binding or pharmacokinetics. These include engineering of “thiomabs” comprising cysteine substitutions at positions on the heavy and light chains that provide reactive thiol groups and do not disrupt immunoglobulin folding and assembly or alter

antigen binding (see, e.g., Junutula *et al.*, 2008, *J. Immunol. Meth.* 332: 41-52; and Junutula *et al.*, 2008, *Nature Biotechnol.* 26:925-32). In another method, selenocysteine is cotranslationally inserted into an antibody sequence by recoding the stop codon UGA from termination to selenocysteine insertion, allowing site specific covalent conjugation at the nucleophilic selenol group of selenocysteine in the presence of the other natural amino acids (see, e.g., Hofer *et al.*, 2008, *Proc. Natl. Acad. Sci. USA* 105:12451-56; and Hofer *et al.*, 2009, *Biochemistry* 48(50):12047-57).

#### 4.4 Methods of Using the Antibodies and Compositions

**[0405]** Provided herein are methods of (a) attenuating T cell activity, and/or (b) downregulating PD-1 expression in a subject. In certain embodiments, methods provided herein downregulate PD-1 expression in a cell of a subject. In certain embodiments, methods provided herein attenuate T cell activity in a subject. A non-limiting example of T cell activity is secretion of a cytokine. In certain embodiments, provided herein are methods of inhibiting secretion of a cytokine. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In some embodiments, the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IFN- $\gamma$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . In certain embodiments, the cytokine is IL-1. In other embodiments, the cytokine is IL-6. In yet other embodiments, the cytokine is IL-12. In still other embodiments, the cytokine is IL-22. In certain embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is TNF- $\alpha$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0406]** In one embodiment, provided herein is a method of inhibiting secretion of IL-2. In some embodiments, provided herein is a methods of inhibiting secretion of IL-17. In yet another embodiment, the method of attenuating T cell activity is a method of inhibiting secretion of IFN- $\gamma$ .

**[0407]** In one aspect, provided herein is a method of attenuating activity of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen binding

fragment thereof provided herein. In other embodiments, the maximal percent attenuation of T cell activity is at least about 10%. In another embodiment, the maximal percent attenuation of T cell activity is at least about 20%. In some embodiments, the maximal percent attenuation of T cell activity is at least about 30%. In one embodiment, the maximal percent attenuation of T cell activity is at least about 40%. In another embodiment, the maximal percent attenuation of T cell activity is at least about 45%. In other embodiments, the maximal percent attenuation of T cell activity is at least about 50%. In one embodiment, the maximal percent attenuation of T cell activity is at least about 55%. In another embodiment, the maximal percent attenuation of T cell activity is at least about 60%. In some embodiments, the maximal percent attenuation of T cell activity is at least about 65%. In other embodiments, the maximal percent attenuation of T cell activity is at least about 70%. In another embodiment, the maximal percent attenuation of T cell activity is at least about 75%. In one embodiment, the maximal percent attenuation of T cell activity is at least about 80%. In other embodiments, the maximal percent attenuation of T cell activity is at least about 85%. In another embodiment, the maximal percent attenuation of T cell activity is at least about 90%. In one embodiment, the maximal percent attenuation of T cell activity is at least about 95%. In some embodiments, the maximal percent attenuation of T cell activity is at least about 100%.

**[0408]** In some embodiments, the attenuation of T cell activity is measured by modulation of production of a cytokine. In some embodiments, the attenuation of T cell activity is measured by modulation of secretion of a cytokine. In some embodiments, the attenuation of T cell activity is measured by modulation of expression of a cytokine. In some embodiments, the attenuation of T cell activity is measured by inhibition of production of a cytokine. In some embodiments, the attenuation of T cell activity is measured by inhibition of secretion of a cytokine. In some embodiments, the attenuation of T cell activity is measured by inhibition of expression of a cytokine. In certain embodiments, cytokine production (e.g., cytokine protein production) is modulated. In certain embodiments, cytokine secretion (e.g., cytokine protein secretion) is modulated. In other embodiments, cytokine expression (e.g., cytokine gene expression) is modulated. In some embodiments, the modulation is a decrease, inhibition, or down-regulation of the cytokine. In other embodiments, the modulation is an increase or up-regulation of the cytokine. In certain embodiments, cytokine production (e.g., cytokine protein production) is inhibited. In certain embodiments, cytokine secretion (e.g., cytokine protein secretion) is

inhibited. In other embodiments, cytokine expression (e.g., cytokine gene expression) is inhibited. In certain embodiments, cytokine production from a cell is modulated. In certain embodiments, cytokine secretion from a cell is modulated. In certain embodiments, cytokine expression from a cell is modulated. In certain embodiments, cytokine production from a cell is inhibited. In certain embodiments, cytokine secretion from a cell is inhibited. In certain embodiments, cytokine expression from a cell is inhibited. In certain embodiments, the cell is a T cell. In some embodiments, the cell is not a T cell.

**[0409]** In some embodiments, the cytokine is selected from the group consisting of IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In one embodiment, the cytokine is IL-2. In another embodiment, the cytokine is IL-17. In other embodiments, the cytokine is IFN- $\gamma$ . In some embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In certain embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In other embodiments, the cytokine is IL-6. In yet other embodiments, the cytokine is IL-12. In still other embodiments, the cytokine is IL-22. In certain embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is TNF- $\alpha$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0410]** In some embodiments, the inhibition of cytokine production is concurrent with downregulation of PD-1 expression on the surface of the T cell. In some embodiments, the inhibition of cytokine production is preceded by downregulation of PD-1 expression on the surface of the T cell. In some embodiments, the inhibition of cytokine production precedes downregulation of PD-1 expression on the surface of the T cell. In one embodiment, the downregulation of PD-1 expression on the surface of the T cell occurs as early as 4 hours after contact with the antibody or antigen-binding fragment thereof.

**[0411]** In one aspect, provided herein is a method of modulating PD-1 activity and/or expression in a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof as described herein. In one embodiment, PD-1 activity is modulated. In another embodiment,

PD-1 expression is modulated. In other embodiment, PD-1 activity and PD-1 expression are both modulated. In one embodiment, PD-1 signaling is activated. In another embodiment, PD-1 expression is inhibited. In certain embodiments, the antibody is a PD-1 agonist. In certain embodiments, the antibody provided herein specifically binds to human PD-1, and activates (e.g., partially activates) or otherwise modulates at least one PD-1 activity. In a specific embodiment, the at least one PD-1 activity is inhibition of cytokine production. In certain embodiments, the antibody that specifically binds to PD-1 binds to an ECD of human PD-1, or an epitope of an ECD of human PD-1 thereof. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0412]** In another aspect, provided herein is a method of downregulating PD-1 expression in a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is a PD-1 agonist provided herein. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0413]** In another aspect, provided herein is a method of attenuating T cell activity and down-regulating PD-1 expression in a cell, comprising contacting the cell with an antibody that

specifically binds to PD-1 as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof as described herein. In certain embodiments, the attenuation of T cell activity is inhibition of cytokine production. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0414]** PD-1 activity can relate to any activity of PD-1 such as those known or described in the art. PD-1 activity and PD-1 signaling are used interchangeably herein. In certain aspects, PD-1 activity is induced by PD-1 ligand (*e.g.*, PD-L1) binding to PD-1. Expression levels of PD-1 can be assessed by methods described herein or known to one of skill in the art (*e.g.*, Western blotting, ELISA, immunohistochemistry, or flow cytometry).

**[0415]** Also provided herein is a method of inhibiting cytokine production in a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen binding fragment thereof as described herein. In some embodiments, the antibody binds to an ECD of human PD-1. In some embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific

embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody.

**[0416]** Also provided herein are methods of activating PD-1 signaling in a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof as described herein. In some embodiments, the antibody binds to an ECD of human PD-1. In some embodiments, the antibody binds to an epitope of an ECD of human PD-1. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody. In one embodiment, PD-1 signaling is partially activated.

**[0417]** In one aspect, provided herein are methods of attenuating T cell activity, comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof as described herein. In some embodiments, the antibody binds to an ECD of human PD-1. In some embodiments, the antibody binds to an epitope of an ECD of human PD-1. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L1 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from the PD-L2 binding site. In certain embodiments, the antibody specifically binds to an epitope of an ECD of human PD-1 that is distinct from both the PD-L1 and PD-L2 binding sites. In certain embodiments, binding of PD-L1 to PD-1 is not inhibited by the antibody. In other embodiments, binding of PD-L2 to PD-1 is not inhibited by the antibody. In specific embodiments, neither binding of PD-L1 to PD-1 nor binding of PD-L2 to PD-1 is inhibited by the antibody. In some

embodiments, T cell activity is attenuated by at least about 10%. In some embodiments, T cell activity is attenuated by at least about 15%. In some embodiments, T cell activity is attenuated by at least about 20%. In some embodiments, T cell activity is attenuated by at least about 25%. In some embodiments, T cell activity is attenuated by at least about 30%. In some embodiments, T cell activity is attenuated by at least about 35%. In some embodiments, T cell activity is attenuated by at least about 40%. In some embodiments, T cell activity is attenuated by at least about 45%. In some embodiments, T cell activity is attenuated by at least about 50%. In some embodiments, T cell activity is attenuated by at least about 55%. In some embodiments, T cell activity is attenuated by at least about 60%. In some embodiments, T cell activity is attenuated by at least about 65%. In some embodiments, T cell activity is attenuated by at least about 70%. In some embodiments, T cell activity is attenuated by at least about 75%. In some embodiments, T cell activity is attenuated by at least about 80%. In some embodiments, T cell activity is attenuated by at least about 85%. In some embodiments, T cell activity is attenuated by at least about 90%. In some embodiments, T cell activity is attenuated by at least about 95%. In some embodiments, T cell activity is attenuated by at least about 98%. In some embodiments, T cell activity is attenuated by at least about 99%. In some embodiments, T cell activity is attenuated by at least about 100%. In certain embodiments, T cell activity is attenuated by at least about 25% to about 65%. In specific embodiments, the T cell activity attenuation is assessed by methods described herein. In some embodiments, the T cell activity attenuation is assessed by methods known to one of skill in the art. In certain embodiments, the T cell activity attenuation is relative to T cell activity in a cell that is not contacted with an anti-PD-1 antibody. In certain embodiments, the T cell activity attenuation is relative to T cell activity that is contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0418]** A non-limiting example of T cell activity is secretion of a cytokine. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In some embodiments, the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IFN- $\gamma$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . In certain embodiments, the cytokine is IL-1. In other

embodiments, the cytokine is IL-6. In yet other embodiments, the cytokine is IL-12. In still other embodiments, the cytokine is IL-22. In certain embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is TNF- $\alpha$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0419]** In certain embodiments, cytokine secretion is inhibited as a result of inhibition of cytokine production. In other embodiments, cytokine secretion is inhibited as a result of inhibition of cytokine expression.

**[0420]** In specific embodiments, provided herein are methods of inhibiting cytokine secretion from a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6. In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In some embodiments, the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In certain embodiments, the cytokine is IL-2. In other embodiments, the cytokine is IL-17. In yet other embodiments, the cytokine is IFN- $\gamma$ . In certain embodiments, the cytokine is IL-2 and IL-17. In some embodiments, the cytokine is IL-2 and IFN- $\gamma$ . In yet other embodiments, the cytokine is IL-17 and IFN- $\gamma$ . In still other embodiments, the cytokine is IL-2, IL-17, and IFN- $\gamma$ . In certain embodiments, the cytokine is IL-1. In other embodiments, the cytokine is IL-6. In yet other embodiments, the cytokine is IL-12. In still other embodiments, the cytokine is IL-22. In certain embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is TNF- $\alpha$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0421]** In some embodiments, provided herein are methods of inhibiting IL-2 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific

embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0422]** In one embodiment, IL-2 secretion is inhibited by at least about 5%. In some embodiments, IL-2 secretion is inhibited by at least about 10%. In another embodiment, IL-2 secretion is inhibited by at least about 15%. In other embodiments, IL-2 secretion is inhibited by at least about 20%. In one embodiment, IL-2 secretion is inhibited by at least about 25%. In another embodiment, IL-2 secretion is inhibited by at least about 30%. In some embodiments, IL-2 secretion is inhibited by at least about 35%. In one embodiment, IL-2 secretion is inhibited by at least about 40%. In another embodiment, IL-2 secretion is inhibited by at least about 45%. In other embodiments, IL-2 secretion is inhibited by at least about 50%. In some embodiments, IL-2 secretion is inhibited by at least about 55%. In another embodiment, IL-2 secretion is inhibited by at least about 60%. In one embodiment, IL-2 secretion is inhibited by at least about 65%. In one embodiment, IL-2 secretion is inhibited by at least about 70%. In another embodiment, IL-2 secretion is inhibited by at least about 75%. In some embodiments, IL-2 secretion is inhibited by at least about 80%. In other embodiments, IL-2 secretion is inhibited by at least about 85%. In another embodiment, IL-2 secretion is inhibited by at least about 90%. In one embodiment, IL-2 secretion is inhibited by at least about 95%. In some embodiments, IL-2 secretion is inhibited by at least about 98%. In another embodiment, IL-2 secretion is inhibited by at least about 99%. In specific embodiments, IL-2 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-2 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-2 secretion is assessed by methods known to one of skill in the art (*e.g.*, MSD multiplex assay). In a specific embodiment, the IL-2 secretion is inhibited relative to IL-2 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-2 secretion is inhibited relative to IL-2 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0423]** In certain embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-2 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (*e.g.*, MSD multiplex assay). In a specific embodiment, the IL-2 secretion is inhibited relative to IL-2 secretion from a cell that is

not contacted with an anti-PD-1 antibody. In other embodiments, the IL-2 secretion is inhibited relative to IL-2 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0424]** In some embodiments, provided herein are methods of inhibiting IL-17 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0425]** In one embodiment, IL-17 secretion is inhibited by at least about 5%. In some embodiments, IL-17 secretion is inhibited by at least about 10%. In another embodiment, IL-17 secretion is inhibited by at least about 15%. In other embodiments, IL-17 secretion is inhibited by at least about 20%. In one embodiment, IL-17 secretion is inhibited by at least about 25%. In another embodiment, IL-17 secretion is inhibited by at least about 30%. In some embodiments, IL-17 secretion is inhibited by at least about 35%. In one embodiment, IL-17 secretion is inhibited by at least about 40%. In another embodiment, IL-17 secretion is inhibited by at least about 45%. In other embodiments, IL-17 secretion is inhibited by at least about 50%. In some embodiments, IL-17 secretion is inhibited by at least about 55%. In another embodiment, IL-17 secretion is inhibited by at least about 60%. In one embodiment, IL-17 secretion is inhibited by at least about 65%. In one embodiment, IL-17 secretion is inhibited by at least about 70%. In another embodiment, IL-17 secretion is inhibited by at least about 75%. In some embodiments, IL-17 secretion is inhibited by at least about 80%. In other embodiments, IL-17 secretion is inhibited by at least about 85%. In another embodiment, IL-17 secretion is inhibited by at least about 90%. In one embodiment, IL-17 secretion is inhibited by at least about 95%. In some embodiments, IL-17 secretion is inhibited by at least about 98%. In another embodiment, IL-17 secretion is inhibited by at least about 99%. In specific embodiments, IL-17 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-17 secretion is assessed by methods described herein. In other embodiments, the

inhibition of IL-17 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-17 secretion is inhibited relative to IL-17 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-17 secretion is inhibited relative to IL-17 secretion in a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0426]** In certain embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM.

embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-17 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-17 secretion is inhibited relative to IL-17 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-17 secretion is inhibited relative to IL-17 secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0427]** In some embodiments, provided herein are methods of inhibiting IFN- $\gamma$  secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0428]** In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 5%. In some embodiments, IFN- $\gamma$  secretion is inhibited by at least about 10%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 15%. In other embodiments, IFN- $\gamma$  secretion is inhibited by at least about 20%. In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 25%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 30%. In some embodiments, IFN- $\gamma$  secretion is inhibited by at least about 35%. In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 40%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 45%. In other embodiments, IFN- $\gamma$  secretion is inhibited by at least about 50%. In some embodiments, IFN- $\gamma$  secretion is inhibited by at least about 55%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 60%. In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 65%. In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 70%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 75%. In some embodiments, IFN- $\gamma$  secretion is inhibited by at least about 80%. In other embodiments, IFN- $\gamma$  secretion is inhibited by at least about 85%. In another embodiment, IFN- $\gamma$  secretion is

inhibited by at least about 90%. In one embodiment, IFN- $\gamma$  secretion is inhibited by at least about 95%. In some embodiments, IFN- $\gamma$  secretion is inhibited by at least about 98%. In another embodiment, IFN- $\gamma$  secretion is inhibited by at least about 99%. In specific embodiments, IFN- $\gamma$  secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IFN- $\gamma$  secretion is assessed by methods described herein. In other embodiments, the inhibition of IFN- $\gamma$  secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0429]** In certain embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another

embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IFN- $\gamma$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IFN- $\gamma$  secretion is inhibited relative to IFN- $\gamma$  secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0430]** In some embodiments, provided herein are methods of inhibiting IL-1 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0431]** In one embodiment, IL-1 secretion is inhibited by at least about 5%. In some embodiments, IL-1 secretion is inhibited by at least about 10%. In another embodiment, IL-1 secretion is inhibited by at least about 15%. In other embodiments, IL-1 secretion is inhibited by at least about 20%. In one embodiment, IL-1 secretion is inhibited by at least about 25%. In another embodiment, IL-1 secretion is inhibited by at least about 30%. In some embodiments, IL-1 secretion is inhibited by at least about 35%. In one embodiment, IL-1 secretion is inhibited by at least about 40%. In another embodiment, IL-1 secretion is inhibited by at least about 45%. In other embodiments, IL-1 secretion is inhibited by at least about 50%. In some embodiments,

IL-1 secretion is inhibited by at least about 55%. In another embodiment, IL-1 secretion is inhibited by at least about 60%. In one embodiment, IL-1 secretion is inhibited by at least about 65%. In one embodiment, IL-1 secretion is inhibited by at least about 70%. In another embodiment, IL-1 secretion is inhibited by at least about 75%. In some embodiments, IL-1 secretion is inhibited by at least about 80%. In other embodiments, IL-1 secretion is inhibited by at least about 85%. In another embodiment, IL-1 secretion is inhibited by at least about 90%. In one embodiment, IL-1 secretion is inhibited by at least about 95%. In some embodiments, IL-1 secretion is inhibited by at least about 98%. In another embodiment, IL-1 secretion is inhibited by at least about 99%. In specific embodiments, IL-1 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-1 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-1 secretion is assessed by methods known to one of skill in the art (*e.g.*, MSD multiplex assay). In a specific embodiment, the IL-1 secretion is inhibited relative to IL-1 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-1 secretion is inhibited relative to IL-1 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0432]** In certain embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other

embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-1 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-1 secretion is inhibited relative to IL-1 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-1 secretion is inhibited relative to IL-1 secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0433]** In some embodiments, provided herein are methods of inhibiting IL-6 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0434]** In one embodiment, IL-6 secretion is inhibited by at least about 5%. In some embodiments, IL-6 secretion is inhibited by at least about 10%. In another embodiment, IL-6 secretion is inhibited by at least about 15%. In other embodiments, IL-6 secretion is inhibited by

at least about 20%. In one embodiment, IL-6 secretion is inhibited by at least about 25%. In another embodiment, IL-6 secretion is inhibited by at least about 30%. In some embodiments, IL-6 secretion is inhibited by at least about 35%. In one embodiment, IL-6 secretion is inhibited by at least about 40%. In another embodiment, IL-6 secretion is inhibited by at least about 45%. In other embodiments, IL-6 secretion is inhibited by at least about 50%. In some embodiments, IL-6 secretion is inhibited by at least about 55%. In another embodiment, IL-6 secretion is inhibited by at least about 60%. In one embodiment, IL-6 secretion is inhibited by at least about 65%. In one embodiment, IL-6 secretion is inhibited by at least about 70%. In another embodiment, IL-6 secretion is inhibited by at least about 75%. In some embodiments, IL-6 secretion is inhibited by at least about 80%. In other embodiments, IL-6 secretion is inhibited by at least about 85%. In another embodiment, IL-6 secretion is inhibited by at least about 90%. In one embodiment, IL-6 secretion is inhibited by at least about 95%. In some embodiments, IL-6 secretion is inhibited by at least about 98%. In another embodiment, IL-6 secretion is inhibited by at least about 99%. In specific embodiments, IL-6 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-6 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-6 secretion is assessed by methods known to one of skill in the art (*e.g.*, MesoScale<sup>TM</sup> Discovery (MSD) multiplex assay). In a specific embodiment, the IL-6 secretion is inhibited relative to IL-6 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-6 secretion is inhibited relative to IL-6 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0435]** In certain embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one

embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-6 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-6 secretion is inhibited relative to IL-6 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-6 secretion is inhibited relative to IL-6 secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0436]** In some embodiments, provided herein are methods of inhibiting IL-12 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising

CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0437]** In one embodiment, IL-12 secretion is inhibited by at least about 5%. In some embodiments, IL-12 secretion is inhibited by at least about 10%. In another embodiment, IL-12 secretion is inhibited by at least about 15%. In other embodiments, IL-12 secretion is inhibited by at least about 20%. In one embodiment, IL-12 secretion is inhibited by at least about 25%. In another embodiment, IL-12 secretion is inhibited by at least about 30%. In some embodiments, IL-12 secretion is inhibited by at least about 35%. In one embodiment, IL-12 secretion is inhibited by at least about 40%. In another embodiment, IL-12 secretion is inhibited by at least about 45%. In other embodiments, IL-12 secretion is inhibited by at least about 50%. In some embodiments, IL-12 secretion is inhibited by at least about 55%. In another embodiment, IL-12 secretion is inhibited by at least about 60%. In one embodiment, IL-12 secretion is inhibited by at least about 65%. In one embodiment, IL-12 secretion is inhibited by at least about 70%. In another embodiment, IL-12 secretion is inhibited by at least about 75%. In some embodiments, IL-12 secretion is inhibited by at least about 80%. In other embodiments, IL-12 secretion is inhibited by at least about 85%. In another embodiment, IL-12 secretion is inhibited by at least about 90%. In one embodiment, IL-12 secretion is inhibited by at least about 95%. In some embodiments, IL-12 secretion is inhibited by at least about 98%. In another embodiment, IL-12 secretion is inhibited by at least about 99%. In specific embodiments, IL-12 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-12 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-12 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-12 secretion is inhibited relative to IL-12 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-12 secretion is inhibited relative to IL-12 secretion in a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0438]** In certain embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another

embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-12 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-12 secretion is inhibited relative to IL-12 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-12 secretion is inhibited relative to IL-12 secretion in a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0439]** In some embodiments, provided herein are methods of inhibiting IL-22 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g.,

an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0440]** In one embodiment, IL-22 secretion is inhibited by at least about 5%. In some embodiments, IL-22 secretion is inhibited by at least about 10%. In another embodiment, IL-22 secretion is inhibited by at least about 15%. In other embodiments, IL-22 secretion is inhibited by at least about 20%. In one embodiment, IL-22 secretion is inhibited by at least about 25%. In another embodiment, IL-22 secretion is inhibited by at least about 30%. In some embodiments, IL-22 secretion is inhibited by at least about 35%. In one embodiment, IL-22 secretion is inhibited by at least about 40%. In another embodiment, IL-22 secretion is inhibited by at least about 45%. In other embodiments, IL-22 secretion is inhibited by at least about 50%. In some embodiments, IL-22 secretion is inhibited by at least about 55%. In another embodiment, IL-22 secretion is inhibited by at least about 60%. In one embodiment, IL-22 secretion is inhibited by at least about 65%. In one embodiment, IL-22 secretion is inhibited by at least about 70%. In another embodiment, IL-22 secretion is inhibited by at least about 75%. In some embodiments, IL-22 secretion is inhibited by at least about 80%. In other embodiments, IL-22 secretion is inhibited by at least about 85%. In another embodiment, IL-22 secretion is inhibited by at least about 90%. In one embodiment, IL-22 secretion is inhibited by at least about 95%. In some embodiments, IL-22 secretion is inhibited by at least about 98%. In another embodiment, IL-22 secretion is inhibited by at least about 99%. In specific embodiments, IL-22 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-22 secretion is assessed by methods described herein. In other embodiments, the inhibition of IL-22 secretion is assessed by methods known to one of skill in the art (*e.g.*, MSD multiplex assay). In a specific embodiment, the IL-22 secretion is inhibited relative to IL-22 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-22 secretion is inhibited relative to IL-22 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0441]** In certain embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-22 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (*e.g.*, MSD multiplex assay). In a specific embodiment, the IL-22 secretion is inhibited relative to IL-22 secretion from a cell that

is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-22 secretion is inhibited relative to IL-22 secretion from a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0442]** In some embodiments, provided herein are methods of inhibiting IL-23 secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (*e.g.*, an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0443]** In one embodiment, IL-23 secretion is inhibited by at least about 5%. In some embodiments, IL-23 secretion is inhibited by at least about 10%. In another embodiment, IL-23 secretion is inhibited by at least about 15%. In other embodiments, IL-23 secretion is inhibited by at least about 20%. In one embodiment, IL-23 secretion is inhibited by at least about 25%. In another embodiment, IL-23 secretion is inhibited by at least about 30%. In some embodiments, IL-23 secretion is inhibited by at least about 35%. In one embodiment, IL-23 secretion is inhibited by at least about 40%. In another embodiment, IL-23 secretion is inhibited by at least about 45%. In other embodiments, IL-23 secretion is inhibited by at least about 50%. In some embodiments, IL-23 secretion is inhibited by at least about 55%. In another embodiment, IL-23 secretion is inhibited by at least about 60%. In one embodiment, IL-23 secretion is inhibited by at least about 65%. In one embodiment, IL-23 secretion is inhibited by at least about 70%. In another embodiment, IL-23 secretion is inhibited by at least about 75%. In some embodiments, IL-23 secretion is inhibited by at least about 80%. In other embodiments, IL-23 secretion is inhibited by at least about 85%. In another embodiment, IL-23 secretion is inhibited by at least about 90%. In one embodiment, IL-23 secretion is inhibited by at least about 95%. In some embodiments, IL-23 secretion is inhibited by at least about 98%. In another embodiment, IL-23 secretion is inhibited by at least about 99%. In specific embodiments, IL-23 secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of IL-23 secretion is assessed by methods described herein. In other embodiments, the

inhibition of IL-23 secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-23 secretion is inhibited relative to IL-23 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-23 secretion is inhibited relative to IL-23 secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0444]** In certain embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.0005 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.0001 nM. In another embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.00005 nM.

embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, IL-23 secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the IL-23 secretion is inhibited relative to IL-23 secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-23 secretion is inhibited relative to IL-23 secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0445]** In some embodiments, provided herein are methods of inhibiting GM-CSF secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0446]** In one embodiment, GM-CSF secretion is inhibited by at least about 5%. In some embodiments, GM-CSF secretion is inhibited by at least about 10%. In another embodiment, GM-CSF secretion is inhibited by at least about 15%. In other embodiments, GM-CSF secretion is inhibited by at least about 20%. In one embodiment, GM-CSF secretion is inhibited by at least about 25%. In another embodiment, GM-CSF secretion is inhibited by at least about 30%. In some embodiments, GM-CSF secretion is inhibited by at least about 35%. In one embodiment, GM-CSF secretion is inhibited by at least about 40%. In another embodiment, GM-CSF secretion is inhibited by at least about 45%. In other embodiments, GM-CSF secretion is inhibited by at least about 50%. In some embodiments, GM-CSF secretion is inhibited by at least about 55%. In another embodiment, GM-CSF secretion is inhibited by at least about 60%. In one embodiment, GM-CSF secretion is inhibited by at least about 65%. In one embodiment, GM-CSF secretion is inhibited by at least about 70%. In another embodiment, GM-CSF secretion is inhibited by at least about 75%. In some embodiments, GM-CSF secretion is inhibited by at least about 80%. In other embodiments, GM-CSF secretion is inhibited by at

least about 85%. In another embodiment, GM-CSF secretion is inhibited by at least about 90%. In one embodiment, GM-CSF secretion is inhibited by at least about 95%. In some embodiments, GM-CSF secretion is inhibited by at least about 98%. In another embodiment, GM-CSF secretion is inhibited by at least about 99%. In specific embodiments, GM-CSF secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of GM-CSF secretion is assessed by methods described herein. In other embodiments, the inhibition of GM-CSF secretion is assessed by methods known to one of skill in the art (e.g., MesoScale<sup>TM</sup> Discovery (MSD) multiplex assay). In a specific embodiment, the GM-CSF secretion is inhibited relative to GM-CSF secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-2 secretion is inhibited relative to GM-CSF secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0447]** In certain embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one

embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, GM-CSF secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the GM-CSF secretion is inhibited relative to GM-CSF secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the IL-2 secretion is inhibited relative to GM-CSF secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0448]** In some embodiments, provided herein are methods of inhibiting TNF- $\alpha$  secretion from a cell comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen-binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0449]** In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 5%. In some embodiments, TNF- $\alpha$  secretion is inhibited by at least about 10%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 15%. In other embodiments, TNF- $\alpha$  secretion is inhibited by at least about 20%. In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 25%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 30%. In some embodiments, TNF- $\alpha$  secretion is inhibited by at least about 35%. In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 40%. In another embodiment, TNF- $\alpha$  secretion is

inhibited by at least about 45%. In other embodiments, TNF- $\alpha$  secretion is inhibited by at least about 50%. In some embodiments, TNF- $\alpha$  secretion is inhibited by at least about 55%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 60%. In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 65%. In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 70%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 75%. In some embodiments, TNF- $\alpha$  secretion is inhibited by at least about 80%. In other embodiments, TNF- $\alpha$  secretion is inhibited by at least about 85%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 90%. In one embodiment, TNF- $\alpha$  secretion is inhibited by at least about 95%. In some embodiments, TNF- $\alpha$  secretion is inhibited by at least about 98%. In another embodiment, TNF- $\alpha$  secretion is inhibited by at least about 99%. In specific embodiments, TNF- $\alpha$  secretion is inhibited by at least about 25% or 35%, optionally to about 75%. In some embodiments, the inhibition of TNF- $\alpha$  secretion is assessed by methods described herein. In other embodiments, the inhibition of TNF- $\alpha$  secretion is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0450]** In certain embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 50 nM. In other embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 40 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 30 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 20 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 10 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 5 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 1 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.75 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.5 nM. In other embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.1 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.05 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.01 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.005 nM. In one

embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at most about 0.001 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 50 nM. In other embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 40 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 30 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 20 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 10 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 5 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 1 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.75 nM. In other embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.5 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.1 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.05 nM. In some embodiments, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.01 nM. In another embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.005 nM. In one embodiment, TNF- $\alpha$  secretion is inhibited with an EC<sub>50</sub> of at least about 0.001 nM. In some embodiments, the EC<sub>50</sub> is assessed by methods described herein. In other embodiments, the EC<sub>50</sub> is assessed by methods known to one of skill in the art (e.g., MSD multiplex assay). In a specific embodiment, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion from a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the TNF- $\alpha$  secretion is inhibited relative to TNF- $\alpha$  secretion from a cell contacted with an unrelated antibody (e.g., an antibody that does not specifically bind to PD-1).

**[0451]** In some embodiments, provided herein are methods of downregulating PD-1 expression in a cell, comprising contacting the cell with an antibody that specifically binds to PD-1 (e.g., an ECD of human PD-1 or an epitope of an ECD of human PD-1) as provided herein. In a specific embodiment, the cell is a T cell. In certain embodiments, the cell is contacted with an effective amount of an antibody or antigen binding fragment thereof described herein. In certain embodiments, the antibody is any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 or an antigen-binding fragment thereof, or an antibody comprising CDRs of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6.

**[0452]** In one embodiment, PD-1 expression is downregulated by at least about 5%. In one embodiment, PD-1 expression is downregulated by at least about 10%. In another embodiment, PD-1 expression is downregulated by at least about 15%. In some embodiments, PD-1 expression is downregulated by at least about 20%. In other embodiments, PD-1 expression is downregulated by at least about 25%. In another embodiment, PD-1 expression is downregulated by at least about 30%. In one embodiment, PD-1 expression is downregulated by at least about 35%. In some embodiments, PD-1 expression is downregulated by at least about 40%. In another embodiment, PD-1 expression is downregulated by at least about 45%. In one embodiment, PD-1 expression is downregulated by at least about 50%. In other embodiments, PD-1 expression is downregulated by at least about 55%. In another embodiment, PD-1 expression is downregulated by at least about 60%. In some embodiments, PD-1 expression is downregulated by at least about 65%. In one embodiment, PD-1 expression is downregulated by at least about 70%. In another embodiment, PD-1 expression is downregulated by at least about 75%. In one embodiment, PD-1 expression is downregulated by at least about 80%. In some embodiments, PD-1 expression is downregulated by at least about 85%. In another embodiment, PD-1 expression is downregulated by at least about 90%. In other embodiments, PD-1 expression is downregulated by at least about 95%. In one embodiment, PD-1 expression is downregulated by at least about 98%. In another embodiment, PD-1 expression is downregulated by at least about 99%. In specific embodiments, antibodies provided herein specifically bind to PD-1 and downregulates PD-1 expression by at least about 25% or 35%, optionally to about 75%. In some embodiments, the downregulation of PD-1 expression is assessed by methods described herein. In other embodiments, the downregulation of PD-1 expression is assessed by methods known to one of skill in the art (*e.g.*, flow cytometry, Western blotting, Northern blotting, or RT-PCR). In a specific embodiment, the downregulation of PD-1 expression is assessed by flow cytometry. In another embodiment, the downregulation of PD-1 expression is assessed by Western blotting. In yet another embodiment, the downregulation of PD-1 expression is assessed by Northern blotting. In still another embodiment, the downregulation of PD-1 expression is assessed by RT-PCR. In a specific embodiment, the PD-1 expression is downregulated relative to PD-1 expression in a cell that is not contacted with an anti-PD-1 antibody. In other embodiments, the PD-1 expression is downregulated relative to

PD-1 expression in a cell contacted with an unrelated antibody (*e.g.*, an antibody that does not specifically bind to PD-1).

**[0453]** In one embodiment, provided herein is a method of downregulating PD-1 expression on the surface of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen binding fragment thereof provided herein. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 10%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 20%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 30%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 40%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 45%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 50%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 55%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 60%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 65%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 70%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 75%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 80%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 85%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 90%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 95%. In one embodiment, the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 100%.

**[0454]** In certain embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours after the contact with the antibody or antigen-binding fragment thereof. In other embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, 14 hours, 16 hours, 18 hours, 20 hours, or 22 hours after the contact with the antibody or antigen-binding fragment thereof. In one embodiment, the downregulation occurs as early as 4 hours after the contact. In one embodiment, the downregulation occurs as early as 6 hours after the contact. In another embodiment, the downregulation occurs as early as 8 hours after the contact. In other embodiments, the downregulation occurs as early as 10 hours after the contact. In some embodiments, the downregulation occurs as early as 12 hours after the contact. In one embodiment, the downregulation occurs as early as 14 hours after the contact. In another embodiment, the downregulation occurs as early as 16 hours after the contact. In some embodiments, the downregulation occurs as early as 18 hours after the contact. In other embodiments, the downregulation occurs as early as 20 hours after the contact. In other embodiments, the downregulation occurs as early as 22 hours after the contact. In yet other embodiments, the downregulation of PD-1 expression on the surface of T cells occurs as early as 24 hours after the contact with the antibody or antigen-binding fragment thereof.

**[0455]** In one embodiment, the downregulation of PD-1 expression on the surface of the T cell precedes cytokine inhibition. In another embodiment, the downregulation of PD-1 expression on the surface of the T cell is concurrent with cytokine inhibition. In yet another embodiment, the downregulation of PD-1 expression on the surface of the T cell is preceded by cytokine inhibition. In certain embodiments, the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof. In one embodiment, the cytokine is IL-2. In another embodiment, the cytokine is IL-17. In other embodiments, the cytokine is IFN- $\gamma$ . In certain embodiments, the cytokine is selected from the group consisting of IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, IFN- $\gamma$ , and TNF- $\alpha$ . In certain embodiments, the cytokine is IL-1. In other embodiments, the cytokine is IL-6. In yet other embodiments, the cytokine is IL-12. In still other embodiments, the cytokine is IL-22. In certain embodiments, the cytokine is IL-23. In some embodiments, the cytokine is GM-CSF. In other embodiments, the cytokine is TNF- $\alpha$ . Other combinations of two, three or more of the above-mentioned cytokines are also contemplated.

**[0456]** In other aspects, anti-PD-1 antibodies and fragments thereof of the present disclosure are useful for detecting the presence of PD-1 in a biological sample. Such anti-PD-1 antibodies may include those that bind to human and/or cynomolgus PD-1 but do not induce PD-1 signaling activity. The term “detecting” as used herein encompasses quantitative or qualitative detection. In certain embodiments, a biological sample comprises bodily fluid, a cell, or a tissue.

#### 4.5 Pharmaceutical Compositions

**[0457]** In one aspect, the present disclosure further provides pharmaceutical compositions comprising at least one anti-PD-1 antibody of the present disclosure. In some embodiments, a pharmaceutical composition comprises 1) an anti-PD-1 antibody, and 2) a pharmaceutically acceptable carrier.

**[0458]** Pharmaceutical compositions comprising an antibody are prepared for storage by mixing the antibody having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*see, e.g.*, Remington, Remington's Pharmaceutical Sciences (18th ed. 1980)) in the form of aqueous solutions or lyophilized or other dried forms.

**[0459]** The antibodies of the present disclosure may be formulated in any suitable form for delivery to a target cell/tissue, *e.g.*, as microcapsules or macroemulsions (Remington, *supra*; Park *et al.*, 2005, Molecules 10:146-61; Malik *et al.*, 2007, Curr. Drug. Deliv. 4:141-51), as sustained release formulations (Putney and Burke, 1998, Nature Biotechnol. 16:153-57), or in liposomes (Maclean *et al.*, 1997, Int. J. Oncol. 11:325-32; Kontermann, 2006, Curr. Opin. Mol. Ther. 8:39-45).

**[0460]** An antibody provided herein can also be entrapped in microcapsule prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsule and poly-(methylmethacrylate) microcapsule, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed, for example, in Remington, *supra*.

**[0461]** Various compositions and delivery systems are known and can be used with an antibody that binds to PD-1 as described herein, including, but not limited to, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the antibody, receptor-mediated endocytosis (*see, e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262:4429-32), construction of a nucleic acid as part of a retroviral or other vector, *etc.* In another embodiment,

a composition can be provided as a controlled release or sustained release system. In one embodiment, a pump may be used to achieve controlled or sustained release (see, e.g., Langer, *supra*; Sefton, 1987, *Crit. Ref. Biomed. Eng.* 14:201-40; Buchwald *et al.*, 1980, *Surgery* 88:507-16; and Saudek *et al.*, 1989, *N. Engl. J. Med.* 321:569-74). In another embodiment, polymeric materials can be used to achieve controlled or sustained release of a prophylactic or therapeutic agent (e.g., an antibody that binds to PD-1 as described herein) or a composition of the invention (see, e.g., Medical Applications of Controlled Release (Langer and Wise eds., 1974); Controlled Drug Bioavailability, Drug Product Design and Performance (Smolen and Ball eds., 1984); Ranger and Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61-126; Levy *et al.*, 1985, *Science* 228:190-92; During *et al.*, 1989, *Ann. Neurol.* 25:351-56; Howard *et al.*, 1989, *J. Neurosurg.* 71:105-12; U.S. Pat. Nos. 5,679,377; 5,916,597; 5,912,015; 5,989,463; and 5,128,326; PCT Publication Nos. WO 99/15154 and WO 99/20253). Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In one embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable.

**[0462]** In yet another embodiment, a controlled or sustained release system can be placed in proximity of a particular target tissue, for example, the nasal passages or lungs, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, Medical Applications of Controlled Release Vol. 2, 115-38 (1984)). Controlled release systems are discussed, for example, by Langer, 1990, *Science* 249:1527-33. Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more antibodies that bind to PD-1 as described herein (see, e.g., U.S. Pat. No. 4,526,938, PCT publication Nos. WO 91/05548 and WO 96/20698, Ning *et al.*, 1996, *Radiotherapy & Oncology* 39:179-89; Song *et al.*, 1995, *PDA J. of Pharma. Sci. & Tech.* 50:372-97; Cleek *et al.*, 1997, *Proc. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-54; and Lam *et al.*, 1997, *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-60).

#### 4.6 Kits

**[0463]** Also provided herein are kits comprising an antibody (e.g., an anti- PD-1 antibody) provided herein, or a composition (e.g., a pharmaceutical composition) thereof, packaged into suitable packaging material. A kit optionally includes a label or packaging insert including a description of the components or instructions for use *in vitro*, *in vivo*, or *ex vivo*, of the components therein.

**[0464]** The term “packaging material” refers to a physical structure housing the components of the kit. The packaging material can maintain the components steriley, and can be made of material commonly used for such purposes (e.g., paper, corrugated fiber, glass, plastic, foil, ampoules, vials, tubes, *etc.*).

**[0465]** Kits provided herein can include labels or inserts. Labels or inserts include “printed matter,” e.g., paper or cardboard, separate or affixed to a component, a kit or packing material (e.g., a box), or attached to, for example, an ampoule, tube, or vial containing a kit component. Labels or inserts can additionally include a computer readable medium, such as a disk (e.g., hard disk, card, memory disk), optical disk such as CD- or DVD-ROM/RAM, DVD, MP3, magnetic tape, or an electrical storage media such as RAM and ROM or hybrids of these such as magnetic/optical storage media, FLASH media, or memory type cards. Labels or inserts can include information identifying manufacturer information, lot numbers, manufacturer location, and date.

**[0466]** Kits provided herein can additionally include other components. Each component of the kit can be enclosed within an individual container, and all of the various containers can be within a single package. Kits can also be designed for cold storage. A kit can further be designed to contain antibodies provided herein, or cells that contain nucleic acids encoding the antibodies provided herein. The cells in the kit can be maintained under appropriate storage conditions until ready to use.

**[0467]** Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, suitable methods and materials are described herein.

**[0468]** All applications, publications, patents and other references, GenBank citations and ATCC citations cited herein are incorporated by reference in their entirety. In case of conflict, the specification, including definitions, will control.

**[0469]** As used herein, the singular forms “a,” “and,” and “the” include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to “a peptide sequence” includes a plurality of such sequences and so forth.

**[0470]** As used herein, numerical values are often presented in a range format throughout this document. The use of a range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention unless the context clearly indicates otherwise. Accordingly, the use of a range expressly includes all possible subranges, all individual numerical values within that range, and all numerical values or numerical ranges including integers within such ranges and fractions of the values or the integers within ranges unless the context clearly indicates otherwise. This construction applies regardless of the breadth of the range and in all contexts throughout this patent document. Thus, for example, reference to a range of 90-100% includes 91-99%, 92-98%, 93-95%, 91-98%, 91-97%, 91-96%, 91-95%, 91-94%, 91-93%, and so forth. Reference to a range of 90-100% also includes 91%, 92%, 93%, 94%, 95%, 95%, 97%, *etc.*, as well as 91.1%, 91.2%, 91.3%, 91.4%, 91.5%, *etc.*, 92.1%, 92.2%, 92.3%, 92.4%, 92.5%, *etc.*, and so forth.

**[0471]** In addition, reference to a range of 1-3, 3-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-110, 110-120, 120-130, 130-140, 140-150, 150-160, 160-170, 170-180, 180-190, 190-200, 200-225, 225-250 includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, *etc.* In a further example, reference to a range of 25-250, 250-500, 500-1,000, 1,000-2,500, 2,500-5,000, 5,000-25,000, 25,000-50,000 includes any numerical value or range within or encompassing such values, *e.g.*, 25, 26, 27, 28, 29...250, 251, 252, 253, 254...500, 501, 502, 503, 504..., *etc.*

**[0472]** As also used herein a series of ranges are disclosed throughout this document. The use of a series of ranges include combinations of the upper and lower ranges to provide another range. This construction applies regardless of the breadth of the range and in all contexts throughout this patent document. Thus, for example, reference to a series of ranges such as 5-10, 10-20, 20-30, 30-40, 40-50, 50-75, 75-100, 100-150, includes ranges such as 5-20, 5-30, 5-40,

5-50, 5-75, 5-100, 5-150, and 10-30, 10-40, 10-50, 10-75, 10-100, 10-150, and 20-40, 20-50, 20-75, 20-100, 20-150, and so forth.

**[0473]** For the sake of conciseness, certain abbreviations are used herein. One example is the single letter abbreviation to represent amino acid residues. The amino acids and their corresponding three letter and single letter abbreviations are as follows:

alanine	Ala	(A)
arginine	Arg	(R)
asparagine	Asn	(N)
aspartic acid	Asp	(D)
cysteine	Cys	(C)
glutamic acid	Glu	(E)
glutamine	Gln	(Q)
glycine	Gly	(G)
histidine	His	(H)
isoleucine	Ile	(I)
leucine	Leu	(L)
lysine	Lys	(K)
methionine	Met	(M)
phenylalanine	Phe	(F)
proline	Pro	(P)
serine	Ser	(S)
threonine	Thr	(T)
tryptophan	Trp	(W)
tyrosine	Tyr	(Y)
valine	Val	(V)

**[0474]** The invention is generally disclosed herein using affirmative language to describe the numerous embodiments. The invention also specifically includes embodiments in which particular subject matter is excluded, in full or in part, such as substances or materials, method steps and conditions, protocols, procedures, assays or analysis. Thus, even though the invention is generally not expressed herein in terms of what the invention does not include, aspects that are not expressly included in the invention are nevertheless disclosed herein.

[0475] A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, the following examples are intended to illustrate but not limit the scope of invention described in the claims.

## 5. EXAMPLES

[0476] The examples in this section (*i.e.*, Section 5) are offered by way of illustration, and not by way of limitation.

### 5.1 Example 1: Generation of Anti-PD-1 Antibodies

#### 5.1.1 Generation of anti-PD-1 antibodies

[0477] The parental PD-1-IgG1 mAb was initially generated by mouse immunization methods using human PD-1 extracellular domain (ECD) antigen or CHO-hPD-1 transfected cells. Initial characterization of a hybridoma pool identified a subclone designated PD1Sub1 producing an anti-human PD-1, PD-L1 non-blocking and PD-L2 non-blocking antibody (data not shown) with  $K_D \sim 6$  nM measured by Biacore<sup>®</sup> to soluble antigen. The mouse  $V_H$  and  $V_L$  genes from the PD1Sub1 hybridoma were sequenced and used for human CDR-grafting into human  $\gamma 1$  and  $\kappa$  constant regions of the most homologous human  $V_H$  and  $V_L$  framework genes (IgH1-f and V $\kappa$ 4-1, respectively) utilizing the closest J regions (IgH J6 and Ig $\kappa$  J2, respectively). For human germline HG1  $V_H$  and  $V_L$  regions only, the mouse CDR3 segments were placed into the human  $V_H$  and  $V_L$  framework germline genes, IgH 1-f and V $\kappa$  4-1, respectively.

[0478] Stable HEK-293c18 cell lines expressing either CDR-grafted or germline HG1 antibodies in Deciduous<sup>TM</sup> constructs with stable AID (Activation-Induced Deaminase) were generated for use in a SHM-XEL<sup>TM</sup> affinity maturation platform (AnaptysBio, San Diego, CA). *In situ* generation of genetic diversity in the antibody variable domain resulted in cells expressing higher affinity variants of the parental antibody. These were isolated by flow cytometry using monomeric or dimeric hPD-1. Multiple rounds of affinity purification and selection yielded 6 corridors and 45 clones. Sanger and deep sequencing of these clones, along with additional “*in silico* SHM” events, resulted in rounds of site directed mutagenesis incorporating enriching mutations into  $V_H$  and  $V_L$  CDRs. The highest affinity binding 12 muteins were further characterized biochemically, biophysically for binding kinetics to PD-1, and for binding to full length PD-1 expressed on the surface of CHO cells. Functional characterization of the purified

antibodies was performed in two assays: PD-L1 competition for binding to cell surface PD-1 and inhibitory activity in IL-2 production from reactivation of activated human CD4+ T cells.

[0479] Based on these methods, anti-PD-1 antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 were generated, as shown in **Table 9**.

**Table 9. Characterization of anti-PD-1 antibodies**

Antibody ID	HC/LC TOPO Vector ID	CDR-grafted HG1 HC/LC mutations	K <sub>D</sub> (Biacore) K <sub>D</sub> (KinExA)	PD-L1 Competition (IC <sub>50</sub> )	CD4 <sup>+</sup> T cell IL-2 inhibition (EC <sub>50</sub> )
PD1AB-1	3015/3017	parental/parental	5 nM (n=4) 575 pM (n=2)	>100 nM (n=4)	22 $\pm$ 4 nM (n=5)
PD1AB-2	3015/3193	parental/germline	5 nM (n=2) 425 pM (n=1)	>100 nM (n=2)	27 $\pm$ 4 nM (n=3)
PD1AB-3	3653/3646	D76N/S77N	7 nM (n=2) 350 pM (n=1)	>100 nM (n=1)	21 nM (n=1)
PD1AB-4	3653/3193	D76N/germline	6 nM (n=2) 500 pM (n=1)	>100 nM (n=1)	23 nM (n=1)
PD1AB-5	3650/3193	V24A/germline	6 nM (n=2) 400 pM (n=1)	>100 nM (n=1)	15 nM (n=1)
PD1AB-6	3650/3017	V24A/parental	6 nM (n=2) 450 pM (n=1)	>100 nM (n=1)	16 $\pm$ 3 nM (n=3)

### 5.1.2 CD4+ reactivation assay in human PBMCs or human whole blood

[0480] PD-1 expression was induced on human PBMCs isolated from leukocyte reduction system (LRS) with PHA activation 48 hours at 37 °C. CD4+ T cells were purified from the PBMCs using CD4 isolation kits (Miltenyi Biotec, San Diego, CA) and replated onto 96-wells with immobilized anti-CD3 or anti-CD3 plus titrated anti-PD-1 or hIgG1 isotype control antibodies. Supernatants were collected at 24 and 48 hours for IL-2, IFN- $\gamma$ , and IL-17 cytokine determinations. All six anti-PD-1 clones showed similar inhibition EC<sub>50</sub>s, ranging from 20-36 nM for IL-2, 34-58 nM for IFN- $\gamma$ , and 27-41 nM for IL-17, as shown in **Table 10**.

**Table 10. Comparison of T cell attenuating activity of 6 lead antibodies in PBMC reactivation assay**

	nM	PD1AB-1	PD1AB-3	PD1AB-2	PD1AB-4	PD1AB-5	PD1AB-6	hIgG1
IL-2	EC <sub>50</sub> EC <sub>75</sub> nM	21 $\pm$ 9 40 $\pm$ 9 (n=6)	24 $\pm$ 5 46 $\pm$ 9 (n=2)	24 $\pm$ 8 42 $\pm$ 9 (n=6)	20 $\pm$ 7 39 $\pm$ 8 (n=6)	36 $\pm$ 11 58 $\pm$ 18 (n=2)	24 $\pm$ 7 33 $\pm$ 5 (n=6)	>133

	nM	PD1AB-1	PD1AB-3	PD1AB-2	PD1AB-4	PD1AB-5	PD1AB-6	hIgG1
IFN- $\gamma$	EC <sub>50</sub> EC <sub>75</sub> nM	46 $\pm$ 21 85 $\pm$ 31 (n=3)	ND	58 $\pm$ 14 91 $\pm$ 29 (n=3)	34 $\pm$ 21 64 $\pm$ 29 (n=3)	ND	36 $\pm$ 22 67 $\pm$ 30 (n=3)	>133
IL-17	EC <sub>50</sub> EC <sub>75</sub> nM	41 $\pm$ 9 65 $\pm$ 9 (n=5)	ND	36 $\pm$ 12 59 $\pm$ 10 (n=5)	27 $\pm$ 11 52 $\pm$ 11 (n=5)	ND	37 $\pm$ 11 56 $\pm$ 13 (n=5)	>133

[0481] Inhibition of specific T cell function was then assessed in a human whole-blood matrix. Freshly drawn and heparinized human blood was plated onto wells immobilized with either anti-CD3 or anti-CD3 plus titrated anti-PD-1 or hIgG1 isotype control antibodies. Collected plasma at 24 and 48 hours were measured for IL-2 (24 hours) and IFN- $\gamma$  /IL-17 (48 hours). Compared to three other tested antibody clones, PD1AB-6 showed a 2-3 fold increased potency in specific IL-2 (EC<sub>50</sub> 4.0+0.9 nM, n=4), IFN- $\gamma$  (EC<sub>50</sub> 4.1+2.2 nM, n=2), and IL-17 (EC<sub>50</sub> 3.6+1.2 nM, n=3) inhibition, as shown in **Table 11**.

**Table 11. Comparison of T cell attenuating activity of 4 lead molecules in whole blood assay**

EC <sub>50</sub> nM		PD1AB-1	PD1AB-2	PD1AB-4	PD1AB-6	hIgG1
IL-2	Hu (n=4)	8.3 $\pm$ 2.7	8.4 $\pm$ 1.1	10.2 $\pm$ 2.0	4.0 $\pm$ 0.9	>133
hIFN- $\gamma$	Hu (n=2)	5.9 $\pm$ 0.8	7.6 $\pm$ 0.9	9.2 $\pm$ 1.3	4.1 $\pm$ 2.2	>133
hIL-17	Hu (n=3)	7.2 $\pm$ 1.3	8.9 $\pm$ 2.3	9.9 $\pm$ 3.7	3.6 $\pm$ 1.2	>133

### 5.1.3 Cell-based ligand binding assay

[0482] To assess ligand competition, a cell binding assay was put in-place to evaluate the identified six antibody clones. Briefly, individual antibody clones at semi-log concentrations

from 100 nM to 100 pM were pre-mixed with 10 nM DyL650-PD-L1 before adding to human PD-1-CHO cells ( $2 \times 10^5$  cells) for 45 minutes on ice. Cells were then washed before DyL650-PD-L1 binding was analyzed on a BD FACSArray<sup>TM</sup> and median fluorescence intensity relative to isotype control antibody was plotted at each concentration. As shown in **FIGS. 1A-1B**, PD1AB-6, as well as the other five clones, including the parental clone PD1AB-1, showed no significant competition against DyL650-PD-L1 binding up to 100 nM. In contrast, MDX 4H1 (AnaptysBio, San Diego, CA), an antagonist, ligand-blocking, PD-1 antibody dose-dependently blocked labeled PD-L1 binding, generating a binding EC<sub>50</sub> ~ 5-10 nM.

#### 5.1.4 Epitope mapping

**[0483]** The PD-1 epitope was determined by solving the crystal structure of the PD1AB-6 Fab in complex with the human PD-1 extracellular domain to 1.8 Å resolution. The PD-1:PD1AB-6 Fab interaction site occurs on a distal side of PD-1 relative to the PD-1:PD-L1 interaction site (**FIG. 2**), consistent with the observation that PD-L1 and PD1AB-6 do not compete for PD-1 binding. PD1AB-6 Fab binds against a PD-1 β sheet, with substantial interactions formed with a PD-1 loop composed of residues 100-105 (**FIG. 3**). R104 on PD-1 engages multiple polar interactions with residues on the Fab CDR H1. The adjacent residue G103 also makes a tight polar interaction with the Fab. Both R104 and G103 are mutated in mouse PD-1 (to a histidine and arginine, respectively), providing a structural rationale for the lack of binding of PD1AB-6 to murine PD-1. The PD1AB-6 Fab regions that interact with PD-1 are the CDR H1, H2, H3, L1 and L2. Atomic details of the PD-1:PD1AB-6 Fab interactions are described in the **Table 12**. HC and LC residues that interact with PD-1 epitope are described. Abbreviations are as follows: HB-hydrogen bond, HYD-hydrophobic interaction, ION-ionic interaction.

**Table 12. Atomic details of PD1AB-6:PD-1 interaction**

Type	Chain	Pos	Chain	Pos
HB	PD-1	ASN103.NH2	H	ALA71.O
HB	PD-1	THR51.OY1	L	SER50.O
HB	PD-1	SER57.OY	H	ASP69.OE1
HYD	PD-1	LEU100.CS1	L	PHE55.CY
HYD	PD-1	LEU100.CS1	L	LEU115.CD1
HB	PD-1	ASN103.NH2	L	TYR114.O
HB	PD-1	ASN102.OE1	L	SER117.N
HB	PD-1	ASN102.O	H	TYR50.OH
HB	PD-1	GLY103.O	H	TYR126.OH
HB	PD-1	ASN104.NH2	H	LYS57.O
HB	PD-1	ASN104.NH1	H	ASP49.O
HB	PD-1	ASN104.NH2	H	ASP69.OE
HB	PD-1	ASN104.NH2	H	ASP69.OE
HB	PD-1	ASP105.OE1	H	SER125.OY
HB	PD-1	ASP105.OE1	H	SER125.OY
HB	PD-1	SER109.OY	L	SER50.O

### 5.1.5 Generation of variants of PD1AB-6

**[0484]** PD1AB-6 IgG1 antibody (PD1AB-6-IgG1) and Fc modified IgG4PE antibody (PD1AB-6-4PE) were generated. PD1AB-6-4PE was designed to have significantly lower Fc-mediated effector function. The CH region,  $\gamma 4$  contains two non-standard amino acids substitutions, S228P and L235E (EU numbering systems, Kabat and Wu 1991). Serine 228, a common amino acid type in the hinge of IgG4, was changed to proline, a less commonly observed amino acid type in IgG4 and highly conserved amino acid in IgG1. This change significantly reduced the level of “half-antibody” that is commonly observed in the production of IgG4-subclass antibody. Leucine 235, one of the critical amino acids involved in heavy chain interactions with Fc $\gamma$  receptors was changed to glutamic acid. The L235E substitution significantly reduced the interaction of  $\gamma 4$  chain to Fc $\gamma$ R, eliminating ADCC and Fc-receptor-mediated elimination of PD-1-expressing normal cells. In addition, inherent lack of complement binding by  $\gamma 4$  heavy chain renders the PD1AB-6-4PE molecule devoid of CDC function. Two other variants were generated to minimize binding affinity to C1q for reduced CDC (**FIG. 4**). To generate PD1AB-6-K3, lysine 322 was substituted with alanine in PD1AB-6-IgG1. The K322A substitution is reported to suppress C1q binding on rituximab, a chimeric antibody with a human IgG1 Fc (Idusogie *et al.*, 2000, *J. Immunol.* 164(8):4178-84). PD1AB-6-4P was generated by converting the Fc-backbone of the PD1AB-6-IgG1 to the Fc-backbone of IgG4 with S228P substitution. Serine 228, a common amino acid type in the

hinge of IgG4, was changed to proline, a less commonly observed amino acid type in IgG4 and highly conserved amino acid in IgG1. This change significantly reduces the level of half- antibody that is frequently observed in the production of IgG4-subclass antibody. IgG4 antibody was reported to have attenuated ADCC and CDC function (Overdijk *et al.*, 2012, *J. Immunol.* 189(7):3430-38). All changes were created in the CH region with no changes in the variable regions. The amino acid sequences of the heavy and light chains of PD1AB-6-IgG1 are labeled LC\_PD1AB-6-IgG1 and HC\_PD1AB-6-IgG1, respectively (**FIG. 4**). The two heavy chain variants include HC\_PD1AB-6-IgG1-K322A and HC\_PD1AB-6-IgG4P. The light chain LC\_PD1AB-6-IgG1 is paired with the three individual heavy chains to generate PD1AB-6-IgG1, PD1AB-6-K3, and PD1AB-6-4P, respectively.

### **5.1.6 Cell line development and antibody manufacturing from transient transfection**

#### **5.1.6.1 Molecular Cloning of the Heavy and Light Chains**

**[0485]** IgG LC expression vector pFUSE2ss-CL Ig-hk and IgG HC expression vector pFUSEss-CH Ig-hG1 were purchased from InvivoGen (San Diego, CA).

**[0486]** The amino acid sequence encoding LC\_PD1AB-6-IgG1 (**FIG. 4**) was converted into a codon-optimized gene sequence for protein expression in mammalian cells. Restriction enzyme sites EcoRI at the 5'-end and NheI at the 3'-end were added to the optimized gene. The optimized LC gene with EcoRI and NheI sites were synthesized producing an insert fragment. The IgG LC expression vector pFUSE2ss-CL Ig-hk was digested with EcoRI and NheI producing approximately a 3.5 kb pFUSE2ss-CL Ig-hk-EcoRI/NheI fragment. The insert fragment was ligated into pFUSE2ss-CL Ig-hk-EcoRI/NheI fragment resulting in production of pJS-1 which is pFUSE2ss-CL Ig-hk-LC\_PD1AB-6-IgG1.

**[0487]** The amino acid sequence encoding HC\_PD1AB-6-IgG1, HC\_PD1AB-6-IgG1-K322A, or HC\_PD1AB-6-IgG4P (**FIG. 4**) was converted into a codon-optimized gene sequence for protein expression in mammalian cells. Restriction enzyme sites EcoRI at the 5'-end, a constant region from a stop codon (after the 3'-end of HC sequence in pFUSEss-CH Ig-hG1) to HpaI at the 3'-end were added. The optimized genes with EcoRI and HpaI sites were synthesized producing insert fragments containing the genes encoding HC\_PD1AB-6-IgG1, HC\_PD1AB-6-IgG1-K322A, and HC\_PD1AB-6-IgG4P, respectively. The IgG HC expression vector pFUSEss-CH Ig-hG1 was digested with EcoRI and HpaI producing

approximately a 3.4 kb pFUSEss-CHIg-hG1-EcoRI/HpaI fragment. The insert fragments were ligated into pFUSEss-CHIg-hG1-EcoRI/HpaI fragment resulting in production of pJS-2, pJS-3, and pJS-12, which are pFUSEss-CHIg-hG1-HC\_PD1AB-6-IgG1, pFUSEss-CHIg-hG1-HC\_PD1AB-6-IgG1-K322A, and pFUSEss-CHIg-hG1-HC\_PD1AB-6-IgG4P, respectively.

#### **5.1.6.2 Protein Production**

**[0488]** All three variants of PD1AB-6-IgG1, PD1AB-6-K3, and PD1AB-6-4P were manufactured at the laboratory scale in shake-flasks for *in vitro* and *in vivo* efficacy studies. PD1AB-6-4P and PD1AB-6-K3 antibodies for non-GLP toxicology studies and additional characterization were manufactured in 50 L bioreactors (50 L stirred tanks and 50 L wave bags) using FreeStyle<sup>TM</sup> MAX CHO expression system as well as Expi293<sup>TM</sup> expression system from Life Technologies (Carlsbad, CA). FreeStyle<sup>TM</sup> MAX CHO expression system was used for transient transfection of CHO-S cells using manufacturer's standard protocol. Expi293<sup>TM</sup> expression system was used for transient transfection of Expi293 cells using manufacturer's standard protocol. A 3:2 ratio of light chain versus heavy chain was used for DNA mixture at 1 mg per 1 L of culture during the transfection. Cells were seeded at 0.5 million cells/mL in a 50 L bioreactor at 37 °C and grew over night to reach 1 million cell/mL. Cells then were transfected using manufacturer's standard protocols. On day one post-transfection, 1 mM sodium butyrate plus 1% v/v of feed media (Yeastolate, CHO CD EfficientFeed<sup>TM</sup> A, Glutamax and Glucose) was added to bioreactor, and temperature was dropped to 32 °C. Forty liters of cells plus additives were seeded for a 50 L stirred tank, and 25 L of cells plus additives were seeded for a 50 L wave bioreactor. Cell viability and titer were monitored every day, and batches were harvested when cell viability dropped below 50%. Vi-Cell<sup>TM</sup> instrument was used for viability analysis, and Octet RED equipped with Anti-Human IgG sensor was used for titer analysis using purified antibody for the standard curve. Cells and supernatant were harvested using GE Life Sciences depth filtration and sterilization columns, ULTA Prime GF 5 µm capsules were used for depth filtration followed by ULTA Pure HC 0.6/0.2 µm sterilization capsules. Clarified supernatant were concentrated 5-8 fold using cross flow filtration, 50 Kd cut off Kvick<sup>TM</sup> Lab SCU from GE Life Science were used for TFF. The titer and maximum cell densities obtained for each isotype at harvest are given in **Table 13**.

**Table 13. Productivity in Fed-batch Bioreactor (50 Liter)**

Isotype Cell pool	Cell Line/Volume	Titer (mg/l)	Cell density at harvest (1 x 10 <sup>6</sup> cells/mL)
<b>PD1AB-6-4P</b>	Expi293/25L	22	3.2
<b>PD1AB-6-4P</b>	CHO-S/50L	9	3
<b>PD1AB-6-K3</b>	Expi293/50L	31	4.5
<b>PD1AB-6-K3</b>	CHO-S/50L	15	2

### 5.1.6.3 Protein Purification

**[0489]** Purification of the materials produced was performed by a series of downstream purification steps including protein A affinity chromatography and low pH virus inactivation, followed by IEX interaction (Capto<sup>TM</sup> Adhere & Capto<sup>TM</sup> SP ImpRes) chromatography steps. The purified antibody is bulk formulated by buffer exchanged against (10 mM Succinate pH 5.5, 9% sucrose, 0.05% PS20) buffer, filtered through 0.2 µm filter, and aliquoted.

**[0490]** The protein A affinity chromatography was carried out with MabSelect SuRe<sup>TM</sup>, designed to capture the product and to remove process related impurities. The subsequent virus inactivation step was performed under acidic conditions (pH 3.4 ± 0.1 for 45 min.) followed by conditioning of the inactivation pool to pH 5.5 ± 0.1. After virus inactivation, an anion exchanger was used in a flow-through mode for intermediate polishing step using Capto<sup>TM</sup> Adhere to remove impurities such as aggregates, DNA, host cell protein, and endotoxin. The product pool was conditioned to pH 6.5 ± 0.1, and the conductivity was reduced to 2 mS/cm prior to the next process step. Cation exchanger Capto<sup>TM</sup> SP ImpRes was used as a polishing step, and the product was resolved at 10 mS/cm. The antibody was then buffer exchanged in stock solution (10 mM Succinate, 9% sucrose, 0.05% PS20, pH 5.5) and concentrated to 20 mg/mL. The product pool was then filtered through 0.2 µm filter and aliquoted.

### 5.1.7 Cell-based PD-1 binding assay

**[0491]** PD1AB-6-IgG1 binding was evaluated on CHO cells expressing human PD-1 and cynomolgus PD-1 (**FIGS. 5A-5B**), and on primary human PBMC (**FIG. 6**) and cynomolgus PBMC (**FIG. 7**).

**[0492]** CHO cells expressing human PD-1 and cynomolgus PD-1 were incubated with various concentration of unlabeled PD1AB-6-IgG1 antibody for 30 minutes at 4 °C, washed, and stained with anti-human IgG Fc (eBioscience, San Diego, CA) for 30 minute at 4 °C. Human

IgG1 Fc was used as a negative control. PD1AB-6-IgG1 binds to human PD-1 expressed on CHO cells with an EC<sub>50</sub> = 0.4 nM and binds to cynomolgus PD-1 expressed on CHO cells with an EC<sub>50</sub> = 0.8 nM (**FIGS. 5A-5B**).

**[0493]** Human PBMCs were activated with 1  $\mu$ g/mL plate bound anti-CD3 for 3 days to induce PD-1 expression on T cells. Cells were incubated with various concentration of unlabeled PD1AB-6-IgG1 antibody for 30 minutes at 4 °C, washed, and stained with anti-human IgG Fc (eBioscience, San Diego, CA) for 30 minute at 4 °C. Human IgG1 Fc was used as a negative control. Geometric MFI was determined on CD4+ T cells. Data from 1 of 2 human healthy donors are shown in **FIG. 6**.

**[0494]** Cynomolgus PBMCs were activated with 1  $\mu$ g/mL anti-cynomolgus CD3/CD28 for 2 days to induce PD-1 expression on T cells. Cells were incubated with various concentration of unlabeled PD1AB-6-IgG1 antibody for 30 minutes at 4 °C, washed, and stained with anti-human IgG Fc (eBioscience) for 30 minute at 4 °C. Human IgG1 Fc was used as a negative control. Geometric MFI was determined on CD4+ T cells. Data from 1 of 2 cynomolgus donors are shown in **FIG. 7**.

### 5.1.8 Fc receptor binding assay

**[0495]** To confirm the objectives of variants generation, *i.e.*, decreased Fc $\gamma$ R-mediated effector function, Fc $\gamma$ R binding to the PD1AB-6-K3 and PD1AB-6-4P variants were analyzed by two methodologies. First, binding was tested with displacement Fc $\gamma$ R assays using Cisbio Tag-lite® detection (**FIGS. 8A-8D**). HEK293 cells engineered to express specific Fc $\gamma$ Rs (Fc $\gamma$ RI, Fc $\gamma$ RIIIa, or Fc $\gamma$ RIIb) prelabeled with a terbium (Tb) donor dye were mixed with reference controls or PD1AB-6-IgG1, PD1AB-6-K3, and PD1AB-6-4P antibodies over log concentrations ranging from 10000 nM to 0.1 pM. A second, human-hIgG-d2 (acceptor) was then added to compete for receptor binding. Detection of a Fluorescence Resonance Energy Transfer (FRET) signal generated by Tb-d2 proximity is measured and is inversely proportional to PD1AB-6 variant-bound Fc $\gamma$ R. As shown in **FIGS. 8A-8D**, the PD1AB-6-K3 variant showed decreased binding to Fc $\gamma$ RIIIa (CD16), the low affinity receptor on NK cells responsible for ADCC activity. Binding to Fc $\gamma$ RI (expressed on granulocytes, dendritic cells (DCs), or monocytes) was similar to parental PD1AB-6-IgG1 molecule.

**[0496]** Second, both PD1AB-6-K3 and PD1AB-6-4P variants were tested in FACS-based binding assays (**FIGA. 9A-9C**). Briefly, Fc $\gamma$ RI-CHO or Fc $\gamma$ RIIIaV158-CHO expressing cell

lines were detached and washed prior to mixing with the PD1AB-6-K3 and PD1AB-6-4P variants over different concentrations for 1 hour on ice. PD1AB-6 variant-bound cells were detected with a labeled PE-conjugated F(ab')<sub>2</sub> goat anti-human secondary antibody for an additional hour on ice, washed, and fixed prior to analysis by FACS, and mean fluorescence intensity was plotted at each concentration. The PD1AB-6-4P variant returned significantly higher binding EC<sub>50</sub>s (>15X and >21X) against Fc $\gamma$ RI and Fc $\gamma$ RIIIa lines, respectively (**FIGS. 9A-9C**).

#### 5.1.9 *In vitro* ADCC assay

**[0497]** The ability of PD1AB-6 variants to induce ADCC was evaluated in co-culture assay involving natural killer (NK) cells from healthy donors and PD-1 expressing target cells. Target cells (NCI-OCI-Ly3) pre-treated with PD1AB-6 variants were co-cultured with activated NK cells for 4 hours. Supernatant LDH concentration was used to calculate specific lysis. EC<sub>50</sub> (nM) was calculated using Prism. Error bar represents experimental triplicates. Data are 2 representatives of 4 individual healthy donors. As shown in **FIGS. 10A-10B**, titration of PD1AB-6-IgG1 induced dose dependent ADCC, while PD1AB-6-K3 showed reduced ADCC activity.

#### 5.1.10 *In vitro* CDC assay

**[0498]** The ability of PD1AB-6 variants to induce CDC was evaluated using PD-1 expressing CD20<sup>+</sup> NCI-OCI-Ly3 cells. Target cells (NCI-OCI-Ly3) pre-treated with antibodies were cultured in serum-free media supplemented with 5% rabbit complement for 4 hours. Cell lysis was determined by 7-AAD<sup>+</sup> cells by FACS. Data are representative of 3 independent experiments: (i) CDC activity of PD1AB-6-IgG1 and anti-CD20 IgG1; (ii) CDC activity of PD1AB-6-IgG1 and PD1AB-6-K3; (iii) CDC activity of PD1AB-6-4P and commercial mouse anti-PD-1 IgG1 antibody. As shown in **FIG. 11**, PD1AB-6-K3 consistently did not induce CDC (n=3). Parental PD1AB-6-IgG1 and PD1AB-6-4P also did not induce CDC. This was not due to resistance of the target cell line to complement killing, since anti-CD20 IgG1 repeatedly induced dose dependent CDC on NCI-OCI-Ly3 cells in presence of 5% rabbit complement.

## 5.2 Example 2: Activity Assays

### 5.2.1 Human T cell activation assay

[0499] Functional assessment of PD1AB-6 variants on inhibiting T cell effector function was performed by two methods. In one assay, the peripheral blood mononuclear cells were preactivated to express PD-1 and restimulated in the presence of soluble PD1AB-6-K3 (**FIG. 12**). Peripheral blood mononuclear cells (PBMCs), from healthy donors were preactivated with the mitogen, PHA, for 48 h to upregulate PD-1 expression. These cells were then restimulated using anti-CD3 conjugated Dynabeads<sup>®</sup> (Life Technologies Carlsbad, CA) in the presence of diluted PD1AB-6 variants over a range of 100 nM to 0.1 nM final concentration. T cell activation was measured using IL-2 levels in culture supernatants at 24 h post-stimulation. As shown in **FIG. 12**, PD1AB-6-K3 and the two other variants showed potent T cell inhibitory activity in this assay with an EC<sub>50</sub> of 5-25 nM.

[0500] A second assay was used for direct *ex vivo* measurement of PD1AB-6-K3 in inhibiting T cell function (**FIG. 13**). This was done by directly plating fresh human whole blood in 96-well plates co-coated with anti-CD3 +/- PD1AB-6-K3, and measuring IL-17 and IFN- $\gamma$  levels as readouts of T cell activation. CTLA4Ig (Orencia<sup>®</sup>) was used as a positive control in these assays, and human IgG Fc fragment was used as a negative control. As shown in **FIG. 13**, overall, PD1AB-6-K3 trended towards better efficacy than PD1AB-6-4P. Negative control, hIgG Fc, showed no activity with an EC<sub>50</sub> > 100 nM.

### 5.2.2 Cynomolgus monkey crossreactivity assay

[0501] Determination of functional cynomolgus monkey cross reactivity with lead PD1AB-6-K3 was performed similarly to human samples, using freshly isolated cynomolgus PBMCs, activated with anti-cynomolgus CD3, CD28, and PD1AB-6 variants as indicated. CTLA4Ig was used as a positive control in these assays, and hIgG1 Fc was used as a negative control. Culture supernatants were removed after 48 hours for cytokine determinations using cynomolgus IL-2 MSD assays. As shown in **Table 14**, these assays demonstrated that PD1AB-6-K3 attenuated cynomolgus T cells cytokine secretion to levels comparable to the positive control, CTLA4Ig, and the activity was comparable to that seen in human assays.

**Table 14. PD1AB-6-K3 activity in cynomolgus PBMC assay**

	PD1AB-6-IgG1	PD1AB-6-K3	PD1AB-6-4P	CTLA4Ig
	IL-2 EC50 (nM)			
<b>NHP1</b>	0.28	0.24	<0.1	Not done
<b>NHP2</b>	4.9	2.5	1.24	Not done
<b>NHP3</b>	N.D.	9.24	16.5	14
<b>NHP4</b>	N.D.	1.22	1.23	6

### 5.2.3 *In vitro* mechanism of action

**[0502]** Several antibodies that bind to T cell surface molecules, such as CD3 and CD4, lead to signaling and subsequent downregulation of surface expression of those molecules. Since PD1AB-6 antibody is designed to provide an agonist signal via PD-1, it was of interest to evaluate PD-1 expression after PD1AB-6 treatment *in vitro*.

#### 5.2.3.1 Decreased PD-1 Expression after PD1AB-6 Treatment

**[0503]** Human PBMC from different donors were activated with 1 µg/mL plate bound anti-CD3 + 0.25 µg/mL plate bound anti-CD28 with various concentration of soluble control IgG1 or PD1AB-6-IgG1. After 4 hours to 72 hours incubation, cells were stained for CD3, CD45RO, and PD-1 to assess PD-1 expression on T cells. **FIGS. 14A-14C** show reduced expression of PD-1 on human CD3+ T cells after 48 hours PD1AB-6-IgG1 treatment. Analysis of PD-1 expression showed that PD1AB-6-IgG1 treatment led to downregulation of PD-1 expression on the surface of T cells (representative histogram from one donor in **FIGS. 14A-14B** and analysis of mean fluorescence intensity at 48 hours, across three donors and various antibody concentrations in **FIG. 14C**). PD-1 downregulation was seen as early as 4 hours of incubation with PD1AB-6-IgG1. Downregulation of surface PD-1 expression was likely due to PD1AB-6 induced signaling via PD-1, and was through similar mechanism to what is observed with T cell receptor signaling (San Jose *et al.*, 2000, *Immunity* 12(2):161-70).

## 5.3 Example 3: Physicochemical Characterization of PD1AB-6 Variants

### 5.3.1 Biacore® binding analysis

**[0504]** Purified PD1AB-6 variant antibodies were analyzed on Biacore® T200 for binding to hPD1 antigen using capture method. Fc-specific anti-human IgG was immobilized on Fc2, and Fc1 was left blank as reference channel. Purified PD1 antibodies were captured on anti-Human IgG, and internally produced hPD1 antigen (PD1\_002) was flowed over both channels using two fold dilution series from 100 nM to 200 pM to determine kinetics of binding. The PD-1 used

was the extracellular domain of human PD-1 (residues 32-160) expressed in *E.Coli* as inclusion bodies and refolded. Surface was regenerated between each antigen concentration using 3M Magnesium Chloride. Examples of binding kinetics as well as values of  $k_{on}$ ,  $k_{off}$ , and  $K_D$  for PD1AB-6-IgG1, PD1AB-6-4P, and PD1AB-6-K3 are shown in **FIGS. 15A-15C**. All three variants had similar rates of association and dissociation to the PD-1 antigen with comparable  $K_D$  values of 19-22 nM.

## 6. SEQUENCE LISTING

**[0505]** The present specification is being filed with a computer readable form (CRF) copy of the Sequence Listing. The CRF entitled 10624-294-228\_SEQLIST.txt, which was created on September 22, 2016 and is 50,870 bytes in size, is identical to the paper copy of the Sequence Listing and is incorporated herein by reference in its entirety.

## WHAT IS CLAIMED:

1. An antibody or antigen-binding fragment thereof that
  - (a) binds to an epitope of human PD-1 recognized by an antibody comprising a light chain variable region having an amino acid sequence of SEQ ID NO:8 and a heavy chain variable region having an amino acid sequence of SEQ ID NO:13; or
  - (b) competes for the binding to human PD-1 with an antibody comprising a light chain variable region having an amino acid sequence of SEQ ID NO:8 and a heavy chain variable region having an amino acid sequence of SEQ ID NO:13.
2. An antibody or antigen-binding fragment thereof that binds to PD-1, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a light chain variable region (VL) comprising VL complementarity determining region 1 (CDR1), VL CDR2, and VL CDR3 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in Table 1; and/or
  - (b) a heavy chain variable region (VH) comprising VH complementarity determining region 1 (CDR1), VH CDR2, and VH CDR3 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in Table 2.
3. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a light chain variable region (VL) further comprising VL framework 1 (FR1), VL FR2, VL FR3, and VL FR4 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in Table 3; and/or
  - (b) a heavy chain variable region (VH) further comprising VH framework 1 (FR1), VH FR2, VH FR3, and VH FR4 of any one of antibodies PD1AB-1, PD1AB-2, PD1AB-3, PD1AB-4, PD1AB-5, or PD1AB-6 as set forth in Table 4.

4. The antibody or antigen-binding fragment thereof of claim 2, wherein the VL CDR1, VL CDR2, and VL CDR3 comprise amino acid sequences of SEQ ID NOS:1, 2, and 3, respectively, and the VH CDR1, VH CDR2, and VH CDR3 comprise amino acid sequences of SEQ ID NOS:4, 5, and 6, respectively.
5. The antibody or antigen-binding fragment thereof of claim 2, wherein the VL CDR1, VL CDR2, and VL CDR3 comprise amino acid sequences of SEQ ID NOS:7, 2, and 3, respectively, and the VH CDR1, VH CDR2, and VH CDR3 comprise amino acid sequences of SEQ ID NOS:4, 5, and 6, respectively.
6. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:8.
7. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:9.
8. The antibody or antigen-binding fragment of thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VL comprising an amino acid sequence of SEQ ID NO:10.
9. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:11.
10. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:12.
11. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a VH comprising an amino acid sequence of SEQ ID NO:13.

12. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and
  - (b) a VH comprising an amino acid sequence of SEQ ID NO:11.
13. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and
  - (b) a VH comprising an amino acid sequence of SEQ ID NO:11.
14. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and
  - (b) a VH comprising an amino acid sequence of SEQ ID NO:11.
15. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and
  - (b) a VH comprising an amino acid sequence of SEQ ID NO:12.
16. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and
  - (b) a VH comprising an amino acid sequence of SEQ ID NO:12.
17. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and
- (b) a VH comprising an amino acid sequence of SEQ ID NO:12.

18. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a VL comprising an amino acid sequence of SEQ ID NO:8; and
- (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

19. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a VL comprising an amino acid sequence of SEQ ID NO:9; and
- (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

20. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a VL comprising an amino acid sequence of SEQ ID NO:10; and
- (b) a VH comprising an amino acid sequence of SEQ ID NO:13.

21. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a human IgG1 Fc region or a mutant thereof.

22. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a human IgG1-K322A Fc region.

23. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a human IgG4 Fc region or a mutant thereof.

24. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a human IgG4P Fc region.
25. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a human IgG4PE Fc region.
26. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain Fc region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:36-40.
27. The antibody or antigen-binding fragment thereof of claim 26, wherein the antibody or antigen-binding fragment thereof further comprises a light chain constant region comprising an amino acid sequence of SEQ ID NO:41.
28. The antibody or antigen-binding fragment thereof of any one of claims 1-20, wherein the antibody or antigen-binding fragment thereof comprises:
  - (a) a light chain constant region comprising an amino acid sequence of SEQ ID NO:41; and
  - (b) a heavy chain Fc region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:36-40.
29. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises a light chain comprising an amino acid sequence of SEQ ID NO:31.
30. The antibody or antigen-binding fragment thereof of claim any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:32.
31. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and
- (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:32.

32. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:33.

33. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and
- (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:33.

34. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:34.

35. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and
- (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:34.

36. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain comprising an amino acid sequence of SEQ ID NO:35.

37. The antibody or antigen-binding fragment thereof of any one of claims 1-28, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a light chain comprising an amino acid sequence of SEQ ID NO:31; and
- (b) a heavy chain comprising an amino acid sequence of SEQ ID NO:35.

38. The antibody or antigen-binding fragment thereof of any one of claims 1-37, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to at least one of residues 100-109 within an amino acid sequence of SEQ ID NO:42.
39. The antibody or antigen-binding fragment thereof of claim 38, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to at least one of residues 100-105 within an amino acid sequence of SEQ ID NO:42.
40. The antibody or antigen-binding fragment thereof of any one of claims 1-37, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to at least one residue selected from the group consisting of N33, T51, S57, L100, N102, G103, R104, D105, H107, and S109 within an amino acid sequence of SEQ ID NO:42.
41. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to N33 within an amino acid sequence of SEQ ID NO:42.
42. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to T51 within an amino acid sequence of SEQ ID NO:42.
43. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to S57 within an amino acid sequence of SEQ ID NO:42.
44. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to L100 within an amino acid sequence of SEQ ID NO:42.
45. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to N102 within an amino acid sequence of SEQ ID NO:42.

46. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to G103 within an amino acid sequence of SEQ ID NO:42.
47. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to R104 within an amino acid sequence of SEQ ID NO:42.
48. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to D105 within an amino acid sequence of SEQ ID NO:42.
49. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to H107 within an amino acid sequence of SEQ ID NO:42.
50. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to S109 within an amino acid sequence of SEQ ID NO:42.
51. The antibody or antigen-binding fragment thereof of claim 40, wherein, when bound to PD-1, the antibody or antigen-binding fragment binds to G103 and R104 within an amino acid sequence of SEQ ID NO:42.
52. The antibody or antigen-binding fragment thereof of any one of claims 1-51, wherein the antibody or antigen-binding fragment thereof:
  - (a) attenuates T cell activity; and/or
  - (b) downregulates PD-1 expression on the surface of T cells.
53. The antibody or antigen-binding fragment thereof of any one of claims 1-52, wherein the antibody or antigen-binding fragment thereof specifically binds to human PD-1 and/or monkey PD-1, but not rodent PD-1.

54. The antibody or antigen-binding fragment thereof of any one of claims 1-53, wherein the antibody or antigen-binding fragment thereof has attenuated ADCC activity and/or attenuated CDC activity.
55. The antibody or antigen-binding fragment thereof of claim 52, wherein the attenuation of T cell activity occurs in human PBMC or whole blood samples.
56. The antibody or antigen-binding fragment thereof of claim 52 or 55, wherein the attenuation of T cell activity is measured by inhibition of cytokine production.
57. The antibody or antigen-binding fragment thereof of claim 56, wherein the cytokine that is inhibited by the antibody or antigen-binding fragment thereof comprises IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, TNF- $\alpha$ , and/or IFN- $\gamma$ .
58. The antibody or antigen-binding fragment thereof of claim 52, wherein the downregulation of PD-1 expression on the surface of T cells:
  - (a) occurs as early as 4 hours after the treatment with the antibody or antigen-binding fragment thereof; and/or
  - (b) is concurrent with or precedes cytokine inhibition.
59. The antibody or antigen-binding fragment thereof of claim 53, wherein the  $K_D$  for binding to purified human PD-1 is from about 100 pM to about 10 nM, and the  $K_D$  for binding to human PD-1 expressed on cell surface and monkey PD-1 expressed on cell surface is from about 100 pM to about 10 nM.
60. The antibody or antigen-binding fragment thereof of claim 56, wherein the  $EC_{50}$  for attenuating T cell activity is from about 1 pM to about 10 pM, from about 10 pM to about 100 pM, from about 100 pM to about 1 nM, from about 1 nM to about 10 nM, or from about 10 nM to about 100 nM.
61. The antibody or antigen-binding fragment thereof of claim 56, wherein the maximal percent attenuation of T cell activity is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

62. The antibody or antigen-binding fragment thereof of claim 52 or 58, wherein the maximal percent downregulation of PD-1 expression is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.
63. The antibody or antigen-binding fragment thereof of any one of claims 1-62, wherein the antibody is a monoclonal antibody.
64. The antibody or antigen-binding fragment thereof of any one of claims 1-63, wherein the antibody is a humanized, human, or chimeric antibody.
65. The antibody or antigen-binding fragment thereof of claim 64, wherein the humanized antibody is a deimmunized antibody or a composite human antibody.
66. The antibody or antigen-binding fragment thereof of any one of claims 1-65, wherein the antibody or antigen-binding fragment thereof is a Fab, a Fab', a F(ab')<sub>2</sub>, a Fv, a scFv, a dsFv, a diabody, a triabody, a tetrabody, or a multispecific antibody formed from antibody fragments.
67. The antibody or antigen-binding fragment thereof of any one of claims 1-66, wherein the antibody or antigen-binding fragment thereof is conjugated to an agent.
68. The antibody or antigen-binding fragment thereof of claim 67, wherein the agent is selected from the group consisting of a radioisotope, a metal chelator, an enzyme, a fluorescent compound, a bioluminescent compound, and a chemiluminescent compound.
69. A composition comprising the antibody or antigen-binding fragment thereof of any one of claims 1-68, and a pharmaceutically acceptable carrier.
70. A polynucleotide comprising nucleotide sequences encoding a VH, a VL, or both a VH and a VL of the antibody of any one of claims 1-68.
71. A polynucleotide comprising nucleotide sequences encoding a heavy chain, a light chain, or both a heavy chain and a light chain of the antibody of any one of claims 1-68.

72. The polynucleotide of claim 70 or 71, wherein the polynucleotide is operably linked to a promoter.
73. A vector comprising the polynucleotide of claim 71 or 72.
74. A cell comprising the polynucleotide of claim 71 or 72.
75. A cell comprising the vector of claim 73.
76. An isolated cell producing the antibody or antigen-binding fragment thereof of any one of claims 1-68.
77. A kit comprising the antibody or antigen-binding fragment thereof of any one of claims 1-68.
78. A method of making an antibody or antigen-binding fragment thereof which specifically binds to an epitope of human PD-1, comprising culturing the cell of any one of claims 74 to 76 to express the antibody or antigen-binding fragment thereof.
79. A method of making an antibody or antigen-binding fragment thereof which specifically binds to an epitope of human PD-1, comprising expressing the polynucleotide of any one of claims 70 to 72.
80. A method of attenuating activity of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen binding fragment thereof of any one of claims 1 to 68.
81. The method of claim 80, wherein the maximal percent attenuation of T cell activity is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.
82. The method of claim 80 or 81, wherein the attenuation of T cell activity is measured by inhibition of cytokine production.
83. The method of any one of claims 80 to 82, wherein the cytokine is IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, TNF- $\alpha$ , IFN- $\gamma$ , or any combination thereof.

84. The method of any one of claim 80 to 83, wherein the inhibition of cytokine production is concurrent with or follows downregulation of PD-1 expression on the surface of the T cell.
85. The method of claim 84, wherein the downregulation of PD-1 expression on the surface of the T cell occurs as early as 4 hours after contact with an effective amount of an antibody or antigen-binding fragment thereof.
86. A method of downregulating PD-1 expression on the surface of a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen binding fragment thereof of any one of claims 1 to 68.
87. The method of claim 86, wherein the maximal percent downregulation of PD-1 expression by the antibody or antigen-binding fragment thereof is at least about 10%, 20%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.
88. The method of claim 86 or 87, wherein the downregulation of PD-1 expression on the surface of the T cell occurs as early as 4 hours after contact with the antibody or antigen-binding fragment thereof.
89. The method of any one of claims 86 to 88, wherein the downregulation of PD-1 expression on the surface of the T cell precedes cytokine inhibition.
90. The method of claim 89, wherein the cytokine is IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, TNF- $\alpha$ , IFN- $\gamma$ , or any combination thereof.
91. A method of inhibiting cytokine production by a T cell, comprising contacting the T cell with an effective amount of an antibody or antigen binding fragment thereof of any one of claims 1 to 68.
92. The method of claim 91, wherein the cytokine is IL-1, IL-2, IL-6, IL-12, IL-17, IL-22, IL-23, GM-CSF, TNF- $\alpha$ , IFN- $\gamma$ , or any combination thereof.

93. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-2, IL-17, IFN- $\gamma$ , or any combination thereof.
94. The method of claim 93, wherein the cytokine is IL-2.
95. The method of claim 93, wherein the cytokine is IL-17.
96. The method of claim 93, wherein the cytokine is IFN- $\gamma$ .
97. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-1.
98. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-6.
99. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-12.
100. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-17.
101. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-22.
102. The method of any one of claims 82, 89 or 91, wherein the cytokine is IL-23.
103. The method of any one of claims 82, 89 or 91, wherein the cytokine is GM-CSF.
104. The method of any one of claims 82, 89 or 91, wherein the cytokine is TNF $\alpha$ .

1/15





FIG. 1A

FIG. 1B

2/15




FIG. 2

3/15




FIG. 3

>LC\_PD1AB-6-IgG1  
 DIVMTQSPDSLAVSLGERATINCKSGQSVLYSSNQNK**N**FL**A**WYQQKPGQQPKL**I**Y**W**ASTRESGVPDRFSGSGTDFLTISLQAE  
 DVAVYYCH**Q**YL**Y****W****T**FGQGT**K**LEIKRTVAAPSVFIFPPSDEQLKSGTASW**C**LN**N**YPREAKVQWVKVDNALQSGNSQESVTEQD  
 SKDSTSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

>HC\_PD1AB-6-IgG1  
 EVQLVQSGAEVKKPGATV**K**ISCKAS**G****F****N****I****K****D****T****Y****M****H****W**/QQAPGKGLEWM**G****R****I****D****P****A****N****G****D****R****K****Y****D****P****K****F****Q****G****R****V**/TADTSTDAYMELS  
 SLRSED**T**AVYYCARS**SG****P****V****Y****Y****G****S****S****Y****M****D****Y****W****G****Q****G****T****T****V****V****S****A****S****T****K****G****P****S****V****F****PL****A****P****S****S****K****S****T****S****G****G****T****A****A****L****G****C****L****V****K****D****Y****F****P****E****P****V****T****V****W****N****S****G**  
 AL**T**SGV**H****T****F****P****A****V****L****Q****Q****S****G****L****Y****S****L****S****S****W****V****T****V****P****S****S****L****G****T****Q****T****Y****I****C****N****V****N****H****K****P****S****N****T****K****V****D****K****K****V****E****P****K****S****C****D****K****T****H****T****C****PP****C****P****A****E****L****L****G****G****P****S****V****F****L****F****P****K****P****K****D**  
 TLM**I****S****R****T****P****E****V****T****C****W****V****D****V****S****H****E****D****P****E****V****K****F****N****W****Y****V****D****G****E****V****H****N****A****K****T****K****P****R****E****E****Q****Y****N****S****T****Y****R****W****S****V****L****T****V****L****H****Q****D****W****L****N****G****Y****E****K****K****V****S****N****K****A****L****P****A****P****|**  
 EKTISAKGQPREPQV**T****L****P****P****S****R****D****E****L****T****K****N****Q****V****S****L****T****C****L****V****K****G****F****Y****P****S****D****I****A****E****W****E****S****N****G****Q****P****E****N****N****Y****K****T****T****P****P****V****L****D****S****D****G****S****F****F****L****Y****S****K****L****T****V****D****K****S****R****W****Q**  
 QGNVFSCSV**M****H****E****A****L****H****N****Y****T****Q****K****S****L****S****P****G**

>HC\_PD1AB-6-IgG1-K322A  
 EVQLVQSGAEVKKPGATV**K**ISCKAS**G****F****N****I****K****D****T****Y****M****H****W**/QQAPGKGLEWM**G****R****I****D****P****A****N****G****D****R****K****Y****D****P****K****F****Q****G****R****V**/TADTSTDAYMELS  
 SLRSED**T**AVYYCARS**SG****P****V****Y****Y****G****S****S****Y****M****D****Y****W****G****Q****G****T****T****V****V****S****A****S****T****K****G****P****S****V****F****PL****A****P****S****S****K****S****T****S****G****G****T****A****A****L****G****C****L****V****K****D****Y****F****P****E****P****V****T****V****W****N****S****G**  
 AL**T**SGV**H****T****F****P****A****V****L****Q****Q****S****G****L****Y****S****L****S****S****W****V****T****V****P****S****S****L****G****T****Q****T****Y****I****C****N****V****N****H****K****P****S****N****T****K****V****D****K****K****V****E****P****K****S****C****D****K****T****H****T****C****PP****C****P****A****E****L****L****G****G****P****S****V****F****L****F****P****K****P****K****D**  
 TLM**I****S****R****T****P****E****V****T****C****W****V****D****V****S****H****E****D****P****E****V****K****F****N****W****Y****V****D****G****E****V****H****N****A****K****T****K****P****R****E****E****Q****Y****N****S****T****Y****R****W****S****V****L****T****V****L****H****Q****D****W****L****N****G****Y****E****K****K****V****S****N****K****A****L****P****A****P****|**  
 EKTISAKGQPREPQV**T****L****P****P****S****R****D****E****L****T****K****N****Q****V****S****L****T****C****L****V****K****G****F****Y****P****S****D****I****A****E****W****E****S****N****G****Q****P****E****N****N****Y****K****T****T****P****P****V****L****D****S****D****G****S****F****F****L****Y****S****K****L****T****V****D****K****S****R****W****Q**  
 QGNVFSCSV**M****H****E****A****L****H****N****Y****T****Q****K****S****L****S****P****G**

>HC\_PD1AB-6-IgG4P  
 EVQLVQSGAEVKKPGATV**K**ISCKAS**G****F****N****I****K****D****T****Y****M****H****W**/QQAPGKGLEWM**G****R****I****D****P****A****N****G****D****R****K****Y****D****P****K****F****Q****G****R****V**/TADTSTDAYMELS  
 SLRSED**T**AVYYCARS**SG****P****V****Y****Y****G****S****S****Y****M****D****Y****W****G****Q****G****T****T****V****V****S****A****S****T****K****G****P****S****V****F****PL****A****P****S****S****K****S****T****S****G****G****T****A****A****L****G****C****L****V****K****D****Y****F****P****E****P****V****T****V****W****N****S****G**  
 AL**T**SGV**H****T****F****P****A****V****L****Q****Q****S****G****L****Y****S****L****S****S****W****V****T****V****P****S****S****L****G****T****Q****T****Y****I****C****N****V****N****H****K****P****S****N****T****K****V****D****K****K****V****E****P****K****S****C****D****K****T****H****T****C****PP****C****P****A****E****L****L****G****G****P****S****V****F****L****F****P****K****P****K****D**  
 C**V****H****T****F****P****A****V****L****Q****Q****S****G****L****Y****S****L****S****S****W****V****T****V****P****S****S****L****G****T****Q****T****Y****T****C****N****V****D****K****R****V****E****S****K****Y****G****P****C****P****A****E****F****L****G****G****P****S****V****F****L****F****P****K****D****T****L****M****S****R****T****P****E****V****T**  
 C**V****W****D****V****S****Q****E****D****P****E****V****Q****F****N****W****Y****V****D****G****E****V****H****N****A****K****T****K****P****R****E****E****Q****F****N****S****T****Y****R****W****S****V****L****T****V****L****H****Q****D****W****L****N****G****Y****E****K****K****V****S****N****K****A****L****P****A****P****|**  
 T**L****P****P****S****Q****E****E****M****T****K****N****Q****V****S****L****T****C****L****V****K****G****F****Y****P****S****D****I****A****E****W****E****S****N****G****Q****P****E****N****N****Y****K****T****T****P****P****V****L****D****S****D****G****S****F****F****L****Y****S****R****L****T****D****K****S****R****W****Q****E****G****N****V****F****S****C****S****M****H****E****A****L****H****N****Y****T****Q****K****S****L****S****P****G**

FIG. 4





FIG. 5A

FIG. 5B

6/15




FIG. 6

7/15





FIG. 7

8/15



	EC <sub>50</sub> nM	F <sub>cγ</sub> RI	F <sub>cγ</sub> RIIa (V158)	F <sub>cγ</sub> RIIb
Cetuximab	33.1	915.7	9749.7	
PD1AB-6- gG1	54.7	293.9	2804.5	
PD1AB-6-K3	<b>56.0</b>	<b>622.6</b>	<b>1652.9</b>	
PD1AB-6-4P	868.9	6325.8	7179.1	

FIG. 8D

9/15




FIG. 9B

FIG. 9A

EC <sub>50</sub> nM	PD1AB-6-IgG1	PD1AB-6-K3	PD1AB-6-4P
FcgR1	2.6	1.5	12.4
FcgRIII	31.4	>666	NB

FIG. 9C

10/15




FIG. 10B




FIG. 10A

11/15




FIG. 11

12/15




FIG. 12

13/15




FIG. 13

14/15





FIG. 14A



Gated on CD3+ T cells

FIG. 14B




FIG. 14C

15/15






FIG. 15C

FIG. 15B

FIG. 15A

WO 2017/058859

## INTERNATIONAL SEARCH REPORT

PCT/US2016/054089

International application No.

PCT/US 16/54089

## Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
  - a.  forming part of the international application as filed:
    - in the form of an Annex C/ST.25 text file.
    - on paper or in the form of an image file.
  - b.  furnished together with the international application under PCT Rule 13*ter*.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
  - c.  furnished subsequent to the international filing date for the purposes of international search only:
    - in the form of an Annex C/ST.25 text file (Rule 13*ter*.1(a)).
    - on paper or in the form of an image file (Rule 13*ter*.1(b) and Administrative Instructions, Section 713).
2.  In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

WO 2017/058859

## INTERNATIONAL SEARCH REPORT

PCT/US2016/054089

International application No.

PCT/US 16/54089

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
  
  
  
  
  
  
  
  
  
2.  Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
  
  
  
  
  
  
  
  
3.  Claims Nos.: 29-104 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

-----please see extra sheet-----

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
  
  
  
  
  
  
  
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-4, 6, 11, 18, (21, 26-28)(in part), limited to antibody PD1AB-6-IgG1 (SEQ ID NOs: 1-6, 8, 13, 14-17, 20-22, 24, 36, 41)

## Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/54089

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 39/395, C07K 16/18, C07K 16/28 (2017.01)

CPC - A61K 39/3955, C07K 2317/73, C07K 2317/76

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A61K 39/395, C07K 16/18, C07K 16/28 (2017.01)

CPC - A61K 39/3955, C07K 2317/73, C07K 2317/76

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

CPC - A61K 2039/505, A61K 39/395, C07K 2316/96, C07K 2317/24

(keyword limited; terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatBase, Google Patents, Google Scholar

Search terms: antibody, PD-1, PD1, programmed cell death 1, CD279, IgG1, IgG4, CDR, CDR1, CDR2, CDR3, FR, humanized

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2011/0229461 A1 (TYSON) 22 September 2011 (22.09.2011) para [0043], [0095], [0096]	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)
A	US 2012/0100139 A1 (THOMPSON et al.) 26 April 2012 (26.04.2012) SEQ ID NO: 225	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)
A	US 2014/0212427 A1 (SONG) 31 July 2014 (31.07.2014) SEQ ID N: 47	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)
A	US 7,612,183 B2 (ELLIS et al.) 3 November 2009 (03.11.2009) SEQ ID NO: 16	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)
A	US 2014/0161794 A1 (LUGOVSKOY et al.) 12 June 2014 (12.06.2014) SEQ ID NO: 4	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)
A	US 2006/0029600 A1 (RUBIN et al.) 9 February 2006 (09.02.2006) SEQ ID NO: 2	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)



Further documents are listed in the continuation of Box C.



* Special categories of cited documents:	
“A” document defining the general state of the art which is not considered to be of particular relevance	“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
“E” earlier application or patent but published on or after the international filing date	“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
“O” document referring to an oral disclosure, use, exhibition or other means	
“P” document published prior to the international filing date but later than the priority date claimed	“&” document member of the same patent family

Date of the actual completion of the international search

8 February 2017

Date of mailing of the international search report

27 FEB 2017

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents  
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US 16/54089

**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2006/0110394 A1 (BENDIG et al.) 25 May 2006 (25.05.2006) SEQ ID NO: 11	1-4, 6, 11, 18, 21/(1-4,6,11,18), (26-28)/(1-4,6,11,18)

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US 16/54089

Continuation of Box No. III Observations where unity of invention is lacking

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

**Group I+:** Claims 1-28, drawn to an antibody or antigen-binding fragment thereof that binds to PD-1. The anti-PD-1 antibody or fragment thereof will be searched to the extent that the antibody or fragment thereof encompasses the variable domains of PD1AB-6 and the constant domains of human IgG1 (i.e. antibody PD1AB-6-IgG1), which comprises:

- VL domain of SEQ ID NO: 8 (claims 1, 6, 18)
- VH domain of SEQ ID NO: 13 (claims 1, 11, 18)
- VL CDR1-3 as set forth in Table 1, SEQ ID NOs: 1-3 (claims 2, 4)
- VH CDR1-3 as set forth in Table 2, SEQ ID NOs: 4-6 (claims 2, 4)
- VL FR1-4 as set forth in Table 3, SEQ ID NOs: 14-17 (claim 3)
- VH FR1-4 as set forth in Table 4, SEQ ID NOs: 20-22, 24 (claim 3)
- a human heavy chain IgG1 Fc region of SEQ ID NO: 36 (claims 21, 26, 28)
- a light chain constant region of SEQ ID NO: 41 (claims 27-28)

It is believed that claims 1-4, 6, 11, 18, (21, 26-28)(in part), limited to antibody PD1AB-6-IgG1 (SEQ ID NOs: 1-6, 8, 13, 14-17, 20-22, 24, 36, 41), encompass this first named invention, and thus these claims will be searched without fee to the extent that they encompass an anti-PD-1 antibody comprising antibody variable domains of PD1AB-6 and antibody constant domains of human IgG1 (i.e. antibody PD1AB-6-IgG1). [Note: Claim 22 is not included in the first named invention because it requires mutant IgG1-K322A Fc sequence SEQ ID NO: 37 instead of the non-mutant IgG1 Fc sequence SEQ ID NO: 36]. Additional anti-PD-1 antibodies will be searched upon the payment of additional fees. Applicants must specify the claims that encompass any additionally elected anti-PD-1 antibodies or fragments thereof. Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched. An exemplary election would be an anti-PD-1 antibody that encompasses the variable domains of PD1AB-1 and the constant domains of human IgG1-K322A Fc (i.e. antibody PD1AB-1-IgG1-K322A Fc), which comprises:

- VL domain of SEQ ID NO: 8 (claims 6, 12)
- VH domain of SEQ ID NO: 11 (claims 9, 12)
- VL CDR1-3 as set forth in Table 1, SEQ ID NOs: 1-3 (claims 2, 4)
- VH CDR1-3 as set forth in Table 2, SEQ ID NOs: 4-6 (claims 2, 4)
- VL FR1-4 as set forth in Table 3, SEQ ID NOs: 14-17 (claim 3)
- VH FR1-4 as set forth in Table 4, SEQ ID NOs: 19-22 (claim 3)
- a human heavy chain IgG1-K322A Fc region of SEQ ID NO: 37 (claims 21-22, 26, 28)
- a light chain constant region of SEQ ID NO: 41 (claims 27-28)

It is believed that claims 2-4, 6, 9, 12, (21-22, 26-28)(in part), limited to antibody PD1AB-1-IgG1-K322A Fc (SEQ ID NOs: 1-6, 8, 11, 14-17, 19-22, 37, 41), encompass this exemplary invention.

The inventions listed as Groups I+ do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The technical feature of each of the inventions listed as Group I+ is the specific anti-PD-1 antibody recited therein. Each invention requires an anti-PD-1 antibody comprising a different combination of sequences of light chain variable (VL) domains, heavy chain variable (VH) domains, heavy chain constant domains, and light chain constant domains not required by any of the other inventions.

The feature shared by Groups I+ is an antibody or antigen-binding fragment thereof that binds to PD-1 comprising a light chain variable region and a heavy chain variable region.

However, this shared technical feature does not represent a contribution over prior art, because the shared technical feature is taught by US 2011/0229461 A1 to (Tyson).

Tyson discloses an agonistic humanized antibody that binds to PD-1 (para [0043] "The present invention provides agonistic humanised antibodies having specificity for human PD-1") comprising a light chain variable region and a heavy chain variable region (para [0095] "the antibody according to the present disclosure is provided as PD-1 binding antibody fusion protein"; para [0096] "In one embodiment the fusion protein comprises two domain antibodies, for example as a variable heavy (VH) and variable light (VL) pairing"). Tyson further teaches that the antibody can further comprise human constant region domains of IgG1 or IgG4 (para [0098] "In particular, human IgG constant region domains may be used, especially of the IgG1 and IgG3 isotypes when the antibody molecule is intended for therapeutic uses and antibody effector functions are required. Alternatively, IgG2 and IgG4 isotypes may be used when the antibody molecule is intended for therapeutic purposes and antibody effector functions are not required, e.g. for simply agonising PD-1 activity"). As the technical feature was known in the art at the time of the invention, it cannot be considered a special technical feature that would otherwise unify the groups.

The inventions listed as Groups I+ therefore lack unity of invention under PCT Rule 13 because they do not share a same or corresponding special technical feature.

## 摘要

本文提供涉及特異性結合程式化死亡-1 (PD-1) 以及調節PD-1的表達和/或活性的抗體的組合物、方法和用途。