

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199917081 B2**
(10) Patent No. **772472**

(54) Title
Aza-heterocyclic compounds used to treat neurological disorders
and hair loss

(51)⁶ International Patent Classification(s)
C07D 207/16 C07D 207/12
A61K 031/40 C07D 403/04
A61K 031/41

(21) Application No: 199917081 (22) Application Date: 1998.12.03

(87) WIPO No: W099/62881

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/087788	1998.06.03	US
60/101077	1998.09.18	US

(43) Publication Date : 1999.12.20
(43) Publication Journal Date : 2000.03.02
(44) Accepted Journal Date : 2004.04.29

(71) Applicant(s)
GPI NIL Holdings, Inc.; Amgen, Inc.

(72) Inventor(s)
Gregory S. Hamilton; Mark H. Norman; Yong-Qian
Wu; Jia-He Li

(74) Agent/Attorney
Callinan Lawrie, Private Bag 7, KEW VIC 3101


17081/99



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 207/16, A61K 31/40, C07D 207/12, 403/04, A61K 31/41		(11) International Publication Number: WO 99/62881
A1		(43) International Publication Date: 9 December 1999 (09.12.99)
(21) International Application Number: PCT/US98/25573		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 3 December 1998 (03.12.98)		
(30) Priority Data: 60/087,788 3 June 1998 (03.06.98) US 60/101,077 18 September 1998 (18.09.98) US		
(71) Applicants: <i>Capitol Holdings, Inc. 222 Delaware Drive</i> GUILFORD PHARMACEUTICALS, INC. [US/US]; 6611 Factory Street, Baltimore, MD 21224 (US) AMGEN, INC. [US/US]; One Amgen Center Drive, Patent Operations M/S 27-4-A, Thousand Oaks, CA 91320-1789 (US).		
(72) Inventors: HAMILTON, Gregory, S.; 6501 Frederick Road, Catonsville, MD 21228 (US). NORMAN, Mark, H.; 3038 Shadow Hill Circle, Thousand Oaks, CA 91360 (US). WU, Yong-Qian; 6510 Barley Corn Row, Columbia, MD 21044 (US). <i>STEINER, Joseph, P. 980 Sugar Maple Street</i> <i>Hampstead, MD 21074 (US). Lijia He 27 Warren</i> <i>Manor Court Cockeysville, MD 21031 USA</i>		Published With international search report.
(74) Agent: NATH, Gary, M.; Nath & Associates, 6th floor, 1030 15th Street, N.W., Washington, DC 20005 (US).		
(54) Title: AZA-HETEROCYCLIC COMPOUNDS USED TO TREAT NEUROLOGICAL DISORDERS AND HAIR LOSS		
 (I)		
(57) Abstract		
This invention relates to novel N-heterocyclic carboxylic acids and carboxylic acid isosteres represented by formula (I), their preparation and use for treating neurological disorders including physically damaged nerves and neurodegenerative diseases, and for treating alopecia and promoting hair growth. In said formula, n is 1-3; X is either O or S; R₁ is selected from the group consisting of C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbonyl, or heterocycle; D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl; R is as defined in the application.		

**AZA-HETEROCYCLIC COMPOUNDS USED TO TREAT NEUROLOGICAL
DISORDERS AND HAIR LOSS****Related Application Data**

5 This application is a continuation-in-part of U.S.
patent application serial number 60/087,788 to Hamilton et
al., entitled "Carboxylic Acids and Carboxylic Acid
Isosteres of N-Heterocyclic Compounds", filed June 3,
1998, of U.S. patent application serial number 60/101,077
10 to Hamilton et al., entitled "Carboxylic Acids and
Carboxylic Acid Isosteres of N-Heterocyclic Compounds",
filed September 18, 1998.

BACKGROUND OF THE INVENTION

15

1. Field of the Invention

This invention relates to novel carboxylic acid and
carboxylic acid isosteres of N-heterocyclic compounds,
their preparation and use for treating neurological
20 disorders including physically damaged nerves and
neurodegenerative diseases, and for treating alopecia and
promoting hair growth.

2. Description of the Prior Art

25 It has been found that picomolar concentrations of an
immunosuppressant such as FK506 and rapamycin stimulate
neurite out growth in PC12 cells and sensory nervous,
namely dorsal root ganglion cells (DRGs). Lyons et al.,
Proc. of Natl. Acad. Sci., 1994 vol. 91, pp. 3191-3195.
30 In whole animal experiments, FK506 has been shown to
stimulate nerve regeneration following facial nerve injury
and results in functional recovery in animals with sciatic
nerve lesions.

Several neurotrophic factors effecting specific

neuronal populations in the central nervous system have been identified. For example, it has been hypothesized that Alzheimer's disease results from a decrease or loss of nerve growth factor (NGF). It has thus been proposed to treat Alzheimer's patients with exogenous nerve growth factor or other neurotrophic proteins such as brain derived nerve factor (BDNF), glial derived nerve factor, ciliary neurotrophic factor, and neurotrophin-3 to increase the survival of degenerating neuronal populations.

Clinical application of these proteins in various neurological disease states is hampered by difficulties in the delivery and bioavailability of large proteins to nervous system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display excellent bioavailability and specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, J. Am. Soc. Nephrol. 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, N. Engl. J. Med. 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 N. Engl. J. Med. 321: 1725).

Accordingly, there is a need for small-molecule compounds which are useful for neurotrophic effects and for treating neurodegenerative disorders.

Hair loss occurs in a variety of situations. These situations include male pattern alopecia, alopecia senilis, alopecia areata, diseases accompanied by basic skin lesions or tumors, and systematic disorders such as nutritional disorders and internal secretion disorders. The mechanisms causing hair loss are very complicated, but in some instances can be attributed to aging, genetic

disposition, the activation of male hormones, the loss of blood supply to hair follicles, and scalp abnormalities.

The immunosuppressant drugs FK506, rapamycin and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against graft rejection after organ transplantation. It has been reported that topical, but not oral, application of FK506 (Yamamoto et al., J. Invest. Dermatol., 1994, 102, 160-164; Jiang et al., J. Invest. Dermatol. 1995, 104, 523-525) and cyclosporin (Iwabuchi et al., J. Dermatol. Sci. 1995, 9, 64-69) stimulates hair growth in a dose-dependent manner. One form of hair loss, alopecia areata, is known to be associated with autoimmune activities; hence, topically administered immunomodulatory compounds are expected to demonstrate efficacy for treating that type of hair loss. The hair growth stimulating effects of FK506 have been the subject of an international patent filing covering FK506 and structures related thereto for hair growth stimulation (Honbo et al., EP 0 423 714 A2). Honbo et al. discloses the use of relatively large tricyclic compounds, known for their immunosuppressive effects, as hair revitalizing agents.

The hair growth and revitalization effects of FK506 and related agents are disclosed in many U.S. patents (Goulet et al., U.S. Patent No. 5,258,389; Luly et al., U.S. Patent No. 5,457,111; Goulet et al., U.S. Patent No. 5,532,248; Goulet et al., U.S. Patent No. 5,189,042; and Ok et al., U.S. Patent No. 5,208,241; Rupprecht et al., U.S. Patent No. 5,284,840; Organ et al., U.S. Patent No. 5,284,877). These patents claim FK506 related compounds. Although they do not claim methods of hair revitalization, they disclose the known use of FK506 for affecting hair growth. Similar to FK506 (and the claimed variations in the Honbo et al. patent), the compounds claimed in these patents are relatively large. Further, the cited patents

relate to immunomodulatory compounds for use in autoimmune related diseases, for which FK506's efficacy is well known.


Other U.S. patents disclose the use of cyclosporin and related compounds for hair revitalization (Hauer et al., U.S. Patent No. 5,342,625; Eberle, U.S. Patent No. 5,284,826; Hewitt et al., U.S. Patent No. 4,996,193). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin and related immunosuppressive compounds for hair growth.

However, immunosuppressive compounds by definition suppress the immune system and also exhibit other toxic side effects. Accordingly, there is a need for small molecule compounds which are useful as hair revitalizing compounds.

SUMMARY OF THE INVENTION

The present invention relates to the surprising discovery that N-heterocyclic compounds containing a carboxylic acid or carboxylic acid isostere moiety may be useful for treating neurodegenerative disorders and for treating alopecia and related hair loss disorders. Accordingly, a novel class of compounds containing an acidic moiety or an isostere thereof attached to the 2-carbon of the N-heterocyclic ring is provided. These compounds stimulate neuronal regeneration and outgrowth and as such are useful for treating neurological disorders and neurodegenerative diseases. These compounds also promote hair growth and as such are useful for treating hair loss disorders. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity and/or are non-immunosuppressive.

A preferred embodiment of this invention is a compound of formula (I):



where


n is 1;

X is either O or S;

5 R₁ is selected from the group consisting of C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is selected from



wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and
5 -CONR₃CN;

and

wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or R₂ is optionally substituted with one or more substituents selected from R₃ and Z, where

R₃ and Z are independently hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy,

10 alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, aralkyl, heteroaryl, carbocycle, heterocycle, and CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;

15 or a pharmaceutically acceptable salt, ester, or solvate thereof;

provided that:

when D is a bond, and R₂ is COOH,

then R₁ is not C₁-C₉ straight or branched chain alkyl, C₂-C₉, straight or branched chain alkenyl, C₅-C₇ cycloalkyl, C₅-C₇, cycloalkenyl, 2-(3,4-dichlorophenyl)ethyl, benzyl, or

20 Ar₁, where Ar₁ is 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thiazolyl, 2-thienyl, 3-thienyl, 1-pyridyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, and wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar₁, are optionally substituted with one or more substituents selected from the group consisting of hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C₁-C₉ straight or branched alkyl, C₂-C₉ straight or branched alkenyl, C₁-C₄
25 alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, COOH, and amino;

further provided that:


when D is a bond, and R₂ is -CONZ(R₃), and Z is hydrogen or C₁-C₆ alkyl, and R₃ is

phenyl, or C₂-C₆ straight or branched chain alkyl or alkenyl, wherein said alkyl is

unsubstituted or substituted in one or more positions with Ar₂ as defined below, C₃-C₈

30 cycloalkyl, cycloalkyl connected by methyl or a C₂-C₆ straight or branched chain alkyl or alkenyl chain, C₁-C₄, alkyl ester, or Ar₃ where Ar₃ is selected from the group consisting of 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thiazolyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or


branched alkyl, C₂-C₆ straight or branched alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, and amino; wherein said alkyl ester is optionally substituted with phenyl; or R₃ is the fragment:




- 5 where R₄ is selected from the group consisting of straight or branched chain C₁-C₈ alkyl optionally substituted with C₃-C₉ cycloalkyl, benzyl, or Ar₂ as defined below, and where R₅ is selected from phenyl, benzyl, C₁-C₆ straight or branched chain alkyl, and C₂-C₆ straight chain or branched alkenyl or where said alkyl or alkenyl is optionally substituted with phenyl or where R₂ is COOZ or CONR₆, where R₆ is selected from hydrogen, C₁-C₆
- 10 straight or branched alkyl, and C₂-C₆ straight or branched alkenyl, where said alkyl or alkenyl is optionally substituted with phenyl;
- then R₁, is not C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl or substituted thiophene, wherein said alkyl or alkenyl is optionally substituted in one or more positions with
- 15 -C₁-C₄ alkyl,
- C₁-C₄ alkenyl,
- hydroxy,
- C₃-C₈ cycloalkyl,
- C₅-C₇ cycloalkenyl, or
- 20 -Ar₂, where Ar₂ is 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents selected from hydrogen, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched alkyl, C₂-C₆ straight or branched alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, and amino,
- 25 where said cycloalkyl or cycloalkenyl groups may be optionally substituted with C₁-C₄ alkyl, C₁-C₄, alkenyl, or hydroxy,
- further provided that:
- when D is CH₂, then R₂ is not -OMe, -NHMe, or substituted -NHcyclohexyl;
- further provided that:
- 30 when D is CH₂, and R₂ is -OH,

then R_1 is not phenyl or pyrrolidinemethanol.

Especially preferred embodiments of this invention are where R_2 is selected from the group below:





where the atoms of each ring structure may be optionally substituted at one or more positions with R_3 .

Another preferred embodiment of this invention is where R_2 is selected from the group consisting of $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}_3$, $-\text{PO}_2(\text{R}_3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}_3)_2$, $-\text{OR}_3$, $-\text{SR}_3$, $-\text{NHCOR}_3$, $-\text{N}(\text{R}_3)_2$, $-\text{CON}(\text{R}_3)_2$, $-\text{CONH}(\text{O})\text{R}_3$, $-\text{CONHNHSO}_2\text{R}_3$, $-\text{COHNSO}_2\text{R}_3$, and $-\text{CONR}_3\text{CN}$.

Preferred embodiments of this invention are: (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-hydroxymethyl pyrrolidine; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinetetrazole; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile and (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-aminocarbonyl piperidine.

Another preferred embodiment of this invention is a pharmaceutical composition containing: an effective amount of a compound of formula (I); and a pharmaceutically suitable or acceptable carrier. For neurotrophic compositions a neurotrophic factor different from formula (I) may also be administered or otherwise included in the composition.

Another preferred embodiment of the invention is a method of promoting neuronal regeneration and growth in mammals, comprising administering to a mammal an effective amount of a compound of the invention.

Another preferred embodiment of the invention is a method of treating a neurological disorder in an animal, comprising administering to an animal an effective amount of a compound of the invention to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration.

Yet another preferred embodiment of the invention is a method of preventing neurodegeneration in an animal, comprising administering to an animal an effective amount of a compound of the invention.

Yet another preferred embodiment of the invention is a method of treating alopecia or promoting hair growth in an animal, comprising administering to an animal an effective amount of a compound of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a photograph of C57 Black 6 mice before being shaved for the hair regeneration experiment.

Figure 2 is a photograph of mice treated with a vehicle after six weeks. Figure 2 shows that less than 3% of the shaved area is covered with new hair growth when the vehicle (control) is administered.

Figure 3 is a bar graph illustrating relative hair growth on shaved mice treated with N-heterocyclic carboxylic acids or carboxylic acid isosteres at 1μmole per milliliter three times per week. Hair growth was evaluated after 14 days of treatment.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Alkyl" means a branched or unbranched saturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₁-C₆ straight or branched alkyl hydrocarbon chain contains 1 to 6 carbon atoms, and includes but is not limited to substituents such as methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, and the like. It is also contemplated as within the scope of the present invention that "alkyl" may also refer to a hydrocarbon chain wherein any of the carbon atoms of said alkyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of n-pentyl can be replaced with O to form propyloxymethyl.

"Alkenyl" means a branched or unbranched unsaturated

hydrocarbon chain comprising a designated number of carbon atoms. For example, C₂-C₆ straight or branched alkenyl hydrocarbon chain contains 2 to 6 carbon atoms having at least one double bond, and includes but is not limited to substituents such as ethenyl, propenyl, iso-propenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl, n-hexenyl, and the like. It is also contemplated as within the scope of the present invention that "alkenyl" may also refer to an unsaturated hydrocarbon chain wherein any of the carbon atoms of said alkenyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of 4-pentene can be replaced with O to form (2-propene)oxymethyl.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms.


Specifically, the term "carbocycle" or refers to an organic cyclic moiety in which the cyclic skeleton is comprised of only carbon atoms whereas the term "heterocycle" refers to an organic cyclic moiety in which the cyclic skeleton contains one or more heteroatoms selected from nitrogen, oxygen, or sulfur and which may or may not include carbon atoms.

Thus, the term "carbocycle" refers to a carbocyclic moiety containing the indicated number of carbon atoms. The term "C₃-C₈ cycloalkyl", therefore, refers to an organic cyclic substituent in which three to eight carbon atoms form a three, four, five, six, seven, or eight-membered ring, including, for example, a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, or cyclooctyl ring. As used herein, "carbocycle" may also refer to two or more cyclic ring systems which are fused to form, for example bicyclic, tricyclic, or other similar bridged substituents (e.g. adamantyl).

"Aryl" refers to an aromatic carbocyclic group having

a single ring, for example a phenyl ring; multiple rings, for example biphenyl; or multiple condensed rings in which at least one ring is aromatic, for example naphthyl, 1,2,3,4-tetrahydronaphthyl, anthryl, or phenanthryl, which
5 can be unsubstituted or substituted with one or more other substituents as defined above. The substituents attached to a phenyl ring portion of an aryl moiety in the compounds of Formula (I) may be configured in the ortho-, meta-, or para- orientations.

10 Examples of typical aryl moieties included in the scope of the present invention may include, but are not limited to, the following:




15 "Aralkyl" refers to alkyl or alkylene (alkenyl) chain which is substituted with aryl, heteroaryl, carbocycle or heterocycle, or alternatively one or more aryl, heteroaryl, carbocycle, or heterocycle(s) which is/are substituted with alkyl or alkenyl, i.e. 'Alkyl/alkylene
20 which is substituted with Ar' or 'Ar which is substituted with alkyl/alkylene'.

"Heterocycle" refers to a saturated, unsaturated, or aromatic carbocyclic group having a single ring, multiple rings, or multiple condensed rings, and having at least one hetero atom such as nitrogen, oxygen, or sulfur within at least one of the rings. "Heteroaryl" refers to a heterocycle in which at least one ring is aromatic. Any of the heterocyclic or heteroaryl groups can be unsubstituted or optionally substituted with one or more groups as defined above. Further, bi- or tri-cyclic heteroaryl moieties may comprise at least one ring which is either completely or partially saturated.

As one skilled in the art will appreciate, such heterocyclic moieties may exist in several isomeric forms, all of which are encompassed by the present invention. For example, a 1,3,5-triazine moiety is isomeric to a 1,2,4-triazine group. Such positional isomers are to be considered within the scope of the present invention. Likewise, the heterocyclic or heteroaryl groups can be bonded to other moieties in the compounds of the present invention. The point(s) of attachment to these other moieties is not to be construed as limiting on the scope of the invention. Thus, by way of example, a pyridyl moiety may be bound to other groups through the 2-, 3-, or 4-position of the pyridyl group. All such configurations are to be construed as within the scope of the present invention.

Examples of heterocyclic or heteroaryl moieties included in the scope of the present invention may include, but are not limited to, the following:



"Halo" means at least one fluoro, chloro, bromo, or iodo moiety.

The term "pharmaceutically acceptable salt, ester, or solvate" refers to salt, ester, or solvates of the subject compounds which possess the desired pharmacological activity and which are neither biologically nor otherwise undesirable. The salt, ester, or solvates can be formed with inorganic or organic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, sulfate, thiocyanate, tosylate and undecanoate. Base salt, ester, or solvates include ammonium salts, alkali metal salts such as lithium, sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salt with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as: 1) lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; 2) dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; 3) long chain alkyls such as decyl, lauryl, myristyl and stearyl substituted with one or more halide such as chloride, bromide and iodide; and 4) aralkyl halides like benzyl and phenethyl bromide and others.

The compounds of this invention may possess at least one asymmetric center and thus can be produced as mixtures of stereoisomers or as individual enantiomers or

diastereomers. The individual stereoisomers may be obtained by using an optically active starting material, by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolution of the compound of formula (I). It is understood that the individual stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by the scope of the present invention. The S-stereoisomer at atom 1 of formula I is a most preferred embodiment of the invention.

"Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Isomers" are different compounds that have the same molecular formula and includes cyclic isomers such as (iso)indole and other isomeric forms of cyclic moieties.


"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other.

"Diastereoisomers" are stereoisomers which are not mirror images of each other.

"Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

"Isosteres" are different compounds that have different molecular formulae but exhibit the same or similar properties. For example, tetrazole is an isostere of carboxylic acid because it mimics the properties of carboxylic acid even though they both have very different molecular formulae. Tetrazole is one of many possible isosteric replacements for carboxylic acid. Other carboxylic acid isosteres contemplated by the present invention include $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}^3$, $-\text{PO}_2(\text{R}^3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}^3)_2$, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{NHCOR}^3$, $-\text{N}(\text{R}^3)_2$, $-\text{CON}(\text{R}^3)_2$, $-\text{CONH}(\text{O})\text{R}^3$, $-\text{CONHNHSO}_2\text{R}^3$, $-\text{COHNSO}_2\text{R}^3$, and $-\text{CONR}^3\text{CN}$. In addition, carboxylic acid isosteres can include 5-7 membered

carbocycles or heterocycles containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally substituted in one or more positions. The following structures are non-limiting examples of preferred carbocyclic and heterocyclic isosteres contemplated by this invention.



where the atoms of said ring structure may be optionally substituted at one or more positions with R_3 . The present invention contemplates that when chemical substituents are added to a carboxylic isostere then the inventive compound
5 retains the properties of a carboxylic isostere.

The present invention contemplates that when a carboxylic isostere is optionally substituted with one or more moieties selected from R_3 , then the substitution cannot eliminate the carboxylic acid isosteric properties of the
10 inventive compound. The present invention contemplates that the placement of one or more R_3 substituents upon a carbocyclic or heterocyclic carboxylic acid isostere shall not be permitted at one or more atom(s) which maintain(s) or is/are integral to the carboxylic acid isosteric
15 properties of the inventive compound, if such substituent(s) would destroy the carboxylic acid isosteric properties of the inventive compound.

Other carboxylic acid isosteres not specifically exemplified or described in this specification are also
20 contemplated by the present invention.

It is understood that where chemical substitution is indicated then the chemical substituent chosen would form a sufficiently stable compound.

The term "preventing neurodegeneration" as used
25 herein includes the ability to inhibit or prevent neurodegeneration in patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for inhibiting or preventing further neurodegeneration in patients who are already
30 suffering from or have symptoms of a neurodegenerative disease when the compounds are given concurrently.

The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

35 (i) preventing a disease and/or condition from

occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

- (ii) inhibiting the disease and/or condition, i.e.,
5 arresting its development; or
(iii) relieving the disease and/or condition, i.e.,
causing regression of the disease and/or condition.

The system used in naming the compounds of the present invention is shown below, using a compound of
10 formula I as an example.

A compound of the present invention, especially formula I, wherein n is 1, X is O, D is a bond, R₁ is 1,1-dimethylpropyl, and R₂ is -CN, is named (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile.

15 "Alopecia" refers to deficient hair growth and partial or complete loss of hair, including without limitation androgenic alopecia (male pattern baldness), toxic alopecia, alopecia senilis, alopecia areata, alopecia pelada and trichotillomania. Alopecia results
20 when the pilar cycle is disturbed. The most frequent phenomenon is a shortening of the hair growth or anagen phase due to cessation of cell proliferation. This results in an early onset of the catagen phase, and consequently a large number of hairs in the telogen phase
25 during which the follicles are detached from the dermal papillae, and the hairs fall out. Alopecia has a number of etiologies, including genetic factors, aging, local and systemic diseases, febrile conditions, mental stresses, hormonal problems, and secondary effects of drugs.

30 "Pilar cycle" refers to the life cycle of hair follicles, and includes three phases:

- (1) the anagen phase, the period of active hair growth which, insofar as scalp hair is concerned, lasts about three to five years;
35 (2) the catagen phase, the period when growth stops.

and the follicle atrophies which, insofar as scalp hair is concerned, lasts about one to two weeks; and

- (3) the telogen phase, the rest period when hair progressively separates and finally falls out which, insofar as scalp hair is concerned, lasts about three to four months.

Normally 80 to 90 percent of the follicles are in the anagen phase, less than 1 percent being in the catagen phase, and the rest being in the telogen phase. In the telogen phase, hair is uniform in diameter with a slightly bulbous, non-pigmented root. By contrast, in the anagen phase, hair has a large colored bulb at its root.

"Promoting hair growth" refers to maintaining, inducing, stimulating, accelerating, or revitalizing the germination of hair.

"Treating alopecia" refers to:

- (i) preventing alopecia in an animal which may be predisposed to alopecia; and/or
- (ii) inhibiting, retarding or reducing alopecia; and/or
- (iii) promoting hair growth; and/or
- (iv) prolonging the anagen phase of the hair cycle; and/or
- (v) converting vellus hair to growth as terminal hair. Terminal hair is coarse, pigmented, long hair in which the bulb of the hair follicle is seated deep in the dermis. Vellus hair, on the other hand, is fine, thin, non-pigmented short hair in which the hair bulb is located superficially in the dermis. As alopecia progresses, the hairs change from the terminal to the vellus type.

The term "neurotrophic" as used herein includes without limitation the ability to stimulate neuronal regeneration or growth and/or the ability to prevent or treat neurodegeneration.

The term "non-immunosuppressive" refers to the inability of the compounds of the present invention to trigger an immune response when compared to a control such as FK506 or cyclosporin A. Assays for determining immunosuppression are well known to those of ordinary skill in the art. Specific non-limiting examples of well known assays include PMA and OKT3 assays wherein mitogens are used to stimulate proliferation of human peripheral blood lymphocytes (PBC). Compounds added to such assay systems are evaluated for their ability to inhibit such proliferation.

15

Compounds of the Invention

The present invention relates to the surprising discovery that carboxylic acid or carboxylic acid isostere compounds are neurotrophic and are able to treat alopecia. Accordingly, a novel class of compounds are provided. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

Preferred compounds of the present invention contain carboxylic acid moieties and other isosteric replacements for carboxylic acid moieties, of which several examples

are specified herein. Other isosteric replacements for carboxylic acid moieties, known to those skilled in the art of medicinal chemistry, are within the scope of the invention if not otherwise specified.

5 The compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathic and neurological
10 disorders relating to neurodegeneration. The compounds of this invention can also be administered to mammals other than humans for treatment of various mammalian neurological disorders.

 The novel compounds of the present invention possess
15 an excellent degree of neurotrophic activity. This activity is useful in the stimulation of damaged neurons, the promotion of neuronal regeneration, the prevention of neurodegeneration, and in the treatment of several neurological disorders known to be associated with
20 neuronal degeneration and peripheral neuropathies. The neurological disorders that may be treated include but are not limited to: trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular dystrophy, amyotrophic lateral sclerosis, progressive
25 muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured or prolapsed invertebrate disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, peripheral neuropathic such as those caused by lead, dapsone, ticks,
30 prophyria, or Gullain-Barré syndrome, Alzheimer's disease, and Parkinson's disease.

 The above discussion relating to the utility and administration of the compounds of the present invention also applies to the pharmaceutical compositions of the
35 present invention.

The term "pharmaceutically acceptable carrier" as used herein refers to any carrier, diluent, excipient, suspending agent, lubricating agent, adjuvant, vehicle, delivery system, emulsifier, disintegrant, absorbant, preservative, surfactant, colorant, flavorant, or sweetener.

For these purposes the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion techniques.

For oral administration, the compounds of the present invention may be provided in any suitable dosage form known in the art. For example, the compositions may be incorporated into tablets, powders, granules, beads, chewable lozenges, capsules, liquids, aqueous suspensions or solutions, or similar dosage forms, using conventional equipment and techniques known in the art. Tablet dosage forms are preferred. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient.

When preparing dosage form incorporating the compositions of the invention, the compounds may also be blended with conventional excipients such as binders, including gelatin, pregelatinized starch, and the like; lubricants, such as hydrogenated vegetable oil, stearic

acid, and the like; diluents, such as lactose, mannose, and sucrose; disintegrants, such as carboxymethylcellulose and sodium starch glycolate; suspending agents, such as povidone, polyvinyl alcohol, and the like; absorbents, such as silicon dioxide; preservatives, such as methylparaben, propylparaben, and sodium benzoate; surfactants, such as sodium lauryl sulfate, polysorbate 80, and the like; colorants such as F.D. & C. dyes and lakes; flavorants; and sweeteners.

10 Compositions and methods of the invention also may utilize controlled release technology. Thus, for example, the inventive compounds may be incorporated into a hydrophobic polymer matrix for controlled release over a period of days. Such controlled release films are well known to the art. Particularly preferred are transdermal delivery systems. Other examples of polymers commonly employed for this purpose that may be used in the present invention include nondegradable ethylene-vinyl acetate copolymer and degradable lactic acid-glycolic acid copolymers which may be used externally or internally. Certain hydrogels such as poly(hydroxyethylmethacrylate) or poly(vinylalcohol) also may be useful, but for shorter release cycles than the other polymer releases systems, such as those mentioned above.

25 To be effective therapeutically as central nervous system targets, the compounds of the present invention should readily penetrate the blood-brain barrier when peripherally administered. Compounds which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route or other appropriate delivery system suitable for administration to the brain.

35 The compounds of the present invention may be administered in the form of sterile injectable preparations, for example, as sterile injectable aqueous

or oleaginous suspensions. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be
5 sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.
10 In addition, sterile, fixed oils are conventionally employed as solvents or suspending mediums. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid and its glyceride derivatives, including olive
15 oil and castor oil, especially in their polyoxyethylated versions, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

The compounds of this invention may also be
20 administered rectally in the form of suppositories. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature, but liquid at rectal temperature and, therefore, will melt in the rectum to release the drug.
25 Such materials include cocoa butter, beeswax and polyethylene glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily
30 accessible by topical application, including neurological disorders of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For topical application to the eye, or ophthalmic
35 use, the compounds can be formulated as micronized

suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic
5 uses the compounds may be formulated in an ointment such as petrolatum.

For topical application to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a mixture
10 with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active
15 compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see
20 above) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred
25 levels of about 0.1 mg to about 1,000 mg. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Typically, *in vitro* dosage-effect results
30 provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful. The considerations for determining the proper dose levels are well known in the art.

It is understood, however, that a specific dose level
35 for any particular patient will depend upon a variety of

factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

To effectively treat alopecia or promote hair growth, the compounds used in the inventive methods and pharmaceutical compositions must readily affect the targeted areas. For these purposes, the compounds are preferably administered topically to the skin.


For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The compounds can be administered with other hair revitalizing agents. Specific dose levels for the other hair revitalizing agents will depend upon the factors previously stated and the effectiveness of the drug combination. Other routes of administration known in the pharmaceutical art are also contemplated by this invention.

Specific embodiments of the inventive compounds are presented in Tables I, II, and III. The present invention contemplates employing the compounds of Tables I, II and III, below, for use in compositions and methods to prevent and/or treat a neurological disorder in an animal, and for use in compositions and methods to treat alopecia and

promote hair growth in an animal, and all other uses suggested in this specification.


Table I



D is a bond and R₂ is COOH,

	No.	X	n	R ₁
10	1	O	1	3,4,5-trimethylphenyl
	2	O	2	3,4,5-trimethylphenyl
	3	O	1	tert-butyl
	4	O	3	tert-butyl
15	5	O	1	cyclopentyl
	6	O	2	cyclopentyl
	7	O	3	cyclopentyl
	8	O	1	cyclohexyl
	9	O	2	cyclohexyl
20	10	O	3	cyclohexyl
	11	O	1	cycloheptyl
	12	O	2	cycloheptyl
	13	O	3	cycloheptyl
	14	O	1	2-thienyl
25	15	O	2	2-thienyl
	16	O	3	2-thienyl
	17	O	1	2-furyl
	18	O	2	2-furyl
	19	O	3	2-furyl
30	20	O	3	phenyl
	21	O	1	1,1-dimethylpentyl
	22	O	2	1,1-dimethylhexyl
	23	O	3	ethyl

Table II.



5

10


15

20

25









No.	X	n	R ₁	D	R ₂
24	S	1	1,1-dimethyl propyl	CH ₂	COOH
25	S	1	1,1-dimethyl propyl	bond	COOH
26	O	1	1,1-dimethyl propyl	CH ₂	OH
27	O	1	1,1-dimethyl propyl	bond	SO ₃ H
28	O	1	1,1-dimethyl propyl	CH ₂	CN
29	O	1	1,1-dimethyl propyl	bond	CN
30	O	1	1,1-dimethyl propyl	bond	tetrazolyl
31	S	1	phenyl	(CH ₂) ₂	COOH
32	S	1	phenyl	(CH ₂) ₃	COOH
33	S	2	phenyl	CH ₂	COOH
34	O	1	1,1-dimethyl propyl	bond	CONH ₂
35	O	2	1,1-dimethyl propyl	bond	CONH ₂
36	S	2	2-furyl	bond	PO ₃ H ₂
37	O	2	propyl	(CH ₂) ₂	COOH
38	O	1	propyl	(CH ₂) ₃	COOH
39	O	1	tert-butyl	(CH ₂) ₄	COOH
40	O	1	methyl	(CH ₂) ₅	COOH
41	O	2	phenyl	(CH ₂) ₆	COOH
42	O	2	3,4,5- trimethoxy-phenyl	CH ₂	COOH
43	O	2	3,4,5- trimethoxy-phenyl	CH ₂	tetrazolyl

TABLE III











	No.	n	X	D	R ₂	R ₁
5	44	1	S	bond	COOH	Phenyl
	45	1	O	bond	COOH	α-MethylBenzyl
	46	2	O	bond	COOH	4-MethylBenzyl
	47	1	O	bond	Tetrazole	Benzyl
	48	1	O	bond	SO ₃ H	α-MethylBenzyl
10	49	1	O	CH ₂	COOH	4-MethylBenzyl
	50	1	O	bond	SO ₃ HMe	Benzyl
	51	1	O	bond	CN	α-MethylBenzyl
	52	1	O	bond	PO ₃ H ₂	4-MethylBenzyl
	53	2	O	bond	COOH	Benzyl
15	54	2	O	bond	COOH	α-MethylBenzyl
	55	2	O	bond	COOH	4-MethylBenzyl
	56	2	S	bond	COOH	3,4,5-trimethoxyphenyl
20	57	2	O	bond	COOH	Cyclohexyl
	58	2	O	bond	PO ₃ HEt	i-propyl
	59	2	O	bond	PO ₃ HPropyl	ethyl
	60	2	O	bond	PO ₃ (Et) ₂	Methyl
	61	2	O	bond	OMe	tert-butyl
25	62	1	O	bond	OEt	n-pentyl
	63	2	O	bond	OPropyl	n-hexyl
	64	1	O	bond	OButyl	Cyclohexyl
	65	1	O	bond	OPentyl	cyclopentyl
	66	1	O	bond	OHexyl	n-heptyl
30	67	1	O	bond	SMe	n-octyl
	68	1	O	bond	SEt	n-nonyl
	69	2	O	bond	SPropyl	2-indolyl
	70	2	O	bond	SButyl	2-furyl
	71	2	O	bond	NHCOMe	2-thiazolyl
35	72	2	O	bond	NHCOEt	2-thienyl
	73	1	O	CH ₂	N(Me) ₂	2-pyridyl
	74	1	O	(CH ₂) ₂	N(Me)Et	1,1-dimethylpropyl
	75	1	O	(CH ₂) ₃	CON(Me) ₂	1,1-dimethylpropyl
	76	1	O	(CH ₂) ₄	CONHMe	1,1-dimethylpropyl




	No.	n	X	D	R ₂	R ₁
5	77	1	O	(CH ₂) ₅	CONHEt	1,1-dimethylpropyl
	78	1	O	(CH ₂) ₆	CONHPropyl	1,1-dimethylpropyl
	79	1	O	bond	CONH(O)Me	Benzyl
	80	1	O	bond	CONH(O)Et	α-Methylphenyl
	81	1	O	bond	CONH(O)Propyl	4-Methylphenyl
10	82	1	O	(CH ₂) ₂	COOH	Benzyl
	83	1	O	bond	COOH	α-Methylphenyl
	84	1	O	bond	COOH	4-Methylphenyl
	85	1	O	CH ₂	COOH	1,1-dimethylpropyl
	86	1	O	(CH ₂) ₂	COOH	1,1-dimethylbutyl
15	87	1	O	(CH ₂) ₃	COOH	1,1-dimethylpentyl
	88	1	O	(CH ₂) ₄	COOH	1,1-dimethylhexyl
	89	1	O	(CH ₂) ₅	COOH	1,1-dimethylethyl
	90	1	O	(CH ₂) ₆	COOH	iso-propyl
	91	1	O	(CH ₂) ₇	COOH	tert-butyl
20	92	1	O	(CH ₂) ₈	COOH	1,1-dimethylpropyl
	93	1	O	(CH ₂) ₉	COOH	benzyl
25	94	1	O	(CH ₂) ₁₀	COOH	1,1-dimethylpropyl
	95	1	O	C ₂ H ₂	COOH	cyclohexylmethyl
30	96	1	O	2-OH, Et	COOH	1,1-dimethylpropyl
	97	1	O	2-butylene	COOH	1,1-dimethylpropyl
	98	1	S	i-Pro	COOH	1,1-dimethylpropyl
	99	2	S	tert-Bu	COOH	phenyl
	100	2	O	2-nitro-hexyl	COOH	1,1-dimethylpropyl
35	101	1	O	(CH ₂) ₂	CN	1,1-dimethylpropyl
	102	1	O	(CH ₂) ₃	CN	1,1-dimethylpropyl
	103	3	O	bond	CONHNHSO ₂ Me	Benzyl
	104	3	O	bond	CONHNHSO ₂ Et	α-Methylphenyl
	105	3	O	bond	CONHSO ₂ Me	4-Methylphenyl
40	106	1	O	bond	CONHNHSO ₂ Et	Phenyl
	107	2	O	bond	CON(Me)CN	α-Methylphenyl
	108	1	O	bond	CON(Et)CN	4-Methylphenyl
	109	1	O	(CH ₂) ₂	COOH	methyl
	110	1	O	(CH ₂) ₃	COOH	ethyl
45	111	1	O	(CH ₂) ₄	COOH	n-propyl
	112	1	O	(CH ₂) ₅	COOH	t-butyl
	113	1	O	(CH ₂) ₆	COOH	Pentyl
	114	1	O	(CH ₂) ₇	COOH	Hexyl
	115	1	O	(CH ₂) ₈	COOH	Septyl
	116	1	O	(CH ₂) ₉	COOH	Octyl
	117	1	O	C ₂ H ₂	COOH	Cyclohexyl

No.	n	X	D	R2	R1	
118	2	O	bond		1,1-dimethylpropyl	
5	119	1	O	bond		1,1-dimethylpropyl
120	1	O	bond		1,1-dimethylpropyl	
121	1	O	bond		1,1-dimethylpropyl	
10	122	1	O	bond		1,1-dimethylpropyl
123	1	O	bond		1,1-dimethylpropyl	
124	1	O	bond		1,1-dimethylpropyl	
15	125	1	O	bond		1,1-dimethylpropyl


20

33

No.	n	X	D	R2	R1
126	1	O	bond		1,1-dimethylpropyl
5 127	1	O	bond		1,1-dimethylpropyl
128	1	O	bond		1,1-dimethylpropyl
129	1	O	bond		1,1-dimethylpropyl
10 130	1	O	bond		1,1-dimethylpropyl
131	1	O	bond		1,1-dimethylpropyl
132	1	O	bond		1,1-dimethylpropyl
15 133	1	O	bond		1,1-dimethylpropyl

No.	n	X	D	R2	R1
134	1	O	bond		1,1-dimethylpropyl
135	1	O	bond		1,1-dimethylpropyl
136	1	O	bond		1,1-dimethylpropyl
137	1	O	bond	COOH	1,1-dimethylpropyl
138	2	O	bond	COOH	1,1-dimethylpropyl


Specific embodiments of the present invention also include compound 139:



Compound 139.

Additional claimed or comparative carboxylic acids and isosteres of N-heterocyclic compounds which also show the remarkable neurotrophic and hair growth effects of the present invention are shown below in Table IV:

TABLE IV



Cpd.	n	D	R ₂	L	R ₁
itr/number					
5 A/137	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
B/138	2	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
C	1	bond	COOH	SO ₂	Benzyl
D/26	1	CH ₂	OH	1,2-dioxoethyl	1,1-dimethylpropyl
E/30	1	bond	tetrazole	1,2-dioxoethyl	1,1-dimethylpropyl
10 F/29	1	bond	-CN	1,2-dioxoethyl	1,1-dimethylpropyl
G/35	2	bond	CONH ₂	1,2-dioxoethyl	1,1-dimethylpropyl
where Y and Z are both carbon for compounds A-G,					
H	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
15 I	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
where Z is S for compound H and					
where Y is S for compound I.					

Pharmaceutical Compositions of the Present Invention

20 The present invention relates to a pharmaceutical composition comprising:

- (i) an effective amount of an N-heterocyclic carboxylic acid or carboxylic acid isostere compound; and

25 (ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of an N-heterocyclic carboxylic acid or carboxylic acid isostere

compound for treating neurodegenerative diseases, neurological disorders, and nerve damage, or promoting nerve growth in an animal; and

5 (ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of an N-heterocyclic carboxylic acid or carboxylic acid isostere
10 compound for treating alopecia or promoting hair growth in an animal; and

(ii) a pharmaceutically acceptable carrier.

The compounds can be administered with other neurotrophic agents such as neurotrophic growth
15 factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3 and
20 neurotrophin 4/5. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

Methods of the Present Invention

25 The present invention relates to the use of any of the compounds seen in Tables I, II, III, IV, and other compounds embodied herein, in the preparation of a medicament for the treatment of a disease such as peripheral neuropathy caused by physical injury or disease
30 state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis. The present invention also relates to the use of carboxylic acid and carboxylic acid isostere
35 compounds for treating the above-mentioned neuropathies.

neurological disorders, and neurological damage.

The present invention also relates to a method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal an effective
5 amount of an N-heterocyclic carboxylic acid or carboxylic acid isostere. The present invention also relates to using the inventive compounds and compositions in the preparation of a medicament for the treatment of alopecia or promoting hair growth in an animal.

10 The inventive method is particularly useful for treating male pattern alopecia, alopecia senilis, alopecia areata, alopecia resulting from skin lesions or tumors, alopecia resulting from cancer therapy such as chemotherapy and radiation, and alopecia resulting from
15 systematic disorders such as nutritional disorders and internal secretion disorders.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound
20 employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease or disorder being treated and form of administration.

25 **MPTP Model of Parkinson's Disease in Mice**

MPTP lesioning of dopaminergic neurons in mice was used as an animal model of Parkinson's Disease. Four week old male CD1 white mice were dosed i.p. with 30 mg/kg of MPTP for 5 days. The inventive compounds (4 mg/kg), or
30 vehicle, were administered s.c. along with the MPTP for 5 days, as well as for an additional 5 days following cessation of MPTP treatment. At 18 days following MPTP treatment, the animals were sacrificed and the striata were dissected and homogenized. Immunostaining was
35 performed on sagittal and coronal brain sections using

anti-tyrosine hydroxylase Ig to quantitate survival and recovery of dopaminergic neurons. In animals treated with MPTP and vehicle, a substantial loss of functional dopaminergic terminals was observed as compared to non-lesioned animals. In another protocol, test compounds were administered only subsequent to MPTP-induced lesioning. Thus, after animals were treated with MPTP for 5 days, an additional 3 days passed before beginning oral drug treatment on day 8. Animals were treated with the inventive compounds (0.4 mg/kg), administered orally, once a day for 5 days total. On day 18, the animals were sacrificed and analyzed as described above.

Table V presents the percent recovery of dopaminergic neurons in the first (concurrent dosing) paradigm in animals receiving carboxylic acid or carboxylic acid isostere compounds of the present invention.

Table V, below, shows the remarkable neuroregenerative effects of the inventive carboxylic acid or carboxylic acid isostere related compounds illustrating the neurotrophic capability of carboxylic acid isosteres as a class showing that lesioned animals receiving the carboxylic acid or carboxylic acid isostere compounds provide a remarkable recovery of TH-stained dopaminergic neurons.

Table V - MPTP Neurodegenerative Model

		% Recovery
	Compound A	26.7 %
	Compound B	ND
	Compound C	24.4 %
30	Compound D	23.2 %
	Compound E	19.6 %
	Compound F	34.1 %
	Compound G	46.5 %
	Compound H	14.0 %
35	Compound I	ND

Percent striatal innervation density was quantitated in brain sections with an anti-tyrosine hydroxylase immunoglobulin, which is indicative of functional dopaminergic neurons. The striatal innervation density of 23% for animals pretreated with only a vehicle and administered a vehicle orally during treatment, is indicative of normal non-lesioned striatal tissue. Striatal innervation density is reduced to 5% for animals pretreated with MPTP and administered a vehicle orally during treatment, and is indicative of MPTP-induced lesioning. Surprisingly, striatal innervation density is increased 8-13% for animals pretreated with MPTP and administered 0.4 mg/kg of an inventive compound orally during treatment, indicating substantial neuronal regeneration after induction of MPTP-derived lesions.

The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto. All polymer molecular weights are mean average molecular weights. All percentages are based on the percent by weight of the final delivery system or formulation prepared unless otherwise indicated and all totals equal 100% by weight.


EXAMPLES

The inventive compounds may be prepared by a variety of synthetic sequences that utilize established chemical transformations. A general pathway to the present compounds is described in Scheme I. N-glyoxylproline derivatives may be prepared by reacting L-proline methyl ester with methyl oxalyl chloride as shown in Scheme I. The resulting oxamates may be reacted with a variety of carbon nucleophiles to obtain compounds used in the present invention.

35

40

Scheme I

**EXAMPLE 1 (Compound 137)**

Synthesis of (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

5

a. Synthesis of (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate.

A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methylene chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hr. After filtering to remove solids, the organic phase was washed with water, dried over MgSO_4 , and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to obtain 3.52 g (88%) of the product as a reddish oil.

Mixture of cis-trans amide rotamers; data for trans rotamer given. ^1H NMR (CDCl_3): δ 1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H); 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H, $J = 8.4, 3.3$).

15

20

25

b. Synthesis of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate.


A solution of methyl (2S)-1-(1,2-dioxo-2-

methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90 mmol) in 30 mL of tetrahydrofuran (THF) was cooled to -78°C and treated with 14.2 mL of a 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (100 mL) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil. ¹H NMR (CDCl₃): δ 0.88 (t, 3H); 1.22, 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H, J = 8.4, 3.4).

c. Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid (Compound 137).

A mixture of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 mL), and methanol (50 mL) was stirred at 0°C for 30 min and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 100 mL of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, J = 10.4, 7.3); 4.55 (dd, 1H, J = 8.6, 4.1).

Scheme II



Inventive compounds containing bridged rings may be synthesized using the above synthetic schemes by substituting the substrates containing the N-heterocyclic ring structures with comparable substrates containing bridged ring structures.

EXAMPLE 2

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxamide. (Compound 34)

Isobutyl chloroformate (20 mmol, 2.7 mL) was added to a solution containing (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid (4.89 g, 20 mmol) (from Example 1) in 50 mL methylene chloride at -10°C with stirring. After 5 minutes, ammonia was added dropwise (20 mmol, 10 mL of 2 M ethyl alcohol solution). The reaction was warmed up to room temperature after stirring at -10°C for 30 minutes. The mixture was diluted with water, and extracted into 200 mL methylene chloride. The organic extract was concentrated and further purified by silica gel to give 4.0 g of product as a white solid (81.8% yield). ¹H NMR (CDCl₃): δ 0.91 (t, 3H, J= 7.5); 1.28 (s, 6H, each); 1.63-1.84 (m, 2H); 1.95-2.22 (m, 3H); 2.46 (m, 1H); 3.55-3.67 (m, 2H); 4.67 (t, 1H, J= 7.8);

5.51-5.53 (br, 1H, NH); 6.80 (br, 1H, NH).

EXAMPLE 3

5 Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile. (Compound 29)

To a solution of 0.465 mL DMF (6 mmol) in 10 mL acetonitrile at 0°C was added 0.48 mL (5.5 mmol) of oxalyl chloride. A white precipitate formed immediately and was accompanied by gas evolution. When complete, a solution
10 of 1.2 g (5 mmol) of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxamide (from Example 2) in 2.5 mL acetonitrile was added. When the mixture became homogeneous, 0.9 mL (11 mmol) pyridine was added. After 5 min., the mixture was diluted into water and extracted by
15 200 mL ethyl acetate. The organic layer was concentrated and further purified by silica gel to give 0.8 g product as a white solid (72% yield). ¹H NMR (CDCl₃): δ 0.87 (t, 3H, J= 7.5); 1.22 (s, 3H); 1.24 (s, 3H); 1.80 (m, 2H); 2.03-2.23 (m, 4H); 3.55 (m, 2H); 4.73 (m, 1H).

EXAMPLE 4

20 Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinetetrazole. (Compound 30)

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile (222 mg, 1 mmol) (from Example 3),
25 NaN₃ (81 mg, 1.3 mmol) and NH₄Cl (70 mg, 1.3 mmol) in 3 mL DMF was stirred at 130°C for 16 hours. The mixture was concentrated and purified by silica gel to afford 200 mg product as white solid (75.5% yield). ¹H NMR (CDCl₃): δ
30 0.88 (t, 3H, J= 7.5); 1.22 (s, 6H); 1.68 (m, 2H); 2.05-2.36 (m, 3H); 2.85 (m, 1H); 3.54 (m, 1H); 3.75 (m, 1H); 5.40 (m, 1H).

Example 5

In Vivo Hair Generation Test With C57 Black 6 Mice

35 C57 black 6 mice were used to demonstrate the hair revitalizing properties of the N-heterocyclic carboxylic

acids or carboxylic acid isosteres. Referring now to FIGS. 1 and 2 of the drawings, C57 black 6 mice, approximately 7 weeks old, had an area of about 2 inches by 2 inches on their hindquarters shaved to remove all existing hair. Care was taken not to nick or cause abrasion to the underlying dermal layers. The animals were in anagen growth phase, as indicated by the pinkish color of the skin. Referring now to FIG. 2, four animals per group were treated by topical administration with 20% propylene glycol vehicle (FIG. 2), or neuroimmunophilin FKBP ligands dissolved in the vehicle. The animals were treated with vehicle or neuroimmunophilin ligands every 48 hours (3 applications total over the course of 5 days) and the hair growth was allowed to proceed for 6 weeks. Hair growth was quantitated by the percent of shaved area covered by new hair growth during this time period.

FIG. 2 shows that animals treated with vehicle exhibited only a small amount of hair growth in patches or tufts, with less than 3% of the shaved area covered with new growth.

In contrast, FIG. 3 shows that animals treated for 2 weeks with the N-heterocyclic carboxylic acid compounds i.e. compound A (137), compound B (138), and compound G (35) exhibited dramatic hair growth, covering greater than 25% of the shaved area in all animals for two of the compounds.

FIG. 3 shows the relative hair growth on shaven C57 black 6 mice 14 days after being treated with one of three N-heterocyclic carboxylic acids or carboxylic acid isosteres. The mice had a 2 x 2 inch region on their backside shaved to remove all hair. Care was taken not to nick or cause abrasion to the underlying dermal layers. Compounds at a concentration of 1 μ mole per milliliter were carefully applied to the shaved area of the mice (5 mice per group) three times per week. Hair growth was

evaluated 14 days after initiation of drug treatment. The relative scale for assessing hair growth is as follows:

- 0 = no growth;
1 = beginning of growth in small tufts;
5 2 = hair growth covering over <25% of shaved area;
3 = hair growth covering over >25% of shaved area, but less than 50% of shaved area;
4 = hair growth covering over >50% of shaved area, but less than 75% of shaved area;
10 5 = complete hair growth of shaved area.

Example 6

A lotion comprising the following composition may be prepared.

15		(%)
	95% Ethanol	80.0
	an N-heterocyclic carboxylic acid or carboxylic acid isostere	10.0
	α -Tocopherol acetate	0.01
20	Ethylene oxide (40 mole) adducts of hardened castor oil	0.5
	purified water	9.0
	perfume and dye	q.s.

- 25 Into 95% ethanol are added an N-heterocyclic carboxylic acid or carboxylic acid isostere, α -tocopherol acetate, ethylene oxide (40 mole) adducts of hardened castor oil, perfume and a dye. The resulting mixture is stirred and dissolved, and purified water is added to the
30 mixture to obtain a transparent liquid lotion.

5 mL of the lotion may be applied once or twice per day to a site having marked baldness or alopecia.

Example 7

- A lotion comprising the following composition shown
35 may be prepared.

	(%)
95% Ethanol	80.0
an N-heterocyclic carboxylic acid or carboxylic acid isostere	0.005
Hinokitol	0.01
Ethylene oxide (40 mole) adducts of hardened castor oil	0.5
Purified water	19.0
Perfume and dye	q.s.

Into 95% ethanol are added an N-heterocyclic carboxylic acid or carboxylic acid isostere, hinokitol, ethylene oxide (40 mole) adducts of hardened castor oil, perfume, and a dye. The resulting mixture is stirred, and purified water is added to the mixture to obtain a transparent liquid lotion.

The lotion may be applied by spraying once to 4 times per day to a site having marked baldness or alopecia.

Example 8

An emulsion may be prepared from A phase and B phase having the following compositions.

(A phase)	(%)
Whale wax	0.5
Cetanol	2.0
Petrolatum	5.0
Squalane	10.0
Polyoxyethylene (10 mole) monostearate	2.0
Sorbitan monooleate	1.0
an N-heterocyclic carboxylic acid or carboxylic acid isostere	0.01
(B phase)	(%)
Glycerine	10.0
Purified water	69.0
Perfume, dye, and preservative	q.s.

The A phase and the B phase are respectively heated and melted and maintained at 80°C. Both phases are then mixed and cooled under stirring to normal temperature to obtain an emulsion.

- 5 The emulsion may be applied by spraying once to four times per day to a site having marked baldness or alopecia.

Example 9

- 10 A cream may be prepared from A phase and B phase having the following compositions.

	(A Phase)	(%)
	Fluid paraffin	5.0
	Cetostearyl alcohol	5.5
	Petrolatum	5.5
15	Glycerine monostearate	33.0
	Polyoxyethylene (20 mole) 2-octyldodecyl ether	3.0
	Propylparaben	0.3
	(B Phase)	(%)
20	an N-heterocyclic carboxylic acid or carboxylic acid isostere	0.8
	Glycerine	7.0
	Dipropylene glycol	20.0
	Polyethylene glycol 4000	5.0
25	Sodium Hexametaphosphate	0.005
	Purified water	44.895

- 30 The A phase is heated and melted, and maintained at 70°C. The B phase is added into the A phase and the mixture is stirred to obtain an emulsion. The emulsion is then cooled to obtain a cream.

The cream may be applied once to 4 times per day to a site having marked baldness or alopecia.

Example 10

A liquid comprising the following composition may be prepared.

		(%)
	Polyoxyethylene butyl ether	20.0
5	Ethanol	50.0
	an N-heterocyclic carboxylic acid or carboxylic acid isostere	0.001
	Propylene glycol	5.0
10	Polyoxyethylene hardened castor oil derivative (ethylene oxide 80 mole adducts)	0.4
	Perfume	q.s.
	Purified water	q.s.

15 Into ethanol are added polyoxypropylene butyl ether, propylene glycol, polyoxyethylene hardened castor oil, an N-heterocyclic carboxylic acid or carboxylic acid isostere, and perfume. The resulting mixture is stirred, and purified water is added to the mixture to obtain a
20 liquid.

The liquid may be applied once to 4 times per day to a site having marked baldness or alopecia.

Example 11

25 A shampoo comprising the following composition may be prepared.

		(%)
	Sodium laurylsulfate	5.0
	Triethanolamine laurylsulfate	5.0
	Betaine lauryldimethylaminoacetate	6.0
30	Ethylene glycol distearate	2.0
	Polyethylene glycol	5.0
	an N-heterocyclic carboxylic acid or carboxylic acid isostere	5.0
	Ethanol	2.0

Perfume	0.3
Purified water	69.7

5 Into 69.7 of purified water are added 5.0 g of sodium
laurylsulfate, 5.0 g of triethanolamine laurylsulfate, 6.0
g of betaine lauryldimethyl-aminoacetate. Then a mixture
obtained by adding 5.0 g of an N-heterocyclic carboxylic acid
or carboxylic acid isostere, 5.0 g of polyethylene glycol,
10 and 2.0 g of ethylene glycol distearate to 2.0 g of ethanol,
followed by stirring, and 0.3 g of perfume are successively
added. The resulting mixture is heated and subsequently
cooled to obtain a shampoo.

The shampoo may be used on the scalp once or twice per
day.

15 **Example 12**

A patient is suffering from alopecia senilis. An N-
heterocyclic carboxylic acid or carboxylic acid isostere, or
a pharmaceutical composition comprising the same, may be
administered to the patient. Increased hair growth is
20 expected to occur following treatment.

Example 13

A patient is suffering from male pattern alopecia. An
N-heterocyclic carboxylic acid or carboxylic acid isostere,
or a pharmaceutical composition comprising the same may be
25 administered to the patient. Increased hair growth is
expected to occur following treatment.

Example 14

A patient is suffering from alopecia areata. An N-
heterocyclic carboxylic acid or carboxylic acid isostere, or
30 a pharmaceutical composition comprising the same, may be
administered to the patient. Increased hair growth is
expected to occur following treatment.

Example 15

A patient is suffering from hair loss caused by skin
35 lesions. An N-heterocyclic carboxylic acid or carboxylic

acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 16

5 A patient is suffering from hair loss caused by tumors. An N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 17

10 A patient is suffering from hair loss caused by a systematic disorder, such as a nutritional disorder or an internal secretion disorder. An N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical
15 composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 18

20 A patient is suffering from hair loss caused by chemotherapy. An N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 19

25 A patient is suffering from hair loss caused by radiation. An N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same may, be administered to the patient. Increased hair
30 growth is expected to occur following treatment.

Example 20

 A patient is suffering from a neurodegenerative disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or a pharmaceutical composition comprising

the same is administered. It would be expected that the patient would improve their condition or recover.

Example 21

A patient is suffering from a neurological disorder.
5 A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 22

10 A patient is suffering from stroke. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 23

15 A patient is suffering from Parkinson's Disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient
20 would improve their condition or recover.

Example 24

A patient is suffering from Alzheimer's Disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising
25 same is administered. It would be expected that the patient would improve their condition or recover.

Example 25

A patient is suffering from a peripheral neuropathy. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising
30 same is administered. It would be expected that the patient would improve their condition or recover.

Example 26

A patient is suffering from amyotrophic lateral

sclerosis. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

5 Example 27

A patient is suffering from a spinal injury. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient
10 would improve their condition or recover.

Example 28


A patient is at risk of suffering from a neurodegenerative disease or neurological disorder. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or a pharmaceutical composition comprising
15 the same is prophelactically administered. It would be expected that the patient would be prevented from some or all of the effects of the disease or disorder, or would significantly improve their condition or recover over patients
20 who were not pre-treated.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modification are intended
25 to be included within the scope of the following claims.

Where the terms "comprise", "comprises", "comprised" or "comprising" are used in this specification, they are to be interpreted as specifying the presence of the stated features, integers, steps or components referred to, but not to preclude the presence or addition of one or more other feature, integer, step, component or group thereof.

The claims defining the invention are as follows:-

1. A compound having the formula (I):



5 where


n is 1;


X is either O or S;

R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

10 D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-C₁₀ alkynylene;

R₂ is selected from;





wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,

- COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
 5 -NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and
 -CONR₃CN;

wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl, carbocycle, heterocycle, or R_2 is optionally substituted with one or more substituents selected from R_3 and Z, where

- 10 R_3 and Z are independently hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, aralkyl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉
 15 straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;

or a pharmaceutically acceptable salt, ester or solvate thereof;

provided that:

when D is a bond, and R_2 is COOH,

then R_1 is not:

- 20 -C₁-C₉ straight or branched chain alkyl,
 -C₂-C₉ straight or branched chain alkenyl,
 -C₅-C₇ cycloalkyl,
 -C₅-C₇ cycloalkenyl,

-2-(3,4-dichlorophenyl)ethyl,

-benzyl, or

-Ar₁, where Ar₁ is 1-naphthyl, 2-naphthyl, 2-indolyl,

3-indolyl, 2-furyl, 3-furyl, 2-thiazolyl, 2-thienyl, 3-thienyl, 1-

5 pyridyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl,

and said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar₁ are optionally

substituted with one or more substituents selected from hydrogen, halo,

hydroxyl, nitro, trifluoromethyl, C₁-C₉ straight or branched alkyl, C₂-C₉

straight or branched alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy,

10 benzyloxy, COOH, and amino;

further provided that:

when D is a bond, R₂ is CONZ(R₃), Z is hydrogen or C₁-C₆ alkyl, and R₃ is phenyl or C₂-

C₆ straight or branched chain alkyl or alkenyl,

wherein said alkyl is unsubstituted or substituted in one or more positions with Ar₂

15 as defined below, C₃-C₈ cycloalkyl, cycloalkyl connected by methyl or a C₂-C₆

straight or branched chain alkyl or alkenyl, C₁-C₄ alkyl ester, or Ar₃,

where Ar₃ is selected from 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thiazolyl,

2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one

to three substituents independently selected from hydrogen, halo, hydroxy,


20 nitro, trifluoromethyl, C₁-C₆ straight or branched alkyl, C₂-C₆ straight or

branched alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy,

and amino,

said alkyl ester optionally substituted with phenyl,

or R₃ is the fragment:



25

where R₄ is selected from straight or branched chain C₁-C₈ alkyl optionally substituted with C₃-C₈ cycloalkyl, benzyl, or Ar₂ as defined below,

and where R₅ is selected from phenyl, benzyl, C₁-C₆ straight or branched alkyl, and C₂-C₆ straight or branched alkenyl, where said alkyl or alkenyl is optionally substituted with phenyl,

or where R₂ is COOZ or CONR₆, where R₆ is selected from hydrogen, C₁-C₆ straight or branched alkyl, and C₂-C₆ straight or branched alkenyl, where said alkyl or alkenyl is optionally substituted with phenyl,

then R₁ is not:

- C₁-C₉ straight or branched chain alkyl,
- C₂-C₉ straight or branched chain alkenyl, or
- substituted thiophene,

wherein said alkyl or alkenyl is optionally substituted in one or more positions with:

- C₁-C₄ alkyl,
- C₁-C₄ alkenyl,
- hydroxy,
- C₃-C₈ cycloalkyl,
- C₅-C₇ cycloalkenyl, or

-Ar₂, where Ar₂ is 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents selected from hydrogen, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched alkyl, C₂-C₆ straight or branched alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, and amino,

where said cycloalkyl or cycloalkenyl groups may be optionally substituted with C₁-C₄ alkyl, C₁-C₄ alkenyl, or hydroxy;

further provided that:

when D is CH₂,

then R₂ is not -OMe, -NHMe, or substituted -NHcyclohexyl;

further provided that:


when D is CH₂, and R₂ is -OH,

5 then R₁ is not phenyl or pyrrolidinemethanol;

2. The compound of claim 1, wherein R₂ is selected from:

1
2
3
4

5
6
7
8



wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 .


3. The compound of claim 1, wherein R_2 is selected from:

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -
CONR₃CN.

4. A compound selected from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-hydroxymethylpyrrolidine; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinetetrazole;

(2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-aminocarbonyl piperidine; and compounds 1-25, 27, 28, 31-33, and 35-136.

5. A pharmaceutical composition, comprising:
- an effective amount a compound; and
 - a pharmaceutically acceptable carrier.
- wherein the compound is selected from the compound of formula (I);





where

- n is 1;
- X is either O or S;
- R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;
- D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-C₁₀ alkynylene;
- R₂ is selected from

10
15

20
25
30





wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 .

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and
5 -CONR₃CN; and

wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl, carbocycle, heterocycle, or R₂ is optionally substituted with one or more substituents selected from R₃, where

R₃ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy,
10 alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;


15 or a pharmaceutically acceptable salt, ester or solvate thereof.

6. The pharmaceutical composition of claim 5, further comprising an additional neurotrophic factor.

7. The pharmaceutical composition of claim 6, wherein said additional neurotrophic factor is selected from nerve growth factor, brain derived growth factor, glial cell line derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.

8. The pharmaceutical composition of any one of claims 5 to 7, wherein the compound is non-immunosuppressive.

9. The pharmaceutical composition of claim 5, wherein R₂ is selected from:




wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 .

10. The pharmaceutical composition of claim 5, wherein R_2 is selected from:
- 5 -COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -
CONR₃CN.


11. The pharmaceutical composition of claim 5, wherein the compound is
- 10 selected from any one of the compounds 1-139.

12. Use of a compound
- wherein the compound is selected from a compound of formula (I);



where

- 15 n is 1;
- X is either O or S;
- R_1 is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;
- D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-
- 20 C₁₀ alkynylene;
- R_2 is selected from




- 5 wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,
 $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}_3$, $-\text{PO}_3(\text{R}_3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}_3)_2$, $-\text{OR}_3$, $-\text{SR}_3$,
 $-\text{NHCOR}_3$, $-\text{N}(\text{R}_3)_2$, $-\text{CON}(\text{R}_3)_2$, $-\text{CONH}(\text{O})\text{R}_3$, $-\text{CONHNHSO}_2\text{R}_3$, $-\text{COHNSO}_2\text{R}_3$
and
 $-\text{CONR}_3\text{CN}$; and
10 wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl, carbocycle, heterocycle, or R_2 is optionally substituted with one or more substituents selected from R_3 , where
 R_3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy,

aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;
 5 or a pharmaceutically acceptable salt, ester or solvate thereof, in the preparation of a medicament for treating neurological diseases in an animal.


13. Use of a compound as defined in claim 1 or a pharmaceutically acceptable
 10 salt, ester or solvate thereof, in the preparation of a medicament for treating disease in an animal.

14. Use of a compound in the preparation of a medicament for treating a
 neurological disorder in an animal to stimulate growth of damaged peripheral nerves, to
 15 promote neuronal regeneration, or to prevent neurodegeneration in the animal,
 wherein the compound is selected from a compound of the formula (I);



where

- 20 n is 1-3;
 X is either O or S;
 R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;
 D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-
 25 C₁₀ alkynylene;
 R₂ is selected from



- wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,
- 5 -COOH, -SO₃H, -SO₂HNR₃, -PO₃(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃, -NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -CONR₃CN; and
- 10 wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl, carbocycle, heterocycle, or R₂, is optionally substituted with one or more substituents selected from R₃, where
- R₃ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl,

- sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;
- 5 or a pharmaceutically acceptable salt, ester or solvate thereof.

15. The use of claim 14, wherein the neurological disorder is selected from peripheral neuropathies caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and
- 10 neurological disorders relating to neurodegeneration.

16. The use of claim 14, wherein the neurological disorder is selected from Alzheimer's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular
- 15 dystrophy, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured, or prolapsed invertebrate disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, Gullain-Barré syndrome, and sciatic nerve lesions.

- 20 17. The use of claim 14, wherein the neurological disorder is Alzheimer's disease.


18. The use of claim 14, wherein the neurological disorder is Parkinson's disease.

- 25 19. The use of claim 14, wherein the neurological disorder is amyotrophic lateral sclerosis.

20. The use of any one of claims 14-19, further comprising the use of an
- 30 additional neurotrophic factor.

21. The use of claim 20, wherein said additional neurotrophic factor is selected from nerve growth factor, brain derived growth factor, glial cell line derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.

22. Use of a compound in the preparation of a medicament for treating alopecia or for use in promoting hair growth in an animal wherein the compound is selected from a compound of formula (I);



where


n is 1-3;


X is either O or S;

R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-C₁₀ alkynylene;

R₂ is selected from





wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 .


- 5 $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}_3$, $-\text{PO}_2(\text{R}_3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}_3)_2$, $-\text{OR}_3$, $-\text{SR}_3$,
 $-\text{NHCOR}_3$, $-\text{N}(\text{R}_3)_2$, $-\text{CON}(\text{R}_3)_2$, $-\text{CONH}(\text{O})\text{R}_3$, $-\text{CONHNHSO}_2\text{R}_3$,
 $-\text{COHNSO}_2\text{R}_3$, and
 $-\text{COHNR}_3\text{CN}$; and

10 wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl,
 carbocycle, heterocycle, or R_2 is optionally substituted with one or more substituents
 selected from R_3 , where

15 R_3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy,
 alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl,
 sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 -
 C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle,
 heterocycle, or CO_2R_7 where R_7 is hydrogen or C_1 - C_9 straight or branched chain
 alkyl or C_2 - C_9 straight or branched chain alkenyl;
 or a pharmaceutically acceptable salt, ester or solvate thereof.

20 23. The use of any one of claims 12-22, wherein said compound is non-
 immunosuppressive.

24. The use of claim 22, wherein R_2 is selected from:



wherein the atoms of each ring structure may be optionally substituted at one or more
5 positions with R_3 .


25. The use of claim 22, wherein R_2 is selected from:

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -
10 CONR₃CN.

26. The use of claim 22, wherein the compound is selected from compounds 1-139.

27. A method of treating a neurological disorder in an animal to stimulate growth of damaged peripheral nerves, promoting neuronal regeneration, or preventing neurodegeneration in the animal, comprising:

administering to the animal an effective amount of a compound selected from a compound of formula (I);



10

where


n is 1-3;

X is either O or S;

R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-C₁₀ alkynylene;

R₂ is selected from



- wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,
- 5 -COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃,
and
-CONR₃CN; and
- wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl,
10 carbocycle, heterocycle, or R₂ is optionally substituted with one or more
substituents selected from R₃, where
R₃ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy,
aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl,

thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;

5 or a pharmaceutically acceptable salt, ester or solvate thereof.

28. The method of claim 27, wherein the neurological disorder is selected from peripheral neuropathies caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and
10 neurological disorders relating to neurodegeneration.

29. The method of claim 27, wherein the neurological disorder is selected from Alzheimer's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular
15 dystrophy, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured, or prolapsed intervertebral disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet syndrome syndromes, Guillain-Barré syndrome, and sciatic nerve lesions.

20 30. The method of claim 27, wherein the neurological disorder is Alzheimer's disease.


31. The method of claim 27, wherein the neurological disorder is Parkinson's disease.

25 32. The method of claim 26, wherein the neurological disorder is amyotrophic lateral sclerosis.

33. The method of any one of claims 27-32, further comprising administering
30 an additional neurotrophic factor.

34. The method of claim 33, wherein said additional neurotrophic factor is selected from nerve growth factor, brain derived growth factor, glial cell line derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.

35. A method of treating alopecia or promoting hair growth in an animal, comprising:
administering to the animal an effective amount of a compound of formula (I):



10 where


n is 1-3;

X is either O or S;

R₁ is selected from C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

15 D is a bond, or a C₁-C₁₀ straight or branched chain alkylene, C₂-C₁₀ alkenylene, or C₂-C₁₀ alkynylene;

R₂ is selected from



wherein the atoms of each ring structure may be optionally substituted at one or more positions with R_3 ,

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,

- 5 -NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -CONR₃CN; and

wherein said alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, aryl, heteroaryl, carbocycle, heterocycle, or R₂ is optionally substituted with one or more substituents selected from R₃, where


- 10 R₃ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl,

sulphydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R₇ where R₇ is hydrogen or C₁-C₉ straight or branched chain alkyl or C₂-C₉ straight or branched chain alkenyl;

5 or a pharmaceutically acceptable salt, ester or solvate thereof.

36. The method of any one of claims 27 to 35 wherein the compound is non-immunosuppressive.

10 37. The method of claim 35, wherein R₂ is selected from:



wherein the atoms of each ring structure may be optionally substituted at one or more positions with R₃.

5 38. The method of claim 35, wherein R₂ is selected from:
-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
-NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃, and -
CONR₃CN.

10 39. The method of claim 35, wherein the compound is selected from
compounds 1-139.

40. A compound of any one of claims 1 to 4, substantially as hereinbefore
described.

15 41. A pharmaceutical composition of any one of claims 5 to 11, substantially as
hereinbefore described.

20 42. A use of any one of claims 13 to 26, substantially as hereinbefore described.

43. A method of any one of claims 27 to 39, substantially as hereinbefore
described.

DATED this 26th day of February, 2004

25 **GPI NIL HOLDINGS, INC. and AMGEN, INC.**
By their Patent Attorneys:
CALLINAN LAWRIE





FIG. 1



SUBSTITUTE SHEET (RULE 26)


FIG.2



SUBSTITUTE SHEET (RULE 26)

FIG. 3

Promotion of Hair Growth by Neuroimmunophilin Ligands



SUBSTITUTE SHEET (RULE 26)