

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2004257228 B8**

(54) Title
Method for assaying protein-protein interaction

(51) International Patent Classification(s)
C12Q 1/68 (2006.01) **G01N 33/68** (2006.01)
G01N 33/53 (2006.01) **G01N 33/94** (2006.01)

(21) Application No: **2004257228** (22) Date of Filing: **2004.07.09**

(87) WIPO No: **WO05/007822**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/485,968	2003.07.09	US
60/566,113	2004.04.27	US
60/511,918	2003.10.15	US

(43) Publication Date: **2005.01.27**

(44) Accepted Journal Date: **2008.12.18**

(48) Corrigenda Journal Date: **2009.01.08**

(71) Applicant(s)
Invitrogen Corporation;The Trustees of Columbia University in the City of New York

(72) Inventor(s)
Lee, Kevin J.;Strapps, Walter;Barnea, Gilad;Axel, Richard

(74) Agent / Attorney
Griffith Hack, Level 19 109 St Georges Terrace, Perth, WA, 6000

(56) Related Art
US6455300 (Htun et al) 24 September 2002
AU 2003215660 (WO 2003/076932 A2), Axaron Bioscience AG, published 22 September 2003 and
having a priority date of 13 March 2002
US 2002/0132327 (Hay et al) 19 September 2002
US 2002/106698 A1

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 January 2005 (27.01.2005)

PCT

(10) International Publication Number
WO 2005/007822 A3

- (51) International Patent Classification⁷: G01N 33/53
- (21) International Application Number:
PCT/US2004/021887
- (22) International Filing Date: 9 July 2004 (09.07.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/485,968 9 July 2003 (09.07.2003) US
60/511,918 15 October 2003 (15.10.2003) US
60/566,113 27 April 2004 (27.04.2004) US
- (71) Applicants (for all designated States except US): SENTI-GEN BIOSCIENCES, INC. [US/US]; Audobon Biomedical Center, 3960 Broadway, New York, NY 10031 (US). THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK [US/US]; 254 Engineering Terrace, 1210 Amsterdam Avenue, Mail Code 2205, New York, NY 10027 (US).
- (72) Inventors: LEE, Kevin, J.; 15 West 104th Street, Apt. 3B, New York, NY 10025 (US). AXEL, Richard; 3960 Broadway, New York, NY 10031 (US). STRAPPS, Walter; 3960 Broadway, New York, NY 10031 (US). BARNEA, Gilad; 3960 Broadway, New York, NY 10031 (US).
- (74) Agent: HANSON, Norman, D.; Fulbright & Jaworski L.L.P., 666 Fifth Avenue, New York, NY 10103 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
31 March 2005
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR ASSAYING PROTEIN-PROTEIN INTERACTION

(57) Abstract: The invention relates to a method for determining if a test compound, or a mix of compounds, modulates the interaction between two proteins of interest. The determination is made possible via the use of two recombinant molecules, one of which contains the first protein a cleavage site for a proteolytic molecule, and an activator of a gene. The second recombinant molecule includes the second protein and the proteolytic molecule. If the test compound binds to the first protein, a reaction is initiated whereby the activator is cleaved, and activates a reporter gene.



WO 2005/007822 A3

METHOD FOR ASSAYING PROTEIN-PROTEIN INTERACTION

FIELD OF THE INVENTION

[0001] This invention relates to methods for determining interaction between molecules of interest. More particularly, it relates to determining if a particular substance referred to as the test compound modulates the interaction of two or more specific proteins of interest, via determining activation of a reporter gene in a cell, where the activation, or lack thereof, results from the modulation or its absence. The determination occurs using transformed or transfected cells, which are also a feature of the invention, as are the agents used to transform or transfect them.

BACKGROUND AND RELATED ART

[0002] The study of protein/protein interaction, as exemplified, e.g., by the identification of ligands for receptors, is an area of great interest. Even when a ligand or ligands for a given receptor are known, there is interest in identifying more effective or more selective ligands. GPCRs will be discussed herein as a non-exclusive example of a class of proteins which can be studied in this way.

[0003] The G-protein coupled receptors, or "GPCRs" hereafter, are the largest class of cell surface receptors known for humans. Among the ligands recognized by GPCRs are hormones, neurotransmitters, peptides, glycoproteins, lipids, nucleotides, and ions. They also act as receptors for light, odors, pheromones, and taste. Given these various roles, it is perhaps not surprising that they are the subject of intense research, seeking to identify drugs useful in various conditions. The success rate has been phenomenal. Indeed, Howard, *et al.*, Trends Pharmacol. Sci., 22:132-140 (2001) estimate that over 50% of marketed drugs act on such receptors. "GPCRs" as used herein, refers to any member of the GPCR superfamily of receptors characterized by a seven-transmembrane domain (7TM) structure. Examples of these receptors include, but are not limited to, the class A or "rhodopsin-like" receptors; the class B or "secretin-like" receptors; the class C or "metabotropic glutamate-like" receptors; the Frizzled and Smoothed-related receptors; the adhesion receptor family or EGF-7TM / LNB-7TM receptors; adiponectin receptors and related receptors; and chemosensory receptors including odorant, taste, vomeronasal and pheromone receptors. As examples, the

GPCR superfamily in humans includes but is not limited to those receptor molecules described by Vassilatis, *et al.*, Proc. Natl. Acad. Sci. USA, 100:4903-4908 (2003); Takeda, *et al.*, FEBS Letters, 520:97-101 (2002); Fredricksson, *et al.*, Mol. Pharmacol., 63:1256-1272 (2003); Glusman, *et al.*, Genome Res., 11:685-702 (2001); and Zozulya, *et al.*, Genome Biol., 2:0018.1-0018.12 (2001), all of which are incorporated by reference.

[0004] The mechanisms of action by which GPCRs function has been explicated to some degree. In brief, when a GPCR binds a ligand, a conformational change results, stimulating a cascade of reactions leading to a change in cell physiology. It is thought that GPCRs transduce signals by modulating the activity of intracellular, heterotrimeric guanine nucleotide binding proteins, or "G proteins". The complex of ligand and receptor stimulates guanine nucleotide exchange and dissociation of the G protein heterotrimer into α and $\beta\gamma$ subunits.

[0005] Both the GTP-bound α subunit and the $\beta\gamma$ dimer can act to regulate various cellular effector proteins, including adenylyl cyclase and phospholipase C (PLC). In conventional cell based assays for GPCRs, receptor activity is monitored by measuring the output of a G-protein regulated effector pathway, such as the accumulation of cAMP that is produced by adenylyl cyclase, or the release of intracellular calcium, which is stimulated by PLC activity.

[0006] Conventional G-protein based, signal transduction assays have been difficult to develop for some targets, as a result of two major issues.

[0007] First, different GPCRs are coupled to different G protein regulated signal transduction pathways, and G-protein based assays are dependent on knowing the G-protein specificity of the target receptor, or require engineering of the cellular system, to force coupling of the target receptor to a particular effect or pathway. Second, all cells express a large number of endogenous GPCRs, as well as other signaling factors. As a result, the effector pathways that are measured may be modulated by other endogenous molecules in addition to the target GPCR, potentially leading to false results.

[0008] Regulation of G-protein activity is not the only result of ligand/GPCR binding. Luttrell, *et al.*, J. Cell Sci., 115:455-465 (2002), and Ferguson, Pharmacol. Rev., 53:1-24 (2001), both of which are incorporated by reference, review other activities which lead to termination of the GPCR signal. These termination processes prevent excessive cell

stimulation, and enforce temporal linkage between extracellular signal and corresponding intracellular pathway.

[0009] In the case of binding of an agonist to GPCR, serine and threonine residues at the C terminus of the GPCR molecule are phosphorylated. This phosphorylation is caused by the GPCR kinase, or "GRK," family. Agonist complexed, C-terminal phosphorylated GPCR_s interact with arrestin family members, which "arrest" receptor signaling. This binding inhibits coupling of the receptor to G proteins, thereby targeting the receptor for internalization, followed by degradation and/or recycling. Hence, the binding of a ligand to a GPCR can be said to "modulate" the interaction between the GPCR and arrestin protein, since the binding of ligand to GPCR causes the arrestin to bind to the GPCR, thereby modulating its activity. Hereafter, when "modulates" or any form thereof is used, it refers simply to some change in the way the two proteins of the invention interact, when the test compound is present, as compared to how these two proteins interact, in its absence. For example, the presence of the test compound may strengthen or enhance the interaction of the two proteins, weaken it, inhibit it, or lessen it in some way, manner or form which can then be detected.

[0010] This background information has led to alternate methods for assaying activation and inhibition of GPCRs. These methods involve monitoring interaction with arrestins. A major advantage of this approach is that no knowledge of G-protein pathways is necessary.

[0011] Oakley, *et al.*, Assay Drug Dev. Technol., 1:21-30 (2002) and U.S. Patent Nos. 5,891,646 and 6,110,693, incorporated by reference, describe assays where the redistribution of fluorescently labelled arrestin molecules in the cytoplasm to activated receptors on the cell surface is measured. These methods rely on high resolution imaging of cells, in order to measure arrestin relocalization and receptor activation. It will be recognized by the skilled artisan that this is a complex, involved procedure.

[0012] Various other U.S. patents and patent applications dealing with these points have issued and been filed. For example, U.S. Patent No. 6,528,271 to Bohn, *et al.*, deals with assays for screening for pain controlling medications, where the inhibitor of β -arrestin binding is measured. Published U.S. patent applications, such as 2004/0002119, 2003/0157553, 2003/0143626, and 2002/0132327, all describe different forms of assays

involving GPCRs. Published application 2002/0106379 describes a construct which is used in an example which follows; however, it does not teach or suggest the invention described herein.

5 [0013] It is an object of the invention to develop a simpler assay for monitoring and/or determining modulation of specific protein/protein interactions, where the proteins include but are not limited to, membrane bound proteins, such as receptors, GPCRs in particular. How this
10 is accomplished will be seen in the examples which follow.

[0013a] In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or
15 necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or
addition of further features in various embodiments of the invention.

20 [0013b] It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

25

SUMMARY OF THE INVENTION

[0014] Thus, in accordance with the present invention, there is provided a method for determining if a test compound modulates a specific protein/protein
30 interaction of interest comprising contacting said

compound to a cell which has been transformed or transfected with (a) a nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes said first test protein, (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, and (b) a nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound is to be measured, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site, and determining activity of said reporter gene as a determination of whether said compound modulates said protein/protein interaction.

[0015] The first test protein may be a membrane bound protein, such as a transmembrane receptor, and in particular a GPCR. Particular transmembrane receptors include β 2-adrenergic receptor (ADRB2), arginine vasopressin receptor 2 (AVPR2), serotonin receptor 1a (HTR1A), m2 muscarinic acetylcholine receptor (CHRM2), chemokine (C-C motif) receptor 5 (CCR5), dopamine D2 receptor (DRD2), kappa opioid receptor (OPRK), or α -adregenic receptor (ADRA1A) although it is to be understood that in all cases the invention is not limited to these specific embodiments. For example, molecules such as the insulin growth factor-1 receptor (IGF-1R), which is a tyrosine kinase, and proteins which are not normally membrane bound, like estrogen receptor 1 (ESR1) and estrogen receptors 2 (ESR2). The protease or portion of a protease may be a tobacco etch

virus nuclear inclusion A protease. The protein which activates said reporter gene may be a transcription factor, such as tTA or GAL4. The second protein may be an inhibitory protein, such as an arrestin. The cell may be a eukaryote or a prokaryote. The reporter gene may be an exogenous gene, such as β -galactosidase or luciferase.

[0016] The nucleotide sequence encoding said first test protein may be modified to increase interaction with said second test protein. Such modifications include but are not limited to replacing all or part of the nucleotide sequence of the C-terminal region of said first test protein with a nucleotide sequence which encodes an amino acid sequence which has higher affinity for said second test protein than the original sequence. For example, the C-terminal region may be replaced by a nucleotide sequence encoding the C-terminal region of AVPR2, AGTRL1, GRPR, F2RL1, CXCR2/IL-8b, CCR4, or GRPR.

[0017] The method may comprise contacting more than one test compound to a plurality of samples of cells, each of said samples being contacted by one or more of said test compounds, wherein each of said cell samples have been transformed or transfected with the aforementioned nucleic acid molecules, and determining activity of reporter genes in said plurality of said samples to determine if any of said test compounds modulate a specific, protein/protein interaction. The method may comprise contacting each of said samples with one test compound, each of which differs from all others, or comprise contacting each of said samples with a mixture of said test compounds.

[0018] In another embodiment, there is provided a method for determining if a test compound modulates one or more of a plurality of protein interactions of interest, comprising contacting said test compound to a plurality of samples of cells, each of which has been transformed or transfected with (a) a first nucleic acid molecule comprising, (i) a nucleotide sequence which encodes a first test protein, a nucleotide sequence encoding a cleavage site for a protease, and (ii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, (b) a second nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound of interest is to be measured, (ii) a nucleotide sequence which encodes a protease or a protease which is specific for said cleavage site, wherein said first test protein differs from other first test proteins in each of said plurality of

samples, and determining activity of said reporter gene in at one or more of said plurality of samples as a determination of modulation of one or more protein interactions of interest

[0019] The second test protein may be different in each sample or the same in each sample. All of said samples may be combined in a common receptacle, and each sample comprises a different pair of first and second test proteins. Alternatively, each sample may be tested in a different receptacle. The reporter gene in a given sample may differ from the reporter gene in other samples. The mixture of test compounds may comprise or be present in a biological sample, such as cerebrospinal fluid, urine, blood, serum, pus, ascites, synovial fluid, a tissue extract, or an exudate.

[0020] In yet another embodiment, there is provided a recombinant cell, transformed or transfected with (a) a nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes said first test protein, (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, and (b) a nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound is to be measured, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site.

[0021] One or both of said nucleic acid molecules may be stably incorporated into the genome of said cell. The cell also may have been transformed or transfected with said reporter gene. The first test protein may be a membrane bound protein, such as a transmembrane receptor, and in particular a GPCR. Particular transmembrane receptors include ADRB2, AVPR2, HTR1A, CHRM2, CCR5, DRD2, OPRK, or ADRA1A.

[0022] The protease or portion of a protease may be a tobacco etch virus nuclear inclusion A protease. The protein which activates said reporter gene may be a transcription factor, such as tTA or GAL4. The second protein may be an inhibitory protein. The cell may be a eukaryote or a prokaryote. The reporter gene may be an exogenous gene, such as β -galactosidase or luciferase. The nucleotide sequence encoding said first test protein may be modified to increase interaction with said second test protein, such as by replacing all or part of the nucleotide sequence of the C-terminal region of said first test protein with a nucleotide sequence which encodes an amino acid sequence which has higher affinity for

said second test protein than the original sequence. The C-terminal region may be replaced by a nucleotide sequence encoding the C-terminal region of AVPR2, AGTRLI, GRPR, F2RL1, CXCR2/IL-8B, CCR4, or GRPR.

[0023] In still yet another embodiment, there is provided an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell. The test protein may be a membrane bound protein, such as is a transmembrane receptor. A particular type of transmembrane protein is a GPCR. Particular transmembrane receptors include ADRB2, AVPR2, HTR1A, CHRM2, CCR5, DRD2, OPRK, or ADRA1A. The protease or portion of a protease may be a tobacco etch virus nuclear inclusion A protease. The protein which activates said reporter gene may be a transcription factor, such as tTA or GAL4. As above, the invention is not to be viewed as limited to these specific embodiments.

[0024] In still a further embodiment, there is provided an expression vector comprising an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, and further being operably linked to a promoter.

[0025] In still yet a further embodiment, there is provided an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein whose interaction with another test protein in the presence of a test compound is to be measured, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site. The test protein may be an inhibitory protein, such as an arrestin.

[0026] Also provided is an expression vector comprising an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein whose interaction with another test protein in the presence of a test compound is to be measured, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site, said nucleic acid further being operably linked to a promoter.

[0027] An additional embodiment comprises a fusion protein produced by expression of:

[0028] an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, and further being operably linked to a promoter; or

[0029] an isolated nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a test protein whose interaction with another test protein in the presence of a test compound is to be measured, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site

[0030] In yet another embodiment, there is provided a test kit useful for determining if a test compound modulates a specific protein/protein interaction of interest comprising a separate portion of each of (a) a nucleic acid molecule which comprises, a nucleotide sequence which encodes said first test protein (i) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, (ii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, and (b) a nucleic acid molecule which comprises, (i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound is to be measured, (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site, and container means for holding each of (a) and (b) separately from each other.

[0031] The first test protein may be a membrane bound protein, such as a transmembrane receptor. A particular type of transmembrane receptor is a GPCR. A particular transmembrane protein is a GPCR. Particular transmembrane receptors include ADRB2, AVPR2, HTR1A, CHRM2, CCR5, DRD2, OPRK, or ADRA1A. The protease or portion of a protease may be tobacco etch virus nuclear inclusion A protease. The protein which activates said reporter gene may be a transcription factor, such as tTA or GAL4. The second protein may be an inhibitory protein, such as an arrestin. The kit may further comprise a separate portion of an isolated nucleic acid molecule which encodes a reporter gene. The reporter gene may encode β -galactosidase or luciferase. The nucleotide sequence

encoding said first test protein may be modified to increase interaction with said second test protein, such as by replacing all or part of the nucleotide sequence of the C-terminal region of said first test protein with a nucleotide sequence which encodes an amino acid sequence which has higher affinity for said second test protein than the original sequence. The nucleotide sequence of said C-terminal region may be replaced by a nucleotide sequence encoding the C-terminal region of AVPR2, AGTRL1, GRPR, F2RL1, CXCR2/IL-8B, CCR4, or GRPR.

[0032] It is contemplated that any method or composition described herein can be implemented with respect to any other method or composition described herein. The use of the word "a" or "an" when used in conjunction with the term "comprising" in the claims and/or the specification may mean "one," but it is also consistent with the meaning of "one or more," "at least one," and "one or more than one."

[0033] These, and other, embodiments of the invention will be better appreciated and understood when considered in conjunction with the following description and the accompanying drawings. It should be understood, however, that the following description, while indicating various embodiments of the invention and numerous specific details thereof, is given by way of illustration and not of limitation. Many substitutions, modifications, additions and/or rearrangements may be made within the scope of the invention without departing from the spirit thereof, and the invention includes all such substitutions, modifications, additions and/or rearrangements.

BRIEF DESCRIPTION OF THE FIGURES

[0034] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0035] Figure 1 shows the conceptual underpinnings of the invention, pictorially, using ligand-receptor binding as an example.

[0036] Figures 2a and 2b show that the response of targets in assays in accordance with the invention is dose dependent, both for agonists and antagonists.

[0037] Figure 3 shows that a dose response curve results with a different target and a different agonist as well.

[0038] Figure 4 depicts results obtained in accordance with the invention, using the D2 dopamine receptor.

[0039] Figures 5a and 5b illustrate results of an assay which shows that two molecules can be studied simultaneously.

[0040] Figure 6 sets forth the result of another "multiplex" assay, i.e., one where two molecules are studied simultaneously.

[0041] Figure 7 presents data obtained from assays measuring EGFR activity.

[0042] Figure 8 presents data obtained from assays in accordance with the invention, designed to measure the activity of human type I interferon receptor.

[0043] Figure 9 elaborates on the results in figure 7, showing a dose response curve for IFN- α in the cells used to generate figure 7.

[0044] Figure 10 shows the results of additional experiments where a different transcription factor, and a different cell line, were used.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0045] The present invention relates to methods for determining if a substance of interest modulates interaction of a first test protein, such as a membrane bound protein, like a receptor, e.g., a transmembrane receptor, with a second test protein, like a member of the arrestin family. The methodology involves cotransforming or cotransfecting a cell, which may be prokaryotic or eukaryotic, with two constructs. The first construct includes, a sequence encoding (i) the first test protein, such as a transmembrane receptor, (ii) a cleavage site for a protease, and (iii) a sequence encoding a protein which activates a reporter gene. The second construct includes, (i) a sequence which encodes a second test protein whose interaction with the first test protein is measured and/or determined, and (ii) a nucleotide sequence which encodes a protease or a portion of a protease sufficient to act on the cleavage site that is part of the first construct. In especially preferred embodiments, these constructs become stably integrated into the cells.

[0046] The features of an embodiment of the invention are shown, pictorially, in Figure 1. In brief, first, standard techniques are employed to fuse DNA encoding a

transcription factor to DNA encoding a first test protein, such as a transmembrane receptor molecule, being studied. This fusion is accompanied by the inclusion of a recognition and cleavage site for a protease not expressed endogenously by the host cell being used in the experiments.

[0047] DNA encoding this first fusion protein is introduced into and is expressed by a cell which also contains a reporter gene sequence, under the control of a promoter element which is dependent upon the transcription factor fused to the first test protein, e.g., the receptor. If the exogenous protease is not present, the transcription factor remains tethered to the first test protein and is unable to enter the nucleus to stimulate expression of the reporter gene.

[0048] Recombinant techniques can also be used to produce a second fusion protein. In the depicted embodiment, DNA encoding a member of the arrestin family is fused to a DNA molecule encoding the exogenous protease, resulting in a second fusion protein containing the second test protein, i.e., the arrestin family member.

[0049] An assay is then carried out wherein the second fusion protein is expressed, together with the first fusion protein, and a test compound is contacted to the cells, preferably for a specific length of time. If the test compound modulates interaction of the two test proteins, e.g., by stimulating, promoting or enhancing the association of the first and second test proteins, this leads to release of the transcription factor, which in turn moves to the nucleus, and provokes expression of the reporter gene. The activity of the reporter gene is measured.

[0050] In an alternative system, the two test proteins may interact in the absence of the test compound, and the test compound may cause the two test proteins to dissociate, lessen or inhibit their interaction. In such a case, the level of free, functionally active transcription factor in the cell decreases in the presence of the test compound, leading to a decrease in proteolysis, and a measurable decrease in the activity of the reporter gene.

[0051] In the depicted embodiment, the arrestin protein, which is the second test protein, binds to the receptor in the presence of an agonist; however, it is to be understood that since receptors are but one type of protein, the assay is not dependent upon the use of receptor molecules, nor is agonist binding the only interaction capable of being involved. Any protein will suffice, although the interest in transmembrane proteins is clear. Further,

agonist binding to a receptor is not the only type of binding which can be assayed. One can determine antagonists, *per se* and also determine the relative strengths of different antagonists and/or agonists in accordance with the invention.

[0052] Other details of the invention, include specific methods and technology for making and using the subject matter thereof, are described below.

I. Expression Constructs and Transformation

[0053] The term "vector" is used to refer to a carrier nucleic acid molecule into which a nucleic acid sequence can be inserted for introduction into a cell where it can be replicated. A nucleic acid sequence can be "exogenous," which means that it is foreign to the cell into which the vector is being introduced or that the sequence is homologous to a sequence in the cell but in a position within the host cell nucleic acid in which the sequence is ordinarily not found. Vectors include plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (*e.g.*, YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (see, for example, Maniatis, *et al.*, Molecular Cloning, A Laboratory Manual (Cold Spring Harbor, 1990) and Ausubel, *et al.*, 1994, Current Protocols In Molecular Biology (John Wiley & Sons, 1996), both incorporated herein by reference).

[0054] The term "expression vector" refers to any type of genetic construct comprising a nucleic acid coding for a RNA capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. In other cases, these sequences are not translated, for example, in the production of antisense molecules or ribozymes. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host cell. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleotide sequences that serve other functions as well and are described *infra*.

[0055] In certain embodiments, a plasmid vector is contemplated for use in cloning and gene transfer. In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences

which are capable of providing phenotypic selection in transformed cells. In a non-limiting example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, for example, promoters which can be used by the microbial organism for expression of its own proteins.

[0056] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEMTM-11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as, for example, *E. coli* LE392.

[0057] Bacterial host cells, for example, *E. coli*, comprising the expression vector, are grown in any of a number of suitable media, for example, LB. The expression of the recombinant protein in certain vectors may be induced, as would be understood by those of skill in the art, by contacting a host cell with an agent specific for certain promoters, e.g., by adding IPTG to the media or by switching incubation to a higher temperature. After culturing the bacteria for a further period, generally of between 2 and 24 h, the cells are collected by centrifugation and washed to remove residual media.

[0058] Many prokaryotic vectors can also be used to transform eukaryotic host cells. However, it may be desirable to select vectors that have been modified for the specific purpose of expressing proteins in eukaryotic host cells. Expression systems have been designed for regulated and/or high level expression in such cells. For example, the insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Patents 5,871,986 and 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC[®] 2.0 from INVITROGEN[®] and BACPACKTM BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH[®].

[0059] Other examples of expression systems include STRATAGENE[®]'s COMPLETE CONTROLTM Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System, an *E. coli* expression system. Another example of an inducible expression system is available from INVITROGEN[®], which carries the

T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the *Pichia methanolica* Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast *Pichia methanolica*. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

Regulatory Signals

[0060] The construct may contain additional 5' and/or 3' elements, such as promoters, poly A sequences, and so forth. The elements may be derived from the host cell, i.e., homologous to the host, or they may be derived from distinct source, i.e., heterologous.

[0061] A "promoter" is a control sequence that is a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind, such as RNA polymerase and other transcription factors, to initiate the specific transcription a nucleic acid sequence. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and/or expression of that sequence.

[0062] A promoter generally comprises a sequence that functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as, for example, the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation. Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. To bring a coding sequence "under the control of" a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame "downstream" of (i.e., 3' of) the chosen promoter. The "upstream" promoter stimulates transcription of the DNA and promotes expression of the encoded RNA.

[0063] The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0064] A promoter may be one naturally associated with a nucleic acid molecule, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment and/or exon. Such a promoter can be referred to as "endogenous." Similarly, an enhancer may be one naturally associated with a nucleic acid molecule, located either downstream or upstream of that sequence. Alternatively, certain advantages will be gained by positioning the coding nucleic acid segment under the control of a recombinant or heterologous promoter, which refers to a promoter that is not normally associated with a nucleic acid molecule in its natural environment. A recombinant or heterologous enhancer refers also to an enhancer not normally associated with a nucleic acid molecule in its natural environment. Such promoters or enhancers may include promoters or enhancers of other genes, and promoters or enhancers isolated from any other virus, or prokaryotic or eukaryotic cell, and promoters or enhancers not "naturally occurring," *i.e.*, containing different elements of different transcriptional regulatory regions, and/or mutations that alter expression. For example, promoters that are most commonly used in recombinant DNA construction include the β -lactamase (penicillinase), lactose and tryptophan (trp) promoter systems. In addition to producing nucleic acid sequences of promoters and enhancers synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including PCRTM, in connection with the compositions disclosed herein (see U.S. Patents Nos. 4,683,202 and 5,928,906, each incorporated herein by reference). Furthermore, it is contemplated the control sequences that direct transcription and/or expression of sequences within non-nuclear organelles such as mitochondria, chloroplasts, and the like, can be employed as well.

[0065] Naturally, it will be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the organelle, cell type, tissue, organ, or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression, (see, for example Sambrook, *et al.*, 1989, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, inducible, and/or useful under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins and/or peptides. The promoter may be heterologous or endogenous.

[0066] Additionally any promoter/enhancer combination (as per, for example, the Eukaryotic Promoter Data Base EPDB, www.epd.isb-sib.ch/) could also be used to drive expression. Use of a T3, T7 or SP6 cytoplasmic expression system is another possible embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[0067] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements.

[0068] In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, *Nature*, 334:320-325 (1988)). IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described (Pelletier and Sonenberg, *supra*), as well an IRES from a mammalian message (Macejak and Sarnow, *Nature*, 353:90-94 (1991))1991).

IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Patent Nos. 5,925,565 and 5,935,819, each herein incorporated by reference).

Other Vector Sequence Elements

[0069] Vectors can include a multiple cloning site (MCS), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector (see, for example, Carbonelli, *et al.*, FEMS Microbiol. Lett., 172(1):75-82 (1999), Levenson, *et al.*, Hum. Gene Ther. 9(8):1233-1236 (1998), and Cocea, Biotechniques, 23(5):814-816 (1997)), incorporated herein by reference.) "Restriction enzyme digestion" refers to catalytic cleavage of a nucleic acid molecule with an enzyme that functions only at specific locations in a nucleic acid molecule. Many of these restriction enzymes are commercially available. Use of such enzymes is widely understood by those of skill in the art. Frequently, a vector is linearized or fragmented using a restriction enzyme that cuts within the MCS to enable exogenous sequences to be ligated to the vector. "Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments, which may or may not be contiguous with each other. Techniques involving restriction enzymes and ligation reactions are well known to those of skill in the art of recombinant technology.

[0070] Most transcribed eukaryotic RNA molecules will undergo RNA splicing to remove introns from the primary transcripts. Vectors containing genomic eukaryotic sequences may require donor and/or acceptor splicing sites to ensure proper processing of the transcript for protein expression (see, for example, Chandler, *et al.*, 1997, herein incorporated by reference).

[0071] The vectors or constructs of the present invention will generally comprise at least one termination signal. A "termination signal" or "terminator" comprises a DNA sequence involved in specific termination of an RNA transcript by an RNA polymerase. Thus, in certain embodiments a termination signal that ends the production of an RNA

transcript is contemplated. A terminator may be necessary *in vivo* to achieve desirable message levels.

[0072] In eukaryotic systems, the terminator region may also comprise specific DNA sequences that permit site-specific cleavage of the new transcript so as to expose a polyadenylation site. This signals a specialized endogenous polymerase to add a stretch of about 200 adenosine residues (polyA) to the 3' end of the transcript. RNA molecules modified with this polyA tail appear to more stable and are translated more efficiently. Thus, in other embodiments involving eukaryotes, it is preferred that that terminator comprises a signal for the cleavage of the RNA, and it is more preferred that the terminator signal promotes polyadenylation of the message. The terminator and/or polyadenylation site elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

[0073] Terminators contemplated for use in the invention include any known terminator of transcription described herein or known to one of ordinary skill in the art, including but not being limited to, for example, the termination sequences of genes, such as the bovine growth hormone terminator, viral termination sequences, such as the SV40 terminator. In certain embodiments, the termination signal may be a lack of transcribable or translatable sequence, such as an untranslatable/untranscribable sequence due to a sequence truncation.

[0074] In expression, particularly eukaryotic expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal or the bovine growth hormone polyadenylation signal, both of which are convenient, readily available, and known to function well in various target cells. Polyadenylation may increase the stability of the transcript or may facilitate cytoplasmic transport.

[0075] In order to propagate a vector in a host cell, it may contain one or more origins of replication ~~sites~~ (often termed "ori"), sites, which are specific nucleotide sequences at which replication is initiated. Alternatively, an autonomously replicating sequence (ARS) can be employed if the host cell is yeast.

Transformation Methodology

[0076] Suitable methods for nucleic acid delivery for use with the current invention are believed to include virtually any method by which a nucleic acid molecule (e.g., DNA) can be introduced into a cell as described herein or as would be known to one of ordinary skill in the art. Such methods include, but are not limited to, direct delivery of DNA such as by ex vivo transfection (Wilson, *et al.*, Science, 244:1344-1346 (1989), Nabel *et al.*, Science, 244:1342-1344 (1989), by injection (U.S. Patent Nos. 5,994,624, 5,981,274, 5,945,100, 5,780,448, 5,736,524, 5,702,932, 5,656,610, 5,589,466 and 5,580,859, each incorporated herein by reference), including microinjection (Harlan and Weintraub, J. Cell Biol., 101(3):1094-1099 (1985); U.S. Patent No. 5,789,215, incorporated herein by reference); by electroporation (U.S. Patent No. 5,384,253, incorporated herein by reference; Tur-Kaspa, *et al.*, Mol. Cell Biol., 6:716-718 (1986); Potter, *et al.*, Proc. Natl. Acad. Sci. USA, 81:7161-7165 (1984); by calcium phosphate precipitation (Graham and Van Der Eb, Virology, 52:456-467 (1973); Chen and Okayama, Mol. Cell Biol., 7(8):2745-2752 (1987); Rippe, *et al.*, Mol. Cell Biol., 10:689-695 (1990); by using DEAE-dextran followed by polyethylene glycol (Gopal, Mol. Cell Biol., 5:1188-190 (1985); by direct sonic loading (Fechheimer, *et al.*, Proc. Natl. Acad. Sci. USA, 89(17):8463-8467 (1987); by liposome mediated transfection (Nicolau and Sene, Biochem. & Biophys. Acta, 721:185-190 (1982); Fraley, *et al.*, Proc. Natl. Acad. Sci. USA, 76:3348-3352 (1979); Nicolau, *et al.*, Meth. Enzym., 149:157-176 (1987); Wong, *et al.*, Gene, 10:879-894 (1980); Kaneda, *et al.*, Science, 243:375-378 (1989); Kato, *et al.*, J. Biol. Chem., 266:3361-3364 (1991) and receptor-mediated transfection (Wu and Wu, J. Biol. Chem., 262:4429-4432 (1987); Wu and Wu, 1988); by PEG-mediated transformation of protoplasts (Omirulleh, *et al.*, Plant Mol. Biol., 21(3):415-428 (1987); U.S. Patent Nos. 4,684,611 and 4,952,500, each incorporated herein by reference); by desiccation/inhibition-mediated DNA uptake (Potrykus, *et al.* Mol. Gen. Genet., 199(2):169-177 (1985), and any combination of such methods.

II. Components of the Assay System

[0077] As with the method described herein, the products which are features of the invention have preferred embodiments. For example, in the "three part construct," i.e., that contain sequences encoding a test protein, the cleavage site, and the activator protein, the test

protein is preferably a membrane bound protein, such as a transmembrane receptor, e.g., a member of the GPCR family. These sequences can be modified so that the C terminus of the proteins they encode have better and stronger interactions with the second protein. The modifications can include, e.g., replacing a C-terminal encoding sequence of the test protein, such as a GPCR, with the C terminal coding region for AVPR2, AGTRLI, GRPR, F2PLI, CCR4, CXCR2/IL-8, CCR4, or GRPR, all of which are defined supra.

[0078] The protein which activates the reporter gene may be a protein which acts within the nucleus, like a transcription factor (e.g., tTA, GAL4, etc.), or it may be a molecule that sets a cascade of reactions in motion, leading to an intranuclear reaction by another protein. The skilled artisan will be well versed in such cascades.

[0079] The second construct, as described supra, includes a region which encodes a protein that interacts with the first protein, leading to some measurable phenomenon. The protein may be an activator, an inhibitor, or, more, generically, a "modulator" of the first protein. Members of the arrestin family are preferred, especially when the first protein is a GPCR, but other protein encoding sequences may be used, especially when the first protein is not a GPCR. The second part of these two part constructs encodes the protease, or portion of a protease, which acts to remove the activating molecule from the fusion protein encoded by the first construct.

[0080] However, these preferred embodiments do not limit the invention, as discussed in the following additional embodiments.

Host Cells

[0081] As used herein, the terms "cell," "cell line," and "cell culture" may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. The host cells generally will have been engineered to express a screenable or selectable marker which is activated by the transcription factor that is part of a fusion protein, along with the first test protein.

[0082] In the context of expressing a heterologous nucleic acid sequence, "host cell" refers to a prokaryotic or eukaryotic cell that is capable of replicating a vector and/or expressing a heterologous gene encoded by a vector. When host cells are "transfected" or "transformed" with nucleic acid molecules, they are referred to as "engineered" or

"recombinant" cells or host cells, e.g., a cell into which an exogenous nucleic acid sequence, such as, for example, a vector, has been introduced. Therefore, recombinant cells are distinguishable from naturally-occurring cells which do not contain a recombinantly introduced nucleic acid.

[0083] Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials (www.atcc.org). An appropriate host can be determined by one of skill in the art based on the vector backbone and the desired result. A plasmid or cosmid, for example, can be introduced into a prokaryote host cell for replication of many vectors. Cell types available for vector replication and/or expression include, but are not limited to, bacteria, such as *E. coli* (e.g., *E. coli* strain RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776 (ATCC No. 31537) as well as *E. coli* W3110 (F-, lambda-, prototrophic, ATCC No. 273325), DH5 α , JM109, and KC8, bacilli such as *Bacillus subtilis*; and other enterobacteriaceae such as *Salmonella typhimurium*, *Serratia marcescens*, various *Pseudomonas* specie, as well as a number of commercially available bacterial hosts such as SURE[®] Competent Cells and SOLOPACK[™] Gold Cells (STRATAGENE[®], La Jolla). In certain embodiments, bacterial cells such as *E. coli* LE392 are particularly contemplated as host cells for phage viruses.

[0084] Examples of eukaryotic host cells for replication and/or expression of a vector include, but are not limited to, HeLa, NIH3T3, Jurkat, 293, COS, CHO, Saos, and PC12. Many host cells from various cell types and organisms are available and would be known to one of skill in the art. Similarly, a viral vector may be used in conjunction with either a eukaryotic or prokaryotic host cell, particularly one that is permissive for replication or expression of the vector.

Test Proteins

[0085] The present invention contemplates the use of any two proteins for which a physical interaction is known or suspected. The proteins will exist as fusions proteins, a first test protein fused to a transcription factor, and the second test protein fused to a protease that recognizes a cleavage site in the first fusion protein, cleavage of which releases the transcription factor. The only requirements for the test proteins/fusions are (a) that the first test protein cannot localize to the nucleus prior to cleavage, and (b) that the protease must

remain active following both fusion to the second test protein and binding of the first test protein to the second test protein.

[0086] With respect to the first construct, the first test protein may be, e.g., a naturally membrane bound protein, or one which has been engineered to become membrane bound, via standard techniques. The first test protein may be, e.g., a transmembrane receptor such as any of the GPCRs, or any other transmembrane receptor of interest, including, but not being limited to, receptor tyrosine kinases, receptor serine threonine kinases, cytokine receptors, and so forth. Further, as it is well known that portions of proteins, will function in the same manner as the full length first test protein, such active portions of a first test protein are encompassed by the definition of protein herein.

[0087] As will be evident to the skilled artisan, the present invention may be used to assay for interaction with any protein, and is not limited in its scope to assaying membrane bound receptor, like the GPCRs. For example, the activity of other classes of transmembrane receptors, including but not limited to: receptor tyrosine kinases (RTKs), such as IGF1R, such as the epidermal growth factor receptor (EGFR), ErbB2/HER2/Neu or related RTKs; receptor serine/threonine kinases, such as Transforming Growth Factor-beta (TGF β), activin, or Bone Morphogenetic Protein (BMP) receptors; cytokine receptors, such as receptors for the interferon family for interleukin, erythropoietin, G-CSF, GM-CSF, tumor necrosis factor (TNF) and leptin receptors; and other receptors, which are not necessarily normally membrane bound, such as estrogen receptor 1 (ESR1), and estrogen receptor 2 (ESR2). In each case, the method involves transfecting a cell with a modified receptor construct that directs the expression of a chimeric protein containing the receptor of interest, to which is appended, a protease cleavage site followed by a nucleic acid molecule encoding a transcription factor. The cell is co-transfected with a second construct that directs the expression of a chimeric protein consisting of an interacting protein fused, to the protease that recognizes and cleaves the site described *supra*. In the case of RTKs, such as the EGFR, this interacting protein may consist of a SH2 (Src homology domain 2) containing protein or portion thereof, such as phospholipase C (PLC) or Src homology 2 domain containing transforming protein 1 (SHC1). In the case of receptor serine/threonine kinases, such as TGF β , activin, BMP receptors, this interacting protein may be a Smad protein or portion thereof. In the case of cytokine receptors, such as interferon- α/β or interferon- γ gamma

receptors, this interacting protein may be a signal transducer and activator of transcription (STAT) protein such as, but not being limited to, Stat1, Stat2; Janus kinase (JAK) proteins Jak1, Jak2, or Tyk2; or portions thereof. In each case, the transfected cell contains a reporter gene that is regulated by the transcription factor fused to the receptor. An assay is then performed in which the transfected cells are treated with a test compound for a specific period and the reporter gene activity is measured at the end of the test period. If the test compound activates the receptor of interest, interactions between the receptor of interest and the interacting protein are stimulated, leading to cleavage of the protease site and release of the fused transcription factor, which is in turn measurable as an increase in reporter gene activity.

[0088] Other possible test protein pairs include antibody-ligands, enzyme-substrates, dimerizing proteins, components of signal transduction cascades, and other protein pairs well known to the art.

Reporters

[0089] The protein which activates a reporter gene may be any protein having an impact on a gene, expression or lack thereof which leads to a detectable signal. Typical protein reporters include enzymes such as chloramphenicol acetyl transferase (CAT), β -glucuronidase (GUS) or β -galactosidase. Also contemplated are fluorescent and chemiluminescent proteins such as green fluorescent protein, red fluorescent protein, cyan fluorescent protein luciferase, beta lactamase, and alkaline phosphatase.

Transcriptions Factors and Repressors

[0090] In accordance with the present invention, transcription factors are used to activate expression of a reporter gene in an engineered host cell. Transcription factors are typically classified according to the structure of their DNA-binding domain, which are generally (a) zinc fingers, (b) helix-turn-helix, (c) leucine zipper, (d) helix-loop-helix, or (e) high mobility groups. The activator domains of transcription factors interact with the components of the transcriptional apparatus (RNA polymerase) and with other regulatory proteins, thereby affecting the efficiency of DNA binding.

[0091] The Rel/Nuclear Factor κ B (NF- κ B) and Activating Protein-1 (AP-1) are among the most studied transcription factor families. They have been identified as important components of signal transduction pathways leading to pathological outcomes such as

inflammation and tumorigenesis. Other transcription factor families include the heat shock/E2F family, POU family and the ATF family. Particular transcription factors, such as *tTA* and *GAL4*, are contemplated for use in accordance with the present invention.

[0092] Though transcription factors are one class of molecules that can be used, the assays may be modified to accept the use of transcriptional repressor molecules, where the measurable signal is downregulation of a signal generator, or even cell death.

Proteases and Cleavage Sites

[0093] Proteases are well characterized enzymes that cleave other proteins at a particular site. One family, the Ser/Thr proteases, cleave at serine and threonine residues. Other proteases include cysteine or thiol proteases, aspartic proteases, metalloproteinases, aminopeptidases, di & tripeptidases, carboxypeptidases, and peptidyl peptidases. The choice of these is left to the skilled artisan and certainly need not be limited to the molecules described herein. It is well known that enzymes have catalytic domains and these can be used in place of full length proteases. Such are encompassed by the invention as well. A specific embodiment is the tobacco etch virus nuclear inclusion A protease, or an active portion thereof. Other specific cleavage sites for proteases may also be used, as will be clear to the skilled artisan.

Modification of Test Proteins

[0094] The first test protein may be modified to enhance its binding to the interacting protein in this assay. For example, it is known that certain GPCRs bind arrestins more stably or with greater affinity upon ligand stimulation and this enhanced interaction is mediated by discrete domains, e.g., clusters of serine and threonine residues in the C-terminal tail (Oakley, *et al.*, *J. Biol. Chem.*, 274:32248-32257, 1999 and Oakley, *et al.*, *J. Biol. Chem.*, 276:19452-19460, 2001). Using this as an example, it is clear that the receptor encoding sequence itself may be modified, so as to increase the affinity of the membrane bound protein, such as the receptor, with the protein to which it binds. Exemplary of such modifications are modifications of the C-terminal region of the membrane bound protein, e.g., receptor, such as those described *supra*, which involve replacing a portion of it with a corresponding region of another receptor, which has higher affinity for the binding protein, but does not impact the receptor function. Examples 16 and 20, *supra*, show embodiments of this feature of the invention.

[0095] In addition, the second test protein may be modified to enhance its interaction with the first test protein. For example, the assay may incorporate point mutants, truncations or other variants of the second test protein, e.g., arrestin that are known to bind agonist-occupied GPCRs more stably or in a phosphorylation-independent manner (Kovoor, *et al.*, J. Biol. Chem., 274:6831-6834, 1999).

III. Assay Formats

[0096] As discussed above, the present invention, in one embodiment, offers a straightforward way to assess the interaction of two test proteins when expressed in the same cell. A first construct, as described supra, comprises a sequence encoding a first protein, concatenated to a sequence encoding a cleavage site for a protease or protease portion, which is itself concatenated to a sequence encoding a reporter gene activator. By "concatenated" is meant that the sequences described are fused to produce a single, intact open reading frame, which may be translated into a single polypeptide which contains all the elements. These may, but need not be, separated by additional nucleotide sequences which may or may not encode additional proteins or peptides. A second construct inserted into the recombinant cells is also as described supra, i.e., it contains both a sequence encoding a second protein, and the protease or protease portion. Together, these elements constitute the basic assay format when combined with a candidate agent whose effect on target protein interaction is sought.

[0097] However, the invention may also be used to assay more than one membrane bound protein, such as a receptor, simultaneously by employing different reporter genes, each of which is stimulated by the activation of a protein, such as the classes of proteins described herein. For example, this may be accomplished by mixing cells transfected with different receptor constructs and different reporter genes, or by fusing different transcription factors to each test receptor, and measuring the activity of each reporter gene upon treatment with the test compound. For example, it may be desirable to determine if a molecule of interest activates a first receptor and also determine if side effects should be expected as a result of interaction with a second receptor. In such a case one may, e.g., involve a first cell line encoding a first receptor and a first reporter, such as lacZ, and a second cell line encoding a second receptor and a second reporter, such as GFP. Preferred embodiments of

such a system are seen in Examples 17 and 18. One would mix the two cell lines, add the compound of interest, and look for a positive effect on one, with no effect on the other.

[0098] It is contemplated that the invention relates both to assays where a single pair of interacting test proteins is examined, but more preferably, what will be referred to herein as "multiplex" assays are used. Such assays may be carried out in various ways, but in all cases, more than one pair of test proteins is tested simultaneously. This may be accomplished, e.g., by providing more than one sample of cells, each of which has been transformed or transfected, to test each interacting pair of proteins. The different transformed cells may be combined, and tested simultaneously, in one receptacle, or each different type of transformant may be placed in a different well, and then tested.

[0099] The cells used for the multiplex assays described herein may be, but need not be, the same. Similarly, the reporter system used may, but need not be, the same in each sample. After the sample or samples are placed in receptacles, such as wells of a microarray, one or more compounds may be screened against the plurality of interacting protein pairs set out in the receptacles.

[00100] The fusion proteins expressed by the constructs are also a feature of the invention. Other aspects of the invention which will be clear to the artisan, are antibodies which can identify the fusion proteins as well as various protein based assays for determining the presence of the protein, as well as hybridization assays, such as assays based on PCR, which determine expression of the gene.

IV. Kits

[00101] Any of the compositions described herein may be comprised in a kit. The kits will thus comprise, in suitable container means for the vectors or cells of the present invention, and any additional agents that can be used in accordance with the present invention.

[00102] The kits may comprise a suitably aliquoted compositions of the present invention. The components of the kits may be packaged either in aqueous media or in lyophilized form. The container means of the kits will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which a component may be placed, and preferably, suitably aliquoted. Where there are more than one component in the kit, the

kit also will generally contain a second, third or other additional container into which the additional components may be separately placed. However, various combinations of components may be comprised in a vial. The kits of the present invention also will typically include a means for containing reagent containers in close confinement for commercial sale. Such containers may include injection or blow-molded plastic containers into which the desired vials are retained.

[00103] When the components of the kit are provided in one and/or more liquid solutions, the liquid solution is an aqueous solution, with a sterile aqueous solution being particularly preferred. However, the components of the kit may be provided as dried powder(s). When reagents and/or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container means.

V. Examples

[00104] Specific embodiments describing the invention will be seen in the examples which follow, but the invention should not be deemed as limited thereto.

EXAMPLE 1

[00105] A fusion construct was created, using DNA encoding human $\beta 2$ adrenergic receptor, referred to hereafter as "ADRB2", in accordance with standard nomenclature. Its nucleotide sequence can be found at GenBank, under Accession Number NM_000024 (SEQ ID NO: 1). The tetracycline controlled transactivator tTA, described by Gossen, *et al.*, Proc. Natl. Acad. Sci. USA, 87:5547-5551 (1992), incorporated by reference, was also used. A sequence encoding the recognition and cleavage site for tobacco etch virus nuclear inclusion A protease, described by Parks, *et al.*, Anal. Biochem., 216:413-417 (1994), incorporated by reference, is inserted between these sequences in the fusion coding gene. The CMV promoter region was placed upstream of the ADRB2 coding region, and a poly A sequence was placed downstream of the tTA region.

[00106] A fusion construct was prepared by first generating a form of ADRB2 which lacked internal BamHI and BglII restriction sites. Further, the endogenous stop codon was replaced with a unique BamHI site.

[00107] Overlapping PCR was used to do this. To elaborate, a 5' portion of the coding region was amplified with:

gattgaagat ctgccttctt gctggc (SEQ ID NO: 2),

and

gcagaacttg gaagacctgc ggagtcc (SEQ ID NO: 3),

while a 3' portion of the coding region was amplified with:

ggactcogca ggtcttccaa gttctgc (SEQ ID NO: 4),

and

ttcggatcct agcagtgagt catttgt (SEQ ID NO: 5).

[00108] The resulting PCR products have 27 nucleotides of overlapping sequence and were purified via standard agarose gel electrophoresis. These were mixed together, and amplified with SEQ ID NO: 2, and SEQ ID NO: 5.

[00109] PCR was also used to modify the coding region of tTA so that the endogenous start codon was replaced with a TEV Nla-Pro cleavage site. The cleavage site, defined by the seven amino acid sequence ENLYFQS (SEQ ID NO: 6), is taught by Parks, *et al.*, Anal. Biochem., 216:413-417 (1994), incorporated by reference. The seventh amino acid is known as P1' position, and replacing it with other amino acids is known to reduce the efficiency of cleavage by TEV Nla-Pro. See Kapust, *et al.*, Biochem. Biophys. Res. Commun., 294:949-955 (2002).

[00110] Variants where the seventh amino acid was changed to Tyr, and where it was changed to Leu, were produced. These resulted in intermediate and low efficiency cleavage sites, as compared to the natural high efficiency site.

[00111] A DNA sequence encoding the natural high efficiency site was added to the tTA coding region in two steps. Briefly, BamHI and XbaI restriction sites were added to the 5' end and a XhoI restriction site was added to the 3' end of the tTA coding region by PCR with

ccggatcctc tagattagat aaaagtaaag tg (SEQ ID NO: 7)

and

gactcgagct agcagtatcc tcgcccccc taccc (SEQ ID NO: 8),

and the TEV Nla-Pro cleavage site was added to the 5' end by ligating an oligonucleotide with the sequence

gagaacctgt acttccag (SEQ ID NO: 9)

between the BamHI and XbaI sites.

[00112] This DNA sequence was modified to encode the intermediate and low efficiency cleavage sites by PCR using:

ggatccgaga acctgtactt ccagtacaga tta (SEQ ID NO: 10),

and

ctcgagagat cctcgcgccc cctaccacc (SEQ ID NO: 11).

for ENLYFQY (SEQ ID NO: 12), and

ggatccgaga acctgtactt ccagtaaga tta (SEQ ID NO: 13),

and

ctcgagagat cctcgcgccc cctaccacc (SEQ ID NO: 11) for ENLYFQL (SEQ ID NO: 14).

[00113] These PCR steps also introduced a BamHI restriction site 5' to the sequence encoding each cleavage site, and an XhoI restriction site 3' to tTA stop codon.

[00114] The thus modified ADRB2 coding region was digested with PstI, which cuts at nucleotide position 260 in the coding region, and BamHI. This 3' fragment was ligated with the three variants of tTA modified with the TEV N1a-Pro cleavage sites, that had been digested with BamHI and XhoI, and the resulting complexes were cloned into pBlueScript II, which had been digested with PstI and XhoI.

[00115] A NotI restriction site was introduced 5' to the start codon of the ADRB2 coding region, again via PCR, using

gcggccgcca ccatgaacgg taccgaaggc cca (SEQ ID NO: 15),

and

ctggtgggtg gcccggtacc a (SEQ ID NO: 16).

[00116] The 5' fragment of modified ADRB2 coding region was isolated, via digestion with NotI and PstI and was ligated into each of the constructs of the 3' fragment of ADRB2-TEV-N1a-Pro-cleavage site tTA fusions that had been digested previously, to produce three, full length constructs encoding fusion proteins.

[00117] Each construct was digested with NotI and XhoI, and was then inserted into the commercially available expression vector pcDNA 3, digested with NotI and XhoI.

EXAMPLE 2

[00118] A second construct was also made, whereby the coding sequence for “ β arrestin 2 or ARRB2” hereafter (GenBank, NM_004313) (SEQ ID NO: 17), was ligated to the catalytic domain of the TEV N1a protease (i.e., amino acids 189-424 of mature N1a protease, residues 2040-2279) in the TEV protein. To do this, a DNA sequence encoding ARRB2 was modified, so as to add a BamHI restriction site to its 5' end. Further, the sequence was modified to replace the endogenous stop codon with a BamHI site. The oligonucleotides

caggatcctc tggaatgggg gagaaaccg ggacc (SEQ ID NO: 18),

and

ggatccgcag agttgatcat catagtcgtc (SEQ ID NO: 19)

were used. The resulting PCR product was cloned into the commercially available vector pGEM-T EASY (Promega). The multiple cloning site of the pGEM-T EASY vector includes an EcoRI site 5' to the start codon of ARRB2.

[00119] The TEV N1a-Pro coding region was then modified to replace the endogenous start codon with a BglII site, and to insert at the 3' end a sequence which encodes influenza hemagglutinin epitope YPYDVPDYA (SEQ ID NO: 20) in accordance with Kolodziej, *et al.*, Meth. Enzymol., 194:508-519 (1991), followed by a stop codon, and a NotI restriction site. This was accomplished via PCR, using

agatctagct tgtttaagg accacgtg (SEQ ID NO: 21),

and

gcggccgctc aagcgtaac tggaaatca tatgggtacg agtacaccaa tcattcatg ag
(SEQ ID NO: 22).

[00120] The resulting, modified ARRB2 coding region was digested with EcoRI and BamHI, while the modified TEV coding region was cleaved with BglII and NotI. Both fragments were ligated into a commercially available pcDNA3 expression vector, digested with EcoRI and NotI.

EXAMPLE 3

[00121] Plasmids encoding ADRB2-TEV-N1a-Pro cleavage site-tTA and the ARRB2-TEV-N1a protease fusion proteins were transfected into HEK-293T cells, and into “clone 41,” which is a derivative of HEK-293T, that has a stably integrated β -galactosidase

gene under control of a tTA dependent promoter. About 5×10^4 cells were plated in each well of a 24 well plate, in DMEM medium supplemented with 10% fetal bovine serum, 2mM L-glutamine, 100 units/ml penicillin, 100 μ g/ml G418, and 5 μ g/ml purimycin. Cells were grown to reach 50% confluency the next day, and were then transfected, using 0.4 μ g plasmid DNA, and 2 μ l Fugene (a proprietary transfection reagent containing lipids and other material). The mix was combined in 100 μ l of DMEM medium, and incubated for 15 minutes at room temperature prior to adding cells. Transfected cells were incubated for 8-20 hours before testing by adding drugs which are known agonists for the receptor, and then 16-24 hours after drug addition.

EXAMPLE 4

[00122] The levels of β -galactosidase activity in the cells were first measured by staining the cells with a chromogenic substance, i.e., "X-gal," as taught by MacGregor, *et al.*, Somat. Cell Mol. Genet., 13:253-265 (1987), incorporated by reference. Following culture, cells were washed, twice, in D-PBS with calcium and magnesium, fixed for 5 minutes in 4% paraformaldehyde, and then washed two additional times with D-PBS, calcium and magnesium, for 10 minutes each time. Fixed cells were incubated with 5mM potassium ferricyanide, 5mM potassium ferrocyanide, 2mM $MgCl_2$, 0.1% X-Gal, that had been prepared from a 1:40 dilution of 4% X-Gal stock in dimethylformamide, in D-PBS with calcium and magnesium.

[00123] The reaction was incubated in the dark at room temperature for from 3-4 hours, to overnight. Substrate solution was removed, and cells were mounted under glass coverslips with mowiol mounting medium (10% mowiol, 0.1% 1,4-diazabicyclo[2.2.2]octane, 24% glycerol).

[00124] The results indicated that cells transfected with either the ADRB2-TEV-NIa-Pro cleavage site-tTA plasmid alone or the ARRB2-TEV-NIa protease plasmid alone did not express β -galactosidase. A small fraction of cells transfected with both plasmids did express β -galactosidase, probably due to basal levels of interaction between unstimulated ADRB2 and ARRB2. About 3-5 fold more cells expressed the reporter gene after treatment with either 10 μ M isoproterenol, or 10 μ M epinephrine, both of which are ADRB2 agonists.

[00125] When the cells were pretreated for 5 minutes with the ADRB2 antagonist alprenolol (10 μ M), the agonist induced increase in β -galactosidase expressing cells was blocked, and treatment with alprenolol alone had no apparent effect.

[00126] These results show that one can link agonist binding and GPCR stimulation to transcriptional activation of a reporter gene.

EXAMPLE 5

[00127] A set of experiments were carried out in order to quantify the level of reporter gene activity in the cells more precisely and to maximize the signal-to-background ratio of the assay. This was accomplished by measuring the level of reporter gene induction using a commercially available chemiluminescence assay for β -galactosidase activity. Clone 41 cells were transfected with the ADRB2-tTA fusion constructs, containing either the high, medium or low efficiency cleavage sites, and the ARRB2-TEV-NIa protease expression plasmid described *supra*. Cells were either untreated or treated with 1 μ M isoproterenol 20 hours after the transfection, and the luminescence assay was carried out 24 hours after the drug addition. In brief, following cell culture, the medium was removed, and 50 μ l of lysis buffer (100 mM potassium phosphate, pH7.8, 0.2% Triton X-100) was added to each well. The cells were lysed via incubation for 5 minutes, at room temperature, with mild agitation. Lysates were collected and analyzed via commercially available products.

[00128] In all cases, treatment with agonist increased levels of β -galactosidase activity. However, the background level of reporter gene activity in untreated cells was lowest with the low efficiency cleavage site, relative to the medium and high efficiency sites. Further, agonist treatment resulted in a 4.8-fold stimulation of reporter gene activity in cells transfected with the low efficiency cleavage site, compared to 2.8-fold for the medium efficiency cleavage site and 1.2-fold for the high efficiency cleavage site. Thus, the highest signal-to-background ratio is obtained by using the low efficiency protease cleavage site.

EXAMPLE 6

[00129] These experiments were designed to verify that the agonist stimulated increase in reporter gene expression is dependent on binding and activation of the receptor by the agonist.

[00130] To do this, variants of the ADRB2-tTA fusion constructs were generated following the protocols *supra*, except each contained a mutant form of the receptor with a single amino acid change from D to S at position 113, which results in a greatly reduced affinity for the agonist isoproterenol. See Strader, *et al.*, *J. Biol. Chem.*, 266:5-8 (1991). Three forms of the mutant receptor-tTA fusion construct with each of the different cleavage sites were formed.

[00131] The levels of β -galactosidase activity were measured in clone 41 cells co-transfected with the ADRB2-tTA fusion constructs containing the D113S point mutation and the ARRB2-TEV-N1a protease expression plasmid described previously. The activity tests were carried out exactly as described, *supra*. The results indicated that the agonist isoproterenol did not stimulate reporter gene expression in cells expressing the mutant ADRB2-tTA fusion constructs.

EXAMPLE 7

[00132] These experiments were designed to examine whether the agonist stimulated increase in reporter gene expression is dependent on fusion of TEV N1a-Pro to ARRB2.

[00133] To do this, the levels of β -galactosidase activity were measured in clone 41 cells co-transfected with the ADRB2-tTA fusion construct containing the low efficiency cleavage site and either the ARRB2-TEV-N1a protease expression plasmid described *supra*, or a control TEV-N1a protease fusion to the SH2 domain of phospholipase C. The activity tests were carried out exactly as described, *supra*. The results indicated that agonist-stimulated increase in reporter gene expression was detected only when the TEV protease was fused to ARRB2 and not when fused to an unrelated polypeptide.

EXAMPLE 8

[00134] These experiments were designed to determine if gene expression is induced selectively by agonists of the target receptor, or if it can be stimulated by other molecules.

[00135] ATP is an agonist for G protein coupled receptors P2Y1 and P2Y2, which are expressed endogenously by HEK-293T cells,

[00136] Experiments were carried out using clone 41 cells which were cotransfected with the ADRB2-tTA fusion construct containing the low efficiency cleavage site and the

arrestin-TEV-NIa protease fusion as described supra, which were treated with isoproterenol, ATP, or untreated. The assays were carried out as described, supra.

[00137] The results indicated that induction of reporter gene activity was specific to activation of target receptor. Stimulation of another GPCR pathway was irrelevant.

EXAMPLE 9

[00138] A set of experiments were carried out using clone 41 cells which were cotransfected with the ADRB2-tTA fusion construct containing the low efficiency cleavage site and the ARRB2-TEV-NIa protease fusion as described supra, which were treated with varying amounts of one of the adrenergic receptor agonists isoproterenol and epinephrine. The assays were carried out as described, supra. The results presented in figure 2a show a dose-response curve for the stimulation of reporter gene expression by these two ligands. Each point represents the mean value obtained from three experiments.

[00139] A set of experiments were carried out as described supra, in which the co-transfected clone 41 cells were pretreated with varying concentrations of the adrenergic receptor antagonist alprenolol for 15 minutes, followed by treatment with 1 μ M epinephrine. The results shown in figure 2b indicate a dose-inhibition curve for this antagonist.

EXAMPLE 10

[00140] A similar set of constructs were made to establish an assay for the G protein coupled arginine vasopressin receptor 2 (AVPR2). The AVPR2 coding region (Genbank Accession Number: NM_000054) (SEQ ID NO: 23) was modified to place an EcoRI site at the 5' end and replace the stop codon with a BamHI site using PCR with the primers

gaattcatgc tcatggcgtc caccac (SEQ ID NO: 24)

and

ggatccgat gaagtgctt tggccag (SEQ ID NO: 25).

[00141] The modified AVPR2 coding region was ligated into the three ADRB2-tTA constructs described supra, which had been cut with EcoRI and BamHI. This replaced the entire coding sequence of the ADRB2 with the coding sequence of AVPR2.

[00142] Clone 41 cells were co-transfected with the AVPR2-tTA fusion construct containing the low efficiency cleavage site and the ARRB2-TEV-NIa protease fusion

described supra, and assays were carried out using varying concentrations (1 pM to 2 μM) of [Arg8] vasopressin, an agonist for AVPR2. The data, presented in figure 3, shows a dose-response curve for this agonist, with an EC50 of 3.3 nM, which agrees with previously published data (Oakley, R., *et. al.*, Assay and Drug Development Technologies, 1:21-30, (2002)). The maximal response resulted in an approximately 40-fold induction of reporter gene expression over the background level.

EXAMPLE 11

[00143] A similar set of constructs were made to establish an assay for the G protein coupled serotonin receptor 1a (HTR1A). The HTR1A coding region (Genbank Accession Number: NM_000524) (SEQ ID NO: 26) was modified to place an EcoRI site at the 5' end and replace the stop codon with a BamHI site using PCR with the primers

gaattcatgg atgtgctcag ccctgg (SEQ ID NO: 27)

and

ggatccctgg cggcagaact tacac (SEQ ID NO: 28).

[00144] The modified HTR1A coding region was ligated into the AVPR2- tTA constructs described supra, which had been cut with EcoRI and BamHI. This replaced the entire coding sequence of AVPR2 with the coding sequence of HTR1A. The resulting construct will be referred to as "HTR1A-tTA" hereafter.

[00145] Clone 41 cells were co-transfected with the HTR1A-tTA fusion construct containing the low efficiency cleavage site and the ARRB2-TEV-N1a protease fusion construct described supra, and assays were carried out using 10 μM 8-hydroxy-DPAT HBr (OH-DPAT), an agonist for the HTR1A, as well as with 10 μM serotonin, a natural agonist for HTR1A. The assays were carried out as described, supra. The maximal response to OH-DPAT resulted in a 6.3-fold induction of reporter gene expression over background level and the maximal response to serotonin resulted in a 4.6-fold induction of reporter gene expression over background level.

EXAMPLE 12

[00146] Similar constructs were made to establish an assay for the G protein coupled m2 muscarinic acetylcholine receptor (CHRM2). The CHRM2 coding region (Genbank

Accession Number: NM_000739) (SEQ ID NO: 29) was modified to place an EcoRI site at the 5' end and replace the stop codon with a BglII site using PCR with the primers

gaattcatga ataactcaac aaactcc (SEQ ID NO: 30)

and

agatctcctt gtagcgcta tgttc (SEQ ID NO: 31).

[00147] The modified CHRM2 coding region was ligated into the AVPR2 - tTA constructs described supra, which had been cut with EcoRI and BamHI. This replaced the entire coding sequence of AVPR2 with the coding sequence of CHRM2.

[00148] Clone 41 cells were co-transfected with the CHRM2-tTA fusion construct containing the high efficiency cleavage site and the ARRB2-TEV-N1a protease fusion described supra, where the ARRB2-protease fusion protein was expressed under the control of the Herpes Simplex Virus thymidine kinase (HSV-TK) promoter, and assays were carried out using 10 μ M carbamylcholine Cl (carbochol), an agonist for CHRM2, as described supra. The maximal response to carbochol resulted in a 7.2-fold induction of reporter gene expression over background.

EXAMPLE 13

[00149] α Constructs were also made to establish an assay for the G protein coupled chemokine (C-C motif) receptor 5 (CCR5). The CCR5 coding region (Genbank Accession Number: NM_000579) (SEQ ID NO: 32) was modified to place Not I site at the 5' end and replace the stop codon with a BamHI site using PCR with the primers

gcgccgcat ggattatcaa gtgtcaagtc c (SEQ ID NO: 33)

and

ggatccctgg cggcagaact tacac (SEQ ID NO: 34).

[00150] The CCR5 coding region was also modified to place a BsaI site at the 5' end which, when cut, leaves a nucleotide overhang which is compatible with EcoRI cut DNA using the primers

ggtctccaat tcatggatta tcaagtgca agt (SEQ ID NO: 35)

and

gacgacagcc aggtacatat c (SEQ ID NO: 36).

[00151] The first modified coding region was cut with ClaI and BamHI and the second was cut with BsaI and ClaI. Both fragments were ligated into the AVPR2 - tTA

constructs described supra, which had been cut with EcoRI and BamHI. This replaced the entire coding sequence of AVPR2 with the coding sequence of CCR5.

[00152] The CCR5-tTA fusion construct containing the low efficiency cleavage site was transfected into "clone 34" cells, which are a derivative of the HEK cell line "clone 41" described supra, but which contain a stably integrated ARRB2-TEV-NIa protease fusion gene under the control of the CMV promoter. Assays were carried out using 1 µg/ml "Regulated on Activation, Normal T-Cell Expressed and Secreted" (RANTES), a known agonist for CCR5. The maximal response to RANTES, measured as described supra resulted in an approximately 40-fold induction of reporter gene expression over the background.

EXAMPLE 14

[00153] Next, a set of constructs were made to establish an assay for the G protein coupled dopamine 2 receptor (DRD2). The DRD2 coding region (Genbank Accession Number: NM_000795) (SEQ ID NO: 37) was modified to place an EcoRI site at the 5' end and replace the stop codon with a BglII site using PCR with the primers

gaattcatgg atccactgaa tctgtcc (SEQ ID NO: 38)

and

agatctgcag tggaggatct tcagg (SEQ ID NO: 39).

[00154] The modified DRD2 coding region was ligated into the AVPR2 - tTA constructs described supra, cut with EcoRI and BamHI. This replaced the entire coding sequence of AVPR2 with the coding sequence of DRD2.

[00155] Clone 41 cells were co-transfected with the DRD2-tTA fusion construct containing the medium efficiency cleavage site and the ARRB2-TEV-NIa protease fusion described supra, and assays were carried out using 10 µM dopamine HCl (dopamine), an agonist for DRD2. Results were measured as in the assays described supra. The maximal response to dopamine resulted in a 2.7-fold induction of reporter gene expression over the background.

EXAMPLE 15

[00156] These experiments were designed to demonstrate enhancements of the assay using arrestin variants that bind agonist-occupied GPCRs more stably. First, a fusion of the TEV NIa protease to β-arrestin-1 (ARRB1) was constructed. The coding region of ARRB1

(Genbank Accession Number: NM_004041) (SEQ ID NO: 40) was modified to place an Asp718 site at the 5' end and replace the stop codon with a BamHI site using PCR with the primers

ggatccatgg ggcacaaagg gacgcgagtg (SEQ ID NO: 41)

and

ggatcctctg ttgtgagct gtggagagcc tgtaccatcc tcctcttc (SEQ ID NO: 42).

[00157] The resulting modified ARRB1 coding region was cut with Asp718 and EcoRI and with EcoRI and BamHI, while the modified TEV N1a-Pro coding region described supra was cut with BglII and NotI. All three fragments were ligated into a commercially available pcDNA3 expression vector, which had digested with Asp718 and NotI.

[00158] Clone 41 cells were co-transfected with the DRD2-tTA fusion construct containing the medium efficiency cleavage site and the ARRB1-TEV-N1a protease fusion, and assays were carried out using 10 μ M dopamine HCl (dopamine), an agonist for the D2 receptor, as described supra. The maximal response to dopamine resulted in a 2.1-fold induction of reporter gene expression over the background.

[00159] Truncation of ARRB1 following amino acid 382 has been reported to result in enhanced affinity for agonist-bound GPCRs, independent of GRK-mediated phosphorylation (Kovoor A., *et. al.*, J. Biol. Chem., 274(11):6831-6834 (1999)). To demonstrate the use of such a "constitutively active" arrestin in the present assay, the coding region of β -arrestin-1 was modified to place an Asp718 site at the 5' end and a BamHI site after amino acid 382 using PCR with SEQ ID NO: 41, supra and

ggatccattt gtgcaagtt ctatgag (SEQ ID NO: 43).

[00160] This results in a an ARRB1 coding region which is 36 amino acids shorter than the full-length coding region. The resulting modified ARRB1 coding region, termed "ARRB1 (Δ 383)", was cut with Asp718 and EcoRI and with EcoRI and BamHI, while the modified TEV N1a-Pro coding region described supra was cut with BglII and NotI. All three fragments were ligated into a commercially available pcDNA3 expression vector, digested with Asp718 and NotI.

[00161] Clone 41 cells were co-transfected with the DRD2-tTA fusion construct containing the medium efficiency cleavage site and the ARRB1 (Δ 383)-TEV-NIa protease fusion, and assays were carried out using 10 μ M dopamine HCl (dopamine), an agonist for the DRD2 receptor, as described *supra*. The maximal response to dopamine resulted in an 8.3-fold induction of reporter gene expression over the background.

[00162] To examine the effect of a comparable truncation of the ARRB2 coding region the coding region of ARRB2 was modified to place an Asp718 site at the 5' end and replaced 81 nucleotides at the 3' end with a BamHI site using PCR with the primers

ggtaccatgg gggagaaacc cgggacc (SEQ ID NO: 44)

and

ggatcctgtg gcatagtgg tacc (SEQ ID NO: 45).

[00163] This results in a ARRB2 coding region which is 27 amino acids shorter than the full-length coding region. The resulting modified ARRB2 coding region was cut with Asp718 and BamHI, while the modified TEV NIa-Pro coding region described *supra* was cut with BglII and NotI. Both fragments were ligated into a commercially available pcDNA3 expression vector, digested with Asp718 and NotI.

[00164] Clone 41 cells were co-transfected with the DRD2-tTA fusion construct containing the medium efficiency cleavage site and the ARRB2 (Δ 383)-TEV-NIa protease fusion, and assays were carried out using 10 μ M dopamine HCl (dopamine), an agonist for the DRD2 receptor, as described *supra*. The maximal response to dopamine resulted in a 2.1-fold induction of reporter gene expression over the background.

[00165] These results, presented in figure 4, demonstrate that DRD2 dopamine receptor assay shows the highest signal-to-background ratio using the arrestin variant ARRB1 (Δ 383).

EXAMPLE 16

[00166] This set of experiments was carried out to demonstrate enhancements of the assay using receptor modifications that are designed to increase affinity for the interacting protein. In this example, the C-terminal tail domain of a test receptor was replaced with the corresponding tail domain from AVPR2, a receptor known to bind arrestins with high affinity. In these examples the fusion junction was made 15-18 amino acids after the

conserved NPXXY motif at the end of the seventh transmembrane helix, which typically corresponds to a position immediately after a putative palmitoylation site in the receptor C-terminus.

[00167] First, PCR was used to produce a DNA fragment encoding the C-terminal 29 amino acids from AVPR2, followed by the low efficiency TEV cleavage site and tTA transcription factor. The fragment was also designed such that the first two amino acids (Ala, A and Arg, R) are encoded by the BssHII restriction site GCGCGC. This was accomplished by amplifying the AVPR2-tTA construct with the low efficiency cleavage site described supra, with the primers

tgtgcgcgcg gacgcacccc acccagcctg ggt (SEQ ID NO: 46)

and

ctcgagagat cctcgcgccc cctaccacc (SEQ ID NO: 11).

[00168] Next, the coding region of the DRD2 was modified to place an EcoRI site at the 5' end and to insert a BssHII site after the last amino acid in the coding region (Cys-443). This was done using PCR with the primers

gaattcatgg atccactgaa tctgtcc (SEQ ID NO: 47)

and

tgtgcgcgcg cagtggagga tcttcaggaa ggc (SEQ ID NO: 48).

[00169] The resulting modified D2 coding region was cut with EcoRI and BssHII and the resulting AVPR2 C-terminal tail-low efficiency cleavage site-tTA fragment was cut with BssHII and BamHI. Both fragments were ligated into the AVPR2-low efficiency cleavage site-tTA construct described supra, cut with EcoRI and BamHI.

[00170] Clone 41 cells were co-transfected with the DRD2-AVPR2 Tail-tTA fusion construct containing the low efficiency TEV cleavage site and the ARRB2-TEV-N1a protease fusion described supra, and assays were carried out using 10 μ M dopamine HCl (dopamine), an agonist for the DRD2 receptor. The maximal response to dopamine resulted in an approximately 60-fold induction of reporter gene expression over the background.

[00171] A construct was made which modified the ADRB2 receptor coding region by inserting an Asp718 site at the 5' end and by placing a BssHII site after Cys-341. This was done using PCR with the primers

gcggccgcca ccatgaacgg taccgaaggc cca (SEQ ID NO: 49)

and

tgatgcgacg cacagaagct cctggaagc (SEQ ID NO: 50).

[00172] The modified ADRB2 receptor coding region was cut with EcoRI and BssHII and the AVPR2 C-terminal tail-low efficiency cleavage site-tTA fragment was cut with BssHII and BamHI. Both fragments were ligated into the AVPR2-low efficiency cleavage site-tTA construct described supra cut, with EcoRI and BamHI. The resulting construct is "ADRB2-AVPR2 Tail-tTA." (Also see published application U.S. 2002/0106379, supra, SEQ ID NO: 3 in particular.)

[00173] Clone 41 cells were co-transfected with the ADRB2-AVPR2 Tail-tTA fusion construct containing the low efficiency TEV cleavage site and the ARRB2-TEV-N1a protease fusion described supra, and assays were carried out using 10 μ M isoproterenol, an agonist for the ADRB2 receptor. The maximal response to isoproterenol resulted in an approximately 10-fold induction of reporter gene expression over the background.

[00174] A construct was made which modified the kappa opioid receptor (OPRK; Genbank Accession Number: NM_000912) (SEQ ID NO: 51) coding region by placing a BssHII site after Cys-345. This was done using PCR with the primers

ggtctacttg atgaattcct ggcc (SEQ ID NO: 52)

and

gcgacgacag aagtcgcca aacaccg (SEQ ID NO: 53)

[00175] The modified OPRK receptor coding region was cut with EcoRI and BssHII and AVPR2 C-terminal tail-low efficiency cleavage site-tTA fragment was cut with BssHII and XhoI. Both fragments were ligated into a plasmid containing the modified OPRK receptor sequence, cloned into pcDNA3.1+ at Asp718 (5') and XhoI (3'), which had been digested with EcoRI and XhoI.

[00176] Clone 41 cells were co-transfected with the OPRK-AVPR2 Tail-tTA fusion construct containing the low efficiency cleavage site and the ARRB2-TEV-N1a protease fusion described supra, and assays were carried out using 10 μ M U-69593, an agonist for the OPRK. The maximal response to U-69593 resulted in an approximately 12-fold induction of reporter gene expression over the background.

EXAMPLE 17

[00177] This experiment was designed to demonstrate the use of the assay to measure the activity of two test receptors simultaneously using a multiplex format.

[00178] Clone 41 cells and "clone 1H10" cells, which are cells of an HEK-293T cell line containing a stable integration of the luciferase gene under the control of a tTA-dependent promoter, were each plated on 24-well culture dishes and were transiently transfected with the chimeric ADRB2-AVPR2 Tail-tTA or the DRD2-AVPR2 Tail-tTA fusion constructs described *supra*, respectively. Transient transfections were performed using 100µl of media, 0.4µg of DNA and 2µl of FuGene reagent per well. After 24 hr of incubation, Clone 41 cells expressing ADRB2-AVPR2 Tail-tTA and clone 1H10 cells expressing DRD2-AVPR2 Tail-tTA were trypsinized, mixed in equal amounts, and replated in 12 wells of a 96-well plate. Triplicate wells were incubated without drug addition or were immediately treated with 1µM isoproterenol, 1µM dopamine, or a mixture of both agonists at 1µM. Cells were assayed for reporter gene activity approximately 24 hours after ligand addition. Medium was discarded, cells were lysed in 40µl lysis buffer [100 mM potassium phosphate pH 7.8, 0.2% Triton X-100] and the cell lysate was assayed for beta-galactosidase and for luciferase activity using commercially available luminescent detection reagents.

[00179] The results are presented in Figure 5A and 5B. Treatment with isoproterenol resulted in an approximately seven-fold induction of beta-galactosidase reporter gene activity, whereas luciferase activity remained unchanged. Treatment with dopamine resulted in a 3.5-fold induction of luciferase activity, while beta-galactosidase activity remained unchanged. Treatment with both isoproterenol and dopamine resulted in seven-fold and three-fold induction of beta-galactosidase and luciferase activity, respectively.

EXAMPLE 18

[00180] This experiment was designed to demonstrate the use of the assay to measure the activity of two test receptors simultaneously using a multiplex format.

[00181] "Clone 34.9" cells, which are a derivative of clone 41 cells and containing a stably integrated ARRB2 - TEV N1a protease fusion protein gene, were transiently transfected with the chimeric OPRK-AVPR2 Tail- TEV-N1a-Pro cleavage (Leu)-tTA fusion construct described *supra*. In parallel, "clone HTL 5B8.1" cells, which are an HEK-293T

cell line containing a stable integrated luciferase gene under the control of a tTA-dependent promoter, were transiently transfected with the ADRB-AVPR2 Tail-TEV-N1a-Pro cleavage (Leu)-tTA fusion construct described supra. In each case 5×10^5 cells were plated in each well of a 6-well dish, and cultured for 24 hours in DMEM supplemented with 10% fetal bovine serum, 2 mM L-Glutamine, 100 units/ml penicillin, 500 $\mu\text{g/ml}$ G418, and 3 $\mu\text{g/ml}$ puromycin. Cells were transiently transfected with 100 μl of DMEM, 0.5 μg of OPRK-AVPR2 Tail-TEV-N1a-Pro cleavage (Leu)-tTA DNA, and 2.5 μl Fugene ("clone 34.9 cells") or with 100 μl of DMEM, 0.5 μg of ADRB2-AVPR2 Tail-TEV-N1a-Pro cleavage (Leu)-tTA DNA, 0.5 μg of ARRB2-TEV N1a Protease DNA and 5 μl Fugene ("clone HTL 5B8.1 cells"). Transiently transfected cells were cultured for about 24 hours, and were then trypsinized, mixed in equal amounts and replated in wells of a 96 well plate. Cell were incubated for 24 hours before treatment with 10 μM U-69593, 10 μM isoproterenol or a mixture of both agonists at 10 μM . Sixteen wells were assayed for each experimental condition. After 24 hours, cells were lysed and the activity of both beta-galactosidase and luciferase reporter genes were assayed as described supra. The results are presented in Figure 6. Treatment with U-69593 resulted in an approximately 15-fold induction of beta-galactosidase reporter gene activity, whereas luciferase activity remained unchanged. Treatment with isoproterenol resulted in a 145-fold induction of luciferase activity, while beta-galactosidase activity remained unchanged. Treatment with both U-69593 and isoproterenol resulted in nine-fold and 136-fold induction of beta-galactosidase and luciferase activity, respectively.

EXAMPLE 19

[00182] This experiment was carried out to demonstrate the use of a different transcription factor and promoter in the assay of the invention.

[00183] A fusion construct was created, comprising DNA encoding AVPR2, fused in frame to a DNA sequence encoding the amino acid linker GSENLVYFQLR (SEQ ID NO: 54) which included the low efficiency cleavage site for TEV N1a-Pro described supra, fused in frame to a DNA sequence encoding amino acids 2-147 of the yeast GAL4 protein (GenBank Accession Number P04386) (SEQ ID NO: 55) followed by a linker, i.e., of the sequence PELGSASAELTMVF (SEQ ID NO: 56), followed by amino acids 368-549 of the

murine nuclear factor kappa-B chain p65 protein (GenBank Accession Number A37932) (SEQ ID NO: 57). The CMV promoter was placed upstream of the AVPR2 coding region and a polyA sequence was placed downstream of the GAL4-NFkB region. This construct was designated AVPR2-TEV-N1a-Pro cleavage (Leu)-GAL4.

[00184] HUL 5C1.1 is a derivative of HEK-293T cells, which contain a stably integrated luciferase reporter gene under the control of a GAL4 upstream activating sequence (UAS), commercially available pFR-LUC.

[00185] This AVPR2-TEV-N1a-Pro cleavage (Leu)-GAL4 plasmid was co-transfected along with the β -arrestin2-TEV N1a Protease described supra into HUL 5C1.1 cells. About 2.5×10^4 cells were plated into each well of a 96 well-plate, in DMEM medium supplemented with 10% fetal bovine serum, 2 mM L-Glutamine, 100 units/ml penicillin, 500 μ g/ml G418, and 3 μ g/ml puromycin. Cells were grown to reach 50% confluency the next day and were transfected with 10 μ l per well of a mixture consisting of 85 μ l of DMEM, 0.1 μ g of AVPR2-TEV-N1a-Pro cleavage (Leu)-GAL4 DNA, 0.1 μ g of ARRB2-TEV N1a Protease DNA, and 1 μ l Fugene, which had been incubated for 15 minutes at room temperature prior to addition to the cells. Transfected cells were cultured for about 16 hours before treatment with 10 μ M vasopressin. After six hours, cells were lysed and luciferase activity was assayed as described supra. Under these conditions, treatment with vasopressin resulted in a 180-fold increase in reporter gene activity.

EXAMPLE 20

[00186] This set of experiments were carried out to demonstrate enhancements of the assay using further receptor modifications that are designed to increase the affinity for the interacting protein. In this example, the C-terminal tail domain of the test receptor is replaced with the corresponding tail domain of one of the following receptors: apelin J receptor - AGTRL1 (accession number: NM_005161) (SEQ ID NO: 58), gastrin-releasing peptide receptor - GRPR (accession number: NM_005314) (SEQ ID NO: 59), proteinase-activated receptor 2 - F2RL1 (accession number: NM_005242) (SEQ ID NO: 60), CCR4 (accession number: NM_005508) (SEQ ID NO: 61), chemokine (C-X-C motif) receptor 4 - CXCR4 (accession number: NM_003467) (SEQ ID NO: 62), and interleukin 8 receptor, beta - CXCR2/IL8b (accession number: NM_001557) (SEQ ID NO: 63).

[00187] First PCR was used to produce a DNA fragment encoding the C-terminal tail of the above receptors. These fragments were designed such that the first two amino acids (Ala, A and Arg, R) are encoded by the BssHII restriction site.

[00188] The AGTRL1 C-terminal fragment was amplified with the primers

tgtgcgcgcg gccagagcag gtgcgca (SEQ ID NO: 64)

and

gaggatccgt caaccacaag ggtctc (SEQ ID NO: 65).

[00189] The GRPR C-terminal fragment was amplified with the primers

tgtgcgcgcg gcctgatcat ccggtct (SEQ ID NO: 66)

and

gaggatccga cataccgctc gtgaca (SEQ ID NO: 67).

[00190] The F2RL1 C-terminal fragment was amplified with the primers

tgtgcgcgca gtgtccgcac tgtaaagc (SEQ ID NO: 68)

and

gaggatccat aggaggtctt aacagt (SEQ ID NO: 69).

[00191] The CCR4 C-terminal fragment was amplified with the primers

tgtgcgcgcg gcccttttgt gctctgc (SEQ ID NO: 70)

and

gaggatccca gagcatcatg aagatc (SEQ ID NO: 71).

[00192] The CXCR2/IL8b C-terminal fragment was amplified with the primers

tgtgcgcgcg gcttgatcag caaggac (SEQ ID NO: 72)

and

gaggatccga gagtagtgga agtgtg (SEQ ID NO: 73).

[00193] The CXCR4 C-terminal fragment was amplified with the primers

tgtgcgcgcg ggtccagcct caagatc (SEQ ID NO: 74)

and

gaggatccgc tggagtgaaa acttga (SEQ ID NO: 75).

[00194] The resulting DNA fragments encoding the modified C-terminal tail domains of these receptors were cut with BssHII and BamHI and the fragments were ligated in frame to the OPRK receptor coding region, replacing the AVPR2- C-terminal tail fragment, in the OPRK-AVPR2 Tail- TEV-NIa-Pro cleavage (Leu)-tTA expression construct described supra.

[00195] HTL 5B8.1 cells described supra were co-transfected with each of the above modified OPRK coding region – TEV-NIa-Pro cleavage (Leu)- tTA constructs and the β -arrestin 2 – TEV NIa protease fusion described supra. About 2.5×10^4 cells per well were plated onto a 96 well-plate, in DMEM medium supplemented with 10% fetal bovine serum, 2 mM L-Glutamine, 100 units/ml penicillin, 500 $\mu\text{g/ml}$ G418, and 3 $\mu\text{g/ml}$ puromycin. Cells were grown to reach 50% confluency the next day and were transfected with 10 μl per well of a mixture consisting of 85 μl of DMEM, 0.25 μg of AVPR2-TEV-NIa-Pro cleavage (Leu)-GAL4 DNA, 0.25 μg of ARRB2-TEV NIa protease DNA, and 2.5 μl Fugene (a proprietary transfection reagent containing lipids and other material), which had been incubated for 15 minutes at room temperature prior to addition to the cells. Transfected cells were cultured for about 16 hours before treatment 10 μM U-69593. After six hours, cells were lysed and luciferase activity was assayed as described supra. Under these conditions, treatment with U-69593 resulted in the following relative increases in reporter gene activity for each of the modified OPRK receptors: OPRK-AGTRL1 C-terminal tail – 30 fold; OPRK-GRPR C-terminal tail – 312 fold; OPRK-F2RL1 C-terminal tail – 69.5 fold; OPRK-CCR4 C-terminal tail – 3.5 fold; OPRK-CXCR4 C-terminal tail – 9.3 fold; OPRK-IL8b C-terminal tail – 113 fold.

EXAMPLE 21

[00196] This experiment was designed to produce a cell line that stably expressed the ARRB2-TEV NIa protease fusion protein described supra.

[00197] A plasmid was made which expressed the ARRB2-TEV NIa protease fusion protein under the control of the EF1 α promoter and also expressed the hygromycin resistance gene under the control of the thymidine kinase (TK) promoter.

[00198] This plasmid was transfected into HTL 5B8.1, and clones containing a stable genomic integration of the plasmid were selected by culturing in the presence of 100 $\mu\text{g/ml}$ hygromycin. Resistant clones were isolated and expanded and were screened by transfection of the ADRB2-AVPR2 Tail- TEV-NIa-Pro cleavage (Leu)-tTA plasmid described supra. Three cell lines that were selected using this procedure were designated “HTLA 4C2.10”, “HTLA 2C11.6” and “HTLA 5D4”. About 2.5×10^4 cells per well were plated onto a 96 well-plate, in DMEM medium supplemented with 10% fetal bovine serum, 2

mM L-Glutamine, 100 units/ml penicillin, 500 μ g/ml G418, 3 μ g/ml puromycin, and 100 μ g/ml hygromycin. Cells were grown to reach 50% confluency the next day and were transfected with 10 μ l per well of a mixture consisting of 85 μ l of DMEM, 0.25 μ g of ADRB2-AVPR2-TEV-NIa-Pro cleavage (Leu)-GAL4 DNA and 0.5 μ l Fugene, which had been incubated for 15 minutes at room temperature prior to addition to the cells. Transfected cells were cultured for about 16 hours before treatment 10 μ M isoproterenol. After six hours, cells were lysed and luciferase activity was assayed as described supra. Under these conditions, treatment with isoproterenol resulted in a 112-fold ("HTLA 4C2.10"), 56-fold ("HTLA 2C11.6") and 180-fold ("HTLA 5D4") increase in reporter gene activity in the three cell lines, respectively.

EXAMPLE 22

[00199] This experiment was designed to produce a cell line that stably expressed the ARRB2-TEV NIa protease and the ADRB2-AVPR2 Tail- TEV-NIa-Pro cleavage (Leu)-tTA fusion proteins described supra.

[00200] The ARRB2-TEV NIa protease plasmid containing the hygromycin resistance gene was transfected together with the ADRB2-AVPR2 Tail- TEV-NIa-Pro cleavage (Leu)-tTA fusion protein plasmid described supra into HTL 5B8.1 cells and clones containing stable genomic integration of the plasmids were selected by culturing in the presence of 100 μ g/ml hygromycin. Resistant clones were isolated and expanded, and were screened by treating with 10 μ M isoproterenol and measuring the induction of reporter gene activity as described supra. Three cell lines that were selected using this procedure were designated "HTLAR 1E4", "HTLAR 1C10" and "HTLAR 2G2". Treatment with isoproterenol for 6 hours resulted in a 208-fold ("HTLAR 1E4"), 197-fold ("HTLAR 1C10") and 390-fold ("HTLAR 2G2") increase in reporter gene activity in the three cell lines, respectively.

EXAMPLE 23

[00201] This experiment was designed to demonstrate the use of the assay to measure the activity of the receptor tyrosine kinase epidermal growth factor receptor (EGFR).

[00202] A first fusion construct was created, comprising DNA encoding the human EGFR, which can be found at GenBank under the Accession Number NM_005228 (SEQ ID NO: 76), fused in frame to a DNA sequence encoding amino acids 3-335 of the tetracycline-controlled transactivator tTA, described supra. Inserted between these sequences is a DNA sequence encoding the amino acid sequence GGSGSENLYFQL (SEQ ID NO: 77) which includes the low efficiency cleavage site for TEV N1a-Pro, ENLYFQL (SEQ ID NO: 14), described supra. The CMV promoter was placed upstream of the Epidermal Growth Factor Receptor coding region, and a polyA sequence was placed downstream of the tTA region. This construct is designated EGFR-TEV-N1a-Pro cleavage (Leu)-tTA.

[00203] A second fusion construct was created, comprising DNA encoding the two SH2 domains of human Phospholipase C Gamma 1, corresponding to amino acids 538-759 (GeneBank accession number NP_002651.2) (SEQ ID NO: 78) fused in frame to a DNA sequence encoding the catalytic domain of mature TEV N1a protease, described supra, corresponding to amino acids 2040-2279 (GeneBank accession number AAA47910) (SEQ ID NO: 79). Inserted between these sequences is a linker DNA sequence encoding the amino acids NSSGGNSGS (SEQ ID NO: 80). The CMV promoter was placed upstream of the PLC-Gamma SH2 domain coding sequence and a polyA sequence was placed downstream of the TEV N1a protease sequence. This construct is designated PLC Gamma1-TEV.

[00204] The EGFR-TEV-N1a-Pro cleavage (Leu)-tTA and PLC Gamma1-TEV fusion constructs were transfected into clone HTL5B8.1 cells described supra. About 2.5×10^4 cells were plated into each well of a 96 well-plate, in DMEM medium supplemented with 10% fetal bovine serum, 2 mM L-Glutamine, 100 units/ml penicillin, 500 µg/ml G418, and 3 µg/ml puromycin. Cells were grown to reach 50% confluency the next day and were transfected with 15 µl per well of a mixture consisting of 100 µl of DMEM, 0.4 µg of pcDNA3 DNA ("carrier" vector DNA), 0.04 µg of EGFR-TEV-N1a-Pro cleavage (Leu)-tTA DNA, 0.04 µg of PLC Gamma1-TEV DNA, and 2 µl Fugene (a proprietary transfection reagent containing lipids and other material), which had been incubated for 15 minutes at room temperature prior to addition to the cells. Transfected cells were cultured for about 16 hours before treatment with specified receptor agonists and inhibitors. After six hours, cells were lysed and luciferase activity was assayed as described supra. Results are shown in Figure 7.

[00205] The addition of 2.5 ng/ml human Epidermal Growth Factor (corresponding to the EC80 for this ligand) resulted in a 12.3 fold increase of luciferase reporter gene activity, while addition of 100 ng/ml human Transforming Growth Factor - Alpha resulted in an 18.3 fold increase. Prior treatment with tyrosine kinase inhibitors (70 μ M AG-494; 0.3 μ M AG-1478; 2 mM RG-130022) before addition of human Epidermal Growth Factor blocked the induction of reporter gene activity.

EXAMPLE 24

[00206] This experiment was designed to demonstrate the use of the assay to measure the activity of the human Type I Interferon Receptor.

[00207] A fusion construct was created, comprising DNA encoding human Interferon Receptor I (IFNAR1) (557 amino acids), which can be found in Genbank under Accession Number NM_000629 (SEQ ID NO: 81), fused in frame to a DNA sequence encoding amino acids 3-335 of the tetracycline controlled transactivator tTA, described supra. Inserted between these sequences is a DNA sequence encoding the amino acid sequence GSENLVYFQL (SEQ ID NO: 82) which includes the low efficiency cleavage site for TEV NIa-Pro, ENLVYFQL (SEQ ID NO: 14), described supra. The CMV promoter was placed upstream of the Human Interferon Receptor I (IFNAR1) coding region, and a poly A sequence was placed downstream of the tTA region. This construct is designated IFNAR1-TEV- NIa-Pro cleavage (L)-tTA.

[00208] A second fusion construct was created, using DNA encoding Human Interferon Receptor 2, splice variant 2 (IFNAR2.2) (515 amino acids), which can be found at Genbank, under Accession Number L41942 (SEQ ID NO: 83), fused in frame to a DNA sequence encoding the catalytic domain of the TEV NIa protease, described supra, corresponding to amino acids 2040-2279 (GenBank accession number AAA47910) (SEQ ID NO: 84). Inserted between these sequences is a DNA sequence encoding the amino acid sequence RS (Arg-Ser). The CMV promoter region was placed upstream of the Human Interferon Receptor 2 (IFNAR2.2) coding region, and a poly A sequence was placed downstream of the TEV region. This construct is designated IFNAR2.2-TEV.

[00209] Expression constructs were also generated in which the genes for Human Signal Transducer and Activator of Transcription 1 (STAT1), found in Genbank, under

Accession Number NM_007315 (SEQ ID NO: 85), Human Signal Transducer and Activator of Transcription 2 (STAT2) found in Genbank, under Accession Number NM_005419 (SEQ ID NO: 86), were expressed under the control of the CMV promoter region. These constructs were designated CMV-STAT1 and CMV-STAT2 respectively.

[00210] The IFNAR1- TEV- N1a-Pro cleavage (L)-tTA and IFNAR2.2-TEV fusion constructs, together with CMV-STAT1 and CMV-STAT2 were transiently transfected into HTL5B8.1 cells described supra. About 2.5×10^4 cells were seeded in each well of a 96 well plate and cultured in DMEM medium supplemented with 10% fetal bovine serum, 2mM L-glutamine, 100 units/ml penicillin, 100 μ g/ml G418, and 5 μ g/ml puromycin. After 24 hours of incubation, cells were transfected with 15ng of each IFNAR1- TEV- N1a-Pro cleavage (L)-tTA, IFNAR2.2-TEV, CMV-STAT1 and CMV-STAT2 DNA, or with 60ng control pcDNA plasmid, together with 0.3 μ l Fugene per well. Transfected cells were cultured for 8-20 hours before treatment with 5000 U/ml human interferon-alpha or 5000 U/ml human interferon-beta. At the time of interferon addition, medium was aspirated and replaced with 293 SFM II media supplemented with 2mM L-glutamine, 100 units/ml penicillin, 3 μ g/ml puromycin and 500 μ g/ml of G418. Interferon-treated cells were cultured for an additional 18-20 hours before they were assayed for luciferase reporter gene activity as described supra. Results are shown in Figure 8. Treatment with 5000U/ml IFN- α resulted in 15-fold increase in reporter gene activity, while treatment with 5000U/ml IFN- β resulted in a 10-fold increase. Interferon treatment of HTL5B8.1 cells transfected with the control plasmid pcDNA3 had no effect on reporter gene activity. Figure 9 shows a dose-response curve generated for IFN- α in HTL5B8.1 cells transfected with IFNAR1(ENLYFQ(L)-tTa, IFNAR2.2-TEV, STAT1 and STAT2 expression constructs as described supra.

EXAMPLE 25

[00211] This experiment was designed to demonstrate the use of the assay to measure the activity of the human Type I Interferon Receptor using a different transcription factor and a different cell line.

[00212] A fusion construct was created, using DNA encoding Human Interferon Receptor I (IFNAR1), fused in frame to a DNA sequence encoding the GAL4-NF- κ B-fusion, described supra. Inserted between these sequences is a DNA sequence encoding the amino

acid sequence GSENLVYFQL (SEQ ID NO: 87), which includes the low efficiency cleavage site for TEV NIa-Pro, ENLYFQL (SEQ ID NO: 14), described *supra*. The CMV promoter was placed upstream of the Human Interferon Receptor I (IFNAR1) coding region, and a poly A sequence was placed downstream of the GAL4-NF- κ B region. This construct is designated IFNAR1-TEV- NIa-Pro cleavage (L)-GAL4-NF- κ B.

[00213] CHO-K1 cells were then transiently transfected with a mixture of five plasmids: IFNAR1-TEV-NIa-Pro cleavage (L)-GAL4-NF- κ B, IFNAR2.2-TEV, CMV-STAT1, CMV-STAT2 and pFR-Luc, a luciferase reporter gene plasmid under the control of a GAL4-dependent promoter. About 1.0×10^4 cells per well were seeded in a 96 well plate 24 hours prior to transfections in DMEM medium supplemented with 10% fetal bovine serum, 2mM L-glutamine, 100 units/ml penicillin. Cells were transfected the following day with 10 ng of reporter plasmid (pFR-Luc), plus 20 ng of each of the expression constructs described *supra*, or with 10 ng reporter plasmid plus 80 ng of control pcDNA3 plasmid, together with 0.3 μ l Fugene per well. Transfected cells were cultured for 8-20 hours before treatment with 5000 U/ml human interferon-alpha. At the time of interferon addition, medium was aspirated and replaced with DMEM media supplemented with 2mM L-glutamine, 100 units/ml penicillin. Interferon-treated cells were cultured for an additional 6 hours before they were assayed for luciferase reporter gene activity as described *supra*. Results are shown in Figure 10. IFN- α treatment of CHO-K1 cells transfected with the reporter, IFNAR and STAT constructs resulted in 3-fold increase in reporter gene activity, while interferon treatment of cells transfected with the reporter and control plasmids had no effect on reporter gene activity.

EXAMPLE 26

[00214] This set of experiments was carried out to demonstrate additional enhancements of the assay using receptor modifications designed to increase the affinity of the test receptor for the interacting protein. In these examples, the fusion junction between the test receptor and a C-terminal tail domain of GRPR (Genbank Accession Number: NM_005314) (SEQ ID NO: 59) was made 17-23 amino acids after the conserved NPXXY motif at the end of the seventh transmembrane helix.

[00215] First, PCR was used to produce a DNA fragment encoding the C-terminal 42 amino acids from GRPR beginning 2 amino acids after the putative palmitoylation site (hereafter referred to as GRPR 42aa). The fragment was designed such that the first amino acid of the C-terminal tail is preceded by two amino acids (Ser, S and Arg, R) which are encoded by the XbaI restriction site TCTAGA, and the stop codon is replaced by two amino acids (Gly, G and Ser, S) which are encoded by a BamHI restriction site GGATCC. This was accomplished by amplifying a plasmid containing the GRPR coding region with primers

tctagaggcctgatcatccggtctcac (SEQ ID NO: 88)

and

gaggatccgacataccgctcgtgaca (SEQ ID NO: 67)

[00216] Next the coding region of OPRK (Genbank Accession Number: NM_000912) (SEQ ID NO: 51) was modified to place insert an XbaI site after Pro-347. This was done using PCR with the primers

ggtctacttgatgaattcctggcc (SEQ ID NO: 52)

and

tctagatggaaaacagaagtcccggaaac (SEQ ID NO: 89)

[00217] In addition, the coding region of ADRA1A (Genbank Accession Number: NM_000680) (SEQ ID NO: 90) was modified to insert an XbaI site after Lys-349. This was done using PCR with the primers

ctcgatatctaaacagctgcatcaa (SEQ ID NO: 91)

and

tctagactttctgcagagacactggattc (SEQ ID NO: 92)

[00218] In addition, the coding region of DRD2 (Genbank Accession Number: NM_000795) (SEQ ID NO: 37) was modified to insert two amino acids (Leu and Arg) and an XbaI site after Cys-343. This was done using PCR with the primers

gaattcatggatccactgaatctgtcc (SEQ ID NO: 38)

and

tctagatcgaaggcagtgaggatcttcagg (SEQ ID NO: 93)

[00219] The modified OPRK receptor coding region was cut with EcoRI and XbaI and the GRPR 42aa C-terminal tail fragment was cut with XbaI and BamHI. Both fragments were ligated into a plasmid containing the OPRK receptor with the AVPR2 C-terminal tail-low-efficiency cleavage site-tTA described supra which had been digested with EcoRI and BamHI.

[00220] The modified ADRA1A receptor coding region was cut with EcoRV and XbaI and the OPRK-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site was cut with XbaI and XhoI. Both fragments were ligated into a plasmid containing the ADRA1A receptor which had been digested with EcoRV and XhoI.

[00221] The modified DRD2 receptor coding region was cut with EcoRI and XbaI and the OPRK-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site was cut with XbaI and XhoI. Both fragments were ligated into a pcDNA6 plasmid digested with EcoRI and XhoI

[00222] HTLA 2C11.6 cells, described supra, were transfected with OPRK-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site and assays were carried out using 10 μ M U-69593, an agonist for OPRK. The maximal response to U-69593 resulted in an approximately 200-fold increase in reporter gene activity.

[00223] HTLA 2C11.6 cells were transfected with ADRA1A-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site and assays were carried out using 10 μ M epinephrine, an agonist for ADRA1A. The maximal response to epinephrine resulted in an approximately 14-fold increase in reporter gene activity.

[00224] HTLA 2C11.6 cells were transfected with DRD2-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site and assays were carried out using 10 μ M dopamine, an agonist for DRD2. The maximal response to dopamine resulted in an approximately 30-fold increase in reporter gene activity.

EXAMPLE 27

[00225] This set of experiments were carried out to demonstrate further enhancements of the assay using a different set of test receptor modifications designed to increase the affinity for the interacting protein. In these examples, the C-terminal domain of

the test receptor was replaced with a portion of the endogenous C-terminal tail domain of GRPR.

[00226] First, PCR was used to produce a DNA fragment encoding the truncated GRPR tail, specifically a sequence encoding 23 amino acids from Gly-343 to Asn-365. The fragment was designed such that the first amino acid of the C-terminal tail is preceded by two amino acids (Ser, S and Arg, R) which are encoded by the XbaI restriction site TCTAGA, and the Ser-366 is replaced by two amino acids (Gly, G and Ser, S) which are encoded by a BamHI restriction site GGATCC. This was accomplished by amplifying a plasmid containing the GRPR coding region with primers

tctagaggcctgatcatccgggtctcac (SEQ ID NO: 94)

and

cggatccgttgggtactcttgagg (SEQ ID NO: 95)

[00227] Next the truncated GRPR fragment (hereafter referred to as GRPR 23aa Tail) was cut with XbaI and BamHI and inserted into the OPRK-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site described herein, digested with XbaI and BamHI.

[00228] Similarly, the GRPR 23aa Tail fragment was cut with XbaI and BamHI and inserted into the ADRA1A-GRPR 42aa Tail-tTA fusion construct containing the low efficiency cleavage site described herein, digested with XbaI and BamHI.

[00229] HTLA 2C11.6 cells were transfected with OPRK-GRPR 23aa Tail -tTA fusion construct containing the low efficiency cleavage site and assays were carried out using 10 μ M U-69593, an agonist for OPRK. The maximal response to U-69593 resulted in an approximately 115-fold induction of reporter gene expression over the background.

[00230] HTLA 2C11.6 cells were transfected with ADRA1A-GRPR 23aa Tail-tTA fusion construct containing the low efficiency cleavage site and assays were carried out using 10 μ M epinephrine, an agonist for ADRA1A. The maximal response to epinephrine resulted in an approximately 102-fold induction of reporter gene expression over the background.

EXAMPLE 28

[00231] This experiment was designed to demonstrate the use of the assay to measure the activity of the receptor tyrosine kinase Insulin-like Growth Factor-1 Receptor (IGF1R), specifically by monitoring the ligand-induced recruitment of the intracellular signaling protein SHC1 (Src homology 2 domain-containing transforming protein 1).

[00232] A first fusion construct was created, comprising DNA encoding the human IGF-1R, which can be found at GenBank under the Accession Number NM_000875 (SEQ ID NO: 96), fused in frame to a DNA sequence encoding amino acids 3-335 of the tetracycline-controlled transactivator tTA, described supra. Inserted between these sequences is a DNA sequence encoding the amino acid sequence GSENLVYFQL (SEQ ID NO: 82) which includes the low efficiency cleavage site for TEV Nla-Pro, ENLVYFQL (SEQ ID NO: 14), described supra. The CMV promoter was placed upstream of the IGF1R coding region, and a polyA sequence was placed downstream of the tTA region. This construct is designated IGF1R-TEV-Nla-Pro cleavage (Leu)-tTA.

[00233] A second fusion construct was created, comprising DNA encoding the PTB domain of human SHC1, corresponding to amino acids 1-238 (GeneBank accession number BC014158) (SEQ ID NO: 97) fused in frame to a DNA sequence encoding the catalytic domain of mature TEV Nla protease, described supra, corresponding to amino acids 2040-2279 (GeneBank accession number AAA47910) (SEQ ID NO: 79). Inserted between these sequences is a linker DNA sequence encoding the amino acids NSGS (SEQ ID NO: 98). The CMV promoter was placed upstream of the SHC1 PTB domain coding sequence and a polyA sequence was placed downstream of the TEV Nla protease sequence. This construct is designated SHC1-TEV.

[00234] The IGF1R-TEV-Nla-Pro cleavage (Leu)-tTA and SHC1-TEV fusion constructs were transfected into clone HTL5B8.1 cells described supra. About 2.5×10^4 cells were plated into each well of a 96 well-plate, in DMEM medium supplemented with 10% fetal bovine serum, 2 mM L-Glutamine, 100 units/ml penicillin, 500 μ g/ml G418, and 3 μ g/ml puromycin. Cells were grown to reach 50% confluency the next day and were transfected with 15 μ l per well of a mixture consisting of 100 μ l of DMEM, 0.2 μ g of IGF1R-TEV-Nla-Pro cleavage (Leu)-tTA DNA, 0.2 μ g of SHC1-TEV DNA, and 2 μ l

Fugene (a proprietary transfection reagent containing lipids and other material), which had been incubated for 15 minutes at room temperature prior to addition to the cells. Transfected cells were cultured for about 16 hours before treatment with a specific receptor agonist. After 24 hours, cells were lysed and luciferase activity was assayed as described supra.

[00235] The addition of 1 μ M human Insulin-like Growth Factor 1 resulted in a 90 fold increase of luciferase reporter gene activity.

EXAMPLE 29

[00236] This experiment was designed to demonstrate the use of the assay to measure the interaction of two test proteins that are not normally membrane bound. In this example, the assay was used to measure the ligand-induced dimerization of the nuclear steroid hormone receptors, ESR1 (estrogen receptor 1 or ER alpha) and ESR2 (estrogen receptor 2 or ER beta). In this example, ESR1 is fused to the transcription factor tTA, where the cleavage site for the TEV Nla-Pro protease is inserted between the ESR1 and tTA sequences. This ESR1-tTA fusion is tethered to the membrane by a fusion to the intracellular, C-terminal end of the transmembrane protein CD8. CD8 essentially serves as an inert scaffold that tethers ESR1 to the cytoplasmic side of the cell membrane. The transcription factor fused thereto cannot enter the nucleus until interaction with ESR2 and protease. Any transmembrane protein could be used. This CD8-ESR1-TEV Nla Pro cleavage-tTA fusion protein is expressed together with a second fusion protein comprised of ESR2 and the TEV Nla-Pro protease in a cell line containing a tTA-dependent reporter gene. The estrogen-induced dimerization of ESR1 and ESR2 thereby triggers the release of the tTA transcription factor from the membrane bound fusion, which is detected by the subsequent induction in reporter gene activity.

[00237] A fusion construct was created, comprising DNA encoding human CD8 gene (235 amino acids), which can be found in Genbank under Accession Number NM_001768 (SEQ ID NO: 99), fused in frame to a DNA sequence encoding the human ESR1 (596 amino acids), which can be found in Genbank under Accession Number NM_000125 (SEQ ID NO: 100). Inserted between these sequences is a DNA sequence encoding the amino acid sequence GRA (Gly-Arg-Ala). The resulting construct is then fused in frame to a DNA sequence encoding amino acids 3-335 of the tetracycline controlled

transactivator tTA, described supra. Inserted between these sequences is a DNA sequence encoding the amino acid sequence GSENYLFQL (SEQ ID NO: 82) which includes the low efficiency cleavage site for TEV NIa-Pro, ENLYFQL (SEQ ID NO: 14), described supra. The CMV promoter was placed upstream of the Human CD8 coding region, and a poly A sequence was placed downstream of the tTA region. This construct is designated CD8-ESR1-TEV-NIa-Pro cleavage (L)-tTA.

[00238] A second fusion construct was created, using DNA encoding Human Estrogen Receptor beta (ESR2) (530 amino acids), which can be found at Genbank, under Accession Number NM_001437 (SEQ ID NO: 101), fused in frame to a DNA sequence encoding the catalytic domain of the TEV NIa protease, described supra, corresponding to amino acids 2040-2279 (GenBank accession number AAA47910) (SEQ ID NO: 84). Inserted between these sequences is a DNA sequence encoding the amino acid sequence RS (Arg-Ser). The CMV promoter region was placed upstream of the Human Estrogen Receptor beta (ESR2) coding region, and a poly A sequence was placed downstream of the TEV region. This construct is designated ESR2-TEV.

[00239] The CD8-ESR1-TEV-NIa-Pro cleavage (L)-tTA and ESR2-TEV fusion constructs, together with pCDNA3 were transiently transfected into HTL5B8.1 cells described supra. About 2.0×10^4 cells were seeded in each well of a 96 well plate and cultured in phenol-free DMEM medium supplemented with 10% fetal bovine serum, 2mM L-glutamine, 100 units/ml penicillin, 100µg/ml G418, and 5 µg/ml puromycin. After 24 hours of incubation, cells were transfected with a mixture of 5ng of ESR1-TEV-NIa-Pro cleavage (L)-tTA, 15ng of ESR2-TEV and 40ng of pCDNA3, together with 0.3 µl Fugene per well. 6 hours after transfection, the cells were washed with PBS and incubated in 100µl of phenol-free DMEM without serum for 24 hours before treatment with 50nM 17-β Estradiol. Ligand-treated cells were cultured for an additional 18-20 hours before they were assayed for luciferase reporter gene activity as described supra. Treatment with 50nM 17-β Estradiol resulted in a 16-fold increase in reporter gene activity.

[00240] Other features of the invention will be clear to the skilled artisan and need not be reiterated here.

2004257228 02 Dec 2008

- 58 -

The claims defining the invention are as follows:

1. A method for determining if a test compound modulates
a specific protein/protein interaction of interest
5 comprising contacting said compound to a cell which
has been transformed or transfected with

(a) a nucleic acid molecule which comprises:

(i) a nucleotide sequence which encodes a first
test protein which is a transmembrane receptor,

10 (ii) a nucleotide sequence encoding a cleavage
site for a protease or a portion of a protease, and

(iii) a nucleotide sequence which encodes a
protein which activates a reporter gene in said cell,
wherein the nucleotide sequence which encodes a
15 cleavage site is located between the nucleotide
sequence which encodes a transmembrane receptor and
the nucleotide sequence which encodes an activation
domain; and

(b) a nucleic acid molecule which comprises:

20 (i) a nucleotide sequence which encodes a
second test protein whose interaction with said first
test protein, in the presence of said test compound
is to be measured, and

(ii) a nucleotide sequence which encodes a
25 protease or a portion of a protease which is specific
for said cleavage site,

2004257228 02 Dec 2008

- 59 -

and determining activity of said reporter gene as a determination of whether said compound modulates said protein/protein interaction.

2. A method according to claim 1, wherein said
5 transmembrane receptor is β 2-adrenergic receptor (ADRB2), arginine vasopressin receptor 2 (AVPR2), serotonin receptor 1a (HTR1A), m2 muscarinic acetylcholine receptor (CHRM2), chemokine (C-C motif) receptor 5 (CCR5), dopamine D2 receptor (DRD2), kappa
10 opioid receptor (OPRK), or ADRA1A.
3. A method according to claim 2, wherein said transmembrane receptor is a GPCR.
4. A method according to any one of claims 1 to 3,
15 wherein said protease or portion of a protease is tobacco etch virus nuclear inclusion A protease.
5. A method according to any one of claims 1 to 4, wherein said protein which activates said reporter gene is a transcription factor.
6. A method according to claim 5, wherein said
20 transcription factor is tTA or GAL4.
7. A method according to any one of claims 1 to 6, wherein said second protein is an inhibitory protein.
8. A method according to claim 7, wherein said inhibitory protein is an arrestin.
- 25 9. A method according to any one of claims 1 to 8, wherein said cell is a eukaryote.
10. A method according to any one of claims 1 to 9, wherein said reporter gene is an exogenous gene.

2004257228 02 Dec 2008

- 60 -

11. A method according to claim 10, wherein said exogenous gene encodes β -galactosidase or luciferase.
12. A method according to any one of claims 1 to 11, wherein the nucleotide sequence encoding said first test protein is modified to increase interaction with said second test protein, wherein said modification comprises replacing all or part of the nucleotide sequence of the C-terminal region of said first test protein with a nucleotide sequence which encodes an amino acid sequence which has higher affinity for said second test protein than the original sequence, wherein the nucleotide sequence of said C-terminal region is replaced by a nucleotide sequence encoding all or a part of the C-terminal region of AVPR2, AGTRLI, GRPR, F2PL1, CXCR2/IL-8B, CCR4, or GRPR.
13. A method according to any one of claims 1 to 12, comprising contacting more than one compound to a plurality of samples of cells, each of said samples being contacted by one or more of said compounds, wherein each of said cell samples have been transformed or transfected with (a) and (b), and determining activity of reporter genes in said plurality of said samples to determine if any of said compounds modulates said specific, protein/protein interaction.
14. A method according to claim 12 or claim 13, comprising contacting each of said samples with one compound, each of which differs from all others.
15. A method according to claim 12 or claim 13, comprising contacting each of said samples with a mixture of said compounds.

N:\Perth\Cases\Patent\59000-59999\59482.AU\Specis\59482.AU Specification 2008-12-1.doc 1/12/08

2004257228 02 Dec 2008

- 61 -

16. A method according to claim 1 for determining if a test compound modulates one or more of a plurality of protein interactions of interest, comprising contacting said test compound to a plurality of samples of cells, each of which has been transformed or transfected with

(a) a first nucleic acid molecule comprising:

(i) a nucleotide sequence which encodes a first test protein which is a transmembrane receptor,

(ii) a nucleotide sequence encoding a cleavage site for a protease, and

(iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, wherein the nucleotide sequence which encodes a cleavage site is located between the nucleotide sequence which encodes a transmembrane receptor and the nucleotide sequence which encodes an activation domain; and

(b) a second nucleic acid molecule which comprises:

(i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound of interest is to be measured,

(ii) a nucleotide sequence which encodes a protease or a protease which is specific for said cleavage site,

wherein said first test protein differs from other first test proteins in each of said plurality of samples,

2004257228 02 Dec 2008

- 52 -

and determining activity of said reporter gene in at one or more of said plurality of samples as a determination of modulation of one or more protein interactions of interest.

- 5 17. A method according to claim 16, wherein said second test protein is different in each sample or wherein said second test protein is the same in each sample.
18. A method according to claim 16 or claim 17, wherein all of said samples are combined in a common
10 receptacle, and each samples comprises a different pair of first and second test proteins or wherein each sample is tested in a different receptacle.
19. A method according to any one of claims 16 to 18, wherein the reporter gene in a given sample differs
15 from the reporter gene in other samples.
20. A method according to claim 15, wherein said mixture of compounds comprises a biological sample.
21. A method according to claim 20, wherein said biological sample is cerebrospinal fluid, urine,
20 blood, serum, pus, ascites, synovial fluid, a tissue extract, or an exudate.
22. Recombinant cell, transformed or transfected with:
- (a) a nucleic acid molecule which comprises:
- (i) a nucleotide sequence which encodes a first
25 test protein which is a transmembrane receptor,
- (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and

2004257228 02 Dec 2008

- 63 -

(iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, wherein the nucleotide sequence which encodes a cleavage site is located between the nucleotide sequence which encodes a transmembrane receptor and the nucleotide sequence which encodes an activation domain; and

(b) a nucleic acid molecule which comprises:

(i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound is to be measured, and

(ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site.

23. A recombinant cell according to claim 22, wherein one or both of said nucleic acid molecules are stably incorporated into the genome of said cell.
24. A recombinant cell according to claim 22 or claim 23, wherein said cell has been transformed or transfected with said reporter gene.
25. A recombinant cell according to claim 22, wherein said transmembrane receptor is a GPCR.
26. A recombinant cell according to any one of claims 22 to 25, wherein said protease or portion of a protease is tobacco etch virus nuclear inclusion A protease.
27. A recombinant cell according to any one of claims 22 to 26, wherein said protein which activates said reporter gene is a transcription factor.

N:\Perth\Cases\Patent\59000-59999\IP59482 AU\Specis\IP59482.AU Specification 2008-12-1.doc 1/12/08

2004257228 02 Dec 2008

- 64 -

28. A recombinant cell according to claim 27, wherein said transcription factor is tTA or GAL4.
29. A recombinant cell according to any one of claims 22 to 28, wherein said second protein is an inhibitory protein.
30. A recombinant cell according to claim 29, wherein said inhibitory protein is an arrestin.
31. A recombinant cell according to any one of claims 22 to 30, wherein said cell is a eukaryote.
32. A recombinant cell according to any one of claims 22 to 30, wherein said cell is a prokaryote.
33. A recombinant cell according to any one of claims 22 to 32, wherein said reporter gene is an exogenous gene.
34. A recombinant cell according to claim 33, wherein said exogenous gene encodes β -galactosidase or luciferase.
35. A recombinant cell according to any one of claims 22 to 34, wherein the nucleotide sequence encoding said first test protein is modified to increase interaction with said second test protein, wherein said modification comprises replacing all or part of the nucleotide sequence of the C-terminal region of said first test protein with a nucleotide sequence which encodes an amino acid sequence which has higher affinity for said second test protein than the original sequence, wherein the nucleotide sequence of said C-terminal region is replaced by a nucleotide

2004257228 02 Dec 2008

- 65 -

sequence encoding the C-terminal region of AVPR2, AGTRLI, GRPR, F2PL1, CXCR2/IL-8B, or CCR4.

36. An isolated nucleic acid molecule which comprises, in 5' to 3' order,

5 (i) a nucleotide sequence which encodes a test protein, which is a transmembrane receptor,

(ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease, and

10 (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in a cell.

37. An isolated nucleic acid molecule according to claim 36, wherein said protease or portion of a protease is tobacco etch virus nuclear inclusion A protease.

15 38. An isolated nucleic acid molecule according to claim 36 or claim 37, wherein said protein which activates said reporter gene is a transcription factor.

39. An isolated nucleic acid molecule according to claim 38, wherein said transcription factor is tTA or GAL4.

40. An isolated nucleic acid molecule which comprises:

20 (i) nucleotide sequence which encodes an inhibitory test protein whose interaction with another test protein in the presence of a test compound is to be measured, and

(ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for a 25 cleavage site.

41. An isolated nucleic acid molecule according to claim 40, wherein said inhibitory protein is an arrestin.

2004257228 02 Dec 2008

- 66 -

42. A fusion protein produced by expression of the isolated nucleic acid molecule according to any one of claims 36 to 41, wherein the cleavage site is located between the transmembrane receptor and the activation domain.
- 5
43. A test kit when used for determining if a test compound modulates a specific protein/protein interaction of interest comprising a separate portion of each of:
- 10 (a) a nucleic acid molecule which comprises:
- (i) a nucleotide sequence which encodes a first test protein which is a transmembrane receptor,
- (ii) a nucleotide sequence encoding a cleavage site for a protease or a portion of a protease,
- 15 (iii) a nucleotide sequence which encodes a protein which activates a reporter gene in said cell, wherein the nucleotide sequence which encodes a cleavage site is located between the nucleotide sequence which encodes a transmembrane receptor and the nucleotide sequence which encodes an activation domain; and
- 20
- (b) a nucleic acid molecule which comprises:
- (i) a nucleotide sequence which encodes a second test protein whose interaction with said first test protein in the presence of said test compound is to be measured,
- 25
- (ii) a nucleotide sequence which encodes a protease or a portion of a protease which is specific for said cleavage site, and

2004257228 02 Dec 2008

- 67 -

(c) container means for holding each of (a) and (b) separately from each other.

44. A test kit according to claim 43, wherein said transmembrane receptor is a GPCR.
- 5 45. A test kit according to claim 43 or 44, wherein said protease or portion of a protease is tobacco etch virus nuclear inclusion A protease.
46. A test kit according to any one of claims 43 to 45, wherein said protein which activates said reporter gene is a transcription factor.
- 10 47. A test kit according to claim 46, wherein said transcription factor is tTA or GAL4.
48. A test kit according to any one of claims 43 to 47, wherein said second protein is an inhibitory protein.
- 15 49. A test kit according to claim 48, wherein said inhibitory protein is an arrestin.
50. A test kit according to any one of claims 43 to 49, further comprising a separate portion of an isolated nucleic acid molecule which encodes a reporter gene.
- 20 51. A test kit according to claim 50, wherein said reporter gene encodes β -galactosidase or luciferase.




Fig. 1




Fig
2a





Fig
2b

Vasopressin Dose Response Curve






Figure 4




Figure 5A




Figure 5B




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10

SEQUENCE LISTING

<110> LEE, Kevin
 AXEL, Richard
 STRAPPS, Walter
 BARNEA, Gilad

<120> Method For Assaying Protein-Protein Interaction

<130> SENTI 203.2 CIP

<140>

<141>

<150> 60/566,113

<151> 2004-04-27

<150> 60/511,918

<151> 2003-10-15

<150> 60/485,968

<151> 2003-07-09

<160> 101

<210> 1

<211> 2015

<212> DNA

<213> Homo sapiens

<400> 1

```

actgccaagc ggcttcttca gagcacgggc tggaaactggc aggcaccgcg agcccctagc      60
acccgacaag ctgagtgatg aggacgagtc cccaccacac ccacaccaca gccgctgaat      120
gaggcttcca ggcgtccgct cgcggcccgc agagcccgcg cgtgggtccg cccgctgagg      180
cgccccagc  cagtgcgctt acctgccaga ctgcgcgcca tggggcaacc cgggaacggc      240
agcgccttct tgctggcacc caatagaagc catgcgccgg accacgacgt cacgcagcaa      300
agggacgagg tgtgggtggg gggcatgggc atcgtcatgt ctctcatcgt cctggccatc      360
gtgtttggca atgtgctggg catcacagcc attgccaaat tgcagcgtct gcagacggtc      420
accaactact tcatacattc actggcctgt gctgatctgg tcatgggctt ggcagtggtg      480
ccctttgggg ccgcccatat tcttatgaaa atgtggactt ttggcaactt ctgggtgcgag      540
ttttggactt ccttgatgtg gctgtgcgtc acggccagca ttgagaccct gtgctgcatc      600
gcagtgatgc gctactttgc cactacttca cctttcaagt accagagcct gctgaccaag      660
aataaggccc gggatgatcat tctgatggtg tggatttgtt caggccttac ctctctcttg      720
cccattcaga tgcactggta cggggccacc caccaggaag ccatcaactg ctatgccaat      780
gagacctgct gtgacttctt cacgaaccaa gcctatgcca ttgcctcttc catcgtgtcc      840
ttctacgttc cctgggtgat catggctctc gtctactcca gggctcttca ggaggccaaa      900
agggcagctcc agaagattga caaatctgag ggccgcttcc atgtccagaa ccttagccag      960
gtggagcagg atggggcggc ggggcattgga ctccgcagat ctccaagt tctgcttgaag     1020
gagcacaagc cctcaagac gttaggcatc atcatgggca ctttcaccct ctgctggctg     1080
cccttcttca tggtaacat tgtgcatgtg atccaggata acctcatccg taaggaagtt     1140
tacatcctcc taaattggat aggctatgtc aattctggtt tcaatccctt tatctactgc     1200
cggagcccag atttcaggat tgccttccag gagctctctg gcctgcgcag gtcttctttg     1260
aaggcctatg ggaatggcta ctccagcaac ggcaacacag gggagcagag tggatatcac     1320
gtggaacagg agaagaaaaa taaactgctg tgtgaagacc tcccaggcac ggaagacttt     1380
gtgggccatc aaggtagctg gcctagcgat aacattgatt cacaagggag gaattgtagt     1440
acaaatgact cactgctgta aagcagtttt tctactttta aagaccccc ccccccaac     1500
agaacactaa acagactatt taacttgagg gtaataaact tagaataaaa ttgtaaaaaat     1560
tgtatagaga tatgcagaag gaagggcatc cttctgcctt ttttattttt ttaagctgta     1620
aaaagagaga aaacttattt gagtgattat ttgttatttg tacagttcag ttcctctttg     1680
  
```

catggaattt gtaagtttat gtctaaagag ctttagtcct agaggacctg agtctgctat 1740
 attttcatga cttttccatg tatctacctc actattcaag tattaggggt aatataattgc 1800
 tgctggtaat ttgtatctga aggagatttt ccttctaca cccttgact tgaggatttt 1860
 gagtatctcg gacctttcag ctgtgaacat ggactcttcc cccactcctc ttatttgctc 1920
 acacggggta ttttaggcag ggatttgagg agcagcttca gttgttttcc cgagcaaagg 1980
 tctaaagttt acagtaaata aaatgtttga ccatg 2015

<210> 2
 <211> 26
 <212> DNA
 <213> Homo Sapiens
 <400> 2
 gattgaagat ctgccttctt gctggc 26

<210> 3
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 3
 gcagaacttg gaagacctgc ggagtcc 27

<210> 4
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 4
 ggactccgca ggtcttccaa gttctgc 27

<210> 5
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 5
 ttcggatcct agcagtgagt catttgt 27

<210> 6
 <211> 7
 <212> PRT
 <213> Homo Sapiens
 <400> 6
 Glu Asn Leu Tyr Phe Gln Ser
 5

<210> 7
 <211> 32
 <212> DNA
 <213> Homo Sapiens
 <400> 7
 ccggatcctc tagattagat aaaagtaaag tg 32

<210> 8
 <211> 35
 <212> DNA
 <213> Homo Sapiens
 <400> 8
 gactcgagct agcagtatcc tcgcccccc taccc 35

<210> 9
 <211> 18
 <212> DNA
 <213> Homo Sapiens
 <400> 9
 gagaacctgt acttcag 18

<210> 10
 <211> 33
 <212> DNA
 <213> Homo Sapiens
 <400> 10
 ggatccgaga acctgtactt ccagtacaga tta 33

<210> 11
 <211> 30
 <212> DNA
 <213> Homo Sapiens
 <400> 11
 ctcgagagat cctcgcgccc cctaccacc 30

<210> 12
 <211> 7
 <212> PRT
 <213> Homo Sapiens
 <400> 12
 Glu Asn Leu Tyr Phe Gln Tyr
 5

<210> 13
 <211> 33
 <212> DNA
 <213> Homo Sapiens
 <400> 13
 ggatccgaga acctgtactt ccagctaaga tta 33

<210> 14
 <211> 7
 <212> PRT
 <213> Homo Sapiens
 <400> 14
 Glu Asn Leu Tyr Phe Gln Leu
 5

<210> 15
 <211> 33
 <212> DNA
 <213> Homo Sapiens
 <400> 15
 gcggccgccca ccatgaacgg taccgaaggc cca 33

<210> 16
 <211> 21
 <212> DNA
 <213> Homo Sapiens
 <400> 16
 ctggtgggtg gcccgggtacc a 21

<210> 17
 <211> 1936
 <212> DNA
 <213> Homo sapiens
 <400> 17
 ccccgctgt ctgctaggag agggcgggca gcgcccggc gcgcccgatc cggctgacgc 60
 atctggcccc ggttccccaa gaccagagcg gggcccggag ggagggggaa gaggcgagag 120
 cggggagggc gcgctgctgc attggcgcgg ggaggagcag ggatcttggc agcgggcgag 180
 gaggtgcga gcgagccgcg aaccgagcgg gcggcgggcg cgcgcaccat gggggagaaa 240
 cccgggacca gggctctcaa gaagtcgagc cctaactgca agctcaccgt gtacttgggc 300
 aagcgggact tcgtagatca cctggacaaa gtggaccctg tagatggcgt ggtgcttgtg 360
 gaccctgact acctgaagga ccgcaaagtg tttgtgacce tcacctgcgc cttccgctat 420
 ggccgtgaag acctggatgt gctgggcttg tccttccgca aagacctgtt catcgccacc 480
 taccaggcct tcccccggt gcccaaccca ccccgccccc ccaccgcct gcaggaccgg 540
 ctgctgagga agctgggcca gcatgcccac cccttctctc tcaccatacc ccagaatctt 600
 ccattgctcg tcacactgca gccaggccca gaggatacag gaaaggcctg cggcgtagac 660
 tttgagattc gagccttctg tgctaaatca ctagaagaga aaagccacaa aagggaactct 720
 gtgctggctgg tgatccgaaa ggtgcagttc gccccggaga aacccggccc ccagccttca 780
 gccgaaacca cacgccactt cctcatgtct gaccggctcc tgcacctcga ggcttccctg 840
 gacaaggagc tgtactacca tggggagccc ctcaatgtaa atgtccacgt caccaacaac 900
 tccaccaaga ccgtcaagaa gatcaaagtc tctgtgagac agtacgccga catctgcctc 960
 ttcagcaccg ccagttacaa gtgtcctgtg gctcaactcg aacaagatga ccaggtatct 1020
 cccagctcca cattctgtaa ggtgtacacc ataaccacc tgctcagcga caaccgggag 1080
 aagcggggtc tcgccctgga tgggaaactc aagcacgagg acaccaacct ggcttcagc 1140
 accatcgtga aggaggggtc caacaaggag gtgctgggaa tcctgggtgtc ctacagggtc 1200
 aaggtgaagc tgggtggtgtc tcgaggcggg gatgtctctg tggagctgcc ttttgttctt 1260
 atgcaccca agccccacga ccacatccc ctccccagac cccagtcagc cgtcgggag 1320
 acagatgtcc ctgtggacac caacctcatt gaatttgata ccaactatgc cacagatgat 1380
 gacattgtgt ttgaggactt tgcccggctt cggctgaagg ggatgaagga tgacgactat 1440
 gatgatcaac tctgctagga agcgggggtg gaagaaggga ggggatgggg ttgggagagg 1500
 tgagggcagg attaagatcc ccactgtcaa tgggggattg tcccagcccc tcttcccttc 1560
 cctcaccctg gaagcttctt caaccaatcc cttcacactc tctccccat ccccccaaga 1620
 tacacactgg acctctctt gctgaatgtg ggcattaatt ttttgactgc agctctgctt 1680
 ctccagcccc gccgtgggtg gcaagctgtg ttcataccta aatcttctgg aaggggacag 1740
 tgaaaagagg agtgacagga gggaaaaggg gagacaaaac tctactctc aacctcacac 1800
 caacacctcc cattatcact ctctctgcc cttctcttc aagaggagac ccttggggga 1860
 caaggccgtt tctttgtttc tgagcataaa gaagaaaata aatcttttac taagcatgaa 1920
 aaaaaaaaa aaaaaa 1936

<210> 18
 <211> 35
 <212> DNA
 <213> Homo Sapiens
 <400> 18
 caggatcctc tggaatgggg gagaaaccg ggacc 35

<210> 19
 <211> 30
 <212> DNA
 <213> Homo Sapiens
 <400> 19
 ggatccgcag agttgatcat catagtcgtc 30

<210> 20
 <211> 9
 <212> PRT
 <213> Homo Sapiens
 <400> 20
 Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
 5

<210> 21
 <211> 28
 <212> DNA
 <213> Homo Sapiens
 <400> 21
 agatctagct tgtttaaggg accacgtg 28

<210> 22
 <211> 62
 <212> DNA
 <213> Homo Sapiens
 <400> 22
 gcgccgctc aagcgtaatc tggaacatca tatgggtacg agtacaccaa ttcattcatg 60
 ag 62

<210> 23
 <211> 1809
 <212> DNA
 <213> Homo sapiens
 <400> 23
 agaagatcct gggttctgtg catccgtctg tctgaccatc cctctcaatc ttccctgccc 60
 aggactggcc atactgccac cgcacacgtg cacacacgcc aacaggcacc tgccatgctg 120
 gcatctctat aagggtcca gtccagagac cctgggccat tgaacttgcct cctcaggcag 180
 aggtgagtc cgcacatcac ctccaggccc tcagaacacc tgccccagcc ccaccatgct 240
 catggcgtcc accacttccg ctgtgctggt gcatccctct ctgcccagcc tgcccagcaa 300
 cagcagccag gagaggccac tggacacccg ggaccgctg ctagcccggg eggagctggc 360
 gctgctctcc atagtctttg tggetgtggc cctgagcaat ggcttgggtg tggcggccct 420
 agctcggcgg ggccggcggg gccactgggc accatacac gtcttcattg gccacttgtg 480
 cctggccgac ctggcctgg ctctgttcca agtgcctgcc cagctggcct ggaaggccac 540
 cgaccgcttc cgtgggccag atgcctgtg tggggcctg aagtatctgc agatgggtgg 600

catgtatgcc	tcctcctaca	tgatcctggc	catgacgctg	gaccgccacc	gtgccatctg	660
ccgtcccatg	ctggcgtacc	gccatggaag	tggggctcac	tggaaccggc	cggtgctagt	720
ggcttggggc	ttctcgetcc	ttctcagcct	gccccagetc	ttcatcttcg	cccagcgcga	780
cgtggaaggt	ggcagcgggg	tcactgactg	ctgggcctgc	tttgcggagc	cctggggccg	840
tcgcacctat	gtcacctgga	ttgccctgat	ggtgttcgtg	gcacctacc	tgggtatcgc	900
cgctgccag	gtgctcatct	tccgggagat	tcatgccagt	ctggtgccag	ggccatcaga	960
gaggcctggg	gggcgccgca	ggggacgccc	gacaggcage	cccggtgagg	gagccccact	1020
gtcagcagct	gtggccaaga	ctgtgaggat	gacgctagtg	attgtggtcg	tctatgtgct	1080
gtgctgggca	cccttcttcc	tggtgcagct	gtgggcccgc	tgggaccggg	aggcacctct	1140
ggaagggggc	ccctttgtgc	tactcatggt	gctggccagc	ctcaacagct	gcaccaaccc	1200
ctggatctat	gcatctttca	gcagcagcgt	gtcctcagag	ctgcgaagct	tgctctgctg	1260
tgcccgggga	cgcaccccac	ccagcctggg	tccccagat	gagtcctgca	ccaccgccag	1320
ctcttcctcg	gccaaggaca	cttcctcgtg	aggagctggt	gggtgtcttg	cctctagagg	1380
ctttgagaag	ctcagctgcc	ttcctggggc	tggtcctggg	agccactggg	agggggaccc	1440
gtggagaatt	ggccagagcc	tgtggcccgc	aggctgggac	actgtgtggc	cctggacaag	1500
ccacagcccc	tgcttgggtc	tccacatccc	cagctgtatg	aggagagctt	caggccccag	1560
gactgtgggg	gccccctcagg	tcagctcact	gagctgggtg	taggaggggc	tgccagcagag	1620
gcctgaggag	tggcaggaaa	gagggagcag	gtgccccag	gtgagacagc	ggccccaggg	1680
gcctgaaaag	gaaggaccag	gctggggcca	ggggaccttc	ctgtctccgc	ctttctaate	1740
cctccctcct	cattctctcc	ctaataaaaa	ttggagctct	ttccacatg	gcaaggggct	1800
tccttgaa						1809

<210> 24
 <211> 26
 <212> DNA
 <213> Homo Sapiens
 <400> 24
 gaattcatgc tcatggcgtc caccac

26

<210> 25
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 25
 ggatccccgat gaagtgtcct tggccag

27

<210>	26					
<211>	1266					
<212>	DNA					
<213>	Homo sapiens					
<400>	26					
atggatgtgc	tcagccctgg	tcagggcaac	aacaccacat	caccacccgc	tccttttgag	60
accggcggca	acactactgg	tatctccgac	gtgaccgtca	gctaccaagt	gatcacctct	120
ctgctgctgg	gcacgctcat	cttctgcgcg	gtgctgggca	atgcgtgcgt	ggtggctgcc	180
atcgccttgg	agcgtccct	gcagaacgtg	gccaattatc	ttattggctc	tttggcggtc	240
accgacctca	tggtgtcggg	ggtgggtgctg	cccatggccg	cgctgtatca	ggtgtcctaac	300
aagtggacac	tgggccagg	aacctgcgac	ctgttcacgc	ccctcgacgt	gctgtgctgc	360
accatcatca	tcttgacact	gtgcgccatc	gcgctggaca	ggtactgggc	catcacggac	420
cccactgact	acgtgaacaa	gaggacgccc	cggccgcgctg	cgctcatctc	gctcacttgg	480
cttattggct	tcctcatctc	tatcccgcct	atcctgggct	ggcgcacccc	ggaagaccgc	540
tcggaccccc	acgcattgac	cattagcaag	gatcatggct	acactatcta	ttccaccttt	600
ggagctttct	acatcccgt	gctgctcatg	ctggttctct	atgggcgcat	attccgagct	660
gcgcgcttcc	gcatccgcaa	gacggtcaaa	aaggtggaga	agaccggagc	ggacacccgc	720
catggagcat	ctcccgcctc	gcagcccaag	aagagtgtga	atggagagtc	ggggagcagg	780

aactggaggc	tgggcgtgga	gagcaaggct	gggggtgctc	tgtgcgcaa	tggcgcggtg	840
aggcaagggtg	acgatggcgc	cgccctggag	gtgatcgagg	tgcaccgagt	gggcaactcc	900
aaagagcact	tgccctctgcc	cagcgaggct	ggctctacc	cttgtgcccc	cgccctcttc	960
gagaggaaaa	atgagcgcaa	cgccgaggcg	aagcgcaaga	tggccctggc	ccgagagagg	1020
aagacagtga	agacgctggg	catcatcatg	ggcaccttca	tcctctgctg	gctgccttcc	1080
ttcatcgtgg	ctcttgttct	gcccttctgc	gagagcagct	gccacatgcc	caccctgttg	1140
ggcgccataa	tcaattggct	gggctactcc	aactctctgc	ttaaccccg	catttacgca	1200
tacttcaaca	aggactttca	aaacgcgttt	aagaagatca	ttaagtgtaa	cttctgccc	1260
cagtga						1266

<210> 27

<211> 26

<212> DNA

<213> Homo Sapiens

<400> 27

gaattcatgg atgtgctcag ccctgg 26

<210> 28

<211> 25

<212> DNA

<213> Homo Sapiens

<400> 28

ggatccctgg cggcagaact tacac 25

<210> 29

<211> 1401

<212> DNA

<213> Homo Sapiens

<400> 29

atgaataact	caacaaactc	ctctaacaat	agcctggctc	ttacaagtcc	ttataagaca	60
tttgaagtgg	tgtttattgt	cctgggtggc	ggatccctca	gtttggtgac	cattatcggg	120
aacatcctag	tcattggttc	cattaaagtc	aaccgccacc	tccagaccgt	caacaattac	180
tttttattca	gcttggcctg	tgctgacctt	atcatagggtg	ttttctccat	gaacttgtac	240
accctctaca	ctgtgattgg	ttactggcct	ttgggacctg	tgggtgtgta	cctttggcta	300
gccctggact	atgtggtcag	caatgcctca	gttatgaatc	tgctcatcat	cagctttgac	360
aggctacttct	gtgtcacaaa	acctctgacc	taccagtc	agcggaccac	aaaaatggca	420
ggatgatga	ttgcagctgc	ctgggtcctc	tctttcatcc	tctgggctcc	agccattctc	480
ttctggcagt	tcattgtagg	ggtgagaact	gtggaggatg	gggagtgtca	cattcagttt	540
ttttccaatg	ctgctgtcac	ccttggtagc	gctattgcag	ccttctattt	gccagtgate	600
atcatgactg	tgctatattg	gcacatatcc	cgagccagca	agagcaggat	aaagaaggac	660
aagaaggagc	ctggtgccaa	ccaagacccc	gtttctccaa	gtctggtaca	aggaaggata	720
gtgaagccaa	acaataacaa	catgcccagc	agtgacgatg	gcctggagca	caacaaaate	780
cagaatggca	aagccccag	ggatcctgtg	actgaaaact	gtgttcaggg	agaggagaag	840
gagagctcca	atgactccac	ctcagtcagt	gctgttgctc	ctaataatgag	agatgatgaa	900
ataaccagg	atgaaaacac	agtttccact	tccctgggccc	attccaaaga	tgagaactct	960
aagcaaacat	gcacagaat	tggcaccaag	acccccaaaa	gtgactcatg	taccccaact	1020
aataccaccg	tggaggtagt	ggggtcttca	ggtcagaatg	gagatgaaaa	gcagaatatt	1080
gtagcccgc	agattgtgaa	gatgactaag	cagcctgcaa	aaaagaagcc	tcctccttcc	1140
cgggaaaaga	aagtcaccag	gacaatcttg	gctattctgt	tggccttcat	catcacttgg	1200
gcccataca	atgtcatggt	gctcattaac	accttttgtg	caccttgcat	ccccaacact	1260
gtgtggacaa	ttggttactg	gctttgttac	atcaacagca	ctatcaaccc	tgccctgctat	1320
gcactttgca	atgccacctt	caagaagacc	tttaaacacc	ttctcatgtg	tcattataag	1380
aacataggcg	ctacaaggtg	a				1401

```

<210> 30
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 30
gaattcatga ataactcaac aaactcc 27

<210> 31
<211> 25
<212> DNA
<213> Homo Sapiens
<400> 31
agatctcctt gtagcgcta tgttc 25

<210> 32
<211> 3655
<212> DNA
<213> Homo sapiens
<400> 32
cttcagatag attatatctg gagtgaagga tcctgccacc tacgtatctg gcatagtatt 60
ctgtgtagtg ggatgagcag agaacaaaaa caaataatc cagtgagaaa agcccgtaaa 120
taaaccttca gaccagagat ctattctcca gcttatttta agctcaactt aaaaagaaga 180
actgtttctt gattcttttc gccttcaata cacttaatga tttaactcca ccctccttca 240
aaagaaacag catttctctac ttttatactg tctatatgat tgatttgcac agctcatctg 300
gccagaagag ctgagacatc cgttccccta caagaaactc tccccgggtg gaacaagatg 360
gattatcaag tgtcaagtc aatctatgac atcaattatt atacatcgga gccctgccaa 420
aaaaatcaat tgaagcaaat cgcagcccgc ctctgcctc cgtctactc actgggtgtc 480
atctttgggt ttgtgggcaa catgctggtc atctcatcc tgataaactg caaaaggctg 540
aagagcatga ctgacatcta cctgctcaac ctggccatct ctgacctgtt tttccttctt 600
actgtcccct tctgggctca ctatgctgcc gccagtgagg actttggaaa tacaatgtgt 660
caactcttga cagggtctta ttttataggc ttcttctctg gaatcttctt catcctctc 720
ctgacaatcg ataggtacct ggctgtctgc catgctgtgt ttgctttaa agccaggacg 780
gtcacctttg ggggtgtgac aagtgtgatc acttgggtgg ttggtgtgtt tgcgtctctc 840
ccaggaatca tctttaccag atctcaaaaa gaaggctctc attacacctg cagctctcat 900
ttccataca gtcagtatca attctggaag aatttccaga cattaagat agtcatcttg 960
gggtgtgtcc tgcctgtgct tgtcatggtc atctgctact cgggaatcct aaaaactctg 1020
cttcgggtgc gaaatgagaa gaagaggcac agggctgtga ggcttatctt caccatcatg 1080
attgtttatt ttctctctg ggctccctac aacatgtgct ttctctgaa caccttccag 1140
gaattctttg gcctgaataa ttgcagtagc tctaacaggt tggaccaagc tatgcaggtg 1200
acagagactc ttgggatgac gcactgctgc atcaacccca tcatctatgc ctttgtcggg 1260
gagaagttca gaaactacct cttagtcttc ttccaaaagc acattgccaa acgcttctgc 1320
aaatgctggt ctattttcca gcaagaggct cccgagcgag caagctcagt ttacaccga 1380
tccactgggg agcaggaaat atctgtgggc ttgtgacacg gactcaagtg ggctggtgac 1440
ccagtcagag ttgtgcacat ggcttagttt tcatacacag cctgggctgg ggggtgggtg 1500
ggagaggtct tttttaaag gaagttactg ttatagaggg tctaagattc atccatttat 1560
ttggcatctg tttaaagtag attagatctt ttaagcccat caattataga aagccaaatc 1620
aaaaatggtt gatgaaaaat agcaaccttt ttatctcccc ttcacatgca tcaagttatt 1680
gacaaaactc ccttctactc cgaaagttcc ttatgtatat ttaaaagaaa gcctcagaga 1740
attgctgatt cttgagttta gtgatctgaa cagaaatacc aaaattattt cagaaatgta 1800
caacttttta ctagtaciaa ggcaacatat aggttgtaaa tgtgtttaaa acaggtcttt 1860
gtcttgctat ggggagaaaa gacatgaata tgattagtaa agaaatgaca cttttcatgt 1920
gtgatttccc ctccaaggta tggtaataaa gtttactga cttagaacca ggcgagagac 1980
ttgtggcctg ggagagctgg ggaagcttct taaatgagaa ggaatttgag ttggatcatc 2040
tattgtcggc aaagacagaa gcctcactgc aagcactgca tgggcaagct tggctgtaga 2100

```

```

aggagacaga gctggttggg aagacatggg gaggaaggac aaggctagat catgaagaac 2160
cttgacggca ttgctccgtc taagtcatga gctgagcagg gagatcctgg ttggtggtgc 2220
agaaggttta ctctgtggcc aaaggagggt caggaaggat gagcatttag ggcaaggaga 2280
ccaccaacag ccctcaggtc aggggtgagga tggcctctgc taagctcaag gcgtgaggat 2340
gggaaggagg gaggtatteg taaggatggg aaggagggag gtattcgtgc agcatatgag 2400
gatgcagagt cagcagaact ggggtggatt tggtttgaa gtgagggtca gagaggagtc 2460
agagagaatc cctagtcttc aagcagattg gagaaaccct tgaaaagaca tcaagcacag 2520
aaggaggagg aggaggttta ggtcaagaag aagatggatt ggtgtaaaag gatgggtctg 2580
gtttgcagag cttgaacaca gtctcaccca gactccaggc tgtctttcac tgaatgcttc 2640
tgacttcata gatttccttc ccatcccagc tgaataactg aggggtctcc aggaggagac 2700
tagatztatg aatacacgag gtatgaggtc taggaacata cttcagctca cacatgagat 2760
ctaggtgagg attgattacc tagtagtcat ttcatgggtt gttgggagga ttctatgagg 2820
caaccacagt cagcatttag cacatactac acattcaata agcatcaaac tcttagttac 2880
tcattcaggg atagcactga gcaaagcatt gagcaaagg gtcccatata ggtgagggaa 2940
gcctgaaaaa ctaagatgct gcctgcccag tgcacacaag tgtaggatc attttctgca 3000
ttaaccgtc aataggcaaa ggggggaagg gacatattca tttggaaata agctgccttg 3060
agccttaaaa cccacaaaag tacaatttac cagcctccgt atttcagact gaatgggggt 3120
ggggggggcg ccttaggtac ttattccaga tgccttctcc agacaaacca gaagcaacag 3180
aaaaaatcgt ctctccctcc ctttgaaatg aatatcccc ttagtgtttg ggtatattca 3240
tttcaaaggg agagagagag gtttttttct gttcttctc atatgattgt gcacatactt 3300
gagactgttt tgaatttggg ggatggctaa aaccatcata gtacaggtaa ggtgagggaa 3360
tagtaagtgg tgagaactac tcagggaatg aagggtgtcag aataataaga ggtgctactg 3420
actttctcag cctctgaata tgaacgggtg gcattgtggc tgtcagcagg aagcaacgaa 3480
gggaaatgtc tttccttttg ctotaaagtt gtggagagtg caacagtagc ataggaccct 3540
accctctggg ccaagtcaaa gacattctga catcttagta tttgcatatt cttatgtatg 3600
tgaaagttac aaattgcttg aaagaaaata tgcattctaat aaaaaacacc ttcta 3655

```

```

<210> 33
<211> 31
<212> DNA
<213> Homo Sapiens
<400> 33
gcgcccgcat ggattatcaa gtgtcaagtc c 31

```

```

<210> 34
<211> 25
<212> DNA
<213> Homo Sapiens
<400> 34
ggatccctgg cggcagaact tacac 25

```

```

<210> 35
<211> 33
<212> DNA
<213> Homo Sapiens
<400> 35
ggtctccaat tcatgatta tcaagtgca agt 33

```

```

<210> 36
<211> 21
<212> DNA
<213> Homo Sapiens
<400> 36

```

gacgacagcc aggtacctat c

21

<210> 37

<211> 2643

<212> DNA

<213> Homo sapiens

<400> 37

```

ggcagccgctc cggggccgccc actctcctcg gccggctccct ggetcccgga ggcggccgcg      60
cgtggatgcg gcgggagctg gaagcctcaa gcagccggcg ccgtctctgc cccggggcgc      120
cctatggctt gaagagcctg gccaccagcagg ggtccaccg ccctgatgga tccactgaat      180
ctgtcctggg atgatgatga tctggagagg cagaactgga gccggccctt caacgggtca      240
gacgggaagg cggacagacc ccactacaac tactatgcca cactgctcac cctgctcacc      300
gctgtcatcg tcttcggcaa cgtgctggtg tgcattgctg tgtcccgca gaaggcctg      360
cagaccacca ccaactacct gatcgtcagc ctgcagtgcc cgcacctcct cgtcgccaca      420
ctgggtcatg cctgggttgt ctacctggag gtggtagggt agtggaaatt cagcaggatt      480
cactgtgaca tcttcgtcac tctggacgct atgatgtgca cggcgagcat cctgaacttg      540
tgtgccatca gcatcgacag gtacacagct gtggccatgc ccatgctgta caatacgcgc      600
tacagctcca agcgcggggt caccgtcatg atctccatcg tctgggtcct gtcctcacc      660
atctcctgcc cactcctctt cggactcaat aacgcagacc agaacgagtg catcatggcc      720
aaccggcctc tcgtggtcta ctctccatc gtctccttct acgtgccctt cattgtcacc      780
ctgctggctc acatcaagat ctacattgtc ctccgcagac gccgcaagg agtcaacacc      840
aaacgcagca gccgagcttt cagggccac ctgagggtc cactaaagg caactgtact      900
caccggagg acatgaaact ctgcaccgtt atcatgaagt ctaatgggag tttcccagtg      960
aacaggcggg gagtggaggc tgcccggcga gccaggaggc tggagatgga gatgctctcc     1020
agcaccagcc caccggagag gaccgggtac agccccatcc caccagcca ccaccagctg     1080
actctccccg acccgtcca ccattgctc cacagcactc ccgacagccc cgccaaacca     1140
gagaagaatg ggcattgcaa agaccacccc aagattgcca agatctttga gatccagacc     1200
atgcccaatg gcaaaacccg gacctcctc aagaccatga gccgtaggaa gctctcccag     1260
cagaaggaga agaaagccac tcagatgctc gccattgttc tcggcgtgtt catcatctgc     1320
tggctgcctt tcttcatcac acacatcctg aacatacact gtgactgcaa catcccgcct     1380
gtcctgtaca ggccttcac gtggctgggc tatgtcaaca gcgctgtgaa ccccatcacc     1440
tacaccacct tcaacattga gttccgcaag gcttctctga agatcctcca ctgctgactc     1500
tgetgcctgc ccgcacagca gctgcttcc cacctcctg cccaggccgg ccagcctcac     1560
ccttgcgaac cgtgagcagg aaggcctggg tggatcggcc tcctcttcac cccggcaggc     1620
cctgcagtggt tcgcttggct ccattgctc cactgccgc acaccctcac tctgccaggg     1680
cagtgtagat gagctgggca tggtagcagc cctggggctg ggcctccag ctcaggggca     1740
gctcatagag tccccctcc cacctccagt cccctatcc ttggcacc aaagatgcagcc     1800
gccttctctg acctcctct ggggtctag ggttgcgga gcctgagtc gggccagag     1860
gctgagtttt ctctttgtgg ggcttggcgt ggagcaggcg gtggggagag atggacagtt     1920
cacaccctgc aaggccaca ggaggcaagc aagctctctt gccgaggagc caggcaactt     1980
cagtcctggg agaccatgt aaataccaga ctgcaggtg gaccacagag attccaagc     2040
caaaaacctt agctcctcc cgcacccga tgtggacctc tactttccag gctagtccgg     2100
accacctca ccccgttaca gctccccag tggtttccac atgctctgag aagaggagcc     2160
ctcatcttga agggcccagg aggtctatg gggagaggaa ctcttggcc tagcccacc     2220
tgetgccttc tgacggcct gcaatgtatc cctctcaca gcacatgctg gccagcctgg     2280
ggcctggcag ggaggtcagg ccctggaact ctatctgggc ctgggctagg ggacatcaga     2340
ggttctttga gggactgct ctgccacact ctgacgcaa accacttcc tttctatctc     2400
cttctggcct ttcctctctc ctgtttcct tcccttcac tgctctgccc ttagaggagc     2460
ccacggctaa gaggctgctg aaaacctct ggcctggcct ggcctgccc tgaggaagga     2520
gggaaagctg cagcttggga gagccctgg ggcctagact ctgtaacatc actatccatg     2580
caccaaacta ataaaacttt gacgagtcac ctccaggac cctgggta aaaaaaaaaa     2640
aaa

```

<210> 38

<211> 27

<212> DNA
 <213> Homo Sapiens
 <400> 39
 gaattcatgg atccactgaa tctgtcc

27

<210> 39
 <211> 25
 <212> DNA
 <213> Homo Sapiens
 <400> 39
 agatctgcag tggaggatct tcagg

25

<210> 40
 <211> 1301
 <212> DNA
 <213> Homo sapiens
 <400> 40

atgggacgaca	aagggacgcg	agtgttcaag	aaggccagtc	caaatggaaa	gctcaccgtc	60
tacctgggaa	agcgggactt	tgtggaccac	atcgacctcg	tggaccctgt	ggatgggtgtg	120
gtcctgggtgg	atcctgagta	tctcaaagag	cgagaggtct	atgtgacgct	gacctgcgcc	180
ttccgctatg	gccgggagga	cctggatgtc	ctgggcctga	cctttcgcaa	ggacctgttt	240
gtggccaacg	tacagtctgt	cccaccggcc	cccgaggaca	agaagcccct	gacgaggctg	300
caggaacgcc	tcatcaagaa	gctgggagag	cacgcttacc	ctttcacctt	tgagatccct	360
ccaaaccttc	catgttctgt	gacactgcag	ccggggcccg	aagacacggg	gaaggcttgc	420
ggtgtggact	atgaagtcaa	agccttctgc	gcgagaaatt	tggaggagaa	gatccacaag	480
cggaattctg	tgcgtctggt	catccggaag	gttcagtatg	ccccagagag	gcctggcccc	540
cagcccacag	ccgagaccac	caggcagttc	ctcatgtcgg	acaagccctt	gcacctagaa	600
gcctctctgg	ataaggagat	ctattaccat	ggagaaccca	tcagcgtcaa	cgccacgctc	660
accaacaaca	ccaacaagac	ggtgaagaag	atcaagatct	cagtgcgcca	gtatgcagac	720
atctgccttt	tcaacacagc	tcagtacaag	tgcctgttg	ccatggaaga	ggctgatgac	780
actgtggcac	ccagctcgac	gttctgcaag	gtctacacac	tgacccctt	cctagccaat	840
aaccgagaga	agcggggcct	cgcttggac	gggaagctca	agcacgaaga	cacgaacttg	900
gcctctagca	ccctgttgag	ggaaggtgcc	aaccgtgaga	tcctggggat	cattgtttcc	960
tacaaagtga	aagtgaagct	ggtggtgtct	cggggcggcc	tggtgggaga	tcttgcattcc	1020
agcagctgg	ccgtggaact	gccttcacc	ctaattgcacc	ccaagcccaa	agaggaaccc	1080
ccgcatcggg	aagttccaga	gaacgagacg	ccagtagata	ccaatctcat	agaacttgac	1140
acaaatgatg	acgacattgt	attgaggac	tttgcctgcc	agagactgaa	aggcatgaag	1200
gatgacaagg	aggaagagga	ggatggtacc	ggctctccac	agctcaacaa	cagatagacg	1260
ggcggccct	gcctccacgt	ggctccggct	ccactctcgt	g		1301

<210> 41
 <211> 30
 <212> DNA
 <213> Homo Sapiens
 <400> 41
 ggtaccatgg gcgacaaagg gacgcgagtg

30

<210> 42
 <211> 48
 <212> DNA
 <213> Homo Sapiens
 <400> 42
 ggatcctctg ttggtgagct gtggagagcc tgtaccatcc tectcttc

48

<210> 43
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 43
ggatccattt gtgtcaagtt ctatgag 27

<210> 44
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 44
ggtacatgg gggagaaacc cgggacc 27

<210> 45
<211> 24
<212> DNA
<213> Homo Sapiens
<400> 45
ggatcctgtg gcatagttgg tate 24

<210> 46
<211> 33
<212> DNA
<213> Homo Sapiens
<400> 46
tgtgcgcg gacgcacccc acccagcctg ggt 33

<210> 47
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 47
gaattcatgg atccactgaa tctgtcc 27

<210> 48
<211> 32
<212> DNA
<213> Homo Sapiens
<400> 48
tgtgcgcg cagtggagga tcttcaggaa ggc 32

<210> 49
<211> 33
<212> DNA
<213> Homo Sapiens
<400> 49
gcggcgcca ccatgaacgg taccgaaggc cca 33

<210> 50
 <211> 30
 <212> DNA
 <213> Homo Sapiens
 <400> 50
 tgtgcgcgcg cacagaagct cctggaaggc

30

<210> 51
 <211> 1602
 <212> DNA
 <213> Homo sapiens
 <400> 51

gagctccgtg ctgggagggtg ggaagggggc ttgaccctgg ggactcaggc agtctgggga 60
 cagttccacc agggggccggt gcctagaatt ggtgaggag gcacctcagg ggctgggga 120
 gaaggaacga gcgctcttog cccctctctg gcaccacagc gcgcgctgc tggccgga 180
 ggcagcgaga agtccgttct ccctgtcctg cccccggcga cttgcccggc gggtgggagt 240
 ccgcaggctc cgggtcccca gcgcccgtgg ccaggggcgc ggcaaagtgt gcctctccgc 300
 gtccagcccg ttctttcgct cccgcagcgc cgcagggtgc gcctgtcctc gccttccctgc 360
 tgcaatcgcc ccaccatgga ctccccgatc cagatcttcc gcggggagcc gggccctacc 420
 tgcgccccga gcgctgcct gcccccaac agcagcgcct ggtttccgg ctgggcccag 480
 cccgacagca acggcagcgc cggctcggag gacgcgcagc tggagcccgc gcacatctcc 540
 ccggccatcc cggatcatc cacggcggtc tactccgtag tgttcgtcgt gggcttgggtg 600
 ggcaactcgc tggatcatgt cgtgatcatc cgatacaaa agatgaagac agcaaccaac 660
 atttacatat ttaacctggc tttggcagat gctttagtta ctacaacct gccctttcag 720
 agtacggctc acttgatgaa ttccctggcct tttggggatg tgctgtgcaa gatagtaatt 780
 tccattgatt actacaacat gttcaccagc atcttcaact tgaccatgat gacggtggac 840
 cgctacattg ccgtgtgcca ccccgtagag gctttggact tccgcacacc cttgaaggca 900
 aagatcatca atatctgcat ctggctgctg tcgtcatctg ttggcatctc tgcaatagtc 960
 cttggaggca ccaaagttag ggaagacgtc gatgtcattg agtgcctctt gcagttccca 1020
 gatgatgact actcctgggtg ggacctcttc atgaagatct gcgtcttcat ctttgccctc 1080
 gtgatccctg tcctcatcat catcgtctgc tacacctga tgatcctgcg tctcaagagc 1140
 gtccggctcc tttctggctc ccgagagaaa gatcgcaacc tgcgtaggat caccagactg 1200
 gtccctgggtg tgggtggcagt cttcgtcgtc tgctggactc ccattcacat attcactctg 1260
 gtggaggctc tggggagcac ctcccacagc acagctgctc tctccagcta ttactttctg 1320
 atcgccttag gctataccaa cagtgcctg aatcccattc tctacgcctt tcttgatgaa 1380
 aacttcaagc ggtgtttccg ggacttctgc tttccactga agatgaggat ggagcggcag 1440
 agcactagca ggtccgaaa tacagttcag gatcctgctt acctgagggg catcgatggg 1500
 atgaataaac cagtatgact agtcgtggag atgtcttctt acagttcttc ggaagagag 1560
 gagttcaatg atctaggttt aactcagatc actactgcag tc 1602

<210> 52
 <211> 24
 <212> DNA
 <213> Homo Sapiens
 <400> 52
 ggtctacttg atgaattcct ggcc

24

<210> 53
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 53
 gcgcgcacag aagtcccgga aacaccg

27

				325					330				335		
Gln	Trp	Arg	Gln	Lys	Thr	Asn	Thr	Ser	Tyr	Asn	Phe	His	Ser	Phe	Ser
			340					345					350		
Ile	Arg	Met	Ala	Ile	Ser	Leu	Gly	Leu	Asn	Arg	Asp	Leu	Pro	Ser	Ser
		355					360					365			
Phe	Ser	Asp	Ser	Ser	Ile	Leu	Glu	Gln	Arg	Arg	Arg	Ile	Trp	Trp	Ser
		370				375					380				
Val	Tyr	Ser	Trp	Glu	Ile	Gln	Leu	Ser	Leu	Leu	Tyr	Gly	Arg	Ser	Ile
385					390					395					400
Gln	Leu	Ser	Gln	Asn	Thr	Ile	Ser	Phe	Pro	Ser	Ser	Val	Asp	Asp	Val
				405					410					415	
Gln	Arg	Thr	Thr	Thr	Gly	Pro	Thr	Ile	Tyr	His	Gly	Ile	Ile	Glu	Thr
			420					425					430		
Ala	Arg	Leu	Leu	Gln	Val	Phe	Thr	Lys	Ile	Tyr	Glu	Leu	Asp	Lys	Thr
		435					440					445			
Val	Thr	Ala	Glu	Lys	Ser	Pro	Ile	Cys	Ala	Lys	Lys	Cys	Leu	Met	Ile
	450					455						460			
Cys	Asn	Glu	Ile	Glu	Glu	Val	Ser	Arg	Gln	Ala	Pro	Lys	Phe	Leu	Gln
465						470					475				480
Met	Asp	Ile	Ser	Thr	Thr	Ala	Leu	Thr	Asn	Leu	Leu	Lys	Glu	His	Pro
				485					490					495	
Trp	Leu	Ser	Phe	Thr	Arg	Phe	Glu	Leu	Lys	Trp	Lys	Gln	Leu	Ser	Leu
			500					505					510		
Ile	Ile	Tyr	Val	Leu	Arg	Asp	Phe	Phe	Thr	Asn	Phe	Thr	Gln	Lys	Lys
		515					520						525		
Ser	Gln	Leu	Glu	Gln	Asp	Gln	Asn	Asp	His	Gln	Ser	Tyr	Glu	Val	Lys
		530				535						540			
Arg	Cys	Ser	Ile	Met	Leu	Ser	Asp	Ala	Ala	Gln	Arg	Thr	Val	Met	Ser
545					550						555				560
Val	Ser	Ser	Tyr	Met	Asp	Asn	His	Asn	Val	Thr	Pro	Tyr	Phe	Ala	Trp
				565					570					575	
Asn	Cys	Ser	Tyr	Leu	Phe	Asn	Ala	Val	Leu	Val	Pro	Ile	Lys	Thr	
			580					585					590		
Leu	Leu	Ser	Asn	Ser	Lys	Ser	Asn	Ala	Glu	Asn	Asn	Glu	Thr	Ala	Gln
		595					600					605			
Leu	Leu	Gln	Gln	Ile	Asn	Thr	Val	Leu	Met	Leu	Leu	Lys	Lys	Leu	Ala
		610				615						620			
Thr	Phe	Lys	Ile	Gln	Thr	Cys	Glu	Lys	Tyr	Ile	Gln	Val	Leu	Glu	Glu
625					630						635				640
Val	Cys	Ala	Pro	Phe	Leu	Leu	Ser	Gln	Cys	Ala	Ile	Pro	Leu	Pro	His
				645					650					655	
Ile	Ser	Tyr	Asn	Asn	Ser	Asn	Gly	Ser	Ala	Ile	Lys	Asn	Ile	Val	Gly
			660					665					670		
Ser	Ala	Thr	Ile	Ala	Gln	Tyr	Pro	Thr	Leu	Pro	Glu	Glu	Asn	Val	Asn
		675						680					685		
Asn	Ile	Ser	Val	Lys	Tyr	Val	Ser	Pro	Gly	Ser	Val	Gly	Pro	Ser	Pro
			690			695						700			
Val	Pro	Leu	Lys	Ser	Gly	Ala	Ser	Phe	Ser	Asp	Leu	Val	Lys	Leu	Leu
705					710					715					720
Ser	Asn	Arg	Pro	Pro	Ser	Arg	Asn	Ser	Pro	Val	Thr	Ile	Pro	Arg	Ser
			725						730					735	
Thr	Pro	Ser	His	Arg	Ser	Val	Thr	Pro	Phe	Leu	Gly	Gln	Gln	Gln	Gln
			740					745					750		
Leu	Gln	Ser	Leu	Val	Pro	Leu	Thr	Pro	Ser	Ala	Leu	Phe	Gly	Gly	Ala
		755					760						765		
Asn	Phe	Asn	Gln	Ser	Gly	Asn	Ile	Ala	Asp	Ser	Ser	Leu	Ser	Phe	Thr
		770				775						780			

Phe Thr Asn Ser Ser Asn Gly Pro Asn Leu Ile Thr Thr Gln Thr Asn
 785 790 795 800
 Ser Gln Ala Leu Ser Gln Pro Ile Ala Ser Ser Asn Val His Asp Asn
 805 810 815
 Phe Met Asn Asn Glu Ile Thr Ala Ser Lys Ile Asp Asp Gly Asn Asn
 820 825 830
 Ser Lys Pro Leu Ser Pro Gly Trp Thr Asp Gln Thr Ala Tyr Asn Ala
 835 840 845
 Phe Gly Ile Thr Thr Gly Met Phe Asn Thr Thr Thr Met Asp Asp Val
 850 855 860
 Tyr Asn Tyr Leu Phe Asp Asp Glu Asp Thr Pro Pro Asn Pro Lys Lys
 865 870 875 880
 Glu

<210> 56
 <211> 13
 <212> PRT
 <213> Homo Sapiens
 <400> 56
 Pro Gln Lys Gly Ser Ala Ser Glu Lys Thr Met Val Phe
 5 10

<210> 57
 <211> 549
 <212> PRT
 <213> Homo sapiens
 <400> 57
 Met Asp Asp Leu Phe Pro Leu Ile Phe Pro Ser Glu Pro Ala Gln Ala
 5 10 15
 Ser Gly Pro Tyr Val Glu Ile Ile Glu Gln Pro Lys Gln Arg Gly Met
 20 25 30
 Arg Phe Arg Tyr Lys Cys Glu Gly Arg Ser Ala Gly Ser Ile Pro Gly
 35 40 45
 Glu Arg Ser Thr Asp Thr Thr Lys Thr His Pro Thr Ile Lys Ile Asn
 50 55 60
 Gly Tyr Thr Gly Pro Gly Thr Val Arg Ile Ser Leu Val Thr Lys Asp
 65 70 75 80
 Pro Pro His Arg Pro His Pro His Glu Leu Val Gly Lys Asp Cys Arg
 85 90 95
 Asp Gly Tyr Tyr Glu Ala Asp Leu Cys Pro Asp Arg Ser Ile His Ser
 100 105 110
 Phe Gln Asn Leu Gly Ile Gln Cys Val Lys Lys Arg Asp Leu Glu Gln
 115 120 125
 Ala Ile Ser Gln Arg Ile Gln Thr Asn Asn Asn Pro Phe His Val Pro
 130 135 140
 Ile Glu Glu Gln Arg Gly Asp Tyr Asp Leu Asn Ala Val Arg Leu Cys
 145 150 155 160
 Phe Gln Val Thr Val Arg Asp Pro Ala Gly Arg Pro Leu Leu Leu Thr
 165 170 175
 Pro Val Leu Ser His Pro Ile Phe Asp Asn Arg Ala Pro Asn Thr Ala
 180 185 190
 Glu Leu Lys Ile Cys Arg Val Asn Arg Asn Ser Gly Ser Cys Leu Gly
 195 200 205
 Gly Asp Glu Ile Phe Leu Leu Cys Asp Lys Val Gln Lys Glu Asp Ile
 210 215 220

Glu Val Tyr Phe Thr Gly Pro Gly Trp Glu Ala Arg Gly Ser Phe Ser
 225 230 235 240
 Gln Ala Asp Val His Arg Gln Val Ala Ile Val Phe Arg Thr Pro Pro
 245 250 255
 Tyr Ala Asp Pro Ser Leu Gln Ala Pro Val Arg Val Ser Met Gln Leu
 260 265 270
 Arg Arg Pro Ser Asp Arg Glu Leu Ser Glu Pro Met Glu Phe Gln Tyr
 275 280 285
 Leu Pro Asp Thr Asp Asp Arg His Arg Ile Glu Glu Lys Arg Lys Arg
 290 295 300
 Thr Tyr Glu Thr Phe Lys Ser Ile Met Lys Lys Ser Pro Phe Asn Gly
 305 310 315 320
 Pro Thr Glu Pro Arg Pro Pro Thr Arg Arg Ile Ala Val Pro Thr Arg
 325 330 335
 Asn Ser Thr Ser Val Pro Lys Pro Ala Pro Gln Pro Tyr Thr Phe Pro
 340 345 350
 Ala Ser Leu Ser Thr Ile Asn Phe Asp Glu Phe Ser Pro Met Leu Leu
 355 360 365
 Pro Ser Gly Gln Ile Ser Asn Gln Ala Leu Ala Leu Ala Pro Ser Ser
 370 375 380
 Ala Pro Val Leu Ala Gln Thr Met Val Pro Ser Ser Ala Met Val Pro
 385 390 395 400
 Leu Ala Gln Pro Pro Ala Pro Ala Pro Val Leu Thr Pro Gly Pro Pro
 405 410 415
 Gln Ser Leu Ser Ala Pro Val Pro Lys Ser Thr Gln Ala Gly Glu Gly
 420 425 430
 Thr Leu Ser Glu Ala Leu Leu His Leu Gln Phe Asp Ala Asp Glu Asp
 435 440 445
 Leu Gly Ala Leu Leu Gly Asn Ser Thr Asp Pro Gly Val Phe Thr Asp
 450 455 460
 Leu Ala Ser Val Asp Asn Ser Glu Phe Gln Gln Leu Leu Asn Gln Gly
 465 470 475 480
 Val Ser Met Ser His Ser Thr Ala Glu Pro Met Leu Met Glu Tyr Pro
 485 490 495
 Glu Ala Ile Thr Arg Leu Val Thr Gly Ser Gln Arg Pro Pro Asp Pro
 500 505 510
 Ala Pro Thr Pro Leu Gly Thr Ser Gly Leu Pro Asn Gly Leu Ser Gly
 515 520 525
 Asp Glu Asp Phe Ser Ser Ile Ala Asp Met Asp Phe Ser Ala Leu Leu
 530 535 540
 Ser Gln Ile Ser Ser
 545

<210> 58

<211> 1833

<212> DNA

<213> Homo sapiens

<400> 58

ggaggtggga ggagggagtg acgagtcaag gaggagacag ggacgcagga gggtgcaagg 60
 aagtgtctta actgagacgg gggtaaggca agagaggggtg gaggaaattc tgcaggagac 120
 aggcttcttc cagggtctgg agaaccacaga gccagctcct cctgagtgct ggaaggact 180
 ctgggcatct tcagcccttc ttactctctg aggctcaagc cagaaattca ggctgcttgc 240
 agagtgggtg acagagccac ggagctggtg tccttgggac cctctgcccg tcttctctcc 300
 actcccagc atggaggaag gtggtgattt tgacaactac tatggggcag acaaccagtc 360
 tgagtgtgag tacacagact ggaatcctc gggggccctc atccctgcc a tetacatgtt 420
 ggtcttcttc ctgggcacca cgggcaacgg tctggtgctc tggaccgtgt ttcggagcag 480

ccgggagaag	aggcgtcag	ctgatatctt	cattgctagc	ctggcgggtg	ctgacctgac	540
cttcgtgggtg	acgctgcccc	tgtgggctac	ctacacgtac	cgggactatg	actggccctt	600
tgggaccttc	ttctgcaagc	tcagcagcta	cctcatcttc	gtcaacatgt	acgccagcgt	660
cttctgcctc	accggectca	gcttcgaccg	ctacctggcc	atcgtgaggc	cagtggccaa	720
tgctcggtcg	aggctgcggg	tcagcggggc	cgtggccacg	gcagttcttt	gggtgctggc	780
cgccctcctg	gccatgectg	tcatggtgtt	acgcaccacc	ggggacttgg	agaacaccac	840
taaggtgcag	tgctacatgg	actactgcat	ggtggccact	gtgagctcag	agtgggctg	900
ggaggtgggc	cttggggctc	cgteccaccac	cgtgggcttt	gtggtgccct	tcaccatcat	960
gctgacctgt	tacttcttca	tcgcccacac	catcgctggc	cacttccgca	aggaacgcat	1020
cgagggcctg	cggaaagcggc	gccggctgct	cagcatcatc	gtgggtgctg	tggtgacctt	1080
tgccctgtgc	tggatgccct	accacctggt	gaagacgctg	tacatgctgg	gcagcctgct	1140
gcactggccc	tgtgactttg	acctcttctc	catgaacatc	ttcccctact	gcacctgcat	1200
cagctacgtc	aacagctgcc	tcaaccctt	cctctatgcc	tttttcgacc	cccgttccg	1260
ccaggcctgc	acctccatgc	tctgctgtgg	ccagagcagg	tgcgcaggca	cctcccacag	1320
cagcagtggtg	gagaagtcag	ccagctactc	ttcggggcac	agccaggggc	ccggcccaa	1380
catgggcaag	ggtggagaac	agatgcacga	gaaatccatc	ccctacagcc	aggagacctt	1440
tgtggttgac	tagggctggg	agcagagaga	agcctggcgc	cctcggccct	ccccggcctt	1500
tgcccttgct	ttctgaaaaat	cagagtcacc	tcctctgcc	agagctgtcc	tcaaagcatc	1560
cagtgaacac	tggaagaggc	ttctagaagg	gaagaaattg	tcctctgag	gccgcctgg	1620
gtgacctgca	gagacttctc	gcctggaact	catctgtgaa	ctgggacaga	agcagaggag	1680
gctgctgct	gtgatacccc	cttacctccc	ccagtgcctt	cttcagaata	tctgactgt	1740
cttctgatcc	tgttagtcac	tgtggttcat	caaataaaac	tgtttgtgca	actgtttgtg	1800
ccaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaa			1833

<210> 59
 <211> 1726
 <212> DNA
 <213> Homo sapiens
 <400> 59

aactgcagcc	agggagactc	agactagaat	ggaggtagaa	agaactgatg	cagagtgggt	120
ttaattctaa	gcctttttgt	ggctaagttt	tgttgttgtt	aacttattga	atttagagtt	180
gtattgcact	ggctcatgtga	aagccagagc	agcaccagtg	tcaaaatagt	gacagagag	240
tttgaatacc	atagttagta	tatatgtact	cagagtattt	ttattaaaga	aggcaaagag	300
ccggcatag	atcttatctt	catcttcact	cggttgcaaa	atcaatagtt	aagaaatagc	360
atctaaggga	acttttaggt	gggaaaaaaaa	atctagagat	ggctctaaat	gactgtttcc	420
ttctgaactt	ggaggtggac	catttcatgc	actgcaacat	ctccagtcac	agtgcggatc	480
tccccgtgaa	cgatgactgg	tcccaccggg	ggatcctcta	tgtcatccct	gcagtttatg	540
gggttatcat	tctgataggc	ctcattggca	acatcacttt	gatcaagatc	ttctgtacag	600
tcaagtccat	gcgaaacggt	ccaaacctgt	tcatttccag	tctggctttg	ggagacctgc	660
tcctccta	aacgtgtgct	ccagtggatg	ccagcaggta	cctggctgac	agatggctat	720
ttggcaggat	tggctgcaaa	ctgatcccct	ttatacagct	tacctctgtt	ggggtgtctg	780
tcttcacact	cacggcgctc	tcggcagaca	gatacaaac	cattgtccgg	ccaatggata	840
tccaggcctc	ccatgccctg	atgaagatct	gcctcaaac	cgcctttatc	tggatcatct	900
ccatgctgct	ggccattcca	gaggccgtgt	tttctgacct	ccatcccttc	catgaggaaa	960
gcaccaacca	gaccttcatt	agetgtgccc	catacccaca	ctctaagtga	cttcacccca	1020
aaatccattc	tatggcttcc	tttctggtct	tctacgtcat	cccactgtcg	atcatctctg	1080
tttactacta	cttcattgct	aaaaatctga	tccagagtgc	ttacaatctt	cccgtggaag	1140
ggaatataca	tgtcaagaag	cagattgaa	cccggaaagc	acttgccaag	acagtgtctg	1200
tgtttgtggg	cctgttccgc	ttctgctggc	tccccaatca	tgtcatctac	ctgtaccgct	1260
octaccacta	ctctagagtg	gacacctca	tgtccactt	tgtcaccagc	atctgtgccc	1320
gcctcctggc	cttcaccaac	tcctgcgtga	accctttgc	cctctacctg	ctgagcaaga	1380
gtttcaggaa	acagttcaac	actcagctgc	tctgttgcca	gcctggcctg	atcatccggg	1440
ctcacagcac	tggaaaggag	acaacctgca	tgacctccct	caagagtacc	aacctctccg	1500
tggccacctt	tagcctcatc	aatggaaaca	tctgtcacga	gcgggtatgtc	tagattgacc	1560
cttgattttg	ccccctgagg	gacggttttg	cttlatggct	agacaggaac	ccttgcatcc	1620
attgttgtgt	ctgtgcctc	caaagagcct	tcagaatgct	cctgagtggg	gtaggtgggg	1680

gtggggaggc ccaaatgatg gatcaccatt atatthttgaa agaagc 1726

<210> 60
 <211> 2876
 <212> DNA
 <213> Homo sapiens
 <400> 60

tgaaacctaa	ccgcctctgg	ggaggcgcgc	agcagaggct	ccgattcggg	gcaggtgaga	60
ggctgacttt	ctctcgggtgc	gtccagtggg	gctctgagtt	tcgaatcggc	ggcggcggat	120
tccccgcgcg	cccggcgtcg	gggtctccag	gaggatgcgg	agccccagcg	cggcgtggct	180
gctggggggcc	gccatcctgc	tagcagcctc	tctctcctgc	agtggcacca	tccaaggaaac	240
caatagatcc	tctaaaggaa	gaagccttat	tggtaagggt	gatggcacat	cccacgtcac	300
tggaaaagga	gttacagttg	aaacagtctt	ttctgtggat	gagttttctg	catctgtcct	360
cactggaaaa	ctgaccactg	tcttccttcc	aattgtctac	acaattgtgt	ttgtggtggg	420
tttgccaagt	aacggcatgg	ccctgtgggt	ctttcttttc	cgaactaaga	agaagcaccc	480
tgctgtgatt	tacatggcca	atctggcctt	ggctgacctc	ctctctgtca	tctggttccc	540
cttgaagatt	gcctatcaca	tacatggcaa	caactggatt	tatggggaag	ctctttgtaa	600
tgtgcttatt	ggctttttct	atggcaacat	gtactgttcc	attctcttca	tgacctgcct	660
cagtgtgcag	aggtattggg	tcacgtgaa	ccccatgggg	cactccagga	agaaggcaaa	720
cattggcatt	ggcatctccc	tggcaatatg	gctgtgatt	ctgctggcca	ccatcccctt	780
gtatgtcgtg	aagcagacca	tcttcattcc	tgccctgaac	atcacgacct	gtcatgatgt	840
tttgccctgag	cagctcttgg	tgggagacat	gttcaattac	ttcctctctc	tggccattgg	900
ggctcttctg	ttcccagcct	tectcacagc	ctctgcctat	gtgctgatga	tcagaatgct	960
gcgatcttct	gccatggatg	aaaactcaga	gaagaaaagg	aagagggcca	tcaaacctcat	1020
tgtcactgtc	ctggccatgt	acctgatctg	cttccactct	agtaaccttc	tgcttgtggg	1080
gcattattht	ctgattaaga	gccagggcca	gagccatgtc	tatgccctgt	acattgtagc	1140
cctctgcctc	tctaccctta	acagctgcat	cgacctcttt	gtctattact	ttgtttcaca	1200
tgatttcagg	gatcatgcaa	agaacgctct	cctttgccga	agtgtccgca	ctgtaaaagca	1260
gatgcaagta	tccctcacct	caaagaaaca	ctccaggaaa	tccagctctt	actcttcaag	1320
ttcaaccact	gttaagacct	cctattgagt	tttccaggtc	ctcagatggg	aattgcacag	1380
taggatcttg	aacctgttta	atgthtatgag	gactgtctctg	ttatttccca	atcaaaaagg	1440
tctcaccaca	taccatgtgg	atgcagcacc	tctcaggatt	gctaggagct	cccctgtttg	1500
catgagaaaa	gtagtcccc	aaattaacat	cagtgtctgt	ttcagaatct	ctctactcag	1560
atgaccccag	aaactgaaac	aacagaagca	gacttttcag	aagatgggtga	agacagaaac	1620
ccagtaactt	gcaaaaagta	gacttgggtg	gaagactcac	ttctcagctg	aaattatata	1680
tatacacata	tataatthtt	acatctggga	tcatgataga	cttgttaggg	cttcaaggcc	1740
ctcagagatg	atcagtcocaa	ctgaacgacc	ttacaaatga	ggaaaccaag	ataaatgagc	1800
tgccagaatc	aggtttccaa	tcaacagcag	tgagttggga	ttggacagta	gaatttcaat	1860
gtccagtgag	tgaggttctt	gtaccacttc	atcaaaatca	tggatcttgg	ctgggtgcgg	1920
tgccctcatg	ctgtaatcct	agcactttgg	gaggctgagg	caggcaatca	cttgaggcca	1980
ggagttcgag	accagcctgg	ccatcatggc	gaaacctcat	ctctactaaa	aatacaaaaag	2040
ttaaccaggt	gtgtgggtgca	cgthttgtaat	cccagttact	caggaggctg	aggcacaaga	2100
attgagatc	actthtaactc	aggaggcaga	ggttgacgtg	agccgagatt	gcaccactgc	2160
actccagctt	gggtgatata	ataaaataaa	atagtcgtga	atcttgttca	aaatgcagat	2220
tcctcagatt	caataatgag	agctcagact	gggaacaggg	cccaggaatc	tgtgtggtac	2280
aaacctgcat	ggtgtthtatg	cacacagaga	tttgagaacc	attgttctga	atgctgcttc	2340
catttgacaa	agtgccgtga	taaththttga	aaagagaagc	aaacaatggg	gtctctthta	2400
tgthtcagctt	ataatgaaat	ctgthttgtg	acttattagg	actthtgaatt	atthctthtat	2460
taacctctg	agththttgta	tgtattatta	ttaaagaaaa	atgcaatcag	gattthtaaac	2520
atgtaaatatc	aaaththttgta	taactthttga	tgacttcagt	gaaaththtca	ggtagtctga	2580
gtaaatagatt	gththttgccac	ttagaatagc	atthttgccact	tagtaththta	aaaaataatt	2640
gthttggagat	thattgtcag	ththttgtcac	thgtthattcta	atacaaaatt	ataaagcctt	2700
cagagggthtt	ggaccacatc	tctthttgaaa	atagthttgca	acataththta	gagatacttg	2760
atgccaataat	gactthtatatc	aacgattgtga	ththttgtactt	thaaaaataa	thaththttatt	2820
gtgtaattgga	ththataaata	acaaaaththt	ththttacaact	thaaaaaaaa	aaaaaa	2876

```

<210> 61
<211> 1668
<212> DNA
<213> Homo sapiens
<400> 61
gggagataac tcgtgctcac aggaagccac gcacccttga aaggcaccgg gtccttctta      60
gcatcgtgct tcctgagcaa gcctggcatt gcctcacaga ccttcctcag agccgctttc      120
agaaaagcaa gctgcttctg gttgggcca gacctgcctt gaggagcctg tagagttaaa      180
aatgaacccc cacggatata gcagacacca cctcogatga aagcatatac agcaattact      240
atctgtatga aagtatcccc aagccttgca ccaaagaagg catcaaggca tttggggagc      300
tcttcctgcc ccactgtat tccttgggtt ttgtatttgg tctgcttggg aattctgtgg      360
tggttctggg cctgttcaaaa tacaagcggc tcagggtccat gactgatgtg tacctgctca      420
accttgccat ctcggatctg ctcttctgtt tttccctccc tttttggggc tactatgcag      480
cagaccagtg ggtttttggg ctaggctctg gcaagatgat ttcttggatg tacttgggtg      540
gcttttacag tggcatattc tttgtcatgc tcatgagcat tgatagatac ctggcaattg      600
tgcacgcggg gttttccttg agggcaagga ccttgactta tggggtcac accagtttgg      660
ctacatggtc agtggctgtg ttcgcctccc ttcttggctt tctgttcagc acttggtata      720
ctgagcgcaa ccataacctac tgcaaaacca agtactctct caactccacg acgtggaagg      780
ttctcagctc cctggaaatc aacattctcg gattgggtgat ccccttaggg atcatgctgt      840
tttgcactc catgatcctc aggaccttgc agcattgtaa aatgagaag aagaacaagg      900
cggtgaaagat gatctttgcc gtggtgggtcc tcttccttgg gttctggaca ccttacaaca      960
tagtgctctt cctagagacc ctggtgggagc tagaagtcct tcaggactgc acctttgaaa     1020
gatacttggg ctatgccatc caggccacag aaactctggc ttttgttcac tgctgcctta     1080
atcccatcat ctactttttt ctgggggaga aatttcgcaa gtacatccta cagctcttca     1140
aaacctgcag gggccttttt gtgctctgcc aatactgtgg gctcctccaa atttactctg     1200
ctgacacccc cagctcatct tacacgcagt ccaccatgga tcatgatctt catgatgctc     1260
tgtagaaaaa tgaatgggtg aatgcagag tcaatgaact tccacattc agagcttact     1320
taaaattgta ttttggtaag agatccctga gccagtgctc ggaggaaggc ttacaccac     1380
agtggaaaga cagcttctca tcctgcaggc agctttttct ctcccactag acaagtccag     1440
cctggcaagg gttcacctgg gctcagcctg ccttccctac accaggcttg cctgcaggca     1500
tgagtcagtc tgatgagaac tctgagcagt gcttgaatga agttgtaggt aatattgcaa     1560
ggcaagagact attcccttct aacctgaact gatgggtttc tccagagggg attgcagagt     1620
actggetgat ggagtaaact gctacctttt gctgtggcaa atggggccc     1668

```

```

<210> 62
<211> 1679
<212> DNA
<213> Homo sapiens
<400> 62
gtttgttggc tgccggcagca ggtagcaaag tgacgcccag ggctgagtg ctccagtagc      60
caccgcatct ggagaaccag cggttaccat ggaggggatc agtatataca cttcagataa     120
ctacaccgag gaaatgggct caggggacta tgactccatg aaggaaccct gtttccgtga     180
agaaaatgct aatttcaata aaatcttctt gccaccatc tactccatca tcttcttaac     240
tggcattgtg ggcaatggat tggatcctct ggtcatgggt taccagaaga aactgagaag     300
catgacggac aagtacaggc tgcacctgtc agtggccgac ctctcttttg tcatcacgct     360
tcccttctgg gcagttgatg ccgtggcaaa ctggtacttt gggaaacttc tatgcaaggc     420
agtccatgtc atctacacag tcaaccteta cagcagtgtc ctcatcctgg ccttcatcag     480
tctggcaagc tacttgcca tcgtccacgc caccaacagt cagaggccaa ggaagctgtt     540
ggctgaaaag gtggtctatg ttggcgtctg gatccctgcc ctctgctga ctattccga     600
cttcatcttt gccaacgtca gtgaggcaga tgacagatat atctgtgacc gcttctaccc     660
caatgacttg tgggtggttg tgttccagtt tcagcacatc atgggtggcc ttatcctgcc     720
tggattgtc atcctgtcct gctattgcat tatcatctcc aagctgtcac actccaaggg     780
ccaccagaag cgcaaggccc tcaagaccac agtcacctc atcctggctt tcttcgctg     840
ttggctgctt tactacattg ggatcagcat cgactccttc atcctctggg aatcatcaa     900
gcaaggggtg gagtttgaga aactgtgca caagtggatt tccatcaccg aggccctagc     960

```

tttcttccac	tgttgtctga	accccatcct	ctatgctttc	cttggagcca	aattttaaac	1020
ctctgcccag	cacgcactca	cctctgtgag	cagaggggtcc	agcctcaaga	tcctctccaa	1080
aggaaagcga	ggtggacatt	catctgtttc	cactgagctct	gagtcttcaa	gttttccactc	1140
cagctaacac	agatgtaaaa	gacttttttt	tatacgataa	ataacttttt	tttaagttac	1200
acatttttca	gatataaaag	actgaccaat	attgtacagt	ttttattgct	tgttggattt	1260
ttgtcttgtg	tttctttagt	ttttgtgaag	tttaattgac	ttatttataat	aaattttttt	1320
tgtttcataat	tgatgtgtgt	ctaggcagga	cctgtggcca	agttcttagt	tgctgtatgt	1380
ctcgtggtag	gactgtagaa	aagggaaactg	aacattccag	agcgtgtagt	gaatcacgta	1440
aagctagaaa	tgatccccag	ctgtttatgc	atagataatc	tctccattcc	cgtggaacgt	1500
ttttcctggt	cttaagacgt	gattttgctg	tagaagatgg	cacttataac	caaagcccaa	1560
agtggatatag	aaatgctggg	ttttcagttt	tcaggagtgg	gttgatttca	gcacctacag	1620
tgtaacagct	tgattaagt	tgtaataaaa	agtacatggt	aaacttactt	agtgttatg	1679

<210> 63

<211> 2859

<212> DNA

<213> Homo sapiens

<400> 63

cattcagaga	cagaaggtgg	atagacaaat	ctccaccttc	agactggtag	gctcctccag	60
aagccatcag	acaggaagat	gtgaaaatcc	ccagcactca	tcccagaatc	actaagtggc	120
acctgtcctg	ggccaaagtc	ccaggacaga	cctcattggt	cctctgtggg	aatacctccc	180
caggagggca	tcctggattt	cccccttgca	accagggtca	gaagtttcat	cgtaaggtt	240
gtttcatctt	tttttccctg	tctaacagct	ctgactacca	cccaaccttg	aggcacagt	300
aagacatcgg	tggccactcc	aataacagca	ggtcacagct	gctcttctgg	agggtgctcta	360
caggtgaaaa	gcccagcgac	ccagtcagga	tttaagttta	cctcaaaaat	ggaagatttt	420
aacatggaga	gtgacagctt	tgaagatttc	tggaaagggtg	aagatcttag	taattacagt	480
tacagctcta	ccctgcccc	ttttctacta	gatgccgcc	catgtgaacc	agaatccctg	540
gaaatcaaca	agtattttgt	ggtcattatc	tatgccctgg	tattcctgct	gagcctgctg	600
ggaaactccc	tcgtgatgct	ggtcatctta	tacagcagg	tcggccgctc	cgtaactgat	660
gctacctgc	tgaacctagc	cttggccgac	ctactctttg	ccctgacctt	gccccctctg	720
gcccctcca	aggtgaatgg	ctggattttt	ggcacattcc	tgtgcaaggt	ggtctcactc	780
ctgaaggaag	tcaacttcta	tagtggcatc	ctgctactgg	cctgcatcag	tgtggaccctg	840
tacotggcca	ttgtccatgc	cacacgcaca	ctgaccaga	agcgtactt	ggtcaaatc	900
atatgtctca	gcactctggg	tctgtccttg	ctcctggccc	tgctctctt	acttttccga	960
aggaccgtct	actcatccaa	tgtagccca	gcctgctatg	aggacatggg	caacaataca	1020
gcaaactggc	ggatgctggt	acggatcctg	cccagtcct	ttggcttcat	cgtaaccctg	1080
ctgatcatgc	tgttctgcta	cggatccacc	ctgctacgc	tgtttaaggc	ccacatgggg	1140
cagaagcacc	gggcatgcg	ggtcatcttt	gctgtcgtcc	tcactctcct	gctctgctgg	1200
ctgccctaca	acctggctct	gctggcagac	acctcatga	ggaccaggt	gatccaggag	1260
acctgtgagc	gcccgaatca	catcgaccgg	gctctggatg	ccaccgagat	tctgggcatc	1320
cttcacagct	gctcaacct	cctcatctac	gccttcattg	gccagaagtt	tcgcatgga	1380
ctcctcaaga	ttctagctat	acatggcttg	atcagcaagg	actcctgccc	caaagacagc	1440
aggccttct	ttgttggtc	ttcttcaggg	caacttcca	ctactctcta	agacctctg	1500
cctaagtgca	gcccctggg	gttctctcct	tctcttcaca	gtcacattcc	aagcctcatg	1560
tccactggtt	cttcttggtc	tcagtgtcaa	tgcagcccc	attgtgggtca	caggaagtag	1620
aggaggccac	gttcttacta	gtttcccttg	catggtttag	aaagcttgcc	ctgggtgctc	1680
acccttggc	ataattacta	tgatatttgc	tggagctctg	cccatcctgc	ccctgagccc	1740
atggcactct	atgttctaag	aagtgaaaat	ctacactcca	gtgagacagc	tctgcatact	1800
cattaggatg	gctagtatca	aaagaaagaa	aatcaggctg	gccaacgggg	tgaaacctctg	1860
tctctactaa	aaatacaaaa	aaaaaaaaaa	attagccggg	cgtgggtggg	agtgcctgta	1920
atcacagcta	cttgggaggg	tgagatggga	gaatcacttg	aaccggggag	gcagaggttg	1980
cagtgaaccg	agattgtgcc	cctgcactcc	agcctgagcg	acagtgagac	tctgtctcag	2040
tccatgaaga	tgtagaggag	aaactggaac	tctcgagcgt	tgctgggggg	gattgtaaaa	2100
tggtgtgacc	actgcagaag	acagtatggc	agcttctctc	aaaacttcag	acatagaatt	2160
aacacatgat	cttgcatttc	cacttatagg	aattgaccca	caagaaatga	aagcagggac	2220
ttgaacccat	attgtacac	caatattcat	agcagcttat	tcacaagacc	caaaaggcag	2280

aagcaaccca	aatgttcatc	aatgaatgaa	tgaatggcta	agcaaaatgt	gatatgtacc	2340
taacgaagta	tccttcagcc	tgaagagga	atgaagtact	catacatggt	acaacacgga	2400
cgaaccttga	aaactttatg	ctaagtgaaa	taagccagac	atcaacagat	aaatagttta	2460
tgattccacc	tacatgaggt	actgagagtg	aacaaattta	cagagacaga	aagcagaaca	2520
gtgattacca	gggactgagg	ggaggggagc	atgggaagtg	acggtttaat	gggcacaggg	2580
tttatgttta	ggatgttgaa	aaagtctctg	agataaacag	tagtgatagt	tgtaccgcaa	2640
tgtgacttaa	tgccactaaa	ttgacactta	aaaatggttt	aaatggtaa	ttttgttatg	2700
tatattttat	atcaatttaa	aaaaaacct	gagcccaaaa	aggtatttta	atcaccaagg	2760
ctgattaaac	caaggctaga	accacctgcc	tatatttttt	gttaaataat	ttcattcaat	2820
atcttttttt	taataaacca	tttttacttg	ggtgtttat			2859

<210> 64
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 64
 tgtgcgcgcg gccagagcag gtgcgca

27

<210> 65
 <211> 26
 <212> DNA
 <213> Homo Sapiens
 <400> 65
 gaggatccgt caaccacaag ggtctc

26

<210> 66
 <211> 27
 <212> DNA
 <213> Homo Sapiens
 <400> 66
 tgtgcgcgcg gctgatcat ccggtct

27

<210> 67
 <211> 26
 <212> DNA
 <213> Homo Sapiens
 <400> 67
 gaggatccga cataccgctc gtgaca

26

<210> 68
 <211> 28
 <212> DNA
 <213> Homo Sapiens
 <400> 68
 tgtgcgcgca gtgtccgcac tgtaaagc

28

<210> 69
 <211> 26
 <212> DNA
 <213> Homo Sapiens

<400> 69
gaggatccat aggaggtcct aacagt 26

<210> 70
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 70
tgtgcgcgcg gcctttttgt gctctgc 27

<210> 71
<211> 26
<212> DNA
<213> Homo Sapiens
<400> 71
gaggatccca gagcatcatg aagatc 26

<210> 72
<211> 28
<212> DNA
<213> Homo Sapiens
<400> 72
tgtgcgcgcg gcttgatcag caaggac 28

<210> 73
<211> 26
<212> DNA
<213> Homo Sapiens
<400> 73
gaggatccga gagtagtgga agtgtg 26

<210> 74
<211> 27
<212> DNA
<213> Homo Sapiens
<400> 74
tgtgcgcgcg ggtccagcct caagatc 27

<210> 75
<211> 26
<212> DNA
<213> Homo Sapiens
<400> 75
gaggatccgc tggagtgaaa acttga 26

<210> 76
<211> 5616
<212> DNA
<213> Homo Sapiens
<400> 76

ccccggcgca	gcgcggccgc	agcagcctcc	gccccccgca	cggtgtgagc	gcccgacgcg	60
gccgaggcgg	ccggagtcce	gagctagccc	cgggcgccgc	cgccgcccag	accggacgac	120
aggccacctc	gtcggcgtcc	gcccgagtcc	ccgcctcgcc	gccaacgcca	caaccaccgc	180
gcacggcccc	ctgactccgt	ccagtattga	tggggagagc	cggagcgagc	tcttcgggga	240
gcagcgatgc	gacctccgg	gacggccggg	gcagcgctcc	tggcgctgct	ggctgcgtc	300
tgcccggcga	gtcgggctct	ggaggaaaag	aaagtttgcc	aaggcacgag	taacaagctc	360
acgcagttgg	gcacttttga	agatcatttt	ctcagcctcc	agaggatggt	caataactgt	420
gaggtggctc	ttgggaattt	ggaaattacc	tatgtgcaga	ggaattatga	tctttccttc	480
ttaaagacca	tccaggaggt	ggctggttat	gtcctcattg	ccctcaacac	agtggagcga	540
attcctttgg	aaaacctgca	gatcatcaga	ggaaatatgt	actacgaaaa	ttcctatgcc	600
ttagcgatcc	tatctaacta	tgatgcaaat	aaaaccggac	tgaaggagct	gcccagtaga	660
aatttacagg	aaatcctgca	tggcgccgtg	cggttcagca	acaaccctgc	cctgtgcaac	720
gtggagagca	tccagtgccg	ggacatagtc	agcagtgact	ttctcagcaa	catgtcgatg	780
gacttccaga	accacctggg	cagctgccaa	aagtgtgatc	caagctgtcc	caatgggagc	840
tgctgggggtg	caggagagga	gaactgccag	aaactgacca	aatcatctg	tgcccagcag	900
tgctccgggc	gctgcctggt	caagtcccc	agtgactgct	gccacaacca	gtgtgctgca	960
ggctgcacag	gcccccgggg	gagcgactgc	ctggtctgcc	gcaaattccg	agcgaagcc	1020
acgtgcaagg	acacctgccc	cccactcatg	ctctacaacc	ccaccacgta	ccagatggat	1080
gtgaaccccg	agggcaata	cagctttggt	gccacctgcg	tgaagaagtg	tccccgtaat	1140
tatgtgggtg	cagatcacgg	ctcgtgcgtc	cgagcctgtg	gggcccagag	ctatgagatg	1200
gaggaagacg	gctgcccga	gtgtaagaag	tgcgaagggc	cttgccgcaa	agtgtgtaac	1260
ggaataggta	ttggtgaatt	taaagactca	ctctccataa	atgctacgaa	tattaaacac	1320
ttcaaaaact	gcactccat	cagtggcgat	ctccacatcc	tgccggtggc	athtaggggt	1380
gactccttca	cacatactcc	tctctggat	ccacagaaac	tggatattct	gaaaaccgta	1440
aaggaaatca	cagggttttt	gctgattcag	gcttggcctg	aaaacaggac	ggacctccat	1500
gcctttgaga	acctagaaat	catacgccgc	aggaccaagc	aacatggtca	gttttctctt	1560
gcagtcgtca	gcctgaacat	aacatccttg	ggattacgct	ccctcaagga	gataagtgat	1620
ggagatgtga	taatttcagg	aaacaaaaat	ttgtgctatg	caaatacaat	aaactggaaa	1680
aaactgtttg	ggacctccgg	tcagaaaacc	aaaattataa	gcaacagagg	tgaaaacagc	1740
tgcaaggcca	caggccaggt	ctgccatgcc	ttgtgctccc	ccgagggctg	ctggggcccc	1800
gagcccaggg	actgcgtctc	ttgccggaat	gtcagccgag	gcagggaatg	cgtggacaa	1860
tgcaaccttc	tggagggtga	gccaagggag	tttgtggaga	actctgagtg	catacagtg	1920
caccagaggt	gcctgcctca	ggccatgaac	atcacctgca	caggacgggg	accagacaac	1980
tgtatccagt	gtgcccacta	cattgacggc	cccactgcg	tcaagacctg	cccggcagga	2040
gtcatgggg	aaaacaacac	cctggtctgg	aagtacgcag	acgccggcca	tgtgtgccac	2100
ctgtgccatc	caaatctgac	ctacggatgc	actggggccag	gtcctgaagg	ctgtccaacg	2160
aatggggccta	agatcccgtc	catcgccact	gggatgggtg	gggcccctct	cttgcctg	2220
gtggtggccc	tggggatcgg	cctcttcctg	cgaaaggccc	acatcgttct	gaagcgacg	2280
ctgcccaggg	tgctgcagga	gagggagctt	gtggagcctc	ttacaccag	tggagaagct	2340
ccaaccaag	ctctcttgag	gatcttgaag	gaaactgaat	tcaaaaagat	caaagtgtct	2400
ggctccgggtg	cgttccggcac	ggtgtataag	ggactctgga	tcccagaagg	tgagaaagt	2460
aaaattcccc	tcgctatcaa	ggaattaaga	gaagcaacat	ctccgaaagc	caacaaggaa	2520
atctcctgat	aagcctacgt	gatggccagc	gtggacaacc	cccacgtgtg	ccgcctgctg	2580
ggcatctgcc	tcacctccac	cgtgcagctc	atcacgcagc	tcatgccctt	cggtgcctc	2640
ctggactatg	tccgggaaca	caaagacaat	attggctccc	agtacctgct	caactggtgt	2700
gtgcagatcg	caaagggcat	gaactacttg	gaggaccgtc	gcttgggtgca	ccgcgacctg	2760
gcagccagga	acgtactggg	gaaaacaccg	cagcatgtca	agatcacaga	ttttgggctg	2820
gccaactgca	tgggtgcgga	agagaaaaga	taccatgcag	aaggaggcaa	agtgcctatc	2880
aagtggatgg	cattggaatc	aattttacac	agaatctata	cccaccagag	tgatgtctgg	2940
agctacgggg	tgaccgtttg	ggagttgatg	acctttggat	ccaagccata	tgacggaatc	3000
cctgccagcg	agatctcctc	catcctggag	aaaggagaac	gcctccctca	gccaccata	3060
tgtaccatcg	atgtctacat	gatcatggtc	aagtgtgga	tgatagacgc	agatagtgc	3120
ccaaagttcc	gtgagttgat	catogaatc	tccaaaatgg	cccagacctc	ccagcgctac	3180
cttgtcattc	agggggatga	aagaatgcat	ttgccaaagc	ctacagactc	caacttctac	3240
cgtgccttga	tggatgaaga	agacatggac	gacgtgggtg	atgccgacga	gtacctcatc	3300
ccacagcagg	gcttcttcag	cagcccctcc	acgtcacgga	ctcccctctc	gagctctctg	3360
agtgcacaaca	gcaacaatcc	caccgtggct	tgcattgata	gaaatgggct	gcaaagctgt	3420

```

cccatcaagg aagacagctt cttgcagcga tacagctcag accccacagg cgccttgact 3480
gaggacagca tagacgacac cttcctccca gtgcctgaat acataaacca gtccgttccc 3540
aaaaggcccg ctggctctgt gcagaatcct gtctatcaca atcagcctct gaaccccgcg 3600
cccagcagag acccactacta ccaggacccc cacagcactg cagtgggcaa ccccgagtat 3660
ctcaaacctg tccagcccaac ctgtgtcaac agcacattcg acagccctgc ccaactgggce 3720
cagaaaggca gccaccaaat tagcctggac aacctgact accagcagga cttctttccc 3780
aaggaagcca agccaaatgg catctttaag ggctccacag ctgaaaatgc agaataccta 3840
agggctcggc cacaaagcag tgaatttatt ggagcatgac cacggaggat agtatgagcc 3900
ctaaaaatcc agactctttc gataccagg accaagccac agcaggctct ccatcccaac 3960
agccatgccc gcattagctc ttagaccac agactggttt tgcaacgttt acaccgacta 4020
ggcaggaagt acttccacct cgggcacatt ttgggaagtt gcattccttt gtcttcaaac 4080
tgtgaagcat ttacagaaac gcattccagca agaattttgt ccttttgagc agaaatttat 4140
ctttcaaga ggtatatttg aaaaaaaaaa aaagtatatg tgaggatttt tattgattgg 4200
ggatcttggg gtttttcatt gtcgctattg atttttactt caatgggctc ttccaacaag 4260
gaagaagctt gctggtagca cttgctacc tgagttcatc caggccaac tgtgagcaag 4320
gagcacaagc cacaaagtct ccagaggatg cttgattcca gtggttctgc tcaaggctt 4380
ccactgcaa acactaaaga tccaagaagg cttcatggc ccagcaggc cggatcggta 4440
ctgtatcaag tcatggcagg tacagtagga taagccactc tgtcccttcc tgggcaaaga 4500
agaaacggag gggatggaat tcttcttag acttactttt gtaaaaatgt cccacggta 4560
cttactcccc actgatggac cagtggtttc cagtcatgag cgttagactg acttgtttgt 4620
cttccattcc attgttttga aactcagtat gctgcccctg tcttgctgtc atgaaatcag 4680
caagagagga tgacacatca aataataact cggattccag ccacattgg attcatcagc 4740
atitggacca atagcccaca gctgagaatg tggaaatcct aaggatagca ccgcttttgt 4800
tctcgcaaaa acgtatctcc taatttgagg ctccagatgaa atgcattcagg tcctttgggg 4860
catagatcag aagactacaa aaatgaagct gctctgaaat ctcttttagc catcacccca 4920
acccccaaa attagtttgt gttacttatg gaagatagtt ttctctttt acttcacttc 4980
aaaagctttt tactcaaga gtatatgttc cctccaggtc agctgcccc aaaccccctc 5040
cttacgcttt gtcacacaaa aagtgtctct gccttgagtc atctattcaa gcaattacag 5100
ctctggccac aacagggcat ttacagggtg cgaatgacag tagcattatg agtagtgtgg 5160
aattcaggta gtaaatatga aactagggtt tgaatttgat aatgctttca caacatttgc 5220
agatgtttta gaaggaaaa agttccttcc taaaataatt tctctacaat tgggaagttg 5280
gaagattcag ctagttagga gccaccttt ttctctaate tgtgtgtgcc ctgtaacctg 5340
actggttaac agcagtoctt tgtaaacagt gttttaaact ctctagtca atatccacc 5400
catccaattt atcaaggaag aaatggttca gaaaatattt tcagcctaca gttatgttca 5460
gtcacacaca catacaaaat gtctcttttg cttttaaagt aatttttgac tcccagatca 5520
gtcagagccc ctacagcatt gttaagaaag tatttgattt ttgtctcaat gaaaataaaa 5580
ctatattcat ttccactcta aaaaaaaaaa aaaaaa 5616

```

```

<210> 77
<211> 12
<212> PRT
<213> Homo Sapiens
<400> 78
Gly Gly Ser Gly Ser Glu Asn Leu Tyr Phe Gln Leu
          5                      10

```

```

<210> 78
<211> 1291
<212> PRT
<213> Homo sapiens
<400> 78
Met Ala Gly Ala Ala Ser Pro Cys Ala Asn Gly Cys Gly Pro Gly Ala
          5                      10                      15
Pro Ser Asp Ala Glu Val Leu His Leu Cys Arg Ser Leu Glu Val Gly
          20                      25                      30

```

Thr Val Met Thr Leu Phe Tyr Ser Lys Lys Ser Gln Arg Pro Glu Arg
 35 40 45
 Lys Thr Phe Gln Val Lys Leu Glu Thr Arg Gln Ile Thr Trp Ser Arg
 50 55 60
 Gly Ala Asp Lys Ile Glu Gly Ala Ile Asp Ile Arg Glu Ile Lys Glu
 65 70 75 80
 Ile Arg Pro Gly Lys Thr Ser Arg Asp Phe Asp Arg Tyr Gln Glu Asp
 85 90 95
 Pro Ala Phe Arg Pro Asp Gln Ser His Cys Phe Val Ile Leu Tyr Gly
 100 105 110
 Met Glu Phe Arg Leu Lys Thr Leu Ser Leu Gln Ala Thr Ser Glu Asp
 115 120 125
 Glu Val Asn Met Trp Ile Lys Gly Leu Thr Trp Leu Met Glu Asp Thr
 130 135 140
 Leu Gln Ala Pro Thr Pro Leu Gln Ile Glu Arg Trp Leu Arg Lys Gln
 145 150 155 160
 Phe Tyr Ser Val Asp Arg Asn Arg Glu Asp Arg Ile Ser Ala Lys Asp
 165 170 175
 Leu Lys Asn Met Leu Ser Gln Val Asn Tyr Arg Val Pro Asn Met Arg
 180 185 190
 Phe Leu Arg Glu Arg Leu Thr Asp Leu Glu Gln Arg Ser Gly Asp Ile
 195 200 205
 Thr Tyr Gly Gln Phe Ala Gln Leu Tyr Arg Ser Leu Met Tyr Ser Ala
 210 215 220
 Gln Lys Thr Met Asp Leu Pro Phe Leu Glu Ala Ser Thr Leu Arg Ala
 225 230 235 240
 Gly Glu Arg Pro Glu Leu Cys Arg Val Ser Leu Pro Glu Phe Gln Gln
 245 250 255
 Phe Leu Leu Asp Tyr Gln Gly Glu Leu Trp Ala Val Asp Arg Leu Gln
 260 265 270
 Val Gln Glu Phe Met Leu Ser Phe Leu Arg Asp Pro Leu Arg Glu Ile
 275 280 285
 Glu Glu Pro Tyr Phe Phe Leu Asp Glu Phe Val ThrP he Leu Phe Ser
 290 295 300
 Lys Glu Asn Ser Val Trp Asn Ser Gln Leu Asp Ala Val Cys Pro Asp
 305 310 315 320
 Thr Met Asn Asn Pro Leu Ser His Tyr Trp Ile Ser Ser Ser His Asn
 325 330 335
 Thr Tyr Leu Thr Gly Asp Gln Phe Ser Ser Glu Ser Ser Leu Glu Ala
 340 345 350
 Tyr Ala Arg Cys Leu Arg Met Gly Cys Arg Cys Ile Glu Leu Asp Cys
 355 360 365
 Trp Asp Gly Pro Asp Gly Met Pro Val Ile Tyr His Gly His Thr Leu
 370 375 380
 Thr Thr Lys Ile Lys Phe Ser Asp Val Leu His Thr Ile Lys Glu His
 385 390 395 400
 Ala Phe Val Ala Ser Glu Tyr Pro Val Ile Leu Ser Ile Glu Asp His
 405 410 415
 Cys Ser Ile Ala Gln Gln Arg Asn Met Ala Gln Tyr Phe Lys Lys Val
 420 425 430
 Leu Gly Asp Thr Leu Leu Thr Lys Pro Val Glu Ile Ser Ala Asp Gly
 435 440 445
 Leu Pro Ser Pro Asn Gln Leu Lys Arg Lys Ile Leu Ile Lys His Lys
 450 455 460
 Lys Leu Ala Glu Gly Ser Ala Tyr Glu Glu Val Pro Thr Ser Met Met
 465 470 475 480
 Tyr Ser Glu Asn Asp Ile Ser Asn Ser Ile Lys Asn Gly Ile Leu Tyr

				485					490					495			
Leu	Glu	Asp	Pro	Val	Asn	His	Glu	Trp	Tyr	Pro	His	Tyr	Phe	Val	Leu		
			500						505					510			
Thr	Ser	Ser	Lys	Ile	Tyr	Tyr	Ser	Glu	Glu	Thr	Ser	Ser	Asp	Gln	Gly		
			515						520					525			
Asn	Glu	Asp	Glu	Glu	Glu	Pro	Lys	Glu	Val	Ser	Ser	Ser	Thr	Glu	Leu		
			530				535						540				
His	Ser	Asn	Glu	Lys	Trp	Phe	His	Gly	Lys	Leu	Gly	Ala	Gly	Arg	Asp		
545					550					555					560		
Gly	Arg	His	Ile	Ala	Glu	Arg	Leu	Leu	Thr	Glu	Tyr	Cys	Ile	Glu	Thr		
				565						570					575		
Gly	Ala	Pro	Asp	Gly	Ser	Phe	Leu	Val	Arg	Glu	Ser	Glu	Thr	Phe	Val		
			580						585					590			
Gly	Asp	Tyr	Thr	Leu	Ser	Phe	Trp	Arg	Asn	Gly	Lys	Val	Gln	His	Cys		
			595					600					605				
Arg	Ile	His	Ser	Arg	Gln	Asp	Ala	Gly	Thr	Pro	Lys	Phe	Phe	Leu	Thr		
						615						620					
Asp	Asn	Leu	Val	Phe	Asp	Ser	Leu	Tyr	Asp	Leu	Ile	Thr	His	Tyr	Gln		
625					630					635					640		
Gln	Val	Pro	Leu	Arg	Cys	Asn	Glu	Phe	Glu	Met	Arg	Leu	Ser	Glu	Pro		
				645						650					655		
Val	Pro	Gln	Thr	Asn	Ala	His	Glu	Ser	Lys	Glu	Trp	Tyr	His	Ala	Ser		
			660							665					670		
Leu	Thr	Arg	Ala	Gln	Ala	Glu	His	Met	Leu	Met	Arg	Val	Pro	Arg	Asp		
								680					685				
Gly	Ala	Phe	Leu	Val	Arg	Lys	Arg	Asn	Glu	Pro	Asn	Ser	Tyr	Ala	Ile		
						695							700				
Ser	Phe	Arg	Ala	Glu	Gly	Lys	Ile	Lys	His	Cys	Arg	Val	Gln	Gln	Glu		
705					710					715					720		
Gly	Gln	Thr	Val	Met	Leu	Gly	Asn	Ser	Glu	Phe	Asp	Ser	Leu	Val	Asp		
					725					730					735		
Leu	Ile	Ser	Tyr	Tyr	Glu	Lys	His	Pro	Leu	Tyr	Arg	Lys	Met	Lys	Leu		
			740						745						750		
Arg	Tyr	Pro	Ile	Asn	Glu	Glu	Ala	Leu	Glu	Lys	Ile	Gly	Thr	Ala	Glu		
							760						765				
Pro	Asp	Tyr	Gly	Ala	Leu	Tyr	Glu	Gly	Arg	Asn	Pro	Gly	Phe	Tyr	Val		
						775							780				
Glu	Ala	Asn	Pro	Met	Pro	Thr	Phe	Lys	Cys	Ala	Val	Lys	Ala	Leu	Phe		
785					790					795					800		
Asp	Tyr	Lys	Ala	Gln	Arg	Glu	Asp	Glu	Leu	Thr	Phe	Ile	Lys	Ser	Ala		
				805						810					815		
Ile	Ile	Gln	Asn	Val	Glu	Lys	Gln	Glu	Gly	Gly	Trp	Trp	Arg	Gly	Asp		
			820							825					830		
Tyr	Gly	Gly	Lys	Lys	Gln	Leu	Trp	Phe	Pro	Ser	Asn	Tyr	Val	Glu	Glu		
			835					840							845		
Met	Val	Asn	Pro	Val	Ala	Leu	Glu	Pro	Glu	Arg	Glu	His	Leu	Asp	Glu		
						855						860					
Asn	Ser	Pro	Leu	Gly	Asp	Leu	Leu	Arg	Gly	Val	Leu	Asp	Val	Pro	Ala		
865					870							875			880		
Cys	Gln	Ile	Ala	Ile	Arg	Pro	Glu	Gly	Lys	Asn	Asn	Arg	Leu	Phe	Val		
				885						890					895		
Phe	Ser	Ile	Ser	Met	Ala	Ser	Val	Ala	His	Trp	Ser	Leu	Asp	Val	Ala		
				900						905					910		
Ala	Asp	Ser	Gln	Glu	Glu	Leu	Gln	Asp	Trp	Val	Lys	Lys	Ile	Arg	Glu		
			915					920							925		
Val	Ala	Gln	Thr	Ala	Asp	Ala	Arg	Leu	Thr	Glu	Gly	Lys	Ile	Met	Glu		
						935									940		

Arg Arg Lys Lys Ile Ala Leu Glu Leu Ser Glu Leu Val Val Tyr Cys
 945 950 955 960
 Arg Pro Val Pro Phe Asp Glu Glu Lys Ile Gly Thr Glu Arg Ala Cys
 965 970 975
 Tyr Arg Asp Met Ser Ser Phe Pro Glu Thr Lys Ala Glu Lys Tyr Val
 980 985 990
 Asn Lys Ala Lys Gly Lys Lys Phe Leu Gln Tyr Asn Arg Leu Gln Leu
 995 1000 1005
 Ser Arg Ile Tyr Pro Lys Gly Gln Arg Leu Asp Ser Ser Asn Tyr Asp
 1010 1015 1020
 Pro Leu Pro Met Trp Ile Cys Gly Ser Gln Leu Val Ala Leu Asn Phe
 1025 1030 1035 1040
 Gln Thr Pro Asp Lys Pro Met Gln Met Asn Gln Ala Leu Phe Met Thr
 1045 1050 1055
 Gly Arg His Cys Gly Tyr Val Leu Gln Pro Ser Thr Met Arg Asp Glu
 1060 1065 1070
 Ala Phe Asp Pro Phe Asp Lys Ser Ser Leu Arg Gly Leu Glu Pro Cys
 1075 1080 1085
 Ala Ile Ser Ile Glu Val Leu Gly Ala Arg His Leu Pro Lys Asn Gly
 1090 1095 1100
 Arg Gly Ile Val Cys Pro Phe Val Glu Ile Glu Val Ala Gly Ala Glu
 1105 1110 1115 1120
 Tyr Asp Ser Thr Lys Gln Lys Thr Glu Phe Val Val Asp Asn Gly Leu
 1125 1130 1135
 Asn Pro Val Trp Pro Ala Lys Pro Phe His Phe Gln Ile Ser Asn Pro
 1140 1145 1150
 Glu Phe Ala Phe Leu Arg Phe Val Val Tyr Glu Glu Asp Met Phe Ser
 1155 1160 1165
 Asp Gln Asn Phe Leu Ala Gln Ala Thr Phe Pro Val Lys Gly Leu Lys
 1170 1175 1180
 Thr Gly Tyr Arg Ala Val Pro Leu Lys Asn Asn Tyr Ser Glu Asp Leu
 1185 1190 1195 1200
 Glu Leu Ala Ser Leu Leu Ile Lys Ile Asp Ile Phe Pro Ala Lys Gln
 1205 1210 1215
 Glu Asn Gly Asp Leu Ser Pro Phe Ser Gly Thr Ser Leu Arg Glu Arg
 1220 1225 1230
 Gly Ser Asp Ala Ser Gly Gln Leu Phe His Gly Arg Ala Arg Glu Gly
 1235 1240 1245
 Ser Phe Glu Ser Arg Tyr Gln Gln Pro Phe Glu Asp Phe Arg Ile Ser
 1250 1255 1260
 Gln Glu His Leu Ala Asp His Phe Asp Ser Arg Glu Arg Arg Ala Pro
 1265 1270 1275 1280
 Arg Arg Thr Arg Val Asn Gly Asp Asn Arg Leu
 1285 1290

<210> 79
 <211> 3054
 <212> PRT
 <213> Homo sapiens
 <400> 79

Met Ala Leu Ile Phe Gly Thr Val Asn Ala Asn Ile Leu Lys Glu Val
 5 10 15
 Phe Gly Gly Ala Arg Met Ala Cys Val Thr Ser Ala His Met Ala Gly
 20 25 30
 Ala Asn Gly Ser Ile Leu Lys Lys Ala Glu Glu Thr Ser Arg Ala Ile
 35 40 45

Met His Lys Pro Val Ile Phe Gly Glu Asp Tyr Ile Thr Glu Ala Asp
50 55 60
Leu Pro Tyr Thr Pro Leu His Leu Glu Val Asp Ala Glu Met Glu Arg
65 70 75 80
Met Tyr Tyr Leu Gly Arg Arg Ala Leu Thr His Gly Lys Arg Arg Lys
85 90 95
Val Ser Val Asn Asn Lys Arg Asn Arg Arg Arg Lys Val Ala Lys Thr
100 105 110
Tyr Val Gly Arg Asp Ser Ile Val Glu Lys Ile Val Val Pro His Thr
115 120 125
Glu Arg Lys Val Asp Thr Thr Ala Ala Val Glu Asp Ile Cys Asn Glu
130 135 140
Ala Thr Thr Gln Leu Val His Asn Ser Met Pro Lys Arg Lys Lys Gln
145 150 155 160
Lys Asn Phe Leu Pro Ala Thr Ser Leu Ser Asn Val Tyr Ala Gln Thr
165 170 175
Trp Ser Ile Val Arg Lys Arg His Met Gln Val Glu Ile Ile Ser Lys
180 185 190
Lys Ser Val Arg Ala Arg Val Lys Arg Phe Glu Gly Ser Val Gln Leu
195 200 205
Phe Ala Ser Val Arg His Met Tyr Gly Glu Arg Lys Arg Val Asp Leu
210 215 220
Arg Ile Asp Asn Trp Gln Gln Glu Thr Leu Leu Asp Leu Ala Lys Arg
225 230 235 240
Phe Lys Asn Glu Arg Val Asp Gln Ser Lys Leu Thr Phe Gly Ser Ser
245 250 255
Gly Leu Val Leu Arg Gln Gly Ser Tyr Gly Pro Ala His Trp Tyr Arg
260 265 270
His Gly Met Phe Ile Val Arg Gly Arg Ser Asp Gly Met Leu Val Asp
275 280 285
Ala Arg Ala Lys Val Thr Phe Ala Val Cys His Ser Met Thr His Tyr
290 295 300
Ser Asp Lys Ser Ile Ser Glu Ala Phe Phe Ile Pro Tyr Ser Lys Lys
305 310 315 320
Phe Leu Glu Leu Arg Pro Asp Gly Ile Ser His Glu Cys Thr Arg Gly
325 330 335
Val Ser Val Glu Arg Cys Gly Glu Val Ala Ala Ile Leu Thr Gln Ala
340 345 350
Leu Ser Pro Cys Gly Lys Ile Thr Cys Lys Arg Cys Met Val Glu Thr
355 360 365
Pro Asp Ile Val Glu Gly Glu Ser Gly Glu Ser Val Thr Asn Gln Gly
370 375 380
Lys Leu Leu Ala Met Leu Lys Glu Gln Tyr Pro Asp Phe Pro Met Ala
385 390 395 400
Glu Lys Leu Leu Thr Arg Phe Leu Gln Gln Lys Ser Leu Val Asn Thr
405 410 415
Asn Leu Thr Ala Cys Val Ser Val Lys Gln Leu Ile Gly Asp Arg Lys
420 425 430
Gln Ala Pro Phe Thr His Val Leu Ala Val Ser Glu Ile Leu Phe Lys
435 440 445
Gly Asn Lys Leu Thr Gly Ala Asp Leu Glu Glu Ala Ser Thr His Met
450 455 460
Leu Glu Ile Ala Arg Phe Leu Asn Asn Arg Thr Glu Asn Met Arg Ile
465 470 475 480
Gly His Leu Gly Ser Phe Arg Asn Lys Ile Ser Ser Lys Ala His Val
485 490 495
Asn Asn Ala Leu Met Cys Asp Asn Gln Leu Asp Gln Asn Gly Asn Phe

			500						505						510
Ile	Trp	Gly	Leu	Arg	Gly	Ala	His	Ala	Lys	Arg	Phe	Leu	Lys	Gly	Phe
		515						520						525	
Phe	Thr	Glu	Ile	Asp	Pro	Asn	Glu	Gly	Tyr	Asp	Lys	Tyr	Val	Ile	Arg
		530				535					540				
Lys	His	Ile	Arg	Gly	Ser	Arg	Lys	Leu	Ala	Ile	Gly	Asn	Leu	Ile	Met
545					550						555				560
Ser	Thr	Asp	Phe	Gln	Thr	Leu	Arg	Gln	Gln	Ile	Gln	Gly	Glu	Thr	Ile
				565						570					575
Glu	Arg	Lys	Glu	Ile	Gly	Asn	His	Cys	Ile	Ser	Met	Arg	Asn	Gly	Asn
				580				585						590	
Tyr	Val	Tyr	Pro	Cys	Cys	Cys	Val	Thr	Leu	Glu	Asp	Gly	Lys	Ala	Gln
		595						600						605	
Tyr	Ser	Asp	Leu	Lys	His	Pro	Thr	Lys	Arg	His	Leu	Val	Ile	Gly	Asn
		610				615					620				
Ser	Gly	Asp	Ser	Lys	Tyr	Leu	Asp	Leu	Pro	Val	Leu	Asn	Glu	Glu	Lys
625					630					635					640
Met	Tyr	Ile	Ala	Asn	Glu	Gly	Tyr	Cys	Tyr	Met	Asn	Ile	Phe	Phe	Ala
				645						650					655
Leu	Leu	Val	Asn	Val	Lys	Glu	Glu	Asp	Ala	Lys	Asp	Phe	Thr	Lys	Phe
			660					665							670
Ile	Arg	Asp	Thr	Ile	Val	Pro	Lys	Leu	Gly	Ala	Trp	Pro	Thr	Met	Gln
		675					680							685	
Asp	Val	Ala	Thr	Ala	Cys	Tyr	Leu	Leu	Ser	Ile	Leu	Tyr	Pro	Asp	Val
		690				695						700			
Leu	Arg	Ala	Glu	Leu	Pro	Arg	Ile	Leu	Val	Asp	His	Asp	Asn	Lys	Thr
705					710					715					720
Met	His	Val	Leu	Asp	Ser	Tyr	Gly	Ser	Arg	Thr	Thr	Gly	Tyr	His	Met
				725					730						735
Leu	Lys	Met	Asn	Thr	Thr	Ser	Gln	Leu	Ile	Glu	Phe	Val	His	Ser	Gly
			740						745						750
Leu	Glu	Ser	Glu	Met	Lys	Thr	Tyr	Asn	Val	Gly	Gly	Met	Asn	Arg	Asp
		755					760					765			
Val	Val	Thr	Gln	Gly	Ala	Ile	Glu	Met	Leu	Ile	Lys	Ser	Ile	Tyr	Lys
		770					775					780			
Pro	His	Leu	Met	Lys	Gln	Leu	Leu	Glu	Glu	Glu	Pro	Tyr	Ile	Ile	Val
785					790					795					800
Leu	Ala	Ile	Val	Ser	Pro	Ser	Ile	Leu	Ile	Ala	Met	Tyr	Asn	Ser	Gly
				805						810					815
Thr	Phe	Glu	Gln	Ala	Leu	Gln	Met	Trp	Leu	Pro	Asn	Thr	Met	Arg	Leu
			820					825							830
Ala	Asn	Leu	Ala	Ala	Ile	Leu	Ser	Ala	Leu	Ala	Gln	Lys	Leu	Thr	Leu
		835					840								845
Ala	Asp	Leu	Phe	Val	Gln	Gln	Arg	Asn	Leu	Ile	Asn	Glu	Tyr	Ala	Gln
		850					855					860			
Val	Ile	Leu	Asp	Asn	Leu	Ile	Asp	Gly	Val	Arg	Val	Asn	His	Ser	Leu
865					870					875					880
Ser	Leu	Ala	Met	Glu	Ile	Val	Thr	Ile	Lys	Leu	Ala	Thr	Gln	Glu	Met
				885						890					895
Asp	Met	Ala	Leu	Arg	Glu	Gly	Gly	Tyr	Ala	Val	Thr	Ser	Glu	Lys	Val
			900					905							910
His	Glu	Met	Leu	Glu	Lys	Asn	Tyr	Val	Lys	Ala	Leu	Lys	Asp	Ala	Trp
		915					920								925
Asp	Glu	Leu	Thr	Trp	Leu	Glu	Lys	Phe	Ser	Ala	Ile	Arg	His	Ser	Arg
		930				935						940			
Lys	Leu	Leu	Lys	Phe	Gly	Arg	Lys	Pro	Leu	Ile	Met	Lys	Asn	Thr	Val
945					950						955				960

Asp Cys Gly Gly His Ile Asp Leu Ser Val Lys Ser Leu Phe Lys Phe
 965 970 975
 His Leu Glu Leu Leu Lys Gly Thr Ile Ser Arg Ala Val Asn Gly Gly
 980 985 990
 Ala Arg Lys Val Arg Val Ala Lys Asn Ala Met Thr Lys Gly Val Phe
 995 1000 1005
 Leu Lys Ile Tyr Ser Met Leu Pro Asp Val Tyr Lys Phe Ile Thr Val
 1010 1015 1020
 Ser Ser Val Leu Ser Leu Leu Leu Thr Phe Leu Phe Gln Ile Asp Cys
 1025 1030 1035 1040
 Met Ile Arg Ala His Arg Glu Ala Lys Val Ala Ala Gln Leu Gln Lys
 1045 1050 1055
 Glu Ser Glu Trp Asp Asn Ile Ile Asn Arg Thr Phe Gln Tyr Ser Lys
 1060 1065 1070
 Leu Glu Asn Pro Ile Gly Tyr Arg Ser Thr Ala Glu Glu Arg Leu Gln
 1075 1080 1085
 Ser Glu His Pro Glu Ala Phe Glu Tyr Tyr Lys Phe Cys Ile Gly Lys
 1090 1095 1100
 Glu Asp Leu Val Glu Gln Ala Lys Gln Pro Glu Ile Ala Tyr Phe Glu
 1105 1110 1115 1120
 Lys Ile Ile Ala Phe Ile Thr Leu Val Leu Met Ala Phe Asp Ala Glu
 1125 1130 1135
 Arg Ser Asp Gly Val Phe Lys Ile Leu Asn Lys Phe Lys Gly Ile Leu
 1140 1145 1150
 Ser Ser Thr Glu Arg Glu Ile Ile Tyr Thr Gln Ser Leu Asp Asp Tyr
 1155 1160 1165
 Val Thr Thr Phe Asp Asp Asn Met Thr Ile Asn Leu Glu Leu Asn Met
 1170 1175 1180
 Asp Glu Leu His Lys Thr Ser Leu Pro Gly Val Thr Phe Lys Gln Trp
 1185 1190 1195 1200
 Trp Asn Asn Gln Ile Ser Arg Gly Asn Val Lys Pro His Tyr Arg Thr
 1205 1210 1215
 Glu Gly His Phe Met Glu Phe Thr Arg Asp Thr Ala Ala Ser Val Ala
 1220 1225 1230
 Ser Glu Ile Ser His Ser Pro Ala Arg Asp Phe Leu Val Arg Gly Ala
 1235 1240 1245
 Val Gly Ser Gly Lys Ser Thr Gly Leu Pro Tyr His Leu Ser Lys Arg
 1250 1255 1260
 Gly Arg Val Leu Met Leu Glu Pro Thr Arg Pro Leu Thr Asp Asn Met
 1265 1270 1275 1280
 His Lys Gln Leu Arg Ser Glu Pro Phe Asn Cys Phe Pro Thr Leu Arg
 1285 1290 1295
 Met Arg Gly Lys Ser Thr Phe Gly Ser Ser Pro Ile Thr Val Met Thr
 1300 1305 1310
 Ser Gly Phe Ala Leu His His Phe Ala Arg Asn Ile Ala Glu Val Lys
 1315 1320 1325
 Thr Tyr Asp Phe Val Ile Ile Asp Glu Cys His Val Asn Asp Ala Ser
 1330 1335 1340
 Ala Ile Ala Phe Arg Asn Leu Leu Phe Glu His Glu Phe Glu Gly Lys
 1345 1350 1355 1360
 Val Leu Lys Val Ser Ala Thr Pro Pro Gly Arg Glu Val Glu Phe Thr
 1365 1370 1375
 Thr Gln Phe Pro Val Lys Leu Lys Ile Glu Glu Ala Leu Ser Phe Gln
 1380 1385 1390
 Glu Phe Val Ser Leu Gln Gly Thr Gly Ala Asn Ala Asp Val Ile Ser
 1395 1400 1405
 Cys Gly Asp Asn Ile Leu Val Tyr Val Ala Ser Tyr Asn Asp Val Asp

1410		1415		1420
Ser Leu Gly Lys Leu Leu Val Gln Lys Gly Tyr Lys Val Ser Lys Ile				
1425		1430		1440
Asp Gly Arg Thr Met Lys Ser Gly Gly Thr Glu Ile Ile Thr Glu Gly				
	1445		1450	1455
Thr Ser Val Lys Lys His Phe Ile Val Ala Thr Asn Ile Ile Glu Asn				
	1460		1465	1470
Gly Val Thr Ile Asp Ile Asp Val Val Val Asp Phe Gly Thr Lys Val				
	1475		1480	1485
Val Pro Val Leu Asp Val Asp Asn Arg Ala Val Gln Tyr Asn Lys Thr				
	1490		1495	1500
Val Val Ser Tyr Gly Glu Arg Ile Gln Lys Leu Gly Arg Val Gly Arg				
1505		1510		1520
His Lys Glu Gly Val Ala Leu Arg Ile Gly Gln Thr Asn Lys Thr Leu				
	1525		1530	1535
Val Glu Ile Pro Glu Met Val Ala Thr Glu Ala Ala Phe Leu Cys Phe				
	1540		1545	1550
Met Tyr Asn Leu Pro Val Thr Thr Gln Ser Val Ser Thr Thr Leu Leu				
	1555		1560	1565
Glu Asn Ala Thr Leu Leu Gln Ala Arg Thr Met Ala Gln Phe Glu Leu				
	1570		1575	1580
Ser Tyr Phe Tyr Thr Ile Asn Phe Val Arg Phe Asp Gly Ser Met His				
1585		1590		1600
Pro Val Ile His Asp Lys Leu Lys Arg Phe Lys Leu His Thr Cys Glu				
	1605		1610	1615
Thr Phe Leu Asn Lys Leu Ala Ile Pro Asn Lys Gly Leu Ser Ser Trp				
	1620		1625	1630
Leu Thr Ser Gly Glu Tyr Lys Arg Leu Gly Tyr Ile Ala Glu Asp Ala				
	1635		1640	1645
Gly Ile Arg Ile Pro Phe Val Cys Lys Glu Ile Pro Asp Ser Leu His				
	1650		1655	1660
Glu Glu Ile Trp His Ile Val Val Ala His Lys Gly Asp Ser Gly Ile				
1665		1670		1680
Gly Arg Leu Thr Ser Val Gln Ala Ala Lys Val Val Tyr Thr Leu Gln				
	1685		1690	1695
Thr Asp Val His Ser Ile Ala Arg Thr Leu Ala Cys Ile Asn Arg Arg				
	1700		1705	1710
Ile Ala Asp Glu Gln Met Lys Gln Ser His Phe Glu Ala Ala Thr Gly				
	1715		1720	1725
Arg Ala Phe Ser Phe Thr Asn Tyr Ser Ile Gln Ser Ile Phe Asp Thr				
	1730		1735	1740
Leu Lys Ala Asn Tyr Ala Thr Lys His Thr Lys Glu Asn Ile Ala Val				
1745		1750		1760
Leu Gln Gln Ala Lys Asp Gln Leu Leu Glu Phe Ser Asn Leu Ala Lys				
	1765		1770	1775
Asp Gln Asp Val Thr Gly Ile Ile Gln Asp Phe Asn His Leu Glu Thr				
	1780		1785	1790
Ile Tyr Leu Gln Ser Asp Ser Glu Val Ala Lys His Leu Lys Leu Lys				
	1795		1800	1805
Ser His Trp Asn Lys Ser Gln Ile Thr Arg Asp Ile Ile Ile Ala Leu				
1810		1815		1820
Ser Val Leu Ile Gly Gly Gly Trp Met Leu Ala Thr Tyr Phe Lys Asp				
1825		1830		1840
Lys Phe Asn Glu Pro Val Tyr Phe Gln Gly Lys Lys Asn Gln Lys His				
	1845		1850	1855
Lys Leu Lys Met Arg Glu Ala Arg Gly Ala Arg Gly Gln Tyr Glu Val				
	1860		1865	1870

Ala Ala Glu Pro Glu Ala Leu Glu His Tyr Phe Gly Ser Ala Tyr Asn
 1875 1880 1885
 Asn Lys Gly Lys Arg Lys Gly Thr Thr Arg Gly Met Gly Ala Lys Ser
 1890 1895 1900
 Arg Lys Phe Ile Asn Met Tyr Gly Phe Asp Pro Thr Asp Phe Ser Tyr
 1905 1910 1915 1920
 Ile Arg Phe Val Asp Pro Leu Thr Gly His Thr Ile Asp Glu Ser Thr
 1925 1930 1935
 Asn Ala Pro Ile Asp Leu Val Gln His Glu Phe Gly Lys Val Arg Thr
 1940 1945 1950
 Arg Met Leu Ile Asp Asp Glu Ile Glu Pro Gln Ser Leu Ser Thr His
 1955 1960 1965
 Thr Thr Ile His Ala Tyr Leu Val Asn Ser Gly Thr Lys Lys Val Leu
 1970 1975 1980
 Lys Val Asp Leu Thr Pro His Ser Ser Leu Arg Ala Ser Glu Lys Ser
 1985 1990 1995 2000
 Thr Ala Ile Met Gly Phe Pro Glu Arg Glu Asn Glu Leu Arg Gln Thr
 2005 2010 2015
 Gly Met Ala Val Pro Val Ala Tyr Asp Gln Leu Pro Pro Lys Asn Glu
 2020 2025 2030
 Asp Leu Thr Phe Glu Gly Glu Ser Leu Phe Lys Gly Pro Arg Asp Tyr
 2035 2040 2045
 Asn Pro Ile Ser Ser Thr Ile Cys His Leu Thr Asn Glu Ser Asp Gly
 2050 2055 2060
 His Thr Thr Ser Leu Tyr Gly Ile Gly Phe Gly Pro Phe Ile Ile Thr
 2065 2070 2075 2080
 Asn Lys His Leu Phe Arg Arg Asn Asn Gly Thr Leu Leu Val Gln Ser
 2085 2090 2095
 Leu His Gly Val Phe Lys Val Lys Asn Thr Thr Thr Leu Gln Gln His
 2100 2105 2110
 Leu Ile Asp Gly Arg Asp Met Ile Ile Ile Arg Met Pro Lys Asp Phe
 2115 2120 2125
 Pro Pro Phe Pro Gln Lys Leu Lys Phe Arg Glu Pro Gln Arg Glu Glu
 2130 2135 2140
 Arg Ile Cys Leu Val Thr Thr Asn Phe Gln Thr Lys Ser Met Ser Ser
 2145 2150 2155 2160
 Met Val Ser Asp Thr Ser Cys Thr Phe Pro Ser Ser Asp Gly Ile Phe
 2165 2170 2175
 Trp Lys His Trp Ile Gln Thr Lys Asp Gly Gln Cys Gly Ser Pro Leu
 2180 2185 2190
 Val Ser Thr Arg Asp Gly Phe Ile Val Gly Ile His Ser Ala Ser Asn
 2195 2200 2205
 Phe Thr Asn Thr Asn Asn Tyr Phe Thr Ser Val Pro Lys Asn Phe Met
 2210 2215 2220
 Glu Leu Leu Thr Asn Gln Glu Ala Gln Gln Trp Val Ser Gly Trp Arg
 2225 2230 2235 2240
 Leu Asn Ala Asp Ser Val Leu Trp Gly Gly His Lys Val Phe Met Ser
 2245 2250 2255
 Lys Pro Glu Glu Pro Phe Gln Pro Val Lys Glu Ala Thr Gln Leu Met
 2260 2265 2270
 Asn Glu Leu Val Tyr Ser Gln Gly Glu Lys Arg Lys Trp Val Val Glu
 2275 2280 2285
 Ala Leu Ser Gly Asn Leu Arg Pro Val Ala Glu Cys Pro Ser Gln Leu
 2290 2295 2300
 Val Thr Lys His Val Val Lys Gly Lys Cys Pro Leu Phe Glu Leu Tyr
 2305 2310 2315 2320
 Leu Gln Leu Asn Pro Glu Lys Glu Ala Tyr Phe Lys Pro Met Met Gly

				2325					2330					2335	
Ala	Tyr	Lys	Pro	Ser	Arg	Leu	Asn	Arg	Glu	Ala	Phe	Leu	Lys	Asp	Ile
				2340					2345					2350	
Leu	Lys	Tyr	Ala	Ser	Glu	Ile	Glu	Ile	Gly	Asn	Val	Asp	Cys	Asp	Leu
				2355					2360					2365	
Leu	Glu	Leu	Ala	Ile	Ser	Met	Leu	Val	Thr	Lys	Leu	Lys	Ala	Leu	Gly
				2370					2375					2380	
Phe	Pro	Thr	Val	Asn	Tyr	Ile	Thr	Asp	Pro	Glu	Glu	Ile	Phe	Ser	Ala
				2385					2390					2395	2400
Leu	Asn	Met	Lys	Ala	Ala	Met	Gly	Ala	Leu	Tyr	Lys	Gly	Lys	Lys	Lys
				2405					2410					2415	
Glu	Ala	Leu	Ser	Glu	Leu	Thr	Leu	Asp	Glu	Gln	Glu	Ala	Met	Leu	Lys
				2420					2425					2430	
Ala	Ser	Cys	Leu	Arg	Leu	Tyr	Thr	Gly	Lys	Leu	Gly	Ile	Trp	Asn	Gly
				2435					2440					2445	
Ser	Leu	Lys	Ala	Glu	Leu	Arg	Pro	Ile	Glu	Lys	Val	Glu	Asn	Asn	Lys
				2450					2455					2460	
Thr	Arg	Thr	Phe	Thr	Ala	Ala	Pro	Ile	Asp	Thr	Leu	Leu	Ala	Gly	Lys
				2465					2470					2475	2480
Val	Cys	Val	Asp	Asp	Phe	Asn	Asn	Gln	Phe	Tyr	Asp	Leu	Asn	Ile	Lys
				2485					2490					2495	
Ala	Pro	Trp	Thr	Val	Gly	Met	Thr	Lys	Phe	Tyr	Gln	Gly	Trp	Asn	Glu
				2500					2505					2510	
Leu	Met	Glu	Ala	Leu	Pro	Ser	Gly	Trp	Val	Tyr	Cys	Asp	Ala	Asp	Gly
				2515					2520					2525	
Ser	Gln	Phe	Asp	Ser	Ser	Leu	Thr	Pro	Phe	Leu	Ile	Asn	Ala	Val	Leu
				2530					2535					2540	
Lys	Val	Arg	Leu	Ala	Phe	Met	Glu	Glu	Trp	Asp	Ile	Gly	Glu	Gln	Met
				2545					2550					2555	2560
Leu	Arg	Asn	Leu	Tyr	Thr	Glu	Ile	Val	Tyr	Thr	Pro	Ile	Leu	Thr	Pro
				2565					2570					2575	
Asp	Gly	Thr	Ile	Lys	Lys	His	Lys	Gly	Asn	Asn	Ser	Gly	Gln	Pro	
				2580					2585					2590	
Ser	Thr	Val	Val	Asp	Asn	Thr	Leu	Met	Val	Ile	Ile	Ala	Met	Leu	Tyr
				2595					2600					2605	
Thr	Cys	Glu	Lys	Cys	Gly	Ile	Asn	Lys	Glu	Glu	Ile	Val	Tyr	Tyr	Val
				2610					2615					2620	
Asn	Gly	Asp	Asp	Leu	Leu	Ile	Ala	Ile	His	Pro	Asp	Lys	Ala	Glu	Arg
				2625					2630					2635	2640
Leu	Ser	Arg	Phe	Lys	Glu	Ser	Phe	Gly	Glu	Leu	Gly	Leu	Lys	Tyr	Glu
				2645					2650					2655	
Phe	Asp	Cys	Thr	Thr	Arg	Asp	Lys	Thr	Gln	Leu	Trp	Phe	Met	Ser	His
				2660					2665					2670	
Arg	Ala	Leu	Glu	Arg	Asp	Gly	Met	Tyr	Ile	Pro	Lys	Leu	Glu	Glu	Glu
				2675					2680					2685	
Arg	Ile	Val	Ser	Ile	Leu	Glu	Trp	Asp	Arg	Ser	Lys	Glu	Pro	Ser	His
				2690					2695					2700	
Arg	Leu	Glu	Ala	Ile	Cys	Ala	Ser	Met	Ile	Glu	Ala	Trp	Gly	Tyr	Asp
				2705					2710					2715	2720
Lys	Leu	Val	Glu	Glu	Ile	Arg	Asn	Phe	Tyr	Ala	Trp	Val	Leu	Glu	Gln
				2725					2730					2735	
Ala	Pro	Tyr	Ser	Gln	Leu	Ala	Glu	Glu	Gly	Lys	Ala	Pro	Tyr	Leu	Ala
				2740					2745					2750	
Glu	Thr	Ala	Leu	Lys	Phe	Leu	Tyr	Thr	Ser	Gln	His	Gly	Thr	Asn	Ser
				2755					2760					2765	
Glu	Ile	Glu	Glu	Tyr	Leu	Lys	Val	Leu	Tyr	Asp	Tyr	Asp	Ile	Pro	Thr
				2770					2775					2780	

Thr Glu Asn Leu Tyr Phe Gln Ser Gly Thr Val Asp Ala Gly Ala Asp
 2785 2790 2795 2800
 Ala Gly Lys Lys Lys Asp Gln Lys Asp Asp Lys Val Ala Glu Gln Ala
 2805 2810 2815
 Ser Lys Asp Arg Asp Val Asn Ala Gly Thr Ser Gly Thr Phe Ser Val
 2820 2825 2830
 Pro Arg Ile Asn Ala Met Ala Thr Lys Leu Gln Tyr Pro Arg Met Arg
 2835 2840 2845
 Gly Glu Val Val Val Asn Leu Asn His Leu Leu Gly Tyr Lys Pro Gln
 2850 2855 2860
 Gln Ile Asp Leu Ser Asn Ala Arg Ala Thr His Glu Gln Phe Ala Ala
 2865 2870 2875 2880
 Trp His Gln Ala Val Met Thr Ala Tyr Gly Val Asn Glu Glu Gln Met
 2885 2890 2895
 Lys Ile Leu Leu Asn Gly Phe Met Val Trp Cys Ile Glu Asn Gly Thr
 2900 2905 2910
 Ser Pro Asn Leu Asn Gly Thr Trp Val Met Met Asp Gly Glu Asp Gln
 2915 2920 2925
 Val Ser Tyr Pro Leu Lys Pro Met Val Glu Asn Ala Gln Pro Thr Leu
 2930 2935 2940
 Arg Gln Ile Met Thr His Phe Ser Asp Leu Ala Glu Ala Tyr Ile Glu
 2945 2950 2955 2960
 Met Arg Asn Arg Glu Arg Pro Tyr Met Pro Arg Tyr Gly Leu Gln Arg
 2965 2970 2975
 Asn Ile Thr Asp Met Ser Leu Ser Arg Tyr Ala Phe Asp Phe Tyr Glu
 2980 2985 2990
 Leu Thr Ser Lys Thr Pro Val Arg Ala Arg Glu Ala His Met Gln Met
 2995 3000 3005
 Lys Ala Ala Ala Val Arg Asn Ser Gly Thr Arg Leu Phe Gly Leu Asp
 3010 3015 3020
 Gly Asn Val Gly Thr Ala Glu Glu Asp Thr Glu Arg His Thr Ala His
 3025 3030 3035 3040
 Asp Val Asn Arg Asn MET et His Thr Leu Leu Gly Val Arg Gln
 3045 3050

<210> 80
 <211> 9
 <212> PRT
 <213> Homo Sapiens
 <400> 80
 Asn Ser Ser Gly Gly Asn Ser Gly Ser
 5

<210> 81
 <211> 2755
 <212> DNA
 <213> Homo sapiens
 <400> 81
 ttaggacggg gcgatggcgg ctgagaggag ctgcgcgtgc gccaacatgt aactgggtggg 60
 atctgcggcg gctcccagat gatggctcgtc ctctctggcg cgacgacct agtgctcgtc 120
 gccgtgggcc catgggtggt gtccgcagcc gcaggtggaa aaaatctaaa atctcctcaa 180
 aaagtagagg tcgacatcat agatgacaac tttatcctga ggtggaacag gagcgatgag 240
 tctgtcggga atgtgacttt ttcattcgat tatcaaaaaa ctgggatgga taattggata 300
 aaattgtctg ggtgtcagaa tattactagt accaaatgca acttttcttc actcaagctg 360
 aatgtttatg aagaaattaa attgcgata agagcagaaa aagaaaacac ttcttcatgy 420
 tatgaggttg actcatttac accatttcgc aaagctcaga ttggtcctcc agaagtacat 480

ttagaagctg	aagataaggc	aatagtgata	cacatctctc	ctggaacaaa	agatagtggt	540
atgtgggctt	tggatggttt	aagctttaca	tatagcttac	ttatctggaa	aaactcttca	600
ggtgtagaag	aaaggattga	aatatattat	tccagacata	aaatttataa	actctcacca	660
gagactactt	attgtctaaa	agttaaagca	gcactactta	cgtcattgaa	aattgggtgc	720
tatagttccag	tacattgtat	aaagaccaca	gttgaaaatg	aactacctcc	accagaaaat	780
atagaagtca	gtgtccaaaa	tcagaactat	gttcttaaat	gggattatac	atatgcaaac	840
atgacctttc	aagttcagtg	gctccacgcc	tttttaaaaa	ggaatcctgg	aaaccatttg	900
tataaatgga	aacaaatacc	tgactgtgaa	aatgtcaaaa	ctacctcagtg	tgtctttcct	960
caaacgcttt	tccaaaaagg	aatttacctt	ctccgcgtac	aagcatctga	tggaaataac	1020
acatcttttt	ggtctgaaga	gataaagttt	gatactgaaa	tacaagcttt	cctacttcct	1080
ccagtcttta	acattagatc	ccttagtgat	tcattccata	tctatatcgg	tgctccaaaa	1140
cagtctggaa	acacgcctgt	gatccaggat	tatccactga	tttatgaaat	tattttttgg	1200
gaaaacactt	caaatgctga	gagaaaaat	atcgagaaaa	aaactgatgt	tacagttcct	1260
aatttgaaac	cactgactgt	atattgtgtg	aaagccagag	cacacaccat	ggatgaaaag	1320
ctgaataaaa	gcagtggttt	tagtgacgct	gtatgtgaga	aaacaaaacc	aggaaatacc	1380
tctaaaattt	ggcttatagt	tggaatttgt	attgcattat	tgctctccc	gtttgtcatt	1440
tatgtgcga	aagtcttctt	gagatgcac	aattatgtct	tctttccatc	acttaaacct	1500
tctccagta	tagatgagta	tttctctgaa	cagccattga	agaatcttct	gctttcaact	1560
tctgaggaac	aaatcgaaaa	atgtttcata	attgaaaaata	taagcacaat	tgctacagta	1620
gaagaaacta	atcaaaactga	tgaagatcat	aaaaaatata	gttcccaaac	tagccaagat	1680
tcaggaaatt	attctaataga	agatgaaagc	gaaagtaaaa	caagtgaaga	actacagcag	1740
gactttgtat	gaccagaaat	gaactgtgtc	aagtataagg	tttttcagca	ggagttacac	1800
tgggagcctg	aggctctcac	cttctctca	gtaactacag	agaggacggt	tcctgtttag	1860
ggaaagaaaa	aacatcttca	gatcataggt	cctaaaaata	cgggcaagct	cttaactatt	1920
taaaaatgaa	attacaggcc	cgggcacggg	ggctcacacc	tgtaatccca	gcactttggg	1980
aggctgaggc	aggcagatca	tgaggccaag	agatcgagac	cagcctggcc	aacgtggtga	2040
aaccccatct	ctactaaaaa	tacaaaaatt	agccgggtag	taggtaggcg	cgcgcctggt	2100
gtcttagcta	ctcaggaggc	tgaggcagga	gaatcgcttg	aaaacaggag	gtggaggttg	2160
cagtgagccg	agatcacgcc	actgcactcc	agcctggtga	cagcgtgaga	ctcttataaa	2220
aaagaaatta	aaagagttga	gacaaacggt	tcctacattc	ttttccatgt	gtaaaatcat	2280
gaaaagcct	gtcaccggac	ttgcattgga	tgagatgagt	cagacccaaa	cagtggccac	2340
ccgtcttctc	cctgtgagcc	taagtgcagc	cgtgctagct	gcgcaccgtg	gctaaggatg	2400
acgtctgtgt	tcctgtccat	cactgatgct	gctggctact	gcatgtgcca	cacctgtctg	2460
ttcgccattc	ctaacattct	gtttcattct	tcctcgggag	atatttcaaa	catttggtct	2520
tttcttttaa	cactgagggt	aggcccttag	gaaatattat	taggaaagtc	tgaacacggt	2580
atcacttggg	tttctggaaa	gtagcttacc	ctagaaaaaca	gctgcaaatg	ccagaaagat	2640
gatccctaaa	aatgttgagg	gacttctggt	cattcatccc	gagaacattg	gcttccacat	2700
cacagtatct	acccttacct	ggtttaggat	taaagccagg	caatctttta	ctatg	2755

<210> 82
 <211> 9
 <212> PRT
 <213> Homo Sapiens
 <400> 82
 Gly Ser Glu Asn Leu Tyr Phe Gln Leu
 5

<210> 83
 <211> 2897
 <212> DNA
 <213> Homo sapiens
 <400> 83
 cccgcactaa agacgcttct tcccggcggg taggaatccc gccggcgagc cgaacagttc 60
 cccgagcgca gcccgcgac caccaccgg cgcacgggc cgcttttgtc ccccgcccgc 120
 cgcttctgtc cgagaggccg cccgcgaggc gcatcctgac cgcgagcgtc ggttcccaga 180

gccgggcgcg	gctggggccc	gaggctagca	tctctcggga	gccgcaaggc	gagagctgca	240
aagtttaatt	agacacttca	gaattttgat	cacctaattgt	tgatttcaga	tgtaaaagtc	300
aagagaagac	tctaaaaata	gcaaagatgc	ttttgagcca	gaatgccttc	atcttcagat	360
cacttaattt	ggttctcatg	gtgtatatca	gcctcgtggt	tggtatttca	tatgattcgc	420
ctgattacac	agatgaatct	tgcactttca	agatatacatt	gcgaaatttc	cggtccatct	480
tatcatggga	tttaaaaaac	cactccattg	taccaactca	ctatacattg	ctgtatacaa	540
tcatgagtaa	accagaagat	ttgaagggtg	ttaagaactg	tgcaaatacc	acaagatcat	600
tttgtgacct	cacagatgag	tggagaagca	cacacgaggc	ctatgtcacc	gtcctagaag	660
gattcagcgg	gaacacaacg	ttgttcagtt	gctcacacaa	tttctggctg	gccatagaca	720
tgtcttttga	accaccagag	tttgagattg	ttggttttac	caaccacatt	aatgtgatgg	780
tgaattttcc	atctattggt	gaggaagaat	tacagtttga	tttatctctc	gtcattgaag	840
aacagtcaga	gggaattggt	aagaagcata	aacccgaaat	aaaagggaaac	atgagtggaag	900
atctcaccta	tatcattgac	aagttaattc	caaacacgaa	ctactgtgta	tctgtttatt	960
tagagcacag	tgatgagcaa	gcagtaataa	agtctccctt	aaaatgcacc	ctccttcac	1020
ctggccagga	atcagaatca	gcagaatctg	ccaaaatagg	aggaataaatt	actgtgtttt	1080
tgatagcatt	ggtcttgaca	agcaccatag	tgacactgaa	atggattggt	tatatatgct	1140
taagaaatag	cctccccaaa	gtcttgaatt	ttcataactt	tttagcctgg	ccatttccta	1200
acctgccacc	gttgggaagc	atggatatgg	tggaggtcat	ttacatcaac	agaaagaaga	1260
aagtgtggga	ttataattat	gatgatgaaa	gtgatagcga	tactgaggca	gcgcccagga	1320
caagtggcgg	tggctatacc	atgcatggac	tgactgtcag	gcctctgggt	caggcctctg	1380
ccacctctac	agaatcccag	ttgatagacc	cggagtccga	ggaggagcct	gacctgcctg	1440
aggttgatgt	ggagctcccc	acgatgccaa	aggacagccc	tcagcagttg	gaactcttga	1500
gtgggcccctg	tgagaggaga	aagagtcacc	tccaggacce	ttttcccga	gaggactaca	1560
gctccacgga	ggggtctggg	ggcagaatta	ccttcaatgt	ggacttaaac	tctgtgtttt	1620
tgagagttct	tgatgacgag	gacagtgcag	acttagaagc	ccctctgatg	ctatcgtctc	1680
atctggaaga	gatggttgac	ccagaggatc	ctgataatgt	gcaatcaaac	catttgctgg	1740
ccagcgggga	agggacacag	ccaacctttc	ccagcccctc	ttcagagggc	ctgtggtccg	1800
aagatgctcc	atctgatcaa	agtgacactt	ctgagtcaga	tgttgacctt	ggggatgggt	1860
atataatgag	atgactccaa	aactattgaa	tgaacttggg	cagacaagca	cctacagggg	1920
tctttgtctc	tgcatcctaa	cttgctgcct	tatcgtctgc	aagtgttctc	caagggaaag	1980
aggaggaaac	tgtggtgttc	ctttcttcca	ggtgacatca	cctatgcaca	ttcccagtat	2040
ggggaccata	gtatgagaca	gtgcattggt	tacatattca	aagtgggtgca	ctttgaagga	2100
agcacatgtg	cacctttcct	ttacactaat	gcacttagga	tgtttctgca	tcatgtctac	2160
cagggagcag	ggttccccac	agtttcagag	gtgggtccagg	accctatgat	atctctcttc	2220
tttctgtctt	ttttttttt	ttttgagaca	gagtctcgtt	ctgtcgccca	agctggagcg	2280
caatgggtgtg	atcttggtctc	actgcaacat	ccgctcccg	ggttcagggtg	attctcctgc	2340
ctcagcctcc	ctcgcaagta	gctgggatta	cagggcctg	ccaccatgcc	tagcaaat	2400
ttgtattttt	agtggagaca	ggattttacc	atggtggcca	ggctggtctc	gaactcctga	2460
cctcaagtga	tctgccctcc	tcagcctcgt	aaagtctgg	gattacaggg	gtgagccgct	2520
gtgcctggct	ggccctgtga	tatttctgtg	aaataaatg	ggccaggggtg	ggagcagggg	2580
aagaaaagga	aaatagtagc	aagagctgca	aagcaggcag	gaagggagga	ggagagccag	2640
gtgagcagtg	gagagaaggg	gggcccctgca	caaggaaca	gggaagagcc	atcgaagttt	2700
cagtcgggtga	gccttgggca	cctcacccat	gtcacatcct	gtctcctgca	attggaatc	2760
caccttgtcc	agccctcccc	agttaaagtg	gggaagacag	actttaggat	cacgtgtgtg	2820
actaatacag	aaaggaaca	tggcgtcggg	gagagggata	aaacctgaat	gccatatttt	2880
aagttaaaaa	aaaaaaa					2897

<210> 84

<211> 3054

<212> PRT

<213> Homo sapiens

<400> 84

Met Ala Leu Ile Phe Gly Thr Val Asn Ala Asn Ile Leu Lys Glu Val
 1 5 10 15
 Phe Gly Gly Ala Arg Met Ala Cys Val Thr Ser Ala His Met Ala Gly
 20 25 30

Ala Asn Gly Ser Ile Leu Lys Lys Ala Glu Glu Thr Ser Arg Ala Ile
 35 40 45
 Met His Lys Pro Val Ile Phe Gly Glu Asp Tyr Ile Thr Glu Ala Asp
 50 55 60
 Leu Pro Tyr Thr Pro Leu His Leu Glu Val Asp Ala Glu Met Glu Arg
 65 70 75 80
 Met Tyr Tyr Leu Gly Arg Arg Ala Leu Thr His Gly Lys Arg Arg Lys
 85 90 95
 Val Ser Val Asn Asn Lys Arg Asn Arg Arg Arg Lys Val Ala Lys Thr
 100 105 110
 Tyr Val Gly Arg Asp Ser Ile Val Glu Lys Ile Val Val Pro His Thr
 115 120 125
 Glu Arg Lys Val Asp Thr Thr Ala Ala Val Glu Asp Ile Cys Asn Glu
 130 135 140
 Ala Thr Thr Gln Leu Val His Asn Ser Met Pro Lys Arg Lys Lys Gln
 145 150 155 160
 Lys Asn Phe Leu Pro Ala Thr Ser Leu Ser Asn Val Tyr Ala Gln Thr
 165 170 175
 Trp Ser Ile Val Arg Lys Arg His Met Gln Val Glu Ile Ile Ser Lys
 180 185 190
 Lys Ser Val Arg Ala Arg Val Lys Arg Phe Glu Gly Ser Val Gln Leu
 195 200 205
 Phe Ala Ser Val Arg His Met Tyr Gly Glu Arg Lys Arg Val Asp Leu
 210 215 220
 Arg Ile Asp Asn Trp Gln Gln Glu Thr Leu Leu Asp Leu Ala Lys Arg
 225 230 235 240
 Phe Lys Asn Glu Arg Val Asp Gln Ser Lys Leu Thr Phe Gly Ser Ser
 245 250 255
 Gly Leu Val Leu Arg Gln Gly Ser Tyr Gly Pro Ala His Trp Tyr Arg
 260 265 270
 His Gly Met Phe Ile Val Arg Gly Arg Ser Asp Gly Met Leu Val Asp
 275 280 285
 Ala Arg Ala Lys Val Thr Phe Ala Val Cys His Ser Met Thr His Tyr
 290 295 300
 Ser Asp Lys Ser Ile Ser Glu Ala Phe Phe Ile Pro Tyr Ser Lys Lys
 305 310 315 320
 Phe Leu Glu Leu Arg Pro Asp Gly Ile Ser His Glu Cys Thr Arg Gly
 325 330 335
 Val Ser Val Glu Arg Cys Gly Glu Val Ala Ala Ile Leu Thr Gln Ala
 340 345 350
 Leu Ser Pro Cys Gly Lys Ile Thr Cys Lys Arg Cys Met Val Glu Thr
 355 360 365
 Pro Asp Ile Val Glu Gly Glu Ser Gly Glu Ser Val Thr Asn Gln Gly
 370 375 380
 Lys Leu Leu Ala Met Leu Lys Glu Gln Tyr Pro Asp Phe Pro Met Ala
 385 390 395 400
 Glu Lys Leu Leu Thr Arg Phe Leu Gln Gln Lys Ser Leu Val Asn Thr
 405 410 415
 Asn Leu Thr Ala Cys Val Ser Val Lys Gln Leu Ile Gly Asp Arg Lys
 420 425 430
 Gln Ala Pro Phe Thr His Val Leu Ala Val Ser Glu Ile Leu Phe Lys
 435 440 445
 Gly Asn Lys Leu Thr Gly Ala Asp Leu Glu Glu Ala Ser Thr His Met
 450 455 460
 Leu Glu Ile Ala Arg Phe Leu Asn Asn Arg Thr Glu Asn Met Arg Ile
 465 470 475 480
 Gly His Leu Gly Ser Phe Arg Asn Lys Ile Ser Ser Lys Ala His Val

				485					490					495	
Asn	Asn	Ala	Leu	Met	Cys	Asp	Asn	Gln	Leu	Asp	Gln	Asn	Gly	Asn	Phe
				500					505					510	
Ile	Trp	Gly	Leu	Arg	Gly	Ala	His	Ala	Lys	Arg	Phe	Leu	Lys	Gly	Phe
		515					520							525	
Phe	Thr	Glu	Ile	Asp	Pro	Asn	Glu	Gly	Tyr	Asp	Lys	Tyr	Val	Ile	Arg
		530				535					540				
Lys	His	Ile	Arg	Gly	Ser	Arg	Lys	Leu	Ala	Ile	Gly	Asn	Leu	Ile	Met
545					550						555				560
Ser	Thr	Asp	Phe	Gln	Thr	Leu	Arg	Gln	Gln	Ile	Gln	Gly	Glu	Thr	Ile
				565						570					575
Glu	Arg	Lys	Glu	Ile	Gly	Asn	His	Cys	Ile	Ser	Met	Arg	Asn	Gly	Asn
				580				585						590	
Tyr	Val	Tyr	Pro	Cys	Cys	Cys	Val	Thr	Leu	Glu	Asp	Gly	Lys	Ala	Gln
		595					600						605		
Tyr	Ser	Asp	Leu	Lys	His	Pro	Thr	Lys	Arg	His	Leu	Val	Ile	Gly	Asn
		610				615						620			
Ser	Gly	Asp	Ser	Lys	Tyr	Leu	Asp	Leu	Pro	Val	Leu	Asn	Glu	Glu	Lys
625					630					635					640
Met	Tyr	Ile	Ala	Asn	Glu	Gly	Tyr	Cys	Tyr	Met	Asn	Ile	Phe	Phe	Ala
				645						650					655
Leu	Leu	Val	Asn	Val	Lys	Glu	Glu	Asp	Ala	Lys	Asp	Phe	Thr	Lys	Phe
			660					665							670
Ile	Arg	Asp	Thr	Ile	Val	Pro	Lys	Leu	Gly	Ala	Trp	Pro	Thr	Met	Gln
		675					680								685
Asp	Val	Ala	Thr	Ala	Cys	Tyr	Leu	Leu	Ser	Ile	Leu	Tyr	Pro	Asp	Val
		690				695						700			
Leu	Arg	Ala	Glu	Leu	Pro	Arg	Ile	Leu	Val	Asp	His	Asp	Asn	Lys	Thr
705					710						715				720
Met	His	Val	Leu	Asp	Ser	Tyr	Gly	Ser	Arg	Thr	Thr	Gly	Tyr	His	Met
				725						730					735
Leu	Lys	Met	Asn	Thr	Thr	Ser	Gln	Leu	Ile	Glu	Phe	Val	His	Ser	Gly
			740					745						750	
Leu	Glu	Ser	Glu	Met	Lys	Thr	Tyr	Asn	Val	Gly	Gly	Met	Asn	Arg	Asp
			755				760								765
Val	Val	Thr	Gln	Gly	Ala	Ile	Glu	Met	Leu	Ile	Lys	Ser	Ile	Tyr	Lys
		770				775						780			
Pro	His	Leu	Met	Lys	Gln	Leu	Leu	Glu	Glu	Glu	Pro	Tyr	Ile	Ile	Val
785					790						795				800
Leu	Ala	Ile	Val	Ser	Pro	Ser	Ile	Leu	Ile	Ala	Met	Tyr	Asn	Ser	Gly
				805						810					815
Thr	Phe	Glu	Gln	Ala	Leu	Gln	Met	Trp	Leu	Pro	Asn	Thr	Met	Arg	Leu
			820					825							830
Ala	Asn	Leu	Ala	Ala	Ile	Leu	Ser	Ala	Leu	Ala	Gln	Lys	Leu	Thr	Leu
		835					840								845
Ala	Asp	Leu	Phe	Val	Gln	Gln	Arg	Asn	Leu	Ile	Asn	Glu	Tyr	Ala	Gln
		850				855						860			
Val	Ile	Leu	Asp	Asn	Leu	Ile	Asp	Gly	Val	Arg	Val	Asn	His	Ser	Leu
865					870						875				880
Ser	Leu	Ala	Met	Glu	Ile	Val	Thr	Ile	Lys	Leu	Ala	Thr	Gln	Glu	Met
				885						890					895
Asp	Met	Ala	Leu	Arg	Glu	Gly	Gly	Tyr	Ala	Val	Thr	Ser	Glu	Lys	Val
			900					905							910
His	Glu	Met	Leu	Glu	Lys	Asn	Tyr	Val	Lys	Ala	Leu	Lys	Asp	Ala	Trp
		915					920								925
Asp	Glu	Leu	Thr	Trp	Leu	Glu	Lys	Phe	Ser	Ala	Ile	Arg	His	Ser	Arg
		930				935									940

Lys Leu Leu Lys Phe Gly Arg Lys Pro Leu Ile Met Lys Asn Thr Val
 945 950 955 960
 Asp Cys Gly Gly His Ile Asp Leu Ser Val Lys Ser Leu Phe Lys Phe
 965 970 975
 His Leu Glu Leu Leu Lys Gly Thr Ile Ser Arg Ala Val Asn Gly Gly
 980 985 990
 Ala Arg Lys Val Arg Val Ala Lys Asn Ala Met Thr Lys Gly Val Phe
 995 1000 1005
 Leu Lys Ile Tyr Ser Met Leu Pro Asp Val Tyr Lys Phe Ile Thr
 1010 1015 1020
 Val Ser Ser Val Leu Ser Leu Leu Leu Thr Phe Leu Phe Gln Ile
 1025 1030 1035
 Asp Cys Met Ile Arg Ala His Arg Glu Ala Lys Val Ala Ala Gln
 1040 1045 1050
 Leu Gln Lys Glu Ser Glu Trp Asp Asn Ile Ile Asn Arg Thr Phe
 1055 1060 1065
 Gln Tyr Ser Lys Leu Glu Asn Pro Ile Gly Tyr Arg Ser Thr Ala
 1070 1075 1080
 Glu Glu Arg Leu Gln Ser Glu His Pro Glu Ala Phe Glu Tyr Tyr
 1085 1090 1095
 Lys Phe Cys Ile Gly Lys Glu Asp Leu Val Glu Gln Ala Lys Gln
 1100 1105 1110
 Pro Glu Ile Ala Tyr Phe Glu Lys Ile Ile Ala Phe Ile Thr Leu
 1115 1120 1125
 Val Leu Met Ala Phe Asp Ala Glu Arg Ser Asp Gly Val Phe Lys
 1130 1135 1140
 Ile Leu Asn Lys Phe Lys Gly Ile Leu Ser Ser Thr Glu Arg Glu
 1145 1150 1155
 Ile Ile Tyr Thr Gln Ser Leu Asp Asp Tyr Val Thr Thr Phe Asp
 1160 1165 1170
 Asp Asn Met Thr Ile Asn Leu Glu Leu Asn Met Asp Glu Leu His
 1175 1180 1185
 Lys Thr Ser Leu Pro Gly Val Thr Phe Lys Gln Trp Trp Asn Asn
 1190 1195 1200
 Gln Ile Ser Arg Gly Asn Val Lys Pro His Tyr Arg Thr Glu Gly
 1205 1210 1215
 His Phe Met Glu Phe Thr Arg Asp Thr Ala Ala Ser Val Ala Ser
 1220 1225 1230
 Glu Ile Ser His Ser Pro Ala Arg Asp Phe Leu Val Arg Gly Ala
 1235 1240 1245
 Val Gly Ser Gly Lys Ser Thr Gly Leu Pro Tyr His Leu Ser Lys
 1250 1255 1260
 Arg Gly Arg Val Leu Met Leu Glu Pro Thr Arg Pro Leu Thr Asp
 1265 1270 1275
 Asn Met His Lys Gln Leu Arg Ser Glu Pro Phe Asn Cys Phe Pro
 1280 1285 1290
 Thr Leu Arg Met Arg Gly Lys Ser Thr Phe Gly Ser Ser Pro Ile
 1295 1300 1305
 Thr Val Met Thr Ser Gly Phe Ala Leu His His Phe Ala Arg Asn
 1310 1315 1320
 Ile Ala Glu Val Lys Thr Tyr Asp Phe Val Ile Ile Asp Glu Cys
 1325 1330 1335
 His Val Asn Asp Ala Ser Ala Ile Ala Phe Arg Asn Leu Leu Phe
 1340 1345 1350
 Glu His Glu Phe Glu Gly Lys Val Leu Lys Val Ser Ala Thr Pro
 1355 1360 1365
 Pro Gly Arg Glu Val Glu Phe Thr Thr Gln Phe Pro Val Lys Leu

1370		1375		1380
Lys Ile Glu Glu Ala Leu Ser Phe Gln Glu Phe Val Ser Leu Gln				
1385		1390		1395
Gly Thr Gly Ala Asn Ala Asp Val Ile Ser Cys Gly Asp Asn Ile				
1400		1405		1410
Leu Val Tyr Val Ala Ser Tyr Asn Asp Val Asp Ser Leu Gly Lys				
1415		1420		1425
Leu Leu Val Gln Lys Gly Tyr Lys Val Ser Lys Ile Asp Gly Arg				
1430		1435		1440
Thr Met Lys Ser Gly Gly Thr Glu Ile Ile Thr Glu Gly Thr Ser				
1445		1450		1455
Val Lys Lys His Phe Ile Val Ala Thr Asn Ile Ile Glu Asn Gly				
1460		1465		1470
Val Thr Ile Asp Ile Asp Val Val Val Asp Phe Gly Thr Lys Val				
1475		1480		1485
Val Pro Val Leu Asp Val Asp Asn Arg Ala Val Gln Tyr Asn Lys				
1490		1495		1500
Thr Val Val Ser Tyr Gly Glu Arg Ile Gln Lys Leu Gly Arg Val				
1505		1510		1515
Gly Arg His Lys Glu Gly Val Ala Leu Arg Ile Gly Gln Thr Asn				
1520		1525		1530
Lys Thr Leu Val Glu Ile Pro Glu Met Val Ala Thr Glu Ala Ala				
1535		1540		1545
Phe Leu Cys Phe Met Tyr Asn Leu Pro Val Thr Thr Gln Ser Val				
1550		1555		1560
Ser Thr Thr Leu Leu Glu Asn Ala Thr Leu Leu Gln Ala Arg Thr				
1565		1570		1575
Met Ala Gln Phe Glu Leu Ser Tyr Phe Tyr Thr Ile Asn Phe Val				
1580		1585		1590
Arg Phe Asp Gly Ser Met His Pro Val Ile His Asp Lys Leu Lys				
1595		1600		1605
Arg Phe Lys Leu His Thr Cys Glu Thr Phe Leu Asn Lys Leu Ala				
1610		1615		1620
Ile Pro Asn Lys Gly Leu Ser Ser Trp Leu Thr Ser Gly Glu Tyr				
1625		1630		1635
Lys Arg Leu Gly Tyr Ile Ala Glu Asp Ala Gly Ile Arg Ile Pro				
1640		1645		1650
Phe Val Cys Lys Glu Ile Pro Asp Ser Leu His Glu Glu Ile Trp				
1655		1660		1665
His Ile Val Val Ala His Lys Gly Asp Ser Gly Ile Gly Arg Leu				
1670		1675		1680
Thr Ser Val Gln Ala Ala Lys Val Val Tyr Thr Leu Gln Thr Asp				
1685		1690		1695
Val His Ser Ile Ala Arg Thr Leu Ala Cys Ile Asn Arg Arg Ile				
1700		1705		1710
Ala Asp Glu Gln Met Lys Gln Ser His Phe Glu Ala Ala Thr Gly				
1715		1720		1725
Arg Ala Phe Ser Phe Thr Asn Tyr Ser Ile Gln Ser Ile Phe Asp				
1730		1735		1740
Thr Leu Lys Ala Asn Tyr Ala Thr Lys His Thr Lys Glu Asn Ile				
1745		1750		1755
Ala Val Leu Gln Gln Ala Lys Asp Gln Leu Leu Glu Phe Ser Asn				
1760		1765		1770
Leu Ala Lys Asp Gln Asp Val Thr Gly Ile Ile Gln Asp Phe Asn				
1775		1780		1785
His Leu Glu Thr Ile Tyr Leu Gln Ser Asp Ser Glu Val Ala Lys				
1790		1795		1800

His	Leu	Lys	Leu	Lys	Ser	His	Trp	Asn	Lys	Ser	Gln	Ile	Thr	Arg
1805						1810					1815			
Asp	Ile	Ile	Ile	Ala	Leu	Ser	Val	Leu	Ile	Gly	Gly	Gly	Trp	Met
1820						1825					1830			
Leu	Ala	Thr	Tyr	Phe	Lys	Asp	Lys	Phe	Asn	Glu	Pro	Val	Tyr	Phe
1835						1840					1845			
Gln	Gly	Lys	Lys	Asn	Gln	Lys	His	Lys	Leu	Lys	Met	Arg	Glu	Ala
1850						1855					1860			
Arg	Gly	Ala	Arg	Gly	Gln	Tyr	Glu	Val	Ala	Ala	Glu	Pro	Glu	Ala
1865						1870					1875			
Leu	Glu	His	Tyr	Phe	Gly	Ser	Ala	Tyr	Asn	Asn	Lys	Gly	Lys	Arg
1880						1885					1890			
Lys	Gly	Thr	Thr	Arg	Gly	Met	Gly	Ala	Lys	Ser	Arg	Lys	Phe	Ile
1895						1900					1905			
Asn	Met	Tyr	Gly	Phe	Asp	Pro	Thr	Asp	Phe	Ser	Tyr	Ile	Arg	Phe
1910						1915					1920			
Val	Asp	Pro	Leu	Thr	Gly	His	Thr	Ile	Asp	Glu	Ser	Thr	Asn	Ala
1925						1930					1935			
Pro	Ile	Asp	Leu	Val	Gln	His	Glu	Phe	Gly	Lys	Val	Arg	Thr	Arg
1940						1945					1950			
Met	Leu	Ile	Asp	Asp	Glu	Ile	Glu	Pro	Gln	Ser	Leu	Ser	Thr	His
1955						1960					1965			
Thr	Thr	Ile	His	Ala	Tyr	Leu	Val	Asn	Ser	Gly	Thr	Lys	Lys	Val
1970						1975					1980			
Leu	Lys	Val	Asp	Leu	Thr	Pro	His	Ser	Ser	Leu	Arg	Ala	Ser	Glu
1985						1990					1995			
Lys	Ser	Thr	Ala	Ile	Met	Gly	Phe	Pro	Glu	Arg	Glu	Asn	Glu	Leu
2000						2005					2010			
Arg	Gln	Thr	Gly	Met	Ala	Val	Pro	Val	Ala	Tyr	Asp	Gln	Leu	Pro
2015						2020					2025			
Pro	Lys	Asn	Glu	Asp	Leu	Thr	Phe	Glu	Gly	Glu	Ser	Leu	Phe	Lys
2030						2035					2040			
Gly	Pro	Arg	Asp	Tyr	Asn	Pro	Ile	Ser	Ser	Thr	Ile	Cys	His	Leu
2045						2050					2055			
Thr	Asn	Glu	Ser	Asp	Gly	His	Thr	Thr	Ser	Leu	Tyr	Gly	Ile	Gly
2060						2065					2070			
Phe	Gly	Pro	Phe	Ile	Ile	Thr	Asn	Lys	His	Leu	Phe	Arg	Arg	Asn
2075						2080					2085			
Asn	Gly	Thr	Leu	Leu	Val	Gln	Ser	Leu	His	Gly	Val	Phe	Lys	Val
2090						2095					2100			
Lys	Asn	Thr	Thr	Thr	Leu	Gln	Gln	His	Leu	Ile	Asp	Gly	Arg	Asp
2105						2110					2115			
Met	Ile	Ile	Ile	Arg	Met	Pro	Lys	Asp	Phe	Pro	Pro	Phe	Pro	Gln
2120						2125					2130			
Lys	Leu	Lys	Phe	Arg	Glu	Pro	Gln	Arg	Glu	Glu	Arg	Ile	Cys	Leu
2135						2140					2145			
Val	Thr	Thr	Asn	Phe	Gln	Thr	Lys	Ser	Met	Ser	Ser	Met	Val	Ser
2150						2155					2160			
Asp	Thr	Ser	Cys	Thr	Phe	Pro	Ser	Ser	Asp	Gly	Ile	Phe	Trp	Lys
2165						2170					2175			
His	Trp	Ile	Gln	Thr	Lys	Asp	Gly	Gln	Cys	Gly	Ser	Pro	Leu	Val
2180						2185					2190			
Ser	Thr	Arg	Asp	Gly	Phe	Ile	Val	Gly	Ile	His	Ser	Ala	Ser	Asn
2195						2200					2205			
Phe	Thr	Asn	Thr	Asn	Asn	Tyr	Phe	Thr	Ser	Val	Pro	Lys	Asn	Phe
2210						2215					2220			
Met	Glu	Leu	Leu	Thr	Asn	Gln	Glu	Ala	Gln	Gln	Trp	Val	Ser	Gly

2225	2230	2235
Trp Arg Leu Asn Ala Asp Ser Val Leu Trp Gly Gly His Lys Val		
2240	2245	2250
Phe Met Ser Lys Pro Glu Glu Pro Phe Gln Pro Val Lys Glu Ala		
2255	2260	2265
Thr Gln Leu Met Asn Glu Leu Val Tyr Ser Gln Gly Glu Lys Arg		
2270	2275	2280
Lys Trp Val Val Glu Ala Leu Ser Gly Asn Leu Arg Pro Val Ala		
2285	2290	2295
Glu Cys Pro Ser Gln Leu Val Thr Lys His Val Val Lys Gly Lys		
2300	2305	2310
Cys Pro Leu Phe Glu Leu Tyr Leu Gln Leu Asn Pro Glu Lys Glu		
2315	2320	2325
Ala Tyr Phe Lys Pro Met Met Gly Ala Tyr Lys Pro Ser Arg Leu		
2330	2335	2340
Asn Arg Glu Ala Phe Leu Lys Asp Ile Leu Lys Tyr Ala Ser Glu		
2345	2350	2355
Ile Glu Ile Gly Asn Val Asp Cys Asp Leu Leu Glu Leu Ala Ile		
2360	2365	2370
Ser Met Leu Val Thr Lys Leu Lys Ala Leu Gly Phe Pro Thr Val		
2375	2380	2385
Asn Tyr Ile Thr Asp Pro Glu Glu Ile Phe Ser Ala Leu Asn Met		
2390	2395	2400
Lys Ala Ala Met Gly Ala Leu Tyr Lys Gly Lys Lys Lys Glu Ala		
2405	2410	2415
Leu Ser Glu Leu Thr Leu Asp Glu Gln Glu Ala Met Leu Lys Ala		
2420	2425	2430
Ser Cys Leu Arg Leu Tyr Thr Gly Lys Leu Gly Ile Trp Asn Gly		
2435	2440	2445
Ser Leu Lys Ala Glu Leu Arg Pro Ile Glu Lys Val Glu Asn Asn		
2450	2455	2460
Lys Thr Arg Thr Phe Thr Ala Ala Pro Ile Asp Thr Leu Leu Ala		
2465	2470	2475
Gly Lys Val Cys Val Asp Asp Phe Asn Asn Gln Phe Tyr Asp Leu		
2480	2485	2490
Asn Ile Lys Ala Pro Trp Thr Val Gly Met Thr Lys Phe Tyr Gln		
2495	2500	2505
Gly Trp Asn Glu Leu Met Glu Ala Leu Pro Ser Gly Trp Val Tyr		
2510	2515	2520
Cys Asp Ala Asp Gly Ser Gln Phe Asp Ser Ser Leu Thr Pro Phe		
2525	2530	2535
Leu Ile Asn Ala Val Leu Lys Val Arg Leu Ala Phe Met Glu Glu		
2540	2545	2550
Trp Asp Ile Gly Glu Gln Met Leu Arg Asn Leu Tyr Thr Glu Ile		
2555	2560	2565
Val Tyr Thr Pro Ile Leu Thr Pro Asp Gly Thr Ile Ile Lys Lys		
2570	2575	2580
His Lys Gly Asn Asn Ser Gly Gln Pro Ser Thr Val Val Asp Asn		
2585	2590	2595
Thr Leu Met Val Ile Ile Ala Met Leu Tyr Thr Cys Glu Lys Cys		
2600	2605	2610
Gly Ile Asn Lys Glu Glu Ile Val Tyr Tyr Val Asn Gly Asp Asp		
2615	2620	2625
Leu Leu Ile Ala Ile His Pro Asp Lys Ala Glu Arg Leu Ser Arg		
2630	2635	2640
Phe Lys Glu Ser Phe Gly Glu Leu Gly Leu Lys Tyr Glu Phe Asp		
2645	2650	2655

Cys Thr Thr Arg Asp Lys Thr Gln Leu Trp Phe Met Ser His Arg
2660 2665 2670
Ala Leu Glu Arg Asp Gly Met Tyr Ile Pro Lys Leu Glu Glu Glu
2675 2680 2685
Arg Ile Val Ser Ile Leu Glu Trp Asp Arg Ser Lys Glu Pro Ser
2690 2695 2700
His Arg Leu Glu Ala Ile Cys Ala Ser Met Ile Glu Ala Trp Gly
2705 2710 2715
Tyr Asp Lys Leu Val Glu Glu Ile Arg Asn Phe Tyr Ala Trp Val
2720 2725 2730
Leu Glu Gln Ala Pro Tyr Ser Gln Leu Ala Glu Glu Gly Lys Ala
2735 2740 2745
Pro Tyr Leu Ala Glu Thr Ala Leu Lys Phe Leu Tyr Thr Ser Gln
2750 2755 2760
His Gly Thr Asn Ser Glu Ile Glu Glu Tyr Leu Lys Val Leu Tyr
2765 2770 2775
Asp Tyr Asp Ile Pro Thr Thr Glu Asn Leu Tyr Phe Gln Ser Gly
2780 2785 2790
Thr Val Asp Ala Gly Ala Asp Ala Gly Lys Lys Lys Asp Gln Lys
2795 2800 2805
Asp Asp Lys Val Ala Glu Gln Ala Ser Lys Asp Arg Asp Val Asn
2810 2815 2820
Ala Gly Thr Ser Gly Thr Phe Ser Val Pro Arg Ile Asn Ala Met
2825 2830 2835
Ala Thr Lys Leu Gln Tyr Pro Arg Met Arg Gly Glu Val Val Val
2840 2845 2850
Asn Leu Asn His Leu Leu Gly Tyr Lys Pro Gln Gln Ile Asp Leu
2855 2860 2865
Ser Asn Ala Arg Ala Thr His Glu Gln Phe Ala Ala Trp His Gln
2870 2875 2880
Ala Val Met Thr Ala Tyr Gly Val Asn Glu Glu Gln Met Lys Ile
2885 2890 2895
Leu Leu Asn Gly Phe Met Val Trp Cys Ile Glu Asn Gly Thr Ser
2900 2905 2910
Pro Asn Leu Asn Gly Thr Trp Val Met Met Asp Gly Glu Asp Gln
2915 2920 2925
Val Ser Tyr Pro Leu Lys Pro Met Val Glu Asn Ala Gln Pro Thr
2930 2935 2940
Leu Arg Gln Ile Met Thr His Phe Ser Asp Leu Ala Glu Ala Tyr
2945 2950 2955
Ile Glu Met Arg Asn Arg Glu Arg Pro Tyr Met Pro Arg Tyr Gly
2960 2965 2970
Leu Gln Arg Asn Ile Thr Asp Met Ser Leu Ser Arg Tyr Ala Phe
2975 2980 2985
Asp Phe Tyr Glu Leu Thr Ser Lys Thr Pro Val Arg Ala Arg Glu
2990 2995 3000
Ala His Met Gln Met Lys Ala Ala Ala Val Arg Asn Ser Gly Thr
3005 3010 3015
Arg Leu Phe Gly Leu Asp Gly Asn Val Gly Thr Ala Glu Glu Asp
3020 3025 3030
Thr Glu Arg His Thr Ala His Asp Val Asn Arg Asn Met His Thr
3035 3040 3045
Leu Leu Gly Val Arg Gln
3050

<211> 4157
 <212> DNA
 <213> Homo sapiens
 <400> 85

```

agcggggcgg ggcgccagcg ctgccttttc tectgccggg tagtttcgct ttctctgcgca      60
gagtctgcgg aggggctcgg ctgcaccggg gggatcgcgc ctggcagacc ccagaccgag      120
cagagggcgac ccagcgcgct cgggagaggg tgcaccgccc cgccccgcc tagcccttcc      180
ggatcctgcg cgcagaaaag tttcatttgc tgtatgccat cctcgagagc tgtctaggtt      240
aacgttcgca ctctgtgtat ataacctcga cagtcttggc acctaacgtg ctgtgcgtag      300
ctgctccttt ggttgaatcc ccaggccctt gttggggcac aagggtggcag gatgtctcag      360
tggtagcaac ttcagcagct tgactcaaaa ttcctggagc aggttcacca gctttatgat      420
gacagttttc ccaatgaaat cagacagtac ctggcacagt ggttagaaaa gcaagactgg      480
gagcacgctg caatgatgt ttcatttggc accatccggt ttcattgacct cctgtcacag      540
ctggatgatc aatatagtcg cttttctttg gagaataact tcttgetaca gcataacata      600
aggaaaagca agcgtaatct tcaggataat tttcaggaag acccaatcca gatgtctatg      660
atcattttaca gctgtctgaa ggaagaaagg aaaattctgg aaaacgcca gagatttaat      720
caggctcagt cggggaatat tcagagcaca gtgatgttag acaaacagaa agagcttgac      780
agtaaagtca gaaatgtgaa ggacaaggtt atgtgtatag agcatgaaat caagagcctg      840
gaagatttac aagatgaata tgacttcaaa tgcaaaacct tgcagaacag agaacacgag      900
accaatggtg tggcaaagag tgatcagaaa caagaacagc tgttactcaa gaagatgtat      960
ttaatgcttg acaataagag aaaggaagta gttcacaaaa taatagagtt gctgaatgtc     1020
actgaactta cccagaatgc cctgattaat gatgaactag tggagtggaa gcggagacag     1080
cagagcgcct gtattggggg gccgcccaat gcttgettgg atcagctgca gaactggttc     1140
actatagttg cggagagtct gcagcaagtt cggcagcagc ttaaaaagt tggaggaattg     1200
gaacagaaat acacctcga acatgaccct atcacaaaa acaacaagt gttatgggac     1260
cgcaccttca gtcttttcca gcagctcatt cacagctcgt ttgtggtgga aagacagccc     1320
tgcattgcaa cgcacctca gaggccgctg gtcttgaaga caggggtcca gttcactgtg     1380
aagttgagac tgttggtgaa attgcaagag ctgaattata atttgaaagt caaagtctta     1440
tttgataaag atgtgaatga gagaaataca gtaaaaggat ttaggaaagt caacattttg     1500
ggcacgcaca caaaagtgat gaacatggag gagtccacca atggcagtct ggcggctgaa     1560
tttcggcacc tgcaattgaa agaacagaaa aatgctggca ccagaacgaa tgagggtcct     1620
ctcatcgtta ctgaagagct tcactcctt agttttgaaa cccaattgtg ccagcctggg     1680
ttggtaattg acctcgagac gacctctctg cccgttgtgg tgatctccaa cgtcagccag     1740
ctcccagagc gttgggcctc catcctttgg tacaacatgc tgggtggcga acccaggaat     1800
ctgtccttct tcoctgactcc accatgtgca cgatgggctc agctttcaga agtgctgagt     1860
tggcagtttt ctctgtcac caaaagaggt ctcaatgtgg accagctgaa catgttggga     1920
gagaagcttc ttggtcetaa cgcagcccc gatggctca ttcctgggac gaggttttgt     1980
aaggaaaaata taaatgataa aaattttccc tctctgcttt ggattgaaag catcctagaa     2040
ctcattaaaa aacacctgct cctctcttgg aatgatgggt gcatcatggg cttcatcagc     2100
aaggagcgag agcgtgccct gttgaaggac cagcagccgg ggacctcct gctgcggttc     2160
agtgagagct cccgggaagg ggccatcaca ttcacatggg tggagcggtc ccagaacgga     2220
ggcgaacctg acttccatgc ggttgaacct tacacgaaga aagaacttct tgctgttact     2280
ttccctgaca tcattcgcaa ttacaaagtc atggctgctg agaatattcc tgagaatccc     2340
ctgaagtatc tgtatccaaa tattgacaaa gaccatgcct ttggaagta ttactccagg     2400
ccaaaggaag caccagagcc aatggaactt gatggcccta aaggaactgg atatatcaag     2460
actgagttga tttctgtgtc tgaagttcac cctcttagac ttcagaccac agacaacctg     2520
ctocccatgt ctectgagga gtttgacgag gtgtctcggg tagtgggctc tgtagaatc     2580
gacagtatga tgaacacagt atagagcatg aattttttcc atcttctctg gcgacagttt     2640
tccttctcat ctgtgattcc ctectgctac tctgttccct cacatcctgt gtttctaggg     2700
aatgaaaga aaggccagca aattcgtctg aacctgttga tagcaagtga attttctct     2760
aactcagaaa catcagttac tctgaaggg atcctgcatc ttaactgaag taaaattgaa     2820
aggcattctc tgaagagtgg gtttcacaag tgaaaaacat ccagatacac ccaaagtatc     2880
aggacgagaa tgagggtcct ttgggaaagg agaagtttaag caacatctag caaatgttat     2940
gcataaagtc agtgcaccaac tgttataggt tgttgataa atcagtggtt atttagggaa     3000
ctgcttgacg taggaacggt aaatttctgt gggagaattc ttacatgttt tctttgcttt     3060
aagtgtaact ggcagttttc cattggttta cctgtgaaat agttcaaagc caagtttata     3120
tacaattata tcagtctctt ttcaaaggta gccatcatgg atctggtagg gggaaaatgt     3180
    
```

gtattttatt	acatccttca	cattggctat	ttaaagacaa	agacaaattc	tgtttcttga	3240
gaagagaata	ttagctttac	tgtttgttat	ggcttaatga	cactagctaa	tatcaataga	3300
aggatgtaca	tttccaaatt	cacaagttgt	gtttgatata	caaagctgaa	tacattctgc	3360
tttcatcttg	gtcacatata	attattttta	cagttctccc	aaggagttta	ggctattcac	3420
aaccctcat	tcaaaagttg	aaattaacca	tagatgtaga	taaactcaga	aatttaattc	3480
atgltttcta	aatgggctac	tttgcctttt	ttgttattag	ggtgglatit	agtctattag	3540
ccacaaaatt	gggaaaaggag	tagaaaaagc	agtaactgac	aacttgaata	atacaccaga	3600
gataatatga	gaatcagatc	atltcaaaac	tcatttccta	tgtaactgca	ttgagaactg	3660
catatgtttc	gctgatatat	gtgtttttca	catttgcgaa	tggttccatt	ctctctcctg	3720
tactttttcc	agacactttt	ttgagtggat	gatgtttcgt	gaagtatact	gtatttttac	3780
ctttttcctt	ccttatcact	gacacaaaaa	gtagattaag	agatgggttt	gacaaggttc	3840
ttccctttta	catactgctg	tctatgtggc	tgtatcttgt	ttttccacta	ctgctaccac	3900
aactatatta	tcattgcaaat	gctgtattct	tctttggtag	agataaagat	ttcttgagtt	3960
ttgttttaaa	atataagcta	aagtatctgt	attgcattaa	atataatatg	cacacagtgc	4020
tttccgtggc	actgcataca	atctgaggcc	tcctctctca	gtttttatat	agatggcgag	4080
aacctaagtt	tcagttgatt	ttacaattga	aatgactaaa	aaacaaagaa	gacaacatta	4140
aaacaatatt	gttttcta					4157

<210> 86
 <211> 4451
 <212> DNA
 <213> Homo sapiens
 <400> 86

gctcactacta	gggacgggaa	gtcgcgacca	gagccattgg	agggcgcggg	gactgcaacc	60
ctaatacagca	gagcccaaat	ggcgcagtgg	gaaatgctgc	agaatcttga	cagccccttt	120
caggatcagc	tgaccagct	ttactcgcac	agcctcctgc	ctgtggacat	tcgacagtac	180
ttggctgtct	ggattgaaga	ccagaactgg	caggaagctg	cacttgggag	tgatgattcc	240
aaggctacca	tgctattctt	ccacttcttg	gatcagctga	actatgagtg	tgccctgtgc	300
agccaggacc	cagagtcctt	gttgctgcag	cacaatttgc	ggaaattctg	ccgggacatt	360
cagccccttt	cccaggatcc	taccagttg	gctgagatga	tctttaacct	ccttctggaa	420
gaaaaaagaa	ttttgatcca	ggctcagagg	gcccatttgg	aacaaggaga	gccagttctc	480
gaaacacctg	tggagagcca	gcaacatgag	attgaaatccc	ggatcctgga	tttaagggct	540
atgatggaga	agctggtaaa	atccatcagc	caactgaaag	accagcagga	tgtcttctgc	600
ttccgatata	agatccaggc	caaaggggag	acaccctctc	tggaccccca	tcagaccaaa	660
gagcagaaga	ttctgcagga	aactctcaat	gaactggaca	aaaggagaaa	ggaggtgctg	720
gatgcctcca	aagcactgct	aggccgatta	actaccctaa	tcgagctact	gctgccaaag	780
ttggaggagt	ggaaggccca	gcagcaaaaa	gcctgcatca	gagctcccat	tgaccacggg	840
ttggaacagc	tggagacatg	gttcacagct	ggagcaaaagc	tgttgtttca	cctgagcgag	900
ctgctgaagg	agctgaaggg	actgagttgc	ctggttagct	atcaggatga	ccctctgacc	960
aaaggggtgg	acctacgcaa	cgcccaggtc	acagagttgc	tacagcgtct	gctccacaga	1020
gcctttgtgg	tagaaaccca	gccctgcatg	ccccaaactc	cccategacc	cctcatcctc	1080
aagactggca	gcaagttcac	cgtccgaaca	aggctgctgg	tgagactcca	ggaaggcaat	1140
gagtcactga	ctgtggaagt	ctccattgac	aggaatcctc	ctcaattaca	aggcttccgg	1200
aagttaaca	ttctgacttc	aaaccagaaa	actttgacc	ccgagaaggg	gcagagtcat	1260
ggtttgattt	gggacttttg	ttacctgact	ctgggtggagc	aacgttcagg	tggttcagga	1320
aagggcagca	ataagggggc	actaggtgtg	acagaggaac	tgacatcat	cagcttcaag	1380
gtcaaatata	cctaccaggg	tctgaagcag	gagctgaaaa	cggacaccct	ccctgtggtg	1440
attatltcca	acatgaacca	gctctcaatt	gcctgggctt	cagttctctg	gttcaatttg	1500
ctcagcccaa	accttcagaa	ccagcagttc	ttctccaacc	cccccaaggc	cccctggagc	1560
ttgctggggc	ctgctctcag	ttggcagttc	tctctctatg	ttggccgagg	cctcaactca	1620
gaccagctga	gcatctgag	aaacaagctg	ttcgggcaga	actgtaggac	tgaggatcca	1680
ttattgtcct	gggctgactt	cactaagcga	gagagccctc	ctggcaagtt	accattctgg	1740
acatggctgg	acaaaattct	ggagttggta	catgaccacc	tgaaggatct	ctggaatgat	1800
ggacgcacat	tgggctttgt	gagtcggagc	caggagcgcc	ggctgctgaa	gaagaccatg	1860
tctggcacct	ttctactgag	cttcagttaa	tcgtcagaag	ggggcattac	ctgctcctgg	1920
gtggagcacc	aggatgatga	caaggtgctc	atctactctg	tgcaaccgta	cacgaaggag	1980

```

gtgctgcagt cactcccgct gactgaaatc atccgccatt accagttgct cactgaggag 2040
aatataacctg aaaacccact gcgcttctct tatccccgaa tcccccgga tgaagctttt 2100
gggtgctact accaggagaa agttaatctc caggaacgga ggaaatacct gaaacacagg 2160
ctcattgtgg tctctaatag acaggtggat gaactgcaac aaccgctgga gcttaagcca 2220
gagccagagc tggagtccat agagctggaa ctagggctgg tgccagagcc agagctcagc 2280
ctggacttag agccactgct gaaggcaggg ctggatctgg ggccagagct agagtctgtg 2340
ctggagtcca ccttggagcc tgtgatagag cccacactat gcatggtatc acaaacagtg 2400
ccagagccag accaaggacc tgtatcacag ccagtgccag agccagattt gccctgtgat 2460
ctgagacatt tgaactga gccaatggaa atcttcagaa actgtgtaaa gattgaagaa 2520
atcatgccga atggtgacct actgttggct ggccagaaca ccgtggatga ggtttacgtc 2580
tccgccccca gccacttcta cactgatgga cccttgatgc cttctgactt ctaggaacca 2640
catttcctct gttcttttca tatctcttgc ccttctact cctcatagca tgatattgtt 2700
ctccaaggat ggaatcagg catgtgtccc ttccaagctg tgttaactgt tcaaactcag 2760
gctgtgtga cctcattggg gtgagaggtg aaagcataac atgggtacag aggggacaac 2820
aatgaatcag aacagatgct gagccatagg tctaaatagg atcctggagg ctgcctgctg 2880
tgctgggagg tataggggtc ctggggcag gccaggcag tgacaggta ctggagggc 2940
tcagggcagt ggcttcttcc cagtatggaa ggatttcaac attttaatag ttgggttaggc 3000
taaactggtg catactggca ttggcccttg gtggggagca cagacacagg ataggactcc 3060
atctcttct tccattcctt catgtctagg ataacttget ttcttcttcc ctttactcct 3120
ggctcaagcc ctgaatttct tcttttctg caggggttga gagcttctg ccttagccta 3180
ccatgtgaaa ctctaccctg aagaaagga tggataggaa gtagacctct ttttcttacc 3240
agtctcctcc cctactctgc ccctaagctg gctgtacctg ttctcccc ataaaatgat 3300
cctgccaatc taatgtgagt gtgaagcttt gcacactagt ttatgctacc tagtctccac 3360
tttctcaatg cttaggagac agatcactcc tggaggctgg ggatggtagg attgctgggg 3420
atttttttt ttttaaacag ggtctcactc tgttgcccag gctagagtgc aatggtgcaa 3480
tcacagctca ctgcagctc aacctcctgg gttcaagcaa tctctctacc tcagcctcct 3540
gggtagctag caccatggca tgcgccacca tgcctatth tttttttta aagacagggt 3600
cttgctatat tgcccaggct ggtcttgaac tgggctcaag tgatcctcac gccttggcct 3660
ccaaaagtgc tgggattata ggcatgagcc actgtgcttg gcaggattt ttttttttt 3720
ttttttgaga tggagtttct ctcttgttgt ccaggctgga gtgcaatggt gtgatctcgg 3780
ctcactgcaa cctccgcctt cggggtcaa gtgactctcc tgctcagcc tcccagtag 3840
ctgggattac agatctgcac caccatgcc agctaattt gtatttttag tagagacggg 3900
gtttctccat gttggtcagg ctggtctcga actcctgacc tcaagtgatc tgtccacctc 3960
ggcctcccag agtgctggga ttacaggcgt gagccactgt tcccagcagg aatttctttt 4020
ttatagtatt ggataaagtt tgggtgtttt acagaggaga agcaatgggt cttagctctt 4080
tctctattat gttatcatcc tcccttttt gtacaatatg ttgtttacct gaaaggaagg 4140
ttctatattc tgggtgtgga acctggacaa agtccaagtc tgtggaactt aaaaccttga 4200
aggctctgta taggactctg gacaatctca caccttagct attcccaggg aaccccaggg 4260
ggcaactgac attgctccaa gatgtctccc tgatgtagct tgagatataa aggaaaggcc 4320
ctgcacaggt ggctgtttct tgtctgttat gtcagaggaa cagtctgtt cagaaagggg 4380
ctctctgag cagaaatggc taataaactt tgtgctgatc tggaaaaaaa aaaaaaaaaa 4440
aaaaaaaaa a 4451

```

```

<210> 87
<211> 9
<212> PRT
<213> Homo sapiens
<400> 87
Gly Ser Glu Asn Leu Tyr Phe Gln Leu
5

```

```

<210> 88
<211> 27
<212> DNA
<213> homo sapiens
<400> 88

```

tctagaggcc tgatcatccg gtctcac 27

<210> 89

<211> 29

<212> DNA

<213> homo sapiens

<400> 89

tctagatgga aaacagaagt cccggaaac 29

<210> 90

<211> 2290

<212> DNA

<213> Homo sapiens

<400> 90

gaattccgaa tcatgtgcag aatgctgaat cttccccag ccaggacgaa taagacagcg 60
 cggaaaagca gattctcgta attctggaat tgcattgtgc aaggagtctc ctggatcttc 120
 gcaccacagct tcgggtaggg agggagtccg ggtcccgggc taggccagcc cggcaggtgg 180
 agaggggtccc cggcagcccc gcgcgcccct ggccatgtct ttaatgccct gcccttcat 240
 gtggccttct gagggttccc agggctggcc agggttgttt cccaccgagc cgcgcgctct 300
 cacccccagc caaacccacc tggcagggct ccctccagcc gagacctttt gattcccggc 360
 tcccgcgctc ccgcctccgc gccagcccgg gaggtggccc tggacagccg gacctcgccc 420
 ggccccggct gggaccatgg tgtttctctc gggaaatgct tccgacagct ccaactgcac 480
 ccaaccgccc gcaccgggtga acatttcaa ggcattctct ctccgggtga tcttgggggg 540
 cctcattctt ttcgggtgct tgggtaacat cctagtgate ctctccgtag cctgtcaccg 600
 acacctgcac tcagtacgc actactacat cgtcaacctg gcggtggccg acctcctgct 660
 cacctccagc gtgctgccct tctccgceat cttcgaggtc ctaggctact gggccttcgg 720
 cagggctctc tgcaacatct gggcggcagt ggatgtgctg tgctgcaccg cgtccatcat 780
 gggcctctgc atcatctcca tcgaccgcta catcggcgtg agctaccgac tgcgctaccc 840
 aaccatcgct acccagagga ggggtctcat ggctctgctc tgcgctctggg cactctccct 900
 ggtcatatcc attggacccc tgttcggctg gaggcagccg gccccgagg acgagaccat 960
 ctgccagatc aacgaggagc cgggctacgt gctctctca gcgctgggct ccttctacct 1020
 gcctctggcc atcatcctgg tcatgtactg ccgctctac gtggtggcca agagggagag 1080
 ccggggcctc aagtctggcc tcaagaccga caagtcggac tcggagcaag tgacgctccg 1140
 catccatcgg aaaaacgccc cggcaggagg cagcgggatg gccagcggca agaccaagac 1200
 gcacttctca gtgaggctcc tcaagttctc ccgggagaag aaagcggcca aaacgctggg 1260
 catcgtggctc ggctgctctg tccctctgctg gctgcctttt ttcttagtca tgcccattgg 1320
 gtctttcttc cctgatttca agccctctga aacagttttt aaaatagtat tttggctcgg 1380
 atatctaaac agctgcatca accccatcat ataccatgc tccagccaag agttcaaaaa 1440
 ggcctttcag aatgtcttga gaatccagtg tctccgcaga aagcagtctt ccaaaccatgc 1500
 cctgggctac accctgcacc cgcccagcca ggccgtggaa gggcaacaca aggacatggg 1560
 gcgcatcccc gtgggatcaa gagagacctt ctacaggatc tccaagacgg atggcgtttg 1620
 tgaatggaat tttttctctt ccattgcccc tggatctgcc aggattacag tgtccaaaga 1680
 ccaatcctcc tgtaccacag cccgggtgag aagtaaaagc tttttggagg tctgctgctg 1740
 tgtagggccc tcaaccccca gccttgaaa gaaccatcaa gttccaacca ttaagggtcca 1800
 caccatctcc ctcagtgaga acggggagga agtctaggac aggaaagatg cagaggaag 1860
 gggataatc ttaggtacct accccacttc cttctcggaa ggccagctct tcttggagga 1920
 caagacagga ccaatcaaag aggggacctg ctgggaatgg ggtgggtggg agaccctact 1980
 catcaggcag cgggtagggc acaggggaaga gggaggggtg ctcaacaacca accagttcag 2040
 aatgatacgg aacagcattt ccctgcagct aatgctttct tggctactct gtgccactt 2100
 caacgaaaac caccatggga aacagaattt catgcacaat ccaaaaagact ataaatatag 2160
 gattatgatt tcatcatgaa ttttttgagc acacactcta agtttggagc tttttcttga 2220
 tggaggtgag gggattttat tttcaggctc aacctactga cagccacatt tgacatttat 2280
 gccggaattc 2290

<210> 91
 <211> 26
 <212> DNA
 <213> homo sapiens
 <400> 91
 ctcggatatc taaacagctg catcaa 26

<210> 92
 <211> 29
 <212> DNA
 <213> homo sapiens
 <400> 92
 tctagacttt ctgcagagac actggattc 29

<210> 93
 <211> 31
 <212> DNA
 <213> homo sapiens
 <400> 93
 tctagatcga aggcagtgga ggatcttcag g 31

<210> 94
 <211> 27
 <212> DNA
 <213> homo sapiens
 <400> 94
 tctagaggcc tgatcatccg gtctcac 27

<210> 95
 <211> 23
 <212> DNA
 <213> homo sapiens
 <400> 95
 cggatccggtt ggtactcttg agg 23

<210> 96
 <211> 4989
 <212> DNA
 <213> homo sapiens
 <400> 96
 tttttttttt ttttgagaaa gggaatttca tcccaaataa aaggaatgaa gtctggtcc 60
 ggaggagggt ccccgacctc gctgtggggg ctctgtttc tctcgcgcgc gctctcgctc 120
 tggccgacga gtggagaaat ctgcgggcca ggcacgcaca tccgcaacga ctatcagcag 180
 ctgaagcgc tggagaactg cacgggtatc gagggctacc tccacatcct gctcatctcc 240
 aaggccgagg actaccgcag ctaccgcttc cccaagctca cggtcattac cgagtacttg 300
 ctgctgttcc gagtggctgg cctcgagagc ctcgagacc tcttccccaa cctcacggte 360
 atccgcggtt ggaaactctt ctacaactac gccctggtca tcttcgagat gaccaatctc 420
 aaggatattg ggctttacaa cctgaggaac attactcggg gggccatcag gattgagaaa 480
 aatgctgacc tctgttacct ctccactgtg gactggtccc tgatcctgga tgcggtgtcc 540
 aataactaca ttgtggggaa taagcccca aaggaatgtg gggacctgtg tccagggacc 600
 atggaggaga agccgatgtg tgagaagacc accatcaaca atgagtacaa ctaccgctgc 660
 tggaccacaa accgctgcca gaaaatgtgc ccaagcagct gtgggaagcg ggcgtgcacc 720

gagaacaatg	agtgctgcca	ccccgagtgc	ctgggcagct	gcagcgcgcc	tgacaacgac	780
acggcctgtg	tagcttgccg	ccactactac	tatgcoggtg	tctgtgtgcc	tgccctgccg	840
cccaacacct	acaggtttga	gggctggcgc	tgtgtggacc	gtgacttctg	cgccaacatc	900
ctcagcgcog	agagcagcga	ctccgagggg	tttgtgatcc	acgacggcga	gtgcatgcag	960
gagtgccect	cgggcttcat	cgcgaacggc	agccagagca	tgtactgcat	cccttgtgaa	1020
ggtccttgcc	cgaaggtctg	tgaggaagaa	aagaaaaaa	agaccattga	ttctgttact	1080
tctgctcaga	tgctccaagg	atgcaccatc	ttcaagggca	at ttgctcat	taacatccga	1140
cgggggaata	acattgcttc	agagctggag	aacttcatgg	ggctcatcga	ggtggtgacg	1200
ggctacgtga	agatccgcc	ttctcatgcc	ttggtctcct	tgtccttctc	aaaaaacctt	1260
cgctcatcc	taggagagga	gcagctagaa	gggaattact	ccttctacgt	cctcgacaac	1320
cagaacttgc	agcaactgtg	ggactgggac	caccgcgaac	tgaccatcaa	agcagggaaa	1380
atgtactttg	ctttcaatcc	caaattatgt	gtttccgaaa	tttaccgcat	ggaggaagtg	1440
acggggacta	aagggcgcca	aagcaaaagg	gacataaaac	ccaggaacaa	cggggagaga	1500
gcctcctgtg	aaagtgacgt	cctgcatttc	acctccacca	ccacgtcgaa	gaatcgatc	1560
atcataacct	ggcaccggta	cgggccctct	gactacaggg	atctcatcag	cttcaccgtt	1620
tactacaagg	aagcaccctt	taagaatgtc	acagagtatg	atgggcagga	tgccctgcggc	1680
tccaacagct	ggaacatggt	ggacgtggac	ctcccgccca	acaaggacgt	ggagcccggc	1740
atcttactac	atgggctgaa	gccctggact	cagtacgcgg	tttacgtcaa	ggctgtgacc	1800
ctcaccatgg	tggagaacga	ccatattcgt	ggggccaaga	gtgagatcct	gtacattcgc	1860
accaatgctt	ccattccttc	cattcccttg	gacgttcttt	cagcatcgaa	ctcctcttct	1920
cagttaatcg	tgaagtggaa	ccctccctct	ctgcccaacg	gcaacctgag	ttactacatt	1980
gtgcgctggc	agcggcagcc	tcaggacggc	tacctttacc	ggcacaatta	ctgctccaaa	2040
gacaaaatcc	ccatcaggaa	gtatgccgac	ggcaccatcg	acattgagga	ggtcacagag	2100
aaccccaaga	ctgaggtgtg	tgggtggggag	aaagggcctt	gctgcgcctg	ccccaaaact	2160
gaagcgcgag	agcagggcga	gaaggaggag	gctgaatacc	gcaaagtctt	tgagaatttc	2220
ctgcacaact	ccatctcgtt	gcccagacct	gaaaggaagc	ggagagatgt	catgcaagtg	2280
gccaacacca	ccatgtccag	ccgaagcagg	aacaccacgg	cgcgagacac	ctacaacatc	2340
accgaccggg	aagagctgga	gacagagtac	cctttctttg	agagcagagt	ggataacaag	2400
gagagaactg	tcattttctaa	ccttcggcct	ttcacattgt	accgcatcga	tatccacagc	2460
tgcaaccacg	aggctgagaa	gctgggctgc	agcgcctcca	acttcgtctt	tgcaaggact	2520
atgcccgag	aaggagcaga	tgacattcct	ggggcagtg	cctgggagcc	aaggcctgaa	2580
aactccatct	ttttaaagtg	gcccgaacct	gagaatccca	atggattgat	tctaattgat	2640
gaaataaaat	acggatcaca	agttgaggat	cagcgagaat	gtgtgtccag	acaggaatac	2700
aggaagtatg	gagggggcaa	gctaaaccgg	ctaaaccggg	ggaactacac	agcccggatt	2760
cagggccacat	ctctctctgg	gaatgggtcg	tggacagatc	ctgtgttctt	ctatgtccag	2820
gccaacacag	gatatgaaaa	cttcatccat	ctgatcatcg	ctctgccctg	cgctgtcctg	2880
ttgatcgtgg	gaggggttgg	gattatgctg	tacgtcttcc	atagaaagag	aaataacagc	2940
aggctggggg	atggagtgtc	gtatgcctct	gtgaaccogg	agtacttcag	cgctgtctgat	3000
gtgtacgttc	ctgatgagtg	ggaggtggct	cgggagaaga	tcacctatgag	ccgggaactt	3060
gggcaggggt	cgtttgggat	ggtctatgaa	ggagttgcca	aggggtgtgt	gaaagatgaa	3120
cctgaaacca	gagtggccat	taaaacagtg	aacgaggccg	caagcatgcg	tgagaggatt	3180
gagtttctca	acgaagcttc	tgtgatgaag	gagttcaatt	gtcaccatgt	ggtgcgattg	3240
ctgggtgtgg	tgtcccagg	ccagccaaca	ctggtcatca	tggaaactgat	gacacggggc	3300
gatctcaaaa	gttatctcgg	gtctctgagg	ccagaaatgg	agaataatcc	agtccctagca	3360
cctccaagcc	tgagccaagat	gattcagatg	gcccggagaga	ttgcagacgg	catggcatac	3420
ctcaacgcca	ataagttcgt	ccacagagac	cttgcctgcc	ggaattgcat	ggtagccgaa	3480
gatttcacag	tcaaaatcgg	agattttggg	atgacgcgag	atatctatga	gacagactat	3540
taccggaaag	gagggcaaagg	gctgctgccc	gtgctctgga	tgtctcctga	gtccctcaag	3600
gatggagtct	tcaccactta	ctcggacgtc	tggctcctcg	gggtcgtcct	ctgggagatc	3660
gccacactgg	ccgagcagcc	ctaccagggc	ttgtccaacg	agcaagtctc	tcgcttcgct	3720
atggagggcg	gcttcttgg	caagccagac	aactgtcctg	acatgctgtt	tgaactgatg	3780
cgcatgtgct	ggcagataaa	ccccaaagatg	aggccttctc	tcctggagat	catcagcagc	3840
atcaaagagg	agatggagcc	tggcttccgg	gaggtctcct	tctactacag	cgaggagaac	3900
aagctgcccg	agcgggagga	gctggacctg	gagccagaga	acatggagag	cgtccccctg	3960
gacccctcgg	cctcctcgtc	ctccctgcca	ctgcccagaca	gacactcagg	acacaaggcc	4020
gagaacggcc	ccggccctgg	ggtgctggtc	ctccgcgcca	gcttcgacga	gagacagcct	4080
tacgcccaca	tgaaacggggg	ccgcaagaac	gagcggggcct	tgccgctgcc	ccagtcttgc	4140

acctgctgat ccttggatcc tgaatctgtg caaacagtaa cgtgtgcgca cgcgcagcgg 4200
 ggtggggggg gagagagagt ttaacaatc cattcacaag cctcctgtac ctcagtggat 4260
 cttcagttct gcccttgctg cccgcgggag acagcttctc tgcagtataa cacatttggg 4320
 atgttccttt tttcaatatg caagcagctt tttattccct gcccaaacc ttaactgaca 4380
 tgggccttta agaaccctaa tgacaacact taatagcaac agagcacttg agaaccagtc 4440
 tctcactct gtcctgtcc ttcctgttc tcccttctc tctcctctc gcttcataac 4500
 ggaaaaataa ttgccacaag tccagctggg aagccctttt tatcagtttg aggaagtggc 4560
 tgtcctgtg gccccatcca accactgtac acaccgcct gacaccgtgg gtcattacaa 4620
 aaaaacacgt ggagatggaa atttttacct ttatctttca cctttctagg gacatgaaat 4680
 ttacaaaggg ccatcgttca tccaaggctg ttaccatttt aacgctgcct aattttgcca 4740
 aatcctgaa ctttctccct catcggcccg gcgctgattc ctcgtgtccg gaggcattgg 4800
 tgagcatggc agctgggtgc tccatttgag agacacgctg gcgacacact ccgtccatcc 4860
 gactgccctt gctgtgctgc tcaaggccac aggcacacag gtctcattgc ttctgactag 4920
 attattattt gggggaaactg gacacaatag gtctttctct cagtgaaggt ggggagaagc 4980
 tgaaccggc 4989

<210> 97

<211> 3076

<212> DNA

<213> homo sapiens

<400> 97

gtttctccag ggaggcaggc cccggggaga aagttggagc ggtaacctaa gctggcagtg 60
 gcgatgatccg gcaccaaatc ggcccgcggt gcggtgcgga gactccatga ggccctggac 120
 atgaacaagc tgagtggagg cggcggggcg aggactcggg tggaggggg ccagcttggg 180
 ggcgaggagt ggaccgcca cgggagcttt gcaataagc ccacgcgggg ctggctgcat 240
 cccaacgaca aagtcattggg acccgggggt tctacttgg ttcgggtacat gggttgtgtg 300
 gaggtcctcc agtcaatgag tgccctggac tcaaacacc ggactcaggt caccagggag 360
 gccatcagtc tgggtgtgtg ggctgtgccc ggtgctaagg gggcgacaag gaggagaaaag 420
 cctctgtagc gcccgctcag ctctatcctg gggaggagta acctgaaat tgctggaatg 480
 ccaatcactc tcaccgtctc caccagcagc ctcaacctca tggccgcaga ctgcaaacag 540
 atcatcgcca accaccat gcaatctatc tcatttgcac ccggcgggga tccggacaca 600
 gccgagtagt tgcctatgt tgccaaagac cctgtgaatc agagagcctg ccacattctg 660
 gagtgtcccg aagggttgc ccaggatgtc atcagcacca ttggccaggc cttcgagttg 720
 cgcttcaaac aatacctcag gaaccacacc aaactggcca cccctcatga caggatggct 780
 ggctttgatg gctcagcatg ggatgaggag gaggaagagc cacctgacca tcagtactat 840
 aatgacttcc cggggaagga accccccttg ggggggggtg tagacatgag gcttcgggaa 900
 ggagccgctc caggggctgc tcgaccact gcacccaatg ccagacccc cagccacttg 960
 ggagctacat tgcctgtagg acagcctgtt gggggagatc cagaagtccg caaacagatg 1020
 ccacctccac caccctgtcc agcaggcaga gagctttttg atgatccctc ctatgtcaac 1080
 gtccagaacc tagacaaggc ccggcaagca gtgggtggtg ctgggcccc caatcctgct 1140
 atcaatggca gtgcacccc ggacctgtt gacatgaagc ccttcgaaga tgetcttcgc 1200
 gtgctccac ctcccagtc ggtgtccatg gctgagcagc tccgagggga gccctggttc 1260
 catgggaagc tgagccggcg ggaggctgag gcactgctgc agctcaatgg ggacttctg 1320
 gtacgggaga gcacgaccac acctggccag tatgtgctca ctggcttgc gagtgggcag 1380
 cctaagcatt tgctactggt ggacctgag ggtgtggttc ggactaagga tcaccgcttt 1440
 gaaagtgtca gtcacctat cagctaccac atggacaatc acttgccat catctctgag 1500
 ggcagogaac tgtgtctaca gcaacctgtg gagcggaaac tgtgatctgc cctagcgtc 1560
 tcttccagaa gatgccctcc aatcctttcc acctatctc ctaactctcg ggacctcgtt 1620
 tgggagtgtt ctgtgggctt ggcttgtgt cagagctggg agtagcatgg actctgggtt 1680
 tcatatccag ctgagtgaga gggtttgagt caaaagcctg ggtgagaatc ctgctctcc 1740
 ccaaacatta atcaccaag tattaatgta cagagtggcc cctcacctgg gccttctctg 1800
 tgccaacctg atgcccctc cccaagaagg tgagtgtctg tcatggaaaa tgtcctgtgg 1860
 tgacaggccc agtggaacag tcaccttct gggcaagggg gaacaaatca cacctctggg 1920
 cttcagggta tccagacc ctctcaacac ccgcccccc catgtttaaa ctttgtgcct 1980
 ttgaccatct cttaggtcta atgatatttt atgcaaacag ttcttgagc cctgaattca 2040
 atgacagga tgccaacacc ttcttgctt ctgggacctg tgttcttgc gagcaccctc 2100

tccggtttgg	gttgggataa	cagaggcagg	agtggcagct	gtcccctctc	cctgggggata	2160
tgcaaccott	agagattgcc	ccagagcccc	actcccggcc	aggcgggaga	tggaaccctc	2220
ccttgctcag	tgccctcctgg	ccggggcccc	tcaccccaag	gggtctgtat	atacatttca	2280
taaggcctgc	cctcccatgt	tgcatgccta	tgtactctac	gccaaagtgc	agcccttcc	2340
cctgaagcct	ctgcctgccc	tcctttctg	ggaggggggg	gtgggggtga	ctgaatttgg	2400
gcctcttgta	cagttaactc	tcccaggtgg	atlttggtga	ggtgagaaaa	ggggcattga	2460
gactataaag	cagtagacaa	tcccacata	ccatctgtag	agttggaact	gcattctttt	2520
aaagttttat	atgcataat	tttagggctg	tagacttact	ttcctatfff	cttttccatt	2580
gcttattott	gagcacaana	tgataatcaa	ttattacatt	tatacatcac	ctttttgact	2640
tttccaagcc	cttttacagc	tcttggcatt	ttcctcgcct	aggcctgtga	ggtaaactggg	2700
atcgcacctt	ttataccaga	gacctgaggc	agatgaaatt	tatttccatc	taggactaga	2760
aaaacttggg	tctcttaccg	cgagactgag	aggcagaagt	cagcccgaat	gcctgtcagt	2820
ttcatggagg	ggaaaacgcaa	aacctgcagt	tcttgagtac	cttctacagg	cccggcccag	2880
cctaggcccc	gggtggccac	accacagcaa	gccggcccc	cctcttttgg	ccttggtgat	2940
aagggagagt	tgaccgtttt	catcctggcc	tccttttgc	gtttggatgt	ttccacgggt	3000
ctcacttata	ccaaagggaa	aactcttcat	taaagtccgt	atftcttcta	aaaaaaaaaa	3060
aaaaaaaaaa	aaaaaa					3076

<210> 9811
 <211> 4
 <212> PRT
 <213> homo sapiens
 <400> 9811
 Asn Ser Gly Ser
 1

<210> 99
 <211> 2261
 <212> DNA
 <213> homo sapiens
 <400> 99

gaaatcaggc	tccggggccgg	ccgaagggcg	caactttccc	ccctcggcgc	cccaccggct	60
cccgcgcgcc	tcccctcgcg	cccagccttc	gagccaagca	gcgtcctggg	gagcgcgtca	120
tggccttacc	agtgaccgcc	ttgctcctgc	cgctggcctt	gctgctccac	gccgcaccgc	180
cgagccagtt	ccgggtgtcg	ccgctggatc	ggacctggaa	cctggggcgag	acagtggagc	240
tgaagtgcc	ggtgctgctg	tccaaccoga	cgtcgggctg	ctcgtggctc	ttccagccgc	300
gcggcgcgcg	cgccagtcct	accttctctc	tatacctctc	ccaaaaaag	cccaaggcgg	360
ccgaggggct	ggacaccag	cggttctcgg	gcaagaggtt	gggggacacc	ttcgtcctca	420
coctgagcga	cttccgcoga	gagaacgagg	gctactatft	ctgctcggcc	ctgagcaact	480
ccatcatgta	cttcagccac	ttcgtgccgg	tcttctgccc	agcgaagccc	accacgacgc	540
cagcgcgcgcg	accaccaaca	ccggcgccca	ccatcgcgctc	gcagccctg	tccctgcgcc	600
cagagggcgtg	ccggccagcg	gcggggggcg	cagtgcacac	gagggggctg	gacttcgect	660
gtgatatcta	catctgggcg	cccttggccg	ggacttgtgg	ggtccttctc	ctgtcactgg	720
ttatcaccct	ttactgcaac	cacaggaacc	gaagacgtgt	ttgcaaatgt	ccccggcctg	780
tgggtcaaatc	gggagacaag	cccagccttt	cggcgagata	cgtctaacc	tgtgcaacag	840
ccactacatt	acttcaaaact	gagatccttc	cttttgaggg	agcaagtcct	tccctttcat	900
tttttccagt	cttctctcct	gtgtattcat	tctcatgatt	attatfttag	tgggggcggg	960
gtgggaaaga	ttacttttctc	tttatgtgtt	tgacgggaaa	caaaactagg	taaaatctac	1020
agtacaccac	aagggtcaca	atactgttgt	gcgcacatcg	cggtagggcg	tggaaagggg	1080
caggccagag	ctaccgcag	agttctcaga	atcatgctga	gagagctgga	ggcaccatg	1140
ccatctcaac	ctcttccccg	cccgttttac	aaagggggag	gctaaagccc	agagacagct	1200
tgatcaaagg	cacacagcaa	gtcagggttg	gagcagtagc	tggagggacc	ttgtctcca	1260
gctcagggct	ctttctctca	caccattcag	gtctttcttt	ccgaggcccc	tgtctcaggg	1320
tgaggtgctt	gagtetccaa	cggaagggga	acaagtactt	cttgatacct	gggatactgt	1380
gcccagagcc	tcgaggagggt	aatgaattaa	agaagagaac	tgcctttggc	agagttctat	1440

aatgtaaaca	atatcagact	tttttttttt	ataatcaagc	ctaaaattgt	atagacctaa	1500
aataaaatga	agtgggtgagc	ttaaccctgg	aaaatgaatc	cctctatctc	taaagaaaaat	1560
ctctgtgaaa	cccctatgtg	gaggcggaat	tgctctccca	gcccttgcac	tcagagagggg	1620
cccatgaaag	aggacaggct	accocctttac	aaatagaatt	tgagcatcag	tgagggttaa	1680
ctaaggccct	cttgaatctc	tgaattttgag	atacaaacat	gttcctggga	tcactgatga	1740
ctttttatac	tttgtaaaga	caattgttgg	agagcccctc	acacagccct	ggcctctgct	1800
caactagcag	atacagggat	gaggcagacc	tgactctctt	aaggaggctg	agagcccaaa	1860
ctgctgtccc	aaacatgcac	ttccttgcct	aaggtatggt	acaagcaatg	cctgcccatt	1920
ggagagaaaa	aacttaagta	gataaggaaa	taagaaccac	tcataattct	tcaccttagg	1980
aataatctcc	tgtaaatatg	gtgtacattc	ttcctgatta	ttttctacac	atacatgtaa	2040
aatatgtctt	tcttttttaa	atagggttgt	actatgctgt	tatgagtggc	tttaatgaat	2100
aaacatttgt	agcatcctct	ttaatgggta	aacagcaaaa	aaaaaaaaaa	aaaaaaaaaa	2160
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	2220
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a		2261

<210> 100

<211> 6450

<212> DNA

<213> homo sapiens

<400> 100

gagttgtgcc	tggagtgatg	tttaagccaa	tgtcagggca	aggcaacagt	ccctggccgt	60
cctccagcac	ctttgtaatg	catatgagct	cgggagacca	gtacttaaag	ttggaggccc	120
gggagcccag	gagctggcgg	agggcgttcg	tcctgggagc	tgcaacttgc	ccgtcgggtc	180
gccggcttca	ccggaccgca	ggctcccggg	gcagggccgg	ggccagagct	cgcgtgtcgg	240
cgggacatgc	gctgcgtcgc	ctctaaccctc	gggtgtgct	ctttttccag	gtggcccgc	300
ggtttctgag	ccttctgccc	tgccgggaca	cggctgcac	cctgcccggc	gccaccggac	360
atgaccatga	ccctccacac	caaagcatct	gggatggccc	tactgcatca	gatccaaggg	420
aacgagctgg	agcccctgaa	ccgtccgcag	ctcaagatcc	ccctggagcg	gcccctgggc	480
gaggtgtacc	tggacagcag	caagcccggc	gtgtacaact	accccagagg	cgccgcctac	540
gagttcaacg	ccgcccggcg	cgccaacgcg	caggtctacg	gtcagaccgg	cctcccctac	600
ggcccggggt	ctgaggtctg	ggcgttcggc	tccaacggcc	tgggggggtt	ccccccactc	660
aacagcgtgt	ctccgagccc	gctgatgcta	ctgcaccgc	cgcgcagct	gtcgcctttc	720
ctgcagcccc	acggccagca	ggtgccctac	tacctggaga	acgagcccag	cggctacacg	780
gtgcgcgagg	ccggcccggc	ggcattctac	aggccaaatt	cagataatcg	acgccagggg	840
ggcagagaaa	gattggccag	taccaatgac	aagggaaagta	tggctatgga	atctgccaaag	900
gagactcgct	actgtgcagt	gtgcaatgac	tatgcttcag	gctaccatta	tggagtctgg	960
tcctgtgagg	gctgcaaggg	cttcttcaag	agaagtattc	aaggacataa	cgactatatg	1020
tgccagccca	ccaaccagtg	caccattgat	aaaaacagga	ggaagagctg	ccaggcctgc	1080
cggctccgca	aatgctacga	agtgggaatg	atgaaaggtg	ggatacgaaa	agaccgaaag	1140
ggagggagaa	tgttgaaaca	caagcgccag	agagatgatg	gggagggcag	gggtgaagtg	1200
gggtctgctg	gagacatgag	agctgccaac	ctttggccaa	gcccgcctcat	gatcaaacgc	1260
tctaagaaga	acagcctggc	cttgtccctg	acggccgacc	agatgggtcag	tgccctgttg	1320
gatgctgagc	ccccatact	ctattccgag	tatgatccta	ccagaccctt	cagtgaagct	1380
tcgatgatgg	gcttactgac	caacctggca	gacagggagc	tggttcacat	gatcaactgg	1440
gcgaagaggg	tgccaggctt	tgtggatttg	accctccatg	atcagggtcca	ccttctagaa	1500
tgtgcctggc	tagagatcct	gatgattggt	ctcgtctggc	gctccatgga	gcaccagtg	1560
aagctactgt	ttgctcctaa	cttgotcttg	gacaggaacc	agggaaaatg	tgtagagggc	1620
atggtggaga	tcttcgacat	gctgctggct	acatcatctc	ggttccgcat	gatgaatctg	1680
cagggagagg	agtttgtgtg	cctcaaactc	attatthttgc	ttaatctctgg	agtgtacaca	1740
ttctgtcca	gcaccctgaa	gtctctggaa	gagaaggacc	atatccaccg	agtccctggc	1800
aagatcacag	acactttgat	ccacctgatg	gccaaaggcag	gocctgacct	gcagcagcag	1860
caccagcggc	tggcccagct	cctcctcatc	ctctcccaca	tcaggcacat	gagtaacaaa	1920
ggcatggagc	atctgtacag	catgaagtgc	aagaacgtgg	tgcccctcta	tgacctgctg	1980
ctggagatgc	tggacgccc	ccgcctacat	gcgcccacta	gocgtggagg	ggcatccgtg	2040
gagggagcgg	accaaagcca	cttggccact	gcgggtctta	cttcactgca	ttccttgcaa	2100
aagtattaca	tcacggggga	ggcagagggg	ttccctgcca	cagtctgaga	gctccctggc	2160

tcccacacgg	ttcagataat	ccctgctgca	ttttaccctc	atcatgcacc	acttttagcca	2220
aattctgtct	cctgcataca	ctccggcatg	catccaacac	caatggcttt	ctagatgagt	2280
ggccattcat	ttgcttgctc	agttcttagt	ggcacatctt	ctgtcttctg	ttgggaacag	2340
ccaaagggat	tccaaggcta	aatctttgta	acagctctct	ttcccccttg	ctatgttact	2400
aagcgtgagg	attcccgtag	ctcttcacag	ctgaactcag	tctatgggtt	ggggctcaga	2460
taactctgtg	catttaagct	acttgtagag	accaggcct	ggagagtaga	cattttgctt	2520
ctgataagca	ctttttaaat	ggctctaaga	ataagccaca	gcaaagaatt	taaagtggct	2580
cctttaattg	gtgacttggg	gaaagctagg	tcaagggttt	attatagcac	cctcttgat	2640
tcctatggca	atgcatcctt	ttatgaaagt	ggtagacett	aaagctttaa	tatgactgta	2700
gcagagtatc	tgggtgattgt	caattcactt	ccccctatag	gaatacaagg	ggccacacag	2760
ggaaggcaga	tcccctagtt	ggccaagact	tattttaact	tgatacactg	cagattcaga	2820
gtgtcctgaa	gctctgcctc	tggctttccg	gtcactgggtt	ccagttaatt	catgcctccc	2880
atggacctat	ggagagcaac	aagttgatct	tagttaagtc	tcctatatg	aggataagt	2940
tcctgatttt	tggttttatt	tttgtgttac	aaaagaaagc	cctccctccc	tgaacttgca	3000
gtaagggtcag	cttcaggacc	tgttccagtg	ggcactgtac	ttggatcttc	ccggcgtgtg	3060
tgtgccttac	acaggggtga	actgttccact	gtgggtgatgc	atgatgaggg	taaattggtag	3120
ttgaaaggag	caggggcccct	gggtgttgc	ttagcccctgg	ggcatggagc	tgaacagtac	3180
ttgtgcagga	ttgttgtggc	tactagagaa	caagagggaa	agtagggcag	aaactggata	3240
cagttctgag	cacagccaga	cttgctcagg	tggccctgca	caggctgcag	ctacctagga	3300
acattccttg	cagaccccgc	attgcctttg	ggggtgcccct	gggatccctg	gggtagcca	3360
gctcttattc	atctccagc	gtggcccctgg	ttggaagaag	cagctgtcaa	gttgtagaca	3420
gctgtgttcc	tacaattggc	ccagcaccct	ggggcacggg	agaaggggtgg	ggaccgttgc	3480
tgtcactact	caggctgact	ggggcctggt	cagattacgt	atgcccttgg	tggtttagag	3540
ataatccaaa	atcagggttt	ggtttgggga	agaaaatcct	cccccttcc	cccccgccc	3600
gttccctacc	gcccctcact	ctgccagctc	atcttcttca	atcttctttg	acctatagge	3660
taaaaaagaa	aggctcattc	cagccacagg	cgagccttcc	ctgggcccctt	gcttctctag	3720
cacaattatg	ggttacttcc	tttttcttaa	caaaaaagaa	tgtttgattt	cctctgggtg	3780
accttattgt	ctgtaattga	aaccttattg	agaggtgatg	tctgtgttag	ccaatgacct	3840
aggtagctgc	tcgggcttct	cttgggtatgt	cttgttttgg	aaagtggatt	tcattcattt	3900
ctgattgtcc	agtttaagtga	tcaccaaagg	actgagaatc	tgggagggca	aaaaaaaaaa	3960
aaaaagtttt	tatgtgcact	taaatttggg	gacaatttta	tgtatctgtg	ttaaggatat	4020
gcttaagaac	ataattcttt	tgttgcgttt	tgtttaagaa	gcaccttagt	ttgtttaaga	4080
agcaccttat	atagtataat	atatattttt	tgaaattac	attgcttgtt	tatcagacaa	4140
ttgaatgtag	taattctggt	ctggatttaa	tttgactggg	ttaacatgca	aaaaccaagg	4200
aaaaatattt	agtttttttt	tttttttttg	tatacttttc	aagctacctt	gtcatgtata	4260
cagtcattta	tgccctaaagc	ctgggtgatta	ttcattttaa	tgaagatcac	atctcatatc	4320
aacttttgta	tccacagtag	acaaaatagc	actaatccag	atgcctattg	ttggatattg	4380
aatgacagac	aatccttagt	agcaaagatt	atgcctgaaa	aggaaaatta	ttcagggcag	4440
ctaattttgc	ttttaccaaa	atatacgtag	taataatttt	ggacagtagc	taatgggtca	4500
gtgggttctt	tttaattggtt	atacttagat	tttcttttaa	aaaaattaaa	ataaaacaaa	4560
aaaaatttct	aggactagac	gatgtaatac	cagctaaagc	caaacaatta	tacagtggaa	4620
ggttttacat	tattcatcca	atgtgtttct	atcatgttta	agatactact	acatttgaag	4680
tgggcagaga	acatcagatg	attgaaatgt	tcgcccaggg	gtctccagca	actttggaaa	4740
tctctttgta	tttttacttg	aagtgccact	aatggacagc	agatattttc	tggctgatgt	4800
tggatttggg	tgtaggaaca	tgatttaaaa	aaaaaactct	tgccctctgct	ttccccact	4860
ctgaggcaag	ttaaaatgta	aaagatgtga	tttatctggg	gggctcaggt	atgggtggga	4920
agtggattca	ggaatctggg	gaatggcaaa	tatatthaaga	agagtattga	aagtatttgg	4980
aggaaaatgg	ttaattctgg	gtgtgcacca	aggttcagta	gagtcacctt	ctgccctgga	5040
gaccacaaat	caactagctc	catttacagc	catttctaaa	atggcagctt	cagttctaga	5100
gaagaaagaa	caacatcagc	agtaaagtcc	atggaatagc	tagtggctctg	tgtttctttt	5160
cgccattgct	taacttgccg	taatgatctt	ataatggcat	catgcagcaa	ttatgagagg	5220
ctaggtcatc	caaagagaag	accctatcaa	tgtaggttgc	aaaatctaac	ccctaaggaa	5280
gtgcagtctt	tgatttgatt	tccttagtaa	ccttgcagat	atgttthaacc	aagccatagc	5340
ccatgccttt	tgagggctga	acaaataagg	gacttactga	taatttactt	ttgatcacat	5400
taaggtgttc	tcaccttgaa	atcttataca	ctgaaatggc	cattgattta	ggccactggc	5460
ttagagtact	ccttcccctg	catgacactg	attacaaata	ctttcctatt	catactttcc	5520
aattatgaga	tggactgtgg	gtactgggag	tgatcactaa	caccatagta	atgtctaata	5580

ttcacaggca	gatctgcttg	gggaagctag	ttatgtgaaa	ggcaaataaa	gtcatacagt	5640
agctcaaaaag	gcaaccataa	ttctctttgg	tgcaagtctt	gggagcgtga	tctagattac	5700
actgcacccat	tcccaagtta	atcccctgaa	aaacttactct	caactggagc	aaatgaactt	5760
tgggtcccaaa	tatoccatctt	ttcagtagcgt	ttaattatgc	tctgtttcca	actgcatttc	5820
ctttccaatt	gaattaaagt	gtggcctcgt	ttttagtcct	ttaaaattgt	tttctaagta	5880
attgctgcct	ctattatggc	acttcaattt	tgcactgtct	tttgagattc	aagaaaaatt	5940
tctattcatt	tttttgcac	caattgtgcc	tgaactttta	aaatatgtaa	atgctgccat	6000
gttccaaacc	catcgtcagt	gtgtgtgttt	agagctgtgc	accctagaaa	caacatactt	6060
gtcccatgag	cagggtgcctg	agacacagac	ccctttgcat	tcacagagag	gtcattgggt	6120
atagagactt	gaattaataa	gtgacattat	gccagtttct	gttctctcac	agggtataaa	6180
caatgctttt	tgtgcactac	atactcttca	gtgtagagct	cttgttttat	gggaaaaggc	6240
tcaaagtcca	aattgtgttt	gatggattaa	tatgcccttt	tgccgatgca	tactattact	6300
gatgtgactt	ggttttgtcg	cagctttgct	ttgtttaatg	aaacacactt	gtaaacctct	6360
tttgcacttt	gaaaaagaat	ccagcgggat	gctcgagcac	ctgtaaacaa	ttttctcaac	6420
ctatttgatg	ttcaaataaa	gaattaaact				6450

<210> 101
 <211> 2011
 <212> DNA
 <213> homo sapiens
 <400> 101

tttcagtttc	tccagctgct	ggctttttgg	acaccactc	ccccgccagg	aggcagttgc	60
aagcgcggag	gctgcgagaa	ataactgcct	cttgaaactt	gcagggcgaa	gagcaggcgg	120
cgagcgtctgg	gccggggagg	gaccaccoga	gctgcgacgg	gctctggggc	tgcggggcag	180
ggctggcgcc	cggagcctga	gctgcagcgc	gtgcgctcgc	tttctcaac	agggtggcggc	240
ggggcgcgcg	ccgggagacc	ccccetaatg	cgggaaaagc	acgtgtccgc	attttagaga	300
aggcaaggcc	ggtgtgttta	tctgcaagcc	attatacttg	cccacgaatc	tttgagaaca	360
ttataatgac	ctttgtgcct	cttcttgcaa	gggtgtttct	cagctgttat	ctcaagacat	420
ggatataaaa	aaactcaccat	ctagccttaa	ttctccttcc	tcctacaact	gcagtcactc	480
catcttacc	ctggagcagc	gctccatata	cataccttcc	tcctatgtag	acagccacca	540
tgaatatcca	gccatgacat	tctatagccc	tgetgtgatg	aattacagca	ttcccagcaa	600
tgtcactaac	ttggaaggctg	ggcctggctg	gcagaccaca	agcccaaatg	tgttgtggcc	660
aacacctggg	cacctttctc	ctttagtggt	ccatcgccag	ttatcacatc	tgtatgcgga	720
acctcaaaaag	agtccctggg	gtgaagcaag	atcgctagaa	cacaccttac	ctgtaaacag	780
agagacactg	aaaaggaagg	ttagtgggaa	ccgttgccgc	agccctgtta	ctgggtccagg	840
ttcaaagagg	gatgctcact	tctgcgctgt	ctgcagcgat	tacgcacatg	gatatcacta	900
tggagtctgg	tctgtgtgaag	gatgtaaggc	cttttttaaa	agaagcattc	aaggacataa	960
tgattatatt	tgtccagcta	caaatcagtg	tacaatcgat	aaaaaccggc	gcaagagctg	1020
ccaggcctgc	cgacttcgga	agtgttacga	agtgggaatg	gtgaagtgtg	gctcccggag	1080
agagagatgt	gggtaccgcc	ttgtgctggg	acagagaagt	gccgacgagc	agctgcactg	1140
tgccggcaag	gccaagagaa	gtggcggcca	cgcgccccga	gtgcgggagc	tgctgctgga	1200
cgccctgagc	cccagcagc	tagtgctcac	cctcctggag	gctgagccgc	cccatgtgct	1260
gatcagccgc	cccagtgccg	ccttcaccga	ggcctccatg	atgatgtccc	tgaccaagtt	1320
ggccgacaag	gagttgggtac	acatgatcag	ctgggccaag	aagattcccg	gctttgtgga	1380
gctcagcctg	ttcgaccaag	tgcggctctt	ggagagctgt	tggatggagg	tgttaatgat	1440
ggggctgatg	tggcgtctca	ttgaccacc	cggcaagctc	atctttgctc	cagatcttgt	1500
tctggacagg	gatgagggga	aatgcgtaga	aggaattctg	gaaatctttg	acatgctcct	1560
ggcaactact	tcaaggtttc	gagagttaaa	actccaacac	aaagaatatc	tctgtgtcaa	1620
ggccatgate	ctgctcaatt	ccagtatgta	ccctctggtc	acagcgacct	aggatgctga	1680
cagcagccgg	aagctggctc	acttgcctga	cgccgtgacc	gatgctttgg	tttgggtgat	1740
tgcacaagag	ggcatctcct	cccagcagca	atccatgccc	ctggctaacc	tcctgatgct	1800
cctgtcccac	gtcagggcatg	cgagtaacaa	gggcatggaa	catctgctca	acatgaagtg	1860
caaaaatgtg	gtcccagtg	atgacctgct	gctggagatg	ctgaatgccc	acgtgcttcg	1920
cgggtgcaag	tcctccatca	cggggctccga	gtgcagccc	gcagaggaca	gtaaaagcaa	1980
agagggctcc	cagaaccac	agtctcagtg	a			2011