



(51) International Patent Classification:

A61K 31/519 (2006.01) A61P 35/00 (2006.01)
A61K 31/437 (2006.01)

(21) International Application Number:

PCT/US2020/065415

(22) International Filing Date:

16 December 2020 (16.12.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/952,020 20 December 2019 (20.12.2019) US
63/009,788 14 April 2020 (14.04.2020) US

(71) Applicant: **RECURIUM IP HOLDINGS, LLC** [US/US];
10835 Road to the Cure, Suite 205, San Diego, California
92121 (US).

(72) Inventors: **SAMATAR, Ahmed, Abdi**; 10835 Road to the
Cure, Suite 205, San Diego, California 92121 (US). **LI,
Jiali**; 10835 Road to the Cure, Suite 205, San Diego, Cali-
fornia 92121 (US). **MA, Jianhui**; 10835 Road to the Cure,
Suite 205, San Diego, California 92121 (US). **HUANG,
Peter, Qinhua**; 10835 Road to the Cure, Suite 205, San


Diego, California 92121 (US). **HEGDE, Sayee, Gajanan**;
10835 Road to the Cure, Suite 205, San Diego, Califor-
nia 92121 (US). **BOREN, Brant, Clayton**; 10835 Road
to the Cure, Suite 205, San Diego, California 92121 (US).
BUNKER, Kevin, Duane; 10835 Road to the Cure, Suite
205, San Diego, California 92121 (US). **DONATE, Fer-
nando**; 10835 Road to the Cure, Suite 205, San Diego, Cal-
ifornia 92121 (US).

(74) Agent: **MILLER, Kimberly, J.**; Knobbe, Martens, Olson
& Bear, LLP, 2040 Main Street, 14th Floor, Irvine, Cali-
fornia 92614 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,
NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW,
SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(54) Title: COMBINATIONS

Figure 3



(57) Abstract: Disclosed herein are combinations of compounds for treating a disease or condition, such as cancer. A combination of compounds for treating a disease or condition can include a SERD inhibitor and a WEE1 inhibitor, along with pharmaceutically acceptable salts of any of the foregoing.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

COMBINATIONS

INCORPORATION BY REFERENCE TO ANY PRIORITY APPLICATIONS

[0001] Any and all applications for which a foreign or domestic priority claim is identified, for example, in the Application Data Sheet or Request as filed with the present application, are hereby incorporated by reference under 37 CFR 1.57, and Rules 4.18 and 20.6, including U.S. Provisional Application Nos. 62/952,020, filed December 20, 2019 and 63/009,788, filed April 14, 2020.

Field

[0002] The present application relates to the fields of chemistry, biochemistry and medicine. More particularly, disclosed herein are combination therapies, and methods of treating diseases and/or conditions with a combination therapies described herein.

Description

[0003] Cancers are a family of diseases that involve abnormal cell growth with the potential to invade or spread to other parts of the body. Cancer treatments today include surgery, hormone therapy, radiation, chemotherapy, immunotherapy, targeted therapy and combinations thereof. Survival rates vary by cancer type and by the stage at which the cancer is diagnosed. In 2019, roughly 1.8 million people will be diagnosed with cancer, and an estimated 606,880 people will die of cancer in the United States. Thus, there still exists a need for effective cancer treatments.

SUMMARY

[0004] Some embodiments described herein relate to a combination of compounds that can include an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof.

[0005] Other embodiments described herein relate to a combination of compounds that can include an effective amount of Compound (C), or a pharmaceutically acceptable salt

thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof.

[0006] Some embodiments described herein relate to the use of a combination of compounds for treating a disease or condition, wherein the combination includes an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof. Other embodiments described herein relate to the use of a combination of compounds in the manufacture of a medicament for treating a disease or condition, wherein the combination includes an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof.

[0007] Some embodiments described herein relate to the use of a combination of compounds for treating a disease or condition, wherein the combination includes an effective amount of Compound (C), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof. Other embodiments described herein relate to the use of a combination of compounds in the manufacture of a medicament for treating a disease or condition, wherein the combination includes an effective amount of Compound (C), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof.

[0008] In some embodiments, the disease or condition can be a cancer described herein.

DRAWINGS

[0009] Figure 1 provides examples of Compound (B).

[0010] Figure 2 shows the results of a combination study of Compound (A) with Compound 1 in a ZR-75-1-R xenograft tumor model.

[0011] Figure 3 shows the results of a combination study of Compound (A) with Compound 1 in a MCF-7 xenograft tumor model.


Definitions for Compound (A), and pharmaceutically acceptable salts thereof

[0012] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art. All patents, applications, published applications and other publications referenced herein are incorporated by reference in their entirety unless stated otherwise. In the event that there are a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

[0013] Whenever a group is described as being “optionally substituted” that group may be unsubstituted or substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) may be selected from one or more the indicated substituents. If no substituents are indicated, it is meant that the indicated “optionally substituted” or “substituted” group may be substituted with one or more group(s) individually and independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), cycloalkyl(alkyl), heteroaryl(alkyl), heterocyclyl(alkyl), hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, nitro, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, an amino, a mono-substituted amino group and a di-substituted amino group.

[0014] As used herein, “C_a to C_b” in which “a” and “b” are integers refer to the number of carbon atoms in a group. The indicated group can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C₁ to C₄ alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, CH₃-, CH₃CH₂-, CH₃CH₂CH₂-, (CH₃)₂CH-, CH₃CH₂CH₂CH₂-, CH₃CH₂CH(CH₃)- and (CH₃)₃C-. If no “a” and “b” are designated, the broadest range described in these definitions is to be assumed.

[0015] If two “R” groups are described as being “taken together” the R groups and the atoms they are attached to can form a cycloalkyl, cycloalkenyl, aryl, heteroaryl or heterocycle. For example, without limitation, if R^a and R^b of an NR^aR^b group are indicated to be “taken together,” it means that they are covalently bonded to one another to form a ring:



[0016] As used herein, the term “alkyl” refers to a fully saturated aliphatic hydrocarbon group. The alkyl moiety may be branched or straight chain. Examples of branched

alkyl groups include, but are not limited to, iso-propyl, sec-butyl, t-butyl and the like. Examples of straight chain alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl and the like. The alkyl group may have 1 to 30 carbon atoms (whenever it appears herein, a numerical range such as “1 to 30” refers to each integer in the given range; *e.g.*, “1 to 30 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 30 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 12 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 6 carbon atoms. An alkyl group may be substituted or unsubstituted.

[0017] The term “alkenyl” used herein refers to a monovalent straight or branched chain radical of from two to twenty carbon atoms containing a carbon double bond(s) including, but not limited to, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like. An alkenyl group may be unsubstituted or substituted.

[0018] The term “alkynyl” used herein refers to a monovalent straight or branched chain radical of from two to twenty carbon atoms containing a carbon triple bond(s) including, but not limited to, 1-propynyl, 1-butylnyl, 2-butylnyl and the like. An alkynyl group may be unsubstituted or substituted.

[0019] As used herein, “cycloalkyl” refers to a completely saturated (no double or triple bonds) mono- or multi- cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. As used herein, the term “fused” refers to two rings which have two atoms and one bond in common. As used herein, the term “bridged cycloalkyl” refers to compounds wherein the cycloalkyl contains a linkage of one or more atoms connecting non-adjacent atoms. As used herein, the term “spiro” refers to two rings which have one atom in common and the two rings are not linked by a bridge. Cycloalkyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). A cycloalkyl group may be unsubstituted or substituted. Typical mono-cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Examples of fused cycloalkyl groups are decahydronaphthalenyl, dodecahydro-1H-phenalenyl and tetradecahydroanthracenyl; examples of bridged cycloalkyl groups are bicyclo[1.1.1]pentyl,

adamantanyl, and norbornanyl; and examples of spiro cycloalkyl groups include spiro[3.3]heptane and spiro[4.5]decane.

[0020] As used herein, “cycloalkenyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all the rings (otherwise the group would be “aryl,” as defined herein). Cycloalkenyl groups can contain 3 to 10 atoms in the ring(s) or 3 to 8 atoms in the ring(s). When composed of two or more rings, the rings may be connected together in a fused, bridged or spiro fashion. A cycloalkenyl group may be unsubstituted or substituted.

[0021] As used herein, “cycloalkynyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more triple bonds in at least one ring. If there is more than one triple bond, the triple bonds cannot form a fully delocalized pi-electron system throughout all the rings. Cycloalkynyl groups can contain 6 to 10 atoms in the ring(s) or 6 to 8 atoms in the ring(s). When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. A cycloalkynyl group may be unsubstituted or substituted.

[0022] As used herein, “aryl” refers to a carbocyclic (all carbon) monocyclic or polycyclic aromatic ring system (including fused ring systems where two carbocyclic rings share a chemical bond) that has a fully delocalized pi-electron system throughout all the rings. The number of carbon atoms in an aryl group can vary. For example, the aryl group can be a C₆-C₁₄ aryl group, a C₆-C₁₀ aryl group, or a C₆ aryl group. Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted.

[0023] As used herein, “heteroaryl” refers to a monocyclic or polycyclic aromatic ring system (a ring system with fully delocalized pi-electron system) that contain(s) one or more heteroatoms (for example, 1, 2 or 3 heteroatoms), that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur. The number of atoms in the ring(s) of a heteroaryl group can vary. For example, the heteroaryl group can contain 4 to 14 atoms in the ring(s), 5 to 10 atoms in the ring(s) or 5 to 6 atoms in the ring(s). Furthermore, the term “heteroaryl” includes fused ring systems where two rings, such as at least one aryl ring and at least one heteroaryl ring, or at least two heteroaryl rings, share at least one chemical bond. Examples of heteroaryl rings include, but are not limited to, furan, furazan, thiophene,

benzothiophene, phthalazine, pyrrole, oxazole, benzoxazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, thiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, benzothiazole, imidazole, benzimidazole, indole, indazole, pyrazole, benzopyrazole, isoxazole, benzoisoxazole, isothiazole, triazole, benzotriazole, thiadiazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, purine, pteridine, quinoline, isoquinoline, quinazoline, quinoxaline, cinnoline and triazine. A heteroaryl group may be substituted or unsubstituted.

[0024] As used herein, “heterocyclyl” or “heteroalicyclyl” refers to three-, four-, five-, six-, seven-, eight-, nine-, ten-, up to 18-membered monocyclic, bicyclic and tricyclic ring system wherein carbon atoms together with from 1 to 5 heteroatoms constitute said ring system. A heterocycle may optionally contain one or more unsaturated bonds situated in such a way, however, that a fully delocalized pi-electron system does not occur throughout all the rings. The heteroatom(s) is an element other than carbon including, but not limited to, oxygen, sulfur and nitrogen. A heterocycle may further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo-systems and thio-systems such as lactams, lactones, cyclic imides, cyclic thioimides and cyclic carbamates. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. As used herein, the term “fused” refers to two rings which have two atoms and one bond in common. As used herein, the term “bridged heterocyclyl” or “bridged heteroalicyclyl” refers to compounds wherein the heterocyclyl or heteroalicyclyl contains a linkage of one or more atoms connecting non-adjacent atoms. As used herein, the term “spiro” refers to two rings which have one atom in common and the two rings are not linked by a bridge. Heterocyclyl and heteroalicyclyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). Additionally, any nitrogens in a heteroalicyclic may be quaternized. Heterocyclyl or heteroalicyclic groups may be unsubstituted or substituted. Examples of such “heterocyclyl” or “heteroalicyclyl” groups include but are not limited to, 1,3-dioxin, 1,3-dioxane, 1,4-dioxane, 1,2-dioxolane, 1,3-dioxolane, 1,4-dioxolane, 1,3-oxathiane, 1,4-oxathiin, 1,3-oxathiolane, 1,3-dithiole, 1,3-dithiolane, 1,4-oxathiane, tetrahydro-1,4-thiazine, 2H-1,2-oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouracil, trioxane, hexahydro-1,3,5-triazine, imidazoline, imidazolidine, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, morpholine, oxirane, piperidine N-Oxide, piperidine, piperazine, pyrrolidine, azepane,


pyrrolidone, pyrrolidione, 4-piperidone, pyrazoline, pyrazolidine, 2-oxopyrrolidine, tetrahydropyran, 4H-pyran, tetrahydrothiopyran, thiamorpholine, thiamorpholine sulfoxide, thiamorpholine sulfone and their benzo-fused analogs (e.g., benzimidazolidinone, tetrahydroquinoline and/or 3,4-methylenedioxyphenyl). Examples of spiro heterocyclyl groups include 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 2-oxa-6-azaspiro[3.3]heptane, 2,6-diazaspiro[3.3]heptane, 2-oxaspiro[3.4]octane and 2-azaspiro[3.4]octane.

[0025] As used herein, “aralkyl” and “aryl(alkyl)” refer to an aryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, 2-phenylalkyl, 3-phenylalkyl and naphthylalkyl.

[0026] As used herein, “heteroaralkyl” and “heteroaryl(alkyl)” refer to a heteroaryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and heteroaryl group of heteroaralkyl may be substituted or unsubstituted. Examples include but are not limited to 2-thienylalkyl, 3-thienylalkyl, furylalkyl, thienylalkyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl and imidazolylalkyl and their benzo-fused analogs.

[0027] A “heteroalicyclyl(alkyl)” and “heterocyclyl(alkyl)” refer to a heterocyclic or a heteroalicyclic group connected, as a substituent, via a lower alkylene group. The lower alkylene and heterocyclyl of a (heteroalicyclyl)alkyl may be substituted or unsubstituted. Examples include but are not limited tetrahydro-2H-pyran-4-yl(methyl), piperidin-4-yl(ethyl), piperidin-4-yl(propyl), tetrahydro-2H-thiopyran-4-yl(methyl) and 1,3-thiazinan-4-yl(methyl).

[0028] As used herein, “lower alkylene groups” are straight-chained -CH₂- tethering groups, forming bonds to connect molecular fragments via their terminal carbon atoms. Examples include but are not limited to methylene (-CH₂-), ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-) and butylene (-CH₂CH₂CH₂CH₂-). A lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene group and/or by substituting both

hydrogens on the same carbon with a cycloalkyl group (e.g., ).

[0029] As used herein, the term “hydroxy” refers to a -OH group.

[0030] As used herein, “alkoxy” refers to the Formula -OR wherein R is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl) is defined herein. A non-limiting list of

alkoxys are methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, phenoxy and benzoxy. An alkoxy may be substituted or unsubstituted.

[0031] As used herein, “acyl” refers to a hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl) and heterocyclyl(alkyl) connected, as substituents, via a carbonyl group. Examples include formyl, acetyl, propanoyl, benzoyl and acryl. An acyl may be substituted or unsubstituted.

[0032] A “cyano” group refers to a “-CN” group.

[0033] The term “halogen atom” or “halogen” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, such as, fluorine, chlorine, bromine and iodine.

[0034] A “thiocarbonyl” group refers to a “-C(=S)R” group in which R can be the same as defined with respect to O-carboxy. A thiocarbonyl may be substituted or unsubstituted.

[0035] An “O-carbamyl” group refers to a “-OC(=O)N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-carbamyl may be substituted or unsubstituted.

[0036] An “N-carbamyl” group refers to an “ROC(=O)N(R_A-)” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-carbamyl may be substituted or unsubstituted.

[0037] An “O-thiocarbamyl” group refers to a “-OC(=S)-N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-thiocarbamyl may be substituted or unsubstituted.

[0038] An “N-thiocarbamyl” group refers to an “ROC(=S)N(R_A-)” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-thiocarbamyl may be substituted or unsubstituted.

[0039] A “C-amido” group refers to a “-C(=O)N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl,

aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A C-amido may be substituted or unsubstituted.

[0040] An “N-amido” group refers to a “RC(=O)N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-amido may be substituted or unsubstituted.

[0041] An “S-sulfonamido” group refers to a “-SO₂N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An S-sulfonamido may be substituted or unsubstituted.

[0042] An “N-sulfonamido” group refers to a “RSO₂N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-sulfonamido may be substituted or unsubstituted.

[0043] An “O-carboxy” group refers to a “RC(=O)O-” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. An O-carboxy may be substituted or unsubstituted.

[0044] The terms “ester” and “C-carboxy” refer to a “-C(=O)OR” group in which R can be the same as defined with respect to O-carboxy. An ester and C-carboxy may be substituted or unsubstituted.

[0045] A “nitro” group refers to an “-NO₂” group.

[0046] A “sulfenyl” group refers to an “-SR” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A sulfenyl may be substituted or unsubstituted.

[0047] A “sulfinyl” group refers to an “-S(=O)-R” group in which R can be the same as defined with respect to sulfenyl. A sulfinyl may be substituted or unsubstituted.

[0048] A “sulfonyl” group refers to an “SO₂R” group in which R can be the same as defined with respect to sulfenyl. A sulfonyl may be substituted or unsubstituted.

[0049] As used herein, “haloalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkyl, di-haloalkyl and tri-haloalkyl). Such groups include but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 1-chloro-2-fluoromethyl and 2-fluoroisobutyl. A haloalkyl may be substituted or unsubstituted.

[0050] As used herein, “haloalkoxy” refers to an alkoxy group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkoxy, di-haloalkoxy and tri-haloalkoxy). Such groups include but are not limited to, chloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 1-chloro-2-fluoromethoxy and 2-fluoroisobutoxy. A haloalkoxy may be substituted or unsubstituted.

[0051] The term “amino” as used herein refers to a $-NH_2$ group.

[0052] A “mono-substituted amino” group refers to a “-NHR” group in which R can be an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. A mono-substituted amino may be substituted or unsubstituted. Examples of mono-substituted amino groups include, but are not limited to, $-NH(\text{methyl})$, $-NH(\text{phenyl})$ and the like.

[0053] A “di-substituted amino” group refers to a “-NR_AR_B” group in which R_A and R_B can be independently an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. A di-substituted amino may be substituted or unsubstituted. Examples of di-substituted amino groups include, but are not limited to, $-N(\text{methyl})_2$, $-N(\text{phenyl})(\text{methyl})$, $-N(\text{ethyl})(\text{methyl})$ and the like.

[0054] Where the numbers of substituents is not specified (e.g. haloalkyl), there may be one or more substituents present. For example “haloalkyl” may include one or more of the same or different halogens. As another example, “C₁-C₃ alkoxyphenyl” may include one or more of the same or different alkoxy groups containing one, two or three atoms.

[0055] As used herein, a radical indicates species with a single, unpaired electron such that the species containing the radical can be covalently bonded to another species. Hence, in this context, a radical is not necessarily a free radical. Rather, a radical indicates a specific portion of a larger molecule. The term “radical” can be used interchangeably with the term “group.”

[0056] As used herein, when a chemical group or unit includes an asterisk (*), that asterisk indicates a point of attachment of the group or unit to another structure.

[0057] As used herein, “linking groups” are chemical groups that are indicated as having multiple open valencies for connecting to two or more other groups. For example, lower alkylene groups of the general formula $-(CH_2)_n-$ where n is in the range of 1 to 10, are examples of linking groups that are described elsewhere herein as connecting molecular fragments via their terminal carbon atoms. Other examples of linking groups include $-(CH_2)_nO-$, $-(CH_2)_nNH-$, $-(CH_2)_nN(C_1-C_6alkyl)-$, and $-(CH_2)_nS-$, wherein each n is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. Those skilled in the art will recognize that n can be zero for some linking groups such as $-(CH_2)_nO-$, in which case the linking group is simply $-O-$. Those skilled in the art will also recognize that reference herein to an asymmetrical linking group will be understood as a reference to all orientations of that group (unless stated otherwise). For example, reference herein to $-(CH_2)_nO-$ will be understood as a reference to both $-(CH_2)_nO-$ and $-O-(CH_2)_n-$.

[0058] The term “pharmaceutically acceptable salt” refers to a salt of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, the salt is an acid addition salt of the compound. Pharmaceutical salts can be obtained by reacting a compound with inorganic acids such as hydrohalic acid (e.g., hydrochloric acid or hydrobromic acid), a sulfuric acid, a nitric acid and a phosphoric acid (such as 2,3-dihydroxypropyl dihydrogen phosphate). Pharmaceutical salts can also be obtained by reacting a compound with an organic acid such as aliphatic or aromatic carboxylic or sulfonic acids, for example formic, acetic, succinic, lactic, malic, tartaric, citric, ascorbic, nicotinic, methanesulfonic, ethanesulfonic, p-toluenesulfonic, trifluoroacetic, benzoic, salicylic, 2-oxopentanedioic, or naphthalenesulfonic acid. Pharmaceutical salts can also be obtained by reacting a compound with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a sodium, a potassium or a lithium salt, an alkaline earth metal salt, such as a calcium or a magnesium salt, a salt of a carbonate, a salt of a bicarbonate, a salt of organic bases such as dicyclohexylamine, N-methyl-D-glucamine, tris(hydroxymethyl)methylamine, C_1-C_7 alkylamine, cyclohexylamine, triethanolamine, ethylenediamine, and salts with amino acids such as arginine and lysine. For compounds of Formulae (A) and/or (B), those skilled in the art understand that when a salt is formed by protonation of a nitrogen-based group (for example, NH_2), the nitrogen-based group can be

associated with a positive charge (for example, NH_2 can become NH_3^+) and the positive charge can be balanced by a negatively charged counterion (such as Cl^-).

[0059] It is understood that, in any compound described herein having one or more chiral centers, if an absolute stereochemistry is not expressly indicated, then each center may independently be of R-configuration or S-configuration or a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, enantiomerically enriched, racemic mixture, diastereomerically pure, diastereomerically enriched, or a stereoisomeric mixture. In addition, it is understood that, in any compound described herein having one or more double bond(s) generating geometrical isomers that can be defined as E or Z, each double bond may independently be E or Z a mixture thereof. Likewise, it is understood that, in any compound described, all tautomeric forms are also intended to be included.

[0060] It is to be understood that where compounds disclosed herein have unfilled valencies, then the valencies are to be filled with hydrogens or isotopes thereof, e.g., hydrogen-1 (protium) and hydrogen-2 (deuterium).

[0061] It is understood that the compounds described herein can be labeled isotopically. Substitution with isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements. Each chemical element as represented in a compound structure may include any isotope of said element. For example, in a compound structure a hydrogen atom may be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom may be present, the hydrogen atom can be any isotope of hydrogen, including but not limited to hydrogen-1 (protium) and hydrogen-2 (deuterium). Thus, reference herein to a compound encompasses all potential isotopic forms unless the context clearly dictates otherwise.

[0062] It is understood that the methods and combinations described herein include crystalline forms (also known as polymorphs, which include the different crystal packing arrangements of the same elemental composition of a compound), amorphous phases, salts, solvates, and hydrates. In some embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, ethanol, or the like. In other embodiments, the compounds described herein exist in unsolvated form. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process

of crystallization with pharmaceutically acceptable solvents such as water, ethanol, or the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

[0063] Where a range of values is provided, it is understood that the upper and lower limit, and each intervening value between the upper and lower limit of the range is encompassed within the embodiments.


[0064] Terms and phrases used in this application, and variations thereof, especially in the appended claims, unless otherwise expressly stated, should be construed as open ended as opposed to limiting. As examples of the foregoing, the term 'including' should be read to mean 'including, without limitation,' 'including but not limited to,' or the like; the term 'comprising' as used herein is synonymous with 'including,' 'containing,' or 'characterized by,' and is inclusive or open-ended and does not exclude additional, unrecited elements or method steps; the term 'having' should be interpreted as 'having at least;' the term 'includes' should be interpreted as 'includes but is not limited to;' the term 'example' is used to provide exemplary instances of the item in discussion, not an exhaustive or limiting list thereof; and use of terms like 'preferably,' 'preferred,' 'desired,' or 'desirable,' and words of similar meaning should not be understood as implying that certain features are critical, essential, or even important to the structure or function, but instead as merely intended to highlight alternative or additional features that may or may not be utilized in a particular embodiment. In addition, the term "comprising" is to be interpreted synonymously with the phrases "having at least" or "including at least". When used in the context of a process, the term "comprising" means that the process includes at least the recited steps, but may include additional steps. When used in the context of a compound, composition or device, the term "comprising" means that the compound, composition or device includes at least the recited features or components, but may also include additional features or components.

[0065] With respect to the use of substantially any plural and/or singular terms herein, those having skill in the art can translate from the plural to the singular and/or from the singular to the plural as is appropriate to the context and/or application. The various singular/plural permutations may be expressly set forth herein for sake of clarity. The indefinite

article “a” or “an” does not exclude a plurality. The mere fact that certain measures are recited in mutually different dependent claims does not indicate that a combination of these measures cannot be used to advantage. Any reference signs in the claims should not be construed as limiting the scope.

Compound (A)

[0066] Some embodiments disclosed herein relate to the use of a combination of compounds for treating a disease or condition, wherein the combination can include an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein: the Compound (A) has the structure:




[0067] Compound (A) can be a salt. For example, in some embodiments, Compound (A) can be a hydrogen sulfate salt. Those skilled in the art understand that the hydrosulfate salt of Compound (A) has a single molecule of Compound (A) for a single molecule of hydrogen sulfate. In other embodiments, Compound (A) can be a sulfate salt. Those skilled in the art understand that the sulfate salt of Compound (A) has two molecules of Compound (A) for a single molecule of sulfate. Further, those skilled in the art understand that hydrogen sulfate and sulfate salts of Compound (A) are where the nitrogen of Compound (A) can be protonated.

[0068] In some embodiments, Compound (A) can be a pharmaceutically acceptable salt form of Compound (A) that can include the hydrosulfate salt of Compound A and the sulfate salt of Compound (A). As an example, a pharmaceutically acceptable salt form of Compound (A) can be a pharmaceutically acceptable salt form of Compound (A) that consists essentially of the hydrosulfate salt of Compound (A) and the sulfate salt of Compound (A). Exemplary salt forms of Compound (A) include Form A and Form C. In some embodiments, Compound (A), or a pharmaceutically acceptable salt thereof, can be Form A. In some embodiments, Compound


(A), or a pharmaceutically acceptable salt thereof, can be Form C. In some embodiments, Compound (A), or a pharmaceutically acceptable salt thereof, can include Form A and Form C. Additional details regarding Form A and Form C of Compound (A) are provided in International Application No. PCT/US2020/058526, filed November 2, 2020, which is hereby incorporated by reference in its entirety.


[0069] Other embodiments disclosed herein relate to the use of a combination of compounds for treating a disease or condition, wherein the combination can include an effective amount of Compound (C), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein: the Compound (C) has the structure:



wherein: X¹, Y¹ and Z¹ can be each independently C or N; with the first proviso that at least one of X¹, Y¹ and Z¹ is N; with the second proviso that each of X¹, Y¹ and Z¹ is uncharged; with third proviso that two of the dotted lines indicate double bonds; with the fourth proviso that the valencies of X¹, Y¹ and Z¹ can be each independently satisfied by attachment to a substituent selected from H and R¹²; X² can be O; A¹ can be selected from an optionally substituted cycloalkyl, an optionally substituted aryl, an optionally substituted heteroaryl and an optionally substituted heterocyclyl; R¹ can be selected from an optionally substituted C₁₋₆ alkyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted cycloalkyl(C₁₋₆ alkyl), an optionally substituted cycloalkenyl(C₁₋₆ alkyl), an optionally substituted aryl(C₁₋₆ alkyl), an optionally substituted heteroaryl(C₁₋₆ alkyl) and an

optionally substituted heterocyclyl(C₁₋₆ alkyl); R² and R³ can be each independently selected from hydrogen, halogen, an optionally substituted C₁₋₆ alkyl and an optionally substituted C₁₋₆ haloalkyl; or R² and R³ together with the carbon to which R² and R³ are attached can form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl; R⁴ and R⁵ can be each independently selected from hydrogen, halogen, an optionally substituted C₁₋₆ alkyl and an optionally substituted C₁₋₆ haloalkyl; or R⁴ and R⁵ together with the carbon to which R⁴ and R⁵ are attached can form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl; R⁶, R⁷, R⁸ and R⁹ can be each independently selected from hydrogen, halogen, hydroxy, an optionally substituted alkyl, an optionally substituted alkoxy, an optionally substituted haloalkyl, an optionally substituted mono-substituted amine, and an optionally substituted di-substituted amine; R¹⁰ can be hydrogen, halogen, an optionally substituted alkyl, or an optionally substituted cycloalkyl; R¹¹ can be hydrogen; R¹² can be hydrogen, halogen, an optionally substituted C₁₋₃ alkyl, an optionally substituted C₁₋₃ haloalkyl or an optionally substituted C₁₋₃ alkoxy; and



provided that the Compound (C) cannot be , or a pharmaceutically acceptable salt thereof.

[0070] In some embodiments, for Compound (C), or a pharmaceutically acceptable salt thereof, when X¹ is NH; Y¹ and Z¹ are each C; A¹ is a phenyl, 2-fluorophenyl or 2,6-difluorophenyl; R² and R³ are each methyl or one of R² and R³ is hydrogen and the other of R² and R³ is methyl; and R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each hydrogen; then R¹ cannot be 2-hydroxyethyl, 2-methylpropyl, 2-fluoro-2-methylpropyl, 3-fluoro-2-methylpropyl, 3-hydroxy-2-methylpropyl or 2-fluoro-3-hydroxy-2-methylpropyl. In other embodiments, for Compound (C), or a pharmaceutically acceptable salt thereof, when R¹⁰ is hydrogen, R¹¹ is hydrogen, X¹ is NH, Y¹ and Z¹ are each C, A¹ is an optionally substituted phenyl, one of R² and R³ is hydrogen or an optionally substituted C₁₋₆ alkyl and the other of R² and R³ is an optionally substituted C₁₋₆

alkyl, then R¹ cannot be a substituted C₁₋₆ alkyl substituted with one or more substituents selected from the group consisting of halogen and hydroxy.

[0071] In some embodiments, A¹ can be an optionally substituted aryl. For example, A¹ can be an optionally substituted phenyl. Thus, A¹ can be a substituted phenyl or an unsubstituted phenyl. In other embodiments, A¹ can be an optionally substituted cycloalkyl, such as an optionally substituted bicyclopentyl.

[0072] In some embodiments, R¹ can be selected from an optionally substituted C₁₋₆ alkyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkyl(C₁₋₆ alkyl), an optionally substituted heterocyclyl and an optionally substituted heterocyclyl(C₁₋₆ alkyl).

[0073] In some embodiments, R¹ can be a substituted cycloalkyl. In some embodiments, R¹ is substituted cycloalkyl that can be substituted with one or more substituents selected from halogen, hydroxy, haloalkyl, an optionally substituted alkyl, an optionally substituted cycloalkyl, a substituted alkoxy, a substituted mono-substituted amine and a substituted di-substituted amine. In some embodiments, R¹ can be an optionally substituted cycloalkyl selected from unsubstituted cyclobutyl, unsubstituted difluorocyclobutyl, unsubstituted cyclopentyl and unsubstituted bicyclopentyl. In other embodiments, R¹ can be an optionally substituted cycloalkyl(C₁₋₆ alkyl) selected from unsubstituted cyclopropylmethyl, unsubstituted bicyclopentylmethyl, unsubstituted fluorocyclopropylmethyl, unsubstituted fluorocyclobutylmethyl, unsubstituted methoxycyclopropylmethyl and unsubstituted trifluoromethylcyclopropylmethyl. In still other embodiments, R¹ can be an optionally substituted heterocyclyl selected from unsubstituted tetrahydropyranyl, unsubstituted tetrahydrofuranlyl, and unsubstituted oxetanyl. In yet still other embodiments, R¹ is an optionally substituted heterocyclyl(C₁₋₆ alkyl) can be selected from unsubstituted oxetanylmethyl and unsubstituted fluorooxetanylmethyl

[0074] In some embodiments, R¹ can be a substituted alkyl. In some embodiments, R¹ can be a substituted alkyl that is substituted with one or more substituents selected from halogen, hydroxy, haloalkyl, an optionally substituted cycloalkyl, a substituted alkoxy, a substituted mono-substituted amine and a substituted di-substituted amine. For example, R¹ can be a substituted alkyl that is a haloalkyl. In some embodiments, R¹ can be an optionally substituted C₁₋₆ alkyl selected from C₄ alkyl, fluoro(C₄ alkyl), and trifluoro(C₂ alkyl).


[0075] In some embodiments, R^2 and R^3 can be each independently selected from hydrogen, halogen, an optionally substituted C_{1-6} alkyl and an optionally substituted C_{1-6} haloalkyl. In other embodiments, R^2 and R^3 together with the carbon to which R^2 and R^3 are attached can form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl. In some embodiments, R^2 can be selected from hydrogen, methyl, fluoromethyl and difluoromethyl.


[0076] In some embodiments R^4 and R^5 can be each independently selected from hydrogen, halogen, an optionally substituted C_{1-6} alkyl and an optionally substituted C_{1-6} haloalkyl. In other embodiments, R^4 and R^5 together with the carbon to which R^4 and R^5 are attached can form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl.


[0077] In some embodiments, R^7 can be selected from halogen, hydroxy and unsubstituted alkoxy. For example, in some embodiments, R^7 can be selected from fluoro and methoxy.
















[0078] In some embodiments, R^{12} can be hydrogen. In other embodiments, R^{12} can be not hydrogen.


[0079] Examples of Compound (C) include the following:














[0080] Compound (A) and Compound (C), along with pharmaceutically acceptable salts of any of the foregoing, can be prepared as described herein and in WO 2017/172957, which is hereby incorporated by reference in its entirety. As described in WO 2017/172957, Compound (A) is an estrogen receptor alpha (ER α) inhibitor.

Definitions for Compound (B), and pharmaceutically acceptable salts thereof


[0081] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art. All patents, applications, published applications and other publications referenced herein are incorporated by reference in their entirety unless stated otherwise. In the event that there are a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

[0082] Whenever a group is described as being “optionally substituted” that group may be unsubstituted or substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) may be selected from one or more the indicated substituents. If no substituents are indicated, it is meant that the indicated “optionally substituted” or “substituted” group may be substituted with one or more group(s) individually and independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), cycloalkyl(alkyl), heteroaryl(alkyl), heterocyclyl(alkyl), hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-

carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, nitro, sulfenyl, sulfinyl, sulfonyl, haloalkyl, hydroxyalkyl, haloalkoxy, an amino, a mono-substituted amine group, a di-substituted amine group and an amine(C₁-C₆ alkyl).

[0083] As used herein, “C_a to C_b” in which “a” and “b” are integers refer to the number of carbon atoms in a group. The indicated group can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C₁ to C₄ alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, CH₃-, CH₃CH₂-, CH₃CH₂CH₂-, (CH₃)₂CH-, CH₃CH₂CH₂CH₂-, CH₃CH₂CH(CH₃)- and (CH₃)₃C-. If no “a” and “b” are designated, the broadest range described in these definitions is to be assumed.

[0084] If two “R” groups are described as being "taken together" the R groups and the atoms they are attached to can form a cycloalkyl, cycloalkenyl, aryl, heteroaryl or heterocycle. For example, without limitation, if R^a and R^b of an NR^aR^b group are indicated to be "taken together," it means that they are covalently bonded to one another to form a ring:



[0085] As used herein, the term “alkyl” refers to a fully saturated aliphatic hydrocarbon group. The alkyl moiety may be branched or straight chain. Examples of branched alkyl groups include, but are not limited to, iso-propyl, sec-butyl, t-butyl and the like. Examples of straight chain alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl and the like. The alkyl group may have 1 to 30 carbon atoms (whenever it appears herein, a numerical range such as “1 to 30” refers to each integer in the given range; *e.g.*, “1 to 30 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 30 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 12 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 6 carbon atoms. An alkyl group may be substituted or unsubstituted.

[0086] The term “alkenyl” used herein refers to a monovalent straight or branched chain radical of from two to twenty carbon atoms containing a carbon double bond(s) including,

but not limited to, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like. An alkenyl group may be unsubstituted or substituted.

[0087] The term “alkynyl” used herein refers to a monovalent straight or branched chain radical of from two to twenty carbon atoms containing a carbon triple bond(s) including, but not limited to, 1-propynyl, 1-butylnyl, 2-butylnyl and the like. An alkynyl group may be unsubstituted or substituted.

[0088] As used herein, “cycloalkyl” refers to a completely saturated (no double or triple bonds) mono- or multi- cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. As used herein, the term “fused” refers to two rings which have two atoms and one bond in common. As used herein, the term “bridged cycloalkyl” refers to compounds wherein the cycloalkyl contains a linkage of one or more atoms connecting non-adjacent atoms. As used herein, the term “spiro” refers to two rings which have one atom in common and the two rings are not linked by a bridge. Cycloalkyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). A cycloalkyl group may be unsubstituted or substituted. Examples of mono-cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. Examples of fused cycloalkyl groups are decahydronaphthalenyl, dodecahydro-1H-phenalenyl and tetradecahydroanthracenyl; examples of bridged cycloalkyl groups are bicyclo[1.1.1]pentyl, adamantanyl and norbornanyl; and examples of spiro cycloalkyl groups include spiro[3.3]heptane and spiro[4.5]decane.

[0089] As used herein, “cycloalkenyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all the rings (otherwise the group would be “aryl,” as defined herein). Cycloalkenyl groups can contain 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). When composed of two or more rings, the rings may be connected together in a fused, bridged or spiro fashion. A cycloalkenyl group may be unsubstituted or substituted.

[0090] As used herein, “carbocyclyl” refers to a non-aromatic a mono- or multi-cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion, as described herein. Carbocyclyl groups can

contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). A carbocyclyl group may be unsubstituted or substituted. Examples of carbocyclyl groups include, but are in no way limited to, cycloalkyl groups and cycloalkenyl groups, as defined herein, and the non-aromatic portions of 1,2,3,4-tetrahydronaphthalene, 2,3-dihydro-1H-indene, 5,6,7,8-tetrahydroquinoline and 6,7-dihydro-5H-cyclopenta[b]pyridine.

[0091] As used herein, “aryl” refers to a carbocyclic (all carbon) monocyclic or polycyclic aromatic ring system (including fused ring systems where two carbocyclic rings share a chemical bond) that has a fully delocalized pi-electron system throughout all the rings. The number of carbon atoms in an aryl group can vary. For example, the aryl group can be a C₆-C₁₄ aryl group, a C₆-C₁₀ aryl group or a C₆ aryl group. Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted.

[0092] As used herein, “heteroaryl” refers to a monocyclic or polycyclic aromatic ring system (a ring system with fully delocalized pi-electron system) that contain(s) one or more heteroatoms (for example, 1, 2 or 3 heteroatoms), that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur. The number of atoms in the ring(s) of a heteroaryl group can vary. For example, the heteroaryl group can contain 4 to 14 atoms in the ring(s), 5 to 10 atoms in the ring(s) or 5 to 6 atoms in the ring(s), such as nine carbon atoms and one heteroatom; eight carbon atoms and two heteroatoms; seven carbon atoms and three heteroatoms; eight carbon atoms and one heteroatom; seven carbon atoms and two heteroatoms; six carbon atoms and three heteroatoms; five carbon atoms and four heteroatoms; five carbon atoms and one heteroatom; four carbon atoms and two heteroatoms; three carbon atoms and three heteroatoms; four carbon atoms and one heteroatom; three carbon atoms and two heteroatoms; or two carbon atoms and three heteroatoms. Furthermore, the term “heteroaryl” includes fused ring systems where two rings, such as at least one aryl ring and at least one heteroaryl ring or at least two heteroaryl rings, share at least one chemical bond. Examples of heteroaryl rings include, but are not limited to, furan, furazan, thiophene, benzothiophene, phthalazine, pyrrole, oxazole, benzoxazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, thiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, benzothiazole, imidazole, benzimidazole, indole, indazole, pyrazole, benzopyrazole, isoxazole, benzoisoxazole, isothiazole, triazole, benzotriazole, thiadiazole, tetrazole, pyridine, pyridazine,

pyrimidine, pyrazine, purine, pteridine, quinoline, isoquinoline, quinazoline, quinoxaline, cinnoline and triazine. A heteroaryl group may be substituted or unsubstituted.

[0093] As used herein, “heterocyclyl” or “heteroalicyclyl” refers to three-, four-, five-, six-, seven-, eight-, nine-, ten-, up to 18-membered monocyclic, bicyclic and tricyclic ring system wherein carbon atoms together with from 1 to 5 heteroatoms constitute said ring system. A heterocycle may optionally contain one or more unsaturated bonds situated in such a way, however, that a fully delocalized pi-electron system does not occur throughout all the rings. The heteroatom(s) is an element other than carbon including, but not limited to, oxygen, sulfur and nitrogen. A heterocycle may further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo-systems and thio-systems such as lactams, lactones, cyclic imides, cyclic thioimides and cyclic carbamates. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. As used herein, the term “fused” refers to two rings which have two atoms and one bond in common. As used herein, the term “bridged heterocyclyl” or “bridged heteroalicyclyl” refers to compounds wherein the heterocyclyl or heteroalicyclyl contains a linkage of one or more atoms connecting non-adjacent atoms. As used herein, the term “spiro” refers to two rings which have one atom in common and the two rings are not linked by a bridge. Heterocyclyl and heteroalicyclyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). For example, five carbon atoms and one heteroatom; four carbon atoms and two heteroatoms; three carbon atoms and three heteroatoms; four carbon atoms and one heteroatom; three carbon atoms and two heteroatoms; two carbon atoms and three heteroatoms; one carbon atom and four heteroatoms; three carbon atoms and one heteroatom; or two carbon atoms and one heteroatom. Additionally, any nitrogens in a heteroalicyclic may be quaternized. Heterocyclyl or heteroalicyclic groups may be unsubstituted or substituted. Examples of such “heterocyclyl” or “heteroalicyclyl” groups include but are not limited to, 1,3-dioxin, 1,3-dioxane, 1,4-dioxane, 1,2-dioxolane, 1,3-dioxolane, 1,4-dioxolane, 1,3-oxathiane, 1,4-oxathiin, 1,3-oxathiolane, 1,3-dithiole, 1,3-dithiolane, 1,4-oxathiane, tetrahydro-1,4-thiazine, 2H-1,2-oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouracil, trioxane, hexahydro-1,3,5-triazine, imidazoline, imidazolidine, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, morpholine, oxirane, piperidine N-Oxide, piperidine, piperazine, pyrrolidine, azepane,


pyrrolidone, pyrrolidione, 4-piperidone, pyrazoline, pyrazolidine, 2-oxopyrrolidine, tetrahydropyran, 4H-pyran, tetrahydrothiopyran, thiamorpholine, thiamorpholine sulfoxide, thiamorpholine sulfone and their benzo-fused analogs (e.g., benzimidazolidinone, tetrahydroquinoline and/or 3,4-methylenedioxyphenyl). Examples of spiro heterocyclyl groups include 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 2-oxa-6-azaspiro[3.3]heptane, 2,6-diazaspiro[3.3]heptane, 2-oxaspiro[3.4]octane and 2-azaspiro[3.4]octane.

[0094] As used herein, “aralkyl” and “aryl(alkyl)” refer to an aryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, 2-phenylalkyl, 3-phenylalkyl and naphthylalkyl.

[0095] As used herein, “heteroaralkyl” and “heteroaryl(alkyl)” refer to a heteroaryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and heteroaryl group of heteroaralkyl may be substituted or unsubstituted. Examples include but are not limited to 2-thienylalkyl, 3-thienylalkyl, furylalkyl, thienylalkyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl and imidazolylalkyl and their benzo-fused analogs.

[0096] A “heteroalicyclyl(alkyl)” and “heterocyclyl(alkyl)” refer to a heterocyclic or a heteroalicyclic group connected, as a substituent, via a lower alkylene group. The lower alkylene and heterocyclyl of a (heteroalicyclyl)alkyl may be substituted or unsubstituted. Examples include but are not limited tetrahydro-2H-pyran-4-yl(methyl), piperidin-4-yl(ethyl), piperidin-4-yl(propyl), tetrahydro-2H-thiopyran-4-yl(methyl) and 1,3-thiazinan-4-yl(methyl).

[0097] As used herein, “lower alkylene groups” are straight-chained -CH₂- tethering groups, forming bonds to connect molecular fragments via their terminal carbon atoms. Examples include but are not limited to methylene (-CH₂-), ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-) and butylene (-CH₂CH₂CH₂CH₂-). A lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene group and/or by substituting both

hydrogens on the same carbon with a cycloalkyl group (e.g., ).

[0098] As used herein, the term “hydroxy” refers to a -OH group.

[0099] As used herein, “alkoxy” refers to the Formula -OR wherein R is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl) is defined herein. A non-limiting list of

alkoxys are methoxy, ethoxy, n-propoxy, 1-methylethoxy (iso-propoxy), n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, phenoxy and benzyloxy. An alkoxy may be substituted or unsubstituted.

[0100] As used herein, “acyl” refers to a hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl) and heterocyclyl(alkyl) connected, as substituents, via a carbonyl group. Examples include formyl, acetyl, propanoyl, benzoyl and acryl. An acyl may be substituted or unsubstituted.

[0101] A “cyano” group refers to a “-CN” group.

[0102] The term “halogen atom” or “halogen” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, such as, fluorine, chlorine, bromine and iodine.

[0103] A “thiocarbonyl” group refers to a “-C(=S)R” group in which R can be the same as defined with respect to O-carboxy. A thiocarbonyl may be substituted or unsubstituted.

[0104] An “O-carbamyl” group refers to a “-OC(=O)N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-carbamyl may be substituted or unsubstituted.

[0105] An “N-carbamyl” group refers to an “ROC(=O)N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-carbamyl may be substituted or unsubstituted.

[0106] An “O-thiocarbamyl” group refers to a “-OC(=S)-N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-thiocarbamyl may be substituted or unsubstituted.

[0107] An “N-thiocarbamyl” group refers to an “ROC(=S)N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-thiocarbamyl may be substituted or unsubstituted.

[0108] A “C-amido” group refers to a “-C(=O)N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl,

aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A C-amido may be substituted or unsubstituted.

[0109] An “N-amido” group refers to a “RC(=O)N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-amido may be substituted or unsubstituted.

[0110] An “S-sulfonamido” group refers to a “-SO₂N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An S-sulfonamido may be substituted or unsubstituted.

[0111] An “N-sulfonamido” group refers to a “RSO₂N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-sulfonamido may be substituted or unsubstituted.

[0112] An “O-carboxy” group refers to a “RC(=O)O-” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. An O-carboxy may be substituted or unsubstituted.

[0113] The terms “ester” and “C-carboxy” refer to a “-C(=O)OR” group in which R can be the same as defined with respect to O-carboxy. An ester and C-carboxy may be substituted or unsubstituted.

[0114] A “nitro” group refers to an “-NO₂” group.

[0115] A “sulfenyl” group refers to an “-SR” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A sulfenyl may be substituted or unsubstituted.

[0116] A “sulfinyl” group refers to an “-S(=O)-R” group in which R can be the same as defined with respect to sulfenyl. A sulfinyl may be substituted or unsubstituted.

[0117] A “sulfonyl” group refers to an “SO₂R” group in which R can be the same as defined with respect to sulfenyl. A sulfonyl may be substituted or unsubstituted.

[0118] As used herein, “haloalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkyl, di-haloalkyl, tri-haloalkyl and polyhaloalkyl). Such groups include but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 1-chloro-2-fluoromethyl, 2-fluoroisobutyl and pentafluoroethyl. A haloalkyl may be substituted or unsubstituted.

[0119] As used herein, “haloalkoxy” refers to an alkoxy group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkoxy, di-haloalkoxy and tri-haloalkoxy). Such groups include but are not limited to, chloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 1-chloro-2-fluoromethoxy and 2-fluoroisobutoxy. A haloalkoxy may be substituted or unsubstituted.

[0120] The term “amino” as used herein refers to a $-NH_2$ group.

[0121] A “mono-substituted amine” group refers to a “ $-NHR_A$ ” group in which R_A can be an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. The R_A may be substituted or unsubstituted. Examples of mono-substituted amino groups include, but are not limited to, $-NH(\text{methyl})$, $-NH(\text{phenyl})$ and the like.

[0122] A “di-substituted amine” group refers to a “ $-NR_AR_B$ ” group in which R_A and R_B can be independently an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. R_A and R_B can independently be substituted or unsubstituted. Examples of di-substituted amino groups include, but are not limited to, $-N(\text{methyl})_2$, $-N(\text{phenyl})(\text{methyl})$, $-N(\text{ethyl})(\text{methyl})$ and the like.

[0123] As used herein, “amine(alkyl)” group refers to an $-(\text{alkylene})-NR'R''$ radical where R' and R'' are independently hydrogen or alkyl as defined herein. An amine(alkyl) may be substituted or unsubstituted. Examples of amine(alkyl) groups include, but are not limited to, $-CH_2NH(\text{methyl})$, $-CH_2NH(\text{phenyl})$, $-CH_2CH_2NH(\text{methyl})$, $-CH_2CH_2NH(\text{phenyl})$, $-CH_2N(\text{methyl})_2$, $-CH_2N(\text{phenyl})(\text{methyl})$, $-NCH_2(\text{ethyl})(\text{methyl})$, $-CH_2CH_2N(\text{methyl})_2$, $-CH_2CH_2N(\text{phenyl})(\text{methyl})$, $-NCH_2CH_2(\text{ethyl})(\text{methyl})$ and the like.

[0124] Where the number of substituents is not specified (e.g. haloalkyl), there may be one or more substituents present. For example, “haloalkyl” may include one or more of the

same or different halogens. As another example, “C₁-C₃ alkoxyphenyl” may include one or more of the same or different alkoxy groups containing one, two or three atoms.

[0125] As used herein, a radical indicates species with a single, unpaired electron such that the species containing the radical can be covalently bonded to another species. Hence, in this context, a radical is not necessarily a free radical. Rather, a radical indicates a specific portion of a larger molecule. The term “radical” can be used interchangeably with the term “group.”

[0126] The term “pharmaceutically acceptable salt” refers to a salt of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, the salt is an acid addition salt of the compound. Pharmaceutical salts can be obtained by reacting a compound with inorganic acids such as hydrohalic acid (e.g., hydrochloric acid or hydrobromic acid), a sulfuric acid, a nitric acid and a phosphoric acid (such as 2,3-dihydroxypropyl dihydrogen phosphate). Pharmaceutical salts can also be obtained by reacting a compound with an organic acid such as aliphatic or aromatic carboxylic or sulfonic acids, for example formic, acetic, succinic, lactic, malic, tartaric, citric, ascorbic, nicotinic, methanesulfonic, ethanesulfonic, p-toluenesulfonic, trifluoroacetic, benzoic, salicylic, 2-oxopentanedioic or naphthalenesulfonic acid. Pharmaceutical salts can also be obtained by reacting a compound with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a sodium, a potassium or a lithium salt, an alkaline earth metal salt, such as a calcium or a magnesium salt, a salt of a carbonate, a salt of a bicarbonate, a salt of organic bases such as dicyclohexylamine, N-methyl-D-glucamine, tris(hydroxymethyl)methylamine, C₁-C₇ alkylamine, cyclohexylamine, triethanolamine, ethylenediamine and salts with amino acids such as arginine and lysine. Those skilled in the art understand that when a salt is formed by protonation of a nitrogen-based group (for example, NH₂), the nitrogen-based group can be associated with a positive charge (for example, NH₂ can become NH₃⁺) and the positive charge can be balanced by a negatively charged counterion (such as Cl⁻).

[0127] It is understood that, in any compound described herein having one or more chiral centers, if an absolute stereochemistry is not expressly indicated, then each center may independently be of R-configuration or S-configuration or a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, enantiomerically enriched, racemic

mixture, diastereomerically pure, diastereomerically enriched or a stereoisomeric mixture. In addition, it is understood that, in any compound described herein having one or more double bond(s) generating geometrical isomers that can be defined as E or Z, each double bond may independently be E or Z a mixture thereof. Likewise, it is understood that, in any compound described, all tautomeric forms are also intended to be included.

[0128] It is to be understood that where compounds disclosed herein have unfilled valencies, then the valencies are to be filled with hydrogens or isotopes thereof, e.g., hydrogen-1 (protium) and hydrogen-2 (deuterium).

[0129] It is understood that the compounds described herein can be labeled isotopically. Substitution with isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements. Each chemical element as represented in a compound structure may include any isotope of said element. For example, in a compound structure a hydrogen atom may be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom may be present, the hydrogen atom can be any isotope of hydrogen, including but not limited to hydrogen-1 (protium) and hydrogen-2 (deuterium). Thus, reference herein to a compound encompasses all potential isotopic forms unless the context clearly dictates otherwise.

[0130] It is understood that the methods and combinations described herein include crystalline forms (also known as polymorphs, which include the different crystal packing arrangements of the same elemental composition of a compound), amorphous phases, salts, solvates and hydrates. In some embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, ethanol or the like. In other embodiments, the compounds described herein exist in unsolvated form. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol or the like. Hydrates are formed when the solvent is water or alcoholates are formed when the solvent is alcohol. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.


[0131] Where a range of values is provided, it is understood that the upper and lower limit, and each intervening value between the upper and lower limit of the range is encompassed within the embodiments.

[0132] Terms and phrases used in this application, and variations thereof, especially in the appended claims, unless otherwise expressly stated, should be construed as open ended as opposed to limiting. As examples of the foregoing, the term 'including' should be read to mean 'including, without limitation,' 'including but not limited to,' or the like; the term 'comprising' as used herein is synonymous with 'including,' 'containing,' or 'characterized by,' and is inclusive or open-ended and does not exclude additional, unrecited elements or method steps; the term 'having' should be interpreted as 'having at least;' the term 'includes' should be interpreted as 'includes but is not limited to;' the term 'example' is used to provide exemplary instances of the item in discussion, not an exhaustive or limiting list thereof; and use of terms like 'preferably,' 'preferred,' 'desired,' or 'desirable,' and words of similar meaning should not be understood as implying that certain features are critical, essential, or even important to the structure or function, but instead as merely intended to highlight alternative or additional features that may or may not be utilized in a particular embodiment. In addition, the term "comprising" is to be interpreted synonymously with the phrases "having at least" or "including at least". When used in the context of a compound, composition or device, the term "comprising" means that the compound, composition or device includes at least the recited features or components, but may also include additional features or components.



[0133] With respect to the use of substantially any plural and/or singular terms herein, those having skill in the art can translate from the plural to the singular and/or from the singular to the plural as is appropriate to the context and/or application. The various singular/plural permutations may be expressly set forth herein for sake of clarity. The indefinite article "a" or "an" does not exclude a plurality. The mere fact that certain measures are recited in mutually different dependent claims does not indicate that a combination of these measures cannot be used to advantage. Any reference signs in the claims should not be construed as limiting the scope.


Compound (B)

[0134] As described herein, some embodiments disclosed herein relate to the use of a combination of compounds for treating a disease or condition, wherein the combination can include an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof (as described herein), and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein: Compound (B) has the structure:



wherein: R^{1a} can be selected from hydrogen, halogen and a substituted or unsubstituted C₁-C₆ alkyl; Ring A-a can be selected from a substituted or unsubstituted phenyl and a substituted or unsubstituted 5-6 membered monocyclic heteroaryl; Ring B-a can be selected from a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl and a substituted or unsubstituted 5-7


membered monocyclic heterocyclyl; R^{2a} can be selected from ,  and




; m-a can be 0, 1, 2 or 3; R^{3a} can be selected from halogen and a substituted or unsubstituted C₁-C₆ alkyl; X-a can be selected from hydrogen, halogen, hydroxy, cyano, a substituted or unsubstituted 4-6 membered monocyclic heterocyclyl, a substituted or unsubstituted amine(C₁-C₆ alkyl), a substituted or unsubstituted -NH-(CH₂)₁₋₆-amine, a mono-substituted amine, a di-substituted amine, an amino, a substituted or unsubstituted C₁-C₆ alkyl, a substituted or unsubstituted C₁-C₆ alkoxy, a substituted or unsubstituted C₃-C₆ cycloalkoxy, a substituted or unsubstituted (C₁-C₆ alkyl)acyl, a substituted or unsubstituted C-amido, a substituted or unsubstituted N-amido, a substituted or unsubstituted C-carboxy, a substituted or unsubstituted O-carboxy, a substituted or unsubstituted O-carbamyl and a substituted or unsubstituted N-carbamyl; Y-a can be CH or N; Y^{1-a} can be CR^{4A-a} or N; Y^{2-a} can be CR^{4B-a} or N;

Ring C-a can be selected from a substituted or unsubstituted C₆-C₁₀ aryl, a substituted or unsubstituted monocyclic 5-10 membered heteroaryl, a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl, a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl and a substituted or unsubstituted 7-10 membered bicyclic heterocyclyl; R^{4A-a} and R^{4B-a} can be independently selected from hydrogen, halogen and an unsubstituted C₁₋₄ alkyl; and R^{5-a} can be a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl.

[0135] In some embodiments, R^{1a} can be selected from hydrogen, halogen and a substituted or unsubstituted C₁-C₆ alkyl. In some embodiments, Ring A-a can be selected from a substituted or unsubstituted phenyl and a substituted or unsubstituted 5-6 membered monocyclic heteroaryl. In some embodiments, Ring B-a can be selected from a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl and a substituted or unsubstituted 5-7 membered



monocyclic heterocyclyl. In some embodiments, R^{2a} can be selected from



. In some embodiments, m-a can be 0, 1, 2 or 3. In some embodiments, R^{3a} can be selected from halogen and a substituted or unsubstituted C₁-C₆ alkyl. In some embodiments, X-a can be selected from hydrogen, halogen, hydroxy, cyano, a substituted or unsubstituted 4-6 membered monocyclic heterocyclyl, a substituted or unsubstituted amine(C₁-C₆ alkyl), a substituted or unsubstituted -NH-(CH₂)₁₋₆-amine, a mono-substituted amine, a di-substituted amine, an amino, a substituted or unsubstituted C₁-C₆ alkyl, a substituted or unsubstituted C₁-C₆ alkoxy, a substituted or unsubstituted C₃-C₆ cycloalkoxy, a substituted or unsubstituted (C₁-C₆ alkyl)acyl, a substituted or unsubstituted C-amido, a substituted or unsubstituted N-amido, a substituted or unsubstituted C-carboxy, a substituted or unsubstituted O-carboxy, a substituted or unsubstituted O-carbamyl and a substituted or unsubstituted N-carbamyl. In some embodiments, Y-a can be CH or N. In some embodiments, Y^{1-a} can be CR^{4A-a} or N. In some embodiments, Y^{2-a} can be CR^{4B-a} or N. In some embodiments, Ring C-a can be selected from a substituted or unsubstituted C₆-C₁₀ aryl, a substituted or unsubstituted monocyclic 5-10 membered heteroaryl, a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl, a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl and a substituted or unsubstituted 7-10

membered bicyclic heterocyclyl. In some embodiments, R^{4A-a} and R^{4B-a} are independently selected from hydrogen, halogen and an unsubstituted C₁₋₄ alkyl.


[0136] In some embodiments, R^{1a} can be selected from hydrogen, halogen and C_{1-C6} alkyl. In some embodiments, R^{1a} can be hydrogen. In other embodiments, R^{1a} can be halogen. In some embodiments, R^{1a} can be fluoro. In still other embodiments, R^{1a} can be an unsubstituted C_{1-C6} alkyl (such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, pentyl (straight chain or branched) or hexyl (straight chain or branched)). In some embodiments, R^{1a} can be an unsubstituted methyl. In some embodiments, R^{1a} can be a substituted C_{1-C6} alkyl, such as those described herein. In some embodiments, R^{1a} can be an unsubstituted C_{1-C6} haloalkyl (such as a C_{1-C6} fluoroalkyl, a C_{1-C6} chloroalkyl or a C_{1-C6} chlorofluoroalkyl). In some embodiments, R^{1a} can be -CHF₂, -CF₃, -CF₂CH₃ or -CH₂CF₃.



[0137] In some embodiments, Ring A-a can be selected from a substituted or unsubstituted phenyl and a substituted or unsubstituted 5-6 membered monocyclic heteroaryl.



[0138] In some embodiments, Ring A-a can be a substituted phenyl. In other embodiments, Ring A can be an unsubstituted phenyl.


[0139] In some embodiments, Ring A-a can be a substituted 5-6 membered monocyclic heteroaryl. In some embodiments, Ring A-a can be an unsubstituted 5-6 membered monocyclic heteroaryl. In some embodiments, Ring A-a can be selected from a substituted or unsubstituted pyrrole, a substituted or unsubstituted furan, a substituted or unsubstituted thiophene, a substituted or unsubstituted imidazole, a substituted or unsubstituted pyrazole, a substituted or unsubstituted oxazole, a substituted or unsubstituted thiazole, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine and a substituted or unsubstituted pyridazine.



[0140] When substituted, Ring A-a can be substituted with one or more substituents selected from halogen, an unsubstituted C_{1-C4} haloalkyl and an unsubstituted C_{1-C4} alkyl. In some embodiments, Ring A-a is mono-substituted with a halogen (for example, fluoro).







In some embodiments,  can be a substituted or unsubstituted . In some

embodiments,  can be a substituted or unsubstituted , wherein the Ring A-a is





unsubstituted. In other embodiments,  can be selected from a substituted or

unsubstituted , a substituted or unsubstituted  and a substituted or unsubstituted




















. As described herein, the Ring A-a portion of , a  and  can be unsubstituted.


[0142] In some embodiments, Ring B-a can be selected from a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl and a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl.

[0143] In some embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl. In some embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 5 membered carbocyclyl. In other embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 6 membered carbocyclyl. In still other embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 7 membered carbocyclyl.



[0144] In some embodiments,  can be selected from: , , and ; wherein each of the aforementioned groups are substituted or unsubstituted.

[0145] In some embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 5-7 membered heterocyclyl. In some embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 5 membered heterocyclyl. In other embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 6 membered heterocyclyl. In still other embodiments, Ring B-a can be a substituted or unsubstituted monocyclic 7 membered heterocyclyl.


[0146] In some embodiments,  can be selected from: , , , , , , , , , , , , , , , ,  and ; wherein each of the aforementioned groups are substituted or unsubstituted, including any -NH group.



[0148] In some embodiments, when Ring B-a is substituted, Ring B-a can be substituted with 1, 2 or 3 substituents independently selected from halogen, hydroxy, amino, an unsubstituted N-linked amido (for example, –NHC(O)C₁-C₆ alkyl), an unsubstituted C₁-C₆ haloalkyl (such as those described herein) and a substituted or unsubstituted C₁-C₆ alkyl (such as those described herein). In some embodiments, when Ring B-a is substituted, Ring B-a can be substituted with 1, 2 or 3 substituents independently selected from halogen, hydroxy, amino, an unsubstituted N-linked amido (for example, –NHC(O)C₁-C₆ alkyl) and a substituted or unsubstituted C₁-C₆ alkyl (such as those described herein). In some embodiments, Ring B-a can be substituted with 1, 2 or 3 substituents independently selected from fluoro, hydroxy, amino, an unsubstituted –NHC(O)C₁-C₆ alkyl, an unsubstituted C₁-C₆ haloalkyl (such as those described herein) and an unsubstituted C₁-C₆ alkyl (such as those described herein). In some embodiments, Ring B-a can be substituted with 1 or 2 substituents independently selected from fluoro, hydroxy, –CF₃, –CHF₂, –CF₂CH₃, an unsubstituted methyl, an unsubstituted ethyl and –NHC(O)CH₃.




[0149] In some embodiments, can be selected from:





; wherein each of the aforementioned groups are substituted or unsubstituted, including any -NH group.

[0150]


In some embodiments,



can be selected from:




and ; wherein each of the




aforementioned groups are substituted or unsubstituted. In some embodiments,

can be




selected from:




and ; wherein each of the aforementioned groups are substituted or

unsubstituted. In some embodiments,




can be a substituted or unsubstituted




. In


some embodiments,





can be a substituted or






[0151] Both Ring A-a and Ring B-a can be substituted or unsubstituted. In some


embodiments, Ring A-a and Ring B-a of  can be independently substituted or



unsubstituted. In some embodiments, Ring A-a and Ring B-a of  can be both


unsubstituted. In some embodiments, Ring A-a and Ring B-a of  can be both


independently substituted. In some embodiments, Ring A-a of  can be substituted and


Ring B of  can be unsubstituted. In some embodiments, Ring A-a of  can be


unsubstituted and Ring B-a of  can be substituted. In some embodiments, Ring A of



 can be unsubstituted and Ring B-a of  can be substituted with 1, 2 or 3 substituents independently selected from halogen, hydroxy and a substituted or unsubstituted C₁-

C₆ alkyl (such as those described herein). In some embodiments, Ring A-a of  can be




unsubstituted and Ring B-a of  can be substituted with 1, 2 or 3 substituents independently selected from fluoro, hydroxy, amino, an unsubstituted N-linked amido (for example, $-\text{NHC(O)C}_1\text{-C}_6$ alkyl), an unsubstituted $\text{C}_1\text{-C}_6$ haloalkyl (such as those described herein) and an unsubstituted $\text{C}_1\text{-C}_6$ alkyl (such as those described herein). In some embodiments,




Ring A-a of  can be unsubstituted and Ring B-a of  can be substituted with 1 or 2 substituents independently selected from fluoro, hydroxy, amino, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CF}_2\text{CH}_3$, an unsubstituted methyl, an unsubstituted ethyl and $-\text{NHC(O)CH}_3$.

[0152]


In some embodiments, R^{2a} can be selected from



In some embodiments, R^{2a} can be



. In some embodiments, R^{2a} can be



[0153]

In some embodiments, Y-a can be CH or N (nitrogen). In some embodiments, Y-a can be CH. In some embodiments, Y-a can be N (nitrogen).

[0154]

In some embodiments, R^{3a} can be selected from halogen and a substituted or unsubstituted $\text{C}_1\text{-C}_6$ alkyl (such as those described herein). In some embodiments, R^{3a} can be halogen. In some embodiments, R^{3a} can be a substituted $\text{C}_1\text{-C}_6$ alkyl (such as those described herein). In some embodiments, R^{3a} can be an unsubstituted $\text{C}_1\text{-C}_6$ alkyl (such as those described herein).



[0155]


In some embodiments, m-a can be 0, 1, 2 or 3. In some embodiments, m-a can be 0. In some embodiments, m-a can be 1. In some embodiments, m-a can be 2. In some embodiments, m-a can be 3. When m-a is 2 or 3, the R^{3a} groups can be the same or different from each other.

[0156] In some embodiments, X-a can be selected from hydrogen, halogen, hydroxy, cyano, a substituted or unsubstituted 4-6 membered monocyclic heterocyclyl, a substituted or unsubstituted amine(C₁-C₆ alkyl), a substituted or unsubstituted -NH-(CH₂)₁₋₆-amine, a mono-substituted amine, a di-substituted amine, an amino, a substituted or unsubstituted C₁-C₆ alkyl (such as those described herein), a substituted or unsubstituted C₁-C₆ alkoxy (such as methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-butoxy, t-butoxy, pentoxy (straight chain or branched) or hexoxy (straight chain or branched)), a substituted or unsubstituted C₃-C₆ cycloalkoxy (such as cyclopropoxy, cyclobutoxy, cyclopentoxy or cyclohexoxy), a substituted or unsubstituted (C₁-C₆ alkyl)acyl, a substituted or unsubstituted C-amido, a substituted or unsubstituted N-amido, a substituted or unsubstituted C-carboxy, a substituted or unsubstituted O-carboxy, a substituted or unsubstituted O-carbamyl and a substituted or unsubstituted N-carbamyl.


[0157] In some embodiments, X-a can be hydrogen. In other embodiments, X-a can be halogen. In some embodiments, X-a can be fluoro. In some embodiments, X-a can be chloro. In still other embodiments, X-a can be hydroxy. In yet still other embodiments, X-a can be cyano. In some embodiments, X-a can be an amino.

[0158] In some embodiments, X-a can be an unsubstituted C₁-C₆ alkyl (such as those described herein). In some embodiments, X-a can be an unsubstituted methyl, an unsubstituted ethyl or an unsubstituted iso-propyl. In some embodiments, X-a can be a substituted C₁-C₆ alkyl (such as those described herein). In some embodiments, X-a can be an unsubstituted C₁-C₆ haloalkyl (such as a C₁-C₆ fluoroalkyl, a C₁-C₆ chloroalkyl or a C₁-C₆ chlorofluoroalkyl). In some embodiments, X-a can be selected from -CHF₂, -CF₃, -CF₂CH₃ and -CH₂CF₃. In some embodiments, X-a can be an unsubstituted C₁-C₆ hydroxyalkyl (such as a C₁-C₆ mono-hydroxyalkyl or a C₁-C₆ di-hydroxyalkyl). In some embodiments, X-a can be selected from -CH₂OH, -CH₂CH₂OH, -CH(OH)CH₃ and -C(OH)(CH₃)₂. In some embodiments, X-a can be an unsubstituted C₁-C₆ cyanoalkyl (such as a C₁-C₆ mono-cyanoalkyl or a C₁-C₆ di-cyanoalkyl). In

some embodiments, X-a can be selected from . In some embodiments, X-a can be an unsubstituted C₁-C₆ alkoxyalkyl (such as a C₁-C₆ mono-alkoxyalkyl or a C₁-C₆ di-alkoxyalkyl). In some embodiments, X-a can be selected from .




[0159] In some embodiments, X-a can be an unsubstituted C₁-C₆ alkoxy (such as those described herein). In some embodiments, X-a can be an unsubstituted methoxy, an unsubstituted ethoxy or an unsubstituted iso-propoxy. In some embodiments, X-a can be a substituted C₁-C₆ alkoxy (such as those described herein). In some embodiments, X-a can be a C₁-C₆ alkoxy substituted with 1, 2 or 3 substituents independently selected from halogen, an amino, a mono-substituted amine (such as those described herein) and a di-substituted amine (such as those described herein). In some embodiments, X-a can be a C₁-C₆ alkoxy substituted with 1 substituent selected from halogen, an amino, a mono-substituted amine (such as those described herein) and a di-substituted amine (such as those described herein).





[0161] In some embodiments, X-a can be a substituted C₃-C₆ cycloalkoxy (such as those described herein). In some embodiments, X-a can be an unsubstituted C₃-C₆ cycloalkoxy (such as those described herein).









[0162] In some embodiments, X-a can be a substituted (C₁-C₆ alkyl)acyl, such as a substituted -(CO)-CH₃. In some embodiments, X-a can be an unsubstituted (C₁-C₆ alkyl)acyl, such as an unsubstituted -(CO)-CH₃.







[0163] In some embodiments, X-a can be a substituted 4-6 membered monocyclic heterocyclyl. In some embodiments, X-a can be an unsubstituted 4-6 membered monocyclic heterocyclyl. In some embodiments, X-a can be selected from azetidine, oxetane, diazetidene, azaoxetane, pyrrolidine, tetrahydrofuran, imidazoline, pyrazolidine, piperidine, tetrahydropyran, piperazine, morpholine and dioxane; wherein each of the aforementioned groups are substituted or unsubstituted, including any -NH group. . In some embodiments, X-a can be selected from









 and  ; wherein each of the aforementioned groups are substituted or unsubstituted, including any –NH group.

[0164] In some embodiments, X-a can be a 4-6 membered monocyclic heterocyclyl (such as those described herein) substituted with 1 or 2 substituents independently selected from halogen, a substituted or unsubstituted C₁-C₆ alkyl (such as those described herein), a mono-substituted amine (such as those described herein), a di-substituted amine (such as those described herein), an amino, substituted or unsubstituted amine(C₁-C₆ alkyl) and a substituted or unsubstituted (C₁-C₆ alkyl)acyl. In some embodiments, X-a can be a 4-6 membered monocyclic heterocyclyl substituted with 1 or 2 substituents independently selected from fluoro, an unsubstituted methyl, an unsubstituted ethyl, an unsubstituted iso-propyl, –CH₂OH and –


N(CH₃)₂. In some embodiments, X-a can be selected from , , , , , ,  and .

[0165] In some embodiments, X-a can be a substituted amine(C₁-C₆ alkyl). In some embodiments, X-a can be an unsubstituted amine(C₁-C₆ alkyl). In some embodiments, X-a can be selected from , , , ,  and  ; wherein each of the aforementioned groups are substituted or unsubstituted, including any –NH group. .

[0166] In some embodiments, X-a can be a substituted –NH-(CH₂)₁₋₆-amine. In some embodiments, X-a can be an unsubstituted –NH-(CH₂)₁₋₆-amine. In some embodiments, X-a can be selected from , , , ,  and  ; wherein each of the aforementioned groups are substituted or unsubstituted, including any –NH group.

[0167] In some embodiments, X-a can be a mono-substituted amine. In some embodiments, the substituent of the mono-substituted amine is an unsubstituted C₁-C₆ alkyl (such as those as described herein) or an unsubstituted C₃-C₆ cycloalkyl (such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl).

[0168] In some embodiments, X-a can be a di-substituted amine. In some embodiments, the two substituents of the di-substituted amine are independently selected from an unsubstituted C₁-C₆ alkyl (such as those as described herein) and an unsubstituted C₃-C₆ cycloalkyl (such as those as described herein).



[0170] In some embodiments, X-a can be a substituted or unsubstituted C-amido. In some embodiments, X-a can be a substituted or unsubstituted N-amido. In some embodiments, X-a can be a substituted or unsubstituted C-carboxy. In some embodiments, X-a can be a substituted or unsubstituted O-carboxy. In some embodiments, X-a can be a substituted or unsubstituted O-carbamyl. In some embodiments, X-a can be a substituted or unsubstituted N-carbamyl. In some embodiments, X-a can be mono-substituted with an unsubstituted C₁-C₆ hydroxyalkyl (such as those described herein).

[0171] In some embodiments, Y^{1-a} can be CR^{4A-a} or N (nitrogen). In some embodiments, Y^{1-a} can be CR^{4A-a}. In some embodiments, Y^{1-a} can be N (nitrogen).

[0172] In some embodiments, Y^{2-a} can be CR^{4B-a} or N (nitrogen). In some embodiments, Y^{2-a} can be CR^{4B-a}. In some embodiments, Y^{2-a} can be N (nitrogen).

[0173] In some embodiments, Y^{1-a} and Y^{2-a} can each be N (nitrogen). In some embodiments, Y^{1-a} can be CR^{4A-a} and Y^{2-a} can be CR^{4B-a}. In some embodiments, Y^{1-a} can be CR^{4A-a} and Y^{2-a} can be N (nitrogen). In some embodiments, Y^{1-a} can be N (nitrogen) and Y^{2-a} can be CR^{4B-a}.



[0174] In some embodiments, R^{4A-a} can be hydrogen. In some embodiments, R^{4A-a} can be halogen. In some embodiments, R^{4A-a} can be an unsubstituted C₁₋₄ alkyl (such as those described herein).

[0175] In some embodiments, R^{4B-a} can be hydrogen. In some embodiments, R^{4B-a} can be halogen. In some embodiments, R^{4B-a} can be an unsubstituted C_{1-4} alkyl (such as those described herein).

[0176] In some embodiments, R^{4A-a} and R^{4B-a} can each be hydrogen. In some embodiments, R^{4A-a} and R^{4B-a} can each be halogen (wherein the halogens can be the same or different from each other). In some embodiments, R^{4A-a} and R^{4B-a} can each be an unsubstituted C_{1-4} alkyl (such as those described herein, and wherein the C_{1-4} alkyls can be the same or different from each other). In some embodiments, one of R^{4A-a} and R^{4B-a} can be hydrogen and the other of R^{4A-a} and R^{4B-a} can be halogen. In some embodiments, one of R^{4A-a} and R^{4B-a} can be hydrogen and the other of R^{4A-a} and R^{4B-a} can be an unsubstituted C_{1-4} alkyl (such as those described herein). In some embodiments, one of R^{4A-a} and R^{4B-a} can be halogen and the other of R^{4A-a} and R^{4B-a} can be an unsubstituted C_{1-4} alkyl (such as those described herein).



[0177] In some embodiments, R^{2a} can be R^{5a} . For example, R^{2a} can be



. When R^{2a} is R^{5a} , in some embodiments, R^{5-a} can be a substituted 5-7 membered monocyclic heterocyclyl. In other embodiments, R^{5-a} can be an unsubstituted 5-7 membered monocyclic heterocyclyl. Examples of R^{5-a} groups include a substituted or unsubstituted piperidinyl, a substituted or unsubstituted pyrrolidinyl and a substituted or unsubstituted azepanyl. When substituted the R^{5-a} group, possible substituents include an unsubstituted C_{1-4} alkyl, halogen, hydroxy and unsubstituted C_{1-4} haloalkyl.

[0178] In some embodiments, Ring C-a can be selected from a substituted or unsubstituted C_6-C_{10} aryl, a substituted or unsubstituted monocyclic 5-10 membered heteroaryl, a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl, a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl and a substituted or unsubstituted 7-10 membered bicyclic heterocyclyl.

[0179] In some embodiments, Ring C-a can be a substituted C_6-C_{10} aryl. In some embodiments, Ring C-a can be an unsubstituted C_6-C_{10} aryl. In some embodiments, Ring C-a can be a substituted C_6 aryl. In some embodiments, Ring C-a can be an unsubstituted C_6 aryl.

[0180] In some embodiments, Ring C-a can be a substituted 5-10 membered heteroaryl. In some embodiments, Ring C-a can be an unsubstituted 5-10 membered heteroaryl. In some embodiments, Ring C-a can be a substituted 5-6 membered heteroaryl. In some embodiments, Ring C-a can be an unsubstituted 5-6 membered heteroaryl. In some embodiments, Ring C-a can be selected from furan, thiophene, pyrrole, oxazole, thiazole, imidazole, benzimidazole, indole, pyrazole, isoxazole, pyridine, pyridazine, pyrimidine, pyrazine, purine, quinoline, isoquinoline, quinazoline and quinoxaline; wherein each of the aforementioned groups are substituted or unsubstituted, including any –NH group.


[0181] In some embodiments, Ring C-a can be a substituted or unsubstituted monocyclic 5 membered carbocyclyl. In some embodiments, Ring C-a can be a substituted or unsubstituted monocyclic 6 membered carbocyclyl. In some embodiments, Ring C-a can be a substituted or unsubstituted monocyclic 7 membered carbocyclyl.

[0182] In some embodiments, Ring C-a can be a substituted or unsubstituted 5 membered monocyclic heterocyclyl. In some embodiments, Ring C-a can be a substituted or unsubstituted 6 membered monocyclic heterocyclyl. In some embodiments, Ring C-a can be a substituted or unsubstituted 7 membered monocyclic heterocyclyl. In some embodiments, Ring C can be selected from imidazoline, imidazolidine, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, morpholine, piperidine, piperazine, pyrrolidine, pyrrolidone, 4-piperidone, pyrazoline, pyrazolidine, tetrahydropyran, azepine, oxepine and diazepine; wherein each of the aforementioned groups are substituted or unsubstituted, including any –NH group.

[0183] In some embodiments, Ring C-a can be a substituted or unsubstituted 7 membered bicyclic heterocyclyl (for example, a fused, a bridged or a spiro heterocyclyl). In some embodiments, Ring C-a can be a substituted or unsubstituted 8 membered bicyclic heterocyclyl, such as, a fused, a bridged or a spiro heterocyclyl. In some embodiments, Ring C-a can be a substituted or unsubstituted 9 membered bicyclic heterocyclyl (for example, a fused, a bridged or a spiro heterocyclyl). In some embodiments, Ring C-a can be a substituted or unsubstituted 10 membered bicyclic heterocyclyl, such as, a fused, a bridged or a spiro heterocyclyl. In some embodiments, Ring C-a can be selected from pyrrolizidine, indoline, 1,2,3,4 tetrahydroquinoline, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 2-oxa-6-azaspiro[3.3]heptane, 2,6-diazaspiro[3.3]heptane, 2-oxaspiro[3.4]octane and 2-


azaspiro[3.4]octane; wherein each of the aforementioned groups are substituted or unsubstituted, including any -NH group.


[0184] In some embodiments, Ring C-a can be substituted with one or more substituents independently selected from an unsubstituted C₁-C₆ alkyl (as described herein) and an unsubstituted (C₁-C₆ alkyl)acyl. In some embodiments, Ring C-a can be substituted with one substituent selected from an unsubstituted C₁-C₆ alkyl (as described herein) and an unsubstituted (C₁-C₆ alkyl)acyl.





[0186] A non-limiting list of WEE1 inhibitors of Compound (B) are described herein, and include those provided in Figure 1.


[0187] Examples of Compound (B) include the following:















[0188] Compound (B), along with pharmaceutically acceptable salts thereof, can be prepared as described herein and in WO 2019/173082, which is hereby incorporated by reference in its entirety. As described in WO 2019/173082, Compound (B) is a WEE1 inhibitor.

[0189] Embodiments of combinations of Compound (A), including pharmaceutically acceptable salts and salt forms thereof (such as Form A and/or Form C), and Compound (B), including pharmaceutically acceptable salts thereof, are provided in Table 1. In Table 1, “A” represents Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and the numbers represent a compound as provided in Figure 1, including pharmaceutically acceptable salts thereof. For example, in Table 1, a combination represented by 1:A corresponds



to a combination of and Compound (A), including pharmaceutically acceptable salts of any of the foregoing.

Table 1

Cmpd:Cmpd	Cmpd:Cmpd	Cmpd:Cmpd	Cmpd:Cmpd
1:A	4:A	7:A	10:A
2:A	5:A	8:A	11:A
3:A	6:A	9:A	12:A

[0190] The order of administration of compounds in a combination described herein can vary. In some embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof, can be administered prior to all of Compound (B), or a pharmaceutically acceptable salt thereof. In other embodiments, Compound (A), including pharmaceutically acceptable salts and

salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof, can be administered prior to at least one Compound (B), or a pharmaceutically acceptable salt thereof. In still other embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof, can be administered concomitantly with Compound (B), or a pharmaceutically acceptable salt thereof. In yet still other embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof, can be administered subsequent to the administration of at least one Compound (B), or a pharmaceutically acceptable salt thereof. In some embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof, can be administered subsequent to the administration of all Compound (B), or a pharmaceutically acceptable salt thereof.

[0191] There may be several advantages for using a combination of compounds described herein. For example, combining compounds that attack multiple pathways at the same time, can be more effective in treating a cancer, such as those described herein, compared to when the compounds of combination are used as monotherapy.

[0192] In some embodiments, a combination as described herein of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can decrease the number and/or severity of side effects that can be attributed to a compound described herein, such as Compound (B), or a pharmaceutically acceptable salt thereof. In other embodiments, a combination as described herein of Compound (C), including pharmaceutically acceptable salts thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can decrease the number and/or severity of side effects that can be attributed to Compound (B), or a pharmaceutically acceptable salt thereof.

[0193] Using a combination of compounds described herein can result in additive, synergistic or strongly synergistic effect. A combination of compounds described herein can result in an effect that is not antagonistic.

[0194] In some embodiments, a combination as described herein of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and one or more of

Compound (B), or pharmaceutically acceptable salts thereof, can result in an additive effect. In other embodiments, a combination as described herein of Compound (C), including pharmaceutically acceptable salts thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can result in an additive effect.

[0195] In some embodiments, a combination as described herein of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can result in a synergistic effect. In other embodiments, a combination as described herein of Compound (C), including pharmaceutically acceptable salts thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can result in a synergistic effect.

[0196] In some embodiments, a combination as described herein of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can result in a strongly synergistic effect. In other embodiments, a combination as described herein of Compound (C), including pharmaceutically acceptable salts thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, can result in a strongly synergistic effect.

[0197] In some embodiments, a combination as described herein of Compound (A), including pharmaceutically acceptable salts and salt forms, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, is not antagonistic. In other embodiments, a combination as described herein of Compound (C), including pharmaceutically acceptable salts thereof, and one or more of Compound (B), or pharmaceutically acceptable salts thereof, is not antagonistic.

[0198] As used herein, the term “antagonistic” means that the activity of the combination of compounds is less compared to the sum of the activities of the compounds in combination when the activity of each compound is determined individually (i.e., as a single compound). As used herein, the term “synergistic effect” means that the activity of the combination of compounds is greater than the sum of the individual activities of the compounds in the combination when the activity of each compound is determined individually. As used herein, the term “additive effect” means that the activity of the combination of compounds is about equal to the sum of the individual activities of the compounds in the combination when the activity of each compound is determined individually.

[0199] A potential advantage of utilizing a combination as described herein may be a reduction in the required amount(s) of the compound(s) that is effective in treating a disease condition disclosed herein compared to when each compound is administered as a monotherapy. For example, the amount of Compound (B), or a pharmaceutically acceptable salt thereof, used in a combination described herein can be less compared to the amount of Compound (B), or a pharmaceutically acceptable salt thereof, needed to achieve the same reduction in a disease marker (for example, tumor size) when administered as a monotherapy. Another potential advantage of utilizing a combination as described herein is that the use of two or more compounds having different mechanisms of action can create a higher barrier to the development of resistance compared to when a compound is administered as monotherapy. Additional advantages of utilizing a combination as described herein may include little to no cross resistance between the compounds of a combination described herein; different routes for elimination of the compounds of a combination described herein; and/or little to no overlapping toxicities between the compounds of a combination described herein.

Pharmaceutical Compositions

[0200] Compound (A), including pharmaceutically acceptable salts and salt forms thereof, can be provided in a pharmaceutical composition. Compound (B), including pharmaceutically acceptable salts thereof, can be provided in a pharmaceutical composition. Similarly, Compound (C), including pharmaceutically acceptable salts thereof, can be provided in a pharmaceutical composition.

[0201] The term “pharmaceutical composition” refers to a mixture of one or more compounds and/or salts disclosed herein with other chemical components, such as diluents, carriers and/or excipients. The pharmaceutical composition facilitates administration of the compound to an organism. Pharmaceutical compositions can also be obtained by reacting compounds with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, and salicylic acid. Pharmaceutical compositions will generally be tailored to the specific intended route of administration.

[0202] As used herein, a “carrier” refers to a compound that facilitates the incorporation of a compound into cells or tissues. For example, without limitation, dimethyl

sulfoxide (DMSO) is a commonly utilized carrier that facilitates the uptake of many organic compounds into cells or tissues of a subject.

[0203] As used herein, a “diluent” refers to an ingredient in a pharmaceutical composition that lacks appreciable pharmacological activity but may be pharmaceutically necessary or desirable. For example, a diluent may be used to increase the bulk of a potent drug whose mass is too small for manufacture and/or administration. It may also be a liquid for the dissolution of a drug to be administered by injection, ingestion or inhalation. A common form of diluent in the art is a buffered aqueous solution such as, without limitation, phosphate buffered saline that mimics the pH and isotonicity of human blood.

[0204] As used herein, an “excipient” refers to an essentially inert substance that is added to a pharmaceutical composition to provide, without limitation, bulk, consistency, stability, binding ability, lubrication, disintegrating ability etc., to the composition. For example, stabilizers such as anti-oxidants and metal-chelating agents are excipients. In an embodiment, the pharmaceutical composition comprises an anti-oxidant and/or a metal-chelating agent. A “diluent” is a type of excipient.

[0205] In some embodiments, Compounds (B), along with pharmaceutically acceptable salts thereof, can be provided in a pharmaceutical composition that includes Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and/or Compound (C), including pharmaceutically acceptable salts thereof. In other embodiments, Compound (B), along with pharmaceutically acceptable salts thereof, can be administered in a pharmaceutical composition that is separate from a pharmaceutical composition that includes Compound (A), including pharmaceutically acceptable salts and salt forms thereof. In still other embodiments, Compounds (B), along with pharmaceutically acceptable salts thereof, can be administered in a pharmaceutical composition that is separate from a pharmaceutical composition that includes Compound (C), including pharmaceutically acceptable salts thereof.

[0206] The pharmaceutical compositions described herein can be administered to a human patient *per se*, or in pharmaceutical compositions where they are mixed with other active ingredients, as in combination therapy, or carriers, diluents, excipients or combinations thereof. Proper formulation is dependent upon the route of administration chosen. Techniques for formulation and administration of the compounds described herein are known to those skilled in the art.

[0207] The pharmaceutical compositions disclosed herein may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or tableting processes. Additionally, the active ingredients are contained in an amount effective to achieve its intended purpose. Many of the compounds used in the pharmaceutical combinations disclosed herein may be provided as salts with pharmaceutically compatible counterions.

[0208] Multiple techniques of administering a compound, salt and/or composition exist in the art including, but not limited to, oral, rectal, pulmonary, topical, aerosol, injection, infusion and parenteral delivery, including intramuscular, subcutaneous, intravenous, intramedullary injections, intrathecal, direct intraventricular, intraperitoneal, intranasal and intraocular injections. In some embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, can be administered orally. In some embodiments, Compound (C), including pharmaceutically acceptable salts thereof, can be administered orally. In some embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, can be provided to a subject by the same route of administration as Compound (B), along with pharmaceutically acceptable salts thereof. In other embodiments, Compound (A), including pharmaceutically acceptable salts and salt forms thereof, can be provided to a subject by a different route of administration as Compound (B), along with pharmaceutically acceptable salts thereof. In still other embodiments, Compound (C), including pharmaceutically acceptable salts thereof, can be provided to a subject by the same route of administration as Compound (B), along with pharmaceutically acceptable salts thereof. In yet still other embodiments, Compound (C), including pharmaceutically acceptable salts thereof, can be provided to a subject by a different route of administration as Compound (B), along with pharmaceutically acceptable salts thereof.

[0209] One may also administer the compound, salt and/or composition in a local rather than systemic manner, for example, via injection or implantation of the compound directly into the affected area, often in a depot or sustained release formulation. Furthermore, one may administer the compound in a targeted drug delivery system, for example, in a liposome coated with a tissue-specific antibody. The liposomes will be targeted to and taken up selectively by the organ. For example, intranasal or pulmonary delivery to target a respiratory disease or condition may be desirable.

[0210] The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, may be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions that can include a compound and/or salt described herein formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

Uses and Methods of Treatment

[0211] As provided herein, in some embodiments, a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, can be used to treat a disease or condition. In some embodiments, a combination of compounds that includes an effective amount of Compound (C), including pharmaceutically acceptable salts thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, can be used to treat a disease or condition.

[0212] In some embodiments, the disease or condition can be selected from a breast cancer, a cervical cancer, an ovarian cancer, an uterine cancer, a vaginal cancer, a vulvar cancer, a brain cancer, a cervicocerebral cancer, an esophageal cancer, a thyroid cancer, a small cell cancer, a non-small cell cancer, a lung cancer, a stomach cancer, a gallbladder/bile duct cancer, a liver cancer, a pancreatic cancer, a colon cancer, a rectal cancer, a choriocarcinoma, an uterus body cancer, an uterocervical cancer, a renal pelvis/ureter cancer, a bladder cancer, a prostate cancer, a penis cancer, a testicular cancer, a fetal cancer, a Wilms' cancer, a skin cancer, a malignant melanoma, a neuroblastoma, an osteosarcoma, an Ewing's tumor, a soft part sarcoma, an acute leukemia, a chronic lymphatic leukemia, a chronic myelocytic leukemia, polycythemia

vera, a malignant lymphoma, multiple myeloma, a Hodgkin's lymphoma, and a non-Hodgkin's lymphoma. In other embodiments, the disease or condition can be selected from a breast cancer, a cervical cancer, an ovarian cancer, an uterine cancer, a vaginal cancer, and a vulvar cancer.

[0213] As used herein, a “subject” refers to an animal that is the object of treatment, observation or experiment. “Animal” includes cold- and warm-blooded vertebrates and invertebrates such as fish, shellfish, reptiles and, in particular, mammals. “Mammal” includes, without limitation, mice, rats, rabbits, guinea pigs, dogs, cats, sheep, goats, cows, horses, primates, such as monkeys, chimpanzees, and apes, and, in particular, humans. In some embodiments, the subject can be human. In some embodiments, the subject can be a child and/or an infant, for example, a child or infant with a fever. In other embodiments, the subject can be an adult.

[0214] As used herein, the terms “treat,” “treating,” “treatment,” “therapeutic,” and “therapy” do not necessarily mean total cure or abolition of the disease or condition. Any alleviation of any undesired signs or symptoms of the disease or condition, to any extent can be considered treatment and/or therapy. Furthermore, treatment may include acts that may worsen the subject's overall feeling of well-being or appearance.

[0215] The term “effective amount” is used to indicate an amount of an active compound, or pharmaceutical agent, that elicits the biological or medicinal response indicated. For example, an effective amount of compound, salt or composition can be the amount needed to prevent, alleviate or ameliorate symptoms of the disease or condition, or prolong the survival of the subject being treated. This response may occur in a tissue, system, animal or human and includes alleviation of the signs or symptoms of the disease or condition being treated. Determination of an effective amount is well within the capability of those skilled in the art, in view of the disclosure provided herein. The effective amount of the compounds disclosed herein required as a dose will depend on the route of administration, the type of animal, including human, being treated and the physical characteristics of the specific animal under consideration. The dose can be tailored to achieve a desired effect, but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

[0216] For example, an effective amount of a compound, or radiation, is the amount that results in: (a) the reduction, alleviation or disappearance of one or more symptoms caused by

the cancer, (b) the reduction of tumor size, (c) the elimination of the tumor, and/or (d) long-term disease stabilization (growth arrest) of the tumor.

[0217] Various types of breast cancer are known. In some embodiments, the breast cancer can be ER positive breast cancer. In some embodiments, the breast cancer can be ER positive, HER2-negative breast cancer. In some embodiments, the breast cancer can be local breast cancer (as used herein, “local” breast cancer means the cancer has not spread to other areas of the body). In other embodiments, the breast cancer can be metastatic breast cancer. A subject can have a breast cancer that has not been previously treated.

[0218] In some cases, following breast cancer treatment, a subject can relapse or have reoccurrence of breast cancer. As used herein, the terms “relapse” and “reoccurrence” are used in their normal sense as understood by those skilled in the art. Thus, the breast cancer can be recurrent breast cancer. In some embodiments, the subject has relapsed after a previous treatment for breast cancer. For example, the subject has relapsed after receiving one or more treatments with a SERM, a SERD and/or aromatase inhibitor, such as those described herein.

[0219] Within ESR1, several amino acid mutations have been identified. Mutations in ESR1 have been proposed as playing a role in resistance. There are several therapies for inhibiting estrogen receptors, including selective ER modulators (SERM), selective ER degraders (SERD) and aromatase inhibitors. One issue that can arise from the aforementioned cancer therapies is the development of resistance to the cancer therapy. Acquired resistance to cancer therapy, such as endocrine therapy, has been noted in nearly one-third of women treated with tamoxifen and other endocrine therapies. See Alluri et al., “Estrogen receptor mutations and their role in breast cancer progression” *Breast Cancer Research* (2014) 16:494. Researchers have suspected mutations in the estrogen receptor as one of the reasons for acquired resistance to cancer therapy, such as endocrine therapy. Thus, there is a need for compounds that can treat breast cancer wherein the cancer has one or more mutations within ESR1.

[0220] Some embodiments disclosed herein are relate to the use of a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, in the manufacture for a medicament for treating breast cancer in a subject in need thereof, wherein the breast cancer has at least one point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α).

Other embodiments relate herein are directed to the use of a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, for treating breast cancer in a subject in need thereof, wherein the breast cancer has at least one point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α). Still other embodiments disclosed herein are relate to a method of treating breast cancer in a subject in need thereof with a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein the breast cancer has at least one point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α).

[0221] In some embodiments, the mutation can be in the ligand binding domain (LBD) of ESR1. In some embodiments, one or more mutations can be at an amino acid selected from: A593, S576, G557, R555, L549, A546, E542, L540, D538, Y537, L536, P535, V534, V533, N532, K531, C530, H524, E523, M522, R503, L497, K481, V478, R477, E471, S463, F461, S432, G420, V418, D411, L466, S463, L453, G442, M437, M421, M396, V392, M388, E380, G344, S338, L370, S329, K303, A283, S282, E279, G274, K252, R233, P222, G160, N156, P147, G145, F97, N69, A65, A58 and S47. In some embodiments, one or more mutations can be at an amino acid selected from: D538, Y537, L536, P535, V534, S463, V392 and E380. In some embodiments, one or more mutations can be at an amino acid selected from: D538 and Y537.

[0222] In some embodiments, one or more mutations can be selected from: K303R, D538G, Y537S, E380Q, Y537C, Y537N, A283V, A546D, A546T, A58T, A593D, A65V, C530L, D411H, E279V, E471D, E471V, E523Q, E542G, F461V, F97L, G145D, G160D, G274R, G344D, G420D, G442R, G557R, H524L, K252N, K481N, K531E, L370F, L453F, L466Q, L497R, L536H, L536P, L536Q, L536R, L540Q, L549P, M388L, M396V, M421V, M437I, M522I, N156T, N532K, N69K, P147Q, P222S, P535H, R233G, R477Q, R503W, R555H, S282C, S329Y, S338G, S432L, S463P, S47T, S576L, V392I, V418E, V478L, V533M, V534E, Y537D and Y537H.

[0223] Some embodiments disclosed herein are relate to the use of a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, in the manufacture for a medicament for treating breast cancer in a subject in need thereof, wherein the breast cancer does not include at least one point mutation (for example, a point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α)). Other embodiments relate herein are directed to the use of a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, for treating breast cancer in a subject in need thereof, wherein the breast cancer does not include has at least one point mutation, such as a point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α). Still other embodiments disclosed herein are relate to a method of treating breast cancer in a subject in need thereof with a combination of compounds that includes an effective amount of Compound (A), including pharmaceutically acceptable salts and salt forms thereof , and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein the breast cancer does not include has at least one point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α) (for example, a point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α)).

[0224] As provided herein, several studies have shown that a potential cause of resistance in ER-positive breast cancer is due to acquired mutations in ESR1 due to endocrine therapy. In some embodiments, the subject had been previously treated with one or more selective ER modulators. For example, subject had been treated previously with one or more selected ER modulators selected from tamoxifen, raloxifene, ospemifene, bazedoxifene, toremifene and lasofoxifene, or a pharmaceutically acceptable salt of any of the foregoing. In some embodiments, the subject had been treated previously with one or more selective ER degraders, such as fulvestrant, (E)-3-[3,5-Difluoro-4-[(1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenyl]prop-2-enoic acid (AZD9496), (R)-6-(2-(ethyl(4-(2-(ethylamino)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (elacestrant, RAD1901), (E)-3-(4-((E)-2-(2-chloro-4-fluorophenyl)-1-

(1H-indazol-5-yl)but-1-en-1-yl)phenyl)acrylic acid (Brilanestrant, ARN-810, GDC-0810), (E)-3-(4-((2-(2-(1,1-difluoroethyl)-4-fluorophenyl)-6-hydroxybenzo[b]thiophen-3-yl)oxy)phenyl)acrylic acid (LSZ102), (E)-N,N-dimethyl-4-((2-((5-((Z)-4,4,4-trifluoro-1-(3-fluoro-1H-indazol-5-yl)-2-phenylbut-1-en-1-yl)pyridin-2-yl)oxy)ethyl)amino)but-2-enamide (H3B-6545), (E)-3-(4-((2-(4-fluoro-2,6-dimethylbenzoyl)-6-hydroxybenzo[b]thiophen-3-yl)oxy)phenyl)acrylic acid (rintodestrant, GIT48), D-0502, SHR9549, ARV-471, 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetid-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (giredestrant, GDC-9545), (S)-8-(2,4-dichlorophenyl)-9-(4-((1-(3-fluoropropyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylic acid (SAR439859), N-[1-(3-fluoropropyl)azetid-3-yl]-6-[(6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl]pyridin-3-amine (AZD9833), OP-1250 and LY3484356, or a pharmaceutically acceptable salt of any of the foregoing. In some embodiments, the subject had been treated previously with one or more aromatase inhibitors. The aromatase inhibitors can be a steroidal aromatase inhibitor or a non-steroidal aromatase inhibitor. For example, the one or more aromatase inhibitors can be selected from (exemestane (steroidal aromatase inhibitor), testolactone (steroidal aromatase inhibitor); anastazole (non-steroidal aromatase inhibitor) and letrozole (non-steroidal aromatase inhibitor), including pharmaceutically acceptable salts of any of the foregoing.

[0225] In some embodiments, the breast cancer can be present in subject, wherein the subject can be a woman. As women approach middle-age, a woman can be in a stage of menopause. In some embodiments, the subject can be a premenopausal woman. In other embodiments, the subject can be a perimenopausal woman. In still other embodiments, the subject can be a menopausal woman. In yet still other embodiments, the subject can be a postmenopausal woman. In other embodiments, the breast cancer can be present in a subject, wherein the subject can be a man. The serum estradiol level of the subject can vary. In some embodiments, the serum estradiol level (E2) of the subject can be in the range of >15 pg/mL to 350 pg/mL. In other embodiments, the serum estradiol level (E2) of the subject can be ≤ 15 pg/mL. In other embodiments, the serum estradiol level (E2) of the subject can be ≤ 10 pg/mL.

[0226] The amount of compound, salt and/or composition required for use in treatment will vary not only with the particular compound or salt selected but also with the route

of administration, the nature and/or symptoms of the disease or condition being treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician. In cases of administration of a pharmaceutically acceptable salt, dosages may be calculated as the free base. As will be understood by those of skill in the art, in certain situations it may be necessary to administer the compounds disclosed herein in amounts that exceed, or even far exceed, the dosage ranges described herein in order to effectively and aggressively treat particularly aggressive diseases or conditions.

[0227] As will be readily apparent to one skilled in the art, the useful *in vivo* dosage to be administered and the particular mode of administration will vary depending upon the age, weight, the severity of the affliction, the mammalian species treated, the particular compounds employed and the specific use for which these compounds are employed. The determination of effective dosage levels, that is the dosage levels necessary to achieve the desired result, can be accomplished by one skilled in the art using routine methods, for example, human clinical trials, *in vivo* studies and *in vitro* studies. For example, useful dosages of compounds (A), (B) and/or (C), or pharmaceutically acceptable salts of any of the foregoing, can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Such comparison can be done by comparison against an established drug, such as cisplatin and/or gemcitabine)

[0228] Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vivo* and/or *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations. Dosage intervals can also be determined using MEC value. Compositions should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

[0229] It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust administration due to toxicity or organ dysfunctions. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered

dose in the management of the disorder of interest will vary with the severity of the disease or condition to be treated and to the route of administration. The severity of the disease or condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency, will also vary according to the age, body weight and response of the individual patient. A program comparable to that discussed above may be used in veterinary medicine.

[0230] Compounds, salts and compositions disclosed herein can be evaluated for efficacy and toxicity using known methods. For example, the toxicology of a particular compound, or of a subset of the compounds, sharing certain chemical moieties, may be established by determining *in vitro* toxicity towards a cell line, such as a mammalian, and preferably human, cell line. The results of such studies are often predictive of toxicity in animals, such as mammals, or more specifically, humans. Alternatively, the toxicity of particular compounds in an animal model, such as mice, rats, rabbits, dogs or monkeys, may be determined using known methods. The efficacy of a particular compound may be established using several recognized methods, such as *in vitro* methods, animal models, or human clinical trials. When selecting a model to determine efficacy, the skilled artisan can be guided by the state of the art to choose an appropriate model, dose, route of administration and/or regime.

EXAMPLES

[0231] Additional embodiments are disclosed in further detail in the following examples, which are not in any way intended to limit the scope of the claims.

Xenograft Tumor Model

[0232] ZR-75-1-R breast cancer tumor cells (Tamoxifen resistant) were maintained *in vitro* as monolayer culture in RPMI1640 Medium supplemented with 10% fetal bovine serum and 10 μ M tamoxifen at 37°C in an atmosphere of 5% CO₂ in air. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation. BALB/c nude mice were implanted subcutaneously in the right flank with ZR-75-1-R tumor cells (1×10^7) in 100 μ l PBS:Matrigel (1:1). When tumors reached approximately 191 mm³, animals were randomly distributed into treatment groups of 10 animals each and dosed orally, daily with vehicle, freebase Compound (A) at 30 mg/kg, Compound 1 at 80 mg/kg, or compound A at 30 mg/kg in

combination with Compound 1 at 80 mg/kg for 28 days. In addition, estradiol benzoate injections were delivered by s.c. (40 µg/ 20 µL, twice weekly). Tumor volumes were evaluated twice per week to calculate tumor volume over time, and mice were weighed twice per week as a surrogate for signs of toxicity. Tumor growth inhibition (TGI) was calculated using the following equation $TGI = (1 - (T_d - T_0) / (C_d - C_0)) \times 100\%$. T_d and C_d are the mean tumor volumes of the treated and control animals, and T_0 and C_0 are the mean tumor volumes of the treated and control animals at the start of the experiment.

[0233] In Figure 2, the second from the bottom line (indicated with triangles) represents the data for Compound 1 (80 mg/kg), and the third line from the bottom (indicated with triangles) represents data for Compound (A) (30 mg/kg). As shown in Figure 2, freebase Compound (A) at 30 mg/kg and Compound 1 exhibited antitumor activity with TGI values of 43.0% and 69.9% respectively. Freebase Compound (A) at 30 mg/kg in combination with Compound 1 at 80 mg/kg, showed significant antitumor activity with a TGI of 80.8%.

[0234] Example 2: MCF-7 breast cancer tumor cells were cultured *in vitro* in DMEM Medium supplemented with 15% fetal bovine serum at 37 °C in an atmosphere of 5% CO₂ in air. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation. BALB/c nude mice were implanted subcutaneously on the 2nd right mammary fat pad with MCF-7 tumor cells (1.5×10^7) in 100µl DMEM:Matrigel (1:1). When tumors reached approximately 203 mm³, animals were randomly distributed into treatment groups of 8 animals each and dosed orally, daily with vehicle, freebase Compound (A) at 10 mg/kg, Compound 1 at 80 mg/kg, or Compound (A) at 10 mg/kg in combination with Compound 1 at 80 mg/kg for the duration noted in Figure 3. In addition, estradiol benzoate injections were delivered by s.c. (40 µg/ 20 µL, twice weekly). Tumor volumes were evaluated twice per week to calculate tumor volume over time, and mice were weighed twice per week as a surrogate for signs of toxicity. TGI values were calculated using the equation provide in Example 1.

[0235] In Figure 3, the bottom line (indicated with squares) represents the data for Compound 1 (80 mg/kg) in combination with Compound (A) (10 mg/kg), the second line from the bottom (indicated with circles) represents data for Compound 1 (80 mg/kg), the third line from the bottom (indicated with squares) represents data for Compound (A) (10 mg/kg) and the top line (indicated with circles) represents data for Vehicle. As shown in Figure 3, freebase Compound (A) at 10 mg/kg and Compound 1 exhibited antitumor activity with TGI values of


128.3%. and 132.6%, respectively. Freebase Compound (A) at 10 mg/kg in combination with Compound 1 at 80 mg/kg, showed significant antitumor activity with a TGI of 158.7%. The data provided herein demonstrates that a combination of a SERD inhibitor and a WEE1 inhibitor described herein can be used to treat a disease or condition described herein.

[0236] Furthermore, although the foregoing has been described in some detail by way of illustrations and examples for purposes of clarity and understanding, it will be understood by those of skill in the art that numerous and various modifications can be made without departing from the spirit of the present disclosure. Therefore, it should be clearly understood that the forms disclosed herein are illustrative only and are not intended to limit the scope of the present disclosure, but rather to also cover all modification and alternatives coming with the true scope and spirit of the disclosure.


WHAT IS CLAIMED IS:

1. Use of a combination of compounds for treating a disease or condition, wherein the combination includes an effective amount of Compound (A), or a pharmaceutically acceptable salt thereof, and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein:

the Compound (A) has the structure:



the one or more of Compound (B) has the structure




wherein:

R^{1a} is selected from the group consisting of hydrogen, halogen and a substituted or unsubstituted C₁-C₆ alkyl;

Ring A-a is selected from the group consisting of a substituted or unsubstituted phenyl and a substituted or unsubstituted 5-6 membered monocyclic heteroaryl;

Ring B-a is selected from the group consisting of a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl and a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl;



m-a is 0, 1, 2 or 3;

R^{3a} is selected from the group consisting of halogen and a substituted or unsubstituted C₁-C₆ alkyl;

X-a is selected from the group consisting of hydrogen, halogen, hydroxy, cyano, a substituted or unsubstituted 4-6 membered monocyclic heterocyclyl, a substituted or unsubstituted amine(C₁-C₆ alkyl), a substituted or unsubstituted -NH-(CH₂)₁₋₆-amine, a mono-substituted amine, a di-substituted amine, an amino, a substituted or unsubstituted C₁-C₆ alkyl, a substituted or unsubstituted C₁-C₆ alkoxy, a substituted or unsubstituted C₃-C₆ cycloalkoxy, a substituted or unsubstituted (C₁-C₆ alkyl)acyl, a substituted or unsubstituted C-amido, a substituted or unsubstituted N-amido, a substituted or unsubstituted C-carboxy, a substituted or unsubstituted O-carboxy, a substituted or unsubstituted O-carbamyl and a substituted or unsubstituted N-carbamyl;

Y-a is CH or N;

Y^{1-a} is CR^{4A-a} or N;

Y^{2-a} is CR^{4B-a} or N;


Ring C-a is selected from the group consisting of a substituted or unsubstituted C₆-C₁₀ aryl, a substituted or unsubstituted monocyclic 5-10 membered heteroaryl, a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl, a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl and a substituted or unsubstituted 7-10 membered bicyclic heterocyclyl;

R^{4A-a} and R^{4B-a} are independently selected from the group consisting of hydrogen, halogen and an unsubstituted C₁₋₄ alkyl; and

R^{5-a} is a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl.

2. Use of a combination of compounds for treating a disease or condition, wherein the combination includes an effective amount of Compound (C) and an effective amount of one or more of Compound (B), or a pharmaceutically acceptable salt thereof, wherein:

the Compound (C) has the structure:



wherein:

X^1 , Y^1 and Z^1 are each independently C or N;

with the first proviso that at least one of X^1 , Y^1 and Z^1 is N;

with the second proviso that each of X^1 , Y^1 and Z^1 is uncharged;

with third proviso that two of the dotted lines indicate double bonds;

with the fourth proviso that the valencies of X^1 , Y^1 and Z^1 can be each independently satisfied by attachment to a substituent selected from H and R^{12} ;

X^2 is O;

A^1 is selected from the group consisting of an optionally substituted cycloalkyl, an optionally substituted aryl, an optionally substituted heteroaryl and an optionally substituted heterocyclyl;

R^1 is selected from the group consisting of an optionally substituted C_{1-6} alkyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted cycloalkyl(C_{1-6} alkyl), an optionally substituted cycloalkenyl(C_{1-6} alkyl), an optionally substituted aryl(C_{1-6} alkyl), an optionally substituted heteroaryl(C_{1-6} alkyl) and an optionally substituted heterocyclyl(C_{1-6} alkyl);

R^2 and R^3 are each independently selected from the group consisting of hydrogen, halogen, an optionally substituted C_{1-6} alkyl and an optionally substituted C_{1-6} haloalkyl; or R^2

and R³ together with the carbon to which R² and R³ are attached form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, halogen, an optionally substituted C₁₋₆ alkyl and an optionally substituted C₁₋₆ haloalkyl; or R⁴ and R⁵ together with the carbon to which R⁴ and R⁵ are attached form an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl or an optionally substituted heterocyclyl;


R⁶, R⁷, R⁸ and R⁹ are each independently selected from the group consisting of hydrogen, halogen, hydroxy, an optionally substituted alkyl, an optionally substituted alkoxy, an optionally substituted haloalkyl, an optionally substituted mono-substituted amine, and an optionally substituted di-substituted amine;

R¹⁰ is hydrogen, halogen, an optionally substituted alkyl, or an optionally substituted cycloalkyl;

R¹¹ is hydrogen; and

R¹² is hydrogen, halogen, an optionally substituted C₁₋₃ alkyl, an optionally substituted C₁₋₃ haloalkyl or an optionally substituted C₁₋₃ alkoxy; and

the one or more of Compound (B) has the structure






wherein:

R^{1a} is selected from the group consisting of hydrogen, halogen and a substituted or unsubstituted C₁-C₆ alkyl;

Ring A-a is selected from the group consisting of a substituted or unsubstituted phenyl and a substituted or unsubstituted 5-6 membered monocyclic heteroaryl;

Ring B-a is selected from the group consisting of a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl and a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl;

R^{2a} is selected from the group consisting of ,  and  ;

$m-a$ is 0, 1, 2 or 3;

R^{3a} is selected from the group consisting of halogen and a substituted or unsubstituted C_1 - C_6 alkyl;

$X-a$ is selected from the group consisting of hydrogen, halogen, hydroxy, cyano, a substituted or unsubstituted 4-6 membered monocyclic heterocyclyl, a substituted or unsubstituted amine(C_1 - C_6 alkyl), a substituted or unsubstituted $-NH-(CH_2)_{1-6}$ -amine, a mono-substituted amine, a di-substituted amine, an amino, a substituted or unsubstituted C_1 - C_6 alkyl, a substituted or unsubstituted C_1 - C_6 alkoxy, a substituted or unsubstituted C_3 - C_6 cycloalkoxy, a substituted or unsubstituted $(C_1$ - C_6 alkyl)acyl, a substituted or unsubstituted C-amido, a substituted or unsubstituted N-amido, a substituted or unsubstituted C-carboxy, a substituted or unsubstituted O-carboxy, a substituted or unsubstituted O-carbamyl and a substituted or unsubstituted N-carbamyl;

$Y-a$ is CH or N;

Y^{1-a} is CR^{4A-a} or N;


Y^{2-a} is CR^{4B-a} or N;














Ring C-a is selected from the group consisting of a substituted or unsubstituted C_6 - C_{10} aryl, a substituted or unsubstituted monocyclic 5-10 membered heteroaryl, a substituted or unsubstituted monocyclic 5-7 membered carbocyclyl, a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl and a substituted or unsubstituted 7-10 membered bicyclic heterocyclyl;


R^{4A-a} and R^{4B-a} are independently selected from the group consisting of hydrogen, halogen and an unsubstituted C_{1-4} alkyl; and
















R^{5-a} is a substituted or unsubstituted 5-7 membered monocyclic heterocyclyl.
















3. The use of Claim 1 or 2, wherein the Compound (C) is selected from the group consisting of:























4. The use of any one of Claims 1-3, wherein the Compound (B) is selected from the group consisting of:





and





, or a pharmaceutically acceptable salt of any of the foregoing.

5. The use of Claim 1 or 2, wherein the Compound (B) is selected from the group consisting of:









6. The use of any one of Claims 1-5, wherein the disease or condition is selected from the group consisting of a breast cancer, a cervical cancer, an ovarian cancer, an uterine cancer, a vaginal cancer, a vulvar cancer, a brain cancer, a cervicocerebral cancer, an esophageal cancer, a thyroid cancer, a small cell cancer, a non-small cell cancer, a lung cancer , a stomach cancer, a gallbladder/bile duct cancer, a liver cancer, a pancreatic cancer, a colon cancer, a rectal cancer, a choriocarcinoma, an uterus body cancer, an uterocervical cancer, a renal pelvis/ureter cancer, a bladder cancer, a prostate cancer, a penis cancer, a testicular cancer, a fetal cancer, a Wilms' cancer, a skin cancer, a malignant melanoma, a neuroblastoma, an osteosarcoma, an Ewing's tumor, a soft part sarcoma, an acute leukemia, a chronic lymphatic leukemia, a chronic myelocytic leukemia, polycythemia vera, a malignant lymphoma, multiple myeloma, a Hodgkin's lymphoma, and a non-Hodgkin's lymphoma.

7. The use of any one of Claims 1-5, wherein the disease or condition is selected from the group consisting of a breast cancer, a cervical cancer, an ovarian cancer, an uterine cancer, a vaginal cancer, and a vulvar cancer.

8. The use of Claim 7, wherein the disease or condition is a breast cancer.

9. The use of any one of Claims 6-8, wherein the breast cancer that does not include any point mutations ER mutations.

10. The use of any one of Claims 6-8, wherein the disease or condition is breast cancer that has at least one point mutation within the Estrogen Receptor 1 (ESR1) that encodes Estrogen receptor alpha (ER α), wherein the mutation is selected from the group consisting of: K303R, D538G, Y537S, E380Q, Y537C, Y537N, A283V, A546D, A546T, A58T, A593D, A65V, C530L, D411H, E279V, E471D, E471V, E523Q, E542G, F461V, F97L, G145D, G160D, G274R, G344D, G420D, G442R, G557R, H524L, K252N, K481N, K531E, L370F, L453F, L466Q, L497R, L536H, L536P, L536Q, L536R, L540Q, L549P, M388L, M396V, M421V, M437I, M522I, N156T, N532K, N69K, P147Q, P222S, P535H, R233G, R477Q, R503W,

R555H, S282C, S329Y, S338G, S432L, S463P, S47T, S576L, V392I, V418E, V478L, V533M, V534E, Y537D and Y537H.

11. The use of any one of Claims 6-10, wherein the breast cancer is ER positive breast cancer.

12. The use of any one of Claims 6-10, wherein the breast cancer is ER positive/HER2-negative breast cancer.

13. The use of any one of Claims 6-12, wherein the breast cancer is local breast cancer.

14. The use of any one of Claims 6-12, wherein the breast cancer is metastatic breast cancer.

15. The use of any one of Claims 6-14, wherein the breast cancer is recurrent breast cancer.

16. The use of any one of Claims 6-15, wherein the breast cancer has been previously treated with an endocrine therapy.

17. The use of Claim 16, wherein the treatment was with a selective ER modulator (SERM).

18. The use of Claim 17, wherein the selective ER modulator is selected from the group consisting of tamoxifen, raloxifene, ospemifene, bazedoxifene, toremifene and lasofoxifene, or a pharmaceutically acceptable salt of any of the foregoing.

19. The use of Claim 16, wherein the treatment was with a selective ER degrader (SERD).

20. The use of Claim 19, wherein the selective ER degrader is selected from the group consisting of fulvestrant, (E)-3-[3,5-Difluoro-4-[(1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenyl]prop-2-enoic acid (AZD9496), (R)-6-(2-(ethyl(4-(2-(ethylamino)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (elacestrant, RAD1901), (E)-3-(4-((E)-2-(2-chloro-4-fluorophenyl)-1-(1H-indazol-5-yl)but-1-en-1-yl)phenyl)acrylic acid (Brilanestrant, ARN-810, GDC-0810), (E)-3-(4-((2-(2-(1,1-difluoroethyl)-4-fluorophenyl)-6-hydroxybenzo[b]thiophen-3-yl)oxy)phenyl)acrylic acid (LSZ102), (E)-N,N-dimethyl-4-((2-((5-((Z)-4,4,4-trifluoro-1-(3-fluoro-1H-indazol-5-yl)-2-phenylbut-1-en-1-yl)pyridin-2-yl)oxy)ethyl)amino)but-2-enamide (H3B-6545), (E)-3-(4-((2-(4-fluoro-2,6-dimethylbenzoyl)-6-hydroxybenzo[b]thiophen-3-

yl)oxy)phenyl)acrylic acid (rintodestrant, G1T48), D-0502, SHR9549, ARV-471, 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (giredestrant, GDC-9545), (S)-8-(2,4-dichlorophenyl)-9-(4-((1-(3-fluoropropyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylic acid (SAR439859), N-[1-(3-fluoropropyl)azetidin-3-yl]-6-[(6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl]pyridin-3-amine (AZD9833), OP-1250 and LY3484356, or a pharmaceutically acceptable salt of any of the foregoing.

21. The use of Claim 16, wherein the treatment was with an aromatase inhibitor.

22. The use of Claim 21, wherein the aromatase inhibitor is a steroidal aromatase inhibitor.

23. The use of Claim 22, wherein the steroidal aromatase inhibitor is selected from the group consisting of exemestane and testolactone, or a pharmaceutically acceptable salt of any of the foregoing.

24. The use of Claim 21, wherein the aromatase inhibitor is a non-steroidal aromatase inhibitor.

25. The use of Claim 24, wherein the non-steroidal aromatase inhibitor is selected from the group consisting of anastazole and letrozole, or a pharmaceutically acceptable salt of any of the foregoing.

26. The use of any one of Claims 6-14, wherein the breast cancer has not been previously treated.

27. The use of any one of Claim 6-26, wherein the breast cancer is present in a woman.

28. The use of Claim 27, wherein the subject is a premenopausal woman.

29. The use of Claim 27, wherein the subject is a perimenopausal woman.

30. The use of Claim 27, wherein the subject is a menopausal woman.

31. The use of Claim 27, wherein the breast cancer is present in a postmenopausal woman.

32. The use of any one of Claim 6-26, wherein the breast cancer is present a man.

33. The use of any one of Claim 6-32, wherein the breast cancer is present in a subject that has a serum estradiol level in the range of >15 pg/mL to 350 pg/mL.

34. The use of any one of Claim 6-32, wherein the breast cancer is present in a subject that has a serum estradiol level ≤ 15 pg/mL.

35. The use of any one of Claim 6-32, wherein the breast cancer is present in a subject that has a serum estradiol level ≤ 10 pg/mL.

Figure 1







Compound No.	Structure
1	
2	
3	
4	
5	
6	

Figure 1 (cont.)







Compound No.	Structure
7	
8	
9	
10	
11	
12	

Figure 2




Figure 3

