



(11)

EP 3 072 883 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
28.11.2018 Bulletin 2018/48

(51) Int Cl.:
C07C 403/24 ^(2006.01) **A23L 33/10** ^(2016.01)
A23K 20/00 ^(2016.01)

(21) Application number: **14848508.9**

(86) International application number:
PCT/CN2014/000862

(22) Date of filing: **23.09.2014**

(87) International publication number:
WO 2015/043112 (02.04.2015 Gazette 2015/13)

(54) **CAROTENOID COMPOUND COMING FROM PLANT AND CONTAINING NATURAL
ASTAXANTHIN, PREPARATION METHOD THEREFOR, AND COMPOSITION**

CAROTINOIDVERBINDUNG PFLANZLICHEN URSPRUNGS MIT NATÜRLICHEM ASTAXANTHIN,
HERSTELLUNGSVERFAHREN DAFÜR UND ZUSAMMENSETZUNG

COMPOSITION DE CAROTÉNOÏDES PROVENANT D'UNE PLANTE ET CONTENANT DE
L'ASTAXANTHINE NATURELLE, PROCÉDÉ DE PRÉPARATION S'Y RAPPORTANT ET
COMPOSITION

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

• **WANG, Shengnan**
Xinchang County
Zhejiang 312500 (CN)

(30) Priority: **24.09.2013 CN 201310440119**

(74) Representative: **Pallini Gervasi, Diego et al**
Notarbartolo & Gervasi GmbH
Bavariaring 21
80336 Munich (DE)

(43) Date of publication of application:
28.09.2016 Bulletin 2016/39

(73) Proprietor: **Zhejiang Medicine Co., Ltd. Xinchang
Pharmaceutical Factory**
Zhejiang 312500 (CN)

(56) References cited:
WO-A1-01/55100 WO-A1-97/23436
WO-A1-2010/138210 WO-A1-2011/145112
CN-A- 1 472 201 CN-A- 101 863 812
CN-A- 103 012 229 US-A- 5 453 565
US-A1- 2006 260 010

(72) Inventors:
• **XU, Xinde**
Xinchang County
Zhejiang 312500 (CN)
• **SHAO, Bin**
Xinchang County
Zhejiang 312500 (CN)
• **PENG, Yongjian**
Xinchang County
Zhejiang 312500 (CN)

• **KAMATA T ET AL: "Study of astaxanthin diester
extracted from Adonis aestivalis",
COMPARATIVE BIOCHEMISTRY AND
PHYSIOLOGY. B.
COMPARATIVEBIOCHEMISTRY, PERGAMON
PRESS, LONDON, GB, vol. 86, no. 3, 1 January
1987 (1987-01-01), pages 587-591, XP025753421,
ISSN: 0305-0491, DOI:
10.1016/0305-0491(87)90453-6 [retrieved on
1987-01-01]**


Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD OF THE INVENTION**

[0001] The present invention relates to a carotenoid mixture coming from plant extractive with less impurity containing natural astaxanthin with high biological activities, and its preparation method and composition, and belongs to fields of biochemistry.

BACKGROUND OF THE INVENTION

[0002] Astaxanthin's formula is $C_{40}H_{52}O_4$ and is referred to as 3,3'-dihydroxy-4,4'-diketo- β , β' -carotene widely existed in nature, and is an important substance of carotenoid, has a higher physiological activity than other carotenoid. The molecular formula is as follows:

*trans-astaxanthin*

[0003] As for a kind of natural preparation, astaxanthin has broader uses and development prospects in various industries such as pharmaceuticals, aquacultures, foods, chemistries.

[0004] Astaxanthin plays a role in the body without being transformed into vitamin A, and has various excellent biological functions such as protecting retinas and central nervous systems, preventing UV radiation, preventing cardiovascular disease and enhancing immunity of organism etc. as for a kind of non-vitamin A source carotenoid

[0005] First of all, astaxanthin is referred to as "super VE" and is substances with the most oxidizability so far, and is 1000 times as much as that of natural VE. Its molecular structure has a long conjugated double bonds, and is coupled with a α -ketone hydroxyl composed of unsaturated keto group and hydroxyl at the end of carbon chain, and has lively electronic effects, to make astaxanthin easily react with radicals and clear it, and has strong antioxidant properties. Double blind human assays have proven that astaxanthin contributes to improve antioxygenic property of erythrocyte and reduce levels of phospholipid hydroperoxide.

[0006] At the same time, it is shown from immunological researches that astaxanthin has a stronger ability of inhibiting cancer than that of β -carotene, can directly act on immune systems and induces tumor shrink. Astaxanthin has functions of improving anti-tumor immune responses and reduces incidences of various cancers of mice such as gastric cancer, colon cancer, oral cancer and bladder cancer in a certain extent, reduces carcinogenicity of aflatoxin, and prevents from skin aging and skin cancer caused by light radiation as for a light protective agent.

[0007] Furthermore, astaxanthin has high immunomodulatory activities and promotes production of human immunoglobulins. Astaxanthin can effectively prevent from occurrence and spread of diseases after combining with its antioxidant function. Feeding astaxanthin can enhance antibody responses and humoral immunity, slow down reductions of immune ability, enhance functions of T cells in vivo, increase numbers of neutrophils, and participate in organism cellular immunity. Assays shows that feeding astaxanthin can enhance antibody responses and humoral immunity, improve immunity of animals; enhance cellular immunity responses and humoral immune responses of dogs and reduce DNA damage and inflammation.




[0008] In addition, astaxanthin itself is bright red, and is the end of carotenoid synthesis in vivo, and has strong abilities of pigmentation. Astaxanthin can be directly stored and deposited in tissues without modification or biochemical conversion, and can be used as a functional pigment in various industrial fields after entering animal body. It is shown from a large number of animal assays that adding appropriate astaxanthin in baits can improve colors of skin and muscle, and enhance ornamental values and commodity values of aquatic animals. It may be found from feeding astaxanthin of euphausia superba and opossum shrimp into the red carp that the pigmentation effect on fish skin is obvious and maintained.

[0009] Furthermore, astaxanthin also can enhance survival rates of fishes and shrimps, reduce infectious diseases, promote early embryonic developments; and have functions such as keeping fresh and preventing discoloration, off-flavor and metamorphism in food industry. It also can be applied in fields of cosmetics, because of its effects such as anti-oxidation, anti-wrinkle and anti-ultraviolet radiation. Astaxanthin is remarkably the only substance to effectively prevent from diabetic kidney damages. It has been found from several studies that astaxanthin also can prevent from

diseases such as arteriosclerosis, cataracts, cardiovascular disease, and used as nutritional supplements.

[0010] Astaxanthin has multiple double bonds in its molecular structural formula, correspondingly has cis- and trans-isomers. Generally speaking, all trans-isomers astaxanthin has the highest activity relative to cis- isomers, astaxanthin has two isomers as 9- cis and 13 - cis, because of influences of spatial structure.

[0011] In addition, due to four of chiral centers existed in its molecular structural formula, astaxanthin also has optical isomers. The main optical isomer includes three forms such as (3 R, 3 R) ⁻, (3 R, 3 S, ['] meso) - and (3 S, 3 'S)-, and the molecular structure of the three isomers are shown below. Different source's astaxanthin has different optical isomers. Different optical isomer has different physiological activities. Generally speaking, the bioavailability of (3 S, 3 'S)- configuration in animal body is relatively high and is 15wt% higher than the bioavailability of (3 R, 3 R)'-configuration. The molecular structural formula of astaxanthin of salmon is (3 S, 3 'S) - configuration.



[0012] At present, astaxanthin is mainly produced in two ways such as chemical synthesis and natural extraction. Astaxanthin prepared by chemical synthesis is expensive and has significant differences in molecular structures, biological functions, application effects and biological safeties. Synthetic astaxanthin is a mixture of three different configurations wherein most of them is (3R and 3'S). Synthetic astaxanthin obviously has stabilities and antioxidant activities lower than that of natural astaxanthin, and the effects of biological absorption and pigmentation of synthetic astaxanthin are also worse than that of natural astaxanthin. And synthetic astaxanthin can not be converted into natural astaxanthin in animal bodies. On the other hand, pollutions are inevitably produced in process of synthesis, and then leads to generation of some potential safety hazard and reduction of safety performance.

[0013] Natural astaxanthin often combines with protein in vivo to show different colors such as green, brown. Astaxanthin released is bright red after protein denaturation by heating. Its molecular formula has a hydroxyl group which easily combines with carboxyl to generate stable astaxanthin esters. Therefore most of natural astaxanthin exists in the states of astaxanthin esters, such as monadic ester, dibasic ester.

[0014] There are six sources from extraction of crustaceans animals and plants such as bacteria, protozoa, fungi, microalgae in natural astaxanthin. Wherein the bacteria thalli has basically no use value because of its slow growth and low synthesis. And astaxanthin of protozoa is not synthesized by itself, and consequently is not considered.

[0015] It have been reported from domestic and international literatures that producing astaxanthin from microalgae is to culture product of haematococcus pluvialis. Natural astaxanthin obtained by the method has with predominance of (3S, 3'S) - configuration, has basically the same molecular structure of astaxanthin as that of salmon organisms, and has higher activity. However, it is difficult to establish a stable and effecient production technology system, because the Haematococcus pluvialis is sensitive to environment, and susceptible to protozoa and bacterial contamination.

[0016] Astaxanthin can be accumulated and cultured from products of some fungi such as Phaffia rhodozyma. It would

be required to add a lot of carbon source and nitrogen source, especially has high price organic nitrogen source (yeast extract and peptone, etc.) so as to improve astaxanthin yields of *Phaffia rhodozyma* and overcome disadvantages of low content. Consequently it increases production costs and is adverse to commercial production. At present, studies on industrial production of astaxanthin by uses of fungi focus on breeding of high-yield strain, exploration on fermentation process and regularity, studies on cheap fermentation substrates and its physiological function, etc.

[0017] Many factors restrict astaxanthin extraction including shellfish waste such as shrimp shell being perishable, big volume, relatively high transportation cost, low content levels of astaxanthin with large changes, binding state of astaxanthin, particle size shell etc.. No better method with industrial scales of producing astaxanthin from these waste has been found at present.


[0018] Nowadays, natural astaxanthin is mainly obtained by culture of *Haematococcus pluvialis*, fermentation of *Phaffia rhodozyma*, and extraction of aquatic products such as shrimp and crab. However, it would be difficult to realize large-scale production because of some deficiencies such as culture conditions, high costs, hard extraction conditions. It should be noted that some plants also contain astaxanthin and carotene ketone, in particular, for the genus *Ranunculaceae* plants such as *Adonis amurensis*, *Adonis vernalis*, *Adonis aestivalis*, *Adonis palaestina*, besides animals and microorganisms. Petals of *Ranunculaceae* *Adonis* genus also contain a large amount of astaxanthin. To extract natural astaxanthin by planting a large amount of *Adonis amurensis* with high-content of natural astaxanthin is a up-and-coming method.

[0019] Production of natural astaxanthin by extracting from plant tissue such as *Adonis amurensis* (also called "adonis") is a up-and-coming method and is different from production method by culture of algae, fermentation of fungus, and extraction of shellfish. On the one hand, natural astaxanthin obtained by this method has high activities, and its molecular structure of astaxanthin is basically consistent with that of astaxanthin in organisms of salmon. On the other hand, cultivation condition of *Adonis amurensis* is simple, to be easy to realize large-scale cultivation and realize industrial production.


[0020] *Adonis amurensis* (scientific name: *Adonis amurensis*), another name false hellebore or *adonis aestivalis*, is a perennial herbaceous plant of *Adonis* genus, and mainly distributes in northeast China, South Korea, Japan, Siberia. Flowering phase is a cold period of early spring from February to April. Most of *Adonis amurensis* are basically yellow flowers with 3cm to 4cm, flower diameter modified by gene may have an increased diameter. Most of *Adonis amurensis* of European varieties are red flowers, other varieties are green, white, orange and double flowers. *Adonis amurensis* contains natural astaxanthin, wherein the content of astaxanthin in the dried flower is about 2 wt %.

[0021] Carotenoid of *Adonis amurensis* contains 75-90wt% of monoester, diester or free astaxanthin, 5~10wt% of lutein and adonirubin, and less than 1wt% of 3-hydroxyechinenone, 4-hydroxyechinenone and β -carotene, etc.

[0022] Generally carotenoid is extracted from *Adonis amurensis* by organic solvents or super critical method to obtain *adonis oleoresin* (also called a "adonis extractum") containing less than 10wt% of astaxanthin, and astaxanthin in *oleoresin* also exists in the form of diester or monoester. As mentioned above, the bioavailability of astaxanthin in esterified state can be affected in animal body. It is necessary to obtain free natural astaxanthin by saponifying astaxanthin. Astaxanthin is a very unstable substance, with high antioxidant activity, and very easily oxidized to semi-astacene under the conditions of alkaline environment or oxygen or heating, thereby affect its bioactivity. The molecular structural formula of semi-astacene and astacene are shown in the figures below:





Semi-astacene ($C_{40}H_{50}O_4$)

Astacene ($C_{40}H_{48}O_4$)

[0023] Semi-astacene and astacene also belong to carotenoid, some aquatic animals such as shrimp and crab show cyan when alive and show red after cooking. It is because astaxanthin of shrimp and crab is changed to semi-astacene and astacene after heating. There are a large number of semi-astacene and astacene in human diet, and a relatively large number of semi-astacene and astacene exists in human blood without security problem. Its antioxidant activity is reduced when astaxanthin is oxidized to semi-astacene and atacene. It should try to prevent from astaxanthin to semi-astacene or atacene during processing if astaxanthin is used as a dietary supplement or a coloring agent.

[0024] So some conditions such as appropriate solvent, alkali concentration, saponification time, reaction temperature and substrate concentration must be controled in the process of hydrolyzing astaxanthin diester into free astaxanthin, to maintain a balance of reaction, and reduce generation of semiastacene and atacene in the reaction process, to keep strong antioxidant activity of astaxanthin products.

[0025] Alternately, there is other carotenoid such as adonirubin, lutein, echinenone besides astaxanthin in carotenoid of adonis amurensis. These carotenoids are found in daily dietary composition and in human blood. But these carotenoids should be removed in the process of extraction and purification of astaxanthin, in order to improve proportion of astaxanthin in the total carotenoid.

Adonirubin (Adonirubin, 3-hydroxy- β,β' -carotene-4,4'-dinone, $C_{40}H_{52}O_3$)Echinenone (Echinenone, β,β -Carotene-1-one, $C_{40}H_{54}O$)

[0026] In previous literature, it was reported that natural astaxanthin was extracted from Adonis amurensis. US2006/0260010A1 discloses plantation of new modified plant of Adonis palaestina to increase astaxanthin content of flowers, and methods of extracting astaxanthin from flowers. The application more detailedly describes variety improvement of Adonis amurensis, flower characteristics, contents of harmful material composition of cardenolide, forms of astaxanthin in flowers and compositions of other carotenoid, and wherein describes that contents of Adonis oleoresin containing astaxanthin obtained by supercritical CO_2 extraction is 5wt%~10wt% and the purity of product is not high. However it was not mentioned how to remove impurities in Adonis oleoresin or Astaxanthin crystal containing Astaxanthin.

[0027] US Patent 5453565 revealed a method of extracting astaxanthin from Adonis amurensis by cultivating Adonis amurensis genus having at least 16 petals per flower head. The product obtained by the method is in Adonis amurensis oleoresin form, and the Adonis amurensis oleoresin containing low levels of astaxanthin are purified by silica gel column chromatography to obtain 20wt% of the total carotenoid containing 80wt% astaxanthin in the form of fatty acid ester in

the product.

[0028] The Chinese patent CN101863812A also disclosed a method of preparing ointments containing astaxanthin by extracting Adonis dried flower with a mixture solvent of propane and butane and purifying the ointments by silica gel column to obtain a product with high purity. The astaxanthin of the product is also in the form of fatty acid ester.

[0029] The Chinese patent CN 1234687C described a method of extracting astaxanthin from Adonis amurensis to obtain 5wt% of the total carotenoid in the final dried product, and 80wt% of astaxanthin fatty acid ester in the total carotenoid.


[0030] The above-mentioned published applications or patents mainly relate to improvement of plant species to improve astaxanthin contents in the flowers, or processes of extracting Adonis amurensis oleoresin with lower levels of astaxanthin from Adonis amurensis, purifying Adonis amurensis oleoresin by silica gel column chromatography to finally obtain low levels of the total carotenoid with content of 5~10wt%, without research or definition for carotenoid with lower activity.

[0031] On the other hand, the bioavailability of free astaxanthin is higher than that of astaxanthin in the form of fatty acid ester in vivo, is generally higher than 15% (Johnson and the an. Crit. Rev. Biotechnol, 1991). So it would improve the bioavailability of astaxanthin if the astaxanthin fatty acid ester is saponified to free astaxanthin. On the other hand, these impurities including other carotenoid should be removed in the process of saponification of astaxanthin fatty acid and subsequent purification. Especially the appropriate reaction conditions should be selected to reduce semi-astacene and Atacene, to maintain antioxidant activities of astaxanthin.

SUMMARY OF THE INVENTION

[0032] The present invention is as defined in the claims and provides a carotenoid mixture coming from plants containing natural astaxanthin and a preparation method thereof to overcome the deficiencies of the prior art.

[0033] According to one aspect of the present invention, the present invention provides a high-content carotenoid mixture from flower, stem and fruit of adonis amurensis, the content of the total carotenoid in the high-content carotenoid mixture is higher than 95wt%, wherein the proportion of all-trans (3S, 3'S)-astaxanthin in the total carotenoid is higher than 80wt%, the all-trans (3S, 3'S) - astaxanthin has the molecular structure as shown in formula (I):



(I) .


[0034] Preferably, the high-content carotenoid comprises semi-astacene, astacene and adonirubin.

[0035] Preferably, in the high-content carotenoid mixture, the proportion of the total content of semi-astacene and astacene in the total carotenoid is less than 10wt%, and the proportion of adonirubin in the total carotenoid is less than 10 wt %.

[0036] According to another aspect of the present invention, the present invention provides a method of preparing a high content carotenoid mixture by using adonis amurensis extract as raw materials, and the method comprises the following steps:

- (a) mixing adonis amurensis extract with alcohols solvent and fully dissolving, to obtain a mixed solution;
- (b) heating the mixed solution of the step (a) to 25~75 °C, and dropwise adding an alkali solution to the reaction, and the reaction time is 0.5 ~ 6.0hr;
- (c) adding a mixture of alcohols solvent and water to obtain a reaction solution, after completion of the step (b) ;
- (d) filtering the reaction solution of the step (c) to obtain a filter cake, and then washing the filter cake with a mixture of alcohols solvent and water; and
- (e) drying the filter cake of the step (d) in vacuum to obtain a carotenoid mixture;

wherein in the high-content carotenoid mixture, the content of the total carotenoid is higher than 95 wt %, the proportion of the all-trans (3S, 3'S) astaxanthin in the carotenoid is higher than 80wt%, the all-trans (3S, 3'S) - astaxanthin has the molecular structure as shown in formula (I):



(I)。

[0037] Preferably, the alcohols solvent of the step (a) is ethanol, propylene glycol or glycerol. The alcohols solvent is beneficial to dissolution of astaxanthin fatty acid ester of the reaction substrates, the astaxanthin of the reaction product has low solubility in it, in order to facilitate the reaction and precipitate carotenoid crystals in the reaction process after completion of the reaction.

[0038] The ratio of volume of the alcohols solvent to the weight of Adonis amurensis extract in the step (a) is 5~50 times. More preferably, the ratio of volume of the alcohols solvent to the weight of Adonis amurensis extract in the step (a) is 25~40 times. If an amount of the solvent is too large, it is adverse to the reaction and prolong the reaction time and increase amounts of alkali. If an amount of the solvent is too small, a concentration of the reaction is too high. More importantly, increasing concentration of alkali in the reaction system makes more impurities generate, and then it is adverse to improve the purity of final products.

[0039] Preferably, the alkali solution in the step (b) is sodium hydroxide solution, potassium hydroxide solution or sodium methoxide solution.

[0040] The concentration of the alkali solution of the reaction mixture in the step (b) is 0.01~2.50 mol/L after completion of adding dropwise the alkali solution. More preferably, the concentration of the alkali solution of the reaction mixture in the step (b) is 1.00~2.00 mol/L after completion of adding dropwise the alkali solution. The alkali concentration of the reaction system is critical, because firstly the saponification is not complete if alkali concentration is low, and it takes a long reaction time, secondly, it is easy to produce oxidation of astaxanthin to semi-astacene and atacene, to reduce purities of astaxanthin in final products if alkali concentration is too high. Consequently, the alkali concentration of the reaction system of the present invention is 0.01~2.50mol/L, preferably, 1.00~2.00 mol/L. Some general separation measures such as using filter aids, changing the acidity-basicity and other measures in the process are used for increasing separation efficiency.

[0041] The volume ratio of alcohols solvent to water in the step (c) is 1:5~10. The volume ratio of alcohols solvent to water in the step (d) is 2~3:1. The content of the total carotenoid and proportion of astaxanthin therein in the final product may be adjusted by adjusting the proportion of alcohols solvent to water and their total amount. The more the usage of the mixture solution is, the higher the percentage of alcohols solvent is, and then the higher the total carotenoid content is in the final product, and the higher the proportion of astaxanthin is. But it affects yields of final products.

[0042] According to another aspect of the present invention, the present invention provides a composition comprising the high content carotenoid mixture for the preparation of foods, feed additives, dietary supplements, and cosmetics. Preferably, the composition further comprises of the group consist of vegetable oils, modified starch, sucrose, dextrin, and other food accessories.

[0043] Application form of astaxanthin may be in the form of oil, that is, the astaxanthin is dispersed in vegetable oil. And the application form of astaxanthin also may be in the form of powder, that is, after mixing with one or more selected from gelatin, modified starch, dextrin, sucrose and other auxiliary materials, the astaxanthin is embedded by a microcapsule emulsion technology in order to obtain astaxanthin in the form of microcapsule powder with good stability.

[0044] It may be seen from it that the preparation method of the high content carotenoid mixture by using Adonis amurensis as raw material of the present invention only relates to a saponification, and then after the saponification, washing with a mixture of water and ethanol for several times to remove impurities in amorphous state to obtain a carotenoid mixture containing astaxanthin, and further obtains a high content carotenoid mixture by vacuum drying. No additional crystallization and purification steps is used in the whole process and no silica gel column chromatography process reported by the prior art is used. Most of another carotenoid impurities is removed, the content of semi-astacene and atacene is little and the final product maintains high biological activity. So the process is suitable for industrial production.

[0045] Thus the content determined by UV spectrophotometry is higher than 95 wt %, often higher than 97 wt %, and the content of all trans (3 S, 3 S)-astaxanthin analyzed by HPLC is higher than 80 wt %, and the remainders are calendulin, semi-astacene, astacene, echinenone or hydroxyechinenone, in the carotenoid mixture obtained by the method of the present invention. These carotenoids are dietary sources and can be detected in human blood, and no security risk is present in the content. Also, no safety issues is present in the products, because ethanol is only used as solvents of the process, without uses of any other toxic and harmful organic solvent, and without other chemical impurities. Besides, the method is convenient to operate, and suitable for large-scale commercial production, because no additional purifi-

cation steps such as column separation, multiple recrystallization are used in the process besides one step reaction .

DETAILED DESCRIPTION OF THE PRESENT INVENTION AND PREFERRED EMBODIMENTS THEREOF

[0046] Hereafter, the present invention will be described specifically with reference to examples. The examples are given only for illustration of the technical solution of the present invention and should not be construed to limit the present invention.

[0047] Firstly, the detection method of the total carotenoid content in crystals of the present invention is that determining the content of the total carotenoid by ultraviolet-visible spectrophotometry (UV). 20 mg of samples to be tested are weighed and added into a 100 ml brown volumetric flask, and 10 ml of chloroform are added to the brown volumetric flask to dissolve, and then n-hexane is added to constant volume, shaken well to obtain a mixture solution. 1 ml of the mixture solution is added to a 100 ml volumetric flask, and 4 ml of chloroform is added, then hexane is added to constant volume to a scale. The absorbance value of the solution is determined at 470 nm by ultraviolet-visible absorption photometer, and n-hexane is used as a blank.

[0048] The content C of the total carotenoid is $C = A * 10000 / (2100 * W)$, wherein A is an absorbance, W is sample weight.

[0049] In addition, the condition of detecting a proportion of astaxanthin in the total carotenoid of the present invention is as follows: HPLC chromatographic column: Eclipse XDB - C18, 5 μ m, 4.6 x 150 mm; Mobile phase: methanol/water = 95/5 (V/V); Flow rate: 1 ml/min; Column temperature : 25 °C; Detection wavelength: 478 nm.

[0050] The condition of determining chiral of astaxanthin of the present invention is as follows: chromatographic column: CHIRALPAK IC (0.46 cm $\times$ 25 cm $\times$ 5 μ m); Mobile phase: ACN/MtBE = 65/35 (v/v); Flow rate: 1 ml/min; Detection wavelength: 478 nm; Column temperature: 25°C.

Example 1

[0051] 50 g (UV content is 10.5wt%) of Adonis amurensis extract (Adonis amurensis extract is obtained by extracting Adonis amurensis dried granules with a mixture of propane and butane and then is volatilized.) is added to 2500 ml of absolute ethanol, heated to 75°C under stirring, kept for 1.0hr to fully dissolve the Adonis amurensis extract. 432 ml of NaOH solution with 45% of the mass concentration (w/w) is added dropwise to form a reaction mixture solution, and the alkali concentration of the reaction mixture solution is 2.0 mol/L after completion of adding, and stirred at 75 °C for 4.0 hr. 5000 ml of deionized water and 1000 ml of ethanol are added to it, and then a dark red crystal is precipitated after slowly stirring for 1.0 hr. After filtration, a filter cake is washed with a mixture of ethanol and deionized water (the ratio of ethanol to deionized water is 2:1) for twice, 600 ml of the mixture of ethanol and deionized water per time, to obtain a crystal, the crystal is dried in vacuum, to eventually obtain 3.5 g of a dark purple carotenoid crystal.

[0052] Its maximum absorption wavelength of the carotenoid crystal by UV detection is at 470 nm, its content is 97.5 wt%, the ratio of astaxanthin analyzed by HPLC is 82.5 wt %, and, 98 wt % of the astaxanthin according to chiral HPLC analysis is (3 S, 3 'S) configuration, 2 wt % of the astaxanthin is (3S, 3'R)' configuration.

[0053] At the same time, HPLC/MS analysis shows that there are all-trans atacene (6.81 wt %), and 9-cis semi-astacene and 13-cis semi-astacene (respectively 1.65 wt% and 1.65 wt%), as well as adonirubin (7.51 wt %), hydroxyechinenone (0.85wt%) in the carotenoid crystal, besides all-trans astaxanthin. The total carotenoid composition in the crystal is shown in Table 1.

Table 1 Various of carotenoid composition in the carotenoid crystal obtained from Example 1

Carotenoid composition (wt%) in carotenoid mixture from adonis amurensis		
	all- <i>trans</i> -(3S,3'S)-astaxanthin	80.85
	all- <i>trans</i> -(3S,3'R)-astaxanthin	1.65
	Total of all- <i>trans</i> -astaxanthin	82.5
	all- <i>trans</i> -semiastacene	6.81
	9- <i>cis</i> -semiastacene	1.65
	13- <i>cis</i> -semiastacene	0.46
	Adonirubin	7.51
	Hydroxyechinenone	0.85
	other carotenoid	0.22

[0054] Also, as shown in Table 2, a relative content of each composition in carotenoid crystal of the final product is

changed by changing a proportion of solvent to amount of adding alkali, without changes of other conditions.

Table 2 Influences of solvent ratio and alkali concentration on various of carotenoid

Comparative examples	Alkali concentration of the final reaction solution (mol/L)	Proportion of alcohols solvent to raw materials	Proportion of various carotenoid composition of Crystals (wt%)			
			all- <i>trans</i> -astaxanthin	Semi-astacene	astacene	adonirubin
1	2.5	25	78.8	5.3	3.4	8.5
2	1.0	40	83.2	4.7	/	5.2
3	0.01	5	86.3	2.5	/	4.8

[0055] In comparative example 1, due to high concentration of alkali, part of semi-astacene are oxidized into astacene, and proportion of adonirubin is also increased at the same time. It would be a reason to result in increasing relative contents of other carotenoid after degradation of parts of astaxanthin. In comparative example 3, the proportion of impurities such as semi-astacene, adonirubin are lower, and the proportion of all-*trans* astaxanthin is higher, because the alkali concentration is too low, there are still a large part of raw material not to have reacted completely after 6 hr and the yield of the final product is low.

Example 2

[0056] 100 g of Adonis amurensis extract (UV content is 10.5 wt %) is added to 3000 ml of glycerol, heated to 60°C under stirring, and kept for 1.0 hr to fully dissolve the Adonis amurensis extract. 1000 ml of CH₃ONa solution with the mass concentration of 25% (w/w) is added dropwise to form a reaction mixture solution, and the alkali concentration of the reaction mixture solution is 1.16 mol/L, and stirred at 60°C for 0.5 hr. 6000 ml of deionized water and 1000 ml glycerol is added to it, and then a dark red crystal is precipitated after slowly stirring for 1.0 hr. After filtration, a filter cake is washed with a mixture of glycerol and deionized water (the ratio of glycerine and deionized water is 3:1) for twice, 500 ml of the mixture of glycerol and deionized water per time, to obtain a crystal, the crystal is dried in vacuum, to eventually obtain 4.8 g of a dark purple carotenoid crystal.

[0057] Its maximum absorption wavelength of the carotenoid crystal by UV detection is at 470 nm, its content is 98.2 wt %, the ratio of astaxanthin analyzed by HPLC is 87.1 wt % and 97 wt % of the astaxanthin according to chiral HPLC analysis is (3S, 3'S) configuration, 3 wt % of the astaxanthin is (3S, 3'R)' configuration.

[0058] At the same time, HPLC/MS analysis shows that there are all-*trans* atacene (3.81 wt %), and 9-*cis* semiastacene and 13-*cis* semiastacene (respectively 1.01wt% and 0.26wt %), as well as adonirubin (5.81 wt %), hydroxy echinenone (0.75wt%) in the carotenoid crystal, besides all-*trans* astaxanthin. The total carotenoid composition in the crystal is shown in Table 3.

Table 3 Various of carotenoid composition in the carotenoid crystals obtained from Example 2

Carotenoid composition (wt%) in carotenoid mixtures from adonis amurensis	
all- <i>trans</i> -(3S,3'S)-astaxanthin	84.49
all- <i>trans</i> -(3S,3'R)-astaxanthin	2.61
Total of all- <i>trans</i> -astaxanthin	87.1
all- <i>trans</i> -astacene	3.81
9- <i>cis</i> -semiastacene	1.01
13- <i>cis</i> -semiastacene	0.26
Adonirubin	5.81
Hydroxyechinenone	0.75
other carotenoid	1.26

Example 3

[0059] 100 kg adonis amurensis extract (UV content is 10.5 wt %) is added to 5000L of propylene glycol, and is fully dispersed under stirring at 25 °C. 123 L of KOH solution with a mass concentration of 25% (w/w) is added dropwise, and is stirred at 25°C for 2.5 hr. After sampling and analysis with HPLC, it is found that the reaction is not complete, and

the reaction is continued for 3.5 hr. After completion of the reaction monitored by HPLC, 5000L of deionized water and 500L of propylene glycol are added to it to terminate the reaction. After stirring slowly for 1.0 hr, a dark red crystal is precipitated. After filtration by pressure, a filter cake is washed with a mixture of propylene alcohol and deionized water (the ratio of propylene alcohol to water is 2:1) for twice, 600 ml of the mixture of propylene alcohol and deionized water per time, to obtain a crystal, the crystal is dried in vacuum, to eventually obtain 6.8 Kg of a dark purple carotenoid crystal.

[0060] Its maximum absorption wavelength of the carotenoid crystal by UV detection is at 470 nm, its content is 94.9 wt %, the ratio of astaxanthin analyzed by HPLC is 87.4 wt %, and the 91 wt % of astaxanthin according to chiral HPLC analysis is (3S, 3'S) configuration, the 9 wt % of astaxanthin is (3S, 3'R)' configuration.

[0061] At the same time, HPLC/MS analysis shows that there are all-trans atacene (5.23 wt %), and 9-cis semiastacene and 13 - cis semiastacene (separately 0.85 wt % and 0.23 wt %), as well as adonirubin (4.32 wt %), hydroxy echinenone (0.68wt%) in the carotenoid crystal, besides all-trans astaxanthin. The total carotenoid composition in the crystal is shown in Table 4.

Table 4 Various of carotenoid composition in the carotenoid crystals obtained from Example 3

Carotenoid composition (wt%) in carotenoid mixtures from adonis amurensis	
all- <i>trans</i> -(3S,3'S)-astaxanthin	79.53
all- <i>trans</i> -(3S,3'R)-astaxanthin	7.87
Total of all- <i>trans</i> -astaxanthin	87.4
all- <i>trans</i> -astacene	5.23
9- <i>cis</i> -semiastacene	0.85
13- <i>cis</i> -semiastacene	0.23
Adonirubin	4.32
Hydroxyechinenone	0.68
other carotenoid	1.29

[0062] It may be seen from it that a mixture solution of alcohols solvent and water is added to terminate the reaction after completion of the reaction of the present invention, and the content of the total carotenoid and the proportion of astaxanthin of the final products can be adjusted by adjusting the proportion and the total amount of alcohols solvent and water. The more the usage of the mixture solution is, the higher the percentage of the alcohols solvent is, and then the higher the total carotenoid content in the final product is, and the higher the proportion of astaxanthin is. But it affects the yield of final products.

Example 4

Preparation of oral preparation of natural astaxanthin

[0063] Carotenoid is a kind of liposoluble dietary component and is often combined with lipoproteins in human blood. So carotenoid is mixed with a small amount of vegetable fat when oral administration, or is used with food containing a certain amount of grease when oral administration. It will greatly enhance absorption and bioavailability of the carotenoid . Carotenoid is mixed with modified starch and grease, and then emulsified and dispersed to obtain nanoscale particles, and afterwards embedded by common microcapsule emulsion technology in order to obtain astaxanthin. It can not only improve the stability of carotenoid , but also improve the absorption and bioavailability in vivo.

[0064] We have developed the following two dosage forms as follows by using the carotenoid crystal containing natural astaxanthin obtained from Example 2.

(1) 55 g of the carotenoid mixture crystal containing natural astaxanthin are adequately grinded to make particles with 3 μ m of particle size , and then mixed with 200 g of olive oil under stirring to make particles uniformly disperse to obtain an oil suspension. The content of the total carotenoid of the oil suspension is 20 wt%, wherein natural astaxanthin is 175 mg/g. After filling with nitrogen and packaging closely, the change of the total carotenoid and astaxanthin in the oil suspension is monitored by UV 和 HPLC. The content of the effective ingredient of the oil suspension is stability after one month. This oil suspension containing natural astaxanthin can be administered in the form of soft capsules.


(2) 200 g of the carotenoid mixture crystal containing natural astaxanthin is mixed with modified starch and sucrose to prepare water-soluble microcapsule particles containing astaxanthin according to the method of CN101177540B. It may be found that the microcapsule particles have good stability, and the retention rate is higher than 95% in case

of airtight package and stored at 40°C for two years. It is shown from assays in vitro that the bioavailability of the microcapsule particles is increased nearly tenfold in comparison with the astaxanthin crystal. This microcapsule particles containing natural astaxanthin can be administered in the form of tablets or capsules.

[0065] It can be seen from the above Examples that the high content carotenoid crystal from *Adonis amurensis* are firstly illustrated in the present invention, wherein the high content carotenoid crystal contains a large amount of all trans (3 S, 3 S) - astaxanthin and a small amount of semi-astacene, astacene, adonirubin and trace amounts of other carotenoids. The carotenoid may be exist in dietary ingredients, and can be detected in human blood. This kind of carotenoid crystal has high purity, and can be used in multiple forms in the fields of dietary supplements of human being, food additives, feed additives and cosmetic products. The present invention also provides a method of preparing the high-content carotenoid mixture from *Adonis amurensis* with simple and reliable, suitable for industrial production.


Claims

1. A high-content carotenoid mixture prepared from flower, stem and fruit of *Adonis amurensis*, **characterized in that**, the content of the total carotenoid in the high-content carotenoid mixture is higher than 95wt%, wherein the proportion of all-trans (3S, 3'S)-astaxanthin in the total carotenoid is higher than 80wt%, the all-trans (3S, 3'S) - astaxanthin has the molecular structure as shown in formula (I):



2. The high-content carotenoid compound according to claim 1, **characterized in that**, the total carotenoid comprises semi-astacene, astacene and adonirubin.
3. The high-content of carotenoid compound according to claim 2, **characterized in that**, the proportion of the total content of semi-astacene and astacene in the total carotenoid is less than 10 wt %, the proportion of adonirubin in the total carotenoid is less than 10 wt %.
4. A method of preparing a high-content carotenoid mixture by using *Adonis amurensis* extract as raw material, **characterized in that**, comprising the following steps:
 - (a) mixing *Adonis amurensis* extract with alcohols solvent and fully dissolving, to obtain a mixed solution; wherein a ratio of the volume of the alcohols solvent to the weight of *Adonis amurensis* extract is 5~50 times;
 - (b) heating the mixed solution of the step (a) to 25~75°C, and dropwise adding an alkali solution to the reaction, and the reaction time is 0.5 - 6.0hr; wherein a concentration of the alkali solution of the reaction is 0.01~2.50 mol/L after adding dropwise the alkali solution;
 - (c) adding a mixture of alcohols solvent and water to obtain a reaction solution, after completion of the step (b) ; wherein a volume ratio of the alcohols solvent to water in the step (c) is 1:5 ~ 10;
 - (d) filtering the reaction solution of the step (c) to obtain a filter cake, and then washing the filter cake with a mixture of alcohols solvent and water; wherein a volume ratio of the alcohols solvent to water in the step (d) is 2 ~ 3:1; and
 - (e) drying the filter cake of the step (d) in vacuum to obtain a carotenoid mixture;

wherein, in the high-content carotenoid mixture, the content of the total carotenoid is higher than 95 wt %, the proportion of the all-trans (3S, 3'S) astaxanthin in the carotenoid is higher than 80wt%, the all-trans (3S, 3'S) - astaxanthin has the molecular structure as shown in formula (I):




(I) .

5. The method according to claim 4, **characterized in that**, in the step (a), the ratio of the volume of the alcohols solvent to the weight of Adonis amurensis extract is 25 ~ 40 times.
6. The method according to claim 4, **characterized in that**, in the step (b), the concentration of the alkali solution of the reaction is 1.00 ~ 2.00 mol/L after adding dropwise the alkali solution.
7. The method according to claim 5 or 6, **characterized in that**, the alcohols solvent in the step (a) is ethanol, propylene glycol or glycerol; the alkali solution in step (b) is sodium hydroxide solution, potassium hydroxide solution or sodium methoxide solution.
8. A composition comprising the high-content carotenoid mixture according to any one of claims 1 ~ 3.
9. The composition according to claim 8, **characterized in that** it further comprises one or more of vegetable oil, modified starch, sucrose, dextrin and gelatin.

Patentansprüche

1. Mischung mit hohem Carotinoid-Gehalt, hergestellt aus Blüte, Stil und Frucht von Adonis amurensis, **dadurch gekennzeichnet, dass** der Gesamtgehalt von Carotinoid in der Mischung mit hohem Carotinoid-Gehalt größer als 95 Gew.-% ist, wobei der Anteil von all-trans-(3S,3')-Astaxanthin am Gesamtcarotinoid größer als 80 Gew.-% ist, wobei das all-trans-(3S,3')-Astaxanthin die molekulare Struktur aufweist, wie in Formel (I) gezeigt:




(I) .

2. Verbindung mit hohem Carotinoid-Gehalt gemäß Anspruch 1, **dadurch gekennzeichnet, dass** das Gesamtcarotinoid Semiastacen, Astacen und Adonirubin umfasst.
3. Verbindung mit hohem Carotinoid-Gehalt gemäß Anspruch 2, **dadurch gekennzeichnet, dass** der Anteil des Gesamtgehalts von Semiastacen und Astacen am Gesamtcarotinoid weniger als 10 Gew.-% beträgt und der Anteil von Adonirubin am Gesamtcarotinoid weniger als 10 Gew.-% beträgt.
4. Verfahren zum Herstellen einer Mischung mit hohem Carotinoid-Gehalt unter Verwendung eines Extrakts von Adonis amurensis als Rohmaterial, **dadurch gekennzeichnet, dass** es die folgenden Schritte umfasst:
 - (a) Mischen des Adonis-amurensis-Extrakts mit Alkohollösungsmittel und vollständiges Auflösen, um eine gemischte Lösung zu erhalten, wobei ein Verhältnis des Volumens des Alkohollösungsmittels zum Gewicht von Adonis-amurensis-Extrakt das 5-50-Fache beträgt;
 - (b) Erhitzen der gemischten Lösung aus Schritt (a) auf 25-75 °C und tropfenweise Zugabe einer alkalischen Lösung zu der Reaktion, wobei die Reaktionsdauer 0,5-6,0 h beträgt, wobei eine Konzentration der alkalischen Lösung der Reaktion 0,01-2,50 mol/l nach der tropfenweisen Zugabe der alkalischen Lösung beträgt;

(c) nach Abschluss von Schritt (b) Zugabe einer Mischung von Alkohollösungsmittel und Wasser, um eine Reaktionslösung zu erhalten, wobei ein Volumenverhältnis des Alkohollösungsmittels zu Wasser in dem Schritt (c) 1:5-10 beträgt;

(d) nach Abschluss von Schritt (c) Filtrieren, um einen Filterkuchen zu erhalten, und dann Waschen des Filterkuchens mit einer Mischung von Alkohollösungsmittel und Wasser, um eine Reaktionslösung zu erhalten, wobei ein Volumenverhältnis des Alkohollösungsmittels zu Wasser in dem Schritt (d) 2-3:1 beträgt;

(e) Trocknen des Filterkuchens von Schritt (d) im Vakuum, um eine Carotinoid-Mischung zu erhalten, wobei der Gehalt an Gesamtcarotinoid in der Mischung mit hohem Carotinoid-Gehalt größer als 95 Gew.-% ist, wobei der Anteil von all-trans-(3S,3')-Astaxanthin am Gesamtcarotinoid größer als 80 Gew.-% ist, wobei das all-trans-(3S,3')-Astaxanthin die molekulare Struktur aufweist, wie in Formel (I) gezeigt:




(I) .

5. Verfahren gemäß Anspruch 4, **dadurch gekennzeichnet, dass** in Schritt (a) das Verhältnis des Volumens des Alkohollösungsmittels zum Gewicht von Adonis-amurensis-Extrakt das 25-40-Fache beträgt.
6. Verfahren gemäß Anspruch 4, **dadurch gekennzeichnet, dass** die Konzentration der alkalischen Lösung der Reaktion in Schritt (b) 1,00-2,00 mol/l nach tropfenweiser Zugabe der alkalischen Lösung beträgt.
7. Verfahren gemäß Anspruch 5 oder 6, **dadurch gekennzeichnet, dass** das Alkohollösungsmittel in Schritt (a) Ethanol, Propylenglykol oder Glycerin ist, und die alkalische Lösung in Schritt (b) eine Natriumhydroxidlösung, Kaliumhydroxidlösung oder Natriummethoxidlösung ist.
8. Zusammensetzung, umfassend die Mischung mit hohem Carotinoid-Gehalt gemäß einem der Ansprüche 1-3.
9. Zusammensetzung gemäß Anspruch 8, **dadurch gekennzeichnet, dass** sie des Weiteren eines oder mehrere aus Pflanzenöl, modifizierter Stärke, Saccharose, Dextrin und Gelatine umfasst.

Revendications

1. Mixture de caroténoïdes à haute teneur préparée à partir de fleurs, tiges et fruits de l'adonide de l'Amour (Adonis amurensis), **caractérisée en ce que**, la teneur des caroténoïdes totaux dans la mixture de caroténoïdes à haute teneur est supérieure à 95 % en masse, où le pourcentage de l'astaxanthine (3S, 3'S)-tout-trans dans les caroténoïdes totaux est supérieur à 80 % en masse, l'astaxanthine (3S, 3'S)-tout-trans a la structure moléculaire comme le montre la formule (I):



(I) .

2. Composé de caroténoïdes à haute teneur selon la revendication 1, **caractérisé en ce que**, les caroténoïdes totaux comprennent semi-astacène, astacène et adonirubine.

3. Composé de caroténoïdes à haute teneur selon la revendication 2, **caractérisé en ce que**, le pourcentage de la teneur totale en semi-astacène et astacène dans les caroténoïdes totaux est inférieur à 10 % en masse, le pourcentage d'adonirubine dans les caroténoïdes totaux est inférieur à 10 % en masse.

4. Méthode pour la préparation d'une mixture de caroténoïdes à haute teneur en utilisant un extrait de l'adonide de l'Amour comme matière première, **caractérisée en ce qu'elle** comprend les étapes suivantes :

(a) le mélange d'un extrait de l'adonide de l'Amour avec un solvant d'alcools et la dissolution complète, pour obtenir une solution mélangée ; où un rapport du volume du solvant d'alcools sur poids de l'extrait de l'adonide de l'Amour est de 5~50 fois ;


(b) le chauffage de la solution mélangée de l'étape (a) à 25~75°C, et l'ajout par gouttes d'une solution alcaline à la réaction, et le temps de réaction est de 0,5~6,0 h ; où une concentration de la solution alcaline de la réaction est de 0,01~2,50 mol/L après l'ajout par gouttes de la solution alcaline ;

(c) l'ajout d'une mixture d'un solvant d'alcools et d'eau pour obtenir une solution de réaction, après la réalisation de l'étape (b) ; où un rapport de volume du solvant d'alcools sur eau dans l'étape (c) est de 1:5~10 ;

(d) la filtration de la solution de réaction de l'étape (c) pour obtenir un gâteau de filtration, et ensuite le lavage du gâteau de filtration avec une mixture de solvant d'alcools et d'eau ; où un rapport de volume du solvant d'alcools sur eau dans l'étape (d) est de 2~3:1 ; et

(e) le séchage du gâteau de filtration de l'étape (d) sous vide pour obtenir une mixture de caroténoïdes ;

où dans la mixture de caroténoïdes à haute teneur, la teneur des caroténoïdes totaux est supérieure à 95 % en masse, le pourcentage de l'astaxanthine (3S, 3'S)-tout-trans dans les caroténoïdes totaux est supérieur à 80 % en masse, l'astaxanthine (3S, 3'S)-tout-trans a la structure moléculaire comme le montre la formule (I) :



(I) .

5. Méthode selon la revendication 4, **caractérisée en ce que**, dans l'étape (a), le rapport du volume du solvant d'alcools sur poids de l'extrait de l'adonide de l'Amour est de 25~40 fois.
6. Méthode selon la revendication 4, **caractérisée en ce que**, dans l'étape (b), la concentration de la solution alcaline de la réaction est de 1,00~2,00 mol/L après l'ajout par gouttes de la solution alcaline.
7. Méthode selon la revendication 5 ou 6, **caractérisée en ce que**, le solvant d'alcools dans l'étape (a) est éthanol, propylène glycol ou glycérol ; la solution alcaline dans l'étape (b) est une solution d'hydroxyde de sodium, une solution d'hydroxyde de potassium ou une solution de méthanolate de sodium.
8. Composition comprenant la mixture de caroténoïdes à haute teneur selon l'une quelconque des revendications 1~3.
9. Composition selon la revendication 8, **caractérisée en ce qu'elle** comprend en outre l'un ou plus d'une huile végétale, d'un amidon modifié, d'un sucrose, d'une dextrine ou d'une gélatine.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20060260010 A1 [0026]
- US 5453565 A [0027]
- CN 101863812 A [0028]
- CN 1234687 C [0029]
- CN 101177540 B [0064]

Non-patent literature cited in the description

- JOHNSON. *Crit. Rev. Biotechnol*, 1991 [0031]