

**(12) STANDARD PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. **AU 2005233160 B2**

(54) Title  
**CETP inhibitors**

(51) International Patent Classification(s)  
**C07C 205/00** (2006.01) **C07D 213/68** (2006.01)  
**C07C 229/00** (2006.01) **C07D 213/89** (2006.01)  
**C07C 233/18** (2006.01) **C07D 215/12** (2006.01)  
**C07C 271/16** (2006.01) **C07D 215/26** (2006.01)  
**C07C 271/20** (2006.01) **C07D 217/02** (2006.01)  
**C07C 271/22** (2006.01) **C07D 217/14** (2006.01)  
**C07C 279/08** (2006.01) **C07D 261/08** (2006.01)  
**C07C 311/04** (2006.01) **C07D 271/06** (2006.01)  
**C07C 311/08** (2006.01) **C07D 307/52** (2006.01)  
**C07C 317/32** (2006.01) **C07D 307/79** (2006.01)  
**C07D 209/08** (2006.01) **C07D 333/20** (2006.01)  
**C07D 213/40** (2006.01) **C07D 333/22** (2006.01)  
**C07D 213/64** (2006.01) **C07D 333/28** (2006.01)  
**C07D 213/65** (2006.01)

(21) Application No: **2005233160** (22) Date of Filing: **2005.04.08**

(87) WIPO No: **WO05/100298**

(30) Priority Data

(31) Number (32) Date (33) Country  
**60/561,611** **2004.04.13** **US**

(43) Publication Date: **2005.10.27**

(44) Accepted Journal Date: **2011.06.02**

(71) Applicant(s)  
**Merck Sharp & Dohme Corp.**

(72) Inventor(s)  
**Thompson, Christopher F.;Taylor, Gayle E.;Quraishi, Nazia;Deng, Qiaolin;Bohn, Joann;Sinclair, Peter J.;Ali, Amjad;Lu, Zhijian**

(74) Agent / Attorney  
**Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000**

(56) Related Art  
**WO 2004/092136 A1**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
27 October 2005 (27.10.2005)

PCT

(10) International Publication Number  
**WO 2005/100298 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 205/00**,  
229/00

**Nazia** [US/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US).

(21) International Application Number:  
PCT/US2005/012196

(74) Common Representative: **MERCK & CO., INC.**; 126  
East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 8 April 2005 (08.04.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/561,611 13 April 2004 (13.04.2004) US

(71) Applicant (for all designated States except US): **MERCK  
& CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ALI, Amjad**  
[GB/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **BOHN, Joann** [US/US]; 126 East  
Lincoln Avenue, Rahway, NJ 07065-0907 (US). **DENG,  
Qiaolin** [CN/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **LU, Zhijian** [US/US]; 126 East Lincoln  
Avenue, Rahway, NJ 07065-0907 (US). **SINCLAIR,  
Peter, J.** [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US). **TAYLOR, Gayle, E.** [US/US];  
126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).  
**THOMPSON, Christopher, F.** [US/US]; 126 East Lin-  
coln Avenue, Rahway, NJ 07065-0907 (US). **QURASHI,**

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,  
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,  
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,  
ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,  
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: CETP INHIBITORS



(57) Abstract: Compounds of Formula (I), including pharmaceutically acceptable salts of the compounds, are CETP inhibitors, and are useful for raising HDL-cholesterol, reducing LDL-cholesterol, and for treating or preventing atherosclerosis. In the compounds of Formula 1, A1 and A2 are each an aromatic ring, a 5-6-membered heterocyclic ring, an aromatic ring fused to a heterocyclic ring, a phenyl ring fused to a heterocyclic ring, or a cycloalkyl ring.

WO 2005/100298 A1

TITLE OF THE INVENTION  
CETP INHIBITORS

5 FIELD OF THE INVENTION

This invention relates to a class of chemical compounds that inhibit cholesterol ester transfer protein (CETP) and therefore may have utility in the treatment and prevention of atherosclerosis.

BACKGROUND OF THE INVENTION

10 Atherosclerosis and its clinical consequences, coronary heart disease (CHD), stroke and peripheral vascular disease, represent a truly enormous burden to the health care systems of the industrialized world. In the United States alone, approximately 13 million patients have been diagnosed with CHD, and greater than one half million deaths are attributed to CHD each year. Further, this toll is expected to grow over the next quarter century as an epidemic in obesity and diabetes continues to grow.

15 It has long been recognized that in mammals, variations in circulating lipoprotein profiles correlate with the risk of atherosclerosis and CHD. The clinical success of HMG-CoA Reductase inhibitors, especially the statins, in reducing coronary events is based on the reduction of circulating Low Density Lipoprotein cholesterol (LDL-C), levels of which correlate directly with increased risk for atherosclerosis. More recently, epidemiologic studies have demonstrated an inverse  
20 relationship between High Density Lipoprotein cholesterol (HDL-C) levels and atherosclerosis, leading to the conclusion that low serum HDL-C levels are associated with an increased risk for CHD.

Metabolic control of lipoprotein levels is a complex and dynamic process involving many factors. One important metabolic control in man is the cholesteryl ester transfer protein (CETP), a plasma glycoprotein that catalyzes the movement of cholesteryl esters from HDL to the apoB containing  
25 lipoproteins, especially VLDL (see Hesler, C.B., et. al. (1987) *Purification and characterization of human plasma cholesteryl ester transfer protein. J. Biol. Chem.* **262**(5), 2275-2282)). Under physiological conditions, the net reaction is a heteroexchange in which CETP carries triglyceride to HDL from the apoB lipoproteins and transports cholesterol ester from HDL to the apoB lipoprotein.

In humans, CETP plays a role in reverse cholesterol transport, the process whereby  
30 cholesterol is returned to the liver from peripheral tissues. Intriguingly, many animals do not possess CETP, including animals that have high HDL levels and are known to be resistant to coronary heart disease, such as rodents (see Guyard-Dangremont, V., et. al., (1998) *Phospholipid and cholesteryl ester transfer activities in plasma from 14 vertebrate species. Relation to atherogenesis susceptibility, Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **120**(3), 517-525). Numerous epidemiologic studies correlating


the effects of natural variation in CETP activity with respect to coronary heart disease risk have been performed, including studies on a small number of known human null mutations (see Hirano, K.-I., Yamashita, S. and Matsuzawa, Y. (2000) *Pros and cons of inhibiting cholesteryl ester transfer protein*, *Curr. Opin. Lipidol.* **11**(6), 589-596). These studies have clearly demonstrated an inverse correlation  
5 between plasma HDL-C concentration and CETP activity (see Inazu, A., et. al. (2000) *Cholesteryl ester transfer protein and atherosclerosis*, *Curr. Opin. Lipidol.* **11**(4), 389-396), leading to the hypothesis that pharmacologic inhibition of CETP lipid transfer activity may be beneficial to humans by increasing levels of HDL-C while lowering those of LDL.

Despite the significant therapeutic advance that statins such as simvastatin (ZOCOR®)  
10 represent, statins only achieve a risk reduction of approximately one-third in the treatment and prevention of atherosclerosis and ensuing atherosclerotic disease events. Currently, few pharmacologic therapies are available that favorably raise circulating levels of HDL-C. Certain statins and some fibrates offer modest HDL-C gains. Niacin, which provides the most effective therapy for raising HDL-C that has been clinically documented, suffers from patient compliance issues, due in part to side effects such as  
15 flushing. An agent that safely and effectively raises HDL cholesterol levels can answer a significant, but as yet unmet medical need by offering a means of pharmacologic therapy that can significantly improve circulating lipid profiles through a mechanism that is complementary to existing therapies.

New classes of chemical compounds that inhibit CETP are being investigated at several pharmaceutical companies. No CETP inhibitors are currently being marketed. One CETP inhibitor,  
20 torcetrapib, is currently in clinical trials, and is being developed for use in combination with atorvastatin. It is not currently being developed as a drug for monotherapy. New compounds are needed so that additional pharmaceutical compounds can be found that are safe and effective, either alone or in combination with other drugs that are used for treatment of lipid disorders. The compounds described herein are very potent CETP inhibitors and may be suitable for use in monotherapy and/or combination  
25 therapy.

## SUMMARY OF THE INVENTION

A first aspect of the invention provides for a compound having Formula Ib, or a pharmaceutically acceptable salt thereof, wherein



5           A<sup>1</sup> is selected from the group consisting of phenyl, thienyl, furyl, pyridyl, 1-oxidopyridinyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl, oxazolyl, isoxazolyl, and oxadiazolyl;

          wherein A<sup>1</sup> is optionally substituted with 1-4 substituent groups independently selected from -C<sub>1</sub>-C<sub>5</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub>alkyl, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -CO<sub>2</sub>H, halogen, -NR<sup>10</sup>R<sup>11</sup>,  
 10       -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)H, -C(=O)NR<sup>10</sup>R<sup>11</sup>, -SC<sub>1</sub>-C<sub>3</sub>alkyl, -C<sub>2</sub>-C<sub>3</sub>alkenyl, -NO<sub>2</sub>, -C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds, wherein -C<sub>1</sub>-C<sub>3</sub> alkyl and -C<sub>1</sub>-C<sub>5</sub> alkyl in all occurrences are optionally substituted with 1-6 substituents independently selected from 1-5 halogens and one -OH- group; and  
 15       -C<sub>3</sub>-C<sub>6</sub> cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and -C<sub>1</sub>-C<sub>3</sub>alkyl; and -C<sub>2</sub>-C<sub>3</sub> alkenyl is optionally substituted with 1-3 halogens:

          A<sup>2</sup> is selected from the group consisting of phenyl, thienyl, furyl, pyridyl, 1-oxidopyridinyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl,  
 20       dihydroindolyl, oxazolyl, isoxazolyl, oxadiazolyl, and C<sub>3</sub>-C<sub>6</sub> cycloalkyl;

wherein  $A^2$  is optionally substituted with 1-2 substituent groups independently selected from  $-C_1-C_4$  alkyl,  $-OC_1-C_3$  alkyl,  $-C(=O)C_1-C_3$ alkyl,  $-C(=O)H$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)_xC_1-C_3$  alkyl,  $-NHS(O)_2C_1-C_3$  alkyl,  $-NR^{10}R^{11}$ ,  $-NR^{10}(C=O)R^{11}$ ,  $-C_2-C_3$  alkenyl,  $-C(=O)NR^{10}R^{11}$ , halogen,  $-C_3-C_6$  cycloalkyl, and a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds, wherein  $C_1-C_3$  alkyl,  $C_1-C_4$  alkyl, and  $C_2-C_3$ alkenyl in all instances are optionally substituted with 1-3 halogens, and  $-C_3-C_6$  cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and  $-C_1-C_3$  alkyl;

10 x is an integer selected from 0, 1, and 2;

n is an integer selected from 0 and 1;

Z is selected from the group consisting of  $-S(O)_xC_1-C_6$  alkyl,  $-S(O)_2NR^{17}R^{18}$ ,  $-C(=S)OC_1-C_6$ alkyl, and  $-C(=O)X$ , wherein X is selected from the group consisting of H,  $-C_1-C_6$  alkyl,  $-OC_1-C_6$  alkyl,  $-SC_1-C_6$  alkyl, and  $-NR^{10}R^{11}$ ; wherein  $-C_1-C_6$  alkyl in all instances is optionally substituted with 1-13 halogens and 1-2 substituents independently selected from  $-OC_1-C_3$ alkyl,  $-CN$ , and  $-NO_2$ , wherein  $-OC_1-C_3$ alkyl is optionally substituted with 1-7 halogens and is optionally also substituted with 1-2  $-OC_1-C_2$  alkyl;

15  $R^1$  is selected from the group consisting of H,  $-OH$ ,  $-C_1-C_3$  alkyl, and  $-OC_1-C_3$  alkyl, wherein  $C_1-C_3$  alkyl, and  $-OC_1-C_3$  alkyl are each optionally substituted with 1-3 halogens and are also optionally substituted with one  $-OC_1-C_2$  alkyl;


20  $R^{17}$  and  $R^{18}$  are each independently selected from the group consisting of H,  $-C_1-C_5$  alkyl, and  $-C_3-C_7$  cycloalkyl, wherein  $-C_1-C_5$  alkyl, and  $-C_3-C_7$  cycloalkyl are optionally substituted with 1-13 halogens;

25  $R^2$  and  $R^3$  are each independently selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-C_1-C_3$  alkyl,  $-OC_1-C_3$ alkyl,  $-C_2-C_3$ alkenyl,  $-C_3-C_6$ cycloalkyl optionally having a double bond,  $-OC_3-C_6$ cycloalkyl optionally having a double bond,  $-C(=O)C_1-C_3$ alkyl,  $-C(=O)C_3-C_6$ cycloalkyl,  $-C(=O)H$ ,  $-CO_2H$ ,  $-CO_2C_1-C_3$ alkyl,  $-C(=O)NR^{10}R^{11}$ ,  $-CN$ ,  $-NO_2$ , and a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, and optionally 1-3 double bonds, wherein  $C_1-C_3$  alkyl and  $-C_2-C_3$  alkenyl in all instances are optionally substituted with 1-5 halogens, and  $-C_3-C_6$  cycloalkyl and the 5-6-membered heterocyclic ring are in all occurrences optionally substituted with 1-3 substituents independently selected from halogen,  $C_1-C_3$  alkyl,  $OC_1-C_3$  alkyl,  $CF_3$ , and  $-OCF_3$ ; and

3b

$R^{10}$  and  $R^{11}$  are each independently selected from H and  $-C_1-C_3$  alkyl.

A second aspect of the invention provides for a compound having formula II, or a pharmaceutically acceptable salt thereof, wherein



5  $R^1$  is selected from the group consisting of H, halogen, and  $C_1-C_3$  alkyl, wherein  $C_1-C_3$  alkyl is optionally substituted with 1-3 halogens;

$R^2$  is selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-OC_1-C_3$  alkyl,  $C_1-C_3$  alkyl,  $-CN$ ,  $-NO_2$ , and pyridyl, wherein  $C_1-C_3$  alkyl in all occurrences is optionally substituted with 1-3 halogens;

10  $R^4$  is selected from the group consisting of H, halogen,  $-C_1-C_3$  alkyl,  $-OC_1-C_2$  alkyl,  $-SC_1-C_2$  alkyl and  $-CN$ , wherein  $-C_1-C_3$  alkyl,  $-SC_1-C_3$  alkyl, and  $-OC_1-C_3$  alkyl are optionally substituted with 1-3 F;

$R^5$  and  $R^6$  are independently selected from the group consisting of H, halogen,  $-CH_3$  and  $-OCH_3$ , wherein  $-CH_3$  and  $-OCH_3$  are optionally substituted with 1-3 F;

15  $R^7$  is selected from the group consisting of H,  $-C_1-C_5$  alkyl,  $-OC_1-C_3$  alkyl,  $-C_2-C_3$  alkenyl, halogen,  $-CN$ ,  $-CO_2H$ ,  $-CO_2C_1-C_3$  alkyl,  $-SC_1-C_3$  alkyl,  $-C(=O)NR^{10}R^{11}$ ,  $-C(=O)H$ ,  $-C(=O)C_1-C_3$  alkyl,  $C_3-C_6$  cycloalkyl, and 5-(1,2,4-oxadiazolyl), wherein  $-C_1-C_3$  alkyl and  $-C_1-C_5$  alkyl in all occurrences is optionally substituted with 1-6 substituent groups independently selected from 1-5 halogens and one  $-OH$ ,  
20  $-C_2-C_3$  alkenyl is optionally substituted with 1-3 halogens, and 1,2,4-oxadiazolyl and  $C_3-C_6$  cycloalkyl are optionally substituted with 1-2 substituent groups independently selected from halogen and  $C_1-C_3$  alkyl;

$R^8$  and  $R^9$  are each independently selected from the group consisting of H,  $-C_1-C_3$  alkyl, halogen,  $-S(O)_x C_1-C_3$  alkyl,  $-NR^{10}R^{11}$ ,  $-OC_1-C_3$  alkyl,  $C_2-C_3$  alkenyl,  $-NO_2$ ,  $-CN$ ,  $-C(O)NR^{10}R^{11}$ ,  $-C(=O)H$ ,  $-NHC(=O)C_1-C_3$  alkyl,  $-NHS(O)_2 C_1-C_3$  alkyl,  $CO_2H$ ,  $CO_2C_1-C_3$  alkyl,  $C_3-C_6$  cycloalkyl, and pyridyl, wherein  $C_1-C_3$  alkyl in all occurrences is  
5 optionally substituted with 1-3 halogens,  $C_2-C_3$  alkenyl is optionally substituted with 1-3 halogens, and  $C_3-C_6$  cycloalkyl and pyridyl are optionally substituted with 1-2 substituent groups independently selected from halogen and  $C_1-C_3$  alkyl;

$R^{10}$  and  $R^{11}$  are each H or  $C_1-C_3$  alkyl;

$Z$  is selected from the group consisting of  $-C(=O)C_1-C_3$  alkyl,  
10  $-C(=O)OC_1-C_3$  alkyl,  $-S(O)_y C_1-C_3$  alkyl,  $-C(=O)H$ ,  $-C(O)NR^{10}R^{11}$ ,  $-C(=O)SC_1-C_3$  alkyl, and  $-C(=S)OC_1-C_3$  alkyl;

$n$  is an integer selected from 0 and 1;

$x$  is an integer selected from 0, 1 and 2; and

$y$  is an integer selected from 0 and 1.

15 A third aspect of the invention provides for a method of treating atherosclerosis in a patient in need of treatment comprising the administration of a therapeutically effective amount of the compound of the first or second aspect of the invention, or a pharmaceutically acceptable salt thereof to said patient.

A fourth aspect of the invention provides for a method of raising HDL-C in a  
20 patient in need of treatment comprising the administration of a therapeutically effective amount of the compound of the first or second aspect of the invention, or a pharmaceutically acceptable salt thereof to said patient.

A fifth aspect of the invention provides for a pharmaceutical composition comprising a compound of the first or second aspect of the invention, or a  
25 pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

A sixth aspect of the invention provides for the use of a compound of the first or second aspect of the invention, or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of atherosclerosis.


A seventh aspect of the invention provides for a pharmaceutical composition  
30 comprising a compound of the first or second aspect of the invention, or a pharmaceutically acceptable salt thereof, and one or more active ingredients selected from the group consisting of:

- (a) PPAR gamma agonists and partial agonists;
- (b) biguanides;



- (c) protein tyrosine phosphatase-1B (PTP-1B) inhibitors,  
 (d) dipeptidyl peptidase IV (DP-IV) inhibitors;  
 (e) insulin or insulin mimetics;  
 (f) sulfonylureas;  
 5 (g)  $\alpha$ -glucosidase inhibitors;  
 (h) one or more compounds selected from the group consisting of  
 (a) HMG-CoA reductase inhibitors; (b) bile acid sequestrants; (c) niacin, nicotinyl  
 alcohol, nicotinamide, and nicotinic acid or a salt thereof; (d) PPAR $\alpha$  agonists;  
 (e) cholesterol absorption inhibitors; (f) acyl CoA:cholesterol acyltransferase (ACAT)  
 10 inhibitors; (g) phenolic anti-oxidants, such as probucol, and (h) a microsomal triglyceride  
 transfer protein (MTP)/ApoB secretion inhibitor;  
 (i) PPAR $\alpha$ / $\gamma$ dual agonists;  
 (j) PPAR $\delta$  agonists;  
 (k) antiobesity compounds  
 15 (l) ileal bile acid transporter inhibitors;  
 (m) anti-inflammatory agents;  
 (n) glucagon receptor antagonists;  
 (o) GLP-1,  
 (p) GIP-1, and  
 20 (q) GLP-1 analogs.

Disclosed herein are compounds having Formula I, including pharmaceutically acceptable salts of the compounds, which are CETP inhibitors, having the utilities described below:



In the compounds of Formula I:

A<sup>1</sup> is selected from the group consisting of:

- (a) an aromatic ring selected from phenyl and naphthyl;
- (b) a phenyl ring fused to a 5-7 membered non-aromatic cycloalkyl  
5 ring, which optionally comprises 1-2 double bonds;
- (c) a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds and a carbonyl group or -N(O)- group, wherein the point of attachment of A<sup>1</sup> to the attached phenyl ring is a carbon atom; and
- 10 (d) a benzoheterocyclic ring comprising a phenyl ring fused to a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from O, N, and S, and optionally 1-2 double bonds, wherein the point of attachment of A<sup>1</sup> to the attached phenyl ring is a carbon atom;

A<sup>2</sup> is selected from the group consisting of:

- 15 (a) an aromatic ring selected from phenyl and naphthyl;
- (b) a phenyl ring fused to a 5-7 membered non-aromatic cycloalkyl ring, which optionally comprises 1-2 double bonds;
- (c) a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double  
20 bonds and a carbonyl group or -N(O)- group;

(d) a benzoheterocyclic ring comprising a phenyl ring fused to a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from O, N, and S, and optionally 1-2 double bonds; and

(e) a -C<sub>3</sub>-C<sub>8</sub> cycloalkyl ring optionally having 1-3 double bonds;

5                    wherein A<sup>1</sup> and A<sup>2</sup> are each optionally substituted with 1-5 substituent groups independently selected from R<sup>a</sup>;

Each R<sup>a</sup> is independently selected from the group consisting of -C<sub>1</sub>-C<sub>6</sub> alkyl, -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -C<sub>3</sub>-C<sub>8</sub> cycloalkyl optionally having 1-3 double bonds, -OC<sub>1</sub>-C<sub>6</sub>alkyl, -OC<sub>2</sub>-C<sub>6</sub> alkenyl, -OC<sub>2</sub>-C<sub>6</sub> alkynyl, -OC<sub>3</sub>-C<sub>8</sub> cycloalkyl optionally having 1-3 double bonds, -C(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, -C(=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -C(=O)H, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, -C(=O)SC<sub>1</sub>-C<sub>6</sub>alkyl, -NR<sup>10</sup>R<sup>11</sup>, -C(=O)NR<sup>10</sup>R<sup>11</sup>, -NR<sup>10</sup>C(=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, -NR<sup>10</sup>C(=O)NR<sup>10</sup>R<sup>11</sup>, -S(O)<sub>x</sub>C<sub>1</sub>-C<sub>6</sub> alkyl, -S(O)<sub>y</sub>NR<sup>10</sup>R<sup>11</sup>, -NR<sup>10</sup>S(O)<sub>y</sub>NR<sup>10</sup>R<sup>11</sup>, halogen, -CN, -NO<sub>2</sub>, and a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, said heterocyclic ring optionally also comprising a carbonyl group and optionally also comprising 1-3 double bonds,

15                    wherein for compounds in which R<sup>a</sup> is selected from the group consisting of a heterocyclic ring, -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -OC<sub>3</sub>-C<sub>8</sub> cycloalkyl, and -C(=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl, the heterocyclic ring and -C<sub>3</sub>-C<sub>8</sub> cycloalkyl groups of R<sup>a</sup> are optionally substituted with 1-5 substituent groups independently selected from halogen, -C<sub>1</sub>-C<sub>3</sub> alkyl, and -OC<sub>1</sub>-C<sub>3</sub> alkyl, wherein -C<sub>1</sub>-C<sub>3</sub> alkyl, and -OC<sub>1</sub>-C<sub>3</sub> alkyl are optionally substituted with 1-7 halogens,

20                    wherein for compounds in which R<sup>a</sup> is selected from the group consisting of -C<sub>1</sub>-C<sub>6</sub> alkyl, -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -OC<sub>1</sub>-C<sub>6</sub>alkyl, -OC<sub>2</sub>-C<sub>6</sub> alkenyl, -OC<sub>2</sub>-C<sub>6</sub> alkynyl, -C(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, -CO<sub>2</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, -C(=O)SC<sub>1</sub>-C<sub>6</sub>alkyl, -NR<sup>10</sup>C(=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, and -S(O)<sub>x</sub>C<sub>1</sub>-C<sub>6</sub> alkyl, the alkyl, alkenyl, and alkynyl groups of R<sup>a</sup> are optionally substituted with 1-15 halogens and are optionally also substituted with 1-3 substituent groups independently selected from (a) -OH, (b) -CN, (c) -NR<sup>10</sup>R<sup>11</sup>, (d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl optionally having 1-3 double bonds and optionally substituted with 1-15 halogens, (e) -OC<sub>1</sub>-C<sub>4</sub>alkyl optionally substituted with 1-9 halogens and optionally also substituted with 1-2 substituent groups independently selected from -OC<sub>1</sub>-C<sub>2</sub> alkyl, (f) -OC<sub>3</sub>-C<sub>8</sub> cycloalkyl optionally having 1-3 double bonds and optionally substituted with 1-15 halogens, (g) -CO<sub>2</sub>H, (h) -C(=O)CH<sub>3</sub>, and (i) -CO<sub>2</sub>C<sub>1</sub>-C<sub>4</sub>alkyl which is optionally substituted with 1-9 halogens;

30

n is an integer selected from 0 and 1;

p is an integer from 0-4;

x is an integer selected from 0, 1, and 2;

y is an integer selected from 1 and 2;

Z is selected from the group consisting of  $-S(O)_x C_1-C_6$  alkyl,  $-S(O)_2 NR^{17}R^{18}$ ,  $-C(=S)OC_1-C_6$  alkyl, and  $-C(=O)X$ , wherein X is selected from the group consisting of H,  $-C_1-C_6$  alkyl,  $-OC_1-C_6$  alkyl,  $-SC_1-C_6$  alkyl, and  $-NR^{10}R^{11}$ ; wherein  $-C_1-C_6$  alkyl in all instances is optionally substituted with 1-13 halogens and 1-2 substituents independently selected from  $-OC_1-C_3$  alkyl,  $-CN$ , and  $-NO_2$ , wherein  $-OC_1-C_3$  alkyl is optionally substituted with 1-7 halogens and is optionally also substituted with 1-2  $-OC_1-C_2$  alkyl;


$R^1$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ , and  $R^{16}$  are each independently selected from the group consisting of H,  $-OH$ , halogen,  $-C_1-C_4$  alkyl,  $-C_3-C_6$  cycloalkyl,  $-OC_1-C_4$  alkyl, and  $-NR^{10}R^{11}$ , wherein  $-C_1-C_4$  alkyl,  $-C_3-C_6$  cycloalkyl, and  $-OC_1-C_4$  alkyl are each optionally substituted with 1-9 halogens and are each optionally also substituted with 1-2 groups independently selected from  $-OH$ ,  $-C(=O)CH_3$ ,  $-OC(=O)CH_3$ ,  $-OC_1-C_2$  alkyl, and  $-OC_1-C_2$  alkylene( $OC_1-C_2$  alkyl);

$R^{10}$  and  $R^{11}$  are each independently selected from H,  $-C_1-C_5$  alkyl,  $-C(=O)C_1-C_5$  alkyl and  $-S(O)_y C_1-C_5$  alkyl, wherein  $-C_1-C_5$  alkyl in all instances is optionally substituted with 1-11 halogens; and

$R^{17}$  and  $R^{18}$  are each independently selected from the group consisting of H,  $-C_1-C_5$  alkyl, and  $-C_3-C_7$  cycloalkyl, wherein  $-C_1-C_5$  alkyl, and  $-C_3-C_7$  cycloalkyl are optionally substituted with 1-13 halogens.

#### DETAILED DESCRIPTION OF THE INVENTION

A subgroup of the compounds of this invention has the structure of Formula Ia, written below, or a pharmaceutically acceptable salt thereof:



In the compounds of Formula Ia:

A<sup>1</sup> is selected from the group consisting of:

- (a) an aromatic ring selected from phenyl and naphthyl;
  - (b) a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds and a carbonyl group or -N(O)- group, wherein the point of attachment of A<sup>1</sup> to the attached phenyl ring is a carbon atom; and
  - (c) a bicyclic ring comprising a phenyl ring fused to a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from O, N, and S, and optionally 1-2 double bonds, wherein the point of attachment of A<sup>1</sup> to the attached phenyl ring is a carbon atom;
- wherein A<sup>1</sup> is optionally substituted with 1-4 substituent groups independently selected from -C<sub>1</sub>-C<sub>5</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub>alkyl, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -CO<sub>2</sub>H, halogen, -NR<sup>10</sup>R<sup>11</sup>, -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)H, -C(=O)NR<sup>10</sup>R<sup>11</sup>, -SC<sub>1</sub>-C<sub>3</sub> alkyl, -C<sub>2</sub>-C<sub>3</sub> alkenyl, -CN, -NO<sub>2</sub>, -C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds, wherein -C<sub>1</sub>-C<sub>3</sub> alkyl and -C<sub>1</sub>-C<sub>5</sub> alkyl in all occurrences are optionally substituted with 1-6 substituents independently selected from 1-5 halogens and one -OH group; and -C<sub>3</sub>-C<sub>6</sub> cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and -C<sub>1</sub>-C<sub>3</sub> alkyl; and -C<sub>2</sub>-C<sub>3</sub> alkenyl is optionally substituted with 1-3 halogens;

20

A<sup>2</sup> is selected from the group consisting of phenyl, naphthyl, -C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and a heterocyclic 5-6 membered ring having 1-3 heteroatoms independently selected from O, N, and S, and optionally also comprising 1-3 double bonds and a carbonyl group or -N(O)- group, wherein A<sup>2</sup> is

optionally substituted with 1-2 substituent groups independently selected from -C<sub>1</sub>-C<sub>4</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)H, -NO<sub>2</sub>, -CN, -S(O)<sub>x</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -NHS(O)<sub>2</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -NR<sup>10</sup>R<sup>11</sup>, -NR<sup>10</sup>C(=O)R<sup>11</sup>, -C<sub>2</sub>-C<sub>3</sub> alkenyl, -C(=O)NR<sup>10</sup>R<sup>11</sup>, halogen, -C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds, wherein C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl, and C<sub>2</sub>-C<sub>3</sub>alkenyl in all instances are optionally substituted with 1-3 halogens, and -C<sub>3</sub>-C<sub>6</sub> cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and -C<sub>1</sub>-C<sub>3</sub> alkyl;

x is an integer selected from 0, 1, and 2;

n is an integer selected from 0 and 1;

R<sup>1</sup> is selected from the group consisting of H, OH, C<sub>1</sub>-C<sub>3</sub> alkyl, and -OC<sub>1</sub>-C<sub>3</sub> alkyl, wherein C<sub>1</sub>-C<sub>3</sub> alkyl and -OC<sub>1</sub>-C<sub>3</sub> alkyl are each optionally substituted with 1-3 halogens and also are optionally substituted with one -OC<sub>1</sub>-C<sub>2</sub>alkyl;


R<sup>2</sup> and R<sup>3</sup> are each independently selected from the group consisting of H, halogen, -NR<sup>10</sup>R<sup>11</sup>, -C<sub>1</sub>-C<sub>3</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -C<sub>2</sub>-C<sub>3</sub> alkenyl, -C<sub>3</sub>-C<sub>6</sub> cycloalkyl optionally having a double bond, -OC<sub>3</sub>-C<sub>6</sub> cycloalkyl optionally having a double bond, -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)C<sub>3</sub>-C<sub>6</sub> cycloalkyl, -C(=O)H, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)NR<sup>10</sup>R<sup>11</sup>, -CN, -NO<sub>2</sub>, and a 5-6-membered heterocyclic ring having 1-4 heteroatoms independently selected from N, S, and O, and optionally 1-3 double bond, wherein C<sub>1</sub>-C<sub>3</sub> alkyl and -C<sub>2</sub>-C<sub>3</sub> alkenyl in all instances are optionally substituted with 1-5 halogens, and -C<sub>3</sub>-C<sub>6</sub> cycloalkyl and the 5-6-membered heterocyclic ring are in all occurrences optionally substituted with 1-3 substituents independently selected from halogen, -C<sub>1</sub>-C<sub>3</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -CF<sub>3</sub>, and -OCF<sub>3</sub>; and

R<sup>10</sup> and R<sup>11</sup> are each independently selected from H and -C<sub>1</sub>-C<sub>3</sub> alkyl.

In a subgroup of Formula Ia,

R<sup>2</sup> and R<sup>3</sup> are each independently selected from the group consisting of H, halogen, -NR<sup>10</sup>R<sup>11</sup>, -C<sub>1</sub>-C<sub>3</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -CN, -NO<sub>2</sub>, and pyridyl, wherein C<sub>1</sub>-C<sub>3</sub> alkyl in all instances is optionally substituted with 1-3 halogens.

A subgroup of Formula Ia has Formula Ib:



where the definitions of the groups are as defined previously.

In subgroups of compounds having Formula Ia and Formula Ib, R<sup>2</sup> is selected from H,  
 5 halogen, -NR<sup>10</sup>R<sup>11</sup>, -C<sub>1</sub>-C<sub>3</sub> alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -CN, -NO<sub>2</sub>, and pyridyl, wherein C<sub>1</sub>-C<sub>3</sub> alkyl in all  
 instances is optionally substituted with 1-3 halogens; and

R<sup>3</sup> is selected from H, halogen, -CH<sub>3</sub>, -CF<sub>3</sub>, -OCH<sub>3</sub>, and -OCF<sub>3</sub>.

10 In subgroups of Formula I, Ia, and Ib, R<sup>1</sup> is H, halogen, or C<sub>1</sub>-C<sub>3</sub> alkyl, wherein C<sub>1</sub>-C<sub>3</sub>  
 alkyl is optionally substituted with 1-3 halogens.

In subgroups of Formula I, Ia, and Ib, A<sup>1</sup> may be phenyl, thienyl, furyl, pyridyl,  
 quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl, oxazolyl, isoxazolyl,  
 15 or oxadiazolyl.

In subgroups of Formula I, Ia, and Ib, A<sup>2</sup> may be phenyl, thienyl, furyl, pyridyl, 1-  
 oxidopyridinyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl,  
 oxazolyl, isoxazolyl, oxadiazolyl, and C<sub>3</sub>-C<sub>6</sub> cycloalkyl.

20 In subgroups of Formula I, Ia, and Ib, A<sup>2</sup> may be phenyl, pyridyl, thienyl, 1-  
 oxidopyridinyl, or cyclohexyl.


In subgroups of Formula I, Ia and Ib, A<sup>1</sup> may be one of the following: 2-thienyl, 3-  
 25 thienyl, 2-furyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 3-quinolyl, 5-quinolyl, 6-quinolyl, 8-quinolyl, 4-

isoquinolyl, 5-isoquinolyl, 4-isoxazolyl, 5-1,2,4-oxadiazolyl, 5-(2,3-dihydrobenzofuranyl), or 5-indolyl, where A<sup>1</sup> is optionally substituted with 1-2 substituent groups independently selected from -CH<sub>3</sub>, -OCH<sub>3</sub>, acetyl, and halogen.

5 In subgroups of Formula I, Ia and Ib, A<sup>2</sup> may be one of the following: 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-thienyl, 3-thienyl, 1-oxidopyridin-3-yl, and cyclohexyl, where A<sup>2</sup> is optionally substituted with 1-2 substituents which are independently selected from -CH<sub>3</sub>, -OCH<sub>3</sub>, acetyl, and halogen.

10 In subgroups of Formula I, Ia, and Ib, Z is selected from the group consisting of -C(=O)C<sub>1</sub>-C<sub>3</sub> alkyl, -C(=O)OC<sub>1</sub>-C<sub>3</sub> alkyl, -S(O)<sub>y</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -C(=O)H, -C(O)NR<sup>10</sup>R<sup>11</sup>, -C(=O)SC<sub>1</sub>-C<sub>3</sub> alkyl, and -C(=S)OC<sub>1</sub>-C<sub>3</sub> alkyl.

A subgroup of the compounds of this invention has Formula II:



15 In the compounds of Formula II, R<sup>2</sup> is selected from the group consisting of H, halogen, -NR<sup>10</sup>R<sup>11</sup>, -OC<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkyl, -CN, -NO<sub>2</sub>, and 3-pyridyl, wherein C<sub>1</sub>-C<sub>3</sub> alkyl in all occurrences is optionally substituted with 1-3 halogens;

20



R<sup>4</sup> is selected from the group consisting of H, halogen, -C<sub>1</sub>-C<sub>3</sub> alkyl, -OC<sub>1</sub>-C<sub>2</sub> alkyl, -SC<sub>1</sub>-C<sub>2</sub> alkyl and -CN, wherein -C<sub>1</sub>-C<sub>3</sub> alkyl, -SC<sub>1</sub>-C<sub>2</sub> alkyl, and -OC<sub>1</sub>-C<sub>2</sub> alkyl are optionally substituted with 1-3 F;

5 R<sup>5</sup> and R<sup>6</sup> are each independently H, halogen, -CH<sub>3</sub> or -OCH<sub>3</sub>, wherein -CH<sub>3</sub> and -OCH<sub>3</sub> are optionally substituted with 1-3 F;

R<sup>7</sup> is selected from the group consisting of H, -C<sub>1</sub>-C<sub>5</sub>alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -C<sub>2</sub>-C<sub>3</sub> alkenyl, halogen, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -SC<sub>1</sub>-C<sub>3</sub> alkyl, -C(=O)NR<sup>10</sup>R<sup>11</sup>, -C(=O)H,  
10 -C(=O)C<sub>1</sub>-C<sub>3</sub> alkyl, and 5-(1,2,4-oxadiazolyl), wherein -C<sub>1</sub>-C<sub>3</sub> alkyl and -C<sub>1</sub>-C<sub>5</sub> alkyl in all occurrences is optionally substituted with 1-6 substituent groups independently selected from 1-5 halogens and one -OH, and -C<sub>2</sub>-C<sub>3</sub> alkenyl is optionally substituted with 1-3 halogens;

R<sup>8</sup> and R<sup>9</sup> are each independently selected from the group consisting of H, -C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -S(O)<sub>x</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -NR<sup>10</sup>R<sup>11</sup>, -OC<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>2</sub>-C<sub>3</sub> alkenyl, -NO<sub>2</sub>, -CN, -C(O)NR<sup>10</sup>R<sup>11</sup>,  
15 -NHC(=O)C<sub>1</sub>-C<sub>3</sub> alkyl, -NHS(O)<sub>2</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1</sub>-<sub>3</sub>alkyl, and pyridyl, wherein C<sub>1</sub>-C<sub>3</sub> alkyl in all occurrences is optionally substituted with 1-3 halogens, and C<sub>2</sub>-C<sub>3</sub> alkenyl is optionally substituted with 1-3 halogens; and

20 R<sup>10</sup> and R<sup>11</sup> are each H or C<sub>1</sub>-C<sub>3</sub> alkyl.

In subgroups of Formula Ia, Ib, and II, R<sup>1</sup> is H or CH<sub>3</sub>.

In subgroups of Formula Ia, Ib, and II, R<sup>2</sup> is H, -CF<sub>3</sub>, -OCH<sub>3</sub>, -NO<sub>2</sub>, -CN, halogen, -NH<sub>2</sub>  
25 or 3-pyridyl.

In subgroups of Formula Ia, Ib, and II, R<sub>2</sub> is a group other than H.

In subgroups of Formula II, R<sup>4</sup> is selected from H, halogen, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -OCH<sub>3</sub>,  
30 -OC<sub>2</sub>H<sub>5</sub>, -SCH<sub>3</sub>, and -CN.

In subgroups of Formula II, R<sup>5</sup> is H or F.

In subgroups of Formula II, R<sup>6</sup> is H, F, -CH<sub>3</sub>, or -OCH<sub>3</sub>.

In subgroups of Formula II, R<sup>7</sup> is selected from H, C<sub>1</sub>-C<sub>4</sub>alkyl, -C(=O)H, -C(=O)CH<sub>3</sub>, -CH=CH<sub>2</sub>, -CN, Cl, F, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -OCH<sub>3</sub>, -SCH<sub>3</sub>, -C(=O)NR<sup>10</sup>R<sup>11</sup>, C<sub>3</sub>-C<sub>6</sub>cycloalkyl and 3-methyl-5-(1,2,4-oxadiazolyl), wherein C<sub>1</sub>-C<sub>4</sub>alkyl and C<sub>3</sub>-C<sub>6</sub>cycloalkyl are optionally substituted with 1-6 substituents which are independently selected from 1-5 F and one -OH.

In subgroups of Formula II, R<sup>8</sup> and R<sup>9</sup> each are independently selected from H, C<sub>1</sub>-C<sub>2</sub>alkyl, which is optionally substituted with 1-3 F; halogen; -CN; -NO<sub>2</sub>; -S(O)<sub>x</sub>CH<sub>3</sub>, which is optionally substituted with 1-3F; -OCH<sub>3</sub>, which is optionally substituted with 1-3 F; -CH=CH<sub>2</sub>; -C(=O)H; -C(=O)NR<sup>10</sup>R<sup>11</sup>; -CO<sub>2</sub>H; -NR<sup>10</sup>R<sup>11</sup>; -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl; -NHC(=O)CH<sub>3</sub>; -NHS(O)<sub>2</sub>CH<sub>3</sub>; and 4-pyridyl, wherein x is 0, 1, or 2.

In subgroups of Formula I, Ia, Ib, and II, R<sup>10</sup> and R<sup>11</sup> are each H or -CH<sub>3</sub>.

In subgroups of Formula I, Ia, Ib, and II, n is 0. In subgroups of Formula I, Ia, Ib, and II, n is 1.

In subgroups of Formula II, R<sup>4</sup> is -OCH<sub>3</sub>.

In subgroups of Formula II, R<sup>7</sup> is -CH(CH<sub>3</sub>)<sub>2</sub>; -CH(OH)CH<sub>3</sub>; -CH(OH)CF<sub>3</sub>; -CH(CH<sub>3</sub>)CF<sub>3</sub>; -C(OH)(CH<sub>3</sub>)CF<sub>3</sub>; -CH<sub>3</sub>, which is optionally substituted with -OH or 1-3 F; -C<sub>2</sub>H<sub>5</sub>, which is optionally substituted with 1-5 F; n-C<sub>3</sub>H<sub>7</sub>; tert-C<sub>4</sub>H<sub>9</sub>; -C(=O)H, -C(=O)CH<sub>3</sub>, -CH=CH<sub>2</sub>, -CN, Cl, F, -CO<sub>2</sub>H, -CO<sub>2</sub>CH<sub>3</sub>; -OCH<sub>3</sub>; -SCH<sub>3</sub>; -C(=O)NR<sup>10</sup>R<sup>11</sup>; 3-methyl-5-(1,2,4-oxadiazolyl), or C<sub>3</sub>-C<sub>6</sub>cycloalkyl.

In subgroups of Formula II, R<sup>5</sup> and R<sup>6</sup> are H.

In subgroups of Formula II, R<sup>8</sup> and R<sup>9</sup> are each independently selected from H, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -NO<sub>2</sub>, -OCH<sub>3</sub>, -CN, -NO<sub>2</sub>, halogen, S(O)<sub>x</sub>CH<sub>3</sub>, -C(=O)NR<sup>10</sup>R<sup>11</sup>, and -CH=CH<sub>2</sub>, where -CH<sub>3</sub> and -OCH<sub>3</sub> are each independently substituted with 1-3 F.

In subgroups of Formula II, R<sup>8</sup> is -CF<sub>3</sub>, -NO<sub>2</sub>, Cl, -CH<sub>3</sub>, -OCF<sub>3</sub>, or -OCF<sub>2</sub>H; and R<sup>9</sup> is H, -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, halogen, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -OCH<sub>3</sub>, -S(O)<sub>x</sub>CH<sub>3</sub>, -CF<sub>2</sub>H, -C(=O)NR<sup>10</sup>R<sup>11</sup>, or -CH=CH<sub>2</sub>.

In subgroups of Formula II, at least one of  $R^4$  and  $R^7$  is a substituent group other than H.  
 In subgroups of Formula II, both of  $R^4$  and  $R^7$  are substituent groups other than H.

In subgroups of Formula I, Ia, Ib, and II, Z is selected from  $-S(O)_2CH_3$  and  $-C(=O)X$ ,  
 5 where X is selected from H,  $-OC_1-C_3$  alkyl,  $-CH_3$ , and  $-NR^{10}R^{11}$ .


Specific examples of the compounds of this invention are provided in the examples and  
 are summarized in Table 1 below and Tables 2-5 in the examples. The specific embodiments include the  
 compounds and pharmaceutically acceptable salts of the compounds.

10


**Table 1**

Example	Structure
Ex. 6	Ex. 7


Ex. 8




Ex. 9




Ex. 10




Ex. 11




Ex. 12




Ex. 13




Ex. 14




Ex. 19




Ex. 22




Ex. 28




Ex. 29




Ex.30




Ex. 31




Ex. 32




Ex. 33




Ex. 34




Ex. 35




Ex. 36




Ex. 37




Ex. 38




Ex. 39




Ex. 40




Ex. 41




Ex. 42




Ex. 43




Ex 44




Ex. 45




Ex 46



Ex. 47




Ex 48







Ex. 134




Ex 135




Ex. 136




Ex 137




Ex. 138




Ex 139




Ex. 140




Ex 141




Ex. 142




Ex 143




Ex. 144




Ex 145




Ex. 146




Ex 147




Ex. 148



Ex 149



Ex. 150

Definitions

“Ac” is acetyl, which is CH<sub>3</sub>C(=O)-.

"Alkyl" means saturated carbon chains which may be linear or branched or combinations thereof, unless the carbon chain is defined otherwise. Other groups having the prefix "alk", such as alkoxy and alkanoyl, also may be linear or branched or combinations thereof, unless the carbon chain is defined otherwise. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

"Alkylene" groups are alkyl groups that are difunctional rather than monofunctional. For example, methyl is an alkyl group and methylene (-CH<sub>2</sub>-) is the corresponding alkylene group.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

"Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

"Cycloalkyl" means a saturated carbocyclic ring having from 3 to 8 carbon atoms, unless otherwise stated. The term also includes a cycloalkyl ring fused to an aryl group. Examples of cycloalkyl include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like. "Cycloalkenyl" means a non-aromatic carbocyclic ring having one or more double binds.

"Aryl" (and "arylene") when used to describe a substituent or group in a structure means a monocyclic or bicyclic compound in which the rings are aromatic and which contains only carbon ring atoms. The term "aryl" can also refer to an aryl group that is fused to a cycloalkyl or heterocycle. Preferred "aryls" are phenyl and naphthyl. Phenyl is generally the most preferred aryl group.

"Heterocyclyl," "heterocycle," and "heterocyclic" means a fully or partially saturated or aromatic 5-6 membered ring containing 1-4 heteroatoms in the ring independently selected from N, S and O, unless otherwise stated. The heterocyclic ring may also be defined to include an optional carbonyl group or -N(O)-group as part of the ring structure. An example of the latter is pyridine N-oxide.

"Benzoheterocycle" represents a phenyl ring fused to a 5-6-membered heterocyclic ring having 1-2 heteroatoms, each of which is O, N, or S, where the heterocyclic ring may be saturated or unsaturated (i.e. the heterocyclic ring may have 1-2 double bonds in addition to the double bond of the phenyl ring). Examples include indole, 2,3-dihydroindole, benzofuran, 2,3-dihydrobenzofuran, quinoline, and isoquinoline. When the fused heterocycle is aromatic, the benzoheterocycle may also be referred to as benzoheteroaromatic or benzheteroaryl.

"Halogen" includes fluorine, chlorine, bromine and iodine. Halogen substituents are most often fluorine or chlorine.

"Me" represents methyl.

The term "composition," as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutically acceptable carrier.

The substituent "tetrazole" means a 2*H*-tetrazol-5-yl substituent group and tautomers thereof.

#### Optical Isomers - Diastereomers - Geometric Isomers - Tautomers

Compounds of Formula I may contain one or more asymmetric centers and can thus occur as racemates, racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

Some of the compounds described herein may contain olefinic double bonds, and unless specified otherwise, are meant to include both E and Z geometric isomers.

Some of the compounds described herein may exist as tautomers. An example is a ketone and its enol form, known as keto-enol tautomers. The individual tautomers as well as mixtures thereof are encompassed with compounds of Formula I.

Compounds of the Formula I having one or more asymmetric centers may be separated into diastereoisomers, enantiomers, and the like by methods well known in the art.

Alternatively, enantiomers and other compounds with chiral centers may be synthesized by stereospecific synthesis using optically pure starting materials and/or reagents of known configuration.

Some of the crystalline forms of compounds of the present invention may exist as polymorphs, and as such are intended to be included in the present invention. In addition, some of the compounds of the instant invention may form solvates with water or common organic solvents. Such solvates and hydrates are likewise encompassed within the scope of this invention.

Some of the biphenyl and biaryl compounds herein are observed as mixtures of atropisomers (rotamers) in the NMR spectra. The individual atropisomers as well as the mixtures are encompassed with the compounds of this invention.

### Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts in the solid form may exist in more than one crystal structure, and may also be in the form of hydrates. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

### Metabolites – Prodrugs

Therapeutically active metabolites, where the metabolites themselves fall within the scope of the claimed invention, are also compounds of the current invention. Prodrugs, which are compounds that are converted to the claimed compounds as they are being administered to a patient or after they have been administered to a patient, are also compounds of this invention.

Prodrugs which yield compounds of Formula I *in vivo* in which Z is -CO<sub>2</sub>H are also compounds of this invention. Compounds of Formula I in which Z is -CO<sub>2</sub>H are generally unstable to make and use in pharmaceutical compositions.

### Utilities

Compounds of the current invention are potent inhibitors of CETP. They are therefore useful in treating diseases and conditions that are treated by inhibitors of CETP.

5 One aspect of the present invention provides a method for treating or reducing the risk of developing a disease or condition that may be treated or prevented by inhibition of CETP by administering a therapeutically effective amount of a compound of this invention to a patient in need of treatment. A patient is a human or mammal, and is most often a human. A "therapeutically effective amount" is the amount of compound that is effective in obtaining a desired clinical outcome in the  
10 treatment of a specific disease.

Diseases or conditions that may be treated with compounds of this invention, and diseases which the patient may have a reduced risk of developing as a result of being treated with the compounds of this invention, include: atherosclerosis, peripheral vascular disease, dyslipidemia, hyperbetalipoproteinemia, hypoalphalipoproteinemia, hypercholesterolemia, hypertriglyceridemia,  
15 familial-hypercholesterolemia, cardiovascular disorders, angina, ischemia, cardiac ischemia, stroke, myocardial infarction, reperfusion injury, angioplastic restenosis, hypertension, vascular complications of diabetes, obesity and endotoxemia.

The compounds of this invention are expected to be particularly effective in raising HDL-C and/or increasing the ratio of HDL-C to LDL-C. These changes in HDL-C and LDL-C may be  
20 beneficial in treating atherosclerosis, reducing or reversing the development of atherosclerosis, reducing the risk of developing atherosclerosis, or preventing atherosclerosis.

### Administration and Dose Ranges

Any suitable route of administration may be employed for providing a mammal,  
25 especially a human, with an effective dose of a compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like. Preferably compounds of Formula I are administered orally.

The effective dosage of active ingredient employed may vary depending on the particular  
30 compound employed, the mode of administration, the condition being treated and the severity of the condition being treated. Such dosage may be ascertained readily by a person skilled in the art.

When treating the diseases for which compounds of Formula I are indicated, generally satisfactory results are obtained when the compounds of the present invention are administered at a daily dosage of from about 0.01 milligram to about 100 milligram per kilogram of animal or human body

weight, preferably given as a single daily dose or in divided doses two to six times a day, or in sustained release form. In the case of a 70 kg adult human, the total daily dose will generally be from about 0.5 milligram to about 500 milligrams. For a particularly potent compound, the dosage for an adult human may be as low as 0.1 mg. The dosage regimen may be adjusted within this range or even outside of this range to provide the optimal therapeutic response.

Oral administration will usually be carried out using tablets. Examples of doses in tablets are 0.5 mg, 1 mg, 2 mg, 5 mg, 10 mg, 25 mg, 50 mg, 100 mg, 250 mg, and 500 mg. Other oral forms can also have the same dosages (e.g. capsules).

#### 10 Pharmaceutical Compositions

Another aspect of the present invention provides pharmaceutical compositions which comprise a compound of Formula I and a pharmaceutically acceptable carrier. The pharmaceutical compositions of the present invention comprise a compound of Formula I or a pharmaceutically acceptable salt as an active ingredient, as well as a pharmaceutically acceptable carrier and optionally other therapeutic ingredients. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic bases or acids and organic bases or acids. A pharmaceutical composition may also comprise a prodrug, or a pharmaceutically acceptable salt thereof, if a prodrug is administered. Pharmaceutical compositions may also consist essentially of a compound of Formula I and a pharmaceutically acceptable carrier.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular (ophthalmic), pulmonary (nasal or buccal inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

In practical use, the compounds of Formula I can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents



and the like in the case of oral solid preparations such as, for example, powders, hard and soft capsules and tablets, with the solid oral preparations being preferred over the liquid preparations.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed.

5 If desired, tablets may be coated by standard aqueous or nonaqueous techniques. Such compositions and preparations should contain at least 0.1 percent of active compound. The percentage of active compound in these compositions may, of course, be varied and may conveniently be between about 2 percent to about 60 percent of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that an effective dosage will be obtained. The active compounds can also be  
10 administered intranasally as, for example, liquid drops or spray.

The tablets, pills, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin. When a dosage unit form is a capsule, it may contain, in  
15 addition to materials of the above type, a liquid carrier such as a fatty oil.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

20 Compounds of formula I may also be administered parenterally. Solutions or suspensions of these active compounds can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

25 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or  
30 dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

#### Combination Therapy

Compounds of the invention (e.g. Formula I) may be used in combination with other drugs that may also be useful in the treatment or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of Formula I. When a  
5 compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the compound of Formula I is preferred. However, the combination therapy also includes therapies in which the compound of Formula I and one or more other drugs are administered on different schedules.

When oral formulations are used, the drugs may be combined into a single combination  
10 tablet or other oral dosage form, or the drugs may be packaged together as separate tablets or other oral dosage forms. It is also contemplated that when used in combination with one or more other active ingredients, the compound of the present invention and the other active ingredients may be used in lower doses than when each is used singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a compound of  
15 Formula I.

Examples of other active ingredients that may be administered in combination with a compound of this invention (e.g. Formula I), and either administered separately or in the same pharmaceutical composition, include, but are not limited to, other compounds which improve a patient's lipid profile, such as (i) HMG-CoA reductase inhibitors, (which are generally statins, including  
20 lovastatin, simvastatin, rosuvastatin, pravastatin, fluvastatin, atorvastatin, rivastatin, itavastatin, ZD-4522 and other statins), (ii) bile acid sequestrants (cholestyramine, colestipol, and dialkylaminoalkyl derivatives of a cross-linked dextran), (iii) niacin and related compounds, such as nicotinyl alcohol, nicotinamide, and nicotinic acid or a salt thereof, (iv) PPAR $\alpha$  agonists, such as gemfibrozil and fenofibric acid derivatives (fibrates), including clofibrate, fenofibrate and bezafibrate, (v) cholesterol  
25 absorption inhibitors, such as ezetimibe, (vi) acyl CoA:cholesterol acyltransferase (ACAT) inhibitors, such as avasimibe, (vii) phenolic anti-oxidants, such as probucol, and (viii) a microsomal triglyceride transfer protein (MTP)/ApoB secretion inhibitor.

Preferred classes of therapeutic compounds that can be used with the compounds of this invention for use in improving a patient's lipid profile (i.e. raising HDL-C and lowering LDL-C) include  
30 one or both of statins and cholesterol absorption inhibitors. Particularly preferred are combinations of compounds of this invention with simvastatin, ezetimibe, or both simvastatin and ezetimibe. Also preferred are combinations with atorvastatin, ezetimibe, or both compounds.

Finally compounds of this invention can be used with compounds that are useful for treating other disease, such as diabetes and obesity, as well as other anti-atherosclerotic compounds.

Examples of other active ingredients that may be administered in combination with a compound of this invention include, but are not limited to:

(a) PPAR gamma agonists and partial agonists, including glitazones and non-glitazones (e.g. pioglitazone, englitazone, MCC-555, rosiglitazone, balaglitazone, netoglitazone, T-131, LY-300512, and LY-818;

(b) biguanides such as metformin and phenformin;

(c) protein tyrosine phosphatase-1B (PTP-1B) inhibitors,

(d) dipeptidyl peptidase IV (DP-IV) inhibitors;

(e) insulin or insulin mimetics;

(f) sulfonylureas, such as tolbutamide and glipizide, or related materials;

(g)  $\alpha$ -glucosidase inhibitors (such as acarbose);

(h) agents which improve a patient's lipid profile, as described previously;

(i) PPAR $\alpha/\gamma$  dual agonists, such as muraglitazar, tesaglitazar, farglitazar, and JT-501;

(j) PPAR $\delta$  agonists such as those disclosed in WO97/28149;

(k) antiobesity compounds, including 5-HT(serotonin) inhibitors, neuropeptide Y5 (NPY5) inhibitors, melanocortin 4 receptor (Mc4r) agonists, cannabinoid receptor 1 (CB-1) antagonists/inverse agonists, and  $\beta$ 3 adrenergic receptor agonists;

(l) ileal bile acid transporter inhibitors;

(m) agents intended for use in inflammatory conditions such as aspirin, non-steroidal anti-inflammatory drugs, glucocorticoids, azulfidine, and cyclo-oxygenase 2 selective inhibitors, including etoricoxib and rofecoxib;

(n) glucagon receptor antagonists;

(o) GLP-1,

(p) GIP-1, and

(q) GLP-1 analogs, such as exendins, for example exenitide.

The combination therapies described above which use the compounds of this invention may also be useful in the treatment of the metabolic syndrome. According to one widely used definition, a patient having metabolic syndrome is characterized as having three or more symptoms selected from the following group of five symptoms: (1) abdominal obesity; (2) hypertriglyceridemia; (3) low high-density lipoprotein cholesterol (HDL); (4) high blood pressure; and (5) elevated fasting glucose, which may be in the range characteristic of Type 2 diabetes if the patient is also diabetic. Each of these symptoms is defined clinically in the recently released Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in




Data was evaluated by obtaining an initial rate, expressed in relative fluorescence units per second, for the pseudolinear portion of the curve, often 0-500 or 1000 sec. Comparison of the rates of samples with inhibitors to an uninhibited (DMSO only) positive control yielded a percent inhibition. A plot of percent inhibition vs. log of inhibitor concentration, fit to a Sigmoidal 4 parameter equation was used to calculate IC<sub>50</sub>.

## EXAMPLES

The following examples are provided so that the invention will be more fully appreciated and understood. They should not be construed as limiting the invention in any way. The scope of the invention is defined by the appended claims. Examples 1-5, 15-18, 20, 21, 23-27, 49, and 50 are examples of synthetic intermediates. The remaining examples are compounds of the invention. Compounds of this invention have an IC<sub>50</sub> value as measured using the assay described above of less than or equal to 50 μM.


The following Schemes are provided to further teach how compounds that are not specifically disclosed herein can be synthesized by one of ordinary skill in the art. Starting materials are made using known procedures or as illustrated. Some starting materials may also be available for purchase.

## SCHEME 1



Intermediates 1-2, 1-3 and 1-4 utilized in the present invention can be purchased or prepared as shown in Scheme 1. An appropriately substituted 2-haloaniline wherein R<sup>a</sup> and p are as defined in the claims and where the halogen is preferably iodo or bromo is treated with CuCN in DMF at elevated temperature to afford the corresponding 2-cyanoaniline. Alternatively, the nitrile can be prepared by treatment of 1-1 with KCN and CuI in the presence of a palladium (II) salt or in the presence of certain copper or nickel complexes ( See: Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 867 (2001) and references therein). Iodides 1-3 are prepared by treatment of 1-2 with isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite or the like in diiodomethane (see for example: Smith et al., *J. Org. Chem.* 55, 2543, (1990) and references cited therein). Alternatively, the iodide can be prepared first by diazonium formation using isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite, sodium nitrite, nitrous acid or the like followed by heating in the presence of iodine or an iodide salt such as copper iodide, sodium iodide, potassium iodide, tetrabutylammonium iodide or the like. Reduction of nitrile 1-2 is carried out using Raney nickel under hydrogen atmosphere in water, methanol, ethanol or the like to afford 2-aminomethyl aniline 1-4. Alternatively, the nitrile can be reduced with Palladium on carbon under hydrogen atmosphere in methanol, ethanol or the like or with lithium aluminum hydride, lithium borohydride, borane or the like in ether, tetrahydrofuran, dimethoxyethane or the like to afford 1-4. Other methods for reduction of a nitrile to an aminomethyl group can be found in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 1204 (2001) and references therein.

SCHEME 2




- 5 Intermediates 2-4 utilized in the present invention wherein  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in Scheme 2. Aminomethylaniline 2-1 can be purchased or prepared according to the procedure outlined in Scheme 1. Alkylation of 2-1 can be carried out by treatment with an appropriately substituted alkyl halide, mesylate or tosylate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a base such as triethylamine, diisopropylethylamine, N-methylmorpholine, lithium diisopropylamide or lithium-, sodium-, or potassium bis(trimethylsilyl)amide or the like to afford disubstituted amine 2-2.
- 10 Alternatively, treatment of amine 2-1 with an appropriately substituted aldehyde in the presence of a reducing agent such as sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride or the like in methanol, ethanol, dichloroethane, tetrahydrofuran or the like or according to methods

described in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 1187-1189 (2001) and references cited therein affords dialkylamine 2-2. Alternatively, amine 2-1 can be treated with an appropriately substituted carboxylic acid halide, carboxylic acid activated ester or carboxylic acid in the presence of a coupling reagent such as EDC, 5 DCC, BOP, HATU or the like and a hindered base such as triethylamine, diisopropylethylamine, N-methyl morpholine or the like in a suitable solvent such as dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like to give the corresponding amide which can be reduced with LAH, borane or the like to afford dialkylamine 2-2. Acylation of 2-2 with a carboxylic acid and coupling reagent such as EDC EDC, DCC, BOP, HATU or the like, or a carboxylic acid activated ester or an acid 10 halide, alkylchloroformate, alkylisocyanate, alkylsulfonyl chloride, chlorosulfonyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like affords 2-3. Iodide 2-4 can be prepared by treatment of 2-3 with isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite or the like in diiodomethane (see for example: Smith et al., *J. Org. Chem.* 55, 2543, (1990) and 15 references cited therein). Alternatively, the iodide can be prepared first by diazonium formation using isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite, sodium nitrite, nitrous acid or the like followed by heating in the presence of iodine or an iodide salt such as copper iodide, sodium iodide, potassium iodide, tetrabutylammonium iodide or the like.




SCHEME 3



- 5 Haloaryl intermediates 3-4 of the present invention wherein  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in scheme 3. 2-halobenzylbromides 3-2 wherein the halo is preferably iodo or bromo can be purchased or prepared from 2-halotoluenes via benzylic bromination using N-bromosuccinimide in carbon tetrachloride. Treatment of 3-2 with an appropriately substituted benzylamine affords dialkyl amine 3-3. Acylation of 3-3 with a carboxylic acid and coupling reagent
- 10 such as EDC EDC, DCC, BOP, HATU or the like, or a carboxylic acid activated ester or an acid halide, alkylchloroformate, alkylisocyanate, alkylsulfonyl chloride, chlorosulfonyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like affords 3-4.


15

SCHEME 4



- 5 Biaryl compounds 4-2 of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared from aryl halide 4-1 where the halide is preferably bromo or iodo via a Suzuki or Stille reaction wherein 4-1 is coupled with an appropriately substituted aryl boronic acid, aryl boronate ester or aryl trialkyl tin in the presence of a palladium catalyst. The coupling reaction may be carried out using Pd(II)acetate and potassium carbonate in aqueous acetone at reflux. Alternatively the reaction may
- 10 employ tetrakis(triphenylphosphine)palladium in an ethanol/toluene mix in the presence of sodium carbonate. Alternatively, as practiced by those skilled in the art the reaction can employ a number of Palladium (0) compounds and Palladium (II) salts in a number of solvents and in the presence of a variety of ligands, bases, and promoters, generally but not exclusively, with heating and/or microwave irradiation. Some appropriate reaction conditions can be found described in Miyaura et al., *Chem. Rev.* 95, 2457 (1995) and references cited within and as described in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 868-869 (2001) and references cited therein. Compounds 4-1 are prepared as shown in Schemes 2 and 3.
- 15


SCHEME 5



- 5 Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in Scheme 5. 2-cyano iodobenzenes 5-1 can be purchased or prepared according to the procedures outlined in Scheme 1. Biphenyl compounds 5-2 are prepared via a Suzuki or Stille reaction or variation thereof employing palladium catalyzed cross coupling of iodide 5-1 with an appropriately substituted aryl boronic acid, aryl boronate ester or aryl trialkyl tin as described in Miyaua et al., *Chem. Rev.* 95, 2457 (1995) and references cited within and as described in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 868-869 (2001) and references cited therein. Reduction of nitrile 5-2 is accomplished with Lithium aluminum hydride in diethyl ether to afford 2-aminomethyl aniline 5-3. Alternatively, the nitrile can be reduced with Palladium on carbon or Raney nickel under hydrogen atmosphere in methanol, ethanol or the like. Other methods for reduction of a nitrile to an aminomethyl group can be found in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 1204 (2001) and references therein. Alkylation of 5-3 is achieved by treatment with an appropriately substituted benzyl halide, mesylate or tosylate or the like in dichloromethane, dichloroethane,

tetrahydrofuran, dimethoxyethane or the like in the presence of a base such as triethylamine, diisopropylethylamine, N-methylmorpholine, lithium diisopropylamide or lithium-, sodium-, or potassium bis(trimethylsilyl)amide or the like to afford disubstituted amine 5-4. Alternatively, treatment of amine 5-3 with an appropriately substituted aldehyde in the presence of a reducing agent such as sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride or the like in methanol, ethanol, dichloroethane, tetrahydrofuran or the like or according to methods described in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 1187-1189 (2001) and references cited therein affords disubstituted amine 5-4. Alternatively, amine 5-3 can be treated with an appropriately substituted carboxylic acid halide, carboxylic acid activated ester or carboxylic acid in the presence of a coupling reagent such as EDC, DCC, BOP, HATU or the like and a hindered base such as triethylamine, diisopropylethylamine, N-methyl morpholine or the like in a suitable solvent such as dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like to give the corresponding amide which can be reduced with LAH, borane or the like to afford disubstituted amine 5-4. Acylation of 5-4 with a carboxylic acid and coupling reagent such as EDC EDC, DCC, BOP, HATU or the like, or a carboxylic acid activated ester or an acid halide, alkylchloroformate, alkylisocyanate, alkylsulfonyl chloride, chlorosulfonyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like affords 5-5.

SCHEME 6




- 5 Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared by the procedure described in Scheme 6. Base mediated hydrolysis of 2-cyanobenzenes 6-1 can be achieved by treatment with aqueous potassium hydroxide or the like in the presence of protic solvents such as isopropanol to afford the benzoic acid 6-2. Reduction to the benzyl alcohol 6-3 can be achieved with borane in tetrahydrofuran as described in Smith, M.B. and March, J. "March's Advanced
- 10 Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New York, pp. 1549 (2001) and references therein. Conversion of alcohol 6-3 to bromide 6-4 can be achieved with carbon tetrabromide and triphenylphosphine in dichloromethane (see for example: Iwasaki et al., *Tetrahedron*, 52, 13327 (1996) and references cited therein). Treatment of 6-4 with an appropriately substituted amine followed by acylation with a carboxylic acid and coupling reagent such as EDC EDC, DCC, BOP, HATU or the like ,
- 15 or a carboxylic acid activated ester or an acid halide, alkylchloroformate, alkylisocyanate, alkylsulfonyl chloride, chlorosulfonyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran,

dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like affords 6-5.

SCHEME 7


5



Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared by acylation of benzylamine 7-1 with a carboxylic acid and coupling reagent such as EDC

- 10 EDC, DCC, BOP, HATU or the like, or a carboxylic acid activated ester or an acid halide, alkylchloroformate, alkylisocyanate, alkylsulfonyl chloride, chlorosulfonyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like affords 7-2. Alkylation of 7-2 can be carried out by treatment with an appropriately substituted alkyl
- 15 halide, mesylate or tosylate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a base such as triethylamine, diisopropylethylamine, N-methylmorpholine, lithium diisopropylamide or lithium-, sodium-, or potassium bis(trimethylsilyl)amide or the like to afford compounds 7-3.

SCHEME 8




5

Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in scheme 8. Nitro compounds 8-1 can be reduced with catalysts such as platinum oxide and the like under a hydrogen atmosphere in solvents such as tetrahydrofuran and the like to afford aryl amines 8-2. Acylation of amine 8-2 with an acid chloride, alkylchloroformate, sulfonyl chloride, alkylisocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like in the presence of a hindered base such as triethylamine, diisopropylethylamine, N-methyl morpholine, pyridine or the like affords compounds 8-3. Alternatively, alkylation of aniline 8-2 with an alkyl halide, mesylate or tosylate or the like in the presence of organic bases such as potassium hexamethyldisilazide and the like in solvents such as tetrahydrofuran, diethyl ether and the like affords compounds 8-3.

15


SCHEME 9



- 5 Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in scheme 9. Aryl amine 9-1 can be converted to the corresponding aryl iodide 9-2 by treatment with isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite or the like in diiodomethane (see for example: Smith et al., *J. Org. Chem.* 55, 2543, (1990) and references cited therein). Alternatively, the iodide can be prepared first by diazonium formation using isoamyl nitrite, n-pentyl nitrite, t-butyl nitrite, sodium nitrite, nitrous acid or the like followed by heating in the presence of iodine or an iodide salt such as copper iodide, sodium iodide, potassium iodide, tetrabutylammonium iodide or the like. to afford compounds 9-1. Compounds 9-2 can be converted to compounds 9-3 via Suzuki or Stille reaction by coupling with an appropriately substituted aryl boronic acid, aryl boronate ester or aryl trialkyl tin in the presence of a palladium catalyst (see for example: Miyaura et al., *Chem Rev.* 95, 2457 and references cited therein).
- 10
- 15




SCHEME 10




- 5 Compounds of the present invention wherein  $A^1$ ,  $A^2$ ,  $R^1$ ,  $R^a$ ,  $Z$ ,  $n$  and  $p$  are as defined in the claims can be prepared as shown in scheme 10. Treatment of nitriles 10-1 with aqueous formic acid in the presence of catalysts such as platinum oxide and the like affords aldehyde 10-2 which can be further modified by reaction with organophosphorane reagents such as methyltriphenylphosphonium bromide as described in Smith, M. B. and March, J. "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., John Wiley and Sons, New
- 10 York, pp. 1231-1237 (2001) to give compounds 10-3.

SCHEME 11



- 5 Phenethyl substituted compounds 11-4 of the present invention can be prepared as described in scheme 11. Intermediate 11-2 can be prepared by treating benzylamine 11-1 with a appropriately substituted phenyl acetic acid halide, phenyl acetic acid activated ester or phenyl acetic acid in the presence of a coupling reagent such as EDC, DCC, BOP, HATU or the like in a suitable solvent such as dichloromethane, dichloroethane and the like. Amide 11-2 can be reduced with lithium aluminum
- 10 hydride, borane or the like in ether, tetrahydrofuran or the like to afford 11-3. Acylation of 11-3 with an acid chloride, alkyl chloroformate, alkyl isocyanate or the like in dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane or the like affords compounds 11-4.

## EXAMPLE 1




5

2-Amino-5-(trifluoromethyl)benzonitrile

In a 2-liter flask equipped with a septum inlet, magnetic stirring bar, and condenser leading to a mercury bubbler was placed 100g (0.348 mol) of 4-amino-3-iodobenzotrifluoride in 1 liter DMF. To this reaction  
10 flask with stirring was added 37.4 g (0.416mol) of copper cyanide and the contents were heated to 100 °C under nitrogen for fourteen hours. Afterwards, TLC analysis revealed minor starting material (20/80) ethyl acetate/hexane. The contents were cooled to 40°C and filtered through a pad of Celite. The solids were washed with ether and the filtrate transferred to a separatory funnel and washed with about 5% aqueous ammonium hydroxide. The aqueous layer was back-extracted with ether and the organics were  
15 combined and washed with brine, dried over magnesium sulfate, filtered and concentrated to yield a black oil. The oil was chromatographed on a Biotage 75L cartridge with ethyl acetate/hexane to obtain 68 g 2-amino-5-(trifluoromethyl)benzonitrile as a black solid.

## EXAMPLE 2

20


[2-(Aminomethyl)-4-(trifluoromethyl)phenyl]amine

In a rocker shaker was charged 37 g (0.2 mol) of 2-amino-5-(trifluoromethyl)benzonitrile in 500 mL of ethanol and 76 mL of ammonium hydroxide. The contents were degassed/blanketed with nitrogen, 25 g  
25 of Raney nickel were added, then the shaker was pressurized with 40 psi hydrogen gas. At 15 hours TLC analysis (20/80 ethyl acetate/hexane) revealed no starting material. The contents were filtered through

Celite and washed with ether and concentrated to 37 g black oil. The oil was chromatographed on a Biotage 75L cartridge with methanol/methylene chloride and ammonium hydroxide/methanol/methylene chloride to give [2-(aminomethyl)-4-(trifluoromethyl)phenyl]amine as a green oil.

5


## EXAMPLE 3

[2-Amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]amine

- 10 In a 500-mL flask equipped with a septum inlet, magnetic stirring bar, and condenser leading to a mercury bubbler was placed 22.8 g (0.12 mol) of [2-(aminomethyl)-4-(trifluoromethyl)phenyl]amine in 218 mL dimethoxyethane. To this reaction flask with stirring was added 26 mL (0.24 mol) N-methylmorpholine and 24 mL (0.13 mol) 3,5-bis(trifluoromethyl)benzyl bromide. The contents were heated to and maintained at reflux under nitrogen for 12 hours. Afterwards, TLC analysis revealed a
- 15 minor amount of starting material remaining. The contents were cooled to room temperature, diluted with ether and transferred to a separatory funnel and washed with water and brine, dried over magnesium sulfate, filtered and concentrated to afford an oil. The oil was dissolved in hexane, cooled in an ice bath and acetic acid was added slowly while stirring to precipitate solids. The solids were filtered, dissolved in ether, and cooled in an ice bath and stirred while saturated sodium bicarbonate was added slowly until
- 20 basic. The contents were transferred to a separatory funnel and washed with brine, dried over magnesium sulfate, filtered, and concentrated to give 12 g [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]amine as an orange oil. In addition the hexane mother liquor contained significant amounts of product and byproducts. This liquor was concentrated to yield dark yellow oil that was used without further purification in the following step.


25

## EXAMPLE 4

5 Methyl [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]carbamate


In a 250-mL flask equipped with a septum inlet, magnetic stirring bar, and connecting tube leading to a mercury bubbler was placed 12 g (0.029 mol) of [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]amine in 120 mL DME. To the reaction flask with stirring was added 6.4 mL (0.058 mol) N-methylmorpholine and the mixture was cooled in an ice bath under nitrogen. To this mixture was added dropwise over ten minutes, 2.7 mL (0.035 mol) of methyl chloroformate. Afterwards, the mixture was allowed to stir for 15 minutes and HPLC showed no starting material present. Water was added to the mixture and the contents were diluted with hexane and transferred to a separatory funnel. The layers were separated, then washed with brine, dried over magnesium sulfate, filtered and concentrated to give 13 g crude oil. The oil was chromatographed on a Biotage 40M cartridge with methylene chloride/hexane to give methyl [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]carbamate as a yellow oil.

## EXAMPLE 5

20 Methyl [3,5-bis(trifluoromethyl)benzyl][2-iodo-5-(trifluoromethyl)benzyl]carbamate

In a 250-mL flask equipped with a septum inlet, magnetic stirring bar, and condenser leading to a mercury bubbler was placed 9 g (0.019 mol) of methyl [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl]carbamate in 90 mL  $\text{CHCl}_3$ . To this stirred solution was added 3.6 mL (0.027 mol) of isoamyl nitrite (tech. grade). The mixture was stirred at room temperature for 5 minutes under a nitrogen blanket. To this was added 9.7 g (0.038 mol) of iodine and the mixture was heated to and maintained at reflux for 2 hrs. The mixture was cooled and diluted with methylene chloride, transferred to a separatory funnel and washed with saturated aqueous sodium thiosulfate and brine; dried over magnesium sulfate, filtered, concentrated and chromatographed on a Biotage 40M cartridge with methylene chloride/hexane to give methyl [3,5-bis(trifluoromethyl)benzyl][2-iodo-5-(trifluoromethyl)benzyl]carbamate.

## EXAMPLE 6




15 Methyl [3,5-bis(trifluoromethyl)benzyl][2'-methoxy-5'-methyl-4-(trifluoromethyl)biphenyl-2-yl]methyl carbamate

A mixture of methyl [3,5-bis(trifluoromethyl)benzyl][2-iodo-5-(trifluoromethyl)benzyl]carbamate (0.10 g, 0.17 mmol) from Example 5, and 2-methoxy-5-methylphenyl boronic acid (0.034 g, 0.2 mmol), palladium acetate (0.0038 g, 0.017 mmol) and potassium carbonate (0.047 g, 0.34 mmol) in 4:1 acetone/water (10 ml) was heated and maintained at reflux for 1 h. The color of the solution turned dark. Acetone was removed and the residue was extracted with methylene chloride (3 x 10 ml). The combined organic layers were washed with water, then brine, and dried over sodium sulfate. The title compound was obtained as a colorless oil by preparative thin layer chromatography using acetone/hexane (5:95) as the eluant.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz):  $\delta$  7.73 (s, 1H), 7.58 (d,  $J$  = 8.5 Hz, 1H), 7.47 (s, 2H), 7.35 (m,

1H), 7.32 (d,  $J = 8.0$  Hz, 1H), 7.19 (m, 1H), 6.85 (m, 2H), 4.23-4.36 (m, 4H), 3.79 (m, 3H), 3.68 (s, 3H), 2.25 (s, 3H). LC-MS ( $M+1$ ): 580.0 (4.61 min).

## EXAMPLE 7



Methyl [3,5-bis(trifluoromethyl)benzyl][(4-fluoro-2'-methoxy-5'-methylbiphenyl-2-yl)methyl]carbamate

Step A: [3,5-Bis(trifluoromethyl)benzyl](2-bromo-5-fluorobenzyl)amine

- 10 To a solution of 2-bromo-5-fluoro benzyl bromide (0.5 g, 1.86 mmol) and 3,5-bis(trifluoromethyl)benzylamine (0.45 g, 1.86 mmol) in methylene chloride (10 ml) at 0 °C, triethyl amine (0.39 ml, 2.80 mmol) was added. The solution was slowly warmed to room temperature and stirred for 18 h. The solvent was removed under reduced pressure. The title compound was obtained by flash column chromatography using EtOAc/hexane (2:8) as the eluant.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.88 (s, 2H), 7.80 (s, 1H), 7.53 (dd,  $J = 8.8, 5.3$  Hz, 1H), 7.18 (dd,  $J = 9, 3.3$  Hz, 1H), 6.91 (dt,  $J = 8.5, 3.3$  Hz, 1H), 3.98 (s, 2H), 3.90 (s, 2H).
- 15

Step B: Methyl [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-fluorobenzyl)carbamate


- To a solution of [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-fluorobenzyl)amine from Step A (0.14 g) in methylene chloride at room temperature, excess methyl chloroformate (0.1 ml) and triethyl amine (0.1 ml) were added. The solution was stirred at room temperature for 2 h and the solvent was removed under reduced pressure. The title compound was obtained by flash column chromatography using EtOAc/hexane (1:9) as the eluant.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.80 (s, 1H), 7.68 (s, 1H), 7.60 (s, 1H), 7.48(m, 1H), 6.88 (m, 2H), 4.60 (m, 4H), 3.81 (s, 3H).
- 20

Step C: Methyl [3,5-bis(trifluoromethyl)benzyl][(4-fluoro-2'-methoxy-5'-methylbiphenyl-2-yl)methyl]carbamate

25

A mixture of the Methyl [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-fluorobenzyl)carbamate from Step B (0.02g, 0.04 mmol), potassium carbonate (0.011 g, 0.08 mmol), 2-methoxy-5-methyl phenyl boronic acid (0.008 g, 0.05 mmol) and catalytic amount of palladium acetate (1 mg) in 4:1 acetone/water (2.5 ml) was heated to and maintained at reflux for 1 h. TLC analysis (acetone/hexane = 5:95) showed no starting material. The solvent was removed under reduced pressure and the residue was extracted with methylene chloride (3 x 10 ml). The combined organic layers were washed with brine and dried over sodium sulfate. The title compound was obtained by preparative thin layer chromatography using acetone:hexane (5:95) as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.75 (s, 1H), 7.50 (s, 1H), 7.38 (s, 1H), 7.20 (m, 2H), 7.04 (m, 2H), 6.84 (s, 1H), 6.80 (m, 1H), 4.18-4.60 (m, 4H), 3.80 (s, 3H), 3.68 (s, 3H), 2.32 (s, 3H). LC-MS (M+1) 530.4 (4.43 min).

## EXAMPLE 8



5

Methyl [3,5-bis(trifluoromethyl)benzyl] [(5'-isopropyl-2'-methoxy-4-nitrobiphenyl-2-yl)methyl]carbamate  
 Step A: (2-Bromo-5-nitrophenyl)methanol

To a solution of methyl 2-bromo-5-nitro benzoate (0.52 g, 2.00 mmol) in THF (10 ml) at 0°C, a 1.0 M solution of DIBAL in THF (2.4 ml, 2.4 mmol) was added dropwise. The mixture was stirred at 0° C for 4 h. The reaction was quenched with water extracted with ethyl acetate (3 x 20 ml). The combined organic layers were washed with brine, and dried over sodium sulfate. The title compound was obtained as a white solid following flash column chromatography using 8:92 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.46 (d, *J* = 2.5 Hz, 1H), 8.05 (dd, *J* = 8.5, 3.0 Hz, 1H), 7.74 (d, *J* = 8.5 Hz, 1H), 4.84 (s, 2H).



Step B: 2-Bromo-5-nitrobenzyl methanesulfonate

To a solution of the (2-bromo-5-nitrophenyl)methanol from Step A (0.33 g, 1.42 mmol) in methylene chloride (20 ml) at 0° C, methanesulfonyl chloride (0.13 ml, 1.70 mmol) and triethylamine (0.3 ml, 2.1 mmol) were added. The solution was stirred at 0° C for 1h. The reaction was diluted with methylene chloride (30 ml). The solution was washed with water, brine, and dried over sodium sulfate. The title compound was obtained by flash column chromatography using 10:90 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.40 (d, *J* = 2.5 Hz, 1H), 8.17 (dd, *J* = 8.5, 3.0 Hz, 1H), 7.82 (d, *J* = 8.5 Hz, 1H), 5.40 (s, 2H), 3.20 (s, 3H).

Step C: [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-nitrobenzyl)amine

To a solution of 2-bromo-5-nitrobenzyl methanesulfonate from Step B (0.42 g, 1.35 mmol) and 3,5-bis-trifluoromethyl benzylamine (0.33 g, 1.35 mmol) in DMF (10 ml) at 0° C, potassium carbonate (0.37 g, 2.69 mmol) was added. The mixture was stirred at 0° C for 2h. TLC analysis (EtOAc/hexane = 2:8) showed completion of the reaction. Water (50 ml) was added. The mixture was extracted with EtOAc (3 x 40 ml). The combined organic layers were washed with brine, and dried over sodium sulfate. The title compound was obtained by flash column chromatography using 2:8 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.35 (d, *J* = 2.5 Hz, 1H), 8.03 (dd, *J* = 8.5, 2.5 Hz, 1H), 7.88 (s, 2H), 7.81 (s, 1H), 8.76 (d, *J* = 8.5 Hz, 1H), 4.00 (s, 4H).

Step D: Methyl [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-nitrobenzyl)carbamate

To a solution of [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-nitrobenzyl)amine from Step C (0.47 g, 1.0 mmol) in methylene chloride (20 ml) at room temperature, methyl chloroformate (0.12 ml, 1.54 mmol) and triethylamine (0.29 ml, 2.05 mmol) were added. The solution was stirred at room temperature for 18 h. The solvent was removed under reduced pressure. The title compound was obtained by flash column chromatography using 1:9 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.11 (m, 1H), 8.02 (dd, *J* = 9.0, 2.5 Hz, 1H), 7.81 (s, 1H), 7.74 (d, *J* = 9.0 Hz, 1H), 7.70 (m, 1H), 7.65 (m, 1H), 4.67 (m, 4H), 3.89 (s, 3H).


Step E: Methyl [3,5-bis(trifluoromethyl)benzyl][(5'-isopropyl-2'-methoxy-4-nitrobiphenyl-2-yl)methyl]carbamate

To a mixture of Methyl [3,5-bis(trifluoromethyl)benzyl](2-bromo-5-nitrobenzyl)carbamate from Step D (0.39 g, 0.75 mmol), 2-methoxy-5-isopropylphenyl boronic acid (0.175 g, 0.90 mmol), and potassium carbonate (0.21 g, 1.50 mmol) in 4:1 acetone/water (10 ml), a catalytic amount of palladium acetate (10 mg) was added. The mixture was heated to and maintained at reflux for 2 h. TLC analysis (acetone/hexane = 5:95) showed completion of the reaction. The acetone was removed under reduced pressure and the organic was extracted with methylene chloride (3 x 25 ml). The combined organic layers were washed with brine and dried over sodium sulfate. The title compound was obtained by flash

column chromatography using 5:95 acetone/hexane as the eluant.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  8.20 (m, 2H), 7.75 (s, 1H), 7.48 (M, 1H), 7.40 (d,  $J = 8.0$  Hz, 1H), 7.29 (m, 1H), 6.93 (m, 2H), 4.25-4.60 (m, 4H), 3.80 (m, 3H), 3.69 (s, 3H), 2.89 (m, 1H), 1.20 (s, 6H). LC-MS ( $M+1$ ) 585.4 (4.49 min).

5

## EXAMPLE 9



Methyl [(4-amino-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl][3,5-bis(trifluoromethyl)benzyl]carbamate

10


A solution of the methyl [3,5-bis(trifluoromethyl)benzyl][(5'-isopropyl-2'-methoxy-4-nitrobiphenyl-2-yl)methyl]carbamate from Example 8 (0.31 g, 0.53 mmol) in MeOH (10 ml) was charged with a catalytic amount Pd/C and hydrogen at 40 psi for 1 h. The mixture was filtered through Celite and the filtrate was concentrated. The title compound was obtained by flash column chromatography using 3:7

15

EtOAc/hexane as the eluant.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.72 (m, 1H), 7.50 (s, 1H), 7.30 (s, 1H), 7.16 (m, 1H), 7.03 (d,  $J = 8.0$  Hz, 1H), 6.94 (s, 1H), 6.82 (m, 1H), 6.69 (dd,  $J = 8.5, 2.5$  Hz, 1H), 6.58 (m, 1H), 4.18-4.30 (m, 4H), 3.75 (s, 3H), 3.65 (s, 3H), 2.83 (m, 1H), 1.20 (s, 6H). LC-MS ( $M+1$ ): 555.3 (3.58 min).

20

## EXAMPLE 10




5 Methyl [(4-iodo-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl][3,5-

To a solution of the methyl [(4-amino-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl][3,5-bis(trifluoromethyl)benzyl]carbamate from Example 9 (0.31 g, 0.56 mmol) in chloroform (10 ml), n-amyl nitrite (0.11 ml, 0.84 mmol) and iodine (0.28 g, 1.1 mmol) were added. The mixture was heated to and maintained at reflux for 2 h. TLC analysis (5:95 EtOAc/hexane) showed completion of the reaction. The mixture was diluted with methylene chloride (20 ml) and washed with saturated sodium thiosulfate solution, brine. The light yellow solution was dried over sodium sulfate. The title compound was

10 obtained by flash column chromatography using 5:95 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.74 (s, 1H), 7.67 (dd, *J* = 8.0, 2.0 Hz, 1H), 7.47 (m, 1H), 7.31 (dd, *J* = 8.0, 2.0 Hz, 1H), 7.21 (d, *J* = 8.0 Hz, 1H), 7.16 (d, *J* = 8.5 Hz, 1H), 6.95 (d, *J* = 8.0 Hz, 1H), 6.91 (dd, *J* = 4.0, 2.5 Hz, 1H), 6.87 (d, *J* = 8.5 Hz, 1H), 4.20-4.48 (m, 4H), 3.78 (s, 3H), 3.70 (s, 3H), 2.92 (m, 1H), 1.23 (s, 3H), 1.21 (s, 3H). LC-MS (M+1) 666.0 (4.74 min).

## EXAMPLE 11




5 Methyl [3,5-bis(trifluoromethyl)benzyl][(5'-isopropyl-2'-methoxy-4-pyridin-3-yl)biphenyl-2-yl)methyl]carbamate

A mixture of methyl [3,5-bis(trifluoromethyl)benzyl][(4-iodo-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl]carbamate from Example 10 (0.03 g, 0.045 mmol), potassium carbonate (0.012 g, 0.09 mmol), 3-pyridyl boronic acid (0.0066 g, 0.054 mmol) and a catalytic amount of palladium acetate (1 mg) in 4:1  
 10 acetone/water (2 ml) was heated to and maintained at reflux for 5 h. Acetone was removed and the organic was extracted with methylene chloride (3 x 10 ml). The combined organic layers were washed with brine and dried over sodium sulfate. The title compound was obtained after preparative thin layer chromatography using 3:7 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.90 (m, 1H), 8.65 (m, 1H), 7.90 (m, 1H), 7.72 (s, 1H), 7.70 (m, 1H), 7.60 (dd, *J* = 8.0, 2.5 Hz, 1H), 7.48 (m, 2H), 7.39 (d, *J* = 8.0 Hz, 1H), 7.35 (s, 1H), 7.25 (m, 1H), 7.00 (s, 1H), 6.93 (d, *J* = 8.0 Hz, 1H), 4.20-4.62 (m, 4H), 3.77 (s, 3H), 3.72 (s, 3H), 2.90 (m, 1H), 1.22 (s, 6H). LC-MS (M+1) 617.0 (3.76 min).

15

## EXAMPLE 12




5 Methyl [3,5-bis(trifluoromethyl)benzyl][(4-cyano-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl]carbamate

A mixture of methyl [3,5-bis(trifluoromethyl)benzyl][(4-iodo-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl]carbamate from Example 10 (0.03 g, 0.045 mmol) and CuCN (0.008 g, 0.09 mmol) in DMF (1 ml) was stirred at 100°C overnight. Water (10 ml) was added and the mixture was extracted with EtOAc (3 x 15 ml). The combined EtOAc layers were washed with brine, and dried over sodium sulfate. The title compound was obtained after preparative thin layer chromatography using 2:8 EtOAc/hexane as the eluant. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.72 (s, 1H), 7.74 (m, 1H), 7.64 (d, *J* = 8.0 Hz, 1H), 7.49 (m, 1H), 7.38 (m, 1H), 7.36 (d, *J* = 8.0 Hz, 1H), 7.28 (m, 1H), 6.94 (d, *J* = 2.5 Hz, 1H), 6.91 (m, 1H), 4.26-4.49 (m, 4H), 3.81 (m, 3H), 3.70 (s, 3H), 2.92 (m, 1H), 1.22 (s, 6H). LC-MS (*M*+1) 565.3 (4.42 min).

15

## EXAMPLE 13




5 Methyl [3,5-bis(trifluoromethyl)benzyl][(4-chloro-5'-isopropyl-2'-methoxybiphenyl-2-yl)methyl]carbamate

The title compound was prepared following the procedures from Example 8, Steps A, B, C, D, and E starting from methyl 2-bromo-5-chlorobenzoate. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.74 (s, 1H), 7.47 (s, 1H), 7.38 (m, 1H), 7.32 (s, 1H), 7.31 (dd, *J* = 8.5, 2.5 Hz, 1H), 7.24 (d, *J* = 2.5 Hz, 1H), 7.22 (m, 1H), 6.94 (d, *J* = 2.5 Hz, 1H), 6.87 (d, *J* = 8.5 Hz, 1H), 4.18-4.50 (m, 4H), 3.75 (s, 3H), 3.70 (s, 3H), 2.83 (m, 1H), 1.20 (s, 6H). LC-MS (M+1) 574.3 (4.66 min).

10

## EXAMPLE 14



5 Methyl [3,5-bis(trifluoromethyl)benzyl][(4-chloro-5'-formyl-2'-methoxybiphenyl-2-yl)methyl]carbamate

The title compound was prepared followed the procedures from Example 8, Step A, B, C, D, and E


starting from methyl 2-bromo-5-chlorobenzoate and using 2-methoxy-5-formylphenyl boronic acid for

Step E. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 9.95 (s, 1H), 7.92 (d, *J* = 9.0 Hz, 1H), 7.74 (s, 1H), 7.65 (d, *J* = 2.0

Hz, 1H), 7.48 (m, 1H), 7.38 (dd, *J* = 8.0, 2.0 Hz, 1H), 7.34 (m, 1H), 7.23 (d, *J* = 2.0 Hz, 1H), 7.20 (m,

10 1H), 7.06 (d, *J* = 9.0 Hz, 1H), 4.10-4.58 (m, 4H), 3.82 (s, 3H), 3.70 (s, 3H). LC-MS (M+1) 560.1 (4.13 min).

## EXAMPLE 15



15 2-Iodo-5-(trifluoromethyl)benzonitrile

2-Amino-5-(trifluoromethyl)benzonitrile (3.06 g, 16.45 mmol) was suspended in CH<sub>2</sub>I<sub>2</sub> (36 mL) and *t*-


butyl nitrite (3.9 mL, 32.9 mmol) was added dropwise by syringe. The reaction was heated slowly to 100

20 °C and was maintained at this temperature for 30 minutes. The reaction was then cooled to room

temperature, diluted with hexanes (200 mL), loaded on a silica gel column, and purified with 100%

hexanes to 15% EtOAc/hexanes. The resulting product, 2-iodo-5-(trifluoromethyl)benzonitrile was contaminated with minor impurities which were removed by silica gel chromatography with 25% CH<sub>2</sub>Cl<sub>2</sub>/hexanes. 2-Iodo-5-(trifluoromethyl)benzonitrile was obtained as a white solid.  $R_f = 0.44$  (15% EtOAc/hexanes). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz)  $\delta$  8.10 (d,  $J = 8.5$  Hz, 1H), 7.85 (d,  $J = 1.8$  Hz, 1H), 7.52 (dd,  $J = 8.5, 1.8$  Hz, 1H).

## EXAMPLE 16




10

5'-Isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-carbonitrile

To a solution of 2-iodo-5-(trifluoromethyl)benzonitrile (2.0 g, 6.7 mmol) and (5-isopropyl-2-methoxyphenyl)boronic acid (1.6 g, 8.4 mmol) in dimethyl ethylene glycol (30.4 mL) was added 2M Na<sub>2</sub>CO<sub>3</sub> (6.8 mL), ethanol (9.6 mL), and water (10 mL). The solution was degassed with nitrogen for 2 minutes. Pd(PPh<sub>3</sub>)<sub>4</sub> (774 mg, 0.67 mmol) was added and the solution was degassed with nitrogen again for 2 minutes. The solution was divided equally into two 40 mL microwave tubes. Each tube was degassed with nitrogen for 1 minute, sealed, and placed in a microwave reactor. The wattage was set for 200 W until the temperature reached 150°C and then the temperature was held at 150°C for ten minutes. The tubes were then cooled to room temperature, combined, poured into H<sub>2</sub>O (50 mL), and extracted with EtOAc (100 mL). The organic layer was washed with brine (50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 15% CH<sub>2</sub>Cl<sub>2</sub>/hexanes afforded 5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-carbonitrile as a light yellow oil.  $R_f = 0.65$  (25% EtOAc/hexanes). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz)  $\delta$  7.97 (s, 1H), 7.85 (d,  $J = 8.0$  Hz, 1H), 7.63 (d,  $J = 8.0$  Hz, 1H), 7.31 (dd,  $J = 8.5, 2.0$  Hz, 1H), 7.12 (d,  $J = 2.0$  Hz, 1H), 6.97 (d,  $J = 8.5$  Hz, 1H), 3.82 (s, 3H), 2.93 (m, 1H), 1.27 (d,  $J = 7.0$  Hz, 6H).




## EXAMPLE 17

5 1-[5'-Isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine

5'-Isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-carbonitrile (996.2 mg, 3.12 mmol) was dissolved in Et<sub>2</sub>O (33 mL) and cooled to 0 °C. LAH (12.49 mL of a 1 M solution in Et<sub>2</sub>O, 12.49 mmol) was added dropwise by syringe. After stirring at 0 °C for 10 minutes, the reaction was warmed to room temperature and stirred at room temperature for 6 hours. The reaction was then quenched by slow dropwise addition of 1.5 mL of H<sub>2</sub>O (vigorous evolution of gas), followed by 1.5 mL of 30% NaOH, followed by 3.0 mL of H<sub>2</sub>O. The resulting gelatinous precipitate was washed with 5 x 20 mL of CH<sub>2</sub>Cl<sub>2</sub>; the organic washes were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification of the residue by flash chromatography with 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub> containing 0.1% Et<sub>3</sub>N afforded 1-[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine. R<sub>f</sub> = 0.30 (10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>). LCMS = 324.3 (M+1)<sup>+</sup>.  
 15 <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.77 (s, 1H), 7.55 (d, *J* = 6.8 Hz, 1H), 7.32 (d, *J* = 7.8 Hz, 1H), 7.25 (dd, *J* = 8.3, 2.1 Hz, 1H), 7.00 (d, *J* = 2.1 Hz, 1H), 6.92 (d, *J* = 8.4 Hz, 1H), 3.66-3.74 (m, 5H), 2.91 (m, 1H), 1.26 (d, *J* = 6.9 Hz, 6H).

## 20 EXAMPLE 18




Methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

To a solution of 1-[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine (166 mg, 0.51 mmol) and methyl chloroformate (107  $\mu$ L, 1.39 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added N,N-diisopropylethylamine (485  $\mu$ L, 2.79 mmol). The reaction was stirred at room temperature for twenty minutes and then was poured into  $\text{H}_2\text{O}$  (50 mL). The mixture was extracted with EtOAc (100 mL), and the organic layer was washed with brine (50 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated.

Purification by flash chromatography with 2% to 20% EtOAc/hexanes afforded methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate.  $R_f$  = 0.29 (15% EtOAc/hexanes). LCMS = 382.3 ( $\text{M}+1$ )<sup>+</sup>.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.68 (s, 1H), 7.57 (d,  $J$  = 8.0 Hz, 1H), 7.31 (d,  $J$  = 8.0 Hz, 1H), 7.25 (dd,  $J$  = 8.5, 2.5 Hz, 1H), 6.97 (d,  $J$  = 2.5 Hz, 1H), 6.91 (d,  $J$  = 8.5 Hz, 1H), 4.17 (m, 2H), 3.75 (s, 3H), 3.65 (s, 3H), 2.89 (m, 1H), 1.25 (d,  $J$  = 7.0 Hz, 6H).

## EXAMPLE 19

Methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(trifluoromethyl)benzyl]carbamate


Methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (20 mg, 0.053 mmol) was dissolved in THF (1 mL). 3-(Trifluoromethyl) benzyl bromide (122  $\mu$ L, 0.79 mmol) was added followed by potassium bis(trimethylsilyl)amide (320  $\mu$ L of a 0.5 M solution in toluene, 0.160 mmol). The reaction was stirred for two hours and then poured into  $\text{H}_2\text{O}$  (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 2% to 25%

EtOAc/hexanes afforded methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-

(trifluoromethyl)benzyl]carbamate.  $R_f = 0.67$  (15% EtOAc/hexanes). LCMS = 540.3 ( $M+1$ )<sup>+</sup>. <sup>1</sup>H NMR ( $C_6D_6$ , 500 MHz, 70°C)  $\delta$  7.72 (s, 1H), 7.36 (dd,  $J = 8.0, 1.0$  Hz, 1H), 7.19-7.23 (m, 2H), 7.03-7.07 (m, 3H), 6.86 (m, 2H), 6.57 (d,  $J = 8.5$  Hz, 1H), 4.41 (s, 2H), 4.05 (s, 2H), 3.36 (s, 3H), 3.22 (s, 3H), 2.73 (m, 1H), 1.25 (d,  $J = 6.5$  Hz, 6H).

5

## EXAMPLE 20




10

[3,5-Bis(trifluoromethyl)benzyl]{{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine

1-[5'-Isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine (94.1 mg, 0.291 mmol) was dissolved in THF (4 mL). 3,5-Bis(trifluoromethyl)benzyl bromide (320  $\mu$ L, 1.75 mmol) was added followed by potassium bis(trimethylsilyl)amide (2.33 mL of a 0.5 M solution in toluene, 1.164 mmol). The reaction was stirred at room temperature for 72 hours and then poured into  $H_2O$  (30 mL). The mixture was extracted with EtOAc (75 mL). The organic layer was washed with brine (25 mL), dried over  $Na_2SO_4$ , filtered, and concentrated. Purification by flash chromatography with 0 to 15% EtOAc/hexanes, then a second column with 50%  $CH_2Cl_2$ , then a third column with 25% EtOAc/hexanes afforded [3,5-bis(trifluoromethyl)benzyl]{{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine as a light yellow oil.  $R_f = 0.18$  (40%  $CH_2Cl_2$ /hexanes). LCMS = 550.3 ( $M+1$ )<sup>+</sup>. <sup>1</sup>H NMR ( $CDCl_3$ , 600 MHz)  $\delta$  7.75 (s, 1H), 7.74 (s, 1H), 7.70 (s, 2H), 7.58 (d,  $J = 7.9$  Hz, 1H), 7.34 (d,  $J = 7.8$  Hz, 1H), 7.24 (dd,  $J = 8.5, 2.2$  Hz, 1H), 6.99 (d,  $J = 2.2$  Hz, 1H), 6.90 (d,  $J = 8.5$  Hz, 1H), 3.68-3.76 (m, 4H), 3.68 (s, 3H), 2.88 (m, 1H), 1.22 (d,  $J = 6.4$  Hz, 6H).

20

## EXAMPLE 21




5

1-([5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl)-3-nitro-5-(trifluoromethyl)benzylamine

To a solution of 3-nitro-5-(trifluoromethyl)benzaldehyde (366 mg, 1.67 mmol) in MeOH (4 mL) was added a solution of 1-([5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine (270 mg, 0.84 mmol) in MeOH (4 mL) followed by addition of NaCNBH<sub>3</sub> (105 mg, 1.67 mmol) and one drop of acetic acid. The reaction was stirred for 24 hours at room temperature and then poured into H<sub>2</sub>O (30 mL). The mixture was extracted with EtOAc (60 mL), and the organic extracts were washed with brine (20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 2% to 25% EtOAc/hexanes afforded 1-([5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl)-3-nitro-5-(trifluoromethyl)benzylamine as a yellow oil. R<sub>f</sub> = 0.45 (15% EtOAc/hexanes). LCMS = 527.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.33 (s, 1H), 8.27 (s, 1H), 7.85 (s, 1H), 7.74 (s, 1H), 7.56 (d, J = 8.0 Hz, 1H), 7.33 (d, J = 8.0 Hz, 1H), 7.22 (dd, J = 8.5 Hz, 2.0 Hz, 1H), 6.97 (d, J = 2.0 Hz, 1H), 6.89 (d, J = 8.5 Hz, 1H), 3.67-3.78 (m, 4H), 3.70 (s, 3H), 2.87 (m, 1H), 1.25 (d, J = 7.0 Hz, 6H).

15

## EXAMPLE 22




5 Methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}{[3-nitro-5-(trifluoromethyl)benzyl]carbamate

To a solution of {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}{[3-nitro-5-(trifluoromethyl)benzyl]amine (428 mg, 0.81 mmol) and methyl chloroformate (126  $\mu$ L, 1.39 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added N,N-diisopropylethylamine (566  $\mu$ L, 3.25 mmol). The reaction was stirred at

10 room temperature for twenty minutes and then was poured into  $\text{H}_2\text{O}$  (25 mL). The mixture was extracted with EtOAc (70 mL), and the organic extracts were washed with brine (25 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 2% to 20% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}{[3-nitro-5-(trifluoromethyl)benzyl]carbamate.  $R_f$  = 0.37 (25% EtOAc/hexanes). LCMS = 585.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR

15 ( $\text{C}_6\text{D}_6$ , 500 MHz, 70°C)  $\delta$  8.03 (s, 1H), 7.78 (s, 1H), 7.64 (s, 1H), 7.37 (s, 1H), 7.33 (d,  $J$  = 8.0 Hz, 1H), 7.04-7.06 (m, 2H), 6.83 (s, 1H), 6.60 (d,  $J$  = 8.4 Hz, 1H), 4.37 (bs, 2H), 3.93 (bs, 2H), 3.36 (s, 3H), 3.26 (s, 3H), 2.73 (m, 1H), 1.15 (d,  $J$  = 6.8 Hz, 6H).

## EXAMPLE 23


5 3-Nitro-5-(trifluoromethyl)benzaldehydeStep A: [3-Nitro-5-(trifluoromethyl)phenyl]methanol

To a solution of 3-nitro-5-(trifluoromethyl)benzoic acid (2.0 g, 8.5 mmol) in THF (100 mL) was added borane-THF (17 mL of a 1 M solution in THF, 17 mmol). The reaction was stirred at 60°C for 24 hours and then poured into H<sub>2</sub>O (50 mL). The resultant mixture was extracted with EtOAc (100 mL) and the  
10 organic layer was washed with brine (50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 50% EtOAc/hexanes afforded [3-nitro-5-(trifluoromethyl)phenyl]methanol. *R<sub>f</sub>* = 0.35 (75% EtOAc/hexanes). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) 8.44 (s, 1H), 8.40 (s, 1H), 7.98 (s, 1H), 4.91 (s, 2H), 2.07 (bs, 1H).

Step B: 3-Nitro-5-(trifluoromethyl)benzaldehyde

15 A solution of [3-nitro-5-(trifluoromethyl)phenyl]methanol (400 mg, 1.88 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was cooled to 0°C and then Dess-Martin periodinane (1.59 g, 3.76 mmol) was added. The reaction was slowly warmed to room temperature. After stirring at room temperature for twenty minutes, the reaction was poured into 1N NaOH (25 mL). The mixture was extracted with EtOAc (50 mL) and the organic  
20 extracts were washed with brine (25 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 0% to 15% EtOAc/hexanes afforded 3-nitro-5-(trifluoromethyl)benzaldehyde. *R<sub>f</sub>* = 0.39 (15% EtOAc/hexanes). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 10.18 (s, 1H), 8.91 (s, 1H), 8.74 (s, 1H), 8.48 (s, 1H).

## EXAMPLE 24

5    3-Iodo-5-(trifluoromethyl)benzaldehydeStep A: [3-Amino-5-(trifluoromethyl)phenyl]methanol

To a solution of [3-nitro-5-(trifluoromethyl)phenyl]methanol (1.8 g, 8.0 mmol) (Step A, Example 23) in THF (36 mL) was added concentrated HCl (18 mL) and  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$  (8.7 g, 38.8 mmol). The reaction was stirred at room temperature for 24 hours and then poured slowly into saturated  $\text{NaHCO}_3$  (250 mL).

- 10    After neutralization was complete, the reaction was extracted with EtOAc (3 x 50 mL). The organic extracts were washed with brine (75 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 50% EtOAc/hexanes afforded 1.5 g (quantitative) of [3-amino-5-(trifluoromethyl)phenyl]methanol.  $R_f = 0.35$  (50% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  6.98 (s, 1H), 6.84 (s, 1H), 6.81 (s, 1H), 4.65 (s, 2H), 3.87 (bs, 1H).

15    Step B: [3-Iodo-5-(trifluoromethyl)phenyl]methanol

[3-amino-5-(trifluoromethyl)phenyl]methanol (500 mg, 5.2 mmol) was suspended in  $\text{CH}_2\text{I}_2$  (5 mL) and *t*-butyl nitrite (622  $\mu\text{L}$ , 5.2 mmol) was added dropwise by syringe. The reaction was heated slowly to  $100^\circ\text{C}$  and was maintained at this temperature for 30 minutes. The reaction was then cooled to room temperature, diluted with hexanes (50 mL), loaded on a silica gel column, and purified with 100% hexanes to 25% EtOAc/hexanes to afford [3-iodo-5-(trifluoromethyl)phenyl]methanol.  $R_f = 0.19$  (15% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.91 (s, 1H), 7.87 (s, 1H), 7.59 (s, 1H), 4.73 (d,  $J = 4.9$  Hz, 2H), 1.86 (t,  $J = 5.6$  Hz, 1H).


Step C: 3-Iodo-5-(trifluoromethyl)benzaldehyde

- 25    A solution of [3-iodo-5-(trifluoromethyl)phenyl]methanol (287 mg, 0.95 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was cooled to  $0^\circ\text{C}$  and then Dess-Martin periodinane (604 mg, 1.42 mmol) was added. The reaction was slowly warmed to room temperature. After stirring at room temperature for twenty minutes, the reaction was poured into 1N NaOH (25 mL). The mixture was extracted with EtOAc (50 mL) and the organic extracts were washed with brine (25 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 0% to 10% EtOAc/hexanes afforded 3-iodo-5-(trifluoromethyl)benzaldehyde.

$R_f = 0.54$  (15% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 600 MHz)  $\delta$  9.98 (s, 1H), 8.39 (s, 1H), 8.19 (s, 1H), 8.09 (s, 1H).

## EXAMPLE 25

5

3-Bromo-5-(trifluoromethyl)benzaldehydeStep A: [3-Bromo-5-(trifluoromethyl)-phenyl]methanol


- 10 [3-amino-5-(trifluoromethyl)phenyl]methanol (900 mg, 4.71 mmol) (Step A, Example 24) was suspended in  $\text{CHBr}_3$  (9 mL), and *t*-butyl nitrite (600  $\mu\text{L}$ , 5.04 mmol) was added dropwise by syringe. The reaction was heated slowly to  $80^\circ\text{C}$  and was maintained at this temperature for 10 minutes. The reaction was then cooled to room temperature, diluted with hexanes (50 mL), loaded on a silica gel column, and purified with 100% hexanes to 20% EtOAc/hexanes (2 columns) to afford [3-bromo-5-
- 15 (trifluoromethyl) phenyl]methanol.  $R_f = 0.31$  (25% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.71 (s, 1H), 7.68 (s, 1H), 7.56 (s, 1H), 4.76 (d,  $J = 5.5$  Hz, 2H), 1.86 (t,  $J = 5.7$  Hz, 1H).

Step B: 3-Bromo-5-(trifluoromethyl)benzaldehyde

- A solution of [3-bromo-5-(trifluoromethyl)phenyl]methanol (409 mg, 1.61 mmol) in  $\text{CH}_2\text{Cl}_2$  (25 mL) was cooled to  $0^\circ\text{C}$  and then Dess-Martin periodinane (1.02 g, 2.41 mmol) was added. The reaction was
- 20 stirred at  $0^\circ\text{C}$  for 30 minutes and then warmed to room temperature. After stirring at room temperature for thirty minutes, the reaction was poured into 1N NaOH (25 mL). The mixture was extracted with EtOAc (100 mL), and the organic extracts were washed with 1N NaOH (25 mL), then brine (25 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 25% EtOAc/hexanes afforded 3-bromo-5-(trifluoromethyl)benzaldehyde.  $R_f = 0.60$  (25% EtOAc/hexanes).  $^1\text{H}$
- 25 NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  10.02 (s, 1H), 8.20 (s, 1H), 8.07 (s, 1H), 8.02 (s, 1H).



## EXAMPLE 26

5     3-(Methylthio)-5-(trifluoromethyl)benzaldehydeStep A: [3-(Methylthio)-5-(trifluoromethyl)phenyl] methanol


To a solution of [3-amino-5-(trifluoromethyl)phenyl]methanol (319 mg, 1.67 mmol) (Step A, Example 24) in  $\text{CHCl}_3$  (5 mL) was added dimethyl disulfide (296  $\mu\text{L}$ , 3.34 mmol). Next, *t*-butyl nitrite (285  $\mu\text{L}$ , 2.4 mmol) was added dropwise by syringe. The reaction was stirred at room temperature for two hours  
10     and then poured into  $\text{H}_2\text{O}$  (15 mL). The mixture was extracted with EtOAc (50 mL), washed with brine (15 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 0% to 50% EtOAc/hexanes afforded [3-(methylthio)-5-(trifluoromethyl)phenyl]-methanol. The [3-(methylthio)-5-(trifluoromethyl)phenyl]methanol was contaminated with minor impurities that were removed after the next step.  $R_f = 0.53$  (50% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.63 (s, 1H), 7.40 (s, 1H), 7.35 (s, 1H), 4.71 (s, 2H), 2.51 (s, 3H).  
15

Step B: 3-(Methylthio)-5-(trifluoromethyl)benzaldehyde

A solution of [3-(methylthio)-5-(trifluoromethyl)phenyl]methanol (200 mg, 0.99 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) was cooled to  $0^\circ\text{C}$  and then Dess-Martin periodinane (573 mg, 1.35 mmol) was added. The reaction was slowly warmed to room temperature. After stirring at room temperature for twenty minutes, the  
20     reaction was poured into 1N NaOH (25 mL). The mixture was extracted with EtOAc (75 mL). The organic extracts were washed with brine (25 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 1% to 10% EtOAc/hexanes afforded 3-(methylthio)-5-(trifluoromethyl)benzaldehyde.  $R_f = 0.55$  (25% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  10.02 (s, 1H), 7.87 (s, 1H), 7.84 (s, 1H), 7.68 (s, 1H), 2.58 (s, 3H).

25

## EXAMPLE 27

5 3-Methoxy-5-(trifluoromethyl)benzaldehydeStep A: 1-Iodo-3-methoxy-5-(trifluoromethyl)benzene

3-methoxy-5-(trifluoromethyl)aniline (100 mg, 0.52 mmol) was dissolved in  $\text{CHCl}_3$  (5.2 mL) and *t*-butyl nitrite (124  $\mu\text{L}$ , 1.05 mmol) was added dropwise by syringe. Iodine (266 mg, 1.05 mmol) was added and then the reaction was slowly heated to  $50^\circ\text{C}$  and was maintained at this temperature for an hour and thirty  
10 minutes. The reaction was then cooled to room temperature and poured into aq.  $\text{NaHSO}_3$  (50 mL). The mixture was extracted with EtOAc (50 mL) and the organic extracts were washed with aq.  $\text{NaHSO}_3$  (3 X 50 mL) and brine (20 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 1% to 15% EtOAc/hexanes afforded 1-iodo-3-methoxy-5-(trifluoromethyl)benzene.  $R_f = 0.75$  (25% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.53 (s, 1H), 7.41 (s, 1H), 7.09 (s, 1H),  
15 3.82 (s, 3H).

Step B: 3-Methoxy-5-(trifluoromethyl)benzonitrile


To a solution of 1-iodo-3-methoxy-5-(trifluoromethyl)benzene (200 mg, 0.66 mmol) in DMF (2 mL) was added  $\text{CuCN}$  (300 mg, 5.0 mmol). The reaction was stirred at  $100^\circ\text{C}$  for 24 hours and then poured into aq.  $\text{NH}_3$  (40 mL). The mixture was extracted with EtOAc (70 mL) and the organic extracts were washed  
20 with brine (25 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 5% to 80% EtOAc/hexanes afforded 3-methoxy-5-(trifluoromethyl)benzonitrile.  $R_f = 0.64$  (50% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 600 MHz)  $\delta$  7.47 (s, 1H), 7.34 (s, 1H), 7.31 (s, 1H), 3.89 (s, 3H).

Step C: 3-Methoxy-5-(trifluoromethyl)benzaldehyde

A mixture of 3-methoxy-5-(trifluoromethyl)benzonitrile (88 mg, 0.43 mmol) and  $\text{PtO}_2$  (9.8 mg, 0.043  
25 mmol) in 88% formic acid (651  $\mu\text{L}$ ) was heated to  $60^\circ\text{C}$ . The reaction was stirred at this temperature for thirty minutes. The reaction was then cooled to room temperature, diluted with hexanes (5 mL), loaded on a silica gel column, and purified with 1% to 15% EtOAc/hexanes to afford 3-methoxy-5-(trifluoromethyl)benzaldehyde.  $R_f = 0.56$  (25% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  10.01 (s, 1H), 7.71 (s, 1H), 7.56 (s, 1H), 7.39 (s, 1H), 3.92 (s, 3H).

30


## EXAMPLE 28



5 *N*-[3,5-Bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}methanesulfonamide

To a solution of [3,5-bis(trifluoromethyl)benzyl]{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine (8.3 mg, 0.015 mmol) (Example 20) and methanesulfonyl chloride (10.5  $\mu$ L, 0.135 mmol) in  $\text{CH}_2\text{Cl}_2$  (500  $\mu$ L) was added *N,N*-diisopropylethylamine (47.4  $\mu$ L, 0.273 mmol). The reaction  
 10 was stirred at room temperature for twenty minutes and then was poured into  $\text{H}_2\text{O}$  (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 50%  $\text{CH}_2\text{Cl}_2$ /hexanes afforded *N*-[3,5-bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}methanesulfonamide.  $R_f$  = 0.20 (50%  $\text{CH}_2\text{Cl}_2$ /hexanes). LCMS = 628.3 ( $\text{M}+1$ )<sup>+</sup>.  $^1\text{H}$  NMR  
 15 ( $\text{CDCl}_3$ , 600 MHz)  $\delta$  7.69 (s, 1H), 7.55 (s, 1H), 7.50 (d,  $J$  = 8.4 Hz, 1H), 7.47 (s, 2H), 7.28-7.30 (m, 2H), 6.94 (d,  $J$  = 9.0 Hz, 1H), 6.92 (d,  $J$  = 1.8 Hz, 1H), 4.44 (d,  $J$  = 15.0, 1H), 4.32 (d,  $J$  = 15.0 Hz, 1H), 4.19 (d,  $J$  = 15.6 Hz, 1H), 4.14 (d,  $J$  = 15.6 Hz, 1H), 3.72 (s, 3H), 2.89 (m, 1H), 2.69 (s, 3H), 1.21-1.25 (m, 6H).


## EXAMPLE 29



5 [3,5-Bis(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} formamide

A solution of [3,5-bis(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} amine (88 mg, 0.43 mmol) (Example 20) in 80% formic acid (3 mL) was heated to and maintained at reflux for 24 hours. The reaction was then cooled to room temperature, poured into  
 10 saturated NaHCO<sub>3</sub> (30 mL), and, after neutralization was complete, extracted with EtOAc (25 mL). The organic extracts were washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 15% EtOAc/hexanes afforded [3,5-bis(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} formamide. R<sub>f</sub> = 0.20 (15% EtOAc/hexanes). LCMS = 578.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>, 600  
 15 MHz, rotamers present, ratio of major:minor ~3:1. Data for major rotamer given.) δ 6.86-8.31 (m, 10H), 4.24-4.47 (m, 4H), 3.71 (s, 3H), 2.87 (m, 1H), 1.27 (m, 6H).


## EXAMPLE 30



5 *O*-Methyl [3,5-bis(trifluoromethyl)benzyl]{{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}thiocarbamate

In a dry flask was placed CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and thiophosgene (20 μL, 0.262 mmol). [3,5-Bis(trifluoromethyl)benzyl]{{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine (15 mg, 0.027 mmol) (Example 20) was added to the flask as a solution in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). The reaction was stirred for several minutes at room temperature, and then a slurry of NaOMe (133 mg, 2.46 mmol) in MeOH (2 mL) was added. The reaction was stirred at room temperature for 20 minutes and then poured into saturated NH<sub>4</sub>Cl (25 mL). The mixture was extracted with EtOAc (50 mL) and the organic extract was washed with brine (25 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified by preparative thin layer chromatography (40% CH<sub>2</sub>Cl<sub>2</sub>/hexanes) to yield *O*-methyl[3,5-bis(trifluoromethyl)benzyl]{{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}thiocarbamate. *R*<sub>f</sub> = 0.41 (40% CH<sub>2</sub>Cl<sub>2</sub>/hexanes). LCMS = 624.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) (rotamers present; doubling of some peaks; partial data given) δ 7.55-7.73 (m, 3 H), 7.22-7.36 (m, 4 H), 6.87-6.93 (m, 2H), 5.11 (s), 5.02 (d, *J* = 15.6 Hz), 4.85 (d, *J* = 15.6 Hz), 4.70 (d, *J* = 15.8 Hz), 4.50 (s), 4.49 (d, *J* = 15.8 Hz), 4.09 (s, minor rotamer), 4.01 (s, major rotamer), 3.68 (s, minor rotamer), 3.67 (s, major rotamer), 2.87 (m, 1H), 1.19-1.22 (m, 6H).


## EXAMPLE 31



5 *N*-[3,5-Bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}-*N'*-methylurea

To a solution of [3,5-bis(trifluoromethyl)benzyl]{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine (17.4 mg, 0.031 mmol) (Example 20) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added methyl isocyanate (3 drops added by pipette, ~ 15 μL). The reaction was stirred at room temperature for 30 minutes, and  
 10 then MeOH (1 mL) was added and the reaction was stirred for 5 minutes. The reaction was diluted with EtOAc (30 mL) and washed with water and brine (15 mL each). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography (40% EtOAc/hexanes) gave *N*-[3,5-bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}-*N'*-methylurea. *R<sub>f</sub>* = 0.32 (40% EtOAc/hexanes). LCMS = 607.4 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz)  
 15 δ 7.74 (s, 1H), 7.66 (s, 2H), 7.59 (d, *J* = 7.7 Hz, 1H), 7.43 (s, 1H), 7.32 (d, *J* = 7.8 Hz, 1H), 7.24 (dd, *J* = 8.7, 2.3 Hz, 1H), 6.90-6.92 (m, 2H), 4.85 (d, *J* = 15.8 Hz, 1H), 4.50 (d, *J* = 15.8 Hz, 1H), 4.45 (m, 1H), 4.39 (d, *J* = 17.9 Hz, 1H), 4.02 (d, *J* = 17.8 Hz, 1H), 3.71 (s, 3H), 2.87 (m, 1H), 2.76 (d, *J* = 4.3 Hz, 3H), 1.20-1.22 (m, 6H).

## EXAMPLE 32



5


*N*-[3,5-Bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)-biphenyl-2-yl]methyl}urea

[3,5-Bis(trifluoromethyl)benzyl]{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine (40.3 mg, 0.073 mmol) (Example 20) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) and the solution was cooled to -78 °C. Chlorosulfonyl isocyanate (32 μL, 0.367 mmol) was added dropwise by syringe. The reaction was stirred at -78 °C for 15 minutes, and then at 0 °C for 20 minutes. The reaction was then poured into H<sub>2</sub>O (15 mL) and saturated NaHCO<sub>3</sub> (15 mL) was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 25 mL). The organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

Purification of the residue by flash chromatography with 50% EtOAc/hexanes afforded *N*-[3,5-bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}urea.

*R*<sub>f</sub> = 0.24 (40% EtOAc/hexanes). LCMS = 593.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.75 (s, 1H), 7.66 (s, 2H), 7.60 (d, *J* = 7.7 Hz, 1H), 7.47 (s, 1H), 7.33 (d, *J* = 8.0 Hz, 1H), 7.25 (dd, *J* = 8.5, 2.3 Hz, 1H), 6.92 (d, *J* = 2.4 Hz, 1H), 6.90 (d, *J* = 8.5 Hz, 1H), 4.80 (d, *J* = 15.8 Hz, 1H), 4.53 (s, 2H), 4.48 (d, *J* = 15.8 Hz, 1H), 4.45 (d, *J* = 17.8 Hz, 1H), 4.07 (d, *J* = 17.8 Hz, 1H), 3.69 (s, 3H), 2.87 (m, 1H), 1.20-1.22 (m, 6H).

## EXAMPLE 33



5 Methyl[3-amino-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate


A solution of methyl {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} [3-nitro-5-(trifluoromethyl)benzyl] carbamate (174 mg, 0.03 mmol) (Example 22) and PtO<sub>2</sub> (10 mg) in THF (10 mL) was placed under hydrogen. The reaction was stirred at room temperature under hydrogen for two hours.

- 10 The reaction was then diluted with hexanes (25 mL), loaded on a silica gel column, and purified with 15% EtOAc/hexanes to afford methyl[3-amino-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate. *R<sub>f</sub>* = 0.31 (25% EtOAc/hexanes). LCMS = 555.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.59-6.28 (m, 9H), 4.48-3.62 (m, 12H), 2.87 (m, 1H), 1.22 (broad singlet, 6H).

15



## EXAMPLE 34



5

Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-[(methylsulfonyl)amino]-5-(trifluoromethyl)benzyl]carbamate


To a solution of methyl[3-amino-5-(trifluoromethyl)benzyl][[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]carbamate (8 mg, 0.014 mmol) (Example 33) and

10 methanesulfonyl chloride (3.3  $\mu$ L, 0.043 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 mL) was added N,N-diisopropylethylamine (15  $\mu$ L, 0.086 mmol). The reaction was stirred at room temperature for thirty minutes and then was poured into  $\text{H}_2\text{O}$  (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by

15 preparative thin layer chromatography with 25% EtOAc/hexanes followed by a preparative thin layer chromatography in 2:2:1 ( $\text{CH}_2\text{Cl}_2$ : Hex:  $\text{Et}_2\text{O}$ ) and then a third and final purification by silica gel chromatography with 2% to 25% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-[(methylsulfonyl)amino]-5-(trifluoromethyl)benzyl]carbamate.  $R_f = 0.40$  (25% EtOAc/hexanes).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 500 MHz,  $70^\circ\text{C}$ )  $\delta$  7.69 (s, 1H), 7.52 (s, 1H), 7.38-7.25 (m, 3H), 7.08-7.04 (m, 2H), 6.86 (s, 1H), 6.30 (d,  $J = 8.5$  Hz, 1H), 4.40 (broad singlet, 2H), 4.04-3.99 (m, 2H), 3.33 (s, 3H), 3.26 (s, 3H), 2.74 (m, 1H), 2.67 (s, 3H), 1.22 (d,  $J = 6.5$  Hz, 6H).

20


## EXAMPLE 35



5 Methyl[3-(dimethylamino)-5-(trifluoromethyl)benzyl][[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

Methyl[3-amino-5-(trifluoromethyl)benzyl][[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (8 mg, 0.014 mmol) (Example 33) was dissolved in THF (1 mL). MeI (3.6  $\mu$ L, 0.056 mmol) was added followed by potassium bis(trimethylsilyl)amide (112  $\mu$ L of a 0.5 M solution in toluene, 0.056 mmol). The reaction was stirred at room temperature for two hours and then poured into H<sub>2</sub>O (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 5% to 25% EtOAc/hexanes afforded methyl[3-(dimethylamino)-5-(trifluoromethyl)benzyl][[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate.  $R_f$  = 0.53 (25% EtOAc/hexanes). LCMS = 583.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz)  $\delta$  7.56-6.44 (m, 9H), 4.49-4.13 (m, 4H), 3.73-3.65 (m, 6H), 2.92-2.85 (m, 7H), 1.21(broad singlet, 6H).

## EXAMPLE 36




5 Methyl[3-cyano-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

To a solution of methyl[3-iodo-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Example 119, 508 mg, 0.76 mmol) in DMF (40 mL) was added CuCN (1.7 g, 19.0 mmol). The reaction was stirred at 100°C for 30 hours. The reaction was  
 10 poured into aq. NH<sub>3</sub> (40 mL) and extracted with EtOAc (3 x 40 mL). All the collected organic extracts were washed with brine (40 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 5% to 80% EtOAc/hexanes afforded methyl[3-cyano-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate. R<sub>f</sub> = 0.55 (20% EtOAc/hexanes). LCMS = 565.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C) δ 7.63 (s, 1H), 7.43-7.38 (m, 3H), 7.17-7.13 (m, 3H), 6.86 (s, 1H), 6.74 (d, J = 8.5 Hz, 1H), 4.37 (broad singlet, 2H), 4.08 (broad singlet, 2H), 3.45 (s, 3H), 3.40 (s, 3H), 2.80 (m, 1H), 1.18 (d, J = 7.0 Hz, 6H).

15


## EXAMPLE 37



5 Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}{[3-pyridin-4-yl-5-(trifluoromethyl)benzyl]carbamate

Methyl[3-iodo-5-(trifluoromethyl)benzyl]{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (Example 119, 20 mg, 0.03 mmol) and pyridine-4-boronic acid (4.6 mg, 0.04 mmol) were placed in a microwave tube in a microwave tube, and dissolved with dimethyl ethylene glycol (170  $\mu$ L). Next, 2M Na<sub>2</sub>CO<sub>3</sub> (38  $\mu$ L), ethanol (42  $\mu$ L), and water (56  $\mu$ L) were added. The solution was degassed with nitrogen for 2 minutes. Pd(PPh<sub>3</sub>)<sub>4</sub> (3.5 mg, 0.003 mmol) was added and the solution was degassed with nitrogen again for 2 minutes. The solution was sealed and placed in a microwave reactor. The wattage was set for 60 W until the temperature reached 150°C and then the temperature was held at 150°C for twenty minutes. The mixture was then cooled to room temperature, poured into H<sub>2</sub>O (10 mL), and extracted with EtOAc (30 mL). The organic layer was washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 5% to 80% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-pyridin-4-yl-5-(trifluoromethyl)benzyl]carbamate. R<sub>f</sub> = 0.40 (75% EtOAc/hexanes). LCMS = 617.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C)  $\delta$  8.54 (broad singlet, 2H), 7.70 (s, 1H), 7.51 (s, 1H), 7.38 (s, 1H), 7.35 (d, J = 8.0, 1H), 7.28 (s, 1H), 7.06 (d, J = 8.0, 1H), 7.02 (dd, J = 2.0 Hz, 8.5, 1H), 6.89 (m, 2H), 6.83 (d, J = 2.0, 1H), 6.58 (d, J = 8.5, 1H), 4.45 (broad singlet, 2H), 4.13 (broad singlet, 2H), 3.38 (s, 3H), 3.23 (s, 3H), 2.65 (m, 1H), 1.12 (d, J = 7.0 Hz, 6H).

## EXAMPLE 38




5 Methyl[3-chloro-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate

To a solution of  $\text{CuCl}_2$  (11.9 mg, 0.09 mmol) in acetonitrile (1 mL) was added *t*-butyl nitrite (11  $\mu\text{L}$ , 0.092 mmol). Next, the solution was cooled to  $0^\circ\text{C}$  and methyl[3-amino-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate (41 mg, 0.074 mmol)

- 10 (Example 33) was added as a solution in acetonitrile (1.5 mL). The reaction was slowly warmed to room temperature. After stirring at room temperature for 2 hours, the mixture was diluted with EtOAc (30 mL) and poured into 20% HCl (5 mL). The organic extracts were washed with 20% HCl (2 x 5 mL),  $\text{H}_2\text{O}$  (10 mL), brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 5% to 80% EtOAc/hexanes afforded methyl[3-chloro-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-
- 15 methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate.  $R_f = 0.65$  (15% EtOAc/hexanes). LCMS = 574.3 ( $\text{M}+1$ ) $^+$ .  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 500 MHz,  $70^\circ\text{C}$ )  $\delta$  7.68 (s, 1H), 7.36 (d,  $J = 7.5$  Hz, 1H), 7.30 (s, 1H), 7.08-7.05 (m, 4H), 6.83 (d,  $J = 2.5$  Hz, 1H), 6.60 (d,  $J = 8.5$  Hz, 1H), 4.35 (broad singlet, 3H), 4.01 (broad singlet, 1H), 3.36 (s, 3H), 3.26 (s, 3H), 2.74 (m, 1H), 1.16 (d,  $J = 7.0$  Hz, 6H).


## EXAMPLE 39



5 Methyl[3-(aminocarbonyl)-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

A solution of methyl[3-cyano-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (35.0 mg, 0.06 mmol) (Example 36) in DMSO (248  $\mu$ L) was cooled to 0°C. Next, K<sub>2</sub>CO<sub>3</sub> (17.4 mg, 0.05 mmol) was added followed by 30% H<sub>2</sub>O<sub>2</sub> (42.2  $\mu$ L), and the reaction was slowly warmed to room temperature. After stirring at room temperature for 15 minutes, the mixture was diluted with EtOAc (50 mL) and poured into H<sub>2</sub>O (15 mL). The organic extract was washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 5% to 80% EtOAc/hexanes afforded methyl[3-(aminocarbonyl)-5-(trifluoromethyl)benzyl] {[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate. R<sub>f</sub> = 0.45 (20% EtOAc/hexanes). LCMS = 583.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C)  $\delta$  7.75 (s, 1H), 7.68 (s, 2H), 7.35 (s, 1H), 7.34 (s, 1H), 7.08-7.04 (m, 2H), 6.87 (d, *J* = 2.5 Hz, 1H), 6.62 (d, *J* = 8.5 Hz, 1H), 5.08 (broad singlet, 2H), 4.41 (broad singlet, 2H), 4.07 (broad singlet, 2H), 3.37 (s, 3H), 3.29 (s, 3H), 2.75 (m, 1H), 1.16 (d, *J* = 7.5 Hz, 6H).


## EXAMPLE 40



5 Methyl[3-formyl-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate

A mixture of methyl[3-cyano-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate (150 mg, 0.26 mmol) (Example 36) and PtO<sub>2</sub> (30 mg, 0.132 mmol) in 88% formic acid (2 mL) was heated to 60°C. The reaction was stirred at this temperature  
 10 for 4 hours. The reaction was then cooled to room temperature, diluted with hexanes (5 mL), loaded on a silica gel column, and purified with 5% to 25% EtOAc/hexanes to afford methyl[3-formyl-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate. R<sub>f</sub> = 0.34 (25% EtOAc/hexanes). LCMS = 568.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C) δ 9.40 (s, 1H), 7.67 (broad singlet, 1H), 7.64 (s, 1H), 7.50 (broad singlet, 1H), 7.39 (broad singlet, 1H), 7.35 (d, J = 7.5 Hz, 1H), 7.07-7.03 (m, 2H), 6.85 (d, J = 2.0 Hz, 1H), 6.60 (d, J = 8.5 Hz, 1H), 4.40 (broad singlet, 2H), 4.03 (broad singlet, 2H), 3.38 (s, 3H), 3.26 (s, 3H), 2.73 (m, 1H), 1.15 (d, J = 7.0 Hz, 6H).

## EXAMPLE 41




5 3-[[[5'-Isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]-  
(methoxycarbonyl)amino]methyl]-5-(trifluoromethyl)benzoic acid

To a solution of methyl[3-formyl-5-(trifluoromethyl)benzyl]{{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl} carbamate (37.0 mg, 0.065 mmol) (Example 40) in THF (123  $\mu$ L) was added *t*-butanol (360  $\mu$ L), 2-methyl 2-butene (123  $\mu$ L), and a solution of NaClO<sub>2</sub> (12.9 mg, 0.014 mmol) and NaH<sub>2</sub>PO<sub>4</sub> (19.7 mg, 0.014 mmol) in H<sub>2</sub>O (152  $\mu$ L). The reaction was stirred vigorously for  
 10 two hours at room temperature, diluted with EtOAc (30 mL), and poured into 1N HCl (10 mL). The aqueous layer was extracted with EtOAc (3 x 20 mL) and the combined organic extracts were washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 0.1% acetic acid in 8% MeOH/CH<sub>2</sub>Cl<sub>2</sub> afforded 3-[[[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]-(methoxycarbonyl)amino]methyl]-5-(trifluoromethyl)benzoic  
 15 acid. *R*<sub>f</sub> = 0.35 (10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>). LCMS = 584.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C)  $\delta$  8.28 (broad singlet, 1H), 8.04 (broad singlet, 1H), 7.68 (broad singlet, 1H), 7.37-7.34 (m, 2H), 7.07-7.05 (m, 2H), 6.87 (s, 1H), 6.62 (d, *J* = 8.0 Hz, 1H), 4.43 (broad singlet, 2H), 4.41 (broad singlet, 2H), 3.40 (s, 3H), 3.29 (s, 3H), 2.75 (m, 1H), 1.17 (d, *J* = 6.5 Hz, 6H).

20



## EXAMPLE 42




5 Methyl[3-(difluoromethyl)-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate

A solution of methyl[3-formyl-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate (24.7 mg, 0.044 mmol) (Example 40) in CH<sub>2</sub>Cl<sub>2</sub> (300 μL) was cooled to 0°C. DAST (92.8 μL, 9.6 mmol) was added and the reaction was slowly warmed to  
 10 room temperature. After stirring at room temperature for 24 hours, the reaction was diluted with EtOAc (30 mL), and poured into NaHCO<sub>3</sub> (10 mL). The organic extract was washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 5% to 25% EtOAc/hexanes afforded methyl[3-(difluoromethyl)-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate. R<sub>f</sub> = 0.42 (25% EtOAc/hexanes) LCMS = 590.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C) δ 7.67 (broad singlet, 1H), 7.40 (s, 1H), 7.34 (d, J = 7.5 Hz, 1H), 7.29 (broad singlet, 1H), 7.22 (broad singlet, 1H), 7.07-7.03 (m, 2H), 6.84 (d, J = 2.0 Hz, 1H), 6.59 (d, J = 8.5 Hz, 1H), 5.97 (t, J = 56 Hz, 1H), 4.39 (broad singlet, 2H), 4.01 (broad singlet, 2H), 3.34 (s, 3H), 3.24 (s, 3H), 2.73 (m, 1H), 1.15 (d, J = 7.0 Hz, 6H).

20


## EXAMPLE 43



5 Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(trifluoromethyl)-5-vinylbenzyl]carbamate

A suspension of  $\text{Ph}_3\text{PCH}_2\text{Br}$  (53.4 mg, 0.15 mmol) in THF (500  $\mu\text{L}$ ) was cooled to  $0^\circ\text{C}$ . potassium bis(trimethylsilyl)amide (256  $\mu\text{L}$  of a 0.5 M solution in toluene, 0.128 mmol) was added slowly and the reaction was kept at  $0^\circ\text{C}$ . After fifteen minutes, methyl[3-formyl-5-(trifluoromethyl)benzyl][5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl carbamate (24.2 mg, 0.043 mmol) (Example 40) was dissolved in THF (1 mL) and was added via cannula to the reaction which was then slowly warmed to room temperature. After fifteen minutes at room temperature, the reaction was diluted with EtOAc (30 mL), and poured into  $\text{NaHCO}_3$  (10 mL). The organic extract was washed with brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 5% to 25% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(trifluoromethyl)-5-vinylbenzyl]carbamate.  $R_f = 0.42$  (25% EtOAc/hexanes) LCMS = 566.2 ( $\text{M}+1$ )<sup>+</sup>.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 500 MHz,  $70^\circ\text{C}$ )  $\delta$  7.68 (broad singlet, 1H), 7.36 (s, 1H), 7.34 (d,  $J = 8.0$  Hz, 1H), 7.15 (broad singlet, 1H), 7.10 (d,  $J = 10.5$  Hz, 1H), 7.03 (d,  $J = 8.0$  Hz, 1H), 7.01 (dd,  $J = 8.5$ , 2.0 Hz, 1H), 6.81 (d,  $J = 2.0$  Hz, 1H), 6.54 (d,  $J = 8.5$  Hz, 1H), 6.32-6.27 (m, 1H), 5.40 (d,  $J = 18$  Hz, 1H), 4.97 (d,  $J = 11$  Hz, 1H), 4.37 (broad singlet, 2H), 4.04 (broad singlet, 2H), 3.34 (s, 3H), 3.20 (s, 3H), 2.69 (m, 1H), 1.12 (d,  $J = 7.0$  Hz, 6H).


## EXAMPLE 44



5 Methyl[3-ethyl-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate


A solution of methyl[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl[3-(trifluoromethyl)-5-vinylbenzyl]carbamate (14 mg, 0.025 mmol) (Example 44) and Pd/C (5 mg) in THF (1 mL) and MeOH (500  $\mu$ L) was placed under hydrogen. The reaction was stirred at room temperature  
 10 under hydrogen for two hours. The reaction was then diluted with hexanes (25 mL), loaded on a silica gel column, and purified with 20% EtOAc/hexanes to afford methyl[3-ethyl-5-(trifluoromethyl)benzyl][{5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl}methyl]carbamate.  $R_f$  = 0.31 (25% EtOAc/hexanes). LCMS = 568.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C)  $\delta$  7.72 (broad singlet, 1H), 7.34 (d,  $J$  = 7.5 Hz, 1H), 7.20 (s, 1H), 7.10 (s, 1H), 7.07 (d,  $J$  = 8.0 Hz, 1H), 7.01 (dd,  $J$  = 2.0 Hz, 8.5, 1H), 7.01 (broad singlet, 1H), 6.86 (d,  $J$  = 2.5 Hz, 1H), 6.59 (d,  $J$  = 8.5 Hz, 1H), 4.44 (broad singlet, 2H), 4.11 (broad singlet, 2H), 3.39 (s, 3H), 3.24 (s, 3H), 2.73 (m, 1H), 2.25 (q,  $J$  = 7.5 Hz, 2H), 1.15 (d,  $J$  = 7.0 Hz, 6H), 0.92 (t,  $J$  = 7.5 Hz, 3H).

## EXAMPLE 45




- 5 Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylsulfinyl)-5-(trifluoromethyl)benzyl]carbamate
- Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylthio)-5-(trifluoromethyl)benzyl]carbamate (21.7 mg, 0.037 mmol) (Example 129) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was cooled to -78°C. Next, *m*-CPBA (8.3 mg, 0.037 mmol) was added. The reaction was stirred at -78°C for fifteen
- 10 minutes, then was warmed to -45°C, and after five minutes at -45°C was poured into NaHSO<sub>3</sub> (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with water (10mL), brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 50% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylsulfinyl)-5-(trifluoromethyl)benzyl]carbamate. *R<sub>f</sub>*=
- 15 0.20 (50% EtOAc/hexanes). LCMS = 602.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C) δ 7.63 (broad singlet, 1H), 7.54 (broad singlet, 1H), 7.48 (broad singlet, 1H), 7.33 (s, 1H), 7.32 (s, 1H), 7.07-7.05 (m, 2H), 6.89 (s, 1H), 6.62 (d, *J* = 8.0 Hz, 1H), 4.41 (broad singlet, 2H), 4.04 (broad singlet, 2H), 3.37 (s, 3H), 3.30 (s, 3H), 2.75 (m, 1H), 1.93 (s, 3H), 1.16 (d, *J* = 6.5 Hz, 6H).

## EXAMPLE 46



- 5 Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylsulfonyl)-5-(trifluoromethyl)benzyl]carbamate
- Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylthio)-5-(trifluoromethyl)benzyl]carbamate (18.0 mg, 0.031 mmol) (Example 129) in  $\text{CH}_2\text{Cl}_2$  (1 mL) was cooled to  $0^\circ\text{C}$ . Next, *m*-CPBA (17.3 mg, 0.078 mmol) was added. The reaction was stirred at  $0^\circ\text{C}$  for fifteen
- 10 minutes and then was poured into  $\text{NaHSO}_3$  (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with water (10 mL), brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification by flash chromatography with 50% EtOAc/hexanes afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[3-(methylsulfonyl)-5-(trifluoromethyl)benzyl]carbamate.  $R_f = 0.61$  (50% EtOAc/hexanes). LCMS = 618.2 ( $\text{M}+1$ )<sup>+</sup>.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 500 MHz,  $70^\circ\text{C}$ )  $\delta$  8.01 (s, 1H), 7.73 (s, 1H), 7.59 (broad singlet, 1H), 7.44 (broad singlet, 1H), 7.32 (d,  $J = 8.0$  Hz, 1H), 7.06 (s, 1H), 7.05 (s, 1H), 6.87 (d,  $J = 2.5$  Hz, 1H), 6.63 (d,  $J = 8.5$  Hz, 1H), 4.39 (broad singlet, 2H), 3.96 (broad singlet, 2H), 3.36 (s, 3H), 3.31 (s, 3H), 2.75 (m, 1H), 2.20 (s, 3H), 1.16 (d,  $J = 7.0$  Hz, 6H).
- 15

## EXAMPLE 47




5 Methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[(1-oxidopyridin-3-yl)methyl]carbamate

To a solution of methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}(pyridin-3-ylmethyl)carbamate (12 mg, 0.025 mmol) (Example 123) in CH<sub>2</sub>Cl<sub>2</sub> (400 μL) was added m-CPBA (28 mg, 0.13 mmol). The reaction was stirred at room temperature for two hours and then was poured into

10 NaHSO<sub>3</sub> (10 mL). The mixture was extracted with EtOAc (50 mL), and the organic extracts were washed with water (10mL), brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 10 to 20% MeOH/EtOAc afforded methyl{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}[(1-oxidopyridin-3-yl)methyl]carbamate. R<sub>f</sub> = 0.36 (20% MeOH/EtOAc). LCMS = 489.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz, 70°C) δ 6.21-8.22 (m, aromatics, 10H), 4.32 (s, 2H), 3.77 (s, 2H), 3.37 (s, 3H), 3.17 (s, 3H), 2.89 (m, 1H), 1.19 (d, J = 7.0 Hz, 6H).

15

## EXAMPLE 48



Methyl{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}[[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

Step A: 2-[3,5-Bis(trifluoromethyl)phenyl]-N-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}acetamide

A solution of 3,5-bis(trifluoromethyl)phenylacetic acid (105.3 mg, 0.387 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was cooled to 0 °C. N,N-diisopropylethylamine (169  $\mu\text{L}$ , 0.968 mmol) was added, followed by HATU (139 mg, 0.3667 mmol). The reaction was stirred at 0 °C for 2 minutes and then a solution of 1-[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methanamine (62.5 mg, 0.193 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was added. The reaction was stirred for 1 hour at 0 °C and then diluted with EtOAc (75 mL). The organics were washed with saturated  $\text{NaHCO}_3$  and brine (25 mL each), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification of the residue by flash chromatography with 100%  $\text{CH}_2\text{Cl}_2$  gave 2-[3,5-bis(trifluoromethyl)phenyl]-N-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}acetamide.  $R_f$  = 0.26 (25% EtOAc/hexanes). LCMS = 578.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.79 (s, 1H), 7.70 (s, 2H), 7.59 (m, 2H), 7.33 (d,  $J$  = 8.5 Hz, 1H), 7.25 (m, 1H), 6.97 (d,  $J$  = 2.0 Hz, 1H), 6.91 (d,  $J$  = 8.5 Hz, 1H), 5.84 (bs, 1H), 4.48 (dd,  $J$  = 15.1, 6.6 Hz, 1H), 4.20 (dd,  $J$  = 14.9, 4.3 Hz, 1H), 3.69 (s, 3H), 3.55 (s, 2H), 2.88 (m, 1H), 1.24 (d,  $J$  = 6.8 Hz, 6H).

Step B: {2-[3,5-Bis(trifluoromethyl)phenyl]ethyl}[[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine

To a solution of 2-[3,5-bis(trifluoromethyl)phenyl]-N-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}acetamide (77.6 mg, 0.134 mmol) in THF (4.4 mL) was added

BH<sub>3</sub> (1.34 mL of a 1M solution in THF, 1.34 mmol). The reaction was stirred overnight at room temperature and then 1 N HCl (10 mL) was added. The mixture was stirred for 5 minutes and then adjusted to a slightly basic pH with 1 N NaOH. The mixture was then extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 x 20 mL). The organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification of the residue by flash chromatography with 25 to 40% EtOAc/hexanes gave {2-[3,5-bis(trifluoromethyl)phenyl]ethyl}{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine. *R<sub>f</sub>* = 0.17 (25% EtOAc/hexanes). LCMS = 564.3 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>, 500 MHz) δ 7.75 (s, 1H), 7.73 (s, 1H), 7.64 (s, 2H), 7.53 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 7.24 (dd, *J* = 8.2, 2.0 Hz, 1H), 6.98 (d, *J* = 2.3 Hz, 1H), 6.91 (d, *J* = 8.5 Hz, 1H), 3.58-3.73 (m, 5H), 2.88 (m, 1H), 2.74-2.81 (m, 4H), 1.22 (d, *J* = 6.9 Hz, 6H).

Step C: Methyl{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

To a solution of amine {2-[3,5-bis(trifluoromethyl)phenyl]ethyl}{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}amine (16.6 mg, 0.0295 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added methyl chloroformate (5 μL, 0.059 mmol) followed by N,N-diisopropylethylamine (20 μL, 0.116 mmol). The reaction was stirred at room temperature for 30 minutes and then poured into saturated NaHCO<sub>3</sub> (15 mL). The mixture was extracted with EtOAc (40 mL) and the organic layer was washed with brine (20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification by flash chromatography with 25% EtOAc/hexanes yielded methyl{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate. *R<sub>f</sub>* = 0.24 (25% EtOAc/hexanes). LCMS = 622.2 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub> at 70 °C, 600 MHz) δ 7.74 (bs, 1H), 7.62 (s, 1H), 7.36 (d, *J* = 7.0 Hz, 1H), 7.22 (s, 2H), 7.07 (m, 2H), 6.88 (s, 1H), 6.63 (d, *J* = 8.1 Hz, 1H), 4.30 (bs, 2H), 3.34 (s, 3H), 3.25 (s, 3H), 2.97-3.04 (m, 2H), 2.75 (m, 1H), 2.33 (bs, 2H), 1.17 (d, *J* = 6.4 Hz, 6H).

## EXAMPLE 49

Methyl[3,5-bis(trifluoromethyl)benzyl][2-cyano-5-(trifluoromethyl)benzyl]carbamate

To a solution of Example 5 dissolved in DMF (1.5 mL) was added CuCN (20 mg, 0.22 mmol) and the resultant mixture was heated at 100 °C for 16 h. The reaction was cooled to room temperature, poured into NH<sub>4</sub>OH (10 mL) and extracted with EtOAc (4 x 30 mL). The combined organic extracts were washed with H<sub>2</sub>O and brine (10 mL each), dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. Purification by flash chromatography (0 to 25% EtOAc/hexanes) afforded the title compound as a colorless oil. LCMS = 485 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): δ 7.81 (s, 1H), 7.78 (d, *J* = 7.7 Hz, 1H), 7.67-7.31 (m, 4H), 4.82 (br s, 2H), 4.72 (s, 2H), 3.91 (s, 3H).




## EXAMPLE 50

Methyl[3,5-bis(trifluoromethyl)benzyl][2-formyl-5-(trifluoromethyl)benzyl]carbamate

- 5 Example 49 (85 mg, 0.18 mmol) was dissolved in 80% aqueous formic acid (5 mL) and platinum oxide (80% hydrate, 4.0 mg, 0.018 mmol) was added in one portion. The reaction was heated at 60 °C for 16 h and the platinum removed by filtration. H<sub>2</sub>O (15 mL) was added and the mixture extracted with diethyl ether (4 x 30 mL). The combined organic extracts were washed with H<sub>2</sub>O and brine (10 mL each), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. Purification by flash chromatography (0 to 25%  
10 EtOAc/hexanes) gave the title compound as a colorless oil. LCMS = 488 (M+1)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): δ 10.21 (s, 1H), 7.96 (d, *J*= 8.0 Hz, 1H), 7.81 (s, 1H), 7.79 (d, *J*= 8.0 Hz, 1H), 7.72-7.55 (m, 3H), 5.18 (br, s, 2H), 4.78 (br s, 2H), 3.85 (s, 3H)












Following the procedures outlined for Examples 1-50 the compounds listed in Tables 2-5 were prepared.











Table 2
























5

Example	A <sup>1</sup>	LC/MS Data
51		566.3
52		564.1
53		600.3
54		550.3
55		570.2
56		604.0
57		594.3

58		584.3
59		602.0
60		602.0
61		596.3
62		608.3
63		588. (M+Na)
64		584.4
65		602.4 (M+Na)
66		618. (M+Na)
67		564.4
68		634

69	 <chem>CCOC1=CC=C(CC)C=C1</chem>	608.1
70	 <chem>CCOC1=CC=C(C(C)(C)C)C=C1</chem>	622.0
71	 <chem>CC1=CC=C(C)C=C1</chem>	578.1
72	 <chem>CS1=CC=C(C=C1)</chem>	604.3
73	 <chem>CCOC1=CC=C(C=O)C=C1</chem>	594.3
74	 <chem>CCOC1=CC=C(C(F)C(C)C)C=C1</chem>	626.1
75	 <chem>CCOC1=CC=C(C(F)C)C=C1</chem>	598.0
76	 <chem>CC1=CC=C2C(=C1)N=CC=C2</chem>	601.3
77	 <chem>C1=CC=C2C(=C1)N=CC=C2</chem>	587.2
78	 <chem>COC1=CC=C2C(=C1)N=CC=C2</chem>	617.2

79		567.2
79a		635.0
80		587.2
81		587.2
82		602.1
83		587.2
84		587.2
85		597.2
86		600.2
87		542

88	 <chem>Clc1cc(s1)C</chem>	576
89	 <chem>c1ccoc1C</chem>	526
90	 <chem>Cc1cc(s1)C</chem>	556
91	 <chem>c1ccccc1C</chem>	536.2
92	 <chem>Cc1cc(s1)C</chem>	556
93	 <chem>c1ccncc1C</chem>	537.2
94	 <chem>COc1ccncc1C</chem>	567.2
95	 <chem>Cc1cc(s1)C</chem>	556
96	 <chem>CCc1ccccc1C</chem>	564.4
97	 <chem>COc1ccc(C=C)cc1C</chem>	592.2
98	 <chem>c1cc(s1)C</chem>	542








99		555
100		537.2
101		581.2
101a		609.2
102		584

Table 3



EXAMPLE	A <sup>1</sup>	R <sup>2</sup>	LC/MS
103		OMe	564.4 (M+Na) (4.40)










104		OMe	592.4 (M+Na) (4.60)
105		F	558.4 (M+1) (4.62)
106		F	558.4 (M+1) (4.64)
107		F	572.5 (M+1) (4.68)
108		H	512.2 (M+1) (4.56)
108a		Cl	564.0 (M+1) (4.57)
109		H	540.2 (M+1) (4.76)
110		Cl	592.2 (M+1) (4.74)











Table 4



5

EXAMPLE	Q	LC/MS Data
111		540.3 (M+1)
112		622.3 (M+1)
113		558.3 (M+1)
114		556.3 (M+1)
115		538.3 (M+1)
116		517.3 (M+1)

117		562.2 (M+1)
118		472.4 (M+1)
119		666.2 (M+1)
120		540.2 (M+1)
121		597.4 (M+1)
122		620.2 (M+1)
123		473.2 (M+1)
124		473.3 (M+1)









125		473.3 (M+1)
126		500.3 (M+1)
127		478.2 (M+1)
128		478.3 (M+1)
129		586.2 (M+1)
130		570.2 (M+1)

Table 5



EXAMPLE	Z	LCMS (M+1) <sup>+</sup>
131		606.3
132		621.4
133		624.3


## EXAMPLE 134



5 Methyl [3,5-bis(trifluoromethyl)benzyl][2-isoquinolin-4-yl-5-(trifluoromethyl)benzyl]carbamate

To a solution of methyl [3,5-bis(trifluoromethyl)benzyl](2-iodo-5-trifluoromethyl-benzyl)carbamate (41 mg, 0.070 mmol) (Example 5) in 3:2 EtOH:toluene, 4-isoquinolineboronic acid (18.2 mg, 0.105 mmol) and 2.5 M sodium carbonate (0.175 mmol) were added. The solution was degassed, followed by the addition of a catalytic amount of tetrakis(triphenylphosphine)palladium. The solution was stirred at 95-  
 10 100°C until completion of the reaction. Upon cooling, the solvent was removed in vacuo. The desired product was obtained by preparative thin layer chromatography using 25% acetone/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 9.36 (s, 1H), 8.38 (s, 1H), 8.10-8.13 (m, 1H), 7.68-7.80 (m, 4H), 7.26-7.64 (m, 5H), 4.22-4.48 (m, 4H), 3.64-3.78 (m, 3H). LC-MS (M+1) 587.2 (3.48 min).

## EXAMPLE 135



5 N-[3,5-Bis(trifluoromethyl)benzyl]-N-[[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]acetamide

Step A: N-[2-amino-5-(trifluoromethyl)benzyl]-N-[3,5-bis(trifluoromethyl)benzyl]-acetamide To a solution of [2-amino-5-(trifluoromethyl)benzyl][3,5-bis(trifluoromethyl)benzyl] amine (Example 3) (270 mg, 0.65 mmol) in methylene chloride at 0 °C, acetic anhydride (0.061 mL, 0.65 mmol) in methylene chloride and then triethylamine were added (0.018 mL, 0.13 mmol). The solution was stirred for 0.5 hours at 0 °C. This solution was washed with saturated sodium bicarbonate, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was purified by column chromatography using 40% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.80 (s, 1H), 7.56 (s, 2H), 7.30 (d, J = 8.5 Hz, 1H), 7.04 (s, 1H), 6.65 (d, J = 8.5 Hz, 1H), 4.67 (s, 2H), 4.65 (s, 2H), 2.22 (s, 3H).


15 Step B: N-[3,5-bis(trifluoromethyl)benzyl]-N-[2-iodo-5-(trifluoromethyl)benzyl] acetamide

To a solution of the title compound from Step A (58 mg, 0.127 mmol) in chloroform at room temperature, n-amyl nitrite (0.025 mL, 0.191 mmol) was added. The solution was stirred for 5 minutes and then iodine (65 mg, 0.254 mmol) was added. This solution was stirred at 70 °C for 1 hour. Upon cooling to room temperature, the organic solution was washed with saturated sodium thiosulfate and brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was purified by column chromatography using 25% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): a 1:2 mixture of rotamers δ 8.01 (d, J = 8.0 Hz, 2/3H), 7.93 (d, J = 8.0 Hz, 1/3H), 7.84 (s, 1/3H), 7.79 (s, 2/3H), 7.67 (s, 4/3H), 7.60 (s, 2/3H), 7.43 (s, 1/3H), 7.29 (d, J = 8.0 Hz, 2/3H), 7.23 (m, 1H), 4.78 (s, 2/3H), 4.72 (s, 4/3H), 4.68 (s, 2/3H), 4.55 (s, 4/3H), 2.32 (s, 1H), 2.27 (s, 2H).

Step C: *N*-[3,5-Bis(trifluoromethyl)benzyl]-*N*-{[5'-isopropyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}acetamide

The title compound was prepared according to the procedure described in Example 6, but using (5-isopropyl-2-methoxyphenyl)boronic acid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): a mixture of 1:3 rotamers δ 7.76 (m, 1H), 7.58-7.63 (m, 2H), 7.24-7.37 (m, 3H), 6.8-6.94 (m, 3H), 3.69 (s, 9/4H), 3.62 (s, 3/4H), 2.79-2.91 (m, 1H), 2.09 (s, 3/4H), 2.01 (s, 9/4H), 1.15-1.26 (m, 6H). LC-MS (M+1) 592.0 (4.60 min).


#### EXAMPLE 136



Methyl [3,5-bis(trifluoromethyl)benzyl]{[5'-(hydroxymethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

Sodium borohydride (9.9 mg, 0.262 mmol) was added to a solution of methyl [3,5-bis(trifluoromethyl)benzyl]{[5'-formyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (Example 73, 78 mg, 0.131 mmol) in methanol at room temperature. The reaction was stirred for one hour, then solvent was removed in vacuo. The title compound was obtained by column chromatography using 3:7 acetone/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.25 (s, 1H), 7.30-7.62 (m, 6H), 7.08 (m, 1H), 6.94 (m, 1H), 4.64 (s, 2H), 4.15-4.40 (m, 4H), 3.64-3.74 (m, 6H). LC-MS (M-17) 578.0 (4.09 min).

## EXAMPLE 137




5 Methyl [3,5-bis(trifluoromethyl)benzyl][{5'-(fluoromethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl)methyl}carbamate

Diethylaminosulfur trifluoride (0.014 mL, 0.103 mmol) was added dropwise to a solution of methyl [3,5-bis(trifluoromethyl)benzyl][{5'-(hydroxymethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl)methyl}carbamate (Example 136, 41 mg, 0.069 mmol) in anhydrous methylene chloride at 0 °C. The  
 10 reaction was gradually warmed to room temperature until the reaction was complete. At this time, the reaction was re-cooled to 0 °C and water was added. The organic layer was diluted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was purified by preparative thin layer chromatography using 15% acetone/hexane.




## EXAMPLE 138



5 Methyl [3,5-bis(trifluoromethyl)benzyl][{[5'-(difluoromethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

Diethylaminosulfur trifluoride (0.024 mL, 0.179 mmol) was added dropwise to a solution of methyl [3,5-bis(trifluoromethyl)benzyl][{[5'-formyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (Example 73, 53 mg, 0.089 mmol) in anhydrous methylene chloride at 0 °C. The reaction was gradually  
 10 warmed to room temperature until the reaction was complete. At this time, the reaction was re-cooled to 0 °C, and water was added. The organic layer was diluted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was purified by column chromatography using 1:3 EtOAc:hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.77 (s, 1H), 7.63 (d, *J* = 7.5 Hz, 1H), 7.36-7.60 (m, 4H), 7.35 (d, *J* = 8.0 Hz, 1H), 7.26 (s, 1H), 7.04 (d, *J* = 7.5 Hz, 1H), 6.65 (t, *J* = 56.5 Hz, 1H), 4.20-4.60 (m, 4H), 3.75 (m, 6H). LC-MS (M+Na) 638.3 (4.34 min).  
 15


## EXAMPLE 139



5 Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(2,2,2-trifluoro-1-hydroxyethyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

To a solution of methyl [3,5-bis(trifluoromethyl)benzyl] {[5'-formyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Example 73, 184 mg, 0.31 mmol) in anhydrous tetrahydrofuran at 0 °C, 0.5 M trimethyl(trifluoromethyl) silane (2.5 mL, 1.24 mmol) and a catalytic  
 10 amount of tetrabutylammonium fluoride were added. The reaction was stirred with gradual warming to room temperature. Once the reaction was complete, it was quenched with saturated ammonium chloride solution. The organic was extracted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by preparative thin layer chromatography using 1:4 acetone:hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.74 (s, 1H), 7.61 (m, 1H),  
 15 7.15-7.56 (m, 6H), 6.99 (m, 1H), 4.97 (m, 1H), 4.19-4.38 (m, 4H), 3.61-3.76 (m, 6H). LC-MS (M+Na) 686.9 (4.28 min).

## EXAMPLE 140



5


Methyl [3,5-bis(trifluoromethyl)benzyl] {[5'-(1-hydroxyethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

Methylmagnesium bromide (0.59 mL, 0.83 mmol) was added dropwise to a solution of methyl [3,5-bis(trifluoromethyl)benzyl] {[5'-formyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Example 73, 289 mg, 0.49 mmol) in anhydrous tetrahydrofuran at 0 °C. The solution was allowed to warm to room temperature. Once the reaction was complete, saturated ammonium chloride solution was added to quench the reaction. The mixture was extracted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by flash chromatography using 30% acetone/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.73 (s, 1H), 7.58 (m, 1H), 7.32-7.51 (m, 5H), 7.11 (m, 1H), 6.94 (m, 1H), 4.87 (m, 1H), 4.25-4.43 (m, 4H), 3.69 (m, 6H), 1.48 (d, *J* = 4.2 Hz, 3H). LC-MS (M+Na) 632.0 (4.18 min).

10

15


## EXAMPLE 141



5 Methyl {[5'-acetyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}{[3,5-  
bis(trifluoromethyl)benzyl]carbamate

Methyl [3,5-bis(trifluoromethyl)benzyl]{[5'-(1-hydroxyethyl)-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate (Example 140, 29 mg, 0.048 mmol) was dissolved in methylene chloride, and Dess-Martin periodinane (31mg, 0.072 mmol) was added. Once the reaction was complete, the solution was  
 10 diluted in diethyl ether and washed sequentially with saturated sodium thiosulfate, saturated sodium bicarbonate and water. The aqueous extracts were combined and back extracted with diethyl ether then recombined with the previous organic layers. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by preparative thin layer chromatography using 2:3 EtOAc:hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 8.02 (d, *J* = 7.8 Hz, 1H), 7.76  
 15 (d, *J* = 2.4 Hz, 1H), 7.73 (s, 1H), 7.61 (d, *J* = 7.8 Hz, 1H), 7.32-7.55 (m, 4H), 6.99 (m, 1H), 4.18-4.58 (m, 4H), 3.79 (s, 3H), 3.69-3.76 (m, 3H), 2.57 (s, 3H). LC-MS (M+1) 608.3 (4.19 min).

## EXAMPLE 142



5 Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(pentafluoroethyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

Step A: Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(trifluoroacetyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(2,2,2-trifluoro-1-hydroxyethyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Example 139, 29 mg, 0.048 mmol) was dissolved in methylene chloride, and Dess-Martin periodinane (31 mg, 0.072 mmol) was added. Once the reaction was complete, the solution was diluted in diethyl ether and washed sequentially with saturated sodium thiosulfate, saturated sodium bicarbonate and water. The aqueous extracts were combined and back extracted with diethyl ether, then recombined with the previous organic layers. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was

10 (trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Example 139, 29 mg, 0.048 mmol) was dissolved in methylene chloride, and Dess-Martin periodinane (31 mg, 0.072 mmol) was added. Once the reaction was complete, the solution was diluted in diethyl ether and washed sequentially with saturated sodium thiosulfate, saturated sodium bicarbonate and water. The aqueous extracts were combined and back extracted with diethyl ether, then recombined with the previous organic layers. The combined organic

15 layers were dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by preparative thin layer chromatography using 1:3 acetone:hexane.

Step B: Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(pentafluoroethyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

Diethylaminosulfur trifluoride (0.008 mL, 0.058 mmol) was added dropwise to a solution of Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(trifluoroacetyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Step A) (19 mg, 0.029 mmol) in anhydrous methylene chloride at 0 °C. The reaction was gradually warmed to room temperature until the reaction was complete. At this time, the reaction was re-cooled to 0 °C, and water was added to quench. The organic layer was diluted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated.


20 [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(trifluoroacetyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate (Step A) (19 mg, 0.029 mmol) in anhydrous methylene chloride at 0 °C. The reaction was gradually warmed to room temperature until the reaction was complete. At this time, the reaction was re-cooled to 0 °C, and water was added to quench. The organic layer was diluted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated.

25 The desired product was purified by column chromatography using 30% EtOAc/hexane. <sup>1</sup>H NMR

(CDCl<sub>3</sub>, 600 MHz)  $\delta$  7.74 (s, 1H), 7.60-7.66 (m, 2H), 7.28-7.54 (m, 5H), 7.06 (d,  $J$  = 9.0 Hz, 1H), 4.16-4.59 (m, 4H), 3.80 (s, 3H), 3.68 (s, 3H).

## EXAMPLE 143


5



Methyl [3,5-bis(trifluoromethyl)benzyl]{{[2'-methoxy-5'-(2,2,2-trifluoro-1-hydroxy-1-methylethyl)-4-(trifluoromethyl)biphenyl-2-yl]methyl}carbamate

- 10 The title compound from Example 142, Step A (7 mg, 0.027 mmol) was dissolved in anhydrous THF at 0 °C. Methylmagnesium bromide (0.023 mL, 0.032 mmol) was added dropwise, and the mixture was slowly warmed to room temperature. Once the reaction was complete, saturated ammonium chloride solution was added to quench the reaction. The mixture was extracted with methylene chloride, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was
- 15 obtained by flash chromatography using 30% acetone/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz)  $\delta$  7.73 (s, 1H), 7.30-7.61 (m, 7H), 6.98 (m, 1H), 4.19-4.62 (m, 4H), 3.75 (s, 3H), 3.64-3.71 (m, 3H), 1.75-1.78 (m, 3H). LC-MS (M+Na) 699.8 (4.35 min).

## EXAMPLE 144




5 Methyl [3,5-bis(trifluoromethyl)benzyl]({2'-methoxy-4-(trifluoromethyl)-5'-(2,2,2-trifluoro-1-methylethyl)biphenyl-2-yl}methyl)carbamate

Step A: Methyl [3,5-bis(trifluoromethyl)benzyl]({2'-methoxy-4-(trifluoromethyl)-5'-[1-(trifluoromethyl)vinyl]biphenyl-2-yl}methyl)carbamate

10 Methanesulfonyl chloride was added to the title compound from Example 143, Step A (45 mg, 0.068 mmol) in methylene chloride at 0 °C. The solution was stirred at the same temperature until starting material was consumed. The solution was washed with water, then brine, dried over anhydrous sodium sulfate, filtered and concentrated. The product was purified by preparative thin layer chromatography using 15% EtOAc/hexane.

15 Step B: The title compound from Step A (10 mg, 0.017 mmol) was hydrogenated at 1 atm in methanol using Pd/C as the catalyst. The solution was filtered through Celite then concentrated. The title compound was purified by preparative thin layer chromatography using 1:4 EtOAc:hexane.

## EXAMPLE 145




5 2'--[[[3,5-Bis(trifluoromethyl)benzyl](methoxycarbonyl)amino]methyl]-6-methoxy-4'-(trifluoromethyl)biphenyl-3-carboxylic acid

Jones Reagent (0.10 mL, 0.382 mmol) was added to a solution of the title compound from Example 136, Step A (113 mg, 0.191 mmol) in acetone. The solution was stirred at room temperature. Upon completion of the reaction, i-PrOH was added to quench the reaction. The solution was filtered, and the  
 10 filtrate was taken up in diethyl ether and water. The organic layer was separated, dried over anhydrous sodium sulfate, filtered and concentrated to give the title compound. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.16 (s, 1H), 7.84 (s, 1H), 7.74 (s, 1H), 7.63 (d, *J* = 8.0 Hz, 1H), 7.45-7.58 (m, 3H), 7.35 (d, *J* = 8.0 Hz, 1H), 7.02 (d, *J* = 8.0 Hz, 1H), 4.20-4.60 (m, 4H), 3.82 (s, 3H), 3.74 (m, 3H). LC-MS (M+1) 609.9 (4.01 min).



## EXAMPLE 146




5 Methyl 2'-{[[[3,5-bis(trifluoromethyl)benzyl](methoxycarbonyl)amino]methyl]}-6-methoxy-4'-(trifluoromethyl)biphenyl-3-carboxylate

(Trimethylsilyl)diazomethane (0.059 mL, 0.117 mmol) was added to a solution of the title compound from Example 135 (32 mg, 0.053 mmol) in diethyl ether at 0 °C. The reaction was allowed to stir at room temperature until completion. It was quenched with a few drops of trifluoroacetic acid then

- 10 concentrated. The title compound was obtained by preparative thin layer chromatography using 30% acetone/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 8.09 (s, 1H), 7.79 (s, 1H), 7.72 (s, 1H), 7.60 (d, *J* = 8.4 Hz, 1H), 7.43-7.58 (m, 3H), 7.32 (d, *J* = 7.8 Hz, 1H), 6.97 (m, 1H), 4.17-4.60 (m, 4H), 3.90 (s, 3H), 3.78 (s, 3H), 3.74 (m, 3H). LC-MS (M+Na) 645.9 (4.40 min).

## EXAMPLE 147




5 Methyl [3,5-bis(trifluoromethyl)benzyl][{[2'-methoxy-5'-[(methylamino)carbonyl]-4-(trifluoromethyl)biphenyl]-2-yl)methyl} carbamate

A mixture of the title compound from Example 145 (22 mg, 0.036 mmol), methylamine hydrochloride (3.6 mg, 0.054 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (10.3 mg, 0.054 mmol), 1-hydroxybenzotriazole hydrate (7.3 mg, 0.054 mmol) and diisopropylethylamine (0.02 mL, 0.108 mmol) were stirred in methylene chloride overnight at room temperature. The solvent was removed in vacuo. The title compound was obtained by preparative thin layer chromatography using 70% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.81 (d, J = 7.2 Hz, 1H), 7.73 (s, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.34-7.56 (m, 4H), 7.32 (d, J = 7.8 Hz, 1H), 6.97 (d, J = 7.8 Hz, 1H), 6.14 (br s, 1H), 4.19-4.42 (m, 4H), 3.76 (s, 3H), 3.65-3.73 (m, 3H), 3.00 (m, 3H). LC-MS (M+1) 622.9 (3.96 min).

15

## EXAMPLE 148




5 Methyl [3,5-bis(trifluoromethyl)benzyl] {[2'-methoxy-5'-(3-methyl-1,2,4-oxadiazol-5-yl)-4-(trifluoromethyl)biphenyl-2-yl]methyl} carbamate

Oxalyl chloride (0.14 mL, 0.28 mmol) and a few drops of dimethylformamide were added to a solution of the title compound from Example 145 (86 mg, 0.14 mmol) in methylene chloride. The reaction mixture was stirred at room temperature for 1 h then concentrated in vacuo. The crude product was re-dissolved in anhydrous dichloroethane, and acetamide oxime (16 mg, 0.21 mmol) was added along with a few drops of dimethylformamide to increase solubility. The reaction was heated to and maintained at reflux overnight. Once the reaction cooled to room temperature, the solvent was removed through evaporation. The oil was taken up in methylene chloride, washed with water, then brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by preparative thin layer chromatography using 50% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 8.15 (s, 1H), 7.86 (s, 1H), 7.71 (s, 1H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.43-7.56 (m, 3H), 7.35 (d, *J* = 7.8 Hz, 1H), 7.07 (m, 1H), 4.20-4.60 (m, 4H), 3.81 (s, 3H), 3.70 (m, 3H), 2.46 (s, 3H). LC-MS (M+1) 647.9 (4.40 min).

10

15

## EXAMPLE 149


5 Methyl [3,5-bis(trifluoromethyl)benzyl][2'-cyano-4-(trifluoromethyl)biphenyl-2-yl]methyl carbamate

A mixture of methyl [3,5-bis(trifluoromethyl)benzyl](2-iodo-5-trifluoromethyl-benzyl)carbamate (29 mg, 0.050 mmol), 2-cyanophenylboronic acid (10.9 mg, 0.074 mmol), potassium fluoride (8.7 mg, 0.149 mmol), 2-dicyclohexylphosphino(biphenyl) ligand (0.7 mg, 0.002 mmol) and a catalytic amount of palladium acetate was stirred in dioxane overnight at 80 °C. The title compound was obtained by

10 preparative thin layer chromatography using 20% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.74 (m, 2H), 7.68 (d, *J* = 7.8 Hz, 1H), 7.64 (t, *J* = 7.8 Hz, 1H), 7.58 (t, *J* = 7.8 Hz, 1H), 7.52 (m, 3H), 7.39 (d, *J* = 7.2 Hz, 1H), 7.24 (m, 1H), 4.30-4.55 (m, 4H), 3.71 (s, 3H). LC-MS (M+1) 561.0 (4.29 min).

## EXAMPLE 150

15



Methyl [3,5-bis(trifluoromethyl)benzyl][[5'-ethyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]carbamate

Step A: 2-Iodo-4-ethylanisole

A mixture of silver sulfate (1.15 g, 3.68 mmol) and iodine (0.93 g, 3.68 mmol) were stirred in methanol at room temperature. A solution of p-ethylanisole (0.50 g, 3.68 mmol) in methanol was added dropwise to the stirring mixture. The mixture was stirred at room temperature until completion of the reaction. It was filtered and the filtrate was concentrated. The residue was dissolved in methylene chloride, washed with 1 N sodium hydroxide, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained without further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz) δ 7.64 (d, J = 2.0 Hz, 1H), 7.15 (dd, J = 8.5, 2.0 Hz, 1H), 6.77 (d, J = 8.0 Hz, 1H), 3.88 (s, 3H), 2.58 (q, J = 15.5, 8.0 Hz, 2H), 1.23 (t, J = 7.5 Hz, 3H).

Step B: 2-(5-Ethyl-2-methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


To a mixture of bispinacolatodiboron (0.70 g, 2.95 mmol), potassium acetate (0.67 g, 6.81 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (55 mg, 0.068 mmol) in dioxane, 4-ethyl-2-iodoanisole (0.595 g, 2.27 mmol) from Step A was added. The reaction was stirred at 80 °C for 4 h then cooled to room temperature. After filtration to remove insoluble material, the solution was diluted in ethyl acetate, washed with water then brine, dried over anhydrous sodium sulfate, filtered and concentrated. The title compound was obtained by flash chromatography using 10% EtOAc/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.50 (d, J = 2.4 Hz, 1H), 7.22 (dd, J = 8.4, 2.4 Hz, 1H), 6.80 (d, J = 8.4 Hz, 1H), 3.82 (s, 3H), 2.59 (q, J = 15.6, 7.8 Hz, 2H), 1.36 (s, 12H), 1.22 (t, J = 7.8 Hz, 3H). LC-MS (M+1) 263.3 (3.90 min).

Step C: Methyl [3,5-bis(trifluoromethyl)benzyl][[5'-ethyl-2'-methoxy-4-(trifluoromethyl)biphenyl-2-yl]methyl]carbamate

A mixture of methyl [3,5-bis(trifluoromethyl)benzyl](2-iodo-5-trifluoromethyl-benzyl)carbamate (105 mg, 0.18 mmol) (Example 5), the title compound from Step B (58 mg, 0.22 mmol), anhydrous potassium carbonate (50 mg, 0.36 mmol), dichloro[1,1'-bis(diphenylphosphino) ferrocene]palladium(II) dichloromethane adduct (14.7 mg, 0.018 mmol), and a catalytic amount of palladium acetate in anhydrous dioxane were stirred at 80 °C for approximately 40 hours. The solvent was removed in vacuo, and the title compound was obtained by preparative thin layer chromatography using 60% CH<sub>2</sub>Cl<sub>2</sub>/hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 600 MHz) δ 7.71 (s, 1H), 7.56 (d, J = 7.8 Hz, 1H), 7.45 (m, 2H), 7.32 (d, J = 7.8 Hz, 1H), 7.25 (m, 1H), 7.21 (m, 1H), 6.88 (d, J = 2.4 Hz, 1H), 6.87 (m, 1H), 4.23-4.56 (m, 4H), 3.74 (m, 3H), 3.68 (s, 3H), 2.60 (m, 2H), 1.24 (m, 3H). LC-MS (M+1) 594.0 (4.76 min).

The claims defining the invention are as follows:

1. A compound having Formula Ib, or a pharmaceutically acceptable salt thereof, wherein



5  $A^1$  is selected from the group consisting of phenyl, thienyl, furyl, pyridyl, 1-oxidopyridinyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl, oxazolyl, isoxazolyl, and oxadiazolyl;

wherein  $A^1$  is optionally substituted with 1-4 substituent groups independently selected from  $-C_1-C_5$  alkyl,  $-OC_1-C_3$ alkyl,  $-CO_2C_1-C_3$ alkyl,  $-CO_2H$ , halogen,  $-NR^{10}R^{11}$ ,  
 10  $-C(=O)C_1-C_3$ alkyl,  $-C(=O)H$ ,  $-C(=O)NR^{10}R^{11}$ ,  $-SC_1-C_3$ alkyl,  $-C_2-C_3$ alkenyl,  $-NO_2$ ,  $-C_3-C_6$ cycloalkyl, and a 5-6-membered heterocyclic ring having 1-3 heteroatoms independently selected from N, S, and O, and optionally also comprising 1-3 double bonds, wherein  $-C_1-C_3$  alkyl and  $-C_1-C_5$  alkyl in all occurrences are optionally substituted with 1-6 substituents independently selected from 1-5 halogens and one  $-OH$ -group; and  
 15  $-C_3-C_6$  cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and  $-C_1-C_3$ alkyl; and  $-C_2-C_3$ alkenyl is optionally substituted with 1-3 halogens:

$A^2$  is selected from the group consisting of phenyl, thienyl, furyl, pyridyl, 1-oxidopyridinyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl,  
 20 dihydroindolyl, oxazolyl, isoxazolyl, oxadiazolyl, and  $C_3-C_6$  cycloalkyl;

wherein  $A^2$  is optionally substituted with 1-2 substituent groups independently selected from  $-C_1-C_4$  alkyl,  $-OC_1-C_3$  alkyl,  $-C(=O)C_1-C_3$ alkyl,  $-C(=O)H$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)_xC_1-C_3$  alkyl,  $-NHS(O)_2C_1-C_3$  alkyl,  $-NR^{10}R^{11}$ ,  $-NR^{10}(C=O)R^{11}$ ,  $-C_2-C_3$  alkenyl,  $-C(=O)NR^{10}R^{11}$ , halogen,  $-C_3-C_6$  cycloalkyl, and a 5-6-membered heterocyclic ring  
 25 having 1-3 heteroatoms independently selected from N, S, and O, and optionally also

comprising 1-3 double bonds, wherein  $C_1$ - $C_3$  alkyl,  $C_1$ - $C_4$  alkyl, and  $C_2$ - $C_3$  alkenyl in all instances are optionally substituted with 1-3 halogens, and  $-C_3$ - $C_6$  cycloalkyl and the 5-6-membered heterocyclic ring are optionally substituted with 1-3 substituents independently selected from halogen and  $-C_1$ - $C_3$  alkyl;

5           x is an integer selected from 0, 1, and 2;

          n is an integer selected from 0 and 1;

          Z is selected from the group consisting of  $-S(O)_xC_1$ - $C_6$  alkyl,  $-S(O)_2NR^{17}R^{18}$ ,  $-C(=S)OC_1$ - $C_6$ alkyl, and  $-C(=O)X$ , wherein X is selected from the group consisting of H,  $-C_1$ - $C_6$  alkyl,  $-OC_1$ - $C_6$  alkyl,  $-SC_1$ - $C_6$  alkyl, and  $-NR^{10}R^{11}$ ; wherein  $-C_1$ - $C_6$  alkyl in all  
10 instances is optionally substituted with 1-13 halogens and 1-2 substituents independently selected from  $-OC_1$ - $C_3$ alkyl,  $-CN$ , and  $-NO_2$ , wherein  $-OC_1$ - $C_3$ alkyl is optionally substituted with 1-7 halogens and is optionally also substituted with 1-2  $-OC_1$ - $C_2$  alkyl;

$R^1$ , is selected from the group consisting of H,  $-OH$ ,  $-C_1$ - $C_3$  alkyl, and  $-OC_1$ - $C_3$  alkyl, wherein  $C_1$ - $C_3$  alkyl, and  $-OC_1$ - $C_3$  alkyl are each optionally substituted with 1-3  
15 halogens and are also optionally substituted with one  $-OC_1$ - $C_2$  alkyl;

$R^{17}$  and  $R^{18}$  are each independently selected from the group consisting of H,  $-C_1$ - $C_5$  alkyl, and  $-C_3$ - $C_7$  cycloalkyl, wherein  $-C_1$ - $C_5$  alkyl, and  $-C_3$ - $C_7$  cycloalkyl are optionally substituted with 1-13 halogens;

$R^2$  and  $R^3$  are each independently selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-C_1$ - $C_3$ alkyl,  $-OC_1$ - $C_3$ alkyl,  $-C_2$ - $C_3$ alkenyl,  $-C_3$ - $C_6$ cycloalkyl optionally having a double bond,  $-OC_3$ - $C_6$ cycloalkyl optionally having a double bond,  $-C(=O)C_1$ - $C_3$ alkyl,  $-C(=O)C_3$ - $C_6$ cycloalkyl,  $-C(=O)H$ ,  $-CO_2H$ ,  $-CO_2C_1$ - $C_3$ alkyl,  $-C(=O)NR^{10}R^{11}$ ,  $-CN$ ,  $-NO_2$ , and a 5-6-membered heterocyclic ring having  
20 1-4 heteroatoms independently selected from N, S, and O, and optionally 1-3 double bond, wherein  $C_1$ - $C_3$  alkyl and  $-C_2$ - $C_3$  alkenyl in all instances are optionally substituted with 1-5 halogens, and  $-C_3$ - $C_6$  cycloalkyl and the 5-6-membered heterocyclic ring are in all occurrences optionally substituted with 1-3 substituents independently selected from halogen,  $C_1$ - $C_3$  alkyl,  $OC_1$ - $C_3$  alkyl,  $CF_3$ , and  $-OCF_3$ ; and

$R^{10}$  and  $R^{11}$  are each independently selected from H and  $-C_1$ - $C_3$  alkyl.

30           2.       The compound of Claim 1, wherein  $R^2$  and  $R^3$  are each independently selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-C_1$ - $C_3$ alkyl,  $-OC_1$ - $C_3$ alkyl,  $-CN$ ,  $NO_2$ , and pyridyl, wherein  $C_1$ - $C_3$ alkyl in all instances is optionally substituted with 1-3 halogens, or a pharmaceutically acceptable salt thereof.

3. The compound of Claim 1, wherein  $R^2$  is selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-C_1-C_3$ alkyl,  $OC_1-C_3$ alkyl,  $-CN$ ,  $-NO_2$  and pyridyl, wherein  $C_1-C_3$ alkyl in all instances is optionally substituted with 1-3 halogens; and


$R^3$  is selected from the group consisting of H, halogen,  $-CH_3$ ,  $CF_3$ ,  $-OCH_3$ , and  $-OCF_3$ , or a pharmaceutically acceptable salt thereof.

4. The compound of any one of Claims 1 to 3, wherein  $A^1$  is selected from the group consisting of phenyl, thienyl, furyl, pyridyl, quinolyl, isoquinolyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl, oxazolyl, and isoxazolyl; and

$A^2$  is selected from phenyl, pyridyl, thienyl, 1-oxidopyridinyl, and cyclohexyl;

wherein  $A^1$  and  $A^2$  are optionally substituted as in Claim 1, or a pharmaceutically acceptable salt thereof.

5. The compound having formula II, or a pharmaceutically acceptable salt thereof, wherein



$R^1$  is selected from the group consisting of H, halogen, and  $C_1-C_3$  alkyl, wherein  $C_1-C_3$  alkyl is optionally substituted with 1-3 halogens;

$R^2$  is selected from the group consisting of H, halogen,  $-NR^{10}R^{11}$ ,  $-OC_1-C_3$  alkyl,  $C_1-C_3$  alkyl,  $-CN$ ,  $-NO_2$ , and pyridyl, wherein  $C_1-C_3$  alkyl in all occurrences is optionally substituted with 1-3 halogens;

$R^4$  is selected from the group consisting of H, halogen,  $-C_1-C_3$  alkyl,  $-OC_1-C_2$  alkyl,  $-SC_1-C_2$  alkyl and  $-CN$ , wherein  $-C_1-C_3$  alkyl,  $-SC_1-C_3$  alkyl, and  $-OC_1-C_3$  alkyl are optionally substituted with 1-3 F;



$R^5$  and  $R^6$  are independently selected from the group consisting of H, halogen, -CH<sub>3</sub> and -OCH<sub>3</sub>, wherein -CH<sub>3</sub> and -OCH<sub>3</sub> are optionally substituted with 1-3 F;

$R^7$  is selected from the group consisting of H, -C<sub>1</sub>-C<sub>5</sub>alkyl, -OC<sub>1</sub>-C<sub>3</sub> alkyl, -C<sub>2</sub>-C<sub>3</sub> alkenyl, halogen, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub> alkyl, -SC<sub>1</sub>-C<sub>3</sub> alkyl, -C(=O)NR<sup>10</sup>R<sup>11</sup>,  
 5 -C(=O)H, -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, and 5-(1,2,4-oxadiazolyl), wherein -C<sub>1</sub>-C<sub>3</sub>alkyl and -C<sub>1</sub>-C<sub>5</sub>alkyl in all occurrences is optionally substituted with 1-6 substituent groups independently selected from 1-5 halogens and one -OH, -C<sub>2</sub>-C<sub>3</sub> alkenyl is optionally substituted with 1-3 halogens, and 1,2,4-oxadiazolyl and C<sub>3</sub>-C<sub>6</sub> cycloalkyl are optionally substituted with 1-2 substituent groups independently  
 10 selected from halogen and C<sub>1</sub>-C<sub>3</sub> alkyl;

$R^8$  and  $R^9$  are each independently selected from the group consisting of H, -C<sub>1</sub>-C<sub>3</sub>alkyl, halogen, -S(O)<sub>x</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -NR<sup>10</sup>R<sup>11</sup>, -OC<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>2</sub>-C<sub>3</sub>alkenyl, -NO<sub>2</sub>, -CN, -C(O)NR<sup>10</sup>R<sup>11</sup>, -C(=O)H, -NHC(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -NHS(O)<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, and pyridyl, wherein C<sub>1</sub>-C<sub>3</sub>alkyl in all occurrences is  
 15 optionally substituted with 1-3 halogens, C<sub>2</sub>-C<sub>3</sub>alkenyl is optionally substituted with 1- halogens, and C<sub>3</sub>-C<sub>6</sub>cycloalkyl and pyridyl are optionally substituted with 1-2 substituent groups independently selected from halogen and C<sub>1</sub>-C<sub>3</sub>alkyl;

$R^{10}$  and  $R^{11}$  are each H or C<sub>1</sub>-C<sub>3</sub>alkyl;

Z is selected from the group consisting of -C(=O)C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)OC<sub>1</sub>-C<sub>3</sub>alkyl, -S(O)<sub>y</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -C(=O)H, -C(O)NR<sup>10</sup>R<sup>11</sup>, -C(=O)SC<sub>1</sub>-C<sub>3</sub>alkyl,  
 20 and -C(=S)OC<sub>1</sub>-C<sub>3</sub> alkyl;

n is an integer selected from 0 and 1;

x is an integer selected from 0, 1 and 2; and

y is an integer selected from 0 and 1.

25 6. A compound of Claim 5, wherein

$R^1$  is H or CH<sub>3</sub>;

$R^2$  is selected from the group consisting of H, -CF<sub>3</sub>, -OCH<sub>3</sub>, -NO<sub>2</sub>, -CN, halogen, -NH<sub>2</sub> and 3-pyridyl;

$R^4$  is selected from the group consisting of H, halogen, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -OCH<sub>3</sub>,  
 30 -OC<sub>2</sub>H<sub>5</sub>, -SCH<sub>3</sub>, and -CN;

$R^5$  is selected from the group consisting of H and F;

$R^6$  is selected from the group consisting of H, F, -CH<sub>3</sub>, and -OCH<sub>3</sub>;

$R^7$  is selected from the group consisting of H, C<sub>1</sub>-C<sub>4</sub>alkyl, -C(=O)H, -C(=O)CH<sub>3</sub>, -CH=CH<sub>2</sub>, -CN, Cl, F, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl, -OCH<sub>3</sub>, -SCH<sub>3</sub>, -C(=O)NR<sup>10</sup>R<sup>11</sup>, and


3-methyl-5-(1,2,4-oxadiazolyl), wherein  $C_1$ - $C_4$ alkyl is optionally substituted with 1-6 substituents which are independently selected from 1-5 F and one -OH;

$R^8$  and  $R^9$  are each independently selected from the group consisting of H,  $C_1$ - $C_2$ alkyl, which is optionally substituted with 1-3 F; halogen; -CN; -NO<sub>2</sub>; -S(O)<sub>x</sub>CH<sub>3</sub>, which is optionally substituted with 1-3 F; -OCH<sub>3</sub>, which is optionally substituted with 1-3 F; -CH=CH<sub>2</sub>; -C(=O)H; -C(=O)NR<sup>10</sup>R<sup>11</sup>; -CO<sub>2</sub>H; -NR<sup>10</sup>R<sup>11</sup>; -CO<sub>2</sub>C<sub>1</sub>-C<sub>3</sub>alkyl; -NHC(O)CH<sub>3</sub>; -NHS(O)<sub>2</sub>CH<sub>3</sub>; and 4-pyridyl; and


$R^{10}$  and  $R^{11}$  are each independently selected from H and CH<sub>3</sub>, or a pharmaceutically acceptable salt thereof.

7. A compound of Claim 6, wherein n is 0, or a pharmaceutically acceptable salt thereof.


8. A compound of Claim 1, which is selected from the following compounds, or a pharmaceutically acceptable salt thereof:




Ex. 10




Ex. 11




Ex. 12




Ex. 13




Ex. 14




Ex. 19




Ex. 22




Ex. 28




Ex. 29




Ex.30




Ex. 31




Ex. 32




Ex. 33




Ex. 34




Ex. 35




Ex. 36




Ex. 37




Ex. 38




Ex. 39




Ex. 40




Ex. 41




Ex. 42




Ex. 43




Ex. 44




Ex. 45




Ex. 46




Ex. 47




Ex. 48




Ex. 134




Ex. 135









Ex. 136









Ex. 137










<p>Ex. 138</p> 	<p>Ex. 139</p> 
<p>Ex. 140</p> 	<p>Ex. 141</p> 
<p>Ex. 142</p> 	<p>Ex. 143</p> 


<p>Ex. 144</p> 	<p>Ex. 145</p> 
<p>Ex. 146</p> 	<p>Ex. 147</p> 
<p>Ex. 148</p> 	<p>Ex. 149</p> 


Ex. 150

































A <sup>1</sup>	R <sup>2</sup>
	OMe
	OMe
	F
	F
	F
	H
	Cl
	H
	Cl









9. A method of treating atherosclerosis in a patient in need of treatment comprising the administration of a therapeutically effective amount of the compound of any one of Claims 1 to 8, or a pharmaceutically acceptable salt thereof to said patient.

5 10. A method of raising HDL-C in a patient in need of treatment comprising the administration of a therapeutically effective amount of the compound of any one of Claims 1 to 8, or a pharmaceutically acceptable salt thereof to said patient.

11. A pharmaceutical composition comprising a compound of any one of Claims 1 to 8, or a pharmaceutically acceptable salt thereof, and a pharmaceutically  
10 acceptable carrier.

12. The use of a compound of any one of Claims 1 to 8 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of atherosclerosis.

13. A pharmaceutical composition comprising a compound of any one of  
15 Claims 1 to 8, or a pharmaceutically acceptable salt thereof, and one or more active ingredients selected from the group consisting of:

- (a) PPAR gamma agonists and partial agonists;
- (b) biguanides;
- (c) protein tyrosine phosphatase-1B (PTP-1B) inhibitors,
- 20 (d) dipeptidyl peptidase IV (DP-IV) inhibitors;
- (e) insulin or insulin mimetics;
- (f) sulfonylureas;
- (g)  $\alpha$ -glucosidase inhibitors;
- (h) one or more compounds selected from the group consisting of
- 25 (a) HMG-CoA reductase inhibitors; (b) bile acid sequestrants; (c) niacin, nicotiny alcohol, nicotinamide, and nicotinic acid or a salt thereof; (d) PPAR $\alpha$  agonists; (e) cholesterol absorption inhibitors; (f) acyl CoA:cholesterol acyltransferase (ACAT) inhibitors; (g) phenolic anti-oxidants, such as probucol, and (h) a microsomal triglyceride transfer protein (MTP)/ApoB secretion inhibitor;

- (i) PPAR $\alpha$ / $\gamma$ dual agonists;
- (j) PPAR $\delta$  agonists;
- (k) antiobesity compounds
- (l) ileal bile acid transporter inhibitors;
- 5 (m) anti-inflammatory agents;
- (n) glucagon receptor antagonists;
- (o) GLP-1,
- (p) GIP-1, and
- (q) GLP-1 analogs.

10

**Dated 4 May, 2011  
Merck Sharp & Dohme Corp.**

**Patent Attorneys for the Applicant/Nominated Person  
SPRUSON & FERGUSON**