



- (51) **International Patent Classification:**
C12N 9/20 (2006.01) *C11D 3/386* (2006.01)
- (21) **International Application Number:**
PCT/EP2015/080463
- (22) **International Filing Date:**
18 December 2015 (18.12.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
14199760.1 22 December 2014 (22.12.2014) EP
- (71) **Applicant:** NOVOZYMES A/S [DK/DK]; Krogshoejvej 36, 2880 Bagsvaerd (DK).
- (72) **Inventors:** BORCH, Kim; Krogshoejvej 36, 2880 Bagsvaerd (DK). VIND, Jesper; Krogshoejvej 36, 2880 Bagsvaerd (DK).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
— with sequence listing part of description (Rule 5.2(a))



WO 2016/102356 A1

(54) **Title:** DETERGENT COMPOSITIONS, LIPASE VARIANTS AND POLYNUCLEOTIDES ENCODING SAME

(57) **Abstract:** The present invention relates to detergent compositions comprising lipase variants. The present invention also relates to lipase variants and polynucleotides encoding the variants; nucleic acid constructs, vectors, and host cells comprising the polynucleotides; and methods of using the variants.

**DETERGENT COMPOSITIONS,
LIPASE VARIANTS AND POLYNUCLEOTIDES ENCODING SAME**

REFERENCE TO A SEQUENCE LISTING

5 This application contains a Sequence Listing in computer readable form, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION**Field of the Invention**

10 The present invention relates to lipase variants, polynucleotides encoding the variants, methods of producing the variants, and methods of using the variants.

Description of the Related Art

15 Lipases are important biocatalysts which have shown to be useful for various applications and a large number of different lipases have been identified and many commercialized. However, new lipases suitable for use in various compositions adapted to conditions currently used are desirable.

20 Lipases have been employed in compositions for the removal of lipid stains by hydrolyzing triglycerides to generate fatty acids. Current detergent, cleaning and/or fabric care compositions comprise many active ingredients which are interfering with the ability of lipases to remove lipid stains and under certain circumstances be detrimental for lipase activity. Stability of lipase to resist the interference from such ingredients is desirable. Furthermore, such compositions are not used immediately after production and as a consequence the stability of the lipases may be affected during storage. Thus, a need exists for lipases that are active and
25 stable in the harsh environment of detergent compositions.

30 The present invention provides lipase variants with improved properties compared to its parent. In particular the invention relates to variants comprising a cysteine-cysteine (C-C) bridge. WO99/42566 describes C-C variants wherein an N-terminal amino acid extension comprising cystein was added which could form a C-C bridge with a cystein comprised in the lipase amino acid sequence.

SUMMARY OF THE INVENTION

35 The present invention relates to a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of SEQ ID NO: 2.

The present invention also relates to polynucleotides encoding the variants; nucleic acid constructs, vectors, and host cells comprising the polynucleotides; and methods of producing the variants. The present invention also relates to methods of using the variants.

5 Definitions

Lipase: The terms “lipase”, “lipase enzyme”, “lipolytic enzyme”, “lipid esterase”, “lipolytic polypeptide”, and “lipolytic protein” refers to an enzyme in class EC3.1.1 as defined by Enzyme Nomenclature. It may have lipase activity (triacylglycerol lipase, EC3.1.1.3), cutinase activity (EC3.1.1.74), sterol esterase activity (EC3.1.1.13) and/or wax-ester hydrolase activity (EC3.1.1.50). For purposes of the present invention, lipase activity is determined according to the procedure described in the Example section. In one aspect, the variants of the present invention have at least 20%, *e.g.*, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or 100% of the lipase activity of the mature polypeptide of SEQ ID NO: 2.

Allelic variant: The term “allelic variant” means any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

cDNA: The term “cDNA” means a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic or prokaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps, including splicing, before appearing as mature spliced mRNA.

Coding sequence: The term “coding sequence” means a polynucleotide, which directly specifies the amino acid sequence of a variant. The boundaries of the coding sequence are generally determined by an open reading frame, which begins with a start codon such as ATG, GTG or TTG and ends with a stop codon such as TAA, TAG, or TGA. The coding sequence may be a genomic DNA, cDNA, synthetic DNA, or a combination thereof.

Control sequences: The term “control sequences” means nucleic acid sequences necessary for expression of a polynucleotide encoding a variant of the present invention. Each control sequence may be native (*i.e.*, from the same gene) or foreign (*i.e.*, from a different gene) to the polynucleotide encoding the variant or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The

control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the polynucleotide encoding a variant.

Expression: The term “expression” includes any step involved in the production of a variant including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

Expression vector: The term “expression vector” means a linear or circular DNA molecule that comprises a polynucleotide encoding a variant and is operably linked to control sequences that provide for its expression.

Fragment: The term “fragment” means a polypeptide having one or more (e.g., several) amino acids absent from the amino and/or carboxyl terminus of a polypeptide; wherein the fragment has lipase activity. In one aspect, a fragment contains at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% but less than 100% of the number of amino acids 1 to 269 of the mature polypeptide of a parent lipase. In one aspect the parent lipase is SEQ ID NO: 2; SEQ ID NO: 4; or SEQ ID NO: 6.

High stringency conditions: The term “high stringency conditions” means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 65°C.

Host cell: The term “host cell” means any cell type that is susceptible to transformation, transfection, transduction, or the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention. The term “host cell” encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication.

Improved property: The term “improved property” means a characteristic associated with a variant that is improved compared to the parent. The improved property of the variants of the present invention is improved stability as compared to the parent enzyme or the mature polypeptide of SEQ ID NO: 2. In one aspect the parent enzyme is SEQ ID NO: 2; SEQ ID NO: 4; or SEQ ID NO: 6. In one aspect the stability of the present invention may be thermostability, stability in the presence of proteolytic enzymes, stability in the presence of surfactant, stability in the presence of a reducing agent, stability in detergent compositions, stability under storage conditions, stability under storage conditions in the presence of protease, stability under storage conditions in the presence of a reducing agent, or detergent stability. The property is determined by the assay described in the section “Examples”.

Isolated: The term “isolated” means a substance in a form or environment which does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally

occurring substance, (2) any substance including, but not limited to, any enzyme, variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated (e.g., multiple copies of a gene encoding the substance; use of a stronger promoter than the promoter naturally associated with the gene encoding the substance). An isolated substance may be present in a fermentation broth sample.

Low stringency conditions: The term "low stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 50°C.

Mature polypeptide: The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 1 to 269 of SEQ ID NO: 2, 1 to 274 of SEQ ID NO: 4, or 1 to 269 of SEQ ID NO: 6. It is known in the art that a host cell may produce a mixture of two or more different mature polypeptides (i.e., with a different C-terminal and/or N-terminal amino acid) expressed by the same polynucleotide.

Mature polypeptide coding sequence: The term "mature polypeptide coding sequence" means a polynucleotide that encodes a mature polypeptide having lipase activity. In one aspect, the mature polypeptide coding sequence is nucleotides 67 to 873 of SEQ ID NO: 1; 67 to 888 of SEQ ID NO: 3; or 67 to 873 of SEQ ID NO: 5.

Medium stringency conditions: The term "medium stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 55°C.

Medium-high stringency conditions: The term "medium-high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 60°C.

Mutant: The term "mutant" means a polynucleotide encoding a variant.

Nucleic acid construct: The term "nucleic acid construct" means a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or

is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic, which comprises one or more control sequences.

Operably linked: The term “operably linked” means a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide such that the control sequence directs expression of the coding sequence.

Parent or parent lipase: The term “parent” or “parent lipase” means a lipase to which an alteration is made to produce the enzyme variants of the present invention. The parent may be a naturally occurring (wild-type) polypeptide or a variant or fragment thereof.

Reducing Agent: The term “Reducing agent” means any agent or substance that can reduce i.e. break a disulfide bond present in a lipase according to the invention. Examples of reducing agents that can break a cysteine bridge (cys-bridge, C-C bridge) is such as e.g. tris(2-carboxyethyl)phosphine (TCEP), dithiothreitol (DTT), 2-mercaptoethanol (beta-ME), tris(3-hydroxypropyl)phosphine (THPP) and sulphites (such as sodium sulphite).

Sequence identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter “sequence identity”.

For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends Genet.* 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled “longest identity” (obtained using the –nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

For purposes of the present invention, the sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *supra*), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled “longest identity” (obtained using the –nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

Subsequence: The term “subsequence” means a polynucleotide having one or more (e.g., several) nucleotides absent from the 5' and/or 3' end of a mature polypeptide coding sequence; wherein the subsequence encodes a fragment having lipase activity. In one aspect, a subsequence contains at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at

least 75%, at least 80%, at least 85%, at least 90%, or at least 95% but less than 100% of the number of nucleotides 67 to 873 of SEQ ID NO: 1, 67 to 888 of SEQ ID NO: 3, or 67 to 873 of SEQ ID NO: 5.

Variant: The term “variant” means a polypeptide having lipase activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (*e.g.*, several) positions. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position. The variants of the present invention have at least 20%, *e.g.*, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 100% of the lipase activity of the mature polypeptide of SEQ ID NO: 2.

Very high stringency conditions: The term “very high stringency conditions” means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 70°C.

Very low stringency conditions: The term “very low stringency conditions” means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 45°C.

Wash performance: In the present context the term “wash performance” is used as an enzyme’s ability to remove lipid or lipid-containing stains present on the object to be cleaned. The wash performance may be quantified by calculating the so-called G/Int value defined in the description of AMSA in the Methods section below. The term “wash performance” includes cleaning in general *e.g.* hard surface cleaning as in dish wash, but also wash performance on textiles such as laundry, and also industrial and institutional cleaning.

Wild-type lipase: The term “wild-type” lipase means a lipase expressed by a naturally occurring microorganism, such as a bacterium, yeast, or filamentous fungus found in nature.

Conventions for Designation of Variants

For purposes of the present invention, the mature polypeptide disclosed in SEQ ID NO: 2 is used to determine the corresponding amino acid residue in another lipase. The amino acid sequence of another lipase is aligned with the mature polypeptide disclosed in SEQ ID NO: 2, and based on the alignment, the amino acid position number corresponding to any amino acid residue in the mature polypeptide disclosed in SEQ ID NO: 2 is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European

Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends Genet.* 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix.

Identification of the corresponding amino acid residue in another lipase can be determined by an alignment of multiple polypeptide sequences using several computer programs including, but not limited to, MUSCLE (multiple sequence comparison by log-expectation; version 3.5 or later; Edgar, 2004, *Nucleic Acids Research* 32: 1792-1797), MAFFT (version 6.857 or later; Katoh and Kuma, 2002, *Nucleic Acids Research* 30: 3059-3066; Katoh *et al.*, 2005, *Nucleic Acids Research* 33: 511-518; Katoh and Toh, 2007, *Bioinformatics* 23: 372-374; Katoh *et al.*, 2009, *Methods in Molecular Biology* 537: 39-64; Katoh and Toh, 2010, *Bioinformatics* 26: 1899-1900), and EMBOSS EMMA employing ClustalW (1.83 or later; Thompson *et al.*, 1994, *Nucleic Acids Research* 22: 4673-4680), using their respective default parameters.

When the other enzyme has diverged from the mature polypeptide of SEQ ID NO: 2 such that traditional sequence-based comparison fails to detect their relationship (Lindahl and Elofsson, 2000, *J. Mol. Biol.* 295: 613-615), other pairwise sequence comparison algorithms can be used. Greater sensitivity in sequence-based searching can be attained using search programs that utilize probabilistic representations of polypeptide families (profiles) to search databases. For example, the PSI-BLAST program generates profiles through an iterative database search process and is capable of detecting remote homologs (Atschul *et al.*, 1997, *Nucleic Acids Res.* 25: 3389-3402). Even greater sensitivity can be achieved if the family or superfamily for the polypeptide has one or more representatives in the protein structure databases. Programs such as GenTHREADER (Jones, 1999, *J. Mol. Biol.* 287: 797-815; McGuffin and Jones, 2003, *Bioinformatics* 19: 874-881) utilize information from a variety of sources (PSI-BLAST, secondary structure prediction, structural alignment profiles, and solvation potentials) as input to a neural network that predicts the structural fold for a query sequence. Similarly, the method of Gough *et al.*, 2000, *J. Mol. Biol.* 313: 903-919, can be used to align a sequence of unknown structure with the superfamily models present in the SCOP database. These alignments can in turn be used to generate homology models for the polypeptide, and such models can be assessed for accuracy using a variety of tools developed for that purpose.

For proteins of known structure, several tools and resources are available for retrieving and generating structural alignments. For example the SCOP superfamilies of proteins have been structurally aligned, and those alignments are accessible and downloadable. Two or more protein structures can be aligned using a variety of algorithms such as the distance alignment matrix (Holm and Sander, 1998, *Proteins* 33: 88-96) or combinatorial extension (Shindyalov and Bourne, 1998, *Protein Engineering* 11: 739-747), and implementation of these algorithms can

additionally be utilized to query structure databases with a structure of interest in order to discover possible structural homologs (e.g., Holm and Park, 2000, *Bioinformatics* 16: 566-567).

In describing the variants of the present invention, the nomenclature described below is adapted for ease of reference. The accepted IUPAC single letter or three letter amino acid abbreviation is employed.

Substitutions. For an amino acid substitution, the following nomenclature is used: Original amino acid, position, substituted amino acid. Accordingly, the substitution of threonine at position 226 with alanine is designated as "Thr226Ala" or "T226A". Multiple mutations are separated by addition marks ("+"), e.g., "Gly205Arg + Ser411Phe" or "G205R + S411F", representing substitutions at positions 205 and 411 of glycine (G) with arginine (R) and serine (S) with phenylalanine (F), respectively.

Deletions. For an amino acid deletion, the following nomenclature is used: Original amino acid, position, *. Accordingly, the deletion of glycine at position 195 is designated as "Gly195*" or "G195*". Multiple deletions are separated by addition marks ("+"), e.g., "Gly195* + Ser411*" or "G195* + S411*".

Insertions. For an amino acid insertion, the following nomenclature is used: Original amino acid, position, original amino acid, inserted amino acid. Accordingly the insertion of lysine after glycine at position 195 is designated "Gly195GlyLys" or "G195GK". An insertion of multiple amino acids is designated [Original amino acid, position, original amino acid, inserted amino acid #1, inserted amino acid #2; etc.]. For example, the insertion of lysine and alanine after glycine at position 195 is indicated as "Gly195GlyLysAla" or "G195GKA".

In such cases the inserted amino acid residue(s) are numbered by the addition of lower case letters to the position number of the amino acid residue preceding the inserted amino acid residue(s). In the above example, the sequence would thus be:

<u>Parent:</u>	<u>Variant:</u>
195	195 195a 195b
G	G - K - A

Multiple alterations. Variants comprising multiple alterations are separated by addition marks ("+"), e.g., "Arg170Tyr+Gly195Glu" or "R170Y+G195E" representing a substitution of arginine and glycine at positions 170 and 195 with tyrosine and glutamic acid, respectively.

Different alterations. Where different alterations can be introduced at a position, the different alterations are separated by a comma, e.g., "Arg170Tyr,Glu" represents a substitution of arginine at position 170 with tyrosine or glutamic acid. Thus, "Tyr167Gly,Ala + Arg170Gly,Ala" designates the following variants:

"Tyr167Gly+Arg170Gly", "Tyr167Gly+Arg170Ala", "Tyr167Ala+Arg170Gly", and "Tyr167Ala+Arg170Ala".

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides lipase variants with improved stability as compared to the parent lipase or to the mature polypeptide of SEQ ID NO: 2.

5 Variants

The present invention relates to a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase.

10 In one aspect the parent lipase is a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90 % identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a
15 peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to
20 positions 90-101 of said wild-type lipase.

In one aspect the parent lipase is a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2.

25 In one aspect the parent lipase has the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the parent lipase comprises or consists the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the variant is a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at
30 least 60% but less than 100% sequence identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

In one aspect, the variant has sequence identity of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%,
35 but less than 100%, to the amino acid sequence of the parent lipase.

In one aspect, the variant has sequence identity of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at

least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, but less than 100%, to sequence identity to the mature polypeptide of SEQ ID NO: 2.

In one aspect, the number of substitutions in the variants of the present invention is 1-40, *e.g.*, 1-30, 1-20, 1-10 and 1-5, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 substitutions.

In one aspect, the variant comprises or consists of the substitutions E1C +N233C of the mature polypeptide of SEQ ID NO: 2, or of a polypeptide having at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to the mature polypeptide of SEQ ID NO: 2 which has lipase activity, and further the variant has improved stability as compared to the mature lipase of SEQ ID NO: 2. In one aspect the stability of the present invention may be thermostability, stability in the presence of proteolytic enzymes, stability in the presence of surfactant, stability in the presence of a reducing agent, stability in detergent compositions, stability under storage conditions, stability under storage conditions in the presence of protease, stability under storage conditions in the presence of a reducing agent, or detergent stability.

The variants may further comprise one or more additional substitution at one or more (*e.g.*, several) other positions. For example, in one aspect the variant further comprises one or more (*e.g.*, several) substitutions corresponding to any of positions selected from: 2, 4, 8, 11, 15, 27, 33, 38, 43, 48, 51, 54, 56, 57, 58, 60, 69, 71, 83, 86, 91, 92, 94, 96, 97, 98, 99, 101, 111, 123, 150, 152, 163, 176, 179, 187, 188, 189, 198, 199, 200, 210, 216, 220, 224, 225, 227, 228, 229, 231, 236, 238, 239, 246, 249, 254, 255, 256, 257, 260, 263, 264, 265, 266, 267, 269 of SEQ ID NO: 2. In one aspect the variant further comprises one or more (*e.g.*, several) substitutions corresponding to any of positions selected from: V2K, Q4R, Q4V, N8R, N11R, Q15C, D27G, D27R, N33K, N33Q, G38A, E43C, D48C, F51V, S54T, E56K, D57G, S58A, V60K, V60S, L69R, N71C, S83T, I86V, G91A, G91N, G91Q, N92D, N94K, N94R, D96E, D96G, D96L, D96W, L97M, K98E, K98I, K98Q, E99K, E99N, N101D, N101S, D111A, T123V, A150G, A152G, G163K, V176L, R179L, V187Y, V187W, Q188R, T189Y, T189W, H198S, T199R, N200R, E210K, E210Q, S216P, Y220F, S224R, G225R, L227G, L227R, V228R, P229R, T231R, V236R, I238C, E239C, G246C, Q249R, D254S, I255G, P256K, P256T, P256V, A257I, A257V, W260C, G263Q, L264A, I265T, G266D, T267A, L269N, L269V of SEQ ID NO: 2. In one aspect the variant comprises or consists of sets of substitutions at positions corresponding to: E1C N233C; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K N233C D254S P256T; E1C V2K D27G N33K G38A F51V D96E D111A G163K N233C D254S P256T; E1C V2K D27R N33K G38A F51V D96E D111A G163K Q188R N233C D254S P256T; E1C D27R G38A G91A N92D D96L K98Q D111A G163K N233C D254S P256T; E1C D27R G38A G91N N94R D96E D111A G163K S216P L227G N233C D254S P256T; E1C T231R N233C;

E1C T231R N233C Q249R D254S; E1C G225R T231R N233C; E1C Q15C E43C T231R N233C; E1C L227R T231R N233C; E1C P229R T231R N233C; E1C L227G T231R N233C; E1C E99N N101S T231R N233C; E1C L227G T231R N233C D254S; E1C E210K L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K

5 Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K D57G L69R D96E K98I D111A A152G G163K T231R N233C D254S P256T; E1C V187Y T189Y L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C V60K I86V A150G E210K L227G T231R N233C P256K; E1C V187W T189W L227G T231R N233C; E1C N94K D96L L227G T231R

10 N233C; E1C G91A N92D D96L K98Q L227G T231R N233C; E1C N8R L227G T231R N233C; E1C L227G V228R T231R N233C; E1C Q4R L227G T231R N233C; E1C N11R L227G T231R N233C; E1C S224R L227G T231R N233C; E1C L227G T231R N233C V236R; E1C N200R L227G T231R N233C; E1C T199R L227G T231R N233C; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K

15 D96E K98I D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T A257I; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T A257I;

20 E1C D27R N33K G38A F51V S54T E56K D57G D96E K98I D111A G163K T231R N233C D254S I255G P256T A257V L269V; E1C V2K D27R N33K G38A F51V L69R D96E K98E D111A G163K V176L E210K L227G T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K

25 Y220F T231R N233C D254S P256T; E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T; and E1C N11R D27R N33K D48C F51V L69R N71C E87Q K98E N101R T143A E210K G225R L227G P229R T231R N233C Q249R P250R D254S I255G P256K of SEQ ID NO: 2.

In one aspect the variants according to the invention comprises or consists of one or more (e.g. several) further cystein bridge(s) in addition to the cystein bridge at a position corresponding to E1C N233C of SEQ ID NO: 2. In one aspect the variants comprises or consists of one, two, three, four, five, six, seven, eight, nine, or ten further cystein bridge(s). Such further cystein bridges may link a N- or C-terminal extention comprising a Cystein to the mature part of the lipase as has been described in WO99/42566 and/or the cystein bridge may

35 link different parts of the mature lipase as in naturally occurring proteins. Proteins naturally comprise cysteine bridges and e.g. the lipase of SEQ ID NO: 2 has three cysteine bridges located at positions C22-C268, C36-C41, and C104-C107. It should be understood that the extentions comprising a cystein mentioned in WO99/42566 in one aspect may be added to the

variants of the invention together with a substitution E239C for the purpose of providing an additional cystein bridge. Examples of other cystein bridges to be included in the variants of the invention may be e.g. at positions corresponding to Q15C E43C and/or D48C N71C of SEQ ID NO: 2.

5 The amino acid changes may be of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of 1-30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as
10 a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine,
15 alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

20 Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in a polypeptide can be identified according to procedures known
25 in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for lipase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton *et al.*, 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other
30 biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Lett.* 309: 59-64. The identity of essential amino acids can
35 also be inferred from an alignment with a related polypeptide.

In one aspect the variant has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 100% of the activity of the parent lipase.

In one aspect, the variant has improved stability.

In one aspect, the variant has improved thermostability compared to the parent enzyme.

In one aspect the variant is more stable at increased temperature compared to the parent lipase. Thermostability is determined by using the DSC assay as described in Example 1.

5 In one aspect, the variant has improved stability in the presence of proteolytic enzymes compared to the parent enzyme. In one aspect, the variant is more stable in the presence of proteolytic enzymes compared to the parent lipase. In one aspect the variant is more stable in the presence of proteolytic enzymes at increased temperature compared to the parent lipase. Improved stability in the presence of proteolytic enzymes may be determined by using the DSC
10 assay as described in Example 1.

In one aspect, the variant has improved stability in the presence of surfactant compared to the parent enzyme. In one aspect, the variant is more stable in the presence of surfactant compared to the parent lipase. In one aspect the variant is more stable in the presence of surfactant at increased temperature compared to the parent lipase. Improved stability in the
15 presence of surfactant may be determined by using the DSC assay as described in Example 1.

In one aspect, the variant has improved stability in the presence of a reducing agent compared to the parent enzyme. In one aspect, the variant is more stable in the presence of a reducing agent compared to the parent lipase. In one aspect the variant is more stable in the presence of a reducing agent at increased temperature compared to the parent lipase. Improved stability in the presence of a reducing agent may be determined by using the
20 NanoDSF assay as described in Example 1.

In one aspect, the variant has improved stability in detergent compositions compared to the parent enzyme. In one aspect, the variant is more stable in detergent compositions compared to the parent lipase. In one aspect the variant is more stable in detergent
25 compositions at increased temperature compared to the parent lipase. Improved stability in detergent compositions may be determined by using the DSC assay as described in Example 1.

In one aspect, the variant has improved stability under storage conditions (i.e. has improved storage stability) compared to the parent enzyme. In one aspect, the variant is more stable under storage conditions compared to the parent lipase. In one aspect the variant is more
30 stable under storage conditions at increased temperature compared to the parent lipase. Improved stability under storage conditions may be determined by using the Storage stability assay protocol A as described in Example 1.

In one aspect, the variant has improved stability under storage conditions (i.e. has improved storage stability) in the presence of protease compared to the parent enzyme. In one
35 aspect, the variant is more stable under storage conditions in the presence of protease compared to the parent lipase. In one aspect the variant is more stable under storage conditions in the presence of protease at increased temperature compared to the parent lipase. Improved

stability under storage conditions in the presence of protease may be determined by using the Storage stability assay protocol B as described in Example 1.

In one aspect, the variant has improved stability under storage conditions (i.e. has improved storage stability) in the presence of a reducing agent compared to the parent enzyme.

5 In one aspect, the variant is more stable under storage conditions in the presence of a reducing agent compared to the parent lipase. In one aspect the variant is more stable under storage conditions in the presence of a reducing agent at increased temperature compared to the parent lipase. Improved stability under storage conditions in the presence of a reducing agent may be determined by using a modified version of the Storage stability assay as described in Example
10 1.

In one aspect, the variant has improved detergent stability compared to the parent enzyme. In one aspect, the variant is more stable in detergent compared to the parent lipase. In one aspect the variant is more stable in detergent at increased temperature compared to the parent lipase. Improved stability in detergent may be determined by using the DSC assay or the
15 NanoDSF assay as described in Example 1.

Parent lipases

The parent lipase is (a) a polypeptide having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6; (b) a polypeptide
20 encoded by a polynucleotide that hybridizes under low stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, or (ii) the full-length complement of (i); or (c) a polypeptide encoded by a polynucleotide having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5.

25 In one aspect, the parent has a sequence identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have lipase activity. In one aspect, the amino acid sequence of the parent differs by up to 40 amino
30 acids, e.g., 1-40, 1-30, 1-20, 1-10, 1-5, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 from the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect, the parent comprises or consists of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect, the parent comprises or consists of the mature polypeptide of
35 SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect, the parent comprises or consists of amino acids 1 to 269 of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

In one aspect, the parent is a fragment of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 containing at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%,

at least 80%, at least 85%, at least 90%, or at least 95% but less than 100% of the number of amino acids 1 to 269 of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect, the parent is an allelic variant of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

5 In one aspect, the parent is encoded by a polynucleotide that hybridizes under very low stringency conditions, low stringency conditions, medium stringency conditions, medium-high stringency conditions, high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, or (ii) the full-length complement of (i) (Sambrook *et al.*, 1989, *Molecular Cloning, A Laboratory*
10 *Manual*, 2d edition, Cold Spring Harbor, New York).

The polynucleotide of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5 or a subsequence thereof, as well as the polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 or a fragment thereof, may be used to design nucleic acid probes to identify and clone DNA encoding a parent from strains of different genera or species according to methods well known
15 in the art. In particular, such probes can be used for hybridization with the genomic DNA or cDNA of a cell of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 15, *e.g.*, at least 25, at least 35, or at least 70 nucleotides in length. Preferably, the nucleic acid probe is at least 100 nucleotides in length,
20 *e.g.*, at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, or at least 800 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

25 A genomic DNA or cDNA library prepared from such other strains may be screened for DNA that hybridizes with the probes described above and encodes a parent. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In
30 order to identify a clone or DNA that hybridizes with SEQ ID NO: 1 or a subsequence thereof, the carrier material is used in a Southern blot.

For purposes of the present invention, hybridization indicates that the polynucleotide hybridizes to a labeled nucleic acid probe corresponding to (i) SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5; (ii) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or
35 SEQ ID NO: 5; (iii) the full-length complement thereof; or (iv) a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film or any other detection means known in the art.

In one aspect, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5. In one aspect, the nucleic acid probe is nucleotides 67 to 873 of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5. In one aspect, the nucleic acid probe is a polynucleotide that encodes the polypeptide of SEQ ID NO: 2; the mature polypeptide thereof; or a fragment thereof. In one aspect, the nucleic acid probe is SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5.

In one aspect, the parent is encoded by a polynucleotide having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%.

In one aspect the parent lipase is a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90 % identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase.

In one aspect the parent lipase is a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2.

In one aspect the parent is the polypeptide with the sequence of SEQ ID NO: 2. In one aspect the parent is the polypeptide with the sequence of SEQ ID NO: 4. In one aspect the parent is the polypeptide with the sequence of SEQ ID NO: 6.

The polypeptide may be a hybrid polypeptide in which a region of one polypeptide is fused at the N-terminus or the C-terminus of a region of another polypeptide.

The parent may be a fusion polypeptide or cleavable fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of the present invention. A fusion polypeptide is produced by fusing a polynucleotide encoding another polypeptide to a polynucleotide of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fusion polypeptide is under control of the same promoter(s) and terminator. Fusion polypeptides may also be constructed

using intein technology in which fusion polypeptides are created post-translationally (Cooper *et al.*, 1993, *EMBO J.* 12: 2575-2583; Dawson *et al.*, 1994, *Science* 266: 776-779).

A fusion polypeptide can further comprise a cleavage site between the two polypeptides. Upon secretion of the fusion protein, the site is cleaved releasing the two polypeptides.

5 Examples of cleavage sites include, but are not limited to, the sites disclosed in Martin *et al.*, 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-576; Svetina *et al.*, 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson *et al.*, 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward *et al.*, 1995, *Biotechnology* 13: 498-503; and Contreras *et al.*, 1991, *Biotechnology* 9: 378-381; Eaton *et al.*, 1986, *Biochemistry* 25: 505-512; Collins-Racie *et al.*, 1995, *Biotechnology* 13: 982-987; Carter
10 *et al.*, 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248; and Stevens, 2003, *Drug Discovery World* 4: 35-48.

The parent may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the parent encoded by a polynucleotide is produced by the source or by a strain
15 in which the polynucleotide from the source has been inserted. In one aspect, the parent is secreted extracellularly.

The parent may be a bacterial lipase. For example, the parent may be a Gram-positive bacterial polypeptide such as a *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, or *Streptomyces* lipase, or a
20 Gram-negative bacterial polypeptide such as a *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, or *Ureaplasma* lipase.

In one aspect, the parent is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*,
25 *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* lipase.

In one aspect, the parent is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* lipase.

In one aspect, the parent is a *Streptomyces achromogenes*, *Streptomyces avermitilis*,
30 *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* lipase.

The parent may be a fungal lipase. For example, the parent may be a yeast lipase such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* lipase; or a filamentous fungal lipase such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*,
35 *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*,

Pseudotrichonympha, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* lipase.

In one aspect, the parent is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*,
 5 *Saccharomyces norbensis*, or *Saccharomyces oviformis* lipase.

In one aspect, the parent is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*,
 10 *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*,
 15 *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Thielavia achromatica*, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia microspora*, *Thielavia ovispora*,
 20 *Thielavia peruviana*, *Thielavia setosa*, *Thielavia spededonium*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* lipase.

In one aspect, the parent is a *Humicola lanuginosa* lipase, e.g., the lipase of SEQ ID NO: 2 or the mature polypeptide thereof.

25 It will be understood that for the aforementioned species, the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

30 Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

The parent may be identified and obtained from other sources including microorganisms
 35 isolated from nature (e.g., soil, composts, water, etc.) or DNA samples obtained directly from natural materials (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms and DNA directly from natural habitats are well known in the art. A polynucleotide encoding a parent may then be obtained by similarly screening a

genomic DNA or cDNA library of another microorganism or mixed DNA sample. Once a polynucleotide encoding a parent has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are known to those of ordinary skill in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

5

Preparation of Variants

The present invention also relates to methods for obtaining a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of SEQ ID NO: 2.; and (b) recovering the variant.

The variants can be prepared using any mutagenesis procedure known in the art, such as site-directed mutagenesis, synthetic gene construction, semi-synthetic gene construction, random mutagenesis, shuffling, etc.

15 Site-directed mutagenesis is a technique in which one or more (e.g., several) mutations are introduced at one or more defined sites in a polynucleotide encoding the parent.

Site-directed mutagenesis can be accomplished *in vitro* by PCR involving the use of oligonucleotide primers containing the desired mutation. Site-directed mutagenesis can also be performed *in vitro* by cassette mutagenesis involving the cleavage by a restriction enzyme at a site in the plasmid comprising a polynucleotide encoding the parent and subsequent ligation of an oligonucleotide containing the mutation in the polynucleotide. Usually the restriction enzyme that digests the plasmid and the oligonucleotide is the same, permitting sticky ends of the plasmid and the insert to ligate to one another. See, e.g., Scherer and Davis, 1979, *Proc. Natl. Acad. Sci. USA* 76: 4949-4955; and Barton *et al.*, 1990, *Nucleic Acids Res.* 18: 7349-4966.

25 Site-directed mutagenesis can also be accomplished *in vivo* by methods known in the art. See, e.g., U.S. Patent Application Publication No. 2004/0171154; Storici *et al.*, 2001, *Nature Biotechnol.* 19: 773-776; Kren *et al.*, 1998, *Nat. Med.* 4: 285-290; and Calissano and Macino, 1996, *Fungal Genet. Newslett.* 43: 15-16.

Any site-directed mutagenesis procedure can be used in the present invention. There are many commercial kits available that can be used to prepare variants.

Synthetic gene construction entails *in vitro* synthesis of a designed polynucleotide molecule to encode a polypeptide of interest. Gene synthesis can be performed utilizing a number of techniques, such as the multiplex microchip-based technology described by Tian *et al.* (2004, *Nature* 432: 1050-1054) and similar technologies wherein oligonucleotides are synthesized and assembled upon photo-programmable microfluidic chips.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988,

Science 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO95/17413; or WO95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman *et al.*, 1991, *Biochemistry* 30: 10832-10837; US5,223,409; WO92/06204) and region-directed mutagenesis (Derbyshire *et al.*, 1986, *Gene* 46: 145; Ner *et al.*, 1988, *DNA* 7: 127).

Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness *et al.*, 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

Semi-synthetic gene construction is accomplished by combining aspects of synthetic gene construction, and/or site-directed mutagenesis, and/or random mutagenesis, and/or shuffling. Semi-synthetic construction is typified by a process utilizing polynucleotide fragments that are synthesized, in combination with PCR techniques. Defined regions of genes may thus be synthesized *de novo*, while other regions may be amplified using site-specific mutagenic primers, while yet other regions may be subjected to error-prone PCR or non-error prone PCR amplification. Polynucleotide subsequences may then be shuffled.

Polynucleotides

The present invention also relates to polynucleotides encoding a variant of the present invention.

Nucleic Acid Constructs

The present invention also relates to nucleic acid constructs comprising a polynucleotide encoding a variant of the present invention operably linked to one or more control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

The polynucleotide may be manipulated in a variety of ways to provide for expression of a variant. Manipulation of the polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides utilizing recombinant DNA methods are well known in the art.

The control sequence may be a promoter, a polynucleotide which is recognized by a host cell for expression of the polynucleotide. The promoter contains transcriptional control sequences that mediate the expression of the variant. The promoter may be any polynucleotide that shows transcriptional activity in the host cell including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a bacterial host cell are the promoters obtained from the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus stearothermophilus* maltogenic
 5 amylase gene (*amyM*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus subtilis* *xylA* and *xylB* genes, *Bacillus thuringiensis* *cryIIIA* gene (Agaisse and Lereclus, 1994, *Molecular Microbiology* 13: 97-107), *E. coli* *lac* operon, *E. coli* *trc* promoter (Egon *et al.*, 1988, *Gene* 69: 301-315), *Streptomyces coelicolor* agarase gene (*dagA*), and prokaryotic beta-lactamase gene (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75: 3727-3731), as well as the *tac*
 10 promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. USA* 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in Gilbert *et al.*, 1980, *Scientific American* 242: 74-94; and in Sambrook *et al.*, 1989, *supra*. Examples of tandem promoters are disclosed in WO99/43835.

Examples of suitable promoters for directing transcription of the nucleic acid constructs
 15 of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus nidulans* acetamidase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Aspergillus oryzae* TAKA amylase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Fusarium oxysporum* trypsin-like protease (WO96/00787),
 20 *Fusarium venenatum* amyloglucosidase (WO00/56900), *Fusarium venenatum* Daria (WO00/56900), *Fusarium venenatum* Quinn (WO00/56900), *Rhizomucor miehei* lipase, *Rhizomucor miehei* aspartic proteinase, *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III,
 25 *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a modified promoter from an *Aspergillus* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus* triose phosphate isomerase gene; non-limiting examples include modified promoters
 30 from an *Aspergillus niger* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus nidulans* or *Aspergillus oryzae* triose phosphate isomerase gene); and mutant, truncated, and hybrid promoters thereof.

In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces*
 35 *cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase.

Other useful promoters for yeast host cells are described by Romanos *et al.*, 1992, *Yeast* 8: 423-488.

The control sequence may also be a transcription terminator, which is recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3'-terminus of the polynucleotide encoding the variant. Any terminator that is functional in the host cell may be used.

Preferred terminators for bacterial host cells are obtained from the genes for *Bacillus clausii* alkaline protease (*aprH*), *Bacillus licheniformis* alpha-amylase (*amyL*), and *Escherichia coli* ribosomal RNA (*rrnB*).

Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

The control sequence may also be an mRNA stabilizer region downstream of a promoter and upstream of the coding sequence of a gene which increases expression of the gene.

Examples of suitable mRNA stabilizer regions are obtained from a *Bacillus thuringiensis cryIIIA* gene (WO94/25612) and a *Bacillus subtilis* SP82 gene (Hue *et al.*, 1995, *Journal of Bacteriology* 177: 3465-3471).

The control sequence may also be a leader, a nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is operably linked to the 5'-terminus of the polynucleotide encoding the variant. Any leader that is functional in the host cell may be used.

Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3'-terminus of the variant-encoding sequence and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell may be used.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase,

Aspergillus niger alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Mol. Cellular Biol.* 15: 5983-5990.

5 The control sequence may also be a signal peptide coding region that encodes a signal peptide linked to the N-terminus of a variant and directs the variant into the cell's secretory pathway. The 5'-end of the coding sequence of the polynucleotide may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the variant. Alternatively, the 5'-end of the coding sequence
10 may contain a signal peptide coding sequence that is foreign to the coding sequence. A foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, a foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the variant. However, any signal peptide coding sequence that directs the
15 expressed variant into the secretory pathway of a host cell may be used.

Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus stearothermophilus* neutral proteases (*nprT*, *nprS*,
20 *nprM*), and *Bacillus subtilis* *prsA*. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, *Humicola insolens* cellulase,
25 *Humicola insolens* endoglucanase V, *Humicola lanuginosa* lipase, and *Rhizomucor miehei* aspartic proteinase.

Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos *et al.*, 1992, *supra*.

30 The control sequence may also be a propeptide coding sequence that encodes a propeptide positioned at the N-terminus of a variant. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to an active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from
35 the genes for *Bacillus subtilis* alkaline protease (*aprE*), *Bacillus subtilis* neutral protease (*nprT*), *Myceliophthora thermophila* laccase (WO 95/33836), *Rhizomucor miehei* aspartic proteinase, and *Saccharomyces cerevisiae* alpha-factor.

Where both signal peptide and propeptide sequences are present, the propeptide sequence is positioned next to the N-terminus of the variant and the signal peptide sequence is positioned next to the N-terminus of the propeptide sequence.

It may also be desirable to add regulatory sequences that regulate expression of the variant relative to the growth of the host cell. Examples of regulatory systems are those that cause expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the *Aspergillus niger* glucoamylase promoter, *Aspergillus oryzae* TAKA alpha-amylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the polynucleotide encoding the variant would be operably linked with the regulatory sequence.

Expression Vectors

The present invention also relates to recombinant expression vectors comprising a polynucleotide encoding a variant of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleotide and control sequences may be joined together to produce a recombinant expression vector that may include one or more convenient restriction sites to allow for insertion or substitution of the polynucleotide encoding the variant at such sites. Alternatively, the polynucleotide may be expressed by inserting the polynucleotide or a nucleic acid construct comprising the polynucleotide into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the polynucleotide. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vector may be a linear or closed circular plasmid.

The vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector

or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

The vector preferably contains one or more selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

Examples of bacterial selectable markers are *Bacillus licheniformis* or *Bacillus subtilis* *dal* genes, or markers that confer antibiotic resistance such as ampicillin, chloramphenicol, kanamycin, neomycin, spectinomycin or tetracycline resistance. Suitable markers for yeast host cells include, but are not limited to, ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are *Aspergillus nidulans* or *Aspergillus oryzae* *amdS* and *pyrG* genes and a *Streptomyces hygroscopicus* *bar* gene.

The vector preferably contains an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the variant or any other element of the vector for integration into the genome by homologous or non-homologous recombination. Alternatively, the vector may contain additional polynucleotides for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, 400 to 10,000 base pairs, and 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding polynucleotides. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" means a polynucleotide that enables a plasmid or vector to replicate *in vivo*.

Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*.

5 Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems *et al.*, 1991, *Gene* 98: 61-67; Cullen *et al.*, 1987, *Nucleic Acids Res.* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising
10 the gene can be accomplished according to the methods disclosed in WO 00/24883.

More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of a variant. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the
15 polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art
20 (see, e.g., Sambrook *et al.*, 1989, *supra*).

Host Cells

The present invention also relates to recombinant host cells, comprising a polynucleotide encoding a variant of the present invention operably linked to one or more control sequences
25 that direct the production of a variant of the present invention. A construct or vector comprising a polynucleotide is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent
30 depend upon the gene encoding the variant and its source.

The host cell may be any cell useful in the recombinant production of a variant, e.g., a prokaryote or a eukaryote.

The prokaryotic host cell may be any Gram-positive or Gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*,
35 *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*. Gram-negative bacteria include, but are not limited to, *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

The bacterial host cell may be any *Bacillus* cell including, but not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and
5 *Bacillus thuringiensis* cells.

The bacterial host cell may also be any *Streptococcus* cell including, but not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi* subsp. *Zooepidemicus* cells.

The bacterial host cell may also be any *Streptomyces* cell, including, but not limited to,
10 *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

The introduction of DNA into a *Bacillus* cell may be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Mol. Gen. Genet.* 168: 111-115), competent cell transformation (see, e.g., Young and Spizizen, 1961, *J. Bacteriol.* 81: 823-829, or Dubnau and
15 Davidoff-Abelson, 1971, *J. Mol. Biol.* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, *J. Bacteriol.* 169: 5271-5278). The introduction of DNA into an *E. coli* cell may be effected by protoplast transformation (see, e.g., Hanahan, 1983, *J. Mol. Biol.* 166: 557-580) or electroporation (see, e.g., Dower *et al.*, 1988, *Nucleic Acids Res.* 16: 6127-6145). The
20 introduction of DNA into a *Streptomyces* cell may be effected by protoplast transformation, electroporation (see, e.g., Gong *et al.*, 2004, *Folia Microbiol. (Praha)* 49: 399-405), conjugation (see, e.g., Mazodier *et al.*, 1989, *J. Bacteriol.* 171: 3583-3585), or transduction (see, e.g., Burke *et al.*, 2001, *Proc. Natl. Acad. Sci. USA* 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may be effected by electroporation (see, e.g., Choi *et al.*, 2006, *J. Microbiol.*
25 *Methods* 64: 391-397), or conjugation (see, e.g., Pinedo and Smets, 2005, *Appl. Environ. Microbiol.* 71: 51-57). The introduction of DNA into a *Streptococcus* cell may be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, *Infect. Immun.* 32: 1295-1297), protoplast transformation (see, e.g., Catt and Jollick, 1991, *Microbios* 68: 189-207), electroporation (see, e.g., Buckley *et al.*, 1999, *Appl. Environ. Microbiol.* 65: 3800-3804) or
30 conjugation (see, e.g., Clewell, 1981, *Microbiol. Rev.* 45: 409-436). However, any method known in the art for introducing DNA into a host cell can be used.

The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

The host cell may be a fungal cell. "Fungi" as used herein includes the phyla
35 Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota as well as the Oomycota and all mitosporic fungi (as defined by Hawksworth *et al.*, In, *Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK).

The fungal host cell may be a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of*
 5 *Yeast* (Skinner, Passmore, and Davenport, editors, Soc. App. Bacteriol. Symposium Series No. 9, 1980).

The yeast host cell may be a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell such as a *Kluyveromyces lactis*, *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*,
 10 *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, *Saccharomyces oviformis*, or *Yarrowia lipolytica* cell.

The fungal host cell may be a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of
 15 chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

The filamentous fungal host cell may be an *Acremonium*, *Aspergillus*, *Aureobasidium*,
 20 *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

For example, the filamentous fungal host cell may be an *Aspergillus awamori*,
 25 *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermisporea*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium*
 30 *pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium*
 35 *sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*,

Trichoderma koningii, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*.

5 Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238023, Yelton *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81: 1470-1474, and Christensen *et al.*, 1988, *Bio/Technology* 6: 1419-1422. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and
10 Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, 1983, *J. Bacteriol.* 153: 163; and Hinnen *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75: 1920.

Methods of Production

15 The present invention also relates to methods of producing a variant, comprising: (a) cultivating a host cell of the present invention under conditions suitable for expression of the variant; and (b) recovering the variant.

The host cells are cultivated in a nutrient medium suitable for production of the variant using methods known in the art. For example, the cell may be cultivated by shake flask
20 cultivation, or small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the variant to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or
25 may be prepared according to published compositions (*e.g.*, in catalogues of the American Type Culture Collection). If the variant is secreted into the nutrient medium, the variant can be recovered directly from the medium. If the variant is not secreted, it can be recovered from cell lysates.

The variant may be detected using methods known in the art that are specific for the
30 variants. These detection methods include, but are not limited to, use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the variant. Lipase activity may be determined by using the p-nitrophenyl (pNP) assay as described in the section "Examples".

The variant may be recovered using methods known in the art. For example, the variant
35 may be recovered from the nutrient medium by conventional procedures including, but not limited to, collection, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

The variant may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., *Protein Purification*, Janson and Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure variants.

In one aspect, the variant is not recovered, but rather a host cell of the present invention expressing the variant is used as a source of the variant.

Compositions

Compositions comprising the polypeptide of the present inventions are contemplated. In certain aspects the present invention relates to detergent composition comprising a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase.

In one aspect the parent lipase is a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90 % identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase.

In one aspect the parent lipase is a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2.

In one aspect the parent lipase has the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the parent lipase comprises or consists the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

In one aspect the invention relates to compositions comprising lipase variants further comprising substitution at a position corresponding to any of positions; 2, 4, 8, 11, 15, 27, 33, 38, 43, 48, 51, 54, 56, 57, 58, 60, 69, 71, 83, 86, 91, 92, 94, 96, 97, 98, 99, 101, 111, 123, 150, 152, 163, 176, 179, 187, 188, 189, 198, 199, 200, 210, 216, 220, 224, 225, 227, 228, 229, 231, 236, 238, 239, 246, 249, 254, 255, 256, 257, 260, 263, 264, 265, 266, 267, 269 of SEQ ID NO:

2. In one aspect the substitutions is selected from: V2K, Q4R, Q4V, N8R, N11R, Q15C, D27G, D27R, N33K, N33Q, G38A, E43C, D48C, F51V, S54T, E56K, D57G, S58A, V60K, V60S, L69R, N71C, S83T, I86V, G91A, G91N, G91Q, N92D, N94K, N94R, D96E, D96G, D96L, D96W, L97M, K98E, K98I, K98Q, E99K, E99N, N101D, N101S, D111A, T123V, A150G, A152G, G163K, V176L, R179L, V187Y, V187W, Q188R, T189Y, T189W, H198S, T199R, N200R, E210K, E210Q, S216P, Y220F, S224R, G225R, L227G, L227R, V228R, P229R, T231R, V236R, I238C, E239C, G246C, Q249R, D254S, I255G, P256K, P256T, P256V, A257I, A257V, W260C, G263Q, L264A, I265T, G266D, T267A, L269N, L269V of SEQ ID NO: 2. In one aspect a set of substitutions is selected from: E1C N233C; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K N233C D254S P256T; E1C V2K D27G N33K G38A F51V D96E D111A G163K N233C D254S P256T; E1C V2K D27R N33K G38A F51V D96E D111A G163K Q188R N233C D254S P256T; E1C D27R G38A G91A N92D D96L K98Q D111A G163K N233C D254S P256T; E1C D27R G38A G91N N94R D96E D111A G163K S216P L227G N233C D254S P256T; E1C T231R N233C; E1C T231R N233C Q249R D254S; E1C G225R T231R N233C; E1C Q15C E43C T231R N233C; E1C L227R T231R N233C; E1C P229R T231R N233C; E1C L227G T231R N233C; E1C E99N N101S T231R N233C; E1C L227G T231R N233C D254S; E1C E210K L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K D57G L69R D96E K98I D111A A152G G163K T231R N233C D254S P256T; E1C V187Y T189Y L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C V60K I86V A150G E210K L227G T231R N233C P256K; E1C V187W T189W L227G T231R N233C; E1C N94K D96L L227G T231R N233C; E1C G91A N92D D96L K98Q L227G T231R N233C; E1C N8R L227G T231R N233C; E1C L227G V228R T231R N233C; E1C Q4R L227G T231R N233C; E1C N11R L227G T231R N233C; E1C S224R L227G T231R N233C; E1C L227G T231R N233C V236R; E1C N200R L227G T231R N233C; E1C T199R L227G T231R N233C; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T A257I; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T A257I; E1C D27R N33K G38A F51V S54T E56K D57G D96E K98I D111A G163K T231R N233C D254S I255G P256T A257V L269V; E1C V2K D27R N33K G38A F51V L69R D96E K98E D111A G163K V176L E210K L227G T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R G38A

F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T; and E1C N11R D27R N33K D48C F51V L69R N71C E87Q K98E N101R T143A E210K G225R L227G P229R T231R N233C Q249R P250R D254S I255G P256K.

In one aspect said variant has increased stability compared to the parent lipase.

5 The non-limiting list of composition components illustrated hereinafter are suitable for use in the compositions and methods herein may be desirably incorporated in certain embodiments of the invention, e.g. to assist or enhance cleaning performance, for treatment of the substrate to be cleaned, or to modify the aesthetics of the composition as is the case with perfumes, colorants, dyes or the like. The levels of any such components incorporated in any
10 compositions are in addition to any materials previously recited for incorporation. The precise nature of these additional components, and levels of incorporation thereof, will depend on the physical form of the composition and the nature of the cleaning operation for which it is to be used. Although components mentioned below are categorized by general header according to a particular functionality, this is not to be construed as a limitation, as a component may comprise
15 additional functionalities as will be appreciated by the skilled artisan.

 Unless otherwise indicated the amounts in percentage is by weight of the composition (wt%). Suitable component materials include, but are not limited to, surfactants, builders, chelating agents, dye transfer inhibiting agents, dispersants, enzymes, and enzyme stabilizers, catalytic materials, bleach activators, hydrogen peroxide, sources of hydrogen peroxide,
20 preformed peracids, polymeric dispersing agents, clay soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, hueing dyes, perfumes, perfume delivery systems, structure elasticizing agents, fabric softeners, carriers, hydrotropes, processing aids, solvents and/or pigments. In addition to the disclosure below, suitable examples of such other components and levels of use are found in US5576282, US6306812, and US6326348 hereby
25 incorporated by reference.

 Thus, in certain embodiments the invention do not contain one or more of the following adjuncts materials: surfactants, soaps, builders, chelating agents, dye transfer inhibiting agents, dispersants, additional enzymes, enzyme stabilizers, catalytic materials, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, preformed peracids, polymeric dispersing
30 agents, clay soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, perfumes, perfume delivery systems, structure elasticizing agents, fabric softeners, carriers, hydrotropes, processing aids, solvents and/or pigments. However, when one or more components are present, such one or more components may be present as detailed below:

Surfactants - The compositions according to the present invention may comprise a
35 surfactant or surfactant system wherein the surfactant can be selected from nonionic surfactants, anionic surfactants, cationic surfactants, ampholytic surfactants, zwitterionic surfactants, semi-polar nonionic surfactants and mixtures thereof. When present, surfactant is typically present at a level of from 0.1 to 60wt%, from 0.2 to 40wt%, from 0.5 to 30wt%, from 1

to 50wt%, from 1 to 40wt%, from 1 to 30wt%, from 1 to 20wt%, from 3 to 10wt%, from 3 to 5wt%, from 5 to 40wt%, from 5 to 30wt%, from 5 to 15wt%, from 3 to 20wt%, from 3 to 10wt%, from 8 to 12wt%, from 10 to 12wt%, from 20 to 25wt% or from 25-60%.

5 Suitable anionic deterative surfactants include sulphate and sulphonate deterative surfactants.

Suitable sulphonate deterative surfactants include alkyl benzene sulphonate, in one aspect, C₁₀₋₁₃ alkyl benzene sulphonate. Suitable alkyl benzene sulphonate (LAS) may be obtained, by sulphonating commercially available linear alkyl benzene (LAB); suitable LAB includes low 2-phenyl LAB, such as Isochem® or Petrelab®, other suitable LAB include high 2-phenyl LAB, such as Hyblene®. A suitable anionic deterative surfactant is alkyl benzene sulphonate that is obtained by DETAL catalyzed process, although other synthesis routes, such as HF, may also be suitable. In one aspect a magnesium salt of LAS is used.

Suitable sulphate deterative surfactants include alkyl sulphate, in one aspect, C₈₋₁₈ alkyl sulphate, or predominantly C₁₂ alkyl sulphate.

15 Another suitable sulphate deterative surfactant is alkyl alkoxyated sulphate, in one aspect, alkyl ethoxyated sulphate, in one aspect, a C₈₋₁₈ alkyl alkoxyated sulphate, in one aspect, a C₈₋₁₈ alkyl ethoxyated sulphate, typically the alkyl alkoxyated sulphate has an average degree of alkoxylation of from 0.5 to 20, or from 0.5 to 10, typically the alkyl alkoxyated sulphate is a C₈₋₁₈ alkyl ethoxyated sulphate having an average degree of ethoxylation of from 20 0.5 to 10, from 0.5 to 7, from 0.5 to 5 or from 0.5 to 3.

The alkyl sulphate, alkyl alkoxyated sulphate and alkyl benzene sulphonates may be linear or branched, substituted or un-substituted.

The deterative surfactant may be a mid-chain branched deterative surfactant, in one aspect, a mid-chain branched anionic deterative surfactant, in one aspect, a mid-chain branched alkyl sulphate and/or a mid-chain branched alkyl benzene sulphonate, e.g. a mid-chain branched alkyl sulphate. In one aspect, the mid-chain branches are C₁₋₄ alkyl groups, typically methyl and/or ethyl groups.

Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), 30 phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated 35 fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodeceny/tetradecenyl succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or soap, and combinations thereof.

Suitable non-ionic deterative surfactants are selected from the group consisting of: C₈-C₁₈ alkyl ethoxylates, such as, NEODOL®; C₆-C₁₂ alkyl phenol alkoxyates wherein the alkoxyate units may be ethyleneoxy units, propyleneoxy units or a mixture thereof; C₁₂-C₁₈ alcohol and C₆-C₁₂ alkyl phenol condensates with ethylene oxide/propylene oxide block polymers such as Pluronic®; C₁₄-C₂₂ mid-chain branched alcohols; C₁₄-C₂₂ mid-chain branched alkyl alkoxyates, typically having an average degree of alkoxylation of from 1 to 30; alkylpolysaccharides, in one aspect, alkylpolyglycosides; polyhydroxy fatty acid amides; ether capped poly(oxyalkylated) alcohol surfactants; and mixtures thereof.

Suitable non-ionic deterative surfactants include alkyl polyglucoside and/or an alkyl alkoxyated alcohol.

In one aspect, non-ionic deterative surfactants include alkyl alkoxyated alcohols, in one aspect C₈₋₁₈ alkyl alkoxyated alcohol, e.g. a C₈₋₁₈ alkyl ethoxylated alcohol, the alkyl alkoxyated alcohol may have an average degree of alkoxylation of from 1 to 50, from 1 to 30, from 1 to 20, or from 1 to 10. In one aspect, the alkyl alkoxyated alcohol may be a C₈₋₁₈ alkyl ethoxylated alcohol having an average degree of ethoxylation of from 1 to 10, from 1 to 7, more from 1 to 5 or from 3 to 7. The alkyl alkoxyated alcohol can be linear or branched, and substituted or unsubstituted. Suitable nonionic surfactants include Lutensol®.

Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxyated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxyated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or *N*-acyl *N*-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof.

Suitable cationic deterative surfactants include alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, alkyl ternary sulphonium compounds, and mixtures thereof.

Suitable cationic deterative surfactants are quaternary ammonium compounds having the general formula: (R)(R₁)(R₂)(R₃)N⁺ X⁻, wherein, R is a linear or branched, substituted or unsubstituted C₆₋₁₈ alkyl or alkenyl moiety, R₁ and R₂ are independently selected from methyl or ethyl moieties, R₃ is a hydroxyl, hydroxymethyl or a hydroxyethyl moiety, X is an anion which provides charge neutrality, suitable anions include: halides, e.g. chloride; sulphate; and sulphonate. Suitable cationic deterative surfactants are mono-C₆₋₁₈ alkyl mono-hydroxyethyl dimethyl quaternary ammonium chlorides. Highly suitable cationic deterative surfactants are mono-C₈₋₁₀ alkyl mono-hydroxyethyl dimethyl quaternary ammonium chloride, mono-C₁₀₋₁₂ alkyl

mono-hydroxyethyl di-methyl quaternary ammonium chloride and mono-C₁₀ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride.

Non-limiting examples of cationic surfactants include alkyl dimethylethanolamine quat (ADMEAQ), cetyltrimethylammonium bromide (CTAB), dimethyldistearyl ammonium chloride (DSDMAC), and alkylbenzyl dimethyl ammonium, alkyl quaternary ammonium compounds, alkoxyated quaternary ammonium (AQA) compounds, ester quats, and combinations thereof.


Suitable amphoteric/zwitterionic surfactants include amine oxides and betaines such as alkyl dimethyl betaines, sulfobetaines, or combinations thereof. Amine-neutralized anionic surfactants - Anionic surfactants of the present invention and adjunct anionic cosurfactants, may exist in an acid form, and said acid form may be neutralized to form a surfactant salt which is desirable for use in the present detergent compositions. Typical agents for neutralization include the metal counterion base such as hydroxides, eg, NaOH or KOH. Further preferred agents for neutralizing anionic surfactants of the present invention and adjunct anionic surfactants or cosurfactants in their acid forms include ammonia, amines, or alkanolamines.

Alkanolamines are preferred. Suitable non-limiting examples including monoethanolamine, diethanolamine, triethanolamine, and other linear or branched alkanolamines known in the art; e.g., highly preferred alkanolamines include 2-amino-1-propanol, 1-aminopropanol, monoisopropanolamine, or 1-amino-3-propanol. Amine neutralization may be done to a full or partial extent, e.g. part of the anionic surfactant mix may be neutralized with sodium or potassium and part of the anionic surfactant mix may be neutralized with amines or alkanolamines.

Non-limiting examples of semipolar surfactants include amine oxides (AO) such as alkyl dimethylamine oxide

Surfactant systems comprising mixtures of one or more anionic and in addition one or more nonionic surfactants optionally with an additional surfactant such as a cationic surfactant, may be preferred. Preferred weight ratios of anionic to nonionic surfactant are at least 2:1, or at least 1:1 to 1:10.

In one aspect a surfactant system may comprise a mixture of isoprenoid surfactants represented by formula A and formula B:



where Y is CH₂ or null, and Z may be chosen such that the resulting surfactant is selected from the following surfactants: an alkyl carboxylate surfactant, an alkyl polyalkoxy surfactant, an alkyl anionic polyalkoxy sulfate surfactant, an alkyl glycerol ester sulfonate surfactant, an alkyl dimethyl amine oxide surfactant, an alkyl polyhydroxy based surfactant, an alkyl phosphate ester surfactant, an alkyl glycerol sulfonate surfactant, an alkyl polygluconate surfactant, an alkyl polyphosphate ester surfactant, an alkyl phosphonate surfactant, an alkyl polyglycoside

surfactant, an alkyl monoglycoside surfactant, an alkyl diglycoside surfactant, an alkyl sulfosuccinate surfactant, an alkyl disulfate surfactant, an alkyl disulfonate surfactant, an alkyl sulfosuccinamate surfactant, an alkyl glucamide surfactant, an alkyl taurinate surfactant, an alkyl sarcosinate surfactant, an alkyl glycinate surfactant, an alkyl isethionate surfactant, an

5 alkyl dialkanolamide surfactant, an alkyl monoalkanolamide surfactant, an alkyl monoalkanolamide sulfate surfactant, an alkyl diglycolamide surfactant, an alkyl diglycolamide sulfate surfactant, an alkyl glycerol ester surfactant, an alkyl glycerol ester sulfate surfactant, an alkyl glycerol ether surfactant, an alkyl glycerol ether sulfate surfactant, alkyl methyl ester sulfonate surfactant, an alkyl polyglycerol ether surfactant, an alkyl polyglycerol ether sulfate

10 surfactant, an alkyl sorbitan ester surfactant, an alkyl ammonioalkanesulfonate surfactant, an alkyl amidopropyl betaine surfactant, an alkyl allylated quat based surfactant, an alkyl monohydroxyalkyl-di-alkylated quat based surfactant, an alkyl di-hydroxyalkyl monoalkyl quat based surfactant, an alkylated quat surfactant, an alkyl trimethylammonium quat surfactant, an alkyl polyhydroxyalkyl oxypropyl quat based surfactant, an alkyl glycerol ester quat surfactant, an

15 alkyl glycol amine quat surfactant, an alkyl monomethyl dihydroxyethyl quaternary ammonium surfactant, an alkyl dimethyl monohydroxyethyl quaternary ammonium surfactant, an alkyl trimethylammonium surfactant, an alkyl imidazoline-based surfactant, an alken-2-yl- succinate surfactant, an alkyl α -sulfonated carboxylic acid surfactant, an alkyl α -sulfonated carboxylic acid alkyl ester surfactant, an α olefin sulfonate surfactant, an alkyl phenol ethoxylate surfactant,

20 an alkyl benzenesulfonate surfactant, an alkyl sulfobetaine surfactant, an alkyl hydroxysulfobetaine surfactant, an alkyl ammoniocarboxylate betaine surfactant, an alkyl sucrose ester surfactant, an alkyl alkanolamide surfactant, an alkyl di(polyoxyethylene) monoalkyl ammonium surfactant, an alkyl mono(polyoxyethylene) dialkyl ammonium surfactant, an alkyl benzyl dimethylammonium surfactant, an alkyl aminopropionate surfactant, an alkyl

25 amidopropyl dimethylamine surfactant, or a mixture thereof; and if Z is a charged moiety, Z is charge-balanced by a suitable metal or organic counter ion. Suitable counter ions include a metal counter ion, an amine, or an alkanolamine, e.g., C1-C6 alkanolammonium. More specifically, suitable counter ions include Na⁺, Ca⁺, Li⁺, K⁺, Mg⁺, e.g., monoethanolamine (MEA), diethanolamine (DEA), triethanolamine (TEA), 2- amino-1-propanol, 1-aminopropanol,

30 methyldiethanolamine, dimethylethanolamine, monoisopropanolamine, triisopropanolamine, 1-amino-3-propanol, or mixtures thereof. In one aspect, the compositions contain from 5% to 97% of one or more non- isoprenoid surfactants; and one or more adjunct cleaning additives; wherein the weight ratio of surfactant of formula A to surfactant of formula B is from 50:50 to 95:5.

Soap - The compositions herein may contain soap. Without being limited by theory, it

35 may be desirable to include soap as it acts in part as a surfactant and in part as a builder and may be useful for suppression of foam and may furthermore interact favorably with the various cationic compounds of the composition to enhance softness on textile fabrics treaded with the inventive compositions. Any soap known in the art for use in laundry detergents may be utilized.

In one aspect, the compositions contain from 0wt% to 20wt%, from 0.5wt% to 20wt%, from 4wt% to 10wt%, or from 4wt% to 7wt% of soap.

Examples of soap useful herein include oleic acid soaps, palmitic acid soaps, palm kernel fatty acid soaps, and mixtures thereof. Typical soaps are in the form of mixtures of fatty acid soaps having different chain lengths and degrees of substitution. One such mixture is topped palm kernel fatty acid.

In one aspect, the soap is selected from free fatty acid. Suitable fatty acids are saturated and/or unsaturated and can be obtained from natural sources such a plant or animal esters (e.g., palm kernel oil, palm oil, coconut oil, babassu oil, safflower oil, tall oil, castor oil, tallow and fish oils, grease, and mixtures thereof), or synthetically prepared (e.g., via the oxidation of petroleum or by hydrogenation of carbon monoxide via the Fisher Tropsch process).

Examples of suitable saturated fatty acids for use in the compositions of this invention include capric, lauric, myristic, palmitic, stearic, arachidic and behenic acid. Suitable unsaturated fatty acid species include: palmitoleic, oleic, linoleic, linolenic and ricinoleic acid. Examples of preferred fatty acids are saturated C_n fatty acid, saturated C_{12} - C_{14} fatty acids, and saturated or unsaturated C_n to C_{18} fatty acids, and mixtures thereof.

When present, the weight ratio of fabric softening cationic cosurfactant to fatty acid is preferably from about 1:3 to about 3: 1, more preferably from about 1:1.5 to about 1.5:1, most preferably about 1:1.

Levels of soap and of nonsoap anionic surfactants herein are percentages by weight of the detergent composition, specified on an acid form basis. However, as is commonly understood in the art, anionic surfactants and soaps are in practice neutralized using sodium, potassium or alkanolammonium bases, such as sodium hydroxide or monoethanolamine.

Hydrotropes – The compositions of the present invention may comprise one or more hydrotropes. A hydrotrope is a compound that solubilises hydrophobic compounds in aqueous solutions (or oppositely, polar substances in a non-polar environment). Typically, hydrotropes have both hydrophilic and a hydrophobic character (so-called amphiphilic properties as known from surfactants); however the molecular structure of hydrotropes generally do not favor spontaneous self-aggregation, see e.g. review by Hodgdon and Kaler (2007), Current Opinion in Colloid & Interface Science 12: 121-128. Hydrotropes do not display a critical concentration above which self-aggregation occurs as found for surfactants and lipids forming micellar, lamellar or other well defined meso-phases. Instead, many hydrotropes show a continuous-type aggregation process where the sizes of aggregates grow as concentration increases. However, many hydrotropes alter the phase behavior, stability, and colloidal properties of systems containing substances of polar and non-polar character, including mixtures of water, oil, surfactants, and polymers. Hydrotropes are classically used across industries from pharma, personal care, food, to technical applications. Use of hydrotropes in detergent compositions allow for example more concentrated formulations of surfactants (as in the process of

compacting liquid detergents by removing water) without inducing undesired phenomena such as phase separation or high viscosity.

The detergent may contain from 0 to 10wt%, such as from 0 to 5wt%, 0.5 to 5wt%, or from 3% to 5wt%, of a hydrotrope. Any hydrotrope known in the art for use in detergents may be utilized. Non-limiting examples of hydrotropes include sodium benzenesulfonate, sodium p-toluene sulfonate (STS), sodium xylene sulfonate (SXS), sodium cumene sulfonate (SCS), sodium cymene sulfonate, amine oxides, alcohols and polyglycoethers, sodium hydroxynaphthoate, sodium hydroxynaphthalene sulfonate, sodium ethylhexyl sulfate, and combinations thereof.

Builders - The compositions of the present invention may comprise one or more builders, co-builders, builder systems or a mixture thereof. When a builder is used, the cleaning composition will typically comprise from 0 to 65wt%, at least 1wt%, from 2 to 60wt% or from 5 to 10wt% builder. In a dish wash cleaning composition, the level of builder is typically 40 to 65wt% or 50 to 65wt%. The composition may be substantially free of builder; substantially free means "no deliberately added" zeolite and/or phosphate. Typical zeolite builders include zeolite A, zeolite P and zeolite MAP. A typical phosphate builder is sodium tri-polyphosphate.

The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), ethanolamines such as 2-aminoethan-1-ol (MEA), iminodiethanol (DEA) and 2,2',2''-nitrilotriethanol (TEA), and carboxymethylinulin (CMI), and combinations thereof.

The cleaning composition may include a co-builder alone, or in combination with a builder, e.g. a zeolite builder. Non-limiting examples of co-builders include homopolymers of polyacrylates or copolymers thereof, such as poly(acrylic acid) (PAA) or copoly(acrylic acid/maleic acid) (PAA/PMA). Further non-limiting examples include citrate, chelators such as aminocarboxylates, aminopolycarboxylates and phosphonates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2''-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-N,N'-disuccinic acid (EDDS), methylglycinediacetic acid (MGDA), glutamic acid-N,N'-diacetic acid (GLDA), 1-hydroxyethane-1,1-diylbis(phosphonic acid) (HEDP), ethylenediaminetetrakis(methylene)tetrakis(phosphonic acid) (EDTMPA), diethylenetriaminepentakis(methylene)pentakis(phosphonic acid) (DTPMPA), N-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N'-diacetic acid (ASDA), aspartic acid-N- monopropionic acid (ASMP), iminodisuccinic acid (IDA), N- (2-sulfomethyl) aspartic acid (SMAS), N- (2-sulfoethyl) aspartic acid (SEAS), N- (2-sulfomethyl) glutamic acid (SMGL), N- (2- sulfoethyl) glutamic acid (SEGL), N- methyliminodiacetic

acid (MIDA), α -alanine-N,N-diacetic acid (α -ALDA), serine-N,N-diacetic acid (SEDA), isoserine-N,N-diacetic acid (ISDA), phenylalanine-N,N-diacetic acid (PHDA), anthranilic acid-N,N-diacetic acid (ANDA), sulfanilic acid-N,N-diacetic acid (SLDA), taurine-N,N-diacetic acid (TUDA) and sulfomethyl-N,N-diacetic acid (SMDA), N-(hydroxyethyl)-ethylenediaminetriacetate (HEDTA),
5 diethanolglycine (DEG), Diethylenetriamine Penta (Methylene Phosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), and combinations and salts thereof. Further exemplary builders and/or co-builders are described in, e.g., WO09/102854, US5977053.

In one aspect, the invention relates to compositions comprising a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C
10 of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of SEQ ID NO: 2, the composition comprising up to 10wt% or 15wt% aluminosilicate (anhydrous basis) and/or phosphate builder, the composition having a reserve alkalinity of greater than 4 or 7.5. As used herein the term
15 "reserve alkalinity" is a measure of the buffering capacity of the composition (g/NaOH/100g composition) determined by titrating a 1% (w/v) solution of composition with hydrochloric acid to pH 7.5 i.e. in order to calculate reserve alkalinity. Reserve alkalinity may be calculated as disclosed on page 9 in WO2006/090335. In one aspect the parent lipase is a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90 % identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type
20 lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions
25 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase. In one aspect the parent lipase is a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to
30 T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2. In one aspect the parent lipase has the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the parent lipase comprises or consists the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the invention relates to compositions comprising lipase variants further
35 comprising substitution at a position corresponding to any of positions; 2, 4, 8, 11, 15, 27, 33, 38, 43, 48, 51, 54, 56, 57, 58, 60, 69, 71, 83, 86, 91, 92, 94, 96, 97, 98, 99, 101, 111, 123, 150, 152, 163, 176, 179, 187, 188, 189, 198, 199, 200, 210, 216, 220, 224, 225, 227, 228, 229, 231, 236, 238, 239, 246, 249, 254, 255, 256, 257, 260, 263, 264, 265, 266, 267, 269 of SEQ ID NO:

2. In one aspect the substitutions is selected from: V2K, Q4R, Q4V, N8R, N11R, Q15C, D27G, D27R, N33K, N33Q, G38A, E43C, D48C, F51V, S54T, E56K, D57G, S58A, V60K, V60S, L69R, N71C, S83T, I86V, G91A, G91N, G91Q, N92D, N94K, N94R, D96E, D96G, D96L, D96W, L97M, K98E, K98I, K98Q, E99K, E99N, N101D, N101S, D111A, T123V, A150G, A152G, G163K, V176L, R179L, V187Y, V187W, Q188R, T189Y, T189W, H198S, T199R, N200R, E210K, E210Q, S216P, Y220F, S224R, G225R, L227G, L227R, V228R, P229R, T231R, V236R, I238C, E239C, G246C, Q249R, D254S, I255G, P256K, P256T, P256V, A257I, A257V, W260C, G263Q, L264A, I265T, G266D, T267A, L269N, L269V of SEQ ID NO: 2. In one aspect a set of substitutions is selected from: E1C N233C; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K N233C D254S P256T; E1C V2K D27G N33K G38A F51V D96E D111A G163K N233C D254S P256T; E1C V2K D27R N33K G38A F51V D96E D111A G163K Q188R N233C D254S P256T; E1C D27R G38A G91A N92D D96L K98Q D111A G163K N233C D254S P256T; E1C D27R G38A G91N N94R D96E D111A G163K S216P L227G N233C D254S P256T; E1C T231R N233C; E1C T231R N233C Q249R D254S; E1C G225R T231R N233C; E1C Q15C E43C T231R N233C; E1C L227R T231R N233C; E1C P229R T231R N233C; E1C L227G T231R N233C; E1C E99N N101S T231R N233C; E1C L227G T231R N233C D254S; E1C E210K L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K D57G L69R D96E K98I D111A A152G G163K T231R N233C D254S P256T; E1C V187Y T189Y L227G T231R N233C; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C V60K I86V A150G E210K L227G T231R N233C P256K; E1C V187W T189W L227G T231R N233C; E1C N94K D96L L227G T231R N233C; E1C G91A N92D D96L K98Q L227G T231R N233C; E1C N8R L227G T231R N233C; E1C L227G V228R T231R N233C; E1C Q4R L227G T231R N233C; E1C N11R L227G T231R N233C; E1C S224R L227G T231R N233C; E1C L227G T231R N233C V236R; E1C N200R L227G T231R N233C; E1C T199R L227G T231R N233C; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I; E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T A257I; E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T A257I; E1C D27R N33K G38A F51V S54T E56K D57G D96E K98I D111A G163K T231R N233C D254S I255G P256T A257V L269V; E1C V2K D27R N33K G38A F51V L69R D96E K98E D111A G163K V176L E210K L227G T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T; E1C D27R G38A


F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T; and E1C N11R D27R N33K D48C F51V L69R N71C E87Q K98E N101R T143A E210K G225R L227G P229R T231R N233C Q249R P250R D254S I255G P256K.

Chelating Agents and Crystal Growth Inhibitors - The compositions herein may contain a
 5 chelating agent and/or a crystal growth inhibitor. Suitable molecules include copper, iron and/or manganese chelating agents and mixtures thereof. Suitable molecules include DTPA (Diethylene triamine pentaacetic acid), HEDP (Hydroxyethane diphosphonic acid), DTPMP (Diethylene triamine penta(methylene phosphonic acid)), 1,2-Dihydroxybenzene-3,5-disulfonic acid disodium salt hydrate, ethylenediamine, diethylene triamine, ethylenediaminedisuccinic
 10 acid (EDDS), N-hydroxyethylethylenediaminetri-acetic acid (HEDTA), triethylenetetraaminehexaacetic acid (TTA), N-hydroxyethyliminodiacetic acid (HEIDA), dihydroxyethylglycine (DHEG), ethylenediaminetetrapropionic acid (EDTP), carboxymethyl inulin and 2-Phosphonobutane 1,2,4-tricarboxylic acid (Bayhibit® AM) and derivatives thereof. Typically the composition may comprise from 0.005 to 15wt% or from 3.0 to 10wt% chelating
 15 agent or crystal growth inhibitor.

Bleach Component - The bleach component suitable for incorporation in the methods and compositions of the invention comprise one or a mixture of more than one bleach component. Suitable bleach components include bleaching catalysts, photobleaches, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, pre-formed peracids and mixtures
 20 thereof. In general, when a bleach component is used, the compositions of the present invention may comprise from 0 to 30wt%, from 0.00001 to 90wt%, 0.0001 to 50wt%, from 0.001 to 25wt% or from 1 to 20wt%. Examples of suitable bleach components include:

(1) Pre-formed peracids: Suitable preformed peracids include, but are not limited to, compounds selected from the group consisting of pre-formed peroxyacids or salts thereof,
 25 typically either a peroxycarboxylic acid or salt thereof, or a peroxysulphonic acid or salt thereof.


The pre-formed peroxyacid or salt thereof is preferably a peroxycarboxylic acid or salt thereof, typically having a chemical structure corresponding to the following chemical formula:



wherein: R¹⁴ is selected from alkyl, aralkyl, cycloalkyl, aryl or heterocyclic groups; the R¹⁴ group
 30 can be linear or branched, substituted or unsubstituted; and Y is any suitable counter-ion that achieves electric charge neutrality, preferably Y is selected from hydrogen, sodium or potassium. Preferably, R¹⁴ is a linear or branched, substituted or unsubstituted C₆₋₉ alkyl. Preferably, the peroxyacid or salt thereof is selected from peroxyhexanoic acid, peroxyheptanoic acid, peroxyoctanoic acid, peroxydecanoic acid, any salt
 35 thereof, or any combination thereof. Particularly preferred peroxyacids are phthalimido-peroxy-

alkanoic acids, in particular ϵ -phthahlimido peroxy hexanoic acid (PAP). Preferably, the peroxyacid or salt thereof has a melting point in the range of from 30°C to 60°C.

The pre-formed peroxyacid or salt thereof can also be a peroxysulphonic acid or salt thereof, typically having a chemical structure corresponding to the following chemical formula:



wherein: R^{15} is selected from alkyl, aralkyl, cycloalkyl, aryl or heterocyclic groups; the R^{15} group can be linear or branched, substituted or unsubstituted; and Z is any suitable counter-ion that achieves electric charge neutrality, preferably Z is selected from hydrogen, sodium or potassium. Preferably R^{15} is a linear or branched, substituted or unsubstituted C_{6-9} alkyl. Preferably such bleach components may be present in the compositions of the invention in an amount from 0.01 to 50wt% or from 0.1 to 20wt%.

(2) Sources of hydrogen peroxide include e.g., inorganic perhydrate salts, including alkali metal salts such as sodium salts of perborate (usually mono- or tetra-hydrate), percarbonate, persulphate, perphosphate, persilicate salts and mixtures thereof. In one aspect of the invention the inorganic perhydrate salts such as those selected from the group consisting of sodium salts of perborate, percarbonate and mixtures thereof. When employed, inorganic perhydrate salts are typically present in amounts of 0.05 to 40wt% or 1 to 30wt% of the overall composition and are typically incorporated into such compositions as a crystalline solid that may be coated. Suitable coatings include: inorganic salts such as alkali metal silicate, carbonate or borate salts or mixtures thereof, or organic materials such as water-soluble or dispersible polymers, waxes, oils or fatty soaps. Preferably such bleach components may be present in the compositions of the invention in an amount of 0.01 to 50wt% or 0.1 to 20wt%.

(3) The term bleach activator is meant herein as a compound which reacts with hydrogen peroxide to form a peracid via perhydrolysis. The peracid thus formed constitutes the activated bleach. Suitable bleach activators to be used herein include those belonging to the class of esters, amides, imides or anhydrides. Suitable bleach activators are those having $\text{R}-(\text{C}=\text{O})-\text{L}$ wherein R is an alkyl group, optionally branched, having, when the bleach activator is hydrophobic, from 6 to 14 carbon atoms, or from 8 to 12 carbon atoms and, when the bleach activator is hydrophilic, less than 6 carbon atoms or less than 4 carbon atoms; and L is leaving group. Examples of suitable leaving groups are benzoic acid and derivatives thereof - especially benzene sulphonate. Suitable bleach activators include dodecanoyl oxybenzene sulphonate, decanoyl oxybenzene sulphonate, decanoyl oxybenzoic acid or salts thereof, 3,5,5-trimethyl hexanoyloxybenzene sulphonate, tetraacetyl ethylene diamine (TAED), sodium 4-[(3,5,5-trimethylhexanoyl)oxy]benzene-1-sulfonate (ISONOBS), 4-(dodecanoyloxy)benzene-1-sulfonate

(LOBS), 4-(decanoyloxy)benzene-1-sulfonate, 4-(decanoyloxy)benzoate (DOBS or DOBA), 4-(nonanoyloxy)benzene-1-sulfonate (NOBS), and/or those disclosed in WO98/17767. A family of bleach activators is disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like triacetin has the advantage that it is environmentally friendly. Furthermore acetyl triethyl citrate and triacetin have good hydrolytical stability in the product upon storage and are efficient bleach activators. Finally ATC is multifunctional, as the citrate released in the perhydrolysis reaction may function as a builder. Alternatively, the bleaching system may comprise peroxyacids of, for example, the amide, imide, or sulfone type. The bleaching system may also comprise peracids such as 6-(phthalimido)peroxyhexanoic acid (PAP). Suitable bleach activators are also disclosed in WO98/17767. While any suitable bleach activator may be employed, in one aspect of the invention the subject cleaning composition may comprise NOBS, TAED or mixtures thereof. When present, the peracid and/or bleach activator is generally present in the composition in an amount of 0.1 to 60wt%, 0.5 to 40wt% or 0.6 to 10wt% based on the fabric and home care composition. One or more hydrophobic peracids or precursors thereof may be used in combination with one or more hydrophilic peracid or precursor thereof. Preferably such bleach components may be present in the compositions of the invention in an amount of 0.01 to 50wt%, or 0.1 to 20wt%.

The amounts of hydrogen peroxide source and peracid or bleach activator may be selected such that the molar ratio of available oxygen (from the peroxide source) to peracid is from 1:1 to 35:1, or even 2:1 to 10:1.

(4) Diacyl peroxides – preferred diacyl peroxide bleaching species include those selected from diacyl peroxides of the general formula: $R^1-C(O)-OO-(O)C-R^2$, in which R^1 represents a C_6-C_{18} alkyl, preferably C_6-C_{12} alkyl group containing a linear chain of at least 5 carbon atoms and optionally containing one or more substituents (e.g. $-N^+(CH_3)_3$, $-COOH$ or $-CN$) and/or one or more interrupting moieties (e.g. $-CONH-$ or $-CH=CH-$) interpolated between adjacent carbon atoms of the alkyl radical, and R^2 represents an aliphatic group compatible with a peroxide moiety, such that R^1 and R^2 together contain a total of 8 to 30 carbon atoms. In one preferred aspect R^1 and R^2 are linear unsubstituted C_6-C_{12} alkyl chains. Most preferably R^1 and R^2 are identical. Diacyl peroxides, in which both R^1 and R^2 are C_6-C_{12} alkyl groups, are particularly preferred. Preferably, at least one of, most preferably only one of, the R groups (R_1 or R_2), does not contain branching or pendant rings in the alpha position, or preferably neither in the alpha nor beta positions or most preferably in none of the alpha or beta or gamma positions. In one further preferred embodiment the DAP may be asymmetric, such that preferably the hydrolysis of R_1 acyl group is rapid to generate peracid, but the hydrolysis of R_2 acyl group is slow.

The tetraacyl peroxide bleaching species is preferably selected from tetraacyl peroxides of the general formula: $R^3-C(O)-OO-C(O)-(CH_2)_n-C(O)-OO-C(O)-R^3$, in which R^3 represents a C_1-C_9 alkyl, or C_3-C_7 , group and n represents an integer from 2 to 12, or 4 to 10 inclusive.

Preferably, the diacyl and/or tetraacyl peroxide bleaching species is present in an amount sufficient to provide at least 0.5ppm, at least 10ppm, or at least 50ppm by weight of the wash liquor. In a preferred embodiment, the bleaching species is present in an amount sufficient to provide from 0.5 to 300ppm, from 30 to 150ppm by weight of the wash liquor.

Preferably the bleach component comprises a bleach catalyst (5 and 6).

(5) Preferred are organic (non-metal) bleach catalysts include bleach catalyst capable of accepting an oxygen atom from a peroxyacid and/or salt thereof, and transferring the oxygen atom to an oxidizable substrate. Suitable bleach catalysts include, but are not limited to: iminium cations and polyions; iminium zwitterions; modified amines; modified amine oxides; N-sulphonyl imines; N-phosphonyl imines; N-acyl imines; thiadiazole dioxides; perfluoroimines; cyclic sugar ketones and mixtures thereof.

Suitable iminium cations and polyions include, but are not limited to, N-methyl-3,4-dihydroisoquinolinium tetrafluoroborate, prepared as described in Tetrahedron (1992), 49(2), 423-38 (e.g. compound 4, p.433); N-methyl-3,4-dihydroisoquinolinium p-toluene sulphonate, prepared as described in US5360569 (e.g. Column 11, Example 1); and N-octyl-3,4-dihydroisoquinolinium p-toluene sulphonate, prepared as described in US5360568 (e.g. Column 10, Ex. 3).

Suitable iminium zwitterions include, but are not limited to, N-(3-sulfopropyl)-3,4-dihydroisoquinolinium, inner salt, prepared as described in US5576282 (e.g. Column 31, Ex. II); N-[2-(sulphooxy)dodecyl]-3,4-dihydroisoquinolinium, inner salt, prepared as described in US5817614 (e.g. Column 32, Ex. V); 2-[3-[(2-ethylhexyl)oxy]-2-(sulphooxy)propyl]-3,4-dihydroisoquinolinium, inner salt, prepared as described in WO05/047264 (e.g. p.18, Ex. 8), and 2-[3-[(2-butyloctyl)oxy]-2-(sulphooxy)propyl]-3,4-dihydroisoquinolinium, inner salt.

Suitable modified amine oxygen transfer catalysts include, but are not limited to, 1,2,3,4-tetrahydro-2-methyl-1-isoquinolinol, which can be made according to the procedures described in Tetrahedron Letters (1987), 28(48), 6061-6064. Suitable modified amine oxide oxygen transfer catalysts include, but are not limited to, sodium 1-hydroxy-N-oxy-N-[2-(sulphooxy)decyl]-1,2,3,4-tetrahydroisoquinoline.

Suitable N-sulphonyl imine oxygen transfer catalysts include, but are not limited to, 3-methyl-1,2-benzisothiazole 1,1-dioxide, prepared according to the procedure described in the Journal of Organic Chemistry (1990), 55(4), 1254-61.

Suitable N-phosphonyl imine oxygen transfer catalysts include, but are not limited to, [R(E)]-N-[(2-chloro-5-nitrophenyl)methylene]-P-phenyl-P-(2,4,6-trimethylphenyl)-phosphinic

amide, which can be made according to the procedures described in the Journal of the Chemical Society, Chemical Communications (1994), (22), 2569-70.

Suitable N-acyl imine oxygen transfer catalysts include, but are not limited to, [N(E)]-N-(phenylmethylene)acetamide, which can be made according to the procedures described in Polish Journal of Chemistry (2003), 77(5), 577-590.

Suitable thiadiazole dioxide oxygen transfer catalysts include but are not limited to, 3-methyl-4-phenyl-1,2,5-thiadiazole 1,1-dioxide, which can be made according to the procedures described in US5753599 (Column 9, Ex. 2).


Suitable perfluoroimine oxygen transfer catalysts include, but are not limited to, (Z)-2,2,3,3,4,4,4-heptafluoro-N-(nonafluorobutyl)butanimidoyl fluoride, which can be made according to the procedures described in Tetrahedron Letters (1994), 35(34), 6329-30.

Suitable cyclic sugar ketone oxygen transfer catalysts include, but are not limited to, 1,2:4,5-di-O-isopropylidene-D-erythro-2,3-hexodiuro-2,6-pyranose as prepared in US6649085 (Column 12, Ex. 1).

Preferably, the bleach catalyst comprises an iminium and/or carbonyl functional group and is typically capable of forming an oxaziridinium and/or dioxirane functional group upon acceptance of an oxygen atom, especially upon acceptance of an oxygen atom from a peroxyacid and/or salt thereof. Preferably, the bleach catalyst comprises an oxaziridinium functional group and/or is capable of forming an oxaziridinium functional group upon acceptance of an oxygen atom, especially upon acceptance of an oxygen atom from a peroxyacid and/or salt thereof. Preferably, the bleach catalyst comprises a cyclic iminium functional group, preferably wherein the cyclic moiety has a ring size of from five to eight atoms (including the nitrogen atom), preferably six atoms. Preferably, the bleach catalyst comprises an aryliminium functional group, preferably a bi-cyclic aryliminium functional group, preferably a 3,4-dihydroisoquinolinium functional group. Typically, the imine functional group is a quaternary imine functional group and is typically capable of forming a quaternary oxaziridinium functional group upon acceptance of an oxygen atom, especially upon acceptance of an oxygen atom from a peroxyacid and/or salt thereof. In one aspect, the detergent composition comprises a bleach component having a $\log P_{o/w}$ no greater than 0, no greater than -0.5, no greater than -1.0, no greater than -1.5, no greater than -2.0, no greater than -2.5, no greater than -3.0, or no greater than -3.5. The method for determining $\log P_{o/w}$ is described in more detail below.


Typically, the bleach ingredient is capable of generating a bleaching species having a X_{SO} of from 0.01 to 0.30, from 0.05 to 0.25, or from 0.10 to 0.20. The method for determining X_{SO} is described in more detail below. For example, bleaching ingredients having an isoquinolinium structure are capable of generating a bleaching species that has an oxaziridinium structure. In this example, the X_{SO} is that of the oxaziridinium bleaching species.

Preferably, the bleach catalyst has a chemical structure corresponding to the following chemical formula:



wherein: n and m are independently from 0 to 4, preferably n and m are both 0; each R¹ is independently selected from a substituted or unsubstituted radical selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, fused aryl, heterocyclic ring, fused heterocyclic ring, nitro, halo, cyano, sulphonato, alkoxy, keto, carboxylic, and carboalkoxy radicals; and any two vicinal R¹ substituents may combine to form a fused aryl, fused carbocyclic or fused heterocyclic ring; each R² is independently selected from a substituted or unsubstituted radical independently selected from the group consisting of hydrogen, hydroxy, alkyl, cycloalkyl, alkaryl, aryl, aralkyl, alkenes, heterocyclic ring, alkoxys, arylcarbonyl groups, carboxyalkyl groups and amide groups; any R² may be joined together with any other of R² to form part of a common ring; any geminal R² may combine to form a carbonyl; and any two R² may combine to form a substituted or unsubstituted fused unsaturated moiety; R³ is a C₁ to C₂₀ substituted or unsubstituted alkyl; R⁴ is hydrogen or the moiety Q_t-A, wherein: Q is a branched or unbranched alkylene, t = 0 or 1 and A is an anionic group selected from the group consisting of OSO₃⁻, SO₃⁻, CO₂⁻, OCO₂⁻, OPO₃²⁻, OPO₃H⁻ and OPO₂⁻; R⁵ is hydrogen or the moiety -CR¹¹R¹²-Y-G_b-Y_c-[(CR⁹R¹⁰)_y-O]_k-R⁸, wherein: each Y is independently selected from the group consisting of O, S, N-H, or N-R⁸; and each R⁸ is independently selected from the group consisting of alkyl, aryl and heteroaryl, said moieties being substituted or unsubstituted, and whether substituted or unsubstituted said moieties having less than 21 carbons; each G is independently selected from the group consisting of CO, SO₂, SO, PO and PO₂; R⁹ and R¹⁰ are independently selected from the group consisting of H and C₁-C₄ alkyl; R¹¹ and R¹² are independently selected from the group consisting of H and alkyl, or when taken together may join to form a carbonyl; b = 0 or 1; c can = 0 or 1, but c must = 0 if b = 0; y is an integer from 1 to 6; k is an integer from 0 to 20; R⁶ is H, or an alkyl, aryl or heteroaryl moiety; said moieties being substituted or unsubstituted; and X, if present, is a suitable charge balancing counterion, preferably X is present when R⁴ is hydrogen, suitable X, include but are not limited to: chloride, bromide, sulphate, methosulphate, sulphonate, p-toluenesulphonate, borontetrafluoride and phosphate.

In one aspect of the present invention, the bleach catalyst has a structure corresponding to general formula below:



wherein R¹³ is a branched alkyl group containing from three to 24 carbon atoms (including the branching carbon atoms) or a linear alkyl group containing from one to 24 carbon atoms; preferably R¹³ is a branched alkyl group containing from eight to 18 carbon atoms or linear alkyl group containing from eight to eighteen carbon atoms; preferably R¹³ is selected from the group consisting of 2-propylheptyl, 2-butyloctyl, 2-pentylononyl, 2-hexyldecyl, n-dodecyl, n-tetradecyl, n-hexadecyl, n-octadecyl, iso-nonyl, iso-decyl, iso-tridecyl and iso-pentadecyl; preferably R¹³ is selected from the group consisting of 2-butyloctyl, 2-pentylononyl, 2-hexyldecyl, iso-tridecyl and iso-pentadecyl.

Preferably the bleach component comprises a source of peracid in addition to bleach catalyst, particularly organic bleach catalyst. The source of peracid may be selected from (a) pre-formed peracid; (b) percarbonate, perborate or persulfate salt (hydrogen peroxide source) preferably in combination with a bleach activator; and (c) perhydrolase enzyme and an ester for forming peracid in situ in the presence of water in a textile or hard surface treatment step.

When present, the peracid and/or bleach activator is generally present in the composition in an amount of from 0.1 to 60wt%, from 0.5 to 40wt% or from 0.6 to 10wt% based on the composition. One or more hydrophobic peracids or precursors thereof may be used in combination with one or more hydrophilic peracid or precursor thereof.

The amounts of hydrogen peroxide source and peracid or bleach activator may be selected such that the molar ratio of available oxygen (from the peroxide source) to peracid is from 1:1 to 35:1, or 2:1 to 10:1.

(6) Metal-containing Bleach Catalysts – The bleach component may be provided by a catalytic metal complex. One type of metal-containing bleach catalyst is a catalyst system comprising a transition metal cation of defined bleach catalytic activity, such as copper, iron, titanium, ruthenium, tungsten, molybdenum, or manganese cations, an auxiliary metal cation having little or no bleach catalytic activity, such as zinc or aluminum cations, and a sequester having defined stability constants for the catalytic and auxiliary metal cations, particularly ethylenediaminetetraacetic acid, ethylenediaminetetra(methylenephosphonic acid) and water-soluble salts thereof. Such catalysts are disclosed in US4430243. Preferred catalysts are described in WO09/839406, US6218351 and WO00/012667. Particularly preferred are transition metal catalyst or ligands therefore that are cross-bridged polydentate N-donor ligands.

If desired, the compositions herein can be catalyzed by means of a manganese compound. Such compounds and levels of use are well known in the art and include, e.g., the manganese-based catalysts disclosed in US5576282.

Cobalt bleach catalysts useful herein are known, and are described e.g. in US5597936; US5595967. Such cobalt catalysts are readily prepared by known procedures, such as taught e.g. in US5597936 and US5595967.

Compositions herein may also suitably include a transition metal complex of ligands such as bispidones (US7501389) and/or macropolycyclic rigid ligands - abbreviated as "MRLs".

As a practical matter, and not by way of limitation, the compositions and processes herein can be adjusted to provide on the order of at least one part per hundred million of the active MRL species in the aqueous washing medium, and will typically provide from 0.005 to 25ppm, from 0.05 to 10ppm, or from 0.1 to 5ppm, of the MRL in the wash liquor.

5 Suitable transition-metals in the instant transition-metal bleach catalyst include e.g. manganese, iron and chromium. Suitable MRLs include 5,12-diethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane. Suitable transition metal MRLs are readily prepared by known procedures, such as taught e.g. in US6225464 and WO00/32601.

10 (7) Photobleaches – suitable photobleaches include e.g. sulfonated zinc phthalocyanine sulfonated aluminium phthalocyanines, xanthene dyes and mixtures thereof. Preferred bleach components for use in the present compositions of the invention comprise a hydrogen peroxide source, bleach activator and/or organic peroxyacid, optionally generated in situ by the reaction of a hydrogen peroxide source and bleach activator, in combination with a bleach catalyst. Preferred bleach components comprise bleach catalysts, preferably organic bleach catalysts, as
15 described above.

Particularly preferred bleach components are the bleach catalysts in particular the organic bleach catalysts.

Exemplary bleaching systems are also described, e.g. in WO2007/087258, WO2007/087244, WO2007/087259 and WO2007/087242.

20 Fabric Hueing Agents - The composition may comprise a fabric hueing agent. Suitable fabric hueing agents include dyes, dye-clay conjugates, and pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Color Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic
25 Violet and Basic Red, or mixtures thereof.


In one aspect, suitable small molecule dyes include small molecule dyes selected from the group consisting of Color Index (Society of Dyers and Colorists, Bradford, UK) numbers Direct Violet 9, Direct Violet 35, Direct Violet 48, Direct Violet 51, Direct Violet 66, Direct Violet 99, Direct Blue 1, Direct Blue 71, Direct Blue 80, Direct Blue 279, Acid Red 17, Acid Red 73,
30 Acid Red 88, Acid Red 150, Acid Violet 15, Acid Violet 17, Acid Violet 24, Acid Violet 43, Acid Red 52, Acid Violet 49, Acid Violet 50, Acid Blue 15, Acid Blue 17, Acid Blue 25, Acid Blue 29, Acid Blue 40, Acid Blue 45, Acid Blue 75, Acid Blue 80, Acid Blue 83, Acid Blue 90 and Acid Blue 113, Acid Black 1, Basic Violet 1, Basic Violet 3, Basic Violet 4, Basic Violet 10, Basic Violet 35, Basic Blue 3, Basic Blue 16, Basic Blue 22, Basic Blue 47, Basic Blue 66, Basic Blue
35 75, Basic Blue 159 and mixtures thereof. In one aspect, suitable small molecule dyes include small molecule dyes selected from the group consisting of Color Index (Society of Dyers and Colorists, Bradford, UK) numbers Acid Violet 17, Acid Violet 43, Acid Red 52, Acid Red 73, Acid Red 88, Acid Red 150, Acid Blue 25, Acid Blue 29, Acid Blue 45, Acid Blue 113, Acid Black 1,

Direct Blue 1, Direct Blue 71, Direct Violet 51 and mixtures thereof. In one aspect, suitable small molecule dyes include small molecule dyes selected from the group consisting of Color Index (Society of Dyers and Colorists, Bradford, UK) numbers Acid Violet 17, Direct Blue 71, Direct Violet 51, Direct Blue 1, Acid Red 88, Acid Red 150, Acid Blue 29, Acid Blue 113 or mixtures thereof.

Suitable polymeric dyes include polymeric dyes selected from the group consisting of polymers containing conjugated chromogens (dye-polymer conjugates) and polymers with chromogens co-polymerized into the backbone of the polymer and mixtures thereof.

In one aspect, suitable polymeric dyes include polymeric dyes selected from the group consisting of fabric-substantive colorants sold under the name of Liquitint® (Milliken), dye-polymer conjugates formed from at least one reactive dye and a polymer selected from the group consisting of polymers comprising a moiety selected from the group consisting of a hydroxyl moiety, a primary amine moiety, a secondary amine moiety, a thiol moiety and mixtures thereof. In still one aspect, suitable polymeric dyes include polymeric dyes selected from the group consisting of Liquitint® Violet CT, carboxymethyl cellulose (CMC) conjugated with a reactive blue, reactive violet or reactive red dye such as CMC conjugated with C.I. Reactive Blue 19, sold by Megazyme, Wicklow, Ireland under the product name AZO-CM-CELLULOSE, product code S-ACMC, alkoxylated triphenyl-methane polymeric colorants, alkoxylated thiophene polymeric colorants, and mixtures thereof.

Preferred hueing dyes include the whitening agents found in WO08/87497. These whitening agents may be characterized by the following structure (I):



(I)

wherein R₁ and R₂ can independently be selected from:



wherein R' is selected from the group consisting of H, CH₃, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein R'' is selected from the group consisting of H, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein x + y ≤ 5; wherein y ≥ 1; and wherein z = 0 to 5;



wherein R' is selected from the group consisting of H, CH₃, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein R'' is selected from the group consisting of H, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein x + y ≤ 10; wherein y ≥ 1; and wherein z = 0 to 5;


c) R₁ = [CH₂CH₂(OR₃)CH₂OR₄] and R₂ = [CH₂CH₂(OR₃)CH₂OR₄]

5 wherein R₃ is selected from the group consisting of H, (CH₂CH₂O)_zH, and mixtures thereof; and wherein z = 0 to 10;

wherein R₄ is selected from the group consisting of (C₁-C₁₆)alkyl, aryl groups, and mixtures thereof; and

d) wherein R₁ and R₂ can independently be selected from the amino addition
10 product of styrene oxide, glycidyl methyl ether, isobutyl glycidyl ether, isopropylglycidyl ether, t-butyl glycidyl ether, 2-ethylhexylglycidyl ether, and glycidylhexadecyl ether, followed by the addition of from 1 to 10 alkylene oxide units.


A preferred whitening agent of the present invention may be characterized by the following structure (II):



15 (II)

wherein R' is selected from the group consisting of H, CH₃, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein R'' is selected from the group consisting of H, CH₂O(CH₂CH₂O)_zH, and mixtures thereof; wherein x + y ≤ 5; wherein y ≥ 1; and wherein z = 0 to 5.

20 A further preferred whitening agent of the present invention may be characterized by the following structure (III):



(III)

typically comprising a mixture having a total of 5 EO groups. Suitable preferred molecules are those in Structure I having the following pendant groups in "part a" above.

TABLE 1

	R1				R2			
	R'	R''	X	y	R'	R''	x	y
A	H	H	3	1	H	H	0	1
B	H	H	2	1	H	H	1	1
c = b	H	H	1	1	H	H	2	1
d = a	H	H	0	1	H	H	3	1

Further whitening agents of use include those described in US2008/34511 (Unilever). A preferred agent is "Violet 13".

5 Suitable dye clay conjugates include dye clay conjugates selected from the group comprising at least one cationic/basic dye and a smectite clay, and mixtures thereof. In one aspect, suitable dye clay conjugates include dye clay conjugates selected from the group consisting of one cationic/basic dye selected from the group consisting of C.I. Basic Yellow 1 through 108, C.I. Basic Orange 1 through 69, C.I. Basic Red 1 through 118, C.I. Basic Violet 1 through 51, C.I. Basic Blue 1 through 164, C.I. Basic Green 1 through 14, C.I. Basic Brown 1 through 23, CI Basic Black 1 through 11, and a clay selected from the group consisting of Montmorillonite clay, Hectorite clay, Saponite clay and mixtures thereof. In still one aspect, suitable dye clay conjugates include dye clay conjugates selected from the group consisting of: Montmorillonite Basic Blue B7 C.I. 42595 conjugate, Montmorillonite Basic Blue B9 C.I. 52015 conjugate, Montmorillonite Basic Violet V3 C.I. 42555 conjugate, Montmorillonite Basic Green G1 C.I. 42040 conjugate, Montmorillonite Basic Red R1 C.I. 45160 conjugate, Montmorillonite C.I. Basic Black 2 conjugate, Hectorite Basic Blue B7 C.I. 42595 conjugate, Hectorite Basic Blue B9 C.I. 52015 conjugate, Hectorite Basic Violet V3 C.I. 42555 conjugate, Hectorite Basic Green G1 C.I. 42040 conjugate, Hectorite Basic Red R1 C.I. 45160 conjugate, Hectorite C.I. Basic Black 2 conjugate, Saponite Basic Blue B7 C.I. 42595 conjugate, Saponite Basic Blue B9 C.I. 52015 conjugate, Saponite Basic Violet V3 C.I. 42555 conjugate, Saponite Basic Green G1 C.I. 42040 conjugate, Saponite Basic Red R1 C.I. 45160 conjugate, Saponite C.I. Basic Black 2 conjugate and mixtures thereof.

25 Suitable pigments include pigments selected from the group consisting of flavanthrone, indanthrone, chlorinated indanthrone containing from 1 to 4 chlorine atoms, pyranthrone, dichloropyranthrone, monobromodichloropyranthrone, dibromodichloropyranthrone, tetrabromopyranthrone, perylene-3,4,9,10-tetracarboxylic acid diimide, wherein the imide groups may be unsubstituted or substituted by C1-C3 -alkyl or a phenyl or heterocyclic radical, and wherein the phenyl and heterocyclic radicals may additionally carry substituents which do not confer solubility in water, anthrapyrimidinecarboxylic acid amides, violanthrone, isoviolanthrone, dioxazine pigments, copper phthalocyanine which may contain up to 2 chlorine

atoms per molecule, polychloro-copper phthalocyanine or polybromochloro-copper phthalocyanine containing up to 14 bromine atoms per molecule and mixtures thereof.

In one aspect, suitable pigments include pigments selected from the group consisting of Ultramarine Blue (C.I. Pigment Blue 29), Ultramarine Violet (C.I. Pigment Violet 15) and mixtures thereof.

The aforementioned fabric hueing agents can be used in combination (any mixture of fabric hueing agents can be used). Suitable hueing agents are described in more detail in US7208459. Preferred levels of dye in compositions of the invention are 0.00001 to 0.5wt%, or 0.0001 to 0.25wt%. The concentration of dyes preferred in water for the treatment and/or cleaning step is from 1ppb to 5ppm, 10ppb to 5ppm or 20ppb to 5ppm. In preferred compositions, the concentration of surfactant will be from 0.2 to 3g/l.

Encapsulates - The composition may comprise an encapsulate. In one aspect, an encapsulate comprising a core, a shell having an inner and outer surface, said shell encapsulating said core.

In one aspect of said encapsulate, said core may comprise a material selected from the group consisting of perfumes; brighteners; dyes; insect repellants; silicones; waxes; flavors; vitamins; fabric softening agents; skin care agents in one aspect, paraffins; enzymes; anti-bacterial agents; bleaches; sensates; and mixtures thereof; and said shell may comprise a material selected from the group consisting of polyethylenes; polyamides; polyvinylalcohols, optionally containing other co-monomers; polystyrenes; polyisoprenes; polycarbonates; polyesters; polyacrylates; aminoplasts, in one aspect said aminoplast may comprise a polyureas, polyurethane, and/or polyureaurethane, in one aspect said polyurea may comprise polyoxymethyleneurea and/or melamine formaldehyde; polyolefins; polysaccharides, in one aspect said polysaccharide may comprise alginate and/or chitosan; gelatin; shellac; epoxy resins; vinyl polymers; water insoluble inorganics; silicone; and mixtures thereof.

In one aspect of said encapsulate, said core may comprise perfume.

In one aspect of said encapsulate, said shell may comprise melamine formaldehyde and/or cross linked melamine formaldehyde.

In a one aspect, suitable encapsulates may comprise a core material and a shell, said shell at least partially surrounding said core material, is disclosed. At least 75%, 85% or 90% of said encapsulates may have a fracture strength of from 0.2 to 10 MPa, from 0.4 to 5MPa, from 0.6 to 3.5 MPa, or from 0.7 to 3MPa; and a benefit agent leakage of from 0 to 30%, from 0 to 20%, or from 0 to 5%.

In one aspect, at least 75%, 85% or 90% of said encapsulates may have a particle size from 1 to 80 microns, from 5 to 60 microns, from 10 to 50 microns, or from 15 to 40 microns.

In one aspect, at least 75%, 85% or 90% of said encapsulates may have a particle wall thickness from 30 to 250nm, from 80 to 180nm, or from 100 to 160nm.

In one aspect, said encapsulates' core material may comprise a material selected from the group consisting of a perfume raw material and/or optionally a material selected from the group consisting of vegetable oil, including neat and/or blended vegetable oils including castor oil, coconut oil, cottonseed oil, grape oil, rapeseed, soybean oil, corn oil, palm oil, linseed oil, safflower oil, olive oil, peanut oil, coconut oil, palm kernel oil, castor oil, lemon oil and mixtures thereof; esters of vegetable oils, esters, including dibutyl adipate, dibutyl phthalate, butyl benzyl adipate, benzyl octyl adipate, tricresyl phosphate, trioctyl phosphate and mixtures thereof; straight or branched chain hydrocarbons, including those straight or branched chain hydrocarbons having a boiling point of greater than about 80°C; partially hydrogenated terphenyls, dialkyl phthalates, alkyl biphenyls, including monoisopropylbiphenyl, alkylated naphthalene, including dipropyl naphthalene, petroleum spirits, including kerosene, mineral oil and mixtures thereof; aromatic solvents, including benzene, toluene and mixtures thereof; silicone oils; and mixtures thereof.

In one aspect, said encapsulates' wall material may comprise a suitable resin including the reaction product of an aldehyde and an amine, suitable aldehydes include, formaldehyde. Suitable amines include melamine, urea, benzoguanamine, glycoluril, and mixtures thereof. Suitable melamines include methylol melamine, methylated methylol melamine, imino melamine and mixtures thereof. Suitable ureas include dimethylol urea, methylated dimethylol urea, urea-resorcinol, and mixtures thereof.

In one aspect, suitable formaldehyde scavengers may be employed with an encapsulate e.g. in a capsule slurry and/or added to a composition before, during or after the encapsulate is added to such composition. Suitable capsules may be made by the following teaching of US2008/0305982; and/or US2009/0247449.

In a preferred aspect the composition can also comprise a deposition aid, preferably consisting of the group comprising cationic or nonionic polymers. Suitable polymers include cationic starches, cationic hydroxyethylcellulose, polyvinylformaldehyde, locust bean gum, mannans, xyloglucans, tamarind gum, polyethyleneterephthalate and polymers containing dimethylaminoethyl methacrylate, optionally with one or monomers selected from the group comprising acrylic acid and acrylamide.

Perfumes - In one aspect the composition comprises a perfume that comprises one or more perfume raw materials selected from the group consisting of 1,1'-oxybis-2-propanol; 1,4-cyclohexanedicarboxylic acid, diethyl ester; (ethoxymethoxy)cyclododecane; 1,3-nonanediol, monoacetate; (3-methylbutoxy)acetic acid, 2-propenyl ester; beta-methyl cyclododecaneethanol; 2-methyl-3-[(1,7,7-trimethylbicyclo[2.2.1]hept-2-yl)oxy]-1-propanol; oxacyclohexadecan-2-one; alpha-methyl-benzenemethanol acetate; trans-3-ethoxy-1,1,5-trimethylcyclohexane; 4-(1,1-dimethylethyl)cyclohexanol acetate; dodecahydro-3a,6,6,9a-tetramethylnaphtho[2,1-b]furan; beta-methyl benzenepropanal; beta-methyl-3-(1-methylethyl)benzenepropanal; 4-phenyl-2-butanone; 2-methylbutanoic acid, ethyl ester;

benzaldehyde; 2-methylbutanoic acid, 1-methylethyl ester; dihydro-5-pentyl-2(3H)furanone; (2E)-1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-2-buten-1-one; dodecanal; undecanal; 2-ethyl- α , α -dimethylbenzenepropanal; decanal; α , α -dimethylbenzeneethanol acetate; 2-(phenylmethylene)octanal; 2-[[3-[4-(1,1-dimethylethyl)phenyl]-2-methylpropylidene]amino]benzoic acid, methyl ester; 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-2-buten-1-one; 2-pentylcyclopentanone; 3-oxo-2-pentyl cyclopentaneacetic acid, methyl ester; 4-hydroxy-3-methoxybenzaldehyde; 3-ethoxy-4-hydroxybenzaldehyde; 2-heptylcyclopentanone; 1-(4-methylphenyl)ethanone; (3E)-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3-buten-2-one; (3E)-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one; benzeneethanol; 2H-1-benzopyran-2-one; 4-methoxybenzaldehyde; 10-undecenal; propanoic acid, phenylmethyl ester; beta-methylbenzenepentanol; 1,1-diethoxy-3,7-dimethyl-2,6-octadiene; α , α -dimethylbenzeneethanol; (2E)-1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-buten-1-one; acetic acid, phenylmethyl ester; cyclohexanepropanoic acid, 2-propenyl ester; hexanoic acid, 2-propenyl ester; 1,2-dimethoxy-4-(2-propenyl)benzene; 1,5-dimethyl-bicyclo[3.2.1]octan-8-one oxime; 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde; 3-buten-2-ol; 2-[[[2,4(or 3,5)-dimethyl-3-cyclohexen-1-yl]methylene]amino]benzoic acid, methyl ester; 8-cyclohexadecen-1-one; methyl ionone; 2,6-dimethyl-7-octen-2-ol; 2-methoxy-4-(2-propenyl)phenol; (2E)-3,7-dimethyl-2,6-Octadien-1-ol; 2-hydroxy-Benzoic acid, (3Z)-3-hexenyl ester; 2-tridecenitrile; 4-(2,2-dimethyl-6-methylenecyclohexyl)-3-methyl-3-buten-2-one; tetrahydro-4-methyl-2-(2-methyl-1-propenyl)-2H-pyran; Acetic acid, (2-methylbutoxy)-, 2-propenyl ester; Benzoic acid, 2-hydroxy-, 3-methylbutyl ester; 2-Buten-1-one, 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (Z)-; Cyclopentanecarboxylic acid, 2-hexyl-3-oxo-, methyl ester; Benzenepropanal, 4-ethyl- α , α -dimethyl-; 3-Cyclohexene-1-carboxaldehyde, 3-(4-hydroxy-4-methylpentyl)-; Ethanone, 1-(2,3,4,7,8,8a-hexahydro-3,6,8,8-tetramethyl-1H-3a,7-methanoazulen-5-yl)-, [3R-(3. α .,3a. β .,7. β .,8a. α .)]-; Undecanal, 2-methyl-2H-Pyran-2-one, 6-butyltetrahydro-; Benzenepropanal, 4-(1,1-dimethylethyl)- α -methyl-; 2(3H)-Furanone, 5-heptyldihydro-; Benzoic acid, 2-[(7-hydroxy-3,7-dimethyloctylidene)amino]-, methyl; Benzoic acid, 2-hydroxy-, phenylmethyl ester; Naphthalene, 2-methoxy-; 2-Cyclopenten-1-one, 2-hexyl-; 2(3H)-Furanone, 5-hexyldihydro-; Oxiranecarboxylic acid, 3-methyl-3-phenyl-, ethyl ester; 2-Oxabicyclo[2.2.2]octane, 1,3,3-trimethyl-; Benzenepentanol, γ -methyl-; 3-Octanol, 3,7-dimethyl-; 3,7-dimethyl-2,6-octadienenitrile; 3,7-dimethyl-6-octen-1-ol; Terpeneol acetate; 2-methyl-6-methylene-7-Octen-2-ol, dihydro derivative; 3a,4,5,6,7,7a-hexahydro-4,7-Methano-1H-inden-6-ol propanoate; 3-methyl-2-buten-1-ol acetate; (Z)-3-Hexen-1-ol acetate; 2-ethyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-buten-1-ol; 4-(octahydro-4,7-methano-5H-inden-5-ylidene)-butanal; 3-2,4-dimethyl-cyclohexene-1-carboxaldehyde; 1-(1,2,3,4,5,6,7,8-octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)-ethanone; 2-hydroxy-benzoic acid, methyl ester; 2-hydroxy-benzoic acid, hexyl ester; 2-phenoxy-ethanol; 2-hydroxy-benzoic acid, pentyl ester; 2,3-heptanedione; 2-hexen-1-ol; 6-Octen-2-ol, 2,6-dimethyl-; damascone (α , β , γ or

delta or mixtures thereof), 4,7-Methano-1H-inden-6-ol, 3a,4,5,6,7,7a-hexahydro-, acetate; 9-Undecenal; 8-Undecenal; Isocyclocitral; Ethanone, 1-(1,2,3,5,6,7,8,8a-octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)-; 3-Cyclohexene-1-carboxaldehyde, 3,5-dimethyl-; 3-Cyclohexene-1-carboxaldehyde, 2,4-dimethyl-; 1,6-Octadien-3-ol, 3,7-dimethyl-; 1,6-Octadien-3-ol, 3,7-dimethyl-, acetate; Lilial (p-t-Bucinal), and Cyclopentanone, 2-[2-(4-methyl-3-cyclohexen-1-yl)propyl]- and 1-methyl-4-(1-methylethenyl)cyclohexene and mixtures thereof.

In one aspect the composition may comprise an encapsulated perfume particle comprising either a water-soluble hydroxylic compound or melamine-formaldehyde or modified polyvinyl alcohol. In one aspect the encapsulate comprises (a) an at least partially water-soluble solid matrix comprising one or more water-soluble hydroxylic compounds, preferably starch; and (b) a perfume oil encapsulated by the solid matrix.

In a further aspect the perfume may be pre-complexed with a polyamine, preferably a polyethylenimine so as to form a Schiff base.

Polymers - The composition may comprise one or more polymers. Examples are carboxymethylcellulose, poly(vinyl-pyrrolidone), poly (ethylene glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid co-polymers.

The composition may comprise one or more amphiphilic cleaning polymers such as the compound having the following general structure: bis((C₂H₅O)(C₂H₄O)_n)(CH₃)-N⁺-C_xH_{2x}-N⁺-(CH₃)-bis((C₂H₅O)(C₂H₄O)_n), wherein n = from 20 to 30, and x = from 3 to 8, or sulphated or sulphonated variants thereof.

The composition may comprise amphiphilic alkoxyated grease cleaning polymers which have balanced hydrophilic and hydrophobic properties such that they remove grease particles from fabrics and surfaces. Specific embodiments of the amphiphilic alkoxyated grease cleaning polymers of the present invention comprise a core structure and a plurality of alkoxyate groups attached to that core structure. These may comprise alkoxyated polyalkylenimines, preferably having an inner polyethylene oxide block and an outer polypropylene oxide block.


Alkoxyated polycarboxylates such as those prepared from polyacrylates are useful herein to provide additional grease removal performance. Such materials are described in WO91/08281 and PCT90/01815. Chemically, these materials comprise polyacrylates having one ethoxy side-chain per every 7-8 acrylate units. The side-chains are of the formula - (CH₂CH₂O)_m (CH₂)_nCH₃ wherein m is 2-3 and n is 6-12. The side-chains are ester-linked to the polyacrylate "backbone" to provide a "comb" polymer type structure. The molecular weight can vary, but is typically in the range of 2000 to 50,000. Such alkoxyated polycarboxylates can comprise from 0.05wt% to 10wt% of the compositions herein.

The isoprenoid-derived surfactants of the present invention, and their mixtures with other cosurfactants and other adjunct ingredients, are particularly suited to be used with an amphiphilic graft co-polymer, preferably the amphiphilic graft co-polymer comprises (i) polyethylene glycol

backbone; and (ii) and at least one pendant moiety selected from polyvinyl acetate, polyvinyl alcohol and mixtures thereof. A preferred amphilic graft co-polymer is Sokalan HP22, supplied from BASF. Suitable polymers include random graft copolymers, preferably a polyvinyl acetate grafted polyethylene oxide copolymer having a polyethylene oxide backbone and multiple polyvinyl acetate side chains. The molecular weight of the polyethylene oxide backbone is preferably 6000 and the weight ratio of the polyethylene oxide to polyvinyl acetate is 40 to 60 and no more than 1 grafting point per 50 ethylene oxide units.

Carboxylate polymer - The composition of the present invention may also include one or more carboxylate polymers such as a maleate/acrylate random copolymer or polyacrylate homopolymer. In one aspect, the carboxylate polymer is a polyacrylate homopolymer having a molecular weight of from 4,000 to 9,000Da, or from 6,000 to 9,000Da.

Soil release polymer - The composition of the present invention may also include one or more soil release polymers having a structure as defined by one of the following structures (I), (II) or (III):



wherein:

a, b and c are from 1 to 200;

d, e and f are from 1 to 50;

Ar is a 1,4-substituted phenylene;

sAr is 1,3-substituted phenylene substituted in position 5 with SO₃Me;

Me is Li, K, Mg/2, Ca/2, Al/3, ammonium, mono-, di-, tri-, or tetraalkylammonium wherein the alkyl groups are C₁-C₁₈ alkyl or C₂-C₁₀ hydroxyalkyl, or mixtures thereof;

R¹, R², R³, R⁴, R⁵ and R⁶ are independently selected from H or C₁-C₁₈ n- or iso-alkyl; and

R⁷ is a linear or branched C₁-C₁₈ alkyl, or a linear or branched C₂-C₃₀ alkenyl, or a cycloalkyl group with 5 to 9 carbon atoms, or a C₈-C₃₀ aryl group, or a C₆-C₃₀ arylalkyl group.

Suitable soil release polymers are polyester soil release polymers such as Repel-o-tex polymers, including Repel-o-tex, SF-2 and SRP6 supplied by Rhodia. Other suitable soil release polymers include Texcare polymers, including Texcare SRA100, SRA300, SRN100, SRN170, SRN240, SRN300 and SRN325 supplied by Clariant. Other suitable soil release polymers are Marloquest polymers, such as Marloquest SL supplied by Sasol.

Cellulosic polymer - The composition of the present invention may also include one or more cellulosic polymers including those selected from alkyl cellulose, alkyl alkoxyalkyl cellulose, carboxyalkyl cellulose, alkyl carboxyalkyl cellulose. In one aspect, the cellulosic polymers are selected from the group comprising carboxymethyl cellulose, methyl cellulose, methyl hydroxyethyl cellulose, methyl carboxymethyl cellulose, and mixtures thereof. In one

aspect, the carboxymethyl cellulose has a degree of carboxymethyl substitution from 0.5 to 0.9 and a molecular weight from 100,000 to 300,000Da.

Enzymes - The composition may comprise one or more enzymes which provide cleaning performance and/or fabric care benefits. Examples of suitable enzymes include, but are not limited to, hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, mannanases, pectate lyases, keratinases, reductases, oxidases, phenoloxidases, lipxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, chlorophyllases, amylases, or mixtures thereof. A typical combination is an enzyme cocktail that may comprise e.g. a protease and lipase in conjunction with amylase. When present in a composition, the aforementioned additional enzymes may be present at levels from 0.00001 to 2wt%, from 0.0001 to 1wt% or from 0.001 to 0.5wt% enzyme protein by weight of the composition.

In general the properties of the selected enzyme(s) should be compatible with the selected detergent, (i.e., pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

In one aspect preferred enzymes would include a cellulase. Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US4435307, US5648263, US5691178, US5776757 and WO89/09259.

Especially suitable cellulases are the alkaline or neutral cellulases having colour care benefits. Examples of such cellulases are cellulases described in EP0495257, EP0531372, WO96/11262, WO96/29397, WO98/08940. Other examples are cellulase variants such as those described in WO94/07998, EP0531315, US5457046, US5686593, US5763254, WO95/24471, WO98/12307 and PCT/DK98/00299.

Commercially available cellulases include Celluzyme™, and Carezyme™ (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

In one aspect preferred enzymes would include a protease. Suitable proteases include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloproteases protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.

The term "subtilases" refers to a sub-group of serine protease according to Siezen et al., Protein Engng. 4 (1991) 719-737 and Siezen et al. Protein Science 6 (1997) 501-523. Serine proteases are a subgroup of proteases characterized by having a serine in the active site, which forms a covalent adduct with the substrate. The subtilases may be divided into 6 sub-divisions, i.e. the Subtilisin family, the Thermitase family, the Proteinase K family, the Lantibiotic peptidase family, the Kexin family and the Pyrolysins family.

Examples of subtilases are those derived from *Bacillus* such as *Bacillus lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US7262042 and WO09/021867, and subtilisin *lentus*, subtilisin *Novo*, subtilisin *Carlsberg*, *Bacillus licheniformis*, subtilisin *BPN'*, subtilisin 309, subtilisin 147 and subtilisin 168 described in WO89/06279 and protease PD138 described in (WO93/18140). Other useful proteases may be those described in WO92/175177, WO01/016285, WO02/026024 and WO02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO89/06270, WO94/25583 and WO05/040372, and the chymotrypsin proteases derived from *Cellulomonas* described in WO05/052161 and WO05/052146.

A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in WO95/23221, and variants thereof which are described in WO92/21760, WO95/23221, EP1921147 and EP1921148.

Examples of metalloproteases are the neutral metalloprotease as described in WO07/044993 (Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*.

Examples of useful proteases are the variants described in: WO92/19729, WO96/034946, WO98/20115, WO98/20116, WO99/011768, WO01/44452, WO03/006602, WO04/03186, WO04/041979, WO07/006305, WO11/036263, WO11/036264, especially the variants with substitutions in one or more of the following positions: 3, 4, 9, 15, 27, 36, 57, 68, 76, 87, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 118, 120, 123, 128, 129, 130, 160, 167, 170, 194, 195, 199, 205, 206, 217, 218, 222, 224, 232, 235, 236, 245, 248, 252 and 274 using the BPN' numbering. More preferred the subtilase variants may comprise the mutations: S3T, V4I, S9R, A15T, K27R, *36D, V68A, N76D, N87S,R, *97E, A98S, S99G,D,A, S99AD, S101G,M,R S103A, V104I,Y,N, S106A, G118V,R, H120D,N, N123S, S128L, P129Q, S130A, G160D, Y167A, R170S, A194P, G195E, V199M, V205I, L217D, N218D, M222S, A232V, K235L, Q236H, Q245R, N252K, T274A (using BPN' numbering).

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Blaze®, Duralase™, Durazym™, Relase®, Relase® Ultra, Savinase®, Savinase® Ultra, Primase®, Polarzyme®, Kannase®, Liquanase®, Liquanase® Ultra, Ovozime®, Coronase®, Coronase® Ultra, Neutrase®, Everlase® and Esperase® all could be sold as Ultra® or Evity® (Novozymes A/S), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Purafect®, Purafect Prime®, Preferenz™, Purafect MA®, Purafect Ox®,

Purafect OxP®, Puramax®, Properase®, Effectenz™, FN2®, FN3®, FN4®, Excellase®, Opticlean® and Optimase® (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and variants hereof (Henkel AG) and KAP (*Bacillus alkalophilus subtilisin*) from Kao.

5 In one aspect preferred enzymes would include an amylase. Suitable amylases may be an alpha-amylase or a glucoamylase and may be of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described in more detail in GB1296839.

10 Suitable amylases include amylases having SEQ ID NO: 3 in WO95/10603 or variants having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred variants are described in WO94/02597, WO94/18314, WO97/43424 and SEQ ID NO: 4 of WO99/019467, such as variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305,
15 391, 408, and 444.

Different suitable amylases include amylases having SEQ ID NO: 6 in WO02/010355 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193.

Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33
20 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO2006/066594 or variants having 90% sequence identity thereof. Preferred variants of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one of more of the following positions: G48, T49, G107, H156, A181, N190, M197, I201, A209 and
25 Q264. Most preferred variants of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO2006/066594 and residues 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;

H156Y+A181T+N190F+A209V+Q264S; or

30 G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S.

Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO99/019467 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269.
35 Particularly preferred amylases are those having deletion in positions R181 and G182, or positions H183 and G184.

Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 2 or SEQ ID NO: 7 of WO96/023873 or variants thereof having 90% sequence

identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred variants of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476. More preferred variants are those having a deletion in positions 181 and 182 or positions 183 and 184. Most preferred amylase variants of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476.

Other amylases which can be used are amylases having SEQ ID NO: 2 of WO08/153815, SEQ ID NO: 10 in WO01/66712 or variants thereof having 90% sequence identity to SEQ ID NO: 2 of WO08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO01/66712. Preferred variants of SEQ ID NO: 10 in WO01/66712 are those having a substitution, a deletion or an insertion in one of more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264.

Further suitable amylases are amylases having SEQ ID NO: 2 of WO09/061380 or variants having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred variants of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or an insertion in one of more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred variants of SEQ ID NO: 2 are those having the substitution in one of more of the following positions: Q87E,R, Q98R, S125A, N128C, T131I, T165I, K178L, T182G, M201L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase variants of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;

N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;

S125A+N128C+K178L+T182G+Y305R+G475K; or

S125A+N128C+T131I+T165I+K178L+T182G+Y305R+G475K wherein the variants are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

Other suitable amylases are the alpha-amylase having SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase variants are those having a substitution, a deletion or an insertion in one of more of the following positions of SEQ ID NO: 12 in WO01/66712: R28, R118, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, K446, Q449, R458, N471, N484. Particular preferred amylases include variants having a deletion of D183 and G184 and having the substitutions R118K, N195F, R320K and R458K, and a variant additionally having substitutions in one or

more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions.

Other examples are amylase variants such as those described in WO2011/098531,
5 WO2013/001078 and WO2013/001087.

Commercially available amylases are Duramyl™, Termamyl™, Termamyl Ultra™, Fungamyl™, Ban™, Stainzyme™, Stainzyme Plus™, Amplify®, Supramyl™, Natalase™, Liquozyme X and BAN™ (from Novozymes A/S), KEMZYM® AT 9000 Biozym Biotech Trading GmbH Wehlstrasse 27b A-1200 Wien Austria, and Rapidase™, Purastar™/Effectenz™,
10 Powerase, Preferenz S100, Preferenx S110, ENZYsize®, OPTISIZE HT PLUS®, and PURASTAR OXAM® (Danisco/DuPont) and KAM® (Kao).

Suitable lipases and cutinases include those of bacterial or fungal origin. Chemically modified or protein engineered mutant enzymes are included. Examples include lipase from *Thermomyces*, e.g. from *T. lanuginosus* (previously named *Humicola lanuginosa*) as described
15 in EP258068 and EP305216, cutinase from *Humicola*, e.g. *H. insolens* (WO96/13580), lipase from strains of *Pseudomonas* (some of these now renamed to *Burkholderia*), e.g. *P. alcaligenes* or *P. pseudoalcaligenes* (EP218272), *P. cepacia* (EP331376), *P. sp.* strain SD705 (WO95/06720 & WO96/27002), *P. wisconsinensis* (WO96/12012), GDSL-type *Streptomyces* lipases (WO10/065455), cutinase from *Magnaporthe grisea* (WO10/107560), cutinase from
20 *Pseudomonas mendocina* (US5,389,536), lipase from *Thermobifida fusca* (WO11/084412, WO13/033318), *Geobacillus stearothermophilus* lipase (WO11/084417), lipase from *Bacillus subtilis* (WO11/084599), and lipase from *Streptomyces griseus* (WO11/150157) and *S. pristinaespiralis* (WO12/137147).

Other examples are lipase variants such as those described in EP407225, WO92/05249,
25 WO94/01541, WO94/25578, WO95/14783, WO95/30744, WO95/35381, WO95/22615, WO96/00292, WO97/04079, WO97/07202, WO00/34450, WO00/60063, WO01/92502, WO07/87508 and WO09/109500.

Preferred commercial lipase products include Lipolase™, Lipex™, Lipex Evity™, Lipex 105T™, Lipolex™ and Lipoclean™ (Novozymes A/S), Lumafast (originally from Genencor) and
30 Lipomax (originally from Gist-Brocades).

Still other examples are lipases sometimes referred to as acyltransferases or perhydrolases, e.g. acyltransferases with homology to *Candida antarctica* lipase A (WO10/111143), acyltransferase from *Mycobacterium smegmatis* (WO05/56782), perhydrolases from the CE 7 family (WO09/67279), and variants of the *M. smegmatis*
35 perhydrolase in particular the S54V variant used in the commercial product Gentle Power Bleach from Huntsman Textile Effects Pte Ltd (WO10/100028).

In one aspect, other preferred enzymes include microbial-derived endoglucanases exhibiting endo-beta-1,4-glucanase activity (EC3.2.1.4), including a bacterial polypeptide

endogenous to a member of the genus *Bacillus* which has a sequence of at least 90%, 94%, 97% or 99% identity to the amino acid sequence SEQ ID NO:2 in US7141403 and mixtures thereof. Suitable endoglucanases are sold under the tradenames Celluclean® and Whitezyme® (Novozymes).

5 Other preferred enzymes include pectate lyases sold under the tradenames Pectawash®, Pectaway®, Xpect® and mannanases sold under the tradenames Mannaway® (Novozymes), and Purabrite® (Danisco/DuPont).


The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of
10 these enzymes. A detergent additive of the invention, *i.e.*, a separate additive or a combined additive, can be formulated, for example, as granulate, liquid, slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, *e.g.* as disclosed in US4106991 and
15 US4661452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty
20 acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP238216.

25 Dye Transfer Inhibiting Agents - The compositions of the present invention may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in a composition, the dye transfer
30 inhibiting agents may be present at levels from 0.0001 to 10wt%, from 0.01 to 5wt% or from 0.1 to 3wt%.

Brighteners - The compositions of the present invention can also contain additional components that may tint articles being cleaned, such as fluorescent brighteners.

The composition may comprise C.I. fluorescent brightener 260 in alpha-crystalline form
35 having the following structure:




In one aspect, the brightener is a cold water soluble brightener, such as the C.I. fluorescent brightener 260 in alpha-crystalline form. In one aspect the brightener is predominantly in alpha-crystalline form, which means that typically at least 50wt%, at least 75wt%, at least 90wt%, at least 99wt%, or even substantially all, of the C.I. fluorescent brightener 260 is in alpha-crystalline form.

The brightener is typically in micronized particulate form, having a weight average primary particle size of from 3 to 30 micrometers, from 3 micrometers to 20 micrometers, or from 3 to 10 micrometers.

The composition may comprise C.I. fluorescent brightener 260 in beta-crystalline form, and the weight ratio of: (i) C.I. fluorescent brightener 260 in alpha-crystalline form, to (ii) C.I. fluorescent brightener 260 in beta-crystalline form may be at least 0.1, or at least 0.6. BE680847 relates to a process for making C.I. fluorescent brightener 260 in alpha-crystalline form.

Commercial optical brighteners which may be useful in the present invention can be classified into subgroups, which include, but are not necessarily limited to, derivatives of stilbene, pyrazoline, coumarin, carboxylic acid, methinecyanines, dibenzothiophene-5,5-dioxide, azoles, 5- and 6-membered-ring heterocycles, and other miscellaneous agents. Examples of such brighteners are disclosed in "The Production and Application of Fluorescent Brightening Agents", M. Zahradnik, Published by John Wiley & Sons, New York (1982). Specific nonlimiting examples of optical brighteners which are useful in the present compositions are those identified in US4790856 and US3646015.

A further suitable brightener has the structure below:



Suitable fluorescent brightener levels include lower levels of from 0.01wt%, from 0.05wt%, from 0.1wt% or from 0.2wt% to upper levels of 0.5wt% or 0.75wt%.

In one aspect the brightener may be loaded onto a clay to form a particle. Silicate salts - The compositions of the present invention can also contain silicate salts, such as sodium or potassium silicate. The composition may comprise of from 0wt% to less than 10wt% silicate salt, to 9wt%, or to 8wt%, or to 7wt%, or to 6wt%, or to 5wt%, or to 4wt%, or to 3wt%, or even to 2wt%, and from above 0wt%, or from 0.5wt%, or from 1wt% silicate salt. A suitable silicate salt is sodium silicate.

Dispersants - The compositions of the present invention can also contain dispersants. Suitable water-soluble organic materials include the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

Enzyme Stabilizers - Enzymes for use in compositions can be stabilized by various techniques. The enzymes employed herein can be stabilized by the presence of water-soluble sources of calcium and/or magnesium ions. Examples of conventional stabilizing agents are, e.g. a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, a peptide aldehyde, lactic acid, boric acid, or a boric acid derivative, e.g. an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in, for example, WO92/19709 and WO92/19708. In case of aqueous compositions comprising protease, a reversible protease inhibitor, such as a boron compound including borate, 4-formyl phenylboronic acid, phenylboronic acid and derivatives thereof, or compounds such as calcium formate, sodium formate and 1,2-propane diol can be added to further improve stability. The peptide aldehyde may be of the formula $B_2-B_1-B_0-R$ wherein: R is hydrogen, CH_3 , CX_3 , CHX_2 , or CH_2X , wherein X is a halogen atom; B_0 is a phenylalanine residue with an OH substituent at the *p*-position and/or at the *m*-position; B_1 is a single amino acid residue; and B_2 consists of one or more amino acid residues, optionally comprising an N-terminal protection group. Preferred peptide aldehydes include but are not limited to: Z-RAY-H, Ac-GAY-H, Z-GAY-H, Z-GAL-H, Z-GAF-H, Z-GAV-H, Z-RVY-H, Z-LVY-H, Ac-LGAY-H, Ac-FGAY-H, Ac-YGAY-H, Ac-FGVY-H or Ac-WLVY-H, where Z is benzyloxycarbonyl and Ac is acetyl.

Solvents - Suitable solvents include water and other solvents such as lipophilic fluids. Examples of suitable lipophilic fluids include siloxanes, other silicones, hydrocarbons, glycol ethers, glycerine derivatives such as glycerine ethers, perfluorinated amines, perfluorinated and hydrofluoroether solvents, low-volatility nonfluorinated organic solvents, diol solvents, other environmentally-friendly solvents and mixtures thereof.

Structurant/Thickeners - Structured liquids can either be internally structured, whereby the structure is formed by primary ingredients (e.g. surfactant material) and/or externally structured by providing a three dimensional matrix structure using secondary ingredients (e.g. polymers, clay and/or silicate material). The composition may comprise a structurant, from 0.01 to 5wt%, or from 0.1 to 2.0wt%. The structurant is typically selected from the group consisting of diglycerides and triglycerides, ethylene glycol distearate, microcrystalline cellulose, cellulose-

based materials, microfiber cellulose, hydrophobically modified alkali-swellaable emulsions such as Polygel W30 (3VSigma), biopolymers, xanthan gum, gellan gum, and mixtures thereof. A suitable structurant includes hydrogenated castor oil, and non-ethoxylated derivatives thereof. A suitable structurant is disclosed in US6855680. Such structurants have a thread-like structuring system having a range of aspect ratios. Other suitable structurants and the processes for making them are described in WO10/034736.

Conditioning Agents - The composition of the present invention may include a high melting point fatty compound. The high melting point fatty compound useful herein has a melting point of 25°C or higher, and is selected from the group consisting of fatty alcohols, fatty acids, fatty alcohol derivatives, fatty acid derivatives, and mixtures thereof. Such compounds of low melting point are not intended to be included in this section. Non-limiting examples of the high melting point compounds are found in International Cosmetic Ingredient Dictionary, Fifth Edition, 1993, and CTFA Cosmetic Ingredient Handbook, Second Edition, 1992.

The high melting point fatty compound is included in the composition at a level of from 0.1 to 40wt%, from 1 to 30wt%, from 1.5 to 16wt%, from 1.5 to 8wt% in view of providing improved conditioning benefits such as slippery feel during the application to wet hair, softness and moisturized feel on dry hair.


The compositions of the present invention may contain a cationic polymer. Concentrations of the cationic polymer in the composition typically range from 0.05 to 3wt%, from 0.075 to 2.0wt%, or from 0.1 to 1.0wt%. Suitable cationic polymers will have cationic charge densities of at least 0.5 meq/gm, at least 0.9 meq/gm, at least 1.2 meq/gm, at least 1.5 meq/gm, or less than 7 meq/gm, and less than 5 meq/gm, at the pH of intended use of the composition, which pH will generally range from pH3 to pH9, or between pH4 and pH8. Herein, "cationic charge density" of a polymer refers to the ratio of the number of positive charges on the polymer to the molecular weight of the polymer. The average molecular weight of such suitable cationic polymers will generally be between 10,000 and 10 million, between 50,000 and 5 million, or between 100,000 and 3 million.

Suitable cationic polymers for use in the compositions of the present invention contain cationic nitrogen-containing moieties such as quaternary ammonium or cationic protonated amino moieties. Any anionic counterions can be used in association with the cationic polymers so long as the polymers remain soluble in water, in the composition, or in a coacervate phase of the composition, and so long as the counterions are physically and chemically compatible with the essential components of the composition or do not otherwise unduly impair composition performance, stability or aesthetics. Nonlimiting examples of such counterions include halides (e.g., chloride, fluoride, bromide, iodide), sulfate and methylsulfate.

Nonlimiting examples of such polymers are described in the CTFA Cosmetic Ingredient Dictionary, 3rd edition, edited by Estrin, Crosley, and Haynes, (The Cosmetic, Toiletry, and Fragrance Association, Inc., Washington, D.C. (1982)).

Other suitable cationic polymers for use in the composition include polysaccharide polymers, cationic guar gum derivatives, quaternary nitrogen-containing cellulose ethers, synthetic polymers, copolymers of etherified cellulose, guar and starch. When used, the cationic polymers herein are either soluble in the composition or are soluble in a complex
 5 coacervate phase in the composition formed by the cationic polymer and the anionic, amphoteric and/or zwitterionic surfactant component described hereinbefore. Complex coacervates of the cationic polymer can also be formed with other charged materials in the composition. Suitable cationic polymers are described in US3962418; US3958581; and US2007/0207109.

10 The composition of the present invention may include a nonionic polymer as a conditioning agent. Polyalkylene glycols having a molecular weight of more than 1000 are useful herein. Useful are those having the following general formula:



wherein R^{95} is selected from the group consisting of H, methyl, and mixtures thereof.

15 Conditioning agents, and in particular silicones, may be included in the composition. The conditioning agents useful in the compositions of the present invention typically comprise a water insoluble, water dispersible, non-volatile, liquid that forms emulsified, liquid particles. Suitable conditioning agents for use in the composition are those conditioning agents characterized generally as silicones (e.g., silicone oils, cationic silicones, silicone gums, high
 20 refractive silicones, and silicone resins), organic conditioning oils (e.g., hydrocarbon oils, polyolefins, and fatty esters) or combinations thereof, or those conditioning agents which otherwise form liquid, dispersed particles in the aqueous surfactant matrix herein. Such conditioning agents should be physically and chemically compatible with the essential components of the composition, and should not otherwise unduly impair composition stability,
 25 aesthetics or performance.

The concentration of the conditioning agent in the composition should be sufficient to provide the desired conditioning benefits. Such concentration can vary with the conditioning agent, the conditioning performance desired, the average size of the conditioning agent particles, the type and concentration of other components, and other like factors.

30 The concentration of the silicone conditioning agent typically ranges from 0.01 to 10wt%. Non-limiting examples of suitable silicone conditioning agents, and optional suspending agents for the silicone, are described in U.S. Reissue Pat. No. 34,584; US5104646; US5106609; US4152416; US2826551; US3964500; US4364837; US6607717; US6482969; US5807956; US5981681; US6207782; US7465439; US7041767; US7217777; US2007/0286837A1;
 35 US2005/0048549A1; US2007/0041929A1; GB849433; DE10036533, which are all incorporated herein by reference; Chemistry and Technology of Silicones, New York: Academic Press (1968); General Electric Silicone Rubber Product Data Sheets SE 30, SE 33, SE 54 and SE 76;

Silicon Compounds, Petrarch Systems, Inc. (1984); and in Encyclopedia of Polymer Science and Engineering, vol. 15, 2d ed., pp 204-308, John Wiley & Sons, Inc. (1989).

The compositions of the present invention may also comprise from 0.05 to 3wt% of at least one organic conditioning oil as the conditioning agent, either alone or in combination with other conditioning agents, such as the silicones (described herein). Suitable conditioning oils include hydrocarbon oils, polyolefins, and fatty esters. Also suitable for use in the compositions herein are the conditioning agents described in US5674478 and US5750122 or in US4529586; US4507280; US4663158; US4197865; US4217914; US4381919; and US4422853.

Hygiene and malodour – The compositions of the present invention may also comprise one or more of zinc ricinoleate, thymol, quaternary ammonium salts such as Bardac®, polyethylenimines (such as Lupasol® from BASF) and zinc complexes thereof, silver and silver compounds, especially those designed to slowly release Ag⁺ or nano-silver dispersions.

Probiotics – The compositions may comprise probiotics such as those described in WO09/043709.

Suds Boosters - If high sudsing is desired, suds boosters such as the C₁₀-C₁₆ alkanolamides or C₁₀-C₁₄ alkyl sulphates can be incorporated into the compositions, typically at 1 to 10wt% levels. The C₁₀-C₁₄ monoethanol and diethanol amides illustrate a typical class of such suds boosters. Use of such suds boosters with high sudsing adjunct surfactants such as the amine oxides, betaines and sultaines noted above is also advantageous. If desired, water-soluble magnesium and/or calcium salts such as MgCl₂, MgSO₄, CaCl₂, CaSO₄ and the like, can be added at levels of, typically, 0.1 to 2wt%, to provide additional suds and to enhance grease removal performance.

Suds Suppressors - Compounds for reducing or suppressing the formation of suds can be incorporated into the compositions of the present invention. Suds suppression can be of particular importance in the so-called "high concentration cleaning process" as described in US4489455 and US4489574, and in front-loading -style washing machines. A wide variety of materials may be used as suds suppressors, and suds suppressors are well known to those skilled in the art. See e.g. Kirk Othmer Encyclopedia of Chemical Technology, Third Edition, Volume 7, p.430-447 (John Wiley & Sons, Inc., 1979). Examples of suds suppressors include monocarboxylic fatty acid and soluble salts therein, high molecular weight hydrocarbons such as paraffin, fatty acid esters (e.g., fatty acid triglycerides), fatty acid esters of monovalent alcohols, aliphatic C₁₈-C₄₀ ketones (e.g., stearone), N-alkylated amino triazines, waxy hydrocarbons preferably having a melting point below about 100°C, silicone suds suppressors, and secondary alcohols. Suds suppressors are described in US2954347; US4265779; US4265779; US3455839; US3933672; US4652392; US4978471; US4983316; US5288431; US4639489; US4749740; US4798679; US4075118; EP89307851.9; EP150872; and DOS 2,124,526.

For any detergent compositions to be used in automatic laundry washing machines, suds should not form to the extent that they overflow the washing machine. Suds suppressors, when utilized, are preferably present in a "suds suppressing amount. By "suds suppressing amount" is meant that the formulator of the composition can select an amount of this suds
5 controlling agent that will sufficiently control the suds to result in a low-sudsing laundry detergent for use in automatic laundry washing machines.

The compositions herein will generally comprise from 0 to 10wt% of suds suppressor. When utilized as suds suppressors, monocarboxylic fatty acids, and salts therein, will be present typically in amounts up to 5wt%. Preferably, from 0.5 to 3wt% of fatty
10 monocarboxylate suds suppressor is utilized. Silicone suds suppressors are typically utilized in amounts up to 2.0wt%, although higher amounts may be used. Monostearyl phosphate suds suppressors are generally utilized in amounts ranging from 0.1 to 2wt%. Hydrocarbon suds suppressors are typically utilized in amounts ranging from 0.01 to 5.0wt%, although higher levels can be used. The alcohol suds suppressors are typically used at 0.2 to 3wt%.

The compositions herein may have a cleaning activity over a broad range of pH. In
15 certain embodiments the compositions have cleaning activity from pH4 to pH11.5. In other embodiments, the compositions are active from pH6 to pH11, from pH7 to pH11, from pH8 to pH11, from pH9 to pH11, or from pH10 to pH11.5.

The compositions herein may have cleaning activity over a wide range of temperatures,
20 e.g., from 10°C or lower to 90°C. Preferably the temperature will be below 50°C or 40°C or even 30°C. In certain embodiments, the optimum temperature range for the compositions is from 10°C to 20°C, from 15°C to 25°C, from 15°C to 30°C, from 20°C to 30°C, from 25°C to 35°C, from 30°C to 40°C, from 35°C to 45°C, or from 40°C to 50°C.

25 **Form of the composition**

The compositions described herein are advantageously employed for example, in laundry applications, hard surface cleaning, dishwashing applications, as well as cosmetic applications such as dentures, teeth, hair and skin. The compositions of the invention are in particular solid or liquid cleaning and/or treatment compositions. In one aspect the invention
30 relates to a composition, wherein the form of the composition is selected from the group consisting of a regular, compact or concentrated liquid; a gel; a paste; a soap bar; a regular or a compacted powder; a granulated solid; a homogenous or a multilayer tablet with two or more layers (same or different phases); a pouch having one or more compartments; a single or a multi-compartment unit dose form; or any combination thereof.

The form of the composition may separate the components physically from each other in compartments such as e.g. water dissolvable pouches or in different layers of tablets. Thereby
35 negative storage interaction between components can be avoided. Different dissolution profiles of each of the compartments can also give rise to delayed dissolution of selected components in the

wash solution.

Pouches can be configured as single or multicompartments. It can be of any form, shape and material which is suitable for hold the composition, e.g. without allowing the release of the composition to release of the composition from the pouch prior to water contact. The pouch is made from water soluble film which encloses an inner volume. Said inner volume can be divided into compartments of the pouch. Preferred films are polymeric materials preferably polymers which are formed into a film or sheet. Preferred polymers, copolymers or derivatives thereof are selected polyacrylates, and water soluble acrylate copolymers, methyl cellulose, carboxy methyl cellulose, sodium dextrin, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, malto dextrin, poly methacrylates, most preferably polyvinyl alcohol copolymers and, hydroxypropyl methyl cellulose (HPMC). Preferably the level of polymer in the film for example PVA is at least about 60%. Preferred average molecular weight will typically be about 20,000 to about 150,000. Films can also be of blended compositions comprising hydrolytically degradable and water soluble polymer blends such as polylactide and polyvinyl alcohol (known under the Trade reference M8630 as sold by MonoSol LLC, Indiana, USA) plus plasticisers like glycerol, ethylene glycerol, propylene glycol, sorbitol and mixtures thereof. The pouches can comprise a solid laundry cleaning composition or part components and/or a liquid cleaning composition or part components separated by the water soluble film. The compartment for liquid components can be different in composition than compartments containing solids (US2009/0011970 A1).

Lipase particles

The lipase variants of the present invention comprised in a water-soluble film may be present as lipase particles. The lipase particles may even contain one or more additional enzymes, as described below.

Lipase particles are any form of lipase variant in a solid particulate form. That can be as lipase crystals, lipase precipitate, spray or freeze-dried lipase or any form of granulated lipase, either as a powder or a suspension in liquid. Typically the particle size, measured as equivalent spherical diameter (volume based average particle size), of the lipase particles is below 2mm, preferably below 1mm, below 0.5mm, below 0.25mm, or below 0.1mm; and above 0.05um, preferably above 0.1um, above 0.5um, above 1um, above 5um or above 10um. In a preferred embodiment, the particle size of the lipase particles is from 0.5um to 100um.

The lipase particles contain at least 1% w/w lipase protein, preferably at least 5% w/w lipase protein, at least 10% w/w lipase protein, at least 20% w/w lipase protein, at least 30% w/w lipase protein, at least 40% w/w lipase protein, at least 50% w/w lipase protein, at least 60% w/w lipase protein, at least 70% w/w lipase protein, at least 80% w/w lipase protein, or at least 90% w/w lipase protein.

In a preferred embodiment, the lipase particles are lipase crystals, or the lipase protein is on a crystalline form.

Enzyme crystallization may be carried out in a number of ways, as known in the art (e.g., as described in WO91/09943 or WO94/22903).

The lipase may be formulated in the lipase particle as known in the art for solid enzyme formulations, such as formulations for reducing dust, improving stability and/or modifying release rate of the enzyme. The lipase particle may also be formulated in a matrix or coated with agents suppressing dissolution of the enzyme particle in the PVOH/film solution used for preparing the water-soluble film.

The lipase molecules on the surface of the lipase particles may also be cross-linked, like CLECs (Cross-Linked Enzyme Crystals) or CLEA (Cross-Linked Enzyme Aggregate).

Water-soluble film

Water-soluble films, optional ingredients for use therein, and methods of making the same are well known in the art. In one class of embodiments, the water-soluble film includes PVOH. PVOH is a synthetic resin generally prepared by the alcoholysis, usually termed hydrolysis or saponification, of polyvinyl acetate. Fully hydrolyzed PVOH, wherein virtually all the acetate groups have been converted to alcohol groups, is a strongly hydrogen-bonded, highly crystalline polymer which dissolves only in hot water - greater than about 140°F (60°C). If a sufficient number of acetate groups are allowed to remain after the hydrolysis of polyvinyl acetate, then the PVOH polymer, being known as partially hydrolyzed, it is more weakly hydrogen-bonded and less crystalline and are soluble in cold water - less than about 50°F (10°C). An intermediate cold/hot water-soluble film can include, for example, intermediate partially-hydrolyzed PVOH (e.g., with degrees of hydrolysis of about 94% to about 98%), and is readily soluble only in warm water - e.g., rapid dissolution at temperatures of about 40°C and greater. Both fully and partially hydrolyzed PVOH types are commonly referred to as PVOH homopolymers although the partially hydrolyzed type is technically a vinyl alcohol-vinyl acetate copolymer.

The degree of hydrolysis of the PVOH included in the water-soluble films of the present disclosure can be about 75% to about 99%. As the degree of hydrolysis is reduced, a film made from the resin will have reduced mechanical strength but faster solubility at temperatures below about 20°C. As the degree of hydrolysis increases, a film made from the resin will tend to be mechanically stronger and the thermoformability will tend to decrease. The degree of hydrolysis of the PVOH can be chosen such that the water-solubility of the resin is temperature dependent, and thus the solubility of a film made from the resin, compatibilizing agent, and additional ingredients is also influenced. In one class of embodiments the film is cold water-soluble. A cold water-soluble film, soluble in water at a temperature of less than 10°C, can include PVOH with a degree of hydrolysis in a range of about 75% to about 90%, or in a range of about 80% to about 90%, or in a range of about 85% to about 90%. In another class of embodiments the film is hot water-soluble. A hot water-soluble film, soluble in water at a temperature of at least about 60°C, can include PVOH with a degree of hydrolysis of at least about 98%.

Other film-forming resins for use in addition to or in an alternative to PVOH can include, but are not limited to, modified polyvinyl alcohols, polyacrylates, water-soluble acrylate copolymers, polyacrylates, polyacrylamides, polyvinyl pyrrolidone, pullulan, water-soluble natural polymers including, but not limited to, guar gum, xanthan gum, carrageenan, and starch, water-soluble polymer derivatives including, but not limited to, ethoxylated starch and hydroxypropylated starch, poly(sodium acrylamido-2-methylpropane sulfonate), polymonomethylmaleate, copolymers thereof, and combinations of any of the foregoing. In one class of embodiments, the film-forming resin is a terpolymer consisting of vinyl alcohol, vinyl acetate, and sodium acrylamido-2-methylpropanesulfonate. Unexpectedly, water-soluble films based on a vinyl alcohol, vinyl acetate, and sodium acrylamido-2-methylpropanesulfonate terpolymer have demonstrated a high percent recovery of enzyme.

The water-soluble resin can be included in the water-soluble film in any suitable amount, for example an amount in a range of about 35 wt% to about 90 wt%. The preferred weight ratio of the amount of the water-soluble resin as compared to the combined amount of all enzymes, enzyme stabilizers, and secondary additives can be any suitable ratio, for example a ratio in a range of about 0.5 to about 5, or about 1 to 3, or about 1 to 2.

Water-soluble resins for use in the films described herein (including, but not limited to PVOH resins) can be characterized by any suitable viscosity for the desired film properties, optionally a viscosity in a range of about 5.0 to about 30.0 cP, or about 10.0 cP to about 25 cP. The viscosity of a PVOH resin is determined by measuring a freshly made solution using a Brookfield LV type viscometer with UL adapter as described in British Standard EN ISO 15023-2:2006 Annex E Brookfield Test method. It is international practice to state the viscosity of 4% aqueous polyvinyl alcohol solutions at 20°C. All PVOH viscosities specified herein in cP should be understood to refer to the viscosity of 4% aqueous polyvinyl alcohol solution at 20°C, unless specified otherwise.

It is well known in the art that the viscosity of a PVOH resin is correlated with the weight average molecular weight ($\overline{M}_w$) of the same PVOH resin, and often the viscosity is used as a proxy for $\overline{M}_w$. Thus, the weight average molecular weight of the water-soluble resin optionally can be in a range of about 35,000 to about 190,000, or about 80,000 to about 160,000. The molecular weight of the resin need only be sufficient to enable it to be molded by suitable techniques to form a thin plastic film.

The water-soluble films according to the present disclosure may include other optional additive ingredients including, but not limited to, plasticizers, surfactants, defoamers, film formers, antiblocking agents, internal release agents, anti-yellowing agents and other functional ingredients, for example in amounts suitable for their intended purpose.

Water is recognized as a very efficient plasticizer for PVOH and other polymers; however, the volatility of water makes its utility limited since polymer films need to have at least

some resistance (robustness) to a variety of ambient conditions including low and high relative humidity. Glycerin is much less volatile than water and has been well established as an effective plasticizer for PVOH and other polymers. Glycerin or other such liquid plasticizers by themselves can cause surface "sweating" and greasiness if the level used in the film formulation is too high. This can lead to problems in a film such as unacceptable feel to the hand of the consumer and even blocking of the film on the roll or in stacks of sheets if the sweating is not mitigated in some manner, such as powdering of the surface. This could be characterized as over plasticization. However, if too little plasticizer is added to the film the film may lack sufficient ductility and flexibility for many end uses, for example to be converted into a final use format such as pouches.

Plasticizers for use in water-soluble films of the present disclosure include, but are not limited to, sorbitol, glycerol, diglycerol, propylene glycol, ethylene glycol, diethyleneglycol, triethylene glycol, tetraethyleneglycol, polyethylene glycols up to MW 400, 2 methyl 1, 3 propane diol, lactic acid, monoacetin, triacetin, triethyl citrate, 1,3-butanediol, trimethylolpropane (TMP), polyether triol, and combinations thereof. Polyols, as described above, are generally useful as plasticizers. As less plasticizer is used, the film can become more brittle, whereas as more plasticizer is used the film can lose tensile strength. Plasticizers can be included in the water-soluble films in an amount in a range of about 25 phr to about 50 phr, or from about 30 phr to about 45 phr, or from about 32 phr to about 42 phr, for example.

Surfactants for use in water-soluble films are well known in the art. Optionally, surfactants are included to aid in the dispersion of the resin solution upon casting. Suitable surfactants for water-soluble films of the present disclosure include, but are not limited to, dialkyl sulfosuccinates, lactylated fatty acid esters of glycerol and propylene glycol, lactic esters of fatty acids, sodium alkyl sulfates, polysorbate 20, polysorbate 60, polysorbate 65, polysorbate 80, alkyl polyethylene glycol ethers, lecithin, acetylated fatty acid esters of glycerol and propylene glycol, sodium lauryl sulfate, acetylated esters of fatty acids, myristyl dimethylamine oxide, trimethyl tallow alkyl ammonium chloride, quaternary ammonium compounds, salts thereof and combinations of any of the foregoing. Thus, surfactants can be included in the water-soluble films in an amount of less than about 2 phr, for example less than about 1 phr, or less than about 0.5 phr, for example.

One type of secondary component contemplated for use is a defoamer. Defoamers can aid in coalescing of foam bubbles. Suitable defoamers for use in water-soluble films according to the present disclosure include, but are not limited to, hydrophobic silicas, for example silicon dioxide or fumed silica in fine particle sizes, including Foam Blast® defoamers available from Emerald Performance Materials, including Foam Blast® 327, Foam Blast® UVD, Foam Blast® 163, Foam Blast® 269, Foam Blast® 338, Foam Blast® 290, Foam Blast® 332, Foam Blast® 349, Foam Blast® 550 and Foam Blast® 339, which are proprietary, non-mineral oil defoamers. In embodiments, defoamers can be used in an amount of 0.5 phr, or less, for example, 0.05 phr,

0.04 phr, 0.03 phr, 0.02 phr, or 0.01 phr. Preferably, significant amounts of silicon dioxide will be avoided, in order to avoid stress whitening.

Processes for making water-soluble articles, including films, include casting, blow-molding, extrusion and blown extrusion, as known in the art. One contemplated class of
5 embodiments is characterized by the water-soluble film described herein being formed by casting, for example, by admixing the ingredients described herein with water to create an aqueous mixture, for example a solution with optionally dispersed solids, applying the mixture to a surface, and drying off water to create a film. Similarly, other compositions can be formed by drying the mixture while it is confined in a desired shape.

10 In one contemplated class of embodiments, the water-soluble film is formed by casting a water-soluble mixture wherein the water-soluble mixture is prepared according to the steps of:

(a) providing a mixture of water-soluble resin, water, and any optional additives excluding plasticizers;

(b) boiling the mixture for 30 minutes;

15 (c) degassing the mixture in an oven at a temperature of at least 40°C; optionally in a range of 40°C to 70°C, e.g., about 65°C;

(d) adding one or more enzymes, plasticizer, and additional water to the mixture at a temperature of 65°C or less; and

(e) stirring the mixture without vortex until the mixture appears substantially uniform in color and
20 consistency; optionally for a time period in a range of 30 minutes to 90 minutes, optionally at least 1 hour; and

(f) casting the mixture promptly after the time period of stirring (e.g., within 4 hours, or 2 hours, or 1 hour).

If the enzyme is added to the mixture too early, e.g., with the secondary additives or
25 resin, the activity of the enzyme may decrease. Without intending to be bound by any particular theory, it is believed that boiling of the mixture with the enzyme leads to the enzyme denaturing and storing in solution for extended periods of time also leads to a reduction in enzyme activity.

In one class of embodiments, high enzyme activity is maintained in the water-soluble films according to the present disclosure by drying the films quickly under moderate to mild
30 conditions. As used herein, drying quickly refers to a drying time of less than 24 hours, optionally less than 12 hours, optionally less than 8 hours, optionally less than 2 hours, optionally less than 1 hour, optionally less than 45 minutes, optionally less than 30 minutes, optionally less than 20 minutes, optionally less than 10 minutes, for example in a range of about 6 minutes to about 10 minutes, or 8 minutes. As used herein, moderate to mild conditions refer
35 to drying temperatures of less than 170°F (77°C), optionally in a range of about 150°F to about 170°F (about 66°C to about 77°C), e.g., 165°F (74°C). As the drying temperature increases, the enzymes tend to denature faster, whereas as the drying temperature decreases, the drying time increases, thus exposing the enzymes to solution for an extended period of time.

The film is useful for creating a packet to contain a composition, for example laundry or dishwashing compositions, thereby forming a pouch. The film described herein can also be used to make a packet with two or more compartments made of the same film or in combination with films of other polymeric materials. Additional films can, for example, be obtained by casting, 5 blow-molding, extrusion or blown extrusion of the same or a different polymeric material, as known in the art. In one type of embodiment, the polymers, copolymers or derivatives thereof suitable for use as the additional film are selected from polyvinyl alcohols, polyvinyl pyrrolidone, polyalkylene oxides, polyacrylic acid, cellulose, cellulose ethers, cellulose esters, cellulose amides, polyvinyl acetates, polycarboxylic acids and salts, polyaminoacids or peptides, 10 polyamides, polyacrylamide, copolymers of maleic/acrylic acids, polysaccharides including starch and gelatin, natural gums such as xanthan, and carrageenans. For example, polymers can be selected from polyacrylates and water-soluble acrylate copolymers, methylcellulose, carboxymethylcellulose sodium, dextrin, ethylcellulose, hydroxyethyl cellulose, hydroxypropyl methylcellulose, maltodextrin, polymethacrylates, and combinations thereof, or selected from 15 polyvinyl alcohols, polyvinyl alcohol copolymers and hydroxypropyl methyl cellulose (HPMC), and combinations thereof.

The pouches and/or packets of the present disclosure comprise at least one sealed compartment. Thus the pouches may comprise a single compartment or multiple compartments. The pouches may have regions with and without enzymes. In embodiments including multiple 20 compartments, each compartment may contain identical and/or different compositions. In turn, the compositions may take any suitable form including, but not limited to liquid, solid and combinations thereof (e.g., a solid suspended in a liquid). In some embodiments, the pouches comprises a first, second and third compartment, each of which respectively contains a different first, second and third composition. In some embodiments, the compositions may be visually 25 distinct as described in EP 2258820.

The compartments of multi-compartment pouches and/or packets may be of the same or different size(s) and/or volume(s). The compartments of the present multi-compartment pouches can be separate or conjoined in any suitable manner. In some embodiments, the second and/or third and/or subsequent compartments are superimposed on the first compartment. In one 30 aspect, the third compartment may be superimposed on the second compartment, which is in turn superimposed on the first compartment in a sandwich configuration. Alternatively the second and third compartments may be superimposed on the first compartment. However it is also equally envisaged that the first, second and optionally third and subsequent compartments may be attached to one another in a side by side relationship. The compartments may be 35 packed in a string, each compartment being individually separable by a perforation line. Hence each compartment may be individually torn-off from the remainder of the string by the end-user.

In some embodiments, multi-compartment pouches and/or packets include three compartments consisting of a large first compartment and two smaller compartments. The

second and third smaller compartments are superimposed on the first larger compartment. The size and geometry of the compartments are chosen such that this arrangement is achievable. The geometry of the compartments may be the same or different. In some embodiments the second and optionally third compartment each has a different geometry and shape as compared to the first compartment. In these embodiments, the second and optionally third compartments are arranged in a design on the first compartment. The design may be decorative, educative, or illustrative, for example to illustrate a concept or instruction, and/or used to indicate origin of the product. In some embodiments, the first compartment is the largest compartment having two large faces sealed around the perimeter, and the second compartment is smaller covering less than about 75%, or less than about 50% of the surface area of one face of the first compartment. In embodiments in which there is a third compartment, the aforementioned structure may be the same but the second and third compartments cover less than about 60%, or less than about 50%, or less than about 45% of the surface area of one face of the first compartment.

The pouches and/or packets of the present disclosure may comprise one or more different films. For example, in single compartment embodiments, the packet may be made from one wall that is folded onto itself and sealed at the edges, or alternatively, two walls that are sealed together at the edges. In multiple compartment embodiments, the packet may be made from one or more films such that any given packet compartment may comprise walls made from a single film or multiple films having differing compositions. In one aspect, a multi-compartment pouch comprises at least three walls: an outer upper wall; an outer lower wall; and a partitioning wall. The outer upper wall and the outer lower wall are generally opposing and form the exterior of the pouch. The partitioning wall is interior to the pouch and is secured to the generally opposing outer walls along a seal line. The partitioning wall separates the interior of the multi-compartment pouch into at least a first compartment and a second compartment. In one class of embodiments, the partitioning wall may be the only enzyme containing film thereby minimizing the exposure of the consumer to the enzymes.

Pouches and packets may be made using any suitable equipment and method. For example, single compartment pouches may be made using vertical form filling, horizontal form filling, or rotary drum filling techniques commonly known in the art. Such processes may be either continuous or intermittent. The film may be dampened, and/or heated to increase the malleability thereof. The method may also involve the use of a vacuum to draw the film into a suitable mold. The vacuum drawing the film into the mold can be applied for about 0.2 to about 5 seconds, or about 0.3 to about 3, or about 0.5 to about 1.5 seconds, once the film is on the horizontal portion of the surface. This vacuum can be such that it provides an under-pressure in a range of 10 mbar to 1000 mbar, or in a range of 100 mbar to 600 mbar, for example.

The molds, in which packets may be made, can have any shape, length, width and depth, depending on the required dimensions of the pouches. The molds may also vary in size

and shape from one to another, if desirable. For example, the volume of the final pouches may be about 5 ml to about 300 ml, or about 10 to 150 ml, or about 20 to about 100 ml, and that the mold sizes are adjusted accordingly.

5 In one aspect, the packet includes a first and a second sealed compartment. The second compartment is in a generally superposed relationship with the first sealed compartment such that the second sealed compartment and the first sealed compartment share a partitioning wall interior to the pouch.

10 In one aspect, the packet including a first and a second compartment further includes a third sealed compartment. The third sealed compartment is in a generally superposed relationship with the first sealed compartment such that the third sealed compartment and the first sealed compartment share a partitioning wall interior to the pouch.

In various aspects, the first composition and the second composition are selected from one of the following combinations: liquid, liquid; liquid, powder; powder, powder; and powder, liquid.

15 In various aspects, the first, second and third compositions are selected from one of the following combinations: solid, liquid, liquid and liquid, liquid, liquid.

In one aspect, the single compartment or plurality of sealed compartments contains a composition. The plurality of compartments may each contain the same or a different composition. The composition is selected from a liquid, solid or combination thereof.

20 Heat can be applied to the film in the process commonly known as thermoforming. The heat may be applied using any suitable means. For example, the film may be heated directly by passing it under a heating element or through hot air, prior to feeding it onto a surface or once on a surface. Alternatively, it may be heated indirectly, for example by heating the surface or applying a hot item onto the film. The film can be heated using an infrared light. The film may be
25 heated to a temperature of at least 50°C, for example about 50 to about 150°C, about 50 to about 120°C, about 60 to about 130°C, about 70 to about 120°C, or about 60 to about 90°C.

Alternatively, the film can be wetted by any suitable means, for example directly by spraying a wetting agent (including water, a solution of the film composition, a plasticizer for the film composition, or any combination of the foregoing) onto the film, prior to feeding it onto the
30 surface or once on the surface, or indirectly by wetting the surface or by applying a wet item onto the film.

Once a film has been heated and/or wetted, it may be drawn into an appropriate mold, preferably using a vacuum. The film can be thermoformed with a draw ratio of at least about 1.5, for example, and optionally up to a draw ratio of 2, for example. The filling of the molded film
35 can be accomplished by utilizing any suitable means. In some embodiments, the most preferred method will depend on the product form and required speed of filling. In some embodiments, the molded film is filled by in-line filling techniques. The filled, open packets are then closed forming the pouches, using a second film, by any suitable method. This may be accomplished while in

horizontal position and in continuous, constant motion. The closing may be accomplished by continuously feeding a second film, preferably water-soluble film, over and onto the open packets and then preferably sealing the first and second film together, typically in the area between the molds and thus between the packets.

5 Any suitable method of sealing the packet and/or the individual compartments thereof may be utilized. Non-limiting examples of such means include heat sealing, solvent welding, solvent or wet sealing, and combinations thereof. The water-soluble packet and/or the individual compartments thereof can be heat sealed at a temperature of at least 200°F (93°C), for example in a range of about 220°F (about 105°C) to about 290°F (about 145°C), or about 230°F
10 (about 110°C) to about 280°F (about 140°C). Typically, only the area which is to form the seal is treated with heat or solvent. The heat or solvent can be applied by any method, typically on the closing material, and typically only on the areas which are to form the seal. If solvent or wet sealing or welding is used, it may be preferred that heat is also applied. Preferred wet or solvent sealing/welding methods include selectively applying solvent onto the area between the molds,
15 or on the closing material, by for example, spraying or printing this onto these areas, and then applying pressure onto these areas, to form the seal. Sealing rolls and belts as described above (optionally also providing heat) can be used, for example.

The formed pouches may then be cut by a cutting device. Cutting can be accomplished using any known method. It may be preferred that the cutting is also done in continuous
20 manner, and preferably with constant speed and preferably while in horizontal position. The cutting device can, for example, be a sharp item, or a hot item, or a laser, whereby in the latter cases, the hot item or laser 'burns' through the film/ sealing area.

The different compartments of a multi-compartment pouches may be made together in a side-by-side style wherein the resulting, cojoined pouches may or may not be separated by
25 cutting. Alternatively, the compartments can be made separately.

In some embodiments, pouches may be made according to a process including the steps of:

- a) forming a first compartment (as described above);
- b) forming a recess within some or all of the closed compartment formed in step (a), to
30 generate a second molded compartment superposed above the first compartment;
- c) filling and closing the second compartments by means of a third film;
- d) sealing the first, second and third films; and
- e) cutting the films to produce a multi-compartment pouch.

The recess formed in step (b) may be achieved by applying a vacuum to the
35 compartment prepared in step (a).

In some embodiments, second, and/or third compartment(s) can be made in a separate step and then combined with the first compartment as described in EP 2088187 or WO 2009/152031.

In other embodiments, pouches may be made according to a process including the steps of:

- a) forming a first compartment, optionally using heat and/or vacuum, using a first film on a first forming machine;
- 5 b) filling the first compartment with a first composition;
- c) on a second forming machine, deforming a second film, optionally using heat and vacuum, to make a second and optionally third molded compartment;
- d) filling the second and optionally third compartments;
- e) sealing the second and optionally third compartment using a third film;
- 10 f) placing the sealed second and optionally third compartments onto the first compartment;
- g) sealing the first, second and optionally third compartments; and
- h) cutting the films to produce a multi-compartment pouch.

The first and second forming machines may be selected based on their suitability to perform the above process. In some embodiments, the first forming machine is preferably a horizontal forming machine, and the second forming machine is preferably a rotary drum forming machine, preferably located above the first forming machine.

It should be understood that by the use of appropriate feed stations, it may be possible to manufacture multi-compartment pouches incorporating a number of different or distinctive compositions and/or different or distinctive liquid, gel or paste compositions.

Processes of Making the Compositions

The compositions of the present invention can be formulated into any suitable form and prepared by any process chosen by the formulator, non-limiting examples of which are described in Applicants' examples and in US4990280; US20030087791A1; US20030087790A1; 25 US20050003983A1; US20040048764A1; US4762636; US6291412; US20050227891A1; EP1070115A2; US5879584; US5691297; US5574005; US5569645; US5565422; US5516448; US5489392; US5486303 all of which are incorporated herein by reference. The compositions of the invention or prepared according to the invention comprise cleaning and/or treatment composition including, but not limited to, compositions for treating fabrics, hard surfaces and 30 any other surfaces in the area of fabric and home care, including: air care including air fresheners and scent delivery systems, car care, dishwashing, fabric conditioning (including softening and/or freshening), laundry detergency, laundry and rinse additive and/or care, hard surface cleaning and/or treatment including floor and toilet bowl cleaners, granular or powder-form all-purpose or "heavy-duty" washing agents, especially cleaning detergents; liquid, gel or 35 paste-form all-purpose washing agents, especially the so-called heavy-duty liquid types; liquid fine-fabric detergents; hand dishwashing agents or light duty dishwashing agents, especially those of the high-foaming type; machine dishwashing agents, including the various tablet, granular, liquid and rinse-aid types for household and institutional use: car or carpet shampoos,

bathroom cleaners including toilet bowl cleaners; as well as cleaning auxiliaries such as bleach additives and "stain-stick" or pre-treat types, substrate-laden compositions such as dryer added sheets. Preferred are compositions and methods for cleaning and/or treating textiles and/or hard surfaces, most preferably textiles. The compositions are preferably compositions used in a pre-treatment step or main wash step of a washing process, most preferably for use in textile washing step.

As used herein, the term "fabric and/or hard surface cleaning and/or treatment composition" is a subset of cleaning and treatment compositions that includes, unless otherwise indicated, granular or powder-form all-purpose or "heavy-duty" washing agents, especially cleaning detergents; liquid, gel or paste-form all-purpose washing agents, especially the so-called heavy-duty liquid types; liquid fine-fabric detergents; hand dishwashing agents or light duty dishwashing agents, especially those of the high-foaming type; machine dishwashing agents, including the various tablet, granular, liquid and rinse-aid types for household and institutional use; liquid cleaning and disinfecting agents, car or carpet shampoos, bathroom cleaners including toilet bowl cleaners; fabric conditioning compositions including softening and/or freshening that may be in liquid, solid and/or dryer sheet form; as well as cleaning auxiliaries such as bleach additives and "stain-stick" or pre-treat types, substrate-laden compositions such as dryer added sheets. All of such compositions which are applicable may be in standard, concentrated or even highly concentrated form even to the extent that such compositions may in certain aspect be non-aqueous.

Method of Use

The present invention includes a method for cleaning any surface including treating a textile or a hard surface or other surfaces in the field of fabric and/or home care. It is contemplated that cleaning as described may be both in small scale as in e.g. family house hold as well as in large scale as in e.g. industrial and professional settings. In one aspect of the invention, the method comprises the step of contacting the surface to be treated in a pre-treatment step or main wash step of a washing process, most preferably for use in a textile washing step or alternatively for use in dishwashing including both manual as well as automated/mechanical dishwashing. In one aspect of the invention the lipase variant and other components are added sequentially into the method for cleaning and/or treating the surface. Alternatively, the lipase variant and other components are added simultaneously.

As used herein, washing includes but is not limited to, scrubbing, and mechanical agitation. Washing may be conducted with a foam composition as described in WO08/101958 and/or by applying alternating pressure (pressure/vacuum) as an addition or as an alternative to scrubbing and mechanical agitation. Drying of such surfaces or fabrics may be accomplished by any one of the common means employed either in domestic or industrial settings. The cleaning compositions of the present invention are ideally suited for use in laundry as well as

dishwashing applications. Accordingly, the present invention includes a method for cleaning an object including but not limiting to fabric, tableware, cutlery and kitchenware. The method comprises the steps of contacting the object to be cleaned with a said cleaning composition comprising at least one aspect of Applicants' cleaning composition, cleaning additive or mixture thereof. The fabric may comprise most any fabric capable of being laundered in normal consumer or institutional use conditions. The solution may have a pH from 8 to 10.5. The compositions may be employed at concentrations from 500 to 15.000ppm in solution. The water temperatures typically range from 5°C to 90°C. The water to fabric ratio is typically from 1:1 to 30:1.

In one aspect the invention relates to a method of using a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase.

In one aspect the invention relates to use of the composition comprising a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase for cleaning an object.

In one aspect the parent lipase is a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90 % identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase. In one aspect the parent lipase is a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2. In one aspect the parent lipase has the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. In one aspect the parent lipase comprises or consists the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

In one aspect the invention relates to a method of producing the composition, comprising adding a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase

activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase, and a surfactant. In one aspect the invention relates to a method for cleaning a surface, comprising contacting a lipid stain present on the surface to be cleaned with the cleaning composition. In one aspect the invention relates to a method for hydrolyzing a lipid present in a soil and/or a stain on a surface, comprising contacting the soil and/or the stain with the cleaning composition. In one aspect the invention relates to use of the composition in the hydrolysis of a carboxylic acid ester. In one aspect the invention relates to use of the composition in the hydrolysis, synthesis or interesterification of an ester. In one aspect the invention relates to use of the composition for the manufacture of a stable formulation.

In one aspect the invention relates to use of a variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase. In one aspect the parent lipase is consisting or comprising SEQ ID NO: 2; SEQ IN NO: 4; SEQ ID NO: 6; or any fragment thereof having lipase activity. In one aspect the variant according to the invention is used for hydrolysis, synthesis or interesterification of an ester; hydrolysis of a carboxylic acid ester; hydrolyzing a lipid, cleaning a surface; and/or producing a detergent composition.

EXAMPLES

Example 1: Assays

p-nitrophenyl (pNP) assay:

The hydrolytic activity of a lipase may be determined by a kinetic assay using p-nitrophenyl acyl esters as substrate.

A 100mM stock solution in DMSO of the substrates: p-Nitrophenyl butyrate (C4), p-Nitrophenyl caproate (C6), p-Nitrophenyl caprate (C10), p-Nitrophenyl laurate (C12) and p-Nitrophenyl palmitate (C16) (all from Sigma-Aldrich Danmark A/S, Kirkebjerg Allé 84, 2605 Brøndby; Cat.no.: C4:N-9876, C6: N-0502, C10: N-0252, C12: N-2002, C16: N-2752) may be diluted to a final concentration of 1mM into assay buffer (50mM Tris; pH 7.7; 0.4% TritonX-100).

The lipase of the invention, the parent lipase and appropriate controls e.g. Buffer (negative), LipolaseTM & LipexTM (positive) in 50mM Hepes; pH 8.0; 10ppm TritonX-100; +/- 20mM CaCl₂ may be added to the substrate solution in the following final concentrations: 0.01 mg/ml; 5x10⁻³ mg/ml; 2.5x10⁻⁴ mg/ml; and 1.25x10⁻⁴ mg/ml in 96-well NUNC plates (Cat. No:260836, Kamstrupvej 90, DK-4000, Roskilde). Release of p-nitrophenol by hydrolysis of p-nitrophenyl acyl may be monitored at 405nm for 5 minutes in 10 second intervals on a Spectra max 190 (Molecular Devices GmbH, Bismarckring 39, 88400 Biberach an der Riss, GERMANY). The hydrolytic activity towards one or more substrates of a variant may be compared to that of the parent lipase.

Differential Scanning Calorimetry (DSC) Assay:

The thermostability of the lipase variants were determined by Differential Scanning Calorimetry (DSC) using a VP-Capillary Differential Scanning Calorimeter (MicroCal Inc., Piscataway, NJ, USA). The thermal denaturation temperature, T_d ($^{\circ}\text{C}$), was taken as the top of denaturation peak (major endothermic peak) in thermograms (C_p vs. T) obtained after heating enzyme solutions (approx. 0.5mg/mL) in buffer (50mM Hepes pH 8) at a constant programmed heating rate of 200 K/hr.

Sample- and reference-solutions (approx. 0.2mL) were loaded into the calorimeter (reference: buffer without enzyme) from storage conditions at 10°C and thermally pre-equilibrated for 20 minutes at 20°C prior to DSC scan from 20°C to 100°C . Denaturation temperatures were determined at an accuracy of approximately $\pm 1^{\circ}\text{C}$.

DSC were also conducted by adding 0.05g/L Relase (Novozymes A/S) and/or 1mM LAS.

Storage Stability Assays:

Protocol A: *Aspergillus oryzae* strains producing the lipase variant were grown for 5 days at 37°C in 2xSC medium with 2% maltose without shaking. 2xSC medium is 15g yeast nitrogen base without amino acids (Difco 291920), 22.6g succinic acid (Merck 822260), 13.6g +sodium hydroxide (Merck 106498), 11.2g casamino acids (vitamin assay, Difco 228830) and 0.2g L-tryptophan (Merck 108374) dissolved in 1L deionized water.

10uL of the culture broth were added to 90uL of detergent composition, stirred for 10 minutes and sealed in small plastic containers. For stress conditions C1 samples with detergent composition D0001 were stored at -20°C in detergent D001 with 0.02% calcium chloride (unstressed) and in detergent D001 with 0.02% calcium chloride and 1.35% Relase 16L EXI (Novozymes) at 48°C (stressed). For stress conditions C2 samples with detergent composition D0001 were stored at -20°C in detergent D001 with 0.02% calcium chloride (unstressed) and in detergent D001 with 0.02% calcium chloride and 1.35% Savinase Ultra 16L (Novozymes) at 35°C (stressed). For stress conditions C3 samples with detergent composition D0002 samples were stored at -20°C in detergent D002 with 0.02% calcium chloride (unstressed) and in detergent D002 with 0.02% calcium chloride and 1.35% Relase 16L EXI (Novozymes) at 55°C (stressed). For stress conditions C4 samples with detergent composition D0002 samples were stored at -20°C in detergent D002 with 0.02% calcium chloride (unstressed) and in detergent D002 with 0.02% calcium chloride and 1.35% Savinase 16L (Novozymes) at 35°C (stressed). Storage time was 19 hours.

After storage possible condensation liquid was collected by centrifugation. To the 100uL stressed or unstressed sample 230uL of buffer (0.1M Tris-HCl; 9mM CaCl_2 ; 0.0225% Brij-30; pH8.0 +0.85% 4-FBPA (31.5g/l)) were added corresponding to a 3.3-fold dilution. After 10 minutes stirring 5uL sample aliquots were further diluted with the same buffer 60-fold. Then one

part of this lipase dilution was mixed with four parts of 0.5mM pNP-palmitate, 1mM calcium chloride, 100mM Tris (pH8.0), 6.5mM Deoxycholate, 1.4g/L AOS and for 30 minutes release of the pNP chromophore was measured spectrophotometrically. This was used to determine activity via the initial linear slope of the reaction.

- 5 Residual activity (RA) was calculated as the ratio of the measured velocities of stressed versus unstressed sample. Half life ($T_{1/2}$) was calculated based on the following formula:

$$\text{Half life} = \text{Stress time} * \ln(0,5)/\ln(\text{residual activity})$$

The median value of the residual activity and the half-lives was calculated based on two to four replicates.

- 10 The half life improvement factor (HIF) of the specific mutations was calculated by dividing the half life of the lipase variant with the half life of the parent lipase with the sequence of SEQ ID NO: 2.

Protocol B (purified enzyme in detergent with protease): After active site titration purified lipase variants were diluted with a buffer (10mM Succinic acid + 2mM CaCl₂ + 0.02% Brij 35 adjusted to pH6.5) to a the specified concentration. 10uL of the 100ppm lipase solution was added to a 90uL of detergent composition, stirred for 10 minutes and sealed. For stress conditions C1 samples with detergent composition D0001 were stored at -20°C in detergent D001 with 0.02% calcium chloride (unstressed) and in detergent D001 with 0.02% calcium chloride and 1.35% Relase 16L EXI (Novozymes) at 48°C (stressed). For stress conditions C2 samples with detergent composition D0001 were stored at -20°C in detergent D001 with 0.02% calcium chloride (unstressed) and in detergent D001 with 0.02% calcium chloride and 1.35% Savinase Ultra 16L (Novozymes) at 35°C (stressed). For stress conditions C3 samples with detergent composition D0002 samples were stored at -20°C in detergent D002 with 0.02% calcium chloride (unstressed) and in detergent D002 with 0.02% calcium chloride and 1.35% Relase 16L EXI (Novozymes) at 55°C (stressed). For stress conditions C4 samples with detergent composition D0002 samples were stored at -20°C in detergent D002 with 0.02% calcium chloride (unstressed) and in detergent D002 with 0.02% calcium chloride and 1.35% Savinase 16L (Novozymes) at 35°C (stressed). Storage time was 19 hours.

After storage possible condensation liquid was collected by centrifugation. To the 100uL stressed or unstressed sample 230uL of buffer (0.1M Tris-HCl; 9mM CaCl₂; 0.0225% Brij-30; pH8.0 + 0.85% 4-FBPA (31.5g/l)) were added corresponding to a 3.3-fold dilution. After 10 minutes stirring 5uL sample aliquots were further diluted with the same buffer 60-fold. Then one part of this lipase dilution was mixed with four parts of 0.5mM pNP-palmitate, 1mM calcium chloride, 100mM Tris (pH8.0), 6.5mM Deoxycholate, 1.4g/L AOS and for 30 minutes release of the pNP chromophore was measured spectrophotometrically. This was used to determine activity via the initial linear slope of the reaction.

Residual activity (RA) was calculated as the ratio of the measured velocities of stressed versus unstressed sample. Half life ($T_{1/2}$) was calculated based on the following formula:

$$\text{Half life} = \text{Stress time} * \ln(0,5)/\ln(\text{residual activity})$$

The median value of the residual activity and the half-life was calculated based on four replicates. The median value of the residual activity and the half-life was calculated based on three replicates. The half-life improvement factor (HIF) was calculated between a pair of variants being different in a single or a double mutation. The half-life improvement factor is the half-life of the variant with the extra single or double mutation divided by the half-life of the variant without.

Detergents: Composition D001 is a commercially available AVA liquid detergent from Reckitt Benckiser. Composition D002 is a model detergent as listed below.

Composition of D002 (Content of active component in ingredient)	wt%
Soft water	33.64
NaOH, pellets (>99%)	1.75
Linear alkylbenzenesulfonic acid (LAS) (97%)	12.00
Sodium laureth sulfate (SLES) (28%)	17.63
Soy fatty acid (>90%)	2.75
Coco fatty acid (>90%)	2.75
AEO; alcohol ethoxylate with 8 mol EO; Lutensol TO 8 (~100%)	11.00
Triethanol amine (100%)	3.33
Na-citrate, dihydrate (100%)	2.00
DTMPA; diethylenetriaminepentakis(methylene)pentakis(phosphonic acid), heptasodium salt (Dequest 2066 C) (~42% as Na ₇ salt)	0.48
MPG (>98%)	6.00
EtOH, propan-2-ol (90/10%)	3.00
Glycerol (>99.5)	1.71
Sodium formate (>95%)	1.00
PCA (40% as sodium salt)	0.46
Phenoxyethanol	0.50

10 Relative Wash Performance, RP(Wash)

Washing experiments were performed using Automatic Mechanical Stress Assay (AMSA) in order to assess the wash performance in laundry. The AMSA plate has a number of slots for test solutions and a lid firmly squeezing the laundry sample, the textile to be washed against all the slot openings. During the washing time, the plate, test solutions, textile and lid are vigorously shaken to bring the test solution in contact with the textile and apply mechanical stress in a regular, periodic oscillating manner. For further description see WO02/42740 especially the paragraph "Special method embodiments" at page 23-24.

The laundry experiments were conducted using the following experimental conditions:

Detergents:	3.3 g/L Detergent B; or 1.75 g/L Detergent X
Test solution volume:	160 μ L
Wash time:	20 minutes
Temperature:	30°C
Lipase dosage:	0 ppm; or 0.35 ppm
Test material:	Cream annatto stained EMPA221 cotton textile (Detergent B wash); or Cream turmeric stained EMPA221 cotton textile (Detergent X wash)

5 Cream turmeric staining of EMPA221 textile was done according to WO06/125437. The cream annatto stained EMPA221 was produced the same way as cream turmeric stained EMPA221 except for exchanging turmeric with annatto (A-320-WS obtained from Chr. Hansen Natural Colors A/S Boege Allé 10-12, 2970 Hoersholm, Denmark). EMPA221 was obtained from EMPA, Lerchenfeldstrasse 5, CH-9014, St. Gallen, Switzerland.

Water hardness was adjusted to 5°dH or 15°dH by addition of CaCl_2 , MgCl_2 and NaHCO_3 ($\text{Ca}^{2+}:\text{Mg}^{2+}:\text{HCO}_3^- = 4:1:7.5$), when wash with Detergent B and 12°dH ($\text{Ca}^{2+}:\text{Mg}^{2+}:\text{HCO}_3^- = 2:1:4.5$), when wash with Detergent X.

Detergent B composition	Wt%
NaOH, pellets (>99%)	1.05
Linear alkylbenzenesulfonic acid (LAS) (97%)	7.20
Sodium laureth sulfate (SLES) (28%)	10.58
Soy fatty acid (>90%)	2.75
Coco fatty acid (>99%)	2.75
Alcohol ethoxylate (AEO) with 8 mol EO; Lutensol TO 8 (~100%)	6.60
Triethanol amine (100%)	3.33
Na-citrate, dihydrate (100%)	2.00
DTMPA; diethylenetriaminepentakis(methylene)pentakis(phosphonic acid), heptasodium salt (Dequest 2066 C) (~42% as Na_7 salt)	0.48
MPG (>98%)	6.00
EtOH, propan-2-ol (90/10%)	3.00
Glycerol (>99.5)	1.71
Sodium formate (>95%)	1.00
PCA (40% as sodium salt)	0.46
Water up to	100

Detergent X composition	Wt%
Linear alkylbenzenesulfonic acid (LAS) (85%)	17.6

AEO;alcohol ethoxylate with 7 mol EO (99.5%)	2.0
sodium carbonate (99.5%)	20.1
sodium (di)silicate (80%)	12.4
zeolite A (80%)	15.0
sodium sulfate (100%)	31.8
copoly(acrylic acid/maleic acid),sodium salt (92%)	1.1

Detergent B is an example of a liquid detergent composition whereas Detergent X is an example of a powder detergent composition. After washing the textiles were flushed in tap water and excess water was removed from the textiles using filter paper and immediately thereafter the textiles were dried at 100°C for 15 min.

5 The wash performance was measured as the color change of the washed soiled textile. The soil was cream mixed with annatto when washing with Detergent B and cream mixed with turmeric when washing with Detergent X. Annatto contains the colorant norbixin and turmeric contains the colorant curcumin. Norbixin as well as curcumin functions as pH indicators by having pH dependent color change. Lipase activity leads to release of free fatty acids from the cream acylglycerides and this leads to pH decrease and thereby color change of the norbixin or
10 curcumin pH indicators. Lipase wash performance can therefore be expressed as the extent of color change of light reflected-emitted from the washed soiled textile when illuminated with white light.

Color measurements were made with a professional flatbed scanner (EPSON
15 EXPRESSION 11000XL, Atea A/S, Lautrupvang 6, 2750 Ballerup, Denmark), which was used to capture an image of the washed soiled textile. To extract a value for the light intensity from the scanned images, 24-bit pixel values from the image were converted into values for red, green and blue (RGB).

Color change due to lipase activity is measured as the change in the reflection-emitting
20 of green light (G) relative to the light intensity value (Int) calculated by adding the RGB values together as vectors and then taking the length of the resulting vector:

$$\text{Int} = \sqrt{R^2 + G^2 + B^2}$$

The relative wash performance (RP(Wash)) of a lipase relative to a reference lipase is calculated as:

25 $\text{RP(Wash)} = (\text{G/Int(tested lipase)} - \text{G/Int(no enzyme)}) / (\text{G/Int(lipase ref.)} - \text{G/Int(no enzyme)})$.

A lipase is considered to exhibit improved wash performance, if it performs better than the reference ($\text{RP(Wash)} > 1$). In the context of the present invention the reference enzyme is the wildtype lipase or the parent lipase i.e. the control lipase without the substitutions E1C R233C.

Nano Differential Scanning Fluorescence (NanoDSF) Assay

Thermal stability measurements were performed using a nano differential scanning fluorescence instrument (nanoDSF); Prometheus NT.48 (NanoTemper Technologies GmbH, München, Germany). Standard nanoDSF grade capillaries were used (Cat#: PR-C002 NanoTemper Technologies). The purified protein samples were loaded into the capillaries (each sample in triplicate) by capillary action. The emission intensities at 330 and 350 nm were optimized by altering the LED power on the instrument so as to obtain a signal between 3000 and 15000 fluorescence counts. The temperature slope used for thermal unfolding was 3.3 degrees Celsius per minute from 20 to 95 degrees Celsius. The data was analysed using the software PR.Control v1.11.2 supplied by the manufacturer. Typically, the positive (or negative) peak maximum (or minimum) in the first derivative analysis was taken to represent the thermal denaturation temperature, T_d (°C).

Samples were used directly from the purified stocks. To determine the effect of reducing agent on the thermal stabilities, 250mM TCEP in 50mM HEPES, pH 8.0 was added to a final concentration of 0.5 mM to the samples. The samples under reducing and non-reducing conditions were run in parallel.

Example 2: Stability

The stability of a variant according to the invention, the reference lipase and a prior art lipase were determined according to the DSC Assay described in Example 1. The thermal denaturing temperature for each lipase under various conditions is shown in the table below.

Table 2: Thermal denaturing temperature, T_d

Lipase	Substitutions in SEQ ID NO: 2	T _d
Wild type	-	74.5
Prior art lipase	SPPCGRRP +E1* +E239C	72.4
Variant 1	E1C +N233C	75.1
Variant 2	E1C +T231R +N233C	75.4
Variant 3	E1C +L227G +T231R +N233C	78.0

The lipase variants of the invention show increased stability over the prior art lipase at the conditions tested. Stability was also investigated in the presence of 0.05g/L Relase (Novozymes A/S) and/or 1mM LAS. Introduction of the substitutions E1C+N233C resulted in an increased stability in the presence of Relase and/or LAS.

Example 3: Stability

The stability of a variant according to the invention, the reference lipase and a prior art lipase were determined according to the DSC Assay described in Example 1. The thermal denaturing temperature for each lipase under various conditions is shown in the table below.

Table 3A:

Lipase	Substitutions in SEQ ID NO: 2	Td
Wildtype	-	74.5
Variant 7	E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233C D254S P256T	80.6

Table 3B:

Lipase	Substitutions in SEQ ID NO: 2	Td
Control	T231R N233R	74.2
Variant 2	E1C T231R N233C	75.4
Variant 3	E1C L227G T231R N233C	78.1

5

Table 3C:

Lipase	Substitutions in SEQ ID NO: 2	Td
Control	T231R	74.4
Control 8	V2K D27R N33K G38A F51V D96E D111A G163K T231R D254S P256T	75.4
Variant 8	E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T	78.1
Control 9	D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R D254S P256T	81.8
Variant 9	E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T	84.8
Control 4	D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R D254S P256T	78.8
Variant 4	E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T	82.4

Example 4: Storage stability

10 Variants comprising E1C N233C substitutions were tested according to protocol A as described in Example 1. Comparison to a corresponding lipase without E1C N233C substitutions were conducted to show the effect of the double substitution.

Table 4: Improved stability

Stress	HIF	Res Act (%)	Half life (Hour)	Mutations of compared variants
C2	2.2	A: 15 B: 42	A: 7 B: 15	A: T231R N233R B: E1C T231R N233C
C1	>1.2	A: <5 B: 8	A: <4 B: 5	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T A257I
C2	>35.1	A: <5 B: 92	A: <4 B: 154	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V S54T E56K D96E K98I

				D111A G163K T231R N233C D254S P256T A257I
C4	3.3	A: 61 B: 86	A: 26 B: 87	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T A257I
C1	>1.6	A: <5 B: 16	A: <4 B: 7	A: D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I
C2	>36.0	A: <5 B: 92	A: <4 B: 158	A: D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I
C4	2.1	A: 63 B: 80	A: 28 B: 60	A: D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K T231R N233R D254S P256T A257I B: E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L T231R N233C D254S P256T A257I
C1	>11.0	A: <5 B: 76	A: <4 B: 48	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C2	21.3	A: 76 B: 99	A: 48 B: 1024	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C3	>3.3	A: <5 B: 40	A: <4 B: 15	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C4	1.4	A: 87 B: 91	A: 96 B: 138	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C1	>15.6	A: <5 B: 82	A: <4 B: 68	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T
C2	6.9	A: 79 B: 97	A: 56 B: 387	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T
C3	>3.4	A: <5 B: 42	A: <4 B: 15	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T
C4	2.2	A: 82 B: 92	A: 67 B: 150	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S

				P256T
C1	>3.3	A: <5 B: 40	A: <4 B: 14	A: D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233C D254S P256T
C3	>2.2	A: <5 B: 25	A: <4 B: 9	A: D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K Y220F T231R N233C D254S P256T
C1	1.8	A: 55 B: 71	A: 22 B: 39	A: D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233R D254S P256T B: E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T
C3	1.4	A: 46 B: 58	A: 17 B: 24	A: D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233R D254S P256T B: E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T
C1	1.6	A: 56 B: 69	A: 23 B: 35	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T
C3	1.2	A: 43 B: 50	A: 16 B: 19	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T

Example 5: Storage stability

5 Variants comprising E1C N233C substitutions were tested according to protocol B as described in Example 1. Comparison to a corresponding lipase without E1C N233C substitutions were conducted to show the effect of the double substitution.

Table 5: Improved stability

Stress	HIF	Res Act (%)	Half life (Hour)	Mutations of compared variants
C2	1.5	A: 12 B: 25	A: 6 B: 9	A: - B: E1C N233C
C4	1.4	A: 16 B: 27	A: 7 B: 10	A: B: E1C N233C
C2	1.3	A: 5 B: 10	A: 4 B: 6	A: T231R N233R B: E1C T231R N233C
C4	1.2	A: 22 B: 28	A: 9 B: 10	A: T231R N233R B: E1C T231R N233C
C1	4.0	A: 23 B: 69	A: 9 B: 36	A: D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233R D254S P256T B: E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T
C2	8.9	A: 46 B: 92	A: 17 B: 151	A: D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233R D254S P256T B: E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T

C3	1.4	A: 50 B: 60	A: 19 B: 26	A: D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233R D254S P256T B: E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T
C1	>2.1	A: <5 B: 24	A: <4 B: 9	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C3	>3.2	A: <5 B: 39	A: <4 B: 14	A: D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T
C1	1.9	A: 62 B: 78	A: 27 B: 53	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T
C3	1.5	A: 44 B: 57	A: 16 B: 23	A: D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233R D254S P256T B: E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R N233C D254S P256T
C1	>8.0	A: <5 B: 69	A: <4 B: 35	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T
C2	2.6	A: 58 B: 81	A: 24 h B: 62 h	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T
C3	>3.2	A: <5 B: 39	A: <4 h B: 14 h	A: D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233R D254S P256T B: E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S E210K Y220F T231R N233C D254S P256T

Example 6: Wash performance

The relative wash performance, RP(wash), of a prior art lipase and variants according to the invention was studied with the method described in example 1 using Detergent B and compared to RP(wash) of the wildtype lipase.

Table 6: Improved wash performance over wildtype and prior art lipase.

Lipase	Substitutions in SEQ ID NO: 2	RP(wash)
Wildtype	-	1.00
Prior Art	SPPCGRRP +E1* +E239C	0.57
Variant 1	E1C +N233C	1.50
Variant 2	E1C +T231R +N233C	1.72
Variant 3	E1C +L227G +T231R +N233C	1.85

Example 7: Wash performance

The relative wash performance, RP(wash), of variants according to the invention was

studied with the method described in example 1 using a liquid detergent, Detergent B (Det B) or a powder detergent, Detergent X (Det X) and compared to RP(wash) of the corresponding parent lipase i.e. the control lipase without the substitutions E1C+N233C.

Table 7: Improved wash performance over parent lipase.

Lipase	Substitutions in SEQ ID NO: 2	RP(wash) (Det. B 15°dH)	RP(wash) (Det. B 5°dH)	RP(wash) (Det X 12°dH)
Wildtype	-	1.00	-	-
Variant 1	E1C N233C	1.50	-	-
Control 4	D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T	1.00	1.00	1.00
Variant 4	E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T	1.03	3.37	1.88
Control 5	V2K D27R N33K G38A F51V D96E D111A G163K T231R N233R D254S P256T A257I	1.00	1.00	1.00
Variant 5	E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T A257I	1.61	2.12	1.35

5

Example 8: Effect of a reducing agent on stability

The stability of a variant according to the invention, the parent lipase i.e. the control lipase without the substitutions E1C+N233C and a prior art lipase were determined according to the NanoDSF Assay described in Example 1. The thermal denaturing temperature for each lipase in the absence and presence of a reducing agent (TCEP) is shown in the table below.

10

Table 8: Improved stability in the presence and absence of a reducing agent

Lipase	Substitutions in SEQ ID NO: 2	Td (°C)	stdev	Td TCEP (°C)	stdev
Wildtype	-	69.2	1.7	65.3	1.2
Prior art	SPPCGRRP +E1* +E239C	69.7	0.3	60.6	0.8
Wildtype	-	69.2	1.7	65.3	1.2
Variant 1	E1C N233C	72.7	0.7	64.0	0.1
Control 4	D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233R D254S P256T	71.5	0.2	67.4	0.3
Variant 4	E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R N233C D254S P256T	81.0	0.7	71.1	0.9
Control 5	V2K D27R N33K G38A F51V D96E D111A G163K T231R N233R D254S P256T A257I	63.5	0.1	58.5	0.1
Variant 5	E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S P256T A257I	66.2	0.2	60.3	0.1
Control 6	D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R D254S P256T	77.2	0.2	70.5	0.4
Variant 6	E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C D254S P256T	78.6	0.2	71.3	0.2

The invention described and claimed herein is not to be limited in scope by the specific aspects herein disclosed, since these aspects are intended as illustrations of several aspects of the invention. Any equivalent aspects are intended to be within the scope of this invention.

- 5 Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

CLAIMS

1. A variant of a parent lipase, which variant comprises substitutions at the positions corresponding to E1C and N233C of the mature polypeptide of SEQ ID NO: 2, has lipase activity, and has at least 60% but less than 100% sequence identity to the mature polypeptide of the parent lipase.
2. The variant of claim 1, wherein the parent lipase is selected from:
 - a. a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90% identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15 Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) meets the following limitations: (i) comprises a negative amino acid in position E210 of said wild-type lipase; (ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and (iii) comprises a neutral or negative amino acid at a position corresponding to N94 of said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase.
 - b. a lipase which has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 2, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T, D27R and G38A of SEQ ID NO: 2.
 - c. a lipase which has the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6; or comprises or consists the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.
3. The variant of any of claims 1-2 selected from the group consisting of:
 - a. a polypeptide having at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, but less than 100% sequence identity to the mature polypeptide of SEQ ID NO: 2;
 - b. a polypeptide encoded by a polynucleotide that hybridizes under low stringency conditions, medium stringency conditions, medium-high stringency conditions, high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, or (ii) the full-length complement of (i);
 - c. a polypeptide encoded by a polynucleotide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at

least 97%, at least 98%, or at least 99%, or 100% identity to the mature polypeptide coding sequence of SEQ ID NO: 1; and

d. a fragment of the mature polypeptide of SEQ ID NO: 2, which has lipase activity.

5 4. The variant of any of claims 1-3, which further comprises one or more (e.g. several) substitutions at positions corresponding to positions: 2, 4, 8, 11, 15, 27, 33, 38, 43, 48, 51, 54, 56, 57, 58, 60, 69, 71, 83, 86, 91, 92, 94, 96, 97, 98, 99, 101, 111, 123, 150, 152, 163, 176, 179, 187, 188, 189, 198, 199, 200, 210, 216, 220, 224, 225, 227, 228, 229, 231, 236, 238, 239, 246, 249, 254, 255, 256, 257, 260, 263, 264, 265, 266, 267, 269 of SEQ ID NO: 2.

10 5. The variant of the previous claim wherein the substitution is corresponding to V2K, Q4R, Q4V, N8R, N11R, Q15C, D27G, D27R, N33K, N33Q, G38A, E43C, D48C, F51V, S54T, E56K, D57G, S58A, V60K, V60S, L69R, N71C, S83T, I86V, G91A, G91N, G91Q, N92D, N94K, N94R, D96E, D96G, D96L, D96W, L97M, K98E, K98I, K98Q, E99K, E99N, N101D, N101S, D111A,
15 T123V, A150G, A152G, G163K, V176L, R179L, V187Y, V187W, Q188R, T189Y, T189W, H198S, T199R, N200R, E210K, E210Q, S216P, Y220F, S224R, G225R, L227G, L227R, V228R, P229R, T231R, V236R, I238C, E239C, G246C, Q249R, D254S, I255G, P256K, P256T, P256V, A257I, A257V, W260C, G263Q, L264A, I265T, G266D, T267A, L269N, L269V of SEQ ID NO: 2.

20 6. The variant of any of claims 1-5, comprising or consisting of a set of substitutions at positions corresponding to SEQ ID NO: 2 selected from:

- a. E1C N233C;
- 25 b. E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K N233C D254S P256T;
- c. E1C V2K D27G N33K G38A F51V D96E D111A G163K N233C D254S P256T;
- d. E1C V2K D27R N33K G38A F51V D96E D111A G163K Q188R N233C D254S P256T;
- 30 e. E1C D27R G38A G91A N92D D96L K98Q D111A G163K N233C D254S P256T;
- f. E1C D27R G38A G91N N94R D96E D111A G163K S216P L227G N233C D254S P256T;
- g. E1C T231R N233C;
- h. E1C T231R N233C Q249R D254S;
- 35 i. E1C G225R T231R N233C;
- j. E1C Q15C E43C T231R N233C;
- k. E1C L227R T231R N233C;
- l. E1C P229R T231R N233C;

- m. E1C L227G T231R N233C;
- n. E1C E99N N101S T231R N233C;
- o. E1C L227G T231R N233C D254S;
- p. E1C E210K L227G T231R N233C;
- 5 q. E1C D27R N33K G38A F51V D96E K98E N101D D111A G163K H198S E210K
Y220F T231R N233C D254S P256T;
- r. E1C D27R N33K G38A F51V S54T E56K D57G L69R D96E K98I D111A A152G
G163K T231R N233C D254S P256T;
- s. E1C V187Y T189Y L227G T231R N233C;
- 10 t. E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S
E210K Y220F T231R N233C D254S P256T;
- u. E1C V60K I86V A150G E210K L227G T231R N233C P256K;
- v. E1C V187W T189W L227G T231R N233C;
- w. E1C N94K D96L L227G T231R N233C;
- 15 x. E1C G91A N92D D96L K98Q L227G T231R N233C;
- y. E1C N8R L227G T231R N233C;
- z. E1C L227G V228R T231R N233C;
- aa. E1C Q4R L227G T231R N233C;
- bb. E1C N11R L227G T231R N233C;
- 20 cc. E1C S224R L227G T231R N233C;
- dd. E1C L227G T231R N233C V236R;
- ee. E1C N200R L227G T231R N233C;
- ff. E1C T199R L227G T231R N233C;
- gg. E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S
25 P256T;
- hh. E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R
N233C D254S P256T;
- ii. E1C D27R N33K G38A F51V D96E K98I D111A G163K H198S Y220F T231R
N233C D254S P256T;
- 30 jj. E1C D27R N33K G38A F51V E56K L69R D96E K98E D111A G163K R179L
T231R N233C D254S P256T A257I;
- kk. E1C V2K D27R N33K G38A F51V D96E D111A G163K T231R N233C D254S
P256T A257I;
- ll. E1C D27R N33K G38A F51V S54T E56K D96E K98I D111A G163K T231R
35 N233C D254S P256T A257I;
- mm. E1C D27R N33K G38A F51V S54T E56K D57G D96E K98I D111A G163K

T231R N233C D254S I255G P256T A257V L269V;

nn. E1C V2K D27R N33K G38A F51V L69R D96E K98E D111A G163K V176L
E210K L227G T231R N233C D254S P256T;

oo. E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S
5 E210K Y220F T231R N233C D254S P256T;

pp. E1C D27R N33K G38A F51V D96E K98E N101D D111A T123V G163K H198S
E210K Y220F T231R N233C D254S P256T;

qq. E1C D27R G38A F51V L69R D96E K98E D111A G163K E210K T231R N233C
D254S P256T; and

10 rr. E1C N11R D27R N33K D48C F51V L69R N71C E87Q K98E N101R T143A
E210K G225R L227G P229R T231R N233C Q249R P250R D254S I255G P256K.

7. The variant of any of claims 1-6, which has an improved stability relative to the parent
lipase.

8. The variant of claim 7, wherein the stability is selected from the group consisting of
thermostability, stability in the presence of proteolytic enzymes, stability in the presence of
surfactant, stability in detergent compositions, stability under storage conditions.

9. A composition comprising the variant of any of claims 1-8.

10. A method for manufacturing a composition comprising a step of adding the variant of any
of claims 1-8 to the composition.

11. A polynucleotide encoding the variant of any of claims 1-8.

12. A nucleic acid construct comprising the polynucleotide of claim 11.

13. An expression vector comprising the polynucleotide of claim 11.

14. A host cell comprising the polynucleotide of claim 11, the nucleic acid construct of claim
12, or the expression vector of claim 13.

15. A method of producing a lipase variant, comprising: (a) cultivating the host cell of claim
14 under conditions suitable for expression of the variant; and (b) recovering the variant.

16. A method for obtaining a lipase variant, comprising introducing into a parent lipase a
substitution at the positions corresponding to E1C and N233C of SEQ ID NO: 2, wherein the
variant has lipase activity; and recovering the variant.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2015/080463

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed:
 - ☒ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13~~ter~~.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13~~ter~~.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13~~ter~~.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2015/080463

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N9/20 C11D3/386
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBL, FSTA, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6 624 129 B1 (BORCH KIM [DK] ET AL) 23 September 2003 (2003-09-23) column 3, lines 1-15 -----	1-16
Y	WO 00/60063 A1 (NOVO NORDISK AS [DK]; VIND JESPER [DK]; SVENDSEN ALLAN [DK]; PATKAR SH) 12 October 2000 (2000-10-12) page 3, lines 10-12 -----	1-16
Y	WO 2014/186464 A1 (NOVOZYMES AS [DK]; MONOSOL LLC [US]; CASELLA VICTOR [US]) 20 November 2014 (2014-11-20) claim 3 -----	2-16
A	WO 97/04079 A1 (NOVO-NORDISK AS) 6 February 1997 (1997-02-06) claim 3 -----	1-16

☐

Further documents are listed in the continuation of Box C.

☒

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

3 March 2016

Date of mailing of the international search report

17/03/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Obel, Nicolai

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2015/080463

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6624129	B1	23-09-2003	AU 3247699 A 06-09-1999
		EP 1054956 A1 29-11-2000	
		JP 2003526319 A 09-09-2003	
		US 6624129 B1 23-09-2003	
		WO 9942566 A1 26-08-1999	

WO 0060063	A1	12-10-2000	AU 3420100 A 23-10-2000
		EP 1171581 A1 16-01-2002	
		JP 4523178 B2 11-08-2010	
		JP 2003520021 A 02-07-2003	
		MX PA01009700 A 14-05-2002	
		WO 0060063 A1 12-10-2000	

WO 2014186464	A1	20-11-2014	CN 105209612 A 30-12-2015
		EP 2997139 A1 23-03-2016	
		EP 2997140 A1 23-03-2016	
		WO 2014184164 A1 20-11-2014	
		WO 2014186464 A1 20-11-2014	

WO 9704079	A1	06-02-1997	AT 282087 T 15-11-2004
		AU 6414196 A 18-02-1997	
		CN 1193346 A 16-09-1998	
		DE 69633825 D1 16-12-2004	
		DE 69633825 T2 10-11-2005	
		EP 0839186 A1 06-05-1998	
		JP 4307549 B2 05-08-2009	
		JP 2001526523 A 18-12-2001	
		WO 9704078 A1 06-02-1997	
		WO 9704079 A1 06-02-1997	
