(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
21 June 2007 (21.06.2007)

PCT

(10) International Publication Number
WO 2007/068476 A1

(51) International Patent Classification:
C07D 471/08 (2006.01) *A61P 25/00* (2006.01)
A61K 31/439 (2006.01)

(21) International Application Number:
PCT/EP2006/012023

(22) International Filing Date:
14 December 2006 (14.12.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0525672.2 16 December 2005 (16.12.2005) GB

(71) Applicant *(for all designated States except AT, US)*: **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(71) Applicant *(for AT only)*: **NOVARTIS PHARMA GMBH** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **FEUERBACH, Dominik** [DE/DE]; Wehrgasse 3, 79379 Müllheim (DE). **FREDERIKSEN, Mathias** [NO/CH]; Pfeffingerstrasse 18, CH-4053 Basel (CH). **HURTH, Konstanze** [DE/FR]; 4, rue de l'Ancien Golf, F-68300 Saint Louis (FR). **ROY, Bernard, Lucien** [CH/CH]; Route de la Broye 10, CH-1700 Fribourg (CH). **WAGNER, Beatrix** [DE/DE]; Talweg 50, 79540 Lörrach (DE).

(74) Agent: **VOEGELI-LANGE, Regina**; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:
— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: [(1H- INDOL- 5 -YL) -HETEROARYLOXY] - (1-AZA-BICYCLO [3.3.1] NONANES AS CHOLINERGIC LIGANDS OF THE N-ACHR FOR THE TREATMENT OF PSYCHOTIC AND NEURODEGENRATIVE DISORDERS.



(I)


(57) Abstract: The present invention relates to 1-azabicycloalkyl derivatives of Formula (I) wherein the substituents are as defined in the specification and to processes for their production to pharmaceutical compositions comprising them and to their use in the manufacture of a medicament for the treatment and/ or delay of progression of psychotic and nemodegenerative disorders.

WO 2007/068476 A1

[(1H-INDOL-5-YL)-HETEROARYLOXY]-(1-AZA-BICYCLO[3.3.1]NONANES AS CHOLINERGIC LIGANDS OF THE N-ACHR FOR THE TREATMENT OF PSYCHOTIC AND NEURODEGENRATIVE DISORDERS.

The present invention relates to novel 1-aza-bicyclononane derivatives, to processes for their production, their use as pharmaceuticals and to pharmaceutical compositions comprising them.

More particularly the present invention provides in a first aspect, a compound of formula (I)




(I)

wherein


X represents hydrogen or hydroxyl and

Y represents one of the following groups:




in free base or acid addition salt form.

A preferred compound according to the invention is (4S,5R)-4-[5-(1H-indol-5-yl)-pyrimidin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane having the formula shown below.




(II)

A further preferred compound according to the invention is 5-{2-[(4S,5R)-(1-aza-bicyclo[3.3.1]non-4-yl)oxy]-pyrimidin-5-yl}-1,3-dihydro-indol-2-one having the formula shown below.




(III)

A further preferred compound according to the invention is (4S,5R)-4-[6-(1H-indol-5-yl)-pyridin-3-yloxy]-1-aza-bicyclo[3.3.1]nonane having the formula shown below.




(IV)

A further preferred compound according to the invention is (4S,5R)-4-[5-(1H-indol-5-yl)-pyridin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane having the formula shown below.




(V)

A further preferred compound according to the invention is (4S,5R)-4-[6-(1H-indol-5-yl)-pyridazin-3-yloxy]-1-aza-bicyclo[3.3.1]nonane having the formula shown below.



(VI)

A further preferred compound according to the invention is 5-{6-[(4S,5R)-(1-aza-bicyclo[3.3.1]non-4-yl)oxy]-pyridazin-3-yl}-1,3-dihydro-indol-2-one having the formula shown below.



(VII)

Compounds of formula (I) exist in free or acid addition salt form. In this specification, unless otherwise indicated, language such as "compounds of formula (I)" is to be understood as embracing the compounds in any form, for example free base or acid addition salt form. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds of formula (I) , such as picrates or perchlorates, are also included. For therapeutic use, only pharmaceutically acceptable salts or free compounds are employed (where applicable in the form of pharmaceutical preparations), and are therefore preferred.


Compounds of formula (I) may exist in form of various isomers, e.g. keto-enol tautomers. In this specification, unless otherwise indicated, language such as "compounds of formula (I)" is to be understood as embracing the compounds in any form, for example in the keto or in the enol form or any mixture of them

Where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

In a further aspect, the present invention also provides processes for the production of compounds of formula (I).

A first process comprises the steps of


i) reacting a compound of formula (IX)



(IX)

wherein Y is as defined above and Z z represents a leaving group, such as Cl, Br, I, Tosylate

with a compound of formula (X)




(X)

where R represents H , C_1-C_4alkyl or both RO represent together with the B to which they are attached a heterocyclic moiety, X is as defined above, and

ii) recovering the so obtained compound of formula (I)


A second process comprises the steps of

i) reacting a compound of formula (XI)



(XI)

with a compound of formula XII



(XII)

wherein X and Y are as defined above and PG is a suitable protecting group,

ii) subsequent deprotection of the so obtained compound and

iii) recovering the so obtained compound of formula (I) in free base or acid addition salt form.

This process is particular suitable wherein Y represents the 3-pyridyl moiety.

Starting materials are known or may be obtained by well known processes. The synthesis of the starting materials is described e.g in GB 123456, which is hereby incorporated by reference.

The following considerations apply to the individual reaction steps described above:

a) One or more functional groups, for example carboxy, hydroxy, amino, or mercapto, may need to be protected in the starting materials by protecting groups. The protecting groups employed may already be present in precursors and should protect the functional groups concerned against unwanted secondary reactions, such as acylations, etherifications, esterifications, oxidations, solvolysis, and similar reactions. It is a characteristic of protecting groups that they lend themselves readily, i.e. without undesired secondary reactions, to removal, typically by solvolysis, reduction, photolysis or also by enzyme activity, for example under conditions analogous to physiological conditions, and that they are not present in the end-products. The specialist knows, or can easily establish, which protecting groups are suitable with the reactions mentioned hereinabove and hereinafter. The protection of such functional groups by such protecting groups, the protecting groups themselves, and their removal reactions are described for example in standard reference works, such as J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973, in T. W. Greene, "Protective Groups in Organic Synthesis", Wiley, New York 1981, in "The Peptides"; Volume 3 (editors: E. Gross and J. Meienhofer), Academic Press, London and New York 1981, in "Methoden der organischen Chemie" (Methods of organic chemistry), Houben Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974, in H.-D. Jakubke and H. Jescheit, "Aminosäuren, Peptide, Proteine" (Amino acids, peptides, proteins), Verlag Chemie, Weinheim, Deerfield Beach, and Basel 1982, and in Jochen Lehmann, "Chemie der Kohlenhydrate: Monosaccharide und Derivate" (Chemistry of carbohydrates: monosaccharides and derivatives), Georg Thieme Verlag, Stuttgart 1974.

b) Acid addition salts may be produced from the free bases in known manner, and vice-versa. Alternatively, optically pure starting materials can be used. Suitable acid addition salts for use in accordance with the present invention include for example the hydrochloride.

c) Stereoisomeric mixtures, e.g. mixtures of diastereomers, can be separated into their corresponding isomers in a manner known per se by means of suitable separation methods. Diastereomeric mixtures for example may be separated into their individual diastereomers by means of fractionated crystallization, chromatography, solvent distribution, and similar procedures. This separation may take place either at the level of a starting compound or in a compound of formula I itself. Enantiomers may be separated through the formation of diastereomeric salts, for example by salt formation with an enantiomer-pure chiral acid, or by means of chromatography, for example by HPLC, using chromatographic substrates with chiral ligands. Alternatively, optically pure starting materials can be used.

d) Suitable diluents for carrying out the above- described are especially inert organic solvents. These include, in particular, aliphatic, alicyclic or aromatic, optionally halogenated hydrocarbons, such as, for example, benzine, benzene, toluene, xylene, chlorobenzene, dichlorobenzene, petroleum ether, hexane, cyclohexane, dichloromethane, chloroform, carbon tetrachloride; ethers, such as diethyl ether, diisopropyl ether, dioxane, tetrahydrofuran or ethylene glycol dimethyl ether or ethylene glycol diethyl ether; ketones, such as acetone, butanone or methyl isobutyl ketone; nitriles, such as acetonitrile propionitrile or butyronitrile; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methyl-formanilide, N-methyl-pyrrolidone or hexamethylphosphoric triamide; esters, such as methyl acetate or ethyl acetate, sulphoxides, such as dimethyl sulphoxide, alcohols, such as methanol, ethanol, n- or i-propanol, ethylene glycol monomethyl ether, ethylene glycol monoethyl ether, diethyelene glycol monomethyl ether, diethylene glycol monoethyl ether. Further, mixtures of diluents may be employed. Depending on the starting materials, reaction conditions and auxiliaries, water or diluents constaining water may be suitable. It is also possible to use one a starting material as diluent simultaneously.

e) Reaction temperatures can be varied within a relatively wide range. In general, the processes are carried out at temperatures between 0°C and 150°C, preferably between 10°C and 120°C. Deprotonation reactions can be varied within a relatively wide range. In general, the processes are carried out at temperatures between -150°C and +50°C, preferably between -75°C and 0°C.

f) The reactions are generally carried out under atmospheric pressure. However, it is also possible to carry out the processes according to the invention under elevated or reduced pressure – in general between 0.1 bar and 10 bar.

g) Starting materials are generally employed in approximately equimolar amounts. However, it is also possible to use a relatively large excess of one of the components. The reaction is generally carried out in a suitable diluent in the presence of a reaction auxiliary, and the reaction mixture is generally stirred at the required temperature for a number of hours.

h) Working up the reaction mixtures according to the above processes and purification of the compounds thus obtained may be carried out in accordance to known procedures (cf. the Preparation Examples).

The compounds of the invention, exhibit valuable pharmacological properties when tested in vitro and in animals, and are therefore useful as pharmaceuticals.

Thus, the compound of the invention are found to be cholinergic ligands of the nAChR. In addition preferred compound of the invention show selective α7-nAChR activity. The compounds of the present invention may in particular be found to be agonists, partial agonists, antagonists or allosteric modulators of the receptor.

Due to their pharmacological profiles, compound of the invention are anticipated to be useful for the treatment of diseases or conditions as diverse as CNS related diseases, PNS related diseases, diseases related to inflammation, pain and withdrawal symptoms caused by an abuse of chemical substances. Diseases or disorders related to the CNS include general anxiety disorders, cognitive disorders, learning and memory deficits and dysfunctions, Alzheimer`s disease (AD), prodromal AD, mild cognitive impairment in the elderly (MCI), amnestic MCI, age associated memory impairment, attention deficit and hyperactivity disorder (ADHD), Parkinson`s disease, Huntington's disease, ALS, prionic neuro-degenerative disorders such as Creutzfeld-Jacob disease and kuru disease, Gilles de la Tourette's syndrome, psychosis, depression and depressive disorders, mania, manic depression, schizophrenia, the cognitive deficits in schizophrenia, obsessive compulsive disorders, panic disorders, eating disorders, narcolepsy, nociception, AIDS-dementia, senile dementia, mild cognitive dysfunctions related to age, autism, dyslexia, tardive dyskinesia,

epilepsy, and convulsive disorders, post-traumatic stress disorders, transient anoxia, pseudodementia, pre-menstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome and jet lag. Furthermore, compound of the invention may be useful for the treatment of endocrine disorders, such as thyrotoxicosis, pheochromocytoma, hypertension and arrhythmias as well as angina pectoris, hyperkinesia, premature ejaculation and erectile difficulty. Still further, compound of the invention may be useful in the treatment of inflammatory disorders (Wang et al., Nature 2003, 421, 384; de Jonge et al., Nature Immunology 2005, 6, 844; Saeed et al., JEM 2005, 7, 1113), disorders or conditions including inflammatory skin disorders, rheumatoid arthritis, post-operative ileus, Crohn's diesease, inflammatory bowel disease, ulcerative colitis, sepsis, fibromyalgia, pancreatitis and diarrhoea. Compound of the invention may further be useful for the treatment of withdrawal symptoms caused by termination of the use of addictive substances, like heroin, cocaine, tobacco, nicotine, opioids, benzodiazepines and alcohol. Finally, compound of the invention may be useful for the treatment of pain, e.g. caused by migraine, postoperative pain, phantom limb pain or pain associated with cancer. The pain may comprise inflammatory or neuropathic pain, central pain, chronic headache, pain related to diabetic neuropathy, to post therapeutic neuralgia or to peripheral nerve injury.

Furthermore, degenerative ocular disorders which may be treated include ocular diseases which may directly or indirectly involve the degeneration of retinal cells, including ischemic retinopathies in general, anterior ischemic optic neuropathy, all forms of optic neuritis, age-related macular degeneration (AMD), in its dry forms (dry AMD) and wet forms (wet AMD), diabetic retinopathy, cystoid macular edema (CME), retinal detachment, retinitis pigmentosa, Stargardt's disease, Best's vitelliform retinal degeneration, Leber's congenital amaurosis and other hereditary retinal degenerations, pathologic myopia, retinopathy of prematurity,and Leber's hereditary optic neuropathy.

It has been found that the effect of a combination which comprises at least one nicotinic-alpha 7 receptor agonist and at least one compound selected from the group consisting of (a) conventional antipsychotics and (b) atypical antipsychotics is greater than the additive effect of the combined drugs in the treatment of psychiatric disorders. In particular, the combinations disclosed herein can be used to treat schizophrenia which is refractory to monotherapy employing one of the combination partners alone.

Hence, the invention relates to a combination, such as a combined preparation or pharmaceutical composition, which comprises at least one nicotinic-alpha 7 receptor agonist and at least one compound selected from the group consisting of (a) conventional antipsychotics and (b) atypical antipsychotics, in which the active ingredients are present in each case in free form or in the form of a pharmaceutically acceptable salt and optionally at least one pharmaceutically acceptable carrier; for simultaneous, separate or sequential use.

The term "psychiatric disorders" as used herein includes, but is not limited to schizophrenia, anxiety disorders, depression and bipolar disorders. Preferably, the psychiatric disorder to be treated with the combination disclosed herein is schizophrenia, more preferably schizophrenia which is refractory to monotherapy employing one of the combination partners alone. The term "conventional antipsychotics" as used herein includes, but is not limited to haloperidol, fluphenazine, thiotixene and flupentixol.

The term "atypical antipsychotics" as used herein includes, but is not limited to clozaril, risperidone, olanzapine, quetiapine, ziprasidone and aripiprazol.

In another aspect, the compound of the invention are used as diagnostic agents and/or PET ligands, e.g. for the identification and localization of nicotine receptors in various tissues. Properly isotope-labeled agents of the invention exhibit valuable properties as histopathological labeling agents, imaging agents and/or biomarkers, hereinafter "markers", for the selective labeling of the nAChR. More particularly the agents of the invention are useful as markers for labeling the alpha7 nAChR receptors *in vitro* or *in vivo*. In particular, compound of the invention which are properly isotopically labeled are useful as PET markers. Such PET markers are labeled with one or more atoms selected from the group consisting of ^{11}C, ^{13}N, ^{15}O, ^{18}F.

The agents of the invention are therefore useful, for instance, for determining the levels of receptor occupancy of a drug acting at the nAChR, or diagnostic purposes for diseases resulting from an imbalance or dysfunction of nAChR, and for monitoring the effectiveness of pharmacotherapies of such diseases.

In accordance with the above, the present invention provides an agent of the invention for use as a marker for neuroimaging.

In a further aspect, the present invention provides a composition for labeling brain and peripheral nervous system structures involving nAChR*in vivo* and *in vitro* comprising an agent of the invention.

In still a further aspect, the present invention provides a method for labeling brain and peripheral nervous system structures involving nAChR*in vitro* or *in vivo*, which comprises contacting brain tissue with an agent of the invention.

The method of the invention may comprise a further step aimed at determining whether the agent of the invention labeled the target structure. Said further step may be effected by observing the target structure using positron emission tomography (PET) or single photon emission computed tomography (SPECT), or any device allowing detection of radioactive radiations.

In particular, the agents of the invention are α7 nicotinic acetylcholine receptor (αnAChR α7) agonists.

In functional assays, the agents of the invention display high affinity at the nAChR α7 as shown in the following tests:

a) A functional assay for affinity at the nAChR α7 is carried out with a rat pituitary cell line stably expressing the nAChR α7. Briefly, GH3 cells recombinantly expressing the nAChR α7 were seeded 72 h prior to the experiment on black 96-well plates and incubated at 37°C in a humidified atmosphere (5 % CO_2/95 % air). On the day of the experiment medium was removed by flicking the plates and replaced with 100 µl growth medium containing af fluorescent calcium sensitve dye, in the presence of 2.5 mM probenecid (Sigma). The cells were incubated at 37°C in a humidified atmosphere (5 % CO_2/95 % air) for 1 h. Plates were flicked to remove excess of Fluo-4, washed twice with Hepes-buffered salt solution (in mM: NaCl 130, KCl 5.4, $CaCl_2$ 2, $MgSO_4$ 0.8, NaH_2PO_4 0.9, glucose 25, Hepes 20, pH 7.4; HBS) and refilled with 100 µl of HBS containing antagonists when appropriate. The incubation in the presence of the antagonist lasted between 3 and 5 minutes. Plates were then placed into an imaging plate reader and fluorescence signal recordedd In this assay, compound of the

invention exhibit pEC_{50} values of about 5 to about 9. Partial and potent agonists in this test are preferred.

b) To assess the antagonist activity of the compound of the invention on the human neuronal nAChR $\alpha4\beta2$, a similar functional assay is carried out using a human epithelial cell line stably expressing the human $\alpha4\beta2$ subtype (Michelmore et al., Naunyn-Schmiedeberg's Arch. Pharmacol. (2002) 366, 235) In this assay, the preferred compounds of the invention show selectivity for the nAChR $\alpha7$ subtype.

c) To assess the antagonist activity of the compound of the invention on the "ganglionic subtype" ($\alpha3\beta4$), the muscle type of nicotinic receptor ($\alpha1\beta1\gamma\delta$) and the 5-$HT_3$ receptor, similar functional tests as just described under a) are carried out with a human epithelial cell line stably expressing the human ganglionic subtype, a cell line endogenously expressing the human muscle type of nicotinic receptors or a cell line endogenously expressing the murine 5-HT_3 receptor (Michelmore et al., Naunyn-Schmiedeberg's Arch. Pharmacol. (2002) 366, 235). Compounds which display little or no activity on the $\alpha3\beta4$ nAChR, the muscle subtype of nicotinic receptor as well as the 5-HT_3 receptor are especially preferred.

In the model of mice showing sensory gating deficit (DBA/2-mice) described by S. Leonard et al. in Schizophrenia Bulletin 22, 431-445 (1996), the compound of the invention induce significant sensory gating at concentrations of about 10 to about 40 μM.

The compound of the invention may be shown to increase attention in a test of attention for rodents (Robbins, J. Neuropsychiatry Clin. Neurosci. (2001) 13, 326-35), namely the 5-choice serial reaction time test (5-CSRTT). In this test, the rat must observe a wall containing 5 holes. When a light flash appears in one of them, the rat must respond with a nose-poke into the correct hole within 5 sec. in order to receive a food pellet reward, delivered to a feeder in the opposite wall.

Compound of the invention may also show learning/memory enhancing effects in the social recognition and in the object recognition test in mice and rats (Ennaceur and Delacour, Behav. Brain Res. (1988) 31, 47-59).

The compound of the invention are therefore useful for the prevention and treatment (including mitigation and prevention) of various disorders, especially those mentioned above. The usefulness of nAChR α7 agonists in neurodegeneration is documented in the literature, e.g. in Wang et al., J. Biol. Chem. 275, 5626-5632 (2000).

For the treatment of the above and other disorders, the appropriate dosage of a compound (active ingredient) of the invention will, of course, vary depending upon, for example, the host, the mode of administration and the nature and severity of the condition being treated as well as the relative potency of the particular agent of the invention employed. For example, the amount of active agent required may be determined on the basis of known in vitro and in vivo techniques, determining how long a particular active agent concentration in the blood plasma remains at an acceptable level for a therapeutic effect. In general, satisfactory results in animals are indicated to be obtained at daily dosages of from about 0.01 to about 30.0 mg/kg p.o. In humans, an indicated daily dosage is in the range of from about 0.7 to about 1400 mg/day p.o., e.g. from about 50 to 200 mg (70 kg man), conveniently administered once or in divided doses up to 4 x per day or in sustained release form. Oral dosage forms accordingly suitably comprise from about 1.75 or 2.0 to about 700 or 1400 mg of a compound of the invention admixed with an appropriate pharmaceutically acceptable diluent or carrier therefore.

Pharmaceutical compositions contain, for example, from about 0.1 % to about 99.9 %, preferably from about 20 % to about 60 %, of the active ingredient(s).

Examples for compositions comprising a compound of the invention include, for example, a solid dispersion, an aqueous solution, e.g. containing a solubilising agent, a microemulsion and a suspension of, e.g. a salt of a compound of formula I or a free compound of the formula I in the range of from 0.1 to 1 %, e.g. 0.5 %. The composition may be buffered to a pH in the range of, e.g. from 3.5 to 9.5, e.g. to pH 4.5, by a suitable buffer.

The compound of the invention are also commercially useful as research chemicals.

For use according to the invention, a compound of the formula I and/or a pharmaceutically acceptable salt thereof may be administered as single active agent or in combination with one or more other active agents of the formula I and/or a pharmaceutically acceptable salt thereof or especially other active agents commonly employed especially for the treatment of

the disorders mentioned herein or further other disorders, in any customary manner, e.g. orally, for example in the form of tablets, capsules, or as nasal spray, or parenterally, for example in the form of injection solutions or suspensions. The other active agents employed in such combinations are preferably selected from the group consisting of benzodiazepines, selective serotonin reuptake inhibitors (SSRIs), selective serotonin and norepinephrine reuptake inhibitors (SNRIs), conventional antipsychotics, atypical antipsychotics, buspirone, carbamazepine, oxcarbazepine, gabapentin and pregabalin.

An SSRI suitable for the present invention is especially selected from fluoxetine, fuvoxamine, sertraline, paroxetine, citalopram and escitalopram. An SNRI suitable for the present invention is especially selected from venlafaxine and duloxetine. The term "benzodiazepines" as used herein includes, but is not limited to clonazepam, diazepam and lorazepam. The term "conventional antipsychotics" as used herein includes, but is not limited to haloperidol, fluphenazine, thiotixene and flupentixol. The term "atypical antipsychotics" as used herein relates to clozaril, risperidone, olanzapine, quetiapine, ziprasidone and aripiprazol.

Buspirone can be administered in free form or as a salt, e.g. as its hydrochloride, e.g., in the form as marketed, e.g. under the trademark Buspar™ or Bespar™. It can be prepared and administered, e.g., as described in US 3,717,634. Fluoxetine can be administered, e.g., in the form of its hydrochloride as marketed, e.g. under the trademark Prozac™. It can be prepared and administered, e.g., as described in CA 2002182. Paroxetine ((3S,4R)-3-[(1,3-benzodioxol-5-yloxy)methyl]-4-(4-fluorophenyl)piperidine) can be administered, e.g., in the form as marketed, e.g. under the trademark Paxil™. It can be prepared and administered, e.g., as described in US 3,912,743. Sertraline can be administered, e.g., in the form as marketed, e.g. under the trademark Zoloft™. It can be prepared and administered, e.g., as described in US 4,536,518. Clonazepam can be administered, e.g., in the form as marketed, e.g. under the trademark Antelepsin™. Diazepam can be administered, e.g., in the form as marketed, e.g. under the trademark Diazepam Desitin™. Lorazepam can be administered, e.g., in the form as marketed, e.g. under the trademark Tavor™. Citalopram can be administered in free form or as a salt, e.g. as its hydrobromide, e.g., in the form as marketed, e.g. under the trademark Cipramil™. Escitalopram can be administered, e.g., in the form as marketed, e.g. under the trademark Cipralex™. It can be prepared and administered, e.g., as described in AU623144. Venlafaxine can be administered, e.g., in the form as marketed, e.g. under the trademark Trevilor™. Duloxetine can be administered, e.g.,

in the form as marketed, e.g. under the trademark Cymbalta™. It may be prepared and administered, e.g., as described in CA 1302421. Carbamazepine can be administered, e.g., in the form as marketed, e.g. under the trademark Tegretal™ or Tegretol™. Oxcarbazepine can be administered, e.g., in the form as marketed, e.g. under the trademark Trileptal™. Oxcarbazepine is well known from the literature [see for example Schuetz H. et al., Xenobiotica (GB), 16(8), 769-778 (1986)]. Gabapentin can be administered, e.g., in the form as marketed, e.g. under the trademark Neurontin™. Haloperidol can be administered, e.g., in the form as marketed, e.g. under the trademark Haloperidol STADA™. Fluphenazine can be administered, e.g., in the form of its dihydrochloride as marketed, e.g. under the trademark Prolixin™. Thiothixene can be administered, e.g., in the form as marketed, e.g. under the trademark Navane™. It can be prepared, e.g., as described in US 3,310,553. Flupentixol can be administered for instance in the form of its dihydrochloride, e.g., in the form as marketed, e.g. under the trademark Emergil™ or in the form of its decanoate, e.g., in the form as marketed, e.g. under the trademark Depixol™. It can be prepared, e.g., as described in BP 925,538. Clozaril can be administered, e.g., in the form as marketed, e.g. under the trademark Leponex™. It can be prepared, e.g., as described in US 3,539,573. Risperidone can be administered, e.g., in the form as marketed, e.g. under the trademark Risperdal™. Olanzapine can be administered, e.g., in the form as marketed, e.g. under the trademark Zyprexa™. Quetiapine can be administered, e.g., in the form as marketed, e.g. under the trademark Seroquel™. Ziprasidone can be administered, e.g., in the form as marketed, e.g. under the trademark Geodon™. It can be prepared, e.g., as described in GB 281,309. Aripiprazole can be administered, e.g., in the form as marketed, e.g. under the trademark Abilify™. It can be prepared, e.g., as described in US 5,006,528.

The structure of the active ingredients identified by code nos., generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications). The corresponding content thereof is hereby incorporated by reference. Any person skilled in the art is fully enabled to identify the active ingredients and, based on these references, likewise enabled to manufacture and test the pharmaceutical indications and properties in standard test models, both *in vitro* and *in vivo*.

In the case of a combination, the pharmaceutical compositions for separate administration of the combination partners and/or those for administration in a fixed combination, i.e. a single

galenical composition comprising at least two combination partners, according to the invention can be prepared in a manner known per se and are those suitable for enteral, such as oral or rectal, and parenteral administration to mammals, including man, comprising a therapeutically effective amount of at least one pharmacologically active combination partner alone or in combination with one or more pharmaceutically acceptable carriers, especially suitable for enteral or parenteral application. When the combination partners employed are applied in the form as marketed as single drugs, their dosage and mode of administration can take place in accordance with the information provided on the packet leaflet of the respective marketed drug in order to result in the beneficial effect described herein, if not mentioned herein otherwise.

Pharmaceutical preparations for the combination therapy for enteral or parenteral administration are, for example, those in unit dosage forms, such as sugar-coated tablets, tablets, capsules or suppositories, or furthermore ampoules. If not indicated otherwise, these are prepared in a manner known per se, for example by means of conventional mixing, granulating, sugar-coating, dissolving or lyophilizing processes. It will be appreciated that the unit content of a combination partner contained in an individual dose of each dosage form need not in itself constitute an effective amount since the necessary effective amount can instead with a single dosage unit also be reached by administration of a two or more dosage units.

In particular, a therapeutically effective amount of each of the combination partners may be administered simultaneously or sequentially and in any order, and the components may be administered separately (e.g. sequentially after fixed or variable periods of time), or as a fixed combination. For example, the method of treatment (including mitigation) of a disorder according to the invention may comprise (i) administration of the combination partner (a) (a compound of the present invention) in free or pharmaceutically acceptable salt form and (ii) administration of a combination partner (b) (e.g. a different compound of the present invention or an active ingredient of a different formula) in free or pharmaceutically acceptable salt form, simultaneously or sequentially in any order, in jointly therapeutically effective amounts, preferably in synergistically effective amounts, e.g. in daily dosages corresponding to the amounts described herein. The individual combination partners can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. Furthermore, the term "administering" also encompasses the use of a prodrug of a combination partner that convert *in vivo* to the combination partner as such. The

instant invention is therefore to be understood as embracing all such regimes of simultaneous and/ or alternating treatment and the term "administering" is to be interpreted accordingly.

The effective dosage of the combination partners employed may vary, for example depending on the particular compound or pharmaceutical composition employed, the mode of administration, the disorder being treated, and/or the severity of the disorder being treated. Thus, the dosage regimen is selected in accordance with a variety of factors including the route of administration, metabolism by and the renal and hepatic function of the patient. A physician, clinician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the single active ingredients required to prevent, mitigate, counter or arrest the disorder. Optimal precision in achieving concentration of the active ingredients within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the active ingredients' availability to target sites.

In accordance with the foregoing, the present invention also provides:

(1) A compound of the formula I, and/or a salt thereof, for use in the diagnostic or therapeutic treatment of a mammal, especially a human; especially for use as an alpha-7 receptor agonist, for example for use in the treatment (including mitigation) of any one or more disorders, especially of any one or more of the particular disorders set forth hereinbefore and hereinafter.

(2) A pharmaceutical composition comprising a compound of the formula I, and/or a pharmaceutically acceptable salt thereof, as active ingredient together with a pharmaceutically acceptable diluent or carrier.

(2') A pharmaceutical composition for the treatment or prevention of a disorder in the treatment of which alpha-7 receptor activation plays a role or is involved and/or in which alpha-7 receptor activity is involved, especially any one or more of the disorders mentioned hereinbefore or hereinafter, comprising a compound of the formula I, and/or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.

(3) A method for the treatment of a disorder, especially any one or more of the particular disorders set forth hereinbefore, in a subject in need of such treatment, comprising administering a pharmaceutically effective amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof.

(3') A method for treating or preventing a disorder in the treatment of which alpha-7 receptor activation plays a role or is involved and/or in which alpha-7 receptor activity is involved, com-

prising administering to a mammal in need thereof a therapeutically effective amount of a compound of the formula I, and/or a pharmaceutically acceptable salt thereof.

(4) The use of a compound of the formula I, and/or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a disease or condition in the treatment of which alpha-7 receptor activation plays a role or is involved and/or in which alpha-7 receptor activity is involved, especially one or more of the disorders mentioned above.

(5) A method as defined above comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of an alpha-7 agonist of the formula I, and/or a pharmaceutically acceptable salt thereof, and a second pharmaceutically active compound and/or a pharmaceutically acceptable salt thereof, said second pharmaceutically active compound and/or salt thereof being especially for use in the treatment of any one or more of the disorders set forth hereinbefore or hereinafter.

(6) A combination comprising a therapeutically effective amount of an alpha-7 agonist of the formula I, and/or a pharmaceutically acceptable salt thereof, and a second pharmaceutically active compound and/or a pharmaceutically acceptable salt thereof, said second pharmaceutically active compound being especially for use or of use in the treatment of any one or more of the particular disorders set forth hereinbefore.

The Examples which follow serve to illustrate the invention without limiting the scope thereof.

The following abbreviations are used :

AcOEt	ethyl acetate
aq.	aqueous
EtOH	ethanol
FC	flash chromatography
HV	high vacuum
MeOH	MeOH
m.p.	melting point
MTBE	methyl tert-butyl ether
NHMDS	sodium hexamethyl disilazane
rt	room temperature
soln.	solution
THF	tetrahydrofuran

Temperatures are measured in degrees Celsius. Unless indicated otherwise, reactions are carried out at room temperature. The structure of final products, intermediates and starting materials is confirmed by standard analytical methods, e.g. microanalysis and spectroscopic characteristics (e.g. MS, IR, NMR).

example 1 (4S,5R)-4-[5-(1H-Indol-5-yl)-pyrimidin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane

1-Aza-bicyclo[3.3.1]nonan-4-one (11.92 g, 85.6 mmol) is dissolved in 160 ml MeOH and cooled to -10°C. $NaBH_4$ (1.69 g, 42.9 mmol)) is added portionwise so that the inner temperature does not exceed 0°C. The reaction mixture is stirred at -10°C for 1h. Water is added and the solvents are evaporated. The remaining solid is dissolved in MTBE/MeOH, filtered over hyflo and the filtrate is evaporated to give 19.28 g of crude product which is purified by chromatography over aluminum oxide (400g, eluent: MTBE/MeOH 95:5 to 80:20) to give 10.96 g (91%) (4SR,5RS)-1-aza-bicyclo[3.3.1]nonan-4-ol.

To acetic anhydride (150 ml) (4SR,5RS)-(1-aza-bicyclo[3.3.1]nonan-4-ol (21.34 g, 151 mmol) is added portionwise under cooling. The reaction mixture is heated to 120°C for 2.5 h, is evaporated and extracted with THF/sat. K_2CO_3 soln. The aq. layer is reextracted with THF. The combined organic layers are washed with brine, dried over Na_2SO_4, filtered and evaporated. The crude product is purified by bulb-to-bulb distillation (HV, 90°C) to give 24.07 g (87%) of (4SR,5RS)-acetic acid 1-aza-bicyclo[3.3.1]non-4-yl ester.

Acetic acid 1-aza-bicyclo[3.3.1]non-4-yl ester (120.0 g, 655 mmol) is dissolved in 200 ml EtOH abs. and 50 ml water. L(+)-tartaric acid (98.3 g, 655 mmol) is added and the mixture is heated to reflux for 1 minute. The mixture is cooled to rt, then to 4°C. The precipitate is filtered off, washed with EtOH, and 3x recrystallized from EtOH/H_2O 4:1 to give 41.16 g of the tartrate salt ($[\alpha]_D^{rt}$ = -14.72° (c=0.265, MeOH)) which gives after treatment with sat. Na_2CO_3 soln. 22.6 g (123 mmol, 19%) (4S,5R)-acetic acid 1-aza-bicyclo[3.3.1]non-4-yl ester as the free base. Another 10.7 g (32.1 mmol, 5%) of tartrate can be obtained from the mother liquors by another 3 recrystallizations from EtOH/H_2O 4:1.

(4S,5R)-acetic acid 1-aza-bicyclo[3.3.1]non-4-yl ester (5.45 g, 29.7 mmol) is dissolved in 10% aq. NaOH soln. and stirred for 1h at 50°C. After cooling to rt, the mixture is extracted with THF/brine. The organic layer is dried over Na_2SO_4, filtered and the filtrate is evaporated to give 3.83 g (91%) of (4S, 5R)-1-aza-bicyclo[3.3.1]nonan-4-ol, which is dissolved in 50 ml THF and cooled to 0°C. NHMDS (35 ml of 1 M THF soln.) is added dropwise, the reaction mixture is stirred at rt for 0.5h and then added to a precooled soln. (-15°C) of 5-bromo-2-chloro-pyrimidine (5.89 g, 30.5 mmol) in THF. The mixture is stirred for 15h at rt, then diluted

with THF and extracted with 1M aq. NaOH soln. and brine. The aq. layers are 2x reextracted with THF, the combined organic layers are dried over Na_2SO_4, filtered and the filtrate is evaporated. The resulting crude product (10.15 g) is recrystallized from CH_3CN/MeOH to give 5.21 g (65%) of (4S,5R)-4-(5-bromo-pyrimidin-2-yloxy)-1-aza-bicyclo[3.3.1]nonane.

5.19 g (17.4 mmol) of (4S,5R)-4-(5-bromo-pyrimidin-2-yloxy)-1-aza-bicyclo[3.3.1]nonane is dissolved in 200 ml toluene/EtOH 9:1. 5-Indolyl boronic acid (3.47 g, 21.6 mmol), $Pd(PPh_3)_4$ (1.04 g, 0.873 mmol) and a soln. of Na_2CO_3 (7.38 g, 69.6 mmol) in 35 ml H_2O is added. The reaction mixture is stirred at 90°C for 15h. After cooling to rt the mixture is filtered over hyflo, the filtrate is extracted with water and brine. The aq. layers are reextracted with AcOEt and the combined organic layers are dried over Na_2SO_4, filtered and the filtrate is evaporated. The resulting crude product is purified by FC (255 g silica gel, eluent AcOEt/MeOH/Et_3N 70:27:3) and recrystallization from EtOH to give 3.87 g (67%) (4S,5R)-4-[5-(1H-indol-5-yl)-pyrimidin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane. MS (ES^+): m/e = 335 (MH^+), m.p. 195-199°C.

Preparation of the precursor 1-aza-bicyclo[3.3.1]nonan-4-one is done according to M. G. Kim et al., J. Med. Chem. (2003) *46*, 2216.


example 2 (Manufacture of Soft Capsules)

5000 soft gelatin capsules, each comprising as active ingredient 0.05 g of one of the compounds of formula (I) mentioned in the preceding Examples, are prepared as follows:

250 g of the pulverized active ingredient is suspended in 2 litres Lauroglykol® (propylene glycol laurate, Gattefossé S.A., Saint Priest, France) and ground in a wet pulverizer to produce a particle size of about 1 to 3 µm. 0.419 g portions of the mixture are then introduced into soft gelatin capsules using a capsule-filling machine.

Claims:

1. A compound of formula (I)




(I)

wherein

X represents hydrogen or hydroxyl and

Y represents one of the following groups:



in free base or acid addition salt form.

2. (4S,5R)-4-[5-(1H-Indol-5-yl)-pyrimidin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane .

3. 5-{2-[(4S,5R)-(1-Aza-bicyclo[3.3.1]non-4-yl)oxy]-pyrimidin-5-yl}-1,3-dihydro-indol-2-one.

4. (4S,5R)-4-[6-(1H-Indol-5-yl)-pyridin-3-yloxy]-1-aza-bicyclo[3.3.1]nonane .


5. (4S,5R)-4-[5-(1H-Indol-5-yl)-pyridin-2-yloxy]-1-aza-bicyclo[3.3.1]nonane .

6. (4S,5R)-4-[6-(1H-Indol-5-yl)-pyridazin-3-yloxy]-1-aza-bicyclo[3.3.1]nonane .

7. 5-{6-[(4S,5R)-(1-Aza-bicyclo[3.3.1]non-4-yl)oxy]-pyridazin-3-yl}-1,3-dihydro-indol-2-one.


8. A process for preparation of a compound of formula (I) comprising the steps of

i) reacting a compound of formula (IX)



(IX)

wherein Y is as defined above and z represents a leaving group with a compound of formula (X)




(X)

where R represents H , C_1-C_4alkyl or both RO represent together with the B to which they are attached a heterocyclic moiety, X is as defined above, and

ii) recovering the so obtained compound of formula (I).


9. A process for preparation of a compound of formula (I) comprising the steps of

i) reacting a compound of formula (XI)



(XI)

with a compound of formula (XII)



(XII)

wherein X and Y are as defined above and PG is a suitable protecting group,

ii) subsequent deprotection of the so obtained compound and

iii) recovering the so obtained compound of formula (I) in free base or acid addition salt form.

10. The compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, for use as a pharmaceutical.

11. The compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, for use in the prevention, treatment and / or delay of progression of psychotic and neurodegenerative disorders.

12. A pharmaceutical composition comprising a compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, in association with a pharmaceutical carrier or diluent.

13. The use of a compound of claims 1 in free base or pharmaceutically acceptable acid addition salt form, as a pharmaceutical for the prevention and the treatment of psychotic and neurodegenerative disorders.

14. The use of a compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, for the manufacture of a medicament for the prevention, treatment and / or delay of progression of psychotic and neurodegenerative disorders.

15. A method for the prevention, treatment and / or delay of progression of psychotic and neurodegenerative disorders, in a subject in need of such treatment, which comprises administering to such subject a therapeutically effective amount of a compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form.

16. A compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, for use in the prevention, treatment and / or delay of progression of a disease or condition in which nAChR α7 activation plays a role or is implicated.

17. The use of a compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form, as a pharmaceutical for the prevention, treatment and / or delay of progression of a disease or condition in which nAChR α7 activation plays a role or is implicated.

18. A method for treating or preventing a disease or condition in which nAChR α7 activation plays a role or is implicated, in a subject in need of such treatment, which comprises administering to such subject a therapeutically effective amount of compound of claim 1 in free base or pharmaceutically acceptable acid addition salt form.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2006/012023

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D471/08 A61K31/439 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2004/022556 A (NOVARTIS AG [CH]; NOVARTIS PHARMA GMBH [AT]; FEUERBACH DOMINIK [DE]; H) 18 March 2004 (2004-03-18) claim 1	1-18
Y	WO 2004/016608 A (NEUROSEARCH AS [DK]; PETERS DAN [DK]; OLSEN GUNNAR M [DK]; NIELSEN ELS) 26 February 2004 (2004-02-26) page 23; claim 1; compounds D7-D11	1-18
A	US 2005/137398 A1 (JI JIANGUO [US] ET AL) 23 June 2005 (2005-06-23) claim 1	1-18

-/--

[X] Further documents are listed in the continuation of Box C.

[X] See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
5 April 2007	24/04/2007

Name and mailing address of the ISA/	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016	Schuemacher, Anne

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2006/012023

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 2005/123732 A (NOVARTIS AG [CH]; NOVARTIS PHARMA GMBH [AT]; FEUERBACH DOMINIK [DE]; M) 29 December 2005 (2005-12-29) pages 22,25; claim 1; compounds 3G-3M, 3AO, 4AP	1-18
P,Y	WO 2006/005608 A (NOVARTIS AG [CH]; NOVARTIS PHARMA GMBH [AT]; FEUERBACH DOMINIK [DE]; R) 19 January 2006 (2006-01-19) pages 23,24; claim 1; compounds 3H-3R	1-18

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2006/012023

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. [X] Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 13, 15, 17 and 18 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2. [ ] Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. [ ] Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. [ ] As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. [ ] As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. [ ] As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. [ ] No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

[ ] The additional search fees were accompanied by the applicant's protest.

[ ] No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/EP2006/012023

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004022556 A	18-03-2004	AU 2003260486 A1	29-03-2004
		BR 0314325 A	05-07-2005
		CA 2495685 A1	18-03-2004
		CN 1678611 A	05-10-2005
		EP 1537104 A1	08-06-2005
		JP 2005539057 T	22-12-2005
		KR 20050057126 A	16-06-2005
		MX PA05002546 A	27-05-2005
		NZ 538534 A	31-03-2006
		US 2006167002 A1	27-07-2006
WO 2004016608 A	26-02-2004	AU 2003250322 A1	03-03-2004
		BR 0313153 A	28-06-2005
		CA 2493245 A1	26-02-2004
		CN 1675205 A	28-09-2005
		EP 1532144 A1	25-05-2005
		IS 7742 A	14-03-2005
		JP 2005538187 T	15-12-2005
		KR 20050055711 A	13-06-2005
		MX PA05001702 A	19-04-2005
		NZ 538058 A	30-11-2006
		US 2005245567 A1	03-11-2005
US 2005137398 A1	23-06-2005	CA 2549979 A1	21-07-2005
		EP 1735305 A2	27-12-2006
		MX PA06007179 A	04-09-2006
		WO 2005066167 A2	21-07-2005
WO 2005123732 A	29-12-2005	AR 049401 A1	26-07-2006
		AU 2005254689 A1	29-12-2005
		CA 2566155 A1	29-12-2005
		EP 1761533 A1	14-03-2007
WO 2006005608 A	19-01-2006	AU 2005261866 A1	19-01-2006
		CA 2570155 A1	19-01-2006