



(22) Date de dépôt/Filing Date: 1993/07/16

(41) Mise à la disp. pub./Open to Public Insp.: 1994/01/18

(45) Date de délivrance/Issue Date: 2007/04/17

(30) Priorités/Priorities: 1992/07/17 (JP4-212305);
1993/03/04 (JP5-067339)

(51) Cl.Int./Int.Cl. *C12N 15/18* (2006.01),
A01K 67/04 (2006.01), *A61K 38/17* (2006.01),
A61K 39/395 (2006.01), *C07K 14/475* (2006.01),
C07K 14/52 (2006.01), *C07K 16/18* (2006.01),
C12N 15/19 (2006.01), *C12N 5/10* (2006.01),
A61K 38/00 (2006.01)

(72) Inventeurs/Inventors:
MIURA, KENJU, JP;
YAMAGUCHI, NOZOMI, JP;
YAMAICHI, KOZO, JP;
KURIHARA, TATSUYA, JP;
ISHIDA, NOBUHIRO, JP;
...


(73) Propriétaire/Owner:

(54) Titre : FACTEUR DE DIFFERENCIATION MEGACARYOCYTAIRE

(54) Title: MEGAKARYOCYTE DIFFERENTIATION FACTOR

(57) Abrégé/Abstract:

A novel megakaryocyte differentiation factor, for example, consisting essentially of SEQ ID NO: 30; DNA coding for the megakaryocyte differentiation factor, an expression vector comprising the DNA, a host transformed with the expression vector, and a process for production of the megakaryocyte differentiation factor using the host. The megakaryocyte differentiation factor accelerates differentiation of megakaryocytes in the presence of IL-3, and acts as a thrombopoietin, and therefore an effective medicament to various diseases involving a decrease in platelets.



(72) **Inventeurs(suite)/Inventors(continued):** NAKAZATO, HIROSHI, JP; TSURUOKA, NOBUO, JP; IWASA, FUYUKI, JP; TSUJIMOTO, MASAFUMI, JP

(73) **Propriétaires(suite)/Owners(continued):**DAIICHI ASUBIO PHARMA CO., LTD., JP

(74) **Agent:** OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

ABSTRACT OF THE DISCLOSURE

A novel megakaryocyte differentiation factor, for example, consisting essentially of SEQ ID NO: 30; DNA coding for the megakaryocyte differentiation factor, an expression vector comprising the DNA, a host transformed with the expression vector, and a process for production of the megakaryocyte differentiation factor using the host.

The megakaryocyte differentiation factor accelerates differentiation of megakaryocytes in the presence of IL-3, and acts as a thrombopoietin, and therefore an effective medicament to various diseases involving a decrease in platelets.

MEGAKARYOCYTE DIFFERENTIATION FACTOR

BACKGROUND OF THE INVENTION

5

1. Field of the Invention

The present invention relates to a megakaryocyte differentiation factor, gene coding for the factor and a process for production thereof. The megakaryocyte differentiation factor is useful as a hemopoietic stimulating factor for megakaryocyte-platelet lineage.

10

2. Description of Related Art

15

It is well known that various hemopoietic factors inducing the growth and differentiation of blood cells are involved in a process from hemopoietic stem cells to mature blood cells.

20

Although the life time of platelet is as short as 9 to 10 days, concentration of platelets in the blood is maintained rather constant during the stationary state. Moreover, when the number of platelets is artificially reduced by one of various available method in an experimental animals, the number of the platelets recovers in the blood in a few days. From these facts, it is supposed that factors which stimulate formation of platelets are present, and so far a great effort has been made to identifiycation of the factors.

25

It is considered that at least two regulatory factors are involved in the megakaryocyte-platelet hemopoietic lineage. The first factor by itself stimulates formation of megakaryocyte colonies and is called a megakaryocyte colony stimulating factor. The second factor by itself does not have an activity to stimulate formation of megakaryocyte colonies, but in combination with the first factor, increases the number of megakaryocyte colonies and stimulates the differentiation thereof, and is called a megakaryocyte potentiator.

35

The former includes interleukin 3, and granulocyte/macrophage colony stimulating factor, and the latter includes erythropoietin, macrophage colony stimulating factor, interleukin 6, 7 and 11, LIF, and the like. Some of these factors actually exhibit in vivo effect of increasing the number of platelets or shortening the time required to recover the number of platelets (Hideaki Mizoguchi: Tanpakushitsu Kakusan Koso 36, 1195, 1991 in Japanese).

However, most of these factors exhibit a diversity of biological activities other than participation in differentiation of blood cells in various hemopoietic lineages including differentiation in megakaryocyte-platelet lineages. For example, although IL-6 and IL-11 actually exhibit in vivo thrombopoietic action, they stimulate production of acute phase protein, and in severe cases, cause cachexia. Moreover, IL-6 is accompanied with various clinical problems; for example, it is possible for IL-6 to stimulate the growth of mesangium cells in the kidney resulting in renal failure (Tadashi Matsuda et al., Tanpakushitsu Kakusan Koso, 36, 1184, 1991 in Japanese). In addition, since IL-6 does not exhibit a high blood level during a thrombocytopenic phase, it is not considered as a physiological factor.

Platelets play an important role in a hemostatic mechanism. Diseases involving decrease of platelets (Fanconi's syndrome, megakaryocytic thrombocytopenia, aplastic anemia, and the like) are clinically dangerous, and in particular hemorrhaging cannot be controlled. Therefore, it is considered that isolation and identification of a factor which stimulates production of platelets is useful to prevent the above-mentioned danger.

Currently, bone marrow transplantation is becoming a powerful therapeutic means for treating leukemia etc., and the ratio of successful cases is increasing through the use of cytokines such as

erythropoietin (EPO), granulocyte colony stimulating factor (G-CSF) etc. At present a problem in the bone marrow transplantation is a decrease in the number of platelets, and if a thrombopoietic factor is
5 available, it is expected that the ratio of successful cases will increase and a period of hospitalization will be shortened. Not only hemopoietic diseases but also thrombocytopenia in chemotherapy and radio isotopic therapy of cancers may be controlled by
10 thrombopoietin.

The present inventors, considering the various above-mentioned difficulties with known factors, carried out various research to find a factor which stimulates production of platelets and is
15 effective for treatment of patients having thrombocytopenia or insufficient platelet function, and as a result, the present inventors found a novel factor which stimulates differentiation of megakaryocytes, cloned a gene coding for said factor, constructed an expression vector, and succeeded in
20 expressing the gene to produce said factor.

SUMMARY OF THE INVENTION

Accordingly, the present invention provides a megakaryocyte differentiation factor having the
25 following properties;

(1) stimulating differentiation of megakaryocytes;

(2) exhibiting a molecular weight of 55 to 57 kD as determined by gel filtration and SDS-polyacrylamide gel electrophoresis (SDS-PAGE), and having no
30 intermolecular disulfide linkage;

(3) exhibiting an isoelectric point of 6.5 ± 0.5 ; and

(4) having at least one of the amino acid sequences shown in SEQ ID NO: 1 to 9 in the Sequence Listing.
35

The present invention also provides a gene coding for the megakaryocyte differentiation factor.

- 4 -

The present invention further provides an expression vector comprising the gene coding for the megakaryocyte differentiation factor.

5 The present invention moreover provides a host transformed with the expression vector.

The present invention still further provides a process for production of the megakaryocyte differentiation factor using said host.

BRIEF EXPLANATION OF DRAWINGS

10 Figure 1 shows an elution profile of protein (A280 —●—) and a megakaryocyte differentiation factor (acetylcholine esterase activity —o—) from a Q-Sepharose* column, wherein the elution was carried out by NaCl concentration gradient (0 to 1.0M) to
15 obtain fractions 1 to 120.

Fig. 2 shows an elution profile of protein (A280 —●—) and a megakaryocyte differentiation factor (acetylcholine esterase activity —o—) from a Phenyl-Sepharose column, wherein the elution was carried out
20 by ammonium sulfate concentration gradient (30 to 0%) and ethyleneglycol concentration gradient (0 to 50%) to obtain fractions Nos. 1 to 100, followed by 50% ethyleneglycol to obtain fraction Nos. 101 to 120.

Fig. 3 shows an elution profile of protein (A280 —●—) and a megakaryocyte differentiation factor (acetylcholine esterase activity —o—) from an S-Sepharose* column, wherein the elution was carried
25 out by NaCl concentration gradient (0 to 0.5M) to obtain fractions Nos. 1 to 100.

30 Fig. 4 shows an elution profile of protein (A280 —) and megakaryocyte differentiation factor (acetylcholine esterase activity —o—) from a Hiroad 26/60 Superdex 75 Pg column, and a result of analysis of the resulting fractions by SDS-PAGE (lower part of
35 the Figure).

Fig. 5 is an electrophoretic pattern showing a result of SDS-PAGE analysis of the purified megakaryocyte differentiation factor.

- 5 -

Fig. 6 shows a result of an isoelectric focusing of the purified megakaryocyte differentiation factor.

Fig. 7 is an electrophoretic pattern showing a result of sugar chain analysis for the purified megakaryocyte differentiation factor by SDS-PAGE, wherein the column 1 shows a result for a non-treated megakaryocyte differentiation factor, and the column 2 shows a result for an endoglycosidase F-treated megakaryocyte differentiation factor (note, the band near the 35 kD position is derived from the enzyme preparation).

Fig. 8 is a graph comparing acetylcholine esterase activity of megakaryocytes derived from mouse bone marrow cells cultured for 5 days in the presence or absence of purified megakaryocyte differentiation factor (55 kDa protein) and with or without addition of IL-6.

Fig. 9 shows a result of acetylcholine esterase staining of mouse bone marrow cells cultured for 4 days after the addition of IL-3, in the presence (B) or absence (A) of purified megakaryocyte differentiation factor.

Fig. 10 shows a result of May-Gruenwald-Giemsa's staining of mouse bone marrow cells cultured for 4 days after the addition of IL-3, in the presence (B) or absence (A) of purified megakaryocyte differentiation factor.

Fig. 11 shows detection of RNA with a DNA probe (PCR product amplified with KY100 and N1065 described in Example 2.1) wherein the RNA was extracted from cultured Bombyx moli cells infected with recombinant virus (TP055-Bm NPV) comprising a gene coding for megakaryocyte differentiation factor (Lane B), cells infected with wild type virus (B6E) (Lane A) or from non-transformed cells (Lane C).

Fig. 12 is a graph showing an expression of megakaryocyte differentiation factor (TP55) in hemolymph of Bombyx mori after reparation by Matrex*

Blue A column chromatography.

DETAILED DESCRIPTION

As starting materials for isolating the present megakaryocyte differentiation factor, human cells, such as human cancer cells, preferably human epidermoid carcinoma cells A431, particularly preferably human epidermoid carcinoma A431 grown in a protein-free medium can be mentioned.

In addition to the above-defined megakaryocyte differentiation factor, the present invention relates megakaryocyte differentiation factors obtainable from transformants such as cells or animals constructed by gene technology and having the same amino acid sequence as the above-defined megakaryocyte differentiation factor, an amino acid sequence wherein a portion of the above-defined megakaryocyte differentiation factor is deleted, an amino acid sequence wherein a portion of the above-defined megakaryocyte differentiation factor is replaced with other amino acid or amino acid sequence, or an amino acid sequence wherein one or more than one amino acid sequence is added to the above-defined megakaryocyte differentiation factor, or having an amino acid sequence including a combination of said modifications.

Moreover, the present invention relates to megakaryocyte differentiation factor having the amino acid sequence shown in SEQ ID NO: 30, an amino acid sequence wherein a portion of the amino acid sequence shown in SEQ ID NO: 30 is deleted, an amino acid sequence wherein a portion of the amino acid sequence shown in SEQ ID NO: 30 is replaced with an other amino acid or amino acid sequence, or an amino acid sequence wherein one or more than one amino acid sequence is added to the amino acid sequence shown in SEQ ID NO: 30, or having an amino acid sequence including a combination of said modifications.

The present invention also relates to genes

coding for the above-mentioned megakaryocyte differentiation factors. The present invention further relates a process for production of the megakaryocyte differentiation factors using the gene
5 by means of gene recombination technology. The gene recombination technology follows conventional procedures by using a synthetic or natural polynucleotide coding for the amino acid sequence of the native megakaryocyte differentiation factor, an
10 amino acid sequence wherein a portion of the native amino acid sequence is deleted, an amino acid sequence wherein a portion of the native amino acid sequence is replaced with other amino acid or amino acid sequence, or an amino acid sequence wherein one or more than one
15 amino acid is added to the native amino acid sequence, or coding for an amino acid sequence including a combination of said modifications, but not limited to the above.

The above-mentioned various modification can be
20 carried out by a conventional technique such as site-specific a mutagenesis.

The number of amino acids involved in the modification such as addition, deletion or replacement is not limited, but as for addition the number of
25 amino acid depends on the number of amino acids, for example, that of the functional peptide used in a hybrid protein with the megakaryocyte differentiation factor of the present invention or that of a signal peptide added to the present factor, namely depends on
30 purpose of the modification, as for deletion the number of amino acids may be designed or determined so as to maintain megakaryocyte differentiation activity and it is, for example, 1 to 30, preferably 1 to 20 or it can be that of region other than the active region
35 of the present factor, and as for replacement the number of amino acids also may be designed or determined so as to maintain megakaryocyte differentiation activity and it is, for example, 1 to

10, preferably, 1 to 5.

5 An addition or improvement of an signal sequence, choice of host-vector system, and improvement of expression regulatory region may provide efficient expression. In addition, a host may be chosen to provide a glycosylated product. Moreover, a polynucleotide coding for at least one of the amino acid sequences shown in SEQ ID NO: 1 to 9 may be used as a DNA probe for cloning a gene.

10 The present invention further provides a pharmaceutical composition comprising a megakaryocyte differentiation factor as an effective ingredient. The pharmaceutical composition is preferably used as a medicament for thrombocytopenia.

15 In addition, the present megakaryocyte differentiation factors can be used to obtain specific antibodies according to a known procedure.

Now, the present invention is explained in more detail.

20 (1) Starting material

As a starting material for obtaining a novel protein of the present invention, a culture supernatant of cells derived from human epidermoid carcinoma cell A431 (ATCC CRL 1555), rendered to be capable of growing in a protein-free medium according to Yamaguchi et al. method (Yamaguchi N. et al., Cancer Res. 50, 7008, 1991), is mentioned. This cell line was designated as Human epidermoid carcinoma SBM 330, and deposited as FERM BP-3911 with the Fermentation Research Institute Agency of Industrial Science and Technology, 1-3, Higashi 1-chome, Tsukuba-shi Ibraki-ken (307) Japan, on July 1, 1992.

(2) Assay method for megakaryocyte differentiation factor

35 To assay a megakaryocyte differentiation factor, megakaryocyte-series cell lines (for example, CMK cells or cells derived therefrom), or mouse bone marrow cells may be used. For example, activity of

- 9 -

acetylcholine esterase which is known to be specifically detected in murine megakaryocytes is carried out using mouse bone marrow cells according to the Ishibashi et al. method (Ishibashi, T. et al.,
5 Proc. Natl. Acad. Sci. USA 86, 5953, 1989). In addition, histochemical detection of megakaryocytes is carried out by subjecting cultured bone marrow cells to acetylcholine esterase staining and May-Gruenwald-Gremsa's staining and the observing the shape of the
10 stained cells.

(3) Purification of megakaryocyte
differentiation factor

The megakaryocyte differentiation factor can be purified, for example, starting from a culture
15 supernatant of A431 cells cultured in a protein-free medium, concentration by ultrafiltration, and column chromatography using, for example, Matrex* Blue A (Amicon), Q-Sepharose* (Pharmacia), phenyl-Sepharose (Pharmacia), S-Sepharose* (Pharmacia) and Hiload* 26/60
20 Superdex* 75 (Pharmacia) alone or in combination. Protein is monitored by measuring A280 nm.

(4) Determination of partial amino
acid sequence of megakaryocyte
differentiation factor

25 To determine an amino acid sequence, the megakaryocyte differentiation factor purified in the section (3) is digested into fragments with a protease such as Achromobacter Protease I (API) at 37°C for 2 hours. The resulting peptide fragments are
30 separated and recovered by reverse phase HPLC (acetonitrile gradient in 0.1% trifluoroacetic acid) using YMC-Pack AM-303 column. The peptide fragments thus obtained are subjected to a sequencer such as a gas-phase sequencer obtained from, for example,
35 Applied Biosystem. A definite purification process and detailed properties of the megakaryocyte differentiation factor are described in Example 1.

The present invention also provides gene coding

for megakaryocyte differentiation factor. The gene may be cDNA prepared from mRNA, genomic DNA, and synthetic DNA. For example, cDNA can be cloned by a polymerase chain reaction (PCR) using DNA (nucleotide) primers designed on the basis of partial amino acid sequences as shown in Example 1 of a megakaryocyte differentiation factor purified from the above-mentioned human cells, such as human epidermoid carcinoma cells, for example A431 cells. The cloning is described in detail in Examples 1 and 2.

The gene of the present invention further includes DNAs coding for protein or glycoprotein having megakaryocyte differentiation activity and hybridizing with the nucleotide sequence of SEQ ID NO: 30. Nucleotide sequence of DNA cloned in Example 2 and an amino acid sequence predicted from the nucleotide sequence are shown in SEQ ID NO: 30.

Thus once an amino acid sequence is determined, various mutated megakaryocyte differentiation factors, such as a polypeptide, wherein one or more than one amino acid is added to the native amino acid sequence or the amino acid sequence shown in SEQ ID NO: 30 and maintaining megakaryocyte differentiation factor activity, a polypeptide wherein one or more than one amino acid is deleted from the native amino acid sequence or the amino acid sequence shown in SEQ ID NO: 30 and maintaining megakaryocyte differentiation factor activity, a polypeptide wherein one or more than one amino acid is replaced with one or more than one other amino acid, in the native amino acid sequence or the amino acid sequence shown in SEQ ID NO: 30, and maintaining megakaryocyte differentiation factor activity, or a polypeptide including a combination of the above-mentioned modifications such as addition, deletion and/or replacement of amino acids maintaining megakaryocyte differentiation factor activity can be designed and produced.

According to the present invention, although the

nucleotide sequence shown in SEQ ID NO: 30 is disclosed, gene coding for the present megakaryocyte differentiation factor is not limited thereto. Once an amino acid sequence of the present native
5 megakaryocyte differentiation factor or an amino acid sequence of a mutated megakaryocyte differentiation factor is determined, according to the degeneracy of genetic code, various nucleotide sequences coding for the same amino acid sequence can be designed and
10 prepared. In this case, codons used with a high frequency in a chosen host are preferably used.

A gene coding for the present megakaryocyte differentiation factor can be obtained as cDNA according to Example 2, but the gene is not limited to
15 cDNA. Namely, once a nucleotide sequence coding for an amino acid sequence of native megakaryocyte differentiation factor is determined, a gene coding for the native megakaryocyte differentiation factor can be cloned as a cDNA according to a strategy
20 different from the strategy definitely described herein, and moreover a desired gene can be cloned from the genome of a cell producing the megakaryocyte differentiation factor.

Where a desired gene is cloned from the genome, various primer nucleotides or probe nucleotides used
25 in Example 2 can be used as probes for screening genomic DNA fragments. Moreover, other probes designed on the basis of the nucleotide sequence described in SEQ ID NO: 30 can be used. General
30 procedures for cloning a desired DNA from a genome are well known in the art (see Current Protocols In Molecular Biology, John Wiley & Sons, Chapters 5 and 6).

Gene coding for the native megakaryocyte
35 differentiation factor of the present invention can also be prepared by chemical synthesis. It is easy in the art to chemically synthesize DNA using an automated DNA synthesizer, for example, Applied

Biosystems 396 DNA/RNA synthesizer. Accordingly a person with ordinary skill in the art can easily synthesize DNA having the nucleotide sequence shown in SEQ ID NO: 30.

5 A gene coding for the present native megakaryocyte differentiation factor using codons different from native codons, and a gene coding for a mutated megakaryocyte differentiation factor can be prepared by chemical synthesis as described above.
10 Alternatively they can be obtained by site-specific mutagenesis using as a template a DNA or RNA having a nucleotide sequence shown in SEQ ID NO: 30 together with mutagenic primers (for example, see, Current Protocols In Molecular Biology, John Wiley & Sons,
15 Chapter 8).

 Once a gene coding for the present megakaryocyte differentiation factor is obtained, the gene can be used to produce a recombinant megakaryocyte differentiation factor according to a conventional
20 gene recombination technology. Namely, a DNA coding for the present megakaryocyte differentiation factor is inserted into an appropriate expression vector, the vector is introduced to appropriate host cells, the transformed host cells are cultured, and the target
25 megakaryocyte differentiation factor is recovered from the culture (cells or medium). The present megakaryocyte differentiation factor can be biochemically or chemically modified, for example, N-terminal acylated.

30 In addition, on the basis of the nucleotide sequence shown in SEQ ID NO: 30, a protein data base was searched with fasta Program (GCG package). As a result, the megakaryocyte differentiation factor belongs to the super family of serine protease
35 inhibitor. On the other hand, in human leucocyte elastase inhibitor, chicken ovalbumin Y gene product, human plasminogen activator inhibitor 2 and human squamous cell carcinoma antigen, which are similar to

the present megakaryocyte differentiation factor in their expected steric structure, and distribution of hydrophobic and hydrophilic amino acids, the N-terminal portion is not cleaved and forms a signal peptide. Accordingly, there is a possibility that in the present megakaryocyte differentiation factor, the N-terminal portion may function as a signal peptide and the megakaryocyte differentiation factor may be secreted without cleavage of the signal peptide. Further, the present megakaryocyte differentiation factor may be modified so that the first methionine is deleted and the second alanine is acetylated.

As hosts, both the prokaryote and eukaryote can be used. As prokaryote, bacteria such as Escherichia coli the genus Bacillus, for example, B. subtilis and the like can be used. As eukaryote, yeast such as the genus Saccharomyces, for example, S. cerevisiae, insect cells such as Spodoptera frugiperda cells, Cabbage looper cells, Bombyx mori cells, animal cells such as human cells, monkey cells, mouse cells and the like can be used. Moreover, insects per se, such as Bombyx mori may be used.

As expression vectors, plasmid, phage, phagemid, virus such as baculo virus, vaccinia virus or the like can be used. A promoter in an expression vector is selected depending on host used. For example, lac promoter, trp promoter and the like can be used as bacterial promoters, and adh1 promoter, ptk promoter and the like can be used as yeast promoters. On the other hand, baculo virus polyhedrin promoter can be used as insect promoter, and Simian virus 40 early or late promoter can be used for animal cells.

Transformation of a host with an expression vector can be carried out according to conventional procedures well known in the art, and these procedures are described in, for example, Current Protocols in Molecular Biology, John Wiley & Sons. Culturing of a transformant also can be carried out according to a

conventional procedure.

Purification of a megakaryocyte differentiation factor from a culture or an insect body can be carried out according to conventional procedures used for isolation and purification of a protein, for example, ultrafiltration, various types of column chromatography such as Q-Sepharose column chromatography and the like.

EXAMPLES

Next, the present invention is further explained by Examples.

Example 1. Purification of megakaryocyte differentiation factor

(1) Culturing of A431 cells

Frozen SBM 330 cells conditioned in a protein-free medium from A431 cells were thawed, and cultured in a primary medium (Ham's F12 medium containing 10% fetal bovine serum). Namely, the cells were plated in 10 T flasks having a culture area of 150 cm², and cultured to confluence at 37°C in the presence of 5% CO₂. Next, the cells were peeled off with a 0.25% trypsin solution (Chiba Kessei) and subcultured in 10 roller bottles having a culture area of 850 cm² at 37°C and 0.5 rpm for about 3 days to recover 1.8 x 10⁹ cells. The cultured cells were attached to a ceramic core (S-451) of an Opti-cell incubator (Charles River Inc. Wilmington, MA) and perfusion culturing was started using 10L of a primary medium.

The perfusion culture was carried out at 37°C with supplying oxygen at 150 mmHg. The primary medium was replaced with a protein-free medium as follow. Namely, after culturing for about 7 days in a primary medium, a protein-free medium was supplied to the culture at a rate of 20L/day, and simultaneously the culture supernatant was recovered from the culture at the same rate. As a result, the primary medium containing serum was substantially completely replaced

- 15 -

with the protein-free medium by supplying about 100L of the protein-free medium. Thereafter, cell culture supernatant was continuously recovered to obtain 1000L of cell culture supernatant. A part (about 300L) of the cell culture supernatant thus obtained was concentrated to 2L using an ultrafiltration membrane (Millipore, Bedford, MA; MW 10,000 cut) and the concentrate was dialyzed against 20 mM Tris/HCl buffer (pH 7.4).

10 (2) Assay of megakaryocyte
differentiation factor using mouse
bone marrow cells

Myeloid cells were pushed out of the femur of female BDF₁ mouse and suspended in an α -MEM medium (Flow Laboratories, Inc. McLean, VA, USA). Percoll^{*} layers having different densities (Pharmacia LKB Biotechnology, Tokyo) were overlaid, and the bone marrow cell suspension was put thereon, followed by centrifugation at 400 x g for 20 minutes. Mononuclear cells collected at the interface of a layer having a density of 1.07 g/ml and a layer having a density of 1.08 g/ml were recovered and washed once with α -MEM containing 10% FBS, and resuspended in the same medium containing 0.5 mM diisopropylfluorophosphate. The suspension was then put into a plastic cell culture dish (Corning, NY, USA) and cultured at 37°C in 5% carbon dioxide and 95% air for 2 hours. During the culturing, at one hour from the start of culturing, the cell culture dish was replaced with a new one. After the culturing, cells were washed with 10% FBS/ α -MEM three times.

The non-adherent bone marrow mononuclear cells thus obtained were suspended in 10% FBS/1% BSA/0.1 mM 2-mercaptoethanol/ α -MEM, and plated in a 96-well microplate (Corning) in an amount of 5×10^4 cell/well. If necessary, to a test sample were added 25 U/ml mouse recombinant IL-3 (Genzyme Corporation, Cambridge, MA, USA) and 1 to 2 μ g/ml

* Trade Mark

anti-IL-6 antibody (Boehringer Mannheim, Mannheim, FRG). Where anti-IL-6 antibody was added, the test sample and the antibody were preincubated at 37°C for an hour prior to seeding the cells.

5 The culturing was carried out at 37°C in 5% CO₂ - 5% O₂ - 90% N₂ for 4 to 5 days. After culturing cells in each well of the microplate they were washed twice with PBS, and lysed with 180 µl of 0.2% (w/v) Triton X-100, 1 mM EDTA, 0.12M NaCl, 50 mM HEPES
10 (pH 7.5), and 20 µl of a substrate, 5.6 mM acetylthiocholine iodide, was added thereon. After culturing with shaking at a room temperature for an hour, 20 µl of the solution was transferred to a microplate for fluorescent assay (Dynatech Micro FLUOR
15 "B" Plate).

 To the microplate were added 20 µl of 0.4 mM 7-diethylamino-3-(4'-maleimidylphenyl)-4-methylcoumarine in acetonitrile and 160 µl of 0.2% (w/v) Triton X-100, 1 mM EDTA, 50 mM sodium acetate
20 (pH 5.0), and fluorescent emission was measured by a fluorometer (excitation 365 nm, emission 450 nm).

 For acetylcholine esterase staining, the cells were centrifuged by a Cytospin to adhere the cells onto a slide glass, and fixed with 5%
25 glutaraldehyde, 10 mM phosphate buffer (pH 6.7) for 15 minutes, and as a substrate acetylthiocholine was used according to Mizoguchi's method (Method for Culturing Hemopoietic Stem Cells, Chugai Igaku, 1986, ed. Y. Miura, pp82-88). Namely, after fixing the
30 cells, the slide glass was washed with 0.1M phosphate buffer (pH 6.0), and to each slide glass was layered a mixture of 1.5 ml of 0.67 mg/ml acetylthiocholine iodide, 0.1M phosphate buffer (pH 6.0), 0.2 ml of 30 mM CuSO₄, 0.2 ml of 5 mM potassium ferricyanide and
35 0.1 ml of 0.1M sodium citrate. The slide glass was incubated at a room temperature for 4 hours, and washed with water. May-Gruenwald-Giemsa's staining is well known in hematology and was carried out using

reagents available from E. Merck (Darmstadt, FRG) wherein May-Gruenwald's staining was carried out for 4 minutes and Giemsa's staining was carried out for 10 minutes.

5 (3) Purification of megakaryocyte
 differentiation factor

 A culture supernatant concentrate of A431
cells was dialyzed and centrifuged to obtain a
supernatant which was then applied to a Matrex Blue
10 A Column equilibrated with 20 mM Tris/HCl buffer
(pH 7.4), and after washing the column with the same
buffer, a bound fraction was eluted with the same
buffer containing 2M NaCl. Megakaryocyte
differentiation activity detected by the above-
15 mentioned method was found in the bound fraction.
Therefore, the bound fraction was dialyzed against
20 mM Tris/HCl buffer (pH 7.4) and applied to a Q-
Sephadex column equilibrated with the same buffer.
The column was thoroughly washed and a megakaryocyte
20 differentiation factor was eluted by NaCl gradient
(see Fig. 1). This factor was eluted near the
position of 0.3 to 0.5M NaCl.

 The active fraction obtained from Q-
Sephadex was made ammonium sulfate 30% saturation,
25 and was applied to a phenyl Sephadex column
equilibrated with 20 mM Tris/HCl buffer (pH 7.4)
containing 30% saturation of ammonium sulfate. The
megakaryocyte differentiation factor was eluted by
simultaneously forming concentration gradients of
30 ammonium sulfate (30% to 0%) and ethyleneglycol (0 to
50%) (see Fig. 2). The megakaryocyte differentiation
factor measured in the presence of anti-IL-6 antibody
was observed over several fractions in the beginning
of the concentration gradient formation.

35 The fractions thus obtained were combined,
thoroughly dialyzed against 50 mM MES/NaOH buffer
(pH 6.0), and applied to an S-Sephadex column
equilibrated with the same buffer. A bound fraction

- 18 -

was eluted by a 0 to 0.5M NaCl concentration gradient (see Fig. 3). Although activity was widely distributed, relatively high activity was found in the beginning of the concentration gradient formation.

5 The fractions obtained from S-Sepharose were applied to a Hilord^{*} 26/60 Super Dex 75 (Pharmacia) column for gel filtration. The column used had be previously equilibrated with the same buffer, and elution was carried out with the same buffer (see Fig. 4).

10 Activity of megakaryocyte differentiation activity was eluted near the position corresponding to a molecular weight of 55 to 57 kDa.

According to the above-mentioned steps, about 80 µg of a fraction showing two bands near 55 to 57 kDa as analyzed by SDS-PAGE was obtained from 300L of a culture supernatant of A431 derived cells (see Fig. 5). The two bands were correlated with the activity (Fig. 1). Accordingly, it was concluded that the two bands observed in said fraction correspond to a desired megakaryocyte differentiation factor.

(4) Properties of megakaryocyte differentiation factor

The present megakaryocyte differentiation factor has the following properties.

25 1) Molecular weight: about 55 kDa (gel filtration and SDS-PAGE) (Figs. 4 and 5)

The present factor exhibits two bands in SDS-PAGE, and there is no difference in mobility between reducing condition and a non-reducing condition. Therefore the factor does not have intermolecular disulfide linkage.

2) Isoelectric point: 6.5 ± 0.5 (Fig. 6)

A several bands are observed in the above-mentioned range.

35 3) The above-mentioned heterogeneity of the present factor can be explained as heterogeneity in a sugar chain structure of glycoprotein. Namely, where the present factor is treated with

* Trade Mark

endoglycosidase F which is an asparagine linked sugar removing enzyme, the molecular weight of the present factor decreased to a molecular weight of about 40 kDa, and the heterogeneity also decreased (Fig. 7), in SDS-PAGE. In addition, where a fraction exhibiting a single band and a fraction exhibiting two bands were digested with API and a peptide map was prepared by fractionation by a reversed phase HPLC, the difference was not observed between the two fractions.

4) The present factor contains at least one of the amino acid sequences shown in SEQ ID NO: 1 to 9.

5) Biological activity

Where mouse bone marrow cells were cultured in the presence of a purified megakaryocyte differentiation factor and IL-3, the proliferation and differentiation of megakaryocytes are observed (Figs. 8, 9 and 10). Fig. 8 shows the result of measurement for acetylcholine esterase activity of megakaryocytes; Fig. 9 shows the result of acetylcholine esterase staining of the cultured cells ($\times 20$); and Fig. 10 shows the result of May-Gruenwald-Giemsa's staining of the cultured cells ($\times 100$). In both the Figs. 9 and 10, it is seen that megakaryocytes increased in the presence of a megakaryocyte differentiation factor (B) in comparison with in the absence of the same (A).

(5) Structure of megakaryocyte differentiation factor

To characterize the structure of a purified megakaryocyte differentiation factor, the factor was digested with API and the structures of the resulting fragments were determined. After the digestion of the factor with API, each fragment was recovered by reversed phase HPLC, and the structures for appropriate fractions were determined. As a result, the peptide fragments had amino acid sequences shown in SEQ ID NO: 1 to 9.

- 20 -

Example 2. Determination of Structure of CDNA
for megakaryocyte differentiation factor

1. Analysis of cDNA nucleotide sequence of
megakaryocyte differentiation factor by PCR (1)

5 Oligonucleotides NI065 (SEQ ID NO: 10;
corresponding to 449-486 of SEQ ID NO: 30) and NI067
(SEQ ID NO: 11; corresponding to 1049-1080 of SEQ ID
NO: 30) were synthesized by designing nucleotide
10 sequences on the basis of the amino acid sequences
shown in SEQ ID NO: 3 and 4 respectively.

 Total RNA was purified from A431 cells using
ISOGEN* (Wako Pure Chemical) according to the
manufacturer's instructions. RNA having poly A was
purified from the total RNA, and a reaction was
15 carried out using a 3'-RACE Kit (Gibco BRL). Namely,
the above-mentioned oligomer NI065 and oligomer 3'-
RACE adaptor primer (SEQ ID NO: 12) attached to the
3'-RACE* Kit (Gibco BRL) were used to carry out
polymerase chain reaction (PCR) according to
20 instructions included with the kit.

 The reaction product was then subjected to
second PCR using the primer NI067 and the oligomer 3'-
RACE adapter primer included in the 3'-RACE Kit (Gibco
BRL) to obtain a DNA fragment of about 900 base pairs.
25 Next, using a direct nucleotide sequence determination
method for a PCR product, according to U. Gyllensten
et. al., Proc. Natl. Acad. Sci. USA 85: 7652 (1988),
the DNA fragment of about 900 base pairs was directly
used as a reaction substrate to determine a nucleotide
30 sequence of a portion representing protein and a
portion downstream of the protein portion using a Taqu
Dye Deoxy Terminator Cycle Sequencing kit available
from Applied Biosystem and a fluorescent nucleotide
sequencer (Applied Biosystem, Type 370A) according to
35 a manufacturers instruction. As a result, a sequence
from nucleotide number 1081 to 1950 of SEQ ID NO: 30
was shown.

On the basis of this sequence, oligomer

KY100 (SEQ ID NO: 13; corresponding to 1255-1236 of
SEQ ID NO: 30) was synthesized. The reaction product
obtained by the PCR using NI065 and the oligomer 3'-
RACE adapter primer attached to the 3'-RACE Kit (Gibco
5 BRL) was used as a reaction substrate to carry out a
further PCR using NI065 and KY100. As a result, a DNA
fragment of 807 base pairs was obtained.

This DNA fragment of 807 base pairs was
directly used as a reaction substrate to determine its
10 nucleotide sequence using Taq Dye Deoxy Terminator
Cycle Sequencing kit available from Applied Biosystem
and a fluorescent nucleotide sequencer according to a
manufactures instruction. As a result, a nucleotide
sequence from nucleotide number 487 to 1080 of SEQ ID
15 NO: 30 was shown. On the basis of this sequence,
oligomers NI073 (SEQ ID NO: 14; corresponding to 864-
886 of SEQ ID NO: 30), NI074 (SEQ ID NO: 15;
corresponding to 1012 to 992 of SEQ ID NO: 30), and
NI075 (SEQ ID No: 16; corresponding to 802-782 of SEQ
20 ID NO: 30) were synthesized.

2. Analysis of cDNA nucleotide sequence of
megakaryocyte differentiation factor by PCR (2)

A. Preparation of mRNA from megakaryocyte
differentiation factor expressing cell line (A431)

25 From 1.1g of frozen cells of human
epidermoid carcinoma cell line (A431), 25 µg of mRNA
was extracted and purified using an RNA extraction kit
and an mRNA purification kit available from Pharmacia-
LKB.

30 B. Preparation of DNA phage library from
megakaryocyte differentiation factor expressing cell
line (A431)

(1) Synthesis of cDNA

35 From 5 µg of the mRNA derived from
A431, cDNA was synthesized using a cDNA synthesis kit
Time Saver available from Pharmacia-LKB. First 5 µg
of mRNA dissolved in 20 µl of distilled water treated
with diethylpyrocarbonate (DEPC) was heated at 65°C

for 10 minutes and cooled on ice. 11 μ l of a first strand reaction mixture, 1 μ l of DTT solution and 1 μ l of 130 U/ml NotI/oligomer 18 primer solution (Pharmacia-LKB) were added thereto, and the mixture
5 was incubated at 37°C for an hour.

The reaction mixture was added to a second strand reaction mixture, and the mixture was incubated at 12°C for 30 minutes and 22°C for an hour, and heated at 65°C for 10 minutes. 100 μ l of a
10 mixture of phenol/chloroform/isoamylalcohol (25:24:1; abbreviated as PC hereinafter) was added thereto, and the mixture was vigorously stirred, centrifuged at 14,000 $\times$ g for one minute to obtain a supernatant, which was then fractionated by a Sephacryl S-400 spin
15 column (Pharmacia LKB) to obtain 100 μ l of cDNA solution.

(2) Addition of EcoRI adaptor

To 100 μ l of the cDNA solution were added 5 μ l of 10 U/ml EcoRI adaptor (Pharmacia
20 LKB), 30 μ l of polyethylene glycol buffer, 1 μ l of 1/5 diluted ATP solution and 1 μ l of T4 DNA ligase, and the mixture was incubated at 37°C for an hour. After heating at 65°C for 10 minutes, 1.5 μ l of ATP solution and 1 μ l of T4 polynucleotide kinase were added
25 thereto and the mixture was incubated at 37°C for 30 minutes. After heating at 65°C for 10 minutes, 2 μ l of 20 U/ μ l NotI was added to the mixture, which was then incubated at 37°C for an hour. 150 μ l of PC was added to the mixture, which was then vigorously
30 stirred and centrifuged at 14,000 $\times$ g for a minute, and the supernatant was fractionated on Sephadex S-400 spin column to obtain 150 μ l of cDNA solution.

(3) Incorporation of cDNA into phage vector and in vitro packaging

35 After digesting with EcoRI and NotI, to 15 μ l of the cDNA solution were added 2 μ g of dephosphorylated λ gt11D (Pharmacia LKB). After ethanol precipitation, the precipitate was dissolved

- 23 -

in 8 μ l of ligase buffer solution, and 1 μ l of 1/75-diluted ATP solution and 1 μ l of T4 DNA ligase were added to the solution, which was then incubated at 16°C for 30 minutes and stored on ice.

5 An in vitro packaging reaction was carried out using Gigapack II Gold* (Stratagene), and a library of 3.22×10^6 pfu of recombinant phages was obtained from said 3 ligase reaction products. Said library was amplified in an E. coli Y1090 r⁻ host to
10 obtain 6.0×10^{10} pfu/ml of an A431 phage library stock.

C. Identification and isolation of cDNA fragment for megakaryocyte differentiation factor by PCR

15 (1) Amplification of cDNA insert fragment in A431 phage library by PCR

10 μ l of a stock solution of 6.0×10^{10} pfu/ml A431 phage library (corresponding to 6.0×10^8 pfu) was used as a template DNA for PCR
20 reaction, and 5 μ l of 10 $\times$ PCR buffer, 8 μ l of 1.25 mM 4 dNTPs, 2 μ l of 1 OD/ml λ gt11-forward primer (λ gt 11F) (SEQ ID NO: 17), 2 μ l of 1 OD/ml λ gt11-reverse primer (λ gt 11R) (SEQ ID NO: 18), and 1 μ l of 5 U/ μ l Taq DNA polymerase (Perkin Elmer Cetus) were added,
25 and the total volume of the mixture was made to be 50 μ l with DEPC-treated distilled water. 30 reaction cycles of 93°C for a minute, 55°C for 2 minutes and 72°C for 3 minutes were carried out, and the reaction mixture was incubated at 72°C for 10 minutes. As a
30 result of an analysis by 1% Agarose gel electrophoresis, a smeary pattern ranging 0.8 to 6 kb was shown.

(2) PCR analysis using cDNA insert DNA amplification fragment mixture as a template and using
35 TP7 (SEQ ID NO: 20; corresponding to 683-703 of SEQ ID NO: 30), /TP10, TP7/TP6 (SEQ ID NO: 19; corresponding to 1036-1001 of SEQ ID NO: 30), TP8 (SEQ ID NO: 21; corresponding to 941-964 of SEQ ID NO: 30), TP10 (SEQ

- 24 -

ID NO: 22; corresponding to 1036-986 of SEQ ID NO: 30) and TP8/TP6 as primers

1 μ l of a 1/5000-diluted solution of the above-mentioned PCR reaction product was used as a template DNA for PCR reaction, and 5 μ l of 10 x PCR buffer, 8 μ l of 1.25 mM 4 dNTPs, 2 μ l each of 1 OD/ml primers in combination as described hereinafter and 1 μ l of Perfect Match (Stratagene) were added, and a total volume of the mixture was made to 49 μ l with DEPC-treated distilled water.

The reaction mixture was heated at 95°C for 5 minutes and 60°C for 5 minutes, 1 μ l of 5 U/ μ l Taq DNA polymerase (Perkin Elmer Cetus) was added thereon, and 30 cycles of 94°C for a minute, 60°C for 2 minutes and 72°C for 3 minutes were carried out, followed by an incubation at 72°C for 10 minutes. As primers, TP7/TP10, TP7/TP6, TP8/TP10, and TP8/TP6 were used.

As a result of an analysis of the PCR reaction product by 2% agarose gel electrophoresis, bands of 354 bp, 354 bp, 96 bp and 96 bp were obtained corresponding to primers respectively.

(3) Analysis for primary sequence of PCR amplification products (354 bp) obtained by using primers TP7/TP10 and TP7/TP6

The bands of said PCR amplification products (354 bp) obtained using primers TP7/TP10 and TP7/TP6 were excised from the 2% agarose gel after the electrophoresis, and to the excised agarose gel pieces was added 50 μ l of DEPC-treated distilled water, and the mixture was heated at 45°C for 30 minutes. To 2 μ l of this solution as a template DNA, were added 5 μ l of 10 x PCR buffer, 8 μ l of 1.25 mM 4 dNTPs, 2 μ l each of 1 OD/ml primers in combination and 1 μ l of Perfect Match* (Stratagene), and a total volume of the reaction mixture was made to 49 μ l with DEPC-treated distilled water. The mixture

* Trade Mark

was heated at 95°C for 5 minutes and 60°C for 5 minutes, 1 µl of 5 U/µl Taq DNA polymerase (Perkin Elmer Cetus) was added, and 30 reaction cycles of 94°C for a minutes, 60°C for 2 minutes and 72°C 3 minutes were carried out, followed by incubation at 72°C for 10 minutes. As primes for the above-mentioned PCR reaction, TP7/TP10 and TP7/TP6 were used. Bands of the PCR reaction products (each 354 bp) were excised from 2% agarose gel of the electrophoresis, extracted and purified, and the product was inserted into pCR II (Invitrogen), which was then used to transform *E. coli* INVαF' (Invitrogen). Plasmid DNA was extracted and purified, and it was confirmed by EcoRI digestion that DNA fragment of 354 bp had been inserted.

Primary sequence of the DNA insert fragment was analyzed using M13 forward primer (M13F) (SEQ ID NO: 23) and M13 reverse primer (M13R) (SEQ ID NO: 24) (Applied Biosystem's automated sequencer, Model 370A). As a result, a 296 bp sequence corresponding to the nucleotide number 704 to 999 of SEQ ID NO: 30 was found, and this sequence contained C-terminal 3 amino acids (XRK; but ERK from DNA nucleotide sequence) of SEQ ID NO: 9 corresponding to the downstream portion of primer TP7, N-terminal 5 amino acids (ADLSG) of SEQ ID NO: 6 corresponding to the upstream portion of primer TP6, and 8 amino acids (YLRALGLK) of SEQ ID NO: 5 corresponding to primer TP8, revealing that the PCR reaction products (each 354 bp) is a part of cDNA coding for megakaryocyte differentiation factor.

3. Screening of cDNA coding for megakaryocyte differentiation factor

A. Preparation of cDNA plasmid library from megakaryocyte differentiation factor expressing cell line (A431)

(1) Synthesis of first strand cDNA

From 5 µg of mRNA derived from A431 cell line, cDNA was synthesized using a Super

Script plasmid system available from GIBCO. First
2 μ l of NotI primer adaptor was added to 5 μ g of mRNA
dissolved in 5 μ l of diethylpyrocarbonate (DEPC)-
treated distilled water, and the mixture was heated at
5 70°C for 10 minutes and cooled on ice. 4 μ l of
5 x first strand buffer, 2 μ l of 0.1 M DTT solution,
1 μ l of 10 mM 4dNTPs and 1 μ l of DEPC-treated
distilled water were added thereto, and the mixture
was incubated at 37°C for 2 minutes. 5 μ l of a Super
10 Script reverse transcriptase was added to the reaction
mixture, which was then incubated at 37°C for one hour
and then put on ice to stop the reaction.

(2) Synthesis of second strand cDNA

To 18 μ l of 20 μ l reaction mixture
15 for the first strand cDNA synthesis were added 93 μ l
of DEPC-treated distilled water, 30 μ l of 5 x second
strand buffer, 3 μ l of 10 mM 4 NTPs, 1 μ l of 10 U/ μ l
E. coli DNA ligase, 4 μ l of 10 U/ μ l E. coli DNA
polymerase and 1 μ l of 2 U/ μ l E. coli RNase H, and the
20 mixture was incubated at 16°C for 2 hours. 2 μ l
(10 U) of T4 DNA polymerase was added to the reaction
mixture, which was then incubated at 16°C for
5 minutes.

The reaction mixture was put on
25 ice, and after adding 10 μ l of 0.5M EDTA and 150 μ l of
PC thereto, was vigorously stirred and centrifuged at
14,000 x g for 10 minutes, and 140 μ l of the
supernatant was transferred to a fresh centrifuge
tube. 70 μ l of 7.5M ammonium acetate and 0.5 ml of
30 ethanol were added to the supernatant, which was then
allowed to stand at -80°C for 30 minutes. The mixture
was centrifuged at 14,000 x g for 10 minutes, and
after removing the supernatant, the precipitate was
washed with 0.5 ml of 70% ethanol and dried under a
35 reduced pressure.

(3) Addition of BstXI adapter

The above-mentioned cDNA
precipitate was dissolved in 25 μ l of DEPC-treated

distilled water, 10 μ l of 5 $\times$ T4 DNA ligase buffer, 10 μ l of BstXI adapter (Invitrogen) and 5 μ l of T4 DNA ligase were added to the solution, which were then incubated at 16°C for 16 hours. 50 μ l of PC was added to the mixture, which was then vigorously stirred, and centrifuged at 14,000 $\times$ g for 5 minutes. 45 μ l of the supernatant was transferred to a fresh centrifuge tube. 25 μ l of 7.5M ammonium acetate and 150 μ l of ethanol were added to the tube, which was stirred and allowed to stand at -80°C for 30 minutes. After centrifuging at 14,000 $\times$ g for 10 minutes to remove supernatant, the precipitate was washed with 0.5 ml of 70% ethanol and dried under a reduced pressure.

(4) NotI digestion

The above-mentioned cDNA precipitate was dissolved in 41 μ l of DEPC-treated distilled water, and 5 μ l of REAct 7 buffer and 4 μ l of NotI were added to the solution, which was incubated at 37°C for 2 hours. 50 μ l of PC was added to the mixture, which was then vigorously stirred and centrifuged at 14,000 $\times$ g for 10 minutes, and 45 μ l of the supernatant was transferred into a centrifuge tube.

(5) Elimination of adapter and size fractionation of partial cDNA

The above-mentioned cDNA solution was fractionated using a Quick Spin Column Linker 5 (Boehringer Mannheim). 50 μ l of 40 μ g/ μ l cDNA was obtained.

(6) Incorporation of cDNA into phagevector and transformation of E. coli

To 37.5 μ l of the above-mentioned cDNA solution were added 12.5 μ l of pCR/CMV (Invitrogen) vector (29 μ g/ μ l) digested with Not I and BstXI, and further added were 400 μ l of Takara Ligation kit A solution and 50 μ l of B solution, and the mixture was incubated at 16°C for 30 minutes, and 1 ml of Max Efficiency DH5 α competent cells (BRL) were

- 28 -

transformed to obtain 71,550 recombinant clones. All colonies were collected from plate (2.86×10^7 cells/ml) and stored at -80°C in the presence of 20% glycerol.

5 B. Screening of megakaryocyte differentiation factor cDNA by colony hybridization

 Using the cDNA plasmid library derived from A431 cell line, a total of 227,000 (3700/plate) colonies were formed on 60 plates of 9 cm diameter, and the colonies were replicated to nitrocellulose filters. A probe was prepared by carrying out PCR (as described hereinbefore) using primer NI 067 and 3'-RACE adaptor (GIBCO BRL) to obtain a 900 bp PCR product, digesting the PCR product with Bam HI to
10 obtain two DNA fragments (0.5 kb and 0.4 kb) and nick-translating the DNA fragments with $[\alpha\text{-}^{32}\text{P}]\text{dCTP}$.
15

 For the colony hybridization, the filter was incubated in $5 \times \text{SSC}$, 25 mM phosphate buffer (pH 7.4), $5 \times \text{Denhardt's}$ solution, 1% SDS, 100 $\mu\text{g/ml}$ heat denatured salmon sperm DNA and 50%
20 formamide at 42°C for 18 hours, and washed with $5 \times \text{SSC}$, 0.1% SDS at 40°C for 20 minutes and 45°C for 20 minutes. Detection was carried out by exposing BAS 2000 (Fuji Film*) for 18 hours.

25 First, second and third screening was carried out to obtain 4 clones, i.e., TP290, TP308, TP310 and TP317. The length of insert cDNA was 1.2 kb, 1.1 kb, 1.2 kb and 1.2 kb respectively. The TP290, TP310 and TP317 cover a region downstream from
30 the nucleotide number 685 of SEQ ID NO: 30.

4. Analysis of cDNA nucleotide sequence coding for megakaryocyte differentiation factor by PCR (3)

A. Preparation of mRNA from megakaryocyte differentiation factor expressing cell line (HPC-Y11)

35 From 1.1g of frozen cells of human pancreatic cancer cell line (HPC-Y11), 50 μg of mRNA was extracted and purified using an RNA extraction kit and mRNA purification kit available from Pharmacia-

LKB.

B. Preparation of cDNA phage library from megakaryocyte differentiation factor expressing cell line (HPC-Y11)

5 (1) Synthesis of cDNA

From 5 µg of mRNA derived from HPC-Y11, cDNA was synthesized using a Pharmacia-LKB's Time Saver cDNA synthesis kit. First, 5 µg of mRNA was dissolved 20 µl of diethylpyrocarbonate (DEPC)-
10 treated distilled water, and the solution was heated at 65°C for 10 minutes and cooled on ice. 11 µl of first strand reaction mixture, 1 µl of DTT solution and 1 µl of NotI/oligomer 18 primer solution were added to the mixture, which was then incubated at 37°C
15 for an hour.

100 µl of second strand reaction mixture was added to the mixture, which was then incubated at 12°C for 30 minutes and at 22°C for an hour, and heated at 65°C for 10 minutes. 100 µl of
20 phenol- chloroform-isoamyl alcohol (25:24:1, abbreviated as PC) was added to the mixture, which was then vigorously stirred and centrifuged at 14,000 × g for a minute, and the supernatant was fractionated using a Sephacryl S-400 spin column (Pharmacia-LKB) to
25 obtain 100 µl of cDNA solution.

(2) Addition of EcoRI adaptor

To 100 µl of the cDNA solution, were added 5 µl of EcoRI adapter (Pharmacia-LKB), 30 µl of polyethylene glycol, 1 µl of ATP solution and
30 1 µl of T4 DNA ligase, and the mixture was incubated at 37°C for an hour. After heating at 65°C for 10 minutes, 1.5 µl of ATP solution and 1 µl of T4 polynucleotide kinase were added to the mixture, which was then incubated at 37°C for 30 minutes. After
35 heating at 65°C for 10 minutes, 12 µl of Not I was added to the mixture, which was then incubated at 37°C for an hour. 150 µl of PC was added to the mixture, which was then vigorously stirred and centrifuged at

14,000 × g for a minute, and the supernatant was fractionated using a Sephacryl S-400 spin column to obtain 150 µl of cDNA solution.

5 (3) Incorporation of cDNA into phage vector and in vitro packaging

To 15 µl of the cDNA solution, was added 2 µg of λgt 11D (Pharmacia-LKB) which had been digested with EcoRI and Not I and dephosphorylated, and after ethanol precipitation, the precipitate was dissolved in 8 µl of ligase buffer. 1 µl of 1/75-diluted ATP solution and 1 µl of T4 DNA ligase were added to the mixture, which was then incubated at 16°C for 30 minutes and stored on ice. An in vitro packaging reaction was carried out using a Giga Pack II Gold (Stratagene), and 5.34×10^6 pfu of recombinant phages were obtained from the above-mentioned 3 ligase reaction products. The library was amplified in E. coli Y1090 r^{-1} host to obtain a stock of 1.7×10^{11} pfu/ml HPC-Y11 phage library.

20 C. Identification and isolation of 5'-portion of megakaryocyte differentiation factor cDNA

(1) Amplification by PCR of HPC-Y11 phage library cDNA insert DNA fragment primer NI074 upstream portion.

25 To 1 µl (corresponding to 1.7×10^9 pfu) of 6.0×10^{10} pfu/ml HPC-Y11 phage library stock solution as a template DNA for PCR reaction, were added 5 µl of 10 × PCR buffer, 8 µl of 1.25 mM 4 dNTPs, 1 µl of 10 OD/ml λgt11-forward F1 primer (SEQ ID NO: 25), 1 µl of 50 D/ml NI074 primer and 1 µl of Perfect Match (Stratagene), and a total volume was made to be 49 µl with DEPC-treated distilled water.

35 After heating the reaction mixture at 95°C for 5 minutes and 60°C for 5 minutes, 1 µl of 5 U/µl Taq DNA polymerase (Perkin Elmer Cetus) was added thereto, and 35 reaction cycles of 94°C for a minute, 60°C for a minute and 72°C for 2 minutes were

carried out, followed by incubation at 72°C for 10 minutes. A result of analysis by 2% agarose gel electrophoresis showed a smeary pattern ranging from 0.3 to 6 kb.

5 (2) PCR analysis using a template which is a PCR amplification fragment mixture prepared by using λ gt11F1/NI074 primer, and using as primers λ gt11F2 (SEQ ID NO: 26)/NI075, λ gt11F2/TP12 (SEQ ID NO: 28; corresponding to 703-683 of SEQ ID NO: 30),
10 λ gt11F2/TP11 (SEQ ID NO: 27; corresponding to 619-599 of SEQ ID NO: 30), λ gt11F2/TP13 (SEQ ID NO: 29; corresponding to 595-575 of SEQ ID NO: 30), TP7/NI074, TP7/NI075, and NI073/NI074

15 1 μ l of a 1/100 diluted solution of the above-mentioned PCR reaction product was used for a template DNA for PCR reaction, and 5 μ l of 10 $\times$ PCR buffer, 8 μ l of 1.25 mM 4 dNTPs, 0.5 μ l each of 10 OD/ml primers in combination as described hereinafter, and 1 μ l of Perfect Match (Stratagene)
20 were added thereto, and a total of the reaction mixture was made to be 49 μ l with DEPC-treated distilled water. The reaction mixture was heated at 95°C for 5 minutes and at 60°C for 5 minutes, 1 μ l of 5 U/ μ l Taq DNA polymerase (Perkin Elmer Cetus) was
25 added thereto, and 35 reaction cycles of 94°C for a minute, 60°C for 2 minutes and 72°C for 2 minutes were carried out, followed by an incubation at 72°C for 10 minutes.

30 As primers, λ gt11F2/NI075, λ gt11F2/TP12, λ gt11F2/TP11, λ gt11F2/TP13, TP7/NI074, TP7/NI075, and NI073/NI074 were used. As a result of 2% agarose gel electrophoresis, bands of 969 bp, 870 bp, 786 bp, 762 bp, 330 bp, 120 bp and 149 bp were obtained corresponding to the primers.

35 (3) PCR analysis and primary sequence analysis using λ gt11F/TP11 and λ gt11F/TP13 primer, for PCR amplification product (969 bp) prepared using F2/NI075 primers

0.5 μ l of the above mentioned PCR reaction product (969 bp) prepared by using λ gt11F2/NI075 as primers was used as a template DNA for PCR reaction, and 5 μ l of 10 $\times$ PCR buffer, 8 μ l of 1.25 mM 4 dNTPs, 1 μ l of 10 OD/ml λ gt11F primer, 1 μ l of 10 OD/ml TP11 primers or 1 μ l of 10 OD/ml TP13 primers, and 1 μ l of Perfect Match (Stratagene) were added, and the total volume was made to be 49 μ l with DEPC-treated distilled water.

The reaction mixture was heated at 95°C for 5 minutes and 60°C for 5 minutes, 1 μ l of 5 U/ μ l Taq DNA polymerase (Perkin Elmer Setus) was added thereto, and 35 reaction cycles of 94°C for a minute, 60°C for 2 minutes and 72°C for 2 minutes were carried out, followed by an incubation at 72°C for 10 minutes.

Bands of PCR reaction products (678 bp and 654 bp, respectively) were excised from 2% agarose gel after electrophoresis, extracted and purified, and inserted into pCRII (Invitrogen), which was then used to transform E. coli IN V α F' (Introgen). Plasmid DNA was extracted and purified from the transformant, and digested with EcoRI to confirm that a 0.7 kb DNA fragment was inserted. The primary sequence of the inserted DNA fragment was analyzed by using M13 forward primer M13F and M13 reverse primer M13R (Applied Biosystems automated sequencer Model 370A).

As a result, a sequence of 619 bp corresponding to nucleotide number 1 to 619 of SEQ ID NO: 30 was found, and the nucleotide sequence of 133 nucleotides consisting of nucleotide number 487 to 619 of SEQ ID NO: 30 conformed to the N-terminus of the primary sequence found in Example 2.1. In this 619 bp sequence there are 19 amino acids of SEQ ID NO: 3 (VERVDFTNHLEDTR RNINK from DNA nucleotide sequence) and 5 amino acids (LYDAK) of SEQ ID NO: 7, and it was clarified that this PCR reaction products (each

0.7 kb) was a part of cDNA coding for megakaryocyte differentiation factor.

It was considered that the translation starting methionine corresponds to 74th nucleotide, and 5'-non translational region consisted of 73 bp. Accordingly, it was clarified that these PCR reaction products (each 0.7 kb) contain N-terminus of structural gene for megakaryocyte differentiation factor.

5. Analysis of cDNA nucleotide sequence coding for megakaryocyte differentiations factor by PCR (4)

Among an N-terminal portion of structural gene for a megakaryocyte differentiation factor derived from HPC-Y11 and a sequence considered to be a 5'-non-translational region, obtained in the section C(3), the sequence of nucleotide numbers 12 to 31 of SEQ ID NO: 30 which is a sequence part considered to be a 5'-non-translational region was used as a basis to synthesize an oligomer NI 083 (SEQ ID NO: 31).

The RNA having poly A prepared from A431 cells in the section 1., a Preamplification System (Gibco BRL) and random hexamers attached to the System were used to synthesize first strand cDNA according to included instructions, and PCR was carried out by using NI083 and NI074 as well as Ampli Taq (Takara). As a result, a DNA fragment of 1001 bp which is a cDNA fragment for a megakaryocyte differentiation factor was obtained.

This DNA fragment, a PCR product, of 1001 bp was directly used as a substrate for sequencing on a Taq Dye Deoxy Terminator Cycle Sequencing kit (Applied Biosystem) and a fluorescent sequencer (Applied Biosystem Type 370A) according to the included manufacturers instruction. As a result, a sequence of nucleotide numbers 32-486 of SEQ ID NO: 30 was found. In addition, a result obtained for nucleotide numbers 487 to 991 of SEQ ID NO: 30 conformed to the sequence obtained in the section 1.

By combining with the sequence obtained in the section 1., the nucleotide sequence 32-1950 of SEQ ID NO: 30 which is cDNA nucleotide sequence coding for a megakaryocyte differentiation factor of A431 cell was determined.

For this nucleotide sequence, all of possible 3 reading frames were mechanically translated into amino acid sequences and it was found that one of them has a region which can be translated to a continuing amino acid sequence containing all the amino acid sequences shown in SEQ ID NO: 1 to 9, and the reading frame of the megakaryocyte differentiation factor was determined.

It was found that in this reading frame a codon for methionine (nucleotide numbers 74 to 76) found at a translation start position exists, and from this position a region which can be translated to an amino acid sequence containing the amino acid sequences shown in SEQ ID NO: 1 to 9 continues up to the nucleotide position 1213 of SEQ ID NO: 30, and it was determined that the nucleotide sequence from the position 74 to the position 1213 of SEQ ID NO: 30 is the region translated to megakaryocyte differentiation factor.

The nucleotide sequence GCAATGG (nucleotide numbers 71 to 77 of SEQ ID NO: 30) encompassing the methionine codon of nucleotide numbers 74 to 76 of SEQ ID NO: 30 corresponds to a sequence (G/A-N-N-A-T-G-G) encompassing a methionine codon frequently present at a translation start site found by M. Kozak, Nucleic Acids Research (1981) Vol. 9, p5233-5252.

Primary sequence of megakaryocyte differentiation factor was thus clarified, the number of amino acids was 380 in the structural gene, an expected molecular weight was 42904.43, and an expected isoelectric point was 6.79. SEQ ID NO: 1 corresponds to amino acid numbers 188 to 196 of SEQ ID NO: 30; SEQ ID NO: 2 corresponds to amino acid numbers

181 to 187 of SEQ ID NO: 30; SEQ ID NO: 3 corresponds to the amino acid numbers 126 to 144 of SEQ ID NO: 30; SEQ ID NO: 4 corresponds to the amino acid numbers 325 to 341 of SEQ ID NO: 30; SEQ ID NO: 5 corresponds to the amino acid numbers 289 to 297 of SEQ ID NO: 30; SEQ ID NO: 6 corresponds to the amino acid numbers 305 to 324 of SEQ ID NO: 30; SEQ ID NO: 7 corresponds to the amino acid numbers 121 to 125 of SEQ ID NO: 30; and SEQ ID NO: 8 corresponds to the amino acid numbers 284 to 288 of SEQ ID NO: 30; and SEQ ID NO: 9 corresponds to the amino acid numbers 204 to 213 of SEQ ID NO: 30. In addition, A poly A addition signal AATAAA sequence is present at nucleotide numbers 1933 to 1998 of SEQ ID NO: 30.

15 Example 3. Isolation and identification of cDNA coding for megakaryocyte differentiation factor from A431 by PCR and construction of expression vector

20 Oligomers NI078 (SEQ ID NO: 32) and NI079 (SEQ ID NO: 33) were synthesized on the basis of the sequence (SEQ ID NO: 30) obtained in Example 2. Note in the NI078, the sequence of nucleotide numbers 13 to 37 encompassing the translation start methionine codon conforms to the sequence of the nucleotide numbers 74 to 98 of SEQ ID NO: 30 and an EcoRI recognizing site (nucleotide numbers 4 to 9) and an Nru I recognizing site (nucleotide numbers 8 to 13) were artificially added; and in the NI079, the sequence of the nucleotide numbers 17 to 49 conforms to the nucleotide sequence of the nucleotide numbers 1237 to 1269 of SEQ ID NO: 30, and an EcoRI recognizing site (nucleotide numbers 3 to 8) and an Not I recognizing site (nucleotide numbers 9 to 16) were artificially added.

35 The RNA having poly A prepared from A431 cells in Example 2 section 1., a Preamplification System (Gibco BRL), and Oligomers, i.e., random hexamers included in the system were used according to the included manufacturer's instructions to synthesize first strand

cdNA, and PCR was carried out using the synthesized DNA as a template and NI078 and NI079 as primers and using Ampli Taq (Perkin Elmer Cetus). As a result, a DNA fragment of 1224 base pairs which is a cdNA
5 fragment for a megakaryocyte differentiation factor and has all information relating to megakaryocyte differentiation factor, was obtained.

This DNA fragment was treated with EcoRI to generate EcoRI cohesive sites at both the ends of the
10 cdNA coding for a megakaryocyte differentiation factor in virtue of EcoRI recognizing sites artificially added to the oligomers NI078 (SEQ ID NO: 32) and NI079 (SEQ ID NO: 33). This cdNA fragment coding for megakaryocyte differentiation factor was introduced
15 into a mammalian expression vector pdKCR-DHFD at its EcoRI recognizing site to obtain pdKCR-DHFR-TPO55.

The animal cell expression vector pdKCR-dhfr (Oikawa, S. et. al., Biochem. Biophys. Res. Commun. 164, 39, 1989) is a derivative of pKCR (O' Hare et. al., Proc. Natl. Acad. Sci. USA, 78, 1527, 1981) and
20 has SU 40 early promoter and a rabbit β -globin gene and dhfr (dehydrofolate reductase) gene. Note, a host transformed with the expression vector, was designated as Escherichia coli SBM 308, and deposited with the
25 Fermentation Research Institute, Agency of Industrial Science and Technology, 1-3, Higashi 3-chome, Tsukuba-shi, Ibaraki, Japan as FERM P-11506 on June 7, 1990, and transferred to an international deposition under the Budapest treaty as FERM BP-4197 on February 18,
30 1993.

The clone pdKCR-DHFR-TPO55 containing megakaryocyte differentiation factor cdNA which was incorporated in to pdKCR-DHFR was sequenced using a Taq Dye Deoxy Terminator Cycle Sequencing kit (Applied Biosystem) and a fluorescent sequencer (Applied Biosystem Type 370A) according to included
35 instructions. As a result, the determined nucleotide sequence conformed to the sequence of the nucleotide

numbers 99 to 1236 of SEQ ID NO: 30 and oligomers
NI078 and NI079. In addition, it was confirmed by the
sequencing that a megakaryocyte differentiation factor
cDNA inserted into the vector is in correct
5 orientation in relation to an expression vector
promoter.

As shown in the above, once the information of
SEQ ID NO: 30 is provided, it is easy for a person
skilled in the art that the nucleotide sequence is
10 determined by amplifying cDNA coding for megakaryocyte
differentiation factor in total or in a optional
portion on megakaryocyte differentiation factor
expressing cell line (for example, A431) and cloned in
a optional expression vector.

15 Example 4. Expression of megakaryocyte
differentiation factor in Bombyx mori

(1) Construction of Bombyx mori expression
vector

A megakaryocyte differentiation factor cDNA
20 clone pdKCR-DHFR-TP055 was digested with NotI to
cleave the NotI recognizing site artificially added to
the NI079. The NotI cohesive end thus generated was
blunt-ended using a blunting kit available from Takara
Shuzo, and to the blunt end was added an XbaI linker
25 (Takara Shuzo) according to an attached instruction.
The plasmid thus obtained was digested simultaneously
with NruI and XbaI to cleave the NruI recognizing site
artificially added to the NI078 and the XbaI
recognizing site of the XbaI linker introduced to
30 prepare a megakaryocyte differentiation factor cDNA
fragment having an NruI cohesive end and an XbaI
cohesive end. This DNA fragment was inserted at the
Nru I recognizing site into a baculovirus transfer
vector for Bombyx mori nuclear polyhedrosis virus,
35 pBm4 (available from Department of Insect Genetics and
Bleeding National Institute of Sericultural and
Entomological Science, Ohwashi, Tukuba, Ibaraki 305,
Japan) simultaneously digested with NruI and XbaI to

obtain pBm4-TP055.

(2) Construction of TP055 recombinant virus

A cell line derived from Bombyx mori embryonic, BoMol5AIIc (available from Department of Insect Genetics and Breeding National Institute of Sericultural and Entomological Science, Ohwashi, Tukuba, Ibaraki 305, Japan) was subcultured in a medium containing 10% fetal bovine serum (FBS: GIBCO BRL) and 500 µg/ml gentamicin in MGM 448 at 25°C.

TP055 recombinant virus was constructed by co-introducing Bombyx mori nuclear polyhedrosis virus gene DNA and pBm4-TP055 plasmid DNA into Bombyx mori cultured cells by, for example, calcium phosphate co-precipitation method.

Namely, 2 µg of genomic DNA of wild type virus B6E (available from Department of Insect Genetics and Breeding National Institute of Sericultural and Entomological Science, Ohwashi, Tukuba, Ibaraki 305, Japan) and 10 µg of the transfer plasmid pBm4-TP055 were dissolved in 240 µl of sterile purified water, and to the solution was added the same volume of 0.5 M CaCl₂ and 0.1M HEPES, and the mixture was mixed and allowed to stand at a room temperature for 10 minutes. To the mixture was added 480 µl of 0.2M NaCl, 0.05M HEPES, 0.75 mM NaH₂PO₄ and 0.75 mM Na₂HPO₄, and the mixture was stirred for a few second and allowed to stand at a room temperature for 20 to 30 minutes to form calcium phosphate gel containing the genomic DNA and the plasmid.

Next, 960 µl of calcium phosphate gel suspension containing the viral genomic DNA and the transfer vector was added to 4 ml of BoMol5AIIc cells in a 25 cm² T flask (T25, Corning), and the mixture was allowed to stand for 12 hours. The medium was replaced with a fresh MGM448 (containing 10% FBS and the antibiotics), and culturing was carried out for 25°C. On the sixth day the cultured medium was recovered as a viral solution.

The cultured medium was centrifuged to obtain a clear supernatant, which was then diluted, and added to BoMol5AIIc cells cultured on the microtiter plate, and after 8 days a culturing medium in which viral infection was microscopically observed but a polyhedral body was not formed, was selected (by a limited dilution method). The cultured medium was recovered. Contamination with a wild virus in the viral solution factor was not observed.

A recombinant virus thus constructed, containing a DNA coding for megakaryocyte differentiation factor was designated as TP055-BmNPV.

(3) Test for expression of recombinant gene

About 1×10^6 BoMol5A IIc cells were cultured in 4 ml of MGM448 medium containing 10% FBS on the bottom of 25 cm² area of a flask for 2 days by plate culture. To the culture, 0.5 moi of wild type virus B6E or recombinant virus (TP055-BmNPV) containing a gene coding for megakaryocyte differentiation factor were added and infected, BoMol5A IIc cells, and the cells were cultured at 25°C for 3 days, and total RNA was extracted using Isogen (Wako Pure Chemical). Similarly, total RNA was extracted from non infected BoMo 15AIIc cells.

Next, 1 Mg of the RNA thus extracted was size-fractionated by agarose gel electrophoresis, and the separated RNA was transferred to a Zetaprobe membrane by the capillary action. The membrane was soaked in a hybridization buffer containing megakaryocyte differentiation factor cDNA (:PCR product amplified with KY100 and NIO65 described in Example 2.1) (TP055 probe DNA) labeled with digoxigenin (Boehringer Mannheim), and the mixture was incubated at 42°C for 12 hours to allow the formation of specific complex of recombinant megakaryocyte differentiation factor mRNA and the TP055 probe DNA thereof. The complex was then reacted with an alkaline phosphatase-conjugated anti-digoxigenin

antibody (Boehringer Mannheim), and the complexed megakaryocyte differentiation factor mRNA was detected by chemoluminescence generated by hydrolysis of Lumigen PPD (AMPPD) (Boehringer Mannheim) according to manufacture's instructions with alkaline phosphates.

As seen in Fig. 11, recombinant megakaryocyte differentiation factor mRNA was detected in total RNA extracted from TPO55-BmNPV-infected cells, and it was shown that the mRNA was expressed in the TPO55-BmNPV-infected cells. On the other hand expression of mRNA which hybridized with the probe DNA was not observed in the B6E-infected cells and uninfected cells.

(4) Preparation of solution of recombinant virus

About 1×10^6 BoMol5AIIc cells were cultured in 4 ml of MGM448 containing 10% FBS on the bottom of a 25 cm² flask for 2 days, and to this culture was added 10 μ l of the culture medium of BoMol5AIIc cells containing the recombinant virus cloned in the above section (2). After culturing at 25°C for 14 days, the culture medium was centrifuged at 1000 rpm for 5 minutes to obtain a supernatant as a recombinant virus solution.

(5) Preparation of hemolymph of *Bombyx mori*

50 μ l/head of a viral solution of the 10^{-1} -diluted recombinant virus solution obtained in the above section (3) or a 10^{-1} -diluted wild type virus B6E solution was injected to *Bombyx mori* larvae at 5th instar, and the silkworms were fed with commercially available artificial feed (Morus; Katakura Kogyo) at 20°C for 4 to 5 days. The abdomens of 50 silkworms were cut, and an extract containing the hemolymph and the content in the central intestine was taken in a plastic tube cooled with ice, and centrifuged to obtain a supernatant.

(6) Confirmation of activity of megakaryocyte differentiation factor

50 ml of the hemolymph of the silkworms

obtained in the above section (5) was thoroughly dialyzed against a 20 mM Tris/HCl (pH 7.4) buffer, and applied to a Matrex Blue A column ($\phi 2.5 \times 15$ cm) equilibrated with the same buffer. The column was
5 thoroughly washed with the same buffer to eliminate a unbound fraction, and a bound protein was eluted by a concentration gradient of 0 to 1M NaCl. An elution profile for megakaryocyte differentiation activity of the hemolymph obtained from silkworms injected with
10 the recombinant virus was compared with that for a wild type virus.

As seen from Fig. 12, megakaryocyte differentiation activity in the hemolymph from silkworms injected with the recombinant virus was
15 significantly higher than that for the wild type virus.

Although Bombyx mori vacuovirus transfer vector pBm4, Bombyx mori nuclear polyhedrosis virus PbE and Bombyx mori cells BoMol5AIIc were used in
20 Example 4, the present invention is not limited to the use of these materials. Namely, other baculovirus transfer vector (such as pBK283, pBKblue, available from Funakoshi) Bombyx mori nuclear polyhedrosis virus (such as purified DNA available from Funakoshi),
25 Bombyx mori cells (such as BmN4 cells, available from Funakoshi) can be easily used by a person with ordinary skill in the art to obtain a megakaryocyte differentiation factor.

A megakaryocyte differentiation factor of
30 the present invention accelerates formation of megakaryocytes from myeloid cells in the presence of IL-3. The present megakaryocyte differentiation factor plays an important role in differentiation of megakaryocytes and acts in vivo as a thrombopoietin.
35 Accordingly, the present megakaryocyte differentiation factor may be medicaments effective to not only various diseases involving decrease of platelets but also for control of the number of platelets decreasing

2100690

- 42 -

by radiation in the case of bone marrow
transradiation, or for control of the number of
platelets in chemotherapy of cancers.

SEQUENCE LISTING

SEQ ID NO: 1

SEQUENCE TYPE: Amino acid

SEQUENCE LENGTH: 9

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

Xxx Glu Thr Ile Asn Xxx His Phe Lys

1

5

9

SEQ ID NO: 2

SEQUENCE TYPE: Amino acid

SEQUENCE LENGTH: 7

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

Xxx Gln Xxx Ala Phe Thr Lys

1

5

7

SEQ ID NO: 3

SEQUENCE TYPE: Amino acid

SEQUENCE LENGTH: 19

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

Val Glu Xxx Val Asp Phe Thr Asn His Leu Glu Asp Thr Xxx Xxx Asn

1

5

10

15

Ile Asn Lys

19

SEQ ID NO: 4

SEQUENCE TYPE: Amino acid

SEQUENCE LENGTH: 17

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

- 44 -

Xxx Tyr Ile Glu Val Thr Glu Glu Gly Thr Glu Ala Xxx Ala
1 5 10 15
Ala Xxx Gly
17

SEQ ID NO: 5
SEQUENCE TYPE: Amino acid
SEQUENCE LENGTH: 9
TOPOLOGY: Linear
MOLECULE TYPE: Peptide
SEQUENCE
Xxx Tyr Leu Arg Ala Leu Gly Leu Lys
1 5 9

SEQ ID NO: 6
SEQUENCE TYPE: Amino acid
SEQUENCE LENGTH: 20
TOPOLOGY: Linear
MOLECULE TYPE: Peptide
SEQUENCE
Ala Asp Leu Ser Gly Ile Ala Ser Gly Gly Arg Leu Tyr Ile Ser Arg
1 5 10 15
Met Xxx Gly Lys
20

SEQ ID NO: 7
SEQUENCE TYPE: Amino acid
SEQUENCE LENGTH: 5
TOPOLOGY: Linear
MOLECULE TYPE: Peptide
SEQUENCE
Leu Tyr Asp Ala Lys
1 5

SEQ ID NO: 8
SEQUENCE TYPE: Amino acid
SEQUENCE LENGTH: 5

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

Asn Tyr Glu Met Lys

1

5

SEQ ID NO: 9

SEQUENCE TYPE: Amino acid

SEQUENCE LENGTH: 10

TOPOLOGY: Linear

MOLECULE TYPE: Peptide

SEQUENCE

Ala Val Ala Met Met His Gln Xxx Arg Lys

1

5

10

SEQ ID NO: 10

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 38

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

FEATURES: corresponding to amino acid sequence of SEQ ID NO: 3;
I is inosine.

SEQUENCE

GTIGARIIIG TIGAYTTYAC IAAYCAYYTI GARGAYAC

38

SEQ ID NO: 11

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 32

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

FEATURES: corresponding to amino acid sequence of SEQ ID NO: 4; I
is inosine.

SEQUENCE

TACATCGAIG TIACIGARGA RGGIACNGAR GC

32

SEQ ID NO: 12

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 37

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

FEATURES: Oligomer attached to 3'-RACE kit (Gibco BRL).

SEQUENCE

GGCCACGCGT CGACTAGTAC TTTTTTTTTT TTTTTT

34

SEQ ID NO: 13

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 20

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

SEQUENCE

ATGTTGTGGG GACTGCTATA

20

SEQ ID NO: 14

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 23

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

SEQUENCE

CAAGGCGAAT GACCTCTAAG TAT

23

SEQ ID NO: 15

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 21

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

SEQUENCE

CCCCGAAGCA ATCCCAGAGA G

21

SEQ ID NO: 16
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 21
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
CTCAGGCAGC AGAACGTACA T 21

SEQ ID NO: 17
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 21
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
GGCGACGACT CCTGGAGCCC G 21

SEQ ID NO: 18
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 22
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
GACACCAGAC CAACTGGTAA TG 22

SEQ ID NO: 19
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 36
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
CATCCGGGAG ATGTACAGCC GGCCGCCAGA GGCAAT 36

SEQ ID NO: 20

SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 21
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
GCTGTGGCCA TGATGCACCA G

21

SEQ ID NO: 21
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 24
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
TACCTGCGGG CCCTGGGCCT GAAG

24

SEQ ID NO: 22
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 51
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
CATCCGGGAG ATGTACAGCC GGCCGCCAGA GGCAATGCCA GACAGGTCAG C

51

SEQ ID NO: 23
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 17
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
GTTTTCCCAG TCACGAC

17

SEQ ID NO: 24
SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 17
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
CAGGAAACAG CTATGAC

17

SEQ ID NO: 25
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 20
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
AATTATGGCC CACACCAGTG

20

SEQ ID NO: 26
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 20
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
ACTAGCCGCT ACAGTCAACA

20

SEQ ID NO: 27
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 21
STRANDNESS: Single
TOPOLOGY: Linear
MOLECULE TYPE: Synthetic DNA
SEQUENCE
TTGCCACTTG CCTTTGAAGT A

21

SEQ ID NO: 28
SEQUENCE TYPE: Nucleic acid
SEQUENCE LENGTH: 21

- 50 -

STRANDNESS: Single
 TOPOLOGY: Linear
 MOLECULE TYPE: Synthetic DNA
 SEQUENCE
 CTGATGCATC ATGGCGACTG C

21

SEQ ID NO: 29
 SEQUENCE TYPE: Nucleic acid
 SEQUENCE LENGTH: 21
 STRANDNESS: Single
 TOPOLOGY: Linear
 MOLECULE TYPE: Synthetic DNA
 SEQUENCE
 AGCATTACAC AGCACCATTA C

21

SEQUENCE ID NO: 30
 SEQUENCE TYPE: Nucleic acid
 SEQUENCE LENGTH: 1950
 STRANDNESS: Double
 TOPOLOGY: Linear
 MOLECULE TYPE: complimentary DNA (cDNA)
 ORIGINAL SOURCE: Human
 IMMEDIATE SOURCE: A431
 FEATURE: DNA coding for human megakaryocyte differentiation
 factor

SEQUENCE

GGCACGAGAG	GAAGTGAAGC	CCAGCTGTGA	AGGCCGCAGA	CTGCAGTGAG	50
AGGAGGCTGC	ACTCCATTTT	GCA ATG GCC	TCC CTT GCT	GCA GCA AAT	97
Met Ala Ser Leu Ala Ala Ala Asn					
1 5					
GCA GAG TTT	TGC TTC AAC	CTG TTC AGA	GAG ATG GAT	GAC AAT CAA	142
Ala Glu Phe	Cys Phe Asn	Leu Phe Arg	Glu Met Asp	Asp Asn Gln	
10	15	20			
GGA AAT GGA	AAT GTG TTC	TTT TCC TCT	CTG AGC CTC	TTC GCT GCC	187
Gly Asn Gly	Asn Val Phe	Phe Ser Ser	Leu Ser Leu	Phe Ala Ala	
25	30	35			
CTG GCC CTG	GTC CGC TTG	GGC GCT CAA	GAT GAC TCC	CTC TCT CAG	232

Leu	Ala	Leu	Val	Arg	Leu	Gly	Ala	Gln	Asp	Asp	Ser	Leu	Ser	Gln		
40						45					50					
ATT	GAT	AAG	TTG	CTT	CAT	GTT	AAC	ACT	GCC	TCA	GGA	TAT	GGA	AAC	277	
Ile	Asp	Lys	Leu	Leu	His	Val	Asn	Thr	Ala	Ser	Gly	Tyr	Gly	Asn		
55						60					65					
TCT	TCT	AAT	AGT	CAG	TCA	GGG	CTC	CAG	TCT	CAA	CTG	AAA	AGA	GTT	322	
Ser	Ser	Asn	Ser	Gln	Ser	Gly	Leu	Gln	Ser	Gln	Leu	Lys	Arg	Val		
70						75					80					
TTT	TCT	GAT	ATA	AAT	GCA	TCC	CAC	AAG	GAT	TAT	GAT	CTC	AGC	ATT	367	
Phe	Ser	Asp	Ile	Asn	Ala	Ser	His	Lys	Asp	Tyr	Asp	Leu	Ser	Ile		
85						90					95					
GTG	AAT	GGG	CTT	TTT	GCT	GAA	AAA	GTG	TAT	GGC	TTT	CAT	AAG	GAC	412	
Val	Asn	Gly	Leu	Phe	Ala	Glu	Lys	Val	Tyr	Gly	Phe	His	Lys	Asp		
100						105					110					
TAC	ATT	GAG	TGT	GCC	GAA	AAA	TTA	TAC	GAT	GCC	AAA	GTG	GAG	CGA	457	
Tyr	Ile	Glu	Cys	Ala	Glu	Lys	Leu	Tyr	Asp	Ala	Lys	Val	Glu	Arg		
115						120					125					
GTT	GAC	TTT	ACG	AAT	CAT	TTA	GAA	GAC	ACT	AGA	CGT	AAT	ATT	AAT	502	
Val	Asp	Phe	Thr	Asn	His	Leu	Glu	Asp	Thr	Arg	Arg	Asn	Ile	Asn		
130						135					140					
AAG	TGG	GTT	GAA	AAT	GAA	ACA	CAT	GGC	AAA	ATC	AAG	AAC	GTG	ATT	547	
Lys	Trp	Val	Glu	Asn	Glu	Thr	His	Gly	Lys	Ile	Lys	Asn	Val	Ile		
145						150					155					
GGT	GAA	GGT	GGC	ATA	AGC	TCA	TCT	GCT	GTA	ATG	GTG	CTG	GTG	AAT	592	
Gly	Glu	Gly	Gly	Ile	Ser	Ser	Ser	Ala	Val	Met	Val	Leu	Val	Asn		
160						165					170					
GCT	GTG	TAC	TTC	AAA	GGC	AAG	TGG	CAA	TCA	GCC	TTC	ACC	AAG	AGC	637	
Ala	Val	Tyr	Phe	Lys	Gly	Lys	Trp	Gln	Ser	Ala	Phe	Thr	Lys	Ser		
175						180					185					
GAA	ACC	ATA	AAT	TGC	CAT	TTC	AAA	TCT	CCC	AAG	TGC	TCT	GGG	AAG	682	
Glu	Thr	Ile	Asn	Cys	His	Phe	Lys	Ser	Pro	Lys	Cys	Ser	Gly	Lys		
190						195					200					
GCA	GTC	GCC	ATG	ATG	CAT	CAG	GAA	CGG	AAG	TTC	AAT	TTG	TCT	GTT	727	
Ala	Val	Ala	Met	Met	His	Gln	Glu	Arg	Lys	Phe	Asn	Leu	Ser	Val		
205						210					215					
ATT	GAG	GAC	CCA	TCA	ATG	AAG	ATT	CTT	GAG	CTC	AGA	TAC	AAT	GGT	772	

Ile	Glu	Asp	Pro	Ser	Met	Lys	Ile	Leu	Glu	Leu	Arg	Tyr	Asn	Gly			
220						225					230						
GGC	ATA	AAC	ATG	TAC	GTT	CTG	CTG	CCT	GAG	AAT	GAC	CTC	TCT	GAA	817		
Gly	Ile	Asn	Met	Tyr	Val	Leu	Leu	Pro	Glu	Asn	Asp	Leu	Ser	Glu			
235						240					245						
ATT	GAA	AAC	AAA	CTG	ACC	TTT	CAG	AAT	CTA	ATG	GAA	TGG	ACC	AAT	862		
Ile	Glu	Asn	Lys	Leu	Thr	Phe	Gln	Asn	Leu	Met	Glu	Trp	Thr	Asn			
250						255					260						
CCA	AGG	CGA	ATG	ACC	TCT	AAG	TAT	GTT	GAG	GTA	TTT	TTT	CCT	CAG	907		
Pro	Arg	Arg	Met	Thr	Ser	Lys	Tyr	Val	Glu	Val	Phe	Phe	Pro	Gln			
265						270					275						
TTC	AAG	ATA	GAG	AAG	AAT	TAT	GAA	ATG	AAA	CAA	TAT	TTG	AGA	GCC	952		
Phe	Lys	Ile	Glu	Lys	Asn	Tyr	Glu	Met	Lys	Gln	Tyr	Leu	Arg	Ala			
280						285					290						
CTA	GGG	CTG	AAA	GAT	ATC	TTT	GAT	GAA	TCC	AAA	GCA	GAT	CTC	TCT	997		
Leu	Gly	Leu	Lys	Asp	Ile	Phe	Asp	Glu	Ser	Lys	Ala	Asp	Leu	Ser			
295						300					305						
GGG	ATT	GCT	TCG	GGG	GGT	CGT	CTG	TAT	ATA	TCA	AGG	ATG	ATG	CAC	1042		
Gly	Ile	Ala	Ser	Gly	Gly	Arg	Leu	Tyr	Ile	Ser	Arg	Met	Met	His			
310						315					320						
AAA	TCT	TAC	ATA	GAG	GTC	ACT	GAG	GAG	GGC	ACC	GAG	GCT	ACT	GCT	1087		
Lys	Ser	Tyr	Ile	Glu	Val	Thr	Glu	Glu	Gly	Thr	Glu	Ala	Thr	Ala			
325						330					335						
GCC	ACA	GGA	AGT	AAT	ATT	GTA	GAA	AAG	CAA	CTC	CCT	CAG	TCC	ACG	1132		
Ala	Thr	Gly	Ser	Asn	Ile	Val	Glu	Lys	Gln	Leu	Pro	Gln	Ser	Thr			
340						345					350						
CTG	TTT	AGA	GCT	GAC	CAC	CCA	TTC	CTA	TTT	GTT	ATC	AGG	AAG	GAT	1177		
Leu	Phe	Arg	Ala	Asp	His	Pro	Phe	Leu	Phe	Val	Ile	Arg	Lys	Asp			
355						360					365						
GAC	ATC	ATC	TTA	TTC	AGT	GGC	AAA	GTT	TCT	TGC	CCT	TGA			1216		
Asp	Ile	Ile	Leu	Phe	Ser	Gly	Lys	Val	Ser	Cys	Pro	...					
370						375					380						
AAATCCAATT	GGTTTCTGTT	ATAGCAGTCC	CCACAACATC	AAAGAACCAC											1266		
CACAAGTCAA	TAGATTTGAG	TTTAATTGGA	AAAATGTGGT	GTTTCCTTTG											1316		
AGTTTATTTT	TTCCTAACAT	TGGTCAGCAG	ATGACACTGG	TGACTTGACC											1366		
CTTCCTAGAC	ACCTGGTTGA	TTGTCCTGAT	CCCTGCTCTT	AGCATTCTAC											1416		
CACCATGTGT	CTCACCCATT	TCTAATTTC	TTGTCTTTCT	TCCCACGCTC											1466		

ATTCTATCA TTCTCCCCCA TGACCCGTCT GGAAATTATG GAGAGTGCTC	1516
AACTGGTAAG GAGAACGTAG AAGTAGCCCT AGGGATCCTT TTTGAAACTC	1566
TACAGTTATC GCAGATATTC TAGCTTCATT GTAAGCAATC TAGGAAATAA	1616
GCCCTGCTGC TTTCTAGAAA TAAGTGTGAA GGATAAATTT TCTTTGTTGA	1666
CCTATGAAGA TTTTAGAGTT TACCTTCATA TGTTTGATTT TAAATCAGTG	1716
TATAATCTAG ATGGTAAAAA ATGTGAAATT GGGATTAGGG ACCAACC AAA	1766
ATATTTTCATT AATGCTTTCA ATTGACAAAT TTTGGTCTTT CTTTGATAAG	1816
ACAATATGTA CATAGTTTTT TCAAATATTA AAGATCTTTT AACTGTTGGC	1866
AGTTGTTATC TACAGAATCA TATCTCATAT GCTGTGTAGT TTATAAGTTT	1916
TTTCTCTATT TATCAGAATA AAGAAATACA ACAT	1950

SEQ ID NO: 31

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 20

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

ORIGINAL SOURCE: Human

FEATURES: 5'-non-translation region

SEQUENCE

AACTGAAGCC CAGCTGTGAA 20

SEQ ID NO: 32

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 37

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

SEQUENCE

CTCGAATTCG CGATGGCCTC CCTTGCTGCA GCAAATG 37

SEQ ID NO: 33

SEQUENCE TYPE: Nucleic acid

SEQUENCE LENGTH: 49

STRANDNESS: Single

TOPOLOGY: Linear

MOLECULE TYPE: Synthetic DNA

2100690

- 54 -

SEQUENCE

GGGAATTCGC GGCCGCGTGG TGGTTCTTTG ATGTTGTGGG GACTGCTAT

49

CLAIMS:

1. A megakaryocyte differentiation factor having the following properties;
 - (1) stimulating differentiation of megakaryocytes;
 - (2) exhibiting a molecular weight of 55 to 57 kD as determined by gel filtration and SDS-polyacrylamide gel electrophoresis (SDS-PAGE), and having no intermolecular disulphide linkage;
 - (3) exhibiting an isoelectric point of 6.5 +/- 0.5;and
 - (4) having at least one of the amino acid sequences shown in SEQ ID NO: 1 to 9 in the Sequence Listing.
2. A megakaryocyte differentiation factor according to claim 1, produced by human cells.
3. A megakaryocyte differentiation factor according to claim 2, produced by human cancer cells.
4. A megakaryocyte differentiation factor according to claim 3, produced by cells derived from human epidermoid carcinoma cell A431.
5. A megakaryocyte differentiation factor according to claim 4, produced by cells derived from human epidermoid carcinoma cell A431, cultured in a protein-free medium.
6. A megakaryocyte differentiation factor having an amino acid sequence the same as the native amino acid sequence of a megakaryocyte differentiation factor according to claim 1, an amino acid sequence wherein 1 to 30 amino acid residues in said native amino acid sequence is deleted, an amino acid sequence wherein 1 to 30 amino acid residues in said native amino acid sequence is

replaced with other amino acids, an amino acid sequence wherein 1 to 30 amino acid residues is added to said native amino acid sequence, or an amino acid sequence including a combination of said amino acid modifications, wherein any one of the amino acid sequence having amino acid residues deleted, replaced or added retains the activity of the native amino acid sequence.

7. A megakaryocyte differentiation factor essentially consisting of the amino acid sequence shown in SEQ ID NO: 30.

8. A megakaryocyte differentiation factor having an amino acid sequence wherein 1 to 30 amino acid residues in the amino acid sequence shown in SEQ ID NO: 30 is deleted, an amino acid sequence wherein 1 to 30 amino acid residues in the amino acid sequence shown in SEQ ID NO: 30 is replaced with other amino acids, or an amino acid sequence wherein 1 to 30 amino acid residues is added to the amino acid sequence shown in SEQ ID NO: 30, or an amino acid sequence including a combination of said amino acid modification, wherein any one of said amino acid sequence having amino acid residues deleted, replaced or added retains the activity of the native amino acid sequence.

9. A megakaryocyte differentiation factor according to claim 6, which is glycosylated.

10. A megakaryocyte differentiation factor according to claim 7, which is glycosylated.

11. A megakaryocyte differentiation factor according to claim 8, which is glycosylated.

12. A method of screening for a nucleic acid coding for a protein having megakaryocyte differentiation factor activity, said method comprising the steps of:

(1) hybridizing nucleic acids obtained from a culture of transformed cells constructed by a gene recombinant technique with a polynucleotide coding for at least one of the amino acid sequences shown in SEQ ID NOS: 1 to 9, under stringent conditions; and

(2) obtaining a nucleic acid which has hybridized with said polynucleotide.

13. A megakaryocyte differentiation factor according to claim 6, wherein the N-terminus of the factor is biochemically or chemically modified.

14. A megakaryocyte differentiation factor according to claim 13, wherein the first methionine is deleted and/or the second alanine is acetylated.

15. A megakaryocyte differentiation factor according to claim 8, wherein the N-terminus of the factor is biochemically or chemically modified.

16. A megakaryocyte differentiation factor according to claim 15, wherein the first methionine is deleted and/or the second alanine is acetylated.

17. The method of claim 12, wherein the N-terminus of the protein is biochemically or chemically modified.


18. The method of claim 17, wherein the protein has a methionine in the first position and/or an alanine in the second position, said methionine in the first position being deleted or said alanine in the second position being acetylated.

19. An isolated DNA coding for a megakaryocyte differentiation factor according to claim 1.
20. An isolated DNA coding for a megakaryocyte differentiation factor according to claim 6.
21. An expression vector comprising DNA according to claim 19.
22. An expression vector comprising DNA according to claim 20.
23. A host cell transformed with an expression vector according to claim 21.
24. A host cell transformed with an expression vector according to claim 22.
25. A pharmaceutical composition comprising a megakaryocyte differentiation factor according to claim 1 and a pharmaceutically acceptable carrier.
26. A pharmaceutical composition comprising a megakaryocyte differentiation factor according to claim 6 and a pharmaceutically acceptable carrier.
27. A pharmaceutical composition according to claim 25, for treating thrombocytopenia.
28. A pharmaceutical composition according to claim 26, for treating thrombocytopenia.
29. An antibody that specifically reacts with megakaryocyte differentiation factor according to claim 1.
30. An antibody that specifically reacts with megakaryocyte differentiation factor according to claim 6.

31. A process for production of a megakaryocyte differentiation factor according to claim 1, comprising the steps of culturing a host according to claim 23 and recovering the megakaryocyte differentiation factor.

32. A process according to claim 31, wherein the host is silkworm Bombyx mori.


Fig. 1



PATENT AGENTS


Swabe, Ogilvy, Renault

Fig. 2



PATENT AGENTS


Awakey Ogilvy Renault

Fig. 3

PATENT AGENTS

Swabe, Ogilvy & Renault

Fig. 12



PATENT AGENTS

Swaney Ogilvy Renault

UNSCANNABLE ITEM

RECEIVED WITH THIS APPLICATION

(ITEM ON THE 10TH FLOOR ZONE 5 IN THE FILE PREPARATION SECTION)

2100690

DOCUMENT REÇU AVEC CETTE DEMANDE

NE POUVANT ÊTRE BALAYÉ

(DOCUMENT AU 10 IÈME ÉTAGE AIRE 5 DANS LA SECTION DE LA
PRÉPARATION DES DOSSIERS)
