




- (51) **International Patent Classification:**
C07K 16/28 (2006.01) C07K 16/32 (2006.01)
- (21) **International Application Number:**
PCT/EP2016/065914
- (22) **International Filing Date:**
6 July 2016 (06.07.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
1511787.2 6 July 2015 (06.07.2015) GB
- (71) **Applicant:** UCB BIOPHARMA SPRL [BE/BE]; 60, Allée de la Recherche, 1070 Brussels (BE).
- (72) **Inventors:** GRIFFIN, Robert Anthony; c/o IPD, UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). HUMPHREYS, David Paul; c/o IPD, UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB). PETERS, Shirley Jane; c/o IPD, UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).
- (74) **Agent:** UCB INTELLECTUAL PROPERTY; c/o UCB Celltech, 208 Bath Road, Slough, Berkshire SL1 3WE (GB).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) **Title:** FUSION PROTEINS WHICH BIND TO HUMAN FC RECEPTORS

Figure 1(a)
Hinge-Fc-tailpiece-C575S



(57) **Abstract:** The invention relates to fusion proteins which bind to human Fc-receptors. The fusion proteins are initially produced as monomers, and are capable of assembly into multimers at a target site of interest. The invention also relates to therapeutic compositions comprising the fusion proteins, and their widespread use in the treatment of diseases.



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

— *of inventorship (Rule 4.17(iv))*

Declarations under Rule 4.17:

Published:

— *as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii))*

— *with international search report (Art. 21(3))*

Fusion proteins which bind to human Fc receptors

The invention relates to fusion proteins which bind to human Fc-receptors. The fusion proteins are initially produced as monomers, and are capable of assembly into multimers at a target site of interest. The invention also relates to therapeutic compositions comprising the fusion proteins, and their widespread use in the treatment of diseases.

BACKGROUND

Monoclonal antibodies (mAb) represent one of the fastest growing sectors of medical treatment. They are now successfully employed in the treatment of many diseases, ranging from cancer to autoimmunity. These diseases are rapidly increasing in number due to our aging population and so new and more effective treatments are required. To maintain this growth in mAb-based treatments and increase their effectiveness we need to better understand the mechanisms of action of mAb and develop new technologies and formats that can augment even further their native therapeutic capabilities.

Almost all mAb depend on their Fc region for function which interfaces with key elements of the immune system; most notably complement and immune effector cells. Key to this interaction are the structural relationships between the mAb Fc region and the immune receptor molecules; the first component of complement (C1) and various Fc gamma Receptors (FcγRs). However, these relationships remain only partially understood.

It has recently been found that complement may be activated by IgG hexamers assembled at the cell surface. Specific noncovalent interactions between Fc segments of IgG antibodies were observed to result in the formation of ordered antibody hexamers after antigen binding on cells. The hexamers recruited and activated C1, triggering the complement cascade. (Diebold C.A. et al., Science 343, 1260-1263, (2014).

Advances in antibody therapies have largely focussed on making the mAb more potent. In the case of enhanced effector functions this can often result in a less well tolerated drug with greater side-effects and dose-limiting toxicity. Thus, there remains a significant clinical need for improved antibody therapies with better safety profiles, which are highly effective and potent, and yet have fewer adverse side effects and low toxicity.

In the present invention we therefore provide improved fusion proteins which bind to human Fc-receptors. The fusion proteins are initially produced as monomers, and are capable of assembly into multimers at a target site of interest. The fusion proteins may be used to design a new class of antibody therapeutics, that are administered as relatively inert monomers, and assemble into highly potent multimers only when they reach a desired location at a target site of interest. The fusion proteins may thus provide improved therapeutic compositions with greater safety, which combine enhanced efficacy and potency with fewer adverse side effects and low toxicity.

DESCRIPTION OF THE INVENTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of skill in the art to which this invention belongs. All publications and patents referred to herein are incorporated by reference.

It will be appreciated that any of the embodiments described herein may be combined.

In the present specification the EU numbering system is used to refer to the residues in antibody domains, unless otherwise specified. This system was originally devised by Edelman et al, 1969 and is described in detail in Kabat et al, 1987.

Edelman *et al.*, 1969; "The covalent structure of an entire γ G immunoglobulin molecule," PNAS Biochemistry Vol.63 pp78-85.

Kabat *et al.*, 1987; in Sequences of Proteins of Immunological Interest, US Department of Health and Human Services, NIH, USA.

Where a position number and/or amino acid residue is given for a particular antibody isotype, it is intended to be applicable to the corresponding position and/or amino acid residue in any other antibody isotype, as is known by a person skilled in the art. When referring to an amino acid residue in a tailpiece derived from IgM or IgA, the position number given is the position number of the residue in naturally occurring IgM or IgA, according to conventional practice in the art.

The present invention provides a monomeric fusion protein comprising an antibody Fc-domain having two heavy chain Fc-regions derived from IgG; wherein one or each heavy chain Fc-region is fused at its C-terminal to an antibody tailpiece; wherein the tailpiece cysteine residue is modified to prevent disulphide bond formation.

In one embodiment, the tailpiece cysteine residue is deleted, substituted, or blocked with a thiol capping agent.

In one embodiment, the fusion protein of the present invention further comprises an antigen binding region. The antigen binding region may comprise any suitable antigen binding domain. In one example the antigen binding region may be derived from an antibody and may for example comprise a VH and/or a VL antigen binding domain. In one embodiment, the antigen binding region is selected from the group consisting of Fab, scFv, single domain antibody (dAb), and DARPin. In one example a single domain antibody may be a VH, VL or VHH domain. In one embodiment, the antigen binding region is fused to the N-terminus of the heavy chain Fc-region. The antigen binding region may be fused directly to the N-terminus of the heavy chain Fc-region. Alternatively it may be fused indirectly by means of an intervening amino acid sequence. For example, a short peptide linker or a hinge sequence may be provided between the fusion partner and the heavy chain Fc-region.

In one embodiment, the fusion protein of the present invention further comprises a fusion partner. Typically the term 'fusion partner' refers to an antigen, pathogen-associated molecular pattern (PAMP), drug, ligand, receptor, cytokine or chemokine. In one example the 'fusion partner' does not include an antibody or a variable domain
5 derived from an antibody. In one embodiment, the fusion partner is fused to the N-terminus of the heavy chain Fc-region. The fusion partner may be fused directly to the N-terminus of the heavy chain Fc-region. Alternatively it may be fused indirectly by means of an intervening amino acid sequence. For example, a short peptide linker or a hinge sequence may be provided between the fusion partner and the
10 heavy chain Fc-region.

The antibody Fc-domain component of the monomers of the present invention may be derived from any suitable species. In one embodiment the antibody Fc-domain is derived from a human Fc-domain.

15 The antibody Fc-domain may be derived from any suitable class of antibody, including IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3 and IgG4), and IgM. Typically, the antibody Fc-domain is derived from IgG. In one embodiment the antibody Fc-domain is derived from
20 IgG1. In one embodiment the antibody Fc-domain is derived from IgG2. In one embodiment the antibody Fc domain is derived from IgG3. In one embodiment the antibody Fc domain is derived from IgG4.

The antibody Fc-domain comprises two individual polypeptide chains, each referred
25 to as a heavy chain Fc-region. The two heavy chain Fc-regions dimerise to create the antibody Fc-domain. Whilst the two heavy chain Fc-regions that together form the antibody Fc domain may be different from one another it will be appreciated that these will usually be the same as one another.

30 Typically each heavy chain Fc-region comprises or consists of two or three heavy chain constant domains.

In native antibodies, the heavy chain Fc-region of IgA, IgD and IgG is composed of two heavy chain constant domains (CH2 and CH3) and that of IgE and IgM is

composed of three heavy chain constant domains (CH2, CH3 and CH4). These dimerise to create the antibody Fc domain.

5 In the present invention, the heavy chain Fc-region may comprise heavy chain constant domains from one or more different classes of antibody, for example one, two or three different classes.

10 In one embodiment the heavy chain Fc-region comprises CH2 and CH3 domains derived from IgG1.

In one embodiment the heavy chain Fc-region comprises CH2 and CH3 domains derived from IgG2.

15 In one embodiment the heavy chain Fc-region comprises CH2 and CH3 domains derived from IgG3.

In one embodiment the heavy chain Fc-region comprises CH2 and CH3 domains derived from IgG4.

20 In our earlier application, PCT/EP2015/054687, we disclose that the CH3 domain plays a significant role in the polymerisation of fusion proteins comprising an antibody Fc-domain and a tail-piece sequence. The amino acid at position 355 of the CH3 domain was found to have a particularly strong effect.

25 Thus in one embodiment, the heavy chain Fc-region comprises a CH3 domain derived from IgG1.

In one embodiment, the heavy chain Fc-region comprises a CH2 domain derived from IgG4 and a CH3 domain derived from IgG1.

30 In one embodiment, the heavy chain Fc-region comprises an arginine residue at position 355.

In one embodiment the heavy chain Fc-region comprises a CH4 domain from IgM. The IgM CH4 domain is typically located between the CH3 domain and the tailpiece.

5 In one embodiment the heavy chain Fc-region comprises CH2 and CH3 domains derived from IgG and a CH4 domain derived from IgM.

10 It will be appreciated that the heavy chain constant domains for use in producing a heavy chain Fc-region of the present invention may include variants of the naturally occurring constant domains described above. Such variants may comprise one or more amino acid variations compared to wild type constant domains. In one example the heavy chain Fc-region of the present invention comprises at least one constant domain which varies in sequence from the wild type constant domain. It will be appreciated that the variant constant domains may be longer or shorter than the wild type constant domain. Preferably the variant constant domains are at least 50% 15 identical or similar to a wild type constant domain. The term "identity", as used herein, indicates that at any particular position in the aligned sequences, the amino acid residue is identical between the sequences. The term "similarity", as used herein, indicates that, at any particular position in the aligned sequences, the amino acid residue is of a similar type between the sequences. For example, leucine may 20 be substituted for isoleucine or valine. Other amino acids which can often be substituted for one another include but are not limited to:

- phenylalanine, tyrosine and tryptophan (amino acids having aromatic side chains);
- lysine, arginine and histidine (amino acids having basic side chains);
- aspartate and glutamate (amino acids having acidic side chains);
- 25 – asparagine and glutamine (amino acids having amide side chains); and
- cysteine and methionine (amino acids having sulphur-containing side chains).

Degrees of identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing. 30 Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M.

and Devereux, J., eds., M Stockton Press, New York, 1991). In one example the variant constant domains are at least 60% identical or similar to a wild type constant domain. In another example the variant constant domains are at least 70% identical or similar. In another example the variant constant domains are at least 80% identical or similar. In another example the variant constant domains are at least 90% identical or similar. In another example the variant constant domains are at least 95% identical or similar.

In the fusion proteins of the invention, one or each heavy chain Fc-region is fused at its C-terminus to an antibody tailpiece, wherein the tailpiece cysteine residue is modified to prevent disulphide bond formation.

The antibody tailpiece of the invention may be derived from any suitable species. Antibody tailpieces are evolutionarily conserved and are found in most species, including primitive species such as teleosts. In one embodiment, the antibody tailpiece is derived from a human antibody.

In humans, IgM and IgA occur naturally as covalent multimers of the common H₂L₂ antibody unit. IgM occurs as a pentamer when it has incorporated a J-chain, or as a hexamer when it lacks a J-chain. IgA occurs as monomer and dimer forms. The heavy chains of IgM and IgA possess an 18 amino acid extension to the C-terminal constant domain, known as a tailpiece. This tailpiece naturally includes a cysteine residue that forms a disulphide bond with adjacent heavy chains in the polymer, and is believed to have an important role in polymerisation. The tailpiece also contains a glycosylation site.

In our earlier application, PCT/EP2015/054687, we disclosed that recombinant fusion proteins comprising an antibody Fc-domain derived from IgG and a tailpiece, are synthesised predominantly as hexamers.

The present inventors have unexpectedly found that modification of the tailpiece cysteine residue results in fusion proteins with very unusual multimerisation properties. The fusion proteins of the present invention are synthesized predominantly as monomers, and are capable of assembly into multimers at high

concentrations, as shown in Figure 1. The fusion proteins may be used to design a new class of antibody therapeutics, that are administered as relatively inert monomers, and assemble into highly potent multimers only when they reach a desired location at a target site of interest. The fusion proteins may thus provide improved therapeutic compositions with greater safety, which combine enhanced efficacy and potency with fewer adverse side effects and low toxicity.

In one example the monomers of the present invention are directed to a target site of interest via the Fc domain and/or, where present, the antigen binding region and/or fusion partner. For example, the monomer of the present invention may comprise an antigen binding region which may bind an antigen expressed on the surface of a cell, such as a tumor cell or an immune cell, such that assembly of the monomer into a highly potent multimer may occur on the tumor cell or immune cell surface. Examples of suitable antigens include HER2/neu or CD20.

In the fusion proteins of the present invention, the tailpiece cysteine residue is modified to prevent disulphide bond formation. The cysteine residue may be modified using any any suitable method that prevents disulphide bond formation.

In one embodiment, the tailpiece cysteine residue is mutated. In one embodiment, the tailpiece cysteine residue is deleted. In one embodiment, the tailpiece cysteine residue is substituted with another amino acid residue.

In one embodiment, the antibody tailpiece is derived from human IgM or IgA, wherein the tailpiece cysteine residue normally found at position 575 of IgM or position 495 of IgA has been deleted or substituted with another amino acid residue.

In one embodiment, the tailpiece cysteine residue normally found at position 575 of IgM is substituted with a serine, threonine or alanine residue (C575S, C575T, or C575A).

In one embodiment, the tailpiece cysteine residue normally found at position 495 of IgA is substituted with a serine, threonine or alanine residue (C495S, C495T, or C495A).

In one embodiment, the tailpiece cysteine residue is blocked with a thiol capping agent. A thiol capping agent is a compound that reacts with sulphhydryl groups in reduced cysteine residues, preventing them from forming disulphide bonds.

5 Examples of suitable thiol capping agents include N-ethylmaleimide, iodoacetic acid, or iodoacetamide.

In one embodiment, the thiol capping agent used to block the tailpiece cysteine residue is conjugated to a drug. The capping reaction thus effectively forms a bridge
10 between the drug and the cysteine residue in the target protein, conjugating the drug to the tailpiece.

In one embodiment, the tailpiece comprises all or part of an 18 amino acid tailpiece sequence from human IgM or IgA as shown in Table 1, wherein the cysteine residue
15 normally found at position 575 of IgM, or position 495 of IgA, is deleted, substituted, or blocked with a thiol capping agent.

The tailpiece may be fused directly to the C-terminus of the heavy chain Fc-region. Alternatively, it may be fused indirectly by means of an intervening amino acid
20 sequence. For example, a short linker sequence may be provided between the tailpiece and the heavy chain Fc-region.

The tailpiece of the present invention may include variants or fragments of the tailpiece sequences described above. A variant of an IgM or IgA tailpiece typically
25 has an amino acid sequence which is identical to the described sequence in 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17 of the 18 amino acid positions shown in Table 1. A fragment typically comprises 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17 amino acids. The tailpiece may be a hybrid IgM/IgA tailpiece. Fragments of variants are also envisaged.

30

Table 1 Tailpiece sequences

<u>Tailpiece</u>	<u>Sequence</u>
------------------	-----------------

IgM	PTLYNVSLVMSDTAGTCY	SEQ ID NO: 1
IgA	PTHVNVSVVMAEVDGTCY	SEQ ID NO: 2

Each heavy chain Fc-region of the present invention may optionally possess a native or a modified hinge region at its N-terminus.

5

A native hinge region is the hinge region that would normally be found between Fab and Fc domains in a naturally occurring antibody. A modified hinge region is any hinge that differs in length and/or composition from the native hinge region. Such hinges can include hinge regions from other species, such as human, mouse, rat, rabbit, shark, pig, hamster, camel, llama or goat hinge regions. Other modified hinge regions may comprise a complete hinge region derived from an antibody of a different class or subclass from that of the heavy chain Fc-region. Alternatively, the modified hinge region may comprise part of a natural hinge or a repeating unit in which each unit in the repeat is derived from a natural hinge region. In a further alternative, the natural hinge region may be altered by converting one or more cysteine or other residues into neutral residues, such as serine or alanine, or by converting suitably placed residues into cysteine residues. By such means the number of cysteine residues in the hinge region may be increased or decreased. Other modified hinge regions may be entirely synthetic and may be designed to possess desired properties such as length, cysteine composition and flexibility.

A number of modified hinge regions have already been described for example, in US5677425, WO9915549, WO2005003170, WO2005003169, WO2005003170, WO9825971 and WO2005003171 and these are incorporated herein by reference.

25

Examples of suitable hinge sequences are shown in Table 2.

In one embodiment, the heavy chain Fc-region possesses an intact hinge region at its N-terminus.

30

In one embodiment the heavy chain Fc-region and hinge region are derived from IgG4 and the hinge region comprises the mutated sequence CPPC (SEQ ID NO: 11). The core hinge region of human IgG4 naturally contains the sequence CPSC (SEQ ID NO: 12), compared to IgG1 which contains the sequence CPPC. The serine residue present in the IgG4 sequence leads to increased flexibility in this region, and therefore a proportion of molecules form disulphide bonds within the same protein chain (an intrachain disulphide) rather than bridging to the other heavy chain in the IgG molecule to form the interchain disulphide. (Angal S. et al, Mol Immunol, Vol 30(1), p105-108, 1993). Changing the serine residue to proline to give the same core sequence as IgG1 allows complete formation of inter-chain disulphides in the IgG4 hinge region, thus reducing heterogeneity in the purified product. This altered isotype is termed IgG4P.

Table 2. Hinge sequences

<u>Hinge</u>	<u>Sequence</u>	
Human IgA1	VPSTPPTPSPSTPPTSPSPS	SEQ ID NO: 3
Human IgA2	VPPPPP	SEQ ID NO: 4
Human IgD	ESPKAQASSVPTAQPPQAEGLAKATTAPATTRNTGRGGEE KKKEKEKEEQEERETKTP	SEQ ID NO: 5
Human IgG1	EPKSCDKTHTCPPCP	SEQ ID NO: 6
Human IgG2	ERKCCVECPPCP	SEQ ID NO: 7
Human IgG3	ELKTPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTP PPCPRCPEPKSCDTPPPCPRCP	SEQ ID NO: 8
Human IgG4	ESKYGPPCPSCP	SEQ ID NO: 9
Human IgG4(P)	ESKYGPPCPPCP	SEQ ID NO: 10
Recombinant v1	CPPC	SEQ ID NO: 11
Recombinant v2	CPSC	SEQ ID NO: 12
Recombinant v3	CPRC	SEQ ID NO: 13
Recombinant v4	SPPC	SEQ ID NO: 14

Recombinant v5	CPPS	SEQ ID NO: 15
Recombinant v6	SPPS	SEQ ID NO: 16
Recombinant v7	DKHTTCAA	SEQ ID NO: 17
Recombinant v8	DKHTTCPPCPA	SEQ ID NO: 18
Recombinant v9	DKHTTCPPCPATCPPCPA	SEQ ID NO: 19
Recombinant v10	DKHTTCPPCPATCPPCPATCPPCPA	SEQ ID NO: 20
Recombinant v11	DKHTTCPPCPAGKPTLYNSLVMSDTAGTCY	SEQ ID NO: 21
Recombinant v12	DKHTTCPPCPAGKPTHVNVSVVMAEVDGTCY	SEQ ID NO: 22
Recombinant v13	DKHTTCCVECPCPA	SEQ ID NO: 23
Recombinant v14	DKHTTCPRCPEPKSCDTPPPCPRCPA	SEQ ID NO: 24
Recombinant v15	DKHTTCPSCPA	SEQ ID NO: 25

In one embodiment, a monomeric fusion protein of the invention comprises an amino acid sequence as provided in Figure 2, optionally with an alternative hinge or tailpiece sequence.

Accordingly in one example the present invention provides a monomeric fusion protein comprising two identical polypeptide chains, each polypeptide chain comprising or consisting of the sequence given in any one of SEQ ID NOs 26 to 37.

In one example where the hinge may be varied from the sequences given in SEQ ID NOs 26 to 37, the present invention provides a monomeric fusion protein comprising two identical polypeptide chains, each polypeptide chain comprising or consisting of the sequence given in amino acids 6 to 222 of any one of SEQ ID NOs 26 to 29, or the sequence given in amino acids 6 to 333 of SEQ ID NOs 30 or 31, or the sequence given in amino acids 6 to 221 of any one of SEQ ID NOs 32 to 35, or the sequence given in amino acids 6 to 332 of SEQ ID NOs 36 or 37.

In one embodiment, a monomeric fusion protein of the invention comprises an amino acid sequence as provided in Figure 3. Accordingly in one example the present invention provides a monomeric fusion protein comprising or consisting of an amino acid sequence given in any one of SEQ ID Nos 38 to 69.

5

The monomeric fusion protein of the invention is capable of concentration-dependent assembly into multimers. The multimer may comprise two, three, four, five, six, seven, eight, nine, ten, eleven or twelve or more monomer units. Such a multimer may alternatively be referred to as a dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer, decamer, undecamer, dodecamer, etc., respectively. In one example, the monomeric fusion protein assembles into a hexamer.

10

Thus in one embodiment, the present invention provides a mixture comprising a monomeric fusion protein of the invention and a multimer, said multimer comprising two or more monomer units.

15

In one embodiment, the mixture comprises a monomeric fusion protein of the invention and a hexamer.

In one embodiment, the mixture comprises greater than 55% monomer, for example, greater than 65%, greater than 75%, greater than 85%, greater than 90% or greater than 95% monomer.

20

In one embodiment, the mixture is enriched for the monomeric form of the fusion protein of the invention. In one example, the term "enriched" means that greater than 80% of the fusion protein of the invention is present in the mixture in monomeric form, such as greater than 90% or greater than 95%. It will be appreciated that the proportion of monomer in a sample can be determined using any suitable method such as analytical size exclusion chromatography, as described herein below.

25

30

In one embodiment, the monomeric fusion protein of the invention is produced or formulated using a component that stabilises the monomeric form of the protein. The component may increase the ratio of monomer to multimer in a mixture, so that a higher proportion of the protein is present in monomeric form than would otherwise

be the case. Examples of suitable components include pharmaceutical excipients, diluents or carriers which enable the monomeric fusion protein of the invention to be formulated at high concentration prior to administration.

- 5 The monomeric fusion proteins of the present invention may comprise one or more mutations that alter the functional properties of the proteins, for example, binding to Fc-receptors such as leukocyte receptors, complement or FcRn. Examples of mutations that alter the functional properties of the proteins are described in detail in our earlier application, PCT/EP2015/054687. It will be appreciated that any of these
- 10 mutations may be combined in any suitable manner to achieve the desired functional properties, and/or combined with other mutations to alter the functional properties of the proteins.

- The present invention also provides an isolated DNA sequence encoding a
- 15 polypeptide chain of the present invention, or a component part thereof. The DNA sequence may comprise synthetic DNA, for instance produced by chemical processing, cDNA, genomic DNA or any combination thereof.

- DNA sequences which encode a polypeptide chain of the present invention can be
- 20 obtained by methods well known to those skilled in the art. For example, DNA sequences coding for part or all of a polypeptide chain may be synthesised as desired from the determined DNA sequences or on the basis of the corresponding amino acid sequences.

- 25 In one example, a monomeric fusion protein of the invention is encoded by a DNA sequence as provided in Figure 3. Accordingly in one example the present invention provides a DNA sequence given in any one of SEQ ID Nos 38 to 69.

- Standard techniques of molecular biology may be used to prepare DNA sequences
- 30 coding for a polypeptide chain of the present invention. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate.

The present invention also relates to a cloning or expression vector comprising one or more DNA sequences of the present invention. Accordingly, provided is a cloning or expression vector comprising one or more DNA sequences encoding a polypeptide chain of the present invention, or a component part thereof.

5

General methods by which the vectors may be constructed, transfection methods and culture methods are well known to those skilled in the art. In this respect, reference is made to "Current Protocols in Molecular Biology", 1999, F. M. Ausubel (ed), Wiley Interscience, New York and the Maniatis Manual produced by Cold
10 Spring Harbor Publishing.

15

Also provided is a host cell comprising one or more cloning or expression vectors comprising one or more DNA sequences encoding a monomeric fusion protein of the present invention. Any suitable host cell/vector system may be used for expression
of the DNA sequences encoding the monomeric fusion protein of the present
invention. Bacterial, for example *E. coli*, and other microbial systems such as
Saccharomyces or Pichia may be used or eukaryotic, for example mammalian, host
cell expression systems may also be used. Suitable mammalian host cells include
CHO cells. Suitable types of chinese hamster ovary (CHO cells) for use in the
20 present invention may include CHO and CHO-K1 cells, including dhfr- CHO cells,
such as CHO-DG44 cells and CHO-DXB11 cells, which may be used with a DHFR
selectable marker, or CHOK1-SV cells which may be used with a glutamine
synthetase selectable marker. Other suitable host cells include NSO cells.

20

25

The present invention also provides a process for the production of a monomeric fusion protein according to the present invention, comprising culturing a host cell containing a vector of the present invention under conditions suitable for expression of the monomeric fusion protein, and isolating and optionally purifying the monomeric fusion protein.

30

The monomeric fusion proteins of the present invention are expressed at good levels from host cells. Thus the properties of the monomeric fusion protein are conducive to commercial processing.

The monomeric fusion proteins of the present invention may be made using any suitable method. In one embodiment, the monomeric fusion protein of the invention may be produced under conditions which minimise aggregation. In one example, aggregation may be minimised by the addition of preservative to the culture media, culture supernatant, or purification media. Examples of suitable preservatives include ethylenediaminetetraacetic acid (EDTA), ethyleneglycoltetraacetic acid (EGTA), or acidification to below pH 6.0.

In one embodiment there is provided a process for purifying a monomeric fusion protein of the present invention comprising the steps: performing anion exchange chromatography in non-binding mode such that the impurities are retained on the column and the monomeric fusion protein is eluted.

In one embodiment the purification employs affinity capture on an FcRn, FcγR or C-reactive protein column.

In one embodiment the purification employs protein A.

Suitable ion exchange resins for use in the process include Q.FF resin (supplied by GE-Healthcare). The step may, for example be performed at a pH about 8.

The process may further comprise an initial capture step employing cation exchange chromatography, performed for example at a pH of about 4 to 5, such as 4.5. The cation exchange chromatography may, for example employ a resin such as CaptoS resin or SP sepharose FF (supplied by GE-Healthcare). The monomeric fusion protein can then be eluted from the resin with an ionic salt solution such as sodium chloride, for example at a concentration of 200mM.

The chromatography step or steps may include one or more washing steps, as appropriate.

The purification process may also comprise one or more filtration steps, such as a diafiltration step.

The monomeric fusion proteins of the invention can be purified according to molecular size, for example by size exclusion chromatography.

Thus in one embodiment there is provided a purified monomeric fusion protein
5 according to the invention, in substantially purified form, in particular free or substantially free of endotoxin and/or host cell protein or DNA.

Purified form as used herein is intended to refer to at least 90% purity, such as 91, 92, 93, 94, 95, 96, 97, 98, 99% w/w or more pure.

10

Substantially free of endotoxin is generally intended to refer to an endotoxin content of 1 EU per mg protein product or less, such as 0.5 or 0.1 EU per mg product.

15

Substantially free of host cell protein or DNA is generally intended to refer to a host cell protein and/or DNA content of 400µg per mg of protein product or less, such as 100µg per mg product or less, in particular 20µg per mg product.

20

As the monomeric fusion proteins of the present invention are useful in the treatment and/or prophylaxis of a pathological condition, the present invention also provides a pharmaceutical or diagnostic composition comprising a monomeric fusion protein of the present invention in combination with one or more of a pharmaceutically acceptable excipient, diluent or carrier.

25

The present invention also provides a process for preparation of a pharmaceutical or diagnostic composition comprising adding and mixing the monomeric fusion protein of the present invention together with one or more of a pharmaceutically acceptable excipient, diluent or carrier.

30

In one embodiment, the pharmaceutical composition comprises a component that stabilises the monomeric form of the protein or increases the ratio of monomer to multimer in a mixture.

The monomeric fusion protein may be the sole active ingredient in the pharmaceutical or diagnostic composition, or may be accompanied by other active

ingredients including antibody ingredients or non-antibody ingredients such as other drug molecules.

- The pharmaceutical compositions suitably comprise a therapeutically effective amount of the monomeric fusion protein of the invention. The term “therapeutically effective amount” as used herein refers to an amount of a therapeutic agent needed to treat, ameliorate or prevent a targeted disease or condition, or to exhibit a detectable therapeutic or preventative effect. For any medicine, the therapeutically effective amount can be estimated initially either in cell culture assays or in animal models, usually in rodents, rabbits, dogs, pigs or primates. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.
- 15 The precise therapeutically effective amount for a human subject will depend upon the severity of the disease state, the general health of the subject, the age, weight and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities and tolerance/response to therapy. This amount can be determined by routine experimentation and is within the judgement of the clinician. Generally, a therapeutically effective amount will be from 0.01 mg/kg to 500 mg/kg, for example 0.1 mg/kg to 200 mg/kg, such as 100mg/kg. Pharmaceutical compositions may be conveniently presented in unit dose forms containing a predetermined amount of a monomeric fusion protein of the invention per dose.
- 20
- 25 Compositions may be administered individually to a patient or may be administered in combination (*e.g.* simultaneously, sequentially or separately) with other agents, drugs or hormones.

- The dose at which the monomeric fusion protein of the present invention is administered depends on the nature of the condition to be treated, the extent of the disease present, and on whether the monomeric fusion protein is being used prophylactically or to treat an existing condition.
- 30

The frequency of dose will depend on the half-life of the monomeric fusion protein and the duration of its effect. If the monomeric fusion protein has a short half-life (e.g. 2 to 10 hours) it may be necessary to give one or more doses per day.

- Alternatively, if the monomeric fusion protein has a long half-life (e.g. 2 to 15 days),
5 and/or long lasting pharmacodynamic effects, it may only be necessary to give a dosage once per day, once per week or even once every 1 or 2 months.

In one embodiment the dose is delivered bi-weekly, i.e. twice a month.

- 10 Half-life as employed herein is intended to refer to the duration of the monomeric fusion protein in circulation, for example in serum or plasma.

Pharmacodynamics as employed herein refers to the profile and, in particular, duration of the biological action of the monomeric fusion protein.

15

The pharmaceutically acceptable carrier should not itself induce the production of antibodies harmful to the individual receiving the composition and should not be toxic. Suitable carriers may be large, slowly metabolised macromolecules such as proteins, polypeptides, liposomes, polysaccharides, polylactic acids, polyglycolic
20 acids, polymeric amino acids, amino acid copolymers and inactive virus particles.

Pharmaceutically acceptable salts can be used, for example mineral acid salts, such as hydrochlorides, hydrobromides, phosphates and sulphates, or salts of organic acids, such as acetates, propionates, malonates and benzoates.

- 25 Pharmaceutically acceptable carriers in therapeutic compositions may additionally contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents or pH buffering substances, may be present in such compositions.

- 30 A thorough discussion of pharmaceutically acceptable carriers is available in Remington's Pharmaceutical Sciences (Mack Publishing Company, N.J. 1991).

The therapeutic suspensions or solution formulations can also contain one or more excipients. Excipients are well known in the art and include buffers (e.g., citrate

buffer, phosphate buffer, acetate buffer and bicarbonate buffer), amino acids, urea, alcohols, ascorbic acid, phospholipids, proteins (e.g., serum albumin), EDTA, sodium chloride, liposomes, mannitol, sorbitol, and glycerol. Solutions or suspensions can be encapsulated in liposomes or biodegradable microspheres.

5

The formulation will generally be provided in a substantially sterile form employing sterile manufacture processes. This may include production and sterilization by filtration of the buffered solvent/solution used for the formulation, aseptic suspension of the protein in the sterile buffered solvent solution, and dispensing of the formulation into sterile receptacles by methods familiar to those of ordinary skill in the art.

10

Suitable forms for administration include forms suitable for parenteral administration, e.g. by injection or infusion, for example by bolus injection or continuous infusion.

15

Where the product is for injection or infusion, it may take the form of a suspension, solution or emulsion in an oily or aqueous vehicle and it may contain formulatory agents, such as suspending, preservative, stabilising and/or dispersing agents. The monomeric fusion protein may be in the form of nanoparticles. Alternatively, the monomeric fusion protein may be in dry form, for reconstitution before use with an appropriate sterile liquid.

20

Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals. However, in one or more embodiments the compositions are adapted for administration to human subjects.

25

The pharmaceutical compositions of this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, transcutaneous (for example, see WO98/20734), subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, intravaginal or rectal routes. Typically, the therapeutic compositions may be prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared.

30

Direct delivery of the compositions will generally be accomplished by injection, subcutaneously, intraperitoneally, intravenously or intramuscularly, or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Dosage treatment may be a single dose schedule or a multiple dose schedule

5

It will be appreciated that the active ingredient in the composition will be a protein molecule. As such, it will be susceptible to degradation in the gastrointestinal tract. Thus, if the composition is to be administered by a route using the gastrointestinal tract, the composition will need to contain agents which protect the protein from degradation but which release the protein once it has been absorbed from the gastrointestinal tract.

10

In one embodiment the formulation is provided as a formulation for topical administrations including inhalation.

15

In one embodiment we provide the monomeric fusion protein of the invention for use in therapy. In one embodiment we provide the use of a monomeric fusion protein of the invention for the manufacture of a medicament.

20

In one embodiment we provide the monomeric fusion protein of the invention for use in the treatment of cancer.

In one embodiment we provide the use of a monomeric fusion protein of the invention for the manufacture of a medicament for the treatment of cancer.

25

Examples of cancers which may be treated using the monomeric fusion protein of the invention include colorectal cancer, liver cancer, prostate cancer, pancreatic cancer, breast cancer, ovarian cancer, thyroid cancer, renal cancer, bladder cancer, head and neck cancer or lung cancer. In one embodiment the cancer is skin cancer, such as melanoma. In one embodiment the cancer is Leukemia. In one embodiment the cancer is glioblastoma, medulloblastoma or neuroblastoma. In one embodiment the cancer is a neuroendocrine cancer. In one embodiment the cancer is Hodgkin's or non-Hodgkins lymphoma.

30

In one embodiment we provide the monomeric fusion protein of the invention for use in the treatment of immune disorders.

In one embodiment we provide the use of the monomeric fusion protein of the invention for the preparation of a medicament for the treatment of immune disorders.

Examples of immune disorders which may be treated using the monomeric fusion protein of the invention include autoimmune diseases, for example Acute Disseminated Encephalomyelitis (ADEM), Acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, ANCA-associated vasculitis, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome (APS), Autoimmune angioedema, Autoimmune aplastic anemia, Autoimmune dysautonomia, Autoimmune hepatitis, Autoimmune hyperlipidemia, Autoimmune immunodeficiency, Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura (ATP), Autoimmune thyroid disease, Autoimmune urticarial, Axonal & nal neuropathies, Balo disease, Behcet's disease, Bullous pemphigoid, Cardiomyopathy, Castleman disease, Celiac disease, Chagas disease, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, Cicatricial pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogans syndrome, Cold agglutinin disease, Congenital heart block, Coxsackie myocarditis, CREST disease, Essential mixed cryoglobulinemia, Demyelinating neuropathies, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Dilated cardiomyopathy, Discoid lupus, Dressler's syndrome, Endometriosis, Eosinophilic angiocentric fibrosis, Eosinophilic fasciitis, Erythema nodosum, Experimental allergic encephalomyelitis, Evans syndrome, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (GPA) see Wegener's, Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Hemolytic anemia, Henoch-Schonlein purpura, Herpes gestationis, Hypogammaglobulinemia, Idiopathic hypocomplementemic tubulointerstitial nephritis, Idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related disease, IgG4-related sclerosing disease, Immunoregulatory lipoproteins, Inflammatory aortic aneurysm, Inflammatory

pseudotumour, Inclusion body myositis, Insulin-dependent diabetes (type1), Interstitial cystitis, Juvenile arthritis, Juvenile diabetes, Kawasaki syndrome, Kuttner's tumour, Lambert-Eaton syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus (SLE), Lyme disease, chronic, Mediastinal fibrosis, Meniere's disease, Microscopic polyangiitis, Mikulicz's syndrome, Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multifocal fibrosclerosis, Multiple sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neuromyelitis optica (Devic's), Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Ormond's disease (retroperitoneal fibrosis), Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paraproteinemic polyneuropathies, Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis (peripheral uveitis), Pemphigus vulgaris, Periaortitis, Periarthritis, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia, POEMS syndrome, Polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, Polymyalgia rheumatic, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Progesterone dermatitis, Primary biliary cirrhosis, Primary sclerosing cholangitis, Psoriasis, Psoriatic arthritis, Idiopathic pulmonary fibrosis, Pyoderma gangrenosum, Pure red cell aplasia, Raynauds phenomenon, Reflex sympathetic dystrophy, Reiter's syndrome, Relapsing polychondritis, Restless legs syndrome, Retroperitoneal fibrosis (Ormond's disease), Rheumatic fever, Rheumatoid arthritis, Riedel's thyroiditis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Sjogren's syndrome, Sperm & testicular autoimmunity, Stiff person syndrome, Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombotic, thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome, Transverse myelitis, Ulcerative colitis, Undifferentiated connective tissue disease (UCTD), Uveitis, Vasculitis, Vesiculobullous dermatosis, Vitiligo, Waldenstrom Macroglobulinaemia, Warm idiopathic haemolytic anaemia and Wegener's granulomatosis (now termed Granulomatosis with Polyangiitis (GPA)).

In one embodiment the autoimmune disease is selected from immune thrombocytopenia (ITP), chronic inflammatory demyelinating polyneuropathy (CIDP), Kawasaki disease and Guillain-Barre syndrome (GBS).

- 5 In one embodiment the monomeric fusion proteins of the invention are employed in the treatment or prophylaxis of epilepsy or seizures.

In one embodiment the monomeric fusion proteins and fragments according to the disclosure are employed in the treatment or prophylaxis of multiple sclerosis.

10

The monomeric fusion protein according to the present disclosure may be employed in treatment or prophylaxis.

- The monomeric fusion protein of the present invention may also be used in
15 diagnosis, for example in the *in vivo* diagnosis and imaging of disease states involving Fc-receptors, such as B-cell related lymphomas.

FIGURE LEGENDS

20

Figure 1

- 1(a)** Fusion proteins in which the tailpiece cysteine residue has been deleted or substituted with another amino acid residue are expressed predominantly as monomers and are capable of concentration-dependent assembly into hexamers.
25 **1(b)** Control proteins lacking the modified tailpiece are unable to assemble into hexamers, even at the highest concentrations tested.
1(c) Control proteins comprising an unmodified tailpiece with an intact cysteine residue are expressed predominantly in hexameric form.

Figure 2

30 Example amino acid sequences of a polypeptide chain of a monomeric fusion protein. In each sequence, the tailpiece sequence is underlined, and any mutations are shown in bold and underlined. The hinge is in bold. In constructs comprising a CH4 domain from IgM, this region is shown in italics.

Figure 3

Example amino acid and DNA sequences for antibody fusion proteins of the invention. The tailpiece sequence is underlined.

5

Figure 4

Chromatographs demonstrating that full length antibody fusion proteins comprising the modified tailpiece of the invention exhibit concentration-dependent multimerisation into a high molecular weight species (HMWS).

10 **4(a)** Rituximab IgG1 FL IgM tp C575S

4(b) Trastuzumab IgG1 FL IgM tp C575S

Figure 5

15 Complement killing of cells by a fusion protein of the invention. The graphs show complement killing of CD20 positive Raji cells and CD20 positive Ramos cells. The results demonstrate that the anti-CD20 antibody rituximab shows enhanced CDC when modified with the tailpiece of the invention.

20 **EXAMPLES**

Example 1: Molecular Biology

DNA sequences were assembled using standard molecular biology methods, including PCR, restriction-ligation cloning, point mutagenesis (Quikchange) and Sanger sequencing. Expression constructs were cloned into expression plasmids (pNAFL, pNAFH) suitable for both transient and stable expression in CHO cells. Other examples of suitable expression vectors include pCDNA3 (Invitrogen).

30 Diagrams showing example amino acid sequences of a polypeptide chain of a monomeric fusion protein are provided in Figure 2. In each sequence, the tailpiece sequence is underlined, and any mutations are shown in bold and underlined. The hinge is in bold. In constructs comprising a CH4 domain from IgM, this region is shown in italics.

Example 2: Expression

- 5 Small scale expression was performed using 'transient' expression of HEK293 or CHO cells transfected using lipofectamine or electroporation. Cultures were grown in shaking flasks or agitated bags in CD-CHO (Lonza) or ProCHO5 (Life Technologies) media at scales ranging from 50 – 2000ml for 5-10 days. Cells were removed by centrifugation and culture supernatants were stored at 4°C until purified.
- 10 Preservatives were added to some cultures after removal of cells.

Example 3: Purification and analysis

- 15 Proteins were purified from culture supernatants after checking / adjusting pH to be ≥ 6.5 , by protein A chromatography with step elution using a pH3.4 buffer. Eluate was immediately neutralised to $\sim$ pH7.0 using 1M Tris pH8.5.

Samples were concentrated in 0.1M sodium citrate buffer pH7.5 using a pressure stirred cell, and 10kDa cut-off membrane to $>100\text{mg/ml}$. Samples were then diluted in PBS to a range of different final concentrations as shown in Table 3, before SE-HPLC analysis of multimeric form as described below. Endotoxin was tested using the limulus amoebocyte lysate (LAL) assay. Samples used in assays were $<1\text{EU/mg}$.

25

SE-HPLC analysis of multimeric form**TSK-G3000**

- Proteins were analysed using size exclusion HPLC, the column used was 15 ml TSKgel -G3000SW (Tosoh) on system Agilent 1100 Series. The mobile phase was 0.2 M sodium phosphate, pH7.0, flow rate 1ml/minute, 50 μg protein injected. Signal was detected using a UV absorbance detector at 280 nm wavelength.
- 30

uPLC

Proteins were analysed using size exclusion HPLC, the column used was 2.5ml BEH 200 (Waters) on system Agilent 1100 Series. The mobile phase was 0.2 M sodium phosphate, pH7.0, flow rate 0.4 ml/minute, 2.5 and 5µg protein injected. Signal was detected using a Fluorescence detector Excitation: 350 nm, Emission: 390nm.

5

Superdex 200 SEC-MALS

Proteins were analysed using size exclusion HPLC, the column used was 24ml Superdex 200 10/300 (GE) on system Agilent 1100 Series. The mobile phase was 10mM HEPES, 100mM NaCl pH7.5, flow rate 0.5 ml/minute, 100µg protein injected. Signal was detected using a UV absorbance detector at 280 nm wavelength and additional refractive index detector (Viscotek VE3580) and MALS detector (Viscotek SEC-MALS 20, Malvern)

10

Results

15

The results demonstrated that the fusion proteins of the invention are synthesised predominantly as monomers, and are capable of assembly into multimers at high concentrations, as shown in Figure 1.

20 Hinge-Fc-tailpiece-C575S:

Fusion proteins in which the tailpiece cysteine residue has been deleted or substituted with another amino acid residue are expressed predominantly as monomers and are capable of concentration-dependent assembly into hexamers. Figure 1(a).

25

Control protein: Hinge-Fc (no tailpiece):

Control proteins lacking the modified tailpiece are unable to assemble into hexamers, even at the highest concentrations tested. Figure 1(b).

30 Control protein: Hinge-Fc-tailpiece (C575):

Control proteins comprising an unmodified tailpiece with an intact cysteine residue are expressed predominantly in hexameric form. Figure 1(c).

Example 5: Complement Dependent Cytotoxicity (CDC)

Full-length antibody constructs were prepared as shown in Table 3 below. Rituximab and trastuzumab are well-known antibodies that recognise CD20 and HER2/neu respectively. CA1151 recognises a clostridium difficile exotoxin and was included in the present study for comparison as a negative control antibody. Amino acid and DNA sequences for the antibody constructs are provided in Figure 3.

Table 3

Antibody construct
rituximab IgG1 IgM tailpiece C575
rituximab IgG1 IgM tailpiece C575S
rituximab IgG1 wild type
CA1151 IgG1 IgM tailpiece C575
CA1151 IgG1 IgM tailpiece C575S
CA1151 IgG1 wild type
trastuzumab IgG1 IgM tailpiece C575
trastuzumab IgG1 IgM tailpiece C575S
trastuzumab IgG1 wild type

Concentration and analytical HPLC

Proteins were concentrated by centrifugation using Amicon Ultra-15 Centrifugal Filter Units. Samples were centrifuged at 4000 RPM until desired concentration was reached. Proteins were analysed using size exclusion HPLC, the column used was 15 ml TSKgel -G3000SW (Tosoh) on system Agilent 1100 Series. The mobile phase was 0.2 M sodium phosphate, pH7.0, flow rate 1ml/minute, 50µg protein injected. Signal was detected using a UV absorbance detector at 280 nm wavelength.

The results demonstrated that full length antibody fusion proteins comprising the modified tailpiece of the invention exhibit concentration-dependent multimerisation into a high molecular weight species (HMWS). Figure 4.

CDC assay

The biological activity of the modified antibodies was examined using a complement dependent cytotoxicity assay. Target cells (50×10^5) were incubated with an antibody concentration series in a flat-bottom 96-well plate at a final volume of 100 μ l, and allowed to opsonise for 15 minutes at room temperature. Opsonised target cells
5 were incubated with normal human serum at a final concentration of 30% (42 μ l added) and incubated in a 37°C incubator for 30 minutes. Cell lysis measured as a fraction of PI+ cells determined by BD FACSCalibur flow cytometer. Graphs were plotted using GraphPad Prism software.

- 10 Results for a fusion protein comprising the anti-CD20 antibody rituximab are shown in Figure 5. The graphs show complement killing of CD20 positive Raji cells and CD20 positive Ramos cells. The results demonstrated that rituximab shows enhanced CDC when modified with the tailpiece of the invention.

15

CLAIMS

1. A monomeric fusion protein comprising an antibody Fc-domain comprising two heavy chain Fc-regions derived from IgG;
- 5 wherein one or each heavy chain Fc-region is fused at its C-terminal to an antibody tailpiece; and
wherein the tailpiece cysteine residue is modified to prevent disulphide bond formation.
- 10 2. The monomeric fusion protein of claim 1, wherein the cysteine residue is deleted, substituted, or blocked with a thiol capping agent.
3. The monomeric fusion protein of claim 1 or claim 2, further comprising an antigen binding region.
- 15 4. The monomeric fusion protein of claim 3, wherein the antigen binding region is a VH or a VL antigen binding region.
5. The monomeric fusion protein of claim 3 or claim 4, wherein the antigen binding
20 region is selected from the group consisting of Fab, scFv, dAb, VHH, and DARPin.
6. The monomeric fusion protein of any one of claims 1 to 5, further comprising a fusion partner.
- 25 7. The monomeric fusion protein of claim 6, wherein the fusion partner is selected from the group consisting of antigen, pathogen-associated molecular pattern (PAMP), drug, ligand, receptor, cytokine or chemokine.
8. The monomeric fusion protein of any preceding claim, wherein the heavy chain
30 Fc-region comprises CH2 and CH3 domains derived from IgG1, IgG2, IgG3, or IgG4.
9. The monomeric fusion protein of any preceding claim, wherein the tailpiece is derived from human IgM or IgA.

10. The monomeric fusion protein of any preceding claim, wherein each heavy chain Fc-region possesses a hinge region at its N-terminus.

11. The monomeric fusion protein of claim 10, wherein the hinge region
5 comprises the mutated sequence CPPC.

12. The monomeric fusion protein of any preceding claim, comprising one or more mutations which alter its Fc-receptor binding profile.

10 13. The monomeric fusion protein of claim 1 wherein each heavy chain Fc-region comprises or consists of the sequence given in amino acids 6 to 222 of any one of SEQ ID NOs 26 to 29, or the sequence given in amino acids 6 to 333 of SEQ ID NOs 30 or 31, or the sequence given in amino acids 6 to 221 of any one of SEQ ID NOs 32 to 35, or the sequence given in amino acids 6 to 332 of SEQ ID NOs 36 or 37.

15

14. The monomeric fusion protein of claim 13 wherein each heavy chain Fc-region further comprises a hinge region having a sequence given in any one of SEQ ID NOs: 3 to 25.

20 15. The monomeric fusion protein of claim 1 wherein each polypeptide monomer unit comprises or consists of two identical polypeptide chains each polypeptide chain comprising or consisting of the sequence given in any one of SEQ ID NOs 26 to 37.

25 16. The monomeric fusion protein of any preceding claim which is a purified monomer.

17. A mixture comprising a monomeric fusion protein according to any preceding claim and a multimer, said multimer comprising two or more monomer units.

30 18. The mixture of claim 17, comprising greater than 55% monomer.

19. An isolated DNA sequence encoding a polypeptide chain of a monomeric fusion protein according to any of claims 1 to 16, or a component part thereof.

20. A cloning or expression vector comprising one or more DNA sequences according to claim 18 or claim 19.

21. A host cell comprising one or more cloning or expression vectors according to claim 20.

22. A process for the production of a monomeric fusion protein according to any of claims 1-16, comprising culturing a host cell according to claim 21 under conditions suitable for protein expression, and isolating and optionally purifying the monomeric fusion protein.

23. A pharmaceutical composition comprising a monomeric fusion protein of any one of claims 1 to 16, in combination with a pharmaceutically acceptable excipient, diluent or carrier.

24. The pharmaceutical composition of claim 23, comprising a component that stabilises the monomeric form of the protein or increases the ratio of monomer to multimer in a mixture.

25. The pharmaceutical composition of claim 24, additionally comprising other active ingredients.

26. The monomeric fusion protein of any one of claims 1 to 16 or the pharmaceutical composition of any one of claims 23 to 25 for use in therapy.

27. Use of the monomeric fusion protein of any one of claims 1 to 16 for the preparation of a medicament.

28. The monomeric fusion protein of any one of claims 1 to 16 or the pharmaceutical composition of any one of claims 23 to 25 for use in the treatment of cancer.

29. Use of the monomeric fusion protein of any one of claims 1 to 16 for the preparation of a medicament for the treatment of cancer.

30. The monomeric fusion protein of any one of claims 1 to 16 or the pharmaceutical composition of any one of claims 23 to 25, for use in the treatment of immune disorders.

5

31. Use of the monomeric fusion protein of any one of claims 1 to 16 for the preparation of a medicament for the treatment of immune disorders.

Figure 1(a)

Hinge-Fc-tailpiece-C575S





Figure 1(a) continued

Hinge-Fc-tailpiece-C575S





Figure 1(a) continued

Hinge-Fc-tailpiece-C575S





Figure 1(a) continued
Hinge-Fc-tailpiece-C575S





Figure 1(b)

Control protein: Hinge-Fc (no tailpiece)





Figure 1(b) continued

Control protein: Hinge-Fc (no tailpiece)






Figure 1(b) continued

Control protein: Hinge-Fc (no tailpiece)

**Figure 1(c)**

Control protein: Hinge-Fc-tailpiece (C575)




Figure 2**1. Human IgG1 Fc-protein IgM tp C575X SEQ ID NO: 26**

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVVFSCSVMHEALHNHYTQKSLSLSPGKPTLYNVSLVMSDTAGTXY

2. Human IgG4 Fc-protein IgM tp C575X SEQ ID NO: 27

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
VDKSRWQEGNVVFSCSVMHEALHNHYTQKSLSLGLGKPTLYNVSLVMSDTAGTXY

3. Human IgG1 Fc-protein IgA tp C495X SEQ ID NO: 28

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVVFSCSVMHEALHNHYTQKSLSLSPGKPTHVNVSVVMAEVDGTXY

4. Human IgG4 Fc-protein IgA tp C495X SEQ ID NO: 29

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
VDKSRWQEGNVVFSCSVMHEALHNHYTQKSLSLGLGKPTHVNVSVVMAEVDGTXY

5. Human IgG1 Fc-protein Cμ4-IgM tp C575X SEQ ID NO: 30

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVVFSCSVMHEALHNHYTQKSLSLSPGKVALHRPDVYLLPPAREQLNLRESATI
TCLVTGFSPADV FVQWMQRGQPLSPEKYVTSAPMPEPQAPGRYFAHSILTVSEEEWNTGETY
TCVAHEALPNRV TERTV D KSTGKPTLYNVSLVMSDTAGTXY

Figure 2 continued**6. Human IgG4 Fc-protein C μ 4-IgM tp C575X SEQ ID NO: 31**

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
 KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
 TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
 VDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLKVALHRPDVYLLPPAREQLNLRESATI
 TCLVTGFSPADV FVQWMQRGQPLSPEKYVTSAPMPEPQAPGRYFAHSILTVSEEWNTGETY
 TCVAHEALPNRV TERTVDKSTGKPTLYNVSLVMSDTAGTXY

7. Human IgG1 Fc-protein IgM tp Δ C575 SEQ ID NO: 32

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
 KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
 TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
 VDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGKPTLYNVSLVMSDTAGTY

8. Human IgG4 Fc-protein IgM tp Δ C575 SEQ ID NO: 33

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
 KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
 TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
 VDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLKPTLYNVSLVMSDTAGTY

9. Human IgG1 Fc-protein IgA tp Δ C495 SEQ ID NO: 34

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
 KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
 TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
 VDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGKPTHVNVSVVMAEVDGTY

10. Human IgG4 Fc-protein IgA tp Δ C495 SEQ ID NO: 35

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
 KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
 TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
 VDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLKPTHVNVSVVMAEVDGTY

Figure 2 continued**11. Human IgG1 Fc-protein C μ 4-IgM tp Δ C575 SEQ ID NO: 36**

CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY
TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGKVALHRPDVYLLPPAREQLNLRESATI
TCLVTGFSFSPADVFFVQWMQRGQPLSPEKYVTSAPMPEPQAPGRYFAHSILTVSEEWNTGETY
TCVAHEALPNRVTERTVDKSTGKPTLYNVSLVMSDTAGTY

12. Human IgG4 Fc-protein C μ 4-IgM tp Δ C575 SEQ ID NO: 37

CPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
TLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLT
VDKSRWQEGNVFSQSVMEALHNHYTQKSLSLSPGKVALHRPDVYLLPPAREQLNLRESATI
TCLVTGFSFSPADVFFVQWMQRGQPLSPEKYVTSAPMPEPQAPGRYFAHSILTVSEEWNTGETY
TCVAHEALPNRVTERTVDKSTGKPTLYNVSLVMSDTAGTY

Figure 3**Rituximab****Leader sequence DNA Heavy chain SEQ ID NO: 38**

atgggttgagcttgattctcctgttccttggtggcggtggcaacgcgggtgctgagc

Leader sequence protein Heavy chain SEQ ID NO: 39

MGWSLILLFLVAVATRVLS

Leader sequence DNA Light Chain SEQ ID NO: 40

atggattccaagtgcagatcatctcgttcctcctcatctccgcctccgtcatcatgtctaggggc

Leader sequence protein Light Chain SEQ ID NO: 41

MDFQVQIISFLLISASVLM SRG

Figure 3 continued**Rituximab IgG1 DNA****heavy chain SEQ ID NO: 42**

caagtgcagctgcagcagccaggagccgaactcgtgaagcccggtgcttcagtgaagatgtcctgcaaagcctccgggtacactttc
acctcgtacaacatgcactgggtcaagcagacccccggtagaggactggagtggatcggcgccatctaccctgggaacggagacac
ctctacaacaaaagttcaagggaaaggccacactgaccgtgacaagtcgtcctcgaccggtacatgcagctgtcctctctgac
ttccgaggacagcgctgtgtactactgtgcccggtccacttactacggcgagattgggtactttaacgtctggggagccggcaccacc
gtgctcgtgactgtctcgagtgttctacaaagggcccatcggtcttccccctggcaccctctccaagagcacctctgggggcacag
cggccctggggtgctgtgtaaggactacttccccgaacctgtgacgggtgtcgtggaactcaggcgccctgaccagcggcgtgcaca
ccttcccggtgtcctacagtcctcaggactctactccctcagcagcgtggtgaccgtgccctccagcagcttgggcaccagacctac
atctgcaacgtgaatcacaagcccagcaacaccaaggctcgataagaaagttgagcccaaattctgtgacaaaactcacacatgcc
accgtgccagcacctgaactcctggggggaccgtcagcttctcttcccccaaaaaccaaggacacctcatgatctcccgacc
cctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgca
taatgccaagacaaaagccgcgaggagcagtagacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggc
tgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacctctccaaagccaaagggca
gccccgagaaccacaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctgtgtaag
gcttctatcccagcgacatcgcggtggagtgggagagcaatgggcagccggagaaactacaagaccagcctccctgctggac
tcgacggctccttctctctacagcaagctcacgtggacaagagcaggtggcagcaggggaacgtcttctcatgctcgtgatgc
atgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaa

light chain SEQ ID NO: 43

cagattgtgtgtcgcaatcccctgcgatcctgtcggcctcaccggagaaaaggtcacgatgacttgccgcgttctctcctcgtgc
atacattcactggtttcagcagaagccgggaagcagcccaaagccttggatctatgccacctccaacctggcatcgggtgtccccgtg
cggttcagcggtagcgggtccggaaccagctactccttgaccatttcgagagtggaaagccgaggacgcggccacctactactgtcag
cagtggttagcaaacccgccgaccttcggagggggcactaagctggagatcaaactacggttagcggccccatctgtcttcatctt
ccgcatctgatgagcagttgaaatctggaactgcctctgttgtgtcctgctgaataacttctatcccagagaggccaaagtacagt
ggaagggtggataacgccctccaatcgggtaactcccaggagagtgacagagcaggacagcaaggacagcacctacagcctcag
cagcacctgacgctgagcaaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgccc
gtcacaagagcttcaacaggggagagtgt

Figure 3 continued**Rituximab IgG1 Protein****heavy chain** SEQ ID NO: 44

QVQLQQPGAELVKPGASVKMSCKASGYTFTSYNMHWVKQTPGRGLEWIGAIYPGNGDTSYNQKFKGK
ATLTADKSSSTAYMQLSSLTSEDSAVYYCARSTYYGGDWYFNVWGAGTTVLTVSSASTKGPSVFPLAPS
SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ
PREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGSSFFLYSKLTVD
KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

light chain SEQ ID NO: 45

QIVLSQSPAILSASPGEKVTMTCRASSSVSYIHWFAQKPGSSPKPWYATSNLASGVPVRFSGSGSGTSYSL
TISRVEAEDAATYYCQQWTSNPPTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 3 continued**Rituximab IgG1 IgM tp DNA****heavy chain SEQ ID NO: 46**

caagtgcagctgcagcagccaggagccgaactcgtgaagcccggtgcttcagtgaagatgtcctgcaaagcctccgggtacactttc
acctcgtacaacatgcactgggtcaagcagacccccggtagaggactggagtggatcggcgccatctaccctgggaacggagacac
ctctacaacaaaagttcaagggaaaggccacactgaccgtgacaagtcgtcctcgaccggtacatgcagctgtcctctctgac
ttccgaggacagcgtgtgtactactgtgcccggtccacttactacggcggagattgggtactttaacgtctggggagccggcaccacc
gtgctcgtgactgtctcgagtgttctacaaagggcccatcggtcttccccctggcacccctcctccaagagcacctctgggggcacag
cggccctggggtgcttggtcaaggactacttccccgaacctgtgacgggtgctgtggaactcaggcgcctgaccagcggcgtgcaca
ccttccgggtgtcctacagtctcaggactctactccctcagcagcgtggtgaccgtgcctccagcagcttgggcaccagacctac
atctgcaacgtgaatcacaagcccagcaacaccaaggtcgataagaaagttgagccaaatcttgtgacaaaactcacacatgccc
accgtgccagcacctgaactcctggggggaccgtcagcttcttcttccccccaaaacccaaggacacccctcatgatctcccgacc
cctgaggtcacatgcgtggtgggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgca
taatgccaagacaaagccgcgaggaggagcagtaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggc
tgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacatctccaaagccaaagggca
gccccgagaaccacaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaa
gcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgctcccggtgctggac
tccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
atgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaaccgaccctgtataacgtgagcctggtgatga
gcgataccgcgggcacctgctat

light chain SEQ ID NO: 47

cagattgtgtgtcgcaatcccctgcgatcctgtcggcctcacccggagaaaaggtcacgatgacttgccgcgttctcctccgtgtc
atacattcactggtttcagcagaagccgggaagcagcccaaagccttgatctatgccacctccaacctggcatccggtgtccccgtg
cggttcagcggtagcgggtccggaaccagctactccttgaccatttcgagagtggaaagccgaggacgcggccacctactactgtcag
cagtggtactagcaaccccgaccttcggagggggcactaagctggagatcaaactacggttagcggccccatctgtcttcatcttc
ccgcatctgatgagcagttgaaatctggaactgcctctgtgtgtgcctgtgaataacttctatcccagagaggccaaagtacagt
ggaaggtggataacgcctccaatcgggtaactcccaggagagtgacagagcaggacagcaaggacagcacctacagcctcag
cagcacctgacgctgagcaaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgccc
gtcacaaagagcttcaacaggggagagtg

Figure 3 continued**Rituximab IgG1 IgM tp Protein****heavy chain** SEQ ID NO: 48

QVQLQQPGAELVKPGASVKMSCKASGYTFTSYNMHWVKQTPGRGLEWIGAIYPGNGDTSYNQKFKGK
ATLTADKSSSTAYMQLSSLTSEDSAVYYCARSTYYGGDWYFNVWGAGTTVLTVSSASTKGPSVFPLAPS
SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ
PREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVD
KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKPTLYNVSLVMSDTAGTCY

light chain SEQ ID NO: 49

QIVLSQSPAILSASPGEKVTMTCRASSSVSYIHWFQQKPGSSPKPWYATSNLASGVPVRFSGSGSGTSYSL
TISRVEAEDAATYYCQQWTSNPPTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 3 continued**Rituximab IgG1 IgM tp C575S DNA****heavy chain SEQ ID NO: 50**

caagtgcagctgcagcagccaggagccgaactcgtgaagcccggtcctcagtgaagatgtcctgcaaagcctccgggtacactttc
acctcgtacaacatgcactgggtcaagcagacccccggtagaggactggagtggatcggcgccatctaccctgggaacggagacac
ctcctacaacaaaagttcaagggaaaggccacactgaccgtgacaagtcgtcctcgaccggtacatgcagctgtcctctgtac
ttccgaggacagcgtgtgtactactgtgcccggtccacttactacggcggagattgggtactttaacgtctggggagccggcaccacc
gtgctcgtgactgtctcgagtgttctacaaagggcccatcggtcttccccctggcacctcctccaagagcacctctgggggcacag
cggccctggggtcctgggtcaaggactacttccccgaacctgtgacgggtgctgtggaactcaggcgcctgaccagcggcgtgcaca
ccttccgggtgtcctacagtctcaggactctactccctcagcagcgtggtgaccgtgccctccagcagcttgggcacccagacctac
atctgcaacgtgaatcacaagcccagcaacaccaaggctgataagaaagttgagcccaaacttgtgacaaaactcacacatgcc
accgtgccagcacctgaactcctggggggaccgtcagcttctcttccccccaaaacccaaggacacccctcatgatctcccgacc
cctgaggtcacatgcgtgggtgggtgacgtgagccacgaagaccctgaggtcaagttcaactggtagctggacggcgtggaggtgca
taatgccaagacaaaagccgcgaggagagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggc
tgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggca
gccccgagaaccacaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaa
gttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgctctccgtgctggac
tccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
atgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaaccgaccctgtataacgtgagcctgggtgatga
gcgataccgcgggcaccagctat

light chain SEQ ID NO: 51

cagattgtgtgtcgcaatcccctgcgatcctgtcggcctcaccggagaaaaaggtcacgatgacttgccgcgttctcctccgtgtc
atacattcactggtttcagcagaagccgggaagcagcccaaagccttggatctatgccacctccaacctggcatccggtgtccccgtg
cggttcagcggtagcgggtccggaaccagctactccttgaccatttcgagagtggaaagccgaggacgcggccacctactactgtcag
cagtgactagcaaccccgaccttcggagggggcactaagctggagatcaaactacggttagcggccccatctgtcttcatcttc
ccgcatctgatgagcagttgaaatctggaactgcctctgttgtgtgcctgtgaataacttctatcccagagaggccaaagtacagt
ggaaggtggataacgcctccaatcgggttaactcccaggagagtgctacagagcaggacagcaaggacagcacctacagcctcag
cagcacctgacgctgagcaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgccc
gtcacaaagagcttcaacaggggagagtg

Figure 3 continued**Rituximab IgG1 IgM tp C575S Protein****heavy chain** SEQ ID NO: 52

QVQLQQPGAELVKPGASVKMSCKASGYTFTSYNMHWVKQTPGRGLEWIGAIYPGNGDTSYNQKFKGK
ATLTADKSSSTAYMQLSSLTSEDSAVYYCARSTYYGGDWYFNVWGAGTTVLTVSSASTKGPSVFPLAPS
SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ
PREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD
KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKPTLYNVSLVMSDTAGTSY

light chain SEQ ID NO: 53

QIVLSQSPAILSASPGEKVTMTCRASSSVSYIHWFQQKPGSSPKPWYATSNLASGVPVRFSGSGSGTSYSL
TISRVEAEDAATYYCQQWTSNPPTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 3 continued**Herceptin**

Leader sequence herceptin DNA Heavy chain SEQ ID NO: 54

atggagtggagctgggtctttctgttcttcctctccgtgaccactggggttcatagc

Leader sequence herceptin protein Heavy chain SEQ ID NO: 55

MEWSWVFLFFLSVTTGVHS

Leader sequence Herceptin DNA Light chain SEQ ID NO: 56

atgatgtcctctgctcagttccttggctcctgttgctctgtttcaaggtaccagatgt

Leader sequence Herceptin protein Light chain SEQ ID NO: 57

MMSSAQFLGLLLLCFQTRC

Figure 3 continued**Herceptin IgG1 DNA****heavy chain SEQ ID NO: 58**

gaagtgcagctggtggaatcaggtgggggattggttcaacctggaggaagtctccgtctgtcttgagcagcctctgggttcaacatca
aggatacctacatacactgggtgagacaggcaccaggtaaaggattggagtgggtggcaaggatctacctaccaatggctacacc
cgatatgccgacagtgtgaaaggcaggttcacgatttccgctgacacaagcaagaacacagcctacctgcagatgaactactgcg
agcagaggatacggctgtctattactgctctagatggggaggggacggtttctacgctatggactattggggacagggcactcttg
acagtctcgagtgttctacaaagggccatcggtcttccccctggcaccctcctcaagagcacctctgggggcacagcggccctgg
gctgcctggtcaaggactacttccccgaacctgtgacgggtgctgtggaactcaggcgccctgaccagcggcgtgcacaccttccgg
ctgtcctacagtcctcaggactctactccctcagcagcgtggtgaccgtgccctccagcagcttgggcacccagacctacatctgcaa
cgtgaatcacaagcccagcaacaccaaggtcgataagaaagttgagccaaatcttgtagaaaaactcacatgccaccgtgcc
cagcacctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacacccctcatgatctccggacccctgaggt
cacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgcca
agacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcacctcctgcaccaggactggctgaatggc
aaggagtacaagtgcaaggtctcaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaagggcagccccgag
aaccacaggtgtacacctgcccccatcccggtgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatc
ccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccggtgctggactccgacgg
ctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggct
ctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaa

light chain SEQ ID NO: 59

gatatccagatgactcagagtccaagtagtctcagtgccagtgtaggtgatagggttaactatcacttgtcgtgccagtcaggatgtta
atactgctgtagcctggtatcagcagaaaccaggtaaagccccaaaactactcatctattcggcctccttcttactctggagtacca
tctagattcagtggttagcagaagtggtagtattcactctgactatcagtagtctccagccagaagatttcgccacttactactgcca
gcaacattatactactcctcccacgttcggtcagggtagtaagtagaaatcaaactacggtacggccccatctgtcttcatcttcc
cgccatctgatgagcagttgaaatctggaactgcctctgttggtgctgtaataacttctatcccagagaggccaaagtacagtg
gaaggtggataacgccctccaatcgggtaactcccaggagagtgacacagagcaggacagcaaggacagcacctacagcctcagc
agcacctgacgtgagcaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgccg
tcacaaagagcttcaacaggggagagtg

Figure 3 continued**Herceptin IgG1 Protein****heavy chain** SEQ ID NO: 60

EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGYTRYADSVKGRFTI
SADTSKNTAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDYWGQGTLLTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

light chain SEQ ID NO: 61

DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSRSGTDF
TLTISSLQPEDFATYYCQQHYTTPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 3 continued**Herceptin IgG1 IgM tp DNA****heavy chain SEQ ID NO: 62**

gaagtgcagctggtggaatcaggtgggggattggttcaacctggaggaagtctccgtctgtcttgtagcagcctctgggttcaacatca
aggatacctacatacactgggtgagacaggcaccaggtaaaggattggagtggtggcaaggatctacctaccaatggctacacc
cgatatgccgacagtgtgaaaggcaggttcacgatttccgctgacacaagcaagaacacagcctacctgcagatgaactactgcg
agcagaggatacggctgtctattactgtcttagatggggaggggacggtttctacgctatggactattggggacagggcactcttg
acagtctcgagtgttctacaaaggcccatcggcttccccctggcaccctcctcaagagcacctctgggggcacagcggccctgg
gctgcctggtcaaggactacttccccgaacctgtgacggtgtcgtggaactcaggcgcctgaccagcggcgtgcacaccttccgg
ctgtcctacagtcctcaggacttactcctcagcagcgtggtgaccgtgccctccagcagcttgggcaccagacctacatctgcaa
cgtgaatcacaagcccagcaacaccaaggtcgataagaaagttgagccaaatcttgtagacaaaactcacatgcccaccgtgcc
cagcacctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacacccctcatgatctcccgaccctgaggt
cacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgcca
agacaaagcccggggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggc
aaggagtacaagtgaagggtctccaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaagggcagccccgag
aaccacaggtgtacacctgcccccatcccggtgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatc
ccagcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccggtgctggactccgacgg
ctcttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggct
ctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaaccgacctgtataacgtgagcctggtgatgagcgatacc
gcgggcacctgctat

light chain SEQ ID NO: 63

gatatccagatgactcagagtccaagtagtctcagtgccagtgtaggtgataggtaactatcacttgtcgtgccagtcaggatgtta
atactgtgtagcctggtatcagcagaaaccaggtaaagccccaaaactactcatctattcggcctccttctactctggagtacca
tctagattcagtggttagcagaagtgtgactgatttactctgactatcagtagtctccagccagaagatttcgccacttactactgcca
gcaacattatactactctccacgttcggtcaggttactaaagtagaaatcaaactacggttagcggccccatctgtcttcatcttcc
cgccatctgatgagcagttgaaatctggaactgcctctgttgtgtgcctgtgaataacttctatccagagaggccaaagtacagtg
gaaggtggataacgcctccaatcgggtaactcccaggagagtgacacagagcaggacagcaaggacagcacctacagcctcagc
agcacctgacgtgagcaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcggccg
tcacaaagagcttcaacaggggagagtg

Figure 3 continued**Herceptin IgG1 IgM tp Protein****heavy chain** SEQ ID NO: 64

EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGYTRYADSVKGRFTI
SADTSKNTAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDYWGQGTLLTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK PTLYNVSLVMSDTAGTCY

light chain SEQ ID NO: 65

DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSRSGTDF
TLTISSLQPEDFATYYCQQHYYTTPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 3 continued**Herceptin IgG1 IgM tp C575S DNA****heavy chain** SEQ ID NO: 66

gaagtgcagctggtggaatcaggtgggggattggttcaacctggaggaagtctccgtctgtcttgtgcagcctctgggttcaacatca
aggatacctacatacactgggtgagacaggcaccaggtaaaggattggagtgggtggcaaggatctacctaccaatggctacacc
cgatatgccgacagtgtgaaaggcaggttcacgatttccgctgacacaagcaagaacacagcctacctgcagatgaactcactgcg
agcagaggatacggctgtctattactgctctagatggggaggggacggtttctacgctatggactattggggacagggcactcttgtg
acagtctcagtgcttctacaaagggccatcggtcttccccctggcaccctcctcaagagcacctctgggggcacagcgccctgg
gtgcctgtgaaggactacttccccgaacctgtgacgggtgctgtggaactcaggcgccctgaccagcggtgcacaccttccgg
ctgtcctacagtctcaggacttactccctcagcagcgtggtgaccgtgccctccagcagcttgggcacccagacctacatctgcaa
cgtgaatcacaagcccagcaacaccaaggctgataagaaagttgagccaaatcttgtgacaaaactcacacatgccaccgtgcc
cagcacctgaactcctggggggaccgtcagtccttcttccccccaaaacccaaggacacctcatgatctccggaccctgaggct
cacatgcgtggtggtggacgtgagccacgaagacctgagggtcaagttcaactggtacgtggacggcgtggaggtgcataatgcca
agacaaagccgaggaggagcagtaaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggc
aaggagtacaagtgaaggcttccaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaagggcagccccgag
aaccacaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctgtgtaaaggcttctatc
ccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacgg
ctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttcttcatgctccgtgatgcatgaggct
ctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaaccgacctgtataacgtgagcctggtgatgagcgataacc
gcgggacaccagctat

light chain SEQ ID NO: 67

gatatccagatgactcagagtccaagtagtctcagtgccagtgtaggtgataggtaactatcacttgtcgtgccagtcaggatgtta
atactgctgtagcctggtatcagcagaaaccaggtaaagccccaaaactactcatctattcggcctccttctactctggagtacca
tctagattcagtggttagcagaagtgtgactgatttactctgactatcagtagtctccagccagaagatttcgccacttactactgcca
gcaacattatactactctcccacgttcgggtcagggtactaaagtagaaatcaaactacggttagcgccccatctgttcttcatctcc
cgccatctgatgagcagttgaaatctggaactgcctctgttgtgtgcctgtgaataacttctatcccagagaggccaaagtacagtg
gaaggtggataacgccctccaatcgggtaactcccaggagagtggtcacagagcaggacagcaaggacagcacctacagcctcagc
agcacctgacgctgagcaaagcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgcccc
tcacaaagagcttcaacaggggagagtggt

Figure 3 continued**Herceptin IgG1 IgM tp C575S DNA****heavy chain** SEQ ID NO: 68

EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGYTRYADSVKGRFTI
SADTSKNTAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDYWGQGLTVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK PTLYNVSLVMSDTAGTSY

light chain SEQ ID NO: 69

DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSRSGTDF
TLTISSLQPEDFATYYCQQHYTTPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Figure 4(a)

Rituximab IgG1 FL IgM tp C575S




Figure 4(a) continued

Rituximab IgG1 FL IgM tp C575S




Figure 4(a) continued

Rituximab IgG1 FL IgM tp C575S





Figure 4(b)

Trastuzumab IgG1 FL IgM tp C575S




Figure 4(b) continued

Trastuzumab IgG1 FL IgM tp C575S




Figure 4(b) continued

Trastuzumab IgG1 FL IgM tp C575S





Figure 5

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/065914

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/28 C07K16/32
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SITIA R ET AL: "Developmental regulation of IgM secretion: The role of the carboxy-terminal cysteine", CELL, CELL PRESS, US, vol. 60, no. 5, 9 March 1990 (1990-03-09), pages 781-790, XP024245106, ISSN: 0092-8674, DOI: 10.1016/0092-8674(90)90092-S [retrieved on 1990-03-09] abstract; page 783-784; fig. 4 ----- -/--	1-31



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 September 2016

Date of mailing of the international search report

04/10/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bernhardt, Wiebke

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2016/065914

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHRISTINA BRUNKE ET AL: "Effect of a tail piece cysteine deletion on biochemical and functional properties of an epidermal growth factor receptor-directed IgA2 m(1) antibody", MABS, vol. 5, no. 6, 1 November 2013 (2013-11-01), pages 936-945, XP055300857, US ISSN: 1942-0862, DOI: 10.4161/mabs.26396 abstract p. 939, left-hand col. -----	1-31
Y	SMITH R ET AL: "Addition of a mu-tailpiece to IgG results in polymeric antibodies with enhanced effector functions including complement-mediated cytolysis by IgG4", THE JOURNAL OF IMMUNOLOGY, THE AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US, vol. 154, no. 5, 1 March 1995 (1995-03-01), pages 2226-2236, XP002170004, ISSN: 0022-1767 p. 2228, left-hand col., fig. 1 -----	1-31
X,P	WO 2015/132364 A1 (UCB BIOPHARMA SPRL [BE]) 11 September 2015 (2015-09-11) -----	19-21
Y,P	pages 36-38 claims	1-18, 22-31
X,P	WO 2015/132365 A1 (UCB BIOPHARMA SPRL [BE]) 11 September 2015 (2015-09-11) -----	19-21
Y,P	pages 36-38 claims	1-18, 22-31
Y	SORENSEN V ET AL: "Effect of the IgM and IgA secretory tailpieces on polymerization and secretion of IgM and IgG", THE JOURNAL OF IMMUNOLOGY, THE AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US, vol. 156, no. 8, 15 April 1996 (1996-04-15), pages 2858-2865, XP002111658, ISSN: 0022-1767 abstract page 2858 figures tables -----	1-31
Y	US 2004/175786 A1 (CHOI INGRID [DE] ET AL) 9 September 2004 (2004-09-09) paragraph [0010] figures 2b and 3 -----	1-31

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2016/065914

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2015132364 A1	11-09-2015	AU 2015226100 A1 TW 201619188 A UY 36021 A WO 2015132364 A1	25-08-2016 01-06-2016 30-09-2015 11-09-2015
WO 2015132365 A1	11-09-2015	AU 2015226101 A1 WO 2015132365 A1	25-08-2016 11-09-2015
US 2004175786 A1	09-09-2004	DE 10001372 A1 EP 1250443 A2 US 2004175786 A1 WO 0151644 A2	02-08-2001 23-10-2002 09-09-2004 19-07-2001