



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2203652 C 2008/03/25

(11)(21) **2 203 652**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1995/10/17
(87) Date publication PCT/PCT Publication Date: 1996/05/02
(45) Date de délivrance/Issue Date: 2008/03/25
(85) Entrée phase nationale/National Entry: 1997/04/24
(86) N° demande PCT/PCT Application No.: US 1995/013069
(87) N° publication PCT/PCT Publication No.: 1996/012497
(30) Priorité/Priority: 1994/10/25 (US08/328,520)

(51) Cl.Int./Int.Cl. *A61K 31/7125* (2006.01),
A61K 48/00 (2006.01), *C07H 21/00* (2006.01)

(72) Inventeurs/Inventors:
AGRAWAL, SUDHIR, US;
DIASIO, ROBERT B., US;
ZHANG, RUIWEN, US

(73) Propriétaire/Owner:
IDERA PHARMACEUTICALS, INC., US

(74) Agent: MBM & CO.

(54) Titre : UTILISATION D'OLIGONUCLEOTIDES SUBSTITUES EN 2' POUR FREINER L'EXPRESSION D'UN GENE
(54) Title: A METHOD OF DOWN-REGULATING GENE EXPRESSION

(57) **Abrégé/Abstract:**

Disclosed is a method of down-regulating the expression of a gene in an animal, wherein a pharmacological formulation comprising an oligonucleotide complementary to the gene is orally administered to an animal. The oligonucleotide administered has non-phosphodiester internucleotide linkages and includes at least one 2'-substituted ribonucleotide, the oligonucleotide inhibiting the expression of a product of the gene, thereby down-regulating the expression of the gene.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ :**A61K 31/70****A1**

(11) International Publication Number:

WO 96/12497

(43) International Publication Date:

2 May 1996 (02.05.96)

(21) International Application Number: PCT/US95/13069

(22) International Filing Date: 17 October 1995 (17.10.95)

(30) Priority Data:

08/328,520

25 October 1994 (25.10.94)

US

(71) Applicant: HYBRIDON, INC. [US/US]; One Innovation Drive,
Worcester, MA 01605 (US).(72) Inventors: AGRAWAL, Sudhir; 61 Lamplighter Drive,
Shrewsbury, MA 01545 (US). DIASIO, Robert, B.; 1225
Branchwater Lane, Birmingham, AL 35216 (US). ZHANG,
Ruiwen; 1824 Russet Woods Lane, Birmingham, AL 35244
(US).(74) Agents: KERNER, Ann-Louise et al.; Lappin & Kusmer, Two
Hundred State Street, Boston, MA 02109 (US).(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA,
CH, CN, CZ, DE, DK, EE, FI, GB, GE, HU, IS, JP, KE,
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK,
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,
SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT,
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD,
SZ, UG).**Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: USE OF 2'-SUBSTITUTED OLIGONUCLEOTIDES TO DOWN-REGULATING GENE EXPRESSION

(57) Abstract

Disclosed is a method of down-regulating the expression of a gene in an animal, wherein a pharmaceutical formulation comprising an oligonucleotide complementary to the gene is orally administered to an animal. The oligonucleotide administered has non-phosphodiester internucleotide linkages and includes at least one 2'-substituted ribonucleotide, the oligonucleotide inhibiting the expression of a product of the gene, thereby down-regulating the expression of the gene.

-1-

A METHOD OF DOWN-REGULATING GENE EXPRESSION

BACKGROUND OF THE INVENTION

The present invention relates to the control of gene expression. More particularly, this invention relates to the use of synthetic oligonucleotides to down-regulate the expression of a gene in an animal.

The potential for the development of an antisense oligonucleotide therapeutic approach was first suggested in three articles published in 1977 and 1978. Paterson et al. (*Proc. Natl. Acad. Sci. (USA)* (1977) 74:4370-4374) discloses that cell-free translation of mRNA can be inhibited by the binding of an oligonucleotide complementary to the mRNA. Zamecnik et al. (*Proc. Natl. Acad. Sci. (USA)* (1978) 75:280-284 and 285-288) discloses that a 13mer synthetic oligonucleotide that is complementary to a part of the Rous sarcoma virus (RSV) genome inhibits RSV replication in infected chicken fibroblasts and inhibits RSV-mediated transformation of primary chick fibroblasts into malignant sarcoma cells.

These early indications that synthetic oligonucleotides can be used to inhibit virus propagation and neoplasia have been followed by the use of synthetic oligonucleotides to inhibit a wide variety of viruses, such as HIV (see, e.g., U.S. Patent No. 4,806,463); influenza (see, e.g., Leiter et al. (1990) (*Proc. Natl. Acad. Sci. (USA)* 87:3430-3434); vesicular stomatitis virus (see,

-2-

e.g., Agris et al. (1986) *Biochem.* 25:6268-6275);
herpes simplex (see, e.g., Gao et al. (1990)
Antimicrob. Agents Chem. 34:808-812); SV40 (see, e.g.,
Birg et al. (1990) (*Nucleic Acids Res.* 18:2901-2908);
5 and human papilloma virus (see, e.g., Storey et
al. (1991) (*Nucleic Acids Res.* 19:4109-4114). The use
of synthetic oligonucleotides and their analogs as
antiviral agents has recently been extensively
reviewed by Agrawal (*Trends in Biotech.* (1992) 10:152-
10 158).

In addition, synthetic oligonucleotides have
been used to inhibit a variety of non-viral
pathogens, as well as to selectively inhibit the
15 expression of certain cellular genes. Thus, the
utility of synthetic oligonucleotides as agents to
inhibit virus propagation, propagation of non-
viral, pathogens and selective expression of
cellular genes has been well established.

20 Improved oligonucleotides have more recently
been developed that have greater efficacy in
inhibiting such viruses, pathogens and selective
gene expression. Some of these oligonucleotides
25 having modifications in their internucleotide
linkages have been shown to be more effective than
their unmodified counterparts. For example,
Agrawal et al. (*Proc. Natl. Acad. Sci. (USA)* (1988)
85:7079-7083) teaches that oligonucleotide
30 phosphorothioates and certain oligonucleotide
phosphoramidates are more effective at inhibiting
HIV-1 than conventional phosphodiester-linked
oligodeoxynucleotides. Agrawal et al. (*Proc. Natl.*

-3-

Acad. Sci. (USA) (1989) 86:7790-7794) discloses the advantage of oligonucleotide phosphorothioates in inhibiting HIV-1 in early and chronically infected cells.

5

In addition, chimeric oligonucleotides having more than one type of internucleotide linkage within the oligonucleotide have been developed. Pederson et al. (U.S. Patent Nos. 5,149,797 and 5,220,007 discloses chimeric oligonucleotides having an oligonucleotide phosphodiester or oligonucleotide phosphorothioate core sequence flanked by nucleotide methylphosphonates or phosphoramidates. Furdon et al. (*Nucleic Acids Res.* (1989) 17:9193-9204) discloses chimeric oligonucleotides having regions of oligonucleotide phosphodiesters in addition to either oligonucleotide phosphorothioate or methylphosphonate regions. Quartin et al. (*Nucleic Acids Res.* (1989) 17:7523-7562) discloses chimeric oligonucleotides having regions of oligonucleotide phosphodiesters and oligonucleotide methylphosphonates. Inoue et al. (*FEBS Lett.* (1987) 215(2):327-330) discloses chimeric oligonucleotides having regions of deoxyribonucleotides and 2'-O-methyl-ribonucleotides.

25

Many of these modified oligonucleotides have contributed to improving the potential efficacy of the antisense oligonucleotide therapeutic approach. However, certain deficiencies remain in the known oligonucleotides, and these deficiencies can limit the effectiveness of such

30

-4-

oligonucleotides as therapeutic agents. For example, Wickstrom (*J. Biochem. Biophys. Meth.* (1986) 13:97-102) teaches that oligonucleotide phosphodiester are susceptible to nuclease-mediated degradation, thereby limiting their bioavailability *in vivo*. Agrawal et al. (*Proc. Natl. Acad. Sci. (USA)* (1990) 87:1401-1405) teaches that oligonucleotide phosphoramidates or methylphosphonates when hybridized to RNA do not activate RNase H, the activation of which can be important to the function of antisense oligonucleotides. Thus, a need for methods of controlling gene expression exists which uses oligonucleotides with improved therapeutic characteristics.

Several reports have been published on the development of phosphorothioate-linked oligonucleotides as potential anti-AIDS therapeutic agents. Although extensive studies on chemical and molecular mechanisms of oligonucleotides have demonstrated the potential value of this novel therapeutic strategy, little is known about the pharmacokinetics and metabolism of these compounds *in vivo*.

Recently, several preliminary studies on this topic have been published. Agrawal et al. (*Proc. Natl. Acad. Sci. (USA)* (1991) 88:7595-7599) describes the intravenously and intraperitoneally administration to mice of a 20-mer phosphorothioate linked-oligonucleotide. In this study, approximately 30% of the administered dose

-5-

was excreted in the urine over the first 24 hours with accumulation preferentially in the liver and kidney. Plasma half-lives ranged from about 1 hour ($t_{1/2\alpha}$) and 40 hours ($t_{1/2\beta}$), respectively.

5 Similar results have been reported in subsequent studies (Iversen (1991) *Anti-Cancer Drug Design* 6:531-538; Iversen (1994) *Antisense Res. Devel.* 4:43-52; and Sands (1994) *Mol. Pharm.* 45:932-943). However, stability problems may exist when oligonucleotides
10 are administered intravenously and intraperitoneally.

Thus, there remains a need to develop more effective therapeutic methods of down-regulating
15 the expression of genes which can be easily manipulated to fit the animal and condition to be treated, and the gene to be targeted. Preferably, these methods should be simple, painless, and precise in effecting the target gene.

20 SUMMARY OF THE INVENTION

The present invention provides a method of down-regulating the expression of a gene in an
25 animal which involves the administration of an oligonucleotide complementary to the gene via an oral route, thereby bypassing the complications which may be experienced during intravenous and other modes of in vivo administration.

30 It has been discovered that certain oligonucleotides (with other than phosphodiester bonds and having at least one 2'-substituted ribonucleotide) are relatively stable *in vivo*

-6-

following oral administration to an animal, and that these molecules are successfully absorbed from the gastrointestinal tract and distributed to various body tissues. This discovery has been
5 exploited to develop the present invention, which is a method of down-regulating the expression of a gene in an animal.

10 This method is also a means of examining the function of various genes in an animal, including those essential to animal development. Presently, gene function can only be examined by the arduous task of making a "knock out" animal such as a mouse. This task is difficult, time-consuming and
15 cannot be accomplished for genes essential to animal development since the "knock out" would produce a lethal phenotype. The present invention overcomes the shortcomings of this model.

20 In the method of the invention, a pharmaceutical formulation containing an oligonucleotide complementary to the targeted gene is orally administered in a pharmaceutically acceptable carrier to the animal harboring the
25 gene. The oligonucleotide inhibits the expression of the gene, thereby down-regulating its expression.

30 For purposes of the invention, the term "animal" is meant to encompass humans as well as other mammals, as well as reptiles amphibians, and insects. The term "oral administration" refers to the provision of the formulation via the mouth through ingestion, or via some other part of

-7-

the gastrointestinal system including the esophagus.

As used herein, the term "oligonucleotide" is meant to include polymers of two or more nucleotides or nucleotide analogs connected together via 5' to 3' internucleotide linkages which may include any linkages that are known in the antisense art. Such molecules have a 3' terminus and a 5' terminus. The particular oligonucleotide being administered include at least one 2'-substituted ribonucleotide, and are connected via non-phosphodiester internucleotide linkages.

For purposes of the invention, the term "2'-substituted oligonucleotide" refers to an oligonucleotide having a sugar attached to a chemical group other than a hydroxyl group at its 2' position. The 2'-OH of the ribose molecule can be substituted with -O-lower alkyl containing 1-6 carbon atoms, aryl or substituted aryl or allyl having 2-6 carbon atoms, e.g., 2'-O-allyl, 2'-O-aryl, 2'-O-alkyl (such as a 2'-O-methyl), 2'-halo, or 2'-amino, but not with 2'-H, wherein allyl, aryl, or alkyl groups may be unsubstituted or substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carbalkoxyl or amino groups.

The term "non-phosphodiester-linked oligonucleotide" as used herein is an oligonucleotide in which all of its nucleotides are covalently linked via a synthetic linkage,

-8-

i.e., a linkage other than a phosphodiester between the 5' end of one nucleotide and the 3' end of another nucleotide in which the 5' nucleotide phosphate has been replaced with any number of chemical groups. Preferable synthetic linkages include alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, phosphoramidates, phosphoramidites, phosphate esters, carbamates, carbonates, phosphate triesters, acetamdate, and carboxymethyl esters. In one preferred embodiment of the invention, the all of the nucleotides of the oligonucleotide comprises are linked via phosphorothioate and/or phosphorodithioate linkages.

In some embodiments of the invention, the oligonucleotides administered are modified. As used herein, the term "modified oligonucleotide" encompasses oligonucleotides with modified nucleic acid(s), base(s), and/or sugar(s) other than those found in nature. For example, a 3', 5'-substituted oligonucleotide is an oligonucleotide having a sugar which, at both its 3' and 5' positions is attached to a chemical group other than a hydroxyl group (at its 3' position) and other than a phosphate group (at its 5' position).

A modified oligonucleotide may also be one with added substituents such as diamines, cholesteryl, or other lipophilic groups, or a capped species. In addition, unoxidized or partially oxidized oligonucleotides having a

-9-

substitution in one nonbridging oxygen per nucleotide in the molecule are also considered to be modified oligonucleotides. Also considered as modified oligonucleotides are oligonucleotides having nuclease resistance-conferring bulky substituents at their 3' and/or 5' end(s) and/or various other structural modifications not found *in vivo* without human intervention are also considered herein as modified.

In one preferred embodiment of the invention, the oligonucleotide administered includes at least one 2'-substituted ribonucleotide at its 3' terminus. In some embodiments, all but four or five nucleotides at its 5' terminus are 2'-substituted ribonucleotides, and in some embodiments, these four or five unsubstituted 5' nucleotides are deoxyribonucleotides. In other aspects, the oligonucleotide has at least one 2'-substituted ribonucleotide at both its 3' and 5' termini, and in yet other embodiments, the oligonucleotide is composed of 2'-substituted ribonucleotides in all positions with the exception of at least four or five contiguous deoxyribonucleotide nucleotides in any interior position. Another aspect of the invention includes the administration of an oligonucleotide composed of nucleotides that are all 2'-substituted ribonucleotides. Particular embodiments include oligonucleotides having a 2'-O-alkyl-ribonucleotide such as a 2'-O methyl.

In another embodiment of the invention, the oligonucleotide administered has at least one

-10-

deoxyribonucleotide, and in a preferred embodiment, the oligonucleotide has at least four or five contiguous deoxyribonucleotides capable of activating RNase H.

5

The oligonucleotide administered is complementary to a gene of a virus, pathogenic organism, or a cellular gene in some embodiments of the invention. In some embodiments, the
10 oligonucleotide is complementary to a gene of a virus involved in AIDS, oral or genital herpes, papilloma warts, influenza, foot and mouth disease, yellow fever, chicken pox, shingles, adult T-cell leukemia, Burkitt's lymphoma,
15 nasopharyngeal carcinoma, or hepatitis. In one particular embodiment, the oligonucleotide is complementary to an HIV gene and includes about 15 to 26 nucleotides linked by phosphorothioate internucleotide linkages, at least one of the
20 nucleotides at the 3' terminus being a 2'-substituted ribonucleotide, and at least four contiguous deoxyribonucleotides.

In another embodiment, the oligonucleotide is
25 complementary to a gene encoding a protein in associated with Alzheimer's disease.

In yet other embodiments, the oligonucleotide is complementary to a gene encoding a protein
30 expressed in a parasite that causes a parasitic disease such as amebiasis, Chagas' disease, toxoplasmosis, pneumocytosis, giardiasis, cryptosporidiosis, trichomoniasis, malaria,

WO 96/12497

PCT/US95/13069

-11-

ascariasis, filariasis, trichinosis, or
schistosomiasis infections.

-12-

BRIEF DESCRIPTION OF THE DRAWINGS

5 The foregoing and other objects of the present invention, the various features thereof, as well as the invention itself may be more fully understood from the following description, when read together with the accompanying drawings in which:

10 FIG. 1 is a graphic representation showing the time course of radiolabelled, modified PS-oligonucleotide 1 in liver, kidney and plasma following the oral administration of radiolabelled, oligonucleotide.

15 FIG. 2A is an HPLC chromatograph of radiolabelled oligonucleotide standard;

20 FIG. 2B is an HPLC chromatogram of oligonucleotides extracted from plasma samples taken 12 hours after the administration of radiolabelled oligonucleotide;

25 FIG. 3A is an HPLC chromatogram of radiolabelled oligonucleotide standard;

30 FIG. 3B is an HPLC chromatogram of oligonucleotides extracted from rat liver 6 hours after the administration of radiolabelled oligonucleotide;

-13-

FIG. 3C is an HPLC chromatogram of oligonucleotides extracted from rat liver 24 hours after the administration of radiolabelled oligonucleotide;

5

FIG. 4 is a graphic representation demonstrating the course of urinary excretion of radioactivity in rats following the oral administration of radiolabelled oligonucleotide;

10

FIG. 5A is an HPLC chromatogram of radiolabelled oligonucleotide standard;

15

FIG. 5B is an HPLC chromatogram of oligonucleotides extracted from rat urine 6 hours after the administration of radiolabelled oligonucleotide;

20

FIG. 5C is an HPLC chromatogram of oligonucleotides extracted from rat urine 12 hours after the administration of radiolabelled oligonucleotide;

25

FIG. 6 is a graphic representation showing the course of radioactivity in the gastrointestinal tract and feces in rats following the oral administration of radiolabelled oligonucleotide;

30

FIG. 7 is an HPLC chromatogram of oligonucleotides extracted from rat stomach 1 hour, 3 hours, and 6 hours after the administration of radiolabelled oligonucleotide; and

WO 96/12497

PCT/US95/13069

-14-

FIG. 8 is an HPLC chromatogram of
oligonucleotides extracted from rat large
intestine
3 hours, 6 hours, and 12 hours after the
5 administration of radiolabelled oligonucleotide.

-15-

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 The patent and scientific literature referred to herein establishes the knowledge that is available to those with skill in the art.

10 This invention provides a method of down-regulating the expression of a gene in an animal by the oral administration of an oligonucleotide whose nucleotide sequence is complementary to the targeted gene.

15 It is known that an oligonucleotide, called an "antisense oligonucleotide," can bind to a target single-stranded nucleic acid molecule according to the Watson-Crick or the Hoogsteen
20 rule of base pairing, and in doing so, disrupt the function of the target by one of several mechanisms: by preventing the binding of factors required for normal transcription, splicing, or translation; by triggering the enzymatic
25 destruction of mRNA by RNase H if a contiguous region of deoxyribonucleotides exists in the oligonucleotide, and/or by destroying the target via reactive groups attached directly to the antisense oligonucleotide.

30 Thus, because of the properties described above, such oligonucleotides are useful therapeutically by their ability to control or down-regulate the expression of a particular gene

-16-

in an animal, according to the method of the present invention.

5 The oligonucleotides useful in the method of the invention are at least 6 nucleotides in length, but are preferably 6 to 50 nucleotides long, with 15 to 30mers being the most common. They are composed of deoxyribonucleotides, ribonucleotides, or a combination of both, with
10 the 5' end of one nucleotide and the 3' end of another nucleotide being covalently linked by non-phosphodiester internucleotide linkages. Such linkages include alkylphosphonates, phosphorothioates, phosphorodithioates,
15 alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamdate, carboxymethyl esters, carbonates, and phosphate triesters. Oligonucleotides with these linkages can be prepared according to known
20 methods such as phosphoramidate or H-phosphonate chemistry which can be carried out manually or by an automated synthesizer as described by Brown (*A Brief History of Oligonucleotide Synthesis. Protocols for Oligonucleotides and Analogs, Methods in Molecular Biology* (1994)
25 20:1-8). (See also, e.g., Sonveaux "Protecting Groups in Oligonucleotides Synthesis" in Agrawal (1994) *Methods in Molecular Biology* 26:1-72; Uhlmann et al. (1990) *Chem. Rev.* 90:543-583).

30 The oligonucleotides of the composition may also be modified in a number of ways without compromising their ability to hybridize to the target nucleic acid. Such modifications include,

-17-

for example, those which are internal or at the end(s) of the oligonucleotide molecule and include additions to the molecule of the internucleoside phosphate linkages, such as cholesteryl or diamine compounds with varying numbers of carbon residues between the amino groups and terminal ribose, deoxyribose and phosphate modifications which cleave, or crosslink to the opposite chains or to associated enzymes or other proteins which bind to the viral genome. Examples of such modified oligonucleotides include oligonucleotides with a modified base and/or sugar such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide having a sugar which, at both its 3' and 5' positions is attached to a chemical group other than a hydroxyl group (at its 3' position) and other than a phosphate group (at its 5' position). Other modified oligonucleotides are capped with a nuclease resistance-conferring bulky substituent at their 3' and/or 5' end(s), or have a substitution in one nonbridging oxygen per nucleotide. Such modifications can be at some or all of the internucleoside linkages, as well as at either or both ends of the oligonucleotide and/or in the interior of the molecule. For the preparation of such modified oligonucleotides, see, e.g., Agrawal (1994) *Methods in Molecular Biology* 26; Uhlmann et al. (1990) *Chem. Rev.* 90:543-583).

Oligonucleotides which are self-stabilized are also considered to be modified oligonucleotides useful in the methods of the invention (Tang et al. (1993) *Nucleic Acids Res.*

-18-

20:2729-2735). These oligonucleotides comprise two regions: a target hybridizing region; and a self-complementary region having an oligonucleotide sequence complementary to a nucleic acid sequence that is within the self-stabilized oligonucleotide.

The preparation of these unmodified and modified oligonucleotides is well known in the art (reviewed in Agrawal et al. (1992) *Trends Biotechnol.* 10:152-158) (see, e.g., Uhlmann et al. (1990) *Chem. Rev.* 90:543-584; and (1987) *Tetrahedron. Lett.* 28:(31):3539-3542); Agrawal (1994) *Methods in Molecular Biology* 20:63-80).

These oligonucleotides are provided with additional stability by having non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, phosphoramidates, phosphoramidites, phosphate esters, carbamates, carbonates, phosphate triesters, acetamidate, and carboxymethyl esters. Particularly useful oligonucleotides are linked with phosphorothioate and/or phosphorodithioate internucleoside linkages.

The oligonucleotides administered to the animal may be hybrid oligonucleotides in that they contain both deoxyribonucleotides and at least one 2' substituted ribonucleotide. For purposes of the invention, the term "2'-substituted" means substitution of the 2'-OH of the ribose molecule with, e.g., 2'-O-allyl, 2'-O-alkyl, 2'-halo, or

-19-

2'-amino, but not with 2'-H, wherein allyl, aryl, or alkyl groups may be unsubstituted or substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carbalkoxyl or amino groups.

The hybrid DNA/RNA oligonucleotides useful in the method of the invention resist nucleolytic degradation, form stable duplexes with RNA or DNA, and preferably activate RNase H when hybridized with RNA. They may additionally include at least one unsubstituted ribonucleotide. For example, an oligonucleotide useful in the method of the invention may contain all deoxyribonucleotides with the exception of one 2' substituted ribonucleotide at the 3' terminus of the oligonucleotide.

Alternatively, the oligonucleotide may have at least one substituted ribonucleotide at both its 3' and 5' termini.

One preferred class of oligonucleotides useful in the method of the invention contains four or more deoxyribonucleotides in a contiguous block, so as to provide an activating segment for RNase H. In certain cases, more than one such activating segment will be present at any location within the oligonucleotide. There may be a majority of deoxyribonucleotides in oligonucleotides according to the invention. In fact, such oligonucleotides may have as many as all but one nucleotide being deoxyribonucleotides. Thus, a preferred oligonucleotide having from

-20-

about 2 to about 50 nucleotides or most preferably
from about 12 to about 25 nucleotides, the number
of deoxyribonucleotides present ranges from 1 to
about 24. Other useful oligonucleotides may
5 consist only of 2'-substituted ribonucleotides.

TABLE 1 lists some representative species of
oligonucleotides which are useful in the method of
the invention. 2'-substituted nucleotides are
10 underscored.

TABLE 1

	<u>NO.</u>	<u>OLIGONUCLEOTIDE</u>
15	1	CTCTCGCACCCATCTCTCTCCTTCU
	2	CTCTCGCACCCATCTCTCTCCTUCU
	3	CTCTCGCACCCATCTCTCTCCUUCU
	4	CTCTCGCACCCATCTCUCUCCUUCU
	5	CTCTCGCACCCAUCUCUCUCCUUCU
20	6	CTCTCGCACCCAUCUCUCUCCUUCU
	7	CTCTCGCACCCAUCUCUCUCCUUCU
	8	CUCUCGCACCCAUCUCUCUCCUUCU
	9	CTCTCGCACCCATCTCTCTCCTTCU
	10	CUCTCGCACCCATCTCTCTCCTTCU
25	11	CUCUCGCACCCATCTCTCTCCUUCU
	12	CUCUCGCACCCATCTCUCUCCUUCU
	13	CUCUCGCACCCAUCUCUCUCCUUCU
	14	CUCUCGCACCCATCTCTCUCUCCUUCU
	15	CTCTCGCACCCAUCUCUCUCCUUCU
30	16	CUCUCGCACCCAUCTCTCTCCUUCU
	17	CUCUCGCACCCATCTCTCTCCUUCU
	18	CUCTCGCACCCAUCUCUCUCCUUCU
	19	CUCTCGCACCCATCTCTCUCUCCUUCU

-21-

The 2' substituted ribonucleotide(s) in the oligonucleotide may contain at the 2' position of the ribose, a -O-lower alkyl containing 1-6 carbon atoms, aryl or substituted aryl or allyl having 2-
5 6 carbon atoms e.g., 2'-O-allyl, 2'-O-aryl, 2'-O-alkyl, 2'-halo, or 2'-amino, but not with 2'-H, wherein allyl, aryl, or alkyl groups may be unsubstituted or substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl,
10 acyloxy, alkoxy, carboxyl, carbalkoxyl or amino groups. Useful substituted ribonucleotides are 2'-O-alkyls such as 2'-O-methyl.

Preferably, oligonucleotides according to the
15 invention will range from about 2 to about 50 nucleotides in length, and most preferably from about 15 to about 25 nucleotides in length. Thus, in this preferred case, oligonucleotides according to the invention will have from 14 to 24
20 non-phosphodiester internucleotide linkages.

The oligonucleotides according to the invention are effective in inhibiting the expression of various genes in viruses, pathogenic
25 organisms, or in inhibiting the expression of cellular genes. The ability to inhibit such agents is clearly important to the treatment of a variety of disease states. Thus, oligonucleotides according to the method of the invention have a
30 nucleotide sequence which is complementary to a nucleic acid sequence that is from a virus, a pathogenic organism or a cellular gene. Preferably such oligonucleotides are from about 6 to about 50 nucleotides in length.

-22-

For purposes of the invention, the term "oligonucleotide sequence that is complementary to a nucleic acid sequence" is intended to mean an oligonucleotide sequence that binds to the target nucleic acid sequence under physiological conditions, e.g., by Watson-Crick base pairing (interaction between oligonucleotide and single-stranded nucleic acid) or by Hoogsteen base pairing (interaction between oligonucleotide and double-stranded nucleic acid) or by any other means including in the case of a oligonucleotide binding to RNA, pseudoknot formation. Such binding (by Watson Crick base pairing) under physiological conditions is measured as a practical matter by observing interference with the function of the nucleic acid sequence.

The nucleic acid sequence to which an oligonucleotide according to the invention is complementary will vary, depending upon the gene to be down-regulated. In some cases, the target gene or nucleic acid sequence will be a virus nucleic acid sequence. The use of antisense oligonucleotides to inhibit various viruses is well known (reviewed in Agrawal (1992) *Trends in Biotech.* 10:152-158). Viral nucleic acid sequences that are complementary to effective antisense oligonucleotides have been described for many viruses, including human immunodeficiency virus type 1 (HIV-1) (U.S. Patent No. 4,806,463), herpes simplex virus (U.S. Patent No. 4,689,320), influenza virus (U.S. Patent No. 5,194,428), and human papilloma virus (Storey et al. (1991) *Nucleic Acids Res.* 19:4109-4114). Sequences complementary

-23-

to any of these nucleic acid sequences can be used for oligonucleotides according to the invention, as can be oligonucleotide sequences complementary to nucleic acid sequences from any other virus.

5 Additional viruses that have known nucleic acid sequences against which antisense oligonucleotides can be prepared include, but are not limited to, foot and mouth disease virus (see, Robertson et al. (1985) *J. Virol.* 54:651; Harris et al. (1980) *Virol.* 10 36:659), yellow fever virus (see Rice et al. (1985) *Science* 229:726), varicella-zoster virus (see, Davison and Scott (1986) *J. Gen. Virol.* 67:2279), Epstein-Barr virus, cytomegalovirus, respiratory syncytial virus (RSV), and cucumber 15 mosaic virus (see Richards et al. (1978) *Virol.* 89:395).

For example, an oligonucleotide has been designed which is complementary to a portion of 20 the HIV-1 gene, and as such, has significant anti-HIV effects (Agrawal (1992) *Antisense Res. Development* 2:261-266). The target of this oligonucleotide has been found to be conserved among various HIV-1 isolates. It is 56% G + C rich, water soluble, 25 and relatively stable under physiological conditions. This oligonucleotide binds to a complementary RNA target under physiological conditions, with the T of the duplex approximately being 56°C. The antiviral activity of this 30 oligonucleotide has been tested in several models, including acutely and chronically infected CEM cells, long-term cultures mimicking *in vivo* conditions, human peripheral blood lymphocytes and

-24-

macrophages, and isolates from HIV-1 infected patients (Lisiewicz et al. (*Proc. Natl. Acad. Sci. (USA)* (1992) 89:11209-11213); Lisiewicz et al. (*Proc. Natl. Acad. Sci. (USA)* (1993) 90:3860-3864); Lisiewicz et al. (*Proc. Natl. Acad. Sci. (USA)* (1994) 91:7942-7946); Agrawal et al. (*Trends Biotechnol.* (1992) 10(5):152-158).

The oligonucleotides according to the invention alternatively can have an oligonucleotide sequence complementary to a nucleic acid sequence of a pathogenic organism. The nucleic acid sequences of many pathogenic organisms have been described, including the malaria organism, *Plasmodium falciparum*, and many pathogenic bacteria. Oligonucleotide sequences complementary to nucleic acid sequences from any such pathogenic organism can be used in oligonucleotides according to the invention. Examples of pathogenic eucaryotes having known nucleic acid sequences against which antisense oligonucleotides can be prepared include *Trypanosoma brucei gambiense* and *Leishmania* (See Campbell et al., *Nature* 311:350 (1984)), *Fasciola hepatica* (See Zurita et al., *Proc. Natl. Acad. Sci. USA* 84:2340 (1987)).

Antifungal oligonucleotides can be prepared using a target hybridizing region having an oligonucleotide sequence that is complementary to a nucleic acid sequence from, e.g., the chitin synthetase gene, and antibacterial oligonucleotides can be prepared using, e.g., the alanine racemase gene. Among fungal diseases that may be treatable by the method of treatment

-25-

according to the invention are candidiasis, histoplasmosis, cryptococcocis, blastomycosis, aspergillosis, sporotrichosis, chromomycosis, dermatophytosis, and coccidioidomycosis. The method might also be used to treat rickettsial diseases (e.g., typhus, Rocky Mountain spotted fever), as well as sexually transmitted diseases caused by Chlamydia trachomatis or Lymphogranuloma venereum. A variety of parasitic diseases may be treated by the method according to the invention, including amebiasis, Chagas' disease, toxoplasmosis, pneumocystosis, giardiasis, cryptosporidiosis, trichomoniasis, and Pneumocystis carini pneumonia; also worm (helminthic) diseases such as ascariasis, filariasis, trichinosis, schistosomiasis and nematode or cestode infections. Malaria may be treated by the method of treatment of the invention regardless of whether it is caused by P. falciparum, P. vivax, P. ovale, or P. malariae.

The infectious diseases identified above may all be treated by the method of treatment according to the invention because the infectious agents for these diseases are known and thus oligonucleotides according to the invention can be prepared, having oligonucleotide sequence that is complementary to a nucleic acid sequence that is an essential nucleic acid sequence for the propagation of the infectious agent, such as an essential gene.

Other disease states or conditions that may be treatable by the method according to the

-26-

invention are those which result from an abnormal expression or product of a cellular gene. These conditions may be treated by administration of oligonucleotides according to the invention, and have been discussed earlier in this disclosure.

Other oligonucleotides according to the invention can have a nucleotide sequence complementary to a cellular gene or gene transcript, the abnormal expression or product of which results in a disease state. The nucleic acid sequences of several such cellular genes have been described, including prion protein (Stahl et al. (1991) *FASEB J.* 5:2799-2807), the amyloid-like protein associated with Alzheimer's disease (U.S. Patent No. 5,015,570), and various well-known oncogenes and proto-oncogenes, such as *c-myb*, *c-myc*, *c-abl*, and *n-ras*. In addition, oligonucleotides that inhibit the synthesis of structural proteins or enzymes involved largely or exclusively in spermatogenesis, sperm motility, the binding of the sperm to the egg or any other step affecting sperm viability may be used as contraceptives. Similarly, contraceptives for women may be oligonucleotides that inhibit proteins or enzymes involved in ovulation, fertilization, implantation or in the biosynthesis of hormones involved in those processes.

Hypertension may be controlled by oligonucleotides that down-regulate the synthesis of angiotensin converting enzyme or related enzymes in the renin/angiotensin system. Platelet aggregation may be controlled by suppression of

-27-

the synthesis of enzymes necessary for the synthesis of thromboxane A₂ for use in myocardial and cerebral circulatory disorders, infarcts, arteriosclerosis, embolism and thrombosis.

5 Deposition of cholesterol in arterial wall may be inhibited by suppression of the synthesis of fatty acid co-enzyme A: cholesterol acyl transferase in arteriosclerosis. Inhibition of the synthesis of cholinephosphotransferase may be useful in
10 hypolipidemia.

There are numerous neural disorders in which hybridization arrest may be used to reduce or eliminate adverse effects of the disorder. For
15 example, suppression of the synthesis of monoamine oxidase may be used in Parkinson's disease. Suppression of catechol o-methyl transferase may be used to treat depression; and suppression of indole N-methyl transferase may be used in
20 treating schizophrenia.

Suppression of selected enzymes in the arachidonic acid cascade which leads to prostaglandins and leukotrienes may be useful in
25 the control of platelet aggregation, allergy, inflammation, pain and asthma.

Suppression of the protein expressed by the multidrug resistance (*mdr-1*) gene, which can be
30 responsible for development of resistance of tumors to a variety of anti-cancer drugs and is a major impediment in chemotherapy may prove to be beneficial in the treatment of cancer. Oligonucleotide sequences complementary to nucleic

-28-

acid sequences from any of these genes can be used for oligonucleotides according to the invention, as can be oligonucleotide sequences complementary to any other cellular gene transcript, the
5 abnormal expression or product of which results in a disease state.

The oligonucleotides described herein are administered orally or enterally to the animal
10 subject in the form of therapeutic pharmaceutical formulations that are effective for treating virus infection, infections by pathogenic organisms, or disease resulting from abnormal gene expression or from the expression of an abnormal gene product.
15 In some aspects of the method according to the invention, the oligonucleotides are administered in conjunction with other therapeutic agents, e.g., AZT in the case of AIDS.

20 The therapeutic pharmaceutical formulation containing the oligonucleotide includes a physiologically acceptable carrier, such as an inert diluent or an assimilable edible carrier with which the peptide is administered. Suitable
25 formulations that include pharmaceutically acceptable excipients for introducing compounds to the bloodstream by other than injection routes can be found in *Remington's Pharmaceutical Sciences* (18th ed.) (Genarro, ed. (1990) Mack Publishing Co., Easton,
30 PA). The oligonucleotide and other ingredients may be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the individual's diet. The oligonucleotide may be incorporated with

-29-

excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. When the oligonucleotide is administered orally, it may be mixed with other food forms and pharmaceutically acceptable flavor enhancers. When the oligonucleotide is administered enterally, they may be introduced in a solid, semi-solid, suspension, or emulsion form and may be compounded with any number of well-known, pharmaceutically acceptable additives. Sustained release oral delivery systems and/or enteric coatings for orally administered dosage forms are also contemplated such as those described in U.S. Patent Nos. 4,704,295, 4,556,552, 4,309,404, and 4,309,406.

The amount of oligonucleotide in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit contains between from about 50 micrograms to about 200 mg per kg body weight of the animal, with 10 mg to 100 mg per kg being most preferable.

It will be appreciated that the unit content of active ingredient or ingredients contained in an individual dose of each dosage form need not in itself constitute an effective amount since the necessary effective amount can be reached by administration of a plurality of dosage units

-30-

(such as capsules or tablets or combinations thereof).

5 In order to determine if the oligonucleotide administered according to the method of the invention is absorbed into body tissues, and if so, in which tissues absorption occurs, the following study was performed. Samples of various body tissues were analyzed for radioactivity at increasing hours after oral administration of a radioactively labelled oligonucleotide. FIG. 1 illustrates the plasma, liver, and kidney concentration-time course of an oligonucleotide equivalents after oral administration of the radiolabelled oligonucleotide. These results demonstrate that the drug is absorbed through gastrointestinal tract and accumulated in the kidney and the liver.

20 The chemical form of radioactivity in plasma was further evaluated by HPLC, demonstrating the presence of both intact oligonucleotide (A) as well as metabolites (B) 12 hours after oral administration (FIG. 2B). Intact oligonucleotide was also detected in liver 6 hour (FIG. 3B) and 12 hours (FIG. 3C) after administration. Radioactivity in brain, thymus, heart, lung, liver, kidney, adrenals, stomach, small intestine, large intestine, skeletal muscle, testes, thyroid, epidermis, whole eye, and bone marrow was detectable 48 hours after oral administration of the radiolabelled oligonucleotide.

-31-

Further evidence to support the absorption of the oligonucleotide comes from urine sample analysis after radioactively labelled oligonucleotide was orally administered. FIG. 4 shows the cumulative excretion of the radioactively labelled oligonucleotide into the urine over 48 hr following the administration of radiolabelled oligonucleotide. That the oligonucleotide continues to be excreted in the urine over time implies that other tissues had absorbed it, and that the body was capable of absorption for an extended period of time. FIGS. 5B and 5C demonstrate that although the majority of radioactivity in urine was present as degradative products, intact oligonucleotide was also detected, demonstrating that this oligonucleotide is absorbed intact.

To determine the level of bioavailability of oligonucleotides following oral administration the level of the oligonucleotide in the gastrointestinal tract (stomach and intestine) and feces was measured. FIG. 6 shows that approximately 80% of administered oligonucleotide remained or was excreted in feces, indicating that 20% of administered oligonucleotide was absorbed. This oligonucleotide was stable in stomach; no significant degradative products in stomach contents were detected six hours after oral administration (FIG. 7). The majority of administered oligonucleotide in the contents of the large intestine were also present as the intact compound (FIG. 8).

-32-

Thus, using the method of the invention, successful absorption of oligonucleotides was accomplished through the gastrointestinal tract and distributed throughout the body. Intact
5 oligonucleotides were detected in plasma and various tissues and excreted into the urine. These results demonstrate that oral administration is a potential means for delivery of oligonucleotides as therapeutic agents.

10 The following examples illustrate the preferred modes of making and practicing the present invention, but are not meant to limit the scope of the invention since alternative methods
15 may be utilized to obtain similar results.

EXAMPLES

1. Synthesis and Analysis of Oligonucleotide

20 A hybrid 25-mer phosphorothioate-linked oligonucleotide having SEQ ID NO:1 and containing 2'-O-methyl ribonucleotide 3' and 5' sequences and a deoxyribonucleotide interior was synthesized,
25 purified, and analyzed as follows.

30 Unmodified phosphorothioate deoxynucleosides were synthesized on CPG on a 5-6 μ mole scale on an automated synthesizer (model 8700, Millipore, Bedford, MA) using the H-phosphonate approach described in U.S. Patent No. 5,149,798. Deoxynucleoside H-phosphonates were obtained from Millipore (Bedford, MA).

-33-

2'-O-methyl ribonucleotide H-phosphonates or phosphorothioates were synthesized by standard procedures (see, e.g., "Protocols for Oligonucleotides and Analogs" in *Meth. Mol. Biol.*

5 (1993) volume 20) or commercially obtained (e.g., from Glenn Research, Sterling, VA and Clontech, Palo Alto, CA). Segments of oligonucleotides containing 2'-O-methyl nucleoside(s) were

10 assembled by using 2'-O-methyl ribonucleoside H-phosphonates or phosphorothioates for the desired cycles. Similarly, segments of oligonucleotides containing deoxyribonucleosides were assembled by using deoxynucleoside H-phosphonates for the
15 desired cycles. After assembly, CPG bound oligonucleotide H-phosphonate was oxidized with sulfur to generate the phosphorothioate linkage. Oligonucleotides were then deprotected in concentrated NH_4OH at 40°C for 48 hours.

20 Crude oligonucleotide (about 500 A_{260} units) was analyzed on reverse low pressure chromatography on a C_{18} reversed phase medium. The DMT group was removed by treatment with 80% aqueous acetic acid, then the oligonucleotides
25 were dialyzed against distilled water and lyophilized.

2. Radioactive Labelling of Oligonucleotide

30 To obtain ^{35}S -labelled oligonucleotide, synthesis was carried out in two steps. The first 19 nucleotides of the sequence SEQ ID NO:1) from its 3'-end were assembled using the β -cyanoethyl-phosphoramidite approach (see, Beaucage in *Protocols*

-34-

for Oligonucleotides and Analogs (Agrawal, ed.), Humana Press, (1993), pp. 33-61). The last six nucleotides were assembled using the H-phosphonate approach (see, Froehler in *Protocols for Oligonucleotides and Analogs* (Agrawal, ed.) Humana Press, 1993, pp. 5 63-80). Controlled pore glass (CPG) support-bound oligonucleotide (30 mg of CPG; approximately 1 μ M) containing five H-phosphonate linkage was oxidized with $^{35}\text{S}_8$ (4 mCi, 1 Ci/mg, Amersham; 1 Ci = 37 GBq) 10 in 60 ml carbon disulfide/pyridine/triethylamine (10:10:1). The oxidation reaction was performed at room temperature for 1 hr with occasional shaking. Then 2 μ l, 5 μ l, and 200 μ l of 5% cold sulfur ($^{32}\text{S}_8$) in same solvent mixture was added 15 every 30 min to complete the oxidation. The solution was removed and the CPG support was washed with carbon disulfide/pyridine/triethylamine (10:10:1) (3 x 500 μ l) and with acetonitrile (3 x 700 μ l). The product was 20 deprotected in concentrated ammonium hydroxide (55°C, 14 hr) and evaporated. The resultant product was purified by polyacrylamide gel electrophoresis (20% polyacrylamide containing 7 M urea). The desired band was excised under UV 25 shadowing and the PS-oligonucleotide was extracted from the gel and desalted with a Sep-Pak C18 cartridge (Waters) and Sephadex G-15 column. The yield was 20 A_{260} units (600 μ g; specific activity, 1 μ Ci/ μ g).

30

3. Animals and Treatment

Male Sprague-Dawley rats (100-120 g, Harlan Laboratories, Indianapolis, IN) were used in the

-35-

study. The animals were fed with commercial diet and water *ad libitum* for 1 week prior to the study.

Animals were dosed via gavage at a dose of 50 mg/kg. Unlabelled and ³⁵S-labelled oligonucleotides were dissolved in physiological saline (0.9% NaCl) in a concentration of 25 mg/ml. Doses were based on the pretreatment body weight and rounded to the nearest 0.01 ml. After dosing, each animal was placed in a metabolism cage and fed with commercial diet and water *ad libitum*. Total voided urine was collected and each metabolism cage was then washed following the collection intervals. Total excreted feces was collected from each animal at various timepoints and feces samples were homogenized prior to quantitation of radioactivity. Blood samples were collected in heparinized tubes from animals at the various timepoints. Plasma was separated by centrifugation. Animals were euthanized by exsanguination under sodium pentobarbital anesthesia. Following euthanasia, the tissues were collected from each animal. All tissues/organs were trimmed of extraneous fat or connective tissue, emptied and cleaned of all contents, individually weighed, and the weights recorded.

4. Total Radioactivity Measurements

The total radioactivities in tissues and body fluids were determined by liquid scintillation spectrometry (LS 6000TA, Beckman, Irvine, CA). In brief, biological fluids (plasma, 50-100 µl;

-36-

urine, 50-100 μ l) were mixed with 6 ml
scintillation solvent (Budget-Solve, RPI, Mt.
Prospect, IL) to determine total radioactivity.
Feces were ground and weighed prior to being
5 homogenized in a 9-fold volume of 0.9% NaCl
saline. An aliquot of the homogenate (100 μ l) was
mixed with solubilizer (TS-2, RPI, Mt. Prospect,
IL) and then with scintillation solvent (6 ml) to
permit quantitation of total radioactivity.

10

Following their removal, tissues were
immediately blotted on Whatman No. 1 filter paper
and weighed prior to being homogenized in 0.9%
NaCl saline (3-5 ml per gram of wet weight). The
15 resulting homogenate (100 μ l) was mixed with
solubilizer (TS-2, RPI, Mt. Prospect, IL) and then
with scintillation solvent (6 ml) to determine
total radioactivity. The volume of 0.9% NaCl
saline added to each tissue sample was recorded.
20 The homogenized tissues/organs were kept frozen at
 $\leq -70^{\circ}\text{C}$ until the use for further analysis.

5. HPLC Analysis

25 The radioactivity in urine was analyzed by
paired-ion HPLC using a modification of the method
described essentially by Sands et al. (*Mol. Pharm.*
(1994) 45:932-943). Urine samples were
centrifuged and passed through a 0.2- μ m Acro
30 filter (Gelman, Ann Arbor, MI) prior to analysis.
Hybrid oligonucleotide and metabolites in plasma
samples were extracted using the above methods in
sample preparation for PAGE. A Microsorb MV-C4
column (Rainin Instruments, Woburn, MA) was

WO 96/12497

PCT/US95/13069

-37-

employed in HPLC using a Hewlett Packard 1050 HPLC with a quaternary pump for gradient making.

Mobile phase included two buffers; Buffer A was 5 mM-A reagent (Waters Co., Bedford, MA) in water and Buffer B was 4:1 (v/v) Acetonitrile

(Fisher)/water. The column was eluted at a flow rate of 1.5 ml/min, using the following gradient:

(1) 0-4 min, 0% buffer B; (2) 4-15 min 0-35%

Buffer B; and (3) 15-70 min 35%-80% Buffer B. The

column was equilibrated with Buffer A for at least 30 min prior to the next run. By using a RediFrac

fraction collector (Pharmacia LKB Biotechnology,

Piscataway, NJ), 1-min fractions (1.5 ml) were

collected into 7-ml scintillation vials and mixed

with 5 ml scintillation solvent to determine

radioactivity in each fraction.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are considered to be within the scope of this invention, and are covered by the following claims.

WO 96/12497

PCT/US95/13069

-38-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5 (i) APPLICANT: Hybridon, Inc.

(ii) TITLE OF INVENTION: A Method Of Down-Regulating Gene
Expression

10 (iii) NUMBER OF SEQUENCES: 1

(iv) CORRESPONDENCE ADDRESS:

15 (A) ADDRESSEE: Lappin & Kusmer
(B) STREET: 200 State Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: USA
(F) ZIP: 02109

20 (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

30

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Kerner, Ann-Louise
(B) REGISTRATION NUMBER: 33,523
(C) REFERENCE/DOCKET NUMBER: HYZ-030PCT

35

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 617-330-1300
(B) TELEFAX: 617-330-1311

40

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA/RNA

50

(iii) HYPOTHETICAL: NO

WO 96/12497

PCT/US95/13069

-39-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

5
CUCUCGCACC CATCTCTCTC CUUCU

25

THE EMBODIMENTS OF THE INVENTION FOR WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A pharmaceutical formulation for oral administration comprising an oligonucleotide that inhibits or down-regulates expression of a gene in a mammal,

wherein said oligonucleotide is complementary to a partial sequence of the gene or of the gene transcript,

wherein said oligonucleotide has non-phosphodiester internucleotide linkages and comprises at least one 2'-substituted ribonucleotide

and wherein the oligonucleotide has a 2'-substituted ribonucleotide at the 3' terminus.

2. The pharmaceutical formulation of Claim 1 wherein the oligonucleotide further comprises at least one 2'-substituted ribonucleotide at the 5' terminus.

3. The pharmaceutical formulation of Claim 1 wherein all the ribonucleotides in the oligonucleotide are 2'-substituted ribonucleotides.

4. The pharmaceutical formulation of any one of Claims 1 to 3 wherein the 2'-substituted ribonucleotide is a 2'-O-alkyl-ribonucleotide.

5. The pharmaceutical formulation of Claim 1, 2 or 4 wherein the oligonucleotide comprises at least one deoxyribonucleotide.

6. The pharmaceutical formulation of Claim 5 wherein the oligonucleotide comprises a region of at least four contiguous deoxyribonucleotides capable of activating RNase H activity.

7. The pharmaceutical formulation of any one of Claims 1 to 6 wherein the oligonucleotide comprises one or more internucleotide linkages selected from alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, phosphoramidates, phosphoramidites, phosphate esters, carbamates, carbonates, phosphate triesters, acetamides, or carboxymethyl esters.

8. The pharmaceutical formulation of Claim 7 wherein essentially all of the oligonucleotides are linked via phosphorothioate or phosphorodithioate internucleotide linkages.

9. The pharmaceutical formulation of any one of Claims 1 to 8 wherein the oligonucleotide is modified.

10. The pharmaceutical formulation of Claim 8 wherein all of the oligonucleotides are linked via phosphorothioate linkages.

11. The pharmaceutical formulation of Claim 10 further comprising at least two 2'-O-methylribonucleotides.

12. The pharmaceutical formulation of Claim 11 comprising at least one 2'-O-methyl-ribonucleotide at each terminus of the oligonucleotide.

13. The pharmaceutical formulation of any one of Claims 1 to 12 wherein the oligonucleotide is complementary to a partial sequence of a gene of a virus, a pathogenic organism, or a cellular gene, or a gene transcript.

14. The pharmaceutical formulation of Claim 13 wherein the oligonucleotide is complementary to a partial sequence of a gene or a gene transcript of a virus involved in AIDS, oral herpes, genital herpes, papilloma warts, influenza, foot and mouth disease, yellow fever, chicken pox, shingles, adult T-cell leukemia, Burkitt's lymphoma, nasopharyngeal carcinoma, or hepatitis.

15. The pharmaceutical formulation of any one of Claims 1 to 12 wherein the oligonucleotide is complementary to a partial sequence of a gene or a gene transcript encoding a protein associated with Alzheimer's disease.



16. The pharmaceutical formulation of any one of Claims 1 to 12 wherein the oligonucleotide is complementary to a gene encoding a protein in a parasite causing a parasitic disease selected from amebiasis, Chagas' disease, toxoplasmosis, pneumocytosis, giardiasis, cryptosporidiosis, trichomoniasis, malaria,

ascariasis, filariasis, trichinosis, or schistosomiasis infections.



17. The pharmaceutical formulation of Claim 1, 2 or 9 wherein the oligonucleotide is complementary to a partial sequence of an HIV gene or gene transcript and comprises about 15 to 26 nucleotides linked by phosphorothioate internucleotide linkages, at least one of the nucleotides at the 3' terminus being a 2'-substituted ribonucleotide.

18. The use of the pharmaceutical formulation according to any one of Claims 1 to 17 for the manufacture of a medicament, for oral administration, for inhibiting or down regulating the expression of the gene in the mammal.




19. The use of the pharmaceutical formulation according to any one of Claims 1 to 17 for inhibition or down regulation of the expression of the gene in a mammal in need thereof.




**FIG. 1****FIG. 4**

2/7


**FIG. 2A****FIG. 2B**

3/7


**FIG. 3A****FIG. 3B****FIG. 3C**

**FIG. 5A****FIG. 5B****FIG. 5C**


5/7

**FIG. 6**

6/7

**FIG. 7**

7/7

**FIG. 8**