

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2010277350 B2**

(54) Title
Herbicidally active heteroaryl-substituted cyclic diones or derivatives thereof

(51) International Patent Classification(s)
C07D 401/08 (2006.01) **C07D 409/08** (2006.01)
A01N 43/78 (2006.01) **C07D 417/08** (2006.01)
C07D 405/08 (2006.01) **C07D 417/14** (2006.01)

(21) Application No: **2010277350** (22) Date of Filing: **2010.07.29**

(87) WIPO No: **WO11/012862**

(30) Priority Data

(31) Number	(32) Date	(33) Country
0913436.2	2009.07.31	GB
1012582.1	2010.07.27	GB

(43) Publication Date: **2011.02.03**

(44) Accepted Journal Date: **2015.09.24**

(71) Applicant(s)
Syngenta Limited

(72) Inventor(s)
Jeanmart, Stephane Andre Marie; Longstaff, Adrian; Mathews, Christopher John; Viner, Russell Colin; Wood, Frances Kathryn

(74) Agent / Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000

(56) Related Art
US 2007/0135630 (& WO 2002/088098)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 February 2011 (03.02.2011)

(10) International Publication Number
WO 2011/012862 A1

(51) International Patent Classification:

C07D 401/08 (2006.01) C07D 417/08 (2006.01)
C07D 405/08 (2006.01) C07D 417/14 (2006.01)
C07D 409/08 (2006.01) A01N 43/78 (2006.01)

(21) International Application Number:

PCT/GB2010/001449

(22) International Filing Date:

29 July 2010 (29.07.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0913436.2 31 July 2009 (31.07.2009) GB
1012582.1 27 July 2010 (27.07.2010) GB

(71) Applicant (for all designated States except US): SYNGENTA LIMITED [GB/GB]; European Regional Centre, Priestley Road, Surrey Research Park, Guildford, Surrey GU2 7YH (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JEANMART, Stephane André Marie [BE/CH]; c/o Syngenta Crop Protection Muenchwil AG, Schaffhauserstrasse, CH-4332 Stein (CH). LONGSTAFF, Adrian [GB/GB]; c/o Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY (GB). MATH- EWS, Christopher John [GB/GB]; c/o Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY (GB). RUSSELL, Claire Janet [GB/GB]; c/o Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY (GB). VINER, Russell Colin [GB/GB]; c/o Syngenta

Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY (GB). WOOD, Frances Kathryn [GB/GB]; c/o Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY (GB).

(74) Agents: SYNGENTA LIMITED et al.; P O Box 3538, Jealott's Hill International Research Centre, Bracknell, Berks RG42 6YA (GB).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: HERBICIDALLY ACTIVE HETEROARYL-SYBSTITYTED CYCLIC DIONES OR DERIVATIVES THEREOF



(I)

(57) Abstract: The invention relates to a compound of formula (I), which is suitable for use as a herbicide wherein G is hydrogen or an agriculturally acceptable metal, sulfonium, ammonium or latentating group; Q is a unsubstituted or substituted C₃-C₈ saturated or mono-unsaturated heterocyclyl containing at least one heteroatom selected from O, N and S, or Q is heteroaryl or substituted heteroaryl; m is 1, 2 or 3; and Het is an optionally substituted monocyclic or bicyclic heteroaromatic ring; and wherein the compound is optionally an agronomically acceptable salt thereof.

Herbicidally active heteroaryl-substituted cyclic diones or derivatives thereof

The present invention relates to novel, herbicidally active heteroaryl diones or derivatives thereof, specifically herbicidally active heteroaryl-substituted cyclic diones or derivatives thereof, more specifically herbicidally active 2-heteroaryl-cyclopentane-1,3-diones or derivatives thereof; to processes for the preparation of these compounds or derivatives; to compositions comprising these compounds or derivatives; and to their use in controlling weeds, especially in crops of useful plants, or in inhibiting undesired plant growth.

Heteroaryl cyclic dione compounds having herbicidal action are described in US 4,678,501. WO 96/16061 A1 (Bayer AG) discloses thiophene-substituted cyclic diones, wherein the cyclic dione is selected from one of eight heterocyclic or carbocyclic classes such as cyclopentanedione or cyclohexanedione, and the use of these diones as pesticides and herbicides. WO 02/088098 A1 (Bayer AG) discloses thiazolyl-substituted carbocyclic 1,3-diones, specifically 2-(thiazolyl)-cyclopentane-1,3-diones and 2-(thiazolyl)-cyclohexane-1,3-diones, and derivatives thereof, and their uses as pesticidal agents, herbicides, and fungicides. WO 03/035643 A1 (Bayer Cropscience AG) discloses 5-membered-heterocycles substituted by oxo and also substituted by either N-linked-pyrazolyl or C-linked-pyrazolyl, and their uses as phytosanitary products, microbicides, and herbicides. WO 2009/000533 A1 (Syngenta Limited) discloses *inter alia* pyrandione, thiopyrandione and cyclohexanetrione compounds, which are substituted by an optionally substituted monocyclic or bicyclic heteroaromatic ring such as thiophene or thiazolyl, and their use as herbicides. WO 2009/015877 A1 (Syngenta Limited) discloses bicyclic (bridged carbocyclic) diones, which are substituted by an optionally substituted monocyclic or bicyclic heteroaromatic ring such as thiophene or thiazolyl, and their use as herbicides. WO 2009/086041 A1 (E.I. DuPont de Nemours & Co.) discloses herbicidal pyridazinone derivatives substituted by -G-J where G and J are each independently an optionally substituted phenyl ring or an optionally substituted 5- or 6-membered heteroaromatic ring; in WO 2009/086041, G can for example be substituted 1H-pyrazol-1-yl.

US 4,338,122 (Union Carbide Corp.) discloses 2-aryl-1,3-cyclopentanedione compounds exhibiting acaricidal and herbicidal activity. WO 96/01798 (Bayer AG) and its derived patent US 5,840,661 disclose 2-aryl-cyclopentane-1,3-dione derivatives and their use as pesticides and herbicides. WO 01/74770 (Bayer AG), its equivalent US 2003/0216260 A1, and its derived AU patent 782557 (AU 200144215C) disclose C₂-phenyl-substituted cyclic (heterocyclic or carbocyclic) ketoenols and their use as pesticides and herbicides. WO

- 2 -


2008/071405 A1 (Syngenta Limited *et al.*) discloses *inter alia* pyrandione, thiopyrandione and cyclohexanetrione compounds, which are substituted by a phenyl ring which is substituted by optionally substituted aryl or optionally substituted heteroaryl, and their use as herbicides.

WO 2008/145336 A1 (Syngenta Limited) discloses bicyclic (bridged carbocyclic) diones, which are substituted by a substituted phenyl ring, and their use as herbicides.

Copending patent application PCT/EP2009/058250, filed on 1 July 2009 and published on 7 January 2010 as WO 2010/000773 A1 (Syngenta Limited), discloses 5-(heterocyclylalkyl)-3-hydroxy-2-phenyl-cyclopent-2-enones, and their 2-phenyl-4-(heterocyclylalkyl)-cyclopentane-1,3-dione tautomers, and derivatives thereof, as herbicides. Copending patent application PCT/EP2009/066712, filed on 9 December 2009 and published on 24 June 2010 as WO 2010/069834 A1 (Syngenta Participations AG and Syngenta Limited), discloses 2-phenyl-4-(heteroaryl-methyl)-cyclopentane-1,3-diones, and derivatives thereof as herbicides.

Novel heteroaryl dione compounds and derivatives thereof, having herbicidal and/or plant-growth-inhibiting properties, have now been found.

The present invention accordingly relates to a compound of formula (I)



wherein:

G is hydrogen or an agriculturally acceptable metal, sulfonium, ammonium or latentating group; and

R¹, R², R³, R⁴ and R⁵ are independently hydrogen, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆alkenyloxy, C₃-C₆haloalkenyloxy, C₃-C₆alkynyloxy, C₃-C₆cycloalkyl, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkoxysulfonyl, C₁-C₆haloalkoxysulfonyl, cyano, nitro, phenyl, phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, or heteroaryl or

- 3 -

heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, or benzyl or benzyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, or C₃-C₆cycloalkylC₁-C₃alkyl in which a ring or chain methylene group is optionally replaced by an oxygen or sulfur atom; and/or

R² and R³ or R⁴ and R⁵ together with the carbon atoms to which they are attached form an optionally substituted 3- to 8-membered ring, optionally containing an oxygen, sulphur or nitrogen atom; and/or

R¹ and R⁴ together form a bond; and

Q is C₃-C₈ saturated or mono-unsaturated heterocyclyl containing at least one heteroatom selected from O, N and S, unsubstituted or substituted by a residue of formula =O, =N-R¹⁰, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxyC₁-C₂alkyl, C₃-C₆cycloalkyl, phenyl or phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, where R¹⁰ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₁-C₆haloalkylsulfinyl or C₁-C₆haloalkylsulfonyl; or

Q is an heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; and

m is 1, 2 or 3; and

Het is an optionally substituted monocyclic or bicyclic heteroaromatic ring;

wherein, in the optionally substituted monocyclic or bicyclic heteroaromatic ring which is Het, the optional substituents are selected, independently, from halogen, nitro, cyano, rhodano, isothiocyanato, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy(C₁-C₆)alkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₇cycloalkyl (itself optionally substituted with C₁-C₆alkyl or halogen), C₅₋₇cycloalkenyl (itself optionally substituted with C₁-C₆alkyl or halogen), hydroxy, C₁-C₁₀alkoxy, C₁-C₁₀alkoxy(C₁-C₁₀)alkoxy, tri(C₁-C₄)alkylsilyl(C₁-C₆)alkoxy, C₁-C₆alkoxycarbonyl(C₁-C₁₀)alkoxy, C₁-C₁₀haloalkoxy, aryl(C₁-C₄)alkoxy (where the aryl group is optionally substituted with halogen or C₁-C₆alkyl), C₃-C₇cycloalkyloxy (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), C₃-C₁₀alkenyloxy, C₃-

- 4 -

C₁₀alkynyloxy, mercapto, C₁-C₁₀alkylthio, C₁-C₁₀haloalkylthio, aryl(C₁-C₄)alkylthio, C₃-C₇cycloalkylthio (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), tri(C₁-C₄)-alkylsilyl(C₁-C₆)alkylthio, arylthio, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkylsulfinyl, C₁-C₆haloalkylsulfinyl, arylsulfonyl, tri(C₁-C₄)alkylsilyl, aryl-di(C₁-C₄)alkylsilyl, C₁-C₄alkyldiarylsilyl, triarylsilyl, C₁-C₁₀alkylcarbonyl, HO₂C, C₁-C₁₀alkoxycarbonyl, aminocarbonyl, C₁-C₆alkylaminocarbonyl, di(C₁-C₆alkyl)-aminocarbonyl, *N*-(C₁-C₃ alkyl)-*N*-(C₁-C₃alkoxy)aminocarbonyl, C₁-C₆alkylcarbonyloxy, arylcarbonyloxy, di(C₁-C₆)alkylaminocarbonyloxy, aryl (itself optionally substituted with C₁-C₆alkyl or halogen), heteroaryl (itself optionally substituted with C₁-C₆alkyl or halogen), heterocyclyl (itself optionally substituted with C₁-C₆alkyl or halogen), aryloxy (where the aryl group is optionally substituted with C₁-C₆alkyl or halogen), heteroaryloxy (where the heteroaryl group is optionally substituted with C₁-C₆alkyl or halogen), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C₁-C₆alkyl or halogen), amino, C₁-C₆alkylamino, di(C₁-C₆)alkylamino, C₁-C₆alkylcarbonylamino, *N*-(C₁-C₆)alkylcarbonyl-*N*-(C₁-C₆)alkylamino, and arylcarbonyl (where the aryl group is itself optionally substituted with halogen or C₁-C₆alkyl);

or two adjacent positions on the Het heteroaromatic system are optionally cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen or C₁-C₆alkyl;

or, in the optionally substituted monocyclic or bicyclic heteroaromatic ring which is Het, the optional substituents are selected from arylcarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), (C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkoxycarbonyl-*N*-(C₁-C₆)alkylamino, aryloxy carbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryloxy carbonyl-*N*-(C₁-C₆)alkylamino, (where the aryl group is substituted by C₁-C₆alkyl or halogen), arylsulphonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), arylsulphonyl-*N*-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryl-*N*-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), arylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), heteroaryl amino (where the heteroaryl group is substituted by C₁-C₆alkyl or halogen), heterocyclylamino (where the heterocyclyl group is substituted by C₁-C₆alkyl or halogen), aminocarbonylamino, C₁-C₆alkylaminocarbonylamino, di(C₁-C₆)alkylaminocarbonylamino, arylaminocarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryl-*N*-(C₁-C₆)alkylaminocarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), C₁-C₆alkylaminocarbonyl-*N*-(C₁-C₆)alkylamino, di(C₁-C₆)alkylaminocarbonyl-*N*-(C₁-C₆)alkylamino, arylaminocarbonyl-*N*-(C₁-C₆)alkylamino (where

the aryl group is substituted by C₁-C₆alkyl or halogen), and aryl-*N*-(C₁-C₆)alkylaminocarbonyl-*N*-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen);

and wherein, when R² and R³ or R⁴ and R⁵ together with the carbon atoms to which they are attached form an optionally substituted 3- to 8-membered ring, optionally containing an oxygen, sulphur or nitrogen atom, then:

when R² and R³ or R⁴ and R⁵ together with the carbon atoms to which they are attached form a carbocycle, then the optional substituents on the carbocycle are selected, independently, from halogen, nitro, cyano, rhodano, isothiocyanato, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy(C₁-C₆)alkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₇cycloalkyl (itself optionally substituted with C₁-C₆alkyl or halogen), C₅₋₇cycloalkenyl (itself optionally substituted with C₁-C₆alkyl or halogen), hydroxy, C₁-C₁₀alkoxy, C₁-C₁₀alkoxy(C₁-C₁₀)alkoxy, tri(C₁-C₄)alkylsilyl(C₁-C₆)alkoxy, C₁-C₆alkoxycarbonyl(C₁-C₁₀)alkoxy, C₁-C₁₀haloalkoxy, aryl(C₁-C₄)alkoxy (where the aryl group is optionally substituted with halogen or C₁-C₆alkyl), C₃-C₇cycloalkyloxy (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), C₃-C₁₀alkenyloxy, C₃-C₁₀alkynyloxy, mercapto, C₁-C₁₀alkylthio, C₁-C₁₀haloalkylthio, aryl(C₁-C₄)alkylthio, C₃-C₇cycloalkylthio (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), tri(C₁-C₄)-alkylsilyl(C₁-C₆)alkylthio, arylthio, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkylsulfinyl, C₁-C₆haloalkylsulfinyl, arylsulfonyl, tri(C₁-C₄)alkylsilyl, aryl-di(C₁-C₄)alkylsilyl, C₁-C₄alkyldiarylsilyl, triarylsilyl, C₁-C₁₀alkylcarbonyl, HO₂C, C₁-C₁₀alkoxycarbonyl, aminocarbonyl, C₁-C₆alkylaminocarbonyl, di(C₁-C₆alkyl)-aminocarbonyl, *N*-(C₁-C₃alkyl)-*N*-(C₁-C₃alkoxy)aminocarbonyl, C₁-C₆alkylcarbonyloxy, arylcarbonyloxy, di(C₁-C₆)alkylaminocarbonyloxy, aryl (itself optionally substituted with C₁-C₆alkyl or halogen), heteroaryl (itself optionally substituted with C₁-C₆alkyl or halogen), heterocyclyl (itself optionally substituted with C₁-C₆alkyl or halogen), aryloxy (where the aryl group is optionally substituted with C₁-C₆alkyl or halogen), heteroaryloxy (where the heteroaryl group is optionally substituted with C₁-C₆alkyl or halogen), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C₁-C₆alkyl or halogen), amino, C₁-C₆alkylamino, di(C₁-C₆)alkylamino, C₁-C₆alkylcarbonylamino, *N*-(C₁-C₆)alkylcarbonyl-*N*-(C₁-C₆)alkylamino, and arylcarbonyl (where the aryl group is itself optionally substituted with halogen or C₁-C₆alkyl);

and when R² and R³ or R⁴ and R⁵ together with the carbon atoms to which they are attached form an optionally substituted 3- to 8-membered ring which contains an oxygen, sulphur or nitrogen atom, then, in the resulting heterocyclyl group, the one or more optional substituents are independently selected from halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-

- 6 -

C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, nitro and cyano;

and wherein the latentating group G is selected from the groups C₁-C₈alkyl, C₂-C₈haloalkyl, phenylC₁-C₈alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroarylC₁-C₈alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₈alkenyl, C₃-C₈haloalkenyl, C₃-C₈alkynyl, C(X^a)-R^a, C(X^b)-X^c-R^b, C(X^d)-N(R^c)-R^d, -SO₂-R^e, -P(X^e)(R^f)-R^g and CH₂-X^f-R^h;

wherein X^a, X^b, X^c, X^d, X^e and X^f are independently of each other oxygen or sulfur;

and wherein R^a is H, C₁-C₁₈alkyl (e.g. C₁-C₆alkyl or C₁-C₄alkyl such as *tert*-butyl or isopropyl), C₂-C₁₈alkenyl, C₂-C₁₈alkynyl, C₁-C₁₀haloalkyl (e.g. C₁-C₁₀fluoroalkyl), C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro;

R^b is C₁-C₁₈alkyl (e.g. C₁-C₆alkyl or C₁-C₄alkyl such as C₁-C₂alkyl), C₃-C₁₈alkenyl, C₃-C₁₈alkynyl, C₂-C₁₀haloalkyl (e.g. C₂-C₁₀fluoroalkyl), C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₂-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-

- 7 -

C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl, (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl, or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro;

R^c and R^d are each independently of each other hydrogen, C₁-C₁₀alkyl (e.g. C₁-C₆alkyl or C₁-C₄alkyl such as C₁-C₂alkyl), C₃-C₁₀alkenyl, C₃-C₁₀alkynyl, C₂-C₁₀haloalkyl (e.g. C₂-C₁₀fluoroalkyl), C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₂-C₅)alkylaminoalkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or heteroaryl amino or heteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or diheteroaryl amino or diheteroaryl amino

- 8 -

substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro, or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro, or C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino or C₃-C₇cycloalkoxy; or R^c and R^d join together to form a 3-7 membered ring, optionally containing one heteroatom selected from O or S;

R^e is C₁-C₁₀alkyl (e.g. C₁-C₆alkyl or C₁-C₄alkyl such as C₁-C₂alkyl), C₂-C₁₀alkenyl, C₂-C₁₀alkynyl, C₁-C₁₀haloalkyl (e.g. C₁-C₁₀fluoroalkyl), C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro, or heteroaryl amino or heteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro, or diheteroaryl amino or diheteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro, or C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino or C₂-C₈dialkylamino;

- 9 -

R^f and R^g are each independently of each other C_1 - C_{10} alkyl (e.g. C_1 - C_6 alkyl or C_1 - C_4 alkyl such as C_1 - C_2 alkyl), C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkoxy, C_1 - C_{10} haloalkyl (e.g. C_1 - C_{10} fluoroalkyl), C_1 - C_{10} cyanoalkyl, C_1 - C_{10} nitroalkyl, C_1 - C_{10} aminoalkyl, C_1 - C_5 alkylamino(C_1 - C_5)alkyl, C_2 - C_8 dialkylamino(C_1 - C_5)alkyl, C_3 - C_7 cycloalkyl(C_1 - C_5)alkyl, C_1 - C_5 alkoxy(C_1 - C_5)alkyl, C_3 - C_5 alkenyloxy(C_1 - C_5)alkyl, C_3 - C_5 alkynyloxy(C_1 - C_5)alkyl, C_1 - C_5 alkylthio(C_1 - C_5)alkyl, C_1 - C_5 alkylsulfinyl(C_1 - C_5)alkyl, C_1 - C_5 alkylsulfonyl(C_1 - C_5)alkyl, C_2 - C_8 alkylideneaminoxy(C_1 - C_5)alkyl, C_1 - C_5 alkylcarbonyl(C_1 - C_5)alkyl, C_1 - C_5 alkoxycarbonyl(C_1 - C_5)alkyl, aminocarbonyl(C_1 - C_5)alkyl, C_1 - C_5 alkylaminocarbonyl(C_1 - C_5)alkyl, C_2 - C_8 dialkylaminocarbonyl(C_1 - C_5)alkyl, C_1 - C_5 alkylcarbonylamino(C_1 - C_5)alkyl, N -(C_1 - C_5)alkylcarbonyl- N -(C_2 - C_5)alkylaminoalkyl, C_3 - C_6 trialkylsilyl(C_1 - C_5)alkyl, phenyl(C_1 - C_5)alkyl (wherein the phenyl is optionally substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, C_1 - C_3 alkylthio, C_1 - C_3 alkylsulfinyl, C_1 - C_3 alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C_1 - C_5)alkyl (wherein the heteroaryl is optionally substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, C_1 - C_3 alkylthio, C_1 - C_3 alkylsulfinyl, C_1 - C_3 alkylsulfonyl, halogen, cyano, or by nitro), C_2 - C_5 haloalkenyl, C_3 - C_6 cycloalkyl, or phenyl or phenyl substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or nitro, or heteroaryl or heteroaryl substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or by nitro, or heteroaryl amino or heteroaryl amino substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or by nitro, or diheteroaryl amino or diheteroaryl amino substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or nitro, or phenyl amino or phenyl amino substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or nitro, or diphenyl amino or diphenyl amino substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or nitro, or C_3 - C_7 cycloalkyl amino, di- C_3 - C_7 cycloalkyl amino, C_3 - C_7 cycloalkoxy, C_1 - C_{10} haloalkoxy, C_1 - C_5 alkyl amino or C_2 - C_8 dialkyl amino, or benzyloxy or phenoxy, wherein the benzyl and phenyl groups are in turn optionally substituted by C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, halogen, cyano or nitro; and

R^h is C_1 - C_{10} alkyl (e.g. C_1 - C_6 alkyl or C_1 - C_4 alkyl such as C_1 - C_2 alkyl), C_3 - C_{10} alkenyl, C_3 - C_{10} alkynyl, C_1 - C_{10} haloalkyl (e.g. C_1 - C_{10} fluoroalkyl), C_1 - C_{10} cyanoalkyl, C_1 - C_{10} nitroalkyl, C_2 - C_{10} aminoalkyl, C_1 - C_5 alkylamino(C_1 - C_5)alkyl, C_2 - C_8 dialkylamino(C_1 - C_5)alkyl, C_3 - C_7 cycloalkyl(C_1 - C_5)alkyl, C_1 - C_5 alkoxy(C_1 - C_5)alkyl, C_3 - C_5 alkenyloxy(C_1 - C_5)alkyl, C_3 - C_5 alkynyloxy(C_1 - C_5)alkyl, C_1 - C_5 alkylthio(C_1 - C_5)alkyl, C_1 - C_5 alkylsulfinyl(C_1 - C_5)alkyl, C_1 - C_5 alkylsulfonyl(C_1 - C_5)alkyl, C_2 - C_8 alkylideneaminoxy(C_1 - C_5)alkyl, C_1 - C_5 alkylcarbonyl(C_1 - C_5)alkyl, C_1 - C_5 alkoxycarbonyl(C_1 - C_5)alkyl, aminocarbonyl(C_1 - C_5)alkyl, C_1 -


2010277350 04 May 2012

- 10 -

C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), phenoxy(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryloxy(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl, or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen or by nitro, or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro;

and wherein the compound is optionally an agronomically acceptable salt thereof.

In one aspect the present invention provides a compound of formula I



(I),

wherein:

G is hydrogen or an agriculturally acceptable metal, sulfonium, ammonium or latentating group; and

R¹ is hydrogen, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆alkenyloxy, C₃-C₆haloalkenyloxy, C₃-

2010277350 04 May 2012

- 10A -

C₆alkynyloxy, C₃-C₆cycloalkyl, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkoxysulfonyl, C₁-C₆haloalkoxysulfonyl, cyano, nitro, phenyl; phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or benzyl or benzyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or C₃-C₆cycloalkylC₁-C₃alkyl in which a ring or chain methylene group is optionally replaced by an oxygen or sulfur atom;

R² and R³ independently are hydrogen or methyl; and

R⁴ and R⁵ independently are hydrogen or methyl;

and/or R¹ and R⁴ together form a bond; and

Q is C₃-C₈ saturated or mono-unsaturated heterocyclyl containing at least one heteroatom selected from O, N and S, unsubstituted or substituted by a residue of formula =O, =N-R¹⁰, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxyC₁-C₂alkyl, C₃-C₆cycloalkyl, phenyl or phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, where R¹⁰ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₁-C₆haloalkylsulfinyl or C₁-C₆haloalkylsulfonyl; or












Q is an heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; and

m is 1; and

Het is a group selected from the formulae Het₁ to Het₁₂:

2010277350 04 May 2012

- 10B -

Het₁Het₂Het₃Het₄Het₅Het₆Het₇Het₈Het₉Het₁₀Het₁₁Het₁₂

wherein:

B designates the point of attachment to the ketoenol moiety;

W¹ is N or CR⁹;

W² and W³ are independently of each other N or CR⁷;

W⁴ is N or CR¹⁰;

with the proviso that at least one of W¹, W², W³ or W⁴ is N;

X is O, S, or NR¹²;

2010277350 04 May 2012

- 10C -

Z is N or CR¹³;

wherein

R⁶ is halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, nitro or cyano;

R⁷ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆ cycloalkyl, C₅-C₆cycloalkenyl, halogen, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, or C₁-C₆haloalkoxy; or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, cyano or nitro; or phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano;

R⁸ is hydrogen, C₁-C₄ alkyl, C₂-C₃alkenyl, C₂-C₃alkynyl, C₁-C₄ haloalkyl, or C₂-C₃ haloalkenyl;

R⁹ is hydrogen, methyl, halomethyl, or halogen;

R¹⁰ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄ alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl or cyano;

R¹¹ is hydrogen, methyl, ethyl, halomethyl, or haloethyl; or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, cyano or nitro; or phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano;

R¹² is hydrogen, methyl, ethyl, or halomethyl; and

R¹³ is hydrogen, methyl, ethyl, halomethyl, haloethyl, halogen, cyano or nitro;

2010277350 04 May 2012

- 10D -

and wherein the latentating group G is selected from the groups C₁-C₈alkyl, C₂-C₈haloalkyl, phenylC₁-C₈alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroarylC₁-C₈alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₈alkenyl, C₃-C₈haloalkenyl, C₃-C₈alkynyl, C(X^a)-R^a, C(X^b)-X^c-R^b, C(X^d)-N(R^c)-R^d, -SO₂-R^e, -P(X^e)(R^f)-R^g and CH₂-X^f-R^h;

wherein X^a, X^b, X^c, X^d, X^e and X^f are independently of each other oxygen or sulfur;

and wherein R^a is H, C₁-C₁₈alkyl, C₂-C₁₈alkenyl, C₂-C₁₈alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, N-(C₁-C₅)alkylcarbonyl-N-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro;

R^b is C₁-C₁₈alkyl, C₃-C₁₈alkenyl, C₃-C₁₈alkynyl, C₂-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₂-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-

2010277350 04 May 2012

- 10E -

C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; and

R^c and R^d are each independently of each other hydrogen, C₁-C₁₀alkyl, C₃-C₁₀alkenyl, C₃-C₁₀alkynyl, C₂-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₂-C₅)alkylaminoalkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl-amino or heteroaryl-amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or diheteroaryl-amino or diheteroaryl-amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or

2010277350 04 May 2012

- 10F -

phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino or C₃-C₇cycloalkoxy;

or R^c and R^d join together to form a 3-7 membered ring, optionally containing one heteroatom selected from O or S; and

R^e is C₁-C₁₀alkyl, C₂-C₁₀alkenyl, C₂-C₁₀alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or heteroaryl amino or heteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diheteroaryl amino or diheteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino or C₂-C₈dialkylamino;

2010277350 04 May 2012

- 10G -

R^f and R^g are each independently of each other C₁-C₁₀alkyl, C₂-C₁₀alkenyl, C₂-C₁₀alkynyl, C₁-C₁₀alkoxy, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₂-C₅)alkylaminoalkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or heteroarylamino or heteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diheteroarylamino or diheteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or C₃-C₇cycloalkylamino, diC₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino or C₂-C₈dialkylamino; or benzyloxy or phenoxy, wherein the benzyl and phenyl groups are in turn optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; and

R^h is C₁-C₁₀alkyl, C₃-C₁₀alkenyl, C₃-C₁₀alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₂-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-

2010277350 04 May 2012

- 10H -

C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), phenoxy(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryloxy(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen or by nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro;

and wherein the term "heteroaryl" means an aromatic ring system containing at least one heteroatom and consisting either of a single ring or of two fused rings;

and wherein the compound is optionally an agronomically acceptable salt thereof.

In the substituent definitions of the compounds of the formula I, each alkyl moiety either alone or as part of a larger group (such as alkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl or cycloalkylalkyl) is a straight or branched chain and is, for example, independently methyl, ethyl, *n*-propyl, *n*-butyl, *n*-pentyl, *n*-hexyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl or neopentyl. The alkyl groups are suitably C₁-C₆alkyl groups, but are preferably C₁-C₄alkyl or C₁-C₃alkyl groups, and, more preferably, C₁-C₂alkyl groups.

Alkenyl and alkynyl moieties can be in the form of straight or branched chains, and the alkenyl moieties, where appropriate, can be of either the (*E*)- or (*Z*)-configuration. Examples are vinyl, allyl and propargyl. Alkenyl and alkynyl moieties can contain one or

2010277350 04 May 2012

- 10I -

more double and/or triple bonds in any combination. It is understood, that allenyl is included in these terms.

Halogen is fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine.

- 11 -

Haloalkyl (e.g. fluoroalkyl) groups are alkyl groups which are substituted with one or more of the same or different halogen (e.g. fluorine) atoms and are suitably, independently, CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, (CH₃)₂CF, CF₃CH₂ or CHF₂CH₂; preferably, independently, CF₃, CF₂H, FCH₂, CH₃CHF, (CH₃)₂CF, CF₃CH₂ or CHF₂CH₂.

The terms "heteroaryl" and/or "heteroaromatic" preferably refer to an aromatic ring system containing at least one heteroatom and consisting either of a single ring or of two or more fused rings. Preferably, single rings will contain up to three heteroatoms and bicyclic systems up to four heteroatoms which will preferably be chosen from nitrogen, oxygen and sulphur. Examples of such groups include independently furyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, benzofuryl, benzisofuryl, benzothienyl, benzisothienyl, indolyl, isoindolyl, indazolyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, 2,1,3-benzoxadiazole, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, benzotriazinyl, purinyl, pteridinyl or indolizinyl. Preferred examples of heteroaromatic radicals include independently pyridyl, pyrimidinyl, triazinyl, thienyl, furyl, oxazolyl, isoxazolyl, 2,1,3-benzoxadiazolyl or thiazolyl. Another group of preferred heteroaryls comprises independently pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyridazinyl, pyrazinyl, quinolinyl, isoquinolinyl, cinnolinyl, quinazolinyl or quinoxalinyl.

The term "heterocyclyl" preferably refers to a non-aromatic preferably monocyclic or bicyclic ring systems containing up to 7 atoms including one or more (preferably one or two) heteroatoms selected from O, S and N. Examples of such rings include 1,3-dioxolane, oxetane, tetrahydrofuran, tetrahydropyran, morpholine, thiomorpholin and piperazine. When present, the optional substituents on heterocyclyl include C₁-C₆alkyl and C₁-C₆haloalkyl as well as those optional substituents given above for an alkyl moiety.

Cycloalkyl includes preferably independently cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl. Cycloalkylalkyl is preferentially cyclopropylmethyl.

Carbocyclic rings include independently aryl, cycloalkyl or carbocyclic groups, and cycloalkenyl groups. In the compounds of the formula I, each aryl group, either alone or as part of a larger group (e.g. aryloxy etc.), is independently preferably phenyl.

When present, the optional substituents on aryl (preferably phenyl), heteroaryl and/or carbocycles are preferably selected, independently, from halogen, nitro, cyano, rhodano, isothiocyanato, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy(C₁-C₆)alkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₇cycloalkyl (itself optionally substituted with C₁-C₆alkyl or halogen), C₅₋₇cycloalkenyl (itself optionally substituted with C₁-C₆alkyl or halogen), hydroxy, C₁-C₁₀alkoxy, C₁-C₁₀alkoxy(C₁-C₁₀)alkoxy, tri(C₁-C₄)alkylsilyl(C₁-C₆)alkoxy, C₁-C₆alkoxycarbonyl(C₁-C₁₀)alkoxy, C₁-C₁₀haloalkoxy, aryl(C₁-C₄)alkoxy (where the aryl group is optionally substituted with halogen or C₁-C₆alkyl), C₃-C₇cycloalkyloxy (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), C₃-C₁₀alkenyloxy, C₃-C₁₀alkynyloxy, mercapto, C₁-C₁₀alkylthio, C₁-C₁₀haloalkylthio, aryl(C₁-C₄)alkylthio, C₃-C₇cycloalkylthio (where the cycloalkyl group is optionally substituted with C₁-C₆alkyl or halogen), tri(C₁-C₄)-alkylsilyl(C₁-C₆)alkylthio, arylthio, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkylsulfinyl, C₁-C₆haloalkylsulfinyl, arylsulfonyl, tri(C₁-C₄)alkylsilyl, aryl-di(C₁-C₄)alkylsilyl, C₁-C₄alkyldiarylsilyl, triarylsilyl, C₁-C₁₀alkylcarbonyl, HO₂C, C₁-C₁₀alkoxycarbonyl, aminocarbonyl, C₁-C₆alkylaminocarbonyl, di(C₁-C₆alkyl)-aminocarbonyl, N-(C₁-C₃alkyl)-N-(C₁-C₃alkoxy)aminocarbonyl, C₁-C₆alkylcarbonyloxy, arylcarbonyloxy, di(C₁-C₆)alkylaminocarbonyloxy, aryl (itself optionally substituted with C₁-C₆alkyl or halogen), heteroaryl (itself optionally substituted with C₁-C₆alkyl or halogen), heterocyclyl (itself optionally substituted with C₁-C₆alkyl or halogen), aryloxy (where the aryl group is optionally substituted with C₁-C₆alkyl or halogen), heteroaryloxy (where the heteroaryl group is optionally substituted with C₁-C₆alkyl or halogen), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C₁-C₆alkyl or halogen), amino, C₁-C₆alkylamino, di(C₁-C₆)alkylamino, C₁-C₆alkylcarbonylamino, N-(C₁-C₆)alkylcarbonyl-N-(C₁-C₆)alkylamino, and arylcarbonyl (where the aryl group is itself optionally substituted with halogen or C₁-C₆alkyl); or two adjacent positions on an aryl or heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen or C₁-C₆alkyl. Further substituents for aryl or heteroaryl include arylcarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), (C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkoxycarbonyl-N-(C₁-C₆)alkylamino, aryloxy carbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryloxy carbonyl-N-(C₁-C₆)alkylamino, (where the aryl group is substituted by C₁-C₆alkyl or halogen), arylsulphonylamino (where the aryl group is

substituted by C₁-C₆alkyl or halogen), arylsulphonyl-N-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryl-N-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), arylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), heteroaryl amino (where the heteroaryl group is substituted by C₁-C₆alkyl or halogen), heterocyclamino (where the heterocycl group is substituted by C₁-C₆alkyl or halogen), aminocarbonylamino, C₁-C₆alkylaminocarbonylamino, di(C₁-C₆)alkylaminocarbonylamino, arylaminocarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), aryl-N- (C₁-C₆)alkylaminocarbonylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen), C₁-C₆alkylaminocarbonyl-N-(C₁-C₆)alkylamino, di(C₁-C₆)alkylaminocarbonyl-N-(C₁-C₆)alkylamino, arylaminocarbonyl-N-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen) and aryl-N-(C₁-C₆)alkylaminocarbonyl-N-(C₁-C₆)alkylamino (where the aryl group is substituted by C₁-C₆alkyl or halogen).

For substituted heterocycl groups it is preferred that one or more substituents are independently selected from halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, nitro and cyano. It is to be understood that dialkylamino substituents include those where the dialkyl groups together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which is optionally substituted by one or two independently selected C₁-C₆alkyl groups. When heterocyclic rings are formed by joining two groups on an N atom, the resulting rings are suitably pyrrolidine, piperidine, thiomorpholine and morpholine each of which may be substituted by one or two independently selected C₁-C₆alkyl groups.

The invention relates also to the agriculturally acceptable salts which the compounds of formula I are able to form with transition metal, alkali metal or alkaline earth metal bases, amines, quaternary ammonium bases or tertiary sulfonium bases.

Among the transition metal, alkali metal or alkaline earth metal salt capable of forming transition metal, alkali metal or alkaline earth metal salts (i.e. where G is a metal), special mention should be made of the hydroxides of copper, iron, lithium, sodium, potassium, magnesium or calcium; and preferably the hydroxides, bicarbonates or carbonates of sodium or potassium.

- 14 -

Examples of amines suitable for ammonium salt formation (i.e. where G is an ammonium) include ammonia, or primary, secondary or tertiary C₁-C₁₈alkylamines, C₁-C₄hydroxyalkylamines or C₂-C₄alkoxyalkyl-amines, for example methylamine, ethylamine, *n*-propylamine, isopropylamine, the four butylamine isomers, *n*-amylamine, isoamylamine, hexylamine, heptylamine, octylamine, nonylamine, decylamine, pentadecylamine, hexadecylamine, heptadecylamine, octadecylamine, methylethylamine, methylisopropylamine, methylhexylamine, methylnonylamine, methylpentadecylamine, methyloctadecylamine, ethylbutylamine, ethylheptylamine, ethyloctylamine, hexylheptylamine, hexyloctylamine, dimethylamine, diethylamine, di-*n*-propylamine, di-isopropylamine, di-*n*-butylamine, di-*n*-amylamine, di-isoamylamine, dihexylamine, diheptylamine, dioctylamine, ethanolamine, *n*-propanolamine, isopropanolamine, *N,N*-diethanolamine, *N*-ethylpropanolamine, *N*-butylethanolamine, allylamine, *n*-but-2-enylamine, *n*-pent-2-enylamine, 2,3-dimethylbut-2-enylamine, dibut-2-enylamine, *n*-hex-2-enylamine, propylenediamine, trimethylamine, triethylamine, tri-*n*-propylamine, tri-isopropylamine, tri-*n*-butylamine, tri-isobutylamine, tri-sec-butylamine, tri-*n*-amylamine, methoxyethylamine or ethoxyethylamine; or heterocyclic amines, for example pyridine, quinoline, isoquinoline, morpholine, piperidine, pyrrolidine, indoline, quinuclidine or azepine; or primary arylamines, for example anilines, methoxyanilines, ethoxyanilines, *o*-, *m*- or *p*-toluidines, phenylenediamines, benzidines, naphthylamines or *o*-, *m*- or *p*-chloroanilines; but especially triethylamine, isopropylamine or di-isopropylamine.

Preferred quaternary ammonium bases suitable for salt formation (i.e. where G is an ammonium) correspond, for example, to the formula [N(R_a R_b R_c R_d)]OH, wherein R_a, R_b, R_c and R_d are each independently of the others hydrogen or C₁-C₄alkyl. Further suitable tetraalkylammonium bases with other anions can be obtained, for example, by anion exchange reactions.

Preferred tertiary sulfonium bases suitable for salt formation (i.e. where G is a sulfonium) correspond, for example, to the formula [SR_eR_fR_g]OH, wherein R_e, R_f and R_g are each independently of the others C₁-C₄ alkyl. Trimethylsulfonium hydroxide is especially preferred. Suitable sulfonium bases may be obtained from the reaction of thioethers, in particular dialkylsulfides, with alkylhalides, followed by conversion to a suitable base, for example a hydroxide, by anion exchange reactions.

- 15 -

It should be understood that in those compounds of formula I, where G is a metal, ammonium or sulfonium as mentioned above and as such represents a cation, the corresponding negative charge is largely delocalised across the O-C=C-C=O unit.

The compounds of formula I according to the invention also include hydrates, for example hydrates which may be formed during salt formation.

The latentating group G is selected to allow its removal by one or a combination of biochemical, chemical or physical processes to afford compounds of formula I where G is H before, during or following (preferably during or following) application to the treated area or plants. Examples of these processes include enzymatic cleavage (e.g. enzymatic cleavage of esters), chemical hydrolysis and photolysis. Compounds bearing such groups G may, in some cases, offer certain advantage(s), such as: improved penetration of the cuticula of the plants treated; increased tolerance of crops; improved compatibility or stability in formulated mixtures containing other herbicides, herbicide safeners, plant growth regulators, fungicides and/or insecticides; and/or reduced leaching in soils; in particular improved penetration of the cuticula of the plants treated.


In the latentating group G, preferably, X^a , X^b , X^c , X^d , X^e and/or X^f are oxygen. More preferably, all of X^a , X^b , X^c , X^d , X^e and X^f are oxygen.

Preferably, the latentating group G is a group $-C(X^a)-R^a$ or $-C(X^b)-X^c-R^b$.

More preferably, the latentating group G is a group $-C(X^a)-R^a$ or $-C(X^b)-X^c-R^b$, wherein R^a is hydrogen or C_1 - C_{18} alkyl (more preferably, hydrogen or C_1 - C_6 alkyl, still more preferably C_1 - C_6 alkyl, most preferably C_1 - C_4 alkyl such as *tert*-butyl or isopropyl), R^b is C_1 - C_{18} alkyl (more preferably, C_1 - C_6 alkyl, still more preferably C_1 - C_4 alkyl such as C_1 - C_2 alkyl), and the meanings of X^a , X^b and X^c are as defined above (more preferably, X^a , X^b and X^c are oxygen).

It is preferred that G is hydrogen, an alkali metal or alkaline earth metal, where hydrogen is especially preferred.

Depending on the nature of the substituents, compounds of formula I may exist in different isomeric forms. When G is hydrogen, for example, compounds of formula I may exist in different tautomeric forms (one dione tautomer and two different keto-enol tautomers), as shown in the following scheme:




This invention covers all such isomers and tautomers and mixtures thereof in all proportions. Also, when substituents contain double bonds, *cis*- and *trans*-isomers can exist. These isomers, too, are within the scope of the claimed compounds of the formula I.

In a preferred group of compounds of the formula (I), R^1 is hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy, and, more preferably, R^1 is hydrogen or methyl.


In another preferred group of the compounds of the formula (I), R^2 and R^3 independently are hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy, and, more preferably, R^2 and R^3 independently are hydrogen or methyl.


Preferably, in the compounds of the formula (I), R^4 and R^5 independently are hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy, and, more preferably, R^4 and R^5 independently are hydrogen or methyl.

Preferred saturated or mono-unsaturated rings Q are those of formulae Q_1 to Q_{107} :




- 17 -





- 19 -



wherein:

R is hydrogen, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆alkenyloxy, C₃-C₆haloalkenyloxy, C₃-C₆alkynyloxy, C₃-C₆cycloalkyl, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkoxysulfonyl, C₁-C₆haloalkoxysulfonyl, cyano, nitro, phenyl, phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, or heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl;

R' is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₆-C₁₀arylsulfonyl, C₆-C₁₀arylcarbonyl, C₆-C₁₀arylaminocarbonyl, C₇-C₁₆arylalkylaminocarbonyl, C₁-C₉heteroarylsulfonyl, C₁-C₉heteroarylcarbonyl, C₁-C₉heteroarylaminocarbonyl, C₂-C₁₅heteroarylalkylaminocarbonyl;

R'' is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₁-C₆haloalkylsulfinyl or C₁-C₆haloalkylsulfonyl;

n is 0, 1, 2, 3 or 4; and

A denotes the position of attachment to the $-(CR^4R^5)_m-$ moiety.

- 20 -

Groups Q₁, Q₂, Q₃, Q₄, Q₅, Q₆, Q₇, Q₂₅, Q₂₆, Q₂₇, Q₂₈, Q₂₉, Q₈₆, Q₈₇, Q₈₈, Q₈₉, Q₉₀ are more preferred, and groups Q₁ to Q₇ are particularly preferred.

Preferably, R and R' are independently hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy, and R'' is hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy or C₁-C₆haloalkylcarbonyl.

Preferably, n is 0, 1 or 2. More preferably, n is 0.

In another preferred group of the compounds of the formula (I), Q is a 5- or 6-membered heteroaryl or is a 5- or 6-membered heteroaryl which is substituted one to three times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.

Preferably, the heteroatoms in these Q moieties are selected from 1 or 2 nitrogen, oxygen or sulphur atoms.

In particular, Q is thienyl, furyl, oxazolyl, isoxazolyl, benzofuryl, thiazolyl, oxazolyl, isothiazolyl, benzothienyl, benzoisothienyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl or benzisoxazolyl, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.













In particular, Q is pyridyl, pyrimidinyl, pyridazinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl or quinoxalinyl, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.

Preferably, in the compounds of the formula (I), m is 1 or 2, and most preferably m is 1.

Preferably, Het is an optionally substituted monocyclic 6-membered nitrogen containing heteroaromatic ring, or, preferably, is an optionally substituted monocyclic 5-membered sulfur or (preferably) nitrogen containing heteroaromatic ring. More preferably, Het is a monocyclic 5-membered sulfur and nitrogen containing heteroaromatic ring.

Even more preferably, Het is a group selected from the formulae Het₁ to Het₁₂ :

- 21 -

Het₁Het₂Het₃Het₄Het₅Het₆Het₇Het₈Het₉Het₁₀Het₁₁Het₁₂

wherein:

B designates the point of attachment to the ketoenol moiety;

W¹ is N or CR⁹;

W² and W³ are independently of each other N or CR⁷;

W⁴ is N or CR¹⁰;

with the proviso that at least one of W¹, W², W³ or W⁴ is N;

X is O, S, or NR¹²;

Z is N or CR¹³;

wherein

R⁶ is hydrogen, halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, nitro or cyano; preferably halogen, C₁-C₂alkyl, C₁-C₂haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H,

- 22 -

FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), vinyl, ethynyl, or methoxy; and even more preferably methyl or ethyl;

R⁷ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆cycloalkyl, C₅-C₆cycloalkenyl, halogen, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆haloalkoxy, optionally substituted aryl (e.g. optionally substituted phenyl), optionally substituted aryloxy (e.g. optionally substituted phenoxy), optionally substituted heteroaryl or optionally substituted heteroaryloxy; preferably optionally substituted aryl (e.g. optionally substituted phenyl) or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₁-C₂ haloalkoxy, cyano or nitro; and even more preferably phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₁-C₂ haloalkoxy or cyano;

R⁸ is hydrogen, C₁-C₄ alkyl, C₂-C₃alkenyl, C₂-C₃alkynyl, C₁-C₄ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), or C₂-C₃ haloalkenyl; preferably methyl or ethyl;

R⁹ is hydrogen, methyl, halomethyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, or BrCH₂), or halogen; preferably hydrogen;

R¹⁰ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₂-C₄ alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl or cyano; preferably hydrogen, halogen, methyl or ethyl;

R¹¹ is hydrogen, methyl, ethyl, halomethyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, or BrCH₂), haloethyl (e.g. CH₃CHF, CF₃CH₂ or CHF₂CH₂), optionally substituted aryl (e.g. optionally substituted phenyl) or optionally substituted heteroaryl; preferably optionally substituted aryl (e.g. optionally substituted phenyl) or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₁-C₂ haloalkoxy, cyano or nitro; even more preferably phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, BrCH₂, CH₃CHF, CF₃CH₂ or CHF₂CH₂), C₁-C₂ haloalkoxy or cyano;

R¹² is hydrogen, methyl, ethyl, or halomethyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, or BrCH₂); and

- 23 -

R¹³ is hydrogen, methyl, ethyl, halomethyl (e.g. CF₃, CF₂Cl, CF₂H, CCl₂H, FCH₂, ClCH₂, or BrCH₂), haloethyl (e.g. CH₃CHF, CF₃CH₂ or CHF₂CH₂), halogen, cyano or nitro; preferably hydrogen.

More preferably, Het is a group of the formula Het₂, wherein X is S and Z is N and R⁶ and R⁷ are as defined above. Preferably, in this embodiment, R⁶ is methyl or ethyl. Preferably, in this embodiment, R⁷ is phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano; more preferably, R⁷ is phenyl substituted once, twice or three times (e.g. once) by halogen (e.g. chlorine); even more preferably, R⁷ is 4-chlorophenyl.

It is also more preferred that Het is a group of the formula Het₁₀, wherein Z is CR¹³ and R⁶, R⁷ and R¹³ are as defined above. Preferably, in this embodiment, R¹³ is hydrogen or methyl; more preferably hydrogen. Preferably, in this embodiment, R⁶ is methyl or ethyl. Preferably, in this embodiment, R⁷ is phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano; more preferably, R⁷ is phenyl substituted once, twice or three times (e.g. once) by halogen (e.g. chlorine); even more preferably, R⁷ is 4-chlorophenyl.

It is also preferred that Het is a group of the formula Het₂, wherein X is S and Z is CR¹³ and R⁶, R⁷ and R¹³ are as defined above. Preferably, in this embodiment, R¹³ is hydrogen. Preferably, in this embodiment, R⁶ is methyl or ethyl. Preferably, in this embodiment, R⁷ is phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano.

It is also preferred that Het is a group of the formula Het₁, wherein W¹ is CR⁹, W² is N, W³ is CR⁷, W⁴ is N, and R⁶, R⁷ and R⁹ are as defined above. Preferably, in this embodiment, R⁹ is hydrogen. Preferably, in this embodiment, R⁶ is methyl or ethyl. Preferably, in this embodiment, R⁷ is phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano.

In a group of preferred compounds of formula I,

R¹ is hydrogen or methyl;

R² and R³ independently are hydrogen or methyl;


R⁴ and R⁵ independently are hydrogen or methyl; and

- 24 -

Q is pyridyl, pyrimidinyl, pyridazinyl, quinoliny, isoquinoliny, cinnoliny, phthalazinyl, quinazolinyl or quinoxaliny, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl; or Q is thienyl, furyl, oxazolyl, isoxazolyl, benzofuryl, thiazolyl, oxazolyl, isothiazolyl, benzothienyl, benzoisothienyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl or benzisoxazolyl, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl; and

m is 1;

Het is a group Het₂




Het₂

wherein X is S, Z is N, R⁶ is methyl or ethyl, R⁷ is 4-chlorophenyl or 4-bromophenyl, and B designates the point of attachment to the ketoenol moiety; and


G is hydrogen, an alkali metal or alkaline earth metal. Within this group of preferred compounds of formula I, a particularly preferred group of compounds of formula I are those wherein R¹ to R⁵ and G are hydrogen and Q, Het and m are as defined.

In a particularly preferred group of the compounds of formula I, R¹ to R⁵ are hydrogen and G is hydrogen, an alkali or alkaline earth metal, m is 1, Q is group selected from




- 25 -

wherein n is 0 and A designates the point of attachment to the $-(CR^4R^5)_m$ moiety, or Q is pyridyl, and Het is a group of the formula

Het₂


wherein X is S, Z is N, R⁶ is methyl or ethyl, R⁷ is phenyl substituted by halogen, and B designates the point of attachment to the ketoenol moiety, or Het is a group of the formula

Het₁₀

wherein Z is CH or C-CH₃, R⁶ is methyl or ethyl, R⁷ is phenyl substituted by halogen, and B designates the point of attachment to the ketoenol moiety.

Even more preferably, Q is a group of the formula Q⁷ or pyridine-2-yl, and Het is a group of the formula Het₂ or Het₁₀, wherein R⁷ is 4-chlorophenyl.

In an alternative suitable embodiment, Het is a group of formula Het₁₃:

Het₁₃


wherein:

B designates the point of attachment to the ketoenol moiety;


R^{6a} is halogen, C₁-C₂alkyl, C₁-C₂fluoroalkyl, vinyl, ethynyl, or methoxy; preferably methyl or ethyl; and

R¹⁴ is hydrogen or C₁-C₄alkyl.

Preferably, the compound of formula I is one of compounds A1 to A25 as shown below, or an agriculturally or agronomically acceptable salt thereof (e.g. agriculturally acceptable metal or ammonium salt, e.g. an alkali metal or alkaline earth metal salt thereof):



- 27 -




More preferably, the compound of formula I is one of compounds A7, A12, A16, A17, A18, A20, A23, A24 or A25 as shown above, or an agriculturally or agronomically acceptable salt

- 28 -

thereof (e.g. agriculturally acceptable metal or ammonium salt, e.g. an alkali metal or alkaline earth metal salt thereof).

Certain compounds of formula (I) are alkenes, and as such undergo further reactions typical of alkenes to give additional compounds of formula (I) according to known procedures. Example of such reaction include, but are not restricted to, halogenation or hydrogenation




wherein R^1 and R^4 form a bond

Compounds of formula (I) wherein R^1 and R^4 form a bond and R^5 is halogen (preferably chloride or bromide) or R^5 is C_1 - C_6 alkylsulfonate (preferably mesylate) or C_1 - C_6 haloalkylsulfonate (preferably triflate) or an arylsulfonate (preferable tosylate) may undergo a cross-coupling reaction with a suitable coupling partner under conditions described in the literature for Suzuki-Miyaura, Sonogashira and related cross-coupling reactions to give additional compounds of formula (I) (see, for example, O'Brien, C. J. and Organ, M. G. *Angew. Chem. Int. Ed.* (2007), 46, 2768-2813; Suzuki, A. *Journal of Organometallic Chemistry* (2002), 653, 83; Miyaura N. and Suzuki, A. *Chem. Rev.* (1995), 95, 2457-2483).

Those skilled in the art will appreciate that compounds of formula (I) may contain an heteroaromatic moiety bearing one or more substituents capable of being transformed into alternative substituents under known conditions, and that these compounds may themselves serve as intermediates in the preparation of additional compounds of formula (I).

Additional compounds of formula (I) may be prepared by selectively reducing compounds of formula (I) where Q is a Heteroaromatic ring preferably furan or pyrrole in the presence of a suitable catalyst, and in a suitable solvent.




For example, compounds of formula (I) wherein R⁶, R⁷, R⁸ or R¹⁰ is alkenyl or alkynyl, may be reduced to compounds of formula (I) wherein R⁶, R⁷, R⁸ or R¹⁰ is alkyl under known conditions and compounds of formula (I) wherein R⁷ is halogen, preferably bromide or iodine, may undergo a cross-coupling reaction with a suitable coupling partner under conditions described in the literature for Suzuki-Miyaura, Sonogashira and related cross-coupling reactions to give additional compounds of formula (I) (see, for example, O'Brien, C. J. and Organ, M. G. *Angew. Chem. Int. Ed.* (2007), 46, 2768-2813; Suzuki, A. *Journal of Organometallic Chemistry* (2002), 653, 83; Miyaura N. and Suzuki, A. *Chem. Rev.* (1995), 95, 2457-2483).

Compounds of formula (I) wherein G is C₁-C₈alkyl, C₂-C₈haloalkyl, phenylC₁-C₈alkyl (wherein the phenyl may optionally be substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsufinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroarylC₁-C₈alkyl (wherein the heteroaryl may optionally be substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsufinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₈ alkenyl, C₃-C₈haloalkenyl, C₃-C₈alkynyl, C(X^a)-R^a, C(X^b)-X^c-R^b, C(X^d)-N(R^c)-R^d, -SO₂-R^e, -P(X^e)(R^f)-R^g or CH₂-X^f-R^h where X^a, X^b, X^c, X^d, X^e, X^f, R^a, R^b, R^c, R^d, R^e, R^f, R^g and R^h are as defined above may be prepared by treating compounds of formula (A), which are compounds of formula (I) wherein G is H, with a reagent G-Z, wherein G-Z is alkylating agent such as an alkyl halide (the definition of alkyl halides includes simple C₁-C₈ alkyl halides such as methyl iodide and ethyl iodide, substituted alkyl halides such as chloromethyl alkyl ethers, Cl-CH₂-X^f-R^h, wherein X^f is oxygen, and chloromethyl alkyl sulfides Cl-CH₂-X^f-R^h, wherein X^f is sulfur), a C₁-C₈alkyl

- 30 -

sulfonate, or a di-C₁-C₈alkyl sulfate, or with a C₃-C₈alkenyl halide, or with a C₃-C₈alkynyl halide, or with an acylating agent such as a carboxylic acid, HO-C(X^a)R^a, wherein X^a is oxygen, an acid chloride, Cl-C(X^a)R^a, wherein X^a is oxygen, or acid anhydride, [R^aC(X^a)]₂O, wherein X^a is oxygen, or an isocyanate, R^cN=C=O, or a carbamoyl chloride, Cl-C(X^d)-N(R^c)-R^d (wherein X^d is oxygen and with the proviso that neither R^c or R^d is hydrogen), or a thiocarbamoyl chloride Cl-C(X^d)-N(R^c)-R^d (wherein X^d is sulfur and with the proviso that neither R^c nor R^d is hydrogen) or a chloroformate, Cl-C(X^b)-X^c-R^b, (wherein X^b and X^c are oxygen), or a chlorothioformate Cl-C(X^b)-X^c-R^b (wherein X^b is oxygen and X^c is sulfur), or a chlorodithioformate Cl-C(X^b)-X^c-R^b, (wherein X^b and X^c are sulfur), or an isothiocyanate, R^cN=C=S, or by sequential treatment with carbon disulfide and an alkylating agent, or with a phosphorylating agent such as a phosphoryl chloride, Cl-P(X^e)(R^f)-R^g or with a sulfonylating agent such as a sulfonyl chloride Cl-SO₂-R^e, preferably in the presence of at least one equivalent of base.

Isomeric compounds of formula (I) may be formed. For example, compounds of formula (A) may give rise to two isomeric compounds of formula (I), or to isomeric mixtures of compounds of formula (I). This invention covers both isomeric compounds of formula (I), together with mixtures of these compounds in any ratio.



The O-alkylation of cyclic 1,3-diones is known; suitable methods are described, for example, in US4436666. Alternative procedures have been reported by Pizzorno, M. T. and Albonico, S. M. *Chem. Ind. (London)* (1972), 425; Born, H. *et al.* *J. Chem. Soc.* (1953), 1779; Constantino, M. G. *et al.* *Synth. Commun.* (1992), 22 (19), 2859; Tian, Y. *et al.* *Synth. Commun.* (1997), 27 (9), 1577; Chandra Roy, S. *et al.*, *Chem. Lett.* (2006), 35 (1), 16; Zubaidha, P. K. *et al.* *Tetrahedron Lett.* (2004), 45, 7187 and by Zwanenburg, B. *et al.* *Tetrahedron* (2005), 45 (22), 7109.


The acylation of cyclic 1,3-diones may be effected by procedures similar to those described, for example, in US4551547, US4175135, US4422870, US4659372 and US4436666. Typically diones of formula (A) may be treated with the acylating agent in the presence of at least one equivalent of a suitable base, optionally in the presence of a suitable solvent. The base may be inorganic, such as an alkali metal carbonate or hydroxide, or a metal hydride, or an organic base such as a tertiary amine or metal alkoxide. Examples of suitable inorganic bases include sodium carbonate, sodium or potassium hydroxide, sodium hydride, and suitable organic bases include trialkylamines, such as trimethylamine and triethylamine, pyridines or other amine bases such as 1,4-diazobicyclo[2.2.2]octane and 1,8-diazabicyclo[5.4.0]undec-7-ene. Preferred bases include triethylamine and pyridine. Suitable solvents for this reaction are selected to be compatible with the reagents and include ethers such as tetrahydrofuran and 1,2-dimethoxyethane and halogenated solvents such as dichloromethane and chloroform. Certain bases, such as pyridine and triethylamine, may be employed successfully as both base and solvent. For cases where the acylating agent is a carboxylic acid, acylation is preferably effected in the presence of a coupling agent such as 2-chloro-1-methylpyridinium iodide, *N,N'*-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and *N,N'*-carbodiimidazole, and optionally a base such as triethylamine or pyridine in a suitable solvent such as tetrahydrofuran, dichloromethane or acetonitrile. Suitable procedures are described, for example, by Zhang, W. and Pugh, G. *Tetrahedron Lett.* (1999), 40 (43), 7595 and Isobe, T. and Ishikawa, T. *J. Org. Chem.* (1999), 64 (19) 6984.

Phosphorylation of cyclic-1,3-diones may be effected using a phosphoryl halide or thiophosphoryl halide and a base by procedures analogous to those described in US4409153.


Sulfonylation of compounds of formula (A) may be achieved using an alkyl or aryl sulfonyl halide, preferably in the presence of at least one equivalent of base, for example by the procedure of Kowalski, C. J. and Fields, K. W. *J. Org. Chem.* (1981), 46, 197.

Compounds of formula (A) may be prepared from a compounds of formula (I) by hydrolysis, preferably in the presence of an acid catalyst such as hydrochloric acid and optionally in the presence of a suitable solvent such as tetrahydrofuran or acetone preferably between 25 °C and 150 °C under conventional heating or under microwave irradiation.

- 32 -



In a further approach, compounds of formula (A) may be prepared by the cyclisation of a compound of formula (B) or a compound of formula (C), wherein R''' is hydrogen or an alkyl group, preferably in the presence of an acid or base, and optionally in the presence of a suitable solvent, by analogous methods to those described by T. N. Wheeler, US4209532. Compounds of formula (B) or compounds of formula (C) wherein R''' is hydrogen may be cyclised under acidic conditions, preferably in the presence of a strong acid such as sulfuric acid, polyphosphoric acid or Eaton's reagent, optionally in the presence of a suitable solvent such as acetic acid, toluene or dichloromethane.




Compounds of formula (B) or compounds of formula (C) wherein R''' is alkyl (preferably methyl or ethyl), may be cyclised under acidic or basic conditions, preferably in the presence of at least one equivalent of a strong base such as potassium *tert*-butoxide, lithium

diisopropylamide or sodium hydride and in a solvent such as tetrahydrofuran, toluene, dimethylsulfoxide or *N,N*-dimethylformamide.


Compounds of formula (B) and compounds of formula (C), wherein R''' is H, may be esterified to, respectively, compounds of formula (B) and compounds of formula (C), wherein R''' is alkyl, under standard conditions, for example by heating with an alkyl alcohol, ROH, in the presence of an acid catalyst.

Compounds of formula (B) and compounds of formula (C), wherein R''' is H, may be prepared, respectively, by saponification of a compounds of formula (D) and compounds of formula (E) wherein R''' is alkyl (preferably methyl or ethyl), under standard conditions, followed by acidification of the reaction mixture to effect decarboxylation, by similar processes to those described, for example, by T. N. Wheeler, US4209532.



Compounds of formula (D) and compounds of formula (E), wherein R''' is alkyl, may be prepared by treating, respectively, compounds of formula (F) with suitable carboxylic acid chlorides of formula (G) or suitable carboxylic acid chlorides of formula (H) under basic conditions. Suitable bases include potassium *tert*-butoxide, sodium bis(trimethylsilyl)amide and lithium diisopropylamide and the reaction is preferably conducted in a suitable solvent (such as tetrahydrofuran or toluene) at a temperature of between -80 °C and 30 °C. Alternatively, compounds of formula (D) and compounds of formula (E), wherein R''' is H, may be prepared by treating a compound of formula (F) with a suitable base (such as


potassium *tert*-butoxide, sodium bis(trimethylsilyl)amide and lithium diisopropylamide) in a suitable solvent (such as tetrahydrofuran or toluene) at a suitable temperature (between -80 °C and 30 °C) and reacting the resulting anion with a suitable anhydride of formula (J):



Compounds of formula (F) are known compounds, or may be prepared from known compounds by known methods.


Compounds of formula (J) may be prepared, for example, by analogous methods to those described by Ballini, R. *et al.* *Synthesis* (2002), (5), 681-685; Bergmeier, S. C. and Ismail, K. A. *Synthesis* (2000), (10), 1369-1371; Groutas, W. C. *et al.* *J. Med. Chem.* (1989), 32 (7), 1607-11 and Bernhard, K. and Lincke, H. *Helv. Chim. Acta* (1946), 29, 1457-1466.

Compounds of formula (G) or compounds of formula (H) may be prepared from a compound of formula (J) by treatment with an alkyl alcohol, $R'''-OH$, in the presence of a base, such as dimethylaminopyridine or an alkaline metal alkoxide (see, for example, Buser, S. and Vasella, A. *Helv. Chim. Acta*, (2005), 88, 3151 and M. Hart *et al.* *Bioorg. Med. Chem. Letters*, (2004), 14, 1969), followed by treatment of the resulting acid with a chlorinating reagent such as oxalyl chloride or thionyl chloride under known conditions (see, for example, Santelli-Rouvier, C. *Tetrahedron Lett.* (1984), 25 (39), 4371; Walba D. and Wand, M. *Tetrahedron Lett.* (1982), 23 (48), 4995; Cason, J. *Org. Synth. Coll. Vol. III*, (169), 1955).



Compounds of formula (G) and compounds of formula (H) may be made from known compounds by known methods. For example, analogous methods to obtain compounds of formula (G) and compounds of formula (H) are described by Bergmeier, S. C. and Ismail, K. A. *Synthesis* (2000), (10), 1369-1371.

In a further approach to compounds of formula (I) may be prepared by treating compounds of formula (K) with compounds of formula (L) wherein LG is a leaving group such as halogen (preferably iodide, bromide or chloride) or an activated alcohol (preferably mesylate or tosylate) under basic conditions. Suitable bases include lithium diisopropylamide, sodium hexamethyldisilazide, potassium *tert*-butoxide and the reaction is preferably conducted in a suitable solvent (such as tetrahydrofuran) at a temperature between -80 °C and 30 °C




- 36 -

Compounds of formula (L) are known, or may be made known compounds by known methods (see for example: WO2006016178; Ueno, H. *et al.* J. Med. Chem. (2005), 48(10), 3586-3604; Kanoh, S. *et al.* Tetrahedron (2002), 58(35), 7049-7064; Strachan, J.-P. *et al.* J. Org. Chem. (2006), 71(26), 9909-9911).

Compounds of formula (K) are known compounds or may be made from known compounds by known methods (see, for example, Song, Y. S. S. *et al.* Tetrahedron Lett. (2005), 46 (46), 5987-5990 and WO09030450)

Alternatively, compounds of formula (K) wherein G is C₁-C₆alkyl may be prepared by alkylation of compounds of formula (K), wherein G is hydrogen under known conditions. Compounds of formula (K), wherein G is hydrogen, are known, or may be prepared from known compounds by known methods (see, for example, DE10118310).


Alternatively, in a further approach to compounds of formula (K), compounds of formula (M), which are compounds of formula (K) wherein G is hydrogen and Het is (Het₂) when R⁶ is CH₂R^{''''} and R^{''''} is hydrogen or methyl, may be prepared by thermal rearrangement of compounds of formula (AN), optionally in the presence of a suitable solvent and optionally under microwave irradiation.




Preferably, the rearrangement is effected by heating compounds of formula (AN) at temperatures of between 120-300 °C, optionally in a suitable solvent such as 1,2-dimethoxyethane, diethylene glycol methyl ether, triglyme, tetraglyme, xylene, mesitylene or Dowtherm®, and optionally under microwave irradiation.

Similarly, compounds of formula (O), which are compounds of formula (K) wherein G is hydrogen and Het is (Het₃) when R⁶ is CH₂R^{''''} and R^{''''} is hydrogen or methyl, may be prepared from compounds of formula (P) using similar methods.


- 37 -




Compounds of formula (N) may be prepared from compounds of formula (Q) by alkylation with compounds of formula (R), wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, optionally in the presence of a suitable base and optionally in a suitable solvent as described above for the alkylation of compounds of formula (A)



Similarly, compounds of formula (P) may be prepared from compounds of formula (Q) by alkylation with compounds of formula (S), wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, under similar conditions.




In an alternative approach, compounds of formula (N) may be prepared from compounds of formula (Q) by condensation with alcohols of formula (T), optionally in the presence of a suitable acid catalyst such as *p*-toluenesulfonic acid, or a Lewis acid catalyst, for example, ytterbium (III) trifluoromethanesulfonate, lanthanum (III) trifluoromethanesulfonate, sodium tetrachloroaurate (III) dihydrate, titanium (IV) chloride, indium (III) chloride or aluminium chloride, and optionally in a suitable solvent. Suitable solvents are selected to be compatible with the reagents used, and include, for example, toluene, ethanol or acetonitrile. Similar approaches have been described by, for example, M. Curini; F. Epifano, S. Genovese, *Tetrahedron Lett.* (2006), 47, 4697-700; A. Arcadi, G. Bianchi, S. Di Giuseppe, F. Marinelli, *Green Chemistry* (2003), 5, 64-7.




Alternatively, the condensation may be effected in the presence of suitable coupling agents such as 2-chloro-1-methylpyridinium iodide, N,N'-dicyclohexylcarbodiimide, 1, (3-dimethylaminopropyl)-3-ethylcarbodiimide and N,N-carbodiimidazole and a suitable base such as triethylamine or pyridine in a suitable solvent such as tetrahydrofuran, acetonitrile or dichloromethane, or in the presence of a triarylphosphine (such as triphenylphosphine) and a dialkyl azidodicarboxylate (preferably diethyl azidodicarboxylate or diisopropyl azidodicarboxylate) and in a suitable solvent such as diethyl ether, tetrahydrofuran or 1,4-dioxane as described, for example, by O. Mitsunobu, *Synthesis* (1981), 1, 1-28.

Using similar processes, compounds of formula (P) may be prepared by reaction of compounds of formula (Q) with compounds of formula (U).

- 39 -




Additional compounds of formula (N) wherein R^7 is an aromatic or heteroaromatic moiety, or is an alkyl, alkenyl or alkynyl group, may be prepared by the reaction of compounds of formula (V), wherein A is an atom or group suitable for undergoing cross-coupling reactions (for example A is chlorine, bromine or iodine, or a haloalkylsulfonate such as trifluoromethanesulfonate), and R'''' is as defined for compound of formula (O), with a suitable coupling partner under conditions described in the literature for Suzuki-Miyaura, Sonogashira, Stille and related cross-coupling reactions.



For example, compounds of formula (V) may be treated with aryl-, heteroaryl-, alkyl-, alkenyl- or alkynylboronic acids, $R^7\text{-B(OH)}_2$, boronate esters, $R^7\text{-B(OR}'')_2$, wherein R'' is C_1 - C_6 alkyl or $R^7\text{-B(OR}'')_2$ represents cyclic boronate esters derived from a C_1 - C_6 diol (especially preferred are cyclic boronate esters derived from pinacol), or a metal (especially potassium) aryl-, heteroaryl-, alkyl-, alkenyl- and alkynyltrifluoroborate salts, $M^+[\text{R}^7\text{-BF}_3]^-$ in the presence of a suitable palladium catalyst, a suitable ligand and a suitable base in the presence of a suitable solvent, under Suzuki-Miyaura conditions (see, for example K. Billingsley and S. Buchwald, J. Am. Chem. Soc. (2007), 129, 3358-3366; H. Stefani, R. Cella and A. Vieira, Tetrahedron (2007), 63, 3623-3658; N. Kudo, M. Perseghini and G. Fu, Angew. Chem. Int. Ed. (2006), 45, 1282-1284; A. Roglans, A. Pla-Quintana and M. Moreno-

Mañas, Chem. Rev. (2006), 106, 4622-4643; J-H Li, Q-M Zhu and Y-X Xie, Tetrahedron (2006), 10888-10895; S. Nolan *et al.*, J. Org. Chem. (2006), 71, 685-692; M. Lysén and K. Köhler, Synthesis (2006), 4, 692-698; K. Anderson and S. Buchwald, Angew. Chem. Int. Ed. (2005), 44, 6173-6177; Y. Wang and D. Sauer, Org. Lett. (2004), 6 (16), 2793-2796; I. Kondolff, H. Doucet and M. Santelli, Tetrahedron, (2004), 60, 3813-3818; F. Bellina, A. Carpita and R. Rossi, Synthesis (2004), 15, 2419-2440; H. Stefani, G. Molander, C-S Yun, M. Ribagorda and B. Biolatto, J. Org. Chem. (2003), 68, 5534-5539; A. Suzuki, Journal of Organometallic Chemistry (2002), 653, 83; G. Molander and C-S Yun, Tetrahedron (2002), 58, 1465-1470; G. Zou, Y. K. Reddy and J. Falck, Tetrahedron Lett. (2001), 42, 4213-7215; S. Darses, G. Michaud and J-P, Genêt, Eur. J. Org. Chem. (1999), 1877-1883).




Alternatively, compounds of formula (N), wherein R^7 is an optionally substituted acetylene, may be prepared from compounds of formula (V) by reacting with a terminal alkyne, $\text{R}^7\text{-H}$, in the presence of a suitable palladium catalyst and optionally in the presence of a suitable copper co-catalyst, a suitable ligand, a suitable base and a suitable additive under conditions known to effect the Sonogashira coupling (see, for example, U. Sorenson and E Pombo-Villar, Tetrahedron (2005), 2697-2703; N. Leadbeater and B. Tominack, Tetrahedron Lett. (2003), 44, 8653-8656; K. Sonogashira, J. Organomet. Chem. (2002), 653, 46-49).


In a further approach, compounds of formula (N), wherein R^7 is alkyl, optionally substituted vinyl, optionally substituted ethynyl, optionally substituted aryl or optionally substituted heteroaryl, may be prepared from compounds of formula (V) by reaction with a suitable organostannane under Stille conditions (see, for example, R. Bedford, C. Cazin and S. Hazlewood (2002), 22, 2608-2609; S. Ley *et al.*, Chem. Commun. (2002), 10, 1134-1135; G. Grasa and S. Nolan, Org. Lett. (2001), 3 (1), 119-122; T. Weskamp, V. Boehm, J. Organomet. Chem. (1999), 585 (2), 348-352; A. Littke and G. Fu, Angew. Chem. Int. Ed. (1999), 38 (16), 2411-2413; J. Stille *et al.*, Org. Synth. (1992), 71, 97).

- 41 -


Compounds of formula (P) may be prepared from compounds of formula (W), wherein A and R^{''''} are as defined for compounds of formula (V), by analogous methods using appropriate starting materials.



Compounds of formula (V) may be prepared from compounds of formula (Q), by reaction with compounds of formula (Z) wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, by processes analogous to those described above for the preparation of compounds of formula (N) from compounds of formula (Q). Alternatively, compounds of formula (V) may be prepared by reaction of compounds of formula (Q) with compounds of formula (AA) by processes analogous to those described above for the preparation of compounds of formula (N) from compounds of formula (Q).




Compounds of formula (W) may be prepared from compounds of formula (Q), by reaction with compounds of formula (AB) wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, by processes analogous to those described above for the preparation of compounds of formula (N) from compounds of formula (Q). Alternatively, compounds of formula (W) may be prepared by reaction of compounds of formula (Q) with compounds of formula (AB) by processes analogous to those described above for the preparation of compounds of formula (N) from compounds of formula (Q).



In an alternative approach, compounds of formula (Q) may be treated with a halogenating agent such as phosphorus oxychloride, phosphorus pentachloride, phosphorus pentabromide, phosphorus oxybromide, oxalyl chloride or oxalyl bromide, optionally in a suitable solvent such as toluene, chloroform, dichloromethane with optionally the presence of dimethylformamide, and the resulting vinyl halides of formula (AD), wherein Hal is chlorine or bromine may be converted by reaction with alcohols of formula (T), or of formula (U), or of formula (AA) or of formula (AC) optionally in the presence of a suitable base such as sodium hydride, sodium *tert*-butoxide, potassium *tert*-butoxide and a suitable solvent such as tetrahydrofuran, 1,4-dioxane, diethylene glycol dimethyl ether to give compounds of formula (N), formula (P), formula (V) and formula (W) respectively:


- 43 -




Compounds of formula (Q) are known compounds or may be made from known compounds by known methods.

Those skilled in the art will appreciate that compounds of formula (K) may contain a heteroaromatic moiety bearing one or more substituents capable of being transformed into alternative substituents under known conditions, and that these compounds may themselves serve as intermediates in the preparation of additional compounds of formula (K). For example, a heterocycle of formula (M) wherein R⁷ is alkenyl or alkynyl, may be reduced to compounds of formula (M) wherein R⁷ is alkyl under known conditions.

- 44 -




In a further approach to compounds of formula (M), wherein Het is a group of formula (Het₂), X is S, and Z is N, compounds of formula (AE) wherein L is a suitable leaving group such as a halogen or an alkyl- or haloalkylsulfonate, may be treated with compounds of formula (AF) in the presence of a suitable base (such as triethylamine or pyridine), and optionally in a suitable solvent (such as water, acetone, ethanol or isopropanol) according to known procedures, (see, for example, E. Knott, J. Chem. Soc. (1945), 455; H. Brederick, R. Gompper, Chem. Ber. (1960), 93, 723; B. Friedman, M. Sparks and R. Adams, J. Am. Chem. Soc. (1937), 59, 2262).



Alternatively, compounds of formula (AE) may be treated with thiourea, by known procedures (see, for example, V. Pshenichniya, O. Gulyakevich and V. Kripach, Chemistry of Heterocyclic Compounds (1990), 10, 1409-1412), and the resulting products of formula (AG) may be converted into additional compounds of formula (K) by conversion to halides of formula (AH), wherein Hal is chlorine, bromine or iodine, under Sandmeyer conditions, and compounds of formula (AH) may be converted to compounds of formula (K) by cross-coupling under known conditions for the Suzuki-Miyaura, Sonogashira, Stille and related reactions, as described previously.

- 45 -




Compounds of formula (AE) may be prepared from compounds of formula (Q) under known conditions (see, for example, V. Pshenichniya, O. Gulyakevich and V. Kripach, *Chemistry of Heterocyclic Compounds* (1990), 10, 1409-1412; V. Pshenichniya, O. Gulyakevich and V. Kripach, *Russian Journal of Organic Chemistry* (1989), 25 (9), 1882-1888).

Compounds of formula (R), formula (S), formula (T), formula (U), formula (Z), formula (AA), formula (AB) and formula (AC) are known or may be prepared by known methods from known compounds (see, for example T. T. Denton, X. Zhang, J. R. Cashman, *J. Med. Chem.* (2005), 48, 224-239; J. Reinhard, W. E. Hull, C.-W. von der Lieth, U. Eichhorn, H.-C. Kliem, *J. Med. Chem.* (2001), 44, 4050-4061; H. Kraus and H. Fiege, DE19547076; M. L. Boys, L. A. Schretzman, N. S. Chandrakumar, M. B. Tollefson, S. B. Mohler, V. L. Downs, T. D. Penning, M. A. Russell, J. A. Wendt, B. B. Chen, H. G. Stenmark, H. Wu, D. P. Spangler, M. Clare, B. N. Desai, I. K. Khanna, M. N. Nguyen, T. Duffin, V. W. Engleman, M. B. Finn, S. K. Freeman, M. L. Hanneke, J. L. Keene, J. A. Klover, G. A. Nickols, M. A. Nickols, C. N. Steininger, M. Westlin, W. Westlin, Y. X. Yu, Y. Wang, C. R. Dalton, S. A. Norring, *Bioorg.*

- 46 -

Med. Chem. Lett. (2006), 16, 839-844; A. Silberg, A. Benko, G. Csavassy, Chem. Ber. (1964), 97, 1684-1687; K. Brown and R. Newbury, Tetrahedron Lett. (1969), 2797; A. Jansen and M. Szelke, J. Chem. Soc. (1961), 405; R. Diaz-Cortes, A. Silva and L. Maldonado, Tetrahedron Lett. (1997), 38(13), 2007-2210; M. Friedrich, A. Waechtler and A. De Meijure, Synlett. (2002), 4, 619-621; F. Kerdesky and L. Seif, Synth. Commun. (1995), 25 (17), 2639-2645; Z. Zhao, G. Scarlato and R. Armstrong, Tetrahedron Lett. (1991), 32 (13), 1609-1612; K-T. Kang and S. Jong, Synth. Commun. (1995), 25 (17), 2647-2653; M. Altamura and E. Perrotta, J. Org. Chem. (1993), 58 (1), 272-274).


Alternatively, compounds of formula (I) where R¹ and R⁴ form a bond can be prepared from compounds of formula (AJ) by known methods (see for example Nagaoka, H. *et al.* Tetrahedron Letters (1985), 26 (41), 5053-5056; Nagaoka, H. *et al.* J. Am. Chem. Soc. (1986), 108 (16), 5019-5021; Zuki, M. *et al.* Bull. Chem. Soc. Japan (1988), 61(4), 1299-1312; Enholm, E. J. *et al.* J. Org. Chem. (1996), 61 (16), 5384-5390; Clive, D. L. J. *et al.* Tetrahedron (2001), 57 (18), 3845-3858; Bartoli, G. *et al.* J. Org. Chem. (2002), 67 (25), 9111-9114. Jung, M. E. *et al.* Chem. Comm. (2003), (2), 196-197; EP1433772; JP2004203844; IN194295)



wherein R¹ and R⁴ form a bond


Compounds of formula (AJ) may be prepared by treating compounds of formula (K) with compounds of formula (AK) under basic conditions. Suitable bases include lithium diisopropylamide, sodium hexamethyldisilazide, potassium *tert*-butoxide and the reaction is preferably conducted in a suitable solvent (such as tetrahydrofuran) at a temperature between -80 °C and 30 °C

- 47 -




Compounds of formula (AK) are known, or may be made from known compounds by known methods.

Compounds of formula (I) (wherein G is C₁-C₄alkyl) may be prepared by reacting a compounds of formula (AL) (wherein G is C₁-C₄alkyl, and Hal is a halogen, preferably bromine or iodine), with heteroaryl boronic acids, Het-B(OH)₂ of formula (AM) or heteroaryl boronate esters in the presence of a suitable palladium catalyst (for example 0.001-50% palladium(II) acetate with respect to compound (AL)) and a base (for example 1 to 10 equivalents potassium phosphate with respect to compound (AL)) and preferably in the presence of a suitable ligand (for example 0.001-50% (2-dicyclohexylphosphino)-2',6'-dimethoxybiphenyl with respect to compound (AL)), and in a suitable solvent (for example toluene or 1,2-dimethoxyethane), preferably between 25 °C and 200 °C under conventional heating or under microwave irradiation (see, for example, Song, Y. S. S. *et al.* Tetrahedron Lett. (2005), 46 (46), 5987-5990).



Preferred coupling partners include heteroarylboronic acids, (P₁) to (P₈), wherein R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, X, W¹, W², W³, W⁴ and Z are as defined above.


- 48 -




Heteroarylboronic acids of formula (AM) are known compounds, or may be prepared from known compounds by known methods (see for example A. Voisin *et al.*, *Tetrahedron* (2005), 1417-1421; A. Thompson *et al.*, *Tetrahedron* (2005), 61, 5131-5135; K. Billingsley and S. Buchwald, *J. Am. Chem. Soc.* (2007), 129, 3358-3366; N. Kudo, M. Pauro and G. Fu, *Angew. Chem. Int. Ed.* (2006), 45, 1282-1284; A. Ivachtchenko *et al.*, *J. Heterocyclic Chem.* (2004), 41(6), 931-939; H. Matondo *et al.*, *Synth. Commun.* (2003), 33 (5) 795-800; A. Bouillon *et al.*, *Tetrahedron* (2003), 59, 10043-10049; W. Li *et al.*, *J. Org. Chem.* (2002), 67, 5394-5397; C. Enguehard *et al.*, *J. Org. Chem.* (2000), 65, 6572-6575; H-N Nguyen, X. Huang and S. Buchwald, *J. Am. Chem. Soc.* (2003), 125, 11818-11819, and references therein).

Compounds of formula (AL) may be prepared by halogenating compounds of formula (AN), followed by alkylation of the resulting halide of formula (AO) with a C₁-C₄alkyl halide or tri-C₁-C₄alkylorthoformate under known conditions, for example by the procedures of Shepherd R. G. *et al.* *J. Chem. Soc. Perkin Trans. 1* (1987), 2153-2155 and Lin Y. -L. *et al.* *Bioorg. Med. Chem.* (2002), 10, 685-690. Alternatively, compounds of formula (AL) may be prepared by alkylating a compound of formula (AN) with a C₁₋₄ alkyl halide or a tri-C₁₋₄-alkylorthoformate, and halogenating the resulting enone of formula (AP) under known conditions (see for example Song, Y. S. *et al.* *Tetrahedron Lett.* (2005), 46 (36), 5987-5990; Kuethe, J. T. *et al.* *J. Org. Chem.* (2002), 67(17), 5993-6000; Belmont, D. T. *et al.* *J. Org. Chem.* 1985, 50 (21), 4102-4107).

- 49 -




Compounds of formula (AP) may be prepared by treating compounds of formula (AQ) with compounds of formula (L) wherein LG is a leaving group such as halogen (preferably iodide or bromide) or an activated alcohol (preferably mesylate or tosylate) under basic conditions. Suitable bases include lithium diisopropylamide, sodium hexamethyldisilazide, potassium *tert*-butoxide and the reaction is preferably conducted in a suitable solvent (such as tetrahydrofuran) at a temperature between -80 °C and 30 °C (see, for example, Gulias, M. *et al.* Org. Lett. (2003), 5(11), 1975-1977; Altenbach, R. J. *et al.* J. Med. Chem. (2006), 49 (23), 6869-6887; Snowden, R. L. Tetrahedron (1986), 42 (12), 3277-90; Oppolzer, W. *et al.* Helv. Chim. Acta (1980), 63 (4), 788-92; Mellor, M. *et al.* Synth. Commun. 1979, 9 (1), 1-4).




- 50 -

Compounds of formula (AQ) are known, or may be made from known compounds by known methods.


Alternatively compounds of formula (AP) where R^1 and R^4 form a bond can be prepared from compounds of formula (AR) by known methods (see, for example, Nagaoka, H. *et al.* Tetrahedron Letters (1985), 26 (41), 5053-5056; Nagaoka, H. *et al.* J. Am. Chem. Soc. (1986), 108 (16), 5019-5021; zuki, M. *et al.* Bull. Chem. Soc. Japan (1988), 61(4), 1299-1312; Enholm, E. J. *et al.* J. Org. Chem. (1996), 61 (16), 5384-5390; Clive, D. L. J. *et al.* Tetrahedron (2001), 57 (18), 3845-3858; Bartoli, G. *et al.* J. Org. Chem. (2002), 67 (25), 9111-9114. Jung, M. E. *et al.* Chem. Comm. (2003), (2), 196-197; EP1433772; JP2004203844; IN194295).



Compounds of formula (AR) may be prepared by treating compounds of formula (AQ where in R^1 is hydrogen) with compounds of formula (AK) under basic conditions. Suitable bases include lithium diisopropylamide, sodium hexamethyldisilazide, potassium *tert*-butoxide and the reaction is preferably conducted in a suitable (such as tetrahydrofuran) at a temperature between -80 °C and 30 °C (see, for example, Aleman, J. *et al.* Chem. Comm. (2007), (38), 3921-3923).



Compounds of formula (AN) may be prepared from compounds of formula (AP) by hydrolysis, preferably in the presence of an acid catalyst such as hydrochloric acid and optionally in the presence of a suitable solvent such as tetrahydrofuran or acetone preferably between 25 °C and 150 °C under conventional heating or under microwave irradiation.




Alternatively, compounds of formula (AN) can be made from known compounds by known methods (see for example Manukina, T. A. *et al.* Zhurnal Organicheskoi Khimii (1986), 22(4), 873-4; Mellor, M. *et al.* Synth. Commun. 1979, 9 (1), 1-4).

In a further approach, compounds of formula (A) may be prepared by reacting compounds of formula (AN) with suitable heteroaryl halides (Het-Hal where Hal is, for example, an iodide or bromide), in the presence of a suitable palladium catalyst (for example 0.001-50% palladium(II) acetate with respect to compound (AN)) and a base (for example 1 to 10 equivalents potassium phosphate with respect to compound (AN)) and preferably in the presence of a suitable ligand (for example 0.001-50% (2-dicyclohexylphosphino)-2',4',6'-triisopropylbiphenyl with respect to compound (AN)), and in a suitable solvent (for example 1,4-dioxane), preferably between 25°C and 200°C. Similar couplings are known in the literature (see for example, J. M. Fox, X. Huang, A. Chieffi, and S. L. Buchwald, J. Am. Chem. Soc. (2000), 122, 1360-1370; B. Hong *et al.* WO 2005/000233).


Alternatively, compounds of formula (A) may be prepared by reacting compounds of formula (AN) with suitable heteroaryl halides (Het-Hal where Hal is, for example, an iodide or bromide) in the presence of a suitable copper catalyst (for example 0.001-50% copper(I) iodide with respect to compound (AN)) and a base (for example 1 to 10 equivalents potassium carbonate with respect to compound (AN)) and preferably in the presence of a suitable ligand (for example 0.001-50% L-proline with respect to compound (AN)), and in a suitable solvent (for example dimethylsulfoxide), preferably between 25°C and 200°C. Similar

- 52 -

couplings are known in the literature for aryl halides (see for example, Y. Jiang, N. Wu, H. Wu, and M. He, *Synlett* (2005), 18, 2731-2734).




In a further approach, compounds of formula (A) may be prepared by reaction of compounds of formula (AN) with a heteroaryl lead tricarboxylates under conditions described in the literature (for example see, J. T. Pinhey, B. A. Rowe, *Aust. J. Chem.* (1979), 32, 1561-6; J. Morgan, J. T. Pinhey, *J. Chem. Soc. Perkin Trans. 1* (1990), 3, 715-20; J. T. Pinhey, Roche, E. G. *J. Chem. Soc. Perkin Trans. 1* (1988), 2415-21). Preferably the heteroaryl lead tricarboxylates are heteroaryl triacetates of formula (AS) and the reaction is conducted in the presence of a suitable ligand (for example N,N-dimethylaminopyridine, pyridine, imidazole, bipyridine, and 1,10-phenanthroline, preferably one to ten equivalents of N,N-dimethylaminopyridine with respect to compound (AN)) and in a suitable solvent (for example chloroform, dichloromethane and toluene, preferably chloroform and optionally in the presence of a co-solvent such as toluene) at 25°C to 100°C (preferably 60-90°C).




Compounds of formula (AS) may be prepared from compounds of formula (AM) by treatment with lead tetraacetate in a suitable solvent (for example chloroform) at 25°C to 100°C (preferably 25-50°C), optionally in the presence of a catalyst such as mercury diacetate, according to procedures described in the literature (for example see, K. Shimi, G. Boyer, J-P. Finet and J-P. Galy, *Letters in Organic Chemistry* (2005), 2, 407-409; J. Morgan and J. T. Pinhey, *J. Chem. Soc. Perkin Trans. 1* (1990), 3, 715-20).

- 53 -



Additional compounds of formula (A) may be prepared by reacting an iodonium ylide of formula (AT), wherein Ar is an optionally substituted phenyl group, and an aryl boronic acid of formula (AM), in the presence of a suitable palladium catalyst, a base and in a suitable solvent.




Suitable palladium catalysts are generally palladium(II) or palladium(0) complexes, for example palladium(II) dihalides, palladium(II) acetate, palladium(II) sulfate, bis(triphenylphosphine)-palladium(II) dichloride, bis(tricyclopentylphosphine)palladium(II) dichloride, bis(tricyclohexylphosphine)palladium(II) dichloride, *bis(dibenzylideneacetone)palladium(0)* or *tetrakis-(triphenylphosphine)palladium(0)*. The palladium catalyst can also be prepared *in situ* from palladium(II) or palladium(0) compounds by complexing with the desired ligands, by, for example, combining the palladium(II) salt to be complexed, for example palladium(II) dichloride (PdCl₂) or palladium(II) acetate (Pd(OAc)₂), together with the desired ligand, for example triphenylphosphine (PPh₃), tricyclopentylphosphine, tricyclohexylphosphine, 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl or 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl and the selected solvent, with a compound of formula (AT), the arylboronic acid of formula (AM), and a base. Also suitable are bidendate ligands, for example 1,1'-bis(diphenylphosphino)ferrocene or 1,2-bis(diphenylphosphino)ethane. By heating the reaction medium, the palladium(II) complex or palladium(0) complex desired for the C-C coupling reaction is thus formed *in situ*, and then initiates the C-C coupling reaction.


The palladium catalysts are used in an amount of from 0.001 to 50 mol %, preferably in an amount of from 0.1 to 15 mol %, based on the compound of formula (AT). The reaction may

also be carried out in the presence of other additives, such as tetralkylammonium salts, for example, tetrabutylammonium bromide. Preferably the palladium catalyst is palladium acetate, the base is lithium hydroxide and the solvent is aqueous 1,2-dimethoxyethane.

A compound of formula (AT) may be prepared from a compound of formula (AN) by treatment with a hypervalent iodine reagent such as a (diacetoxy)iodobenzene or an iodosylbenzene and a base such as aqueous sodium carbonate, lithium hydroxide or sodium hydroxide in a solvent such as water or an aqueous alcohol such as aqueous ethanol according to the procedures of Schank K. *et al.* Synthesis (1983), 392, Moriarty R. M. *et al.* J. Am. Chem. Soc. (1985), 107, 1375 or of Yang Z. *et al.* Org. Lett. (2002), 4 (19), 3333.




In a further approach to compounds of formula (A), compounds of formula (AU), which are compounds of formula (I) wherein G is hydrogen and Het is (Het₂) when R⁶ is CH₂R^{''''} and R^{''''} is hydrogen or methyl, may be prepared by thermal rearrangement of compounds of formula (AV), optionally in the presence of a suitable solvent and optionally under microwave irradiation.




Preferably, the rearrangement is effected by heating compounds of formula (AT) at temperatures of between 120-300 °C, optionally in a suitable solvent such as 1,2-dimethoxyethane, diethylene glycol methyl ether, triglyme, tetraglyme, xylene, mesitylene or Dowtherm®, and optionally under microwave irradiation.

Similarly, compounds of formula (AW), which are compounds of formula (I) wherein G is hydrogen and Het is (Het₃) when R⁶ is CH₂R^{''''} and R^{''''} is hydrogen or methyl, may be prepared from compounds of formula (AX) using similar methods.




Compounds of formula (AV) may be prepared from compounds of formula (AN) by alkylation with compounds of formula (R), wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, optionally in the presence of a suitable base and optionally in a suitable solvent as described above for the alkylation of compounds of formula (A)




Similarly, compounds of formula (AX) may be prepared from compounds of formula (AN) by alkylation with compounds of formula (S), wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, under similar conditions.

- 56 -



In an alternative approach, compounds of formula (AV) may be prepared from compounds of formula (AN) by condensation with alcohols of formula (T), optionally in the presence of a suitable acid catalyst such as *p*-toluenesulfonic acid, or a Lewis acid catalyst, for example, ytterbium (III) trifluoromethanesulfonate, lanthanum (III) trifluoromethanesulfonate, sodium tetrachloroaurate (III) dihydrate, titanium (IV) chloride, indium (III) chloride or aluminium chloride, and optionally in a suitable solvent. Suitable solvents are selected to be compatible with the reagents used, and include, for example, toluene, ethanol or acetonitrile. Similar approaches have been described by, for example, M. Curini; F. Epifano, S. Genovese, *Tetrahedron Lett.* (2006), 47, 4697-700; A. Arcadi, G. Bianchi, S. Di Giuseppe, F. Marinelli, *Green Chemistry* (2003), 5, 64-7.




Alternatively, the condensation may be effected in the presence of suitable coupling agents such as 2-chloro-1-methylpyridinium iodide, *N,N'*-dicyclohexylcarbodiimide, 1, (3-dimethylaminopropyl)-3-ethylcarbodiimide and *N,N*-carbodiimidazole and a suitable base such as triethylamine or pyridine in a suitable solvent such as tetrahydrofuran, acetonitrile or


- 57 -

dichloromethane, or in the presence of a triarylphosphine (such as triphenylphosphine) and a dialkyl azidodicarboxylate (preferably diethyl azidodicarboxylate or diisopropyl azidodicarboxylate) and in a suitable solvent such as diethyl ether, tetrahydrofuran or 1,4-dioxane as described, for example, by O. Mitsunobu, *Synthesis* (1981), 1, 1-28.


Using similar processes, compounds of formula (AX) may be prepared by reaction of compounds of formula (AN) with compounds of formula (U).



Additional compounds of formula (AV) wherein R⁷ is an aromatic or heteroaromatic moiety, or is an alkyl, alkenyl or alkynyl group, may be prepared by the reaction of compounds of formula (AY), wherein A is an atom or group suitable for undergoing cross-coupling reactions (for example A is chlorine, bromine or iodine, or a haloalkylsulfonate such as trifluoromethanesulfonate), and R¹¹ is as defined for compound of formula (O), with a suitable coupling partner under conditions described in the literature for Suzuki-Miyaura, Sonogashira, Stille and related cross-coupling reactions.



For example, compounds of formula (AY) may be treated with aryl-, heteroaryl-, alkyl-, alkenyl- or alkynylboronic acids, $R^7\text{-B(OH)}_2$, boronate esters, $R^7\text{-B(OR}'')_2$, wherein R'' is $C_1\text{-}C_6$ alkyl or $R^7\text{-B(OR}'')_2$ represents cyclic boronate esters derived from a $C_1\text{-}C_6$ diol (especially preferred are cyclic boronate esters derived from pinacol), or a metal (especially potassium) aryl-, heteroaryl, alkyl-, alkenyl- and alkynyltrifluoroborate salts, $M^+[\text{R}^7\text{-BF}_3]^-$ in the presence of a suitable palladium catalyst, a suitable ligand and a suitable base in the presence of a suitable solvent, under Suzuki-Miyaura conditions (see, for example K. Billingsley and S. Buchwald, *J. Am. Chem. Soc.* (2007), 129, 3358-3366; H. Stefani, R. Cella and A. Vieira, *Tetrahedron* (2007), 63, 3623-3658; N. Kudo, M. Perseghini and G. Fu, *Angew. Chem. Int. Ed.* (2006), 45, 1282-1284; A. Roglans, A. Pla-Quintana and M. Moreno-Mañas, *Chem. Rev.* (2006), 106, 4622-4643; J-H Li, Q-M Zhu and Y-X Xie, *Tetrahedron* (2006), 10888-10895; S. Nolan *et al.*, *J. Org. Chem.* (2006), 71, 685-692; M. Lysén and K. Köhler, *Synthesis* (2006), 4, 692-698; K. Anderson and S. Buchwald, *Angew. Chem. Int. Ed.* (2005), 44, 6173-6177; Y. Wang and D. Sauer, *Org. Lett.* (2004), 6 (16), 2793-2796; I. Kondolff, H. Doucet and M. Santelli, *Tetrahedron*, (2004), 60, 3813-3818; F. Bellina, A. Carpita and R. Rossi, *Synthesis* (2004), 15, 2419-2440; H. Stefani, G. Molander, C-S Yun, M. Ribagorda and B. Biolatto, *J. Org. Chem.* (2003), 68, 5534-5539; A. Suzuki, *Journal of Organometallic Chemistry* (2002), 653, 83; G. Molander and C-S Yun, *Tetrahedron* (2002), 58, 1465-1470; G. Zou, Y. K. Reddy and J. Falck, *Tetrahedron Lett.* (2001), 42, 4213-7215; S. Darses, G. Michaud and J-P, Genêt, *Eur. J. Org. Chem.* (1999), 1877-1883).




Alternatively, compounds of formula (AV), wherein R^7 is an optionally substituted acetylene, may be prepared from compounds of formula (AY) by reacting with a terminal alkyne, $R^7\text{-H}$, in the presence of a suitable palladium catalyst and optionally in the presence of a suitable copper co-catalyst, a suitable ligand, a suitable base and a suitable additive under conditions known to effect the Sonogashira coupling (see, for example, U. Sorenson and E Pombo-

Villar, *Tetrahedron* (2005), 2697-2703; N. Leadbeater and B. Tominack, *Tetrahedron Lett.* (2003), 44, 8653-8656; K. Sonogashira, *J. Organomet. Chem.* (2002), 653, 46-49).


In a further approach, compounds of formula (AV), wherein R^7 is alkyl, optionally substituted vinyl, optionally substituted ethynyl, optionally substituted aryl or optionally substituted heteroaryl, may be prepared from compounds of formula (AY) by reaction with a suitable organostannane under Stille conditions (see, for example, R. Bedford, C. Cazin and S. Hazlewood (2002), 22, 2608-2609; S. Ley *et al.*, *Chem. Commun.* (2002), 10, 1134-1135; G. Grasa and S. Nolan, *Org. Lett.* (2001), 3 (1), 119-122; T. Weskamp, V. Boehm, J. *Organomet. Chem.* (1999), 585 (2), 348-352; A. Littke and G. Fu, *Angew. Chem. Int. Ed.* (1999), 38 (16), 2411-2413; J. Stille *et al.*, *Org. Synth.* (1992), 71, 97).

Compounds of formula (AX) may be prepared from compounds of formula (AZ), wherein A and R'''' are as defined for compounds of formula (AY), by analogous methods using appropriate starting materials.




Compounds of formula (AY) may be prepared from compounds of formula (AN), by reaction with compounds of formula (Z) wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, by processes analogous to those described above for the preparation of compounds of formula (AV) from compounds of formula (AN). Alternatively, compounds of formula (AY) may be prepared by reaction of compounds of formula (AN) with compounds of formula (AA) by processes analogous to those described above for the preparation of compounds of formula (AV) from compounds of formula (AN).

- 60 -




Compounds of formula (AZ) may be prepared from compounds of formula (AN), by reaction with compounds of formula (AB) wherein L is a suitable leaving group such as a halogen or an alkyl- or aryl-sulfonate, by processes analogous to those described above for the preparation of compounds of formula (AV) from compounds of formula (AN). Alternatively, compounds of formula (AZ) may be prepared by reaction of compounds of formula (AN) with compounds of formula (AB) by processes analogous to those described above for the preparation of compounds of formula (AV) from compounds of formula (AN).

- 61 -




In an alternative approach, compounds of formula (AN) may be treated with a halogenating agent such as phosphorus oxychloride, phosphorus pentachloride, phosphorus pentabromide, phosphorus oxybromide, oxalyl chloride or oxalyl bromide, optionally in a suitable solvent such as toluene, chloroform, dichloromethane with optionally the presence of dimethylformamide, and the resulting vinyl halides of formula (BA), wherein Hal is chlorine or bromine may be converted by reaction with alcohols of formula (T), or of formula (U), or of formula (AA) or of formula (AC) optionally in the presence of a suitable base such as sodium hydride, sodium *tert*-butoxide, potassium *tert*-butoxide and a suitable solvent such as tetrahydrofuran, 1,4-dioxane, diethylene glycol dimethyl ether to give compounds of formula (AV), formula (AX), formula (AY) and formula (AZ) respectively:


- 62 -



- 63 -




Compounds of formula (BB) may be prepared from compounds of formula (BC), by reaction 2-chloro-3-iodopropene derivatives under basic conditions. Suitable bases include sodium bis(trimethylsilyl)amide, lithium diisopropylamide, n-butyl lithium and the reaction is preferably conducted in a suitable solvent (such as tetrahydrofuran or toluene) at a temperature of between -80°C and 30°C (see, for example, Y. Xu *et al.* J. Org. Chem. 2009, DOI: 10.1021/jo900696k).



Compounds of formula (BC) are known or may be made from known compounds by known methods.


In a further approach to compounds of formula (A), wherein Het is a group of formula (Het₂), X is S, and Z is N, compounds of formula (BD) wherein L is a suitable leaving group such as a halogen or an alkyl- or haloalkylsulfonate, may be treated with compounds of formula (BE) in the presence of a suitable base (such as triethylamine or pyridine), and optionally in a suitable solvent (such as water, acetone, ethanol or isopropanol) according to known procedures, (see, for example, E. Knott, J. Chem. Soc. (1945), 455; H. Brederick, R. Gompper, Chem. Ber. (1960), 93, 723; B. Friedman, M. Sparks and R. Adams, J. Am. Chem. Soc. (1937), 59, 2262).

- 64 -



Alternatively, compounds of formula (BD) may be treated with thiourea, by known procedures (see, for example, V. Pshenichniya, O. Gulyakevich and V. Kripach, *Chemistry of Heterocyclic Compounds* (1990), 10, 1409-1412), and the resulting products of formula (BF) may be converted into additional compounds of formula (I) by conversion to halides of formula (BG), wherein Hal is chlorine, bromine or iodine, under Sandmeyer conditions, and compounds of formula (BG) may be converted to compounds of formula (I) by cross-coupling under known conditions for the Suzuki-Miyaura, Sonogashira, Stille and related reactions, as described previously.


- 65 -




Compounds of formula (BD) may be prepared from compounds of formula (AN) under known conditions (see, for example, V. Pshenichniya, O. Gulyakevich and V. Kripach, *Chemistry of Heterocyclic Compounds* (1990), 10, 1409-1412; V. Pshenichniya, O. Gulyakevich and V. Kripach, *Russian Journal of Organic Chemistry* (1989), 25 (9), 1882-1888).

Additional compounds of formula (A) may be prepared by the pinacol rearrangement of compounds of formula (BG) or compounds of formula (BJ) wherein $\text{R}^{\text{''''''}}$ is C_1 - C_4 alkyl (preferably methyl) under acidic conditions (see, for example, Eberhardt, U. *et. al.* *Chem. Ber.* (1983), 116(1), 119-35 and Wheeler, T. N. US4283348)

- 66 -




Compounds of formula (BH) and compounds of formula (BJ) may be prepared by treating compounds of formula (BK) with compounds of formula (BL) in the presence of an acid (such as titanium tetrachloride or magnesium iodide) optionally in a suitable solvent (such as dichloromethane) at a temperature between -80 °C and 30 °C (see, for example, Li, W.-D. Z. and Zhang, X.-X. *Org. Lett.* (2002), 4(20), 3485-3488; Shimada, J. *et al.* *J. Am. Chem. Soc.* (1984), 106(6), 1759-73; Eberhardt, U. *et al.* *Chem. Ber.* (1983), 116(1), 119-35 and Wheeler, T. N. US4283348).




Compounds of formula (BK) are known or may be made from known compounds by known methods.

Compounds of formula (BL) may be prepared from compounds of formula (BM) where in R''' is an alkyl group (preferably methyl) in the presence of chloro tri-C₁-C₄alkyl silyl and a metal (preferably sodium) in a suitable solvent (such as toluene or diethyl ether) at a temperature between 20 °C and 150 °C (see, for example, Blanchard, A. N. and Burnell, D. J. *Tetrahedron Lett.* (2001), 42(29), 4779-4781 and Salaun, J. *et al.* *Tetrahedron* (1989), 45(10), 3151-62).



Compounds of formula (BM) are analogous to compounds of formula (H) and compounds of formula (G) and may be prepared by known methods analogous to those described for compounds of formula (H) and compounds of formula (G).

Additional compounds of formula (I) may be prepared wherein R¹ and R⁴ form a bond and R⁵ is C₁-C₆alkylsulfonate (preferably mesylate) or C₁-C₆haloalkylsulfonate (preferably triflate) or an arylsulfonate (preferably tosylate) may be prepared from compounds of formula (BN) following known procedures (Specklin *et al.* *J. Org. Chem.* 2008, 73(19), 7845-7848).




wherein R¹ and R⁴ form a bond

Compounds of formula (BM) may be prepared from compounds of formula (BO) under basic or acidic conditions. For example of a procedure see G. Quinkert *et al.* *Helv. Chim. Acta*, 1986, 69(3), 469-537.

- 68 -




Compounds of formula (BO) may be prepared by reaction of compounds of formula (K) wherein R^5 is hydrogen with acids chloride of formula (BP) in the presence of a base.



Compounds of formula (BP) are known or may be made by known methods from known compounds.

Alternatively, compounds of formula (BN) can be prepared from compounds of formula (AJ) using known oxidative procedures (see for example D. B. Dess and J. C. Martin J. Org. Chem. 1983, 48 (22), 4155–4156).

- 69 -



wherein R⁵ is hydrogen

The compounds of formula I according to the invention can be used as crop protection agents in unmodified form, as obtained in the synthesis, but they are generally formulated into crop protection compositions in a variety of ways using formulation adjuvants, such as carriers, solvents and surface-active substances.

Hence, the invention also provides a herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined herein.

The formulations (compositions) can be in various physical forms, for example in the form of dusting powders, gels, wettable powders, coated or impregnated granules for manual or mechanical distribution on target sites, water-dispersible granules, water-soluble granules, emulsifiable granules, water-dispersible tablets, effervescent compressed tablets, water-soluble tapes, emulsifiable concentrates, microemulsifiable concentrates, oil-in-water (EW) or water-in-oil (WO) emulsions, other multiphase systems such as oil/water/oil and water/oil/water products, oil flowables, aqueous dispersions, oily dispersions, suspoemulsions, capsule suspensions, soluble liquids, water-soluble concentrates (with water or a water-miscible organic solvent as carrier), impregnated polymer films or in other forms known, for example, from the Manual on Development and Use of FAO Specifications for Plant Protection Products, 5th Edition, 1999. The active ingredient may be incorporated into microfibers or micro-rods formed of polymers or polymerizable monomers and having diameter of about 0.1 to about 50 microns and aspect ratio of between about 10 and about 1000.

Such formulations can either be used directly or are diluted prior to use. They can then be applied through suitable ground or aerial application spray equipment or other ground application equipment such as central pivot irrigation systems or drip/trickle irrigation means.

- 70 -

Diluted formulations can be prepared, for example, with water, liquid fertilisers, micro-nutrients, biological organisms, oil or solvents.

The formulations can be prepared, for example, by mixing the active ingredient with formulation adjuvants in order to obtain compositions in the form of finely divided solids, granules, solutions, dispersions or emulsions. The active ingredients can also be contained in fine microcapsules consisting of a core and a polymeric shell. Microcapsules usually have a diameter of from 0.1 to 500 microns. They contain active ingredients in an amount of about from 25 to 95 % by weight of the capsule weight. The active ingredients can be present in the form of liquid technical material, in the form of a suitable solution, in the form of fine particles in solid or liquid dispersion or as a monolithic solid. The encapsulating membranes comprise, for example, natural and synthetic gums, cellulose, styrene-butadiene copolymers or other similar suitable membrane forming material, polyacrylonitrile, polyacrylate, polyester, polyamides, polyureas, polyurethane, aminoplast resins or chemically modified starch or other polymers that are known to the person skilled in the art in this connection.

Alternatively it is possible for fine so called "microcapsules" to be formed wherein the active ingredient is present in the form of finely divided particles in a solid matrix of a base substance, but in that case the microcapsule is not encapsulated with a diffusion limiting membrane as outlined in the preceding paragraph.

The active ingredients may be adsorbed on a porous carrier. This may enable the active ingredients to be released into their surroundings in controlled amounts (e.g. slow release). Other forms of controlled release formulations are granules or powders in which the active ingredient is dispersed or dissolved in a solid matrix consisting of a polymer, a wax or a suitable solid substance of lower molecular weight. Suitable polymers are polyvinyl acetates, polystyrenes, polyolefins, polyvinyl alcohols, polyvinyl pyrrolidones, alkylated polyvinyl pyrrolidones, copolymers of polyvinyl pyrrolidones and maleic anhydride and esters and half-esters thereof, or chemically modified cellulose esters like carboxymethyl cellulose, methyl cellulose, or hydroxyethyl cellulose. Examples of suitable waxes are polyethylene wax, oxidized polyethylene wax, ester waxes like montan waxes, waxes of natural origin like carnauba wax, candelilla wax, or beeswax etc. Other suitable matrix materials for slow release formulations are starch, stearin, or lignin.

The formulation adjuvants suitable for the preparation of the compositions according to the invention are generally known *per se*.

As liquid carriers there may be used: water, aromatic solvents such as toluene, m-xylene, o-xylene, p-xylene and mixtures thereof, cumene, aromatic hydrocarbon blends with boiling ranges between 140 and 320 °C known under various trademarks like Solvesso[®], Shellsol A[®], Caromax[®], Hydrosol[®], paraffinic and isoparaffinic carriers such as paraffin oils, mineral oils, de-aromatized hydrocarbon solvents with boiling ranges between 50 and 320 °C known for instance under the trademark Exxsol[®], non-dearomatized hydrocarbon solvents with boiling ranges between 100 and 320 °C known under the tradename Varsol[®], isoparaffinic solvents with boiling ranges between 100 and 320 °C known under tradenames like Isopar[®] or Shellsol T[®], hydrocarbons such as cyclohexane, tetrahydronaphthalene (tetralin), decahydronaphthalene, alpha-pinene, d-limonene, hexadecane, isooctane, ester solvents such as ethyl acetate, *n*/*i*-butyl acetate, amyl acetate, *i*-bornyl acetate, 2-ethylhexyl acetate, C₆ – C₁₈ alkyl esters of acetic acid known under the tradename Exxate[®], lactic acid ethylester, lactic acid propylester, lactic acid butylester, benzyl benzoate, benzyl lactate, dipropyleneglycol dibenzoate, dialkyl esters of succinic, maleic and fumaric acid and polar solvents like N-methyl pyrrolidone, N-ethyl pyrrolidone, C₃-C₁₈-alkyl pyrrolidones, gamma-butyrolactone, dimethylsulfoxide, N,N-dimethylformamide, N,N-dimethylacetamide, N,N-dimethyl lactamide, C₄-C₁₈ fatty acid dimethylamides, benzoic acid dimethylamide, acetonitrile, acetone, methyl ethyl ketone, methyl-isobutyl ketone, isoamyl ketone, 2-heptanone, cyclohexanone, isophorone, methyl isobutenyl ketone (mesityl oxide), acetophenone, ethylene carbonate, propylene carbonate, butylene carbonate, alcoholic solvents and diluents such as methanol, ethanol, propanol, *n*/*iso*-butanol, *n*/*iso*-pentanol, 2-ethyl hexanol, *n*-octanol, tetrahydrofurfuryl alcohol, 2-methyl-2,4-pentanediol, 4-hydroxy-4-methyl-2-pentanone, cyclohexanol, benzyl alcohol, ethylene glycol, ethylene glycol butyl ether, ethylene glycol methyl ether, diethylene glycol, diethylene glycol butyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, propylene glycol, dipropylene glycol, dipropylene glycol methyl ether and other similar glycol ether solvents based on ethylene glycol, propylene glycol and butylene glycol feedstocks, triethylene glycol, polyethylene glycol (PEG 400), polypropyleneglycols with molecular masses of 400 - 4000, glycerol, glycerol acetate, glycerol diacetate, glycerol triacetate, 1,4-dioxane, diethylene glycol abietate, chlorobenzene, chlorotoluene, fatty acid esters such as methyl octanoate, isopropyl myristate, methyl laurate, methyl oleate, mixture of C₈-C₁₀ fatty acid methyl esters, rape seed oil methyl and ethyl esters, soy bean oil methyl and ethyl esters, vegetable oils,

- 72 -

fatty acids such as oleic acid, linoleic acid, linolenic acid, esters of phosphoric and phosphonic acid such as triethyl phosphate, C₃-C₁₈-tris-alkyl phosphates, alkylaryl phosphates, bis-octyl-octyl phosphonates.

Water is generally the carrier of choice for the dilution of the concentrates.

Suitable solid carriers are, for example, talc, titanium dioxide, pyrophyllite clay, silica (fumed or precipitated silica and optionally functionalised or treated, for instance silanised), attapulgite clay, kieselguhr, limestone, calcium carbonate, bentonite, calcium montmorillonite, cottonseed husks, wheatmeal, soybean flour, pumice, wood flour, ground walnut shells, lignin and similar materials, as described, for example, in the EPA CFR 180.1001. (c) & (d). Powdered or granulated fertilisers can also be used as solid carriers.

A large number of surface-active substances can advantageously be used both in solid and in liquid formulations, especially in those formulations which can be diluted with a carrier prior to use. Surface-active substances may be anionic, cationic, amphoteric, non-ionic or polymeric and they may be used as emulsifying, wetting, dispersing or suspending agents or for other purposes. Typical surface-active substances include, for example, salts of alkyl sulfates, such as diethanolammonium lauryl sulphate; Sodium lauryl sulphate, salts of alkylarylsulfonates, such as calcium or sodium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol ethoxylates; alcohol-alkylene oxide addition products, such as tridecyl alcohol ethoxylate; soaps, such as sodium stearate; salts of alkylnaphthalenesulfonates, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl)sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride, polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono- and di-alkyl phosphate esters; and also further substances described e.g. in "McCutcheon's Detergents and Emulsifiers Annual", MC Publishing Corp., Ridgewood, New Jersey, 1981.

Further adjuvants which can usually be used in pesticidal formulations include crystallisation inhibitors, viscosity-modifying substances, suspending agents, dyes, anti-oxidants, foaming agents, light absorbers, mixing aids, anti-foams, complexing agents, neutralising or pH-modifying substances and buffers, corrosion-inhibitors, fragrances, wetting agents, absorption improvers, micronutrients, plasticisers, glidants, lubricants, dispersants,

- 73 -

thickeners, anti-freezes, microbiocides, compatibility agents and solubilisers and also liquid and solid fertilisers.

The formulations may also comprise additional active substances, for example further herbicides, herbicide safeners, plant growth regulators, fungicides or insecticides.

Therefore, the invention also provides a herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined herein, and optionally (or preferably) a further herbicide as mixture partner for the compound of formula I, or optionally (or preferably) a safener, or both.

The invention also provides a herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined herein, a safener, and optionally (or preferably) a further herbicide as mixture partner for the compound of formula I,

wherein the safener is benoxacor, cloquintocet-mexyl, cyprosulfamide, mefenpyr-diethyl or N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide.

The compositions according to the invention can additionally include an additive (commonly referred to as an adjuvant), comprising a mineral oil, an oil of vegetable or animal origin, alkyl esters of such oils or mixtures of such oils and oil derivatives. The amount of oil additive used in the composition according to the invention is generally from 0.01 to 10 %, based on the spray mixture. For example, the oil additive can be added to the spray tank in the desired concentration after the spray mixture has been prepared. Preferred oil additives comprise mineral oils or an oil of vegetable origin, for example rapeseed oil, olive oil or sunflower oil, emulsifiable vegetable oil, such as AMIGO® (Loveland Products Inc.), alkyl esters of oils of vegetable origin, for example the methyl derivatives, or an oil of animal origin, such as fish oil or beef tallow. A preferred additive contains, for example, as active components essentially 80 % by weight alkyl esters of fish oils and 15 % by weight methylated rapeseed oil, and also 5 % by weight of customary emulsifiers and pH modifiers. Especially preferred oil additives comprise alkyl esters of C₈-C₂₂ fatty acids, especially the methyl derivatives of C₁₂-C₁₈ fatty acids, for example the methyl esters of lauric acid, palmitic acid and oleic acid, being important. Those esters are known as methyl laurate (CAS-111-82-0), methyl palmitate (CAS-112-39-0) and methyl oleate (CAS-112-62-9). A preferred fatty acid methyl ester derivative is AGNIQUE ME 18 RD-F® (Cognis). Those and other oil derivatives are also

- 74 -

known from the Compendium of Herbicide Adjuvants, 5th Edition, Southern Illinois University, 2000.

The application and action of the oil additives can be further improved by combining them with surface-active substances, such as non-ionic, anionic, cationic or amphoteric surfactants. Examples of suitable anionic, non-ionic, cationic or amphoteric surfactants are listed on pages 7 and 8 of WO97/34485. Preferred surface-active substances are anionic surfactants of the dodecylbenzylsulfonate type, especially the calcium salts thereof, and also non-ionic surfactants of the fatty alcohol ethoxylate type. Special preference is given to ethoxylated C₁₂-C₂₂ fatty alcohols having a degree of ethoxylation of from 5 to 40. Examples of commercially available surfactants are the Genapol types (Clariant). Also preferred are silicone surfactants, especially polyalkyl-oxide-modified heptamethyltrisiloxanes, which are commercially available e.g. as SILWET L-77®, and also perfluorinated surfactants. The concentration of surface-active substances in relation to the total additive is generally from 1 to 50 % by weight. Examples of oil additives that consist of mixtures of oils or mineral oils or derivatives thereof with surfactants are TURBOCHARGE®, ADIGOR® (both (Syngenta Crop Protection AG), ACTIPRON® (BP Oil UK Limited), AGRI-DEX® (Helena Chemical Company).

The said surface-active substances may also be used in the formulations alone, that is to say without oil additives.

Furthermore, the addition of an organic solvent to the oil additive/surfactant mixture can contribute to a further enhancement of action. Suitable solvents are, for example, SOLVESSO® and AROMATIC® solvents (Exxon Corporation). The concentration of such solvents can be from 10 to 80 % by weight of the total weight. Such oil additives, which may be in admixture with solvents, are described, for example, in US 4 834 908. A commercially available oil additive disclosed therein is known by the name MERGE® (BASF). Further oil additives that are preferred according to the invention are SCORE® and ADIGOR® (both Syngenta Crop Protection AG).

In addition to the oil additives listed above, in order to enhance the activity of the compositions according to the invention it is also possible for formulations of alkylpyrrolidones, (e.g. AGRIMAX® from ISP) to be added to the spray mixture. Formulations of synthetic latices,

- 75 -

such as, for example, polyacrylamide, polyvinyl compounds or poly-1-p-menthene (e.g. BOND®, COURIER® or EMERALD®) can also be used.

Such adjuvant oils as described in the preceding paragraphs may be employed as the carrier liquid in which an active compound is dissolved, emulsified or dispersed as appropriate to the physical form of the active compound.

The pesticidal (e.g. herbicidal) formulations generally contain from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of a compound of formula I, and preferably from 1 to 99.9 % by weight of a formulation adjuvant, which preferably includes from 0 to 25 % by weight of a surface-active substance. Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The rate of application of the compounds of formula I may vary within wide limits and depends upon the nature of the soil, the method of application (pre- or post-emergence; seed dressing; application to the seed furrow; no tillage application etc.), the crop plant, the weed or grass to be controlled, the prevailing climatic conditions, and other factors governed by the method of application, the time of application and the target crop. The compounds of formula I according to the invention are generally applied at a rate of 1 to 2000 g/ha, preferably 1 to 1000 g / ha, most preferably at 1 to 500 g / ha, and most preferably at 10 to 250 g / ha (in particular at 10, 15, 16, 20, 30, 50, 60, 62.5, 100, 125 or 250 g / ha).

Preferred formulations have especially the following representative compositions:
(% = percent by weight):

Emulsifiable concentrates:

active ingredient:	1 to 95 %, preferably 60 to 90 %
surface-active agents:	1 to 30 %, preferably 5 to 20 %
solvents as liquid carrier:	1 to 80 %, preferably 1 to 35 %

Dusts:

active ingredient:	0.1 to 10 %, preferably 0.1 to 5 %
solid carriers:	99.9 to 90 %, preferably 99.9 to 99 %

Suspension concentrates:

active ingredient:	5 to 75 %, preferably 10 to 50 %
water:	94 to 24 %, preferably 88 to 30 %
surface-active agents:	1 to 40 %, preferably 2 to 30 %

- 76 -

Wettable powders:

active ingredient:	0.5 to 90 %, preferably 1 to 80 %
surface-active agents:	0.5 to 20 %, preferably 1 to 15 %
solid carriers:	5 to 95 %, preferably 15 to 90 %

Granules:

active ingredient:	0.1 to 30 %, preferably 0.1 to 15 %
solid carriers:	99.5 to 70 %, preferably 97 to 85 %

Waterdispersible granules:

active ingredient:	1 to 90 %, preferably 10 to 80 %
surface-active agents:	0.5 to 80 %, preferably 5 to 30 %
solid carriers:	90 to 10 %, preferably 70 to 30 %

The following Examples further illustrate, but do not limit, the invention.

F1. Emulsifiable concentrates

	a)	b)	c)	d)
active ingredient	5 %	10 %	25 %	50 %
calcium dodecylbenzene-sulfonate	6 %	8 %	6 %	8 %
castor oil polyglycol ether (36 mol of ethylene oxide)	4 %	-	4 %	4 %
octylphenol polyglycol ether (7-8 mol of ethylene oxide)	-	4 %	-	2 %
NMP	-	10 %		20 %
arom. hydrocarbon mixture C ₉ -C ₁₂	85 %	68 %	65 %	16 %

Emulsions of any desired concentration can be prepared from such concentrates by dilution with water.

F2. Solutions

	a)	b)	c)	d)
active ingredient	5 %	10 %	50 %	90 %
1-methoxy-3-(3-methoxy-propoxy)-propane	40 %	50 %	-	
polyethylene glycol MW 400	20 %	10 %	-	-
NMP	-		50 %	10 %

- 77 -

arom. hydrocarbon	35 %	30 %	-	-
mixture C ₉ -C ₁₂				

The solutions are suitable for application undiluted or after dilution with water.

<u>F3. Wettable powders</u>	a)	b)	c)	d)
active ingredient	5 %	25 %	50 %	80 %
sodium lignosulfonate	4 %	-	3 %	-
sodium lauryl sulfate	2 %	3 %	-	4 %
sodium diisobutyl-naphthalene-sulfonate	-	6 %	5 %	6 %
octylphenol polyglycol ether (7-8 mol of ethylene oxide)	-	1 %	2 %	-
highly disperse silicic acid	1 %	3 %	5 %	10 %
kaolin	88 %	62 %	35 %	-

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, yielding wettable powders which can be diluted with water to give suspensions of any desired concentration.

<u>F4. Coated granules</u>	a)	b)	c)
active ingredient	0.1 %	5 %	15 %
highly dispersed silica	0.9 %	2 %	2 %
inorg. carrier	99.0 %	93 %	83 %
(diameter 0.1 - 1 mm)			
e.g. CaCO ₃ or SiO ₂			

The active ingredient is dissolved in methylene chloride, the solution is sprayed onto the carrier and the solvent is subsequently evaporated off *in vacuo*.

<u>F5. Coated granules</u>	a)	b)	c)
active ingredient	0.1 %	5 %	15 %
polyethylene glycol MW 200	1.0 %	2 %	3 %
highly dispersed silica	0.9 %	1 %	2 %
inorg. carrier	98.0 %	92 %	80 %
(diameter 0.1 - 1 mm)			
e.g. CaCO ₃ or SiO ₂			

- 78 -

The finely ground active ingredient is applied uniformly, in a mixer, to the carrier moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

<u>F6. Extruded granules</u>	a)	b)	c)	d)
active ingredient	0.1 %	3 %	5 %	15 %
sodium lignosulfonate	1.5 %	2 %	3 %	4 %
carboxymethylcellulose	1.4 %	2 %	2 %	2 %
kaolin	97.0 %	93 %	90 %	79 %

The active ingredient is mixed and ground with the adjuvants and the mixture is moistened with water. The resulting mixture is extruded and then dried in a stream of air.

<u>F7. Water-dispersible granules</u>	a)	b)	c)	d)
active ingredient	5 %	10 %	40 %	90 %
sodium lignosulfonate	20 %	20 %	15 %	7 %
dibutyl naphthalene sulfonate	5 %	5 %	4 %	2 %
Gum arabic	2 %	1 %	1 %	1 %
Diatomaceous earth	20 %	30 %	5 %	
Sodium sulphate		4 %	5 %	
kaolin	48 %	30 %	30 %	

The active ingredient is mixed and ground with the adjuvants and the mixture is moistened with water. The resulting mixture is extruded and then dried in a stream of air.

<u>F8. Dusts</u>	a)	b)	c)
active ingredient	0.1 %	1 %	5 %
talcum	39.9 %	49 %	35 %
kaolin	60.0 %	50 %	60 %

Ready-to-use dusts are obtained by mixing the active ingredient with the carriers and grinding the mixture in a suitable mill.

<u>F9. Suspension concentrates</u>	a)	b)	c)	d)
active ingredient	3 %	10 %	25 %	50 %
propylene glycol	5 %	5 %	5 %	5 %
nonylphenol polyglycol ether (15 mol of ethylene oxide)	-	1 %	2 %	-
sodium lignosulfonate	3 %	3 %	7 %	6 %

- 79 -

heteropolysaccharide (Xanthan)	0.2 %	0.2 %	0.2 %	0.2 %
1,2-Benzisothiazolin-3-on	0.1 %	0.1 %	0.1 %	0.1 %
silicone oil emulsion	0.7 %	0.7 %	0.7 %	0.7 %
water	87 %	79 %	62 %	38 %

The finely ground active ingredient is intimately mixed with the adjuvants, yielding a suspension concentrate from which suspensions of any desired concentration can be prepared by dilution with water.

The invention also provides a method of controlling grasses and weeds in crops of useful plants, which comprises applying a herbicidally effective amount of a compound of formula I as defined herein, or of a composition comprising such a compound, to the plants or to the locus thereof.

Crops of useful plants, in which the compositions and/or the methods of controlling grasses and weeds according to the invention can be used, are typically cereals (in particular wheat, barley, rye or triticale; preferably wheat or barley), rice, corn (i.e. maize), rape, sugarbeet, sugarcane, soybean, cotton, sunflower, peanut, or plantation crops. Alternatively, the crops of useful plants can be oats (e.g. *Avena sativa*, the common oat). The crops of useful plants are preferably cereals (e.g. wheat, barley, rye or triticale), rice, corn or soybean; or more preferably are wheat, barley, rice, corn or soybean; or most preferably are rice.

The term "crops" is to be understood as also including crops that have been rendered tolerant to herbicides or classes of herbicides (for example ALS, GS, EPSPS, PPO and HPPD inhibitors) as a result of conventional methods of breeding or genetic engineering. An example of a crop that has been rendered tolerant e.g. to imidazolinones, such as imazamox, by conventional methods of breeding is Clearfield® summer rape (Canola). Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady® and LibertyLink®.

The weeds to be controlled may be monocotyledonous and/or dicotyledonous weeds, such as, for example, *Stellaria*, *Nasturtium*, *Agrostis*, *Digitaria* (e.g. *Digitaria sanguinalis* (DIGSA)), *Avena* (e.g. *Avena* species other than *Avena sativa* (the common oat); preferably *Avena fatua* (AVEFA), also known as the common wild oat), *Setaria* (e.g. *Setaria faberi* (SETFA)), *Sinapis*, *Lolium* (e.g. *Lolium perenne* (LOLPE)), *Solanum*, *Echinochloa* (e.g. *Echinochloa crus-galli* (ECHCG)), *Scirpus*, *Monochoria*, *Sagittaria*, *Bromus*, *Alopecurus* (e.g. *Alopecurus*

- 80 -

myosuroides (ALOMY)), Sorghum, Rottboellia, Cyperus, Abutilon, Sida, Xanthium, Amaranthus, Chenopodium, Ipomoea, Chrysanthemum, Galium, Viola and/or Veronica. The weeds to be controlled may alternatively be Phalaris, Apera, Leptochloa, Geranium, Beta, Brassica, Kochia, Poa, Sinapis, Polygonum, Brachiaria, Eriochloa, Bidens, Euphorbia, and/or Panicum.

Control of monocotyledonous weeds (e.g. weedy grasses) is preferred; in particular Agrostis, Avena (e.g. Avena species other than *Avena sativa* (the common oat); preferably *Avena fatua* (AVEFA), also known as the common wild oat), Setaria (e.g. *Setaria faberi* (SETFA)), Lolium (e.g. *Lolium perenne* (LOLPE)), Echinochloa (e.g. *Echinochloa crus-galli* (ECHCG)), Bromus, Alopecurus (e.g. *Alopecurus myosuroides* (ALOMY)), and/or Sorghum.

Alternatively, the monocotyledonous weeds to be controlled are, in particular, Phalaris, Apera, Panicum, Digitaria, Brachiaria, Poa, Eriochloa, Rottboellia, and/or Leptochloa; and/or can be volunteer (non-crop) cereals and/or volunteer (non-crop) maize. The monocotyledonous weeds, to be controlled by compound of formula I, can be either sensitive to or partially or wholly resistant to one or more herbicides, not being a compound of formula I, which are already approved and commercially-available for herbicidal use (and/or which are already used in agriculture as herbicides).

Crops are also to be understood as being those which have been rendered resistant to harmful insects by genetic engineering methods, for example Bt maize (resistant to European corn borer), Bt cotton (resistant to cotton boll weevil) and also Bt potatoes (resistant to Colorado beetle). Examples of Bt maize are the Bt-176 maize hybrids of NK® (Syngenta Seeds). The Bt toxin is a protein that is formed naturally by *Bacillus thuringiensis* soil bacteria. Examples of toxins and transgenic plants able to synthesise such toxins are described in EP-A-451 878, EP-A-374 753, WO 93/07278, WO 95/34656, WO 03/052073 and EP-A-427 529. Examples of transgenic plants that contain one or more genes which code for an insecticidal resistance and express one or more toxins are KnockOut® (maize), Yield Gard® (maize), NuCOTIN33B® (cotton), Bollgard® (cotton), NewLeaf® (potatoes), NatureGard® and Protexcta®. Plant crops and their seed material can be resistant to herbicides and at the same time also to insect feeding ("stacked" transgenic events). Seed can, for example, have the ability to express an insecticidally active Cry3 protein and at the same time be glyphosate-tolerant. The term "crops" is to be understood as also including crops obtained as a result of conventional methods of breeding or genetic engineering which contain so-called output traits (e.g. improved flavour, storage stability, nutritional content).

Areas under cultivation are to be understood as including land where the crop plants are already growing as well as land intended for the cultivation of those crop plants.

The compounds of formula I according to the invention can also be used in combination with further herbicides. Preferably, in these mixtures, the compound of the formula I is one of those compounds listed in Tables 1 to 54 and/or in Table A1 hereinbelow. The following mixtures of the compound of formula I may be important:

compound of formula I + acetochlor, compound of formula I + acifluorfen, compound of formula I + acifluorfen-sodium, compound of formula I + aclonifen, compound of formula I + acrolein, compound of formula I + alachlor, compound of formula I + alloxymid, compound of formula I + allyl alcohol, compound of formula I + ametryn, compound of formula I + amicarbazone, compound of formula I + amidosulfuron, compound of formula I + aminopyralid, compound of formula I + amitrole, compound of formula I + ammonium sulfamate, compound of formula I + anilofos, compound of formula I + asulam, compound of formula I + atraton, compound of formula I + atrazine, compound of formula I + azimsulfuron, compound of formula I + BCPC, compound of formula I + beflubutamid, compound of formula I + benazolin, compound of formula I + benfluralin, compound of formula I + benfuresate, compound of formula I + bensulfuron, compound of formula I + bensulfuron-methyl, compound of formula I + bensulide, compound of formula I + bentazone, compound of formula I + benzfendizone, compound of formula I + benzobicyclon, compound of formula I + benzofenap, compound of formula I + bifenox, compound of formula I + bilanafos, compound of formula I + bispyribac, compound of formula I + bispyribac-sodium, compound of formula I + borax, compound of formula I + bromacil, compound of formula I + bromobutide, compound of formula I + bromoxynil, compound of formula I + butachlor, compound of formula I + butafenacil, compound of formula I + butamifos, compound of formula I + butralin, compound of formula I + butroxydim, compound of formula I + butylate, compound of formula I + cacodylic acid, compound of formula I + calcium chlorate, compound of formula I + cafenstrole, compound of formula I + carbetamide, compound of formula I + carfentrazone, compound of formula I + carfentrazone-ethyl, compound of formula I + CDEA, compound of formula I + CEPC, compound of formula I + chlorflurenol, compound of formula I + chlorflurenol-methyl, compound of formula I + chloridazon, compound of formula I + chlorimuron, compound of formula I + chlorimuron-ethyl, compound of formula I + chloroacetic acid, compound of formula I + chlorotoluron, compound of formula I + chlorpropham, compound of formula I + chlorsulfuron, compound of formula I + chlorthal,

- 82 -

compound of formula I + chlorthal-dimethyl, compound of formula I + cinidon-ethyl, compound of formula I + cinmethylin, compound of formula I + cinosulfuron, compound of formula I + cisanilide, compound of formula I + clethodim, compound of formula I + clodinafop, compound of formula I + clodinafop-propargyl, compound of formula I + clomazone, compound of formula I + clomeprop, compound of formula I + clopyralid, compound of formula I + cloransulam, compound of formula I + cloransulam-methyl, compound of formula I + CMA, compound of formula I + 4-CPB, compound of formula I + CPMF, compound of formula I + 4-CPP, compound of formula I + CPPC, compound of formula I + cresol, compound of formula I + cumyluron, compound of formula I + cyanamide, compound of formula I + cyanazine, compound of formula I + cycloate, compound of formula I + cyclosulfamuron, compound of formula I + cycloxydim, compound of formula I + cyhalofop, compound of formula I + cyhalofop-butyl, compound of formula I + 2,4-D, compound of formula I + 3,4-DA, compound of formula I + daimuron, compound of formula I + dalapon, compound of formula I + dazomet, compound of formula I + 2,4-DB, compound of formula I + 3,4-DB, compound of formula I + 2,4-DEB, compound of formula I + desmedipham, compound of formula I + dicamba, compound of formula I + dichlobenil, compound of formula I + ortho-dichlorobenzene, compound of formula I + para-dichlorobenzene, compound of formula I + dichlorprop, compound of formula I + dichlorprop-P, compound of formula I + diclofop, compound of formula I + diclofop-methyl, compound of formula I + diclosulam, compound of formula I + difenzoquat, compound of formula I + difenzoquat metilsulfate, compound of formula I + diflufenican, compound of formula I + diflufenzopyr, compound of formula I + dimefuron, compound of formula I + dimepiperate, compound of formula I + dimethachlor, compound of formula I + dimethametryn, compound of formula I + dimethenamid, compound of formula I + dimethenamid-P, compound of formula I + dimethipin, compound of formula I + dimethylarsinic acid, compound of formula I + dinitramine, compound of formula I + dinoterb, compound of formula I + diphenamid, compound of formula I + diquat, compound of formula I + diquat dibromide, compound of formula I + dithiopyr, compound of formula I + diuron, compound of formula I + DNOC, compound of formula I + 3,4-DP, compound of formula I + DSMA, compound of formula I + EBEP, compound of formula I + endothal, compound of formula I + EPTC, compound of formula I + esprocarb, compound of formula I + ethalfluralin, compound of formula I + ethametsulfuron, compound of formula I + ethametsulfuron-methyl, compound of formula I + ethofumesate, compound of formula I + ethoxyfen, compound of formula I + ethoxysulfuron, compound of formula I + etobenzanid, compound of formula I + fenoxaprop-P, compound of formula I + fenoxaprop-P-ethyl, compound of formula I + fentrazamide, compound of formula

I + ferrous sulfate, compound of formula I + flamprop-M, compound of formula I + flazasulfuron, compound of formula I + florasulam, compound of formula I + fluazifop, compound of formula I + fluazifop-butyl, compound of formula I + fluazifop-P, compound of formula I + fluazifop-P-butyl, compound of formula I + flucarbazone, compound of formula I + flucarbazone-sodium, compound of formula I + flucetosulfuron, compound of formula I + fluchloralin, compound of formula I + flufenacet, compound of formula I + flufenpyr, compound of formula I + flufenpyr-ethyl, compound of formula I + flumetsulam, compound of formula I + flumiclorac, compound of formula I + flumiclorac-pentyl, compound of formula I + flumioxazin, compound of formula I + fluometuron, compound of formula I + fluoroglycofen, compound of formula I + fluoroglycofen-ethyl, compound of formula I + flupropanate, compound of formula I + flupyrsulfuron, compound of formula I + flupyrsulfuron-methyl-sodium, compound of formula I + flurenol, compound of formula I + fluridone, compound of formula I + flurochloridone, compound of formula I + fluroxypyr, compound of formula I + flurtamone, compound of formula I + fluthiacet, compound of formula I + fluthiacet-methyl, compound of formula I + fomesafen, compound of formula I + foramsulfuron, compound of formula I + fosamine, compound of formula I + glufosinate, compound of formula I + glufosinate-ammonium, compound of formula I + glyphosate, compound of formula I + halosulfuron, compound of formula I + halosulfuron-methyl, compound of formula I + haloxyfop, compound of formula I + haloxyfop-P, compound of formula I + HC-252, compound of formula I + hexazinone, compound of formula I + imazamethabenz, compound of formula I + imazamethabenz-methyl, compound of formula I + imazamox, compound of formula I + imazapic, compound of formula I + imazapyr, compound of formula I + imazaquin, compound of formula I + imazethapyr, compound of formula I + imazosulfuron, compound of formula I + indanofan, compound of formula I + iodomethane, compound of formula I + iodosulfuron, compound of formula I + iodosulfuron-methyl-sodium, compound of formula I + ioxynil, compound of formula I + isoproturon, compound of formula I + isouron, compound of formula I + isoxaben, compound of formula I + isoxachlortole, compound of formula I + isoxaflutole, compound of formula I + karbutilate, compound of formula I + lactofen, compound of formula I + lenacil, compound of formula I + linuron, compound of formula I + MAA, compound of formula I + MAMA, compound of formula I + MCPA, compound of formula I + MCPA-thioethyl, compound of formula I + MCPB, compound of formula I + mecoprop, compound of formula I + mecoprop-P, compound of formula I + mefenacet, compound of formula I + mefluidide, compound of formula I + mesosulfuron, compound of formula I + mesosulfuron-methyl, compound of formula I + mesotrione, compound of formula I + metam, compound of formula I + metamifop, compound of formula I + metamitron,

- 84 -

compound of formula I + metazachlor, compound of formula I + methabenzthiazuron, compound of formula I + methylarsonic acid, compound of formula I + methylmymron, compound of formula I + methyl isothiocyanate, compound of formula I + metobenzuron, compound of formula I + metolachlor, compound of formula I + S-metolachlor, compound of formula I + metosulam, compound of formula I + metoxuron, compound of formula I + metribuzin, compound of formula I + metsulfuron, compound of formula I + metsulfuron-methyl, compound of formula I + MK-616, compound of formula I + molinate, compound of formula I + monolinuron, compound of formula I + MSMA, compound of formula I + naproanilide, compound of formula I + napropamide, compound of formula I + naptalam, compound of formula I + neburon, compound of formula I + nicosulfuron, compound of formula I + nonanoic acid, compound of formula I + norflurazon, compound of formula I + oleic acid (fatty acids), compound of formula I + orbencarb, compound of formula I + orthosulfamuron, compound of formula I + oryzalin, compound of formula I + oxadiargyl, compound of formula I + oxadiazon, compound of formula I + oxasulfuron, compound of formula I + oxaziclomefone, compound of formula I + oxyfluorfen, compound of formula I + paraquat, compound of formula I + paraquat dichloride, compound of formula I + pebulate, compound of formula I + pendimethalin, compound of formula I + penoxsulam, compound of formula I + pentachlorophenol, compound of formula I + pentanochlor, compound of formula I + pentoxazone, compound of formula I + pethoxamid, compound of formula I + petroleum oils, compound of formula I + phenmedipham, compound of formula I + phenmedipham-ethyl, compound of formula I + picloram, compound of formula I + picolinafen, compound of formula I + pinoxaden, compound of formula I + piperophos, compound of formula I + potassium arsenite, compound of formula I + potassium azide, compound of formula I + pretilachlor, compound of formula I + primisulfuron, compound of formula I + primisulfuron-methyl, compound of formula I + prodiamine, compound of formula I + profluazol, compound of formula I + profoxydim, compound of formula I + prometon, compound of formula I + prometryn, compound of formula I + propachlor, compound of formula I + propanil, compound of formula I + propaquizafop, compound of formula I + propazine, compound of formula I + propham, compound of formula I + propisochlor, compound of formula I + propoxycarbazone, compound of formula I + propoxycarbazone-sodium, compound of formula I + propyzamide, compound of formula I + prosulfocarb, compound of formula I + prosulfuron, compound of formula I + pyraclonil, compound of formula I + pyraflufen, compound of formula I + pyraflufen-ethyl, compound of formula I + pyrazolynate, compound of formula I + pyrazosulfuron, compound of formula I + pyrazosulfuron-ethyl, compound of formula I + pyrazoxyfen, compound of formula I + pyribenzoxim, compound of formula I +

pyributicarb, compound of formula I + pyridafol, compound of formula I + pyridate, compound of formula I + pyriftalid, compound of formula I + pyriminobac, compound of formula I + pyriminobac-methyl, compound of formula I + pyrimisulfan, compound of formula I + pyrithiobac, compound of formula I + pyrithiobac-sodium, compound of formula I + quinclorac, compound of formula I + quinmerac, compound of formula I + quinoctamine, compound of formula I + quizalofop, compound of formula I + quizalofop-P, compound of formula I + rimsulfuron, compound of formula I + sethoxydim, compound of formula I + siduron, compound of formula I + simazine, compound of formula I + simetryn, compound of formula I + SMA, compound of formula I + sodium arsenite, compound of formula I + sodium azide, compound of formula I + sodium chlorate, compound of formula I + sulcotrione, compound of formula I + sulfentrazone, compound of formula I + sulfometuron, compound of formula I + sulfometuron-methyl, compound of formula I + sulfosate, compound of formula I + sulfosulfuron, compound of formula I + sulfuric acid, compound of formula I + tar oils, compound of formula I + 2,3,6-TBA, compound of formula I + TCA, compound of formula I + TCA-sodium, compound of formula I + tebuthiuron, compound of formula I + tepraloxym, compound of formula I + terbacil, compound of formula I + terbumeton, compound of formula I + terbuthylazine, compound of formula I + terbutryn, compound of formula I + thenylchlor, compound of formula I + thiazopyr, compound of formula I + thifensulfuron, compound of formula I + thifensulfuron-methyl, compound of formula I + thiobencarb, compound of formula I + tiocarbazil, compound of formula I + topramezone, compound of formula I + tralkoxydim, compound of formula I + tri-allate, compound of formula I + triasulfuron, compound of formula I + triaziflam, compound of formula I + tribenuron, compound of formula I + tribenuron-methyl, compound of formula I + tricamba, compound of formula I + triclopyr, compound of formula I + trietazine, compound of formula I + trifloxysulfuron, compound of formula I + trifloxysulfuron-sodium, compound of formula I + trifluralin, compound of formula I + triflusulfuron, compound of formula I + triflusulfuron-methyl, compound of formula I + trihydroxytriazine, compound of formula I + tritosulfuron, compound of formula I + [3-[2-chloro-4-fluoro-5-(1-methyl-6-trifluoromethyl-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-3-yl)phenoxy]-2-pyridyloxy]acetic acid ethyl ester (CAS RN 353292-31-6), compound of formula I + 4-[(4,5-dihydro-3-methoxy-4-methyl-5-oxo)-1H-1,2,4-triazol-1-yl]carbonylsulfamoyl]-5-methylthiophene-3-carboxylic acid (BAY636), compound of formula I + BAY747 (CAS RN 335104-84-2), compound of formula I + topramezone (CAS RN 210631-68-8), compound of formula I + 4-hydroxy-3-[[2-[(2-methoxyethoxy)methyl]-6-(trifluoromethyl)-3-pyridinyl]carbonyl]-bicyclo[3.2.1]oct-3-en-2-one (CAS RN 352010-68-5), and

- 86 -

compound of formula I + 4-hydroxy-3-[[2-(3-methoxypropyl)-6-(difluoromethyl)-3-pyridinyl]carbonyl]-bicyclo[3.2.1]oct-3-en-2-one.

The following alternative mixtures of the compound of formula I may be important (preferably, in these mixtures, the compound of the formula I is one of those compounds listed in Tables 1 to 54 and/or in Table A1 hereinbelow):

compound of formula I + one of the herbicidal compounds disclosed in WO2010/059676 (Dow, e.g. for use with cereal crops, e.g. can be plus cloquintocet-mexyl),

compound of formula I + one of the herbicidal compounds disclosed in WO2010/059680 (Dow, e.g. for use with cereal crops, e.g. can be plus a safener other than cloquintocet-mexyl), and

compound of formula I + one of the herbicidal compounds disclosed in WO2010/059671 (Dow, e.g. for use with rice crops, e.g. can be plus a safener).

The mixing partners for the compound of formula I may also be in the form of esters or salts, as mentioned e.g. in The Pesticide Manual, 12th Edition (BCPC) 2000.

The compounds of formula I according to the invention can also be used in combination with safeners. Preferably, in these mixtures, the compound of the formula I is one of those compounds listed in Tables 1 to 54 below. The following mixtures with safeners, especially, come into consideration:

compound of formula I + cloquintocet-mexyl, compound of formula I + cloquintocet acid and salts thereof, compound of formula I + fenclorazole-ethyl, compound of formula I + fenclorazole acid and salts thereof, compound of formula I + mefenpyr-diethyl, compound of formula I + mefenpyr diacid, compound of formula I + isoxadifen-ethyl, compound of formula I + isoxadifen acid, compound of formula I + furilazole, compound of formula I + furilazole R isomer, compound of formula (I) + N-(2-methoxybenzoyl)-4-[[methylaminocarbonyl]amino]benzenesulfonamide, compound of formula I + benoxacor, compound of formula I + dichlormid, compound of formula I + AD-67, compound of formula I + oxabetrinil, compound of formula I + cyometrinil, compound of formula I + cyometrinil Z-isomer, compound of formula I + fenclorim, compound of formula I + cyprosulfamide, compound of formula I + naphthalic anhydride, compound of formula I + flurazole, compound of formula I + CL 304,415, compound of formula I + dicyclonon, compound of formula I + fluxofenim, compound of formula I + DKA-24, compound of formula I + R-29148 and compound of formula I + PPG-1292. A safening effect can also be observed for the mixtures

- 87 -

compound of the formula I + dymron, compound of the formula I + MCPA, compound of the formula I + mecoprop and compound of the formula I + mecoprop-P.

The above-mentioned safeners and herbicides are described, for example, in the Pesticide Manual, Twelfth Edition, British Crop Protection Council, 2000. R-29148 is described, for example by P.B. Goldsbrough *et al.*, Plant Physiology, (2002), Vol. 130 pp. 1497-1505 and references therein, PPG-1292 is known from WO09211761 and N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide is known from EP365484.

Benoxacor, cloquintocet-mexyl, cyprosulfamide, mefenpyr-diethyl and N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide are especially preferred safeners. Cloquintocet-mexyl is particularly valuable and is the most preferred safener.

Therefore, the invention also provides a herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined herein, and optionally (or preferably) a further herbicide as mixture partner for the compound of formula I, or optionally (or preferably) a safener, or both.

The invention also provides a herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined herein, a safener, and optionally (or preferably) a further herbicide as mixture partner for the compound of formula I,

wherein the safener is benoxacor, cloquintocet-mexyl, cyprosulfamide, mefenpyr-diethyl or N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide.

The rate of application of safener, relative to the herbicide (e.g. the compound of formula I), is largely dependent upon the mode of application. In the case of field treatment:

a) generally from 0.001 to 5.0 kg of safener / ha, preferably from 0.001 to 0.5 kg of safener / ha, more preferably 2 to 100 g of safener / ha (in particular at 2.5, 5, 7.5, 10, 20 or 50 g of safener / ha); and

b) generally from 0.001 to 2 kg of herbicide / ha, but preferably from 0.005 to 1 kg of herbicide / ha, more preferably 5 to 500 g of herbicide / ha, and most preferably at 10 to 250 g of herbicide / ha (in particular at 10, 15, 16, 20, 30, 50, 60, 62.5, 100, 125 or 250 g of herbicide / ha);

are applied.

The safener and the herbicide (e.g. the compound of formula I) can for example be used (e.g. when together in a mixture formulation) at a herbicide : safener ratio of from 16:1 to 1:1, such as 8:1, 4:1 or 2:1, measured on the basis of the rates of application of herbicide and safener in g / ha; in particular where the safener is cloquintocet-mexyl.

The herbicidal compositions according to the invention are suitable for all methods of application customary in agriculture, such as, for example, pre-emergence application, post-emergence application and seed dressing. Depending upon the intended use, the safeners can be used for pretreating the seed material of the crop plant (dressing the seed or seedlings) or introduced into the soil before or after sowing, followed by the application of the (unsafened) compound of the formula (I), optionally in combination with a co-herbicide. It can, however, also be applied alone or together with the herbicide before or after emergence of the plants. The treatment of the plants or the seed material with the safener can therefore take place in principle independently of the time of application of the herbicide. The treatment of the plant by simultaneous application of herbicide and safener (e.g. in the form of a tank mixture) is generally preferred. The rate of application of safener relative to herbicide is largely dependent upon the mode of application. In the case of field treatment, generally from 0.001 to 5.0 kg of safener/ha, preferably from 0.001 to 0.5 kg of safener/ha, are applied. In the case of seed dressing, generally from 0.001 to 10 g of safener/kg of seed, preferably from 0.05 to 2 g of safener/kg of seed, are applied. When the safener is applied in liquid form, with seed soaking, shortly before sowing, it is advantageous to use safener solutions which contain the active ingredient in a concentration of from 1 to 10 000 ppm, preferably from 100 to 1000 ppm.

It is preferred to apply the mixture partner of the compound of formula I together with one of the safeners mentioned above.

The following Examples illustrate the invention further but do not limit the invention.


Preparation Examples:

Those skilled in the art will appreciate that certain compounds described below are β -ketoenols, and as such may exist as a single tautomer or as a mixture of keto-enol and diketone tautomers, as described, for example by J. March, Advanced Organic Chemistry, third edition, John Wiley and Sons. The compounds shown below, and in Table A1 are drawn as an arbitrary single enol tautomer, but it should be inferred that this description covers both the diketone form and any possible enols which could arise through tautomerism. Where

more than one tautomer is observed in proton NMR, the data shown are for the mixture of tautomers. Furthermore, some of the compounds shown below are drawn as single enantiomers for the purposes of simplicity, but unless specified as single enantiomers, these structures should be construed as representing a mixture of enantiomers. Additionally, some of the compounds can exist as diastereoisomers, and it should be inferred that these can be present as a mixture of diastereoisomers or as any possible single diastereoisomer.


Example 1:

Preparation of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione



Step 1:


Preparation of 5-bromo-4-oxo-pentanoic acid methyl ester



30 ml of trimethylsilyldiazomethane (2 M solution in diethyl ether) was added drop wise to a solution of 3-carbomethoxy propionylchloride (3.75 g, 30.5 mmol) in acetonitrile (75 ml) over a period of 10 mins at room temperature. The mixture was then stirred for 1 h. The reaction was cooled to 0 °C and 10 g of 33% HBr in acetic acid was added slowly over 20 min, then the reaction was warmed to room temperature and stirred for 16 h. The solvent was removed *in vacuo* without heating and the residue was dissolved in ethyl acetate (100 ml), washed with 50 ml of saturated sodium bicarbonate solution and then 50 ml of brine. The organics were passed through a phase separator to remove any water and the solvent removed *in vacuo*. The resultant crude material was purified by kuglerhor distillation under reduced pressure (~3 mbar, 150 °C) to give 5-bromo-4-oxo-pentanoic acid methyl ester (7 g).

Step 2:


Preparation of 5-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-oxo-pentanoic acid methyl ester



3-(4-Chloro-phenyl)-5-methyl-1H-pyrazole (1.84 g, 9.6 mmol) was dissolved in dimethylformamide (20 ml) followed by the addition of sodium hydride (60% dispersion in oil, 384 mg, 9.6 mmol). The reaction was stirred at room temperature for 15 min, followed by the addition of 5-bromo-4-oxo-pentanoic acid methyl ester (2 g, 9.6 mmol). The reaction was stirred at room temperature for a further 30 min, before the addition of water (200 ml). The solution was then extracted with ethyl acetate (2 x 150 ml). The organics were dried over magnesium sulfate and concentrated *in vacuo*. The crude product was then purified by column chromatography (120 g column, DCM to 90% DCM/EtOAc over 50 mins) to give 5-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-oxo-pentanoic acid methyl ester (1.84 g).

Step 3:

Preparation of 2-[3-(4-Chloro-phenyl)-5-methyl-pyrazol-1-yl]-cyclopentane-1,3-dione




5-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-oxo-pentanoic acid methyl ester (1.7 g, 5.5 mmol) was dissolved in dimethylformamide (16 ml) and anhydrous potassium carbonate (3 g, 22 mmol) added. The mixture was then heated to 200 °C for 10 mins in the microwave (using "normal" absorbion). The reaction was reduced *in vacuo*, and partitioned between 2 M hydrochloric acid (50 ml) and dichloromethane (100 ml). The layers were separated, and the aqueous layer saturated with sodium chloride and extracted with dichloromethane (100 ml). The organics were combined, dried over magnesium sulfate and the solvent removed *in vacuo*. The reaction was then purified by column chromatography (using DCM:EtOAc:AcOH

- 91 -

(9:1:0.01%) to (3:7:0.01%)) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-cyclopentane-1,3-dione (0.8 g).

Step 4:


Preparation of 2-[3-(4-Chloro-phenyl)-5-methyl-pyrazol-1-yl]-3-methoxy-cyclopent-2-enone



Neat dimethyl sulfate (0.12 ml, 1.3 mmol) was added at room temperature to a vigorously stirred suspension of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-cyclopentane-1,3-dione (374 mg, 1.3 mmol) and potassium carbonate (180 mg, 1.3 mmol) in acetone (10 ml). The mixture was then heated to 60 °C for 3 h with continued stirring, then cooled and the solvent removed *in vacuo*. The residue was partitioned between ethyl acetate (30 ml) and 1M sodium hydroxide solution (20 ml), the layers were separated, and the aqueous phase was further extracted with ethyl acetate (2 x 20 ml). The combined organics were washed with brine (20 ml), dried over magnesium sulfate and concentrated. The crude product was then purified by flash chromatography (using 20% to 100% EtOAc in DCM) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-3-methoxy-cyclopent-2-enone (253.2 mg).

Step 5:

Preparation of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione




A solution of lithium diisopropylamide (1.8 M in tetrahydrofuran/heptane/ethylbenzene, 0.25 ml, 0.45 mmol) was added drop wise to a solution of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-3-methoxy-cyclopent-2-enone (124 mg, 0.41 mmol) in anhydrous tetrahydrofuran (1 ml) at -78 °C, under nitrogen, and allowed to stir at this temperature for 30 minutes.

- 92 -

Tetrahydropyranyl-4-carbaldehyde (51.5 mg, 0.45 mmol) was added drop wise and the reaction stirred at -78 °C for a further 30 minutes. The reaction was allowed to warm to room temperature over 30 mins and quenched by the addition of 1 M hydrochloric acid (5 ml) and extracted with dichlormethane (3 x 10 ml) to give 223.5 mg of crude material. The crude material was dissolved in acetone (2 ml) and 2 M hydrochloric acid (2 ml) added. This solution was heated in the microwave (using "normal" absorbtion) for 1 h at 120 °C. This solution was extracted with dichloromethane (2 x 10 ml) and the organics dried over magnesium sulfate and reduced *in vacuo*. The crude material was purified by reverse phase preparative HPLC (eluting with acetonitrile/water) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (32.2 mg).


Preparation of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione



To a solution of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (9 mg, 0.02 mmol) in methanol (0.1 ml) was added 5% palladium on activated charcoal (1 mg) and the reaction stirred for 6 hours under 1.5 bar of hydrogen. The solution was then filtered through a pad of Celite, washed with methanol (10 ml) and concentrated *in vacuo* to give a white solid, which was purified by reverse phase preparative HPLC (eluting with acetonitrile/water) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione (4 mg).

Preparation of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(Z)-ylidene]-cyclopentane-1,3-dione and 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione


- 93 -



A solution of lithium diisopropylamide (1.8 M in tetrahydrofuran/heptane/ethylbenzene, 0.25 ml, 0.45 mmol) was added drop wise to a solution of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-3-methoxy-cyclopent-2-enone (124 mg, 0.41 mmol) in anhydrous tetrahydrofuran (1 ml) at -78 °C, under nitrogen, and allowed to stir at this temperature for 30 minutes. 5-Fluoro-pyridine-2-carbaldehyde (56 mg, 0.45 mmol) was then added drop wise and the reaction stirred at -78 °C for a further 30 minutes. The reaction was then allowed to warm to room temperature over 30 mins and quenched by the addition of 1 M hydrochloric acid (5 ml) and extracted with dichloromethane (3 x 10 ml) to give 186 mg of crude material. The crude material was dissolved in acetone (2 ml) and 2 M hydrochloric acid (2 ml) added. This solution was heated in the microwave (using "normal" absorption) for 1 h at 120 °C. This solution was extracted with dichloromethane (2 x 10 ml) and the organics dried over magnesium sulfate and reduced *in vacuo*. The crude material was purified by reverse phase preparative HPLC (eluting with acetonitrile/water) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(Z)-ylidene]-cyclopentane-1,3-dione (22 mg) and 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (5 mg).


- 94 -

Preparation of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-(5-fluoro-pyridin-2-ylmethyl)-cyclopentane-1,3-dione




To a solution of 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(Z)-ylidene]-cyclopentane-1,3-dione (17 mg, 0.04 mmol) in methanol (0.4 ml) was added 5% palladium on activated charcoal (2 mg) and the reaction stirred for 6 hours under 1.5 bar of hydrogen. The solution was then filtered through a pad of Celite, washed with methanol (10 ml) and concentrated *in vacuo* to give a white solid, which was purified by column chromatography using DCM:EtOAc:AcOH (9:1:0.01%) to (3:7:0.01%) to give 2-[3-(4-chloro-phenyl)-5-methyl-pyrazol-1-yl]-4-(5-fluoro-pyridin-2-ylmethyl)-cyclopentane-1,3-dione (15.7 mg).

Preparation of 2-(1,5-Dimethyl-1H-indazol-4-yl)-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione



Step 1:

Preparation of 5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone




A solution of 3-methoxy-cyclopent-2-enone (18.8 g, 0.17 mol) in THF (200 ml) was added drop wise over 30 minutes (using an addition funnel) to a solution of lithium diisopropylamide solution (1.8 M in tetrahydrofuran/heptane/ethylbenzene, 103 ml, 0.18 mol) in THF (150 ml) at -78 °C, under nitrogen and allowed to stir at this temperature for 30 minutes.

- 95 -

Tetrahydropyranyl-4-carbaldehyde (21 g, 0.18 mol) was then added drop wise and the reaction stirred at -78 °C for 30 minutes, before being allowed to warm to room temperature overnight (18 h). The reaction was then quenched by the addition of 1 M hydrochloric acid (500 ml) and extracted with ethyl acetate (3 x 500 ml). The aqueous layer was then saturated with sodium chloride and extracted with ethyl acetate (2 x 250 ml). The combined organics were evaporated and the crude product was purified using column chromatography (EtOAc: hexane, 3:7 to 100% EtOAc) to give 5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (20.8 g).


Step 2:

Preparation of 3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-methylidene]-cyclopent-2-enone




5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (1.1 g, 5 mmol) was dissolved in dichloromethane (25 ml) in a 50 ml round bottom flask fitted with a reflux condenser. Triethylamine (1.4 ml, 10 mmol) and then methane sulfonyl chloride (0.8 ml, 10 mmol) was added dropwise which caused the reaction to warm to reflux. After 10 minutes the reaction was quenched by the addition of 1 M hydrochloric acid (10 ml). The layers were separated and the organics collected, passed through a phase separation cartridge and the solvent removed *in vacuo*. The crude product was dissolved in methanol (25 ml) and anhydrous potassium carbonate (1.4 g, 10 mmol) added and the mixture was stirred at room temperature for 72 h. The solvent was then removed *in vacuo* and the residue dissolved in ethyl acetate (25 ml) and filtered to remove any solid. The filtrate was evaporated to dryness and the residue purified on by column chromatography (Hexane/EtOAc (7:3) to 100% EtOAc) to give 3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-methylidene]-cyclopent-2-enone (938 mg).

- 96 -

Step 3:Preparation of 3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone


A solution of 3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-methylidene]-cyclopent-2-enone (938 mg, 4.5 mmol) in ethanol (45 ml) was added to 5% palladium on activated charcoal (94 mg) and the reaction stirred for 7.5 hours under 3 bar of hydrogen. The reaction was then filtered through a pad of Celite, washed with ethanol and concentrated *in vacuo* to give 3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone (906 mg).

Step 4:Preparation of 2-bromo-3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone

To a stirred solution of 3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone (906 mg, 4.7 mmol) in 1,2-dichloroethane (15 ml) at 0 °C in an amber flask was added portion wise over one hour *N*-bromosuccinimide (890 mg, 5 mmol). The reaction was stirred at 0 °C for a further 90 minutes and then any remaining solid was removed by filtration. The filtrate was evaporated to dryness under reduced pressure, the resultant solid dissolved in warm toluene (50 ml) and washed quickly with ice-cold water (2 x 20 ml). The organic phase was dried over magnesium sulfate and the solvent evaporated *in vacuo*. The residue was purified using column chromatography (40% to 80% EtOAc in hexane) to give 2-bromo-3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone (1.1 g).

Step 5:Preparation of 2-(1,5-dimethyl-1H-indazol-4-yl)-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione


- 97 -




To a stirred suspension of the 2-bromo-3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone (150 mg, 0.5 mmol), 1,5-dimethyl-1H-indazole-4-boronic acid (148 mg, 0.78 mmol) and freshly powdered potassium phosphate tribasic (221 mg, 1 mmol) in anhydrous, de-gassed toluene (1.5 ml) under a nitrogen atmosphere were added palladium acetate (6 mg, 0.03 mmol) and 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (21 mg, 0.05 mmol). The reaction was heated to 140 °C for 30 minutes in the microwave (using "normal" absorbtion). The reaction was diluted with water (10 ml) and extracted with ethyl acetate (3 x 15 ml). The combined organic extracts were washed with brine (10 ml), dried over magnesium sulfate and evaporated to dryness under reduced pressure to give a brown oil. The crude product was purified by column chromatography (30% EtOAc/hexane to 100% EtOAc as eluent) to give 2-(1,5-Dimethyl-1H-indazol-4-yl)-3-methoxy-5-(tetrahydro-pyran-4-ylmethyl)-cyclopent-2-enone (41 mg). This material was dissolved in acetone (2 ml) and 2 M hydrochloric acid (2 ml) added. This solution was heated in the microwave (using "normal" absorbtion) for 1 h at 120 °C. This solution was then extracted with dichloromethane (2 x 10 ml) and the organics dried over magnesium sulfate and reduced *in vacuo*. The crude material was purified by reverse phase preparative HPLC (eluting with acetonitrile/water) to give 2-(1,5-dimethyl-1H-indazol-4-yl)-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione (4 mg).


Example2:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione



Step 1:Preparation of 2-(4-Chloro-phenyl)-5-methyl-3H-pyrimidin-4-one


Sodium metal (276 mg, 12 mmol) was added portion wise to ethanol (15 ml) under a nitrogen atmosphere and the mixture allowed to stir at room temperature for 30 minutes until the reaction became homogeneous. 4-Chloro-benzamidine (hydroiodide salt) (1.41 g, 5 mmol) was then added in one portion and the reaction stirred at room temperature for a further 60 minutes. 2-Methyl-3-oxo-propionic acid ethyl ester (716 mg, 5.5 mmol) was finally added drop wise. The reaction was left to stir at room temperature overnight before quenching the reaction slowly with 2M Hydrochloric acid until reaction pH measured pH 1. The resulting precipitate was filtered and air dried, to give 2-(4-Chloro-phenyl)-5-methyl-3H-pyrimidin-4-one as a white solid (570 mg, 2.58 mmol, 52% yield).

Step 2:Preparation of 6-Chloro-2-(4-chloro-phenyl)-5-methyl-1,6-dihydro-pyrimidine

2-(4-Chloro-phenyl)-5-methyl-3H-pyrimidin-4-one (570 mg, 2.58 mmol) was suspended in phosphorus oxychloride (2 ml) and heated to reflux for 2 hours. The mixture was cooled before being stripped to dryness *in vacuo*. The resulting crude product was solubilised in water (10 ml) and ethyl acetate (10 ml) and the biphasic mixture separated. The aqueous layer was further extracted with ethyl acetate (10 ml) before the combined organics were dried with magnesium sulphate, filtered and stripped to dryness *in vacuo* to give 6-Chloro-2-(4-chloro-phenyl)-5-methyl-1,6-dihydro-pyrimidine as a white solid (567 mg, 2.38 mmol, 92% yield).

Step 3:Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-cyclopentane-1,3-dione


- 99 -



To a microwave vial was added 1,3-cyclopentadione (0.5 g, 5.1 mmol), 6-Chloro-2-(4-chloro-phenyl)-5-methyl-1,6-dihydro-pyrimidine (1.04 g, 4.3 mmol), palladium acetate (58 mg, 0.26 mmol), 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (183 mg, 0.38 mmol), potassium phosphate tribasic (2.49 g, 11.7 mmol) and 1,4-dioxane (10 ml). This was heated in the microwave for 30 minutes at 150°C on a normal setting with a pre-stir of 25 seconds. The resulting slurry was diluted with ethyl acetate (10 ml) and 2M hydrochloric acid (10 ml) before filtering through celite. The filter cake was washed with further ethyl acetate (10 ml) and 2M hydrochloric acid (10 ml). The resulting biphasic mixture was separated and the aqueous layer was extracted with further ethyl acetate (2 x 10 ml). The combined organic layers were then dried with magnesium sulphate, filtered and stripped to dryness to yield a dark yellow solid. This was purified by normal phase chromatography (gradient system of 100% dichloromethane – 5% methanol: dichloromethane) to give 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-cyclopentane-1,3-dione as a yellow solid (348 mg, 1.16 mmol, 22% yield).

Step 4:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-3-methoxy-cyclopent-2-enone




2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-cyclopentane-1,3-dione (650 mg, 2.16 mmol) was slurried in acetone (20 ml) and potassium carbonate (300 mg, 2.16 mmol) was added. After stirring at room temperature for 5 minutes, dimethyl sulphate (0.20 ml, 2.16 ml) was added in one portion and the reaction was heated to reflux. After 3 hours at reflux the reaction was complete and subsequently cooled to room temperature, before stripping to dryness. The resulting orange solid was re-solubilised in ethyl acetate (20 ml) and 1M sodium hydroxide (20 ml) and the biphasic mixture separated. The aqueous layer was further extracted with ethyl acetate (2 x 10 ml) before combining the organic layers, drying over magnesium sulphate, filtering and stripping to dryness to isolate a brown solid. This was

- 100 -

purified by normal phase chromatography (gradient system of 100% dichloromethane – 5% methanol: dichloromethane) to isolate 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-3-methoxy-cyclopent-2-enone as a yellow foam (526 mg, 1.67 mmol, 77% yield).

Step 5:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone




A solution of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-3-methoxy-cyclopent-2-enone (69.5 mg, 0.22 mmol) in anhydrous tetrahydrofuran (3 ml), under nitrogen, was cooled to -78°C and stirred for 10 minutes before the drop wise addition of lithium diisopropylamide (as a 2.0M solution in hexane/tetrahydrofuran/ethylbenzene) (0.14 ml, 0.28 mmol). The resulting brown solution was stirred at -78°C for a further 45 minutes before the drop wise addition of 4-formyltetrahydropyran (34 mg, 0.30 mmol). The orange solution was then warmed to room temperature. After 2.5h the reaction was shown to have reached a steady state and was therefore quenched with saturated ammonium chloride (5ml), ethyl acetate (5 ml) and water (1 ml). The biphasic mixture was separated and the aqueous layer further extracted with ethyl acetate (2 x 5 ml). Combined organics were dried over magnesium sulphate, filtered and stripped to dryness to yield a yellow oil. This was purified by normal phase chromatography (100% dichloromethane – 5% methanol: dichloromethane) to give 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone as a dark yellow oil (50.3 mg, 0.12 mmol, 53% yield).

Step 6:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene-cyclopentane-1,3-dione]


- 101 -



2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (50 mg, 0.12 mmol) was solubilised in 1,4-dioxane (2.0 ml) to form a yellow solution. Concentrated hydrochloric acid (20 μ l) was then added in one portion and the resulting orange solution was heated to 60°C. It was stirred at this temperature for 6 hours, adding more concentrated hydrochloric acid after 3 hours (20 μ l) and 4 hours (20 μ l), before reaction completion was seen. Reaction was cooled to room temperature and then stripped to dryness to yield 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene-cyclopentane-1,3-dione as a dark orange oil (88 mg, 0.22 mmol, 185% yield).

Step 7:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-(tetrahydro-pyran-4-yl)methyl)-cyclopentane-1,3-dione




2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene-cyclopentane-1,3-dione (88 mg, 0.22 mmol) was solubilised in ethanol (2.5 ml) and 5% palladium on activated charcoal (4.4 mg, 5% weight relative to starting material) was added. The reaction was stirred at room temperature under 1.5 bar pressure of hydrogen for 14.5 hours, adding more 5% palladium on activated charcoal (4.4 mg, 5% weight relative to starting material) after 7.5 hours. Reaction was then filtered through celite, washing the filter

- 102 -

pad with ethanol (10 ml). The resulting solution was stripped to dryness to yield a yellow oil which was purified by reverse phase preparative HPLC (using water, acetonitrile) to give 2-[2-(4-Chloro-phenyl)-5-methyl-pyrimidin-4-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione as a pale yellow oil (9.8 mg, 0.025 mmol, 11% yield).


Example 3:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione



Step 1:

Preparation of 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde




To a suspension of 4-chlorophenyl thioamide (10g, 58.2mmol) in 1,2-dimethoxyethane (80ml) was added 2-chloromalonaldehyde (11.3g, 87.3mmol), followed by magnesium carbonate hexahydrate (14.12g, 30.8mmol), and the reaction heated to 60°C under nitrogen for 3 hours. The crude reaction mixture was filtered through a plug of silica (50g), washed with ethyl acetate, and the combined filtrate concentrated under reduced pressure to give 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde (7.66g).

Step 2:

Preparation of [2-(4-Chloro-phenyl)-thiazol-5-yl]-methanol


- 103 -



To a solution of 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde (5.59g, 25mmol) in methanol (65ml) at 0°C was added sodium borohydride (1.04g, 27.5mmol) portionwise over a period of 5 minutes. The reaction was then allowed to warm to room temperature and stirred under nitrogen for 3 hours. The reaction was quenched with saturated aqueous ammonium chloride solution (100ml) and stirred for 5 minutes, before the methanol was removed under reduced pressure. The resulting heterogeneous mixture was then extracted with ethyl acetate (2 x 100ml), the combined organics dried over magnesium sulphate and the solvent removed under reduced pressure to give [2-(4-Chloro-phenyl)-thiazol-5-yl]-methanol (5.27g).

Step 3:

Preparation of 3-[2-(4-Chloro-phenyl)-thiazol-5-ylmethoxy]-cyclopent-2-enone




To a solution of [2-(4-Chloro-phenyl)-thiazol-5-yl]-methanol (2g, 8.8mmol) in anhydrous THF (50ml), at 0°C, under an atmosphere of nitrogen was added cyclopentane-1,3-dione (1.12g, 11.4mmol), followed by triphenylphosphine (2.99g, 11.4mmol). Diisopropylazodicarboxylate (2.2ml, 11.4mmol) was then added dropwise over a period of 5 minutes and the reaction allowed to warm to room temperature and stirred for 2 hours. The crude reaction mixture was dry loaded onto silica and purified by flash chromatography to give 3-[2-(4-Chloro-phenyl)-thiazol-5-ylmethoxy]-cyclopent-2-enone (2.12g).

Step 4:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-cyclopentane-1,3-dione


- 104 -



To a solution of 3-[2-(4-Chloro-phenyl)-thiazol-5-ylmethoxy]-cyclopent-2-enone (1.29g, 4.22mmol) in ethylene glycol dimethyl ether (10ml) was added 1-butyl-3-methylimidazolium bis(trifluoromethylsulphonyl) imide (100 μ l) and the reaction heated to 230°C by microwave irradiation for 30 minutes. The crude reaction mixture was dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-cyclopentane-1,3-dione (1.12g).

Step 5:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone




To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-cyclopentane-1,3-dione (1.12g, 3.66mmol) in acetone (20ml) was added potassium carbonate (1.04g, 7.32mmol), followed by iodomethane (455 μ l, 7.32mmol), and the reaction heated to 40°C for 6 hours. The solvent was then removed from the crude reaction mixture under reduced pressure and the residue partitioned between water (100ml) and ethyl acetate (100ml). The organic layer was separated, dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone (780mg).

Step 6:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopent-2-enone


- 105 -



To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone (640mg, 2mmol) in anhydrous tetrahydrofuran (12ml) at -78°C under an atmosphere of nitrogen was added lithium diisopropylamide (1.8M in THF/heptanes/ethylbenzene; 1.1ml, 2mmol) dropwise over a period of 5 minutes, and the reaction allowed to stir at 78°C for 30 minutes. A solution of tetrahydro-pyran-4-carbaldehyde (228mg, 2mmol) in anhydrous tetrahydrofuran (1ml) was then added dropwise over a period of 5 minutes before the reaction is allowed to warm to room temperature and stirred for a further 30 minutes. Potassium *tert*-butoxide (337mg, 3 mmol) was then added in one portion and the reaction stirred for a further 90 minutes. The reaction was quenched by the addition of saturated aqueous ammonium chloride solution (50ml) and extracted with ethyl acetate (50ml) and the organic later separated, dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopent-2-enone (195mg).

Step 7:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione




To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopent-2-enone (190mg, 0.45mmol) in acetone (2ml) was added 2N hydrochloric acid (2ml) and the reaction heated to 120°C for 30 minutes by microwave irradiation. The crude reaction was diluted with ethyl acetate (25ml) and washed with saturated aqueous

- 106 -

ammonium chloride solution (25ml), brine (25ml) and the organic later separated, dried over magnesium sulphate and the solvent removed under reduced pressure to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (168mg).


Step 8:

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione




To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (130mg, 0.32mmol) in ethanol (2ml) was added 5% palladium on carbon (13mg), and the reaction stirred under an atmosphere hydrogen at a pressure of 2 bar for 4 hours. The crude reaction mixture was filtered through a pad of Celite and purified by preparative mass-directed HPLC to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione (30mg).

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(5-fluoro-pyridin-2-ylmethyl)-cyclopentane-1,3-dione

**Step 1:**

- 107 -

Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-5-[(5-fluoro-pyridin-2-yl)-hydroxy-methyl]-3-methoxy-cyclopent-2-enone




To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone (400mg, 1.25mmol) in anhydrous tetrahydrofuran (10ml) at -78°C under an atmosphere of nitrogen was added lithium diisopropylamide (1.8M in THF/heptanes/ethylbenzene ; 0.76ml, 1.37mmol) dropwise over a period of 5 minutes, and the reaction allowed to stir at 78°C for 30 minutes. A solution of 5-fluoro-pyridine-2-carbaldehyde (171mg, 1.37mmol) in anhydrous tetrahydrofuran (1ml) was then added dropwise over a period of 5 minutes before the reaction was allowed to warm to room temperature and stirred for a further 30 minutes. The reaction was quenched by the addition of saturated aqueous ammonium chloride solution (50ml) and extracted with ethyl acetate (50ml) and the organic later separated, dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-5-[(5-fluoro-pyridin-2-yl)-hydroxy-methyl]-3-methoxy-cyclopent-2-enone (400mg).

- 108 -

Step 2:


Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione



To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopent-2-enone (190mg, 0.45mmol) in acetone (2ml) was added 2N hydrochloric acid (2ml) and the reaction heated to 130°C for 90 minutes by microwave irradiation. The crude reaction is then dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (93mg).

Step 3:


Preparation of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(5-fluoro-pyridin-2-ylmethyl)-cyclopentane-1,3-dione



To a solution of 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-[1-(5-fluoro-pyridin-2-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (70mg, 0.17mmol) in glacial acetic acid (2ml) was added zinc powder (10mg), and the reaction heated to 80°C for 17 hours. The crude reaction mixture was then dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-methyl-thiazol-4-yl]-4-(5-fluoro-pyridin-2-ylmethyl)-cyclopentane-1,3-dione (55mg).


- 109 -

Preparation of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione



Step 1:


Preparation of 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde



To a suspension of 4-chlorophenyl thioamide (10g, 58.2mmol) in 1,2-dimethoxyethane (80ml) was added 2-chloromalonaldehyde (11.3g, 87.3mmol), followed by magnesium carbonate hexahydrate (14.12g, 30.8mmol), and the reaction heated to 60°C under an atmosphere of nitrogen for 3 hours. The crude reaction mixture is filtered through a plug of silica (50g), which was washed with ethyl acetate, and the combined filtrate concentrated under reduced pressure to give 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde (7.66g).

Step 2:


Preparation of 1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethanol




To a solution of 2-(4-Chloro-phenyl)-thiazole-5-carbaldehyde (4.95g, 22.14mmol) in THF (65ml) at 0°C under an atmosphere of nitrogen was added methyl magnesium chloride (3M in THF;

- 110 -

8.11ml, 24.35mmol) dropwise over a period of 5 minutes. The reaction was then allowed to warm to room temperature and stirred for a further 2 hours. The reaction was quenched with saturated aqueous ammonium chloride solution (100ml) and stirred for 5 minutes, before the methanol is removed under reduced pressure. The resulting heterogeneous mixture was then extracted with ethyl acetate (2 x 100ml), the combined organics dried over magnesium sulphate and the solvent removed under reduced pressure to give 1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethanol (4.71g).

Step 3:Preparation of 3-{1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethoxy}-cyclopent-2-enone

To a solution of 1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethanol (4.1g, 17.1mmol) in anhydrous THF (100ml), at 0°C under an atmosphere of nitrogen was added cyclopentane-1,3-dione (2.18g, 22.23mmol), followed by triphenylphosphine (5.83g, 22.23mmol). Diisopropylazodicarboxylate (4.31ml, 22.23mmol) was then added dropwise over a period of 5 minutes and the reaction allowed to warm to room temperature and stirred for 2 hours. The crude reaction mixture was dry loaded onto silica and purified by flash chromatography to give 3-{1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethoxy}-cyclopent-2-enone (2.29g).

Step 4:Preparation of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-cyclopentane-1,3-dione


To a solution of 3-{1-[2-(4-Chloro-phenyl)-thiazol-5-yl]-ethoxy}-cyclopent-2-enone (2.29g, 7.16mmol) in ethylene glycol dimethyl ether (10ml) was added 1-butyl-3-methylimidazolium bis(trifluoromethylsulphonyl) imide (100μl) and the reaction heated to 230°C by microwave irradiation for 30 minutes. The crude reaction mixture was dry loaded onto silica and purified by

- 111 -

flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-cyclopentane-1,3-dione (681mg).

Step 5:


Preparation of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone



To a solution of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-cyclopentane-1,3-dione (650mg, 2.03mmol) in acetone (10ml) was added potassium carbonate (553mg, 4mmol), followed by iodomethane (248 μ l, 4mmol), and the reaction heated to 40°C for 6 hours. The solvent was then removed from the crude reaction mixture under reduced pressure and the residue partitioned between water (100ml) and ethyl acetate (100ml). The organic layer was separated, dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone (395mg).

Step 6:

Preparation of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone




To a solution of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-3-methoxy-cyclopent-2-enone (175mg, 0.52mmol) in anhydrous tetrahydrofuran (5ml) at -78°C under an atmosphere of nitrogen was added lithium diisopropylamide (1.8M in THF/heptanes/ethylbenzene; 1.1ml,

- 112 -

2mmol) dropwise over a period of 1 minute and the reaction allowed to stir at 78°C for 40 minutes. A solution of tetrahydro-pyran-4-carbaldehyde (228mg, 2mmol) in anhydrous tetrahydrofuran (1ml) was then added dropwise over a period of 30 seconds before the reaction was allowed to warm to room temperature and stirred for a further 30 minutes. The reaction was quenched by the addition of saturated aqueous ammonium chloride solution (50ml) and extracted with ethyl acetate (50ml) and the organic layer separated, dry loaded onto silica and purified by flash chromatography to give 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (86mg).

Step 7:

Preparation of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione





To a stirred solution of 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-3-methoxy-5-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopent-2-enone (85mg, 0.19mmol) in 1,4-dioxane (3ml) was added 12M hydrochloric acid (32μl, 0.38mmol) and the reaction heated to 60°C for 12 hours. The solvent was removed under reduced pressure and the residue purified by mass-directed preparative HPLC to give 2-[2-(4-Chloro-phenyl)-5-ethyl-thiazol-4-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (4mg).


Example 4:

Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione

- 113 -

Step 1:Preparation of [5-(4-Chloro-phenyl)-thiophen-2-yl]-methanol

To a solution of 5-(4-Chloro-phenyl)-thiophene-2-carbaldehyde (5g, 22.45mmol) in methanol (250ml) at 0°C under an atmosphere of nitrogen was added sodium borohydride (1.02g, 26.94mmol) portionwise over a period of 5 minutes. The reaction was then allowed to warm to room temperature and stirred for 3 hours. The reaction was quenched with saturated aqueous ammonium chloride solution (100ml) and stirred for 5 minutes, before the methanol is removed under reduced pressure. The resulting heterogeneous mixture was then extracted with ethyl acetate (2 x 100ml), the combined organics dried over magnesium sulphate and the solvent removed under reduced pressure to give [5-(4-Chloro-phenyl)-thiophen-2-yl]-methanol (4.89g).


Step 2:Preparation of 3-[5-(4-Chloro-phenyl)-thiophen-2-ylmethoxy]-cyclopent-2-en-1-one

- 114 -

To a solution of [2-(4-Chloro-phenyl)-thiazol-5-yl]-methanol (4.89g, 21.75mmol) in anhydrous THF (100ml), at 0°C under an atmosphere of nitrogen was added cyclopentane-1,3-dione (2.78g, 28.27mmol) followed by triphenylphosphine (7.41g, 28.27mmol). Diisopropylazodicarboxylate (5.48ml, 28.27mmol) was then added dropwise over a period of 5 minutes and the reaction allowed to warm to room temperature and stirred for 2 hours. The crude reaction mixture was dry loaded onto silica and purified by flash chromatography to give 3-[5-(4-Chloro-phenyl)-thiophen-2-ylmethoxy]-cyclopent-2-enone (5.1g).

Step 3:


Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-cyclopentane-1,3-dione



To a solution of 3-[5-(4-Chloro-phenyl)-thiophen-2-ylmethoxy]-cyclopent-2-enone (4.89, 4.22mmol) in 1,2-dimethoxyethane (10ml) was added 1-butyl-3-methylimidazolium bis(trifluoromethylsulphonyl) imide (100μl) and the reaction heated to 180°C by microwave irradiation for 30 minutes. The crude reaction mixture was dry loaded onto silica and purified by flash chromatography to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-cyclopentane-1,3-dione (2.21g).

Step 4:

Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-cyclopent-2-enone




To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-cyclopentane-1,3-dione (2.21g, 7.25mmol) in acetone (50ml) was added potassium carbonate (1514g, 10.87mmol) followed by iodomethane (670μl, 10.87mmol) and the reaction heated to 40°C for 6 hours. The solvent was then removed from the crude reaction mixture under reduced pressure and the residue partitioned between water (100ml) and ethyl acetate (100ml). The organic layer was separated,

- 115 -

dry loaded onto silica and purified by flash chromatography to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-cyclopent-2-enone (818mg).

Step 5:


Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone



To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-cyclopent-2-enone (255mg, 0.8mmol) in anhydrous tetrahydrofuran (6ml) at -78°C under an atmosphere of nitrogen was added lithium diisopropylamide (1.8M in THF/heptanes/ethylbenzene; 0.49ml, 0.89mmol) dropwise over a period of 5 minutes and the reaction allowed to stir at 78°C for 30 minutes. A solution of tetrahydro-pyran-4-carbaldehyde (102mg, 0.89mmol) in anhydrous tetrahydrofuran (1ml) was then added dropwise over a period of 5 minutes before the reaction was allowed to warm to room temperature and stirred for a further 30 minutes. The reaction was quenched by the addition of saturated aqueous ammonium chloride solution (50ml) and extracted with ethyl acetate (50ml) and the organic later separated, dry loaded onto silica and purified by flash chromatography to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (292mg).

Step 6:

Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione




- 116 -

To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-5-[hydroxy-(tetrahydro-pyran-4-yl)-methyl]-3-methoxy-cyclopent-2-enone (280mg, 0.65mmol) in acetone (2ml) was added 2N hydrochloric acid (2ml) and the reaction heated to 120°C for 60 minutes by microwave irradiation. The crude reaction was diluted with ethyl acetate (25ml) and washed with saturated aqueous ammonium chloride solution (25ml), brine (25ml) and the organic layer dry loaded onto silica and purified by flash chromatography to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (88mg).


Step 7:


Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione




To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-[1-(tetrahydro-pyran-4-yl)-meth-(E)-ylidene]-cyclopentane-1,3-dione (88mg, 0.22mmol) in ethanol (2ml) was added 5% palladium on carbon (9mg) and the reaction stirred under an atmosphere of hydrogen at a pressure of 2 bar for 4 hours. The crude reaction mixture was filtered through a pad of Celite and purified by preparative mass-directed HPLC to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-pyran-4-ylmethyl)-cyclopentane-1,3-dione (43mg).

Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-furan-3-ylmethyl)-cyclopentane-1,3-dione



Step 1:Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-5-(tetrahydro-furan-3-ylmethyl)-cyclopent-2-enone

To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-cyclopent-2-enone (285mg, 0.89mmol) in anhydrous tetrahydrofuran (6ml) at -78°C under an atmosphere of nitrogen is added lithium diisopropylamide (1.8M in THF/heptanes/ethylbenzene; 0.55ml, 1.0mmol) dropwise over a period of 5 minutes and the reaction allowed to stir at -78°C for 30 minutes. A solution of 3-iodomethyl tetrahydrofuran (212mg, 1mmol) in anhydrous tetrahydrofuran (1ml) was then added dropwise over a period of 5 minutes before the reaction was allowed to warm to room temperature and stirred for a further 30 minutes. The reaction was quenched by the addition of saturated aqueous ammonium chloride solution (50ml) and extracted with ethyl acetate (50ml) and the organic later separated, dry loaded onto silica and purified by flash chromatography to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-5-(tetrahydro-furan-3-ylmethyl)-cyclopent-2-enone (150mg)

Step 2:Preparation of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-furan-3-ylmethyl)-cyclopentane-1,3-dione

To a solution of 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-3-methoxy-5-(tetrahydro-furan-3-ylmethyl)-cyclopent-2-enone (150mg, 0.37mmol) in acetone (2ml) was added 2N hydrochloric

acid (2ml) and the reaction heated to 120°C for 60 minutes by microwave irradiation. The crude reaction was diluted with ethyl acetate (25ml) and washed with saturated aqueous ammonium chloride solution (25ml), brine (25ml). The solvent was removed under reduce pressure and the residue purified by mass-directed preparative HPLC to give 2-[5-(4-Chloro-phenyl)-2-methyl-thiophen-3-yl]-4-(tetrahydro-furan-3-ylmethyl)-cyclopentane-1,3-dione (15mg).

Compounds characterised by HPLC-MS were analysed using one of three methods described below.

Method A

Compounds characterised by HPLC-MS were analysed using a Waters 2795 HPLC equipped with a Waters Atlantis dC18 column (column length 20 mm, internal diameter of column 3 mm, particle size 3 micron, temperature 40 °C), Waters photodiode array and Micromass ZQ2000. The analysis was conducted using a three minutes run time, according to the following gradient table:

Time (mins)	Solvent A (%)	Solvent B (%)	Flow (ml / mn)
0.00	90.0	10.0	2.00
0.25	90.0	10.0	2.00
2.00	10.0	90.0	2.00
2.50	10.0	90.0	2.00
2.60	90.0	10.0	2.00
3.0	90.0	10.0	2.00

Solvent A: H₂O containing 0.1% HCOOH

Solvent B: CH₃CN containing 0.1% HCOOH

Method B

Compounds characterised by HPLC-MS were analysed using an Waters 2777 injector with a 1525 micro pump HPLC equipped with a Waters Atlantis dC18 IS column (column length 20 mm,

- 119 -

internal diameter of column 3 mm, particle size 3 micron), Waters 2996 photodiode array, Waters 2420 ELSD and Micromass ZQ2000. The analysis was conducted using a three minutes run time, according to the following gradient table:

Time (mins)	Solvent A (%)	Solvent B (%)	Flow (ml / mn)
0.00	95.0	5	1.300
2.50	0.00	100	1.300
2.80	0.00	100	1.300
2.90	95.0	5	1.300

Solvent A: H₂O with 0.05% TFA

Solvent B: CH₃CN with 0.05% TFA

Method C:

Compounds characterised by HPLC-MS were analysed using a Finnigan Surveyor MSQ Plus equipped with a Waters Xterra column (column length 50 mm, internal diameter of column 4.6 mm, particle size 3.5 micron, temperature 40 °C), Waters photodiode array and Micromass ZQ2000. The analysis was conducted using a six minutes run time, according to the following gradient table:

Time (mins)	Solvent A (%)	Solvent B (%)	Flow (ml / mn)
0.00	90.0	10.0	1.30
3.80	0.00	100	1.30
4.80	0.00	100	1.30
5.00	90.0	10.0	1.30
6.00	90.0	10.0	1.30

Solvent A: H₂O containing 0.05% HCOOH

Solvent B: CH₃CN containing 0.05% HCOOH

Method D








Compounds characterised by UPLC-MS were analysed using a Waters Acquity UPLC equipped with a Waters Atlantis dC18 column (column length 20 mm, internal diameter of column 3 mm, particle size 3 micron, temperature 40 °C), Waters photodiode array and Micromass ZQ2000. The analysis was conducted using a two minute run time, according to the following gradient table:

Time (mins)	Solvent A (%)	Solvent B (%)	Flow (ml / mn)
0.00	90.0	10.0	2.00
1.50	10.0	90.0	2.00
1.75	10.0	90.0	2.00
1.9	90.0	10.0	2.00
2.00	90.0	10.0	2.00








Solvent A: H₂O containing 0.1% HCOOH









Solvent B: CH₃CN containing 0.1% HCOOH

Table A1

Compound Number	Structure	LC/MS, NMR or other physical data
A1		LC/MS (Method D) ES ⁺ : MH ⁺ = 410 rt= 1.43 min
A2		LC/MS (Method D) ES ⁺ : MH ⁺ = 434, 436 rt= 1.03 min
A3		LC/MS (Method D) ES ⁺ : MH ⁺ = 432, 434 rt= 1.16 min
A4		LC/MS (Method D) ES ⁺ : MH ⁺ = 447, 449 rt= 1.12 min
A5		LC/MS (Method D) ES ⁺ : MH ⁺ = 389, 391 rt= 0.94 min
A6		LC/MS (Method D) ES ⁺ : MH ⁺ = 403, 405 rt= 0.99 min
A7		LC/MS (Method D) ES ⁺ : MH ⁺ = 416, 418 rt= 1.35 min

- 122 -

A8		LC/MS (Method D) ES ⁺ : MH ⁺ = 399, 401 rt= 1.00 min
A9		LC/MS (Method D) ES ⁺ : MH ⁺ = 397, 399 rt= 1.04 min
A10		LC/MS (Method D) ES ⁺ : MH ⁺ = 401, 403 rt= 1.01 min
A11		LC/MS (Method D) ES ⁺ : MH ⁺ = 384, 386 rt= 1.36 min
A12		LC/MS (Method D) ES ⁺ : MH ⁺ = 415, 417 rt= 1.03 min
A13		LC/MS (Method D) ES ⁺ : MH ⁺ = 413, 415 rt= 1.03 min
A14		LC/MS (Method D) ES ⁺ : MH ⁺ = 413, 415 rt= 1.41 min

A15		LC/MS (Method D) ES ⁺ : MH ⁺ = 395, 397 rt= 1.24 min
A16		LC/MS (Method D) ES ⁺ : MH ⁺ = 397, 399 rt= 0.79 min
A17		LC/MS (Method D) ES ⁺ : MH ⁺ = 370 rt= 1.02 min
A18		LC/MS (Method D) ES ⁺ : MH ⁺ = 404, 406 rt= 1.12 min
A19		LC/MS (Method A) ES ⁺ : MH ⁺ = 341, 342 rt = 1.14 min
A20		LC/MS (Method A) ES ⁺ : MH ⁺ = 398, 400 rt = 1.41 min
A21		LC/MS (Method A) ES ⁺ : MH ⁺ = 396, 398 rt= 1.45 min
A22		LC/MS (Method A) ES ⁺ : MH ⁺ = 396, 398 rt= 1.59 min

- 124 -





A23		LC/MS (Method A) ES ⁺ : MH ⁺ = 385, 387 rt= 1.40 min
A24		LC/MS (Method D) ES ⁺ : MH ⁺ = 402, 404 rt= 1.27 min
A25		LC/MS (Method A) ES ⁺ : MH ⁺ = 387, 389 rt= 1.35 min

Table 1 covers 98 compounds of the following type:



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1 below:

Compound d Number	R ⁶	R ⁷
1.001	CH ₃	H
1.002	CH ₃	F
1.003	CH ₃	Cl
1.004	CH ₃	Br
1.005	CH ₃	I
1.006	CH ₃	CH ₃
1.007	CH ₃	CH ₂ CH ₃
1.008	CH ₃	<i>n</i> -Pr
1.009	CH ₃	<i>i</i> -Pr


1.010	CH ₃	<i>n</i> -Bu
1.011	CH ₃	<i>i</i> -Bu
1.012	CH ₃	<i>t</i> -Bu
1.013	CH ₃	OCH ₃
1.014	CH ₃	OCH ₂ CH ₃
1.015	CH ₃	<i>O</i> <i>n</i> -Pr
1.016	CH ₃	<i>O</i> <i>i</i> -Pr
1.017	CH ₃	-CCH
1.018	CH ₃	-CH=CH ₂
1.019	CH ₃	Phenyl
1.020	CH ₃	2-fluorophenyl
1.021	CH ₃	2-chlorophenyl
1.022	CH ₃	2-trifluoromethylphenyl
1.023	CH ₃	2-nitrophenyl
1.024	CH ₃	2-methylphenyl
1.025	CH ₃	2-methanesulfonylphenyl
1.026	CH ₃	2-cyanophenyl
1.027	CH ₃	3-fluorophenyl
1.028	CH ₃	3-chlorophenyl
1.029	CH ₃	3-trifluoromethylphenyl
1.030	CH ₃	3-nitrophenyl
1.031	CH ₃	3-methylphenyl
1.032	CH ₃	3-methanesulfonylphenyl
1.033	CH ₃	3-cyanophenyl
1.034	CH ₃	4-fluorophenyl
1.035	CH ₃	4-chlorophenyl
1.036	CH ₃	4-bromophenyl
1.037	CH ₃	4-difluoromethoxyphenyl
1.038	CH ₃	2-fluoro-4-chlorophenyl
1.039	CH ₃	3-fluoro-4-chlorophenyl
1.040	CH ₃	2-chloro-4-chlorophenyl
1.041	CH ₃	2-chloro-4-fluorophenyl

1.042	CH ₃	3-chloro-4-chlorophenyl
1.043	CH ₃	3-chloro-4-fluorophenyl
1.044	CH ₃	2-methyl-4-chlorophenyl
1.045	CH ₃	4-trifluoromethylphenyl
1.046	CH ₃	4-nitrophenyl
1.047	CH ₃	4-methylphenyl
1.048	CH ₃	4-methanesulfonylphenyl
1.049	CH ₃	4-cyanophenyl
1.050	CH ₂ CH ₃	H
1.051	CH ₂ CH ₃	F
1.052	CH ₂ CH ₃	Cl
1.053	CH ₂ CH ₃	Br
1.054	CH ₂ CH ₃	I
1.055	CH ₂ CH ₃	CH ₃
1.056	CH ₂ CH ₃	CH ₂ CH ₃
1.057	CH ₂ CH ₃	<i>n</i> -Pr
1.058	CH ₂ CH ₃	<i>i</i> -Pr
1.059	CH ₂ CH ₃	<i>n</i> -Bu
1.060	CH ₂ CH ₃	<i>i</i> -Bu
1.061	CH ₂ CH ₃	<i>t</i> -Bu
1.062	CH ₂ CH ₃	OCH ₃
1.063	CH ₂ CH ₃	OCH ₂ CH ₃
1.064	CH ₂ CH ₃	<i>On</i> -Pr
1.065	CH ₂ CH ₃	<i>Oi</i> -Pr
1.066	CH ₂ CH ₃	-CCH
1.067	CH ₂ CH ₃	-CH=CH ₂
1.068	CH ₂ CH ₃	Phenyl
1.069	CH ₂ CH ₃	2-fluorophenyl
1.070	CH ₂ CH ₃	2-chlorophenyl
1.071	CH ₂ CH ₃	2-trifluoromethylphenyl
1.072	CH ₂ CH ₃	2-nitrophenyl
1.073	CH ₂ CH ₃	2-methylphenyl
1.074	CH ₂ CH ₃	2-

		methanesulfonylphenyl
1.075	CH ₂ CH ₃	2-cyanophenyl
1.076	CH ₂ CH ₃	3-fluorophenyl
1.077	CH ₂ CH ₃	3-chlorophenyl
1.078	CH ₂ CH ₃	3-trifluoromethylphenyl
1.079	CH ₂ CH ₃	3-nitrophenyl
1.080	CH ₂ CH ₃	3-methylphenyl
1.081	CH ₂ CH ₃	3-methanesulfonylphenyl
1.082	CH ₂ CH ₃	3-cyanophenyl
1.083	CH ₂ CH ₃	4-fluorophenyl
1.084	CH ₂ CH ₃	4-chlorophenyl
1.085	CH ₂ CH ₃	4-bromophenyl
1.086	CH ₂ CH ₃	4-difluoromethoxyphenyl
1.087	CH ₂ CH ₃	2-fluoro-4-chlorophenyl
1.088	CH ₂ CH ₃	3-fluoro-4-chlorophenyl
1.089	CH ₂ CH ₃	2-chloro-4-chlorophenyl
1.090	CH ₂ CH ₃	2-chloro-4-fluorophenyl
1.091	CH ₂ CH ₃	3-chloro-4-chlorophenyl
1.092	CH ₂ CH ₃	3-chloro-4-fluorophenyl
1.093	CH ₂ CH ₃	2-methyl-4-chlorophenyl
1.094	CH ₂ CH ₃	4-trifluoromethylphenyl
1.095	CH ₂ CH ₃	4-nitrophenyl
1.096	CH ₂ CH ₃	4-methylphenyl
1.097	CH ₂ CH ₃	4-methanesulfonylphenyl
1.098	CH ₂ CH ₃	4-cyanophenyl


Table 2 covers 98 compounds of the following type

- 128 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 3 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 4 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 5 covers 98 compounds of the following type

- 129 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 6 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 7 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 8 covers 98 compounds of the following type

- 130 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 8 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 9 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 10 covers 98 compounds of the following type

- 131 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 11 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 12 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 13 covers 98 compounds of the following type

- 132 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 14 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 15 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 16 covers 98 compounds of the following type

- 133 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 17 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 18 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 19 covers 98 compounds of the following type

- 134 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 20 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 21 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 22 covers 98 compounds of the following type

- 135 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 23 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 24 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 25 covers 98 compounds of the following type

- 136 -




where G , is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 26 covers 98 compounds of the following type



where G , is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 26 covers 98 compounds of the following type



where G , is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 28 covers 98 compounds of the following type

- 137 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 29 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 30 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 31 covers 98 compounds of the following type

- 138 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 32 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 33 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 34 covers 98 compounds of the following type

- 139 -




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 35 covers 98 compounds of the following type




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 36 covers 98 compounds of the following type




where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 37 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 38 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.


Table 39 covers 98 compounds of the following type

- 141 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 40 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 41 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 42 covers 98 compounds of the following type

- 142 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 43 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 44 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 44 covers 98 compounds of the following type

- 143 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 45 covers 98 compounds of the following type




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 46 covers 98 compounds of the following type




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 47 covers 98 compounds of the following type




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 48 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 49 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 50 covers 98 compounds of the following type

- 145 -




where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.

Table 51 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 52 covers 98 compounds of the following type



where G, is hydrogen, and R⁶ and R⁷ are as described in Table 1.


Table 54 covers 98 compounds of the following type

- 146 -



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Table 55 covers 98 compounds of the following type



where G, is hydrogen, and R^6 and R^7 are as described in Table 1.

Biological Examples

Example A:

Seeds of a variety of test species are sown in standard soil in pots. After cultivation for one day (pre-emergence) or after 8 days cultivation (post-emergence) under controlled conditions in a glasshouse (at 24/16°C, day/night; 14 hours light; 65 % humidity), the plants are sprayed with an aqueous spray solution derived from the formulation of the technical active ingredient in acetone / water (50:50) solution containing 0.5% Tween 20 (polyoxyethylene sorbitan monolaurate, CAS RN 9005-64-5).

The test plants are then grown in a glasshouse under controlled conditions in a glasshouse (at 24/16°C, day/night; 14 hours light; 65 % humidity) and watered twice daily. After 13 days for pre and post-emergence, the test is evaluated (100 = total damage to plant; 0 = no damage to plant).

Test plants:

Lolium perenne (LOLPE), *Alopecurus myosuroides* (ALOMY), *Echinochloa crus-galli* (ECHCG), *Avena fatua* (AVEFA)

Post-Emergence Activity

Compound Number	Rate g/ha	LOLPE	ALOMY	ECHCG	AVEFA
A1	250	0	10	10	0
A2	250	10	10	20	0
A3	250	0	10	20	0
A4	250	0	20	70	0
A5	250	30	50	80	50
A6	250	0	50	60	0
A7	250	60	90	100	100
A8	250	50	50	90	50
A9	250	40	20	100	60
A10	250	10	20	70	0
A11	250	0	10	80	0
A12	250	100	100	100	90
A13	250	0	10	20	10
A15	250	0	10	20	0
A16	250	100	90	100	100
A17	250	100	90	100	90
A18	250	100	100	100	100
A19	250	0	20	40	10
A20	250	90	80	100	90
A21	250	60	70	90	30
A22	250	50	70	90	60
A23	250	100	90	100	100
A24	250	90	100	100	90
A25	250	100	100	100	100

Pre-Emergence Activity

- 148 -

Compound Number	Rate g/ ha	LOLPE	ALOMY	ECHCG	AVEFA
A1	10	10	50	0	A1
A7	80	70	100	70	A7
A8	80	50	100	60	A8
A9	70	10	100	50	A9
A12	100	100	100	90	A12
A16	90	90	100	80	A16
A17	100	70	100	50	A17
A18	100	80	100	90	A18
A20	70	40	100	20	A20
A21	90	40	90	10	A21
A22	60	40	70	20	A22
A23	100	90	100	90	A23
A24	100	60	100	60	A24
A25	90	70	100	70	A25

EXAMPLE B:

Seeds of the Winter Wheat variety 'Hereward' were sown in standard soil in pots. After 8 days cultivation under controlled conditions in a glasshouse (at 24/16°C, day/night; 14 hours light; 65 % humidity), the plants were sprayed post-emergence with an aqueous spray solution derived from the formulation of the technical active ingredient in acetone / water (50:50) solution containing 0.5% Tween 20 (polyoxyethylene sorbitan monolaurate, CAS RN 9005-64-5).

Seeds of the Winter Wheat variety 'Hereward' were seed treated with a wettable powder formulation of the cereal herbicide safener, cloquintocet mexyl, at a rate of 0.5 grams per kilogram of dry seed prior to the initiation of glasshouse testing. One seed was sown per 1.5 inch plastic pot into a sandy loam soil at a depth of 1cm, 8 days prior to application of the test compounds and was watered and grown under controlled conditions in a glasshouse (at 24/16°C, day/night; 14 hours light; 65 % humidity). The plants were sprayed post-emergence with an aqueous spray solution derived from the formulation of the technical active ingredient in acetone / water (50:50) solution containing 0.5% Tween 20 (polyoxyethylene sorbitan monolaurate, CAS RN 9005-64-5).

2010277350 04 May 2012

- 149 -

The test plants were then grown in a glasshouse under controlled conditions in a glasshouse (at 24/16°C, day/night; 14 hours light; 65 % humidity) and watered twice daily. After 13 days for pre and post-emergence, the test was evaluated (100 = total damage to plant; 0 = no damage to plant).

Compound Number	Rate g/ha	Winter Wheat (Hereward)	Winter Wheat (Hereward) + cloquintocet mexyl
A1	250	20	0
A6	250	10	0
A7	250	90	40
A8	250	20	0
A9	250	50	0
A12	250	70	80
A15	250	10	0
A16	250	60	60
A17	250	60	40
A18	250	80	90
A20	250	70	70
A21	250	20	10
A22	250	60	10
A23	250	80	60
A24	250	70	0
A25	250	80	40

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.


The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

2010277350 20 Aug 2015

- 150 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound of formula I



wherein:

G is hydrogen or an agriculturally acceptable metal, sulfonium, ammonium or latentating group; and

R¹ is hydrogen, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆alkenyloxy, C₃-C₆haloalkenyloxy, C₃-C₆alkynyloxy, C₃-C₆cycloalkyl, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylsulfonyl, C₁-C₆alkoxysulfonyl, C₁-C₆haloalkoxysulfonyl, cyano, nitro, phenyl; phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or benzyl or benzyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; or C₃-C₆cycloalkylC₁-C₃alkyl in which a ring or chain methylene group is optionally replaced by an oxygen or sulfur atom;

R² and R³ independently are hydrogen or methyl; and

R⁴ and R⁵ independently are hydrogen or methyl;









and/or R¹ and R⁴ together form a bond; and

Q is C₃-C₈ saturated or mono-unsaturated heterocyclyl containing at least one heteroatom selected from O, N and S, unsubstituted or substituted by a residue of formula =O, =N-R¹⁰, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxyC₁-C₂alkyl, C₃-C₆cycloalkyl, phenyl or phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, where R¹⁰ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₁-C₆haloalkylsulfinyl or C₁-C₆haloalkylsulfonyl; or

Q is an heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl; and





m is 1; and

Het is a group selected from the formulae Het₁ to Het₁₂:

Het₁Het₂Het₃Het₄Het₅Het₆Het₇Het₈

2010277350 20 Aug 2015

- 152 -

Het₉Het₁₀Het₁₁Het₁₂

wherein:

B designates the point of attachment to the ketoenol moiety;

W¹ is N or CR⁹;

W² and W³ are independently of each other N or CR⁷;

W⁴ is N or CR¹⁰;

with the proviso that at least one of W¹, W², W³ or W⁴ is N;

X is O, S, or NR¹²;

Z is N or CR¹³;

wherein

R⁶ is halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, nitro or cyano;

R⁷ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₃-C₆ cycloalkyl, C₅-C₆cycloalkenyl, halogen, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, or C₁-C₆haloalkoxy; or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, cyano or

2010277350 20 Aug 2015

- 153 -

nitro; or phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano;

R⁸ is hydrogen, C₁-C₄ alkyl, C₂-C₃alkenyl, C₂-C₃alkynyl, C₁-C₄ haloalkyl, or C₂-C₃ haloalkenyl;

R⁹ is hydrogen, methyl, halomethyl, or halogen;

R¹⁰ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄ alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl or cyano;

R¹¹ is hydrogen, methyl, ethyl, halomethyl, or haloethyl; or optionally substituted heteroaryl wherein the optional substituents are selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, cyano or nitro; or phenyl, substituted once, twice or three times, by halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy or cyano;

R¹² is hydrogen, methyl, ethyl, or halomethyl; and

R¹³ is hydrogen, methyl, ethyl, halomethyl, haloethyl, halogen, cyano or nitro;

and wherein the latentating group G is selected from the groups C₁-C₈alkyl, C₂-C₈haloalkyl, phenylC₁-C₈alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroarylC₁-C₈alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₈alkenyl, C₃-C₈haloalkenyl, C₃-C₈alkynyl, C(X^a)-R^a, C(X^b)-X^c-R^b, C(X^d)-N(R^c)-R^d, -SO₂-R^e, -P(X^e)(R^f)-R^g and CH₂-X^f-R^h;

wherein X^a, X^b, X^c, X^d, X^e and X^f are independently of each other oxygen or sulfur;

and wherein R^a is H, C₁-C₁₈alkyl, C₂-C₁₈alkenyl, C₂-C₁₈alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-

2010277350 20 Aug 2015

- 154 -

C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro;

R^b is C₁-C₁₈alkyl, C₃-C₁₈alkenyl, C₃-C₁₈alkynyl, C₂-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₂-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; and

R^c and R^d are each independently of each other hydrogen, C₁-C₁₀alkyl, C₃-C₁₀alkenyl, C₃-C₁₀alkynyl, C₂-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-

C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₂-C₅)alkylaminoalkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl amino or heteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or diheteroaryl amino or diheteroaryl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or phenyl amino or phenyl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diphenyl amino or diphenyl amino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or C₃-C₇cycloalkyl amino, di-C₃-C₇cycloalkyl amino or C₃-C₇cycloalkoxy;

or R^c and R^d join together to form a 3-7 membered ring, optionally containing one heteroatom selected from O or S; and

R^e is C₁-C₁₀alkyl, C₂-C₁₀alkenyl, C₂-C₁₀alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl,

2010277350 20 Aug 2015

- 156 -

C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or heteroarylamino or heteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diheteroarylamino or diheteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or C₃-C₇cycloalkylamino, diC₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino or C₂-C₈dialkylamino;

R^f and R^g are each independently of each other C₁-C₁₀alkyl, C₂-C₁₀alkenyl, C₂-C₁₀alkynyl, C₁-C₁₀alkoxy, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₂-C₅)alkylaminoalkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano, or by nitro), C₂-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or heteroarylamino or heteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro; or diheteroarylamino or diheteroarylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl,





















C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or phenylamino or phenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or diphenylamino or diphenylamino substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; or C₃-C₇cycloalkylamino, diC₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino or C₂-C₈dialkylamino; or benzyloxy or phenoxy, wherein the benzyl and phenyl groups are in turn optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or nitro; and

R^h is C₁-C₁₀alkyl, C₃-C₁₀alkenyl, C₃-C₁₀alkynyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₂-C₁₀aminoalkyl, C₁-C₅alkylamino(C₁-C₅)alkyl, C₂-C₈dialkylamino(C₁-C₅)alkyl, C₃-C₇cycloalkyl(C₁-C₅)alkyl, C₁-C₅alkoxy(C₁-C₅)alkyl, C₃-C₅alkenyloxy(C₁-C₅)alkyl, C₃-C₅alkynyloxy(C₁-C₅)alkyl, C₁-C₅alkylthio(C₁-C₅)alkyl, C₁-C₅alkylsulfinyl(C₁-C₅)alkyl, C₁-C₅alkylsulfonyl(C₁-C₅)alkyl, C₂-C₈alkylideneaminoxy(C₁-C₅)alkyl, C₁-C₅alkylcarbonyl(C₁-C₅)alkyl, C₁-C₅alkoxycarbonyl(C₁-C₅)alkyl, aminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylaminocarbonyl(C₁-C₅)alkyl, C₂-C₈dialkylaminocarbonyl(C₁-C₅)alkyl, C₁-C₅alkylcarbonylamino(C₁-C₅)alkyl, *N*-(C₁-C₅)alkylcarbonyl-*N*-(C₁-C₅)alkylamino(C₁-C₅)alkyl, C₃-C₆trialkylsilyl(C₁-C₅)alkyl, phenyl(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryl(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), phenoxy(C₁-C₅)alkyl (wherein the phenyl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), heteroaryloxy(C₁-C₅)alkyl (wherein the heteroaryl is optionally substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, halogen, cyano or by nitro), C₃-C₅haloalkenyl, C₃-C₈cycloalkyl; or phenyl or phenyl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen or by nitro; or heteroaryl or heteroaryl substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano or by nitro;




































and wherein the term "heteroaryl" means an aromatic ring system containing at least one heteroatom and consisting either of a single ring or of two fused rings;

and wherein the compound is optionally an agronomically acceptable salt thereof.

2. A compound according to claim 1, wherein R^1 is hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy.
3. A compound according to claim 2, wherein R^1 is hydrogen or methyl.
4. A compound according to any one of claims 1 to 3, wherein R^4 and R^5 independently are hydrogen or methyl.
5. A compound according to any one of claims 1 to 4, wherein Q is a group selected from formulae Q_1 to Q_{107} as defined below:

































- 159 -






























2010277350 20 Aug 2015

- 160 -



- 161 -



wherein:

2010277350 20 Aug 2015

- 162 -

R is hydrogen, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆alkenyloxy, C₃-C₆haloalkenyloxy, C₃-C₆alkynyloxy, C₃-C₆cycloalkyl, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkoxysulfonyl, C₁-C₆haloalkoxysulfonyl, cyano, nitro, phenyl, phenyl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl, or heteroaryl or heteroaryl substituted by C₁-C₄alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, cyano, nitro, halogen, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl or C₁-C₃alkylsulfonyl;

R' is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₆-C₁₀arylsulfonyl, C₆-C₁₀arylcarbonyl, C₆-C₁₀arylaminocarbonyl, C₇-C₁₆arylalkylaminocarbonyl, C₁-C₉heteroarylsulfonyl, C₁-C₉heteroarylcarbonyl, C₁-C₉heteroarylaminocarbonyl, C₂-C₁₅heteroarylalkylaminocarbonyl;

R'' is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₇cycloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆alkylcarbonyl, C₁-C₆haloalkylcarbonyl, C₁-C₆alkoxycarbonyl, C₁-C₆alkylaminocarbonyl, C₂-C₈dialkylaminocarbonyl, C₁-C₆haloalkylsulfinyl or C₁-C₆haloalkylsulfonyl;

n is 0, 1, 2, 3 or 4; and

A denotes the position of attachment to the $-(CR^4R^5)_m-$ moiety.

6. A compound according to claim 5, wherein

Q is a group selected from formulae Q₁, Q₂, Q₃, Q₄, Q₅, Q₆, Q₇, Q₂₅, Q₂₆, Q₂₇, Q₂₈, Q₂₉, Q₈₆, Q₈₇, Q₈₈, Q₈₉, and Q₉₀ as defined in claim 5;

R and R' are independently hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy; and

n is 0, 1 or 2.

7. A compound according to claim 6, wherein Q is a group selected from formulae Q₁, Q₂, Q₃, Q₄, Q₅, Q₆ and Q₇.
8. A compound according to claim 6 or claim 7, wherein n is 0.
9. A compound according to any one of claims 1 to 4, wherein Q is a 5- or 6-membered heteroaryl or is a 5- or 6-membered heteroaryl which is substituted one to three times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.
10. A compound according to any one of claims 1 to 4, wherein Q is thienyl, furyl, oxazolyl, isoxazolyl, benzofuryl, thiazolyl, oxazolyl, isothiazolyl, benzothienyl, benzoisothienyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl or benzisoxazolyl, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.
11. A compound according to any one of claims 1 to 4, wherein Q is pyridyl, pyrimidinyl, pyridazinyl, quinoliny, isoquinoliny, cinnoliny, phthalazinyl, quinazolinyl or quinoxaliny, where these rings are optionally substituted one or two times by fluoro, chloro, bromo, methyl, methoxy, cyano or trifluoromethyl.
12. A compound according to any one of claims 1 to 11, wherein Het is an optionally substituted monocyclic 5-membered sulfur or nitrogen containing heteroaromatic ring.
13. A compound according to claim 12, wherein Het is a monocyclic 5-membered sulfur and nitrogen containing heteroaromatic ring.
14. A compound according to any one of claims 1 to 13, wherein, in the formulae Het₁ to Het₁₂ :

B designates the point of attachment to the ketoenol moiety;

W¹ is N or CR⁹;

W² and W³ are independently of each other N or CR⁷;

2010277350 20 Aug 2015

- 164 -

W^4 is N or CR^{10} ;

with the proviso that at least one of W^1 , W^2 , W^3 or W^4 is N;

X is O, S, or NR^{12} ;

Z is N or CR^{13} ;

wherein

R^6 is halogen, C_1 - C_2 alkyl, C_1 - C_2 haloalkyl, vinyl, ethynyl, or methoxy;

R^7 is phenyl, substituted once, twice or three times, by halogen, C_1 - C_2 alkyl, C_1 - C_2 alkoxy, C_1 - C_2 haloalkyl, C_1 - C_2 haloalkoxy or cyano;

R^8 is hydrogen, C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_4 haloalkyl or C_2 - C_3 haloalkenyl;

R^9 is hydrogen, methyl, halomethyl or halogen;

R^{10} is hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 haloalkenyl, C_2 - C_4 alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl or cyano;

R^{11} is phenyl, substituted once, twice or three times, by halogen, C_1 - C_2 alkyl, C_1 - C_2 alkoxy, C_1 - C_2 haloalkyl, C_1 - C_2 haloalkoxy or cyano;

R^{12} is hydrogen, methyl, ethyl or halomethyl; and

R^{13} is hydrogen, methyl, ethyl, halomethyl, haloethyl, halogen, cyano or nitro.

15. A compound according to claim 14, wherein:

2010277350 20 Aug 2015

- 165 -

R⁶ is methyl or ethyl;

R⁸ is methyl or ethyl;

R⁹ is hydrogen; and

R¹⁰ is hydrogen, halogen, methyl or ethyl.

16. A compound according to any one of claims 1 to 15, wherein G is hydrogen, an alkali metal or an alkaline earth metal.

17. A herbicidal composition, which comprises a herbicidally effective amount of a compound of formula I as defined in any one of claims 1 to 16, and optionally a further herbicide as mixture partner for the compound of formula I, or optionally a safener, or both.

18. A herbicidal composition according to claim 17, which comprises the further herbicide as mixture partner for the compound of formula I.

19. A herbicidal composition according to claim 17, which comprises a herbicidally effective amount of a compound of formula I as defined in any one of claims 1 to 16, a safener, and optionally a further herbicide as mixture partner for the compound of formula I, wherein the safener is benoxacor, cloquintocet-mexyl, cyprosulfamide, mefenpyr-diethyl or N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide.

20. A method of controlling grasses and weeds in crops of useful plants, which comprises applying a herbicidally effective amount of a compound of formula I as defined in any one of claims 1 to 16, or of a composition comprising such a compound as defined in any one of claims 17, 18 or 19, to the plants or to the locus thereof.

21. Compound according to claim 1, composition according to claim 17, or method according to claim 20, substantially as hereinbefore described with reference to any one of the examples.