



(86) Date de dépôt PCT/PCT Filing Date: 2019/01/04

(87) Date publication PCT/PCT Publication Date: 2019/07/11

(45) Date de délivrance/Issue Date: 2023/01/03

(85) Entrée phase nationale/National Entry: 2020/07/02

(86) N° demande PCT/PCT Application No.: US 2019/012405

(87) N° publication PCT/PCT Publication No.: 2019/136291

(30) Priorités/Priorities: 2018/01/05 (US62/613,939);
2018/03/15 (US62/643,657)

(51) Cl.Int./Int.Cl. *A61K 31/48* (2006.01),
A61K 31/00 (2006.01), *A61K 9/00* (2006.01),
A61P 25/06 (2006.01)


(72) Inventeurs/Inventors:
HOEKMAN, JOHN D., US;
SATTERLY, KELSEY H., US;
SHREWSBURY, STEPHEN B., US;
YOUNG, SCOTT B., US;
FULLER, CHRISTOPHER, US

(73) Propriétaire/Owner:
IMPEL PHARMACEUTICALS INC., US

(74) Agent: SMART & BIGGAR LP

(54) Titre : ADMINISTRATION INTRANASALE DE DIHYDROERGOTAMINE PAR UN DISPOSITIF OLFACTIF DE PRECISION

(54) Title: INTRANASAL DELIVERY OF DIHYDROERGOTAMINE BY PRECISION OLFACTORY DEVICE



(57) Abrégé/Abstract:

Methods are provided for acutely treating migraine headache with or without aura. The methods comprise administering to a subject with migraine headache an effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or a salt thereof, wherein the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration (C_{max}) of at least 750 pg/ml, (b) with a mean time to C_{max} (T_{max}) of DHE of less than 45 minutes, and (c) a mean plasma AUC_{0-inf} of DHE of at least 2500 pg*hr/ml. Also provided are kits for acutely treating migraine with or without aura in which a liquid pharmaceutical composition comprising DHE or DHE salt is contained within a sealed vial that is attachable to a precision intranasal olfactory delivery device packaged therewith.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
11 July 2019 (11.07.2019)



(10) International Publication Number
WO 2019/136291 A1

(51) International Patent Classification:

A61K 31/48 (2006.01) *A61K 31/00* (2006.01)
A61K 9/00 (2006.01) *A61P 25/06* (2006.01)

(21) International Application Number:

PCT/US2019/012405

(22) International Filing Date:

04 January 2019 (04.01.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/613,939 05 January 2018 (05.01.2018) US
62/643,657 15 March 2018 (15.03.2018) US

(71) Applicant: **IMPEL NEUROPHARMA, INC.** [US/US];
201 Elliott Ave, Suite 260, Seattle, WA 98119 (US).

(72) Inventors: **HOEKMAN, John, D.**; Impel Neuropharma, Inc., 201 Elliott Ave, Suite 260, Seattle, WA 98119 (US). **SATTERLY, Kelsey, H.**; Impel Neuropharma, Inc., 201 Elliott Ave, Suite 260, Seattle, WA 98119 (US). **SHREWSBURY, Stephen, B.**; Impel Neuropharma, Inc., 201 Elliott Ave, Suite 260, Seattle, WA 98119 (US). **YOUMANS, Scott, B.**; Impel Neuropharma, Inc., 201 Elliott Ave, Suite 260, Seattle, WA 98119 (US). **FULLER,**

Christopher; Impel Neuropharma, Inc., 201 Elliott Ave, Suite 260, Seattle, WA 98119 (US).

(74) Agent: **KEIPER, Alison, E.** et al.; Fenwick & West LLP, 801 California Street, Mountain View, CA 94041 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: INTRANASAL DELIVERY OF DIHYDROERGOTAMINE BY PRECISION OLFACTORY DEVICE




FIG. 1

(57) Abstract: Methods are provided for acutely treating migraine headache with or without aura. The methods comprise administering to a subject with migraine headache an effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or a salt thereof, wherein the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration (C_{max}) of at least 750 pg/ml, (b) with a mean time to C_{max} (T_{max}) of DHE of less than 45 minutes, and (c) a mean plasma AUC_{0-inf} of DHE of at least 2500 pg*hr/ml. Also provided are kits for acutely treating migraine with or without aura in which a liquid pharmaceutical composition comprising DHE or DHE salt is contained within a sealed vial that is attachable to a precision intranasal olfactory delivery device packaged therewith.

WO 2019/136291 A1

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

INTRANASAL DELIVERY OF DIHYDROERGOTAMINE BY PRECISION OLFACTORY DEVICE

1. CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to US provisional application nos. 62/643,657, filed March 15, 2018, and 62/613,939, filed January 5, 2018.

2. BACKGROUND

[0002] Dihydroergotamine (DHE), a semisynthetic derivative of the ergot alkaloid ergotamine, has been approved for over 70 years for the treatment of migraines. The exact mechanism of action of DHE is not known, but DHE is known to act as a serotonin receptor agonist, cause vasoconstriction of intracranial blood vessels, and interact centrally with dopamine and adrenergic receptors.

[0003] The oral bioavailability of DHE is poor, and DHE is commonly administered parenterally as the mesylate salt by subcutaneous, intramuscular or intravenous injection, and where approved, by nasal spray. Because migraine headaches are episodic and occur unpredictably, administration by nasal spray is far more convenient for treatment of acute migraine than is administration by injection. However, the previously approved nasal spray drug-device combination product provides only 32% of the bioavailability of the intravenous injection, and variable efficacy (among other factors) has led to its withdrawal from market in the EU and other countries, although it remains available in the United States.

[0004] There is, therefore, a need for an intranasal DHE product that provides increased bioavailability and reduced variability in systemically delivered dose for treatment of migraine.

3. SUMMARY

[0005] We designed a manually actuated, propellant-driven, intranasal administration device that can reproducibly deliver metered doses of liquid pharmaceutical compositions beyond the nasal valve to more distal regions of the nasal cavity. We tested the device in a Phase I clinical trial

that compared the bioavailability of (i) dihydroergotamine (DHE) mesylate administered as a single divided 1.45 mg intranasal dose using our Precision Olfactory Delivery (POD[®]) Device (“INP104”); (ii) a 2.0 mg dose of DHE mesylate administered intranasally using Migranal[®] Nasal Spray according to the US FDA approved product label; and (iii) a 1.0 mg intravenous injection of DHE mesylate for injection (D.H.E. 45[®]), in healthy adult subjects.

[0006] As described in detail in Example 2, INP104 provided 4-fold higher mean maximal plasma concentration, nearly 3-fold higher mean systemic drug exposure, and reached maximal DHE plasma concentration faster than Migranal[®]. The higher maximal plasma concentration and systemic drug exposure were achieved with a lower administered dose of the identical formulation of DHE mesylate, 1.45 mg for INP104 versus 2.0 mg for Migranal[®], and without requiring a 15-minute wait between administration of divided sub-doses, as required for Migranal[®]. In addition, systemic delivery of DHE was more consistent with INP104 than with Migranal[®], with lower coefficient of variation (CV%) in DHE AUC_{0-inf} and C_{max} observed across subjects.

[0007] Accordingly, in a first aspect, methods are provided for acutely treating migraine headache with or without aura. The methods comprise administering to a subject with migraine headache an effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or a salt thereof, wherein the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration (C_{max}) of at least 750 pg/ml, (b) with a mean time to C_{max} (T_{max}) of DHE of less than 45 minutes, and (c) a mean plasma AUC_{0-inf} of DHE of at least 2500 pg*hr/ml.

[0008] In various embodiments, the dose is no more than 2.0 mg DHE or salt thereof, less than 2.0 mg DHE or salt thereof, 1.2-1.8 mg DHE or salt thereof, or 1.4-1.6 mg DHE or salt thereof. In a particular embodiment, the dose is about 1.45 mg DHE or salt thereof.

[0009] In a variety of embodiments, the dose is administered as a plurality of divided doses. In certain embodiments, the dose is administered as two divided doses. In a particular embodiment, one divided dose is administered to each nostril. In typical divided dose embodiments, the dose is administered over no more than 1 minute, over no more than 45 seconds, or over no more than 30 seconds. In various embodiments, the volume of liquid composition administered per divided dose is 140-250 µL, 175 µL-225 µL, about 200 µL, or about 180 µL.

[0010] In typical embodiments, the liquid composition comprises a salt of DHE. In currently preferred embodiments, the liquid composition comprises DHE mesylate. In certain embodiments, the liquid composition comprises DHE mesylate at a concentration of 2.5-7.5 mg/ml, 3.5-6.5 mg/ml, or more particularly, 4.0 mg/ml DHE mesylate.

[0011] In some embodiments, the liquid composition further comprises caffeine. In particular embodiments, the liquid composition comprises caffeine at a concentration of 10 mg/ml. In some embodiments, the liquid composition further comprises dextrose, and in certain embodiments, dextrose at a concentration of 50 mg/ml. In specific embodiments, the liquid composition comprises 4.0 mg/ml DHE mesylate, 10.0 mg/ml caffeine, and 50 mg/ml dextrose.

[0012] In various embodiments, following administration of the dose, the mean C_{max} of DHE is at least 1000 pg/ml, or at least 1200 pg/ml. In various embodiments, following administration of the dose, the mean plasma AUC_{0-inf} of DHE is at least 3000 pg*hr/ml, 4000 pg*hr/ml, 5000 pg*hr/ml, or 6000 pg*hr/ml.

[0013] In some embodiments, following administration of the dose, the mean peak plasma concentration (C_{max}) of 8'-OH-DHE is at least 50 pg/ml or at least 55 pg/ml. In some embodiments, following administration of the dose, the mean plasma AUC_{0-inf} of 8'-OH-DHE is at least 1000 pg*hr/ml.

[0014] In typical embodiments, the intranasal delivery device is a manually actuated, propellant-driven, metered-dose intranasal administration device. In some embodiments, prior to first manual actuation, the liquid pharmaceutical composition and propellant are not in contact within the device. In certain embodiments, the liquid pharmaceutical composition is contained in a vial and the propellant is contained in a canister. The canister may further be a pressurized canister. In currently preferred embodiments, between successive manual actuations, the liquid pharmaceutical composition in the vial and propellant in the canister are not in contact within the device.

[0015] In certain of these embodiments, each manual actuation brings a metered volume of liquid pharmaceutical composition and a separately metered volume of propellant into contact within a dose chamber of the device. In specific embodiments, contact of propellant with liquid pharmaceutical composition within the dose chamber of the device creates a spray of liquid pharmaceutical composition as the formulation is expelled through a nozzle of the device. In

particular embodiments, the nozzle has a plurality of lumens, and the spray is ejected simultaneously through a plurality of nozzle lumens. In some embodiments, the propellant is a hydrofluoroalkane propellant, and in specific embodiments, the propellant is hydrofluoroalkane-134a.

[0016] In various embodiments, prior to first actuation, the vial is nonintegral to the device and is configured to be attachable thereto. In some of these embodiments, the vial is configured to be threadably attachable to the device.

[0017] In some embodiments, the subject has migraine headache with aura. In some embodiments, the subject has migraine headache without aura. In some embodiments, the subject has had onset of at least one prodromal symptom of migraine. In a variety of embodiments, the subject has menstrual-associated migraine. In certain embodiments, the subject has triptan-resistant migraine.

[0018] In typical embodiments, the subject self-administers the liquid pharmaceutical composition.

[0019] In a second, related, aspect, improved methods of acutely treating migraine headache with or without aura by intranasal administration of dihydroergotamine (DHE) or salt thereof are provided. In this aspect, the improvement comprises administering a dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration ($C_{\max}$) of at least 750 pg/ml, (b) with a mean time to $C_{\max}$ ($T_{\max}$) of DHE of less than 45 minutes, and (c) a mean plasma $AUC_{0-\infty}$ of DHE of at least 2500 pg*hr/ml.

[0020] In typical embodiments of this aspect, the intranasal delivery device is a manually actuated, metered-dose, propellant-driven intranasal administration device as used in methods of the first aspect. In certain embodiments, contact of propellant with liquid pharmaceutical composition within a dose chamber of the device ejects a spray of liquid pharmaceutical composition through a nozzle of the device. In specific embodiments, the nozzle has a plurality of lumens, and the spray is ejected simultaneously through a plurality of nozzle lumens.

[0021] In another aspect, kits are provided for acutely treating migraine with or without aura. The kits comprise a vial, within which is sealably contained at least one effective dose of a liquid

pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof, and a device, wherein the vial is configured to be attachable to the device, and wherein the device, upon attachment of the vial, is a manually actuated, metered-dose, propellant-driven intranasal administration device capable of providing, after intranasal administration of a dose of liquid pharmaceutical composition, (a) a mean peak plasma DHE concentration ($C_{\max}$) of at least 750 pg/ml, (b) with a mean time to $C_{\max}$ ($T_{\max}$) of DHE of less than 45 minutes, and (c) a mean plasma $AUC_{0-\infty}$ of DHE of at least 2500 pg*hr/ml.

[0022] In some embodiments, the device within the kit comprises a canister, wherein the canister is a pressurized canister containing propellant.

[0023] In certain of these embodiments, following attachment of the vial to the device and prior to first manual actuation, the liquid pharmaceutical composition and propellant are not in contact within the device. In some embodiments, between successive manual actuations, the liquid pharmaceutical composition in the vial and propellant in the canister are not in contact within the device. In typical embodiments, each manual actuation brings a metered volume of liquid pharmaceutical composition and a separately metered volume of propellant into contact within a dose chamber of the device, and contact of propellant with liquid pharmaceutical composition within the dose chamber of the device creates a spray of liquid pharmaceutical composition as the formulation is expelled through a nozzle of the device.

[0024] In some currently preferred embodiments, the liquid pharmaceutical composition within the vial comprises a salt of DHE. In certain embodiments, the liquid composition comprises DHE mesylate. In particular embodiments, the liquid composition comprises DHE mesylate at a concentration of 2.5-7.5 mg/ml, or about 4.0 mg/ml DHE mesylate. In specific embodiments, the liquid composition comprises 4.0 mg/ml DHE mesylate, 10.0 mg/ml caffeine, and 50 mg/ml dextrose.

[0025] In various kit embodiments, the vial contains no more than 2 ml of liquid pharmaceutical composition. In some embodiments, the vial contains approximately 1 ml of liquid pharmaceutical composition.

[0026] In some embodiments, the pressurized canister contains an amount of propellant sufficient to administer no more than 1 dose of liquid pharmaceutical composition.

[0026a] In particular embodiments, provided are:

- use of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof, for acutely treating migraine headache with or without aura, wherein: an effective dose of the liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof is for intranasal administration to a subject with migraine headache, and the dose comprises about 1.45 mg of DHE or salt thereof, and is for administration to the nose as a plurality of divided doses over a period of no more than 1 minute by a manually actuated, propellant-driven metered-dose intranasal delivery device; and

- a kit for acutely treating migraine with or without aura by intranasal administration of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof, comprising: a vial, within which is sealably contained the liquid pharmaceutical composition, and a device, wherein the vial is configured to be attachable to the device, and wherein the device, upon attachment of the vial, is a manually actuated, metered-dose, propellant-driven intranasal administration device that is suitable for administration of an effective dose of the liquid pharmaceutical composition comprising about 1.45 mg DHE or salt thereof to the nose as a plurality of divided doses over a period of no more than 1 minute.

[0027] Other features and advantages of the present disclosure will become apparent from the following detailed description, including the drawings. It should be understood, however, that the detailed description and the specific examples are provided for illustration only, because various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from the detailed description.

4. BRIEF DESCRIPTION OF THE DRAWINGS

[0028] **FIG. 1** shows a cross section of an embodiment of a handheld, manually actuated, metered-dose, propellant-driven intranasal administration device useful for precision olfactory delivery of dihydroergotamine (DHE).

[0029] **FIG. 2** shows a cross section of the in-line nasal delivery device of **FIG. 1** in the stages of rest and actuation. **FIG. 2A** shows the in-line nasal delivery device at rest with **FIG. 2B** showing the actuation of the pump and **FIG. 2C** showing actuation of the propellant valve.

[0030] **FIG. 3** shows a cross section of another implementation of the in-line nasal delivery device.

[0031] **FIG. 4** shows a cross section of the diffuser as seated within the device.

[0032] **FIG. 5A** shows an exploded view of the dose chamber and the Y-junction unassembled.

[0033] **FIG. 5B** shows an exploded view of the dose chamber and Y-junction in cooperation.

[0034] **FIG. 6** shows arrows representing both dose and propellant flow.

[0035] **FIG. 7** shows the actuator grip and conical spring arrangement.

[0036] **FIG. 8** shows a cross section of the optional nose cone and a side elevation of the optional nose cone.

[0037] **FIGS. 9A and 9B** illustrate the device used in the phase I clinical trial described in Example 2, with further description of the numbered parts set forth in Table 1.

[0038] **FIGS. 10A and 10B** plot plasma concentrations of DHE versus time as measured in the phase I comparative bioavailability clinical trial described in Example 2, with **FIG. 10A** plotting data from 0 to 8 hours and **FIG. 10B** plotting data from 0 to 24 hours.

[0039] **FIGS. 11A and 11B** plot plasma concentrations of the 8'-OH-DHE metabolite of DHE versus time as measured in the phase I comparative bioavailability clinical trial described in Example 2, with **FIG. 11A** plotting data from 0 to 8 hours and **FIG. 11B** plotting data from 0 to 24 hours.

[0040] **FIG. 12A** shows a cross section of an alternate implementation of the in-line nasal delivery device.

[0041] **FIG. 12B** shows a zoomed-in view of the cross section of **FIG. 12A**.

[0042] **FIG. 13A** shows a cross section of the diffuser as seated within the device, according to an additional embodiment.

[0043] **FIG. 13B** shows an exploded view of the nozzle and the Y-junction, according to an additional embodiment.

[0044] **FIG. 14** illustrates the nose cone, according to an additional embodiment.

5. DETAILED DESCRIPTION

5.1. Definitions

[0045] Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs.

5.2. Other interpretational conventions

[0046] Ranges: throughout this disclosure, various aspects of the invention are presented in a range format. Ranges include the recited endpoints. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that

range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. This applies regardless of the breadth of the range.

[0047] Unless specifically stated or apparent from context, as used herein the term "or" is understood to be inclusive.

[0048] Unless specifically stated or apparent from context, as used herein, the terms "a", "an", and "the" are understood to be singular or plural. That is, the articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0049] In this disclosure, "comprises," "comprising," "containing," "having," "includes," "including," and linguistic variants thereof have the meaning ascribed to them in U.S. Patent law, permitting the presence of additional components beyond those explicitly recited.

[0050] Unless specifically stated or otherwise apparent from context, as used herein the term "about" is understood as within a range of normal tolerance in the art, for example within 2 standard deviations of the mean and is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$, even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the stated value.

5.3. Summary of experimental observations

[0051] We designed a manually actuated, propellant-driven, intranasal administration device that can reproducibly deliver metered doses of liquid pharmaceutical compositions beyond the nasal valve to more distal regions of the nasal cavity. We tested the device in a Phase I clinical trial designed to compare the bioavailability of (i) dihydroergotamine (DHE) mesylate administered as a single divided 1.45 mg intranasal dose using this Precision Olfactory Delivery (POD™) Device ("INP104"); (ii) a 2.0 mg dose of DHE mesylate administered intranasally using Migranal® Nasal Spray according to the US FDA approved product label; and (iii) a 1.0 mg intravenous injection of DHE mesylate for injection (D.H.E. 45®), in healthy adult subjects.

[0052] As described in detail in Example 2, INP104 provided nearly 3-fold higher mean systemic drug exposure, nearly 4-fold higher mean maximal plasma concentration, and reached maximal DHE plasma concentration faster than Migranal®. The higher systemic drug exposure

and higher maximal plasma concentration were achieved with a lower administered dose of the identical formulation of DHE mesylate, 1.45 mg for INP104 versus 2.0 mg for Migranal[®], and without requiring a 15-minute wait between administration of divided sub-doses, as required for Migranal[®].

[0053] In addition, systemic delivery of DHE was more consistent with INP104 than with Migranal[®], with lower variation observed across subjects for both AUC_{0-inf} and C_{max} parameters.

[0054] Although bolus intravenous administration of 1 mg DHE mesylate provided greater than 10-fold higher C_{max} than 1.45 mg DHE mesylate administered intranasally by INP104, the high C_{max} achieved with intravenous administration is known to be correlated with adverse events (“AE”s), specifically nausea, and IV DHE mesylate is most commonly administered with an anti-emetic. Within 20-30 minutes following administration, DHE plasma concentrations achieved through INP104 intranasal administration were essentially indistinguishable from concentrations achieved by intravenous administration. Thus, despite a greater than 10-fold higher C_{max}, bolus intravenous administration of 1 mg DHE mesylate provided less than 2-fold greater systemic drug delivery, measured as AUC_{0-inf}, as compared to INP104 intranasal delivery.

[0055] The 8'-OH-DHE metabolite of DHE is known to be active, and to contribute to the long-lasting effect of DHE on migraine. We found that intranasal administration of 1.45 mg DHE mesylate by INP104 provides equivalent systemic exposure to the active metabolite of DHE as bolus intravenous administration of 1.0 mg DHE mesylate. In contrast, the 8'-OH DHE metabolite could be detected in only a minority of subjects administered Migranal[®].

5.4. Methods of treating migraine with or without aura

[0056] Accordingly, in a first aspect, methods are provided for acutely treating migraine headache with or without aura.

[0057] The methods comprise administering to a subject with migraine headache an effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or a salt thereof, wherein the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration (C_{max}) of at least

750 pg/ml, (b) with a mean time to $C_{\max}$ ($T_{\max}$) of DHE of less than 45 minutes, and (c) a mean plasma $AUC_{0-\infty}$ of DHE of at least 2500 pg*hr/ml.

5.4.1. Effective dose

[0058] In various embodiments, the dose is no more than 2.0 mg DHE or salt thereof. In typical embodiments, the dose is less than 2.0 mg DHE or DHE salt.

[0059] In certain embodiments, the dose is 1.2-1.8 mg DHE or salt thereof, 1.4-1.6 mg DHE or salt thereof, or 1.4-1.5 mg DHE or salt thereof. In some embodiments, the dose is about 1.2, 1.25, 1.3, 1.35, 1.4, 1.45, 1.5, 1.55, 1.6, 1.65, or 1.7 mg DHE or salt thereof. In a currently preferred embodiment, the dose is about 1.45 mg DHE or salt thereof.

[0060] In some embodiments, the dose is administered as a single undivided dose. In these embodiments, the dose is administered to either the left or right nostril.

[0061] In other embodiments, the dose is administered as a plurality of divided doses. In some of these embodiments, the dose is administered as 2, 3, or 4 divided doses. In particular embodiments, the dose is administered as 2 divided doses. In currently preferred embodiments, the dose is administered as 2 divided doses, with one divided dose administered to each nostril.

[0062] In embodiments in which the dose is administered as a plurality of divided doses, the entire effective dose is typically administered over no more than 1 minute – that is, all of the plurality of divided doses are administered within 1 minute of administration of the first divided dose. In certain divided dose embodiments, the dose is administered over no more than 45 seconds. In certain divided dose embodiments, the dose is administered over no more than 30 seconds.


[0063] In embodiments in which the dose is administered as a plurality of divided doses, the volume of liquid composition administered per divided dose is typically 140-250 μ L. In certain embodiments, the volume of liquid composition administered per divided dose is 145 μ L-225 μ L. In some embodiments, the volume of liquid composition administered per divided dose is 175 μ L-225 μ L. In particular embodiments, the volume of liquid composition administered per divided dose is about ~ 180 μ L or ~ 200 μ L.

5.4.2. Liquid pharmaceutical composition

[0064] The liquid pharmaceutical composition comprises dihydroergotamine (DHE) or salt thereof.

[0065] In typical embodiments, the liquid pharmaceutical composition comprises a salt of DHE. In preferred embodiments, the liquid composition comprises DHE mesylate.

[0066] Dihydroergotamine mesylate – ergotamine hydrogenated in the 9,10 position as the mesylate salt — is known chemically as ergotaman-3', 6', 18-trione, 9,10-dihydro-12'-hydroxy-2'-methyl-5'-(phenylmethyl)-, (5' α)-, monomethane-sulfonate. Its molecular weight is 679.80 and its empirical formula is $C_{33}H_{37}N_5O_5 \cdot CH_4O_3S$. The structure is shown in formula (I) below:



[0067] In typical embodiments, the liquid pharmaceutical composition comprises DHE mesylate at a concentration of at least 1 mg/ml, 1.5 mg/ml, 2.0 mg/ml, 2.5 mg/ml, 3.0 mg/ml, 3.5 mg/ml, 4.0 mg/ml, 4.5 mg/ml or 5.0 mg/ml. In some embodiments, the liquid pharmaceutical composition comprises DHE mesylate at a concentration of 2.5-7.5 mg/ml. In certain embodiments, the liquid pharmaceutical composition comprises 3.0-5.0 mg/ml or 3.5-6.5 mg/ml DHE mesylate. In particular embodiments, the liquid pharmaceutical composition comprises 4.0 mg/ml DHE mesylate.

[0068] In some embodiments, the composition further comprises caffeine. In particular embodiments, the composition comprises caffeine at a concentration of 1 mg/ml-20 mg/ml, 5 mg/ml-15 mg/ml, or 7.5 mg/ml-12.5 mg/ml. In particular embodiments, the composition comprises 10.0 mg/ml caffeine.

[0069] In some embodiments, the composition further comprises dextrose. In certain embodiments, the composition comprises dextrose at a concentration of 5 mg/ml, 10 mg/ml, 15 mg/ml, 20 mg/ml, 25 mg/ml, 30 mg/ml, 35 mg/ml, 40 mg/ml, 45 mg/ml, or 50 mg/ml. In some embodiments, the composition comprises dextrose at a concentration of at least 50 mg/ml.

[0070] In various currently preferred embodiments, the liquid pharmaceutical composition comprises 4.0 mg/ml DHE mesylate, 10.0 mg/ml caffeine, and 50 mg/ml dextrose.

5.4.3. Systemic delivery

[0071] The methods comprise administering to a subject with migraine headache an effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or a salt thereof, wherein the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration ($C_{\max}$) of at least 750 pg/ml, (b) with a mean time to $C_{\max}$ ($T_{\max}$) of DHE of less than 45 minutes, and (c) a mean plasma $AUC_{0-\infty}$ of DHE of at least 2500 pg*hr/ml.

[0072] In various embodiments, the mean peak plasma DHE concentration ($C_{\max}$) achieved following administration of a dose, whether administered as an undivided dose or a plurality of divided doses, is at least 750 pg/ml, 800 pg/ml, 900 pg/ml, 1000 pg/ml, 1100 pg/ml, or 1200 pg/ml. In some embodiments, the mean DHE $C_{\max}$ achieved following administration of a dose is at least 1250, 1300, 1350, 1400, 1450 or 1500 pg/ml. In certain embodiments, the mean DHE $C_{\max}$ achieved following administration of a dose is at least 750 pg/ml, 800 pg/ml, 900 pg/ml, 1000 pg/ml, 1100 pg/ml, or 1200 pg/ml. In certain embodiments, the mean DHE $C_{\max}$ achieved following administration of a dose is at least 1250, 1300, 1350, 1400, 1450 or 1500 pg/ml. In particular embodiments, the mean DHE $C_{\max}$ achieved following administration of a dose is 1000-1500 pg/ml, 1100-1400 pg/ml, or 1200-1300 pg/ml.

[0073] In various embodiments, the mean time to $C_{\max}$ ($T_{\max}$) of DHE following administration is less than 55 minutes. In typical embodiments, the DHE $T_{\max}$ is less than 50 minutes, 45 minutes, 40 minutes, or 35 minutes. In some embodiments, the $T_{\max}$ of DHE following administration is 30-50 minutes, or 35-45 minutes. In particular embodiments, the DHE $T_{\max}$ is no more than 35 minutes, 40 minutes, or 45 minutes.

[0074] In various embodiments, the mean plasma AUC_{0-inf} of DHE following administration is at least 3000 pg*hr/ml, 4000 pg*hr/ml, 5000 pg*hr/ml, or 6000 pg*hr/ml. In various embodiments, the mean plasma AUC_{0-inf} of DHE following administration is at least 7000 pg*hr/ml, 8000 pg*hr/ml, 9000 pg*hr/ml, or 10,000 pg*hr/ml. In some embodiments, the mean plasma AUC_{0-inf} of DHE following administration is at least 5000, 5100, 5200, 5300, 5400, 5500, 5600, 5700, 5800, 5900, or 6000 pg*hr/ml. In some embodiments, the mean plasma AUC_{0-inf} of DHE following administration is greater than 6000, 5900, 5800, 5700, 5600, 5500, 5400, 5300, 5200, 5100 or 5000 pg*hr/ml.

[0075] In various embodiments, following administration of the dose, the mean peak plasma concentration (C_{max}) of 8'-OH-DHE is at least 50 pg/ml. In certain embodiments, the mean C_{max} of 8'-OH-DHE is at least 55 pg/ml.

[0076] In various embodiments, following administration of the dose, the mean plasma AUC_{0-inf} of 8'-OH-DHE is at least 500 pg*hr/ml. In some embodiments, the mean plasma AUC_{0-inf} of 8'-OH-DHE is at least 600 pg*hr/ml, 700 pg*hr/ml, 800 pg*hr/ml, 900 pg*hr/ml, or even at least 1000 pg*hr/ml. In certain embodiments, the mean plasma AUC_{0-inf} of 8'-OH-DHE is at least 1100 pg*hr/ml, 1200 pg*hr/ml, 1250 pg*hr/ml, 1300 pg*hr/ml, 1400 pg*hr/ml, or 1500 pg*hr/ml.

5.4.4. Migraine

[0077] The methods described herein are used to acutely treat migraine headache, with or without aura.

[0078] In various embodiments, the subject has had onset of at least one prodromal symptom of migraine, without onset of headache pain. In certain embodiments, the subject has had onset of at least one prodromal symptom selected from neck stiffness, facial paresthesia, photosensitivity, acoustic sensitivity, and visual aura.

[0079] In various embodiments, the subject has had onset of at least one symptom associated with acute migraine. In certain embodiments, the subject has had onset of at least one symptom selected from visual aura; headache pain, including dull, throbbing, or pulsing pain; photosensitivity; acoustic sensitivity; nausea; vomiting. Visual aura and headache pain may be unilateral or bilateral, focal or diffuse.

[0080] In various embodiments, administration is performed within 5 minutes, 10 minutes, 15 minutes, or 30 minutes of onset of at least one prodromal symptom. In various embodiments, administration is performed within 5 minutes, 10 minutes, 15 minutes, or 30 minutes of onset of at least one acute symptom.

[0081] In typical embodiments, the subject performs the administration (self-administration). In some embodiments, the administration is performed by another individual, such as a parent, guardian, caregiver, or medical professional.

[0082] In some embodiments, migraine to be treated is associated with menstruation. In some embodiments, migraine to be treated has proven resistant to triptans.

[0083] In various embodiments, the methods are used for acute treatment of cluster headaches rather than migraine.

5.4.5. Device

[0084] In the methods described herein, the dose is administered by an intranasal delivery device that provides, following intranasal administration, (a) a mean peak plasma DHE concentration (C_{max}) of at least 750 pg/ml, (b) with a mean time to C_{max} (T_{max}) of DHE of less than 45 minutes, and (c) a mean plasma AUC_{0-inf} of DHE of at least 2500 pg*hr/ml.

5.4.5.1. Compound delivery device

[0085] In various embodiments, the intranasal administration device is a “compound delivery device” as described in U.S. Pat. No. 9,550,036.

5.4.5.2. Medical unit dose container device

[0086] In various embodiments, the intranasal administration device is a “medical unit dose container” device as described in WO 2014/179228.

5.4.5.3. Manually activated, propellant-driven, metered-dose device

[0087] In typical embodiments, the intranasal delivery device is a manually actuated, propellant-driven, metered-dose intranasal administration device.

[0088] In currently preferred embodiments, the liquid pharmaceutical composition and propellant are not in contact within the device prior to first manual actuation, and, optionally, not in contact within the device between successive manual actuations. In such embodiments, the device typically comprises a vial and a canister, wherein the liquid pharmaceutical composition is contained in the vial and the propellant is contained in the canister. Typically, the canister is a pressurized canister of propellant. In typical embodiments, the propellant is a hydrofluoroalkane propellant suitable for pharmaceutical use. In specific embodiments, the propellant is hydrofluoroalkane-134a.

[0089] In various embodiments, each manual actuation brings a metered volume of liquid pharmaceutical composition and a separately metered volume of propellant into contact within a dose chamber of the device. Contact of propellant with liquid pharmaceutical composition within the dose chamber of the device propels the dose towards the nozzle of the device, creating a spray as the dose is expelled through the nozzle of the device. In particularly preferred embodiments, the nozzle has a plurality of lumens, and the spray is ejected simultaneously through a plurality of nozzle lumens.

[0090] As discussed in further detail below with respect to kits, in some embodiments the vial is nonintegral to the device and is configured to be attachable thereto. In particular embodiments, the vial is configured to be threadably attachable to the device.

5.4.5.3.1. In-line nasal delivery device

[0091] In certain currently preferred embodiments, the manually actuated, propellant-driven metered-dose intranasal administration device is an “in-line nasal delivery device” as described in WO 2017/044897.

[0092] Typically, in these embodiments the device delivers at least a portion of the dose of liquid pharmaceutical composition to the nasal cavity beyond the nasal valve, including delivery to the turbinates and/or the olfactory region. In certain embodiments, the device delivers at least 25%,

30%, 40%, 50%, 60%, or 70% of the dose of liquid pharmaceutical composition beyond the nasal valve. In certain embodiments, the device delivers liquid pharmaceutical composition so that at least 25%, 30%, 40%, 50%, 60%, or 70% of the dose of liquid pharmaceutical composition is brought into contact with the upper third of the nasal cavity (nasal epithelium) of the subject.

[0093] As shown in FIG. 1, the in-line nasal delivery device 1 includes a housing 10, diffuser 20, tip 35, nozzle 40, dose chamber 45, an actuator 50, and a pump 25 to move the liquid pharmaceutical composition into the dose chamber 45. In one series of embodiments, the in-line nasal device 1 is associated and cooperative with a propellant canister 5, a propellant valve 15, and a vial 30 of liquid pharmaceutical composition cooperative with the pump 25 to move the liquid pharmaceutical composition into the dose chamber 45.

[0094] In one series of embodiments, the diffuser 20 is a frit 21 (not shown in FIG. 1). The diffuser provides for the conversion of the liquefied propellant in the propellant canister 5 to gas and/or an increase in temperature of the propellant.

[0095] In one series of embodiments, the propellant valve 15 is a metered dose propellant valve 16.

[0096] In one series of embodiments, the liquid pharmaceutical composition is supplied in the form of a sealed vial 30, e.g., of glass. In one series of embodiments, the vial 30 has a neck 31 (not shown) that is sealed by a removable closure 32 (not shown), for example but not limited to sealed with a plastic cover, crimped metal seal, and rubber stopper (for stability and sterility purposes). When the closure 32 is removed, the device 1 can be engaged with the vial 30. In one series of embodiments, device 1 can be engaged with vial 30 by cooperation with the neck 31 of the vial 30. In a related aspect, further discussed below, sealed vial 30 and device 1 can be co-packaged into a kit to be assembled at time of use.

[0097] In certain embodiments, vial 30 is a 3.5-mL amber glass vial.

[0098] A pump 25 moves the liquid pharmaceutical composition into the dose chamber 45.

[0099] The propellant canister 5 is a canister of a compressed gas or a liquefied propellant. Compressed gases include but are not limited to compressed air and compressed hydrocarbons. In one series of embodiments, the compressed gas is nitrogen or carbon dioxide. Liquefied

propellants include but are not limited to chlorofluorocarbons and hydrofluoroalkanes. In a preferred embodiment, propellant canister 5 contains HFA-134a.

[0100] The canister 5 will generally be provided with a propellant valve 15 by which the gas flow can be controlled.

[0101] The tip 35 includes a nozzle 40. In one series of embodiments, the nozzle 40 has a plurality of nozzle openings 41 (not shown) (synonymously, nozzle lumens). Through the plurality of nozzle openings 41, the liquid pharmaceutical composition and propellant is delivered to the nasal cavity.

[0102] Actuation of the propellant canister 5 is effectively coordinated with actuation of the pump 25 for the vial 30 for the liquid pharmaceutical composition. The arrangement may be such that actuation of the vial 30 for the liquid pharmaceutical composition causes actuation of the propellant canister 5. FIG. 2 shows the device 1 at rest (FIG. 2A) and in actuation (FIG. 2B and 2C).

[0103] As an example, the staging of the device 1 actuation is as follows. The housing 10 is compressed to prime the propellant canister 5. When the housing 10 is compressed, an actuator 50 remains stationary in the housing 10 while the propellant canister 5 and the vial 30 move towards the actuator 50. At this time, the propellant valve 15 associated with the propellant canister 5 is not actuated by compression. The actuator 50 acts upon the pump 25 compressing the pump 25 and the liquid pharmaceutical composition from the vial 30 is moved into the dose chamber 45. After a majority of the liquid pharmaceutical composition has moved into the dose chamber 45, the actuator 50 acts upon the propellant valve 15 and the propellant valve 15 begins to compress. The continued depression of the actuator 50 releases the propellant from the propellant canister 5. The propellant pushes the liquid pharmaceutical composition as it exits the device 1 through the nozzle openings (lumens) 41 (not shown) of the nozzle 40 located in the tip 35. The actuator 50 provides for first actuation of the pump 25, then once the pump 25 bottoms out, the continued depression of the actuator 50 provides for release of the propellant from the canister 5.

[0104] In an alternative implementation of the device 1 (not shown), the device 1 does not include a diffuser 20. In such embodiments, the device typically incorporates another type of dose retaining valve.

[0105] FIG. 3 shows yet another implementation of the device 100. The device 100 can deliver a single or multiple dose from a vial 30 or other container. The device 100 allows for multiple doses to be delivered from the vial 30, or a single dose. For example, the vial 30 may contain a volume of liquid pharmaceutical composition for multiple doses, while the user may decide to only deliver a single dose from the vial 30. The liquid pharmaceutical composition may be a drug, active pharmaceutical ingredient, or a pharmaceutical formulation.

[0106] Initially, the vial 30 may be separate from the rest of the assembled device 100. At the time of use, the device 100 and vial 30 are taken out of their respective packaging. Prior to use, the vial 30 will generally be sealed. In the embodiment where the vial 30 is covered by a plastic cover, metal seal and stopper, the plastic cover and metal seal are pulled away from the top of the vial 30, and the rubber stopper is removed from the vial 30. The vial 30 may be screwed into a pump fitment 180 located at the base of the device 100. For example, but not limitation, the vial 30 may have female threads which can be screwed into male threads on a pump fitment 180, or vice versa. The vial 30 may contain, for example but not limited to, inclusive of end points, 2-3 ml, in another embodiment 2-2.5 ml of liquid pharmaceutical composition.

[0107] As shown in FIG. 3, the device 100 includes a housing 110. The housing 110 contains components of the device 100 including the Y-junction 120. The Y-junction 120 has three branches radiating from a common base. The Y-junction and its three branches may be a molded component. The Y-junction 120 establishes both fluid and gas paths within the device 100, and connects the metered dose pump 130, the dose chamber 150, and the propellant canister 140 when the propellant canister 140 is assembled with the device.

[0108] As shown in FIG. 3, for use of the device 100, the user will generally orient the device 100 with the propellant canister 140 assembled and located at the top and the vial 30 assembled and located at the bottom. Housed within the device's 100 housing 110, the optional check-valve 160 (attached to the metered dose pump 130 stem) press fits into a receiving hub of a first branch of the Y-junction 120. An internal bore provides fluid communication from the metered dose pump 130, through the optional check-valve 160 and to a third branch of the Y-junction 120, which connects to the dose chamber 150. In one series of embodiments, the check valve 160 is an elastomeric component that installs within a plastic housing between the metered dose pump 130 and the Y-junction 120. The optional check valve 160: (a) reduces or eliminates dose

leakage which could occur through the metered dose pump 130 if the pump stem was depressed and the propellant canister 140 was actuated; (b) allows for improved consistency in dose delivery by the device 100; and/or provides that liquid pharmaceutical composition is not pushed back down the internal dose loading channel 230 of the Y-junction 120 and into the metered dose pump 130.

[0109] When oriented as to be used in operation, housed within the device's 100 housing 110, towards the top of the device 100, the propellant canister 140 press fits into a second branch of the Y-junction 120, establishing the gas path through internal bores, through the diffuser 170 and to the dose chamber 150.

[0110] In this implementation of the device 100, the diffuser 170 is annular. As shown in FIG. 4, the annular diffuser 170 sits inside a bore on the back end of the dose chamber 150. The external diameter of the annular diffuser 170 is in a compression fit with the dose chamber 150. In other embodiments, not shown, the annular diffuser is fixed to the dose chamber using means other to or in addition to compression fit.

[0111] An internal dose loading channel 230 which is molded as a portion of the Y-junction 120 fits into the inner bore of the annular diffuser 170 when the dose chamber 150 is installed onto the Y-junction 120. The inner diameter of the annular diffuser 170 is in compression with the internal dose loading channel 230 portion of the Y-junction 120. The annular diffuser 170 is seated between the outer wall of the internal dose loading channel 230 and the inner wall of the dose chamber 150, sealing against both of those surfaces to form the bottom of the dose chamber 150. Additional embodiments of the diffuser 170, dose chamber 150, and Y-junction 120 are discussed with regards to FIGS. 12-13.

[0112] In one series of embodiments, the diffuser 170 is a frit 171 (not shown). In other embodiments, the diffuser 170 is a component that is homogeneously or heterogeneously porous. In some embodiments, the diffuser 170 may be a disk-shaped member. The diffuser 170: (a) provides for the conversion of the liquefied propellant in the propellant canister 140 to gas; (b) provides an increase in temperature of the propellant; (c) acts to prevent the propellant from flowing back into the device 100; (d) acts to prevent the liquid pharmaceutical composition from flowing back into the device 100; and/or (e) acts to allow gas flow into the dose chamber 150

while preventing the liquid pharmaceutical composition from leaking out. The diffuser may be made of a porous polymer material.

[0113] The relationship in operation of the device 100 between the liquid pharmaceutical composition, the diffuser 170, the inner dose loading tube 230, the dose chamber 150 and the Y-junction 120 are shown at least in FIG. 6. In operation, the liquid pharmaceutical composition being loaded into the dose chamber 150 takes the less restrictive route, flowing out of the vial 30 and filling the dose chamber 150 rather than loading backwards through the diffuser 170 and into the delivery path of the propellant of the Y-junction 120. In operation of the device 100, the staging of operation and the amount of time required for operation of the device allows the diffuser 170 to restrict liquid pharmaceutical composition from flowing back into the Y-junction 120 for the period of time needed, as the propellant canister 140 is activated after liquid pharmaceutical composition loading. During proper device 100 use, the entire actuation of the device 100, including metered dose pump 130 and propellant canister 140, is approximately a second or less than a second. The loaded dose in the dose chamber 150 does not have enough time to flow backwards into the Y-junction 120. Immediately after the dose chamber 150 is full, the propellant expels the liquid pharmaceutical composition from the device 100.

[0114] On the third leg of the Y-junction 120 at a 45-degree angle, the dose chamber 150 press fits into the Y-junction 120, completing the flow paths for both gas and fluid through the device. In one series of embodiments, the angle is 30 degrees, 35 degrees, 40 degrees, 45 degrees, 50 degrees, 55 degrees, 60 degrees, inclusive of endpoints and intervening degrees.

[0115] The Y-junction 120 may contain engagement ribs (not shown) to help secure and position the assembly within the housing 110 of the device 100.

[0116] The device 100 includes a pump fitment 180. The pump fitment 180 secures the metered dose pump 130 to the vial 30 and holds both components in place during device 100 use. One series of embodiments of the pump fitment 180 is that it consists of engagement ribs that retain it within the housing 110, provide vertical displacement, and prevent rotation during installation of the vial 30.

[0117] The device 100 includes a dose chamber 150. The dose chamber 150 receives and stores the liquid pharmaceutical composition that has been pushed out of the inner tube of the Y-junction 120. When the propellant canister 140 is actuated, the Y-junction 120 and dose chamber

150 are pressurized and the propellant gas expels the liquid pharmaceutical composition out of the dose chamber 150. As shown in FIGS. 5A and 5B, the dose chamber 150 is press fit into the Y-junction 120. The nozzle 190 is installed into the end of the dose chamber 150 opposite where it is press fit into the Y-junction 120.

[0118] The nozzle 190 is installed into the distal end (end opposite where the dose chamber 150 is press fit into the Y-junction 120) of the dose chamber 150, forming a liquid and gas-tight seal around the outer diameter. During actuation of the device 100, propellant evacuates liquid pharmaceutical composition from the dose chamber 150, pushing it out the nozzle 190.

[0119] The nozzle 190 forms the narrow plume angle (for example, an angle of 1 to 40 degrees, including endpoints and angles intermittent there between; in one series of embodiments the angle is 5 degrees, 10 degrees, 15 degrees, 20 degrees, 25 degrees, 30 degrees, 35 degrees) multi-stream deposition. The nozzle 190 and resultant angle of the plume produced promotes delivery of the liquid pharmaceutical composition to the olfactory region of the user's nasal cavity.

[0120] In this implementation, as shown in FIG. 8, the device 100 may include an optional nose cone 200. The external geometries of the nose cone 200 assist in providing proper alignment of the device 100 during insertion into the nose. The diametrically opposed flat sides aid with placement against the septum of either naris, with the depth stop providing correct depth of insertion. The nose cone 200 adds redundancy to nozzle 190 retention through mechanical interference incorporated into the design. As shown in FIG. 3 and FIG. 8, there is an opening in the nose cone 200 which aligns with the nozzle 190. The nose cone 200 is not part of the pressurized flow path.

[0121] The housing 110 represents the body of the device 100. The housing 110 includes two different "clamshells" concealing the components of the device 100 and retaining all components to ensure functionality. The housing 110 houses the metered dose pump 130 and pump fitment 180, the actuator grip 210, the Y-junction 120, the propellant canister 140, and the dose chamber 150. The nose cone 200 engages onto the outer geometry of the housing 110, or may be optionally integrated into the design of the clamshells. An additional embodiment of the nose cone 200 is discussed with regards to FIG. 14. The housing 110 is designed to assemble easily through the use of, for example but not limited to, mattel pins, snaps, post or screws, or a combination thereof, molded into the geometry.

[0122] The actuator grip 210 provides for actuation displacement by the user. The actuator grip 210 is composed of two parts, actuator grip A and actuator grip B and surround the Y-junction 120 and reside within the housing 110. FIG. 7 shows two finger grip notches 215 are designed into the actuator grip 210 to allow the user to engage the device 100 with the fingers, for example but not limited to, the index and middle finger. These finger grip notches 215 allow the user to apply downward movement leading to device 100 actuation.

[0123] The metered dose pump 130 draws liquid pharmaceutical composition up from the vial 30 to the Y-junction 120. The metered dose pump 130 may utilize a custom pump fitment 180 to promote functionality within the device 100, and allow attachment of the vial 30 via threads. The metered dose pump 130 may deliver, for example but not limited to, volumes of 130 μ l, 140 μ l, 150 μ l, 160 μ l, 170 μ l, 180 μ l, 190 μ l, 200 μ l, or 230 μ l during actuation. Commercially available metered dose pumps 130 can be used.

[0124] For the device 100 to consistently deliver liquid pharmaceutical composition, the metered dose pump 130 must first deliver liquid pharmaceutical composition, followed by propellant canister 140 actuation to expel the liquid pharmaceutical composition. As shown in FIG. 7, one manner in which to accomplish this is via a conical spring 220 between the propellant canister 140 and Y-junction 120 to create the necessary propellant canister 140 actuation force resulting in the correct order of actuation between the metered dose pump 130 and propellant canister 140. In one implementation, a conical spring 220 is used, although this force is not limited to being produced by a conical spring 220 as other mechanisms can be used. In one series of embodiments, the conical spring 220 has a near zero preload, with a k value of about 25.5 lbf in and a maximum load of 3.2lbf. Selection of the spring or mechanism will include the considerations of: (a) providing for proper device 100 staging; (b) physical space in the device 100; and/or (c) and user feedback regarding how stiff of a conical spring 220 still allows a variety of users to activate the device 100.

[0125] The conical spring 220 is installed inline between the propellant canister 140 and Y-junction 120. The actuator grip 210 physically holds the propellant canister 140. The user activates the device 100 by, for example, applying an in-line force acting down from the actuator grips 210, and up from the vial 30. This force simultaneously acts to activate both the metered dose pump 130 and the propellant canister 140. The conical spring 220 acts in parallel to the

internal propellant canister metering valve spring, increasing the necessary force required to activate the propellant canister 140. By choosing the conical spring 220 such that the necessary force required to actuate the propellant canister 140 is in excess of the maximum necessary force required to completely actuate the metered dose pump 130, the device 100 provides that dose is loaded into the dose chamber 150 before propellant gas begins to expel liquid pharmaceutical composition from the device 100.

[0126] In another embodiment, an extension spring is used in lieu of a conical spring. The extension spring is discussed with regards to FIG. 12A.

[0127] During device 100 actuation, the metered dose pump 130 draws liquid pharmaceutical composition up from the vial 30 at the bottom of the device 100 via the Y-junction 120, through the internal dose loading channel 230 and into the dose chamber 150. The internal dose loading channel 230 provides a clear route for the liquid pharmaceutical composition to be loaded ahead of the diffuser 170, without needed to physically pass through the porous material of the diffuser 170. As shown in FIG. 6, small arrow heads represent the flow of the propellant while large arrow heads represent the flow of the liquid pharmaceutical composition. Priming shots may be required to completely fill the metered dose pump 130 and internal dose loading channel 230 of the Y-junction 120 prior to user dosing. An optional dose cap (not shown) may cover the nose cone 200 of the device 100 and captures the priming shots while also providing a means of visual indication to the user that the device is primed.

[0128] In the second stage of device 100 actuation, once the dose chamber 150 has been filled, the propellant canister 140 releases propellant which enters through the top of the Y-junction 120, following the path shown by smaller arrow heads in FIG. 6. The propellant flows physically through the porous material of the diffuser 170, which promotes the vaporization of the propellant. The diffuser 170 and the path along which the propellant travels (shown by the arrow heads in FIG. 6) convert liquid propellant into gas propellant, resulting in expansion and propulsion of the propellant. The propellant first contacts the liquid pharmaceutical composition at the proximal (distal being closer to the nozzle 190, proximal being farther away from the nozzle 190) face of the diffuser 170 as seated in the device 100. As the propellant continues to expand, it pushes the liquid pharmaceutical composition forward (toward the nozzle 190) in the dose chamber 150, exiting though the nozzle 190 at the end of the dose chamber 150.

[0129] The propellant canister 140 provides the propulsive energy for the device 100. The stem of the propellant valve seats into the top receiver of the Y-junction 120. During use, the user presses down on the actuator grips 210 which pulls the propellant canister 140 body down, actuating the propellant valve. This releases a metered volume of liquid propellant. As the propellant vaporizes and expands, the liquid pharmaceutical composition is forced toward the distal end of dose chamber 150 and out through the nozzle 190.

[0130] As a non-limiting example of propellant, the propellant canister 140 uses HFA 134A as the propellant for the system. Other propellants are envisioned. There are commercially available propellant canisters 140.

[0131] In certain embodiments, the device, propellant canister, and vial containing liquid pharmaceutical composition are provided separately, optionally co-packaged into a kit, and thereafter assembled for use. In certain embodiments, propellant canister 140 is provided assembled within device 100 and the vial containing liquid pharmaceutical composition is provided separately, optionally with the device (with integrated canister) and vial co-packaged into a kit. In some embodiments, the device, propellant canister, and vial containing liquid pharmaceutical composition are provided to the user fully assembled.

5.4.5.3.2. Alternate In-line nasal delivery device

[0132] In certain embodiments, the device comprises the following parts; part numbering is as depicted in FIGS. 9A and 9B.

Table 1 Clinical Trial Device			
COMPONENT	PART ID	PART NAME	MATERIAL
Device	1	Y-Junction	PP
	2	Diffuser	PE
	3	Dose Chamber	PP

Table 1 Clinical Trial Device			
COMPONENT	PART ID	PART NAME	MATERIAL
	4	Metering Pump	POM; PE Medium Density; Chlorobutyl Rubber PP; White Masterbatch Colorant Stainless Steel; PE (HDPE+LDPE)
	5	Finger Grip (right)	ABS
	6	Clamshell (right)	ABS
	7	Clamshell (left)	ABS
	8	Propellant Canister	Propellant: HFA
			Canister: Anodized Aluminum
			HFA Metering Valve: Anodized Aluminum; Polyester; Stainless Steel; EF327 Seat and Gasket
	9	Nozzle	LCP
	10	Check Valve	Silicone
	11	Check Valve Adapter	PP
	12	Finger Grip (left)	ABS
	13	Extension Spring	Stainless Steel
	14	Nose Cone	ABS
Drug	15	Drug Vial	3.5 ml amber glass vial container

Abbreviations

ABS = acrylonitrile butadiene styrene; CMO = contract manufacturing organization;
 HDPE = high density polyethylene; HFA = hydrofluoroalkane-134a; LCP = liquid crystal polymer; LDPE = low density polyethylene; PE = polyethylene; POM = polyacetal copolymer; PP = polypropylene

[0133] The vial contains liquid pharmaceutical composition in an amount sufficient for at least one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses. In particular embodiments, the vial contains liquid pharmaceutical composition in an amount sufficient for at most one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses.

[0134] In various embodiments, the propellant canister contains pressurized propellant in an amount sufficient for optional priming of the device followed by delivery of at least one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses. In particular embodiments, the propellant canister contains pressurized propellant in an amount sufficient for optional priming of the device followed by delivery of at most one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses.

[0135] In some embodiments, with each actuation, a minority of the pressurized liquid hydrofluoroalkane is converted to gaseous hydrofluoroalkane. In certain embodiments, the quantity of pressurized liquid hydrofluoroalkane is sufficient to permit a predetermined number of device actuations. In some of these embodiments, the quantity is sufficient to permit 2, 3, 4, 5, 6, 7 or 8 actuations. In some embodiments, the quantity is sufficient to permit 10, 11, 12, 13, 14, 15, or even 20 actuations. In certain embodiments, a majority of the pressurized liquid hydrofluoroalkane is converted to gaseous hydrofluoroalkanes after 2, 3, 4, 5, 6, 7, or 8 actuations. In certain embodiments, a majority of the pressurized liquid hydrofluoroalkane is converted to gaseous hydrofluoroalkanes after 10, 11, 12, 13, 14, 15, or 20 actuations.

5.4.5.3.3. Alternate In-line nasal delivery device

[0136] FIG. 12A shows a cross section of an alternate implementation of the in-line nasal delivery device 1200. The in-line nasal delivery device 1200 may be an embodiment of the in-line nasal delivery device 100. For example, the device 1200 may use the same or similar components as the device 100, as described with regards to FIGS. 3-9. Additionally, components of device 1200 and device 100 may be used interchangeably or in some combination thereof. In the embodiment of FIG. 12A, the device 1200 includes a housing 12110, a Y-junction 12120, a metered dose pump 12130, a propellant canister 12140, a dose chamber 12150 (shown in FIG.

13A), a check valve 12160, a diffuser 12170 (shown in FIG. 13A), a pump fitment 12180, a nozzle (not shown), a nose cone 12200, and an actuator grip 12210. The housing 12110 includes an upper portion 1205 and a bottom portion 1210. The device 1200 additionally includes an extension spring 1215 and a check valve adapter 1220.

[0137] Similar to the actuator grip 210 described with regards to FIG. 3, the actuator grip 12210 provides for actuation displacement by the user. The actuator grip 12210 surrounds the Y-junction 12120 and resides within the housing 12110. FIG. 12A shows two finger grip notches 12215 that are designed into the actuator grip 12210 to allow the user to engage the device 1200 with the fingers, for example but not limited to, the index and middle finger. The finger grip notches 12215 allow the user to engage or grip the device in order to cause device 1200 actuation.

[0138] More specifically, the actuator grip 12210 includes a guiding feature 1225 that extends along a length of the housing 12110 behind (as illustrated in FIG. 12A) the propellant canister 12140 and captures an end of the propellant canister 12140. In the illustrated example, the end is the bottom of the propellant canister 12140, which is opposite from the end containing the valve for propellant dispersal. The guiding feature 1225 may capture the end of the propellant canister 12140 by folding above or adhering to the end. The propellant canister 12140 is nested within the guiding feature 1225 such that the guiding feature 1225 securely supports the propellant canister 12140. By enveloping a portion of the propellant canister 12140, the guiding feature 1225 is securely coupled to a larger, more rigid surface area of the propellant canister 12140 than when coupled to a narrow surface, such as the propellant valve 15 in the embodiment of device 1. In this configuration, as the user applies downward movement via the finger grip notches 12215 to actuate the device 1200, the guiding feature 1225 transmits the downward force to the propellant canister 12140, thereby actuating the propellant canister 12140. The guiding feature 1225 actuates the propellant canister 12140 in a stable manner and is less likely to lose its physical coupling to the propellant canister 12140.

[0139] In one embodiment, the propellant canister 12140 is entirely enclosed within the housing 12110. In one specific embodiment, the propellant canister 12140 is enclosed by the upper portion of the housing 1205, which may be formed during manufacturing from at least two separate parts. The Y-junction 12120 is fixed in place with the bottom housing portion 1210,

with the guiding feature 1225 extending upward to establish the position of the propellant canister 12140 with respect to the Y-junction 12120. This structure ensures that the propellant canister 12140 moves relative to the Y-junction 12120 during actuation, to which it is fluidly coupled.

[0140] In a similar manner to the conical spring 220 described with regards to FIG. 7, the extension spring 1215 creates an actuation force that ensures a desired order of actuation between the metered dose pump 12130 and the propellant canister 12140. Specifically, during device actuation, the metered dose pump 12130 first delivers liquid pharmaceutical composition to the dose chamber 12150, followed by propellant canister 12140 actuation to expel the liquid pharmaceutical composition. The force of the extension spring 1215 is established to both provide proper order of actuation and enable ease of actuation by users.

[0141] The extension spring 1215 is coupled to the housing upper portion 1205 and the actuator grip 12210. As illustrated in FIG. 12A, a first end of the extension spring 1215 couples to a boss 1230 on the housing upper portion 1205, and a second end of the extension spring 1215 couples to a boss 1235 on the actuator grip 12210. In the embodiment of FIG. 12A, the housing upper portion 1205 and the actuator grip 12210 translate relative to one another during actuation of the device 1200. The extension spring 1215 is coupled to each component such that the extension spring 1215 creates a resisting force when the housing upper portion 1205 and the actuator grip 12210 translate away from each other. As previously described, the user activates the device 1200 by, for example, applying an in-line force acting down from the actuator grips 12210, and up from the vial containing the pharmaceutical composition. This applied force actuates both the metered dose pump 12130 of the vial and the propellant canister 12140. As the applied force on the extension spring 1215 increases, a threshold (higher) force to actuate the propellant canister 12140 is achieved after a threshold (lower) force to actuate the metered dose pump 12130 is achieved, such that the applied force first exceeds the threshold force of the metered dose pump 12130. In this configuration, actuation of the device 1200 first activates the metered dose pump 12130 and then activates the propellant canister 12140 such that dose is loaded into the dose chamber 12150 before propellant begins to expel liquid pharmaceutical composition from the device 1200.

[0142] In some embodiments, the extension spring 1215 may be used in lieu of or in addition to the conical spring 220. The configuration of the extension spring may streamline the assembly process of the device relative to the configuration of the conical spring, as the conical spring may create a resisting force between the propellant canister 140 and Y-junction 120 such that the components are pushed apart during assembly, whereas the extension spring may pull the components towards each other. In addition, the configuration of the extension spring may prolong the shelf life and overall lifetime of the device relative to the configuration of the conical spring. This may be in part due to the press fit between the stem of the propellant canister 140 and Y-junction 120 of the device 100, which may naturally relax over time and which may be propagated by the resisting force of the conical spring between the propellant canister 140 and Y-junction 120, potentially furthering the decrease in durability of the press fit over time.

[0143] The check valve adapter 1220 is an adapter that couples the check valve 12160 and the Y-junction 12120. The check valve 12160 may be an embodiment of check valve 160. In the embodiment of FIGS. 12A-12B, the check valve adapter 1220 is a cylindrical component having a first end that inserts into a channel of the Y-junction 12120 and mates with the check valve 12160 positioned within the channel of the Y-junction 12120 and a second end that mates with the metered dose pump 130. As illustrated in the zoomed-in view in FIG. 12B, an end of the check valve 12160 comprises a flange that is captured at an end of the channel of the Y-junction 12120 and mates with a respective interface of the check valve adapter 1220. The check valve 12160 and/or check valve adapter 1220 may be secured at each end with an adhesive, ultrasonic welding, an interference fit (e.g., press fit, friction fit, or similar), or some combination thereof. The check valve adapter 1220 may augment the function of the check valve 12160 by improving the seal between the check valve 12160 and the Y-junction 12120. As discussed with regards to FIG. 3, a check valve may: (a) reduce or eliminate dose leakage which could occur through the metered dose pump if the pump stem was depressed and the propellant canister was actuated; (b) allow for improved consistency in dose delivery by the device; and/or (c) provide that liquid pharmaceutical composition is not pushed back down an internal dose loading channel of the Y-junction and into the metered dose pump.

[0144] FIG. 13A shows a cross section of a diffuser 12170 as seated within the device 1200, according to an additional embodiment. The diffuser 12170 may be an embodiment of the diffuser 170. In this implementation of the device 1200, the diffuser 12170 is annular. As shown

in FIG. 13A, the diffuser 12170 sits on a shelf 1305 inside a bore 1310 of the Y-junction 12120, and the dose chamber 12150 is inserted into the bore 1310 of the Y-junction 12120. The diffuser 12170 is seated between the shelf of the bore of the Y-junction 12120 and a bottom face of the dose chamber 12150, sealing against both of those surfaces. The diffuser 12170 may further be sealed along its inner diameter to the Y-junction 12120. In this configuration, the diffuser 12170 creates an interference seal along its inner diameter, its upper face, and its lower outer edge (in contact with the shelf 1305). This configuration may allow expansion of the diffuser 12170, for example, as propellant flows through the diffuser 12170 due to changes in temperature or as a result of device assembly. Sealing the diffuser 12170 along its inner diameter may improve the consistency and/or quality of the seal and/or performance of the diffuser 12170 relative to sealing the diffuser 12170 along its top and bottom faces in a compression fit, which could compress the diffusion path within (the path along which propellant travels and is diffused). In this configuration, variations in the manufacturing of the diffuser 12170 may be less likely to affect the performance of the diffuser 12170. For example, the tolerances of the outer diameter of the diffuser 12170 may not need to be as precisely controlled to prevent bending of the diffuser 12170 such that flatness of the diffuser 12170 is maintained to ensure a proper compression fit along its faces. In some instances, the interference seal may or may not be liquid or gas tight.

[0145] FIG. 13B shows an exploded view of the dose chamber 12150 and the Y-junction 12120, according to an additional embodiment. FIG. 13B illustrates the bore 1310 and the shelf 1305 of the Y-junction 12120. The dose chamber 12150 may include a chamfer 1315 around an outer edge of its bottom face such that the dose chamber 12150 may be easily inserted into the bore 1310. In alternate embodiments, the configuration of the dose chamber 12150 and Y-junction 12120 may be reversed such that the dose chamber 12150 includes a bore into which a diffuser and an end of the Y-junction 12120 is inserted.

[0146] FIG. 14 illustrates the nose cone 12200, according to an additional embodiment. The nose cone 12200 may be an embodiment of the nose cone 200. As previously described, the external geometries of the nose cone 12200 assist in providing proper alignment of the device 1200 during insertion into the nose. As shown in FIG. 14, the nose cone 12200 comprises an opening 1405 that aligns with the nozzle (not shown). The dose chamber 12150 (not shown in this view) may be positioned between two bosses 1410a, 1410b that maintain the alignment of the dose chamber 12150 and the nozzle within the nose cone 12200. In the embodiment of FIG. 14, the

nose cone 12200 is integrated into the design of the clamshells. The nose cone 12200 and the clamshells may be molded together during manufacturing, decreasing the overall part count of the device 1200 and enabling easy assembly of the device 1200.

5.5. Kits

[0147] In another aspect, kits are provided for acutely treating migraine with or without aura.

[0148] The kit comprises a vial and a device. The vial is sealed, and sealably contains at least one effective dose of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof. The vial is configured to be attachable to the device. The device is reciprocally configured to receive the vial. Upon attachment of the vial to the device by the user, the device becomes a manually actuated, propellant-driven, metered-dose intranasal administration device capable of providing, after intranasal administration of a dose of liquid pharmaceutical composition, (a) a mean peak plasma DHE concentration ($C_{\max}$) of at least 750 pg/ml, (b) with a mean time to $C_{\max}$ ($T_{\max}$) of DHE of less than 45 minutes, and (c) a mean plasma $AUC_{0-\infty}$ of DHE of at least 2500 pg*hr/ml.

[0149] In typical embodiments, upon attachment of the vial to the device, the device becomes a manually actuated, propellant-driven, metered-dose intranasal administration device as described in Section 5.3.5.3 above. In currently preferred embodiments, upon attachment of the vial to the device, the device becomes a manually actuated, propellant-driven, metered-dose intranasal administration device as particularly described in Section 5.3.5.3.1 above. In currently preferred embodiments, the propellant-containing canister is a pressurized canister that is sealed within the device and is not accessible to the user.

[0150] In various embodiments, the vial is a sealed glass vial. In currently preferred embodiments, the vial is a 3.5-mL amber sealed glass vial.

[0151] In typical embodiments, the liquid pharmaceutical composition that is sealably contained within the vial is a liquid pharmaceutical composition as described in Section 5.3.2 above. In currently preferred embodiments, the vial comprises a liquid pharmaceutical composition having the following composition: a clear, colorless to faintly yellow solution in an amber glass vial containing:

dihydroergotamine mesylate, USP	4.0 mg
caffeine, anhydrous, USP	10.0 mg
dextrose, anhydrous, USP	50.0 mg
carbon dioxide, USP.....	qs
purified water, USP	qs 1.0 mL.

[0152] The vial contains liquid pharmaceutical composition in an amount sufficient for at least one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses. In particular embodiments, the vial contains liquid pharmaceutical composition in an amount sufficient for at most one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses.

[0153] In typical embodiments, the propellant canister within the device that is co-packaged with the vial in the kit contains pressurized propellant in an amount sufficient for optional priming of the device followed by delivery of at least one total dose of DHE, or salt thereof, to be delivered by the device either in a single undivided or a plurality of divided doses. In particular embodiments, the propellant canister contains pressurized propellant in an amount sufficient for optional priming of the device followed by delivery of at most one total dose of DHE, or salt thereof, to be delivered by the device, in a single undivided or a plurality of divided doses.

5.6. Experimental examples

[0154] The invention is further described through reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting.

5.6.1. Example 1: Reproducibility of dose delivery

[0155] Table 2 provides experimental data on one implementation of the in-line device described in Section 5.3.5.1.1 above. As used in Table 2, “dose” refers to a volume delivered in a single device actuation.

TABLE 2

Shot #	Dose Volume [μ L]					
	Device 1	Device 2	Device 3	Device 4	Device 5	Device 6
1	190.6	193.7	185.3	199.2	199.2	145.1
2	181.4	205.5	178.9	167.7	167.7	141.7
3	183.1	188.5	173.3	165.6	165.6	138.5
4	183.2	193.3	145.8	164.6	164.6	136.6
5	183.3	201.5	200.7	162.0	162.0	142.1
6	185.8	207.7	166.3	179.4	179.4	138.9
7	184.3	195.1	180.3	164.8	164.8	140.9
8	183.3	205.4	175.3	164.9	164.9	142.0
9	180.5	178.1	172.0	164.1	164.1	141.8
10	179.7	204.0	178.0	170.6	170.6	143.9
Mean	183.5	197.3	175.6	170.3	170.3	141.2
StDev	3.1	9.3	14.0	11.3	11.3	2.5
Min	179.7	178.1	145.8	162.0	162.0	136.6
Max	190.6	207.7	200.7	199.2	199.2	145.1

185 μ L + 10%	203.5
185 μ L - 10%	166.5
185 μ L + 15%	212.8
185 μ L - 15%	157.3

5.6.2. Example 2: Phase I clinical trial

[0156] A Phase I clinical trial was conducted to compare the bioavailability of dihydroergotamine (DHE) mesylate following (i) single divided dose intranasal administration of INP104, a drug-device combination employing a Precision Olfactory Delivery (POD[®]) Device (Impel NeuroPharma, Seattle); (ii) intranasal administration of Migranal[®] Nasal Spray (Valeant Pharmaceuticals); and (iii) intravenous injection with D.H.E. 45[®] (Valeant Pharmaceuticals) in healthy adult subjects.

5.6.2.1. Study design

[0157] The study was a three-period, three-way, randomized, open-label, single-dose, cross-over, comparative bioavailability study.

[0158] Thirty-six subjects (approximately equal numbers of men and women) were enrolled and randomized into the study. Twenty-eight subjects completed the study. Treatment assignment was randomized in a three-treatment, three-period balanced crossover study of six sequences shown below, with a 7-day washout between treatments:

Table 3			
Sequence	Treatment		
	1	2	3
1	A	B	C
2	B	C	A
3	C	A	B
4	A	C	B
5	B	A	C
6	C	B	A

A = 1.45 mg INP104

B = 1.0 mg D.H.E. 45, IV

C = 2 mg Migranal[®] Nasal Spray.

Subjects all received 10 mg IV metoclopramide 5-10 minutes prior to each treatment.

[0159] INP104 was self-administered using the I123 POD™ Device (Impel NeuroPharma, Seattle). The dose of DHE mesylate was divided, with one spray in each nostril delivering a total target dose of 1.45 mg DHE mesylate.

[0160] The I123 POD Device is a handheld, manually actuated, propellant-driven, metered-dose administration device intended to deliver a drug formulation to the nasal cavity. Drug delivery to the nasal cavity via the I123 POD Device is driven by hydrofluoroalkane-134a (HFA) propellant. The I123 POD Device functions as an intranasal delivery device; the HFA propellant in the I123 POD Device is not intended to deliver drug to the lungs and does not contact the DHE formulation until the time of delivery.

[0161] The INP104 drug component, DHE DP, is a 3.5-mL amber glass vial filled with DHE mesylate 4 mg/mL. The formulation is identical to that in the Migranal[®] Nasal Spray device: a clear, colorless to faintly yellow solution in an amber glass vial containing:

dihydroergotamine mesylate, USP	4.0 mg
caffeine, anhydrous, USP	10.0 mg
dextrose, anhydrous, USP	50.0 mg
carbon dioxide, USP	qs
purified water, USP	qs 1.0 mL.

The DHE DP vial attaches to the I123 POD Device. The I123 POD Device may have a nominal output between 175 μ L/actuation pump and 205 μ L/actuation pump (inclusive). In some embodiments, the I123 POD Device may have a nominal output that is about 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, or 205 μ L/actuation pump.

[0162] A single manual actuation of the device by the user results in the operation of the metering pump to fill the dose chamber with the DHE formulation and subsequent, but almost instantaneous, activation of the propellant canister to expel the formulation through the nozzle, as a spray, resulting in delivery to the nasal cavity of the user. The device is designed to be disposed of after successful single divided-dose drug delivery (1 spray per nostril). Actuation of the I123 POD Device releases approximately 63 μ L of HFA-134a propellant, similar to HFA exposure from metered-dose inhalers.

[0163] D.H.E. 45[®] (Valeant Pharmaceuticals, NDA 005929) was administered in a volume of 1 mL intravenously over 1 minute.

[0164] Migranal[®] (Valeant Pharmaceuticals, NDA 20148) Nasal Spray (2 mg) was self-administered with equal dosing to both nostrils. In accordance with the product label, one spray (0.5 mg) was administered in each nostril initially, followed by an additional spray (0.5 mg) in each nostril 15 minutes later.

5.6.2.2. Pharmacokinetic assessments

Sampling and Processing

[0165] Blood samples for PK analysis were obtained, according to the clinical trial site's standard operating procedures (SOPs), within 15 minutes prior to dosing and at 5, 10, 20, 30, 40 and 50 minutes, and 1, 1.25, 1.5, 1.75, 2, 3, 4, 8, 12, 24, 36 and 48 hours after dosing. For the Migranal[®] Nasal Spray dose, the PK sampling timeclock was started following administration of the first dose of Migranal[®] Nasal Spray.

Pharmacokinetic Analysis

[0166] Individual DHE and 8'-OH-DHE plasma concentration data were listed for each individual and summarized by nominal sampling time-point and administration method with descriptive statistics (sample size [N], arithmetic mean, standard deviation [SD], median, minimum, maximum and geometric mean). Individual and mean DHE and 8'-OH-DHE plasma concentration-time profiles for each administration method were also graphed.

[0167] Pharmacokinetic parameters were computed from the individual plasma DHE and 8'-OH-DHE concentrations using a non-compartmental approach. Appropriate validated PK software (e.g., Phoenix WinNonlin v6.3) was used. The parameters that were determined and their definitions are provided in Table 4 below.

Table 4	
C_{max}	Maximum observed drug concentration.
T_{max}	Time to maximum observed drug concentration. If the maximum value occurs at more than one time-point, T_{max} is defined as the first time point with this value.
AUC_{0-t}	Area under the drug concentration-time curve, calculated using linear-up log-down trapezoidal summation from time zero to the time of the last Measurable concentration.
k_{el}	Apparent terminal elimination rate constant, calculated by linear regression of the terminal linear portion of the log concentration vs. time curve.
AUC_{0-inf}	Area under the drug concentration-time curve from time zero to infinity, calculated as $AUC_{0-t} + C_t/k_{el}$.
$t_{1/2}$	Apparent elimination half-life, calculated as $\ln(2)/k_{el}$.
CL/F (CL for i.v.)	Apparent clearance calculated as $Dose/AUC_{0-inf}$.
V_z/F (V_z for i.v.)	Apparent volume of distribution at the terminal phase, calculated as $Dose/(k_{el} * AUC_{0-inf})$.

Statistical Methods for Pharmacokinetic Analyses

[0168] PK parameters were summarized by administration method using descriptive statistics (arithmetic means, SD, coefficients of variation [CV], sample size [N] minimum, maximum, median and geometric mean). Geometric mean was calculated for AUC_{0-t} , AUC_{0-inf} , and C_{max} .

[0169] No value for k_{el} , $t_{1/2}$, AUC_{0-inf} , CL/F , V_z/F , as appropriate, were reported for cases that did not exhibit a terminal log-linear phase in the concentration versus time profile or did not contain sufficient data during this phase for parameter estimation.

Statistical Analysis

[0170] A comparative bioavailability assessment was undertaken to demonstrate (i) that the lower 90% confidence interval of the DHE after INP104 to DHE after Migranal Nasal Spray geometric mean ratios for C_{max} and AUC (AUC_{0-t} , AUC_{0-inf}) is not less than 80%, and (ii) the upper 90% confidence interval of the DHE after INP104 to D.H.E. 45 Injection (IV) geometric mean ratios for C_{max} and AUC (AUC_{0-t} , AUC_{0-inf}) not greater than 125% — *i.e.*, to demonstrate that exposure is equal to or greater than 80% and equal to or less than 125% range observed between Migranal Nasal Spray and D.H.E. 45 Injection (IV), respectively.

[0171] For each comparator (Migranal Nasal Spray and D.H.E. 45 Injection (IV)), the following analysis methods were performed independently. Analysis of variance (ANOVA) with effects for sequence, subject nested within sequence, period, and treatment were performed on the ln-transformed DHE and 8'OH-DHE AUC_{0-t} , AUC_{0-inf} and C_{max} . Each ANOVA included calculation of least squares mean (LSM), the difference between administration method LSM, and the standard error associated with the difference.

[0172] Only subjects who had completed all three treatments and had sufficient PK sample collection to generate the key PK parameters (AUC_{0-t} , AUC_{0-inf} and C_{max}) for each administration method were included in the ANOVA analysis.

[0173] Ratios of geometric means were calculated using the exponentiation of the difference between treatment LSM from the analyses on the ln-transformed AUC_{0-t} , AUC_{0-inf} and C_{max} . These ratios were expressed as a percentage relative to the reference (comparator) treatment, *i.e.* INP104 [test]/ Comparator [reference]. Consistent with the two one-sided tests for

bioequivalence, 90% confidence intervals were obtained for the ratio of the geometric means for AUC_{0-t} , AUC_{0-inf} and C_{max} .

5.6.2.3. Results: DHE and 8'OH-DHE pharmacokinetics

[0174] The time course of plasma DHE concentrations is plotted in FIGS. 10A and 10B, and initial summary statistics are provided in Table 5 below.

Table 5 DHE pharmacokinetics			
	Migranal[®] (2 mg intranasal)	INP104 (1.45 mg intranasal)	D.H.E. 45[®] (1 mg IV)
AUC_{0-inf} (pg*hr/ml) mean [%CV] (range)	2,248 [68%] (444 - 7247)	6,291 [44%] (978 - 10,445)	10,053 [18%]
C_{max} (pg/ml) mean [%CV] (range)	318 [83%] (25.4 - 1190)	1,258 [54%] (270 - 2660)	14,460 [34%]
T_{max} (min) mean	55	34	5
T_{max} (hr) mean (range)	0.92 (0.5 – 3.08)	0.57 (0.333-2.05)	0.08

[0175] As compared to Migranal[®] Nasal Spray, INP104 provides nearly 3-fold higher mean systemic drug exposure, with an AUC_{0-inf} of 6,291 pg*hr/ml as compared to 2,248 pg*hr/ml for Migranal[®]. INP104 also provides nearly 4-fold higher mean maximal plasma concentration, with a C_{max} of 1,258 pg/ml as compared to 318 pg/ml for Migranal[®]. Maximal DHE plasma concentration is reached faster with INP104, with a mean T_{max} of 34 minutes versus 55 minutes for Migranal[®]. The higher systemic drug exposure and higher maximal plasma concentration were achieved with a lower administered dose of the identical formulation of DHE mesylate, 1.45 mg for INP104 versus 2.0 mg for Migranal[®], and without requiring a 15-minute wait between administration of divided sub-doses, as required for Migranal[®].

[0176] In addition, systemic delivery of DHE was more consistent with INP104 than with Migranal[®], with lower variation observed across subjects for both AUC_{0-inf} and C_{max} parameters (see Table 5 above for coefficients of variation).

[0177] Although bolus intravenous administration of 1 mg DHE mesylate provided greater than 10-fold higher C_{max} than 1.45 mg DHE mesylate administered intranasally by INP104, the high

C_{max} achieved with intravenous administration is known to be correlated with adverse events (“AE”s), specifically nausea, and IV DHE mesylate (D.H.E. 45) is most commonly administered with an anti-emetic. Within 20-30 minutes following administration, DHE plasma concentrations achieved through INP104 intranasal administration were essentially indistinguishable from concentrations achieved by intravenous administration. Thus, despite a greater than 10-fold higher C_{max} , bolus intravenous administration of 1 mg DHE mesylate provided less than 2-fold greater systemic drug delivery, measured as AUC_{0-inf} , as compared to INP104 intranasal delivery.

[0178] The 8'OH-DHE metabolite of DHE is known to be active, and to contribute to the long-lasting effect of DHE on migraine. The time course of plasma 8'-OH-DHE concentrations is plotted in FIGS. 11A and 11B. Initial summary statistics for plasma concentrations of 8'-OH-DHE are provided in Table 6, below.

Table 6 8'-OH-DHE pharmacokinetics			
	Migranal[®] (2 mg intranasal)	INP104 (1.45 mg intranasal)	D.H.E. 45[®] (1 mg IV)
AUC_{0-inf} (pg*hr/ml) [%CV]	1113 [53%] n = 6	1063 [59%] n = 20	924 [63%] n = 28
C_{max} (pg/ml) [%CV]	42 [35%] n = 8	58 [44%] n = 24	392 [26%] n = 28
T_{max} (hr)	2.30 [57%] n = 8	1.43 [53%] n = 24	0.08 [8%] n = 28

[0179] These data demonstrate that intranasal administration of 1.45 mg DHE by INP104 provides equivalent systemic exposure to the active metabolite of DHE as bolus intravenous administration of 1.0 mg DHE. In addition, the metabolite was detected in only 8 subjects after Migranal[®] intranasal delivery, versus 24 subjects following intranasal administration of INP104.

6.

[0180]

7. EQUIVALENTS

[0181] While this invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The appended claims are intended to be construed to include all such embodiments and equivalent variations.

CLAIMS:

1. Use of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof, for acutely treating migraine headache with or without aura, wherein:

an effective dose of the liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof is for intranasal administration to a subject with migraine headache, and

the dose comprises about 1.45 mg of DHE or salt thereof, and is for administration to the nose as a plurality of divided doses over a period of no more than 1 minute by a manually actuated, propellant-driven metered-dose intranasal delivery device.

2. The use according to claim 1, wherein the dose is for administration as two divided doses.

3. The use according to claim 2, wherein one divided dose is for administration to each nostril.

4. The use according to claim 1, wherein the dose is for administration over no more than 45 seconds.

5. The use according to claim 4, wherein the dose is for administration over no more than 30 seconds.

6. The use according to any one of claims 2-5, wherein the volume of liquid composition for administration per divided dose is 140-250 μL .

7. The use according to claim 6, wherein the volume of liquid composition for administration per divided dose is 175 μL -225 μL .

8. The use according to claim 7, wherein the volume of liquid composition for administration per divided dose is about $\sim 200 \mu\text{L}$.

9. The use according to any one of claims 1-8, wherein the liquid composition comprises a salt of DHE.

10. The use according to claim 9, wherein the liquid composition comprises DHE mesylate.
11. The use according to claim 10, wherein the liquid composition comprises DHE mesylate at a concentration of 2.5-7.5 mg/ml.
12. The use according to claim 11, wherein the liquid composition comprises 3.5-6.5 mg/ml DHE mesylate.
13. The use according to claim 12, wherein the liquid composition comprises 4.0 mg/ml DHE mesylate.
14. The use according to any one of claims 1-13, wherein the liquid composition further comprises caffeine.
15. The use according to claim 14, wherein the liquid composition comprises caffeine at a concentration of 10 mg/ml.
16. The use according to any one of claims 1-15, wherein the liquid composition further comprises dextrose.
17. The use according to claim 16, wherein the liquid composition comprises dextrose at a concentration of 50 mg/ml.
18. The use according to any one of claims 1-17, wherein the liquid composition comprises 4.0 mg/ml DHE mesylate, 10.0 mg/ml caffeine, and 50 mg/ml dextrose.
19. The use according to claim 1, wherein, prior to first manual actuation, the liquid pharmaceutical composition and propellant are not in contact within the device.
20. The use according to claim 19, wherein the liquid pharmaceutical composition is contained in a vial and the propellant is contained in a canister, wherein the canister is a pressurized canister.
21. The use according to claim 20, wherein, between successive manual actuations, the liquid pharmaceutical composition in the vial and propellant in the canister are not in contact within the device.

22. The use according to claim 21, wherein each manual actuation brings a metered volume of liquid pharmaceutical composition and a separately metered volume of propellant into contact within a dose chamber of the device.

23. The use according to claim 22, wherein contact of propellant with liquid pharmaceutical composition within the dose chamber of the device creates a spray of liquid pharmaceutical composition as the formulation is expelled through a nozzle of the device.

24. The use according to claim 23, wherein the nozzle has a plurality of lumens, and the spray is ejected simultaneously through a plurality of nozzle lumens.

25. The use according to any one of claims 19-24, wherein the propellant is a hydrofluoroalkane propellant.

26. The use according to claim 25, wherein the propellant is hydrofluoroalkane-134a.

27. The use according to any one of claims 21-24, wherein, prior to first actuation, the vial is nonintegral to the device and is configured to be attachable thereto.

28. The use according to claim 27, wherein the vial is configured to be threadably attachable to the device.

29. The use according to any one of claims 1-28, wherein the subject has migraine headache with aura.

30. The use according to any one of claims 1-28, wherein the subject has migraine headache without aura.

31. The use according to any one of claims 1-30, wherein the subject has had onset of at least one prodromal symptom of migraine.

32. The use according to any one of claims 1-31, wherein the subject has menstrual-associated migraine.

33. The use according to any one of claims 1-32, wherein the subject has migraine that does not respond to triptan drugs.

34. The use according to any one of claims 1-33, wherein the liquid pharmaceutical composition is for self-administration by the subject.

35. A kit for acutely treating migraine with or without aura by intranasal administration of a liquid pharmaceutical composition comprising dihydroergotamine (DHE) or salt thereof, comprising:

a vial, within which is sealably contained the liquid pharmaceutical composition, and
a device,

wherein the vial is configured to be attachable to the device, and

wherein the device, upon attachment of the vial, is a manually actuated, metered-dose, propellant-driven intranasal administration device that is suitable for administration of an effective dose of the liquid pharmaceutical composition comprising about 1.45 mg DHE or salt thereof to the nose as a plurality of divided doses over a period of no more than 1 minute.

36. The kit of claim 35, wherein the device comprises a canister, wherein the canister is a pressurized canister containing propellant.


37. The kit of claim 36, wherein, following attachment of the vial to the device and prior to a first manual actuation, the liquid pharmaceutical composition and propellant are not in contact within the device.


38. The kit of claim 37, wherein, between successive manual actuations, the liquid pharmaceutical composition in the vial and propellant in the canister are not in contact within the device.


39. The kit of claim 37 or claim 38, wherein each manual actuation brings a metered volume of liquid pharmaceutical composition and a separately metered volume of propellant into contact within a dose chamber of the device.


40. The kit of claim 39, wherein contact of propellant with liquid pharmaceutical composition within the dose chamber of the device creates a spray of liquid pharmaceutical composition as the formulation is expelled through a nozzle of the device.


41. The kit of any one of claims 35-40, wherein the liquid pharmaceutical composition within the vial comprises a salt of DHE.
42. The kit of claim 35, wherein the liquid composition comprises DHE mesylate.
43. The kit of claim 42, wherein the liquid composition comprises DHE mesylate at a concentration of 2.5-7.5 mg/ml.
44. The kit of claim 43, wherein the liquid composition comprises 4.0 mg/ml DHE mesylate.
45. The kit of claim 44, wherein the liquid composition comprises 4.0 mg/ml DHE mesylate, 10.0 mg/ml caffeine, and 50 mg/ml dextrose.
46. The kit of any one of claims 35-45, wherein the vial contains no more than 2 ml of liquid pharmaceutical composition.
47. The kit of claim 46, wherein the vial contains approximately 1 ml of liquid pharmaceutical composition.
48. The kit of any one of claims 36-47, wherein the pressurized canister contains an amount of propellant sufficient to administer no more than 1 dose of liquid pharmaceutical composition.
49. The kit of any one of claims 36-47, wherein the pressurized canister contains an amount of propellant sufficient to administer more than 1 dose of liquid pharmaceutical composition.
50. The use according to any one of claims 1-33, wherein the liquid pharmaceutical composition is for administration to the subject by a care-giver of the subject.


**FIG. 1**


**FIG. 2**


**FIG. 3**

**FIG. 4**

**FIG. 5A**

**FIG. 5B**



**FIG. 7**


**FIG. 8**

FIG. 9A





FIG. 9B



SECTION A-A




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 13A




FIG. 13B




FIG. 14

