

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2013314303 B2**

**(54) Title  
Semicfluorinated alkane compositions**

**(51) International Patent Classification(s)**  
**A61K 9/00** (2006.01)      **A61K 47/06** (2006.01)  
**A61K 9/08** (2006.01)      **A61P 27/02** (2006.01)  
**A61K 31/232** (2006.01)

**(21) Application No:** **2013314303**      **(22) Date of Filing:** **2013.09.12**

**(87) WIPO No:** **WO14/041071**

**(30) Priority Data**

<b>(31) Number</b>	<b>(32) Date</b>	<b>(33) Country</b>
<b>12183997.1</b>	<b>2012.09.12</b>	<b>EP</b>
<b>13169399.6</b>	<b>2013.05.27</b>	<b>EP</b>

**(43) Publication Date:** **2014.03.20**

**(44) Accepted Journal Date:** **2018.01.18**

**(71) Applicant(s)**  
**Novaliq GmbH**

**(72) Inventor(s)**  
**Gunther, Bernhard;Scherer, Dieter;Pettigrew, Anthony**

**(74) Agent / Attorney**  
**AJ PARK, L 22 1 Willis St, WELLINGTON, ACT, 6011, AU**

**(56) Related Art**  
**WO 2011/073134 A1**  
**US 2012/0010280 A1**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2014/041071 A1

(43) International Publication Date

20 March 2014 (20.03.2014)

(51) International Patent Classification:

*A61K 9/00* (2006.01) *A61K 31/232* (2006.01)  
*A61K 47/06* (2006.01) *A61P 27/02* (2006.01)  
*A61K 9/08* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2013/068909

(22) International Filing Date:

12 September 2013 (12.09.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12183997.1 12 September 2012 (12.09.2012) EP  
13169399.6 27 May 2013 (27.05.2013) EP

(71) Applicant (for all designated States except US):  
**NOVALIQ GMBH** [DE/DE]; Im Neuenheimer Feld 515,  
69120 Heidelberg (DE).

(72) Inventors; and

(71) Applicants (for US only): **GÜNTHER, Bernhard** [DE/DE]; Schriesheimerstrasse 19, 69221 Dossenheim (DE). **SCHERER, Dieter** [DE/CH]; Im Tschambol 18, CH-4242 Laufen (CH). **PETTIGREW, Anthony** [GB/SE]; Parongatan 5, S-234 37 Lomma (SE).

(74) Agent: **CREMER, Karsten**; Almstadtstr. 7, 10119 Berlin (DE).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2014/041071 A1

(54) Title: SEMIFLUORINATED ALKANE COMPOSITIONS

(57) Abstract: The invention provides novel compositions comprising semifluorinated alkanes and at least one compound sensitive or prone to oxidation. The compositions can be used as medicines that are topically administered to an eye or ophthalmic tissue. The invention further provides kits comprising such compositions.

## TITLE: SEMIFLUORINATED ALKANE COMPOSITIONS

**Description****FIELD**

The present invention is in the field of compositions comprising oxidation-prone pharmaceutically active ingredients, specifically compositions that are based on semifluorinated alkanes. In particular, the compositions of the present invention may be topical ophthalmic compositions useful in the treatment of ocular diseases or conditions.

**BACKGROUND**

5 The stability of an active pharmaceutical compound during storage over a variety of conditions is a generally of concern during development of a formulation for the compound. This aspect is especially relevant for liquid formulations of compounds which are sensitive and/or prone to decomposition via oxidation pathways. It is becoming even more relevant today as countries with hot and  
10 moderately humid climate (i.e. ICH zone IVb countries) are becoming increasingly important to pharmaceutical companies so that new drug formulations are needed which take into account the higher temperatures and humidities of these countries.

Oxidation may be promoted thermally, photolytically, or by chemical means. Compounds comprising aliphatic double bond systems are, in particular, sensitive  
15 towards degradation by oxidation, often via free-radical chain processes with molecular oxygen.

Such processes, known also as auto-oxidation, generally begins via an initiation process in which a free-radical is generated (e.g. via abstraction of a hydrogen atom by an initiator radical), followed by propagation steps involving oxygen and further  
20 compound molecules. Termination may occur when two radical products couple. Compounds comprising free-radical stabilizing aliphatic double bond configurations such as the methylene-interrupted polyene systems of polyunsaturated lipid derivatives are generally more susceptible to auto-oxidation. These and other

5 oxidation processes can readily lead to, over time, the formation and accumulation of undesirable and pharmaceutically ineffective reaction products. Some of the degradants may be harmful or toxic.

The use of antioxidants and/or stabilizer excipients remains one of the foremost strategies for mitigating or preventing the oxidation of such compounds and to increase shelf-life. Typical antioxidants are butylated hydroxytoluene (BHT), ascorbic acid, tocopherol derivatives etc. Many antioxidants will act as free-radical scavengers; and act by terminating free-radicals and inhibiting the chain processes. Excipients which can act as metal ion chelators e.g. ethylene diamine tetraacetic acid (EDTA) may also be used to limit the effect of trace heavy metal impurities which can also catalyse oxidation reactions. The encapsulation or shielding of oxidation-sensitive compounds, such as with liposomes or cyclodextrins are also known methods of preventing degradation.

A further measure to prevent oxidative degradation of such compounds would be the removal of oxygen dissolved in the liquid vehicle through sparging with an inert gas, with for example, nitrogen. This is however process-intensive and not cost-effective, and requires, in addition, special attention during further processing steps and with packaging to prevent the reintroduction of atmosphere.

Polyunsaturated fatty acids (often abbreviated as PUFA) are a class of compounds that are prone to oxidative degradation. These include omega fatty acids such as omega-3 fatty acids (also known as  $\omega$ -3 fatty acids or n-3 fatty acids) and omega-6 fatty acids (also known as  $\omega$ -6 fatty acids or n-6 fatty acids). These are essential fatty acids because they are available only through dietary intake, and because they involved in many human metabolic processes and functions. As such, they have been implicated as beneficial in the treatment or prophylaxis of a wide variety of different health conditions.

For example, these compounds have been found to be useful for the topical treatment and prevention of ocular pathologies such as dry eye disease (DED). DED, also known as keratoconjunctivitis sicca or dysfunctional tear syndrome, is a multifunctional disorder of the tear film and ocular surface which results in

5 discomfort, visual disturbance, and often even in ocular surface damage. The loss in dynamic stability of the structure, composition, volume and distribution, as well as clearance of the tear film can lead to the development of DED.

An inflammation cycle is one of the key processes that maintain and potentially progress the dry eye condition. Omega fatty acids, in relation to their anti-10 inflammatory properties, have been found to reduce the severity of DED and its symptoms and improve tear secretion. Omega-3 fatty acids, in particular, have been associated with the modulation of the production of anti-inflammatory and immunomodulatory eicosanoid prostaglandins such as PGE1. They are also implicated in suppressing the expression of pro-inflammatory cytokines such as IL-115 and TNF- $\alpha$ , which are associated with dry eye disease.

The formulation of such active ingredients prone to oxidation into topical ophthalmic compositions such as eye drops can however be challenging. The choices of excipients, and therefore antioxidants and stabilizers which can be used are more limited due to incompatibility, toxicity or poor tolerance with the ocular surface. At 20 the same time, however, the minimization of degradation (resulting in colour changes of the preparation, precipitation of insoluble materials, etc.) becomes more acute.

25 Compositions comprising omega fatty acids for the topical treatment of the dry eye condition and its symptoms are however known in the art. Gastrointestinal intolerance resulting from the oral ingestion of large quantities of these fatty acids as supplements (and systemic absorption effect) can be avoided with direct and local administration of these active molecules to the surface of the eye.

For example, Remogen® Omega (marketed by TRB Chemedica International S.A.), is a commercially available ophthalmic preparation comprising omega fatty acids that indicated for dry eye conditions. As disclosed in its product information, 30 the preparation is a hypotonic hydrogel containing eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), antioxidant vitamin E (tocopheryl acetate), the emollient substances carbopol 980, glycerol, pemulen, and sodium hydroxide, disodium phosphate, and distilled water. The hydrogel composition itself is described as a microemulsion, with a dispersion of the lipidic molecules (fatty acids and vitamin

5 E). The product is packaged in the form of single-dose containers packed in an aluminium pouch, with the recommendation to discard the product immediately after use. With respect to stability, storage is suggested to be in a refrigerator at 2-8 °C, with a shelf-life of 3 months if stored, at below 25°C, unrefrigerated. Due to its poor thermal stability, the product is shipped in a cold chain, i.e. transported in  
10 refrigerated containers and stored at the warehouse and in the pharmacy under refrigeration. Thus the consignment is expensive and requires a substantial effort in terms of temperature monitoring and logistics.

Similar, if not the same type of ophthalmic hydrogel compositions are also disclosed in US 2012/0010280. It is described that the fatty acids are in solution in  
15 the antioxidant tocopheryl acetate, with the tocopheryl acetate present in amounts of up to 75% by weight of the oily mixture. These are dispersed as oily droplets in the hydrogel. Aqueous emulsions comprising a combination of omega-3 and omega-6 fatty acids, and surfactant (e.g. Tween 80), Glucam E-20, and a drop of vitamin E as an antioxidant were found to be poorly adapted for maintaining stability of the omega  
20 fatty acids are proposed in WO 2006/007510. It should be noted, however, that emulsion systems based on hydrogels, with their generally increased viscosity and bioadhesive properties may incur, upon instillation and also during its contact time to the ocular surface, irritating foreign body sensation and blurred vision. Depending also on the degree of their viscosity, these may be more difficult to dispense and  
25 administer.

In general, one of the major disadvantages of formulations based on emulsions system is that in contrast to single phase systems, emulsion systems may be more complex and difficult to manufacture, especially in sterile form. Frequently, they are not readily sterilisable by thermal treatment without negative impact on their  
30 physical properties. Emulsions are also inherently unstable, and could phase separate with time or fluctuations in storage conditions. They, as with all aqueous based systems, are more prone to microbial contamination during use as well. The aseptic processing of emulsions is complex, costly, and is associated with higher risks of failure, i.e. microbial contamination.

5        Retinoid (Vitamin A) derivatives are another class of hydrocarbon compounds comprising high levels of polyunsaturation, and which are also prone to degradation via oxidation pathways. Retinoid derivatives include retinol, retinal, retinoic acid, tretinoin, isotretinoin, alitretinoin, and related derivatives. Vitamin A derivatives have been used for the treatment of inflammation in relation to the cornea,  
10      conjunctiva and other mucosal and epithelial tissues, including conditions such as dry eye disease.

US2012/0095097 for example discloses aqueous ophthalmic compositions comprising Vitamin A, at least 0.4 w/v% of polyoxyethylene polypropylene glycol and trometamol. Antioxidants such as dibutylhydroxytoluene and  $\alpha$ -tocopherol acetate,  
15      may also be present in these compositions. These compositions are based on micelles of Vitamin A, which is presumably shielded and stabilized by non-ionic surfactant polyoxyethylene polypropylene glycol. These compositions are not preserved as such.

In principle, these preparations would require, if they were to be presented in multi-dose containers which are in principle more cost-efficient and convenient for  
20      patients than single-use vials, preservation in order to ensure their microbiological quality. The same would be applicable to all aqueous-based preparations. At the same time however, preservatives which can be used in ophthalmic formulations are potentially damaging to the eye, in particular to the ocular surface, and should be avoided especially in the context of dry eye disease. At least in earlier years, multi-  
25      dose formulations for ophthalmic administration had to be preserved using a physiologically acceptable preservative in order to reduce the risk of microbial contamination and infection.

Most preservatives are however problematic for DED patients in that they have a potential to negatively affect the ocular surface, thus counteracting the therapeutic intent. This is particularly relevant for patients with moderate to severe dry eye  
30      disease symptoms who may require frequent use for symptom relief, as well as patients who require multiple preserved topical medicaments.

As an alternative, single-dose containers are the main option for the administration of non-preserved formulations such as those used for the Remogen®

5 Omega product. These are however less cost-efficient and convenient to handle for the patient than the conventional multi-dose bottle. Whilst ophthalmic formulations utilizing 'vanishing' preservatives such as sodium chlorite or sodium perborate, which can convert to non-toxic ions and water after instillation and contact with the tear film may also be an option, these may still be irritating to patients especially

10 those with severe disease who may not have sufficient tear volume to effectively degrade the preservatives.

WO 2011/073134 discloses ophthalmic topical pharmaceutical compositions comprising immunosuppressant macrolides such as ciclosporin A and semifluorinated alkanes, for treatment of keratoconjunctivitis sicca. The

15 semifluorinated alkanes in the disclosed compositions serve as suitable liquid vehicles for delivering the therapeutic pharmaceutical agent to the eye, and in particular have a high capacity for dissolving extremely challenging poorly soluble compounds such as ciclosporin. However, there is no mention of the ability of semifluorinated alkanes to stabilize oxidation-sensitive compounds over an extended

20 period of time.

It is therefore an object of the present invention to provide a novel composition which comprises one or more active compounds comprising a polyunsaturated fatty acid or a derivative thereof, and/or which at the same time addresses and overcomes at least one of the various issues and/or at least one of the limitations or

25 disadvantages associated with prior art formulations. In a specific aspect, it is an object of the invention to provide a stable liquid composition for the treatment or prevention of a condition or disease such as dry eye or conditions relating to the inflammation of the ocular tissue; and/or to at least provide the public with a useful choice. In a further aspect, it is an object of the invention to provide a method of

30 stabilizing a polyunsaturated fatty acid selected from omega-3 or omega-6 fatty acid or ester derivatives thereof, or mixtures thereof which does not exhibit one or more of the disadvantages of prior art; and/or which at least provides the public with a useful choice. Further objects of the invention will become clear on the basis of the following description, examples, and patent claims.

5 In this specification where reference has been made to patent specifications, other external documents, or other sources of information, this is generally for the purpose of providing a context for discussing the features of the invention. Unless specifically stated otherwise, reference to such external documents is not to be construed as an admission that such documents, or such sources of information, in  
10 any jurisdiction, are prior art, or form part of the common general knowledge in the art.

In the description in this specification reference may be made to subject matter that is not within the scope of the claims of the current application. That subject matter should be readily identifiable by a person skilled in the art and may assist in  
15 putting into practice the invention as defined in the claims of this application.

#### SUMMARY OF THE INVENTION

In a first aspect, the invention provides a stable liquid composition comprising a polyunsaturated fatty acid or a derivative thereof dissolved in a liquid vehicle comprising a semifluorinated alkane of formula


20  $F(CF_2)_n(CH_2)_mH$

wherein n and m are integers independently selected from the range of 3 to 20, and wherein the liquid vehicle is substantially free of water, wherein the polyunsaturated fatty acid or derivative thereof is an omega-3 or omega-6 fatty acid or derivative thereof, or a mixture of omega-3 or omega-6 fatty acids or derivatives thereof, wherein the derivative(s) are selected from esters, resolvins and  
25 neuroprotectins.

The invention also provides a composition of the invention for use as a medicament.

The invention also relates to a method of stabilising a polyunsaturated fatty acid  
30 selected from an omega-3 or an omega-6 fatty acid or ester derivatives thereof, or mixtures thereof, comprising the step of dissolving, dispersing or suspending the

5 polyunsaturated fatty acid in a liquid vehicle comprising a semifluorinated alkane of formula



or of formula



10 wherein n, m, and o are integers independently selected from the range of 3 to 20 and wherein the liquid vehicle is substantially free of water.

15 The invention also relates to a method of treating a an inflammatory condition or disease of the ophthalmic tissue, or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith, the method comprising topically administering a composition of the invention to an eye or ophthalmic tissue of a subject in need thereof.

20 The invention also relates to use of a stable liquid composition of the invention in the manufacture of a medicament for use in the topical ophthalmic treatment of an inflammatory condition or disease of the ophthalmic tissue or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith.

#### BRIEF DESCRIPTION

25 Described herein is a stable liquid composition comprising an active compound with more than one aliphatic double bond prone to oxidation, wherein the active compound is dissolved, dispersed or suspended in a liquid vehicle comprising a semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$  or of formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ , wherein n, m, and o are integers independently selected from the range of 3 to 20. The liquid vehicle is substantially free of water.

30 In one of the embodiments, the composition described comprises a polyunsaturated fatty acid or derivative thereof, in particular an ester derivative. It is further preferred that the polyunsaturated fatty acid or derivative is selected from an omega-3 or omega-6 fatty acid or derivative, such as an omega-3 or omega-6 fatty

5 acid ester, or a mixture thereof. In a further preferred embodiment, said omega-3 or omega-6 fatty acids are selected from docosahexaenoic acid or eicosapentaenoic acid and ester derivatives thereof.

Also described is the use of such compositions as a medicament, wherein the medicament may be topically administered to the eye or ophthalmic tissue, and/or 10 wherein the medicament may be used for the treatment of a condition or disease of an eye or ophthalmic tissue such as inflammatory conditions of the ophthalmic tissue or keratoconjunctivitis sicca (dry eye).

Also described is a method of stabilizing an active compound with more than one aliphatic double bond prone to oxidation comprising the step of dissolving, 15 dispersing, suspending the active compound in a liquid vehicle comprising a semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$  or of formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ , wherein n, m, and o are integers independently selected from the range of 3 to 20, and wherein the liquid vehicle is substantially free of water. In 20 one of the preferred embodiments, the active compound is an omega-3 or an omega-6 fatty acid or an ester derivative thereof.

#### DETAILED DESCRIPTION OF THE INVENTION

Described is a stable liquid composition comprising an active compound with more than one aliphatic double bond prone to oxidation. The composition is further characterized in that the active compound is dissolved, dispersed or suspended in a 25 liquid vehicle comprising a semifluorinated alkane as defined in claim 1 and further explained below

As used herein, an active compound refers to any type of pharmaceutically, nutraceutically or otherwise health-promoting active compound or derivative that is useful in the prevention, diagnosis, stabilization, treatment, or -generally speaking- 30 management of a condition or disease. A therapeutically effective amount refers to a dose, concentration or strength which is useful for producing a desired pharmacological, nutraceutical or otherwise health-promoting or -supporting effect.

5 Active compounds with more than one aliphatic double bond prone to oxidation are usually susceptible to degradation and deterioration over time and storage. The aliphatic double bonds of these compounds are featured in configurations that are particularly prone to undergo oxidation reactions, leading to the formation of side products and quantitative loss of active compound over time. Described are methods  
10 of stabilizing such active compounds against oxidation, in particular oxidation which may occur and/or be promoted by thermal, photolytic or chemical means, and in the presence of oxygen as may be present in the atmosphere or as introduced to an initially inert environment over time.

As used herein, an aliphatic double bond refers to a carbon-carbon double bond  
15 (C=C). An active compound with more than one such double bond may have at least two aliphatic double bonds, in particular in a configuration wherein the double bonds are separated by a methylene group (-CH<sub>2</sub>-). The one or more aliphatic double bonds of the active compounds of the invention may be in the cis (Z) or trans (E) configuration, or combinations thereof. In one embodiment, the active compound  
20 comprises a structural component of the formula -(HC=CH-CH<sub>2</sub>-HC=CH)<sub>x</sub>-, wherein x is an integer ranging from 2 to 10, in particular from 2 to 5.

Active compounds with more than one aliphatic double bond in which the double bonds are separated by a bridging methylene group (also commonly referred to as compounds possessing methylene-interrupted polyenes) include  
25 polyunsaturated fatty acids. Polyunsaturated fatty acids are linear carboxylic acids possessing carbon chains with more than one carbon-carbon double bond. These lipophilic lipids are prone to oxidation, especially in the presence of atmospheric oxygen via radical initiated auto-oxidation processes. In an embodiment, the compositions described comprise a polyunsaturated fatty acid with number of carbon  
30 atoms ranging from C16 to C24. Derivatives of polyunsaturated fatty acids, such as esters and in particular alkyl esters are also described. It should be understood within the context of the present invention that a reference to any polyunsaturated fatty acid, or group or species thereof, also covers the respective derivatives, in particular the esters, even if not specifically mentioned.

5 Polyunsaturated fatty acids (often abbreviated as PUFA) include omega fatty acids such as omega-3 fatty acids (also known as  $\omega$ -3 fatty acids or n-3 fatty acids) and omega-6 fatty acids (also known as  $\omega$ -6 fatty acids or n-6 fatty acids). Examples of omega-3 fatty acids (also known as  $\omega$ -3 fatty acids or n-3 fatty acids) include eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA),  $\alpha$ -linolenic acid (ALA).

10 Examples of omega-6 fatty acids (also known as  $\omega$ -6 fatty acids or n-6 fatty acids) include gamma-linolenic acid, linoleic acid, dihomo-gamma-linolenic acid. Derivatives of omega-3 fatty acids or omega-6 fatty acids, such as ester or alkyl ester derivatives are also within the scope of the invention. Particularly preferred alkyl ester derivatives are eicosapentaenoic acid ethyl ester or docosahexaenoic acid ethyl ester.

15 Derivatives of eicosapentaenoic acid or docosahexaenoic acid such as resolvins and neuroprotectins are also suitable as active compounds within the scope of the invention.

In a further embodiment, the active compounds with more than one aliphatic double bond prone to oxidation are compounds comprising a structural component with more than one aliphatic double bond in linear conjugation with one another. Active compounds comprising more than one aliphatic double bond in linear conjugation are also commonly referred to as polyenes. Preferably, the active compound comprises at least three, or more preferably four or five linearly conjugated double bonds. Preferably, the active compounds structurally comprise a conjugated triene, tetraene, pentaene, hexaene or heptaene component. In one embodiment, the active compound comprising a polyene component may be a macrocycle, wherein the polyene or the linearly conjugated double bonds form part of the cyclic structure. The one or more linearly conjugated aliphatic double bonds may be in the cis (Z) or trans (E) configuration, or combinations thereof.

30 Examples of active compounds comprising more than one linearly conjugated double bond include, but are not limited to, lipophilic vitamin derivatives such as retinoids and derivatives thereof. This class of compounds are also unstable towards prolonged storage and prone to degradation through oxidative pathways such as auto-oxidation. Retinoids and retinoid derivatives (also often referred to as Vitamin A

5 derivatives) include retinol, retinoic acid and its esters (e.g. retinol palmitate or retinol acetate), retinal, tretinoin, isotretinoin, and alitretinoin.

Further examples of active compounds also comprising linearly conjugated double bond systems sensitive and or prone to radical-mediated oxidative degradation with oxygen include sirolimus (rapamycin), a macrolide 10 immunosuppressant drug compound whose structure contains three contiguous double bonds.

Moreover, polyene compounds are contemplated as oxidation-sensitive active ingredients, in particular polyene antifungals. In one embodiment, the composition comprises a polyene antifungal such as nystatin, natamycin or amphotericin

15 dissolved in a liquid carrier substantially consisting of one or more semifluorinated alkanes as described herein.

The compositions described may optionally further comprise a carotenoid or carotenoid derivative, in particular a xanthophyll. These compounds also possess poly-conjugated double bond systems. Particularly preferred are lutein and 20 zeaxanthin. Derivatives of lutein or zeaxanthin, such as lutein or zeaxanthin esters are also considered. In a particular embodiment, liquid compositions comprising at least two or more semifluorinated alkanes further comprise lutein or a derivative thereof.

In a further embodiment, the composition comprises ciclopirox olamine as active ingredient. Preferably, ciclopirox olamine is incorporated in the dissolved state.

25 As will be discussed in more detail below, some of the key advantages of the compositions described are brought about by the presence of a semifluorinated alkane in the composition as liquid vehicle, or part of the liquid vehicle, for active compounds such as polyunsaturated fatty acids, retinoids, or other active compounds with more than one aliphatic double bond prone to oxidation.

30 Semifluorinated alkanes are linear or branched alkanes some of whose hydrogen atoms have been replaced by fluorine. In a preferred embodiment, the semifluorinated alkanes (SFAs) described are composed of at least one non-fluorinated hydrocarbon segment and at least one perfluorinated hydrocarbon

5 segment. Particularly useful are SFAs which have one non-fluorinated hydrocarbon segment attached to one perfluorinated hydrocarbon segment, according to the general formula  $F(CF_2)_n(CH_2)_mH$ , or two perfluorinated hydrocarbon segments separated by one non-fluorinated hydrocarbon segment, according to the general formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ .

10 Another nomenclature which is used herein refers to the above-mentioned SFAs having two or three segments as RFRH and RFRH<sub>F</sub>, respectively, wherein R<sub>F</sub> designates a perfluorinated hydrocarbon segment, R<sub>H</sub> designates a non-fluorinated segment. Alternatively, the compounds may be referred to as FnHm and FnHmFo, respectively, wherein F means a perfluorinated hydrocarbon segment, H means a  
15 non-fluorinated segment, and n, m and o is the number of carbon atoms of the respective segment. For example, F3H3 is used for perfluoropropylpropane. Moreover, this type of nomenclature is usually used for compounds having linear segments. Therefore, unless otherwise indicated, it should be assumed that F3H3 means 1-perfluoropropylpropane, rather than 2-perfluoropropylpropane, 1-  
20 perfluoroisopropylpropane or 2-perfluoroisopropylpropane.

20 Preferably, the semifluorinated alkanes according to the general formulas  $F(CF_2)_n(CH_2)_mH$  and  $F(CF_2)_n(CH_2)_m(CF_2)_oF$  have segment sizes ranging from 3 to 20 carbon atoms, i.e. n, m and o are integers independently selected from the range of 3 to 20. SFAs which are useful in the context of the present invention are also described  
25 in EP-A 965 334, EP-A 965329 and EP-A 2110126, the disclosure of which documents is incorporated herein.

30 In a further embodiment, the compositions of the invention comprise a semifluorinated alkane according to the formula  $F(CF_2)_n(CH_2)_mH$ , wherein n and m are integers independently selected from the range of 3 to 20. In another particular embodiment, n is an integer from the range of 3 to 8 and m is an integer from the range of 3 to 10. In yet another particular embodiment, the semifluorinated alkane is a compound according to the formula  $F(CF_2)_n(CH_2)_mH$  wherein n is an integer from the range of 6 to 20 and m is an integer from the range of 10 to 20. Most preferably, the semifluorinated alkane is a liquid. Preferred SFAs include, in particular, the  
35 compounds  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,

5  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ . Further preferred SFAs include, in particular,  $F(CF_2)_8(CH_2)_{10}H$  and  $F(CF_2)_{10}(CH_2)_{12}H$ .

In a further embodiment, the composition may comprise more than one SFA. Preferably, compositions comprising more than one SFA comprise at least one of SFAs selected from  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ . In another embodiment, the composition comprises at least two SFAs selected from compounds  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ , and at least one of  $F(CF_2)_8(CH_2)_{10}H$  and  $F(CF_2)_{10}(CH_2)_{12}H$ . In one of the preferred embodiments, the composition comprises a first semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$ , wherein  $n$  is an integer from the range of 3 to 8 and  $m$  is an integer from the range of 3 to 10 and a second semifluorinated alkane of the formula  $F(CF_2)_n(CH_2)_mH$  wherein  $n$  is an integer from the range of 6 to 20 and  $m$  is an integer from the range of 10 to 20.

As mentioned, the compositions comprise an active compound with more than 20 one aliphatic double prone to oxidation, wherein the active compound is dissolved, dispersed or suspended in a liquid vehicle comprising a SFA that is substantially free of water. In some embodiments, the liquid vehicle comprising a SFA may further comprise other organic liquids or other excipients, but is effectively free of water.

According to a particular embodiment, the active compounds useful in the 25 invention may be dissolved, that is, in complete solvation or solution in a liquid vehicle comprising a semifluorinated alkane. Alternatively, the active compounds may be dispersed or suspended in a liquid vehicle comprising a semifluorinated alkane. As used herein, dispersing means the formation of a system having at least one continuous (or coherent) phase and at least one discontinuous (or inner) phase 30 which is dispersed in the continuous phase. The term dispersion is understood to include colloidal systems in which the active compound is finely dispersed in the liquid phase. It is also understood that a suspension is a type of dispersion, in which the dispersed phase is in the solid state. The suspensions useful for practicing the invention are liquids, at least at physiological temperature, which means that the 35 continuous phase is liquid. Typically, suspensions are liquid at room temperature.

5       Liquid SFAs are chemically and physiologically inert, colourless and stable. Their typical densities range from 1.1 to 1.7 g/cm<sup>3</sup>, and their surface tension may be as low as 19 mN/m. SFAs of the  $F(CF_2)_n(CH_2)_mH$  type are insoluble in water but also somewhat amphiphilic, with increasing lipophilicity correlating with an increasing size of the non-fluorinated segment. Liquid SFAs of this type are being used  
10 commercially for unfolding and reapplying a retina, for long-term tamponade as vitreous humor substitute (H. Meinert et al., European Journal of Ophthalmology, Vol. 10(3), pp. 189-197, 2000), and as wash-out solutions for residual silicon oil after vitreo-retinal surgery. This and other applications have established SFAs as physiologically well tolerated compounds.

15       It has also been proposed that SFAs have high solubility for gases such as oxygen, and may act as oxygen carriers (US 6262126). Experimentally, they have been used as blood substitutes, as oxygen-carriers (H. Meinert et al., Biomaterials, Artificial Cells, and Immobilization Biotechnology, Vol. 21(5), pp. 583-95, 1993).

20       In contradiction to such facts and teachings, the inventors have found that liquid compositions comprising SFAs and the oxidation-sensitive active compounds, such as those described above, are surprisingly stable under ambient as well as non-ideal conditions (as illustrated by the examples further below).

25       The inventors have found that a method of stabilising a compound useful in the invention can simply comprise the step of dissolving, dispersing, or suspending an active compound of the invention in a liquid vehicle comprising a semifluorinated alkane, as no additional antioxidants or other active compound stabilizers, as are typically used for the formulation of such compounds are required. This simplification can be advantageous for various medical uses of the composition involving especially, frequent administration to tissues such as ocular or mucosal  
30 tissues. These generally have poor tolerability to a wide variety of antioxidants and stabilizers and generally do not benefit from frequent exposure to such excipients.

      In one embodiment, a method of stabilising a polyunsaturated fatty acid or an ester derivative thereof comprises the step of dissolving, dispersing or suspending the polyunsaturated fatty acid or an ester derivative thereof in a liquid vehicle

5 comprising a semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$  or of formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ , wherein n, m, and o are integers independently selected from the range of 3 to 20 and wherein the liquid vehicle is substantially free of water. In yet a further embodiment, said method of stabilisation comprises the step of dissolving, dispersing, or suspending a polyunsaturated fatty acid selected from an omega-3 or

10 an omega-6 fatty acid or ester derivatives, or mixtures thereof in a liquid vehicle comprising a semifluorinated alkane selected from  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ .

Further embodiments within the scope of the invention are stable liquid

15 compositions consisting of essentially one or more omega-3 fatty acids or derivatives thereof dissolved or dispersed in a liquid semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$  or of formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ , wherein n, m, and o are integers independently selected from the range of 3 to 20 and wherein the liquid vehicle is substantially free of water. Compositions comprising no further active compounds

20 other than omega-3 fatty acid or derivatives thereof are also within the scope of the invention. Preferred are stable liquid compositions consisting of one or more omega-3 fatty acid selected from  $\alpha$ -linolenic acid, docosahexaenoic acid, or eicosapentaenoic acid and ester derivatives thereof, wherein the omega-3 fatty acid or ester derivative thereof is dissolved or dispersed in a semifluorinated alkane useful in the invention, for instance,  $F(CF_2)_6(CH_2)_8H$ . In a further preferred embodiment, the liquid

25 composition consists essentially of docosahexaenoic acid and/or eicosahexaenoic acid or ester derivatives thereof dissolved in or dispersed in  $F(CF_2)_6(CH_2)_8H$ .

In a particular embodiment, liquid compositions comprising at least two or more semifluorinated alkanes further comprise one or more omega-3-fatty acid or

30 omega-3-fatty acid derivative, and/or omega-6 fatty acid or omega-6 fatty acid derivative.

Also described are methods of stabilizing a retinoid selected from retinol, retinoic acid, retinal, tretinoin, isotretinoin, and alitretinoin or derivatives thereof, or sirolimus, or a polyene compound, or ciclopirox olamine, comprising the step of

35 dissolving, dispersing or suspending said compounds in a liquid vehicle comprising a

5 semifluorinated alkane of formula  $F(CF_2)_n(CH_2)_mH$  or of formula  $F(CF_2)_n(CH_2)_m(CF_2)_oF$ , wherein n, m, and o are integers independently selected from the range of 3 to 20 and wherein the liquid vehicle is substantially free of water. In another embodiment, method of stabilising a retinoid selected from retinol, retinoic acid, retinal, tretinoin, isotretinoin, and alitretinoin or derivatives thereof, or

10 sirolimus, comprises the step of dissolving, dispersing or suspending said compounds in a liquid vehicle comprising a semifluorinated alkane selected from  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ .

In a further embodiment of the invention, liquid compositions comprising at least two or more semifluorinated alkanes further comprise at least one solubilized lipophilic vitamin or vitamin derivative.

The liquid compositions comprising an active compound with more than one aliphatic double bond prone to oxidation dissolved, dispersed or suspended in a liquid vehicle comprising a semifluorinated alkane as described previously, wherein the liquid vehicle is substantially free of water may comprise at least about 0.01-10 wt.% of active compound. Preferably, the compositions comprise at least about 1 wt.% of active compound relative to the weight of the composition, such as about 2 wt% or more, or at least 5 wt%.

The compositions of the present invention are useful as medicaments, in particular as medicaments topically administered to an eye or ophthalmic tissue, to the skin, or to the buccal, anal, vaginal, or nasal mucosa.

The SFA component of the liquid vehicle of these compositions exhibit properties rendering them particularly amenable for ophthalmic applications. Many of the SFAs as defined herein have refractive indices close to water. In one of the specific embodiments, the invention is therefore practised with an SFA whose refractive index is from 1.29 to 1.35, and in particular from about 1.30 to about 1.35 at 20°C.

5 Moreover, SFAs according to the invention exhibit a remarkable wetting and spreading behaviour by which they can rapidly and effectively spread over the corneal surface and conjunctiva. Wetting means the ability of a liquid to establish and maintain contact with a solid surface, resulting from intermolecular interactions when the two are brought together. The balance between adhesive and cohesive  
10 forces determines the degree of wetting. The higher the adhesive forces compared to the cohesive forces, the more a drop of liquid will spread across the surface of the solid material. Conversely, very high cohesive forces within the liquid will cause the drop to form a sphere, thus avoiding contact with the surface. Similarly, spreading may also occur at the interface of two liquids which are brought into contact with  
15 each other.

A measure for wetting and spreading is the contact angle  $\theta$ . The contact angle is the angle at which the liquid-vapour interface meets the solid-liquid or liquid-liquid interface. The tendency of a drop to spread out increases as the contact angle decreases. Thus, the contact angle provides an inverse measure of wettability.

20 A low contact angle of less than  $90^\circ$  indicates high wettability and/or spreading, whereas a higher contact angle indicates poor wettability and spreading. Perfect wetting and spreading results in a contact angle of  $0^\circ$ , also reported as no measurable contact angle.

25 SFAs useful according to the invention exhibit excellent wetting of various surfaces. For example, the contact angle of both F4H5 and F6H8 on tablets compressed from either trospium chloride or fenofibrate (150 mg of drug substance compressed at 15-20 kN to tablets of 13 mm in diameter) is not measurable, i.e. there is perfect wetting. It is noted that fenofibrate is an example of a hydrophobic, poorly water-soluble compound, whereas trospium chloride is hydrophilic and water-soluble. For comparison, the contact angle of purified water on the fenofibrate tablet  
30 was determined as  $92.5^\circ$ , i.e. the tablet was poorly wetted by water.

In addition, SFAs useful according to the invention are also capable of forming very small droplets when dispensed from a dropper such as an eye dropper. Without wishing to be bound by theory, it is believed that the small droplet size is a result of

5 an interplay of the SFA's unique properties in terms of their density, viscosity, and surface tension. In any case, it is believed that for topical administration into an eye a small drop or volume of administration is highly advantageous as the capability of the lacrimal sac to accept and hold fluid is extremely limited. In fact, it is very common that the administration of a conventional eye drop formulation based on water or oil

10 immediately leads to a discharge of a substantial fraction of the administered medicine as well as some tear fluid. At the same time, there is a risk that some of the administered dose will be taken up systemically via the nasolacrimal duct.

Also described is a means of formulating non-aqueous ophthalmic compositions which are microbiologically stable. Aqueous ophthalmic compositions are prone to bacterial contamination. In comparison, SFAs useful according to the invention have bacteriostatic properties and do not support microbial growth. Hence, it is possible to formulate preservative-free ophthalmic compositions which are better tolerable for many patients suffering from eye conditions that are readily exacerbated by preservatives or other excipients that may irritated through repeated exposure.

20 The compositions of the invention are thus very well suited for the topical administration to an eye or ophthalmic tissue. Ophthalmic tissue includes any surface of the eye anatomy that is, or can be (i.e. by non-surgical means) topically exposed. Preferably, the compositions are administered to the cornea or conjunctiva. The compositions are also preferably administered to the upper or lower eye lid margins, 25 meibomian gland ducts, eyelashes or any area of the eye or eye lid anatomy.

In particular, it is contemplated that these compositions can be used for the treatment of a condition or disease of an eye or ophthalmic tissue, such as inflammatory conditions of the ophthalmic tissue or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith. According to an embodiment 30 of the invention, a composition consisting of a semifluorinated alkane of the formula  $F(CF_2)_n(CH_2)_mH$ , wherein n is an integer from the range of 3 to 8 and m is an integer from the range of 3 to 10, and optionally one or more further excipients may be used as a medicine in the treatment of keratoconjunctivitis sicca (dry eye) or a symptom or condition associated therewith

5       Keratoconjunctivitis sicca is a complex, multifaceted disease or condition as described above. Aqueous-deficient DED, evaporative DED are within the scope of keratoconjunctivitis sicca and form specific subtypes thereof. Sjögren syndrome, lacrimal gland insufficiency, meibomian gland disease and meibomian gland dysfunction, and other conditions are also associated with keratoconjunctivitis sicca,  
10      being direct or indirect causes thereof, and may be treated with the composition of the invention.

Meibomian gland diseases cover a broad range of meibomian gland disorders including neoplasia and congenital disorders. Meibomian gland dysfunction, on the other hand is understood to be abnormalities of the meibomian glands which are  
15      often characterized by gland duct obstructions and/or changes (qualitative and/or quantitative) to the secretions of the glands. In general, conditions or disease states causing or leading to an abnormal, reduced or increased delivery of lipids to the tear film can give rise to keratoconjunctivitis sicca and the symptoms associated therewith.

20       Symptoms of keratoconjunctivitis sicca include a dry, scratchy, gritty, or sandy feeling in the eye; foreign body sensation; pain or soreness; stinging or burning; itching; increased blinking; eye fatigue; photophobia; blurry vision; redness; mucus discharge; contact lens intolerance; excessive reflex tearing. In addition to the symptoms of keratoconjunctivitis sicca as described, patients with meibomian gland  
25      dysfunction may also experience symptoms including itchiness, redness, swelling, pain or soreness, discharge accumulation or crusting specifically at the lid margins. It is understood that not all patients suffering from keratoconjunctivitis sicca exhibit all symptoms simultaneously. Hence, there is currently no uniform set of criteria for diagnosing the disease. It is also understood that patients may suffer from one or  
30      more subtypes of keratoconjunctivitis sicca, or one or more conditions or disease pathways causing keratoconjunctivitis sicca. It is however important to note that, within the scope of the present invention, any of the aspects, symptoms or pathophysiological consequences of dry eye disease may be addressed.

35       The compositions of the invention may also comprise one or more active compounds and/or further excipients as are required or useful. In an embodiment,

5 the compositions of the invention may further comprise an anti-inflammatory compound selected from the group of NSAIDs, corticosteroids, and immunomodulators. In particular, such compositions may be useful as a medicament for topical administration to an eye or ophthalmic tissue, especially as medicaments for treatment of inflammatory conditions of ophthalmic tissue or keratoconjunctivitis

10 sicca (dry eye), or a symptom or condition associated therewith. Preferred immunomodulators are calcineurin inhibitors (e.g. cyclosporin or tacrolimus) or mTOR inhibitors (e.g. sirolimus). Preferred NSAIDs include flurbiprofen, diclofenac, indometacin, bromfenac, nepafenac, ketoprofen, and ketorolac and salts and derivatives thereof. Preferred corticosteroids include prednisolone, loteprednol,

15 dexamethasone, hydrocortisone, rimexolone, fluorometholone and salts and derivatives thereof. These further active compounds are preferably also dissolved or dispersed in a liquid semifluorinated alkane of the invention.

In another embodiment, the compositions of the invention may also further comprise one or more flavan-3-ols such as catechins. Catechin or catechin isomers (e.g. epicatechin) and derivatives (for example, ester derivatives of catechin) are particularly preferred.

In terms of further excipients, if any, especially preferred are those that are biocompatible and are tolerated by the eye, and which are liquid and/or soluble and miscible in SFAs. In particular, excipients are preferably selected from lipids, oils, lubricants, lipophilic vitamins, viscosity agents, antioxidants, surfactants and mixtures of two or more thereof.

Examples of potentially useful lipids and oily excipients and which may be included in the compositions of the invention include triglyceride oils (e.g. soybean oil, olive oil, sesame oil, cotton seed oil, castor oil, sweet almond oil), mineral oil (e.g. 30 petrolatum and liquid paraffin), medium chain triglycerides (MCT), oily fatty acids, isopropyl myristate, oily fatty alcohols, esters of sorbitol and fatty acids, oily sucrose esters, oily cholesterol esters, oily wax esters, glycerophospholipids, sphingolipids, or any oily substance which is physiologically tolerated by the eye. Any synthetic, semi-synthetic or natural oily excipients which mimic or are structurally analogous or

5 related to the components naturally found in the tear film lipid layer are also within the scope of the invention.

Examples of potentially useful lipophilic vitamin excipients include vitamin E (e.g.  $\alpha$ -tocopherol) and their derivatives (e.g. tocotrienols) and esters (e.g. tocopherol acetate or tocopherol TPGS). In an embodiment, liquid compositions of the invention 10 may further comprise at least one lipophilic vitamin excipient that is completely solubilized.

Examples of potentially useful lubricants and/or viscosity agents include carboxymethyl cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, glycerol, polyvinyl alcohol, polyethylene glycol, propylene glycol, hyaluronic acid,

15 hydroxypropyl guar.

In one of the embodiments, the composition of the invention is free of surfactants. In an alternative embodiment, and depending on the specific active ingredient to be formulated, a small amount of a physiologically acceptable surfactant may be incorporated. Potentially useful surfactant excipients include in particular 20 non-ionic surfactants or amphiphilic lipids. Surfactants which are considered potentially useful include tyloxapol, poloxamers such as Pluronic F68LF or Lutrol F68, Pluronic L-G2LF and Pluronic L62D, polysorbates such as polysorbate 20 and polysorbate 80, polyoxyethylene castor oil derivatives, sorbitan esters, polyoxyl stearates, and mixtures of two or more thereof.

25 The composition may of course comprise further excipients as required or useful such as acids, bases, electrolytes, buffers, solutes, stabilisers, synergists, and - if required in a particular case - a preservative. In one of the embodiments, however, the composition is free of preservatives.

30 The compositions may be formulated to be administered as a liquid solution, gel, suspension, or spray. They may be prepared by commonly known techniques for the manufacture of said liquid solutions, gels, suspensions, or sprays.

Furthermore, described is a pharmaceutical kit comprising the composition as described above and a container holding the composition. Preferably, the container

5 which contains the composition has a dispensing means such as a dropping device adapted for topically administering the composition to the eye of a patient.

The term "comprising" as used in this specification means "consisting at least in part of". When interpreting each statement in this specification that includes the term "comprising", features other than that or those prefaced by the term may also be

10 present. Related terms such as "comprise" and "comprises" are to be interpreted in the same manner.

The following examples serve to illustrate the invention; however, these are not to be understood as restricting the scope of the invention.

## EXAMPLES

### Example 1

Solutions of the omega-3 fatty acid esters, eicosapentaenoic acid ethyl ester (EPA-EE) and docosahexenoic acid ethyl ester (DHA-EE), were prepared in  $F(CF_2)_6(CH_2)_8H$  at concentrations of 1.0 wt% and 5.0 wt%.

Solutions containing a mixture of 1 wt% each of DHA-EE and EPA-EE (total 2.0 wt% of active compound) and 5.0 wt% of a combination of the DHA-EE and EPA-EE, at a ratio of approx. 2:3 in  $F(CF_2)_6(CH_2)_8H$  were also prepared. The solutions were filled in crimped vials and stored at 2-8°C, at 25°C/60% RH, and at 40°C/75% RH for the purpose of conducting a 6 month stability programme. The results obtained after a 13-week period are reported herein.

Samples were taken from the vials at intervals for GC analysis (See Tables 1-3 for result) of the compositions. The remaining percentages of other components detected by GC (FID) but not shown here in the tables are attributed to impurities already present in the original samples of omega-3 fatty ester or the SFA vehicle before formulation. No rise in the level of impurities was observed over the course of the stability program.

No significant observable changes in the content of the omega-3 fatty acid esters were detected under all temperatures tested. Results were more or less also comparable across different batches at all storage conditions. In the following tables,

the content of the active ingredient is presented as percentage relative to the initial content.

Table 1: Content of active compound (%) after storage at 2-8°C

Compound	After 4 weeks
1.0 wt% EPA-EE	99.6
1.0 wt% DHA-EE	99.5
5.0 wt% EPA-EE	99.9
5.0 wt% DHA-EE	101.5
2.0 wt% EPA-EE/DHA-EE	100.2/99.7
5.0 wt% EPA-EE/DHA-EE	100.3/99.5

5 Table 2: Content of active compound (%) after storage at: 25°C/60% RH

Compound	After 4 weeks	After 13 weeks
1.0 wt% EPA-EE	99.6	99.0
1.0 wt% DHA-EE	99.6	99.4
5.0 wt% EPA-EE	99.8	99.5
5.0 wt% DHA-EE	99.8	103.2 <sup>1</sup>
2.0 wt% EPA-EE/DHA-EE	100.2/99.6	100.5/98.6
5.0 wt% EPA-EE/DHA-EE	101.1/98.0	103.5/92.6


<sup>1</sup> Slight increase probably due to impurities with elution times similar to DHA-EE

Table 3: Content of active compound (%) after storage at 40°C/75% RH

Compound	1 week	2 weeks	4 weeks	8 weeks	13 weeks
1.0 wt% EPA-EE	99.6	99.6	99.6	99.4	97.3
1.0 wt% DHA-EE	99.9	99.8	99.8	99.6	99.1
5.0 wt% EPA-EE	99.9	99.9	99.8	99.4	98.7
5.0 wt% DHA-EE	99.9	100.0	100.0	99.9	100.0
2.0 wt% EPA-EE/DHA-EE	101.8/98.2	100.3/99.6	100.5/99.0	100.8/98.3	100.8/98.2
5.0 wt% EPA-EE/DHA-EE	100.9/98.3	101.0/98.0	101.6/96.6	103.1/93.1	105.4/88.4

## Claims

1. A stable liquid composition comprising a polyunsaturated fatty acid or a derivative thereof dissolved in a liquid vehicle comprising a semifluorinated alkane of formula




wherein n and m are integers independently selected from the range of 3 to 20, and wherein the liquid vehicle is substantially free of water, wherein the polyunsaturated fatty acid or derivative thereof is an omega-3 or omega-6 fatty acid or derivative thereof, or a mixture of omega-3 or omega-6 fatty acids or derivatives thereof, wherein the derivative(s) are selected from esters, resolvins and neuroprotectins.


2. The composition of claim 1, wherein the polyunsaturated fatty acid derivative is an ester derivative.
3. The composition of claim 1 or 2, wherein the polyunsaturated fatty acid or derivative thereof is an omega-3 fatty acid or derivative, and wherein the omega-3 fatty acid or derivative is the only active ingredient in the composition.
4. The composition of claim 1 or 2, wherein the omega-3 fatty acid is selected from  $\alpha$ -linolenic acid, docosahexaenoic acid, eicosapentaenoic acid, and ester derivatives thereof.
5. The composition of claim 1 or 2, wherein the omega-3 fatty acid is a derivative of eicosapentaenoic acid or docosahexaenoic acid or is selected from a resolvin or a neuroprotectin.
6. The composition of claim 4, wherein the omega-3 fatty acid ester derivative is selected from docosahexaenoic acid ethyl ester or eicosapentaenoic acid ethyl ester.
7. The composition of claim 3 or 4, consisting essentially of one or more omega-3 fatty acids or ester derivatives thereof dissolved or dispersed in the liquid semifluorinated alkane.

8. The composition of claim 7, consisting essentially of docosahexaenoic acid and /or eicosapentaenoic acid or alkyl ester derivatives thereof dissolved or dispersed in  $F(CF_2)_6(CH_2)_8H$ .
9. The composition of anyone of the preceding claims, comprising about 0.01 -10 wt.-% of the polyunsaturated fatty acid or derivative thereof.
10. The composition of any one of the preceding claims, wherein the semifluorinated alkane is selected from  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ , or wherein the semifluorinated alkane is selected from  $F(CF_2)_8(CH_2)_{10}H$  and  $F(CF_2)_{10}(CH_2)_{12}H$ .
11. The composition of any one of the preceding claims, further comprising a lipid or oily excipient selected from triglyceride oils, mineral oil, medium chain triglycerides, oily fatty acids, isopropyl myristate, oily fatty alcohols, esters of sorbitol and fatty acids, oily sucrose esters, oily cholesterol esters, oily wax esters, glycerophospholipids, and sphingolipids.
12. The composition of any one of claims 1 to 2, 4 or 10, further comprising an anti-inflammatory compound selected from the group of NSAIDs, corticosteroids and immunomodulators.
13. The composition of any one of the preceding claims for use as a medicament.
14. The composition for use as a medicament according to claim 13, wherein the medicament is topically administered to an eye or ophthalmic tissue, to the skin, or to the buccal, anal, vaginal, or nasal mucosa.
15. The composition for use as a medicament according to claim 13 or 14, wherein the medicament is used for the treatment of a condition or disease of an eye or ophthalmic tissue.
16. The composition for use as a medicament according to claim 15, wherein the condition or disease of an eye or ophthalmic tissue is inflammatory conditions of the ophthalmic tissue or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith.

17. A method of stabilising a polyunsaturated fatty acid selected from an omega-3 or an omega-6 fatty acid or ester derivatives thereof, or mixtures thereof, comprising the step of dissolving, dispersing or suspending the polyunsaturated fatty acid in a liquid vehicle comprising a semifluorinated alkane of formula



or of formula



wherein n, m, and o are integers independently selected from the range of 3 to 20 and wherein the liquid vehicle is substantially free of water.

18. The method of claim 17, wherein the semifluorinated alkane is selected from  $F(CF_2)_4(CH_2)_5H$ ,  $F(CF_2)_4(CH_2)_6H$ ,  $F(CF_2)_4(CH_2)_8H$ ,  $F(CF_2)_6(CH_2)_4H$ ,  $F(CF_2)_6(CH_2)_6H$ ,  $F(CF_2)_6(CH_2)_8H$ , and  $F(CF_2)_6(CH_2)_{10}H$ ; or wherein the semifluorinated alkane is selected from  $F(CF_2)_8(CH_2)_{10}H$  and  $F(CF_2)_{10}(CH_2)_{12}H$ .
19. A method of treating a an inflammatory condition or disease of the ophthalmic tissue, or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith, the method comprising topically administering a composition as claimed in any one of claims 1 to 12 to an eye or ophthalmic tissue of a subject in need thereof.
20. Use of a stable liquid composition as claimed in any one of claims 1 to 12 in the manufacture of a medicament for use in the topical ophthalmic treatment of an inflammatory condition or disease of the ophthalmic tissue or keratoconjunctivitis sicca (dry eye) or symptoms or conditions associated therewith.