



US 20080138334A1

(19) **United States**(12) **Patent Application Publication**
SABBADINI et al.(10) **Pub. No.: US 2008/0138334 A1**(43) **Pub. Date: Jun. 12, 2008**(54) **IMMUNE-DERIVED MOIETIES REACTIVE
AGAINST BIOACTIVE LIPIDS, AND
METHODS OF MAKING AND USING SAME****Publication Classification**

- (51) **Int. Cl.**
A61K 39/395 (2006.01)
C07K 16/18 (2006.01)
- (52) **U.S. Cl.** **424/133.1**; 530/387.3; 530/388.1;
424/141.1; 530/387.1

- (76) **Inventors:** **Roger A. SABBADINI**, Lakeside,
CA (US); **William A. Garland**, San
Clemente, CA (US); **Genevieve
Hansen**, San Diego, CA (US)

Correspondence Address:
Biotechnology Law Group
c/o Portfolioip
P.O. Box 52050
Minneapolis, MN 55402


- (21) **Appl. No.:** **11/755,485**
- (22) **Filed:** **May 30, 2007**

Related U.S. Application Data


- (60) Provisional application No. 60/810,185, filed on May 31, 2006, provisional application No. 60/835,569, filed on Aug. 4, 2006, provisional application No. 60/923,644, filed on Apr. 16, 2007.

(57) **ABSTRACT**

Compositions and methods for producing monoclonal antibodies and their derivatives reactive against bioactive lipid targets are described. These compositions include derivatized lipids, each of which comprises a bioactive lipid that having a polar head group and at least one hydrocarbon chain (e.g., a lysolipid such as lysophosphatidic acid or sphingosine-1-phosphate) in which a carbon atom has been derivatized with a pendant reactive group; immunogens made by linking a derivatized lipid to a carrier moiety (e.g., a carrier protein, polyethylene glycol, colloidal gold, alginate, or a silicone bead); monoclonal antibodies and derivatives produced by immunizing an animal with such an immunogen; and therapeutic and diagnostic compositions containing such antibodies and antibody derivatives. Methods for making such derivatized lipids, immunogens, and monoclonal antibodies and derivatives, methods for detecting such antibodies once generated, and therapeutic and diagnostic methods for using such antibodies and derivatives, are also described.




Synthesis of Typical Thiolated S1P-Related Antigen




Synthesis of Typical Thiolated S1P-Related Antigen

FIG. 1a




Synthesis of Typical Thiolated S1P-Related Antigen (Continued)

FIG.1b




Synthesis of Typical Thiolated S1P-Related Antigen (Continued)

FIG. 1c




Synthesis of Typical Protected Thiolated Fatty Acid

FIG.2a




Synthesis of Typical Thiolated Fatty Acid

FIG.2b



Synthesis of Typical Thiolated LPA Hapten

FIG.3a



Synthesis of Typical Thiolated LPA Hapten (Continued)

FIG.3b




FIG.4a




FIG. 4b




FIG.5

Figure 6

V_H

Q A H L Q Q S D A E L V K P G A S V K I S C K V S G F I F I D H
T I H W M K Q R P E Q G L E W I G C I S P R H D I T K Y N E M
F R G K A T L T A D K S S T T A Y I Q V N S L T F E D S A V Y F
C A R G G F Y G S T I W F D F W G Q G T T L T V S

V_L

E T T V T Q S P A S L S M A I G E K V T I R C I T T T D I D D D M
N W F Q Q K P G E P P N L L I S E G N I L R P G V P S R F S S
S G Y G T D F L F T I E N M L S E D V A D Y Y C L Q S D N L P
F T F G S G T K L E I K

Figure 7

pATH50: Nucleotide Sequence: Chimeric Light Chain

```

15  atgattgcct ctgctcagtt ccttggtctc ctgttgctct gttttaaagg taaccagatgt gaacaaactg tgaccacagtc tccagcatcc ctgtccatgg
    Leader
>>.....Sphingomab Chimeric Light Chain.....>
115 ctataggaga aaaagtcacc atcagatgca taaccaccac tgatatgtat gatgatatga actggttcca gcagaagcca ggggaacctc ctacactcct
    CDR1
>.....Sphingomab Chimeric Light Chain.....>
215 tatttccgaa ggcaatattc ttctctctgg agtcccatcc cgattctcca gcagtggcta tggtaacagac ttcttttta caattgaaaa catgctctca
    CDR2
>.....Sphingomab Chimeric Light Chain.....>
315 gaagatgttg cagattacta ctgtttgcag agtgataact taccattcac gtctcgctcg gggacaaaagt tggaaaaataa acgtgagtg
    CDR3
>.....Sphingomab Chimeric Light Chain.....>>

```

pATH50: Chimeric Light Chain protein sequence

```

1  miasaqflgl lllcfqgtrc ettvtqspas lsmaigekvt ircittddid ddmwffqkp geppnllise gnllrpgvps rfsssgygtd flftienmls
101 edvadyyciq sdnlpftfqs gtkleikre

```

Nucleotide alignment of pATH50 plasmid mutations to replace the methionine residue:

```

pATH50      1 agcttgccgccaccatgattgcctctgctcagttcctctctctgttggtctctctgttgcctctgttttcaaggtaaccagatgtgaaacaactgtgacccagttctccagc
pATH52C1    416 .....
pATH53C1    416 .....
pATH54C1    408 .....
pATH55C1    408 .....

pATH50      101 atccctgtccatggctataggagaaaaagtccaccatcagatgcataaccaccactgatattgatgatgatataactgaactgtgtccagcagaagccaggggaa
pATH52C1    516 .....g.....
pATH53C1    516 .....c.t.....
pATH54C1    508 .....ggt.....
pATH55C1    508 .....cc.....

```

PATH50	201	cctcctaacctocttatttccgaaggcaataattcttctgagtgagtcoccatcoccgattctccagcagtggtatggtacagactttcttttacaattg
PATH52C1	616	
PATH53C1	616	
PATH54C1	608	
PATH55C1	608	
PATH50	301	aaacatgctctcagaagatgttgacagattactactgtttgcagagtgataactaccattcaggttcggctcggggacaaaagttggaaaaataaaacgtga
PATH52C1	716	
PATH53C1	716	
PATH54C1	708	
PATH55C1	708	
PATH50	401	gtg
PATH52C1	816	...
PATH53C1	816	...
PATH54C1	808	...
PATH55C1	808	...

Protein alignment of pATH50 plasmid mutations to replace the methionine residue:

PATH50	15	miasaqflgl1lllcfqgtrcettvtgspaslsmaigekvtircitttddiddmnwfgqkpggeppnlllsegnilrpgvprrfsssgygtcdf1ftienmls
PATH52C1	430	v.....
PATH53C1	430	l.....
PATH54C1	422	g.....
PATH55C1	422	p.....
PATH50	315	edvadyyclqsdnlpftfgsgtkleikre
PATH52C1	730	
PATH53C1	730	
PATH54C1	722	
PATH55C1	722	

730 gaagatgttg cagattacta ctgttttgcag atgtataact taccattcac gtctgcgtct gggacaaagt tggaaataaa acgtgagtg
 CDR3 FW4
 e d v a d y y c l q s d n l p f t f g s g t k l e i k r e
 > SoHingomab Chimeric Light Chain >

PATH54

422 atgattgcct ctgctcagtt ccttggtctc ctgttgcct gttttcaagg taaccagatg gaacaactg tgaccagtc tccagcatcc ctgtccatgg
Leader
m i a s a q f l g l l l c f q g t r c e t t v t q s p a s l s m
>>.....Sphingomab Chimeric Light Chain.....>

522 ctataggaga aaaagtcacc atcagatgca taaccaccac tgatattcat gatgatgcta actggttcca gcagaagcca ggggaacctc ctaacctctt
CDR1
a i g e k v t i r c i t t d i d d g n w f q q k p g e p n l
>.....Sphingomab Chimeric Light Chain.....>
M33G >>>

622 tatttcgaa ggcaatattc ttgctcctgg agtcccatcc cgattctcca gcagtggcta tggtaacagac ttttttttta caattgaaaa catgctctca
CDR2
l i s e g n i l r p g v p s r f s s g y g t d f l f t i e n m l s
>.....Sphingomab Chimeric Light Chain.....>

722 gaagatgttg cagattacta ctgtttgcag agtgataact taccattcac gttcggctog gggacaaaagt tggaataaaa acgtgagtg
CDR3
e d v a d y y c l q s d n l p f t f g s g t k l e i k r e
>.....Sphingomab Chimeric Light Chain.....>

PATH55

422 atgattgcct ctgctcagtt ccttggtctc ctgttgcct gttttcaagg taaccagatg gaacaactg tgaccagtc tccagcatcc ctgtccatgg
Leader
m i a s a q f l g l l l c f q g t r c e t t v t q s p a s l s m
>>.....Sphingomab Chimeric Light Chain.....>

522 ctataggaga aaaagtcacc atcagatgca taaccaccac tgatattcat gatgatccta actggttcca gcagaagcca ggggaacctc ctaacctctt
CDR1
a i g e k v t i r c i t t d i d d d p n w f q q k p g e p n l
>.....Sphingomab Chimeric Light Chain.....>
M33P >>>

622 tatttcgaa ggcaatattc ttgctcctgg agtcccatcc cgattctcca gcagtggcta tggtaacagac ttttttttta caattgaaaa catgctctca
CDR3
l i s e g n i l r p g v p s r f s s g y g t d f l f t i e n m l s
>.....Sphingomab Chimeric Light Chain.....>

722 gaagatgttg cagattacta ctgtttgcag agtgataact taccattcac gttcggctog gggacaaaagt tggaataaaa acgtgagtg
CDR3
e d v a d y y c l q s d n l p f t f g s g t k l e i k r e
>.....Sphingomab Chimeric Light Chain.....>

[illegible]

PATH10	101	gagcttcagtgaagata	tctctgaaggtttctgggttcattttcattgaccatactattcactggatgaagcagagggcctgaacagggcctcgaa	tgga
PATH11C1	515			
PATH12C2	515			
PATH13C2	371			
PATH14C1	478			
PATH15C1	514			
PATH16C1	365			
PATH17C1	382			
PATH10	201	cggtatgtatttctccagacatgatatattactaaatacgaatgagatggttcaggggcaaggccaccctgactgcagacaagtctctccactacagcctacata		
PATH11C1	615	...gc.....		
PATH12C2	615	...c.....		
PATH13C2	471	...g.....		
PATH14C1	578	...a.....		
PATH15C1	614	...c.....		
PATH16C1	465	...t.....		
PATH17C1	482	...atg.....		
PATH10	301	caagtcaacagtctgacatttgaaagactctgcagctctatttctgtgcaagaggggggttctacggtagtactatctggtttgacttttggggcccaaggca		
PATH11C1	715			
PATH12C2	715			
PATH13C2	571			
PATH14C1	678			
PATH15C1	714			
PATH16C1	565			
PATH17C1	582			
PATH 10	401	ccactctcacagtctctctcagcctccaccaaggcc		
PATH11C1	815			
PATH12C2	815			
PATH13C2	671			
PATH14C1	778			
PATH15C1	814			
PATH16C1	665			
PATH17C1	682			

Protien alignment of pATH10 plasmid mutations to replace the cysteine residue:

```
PATH10      1 mawswvflfflsvtgvsqahlgqsdaelvkgpaskvsksvsgffidhtihwmkqrpeqglewigci sprhdi ttkynemfrgkatl tadssttayl
PATH11C1    415 .....
PATH12C2    415 .....a.....
PATH13C2    271 .....s.....
PATH14C1    378 .....w.....
PATH15C1    414 .....y.....
PATH16C1    265 .....r.....
PATH17C1    282 .....f.....m.....

PATH10      301 qvnsltfedsavvfcaargfygstiwdfwgggttlvtvssastkg
PATH11C1    715 .....
PATH12C2    715 .....
PATH13C2    571 .....
PATH14C1    678 .....
PATH15C1    714 .....
PATH16C1    565 .....
PATH17C1    582 .....
```

PATH11

```
415 atggcatgga gctgggtctt tctttcttc ctgtcagtaa ctacggcggt ccactccag gctcacctgc aacagctcta cgctgaattg gtgaaacctg
>>.....leader.....
>>.....m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>>.....Sphingomab Chimeric Heavy Chain (path11C1).....>

515 gagcttcagt gaagatatcc tgcaagggtt ctgggttcat ttctattgac catattattc actggatgaa gcagaggcgt gaacagggcc tgaatggat
FW1 CDR1 FW2
>>.....g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>>.....Sphingomab Chimeric Heavy Chain (path11C1).....>

615 cggagctatt totcccagac atgatattac taaataacaat gagatgttca ggggcaaggc cacctgact gcagacaagt cctccactac agcctacata
CDR2
>>.....i g a i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>>.....Sphingomab Chimeric Heavy Chain (path11C1).....>
C51A >>>

715 caagtcaca gtctgacatt tgaagactct gcagctctatt ttgtgcaag aggggggttc tacggtagta ctatctgtt tgacttttg ggccaaggca
FW3 CDR3
>>.....q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>>.....Sphingomab Chimeric Heavy Chain (path11C1).....>
```

815 ccaactctcac agtctcctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>.....> Sphingomab Chimeric Heavy Chain (path11C1)

PATH12

415 atggcatgga.gctgggtctt.tctctcttc.ctgtcagtaa.ctaccggcgt.ccaactccag.gctcaactgc.aacagtctga.cgtgaattg.gtgaacactg
Leader
m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>.....> Sphingomab Chimeric Heavy Chain (path12C2)

515 gagcttcagt gaagatatcc tgcaagggttt ctggttcat ttctattgac catactattc actggatgaa gcagaggcct gaacagggcc tcgaatggat
FW1 CDR1 FW2
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....> Sphingomab Chimeric Heavy Chain (path12C2)

615 cggatctatt tctccacagc atgatattac taaatacaat gagatgttca gggggaaggc caccctgact gcagacaagt cctccactac agcctacata
CDR2
i g s i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....> Sphingomab Chimeric Heavy Chain (path12C2)

C51S >>>

715 caagtcaaca gtctgacatt tgaagactot gcagtctatt tctgtcaag aggggggttc tacggtagta ctatctggtt tgacttttgg ggccaaggca
FW3 CDR3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....> Sphingomab Chimeric Heavy Chain (path12C2)

815 ccaactctcac agtctcctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>.....> Sphingomab Chimeric Heavy Chain (path12C2)

PATH13

271 atggcatgga.gctgggtctt.tctctcttc.ctgtcagtaa.ctaccggcgt.ccaactccag.gctcaactgc.aacagtctga.cgtgaattg.gtgaacactg
Leader
m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>.....> Sphingomab Chimeric Heavy Chain (path13C2)

371 gagcttcagt gaagatatcc tgcaagggttt ctggttcat ttctattgac catactattc actggatgaa gcagaggcct gaacagggcc tcgaatggat
FW1 CDR1 FW2
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....> Sphingomab Chimeric Heavy Chain (path13C2)

471 cggatgatt tctccagac atgatattac taataacaat gagatattca ggggcaagg caccctgact gcagacaagt cctccactac agcctacata
CDR2
i g w i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....Sphingomab Chimeric Heavy Chain (path13C2).....>
C51W >>>

571 caagtcaca gtctgacatt tgaagactct gcagctctatt tctgtgcaag aggggggttc tacggtagta ctatctggtt tgacttttgg ggccaaggca
FW3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....Sphingomab Chimeric Heavy Chain (path13C2).....>

671 ccactctcac agtctctctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>.....Sphingomab Chimeric Heavy Chain (path13C2).....>

pATH14

378 atggcatoga gctgggtctt tctctcttc ctgtcagtaa ctaccggcgt ccactcccag gctcacctgc aacagtctga cgctgaattg gtgaaacctg
Leader
m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>.....Sphingomab Chimeric Heavy Chain.....>

478 gagcttcagt gaagatatcc tgcaaggttt ctggcttcat tttcattgac catactattc actggatgaa gcagagggcct gaacagggcc tcgaatggat
FW1
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....Sphingomab Chimeric Heavy Chain.....>

578 cggatatatt tctccagac atgatattac taataacaat gadatgttca ggggcaaggc caccctgact gcagacaagt cctccactac agcctacata
CDR2
i g y i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....Sphingomab Chimeric Heavy Chain.....>
C51Y >>>

678 caagtcaca gtctgacatt tgaagactct gcagctctatt tctgtgcaag aggggggttc tacggtagta ctatctggtt tgacttttgg ggccaaggca
FW3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....Sphingomab Chimeric Heavy Chain.....>

778 ccactctcac agtctctctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>...Sphingomab Chimeric Heavy Chain...>

PATH15

414 atggaatgga..gctgggtctt..tctctcttc..ctgtcaagtaa..ctaacggcgt..ccactccag gctcacctgc aacagttcga cgtgaattg gtgaacactg
Leader
m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>>.....Sphingomab Chimeric Heavy Chain.....>

514 gagcttcagt gaagatatoc tgoaaggttt ctggcttcct ttctattgac cactactatc actggatgaa gcagaggcct gaacagggcc tcgaatggat
FW1 CDR1 FW2
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....Sphingomab Chimeric Heavy Chain.....>

614 cggacgtatt tctccagac atgatattac taaatacaat gagatgttca ggggcaaggc cacctgact gcagacaagt cctccactac agcctacata
CDR2
i g r i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....Sphingomab Chimeric Heavy Chain.....>
C51R >>>

714 caagtcaaca gtctgacatt tgaagactct gcagctctatt totgtgcaag aggggggttc taggttagta ctatctggtt tgacttttgg ggccaaggca
FW3 CDR3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....Sphingomab Chimeric Heavy Chain.....>

814 ccactctcac agtctctcca gctccacca agggcc
FW4
t t l t v s s a s t k g
>...Sphingomab Chimeric Heavy Chain...>

PATH16

265 atggaatgga..gctgggtctt..tctctcttc..ctgtcaagtaa..ctaacggcgt..ccactccag gctcacctgc aacagttcga cgtgaattg gtgaacactg
Leader
m a w s w v f l f f l s v t t g v h s q a h l q q s d a e l v k p
>>.....Sphingomab Chimeric Heavy Chain.....>

365 gagcttcagt gaagatatoc tgoaaggttt ctggcttcct ttctattgac cactactatc actggatgaa gcagaggcct gaacagggcc tcgaatggat
FW1 CDR1 FW2
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....Sphingomab Chimeric Heavy Chain.....>

465 cggatttatt tctccagac atgatattac taaatacaat gagatgttca ggggcaaggc cacctgact gcagacaagt cctccactac agcctacata
CDR2
i g f i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....Sphingomab Chimeric Heavy Chain.....>
C51F >>>

565 caagtcaaca gtotgacatt tgaagactot gcagtcatt tctgtgcaag aggggggttc tacgtagta ctatctggtt tgacttttg ggccaaggca
FW3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....Sphingomab Chimeric Heavy Chain.....>

665 ccactctcac agtctctctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>...Sphingomab Chimeric Heavy Chain...>

PATH17

282 ataggcataga gctgggtctt ttcttctctt ctgtcagtaa ctacggcgt caactaccag gctcacctgc aacagtcgtga cgctgaattg gtgaacctg
Leader
m a w s w v f l f l s v t t g v h s q a h l q q s d a e l v k p
>.....Sphingomab Chimeric Heavy Chain.....>

382 gagcttcagt gaagatatcc tgcagggtt ctggttcat ttctattgac catactattc actggatgaa gcagaggcct gaacagggcc tcgaatggat
FW1
g a s v k i s c k v s g f i f i d h t i h w m k q r p e q g l e w
>.....Sphingomab Chimeric Heavy Chain.....>

482 cggaatgatt tctccagac atgatattac taaataacaat gagatgttca ggggcaagg caccctgact gcagacaagt cctccactac agcctacata
CDR2
i g m i s p r h d i t k y n e m f r g k a t l t a d k s s t t a y i
>.....Sphingomab Chimeric Heavy Chain.....>
C51M >>>

582 caagtcaaca gtotgacatt tgaagactot gcagtcatt tctgtgcaag aggggggttc tacgtagta ctatctggtt tgacttttg ggccaaggca
FW3
q v n s l t f e d s a v y f c a r g g f y g s t i w f d f w g q g
>.....Sphingomab Chimeric Heavy Chain.....>

682 ccactctcac agtctctctca gcctccacca agggcc
FW4
t t l t v s s a s t k g
>...Sphingomab Chimeric Heavy Chain...>




FIG.8

S1P binding by chimeric and mouse 306D antibodies.




FIG.9

IMMUNE-DERIVED MOIETIES REACTIVE AGAINST BIOACTIVE LIPIDS, AND METHODS OF MAKING AND USING SAME

RELATED APPLICATIONS

[0001] This patent application claims priority to U.S. provisional patent application Ser. No. 60/810,185, filed 31 May 2006 (attorney docket number LPT-3100-PV), U.S. provisional patent application Ser. No. 60/835,569, filed 4 Aug. 2006 (attorney docket number LPT-3100-PV2), and U.S. provisional patent application Ser. No. 60/923,644, filed 16 Apr. 2007 (attorney docket number LPT-3100-PV3). These applications are hereby incorporated by reference in their entirety for any and all purposes.

GOVERNMENT FUNDING

[0002] This invention was funded at least in part by funds supplied by the U.S. government pursuant to grant application NCI 2R44CA110298-2. As a result, the U.S. government may have certain rights in the inventions described herein.

TECHNICAL FIELD

[0003] The present invention relates to monoclonal antibodies, and methods for generating antibodies against immunogens that comprise a bioactive lipid molecule that plays a role in human and/or animal disease as a signaling molecule. One particular class of signaling bioactive lipids that can be addressed in accordance with the invention is lysolipids. Particularly preferred signaling lysolipids are sphingosine-1-phosphate (S1P) and the various lysophosphatidic acids (LPAs). The antibodies of the invention can be further modified to make them suitable for use in a particular animal species, including humans, without eliciting a neutralizing immune response. Such antibodies, and derivatives and variants thereof, can be used in the treatment and/or prevention of various diseases or disorders through the delivery of pharmaceutical compositions that contain such antibodies, alone or in combination with other therapeutic agents and/or treatments. In addition, the antibodies can be also be used to detect bioactive signaling lipids in biologic samples, thereby providing useful information for many purposes including, but not limited to, the diagnosis and/or prognosis of disease and the discovery and development of new treatment modalities that modify the production and or actions of the particular targeted lipid. The diseases or conditions to be affected by the compositions of the invention include, but are not limited to, diseases that have hyperproliferation, angiogenesis, inflammation, fibrosis, and/or apoptosis as part of their underlying pathology.

BACKGROUND OF THE INVENTION

1. Introduction

[0004] The following description includes information that may be useful in understanding the present invention. It is not an admission that any such information is prior art, or relevant, to the presently claimed inventions, or that any publication specifically or implicitly referenced is prior art or even particularly relevant to the presently claimed invention.

2. Background

[0005] A. Bioactive Signaling Lipids

[0006] Lipids and their derivatives are now recognized as important targets for medical research, not as just simple structural elements in cell membranes, solubilizing agents,

feedstock for vitamins or hormones or as a source of energy for β -oxidation, glycolysis or other metabolic processes. In particular, certain bioactive lipids function as signaling mediators important in animal and human disease. Although most of the lipids of the plasma membrane play an exclusively structural role, a small proportion of them are involved in relaying extracellular stimuli into cells. "Lipid signaling" refers to any of a number of cellular signal transduction pathways that use bioactive lipids as first or second messengers, including direct interaction of a lipid signaling molecule with its own specific receptor. Lipid signaling pathways are activated by a variety of extracellular stimuli, ranging from growth factors to inflammatory cytokines, and regulate cell fate decisions such as apoptosis, differentiation and proliferation. Research into bioactive lipid signaling is an area of intense scientific investigation as more and more bioactive lipids are identified and their actions characterized.

[0007] Examples of bioactive lipids include the eicosanoids derived from arachidonic acid (including the eicosanoid metabolites such as the HETEs, cannabinoids, leukotrienes, prostaglandins, lipoxins, epoxyeicosatrienoic acids, and isoeicosanoids), non-eicosanoid cannabinoid mediators, phospholipids and their derivatives such as phosphatidic acid (PA) and phosphatidylglycerol (PG) and cardiolipins as well as lysophospholipids such as lysophosphatidylcholine (LPC) and various lysophosphatidic acids (LPA). Bioactive signaling lipid mediators also include the sphingolipids such as ceramide, ceramide-1-phosphate, sphingosine, sphinganine, sphingosylphosphorylcholine (SPC) and sphingosine-1-phosphate (S1P). Sphingolipids and their derivatives represent a group of extracellular and intracellular signaling molecules with pleiotropic effects on important cellular processes. Other examples of bioactive signaling lipids include phosphatidylinositol (PI), phosphatidylethanolamine (PEA), diacylglyceride (DG), sulfatides, gangliosides, and cerebroside.

[0008] As expected, biological lipids (i.e., lipids that occur in nature, particularly in living organisms) are typically non-immunogenic or very weakly immunogenic. As such, lipids have traditionally been considered to be poor targets for antibody-based therapeutic and diagnostic/prognostic approaches. The literature contains a report of a monoclonal antibody that targets a derivatized form of phosphatidylserine (PS) conjugated to a carrier protein. Phosphatidylserine is a plasma membrane aminophospholipid. Loss of membrane lipid sidedness, in particular the emergence of phosphatidylserine at the cell surface, results in the expression of altered surface properties that modulates cell function and influences the cells interaction with its environment [Zwaal and Schroit, (1997) *Blood*, 89:1121-1132]. For example, PS redistributes from the cell membrane's inner leaflet (its normal location) to the outer leaflet during apoptosis.

[0009] Diaz, Balasubramanian and Schroit [*Bioconj. Chem.* (1998) 9:250-254] disclose production of lipid antigens that elicit specific immune responses against PS. The covalent coupling of PS to a protein carrier (BSA) via the lipid's fatty acyl side chain preserves the PS head group intact as an epitope. Schroit (U.S. Pat. No. 6,300,308, U.S. Pat. No. 6,806,354) discloses antibodies that specifically bind to phosphatidylserine (PS) or a phosphatidylcholine (PC)/polypeptide or a PS/polypeptide conjugate, that are made by administering a PS/polypeptide conjugate or a PC/polypeptide conjugate to an animal. Methods for detecting PS, a PC/polypeptide or a PS/polypeptide conjugate are also dis-

closed. Methods for making an antibody that specifically binds to PS by administering to an animal a pharmaceutical composition comprising a PS/polypeptide conjugate composition are also disclosed, as are methods for treating cancer in the animal to which the conjugate is administered, i.e., as a cancer vaccine. Also disclosed is induction of autoimmunity for the therapy of cancer by immunization of animals with β 2-glycoprotein I/lipid complexes (i.e., non-covalently associated lipid and glycoprotein). The authors assert that several autoimmune responses are directed against β 2-glycoprotein I/lipid complexes (citing Schousboe, (1979) *Biochim. Biophys. Acta*, 579:396-408), and thus the generation of an anti-complex response may represent substantial breakthroughs in the treatment of cancers.

[0010] Thorpe, Schroit et al. describe a monoclonal antibody (3G4) that binds anionic phospholipids in the presence of serum or the serum protein β 2-glycoprotein I (β 2-GPI). Luster et al., *J. Biol. Chem.* 281: 29863-29871. Originally described as specifically targeting anionic phospholipids, this antibody localizes to vascular endothelial cells in tumors in mice. Ran et al. (2005) *Clin. Cancer Res.* 11:1551-1562. Subsequently, the antibody was shown to bind to complexes of anionic phospholipids and β 2-GPI on tumor vessels, so that antibody binding to PS is dependent on β 2-GPI. Huang et al (2005) *Cancer Res.* 65:4408-4416. The antibody enhances binding of β 2-GPI to endothelial cells via dimerization of β 2GPI. In fact, artificial β 2-GPI dimers can bind to endothelial cell membranes even in the absence of antibody. Luster et al., *J. Biol. Chem.* 281: 29863-29871. A humanized version of 3G4 (Tarvacin, Baviximab) is in clinical trials for treatment of cancer and viral diseases.

[0011] Thorpe et al. (WO 2004/006847) disclose antibodies, fragments or immunoconjugates thereof that bind to PS and compete with antibody 3G4 for binding to PS. Thorpe et al (U.S. Pat. No. 6,818,213, U.S. Pat. No. 6,312,294 and U.S. Pat. No. 6,783,760) disclose therapeutic conjugates that bind to aminophospholipids and have an attached therapeutic agent.

[0012] Baldo et al. (U.S. Pat. No. 5,061,626) disclose antibodies to platelet activating factor (PAF), PAF analogues used to generate antibodies and immunoassays using PAF or PAF analogues. PAF is a choline plasmalogen in which the C-2 (sn2) position of glycerol is esterified with an acetyl group instead of a long chain fatty acid.


[0013] Vielhaber et al. report characterization of two antibody reagents supposedly specific for ceramide, one an IgM-enriched polyclonal mouse serum and the other an IgM monoclonal antibody. The monoclonal was found to be specific for sphingomyelin and the antiserum was found to react with various ceramide species in the nanomolar range. Vielhaber, G. et al., (2001) *Glycobiology* 11:451-457. Also citing the deficiencies of commercially available antibody reagents against ceramide, Krishnamurthy et al. recently reported generation of rabbit IgG against ceramide. *J. Lipid Res.* (2007) 48:968-975.


[0014] B. Lysolipids

[0015] Lysolipids are low molecular weight lipids that contain a polar head group and a single hydrocarbon backbone, due to the absence of an acyl group at one or both possible positions of acylation. Relative to the polar head group at sn-3, the hydrocarbon chain can be at the sn-2 and/or sn-1 position(s) (the term "lyso," which originally related to hemolysis, has been redefined by IUPAC to refer to deacylation). See "Nomenclature of Lipids, www.chem.qmul.ac.uk/iupac/lipid/lipln2.html. These lipids are representative of signaling, bioactive lipids, and their biologic and medical importance highlight what can be achieved by targeting lipid signaling molecules for therapeutic, diagnostic/prognostic, or research purposes (Gardell, et al. (2006), *Trends in Molecular Medicine*, vol 12: 65-75). Two particular examples of medically important lysolipids are LPA (glycerol backbone) and S1P (sphingoid backbone). Other lysolipids include sphingosine, lysophosphatidylcholine (LPC), sphingosylphosphorylcholine (lysosphingomyelin), ceramide, ceramide-1-phosphate, sphinganine (dihydrosphingosine), dihydrosphingosine-1-phosphate and N-acetyl-ceramide-1-phosphate. In contrast, the plasmalogens, which contain an O-alkyl ($-\text{O}-\text{CH}_2-$) or O-alkenyl ether at the C-1 (sn1) and an acyl at C-2, are excluded from the lysolipid genus. The structures of selected LPAs, S1P, and dihydro S1P are presented below.

iupac/lipid/lipln2.html. These lipids are representative of signaling, bioactive lipids, and their biologic and medical importance highlight what can be achieved by targeting lipid signaling molecules for therapeutic, diagnostic/prognostic, or research purposes (Gardell, et al. (2006), *Trends in Molecular Medicine*, vol 12: 65-75). Two particular examples of medically important lysolipids are LPA (glycerol backbone) and S1P (sphingoid backbone). Other lysolipids include sphingosine, lysophosphatidylcholine (LPC), sphingosylphosphorylcholine (lysosphingomyelin), ceramide, ceramide-1-phosphate, sphinganine (dihydrosphingosine), dihydrosphingosine-1-phosphate and N-acetyl-ceramide-1-phosphate. In contrast, the plasmalogens, which contain an O-alkyl ($-\text{O}-\text{CH}_2-$) or O-alkenyl ether at the C-1 (sn1) and an acyl at C-2, are excluded from the lysolipid genus.

The structures of selected LPAs, S1P, and dihydro S1P are presented below.





[0016] LPA is not a single molecular entity but a collection of endogenous structural variants with fatty acids of varied lengths and degrees of saturation (Fujiwara, et al. (2005), *J Biol Chem*, vol. 280: 35038-35050). The structural backbone of the LPAs is derived from glycerol-based phospholipids such as phosphatidylcholine (PC) or phosphatidic acid (PA). In the case of lysosphingolipids such as S1P, the fatty acid of the ceramide backbone at sn-2 is missing. The structural backbone of S1P, dihydro S1P (DHS1P) and sphingosylphosphorylcholine (SPC) is based on sphingosine, which is derived from sphingomyelin.

[0017] LPA and S1P regulate various cellular signaling pathways by binding to the same class of multiple transmembrane domain G protein-coupled (GPCR) receptors (Chun J, Rosen H (2006), *Current Pharm Des*, vol. 12: 161-171, and Moolenaar, W H (1999), *Experimental Cell Research*, vol. 253: 230-238). The S1P receptors are designated as S1P₁, S1P₂, S1P₃, S1P₄ and S1P₅ (formerly EDG-1, EDG-5/AGR16, EDG-3, EDG-6 and EDG-8) and the LPA receptors designated as LPA₁, LPA₂, LPA₃ (formerly, EDG-2, EDG-4, and EDG-7). A fourth LPA receptor of this family has been identified for LPA (LPA₄), and other putative receptors for these lysophospholipids have also been reported.

[0018] C. Lysophosphatic Acids (LPA)

[0019] LPA have long been known as precursors of phospholipid biosynthesis in both eukaryotic and prokaryotic cells, but LPA have emerged only recently as signaling molecules that are rapidly produced and released by activated cells, notably platelets, to influence target cells by acting on specific cell-surface receptor (see, e.g., Moolenaar, et al. (2004), *BioEssays*, vol. 26: 870-881, and van Leewen et al. (2003), *Biochem Soc Trans*, vol 31: 1209-1212). Besides being synthesized and processed to more complex phospholipids in the endoplasmic reticulum, LPA can be generated through the hydrolysis of pre-existing phospholipids following cell activation; for example, the sn-2 position is commonly missing a fatty acid residue due to deacylation, leaving only the sn-1 hydroxyl esterified to a fatty acid. Moreover, a key enzyme in the production of LPA, autotoxin (lysoPLD/NPP2), may be the product of an oncogene, as many tumor types up-regulate autotoxin (Brindley, D. (2004), *J Cell Biochem*, vol. 92: 900-12). The concentrations of LPA in human plasma and serum have been reported, including determinations made using a sensitive and specific LC/MS procedure (Baker, et al. (2001), *Anal Biochem*, vol 292: 287-295). For example, in freshly prepared human serum allowed to sit at 25° C. for one hour, LPA concentrations have been estimated to be approximately 1.2 μM, with the LPA analogs 16:0, 18:1,

18:2, and 20:4 being the predominant species. Similarly, in freshly prepared human plasma allowed to sit at 25° C. for one hour, LPA concentrations have been estimated to be approximately 0.7 μM, with 18:1 and 18:2 LPA being the predominant species.

[0020] LPA influences a wide range of biological responses, ranging from induction of cell proliferation, stimulation of cell migration and neurite retraction, gap junction closure, and even slime mold chemotaxis (Goetzl, et al. (2002), *Scientific World Journal*, vol. 2: 324-338). The body of knowledge about the biology of LPA continues to grow as more and more cellular systems are tested for LPA responsiveness. For instance, it is now known that, in addition to stimulating cell growth and proliferation, LPA promote cellular tension and cell-surface fibronectin binding, which are important events in wound repair and regeneration (Moolenaar, et al. (2004), *BioEssays*, vol. 26: 870-881). Recently, anti-apoptotic activity has also been ascribed to LPA, and it has recently been reported that peroxisome proliferation receptor gamma is a receptor/target for LPA (Simon, et al. (2005), *J Biol Chem*, vol. 280: 14656-14662).

[0021] LPA has proven to be difficult targets for antibody production, although there has been a report in the scientific literature of the production of polyclonal murine antibodies against LPA (Chen et al. (2000) *Med Chem Lett*, vol 10: 1691-3).

[0022] D. Sphingosine-1-Phosphate

[0023] S1P is a mediator of cell proliferation and protects from apoptosis through the activation of survival pathways (Maceyka, et al. (2002), *BBA*, vol. 1585: 192-201, and Spiegel, et al. (2003), *Nature Reviews Molecular Cell Biology*, vol. 4: 397-407). It has been proposed that the balance between CER/SPH levels and S1P provides a rheostat mechanism that decides whether a cell is directed into the death pathway or is protected from apoptosis. The key regulatory enzyme of the rheostat mechanism is sphingosine kinase (SPHK) whose role is to convert the death-promoting bioactive signaling lipids (CER/SPH) into the growth-promoting S1P. S1P has two fates: S1P can be degraded by S1P lyase, an enzyme that cleaves S1P to phosphoethanolamine and hexadecanal, or, less common, hydrolyzed by S1P phosphatase to SPH.

[0024] S1P is abundantly generated and stored in platelets, which contain high levels of SPHK and lacks the enzymes for S1P degradation. When platelets are activated, S1P is secreted. In addition, other cell types, for example, mast cells, are also believed to be capable of secreting S1P. Once secreted, S1P is thought to be bound at high concentrations on carrier proteins such as serum albumin and lipoproteins. S1P is found in high concentrations in plasma, with concentrations in the range of 0.5-5 μM having been reported. Intracellular actions of S1P have also been suggested (see, e.g., Spiegel S, Kolesnick R (2002), *Leukemia*, vol. 16: 1596-602; Suomalainen, et al (2005), *Am J Pathol*, vol. 166: 773-81).

[0025] Widespread expression of the cell surface S1P receptors allows S1P to influence a diverse spectrum of cellular responses, including proliferation, adhesion, contraction, motility, morphogenesis, differentiation, and survival. This spectrum of response appears to depend upon the overlapping or distinct expression patterns of the S1P receptors within the cell and tissue systems. In addition, crosstalk between S1P and growth factor signaling pathways, including platelet-derived growth factor (PDGF), vascular endothelial growth factor (VEGF), and basic fibroblastic growth fac-

tor (bFGF), have recently been demonstrated (see, e.g., Baudhuin, et al. (2004), FASEB J, vol. 18: 341-3). The regulation of various cellular processes involving S1P has particular impact on neuronal signaling, vascular tone, wound healing, immune cell trafficking, reproduction, and cardiovascular function, among others. Alterations of endogenous levels of S1P within these systems can have detrimental effects, eliciting several pathophysiologic conditions, including cancer, heart failure, and infectious and autoimmune diseases.

[0026] A recent novel approach to treating cancer invented by Dr. Sabbadini involves reducing the biologically available extracellular levels of S1P, either alone or in combination with conventional anti-cancer treatments, including the administration of chemotherapeutic agents, such as an anthracycline. To this end, the generation of antibodies specific for S1P has been described. See, e.g., commonly owned U.S. patent application Ser. No. 10/820,582. Such antibodies, which can selectively adsorb S1P from serum, act as molecular sponges to neutralize extracellular S1P. See also commonly owned U.S. Pat. Nos. 6,881,546 and 6,858,383 and U.S. patent application Ser. Nos. 10/028,520, 10/029,372, and 11/101,976. Since S1P has also been shown to be pro-angiogenic, an added benefit to the antibody's effectiveness is its ability to starve growing tumors of nutrients and oxygen by limiting blood supply.

[0027] What is particularly unique about the anti-S1P approach is that while sphingolipid-based anti-cancer strategies that target key enzymes of the sphingolipid metabolic pathway, such as SPHK, have been proposed, the lipid mediator S1P itself was not previously emphasized, largely because of difficulties in directly mitigating this lipid target, in particular because of the difficulty first in raising antibodies against a lipid target such as S1P, and second, in detecting antibodies in fact produced against the S1P target. As already noted, similar difficulties exist with respect to treatments and diagnostic approaches directed at other lipid targets. This invention provides an effective solution to both of these dilemmas by providing patentable methods, in particular, the generation of monoclonal antibodies against bioactive lipids.

3. Definitions

[0028] Before describing the instant invention in detail, several terms used in the context of the present invention will be defined. In addition to these terms, others are defined elsewhere in the specification, as necessary. Unless otherwise expressly defined herein, terms of art used in this specification will have their art-recognized meanings.

[0029] An "anti-S1P antibody" refers to any antibody or antibody-derived molecule that binds S1P.

[0030] A "bioactive lipid" refers to a lipid signaling molecule. Bioactive lipids are distinguished from structural lipids (e.g., membrane-bound phospholipids) in that they mediate extracellular and/or intracellular signaling and thus are involved in controlling the function of many types of cells by modulating differentiation, migration, proliferation, secretion, survival, and other processes. In vivo, bioactive lipids can be found in extracellular fluids, where they can be complexed with other molecules, for example serum proteins such as albumin and lipoproteins, or in "free" form, i.e., not complexed with another molecule species. As extracellular mediators, some bioactive lipids alter cell signaling by activating membrane-bound ion channels or GPCRs or enzymes or factors that, in turn, activate complex signaling systems

that result in changes in cell function or survival. As intracellular mediators, bioactive lipids can exert their actions by directly interacting with intracellular components such as enzymes, ion channels or structural elements such as actin. Representative examples of bioactive lipids include LPA and S1P.

[0031] Examples of bioactive lipids include sphingolipids such as ceramide, ceramide-1-phosphate, sphingosine, sphinganine, sphingosylphosphorylcholine (SPC) and sphingosine-1-phosphate (S1P). Sphingolipids and their derivatives and metabolites are characterized by a sphingoid backbone (derived from sphingomyelin). Sphingolipids and their derivatives and metabolites represent a group of extracellular and intracellular signaling molecules with pleiotropic effects on important cellular processes. They include sulfatides, gangliosides and cerebroside. Other bioactive lipids are characterized by a glycerol-based backbone; for example, lysophospholipids such as lysophosphatidyl choline (LPC) and various lysophosphatidic acids (LPA), as well as phosphatidylinositol (PI), phosphatidylethanolamine (PEA), phosphatidic acid, platelet activating factor (PAF), cardiolipin, phosphatidylglycerol (PG) and diacylglyceride (DG). Yet other bioactive lipids are derived from arachidonic acid; these include the eicosanoids (including the eicosanoid metabolites such as the HETEs, cannabinoids, leukotrienes, prostaglandins, lipoxins, epoxyeicosatrienoic acids, and isoeicosanoids), non-eicosanoid cannabinoid mediators. Other bioactive lipids, including other phospholipids and their derivatives, may also be used according to the instant invention.

[0032] In some embodiments of the invention it may be preferable to target glycerol-based bioactive lipids (those having a glycerol-derived backbone, such as the LPAs) for antibody production, as opposed to sphingosine-based bioactive lipids (those having a sphingoid backbone, such as sphingosine and S1P). In other embodiments it may be desired to target arachidonic acid-derived bioactive lipids for antibody generation, and in other embodiments arachidonic acid-derived and glycerol-derived bioactive lipids but not sphingoid-derived bioactive lipids are preferred. Together the arachidonic acid-derived and glycerol-derived bioactive lipids may be referred to in the context of this invention as "non-sphingoid bioactive lipids."

[0033] Specifically excluded from the class of bioactive lipids according to the invention are phosphatidylcholine and phosphatidylserine, as well as their metabolites and derivatives that function primarily as structural members of the inner and/or outer leaflet of cellular membranes.

[0034] A "biomarker" is a specific biochemical in the body which has a particular molecular feature that makes it useful for measuring the progress of disease or the effects of treatment.

For example, S1P is a biomarker for certain hyperproliferative and/or cardiovascular conditions.

[0035] A "carrier" refers to a moiety adapted for conjugation to a hapten, thereby rendering the hapten immunogenic. A representative, non-limiting class of carriers is proteins, examples of which include albumin, keyhole limpet hemocyanin, hemagglutinin, tetanus, and diphtheria toxoid. Other classes and examples of carriers suitable for use in accordance with the invention are known in the art. These, as well as later discovered or invented naturally occurring or synthetic carriers, can be adapted for application in accordance with the invention.

[0036] The term “chemotherapeutic agent” means anti-cancer and other anti-hyperproliferative agents. Put simply, a “chemotherapeutic agent” refers to a chemical intended to destroy cells and tissues. Such agents include, but are not limited to: DNA damaging agents and agents that inhibit DNA synthesis: anthracyclines (doxorubicin, daunorubicin, epirubicin), alkylating agents (bendamustine, busulfan, carboplatin, carmustine, chlorambucil, cyclophosphamide, dacarbazine, hexamethylmelamine, ifosfamide, lomustine, mechlorethamine, melphalan, mitotane, mytomicin, pipobroman, procarbazine, streptozocin, thiotepe, and triethylenemelamine), platinum derivatives (cisplatin, carboplatin, cis diamine-dichloroplatinum), and topoisomerase inhibitors (Camptosar); anti-metabolites such as capecitabine, chlorodeoxyadenosine, cytarabine (and its activated form, ara-CMP), cytosine arabinoside, dacarbazine, floxuridine, fludarabine, 5-fluorouracil, 5-DFUR, gemcitabine, hydroxyurea, 6-mercaptopurine, methotrexate, pentostatin, trimetrexate, 6-thioguanine); anti-angiogenics (bevacizumab, thalidomide, sunitinib, lenalidomide, TNP-470, 2-methoxyestradiol, ranibizumab, sorafenib, erlotinib, bortezomib, pegaptanib, endostatin); vascular disrupting agents (flavonoids/flavones, DMXAA, combretastatin derivatives such as CA4DP, ZD6126, AVE8062A, etc.); biologics such as antibodies (Herceptin, Avastin, Panorex, Rituxin, Zevalin, Mylotarg, Campath, Bexxar, Erbitux); endocrine therapy: aromatase inhibitors (4-hydroandrostendione, exemestane, aminoglutethimide, anastrozole, letrozole), anti-estrogens (Tamoxifen, Toremifene, Raloxifene, Faslodex), steroids such as dexamethasone; immuno-modulators: cytokines such as IFN-beta and IL2), inhibitors to integrins, other adhesion proteins and matrix metalloproteinases); histone deacetylase inhibitors like suberoylanilide hydroxamic acid; inhibitors of signal transduction such as inhibitors of tyrosine kinases like imatinib (Gleevec); inhibitors of heat shock proteins like 17-N-allylamino-17-demethoxygeldanamycin; retinoids such as all trans retinoic acid; inhibitors of growth factor receptors or the growth factors themselves; anti-mitotic compounds and/or tubulin-depolymerizing agents such as the taxoids (paclitaxel, docetaxel, taxotere, BAY 59-8862), navelbine, vinblastine, vincristine, vindesine and vinorelbine; anti-inflammatory such as COX inhibitors and cell cycle regulators, e.g., check point regulators and telomerase inhibitors.

[0037] The term “combination therapy” refers to a therapeutic regimen that involves the provision of at least two distinct therapies to achieve an indicated therapeutic effect. For example, a combination therapy may involve the administration of two or more chemically distinct active ingredients, for example, a fast-acting chemotherapeutic agent and an anti-lipid antibody. Alternatively, a combination therapy may involve the administration of an anti-lipid antibody and/or one or more chemotherapeutic agents, alone or together with the delivery of another treatment, such as radiation therapy and/or surgery. In the context of the administration of two or more chemically distinct active ingredients, it is understood that the active ingredients may be administered as part of the same composition or as different compositions. When administered as separate compositions, the compositions comprising the different active ingredients may be administered at the same or different times, by the same or different routes, using the same or different dosing regimens, all as the particular context requires and as determined by the attending physician. Similarly, when one or more anti-lipid antibody

species, for example, an anti-LPA antibody, alone or in conjunction with one or more chemotherapeutic agents are combined with, for example, radiation and/or surgery, the drug(s) may be delivered before or after surgery or radiation treatment.

[0038] A “derivatized bioactive lipid conjugate” refers to a derivatized bioactive lipid covalently conjugated to a carrier. The carrier may be a protein molecule or may be a moiety such as polyethylene glycol, colloidal gold, adjuvants or silicone beads. A derivatized bioactive lipid conjugate may be used as an immunogen for generating an antibody response according to the instant invention, and the same or a different bioactive lipid conjugate may be used as a detection reagent for detecting the antibody thus produced. In some embodiments the derivatized bioactive lipid conjugate is attached to a solid support when used for detection.

[0039] An “epitope” or “antigenic determinant” refers to that portion of an antigen that reacts with an antibody antigen-binding portion derived from an antibody.

[0040] A “hapten” is a substance that is non-immunogenic but can react with an antibody or antigen-binding portion derived from an antibody. In other words, haptens have the property of antigenicity but not immunogenicity.

[0041] The term “hyperproliferative disorder” refers to diseases and disorders associated with, the uncontrolled proliferation cells, including but not limited to uncontrolled growth of organ and tissue cells resulting in cancers and benign tumors. Hyperproliferative disorders associated with endothelial cells can result in diseases of angiogenesis such as angiomas, endometriosis, obesity, age-related macular degeneration and various retinopathies, as well as the proliferation of endothelial cells and smooth muscle cells that cause restenosis as a consequence of stenting in the treatment of atherosclerosis. Hyperproliferative disorders involving fibroblasts (i.e., fibrogenesis) include but are not limited to disorders of excessive scarring (i.e., fibrosis) such as age-related macular degeneration, cardiac remodeling and failure associated with myocardial infarction, excessive wound healing such as commonly occurs as a consequence of surgery or injury, keloids, and fibroid tumors and stenting.

[0042] An “immunogen” is a molecule capable of inducing a specific immune response, particularly an antibody response in an animal to whom the immunogen has been administered. In the instant invention, the immunogen is a derivatized bioactive lipid conjugated to a carrier, i.e., a “derivatized bioactive lipid conjugate”. The derivatized bioactive lipid conjugate used as the immunogen may be used as capture material for detection of the antibody generated in response to the immunogen. Thus the immunogen may also be used as a detection reagent. Alternatively, the derivatized bioactive lipid conjugate used as capture material may have a different linker and/or carrier moiety from that in the immunogen.

[0043] To “inhibit,” particularly in the context of a biological phenomenon, means to decrease, suppress or delay. For example, a treatment yielding “inhibition of tumorigenesis” may mean that tumors do not form at all, or that they form more slowly, or are fewer in number than in the untreated control.

[0044] In the context of this invention, a “liquid composition” refers to one that, in its filled and finished form as provided from a manufacturer to an end user (e.g., a doctor or nurse), is a liquid or solution, as opposed to a solid. Here, “solid” refers to compositions that are not liquids or solutions.

For example, solids include dried compositions prepared by lyophilization, freeze-drying, precipitation, and similar procedures.

[0045] “Monotherapy” refers to a treatment regimen based on the delivery of one therapeutically effective compound, whether administered as a single dose or several doses over time.

[0046] “Neoplasia” refers to abnormal and uncontrolled cell growth. A “neoplasm”, or tumor, is an abnormal, unregulated, and disorganized proliferation of cell growth, and is generally referred to as cancer. A neoplasm may be benign or malignant. A neoplasm is malignant, or cancerous, if it has properties of destructive growth, invasiveness, and metastasis. Invasiveness refers to the local spread of a neoplasm by infiltration or destruction of surrounding tissue, typically breaking through the basal laminas that define the boundaries of the tissues, thereby often entering the body’s circulatory system. Metastasis typically refers to the dissemination of tumor cells by lymphatic or blood circulating systems. Metastasis also refers to the migration of tumor cells by direct extension through serous cavities, or subarachnoid or other spaces. Through the process of metastasis, tumor cell migration to other areas of the body establishes neoplasms in areas away from the site of initial appearance.

[0047] A “patentable” composition, process, machine, or article of manufacture according to the invention means that the subject matter satisfies all statutory requirements for patentability at the time the analysis is performed. For example, with regard to novelty, non-obviousness, or the like, if later investigation reveals that one or more claims encompass one or more embodiments that would negate novelty, non-obviousness, etc., the claim(s), being limited by definition to “patentable” embodiments, specifically exclude the non-patentable embodiment(s). Also, the claims appended hereto are to be interpreted both to provide the broadest reasonable scope, as well as to preserve their validity. Furthermore, the claims are to be interpreted in a way that (1) preserves their validity and (2) provides the broadest reasonable interpretation under the circumstances, if one or more of the statutory requirements for patentability are amended or if the standards change for assessing whether a particular statutory requirement for patentability is satisfied from the time this application is filed or issues as a patent to a time the validity of one or more of the appended claims is questioned.

[0048] The term “pharmaceutically acceptable salt” refers to salts which retain the biological effectiveness and properties of the agents and compounds of this invention and which are not biologically or otherwise undesirable. In many cases, the agents and compounds of this invention are capable of forming acid and/or base salts by virtue of the presence of charged groups, for example, charged amino and/or carboxyl groups or groups similar thereto. Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids, while pharmaceutically acceptable base addition salts can be prepared from inorganic and organic bases. For a review of pharmaceutically acceptable salts (see Berge, et al. (1977) *J. Pharm. Sci.*, vol. 66, 1-19).

[0049] A “plurality” means more than one.

[0050] The terms “separated”, “purified”, “isolated”, and the like mean that one or more components of a sample contained in a sample-holding vessel are or have been physically removed from, or diluted in the presence of, one or more other sample components present in the vessel. Sample components that may be removed or diluted during a separating or

purifying step include, chemical reaction products, non-reacted chemicals, proteins, carbohydrates, lipids, and unbound molecules.

[0051] The term “species” is used herein in various contexts, e.g., a particular species of chemotherapeutic agent. In each context, the term refers to a population of chemically indistinct molecules of the sort referred in the particular context.

[0052] “Specifically associate,” “specifically bind” and the like refer to a specific, non-random interaction between two molecules, which interaction depends on the presence of structural, hydrophobic/hydrophilic, and/or electrostatic features that allow appropriate chemical or molecular interactions between the molecules. An antibody may be said to “bind” or be “reactive with” (or, equivalently, “reactive against”) the epitope of its target antigen. Antibodies are commonly described in the art as being “against” or “to” their antigens as shorthand for antibody binding to the antigen.

[0053] Herein, “stable” refers to an interaction between two molecules (e.g., a peptide and a TLR molecule) that is sufficiently stable such that the molecules can be maintained for the desired purpose or manipulation. For example, a “stable” interaction between a peptide and a TLR molecule refers to one wherein the peptide becomes and remains associated with a TLR molecule for a period sufficient to achieve the desired effect.

[0054] A “subject” or “patient” refers to an animal in need of treatment that can be effected by molecules of the invention. Animals that can be treated in accordance with the invention include vertebrates, with mammals such as bovine, canine, equine, feline, ovine, porcine, and primate (including humans and non-humans primates) animals being particularly preferred examples.

[0055] A “surrogate marker” refers to laboratory measurement of biological activity within the body that indirectly indicates the effect of treatment on disease state. Examples of surrogate markers for hyperproliferative and/or cardiovascular conditions include SPHK and/or S1PRs.

[0056] A “therapeutically effective amount” (or “effective amount”) refers to an amount of an active ingredient, e.g., an agent according to the invention, sufficient to effect treatment when administered to a subject in need of such treatment. Accordingly, what constitutes a therapeutically effective amount of a composition according to the invention may be readily determined by one of ordinary skill in the art. In the context of cancer therapy, a “therapeutically effective amount” is one that produces an objectively measured change in one or more parameters associated with cancer cell survival or metabolism, including an increase or decrease in the expression of one or more genes correlated with the particular cancer, reduction in tumor burden, cancer cell lysis, the detection of one or more cancer cell death markers in a biological sample (e.g., a biopsy and an aliquot of a bodily fluid such as whole blood, plasma, serum, urine, etc.), induction of induction apoptosis or other cell death pathways, etc. Of course, the therapeutically effective amount will vary depending upon the particular subject and condition being treated, the weight and age of the subject, the severity of the disease condition, the particular compound chosen, the dosing regimen to be followed, timing of administration, the manner of administration and the like, all of which can readily be determined by one of ordinary skill in the art. It will be appreciated that in the context of combination therapy, what constitutes a therapeutically effective amount of a particular active ingredient may

differ from what constitutes a therapeutically effective amount of the active ingredient when administered as a monotherapy (i.e., a therapeutic regimen that employs only one chemical entity as the active ingredient).

[0057] The term “treatment” or “treating” means any treatment of a disease or disorder, including preventing or protecting against the disease or disorder (that is, causing the clinical symptoms not to develop); inhibiting the disease or disorder (i.e., arresting, delaying or suppressing the development of clinical symptoms; and/or relieving the disease or disorder (i.e., causing the regression of clinical symptoms). As will be appreciated, it is not always possible to distinguish between “preventing” and “suppressing” a disease or disorder because the ultimate inductive event or events may be unknown or latent. Accordingly, the term “prophylaxis” will be understood to constitute a type of “treatment” that encompasses both “preventing” and “suppressing”. The term “protection” thus includes “prophylaxis”.

[0058] The term “therapeutic regimen” means any treatment of a disease or disorder using chemotherapeutic and cytotoxic agents, radiation therapy, surgery, gene therapy, DNA vaccines and therapy, siRNA therapy, anti-angiogenic therapy, immunotherapy, bone marrow transplants, aptamers and other biologics such as antibodies and antibody variants, receptor decoys and other protein-based therapeutics.

SUMMARY OF THE INVENTION

[0059] The object of this invention is to provide patentable compositions and methods for generating antibodies, particularly monoclonal antibodies and derivatives thereof, reactive with bioactive lipids correlated, involved, or otherwise implicated in disease processes in animals, particularly in mammals, especially humans.

[0060] Thus, one aspect of the invention concerns patentable intermediates used to produce patentable immunogens that can be used to raise patentable bioactive lipid-reactive antibodies. This patentable class of compounds comprises derivatized bioactive lipids, each of which comprises a bioactive lipid having a polar head group and at least one hydrocarbon chain, wherein a carbon atom within the hydrocarbon chain is derivatized with a pendant reactive group [e.g., a sulfhydryl (thiol) group, a carboxylic acid group, a cyano group, an ester, a hydroxy group, an alkene, an alkyl, an acid chloride group or a halogen atom] that may or may not be protected. Representative bioactive lipids include lysolipids, for example, sphingolipids and sphingolipid metabolites such as ceramide, ceramide-1-phosphate, N-acetyl-ceramide-1-phosphate, sphingosine-1-phosphate (S1P), sphingosine, sphingosylphosphorylcholine (SPC), dihydrosphingosine and dihydrosphingosine-1-phosphate. Other bioactive lipids include lysolipids such as lysophosphatidic acids (LPAs), as well as lysophosphatidic acid metabolites or precursors such as lysophosphatidylinositol (LPI) or lysophosphatidylcholine (LPC). In the context of an LPA, exemplary reactive group positioning includes appending the reactive group to a carbon atom within the hydrocarbon chain or at the sn-1 position of the glycerol backbone of the lysophosphatidic acid moiety. Particularly preferred derivatized bioactive lipids include sulfhydryl derivatives of LPA and S1P.

[0061] A related aspect of the invention relates to immunogens produced from a derivatized bioactive lipid according to the invention. In general, such immunogens comprise a derivatized bioactive lipid covalently linked to a carrier. Examples of suitable carrier moieties include carrier proteins

such as keyhole limpet hemocyanin (KLH) and albumin, polyethylene glycol, colloidal gold, adjuvants or silicone beads. Preferred embodiments of an immunogen according to the invention include a sulfhydryl derivative of LPA covalently linked to KLH or albumin. In the context of sphingolipid-based immunogen, preferred immunogen embodiments include sulfhydryl derivatives of S1P covalently linked to KLH or albumin.

[0062] Immunogens of the invention are prepared by reaction of a derivatized bioactive lipid with a carrier moiety under conditions that allow covalent linkage between the carrier and the bioactive lipid to occur through the pendant reactive group to yield the particular species of bioactive lipid-carrier immunogen. Such immunogens are then preferably isolated or purified prior to administration to a host animal as part of an immunization procedure, which may involve one or several administrations (typically by injection) of the desired immunogen. In preferred embodiments of this aspect, the pendant reactive group of the derivatized bioactive lipid is protected with a suitable protecting group, which is removed and the derivatized bioactive lipid is “deprotected” prior to or as part of the chemistry employed to covalently link the carrier and the bioactive lipid.

[0063] As discussed above, another aspect of the invention concerns methods of making monoclonal antibody reactive against a bioactive lipid. In such methods, an immune competent host animal (e.g., a rodent such as a mouse, a rat, a guinea pig, or rabbit) is immunized with a bioactive lipid immunogen as described herein. Following immunization, the host mounts an antibody response against the bioactive lipid, resulting in the production of antibodies reactive to the particular bioactive lipid species present in the immunogen. The resultant antibodies may be polyclonal or, preferably, monoclonal. With regard to monoclonal antibodies, cell lines that produce a desired antibody are preferably cloned and immortalized to facilitate production of the desired lipid-specific antibody in desired quantities. In preferred embodiments, a desired monoclonal antibody, e.g., a monoclonal antibody reactive against LPA is used to produce antibody derivatives, such as chimeric or humanized antibodies or antibody fragments. In some embodiments, fully humanized antibodies may be produced by immunizing an animal, e.g., a mouse or rat, engineered to contain some or all of a competent human system.

[0064] It is known that lipids are in general a particularly intractable class of molecules for antibody production. One facet of the invention rests on the appreciation that this problem, at least in part, resides in the difficulty in detecting antibodies reactive against a particular target lipid species. However, this problem can be elegantly overcome through the use of the derivatized form of the particular target bioactive lipid, such as a lysolipid or a sphingolipid or sphingolipid metabolite).

[0065] In certain preferred embodiments, such a derivatized bioactive lipid may be used to identify an antibody reactive against an epitope of the particular bioactive lipid present in the immunogen used to generate the antibodies being detected. To perform this role a particular derivatized bioactive lipid or derivatized bioactive lipid conjugate may be attached to a solid support, preferably the solid phase of an assay device, such as an ELISA plate, a Biacore chip, etc. Attachment to a solid support minimizes the likelihood that the bioactive lipid will be washed away during antibody binding and detection.

[0066] Another aspect of the invention concerns pharmaceutical or veterinary compositions that comprise a carrier and an isolated immune-derived moiety according to the invention, for example, a monoclonal antibody or antibody fragment, variant, or derivative. Preferred carriers include those that are pharmaceutically acceptable, particularly when the composition is intended for therapeutic use in humans. For non-human therapeutic applications (e.g., in the treatment of companion animals, livestock, fish, or poultry), acceptable carriers for veterinary use may be employed.

[0067] Related aspects of the invention relate to methods of use or treatment, including preventative or prophylactic treatment, and administration. Such methods typically involve administering to a subject (for example, mammal, particularly a human patient) in need of therapeutic or prophylactic treatment an amount of an immune-derived moiety reactive against a bioactive lipid target, effective to accomplish the desired treatment. In some embodiments the bioactive lipid target is a non-sphingoid bioactive lipid. One preferred example of a therapeutically useful immune-derived moiety is a humanized monoclonal antibody reactive against a lysolipid such as LPA. Routes of administration of an immune-derived moiety according to the invention, preferably as part of a therapeutic composition, may vary depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, ophthalmic and to mucous membranes including vaginal, intrauterine and rectal delivery, pulmonary delivery, intratracheal, intranasal, and epidermal delivery), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration.

[0068] Other aspects of the invention concern various diagnostic, prognostic, and/or research-enabling methods. One such aspect involves use of the derivatized lipid analog to detect the presence of autoantibodies against the natural bioactive lipid in a sample of fluid or tissue from an animal or from an antibody library. Another such aspect concerns methods of detecting target bioactive lipids, other than sphingolipids or metabolites thereof. In general, such methods involve binding of an immune-derived moiety with the target bioactive lipid against which it is reactive. Detection of binding may result, for example, by exposing a sample (e.g., a biopsy or fluid or liquid sample, for instance, blood, serum, plasma, urine, saliva, tears, cerebrospinal fluid, cell culture, etc.) known or suspected to contain the target bioactive lipid with an immune-derived moiety under conditions that allow the immune-derived moiety to bind to the target bioactive lipid, if present in the sample.

[0069] To perform such diagnostic methods, reagents are required, and diagnostic reagents that employ a derivatized lipid according to the invention represent another aspect of the invention. With such reagents in hand, diagnostic assays that utilize such reagents may be prepared.

[0070] These and other aspects and embodiments of the invention are discussed in greater detail in the sections that follow.

BRIEF DESCRIPTION OF THE DRAWINGS

[0071] This patent application contains at least one figure executed in color. Copies of this patent application with color drawing(s) will be provided upon request and payment of the necessary fee.

[0072] FIG. 1. Organic synthesis scheme for making of a typical thiolated-S1P analog that was used as a key component of an immunogen according to the invention, as well as a key component of the laydown material for the ELISA and BiaCore assays.

[0073] FIG. 2. Organic synthesis scheme for making the thiolated-related fatty acid used in the synthesis of the thiolated-LPA analog of FIG. 3.

[0074] FIG. 3. Organic synthesis scheme for making the thiolated-LPA analog that is a key component of an immunogen according to the invention, as well as a key component of the laydown material for the ELISA and other assays.

[0075] FIG. 4. The anti-S1P mAb is specific and sensitive for S1P and does not recognize structurally similar bioactive lipids. Panel A. Competitive ELISA with S1P, SPH, LPA, SPC and other structurally similar biolipids competing for the mAb binding to S1P on the plate. Only free S1P or DH-S1P can compete for binding, demonstrating the specificity of the anti-S1P mAb. SPC only slightly competes for binding. Panel B. Structures of bioactive lipids used in the evaluation of specificity.

[0076] FIG. 5. BiaCore analysis of binding kinetics of anti-S1P mAb to thio-S1P tethered to a Biacore maleimide surface CM5 sensor chip. Various dilutions of anti-S1P mAb were applied to the flow cell for generating sensograms.

[0077] FIG. 6. Amino acid sequences of the mouse V_H and V_L domains of murine Sphingomab™. CDR residues are boxed.

[0078] FIG. 7. Nucleotide and amino acid sequences of the V_H and V_L domains of murine Sphingomab™.

[0079] FIG. 8. Graph showing ELISA results for binding studies of murine Sphingomab™ and chimeric, S1P-binding antibodies derived from murine Sphingomab™.

[0080] FIG. 9. Direct ELISA showing binding of murine and chimeric mAbs to ELISA plates coated with thiolated S1P analog as described in EXAMPLE 6. Data show that the chimeric mAb (α -S1P IgG) has similar, if not greater binding performance compared to the fully murine mAb (μ -S1P IgG).

[0081] As those in the art will appreciate, the following description describes certain preferred embodiments of the invention in detail, and is thus only representative and does not depict the actual scope of the invention. Before describing the present invention in detail, it is understood that the invention is not limited to the particular molecules, systems, and methodologies described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the invention defined by the appended claims.

DETAILED DESCRIPTION OF THE INVENTION

[0082] The present invention relates to compositions and methods for generating and identifying antibodies against bioactive lipid molecules that play a role in human and/or animal disease as a signaling molecule. The invention also relates to these antibodies themselves, and methods of using them therapeutically, diagnostically and as research reagents.

1. Methods for Antibody Production and Identification

[0083] It is known that lipids are in general a particularly intractable class of molecules for antibody production. Anti-

body production can typically be described as a two-part process: a suitable immunogen must be provided which will generate the desired antibody response in an animal, and the resulting antibody, if present, must be detectable.

[0084] As discussed above, effective antibody production requires both antibody generation and antibody detection. As disclosed in the Examples hereinbelow, generation of antibodies targeted to certain bioactive lipids has been achieved using derivatized bioactive lipid as immunogen. In the examples, the thiolated bioactive lipid (e.g., S1P) analog was conjugated to Keyhole Limpet Hemocyanin (KLH) or to fatty-acid free Bovine Serum Albumin (BSA) via SMCC (Pierce, Rockford Ill.) using protocols recommended by the manufacturer. SMCC is a heterobifunctional crosslinker that reacts with primary amines and sulfhydryl groups, and represents a preferred crosslinker. Iodoacetamide (IOA) can also be used for maleimide-activated proteins.

[0085] However, other immunogens and methods of generating antibodies known in the art may also be used. For example, antibodies against phospholipid species have been generated by immunization with liposomes (Maneta-Peyret et al., 1988, 1989; Benerji and Alving, 1990) or by adsorption of monomeric phospholipids to proteins (Tamamura et al., 1971; Maneta-Peyret et al., 1989), to bacteria (Umeda et al., 1989), to acrylamide (Maneta-Peyret et al., 1988, 1989) and to gold [Tomii et al., (1991) *Jpn J. Med Sci. Biol.* 44:75-80]. In many cases, presentation of the bioactive lipid as emulsions or liposomal complexes has resulted in IgMs with limited specificity, sensitivity and/or biological activity in comparison to IgG. For example, two commercially available reagents supposedly specific for ceramide, one an IgM-enriched polyclonal mouse serum and the other an IgM monoclonal antibody, were characterized. The monoclonal was found to be specific for sphingomyelin and the antiserum was found to react with various ceramide species in the nanomolar range. Vielhaber, G. et al., (2001) *Glycobiology* 11:451-457. In a different approach, Ran et al. [(2005) *Clin. Cancer Res.* 11: 1551-1562] used b.End3 endothelial cells that had been treated with peroxide (intended to cause translocation of anionic phospholipids to the external surface of the cells) as an immunogen to elicit generation of antibodies specific for anionic phospholipids. Thus numerous methods are known by which an antibody response to a desired antigenic target may be elicited; any of these may be used in the instant invention as long as the resulting antibodies can be detected and shown to be reactive with the desired bioactive lipid.

[0086] Antibody generation, while of course necessary, is not sufficient if the antibody cannot be detected. Thus one facet of the invention rests on the appreciation that previous failures of others to produce antibodies to bioactive lipids may be attributable at least to shortcomings in the detection step. This problem of detection has been elegantly overcome in the following examples through the use of a derivatized bioactive lipid. The derivatized bioactive lipid is used to detect and identify an antibody reactive against an epitope of the particular bioactive lipid present in the immunogen used to generate the antibodies being detected; the bioactive lipid used for detection in derivatized form contains the same epitope to which antibodies were generated. To perform this role the derivatized lipid may be associated with the solid phase of an assay device, such as an ELISA plate, a BiaCore sensor chip, etc. In some embodiments the derivatized bioactive lipid is covalently conjugated directly to the solid support. By way of example, the derivatized lipid may be

covalently conjugated to an activated BiaCore chip as described in Examples hereinbelow. In other embodiments, the derivatized bioactive lipid is covalently conjugated to a carrier moiety, yielding a "derivatized bioactive lipid conjugate" which is then bound to a solid support. As an example, derivatized lipid covalently conjugated to BSA is used as the laydown material (capture material) for ELISA as described in Examples hereinbelow. In either embodiment, attachment of the derivatized bioactive lipid to the solid support provides a stable detection means which is unlikely to be washed away, as is a risk of some detection methods. Detection of the antibody may be accomplished in a variety of ways. In a preferred embodiment of the invention, the detection is via ELISA, Biacore™ label-free interaction analysis systems, or other solid-support-based routine detection means in which the derivatized bioactive lipid is attached to said solid support. Examples of other solid supports include but are not limited to affinity columns, glass or synthetic beads, multi-well plates and the like.

[0087] The derivatized bioactive lipid conjugate used in the detection step may be the same derivatized bioactive lipid conjugate used as the immunogen, or the derivatized bioactive lipid may be conjugated to a different carrier than in the conjugate used as the immunogen. In some embodiments, e.g. as the laydown for ELISA, it is preferred to use a different derivatized bioactive lipid conjugate in the detection step, than was used as the immunogen, to minimize crossreactivity. By way of examples, the carrier may be BSA (preferably fatty-acid free, particularly in the detection step), KLH or other carriers known in the art. The crosslinker used to conjugate the derivatized bioactive lipid to the protein carrier may be, for example, SMCC or IOA. In one preferred embodiment the immunogen is S1P-IOA-KLH and S1P-SMCC-BSA (fatty acid free BSA) is the capture laydown material in the ELISA, wherein S1P refers to the derivatized S1P that reacts with the crosslinker (IOA or SMCC in this instance) to form a covalent bond with the protein carrier (KLH or BSA in this instance).

2. Compounds

[0088] The term "antibody" ("Ab") or "immunoglobulin" (Ig) refers to any form of a peptide, polypeptide derived from, modeled after or encoded by, an immunoglobulin gene, or fragment thereof, capable of binding an antigen or epitope. See, e.g., IMMUNOBIOLOGY, Fifth Edition, C. A. Janeway, P. Travers, M., Walport, M. J. Shlomchik, ed. Garland Publishing (2001). Antibody molecules or immunoglobulins are large glycoprotein molecules with a molecular weight of approximately 150 kDa, usually composed of two different kinds of polypeptide chain. One polypeptide chain, termed the "heavy" chain (H) is approximately 50 kDa. The other polypeptide, termed the "light" chain (L), is approximately 25 kDa. Each immunoglobulin molecule usually consists of two heavy chains and two light chains. The two heavy chains are linked to each other by disulfide bonds, the number of which varies between the heavy chains of different immunoglobulin isotypes. Each light chain is linked to a heavy chain by one covalent disulfide bond. In any given naturally occurring antibody molecule, the two heavy chains and the two light chains are identical, harboring two identical antigen-binding sites, and are thus said to be divalent, i.e., having the capacity to bind simultaneously to two identical molecules.

[0089] The "light" chains of antibody molecules from any vertebrate species can be assigned to one of two clearly dis-

tinct types, kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains. The ratio of the two types of light chain varies from species to species. As a way of example, the average κ to λ ratio is 20:1 in mice, whereas in humans it is 2:1 and in cattle it is 1:20.

[0090] The “heavy” chains of antibody molecules from any vertebrate species can be assigned to one of five clearly distinct types, called isotypes, based on the amino acid sequences of their constant domains. Some isotypes have several subtypes. The five major classes of immunoglobulin are immunoglobulin M (IgM), immunoglobulin D (IgD), immunoglobulin G (IgG), immunoglobulin A (IgA), and immunoglobulin E (IgE). IgG is the most abundant isotype and has several subclasses (IgG1, 2, 3, and 4 in humans). The Fc fragment and hinge regions differ in antibodies of different isotypes, thus determining their functional properties. However, the overall organization of the domains is similar in all isotypes.

[0091] As used herein, “antibody fragment” and grammatical variants thereof refer to a portion of an intact antibody that includes the antigen binding site or variable regions of an intact antibody, wherein the portion can be free of the constant heavy chain domains (e.g., CH2, CH3, and CH4) of the Fc region of the intact antibody. Alternatively, portions of the constant heavy chain domains (e.g., CH2, CH3, and CH4) can be included in the “antibody fragment”. Examples of antibody fragments are those that retain antigen-binding and include Fab, Fab', F(ab')₂, Fd, and Fv fragments; diabodies; triabodies; single-chain antibody molecules (sc-Fv); minibodies, nanobodies, and multispecific antibodies formed from antibody fragments. By way of example, a Fab fragment also contains the constant domain of a light chain and the first constant domain (CH1) of a heavy chain.

[0092] The term “variable region” refers to N-terminal sequence of the antibody molecule or a fragment thereof. In general, each of the four chains has a variable (V) region in its amino terminal portion, which contributes to the antigen-binding site, and a constant (C) region, which determines the isotype. The light chains are bound to the heavy chains by many noncovalent interactions and by disulfide bonds, and the V regions of the heavy and light chains pair in each arm of antibody molecule to generate two identical antigen-binding sites. Some amino acid residues are believed to form an interface between the light- and heavy-chain variable domains (see Kabat et al., *Sequences of Proteins of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, Md. (1991); Clothia et al., *J. Mol. Biol.*, vol. 186: 651 (1985)).

[0093] Of note, variability is not uniformly distributed throughout the variable domains of antibodies, but is concentrated in three segments called “complementarity-determining regions” (CDRs) or “hypervariable regions” both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the “framework region” (FR). The variable domains of native heavy and light chains each comprise four FR regions connected by three CDRs. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat et al., *Sequences of Proteins of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, Md. (1991)). Collectively, the 6 CDRs contribute to the binding properties of the antibody molecule. However, even a single variable

domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen (see Pluckthun, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994)).

[0094] The terms “constant domain” refers to the C-terminal region of an antibody heavy or light chain. Generally, the constant domains are not directly involved in the binding properties of an antibody molecule to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity. Here, “effector functions” refer to the different physiological effects of antibodies (e.g., opsonization, cell lysis, mast cell, basophil and eosinophil degranulation, and other processes) mediated by the recruitment of immune cells by the molecular interaction between the Fc domain and proteins of the immune system. The isotype of the heavy chain determines the functional properties of the antibody. Their distinctive functional properties are conferred by the carboxy-terminal portions of the heavy chains, where they are not associated with light chains.

[0095] The term “variant” refers to an amino acid sequence which differs from the native amino acid sequence of an antibody by at least one amino acid residue modification. A native or parent or wild-type amino acid sequence refers to the amino acid sequence of an antibody found in nature. “Variant” of the antibody molecule includes, but is not limited to, changes within a variable region or a constant region of a light chain and/or a heavy chain, including in the Fc region, the Fab region, the CH₁ domain, the CH₂ domain, the CH₃ domain, and the hinge region.

[0096] The term “specific” refers to the selective binding of an antibody to its target epitope. Antibody molecules can be tested for specificity of binding by comparing binding to the desired antigen to binding to unrelated antigen or analogue antigen or antigen mixture under a given set of conditions. Preferably, an antibody according to the invention will lack significant binding to unrelated antigens, or even analogs of the target antigen. Here, the term “antigen” refers to a molecule that is recognized and bound by an antibody molecule or immune-derived moiety that binds to the antigen. The specific portion of an antigen that is bound by an antibody is termed the “epitope”. A “hapten” refers to a small molecule that can, under most circumstances, elicit an immune response (i.e., act as an antigen) only when attached to a carrier, for example, a protein, polyethylene glycol (PEG), colloidal gold, silicone beads, and the like. The carrier may be one that also does not elicit an immune response by itself.

[0097] The term “antibody” is used in the broadest sense, and encompasses monoclonal, polyclonal, multispecific (e.g., bispecific, wherein each arm of the antibody is reactive with a different epitope of the same or different antigen), minibody, heteroconjugate, diabody, triabody, chimeric, and synthetic antibodies, as well as antibody fragments that specifically bind an antigen with a desired binding property and/or biological activity.

[0098] The term “monoclonal antibody” (mAb) refers to an antibody, or population of like antibodies, obtained from a population of substantially homogeneous antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, monoclonal antibodies can be made by the hybridoma method first described by Kohler G. and Milstein C. (1975), *Nature*, vol. 256:495-497, or by recombinant DNA methods.

[0099] The term “chimeric” antibody (or immunoglobulin) refers to a molecule comprising a heavy and/or light chain which is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (Cabilly, et al., *infra*; Morrison et al., *Proc. Natl. Acad. Sci. U.S.A.*, vol. 81:6851 (1984)).

[0100] The term “humanized antibody” means human antibodies that also contain selected sequences from non-human (e.g., murine) antibodies in place of the human sequences. A humanized antibody can include conservative amino acid substitutions or non-natural residues from the same or different species that do not significantly alter its binding and/or biologic activity. Such antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulins. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, camel, bovine, goat, or rabbit having the desired properties. Furthermore, humanized antibodies can comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. Thus, in general, a humanized antibody will comprise all of at least one, and in one aspect two, variable domains, in which all or all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), or that of a human immunoglobulin. See, e.g., Cabilly, et al., U.S. Pat. No. 4,816,567; Cabilly, et al., European Patent No. 0,125,023 B1; Boss, et al., U.S. Pat. No. 4,816,397; Boss, et al., European Patent No. 0,120,694 B1; Neuberger, et al., WO 86/01533; Neuberger, et al., European Patent No. 0,194,276 B1; Winter, U.S. Pat. No. 5,225,539; Winter, European Patent No. 0,239,400 B1; Padlan, et al., European Patent Application No. 0,519,596 A1; Queen, et al. (1989), *Proc. Nat'l Acad. Sci. USA*, vol. 86:10029-10033).

[0101] The term “fully human” antibody can refer to an antibody produced in a genetically engineered (ie. Transgenic) mouse (e.g. from Medarex) that, when presented with an immunogen, can produce a human antibody that does not necessarily require CDR grafting. These antibodies are fully human (100% human protein sequences) from animals such as mice in which the non-human antibody genes are suppressed and replaced with human antibody gene expression. The applicants believe that antibodies could be generated against bioactive lipids when presented to these genetically engineered mice or other animals who might be able to produce human frameworks for the relevant CDRs.

[0102] The term “bispecific antibody” can refer to an antibody, or a monoclonal antibody, having binding properties for at least two different epitopes. In one embodiment, the epitopes are from the same antigen. In another embodiment, the epitopes are from two different antigens. Methods for making bispecific antibodies are known in the art. For example, bispecific antibodies can be produced recombi-

nantly using the co-expression of two immunoglobulin heavy chain/light chain pairs. Alternatively, bispecific antibodies can be prepared using chemical linkage. One of skill can produce bispecific antibodies using these or other methods as may be known in the art. Bispecific antibodies include bispecific antibody fragments. One example of a bispecific antibody comprehended by this invention is an antibody having binding properties for an S1P epitope and an LPA epitope, which thus is able to recognize and bind to both S1P and LP1. Another example of a bispecific antibody comprehended by this invention is an antibody having binding properties for an epitope from a bioactive lipid and an epitope from a cell surface antigen. Thus the antibody is able to recognize and bind the bioactive lipid and is able to recognize and bind to cells, e.g., for targeting purposes.

[0103] The term “heteroconjugate antibody” can refer to two covalently joined antibodies. Such antibodies can be prepared using known methods in synthetic protein chemistry, including using crosslinking agents. As used herein, the term “conjugate” refers to molecules formed by the covalent attachment of one or more antibody fragment(s) or binding moieties to one or more polymer molecule(s).

[0104] The term “biologically active” refers to an antibody or antibody fragment that is capable of binding the desired epitope and in some ways exerting a biologic effect. Biological effects include, but are not limited to, the modulation of a growth signal, the modulation of an anti-apoptotic signal, the modulation of an apoptotic signal, the modulation of the effector function cascade, and modulation of other ligand interactions.

[0105] The term “recombinant DNA” refers to nucleic acids and gene products expressed therefrom that have been engineered, created, or modified by man. “Recombinant” polypeptides or proteins are polypeptides or proteins produced by recombinant DNA techniques, for example, from cells transformed by an exogenous DNA construct encoding the desired polypeptide or protein. “Synthetic” polypeptides or proteins are those prepared by chemical synthesis.

[0106] The term “expression cassette” refers to a nucleotide molecule capable of affecting expression of a structural gene (i.e., a protein coding sequence, such as an antibody of the invention) in a host compatible with such sequences. Expression cassettes include at least a promoter operably linked with the polypeptide-coding sequence, and, optionally, with other sequences, e.g., transcription termination signals. Additional regulatory elements necessary or helpful in effecting expression may also be used, e.g., enhancers. Thus, expression cassettes include plasmids, expression vectors, recombinant viruses, any form of recombinant “naked DNA” vector, and the like.

[0107] A “vector” or “plasmid” or “expression vector” refers to a nucleic acid that can be maintained transiently or stably in a cell to effect expression of one or more recombinant genes. A vector can comprise nucleic acid, alone or complexed with other compounds. A vector optionally comprises viral or bacterial nucleic acids and/or proteins, and/or membranes. Vectors include, but are not limited, to replicons (e.g., RNA replicons, bacteriophages) to which fragments of DNA may be attached and become replicated. Thus, vectors include, but are not limited to, RNA, autonomous self-replicating circular or linear DNA or RNA and include both the expression and non-expression plasmids. “Plasmids” can be commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids as

reported with published protocols. In addition, the expression vectors may also contain a gene to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

[0108] The term “promoter” includes all sequences capable of driving transcription of a coding sequence in a cell. Thus, promoters used in the constructs of the invention include cis-acting transcriptional control elements and regulatory sequences that are involved in regulating or modulating the timing and/or rate of transcription of a gene. For example, a promoter can be a cis-acting transcriptional control element, including an enhancer, a promoter, a transcription terminator, an origin of replication, a chromosomal integration sequence, 5' and 3' untranslated regions, or an intronic sequence, which are involved in transcriptional regulation. Transcriptional regulatory regions suitable for use in the present invention include but are not limited to the human cytomegalovirus (CMV) immediate-early enhancer/promoter, the SV40 early enhancer/promoter, the *E. coli* lac or trp promoters, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

[0109] A. Antibodies to Sphingolipids

[0110] The present invention provides methods for preparing antibodies directed against certain bioactive lipids, including sphingolipids. The term “sphingolipid” refers to the sphingolipids as defined by <http://www.lipidmaps.org>, including the following: Sphingoid bases [including sphing-4-enines (sphingosines), sphinganine, 4-hydroxysphinganine (phytosphingosines), sphingoid base homologs and variants, sphingoid base 1-phosphates, lysosphingomyelins and lysoglycosphingolipids; N-methylated sphingoid bases, and sphingoid base analogs]; ceramides [including N-acylsphingosines (ceramides), N-acylsphinganine (dihydroceramides), N-acyl-4-hydroxysphinganine (phytoceramides), acylceramides and ceramide 1-phosphates]; phosphosphingolipids [including ceramide phosphocholines (sphingomyelins), ceramide phosphoethanolamines and ceramide phosphoinositols; phosphosphingolipids; neutral glycosphingolipids [including the simple Glc series (GlcCer, LacCer, etc., GalNAcbl-3Galal-4Galbl-4Glc- (Globo series), GalNAcbl-4Galbl-4Glc- (Ganglio series), Galbl-3GlcNAcbl-3Galbl-4Glc- (Lacto series), Galbl-4GlcNAcbl-3Galbl-4Glc- (Neolacto series), GalNAcbl-3Galal-3Galbl-4Glc- (Isoglobo series), GlcNAcbl-2Manal-3Manbl-4Glc- (Mollu series), GalNAcbl-4GlcNAcbl-3Manbl-4Glc- (Arthro series), Gal- (Gala series) or other neutral glycosphingolipids]; acidic glycosphingolipids [including gangliosides, sulfoglycosphingolipids (sulfatides), glucuronosphingolipids, phosphoglycosphingolipids and other acidic glycosphingolipids; basic glycosphingolipids; amphoteric glycosphingolipids; arsenosphingolipids and other sphingolipids.

[0111] Anti-sphingolipid antibodies are useful for treating or preventing disorders such as hyperproliferative disorders and cardiovascular or cerebrovascular diseases and disorders, as described in greater detail below. In particular embodiments the invention is drawn to methods of preparing antibodies to S1P and its variants which include S1P itself {defined as sphingosine-1-phosphate [sphingene-1-phosphate; D-erythro-sphingosine-1-phosphate; sphing-4-enine-1-phosphate; (E,2S,3R)-2-amino-3-hydroxy-octadec-4-en-1-yl]phosphonic acid} (CAS 26993-30-6)}, or DHS1P {defined as dihydro sphingosine-1-phosphate [sphinganine-

1-phosphate; [(2S,3R)-2-amino-3-hydroxy-octadec-4-en-1-yl]phosphonic acid; D-Erythro-dihydro-D-sphingosine-1-phosphate} (CAS19794-97-9)}. Antibodies to SPC {defined as sphingosylphosphoryl choline, lysosphingomyelin, sphingosylphosphocholine, sphingosine phosphorylcholine, ethanaminium; 2-((((2-amino-3-hydroxy-4-octadecenyl)oxy)hydroxyphosphinyloxy)-N,N,N-trimethyl-, chloride, (R—(R*,S*-(E))), 2-[[[(E,2R,3 S)-2-amino-3-hydroxy-octadec-4-en-1-yl]hydroxy-phosphoryl]oxyethyl 1-trimethyl-azanium chloride (CAS 10216-23-6)]} may also be useful.

[0112] 1. A Preferred Anti-S1P Monoclonal Antibody.

[0113] A specific monoclonal anti-S1P antibody (anti-S1P mAb) is described. This antibody can be used as a therapeutic molecular sponge to selectively absorb S1P and thereby thus lower the effective in vivo extracellular concentrations of this pro-angiogenic, pro-fibrotic and tumor-facilitating factor. This can result in the reduction of tumor volume and metastatic potential, as well as the simultaneous blockage of new blood vessel formation that otherwise can feed the growing tumor. This antibody (and molecules having an equivalent activity) can also be used to treat other hyperproliferative disorders impacted by S1P, including unwanted endothelial cell proliferation, as occurs, for example, in age-related macular degeneration as well as in many cancers. In addition, the ability of S1P to protect cells from apoptosis can be reversed by the agents such as the antibody resulting in an increase in the efficacy of standard pro-apoptotic chemotherapeutic drugs.

[0114] B. Antibodies to Other Bioactive Signaling Lipids

[0115] The methods described herein can be used to prepare monoclonal antibodies against many additional extracellular and intracellular bioactive lipids beyond sphingolipids (e.g., SPC, ceramide, sphingosine, sphinganine, S1P and dihydro-S1P). Other bioactive lipid classes include the leucotrienes, eicosanoids, eicosanoid metabolites such as the HETEs, prostaglandins, lipoxins, epoxyeicosatrienoic acids and isoeicosanoids), non-eicosanoid cannabinoid mediators, phospholipids and their derivatives such as phosphatidic acid (PA) and phosphatidylglycerol (PG), cardiolipins, and lysophospholipids such as lysophosphatidyl choline (LPC) and lysophosphatidic acid (LPA). In short, this invention can be adapted for application to any desired extracellular and/or intracellular signaling bioactive lipid with pleiotropic effects on important cellular processes. Other examples of bioactive lipids include phosphatidylinositol (PI), phosphatidylethanolamine (PEA), diacylglyceride (DG), sulfatides, gangliosides, globosides and cerebroside.

[0116] C. Conjugates.

[0117] A monoclonal antibody, or antigen-binding fragment thereof, described herein can be used alone in vitro or can be administered to a subject, in non-derivatized or non-conjugated forms. In other embodiments, such antibodies, derivatives, and variants can be derivatized or linked to one or more molecular entities. Other molecular entities include naturally occurring, recombinant, or synthetic peptides, polypeptides, and proteins, non-peptide chemical compounds such as isotopes, small molecule therapeutics, etc. Preferred small molecules include radiolabels, fluorescent agents, and small molecule chemotherapeutic agents. Preferred proteins include growth factors, cytokines, and antibodies (including identical antibodies and derivatives or variants of such antibodies). The active ingredients can be linked by any suitable method, taking into account the active ingredients and the intended application, among other factors. For

example, a monoclonal antibody of the invention can be functionally linked to another molecule by chemical coupling, genetic fusion, non-covalent association, or another suitable approach.

[0118] The invention thus envisions conjugates formed between one or more monoclonal antibodies of the invention, or a variant or derivative thereof, with another active ingredient. Such conjugates may be covalent or non-covalent, and may occur via a linker or directly between the active ingredients. Examples of such conjugates include one or more monoclonal antibodies of the invention (or an antigen-binding domain thereof) linked to another therapeutic monoclonal antibody of the same or different class. Alternatively, the monoclonal antibody or antibody derivative or variant of the invention may be linked to a different class of therapeutic agent, for example, a small molecule chemotherapeutic agent or radioisotope. In some embodiments, one or more of each of two or more different therapeutic agents (at least one of which is a compound of the invention) can be linked through a multivalent scaffold.

[0119] As an alternative to conjugates, a monoclonal antibody or antibody derivative or variant of the invention may simply be associated with one or more different therapeutic agents. As an example, a monoclonal antibody of the invention can be combined with one or more other types of therapeutic agents in a delivery vehicle, e.g., a liposome, micelle, nanoparticle, etc., suitable for administration to a subject.

[0120] The invention also envisions conjugating a monoclonal antibody or antibody derivative or variant of the invention, for example, one or more CDRs reactive against a particular target bioactive lipid, with a protein or polypeptide. As an example, one or more CDRs from the variable region of an immunoglobulin heavy or light chain can be grafted into monoclonal antibody.

3. Applications

[0121] The invention is drawn to compositions and methods for treating or preventing hyperproliferative disorders such as cancer, fibrosis and angiogenesis, and cardiovascular, cardiac, and other diseases, disorders or physical trauma, and/or cerebrovascular diseases and disorders, in which therapeutic agents are administered to a patient that alters the activity or concentration of an undesirable, toxic and/or bioactive lipids, or precursors or metabolites thereof. The therapeutic methods and compositions of the invention act by changing the absolute, relative and/or available concentration and/or activities of certain undesirable or toxic lipids. Here, "toxic" refers to a particular lipid's involvement in a disease process, for example, as a signaling molecule.

[0122] Without wishing to be bound by any particular theory, it is believed that inappropriate concentrations of lipids such as LPA and/or their metabolites cause or contribute to the development of various diseases and disorders, including heart disease, neuropathic pain, cancer, angiogenesis, inflammation, and cerebrovascular disease, including stroke-like inner ear pathologies (see, e.g., Scherer, et al. (2006), Cardiovascular Research, vol. 70; 79-87). As such, the instant compositions and methods can be used to treat these diseases and disorders, particularly by decreasing the effective in vivo concentration of a particular target lipid, for example, LPA.

Several classes of diseases that may be treated in accordance with the invention are described below.

A. Hyperproliferative Diseases and Disorders

[0123] i. Cancer

[0124] One cancer therapy strategy is to reduce the biologically available extracellular levels of the tumor-promoter, S1P, either alone or in combination with traditional anti-cancer treatments, including the administration of chemotherapeutic agents, such as an anthracycline. To this end, a monoclonal antibody (mAb) has been developed that is specific for S1P, which can selectively adsorb S1P from the serum, acting as a molecular sponge to neutralize extracellular S1P. Since S1P has been shown to be pro-angiogenic, an added benefit to the antibody's effectiveness can be derived from the antibody's ability to starve the blood supply of the growing tumor. Thus, another sphingolipid-based anti-neoplastic strategy involves combining known activators of CER and SPH production (doxorubicin and related anthracycline glycosides, radiation therapy, etc.) coupled with a strategy to reduce S1P levels.

[0125] While sphingolipid-based anti-cancer strategies that target key enzymes of the sphingolipid metabolic pathway, such as SPHK, have been proposed, S1P itself has not been emphasized, largely because of difficulties in attacking this and related targets. As described herein, a highly specific monoclonal antibody to S1P has been produced that recognizes S1P in the physiological range and is capable of neutralizing S1P by molecular combination. Use of this antibody (and its derivatives) will deprive growing tumor cells of an important growth and survival factor. Moreover, use of such an antibody-based cancer therapy could also be effective when used in combination with conventional cancer treatments, such as surgery, radiation therapy, and/or the administration of cytotoxic anti-cancer agents. Examples of cytotoxic agents include, for example, the anthracycline family of drugs, the vinca alkaloids, the mitomycins, the bleomycins, the cytotoxic nucleosides, the taxanes, the epothilones, discodermolide, the pteridine family of drugs, diynes and the podophyllotoxins. Members of those classes include, for example, doxorubicin, caminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podophyllotoxin derivatives, such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine, paclitaxel and the like. Other antineoplastic agents include estramustine, cisplatin, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabine, idatrexate, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins. Other cytotoxic drugs are well known in the art. An antibody-based combination therapy may improve the efficacy of chemotherapeutic agents by sensitizing cells to apoptosis while minimizing their toxic side effects, although administration of the antibody alone may also have efficacy in delaying the progression of disease. Indeed, the ability of the anti-S1P mAb to retard tumor progression in mouse models of human cancer and in allograft mouse models demonstrates the utility of anti-S1P antibody approaches in treating both human and animal tumors. Moreover, the discovery that several human cancers types (e.g., ovarian, breast, lung, and

melanoma) can be treated in xenograft models demonstrates that the anti-S1P antibody approaches are not limited to one cancer cell or tissue type.

[0126] LPA mediates multiple cellular responses including cell proliferation, differentiation, angiogenesis and motility. A large body of experimental findings suggests that extracellular LPA plays a key role in the progression of several types of human cancer by stimulating tumor cell proliferation, survival, invasion and by inducing angiogenesis and metastasis. In addition, LPA protects a variety of tumor cell types from apoptosis. LPA has long been associated with ovarian and breast cancer [Fang, X., et al., (2002) *Biochim Biophys Acta*, 1582: 257-64]; elevated levels of LPA have been found in both blood and ascites of patients and have been correlated with tumor progression, angiogenesis and metastatic potential. Furthermore, autotoxin (ATX), the enzyme primarily responsible for LPA production, has been correlated with the metastatic and invasive properties of human tumors including melanoma, lung cancer, neuroblastoma, hepatocellular carcinoma, and glioblastoma multiforme. Thus LPA is recognized to be an innovative and promising target for cancer therapy [Mills, G. B. and W. H. Moolenaar (2003) *Nat Rev Cancer*, 3: 582-91].

[0127] It is believed that neutralizing LPA with anti-LPA antibody (such as that disclosed herein) will be a novel anti-angiogenic and anti-metastatic therapeutic approach in the treatment of cancer. Monoclonal antibodies against LPA are believed to act as a "sponge" to selectively bind LPA and thereby lower the effective in vivo extracellular levels of LPA. This is believed to result in the reduction of tumorigenesis and tumor growth as well as the simultaneous blockage of blood vessel formation and the metastatic potential. In addition, the ability of LPA to protect cells from apoptosis is likely to be lost as a result of antibody neutralization, thus increasing the efficacy of standard pro-apoptotic chemotherapeutic drugs.

[0128] ii. Angiogenesis

[0129] Angiogenesis is the process by which new blood vessels are formed from existing blood vessels. The angiogenesis associated with solid and circulating tumors is now considered to be a crucial component of tumorigenesis, as today the view that tumor growth is dependent upon neovascularization is scientifically well accepted. Both S1P and LPA appear important to the angiogenic process.

[0130] LPA is the primary regulator of $GRO\alpha$, an oncogene believed to contribute to tumorigenesis through its pro-angiogenic effect (Lee, et al (2006), *Cancer Res*, vol. 66: 2740-8). LPA also enhances expression of matrix metalloproteinase-2, a recognized player in the cell migration underlying the angiogenic process (Wu, et al. (2005), *Endocrinology*, vol. 146: 3387-3400).

[0131] S1P stimulates DNA synthesis and chemotactic motility of human venous endothelial cells (HUVECs), while inducing differentiation of multicellular structures essential early blood vessel formation. S1P also promotes the migration of bone marrow-derived endothelial cell precursors to neovascularization sites, and cells that over-express S1P receptors are resistant to the anti-angiogenic agents, thalidomide and Neovastat. Thus, S1P, and particularly S1 receptors, are required for angiogenesis and neovascularization. Finally, cross-talk occurs between S1P and other pro-angiogenic growth factors such as VEGF, EGF, PDGF, bFGF, and IL-8. For example, S1P transactivates EGF and VEGF2 receptors, and VEGF up-regulates S1P receptor expression (Igarashi, et al. (2003), *PNAS (USA)*, vol. 100: 10664-10669).

[0132] As will be appreciated, clinical control of angiogenesis is a critical component for the treatment of cancer and other angiogenesis-dependent diseases such as age-related macular degeneration (AMD) and endometriosis. Anti-angiogenic therapeutics are also particularly attractive because the vascular endothelial cells that are involved in tumor angiogenesis do not mutate as easily as do cancer cells; consequently, vascular endothelial cells are less likely than cancer cells to gain resistance to prolonged therapy, making them useful therapeutic targets.

[0133] There are several lines of evidence suggesting that S1P is a potentially significant pro-angiogenic growth factor that may be important in tumor angiogenesis, including that: anti-S1P antibodies can neutralize S1P-induced tube formation, migration of vascular endothelial cells, and protection from cell death in various in vitro assays using HUVECs; injection of breast adenocarcinoma MCF-7 cells expressing elevated S1P levels into mammary fat pads of nude mice results in an increase of angiogenesis-dependent tumors that are both larger and more numerous than when control cells are used; anti-S1P antibodies can dramatically reduce tumor-associated angiogenesis in an orthotopic murine melanoma allograft model; S1P increases new capillary growth into Matrigel plugs implanted in mice, an effect that can be neutralized by the systemic administration of anti-S1P antibodies; in vivo administration of anti-S1P antibodies can completely neutralize pro-angiogenic growth factor-induced angiogenesis (e.g., by bFGF and VEGF) in murine Matrigel plug assays; S1P stimulates the release of bFGF and VEGF from tumor cells in vitro and in vivo, an effect that can be reversed by anti-S1P antibodies; S1P enhances in vitro motility and invasion of a large number of different types of cancer cells, including glioblastoma multiforme cells; and anti-S1P antibodies significantly reduce the neovascularization associated with animal models of AMD.

[0134] The importance of S1P in the angiogenic-dependent tumors makes S1P an excellent target for cancer treatment. Indeed, antibody neutralization of extracellular S1P may result in a marked decrease in cancer progression in mammals, including humans, as a result of inhibition of blood vessel formation with concomitant loss of the nutrients and oxygen needed to support tumor growth. Thus, anti-S1P antibodies have several mechanisms of action, including: (1) direct effects on tumor cell growth; (2) indirect anti-angiogenic effects on vascular endothelial cells; and (3) the indirect anti-angiogenic effects that prevent the release and action of other pro-angiogenic growth factors. Accordingly, anti-S1P antibodies can also serve as anti-metastatic therapeutics, in addition to providing anti-angiogenic therapy.

[0135] Control of angiogenesis is a critical component for the treatment of other angiogenesis-dependent diseases besides cancer, such as age-related macular degeneration, retinopathy of prematurity, diabetic retinopathy, endometriosis, and rheumatoid arthritis (Carmeliet, P. (2005), *Nature*, vol. Vol. 438(15): 932-6).

[0136] Anti-angiogenic therapeutics are also particularly attractive because the vascular endothelial cells that are involved in tumor angiogenesis do not mutate as easily as do cancer cells; consequently, vascular endothelial cells are less likely than cancer cells to gain resistance to prolonged therapy, making them useful therapeutic targets. S1P antibodies, and derivatives thereof, will also be useful in treating other hyperproliferative disorders associated with S1P activ-

ity, such as those cause by aberrant endothelial cell proliferation, as occurs with the angiogenesis associated with AMD.

[0137] iii. Fibrogenesis and Scarring

[0138] (a) S1P, Fibroblasts, and the Remodeling Process

[0139] It is clear that cardiac fibroblasts, particularly myofibroblasts, are key cellular elements in scar formation in response to the cell death and inflammation of a myocardial infarction (MI). Myofibroblast collagen gene expression is a hallmark of remodeling and necessary for scar formation. In addition to its other activities, S1P is also an inflammatory mediator that makes profound contributions to wound healing by activating fibroblast migration and proliferation, in addition to activating platelets, stimulating angiogenesis, and promoting smooth muscle function. Thus, S1P, perhaps produced locally by injured myocardium, could, in part, be responsible for the maladaptive wound healing associated with cardiac remodeling and failure, particularly by activating myofibroblasts in the heart.

[0140] There are three general responses of cells to S1P: protection from cell death; stimulation of proliferation; and the promotion of migratory responses. Accordingly, S1P activity or involvement with a particular disorder, cell line, etc. can be assessed by adapting assays of this sort for this purpose. There is evidence that fibroblasts respond to S1P in all three ways to promote wound healing. For instance, in several of the examples in the Example section below, evidence is presented that demonstrates that S1P contributes to remodeling by promoting cardiac myofibroblast activity (proliferation, migration, and collagen gene expression).

[0141] Anti-S1P antibodies or antibody derivatives will also prevent excess scarring associated with surgical procedures. Excess scarring post injury or surgery, a problem in adult but not fetal skin tissue (Adzick and Lorenz (1994), *Ann Surg*, vol. 220: 10-18), is attributed to excess TGF- β in adult skin tissue post injury. S1P has been implicated as a potent activator the TGF- β signaling system. Accordingly, an antiS1P antibody would be expected to limit excess scarring post injury or surgery.

[0142] (b) Protection from Cell Death by LPA and S1P

[0143] LPA is an agent that protects cancer cells from apoptosis. Thus, as discussed in detail above, an antibody to LPA, for example, will make cancer cells more susceptible to chemotherapy. This has, in fact, been demonstrated in the examples hereinbelow, using newly developed anti-LPA monoclonal antibodies.

[0144] As is the case for many cell types, fibroblasts are directly protected from apoptosis by addition of S1P, and apoptosis is enhanced by inhibitors of SPHK, and S1P blocks cytochrome C release and the resultant caspase activation. Further, fibroblasts transfected with SPHK1 exhibit protection from apoptosis, an effect that may depend upon translocation of SPHK1 to the plasma membrane. It is well-established that SPHK1 up-regulates Akt, thereby regulating Bcl-2 family members and protecting from apoptosis. Also, S1P₃ is required for Akt phosphorylation in mouse embryonic fibroblasts (MEFs). Also, up-regulation of SPHK and resulting increases in S1P levels protect cardiofibroblasts from apoptosis.

[0145] Ceramide, an upstream metabolite of S1P, decreases mitochondrial membrane potential coincident with increasing the transcription of death inducing mitochondrial proteins. Because of the rheostat mechanism, S1P may have the opposite effect and protect cardiac myofibroblasts (i.e., fully differentiated fibroblasts in the heart) from apoptosis. Indeed,

S1P may even activate autophagy as a protection mechanism. These effects could be reversed by the neutralizing anti-S1P antibodies (or other molecules that bind and act to sequester S1P).

B. Pain

[0146] Bioactive lipids are believed to play important roles in the pathogenesis of pain, including neuropathic pain and pain associated with chemotherapy.

[0147] The significant role of LPA signaling in the development of neuropathic pain was established using various pharmacological and genetic approaches, including the use of mice lacking the LPA1 receptor (see e. g., Ueda, et al. (2006), *Pharmacol Ther*, vol. 109: 57-77; Inoue, et al. (2004), *Nat. Med.*, vol. 10: 712-8). Wild-type animals with nerve injury develop behavioral allodynia and hyperalgesia paralleled by demyelination in the dorsal root and increased expression of both the protein kinase C isoform within the spinal cord dorsal horn and the 21 calcium channel subunit in dorsal root ganglia. Intrathecal injection of LPA induced behavioral, morphological and biochemical changes similar to those observed after nerve ligation. In contrast, mice lacking a single LPA receptor (LPA-1, also known as EDG-2) that activates the Rho-Rho kinase pathway do not develop signs of neuropathic pain after peripheral nerve injury. Inhibitors of Rho and Rho kinase also prevented these signs of neuropathic pain. These results imply that receptor-mediated LPA signaling is crucial in the initiation of neuropathic pain and that an antibody to LPA would likely alleviate neuropathic pain in individuals suffering this condition [Moulin, DE (2006), *Pain Res Manag*, vol. 11, Suppl A: 30A-6A].

[0148] In the context of other pain, that associated with chemotherapy is a major dose limiting toxicity of many small molecule chemotherapeutic agents. Indeed, many cases of chemotherapy-induced pain have been reported. For instance, Paclitaxel (Taxol), an anti-neoplastic agent derived from the Pacific yew tree *Taxus brevifolia*, is used to treat a variety of cancers, including ovarian, breast, and non-small cell lung cancer. Paclitaxel's effectiveness, however, is limited by the highly incidental development of severe painful peripheral neuropathy such as numbness and burning pain. A monoclonal antibody against a bioactive lipid correlated with such pain, for example, LPA (or a derivative of such an antibody that contains a lipid-binding portion thereof), could be administered in combination with Paclitaxel in order to reduce the pain associated with the chemotherapeutic agent. As a result of ameliorating this dose-limiting toxicity, the amount of Paclitaxel to be administered could be even higher (and thus even more effective) when used in combination with such a monoclonal antibody or antibody derivative. In some embodiments, the chemotherapeutic agent (or other drug) could be conjugated to or otherwise associated with the antibody or antibody derivative, for example, by covalently linking the small molecule chemotherapeutic agent to the antibody, by linking the small molecule chemotherapeutic to a multivalent scaffold to which is also linked a monoclonal antibody or at least one bioactive lipid binding domain derived from a monoclonal antibody specifically reactive against the target bioactive lipid, etc.

C Cardiovascular Diseases and Disorders

[0149] Ischemic heart disease is the leading cause of death in the U.S. Each year approximately 1.5 million people suffer

heart attacks (myocardial infarctions), of which about one-third (i.e., about 500,000) are fatal. In addition, about 6.75 million Americans suffer from angina pectoris, the most common manifestation of cardiac ischemia. In total, there are more than 13 million patients living with ischemic heart disease in the U.S. alone. "Ischemia" is a condition associated with an inadequate flow of oxygenated blood to a part of the body, typically caused by the constriction or blockage of the blood vessels supplying it. Ischemia occurs any time that blood flow to a tissue is reduced below a critical level. This reduction in blood flow can result from: (i) the blockage of a vessel by an embolus (blood clot); (ii) the blockage of a vessel due to atherosclerosis; (iii) the breakage of a blood vessel (a bleeding stroke); (iv) the blockage of a blood vessel due to acute vasoconstriction; (v) a myocardial infarction (when the heart stops, the flow of blood to organs is reduced, and ischemia results); (vi) trauma; (vii) surgery, during which blood flow to a tissue or organ needs to be reduced or stopped to achieve the aims of surgery (e.g., angioplasty, heart and lung/heart transplants); (viii) exposure to certain agents, e.g., dobutamine or adenosine (Lagerqvist, et al. (1992), *Br. Heart J.*, vol. 68:282-285); or (ix) anti-neoplastic and other chemotherapeutic agents, such as doxorubicin, that are cardiotoxic.

[0150] Even if the flow rate (volume/time) of blood is adequate, ischemia may nonetheless occur due to hypoxia, i.e., a condition in which the oxygen content of blood is insufficient to satisfy normal cellular oxygen requirements of the affected area(s). Hypoxic blood is, by definition, distinct from normoxic blood, i.e., blood in which the oxygen content is sufficient to satisfy normal cellular oxygen requirements. Hypoxic conditions may result from, but are not limited to, forms of heart failure that adversely affect cardiac pumping such as hypertension, arrhythmias, septic shock, trauma, cardiomyopathies, and congestive heart disease.

[0151] Myocardial ischemic disorders occur when cardiac blood flow is restricted (ischemia) and/or when oxygen supply to the heart muscle is compromised (hypoxia) such that the heart's demand for oxygen is not met by the supply. Coronary artery disease (CAD) arising from arteriosclerosis, particularly atherosclerosis, is the most common cause of ischemia, and has symptoms such as stable or unstable angina pectoris. CAD can lead to acute myocardial infarctions (AMI) and sudden cardiac death. The spectrum of ischemic conditions that results in heart failure is referred to as Acute Coronary Syndrome (ACS). Reperfusion injury is often a consequence of ischemia, in particular when anti-coagulants, thrombolytic agents, or anti-anginal medications are used or when the cardiac vasculature is surgically opened by angioplasty or by coronary artery grafting.

[0152] Presently, treatments for acute myocardial infarction and other cardiac diseases include, but are not limited to, mechanical devices and associated procedures therewith, e.g., coronary angioplasty; thrombolytic agents such as streptokinase, tPA, and derivatives thereof. Adjuvants to these therapies include beta-blockers, aspirin and heparin, and glycoprotein (GP) IIb/IIIa inhibitors. GP IIb/IIIa inhibitors decrease platelet aggregation and thrombus formation. Examples include monoclonal antibodies (e.g., abciximab), cyclic peptides (e.g., eptifibatide), and nonpeptide peptidomimetics (e.g., tirofiban, lamifiban, xemilofiban, sibrifiban, and lefradafiban).

[0153] Preventive treatments include those that reduce a patient's cholesterol levels by, e.g., diet management and pharmacological intervention. Statins are one type of agent

used to reduce cholesterol levels. Statins are believed to act by inhibiting the activity of HMG-CoA reductase, which in turn increases the hepatic production of cholesterol receptors. Hepatic cholesterol receptors bind cholesterol and remove it from blood. Such agents include lovastatin, simvastatin, pravastatin, and fluvastatin. These and other statins slow the progression of coronary artery disease, and may induce regression of atherosclerotic lesions in patients, although the range of side effects from the use of such drugs is not fully understood.

[0154] As will be appreciated, monoclonal antibodies and derivatives, and other fragments and variants reactive against a bioactive lipid may be used to effect cardiac therapy, alone or in combination with other therapeutic approaches, including treatment with drugs and/or surgery. Here, "cardiac therapy" refers to the prevention and/or treatment of myocardial diseases, disorders, or physical trauma, including myocardial ischemia, AMI, CAD, and ACS, as well as trauma or cardiac cell and tissue damage that may occur during or as a consequence of interventional cardiology or other surgical or medical procedures or therapies that may cause ischemic or ischemic/reperfusion damage in mammals, particularly humans.

[0155] Besides the heart and brain, an anti-S1P approach can also be applied to other vascular-based, stroke-like conditions such as various inner ear pathologies (Scherer, et al. (2006), *Cardiovasc Res*, vol. 70:79-87).

D. Cerebrovascular Diseases and Disorders

[0156] Patients experiencing cerebral ischemia often suffer from disabilities ranging from transient neurological deficit to irreversible damage (stroke) or death. Cerebral ischemia, i.e., reduction or cessation of blood flow to the central nervous system, can be characterized as either focal or global. Focal cerebral ischemia refers to cessation or reduction of blood flow within the cerebral vasculature resulting from a partial or complete occlusion in the intracranial or extracranial cerebral arteries. Such occlusion typically results in stroke, a syndrome characterized by the acute onset of a neurological deficit that persists for at least 24 hours, reflecting focal involvement of the central nervous system and is the result of a disturbance of the cerebral circulation. Other causes of focal cerebral ischemia include vasospasm due to subarachnoid hemorrhage or iatrogenic intervention.

[0157] Global cerebral ischemia refers to reduction of blood flow within the cerebral vasculature resulting from systemic circulatory failure, which promptly leads to a reduction in oxygen and nutrients to tissues. Thus, global cerebral ischemia results from severe depression of cardiac performance, and is most frequently caused by AMI, although both causes include pump failure resulting from acute myocarditis or depression of myocardial contractility following cardiac arrest or prolonged cardiopulmonary bypass; mechanical abnormalities, such as severe valvular stenosis, massive aortic or mitral regurgitation, and acutely acquired ventricular septal defects; as well as from cardiac arrhythmia, such as ventricular fibrillation, or from interventional procedures, such as carotid angioplasty, stenting, endarterectomy, cardiac catheterization, electrophysiologic studies, and angioplasty.

[0158] Ischemic injury post stroke and/or MI typically leads to cell death by depolarization of critical cells with resulting rise in intracellular Na^+ and Ca^{++} followed by cell death. One channel controlling this process is the Transient

Receptor Potential Protein, a non-voltage dependent channel and recently S1P was identified as an activator of this channel through a GPCR-dependent mechanism. In addition, Transient Receptor Potential Protein, sphingosine kinase 1 and sphingosine kinase 2 share promoter regions with Egr-1, an important master switch believed to regulate cardiovascular pathobiology (Khachigian, L M (2006), *Circ Res*, vol. 98: 186-91) and Sp1, a transcription factor that plays a critical role in the death of neural cells (Simard, et al. (2006), *Nat. Med.*, vol. 12: 433-40). Based on these findings, an antibody to S1P would be expected to mitigate cell death caused by ischemia post hypoxia.

[0159] Those skilled in the art are easily able to identify patients having a stroke or at risk of having a stroke, cerebral ischemia, head trauma, or epilepsy. For example, patients who are at risk of having a stroke include those having hypertension or undergoing major surgery. Traditionally, emergent management of acute ischemic stroke consists of mainly general supportive care, e.g. hydration, monitoring neurological status, blood pressure control, and/or anti-platelet or anti-coagulation therapy. Heparin has been administered to stroke patients with limited and inconsistent effectiveness. In some circumstances, the ischemia resolves itself over a period of time due to the fact that some thrombi get absorbed into the circulation, or fragment and travel distally over a period of a few days. Tissue plasminogen activator (t-PA) or has been approved for treating acute stroke, although such systemic treatment has been associated with increased risk of intracerebral hemorrhage and other hemorrhagic complications. Aside from the administration of thrombolytic agents and heparin, there are no therapeutic options currently on the market for patients suffering from occlusion focal cerebral ischemia. Vasospasm may be partially responsive to vasodilating agents. The newly developing field of neurovascular surgery, which involves placing minimally invasive devices within the carotid arteries to physically remove the offending lesion, may provide a therapeutic option for these patients in the future, although this kind of manipulation may lead to vasospasm itself.

[0160] As will be appreciated, antibodies, antibody-derivatives, and other immune-derived moiety reactive against a bioactive lipid may be used to effect cerebrovascular therapy, alone or in combination with other therapeutic approaches, including treatment with drugs and/or surgery. Here, "cerebrovascular therapy" refers to therapy directed to the prevention and/or treatment of diseases and disorders associated with cerebral ischemia and/or hypoxia. Of particular interest is cerebral ischemia and/or hypoxia resulting from global ischemia resulting from a heart disease, as well as trauma or surgical or medical procedures or therapies that may cause ischemic or ischemic/reperfusion cerebrovascular damage in mammals, particularly humans.

E. Diagnostic and Theranostic Applications for Antibodies that Bind Bioactive Lipids

[0161] As the role of various bioactive lipids in disease is elucidated, new roles for antibody binders of bioactive lipids in diagnostics and theranostics may also be envisioned. According to the instant invention, methods are provided for enhanced detection of bioactive lipids using derivatized lipids bound to a solid support. In addition to use of these detection methods in antibody production and characterization and in research, enhanced detection of bioactive lipids may also provide a valuable diagnostic approach for diseases associated with bioactive lipids. When combined with other tech-

niques, a theranostic approach for designing optimal patient treatment is provided. One nonlimiting example is use of anti-S1P antibodies in diagnostic and theranostic methods relating to the role of S1P as a biomarker for cancer. Diagnostic and theranostic methods using antibodies targeted to LPA or other bioactive lipids, and for other disease indications, are also envisioned.

[0162] Recently, scientific literature has suggested that S1P is a potent tumorigenic growth factor that is likely released from tumor cells, and that S1P may be a novel biomarker for early-stage cancer detection. SPHK, the enzyme which is responsible for the production of S1P, is significantly up-regulated in a variety of cancer types (French, Schrecengost et al. 2003). SPHK activity is up-regulated 2-3 fold in malignant breast, colon, lung, ovarian, stomach, uterine, kidney and rectal cancer when compared to adjacent normal tissue. These workers also showed that SPHK expression varies from patient to patient, suggesting that the tumors of some patients might be more dependent on S1P than those of other patients with the same tumor type. Searching commercially available genomics database (ASCENTA, Genelogic Inc., Gaithersburg Md.) confirms that the relative expression of SPHK is, in general, significantly elevated in a wide variety of malignant tumors.

[0163] Recent publications have also suggested that S1P may be a novel cancer biomarker [Xu, Y. et al., (1998) *JAMA* 280: 719-723; Shen, Z. et al., (2001) *Gynecol Oncol* 83: 25-30; Xiao, Y. J. et al., (2001) *Anal Biochem* 290(2): 302-13; Sutphen (2004) *Cancer Epidemiology* 13(7) 1185-91]. For example, Sutphen et al, have shown that serum S1P levels are elevated in early-stage ovarian cancer patients (Sutphen 2004). One might predict from the data that breast cancer patients might also demonstrate some variability in their dependence on S1P. Taken together, these preliminary observations suggest that the success of an anti-S1P therapeutic, e.g., an anti-S1P mAb therapeutic, might be predicted for an individual patient if that patient's biopsy tissue, blood, urine or other tissue or fluid sample show elevated S1P levels.

[0164] The potential use of S1P in biological fluids has been disclosed in the following patents, all of which are commonly assigned with the instant application. U.S. Pat. No. 6,534,323, U.S. Pat. No. 6,534,322; U.S. Pat. No. 6,210,976; U.S. Pat. No. 6,858,383; U.S. Pat. No. 6,881,546; U.S. Pat. No. 7,169,390 and U.S. Pat. No. 6,500,633.

[0165] Even though humanized antibodies have low toxicity and large therapeutic indices, they are quite costly to the patient and to health care providers. Thus directing utility of the anti-S1P mAb therapeutic to those who would most likely respond to this treatment would lower risks and minimize costs, while providing optimum patient benefit.

[0166] Outlined below are a few proposed applications of biolipid diagnostics and theranostics for improved disease management.

[0167] 1. S1P may be used as a biomarker to predict individual patient therapeutic efficacy especially when combined with sphingolipid-based genomics. Based on recent findings, we would predict that S1P dependent tumors may produce their own S1P in addition to the abundant serum source of S1P. Highly aggressive tumors utilize a strategy of producing their own growth factors, and we suggest that S1P is one of the growth factors. Therefore, serum, plasma or urine measurements of total S1P from individual patients would be one predictor of patient outcome. Moreover, S1P production would be concentrated in the tumor itself and in the tumor

microenvironment (e.g., interstitial fluid). Example 11 hereinbelow describes the use of an anti-S1P mAb in an immunohistochemical method of a tumor section to assess S1P production by the tumor itself. Up-regulation of SPHK may prove useful, but since the kinase is an enzyme, it is believed that the signal as measured by S1P production will be much higher than if one relied on RNA or protein expression of the kinase itself. In addition, it is hypothesized that patients whose tumors have an up-regulation of S1P-receptors and SPHK expression are more likely to have tumors that rely on S1P as a growth factor. It is believed that these patients would benefit most from our putative anti-S1P mAb therapy. Therefore, bioassays from biopsy tissue analyzed by quantitative-PCR for the relative expression of S1P receptors and SPHK would provide a strong theranostic platform. This theranostic platform would consist of serum S1P marker analysis in combination with the genomic or proteomic quantification of S1P-related protein markers as surrogate markers of disease. This novel multi-marker analysis would provide a very strong platform for prediction of individual responsiveness to an anti-S1P mAb (SPHINGOMAB™)-based therapy.

[0168] 2. S1P may be used as a surrogate marker to titrate therapeutic regimen. The concentration of serum S1P from patients being treated with the anti-S1P mAb has the potential to be used as a surrogate marker for evaluating the course of treatment. An ELISA-based platform using patient serum, plasma or urine samples will allow for the accurate measurement of the S1P biomarker levels and to determine more precisely the anti-S1P mAb dosing regimen for individuals. Surrogate marker levels could be used in combination with the standard clinical endpoints to determine efficacy of the medical regimen.

[0169] 3. S1P may be used as a screening tool for the early detection of cancer. The early detection of cancer is of concern due to the strong correspondence of stage of progression and success of therapy. Stage I of ovarian cancer is very difficult to detect due to the fact that majority of patients are asymptomatic. By the time ovarian cancer is diagnosed, most patients are in the later stages of the disease. Detection at an earlier stage has obvious benefits to patient outcome. As described above, ovarian cancer patient serum contains a 2-fold elevation of S1P, and this elevation is easily detectable with our current ELISA platform. Since many solid tumor types, including ovarian cancer, exhibit elevated SPHK expression, it is presumed that many of the patients with these cancers would display elevated blood and/or urine S1P that could allow the clinician to intervene earlier in disease progression.

[0170] Derivatized bioactive lipids described herein can also be used to detect the level of antibodies in a fluid or tissue sample of a patient. Without being limited by the following, such immunoassays that detect the presence of anti-sphingolipid antibodies in blood and can be used to indirectly test for increased sphingolipids in patients with chronic ischemic conditions, cancer or autoimmune disorders such as multiple sclerosis. This assay is based on the assumption that patients produce anti-sphingolipid antibodies as a consequence of elevated blood levels of sphingolipids by analogy to the anti-lactosylsphingosine antibodies observed in patients with colorectal cancer (Jozwiak W. & J. Koscielak, *Eur. J. Cancer Clin. Oncol.* 18:617-621, 1982) and the anti-galactocerebro-

side antibodies detected in the sera of leprosy patients (Vemuri N. et al., *Leprosy Rev.* 67:95-103, 1996).

F. Research

[0171] The bioactive signaling lipid targets of the invention are readily adaptable for use in high-throughput screening assays for screening candidate compounds to identify those which have a desired activity, e.g., inhibiting an enzyme that catalyzes a reaction that produces an undesirable bioactive signaling lipid, or blocking the binding of a bioactive signaling lipid to a receptor therefore. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as therapeutic agents. The methods of screening of the invention comprise using screening assays to identify, from a library of diverse molecules, one or more compounds having a desired activity. A "screening assay" is a selective assay designed to identify, isolate, and/or determine the structure of, compounds within a collection that have a pre-selected activity. The collection can be a traditional combinatorial libraries are prepared according to methods known in the art, or may be purchased commercially and may be a wide-range of organic structures or structures pre-selected for potential bioactive signaling activity. By "identifying" it is meant that a compound having a desirable activity is isolated, its chemical structure is determined (including without limitation determining the nucleotide and amino acid sequences of nucleic acids and polypeptides, respectively) the structure of, and, additionally or alternatively, purifying compounds having the screened activity. Biochemical and biological assays are designed to test for activity in a broad range of systems ranging from protein-protein interactions, enzyme catalysis, small molecule-protein binding, to cellular functions. Such assays include automated, semi-automated assays and high throughput screening assays.

EXAMPLES

[0172] The invention will be further described by reference to the following detailed examples. These Examples are in no way to be considered to limit the scope of the invention in any manner.

Example 1 Synthetic Scheme for Making a Representative Thiolated Analog of S1P

[0173] The synthetic approach described in this example results in the preparation of an antigen by serial addition of structural elements using primarily conventional organic chemistry. A scheme for the approach described in this example is provided in FIG. 1, and the compound numbers in the synthetic description below refer to the numbered structures in FIG. 1.

[0174] This synthetic approach began with the commercially available 15-hydroxyl pentadecyne, 1, and activation by methyl sulphonyl chloride of the 15-hydroxy group to facilitate hydroxyl substitution to produce the sulphonate, 2. Substitution of the sulphonate with t-butyl thiol yielded the protected thioether, 3, which was condensed with Garner's aldehyde to produce 4. Gentle reduction of the alkyne moiety to an alkene (5), followed by acid catalyzed opening of the oxazolidine ring yielded S-protected and N-protected thiol substituted sphingosine, 6. During this last step, re-derivatization with di-t-butyl dicarbonate was employed to mitigate loss of the N-BOC group during the acid-catalyzed ring opening.

[0175] As will be appreciated, compound 6 can itself be used as an antigen for preparing haptens to raise antibodies to sphingosine, or, alternatively, as starting material for two different synthetic approaches to prepare a thiolated S1P analog. In one approach, compound 6 phosphorylation with trimethyl phosphate produced compound 7. Treatment of compound 7 with trimethylsilyl bromide removed both methyl groups from the phosphate and the t-butyloxycarbonyl group from the primary amine, leaving compound 8 with the t-butyl group on the sulfur as the only protecting group. To remove this group, the t-butyl group was displaced by NBS to form the disulfide, 9, which was then reduced to form the thiolated S1P analog, 10.

[0176] Another approach involved treating compound 6 directly with NBSCl to form the disulfide, 11, which was then reduced to form the N-protected thiolated S1P analog, 12. Treatment of this compound with mild acid yielded the thiolated sphingosine analog, 13, which can be phosphorylated enzymatically with, e.g., sphingosine kinase, to yield the thiolated S1P analog, 10.

[0177] Modifications of the presented synthetic approach are possible, particularly with regard to the selection of protecting and de-protecting reagents, e.g., the use of trimethyl disulfide triflate described in Example 3 to de-protect the thiol.

[0178] Compound 2. DCM (400 mL) was added to a 500 mL RB flask charged with 1 (10.3 g, 45.89 mmol), and the resulting solution cooled to 0° C. Next, TEA (8.34 g, 82.60 mmol, 9.5 mL) was added all at once followed by MsCl (7.88 g, 68.84 mmol, 5.3 mL) added drop wise over 10 min. The reaction was allowed to stir at RT for 0.5 h or until the disappearance of starting material (R_f =0.65, 5:1 hexanes: EtOAc). The reaction was quenched with NH_4Cl (300 mL) and extracted (2×200 mL) DCM. The organic layers were dried over MgSO_4 , filtered and the filtrate evaporated to a solid (13.86 g, 99.8% yield). ^1H NMR (CDCl_3) δ 4.20 (t, J =6.5 Hz, 2H), 2.98 (s, 3H), 2.59 (td, J =7 Hz, 3 Hz, 2H), 1.917 (t, J =3 Hz, 1H), 1.72 (quintet, J =7.5 Hz, 2H), 1.505 (quintet, J =7.5 Hz, 2H), 1.37 (br s, 4H), 1.27 (br s, 14H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 85.45, 70.90, 68.72, 46.69, 38.04, 30.22, 30.15, 30.14, 30.07, 29.81, 29.76, 29.69, 29.42, 29.17, 26.09, 19.06, 9.31. The principal ion observed in a HRMS analysis (ES-TOF) of compound 2 was m/z =325.1804 (calculated for $\text{C}_{16}\text{H}_{30}\text{O}_3\text{S}$: $\text{M}+\text{Na}^+$ 325.1808).

[0179] Compound 3. A three-neck IL RB flask was charged with t-butylthiol (4.54 g, 50.40 mmol) and THF (200 mL) and then placed into an ice bath. n-BuLi (31.5 mL of 1.6 M in hexanes) was added over 30 min. Next, compound 2 (13.86 g, 45.82 mmol), dissolved in THF (100 mL), was added over 2 min. The reaction is allowed to stir for 1 hour or until starting material disappeared (R_f =0.7, 1:1 hexanes/EtOAc). The reaction was quenched with saturated NH_4Cl (500 mL) and extracted with EtO_2 (2×250 mL), dried over MgSO_4 , filtered, and the filtrate evaporated to yield a yellow oil (11.67 g, 86% yield). ^1H NMR (CDCl_3) δ 2.52 (t, J =7.5 Hz, 2H), 2.18 (td, J =7 Hz, 2.5 Hz, 2H), 1.93 (t, J =2.5 Hz, 1H), 1.55 (quintet, J =7.5 Hz, 2H), 1.51 (quintet, J =7 Hz, 2H), 1.38 (br s, 4H), 1.33 (s, 9H), 1.26 (s, 14H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 85.42, 68.71, 68.67, 54.07, 42.37, 31.68, 30.58, 30.28, 30.26, 30.19, 30.17, 29.98, 29.78, 29.44, 29.19, 29.02, 19.08.

[0180] Compound 4. A 250 mL Schlenk flask charged with compound 3 (5.0 g, 16.85 mmol) was evacuated and filled with nitrogen three times before dry THF (150 mL) was added. The resulting solution cooled to -78° C. Next, n-BuLi

(10.5 mL of 1.6M in hexanes) was added over 2 min. and the reaction mixture was stirred for 18 min. at -78° C. before the cooling bath was removed for 20 min. The dry ice bath was returned. After 15 min., Garner's aldehyde (3.36 g, 14.65 mmol) in dry THF (10 mL) was then added over 5 min. After 20 min., the cooling bath was removed. Thin layer chromatography (TLC) after 2.7 hr. showed that the Garner's aldehyde was gone. The reaction was quenched with saturated aqueous NH_4Cl (300 mL) and extracted with Et_2O (2×250 mL). The combined Et_2O phases were dried over Na_2SO_4 , filtered, and the filtrate evaporated to give crude compound 4 and its syn diastereomer (not shown in FIG. 1) as a yellow oil (9.06 g). This material was then used in the next step without further purification.

[0181] Compound 5. To reduce the triple bond in compound 4, the oil was dissolved in dry Et_2O (100 mL) under nitrogen. RED-A1 (20 mL, 65% in toluene) was slowly added to the resulting solution at RT to control the evolution of hydrogen gas (H_2). The reaction was allowed to stir at RT overnight or when TLC showed the disappearance of the starting material (R_f =0.6 in 1:1 EtOAc:hexanes) and quenched slowly with cold MeOH or aqueous NH_4Cl to control the evolution of H_2 . The resulting white suspension was filtered through a Celite pad and the filtrate was extracted with EtOAc (2×400 mL). Combined EtOAc extracts were dried over MgSO_4 , filtered, and the filtrate evaporated to leave crude compound 5 and its syn diastereomer (not shown in FIG. 1) as a yellow oil (7.59 g).

[0182] Compound 6. The oil containing compound 5 was dissolved in MeOH (200 mL), PTSA hydrate (0.63 g) was added, and the solution stirred at RT for 1 day and then at 50° C. for 2 days, at which point TLC suggested that all starting material (5) was gone. However, some polar material was present, suggesting that the acid had partially cleaved the BOC group. The reaction was worked up by adding saturated aqueous NH_4Cl (400 mL), and extracted with ether (3×300 mL). The combined ether phases were dried over Na_2SO_4 , filtered, and the filtrate evaporated to dryness, leaving 5.14 g of oil. In order to re-protect whatever amine had formed, the crude product was dissolved in CH_2Cl_2 (150 mL), to which was added BOC_2O (2.44 g) and TEA (1.7 g). When TLC (1:1 hexanes/EtOAc) showed no more material remaining on the baseline, saturated aqueous NH_4Cl (200 mL) was added, and, after separating the organic phase, the mixture was extracted with CH_2Cl_2 (3×200 mL). Combined extracts were dried over Na_2SO_4 , filtered, and the filtrate concentrated to dryness to yield a yellow oil (7.7 g) which was chromatographed on a silica column using a gradient of hexanes/EtOAc (up to 1:1) to separate the diastereomers. By TLC using 1:1 PE/EtOAc, the R_f for the anti isomer, compound 6, was 0.45. For the syn isomer (not shown in FIG. 1) the R_f was 0.40. The yield of compound 6 was 2.45 g (39% overall based on Garner's aldehyde). ^1H NMR of anti isomer (CDCl_3) δ 1.26 (br s, 20H), 1.32 (s, 9H), 1.45 (s, 9H), 1.56 (quintet, 2H, J =8 Hz), 2.06 (q, 2H, J =7 Hz), 2.52 (t, 2H, J =7 Hz), 2.55 (br s, 2H), 3.60 (br s, 1H), 3.72 (ddd, 1H, J =11.5 Hz, 7.0 Hz, 3.5 Hz), 3.94 (dt, 1H, J =11.5 Hz, 3.5 Hz), 4.32 (d, 1H, J =4.5 Hz), 5.28 (br s, 1H), 5.54 (dd, 1H, J =15.5 Hz, 6.5 Hz), 5.78 (dt, 1H, J =15.5 Hz, 6.5 Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 156.95, 134.80, 129.66, 80.47, 75.46, 63.33, 56.17, 42.44, 32.98, 31.70, 30.58, 30.32, 30.31, 30.28, 30.20, 30.16, 30.00, 29.89, 29.80, 29.08, 29.03.

[0183] Anal. Calculated for $\text{C}_{27}\text{H}_{53}\text{NO}_4\text{S}$: C, 66.48; H, 10.95; N, 2.87. Found: C, 65.98; H, 10.46; N, 2.48.

[0184] Compound 7. To a solution of the alcohol compound 6 (609.5 mg, 1.25 mmol) dissolved in dry pyridine (2 mL) was added CBr_4 (647.2 mg, 1.95 mmol, 1.56 equiv). The flask was cooled in an ice bath and $\text{P}(\text{OMe})_3$ (284.7 mg, 2.29 mmol, 1.84 equiv) was added drop wise over 2 min. After 4 min. the ice bath was removed and after 12 hr. the mixture was diluted with ether (20 mL). The resulting mixture washed with aqueous HCl (10 mL, 2 N) to form an emulsion which separated on dilution with water (20 mL). The aqueous phase was extracted with ether (2×10 mL), then EtOAc (2×10 mL). The ether extracts and first EtOAc extract were combined and washed with aqueous HCl (10 mL, 2 N), water (10 mL), and saturated aqueous NaHCO_3 (10 mL). The last EtOAc extract was used to back-extract the aqueous washes. Combined organic phases were dried over MgSO_4 , filtered, and the filtrate concentrated to leave crude product (1.16 g), which was purified by flash chromatography over silica (3×22 cm column) using CH_2Cl_2 , then CH_2Cl_2 - EtOAc (1:20, 1:6, 1:3, and 1:1- product started to elute, 6:4, 7:3). Early fractions contained 56.9 mg of oil. Later fractions provided product (compound 7, 476.6 mg, 64%) as clear, colorless oil.

[0185] Anal. Calculated for $\text{C}_{29}\text{H}_{38}\text{NO}_7\text{PS}$ (595.82): C, 58.46; H, 9.81; N, 2.35. Found: C, 58.09; H, 9.69; N, 2.41.

[0186] Compound 8. A flask containing compound 7 (333.0 mg, 0.559 mmol) and a stir bar was evacuated and filled with nitrogen. Acetonitrile (4 mL, distilled from CaH_2) was injected by syringe and the flask now containing a solution was cooled in an ice bath. Using a syringe, $(\text{CH}_3)_3\text{SiBr}$ (438.7 mg, 2.87 mmol, 5.13 equiv.) was added over the course of 1 min. After 35 min., the upper part of the flask was rinsed with an additional portion of acetonitrile (1 mL) and the ice bath was removed. After another 80 min., an aliquot was removed, the solution dried by blowing nitrogen gas over it, and the residue analyzed by ^1H NMR in CDCl_3 , which showed only traces of peaks ascribed to $\text{P}-\text{OCH}_3$ moieties. After 20 min., water (0.2 mL) was added to the reaction mixture, followed by the CDCl_3 solution used to analyze the aliquot, and the mixture was concentrated to ca. 0.5 mL volume on a rotary evaporator. Using acetone (3 mL) in portions the residue was transferred to a tared test tube, forming a pale brown solution. Water (3 mL) was added in portions. After addition of 0.3 mL, cloudiness was seen. After a total of 1 mL, a gummy precipitate had formed. As an additional 0.6 mL of water was added, more cloudiness and gum separated, but the final portion of water seemed not to change the appearance of the mixture. Overall, this process was accomplished over a period of several hours. The tube was centrifuged and the supernatant removed by pipet. The solid, no longer gummy, was dried over P_4O_{10} in vacuo, leaving compound 8 (258.2 mg, 95%) as a monohydrate.

[0187] Anal. Calculated. for $\text{C}_{22}\text{H}_{46}\text{NO}_5\text{PS}+\text{H}_2\text{O}$ (485.66): C, 54.40; H, 9.96; N, 2.88. Found: C, 54.59; H, 9.84; N, 2.95.

[0188] Compound 9. Compound 8 (202.6 mg, 0.417 mmol) was added in a glove box to a test tube containing a stir bar, dry THF (3 mL) and glacial HOAc (3 mL). NBSCl (90 mg, 0.475 mmol, 1.14 equiv) were added, and after 0.5 hr., a clear solution was obtained. After a total of 9 hr., an aliquot was evaporated to dryness and the residue analyzed by ^1H NMR in CDCl_3 . The peaks corresponding to CH_2StBu and CH_2SSAr suggested that reaction was about 75% complete, and comparison of the spectrum with that of pure NBSCl in CDCl_3 suggested that none of the reagent remained in the reaction. Therefore, an additional portion (24.7 mg, 0.130 mmol, 0.31

equiv) was added, followed 3 hr. later by an additional portion (19.5 mg, 0.103 mmol, 0.25 equiv). After another 1 hr., the mixture was transferred to a new test tube using THF (2 mL) to rinse and water (1 mL) was added.

[0189] Compound 10. PMe_3 (82.4 mg, 1.08 mmol, 1.52 times the total amount of 2-nitrobenzenesulfonyl chloride added) was added to the clear solution compound 9 described above. The mixture grew warm and cloudy, with precipitate forming over time. After 4.5 hr., methanol was added, and the tube centrifuged. The precipitate settled with difficulty, occupying the bottom 1 cm of the tube. The clear yellow supernatant was removed using a pipet. Methanol (5 mL, deoxygenated with nitrogen) was added, the tube was centrifuged, and the supernatant removed by pipet. This cycle was repeated three times. When concentrated, the final methanol wash left only 4.4 mg of residue. The bulk solid residue was dried over P_4O_{10} in vacuo, leaving compound 10 (118.2 mg, 68%) as a monohydrochloride.

[0190] Anal. Calculated for $\text{C}_{18}\text{H}_{38}\text{NO}_5\text{S}+\text{HCl}$ (417.03): C, 51.84; H, 9.43; N, 3.36. Found: C, 52.11; H, 9.12; N, 3.30.

[0191] Compound 11. Compound 6 (1.45 g, 2.97 mmol) was dissolved in AcOH (20 mL), and NBSCl (0.56 g, 2.97 mmol) was added all at once. The reaction was allowed to stir for 3 hr. or until the disappearance of the starting material and appearance of the product was observed by TLC [product $R_f=0.65$, starting material $R_f=0.45$, 1:1 EtOAc /hexanes]. The reaction was concentrated to dryness on a high vacuum line and the residue dissolved in $\text{THF}/\text{H}_2\text{O}$ (100 mL of 10:1).

[0192] Compound 12. Ph_3P (0.233 g, 8.91 mmol) was added all at once to the solution above that contained compound 11 and the reaction was allowed to stir for 3 hr. or until the starting material disappeared. The crude reaction mixture was concentrated to dryness on a high vacuum line, leaving a residue that contained compound 12.

[0193] Compound 13. The residue above containing compound 12 was dissolved in DCM (50 mL) and TFA (10 mL). The mixture was stirred at RT for 5 hr. and concentrated to dryness. The residue was loaded onto a column with silica gel and chromatographed with pure DCM , followed by DCM containing 5% MeOH , then 10% MeOH , to yield the final product, compound 13, as a sticky white solid (0.45 g, 46% yield from 5). ^1H NMR (CDCl_3) δ 1.27 (s), 1.33 (br m.), 1.61 (p, 2H, $J=7.5$ Hz), 2.03 (br d, 2H, $J=7$ Hz), 2.53 (q, 2H, $J=7.5$ Hz), 3.34 (br s, 1H), 3.87 (br d, 2H, $J=12$ Hz), 4.48 (br s, 2H), 4.58 (br s, 2H), 5.42 (dd, 1H, $J=15$ Hz, 5.5 Hz), 5.82 (dt, 1H, $J=15$ Hz, 5.5 Hz), 7.91 (br s, 4H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 136.85, 126.26, 57.08, 34.76, 32.95, 30.40, 30.36, 30.34, 30.25, 30.19, 30.05, 29.80, 29.62, 29.09, 25.34.

Example 2 Synthetic Schemes for Making Thiolated Fatty Acids

[0194] The synthetic approach described in this example details the preparation of a thiolated fatty acid to be incorporated into a more complex lipid structure that could be further complexed to a protein or other carrier and administered to an animal to elicit an immune response. The approach uses conventional organic chemistry. A scheme showing the approach taken in this example is provided in FIG. 2, and the compound numbers in the synthetic description below refer to the numbered structures in FIG. 2.

[0195] Two syntheses are described. The first synthesis, for a C-12 thiolated fatty acid, starts with the commercially available 12-dodecanoic acid, compound 14. The bromine is then displaced with t-butyl thiol to yield the protected C-12 thi-

olated fatty acid, compound 15. The second synthesis, for a C-18 thiolated fatty acid, starts with the commercially available 9-bromo-nonanol (compound 16). The hydroxyl group in compound 16 is protected by addition of a dihydropyran group and the resulting compound, 17, is dimerized through activation of half of the brominated material via a Grignard reaction, followed by addition of the other half. The 18-hydroxy octadecanol (compound 18) produced following acid-catalyzed removal of the dihydropyran protecting group is selectively mono-brominated to form compound 19. During this reaction approximately half of the alcohol groups are activated for nucleophilic substitution by formation of a methane sulfonic acid ester. The alcohol is then oxidized to form the 18-bromocarboxylic acid, compound 20, which is then treated with t-butyl thiol to displace the bromine and form the protected, thiolated C-18 fatty acid, compound 21.

[0196] The protected thiolated fatty acids, each a t-butyl thioether, can be incorporated into a complex lipid and the protecting group removed using, e.g., one of the de-protecting approaches described in Examples 1 and 3. The resulting free thiol then can be used to complex with a protein or other carrier prior to inoculating animal with the hapten.

A. Synthesis of a C-12 Thiolated Fatty Acid

[0197] Compound 15. t-Butyl thiol (12.93 g, 143 mmol) was added to a dry Schlenk flask, and Schlenk methods were used to put the system under nitrogen. Dry, degassed THF (250 mL) was added and the flask cooled in an ice bath. n-BuLi (55 mL of 2.5 M in hexanes, 137.5 mmol) was slowly added over 10 min by syringe. The mixture was allowed to stir at 0° C. for an hour. The bromoacid, compound 14 (10 g, 36 mmol), was added as a solid and the reaction heated and stirred at 60° C. for 24 hr. The reaction was quenched with 2 M HCl (250 mL), and extracted with ether (2×300 mL). The combined ethereal layers were dried with magnesium sulfate, filtered, and the filtrate concentrated by rotary evaporation to yield the thioether acid, compound 15 (10 g, 99% yield) as a beige powder. ¹H NMR (CDCl₃, 500 MHz) δ 1.25-1.35 (br s, 12H), 1.32 (s, 9H), 1.35-1.40 (m, 2H), 1.50-1.60 (m, 2H), 1.60-1.65 (m, 2H), 2.35 (t, 2H, J=7.5 Hz), 2.52 (t, 2H, J=7.5 Hz). Principal ion in HRMS (ES-TOF) was observed at m/z 311.2020, calculated for M+Na⁺ 311.2015.

B. Synthesis of a C-12 Thiolated Fatty Acid

[0198] Compound 17. A dry Schlenk flask was charged with compound 16 (50 g, 224.2 mmol) and dissolved in dry degassed THF (250 mL) distilled from sodium/benzophenone. The flask was cooled in an ice bath and then PTSA (0.5 g, 2.6 mmol) was added. Dry, degassed DHP (36 g, 42.8 mmol) was then added slowly over 5 min. The mixture was allowed to warm up to RT and left to stir overnight and monitored by TLC (10:1 PE: EtOAc) until the reaction was deemed done by the complete disappearance of the spot for the bromoalcohol. TEA (1 g, 10 mmol) was then added to quench the PTSA. The mixture was then washed with cold sodium bicarbonate solution and extracted with EtOAc (3×250 mL). The organic layers were then dried with magnesium sulfate and concentrated to yield 68.2 g of crude product which was purified by column chromatography (10:1 PE: EtOAc) to yield 60 g (99% yield) of a colorless oil. ¹H NMR (CDCl₃, 500 MHz) δ 1.31 (br s, 6H), 1.41-1.44 (m, 2H), 1.51-1.62 (obscured multiplets, 6H), 1.69-1.74 (m, 1H),

1.855 (quintet, J=7.6 Hz, 2H), 3.41 (t, J=7 Hz, 2H), 3.48-3.52 (m, 2H), 3.73 (dt, 2H, J=6.5 Hz), 3.85-3.90 (m, 2H), 4.57 (t, 2H, J=3 Hz).

[0199] Compound 18. Magnesium shavings (2.98 g, 125 mmol) were added to a flame-dried Schlenk flask along with a crystal of iodine. Dry THF (200 mL) distilled from sodium was then added and the system was degassed using Schlenk techniques. Compound 17 (30 g, 97 mmol) was then slowly added to the magnesium over 10 min. and the solution was placed in an oil bath at 65° C. and allowed to stir overnight. The reaction was deemed complete by TLC by quenching an aliquot with acetone and observing the change in RF in a 10:1 PE:EtOAc mixture. The Grignard solution was then transferred by cannula to a three-necked flask under nitrogen containing additional compound 17 (30 g, 97 mmol). The flask containing the resulting mixture was then cooled to 0° C. in an ice bath and a solution of Li₂CuCl₄ (3 mL of 1 M) was then added via syringe. The reaction mixture turned a very dark blue within a few minutes. This mixture was left to stir overnight. The next morning the reaction was deemed complete by TLC (10:1 PE:EtOAc), quenched with a saturated NH₄Cl solution, and then extracted into ether (3×250 mL). The ether layers were dried with magnesium sulfate and concentrated to yield crude product (40 g), which was dissolved in MeOH. Concentrated HCl (0.5 mL) was then added, which resulted in the formation of a white emulsion, which was left to stir for 3 hr. The white emulsion was then filtered to yield 16 g (58% yield) of the pure diol, compound 18. ¹H NMR (CDCl₃, 200 MHz) δ 1.26 (br s, 24H), 1.41-1.42 (m, 4H), 1.51-1.68 (m, 4H), 3.65 (t, 4H, J=6.5 Hz).

[0200] Compound 19. The symmetrical diol, compound 18 (11 g, 38.5 mmol), was added to a dry Schlenk flask under nitrogen, then dry THF (700 mL) distilled from sodium was added. The system was degassed and the flask put in an ice bath. Diisopropylethylamine (6.82 mL, 42.3 mmol) was added via syringe, followed by MsCl (3.96 g, 34.4 mmol) added slowly, and the mixture was left to stir for 1 hr. The reaction was quenched with saturated NaH₂PO₄ solution (300 mL), and then extracted with EtOAc (3×300 mL). The organic layers were then combined, dried with MgSO₄, and concentrated to yield 14 g of a mixture of the diol, monomesylate, and dimesylate. NMR showed a 1:0.8 mixture of CH₂OH: CH₂OMs protons. The mixture was then dissolved in dry THF (500 mL), deoxygenated, and to it was added LiBr (3.5 g, 40.23 mmol). This mixture was allowed reflux overnight, upon which the reaction was quenched with water (150 mL), and extracted with EtOAc (3×250 mL). The organic layer was then dried with MgSO₄, and concentrated to yield a mixture of brominated products that were then purified by flash chromatography (DCM) to yield compound 19 (3.1 g, 25% yield) as a white powder. ¹H NMR (CDCl₃, 500 MHz) δ 1.26 (br s, 26H), 1.38-1.46 (m, 2H), 1.55 (quintet, 2H, J=7.5 Hz), 1.85 (quintet, 2H, J=7.5 Hz), 3.403 (t, 2H, J=6.8 Hz), 3.66 (t, 2H, J=6.8 Hz).

[0201] Compound 20. A round bottom flask was charged with compound 19 (2.01 g, 5.73 mmol) and the solid dissolved in reagent grade acetone (150 mL). Simultaneously, Jones reagent was prepared by dissolving CrO₃ (2.25 g, 22 mmol) in H₂SO₄ (4 mL) and then slowly adding 10 mL of cold water and letting the solution stir for 10 min. The cold Jones reagent was then added to the round bottom flask slowly over 5 min., after which the solution stirred for 1 hr. The resulting orange solution turned green within several minutes. The mixture was then quenched with water (150

mL) extracted twice in ether (3×150 mL). The ether layers were then dried with magnesium sulfate, and concentrated to yield compound 20 (2.08 g, 98% yield) as a white powder. ¹H NMR (CDCl₃, 200 MHz) δ 1.27 (br s, 26H), 1.58-1.71 (m, 2H), 1.77-1.97 (m, 2H), 2.36 (t, 2H, J=7.4 Hz), 3.42 (t, 2H, J=7 Hz).

[0202] Compound 21. t-Butylthiol (11.32 g, 125 mmol) was added to a dry Schlenk flask and dissolved in dry THF (450 mL) distilled from sodium. The solution was deoxygenated by bubbling nitrogen through it before the flask was placed in an ice bath. n-BuLi solution in hexanes (70 mL of 1.6 M) was then added slowly via syringe over 10 min. This mixture was allowed to stir for 1 hr., then compound 20 (5.5 g, 16.2 mmol) was added and the solution was left to reflux at 60°C. overnight. The next morning an aliquot was worked up, analyzed by NMR, and the reaction deemed complete. The reaction was quenched with HCl (200 mL of 2 M) and extracted with ether (3×250 mL). The ethereal layers were then dried with magnesium sulfate, filtered, and the filtrate concentrated to yield the product, compound 21, as a white solid (5 g, 90% yield). ¹H NMR (CDCl₃, 200 MHz) δ 1.26 (br s, 26H), 1.32 (br s, 9H), 1.48-1.70 (m, 4H), 2.35 (t, 2H, J=7.3 Hz), 2.52 (t, 2H, J=7.3 Hz). ¹³C NMR (CDCl₃, 200 MHz) δ 24.69, 28.35, 29.05, 29.21, 29.28, 29.39, 29.55, 29.89, 31.02 (3C), 33.98, 41.75, 179.60.

Example 3 Synthetic Scheme for Making a Thiolated Analog of LPA

[0203] The synthetic approach described in this example results in the preparation of thiolated LPA. The LPA analog can then be further complexed to a carrier, for example, a protein carrier, which can then be administered to an animal to elicit an immunogenic response to LPA. This approach uses both organic chemistry and enzymatic reactions, the synthetic scheme for which is provided in FIG. 3. The compound numbers in the synthetic description below refer to the numbered structures in FIG. 3.

[0204] The starting materials were compound 15 in Example 2 and enantiomerically pure glycerophosphocholine (compound 22). These two chemicals combined to yield the di-acetylated product, compound 23, using DCC to facilitate the esterification. In one synthetic process variant, the resulting di-acetylated glycerophosphocholine was treated first with phospholipase-A2 to remove the fatty acid at the sn-2 position of the glycerol backbone to produce compound 24. This substance was further treated with another enzyme, phospholipase-D, to remove the choline and form compound 26. In another synthetic process variant, the phospholipase-D treatment preceded the phospholipase-A2 treatment to yield compound 25, and treatment of compound 25 with phospholipase-D then yields compound 26. Both variants lead to the same product, the phosphatidic acid derivative, compound 26. The t-butyl protecting group in compound 26 is then removed, first using trimethyl disulfide triflate to produce compound 27, followed by a disulfide reduction to produce the desired LPA derivative, compound 28. As those in the art will appreciate, the nitrobenzyl sulfonyl reaction sequence described in Example 1 can also be used to produce compound 28.

[0205] Compound 23. To a flame-dried Schlenk flask were added the thioether acid, compound 15 (10 g, 35.8 mmol), compound 22 (glycerophosphocholine-CdCl₂ complex, 4.25 g, 8.9 mmol), DCC (7.32 g, 35.8 mmol), and DMAP (2.18 g, 17.8 mmol), after which the flask was evacuated and filled

with nitrogen. A minimal amount of dry, degassed DCM was added (100 mL), resulting in a cloudy mixture. The flask was covered with foil and then left to stir until completed, as by TLC (silica, 10:5:1 DCM: MeOH: concentrated NH₄OH). The insolubility of compound 16 precluded monitoring its disappearance by TLC, but the reaction was stopped when the product spot of R_f 0.1 was judged not to be increasing in intensity. This typically required 3 to 4 days, and in some cases, addition of more DCC and DMAP. Upon completion, the reaction mixture was filtered, and the filtrate concentrated to yield a yellow oil, which was purified using flash chromatography using the solvent system described above to yield 3.6 g (50% yield) of a clear wax containing a mixture of compound 23 and monoacylated products in a ratio of 5 to 1, as estimated from comparing the integrals for the peaks for the (CH₃)₃N—, —CH₂StBu and —CH₂COO— moieties. Analysis of the oil by HRMS (ESI-TOF) produced a prominent ion at m/z 820.4972, calculated for M+Na⁺ = C₄₀H₈₀NNaO₈PS₂⁺ 820.4960.

[0206] A. Synthesis Variant 1—Phospholipase-A2 Treatment

[0207] Compound 24. A mixture of compound 23 and monoacylated products as described above (3.1 g, 3.9 mmol) was dissolved in Et₂O (400 mL) and methanol (30 mL). Borate buffer (100 mL, pH 7.4 0.1M, 0.072 mM in CaCl₂) was added, followed by phospholipase-A2 (from bee venom, 130 units, Sigma). The resulting mixture was left to stir for 10 hr., at which point TLC (silica, MeOH:water 4:1—the previous solvent system 10:5:1 DCM: MeOH: concentrated NH₄OH proved ineffective) showed the absence of the starting material (R_f=0.7) and the appearance of a new spot (R_f=0.2). The organic and aqueous layers were separated and the aqueous layer was washed with ether (2×250 mL). The product was extracted from the aqueous layer with a mixture of DCM:MeOH (2:1, 2×50 mL). The organic layers were then concentrated by rotary evaporation to yield product as a white wax (1.9 g, 86% yield) that NMR showed to be a pure product (compound 24). ¹H NMR (CDCl₃, 500 MHz) δ 1.25-1.27 (br s, 12H), 1.31 (s, 9H), 1.35-1.45 (m, 2H), 1.52-1.60 (m, 4H), 2.31 (t, 2H, J=7.5 Hz), 2.51 (t, 2H, J=7.5 Hz), 3.28 (br s, 9H), 3.25-3.33 (br s, 2H), 3.78-3.86 (m, 1H), 3.88-3.96 (m, 2H), 4.04-4.10 (m, 2H), 4.26-4.34 (m, 2H). Analysis of the wax by HRMS (ESI-TOF) produced a prominent ion at m/z 550.2936, calculated for M+Na⁺ 550.2943 (C₂₄H₅₀NNaO₇PS₂⁺), and an m/z at 528.3115, calculated for MH⁺ 528.3124 (C₂₄H₅₁NO₇PS₂⁺).

[0208] Anal. Calculated. for C₂₄H₅₀NO₇PS + 2H₂O (563.73): C, 51.13; H, 9.66; N, 2.48. Found: C, 50.90; H, 9.37; N, 2.76.

[0209] Compound 26. The lyso compound 24 (1.5 g, 2.7 mmol) was dissolved in a mixture of sec-butanol (5 mL) and Et₂O (200 mL), and the resulting cloudy mixture was sonicated until the cloudiness dissipated. Buffer (200 mL, pH 5.8, 0.2 M NaOAc, 0.08 M CaCl₂) was added, followed by cabbage extract (80 mL of extract from savory cabbage (which contains phospholipase-D), containing 9 mg of protein/mL). The reaction was stirred for 1 day and monitored by TLC (C₁₈RP SiO₂, 5:1 ACN:water), R_f of starting material and product=0.3 and 0.05, respectively. In order to push the reaction to completion, as needed an additional portion of cabbage extract (50 mL) was added and the reaction stirred a further day. This process was repeated twice more, as needed to complete the conversion. When the reaction was complete, the mixture was concentrated on the rotary evaporator to

remove the ether, and then EDTA solution (0.5 M, 25 mL) was added and the product extracted into a 5:4 mixture of MeOH: DCM (300 mL). Concentration of the organic layer followed by recrystallization of the residue from DCM and acetone afforded pure product (0.9 g, 75% yield). ^1H NMR (CDCl_3 , 200 MHz) δ 1.25-1.27 (br s, 12H), 1.33 (s, 9H), 1.52-1.60 (m, 4H), 2.34 (t, 2H, $J=7.5$ Hz), 2.52 (t, 2H, $J=7.5$ Hz), 3.6-3.8 (br s, 1H), 3.85-3.97 (br s, 2H), 4.02-4.18 (m, 2H).

[0210] Compound 27. The protected sample LPA, compound 26 (0.150 g, 0.34 mmol), was methanol washed and added to a vial in the glove box. This was then suspended in a mixture of AcOH:THF (1:1, 10 mL), which never fully dissolved even after 1 hr. of sonication. Solid $[\text{Me}_2\text{SSMe}]\text{OTf}$ (0.114 g, 0.44 mmol) was then added. This was left to stir for 18 hr. The reaction was monitored by removing an aliquot, concentrating it to dryness under vacuum, and re-dissolving or suspending the residue in CD_3OD for observing the ^1H NMR shift of the CH_2 peak closest to the sulfur. The starting material had a peak at 2.52 ppm, whereas the unsymmetrical disulfide formed at this juncture had a peak at around 2.7 ppm. This material (compound 27) was not further isolated or characterized.

[0211] Compound 28. The mixture containing compound 27 was treated with water (100 μL) immediately followed by PMe_3 (0.11 g, 1.4 mmol). After stirring for 3 hr. the solvent was removed by vacuum to yield an insoluble white solid. Methanol (5 mL) was added, the mixture centrifuged, and the mother liquor decanted. Vacuum concentration yielded 120 mg (91% yield) of compound 28, a beige solid. Compound 28 is a thiolated LPA hapten that can be conjugated to a carrier, for example, albumin or KLH, via disulfide bond formation. Characterization of compound 28: ^1H NMR (1:1 CD_3OD : $\text{CD}_3\text{CO}_2\text{D}$, 500 MHz) δ 1.25-1.35 (br s, 12H), 1.32-1.4 (m, 2H), 1.55-1.6 (m, 4H), 2.34 (t, 2H, $J=7$), 2.47 (t, 2H, $J=8.5$), 3.89-3.97 (br s, 2H), 3.98-4.15 (m, 2H), 4.21 (m, 1H). Negative ion ES of the sample dissolved in methanol produced a predominant ion at $m/z=385.1$.

Example 4 Antibodies to S1P

[0212] One type of therapeutic antibody specifically binds undesirable sphingolipids to achieve beneficial effects such as, e.g., (1) lowering the effective concentration of undesirable, toxic sphingolipids (and/or the concentration of their metabolic precursors) that would promote an undesirable effect such as a cardiotoxic, tumorigenic, or angiogenic effect; (2) to inhibit the binding of an undesirable, toxic, tumorigenic, or angiogenic sphingolipids to a cellular receptor therefore, and/or to lower the concentration of a sphingolipid that is available for binding to such a receptor. Examples of such therapeutic effects include, but are not limited to, the use of anti-S1P antibodies to lower the in vivo serum concentration of available S1P, thereby blocking or at least limiting S1P's tumorigenic and angiogenic effects and its role in post-MI heart failure, cancer, or fibrogenic diseases.

[0213] Thiolated S1P (compound 10 of FIG. 1) was synthesized to contain a reactive group (i.e., a sulfhydryl group) capable of cross-linking the essential structural features of S1P to a carrier moiety such as KLH. Prior to immunization, the thio-S1P analog was conjugated via IOA or SMCC cross-linking to protein carriers (e.g., KLH) using standard protocols. SMCC is a heterobifunctional crosslinker that reacts with primary amines and sulfhydryl groups, and represents a preferred crosslinker.

[0214] Swiss Webster or BALB-C mice were immunized four times over a two month period with 50 μg of immunogen (SMCC facilitated conjugate of thiolated-S1P and KLH) per injection. Serum samples were collected two weeks after the second, third, and fourth immunizations and screened by direct ELISA for the presence of anti-S1P antibodies. Spleens from animals that displayed high titers of the antibody were subsequently used to generate hybridomas per standard fusion procedures. The resulting hybridomas were grown to confluency, after which the cell supernatant was collected for ELISA analysis. Of the 55 mice that were immunized, 8 were good responders, showing significant serum titers of antibodies reactive to S1P. Fusions were subsequently carried out using the spleens of these mice and myeloma cells according to established procedures. The resulting 1,500 hybridomas were then screened by direct ELISA, yielding 287 positive hybridomas. Of these 287 hybridomas screened by direct ELISA, 159 showed significant titers. Each of the 159 hybridomas was then expanded into 24-well plates. The cell-conditioned media of the expanded hybridomas were then re-screened to identify stable hybridomas capable of secreting antibodies of interest. Competitive ELISAs were performed on the 60 highest titer stable hybridomas.

[0215] Of the 55 mice and almost 1,500 hybridomas screened, one hybridoma was discovered that displayed performance characteristics that justified limited dilution cloning, as is required to ultimately generate a true monoclonal antibody. This process yielded 47 clones, the majority of which were deemed positive for producing S1P antibodies. Of these 47 clones, 6 were expanded into 24-well plates and subsequently screened by competitive ELISA. From the 4 clones that remained positive, one was chosen to initiate large-scale production of the S1P monoclonal antibody. SCID mice were injected with these cells and the resulting ascites was protein A-purified (50% yield) and analyzed for endotoxin levels (<3 EU/mg). For one round of ascites production, 50 mice were injected, producing a total of 125 mL of ascites. The antibodies were isotyped as IgG1 kappa, and were deemed $>95\%$ pure by HPLC. The antibody was prepared in 20 mM sodium phosphate with 150 mM sodium chloride (pH 7.2) and stored at -70°C .

[0216] The positive hybridoma clone (designated as clone 306D326.26) was deposited with the ATCC (safety deposit storage number SD-5362), and represents the first murine mAb directed against S1P. The clone also contains the variable regions of the antibody heavy and light chains that could be used for the generation of a "humanized" antibody variant, as well as the sequence information needed to construct a chimeric antibody.

[0217] Screening of serum and cell supernatant for S1P-specific antibodies was by direct ELISA using the thiolated S1P analog described in Example 1 (i.e., compound 10) as the antigen. A standard ELISA was performed, as described below, except that 50 μL of sample (serum or cell supernatant) was diluted with an equal volume of PBS/0.1% Tween-20 (PBST) during the primary incubation. ELISAs were performed in 96-well high binding ELISA plates (Costar) coated with 0.1 μg of chemically-synthesized compound 10 conjugated to BSA in binding buffer (33.6 mM Na_2CO_3 , 100 mM NaHCO_3 ; pH 9.5). The thiolated-S1P-BSA was incubated at 37°C for 1 hr. at 4°C overnight in the ELISA plate wells. The plates were then washed four times with PBS (137 mM NaCl, 2.68 mM KCl, 10.14 mM Na_2HPO_4 , 1.76 mM KH_2PO_4 ; pH 7.4) and blocked with PBST for 1 hr. at room

temperature. For the primary incubation step, 75 μ L of the sample (containing the S1P to be measured), was incubated with 25 μ L of 0.1 μ g/mL anti-S1P mAb diluted in PBST and added to a well of the ELISA plate. Each sample was performed in triplicate wells. Following a 1 hr. incubation at room temperature, the ELISA plates were washed four times with PBS and incubated with 100 μ L per well of 0.1 μ g/mL HRP goat anti-mouse secondary (Jackson ImmunoResearch) for 1 hr. at room temperature. Plates were then washed four times with PBS and exposed to tetramethylbenzidine (Sigma) for 1-10 minutes. The detection reaction was stopped by the addition of an equal volume of 1M H_2SO_4 . Optical density of the samples was determined by measurement at 450 nm using an EL- X-800 ELISA plate reader (Bio-Tech).

[0218] For cross reactivity, a competitive ELISA was performed as described above, except for the following alterations (FIG. 4). The primary incubation consisted of the competitor (S1P, SPH, LPA, etc.) and a biotin-conjugated anti-S1P mAb. Biotinylation of the purified monoclonal antibody was performed using the EZ-Link Sulfo-NHS-Biotinylation kit (Pierce). Biotin incorporation was determined as per kit protocol and ranged from 7 to 11 biotin molecules per antibody. The competitor was prepared as follows: lipid stocks were sonicated and dried under argon before reconstitution in DPBS/BSA [1 mg/ml fatty acid-free BSA (Calbiochem) in DPBS (Invitrogen 14040-133)]. Purified anti-S1P mAb was diluted as necessary in PBS/0.5% Triton X-100. Competitor and antibody solutions were mixed together so to generate 3 parts competitor to 1 part antibody. A HRP-conjugated streptavidin secondary antibody (Jackson ImmunoResearch) was used to generate signal.

[0219] Another aspect of the competitive ELISA data shown in FIG. 4 is that it shows that the anti-S1P mAb was unable to distinguish the thiolated-S1P analog (compound 10) from the natural S1P that was added in the competition experiment. It also demonstrates that the antibody does not recognize any oxidation products because the analog was constructed without any double bonds (as is also true for the LPA analog described in Example 3). The anti-S1P mAb was also tested against natural product containing the double bond that was allowed to sit at room temperature for 48 hours. Reverse phase HPLC of the natural S1P was performed according to methods reported previously (Deutschman, et al. (July 2003), *Am Heart J.*, vol. 146(1):62-8), and the results showed no difference in retention time. Further, a comparison of the binding characteristics of the monoclonal antibody to the various lipids shown in FIG. 4 indicates that the epitope recognized by the antibody do not involve the hydrocarbon chain in the region of the double bond of natural S1P. On the other hand, the epitope recognized by the monoclonal antibody is the region containing the amino alcohol on the sphingosine base backbone plus the free phosphate. If the free phosphate is linked with a choline (as is the case with SPC), then the binding was somewhat reduced. If the amino group is esterified to a fatty acid (as is the case with C1P), no antibody binding was observed. If the sphingosine amino alcohol backbone was replaced by a glycerol backbone (as is the case with LPA), then the S1P-specific monoclonal exhibited no binding. These epitope mapping data indicate that there is only one epitope on S1P recognized by the monoclonal antibody, and that this epitope is defined by the unique polar headgroup of S1P.

[0220] In a similar experiment using ELISA measurements, suitable control materials were evaluated to ensure

that this anti-S1P monoclonal antibody did not recognize either the protein carrier or the crosslinking agent. For example, the normal crosslinker SMCC was exchanged for IOA in conjugating the thiolated-S1P to BSA as the laydown material in the ELISA. When IOA was used, the antibody's binding characteristics were nearly identical to when BSA-SMCC-thiolated-S1P was used. Similarly, KLH was exchanged for BSA as the protein that was complexed with thiolated-S1P as the laydown material. In this experiment, there was also no significant difference in the binding characteristics of the antibody.

[0221] Binding kinetics: The binding kinetics of S1P to its receptor or other moieties has, traditionally, been problematic because of the nature of lipids. Many problems have been associated with the insolubility of lipids. For BIAcore measurements, these problems were overcome by directly immobilizing S1P to a BIAcore chip. Antibody was then flowed over the surface of the chip and alterations in optical density were measured to determine the binding characteristics of the antibody to S1P. To circumvent the bivalent binding nature of antibodies, S1P was coated on the chip at low densities. Additionally, the chip was coated with various densities of S1P (7, 20, and 1000 RU) and antibody binding data was globally fit to a 1:1 interaction model. FIG. 5 demonstrates the changes in optical density due to the binding of the monoclonal antibody to S1P at three different densities of S1P. Overall, the affinity of the monoclonal antibody to S1P was determined to be very high, in the range of approximately 88 picomolar (pM) to 99 nM, depending on whether a monovalent or bivalent binding model was used to analyze the binding data.

Example 5 Cloning and Characterization of the Variable Domains of an S1P Monoclonal Antibody

A. Introduction

[0222] The manufacture of biological products is complex, in part because of the complexity associated with the variability of the protein itself. For monoclonal antibodies (mAbs), variability can be localized to the protein backbone or to the carbohydrate moieties appended to these glycosylated proteins. For example, heterogeneity can be attributed to the formation of alternative disulfide pairings, deamidation and the formation of isoaspartyl residues, methionine and cysteine oxidation, cyclization of N-terminal glutamine residues to pyroglutamate and partial enzymatic cleavage of C-terminal lysines by mammalian carboxypeptidases. On the other hand, carbohydrate heterogeneity introduced during cell culture includes differential addition of fucose, alternative mannose branching linkages, and differential presence of terminal sialylation. In addition, mutagenesis can be performed to alter glycosylation patterns. Oxidation is also a source of concern. For instance, the recombinant humanized monoclonal antibody HER2 undergoes oxidation in liquid formulations when exposed to intense light and elevated temperatures. Interestingly, such oxidation was reported to be formulation dependent (Lam, et al. (1997), *Pharm. Sci.*, vol. 86: 1250-1255), and the presence of NaCl in the formulation reportedly caused an increase in oxidation at higher temperatures after contact with stainless steel containers or stainless steel components in the filling process. The methionine residue at position 255 in the heavy chain of the Fc region was determined to be the primary site of oxidation. The oxidation was eliminated by supplementing the media with methionine

and thiosulfate caused by free radicals generated by the presence of metal ions and peroxide impurities in the formulation. For reasons such as these, process engineering is commonly applied to antibody molecules to improve their properties, such as enhanced expression in heterologous systems, resistance to proteases, reduced aggregation, and enhanced stability.

[0223] This example reports the cloning of the murine mAb against S1P. This antibody, termed Sphingomab™, is an IgG1 monoclonal antibody. The overall strategy consisted of cloning the murine variable domains of both the light chain (VL) and the heavy chain (VH). The consensus sequence of 306D VH shows that the constant region fragment is consistent with a gamma 2b isotype. The murine variable domains were cloned together with the constant domain of the light chain (CL) and with the constant domain of the heavy chain (CH1, CH2, and CH3), resulting in a chimeric antibody construct. Also, Sphingomab™ is unique because of the presence of a free cysteine residue in the Fab region at position 50 on the heavy chain in the CDR2 region. Replacing this residue could greatly facilitate formulation and manufacturing processes, as well as improving yields. Indeed, in an effort to improve the biophysical properties of the antibody molecule, substitution of the cysteine residue at position 50 with a panel of amino acid residues was performed by creating a series of constructs containing the desired substitution. These constructs were then expressed in mammalian cells, and the different antibody variants were compared in an ELISA assay for binding to S1P. Compared with the chimeric antibody, the resulting mutants carrying the substitution Cys50Ser and Cys50Arg exhibited a slight decrease in binding to S1P whereas the replacement of Cys with Phe or Ala did not alter the binding to S1P.

B. Materials and Methods

[0224] 1. Antibody Gene Cloning

[0225] A clone from the anti-S1P hybridoma cell line 306D326.1 (ATCC#SD-5362) was grown in DMEM (Dulbecco's Dulbecco's Modified Eagle Medium with GlutaMAX™ I, 4500 mg/L D-Glucose, Sodium Puruvate; Gibco/Invitrogen, Carlsbad, Calif., 111-035-003), 10% FBS (Sterile Fetal Clone I, Perbio Science), and 1× glutamine/ Penicillin/Streptomycin (Gibco/Invitrogen). Total RNA was isolated from 10⁷ hybridoma cells using a procedure based on the RNeasy Mini kit (Qiagen, Hilden Germany). The RNA was used to generate first strand cDNA following the manufacturer's protocol (1st strand synthesis kit, Amersham Biosciences).

[0226] The immunoglobulin heavy chain variable region (VH) cDNA was amplified by PCR using an MHV7 primer (MHV7: 5'-ATGGRATGGAGCKGGRTCTTMTCTT-3' [SEQ ID NO: 1]) in combination with a IgG2b constant region primer MHCG1/2a/2b/3 mixture (MHCG1: 5'-CAGTGGATAGACAGATGGGGG-3' [SEQ ID NO: 2]; MHCG2a: 5'-CAGTGGATAGACCGATGGGGC-3' [SEQ ID NO: 3]; MHCG2b: 5'-CAGTGGATAGACTGATGGGGG-3' [SEQ ID NO: 4]; MHCG3: 5'-CAAGGGATAGACAGATGGGGC-3' [SEQ ID NO: 5]). The product of the reaction was ligated into the pCR2.1®-TOPO® vector (Invitrogen) using the TOPO-TA Cloning® kit and sequence. The variable domain of the heavy chain was then amplified by PCR from this vector and inserted as a Hind III and Apa I fragment and ligated into the expression vector pGID200 (see U.S. Pat. No. 7,060,808) or pG4D200 (id.) containing the

HCMVi promoter, a leader sequence, and the gamma-1 constant region to generate the plasmid pGID200306DVH. The consensus sequence of 306D V_H (FIG. 6; SEQ ID NO: 6) showed that the constant region fragment was consistent with a gamma 2b isotype.

[0227] Similarly, the immunoglobulin kappa chain variable region (VK) was amplified using the MKV 20 primer (5'-GTCTCTGATTCTAGGGCA-3' [SEQ ID NO: 7]) in combination with the kappa constant region primer MKC (5'-ACTGGATGGTGGGAAGATGG-3' [SEQ ID NO: 8]). The product of this reaction was ligated into the pCR2.1®-TOPO vector using the TOPO-TA Cloning® kit and sequence. The variable domain of the light chain was then amplified by PCR and then inserted as a Bam HI and Hind III fragment into the expression vector pKN100 (see U.S. Pat. No. 7,060,808) containing the HCMV promoter, a leader sequence, and the human kappa constant domain, generating plasmid pKN100306DVK.

[0228] The heavy and light chain plasmids pGID200306DVH plus pKN100306DVK were transformed into DH4a bacteria and stocked in glycerol. Large-scale plasmid DNA was prepared as described by the manufacturer (Qiagen, endotoxin-free MAXIPREP™ kit). DNA samples, purified using Qiagen's QIAprep Spin Miniprep Kit or EndoFree Plasmid Mega/Maxi Kit, were sequenced using an ABI 3730x1 automated sequencer, which also translates the fluorescent signals into their corresponding nucleobase sequence. Primers were designed at the 5' and 3' ends so that the sequence obtained would overlap. The length of the primers was 18-24 bases, and preferably they contained 50% GC content and no predicted dimers or secondary structure. The amino acid sequences for the mouse V_H and V_L domains from Sphingomab™ are shown in FIG. 6 (SEQ ID NOS: 6 and 9, respectively). In FIG. 6, the CDR residues (see Kabat, E A (1982), Pharmavol Rev, vol. 34: 23-38) are boxed, and are shown below in Table 1.

TABLE 1

Mouse Sphingomab™ CDR sequences of the mouse V _H and V _L domains		
		CDR
<u>VL CDR</u>		
ITTTDIDDDMN	(SEQ ID NO: 10)	CDR1
EGNILRP	(SEQ ID NO: 11)	CDR2
LQSDNLFPFT	(SEQ ID NO: 12)	CDR3
<u>VH CDR</u>		
DHTIH	(SEQ ID NO: 13)	CDR1
CISPRHDITKYNEMFRG	(SEQ ID NO: 14)	CDR2
GGFYGSTIWFDF	(SEQ ID NO: 15)	CDR3

[0229] The complete nucleotide and amino acid sequences of several chimeric antibody V_H and V_L domains are shown in FIG. 7. In FIG. 7, the amino acid sequences are numbered, and the CDRs identified, according to the Kabat method (Kabat, et al. (1991), NIH National Technical Information Service, pp. 1-3242).

[0230] 2. COS 7 Expression

[0231] For antibody expression in a non-human mammalian system, plasmids were transfected into the African green monkey kidney fibroblast cell line COS 7 by electroporation (0.7 ml at 10^7 cells/ml) using 10 μ g of each plasmid. Transfected cells were plated in 8 ml of growth medium for 4 days. The chimeric 306DH1 $\times$ 306DVK-2 antibody was expressed at 1.5 μ g/ml in transiently co-transfected COS cell conditioned medium. The binding of this antibody to S1P was measured using the S1P ELISA.

[0232] The expression level of the chimeric antibody was determined in a quantitative ELISA as follows. Microtiter plates (Nunc MaxiSorp immunoplate, Invitrogen) were coated with 100 μ l aliquots of 0.4 μ g/ml goat anti-human IgG antibody (Sigma, St. Louis, Mo.) diluted in PBS and incubate overnight at 4° C. The plates were then washed three times with 200 μ l/well of washing buffer (1 $\times$ PBS, 0.1% TWEEN). Aliquots of 200 μ L of each diluted serum sample or fusion supernatant were transferred to the toxin-coated plates and incubated for 37° C. for 1 hr. Following 6 washes with washing buffer, the goat anti-human kappa light chain peroxidase conjugate (Jackson Immuno Research) was added to each well at a 1:5000 dilution. The reaction was carried out for 1 hr at room temperature, plates were washed 6 times with the washing buffer, and 150 μ L of the K-BLUE substrate (Sigma) was added to each well, incubated in the dark at room temperature for 10 min. The reaction was stopped by adding 50 μ l of RED STOP solution (SkyBio Ltd.) and the absorption was determined at 655 nm using a Microplate Reader 3550 (Bio-Rad Laboratories Ltd.). Results from the antibody binding assays are shown in FIG. 8.

[0233] 3. 293F Expression

[0234] For antibody expression in a human system, plasmids were transfected into the human embryonic kidney cell line 293F (Invitrogen) using 293fectin (Invitrogen) and using 293F-FreeStyle Media (Invitrogen) for culture. Light and heavy chain plasmids were both transfected at 0.5 g/mL. Transfections were performed at a cell density of 10^6 cells/mL. Supernatants were collected by centrifugation at 1100 rpm for 5 minutes at 25° C. 3 days after transfection. Expression levels were quantified by quantitative ELISA (see below) and varied from 0.25-0.5 g/mL for the chimeric antibody.

[0235] 4. Quantitative ELISA

[0236] Microtiter ELISA plates (Costar) were coated with rabbit anti-mouse IgG, F(ab')₂ fragment specific (Jackson Immuno Research) or rabbit anti-human, IgG F(ab')₂ fragment specific (Jackson Immuno Research) diluted in 1 M Carbonate Buffer (pH 9.5) at 37° C. for 1 hr. Plates were washed with PBS and blocked with PBS/BSA/Tween-20 for 1 hr at 37° C. For the primary incubation, dilutions of non-specific mouse IgG or human IgG, whole molecule (used for calibration curve) and samples to be measured were added to the wells. Plates were washed and incubated with 100 μ l per well of HRP conjugated goat anti-mouse (H+L) diluted 1:40,000 (Jackson Immuno Research) or HRP conjugated goat anti-human (H+L) diluted 1:50,000 (Jackson Immuno Research) for 1 hr at 37° C. After washing, the enzymatic reaction was detected with Tetramethylbenzidine (Sigma) and stopped by adding 1 M H₂SO₄. The optical density (OD) was measured at 450 nm using a Thermo Multiskan EX. Raw data were transferred to GraphPad software for analysis.

[0237] 5. Direct ELISA

[0238] Microtiter ELISA plates (Costar) were coated overnight with S1P diluted in 1 M Carbonate Buffer (pH 9.5) at

37° C. for 1 hr. Plates were washed with PBS (137 mM NaCl, 2.68 mM KCl, 10.1 mM Na₂HPO₄, 1.76 mM KH₂PO₄; pH 7.4) and blocked with PBS/BSA/Tween-20 for 1 hr at room temp or overnight at 4° C. For the primary incubation (1 hr at room temp.), a standard curve using the anti-S1P mAb and the samples to be tested for binding was built using the following set of dilutions: 0.4 μ g/mL, 0.2 μ g/mL, 0.1 μ g/mL, 0.05 μ g/mL, 0.0125 μ g/mL, and 0 μ g/mL, and 100 μ l added to each well. Plates were washed and incubated with 100 μ l per well of HRP conjugated goat anti-mouse (1:20,000 dilution) (Jackson Immuno Research) or HRP conjugated goat anti-human (H+L) diluted 1:50,000 (Jackson Immuno Research) for 1 hr at room temperature. After washing, the enzymatic reaction was detected with tetramethylbenzidine (Sigma) and stopped by adding 1 M H₂SO₄. The optical density (OD) was measured at 450 nm using a Thermo Multiskan EX. Raw data were transferred to GraphPad software for analysis.

[0239] Table 2, below, shows a comparative analysis of mutants with the chimeric antibody. To generate these results, bound antibody was detected by a second antibody, specific for the mouse or human IgG, conjugated with HRP. The chromogenic reaction was measured and reported as Optical density (OD). The concentration of the panel of antibodies was 0.1 μ g/ml. No interaction of the second antibody with S1P-coated matrix alone was detected.

TABLE 2

Variable Domain	Mutation	Plasmids	Binding
HC	Chimeric	pATH50 + pATH 10	1.5
	CysAla	pATH50 + pATH11C1	2
	CysSer	pATH50 + pATH 12C2	0.6
	CysArg	pATH50 + pATH14C1	0.4
LC	CysPhe	pATH50 + pATH16C1	2
	MetLeu	pATH53C1 + pATH10	1.6

Example 6 Chimeric mAb to S1P

[0240] As used herein, the term “chimeric” antibody (or “immunoglobulin”) refers to a molecule comprising a heavy and/or light chain which is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (Cabilly et al., supra; Morrison et al., Proc. Natl. Acad. Sci. U.S.A. 81:6851 (1984)).

[0241] A chimeric antibody to S1P was generated using the variable regions (Fv) containing the active S1P binding regions of the murine antibody from a particular hybridoma (ATCC safety deposit storage number SD-5362) with the Fc region of a human IgG1 immunoglobulin. The Fc regions contained the CL, ChL, and Ch3 domains of the human antibody. Without being limited to a particular method, chimeric antibodies could also have been generated from Fc regions of human IgG1, IgG2, IgG3, IgG4, IgA, or IgM. As those in the art will appreciate, “humanized” antibodies can be generated by grafting the complementarity determining regions (CDRs, e.g. CDR1-4) of the murine anti-S1P mAb with a human antibody framework regions (e.g., Fr1, Fr4, etc.) such

as the framework regions of an IgG1. FIG. 9 shows the binding of the chimeric and full murine mAbs in a direct ELISA measurement using thiolated-S1P as lay down material.

[0242] For the direct ELISA experiments shown in FIG. 9, the chimeric antibody to S1P had similar binding characteristics to the fully murine monoclonal antibody. ELISAs were performed in 96-well high-binding ELISA plates (Costar) coated with 0.1 μ g of chemically-synthesized, thiolated S1P conjugated to BSA in binding buffer (33.6 mM Na_2CO_3 , 100 mM NaHCO_3 ; pH 9.5). The thiolated S1P-BSA was incubated at 37° C. for 1 hr. or at 4° C. overnight in the ELISA plate. Plates were then washed four times with PBS (137 mM NaCl, 2.68 mM KCl, 10.14 mM Na_2HPO_4 , 1.76 mM KH_2PO_4 ; pH 7.4) and blocked with PBST for 1 hr. at room temperature. For the primary incubation step, 75 μ L of the sample (containing the S1P to be measured), was incubated with 25 μ L of 0.1 μ g/mL anti-S1P monoclonal antibody diluted in PBST and added to a well of the ELISA plate. Each sample was performed in triplicate wells. Following a 1 hr incubation at room temperature, the ELISA plates were washed four times with PBS and incubated with 100 μ L per well of 0.1 μ g/mL HRP goat anti-mouse secondary (Jackson ImmunoResearch) for 1 hr. at room temperature. Plates were then washed four times with PBS and exposed to tetramethylbenzidine (Sigma) for 1-10 minutes. The detection reaction was stopped by the addition of an equal volume of 1M H_2SO_4 . Optical density of the samples was determined by measurement at 450 nm using an EL- X-800 ELISA plate reader (Bio-Tech).

[0243] As was the case with regard to the experiments described in Example 4, the preferred method of measuring either antibody titer in the serum of an immunized animal or in cell-conditioned media (i.e., supernatant) of an antibody-producing cell such as a hybridoma, involves coating the ELISA plate with a target ligand (e.g., a thiolated analog of S1P, LPA, etc.) that has been covalently linked to a protein carrier such as BSA.

[0244] Without being limited to particular method or example, chimeric antibodies could be generated against other lipid targets such as LPA, ceramides, sulfatides, cerebrosides, cardiolipins, phosphatidylserines, phosphatidylinositols, phosphatidic acids, phosphatidylcholines, phosphatidylethanolamines, eicosinoids, and other leukotrienes, etc. Further, many of these lipids could also be glycosylated and/or acetylated, if desired.

Example 7 Antibody-Based Assay for Sphingosine Kinase (SPH Kinase)

[0245] Sphingosine Kinase (SPH kinase or SPHK) catalyzes the conversion of SPH to S1P. A genetic sequence encoding human SPH-kinase has been described (Melendez et al., *Gene* 251:19-26, 2000). Three human homologs of SPH kinase (SKA, SKB, and SKC) have been described (published PCT patent application WO 00/52173). Murine SPH kinase has also been described (Kohama et al., *J. Biol. Chem.* 273:23722-23728, 1998; and published (PCT patent application WO 99/61581). Published PCT patent application WO 99/61581 reports nucleic acids encoding a sphingosine kinase. Published PCT patent application WO 00/52173 reports nucleic acids encoding homologues of sphingosine kinase. Other SPH kinases have also been reported. See, e.g., Pitson et al., *Biochem J.* 350:429-441, 2000; published PCT application WO 00/70028; Liu et al., *J. Biol. Chem.*, 275: 19513-19520, 2000; PCT/AU01/00539, published as WO

01/85953; PCT/US01/04789, published as WO 01/60990; and PCT/EP00/09498, published as WO 01/31029.

[0246] Inhibitors of SPH kinase include, but are not limited to, N,N-dimethylsphingosine (Edsall et al., *Biochem.* 37:12892-12898, 1998); D-threo-dihydrosphingosine (Olivera et al., *Nature* 365:557-560, 1993); and Sphingoid bases (Jonghe et al., "Structure-Activity Relationship of Short-Chain Sphingoid Bases As Inhibitors of Sphingosine Kinase", *Bioorganic & Medicinal Chemistry Letters* 9:3175-3180, 1999)

[0247] Assays of SPH kinase useful for evaluating these and other known or potential SPH kinase inhibitors include those disclosed by Olivera et al., *Methods in Enzymology*, 311:215-223, 1999; Caligan et al., *Analytical Biochemistry*, 281:36-44, 2000.

[0248] Inhibition of SPH kinase is believed to lead to an accumulation of its substrate, SPH, which, like S1P, can be an undesirable sphingolipid in certain conditions. In order to avoid or mitigate these undesirable effects, an agent could be administered that (i) stimulates an enzyme that utilizes SPH as a substrate, provided that the enzyme should not be one that yields S1P as a reaction product (such as, e.g., ceramide synthase; see below); or (ii) inhibits an enzyme that yields SPH as a product.

[0249] Without being limited to a particular method, anti-S1P antibodies (e.g., a monoclonal anti-S1P antibody) could be used as a reagent in an in vitro assay for SPH kinase activity. For example, purified SPHK could be added to the wells of a microtiter plate in the presence of PBS and the substrate for the kinase, SPH (complexed with, for example, fatty-acid free BSA). The resulting product of the reaction, S1P, could then be followed by ELISA using an anti-S1P antibody (e.g., the monoclonal anti-S1P antibody described above in Example 4). In such an assay, inhibition of SPHK by a test compound would result in lower levels of S1P than in a control reaction that did not include an SPHK inhibitory compound. Such an assay could be configured for high throughput, and could thus serve as the basis of a high throughput screening assay for modulators of SPHK activity.

Example 8 Antibody-Based Assay for S1P Lyase or SPP Activities

[0250] The stimulation of enzymes that catalyze reactions that degrade S1P (i.e., reactions that utilize S1P as a reactant) will result in the stimulation of degradation of S1P molecules. Such enzymes include, but are not limited to:

[0251] S-1-P Lyase: S1P lyase catalyzes the conversion of S1P to ethanolamine-P (also known as t-2-hexadecanal) and palmitaldehyde (Veldhoven et al., *Adv. Lipid Res.* 26:67-97, 1993; Van Veldhoven, *Methods in Enzymology*, 311:244-254, 1999). Yeast (Lantermann et al., *Biochem. J.* 332:525-531, 1998), murine (Zhou et al., *Biochem. Biophys. Res. Comm.* 242:502-507, 1998), and human (published PCT patent application WO 99/38983) S1P lyase genes have been reported. Published PCT patent application WO 99/16888 reports S1P lyase DNA and protein sequences. U.S. Pat. No. 6,187,562 and published PCT patent application WO 99/38983 also report an S1P lyase.

[0252] Gain-of-function assays can be developed to discover small molecule compounds that would activate the lyase or increase the expression of the gene encoding it. Without being limited to a particular method, one could use anti-S1P antibodies in an ELISA format to measure the production of S1P from added SPH in in vitro or cell-based

formats. Compounds identified as stimulating S1P lyase activity, either directly at the enzyme or indirectly by elevating the expression level of the gene encoding the enzyme (for example, by gene activation, enhancing S1P lyase mRNA stability, etc.), could be investigated further, as such compounds may prove useful in lowering the extracellular concentration of S1P in patients where S1P levels correlate with toxicity, such as in the treatment of cancer, cardio and cerebrovascular disease, autoimmune disorders, inflammatory disorders, angiogenesis, fibrotic diseases, and age-related macular degeneration.

[0253] S1P Phosphatase: S1P phosphatase (also known as SPP phosphohydrolase) is a mammalian enzyme that catalyzes the conversion of S-1-P to sphingosine (Mandala et al., *Proc. Nat. Acad. Sci.* 95:150-155, 1998; Mandala et al., *Proc. Nat. Acad. Sci.* 97:7859-7864, 2000; Mandala, Prostaglandins & other Lipid Mediators, 64:143-156, 2001; Brindley et al., *Methods in Enzymology*, 311:233-244, 1999). Two S-1-P phosphatases, LBP1 and LBP2, have been isolated from yeast (Mandala et al., *J. Biol. Chem.* 272:32709-32714, 1997); PCT/UW01/03879, published as WO01/57057.

[0254] As with S1P lyase, gain-of-function assays can be developed to discover compounds that would activate S1P phosphatase or increase the expression of the gene encoding it. For example, one can use anti-S1P antibodies in an ELISA format to measure the production of S1P from added SPH in vitro or cell-based formats. Compounds identified as stimulating S1P phosphatase activity, either directly at the enzyme or indirectly by elevating the expression level of the gene encoding the enzyme (for example, by gene activation, enhancing S1P phosphatase mRNA stability, etc.), could be investigated further, as such compounds may prove useful in lowering the extracellular concentration of S1P in patients where S1P levels correlate with toxicity, such as cancer, cardio and cerebrovascular disease, autoimmune disorders, inflammatory disorders, angiogenesis, fibrotic diseases, and age-related macular degeneration.

Example 9 Production and Characterization of Monoclonal Antibodies to LPA

Antibody Production

[0255] Although polyclonal antibodies against naturally-occurring LPA have been reported in the literature (Chen J H, et al., *Bioorg Med Chem Lett.* 2000 Aug. 7; 10(15):1691-3), monoclonal antibodies have not been described. Using an approach similar to that described in Example 4, a C-12 thio-LPA analog (compound 28 in Example 3) as the key component of a hapten formed by the cross-linking of the analog via the reactive SH group to a protein carrier (KLH) via standard chemical cross-linking using either IOA or SMCC as the cross-linking agent, monoclonal antibodies against LPA were generated. To do this, mice were immunized with the thio-LPA-KLH hapten (in this case, thiolated-LPA:SMCC:KLH) using methods described in Example 4 for the generation of anti-S1P monoclonal antibodies. Of the 80 mice immunized against the LPA analog, the five animals that showed the highest titers against LPA (determined using an ELISA in which the same LPA analog (compound 28) as used in the hapten was conjugated to BSA using SMCC and laid down on the ELISA plates) were chosen for moving to the hybridoma phase of development.

[0256] The spleens from these five mice were harvested and hybridomas were generated by standard techniques. Briefly, one mouse yielded hybridoma cell lines (designated 504A). Of all the plated hybridomas of the 504A series, 66 showed positive antibody production as measured by the previously-described screening ELISA.

[0257] Table 3, below, shows the antibody titers in cell supernatants of hybridomas created from the spleens of two of mice that responded to an LPA analog hapten in which the thiolated LPA analog was cross-linked to KLH using heterobifunctional cross-linking agents. These data demonstrate that the anti-LPA antibodies do not react either to the crosslinker or to the protein carrier. Importantly, the data show that the hybridomas produce antibodies against LPA, and not against S1P.

TABLE 3

LPA hybridomas					
mouse #	3rd bleed titer OD at 1:312,500	Supernatants from 24 well	LPA binding OD at 1:20	S1P binding OD at 1:20	Cross reactivity w/ S1P*
1	1.242	1.A.63	1.197	0.231	low
		1.A.65	1.545	0.176	none
2	0.709	2.B.7	2.357	0.302	low
		2.B.63	2.302	0.229	low
		2.B.83	2.712	0.175	none
		2.B.104	2.57	0.164	none
		2.B.1B7	2.387	0.163	none
		2.B.3A6	2.227	0.134	none

*Cross reactivity with S1P from 24 well supernatants

high = OD > 1.0-2.0 at [1:20]

mid = OD 0.4-1.0 at [1:20]

low = OD 0.4-0.2 at [1:20]

none = OD < 0.2 OD at [1:20]

[0258] The development of anti-LPA mAbs in mice was monitored by ELISA (direct binding to 12:0 and 18:1 LPA and competition ELISA). A significant immunological response was observed in at least half of the immunized mice and five mice with the highest antibody titer were selected to initiate hybridoma cell line development following spleen fusion.

[0259] After the initial screening of over 2000 hybridoma cell lines generated from these 5 fusions, a total of 29 anti-LPA secreting hybridoma cell lines exhibited high binding to 18:1 LPA. Of these hybridoma cell lines, 24 were further subcloned and characterized in a panel of ELISA assays. From the 24 clones that remained positive, six hybridoma clones were selected for further characterization. Their selection was based on their superior biochemical and biological properties.

Direct Binding Kinetics

[0260] The binding of 6 anti-LPA mAbs (B3, B7, B58, A63, B3A6, D22) to 12:0 and 18:1 LPA (0.1 uM) was measured by ELISA. EC₅₀ values were calculated from titration curves using 6 increasing concentrations of purified mAbs (0 to 0.4 ug/ml). EC₅₀ represents the effective antibody concentration with 50% of the maximum binding. Max denotes the maximal binding (expressed as OD450). Results are shown in Table 4.

TABLE 4

Direct Binding Kinetics of Anti-LPA mAbs							
		B3	B7	B58	D22	A63	B3A6
12:0 LPA	EC ₅₀ (nM)	1.420	0.413	0.554	1.307	0.280	0.344
	Max (OD450)	1.809	1.395	1.352	0.449	1.269	1.316
18:1 LPA	EC ₅₀ (nM)	1.067	0.274	0.245	0.176	0.298	0.469
	Max (OD450)	1.264	0.973	0.847	0.353	1.302	1.027

[0261] The kinetics parameters k_a (association rate constant), k_d (dissociation rate constant) and K_D (association equilibrium constant) were determined for the 6 lead candidates using the BIAcore 3000 Biosensor machine. In this study, LPA was immobilized on the sensor surface and the anti-LPA mAbs were flowed in solution across the surface. As shown, all six mAbs bound LPA with similar K_D values ranging from 0.34 to 3.8 pM and similar kinetic parameters.

The Anti-LPA Murine mAbs Exhibit High Affinity to LPA

[0262] LPA was immobilized to the sensor chip at densities ranging 150 resonance units. Dilutions of each mAb were passed over the immobilized LPA and kinetic constants were obtained by nonlinear regression of association/dissociation phases. Errors are given as the standard deviation using at least three determinations in duplicate runs. Apparent affinities were determined by $K_D = k_d/k_a$. k_a = Association rate constant in $M^{-1}s^{-1}$, k_d = Dissociation rate constant in s^{-1}

TABLE 5

Affinity of anti-LPA mAb for LPA			
mAbs	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	K_D (pM)
A63	$4.4 \pm 1.0 \times 10^5$	1×10^{-6}	2.3 ± 0.5
B3	$7.0 \pm 1.5 \times 10^5$	1×10^{-6}	1.4 ± 0.3
B7	$6.2 \pm 0.1 \times 10^5$	1×10^{-6}	1.6 ± 0.1
D22	$3.0 \pm 0.9 \times 10^4$	1×10^{-6}	33 ± 10
B3A6	$1.2 \pm 0.9 \times 10^6$	$1.9 \pm 0.4 \times 10^{-5}$	16 ± 1.2

Specificity Profile of Six Anti-LPA mAbs

[0263] Many isoforms of LPA have been identified to be biologically active and it is preferable that the mAb recognize all of them to some extent to be of therapeutic relevance. The specificity of the anti-LPA mAbs was evaluated utilizing a competition assay in which the competitor lipid was added to the antibody-immobilized lipid mixture. Competition ELISA assays were performed with 6 mAbs to assess their specificity. 18:1 LPA was captured on ELISA plates. Each competitor lipid (up to 10 uM) was serially diluted in BSA (1 mg/ml)-PBS and then incubated with the mAbs (3 nM). Mixtures were then transferred to LPA coated wells and the amount of bound antibody was measured with a secondary antibody. Data are normalized to maximum signal (A_{450}) and are expressed as percent inhibition. Assays were performed in triplicate. IC_{50} : Half maximum inhibition concentration; MI: Maximum inhibition (% of binding in the absence of inhibitor); - - - : not estimated because of weak inhibition. A high inhibition result indicates recognition of the competitor lipid by the antibody. As shown in Table 6, all the anti-LPA mAbs recognized the different LPA isoforms.

TABLE 6

Specificity profile of six anti-LPA mAbs.										
	14:0 LPA		16:0 LPA		18:1 LPA		18:2 LPA		20:4 LPA	
	IC_{50} uM	MI %	IC_{50} uM	MI %	IC_{50} uM	MI %	IC_{50} uM	MI %	IC_{50} uM	MI %
504B3	0.02	72.3	0.05	70.3	0.287	83	0.064	72.5	0.02	67.1
504B7	0.105	61.3	0.483	62.9	>2.0	100	1.487	100	0.161	67
504B58-3F8	0.26	63.9	5.698	>100	1.5	79.3	1.240	92.6	0.304	79.8
504B104	0.32	23.1	1.557	26.5	28.648	>100	1.591	36	0.32	20.1
504D22-1	0.164	34.9	0.543	31	1.489	47.7	0.331	31.4	0.164	29.5
504A63-1	1.147	31.9	5.994	45.7	—	—	—	—	0.119	14.5
504B3A6-1	0.108	59.9	1.151	81.1	1.897	87.6	—	—	0.131	44.9

Interestingly, the anti-LPA mAbs were able to discriminate between 12:0 (lauroyl), 14:0 (myristoyl), 16:0 (palmitoyl), 18:1 (oleoyl), 18:2 (linoleoyl) and 20:4 (arachidonoyl) LPAs. The rank order for EC_{50} was for the unsaturated 18:2>18:1>20:4 and for the saturated lipids 14:0>16:0>18:0. mAbs with high specificity are desirable for ultimate drug development. The specificity of the anti-LPA mAbs was assessed for their binding to LPA related biolipids such as distearoyl-phosphatidic acid, lysophosphatidylcholine, S1P, ceramide and ceramide-1-phosphate. None of the six antibodies demonstrated cross-reactivity to distearoyl PA and LPC, the immediate metabolic precursor of LPA.

Example 10 Anti-Cancer Activities of Anti-LPA Monoclonal Antibodies

Cancer Cell Proliferation

[0264] LPA is a potent growth factor supporting cell survival and proliferation by stimulation of G_i , G_q and $G_{12/13}$ via GPCR-receptors and activation of downstream signaling events. Cell lines were tested for their proliferative response to LPA (0.01 mM to 10 mM). Cell proliferation was assayed by using the cell proliferation assay kit from Chemicon (Temecula Calif.) (Panc-1) and the Cell-Blue titer from Pierce (Caki-1). Each data point is the mean of three independent experiments. LPA increased proliferation of 7 human-derived tumor cell lines in a dose dependent manner including SKOV3 and OVCAR3 (ovarian cancer), Panc-1 (pancreatic cancer), Caki-1 (renal carcinoma cell), DU-145 (prostate cancer), A549 (lung carcinoma), and HCT-116 (colorectal adenocarcinoma) cells and one rat-derived tumor cell line, RBL-2H3 (rat leukemia cells). Even though tumor-derived cells normally have high basal levels of proliferation, LPA appears to further augment proliferation in most tumor cell lines. Anti-LPA mAbs (B7 and B58) were assessed for the ability to inhibit LPA-induced proliferation in selected human cancer cell lines. The increase in proliferation induced by LPA was shown to be mitigated by the addition of anti-LPA mAb.

Anti-LPA mAb Sensitizes Tumor Cells to Chemotherapeutic Agents

[0265] The ability of LPA to protect ovarian tumor cells against apoptosis when exposed to clinically-relevant levels of the chemotherapeutic agent, paclitaxel (Taxol) was investigated. SKVO3 cells were treated with 1% FBS (S), Taxol

(0.5 mM), +/-anti-LPA mAbs for 24 h. LPA protected SKVO3 cells from Taxol-induced apoptosis. Apoptosis was assayed by measurement of the caspase activity as recommended by the manufacturer (Promega). As anticipated, LPA protected most of the cancer cell lines tested from taxol-induced cell death. When anti-LPA antibody was added to a selection of the LPA responsive cells, the anti-LPA antibody blocked the ability of LPA to protect cells from death induced by the cytotoxic chemotherapeutic agent. Moreover, the anti-LPA antibody was able to remove the protection provided by serum. Serum is estimated to contain about 5-20 mM LPA. Taxol induced caspase-3,7 activation in SKOV3 cells and the addition of serum to cells protected cells from apoptosis. Taxol-induced caspase activation was enhanced by the addition of all 3 of the anti-LPA mAbs to the culture medium. This suggests that the protective and anti-apoptotic effects of LPA were removed by the selective antibody mediated neutralization of the LPA present in serum.

Anti-LPA mAb Inhibits LPA-Mediated Migration of Tumor Cells

[0266] An important characteristic of metastatic cancers is that the tumor cells escape contact inhibition and migrate away from their tissue of origin. LPA has been shown to promote metastatic potential in several cancer cell types. Accordingly, we tested the ability of anti-LPA mAb to block LPA-dependent cell migration in several human cancer cell lines by using the cell monolayer scratch assay. Cells were seeded in 96 well plates and grown to confluence. After 24 h of starvation, the center of the wells was scratched with a pipette tip. In this art-accepted "scratch assay," the cells respond to the scratch wound in the cell monolayer in a stereotyped fashion by migrating toward the scratch and close the wound. Progression of migration and wound closure are monitored by digital photography at 10 $\times$ magnification at desired timepoints. Cells were not treated (NT), treated with LPA (2.5 mM) with or w/o mAb B7 (10 μ g/ml) or an isotype matching non-specific antibody (NS) (10 μ g/ml). In untreated cells, a large gap remains between the monolayer margins following the scratch. LPA-treated cells in contrast, have only a small gap remaining at the same timepoint, and a few cells are making contact across the gap. In cells treated with both LPA and the anti-LPA antibody B7, the gap at this timepoint was several fold larger than the LPA-only treatment although not as large as the untreated control cells. This shows that the anti-LPA antibody had an inhibitory effect on the LPA-stimulated migration of renal cell carcinoma (Caki-1) cells. Similar data were obtained with mAbs B3 and B58. This indicates that the anti-LPA mAb can reduce LPA-mediated migration of cell lines originally derived from metastatic carcinoma. Anti-LPA mAbs Inhibit Release of Pro-Tumorigenic Cytokines from Tumor Cells

[0267] LPA is involved in the establishment and progression of cancer by providing a pro-growth tumor microenvironment and promoting angiogenesis. In particular, increases of the pro-growth factors such as IL-8 and VEGF have been observed in cancer cells. IL-8 is strongly implicated in cancer progression and prognosis. IL-8 may exert its effect in cancer through promoting neovascularization and inducing chemotaxis of neutrophils and endothelial cells. In addition, overexpression of IL-8 has been correlated to the development of a drug resistant phenotype in many human cancer types.

[0268] Three anti-LPA mAbs (B3, B7 and B58) were tested for their abilities to reduce in vitro IL-8 production compared

to a non-specific antibody (NS). Caki-1 cells were seeded in 96 well plates and grown to confluence. After overnight serum starvation, cells were treated with 18:1 LPA (0.2 mM) with or without anti-LPA mAb B3, B7, B58 or NS (Non-Specific). After 24h, cultured supernatants of renal cancer cells (Caki-1), treated with or without LPA and in presence of increasing concentrations of the anti-LPA mAbs B3, B7 and B58, were collected and analyzed for IL-8 levels using a commercially available ELISA kit (Human Quantikine Kit, R&D Systems, Minneapolis, Minn.). In cells pre-treated with the anti-LPA mAbs, IL-8 expression was significantly reduced in a dose-dependent manner (from 0.1-30 μ g/mL mAb) whereas LPA increased the expression of IL-8 by an average of 100% in non-treated cells. Similar results were obtained with the other well-known pro-angiogenic factor, VEGF. The inhibition of IL-8 release by the anti-LPA mAbs was also observed in other cancerous cell lines such as the pancreatic cell line Panc-1. These data suggest that the blockade of the pro-angiogenic factor release is an additional and potentially important effect of these anti-LPA mAbs.

Anti-LPA mAbs Inhibit Angiogenesis In Vivo

[0269] One of the anti-LPA mAbs (B7) was tested for its ability to mitigate angiogenesis in vivo using the Matrigel Plug assay. This assay utilizes Matrigel, a proprietary mixture of tumor remnants including basement membranes derived from murine tumors. When Matrigel, or its derivative growth factor-reduced (GFR) Matrigel, is injected sc into an animal, it solidifies and forms a 'plug.' If pro-angiogenic factors are mixed with the matrix prior to placement, the plug will be invaded by vascular endothelial cells which eventually form blood vessels. Matrigel can be prepared either alone or mixed with recombinant growth factors (bFGF, VEGF), or tumor cells and then injected sc in the flanks of 6-week old nude (Ncr Nu/Nu) female mice. In this example, Caki-1 (renal carcinoma) cells were introduced inside the Matrigel and are producing sufficient levels of VEGF and/or IL8 and LPA. Matrigel plugs were prepared containing 5 $\times$ 10⁵ Caki-1 cells from mice treated with saline or with 10 mg/kg of anti-LPA mAb-B7, every 3 days starting 1 day prior to Matrigel implantation. Plugs were stained for endothelial CD31, followed by quantitation of the micro-vasculature formed in the plugs. Quantitation data were means +/-SEM of at least 16 fields/section from 3 plugs. The plugs from mice treated with the anti-LPA mAb B7 demonstrated a prominent reduction in blood vessel formation, as assayed by endothelial staining for CD31, compared to the plugs from saline-treated mice. Quantification of stained vessels demonstrates a greater than 50% reduction in angiogenesis in Caki-1-containing plugs from animals treated with mAb B7 compared to saline-treated animals. This was a statistically significant reduction ($p < 0.05$ for mAb B7 vs. Saline as determined by Student's T-test) in tumor cell angiogenesis as a result of anti-LPA mAb treatment.

Anti-LPA mAbs Reduces Tumor Progression in Renal and Pancreatic Xenografts

[0270] The anti-LPA antibodies have been shown (above) to be effective in reducing LPA-induced tumor cell proliferation, migration, protection from cell death and cytokine release in multiple human tumor cell lines. mAbs B58 and B7 were next tested in a xenograft model of renal and pancreatic cancer. Below are preliminary results that demonstrate the potential anti-tumorigenic effects of the anti-LPA antibody approach.

[0271] Tumors were developed by subcutaneous injection of Caki-1 and Panc-1 human tumor cells into the left flank of 4 week old female nude (NCR Nu/Nu) mice using standard protocols. After 10 days for Caki-1 and 30 days for Panc-1, when solid tumors had formed (~200 mm³), mice were randomized into treatment groups. Treatment was initiated by i.p. administration of 25 mg/kg of the anti-LPA mAbs or vehicle (saline solution). Antibodies were administered every three days for the duration of the study. Treatments consisted of 25 mg/kg of the anti-LPA mAb B58 for caki-1 tumors, mAb B7 for Panc-1 or Saline. Data are the mean $\pm$ SEM of 7 saline and 6 B58-treated mice for the caki-1 study and 4 saline and 5 B7-treated mice for the panc-1 study. Tumor volumes were measured every other day using electronic calipers and the tumor volume determined by the formula, $W^2 \times L/2$. Animals were subsequently sacrificed after tumors reached 1500 mm³ in the saline group. Final tumor volumes and weights were recorded.

[0272] In this preliminary experiment, the ability of the anti-LPA mAbs to reduce tumor volume was apparent after the tumors reached approximately 400-500 mm³. At this point, the tumors from the control animals continued to grow, while the tumors from the anti-LPA mAb-treated animals exhibited a slower rate growth in both xenograft models. Data demonstrates that the anti-LPA mAb also reduced the final tumor weights of caki-1 and panc-1 tumors when compared to tumor weights from saline-treated animals.

Anti-LPA mAbs Modulate Levels of Circulating Pro-Angiogenic Cytokines in Animals with Tumors

[0273] The anti-LPA mAbs (B58 and B7) also influenced the levels of circulating pro-angiogenic cytokine. In animals treated with the anti-LPA mAb7 (Panc-1), the serum level of interleukin-8 (IL-8) was not detectable in any antibody-treated animals, whereas IL-8 serum levels were detectable in Panc-1 and Caki-1 xenografts after 85 and 63 days, respectively. More importantly there was a strong correlation ($r=0.98$) between tumor size and IL-8 levels. In the animals bearing Caki-1 tumors the serum levels of human IL-8 were also reduced by the treatment with anti-LPA mAb58 ($r=0.34$) when compared to saline treatment ($r=0.55$). As mentioned above, the reduction of circulating cytokine levels is believed to be due to a direct inhibition of cytokine release from the tumor cells themselves. These data demonstrates the ability of the anti-LPA mAb to reduce tumor progression while also reducing the levels of circulating pro-angiogenic compounds.

[0274] Anti-LPA mAbs Reduces Tumor Progression in a Murine Model of Metastasis

[0275] One important characteristic of tumor progression is the ability of a tumor to metastasize and form secondary tumor nodules at remote sites. In vitro studies described hereinabove have demonstrated the ability of LPA to induce tumor cells to escape contact inhibition and promote migration in a scratch assay for cell motility. In these studies, the anti-LPA mAbs also inhibited LPA's tumor growth promoting effectors. The efficacy of the anti-LPA mAb to inhibit tumor metastasis in vivo. The phenomenon of tumor metastasis has been difficult to mimic in animal models. Many investigators utilize an "experimental" metastasis model in which tumor cells are directly injected into the blood stream.

[0276] Blood vessel formation is an integral process of metastasis because an increase in the number of blood vessels means cells have to travel a shorter distance to reach circulation. It is believed that anti-LPA mAb will inhibit in vivo

tumor cell metastasis, based on the finding that the anti-LPA mAb can block several integral steps in the metastatic process.

[0277] Study: The highly metastatic murine melanoma (B16-F10) was used to examine the therapeutic effect of three anti-LPA mAbs on metastasis in vivo. This model has demonstrated to be highly sensitive to cPA inhibitors of autotaxin. 4 week old female (C57BL/6) mice received an injection of B16-F10 murine melanoma tumor cells (100 μ L of 5×10^4 cells/animal) via the tail vein. Mice (10 per group) were administered 25 mg/kg of the anti-LPA mAb (either B3 or B7) or saline every three days by i.p. injection. After 18 days, lungs were harvested and analyzed. The pulmonary organs are the preferred metastatic site of the melanoma cells, and were therefore closely evaluated for metastatic nodules. The lungs were inflated with 10% buffered formalin via the trachea, in order to inflate and fix simultaneously, so that even small foci could be detectable on histological examination. Lungs were separated into five lobes and tumors were categorized by dimension (large ≥ 5 mm; medium 1-4 mm; small < 1 mm) and counted under a dissecting microscope. Upon examination of the lungs, the number of tumors was clearly reduced in antibody-treated animals. For animals treated with mAb B3, large tumors were reduced by 21%, medium tumors by 17% and small tumors by 22%. Statistical analysis by student's T-test gave a $p < 0.05$ for number of small tumors in animals treated with mAb B3 vs saline.

[0278] As shown in the above examples, it has now been shown that the tumorigenic effects of LPA are extended to renal carcinoma (e.g., Caki-1) and pancreatic carcinoma (Panc-1) cell lines. LPA induces tumor cell proliferation, migration and release of pro-angiogenic and/or pro-metastatic agents, such as VEGF and IL-8, in both cell lines. It has now been shown that three high-affinity and specific monoclonal anti-LPA antibodies demonstrate efficacy in a panel of in vitro cell assays and in vivo tumor models of angiogenesis and metastasis.

Example 11 Immunohistochemistry of Tumor Biopsy Material

[0279] The purpose of this example is to demonstrate that mAbs developed against S1P could be used to detect S1P in biopsy material. This immunohistochemical (IHC) method assesses the level of S1P in the tumor (which is believed to be produced by the tumor itself) and may be more sensitive and specific than measuring protein or RNA expression of sphingosine kinase. In addition, the IHC method would not suffer diminution of the S1P signal as S1P secreted from the tumor is diluted into the extracellular space (e.g., plasma compartment). We analyzed S1P content in U937 human tumor sections (frozen; 10 μ m thick) from a mouse Matrigel/xenograft model. U937 cells (human lymphoma cell line; ATCC cat no# CRL-1593.2) were mixed with Matrigel matrix, at a concentration of 10.5 mg/ml. 600 μ L of Matrigel mix containing U937 (30×10^6 cells/plug in a 600 μ L volume) were implanted into the right flank of 4-6 weeks nu/nu female mice and allowed to grow for 30 days. The animals were sacrificed and the Matrigel plugs were excised and embedded in OTC and flash frozen in dry ice and isopentane. Then were sectioned using a cryostat to 5 μ m sections. Sections were then fixed in 10% neutral buffered formalin, (Sigma, St. Louis Mo.; catalog number: HT 50-1-1; lot#025K4353) for 20 min at room temp and then sections. The sections were washed with 100 mM glycine (pH 7.4) in PBS for 5 min at room temp, washed

2× with PBS/0.1% Tween 20. Sections were blocked in 1% BSA/PBS/0.05% Tween for 20 min at room temp. Primary antibodies (e.g. murine anti-S1P mAb) were diluted (1:25 or at 1:50, as indicated) in 1%BSA/PBS/0.05% Tween and incubated with tumor sections for 3 hr at room temp. Sections were then washed 3× with PBS/0.1% Tween with gentle agitation. Diluted secondary antibodies (FITC-conjugated anti mouse Ab (1:250) and RRX-conjugated anti-rat Ab (1:2500 or 1:500) in 1% BSA/PBS/0.05% Tween were incubated with tumor sections for 1 hr at room temp. Sections were then washed 6× at 5 min intervals with PBS/0.05% Tween. Sections were counterstained with DAPI (4',6-diamidino-2-phenylindole dilactate (DAPI, 10 mg; Sigma, St. Louis Mo.; catalog number D3571, lot 22775) by incubation with DAPI (1:5000) diluted in PBS for 20 min at room temp. Sections were then washed 2× at 5 min intervals with PBS and 1× with DI H₂O and mounted in Gelvitol mounting media and let dry. Primary antibodies used were LT1002 (LH-2; 15 mg/ml) anti-S1P mAb diluted to 1.0 mg/ml and added at a working concentration of 1:25 in 1%BSA/PBS/0.05% Tween. Secondary antibodies used were: Fluorescein (FITC)-conjugated rabbit anti-mouse IgG (H+L) (Jackson ImmunoResearch, West Grove Pa.; catalog #315-095-003; lot number:67031) Ab diluted 1:250 in 1%BSA/PBS/0.05% Tween. Images were captured with a DeltaVision deconvolution microscope system (Applied Precision, Inc., Issaquah, Wash.) The system includes a Photometrics CCD mounted on a Nikon TE-200 inverted epi-fluorescence microscope. In general, 8-10 optical sections spaced by ~0.2 um were taken. Exposure times were set such that the camera response was in the linear range for each fluorophore. Lenses included 20× and 10×. The data sets were deconvolved and analyzed using SoftWorx software (Applied Precision, Inc) on a Silicon Graphics Octane workstation.

[0280] S1P could easily be seen in tumor biopsy images using this IHC method, using the anti-S1P mAb as the primary antibody. In contrast, S1P staining was absent in control samples from which the primary antibody was omitted.

[0281] Without being bound by theory or limited to these examples, it is believed that the measurement of the biomarker S1P could be used in conjunction with measurements of gene expression for S1P receptors and of sphingosine kinase, both of which could serve as surrogate cancer markers. Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, FEBS Lett., 2000, 480, 17 24; Celis, et al., FEBS Lett., 2000, 480, 2 16), SAGE (serial analysis of gene expression) (Madden, et al., Drug Discov. Today, 2000, 5, 415 425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, Methods Enzymol., 1999, 303, 258 72), TOGA (total gene expression analysis) (Sutcliffe, et al.,

Proc. Natl. Acad. Sci. U.S.A., 2000, 97, 1976 81), protein arrays and proteomics (Celis, et al., FEBS Lett., 2000, 480, 2 16; Jungblut, et al., Electrophoresis, 1999, 20, 2100 10), expressed sequence tag (EST) sequencing (Celis, et al., FEBS Lett., 2000, 480, 2 16; Larsson, et al., J. Biotechnol., 2000, 80, 143 57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., Anal. Biochem., 2000, 286, 91 98; Larson, et al., Cytometry, 2000, 41, 203 208), subtractive cloning, differential display (DD) (Jurecic and Belmont, Curr. Opin. Microbiol., 2000, 3, 316 21), comparative genomic hybridization (Carulli, et al., J. Cell Biochem. Suppl., 1998, 31, 286 96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, Eur. J. Cancer, 1999, 35, 1895 904) and mass spectrometry methods (To, Comb. Chem. High Throughput Screen, 2000, 3, 235 41).

[0282] All of the compositions and methods described and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit and scope of the invention as defined by the appended claims.

[0283] All patents, patent applications, and publications mentioned in the specification are indicative of the levels of those of ordinary skill in the art to which the invention pertains. All patents, patent applications, and publications, including those to which priority or another benefit is claimed, are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

[0284] The invention illustratively described herein suitably may be practiced in the absence of any element(s) not specifically disclosed herein. Thus, for example, in each instance herein any of the terms “comprising”, “consisting essentially of”, and “consisting of” may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 41

<210> SEQ ID NO 1

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 1

atggratgga gckggrtctt tmtctt 26

<210> SEQ ID NO 2
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 2

cagtggatag acagatgggg g 21

<210> SEQ ID NO 3
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 3

cagtggatag accgatgggg c 21

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 4

cagtggatag actgatgggg g 21

<210> SEQ ID NO 5
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 5

caagggatag acagatgggg c 21

<210> SEQ ID NO 6
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe Ile Asp His
20 25 30

Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Cys Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn Glu Met Phe

-continued

50	55	60	
Arg Gly Lys Ala Thr	Leu Thr Ala Asp Lys	Ser Ser Thr Thr Ala Tyr	
65	70	75	80
Ile Gln Val Asn Ser	Leu Thr Phe Glu Asp	Ser Ala Val Tyr Phe Cys	
	85	90	95
Ala Arg Gly Gly Phe Tyr	Gly Ser Thr Ile Trp Phe Asp	Phe Trp Gly	
	100	105	110
Gln Gly Thr Thr Leu Thr Val Ser			
115	120		

<210> SEQ ID NO 7
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 7
 gtctctgatt ctaggcca 18

<210> SEQ ID NO 8
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 8
 actggatggt gggaagatgg 20

<210> SEQ ID NO 9
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 9

Glu Thr Thr Val Thr	Gln Ser Pro Ala Ser	Leu Ser Met Ala Ile Gly
1	5	10 15
Glu Lys Val Thr Ile Arg Cys	Ile Thr Thr Thr Asp Ile Asp Asp Asp	
20	25	30
Met Asn Trp Phe Gln Gln Lys	Pro Gly Glu Pro Pro Asn Leu Leu Ile	
35	40	45
Ser Glu Gly Asn Ile Leu Arg	Pro Gly Val Pro Ser Arg Phe Ser Ser	
50	55	60
Ser Gly Tyr Gly Thr Asp Phe	Leu Phe Thr Ile Glu Asn Met Leu Ser	
65	70	75 80
Glu Asp Val Ala Asp Tyr Tyr Cys	Leu Gln Ser Asp Asn Leu Pro Phe	
85	90	95
Thr Phe Gly Ser Gly Thr Lys	Leu Glu Ile Lys	
100	105	

<210> SEQ ID NO 10
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

-continued

Ile Thr Thr Thr Asp Ile Asp Asp Asp Met Asn
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 11

Glu Gly Asn Ile Leu Arg Pro
1 5

<210> SEQ ID NO 12
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

Leu Gln Ser Asp Asn Leu Pro Phe Thr
1 5

<210> SEQ ID NO 13
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 13

Asp His Thr Ile His
1 5

<210> SEQ ID NO 14
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 14

Cys Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn Glu Met Phe Arg
1 5 10 15

Gly

<210> SEQ ID NO 15
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 15

Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp Phe
1 5 10

<210> SEQ ID NO 16
<211> LENGTH: 403
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 16

agcttgccgc caccatgatt gcctctgctc agttccttgg tctcctgttg ctctgttttc 60
aagggtaccag atgtgaaaca actgtgaccc agtctccagc atccctgtcc atggctatag 120
gagaaaaagt caccatcaga tgcataacca ccaactgatat tgatgatgat atgaactggt 180
tccagcagaa gccaggggaa cctcctaacc tccttatttc cgaaggcaat attcttcgtc 240

-continued

```

ctggagtgccc atccccgattc tccagcagtg gctatggtac agactttctt tttacaattg 300
aaaaacatgct ctcagaagat gttgcagatt actactgttt gcagagtgat aacttaccat 360
tcacgttcgg ctcgggggaca aagttggaaa taaaacgtga gtg 403

```

```

<210> SEQ ID NO 17
<211> LENGTH: 129
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 17

```

```

Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
1           5           10           15
Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro Ala Ser Leu Ser
20           25           30
Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Thr Thr Asp
35           40           45
Ile Asp Asp Asp Met Asn Trp Phe Gln Gln Lys Pro Gly Glu Pro Pro
50           55           60
Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro Gly Val Pro Ser
65           70           75           80
Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu Phe Thr Ile Glu
85           90           95
Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln Ser Asp
100          105          110
Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg
115          120          125

```

```

Glu

```

```

<210> SEQ ID NO 18
<211> LENGTH: 403
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (15)..(401)

```

```

<400> SEQUENCE: 18

```

```

agcttgccgc cacc atg att gcc tct gct cag ttc ctt ggt ctc ctg ttg 50
          Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu
          1           5           10
ctc tgt ttt caa ggt acc aga tgt gaa aca act gtg acc cag tct cca 98
Leu Cys Phe Gln Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro
15           20           25
gca tcc ctg tcc atg gct ata gga gaa aaa gtc acc atc aga tgc ata 146
Ala Ser Leu Ser Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile
30           35           40
acc acc act gat att gat gat gat gtg aac tgg ttc cag cag aag cca 194
Thr Thr Thr Asp Ile Asp Asp Asp Val Asn Trp Phe Gln Gln Lys Pro
45           50           55           60
ggg gaa cct cct aac ctc ctt att tcc gaa ggc aat att ctt cgt cct 242
Gly Glu Pro Pro Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro
65           70           75
gga gtc cca tcc cga ttc tcc agc agt ggc tat ggt aca gac ttt ctt 290
Gly Val Pro Ser Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu
80           85           90

```

-continued

```

ttt aca att gaa aac atg ctc tca gaa gat gtt gca gat tac tac tgt      338
Phe Thr Ile Glu Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys
95                      100                      105

```

```

ttg cag agt gat aac tta cca ttc acg ttc ggc tgc ggg aca aag ttg      386
Leu Gln Ser Asp Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu
110                      115                      120

```

```

gaa ata aaa cgt gag tg      403
Glu Ile Lys Arg Glu
125

```

```

<210> SEQ ID NO 19
<211> LENGTH: 403
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (15)..(401)

```

```

<400> SEQUENCE: 19

```

```

agcttgccgc cacc atg att gcc tct gct cag ttc ctt ggt ctc ctg ttg      50
Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu
1                      5                      10

```

```

ctc tgt ttt caa ggt acc aga tgt gaa aca act gtg acc cag tct cca      98
Leu Cys Phe Gln Gly Thr Arg Cys Glu Thr Val Thr Gln Ser Pro
15                      20                      25

```

```

gca tcc ctg tcc atg gct ata gga gaa aaa gtc acc atc aga tgc ata      146
Ala Ser Leu Ser Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile
30                      35                      40

```

```

acc acc act gat att gat gat gat ctt aac tgg ttc cag cag aag cca      194
Thr Thr Thr Asp Ile Asp Asp Asp Leu Asn Trp Phe Gln Gln Lys Pro
45                      50                      55                      60

```

```

ggg gaa cct cct aac ctc ctt att tcc gaa ggc aat att ctt cgt cct      242
Gly Glu Pro Pro Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro
65                      70                      75

```

```

gga gtc cca tcc cga ttc tcc agc agt ggc tat ggt aca gac ttt ctt      290
Gly Val Pro Ser Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu
80                      85                      90

```

```

ttt aca att gaa aac atg ctc tca gaa gat gtt gca gat tac tac tgt      338
Phe Thr Ile Glu Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys
95                      100                      105

```

```

ttg cag agt gat aac tta cca ttc acg ttc ggc tgc ggg aca aag ttg      386
Leu Gln Ser Asp Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu
110                      115                      120

```

```

gaa ata aaa cgt gag tg      403
Glu Ile Lys Arg Glu
125

```

```

<210> SEQ ID NO 20
<211> LENGTH: 403
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (15)..(401)

```

```

<400> SEQUENCE: 20

```

```

agcttgccgc cacc atg att gcc tct gct cag ttc ctt ggt ctc ctg ttg      50
Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu
1                      5                      10

```

```

ctc tgt ttt caa ggt acc aga tgt gaa aca act gtg acc cag tct cca      98
Leu Cys Phe Gln Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro

```

-continued

15	20	25	
gca tcc ctg tcc atg gct ata gga gaa aaa gtc acc atc aga tgc ata			146
Ala Ser Leu Ser Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile			
30	35	40	
acc acc act gat att gat gat gat ggt aac tgg ttc cag cag aag cca			194
Thr Thr Thr Asp Ile Asp Asp Asp Gly Asn Trp Phe Gln Gln Lys Pro			
45	50	55	60
ggg gaa cct cct aac ctc ctt att tcc gaa ggc aat att ctt cgt cct			242
Gly Glu Pro Pro Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro			
65	70	75	
gga gtc cca tcc cga ttc tcc agc agt ggc tat ggt aca gac ttt ctt			290
Gly Val Pro Ser Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu			
80	85	90	
ttt aca att gaa aac atg ctc tca gaa gat gtt gca gat tac tac tgt			338
Phe Thr Ile Glu Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys			
95	100	105	
ttg cag agt gat aac tta cca ttc acg ttc ggc tgc ggg aca aag ttg			386
Leu Gln Ser Asp Asn Met Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu			
110	115	120	
gaa ata aaa cgt gag tg			403
Glu Ile Lys Arg Glu			
125			
<210> SEQ ID NO 21			
<211> LENGTH: 403			
<212> TYPE: DNA			
<213> ORGANISM: Mus musculus			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (15)..(401)			
<400> SEQUENCE: 21			
agcttgccgc cacc	atg att gcc tct gct cag ttc ctt ggt ctc ctg ttg		50
	Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu		
	1 5 10		
ctc tgt ttt caa ggt acc aga tgt gaa aca act gtg acc cag tct cca			98
Leu Cys Phe Gln Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro			
15	20	25	
gca tcc ctg tcc atg gct ata gga gaa aaa gtc acc atc aga tgc ata			146
Ala Ser Leu Ser Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile			
30	35	40	
acc acc act gat att gat gat gat ccg aac tgg ttc cag cag aag cca			194
Thr Thr Thr Asp Ile Asp Asp Asp Pro Asn Trp Phe Gln Gln Lys Pro			
45	50	55	60
ggg gaa cct cct aac ctc ctt att tcc gaa ggc aat att ctt cgt cct			242
Gly Glu Pro Pro Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro			
65	70	75	
gga gtc cca tcc cga ttc tcc agc agt ggc tat ggt aca gac ttt ctt			290
Gly Val Pro Ser Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu			
80	85	90	
ttt aca att gaa aac atg ctc tca gaa gat gtt gca gat tac tac tgt			338
Phe Thr Ile Glu Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys			
95	100	105	
ttg cag agt gat aac tta cca ttc acg ttc ggc tgc ggg aca aag ttg			386
Leu Gln Ser Asp Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu			
110	115	120	
gaa ata aaa cgt gag tg			403
Glu Ile Lys Arg Glu			
125			

-continued

<210> SEQ ID NO 22
<211> LENGTH: 129
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 22
Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
1 5 10 15
Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro Ala Ser Leu Ser
20 25 30
Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Thr Thr Asp
35 40 45
Ile Asp Asp Asp Val Asn Trp Phe Gln Gln Lys Pro Gly Glu Pro Pro
50 55 60
Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro Gly Val Pro Ser
65 70 75 80
Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu Phe Thr Ile Glu
85 90 95
Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln Ser Asp
100 105 110
Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg
115 120 125
Glu

<210> SEQ ID NO 23
<211> LENGTH: 129
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 23
Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
1 5 10 15
Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro Ala Ser Leu Ser
20 25 30
Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Thr Thr Asp
35 40 45
Ile Asp Asp Asp Leu Asn Trp Phe Gln Gln Lys Pro Gly Glu Pro Pro
50 55 60
Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro Gly Val Pro Ser
65 70 75 80
Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu Phe Thr Ile Glu
85 90 95
Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln Ser Asp
100 105 110
Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg
115 120 125
Glu

<210> SEQ ID NO 24
<211> LENGTH: 129
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 24

-continued

Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 1 5 10 15
 Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro Ala Ser Leu Ser
 20 25 30
 Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Thr Asp
 35 40 45
 Ile Asp Asp Asp Gly Asn Trp Phe Gln Gln Lys Pro Gly Glu Pro Pro
 50 55 60
 Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro Gly Val Pro Ser
 65 70 75 80
 Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu Phe Thr Ile Glu
 85 90 95
 Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln Ser Asp
 100 105 110
 Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg
 115 120 125
 Glu

<210> SEQ ID NO 25
 <211> LENGTH: 129
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 25

Met Ile Ala Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 1 5 10 15
 Gly Thr Arg Cys Glu Thr Thr Val Thr Gln Ser Pro Ala Ser Leu Ser
 20 25 30
 Met Ala Ile Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Thr Thr Asp
 35 40 45
 Ile Asp Asp Asp Pro Asn Trp Phe Gln Gln Lys Pro Gly Glu Pro Pro
 50 55 60
 Asn Leu Leu Ile Ser Glu Gly Asn Ile Leu Arg Pro Gly Val Pro Ser
 65 70 75 80
 Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Leu Phe Thr Ile Glu
 85 90 95
 Asn Met Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln Ser Asp
 100 105 110
 Asn Leu Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg
 115 120 125
 Glu

<210> SEQ ID NO 26
 <211> LENGTH: 436
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 26

atggcatgga gctgggtctt tctctcttc ctgtcagtaa ctaccggcgt ccaactcccag 60
 gctcacctgc aacagtctga cgctgaattg gtgaaacctg gagcttcagt gaagatatcc 120
 tgcaagggtt ctggcttcatt tttcattgac catactattc actggatgaa gcagaggcct 180
 gaacagggcc tcgaatggat cggatgtatt tctcccagac atgatattac taaatacaat 240

-continued

```

gagatgttca ggggcaaggc caccctgact gcagacaagt cctccactac agcctacata 300
caagtcaaca gtctgacatt tgaagactct gcagtctatt tctgtgcaag agggggggttc 360
tacggtagta ctatctgggtt tgacttttgg ggccaaggca ccactctcac agtctcctca 420
gcctccacca agggcc 436

```

```

<210> SEQ ID NO 27
<211> LENGTH: 145
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 27

```

```

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1           5           10          15
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20          25          30
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35          40          45
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50          55          60
Glu Trp Ile Gly Cys Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
65          70          75          80
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
85          90          95
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
100         105        110
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
115        120        125
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
130        135        140
Gly
145

```

```

<210> SEQ ID NO 28
<211> LENGTH: 436
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(435)

```

```

<400> SEQUENCE: 28

```

```

atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc 48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1           5           10          15
gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa 96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20          25          30
cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc 144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35          40          45
att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc 192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50          55          60
gaa tgg atc gga gct att tct ccc aga cat gat att act aaa tac aat 240
Glu Trp Ile Gly Ala Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn

```

-continued

65	70	75	80	
gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act				288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr				
	85	90	95	
aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc				336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val				
	100	105	110	
tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac				384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp				
	115	120	125	
ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag				432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys				
	130	135	140	
ggc c				436
Gly				
145				
<210> SEQ ID NO 29				
<211> LENGTH: 436				
<212> TYPE: DNA				
<213> ORGANISM: Mus musculus				
<220> FEATURE:				
<221> NAME/KEY: CDS				
<222> LOCATION: (1)..(435)				
<400> SEQUENCE: 29				
atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc				48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly				
1	5	10	15	
gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa				96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys				
	20	25	30	
cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc				144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe				
	35	40	45	
att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc				192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu				
	50	55	60	
gaa tgg atc gga tct att tct ccc aga cat gat att act aaa tac aat				240
Glu Trp Ile Gly Ser Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn				
65	70	75	80	
gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act				288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr				
	85	90	95	
aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc				336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val				
	100	105	110	
tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac				384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp				
	115	120	125	
ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag				432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys				
	130	135	140	
ggc c				436
Gly				
145				
<210> SEQ ID NO 30				
<211> LENGTH: 436				

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(435)

<400> SEQUENCE: 30

atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc      48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1          5          10          15

gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa      96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20         25         30

cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc     144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35         40         45

att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc     192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50         55         60

gaa tgg atc gga tgg att tct ccc aga cat gat att act aaa tac aat     240
Glu Trp Ile Gly Trp Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
65         70         75         80

gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act     288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
85         90         95

aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc     336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
100        105        110

tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac     384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
115        120        125

ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag     432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
130        135        140

ggc c
Gly
145

<210> SEQ ID NO 31
<211> LENGTH: 436
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(435)

<400> SEQUENCE: 31

atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc      48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1          5          10          15

gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa      96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20         25         30

cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc     144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35         40         45

att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc     192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50         55         60

gaa tgg atc gga tat att tct ccc aga cat gat att act aaa tac aat     240

```

-continued

Glu Trp Ile Gly Tyr Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn	
65 70 75 80	
gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act	288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr	
85 90 95	
aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc	336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val	
100 105 110	
tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac	384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp	
115 120 125	
ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag	432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys	
130 135 140	
ggc c	436
Gly	
145	
 <210> SEQ ID NO 32	
<211> LENGTH: 436	
<212> TYPE: DNA	
<213> ORGANISM: Mus musculus	
<220> FEATURE:	
<221> NAME/KEY: CDS	
<222> LOCATION: (1)..(435)	
 <400> SEQUENCE: 32	
atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc	48
Met Ala Trp Ser Trp Val Phe Leu Phe Leu Ser Val Thr Thr Gly	
1 5 10 15	
gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa	96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys	
20 25 30	
cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc	144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe	
35 40 45	
att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc	192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu	
50 55 60	
gaa tgg atc gga cgt att tct ccc aga cat gat att act aaa tac aat	240
Glu Trp Ile Gly Arg Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn	
65 70 75 80	
gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act	288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr	
85 90 95	
aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc	336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val	
100 105 110	
tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac	384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp	
115 120 125	
ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag	432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys	
130 135 140	
ggc c	436
Gly	
145	
 <210> SEQ ID NO 33	

-continued

```

<211> LENGTH: 436
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(435)

<400> SEQUENCE: 33

atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc      48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1           5           10          15

gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa      96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
          20          25          30

cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc     144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
          35          40          45

att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc     192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
          50          55          60

gaa tgg atc gga ttt att tct ccc aga cat gat att act aaa tac aat     240
Glu Trp Ile Gly Phe Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
        65          70          75          80

gag atg ttc agg ggc aag gcc acc ctg act gca gac aag tcc tcc act     288
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
          85          90          95

aca gcc tac ata caa gtc aac agt ctg aca ttt gaa gac tct gca gtc     336
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
          100         105         110

tat ttc tgt gca aga ggg ggg ttc tac ggt agt act atc tgg ttt gac     384
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
          115         120         125

ttt tgg ggc caa ggc acc act ctc aca gtc tcc tca gcc tcc acc aag     432
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
          130         135         140

ggc c                                                                436
Gly
145

<210> SEQ ID NO 34
<211> LENGTH: 436
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(435)

<400> SEQUENCE: 34

atg gca tgg agc tgg gtc ttt ctc ttc ttc ctg tca gta act acc ggc      48
Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1           5           10          15

gtc cac tcc cag gct cac ctg caa cag tct gac gct gaa ttg gtg aaa      96
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
          20          25          30

cct gga gct tca gtg aag ata tcc tgc aag gtt tct ggc ttc att ttc     144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
          35          40          45

att gac cat act att cac tgg atg aag cag agg cct gaa cag ggc ctc     192
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
          50          55          60

```

<400> SEQUENCE: 35

<400> SEQUENCE: 36

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1 5 10 15
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20 25 30

-continued

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35 40 45
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50 55 60
Glu Trp Ile Gly Ser Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
65 70 75 80
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
85 90 95
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
100 105 110
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
115 120 125
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
130 135 140

Gly
145

<210> SEQ ID NO 37
<211> LENGTH: 145
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 37

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1 5 10 15
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20 25 30
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
35 40 45
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
50 55 60
Glu Trp Ile Gly Trp Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
65 70 75 80
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
85 90 95
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
100 105 110
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
115 120 125
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
130 135 140

Gly
145

<210> SEQ ID NO 38
<211> LENGTH: 145
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 38

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
1 5 10 15
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
20 25 30

-continued

```

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
   35               40               45

Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
   50               55               60

Glu Trp Ile Gly Tyr Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
   65               70               75               80

Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
           85               90               95

Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
   100               105               110

Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
   115               120               125

Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
   130               135               140

```

Gly
145

<210> SEQ ID NO 39
 <211> LENGTH: 145
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 39

```

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
  1               5               10               15

Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
           20               25               30

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
   35               40               45

Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
   50               55               60

Glu Trp Ile Gly Arg Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
   65               70               75               80

Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
           85               90               95

Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
   100               105               110

Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
   115               120               125

Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
   130               135               140

```

Gly
145

<210> SEQ ID NO 40
 <211> LENGTH: 145
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 40

```

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
  1               5               10               15

Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
           20               25               30

```

-continued

```

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
   35           40           45
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
   50           55           60
Glu Trp Ile Gly Phe Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
   65           70           75           80
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
           85           90           95
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
          100          105          110
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
          115          120          125
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
          130          135          140
Gly
145

```

```

<210> SEQ ID NO 41
<211> LENGTH: 145
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 41

```

```

Met Ala Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
 1           5           10           15
Val His Ser Gln Ala His Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
          20           25           30
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Val Ser Gly Phe Ile Phe
   35           40           45
Ile Asp His Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu
   50           55           60
Glu Trp Ile Gly Met Ile Ser Pro Arg His Asp Ile Thr Lys Tyr Asn
   65           70           75           80
Glu Met Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr
           85           90           95
Thr Ala Tyr Ile Gln Val Asn Ser Leu Thr Phe Glu Asp Ser Ala Val
          100          105          110
Tyr Phe Cys Ala Arg Gly Gly Phe Tyr Gly Ser Thr Ile Trp Phe Asp
          115          120          125
Phe Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
          130          135          140
Gly
145

```

We claim:

1. An isolated immune-derived moiety reactive against a bioactive lipid, wherein the moiety is derived from an antibody produced by immunizing an immune competent mammalian host with an immunogen comprising a derivatized bioactive lipid covalently linked to a carrier, wherein the derivatized bioactive lipid comprises a polar head group and at least one hydrocarbon chain, wherein the derivatized bioactive lipid is selected from the group consisting of:

(a) a derivatized lipid wherein a carbon atom within the hydrocarbon chain of the bioactive lipid is derivatized

with an optionally protected pendant sulfhydryl group, wherein the optionally protected pendant sulfhydryl group is optionally appended at a position selected from the group consisting of an SN-1 position and an SN-2 position of the glycerol backbone of the bioactive lipid; and

(b) a derivatized lipid wherein the bioactive lipid is a lyso-lipid and a carbon atom within the hydrocarbon chain is derivatized with an optionally protected pendant reactive group.

2. An isolated immune-derived moiety according to claim 1 is selected from the group consisting of an antibody, an antibody fragment, an antibody variant, and an antibody derivative.

3. An isolated immune-derived moiety according to claim 1, wherein the bioactive lipid is selected from the group consisting of a lysophosphatidic acid, a lysophosphatidic acid precursor, and a lysophosphatidic acid metabolite.

4. An isolated immune-derived moiety according to claim 1 that is an antibody selected from the group consisting of a chimeric monoclonal antibody, a human monoclonal antibody, and a humanized monoclonal antibody.

5. A composition comprising an isolated immune-derived moiety according to claim 1 in a carrier, optionally a pharmaceutically acceptable carrier.

6. A composition according to claim 1, wherein the immune-derived moiety is bispecific.

7. A composition according to claim 6, wherein the immune-derived moiety is reactive against both a lysophosphatidic acid and a sphingolipid.

8. A method of decreasing the effective concentration of a bioactive lipid in a subject, comprising administering to the subject an immune-derived moiety according to claim 1 that is reactive against a bioactive lipid in an amount effective to decrease the effective concentration of said bioactive lipid, wherein the bioactive lipid is a non-sphingoid bioactive lipid.

9. A method according to claim 16, wherein the subject is a mammal, optionally a human.

10. A method of treatment, comprising administering to a subject in need of therapeutic or prophylactic treatment an amount of an immune-derived moiety according to claim 1 effective to accomplish such treatment.

11. A method according to claim 10, wherein the immune-derived moiety is a monoclonal antibody reactive against a lysolipid, optionally selected from the group consisting of a lysophosphatidic acid, a lysophosphatidic acid precursor, and a lysophosphatidic acid metabolite.

12. A method according to claim 10, wherein the subject is a mammal, optionally a human.

13. A method selected from the group consisting of:

(a) a method of inhibiting proliferation of a cancer cell, comprising administering to a subject having a cancer responsive to lysophosphatidic acid signaling an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to inhibit proliferation of the cancer cell, wherein the cancer cell is optionally selected from the group consisting of a renal carcinoma cell, a pancreatic carcinoma cell, a melanoma cell, a lung carcinoma cell, a neuroblastoma cell, a hepatocellular carcinoma cell, a glioblastoma multiforme cell, a breast cancer cell, an ovarian cancer cell, a prostate cancer cell, a colorectal cancer cell, and a leukemia cell;

(b) a method of inhibiting migration of a cancer cell, comprising administering to a subject having a cancer responsive to lysophosphatidic acid signaling an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to inhibit migration of the cancer cell, wherein the cancer cell is optionally selected from the group consisting of a renal carcinoma cell, a pancreatic carcinoma cell, a melanoma cell, a lung carcinoma cell, a neuroblastoma cell, a hepatocellular carcinoma cell, a glioblastoma

multiforme cell, a breast cancer cell, an ovarian cancer cell, a prostate cancer cell, a colorectal cancer cell, and a leukemia cell;

(c) a method of inhibiting tumor metastasis, comprising administering to a subject having a potentially metastatic cancer an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to inhibit metastasis of the tumor, wherein the tumor is optionally selected from the group consisting of renal carcinoma, pancreatic carcinoma, melanoma, lung carcinoma, neuroblastoma, hepatocellular carcinoma, glioblastoma multiforme, breast cancer, ovarian cancer, prostate cancer, colorectal cancer, and leukemia;

(d) a method of inhibiting angiogenesis in a tumor, comprising administering to a subject having a tumor an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to inhibit angiogenesis in the tumor, wherein the tumor is optionally selected from the group consisting of renal carcinoma, pancreatic carcinoma, melanoma, lung carcinoma, neuroblastoma, hepatocellular carcinoma, glioblastoma multiforme, breast cancer, ovarian cancer, prostate cancer, colorectal cancer, and leukemia;

(e) a method of increasing apoptosis of a cancer cell, comprising contacting the cancer cell with an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to increase apoptosis of the cancer cell, wherein the cancer cell is optionally selected from the group consisting of a renal carcinoma cell, a pancreatic carcinoma cell, a melanoma cell, a lung carcinoma cell, a neuroblastoma cell, a hepatocellular carcinoma cell, a glioblastoma multiforme cell, a breast cancer cell, an ovarian cancer cell, a prostate cancer cell, a colorectal cancer cell, and a leukemia cell; and

(f) a method of enhancing the apoptotic effect of a cytotoxic agent against a cancer cell, comprising contacting the cancer cell with an amount of an immune-derived moiety according to claim 3 that is reactive with a lysophosphatidic acid effective to enhance the apoptotic effect of a cytotoxic agent against the cancer cell, wherein the cancer cell is optionally selected from the group consisting of a renal carcinoma cell, a pancreatic carcinoma cell, a melanoma cell, a lung carcinoma cell, a neuroblastoma cell, a hepatocellular carcinoma cell, a glioblastoma multiforme cell, a breast cancer cell, an ovarian cancer cell, a prostate cancer cell, a colorectal cancer cell, and a leukemia cell.

14. A method of treating cancer, comprising administering to an animal having or suspected of having cancer a therapeutically effective amount of an isolated immune-derived moiety reactive against a lysophosphatidic acid to decrease the effective concentration of lysophosphatidic acid in the animal, thereby effecting treatment, wherein the cancer is optionally selected from the group consisting of renal carcinoma, pancreatic carcinoma, melanoma, lung carcinoma, neuroblastoma, hepatocellular carcinoma, glioblastoma multiforme, breast cancer, ovarian cancer, prostate cancer, colorectal cancer, and leukemia.

15. A method according to claim 15 further comprising administering a cytotoxic agent.

16. A method of administration, comprising administering an immune-derived moiety according to claim **1** to a subject in need of treatment with the immune-derived moiety.

17. A method according to claim **16**, wherein the immune-derived moiety is a monoclonal antibody reactive against a lysolipid, optionally a lysophosphatidic acid, and wherein the monoclonal antibody is optionally selected from the group consisting of a chimeric monoclonal antibody, a human monoclonal antibody, and a humanized monoclonal antibody.

18. A method according to claim **16**, wherein the immune-derived moiety is selected from the group consisting of an antibody, an antibody fragment, an antibody variant, and an antibody derivative.

19. A method according to claim **16**, wherein the subject is a mammal, optionally a human, and wherein administration is via a route selected from the group consisting of topical administration (optionally via a route selected from the group consisting of a transdermal route, an epidermal ophthalmic route, an intrauterine route, a vaginal route, a rectal route, a pulmonary route, an intratracheal route, and an intranasal route), oral administration, and parenteral administration (optionally via a route selected from the group consisting of intravenous, intraarterial, subcutaneous, intraperitoneal, intramuscular, and intracranial administration).

* * * * *