(12) STANDARD PATENT (11) Application No. **AU 2011301901 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
Ester pro-drugs of [3-(1-(1H-imidazol-4-yl)ethyl)-2-methylphenyl] methanol for treating retinal diseases

(51) International Patent Classification(s)
A61K 31/4174 (2006.01) ***A61P 27/06*** (2006.01)

(21) Application No: **2011301901** (22) Date of Filing: **2011.09.16**

(87) WIPO No: **WO12/037453**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/383,370	**2010.09.16**	**US**

(43) Publication Date: **2012.03.22**
(44) Accepted Journal Date: **2016.02.11**

(71) Applicant(s)
Allergan, Inc.

(72) Inventor(s)
Dibas, Mohammed I.;Gil, Daniel W.;Chow, Ken;Wang, Liming;Garst, Michael E.;Donello, John E.

(74) Agent / Attorney
Davies Collison Cave, Level 14 255 Elizabeth Street, Sydney, NSW, 2000

(56) Related Art
WHEELER et al. European Journal of Ophthalmology (1999), vol. 9, Suppl. 1, pages S17-S21
WO 2005/034998 A2 (ALLERGAN, INC.) 21 April 2005
MERIN et al. Journal of Ocular Pharmacology and Therapeutics (2008), vol. 24, no. 1, pages 80-86
WO 2009/089132 A1 (ALLERGAN, INC.) 16 July 2009
WO 2010/093930 A1 (ALLERGAN, INC.) 19 August 2010
WO 2006/036480 A1 (ALLERGAN, INC.) 06 April 2006
WO 2010/091209 A1 (ALLERGAN, INC.) 12 August 2010

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
22 March 2012 (22.03.2012)

PCT

(10) International Publication Number
WO 2012/037453 A1

(51) International Patent Classification:
A61K 31/4174 (2006.01) *A61P 27/06* (2006.01)

(21) International Application Number:
PCT/US2011/051926

(22) International Filing Date:
16 September 2011 (16.09.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/383,370 16 September 2010 (16.09.2010) US

(71) Applicant *(for all designated States except US)*: **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, California 92886 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **DIBAS, Mohammed I.** [US/US]; 28871 Niguel Vista, Laguna Niguel, California 92677 (US). **GIL, Daniel W.** [US/US]; 2541 Point Del Mar, Corona Del Mar, California 92625 (US). **CHOW, Ken** [US/US]; 20 Tidal Surf, Newport Coast, California 92657 (US). **WANG, Liming** [US/US]; 16 Trinity, Irvine, California 92612 (US). **GARST, Michael E.** [US/US]; 2627 Raqueta Drive, Newport Beach, California 92660 (US). **DONELLO, John E.** [US/US]; 34041 Pequito Drive, Dana Point, California 92629 (US).

(74) Agents: **ENE, Doina** et al.; Allergan, Inc., 2525 Dupont Drive, Irvine, California 92612 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*

WO 2012/037453 A1

(54) Title: ESTER PRO-DRUGS OF [3-(1-(1H-IMIDAZOL-4-YL)ETHYL)-2-METHYLPHENYL] METHANOL FOR TREATING RETINAL DISEASES

(57) Abstract: The present invention relates to method of treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising a ester pro-drugs of [3-(1-(1H-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals.

ESTER PRO-DRUGS OF [3-(1-(1H-IMIDAZOL-4-YL)ETHYL)-2-METHYLPHENYL] METHANOL FOR TREATING RETINAL DISEASES

By: Mohammed I. Dibas, Daniel W. Gil, Ken Chow, Liming Wang, Michael E. Garst and John E. Donello

RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application No. 61/383,370 filed on September 16, 2010, the entire disclosure of which is incorporated herein by this reference.

BACKGROUND OF THE INVENTION

1. Field of the invention

The present invention relates to a method for treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising ester pro-drugs of [3-(1-(1H-imidazol-4-yl)ethyl)-2-methylphenyl] methanol or of its enantiomers.

2. Summary of the related art


Three alpha-1 and three alpha-2 adrenergic receptors have been characterized by molecular and pharmacological methods. Activation of these alpha receptors evokes physiological responses with useful therapeutic applications.

Compound, 4-[1-(2,3-dimethylphenyl)ethyl]-3*H*-imidazole, generically known as, medetomidine is an alpha 2 adrenergic agonist, for use in the sedation of animals. The hydrochloride salt of the (*S*) enantiomer of medetomidine, generically known as dexmedetomidine, (*S*) 4-[1-(2,3-dimethylphenyl)ethyl]-3*H*-imidazole, is also indicated for use as a sedative or analgesic in cats and dogs.

The metabolite of dexmedetomidine is (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol together with its racemic mixture, compound [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, are described in the literature in *Journal of Chromatography*, (1997), 762(1 + 2), 281-291 by Hui, Y.-H et al.

[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl]methanol is described in "Synthesis of detomidine and medetomidine metabolites: 1,2,3-trisubstituted arenes with 4′(5′)-imidazolylmethyl groups" in *Journal of Heterocyclic Chemistry* (1993), 30(6), (1645-1651) by Stoilov et al.


Kavanagh, et al. describe [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl]methanol in "Synthesis of Possible Metabolites of Medetomidine {1-(2,3-dimethylphenyl)-1-[imidazol-4(5)-yl]ethane" in *Journal of Chemical Research, Synopses* (1993), (4), 152-3.



Medetomidine

4-(1-(2,3-dimethylphenyl) ethyl)-1*H*-imidazole


CAS 86347-14-0



Dexmedetomidine


(*S*)-4-(1-(2,3-dimethylphenyl) ethyl)-1*H*-imidazole

CAS 189255-79-6




(3-(1-(1*H*-imidazol-4-yl)ethyl) -2-methylphenyl)methanol

CAS 128366-50-7



(*R*)-(3-(1-(1*H*-imidazol-4-yl)ethyl) -2-methylphenyl)methanol

CAS 1240244-32-9



(*S*)-(3-(1-(1*H*-imidazol-4-yl)ethyl) -2-methylphenyl)methanol

CAS 189255-79-6

[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl)methanol] is described by Salonen, et al. in "Biotransformation of Medetomidine in the Rat" in *Xenobiotica* (1990), 20(5), 471-80.

PCT Int. Appl. WO 2010093930 A1 discloses [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl]methanol and its (*S*) and (*R*) enantiomers.


2011301901 15 Jan 2016

SUMMARY OF THE INVENTION

Three alpha 1 and three alpha 2 adrenergic receptors have been characterized by molecular and pharmacological methods. Activation of these alpha 2 receptors evokes physiological responses and has useful therapeutic actions.

The present invention relates to a method of treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol. Upon hydrolytic and/or enzymatic cleavage of the ester functionality the parent compound, [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, is released to act as a selective modulator of the alpha 2 adrenergic receptors.

According to a first aspect the present invention provides a method of treating a retinal disease comprising administering a therapeutically effective amount of a composition comprising a compound of Formula I, its individual enantiomers, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula I

wherein

R^1 is H or C_{1-3} alkyl;


R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or

bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl) or -S(C_{1-6} alkyl) groups, to a subject in need thereof.

According to a second aspect the present invention provides a use of a compound of Formula I, its individual enantiomers, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula I

wherein

R^1 is H or C_{1-3} alkyl;


R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl) or -S(C_{1-6} alkyl) groups,

in the manufacture of a medicament for the treatment of a retinal disease.

According to a third aspect the present invention provides a pharmaceutical composition when used for treating a retinal disease in a subject, the composition comprising a compound of Formula II



Formula II

wherein

R^1 is H or C_{1-3} alkyl;

R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl) or -S(C_{1-6} alkyl) groups,

2011301901 15 Jan 2016

In another aspect, the present invention relates to a method treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising ester pro-drugs of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or pharmaceutical compositions containing them. Upon hydrolytic and/or enzymatic cleavage of the ester functionality the parent compound, active metabolite, (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, is released to act as a selective modulator of the alpha 2 adrenergic receptors.

In another aspect the present invention provides relates to a method treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising ester pro-drugs of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or pharmaceutical compositions containing them. Upon hydrolytic and/or enzymatic cleavage of the ester functionality the parent compound (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, is released to act as a selective modulator of the alpha 2 adrenergic receptors.

The ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol are useful for the treatment or prevention of mammals, including humans, in a range of conditions and diseases that are alleviated by alpha 2A, 2B, 2C activation, including but not limited to treating or preventing glaucoma, elevated intraocular pressure, ischemic neuropathies, optic neuropathy, pain, visceral pain, corneal pain, headache pain, migraine, cancer pain, back pain, irritable bowl syndrome pain, muscle pain and pain associated with diabetic neuropathy, the treatment of diabetic retinopathy, other retinal degenerative conditions, stroke, cognitive deficits, neuropsychiatric conditions, drug dependence and addiction, withdrawal of symptoms, obsessive-compulsive disorders, obesity, insulin resistance, stress-related conditions, diarrhea, diuresis, nasal congestion, spasticity, attention deficit disorder, psychoses, anxiety, depression, autoimmune disease, Crohn's disease, gastritis, Alzheimer's, and Parkinson's ALS other neurodegenerative diseases, dermatological conditions, skin erythema (redness) and inflammation, acne, age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic macular edema, tumors, wound healing, inflammation and retinal vein occlusion, enhancing vision in patients with vision loss from conditions including glaucoma, retinitis pigmentosa and neuritis secondary to multiple sclerosis, rosacea (dilation of the blood vessels just under the skin), sunburn, chronic sun damage, discreet erythemas, psoriasis, acne rosacea, menopause-associated hot flashes, hot flashes resulting from orchiectomyatopic dermatitis, photoaging, seborrheic dermatitis, allergic dermatitis, redness of the skin, telangiectasia (dilations of previously existing small blood vessels) of the face, rhinophymia (hypertrophy of the nose with follicular dilation), red bulbous nose, acne-like skin eruptions (may ooze or crust), burning or stinging sensation of the face, irritated and bloodshot and watery eyes, erythema of the skin, cutaneous hyperactivity with dilation of blood vessels of the skin, Lyell's syndrome, Stevens-Johnson syndrome, erythema multiforme minor, erythema multiforme major and other inflammatory skin diseases.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method of treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective

amount of a composition comprising ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol and of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol as alpha-2 agonists with therapeutic utility.

The term "subject", as used herein, refers to a human patient.

In a preferred embodiment the present invention relates to a method of treating retinal diseases in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising esters pro-drugs of (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol as alpha-2 agonists with therapeutic utility. Upon hydrolytic or enzymatic cleavage of the ester functionality the parent compound, active metabolite, (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, is released to act as a selective modulator of the alpha 2 adrenergic receptors.

In one aspect of the invention, there is provided a method for treating retinal diseases in a patient in need thereof which comprises, consists essentially of or consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases in a patient in need thereof which comprises, consists essentially of or consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol , or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases in a patient in need thereof which comprises, consists essentially of or

consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases including but not limited to: age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion. Our compound of interest is also useful for enhancing vision in patients with vision loss from conditions including ocular hypertension, glaucoma, retinitis pigmentosa, nyctalopia, and neuritis secondary to multiple sclerosis.

In another aspect of the invention, there is provided a method for treating retinal diseases including but not limited to: age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion. Our compound of interest is also useful for enhancing vision in patients with vision loss from conditions including ocular hypertension, glaucoma, retinitis pigmentosa, nyctalopia, and neuritis secondary to multiple sclerosis, which comprises, consists essentially of or consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases including but not limited to: including but not limited to age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion. Our compound of interest is also useful for enhancing vision in patients with vision loss from conditions including ocular hypertension, glaucoma, retinitis pigmentosa, nyctalopia, and neuritis secondary to multiple sclerosis, which comprises, consists

essentially of or consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases including but not limited to: including but not limited to age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion. Our compound of interest is also useful for enhancing vision in patients with vision loss from conditions including ocular hypertension, glaucoma, retinitis pigmentosa, nyctalopia, and neuritis secondary to multiple sclerosis, which comprises, consists essentially of or consists of or consists of administering a therapeutically effective amount of a pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, or the tautomers thereof, or pharmaceutically acceptable salts thereof.

In another aspect of the invention, there is provided a method for treating retinal diseases wherein the pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, is selected from topical ocular application, direct injection, applications and formulations that may further enhance the long duration of actions such as a slow releasing pellet, suspension, gel, solution, cream, ointment, foams, emulsions, microemulsions, serums, aerosols, sprays, dispersions, microcapsules, vesicles, microparticles.

In another aspect of the invention, there is provided a method for treating retinal diseases wherein the pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol is selected from topical ocular application, direct injection, applications and formulations that may further enhance


the long duration of actions such as a slow releasing pellet, suspension, gel, solution, cream, ointment, foams, emulsions, microemulsions, serums, aerosols, sprays, dispersions, microcapsules, vesicles, microparticles.

In another aspect of the invention, there is provided a method for treating retinal diseases wherein the pharmaceutical composition comprising, consisting essentially of or consisting of a therapeutically effective amount of ester pro-drugs of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol is selected from topical skin application, direct injection, applications and formulations that may further enhance the long duration of actions such as a slow releasing pellet, suspension, gel, solution, lotion, cream, ointment, foams, emulsions, microemulsions, serums, aerosols, sprays, dispersions, microcapsules, vesicles, microparticles.

"Prodrugs " are frequently referred to by the term " metabolically cleavable derivatives " which refers to compound forms which are rapidly transformed in vivo to the parent compound according to the invention, for example, by hydrolysis in blood. Thus, prodrugs are compounds bearing groups which are removed by biotransformation prior to exhibiting their pharmacological action. *S*uch groups include moieties which are readily cleaved in vivo from the compound bearing it, which compound after cleavage remains or becomes pharmacologically active. *S*uch metabolically cleavable groups form a class well known to practitioners of the art. They include, but are not limited to such groups as alkanoyl (i.e. acetyl, propionyl, butyryl, and the like), unsubstituted and substituted carbocyclic aroyl (such as benzoyl, substituted benzoyl and 1- and 2-naphthoyl), alkoxycarbonyl (such as ethoxycarbonyl), trialklysilyl (such as trimethyl- and triethylsilyl), monoesters formed with dicarboxylic acids (such as succinyl), phosphate, sulfate, sulfonate, sulfonyl, sulfinyl and the like. The compounds bearing the metabolically cleavable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the metabolically cleavable group. (T. Higuchi and V. *S*tella, "Pro-drugs as Novel Delivery *S*ystem", Vol. 14 of the A.C.*S*. *S*ymposium *S*eries; "Bioreversible Carriers in Drug Design", ed.

Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987).

In one aspect, the invention therefore relates to a method of lowering intraocular pressure in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising a compound having Formula I, its individual enantiomers, its individual diastereoisomers, its individual hydrates, its individual solvates, its individual crystal forms, its individual isomers, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula I

wherein


R^1 is H or C_{1-3} alkyl;

R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, heterocycle or aryl; and

R is C_{1-10} alkyl, heterocycle or aryl.

In a preferred aspect, the invention therefore relates to a method of lowering intraocular pressure in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising a compound having Formula II, its individual diastereoisomers, its individual hydrates, its individual solvates, its individual crystal forms, its individual isomers, its individual tautomers or a pharmaceutically acceptable salt thereof,




Formula II

wherein

R^1 is H or C $_{1-3}$ alkyl;
R^2 is H or C $_{1-3}$ alkyl;
R^3 is H, C $_{1-10}$ alkyl, heterocycle or aryl; and
R is C $_{1-10}$ alkyl, heterocycle or aryl.

In another aspect, the invention therefore relates to a method of lowering intraocular pressure in a subject in need of such treatment, which comprises administering a therapeutically effective amount of a composition comprising a compound having Formula III, its individual diastereoisomers, its individual hydrates, its individual solvates, its individual crystal forms, its individual isomers, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula III

wherein
R^1 is H or C $_{1-3}$ alkyl;
R^2 is H or C $_{1-3}$ alkyl;
R^3 is H, C $_{1-10}$ alkyl, heterocycle or aryl; and
R is C $_{1-10}$ alkyl, heterocycle or aryl.

The following paragraphs provide definitions of the various chemical moieties that make up the compounds of the invention and are intended to apply uniformly throughout the specification and claims unless expressly stated otherwise.

The term "alkyl" as used herein, is defined as including a saturated monovalent alkane moiety having straight or branched alkane moieties or combinations thereof and containing 1-10 carbon atoms, preferably 1-8 carbon atoms and more preferably 1-4 carbon atoms. Alkyl moieties can optionally be substituted by, but not limited to, amino groups, aryl groups, halogens. One methylene (-CH_2-) can be replaced by carbonyl, -NH-, carboxyl, amide, sulfur or by oxygen. Examples include, but are not limited to, methyl, ethyl, propyl, butyl, *sec*-butyl, pentyl, *iso*-pentyl, *neo*-pentyl, hexyl,

iso-hexyl, 3-methyl-butyl, 2-amino-N-isobutyl acetamide, *iso*-butyl, *tert*-butyl, *iso*-propyl, ethylphenyl, methylphenyl, 2-amino-3-methyl-butanamide-N-2-methyl-1-propyl, 1-amino-2-methyl-prop-1-yl.

The term "heterocycle" as used herein is defined as an aromatic or non aromatic 5 to 10 membered monocyclic or bicyclic ring containing at least one heteroatom selected from O or N or S or combinations thereof, interrupting the carbocyclic ring structure. Heterocycles can optionally be substituted by, but not limited to, C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups. Examples include, but are not limited to, furyl, pyrryl, pyridyl, pyrimidyl, thienyl, isothiazolyl, imidazolyl, pyrazinyl, benzofuranyl, quinolyl, isoquinolyl, benzothienyl, isobenzofuryl, pyrazolyl, indolyl, isoindolyl, benzimidazolyl, purinyl, carbazolyl, oxazolyl, thiazolyl, isothiazolyl, 1,2,5-thiadiazolyl, 1,2,4-thiadiazolyl, isooxazolyl, quinazolinyl, pyridazinyl, cinnolinyl, phthalazinyl, quinoxalinyl, xanthinyl, hypoxanthinyl, pteridinyl, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, pyrrolidinyl, piperidinyl and piperazinyl.

The term "aryl" as used herein, is defined as including an organic moiety derived from an aromatic hydrocarbon consisting of a monocyclic or bicyclic ring containing 6-10 carbon atoms by removal of one hydrogen atom, such as phenyl or naphtyl. Aryl groups can optionally be substituted by, but not limited to, C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups. Examples include, but are not limited to, phenyl, naphtyl.

The term "H" as used herein refers to a hydrogen atom.

The term "O" as used herein refers to an oxygen atom.

The term "S" as used herein refers to a sulfur atom.

The term "N" as used herein refers to a nitrogen atom.

The term "amino" as used herein refers to a group of formula $-NH_2$.

The term "amide" as used herein refers to a group of formula –C(O)NH- or –NHC(O)-.

The term "halogen", as used herein refers to an atom of chlorine, bromine, iodine or fluorine.

The term "carbonyl" as used herein refers to a group of formula –C=O.

The term "carboxyl", as used herein refers to a group of formula –C(O)O- or –OC(O)-.


Generally R^1 is H or C_{1-3} alkyl. Preferred R^1 is C_{1-3} alkyl. Most preferred R^1 is methyl.

Generally R^2 is H or C_{1-3} alkyl. Preferred R^2 is C_{1-3} alkyl. Most preferred R^2 is methyl.

Generally R^3 is H, C_{1-10} alkyl, heterocycle or aryl. Preferred R^3 is H, phenyl or C_{1-10} alkyl. Most preferred R^3 is H.

Generally R is C_{1-10} alkyl, heterocycle or aryl. Preferred R is methyl, *iso*-butyl, *tert*-butyl, *iso*-propyl, ethylphenyl, phenyl, 2-amino-1-phenylethyl, 2-(2-amino-3-methyl-butyrylamino)-2-methyl-prop-1-yl, 1-amino-2-methyl-prop-1-yl, 2-(2-amino-acetylamino)-2-methyl- prop-1-yl. Most preferred R groups are *tert*-butyl, *iso*-propyl.

As used herein, "tautomer" refers to the migration of protons between adjacent single and double bonds. The tautomerization process is reversible. Compounds described herein can undergo any possible tautomerization that is within the physical characteristics of the compound. The following is a tautomerization example that can occur in compounds described herein:



Compounds of the invention are:

iso-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
Acetic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
Benzoic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
3-Methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
3-Phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-Amino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-(2-Amino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-(2-Amino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-Amino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

Intermediates of the invention are:
iso-Butyric acid 3-[(*S*)-1-(1-*iso*-butyryl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2,2-Dimethyl-propionic acid 3-{(*S*)-1-[1-(2,2-dimethyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;
Acetic acid 3-[(*S*)-1-(1-acetyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
Benzoic acid 3-[(*S*)-1-(1-benzoyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
3-Methyl-butyric acid 2-methyl-3-{(*S*)-1-[1-(3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;
Phenyl-propionic acid 2-methyl-3-{(*S*)-1-[1-(3-phenyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;
2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxy carbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;
2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxycarbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;

2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-(2-*tert*-Butoxycarbonylamino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;
2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

*S*ome compounds of Formula I , Formula II and Formula III and some of their intermediates have at least one stereogenic center in their structure. This stereogenic center may be present in an *(R)* or *(S)* configuration, said *(R)* and *(S)* notation is used in correspondence with the rules described in Pure Appli. Chem. (1976), 45, 11-13.

Compounds according to the present invention may exist in different polymorphic forms. Although not explicitly indicated in the above formula, such forms are intended to be included within the scope of the present invention.

Compounds of Formula I, Formula II or Formula III and their salts can be in the form of a solvate, which is included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

The term "pharmaceutically acceptable salts" refers to salts or complexes that retain the desired biological activity of the above identified compounds and exhibit minimal or no undesired toxicological effects. The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base or acid salt forms, which the compounds of Formula I, Formula II or Formula III are able to form. The acid addition salt form of a compound of Formula I, Formula II or Formula III that occurs in its free form as a base can be obtained by treating the free base with an appropriate acid such as an inorganic acid, for example but not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; or an organic acid such as for example but not limited to, as citric acid, acetic acid, oxalic acid, tartaric acid, succinic acid, malic acid, fumaric acid, ascorbic acid, benzoic acid, tannic acid, palmoic acid, alginic acid, polyglutamic acid, naphthalene-sulfonic acid, napthalenedisulfonic, and polygalacturonic acid as well as base

addition salts such as those formed with alkali- and alkaline earth metals such as sodium, potassium and calcium and the like (Handbook of Pharmaceutical *S*alts, P.Heinrich *S*tahal& Camille G. Wermuth (Eds), Verlag Helvetica Chemica Acta-Zürich, 2002, 329-345).

The compounds can also be administered as pharmaceutically acceptable quaternary salts known by those skilled in the art, which specifically include, but not limiting to the quaternary ammonium salt of the formula $-NY^+Z^-$, wherein Y is hydrogen, alkyl, or benzyl, and Z is a counterion, including but not limited to, chloride, bromide, iodide, -O-alkyl, toluenesulfonate, methylsulfonate, sulfonate, phosphate, or carboxylate (such as fumarate, benzoate, succinate, acetate, glycolate, maleate, malate, fumarate, citrate, tartrate, ascorbate, benzoate, cinnamoate, mandeloate, benzyloate, and diphenylacetate).

In another embodiment of the invention, there are provided pharmaceutical compositions including at least one compound of the invention in a pharmaceutically acceptable carrier thereof. The phrase "pharmaceutically acceptable" means the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Pharmaceutical compositions of the present invention can be used in the form of a solid, a solution, an emulsion, a dispersion, a patch, a micelle, a liposome, and the like, wherein the resulting composition contains one or more compounds of the present invention, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for enteral or parenteral applications. Invention compounds may be combined, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The carriers which can be used include but are not limited to, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. Invention compounds are included in the pharmaceutical

composition in an amount sufficient to produce the desired effect upon the process or disease condition.

Pharmaceutical compositions containing invention compounds may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of a sweetening agent such as sucrose, lactose, or saccharin, flavoring agents such as peppermint, oil of wintergreen or cherry, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets containing invention compounds in admixture with non-toxic pharmaceutically acceptable excipients may also be manufactured by known methods. The excipients used may be, for example, (1) inert diluents such as calcium carbonate, lactose, calcium phosphate or sodium phosphate; (2) granulating and disintegrating agents such as corn starch, potato starch or alginic acid; (3) binding agents such as gum tragacanth, corn starch, gelatin or acacia, and (4) lubricating agents such as magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. In some cases, formulations for oral use may be in the form of hard gelatin capsules wherein the invention compounds are mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the form of soft gelatin capsules wherein the invention compounds are mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

The pharmaceutical compositions may be in the form of a sterile injectable suspension. This suspension may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic

parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides, fatty acids (including oleic acid), naturally occurring vegetable oils like sesame oil, coconut oil, peanut oil, cottonseed oil, etc., or synthetic fatty vehicles like ethyl oleate or the like. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

The present invention concerns also the use of a compound of Formula I, Formula II or Formula III, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the therapeutic application. The present invention concerns also the a method for manufacturing a medicament intended for therapeutic application wherein a compound having general Formula I, Formula II or Formula III, or a pharmaceutically active derivative or salt thereof is used.

Since individual subjects may present a wide variation in severity of symptoms and each drug has its unique therapeutic characteristics, the precise mode of administration and dosage employed for each subject is left to the discretion of the practitioner. The patient will be administered the compound orally in any acceptable form, such as a tablet, liquid, capsule, powder and the like, or other routes may be desirable or necessary, particularly if the patient suffers from nausea. Such other routes may include, without exception, transdermal, parenteral, subcutaneous, intranasal, via an implant stent, intrathecal, intravitreal, topical to the eye, back to the eye, intramuscular, intravenous, and intrarectal modes of delivery. The actual amount of the compound to be administered in any given case will be determined by a physician taking into account the relevant circumstances, such as the severity of the condition, the age and weight of the patient, the patient's general physical condition, the cause of the condition, and the route of administration. Additionally, the formulations may be designed to delay release of the active compound over a given period of time, or to carefully control the amount of drug released at a given time during the course of therapy.


Ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol or of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol and their pharmaceutically-acceptable salts may be administered through different routes, including but not limited to topical eye drops, direct injection, application at the back of the eye or formulations that may further enhance the long duration of actions such as a slow releasing pellet, suspension, gel, or sustained delivery devices such as any suitable drug delivery system (DDS) known in the art. While topical administration is preferred, this compound may also be used in an intraocular implant as described in U.S. U.S. Patent 7,931,909 which is hereby incorporated by reference. Such biocompatible intraocular implants include ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol or of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol and a polymer associated with ester pro-drugs of [3-(1-(*1H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol or of (*R*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol to facilitate release thereof into an eye for an extended period of time.

Ophthalmic formulations of drug products are well known in the art and described in, for example, U.S. Patent Application Publication No. 20050059583; No. 20050277584; U.S. Patent No. 7,297,679; and No. 20070015691; and U.S. Patent Nos. 5,474,979 and 6,582,718, the disclosures of all which are incorporated herein by reference. The ester pro-drugs of [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol, of (*S*) [3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol or of (*R*) [3-(1-(1H-imidazol-4-yl)ethyl)-2-methylphenyl] methanol may be formulated with efficacy enhancing components as disclosed in U.S. Patent Number 7,491,383 B2, which is hereby incorporated by reference in its entirety.

With respect to the present invention reference to a compound or compounds, is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

The present invention also concerns a process for preparing the compounds having general Formula I, Formula II or Formula III. The synthetic scheme set forth below, illustrates how compounds according to the invention can be made. Those skilled in the art will be able to routinely modify and/or adapt the following scheme to synthesize any compounds of the invention covered by Formula I, Formula II or Formula III.

General scheme for synthesizing ester prodrugs of (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol



In a first step (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol (CAS 189255-79-6) can react with the desired acyl chloride, in the presence of *N,N*-dimethyl formamide (DMF), tertahydrofuran (THF), triethylamine (TEA) and 4-dimethyl aminopyridine (DMAP). After a typical work-up by extraction, the residue can be purified by medium pressure liquid chromatography (MPLC) (0% to 40% ethyl acetate in hexanes) to yield the intermediate compound as a solid.

In a second step, the intermediate obtained in the first reaction, can react with methanol (MeOH). The residue can be purified by MPLC (50% ethyl acetate in hexanes then 5% 7N ammonia/ methanol /dichloromethane) to yield the desired compound as a solid.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the

invention claimed. As used herein, the use of the singular includes the plural unless specifically stated otherwise.

The present invention includes all pharmaceutically acceptable isotopically enriched compounds. Any compound of the invention may contain one or more isotopic atoms enriched or different than the natural ratio such as deuterium ^{2}H (or D) in place of protium ^{1}H (or H) or use of ^{13}C enriched material in place of ^{12}C and the like. Similar substitutions can be employed for N, O and S. The use of isotopes may assist in analytical as well as therapeutic aspects of the invention. For example, use of deuterium may increase the *in vivo* half-life by altering the metabolism (rate) of the compounds of the invention. These compounds can be prepared in accord with the preparations described by use of isotopically enriched reagents.

The following examples are for illustrative purposes only and are not intended, nor should they be construed as limiting the invention in any manner. Those skilled in the art will appreciate that variations and modifications of the following examples can be made without exceeding the spirit or scope of the invention.

The IUPAC names of the compounds mentioned in the examples were generated with ACD version 8.

Unless specified otherwise in the examples, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on 300 MHz Varian and acquired at room temperature. Chemical shifts are given in ppm referenced either to internal TMS or to the residual solvent signal.

All the reagents, solvents, catalysts for which the synthesis is not described are purchased from chemical vendors such as Sigma Aldrich, Fluka, Lancaster, however some known reaction intermediates, for which the CAS registry number is mentioned, were prepared in-house following known procedures.
Usually the compounds of the invention were purified by flash column chromatography.

The following abbreviations are used in the examples:

DCM	dichloromethane
MeOH	methanol
CD_3OD	deuterated methanol
NH_3	ammonia
Na_2SO_4	sodium sulfate
DMF	N,N-dimethylformamide
$MgSO_4$	magnesium sulfate
EtOAc	ethylacetate
i-PrOH	*iso*-propanol
$CDCl_3$	deuterated chloroform
MPLC	medium pressure liquid chromatography
DMF	dimethylformamide
TEA	triethylamine
THF	tertahydrofuran
DMAP	4-dimethylaminopyridine
RT	room temperature
Boc-L-Valine	N-(*tert*-Butoxycarbonyl)-L-valine
Boc-Glycine	N-(*tert*-Butoxycarbonyl)glycine
Boc-L-Phenylalanine	N-(*tert*-Butoxycarbonyl)-L-phenylalanine
HCl	hydrochloric acid
H_2O	water
EDCI	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
$NaHCO_3$	sodium bicarbonate

Example 1

Intermediate 1

***iso*-Butyric acid 3-[(*S*)-1-(1-isobutyryl-1*H*-imidazol-4-yl)-ethyl] -2-methyl-benzyl ester**

To a solution of (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol (1.34g, 6.2mmol) in DMF (8ml) and THF (50ml), were added TEA (3.5ml, 24.8mmol), DMAP (780mg, 6.2mmol) and *iso*-butyryl chloride (2.18g, 20.5mmol). The resulting mixture

was stirred at RT for 16 h, quenched with H_2O and extracted with ethyl acetate. The combined organic layers were washed with brine, and dried over Na_2SO_4, and concentrated under reduced pressure. The residue was purified by MPLC (0% to 40% ethyl acetate in hexanes) to yield **Intermediate 1** as solid.

^{1}H-NMR (CD_3OD, δ ppm): 1.15 (d, J=7.03Hz, 6H), 1.26 (d, 6H, J=6.74Hz), 1.56 (d, J=7.03Hz, 3H), 2.34 (s, 3H), 2.58 (hept, J=7.03Hz, 1H), 3.34(hept, J=7.74Hz, 1H), 4.42(q, J=7.03Hz, 1H), 5.15(s, 2H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.31 (s, 1H), 8.35 (s, 1H).

Intermediates 2-6 were prepared in a similar manner to the method described in **Example 1** starting with (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol. The acyl chloride used in each case and the results are tabulated below in **Table 1**.

Table 1


Intermediate number	IUPAC name	Acyl chloride	^{1} NMR (Solvent; δ ppm)
2	**2,2-Dimethyl-propionic acid 3-{(*S*)-1-[1-(2,2-dimethyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester**	Pivaloyl chloride	(CD_3OD): 1.19 (s, 9H), 1.42 (s, 9H), 1.56 (d, J=7.03Hz, 3H), 2.34 (s, 3H), 4.42(q, J=7.03Hz, 1H), 5.15(s, 2H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.33 (s, 1H), 8.40 (s, 1H).
3	**Acetic acid 3-[(*S*)-1-(1-acetyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester**	Acetyl chloride	(CD_3OD): 1.55 (d, J=7.03Hz, 3H), 2.05 (s, 3H), 2.33 (s, 3H), 2.58 (s, 3H), 4.39(q, J=7.03Hz, 1H), 5.15(s, 2H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.30 (s, 1H), 8.29 (s, 1H).
4	**Benzoic acid 3-[(*S*)-1-(1-benzoyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester:**	Benzoyl chloride	(CD_3OD): 1.58 (d, J=7.03Hz, 3H), 2.43 (s, 3H), 4.46(q, J=7.03Hz, 1H), 5.41 (s, 2H), 7.11-7.18

			(m, 2H), 7.27-7.35 (m, 2H), 7.42-7.50 (m, 2H), 7.50-7.63 (m, 3H), 7.65-7.71 (m, 1H), 7.79 (d, J=7.33Hz, 2H), 8.00 (d, J=7.33Hz, 2H), 8/09 (s, 1H).
5	**3-Methyl-butyric acid 2-methyl-3-{(*S*)-1-[1-(3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester**	Methylbutanoyl chloride	(CD_3OD): 0.91 (d, J=6.44Hz, 6H), 1.01 (d, J=6.44Hz, 6H), 1.54 (d, J=7.03Hz, 3H), 2.05 (hept, J=6.44Hz, 1H), 2.15-2.25 (m, 3H), 2.33 (s, 3H), 2.81 (d, J=7.03Hz, 3H), 4.42(q, J=7.03Hz, 1H), 5.14(s, 2H), 7.07-7.19 (m, 3H), 7.28 (s, 1H), 8.32 (s, 1H).
6	**3-Phenyl-propionic acid 2-methyl-3-{(*S*)-1-[1-(3-phenyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester**	Phenylpropanoyl chloride	(CD_3OD): 1.52 (d, J=7.03Hz, 3H), 2.24 (s, 3H), 2.64 (t, J=7.61Hz, 2H), 2.90 (t, J=7.61Hz, 2H), 3.04 (t, J=7.61Hz, 2H),3.24 (t, J=7.61Hz, 2H), 4.34 (q, J=7.03Hz, 1H), 5.13 (s, 2H), 7.08-7.248 (m, 14H), 8.25 (s, 1H).

Example 2

Compound 1

***iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester**








Intermediate 1 was dissolved in MeOH (50ml) and the mixture was stirred at RT for 24 h and then concentrated under reduced pressure. The residue was purified by MPLC (50% ethyl acetate in hexanes then 5% 7N NH_3/ MeOH /DCM) to yield **Compound 1** as a solid.

^{1}H-NMR (CD_3OD; δ ppm): 1.15 (d, J=7.03Hz, 6H), 1.54 (d, J=7.03Hz, 3H), 2.33 (s, 3H), 2.56 (hept, J=7.03Hz, 1H), 4.42(q, J=7.03Hz, 1H), 5.15(s, 2H), 6.70 (s, 1H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.55 (s, 1H).

Compounds 2-6 and of the invention were prepared according to the procedure described in **Example 2**, by reacting the corresponding intermediate with methanol. The results are tabulated below in **Table 2**.

Table 2

Comp. No.	IUPAC name	Inter. No.	^{1}NMR (Solvent, δ ppm)
2	**2,2-Dimethyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester** 	2	(CD_3OD): 1.19 (s, 9H), 1.54 (d, J=7.03Hz, 3H), 2.33 (s, 3H), 4.42 (q, J=7.03Hz, 1H), 5.13 (s, 2H), 6.70 (s, 1H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.55 (s, 1H).
3	**Acetic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl Ester**	3	(CD_3OD): 1.54 (d, J=7.03Hz, 3H), 2.04 (s, 3H), 2.33 (s, 3H), 4.42 (q, J=7.03Hz, 1H), 5.13 (s, 2H), 6.70 (s, 1H), 7.07-7.10 (m,

			2H), 7.12-7.15 (m, 1H), 7.55 (s, 1H).
4	**Benzoic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester** 	4	(CD_3OD): 1.54 (d, J=7.03Hz, 3H), 2.31 (s, 3H), 4.42(q, J=7.03Hz, 1H), 5.13 (s, 2H), 6.70 (s, 1H), 7.07-7.15 (m, 2H), 7.25-7.28 (m, 1H), 7.54-7.47 (m, 2H), 7.55-7.60 (m, 2H), 8.0 (d, J=7.33Hz, 2H).
5	**3-Methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl Ester** 	5	(CD_3OD): 0.93 (d, J=7.03Hz, 6H), 1.54 (d, J=7.03Hz, 3H), 2.07 (hept, J=7.03Hz, 1H), 2.21 (d, J=7.03Hz, 2H), 2.33 (s, 3H), 4.42(q, J=7.03Hz, 1H), 5.15(s, 2H), 6.70 (s, 1H), 7.07-7.10 (m, 2H), 7.12-7.15 (m, 1H), 7.55 (s, 1H).
6	**3-Phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl Ester** 	6	(CD_3OD): 1.54 (d, J=7.03Hz, 3H), 2.23 (s, 3H), 2.65 (t, J=7.61Hz, 2H), 2.91 (t, J=7.61Hz, 2H), 4.40 (q, J=7.03Hz, 1H), 5.13 (s, 2H), 6.70 (s, 1H), 7.08-7.24 (m, 8H), 7.55 (s, 1H).

Example 3

Intermediate 7

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxy carbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester

To a solution of (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol (216mg, 1.0mmol) in DMF (2ml) and THF (12ml) were added EDCI (671mg, 3.5mmol), DMAP (427mg, 3.5mmol) and Boc-L-Valine (651mg, 3.0mmol) . The mixture was stirred at RT for 16 h, quenched with H_2O and extracted with ethyl acetate. The combined organic layers were washed with H_2O, brine, and dried over Na_2SO_4, and concentrated under reduced pressure. The residue was purified by a column

chromatography (30% ethyl acetate in hexanes) to yield **Intermediate 7** as a white solid.

^{1}H-NMR (CD_3OD; δ ppm): 0.85-1.01 (m, 12H), 1.20-1.48 (m, 18H), 1.56 (d, J=7.03Hz, 3H), 2.01-2.20(m, 2H), 2.35 (s, 3H), 4.03(m, 1H), 4.42 (q, J=7.03Hz, 1H), 4.60-4.65 (m, 1H), 5.15-5.29 (m, 2H), 7.10-7.20 (m, 2H), 7.20-7.25 (m, 1H), 7.33 (s, 1H), 8.44 (s, 1H).

Example 4

Intermediate 8

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester


The title compound was prepared from **Intermediate 7** (600mg, 0.98mmol) in 30ml of MeOH according to the procedure described in **Example 2**.

^{1}H-NMR (CD_3OD; δ ppm): 0.85-0.95 (m, 6H), 1.42 (m, 9H), 1.54 (d, J=7.03Hz, 3H), 2.05 (m, 1H), 2.33 (s, 3H), 4.00 (d, J=6.15Hz, 1H), 4.40 (q, J=7.03Hz, 1H), 5.15-5.28 (m, 2H), 6.67 (s, 1H), 7.10-7.20 (m, 2H), 7.20-7.25 (m, 1H), 7.55 (s, 1H).

Example 5

Compound 7

2-Amino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester



To **Intermediate 8** (390mg, 0.94mmol) was added 4N HCl in dioxane (8ml). The resulting solution was stirred at RT for 4 hrs, then quenched with H_2O, neutralized with aqueous saturated $NaHCO_3$ and extracted with 25% isopropyl alcohol in chloroform. The combined organic layers were dried over Na_2SO_4, and concentrated

under reduced pressure. The residue was purified by a column chromatography (5% 7N NH_3/MeOH in DCM) to yield **Compound 7** as a white solid.

^{1}H-NMR (CD_3OD; δ ppm): 0.85 (d, J=6.74Hz, 3H), 0.91 (d, J=6.74Hz, 3H), 1.54 (d, J=7.03Hz, 3H), 1.96 (hept, J=6.74Hz, 1H), 2.33 (s, 3H), 3.28 (d, J=6.74Hz, 2H), 4.42 (q, J=7.03Hz, 1H), 5.20-5.25 (m, 2H), 6.67 (s, 1H), 7.10-7.12 (m, 2H), 7.13-7.20 (m, 1H), 7.55 (s, 1H).

Example 6

Intermediate 9

2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxycarbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester

The title compound was prepared from **Compound 7** (490mg, 1.55mmol), Boc-L-Valine (1.01g, 4.67mmol), EDCI (1.04g, 5.42mmol) and DMAP (671mg, 5.5mmol) according to the procedure described in **Example 3**.

^{1}H-NMR (CD_3OD; δ ppm): 0.85-0.92 (m, 12H), 1.43 (s, 9H), 1.55 (d, J=7.03Hz, 3H), 1.97 (m, 1H), 2.14 (hept, J=6.60Hz, 1H), 2.35 (s, 3H), 3.88 (d, J=7.30Hz,, 1H), 4.35 (d, J=6.90Hz, 1H), 4.42(, d, J=7.03Hz, 1H), 5.18-5.25 (m, 2H), 6.67 (s, 1H), 7.10-7.15 (m, 2H),7/17-7.20 (m, 1H), 7.55 (s, 1H).

Example 7

Intermediate 10

2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester


The title compound was prepared from **Intermediate 9** (750mg, 1.05mmol) in 30ml of MeOH according to the procedure described in **Example 2**.

^{1}H-NMR (CD_3OD; δ ppm): 0.89 (d,d , J=7.03Hz, 6H), 1.44 (s, 9H), 1.54 (d, J=7.33Hz, 3H), 2.14 (hept, J=6.74Hz, 1H), 2.33 (s, 3H), 3.74 (s, 2H), 4.35-4.55 (m, 2H), 5.20 (s, 2H), 6.67 (s, 1H), 7.10-7.17 (m, 2H), 7.19-7.23 (m, 1H), 7.56 (s, 1H).

Example 8

Compound 8

2-(2-Amino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester



The title compound was prepared from **Intermediate 10** (450mg, 0.87mmol) in 8ml of 4N HCl/Dioxane according to the procedure described in **Example 5**.

^{1}H-NMR (CD_3OD; δ ppm): 0.85 (d, J=7.03Hz, 3H), 0.91 (d, J=6.74Hz, 3H), 0.92 (d, J=7.3Hz. 3H), 1.14 (d, J=6.2Hz, 3H), 1.54 (d, J=7.03Hz, 3H), 1.94 (hept, J=5.2Hz, 1H), 2.14 (hept, J=6.2Hz, 1H), 2.33 (s, 3H), 3.18 (d, J=5.2Hz, 1H), 4.34 (d, J=6.2Hz, 1H), 4.42(q, J=7.03Hz, 1H), 5.21-5.26 (m, 2H), 6.67 (s, 1H), 7.10-7.15 (m, 2H), 7.18-7.20 (m, 1H), 7.55 (s, 1H).

Example 9

Intermediate 11

2-(2-*tert*-Butoxycarbonylamino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester


The title compound was prepared from **Compound 8** (405mg, 1.28mmol), Boc-Glycine(675mg, 3.86mmol), EDCI(859mg, 4.48mmol) and DMAP(547mg, 4.48mmol) according to the procedure described in **Example 3**. The title compound was purified by column chromatography using 5% 7N NH_3/MeOH in DCM .

^{1}H-NMR (CD_3OD; δ ppm): 0.89 (d, J=6.74Hz, 3H), 0.91 (d, J=6.74Hz, 3H), 1.55 (d, J=7.30Hz, 3H), 2.14 (hept, J=6.74Hz, 1H), 2.33 (s, 3H), 4.37 (d, J=5.90Hz, 1H), 4.42(q, J=7.03Hz, 1H), 5.20-5.25 (m, 2H), 6.67 (s, 1H), 7.10-7.12 (m, 2H), 7.13-7.20 (m, 1H), 7.55 (s, 1H).

Example 10

Compound 9

2-(2-Amino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester



The title compound was prepared from **Intermediate 11** (320mg, 0.68mmol) with 10ml of 4N HCl/Dioxane according the procedure described in **Example 5**.

^{1}H-NMR (CD_3OD; δ ppm): 0.89 (d, J=6.74Hz, 3H), 0.91 (d, J=6.74Hz, 3H), 2.14 (hept, J=6.74Hz, 1H), 2.33 (s, 3H), 4.37 (d, J=5.90Hz, 1H), 4.42(q, J=7.03Hz, 1H), 5.20-5.25 (m, 2H), 6.67 (s, 1H), 7.10-7.12 (m, 2H), 7.13-7.20 (m, 1H), 7.55 (s, 1H).

Example 11

Intermediate 12

2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester


The title compound was prepared from (*S*)-[3-(1-(1*H*-imidazol-4-yl)ethyl)-2-methylphenyl] methanol (216mg, 1.0mmol), Boc-L-Phenylalanine(795mg, 3.0mmol), EDCI(671mg, 3.5mmol) and DMAP(427mg, 3.5mmol) according to the procedure described in **Example 3**. **Intermediate 12** was purified by a column chromatography using 35-100% ethyl acetate in hexane.

^{1}H-NMR (CD_3OD; δ ppm): 1.36 (s, 9H), 1.55 (d, J=7.03Hz, 3H), 2.28 (s, 3H), 2.85-2.95 (m, 1H), 3.05-3.11(m, 1H), 4.38(m, 1H), 4.40(q, J=7.03Hz, 1H), 5.17(s, 2H), 6.69 (s, 1H), 7.08-7.24 (m, 8H), 7.55 (s, 1H).

Example 12

Compound 10

2-Amino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester



The title compound was prepared from **Intermediate 12** (240mg, 0.52mmol) with 8ml of 4N HCl/Dioxane according to the procedure described in **Example 5**.

^{1}H-NMR (CD_3OD; δ ppm): 1.54 (d, J=7.03Hz, 3H), 2.26 (s, 3H), 2.90-3.00 (m, 2H), 3.73 (t, J=6.40Hz, 1H), 4.40(q, J=7.03Hz, 1H), 5.13-5.18(m, 2H), 6.68 (s, 1H), 7.08-7.12 (m, 5H), 7.13-7.22 (m, 3H), 7.55 (s, 1H).

The following assay was used to demonstrate the potency and selectivity of the compounds according to the invention.

Example 13

Visual enhancement model

Sixteen pigmented (Dutch-Belted) rabbits weighing 2-3 kg are used to evaluate the neuroenhancement effect of pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester. Rabbits are dosed with pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester through intravenous route. Spatial sweep visual evoked potential (sVEP) acuity is assessed with PowerDiva software version 1.8. Recordings were made bilaterally in conscious animals. The results demostrate that pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester enhances visual acuity at 10-30 minutes post-dose in normal DB rabbits.

Example 14

The nerve crush model

This example describes the neuroprotective effect of pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester level in the rat nerve crush model. Sprague Dawley rats weighing 300-350 g were anesthetized with a mixture of ketamine (50mg/kg) and xylazine (0.5 mg/kg). Lateral canthotomy was performed in the right eye and an incision was made in the superior conjunctiva adjacent to the rectus muscle. This was followed by a blunt dissection until optic nerve was exposed. A partial compression was applied to the optic nerve for 30 seconds, 2 to 3 mm distal from the globe, using calibrated cross-acting forceps. Care was taken not to interfere with retinal blood supply. Pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester was administered at 0.03, 0.1, 0.3, 1 mg/kg SC two hours before nerve injury, the vehicle PBS was administered SC as a negative control whereas brimonidine 0.1mg/kg was given by IP injection as a positive control. Control animals receive phosphate-buffered saline (PBS) vehicle. The experiment was terminated 12-15 days later.

Example 15

The chronic ocular hypertension model

Intraocular Pressure (IOP) was elevated in male Witar rats weighing 350-450 g using laser photocoagulation with blue-green argon laser (Coherent, Palo Alto, CA). Rats were anesthetized with a mixture of ketamine (15 mg/kg), acepromazine (1.5 mg/kg), and xylazine (0.3 mg/kg). Laser treatment was done in two parts (1-week interval) on limbal and epsiscleral veins. The amount of energy used was 1 W for 0.2 seconds, delivering a total of 150 spots (50-100 μM). Intraocular pressure was measured using tonometer (TONO-PEN: mentor, Norwell, MA). Rats were sedated with 3.0 mg/kg IM acepromazine during IOP measurements. Proparacaine 0.5% was applied topically on the eyes to anesthetize the cornea. Initial IOP measurements were done before laser treatment to determine baseline IOP and subsequent measurements were done once a week.

Pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester is administered constantly using an osmotic pump (Alzet Osmotic Pumps, Duret Corp., Cupertino, CA) which was inserted subcutaneously on the back at 0.03, 0.1, 0.3, 1 mg/kg/day SC two hours before the first laser treatment (preventive mode) or following the second laser treatment (therapeutic mode). The vehicle PBS was administered by SC osmotic pump as a negative control whereas brimonidine 0.1mg/kg was given by IP injection two hours before the first laser treatment as a positive control. Control animals received phosphate-buffered saline (PBS) vehicle. The experiment was terminated 15-25 days later.

Example 16

The blue light model

Pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester is evaluated in the blue light model of retinal degeneration in rats. The drugs is administered continuously with subcutaneous infusion pumps at a dose of 1 mg/kg/day starting two days before blue light exposure. Twenty 4-month old male Sprague Dawley rats (body weight 470-550 g) were used in this study. The animals were exposed to room light on a 12 hour light/12 hour dark cycle before the experiment. All animals were dark adapted overnight (16-20 hours) before blue light. Under the intensity of 6100-6500 lux, rats were exposed to blue light for 4 hours. After the blue light, rats were placed in the dark for another 3 days before returning to normal 12 hour light/12 hour dark. Ocular Coherence Tomography (OCT) measurement was performed at 7 days post blue light exposure. The results demonstrate that blue light exposure with just saline treatment leads to dramatic reduction of retinal thickness measured by OCT, particularly in the superior retina. Histology studies have shown that the reduction in retinal thickness is attributable to loss of photoreceptors. Brimonidine treatment did not prevent the change in retinal thickness while treatment with pro-drug *iso*-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester significantly reduced the damage caused by blue light.

2011301901 15 Jan 2016


Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

2011301901 15 Jan 2016

The claims defining the invention are as follows:

1. A method of treating a retinal disease comprising administering a therapeutically effective amount of a composition comprising a compound of Formula I, its individual enantiomers, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula I

wherein

R^1 is H or C_{1-3} alkyl;


R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl) or -S(C_{1-6} alkyl) groups,

to a subject in need thereof.

2. A method according to claim 1, wherein the compound is a compound of Formula II, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,




Formula II

wherein

R^1, R^2, R^3 and R are as defined in claim 1.

3. A method according to claim 1, wherein the compound is a compound of Formula III, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula III

wherein

R^1, R^2, R^3 and R are as defined in claim 1.

4. A method according to any one of claims 1 to 3, wherein R^1 is C_{1-3} alkyl, R^2 is C_{1-3} alkyl, R^3 is H and R is C_{1-10} alkyl.

5. A method according to any one of claims 1 to 3, wherein R^1 is methyl, R^2 is methyl, R^3 is H and R is C_{1-4} alkyl.

6. A method according to claim 1 or claim 2, wherein the compound is selected from the group consisting of

iso-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Acetic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Benzoic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-Amino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester; and

2-Amino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

7. A method according to claim 1 or claim 2, wherein the compound is selected from the group consisting of

iso-Butyric acid 3-[(*S*)-1-(1-*iso*-butyryl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-{(*S*)-1-[1-(2,2-dimethyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;

Acetic acid 3-[(*S*)-1-(1-acetyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Benzoic acid 3-[(*S*)-1-(1-benzoyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Methyl-butyric acid 2-methyl-3-{(*S*)-1-[1-(3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;

Phenyl-propionic acid 2-methyl-3-{(*S*)-1-[1-(3-phenyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxy carbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxycarbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;


2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-*tert*-Butoxycarbonylamino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester; and

2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

8. A method according to any one of claims 1 to 7, wherein the retinal disease is selected from the group consisting of age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion.

9. Use of a compound of Formula I, its individual enantiomers, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula I

wherein

R^1 is H or C_{1-3} alkyl;

R^2 is H or C_{1-3} alkyl;


R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -

$C(O)NH(C_{1-6}$ alkyl), $-S(C_{1-6}$ alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, $-O(C_{1-6}$ alkyl), $-OC(O)(C_{1-6}$ alkyl), $-C(O)O(C_{1-6}$ alkyl), $-NHC(O)(C_{1-6}$ alkyl), $-C(O)NH(C_{1-6}$ alkyl) or $-S(C_{1-6}$ alkyl) groups,

in the manufacture of a medicament for the treatment of a retinal disease.

10. Use according to claim 9, wherein the compound is a compound of Formula II, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,




Formula II

wherein

R^1, R^2, R^3 and R are as defined in claim 9.

11. Use according to claim 9, wherein the compound is a compound of Formula III, its individual diastereoisomers, its hydrates, its solvates, its crystal forms, its individual tautomers or a pharmaceutically acceptable salt thereof,



Formula III

wherein

R^1, R^2, R^3 and R are as defined in claim 9.

12. Use according to any one of claims 9 to 11, wherein R^1 is C_{1-3} alkyl, R^2 is C_{1-3} alkyl, R^3 is H and R is C_{1-10} alkyl.

13. Use according to any one of claims 9 to 11, wherein R^1 is methyl, R^2 is methyl, R^3 is H and R is C_{1-4} alkyl.

14. Use according to claim 9 or claim 10, wherein the compound is selected from the group consisting of

iso-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Acetic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Benzoic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-Amino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester; and

2-Amino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

15. Use according to claim 9 or claim 10, wherein the compound is selected from the group consisting of

iso-Butyric acid 3-[(*S*)-1-(1-*iso*-butyryl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-{(*S*)-1-[1-(2,2-dimethyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;

Acetic acid 3-[(*S*)-1-(1-acetyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Benzoic acid 3-[(*S*)-1-(1-benzoyl-1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Methyl-butyric acid 2-methyl-3-{(*S*)-1-[1-(3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;

Phenyl-propionic acid 2-methyl-3-{(*S*)-1-[1-(3-phenyl-propionyl)-1*H*-imidazol-4-yl]-ethyl}-benzyl ester;

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxy carbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;

2-*tert*-Butoxycarbonylamino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-{(*S*)-1-[1-(2-*tert*-butoxycarbonylamino-3-methyl-butyryl)-1*H*-imidazol-4-yl]-ethyl}-2-methyl-benzyl ester;


2-(2-*tert*-Butoxycarbonylamino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-*tert*-Butoxycarbonylamino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester; and

2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.

16. Use according to any one of claims 9 to 15, wherein the retinal disease is selected from the group consisting of age related macular degeneration, wet macular degeneration, dry macular degeneration, geographic atrophy, diabetic retinopathy, diabetic macular edema and retinal vein occlusion.

17. A pharmaceutical composition when used for treating a retinal disease in a subject, the composition comprising a compound of Formula II



Formula II

wherein

R^1 is H or C_{1-3} alkyl;

R^2 is H or C_{1-3} alkyl;

R^3 is H, C_{1-10} alkyl, an aromatic or non-aromatic 5- to 10-membered monocyclic or bicyclic heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof; or a C_{6-10} monocyclic or bicyclic aryl; and

2011301901 15 Jan 2016

R is C_{1-10} alkyl group optionally substituted by amino groups, aryl groups and halogen atoms, wherein one methylene group of the alkyl chain can be replaced by carbonyl, -NH-, carboxyl, amide, -S- or -O-; an aromatic or non-aromatic 5 to 10 membered monocyclic or bicyclic, heterocycle containing at least one heteroatom selected from O, N or S or combinations thereof, wherein the heterocycle is optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl), -S(C_{1-6} alkyl) groups; or a C_{6-10} monocyclic or bicyclic aryl group optionally substituted by C_{1-6} alkyl, amino, halogen, -O(C_{1-6} alkyl), -OC(O)(C_{1-6} alkyl), -C(O)O(C_{1-6} alkyl), -NHC(O)(C_{1-6} alkyl), -C(O)NH(C_{1-6} alkyl) or -S(C_{1-6} alkyl) groups.

18. The pharmaceutical composition of claim 17, wherein the compound is selected from the group consisting of

iso-Butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2,2-Dimethyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Acetic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

Benzoic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

3-Phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-Amino-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-3-methyl-butyrylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester;

2-(2-Amino-acetylamino)-3-methyl-butyric acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester; and

2-Amino-3-phenyl-propionic acid 3-[(*S*)-1-(1*H*-imidazol-4-yl)-ethyl]-2-methyl-benzyl ester.