



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2139028 C 2003/11/18

(11)(21) **2 139 028**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 1994/12/23

(41) Mise à la disp. pub./Open to Public Insp.: 1995/06/29

(45) Date de délivrance/Issue Date: 2003/11/18

(30) Priorité/Priority: 1993/12/28 (HEI-5-334223) JP

(51) Cl.Int.⁵/Int.Cl.⁵ A23K 1/16

(72) Inventeurs/Inventors:
NAKASUGI, SHIGERU, JP;
SHIMADA, KENJIRO, JP;
MORISHITA, KOJI, JP

(73) Propriétaire/Owner:
KYOWA HAKKO KOGYO CO., LTD., JP

(74) Agent: BERESKIN & PARR

(54) Titre : COMPOSITION NUTRITIVE POUR ANIMAUX

(54) Title: NOURISHING COMPOSITION FOR ANIMALS

(57) Abrégé/Abstract:

A nourishing composition for animals which comprises an aqueous solution containing (1) at least one of dextrin and sugar alcohols, and (2) amino acids. When given to animals together with drinking water, the nourishing composition for animals according to the present invention can improve the nutritional conditions of the animals without causing browning in the aqueous solution.



ABSTRACT OF THE DISCLOSURE

A nourishing composition for animals which comprises an aqueous solution containing (1) at least one of dextrin and sugar alcohols, and (2) amino acids. When given to animals together with drinking water, the nourishing composition for animals according to the present invention can improve the nutritional conditions of the animals without causing browning in the aqueous solution.

NOURISHING COMPOSITION FOR ANIMALS

FIELD OF THE INVENTION

This invention relates to a nourishing composition for animals which is useful in improving the nutritional conditions of animals when given together with drinking water.

BACKGROUND OF THE INVENTION

In summer, domestic animals generally drink much water but show decreased appetite, which causes various problems, for example, in the case of pig etc., stagnation in body weight gain and reduction of mating performance, in the case of broiler etc., stagnation in body weight gain, in the case of layer etc., reduction of egg laying performance, decrease in egg shell strength and increase in the ratio of soft/broken shell eggs.

Further, problems such as stagnation in body weight gain may occur at the time of the shipment of broilers, the transfer of large layer chicks, the debeaking of layers, forced moulting of layers, the change of the feed for layers or broilers, the first feeding of the new-born layers or broilers, vaccination of layers or broilers, farrowing and weaning of piglings and the shipment of mother pigs.

To prevent these problems, attempts have been made to add nutritional components such as amino acids to feeds to thereby improve the nutritional conditions of domestic animals.

Even if nutritional components such as amino acids are added to feeds, however, the decreased appetite makes it impossible to achieve the desired effects.

In addition, feed remaining in the digestive organs of pork type pigs, broilers etc. at dressing contravenes the Food Hygiene Law. It is therefore necessary to omit feed and give only water to domestic animals from the day before shipment.

Accordingly, it is expected that when nutritional components including amino acids are added to drinking water, then the nutritional conditions of domestic animals can be maintained or improved even though they show decreased appetite or are fasting.

JP-A-2-255047 has proposed a method for improving the amino acid uptake of an animal by orally administering a liquid composition comprising amino acids together with other nutritional components, for example, carbohydrates, vitamins, minerals and enzymes (the term "JP-A" as used herein means an "unexamined published Japanese patent application").

However, the composition disclosed in the above publication suffers from several disadvantages, for example that a sufficient amount of amino acids can be hardly dissolved in the composition and that glucose, which is one of the carbohydrates employed as an additive, undergoes aminocarbonylation with the amino acids in an aqueous solution resulting in browning or solidification of the feed

aqueous solution.

SUMMARY OF THE INVENTION

The present invention aims at providing a nourishing composition for animals containing (1) at least one of dextrin and sugar alcohols, and (2) amino acids where the amino acids are dissolved in a sufficient amount and never undergo aminocarbonylation even though the composition is formulated into an aqueous solution.

Accordingly, the present invention relates to a nourishing composition for animals which comprises an aqueous solution containing (1) at least one of dextrin and sugar alcohols, and (2) amino acids.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 shows an operation schedule for transferring large layer chicks.

DETAILED DESCRIPTION OF THE INVENTION

Any sugar alcohol may be used in the present invention, so long as it does not undergo aminocarbonylation with amino group of amino acids. Examples thereof include lactitol, isomaltitol, maltitol, mannitol, sorbitol, xylitol and erythritol. As the dextrin, those having a degree of polymerization ranging from 3 to 30, preferably from 5 to 25, may be used.

The amino acid (or acids) used in the present invention is not particularly restricted, so long as it is one required by animals. It is particularly preferable to

use essential amino acids such as lysine, methionine, tryptophan, threonine, alanine, glycine and glutamic acid.

The amino acid content in the nourishing composition for animals according to the present invention ranges from 0.001 to 5.0 % by weight, preferably from 0.003 to 2.0 % by weight (dry basis). For example, the contents of lysine, methionine, tryptophan, threonine, glutamic acid, glycine and alanine preferably range respectively from 0.002 to 2.0 % by weight, from 0.003 to 0.8 % by weight, from 0.001 to 0.3 % by weight, from 0.001 to 0.4 % by weight, from 0.005 to 2.0 % by weight, from 0.005 to 2.0 % by weight and from 0.005 to 2.0 % by weight. The content of dextrin or sugar alcohols ranges from 0.0005 to 10.0 % by weight, preferably 0.001 to 0.8 % by weight.

The nourishing composition for animals of the present invention may contain additional components such as dissolution aids, dispersing agents, stabilizers, preservatives, antibiotics and antioxidants. Also, it may contain proteinous materials such as soluble soybean protein, soluble fish protein, soluble gelatin and defatted milk powder, organic acids such as citric acid, malic acid and succinic acid, vitamins, minerals, ethyl alcohol, colorants and perfumes. It is preferable to use organic acids from among these additives, since they exhibit a buffering action in the solution, improve the appetite and play a role as a metabolite in the TCA cycle *in vivo* so as to serve as an

energy source.

The nourishing composition for animals according to the present invention can be produced by mixing amino acids with at least one of dextrin and sugar alcohols, and then dissolving or suspending the obtained mixture in water or an aqueous solution containing an organic solvent, which exerts no undesirable effects on animals, such as ethyl alcohol. The particle size (diameter) of the amino acids to be used in the formulation ranges from 1.0 to 1000 μm , preferably from 1.0 to 500 μm . For example, the particle sizes of lysine, methionine, tryptophan, threonine, glutamic acid, glycine and alanine preferably range respectively from 1.0 to 500 μm , from 1.0 to 200 μm , from 1.0 to 200 μm , from 1.0 to 500 μm , from 1.0 to 500 μm , from 1.0 to 500 μm and from 1.0 to 500 μm . It is preferable that the dissolution or suspending is carried out immediately before use.

The nourishing composition for animals of the present invention may be given to, for example, poultry such as layer and broiler, monogastric animals such as pig and ruminants such as cattle and sheep. The nourishing composition for animals of the present invention may be given to the animals *ad libitum*. Alternately, it may be orally given to them by force. The dose may range from 10 to 1,000 mg/kg/day in terms of amino acids and from 5 to 1,000 mg/kg/day in terms of dextrin or sugar alcohols. The compositions of the invention may be given as a single or divided administration.

Although the nourishing composition for animals according to the present invention is in the form of an aqueous solution, it never causes aminocarbonylation between sugars and amino acids and the aqueous solution undergoes neither browning nor solidification. Thus the nourishing composition for animals of the present invention makes it possible to increase amino acid uptake thus improving the nutritional conditions of animals.

The nourishing composition for animals of the present invention is useful in, for example, the shipment of broilers, the transfer of large layer chicks, the debeaking of layers, forced moulting of layers, feeding layers or broilers during the summer season, the change of the feed for layers or broilers, the first feeding of the new-born layers or broilers, vaccination of layers or broilers, farrowing, weaning of piglings, weight-maintenance of mother pigs at the shipment, prevention of body weight loss and reduction in mortality.

Next, working examples and test examples of the present invention will be given, but are not to be construed to limit the scope of the present invention.

Lysine, tryptophan, threonine and glutamic acid employed in the following examples were all L-compounds, while methionine and alanine employed therein were all DL-compounds. As the amino acids to be used in the present invention, mention may be made of D-form, L-form and DL-form.

EXAMPLE 1

Lysine, methionine, threonine, tryptophan, glycine, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.) and citric acid were mixed together at each ratio as specified in Table 1 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 15-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 1, 2 and 3 for animals were obtained.

Table 1

<u>Component</u>	<u>Composition 1</u> <u>(g)</u>	<u>Composition 2</u> <u>(g)</u>	<u>Composition 3</u> <u>(g)</u>
lysine	40	80	160
methionine	15	30	60
threonine	20	40	80
tryptophan	15	30	60
glycine	100	200	400
dextrin	300	610	1230
citric acid	10	10	10
total	500	1000	2000

EXAMPLE 2

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed

together at each ratio as specified in Table 2 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 15-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 4, 5 and 6 for animals were obtained. The vitamins used herein was a composition prepared by mixing the vitamins as specified in Table 3 and adding dextrin thereto in such a manner as to give a total amount of 10 g.

Table 2

<u>Component</u>	<u>Composition 4</u> <u>(g)</u>	<u>Composition 5</u> <u>(g)</u>	<u>Composition 6</u> <u>(g)</u>
lysine	10	20	40
methionine	6	15	25
threonine	3	5	10
tryptophan	1	3	5
dextrin	10	17	20
citric acid	5	10	10
vitamin	10	10	10
total	45	80	120

Table 3

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	50.0	biotin	2.0
riboflavin	100.0	choline	9000.0
pantothenic acid	200.0	folic acid	10.0
nicotinic acid	200.0	vitamin B ₁₂	0.2
vitamin B ₆	40.0	linoleic acid	20.0

EXAMPLE 3

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 4 and the resulting mixture was dissolved in 30 l of water. The obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 7 and 8 for animals were obtained. The vitamins used herein was a composition prepared by mixing the vitamins as specified in Table 5 and adding dextrin thereto to give a total amount of 10 g.

Table 4

<u>Component</u>	<u>Composition 7 (g)</u>	<u>Composition 8 (g)</u>
lysine	10	30
methionine	30	65
threonine	5	10
tryptophan	10	30
dextrin	35	45
citric acid	10	10
vitamin	10	10
total	120	200

Table 5

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	80.0	biotin	10.0
riboflavin	220.0	choline	8000.0
pantothenic acid	220.0	folic acid	2.5
nicotinic acid	1000.0	vitamin B ₁₂	0.3
vitamin B ₆	300.0	linoleic acid	10.0

EXAMPLE 4

Lysine, methionine, threonine, tryptophan, glycine, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryō Kogyō Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 6 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 10-fold with the use of a

proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 9 to 11 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 7 and adding dextrin thereto to give a total amount of 10 g.

Table 6

<u>Component</u>	<u>Composition 9</u> <u>(g)</u>	<u>Composition 10</u> <u>(g)</u>	<u>Composition 11</u> <u>(g)</u>
lysine	15	30	60
methionine	10	20	40
threonine	10	20	40
tryptophan	3	5	10
glycine	40	75	150
dextrin	102	230	480
citric acid	10	10	10
vitamin	10	10	10
total	200	400	800

Table 7

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	260.0	biotin	20.0
riboflavin	360.0	choline	4000.0
pantothenic acid	2000.0	folic acid	50.0
nicotinic acid	2200.0	vitamin B ₁₂	1.0
vitamin B ₆	600.0	linoleic acid	160.0

EXAMPLE 5

Lysine, methionine, threonine, tryptophan, glycine, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryō Kogyō Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 8 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 12 to 14 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 9 and adding dextrin thereto to give a total amount of 10 g.

Table 8

<u>Component</u>	<u>Composition 12 (g)</u>	<u>Composition 13 (g)</u>	<u>Composition 14 (g)</u>
lysine	30	60	120
methionine	20	40	80
threonine	20	40	80
tryptophan	5	10	20
glycine	75	150	300
dextrin	230	480	980
citric acid	10	10	10
vitamin	10	10	10
total	400	800	1600

Table 9

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	80.0	biotin	10.0
riboflavin	220.0	choline	7500.0
pantothenic acid	220.0	folic acid	2.5
nicotinic acid	1000.0	vitamin B ₁₂	0.3
vitamin B ₆	300.0	linoleic acid	100.0

EXAMPLE 6

Lysine, methionine, threonine, tryptophan, glycine, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 10 and the

resulting mixture was dissolved in 20 l of water. Thus, nourishing compositions 15 to 17 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 11 and adding dextrin thereto to give a total amount of 2 g.

Table 10

<u>Component</u>	<u>Composition 15 (g)</u>	<u>Composition 16 (g)</u>	<u>Composition 17 (g)</u>
lysine	4	8	16
methionine	2	4	8
threonine	3	6	12
tryptophan	1	2	4
glycine	2	4	8
dextrin	34	72	148
citric acid	2	2	2
vitamin	2	2	2
total	50	100	200

Table 11

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	1.0	biotin	0.2
riboflavin	3.75	choline	1250.0
pantothenic acid	12.0	folic acid	0.3
nicotinic acid	10.0	vitamin B ₁₂	0.02
vitamin B ₆	1.0		

EXAMPLE 7

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 12 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 2-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 18 to 20 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 13 and adding dextrin thereto to give a total amount of 1 g.

Table 12

<u>Component</u>	<u>Composition 18 (g)</u>	<u>Composition 19 (g)</u>	<u>Composition 20 (g)</u>
lysine	4.5	9.0	18.0
methionine	2.2	4.5	9.0
threonine	2.6	5.2	10.5
tryptophan	0.8	1.6	3.2
dextrin	2.9	2.7	7.3
citric acid	1.0	1.0	1.0
vitamin	1.0	1.0	1.0
total	15.0	25.0	50.0

Table 13

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	10.5	biotin	0.5
riboflavin	31.6	choline	4200.0
pantothenic acid	95.0	folic acid	3.2
nicotinic acid	130.2	vitamin B ₁₂	0.2
vitamin B ₆	15.8		

EXAMPLE 8

Lysine, methionine, threonine, tryptophan, glycine, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.) and citric acid were mixed together at each ratio as specified in Table 14 and the resulting mixture was dissolved in 10 l of water. The obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 21 to 23 for animals were obtained.

Table 14

<u>Component</u>	<u>Composition 21</u> <u>(g)</u>	<u>Composition 22</u> <u>(g)</u>	<u>Composition 23</u> <u>(g)</u>
lysine	10	13	15
methionine	5	6.5	8
threonine	7	8	9
tryptophan	2	2.5	3
glycine	20	25	30
dextrin	146	235	325
citric acid	10	10	10
total	200	300	400

EXAMPLE 9

Lysine, methionine, threonine, tryptophan, glycine, glucose, fructose, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryō Kōgyō Co.) and citric acid were mixed together at each ratio as specified in Table 15 and the resulting mixture was dissolved in 1 l of water. Thus nourishing compositions 24 to 27 for animals were obtained.

Table 15

<u>Component</u>	<u>Composition 24</u> <u>(g)</u>	<u>Composition 25</u> <u>(g)</u>	<u>Composition 26</u> <u>(g)</u>	<u>Composition 27</u> <u>(g)</u>
lysine	0.16	0.16	0.16	0.16
methionine	0.06	0.06	0.06	0.06
threonine	0.06	0.06	0.06	0.06
tryptophan	0.11	0.11	0.11	0.11
glycine	0.10	0.10	0.10	0.10
glucose	0.50	-	-	0.40
fructose	-	0.50	-	-
dextrin	-	-	0.50	-
citric acid	0.01	0.01	0.01	0.01

EXAMPLE 10

Lysine, methionine, threonine, tryptophan, sorbitol citric acid and vitamins were mixed together at each ratio as specified in Table 16 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 15-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 28 and 29 for animals were obtained. The vitamins used herein was a composition prepared by mixing the vitamins as specified in Table 3 above and adding sorbitol thereto in such a manner as to give a total amount of 10 g.

Table 16

<u>Component</u>	<u>Composition 28.</u> <u>(g)</u>	<u>Composition 29</u> <u>(g)</u>
lysine	30	60
methionine	15	30
threonine	15	30
tryptophan	5	10
sorbitol	10	20
citric acid	10	20
vitamin	5	10
total	90	180

EXAMPLE 11

Lysine, methionine, threonine, tryptophan, sorbitol, citric acid and vitamins were mixed together at each ratio as specified in Table 17 and the resulting mixture was dissolved in 10 l of water. The obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 30 and 31 for animals were obtained. The vitamins used herein was a composition prepared by mixing the vitamins as specified in Table 18 and adding sorbitol thereto to give a total amount of 10 g.

Table 17

<u>Component</u>	<u>Composition 30</u> <u>(g)</u>	<u>Composition 31</u> <u>(g)</u>
lysine	10	20
methionine	5	10
threonine	5	10
tryptophan	2.5	5
sorbitol	12.5	25
citric acid	2.5	5
vitamin	2.5	5
total	40	80

Table 18

<u>Compound</u>	<u>Content</u> <u>(mg)</u>	<u>Compound</u>	<u>Content</u> <u>(mg)</u>
thiamine	15.0	biotin	0.5
riboflavin	25.0	choline	1500.0
pantothenic acid	50.0	folic acid	2.5
nicotinic acid	50.0	vitamin B ₁₂	0.1
vitamin B ₆	10.0	linoleic acid	5.0

EXAMPLE 12

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.) citric acid and vitamins were mixed together at each ratio as specified in Table 19 and the resulting mixture was dissolved in 30 l of water. The

obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing compositions 32 and 33 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 5 above and adding dextrin thereto to give a total amount of 10 g.

Table 19

<u>Component</u>	<u>Composition 32 (g)</u>	<u>Composition 33 (g)</u>
lysine	20	30
methionine	30	65
threonine	5	10
tryptophan	10	30
dextrin	35	45
citric acid	10	10
vitamin	10	10
total	120	200

EXAMPLE 13

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 20 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 15-fold with the use of a proportional liquid dispenser (Dosatron, a product of

Dosatron International). Thus nourishing composition 34 for animals were obtained. The vitamins used herein was prepared by mixing the vitamins as specified in Table 3 above and adding dextrin thereto to give a total amount of 10 g.

Table 20

<u>Component</u>	<u>Composition 34 (g)</u>
lysine	60
methionine	30
threonine	30
tryptophan	10
dextrin	20
citric acid	20
vitamin	10
total	180

EXAMPLE 14

Lysine, methionine, threonine, tryptophan, dextrin (degree of polymerization: 15.1; NSD 300 manufactured by Nippon Shiryo Kogyo Co.), citric acid and vitamins were mixed together at each ratio as specified in Table 21 and the resulting mixture was dissolved in 20 l of water. The obtained solution was diluted 10-fold with the use of a proportional liquid dispenser (Dosatron, a product of Dosatron International). Thus nourishing composition 35 for animals was obtained. The vitamins used herein was prepared

by mixing the vitamins as specified in Table 22 and adding dextrin thereto to give a total amount of 10 g.

Table 21

<u>Component</u>	<u>Composition 35 (g)</u>
lysine	60
methionine	40
threonine	40
tryptophan	10
dextrin	20
citric acid	10
vitamin	10
total	200

Table 22

<u>Compound</u>	<u>Content (mg)</u>	<u>Compound</u>	<u>Content (mg)</u>
thiamine	260.0	biotin	20.0
riboflavin	360.0	choline	9000.0
pantothenic acid	2000.0	folic acid	50.0
nicotinic acid	2200.0	vitamin B ₁₂	1.0
vitamin B ₆	600.0	linoleic acid	160.0

TEST EXAMPLE 1

Broilers aged 56 days were divided into 4 groups each having 1,000 birds and fasted for 15 hours and 30 minutes before the shipment. During the fasting period, the nourishing compositions 1 to 3 obtained in Example 1 and

water (as a control) were given respectively to the groups.

The average body weight and the number of dead birds of each group were measured at the initiation of the test and after the completion thereof. Thus the body weight loss and mortality were calculated. Table 23 shows the results.

Table 23

	<u>Composition 1</u>	<u>Composition 2</u>	<u>Composition 3</u>	<u>Control</u>
Body weight before test (g)	2958	2983	2978	2977
Body weight after test (g)	2880	2926	2940	2882
Body weight loss (g)	78	57	38	95
Body weight loss (%)	2.63	1.91	1.27	3.19
Mortality (%)	2.0	1.2	0.8	3.2

As Table 23 shows, the nourishing compositions 1 to 3 for animals according to the present invention each suppressed the body weight loss and lowered the mortality, compared with the control.

TEST EXAMPLE 2

When the temperature was high in the summer season, broilers aged 35 days were divided into 4 groups each having 1,000 birds and fed with a common feed for 35 days. During this period, the nourishing compositions 4 to 6 obtained in Example 2 and water (as a control) were given to the birds.

The average body weight and the number of missing

(dead) birds were measured at the initiation of the test and after the completion thereof. Thus the raising rate [(number of birds before test - number of missing birds)/number of birds before test], the feed conversion rate (feed intake/body weight gain), the meat productivity [(average body weight/feed conversion rate) x 100] and the broiler production score (PS) were calculated. PS was calculated in accordance with the following formula.

$$PS = \frac{\text{body weight after test} \times \text{raising rate}}{\text{feed conversion rate} \times \text{test period (days)}} \times 100.$$

Table 24 shows the results.

Table 24

	<u>Composition 4</u>	<u>Composition 5</u>	<u>Composition 6</u>	<u>Control</u>
No. of missing birds	15	10	5	20
Raising rate (%)	98.5	99.0	99.5	98.0
Body weight before test (g)	663	665	638	668
Body weight after test (g)	2869	2892	2955	2837
Body weight gain (g)	2206	2227	2317	2169
Daily body weight gain (g)	63.0	63.6	66.2	62.0
Feed conversion rate (%)	2.34	2.30	2.29	2.38
Meat productivity (%)	123	126	129	119
PS	215	222	229	209

As Table 24 shows, the nourishing compositions 4 to 6 for animals according to the present invention each decreased the number of missing birds and elevated the raising rate, the feed conversion rate and PS, compared with the control.

TEST EXAMPLE 3

When the temperature was high in the summer season, layers aged 30 weeks were divided into 3 groups each having 1,000 birds and fed with a common feed for 8 weeks. During this period, the nourishing compositions 7 and 8 obtained in Example 3 and water (as a control) were given to the birds.

After the completion of the test, the normal egg laying rate, the individual egg weight, the daily egg weight (individual egg weight x normal egg laying rate), the feed intake, the feed conversion rate $[(\text{feed intake}/\text{daily egg weight}) \times 100]$ and the soft/broken egg rate were calculated. Also, regarding the egg qualities, the dense albumen height (the height of dense albumen at breaking) and the yolk diameter of each group were measured and thus the Haugt unit (HU) was calculated.

Tables 25 and 26 show the results.

Table 25

	<u>Composition 7</u>	<u>Composition 8</u>	<u>Control</u>
Normal egg laying rate (%)	87.8	88.8	87.3
Individual egg weight (g)	59.3	59.1	59.6
Daily egg weight (g)	52.1	52.5	52.0
Feed intake (g)	100.0	99.8	100.8
Feed conversion rate (%)	1.92	1.90	1.94
Soft/broken egg rate (%)	2.2	2.0	2.2

Table 26

	<u>Composition 7</u>	<u>Composition 8</u>	<u>Control</u>
Dense albumen height (mm)	5.58	6.28	5.52
Yolk diameter (cm)	4.10	4.16	4.09
HU	71.5	77.6	71.3

As the above Tables 25 and 26 show, the nourishing compositions 7 and 8 for animals according to the present invention each elevated the normal egg laying ratio and the daily egg weight and lowered the soft/broken egg ratio, compared with the control. As the results of the examination on the egg qualities, it was found out that the nourishing compositions 7 and 8 of the present invention each elevated the dense albumen height, the yolk diameter and HU.

TEST EXAMPLE 4

Large layer chicks aged 130 days were divided into 4 groups each having 1,000 birds and then transferred in accordance with the working schedule given in Fig. 1. As shown in Fig. 1, the nourishing compositions 9 to 11 for animals obtained in Example 4 and water as a control were administered to the birds with the use of nipple drinkers attached to gauges both in the brooding and egg-laying houses. The feeding was performed for about 12 hours (from the initiation of fasting to bird trapping/packing) in the brooding house and then for about 2 days from the initiation of feeding of water after transferring the chicks into the egg-laying house.

Table 27 shows the body weights before and after the transfer, the body weight loss and the time (days) required for the recovery of the body weight.

Table 27

	<u>Composition 9</u>	<u>Composition 10</u>	<u>Composition 11</u>	<u>Control</u>
Body weight before transfer (g)	1304	1299	1302	1300
Body weight after transfer (g)	1184	1199	1220	1157
Body weight loss (g)	120	100	92	133
Body weight loss (%)	9.2	7.7	7.1	10.2
Recovery time (days)	4	3	3	5

As Table 27 shows, the nourishing compositions 9 to 11 for animals according to the present invention each suppressed the body weight loss and shortened the time required for the recovery of the body weight, compared with the control.

TEST EXAMPLE 5

Layers aged 12 months were divided into 4 groups each having 1,000 birds. After fasting for 15 days in order to force the bird to moult, water was also not given for 1 day to stress them. Five to 7 days after the initiation of fasting, small feathers fell out. After 10 to 13 days, feathers in the main wings fell out too. After 2 weeks from the initiation of fasting, feeding was initiated. For 3 weeks following the initiation of feeding, the nourishing compositions 12 to 14 for animals obtained in Example 5 and water (as a control) were given to the birds.

Table 28 shows the time (days) required for starting normal egg-laying again after the initiation of feeding (egg-laying starting day after resumption of feeding), the time (days) required for an egg-laying rate achieving 50% of that before moulting (day achieving 50% egg-laying) and the egg-laying peak (the maximum egg-laying rate) of each group.

Table 28

	<u>Composi- tion 12</u>	<u>Composi- tion 13</u>	<u>Composi- tion 14</u>	<u>Control</u>
Egg-laying starting day after resumption of feeding (days)	10	10	9	11
Day achieving 50% egg-laying rate	31	30	30	33
Egg-laying peak (%)	83	85	85	82

As Table 28 shows, the nourishing compositions 12 to 14 for animals according to the present invention each shortened the time required for starting egg-laying again and the time required for an egg-laying rate achieving 50% of that before moulting and improved the egg-laying peak, compared with the control.

TEST EXAMPLE 6

Mother pigs with the third farrowing were divided into 4 groups each having 5 animals. From 5 to 7 days before farrowing to 3 weeks thereafter, the nourishing compositions 15 to 17 for animals obtained in Example 6 and water (as a control) were given respectively to the groups.

The days with puerperal fever of 41 °C or above, the maximum body temperature, the number of piglings, the number of piglings with diarrhea, the number of weanlings and the average body weight at weaning of each group were measured. Table 29 shows the results.

Table 29

	<u>Composi- tion 15</u>	<u>Composi- tion 16</u>	<u>Composi- tion 17</u>	<u>Control</u>
Days with puerperal fever of 41°C or above	2	0	0	3
Maximum body temp. (°C)	41.7	40.7	40.5	42.2
No. of piglings	53	52	54	54
No. of piglings with diarrhea	9	6	4	12
No. of weanlings	47	48	50	47
Average body weight at weaning (kg)	6.8	7.0	7.2	6.4

As Table 29 shows, the nourishing compositions 15 to 17 for animals according to the present invention each improved the days with puerperal fever of 41°C or above, the maximum body temperature, the number of piglings, the number of piglings with diarrhea, the number of weanlings and the average body weight at weaning, compared with the control.

TEST EXAMPLE 7

Piglings farrowed by mother pigs of the third farrowing were divided into 4 groups each having 30 animals. For 40 days starting from 10 days before weaning to 30 days after weaning, the nourishing compositions 18 to 20 for animals obtained in Example 7 and water (as a control) were given respectively to the groups.

The body weight at weaning, the body weight 30 days after weaning, the body weight gain and the average feed

intake of each group were measured and the feed conversion rate was calculated. Table 30 shows the results.

Table 23

	<u>Composi- tion 18</u>	<u>Composi- tion 19</u>	<u>Composi- tion 20</u>	<u>Control</u>
Body weight at weaning (kg)	7.1	7.1	7.1	7.1
Body weight 30 days after weaning (kg)	25.6	26.2	26.4	24.9
Body weight gain (kg)	18.5	19.1	19.3	17.8
Average feed intake (kg)	31.6	32.3	32.4	30.6
Feed conversion rate (%)	1.71	1.69	1.68	1.72

As Table 30 shows, groups fed with the nourishing compositions 18 to 20 for animals of the present invention showed each a large body weight gain after weaning and a decrease in the feed conversion rate, compared with the control.

TEST EXAMPLE 8

Hogs aged 105 days and weighing 110 kg were divided into 4 groups each having 20 animals and fasted for 24 hours before the shipment. Simultaneously with the initiation of fasting, the nourishing compositions 21 to 23 for animals obtained in the above Example 8 and water (as a control) were given respectively to the groups.

Each animal was weighed before and after the test to determine the average body weight of animals of each group

and the body weight loss was calculated. Table 31 shows the results.

Table 31

	<u>Composi- tion 21</u>	<u>Composi- tion 22</u>	<u>Composi- tion 23</u>	<u>Control</u>
Body weight before test (kg)	110.0	110.0	110.0	110.0
Body weight after test (kg)	108.6	108.8	108.0	108.2
Body weight loss	1.4	1.2	1.0	1.8
Body weight loss (%)	1.3	1.1	0.9	1.6

As Table 31 shows, the groups fed with the nourishing compositions 21 to 23 for animals of the present invention showed each a decreased body weight loss, compared with the control.

TEST EXAMPLE 9

The nourishing compositions 24 to 27 for animals obtained in Example 9 were allowed to stand at 40 °C under a humidity of 75 % and the conditions of each solution were observed. Table 32 shows the results.

Table 32

<u>Time</u>	<u>Composi- tion 24</u>	<u>Composi- tion 25</u>	<u>Composi- tion 26</u>	<u>Composi- tion 27</u>
Day 2	Turning to yellow with forming small masses	Becoming powdery	No change	Turning to yellow with forming large masses
Day 5	Enlarging yellow masses	Becoming coarse	No change	Forming a brown mass as a whole
Day 7	The same as day 5	The same as day 5	No change	The same as day 5

As Table 32 shows, the nourishing compositions 24, 25 and 27 containing glucose or fructose as a sugar underwent browning. In contrast, the nourishing composition 26 of the present invention containing dextrin as a sugar suffered from no browning even after 7 days.

TEST EXAMPLE 10

Broilers aged 28 days were divided into 3 groups each having 1,000 birds and fed with a common feed for 5 days. During this period, the nourishing compositions 28 and 29 obtained in Example 10 and water (as a control) were given to the birds.

The average body weight and the number of missing dead birds were measured at the initiation of the test (at the age of 28-days) and at the age of 35 and 42 days. Thus the raising rate was calculated in the same manner as in Test Example 2.

Table 33 shows the results.

Table 33

	<u>Composition 28</u>	<u>Composition 29</u>	<u>Control</u>
No. of dead birds	25	20	36
Raising rate (%)	97.5	98.0	96.4
Body weight at the age of 28 days (g)	972	967	982
Body weight at the age of 35 days (g)	1400	1402	1394
Body weight at the age of 42 days (g)	1798	1828	1768
Body weight gain (g)	826	861	786
Daily body weight gain (g)	59.0	61.5	56.1

As Table 33 shows, the nourishing compositions 28 and 29 for animals according to the present invention each decreased the number of dead birds and elevated the raising rate, compared with the control.

TEST EXAMPLE 11

Healthy new-born chicks hatched from hatchery eggs laid by breeding hens aged 41 weeks and 44 weeks were divided into 3 groups each having 1,000 birds and fed with the nourishing compositions 30 to 31 obtained in Example 11 and water (as a control) were given to the birds for 7 days.

The average body weight and the number of dead birds were measured at the initiation of the test (at the hatch)

and at the age of 7 and 14 days. Thus the raising rate was calculated in the same manner as in Test Example 2.

Table 34 shows the results.

Table 34

	<u>Composition 30</u>	<u>Composition 31</u>	<u>Control</u>
No. of dead birds	10	8	16
Raising rate (%)	99.0	99.2	98.4
Body weight at the hatch (g)	41	40	42
Body weight at the age of 7 days (g)	144	146	142
Body weight at the age of 14 days (g)	412	420	404
Body weight gain (g)	371	380	362
Daily body weight gain (g)	28.5	29.2	27.8

As Table 34 shows, the nourishing composition 30 and 31 for animals according to the present invention each decreased the number of dead birds and elevated the raising rate, compared with the control.

TEST EXAMPLE 12

When the temperature was high in the summer season, breeding hens aged 27 weeks were divided into 3 groups each having 1,000 birds and fed with a common feed for 6 weeks.

During this period, the nourishing compositions 32 and 33 obtained in Example 12 and water (as a control) were given to the birds.

After the completion of the test, the normal egg laying rate, the individual egg weight, the daily egg weight, the feed intake, the feed conversion rate and the soft/broken egg rate were calculated in the same manner as in Test Example 3.

Tables 35 shows the results.

Table 35

	<u>Composition 32</u>	<u>Composition 33</u>	<u>Control</u>
Normal egg laying rate (%)	87.8	88.8	87.3
Individual egg weight (g)	59.3	59.1	59.6
Daily egg weight (g)	52.1	52.5	52.0
Feed intake (g)	100.0	99.8	100.8
Feed conversion rate (%)	1.92	1.90	1.94
Soft/broken egg rate (%)	2.2	2.0	2.2

As Table 35 shows, the nourishing compositions 32 and 33 for animals according to the present invention each elevated the normal egg laying ratio and the daily egg weight and lowered the soft/broken egg ratio, compared with the

control.

TEST EXAMPLE 13

Layers aged 9 weeks were divided into 2 groups each having 1,000 birds and fed with a common feed for 3 days from the day of debeaking. During this period, the nourishing composition 34 obtained in Example 13 and water (as a control) were given to the birds.

The average body weight and the number of dead birds of each group measured at the initiation of the test and at the age of 10, 11 and 12 weeks. Thus the raising rate was calculated in the same manner as in Test Example 2.

Table 36 shows the results.

Table 36

	<u>Composition 34</u>	<u>Control</u>
No. of dead birds	7	12
Raising rate (%)	99.3	98.8
Body weight at the age of 9 weeks (g)	694	710
Body weight at the age of 10 weeks (g)	728	719
Body weight at the age of 11 weeks (g)	812	771
Body weight at the age of 12 weeks (g)	919	867
Body weight gain (g)	225	157

As Table 36 shows, the nourishing composition 34 for animals according to the present invention decreased the number of dead birds and elevated the raising rate, compared with the control.

TEST EXAMPLE 14

Layers aged 16 weeks were divided into 2 groups each having 1,000 birds and infectious coryza vaccine was given to each bird by intramuscular injection in a predetermined dose. For 3 days from the vaccination, the nourishing composition 35 obtained in Example 14 and water (as a control) were given to the birds.

The average body weight and the number of dead birds were measured at the initiation of the test (at the age of 16 weeks) and at the age of 17 and 18 weeks. Thus the raising rate was calculated in the same manner as in Test Example 2.

Table 37 shows the results.

Table 37

	<u>Composition 35</u>	<u>Control</u>
No. of dead birds	4	6
Raising rate (%)	99.6	99.4
Body weight at the age of 16 weeks (g)	1215	1220
Body weight at the age of 17 weeks (g)	1240	1235
Body weight at the age of 18 weeks (g)	1320	1300
Body weight gain (g)	105	80

As Table 37 shows, the nourishing composition 35 for animals according to the present invention decreased the number of dead birds and elevated the raising rate, compared with the control.

The present invention provides a nourishing composition for animals which can improve the nutritional conditions etc. of animals when administered together with drinking water.

While the invention has been described in detail and with reference to specific examples thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A nourishing composition for animals which comprises an aqueous solution containing (1) at least one of dextrin and sugar alcohols, and (2) amino acids.

2. A nourishing composition for animals as claimed in Claim 1 wherein said amino acid is at least one selected from the group consisting of lysine, methionine, tryptophan, threonine, alanine, glycine and glutamic acid.

3. A nourishing composition for animals as claimed in Claim 1 wherein said sugar alcohol is at least one selected from the group consisting of lactitol, isomaltitol, maltitol, mannitol, sorbitol, xylitol and erythritol.

4. A nourishing composition for animals which comprises an aqueous solution containing (1) at least one of dextrin and sugar alcohols, (2) amino acids and (3) organic acids.

5. A nourishing composition for animals as claimed in Claim 4 wherein said organic acid is selected from the group consisting of citric acid, malic acid and succinic acid.

6. A nourishing composition for animals as claimed in Claim 1 in which the amount of dextrin and sugar alcohols ranges from 0.0005 to 10.0 % by weight and the amount of amino acids ranges from 0.001 to 5.0 % by weight.

7. A nourishing composition for animals as claimed in Claim 1 in which the amount of dextrin and sugar alcohols ranges from 0.001 to 0.8 % by weight and the amount of amino

2139028

acid ranges from 0.003 to 2.0 % by weight.

Fig. 1

