

**(12) PATENT
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199870975 B2
(10) Patent No. 741439**

(54) Title
Formulation and method for treating neoplasms by inhalation

(51)⁷ International Patent Classification(s)
A61K 031/00

(21) Application No: **199870975** (22) Application Date: **1997.12.30**

(87) WIPO No: **WO98/29110**

(30) Priority Data

(31) Number **60/033789** (32) Date **1996.12.30** (33) Country **US**

(43) Publication Date : **1998.07.31**
(43) Publication Journal Date : **1998.09.17**
(44) Accepted Journal Date : **2001.11.29**

(71) Applicant(s)
Battelle Memorial Institute

(72) Inventor(s)
Michael E. Placke; Anthony R. Omondi; Michael J. Booker; John E. Frye; Praful K. Shah; Douglas R. Flanagan Jr.; Maureen D. Donovan

(74) Agent/Attorney
PHILLIPS ORMONDE and FITZPATRICK,367 Collins Street,MELBOURNE VIC 3000

(56) Related Art
**US 5565478
EP 709090
WO 94/10175**



IN

(51) International Patent Classification 6 : A61K 31/00		A2	(11) International Publication Number: WO 98/29110 (43) International Publication Date: 9 July 1998 (09.07.98)
(21) International Application Number: PCT/US97/24289		(74) Agents: GOLDSTEIN, Steven, J. et al.; Frost & Jacobs, 2500 PNC Center, 201 East Fifth Street, Cincinnati, OH 45202 (US).	
(22) International Filing Date: 30 December 1997 (30.12.97)		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 60/033,789 30 December 1996 (30.12.96) US		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(71) Applicant (for all designated States except US): BATTELLE MEMORIAL INSTITUTE [US/US]; 505 King Avenue, Columbus, OH 43201-2693 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): PLACKE, Michael, E. [US/US]; 1473 Inglis Avenue, Columbus, OH 43212 (US). OMONDI, Anthony, R. [US/US]; 4515 Ravine Drive, Westerville, OH 43081 (US). BOOKER, Michael, J. [US/US]; 8381 Africa Road, Westerville, OH 43082 (US). FRYE, John, E. [US/US]; 191 Walnut Street, Groveport, OH 43125-1233 (US). SHAH, Praful, K. [US/US]; 4750 Coolbrook Drive, Hilliard, OH 43026 (US). FLANAGAN, Douglas, R., Jr. [US/US]; 1258 Dover Street, Iowa City, IA 52240-2829 (US). DONOVAN, Maureen, D. [US/US]; 2748 Lakeview Drive, N.E., Solon, IA 52333-9563 (US).			

(54) Title: FORMULATION AND METHOD FOR TREATING NEOPLASMS BY INHALATION

(57) Abstract

A formulation, method, and apparatus for treating neoplasms such as cancer by administering a pharmaceutically effective amount of highly toxic composition by inhalation, wherein the composition is a non-encapsulated antineoplastic drug.

FORMULATION AND METHOD FOR TREATING NEOPLASMS BY INHALATION

5 This application claims the benefit of U.S. Provisional Application No. 60/033,789 filed on December 30, 1996.

FIELD OF THE INVENTION

The invention deals with formulations and methods useful for treating 10 neoplasms, particularly neoplasms of the respiratory tract (e.g. lung cancer and cancers of the head and neck), by pulmonary administration of highly toxic or vesicating anticancer drugs. Additionally, several new formulations and methods for treating neoplasms using antineoplastic drugs that are nonvesicants are also disclosed.

15

BACKGROUND OF THE INVENTION

Cancer is one of the leading causes of death worldwide. Lung cancer in particular, is among the top 3 most prevalent cancers and has a very poor survival rate (about 13% five-year survival rate). Despite the availability of 20 many cancer drugs it has been difficult and, in the case of some cancer types, almost impossible to improve cure rates or survival. There are many reasons for this lack of success but one reason is the inability to deliver adequate amounts of the drugs to the tumor without causing debilitating and life-threatening toxicities in the patient. Indeed, most chemotherapeutic drugs 25 used to treat cancer are highly toxic to both normal and tumor tissues.

It is customary in the treatment of cancer to administer the drugs by the intravenous route, which exposes the entire body to the drug. Doses are selected that destroy tumor cells, but these doses also destroy normal cells. As a result, the patient usually experiences severe toxic side effects. For 30 example, severe myelosuppression may result which compromises the ability of the patient to resist infection and allows spread of the tumor. There are other life-threatening effects such as hepatotoxicity, renal toxicity, pulmonary

toxicity, cardiotoxicity, neurotoxicity, and gastrointestinal toxicity caused by anticancer drugs. The anticancer drugs also cause other effects such as alopecia, stomatitis, and cystitis that may not be life threatening, but are serious enough to affect a patient's quality of life. Moreover, it is important to 5 note that these toxicities are not associated to the same extent with all anticancer drugs but are all due to systemic delivery of the drug.

Although myelosuppression is commonly associated with most anticancer drugs, because of differences in the mechanisms by which the various anticancer drugs act or in the ways they are distributed in the body, 10 metabolized and excreted from the body, each drug presents a somewhat different toxicity profile, both quantitatively and qualitatively. For example, anthracyclines such as doxorubicin, epirubicin and idarubicin are known to cause severe cardiotoxicity. Doxorubicin, additionally, is known to cause severe progressive necrosis of tissues when extravasated. Cisplatin therapy 15 is known to cause renal toxicity; vincristine causes neurotoxicity, bleomycin and mitomycin cause pulmonary toxicity, cyclophosphamide causes cystitis; and 5-fluorouracil causes cerebral disjunction (see Cancer Chemotherapy: Principles and Practice, BA Shabner and J.M. Collings, eds. J.B. Lippincott Co., Philadelphia, 1990)

20 The differences in mechanisms of action and pharmacokinetic properties determine, in part, the efficacy of the various anticancer drugs against different tumor types, which exhibit various biological behaviors.

Some attempts have been made to deliver anticancer drugs directly to the tumor or to the region of the tumor to minimize exposure of normal 25 tissues to the drug. This regional therapy, for example has been used to treat liver cancer by delivering drugs directly into the hepatic artery so that the full dose goes to the liver while reducing the amount that goes to the rest of the body. For the treatment of urinary bladder cancer, anticancer drugs are instilled directly into the bladder through the urethra, allowed to remain in 30 contact with the tumor for a period of time and then voided. Other examples of regional therapy include the delivery of anticancer drugs into the peritoneal cavity to treat cancer that has developed in or metastasized to this location.

Other methods of targeting anticancer drugs involve the attachment of the drugs to antibodies that seek out and deliver the drug directly to the cancer cells.

In 1968 Shevchenko, I.T., (Neoplasma 15, 4, 1968) pp.419-426

5 reported on the treatment of advanced bronchial cancer using a combination of inhalation of chemotherapeutic agents, radiotherapy, and oxygen inhalation. The reported chemotherapeutic agents were benzotaph, thiophosphamid, cyclophosphamid and endoxan that were applied as an aerosol by means of an inhaler. For 58 treated patients the combination of three
10 treatments showed tumor disappearance in 8 cases while in 6 the size of the tumor diminished considerably. The study did not include a control group.

In 1970, Sugawa, I. (Ochanoizu Med. J.; Vol. 18; No.3; (1970), pp.103-114, reported on tests using mitomycin-C in the treatment of metastatic lung cancer. One of four patients treated reportedly showed some
15 improvement. Inhalation of mitomycin-C also appeared to reduce tumor growth in IV-inoculated tumors in rabbits; results appeared to be more inconclusive in rats. Tests were conducted to determine the toxic effects to the respiratory tract following intrabronchial infusions of several drugs. The drugs were given to healthy animals and included: thiotepa (rats), Toyomycin
20 (chromomycin A3) (rats,), endoxan (cyclophosphamide) (rats and rabbits), 5-fluorourcil (rats and rabbits), mitomycin-C (rats, rabbits, and dogs). The results of these tests showed that: 5-FU and cyclophosphamide resulted in only mild inflammation; thiotepa produced bronchial obstruction; chromomycin A3 and mitomycin-C produced the most severe results. Toxic
25 effects of mitomycin-C and chromomycin A3 were studied in rabbits and dogs.

In 1983, Tatsumura et al (Jap. J. Cancer Cln., Vol. 29, pp. 765-770) reported that the anticancer drug, fluorouracil (5-FU, MW=130) was effective for the treatment of lung cancer in a small group of human patients when administered directly to the lung by aerosolization. They referred to this as
30 nebulization chemotherapy. It was also noted by Tatsumura et al (1993) (Br. J. Cancer, Vol. 68(6): pp.1146-1149) that the 5-FU did not cause toxicity to the lung. This finding was not totally unexpected because 5-FU has a very

low molecular weight and does not bind tightly to proteins. Therefore, it passes through the lung rapidly lessening the opportunity to cause local toxicity. Moreover 5-FU is considered to be one of the least toxic anticancer drugs when applied directly to tissue. Indeed, 5-FU is used as a topical drug
5 for the treatment of actinic keratosis for which it is applied liberally, twice daily, to lesions on the face. This therapy may continue for up to four weeks. Also, because 5-FU is poorly absorbed from the gastrointestinal tract, there is little concern about the amount of drug that may be inadvertently swallowed and gain access to the blood stream from the gut. It is well known that a
10 large percentage of aerosolized drug intended for the lung is swallowed.

Another report includes the use of β -cytosine arabinoside (Ara-C, cytarabine, MW=243) administered via intratracheal delivery to the respiratory system of rats. Liposome encapsulated and free Ara-C were instilled intratracheally to the rats as a bolus. The encapsulated Ara-C
15 persisted for a long time in the lung while the free Ara-C which is not highly protein bound was rapidly cleared from the lung. The free Ara-C rapidly diffused across the lung mucosa and entered the systemic circulation. The paper suggests that liposome encapsulation of drugs may be a way to produce local pharmacologic effect within the lung without producing adverse
20 side effects in other tissues. However, bolus administration results in multifocal concentrated pockets of drug. See the articles by H.N. MacCullough et al, JNCI, Vol. 63, No. 3, Sept., pp.727-731 (1979) and R.L. Juliano et al, J. Ph. & Exp. Ther., Vol. 214, No.2, pp.381-387 (1980).

An additional report includes the use of cisplatin (MW=300) for
25 inhalation chemotherapy in mice that had been implanted with FM3A cells (murine mammary tumor cells) in the air passages. The cisplatin exposed inhalation group were reported to have statistically smaller lung tumor sizes and survived longer than the untreated control group. See A. Kinoshita, "Investigation of Cisplatin Inhalation Chemotherapy Effects on Mice after Air
30 Passage Implantation of FM3A Cells", J. Jap. Soc. Cancer Ther. 28(4): pp. 705-715 (1993).

In US patent 5,531,219 to Rosenberg, the patent disclosure suggests the use of doxorubicin, 5-FU, vinblastine sulfate, or methotrexate in combination with pulmonary infused liquid fluorocarbons. The patient is suggested to be positioned so that the tumor affected area is at a 5 gravitational low point so that liquid perfluorocarbon having relatively low vapor pressure will pool selectively around the area with the drug then perfused in the pool of liquid perfluorocarbon. The present invention avoids the problems with positioning of the patient and further does not require the liquid fluorocarbons used by Rosenberg.

10 In US patent 5,439,686 to Desai et al there are disclosed compositions where a pharmaceutically active agent is enclosed within a polymeric shell for administration to a patient. One of the routes of administration listed as possible for the compositions of the invention is inhalational. Among the listed pharmaceutically active agents potentially useful in the invention are 15 anticancer agents such as paclitaxel and doxorubicin. No tests using the inhalational route of administration appear to have been made.

Although several antineoplastic drugs have been administered to animals and to humans, for treatment of tumors in the lungs and respiratory system, the differences in the mechanism of action, and toxicity profiles 20 among the broad classes of anticancer drugs, and the heretofore known characterizations have made it impossible to predict whether a particular anticancer drug will be efficacious or toxic based upon previous inhalation results with a different drug of a different type. Further, previous reports used very imprecise means of delivering drugs and were not consistent in 25 delivering measured doses of drugs in an evenly distributed manner to the entire respiratory tract. The present invention provides means for predicting and selecting drugs including the highly toxic chemotherapeutic compounds, amenable for inhalation therapy of neoplastic disease and methods for actually distributing specific measured doses to pre-selected regions of the 30 respiratory tract.

It has now been demonstrated by the applicants that anticancer cytotoxic drugs of multiple classes such as anthracyclines (doxorubicin),

antimicrotubule agents such as the vinca alkaloids (vincristine), and taxanes such as paclitaxel can be given directly by inhalation without causing severe toxicity to the lung or other body organs. This finding is surprising, because it is well known among those who administer cytotoxins such as doxorubicin to patients, that this drug causes severe ulceration of the skin and underlying tissues if allowed to be delivered outside of a vein. After extravasation the drug continues to affect the tissues to such an extent that amputation of limbs in which the extravasation has occurred has been required. So severe is this toxicity that the prescribing information for doxorubicin (and some other similar vesicating drugs) in the Physicians Desk Reference contains a "Box Warning" regarding this danger. The present invention, therefore, provides an effective way to administer chemotherapeutic agents, including highly toxic agents such as doxorubicin, while minimizing the major side effects described above.

The above discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.



BRIEF DESCRIPTION OF THE INVENTION

Broadly, one embodiment of the invention includes a formulation for treating a patient for a neoplasm by inhalation comprising: a safe and effective amount of a vesicant and a pharmaceutically acceptable carrier, preferably the vesicant does not exhibit substantial pulmonary toxicity. In one aspect of the embodiment the vesicant is typically a moderate vesicant such as paclitaxel or carboplatin. A description of such a moderate vesicant would include a non-encapsulated anticancer drug, wherein when 0.2 ml of the drug is injected intradermally to rats, at the clinical concentration for parenteral use in humans: (a) a lesion results that is at least 20 mm² in area fourteen days after the intradermal injection; and (b) at least 50% of the tested rats have this size of lesion. Other aspects of this broad embodiment typically include a vesicant that is a severe vesicant such as doxorubicin, vincristine, and vinorelbine. The neoplasm to be treated is typically a pulmonary neoplasm, a neoplasm of the head and neck, or other systemic neoplasm. The drug may be in the form of a liquid, a powder, a liquid aerosol, or a powdered aerosol. Typically the patient is a mammal such as a



domestic animal or a human. In other aspects the embodiment includes formulations of drugs such as etoposide and a carrier such as DMA. Typically the severe vesicant is an anthracycline such as epirubicin, daunorubicin, methoxymorpholinodoxorubicin, cyanomorpholinyl doxorubicin, doxorubicin, 5 or idarubicin; or a vinca alkaloid such as vincristine, vinorelbine, vinorelbine, vindesine, or vinblastine. In other formulations the drug is typically mechlorethamine, mithramycin, dactinomycin, bisantrene, or amsacrine. Typically the formulation may include a taxane such as paclitaxel, its derivatives and the like. Typical animal and human doses are provided in the 10 tables and text below.

A further broad embodiment of the invention includes a formulation for treating a patient having a neoplasm by inhalation comprising: a safe and effective amount of a non-encapsulated antineoplastic drug having a molecular weight above 350, that does not exhibit substantial pulmonary 15 toxicity; and an effective amount of a pharmaceutically acceptable carrier. The neoplasm treated with the formulation is typically a pulmonary neoplasm, a neoplasm of the head and neck, or a systemic neoplasm. The drug used in the formulation is in the form of a liquid, a powder, a liquid aerosol, or a powdered aerosol. Typically the drug has a protein binding affinity of 25% or 20 50% or more. Further the drug can typically have a higher molecular weights such as above 400, 450, or 500 daltons. Typical animal and human doses are provided in the tables and text below.

In a yet further embodiment of the invention, there is disclosed a formulation for treating a patient for a neoplasm by inhalation comprising: a 25 safe and effective amount of a taxane in an effective amount of vehicle comprising polyethyleneglycol (PEG) and an alcohol. Typically the formulation will also contain an acid, where the acid present in amount effective to stabilize the taxane. Typically the alcohol is ethanol, and the acid is an inorganic acid such as HCl, or an organic acid such as citric acid and the like. 30 In some typical formulations the taxane is paclitaxel and the formulation contains about 8% to 40% polyethyleneglycol, about 90% to 60% alcohol,

and about 0.01% to 2% acid. Typical animal and human doses are provided in the table and text below.

Another embodiment provides for formulations for treating a patient for a neoplasm by inhalation comprising: a safe and effective amount of a drug selected from the group consisting of carmustine, dacarbazine, melphalan, mercaptopurine, mitoxantrone, esorubicin, teniposide, aclacinomycin, plicamycin, streptozocin, and menogaril; and a safe and effective amount of a pharmaceutically effective carrier, wherein the drugs do not exhibit substantial pulmonary toxicity.

10 A yet further embodiment provides for a formulation for treating a patient for a neoplasm by inhalation comprising: a safe and effective amount of a drug selected from the group consisting of estramustine phosphate, geldanamycin, bryostatin, suramin, carboxyamido-triazoles; onconase, and SU101 and its active metabolite SU20; and a safe and effective amount of a 15 pharmaceutically effective carrier, wherein the drugs do not exhibit substantial pulmonary toxicity.

20 A still further embodiment provides for a formulation for treating a patient for a neoplasm by inhalation comprising: a safe and effective amount of etoposide and an effective amount of a DMA carrier. Typical animal and human doses are provided in the tables and text below.

Another embodiment includes a formulation for treating a patient for a neoplasm by inhalation comprising: a safe and effective amount of a microsuspension of 9-aminocamptothecin in an aqueous carrier. Typical animal and human doses are provided in the tables and text below.

25 A further broad embodiment of the invention includes a formulation for treating a patient having a neoplasm comprising: administering to the patient by inhalation, (1) an effective amount of a highly toxic antineoplastic drug; and (2) an effective amount of a chemoprotectant, wherein the chemoprotectant reduces or eliminates toxic effects in the patient that are a 30 result of administering the highly toxic antineoplastic drug. Typically the chemoprotectant reduces or eliminates systemic toxicity in the patient, and/or reduces or eliminates respiratory tract toxicity in the patient.

Typically the formulation includes a chemoprotectant such as dextrazoxane (ICRF-187), mesna (ORG-2766), ethiofos (WR2721), or a mixture thereof. The chemoprotectant may be administered before, after, or during the administration of the antineoplastic drug. The antineoplastic drug used with 5 the chemoprotectant may be a nonvesicant, moderate vesicant, or a severe vesicant. Typical among the drugs with which the chemoprotectant is useful are bleomycin, doxorubicin, and mitomycin-C.

The invention also typically includes a method for treating a patient having a neoplasm comprising: administering to the patient by inhalation, 10 (1) an effective amount of a highly toxic antineoplastic drug; and (2) an effective amount of a chemoprotectant, wherein the chemoprotectant reduces or eliminates toxic effects in the patient that are a result of administering the highly toxic antineoplastic drug. Typically the chemoprotectant reduces or eliminates systemic toxicity in the patient and/or reduces or eliminates 15 respiratory tract toxicity in the patient. Chemoprotectants can typically be dextrazoxane (ICRF-187), mesna (ORG-2766), ethiofos (WR2721), or a mixture thereof. The chemoprotectant may be administered before, after, or during the administration of the antineoplastic drug. Typically the antineoplastic drug is a nonvesicant, a moderate vesicant, or a severe 20 vesicant. Typically the antineoplastic drug comprises bleomycin, doxorubicin, or mitomycin-C.

An additional embodiment of the invention includes a method for treating a patient having a neoplasm comprising: administering a safe and effective amount of a non-encapsulated antineoplastic drug to the patient by 25 inhalation, the drug selected from the group consisting of antineoplastic drugs wherein when 0.2 ml of the drug is injected intradermally to rats, at the clinical concentration for IV use in humans: (a) at least one lesion per rat results which is greater than 20 mm² in area fourteen days after the intradermal injection; and (b) at least 50% of the tested rats have these 30 lesions. In some typical embodiments when the drug is doxorubicin or vinblastine sulfate, the drug is inhaled in the absence of perfluorocarbon. Typical diseases treated include a neoplasm such as a pulmonary neoplasm, a

neoplasm of the head and neck, or other systemic neoplasm. The drug may typically be inhaled as inhaled as a liquid aerosol or as a powdered aerosol. Mammal animals and humans are typical patients treated with the method. The drug may typically be selected from the group consisting of doxorubicin, 5 daunorubicin, methoxymorpholino-doxorubicin, epirubicin, cyanomorpholiny doxorubicin, and idarubicin. When the drug is a vinca alkaloid it is typically selected from the group consisting of vincristine, vinorelbine, vindesine, and vinblastine. Other useful drugs typically include the alkylating agents mechlorethamine, mithramycin and dactinomycin. Still additional useful drugs 10 typically include bisantrene and amsacrine. The drug can typically be a taxane such as doxitaxel or paclitaxel.

Another embodiment of the invention includes a method for treating a patient having a neoplasm comprising: administering an effective amount of a highly toxic non-encapsulated antineoplastic drug to a patient by inhalation, 15 wherein the molecular weight of the drug is above 350, and the drug has no substantial pulmonary toxicity. Typically the neoplasm is a pulmonary neoplasm, a neoplasm of the head and neck, or a systemic neoplasm. The drug may be inhaled as a liquid aerosol or as a powdered aerosol. Typically the drug has a protein binding affinity of 25% , 50% or more. 20 In one aspect the drug is typically selected from the group comprising doxorubicin, epirubicin, daunorubicin, methoxymorpholinodoxorubicin, cyanomorpholiny doxorubicin, and idarubicin. If the drug is doxorubicin or vinca alkaloid it may be typically be administered without the presence of a perfluorocarbon. Typically the vinca alkaloid is selected from the group 25 consisting of vincristine, vinorelbine, vindesine, and vinblastine. Typical alkylating agent type drugs include mechlorethamine, mithramycin, dactinomycin. Other topoisomerase II inhibitors include bisantrene or amsacrine.

An additional embodiment includes a method for treating a patient for 30 a neoplasm by the steps of administering an effective amount of an antineoplastic drug to the patient by inhalation; and administering a pharmaceutically effective amount of the same and/or different antineoplastic

drug to the patient parenterally. The patient may be treated with one or more adjunct therapies including radiotherapy, immunotherapy, gene therapy, chemoprotective drug therapy.

A further embodiment includes a method for treating a patient for a neoplasm including the steps of administering an effective amount of an antineoplastic drug to the patient by inhalation; and administering an effective amount of the same and/or different antineoplastic drug to the patient by isolated organ perfusion. The patient may be treated by one or more adjunct therapies including radiotherapy, immunotherapy, gene therapy, and chemoprotective drug therapy.

An further embodiment includes a method for treating a patient for a pulmonary neoplasm by the steps of (1) selecting one or more antineoplastic drugs efficacious in treating the neoplasm and having a residence time in the pulmonary mucosa sufficient to be efficacious in the treatment of the pulmonary neoplasm; and (2) administering the drug(s) to the patient by inhalation in a non-encapsulated form. Typically when 0.2 ml of at least one of the drugs is injected intradermally to rats, at the clinical concentration for parenteral use in humans: a lesion results which is greater than 20 mm² in area fourteen days after the intradermal injection; and B. at least 50% of the tested rats have these lesions. Typically the molecular weight of at least one of the selected drugs is above 350.

A still further embodiment includes a method of use including the steps of administering one or more non-encapsulated highly toxic anticancer drugs to a mammal by inhalation, wherein at least one of the drugs comprises a severe vesicant.

Another embodiment is an apparatus for treating a patient for a neoplasm by inhalation that is a combination of a nebulizer and a formulation for treating a neoplasm, the formulation including (1) a non-encapsulated anticancer drug, and (2) a pharmaceutically acceptable carrier; wherein when 0.2 ml of the formulation is injected intradermally to rats, at the clinical concentration for parenteral use in humans: (a) a lesion results which is greater than about 20 mm² in area fourteen days after the

intradermal injection; and (b) at least 50% of the tested rats have these lesions. A further embodiment includes a formulation which when injected results in a lesion which is greater than about 10 mm² in area 30 days after the intradermal injection; and at least about 50% of the tested rats have 5 these longer lasting lesions. Typically the formulation includes an anthracycline. Anthracyclines may be selected from the group consisting of epirubicin, daunorubicin, methoxymorpholinodoxorubicin, cyanomorpholinyl doxorubicin, doxorubicin, and idarubicin. The formulation also typically and contain a vinca alkaloid. Vinca alkaloids may be selected from the group 10 consisting of vincristine, vinorelbine, vinorelbine, vindesine, and vinblastine. Alternately, the formulation may contain vesicant selected from the group consisting of mechlorethamine, mithramycin, and dactinomycin; or bisantrene or amsacrine. Typically the formulation can also contain a taxane which is typically a paclitaxel or doxytaxel.

15 Another embodiment of the invention includes an inhalation mask for administering aerosols to an patient comprising: means for enclosing the mouth and nose of the patient, having an open end and a closed end, the open end adapted for placing over the mouth and nose of the patient; upper and lower holes in the closed end adapted for insertion of a nose outlet 20 tube and a mouth inhalation tube; the nose outlet tube attached to the upper hole, adapted to accept exhaled breath from the nose of the patient; a one way valve in the nose tube adapted to allow exhalation but not inhalation; the mouth inhalation tube having an outer and an inner end, partially inserted through the lower hole, the inner end continuing to end at the rear of the 25 patients mouth, the inhalation tube end cut at an angle so that the lower portion extends further into the patients mouth than the upper portion and adapted to fit the curvature of the rear of the patients mouth; and a y-adapter attached to the outer end of the mouth inhalation tube. The mask typically will have a moderate vesicant or a severe vesicant present in the inhalation 30 tube.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the plasma drug concentration time profile for dog #101 having doxorubicin administered intravenously (IV) (circles) and by the pulmonary inhalation route (IH) (squares). The vertical Y scale is the 5 concentration of drug in the circulatory system in ng/ml and the horizontal X scale is time after treatment in hours.

Figure 2 shows the plasma drug concentration time profile for dog #102 having doxorubicin administered intravenously (IV) (circles) and by the pulmonary inhalation route (IH) (squares). The vertical Y scale is the 10 concentration of drug in the circulatory system in ng/ml and the horizontal X scale is time after treatment in hours.

Figure 3 shows the plasma drug concentration time profile for dog #103 having doxorubicin administered intravenously (IV) (circles) and by the pulmonary inhalation route (IH) (squares). The vertical Y scale is the 15 concentration of drug in the circulatory system in ng/ml and the horizontal X scale is time after treatment in hours.

Figure 4 shows a schematic of the pulmonary delivery apparatus arrangement that was used to administer drug to dogs by inhalation for Example 3.

20 Figure 5 shows a schematic of the pulmonary delivery apparatus arrangement that was used to administer high doses and multiple doses of drug to dogs by inhalation for Example 4.

Figure 6 shows a schematic drawing of details of a mask useful for administering drugs by inhalation to a mammal such as a dog.

25 Figure 7 shows a schematic drawing of a portable device for administration of anticancer drugs according to the invention.

DETAILED DESCRIPTION OF THE INVENTION AND BEST MODE

The delivery of antineoplastic drugs by inhalation by the pulmonary 30 route is an attractive alternative to the administration of drugs by various injectable methods, particularly those drugs that are given on a chronic or repeated administration schedule. A cause of concern is the toxic nature of

the drugs particularly those that are cytotoxic such as the classes represented by alkylating agents, taxanes, vinca alkaloids, platinum complexes, anthracyclines and others that are considered particularly toxic especially when administered outside the circulatory system.

5 Broadly, the inventors have discovered that highly toxic, vesicant and previously unknown nonvesicant antineoplastic drugs can be effectively delivered to a patient in need of treatment for neoplasms or cancers by inhalation. This route is particularly effective for treatment of neoplasms or cancers of the pulmonary system because the highly toxic drugs are delivered
10 directly to the site where they are needed, providing regional doses much higher than can be achieved by conventional IV delivery. As used herein the respiratory tract includes the oral and nasal-pharyngeal, tracheo-bronchial, and pulmonary regions. The pulmonary region is defined to include the upper and lower bronchi, bronchioles, terminal bronchioles, respiratory bronchioles,
15 and alveoli.

 An important benefit from inhalation therapy for neoplasms of the head, neck and respiratory tract, is that exposure to the rest of the body is controlled following administration of high doses of drug and consequently is spared much of the adverse side effects often associated with high doses of
20 systemically administered highly toxic antineoplastic drugs, yet significantly increased doses are provided at the site of the tumor. These toxic effects include for example: cardiotoxicity, myelosuppression, thrombocytopenia, renal toxicity, and hepatic toxicity that are often life threatening. The toxic effects are often so severe that it is not uncommon for patients to die from
25 the effects of the systemically administered drugs rather than from the disease for which they are being treated.

 Broadly, vesicants as used herein include chemotherapeutic agents that are toxic and typically cause long lasting damage to surrounding tissue if the drug is extravasated. If inadvertently delivered outside of a vein, a vesicant
30 has the potential to cause pain, cellular damage including cellulitis, tissue destruction (necrosis) with the formation of a long lasting sore or ulcer and sloughing of tissues that may be extensive and require skin grafting. In

extreme cases extravasation of vesicants such as doxorubicin has required surgical excision of the affected area or amputation of the affected limb. Examples of antineoplastic chemotherapeutic agents that are generally accepted vesicants include alkylating agents such as mechlorethamine, 5 dactinomycin, mithramycin; topoisomerase II inhibitors such as bisantrene, doxorubicin (adriamycin), daunorubicin, dactinomycin, amsacrine, epirubicin, daunorubicin, and idarubicin; tubulin inhibitors such as vincristine, vinblastine, and vindesine; and estramustine. A partial list of vesicants is found in Table 1.

10 In another embodiment, vesicants as more narrowly used herein include drugs that produce a lesion in rats, where the average lesion size is greater than about 20 mm² in area, fourteen days after an intradermal injection of 0.2 ml of the drug, and where 50% or more of the animals have this size of lesion. The drug concentration for the intradermal injection is the 15 clinical concentration recommended by the manufacturer for use in humans, the dose recommended in the Physicians Desk Reference, 1997 (or a more current version of this reference), or another drug manual for health specialists. If there is no recommendation by the manufacturer (for example for because the drug is new) and there is no recommendation in the 20 Physicians Desk Reference or similar drug manual for health specialists then other current medical literature may be used. If more than one clinical concentration is recommended, the highest recommended clinical concentration is used. Lesion as used herein means an open sore or ulcer or sloughing off of skin with exposure of underlying tissue.

25 In a yet further embodiment of the invention, 0.2 ml of a highly toxic anticancer drug (vesicant) at a dose recommended for humans (as discussed above) is administered intradermally to rats at a concentration that causes the above mentioned lesion size for a more extended period of time. That is, the lesions remain above about 10 mm² up to at least 30 days in at least 50% or 30 more of the animals.

Nonvesicants typically are also irritating and can cause pain, but do not usually result in long lasting sores or ulcers or sloughing off of tissues except

in exceptional cases. Examples include alkylating agents such as cyclophosphamide, bleomycin (blenoxane), carmustine, and dacarbazine; DNA crosslinking agents such as thiotepa, cisplatin, melphalan (L-PAM); antimetabolites such as cytarabine, fluorouracil (5-FU), methotrexate (MTX),

5 and mercaptourine (6 MP); topoisomerase II inhibitors such as mitoxantrone; epipodophyllotoxins such as etoposide (VP-16) and teniposide (VM-26); hormonal agents such as estrogens, glucocorticosteroids, progestins, and antiestrogens; and miscellaneous agents such as asparaginase, and streptozocin.

10 A listing of materials usually accepted to be vesicants or nonvesicants is provided below as Table 1- Vesicant/Nonvesicant Drug Activity.

Table 1
Vesicant/Non-Vesicant Drug Activity

5

Classification	Vesicant	Non-Vesicant
Alkylating Agents	Mechlorethamine ^{a,c,d,e} * Mitomycin-C ^{a,c,e*} Dactinomycin ^{d,e*} Mithramycin ^d (Plicamycin)	Cyclophosphamide (Cytoxan) ^b Bleomycin (Blenoxane) ^{b,c} Carmustine ^{a,b,d} Mithramycin ^{a,b} (Plicamycin) Dacarbazine ^{a,b,c}
DNA Crosslinking Agents		Thiotepa ^b Cisplatin ^{b,c} Melphalan (L-PAM) ^b
Antimetabolites		Cytarabine (ARA C) ^b Fluorouracil (5 FU) ^{b,d,e} Methotrexate (MTX) ^b Mercaptopurine (6 MP) ^b
Topoisomerase II Inhibitors	Bisantrene ^{c,e*} (Anthracene) Dactinomycin ^{a,c} Doxorubicin ^{a,b,c,d,e*} (Anthracycline) Cyanomorpholinyl Doxorubicin ^{c*} Amsacrine ^{a,c,e*} Epirubicin ^{c,e*} Daunorubicin ^{a,d,e*} Idarubicin ^{a,e*}	Mitoxantrone ^{b,c} (Anthracene) Esorubicin ^c Etoposide (VP-16) ^{a,b,c} (Epipodophyllotoxin) Teniposide (VM-26) ^{a,b,c} (Epipodophyllotoxin) Liposomal anthracyclines ^c
Hormonal Agents		Estrogens ^b Glucocorticosteroids ^b Progestins ^b Antiestrogens ^b
Tubulin Inhibitors	Vinblastine ^{a,d,e*} Vincristine ^{a,d,e*} Vinorelbine ^{c*} Vindesine ^{a,c*} Paclitaxel ^c	Paclitaxel ^{c,f}
Miscellaneous		Asparaginase ^b (Enzyme) Aclacinomycin ^c Streptozocin ^{a,b} Menogaril ^c

a - According to US Patent 5,602,112
b - **Dorr**, R.T. et al, Lack of Experimental Vesicant Activity for the Anticancer Agents Cisplatin, Melphalan, and Mitoxantrone, *Cancer Chemother. Pharmacol.*, Vol. 16, 1986, pp. 91-94
c - According to **Bicher**, A. et al, Infusion Site Soft-Tissue Injury After Paclitaxel
5 Administration, *Cancer*, Vol. 76, No. 1, July 1, 1995, pp. 116-120
d - **Rudolph**, R. et al; Etiology and Treatment of Chemotherapeutic Agent Extravasation Injuries: A Review; *Journal of Clinical Oncology*, Vol. 5; No. 7; July 1987; pp. 1116-1126
e - **Bertelli**, G., Prevention and Management of Extravasation of Cytotoxic Drugs, *Drug Safety*, 12 (4) 1995; pp. 245-255. The listed drugs have been reported in at least one case, either clinically or experimentally, to cause tissue necrosis after accidental extravasation.
10 Symbol: * = vesicants, drugs with the highest potential for localized tissue damage after extravasation
f - **Cancer**, R.T., Communications, Author Reply, *Cancer*, pp. 226

15

Typical embodiments of the invention use highly toxic antineoplastic drugs that have similar or greater vesicating activity than those that have been tested in animals by inhalation to date. One embodiment typically uses severely vesicating toxic antineoplastic drugs having higher vesicating activity than those represented by 5-FU, 5 β -cytosine arabinoside (Ara-C, cytarabine), mitomycin C, and cisplatin. In respect to the latter, it is disclosed that a highly toxic drug represented by the class anthracyclines (of which doxorubicin is among the most toxic), has been administered by inhalation to a patient in need of treatment for neoplasms. In a further embodiment of the invention it is disclosed that vesicants other than doxorubicin can be given to patients by 10 inhalation. In respect to the latter, highly toxic drugs represented by the classes vinca alkaloids, and taxanes, having similar high toxicities have been administered by inhalation to a patient in need of treatment for neoplasms. In a yet further embodiment of the invention there is disclosed that certain antineoplastic drugs that are 15 nonvesicants can be administered by inhalation to a patient in need of treatment for neoplasms. In a further embodiment of the invention there are disclosed formulations and methods for applying the aforementioned highly toxic drugs to a patient in need of treatment for pulmonary neoplasms by inhalation.

Example 1

20 This example illustrates and confirms toxicity and vesicant/nonvesicant activity of several antineoplastic drugs. The vesicant activities of thirteen anticancer drugs were investigated (see the listing in Table 2 below). Doxorubicin has traditionally been considered a vesicant (see Table 1). Paclitaxel has previously been considered a nonvesicant, but recent literature has advocated its classification as a vesicant. Some 25 of the remaining drugs are traditionally considered to be vesicants and others nonvesicants (Table 1). Day fourteen after injection was chosen as the time for comparison for vesicant activity, because lesions caused by nonvesicants should have been significantly reduced while lesions caused by vesicants should still be large. Sterile saline solution (0.9%) for injection USP, pH 4.5-7.0, or sterile water for injection, 30 as appropriate, was used to reconstitute the drugs.

The drugs used for the vesicant activity tests are identified as follows:
doxorubicin (Adriamycin PFS), a red liquid in glass vials, no formulation was necessary;

cisplatin (Platinol-AQTM), a liquid in glass vials, no formulation was necessary; Paclitaxel (TaxolTM), a liquid in glass vials, formulated with saline solution; fluorouracil, a clear yellow liquid in glass vials, no formulation was necessary; cytarabine (Cytosar-UTM), a white powder in glass vials, formulated with water; 9-aminocamptothecin (9-AC 5 colloidal suspension), a yellow powder in glass vials, formulated with water; cyclophosphamide (CytoxanTM), a yellow powder in glass vials, formulated with a saline/water mixture; carboplatin (ParaplatinTM), a white powder in injectable vials, formulated with saline solution; etoposide (VePesidTM), a clear liquid in glass vials, formulated with saline solution; bleomycin (bleomycin sulfate, USP), a lyophilized 10 powder tablet in glass vials, formulated with saline solution; vincristine (vincristine sulfate), an injectable liquid in injection vials, no formulation necessary; vinorelbine tartrate (NavelbineTM), a clear liquid in glass vials, diluted with water per package 15 instructions; and mitomycin (MutamycinTM), a gray crystalline powder in glass amber bottles, formulated with water. All of these drugs were reconstituted following standard and known methods recommended by the manufacturers.

The tests for vesicant activity were conducted using Sprague Dawley rats (7-8 weeks old having 150-200 g of body weight. Each received a single intradermal injection of the test drug at the recommended clinical concentration (listed below in Table 2) in the right dorsum. Approximately 24 hours prior to administration, the hair 20 was removed from the dorsum using clippers and a depilatory agent. Each 0.2 ml injection was given with a 1 ml syringe and 27 gauge needle. All drug solutions were either isotonic or slightly hypertonic.

Table 2
Formulations administered for Vesicant Tests

Test Formulation	Formulation Concentration
1 Doxorubicin	2 mg/ml
2 Platinol	1 mg/ml
3 Paclitaxel	1.2 mg/ml
4 Fluorouracil	50 mg/ml
5 Cytarabine	100 mg/ml
6 9-aminocamptothecin	100 µg/ml
7 Cyclophosphamide	20 mg/ml
8 Carboplatin	10 mg/ml
9 Etoposide	0.4 mg/ml
10 Bleomycin	20 units/ml
11 Vincristine	1 mg/ml
12 Vinorelbine	3 mg/ml
13 Mitomycin-C	0.5 mg/ml

5

Table 3 below is a tabulation of the resultant lesion sizes that developed from intradermal injections of the above drugs. Lesion sizes were measured as more fully 10 discussed below.

Table 3. Individual Lesion Size Measurements (mm²) (part 1 of 3)

(see text for explanation of measurements)

Animal Number	Test Drug	Day of Test (post injection)															
		6	8	10	13	15	17	20	22	24	27	29	31	34	36	38	41
101	Doxorubicin	—	21.4	33.9	57.0	42.9	34.0	35.4	27.2	32.2	31.7	31.7	17.1	8.3	6.3	6.7	4.5
102	Doxorubicin	—	18.8	23.5	10.9	12.9	9.7	10.2	9.9	11.8	10.5	9.9	10.2	2.8	—	—	—
103	Doxorubicin	—	36.5	58.0	82.9	45.5	37.7	28.1	26.9	21.0	23.9	18.6	16.2	12.5	10.3	7.6	6.1
104	Doxorubicin	—	44.6	27.3	33.6	17.7	21.7	28.1	19.5	16.6	16.1	18.9	13.9	9.0	5.1	4.5	4.0
105	Doxorubicin	—	33.9	35.2	33.3	35.1	29.4	30.2	29.7	25.0	24.4	24.8	23.5	24.0	24.5	21.6	22.0
106	Doxorubicin	—	30.6	43.2	32.2	35.2	34.4	29.2	30.2	15.5	16.0	15.4	14.5	16.2	14.8	14.3	5.2
107	Doxorubicin	—	26.1	39.7	38.6	33.8	31.3	25.0	22.0	21.6	19.8	22.4	21.5	20.9	21.0	18.4	18.9
111	Platinol	26.9	18.7	18.0	11.8	21.2	17.1	6.9	1.5	1.0	—	—	—	—	—	—	—
112	Platinol	35.5	20.3	20.8	15.5	16.1	16.2	16.5	4.1	—	—	—	—	—	—	—	—
113	Platinol	15.3	15.8	14.6	10.1	9.1	9.0	8.3	2.9	2.6	1.7	—	—	—	—	—	—
114	Platinol	17.2	11.3	13.2	9.7	9.2	10.3	10.5	9.1	—	—	—	—	—	—	—	—
115	Platinol	26.8	25.0	14.8	21.8	18.0	15.0	16.0	16.0	2.1	1.7	1.4	—	—	—	—	—
116	Platinol	21.8	20.7	12.2	11.8	12.9	12.6	8.4	10.8	8.5	8.4	—	—	—	—	—	—
117	Platinol	24.9	21.3	16.7	15.1	16.4	14.8	14.3	12.2	12.5	2.8	—	—	—	—	—	—
121	Taxol	23.7	21.6	21.2	18.9	3.5	—	—	—	—	—	—	—	—	—	—	—
122	Taxol	37.3	30.1	26.1	25.2	21.8	21.7	5.6	2.1	1.8	—	—	—	—	—	—	—
123	Taxol	7.9	5.9	4.3	1.1	1.2	—	—	—	—	—	—	—	—	—	—	—
124	Taxol	43.2	36.9	32.9	30.6	29.0	28.5	—	—	—	—	—	—	—	—	—	—
125	Taxol	38.4	34.6	28.6	22.1	5.9	—	—	—	—	—	—	—	—	—	—	—
126	Taxol	69.5	59.5	53.3	53.3	42.9	5.2	—	—	—	—	—	—	—	—	—	—
127	Taxol	45.9	23.1	16.1	14.3	8.4	5.0	—	—	—	—	—	—	—	—	—	—
131	Fluorouracil	29.0	19.9	13.5	11.2	14.3	11.6	8.3	2.0	—	—	—	—	—	—	—	—
132	Fluorouracil	17.1	16.2	11.8	3.0	—	—	—	—	—	—	—	—	—	—	—	—
133	Fluorouracil	27.0	23.8	17.4	17.6	17.9	0.5	—	—	—	—	—	—	—	—	—	—
134	Fluorouracil	21.9	18.9	17.0	6.7	—	—	—	—	—	—	—	—	—	—	—	—
135	Fluorouracil	20.5	27.5	21.4	4.5	—	—	—	—	—	—	—	—	—	—	—	—
136	Fluorouracil	23.5	14.0	10.1	9.5	8.0	7.8	1.8	—	—	—	—	—	—	—	—	—
137	Fluorouracil	20.5	7.0	6.2	4.8	4.6	3.8	—	—	—	—	—	—	—	—	—	—

Table 3 (continued - part 2 of 3)

Day of Test

Animal Number	Test Drug	6	8	10	13	15	17	20	22	24	27	29	31	34	36	38	41	
151	9-aminocamptothecin	21.8	15.8	16.0	14.5	9.0	19.9	—	—	—	—	—	—	—	—	—	—	—
152	9-aminocamptothecin	8.6	4.4	5.4	3.7	4.0	3.6	—	—	—	—	—	—	—	—	—	—	—
153	9-aminocamptothecin	4.4	2.6	2.9	1.3	1.1	—	—	—	—	—	—	—	—	—	—	—	—
154	9-aminocamptothecin	23.8	21.9	20.9	19.8	15.5	18.6	—	—	—	—	—	—	—	—	—	—	—
155	9-aminocamptothecin	12.5	7.9	10.0	9.6	9.9	0.6	—	—	—	—	—	—	—	—	—	—	—
156	9-aminocamptothecin	12.6	10.4	5.8	4.6	3.1	—	—	—	—	—	—	—	—	—	—	—	—
157	9-aminocamptothecin	12.5	7.8	5.2	3.7	—	—	—	—	—	—	—	—	—	—	—	—	—
161	Cyclophosphamide	16.4	13.6	11.3	9.4	8.3	8.6	—	—	—	—	—	—	—	—	—	—	—
162	Cyclophosphamide	35.1	33.8	23.2	3.5	1.4	—	—	—	—	—	—	—	—	—	—	—	—
163	Cyclophosphamide	25.8	18.9	21.0	19.3	17.2	17.2	12.1	12.5	14.0	7.8	2.4	—	—	—	—	—	—
165	Cyclophosphamide	19.4	18.2	17.9	17.4	16.6	15.9	13.2	12.2	12.7	7.5	1.8	1.5	—	—	—	—	—
166	Cyclophosphamide	31.8	33.8	25.4	23.9	11.9	2.2	—	—	—	—	—	—	—	—	—	—	—
167	Cyclophosphamide	25.2	19.7	19.1	19.3	18.9	18.9	17.4	14.6	15.6	4.1	2.0	—	—	—	—	—	—
171	Carboplatin	16.2	17.3	12.2	10.9	10.4	8.1	4.6	0.8	—	—	—	—	—	—	—	—	—
172	Carboplatin	9.0	5.1	21.9	17.5	7.6	4.1	5.2	6.2	5.9	5.2	3.2	2.6	—	—	—	—	—
173	Carboplatin	24.8	23.4	17.7	20.5	18.5	16.0	8.6	3.4	0.8	0.6	—	—	—	—	—	—	—
174	Carboplatin	31.9	23.1	18.2	24.2	27.0	19.4	15.5	13.1	11.2	4.0	1.5	—	—	—	—	—	—
175	Carboplatin	20.5	24.5	22.1	13.4	20.4	16.8	5.4	4.9	1.8	1.2	—	—	—	—	—	—	—
177	Carboplatin	42.9	39.1	30.1	31.7	32.7	32.6	35.4	34.7	34.6	23.9	25.2	25.7	19.2	0.6	—	—	—
181	Etoposide	21.1	15.0	11.2	9.2	9.8	9.0	2.9	—	—	—	—	—	—	—	—	—	—
182	Etoposide	—	—	3.8	2.4	2.0	1.7	1.1	—	—	—	—	—	—	—	—	—	—
183	Etoposide	1.3	4.6	3.1	2.9	3.8	1.2	—	—	—	—	—	—	—	—	—	—	—
184	Etoposide	—	9.6	4.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—
185	Etoposide	5.9	6.0	6.0	2.6	2.1	2.0	—	—	—	—	—	—	—	—	—	—	—
186	Etoposide	10.6	14.1	7.7	6.6	8.4	3.8	1.7	—	—	—	—	—	—	—	—	—	—
187	Etoposide	6.5	10.0	9.3	5.3	5.4	5.1	3.5	—	—	—	—	—	—	—	—	—	—

Table 3 (continued - part 3 of 3)
Day of Test

Animal Number	Test Drug	6	8	10	13	15	17	20	22	24	27	29	31	34	36	38	41	
191	Bleomycin	8.2	5.1	8.8	2.2	1.6	—	—	—	—	—	—	—	—	—	—	—	—
192	Bleomycin	21.1	15.3	10.8	16.3	3.8	1.3	—	—	—	—	—	—	—	—	—	—	—
193	Bleomycin	23.5	18.9	15.4	13.8	5.5	1.3	—	—	—	—	—	—	—	—	—	—	—
194	Bleomycin	—	5.0	3.2	1.0	2.3	—	—	—	—	—	—	—	—	—	—	—	—
195	Bleomycin	7.7	6.5	6.7	6.5	7.0	3.2	1.3	—	—	—	—	—	—	—	—	—	—
196	Bleomycin	13.4	7.8	6.8	7.2	6.6	0.7	—	—	—	—	—	—	—	—	—	—	—
197	Bleomycin	27.0	27.0	26.0	25.2	26.0	24.0	1.0	0.6	—	0.4	—	—	—	—	—	—	—
202	Vincristine	—	—	469.0	307.7	227.7	160.5	109.2	93.3	93.6	83.6	67.2	57.9	47.5	40.3	40.2	34.0	
203	Vincristine	—	—	—	165.3	158.5	67.0	29.7	28.6	24.7	21.1	22.0	22.8	27.5	30.6	21.2	13.8	
205	Vincristine	—	—	—	130.4	136.2	111.6	76.1	61.5	58.0	42.0	26.5	18.1	12.6	5.3	4.2	1.3	
206	Vincristine	—	—	145.6	96.9	81.6	96.1	66.7	59.2	51.3	13.0	7.2	—	—	—	—	—	
211	Vinorelbine	—	16.8	421.7	315.2	289.7	274.6	250.8	200.8	170.8	159.1	237.2	243.6	243.1	219.4	180.6	149.0	
212	Vinorelbine	—	436.7	422.1	426.5	408.5	347.6	316.8	298.8	292.4	282.0	251.0	81.3	82.0	83.8	45.8	17.2	
213	Vinorelbine	—	402.2	429.0	352.6	323.4	372.9	366.3	311.6	312.1	299.2	302.3	294.0	102.7	137.7	212.1	192.1	
214	Vinorelbine	—	322.1	261.6	283.6	293.9	241.7	227.0	221.9	227.2	105.0	86.1	72.5	65.3	71.4	52.5	62.0	
215	Vinorelbine	—	297.0	277.8	269.7	225.3	204.2	82.5	69.8	67.8	40.0	28.4	31.9	17.4	19.2	14.0	14.5	
216	Vinorelbine	—	348.3	325.1	308.1	288.9	297.0	278.7	255.9	269.3	255.8	134.9	103.7	61.2	95.7	123.2	108.2	
217	Vinorelbine	—	275.1	309.6	272.1	249.0	217.1	208.1	209.3	190.7	175.5	173.2	172.0	173.4	157.3	187.7	155.5	
221	Mutamycin	45.0	46.8	47.5	77.0	48.2	38.8	45.4	41.6	40.3	28.6	9.6	6.4	4.1	0.7	—	—	
222	Mutamycin	50.4	50.4	49.6	41.9	45.1	34.8	42.0	46.2	9.9	9.3	7.5	—	—	—	—	—	
223	Mutamycin	98.3	73.0	79.1	79.8	71.0	64.6	66.0	28.5	17.6	24.3	28.2	1.1	—	—	—	—	
224	Mutamycin	58.2	82.4	62.6	78.8	73.3	66.1	53.9	36.9	32.9	31.2	19.8	16.8	15.5	16.8	21.0	25.6	
225	Mutamycin	28.1	24.2	28.0	19.8	29.8	23.0	12.8	13.2	11.9	8.5	6.6	7.2	2.0	1.5	—	—	
226	Mutamycin	61.3	53.3	59.9	49.7	48.9	38.0	39.5	42.1	40.6	23.0	5.6	4.8	4.6	4.6	1.2	—	
227	Mutamycin	36.0	35.8	37.8	37.8	39.7	33.8	31.1	13.9	10.9	7.9	8.1	2.9	—	—	—	—	

Results were as follows:

1. Abrasions of the dorsal body were observed in a majority of animals for all drugs except cytarabine.
2. Alopecia of the dorsal body was seen for doxorubicin (3/7), paclitaxel (7/7),
5 and fluorouracil (7/7), etoposide (7/7), bleomycin (7/7), vincristine (2/7), vinorelbine (7/7), and mitomycin-C (mutamycin) (4/7).
3. Discoloration of the skin around the site of injection was seen for doxorubicin, vincristine, vinorelbine, and mitomycin-C.
4. Rough coat was observed in fluorouracil (1/7), vincristine (4/7), and
10 vinorelbine (2/7).
5. Systemic effects were observed only for vincristine. Three animals had to be removed from the tests because of their poor condition.
6. Slight edema was observed for all groups. Moderate edema was observed in doxorubicin, vincristine, vinorelbine, and mitomycin-C treated animals.
- 15 7. Severe edema was observed only for animals treated with vinorelbine and vincristine.
8. Dermal lesions were observed for all drugs except for cisplatin (platinol) and cytarabine.
- 20 25 30 35 40 45 50 55 60 65 70 75 80 85 90 95 100 105 110 115 120 125 130 135 140 145 150 155 160 165 170 175 180 185 190 195 200 205 210 215 220 225 230 235 240 245 250 255 260 265 270 275 280 285 290 295 300 305 310 315 320 325 330 335 340 345 350 355 360 365 370 375 380 385 390 395 400 405 410 415 420 425 430 435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560 565 570 575 580 585 590 595 600 605 610 615 620 625 630 635 640 645 650 655 660 665 670 675 680 685 690 695 700 705 710 715 720 725 730 735 740 745 750 755 760 765 770 775 780 785 790 795 800 805 810 815 820 825 830 835 840 845 850 855 860 865 870 875 880 885 890 895 900 905 910 915 920 925 930 935 940 945 950 955 960 965 970 975 980 985 990 995 1000 1005 1010 1015 1020 1025 1030 1035 1040 1045 1050 1055 1060 1065 1070 1075 1080 1085 1090 1095 1100 1105 1110 1115 1120 1125 1130 1135 1140 1145 1150 1155 1160 1165 1170 1175 1180 1185 1190 1195 1200 1205 1210 1215 1220 1225 1230 1235 1240 1245 1250 1255 1260 1265 1270 1275 1280 1285 1290 1295 1300 1305 1310 1315 1320 1325 1330 1335 1340 1345 1350 1355 1360 1365 1370 1375 1380 1385 1390 1395 1400 1405 1410 1415 1420 1425 1430 1435 1440 1445 1450 1455 1460 1465 1470 1475 1480 1485 1490 1495 1500 1505 1510 1515 1520 1525 1530 1535 1540 1545 1550 1555 1560 1565 1570 1575 1580 1585 1590 1595 1600 1605 1610 1615 1620 1625 1630 1635 1640 1645 1650 1655 1660 1665 1670 1675 1680 1685 1690 1695 1700 1705 1710 1715 1720 1725 1730 1735 1740 1745 1750 1755 1760 1765 1770 1775 1780 1785 1790 1795 1800 1805 1810 1815 1820 1825 1830 1835 1840 1845 1850 1855 1860 1865 1870 1875 1880 1885 1890 1895 1900 1905 1910 1915 1920 1925 1930 1935 1940 1945 1950 1955 1960 1965 1970 1975 1980 1985 1990 1995 2000 2005 2010 2015 2020 2025 2030 2035 2040 2045 2050 2055 2060 2065 2070 2075 2080 2085 2090 2095 2100 2105 2110 2115 2120 2125 2130 2135 2140 2145 2150 2155 2160 2165 2170 2175 2180 2185 2190 2195 2200 2205 2210 2215 2220 2225 2230 2235 2240 2245 2250 2255 2260 2265 2270 2275 2280 2285 2290 2295 2300 2305 2310 2315 2320 2325 2330 2335 2340 2345 2350 2355 2360 2365 2370 2375 2380 2385 2390 2395 2400 2405 2410 2415 2420 2425 2430 2435 2440 2445 2450 2455 2460 2465 2470 2475 2480 2485 2490 2495 2500 2505 2510 2515 2520 2525 2530 2535 2540 2545 2550 2555 2560 2565 2570 2575 2580 2585 2590 2595 2600 2605 2610 2615 2620 2625 2630 2635 2640 2645 2650 2655 2660 2665 2670 2675 2680 2685 2690 2695 2700 2705 2710 2715 2720 2725 2730 2735 2740 2745 2750 2755 2760 2765 2770 2775 2780 2785 2790 2795 2800 2805 2810 2815 2820 2825 2830 2835 2840 2845 2850 2855 2860 2865 2870 2875 2880 2885 2890 2895 2900 2905 2910 2915 2920 2925 2930 2935 2940 2945 2950 2955 2960 2965 2970 2975 2980 2985 2990 2995 3000 3005 3010 3015 3020 3025 3030 3035 3040 3045 3050 3055 3060 3065 3070 3075 3080 3085 3090 3095 3100 3105 3110 3115 3120 3125 3130 3135 3140 3145 3150 3155 3160 3165 3170 3175 3180 3185 3190 3195 3200 3205 3210 3215 3220 3225 3230 3235 3240 3245 3250 3255 3260 3265 3270 3275 3280 3285 3290 3295 3300 3305 3310 3315 3320 3325 3330 3335 3340 3345 3350 3355 3360 3365 3370 3375 3380 3385 3390 3395 3400 3405 3410 3415 3420 3425 3430 3435 3440 3445 3450 3455 3460 3465 3470 3475 3480 3485 3490 3495 3500 3505 3510 3515 3520 3525 3530 3535 3540 3545 3550 3555 3560 3565 3570 3575 3580 3585 3590 3595 3600 3605 3610 3615 3620 3625 3630 3635 3640 3645 3650 3655 3660 3665 3670 3675 3680 3685 3690 3695 3700 3705 3710 3715 3720 3725 3730 3735 3740 3745 3750 3755 3760 3765 3770 3775 3780 3785 3790 3795 3800 3805 3810 3815 3820 3825 3830 3835 3840 3845 3850 3855 3860 3865 3870 3875 3880 3885 3890 3895 3900 3905 3910 3915 3920 3925 3930 3935 3940 3945 3950 3955 3960 3965 3970 3975 3980 3985 3990 3995 4000 4005 4010 4015 4020 4025 4030 4035 4040 4045 4050 4055 4060 4065 4070 4075 4080 4085 4090 4095 4100 4105 4110 4115 4120 4125 4130 4135 4140 4145 4150 4155 4160 4165 4170 4175 4180 4185 4190 4195 4200 4205 4210 4215 4220 4225 4230 4235 4240 4245 4250 4255 4260 4265 4270 4275 4280 4285 4290 4295 4300 4305 4310 4315 4320 4325 4330 4335 4340 4345 4350 4355 4360 4365 4370 4375 4380 4385 4390 4395 4400 4405 4410 4415 4420 4425 4430 4435 4440 4445 4450 4455 4460 4465 4470 4475 4480 4485 4490 4495 4500 4505 4510 4515 4520 4525 4530 4535 4540 4545 4550 4555 4560 4565 4570 4575 4580 4585 4590 4595 4600 4605 4610 4615 4620 4625 4630 4635 4640 4645 4650 4655 4660 4665 4670 4675 4680 4685 4690 4695 4700 4705 4710 4715 4720 4725 4730 4735 4740 4745 4750 4755 4760 4765 4770 4775 4780 4785 4790 4795 4800 4805 4810 4815 4820 4825 4830 4835 4840 4845 4850 4855 4860 4865 4870 4875 4880 4885 4890 4895 4900 4905 4910 4915 4920 4925 4930 4935 4940 4945 4950 4955 4960 4965 4970 4975 4980 4985 4990 4995 5000 5005 5010 5015 5020 5025 5030 5035 5040 5045 5050 5055 5060 5065 5070 5075 5080 5085 5090 5095 5100 5105 5110 5115 5120 5125 5130 5135 5140 5145 5150 5155 5160 5165 5170 5175 5180 5185 5190 5195 5200 5205 5210 5215 5220 5225 5230 5235 5240 5245 5250 5255 5260 5265 5270 5275 5280 5285 5290 5295 5300 5305 5310 5315 5320 5325 5330 5335 5340 5345 5350 5355 5360 5365 5370 5375 5380 5385 5390 5395 5400 5405 5410 5415 5420 5425 5430 5435 5440 5445 5450 5455 5460 5465 5470 5475 5480 5485 5490 5495 5500 5505 5510 5515 5520 5525 5530 5535 5540 5545 5550 5555 5560 5565 5570 5575 5580 5585 5590 5595 5600 5605 5610 5615 5620 5625 5630 5635 5640 5645 5650 5655 5660 5665 5670 5675 5680 5685 5690 5695 5700 5705 5710 5715 5720 5725 5730 5735 5740 5745 5750 5755 5760 5765 5770 5775 5780 5785 5790 5795 5800 5805 5810 5815 5820 5825 5830 5835 5840 5845 5850 5855 5860 5865 5870 5875 5880 5885 5890 5895 5900 5905 5910 5915 5920 5925 5930 5935 5940 5945 5950 5955 5960 5965 5970 5975 5980 5985 5990 5995 6000 6005 6010 6015 6020 6025 6030 6035 6040 6045 6050 6055 6060 6065 6070 6075 6080 6085 6090 6095 6100 6105 6110 6115 6120 6125 6130 6135 6140 6145 6150 6155 6160 6165 6170 6175 6180 6185 6190 6195 6200 6205 6210 6215 6220 6225 6230 6235 6240 6245 6250 6255 6260 6265 6270 6275 6280 6285 6290 6295 6300 6305 6310 6315 6320 6325 6330 6335 6340 6345 6350 6355 6360 6365 6370 6375 6380 6385 6390 6395 6400 6405 6410 6415 6420 6425 6430 6435 6440 6445 6450 6455 6460 6465 6470 6475 6480 6485 6490 6495 6500 6505 6510 6515 6520 6525 6530 6535 6540 6545 6550 6555 6560 6565 6570 6575 6580 6585 6590 6595 6600 6605 6610 6615 6620 6625 6630 6635 6640 6645 6650 6655 6660 6665 6670 6675 6680 6685 6690 6695 6700 6705 6710 6715 6720 6725 6730 6735 6740 6745 6750 6755 6760 6765 6770 6775 6780 6785 6790 6795 6800 6805 6810 6815 6820 6825 6830 6835 6840 6845 6850 6855 6860 6865 6870 6875 6880 6885 6890 6895 6900 6905 6910 6915 6920 6925 6930 6935 6940 6945 6950 6955 6960 6965 6970 6975 6980 6985 6990 6995 7000 7005 7010 7015 7020 7025 7030 7035 7040 7045 7050 7055 7060 7065 7070 7075 7080 7085 7090 7095 7100 7105 7110 7115 7120 7125 7130 7135 7140 7145 7150 7155 7160 7165 7170 7175 7180 7185 7190 7195 7200 7205 7210 7215 7220 7225 7230 7235 7240 7245 7250 7255 7260 7265 7270 7275 7280 7285 7290 7295 7300 7305 7310 7315 7320 7325 7330 7335 7340 7345 7350 7355 7360 7365 7370 7375 7380 7385 7390 7395 7400 7405 7410 7415 7420 7425 7430 7435 7440 7445 7450 7455 7460 7465 7470 7475 7480 7485 7490 7495 7500 7505 7510 7515 7520 7525 7530 7535 7540 7545 7550 7555 7560 7565 7570 7575 7580 7585 7590 7595 7600 7605 7610 7615 7620 7625 7630 7635 7640 7645 7650 7655 7660 7665 7670 7675 7680 7685 7690 7695 7700 7705 7710 7715 7720 7725 7730 7735 7740 7745 7750 7755 7760 7765 7770 7775 7780 7785 7790 7795 7800 7805 7810 7815 7820 7825 7830 7835 7840 7845 7850 7855 7860 7865 7870 7875 7880 7885 7890 7895 7900 7905 7910 7915 7920 7925 7930 7935 7940 7945 7950 7955 7960 7965 7970 7975 7980 7985 7990 7995 8000 8005 8010 8015 8020 8025 8030 8035 8040 8045 8050 8055 8060 8065 8070 8075 8080 8085 8090 8095 8100 8105 8110 8115 8120 8125 8130 8135 8140 8145 8150 8155 8160 8165 8170 8175 8180 8185 8190 8195 8200 8205 8210 8215 8220 8225 8230 8235 8240 8245 8250 8255 8260 8265 8270 8275 8280 8285 8290 8295 8300 8305 8310 8315 8320 8325 8330 8335 8340 8345 8350 8355 8360 8365 8370 8375 8380 8385 8390 8395 8400 8405 8410 8415 8420 8425 8430 8435 8440 8445 8450 8455 8460 8465 8470 8475 8480 8485 8490 8495 8500 8505 8510 8515 8520 8525 8530 8535 8540 8545 8550 8555 8560 8565 8570 8575 8580 8585 8590 8595 8600 8605 8610 8615 8620 8625 8630 8635 8640 8645 8650 8655 8660 8665 8670 8675 8680 8685 8690 8695 8700 8705 8710 8715 8720 8725 8730 8735 8740 8745 8750 8755 8760 8765 8770 8775 8780 8785 8790 8795 8800 8805 8810 8815 8820 8825 8830 8835 8840 8845 8850 8855 8860 8865 8870 8875 8880 8885 8890 8895 8900 8905 8910 8915 8920 8925 8930 8935 8940 8945 8950 8955 8960 8965 8970 8975 8980 8985 8990 8995 9000 9005 9010 9015 9020 9025 9030 9035 9040 9045 9050 9055 9060 9065 9070 9075 9080 9085 9090 9095 9100 9105 9110 9115 9120 9125 9130 9135 9140 9145 9150 9155 9160 9165 9170 9175 9180 9185 9190 9195 9200 9205 9210 9215 9220 9225 9230 9235 9240 9245 9250 9255 9260 9265 9270 9275 9280 9285 9290 9295 9300 9305 9310 9315 9320 9325 9330 9335 9340 9345 9350 9355 9360 9365 9370 9375 9380 9385 9390 9395 9400 9405 9410 9415 9420 9425 9430 9435 9440 9445 9450 9455 9460 9465 9470 9475 9480 9485 9490 9495 9500 9505 9510 9515 9520 9525 9530 9535 9540 9545 9550 9555 9560 9565 9570 9575 9580 9585 9590 9595 9600 9605 9610 9615 9620 9625 9630 9635 9640 9645 9650 9655 9660 9665 9670 9675 9680 9685 9690 9695 9700 9705 9710 9715 9720 9725 9730 9735 9740 9745 9750 9755 9760 9765 9770 9775 9780 9785 9790 9795 9800 9805 9810 9815 9820 9825 9830 9835 9840 9845 9850 9855 9860 9865 9870 9875 9880 9885 9890 9895 9900 9905 9910 9915 9920 9925 9930 9935 9940 9945 9950 9955 9960 9965 9970 9975 9980 9985 9990 9995 9999

A vesicant as determined by the methods used herein is defined as causing a lesion of at least about 20 mm^2 , in at least one half of the animals, two weeks after injection (day 15 in Table 3). Table 3 shows that doxorubicin, paclitaxel, carboplatin, vincristine, vinorelbine, and mitomycin-C 5 fulfill these criteria. Cisplatin, etoposide, bleomycin, cytarabine, cyclophosphamide, fluorouracil, and 9-aminocamptothecin are thus categorized as non-vesicants.

A moderate vesicant as determined by the methods used herein is defined as causing a lesion of at least about 20 mm^2 , in at least one half of 10 the animals, two weeks after injection (day 15 in Table 3), but less than half of the animals will have lesions greater than about 10 mm^2 30 days after injection (day 31 in Table 3). The data from Table 3 shows that paclitaxel, carboplatin, and mitomycin-C fulfill these criteria. Of these, mitomycin-C has been determined to exhibit substantial pulmonary toxicity.

15 A severe vesicant as determined by the methods used herein is defined as causing a lesion of at least about 20 mm^2 , in at least one-half of the animals, two weeks after injection (day 15 in Table 3), and at least one-half of the animals will still have lesions greater than about 10 mm^2 , 30 days after injection (day 31 in Table 3). Table 3 shows that doxorubicin, vincristine, 20 and vinorelbine satisfy these criteria.

Surprisingly it has now been found that moderate to severe vesicants can be used for inhalation therapy of cancer as revealed in the discussion and examples below. Further, other highly toxic drugs, although not having the severity of reaction of moderate to severe vesicants have also been found to 25 be useful in the treatment of cancer by inhalation as further discussed below.

Antineoplastic drugs that are highly toxic and useful in an embodiment of the present invention include the anthracyclines (e.g. doxorubicin, epirubicin, idarubicin, methoxy-morpholinodoxorubicin, daunorubicin, and the like); vinca alkaloids (e.g. vincristine, vinblastine, vindesine, and the like); 30 alkylating agents (e.g. mechlorethamine and the like); carboplatin; nitrogen mustards (e.g. melphalan and the like), topoisomerase I inhibitors (e.g. 9-aminocamptothecin, camptothecin, topotecan, irinotecan, 9-NO-

camptothecin, and the like); topoisomerase II inhibitors (e.g. etoposide, teniposide, and the like); and paclitaxel and the like. These and other useful compounds are further discussed below.

In yet a further embodiment of the invention, there are disclosed

5 formulations and methods for applying an appropriate selection of highly toxic drugs that are efficacious in treating the neoplasm or cancer, that are applied by inhalation and that reside in the pulmonary system for a time sufficient to increase the exposure of the neoplasm to the drug, yet allow a reduction and/or controlled systemic exposure of the drug, and provide a more

10 efficacious treatment for pulmonary neoplasms.

In a further embodiment of the invention, it is disclosed that it is possible to deliver antineoplastic drugs by the pulmonary route as a means to provide systemic treatment of distant tumors. The inventors have shown that for selected drugs inhalation can be used as a noninvasive route of delivery

15 without causing significant toxicity to the respiratory tract. This is in contrast with the prior art that used inhalation for treatment of disease in the respiratory system.

As used herein the term patient includes a mammal including, but not limited to, mice, rats, cats, horses, dogs, cattle, sheep, apes, monkeys, goats, camels, other domesticated animals, and of course humans.

Administration by inhalation as used herein includes the respiratory administration of drugs as either liquid aerosols or powdered aerosols suspended in a gas such as air or other nonreactive carrier gas that is inhaled by a patient. Non-encapsulated drug as used herein means that the

25 antineoplastic drug is not enclosed within a liposome, or within a polymeric matrix, or within an enclosing shell. Where the term encapsulated drug is used herein the term means that the antineoplastic drug is enclosed within a liposome, within a polymeric matrix, or within an enclosing shell. However, in some embodiments the antineoplastic drug may be coupled to various

30 molecules yet is still not enclosed in a liposome, matrix or shell as further discussed below.

In other embodiments of the invention the antineoplastic drugs disclosed herein may be coupled with other molecules through ester bonds. Enzymes present in the respiratory system later cleave the ester bonds. One purpose of coupling the antineoplastic drugs through an ester bond is to

5 increase the residence time of the antineoplastic drug in the pulmonary system. Increased residence time is achieved by: first, an increase in molecular weight due to the attached molecule; second, by appropriate choice of a coupled molecule; third, other factors such as for example charge, solubility, shape, particle size of the delivered aerosol, and protein binding can

10 be modified and used to alter the diffusion of the drug. Molecules useful for esterification with the drug include alpha-hydroxy acids and oligomers thereof, vitamins such as vitamins A, C, E and retinoic acid, other retinoids, ceramides, saturated or unsaturated fatty acids such as linoleic acid and glycerin. Preferred molecules for esterification are those naturally present in

15 the area of deposition of the active drug in the respiratory tract.

As a demonstration of the proof of concept, doxorubicin was used in a series of tests. Doxorubicin was chosen as an initial test agent since it is one of most cytotoxic and potent vesicants of all anti-neoplastic agents considered in the broad embodiment (pulmonary delivery of anti-neoplastic drugs) of the present invention. Based on positive outcome of these proof of concept studies, anticancer drugs from other major classes were simultaneously tested. Results consistently showed that using the approach and methods described in this invention the drug could be safely and effectively delivered by inhalation. In Examples 2 and 3 below, doxorubicin was administered to

20 three dogs (beagles) by both the pulmonary and intravenous route of administration. The dogs were given a clinically effective dosage of the drug and the amount of the drug appearing in the blood system was measured.

25

An anthracycline antineoplastic drug, a salt of doxorubicin, doxorubicin HCl, available from Farmitalia Carlo Erba (now Pharmacia & Upjohn), Milan, Italy, was used in some of the examples herein. The liquid formulation that was administered to the dogs by inhalation of an aerosol was obtained by mixing the doxorubicin hydrochloride with a mixture of ethanol/water at a

doxorubicin concentration of approximately 15 – 25 mg/ml. Typically solutions of 5 – 75% ethanol are preferred. Water/ethanol ratios may be adjusted to select the desired concentration of doxorubicin and the desired particle size of the aerosol.

5

Example 2

Three adult, male, beagle dogs were used in the tests. The dogs (designated dog 101, 102, and 103) had body weights of 10.66, 10.24, and 10.02 kg respectively. As used herein "m²" used alone with reference to dose refers to square meters in terms of the body surface area of a treated animal or patient, at other times it is qualified in terms of lung surface area. The dogs were given a slow IV infusion treatment of the anthracycline drug doxorubicin HCl at the recommended initial clinical dose (for dogs) of 20 mg/m² or 1 mg/kg of body weight. A 1 mg/ml drug solution was administered at a rate of 2.0 ml/kg/hr for 30 minutes. The 30-minute infusion interval simulated the time/dose exposure relationship of the inhalation group in Example 3 below. A series of blood samples were taken to characterize the IV pharmacokinetics at predose, 2, 5, 10, 30, 60, 90 minutes and 2, 4, 6, 12, 18, and 36 hours post dosing. Additional blood samples were collected for clinical pathology evaluations on days 3 and 7 of the IV treatment. Changes in blood chemistry and hematology were as expected with administration of doxorubicin HCl at these doses.

Example 3

25 The three dogs used in Example 2 were allowed a one-week washout period before being subjected to exposure to the anthracycline drug doxorubicin HCl by inhalation. The dogs were acclimated to wearing masks for administration of the aerosol prior to treatment. The dogs were exposed to an aerosol concentration of drug sufficient to deposit a total dose of about 30 10 mg (1 mg/kg). Based on aerosol dosimetry models, approximately one half of this dose was deposited within the respiratory tract. The total dose

was about equal to the dosage administered by IV infusion. The dose was calculated using the following equation:

$$\text{Dose} = \{ \text{Drug Conc. (mg/liter)} \times \text{Mean minute Vol. (liter/min)} \times \text{Exposure Duration (min)} \times \text{Total Deposition Fraction (\%)} \} \div \text{Body Weight (kg)}$$

wherein

Mean Min. Vol. = Tidal volume x minute respiratory rate

Exposure Duration = 30 min

Mean Body Weight = weight in kg for each dog

Total Deposition Fraction = 60% (determined by particle size and respiratory tract deposition models from the published literature such as "Respiratory Tract Deposition of Inhaled Polydisperse Aerosols in Beagle Dogs", R.G. Cuddihy et al, Aerosol Science, Vol 4, pp. 35-45 (1973) and

"Deposition and Retention Models for Internal Dosimetry of the Human Respiratory Tract", Task Group on Lung Dynamics, Health Physics, Vol. 12, pp. 173-207 (1966).

Pulmonary function measurements (respiratory rate, tidal volume, and minute volume (calculated)) were monitored during a 30 minute inhalation exposure session. These data provided an estimate of each animal's inspired volume during exposure, and were used to calculate the mass of drug deposited in the respiratory tract.

A series of blood samples were collected at the end of the exposure to characterize the pharmacokinetics. Clinical pathology evaluations were conducted on the third day. All three dogs were necropsied on the third day.

Referring now to Figure 4, the drug formulation was administered to the dogs of Example 3 with drug exposure system 400. The drug was aerosolized with two Pari LC Jet Plus™ nebulizers 401. The nebulizer was filled with a solution of 15 mg doxorubicin per ml of 50%water/50% ethanol.

The output of each nebulizer 401 was continuous and set to provide the required concentration of aerosol in attached plenum 405. The nebulizers 401 were attached directly to plenum 405 that had a volume of approximately

90 liters . Plenum 405 was connected by four tubes 407 to four venturi 409, respectively, and subsequently connected to four Y-fittings 413 by additional tubing 411. Typical venturi were used to measure the inhaled volume of drug formulation. One end of each of the Y-fittings 411 interfaced with a dog 5 breathing mask 415 while the other end of Y-fitting 411 was connected to tubing 417 leading to an exhaust pump 419. During the tests three dogs 418 were fitted with three of the breathing masks 415. A collection filter 421 was placed in the remaining mask 415. A vacuum pump 423 that drew 1 liter per 10 minute of air for 3 minutes was used in the place of a dog to draw aerosol in order to monitor and measure the amount of drug administered. The vacuum pump was activated four times during the 30-minute administration of drug to the dogs and the amount of drug trapped by the filter set forth in Table 5 below.

A flow of air was supplied to each of the nebulizers 401 from a supply 15 of air 425 via lines 427. Additional air for providing a bias flow of air through the system and for the breathing requirements of the dogs was provided from air supply 425 by supply lines 429 connected to one way valves 431. The one way valves 431 were connected to the upper portion of the nebulizers 401. This additional supply of air provided a continuous flow of air through the 20 system 400 from the air supply 425 to the exhaust pump 417. Alternatively one could eliminate the extra supply of air from supply lines 429 to one way valves 431 and let ambient room air enter the one way valves from the suction action of the nebulizers 401. A Hepa filter 441 mounted to the top of plenum 405 allowed room air to flow in and out of plenum 405 and assured 25 that there was always ambient pressure in the plenum. There was a continuous flow of air containing the aerosol past the masks of the dogs and the dogs were able to breathe air containing the aerosol on demand. An inner tube 621 located within dog breathing mask 415 extended into the mouth of the dogs and was provided with an extension 633 at its lower 30 portion that served to depress the tongue of the dogs to provide an open airway for breathing. See the discussion of Figure 6 below.

Each of the four venturi 409 were connected by line 441 to a pressure transducer 443 (the one shown is typical for the four venturi) that was used to measure pressure differences across the venturi. The pressure transducers 443 were connected by line 445 to an analog amplifier 447 to increase the 5 output signal and prepare the signal sent via line 449 to computer system 451. Computer system 451 is a desk model PC of typical design in the industry and can be used in conjunction with a BUXCO or PO-NE-MAH software program to calculate the uptake of air containing aerosol and thus the drug dosage by each of the dogs.

10 Table 4 below summarizes the exposure data for doxorubicin administration to dogs from Example 3. The total mass for each dog was determined. The total inhaled volume of air for the 30 minute drug administration was measured in liters. The aerosol concentration in mg of drug/liter of air (mg/l) was determined from calibration tests done earlier. A 15 total deposition fraction of 60% was calculated (As calculated 30% for the inhaled dose was deposited in the conducting upper airways and peripheral lung while an additional 30% was deposited in the oral-pharyngeal region) based on the measured doxorubicin aerosol particle size and the published literature (see references cited above).

20 Thus about 25%-30% of the administered doxorubicin was deposited and available to the pulmonary region. Since the drug was administered in its salt form, a correction for the chlorine portion of the molecule was made. As shown in the Table 4 this resulted in an applied dose of 0.51, 0.60, and 0.57 mg/kg to the pulmonary region of dogs 101, 102, and 103 respectively

25 Filter data obtained from analysis of drug deposited on a filter 421 placed in a fourth mask 415 are shown in Table 5 for four different measurements. The drug mass collected on the filter was corrected for the chlorine portion of the doxorubicin salt. Finally, the doxorubicin concentration in the three liters of air drawn into each mask was determined in mg/l. The 30 four figures were averaged to obtain a mean doxorubicin aerosol concentration of 0.218 mg/l.

Table 6 shows data and calculations that verify the figures of Table 4. The dog weight and breath volumes measured for Table 4 are used. However, the mean doxorubicin concentration that was obtained from the filter data shown in Table 5 was used to calculate doxorubicin concentrations.

5 Making calculations with the data as in Table 4, the inhaled dose for each dog was calculated. The inhaled dose was reduced by 40% as before to obtain the total dose deposited, and reduced by 50% again to obtain the total deposited pulmonary dose. The pulmonary doses obtained by this method of 0.47, 0.56, and 0.53 mg/kg for dogs 101, 102, and 103 respectively compare

10 well with the earlier calculated figures in Table 4.

TABLE 4. TOTAL MASS DATA

Dog No.	Dog Weight (kg)	Total Inhaled Vol. (l) For 30 Min.	Inhaled Air Aerosol Conc. (mg/l)	Deposition Fraction	Test Art. Fraction	Inhaled Dose (mg/kg)	Deposited Dose (mg/kg)	Pulmonary Dose (mg/kg)
101	10.66	77.5	0.250	0.60	0.937	1.70	1.02	0.51
102	10.24	86.8	0.250	0.60	0.937	1.99	1.19	0.60
103	10.02	80.8	0.250	0.60	0.937	1.89	1.13	0.57
	A	B	C	D	E			

TABLE 5. FILTER DATA

Sample No.	Sample Vol. (liter)	Weight Gain (mg)	Doxorubicin mass (mg)	Total Conc. (mg/l)	Dox. Conc. (mg/l)	Ratio Dox/Total
1	3	0.78 x .937	0.70	0.260	0.233	0.897
2	3	0.72 x .937	0.61	0.240	0.203	0.847
3	3	0.73 x .937	0.62	0.243	0.207	0.849
4	3	0.77 x .937	0.68	0.257	0.227	0.883
Mean				0.250	0.218	0.869
	A	B	C	D		

TABLE 6. ANALYTICAL DATA

Dog No.	Dog Weight (kg)	Total Inhaled Vol. (l)	Aerosol Conc. (mg/l)	Inhaled Dose (mg/kg)	Deposited Dose (mg/kg)	Pulmonary Dose (mg/kg)
101	10.66	77.5	0.218	1.58	0.95	0.47
102	10.24	86.8	0.218	1.85	1.11	0.56
103	10.02	80.8	0.218	1.76	1.06	0.53
	A	B	C			

Surprisingly it was found that free non-encapsulated doxorubicin administered by the pulmonary route was not rapidly cleared from the lung. Figures 1, 2 and 3 show examples of the type of results achieved when cytotoxic anticancer drugs were given by inhalation. High efficiency 5 nebulization systems as shown in Figures 4 and 5 were used to deliver a large percentage of aerosolized drug to the pulmonary region of the respiratory tract. Doses equal to or greater than those that cause toxicity when given IV, were only moderately absorbed into the blood following pulmonary delivery and caused little to no direct or systemic toxicity after a single exposure at 10 this dose.

As can be seen from Figures 1, 2 and 3, the pulmonary route administered doxorubicin achieved a consistently lower level of doxorubicin in systemic blood, with peak blood levels being over an order of magnitude lower following inhalation exposure. The initial concentration of doxorubicin 15 at 2 minutes was about 1.5 orders of magnitude larger when administered IV than by the pulmonary route. Later, after about 4 hours, the systemic doxorubicin level was about six times higher for the IV administered drug. This suggests that free doxorubicin remained in the lung for an extended period of time and slowly passed through the mucosa into systemic 20 circulation. This reduces the systemic toxic effects of the drug and allows its concentration in the lung for more effective treatment of respiratory tract associated neoplasms while reducing overall systemic toxic effects. It is believed that the toxic effects of doxorubicin to tissues outside the lung are as 25 a result of the aforementioned high levels of systemic drug concentration following IV treatment.

Another surprising finding was that doxorubicin administered by the pulmonary route did not produce the severe toxic effects on the respiratory tract (including the oral and nasal-pharyngeal, tracheo-bronchial, and pulmonary regions). As was noted earlier, doxorubicin belongs to the 30 anthracycline class of drugs that are typically very toxic. In particular doxorubicin is one of the most toxic drugs in the class, yet when the dogs in the test were necropsied, no damage to the respiratory tract was observed.

It is surprising that the doxorubicin was not toxic to the lung when given by inhalation at clinically relevant doses such as 20 to 60 mg/m². Unlike 5-FU and Ara-C, and cisplatin, doxorubicin is well known to generate the production of free radicals (Myers et al, 1977) which are notorious for causing pulmonary toxicity (Knight, 1995). It is this property, in fact, which is held responsible for the cardiotoxicity caused by doxorubicin given by the intravenous route (Myers et al, 1977).

In some typical embodiments, to obtain additional benefits of the disclosed invention for treating pulmonary neoplasms and reducing systemic toxicity, it is important that antineoplastic drugs administered in non-encapsulated form by the pulmonary route be absorbed into and remain in the tumor tissue for an extended period of time and diffuse across the lung mucosa in a relatively slow manner. In general, although solubility, charge and shape have an influence, slow diffusion is obtained by drugs having higher molecular weights while faster diffusion is obtained by those having relatively lower molecular weights. Thus drugs such as doxorubicin having a molecular weight of 543.5, have relatively slow rates of diffusion, drugs such as vincristine (MW=825), vinblastine (MW=811), paclitaxel (MW=854), etoposide (MW=589), having higher molecular weights also diffuse slowly.

Other drugs having somewhat lower molecular weights such as 9-aminocamptothecin, while diffusing more slowly are still included within the invention. It has been demonstrated that significantly higher tissue concentrations can be achieved in the lung by pulmonary delivery compared to conventional parenteral or oral administration. Further, systemic coverage of micrometastases can be provided under these conditions, with the benefit of significantly greater doses of drug delivered to the respiratory tract tumor sites and controlled systemic exposure.

Thus in one embodiment of the invention drugs having a molecular weight above 350 are used. In this regard mitomycin-C (MW of about 334) is thus excluded from this embodiment. While molecular weight is not the sole determinant controlling diffusion through the lung it is one of the important factors for selecting compounds useful in the present invention. This lower

molecular weight limit is about 64% that of doxorubicin. This will help assure that the limited systemic availability of the drug discussed above is maintained. In further embodiments of the invention the molecular weight of the drugs administered is above 400, 450, and 500 respectively.

5 In conjunction with the above discussed molecular weights, protein binding of the antineoplastic agents to be delivered by pulmonary administration should also be considered with respect to diffusion through the lung. Higher rates of protein binding will further slow diffusion through the lung mucosa. In this respect 5-FU and Ara-C in addition to having low
10 molecular weights also have relatively low protein binding affinity of 7% and 13% respectively. That is, when placed into a protein-containing solution, only 7% and 13% of these drugs bind to the protein while the remainder is free in solution. In this respect, cisplatin does not bind to tissues, rather at a later stage it is the platinum in the cisplatin that binds to tissues, thus
15 allowing cisplatin to enter systemic circulation as further discussed below. In comparison doxorubicin, vincristine, vinblastine, paclitaxel, etoposide, and 9-amino-camptothecin have rates of protein binding above 50%. Typically protein-binding affinity above 25% is preferred, more preferred is binding above 50%, with protein binding above 75% being most preferred when lung
20 retention is the objective.

 In a preferred formulation and method for treating neoplasms of the pulmonary system by inhalation, the diffusion characteristics of the particular drug formulation through the pulmonary tissues are chosen to obtain an efficacious concentration and an efficacious residence time in the tissue to be
25 treated. Doses may be escalated or reduced or given more or less frequently to achieve selected blood levels. Additionally the timing of administration and amount of the formulation is preferably controlled to optimize the therapeutic effects of the administered formulation on the tissue to be treated and/or titrate to a specific blood level.

30 Diffusion through the pulmonary tissues can additionally be modified by various excipients that can be added to the formulation to slow or accelerate the absorption of drugs into the pulmonary tissues. For example,

the drug may be combined with surfactants such as the phospholipids, dimyristoylphosphatidyl choline, and dimyristoylphosphatidyl glycerol. The drugs may also be used in conjunction with bronchodilators that can relax the bronchial airways and allow easier entry of the antineoplastic drug to the 5 lung. Albuterol is an example of the latter with many others known in the art. Further, the drug may complexed with biocompatible polymers, micelle forming structures or cyclodextrins

10 Particle size for the aerosolized drug used in the present examples was measured at about 2.0-2.5 μm with a geometric standard deviation (GSD) of about 1.9-2.0. Typically the particles should have a particle size of from about 1.0-5.0 μm with a GSD less than about 2.0 for deposition within the central and peripheral compartments of the lung. As noted elsewhere herein particle sizes are selected depending on the site of desired deposition of the drug particles within the respiratory tract.

15 Aerosols useful in the invention include aqueous vehicles such as water or saline with or without ethanol and may contain preservatives or antimicrobial agents such as benzalkonium chloride, paraben, and the like, and/or stabilizing agents such as polyethyleneglycol.

20 Powders useful in the invention include formulations of the neat drug or formulations of the drug combined with excipients or carriers such as mannitol, lactose, or other sugars. The powders used herein are effectively suspended in a carrier gas for administration. Alternatively, the powder may be dispersed in a chamber containing a gas or gas mixture which is then inhaled by the patient.

25 Further, the invention includes controlling deposition patterns and total dose through careful control of patient inspiratory flow and volume. This may be accomplished using the pulmonary devices described herein and similar devices. The inventors have shown by gamma scintigraphy measurements that drug aerosol deposition is maximized and evenly distributed in the 30 peripheral lung when the patient inhales using slow flow rates and inhales to maximum lung volumes followed by brief breath holds. Central lung deposition is favored when faster inspiratory flow rates and lower inspiratory

volumes are used. Further, total deposited and regionally deposited doses are significantly changed as a patient's inspiratory patterns change. Therefore, the method of treatment and the use of the delivery devices described herein can be modified to target different regions of the respiratory tract and adjusted to deliver different doses of drug. It is the integration of drug molecular weight, protein binding affinity, formulation, aerosol generation condition, particle sized distribution, interface of aerosol delivery to the patient via the device and the control of the patient's inspiratory patterns that permit targeted and controlled delivery of highly toxic anti-cancer drugs to the respiratory tract with the option to minimize or provide controlled systemic availability of drug.

Example 4

The tests for administration of doxorubicin by inhalation referred to in Example 3 were substantially repeated at different dosages using a different drug administration system 500 described below. In the present examples eight dogs were used. The dogs were divided into two dose groups. A first group was the low dose group given a total daily dose of 60 mg/m^2 for three days or a total dose of 180 mg/m^2 . This resulted in a pulmonary deposition of about 90 mg/m^2 .

A high dose group was administered a dose of 180 mg/m^2 daily for three days or a total dose of 540 mg/m^2 . This resulted in a pulmonary deposition of about 270 mg/m^2 .

One half of the animals were necropsied after three days of exposure and the remaining dogs necropsied after a three day recovery period.

The purpose of the tests was to identify the maximum tolerated dose of inhaled drug.

For comparison with the results of Examples 2 and 3, one can convert the data from mg/kg to mg/m^2 (m^2 of body area) by multiplying by 20 (conversion factor for the dog). Thus the exposure of the dogs in Examples 2 and 3 which were the equivalent of a clinical dose (for dogs) was about 20 mg/m^2 . When one compares these dosages to those of Example 4 (180

mg/m² and 540 mg/m²) it is apparent that a significantly higher dose of non-encapsulated drug can be delivered to the lung compared to the known art. Although dogs receiving the lower total dose ranges showed few toxic effects, while dogs receiving the higher total doses had pulmonary toxicity, these 5 doses were 9-27 times higher than those generally given clinically to dogs.

While the present examples used active drug doses of doxorubicin of about 20 mg/m², 180 mg/m², and 270 mg/m², effective amounts of the active anticancer drugs can be from very small amounts to those where toxicity to normal tissue becomes a problem. As used herein, effective amounts and 10 pharmaceutically effective amounts of antineoplastic drug deposited or applied to areas needing treatment are dosages that reduce a neoplasm or tumor mass, stop its growth or eliminate it altogether.

Referring now to Figure 5, the liquid formulation was administered to the dogs by aerosolizing with a nebulizer exposure system 500 comprising a 15 Pari LC Jet Plus™ nebulizer 501. The nebulizer was filled with the solution of drug with which the dogs were to be treated. The output of the nebulizer 501 was pulsed in a series of bursts over time (one pulse every ten seconds). The nebulizer 501 was attached directly to a 460 cc volume plenum 503 and the plenum 503 was connected to a canine mouth only exposure mask 415 via a 20 short piece of anesthesia tubing 505 and Y-fitting 507. The mask 415 was tapered to approximately fit the shape of the dog's snout. There was no bias airflow through the exposure system 500. The test atmosphere was pulled through the exposure system 500 by the inhalation of the dog 511. A one way breathing valve 513 on the top of the nebulizer 501 allowed the dog 511 25 to draw in room air and pull the air through the system 500. The air entrained and transported the aerosolized drug through the plenum 503, tubing 505, Y-fitting 507, and mask 415 to the dog 511. A one way valve 515 connected to the Y-fitting 507 allowed the dog 511 to exhale and the exhaled air exited the system. An air supply 520 provided a flow of air to controller 30 530 via line 521. Air flow to the nebulizer was controlled by controller 530 and supplied to the nebulizer via line 531.

Referring now to Figure 6, details of mask 415 are shown. Means for enclosing the mouth and nose are of flexible material and are preferably held on by straps such as Velcro™ straps or belts. Means for enclosing 601 has one end 603 for inserting the nose and mouth of the dog while the other end 5 605 has two openings 607,609 for attachment of nose outlet tube 611. Nose outlet tube 611 has a one way valve 613 that allows the dog to exhale but not inhale through its nose. Mouth tube 621 is inserted and attached to opening 609 and lies within the means for enclosing 601. An optional Y- connector 623 may be attached and used with mouth tube 621 for providing 10 and receiving inhaled and exhaled gases. Air is generally inhaled through leg 625 of the Y-connector 623. The air passes through the mouth tube 621 and out the inner opening 631 into the respiratory system of the dog. Inner opening 631 is cut at an angle with its lower portion 633 extending further into the mouth of the dog than the upper portion 635. Lower portion 633 15 functions to depress the tongue of the dog and allow more efficient flow of air and aerosol into the dog. When the dog is wearing mask 415 it can only breathe in through its mouth using the mouth tube 621. Means for enclosing 601 effectively seals the dog's mouth and nose from outside air. The use of a nose outlet tube 611 has been found to greatly ease the dog's wearing of the 20 mask. Air exhaled through the mouth exits mouth tube 621 and passes into optionally attached Y-connector or to another tube not shown. Air exits Y connector 623 via outlet tube 627. If desired the Y-connector 623 or other outer tube (e.g. straight tubing) may be made of one piece and simply pass into the enclosing means 601 or may be of separate pieces that fit together. 25 In either case an adapter 637 may be used to hold the mouth tube 621 and or other tubing to which it is connected.

A general device for administering aerosols to a patient includes an inhalation mask for administering aerosols to the patient, including means for enclosing the mouth and nose of the patient, having an open end and a 30 closed end, the open end adapted for placing over the mouth and nose of the patient; upper and lower holes in the closed end adapted for insertion of a nose outlet tube and a mouth inhalation tube; the nose outlet tube attached

to the upper hole, adapted to accept exhaled breath from the nose of the patient; a one way valve in the nose tube adapted to allow exhalation but not inhalation; the mouth inhalation tube having an outer and an inner end, partially inserted through the lower hole, the inner end continuing to end at 5 the rear of the patients mouth, the inhalation tube end cut at an angle so that the lower portion extends further into the patients mouth than the upper portion and adapted to fit the curvature of the rear of the mouth; and a y-adapter attached to the outer end of the mouth inhalation tube.

Pulmonary administration by inhalation may be accomplished by means 10 of producing liquid or powdered aerosols, for example, by the devices disclosed herein or by using any of various devices known in the art. (see e.g. Newman, S.P., 1984, in *Aerosols and the Lung*, Clarke and Davia (Eds.), Butterworths, London, England, pp. 197-224; PCT Publication No. WO 92/16192 dated October 1, 1992; PCT Publication No. WO 91/08760 dated 15 June 27, 1991; NTIS Patent Application 7-504-047 filed April 3, 1990 by Roosdorp and Crystal) including but not limited to nebulizers, metered dose inhalers, and powder inhalers. Various delivery devices are commercially available and can be employed, e.g. Ultravent nebulizer (Mallinckrodt, Inc, St. Louis, MO); Acorn II nebulizer (Marquest Medical Products, Englewood, CO); 20 Ventolin metered dose inhalers (Glaxo Inc., Research Triangle Park, North Carolina); Spinhaler powder inhaler (Fisons Corp., Bedford, MA) or Turbohaler (Astra). Such devices typically entail the use of formulations suitable for dispensing from such a device, in which a propellant material may be present. Ultrasonic nebulizers may also be used.

25 Nebulizer devices such as those in Greenspan et al US patents 5,511,726 and 5,115,971 are useful in the invention. These devices use electrohydrodynamic forces to produce a finely divided aerosol having uniformly sized droplets by electrical atomization. While the Greenspan devices use piezoelectric materials to generate electrical power any power 30 source is acceptable to produce the electrohydrodynamic forces for nebulization.

A nebulizer may be used to produce aerosol particles, or any of various physiologically inert gases may be used as an aerosolizing agent. Other components such as physiologically acceptable surfactants (e.g. glycerides), excipients (e.g. lactose), carriers (e.g. water, alcohol), and diluents may also 5 be included.

As will be understood by those skilled in the art of delivering pharmaceuticals by the pulmonary route, a major criteria for the selection of a particular device for producing an aerosol is the size of the resultant aerosol particles. Smaller particles are needed if the drug particles are mainly or only 10 intended to be delivered to the peripheral lung, i.e. the alveoli (e.g. 0.1-3 μm), while larger drug particles are needed (e.g. 3-10 μm) if delivery is only or mainly to the central pulmonary system such as the upper bronchi. Impact of particle sizes on the site of deposition within the respiratory tract is generally known to those skilled in the art. See for example the discussions 15 and figures in the articles by Cuddihy et al (Aerosol Science; Vol. 4; 1973, pp 35-45) (Fig. 6, 7, and 8 of the article) and The Task Group on Lung Dynamics (Fig. 11 and 14 of the article). As a result primary cancers in the naso-pharyngial or oral-pharyngeal regions and upper tracheo-bronchial regions, often referred to as cancers of the head and neck, are treatable with the 20 present invention. The major metastatic sites (lung and upper respiratory tract) are also readily treated with this invention simultaneously, unlike current methods of treatment.

Referring now to Figure 7, there is disclosed a nebulizer apparatus 700 that is preferably portable for administration of drug to a patient in need of 25 therapy. The nebulizer apparatus 700 is used in combination with the highly toxic drugs of the present invention and with drugs having properties adapted for optimum treatment of neoplasms as discussed elsewhere herein. Figure 7 is a schematic of a nebulizer combination according to the present invention. Nebulizer 701 may be any nebulizer as described earlier herein that is able to 30 produce the particle sizes needed for treatment. In combination with nebulizer 701 there is provided a highly toxic drug formulation 703 for treatment of neoplasms as disclosed herein. An air supply 705 is provided

either as a tank of compressed gas or as a motorized pump or fan for moving air from the room. An optional mouthpiece 707 may be used where it is necessary to provide sealed contact between the nebulizer and the patient. Optionally the mouthpiece 707 may be molded as part of nebulizer 701.

5 Power for use of the nebulizer apparatus 700 may come from the compressed gas from hand manipulation by the user or administrator or by batteries or electrical power not shown but well known by those skilled in the art.

To control environmental contamination resulting from use of a nebulizer, the patient may be placed in a well-ventilated area with exhaust air
10 filtered to remove antineoplastic drug that escapes from the device.

Examples 5 to 11

Examples 5F to 11F show inhalation feasibility and proof of concept tests and Examples 5R to 10R show dose escalation range tests with: vesicant
15 antineoplastic drugs including doxorubicin, paclitaxel, vincristine, vinorelbine; nonvesicant drugs including etoposide, and 9-aminocampothecin (9-AC) and carboplatin. The drugs were delivered to the pulmonary system via aerosol at a particle size of about 2 to about 3 μm . The drugs were delivered in water or other vehicles appropriate for the drug as is known in the art and as
20 exemplified herein.

Table 7 illustrates the dosage schedule for the range-finding studies. A minimum of 7-14 days separated each escalating dose. No range finding tests, only feasibility tests, were performed for mitomycin-C and 9-AC. No feasibility tests, only dose range-finding tests, were performed for vinorelbine.
25 It is important to note that the doses listed in Table 7 are the pulmonary deposited doses not the total doses administered.

The results of the feasibility and dose escalation studies are summarized in Tables 7 to 11.

TABLE 7
 Escalating Dose Regimen for Range-Finding Studies
 Mean Pulmonary Deposited Dose

Example No.	Test Drug	1 st Dose (mg)	2 nd Dose (mg)	3 rd Dose (mg)	4 th Dose (mg)	5 th Dose (mg)	6 th Dose (mg)
5R	Paclitaxel	30	35	40	40	60	-
6R	Doxorubicin	12	15	15	15	18	-
7R	Vincristine	0.55	0.55	0.70	0.70	1.1	1.5
8R	Vinorelbine	6	10	10	15	15	-
9R	Etoposide	25	30	45	55	40	80
10R	9-AC	-	-	-	-	-	-
11R	Carboplatin	30	-	-	-	-	-

Notes: A minimum of 7-14 days separated each escalating dose.
 Animals necropsied after last dosing.

Animals used in Examples 5 to 11 were adult beagle dogs. For the feasibility studies, the dogs were initially given a single intravenous (IV) dose of antineoplastic drug. This dose was given to allow a comparison of how much drug was absorbed into the blood after inhalation compared to IV delivery. The IV dose given was 5 typically the usual human clinical dose that had been scaled down for the dogs based on differences in body mass, or the maximum tolerated dose in the dog, whichever is greater. An average human having a weight of 70kg is considered to have a weight to body surface ratio of 37 kg/m^2 and a lung surface area of $70 - 100 \text{ m}^2$ of lung surface area. The average dog used in the tests was considered to have 10 a weight of 10kg corresponding and a weight to body surface ratio of 20 kg/m^2 and a lung surface area of $40 - 50 \text{ m}^2$ lung surface area (CRC Handbook of Toxicology, 1995, CRC Press Inc.). The single IV dose was used to quantify the plasma kinetics. With most of the cytotoxic agents treated, the single IV dose resulted in a predictable mild decrease in white blood cell counts, with no other measurable 15 toxicities.

After the initial IV and before the inhalation feasibility tests, the dogs were allowed a washout period of at least seven days (until the dogs returned to normal conditions) before they were treated with inhaled antineoplastic drugs. In the inhalation feasibility tests the dogs were generally exposed to a dose of inhaled 20 antineoplastic drug in aerosol form once per day for three consecutive days (except as noted in Tables 8 to 11) and necropsied one day following the last dose with the plasma kinetics characterized after the first and third exposures. With the exception of cisplatin and the high dose of doxorubicin, which caused toxicity to the respiratory tract, the drugs did not exhibit any significant pulmonary toxicity in these repeated 25 exposure inhalation feasibility studies. In the feasibility tests the dogs used the same mask and apparatus used for the earlier examples. In the dose range-finding tests, in order to control the deposited dose, the dogs were fitted with an endotracheal tube and the drug administered as an aerosol directly from the endotracheal tube. This latter procedure made it easier to control the pulmonary 30 deposited dose since the aerosol was released directly into the pulmonary air

passages assuring deep deposition of the drug in the lung. Also use of the endotracheal tube made it possible to do the tests in a shorter time since the dogs needed a four to six week training period to properly acclimate to and use the masks. The calculated deposited doses obtained herein were verified experimentally 5 by pulmonary scintigraphy tests in dogs.

Examples 5F and 5R

Referring now to Table 8, this table shows the details of the feasibility test of paclitaxel. Initially the dogs were administered 120 mg/m² of paclitaxel by IV. After 10 the washout period the dogs were administered a total deposited dose of 120 mg/m² of paclitaxel, by inhalation, three times for a total deposited dose of 360mg/m². This administered dose resulted in a pulmonary deposited dose of about 27 mg each time or a total pulmonary dose of about 81 mg. This represents a total pulmonary deposited dose of about 2.1 mg/m² of lung surface area. The dosages were 15 calculated as follows: the dose of 120 mg/m² was divided by 20 kg/m² to yield a 6 mg/kg dose that was multiplied by 10 kg for the average dog to yield about 60 mg of drug. Since the dogs were using the masks for drug administration, one half or about 30 mg of drug was considered deposited in the deep lung. Since the drug was administered three times the total drug exposure was about 90 mg. The 90 mg 20 of drug was divided by 40 to yield a total dose to the lung of about 2.25 mg/m² lung surface area.

The clinical condition of the dogs was normal. Clinical pathology profiles were normal with only mildly reduced white blood cell counts. The histopathology showed bone marrow and lymphoid depletion, GI villous atrophy and congestion and 25 laryngeal inflammation. These changes indicated that some significant fraction of the deposited drug was absorbed systemically. There was no respiratory tract toxicity found. Bioavailability of the paclitaxel was found to be low to moderate based on plasma kinetic evaluations. The low to moderate bioavailability indicates that most of the paclitaxel remained in the lungs and did not rapidly enter systemic 30 circulation in large amounts. Therefore, given the lack of significant direct

respiratory tract toxicity, the probable dose limiting toxicity is considered to be myelosuppression and/or GI toxicity. Thus factors extrinsic to the lung are expected to limit dosages provided by the pulmonary route.

Referring again to Tables 7 and 8, in the range-finding tests 60 to 120 mg/m² of paclitaxel were administered at weekly intervals for five weeks. The amount of pulmonary deposited dose ranged from about 30 to about 60 mg. This range corresponded to about 0.75 to about 1.50 mg/m² lung surface area. The clinical conditions of these dogs were normal, with clinical pathology changes limited to moderate white blood cell count reduction. The histopathology showed thoracic and mesenteric lymphoid depletion along with GI inflammation and ulceration. The histopathology reflects that normally found in IV administration of paclitaxel particularly GI inflammation and ulceration which is probably associated with systemically administered paclitaxel. Respiratory tract toxicity indicated minimal pulmonary interstitial inflammation. Systemic bioavailability was proportional to dose. The probable dose limiting toxicity is myelosuppression and GI toxicity, and not pulmonary toxicity.

TABLE 8 - Paclitaxel

Summary Results of Dog Feasibility and Dose Range-Finding Studies

Chemotherapy	IV dose	Inhalation Dose	Pulmonary Deposited Dose	Clinical Condition	Clinical Pathology	Histopathology	Respiratory Tract Toxicity	Bioavailability	Probable Dose-Limiting Toxicity
Example 5F Paclitaxel Feasibility	120 mg/m ²	120 mg/m ² x 3 doses (360 mg/m ² total)	30 mg x 3 doses	Normal	↓ WBC	Bone marrow & lymphoid depletion GI villous Atrophy & congestion Laryngeal inflammation	None	Low-moderate	Myelo-suppression GI toxicity
Example 5R Paclitaxel Dose Range-Finding	NA	60-120 mg/m ² (5 wkly Rx)	30-60 mg per dose	Normal	↓↓ WBC	Thoracic and mesenteric lymphoid depletion GI inflammation and ulceration	Minimal pulmonary interstitial inflammation	Proportional to dose	Myelo-suppression GI toxicity

5 * - Divide the pulmonary deposited dose in mg by 40 to get the pulmonary deposited dose in mg/m² of lung surface area.
WBC - white blood cell count

Examples 6F and 6R

Referring now to Table 9, 20 mg of doxorubicin were initially

5 administered by IV. After the washout period three sets of inhalation feasibility tests were made. In the first, a single dose of 20 mg/m² of doxorubicin was administered that gave about a 10 mg body dose, a pulmonary deposited dose of about 5 mg or about 0.125 mg/m² lung surface area. No changes were noted in the animal from this dose. A second set of

10 moderate inhalation dosages of about 40 mg/m² of doxorubicin (about 10 mg deposited within the lung) was administered three times a day for three consecutive days. Total cumulative dose administered was 120 mg/m² corresponding to a about a 60 mg body dose, and a total pulmonary deposited dose of about 30 mg (or about 0.75 mg/m² of lung surface area).

15 A third set of high inhalation dosages of 120 mg/m² of doxorubicin was administered three times per day over a three day period for a total dose of 360 mg/m² corresponding to a 180 mg body dose, a total pulmonary deposited dose of about 90 mg or about 2.25 mg/m² of lung surface area. One half of the low dose group dogs was necropsied the day after the final

20 exposure and the remaining half was necropsied four days later. All high dose dogs were necropsied the day after the final exposure.

Exposure to these extremely high doses resulted in the death of one high dose group dog after three days of exposure with the remaining three dogs euthanized in moderately debilitated to moribund conditions. This

25 dose intensive treatment caused pulmonary edema, a sequela of microscopically recognizable degeneration, necrosis and inflammation of epithelial surfaces lining the bronchials and larynx and the mucosal surfaces of the nose and lips. These lesions were life threatening and more severe in the high dose group, but were considered survivable at the lower dose, based

30 on the clinical condition of the animals. Despite these higher doses, there were no clinical pathology changes indicative of doxorubicin induced myelosuppression. There was microscopic evidence of lymphoid depletion in

the regional lymph nodes of the respiratory and gastrointestinal tracts suggestive of regional drainage of free doxorubicin to the draining lymph nodes of the thoracic and GI systems. WBC values actually increased in the high dose group, a change associated with the inflammatory response

5 observed in the respiratory tract. There were no other clinical pathology changes of note other than increased serum alkaline phosphatase in the high dose group, a nonspecific change, due likely to respiratory tract tissue damage.

Generally, changes noted at the moderate and high dosages were
10 edema, increased white blood cell count and increased respiratory rate. Histopathology revealed thoracic and GI lymphoid depletion for the moderate and higher doses, respectively. Respiratory tract toxicity including airway epithelial degeneration and moderate to severe inflammation was noted at the increased dosages. Bioavailability was low to moderate indicating an
15 absorption rate limiting process in movement of the drug into the systemic circulation. The probable dose limiting toxicity of doxorubicin is expected to be respiratory tract toxicity rather than a systemic toxicity.

In addition, a dose escalation study was conducted on a weekly exposure schedule. Initial doses of 12 mg deposited were delivered via
20 endotracheal tube to the lungs, with a 5th weekly dose of 18 mg deposited within the lungs. This provided a total body dose of 24 to 36 mg/m². The results of this repeated dose trial were similar in character (but not in severity) to the higher dose tests. Animals survived this treatment regimen with minimal clinical evidence of toxicity and no evidence of systemic
25 changes. Histologically, there was no evidence of respiratory tract epithelial degeneration and inflammation.

TABLE 9 - Doxorubicin

Summary Results of Dog Feasibility and Dose Range-Finding Studies

Chemotherapy	IV dose	Inhalation Dose	Pulmonary Deposited Dose*	Clinical Condition	Clinical Pathology	Histopathology	Respiratory Tract Toxicity	Bioavailability	Probable Dose-Limiting Toxicity
Example 6F Doxorubicin Feasibility	20 mg/m ²	20 mg/m ² x 1	5 mg	No change	No change	No change	No change	Low-moderate (absorption rate limited)	Respiratory tract toxicity
		40 mg/m ² x 3 doses (120 mg/m ² total)	10 mg x 3 doses	Mild-moderate pulmonary edema ↑IRR	↑ WBC	Thoracic & GI Lymphoid depletion	Airway epithelial degeneration		
		120 mg/m ² x 3 doses (360 mg/m ² total)	30 mg x 3 doses	Marked edema ↑↑IRR	↑↑ WBC	Thoracic & GI Lymphoid depletion	Moderate-severe inflammation		
Example 6R Doxorubicin Dose Range-Finding	N/A	24-36 mg/m ² (5 wkly Rx)	12-18 mg per dose	↑IRR Mild transient pulmonary edema	↓ WBC	Mild-moderate thoracic and mesenteric lymphoid depletion	Mild-moderate degeneration of airway epithelium Mild-moderate interstitial inflam.	Low-moderate	Respiratory tract toxicity

5

↑ - Increase ↓ - Decrease IRR - increased respiratory rate WBC - white blood cells

* - Divide the pulmonary deposited dose in mg by 40 to get the pulmonary deposited dose in mg/m² of lung surface area.

Plasma levels of doxorubicin were dose dependent and exhibited clear evidence of drug accumulation, including daily increases in Cmax (maximum concentration in blood) and steady state-like profiles, suggesting there was a rate limited absorption from the lung into the blood with significant 5 accumulation of doxorubicin in the lungs following each additional exposure given at a frequency of daily intervals. This accumulation was considered likely responsible for the tissue damage observed.

Referring again to Tables 7 and 9, an inhalation dose range of 20-40 mg/m² was administered in five weekly doses that resulted in a body 10 exposure of about 10 mg to about 20 mg, a pulmonary deposited dose range of about 10 to about 20 mg or a range of about 0.25 mg/m² to about 0.5 mg/m² lung surface area. The clinical condition included increased respiratory rate and mild transient pulmonary edema. A decrease in white blood cell 15 count was noted for the higher dosages. Histopathology revealed mild to moderate thoracic and mesenteric lymphoid depletion. Respiratory tract toxicity noted was mild to moderate degeneration of airway epithelium. A mild to moderate to marked interstitial inflammation was noted with some limited fibrosis. Bioavailability was noted to be low to moderate with absorption being rate limited. The probable dose limiting toxicity appears 20 again to be respiratory tract toxicity.

Example 7F and 7R

Referring now to Table 10, 1.4 mg of vincristine was initially administered by IV. After the washout period one inhalation feasibility test 25 was made. The vincristine was formulated in a 50% water/ 50% ethanol vehicle. A single dose of 2.8 mg/m² of vincristine was administered that gave about a 1.8 mg body dose, a pulmonary deposited dose of about 0.9 mg or about 2.25 mg/m² lung surface area. No changes were noted in the animal from this dose.

TABLE 10 – Vincristine & Vinorelbine

Summary Results of Dog Feasibility and Dose Range-Finding Studies

Chemotherapy	IV dose	Inhalation Dose	Pulmonary Deposited Dose	Clinical Condition	Clinical Pathology	Histopathology	Respiratory Tract Toxicity	Bioavailability	Probable Dose-Limiting Toxicity
Example 7F Vincristine Feasibility	1.4 mg/m ²	2.8 mg/m ² x 1	0.7 mg	Normal	Normal	No change	No change	Undetermined	Undetermined
Example 7R Vincristine Dose Range-Finding	N/A	1.1-3.0 mg/m ² (6 wkly Rx)	0.55-1.5 mg/dose	Normal	↓ WBC	Minimal-mild bone marrow and lymphoid depletion	Minimal interstitial inflammation	Undetermined	Myelosuppression
Example 7R Vinorelbine Dose Range-Finding	N/A	12-30 mg/m ² (5 wkly Rx)	6-15 mg per dose	Normal	↓ WBC	Bone marrow and lymphoid depletion	Minimal pulmonary and airway inflam.	Undetermined	Myelo-suppression

54 ↑ - increased ↓ - decrease IRR - increased respiratory rate WBC - white blood cell

5 * - Divide the pulmonary deposited dose in mg by 40 to get the pulmonary deposited dose in mg/m² of lung surface area.

Referring now to Tables 7 and 10, range finding tests of inhaled vincristine were made in the range of 0.5 to 1.5 mg of pulmonary deposited vincristine administered in six weekly doses. Therefore the amount of 5 pulmonary deposited dose ranged from about 12.5 - 37.5 $\mu\text{g}/\text{m}^2$ lung surface area. This corresponded to a total body dose of 50-150 $\mu\text{g}/\text{kg}$ or 1.0-3.0 mg/m^2 of body surface area. This dose range is near and generally above typical dose ranges for vincristine given IV. But in the examples given here, the entire dose was administered to the lungs. Vincristine is a potent drug 10 and causes significant myelosuppression and neurotoxicity at doses above 1.0 mg/m^2 given systemically. The results of the pilot inhalation studies showed the drug was well tolerated at all doses delivered by pulmonary administration with little to no evidence of respiratory tract toxicity with mild lymphoid depletion/myelosuppression only occurring at the highest doses given (2.0-3.0 15 mg/m^2).

Example 8R

Vinorelbine, which is also a vinca alkaloid was evaluated in a repeated exposure pilot tests. Compared to vincristine, vinorelbine was approximately 20 5-10 times less potent in producing toxicity, but produced similar types of changes. Vinorelbine delivered by pulmonary administration directly into the lungs of dogs by endotracheal tube, on a weekly basis (for 5 weeks) at escalating doses was well tolerated. A dose of 6 mg deposited in the lung was initially selected and escalated to 15 mg deposited within the lung. This 25 represented a lung surface exposure of ~0.15-0.375 mg/m^2 of lung surface area and total body doses of 12-30 mg/m^2 . This treatment regimen produced very minimal effects within the respiratory tract, characterized principally by slight inflammation. At the higher dose levels, inhaled vinorelbine produced sufficient blood levels to cause mild to moderate myelosuppression and 30 lymphoid depletion, both of which were reversible and of a severity, which was not life-threatening.

Examples 9F and 9R

An additional proof of concept, pilot inhalation tests involved etoposide.

Etoposide is a cytotoxic drug, representative of a class of drugs known as topoisomerase II inhibitors. Given orally or IV, etoposide causes typical

5 cytotoxic systemic toxicity, including myelosuppression, severe GI toxicity and alopecia. Etoposide is a highly insoluble drug and therefore difficult to formulate. The vehicle used clinically also causes adverse effects, predominantly anaphylactic type reactions.

In this invention, etoposide was reformulated in a novel vehicle, 10 dimethylacetamide (DMA) which does not cause anaphylactic reactions. While DMA cannot be used for IV administration due to systemic toxicity, it was shown to be a safe delivery vehicle for the pulmonary route of delivery. The etoposide was delivered in a 100% DMA vehicle. This formulation allowed the formation of the appropriate particle sizes. In these tests, escalating 15 doses of etoposide were given to dogs on a weekly schedule. The initial dose used was 25 mg of etoposide deposited in the pulmonary region with a 6th and final dose delivered of 80 mg deposited within the pulmonary region.

This equated to a dose range of 50-160 mg/m² of body surface area. This treatment regimen caused no systemic toxicity and only minimal inflammation 20 of the lung and no overt damage of the respiratory tract. In addition, there was good evidence of lymphoid depletion of the thoracic lymph nodes, in the absence of systemic changes, indicating that the drug was draining directly through the regional lymph system. This would provide additional regional therapeutic effectiveness in dealing with metastatic cells.

25 An additional pharmacokinetic test of inhaled etoposide showed the drug had moderately good bioavailability. A single inhaled total deposited dose of 260 mg/m² (about 65 mg of drug deposited in the pulmonary region) produced blood levels of etoposide similar to an IV dose of 50 mg/m² (see Figures 1-3). In other words, to reach similar blood concentrations 30 approximately 5X more drug was given by inhalation, a dose which caused neither respiratory tract nor systemic toxicity.

Example 10F

Additional proof of concept inhalation studies involved the cytotoxic drug 9-aminocamptothecin (9-AC) which is within the drug class known as 5 camptothecins. Like etoposide, 9-AC is insoluble and difficult to formulate. Supporting the concept and claims of this invention, the inventors generated aerosols of 9-AC formulated as a microsuspension in an aqueous vehicle (100% water).

These aerosols were delivered to dogs at daily doses of 40 mg/m^2 10 body surface area (10 mg of drug deposited within the pulmonary region) for 3 consecutive days. Inhalation treatment produced lower drug plasma levels than an IV dose of 10 mg/m^2 . The daily inhalation dose was 4 times greater than the IV dose and the total cumulative 3 day inhalation dose was 12 times greater than the single IV dose given (which causes mild systemic toxicity). 15 Despite the significantly greater doses given by inhalation, there were no measurable toxic effects (neither local effects within the respiratory tract nor systemic changes). Results from these tests supported the concept of improved overall safety and dose-intensification within the respiratory tract and also demonstrated the concept with aerosolized microsuspensions of 20 chemotherapeutic drugs.

Example 11F

In addition, this feasibility trial was extended to examine another 25 platinum-containing chemotherapeutic, carboplatin. The usual clinical formulation using water was used. Carboplatin is generally considered less toxic than cisplatin at comparable doses, and this appeared consistent with the results seen when the two agents were delivered by inhalation. Inhaled doses of up to 30 mg carboplatin deposited via endotracheal tube into the 30 lungs of dogs (60 mg/m^2 total body dose) caused no evidence of either direct respiratory tract or systemic toxicity.

TABLE 11 – Etoposide, & 9-Aminocamptothecin (9-AC)
Summary Results of Dog Feasibility and Dose Range-Finding Studies

Chemotherapy	IV dose	Inhalation Dose	Pulmonary Deposited Dose*	Clinical Condition	Clinical Pathology	Histopathology	Respiratory Tract Toxicity	Bioavailability	Probable Dose-Limiting Toxicity
Example 9F Etoposide Feasibility	50 mg/m ²	260 mg/m ² x 3 (780 mg/m ² total dose)	65 mg x 3 doses	Normal	No change	Mild thoracic lymphoid depletion	None	Moderate	Undetermined
Example 9R Etoposide Dose Range-Finding	N/A	50-160 mg/m ² (6 wkly Rx)	25-80 mg dose	Normal	No change	Mild-moderate thoracic lymphoid depletion	Mild interstitial inflammation	Moderate	Undetermined
Example 10F 9-AC Feasibility	10 mg/m ²	40 mg/m ² x 3 (120 mg/m ² total)	10 mg x3 doses	Normal	No change	Minimal lymphoid depletion	Minimal interstitial inflammation	Moderate-high	Undetermined

5 58 * - Divide the pulmonary deposited dose in mg by 40 to get the pulmonary deposited dose in mg/m² of lung surface area.

Examples 12 to 20

These examples illustrate results of clinical treatment of dogs having end stage lung cancer where other treatments have failed. For treatment, the dogs were anaesthetized and the inhalation treatment was through an

5 endotracheal tube.

This preliminary trial was performed to determine whether the inhalation chemotherapy treatment could be successfully used in animals with lung tumors. Initially, nine dogs with neoplastic lung disease were studied. Three different drugs were used- doxorubicin, vincristine, cyclophosphamide,

10 cisplatin, and paclitaxel at the doses and schedules summarized in Table 12.

One 16 year old mixed breed dog had no evidence of tumor in the lung following excision of a primary lung tumor, but did have evidence of metastases in the hilar lymph nodes, a sign that metastases would soon appear in the lung. However, the results showed that no metastases

15 developed in the lung for four months during which time the dog received four treatments of inhaled doxorubicin. In six other dogs, there were metastases in the lung and in each of these, the inhaled chemotherapy stopped the growth of the metastases, i.e. there was stable disease (or SD). In two dogs inhalational chemotherapy was not effective and there was

20 progressive disease (or PD). Since no chemotherapy was given to these dogs by the intravenous route, tumors outside of the lung progressed even while the lung tumors were stabilized. Thus, the results demonstrated that inhalational chemotherapy was effective in the local treatment of lung cancer in the dog.

TABLE 12
Summary of Preliminary Clinical Results in Dogs

Ex.	Dog Type and Age	Diagnosis	Inhalation Treatment*	Time of Trial	Results
12	Afghan 10 years old	Advanced lung carcinoma	Dox 5 mg, x 1	1 week	PD extrapulmonary
13	Cocker Spaniel 10-12 years old	Lung metastasis from excised melanoma	Dox 5 mg, x 2 Vincristine 0.5 mg, once	2 mo.	SD lung, died PD extrapulmonary, died
14	Beagle 7 years old	Thyroid carcinoma with lung metastasis	Dox 5 mg, x 4	4 mo.	SD lung PD thyroid & extrapulmonary, died
15	Labrador 8 years old	Thyroid carcinoma with lung metastasis	Dox 5 mg, x 2	2 mo.	SD lung PD thyroid & brain metastasis, died
16	Mixed Breed 16 years old	Excised lung primary, positive hilar lymph nodes	Dox 5 mg, x 4	4 mo.	No lung metastasis Death (CNS metastasis)
17	Rottweiler 3 years old	Excised distal osteosarcoma, lung nodule	Dox 7 mg, x 2 Cisplatin 15 mg, x 1	1 mo.	PD lung Further Rx declined
18	Mixed Breed 14 years old	Lung metastasis (carcinoma)	(Dox 5 mg + CTX 25 mg), x 3 Dox 5 mg, x 1	2-1/2 mo.	SD lung PD visceral & Extrapulmonary, died
19	Flat-coated Retriever 8 years old	Excised salivary adenocarcinoma, lung metastasis	Paclitaxel 22.5 mg, QW x 4	2-1/2 mo.	SD (4 weeks) lung PD lung, Rx discontinued
20	Husky 16 years old	Advanced mammary adenocarcinoma, lung metastasis	Paclitaxel 22.5 mg, x 2 (Paclitaxel 22.5 mg + Dox 5 mg), x 2	2 mo.	SD lung

* - Calculate target dose. Abbreviations: PD = progressive disease, SD = stable disease; Dox = Doxorubicin; CTX = cyclophosphamide; QW = every week;

Examples 21 to 33

Additionally, tests were conducted in dogs using a defined protocol. In these tests, dogs with either gross metastatic disease, micrometastatic hemangiosarcoma or micrometastatic primary lung cancer were randomized 5 to receive either doxorubicin, paclitaxel or both by inhalation via an endotracheal tube in a crossover design. Aerosol particle size was 2 - 3 μm as in the previous tests. The apparatus used was basically that shown in Figure 5 and as described above. Formulations for administration of the drugs were as follows: 16 mg/ml doxorubicin in 70%ethanol/30%water; 75 10 mg paclitaxel in about 30% PEG/70%ethanol. Preferably the paclitaxel is administered with 0.2% of citric acid to prevent degradation of the drug unless it is immediately used after preparation. The treatments were administered once every two weeks, and if a diagnosis of progressive disease was made on two consecutive intervals the dog was crossed over to the 15 alternate drug. At each treatment session, blood was sampled for hematology and biochemical analyses and urine was collected for analysis. The status of the tumors was monitored radiographically.

The results are summarized in Table 13. Pulmonary deposited doses listed in the table are based on scintigraphy studies that relate inhaled doses 20 to deposited doses. Among the 10 dogs that had gross metastatic disease (Examples 21-28), which is regarded as a terminal condition with a very short life expectancy, 4 dogs (in Examples 21, 22, 24, and 27) showed stable disease in the lung indicating that the drug was having a positive effect. In the remaining 6 dogs (see Examples 23, 25, 26, and 28), the lung disease 25 progressed. In two of the dogs with metastatic osteosarcoma (Examples 24 and 25) and in the dog with metastatic melanoma (Example 28), there were partial responses, i.e. there were tumors that decreased in size by more than 50%.

30 Four dogs had splenic hemangiosarcoma (Examples 29 and 30), a disease that invariably metastasizes to the lung and is fatal within two to four months. These dogs were given doxorubicin by inhalation in addition to intravenous chemotherapy to control systemic disease. The results in Table

13 show that each of the four dogs was alive (at least two months at the time of this writing) and that there was no evidence of disease in the lung.

The last group of dogs (Examples 31 - 33) are those that had primary lung tumors which were removed surgically. These dogs had metastases in 5 their thoracic lymph nodes and have a life expectancy measured in weeks. As shown in Table 13, two dogs (Examples 31 and 32) received doxorubicin by inhalation (1.5 mg) and two dogs (Example 33) received paclitaxel (20 mg). The dog that received five treatments of doxorubicin was alive with no evidence of disease 81 days later suggesting that the treatment is having a 10 positive effect. One dog (Example 32) received two doses of doxorubicin and died from metastases outside of the lung. The other two dogs (Example 33) have no evidence of disease but not enough time has passed to determine how effective the treatment will be.

The result of these tests, therefore, confirm those of the preliminary 15 tests that inhalational chemotherapy is effective in the treatment of lung cancer.

Table 13
Efficacy of Inhalational Chemotherapy in Dogs with Lung Cancer

Ex.	Diagnosis	No. of Dogs	Inhalation Treatment*	Results
21	Lung carcinoma	1	DOX 5 mg (5x) then paclitaxel 60 mg (2x)	SD
22	Metastatic	1	DOX 5 mg (2x)	SD
23	hemangiosarcoma	1	DOX 5 mg (1x)	PD
24	Metastatic osteosarcoma	1	DOX 5 mg (5x) + paclitaxel 60 mg (2x)	SD (PR after 3 rd DOX treatment)
25	"	3	DOX 5mg (2x) + paclitaxel 60 mg (1x)	PD (PR in one dog)
26	Metastatic fibrosarcoma	1	DOX 5 mg (2x)	PD
27	Metastatic liposarcoma	1	DOX 5 mg (4x) + paclitaxel 60 mg (1x)	SD
28	Metastatic melanoma	1	paclitaxel 60 mg (2x) + DOX 5 mg (1x)	PD (PR noted in nodules < 2 cm)
29	Splenic hemangiosarcoma	2	DOX 5mg (4x) + systemic chemotherapy	Alive and NED
30	"	2	DOX 1.5 mg(3x) + systemic chemotherapy	Alive and NED
31	Primary lung tumor	1	DOX 1.5 mg (5x)	Alive and NED
32	excised-micrometastatic	1	DOX 1.5 mg (2x)	Dead from extrapleural metastases
33	disease	2	paclitaxel 20 mg (1x)	Alive and NED

* - Deposited pulmonary doses

DOX = doxorubicin; (x) = number of treatments received; SD = stable disease; PD = progressive disease; NED = no evidence of disease; PR = partial response (50% decrease in tumor size)

The safe and effective range of doses of the inhalant antineoplastic drugs in humans and animals (e.g. dogs and similar small animals) are shown in Table 14 below. Larger animal dosages can be calculated by using multiples of the small animal based dose based on the known relationship of

(body weight in kg/m² of body surface area. The exact doses ~~will~~ vary depending upon such factors as the type and location of the tumor, the age and size of the patient, the physical condition of the patient and concomitant therapies that the patient may require. The dosages shown are for doses for 5 one course of therapy. A course of therapy may be given, monthly, weekly, biweekly, triweekly or daily depending on the drug, patient, type of disease, stage of the disease and so on. Exemplary safe and effective amounts of carrier are given for each product have been published by the respective manufacturer and are summarized in the Physicians Desk Reference.

10

Table 14

Drug	Animal Dose* mg/m ²	Human Dose* mg/m ²
Doxorubicin	2 to 90	3 to 130
Paclitaxel	6 to 270	10 to 400
Vincristine	0.06 to 2	0.1 to 3
Vinorelbine	1.3 to 60	2 to 90
Cisplatin	4.6 to 200	7 to 300
Etoposide	4.6 to 200	7 to 300
9-Aminocamptothecin	2.6 to 10	0.04 to 15

* - m² body surface area

Based on the results of the inhalation tests herein with doxorubicin, 15 inhalation treatments with anthracyclines in addition to doxorubicin are also expected to be well tolerated and efficacious when administered by the pulmonary route. Based on the inhalation tests herein with vincristine and vinorelbine, other vinca alkaloids are expected to be well tolerated and efficacious when administered by the pulmonary route. Based on the 20 inhalation tests herein for the vesicants doxorubicin, vincristine, vinorelbine, and paclitaxel, all of which are capable of serious vesicating injuries, other vesicating drugs (e.g. mechlorethamine, dactinomycin, mithramycin, bisantrene, amsacrine, epirubicin, daunorubicin, idarubicin, vinblastine,

vindesine, and so on) are expected to be well tolerated and efficacious when administered by the pulmonary route. The exception, of course, would be vesicant drugs that are known to exhibit significant pulmonary toxicity when administered by IV (e.g. mitomycin-C). In this regard, a safe and effective 5 amount of a particular drug or agent is that amount which based on its potency and toxicity, provides the appropriate efficacy/risk balance when administered via pulmonary means in the treatment of neoplasms. Similarly a safe and effective amount of a vehicle or carrier is that amount based on its solubility characteristics, stability, and aerosol forming characteristics, that 10 provides the required amount of a drug to the appropriate site in the pulmonary system for treatment of the neoplasm.

For the nonvesicant antineoplastic drugs, based on the inhalation tests herein for the vesicating and nonvesicating drugs it is expected that all the nonvesicating drugs that do not exhibit direct pulmonary toxicity when 15 administered intravenously are expected be well tolerated and exhibit efficacy. Bleomycin and mitomycin-C, for example, exhibit sufficient pulmonary toxicity to be excluded except when a chemoprotectant is used. In this regard typically carmustine, dacarbazine, melphalan, methotrexate, mercaptopurine, mitoxantrone, esorubicin, teniposide, aclacinomycin, 20 plicamycin, streptozocin, menogaril are expected to be well tolerated and exhibit efficacy. Similarly, drugs of presently unknown classification such as geldanamycin, bryostatin, suramin, carboxyamido-triazoles such as those in US patent 5,565,478, onconase, and SU101 and its active metabolite SU20 are likewise expected to be well tolerated and exhibit efficacy subject to the 25 limitation on pulmonary toxicity. These drugs would be administered by the same methods disclosed for the tested antineoplastic drugs. They would be formulated with a safe and effective amount of a vehicle and administered in amounts and in a dosing schedule safe and effective for treating the neoplastic disease.

30 Pulmonary toxicity of compounds to be administered by inhalation is an important consideration. As mentioned above one of the major considerations is whether the drug exhibits significant pulmonary toxicity

when injected by IV. While almost all antineoplastic drugs are toxic to the body and thus arguably exhibit pulmonary toxicity if given in a large enough dose, the test for pulmonary toxicity as used herein requires significant pulmonary toxicity at the highest manufacturers recommended dose that is to 5 be administered to a patient. The determination of whether a drug exhibits sufficient pulmonary toxicity by IV so as to exclude it from the group of drugs useful for pulmonary administration can be made from the drug manufacturers recommendations as published in the Physicians Desk Reference (see "Physicians Desk Reference" 1997, (Medical Economics Co.), 10 or later editions thereof), in other drug manuals published for health care providers, publicly available filings of the manufacturer with the FDA, or in literature distributed directly by the manufacturers to physicians, hospitals, and the like. For example in the "Physicians Desk Manual" 1997:

- Doxorubicin (Astra) pp. 531-533 - vesicant, there is no indication of 15 pulmonary toxicity while cardiac toxicity, hematologic toxicity particularly leukopenia and myelosuppression; extravasation injuries are also noted;
- Idarubicin (Pharmacia & Upjohn) pp 2096-2099 - vesicant, primary toxicity appears to be myelosuppression no mention is made of pulmonary toxicity making the drug useful in the present invention;
- Etoposide (Astra) pp539-541 - no indication of pulmonary toxicity, but 20 dose limiting hematologic toxicity is important;
- Paclitaxel (Bristol-Meyers Squibb) pp. 723-727 - vesicant, pulmonary toxicity is not listed for paclitaxel, but dose limiting bone marrow suppression (primarily neutropenia) is important;
- Bleomycin (Blenoxane® Bristol-Meyers Squibb) pp. 697-699 , pulmonary 25 toxicities occur in about 10% of treated patients by IV administered drug, this makes bleomycin unacceptable for pulmonary administration for the present invention;
- Mitomycin C (Mutamycin® Bristol-Meyers Squibb) - vesicant, infrequent 30 but severe life threatening pulmonary toxicity has occurred by IV administration, this although infrequent severe life threatening pulmonary toxicity shows that the drug exhibits substantial pulmonary toxicity;

- Methotrexate (Immunex) pp. 1322-1327 - MW=454, primary toxicity appears to be hepatic and hematologic, signs of pulmonary toxicity should be closely monitored for signs of lesions;
- Dactinomycin (Merck & Co.) - vesicant, primary toxicity appears to be oral, 5 gastrointestinal, hematologic, and dermatologic; no mention is made of pulmonary toxicity making the drug acceptable in the present invention;
- mechlorethamine (Merck & Co.) - vesicant, primary toxicity appears to be renal, hepatic and bone marrow, no mention is made of pulmonary toxicity making the drug acceptable in the present invention;
- 10 • Irinotecan (Camptosar® Pharmacia & Upjohn) - a derivative of camptothecin, primary toxicity appears to be severe diarrhea and neutropenia, no mention is made of pulmonary toxicity making the drug useful in the present invention;
- Vincristine (Oncovin® Lilly) pp. 1521-1523 - extremely toxic with high 15 vesicant activity found in the tests herein, but no pulmonary toxicity noted;
- Vinblastine (Velban® Lilly) pp. 1537-1540 - extremely toxic with high vesicant activity found in the tests herein, but no pulmonary toxicity noted.

The above listing is exemplary only and is not intended to limit the scope of 20 the invention.

An additional embodiment of the invention includes methods and formulations that contain chemoprotectants and are administered by inhalation for preventing toxicity and particularly pulmonary toxicity that may be elicited by antineoplastic drugs. The method would allow the use by 25 inhalation of antineoplastic drugs that exhibit pulmonary toxicity or would reduce the likelihood of pulmonary toxicity. One method would include treating a patient having a neoplasm, via inhalation administration, a pharmaceutically effective amount of a highly toxic antineoplastic drug and a pharmaceutically effective amount of a chemoprotectant, wherein the 30 chemoprotectant reduces or eliminates toxic effects in the patient that are a result of inhaling the highly toxic antineoplastic drug. More narrowly, another embodiment includes a combination of inhaled chemoprotectant and

antineoplastic drug that reduces or eliminates respiratory tract or pulmonary tract toxicity in the patient. The chemoprotectant can be coadministered with the antineoplastic drug by inhalation, or both by inhalation and by IV, or the chemoprotectant can be administered alone.

5 It is known, for example, that dextrazoxane (ICRF-187) when given by intraperitoneal injection to mice is protective against pulmonary damage induced by bleomycin given by subcutaneous injections. See for example Herman, Eugene et al, "Morphologic and morphometric evaluation of the effect of ICRF-187 on bleomycin-induced pulmonary toxicity", Toxicology 98, 10 (1995) pp. 163-175, the text of which is incorporated by reference as if fully rewritten herein. The mice pretreated with intraperitoneal injections of dextrazoxane prior to having bleomycin injected subcutaneously showed reduced pulmonary alterations particularly fibrosis compared to another group of mice that was not pretreated.

15 The following examples illustrate the use of a chemoprotectant by inhalation in conjunction with an antineoplastic drug.

Example 34

Dextrazoxane (ICRF-187) is dissolved in a pharmaceutically acceptable liquid formulation and administered to a patient as an aerosol using the apparatus and methods described herein, at a dose ranging from 10 mg to 20 1000 mg over a period of from one minute to one day prior to giving a chemotherapeutic drug such as doxorubicin by inhalation. The doxorubicin is given in a dose from 1 mg to 50 mg.

25

Example 35

Dextrazoxane (ICRF-187) is administered as described in Example 34 at the same time or up to two hours before giving bleomycin by intravenous injection. The dose of dextrazoxane ranges from about 2 times to about 30 30 times the dose of bleomycin. The dose of bleomycin by IV ranges from about 5 to 40 units/m².

Example 36

Dexrazoxane (ICRF-187) is administered as described in Example 34 at the same time or up to two hours before administering bleomycin by inhalation. The dose of dexrazoxane ranges from about 2 times to about 30 times the 5 dose of bleomycin. The dose of bleomycin by inhalation ranges from 5 to 40 units/m² at intervals of from 1 week to 4 weeks.

Examples 37 and 38

10 Chemoprotectants such as mesna (ORG-2766), and ethiofos (WR2721) may be used in a manner similar to that described in Examples 34 to 36, above.

Combination Therapy

Another embodiment of the invention contemplates drug coadministration by the pulmonary route, and by (1) other local routes, 15 and/or (2) systemically by IV. Results from the clinical tests on dogs indicates that, although the pulmonary route of administration will indeed control neoplastic cells arising in or metastatic to the pulmonary tract, neoplastic cells elsewhere in the body may continue to proliferate. This embodiment provides for effective doses of drug in the lung delivered via the lung and additional 20 drug delivered via (1) other local sites (e.g. liver tumors may also be treated via hepatic artery instillation, ovarian cancer by intraperitoneal administration) and/or additional drug(s) may be provided systemically by IV via the general circulatory system. Administration can be at the same time, or administration followed closely in time by one or more of the other therapeutic routes. 25 Benefits are that much higher dosages can be supplied to affected tissues and effective control of neoplasms can be maintained at multiple critical sites compared to using a single mode of administration.

Also contemplated within the scope of the invention is the combination 30 of drugs for combination chemotherapy treatment. Benefits are those well known in the treatment of cancer using combination chemotherapy by other routes of administration. For example, combining drugs with different mechanisms of action such as an alkylating agent plus a mitotic poison plus a

topoisomerase inhibitor. Such combinations increase the likelihood of destroying tumors that are comprised of cells with many different drug sensitivities. For example, some are easily killed by alkylating agents while mitotic poisons kill others more easily.

5 Also included in the invention are embodiments comprising the method for inhalation therapy disclosed herein and the application of radiotherapy, gene therapy, and/or immunotherapy. Other embodiments include the immediately above method combined with chemotherapy applied by IV and/or local therapy.

10 Also included within the invention are formulations for paclitaxel. In these formulations 100% to 40% ethanol is useful. However, to obtain better control of particle size and stable aerosol generation the addition of polyethylene glycol (PEG) is preferred. Although 1-60% PEG can be used, 8-40% PEG is more preferred, and 10-30% PEG was found to be optimal. A
15 further embodiment also includes the addition of 0.01 to 2% of an organic or inorganic acid, preferably an organic acid such as citric acid and the like. The acid being added to stabilize the formulation. With regard to clinical use in inhalation, citric acid in water has been found to cause tussive and bronchoconstrictive effects. PEG may ameliorate this effect. The formulation
20 contains a safe and effective amount of paclitaxel useful for the treatment of neoplasms.

While the forms of the invention herein disclosed constitute presently preferred embodiments, many others are possible. It is not intended herein to mention all of the possible equivalent forms or ramifications of the
25 invention. It is to be understood that the terms used herein are merely descriptive, rather than limiting, and that various changes may be made without departing from the spirit of the scope of the invention.

Throughout the description and claims of the specification the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.



The claims defining the invention are as follows:

1. A method for treating a patient having a neoplasm comprising: administering a pharmaceutically safe and effective amount of non-encapsulated doxorubicin to said patient by inhalation, wherein said safe and effective amount provides an animal dose of about 2 to 90 mg/m² and a human dose of about 3 to 130 mg/m², wherein both doses are based on body surface area.
2. The method according to claim 1, wherein said doxorubicin is inhaled in the absence of perfluorocarbon.
3. The method according to claim 1, wherein said neoplasm is a pulmonary neoplasm, a neoplasm of the head and neck, or other systemic neoplasm.
4. The method according to claim 1, wherein said drug is inhaled as a liquid aerosol or as a powdered aerosol.
5. A method for treating a patient having a neoplasm comprising: administering a pharmaceutically safe and effective amount of non-encapsulated vincristine or vinorelbine to said patient by inhalation, and wherein said safe and effective amount when vincristine is selected provides an animal dose of about 0.06 to 2 mg/m² and a human dose of about 0.1 to 3 mg/m²; and when vinorelbine is selected said safe and effective amount provides an animal dose of about 1.3 to about 60 mg/m² and a human dose of about 2 to 90 mg/m², wherein all doses are based on body surface area.
6. A method for treating a patient having a neoplasm comprising: administering a pharmaceutically effective amount of nonencapsulated paclitaxel to said patient by inhalation, and wherein said safe and effective amount provides an animal dose of about 6 to 270 mg/m², and a human dose of about 10 to 400 mg/m², wherein both doses are based on body surface area.
7. A method for treating a patient having a neoplasm comprising: administering to said patient by inhalation a formulation comprising doxorubicin in a pharmaceutically acceptable carrier at a safe and effective dose, wherein said administering of said safe and effective dose provides an animal dose of about 2 to 90 mg/m² or a human dose of about 3 to 130 mg/m², and wherein both doses are based on body surface area.
8. A method for treating a patient having a neoplasm comprising:



administering to said patient by inhalation a formulation comprising vincristine or vinorelbine in a pharmaceutically acceptable carrier at a safe and effective dose, wherein said administering of said safe and effective dose when vincristine is selected provides an animal dose of about 0.06 to 2mg/m² and a 5 human dose of about 0.1 to 3 mg/m²; and when vinorelbine is selected an animal dose of about 1.3 to about 60 mg/m² and a human dose of about 2 to 90 mg/m², wherein all doses are based on body surface area.

9. A method for treating a patient having a neoplasm comprising:
administering to said patient by inhalation a formulation comprising paclitaxel in
10 a pharmaceutically acceptable carrier at a safe and effective dose, wherein said
administering of said safe and effective dose provides an animal dose of about
6 to 270 mg/m², and a human dose of about 10 to 400 mg/m², and wherein both
doses are based on body surface area.

10. A method according to any one of the preceding claims substantially as
15 hereinbefore described with reference to any one of the examples.

DATED: 10 September 2001

PHILLIPS ORMONDE & FITZPATRICK

Attorneys for:

20 BATTELLE MEMORIAL INSTITUTE



dog 1 O)




FIG 1




FIG. 2






FIG. 3









FIG. 6



F, G. 7