

(19) **DANMARK**

(10) **DK/EP 3013984 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 Q 1/6853 (2018.01)** **C 12 Q 1/6841 (2018.01)**
- (45) Oversættelsen bekendtgjort den: **2023-06-06**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-03-22**
- (86) Europæisk ansøgning nr.: **14818012.8**
- (86) Europæisk indleveringsdag: **2014-06-25**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-05-04**
- (86) International ansøgning nr.: **US2014044196**
- (87) Internationalt publikationsnr.: **WO2014210225**
- (30) Prioritet: **2013-06-25 US 201361839320 P** **2013-06-25 US 201361839313 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Prognosys Biosciences, Inc., 11125 Flintkote Avenue, Suite A, San Diego, CA 92121, USA**
- (72) Opfinder: **CHEE, Mark, 11125 Flintkote Avenue, Suite A, San Diego, CA 92121, USA**
ROUTENBERG, David, 11125 Flintkote Avenue, Suite A, San Diego, CA 92121, USA
- (74) Fuldmægtig i Danmark: **Gotapatent AB, Box 3127, S-550 03 Jönköping, Sverige**
- (54) Benævnelse: **METODE TIL BESTEMMELSE AF SPATIALE MØNSTRE I BIOLOGISKE TARGETS I EN PRØVE**
- (56) Fremdragne publikationer:
WO-A2-2004/028955
WO-A2-2010/027870
US-A1- 2003 162 216
US-A1- 2011 245 111
US-A1- 2012 195 810
US-A1- 2012 270 748
CHEN L ET AL: "DNA hybridization detection in a microfluidic channel using two fluorescently labelled nucleic acid probes", BIOSENSORS AND BIOELECTRONICS, ELSEVIER BV, NL, vol. 23, no. 12, 15 July 2008 (2008-07-15) , pages 1878-1882, XP022666057, ISSN: 0956-5663, DOI: 10.1016/J.BIOS.2008.02.013 [retrieved on 2008-03-10]
DAVID A. ROUTENBERG ET AL: "Microfluidic probe: a new tool for integrating microfluidic environments and electronic wafer-probing", LAB ON A CHIP: MINIATURISATION FOR CHEMISTRY, PHYSICS, BIOLOGY, MATERIALS SCIENCE AND BIOENGINEERING, vol. 10, no. 1, 1 January 2010 (2010-01-01), pages 123-127, XP055302331, GB ISSN: 1473-0197, DOI: 10.1039/B913912E
ROUTENBERG ET AL.: 'Microfluidic probe: a new tool for integrating microfluidic environments and electronic wafer-probing' LAB CHIP vol. 10, 29 October 2009, pages 123 - 127, XP055302331

DESCRIPTION

Technical Field

[0001] The present disclosure generally relates to assays of biological molecules, and in particular, to methods, compositions, and assay systems for determining spatial patterns of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample.

Background

[0002] In the following discussion, certain articles and methods are described for background and introductory purposes. Nothing contained herein is to be construed as an "admission" of prior art. Applicant expressly reserves the right to demonstrate, where appropriate, that the articles and methods referenced herein do not constitute prior art under the applicable statutory provisions.

[0003] Comprehensive gene expression analysis and protein analysis have been useful tools in understanding mechanisms of biology. Use of these tools has allowed the identification of genes and proteins involved in development and in various diseases such as cancer and autoimmune disease. Conventional methods such as *in situ* hybridization and other multiplexed detection of different transcripts have revealed spatial patterns of gene expression and have helped shed light on the molecular basis of development and disease. Other technologies that have enabled the quantitative analysis of many RNA sequences per sample include microarrays (see Shi et al., *Nature Biotechnology*, 24(9): 1151-61 (2006); and Slonim and Yanai, *Plos Computational Biology*, 5(10):e1000543 (2009)); serial analysis of gene expression (SAGE) (see Velculescu et al., *Science*, 270(5235):484-87 (1995)); high-throughput implementations of qPCR (see Spurgeon et al., *Plos ONE*, 3(2):e1662 (2008)); *in situ* PCR (see Nuovo, *Genome Res.*, 4:151-67 (1995)); and RNA-Seq (see Mortazavi et al., *Nature Methods*, 5(7):621-8 (2008)). As useful as these methods are, however, they do not enable simultaneous measurement of the expression of many genes or the presence and/or activity of multiple proteins at many spatial locations in a sample.

[0004] Laser capture microdissection has permitted the analysis of many genes at a small number of locations, but it is very expensive, laborious, and does not scale well. Certain PCR assays in a 2D format preserve spatial information (see Armani et al., *Lab on a Chip*, 9(24):3526-34 (2009)), but these methods have low spatial resolution because they rely on physically transferring tissues into wells, which also prevents random access to tissue samples and high levels of multiplexing.

[0005] US 2011/245111 discloses a method to provide high-resolution spatial maps of

biological targets in a tissue within a 2D pattern by using a specific combination of identity probes, adapters and tags as well as location tags to define the coordinates of the target.

[0006] At present, there is a need to analyze at high resolution the spatial expression patterns of large numbers of genes, proteins, or other biologically active molecules simultaneously. There is also a need for reproducible, high-resolution spatial maps of biological molecules in tissues. The present disclosure addresses these needs.

Summary

[0007] The invention relates to a method of determining a spatial pattern of abundance, and/or expression of one or more protein targets across multiple sites in a 2D sample, as defined in claim 1.

[0008] In another aspect but not part of the invention is a method of determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample, comprising:

delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises: (1) a target-binding moiety capable of binding to the probe's corresponding biological target; and (2) an identity tag that identifies the probe's corresponding biological target or target-binding moiety;

allowing each probe to bind to its corresponding biological target in the sample;

delivering an address tag to each of the multiple sites in the sample, wherein the address tag is to be coupled to the probe bound to the biological target and identifies the site to which the address tag is delivered;

analyzing the probe/address tag conjugate bound to the one or more biological targets, the analysis comprising: (1) determining abundance, expression, and/or activity of each of the one or more biological targets by assessing the amount of the probe/address tag conjugate bound to the biological target; and (2) determining the identities of the identity tag and the address tag of the probe/address tag conjugate; and

determining a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In some embodiments, the method does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample. In one embodiment, the probe/address tag conjugate bound to the one or more biological targets may be analyzed by sequencing, wherein the amount of a sequencing product indicates abundance, expression, and/or activity of each of the one or more biological targets, and the sequencing product may comprise all or a portion of the address tag sequence and all or a portion of the identity tag sequence.

[0009] In any of the preceding aspects or combinations thereof, the one or more biological targets can be non-nucleic acid molecules. In the invention, the one or more biological targets are a protein. In any of the preceding embodiments, there can be at least two address tags that identify each of the multiple sites in the sample.

[0010] In any of the preceding embodiments, the spatial patterns of abundance, expression, and/or activity of multiple biological targets can be determined in parallel, and the address tag or combination of address tags may be the same for each of the multiple biological targets at a given site of the multiple sites in the sample. In any of the preceding embodiments, the analyzing step may be performed in parallel in the same reaction run.

[0011] In any of the preceding embodiments or combinations thereof, the one or more biological targets may include at least one known marker for the sample, for example, a tissue-specific marker, a cell type marker, a cell lineage marker, a cell morphology marker, a cell cycle marker, a cell death marker, a developmental stage marker, a stem cell or progenitor cell marker, a marker for a differentiated state, an epigenetic marker, a physiological or pathophysiological marker, a marker for a transformed state, a cancer marker, or any combination thereof.

[0012] In yet another aspect but not part of the invention is a method of determining a spatial pattern of abundance, expression, and/or activity of a target protein across multiple sites in a sample, comprising:

delivering a probe for a target protein to multiple sites in a sample, wherein the probe comprises: (1) a target-binding moiety capable of binding to the target protein; (2) a first address tag that identifies each of the multiple sites to which the probe is delivered; and (3) an identity tag that identifies the target protein or the target-binding moiety;

allowing the probe to bind to the target protein in the sample;

analyzing the probe bound to the target protein, the analysis comprising: (1) determining abundance, expression, and/or activity of the target protein by assessing the amount of the probe bound to the target protein; and (2) determining the identities of the identity tag and the first address tag of the probe for the target protein; and

determining a spatial pattern of abundance, expression, and/or activity of the target protein across the multiple sites in the sample based on the analysis.

[0013] In any of the preceding aspects, the method may further comprise:

delivering a probe for a target polynucleotide to each of the multiple sites in the sample, wherein the probe for the target polynucleotide comprises: (1) a sequence that hybridizes to

and identifies the target polynucleotide; (2) a second address tag that identifies each of the multiple sites to which the probe for the target polynucleotide is delivered;

allowing the probe for the target polynucleotide to bind to the target polynucleotide in the sample;

analyzing the probe bound to the target polynucleotide, the analysis comprising: (1) determining abundance, expression, and/or activity of the target polynucleotide by assessing the amount of the probe bound to the target polynucleotide; and (2) determining the identities of the sequence that hybridizes to and identifies the target polynucleotide and the second address tag of the probe for the target polynucleotide; and

determining a spatial pattern of abundance, expression, and/or activity of the target polynucleotide across the multiple sites in the sample based on the analysis of the probe bound to the target polynucleotide at each of the multiple sites in the sample.

[0014] In another aspect but not part of the invention is a method of determining a spatial pattern of abundance, expression, and/or activity of a target protein across multiple sites in a sample, comprising:

delivering a probe for a target protein to multiple sites in the sample, wherein the probe comprises: (1) a target-binding moiety capable of binding to the target protein; and (2) an identity tag that identifies the target protein or the protein-binding moiety;

allowing the probe to bind to the target protein in the sample;

delivering a first address tag to each of the multiple sites in the sample, wherein the first address tag is to be coupled to the probe bound to the target protein and identifies the site to which it is delivered;

analyzing the probe/first address tag conjugate bound to the target protein, the analysis comprising: (1) determining abundance, expression, and/or activity of the target protein by assessing the amount of the probe/first address tag conjugate bound to the target protein; and (2) determining the identities of the identity tag and the first address tag of the probe/first address tag conjugate; and

determining a spatial pattern of abundance, expression, and/or activity of the target protein across the multiple sites in the sample based on the analysis.

[0015] In any of the preceding aspects, the method may further comprise:

delivering a probe for a target polynucleotide to each of the multiple sites in the sample, wherein the probe for the target polynucleotide comprises a sequence that hybridizes to and identifies the target polynucleotide;

allowing the probe for the target polynucleotide to bind to the target polynucleotide in the sample;

delivering a second address tag to each of the multiple sites in the sample, wherein the second address tag is to be coupled to the probe bound to the target polynucleotide and identifies the site to which it is delivered;

analyzing the probe/second address tag conjugate bound to the target polynucleotide, the analysis comprising: (1) determining abundance, expression, and/or activity of the target polynucleotide by assessing the amount of the probe/second address tag conjugate bound to the target polynucleotide; and (2) determining the identities of the sequence that hybridizes to and identifies the target polynucleotide and the second address tag of the probe/second address tag conjugate; and

determining a spatial pattern of abundance, expression, and/or activity of the target polynucleotide across the multiple sites in the sample based on the analysis of the probe/second address tag conjugate bound to the target polynucleotide at each of the multiple sites in the sample.

[0016] In one aspect, the target polynucleotide or the complement thereof may encode all or a portion of the target protein. In some aspects, the step of analyzing the probe or probe/first address tag conjugate bound to the target protein and the step of analyzing the probe or probe/second address tag conjugate bound to the target polynucleotide may be performed in parallel in the same reaction run. In other aspects, the first address tag and the second address tag may be the same for a given site of the multiple sites in the sample. In yet other aspects, the first address tag and the second address tag can be different for a given site of the multiple sites in the sample. In any of the preceding embodiments, the method may further comprise associating abundance, expression, and/or activity of the target protein to abundance, expression, and/or activity of the target polynucleotide at each of the multiple sites in the sample.

[0017] In any of the preceding aspects or any combinations thereof, the biological target or the target protein may comprise an enzyme activity. In certain aspects, the target-binding moiety of the probe in any of the preceding aspects may comprise an antibody or an antigen binding fragment thereof, an aptamer, a small molecule, an enzyme substrate, a putative enzyme substrate, an affinity capture agent, or a combination thereof.

[0018] In any of the preceding aspects or any combinations thereof, the target-binding moiety is conjugated to a polynucleotide comprising the identity tag. In any of the preceding aspects, the target-binding moiety may be conjugated to a polynucleotide capable of specifically hybridizing to a polynucleotide comprising the identity tag. In certain aspects, the probe may comprise a multiplicity of target-binding moieties capable of binding to the same domain or different domains of the target, or capable of binding to different targets.

[0019] In any of the preceding aspects or any combinations thereof, the sample can be a biological sample selected from the group consisting of a freshly isolated sample, a fixed sample, a frozen sample, an embedded sample, a processed sample, or a combination thereof.

[0020] In any of the preceding aspects or any combinations thereof, there can be two address tags that identify each of the multiple sites in the sample. In certain aspects, two probes for each target can be delivered to the sample.

[0021] In any of the preceding aspects or any combinations thereof, the address tag may comprise an oligonucleotide. In another aspect, the identity tag of any of the preceding aspects may comprise an oligonucleotide.

[0022] In any of the preceding aspects or any combinations thereof, the analyzing step may be performed by nucleic acid sequencing. In one aspect, the analyzing step can be performed by high-throughput digital nucleic acid sequencing.

[0023] In any of the preceding aspects or any combinations thereof, the product of the number of the target(s) being assayed and the number of the multiple sites being assayed in the sample can be greater than 20, greater than 50, greater than 75, greater than 100, greater than 1,000, greater than 10,000, greater than 100,000, or greater than 1,000,000.

[0024] In any of the preceding aspects or any combinations thereof, at least one hundred thousand, at least five hundred thousand, or at least one million probes or probe/address tag conjugates bound to the target(s) may be analyzed in parallel.

[0025] In any of the preceding aspects or any combinations thereof, software programmed hardware may perform at least two steps of the delivering step(s), the analyzing step(s) and the determining step(s). In any of the preceding aspects or any combinations thereof, one or more microfluidic devices may be used to perform the delivering step(s).

[0026] In any of the preceding aspects or any combinations thereof, a known percentage of the probe for the biological target, the probe for the target protein, or the probe for the target polynucleotide can be an attenuator probe. In one aspect, the attenuator probe may limit production of an amplifiable product. For example, an attenuator probe may compete with an active probe for binding to the target. While an active probe can lead to the generation of an amplifiable product from the target, an attenuator probe does not or has reduced ability in generating an amplifiable product. In one aspect where a nucleic acid probe is used, the attenuator probe can lack a 5' phosphate.

[0027] In any of the preceding aspects or any combinations thereof, the address tag may be coupled to the probe by ligation, by extension, by ligation following extension, or any combination thereof.

[0028] In any of the preceding aspects or any combinations thereof, the method may further comprise constructing a 3-dimensional pattern of abundance, expression, and/or activity of each target from spatial patterns of abundance, expression, and/or activity of each target of multiple samples. In one aspect, the multiple samples can be consecutive tissue sections of a 3-dimensional tissue sample.

[0029] In yet another aspect but not part of the invention is a system for determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample, comprising:

a first module for delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises: (1) a target-binding moiety capable of binding to the probe's corresponding biological target; and (2) an identity tag that identifies the probe's corresponding biological target or target-binding moiety;

a second module for delivering an address tag to each of the multiple sites in the sample, wherein the address tag is to be coupled to the probe bound to the biological target and identifies the site to which the address tag is delivered;

a third module for analyzing the probe/address tag conjugate bound to the one or more biological targets, the analysis comprising: (1) determining abundance, expression, and/or activity of the one or more biological targets by assessing the amount of the probe/address tag conjugate bound to the biological target; and (2) determining the identities of the identity tag and the address tag of the probe/address tag conjugate; and

a fourth module for determining a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In one aspect, the system does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample.

[0030] In one embodiment, the second module may comprise one or more microfluidic devices for delivering the address tags. In one aspect, the one or more microfluidic devices may comprise a first set of multiple addressing channels, each delivering a different first address tag to the sample. In one aspect, the one or more microfluidic devices may further comprise a second set of multiple addressing channels, each delivering a different second address tag to the sample. In one aspect, the multiple sites in the sample can be chosen by the first and second set of multiple addressing channels cooperatively delivering the first address tags and the second address tags, respectively, to each of the multiple sites, each site identified by a different combination of first and second address tags.

[0031] In another aspect but not part of the invention is a method comprising: delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises a target-binding moiety capable of binding to the probe's corresponding

biological target; allowing each probe to bind to its corresponding biological target in the sample; delivering at least one adaptor to the multiple sites in the sample, wherein the at least one adaptor specifically binds to the probe and comprises an address tag that identifies each of the multiple sites to which the at least one adaptor is delivered, wherein the probe and/or the adaptor comprises an identity tag that identifies the probe's and/or adaptor's corresponding biological target or target-binding moiety; analyzing the at least one adaptor and the probe bound to the one or more biological targets, the analysis comprising: (1) determining abundance, expression, and/or activity of each of the one or more biological targets by assessing the amount of at least one adaptor bound to the probe bound to the biological target; and (2) determining the identities of the identity tag, and the address tag of the at least one adaptor; and determining a spatial pattern of abundance and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In one aspect, the method does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample.

[0032] In another aspect but not part of the invention is a method comprising: delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises a target-binding moiety capable of binding to the probe's corresponding biological target; allowing each probe to bind to its corresponding biological target in the sample; delivering at least one adaptor to the multiple sites in the sample, wherein the at least one adaptor specifically binds to the probe and comprises an address tag that identifies each of the multiple sites to which the at least one adaptor is delivered, wherein the probe and/or the adaptor comprises an identity tag that identifies the probe's and/or adaptor's corresponding biological target or target-binding moiety; analyzing the at least one adaptor and the probe bound to the one or more biological targets by sequencing, wherein the amount of a sequencing product indicates abundance, expression, and/or activity of each of the one or more biological targets, the sequencing product comprising all or a portion of the address tag sequence and all or a portion of the identity tag sequence; and determining a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In one aspect, the method does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample.

[0033] In yet another aspect but not part of the invention is a method comprising: delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises a target-binding moiety capable of binding to the probe's corresponding biological target; allowing each probe to bind to its corresponding biological target in the sample; delivering at least one adaptor to the multiple sites in the sample, wherein the at least one adaptor specifically binds to the probe, wherein the probe and/or the adaptor comprises an identity tag that identifies the probe's and/or adaptor's corresponding biological target or target-binding moiety; delivering an address tag to each of the multiple sites in the sample, wherein the address tag is to be coupled to the at least one adaptor bound to the probe bound to the biological target and identifies the site to which the address tag is delivered; analyzing the adaptor/address tag conjugate, the analysis comprising: (1) determining abundance,

expression, and/or activity of each of the one or more biological targets by assessing the amount of the adaptor/address tag conjugate bound to the probe bound to the biological target; and (2) determining the identities of the identity tag, and the address tag of the adaptor/address tag conjugate; and determining a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In one embodiment, the method does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample.

[0034] In still another aspect but not part of the invention is a method comprising: delivering a probe for each of one or more biological targets to multiple sites in a sample, wherein each probe comprises a target-binding moiety capable of binding to the probe's corresponding biological target; allowing each probe to bind to its corresponding biological target in the sample; delivering at least one adaptor to the multiple sites in the sample, wherein the at least one adaptor specifically binds to the probe, wherein the probe and/or the adaptor comprises an identity tag that identifies the probe's and/or adaptor's corresponding biological target or target-binding moiety; delivering an address tag to each of the multiple sites in the sample, wherein the address tag is to be coupled to the at least one adaptor bound to the probe bound to the biological target and identifies the site to which the address tag is delivered; analyzing the adaptor/address tag conjugate by sequencing, wherein the amount of a sequencing product indicates abundance, expression, and/or activity of each of the one or more biological targets, the sequencing product comprising all or a portion of the address tag sequence and all or a portion of the identity tag sequence; and determining a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample based on the analysis. In one aspect, the method does not depend on an imaging technique for determining spatial information of the one or more biological targets in the sample.

[0035] In any of the preceding aspects, at least two adaptors can be delivered to each of the multiple sites in the sample, wherein the at least two adaptors each specifically binds to one probe that specifically binds to the biological target. In one aspect, the at least two adaptors are joined, for example, by ligation using a portion of the probe sequence as a splint.

[0036] In any of the preceding aspects, the probe for each of the one or more biological targets can comprise an affinity agent for the biological target and an oligonucleotide, and the adaptor can comprise an oligonucleotide.

Brief Description of the Drawings

[0037]

Figure 1 is a flow chart illustrating exemplary steps of a method of determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites

in a sample, according to an aspect of the present disclosure.

Figure 2 is a flow chart illustrating exemplary steps of a method of determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample, according to an aspect of the present disclosure.

Figure 3 illustrates a combinatorial addressing scheme, according to one aspect of the present disclosure.

Figure 4 illustrates combinatorial addressing schemes applied to a sample, according to aspects of the present disclosure.

Figure 5 illustrates a combinatorial addressing scheme applied to a sample, according to one aspect of the present disclosure.

Figure 6 illustrates multiplexable protein detection assays with combinatorial addressing schemes applied to a sample, according to aspects of the present disclosure.

Figure 7 illustrates exemplary antibody-DNA conjugate configurations, according to certain aspects of the present disclosure.

Figure 8 illustrates sequential address tagging schemes, according to aspects of the present disclosure.

Figure 9 illustrates a microfluidic addressing device, according to one aspect of the present disclosure.

Figure 10 provides an immunofluorescence image (Figure 10A) and representative expression maps (Figure 10B-C) generated according to some aspects of the present disclosure.

Figure 11 illustrates a method for reducing random errors during the sequencing step (Figure 11A), and exemplary configurations of probes with integrated X and Y address tags and variable tag region z (Figure 11B), according to some aspects of the present disclosure.

Detailed Description

[0038] A detailed description of one or more aspects of the claimed and/or disclosed subject matter is provided below along with accompanying figures that illustrate the principles of the claimed subject matter. The claimed subject matter is described in connection with embodiments, but is not limited to any embodiment. Numerous specific details are set forth in the following description in order to provide a thorough understanding of the present disclosure. These details are provided for the purpose of example. For the purpose of clarity, technical material that is known in the technical fields related to the claimed subject matter has not been described in detail so that the claimed subject matter is not unnecessarily obscured.

[0039] Unless defined otherwise, all terms of art, notations and other technical and scientific terms or terminology used herein are intended to have the same meaning as is commonly understood by one of ordinary skill in the art to which the claimed subject matter pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. Many of the techniques and procedures described or referenced herein are well understood and commonly employed using conventional methodology by those skilled in the art.

[0040] The practice of the provided embodiments will employ, unless otherwise indicated, conventional techniques and descriptions of organic chemistry, polymer technology, molecular biology (including recombinant techniques), cell biology, biochemistry, and sequencing technology, which are within the skill of those who practice in the art. Such conventional techniques include polymer array synthesis, hybridization and ligation of polynucleotides, and detection of hybridization using a label. Specific illustrations of suitable techniques can be had by reference to the examples herein. However, other equivalent conventional procedures can, of course, also be used. Such conventional techniques and descriptions can be found in standard laboratory manuals such as Green, et al., Eds., *Genome Analysis: A Laboratory Manual Series* (Vols. I-IV) (1999); Weiner, Gabriel, Stephens, Eds., *Genetic Variation: A Laboratory Manual* (2007); Dieffenbach, Dveksler, Eds., *PCR Primer: A Laboratory Manual* (2003); Bowtell and Sambrook, *DNA Microarrays: A Molecular Cloning Manual* (2003); Mount, *Bioinformatics: Sequence and Genome Analysis* (2004); Sambrook and Russell, *Condensed Protocols from Molecular Cloning: A Laboratory Manual* (2006); and Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* (2002) (all from Cold Spring Harbor Laboratory Press); Ausubel et al. eds., *Current Protocols in Molecular Biology* (1987); T. Brown ed., *Essential Molecular Biology* (1991), IRL Press; Goeddel ed., *Gene Expression Technology* (1991), Academic Press; A. Bothwell et al. eds., *Methods for Cloning and Analysis of Eukaryotic Genes* (1990), Bartlett Publ.; M. Kriegler, *Gene Transfer and Expression* (1990), Stockton Press; R. Wu et al. eds., *Recombinant DNA Methodology* (1989), Academic Press; M. McPherson et al., *PCR: A Practical Approach* (1991), IRL Press at Oxford University Press; Stryer, *Biochemistry* (4th Ed.) (1995), W. H. Freeman, New York N.Y.; Gait, *Oligonucleotide Synthesis: A Practical Approach* (2002), IRL Press, London; Nelson and Cox, *Lehninger, Principles of Biochemistry* (2000) 3rd Ed., W. H. Freeman Pub., New York, N.Y.; Berg, et al., *Biochemistry* (2002) 5th Ed., W. H. Freeman Pub., New York, N.Y.; D. Weir & C. Blackwell, eds., *Handbook of Experimental Immunology* (1996), Wiley-Blackwell; A. Abbas et al., *Cellular and Molecular Immunology* (1991, 1994), W.B. Saunders Co.; and J. Coligan et al. eds., *Current Protocols in Immunology* (1991).

[0041] As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. For example, "a" or "an" means "at least one" or "one or more." Thus, reference to "a biological target" refers to one or more biological targets, and reference to "the method" includes reference to equivalent steps and methods disclosed herein and/or known to those skilled in the art, and so forth.

[0042] Throughout this disclosure, various aspects of the claimed subject matter are presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the claimed subject matter. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, where a range of values is provided, it is understood that each intervening value, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the claimed subject matter. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the claimed subject matter, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the claimed subject matter. This applies regardless of the breadth of the range.

[0043] As used herein, an "individual" can be any living organism, including humans and other mammals. A "subject" as used herein can be an organism to which the provided compositions, methods, kits, devices, and systems can be administered or applied. In one aspect, the subject can be a mammal or a cell, a tissue, an organ or a part of the mammal. Mammals include, but are not limited to, humans, and non-human animals, including farm animals, sport animals, rodents and pets.

[0044] As used herein, a "biological sample" can refer to any sample obtained from a living or viral source or other source of macromolecules and biomolecules, and includes any cell type or tissue of a subject from which nucleic acid or protein or other macromolecule can be obtained. The biological sample can be a sample obtained directly from a biological source or a sample that is processed. For example, isolated nucleic acids that are amplified constitute a biological sample. Biological samples include, but are not limited to, body fluids, such as blood, plasma, serum, cerebrospinal fluid, synovial fluid, urine and sweat, tissue and organ samples from animals and plants and processed samples derived therefrom.

[0045] As used herein, a "composition" can be any mixture of two or more products or compounds. It may be a solution, a suspension, liquid, powder, a paste, aqueous, nonaqueous or any combination thereof.

[0046] The terms "polynucleotide," "oligonucleotide," "nucleic acid" and "nucleic acid molecule" are used interchangeably herein to refer to a polymeric form of nucleotides of any length, and may comprise ribonucleotides, deoxyribonucleotides, analogs thereof, or mixtures thereof. This term refers only to the primary structure of the molecule. Thus, the term includes triple-, double- and single-stranded deoxyribonucleic acid ("DNA"), as well as triple-, double- and single-stranded ribonucleic acid ("RNA"). It also includes modified, for example by alkylation, and/or by capping, and unmodified forms of the polynucleotide. More particularly, the terms "polynucleotide," "oligonucleotide," "nucleic acid" and "nucleic acid molecule" include polydeoxyribonucleotides (containing 2-deoxy-D-ribose), polyribonucleotides (containing D-ribose), including tRNA, rRNA, hRNA, and mRNA, whether spliced or unspliced, any other type

of polynucleotide which is an Nor C-glycoside of a purine or pyrimidine base, and other polymers containing normucleotidic backbones, for example, polyamide (e.g., peptide nucleic acids ("PNAs")) and polymorpholino (commercially available from the Anti-Virals, Inc., Corvallis, OR., as Neugene) polymers, and other synthetic sequence-specific nucleic acid polymers providing that the polymers contain nucleobases in a configuration which allows for base pairing and base stacking, such as is found in DNA and RNA. Thus, these terms include, for example, 3'-deoxy-2',5'-DNA, oligodeoxyribonucleotide N3' to P5' phosphoramidates, 2'-O-alkyl-substituted RNA, hybrids between DNA and RNA or between PNAs and DNA or RNA, and also include known types of modifications, for example, labels, alkylation, "caps," substitution of one or more of the nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, *etc.*), with negatively charged linkages (e.g., phosphorothioates, phosphorodithioates, *etc.*), and with positively charged linkages (e.g., aminoalkylphosphoramidates, aminoalkylphosphotriesters), those containing pendant moieties, such as, for example, proteins (including enzymes (e.g. nucleases), toxins, antibodies, signal peptides, poly-L-lysine, *etc.*), those with intercalators (e.g., acridine, psoralen, *etc.*), those containing chelates (of, e.g., metals, radioactive metals, boron, oxidative metals, *etc.*), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, *etc.*), as well as unmodified forms of the polynucleotide or oligonucleotide. A nucleic acid generally will contain phosphodiester bonds, although in some cases nucleic acid analogs may be included that have alternative backbones such as phosphoramidite, phosphorodithioate, or methylphosphoroamidite linkages; or peptide nucleic acid backbones and linkages. Other analog nucleic acids include those with bicyclic structures including locked nucleic acids, positive backbones, non-ionic backbones and non-ribose backbones. Modifications of the ribose-phosphate backbone may be done to increase the stability of the molecules; for example, PNA:DNA hybrids can exhibit higher stability in some environments. The terms "polynucleotide," "oligonucleotide," "nucleic acid" and "nucleic acid molecule" can comprise any suitable length, such as at least 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100, 200, 300, 400, 500, 1,000 or more nucleotides.

[0047] It will be appreciated that, as used herein, the terms "nucleoside" and "nucleotide" will include those moieties which contain not only the known purine and pyrimidine bases, but also other heterocyclic bases which have been modified. Such modifications include methylated purines or pyrimidines, acylated purines or pyrimidines, or other heterocycles. Modified nucleosides or nucleotides can also include modifications on the sugar moiety, e.g., wherein one or more of the hydroxyl groups are replaced with halogen, aliphatic groups, or are functionalized as ethers, amines, or the like. The term "nucleotidic unit" is intended to encompass nucleosides and nucleotides.

[0048] "Nucleic acid probe" refers to a structure comprising a polynucleotide, as defined above, that contains a nucleic acid sequence that can bind to a corresponding target. The polynucleotide regions of probes may be composed of DNA, and/or RNA, and/or synthetic nucleotide analogs.

[0049] The terms "polypeptide", "oligopeptide", "peptide" and "protein" are used interchangeably herein to refer to polymers of amino acids of any length, e.g., at least 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100, 200, 300, 400, 500, 1,000 or more amino acids. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, *etc.*), as well as other modifications known in the art.

[0050] The terms "binding agent" and "target-binding moiety" as used herein may refer to any agent or any moiety thereof that specifically binds to a biological molecule of interest.

[0051] The biological targets or molecules to be detected can be any biological molecules including but not limited to proteins, nucleic acids, lipids, carbohydrates, ions, or multicomponent complexes containing any of the above. Examples of subcellular targets include organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, *etc.* Exemplary nucleic acid targets can include genomic DNA of various conformations (e.g., A-DNA, B-DNA, Z-DNA), mitochondria DNA (mtDNA), mRNA, tRNA, rRNA, hRNA, miRNA, and piRNA.

[0052] As used herein, "biological activity" may include the *in vivo* activities of a compound or physiological responses that result upon *in vivo* administration of a compound, composition or other mixture. Biological activity, thus, may encompass therapeutic effects and pharmaceutical activity of such compounds, compositions and mixtures. Biological activities may be observed *in vitro* systems designed to test or use such activities.

[0053] The term "binding" can refer to an attractive interaction between two molecules which results in a stable association in which the molecules are in close proximity to each other. Molecular binding can be classified into the following types: non-covalent, reversible covalent and irreversible covalent. Molecules that can participate in molecular binding include proteins, nucleic acids, carbohydrates, lipids, and small organic molecules such as pharmaceutical compounds. Proteins that form stable complexes with other molecules are often referred to as receptors while their binding partners are called ligands. Nucleic acids can also form stable complex with themselves or others, for example, DNA-protein complex, DNA-DNA complex, DNA-RNA complex.

[0054] As used herein, the term "specific binding" refers to the specificity of a binder, e.g., an antibody, such that it preferentially binds to a target, such as a polypeptide antigen. When referring to a binding partner (e.g., protein, nucleic acid, antibody or other affinity capture agent, *etc.*), "specific binding" can include a binding reaction of two or more binding partners with high affinity and/or complementarity to ensure selective hybridization under designated assay conditions. Typically, specific binding will be at least three times the standard deviation of

the background signal. Thus, under designated conditions the binding partner binds to its particular target molecule and does not bind in a significant amount to other molecules present in the sample. Recognition by a binder or an antibody of a particular target in the presence of other potential interfering substances is one characteristic of such binding. Preferably, binders, antibodies or antibody fragments that are specific for or bind specifically to a target bind to the target with higher affinity than binding to other non-target substances. Also preferably, binders, antibodies or antibody fragments that are specific for or bind specifically to a target avoid binding to a significant percentage of non-target substances, e.g., non-target substances present in a testing sample. In some aspects, binders, antibodies or antibody fragments of the present disclosure avoid binding greater than about 90% of non-target substances, although higher percentages are clearly contemplated and preferred. For example, binders, antibodies or antibody fragments of the present disclosure avoid binding about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, and about 99% or more of non-target substances. In other aspects, binders, antibodies or antibody fragments of the present disclosure avoid binding greater than about 10%, 20%, 30%, 40%, 50%, 60%, or 70%, or greater than about 75%, or greater than about 80%, or greater than about 85% of non-target substances.

[0055] The term "antibody" as used herein may include an entire immunoglobulin or antibody or any functional fragment of an immunoglobulin molecule which is capable of specific binding to an antigen, such as a carbohydrate, polynucleotide, lipid, polypeptide, or a small molecule, *etc.*, through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule, and can be an immunoglobulin of any class, e.g., IgG, IgM, IgA, IgD and IgE. IgY, which is the major antibody type in avian species such as chicken, is also included. An antibody may include the entire antibody as well as any antibody fragments capable of binding the antigen or antigenic fragment of interest. Examples include complete antibody molecules, antibody fragments, such as Fab, F(ab')₂, CDRs, VL, VH, and any other portion of an antibody which is capable of specifically binding to an antigen. Antibodies used herein are immunoreactive or immunospecific for, and therefore specifically and selectively bind to, for example, proteins either detected (*i.e.*, biological targets) or used for detection (*i.e.*, probes) in the assays of the invention. An antibody as used herein can be specific for any of the biological targets disclosed herein or any combinations thereof. In certain aspects, a biological target itself of the present disclosure can be an antibody or fragments thereof.

[0056] As used herein, a "fragment thereof" "region thereof" and "portion thereof" can refer to fragments, regions and portions that substantially retain at least one function of the full length polypeptide.

[0057] As used herein, the term "antigen" may refer to a target molecule that is specifically bound by an antibody through its antigen recognition site. The antigen may be monovalent or polyvalent, *i.e.*, it may have one or more epitopes recognized by one or more antibodies. Examples of kinds of antigens that can be recognized by antibodies include polypeptides, oligosaccharides, glycoproteins, polynucleotides, lipids, or small molecules, *etc.*

[0058] As used herein, the term "epitope" can refer to a peptide sequence of at least about 3 to 5, preferably about 5 to 10 or 15, and not more than about 1,000 amino acids (or any integer there between), which define a sequence that by itself or as part of a larger sequence, binds to an antibody generated in response to such sequence. There is no critical upper limit to the length of the fragment, which may, for example, comprise nearly the full-length of the antigen sequence, or even a fusion protein comprising two or more epitopes from the target antigen. An epitope for use in the subject invention is not limited to a peptide having the exact sequence of the portion of the parent protein from which it is derived, but also encompasses sequences identical to the native sequence, as well as modifications to the native sequence, such as deletions, additions and substitutions (conservative in nature).

[0059] The terms "complementary" and "substantially complementary" may include the hybridization or base pairing or the formation of a duplex between nucleotides or nucleic acids, for instance, between the two strands of a double-stranded DNA molecule or between an oligonucleotide primer and a primer binding site on a single-stranded nucleic acid. Complementary nucleotides are, generally, A and T (or A and U), or C and G. Two single-stranded RNA or DNA molecules are said to be substantially complementary when the nucleotides of one strand, optimally aligned and compared and with appropriate nucleotide insertions or deletions, pair with at least about 80% of the other strand, usually at least about 90% to about 95%, and even about 98% to about 100%. In one aspect, two complementary sequences of nucleotides are capable of hybridizing, preferably with less than 25%, more preferably with less than 15%, even more preferably with less than 5%, most preferably with no mismatches between opposed nucleotides. Preferably the two molecules will hybridize under conditions of high stringency.

[0060] "Hybridization" as used herein may refer to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide. In one aspect, the resulting double-stranded polynucleotide can be a "hybrid" or "duplex." "Hybridization conditions" typically include salt concentrations of approximately less than 1 M, often less than about 500 mM and may be less than about 200 mM. A "hybridization buffer" includes a buffered salt solution such as 5% SSPE, or other such buffers known in the art. Hybridization temperatures can be as low as 5°C, but are typically greater than 22°C, and more typically greater than about 30°C, and typically in excess of 37°C. Hybridizations are often performed under stringent conditions, *i.e.*, conditions under which a sequence will hybridize to its target sequence but will not hybridize to other, non-complementary sequences. Stringent conditions are sequence-dependent and are different in different circumstances. For example, longer fragments may require higher hybridization temperatures for specific hybridization than short fragments. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents, and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one parameter alone. Generally stringent conditions are selected to be about 5°C lower than the T_m for the specific sequence at a defined ionic strength and pH. The melting temperature T_m can be the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single

strands. Several equations for calculating the T_m of nucleic acids are well known in the art. As indicated by standard references, a simple estimate of the T_m value may be calculated by the equation, $T_m = 81.5 + 0.41 (\% G + C)$, when a nucleic acid is in aqueous solution at 1 M NaCl (see e.g., Anderson and Young, Quantitative Filter Hybridization, in Nucleic Acid Hybridization (1985)). Other references (e.g., Allawi and SantaLucia, Jr., Biochemistry, 36:10581-94 (1997)) include alternative methods of computation which take structural and environmental, as well as sequence characteristics into account for the calculation of T_m .

[0061] In general, the stability of a hybrid is a function of the ion concentration and temperature. Typically, a hybridization reaction is performed under conditions of lower stringency, followed by washes of varying, but higher, stringency. Exemplary stringent conditions include a salt concentration of at least 0.01 M to no more than 1 M sodium ion concentration (or other salt) at a pH of about 7.0 to about 8.3 and a temperature of at least 25°C. For example, conditions of 5 × SSPE (750 mM NaCl, 50 mM sodium phosphate, 5 mM EDTA at pH 7.4) and a temperature of approximately 30°C are suitable for allele-specific hybridizations, though a suitable temperature depends on the length and/or GC content of the region hybridized. In one aspect, "stringency of hybridization" in determining percentage mismatch can be as follows: 1) high stringency: 0.1 × SSPE, 0.1% SDS, 65°C; 2) medium stringency: 0.2 × SSPE, 0.1% SDS, 50°C (also referred to as moderate stringency); and 3) low stringency: 1.0 × SSPE, 0.1% SDS, 50°C. It is understood that equivalent stringencies may be achieved using alternative buffers, salts and temperatures. For example, moderately stringent hybridization can refer to conditions that permit a nucleic acid molecule such as a probe to bind a complementary nucleic acid molecule. The hybridized nucleic acid molecules generally have at least 60% identity, including for example at least any of 70%, 75%, 80%, 85%, 90%, or 95% identity. Moderately stringent conditions can be conditions equivalent to hybridization in 50% formamide, 5 × Denhardt's solution, 5x SSPE, 0.2% SDS at 42°C, followed by washing in 0.2 × SSPE, 0.2% SDS, at 42°C. High stringency conditions can be provided, for example, by hybridization in 50% formamide, 5 × Denhardt's solution, 5 × SSPE, 0.2% SDS at 42°C, followed by washing in 0.1 × SSPE, and 0.1% SDS at 65°C. Low stringency hybridization can refer to conditions equivalent to hybridization in 10% formamide, 5 × Denhardt's solution, 6 × SSPE, 0.2% SDS at 22°C, followed by washing in 1× SSPE, 0.2% SDS, at 37°C. Denhardt's solution contains 1% Ficoll, 1% polyvinylpyrrolidone, and 1% bovine serum albumin (BSA). 20 × SSPE (sodium chloride, sodium phosphate, ethylene diamide tetraacetic acid (EDTA)) contains 3M sodium chloride, 0.2M sodium phosphate, and 0.025 M EDTA. Other suitable moderate stringency and high stringency hybridization buffers and conditions are well known to those of skill in the art and are described, for example, in Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Press, Plainview, N.Y. (1989); and Ausubel et al., Short Protocols in Molecular Biology, 4th ed., John Wiley & Sons (1999).

[0062] Alternatively, substantial complementarity exists when an RNA or DNA strand will hybridize under selective hybridization conditions to its complement. Typically, selective hybridization will occur when there is at least about 65% complementary over a stretch of at least 14 to 25 nucleotides, preferably at least about 75%, more preferably at least about 90%

complementary. See M. Kanehisa, Nucleic Acids Res. 12:203 (1984).

[0063] A "primer" used herein can be an oligonucleotide, either natural or synthetic, that is capable, upon forming a duplex with a polynucleotide template, of acting as a point of initiation of nucleic acid synthesis and being extended from its 3' end along the template so that an extended duplex is formed. The sequence of nucleotides added during the extension process is determined by the sequence of the template polynucleotide. Primers usually are extended by a DNA polymerase.

[0064] "Ligation" may refer to the formation of a covalent bond or linkage between the termini of two or more nucleic acids, e.g., oligonucleotides and/or polynucleotides, in a template-driven reaction. The nature of the bond or linkage may vary widely and the ligation may be carried out enzymatically or chemically. As used herein, ligations are usually carried out enzymatically to form a phosphodiester linkage between a 5' carbon terminal nucleotide of one oligonucleotide with a 3' carbon of another nucleotide.

[0065] "Sequencing," "sequence determination" and the like means determination of information relating to the nucleotide base sequence of a nucleic acid. Such information may include the identification or determination of partial as well as full sequence information of the nucleic acid. Sequence information may be determined with varying degrees of statistical reliability or confidence. In one aspect, the term includes the determination of the identity and ordering of a plurality of contiguous nucleotides in a nucleic acid. "High throughput digital sequencing" or "next generation sequencing" means sequence determination using methods that determine many (typically thousands to billions) of nucleic acid sequences in an intrinsically parallel manner, *i.e.* where DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or alternatively using an ultra-high throughput serial process that itself may be parallelized. Such methods include but are not limited to pyrosequencing (for example, as commercialized by 454 Life Sciences, Inc., Branford, Conn.); sequencing by ligation (for example, as commercialized in the SOLiD™ technology, Life Technologies, Inc., Carlsbad, Calif.); sequencing by synthesis using modified nucleotides (such as commercialized in TruSeq™ and HiSeq™ technology by Illumina, Inc., San Diego, Calif.; HeliScope™ by Helicos Biosciences Corporation, Cambridge, Ma.; and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, Calif.), sequencing by ion detection technologies (such as Ion Torrent™ technology, Life Technologies, Carlsbad, Calif.); sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, Calif.); nanopore-based sequencing technologies (for example, as developed by Oxford Nanopore Technologies, LTD, Oxford, UK), and like highly parallelized sequencing methods.

[0066] "SNP" or "single nucleotide polymorphism" may include a genetic variation between individuals; e.g., a single nitrogenous base position in the DNA of organisms that is variable. SNPs are found across the genome; much of the genetic variation between individuals is due to variation at SNP loci, and often this genetic variation results in phenotypic variation between individuals. SNPs for use in the present invention and their respective alleles may be derived

from any number of sources, such as public databases (U.C. Santa Cruz Human Genome Browser Gateway (genome.ucsc.edu/cgi-bin/hgGateway) or the NCBI dbSNP website (www.ncbi.nlm.nih.gov/SNP/), or may be experimentally determined as described in U.S. Pat. No. 6,969,589; and US Pub. No. 2006/0188875 entitled "Human Genomic Polymorphisms." Although the use of SNPs is described in some of the embodiments presented herein, it will be understood that other biallelic or multi-allelic genetic markers may also be used. A biallelic genetic marker is one that has two polymorphic forms, or alleles. As mentioned above, for a biallelic genetic marker that is associated with a trait, the allele that is more abundant in the genetic composition of a case group as compared to a control group is termed the "associated allele," and the other allele may be referred to as the "unassociated allele." Thus, for each biallelic polymorphism that is associated with a given trait (e.g., a disease or drug response), there is a corresponding associated allele. Other biallelic polymorphisms that may be used with the methods presented herein include, but are not limited to multinucleotide changes, insertions, deletions, and translocations. It will be further appreciated that references to DNA herein may include genomic DNA, mitochondrial DNA, episomal DNA, and/or derivatives of DNA such as amplicons, RNA transcripts, cDNA, DNA analogs, *etc.* The polymorphic loci that are screened in an association study may be in a diploid or a haploid state and, ideally, would be from sites across the genome.

[0067] As used herein, the term "microfluidic device" may generally refer to a device through which materials, particularly fluid borne materials, such as liquids, can be transported, in some embodiments on a micro-scale, and in some embodiments on a nanoscale. Thus, the microfluidic devices described by the presently disclosed subject matter can comprise microscale features, nanoscale features, and combinations thereof.

[0068] Accordingly, an exemplary microfluidic device typically comprises structural or functional features dimensioned on the order of a millimeter-scale or less, which are capable of manipulating a fluid at a flow rate on the order of a $\mu\text{L}/\text{min}$ or less. Typically, such features include, but are not limited to channels, fluid reservoirs, reaction chambers, mixing chambers, and separation regions. In some examples, the channels include at least one cross-sectional dimension that is in a range of from about 0.1 μm to about 500 μm . The use of dimensions on this order allows the incorporation of a greater number of channels in a smaller area, and utilizes smaller volumes of fluids.

[0069] A microfluidic device can exist alone or can be a part of a microfluidic system which, for example and without limitation, can include: pumps for introducing fluids, e.g., samples, reagents, buffers and the like, into the system and/or through the system; detection equipment or systems; data storage systems; and control systems for controlling fluid transport and/or direction within the device, monitoring and controlling environmental conditions to which fluids in the device are subjected, e.g., temperature, current, and the like.

[0070] As used herein, the terms "channel," "micro-channel," "fluidic channel," and "microfluidic channel" are used interchangeably and can mean a recess or cavity formed in a material by imparting a pattern from a patterned substrate into a material or by any suitable material

removing technique, or can mean a recess or cavity in combination with any suitable fluid-conducting structure mounted in the recess or cavity, such as a tube, capillary, or the like. In the present invention, channel size means the cross-sectional area of the microfluidic channel.

[0071] As used herein, the terms "flow channel" and "control channel" are used interchangeably and can mean a channel in a microfluidic device in which a material, such as a fluid, *e.g.*, a gas or a liquid, can flow through. More particularly, the term "flow channel" refers to a channel in which a material of interest, *e.g.*, a solvent or a chemical reagent, can flow through. Further, the term "control channel" refers to a flow channel in which a material, such as a fluid, *e.g.*, a gas or a liquid, can flow through in such a way to actuate a valve or pump.

[0072] As used herein, "chip" may refer to a solid substrate with a plurality of one-, two- or three-dimensional micro structures or micro-scale structures on which certain processes, such as physical, chemical, biological, biophysical or biochemical processes, *etc.*, can be carried out. The micro structures or micro-scale structures such as, channels and wells, electrode elements, electromagnetic elements, are incorporated into, fabricated on or otherwise attached to the substrate for facilitating physical, biophysical, biological, biochemical, chemical reactions or processes on the chip. The chip may be thin in one dimension and may have various shapes in other dimensions, for example, a rectangle, a circle, an ellipse, or other irregular shapes. The size of the major surface of chips of the present invention can vary considerably, *e.g.*, from about 1 mm² to about 0.25 m². Preferably, the size of the chips is from about 4 mm² to about 25 cm² with a characteristic dimension from about 1 mm to about 5 cm. The chip surfaces may be flat, or not flat. The chips with non-flat surfaces may include channels or wells fabricated on the surfaces.

[0073] A microfluidic chip can be used for the methods and assay systems disclosed herein. A microfluidic chip can be made from any suitable materials, such as PDMS (Polydimethylsiloxane), glass, PMMA (polymethylmethacrylate), PET (polyethylene terephthalate), PC (Polycarbonate), *etc.*, or a combination thereof.

[0074] "Multiplexing" or "multiplex assay" herein may refer to an assay or other analytical method in which the presence and/or amount of multiple targets, *e.g.*, multiple nucleic acid target sequences, can be assayed simultaneously by using more than one capture probe conjugate, each of which has at least one different detection characteristic, *e.g.*, fluorescence characteristic (for example excitation wavelength, emission wavelength, emission intensity, FWHM (full width at half maximum peak height), or fluorescence lifetime) or a unique nucleic acid or protein sequence characteristic.

Assays for Determining Spatial Patterns of Biological Targets

[0075] Disclosed herein are spatially-encoded, multiplexed methods and assay systems capable of high levels of multiplexing with an efficient spatial encoding scheme. In one aspect,

provided herein is instrumentation capable of delivering reagents to a sample and thereby spatially encoding multiple sites to which the reagents are delivered. In one aspect, reagents can be delivered to a sample according to a known spatial pattern, for example, a spatial pattern determined by histological features of the sample. In another aspect, reagents are delivered by random-access methods, such as inkjet and pin-spotting. In another aspect, microfluidic devices with addressing channels and the like are used to deliver reagents to a sample, and to spatially encode multiple sites in the sample to which the reagents are delivered. In some aspects, the spatially-encoded ("addressed," or "address tagged"), multiplexed methods and assay systems comprise a decoding feature determined by a readout that is digital in nature. In one aspect, the methods and assay systems disclosed herein detect the presence or absence of a biological target or a biological activity indicative of a biological target. In another aspect, provided herein are methods and assay systems that can detect the amount or abundance of a biological target or biological activity indicative of a biological target at multiple sites in a sample, as well as the location of each of the multiple sites in the sample. Based on the analysis of the amount or abundance and the location information of one or more biological targets or activities, spatial patterns across the multiple sites in the sample can be generated. In any of the preceding aspects, the method or assay system may not depend on an imaging technique for determining spatial or location information of the one or more biological targets in the sample, although the method or assay system may optionally comprise using an imaging technique for other purposes. Imaging techniques may include but are not limited to conventional immunohistochemical (IHC) imaging and immunofluorescence (IF) imaging. Methods and assays systems to determine a spatial pattern of abundance and/or activity of a biological target in a sample are disclosed in detail in U.S. Application Serial No. 13/080,616, entitled "Spatially encoded biological assays" (Pub. No.: US 2011/0245111).

[0076] The present disclosure further provides instrumentation with an ability to deliver reagents to multiple sites in a sample, wherein each of the multiple sites can be identified by the reagents delivered thereto. In one aspect, reagents are delivered in a spatially-defined pattern. The instrumentation, together with software, reagents and protocols, provides a key component of the methods and assay systems of the present disclosure, allowing for measurement of numerous biological targets or activities, including DNA, RNA and/or protein expression, and spatial localization of such biological targets or activities in a sample. In one aspect, the abundance, expression, and/or activity and the location of biological targets in the biological samples are determined after the assay products of the multiplexed assay are removed from the biological sample and pooled for analysis. Determination of the abundance, expression, and/or activity and the location of biological targets can be performed by, e.g., next-generation sequencing, which easily provides millions to trillions of data points at low cost. The assay results such as the amount or activity of biological targets can then be mapped back to a specific location in the biological sample. The methods and assay systems provide tools to analyze the complex spatial patterns of cellular function and regulation in biological samples.

[0077] In one aspect but not part of the invention is a method of determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample is provided in Figure 1. At Step 110, a probe for each of one or more biological

targets is delivered to multiple sites in a sample, each probe comprising a target-binding moiety, an address tag that identifies each site to which the probe is delivered, and an identity tag.

[0078] In any of the aspects of the present disclosure, the sample can be any biological sample or samples that can be affixed to a support or provided essentially in a two-dimensional manner, such that an assayed biological target or activity can be tied back to the location within the biological sample. In certain embodiments, the sample can be a freshly isolated sample, a fixed sample, a frozen sample, an embedded sample, a processed sample, or a combination thereof. Exemplary samples of the present disclosure include tissue sections (e.g., including whole animal sectioning and tissue biopsies), cell populations, or other biological structure disposed upon a support, such as on a slide (e.g., a microscope slide) or culture dish, and the like. In preferred aspects, the methods and assay systems of the present disclosure are compatible with numerous biological sample types, including fresh samples, such as primary tissue sections, and preserved samples including but not limited to frozen samples and formalin-fixed, paraffin-embedded (FFPE) samples. In certain aspects, the sample can be fixed with a suitable concentration of formaldehyde or paraformaldehyde, for example, 4% of formaldehyde or paraformaldehyde in phosphate buffered saline (PBS). In certain aspects, the biological samples are immobilized on a substrate surface having discrete, independently measureable areas.

[0079] In one aspect, the biological sample may contain one or more biological targets of interest. In any of the aspects of the present disclosure, the one or more biological targets can be any biological molecules including but not limited to proteins, nucleic acids, lipids, carbohydrates, ions, or multicomponent complexes containing any of the above. Examples of subcellular targets include organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, *etc.* In some aspects, the one or more biological targets can be nucleic acids, including RNA transcripts, genomic DNA sequences, cDNAs, amplicons, or other nucleic acid sequences. In other aspects, the one or more biological targets can be proteins, enzymes (protein enzymes or ribozymes) and the like.

[0080] At Step 110, the probe for each of the multiple biological targets comprise: (1) a target-binding moiety capable of binding to the probe's corresponding biological target; (2) an address tag that identifies each site to which the probe is delivered; and (3) an identity tag that identifies the probe's corresponding biological target or target-binding moiety. Depending on the nature of the biological target, the target-binding moiety can be a target-specific nucleotide sequence (for example, a sequence complementary to a sequence of a nucleic acid target), small molecule, aptamer, antibody, lipid, carbohydrate, ion, affinity capture agent, or multicomponent complexes containing any of the above. The address tag identifies the position in the sample to which the probe is delivered, and the identity tag identifies the probe's corresponding biological target being assayed or the target-binding moiety. Thus, the identities of the address tag and the identity tag can be used to link assay results to biological targets and locations in the sample. In preferred embodiments, there can be at least two address tags

for a biological target at each of multiple sites in a sample, each address tag identifying a parameter of each of the multiple sites. For example, there can be an X-axis address tag and a Y-axis address tag for each site in a sample placed on an X-Y coordinate plane. Thus, each site can be uniquely identified by its corresponding (X, Y) coordinates. In preferred embodiments of the present disclosure, the address tags and/or the identity tags can be oligonucleotides. In other embodiments, the address tags and/or the identity tags can be mass tags, fluorescent labels, or other moieties.

[0081] In some aspects, the target-binding moiety, address tag, and/or identity tag of the probe are pre-coupled before being delivered to the biological sample. In the case where the probes are oligonucleotides, the target-binding sequence, address tag sequence, and/or identity tag sequence can be synthesized as a single oligonucleotide. Alternatively, the target-binding moiety, address tag, and/or identity tag of the probe can be synthesized or obtained separately and combined before delivery to the biological sample. For example, two separate oligonucleotides can be synthesized and coupled by, e.g., ligation; or an antibody and an oligonucleotide can be prepared separately and conjugated before delivery to the biological sample. In other aspects, the probes and the address tags can be synthesized separately, and delivered to the biological sample at different steps (e.g., probes first and address tags thereafter, or vice versa) in the assay.

[0082] At Step 120, the probe is allowed to bind to its corresponding biological target in the sample and thereby to react or interact with the biological target. For example, conditions are provided to allow oligonucleotides to hybridize to nucleic acid targets, enzymes to catalyze reactions with protein targets, antibodies to bind epitopes within a target, *etc.* In the case where the biological targets are nucleic acids, the probes are typically oligonucleotides and hybridize to the target nucleic acids. In the case that the biological targets are proteins, the probes typically are aptamers, small molecules, or oligonucleotide-conjugated proteins that interact with target proteins by binding to them or by reacting with them (that is, one of the proteins is a substrate for the other). Oligonucleotides may be coupled to the probes or proteins by conjugation, chemical or photocrosslinking via suitable groups and the like.

[0083] In some aspects, after allowing the probes to bind to or interact with the one or more biological targets in the sample, probes bound to the biological targets may be separated from probes delivered to the sample but not bound to the biological targets. In one aspect, in the case where the biological targets are nucleic acids and the probes are oligonucleotides, the separation can be accomplished by, e.g., washing the unhybridized probes from the sample. Similarly, for other assays that are based on affinity binding, including those using aptamer, small molecule, and protein probes, washing steps can be used to remove low affinity binders. In the case where the probe is transformed via interaction with the target, e.g., in the case of a peptide, e.g., via cleavage by a protease or phosphorylation by a kinase, it is convenient to collect all probes, including both probes that have interacted with the biological targets and thus transformed and probes not transformed. After collection or pooling, an antibody or other affinity capture agent can be used to capture probes transformed by addition of a moiety (e.g., a phosphate group in cases of phosphorylation by a kinase). In cases where probes have been

transformed via cleavage, the transformed probes can be separated, *e.g.*, by capturing the non-transformed probes via a tag that is removed from the transformed probes during the transformation (*e.g.*, by cleavage), or by adding a new tag at the site of cleavage.

[0084] In certain other aspects, probes bound to the biological targets may not need to be separated from probes not bound to the biological targets for determining a spatial pattern of abundance, expression, and/or activity of the biological targets. At Step **130**, probes bound to the one or more biological targets are analyzed. In certain aspects, the analysis comprises determining abundance, expression, and/or activity of each biological target and the identities of the identity tag and the address tag for each biological target at each site. Numerous methods can be used to identify the address tags, identity tags and/or target-binding moieties of the probes of the methods and assay systems disclosed herein. The address tags can be detected using techniques such as mass spectroscopy (*e.g.*, matrix-assisted laser desorption/ionization-time of flight mass spectrometry (MALDI-TOF), LC-MS/MS, and TOF/TOF[™] LC/MS/MS), nuclear magnetic resonance imaging, or, preferably, nucleic acid sequencing. Examples of techniques for decoding the probes of the present invention can be found, for example, in US Pub. No. 20080220434. For example, the address tags may be oligonucleotide mass tags (OMTs or massTags). Such tags are described, *e.g.*, in US Pub. No. 20090305237. In yet another aspect, the probes can be amplified and hybridized to a microarray. This would require separate amplification reactions to be carried out, in which each amplification is specific to a particular address tag or subset of tags, accomplished by using tag-specific primers. Each amplification would also incorporate a different resolvable label (*e.g.* fluorophor). Following hybridization, the relative amounts of a particular target mapping to different spatial locations in the sample can be determined by the relative abundances of the resolvable labels. At Step **140**, based on the analysis of probes bound to the one or more biological targets, a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample is determined.

[0085] In a preferred aspect, the probes according to the present disclosure are substrates for high-throughput, next-generation sequencing, and highly parallel next-generation sequencing methods are used to confirm the sequence of the probes (including, for example, the sequence of the target-binding moiety, the address tag, and/or the identity tag). Suitable sequencing technologies include but are not limited to SOLiD[™] technology (Life Technologies, Inc.) or Genome Analyzer (Illumina, Inc.). Such next-generation sequencing methods can be carried out, for example, using a one pass sequencing method or using paired-end sequencing. Next generation sequencing methods include, but are not limited to, hybridization-based methods, such as disclosed in *e.g.*, Drmanac, U.S. Pat. Nos. 6,864,052; 6,309,824; and 6,401,267; and Drmanac et al., U.S. patent publication 2005/0191656; sequencing-by-synthesis methods, *e.g.*, U.S. Pat. Nos. 6,210,891; 6,828,100; 6,969,488; 6,897,023; 6,833,246; 6,911,345; 6,787,308; 7,297,518; 7,462,449 and 7,501,245; US Publication Application Nos. 20110059436; 20040106110; 20030064398; and 20030022207; Ronaghi, et al., Science, 281:363-365 (1998); and Li, et al., Proc. Natl. Acad. Sci., 100:414-419 (2003); ligation-based methods, *e.g.*, U.S. Pat. Nos. 5,912,148 and 6,130,073; and U.S. Pat. Appln Nos. 20100105052, 20070207482 and 20090018024; nanopore sequencing, *e.g.*, U.S. Pat.

Appln Nos. 20070036511; 20080032301; 20080128627; 20090082212; and Soni and Meller, Clin Chem 53:1996-2001 (2007), as well as other methods, e.g., U.S. Pat. Appln Nos. 20110033854; 20090264299; 20090155781; and 20090005252; also, see, McKernan, et al., Genome Res. 19:1527-41 (2009) and Bentley, et al., Nature 456:53-59 (2008).

[0086] In preferred aspects, probes bound to the one or more biological targets are analyzed by sequencing. Such analysis by sequencing comprises determining the amount of a sequencing product, which indicates abundance, expression, and/or activity of each biological target, the sequencing product comprising all or a portion of the address tag sequence and all or a portion of the identity tag sequence identifying each biological target at each site. In one aspect, the address tag sequence of the sequencing product allows the mapping of the assay results back to the multiple sites in the sample.

[0087] In certain aspects, two probes that bind to the same target molecule (for example, two polynucleotide probes that hybridize to adjacent sites on a nucleic acid target) may be assayed by extension followed by ligation (the extension-ligation assay). The extension-ligation assay allows certain target sequence to be determined *de novo*. For example, if the primer and the downstream oligo are separated by 20 bases and reverse transcriptase is used to fill this 20-base gap, 20 bases of sequence of the RNA target can be obtained. In certain aspects, by using the extension-ligation assay in the present methods or assay systems, regions of sequence that are of particular interest may be characterized. For example, these regions may comprise mutations or variations, for example, with implications in cancer, MHC variations, and RNA editing.

[0088] In any of the aspects disclosed herein, an extension assay may also be used and may allow certain target sequences, for example, nucleotide sequences, to be determined *de novo*. In one aspect, an extension assay of the present disclosure may be performed as follows. A first primer may be used to make cDNA from a target sequence. In certain aspects, the first primer can be a random primer (e.g., random hexamer) or a sequence-specific primer. A random primer can be used to make cDNA from the entire transcriptome, while a sequence-specific primer may be used to make cDNA from a specific target sequence. In certain aspects, the first primer may comprise a universal priming site for amplification of the assay products, an adaptor to enable sequence identification by sequencing techniques, and/or an adaptor for attaching address tags. In other aspects, the first primer may be conjugated to an adaptor for attaching address tags. The X and Y address tags as described *infra* can be coupled to the first primer via the adaptor. Note that the X and Y address tags can be coupled to the same or different side relative to the target sequence, and the configuration may be used in any of aspects disclosed herein. The Y address tag can be further linked to a universal priming site or an adaptor for sequencing coupled to biotin. The cDNA with conjugated X and Y address tags are then eluted and captured on a streptavidin bead, and a second primer can then be installed to the cDNA on the opposite side relative to the first primer. In some aspects, capture of the polynucleotide sequence can be based on other hapten-binder combinations other than biotin-streptavidin, or be sequence-based. In certain aspects, the second primer can be a random primer (e.g., random hexamer) or a sequence-specific primer. In certain aspects, the

second primer may comprise a universal priming site for amplification of the assay products, an adaptor to enable sequence identification by sequencing techniques, and/or an adaptor for attaching address tags. In other aspects, the second primer may be conjugated to a universal priming site or an adaptor for sequencing. Together with the priming site or adaptor coupled to biotin, the sequence can be extended from the second primer, amplified, and sequenced.

[0089] The methods and assay systems disclosed herein may comprise an amplification step, and in particular, a nucleic acid amplification step. In certain aspects, the amplification step is performed by PCR. In some aspects, linear amplification (e.g., by T7 RNA polymerase) may be used instead of PCR or as a partial replacement for PCR. In one aspect, linear amplification of polynucleotides may cause less distortion of the relative abundances of different sequences. This can be accomplished by including a T7 RNA pol promoter in one of the universal portions of the sequence. In this case, the promoter itself and regions upstream of the promoter are not copied. In yet other aspects, other amplification methods may be used in the methods and assay systems disclosed herein. For some sequencing methods (e.g., nanopore sequencing), amplification may be optional.

[0090] The T7 RNA polymerase based amplification is a commonly used protocol for mRNA amplification originally described by van Gelder et al., Proc. Natl. Acad. Sci. USA 87, 1663-1667 (1990). The protocol consists of the synthesis of a cDNA complementary to the mRNA ("first strand synthesis"), effected by reverse transcription, followed by second strand synthesis to yield double-stranded cDNA, and *in vitro* transcription using the double-stranded cDNA as template effected with T7 RNA polymerase. The last step provides single-stranded antisense RNA (aRNA), which may be labeled in case labeled nucleotides are provided. The nucleotides may be labeled by radioactive labeling or nonradioactive labeling methods. Eberwine et al. (Proc. Natl. Acad. Sci. USA 89, 3010-3014 (1992)) extended van Gelder's method by adding a second round of amplification using the RNA obtained in the first round as template. Wang et al. (Nature Biotechnol. 18, 457-459 (2000)) provided a variant of the original T7 method, characterized by a modified second strand synthesis method. The second strand synthesis method of Wang *et al.* is known in the art as the SMART™ technology (Clontech) for cDNA synthesis. Baugh et al. (Nucleic Acids Res. 29, E29 (2001)) describe an optimized variant of the method according to van Gelder *et al.* and analyze the performance on Affymetrix DNA chips (GeneChip®). Affymetrix GeneChips are designed to probe the anti-sense strand. Any other DNA chip or microarray probing the anti-sense strand may be envisaged when performing a T7 RNA amplification, wherein labeling occurs during the *in vitro* transcription step.

[0091] In other aspects, amplification techniques such as rolling circle amplification (RCA) and circle to circle amplification (C2CA) may be used for probe, target, tag, and/or signal amplification of the present disclosure. RCA is a linear-isothermal process in the presence of certain DNA polymerases, using an ssDNA mini-circle as a template (Fire and Xu, Proc. Natl. Acad. Sci., 92: 4641-4645 (1995); Daubendiek et al., J. Am. Chem. Soc. 117:77818-7819 (1995)). In certain aspects, a polynucleotide sequence can be replicated about 500 to 1000 times, depending on the amplification time. For example, in a dual targeting assay for a target

protein as discussed *supra*, a linear ligated product is formed (e.g., when two antibodies bind to adjacent domains on a target protein, the antibodies' oligonucleotide tags can be ligated), and can be cut by restriction enzymes and then religated to form a DNA circle by a DNA ligase and a template. In certain aspects, phi29 DNA polymerase can be used to extend the primer, which is also the template, to form a long ssDNA containing a number of sequences complementary to the initial DNA circle. C2CA is based on RCA, and may include three steps: replication, monomerization and ligation (Dahl et al., Proc. Natl. Acad. Sci., 101: 4548-4553 (2004)). The original circular DNA is considered as the positive polarity. After one step of replication (RCA reaction), the product is converted into the opposite polarity. Restriction oligos with the positive polarity (RO^+) can form duplex regions with the RCA product, and the duplex regions can be cleaved by restriction enzymes to generate monomers. Then the monomers can be guided into a ligation step and circularized. These circles serve as the templates for the next round of RCA, primed by the RO^+ . The process can be further repeated to produce around 100-fold higher concentration of target sequences than conventional PCR.

[0092] In another aspect, as shown in Figure 2, a method of determining a spatial pattern of abundance, expression, and/or activity of one or more biological targets across multiple sites in a sample is provided, featuring an efficient implementation of an address tagging scheme for the one or more biological targets across the multiple sites. In one aspect, the address tagging scheme is a combinatorial scheme using at least two address tags for the biological targets for each of the multiple sites in the sample. At Step **210**, a probe for each of one or more biological targets to multiple sites in a sample is delivered, each probe comprising (1) a target-binding moiety capable of binding to the probe's corresponding biological target; and (2) an identity tag that identifies the probe's corresponding biological target or target-binding moiety. Depending on the nature of the biological target, the target-binding moiety can be a target-specific nucleotide sequence (for example, a sequence complementary to a sequence of a nucleic acid target), small molecule, aptamer, antibody, lipid, carbohydrate, ion, affinity capture agent, or multicomponent complexes containing any of the above. At Step **220**, each probe is allowed to interact with or bind to its corresponding biological target in the sample, under appropriate conditions.

[0093] In certain aspects, probes not bound to the biological targets may be removed, and thereby separated from probes bound to the biological targets. Such separation can be performed essentially as discussed above, for example, by washing the sample to remove unhybridized oligonucleotide probes. In certain other aspects, probes bound to the biological targets may not need to be separated from probes not bound to the biological targets for determining a spatial pattern of abundance, expression, and/or activity of the biological targets.

[0094] Next, at Step **230**, an address tag is delivered to each of the multiple sites in the sample, and the address tag is to be coupled to the probe for each biological target and identifies each site to which the address tag is delivered. Note that in this aspect, the probe and address tag are delivered in separate steps. In certain aspects where the probes are oligonucleotides, the address tags may be coupled to the oligonucleotide probes by various

means known to the skilled artisan, for example, by extension, ligation, ligation followed by extension, or any combination thereof. For instance, the information in the address tags can be transferred by using a DNA polymerase to extend a probe oligonucleotide that acts as a primer, and thereby copy and incorporate the sequence of the address tags.

[0095] At Step **240**, probe/address tag conjugates bound to the one or more biological targets are analyzed. In certain aspects, the analysis comprises determining abundance, expression, and/or activity of each biological target and the identities of the identity tag and the address tag for each biological target at each site. In one aspect, the abundance, expression, and/or activity of each biological target can be assessed by determining the amount of the probe or the probe/address tag conjugate bound to the target. Numerous methods can be used to identify the address tags, identity tags and/or target-binding moieties of the probes, as discussed above. In preferred aspects, probes or probe/address tag conjugates bound to the one or more biological targets are analyzed by sequencing. Any suitable sequence techniques and methods as discussed above can be used, including high-throughput, next-generation sequencing, and highly parallel next-generation sequencing methods. Preferably, in any of aspects of the present disclosure, all or a portion of the address tag sequence and all or a portion of the identity tag sequence are determined from the same sequencing product. Preferably, also determined at the same time is the abundance of the sequencing product, for example, the "copy number" or "hits" of the sequencing product. The abundance of the sequencing product may correlate with the amount of the probe or probe/address tag conjugate bound to the target, which in turn can correlate with the abundance, expression, and/or activity of each biological target. In some aspects, the abundance of sequence products reveals the relative quantity of biological targets at the location.

[0096] Based on the analysis of probe/address tag conjugates bound to the one or more biological targets at Step **240**, a spatial pattern of abundance, expression, and/or activity of the one or more biological targets across the multiple sites in the sample is determined at Step **250**, for example, by mapping the assayed abundance, expression, and/or activity of each biological target back to each site of the sample.

[0097] Although individual steps are discussed in a particular order in certain aspects to better explain the claimed subject matter, the precise order of the steps can be varied. For example, Steps **210** and **230** can be combined, so that a mixture of the probes and address tags is delivered. Coupling of the address tag to the probe may be carried out immediately after the combined steps **210** and **230**, or concomitantly with them. It can therefore be appreciated that the address tagging of probe molecules and the separation of probes based on their ability to interact with their corresponding targets can be accomplished with flexibility. Similarly, there is considerable flexibility in the address tagging scheme. As described *infra*, the methods and assay systems disclosed herein are particularly amenable to combinatorial methods.

Spatially Encoded Genomic Assay

[0098] In particular aspects, the methods and assay systems can be used for nucleic acid analysis, for example, for genomic analysis, genotyping, detecting single nucleotide polymorphisms (SNPs), quantitation of DNA copy number or RNA transcripts, localization of particular transcripts within samples, and the like. Figure 3 illustrates an exemplary assay and address tagging scheme. For illustrative purposes, the target is a nucleic acid sequence, and two oligonucleotide probes are provided, and it should be understood that the disclosed methods and assay systems can be used for any suitable target employing one or more suitable probes. Each oligonucleotide probe comprises a target-specific region seen at **305** and **307**, respectively. In certain aspects, for example for detecting SNPs, the two target-specific regions are located on either side of the SNP to be analyzed. Each oligonucleotide probe also comprises a ligation region, seen at **301** and **303**, respectively. The oligonucleotide probes are allowed to hybridize to a target nucleic acid (not shown) in the biological sample. At Step **302**, one or both of the oligonucleotide probes may be extended and ligated to the other probe to form an extended probe comprising target nucleic acid region **309** and ligation regions **301** and **303**. In some aspects, the two probes are immediately adjacent to each other, and only ligation is needed to form an extended probe. In some aspects, Step **302** may be used to incorporate an SNP sequence or other target sequences to be assayed.

[0099] Two address tags, both comprising an address tag region (seen at **315** and **317**), a ligation region (seen at **311** and **313**), and a primer region (seen at **319** and **321**) are combined with and ligated to the extended probe at step **304** to form a target-specific oligonucleotide. In contrast with Figure 1, the probes and address tags are delivered at separate steps. In some aspects, a pair of address tags ligate specifically to one side of the target sequence or the other (*i.e.*, 5' or 3' of the target sequence), respectively. In certain aspects, the ligation and primer regions of the address tags and probes are universal; that is, the set of ligation and primer regions used in constructing the probes and address tags are constant, and only the target-specific regions of the probes and the address tag region of the address tag differ. In alternative aspects, the ligation and primer regions are not universal and each probe and/or address tag may comprise a different ligation and/or primer region.

[0100] Following ligation, the probe/address tag conjugates are eluted, pooled, and, optionally, sequencing adaptors are added to the probe/address tag conjugates via PCR. In alternative aspects, sequencing primers may be ligated to the address tags, or sequencing primer sequences can be included as part of the address tags. As seen in Figure 3, each sequencing adaptor comprises primer region **319** or **321**, compatible with the primer regions **319** and **321** on the address tags. The final construct comprising first adaptor **327**, first primer region **319**, first coding tag **315**, ligation regions **311** and **301**, target region **309**, ligation regions **313** and **303**, second coding tag **317**, second primer region **325** and second adaptor **329** can be subject to sequencing, for example, by input into a digital high-throughput sequencing process.

[0101] A combination of extension and ligation reactions are exemplified in Figure 3, but it should be appreciated that a variety of reactions may be used to couple the address tags to the target-specific probes, including ligation only (*e.g.*, for oligonucleotides that hybridize to contiguous portions of the target nucleic acid sequence). Alternatively, an assay utilizing an

additional oligonucleotide, such as in the GOLDENGATE[®] assay (Illumina, Inc., San Diego, Calif.) (see Fan, et al., Cold Spring Symp. Quant. Biol., 68:69-78 (2003)), may be employed.

[0102] To maximize the efficiency of address tagging, a combinatorial approach using pairs of address tags can be used. By de-coupling the target-specific information and the spatial information in the address tags, the number of oligonucleotides required for determining a spatial pattern of one or more biological targets across multiple sites in a sample is dramatically reduced, with a concomitant decrease in cost.

[0103] Figure 4 illustrates one aspect of a combinatorial address tagging scheme, where nucleic acids in a representative tissue section (shown at **416**) are assayed. Figure 4A shows two probe/address tag conjugates **420** and **422** specifically bound to a biological target **402** of interest. The first probe/address tag conjugate **420** comprises address tag **408**, associated with tag **404**. Tag **404** can be a universal priming site for amplification of the assay products or an adaptor to enable identification of the address tag **408** and/or other regions of probe/address tag conjugates **420**, for example, using sequencing technologies. The second probe/address tag conjugate **422** comprises address tag **406**, associated with tag **410**. Tag **410** can be a universal priming site for amplification of the assay products or an adaptor to enable identification of the address tag **406** and/or other regions of probe/address tag conjugates **422**, for example, using sequencing technologies.

[0104] In other aspects, a biological target **424** is assayed according to the combinatorial address tagging scheme shown in Figure 4D. Two probes **426** and **428** specifically bind to the biological target **424** of interest. In some aspects, a portion of each of probes **426** and **428** specifically binds to the target, while each probe also has a portion that specifically binds to an adaptor **438**, for example, by specific nucleic acid hybridization. In one aspect, the probe or probes specifically hybridize to the adaptor. In cases where the biological target is a nucleic acid and the probes are oligonucleotides, the adaptor can specifically bind to the following combinations: 1) the 5' portion of probe **426** and the 3' portion of probe **428**; 2) the 3' portion of probe **426** and the 5' portion of probe **428**; 3) the 5' portion of probe **426** and the 5' portion of probe **428**; or 4) the 3' portion of probe **426** and the 3' portion of probe **428**. In certain aspects, probe **426** or **428** is a linear molecule, a branched molecule, a circular molecule, or a combination thereof. After binding of the two probes to the biological target and the adaptor to the two probes, address tags can be delivered to the sample and coupled to the adaptor. For example, the adaptor can be tagged with address tag **430**, associated with tag **434**, and/or with address tag **432**, associated with tag **436**. Tags **434** and **436** can be universal priming sites for amplification of the assay products or sequences to enable identification of the address tags and/or other regions of adaptor/address tag conjugates, for example, using sequencing technologies. In certain aspects, the address tags are tagged at the same end of the adaptor, or at different ends of the adaptor. In other aspects, an address tag and/or tag **434** or **436** can be pre-coupled to the adaptor, and the adaptor/address tag or adaptor/tag conjugate or complex is then delivered to the sample in order to bind to the probe bound to the biological target. In certain aspects, the adaptor is a linear molecule, a branched molecule, a circular molecule, or a combination thereof. In some aspects, after an address tag is attached to each

end of the adaptor, the ends can be joined. For example, in Figure 4D, address tags **434** and/or **436** can comprise structures and/or sequences that allow the two ends of the tagged adaptor **438** to be joined to form a circular construct, to facilitate amplification and/or sequencing of the construct.

[0105] In certain aspects, all or a portion of the adaptor/address tag conjugate sequence is determined, for example, by nucleic acid sequencing. In other aspects, all or a portion of the probe sequence, and/or all or a portion of the adaptor/address tag conjugate sequence, is determined. For example, a first address tag can be coupled to probe **426**, and a second address tag can be coupled to adaptor **438**. The duplex formed between probe **426** and adaptor **438** can be subjected to extension and sequencing, to generate a conjugate that comprises sequences of the first address tag, all or a portion of probe **426**, all or a portion of adaptor **438**, and the second address tag.

[0106] The tagging scheme is not limited to the use of two or more probes for the same biological target. For example, in cases where one probe is used, a tag (e.g., an address tag, an adaptor for ligation, or a universal sequencing primer or amplification primer sequence) can be coupled to an adaptor that specifically binds to the probe, rather than to the probe itself.

[0107] In some aspects, at least two adaptors are used. In one aspect, more than one probes are delivered to the sample, and at least one adaptor is provided for each probe that specifically binds to the probe. In one aspect, one or more adaptors are provided for specifically binding to each probe. For example, a pair of adaptors is used to specifically bind to the probe **426** and **428**, respectively. In certain aspects, the adaptors of the pair are DNA molecules that: 1) hybridize or otherwise bind to probe **426** or **428**; 2) have free 3' and/or 5' ends that enable the encoding sequences (e.g., address tags **430** and **432**) to be attached in a subsequent step or steps, for example, by ligation; 3) are in a form where they can be joined if they are co-localized or in proximity to each other. In some aspects, part of probe **426** or **428** acts as a splint to enable ligation, or extension and ligation, of the adaptors in the adaptor pair. Additional tags (e.g., an address tag, an adaptor for ligation, or a universal sequencing primer or amplification primer sequence) can be coupled to the adaptor generated by joining the adaptor pair.

[0108] Figure 4B shows an address tagging scheme that may be used for 100 unique sites in a sample. For example, twenty probe/address tag conjugates a1 through a10 and b1 through b10 can be used, with each of a1 through a10 corresponding to a probe/address tag conjugate **420** (comprising an address tag **408**) and each of b1 through b10 corresponding to a probe/address tag conjugate **422** (comprising an address tag **406**). The address tag comprised in each of a1 through a10 and b1 through b10 may be uniquely identified. Probe/address tag conjugate a1, for example, is delivered to the biological sample through an addressing channel shown as the first horizontal channel in **412**. Probe/address tag conjugate a2 is delivered to the biological sample through the second horizontal channel in **412**. Probe/address tag conjugate a3 is delivered to the biological sample through the third horizontal channel in **412**, and so on. Whereas the "a" probe/address tag conjugates are

delivered in ten horizontal channels, the "b" probe/address tag conjugates are delivered in ten vertical channels as shown in **414**. For example, probe/address tag conjugate b 1 is delivered to the biological sample through the first horizontal channel of **414**, probe/address tag conjugate b2 is delivered to the biological sample through the second horizontal channel of **414**, and so on. In other embodiments, the "a" tags may be referred to as the "X" tags and the "b" tags as "Y" tags. The intersections or junctions between the horizontal and vertical channels are shown as solid squares. Each intersection or junction can be uniquely identified by the combination of the "a" probe/address tag conjugate and the "b" probe/address tag conjugate delivered to the area in the sample corresponding to the intersection or junction.

[0109] Figure 4C shows a representative tissue section **416** coincident with grid **418**. The arrows show how the "a" probe/address tag conjugates and the "b" probe/address tag conjugates are delivered on grid **418** that is coincident with tissue section **416**. If, once analyzed, probe/address tag conjugates a1 and b4, e.g., are associated with a target, then that target is present in the tissue section at location (a1, b4).

[0110] The methods and assay systems disclosed herein are capable of multiplexing. For example, Figure 5 provides an address tagging (or "address coding") scheme used in a multiplexed assay. For clarity, two probes TS01 and TS02, specific for target 1 and target 2, respectively, are shown at **520**. Figure 5 shows address tags **510**, comprising a1, a2, a3, a4 and b1, b2, b3 and b4. A delivery or dispensing scheme is shown at **530**. Like the grid exemplified in Figure 4, a1 through a4 are delivered to the sample through horizontal channels, and b1 through b4 are delivered to the sample through vertical channels. The intersections between the horizontal and vertical channels are shown as solid squares. Each intersection can be uniquely identified by the combination of the "a" probe/address tag conjugate and the "b" probe/address tag conjugate delivered to the area in the sample corresponding to the intersection.

[0111] Probes TS01 and TS02 are delivered to the biological sample and allowed to interact with the entire sample. Probes TS01 and TS02 specifically bind to their corresponding targets if the targets are present in the sample. Unbound probes are then removed, for example, by washing. Address tags **510** are then delivered to the biological sample according to the spatial pattern shown at **530**. The address tags are coupled, for example, by ligation (or by extension followed by ligation), to probes that specifically bind to the biological target 1 or biological target 2 in the sample. The coupled constructs (or "probe/address tag conjugates") are then eluted from the biological sample and pooled. In certain embodiments, sequencing adaptors may be added through, e.g., PCR or ligation, if the sequencing adaptors are not already included in the address tags or probe/address tag conjugates are sequenced by, e.g., high throughput or next generation sequencing.

[0112] The pool of resulting assay products is shown at **540**. For example, presence of the "a1T2b1" product in the pool indicates readout is obtained for TS02 at position (a1, b 1) and therefore target 2 is detected at position (a1, b1). Thus, a sequence readout is obtained for

only TS01 at positions (a4, b1), (a4, b2), (a1, b3), (a2, b3), (a3, b3), (a4, b3) and (a4, b4) (positions shown with horizontal lines in spatial pattern **550**), and a sequence readout is obtained for TS02 only at position (a1, b1) (position shown with vertical lines in spatial pattern **550**). A sequence readout is obtained for both TS01 and TS02 at positions (a2, b1), (a3, b1), (a1, b2), (a2, b2), and (a3, b2) (positions shown with cross-hatching in spatial pattern **550**). No sequence readout is obtained for either TS01 or TS02 at (a1, b4), (a2, b4) or (a3, b4) (positions shown without shading in spatial pattern **550**). Thus, in the biological sample, target 1 is detected in a large portion of the left side and at the bottom of the sample, while target 2 is detected only in the upper left portion of the sample, and neither target is detected in the upper right portion of the biological sample. The differential expression of the two biological targets may be mapped back to the biological sample and to the biological structures or cell types in these locations in the biological sample.

[0113] In addition to location information, relative abundance of the biological targets across the multiple sites in the sample can be obtained. For example, if it is found that there are ten times as many a4T1b1 sequences occurring in the data set as compared to a4T1b2 sequences, this would indicate that target 1 is ten times more abundant at location (a4, b1) than at location (a4, b2).

[0114] In the case of nucleotide analysis as shown in Figure 3, by ligating the address tags directly to the probes, only $2n$ probes are needed for n targets. For example, assaying 100 different targets at 10,000 sites in a sample would require 2×100 probes and 2×100 address tags which are to be coupled to the probes. The total count of assay oligonucleotides would be only 400 (200 probes and 200 address tags), not counting universal primers. In contrast, if the address tags are not decoupled from the probes, the total count of assay oligonucleotides would be $(n \times X \text{ positions}) + (n \times Y \text{ positions})$, or in the above example, 20,000 oligonucleotides, not counting universal primer sequences. In other aspects, for each site in the sample, three, four or more address tags may be used, and attached to the probes or one another by varying means and in varying combinations of steps.

[0115] The methods and assay systems disclosed herein are particularly suitable for generating a large amount of information with even a modest number of assays. For example, five or more biological targets assayed at five or more positions in the sample generates 25 or more combinations. Using digital sequencing as a readout, the optimum number of sequence reads per combination depends on the sensitivity and dynamic range required, and can be adjusted. For example, if for each combination on average 100 reads are sampled, the total for 25 combination is 2,500 reads. If 1,000 targets are assayed at 1,000 locations with an average sampling depth of 1,000, then 10^9 reads are required. These numbers, although large, are within the capacity of intrinsically parallel digital sequencing methods, which can generate datasets of billions or even trillions of reads in a reasonable timeframe and at a very low cost per read. Therefore, by varying the numbers of positions or biological targets assayed, or both, and using digital sequencing, large amounts of information can be obtained. In specific aspects, multiple locations are assayed for two or more biological molecules.

[0116] Thus, provided herein is an ability to look at many different biological targets in many locations of a sample at the same time, for example, in the same reaction run. In some aspects, the product of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than about 20. In other aspects, the product of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than about 50. In other embodiments, the product of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than about 100, greater than about 500, greater than about 1,000, greater than about 10,000, greater than about 25,000, greater than about 100,000, greater than about 500,000, or greater than about 1,000,000. It will be appreciated that even much larger numbers can be contemplated. For example, assaying 10,000 targets per location for 10,000 locations in a sample would generate 10^8 different assays. In some aspects, sufficient numbers of sites in a sample can be assayed to reach a resolution on the order of that of single cells. Further, in aspects where high-throughput digital sequencing is employed, the sequences of at least 1,000 probes or probe/address tag conjugates are typically determined in parallel. More typically, using a digital readout, it is desirable to obtain multiple sequence reads for each assay (defined by a target and a location, *i.e.*, by the identities of an identity tag and an address tag of a target). It is desirable to obtain an average of at least 3 copies per assay, and more typically at least 10 or at least 30 copies per assay, depending on the design of the experiment and requirements of the assay. For a quantitative readout with suitable dynamic range, it may be desirable to obtain at least 1,000 reads per assay. Therefore, if 1,000,000 assays are carried out, the number of sequence reads may be 1 billion or more. With high-throughput digital sequencing, and allowing for redundancy, the sequence of at least 10,000 probes or probe/address tag conjugates can be determined in parallel, or the sequence of at least 100,000, 500,000, 1,000,000, 10,000,000, 100,000,000, 1,000,000,000 or more probes or probe/address tag conjugates can be determined in parallel.

[0117] In certain aspects, disclosed herein are methods and assay systems for evaluating differences in the amount and/or activity of biological targets between different locations in a sample and/or between samples. In one embodiment, the method comprises evaluating the differences in quantity of the biological targets at each location in the biological sample. In another aspect, the method comprises comparing spatial patterns of abundance, expression, and/or activity of one or more biological targets among multiple samples.

Spatially Encoded Protein In Situ Assay

[0118] In certain aspects, it is desirable to correlate spatial patterns of a target polynucleotide expression, for example, mRNA expression patterns within a 2D sample, with histological features of the sample. In certain aspects, the histological features may include the expression pattern of a known marker for the sample, for example, a tissue-specific marker, a cell type marker, a cell lineage marker, a cell morphology marker, a cell cycle marker, a cell death marker, a developmental stage marker, a stem cell or progenitor cell marker, a marker for a

differentiated state, an epigenetic marker, a physiological or pathophysiological marker, a marker for a transformed state, a cancer marker, or any combination thereof. In certain aspects, the histological feature comprises tissue morphology, for example, as indicated by the expression pattern of a protein marker. In certain aspects, in order to obtain spatial information of the sample, e.g., histological features of the sample, expression pattern of a protein marker, and/or tissue morphology, imaging techniques have to be used. For instance, immunohistochemical (IHC) and/or immunofluorescent (IF) imaging may need to be used.

[0119] In certain aspects, provided herein are methods called Spatially Encoded Protein In Situ Assays (SEPIA) for multiplexed *in situ* analysis of proteins. In some aspects, SEPIA and related assay systems can obtain spatial information on the relative abundance of many proteins in tissue sections. In certain aspects, the methods and assay systems of the present disclosure are based on the use of antibodies (or other affinity capture agents capable of specifically binding to a target, other than by nucleotide sequence complementarity) that are labeled with an identity tag that identifies the target protein or the antibody, and one or more address tags that identify the location of each of multiple sites in a sample. In one aspect, there are provided at least two address tags for each site, one of the at least two address tags identifying the location in the tissue section in one dimension (for example, an X coordinate) and the other identifying the location in another dimension (for example, a Y coordinate).

[0120] In any of the aspects disclosed herein, the biological target can be a peptide or a protein, and the methods or assay systems can be used to analyze the presence of antibodies, enzymatic and other protein activities, posttranslational modifications, active and non-active forms of peptides, as well as peptide isoforms in a biological sample. Accordingly, the probes may comprise an active region of an enzyme, a binding domain of an immunoglobulin, defined domains of proteins, whole proteins, synthetic peptides, peptides with introduced mutations, aptamers and the like.

[0121] In any of the aspects disclosed herein, the probes can comprise substrates for enzymes or proenzymes, e.g., kinases, phosphatases, zymogens, proteases, or fragments thereof. In certain aspects, the probes may comprise phosphorylation substrates used to detect proteins involved in one or more signal transduction pathways. In other aspects, the probes can comprise specific protease substrates that associate with specific individual proteases or specific classes of proteases. In other aspects, the probes can comprise different processed forms, isoforms and/or domains of an enzyme. In certain embodiments, a protein-based probe can be conjugated or otherwise linked to an oligonucleotide address tag. In preferred aspects, the oligonucleotide address tag may comprise a nucleotide sequence component that allows for identification of the protein probe.

[0122] In preferred aspects, antibodies that are conjugated to oligonucleotide tags are compatible with the address tagging scheme disclosed herein. In certain aspects, provided herein are methods and assay systems that are highly multiplexed, scalable, and high-throughput for determining a spatial pattern of abundance, expression, and/or activity of a target protein across multiple sites in a sample, using a nucleic acid readout and independent

of imaging for the target protein. In preferred aspects, provided herein are methods and assay systems to correlate nucleic acid expression patterns (e.g., DNA or RNA expression patterns) with cell type-specific protein marker abundance without the need for imaging for the protein marker, for example, by immunohistochemical or immunofluorescent imaging. In preferred aspects, spatial resolution of the present methods and assay systems may approach the scale of individual cells. In certain aspects, correlated 2D and 3D maps of RNA and protein abundance can be generated using the present methods and assay systems.

[0123] As shown in Figure 6, in one aspect, a highly multiplexable protein detection assay is carried out on a sample **616** (shown in Figure 6C). In preferred aspects, sample **616** preserves the spatial organization of cells in a tissue. For example, sample **616** can be a paraffin-embedded or fresh-frozen tissue section fixed to a glass slide. Figure 6A shows two probes **620** and **622** specifically bound to a protein target **602** of interest. The first probe **620** may comprise target-binding moiety **608**, associated with oligonucleotide tag **604**. Target-binding moiety **608** and oligonucleotide tag **604** can be conjugated or covalently linked. Target-binding moiety **608** can comprise any affinity capture agents, e.g., antibodies, that specifically bind to protein target **602**. Probe **620** may further comprise address tag **624** and tag **626**. Tag **626** can be a universal priming site for amplification of the assay products and/or an adaptor to enable identification of the address tag **624** and/or oligonucleotide tag **604** and/or other regions of probe **620**, for example, using sequencing technologies. In certain aspects, tag **626** is conjugated or linked to or otherwise associated with address tag **624**, for example, by ligation, extension, ligation followed by extension, or any combination thereof. In one aspect, conjugated, linked or otherwise associated tag **626** and address tag **624** as a whole are conjugated or linked to or otherwise associated with oligonucleotide tag **604**. In alternative aspects, tag **626** and address tag **624** may be separately conjugated or linked to or otherwise associated with probe **620**, for example, at target-binding moiety **608** and/or oligonucleotide tag **604**.

[0124] Similarly, the second probe **622** may comprise target-binding moiety **606**, associated with oligonucleotide tag **610**. Target-binding moiety **606** and oligonucleotide tag **610** can be conjugated or covalently linked. Target-binding moiety **606** can comprise any affinity capture agents, e.g., antibodies, that specifically bind to protein target **602**. Probe **622** may further comprise address tag **628** and tag **630**. Tag **630** can be a universal priming site for amplification of the assay products and/or an adaptor to enable identification of the address tag **628** and/or oligonucleotide tag **610** and/or other regions of probe **622**, for example, using sequencing technologies. In certain aspects, tag **630** is conjugated or linked to or otherwise associated with address tag **628**, for example, by ligation, extension, ligation followed by extension, or any combination thereof. In one aspect, conjugated, linked or otherwise associated tag **630** and address tag **628** as a whole are conjugated or linked to or otherwise associated with oligonucleotide tag **610**. In alternative aspects, tag **630** and address tag **628** may be separately conjugated or linked to or otherwise associated with probe **622**, for example, at target-binding moiety **606** and/or oligonucleotide tag **610**.

[0125] In certain aspects, target-binding moiety **606** and target-binding moiety **608** bind to

adjacent sites on target **602**, so that two free ends of oligonucleotide tags **604** and **610** are brought close to each other. In one aspect, oligonucleotide tags **604** and **610** may be ligated and the ligation product assayed. In other aspects, one or both of oligonucleotide tags **604** and **610** may be extended and then ligated to the other probe to form an extended probe comprising target target-binding moiety **606** and target-binding moiety **608**. For example, a DNA ligase may be added together with a splint to join two free ends of oligonucleotide tags **604** and **610**, and the DNA ligated product can serve as the template detectable by real-time PCR and/or various sequencing technologies. Such a dual targeting approach may be used to increase assay specificity. Other aspects of the dual targeting approach that converts specific protein detection into nucleic acid analysis, including the proximity ligation assay described in Fredriksson et al., 2002, Nat Biotechnol 20, 473-7, may be used in the methods and assay systems of the present disclosure. It is also within the present disclosure that in certain aspects, target-binding moiety **606** and target-binding moiety **608** may bind to different protein targets. When the protein targets are in close proximity, for example, when the two are in the same complex or brought into contact with each other in a reaction, a ligation product may be formed between oligonucleotide tags **604** and **610** and detected.

[0126] In certain aspects, a primary antibody and a secondary antibody may be used. For example, target-binding moiety **606** and/or target-binding moiety **608**, instead of specifically binding to target **602** directly, may specifically bind to a primary antibody that specifically recognizes target **602**. In this case, target-binding moiety **606** and/or target-binding moiety **608** may be comprised in a secondary antibody. In certain aspects, the approach involving a primary antibody and a secondary antibody may be suitable when target expression is low in a sample, because one molecule of target **602** may be able to bind multiple molecules of a primary antibody, thereby amplifying the signal.

[0127] In other aspects, a biological target **632** is assayed according to the combinatorial address tagging scheme shown in Figure 6D. Two probes **650** and **652** specifically bind to the biological target **632** of interest. In one aspect, the first probe **650** comprises target-binding moiety **638**, associated with oligonucleotide tag **634**, and the second probe **652** comprises target-binding moiety **636**, associated with oligonucleotide tag **640**. Target-binding moiety **638** and oligonucleotide tag **634** (or target-binding moiety **636** and oligonucleotide tag **640**) can be conjugated or covalently linked. In particular aspects, target-binding moiety **638** or **636** comprises an affinity capture agent, e.g., an antibody, that specifically binds to target **632**. In certain aspects, target **632** comprises a protein moiety, an oligosaccharide or polysaccharide moiety, a fatty acid moiety, and/or a nucleic acid moiety. In some aspect, each probe has a portion that specifically binds to an adaptor **662**, for example, by specific nucleic acid hybridization. In one aspect, oligonucleotide tag **634** or **640** (or a portion thereof) specifically hybridizes to the adaptor. The adaptor can specifically bind to the following combinations: 1) the 5' portion of oligonucleotide tag **634** and the 3' portion of oligonucleotide tag **640**; 2) the 3' portion of oligonucleotide tag **634** and the 5' portion of oligonucleotide tag **640**; 3) the 5' portion of oligonucleotide tag **634** and the 5' portion of oligonucleotide tag **640**; or 4) the 3' portion of oligonucleotide tag **634** and the 3' portion of oligonucleotide tag **640**. In certain aspects, oligonucleotide tag **634** or **640** is a linear molecule, a branched molecule, a circular molecule,

or a combination thereof. After binding of the two probes to the biological target and the adaptor to the two probes, address tags can be delivered to the sample and coupled to the adaptor. For example, the adaptor can be tagged with address tag **654**, associated with tag **656**, and/or with address tag **658**, associated with tag **660**. Tags **656** and **660** can be universal priming sites for amplification of the assay products or sequences to enable identification of the address tags and/or other regions of adaptor/address tag conjugates, for example, using sequencing technologies. In certain aspects, the address tags are tagged at the same end of the adaptor, or at different ends of the adaptor. In other aspects, an address tag and/or tag **656** or **660** can be pre-coupled to the adaptor, and the adaptor/address tag or adaptor/tag conjugate or complex is then delivered to the sample in order to bind to the probe bound to the biological target.

[0128] In certain aspects, all or a portion of the adaptor/address tag conjugate sequence is determined, for example, by nucleic acid sequencing. In other aspects, all or a portion of the oligonucleotide tag sequence, and/or all or a portion of the adaptor/address tag conjugate sequence, is determined. For example, a first address tag can be coupled to oligonucleotide tag **634**, and a second address tag can be coupled to adaptor **662**. The duplex formed between oligonucleotide tag **634** and adaptor **662** can be subjected to extension and sequencing, to generate a conjugate that comprises sequences of the first address tag, all or a portion of oligonucleotide tag **634**, all or a portion of adaptor **662**, and the second address tag.

[0129] The tagging scheme is not limited to the use of two or more probes for the same biological target. For example, in cases where one probe is used, a tag (e.g., an address tag, an adaptor for ligation, or a universal sequencing primer or amplification primer sequence) can be coupled to an adaptor that specifically binds to the probe, rather than to the probe itself.

[0130] Additional details of the polynucleotide-protein conjugates used in the present disclosure are disclosed in United States Provisional Patent Application Serial No. 61/902,105, filed November 8, 2013, entitled "Polynucleotide conjugates and methods for analyte detection,".

[0131] In some aspects, more than one adaptor is used. For example, a pair of adaptors is used to specifically bind the oligonucleotide tag **634** and **640**, respectively. In certain aspects, the adaptors of the pair are DNA molecules that: 1) hybridize or otherwise bind to the protein-DNA conjugates, for example, probe **650** or **652**; 2) have free 3' and/or 5' ends that enable the encoding sequences (e.g., address tags **654** and **658**) to be attached in a subsequent step or steps, for example, by ligation; 3) are in a form where they can be joined if they are co-localized or in proximity to each other. In some aspects, part of the oligonucleotide portion of probe **650** or **652** acts as a splint to enable ligation, or extension and ligation, of the adaptors in the adaptor pair. Additional tags (e.g., an address tag, an adaptor for ligation, or a universal sequencing or amplification primer sequence) can be coupled to the adaptor generated by joining the adaptor pair.

[0132] In another aspect, a method disclosed herein makes it easier to carry out protein-based

assays at the same time as nucleic-acid based assays. For example, the adaptors can be designed so that they are compatible with the same encoding oligonucleotides used for the nucleic-acid based assays, *e.g.*, RNA-based assays. Thus, two types of binding assays (*i.e.*, detecting a protein target using a protein-polynucleotide conjugate, and detecting a nucleic acid target using a nucleic acid probe) can be carried out in the same reaction volume or in the same experimental run, and the spatial addressing can be performed on both types of probes simultaneously.

[0133] In yet another aspect, the present disclosure provides a control for assays detecting a protein target or a biological target comprising a protein moiety. For example, the nucleic acid portion of the protein-nucleic acid conjugate is used to hybridize to a nucleic acid in the sample. This anchors an "artificial" protein (known composition and abundance based on the abundance of the hybridizing sequence) in the sample. The "artificial" protein can then be detected using a number of means, including protein-binding spatially-addressed assays disclosed herein. The approach is not limited to proteins. For example, small molecules, such as haptens, can also be used. In one aspect, Figure 6E illustrates the general concept of a method of detecting an RNA with known composition and abundance in the sample, thereby providing a control for the detection of other targets (*e.g.*, protein targets) in the sample. In Figure 6E, conjugates **662** and **664** each comprises a nucleic acid portion and an antibody-binding portion (circle indicates the antibody-binding portion of conjugate **662**, and triangle indicates the antibody-binding portion of conjugate **664**). In certain aspects, RNA **666** with known composition and abundance in the sample is specifically bound by the nucleic acid portions of conjugates **662** and **664**. In some aspects, the composition and/or abundance of RNA **666** are determined experimentally, for example, using a method of the present disclosure, and in specific aspects, simultaneously with the detection of the protein target. In other aspects, the composition and/or abundance of RNA **666** is derived from prior knowledge or knowledge in the art. In particular aspects, the antibody-binding portions can be HA or FLAG, and the antibody portions of probes **650** and **652** can be an anti-HA antibody or an anti-FLAG antibody, for example, polyclonal or monoclonal antibodies. Other protein-antibody binding pairs are known in the art and can be used in the present disclosure.

[0134] Figure 6B shows an address tagging scheme that may be used for 100 unique sites in a sample. For example, twenty probes/address tag conjugates X1 through X10 and Y1 through Y10 can be used, with each of X1 through X10 comprising an address tag **624** and each of Y1 through Y10 comprising an address tag **628**. The address tag comprised in each of X1 through X10 and Y1 through Y10 may be uniquely identified. Probe/address tag conjugate X9, for example, is delivered to the biological sample in the ninth vertical channel in **612**. Whereas the "X" probe/address tag conjugates are delivered in ten vertical channels, the "Y" probe/address tag conjugates are delivered in ten horizontal channels as shown in **614**. For example, probe/address tag conjugate Y1 is delivered to the biological sample in the first horizontal channel of **614**. In other aspects, the "X" tags may be referred to as the "a" tags and the "Y" tags as "b" tags.

[0135] Figure 6C shows a representative tissue section **616** coincident with grid **618**. The

arrows show how the "X" probe/address tag conjugates and the "Y" probe/address tag conjugates are delivered on grid **618** that is coincident with tissue section **616**. If, once analyzed, probe/address tag conjugates X9 and Y1, e.g., are associated with a target, then that target is present in the tissue section at location (X9, Y1).

[0136] Any suitable configuration of the oligonucleotide/antibody (or other target-specific binder) conjugate may be used to convert specific protein detection into nucleic acid analysis. In certain aspects, for example, as shown in Figure 7A, probe **708** specifically binds to protein target **702**. Probe **708** may comprise target-binding moiety **704**, associated with oligonucleotide tag **706**. Target-binding moiety **704** and oligonucleotide tag **706** can be conjugated or covalently linked. Target-binding moiety **704** can comprise any affinity capture agents, e.g., antibodies, that specifically bind to protein target **702**. Probe **708** may further comprise "X" address tag **710** and "Y" address tag **712**. Both address tags **710** and **712** may be conjugated to a universal priming site for amplification of the assay products and/or an adaptor (not shown in Figure 7) to enable identification of the address tags **710** and **712** and/or oligonucleotide tag **706** and/or other regions of probe **708**, for example, using sequencing technologies. Conjugation of the various tags may be accomplished by ligation, extension, ligation followed by extension, or any combination thereof. In some aspects, address tags **710** and **712** are conjugated to one side of oligonucleotide tag **706** or the other (*i.e.*, 5' or 3' of the sequence), respectively. In alternative aspects, both address tags **710** and **712** may be conjugated to either 5' or 3' of oligonucleotide tag **706**. For example, address tags **710** and **712** may be directly or indirectly conjugated, and address tag **710** or **712** may be directly or indirectly conjugated to either 5' or 3' of oligonucleotide tag **706**.

[0137] In other aspects, for example, as shown in Figure 7B, probe **720** specifically binds to protein target **714**. Probe **720** may comprise target-binding moiety **716**, conjugated, linked, or otherwise associated with oligonucleotide tag **718**. Target-binding moiety **716** can comprise any affinity capture agents, e.g., antibodies, that specifically bind to protein target **714**. Probe **720** may further comprise oligonucleotide sequence **722** that specifically hybridizes to oligonucleotide tag **718**. In one embodiment, sequence **722** is complementary to oligonucleotide tag **718**. Sequence **722** may be conjugated to "X" address tag **724** and "Y" address tag **726**. Both address tags **724** and **726** may be conjugated to a universal priming site for amplification of the assay products and/or an adaptor (not shown in Figure 7) to enable identification of the address tags **724** and **726** and/or sequence **722** and/or other regions of probe **720**, for example, using sequencing technologies. Conjugation of the various tags may be accomplished by ligation, extension, ligation followed by extension, or any combination thereof. Similar to Figure 7A, address tags **724** and/or **726** can be conjugated to either side of oligonucleotide sequence **722** (*i.e.*, 5' or 3' of the sequence), either directly or indirectly.

[0138] In further aspects, for example, as shown in Figure 7C, a "2-antibody" format may be used. The "2-antibody" format is similar to the dual targeting approach discussed above, for example, in Figure 6. In this aspect, two antibodies specific for a protein target are conjugated to an oligonucleotide, which can be directly or indirectly conjugated to the "X" and "Y" address tags and a universal priming site for amplification of assay products and/or an adaptor for

sequencing. In some aspects, the two antibodies may bind to different epitopes or sites on the protein target. In preferred aspects, binding of both antibodies to the target is required to generate a signal, thus providing higher specificity than using only one antibody. It is also contemplated that more than two antibodies may be conjugated to an oligonucleotide and used in the methods and assay systems of the present disclosure.

[0139] As disclosed herein, the methods and assay systems permit high levels of multiplexing. In one aspect, the probes can be delivered over the entire surface of a 2D sample in a bulk process, and then address tagged by delivering the address tags in a spatially defined pattern. For example, two sets of address tags (the "X" and "Y" tags) can be used in a combinatorial fashion as discussed *supra*. Once the in situ assay is completed, the assay products are eluted and sequenced. The address tag sequence information identifies the location at which the assay is performed, and the probe sequence information (the identity tag) identifies the protein that is targeted. In one aspect, the frequency of a particular assay product (for example, a sequencing product) in the digital readout can be used to infer the relative abundance of its target in the sample. This information can then be associated with other information, including conventional histological information, and/or transcript abundance obtained via the related Spatially Encoded Genomic Assays (SEGA). In preferred aspects, the methods and assay systems do not depend on imaging techniques for the spatial information of the target protein. Instead, in preferred aspects, the spatial pattern of the target protein abundance, expression, and/or activity can be determined by sequencing.

[0140] In one aspect, in order to integrate the protein and gene expression assays, the same address tagging scheme is compatible with and can be used for both assay types. For example, for each of multiple sites in a sample, the same combination of "X" and "Y" address tags can be tagged to an antibody-DNA conjugate for a target protein, and to a probe for a target polynucleotide sequence. In one aspect, the target polynucleotide or the complement thereof encodes all or a portion of the target protein. Therefore, for each site in the sample, the abundance, expression, and/or activity of the target protein and its corresponding polynucleotide can be detected by assaying for sequencing products with the same set of address tags. In preferred aspects, the step of analyzing probes or probe/address tag conjugates bound to the target protein and the step of analyzing probes or probe/address tag conjugates bound to the target polynucleotide can be performed in parallel in the same reaction run. In alternative aspects, different address tags may be coupled to an antibody-DNA conjugate for a target protein, and to a probe for a target polynucleotide sequence, to determine the abundance, expression, and/or activity of the target protein and the target polynucleotide at a given site. Assay results for the target protein and the target polynucleotide can then be integrated for each site in the sample.

[0141] Various methods can be used to form an amplifiable construct, for example, by using ligation of proximal probes followed by sequential ligation of a pair of spatial encoding adaptors (address tags) as shown in Figure 8A. In one aspect, two DNA probes are hybridized proximal to one another on an RNA target (or template). The probes are subsequently ligated to one another and the quantity of the ligated pair is taken as a measure of the amount of the target

present in the sample. In certain cases, however, the efficiency of T4 DNA ligase is reduced when the ligation reaction occurs on an RNA template as compared to a DNA template. In other cases, the ligation efficiency is dependent on the sequence of the DNA probes that are being joined, the particularly on the identity of the first few bases on either side of the junction. In some aspects, a method disclosed herein mitigates both problems. Figure 8B shows the general principle of the method. In this case, instead of using probes that hybridize in direct proximity on an RNA target, the probes are separated by some distance with non-hybridizing overhanging sequences on their proximal ends. These overhanging sequences are designed to be complementary to a short DNA splint. This splint can be universal for all the probe pairs in a multiplexed assay or can be specific for a given probe pair or subset of probe pairs. The distance between the two probes in a pair can be adjusted to optimize ligation efficiency. There is flexibility in this distance, which provides an additional degree of freedom when designing probes versus the use of proximal probes. Once the probes are hybridized to the RNA target, excess probes are washed away. The splint is hybridized to the overhanging regions at the proximal ends of the probes, and the probes are joined by enzymatic ligation. After ligation, the remaining steps of the assay are performed, for example, ligating the spatial encoding adaptors to each end of the ligated probe pair. In certain aspects, since DNA splinted ligation is more efficient than RNA splinted ligation, a method disclosed herein improves the efficiency at which the two probes are joined. In addition, using a universal splint eliminates the sequence-dependent variation in ligation efficiency between the multiple probe sets in a multiplexed *in situ* assay. In another aspect, probes can be more easily designed, and more suitable probe sets can be designed, due to increased freedom of varying the distance between probes.

Reagent Delivery Systems

[0142] The reagent delivery system of the present disclosure and which does not belong to the invention includes instrumentation that allows the delivery of reagents to discrete portions of the biological sample, maintaining the integrity of the spatial patterns of the addressing scheme. Reagent delivery systems of the present assay systems comprise optional imaging means, reagent delivery hardware and control software. Reagent delivery can be achieved in a number of different ways. It should be noted that reagents may be delivered to many different biological samples at one time. A single tissue section has been exemplified herein; however, multiple biological samples may be manipulated and analyzed simultaneously. For example, serial sections of a tissue sample can be analyzed in parallel and the data combined to build a 3D map.

[0143] Integral to the assay system of the present disclosure is instrumentation that allows for spatial patterning of reagents onto the biological sample. Technologies for formulating and delivering both biological molecules (e.g., oligonucleotides or antibodies) and chemical reagents (e.g., small molecules or dNTPs) are known in the art, and uses of these instrument systems are known to one skilled in the art and easily adaptable to the assay systems of the present disclosure. One example of a suitable reagent delivery system is the Labcyte™ Echo

acoustic liquid handler, which can be used to deliver nanoliter scale droplets containing biological molecules with high precision and reproducibility. One skilled in the art could incorporate this reagent delivery device into the overall system using software to specify the locations to which reagents should be delivered.

[0144] Other instruments that can be used for the delivery of agents and/or coding identifiers onto biological samples include, but are not limited to, ink jet spotting; mechanical spotting by means of pin, pen or capillary; micro contact printing; photochemical or photolithographic methods; and the like. For several applications, it may be preferred to segment or sequester certain areas of the biological samples into one or more assay areas for different reagent distributions and/or biological target determination. The assay areas may be physically separated using barriers or channels.

[0145] In one exemplary aspect, the reagent delivery system may be a flow-based system. The flow-based systems for reagent delivery in the present invention can include instrumentation such as one or more pumps, valves, fluid reservoirs, channels, and/or reagent storage cells. Reagent delivery systems are configured to move fluid to contact a discrete section of the biological sample. Movement of the reagents can be driven by a pump disposed, for example, downstream of the fluid reagents. The pump can drive each fluid reagent to (and past) the reaction compartment. Alternatively, reagents may be driven through the fluid by gravity. US Pub. Nos. 20070166725 and 20050239192 disclose certain general purpose fluidics tools that can be used with the assay systems of the present disclosure, allowing for the precise manipulation of gases, liquids and solids to accomplish very complex analytical manipulations with relatively simple hardware.

[0146] In a more specific example, one or more flow-cells can be attached to the substrate-affixed biological sample from above. The flow-cell can include inlet and outlet tubes connected thereto and optionally an external pump is used to deliver reagents to the flow-cell and across the biological sample. The flow cells are configured to deliver reagents only to certain portions of the biological sample, restricting the amount and type of reagent delivered to any specific section of the biological sample.

[0147] In another aspect, a microfluidic system can be integrated into the substrate upon which the biological sample is disposed or externally attached on top of the substrate. Microfluidic passages for holding and carrying fluid may be formed on and/or above the planar substrate by a fluidics layer abutted to the substrate. Fluid reagents can be selected and delivered according to selective opening and closing of valves disposed between reagent reservoirs.

[0148] Pumps generally include any mechanism for moving fluid and/or reagents disposed in fluid. In some examples, the pump can be configured to move fluid and/or reagents through passages with small volumes (i.e., microfluidic structures). The pump can operate mechanically by exerting a positive or negative pressure on fluid and/or on a structure carrying fluid, electrically by appropriate application of an electric field(s), or both, among other means.

Exemplary mechanical pumps may include syringe pumps, peristaltic pumps, rotary pumps, pressurized gas, pipettors, *etc.* Mechanical pumps may be micromachined, molded, *etc.* Exemplary electrical pumps may include electrodes and may operate by electrophoresis, electroendosmosis, electrocapillarity, dielectrophoresis (including traveling wave forms thereof), and/or the like.

[0149] Valves generally include any mechanism for regulating the passage of fluid through a channel. Valves can include, for example, deformable members that can be selectively deformed to partially or completely close a channel, a movable projection that can be selectively extended into a channel to partially or completely block a channel, an electrocapillary structure, and/or the like.

[0150] An open gasket can be attached to the top of the biological sample and the sample and reagents can be injected into the gasket. Suitable gasket materials include, but are not limited to, neoprene, nitrile, and silicone rubber. Alternatively, a watertight reaction chamber may be formed by a gasket sandwiched between the biological sample on the substrate and a chemically inert, water resistant material such as, but not limited to, black-anodized aluminum, thermoplastics (*e.g.*, polystyrene, polycarbonate, *etc.*), glass, *etc.*

[0151] In an optional embodiment, the assay system comprises imaging means to determine features and organization of the biological sample of interest. The images obtained, *e.g.*, may be used to design the delivery pattern of the reagents. Imaging means are optional, as an individual can instead view the biological sample using, *e.g.*, a microscope, analyze the organization of the biological sample, and specify a spatial pattern for delivery assay reagents. If included, the delivery system can comprise a microcircuit arrangement including an imager, such as a CCD or IGFET-based (*e.g.*, CMOS-based) imager and an ultrasonic sprayer for reagent delivery such as described in US Pub. No. 20090197326. Also, it should be noted that although an X-Y grid configuration is illustrated herein, other configurations can be used, such as, *e.g.*, following the topology of a tissue sample; targeting certain groups of cells, cell layers and/or cell types in a tissue, and the like.

[0152] In yet another alternative, the reagent delivery system controls the delivery of reagents to specific patterns on a biological sample surface using semiconductor techniques such as masking and spraying. Specific areas of a biological sample can be protected from exposure to reagents through use of a mask to protect specific areas from exposure. The reagents may be introduced to the biological sample using conventional techniques such as spraying or fluid flow. The use of masked delivery results in a patterned delivery scheme on the substrate surface.

[0153] In one aspect, the reagent delivery instrumentation is based on inkjet printing technology. There are a variety of different ink jetting mechanisms (*e.g.*, thermal, piezoelectric) and compatibility has been shown with aqueous and organic ink formulations. Sets of independently actuated nozzles can be used to deliver multiple reagents at the same time, and very high resolutions are achieved.

[0154] In order to target specific sites of interest, an informative image of the biological sample to be assayed may be used to assist in the reagent delivery methods and associated encoding scheme. Sample regions of the biological sample can be identified using image processing (e.g., images of cell types differentiated by immunohistochemistry or other staining chemistries) integrated with other features of the assay system. In some aspects, software is used to automatically translate image information into a reagent delivery pattern. In some aspects, a mechanism to register and align very precisely the biological sample for reagent delivery is an important component of the assay systems. Mechanisms such as the use of fiducial markers on slides and/or other very accurate physical positioning systems can be adapted to this purpose.

[0155] The present methods and assay systems may comprise a complete suite of software tailored to the methods or assay systems. Optionally, oligonucleotide design software is used to design the encoding nucleotides (and in aspects where nucleic acids are assayed, the target-specific oligonucleotides) for the specific assay to be run, and may be integrated as a part of the system. Also optionally, algorithms and software for reagent delivery and data analysis (*i.e.*, sequence analysis) may be integrated to determine assay results. Integrated data analysis is particularly useful, as the type of dataset that is generated may be massive as a consequence of scale. Algorithms and software tools that are specifically designed for analysis of the spatially-associated data generated by the assay systems, including pattern-analysis software and visualization tools, enhance the value of the data generated by the assay systems.

[0156] In certain aspects, the assay system comprises processes for making and carrying out the quality control of reagents, e.g., the integrity and sequence fidelity of oligonucleotide pools. In particular, reagents are formulated according to factors such as volatility, stability at key temperatures, and chemical compatibility for compatibility with the reagent delivery instrumentation and may be analyzed by instrumentation integrated within the assay system.

Applications of Assay System

[0157] It will be apparent to one skilled in the art upon reading the present disclosure that there are numerous important areas of biological research, diagnostics, and drug development that will benefit from a high throughput multiplexed assay system that can measure simultaneously the amount and spatial location of a biological target in a biological sample. For example, combining the ability to estimate the relative abundance of different RNA transcripts with the ability to reconstruct an image of spatial patterns of abundance across many locations, which may be as small as or even smaller than individual cells, in a tissue enables many different areas of basic research.

[0158] In one aspect, the assay systems and devices disclosed herein can discriminate different tissue types on the basis of tissue-specific differences in gene expression. In one

aspect, the assay systems and devices disclosed herein can be used to assay and discriminate mRNA and hnRNA, and therefore can be used for parallel analysis of RNA processing *in situ*. In one aspect, probes are designed to target introns and/or exons. In one aspect, intronic probes give signal from hnRNA, but not from mRNA. The gDNA background signal can be measured using selective pretreatments, with DNase and/or RNase. In one aspect, splice-site specific probes that are selective for spliced RNAs may be designed and used. In certain aspects, a combination of intronic probes, exonic probes, and/or splice-site specific probes may be used to identify the relative level of processing intermediates and their differences between different cells in a tissue section. In general, RNA may be bound to proteins of various types, and hnRNA, in particular, is complexed with proteins to form hnRNP (heterogeneous nuclear ribonucleoprotein). In one aspect, the devices and assay systems disclosed herein can be used to perform highly parallel *in situ* footprinting experiments. In certain aspects, instead of targeting 1,000 different RNAs, probes can be tiled densely through a smaller number of RNAs in order to generate a signal profile along the molecule. Relative changes in this profile between cell types would then indicate differences in availability of the RNA, at the specific locations assayed.

In one example, 3-dimensional patterns of gene expression are determined by analyzing a series of tissue sections, in a manner analogous to image reconstruction in CT scanning. Such a method can be used to measure changes in gene expression in disease pathology, e.g., in cancerous

tissue and/or a tissue upon injury, inflammation, or infection. With the assay systems disclosed herein but that are not part of the invention, more detailed information on gene expression and protein localization in complex tissues is obtained, leading to new insights into the function and regulation both in normal and diseased states, and provides new hypotheses that can be tested. For example, an assay system of the invention may enable some of the insights gained from many individual studies and larger programs like ENCODE (Birney, et al., Nature, 447:799-816 (2007)) and modENCODE to be integrated at the tissue level. The assay systems also aid computational efforts to model interacting networks of gene expression in the field of systems biology.

[0159] The assay systems also provide a novel approach to analysis of somatic variation, e.g., somatic mutations in cancer or variability in response to infectious organisms. For example, tumors are typically highly heterogeneous, containing cancer cells as well as genetically normal cells in an abnormal local environment. Cancer cells undergo mutation and selection, and in this process it is not unusual for local clones to develop. Identifying relatively rare somatic mutations in the context of tumors may enable the study of the role of key mutations in the selection of clonal variants. Transcriptional patterns associated with angiogenesis, inflammation, or other cancer-related processes in both cancer and genetically normal cells can be analyzed for insights into cancer biology and assist in the development of new therapeutic agents for the treatment of cancers. In another example, individuals have varying susceptibility to infectious organisms, and the assay systems of the invention can be used to study the interaction between microbes and tissues or the various cell types within the tissue.

[0160] Importantly, in addition to providing spatially-associated information, the invention

allows a great increase in the sensitivity of detecting rare mutations, as signal to noise can be dramatically increased since only a small location is assayed in any given reaction. In a typical assay for rare mutations in a mixed sample, the sample is treated in bulk, *i.e.*, nucleic acids are extracted from many cells into a single pool. Thus, if a mutation is present in one cell in 10,000, it must be detected against a background of normal DNA from ~10,000 cells. In contrast, with the assay systems of the invention many cells can be analyzed, but individual cells or small groups of cells would be identified by the spatial coding system. Therefore, in the assay systems disclosed herein but that are not part of the present invention, background is reduced by orders of magnitude, greatly increasing sensitivity. Furthermore, the spatial organization of mutant cells can be observed, which may be particularly important in detecting key mutations in tissue sections in cancer. Already molecular histological analyses are yielding insights into cancer biology and may have potential for use in diagnostics. The technology of the invention promises to greatly increase the power of such approaches.

[0161] The following exemplary examples are intended to further describe and illustrate various aspects of the invention.

Example 1 Proof of concept of the addressing scheme and scalability

[0162] A model system was developed using a microarray to demonstrate a working multiplexed spatially encoded abundance assays for polynucleotide targets. The basic design validates the concept of the assay, and the addressing scheme, and establishes a working assay prior to addressing issues related to the analysis of a more complicated biological sample.

[0163] A microarray was used as a proxy for a tissue section. The target sequences of the microarray were fully specified, so that the composition of the targets was known and was varied systematically, simplifying analysis by next-generation sequencing. One of skill in the art would appreciate that similar assays can be performed on various samples including tissue sections, and for various targets including polynucleotide or protein targets, as well as other biological targets, according to the present disclosure.

A 16-plex × 8-site assay using 8-section microarray as artificial sample

[0164] This 16-plex × 8-site assay was performed using a custom DNA microarray (Agilent) as an artificial sample. Eight sites were used because of the commercial availability of 8-section microarrays. Sixteen different target sequences were each assayed over a 128-fold range in DNA amount. Differences in DNA amount were obtained by varying the surface area over which each sequence was synthesized. Differences in DNA amount were detected over the entire range for all sixteen targets, using next-generation sequencing as the readout. This example demonstrated a working multiplex assay using a microarray as an artificial sample,

and the spatial encoding accuracy for the model system.

Example 2 A demonstration of spatial encoding using a spotted microarray

[0165] Scalability of both the spatial addressing and assay systems is demonstrated by carrying out a 24-plex × 24-site assay using a microarray model system.

[0166] The amount of biological target, here a DNA target sequence, at each assay location is systematically varied on the microarray substrate. For example, in a microarray with 50 micron spot size (center to center), a 1 mm² area contains ~400 spots. The region around each site is optionally occupied by a region that is devoid of these spots to allow individual resolvability of the target sequences. Alternatively, the spots may be clustered, with two or more directly adjacent spots surrounded by or adjacent to a region that is devoid of target sequences.

[0167] In order to demonstrate that spatial addressing or encoding is accurate, the sites comprise different target compositions to show that the assay readout matches the expected composition of each site. With 24 target sequences, a simple digital pattern is made with each site having a different set of 12 targets present and 12 targets absent, to make a binary code (0 = absent, 1 = present). The assay readout is then determined to show that the detected regions match the expected signal after spatial decoding. In this particular example, the code (address tag) space is large enough (2^{24}) so that even a few errors would not result in different codes being mixed up. Moreover, this design allows identification of errors and allows estimation not only of accuracy of spatial encoding but also of accuracy calling the presence or absence of target sequences.

[0168] The ability to detect quantitative differences is evaluated by generating dose-response curves for each of the 24 assays that are carried out at each site in a 24-site assay. This allows estimation of the limit of detection, dynamic range, and power to detect a given fold-change across the range.

[0169] In one aspect, a latin square design is used to represent individual targets at different ratios by varying the number of features for each target. In other words, with multiple spots in a site, the number of spots allocated to each of the 24 target sequences can be varied and each of the 24 sites can have a different composition. A 1 × 3 inch microarray is sufficiently large to permit multiple replicates. This larger set of 24 sequences will require deconvolution, and this is accomplished by using high throughput techniques such as next-generation sequencing technologies (e.g., SOLiD™ technology (Life Technologies, Inc., Carlsbad, Calif.) or Genome Analyzer (Illumina, Inc., San Diego, Calif.)). The use of the 24-plex assay demonstrates both the accuracy of spatial encoding and decoding, and the quantitative response of the assay system.

Example 3 Assays for preserved samples and biological samples

[0170] Genomic DNA is assayed in order to characterize variation in coding and regulatory sequences, such as single nucleotide polymorphisms (SNPs) or mutations, small insertions and deletions (indels), copy number variants such as gene deletions or amplifications, and genetic rearrangements such as translocations, all of which may be functionally significant in cancer and other diseases. Genomic sequence variation as a function of position in the sample may indicate somatic mosaicism in the sample. In cancer samples, mutations may provide prognostic or diagnostic markers that may be useful in determining the best course of treatment. Mutations may identify regions of the sample that contain cancer cells and assist in distinguishing them from normal cells, or cells in the tumor microenvironment that are genetically normal at the sequence level but perturbed in other ways as a result of the influence of cancer cells. In order to distinguish signal generated from DNA targets from those generated by RNA targets, probes can be designed to hybridize to non-coding sequences that are not transcribed. Alternatively, or in order to confirm the specificity of DNA targeting, RNA may be degraded by treatment with RNase. Genomic DNA is also assayed in order to obtain information about its organization and to provide information on the state of activation of certain genes. For example, the ability of probes to bind to DNA may be used as an indicator of whether DNA is condensed or otherwise inaccessible, or whether DNA is in an open conformation for transcription. This type of determination can benefit from comparative analysis of samples in which genes are differentially active. Similarly it may be useful to relate information about RNA and/or protein abundance to information about the activation state of genes. Other types of information are obtained from analysis of epigenetic markers associated with genomic DNA, such as methylation state and the presence of histones and other proteins and modifications.

[0171] The handling of small absolute numbers of product molecules generated from very small or compromised samples are enhanced to counter the issue of low recovery efficiency; that is, elution is efficient and losses resulting from adsorption of molecules to surfaces are prevented. An approach to addressing the latter issue is to include a carrier material, such as glycogen or carrier nucleic acids.

[0172] In order to adapt the assay to a biological sample and make the tissue section RNA assays as informative as possible, pre-existing information on expression levels in specific tissues to target transcripts across a range of abundances are used in the assay design. Both high abundance transcripts, as well as some medium and low abundance transcripts, are targeted to enable an initial assessment of the quantitative performance characteristics of the assay. In this assay, a control RNA template is immobilized to a solid support in order to create an artificial system. The assay is performed using T4 DNA ligase, which can repair nicks in DNA/RNA hybrids. Assays are carried out on matched slides, or different sections of the same slide, where in one case gDNA is assayed and in the other RNA is assayed. When assaying gDNA the slide can be pretreated with RNase, and when assaying RNA the slide is pretreated with DNase. Results of the assay are confirmed by extracting gDNA or RNA and quantitating the relative amounts by qPCR or RT-qPCR, respectively.

Example 4**Multiplex spatially encoded polynucleotide abundance assays**

[0173] This example describes representative multiplex spatially encoded abundance assays for polynucleotide targets. One of skill in the art would appreciate that similar assays can be performed for protein targets, as well as other biological targets, according to the present disclosures.

A 57-plex assay using formalin-fixed, paraffin-embedded (FFPE) samples

[0174] A scheme using ligation of proximal probes followed by sequential ligation of a pair of spatial encoding adaptors (address tags) was used to form an amplifiable construct. For example, as shown in Figure 8A, two target-specific probe oligos were ligated together following *in situ* hybridization. A unique adaptor or address tag encoding the X position was introduced via a microfluidic channel and ligated to the 5' end of the probes. A second address tag encoding the Y position was similarly installed to the 3' end of the probes. The address tags contained universal priming sites that allowed installation of additional sequencing adaptors via PCR. The final construct is a substrate for next-generation sequencing.

[0175] A 57-plex assay was performed using a pool of probes for 57 targets on commercially sourced FFPE sections of normal human liver and pancreas (Pantomics). The pool included probes for 18 liver specific targets, 19 pancreas specific targets, 4 housekeeping targets, 6 custom-generated negative controls sequences, and 10 pluripotency markers. All liver-specific probes were strongly enriched in liver and all but 3 of the pancreas-specific probes were strongly enriched in pancreas. These 3 probes had very few total counts so it is likely that they were sequences that hybridized or ligated inefficiently and thus were not reporting accurately. The results of this assay were consistent with published data for expression in normal liver and pancreas (BioGPS, available at biogps.org/#goto=welcome).

[0176] A number of different reagent delivery technologies, including random-access methods such as inkjet and pin-spotting, can be used for the multiplex assays. A system using microfluidic flow-channel devices was chosen for several reasons. First, soft-lithographic techniques allow rapid development of such devices at a fraction of the cost and time needed to develop or buy a suitable instrument for printing or spotting reagents. Second, the size of the sampling area can be strictly defined using microfluidic devices, whereas printed droplets of reagent would likely spread non-uniformly on the surface of an FFPE sample and yield sampling areas of varying size and shape. Third, the reagent delivery system using microfluidic devices does not require precise alignment of the sample. This feature allows sequential

ligation of the two encoding positional adaptors (i.e., the two address tags). Compared to simultaneous ligation of the two address tags, sequential ligation minimizes the formation of undesired products. For reagent delivery technologies using inkjet or pin-spotting, the location of each droplet or spot of the first address tag must coincide with a droplet or spot of the second address tag in order to form the full construct during sequential ligation. This would require that precise registration of the sample be preserved throughout both printing steps. In contrast, the microfluidic device based method and system uses a pair of microfluidic devices each having a set of parallel channels, where the first and second devices have their channels oriented perpendicular to one another as shown in Figure 9A.

[0177] A microfluidic addressing device is shown with overlaid layout for a pair of addressing devices in Figure 9A, a poly(dimethylsiloxane) (PDMS) elastomer device with 16×16 channels and $100 \mu\text{m}$ channel width in Figure 9B, and an assembled device with the clamp and peristaltic pump mechanism in Figure 9C.

[0178] The geometry of the devices defines a rectangular array of junctions, each having an area that is defined by the width of the two channels. If each channel receives a different address tag, the result is a unique pair of identifying address tags for each junction or intersection in the array. Fluid flow in microfluidic devices is usually driven by external syringe pumps or vacuum and often requires a complex plumbing setup including connections between the microscopic channels and the macroscopic components of the system. The reagent delivery system used in the example is a self-contained system for loading reagents into the channels. The device is cast out of a PDMS elastomer and includes reagent reservoirs, and microscopic addressing channels, each of which is connected to a larger peristaltic pump channel. The device is applied to the surface of an FFPE sample and clamped in place. A thumb-wheel is applied across all the pump channels and the rolling action draws the liquid from each reservoir through the addressing channel where it contacts the tissue sample and the address tag is ligated onto the hybridized probes. After the first ligation, the device is removed and the sample washed. The second, perpendicular device is used to install the second set of address tags. Only the probes under the area at the intersection of two channels receive both address tags. The devices can be cleaned and reused.

[0179] A set of microfluidic devices with a 5-site $\times$ 5-site layout was fabricated, to match a set of custom-designed tissue microarrays (TMAs) that contained a corresponding 5×5 checkerboard pattern was produced. The TMAs contained the same commercially sourced FFPE sections of normal human liver and pancreas (Pantomics) used above in this example, arranged in a checkerboard pattern. This known pattern of tissue spots on the array was used to verify the accuracy of the spatial encoding system. Figure 10 shows an immunofluorescence image of a TMA as well as expression maps generated by the assay system using the microfluidic reagent delivery system. Figure 10A shows immunofluorescence (IF) image of a custom TMA stained with two liver specific antibodies: PYGL, specific to hepatocytes and Annexin A2, specific to bile-duct cells. The reference was Protein Atlas, available at www.proteinatlas.org. The brightly stained spots are liver tissue and dim spots are pancreas. Figure 10B shows a map of the sum of the 22 most abundant liver-specific genes by

abundance, normalized to housekeeping genes (GAPDH and ActB). Each square corresponds to the signal mapped to one junction, a 500 μm $\times$ 500 μm area centered on one of the tissue cores of the TMA. Figure 10C shows a map of the sum of the 22 most abundant pancreas genes by abundance, normalized by housekeeping genes. The addressing channels of the microfluidic devices used are 500 μm wide at a 2 mm pitch with a depth of 50 μm , which corresponds to a "virtual-volume" of 12.5 nL that encompasses the intersection of the perpendicular channels.

[0180] These results demonstrate that mapped sequencing data using the multiplex system reproduced the expected expression pattern of the tissue sample, and that the multiplex assay is compatible with immunofluorescence imaging, allowing the determination of cell types based on protein markers and correlation with gene expression data.

A 134-plex assay using formalin-fixed, paraffin-embedded (FFPE) samples

[0181] A probe pool and two device layouts were developed. The probe pool consisted of 134 targets representing 69 unique genes shown in Table 1. When reading out expression by sequencing, a few highly expressed genes can account for the majority of the reads, limiting dynamic range of the assay. This issue was mitigated by attenuating some of the most highly expressed genes in the pool. This was accomplished by adding in attenuator probes in known ratios with the active probes. An attenuator probe lacks a 5' phosphate necessary for ligation, preventing production of an amplifiable product and thus decreasing the signal from that target. Table 2 shows the results of attenuation of the top 5 genes. Before attenuation they accounted for 73% of the reads whereas afterwards they accounted for less than 18%. This strategy can be used to achieve very high levels of multiplexing with current sequencing technology while still achieving high dynamic range.

Table 1: List of genes and number of unique targets per gene in 134-plex probe pool.

Pluripotency		Liver		Liver		Pancreas		Pancreas	
AURKB	3	AGXT	2	KRT19	2	AQP8	2	DPEP1	2
HMGB3	2	ALDO	2	KRT7	2	CARS	2	GP2	2
JARID2	3	APOB	2	MCAM	2	CEL	2	PRSS1	2
LIN28A	1	BHMT	2	MYH9	2	CLPS	2	SOX9	2
SOX2	1	CPB2	2	POGZ	2	CPA1	2	WDR38	2
Housekeeping		CYP2A6	2	SMARCA4	2	GCG	2	ASB9	2
ACTB	2	CYP2C8	2	ALB	2	INS	1	CHGA	2
GAPDH	2	HPX	2	ARG1	2	PNLIP	2	GAD2	2
H2AFX	2	SAA4	2	CD14	2	PNLIPRP2	2	INSM1	2
Controls		SERPIND1	2	MBL2	2	PPP4C	2	NCAM1	2
18S	3	VTN	2	PYGL	2	REG1B	2	PAX6	2
Rand Neg	3	CA9	2	SLC27A5	2	SEL1L	2	PPY	2
Other		EPB41L2	2	STOM	2	CA12	2	SV2A	2
FXR1	2	HNF1B	2			CPA2	2	UCHL1	2

Table 2: Attenuation of top 5 assay targets

Probe Name	Fraction of Reads		<u>Atten.</u> Factor
	w/o Atten.	w/ Atten.	
PNLIP_2	0.253	0.048	5.268
PNLIP_1	0.203	0.035	5.871
PRSS1_2	0.114	0.034	3.343
CPA1_2	0.111	0.023	4.841
CLPS_2	0.051	0.037	1.373
Sum	0.732	0.177	

Example 5 Elution and preparation of spatially encoded probes for next generation sequencing

[0182] Using the methods described *supra*, a 134-plex pool of probe pairs was hybridized to an FFPE sample, ligated and spatially encoded with X-positional and Y-positional adaptors. In preparation for elution, a hybridization chamber (Agilent) was applied to the slide and clamped in place to form a leakproof chamber containing the FFPE tissue sample. Using syringes, this chamber was filled with deionized water and the assembly was heated to 80°C for 30 minutes after which time the eluate was removed using a syringe and transferred to a tube.

[0183] The spatially encoded constructs were purified by two rounds of positive selection using magnetic beads to isolate them from any un-encoded probes, leftover positional-encoding adaptors, or malformed constructs. In the first round of purification, the eluate was hybridized to a biotinylated capture probe comprising a sequence that was complementary to a sequence spanning the junction of the X positional adaptor (address tag) and the 5' end of the joined probe pair. This capture probe was then captured on streptavidin functionalized magnetic beads, which were then washed extensively to remove unbound material. Constructs hybridized to capture probes were then eluted by heating in an elution buffer containing a blocking oligonucleotide that was complementary to the capture probe. The eluate was separated from the magnetic beads using a magnet, transferred to a new container, and hybridized with a biotinylated capture probe comprising a sequence that was complementary to a sequence spanning the junction of the 3' end of the joined probe pair and the Y positional adaptor (address tag). This capture probe, together with hybridized constructs, was subsequently captured on streptavidin functionalized magnetic beads and washed.

[0184] The beads were transferred directly into a PCR mix that included primers comprising sequences that enable sequencing of the PCR products on an Illumina MiSeq instrument. The primers also comprised TruSeq barcodes to allow demultiplexing of multiple samples in a single sequencing run. A fraction of the PCR product was analyzed by gel electrophoresis to verify the presence of the amplified spatially encoded constructs. The remaining product was purified using a Qiagen PCR purification Kit. Finally, the spatially encoded constructs were

purified by size selection using either a conventional gel electrophoresis device or the Pippen Prep System (Sage Science).

[0185] The purified encoded construct was sequenced using an Illumina MiSeq instrument and the data were used to generate expression maps.

Example 6 Spatially encoded protein *in situ* assays

[0186] This example describes a spatially encoded protein *in situ* assay. A highly multiplexable protein detection assay was carried out on a tissue microarray like the ones described *supra* containing a checkerboard pattern of liver and pancreas tissue cores. In this case a two-plex assay was encoded using a 5-site × 5-site addressing scheme. The assay was performed by first applying a typical immunostaining procedure with two different primary antibodies, one specific to exocrine cells in the pancreas and one specific to hepatocytes in the liver. Two antibody-DNA conjugates were used as secondary antibodies and were applied to the entire tissue microarray. The conjugates included an oligonucleotide comprising an identity tag as well as an upstream and downstream splint region to allow ligation of X and Y address tags. After applying the primary antibody and secondary antibody conjugate to the entire sample and washing sufficiently, a pair of microfluidic channel devices was used to deliver sequentially the X and Y address tags, which were ligated to the oligonucleotide on the conjugate. The conjugates were eluted from the sample and the combined X and Y tags plus the intervening identity tag formed an amplifiable construct which was amplified, purified and subjected to next generation sequencing to identify the abundance of the antibody targets at each spatially encoded location.

Example 7 Spatially encoded protein *in situ* assays

[0187] This example describes a spatially encoded protein *in situ* assay. As shown in Figure 6A, a highly multiplexable protein detection assay can be carried out on a sample that preserves the spatial organization of cells in a tissue, e.g., a paraffin-embedded or fresh-frozen tissue section fixed to a glass slide. Assay reagents are protein binders (e.g. antibodies) that are identified via linked DNA tags that can be further encoded with tag sequences that encode positional or address information (in this example, indicated as "X" dimension and "Y" dimension). The address tags X and Y are flanked by universal sequences (UP1 and UP2) that can be used as PCR priming sites, adaptors for next-generation sequencing, or both.

[0188] As shown in Figure 6B, the binders, for example, the DNA-labeled antibody probes in this example, are delivered over the entire sample surface in a bulk process. The X and Y address tags are then delivered to the sample and coupled to the probes, so that the probes are encoded by the X and Y address tags in a spatially defined pattern. In this example, two sets of tags (*i.e.*, a set of 10 X address tags, namely X1, X2, X3, ..., X10, and a set of 10 Y

address tags, namely Y1, Y2, Y3, ..., Y10) are used in a combinatorial fashion, and 100 sites in the sample can be uniquely identified by the combinations of X and Y address tags. For example, a site in the sample shown in Figure XB is uniquely identified as (X9, Y1).

[0189] Once the *in situ* assay is completed, the assay products are eluted and sequenced. The address tag sequence information identifies the site at which the assay is performed, and the probe sequence information identifies the protein that is targeted. The frequency of a particular assay product in the digital readout can be used to infer the relative abundance of its target sequence in the sample. This information can then be associated with other information, including conventional histological information, and/or transcript abundance obtained via the related spatially encoded genomic assay.

Example 8 A method to reduce background and increase signal-to-noise ratio

[0190] This example describes a method of detecting rare variant sequences in a mixed population of nucleic acids. The method can be integrated into the methods and assay systems disclosed herein, for example, to reduce the background contributed by random errors and thus to increase the signal to noise ratio (S/N).

[0191] Parallel clonal amplification methods in combination with digital sequencing have permitted large-scale analysis of variation at resolutions in the range of 1% (Druley et al., 2009, Nat. Methods 6: 263-65), but not much below. Although next-generation sequencing enables *de novo* discovery and holds great promise for deep analysis of variation across the genome, a combination of factors at various steps in the sequencing process have made it difficult to obtain very low error rates at readout. These factors include cross-talk between detection channels, incomplete reactions leading to signal loss, increased background as a result of loss in synchronicity of nucleotide addition, and noise and errors in image processing and signal extraction, which worsen significantly at higher sequencing densities. Thus the sequencing readout error rate is far above intrinsic rates exhibited by the high fidelity polymerases used in sequencing reactions. For example, an error rate of 4.4×10^{-7} is estimated for PfuTM polymerase (New England Biolabs, Ipswich, MA). The method described in this example addresses the above technical issues, by using tags to identify target sequences that are "identical by descent." As illustrated in Figure 11A, sequence reads can be partitioned into related groups on this basis.

[0192] Figure 11A shows the concept of the rare variant assay, and Figure 11B provides exemplary configurations of probes that can be used to integrate the rare variant assay in the spatially encoded assays of the present disclosure. The top panel of Figure 11A shows a target sequence of interest flanked by adaptors that contain Illumina adaptor sequences for surface PCR (labeled **a** and **b**). The target can be obtained from a variety of sources, for example, a PCR amplicon. The adaptors contain a variable tag region (labeled **z**). Both strands are shown to illustrate that the Illumina adaptors are asymmetric. The tagged adaptors are used to

construct libraries for sequencing. Single molecules are amplified to form "clusters" on the surface of a flowcell. Sequences are determined for each target region and its associated tag regions. In the final step shown, reads are grouped according to their tag regions, based on the assumption that reads with the same tag sequences are identical by descent, given that *z* is sufficiently long. The groupings are then analyzed to identify rare variant sequences (e.g. targets in the last set numbered 4 are shown in darker color compared to those in sets 1-3 to indicate that the target sequence differs from those in sets 1-3). Similar methods for rare variant sequence detection have been described in Fu et al., 2011, Proc. Natl. Acad. Sci., 108: 9026-9031, and in Schmitt et al., 2012, Proc. Natl. Acad. Sci., 109: 14508-1451.

[0193] With this strategy, the contribution of random sequencing errors can be virtually eliminated. Therefore, barring contamination, the ability to detect a rare variant will be limited in theory by the sample size. Note that although the design shown in Figure 11A references the Illumina adaptors and surface amplification methodology, the method is general and can be used with other sequencing platforms such as the SOLiD platform (Life Technologies), the 454 platform (Roche), and the Pacific Biosciences and Ion Torrent library constructions methods.

[0194] A model system was established to quantitate the improvements in the limit of detection over standard sequencing with the Illumina GAIIx instrument. The model system consisted of a wild-type 100-mer oligo and a mutant sequence containing a unique, single point mutation in the wild-type sequence. Synthetic oligos were cloned into an *E. coli* plasmid vector and individual clones were picked and sequence verified in order to obtain constructs that contained the desired sequences, providing pure, well defined sequence constructs free from oligonucleotide chemical synthesis errors (typically in the range of 0.3-1%). The 100-mer of interest was then excised from the plasmid clone by restriction digestion. Mutant and wild-type oligos were quantitated and mixed at ratios of 1:0, 1:20, 1:1000, 1:10,000, 1:100,000 and 1:1,000,000, and 0:1 to simulate the presence of a rare variant in a wild-type DNA background.

[0195] Next, custom adaptors containing random 10-mer tags were designed and synthesized. Libraries were prepared from the defined oligo mixtures, and sequenced on an Illumina GAIIx instrument according to the constructs and steps outlined in Figure 11A. The data were first analyzed without utilizing the tag information (tag *z* as shown in Figure 11A). This resulted in detection of the point mutation only in the 1:20 sample. A second round of analysis utilizing the tags was done using only high quality reads in which tag1/tag2 pairs were retained if the tags were grouped with each other >99% of the time and had ≥ 2 replicates. In order for a tag group to be scored as a mutation, at least 90% of reads in the group had to agree.

[0196] The mutant allele was also successfully detected in the 1:10,000, 1:100,000, and 1:1,000,000 samples as shown in Table 3. Mutant allele frequencies within a factor of 2 of the expected value were observed, and this difference was accounted for in dilution and pipetting error. The power to observe a mutation in the wild-type (negative control) sample with ~ 7.5 M tag groups is greater than 0.999. Therefore, the difference between the 1:1,000,000 spiked sample and the negative control was highly significant.

Table 3: Demonstration of ability to detect a mutant allele over ~ 6 orders of magnitude

Mutant: WT	Number of Tag Groups Assayed	Number of Mutant Alleles Observed	Estimated Allele Frequency
1:20	3,433	273	0.08
1:1,000	2,539	6	0.0024
1:10,000	157,431	26	1.65E-04
1:100,000	1,752,922	33	1.88E-05
1:1,000,000	4,186,545	5	1.19E-06
(Negative Ctrl) 1:0	7,488,853	0	0

[0197] The power to observe a mutant with frequency f is $1-(1-f)^{\#tags}$, so additional sequencing depth can increase the detection power. The limit of detection in this model system is determined only by sample size and any background contamination that might be present.

[0198] This method can be used to distinguish *in vitro* amplification errors from rare variants present in the original sample. For example, a simple threshold that the mutation frequency within a tag group must be >0.9 can be used to exclude PCR amplification errors from the analysis. This is based on the observation that the expected fraction of copies containing an error at that particular location equals 0.5, conditional on the error occurring in the very first cycle and neglecting the chance of consecutive PCR errors at the same position. No tags in the negative control pass this criterion.

[0199] This method can be integrated into the methods and assay systems for determining a spatial pattern of a target abundance, expression, or activity, in order to reduce the background contributed by random errors and thus to increase the signal to noise ratio (S/N). Non-limiting exemplary configurations of probes that integrate the X and Y address tags and the variable tag region z are shown in Figure 11B.

Example 9 Analysis of brain tissue

[0200] This example describes production of an at least 24-plex protein assay panel and confirmation of its tissue/cell-type specificity by correlation with fluorescent labeling and by analysis of tissue microarrays.

[0201] A set of 26 antigens is selected. These antigens are expressed in neurons, astrocytes, oligodendrocytes, microglia or proliferating cells, and antibodies that have been raised against the antigens are commercially available (Table 4). These antibodies have been successfully used, in conjunction with well-established staining techniques, to mark different cell types and regions within brain sections (Lyck, et al., 2008, J Histochem Cytochem 56, 201-21). For the purpose of the assay, it is necessary to avoid procedures for antibody binding that damage

RNA.

[0202] Antigen accessibility is addressed by exploring systematically a range of "antigen retrieval" protocols and testing their compatibility with RNA. See, MacIntyre, 2001, Br J Biomed Sci. 58,190-6; Kap et al., 2011, PLoS One 6, e27704; Inoue and Wittbrodt, 2011, PLoS One 6, e19713. A panel of antibody assays rather than any individual assay are explored to identify a suitable subset for use in a multiplexed panel.

[0203] The assay system is also validated by using spatially-encoded and conventional IHC fluorescence data and spatially encoded RNA data, applied to brain tissue. High-dimensional protein and mRNA data from 32×32 sites in sections of human brain tissue are generated and compared with published data and brain atlas data.

[0204] The Allen Brain Atlas (www.brain-map.org) can be used to select target genes for production of a panel of gene expression assays with high information content, using the methods and assay systems of the present disclosure. The "Differential Search" tool is used to interrogate the rich spatial expression dataset (generated by *in situ* hybridization), it is identified that ~200 genes are present at a range of abundances in at least one structure/compartments of the brain, and/or are strongly differentially expressed between the different structures/compartments. The selection is reviewed to incorporate any new information or criteria. Probes against the set of ~200 mRNAs are designed and tested for their performance in the multiplexed assay, using the online gene expression data as a reference.

[0205] Protein panels and RNA assay panels are applied simultaneously to analyze sections of normal human brain. For example, the abundance of at least 24 proteins and 192 mRNA analytes over a 32×32 grid of 50 μm pixels from sections of healthy human brain is analyzed. The results are used to generate a rich map of the brain's spatially-organized molecular terrain, and are amenable to analysis in various ways, including those that reveal:

1. 1. The organization of brain into distinct sub-structures: both at the anatomical scale, and at the lower-level multicellular level;
2. 2. Spatial variation in the representation of different cell types across the tissue (e.g. using sets of proteins/mRNAs known to be specific to particular cell types); and
3. 3. The relation between mRNA and protein expression from the same gene at different tissue locations.

Table 4: Candidate proteins to differentiate brain tissues, which have been used in immunohistochemistry and have commercially available antibodies.

Protein	Observed Specificity {Lyck, 2008}
b-tubulin III	Neuropil and neuronal bodies
CD11b	None
CD14	Perivascular macrophages
CD34	Endothelium and white blood cells

Protein	Observed Specificity {Lyck, 2008}
CD39	Endothelium, astroglia, and macrophages
CD45	Microglia, macrophages, and lymphocytes
CD68	Microglia and macrophages
CD169	Endothelial and perivascular macrophages
CNPase	Myelinated fibers and round cell bodies
GFAP	Astroglia in white matter and neocortex
HLA-DR	Microglia/macrophages and lymphocytes
Ki-67	Perivascular space and sub-ventricular zone
MAP-2	Neurons and proximal part of apical dendrites
MBP	Myelinated fibers
Nestin	Endothelial cells/vessel wall
NeuN	Neuronal cell bodies
Neurofilament	Neuronal cell processes
NG2	None
Nkx-2.2	None
NSE	Neuropil
O4 sulfatide	Myelinated fibers
PDGFa-R	None
p25a	Neuropil and round cell bodies
S100b	Astroglia in white matter and neocortex
TOAD-64	Neuropil
Vimentin	Astroglia and endothelial cells/vessel wall

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not

form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US2011245111A](#) [0005]
- [US6969589B](#) [0066]
- [US20060188875A](#) [0066]
- [US080616](#) [0075]
- [US20110245111A](#) [0075]
- [US20080220434A](#) [0084]
- [US20090305237A](#) [0084]
- [US6864052B](#) [0085]
- [US6309824B](#) [0085]
- [US6401267B](#) [0085]
- [US20050191656A](#) [0085]
- [US6210891B](#) [0085]
- [US6828100B](#) [0085]
- [US6969488B](#) [0085]
- [US6897023B](#) [0085]
- [US6833246B](#) [0085]
- [US6911345B](#) [0085]
- [US6787308B](#) [0085]
- [US7297518B](#) [0085]
- [US7462449B](#) [0085]
- [US7501245B](#) [0085]
- [US20110059436A](#) [0085]
- [US20040106110A](#) [0085]
- [US20030064398A](#) [0085]
- [US20030022207A](#) [0085]
- [US5912148A](#) [0085]
- [US6130073A](#) [0085]
- [US20100105052A](#) [0085]
- [US20070207482A](#) [0085]
- [US20090018024A](#) [0085]
- [US20070036511A](#) [0085]
- [US20080032301A](#) [0085]
- [US20080128627A](#) [0085]
- [US20090082212A](#) [0085]
- [US20110033854A](#) [0085]
- [US20090264299A](#) [0085]

- [US20090155781A](#) [0085]
- [US20090005252A](#) [0085]
- [US61902105](#) [0130]
- [US20070166725A](#) [0145]
- [US20050239192A](#) [0145]
- [US20090197326A](#) [0151]

Non-patent literature cited in the description

- **SHI et al.** Nature Biotechnology, 2006, vol. 24, 91151-61 [0003]
- **SLONIMYANAI** Plos Computational Biology, 2009, vol. 5, 10e1000543- [0003]
- **VELCULESCU et al.** Science, 1995, vol. 270, 5235484-87 [0003]
- **SPURGEON et al.** Plos ONE, 2008, vol. 3, 2e1662- [0003]
- **NUOVO** Genome Res., 1995, vol. 4, 151-67 [0003]
- **MORTAZAVI et al.** Nature Methods, 2008, vol. 5, 7621-8 [0003]
- **ARMANI et al.**, Lab on a Chip, 2009, vol. 9, 243526-34 [0004]
- **Genome Analysis: A Laboratory Manual Series** 19990000 vol. I-IV, [0040]
- **Genetic Variation: A Laboratory Manual** 20070000 [0040]
- **PCR Primer: A Laboratory Manual** 20030000 [0040]
- **DNA Microarrays: A Molecular Cloning Manual** 20030000 [0040]
- **MOUNT** Bioinformatics: Sequence and Genome Anazvsis, 2004, [0040]
- **SAMBROOKRUSSELL** Condensed Protocols from Molecular Cloning: A Laboratory Manual, 2006, [0040]
- **SAMBROOKRUSSELL** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press 20020000 [0040]
- **Current Protocols in Molecular Biology** 19870000 [0040]
- **Essential Molecular Biology** IRL Press 19910000 [0040]
- **Gene Expression Technology** Academic Press 19910000 [0040]
- **Methods for Cloning and Analysis of Eukaryotic Genes** Bartlett Publ. 19900000 [0040]
- **M. KRIEGLER** Gene Transfer and Expression Stockton Press 19900000 [0040]
- **Recombinant DNA Methodology** Academic Press 19890000 [0040]
- **M. MCPHERSON et al.** PCR: A Practical Approach IRL Press at Oxford University Press 19910000 [0040]
- **STRYER** Biochemistry W. H. Freeman 19950000 [0040]
- **GAIT** Oligonucleotide Synthesis: A Practical Approach IRL Press 20020000 [0040]
- **NELSONCOX** Lehninger, Principles of Biochemistry W. H. Freeman Pub. 20000000 [0040]
- **BERG et al.** Biochemistry W. H. Freeman Pub. 20020000 [0040]
- **Handbook of Experimental Immunology** Wiley 19960000 [0040]
- **A. ABBAS et al.**, Cellular and Molecular Immunology W.B. Saunders Co. 19910000 [0040]

- Current Protocols in Immunology 19910000 [0040]
- **ANDERSON YOUNG** Quantitative Filter Hybridization, in Nucleic Acid Hybridization, 1985, [0060]
- **ALLAWIS ANTA LUCIA, JR.** Biochemistry, 1997, vol. 36, 10581-94 [0060]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Press 19890000 [0061]
- **AUSUBEL et al.** Short Protocols in Molecular Biology John Wiley & Sons 19990000 [0061]
- **M. KANEHISA** Nucleic Acids Res., 1984, vol. 12, 203- [0062]
- **RONAGHI et al.** Science, 1998, vol. 281, 363-365 [0085]
- **LI et al.** Proc. Natl. Acad. Sci., 2003, vol. 100, 414-419 [0085]
- **SONI MELLER** Clin Chem, 2007, vol. 53, 1996-2001 [0085]
- **MCKERNAN et al.**, Genome Res., 2009, vol. 19, 1527-41 [0085]
- **BENTLEY et al.** Nature, 2008, vol. 456, 53-59 [0085]
- **GELDER et al.** Proc. Natl. Acad. Sci. USA, 1990, vol. 87, 1663-1667 [0090]
- **EBERWINE et al.** Proc. Natl. Acad. Sci. USA, 1992, vol. 89, 3010-3014 [0090]
- **WANG et al.** Nature Biotechnol., 2000, vol. 18, 457-459 [0090]
- **BAUGH et al.** Nucleic Acids Res., 2001, vol. 29, E29- [0090]
- **FIREXU** Proc. Natl. Acad. Sci., 1995, vol. 92, 4641-4645 [0091]
- **DAUBENDIEK et al.** J. Am. Chem. Soc., 1995, vol. 117, 77818-7819 [0091]
- **DAHL et al.** Proc. Natl. Acad. Sci., 2004, vol. 101, 4548-4553 [0091]
- **FAN et al.** Cold Spring Symp. Quant. Biol., 2003, vol. 68, 69-78 [0101]
- **FREDRIKSSON et al.** Nat Biotechnol, 2002, vol. 20, 473-7 [0125]
- **BIRNEY et al.** Nature, 2007, vol. 447, 799-816 [0158]
- **DRULEY et al.** Nat. Methods, 2009, vol. 6, 263-65 [0191]
- **FU et al.** Proc. Natl. Acad. Sci., 2011, vol. 108, 9026-9031 [0192]
- **SCHMITT et al.** Proc. Natl. Acad. Sci., 2012, vol. 109, 14508-1451 [0192]
- **LYCK et al.** J Histochem Cytochem, 2008, vol. 56, 201-21 [0201]
- **MACINTYRE** Br J Biomed Sci., 2001, vol. 58, 190-6 [0202]
- **KAP et al.** PLoS One, 2011, vol. 6, e27704- [0202]
- **INOUE WITT BRODT** PLoS One, 2011, vol. 6, e19713- [0202]

METODE TIL BESTEMMELSE AF SPATIALE MØNSTRE I BIOLOGISKE TARGETS I EN PRØVE

PATENTKRAV

1. En metode til bestemmelse af et spatialt mønster i
5 rigelige mængder og/eller udtryk i et eller flere protein targets over flere steder i en 2D prøve, omfattende:
 - levering af en probe (720) af en eller flere protein targets (714) flere steder i prøven fastgjort til en bærer, hvori hver probe omfatter en target bindende enhed (716) som kan binde
10 til probens tilsvarende protein target og et oligonukleotid mærke (718) med en sekvens som identificerer probens tilsvarende protein target eller target bindende enhed;
 - tillade at hver probe binder til dens tilsvarende protein target i prøven;
 - 15 levering af mindst en adapter til flere steder i prøven fastgjort til bæreren, hvori mindst den ene adaptor specielt bindes til proben, hvori adaptoren omfatter et oligonukleotid identificerings mærke (722) som hybridiserer med oligonukleotid mærket;
 - 20 levering af et oligonukleotid adresse mærke (724, 726) til hvert af flere steder i prøven fastgjort til bæreren, hvori en eller flere mikrofluididiske anordninger med adresse kanaler er anvendt til at udføre levering af oligonukleotid adresse mærket (724, 726);
 - 25 koble oligonukleotid adresse mærket til mindst den ene adaptor bundet til proben bundet til protein target, derved genererende et adaptor/oligonukleotid adresse mærke konjugation, hvori oligonukleotid adresse mærket sekvensen identificerer stedet hvortil oligonukleotid adresse mærket leveres til;
 - 30 eluere adaptor/oligonukleotid adresse mærkets konjugation bundet til proben;
 - analysere det eluerede adaptor/oligonukleotid adresse mærke konjugeret af sekvensering,
 - hvori mængden af et sekvenseret produkt indikerer
35 rigelige mængder og/eller udtryk af hver af det ene eller flere protein targets, det sekvenserede produkt omfatter alle eller en del af oligonukleotid adresse mærke sekvensen og alle eller en del af oligonukleotid identitets mærke sekvensen; og
 - bestemmelse af et spatialt mønster af rigelig mængde
40 og/eller udtryk i det ene eller flere protein targets over flere steder i prøven baseret på analysen.

2. Metoden i henhold til patentkrav 1, hvori mindst to adaptorer er leveret til flere steder i prøven, hvori mindst to adaptorer hver specielt bindes til en probe som specielt binder til protein target.

5 3. Metoden i henhold til patentkrav 2, hvori mindst to adaptorer er forbundne med hinanden, for eksempel gennem ligering ved at bruge en del af oligonukleotid mærket som en skinne.

10 4. Metoden i henhold til patentkravene 1-3, hvori proben for hver af den ene eller flere protein targets omfatter et affinitets middel til protein target og en oligonukleotid, og adaptoren omfatter en oligonukleotid.

15 5. Metoden i henhold til et af patentkravene 1-4, hvori den target bindende del af proben indeholder et antistof eller et antigen som binder fragmenter deraf, en aptamer, et lille molekyle, et enzym substrat, et formodet enzym substrat, et affinitets indfangende middel, eller en kombination heraf.

6. En metode til bestemmelse af et spatialt mønster af rigelig mængde og/eller udtryk af en eller flere protein targets over flere steder i en 2D prøve, omfattende:

20 levering af mindst to prober (620, 622) for hver af en eller flere protein targets (602) til flere steder i prøven bundet til bæreren, hvori hver probe indeholder: (1) en target bindende del (606, 608) som kan binde til probens tilsvarende protein target; og (2) et oligonukleotid identitets mærke (604, 610) med
25 en sekvens som identificerer probens tilsvarende protein target eller target bindende del, hvori den target bindende del af hver probe er konjugeret til en polynukleotid som kan specielt hybridisere til en polynukleotid omfattende oligonukleotidens identitets mærke;

30 tillade hver probe at binde til dets tilsvarende protein target i prøven;

levering af et oligonukleotid adresse mærke (624, 628) til hver af flere steder i prøven fastgjort til mærket, hvori en eller flere mikrofluidiske anordninger med adresserings kanaler er
35 anvendt til at udføre levering af oligonukleotid adresse mærket (624, 628);

kobling af oligonukleotid identitets mærket af mindst to prober bundet til protein target og oligonukleotid adresse mærket, herved genererende en probe/oligonukleotid adresse mærke
40 konjugering, hvori oligonukleotid adresse mærke sekvensen

identificerer stedet hvor oligonukleotid adresse mærket er leveret; eluere probe/oligonukleotid adresse mærke konjugeringen bundet til et eller flere protein targets;

- 5 analysere den eluerede probe/oligonukleotid adresse mærke konjugering ved sekvensering, hvori mængden af det sekvenserede produkt indikerer rigelig mængde og/eller udtryk af hver af et eller flere protein targets, det sekvenserede produkt indeholder alle eller en del af oligonukleotid adresse mærke sekvensen og alle eller en del af oligonukleotid identitets mærke sekvensen; og
- 10 bestemme et spatialt mønster af rigelig mængde og/eller udtryk af et eller flere protein targets over flere steder i prøven baseret på analysen.

7. Metoden i henhold til patentkrav 6, omfatter endvidere bestemmelse af spatialt mønster i rigelig mængde og/eller udtryk
- 15 af en polynukleotid over flere steder i en prøve.

8. Metoden i henhold til patentkrav 7, hvori target polynukleotiden eller komplementet deraf koder alle eller en del af target proteinet.

9. Metoden i henhold til patentkrav 8, hvori analyse af
- 20 target proteinet og analyse af target polynukleotiden bliver udført parallelt i det samme reaktions forløb.

10. Metoden i henhold til et af patentkravene 1-9, hvori en kendt procentdel af proberne til protein targets er en attenuator probe som begrænser produktion af et amplifikationsbart produkt.

- 25 11. Metoden i henhold til patentkrav 10, hvori attenuator proben mangler et 5'fosfat.

DRAWINGS

Figure 1





Figure 2

Figure 4




Figure 4

D.




Figure 5




Figure 6





Figure 6

D



E




Figure 7






Figure 8**A****B**


Figure 9

Figure 10





Figure 11**A.****B.**