

(19)



(11) Publication number:

SG 188361 A1

(43) Publication date:

30.04.2013

(51) Int. Cl:

A61K 31/133, A61P 3/04, C07D

223/16;

(12)

Patent Application

(21) Application number: 2013015581

(71) Applicant:

ARENA PHARMACEUTICALS, INC.
6154 NANCY RIDGE DRIVE SAN DIEGO,
CALIFORNIA 92121 CA US

(22) Date of filing: 31.08.2011

(72) Inventor:

BLACKBURN, ANTHONY C. 12633
CAMINITO ROSITA SAN DIEGO,
CALIFORNIA 92128 US
SHAN, YUN 11429 HONEY RIDGE SAN
DIEGO, CALIFORNIA 92130 US
SHIFRINA, ANNA 11957 MIRO CIRCLE
SAN DIEGO, CALIFORNIA 92131 US
STIRN, SCOTT 14710 CARMEL RIDGE
ROAD SAN DIEGO, CALIFORNIA 92128
US

(30) Priority: US 61/402,611 01.09.2010

(54) Title:

NON-HYGROSCOPIC SALTS OF 5-HT2C AGONISTS

(57) Abstract:

Salts of the 5-HT2C-receptor agonist (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, and dosage forms comprising them that are useful for, inter alia, weight management.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 March 2012 (08.03.2012)

(10) International Publication Number
WO 2012/030957 A2

(51) International Patent Classification:
C07D 223/16 (2006.01)

(74) Agents: **DUFFIELD, Jonathan J.** et al.; Arena Pharmaceuticals, Inc., 6166 Nancy Ridge Drve, San Diego, California 92121 (US).

(21) International Application Number:

PCT/US2011/049960

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

31 August 2011 (31.08.2011)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/402,611 1 September 2010 (01.09.2010) US

(71) Applicant (for all designated States except US): **ARENA PHARMACEUTICALS, INC.** [US/US]; 6166 Nancy Ridge Drive, San Diego, California 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BLACKBURN, Anthony C.** [US/US]; 12633 Caminito Rosita, San Diego, California 92128 (US). **SHAN, Yun** [US/US]; 11429 Honey Ridge, San Diego, California 92130 (US). **SHIFRINA, Anna** [US/US]; 11957 Miro Circle, San Diego, California 92131 (US). **STIRN, Scott** [US/US]; 14710 Carmel Ridge Road, San Diego, California 92128 (US).

[Continued on next page]

(54) Title: NON-HYGROSCOPIC SALTS OF 5-HT2C AGONISTS

PXRD of Compound 1 Hemi-edisylate Salt, Form I




Figure 1

(57) Abstract: Salts of the 5-HT2C-receptor agonist (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, and dosage forms comprising them that are useful for, inter alia, weight management.

WO 2012/030957 A2



Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

NON-HYGROSCOPIC SALTS OF 5-HT_{2C} AGONISTS

FIELD OF THE INVENTION

The present invention relates to salts of the 5-HT_{2C}-receptor agonist (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine, and dosage forms comprising them that are useful for, *inter alia*, weight management.

BACKGROUND OF THE INVENTION

Obesity is a life-threatening disorder in which there is an increased risk of morbidity and mortality arising from concomitant diseases such as type II diabetes, hypertension, stroke, cancer and gallbladder disease.

Obesity is now a major healthcare issue in the Western World and increasingly in some third world countries. The increase in numbers of obese people is due largely to the increasing preference for high fat content foods but also the decrease in activity in most people's lives.

Currently about 30% of the population of the USA is now considered obese.

Whether someone is classified as overweight or obese is generally determined on the basis of their body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m²). Thus, the units of BMI are kg/m² and it is possible to calculate the BMI range associated with minimum mortality in each decade of life. Overweight is defined as a BMI in the range 25-30 kg/m², and obesity as a BMI greater than 30 kg/m² (see table below).

Classification Of Weight By Body Mass Index (BMI)

BMI	CLASSIFICATION
< 18.5	Underweight
18.5-24.9	Normal
25.0-29.9	Overweight
30.0-34.9	Obesity (Class I)
35.0-39.9	Obesity (Class II)
>40	Extreme Obesity (Class III)

As the BMI increases there is an increased risk of death from a variety of causes that are independent of other risk factors. The most common diseases associated with obesity are cardiovascular disease (particularly hypertension), diabetes (obesity aggravates the development of diabetes), gall bladder disease (particularly cancer) and diseases of reproduction. The strength of the link between obesity and specific conditions varies. One of the strongest is the link with type 2 diabetes. Excess body fat underlies 64% of cases of diabetes in men and 77% of cases in women (Seidell, *Semin Vasc Med*, 5:3-14 (2005)). Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk of developing coronary heart disease.

There are problems however with the BMI definition in that it does not take into account the proportion of body mass that is muscle in relation to fat (adipose tissue). To account

for this, obesity can also be defined on the basis of body fat content: greater than 25% in males and greater than 30% in females.

Obesity considerably increases the risk of developing cardiovascular diseases as well. Coronary insufficiency, atheromatous disease, and cardiac insufficiency are at the forefront of 5 the cardiovascular complications induced by obesity. It is estimated that if the entire population had an ideal weight, the risk of coronary insufficiency would decrease by 25% and the risk of cardiac insufficiency and of cerebral vascular accidents would decrease by 35%. The incidence of coronary diseases is doubled in subjects less than 50 years of age who are 30% overweight. The diabetes patient faces a 30% reduced lifespan. After age 45, people with diabetes are about 10 three times more likely than people without diabetes to have significant heart disease and up to five times more likely to have a stroke. These findings emphasize the inter-relations between risks factors for diabetes and coronary heart disease and the potential value of an integrated approach to the prevention of these conditions based on the prevention of obesity (Perry, I. J., *et al.*, *BMJ* 310, 560-564 (1995)).

15 Diabetes has also been implicated in the development of kidney disease, eye diseases and nervous system problems. Kidney disease, also called nephropathy, occurs when the kidney's "filter mechanism" is damaged and protein leaks into urine in excessive amounts and eventually the kidney fails. Diabetes is also a leading cause of damage to the retina at the back of the eye and increases risk of cataracts and glaucoma. Finally, diabetes is associated with 20 nerve damage, especially in the legs and feet, which interferes with the ability to sense pain and contributes to serious infections. Taken together, diabetes complications are one of the nation's leading causes of death.

25 The first line of treatment is to offer diet and life style advice to patients such as reducing the fat content of their diet and increasing their physical activity. However, many patients find this difficult and need additional help from drug therapy to maintain results from these efforts.

30 Most currently marketed products have been unsuccessful as treatments for obesity because of a lack of efficacy or unacceptable side-effect profiles. The most successful drug so far was the indirectly acting 5-hydroxytryptamine (5-HT) agonist d-fenfluramine (ReduxTM) but reports of cardiac valve defects in up to one third of patients led to its withdrawal by the FDA in 1998.


35 In addition, two drugs have been launched in the USA and Europe: Orlistat (XenicalTM), a drug that prevents absorption of fat by the inhibition of pancreatic lipase, and Sibutramine (ReductilTM), a 5-HT/noradrenaline re-uptake inhibitor. However, side effects associated with these products may limit their long-term utility. Treatment with XenicalTM is reported to induce gastrointestinal distress in some patients, while Sibutramine has been associated with raised blood pressure in some patients.

Serotonin (5-HT) neurotransmission plays an important role in numerous physiological processes both in physical and in psychiatric disorders. 5-HT has been implicated in the regulation of feeding behavior. 5-HT is believed to work by inducing a feeling of satiety, such that a subject with enhanced 5-HT stops eating earlier and fewer calories are consumed. It has been shown that a stimulatory action of 5-HT on the 5-HT_{2C} receptor plays an important role in the control of eating and in the anti-obesity effect of d-fenfluramine. As the 5-HT_{2C} receptor is expressed in high density in the brain (notably in the limbic structures, extrapyramidal pathways, thalamus and hypothalamus i.e. PVN and DMH, and predominantly in the choroid plexus) and is expressed in low density or is absent in peripheral tissues, a selective 5-HT_{2C} receptor agonist can be a more effective and safe anti-obesity agent. Also, 5-HT_{2C} knockout mice are overweight with cognitive impairment and susceptibility to seizure.

It is believed that the 5-HT_{2C} receptor may play a role in obsessive compulsive disorder, some forms of depression, and epilepsy. Accordingly, agonists can have anti-panic properties, and properties useful for the treatment of sexual dysfunction.

In sum, the 5-HT_{2C} receptor is a receptor target for the treatment of obesity and psychiatric disorders, and it can be seen that there is a need for selective 5-HT_{2C} agonists which safely decrease food intake and body weight.

The salts and formulations of the present invention comprise the selective 5-HT_{2C}-receptor agonist (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (Compound 1), and are useful for, *inter alia*, weight management, including weight loss and the maintenance of weight loss. Compound 1 is disclosed in PCT patent publication WO2003/086303, which is incorporated herein by reference in its entirety.



1

Various synthetic routes to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine, its related salts, enantiomers, crystalline forms, and intermediates, have been reported in PCT publications, WO 2005/019179, WO 2006/069363, WO 2007/120517, WO 2008/070111, WO 2009/111004, and in United States provisional application 61/396,752 each of which is incorporated herein by reference in its entirety.

Combinations of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine with other agents, including without limitation, phentermine, and uses of such combinations in therapy are described in WO 2006/071740, which is incorporated herein by reference in its entirety.

The following United States provisional applications are related to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine: 61/402,578; 61/403,143; 61/402,580; 61/402,628;

61/403,149; 61/402,589; 61/402,611; 61/402,565; 61/403,185; each of which is incorporated herein by reference in its entirety.

The following applications are related to *(R)*-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine and have the same filing date as the subject application: Attorney Reference 5 Number 178.WO1, a PCT application which claims priority to United States provisional applications 61/402,578 and 61/403,143; Attorney Reference Number 181.WO1, a PCT application which claims priority to United States provisional application 61/402,580; Attorney Reference Number 186.WO1, a PCT application which claims priority to United States 10 provisional applications 61/402,628 and 61/403,149; Attorney Reference Number 187.WO1, a PCT application which claims priority to United States provisional application 61/402,589; and Attorney Reference Number 192.WO1, a PCT application which claims priority to United States 15 provisional applications 61/402,565 and 61/403,185; each of which is incorporated herein by reference in its entirety.

(R)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hydrochloride (lorcaserin hydrochloride) is an agonist of the 5-HT_{2C} receptor and shows effectiveness at reducing obesity in animal models and humans. In December 2009, Arena Pharmaceuticals submitted a New 20 Drug Application, or NDA, for lorcaserin to the FDA. The NDA submission is based on an extensive data package from lorcaserin's clinical development program that includes 18 clinical trials totaling 8,576 patients. The pivotal phase 3 clinical trial program evaluated nearly 7,200 patients treated for up to two years, and showed that lorcaserin consistently produced significant weight loss with excellent tolerability. About two-thirds of patients achieved at least 5% weight 25 loss and over one-third achieved at least 10% weight loss. On average, patients lost 17 to 18 pounds or about 8% of their weight. Secondary endpoints, including body composition, lipids, cardiovascular risk factors and glycemic parameters improved compared to placebo. In addition, heart rate and blood pressure went down. Lorcaserin did not increase the risk of cardiac 30 valvulopathy. Lorcaserin improved quality of life, and there was no signal for depression or suicidal ideation. The only adverse event that exceeded the placebo rate by 5% was generally mild or moderate, transient headache. Based on a normal BMI of 25, patients in the first phase 3 trial lost about one-third of their excess body weight. The average weight loss was 35 pounds or 16% of body weight for the top quartile of patients in the second phase 3 trial.

An immediate-release film-coated 10-mg tablet was developed for the phase 3 clinical trials and commercial launch of lorcaserin, but there remains a need for alternative formulations for oral use. These include formulations characterized by their suitable flowing properties, tabletability, and stability to moisture.

35 In view of the growing demand for compounds useful in the treatment of disorders related to the 5-HT_{2C} receptor, *(R)*-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine has emerged has an important new compound. Accordingly, new formulations of *(R)*-8-chloro-1-

methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine, which are non-hygroscopic and show good solid-state stability under humid conditions are needed. The salts and processes described herein help meet these and other needs.

5

SUMMARY OF THE INVENTION

A priori, it is difficult to predict with confidence which salts of a particular drug will be solid, stable, and readily isolable. *A fortiori*, the hygroscopicity of such salts cannot be predicted with accuracy and must instead be determined empirically. In the course of preparing the salts of the present invention, many counterions commonly used in the pharmaceutical industry (see *e.g.* 10 Berge, *et al.*, *Journal of Pharmaceutical Sciences*, 66:1-19 (1977)) were investigated. Acetate, DL-lactate, ascorbate, D-gluconate, besylate, napsylate, tosylate, isethionate, dichloroacetate, benzoate, esylate, gentisate, hippurate, lactobionate, xinafoate, and sebacate salts of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine were prepared, but all of these failed to crystallize. By contrast, the salts of the present invention are salts of (*R*)-8-chloro-1-methyl-15 2,3,4,5-tetrahydro-1*H*-3-benzazepine which when prepared were discovered to be both crystalline, and non-hygroscopic. Because of their stability to moisture these salts are useful, *inter alia*, for preparing dosage forms of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine.

One aspect of the present invention pertains to certain salts of (*R*)-8-chloro-1-methyl-20 2,3,4,5-tetrahydro-1*H*-3-benzazepine (Compound 1) and pharmaceutically acceptable solvates and hydrates thereof.

One aspect of the present invention pertains to certain salts of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (Compound 1).

One aspect of the present invention pertains to salts selected from: (*R*)-8-chloro-1-methyl-25 2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt; (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate salt; (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt; (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate salt; and (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine succinate salt; and (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt; and 30 pharmaceutically acceptable solvates and hydrates thereof.

One aspect of the present invention pertains to pharmaceutical compositions comprising a salt of the present invention and a pharmaceutically acceptable carrier.

One aspect of the present invention pertains to processes for preparing a pharmaceutical composition comprising admixing a salt of the present invention, and a pharmaceutically acceptable carrier. 35

One aspect of the present invention pertains to bulk pharmaceutical compositions suitable for the manufacture of dosage forms for weight management, comprising a salt of the present invention, and a pharmaceutically acceptable carrier.

5 One aspect of the present invention pertains to processes for preparing bulk pharmaceutical compositions suitable for the manufacture of dosage forms for weight management, comprising admixing a salt of the present invention, and a pharmaceutically acceptable carrier.

10 One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of a salt selected from: a pharmaceutically acceptable salt of (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine and pharmaceutically acceptable solvates and hydrates thereof, wherein the dosage form is a non-hygroscopic dosage form.

One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of a salt of the present invention.

15 One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

One aspect of the present invention pertains to the use of salts of the present invention in the manufacture of a medicament for weight management in an individual.

20 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of treatment of the human or animal body by therapy.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: PXRD of Compound 1 Hemi-edisylate Salt, Form I.

Figure 2: DSC and TGA of Compound 1 Hemi-edisylate Salt, Form I.

25 Figure 3: DMS of Compound 1 Hemi-edisylate Salt, Form I.

Figure 4: PXRD of Compound 1 Phosphate Salt, Form I.

Figure 5: DSC and TGA of Compound 1 Phosphate Salt, Form I.

Figure 6: DMS of Compound 1 Phosphate Salt, Form I.

Figure 7: PXRD of Compound 1 Citrate Salt Hemihydrate, Form I.

30 Figure 8: DSC and TGA of Compound 1 Citrate Salt Hemihydrate, Form I.

Figure 9: DMS of Compound 1 Citrate Salt Hemihydrate, Form I.

Figure 10: PXRD of Compound 1 Hemi-oxalate Salt, Form I.

Figure 11: DSC and TGA of Compound 1 Hemi-oxalate Salt, Form I.

Figure 12: DMS of Compound 1 Hemi-oxalate Salt, Form I.

35 Figure 13: PXRD of Compound 1 Succinate Salt, Form I.

Figure 14: DSC and TGA of Compound 1 Succinate Salt, Form I.

Figure 15: DMS of Compound 1 Succinate Salt, Form I.

Figure 16: PXRD of Compound 1 Oxoglutarate Salt, Form I.

Figure 17: DSC and TGA of Compound 1 Oxoglutarate Salt, Form I.

Figure 18: DMS of Compound 1 Oxoglutarate Salt, Form I.

Figure 19: PXRD of Compound 1 Oxoglutarate Salt Solvate, Form I.

5 Figure 20: DSC and TGA of Compound 1 Oxoglutarate Salt Solvate, Form I.

Figure 21: DMS of Compound 1 Oxoglutarate Salt Solvate, Form I.

DETAILED DESCRIPTION

It should be appreciated that certain features of the invention, which are, for clarity, 10 described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

15 DEFINITIONS

For clarity and consistency, the following definitions will be used throughout this patent document.

The term “agonist” refers to a moiety that interacts with and activates a receptor, such as 20 the 5-HT_{2C} serotonin receptor, and initiates a physiological or pharmacological response characteristic of that receptor.

The term “individual” refers to both humans and non-human mammals. Non-human mammals include but are not limited to rodents such as mice and rats, *etc.* rabbits, dogs, cats, swine, cattle, sheep, horses, and non-human primates such as monkeys and apes, *etc.*

The term “pharmaceutical composition” refers to a composition comprising at least one 25 active ingredient; including but not limited to Compound 1 and pharmaceutically acceptable salts, solvates and hydrates thereof, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, without limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

30 The term “therapeutically effective amount” refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician or caregiver or by an individual, which includes one or more of the following:

(1) Preventing the disease, for example, preventing a disease, condition or disorder in an 35 individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease;

(2) Inhibiting the disease, for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, arresting further development of the pathology and/or symptomatology); and

5 (3) Ameliorating the disease, for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, reversing the pathology and/or symptomatology).

The term “treatment” as used herein refers to one or more of the following:

(1) prevention of a disease, for example, prevention of a disease, condition or disorder 10 in an individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease;

(2) inhibition of a disease, for example, inhibition of a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, arresting further development of the pathology and/or 15 symptomatology); and

(3) amelioration of a disease, for example, amelioration of a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, reversing the pathology and/or symptomatology).

Whether an individual is in need of treatment is a judgment made by a caregiver (*e.g.* 20 nurse practitioner, physician, physician assistant, nurse, *etc.* in the case of humans; veterinarian in the case of animals, including non-human mammals) that an individual or animal requires or will benefit from treatment. This judgment is made based on a variety of factors that are in the realm of a caregiver's expertise, but that includes the knowledge that the individual or animal is ill, or will become ill, as the result of a disease, condition or disorder that is treatable by 25 Compound 1 and pharmaceutically acceptable salts, solvates and hydrates thereof. Accordingly, Compound 1 and pharmaceutically acceptable salts, solvates and hydrates thereof can be used in a protective or preventive manner; or Compound 1 and pharmaceutically acceptable salts, solvates and hydrates thereof can be used to alleviate, inhibit or ameliorate a disease, condition or disorder.

30 The term “weight management” as used herein refers to controlling body weight and in the context of the present invention is directed toward weight loss and the maintenance of weight loss (also called weight maintenance herein). In addition to controlling body weight, weight management includes controlling parameters related to body weight, for example, BMI, percent body fat and waist circumference. For example, weight management for an individual 35 who is overweight or obese can mean losing weight with the goal of keeping weight in a healthier range. Also, for example, weight management for an individual who is overweight or

obese can include losing body fat or circumference around the waist with or without the loss of body weight.

The term “maintenance of weight loss” or “weight maintenance” as used herein refers to preventing, reducing or controlling weight gain after weight loss. It is well known that weight gain often occurs after weight loss. Weight loss can occur, for example, from dieting, exercising, illness, drug treatment, surgery or any combination of these methods, but often an individual that has lost weight will regain some or all of the lost weight. Therefore, weight maintenance in an individual who has lost weight can include preventing weight gain after weight loss, reducing the amount of weight gained after weight loss, controlling weight gain after weight loss or 10 slowing the rate of weight gain after weight loss.

SALTS OF THE INVENTION

The present invention is directed, *inter alia*, to solid, stable, and readily isolable salts of (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine and pharmaceutically acceptable solvates and hydrates thereof. The solid state properties of the crystalline forms of salts the present invention are summarized *infra*.

One aspect of the present invention pertains to salts selected from: (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-edisylate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine phosphate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine citrate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-oxalate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine succinate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine oxoglutarate salt; and pharmaceutically acceptable solvates and hydrates thereof.

One aspect of the present invention pertains to salts selected from: (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-edisylate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine phosphate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-oxalate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine succinate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine oxoglutarate salt; and pharmaceutically acceptable solvates and hydrates thereof.

One aspect of the present invention pertains to salts selected from: (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-edisylate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine phosphate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine citrate salt hemihydrate; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hemi-oxalate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine succinate salt; (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine oxoglutarate salt; and (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine oxoglutarate salt solvate.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt hemihydrate; and (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt solvate.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate salt.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt hemihydrate.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate salt.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine succinate salt.

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt;

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt solvate;

One aspect of the present invention pertains to pharmaceutical compositions comprising a salt of the present invention.

One aspect of the present invention pertains to process for preparing a pharmaceutical composition comprising admixing a salt of the present invention and a pharmaceutically acceptable carrier.

One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt of the present invention.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for weight management in an individual.

One aspect of the present invention pertains to salts of the present invention, for use in a method of treatment of the human or animal body by therapy.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight loss.

One aspect of the present invention pertains to salts of the present invention, for use in a method of maintenance of weight loss.

One aspect of the present invention pertains to salts of the present invention, for use in a method of decreasing food consumption.

One aspect of the present invention pertains to salts of the present invention, for use in a method of increasing meal-related satiety.

One aspect of the present invention pertains to salts of the present invention, for use in a method of reducing pre-meal hunger.

5 One aspect of the present invention pertains to salts of the present invention, for use in a method of reducing intra-meal food intake.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management further comprising a reduced-calorie diet.

10 One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management further comprising a program of regular exercise.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management further comprising a reduced-calorie diet and a program of regular exercise.

15 One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

20 One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 30 \text{ kg/m}^2$.

25 On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$.

On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

30 On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

35 On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$.

On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

5 On aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to salts of the present invention, for use in a method of weight management in combination with phentermine.

10 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention for use in a method of treatment of the human or animal body by therapy.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management.

15 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight loss.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of maintenance of weight loss.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of decreasing food consumption.

20 In some embodiments, the salts and pharmaceutical compositions are for use in a method of increasing meal-related satiety.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of reducing pre-meal hunger.

25 In some embodiments, the salts and pharmaceutical compositions are for use in a method of reducing intra-meal food intake.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management further comprising a reduced-calorie diet.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management further comprising a program of regular exercise.

30 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management further comprising a reduced-calorie diet and a program of regular exercise.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

35 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

5 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 30 \text{ kg/m}^2$.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$.

10 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

15 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$.

20 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

25 In some embodiments, the salts and pharmaceutical compositions are for use in a method of weight management in combination with phentermine.

CRYSTALLINE SALTS

30 Polymorphism is the ability of a substance to exist as two or more crystalline phases that have different arrangements and/or conformations of the molecules in the crystal lattice. Polymorphs show the same properties in the liquid or gaseous state but they may behave differently in the solid state.

35 Besides single-component polymorphs, drugs can also exist as salts and other multicomponent crystalline phases. For example, solvates and hydrates may contain an API host and either solvent or water molecules, respectively, as guests. Analogously, when the guest compound is a solid at room temperature, the resulting form is often called a cocrystal. Salts, solvates, hydrates, and cocrystals may show polymorphism as well. Crystalline phases that share

the same API host, but differ with respect to their guests, may be referred to as pseudopolymorphs of one another.

5 Solvates contain molecules of the solvent of crystallization in a definite crystal lattice. Solvates, in which the solvent of crystallization is water, are termed hydrates. Because water is a constituent of the atmosphere, hydrates of drugs may be formed rather easily.

Recently, polymorph screens of 245 compounds revealed that about 90% of them exhibited multiple solid forms. Overall, approximately half the compounds were polymorphic, often having one to three forms. About one-third of the compounds formed hydrates, and about one-third formed solvates. Data from cocrystal screens of 64 compounds showed that 60% 10 formed cocrystals other than hydrates or solvates. (G. P. Stahly, *Crystal Growth & Design* (2007), 7(6), 1007-1026.)

The present invention is directed, *inter alia*, to crystalline salts of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine and hydrates and solvates thereof. The crystalline forms of the salts of the present invention can be identified by unique solid state signatures with 15 respect to, for example, differential scanning calorimetry (DSC), X-ray powder diffraction (PXRD), and other solid state methods. Further characterization with respect to water or solvent content of the crystalline salts of the present invention can be gauged by any of the following methods for example, thermogravimetric analysis (TGA), DSC and the like. For DSC, it is known that the temperatures observed will depend upon sample purity, the rate of temperature 20 change, as well as sample preparation technique and the particular instrument employed. Thus, the values reported herein relating to DSC thermograms can vary by about ± 6 °C. The values reported herein relating to DSC thermograms can also vary by about ± 20 joules per gram. For PXRD, the relative intensities of the peaks can vary, depending upon the sample preparation technique, the sample mounting procedure and the particular instrument employed. Moreover, 25 instrument variation and other factors can often affect the 2θ values. Therefore, the peak assignments of diffraction patterns can vary by about ± 0.2 ° 2θ . The relative intensities of the reported peaks can also vary. For TGA, the features reported herein can vary by about ± 5 °C. The TGA features reported herein can also vary by about $\pm 2\%$ weight change due to, for example, sample variation. Further characterization with respect to hygroscopicity of the 30 crystalline salt can be gauged by, for example, dynamic moisture sorption (DMS). The DMS features reported herein can vary by about $\pm 5\%$ relative humidity. The DMS features reported herein can also vary by about $\pm 5\%$ weight change. The deliquescence relative humidity (DRH) measurements by water activity meter are sensitive to sample quality and quantity. The DRH measurements reported herein can vary by about $\pm 5\%$ RH.

35 **Compound I Hemi-edisylate Salt.**

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt, Form I (Compound 1 hemi-edisylate salt,

Form I). The physical properties of Compound 1 hemi-edisylate salt, Form I are summarized in Table 1 below.

Table 1

	Compound 1 Hemi-edisylate Salt, Form I
PXRD	Figure 1: Peaks of \geq 22% relative intensity at 6.00, 11.98, 16.07, 19.70, 20.12, 20.99, 22.39, 22.54, 23.12, 26.77, and 27.44 $^{\circ}2\theta$
TGA	Figure 2: < 0.1% weight loss below about 150 $^{\circ}\text{C}$
DSC	Figure 2: extrapolated melting onset at \sim 298 $^{\circ}\text{C}$
DMS	Figure 3: Non-hygroscopic
DRH	99.7% RH at 25 $^{\circ}\text{C}$

Form I of Compound 1 hemi-edisylate salt was an anhydrous crystalline material with a melting onset of \sim 298 $^{\circ}\text{C}$. It was non-hygroscopic by DMS analysis, picking up just 0.14% weight out to and including the 95% RH hold at 25 $^{\circ}\text{C}$. The DRH was determined by water activity measurement of saturated aqueous solution with excess solid to be 99.7% RH at 25 $^{\circ}\text{C}$.

Certain X-ray powder diffraction peaks for Form I of Compound 1 hemi-edisylate salt are shown in Table 2 below.

10

Table 2

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
6.00	76.32	26.77	22.17
10.31	5.18	27.44	42.76
11.98	63.17	27.91	6.99
13.65	15.02	28.78	6.82
14.63	21.05	29.44	10.45
16.07	42.02	29.70	6.33
16.79	9.47	30.18	18.52
17.98	21.48	30.85	2.77
19.70	74.89	31.29	3.71
20.12	53.46	32.65	3.88
20.64	14.33	33.12	5.93
20.99	36.94	33.74	3.51
21.72	3.22	34.32	6.72
22.39	32.48	35.06	5.31
22.54	49.14	35.46	6.63
23.12	100.00	35.86	11.13

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
24.13	42.45	36.23	7.83
24.97	16.43	37.63	2.62
25.51	7.48	38.95	5.33
25.88	15.57	39.27	4.59

One aspect of the present invention is directed to a Compound 1 hemi-edisylate salt having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 23.12° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 6.00° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° and about 6.00° . In some 5 embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° and about 19.70° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° , about 6.00° , and about 10 19.70° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° , about 6.00° , about 19.70° , about 11.98° , about 20.12° , about 22.54° , and about 27.44° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° , about 6.00° , about 19.70° , about 15 11.98° , about 20.12° , about 22.54° , about 27.44° , about 16.07° , about 20.99° , and about 22.39° . One aspect of the present invention is directed to a Compound 1 hemi-edisylate salt having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 2. In some embodiments, the salt has an X-ray powder diffraction pattern substantially as shown in Figure 1, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^{\circ}2\theta$ and also that the relative intensities of the reported peaks can vary.

20 In some embodiments, the Compound 1 hemi-edisylate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 285°C and about 315°C . In some embodiments, the Compound 1 hemi-edisylate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 298°C . In some embodiments, the Compound 1 hemi-edisylate salt has a differential scanning calorimetry thermogram comprising an endotherm with 25 an associated heat flow of about 101 joules per gram. In some embodiments, the Compound 1 hemi-edisylate salt has a thermogravimetric analysis profile substantially as shown in Figure 2, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^{\circ}\text{C}$ and by about $\pm 2\%$ weight change.

In some embodiments, the Compound 1 hemi-edisylate salt has a differential scanning calorimetry thermogram substantially as shown in Figure 2, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

5 In some embodiments, the Compound 1 hemi-edisylate salt has a dynamic moisture sorption profile substantially as shown in Figure 3, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

Form I of Compound 1 hemi-edisylate salt can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I 10 of Compound 1 hemi-edisylate salt can be prepared as described in Example 4. In some embodiments, Form I of Compound 1 hemi-edisylate salt can be prepared by slurrying crystalline Compound 1 hemi-edisylate salt containing one or more crystalline forms other than Form I. In some embodiments, Form I of Compound 1 hemi-edisylate salt can be prepared by recrystallizing crystalline Compound 1 hemi-edisylate salt containing one or more crystalline 15 forms other than Form I.

Compound 1 Phosphate Salt

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate, Form I (Compound 1 phosphate salt, Form I). The 20 physical properties of Form I of Compound 1 phosphate salt are summarized in Table 3 below.

Table 3

	Compound 1 Phosphate salt, Form I
PXRD	Figure 4: Peaks of $\geq 15\%$ relative intensity at 7.45, 14.87, 16.51, 18.16, 19.27, 20.19, 21.05, 23.19, 25.06, 25.77, 28.61, and 29.96 $^{\circ}2\theta$
TGA	Figure 5: $< 0.03\%$ weight loss up to about 178 °C
DSC	Figure 5: extrapolated onset temperature about 208 °C; enthalpy of fusion about 113 J/g
DMS	Figure 6: about 0.14% weight gain at 90% RH
DRH	100% RH at 25 °C

The title salt was a 1:1 salt based on stoichiometry determination. The melting onset by 25 DSC was ~ 208 °C. The TGA result for a crystalline sample prior to the *n*-propanol slurry is consistent with an anhydrous salt. It was non-hygroscopic, picking up 0.14% weight out to and including the 90% RH hold at 25 °C during DMS analysis. The title salt was non-deliquescent; the DRH by water activity measurement of a saturated solution in water with excess solid was 100% RH at 25 °C.

Certain X-ray powder diffraction peaks for Form I of Compound 1 phosphate salt are shown in Table 4 below.

Table 4

Pos. (°2θ)	Rel. Int. (%)	Pos. (°2θ)	Rel. Int. (%)
7.45	30.59	28.96	11.18
10.57	1.00	29.96	15.85
12.37	0.89	30.15	10.60
14.87	54.02	30.61	5.36
16.51	16.28	31.09	2.76
18.16	33.52	31.47	4.27
18.48	4.45	31.68	4.57
19.27	100.00	32.22	6.36
20.19	41.74	32.91	11.02
21.05	20.88	33.36	4.31
22.29	8.70	33.65	3.13
22.78	2.99	34.31	5.55
23.19	24.55	34.80	1.85
23.75	4.80	35.34	5.80
24.42	10.14	36.18	5.37
24.78	11.41	36.62	7.62
25.06	74.60	37.64	2.73
25.77	54.32	38.25	2.15
26.79	1.40	38.48	3.15
27.25	5.21	38.99	3.49
28.61	15.30	39.80	2.28

5 One aspect of the present invention is directed to a Compound 1 phosphate salt having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 19.27 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 25.06 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27 ° and about 25.06 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27 ° and about 25.77 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27 °, about 25.06 °, and about 25.77 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27 °, about 25.06 °, about 25.77 °, about 14.87 °, about 20.19 °, about 18.16 °, and

10

about 7.45 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27 °, about 25.06 °, about 25.77 °, about 14.87 °, about 20.19 °, about 18.16 °, about 7.45 °, about 23.19 °, about 21.05 °, and about 16.51 °. One aspect of the present invention is directed to a Compound 1 phosphate salt having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 4. In some embodiments, the salt has an X-ray powder diffraction pattern substantially as shown in Figure 4, wherein by “substantially” is meant that the reported peaks can vary by about ± 0.2 ° 2θ and also that the relative intensities of the reported peaks can vary.

5 In some embodiments, the Compound 1 phosphate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 190 °C and about 220 °C. In some embodiments, the Compound 1 phosphate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 208 °C. In some embodiments, the Compound 1 phosphate salt has a differential scanning calorimetry thermogram comprising an endotherm with an associated heat flow of about 113 joules per gram. In some embodiments, the Compound 1 phosphate salt has a thermogravimetric analysis profile substantially as shown in Figure 5, wherein by “substantially” is meant that the reported TGA features can vary by about ± 5 °C and by about $\pm 2\%$ weight change.

10 In some embodiments, the Compound 1 phosphate salt has a differential scanning calorimetry thermogram substantially as shown in Figure 5, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

15 In some embodiments, the Compound 1 phosphate salt has a dynamic moisture sorption profile substantially as shown in Figure 6, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

20 Form I of Compound 1 phosphate salt can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 phosphate salt can be prepared as described in Example 9. In some embodiments, Form I of Compound 1 phosphate salt can be prepared by slurring crystalline Compound 1 phosphate salt containing one or more crystalline forms other than Form I. In some 25 embodiments, Form I of Compound 1 phosphate salt can be prepared by recrystallizing crystalline Compound 1 phosphate salt containing one or more crystalline forms other than Form I.

Compound 1 Citrate Salt Hemihydrate

30 One aspect of the present invention pertains to (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine citrate salt hemihydrate, Form I (Compound 1 citrate salt

hemihydrate, Form I). The physical properties of Compound 1 citrate salt hemihydrate, Form I are summarized in Table 5 below.

Table 5

	Compound 1 Citrate Salt Hemihydrate, Form I
PXRD	Figure 7: Peaks of \geq 10% relative intensity at 11.93, 13.01, 17.12, 18.64, 19.11, 19.69, 20.73, 21.74, 24.05, 24.52, 24.81, 26.12, and 26.92 $^{\circ}2\theta$
TGA	Figure 8: 2.6% weight loss up to about 110 $^{\circ}\text{C}$
DSC	Figure 8: onset of dehydration about 80 $^{\circ}\text{C}$
DMS	Figure 9: 0.50% weight gain up to 90% RH
DRH	100% RH at 25 $^{\circ}\text{C}$

5 TGA data for Form I of Compound 1 citrate salt hemihydrate, showed that it was solvated. The mass loss matches closely with a hemihydrate (observed 2.6%, theoretical 2.3%). The onset of dehydration is near 80 $^{\circ}\text{C}$ for the scan rate, 10 $^{\circ}\text{C}/\text{min}$.

10 Form I of Compound 1 citrate salt hemihydrate lost only a small amount of its water of hydration during the drying step at 40 $^{\circ}\text{C}$ and \sim 1% RH for 1 h. It was not hygroscopic, picking up just 0.50% out to and including the 90% RH hold at 25 $^{\circ}\text{C}$, and not deliquescent. The DRH was determined by water activity measurement of saturated aqueous solution with excess solid to be 100% RH at 25 $^{\circ}\text{C}$.

Certain X-ray powder diffraction peaks for Form I of Compound 1 citrate salt hemihydrate are shown in Table 6 below.

Table 6

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
5.21	0.59	26.92	15.24
7.97	2.63	27.45	2.48
10.42	0.78	27.69	4.21
11.93	100.00	28.45	2.64
13.01	21.60	29.62	5.72
13.39	2.56	30.45	6.04
14.59	0.68	30.86	1.49
15.05	3.43	31.21	3.45
15.82	9.09	31.53	2.29
17.12	14.71	31.86	6.20
18.11	6.66	32.39	4.50
18.25	9.72	32.48	4.06

Pos. (°2θ)	Rel. Int. (%)	Pos. (°2θ)	Rel. Int. (%)
18.64	48.80	33.12	4.05
19.11	11.23	33.46	1.21
19.69	10.74	33.80	2.09
20.73	10.25	34.56	1.04
20.90	6.24	35.17	0.49
21.21	4.73	35.86	1.23
21.74	10.41	36.20	1.62
22.05	4.67	36.50	2.29
22.61	0.89	36.78	4.52
23.14	9.91	37.31	1.65
24.05	22.56	37.80	4.73
24.52	26.69	38.26	1.86
24.81	22.12	39.08	2.14
25.39	6.32	39.60	0.99
26.12	22.84		

One aspect of the present invention is directed to a Compound 1 citrate salt hemihydrate having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 11.93 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 18.64 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 11.93 ° and about 18.64 °. In some 5 embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 11.93 ° and about 24.52 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 11.93 °, about 18.64 °, and about 24.52 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising 10 peaks, in terms of 2θ , at about 11.93 °, about 18.64 °, about 24.52 °, about 26.12 °, about 24.05 °, about 24.81 °, and about 13.01 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 11.93 °, about 18.64 °, about 24.52 °, about 26.12 °, about 24.05 °, about 24.81 °, about 13.01 °, about 26.92 °, about 17.12 °, and about 19.11 °. One aspect of the present invention is directed to a Compound 1 citrate salt 15 hemihydrate having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 6. In some embodiments, the salt has an X-ray powder diffraction pattern substantially as shown in Figure 7, wherein by “substantially” is meant that the reported peaks can vary by about ± 0.2 ° 2θ and also that the relative intensities of the reported peaks can vary.

In some embodiments, the Compound 1 citrate salt hemihydrate has a 20 thermogravimetric analysis profile substantially as shown in Figure 8, wherein by

“substantially” is meant that the reported TGA features can vary by about ± 5 °C and by about $\pm 2\%$ weight change.

5 In some embodiments, the Compound 1 citrate salt hemihydrate has a differential scanning calorimetry thermogram substantially as shown in Figure 8, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

10 In some embodiments, the Compound 1 citrate salt hemihydrate has a dynamic moisture sorption profile substantially as shown in Figure 9, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

15 Form I of Compound 1 citrate salt hemihydrate can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 citrate salt hemihydrate can be prepared as described in Example 3. In some embodiments, Form I of Compound 1 citrate salt hemihydrate can be prepared by slurring crystalline Compound 1 citrate salt hemihydrate containing one or more crystalline forms other than Form I. In some embodiments, Form I of Compound 1 citrate salt hemihydrate can be prepared by recrystallizing crystalline Compound 1 citrate salt hemihydrate containing one or more crystalline forms other than Form I.

20 ***Compound 1 Hemi-oxalate Salt***

One aspect of the present invention pertains to (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate, Form I (Compound 1 hemi-oxalate salt, Form I). The physical properties of Compound 1 hemi-oxalate salt, Form I are summarized in Table 7 below.

25

Table 7

Compound 1 Hemi-oxalate Salt, Form I	
PXRD	Figure 10: Peaks of $\geq 10\%$ relative intensity at 6.34, 14.25, 14.51, 16.49, 21.69, 22.03, 24.06, 24.51, 24.92, 25.37, 31.85, 33.49, and 33.91 $^{\circ}2\theta$
TGA	Figure 11: $<0.3\%$ weight loss up to about 150 °C
DSC	Figure 11: extrapolated onset temperature about 212 °C
DMS	Figure 12: non-hygroscopic and non-deliquescent at 25 °C
DRH	100% RH at 25 °C

Form I of Compound 1 hemi-oxalate salt was anhydrous and displayed a melting onset temperature about 212 °C with weight loss by TGA starting just prior to the melting onset. It was determined to be non-hygroscopic and non-deliquescent at 25 °C. The DRH was

determined by water activity measurement of saturated aqueous solution with excess solid to be 100% RH at 25 °C.

Certain X-ray powder diffraction peaks for Form I of Compound 1 hemi-oxalate salt are shown in Table 8 below.

5

Table 8

Pos. (°2θ)	Rel. Int. (%)	Pos. (°2θ)	Rel. Int. (%)
6.34	100.00	25.37	44.12
10.81	0.92	26.60	8.81
11.45	1.22	27.13	6.17
12.65	5.04	27.45	4.50
14.25	23.10	27.81	5.49
14.51	18.93	28.38	4.45
15.43	3.19	29.12	3.46
16.49	10.86	30.44	3.29
17.14	2.19	30.84	3.95
18.31	2.20	31.85	48.33
18.98	4.57	32.76	1.74
19.82	7.58	33.49	15.33
21.03	2.13	33.91	12.30
21.69	51.81	34.53	5.38
22.03	27.18	35.71	1.00
22.56	3.78	36.50	1.20
23.03	1.90	37.60	2.62
24.06	22.77	38.37	1.52
24.51	13.01	39.39	1.31
24.92	13.18		

One aspect of the present invention is directed to a Compound 1 hemi-oxalate salt having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 6.34 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 21.69 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34 ° and about 21.69 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34 ° and about 31.85 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34 °, about 21.69 °, and about 31.85 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34 °, about 21.69 °, about 31.85 °, about 25.37 °, about 22.03 °, about 14.25 °, and

about 24.06 °. In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34 °, about 21.69 °, about 31.85 °, about 25.37 °, about 22.03 °, about 14.25 °, about 24.06 °, about 14.51 °, about 33.49 °, and about 24.92 °. One aspect of the present invention is directed to a Compound 1 hemi-oxalate salt having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 8. In some embodiments, the salt has an X-ray powder diffraction pattern substantially as shown in Figure 10, wherein by “substantially” is meant that the reported peaks can vary by about ± 0.2 ° 2θ and also that the relative intensities of the reported peaks can vary.

10 In some embodiments, the Compound 1 hemi-oxalate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 195 °C and about 225 °C. In some embodiments, the Compound 1 hemi-oxalate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 212 °C. In some embodiments, the Compound 1 hemi-oxalate salt has a thermogravimetric analysis profile substantially as shown in Figure 11, 15 wherein by “substantially” is meant that the reported TGA features can vary by about ± 5 °C and by about $\pm 2\%$ weight change.

20 In some embodiments, the Compound 1 hemi-oxalate salt has a differential scanning calorimetry thermogram substantially as shown in Figure 11, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

In some embodiments, the Compound 1 hemi-oxalate salt has a dynamic moisture sorption profile substantially as shown in Figure 12, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

25 Form I of Compound 1 hemi-oxalate salt can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 hemi-oxalate salt can be prepared as described in Example 4. In some embodiments, Form I of Compound 1 hemi-oxalate salt can be prepared by slurring crystalline Compound 1 hemi-oxalate salt containing one or more crystalline forms other than Form I. In 30 some embodiments, Form I of Compound 1 hemi-oxalate salt can be prepared by recrystallizing crystalline Compound 1 hemi-oxalate salt containing one or more crystalline forms other than Form I.

Compound 1 Succinate Salt

35 One aspect of the present invention pertains to (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine succinate, Form I (Compound 1 succinate salt, Form I). The physical properties of Compound 1 succinate salt, Form I are summarized in Table 9 below.

Table 9

Compound 1 Succinate Salt, Form I	
PXRD	Figure 13: Peaks of $\geq 5\%$ relative intensity at 12.53, 13.63, 15.64, 17.64, 20.65, 21.20, 24.35, 25.11, 26.54, 27.14, and 27.62 $^{\circ}2\theta$
TGA	Figure 14: <0.4% weight loss up to about 125 $^{\circ}\text{C}$
DSC	Figure 14: extrapolated onset temperature about 179 $^{\circ}\text{C}$; enthalpy of fusion 141 J/g
DMS	Figure 15: 0.07% weight gain at 90% RH
DRH	100% RH at 25 $^{\circ}\text{C}$

Compound 1 succinate salt, Form I showed a melting onset by DSC of 179.1 $^{\circ}\text{C}$. TGA showed no residual solvent, but did show apparent loss of significant succinic acid prior to the melting onset. It was non-hygroscopic by DMS analysis, picking up 0.07% weight out to and including the 90% RH hold at 25 $^{\circ}\text{C}$. The DRH was determined by water activity measurement of saturated aqueous solution with excess solid to be 100% RH at 25 $^{\circ}\text{C}$.

Certain X-ray powder diffraction peaks for Form I of Compound 1 succinate salt are shown in Table 10 below.

10

Table 10

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
6.84	1.00	26.05	3.56
11.58	2.35	26.54	14.35
12.53	6.13	27.14	7.58
12.71	3.50	27.62	100.00
13.63	7.75	28.44	2.23
15.07	1.31	30.40	1.02
15.64	34.98	31.73	4.21
16.10	3.90	32.34	0.51
16.41	2.86	32.88	0.89
17.64	13.76	33.60	0.24
19.59	0.24	34.36	0.95
20.34	4.78	35.67	1.66
20.65	38.05	36.34	0.96
21.20	5.87	36.91	2.16
21.61	1.36	37.08	1.53
22.18	1.17	38.00	1.19

Pos. (°2θ)	Rel. Int. (%)	Pos. (°2θ)	Rel. Int. (%)
23.21	2.29	38.61	2.36
24.35	14.11	38.97	0.70
25.11	24.83	39.59	1.01
25.51	4.64		

One aspect of the present invention is directed to a Compound 1 succinate salt having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 27.62° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 20.65° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° and about 20.65° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° and about 15.64° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° , about 20.65° , and about 15.64° . In some 5 embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° , about 20.65° , about 15.64° , about 25.11° , about 26.54° , about 24.35° , and about 17.64° . In some embodiments, the salt has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° , about 20.65° , about 15.64° , about 25.11° , about 26.54° , about 24.35° , about 17.64° , about 13.63° , about 27.14° , and about 12.53° . One 10 aspect of the present invention is directed to a Compound 1 succinate salt having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 10. In some embodiments, the salt has an X-ray powder diffraction pattern substantially as shown in Figure 13, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ 2\theta$ and also that the relative intensities of the reported peaks can vary.

15

20 In some embodiments, the Compound 1 succinate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 160°C and about 190°C . In some embodiments, the Compound 1 succinate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 179°C . In some embodiments, the Compound 1 25 succinate salt has a differential scanning calorimetry thermogram comprising an endotherm with an associated heat flow of about 141 joules per gram. In some embodiments, the Compound 1 succinate salt has a thermogravimetric analysis profile substantially as shown in Figure 14, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$ and by about $\pm 2\%$ weight change.

30 In some embodiments, the Compound 1 succinate salt has a differential scanning calorimetry thermogram substantially as shown in Figure 14, wherein by “substantially” is

meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

In some embodiments, the Compound 1 succinate salt has a dynamic moisture sorption profile substantially as shown in Figure 15, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

Form I of Compound 1 succinate salt can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 succinate salt can be prepared as described in Example 5. In some embodiments, Form I of Compound 1 succinate salt can be prepared by slurring crystalline Compound 1 succinate salt containing one or more crystalline forms other than Form I. In some embodiments, Form I of Compound 1 succinate salt can be prepared by recrystallizing crystalline Compound 1 succinate salt containing one or more crystalline forms other than Form I.

15 **Compound 1 Oxoglutarate Salt**

One aspect of the present invention pertains to a crystalline form of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt (Compound 1 oxoglutarate salt). In some embodiments, the crystalline form of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt is Form I (Compound 1 oxoglutarate salt, Form I). The physical properties of Form I of Compound 1 oxoglutarate salt are summarized in Table 11 below.

Table 11

Compound 1 Oxoglutarate Salt, Form I	
PXRD	Figure 16: Peaks of $\geq 15\%$ relative intensity at 7.86, 13.39, 18.71, 19.10, 20.06, 21.22, 22.84, 23.18, 23.57, 24.67, 25.37, and 26.81 $^{\circ}2\theta$
TGA	Figure 17: negligible weight loss below about 120 °C
DSC	Figure 17: extrapolated onset temperature about 115 °C; enthalpy of fusion 114 J/g
DMS	Figure 18: $\sim 0.106\%$ weight gain at about 90% RH

The title salt was anhydrous material by TGA with a melting onset of ~ 115 °C by DSC. 25 It was non-hygroscopic by DMS. Compound 1 oxoglutarate salt was non-hygroscopic by DMS analysis, picking up about 0.106% out to and including the 90% RH hold at 25 °C.

Certain X-ray powder diffraction peaks for Form I of Compound 1 oxoglutarate salt are shown in Table 12 below.

Table 12

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
7.86	41.21	25.37	16.94
10.09	6.35	25.75	12.57
10.65	3.29	26.81	32.66
11.45	4.60	27.39	2.32
13.39	100.00	27.83	3.40
14.30	14.50	28.41	5.39
15.12	1.59	29.43	5.16
15.69	3.23	29.62	5.28
17.40	2.12	30.62	3.15
18.71	31.16	31.63	2.70
19.10	46.89	32.16	2.09
19.37	5.60	32.76	5.80
20.06	38.48	33.64	4.97
21.22	55.23	34.30	3.31
21.93	3.04	34.66	6.12
22.46	8.90	35.50	2.04
22.84	15.51	35.87	3.24
23.18	48.32	36.20	2.67
23.57	58.00	36.71	1.15
23.82	10.53	37.53	2.34
24.29	9.84	38.13	2.72
24.67	48.09	38.54	4.38
25.12	9.56	38.86	4.03

One aspect of the present invention is directed to a crystalline form of Compound 1 oxoglutarate salt having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 13.39° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 23.57° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° and about 23.57° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° and about 21.22° .
 5 In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° , about 23.57° , and about 21.22° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° and about 21.22° .
 10 In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° , about 23.57° , and about 21.22° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at

about 13.39 °, about 23.57 °, about 21.22 °, about 23.18 °, about 24.67 °, about 19.10 °, and about 7.86 °. In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39 °, about 23.57 °, about 21.22 °, about 23.18 °, about 24.67 °, about 19.10 °, about 7.86 °, about 20.06 °, about 18.71 °, and about 25.37 °. One aspect of the present invention is directed to a crystalline form of Compound 1 oxoglutarate salt having an X-ray powder diffraction pattern comprising one or more peaks listed in Table 12. In some embodiments, the crystalline form has an X-ray powder diffraction pattern substantially as shown in Figure 16, wherein by “substantially” is meant that the reported peaks can vary by about ± 0.2 ° 2θ and also that the relative intensities of the reported peaks can vary.

10 In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 100 °C and about 130 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 115 °C.

15 In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a differential scanning calorimetry thermogram comprising an endotherm with an associated heat flow of about 114 joules per gram. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a thermogravimetric analysis profile substantially as shown in Figure 17, wherein by “substantially” is meant that the reported TGA features can vary by about ± 5 °C and

20 by about $\pm 2\%$ weight change.

In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a differential scanning calorimetry thermogram substantially as shown in Figure 17, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

25 In some embodiments, the crystalline form of Compound 1 oxoglutarate salt has a dynamic moisture sorption profile substantially as shown in Figure 18, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

Form I of Compound 1 oxoglutarate salt can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 oxoglutarate salt can be prepared as described in Example 6. In some embodiments, Form I of Compound 1 oxoglutarate salt can be prepared by slurring crystalline Compound 1 oxoglutarate salt containing one or more crystalline forms other than Form I. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt can be prepared by recrystallizing crystalline Compound 1 oxoglutarate salt containing one or more crystalline forms other than Form I.

Compound 1 Oxoglutarate Salt Solvate

One aspect of the present invention pertains to a crystalline form of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt solvate (Compound 1 oxoglutarate salt solvate). In some embodiments, the crystalline form of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt solvate is Form I (Compound 1 oxoglutarate salt solvate, Form I). The physical properties of Form I of Compound 1 oxoglutarate salt solvate are summarized in Table 13 below.

Table 13

	Compound 1 Oxoglutarate Salt Solvate, Form I
PXRD	Figure 19: Peaks of $\geq 30\%$ relative intensity at 6.29, 11.53, 12.51, 12.97, 14.76, 18.72, 22.65, 23.34, 23.71, 24.83, 25.16, and 26.02 $^{\circ}2\theta$
TGA	Figure 20: about 5.2% weight loss up to about 115 $^{\circ}\text{C}$
DSC	Figure 20: desolvation extrapolated onset temperature about 91 $^{\circ}\text{C}$; second endotherm extrapolated onset temperature about 113 $^{\circ}\text{C}$
DMS	Figure 21: $\sim 1.518\%$ weight gain at about 90% RH

10

The title salt was a solvated crystalline material with a desolvation onset of $\sim 91\text{ }^{\circ}\text{C}$ followed closely by another endotherm at $\sim 113\text{ }^{\circ}\text{C}$; both determined by DSC. Compound 1 oxoglutarate salt solvate, had a weight loss of $\sim 5.2\%$ (desolvation onset $\sim 84\text{ }^{\circ}\text{C}$ by TGA) out to $\sim 110\text{ }^{\circ}\text{C}$. This weight loss was slightly higher than the theoretical value (5.0%) for a monohydrate and slight lower than the theoretical value (5.7%) for a solvate. The desolvation is followed by degradation.

15 The title salt was non-hygroscopic by DMS analysis, losing about 1.518% out to and including the 90% RH hold at 25 $^{\circ}\text{C}$. This type of hysteresis is typically associated with displacement of an organic solvent by water and often results in a form change. However, by 20 PXRD analysis, the crystal form appeared unchanged

Certain X-ray powder diffraction peaks for Form I of Compound 1 oxoglutarate salt solvate are shown in Table 14 below.

Table 14

Pos. ($^{\circ}2\theta$)	Rel. Int. (%)	Pos. ($^{\circ}2\theta$)	Rel. Int. (%)
6.29	43.52	23.34	38.86
7.88	6.60	23.71	32.16
11.53	84.11	24.83	35.84

Pos. (°2θ)	Rel. Int. (%)	Pos. (°2θ)	Rel. Int. (%)
11.88	24.85	25.16	38.94
12.51	59.10	26.02	100.00
12.97	44.83	26.47	14.65
13.30	20.61	27.089	25.48
14.22	15.76	27.69	7.40
14.76	37.04	28.33	9.36
14.87	28.42	28.88	8.66
15.88	3.79	29.54	14.13
17.41	3.32	29.82	9.78
18.47	21.06	30.66	3.22
18.72	33.99	31.30	2.74
19.08	9.33	31.94	6.36
20.03	8.32	33.01	8.23
20.70	17.34	34.03	2.72
21.21	12.16	34.43	5.46
21.50	17.72	36.75	3.46
22.25	15.99	37.35	3.43
22.65	62.16		

One aspect of the present invention is directed to a crystalline form of Compound 1 oxoglutarate salt solvate having an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 26.02° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising a peak, in terms of 2θ , at about 11.53° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° and about 11.53° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° and about 22.65° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° , about 11.53° , and about 22.65° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° , about 11.53° , about 22.65° , about 12.51° , about 12.97° , about 6.29° , and about 25.16° . In some embodiments, the crystalline form has an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° , about 11.53° , about 22.65° , about 12.51° , about 12.97° , about 6.29° , about 25.16° , about 23.34° , about 14.76° , and about 24.83° . One aspect of the present invention is directed to a crystalline form of Compound 1 oxoglutarate salt solvate having an X-ray powder diffraction pattern comprising one or more peaks listed in Table

14. In some embodiments, the crystalline form has an X-ray powder diffraction pattern substantially as shown in Figure 19, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2 ^\circ 2\theta$ and also that the relative intensities of the reported peaks can vary.

In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 75 °C and about 105 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 91 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 100 °C and about 130 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 113 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature between about 75 °C and about 105 °C, and an endotherm with an extrapolated onset temperature between about 100 °C and about 130 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram comprising an endotherm with an extrapolated onset temperature at about 91 °C, and an endotherm with an extrapolated onset temperature at about 113 °C. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a thermogravimetric analysis profile substantially as shown in Figure 20, wherein by “substantially” is meant that the reported TGA features can vary by about ± 5 °C and by about $\pm 2\%$ weight change.

In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a differential scanning calorimetry thermogram substantially as shown in Figure 20, wherein by “substantially” is meant that the reported DSC features can vary by about ± 6 °C and by about ± 20 joules per gram.

In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate has a dynamic moisture sorption profile substantially as shown in Figure 21, wherein by “substantially” is meant that the reported DMS features can vary by about $\pm 5\%$ relative humidity and by about $\pm 5\%$ weight change.

Form I of Compound 1 oxoglutarate salt solvate can be prepared by any of the suitable procedures known in the art for preparing crystalline polymorphs. In some embodiments Form I of Compound 1 oxoglutarate salt solvate can be prepared as described in Example 7. In some embodiments, Form I of Compound 1 oxoglutarate salt solvate can be prepared by slurring crystalline Compound 1 oxoglutarate salt solvate containing one or more crystalline forms other

than Form I. In some embodiments, the crystalline form of Compound 1 oxoglutarate salt solvate can be prepared by recrystallizing crystalline Compound 1 oxoglutarate salt solvate containing one or more crystalline forms other than Form I.

5 One aspect of the present invention pertains to processes for preparing a pharmaceutical composition comprising admixing a crystalline salt of the present invention, and a pharmaceutically acceptable carrier.

One aspect of the present invention pertains to processes for preparing a bulk pharmaceutical composition comprising admixing a crystalline salt of the present invention, and a pharmaceutically acceptable carrier.

10 One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a crystalline salt of the present invention.

One aspect of the present invention pertains to the use of crystalline salts of the present invention, in the manufacture of a medicament for weight management in an individual.

15 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of treatment of the human or animal body by therapy.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management.

20 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight loss.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of maintenance of weight loss.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of decreasing food consumption.

25 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of increasing meal-related satiety.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of reducing pre-meal hunger.

30 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of reducing intra-meal food intake.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management further comprising a reduced-calorie diet.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management further comprising a program of regular exercise.

35 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management further comprising a reduced-calorie diet and a program of regular exercise.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

One aspect of the present invention pertains to crystalline salts of the present invention, 5 for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected 10 from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 30 \text{ kg/m}^2$.

One aspect of the present invention pertains to crystalline salts of the present invention, 15 for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

20 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to crystalline salts of the present invention, 25 for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$.

One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

30 One aspect of the present invention pertains to crystalline salts of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to crystalline salts of the present invention, 35 for use in a method of weight management in combination with phentermine.

HYDRATES AND SOLVATES

It is understood that when the phrase “pharmaceutically acceptable salts, solvates, and hydrates” or the phrase “pharmaceutically acceptable salt, solvate, or hydrate” is used when referring to compounds described herein, it embraces pharmaceutically acceptable solvates and/or hydrates of the compounds, pharmaceutically acceptable salts of the compounds, as well as pharmaceutically acceptable solvates and/or hydrates of pharmaceutically acceptable salts of the compounds. It is also understood that when the phrase “pharmaceutically acceptable solvates and hydrates” or the phrase “pharmaceutically acceptable solvate or hydrate” is used when referring to compounds described herein that are salts, it embraces pharmaceutically acceptable solvates and/or hydrates of such salts.

It will be apparent to those skilled in the art that the dosage forms described herein may comprise, as the active component, either a salts or crystalline form thereof as described herein, or a solvate or hydrate thereof. Moreover, various hydrates and solvates of the salts or crystalline form thereof described herein will find use as intermediates in the manufacture of pharmaceutical compositions. Typical procedures for making and identifying suitable hydrates and solvates, outside those mentioned herein, are well known to those in the art; see for example, pages 202-209 of K.J. Guillory, “Generation of Polymorphs, Hydrates, Solvates, and Amorphous Solids,” in: Polymorphism in Pharmaceutical Solids, ed. Harry G. Britain, Vol. 95, Marcel Dekker, Inc., New York, 1999.

Accordingly, one aspect of the present invention pertains to methods of administering hydrates and solvates of salts or crystalline forms thereof described herein and/or their pharmaceutically acceptable salts, that can be isolated and characterized by methods known in the art, such as, thermogravimetric analysis (TGA), TGA-mass spectroscopy, TGA-Infrared spectroscopy, powder X-ray diffraction (XRPD), Karl Fisher titration, high resolution X-ray diffraction, and the like. There are several commercial entities that provide quick and efficient services for identifying solvates and hydrates on a routine basis. Example companies offering these services include Wilmington PharmaTech (Wilmington, DE), Avantium Technologies (Amsterdam) and Aptuit (Greenwich, CT).

30 ISOTOPES

The present disclosure includes all isotopes of atoms occurring in the present salts and crystalline forms thereof. Isotopes include those atoms having the same atomic number but different mass numbers. One aspect of the present invention includes every combination of one or more atoms in the present salts and crystalline forms thereof that is replaced with an atom having the same atomic number but a different mass number. One such example is the replacement of an atom that is the most naturally abundant isotope, such as ¹H or ¹²C, found in one the present salts and crystalline forms thereof, with a different atom that is not the most

naturally abundant isotope, such as ^2H or ^3H (replacing ^1H), or ^{11}C , ^{13}C , or ^{14}C (replacing ^{12}C). A salt wherein such a replacement has taken place is commonly referred to as being isotopically-labeled. Isotopic-labeling of the present salts and crystalline forms thereof can be accomplished using any one of a variety of different synthetic methods known to those of ordinary skill in the art and they are readily credited with understanding the synthetic methods and available reagents needed to conduct such isotopic-labeling. By way of general example, and without limitation, isotopes of hydrogen include ^2H (deuterium) and ^3H (tritium). Isotopes of carbon include ^{11}C , ^{13}C , and ^{14}C . Isotopes of nitrogen include ^{13}N and ^{15}N . Isotopes of oxygen include ^{15}O , ^{17}O , and ^{18}O . An isotope of fluorine includes ^{18}F . An isotope of sulfur includes ^{35}S . An isotope of chlorine includes ^{36}Cl . Isotopes of bromine include ^{75}Br , ^{76}Br , ^{77}Br , and ^{82}Br . Isotopes of iodine include ^{123}I , ^{124}I , ^{125}I , and ^{131}I . Another aspect of the present invention includes compositions, such as, those prepared during synthesis, preformulation, and the like, and pharmaceutical compositions, such as, those prepared with the intent of using in a mammal for the treatment of one or more of the disorders described herein, comprising one or more of the present salts and crystalline forms thereof, wherein the naturally occurring distribution of the isotopes in the composition is perturbed. Another aspect of the present invention includes compositions and pharmaceutical compositions comprising salts and crystalline forms thereof as described herein wherein the salt is enriched at one or more positions with an isotope other than the most naturally abundant isotope. Methods are readily available to measure such isotope perturbations or enrichments, such as, mass spectrometry, and for isotopes that are radio-isotopes additional methods are available, such as, radio-detectors used in connection with HPLC or GC.

PHARMACEUTICAL COMPOSITIONS

A further aspect of the present invention pertains to pharmaceutical compositions comprising one or more salts according to any of the salt embodiments disclosed herein and one or more pharmaceutically acceptable carriers. Some embodiments pertain to pharmaceutical compositions comprising a salt according to any of the salt embodiments disclosed herein and a pharmaceutically acceptable carrier. Some embodiments pertain to pharmaceutical compositions comprising any subcombination of salts according to any of the salt embodiments disclosed herein.

Another aspect of the present invention pertains to methods of producing pharmaceutical compositions comprising admixing one or more salts according to any of the salt embodiments disclosed herein and one or more pharmaceutically acceptable carriers. Some embodiments pertain to a method of producing a pharmaceutical composition comprising admixing a salt according to any of the salt embodiments disclosed herein and a pharmaceutically acceptable carrier. Some embodiments pertain to a methods of producing

pharmaceutical compositions comprising admixing any subcombination of salts according to any of the salt embodiments disclosed herein and a pharmaceutically acceptable carrier.

5 Salts of the present invention or a solvate, hydrate or physiologically functional derivative thereof can be used as active ingredients in pharmaceutical compositions, specifically as 5-HT_{2C}-receptor modulators. The term “active ingredient” as defined in the context of a “pharmaceutical composition” and is intended to mean a component of a pharmaceutical composition that provides the primary pharmacological effect, as opposed to an “inactive ingredient” which would generally be recognized as providing no pharmaceutical benefit.

10 The dose when using the salts of the present invention can vary within wide limits and as is customary and is known to the physician, it is to be tailored to the individual conditions in each individual case. It depends, for example, on the nature and severity of the illness to be treated, on the condition of the patient, on the salt employed or on whether an acute or chronic disease state is treated or prophylaxis conducted or on whether further active compounds are administered in addition to the salts of the present invention. Representative doses of the present 15 invention include, but are not limited to, about 0.001 mg to about 5000 mg, about 0.001 mg to about 2500 mg, about 0.001 mg to about 1000 mg, 0.001 mg to about 500 mg, 0.001 mg to about 250 mg, about 0.001 mg to 100 mg, about 0.001 mg to about 50 mg and about 0.001 mg to about 25 mg. Multiple doses may be administered during the day, especially when relatively large amounts are deemed to be needed, for example 2, 3 or 4 doses. Depending on the 20 individual and as deemed appropriate from the patient's physician or caregiver it may be necessary to deviate upward or downward from the doses described herein.

25 The amount of active ingredient, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will ultimately be at the discretion of the attendant physician or clinician. In general, one skilled in the art understands how to extrapolate *in vivo* data obtained in a model system, typically an animal model, to another, such as a human. In some circumstances, these 30 extrapolations may merely be based on the weight of the animal model in comparison to another, such as a mammal, preferably a human, however, more often, these extrapolations are not simply based on weights, but rather incorporate a variety of factors. Representative factors include the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, 35 efficacy, pharmacokinetic and toxicology profiles of the particular salt employed, whether a drug delivery system is utilized, on whether an acute or chronic disease state is being treated or prophylaxis conducted or on whether further active compounds are administered in addition to the salts of the present invention and as part of a drug combination. The dosage regimen for treating a disease condition with the salts and/or compositions of this invention is selected in

accordance with a variety factors as cited above. Thus, the actual dosage regimen employed may vary widely and therefore may deviate from a preferred dosage regimen and one skilled in the art will recognize that dosage and dosage regimen outside these typical ranges can be tested and, where appropriate, may be used in the methods of this invention.

5 The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, *e.g.*, into a number of discrete loosely spaced administrations. The daily dose can be divided, especially when relatively large amounts are administered as deemed appropriate, into several, for example 2, 3 or 4 part administrations. If 10 appropriate, depending on individual behavior, it may be necessary to deviate upward or downward from the daily dose indicated.

15 Some embodiments of the present invention include a method of producing a pharmaceutical composition for “combination-therapy” comprising admixing at least one salt according to any of the salt embodiments disclosed herein, together with at least one known pharmaceutical agent as described herein and a pharmaceutically acceptable carrier.

It is noted that when the salts of the present invention are utilized as active ingredients in a pharmaceutical composition, these are not intended for use only in humans, but in other non-human mammals as well. Indeed, recent advances in the area of animal health-care mandate that consideration be given for the use of active agents, such as 5-HT_{2C}-receptor modulators, for 20 the treatment of a 5-HT_{2C}-receptor-associated disease or disorders in companionship animals (*e.g.*, cats, dogs, *etc.*) and in livestock animals (*e.g.*, cows, chickens, fish, *etc.*). Those of ordinary skill in the art are readily credited with understanding the utility of such salts in such settings.

25 One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a pharmaceutical composition of the present invention.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of treatment of the human or animal body by therapy.

30 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight loss.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of maintenance of weight loss.

35 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of decreasing food consumption.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of increasing meal-related satiety.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of reducing pre-meal hunger.

5 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of reducing intra-meal food intake.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management further comprising a reduced-calorie diet.

10 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management further comprising a program of regular exercise.

15 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management further comprising a reduced-calorie diet and a program of regular exercise.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

20 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

25 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

30 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 30 \text{ kg/m}^2$.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$.

35 One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in an individual with an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related co-morbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

One aspect of the present invention pertains to pharmaceutical compositions of the present invention, for use in a method of weight management in combination with phentermine.

20 HYGROSCOPICITY

Many compounds and salts are sensitive to the presence of water vapor or moisture. When compounds and salts interact with moisture, they retain water by either bulk or surface adsorption, capillary condensation, chemical reaction and, in extreme cases, formation of a solution (deliquescence). Deliquescence occurs when a solid dissolves and saturates a thin film of water on its surface. It has been shown that when moisture is absorbed to the extent that deliquescence takes place at a certain critical relative humidity, the liquid film surrounding the solid is saturated. This process is dictated by vapor diffusion and heat transport rates. (Gibson, *Pharmaceutical Preformulation and Formulation: A Practical Guide from Candidate Drug Selection to Commercial Dosage Form*, Informa Health Care, 2001; Kontny *et al.*, *Pharmaceutical Research*, 1987, 4(2), 104-12.)

The opposite of deliquescence is efflorescence, which occurs when a crystal loses water of crystallization below a critical vapor pressure. For example Griesser and Burger (*International Journal of Pharmaceutics*, 1995, 120(1), 83-93) found that caffeine hydrate lost its water of crystallization even at 61% RH. It has also been observed that the three known polymorphs of oxytetracycline have different hygroscopicity profiles. (Burger *et al.*, *Acta Pharmaceutica Technologica*, 1985, 31(4), 230-5.)

Moisture is also an important factor that can affect the stability of candidate drugs and their formulations. Sorption of water molecules onto a candidate drug (or excipient) can often induce hydrolysis (see, e.g., Yoshioka and Carstensen, *Journal of Pharmaceutical Sciences*, 1990, 79(9), 799-801. Other properties such as crystal structure, powder flow, compaction, lubricity, dissolution rate and polymer film permeability may also be effected by moisture adsorption. (Ahlnack and Zografi, *International Journal of Pharmaceutics*, 1990, 62(2-3), 87-95.)

The influence that moisture has on stability depends on how strongly it is bound, *i.e.*, it depends on whether the moisture is in a free or a bound state. Generally, degradation arises as a function of free water, which may be due to its ability to change the pH of the surfaces of drug and excipient. (Monkhouse, *Drug Development and Industrial Pharmacy*, 1984, 10(8-9), 1373-412.) On the other hand, bound water is not available if it is a crystal hydrate, hydrogen bonded, or sorbed or trapped in an amorphous structure

Hygroscopicity can be defined using various parameters. For example, hygroscopicity may be classified as shown in the following table. Callahan *et al.* (*Drug Development and Industrial Pharmacy*, 1982, 8(3), 355-69).

Class 1: Non-hygroscopic	Essentially no moisture increases occur at RH < 90%
Class 2: Slightly hygroscopic	Essentially no moisture increases occur at RH < 90%
Class 3: Moderately hygroscopic	Moisture content increase $\leq 5\%$ after storage for 1 week at RH < 60%
Class 4: Very hygroscopic	Moisture content increase may occur at RH as low as 40-50%

Alternatively, hygroscopicity can be defined using the parameters of the European Pharmacopoeia Technical Guide (1999, p. 86) which has defined hygroscopicity, based on the static method, after storage at 25 °C for 24 hours at 80% RH.

Classification	Increase in mass after 24 h at 25 °C and 80% RH
Slightly hygroscopic	< 2 % and $\geq 0.2\%$
Hygroscopic	< 15 % and $\geq 0.2\%$
Very hygroscopic	$\geq 15\%$
Deliquescent	Forms a liquid

It is highly desirable to have a crystalline form of a therapeutic agent that is neither hygroscopic nor deliquescent. Stable, non-hygroscopic salts facilitate the production of solid pharmaceutical compositions. Hygroscopicity of active pharmaceutical ingredients can cause a number of down-stream problems including a lack of storability, agglomeration, and inadequate flowability during formulating and processing. Hygroscopic formulations may exhibit poor 5 tablettability which can make the manufacture of orally administrable dosage forms problematic.

The salts of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine of the present 10 invention are non-hygroscopic. Because of their stability to moisture they are suitable for active pharmaceutical ingredient storage, for preparing bulk pharmaceutical compositions, and for manufacturing orally administrable solid-dosage forms that are useful for, *inter alia*, weight management.

One aspect of the present invention pertains to dosage forms comprising a 15 therapeutically effective amount of a salt selected from: a pharmaceutically acceptable salt of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine and pharmaceutically acceptable solvates and hydrates thereof, wherein the dosage form is a non-hygroscopic dosage form.

In some embodiments, the salt absorbs less than about 2% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 1% water by weight after about 20 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.6% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.5% water by weight after about 25 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.2% water by weight after about 30 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs less than about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs: less than about 2% water by weight after about 35 2 h at about 90% RH and about 25 °C; less than about 1% water by weight after about 2 h at about 90% RH and about 25 °C; less than about 0.6% water by weight after about 2 h at about 90% RH and about 25 °C; less than about 0.5% water by weight after about 2 h at about 90% RH and about 25 °C; less than about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C;

about 25 °C; less than about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C; less than about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C; or less than about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

5 In some embodiments, the salt absorbs from about 2% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 2% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 2% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

10 In some embodiments, the salt absorbs from about 2% to about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 2% to about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C.

15 In some embodiments, the salt absorbs from about 2% to about 0.5% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 2% to about 0.6% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 2% to about 1% water by weight after about 2 h at about 90% RH and about 25 °C.

20 In some embodiments, the salt absorbs from about 1% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 1% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

25 In some embodiments, the salt absorbs from about 1% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 1% to about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 1% to about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C.

30 In some embodiments, the salt absorbs from about 1% to about 0.5% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 1% to about 0.6% water by weight after about 2 h at about 90% RH and about 25 °C.

35 In some embodiments, the salt absorbs from about 0.6% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.6% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.6% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.6% to about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

5 In some embodiments, the salt absorbs from about 0.6% to about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.6% to about 0.5% water by weight after about 2 h at about 90% RH and about 25 °C.

10 In some embodiments, the salt absorbs from about 0.5% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.5% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.5% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

15 In some embodiments, the salt absorbs from about 0.5% to about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.5% to about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C.

20 In some embodiments, the salt absorbs from about 0.4% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.4% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.4% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

25 In some embodiments, the salt absorbs from about 0.4% to about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.3% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

30 In some embodiments, the salt absorbs from about 0.3% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.3% to about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.2% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

35 In some embodiments, the salt absorbs from about 0.2% to about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.

In some embodiments, the salt absorbs from about 0.1% to about 0.01% water by weight after about 2 h at about 90% RH and about 25 °C.

One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of a salt of the present invention.

5 One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt or a crystalline form thereof.

10 One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate salt or a crystalline form thereof.

One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt hemihydrate or a crystalline form thereof.

15 One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate salt or a crystalline form thereof.

One aspect of the present invention pertains to dosage forms comprising a therapeutically effective amount of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine succinate salt or a crystalline form thereof.

20 One aspect of the present invention pertains to dosage forms for oral administration to an individual in need of weight management.

25 In some embodiments, the weight management comprises weight loss.

In some embodiments, the weight management comprises maintenance of weight loss.

In some embodiments, the weight management comprises decreased food consumption.

In some embodiments, the weight management comprises increasing meal-related satiety.

30 In some embodiments, the weight management comprises reducing pre-meal hunger.

In some embodiments, the weight management comprises reducing intra-meal food intake.

In some embodiments, the weight management further comprises a reduced-calorie diet.

In some embodiments, the weight management further comprises a program of regular exercise.

35 In some embodiments, the weight management further comprises both a reduced-calorie diet and a program of regular exercise.

In some embodiments, the individual in need of weight management is an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

In some embodiments, the individual in need of weight management is an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related 5 comorbid condition.

In some embodiments, the weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the dosage form is for administration in combination with phentermine.

10

INDICATIONS

Obesity is a life-threatening disorder in which there is an increased risk of morbidity and mortality arising from concomitant diseases such as, but not limited to, type II diabetes, hypertension, stroke, certain forms of cancers and gallbladder disease.

15

Obesity has become a major healthcare issue in the Western World and increasingly in some third world countries. The increase in the number of obese people is due largely to the increasing preference for high fat content foods but also, and this can be a more important factor, the decrease in activity in most people's lives. In spite of the growing awareness of the health concerns linked to obesity the percentage of individuals that are overweight or obese 20 continues to increase. The most significant concern, from a public health perspective, is that children who are overweight grow up to be overweight or obese adults, and accordingly are at greater risk for major health problems. Therefore, it appears that the number of individuals that are overweight or obese will continue to increase.

25

Whether someone is classified as overweight or obese is generally determined on the basis of his or her body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m^2). Thus, the units for BMI are kg/m^2 . BMI is more highly correlated with body fat than any other indicator of height and weight. A person is considered overweight when they have a BMI in the range of 25-30 kg/m^2 , whereas a person with a BMI over 30 kg/m^2 is 30 classified as obese. Obesity is further divided into three classes: Class I (BMI of about 30 to about 34.9 kg/m^2), Class II (BMI of about 35 to 39.9 kg/m^2) and Class III (about 40 kg/m^2 or greater); see Table below for complete classifications.

Classification Of Weight By Body Mass Index (BMI)

BMI	CLASSIFICATION
< 18.5	Underweight
18.5-24.9	Normal
25.0-29.9	Overweight

30.0-34.9	Obesity (Class I)
35.0-39.9	Obesity (Class II)
> 40	Extreme Obesity (Class III)

As the BMI increases for an individual there is an increased risk of morbidity and mortality relative to an individual with normal BMI. Accordingly, overweight and obese individuals (BMI of about 25 kg/m^2 and above) are at increased risk for physical ailments such as, but not limited to, high blood pressure, cardiovascular disease (particularly hypertension),
5 high blood cholesterol, dyslipidemia, type II (non-insulin dependent) diabetes, insulin resistance, glucose intolerance, hyperinsulinemia, coronary heart disease, angina pectoris, congestive heart failure, stroke, gallstones, cholecystitis and cholelithiasis, gout, osteoarthritis, obstructive sleep apnea and respiratory problems, some types of cancer (such as endometrial, breast, prostate, and colon), complications of pregnancy, poor female reproductive health (such
10 as menstrual irregularities, infertility, irregular ovulation), diseases of reproduction (such as sexual dysfunction, both male and female, including male erectile dysfunction), bladder control problems (such as stress incontinence), uric acid nephrolithiasis, psychological disorders (such as depression, eating disorders, distorted body image, and low self esteem). Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk
15 of developing other ailments, such as, but not limited to, coronary heart disease.

As mentioned above, obesity increases the risk of developing cardiovascular diseases. Coronary insufficiency, atheromatous disease, and cardiac insufficiency are at the forefront of the cardiovascular complications induced by obesity. The incidence of coronary diseases is doubled in subjects less than 50 years of age who are 30% overweight. The diabetes patient
20 faces a 30% reduced lifespan. After age 45, people with diabetes are about three times more likely than people without diabetes to have significant heart disease and up to five times more likely to have a stroke. These findings emphasize the inter-relations between risks factors for type 2 diabetes and coronary heart disease and the potential value of an integrated approach to the prevention of these conditions based on the prevention of obesity [Perry, I. J., *et al. BMJ*
25 310, 560-564 (1995)]. It is estimated that if the entire population had an ideal weight, the risk of coronary insufficiency would decrease by 25% and the risk of cardiac insufficiency and of cerebral vascular accidents by 35%.

Diabetes has also been implicated in the development of kidney disease, eye diseases and nervous-system problems. Kidney disease, also called nephropathy, occurs when the
30 kidney's "filter mechanism" is damaged and protein leaks into urine in excessive amounts and eventually the kidney fails. Diabetes is also a leading cause of damage to the retina and increases the risk of cataracts and glaucoma. Finally, diabetes is associated with nerve damage, especially in the legs and feet, which interferes with the ability to sense pain and contributes to

serious infections. Taken together, diabetes complications are one of the nation's leading causes of death.

5 The first line of treatment for individuals that are overweight or obese is to offer diet and life style advice, such as, reducing the fat content of their diet and increasing their physical activity. However many patients find these difficult to maintain and need additional help from drug therapy to sustain results from these efforts.

10 Most currently marketed products have been unsuccessful as treatments for obesity owing to a lack of efficacy or unacceptable side-effect profiles. The most successful drug so far was the indirectly acting 5-hydroxytryptamine (5-HT) agonist d-fenfluramine (ReduxTM) but reports of cardiac valve defects in up to one third of the patient population led to its withdrawal by the FDA in 1998.

15 The 5-HT_{2C} receptor is recognized as a well-accepted receptor target for the treatment of obesity, psychiatric, and other disorders. See, for example, Halford *et al.*, *Serotonergic Drugs Effects on Appetite Expression and Use for the Treatment of Obesity*, Drugs 2007; 67 (1): 27-55; Naughton *et al.*, *A Review Of The Role Of Serotonin Receptors In Psychiatric Disorders*. Human Psychopharmacology (2000), 15(6), 397-415.

20 (R)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride (lorcaserin hydrochloride) is an agonist of the 5-HT_{2C} receptor and shows effectiveness at reducing obesity in animal models and humans. In a phase 3 human clinical trial evaluating the safety and efficacy of lorcaserin for weight management, statistical significance ($p < 0.0001$) was achieved on all three of the hierarchically ordered co-primary endpoints for patients treated with lorcaserin versus placebo. Treatment with lorcaserin was generally very well tolerated. An assessment of echocardiograms indicated no apparent drug-related effect on the development of US Food and Drug Administration (FDA)-defined valvulopathy over the two-year treatment period. The hierarchically ordered endpoints were the proportion of patients achieving 5% or greater weight loss after 12 months, the difference in mean weight loss compared to placebo after 12 months, and the proportion of patients achieving 10% or greater weight loss after 12 months. Compared to placebo, using an intent-to-treat last observation carried forward (ITT-LOCF) analysis, treatment with lorcaserin was associated with highly statistically significant ($p < 0.0001$) categorical and average weight loss from baseline after 12 months: 47.5% of lorcaserin patients lost greater than or equal to 5% of their body weight from baseline compared to 20.3% in the placebo group. This result satisfied the efficacy benchmark in the most recent FDA draft guidance. Average weight loss of 5.8% of body weight, or 12.7 pounds, was achieved in the lorcaserin group, compared to 2.2% of body weight, or 4.7 pounds, in the placebo group. 25 Statistical separation from placebo was observed by Week 2, the first post-baseline measurement. 22.6% of lorcaserin patients lost greater than or equal to 10% of their body weight from baseline, compared to 7.7% in the placebo group. Lorcaserin patients who completed 52

weeks of treatment according to the protocol lost an average of 8.2% of body weight, or 17.9 pounds, compared to 3.4%, or 7.3 pounds, in the placebo group (p < 0.0001).

In addition, the 5-HT_{2C} receptor is also involved in other diseases, conditions and disorders, such as, obsessive compulsive disorder, some forms of depression, and epilepsy.

5 Accordingly, 5-HT_{2C} receptor agonists can have anti-panic properties, and properties useful for the treatment of sexual dysfunction. In addition, 5-HT_{2C} receptor agonists are useful for the treatment of psychiatric symptoms and behaviors in individuals with eating disorders such as, but not limited to, anorexia nervosa and bulimia nervosa. Individuals with anorexia nervosa often demonstrate social isolation. Anorexic individuals often present symptoms of being

10 depressed, anxious, obsession, perfectionistic traits, and rigid cognitive styles as well as sexual disinterest. Other eating disorders include, anorexia nervosa, bulimia nervosa, binge eating disorder (compulsive eating) and ED-NOS (*i.e.*, eating disorders not otherwise specified - an official diagnosis). An individual diagnosed with ED-NOS possess atypical eating disorders including situations in which the individual meets all but a few of the criteria for a particular

15 diagnosis. What the individual is doing with regard to food and weight is neither normal nor healthy.

The 5-HT_{2C} receptor plays a role in Alzheimer Disease (AD). Therapeutic agents currently prescribed for Alzheimer's disease (AD) are cholinomimetic agents that act by inhibiting the enzyme acetylcholinesterase. The resulting effect is increased levels of

20 acetylcholine, which modestly improves neuronal function and cognition in patients with AD. Although, dysfunction of cholinergic brain neurons is an early manifestation of AD, attempts to slow the progression of the disease with these agents have had only modest success, perhaps because the doses that can be administered are limited by peripheral cholinergic side effects, such as tremors, nausea, vomiting, and dry mouth. In addition, as AD progresses, these agents

25 tend to lose their effectiveness due to continued cholinergic neuronal loss.

Therefore, there is a need for agents that have beneficial effects in AD, particularly in alleviating symptoms by improving cognition and slowing or inhibiting disease progression, without the side effects observed with current therapies. Therefore, serotonin 5-HT_{2C} receptors, which are exclusively expressed in brain, are attractive targets.

30 Another disease, disorder or condition that can is associated with the function of the 5-HT_{2C} receptor is erectile dysfunction (ED). Erectile dysfunction is the inability to achieve or maintain an erection sufficiently rigid for intercourse, ejaculation, or both. An estimated 20-30 million men in the United States have this condition at some time in their lives. The prevalence of the condition increases with age. Five percent of men 40 years of age report ED. This rate

35 increases to between 15% and 25% by the age of 65, and to 55% in men over the age of 75 years.

Erectile dysfunction can result from a number of distinct problems. These include loss of desire or libido, the inability to maintain an erection, premature ejaculation, lack of emission, and inability to achieve an orgasm. Frequently, more than one of these problems presents themselves simultaneously. The conditions may be secondary to other disease states (typically 5 chronic conditions), the result of specific disorders of the urogenital system or endocrine system, secondary to treatment with pharmacological agents (*e.g.* antihypertensive drugs, antidepressant drugs, antipsychotic drugs, *etc.*) or the result of psychiatric problems. Erectile dysfunction, when organic, is primarily due to vascular irregularities associated with atherosclerosis, diabetes, and hypertension.

10 There is evidence for use of a serotonin 5-HT_{2C} agonist for the treatment of sexual dysfunction in males and females. The serotonin 5-HT_{2C} receptor is involved with the processing and integration of sensory information, regulation of central monoaminergic systems, and modulation of neuroendocrine responses, anxiety, feeding behavior, and cerebrospinal fluid production [Tecott, L. H., *et al. Nature* 374: 542-546 (1995)]. In addition, the serotonin 5-HT_{2C} receptor has been implicated in the mediation of penile erections in rats, monkeys, and humans.

15 In summary, the 5-HT_{2C} receptor is a validated and well-accepted receptor target for the prophylaxis and/or treatment of 5-HT_{2C} mediated receptor diseases and disorders, such as, obesity, eating disorders, psychiatric disorders, Alzheimer Disease, sexual dysfunction and disorders related thereto. It can be seen that there exists a need for selective 5-HT_{2C} receptor agonists that can safely address these needs. The present invention is directed to these, as well as other, important ends.

One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

25 In some embodiments, the weight management comprises weight loss.

In some embodiments, the weight management comprises maintenance of weight loss.

In some embodiments, the weight management comprises decreased food consumption.

In some embodiments, the weight management comprises increasing meal-related satiety.

30 In some embodiments, the weight management comprises reducing pre-meal hunger.

In some embodiments, the weight management comprises reducing intra-meal food intake.

In some embodiments, the weight management further comprises a reduced-calorie diet.

35 In some embodiments, the weight management further comprises a program of regular exercise.

In some embodiments, the weight management further comprises both a reduced-calorie diet and a program of regular exercise.

In some embodiments, the individual in need of weight management is an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

5 In some embodiments, the individual in need of weight management is an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

In some embodiments, the individual in need of weight management is an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

10 In some embodiments, the individual in need of weight management has an initial body mass index $\geq 30 \text{ kg/m}^2$.

In some embodiments, the individual in need of weight management has an initial body mass index $\geq 27 \text{ kg/m}^2$.

15 In some embodiments, the individual in need of weight management has an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

In some embodiments, the individual in need of weight management has an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

20 In some embodiments, the individual in need of weight management has an initial body mass index $\geq 25 \text{ kg/m}^2$.

In some embodiments, the individual in need of weight management has an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

25 In some embodiments, the individual in need of weight management has an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 20 kg/m^2 in the presence of at least one weight related comorbid condition.

30 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 20 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 21 kg/m^2 in the presence of at least one weight related comorbid condition.

35 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 21 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 22 kg/m^2 in the presence of at least one weight related comorbid condition.

5 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 22 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 23 kg/m^2 in the presence of at least one weight related comorbid condition.

10 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 23 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 24 kg/m^2 in the presence of at least one weight related comorbid condition.

15 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 24 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

20 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 25 kg/m^2 in the presence of at least one weight related comorbid condition.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 25 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

25 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 26 kg/m^2 in the presence of at least one weight related comorbid condition.

30 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 26 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 27 kg/m^2 in the presence of at least one weight related comorbid condition.

35 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 27 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 28 kg/m^2 in the presence of at least one weight related comorbid condition.

5 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 28 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 29 kg/m^2 in the presence of at least one weight related comorbid condition.

10 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 29 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 30 kg/m^2 in the presence of at least one weight related comorbid condition.

15 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 30 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

20 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 31 kg/m^2 in the presence of at least one weight related comorbid condition.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 31 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

25 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 32 kg/m^2 in the presence of at least one weight related comorbid condition.

30 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 32 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 33 kg/m^2 in the presence of at least one weight related comorbid condition.

35 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 33 kg/m^2 in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 34 kg/m² in the presence of at least one weight related comorbid condition.

5 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 34 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 35 kg/m² in the presence of at least one weight related comorbid condition.

10 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 35 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 36 kg/m² in the presence of at least one weight related comorbid condition.

15 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 36 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

20 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 37 kg/m² in the presence of at least one weight related comorbid condition.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 37 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

25 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 38 kg/m² in the presence of at least one weight related comorbid condition.

30 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 38 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 39 kg/m² in the presence of at least one weight related comorbid condition.

35 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 39 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual in need of weight management has an initial body mass index \geq about 40 kg/m² in the presence of at least one weight related comorbid condition.

5 In some embodiments, the individual in need of weight management has an initial body mass index \geq about 40 kg/m² in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the method for weight management further comprises administering phentermine to the individual.

10 One aspect of the present invention pertains to methods for the treatment of a disorder related to 5-HT_{2C} receptor activity in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

15 One aspect of the present invention pertains to methods for the treatment of obesity, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

In some embodiments, the method for the treatment of obesity further comprises the administration or prescription of phentermine.

In some embodiments, the method for the treatment of obesity further comprises gastric electrical stimulation.

20 One aspect of the present invention pertains to methods for inducing weight loss, BMI loss, waist circumference loss or body fat percentage loss, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

25 One aspect of the present invention pertains to methods for inducing weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual in preparation of the individual for bariatric surgery, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

30 One aspect of the present invention pertains to methods for maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

35 One aspect of the present invention pertains to methods for maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual following bariatric surgery, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

One aspect of the present invention pertains to methods for inducing satiety in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

5 One aspect of the present invention pertains to methods for decreasing food intake in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

One aspect of the present invention pertains to methods for decreasing hunger in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

10 One aspect of the present invention pertains to methods for decreasing food cravings in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

15 One aspect of the present invention pertains to methods for increasing intermeal interval in an individual, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

20 One aspect of the present invention pertains to methods for the treatment of a disorder selected from: schizophrenia, anxiety, depression, psychoses and alcohol addiction, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

In some embodiments, the disorder is schizophrenia.

In some embodiments, the disorder is anxiety.

In some embodiments, the disorder is depression.

25 In some embodiments, the disorder is psychoses.

In some embodiments, the disorder is alcohol addiction.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for weight management in an individual.

30 In some embodiments, the weight management comprises weight loss.

In some embodiments, the weight management comprises maintenance of weight loss.

In some embodiments, the weight management comprises decreased food consumption.

35 In some embodiments, the weight management comprises increasing meal-related satiety.

In some embodiments, the weight management comprises reducing pre-meal hunger.

In some embodiments, the weight management comprises reducing intra-meal food intake.

In some embodiments, the weight management further comprises a reduced-calorie diet.

In some embodiments, the weight management further comprises a program of regular exercise.

In some embodiments, the weight management further comprises both a reduced-calorie diet and a program of regular exercise.

5 In some embodiments, the individual is an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$.

In some embodiments, the individual is an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

10 In some embodiments, the individual is an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual has an initial body mass index $\geq 30 \text{ kg/m}^2$.

In some embodiments, the individual has an initial body mass index $\geq 27 \text{ kg/m}^2$.

15 In some embodiments, the individual has an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

In some embodiments, the individual has an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

In some embodiments, the individual has an initial body mass index $\geq 25 \text{ kg/m}^2$.

20 In some embodiments, the individual has an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition.

In some embodiments, the individual has an initial body mass index $\geq 25 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

25 In some embodiments, the medicament for weight management is used in combination with phentermine.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for a disorder related to 5-HT_{2C} receptor activity in an individual.

30 One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for the treatment of obesity in an individual.

In some embodiments, the treatment of obesity further comprises the administration or prescription of phentermine.

35 In some embodiments, the treatment of obesity further comprises gastric electrical stimulation.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for inducing weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, 5 in the manufacture of a medicament for inducing weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual in preparation of the individual for bariatric surgery.

One aspect of the present invention pertains to the use of salts of the present invention, 10 in the manufacture of a medicament for maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual following bariatric surgery.

One aspect of the present invention pertains to the use of salts of the present invention, 15 in the manufacture of a medicament for inducing satiety in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for decreasing food intake in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, 20 in the manufacture of a medicament for decreasing hunger in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for decreasing food cravings in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, in the manufacture of a medicament for increasing intermeal interval in an individual.

One aspect of the present invention pertains to the use of salts of the present invention, 25 in the manufacture of a medicament for the treatment of a disorder selected from: schizophrenia, anxiety, depression, psychoses and alcohol addiction in an individual.

In some embodiments, the disorder is schizophrenia.

In some embodiments, the disorder is anxiety.

In some embodiments, the disorder is depression.

30 In some embodiments, the disorder is psychoses.

In some embodiments, the disorder is alcohol addiction.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of treatment of a disorder related to 5-HT_{2C} receptor activity in an individual.

35 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of treatment of obesity in an individual.

In some embodiments, the method of treatment of obesity further comprises the administration or prescription of phentermine.

In some embodiments, the method of treatment of obesity further comprises gastric electrical stimulation.

5 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of inducing weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of inducing weight loss, BMI loss, waist

10 circumference loss or body fat percentage loss in an individual in preparation of the individual for bariatric surgery.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual.

15 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of maintaining weight loss, BMI loss, waist circumference loss or body fat percentage loss in an individual following bariatric surgery.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of inducing satiety in an individual.

20 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of decreasing food intake in an individual.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of decreasing hunger in an individual.

25 One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of decreasing food cravings in an individual.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of increasing intermeal interval in an individual.

One aspect of the present invention pertains to salts and pharmaceutical compositions of the present invention, for use in a method of treatment of a disorder selected from:

30 schizophrenia, anxiety, depression, psychoses and alcohol addiction in an individual.

In some embodiments, the disorder is schizophrenia.

In some embodiments, the disorder is anxiety.

In some embodiments, the disorder is depression.

In some embodiments, the disorder is psychoses.

35 In some embodiments, the disorder is alcohol addiction.

One aspect of the present invention pertains to methods for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt, a pharmaceutical composition, or a dosage form of the present invention.

5 In some embodiments, the weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.

In some embodiments, the weight management is as an adjunct to diet and exercise.

10 In some embodiments, the individual in need of weight management is selected from: an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$; an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein the weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

15 In some embodiments, the method further comprises administering a second anti-obesity agent to the individual.

In some embodiments, the second anti-obesity agent is selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

20 In some embodiments, the method further comprises administering an anti-diabetes agent to the individual.

In some embodiments, the anti-diabetes agent is metformin.

One aspect of the present invention pertains to uses of a salt of the present invention, in the manufacture of a medicament for weight management in an individual.

25 In some embodiments, the weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.

In some embodiments, the medicament is used as an adjunct to diet and exercise.

30 In some embodiments, the individual in need of weight management is selected from: an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$; an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; and an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein the weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

35 In some embodiments, the medicament is used in combination with a second anti-obesity agent.

In some embodiments, the second anti-obesity agent is selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

5 In some embodiments, the medicament is used in combination with an anti-diabetes agent.

In some embodiments, the medicament is used in combination with an anti-diabetes agent; wherein the anti-diabetes agent is metformin.

10 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of treatment of the human or animal body by therapy.

One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management.

15 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management; wherein the weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.

20 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use as an adjunct to diet and exercise for weight management.

25 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management; wherein the individual in need of weight management is selected from: an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$; an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; and an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein the weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.

30 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management in combination with a second anti-obesity agent.

35 One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management in combination with a second anti-obesity agent selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

One aspect of the present invention pertains to salts, pharmaceutical compositions, and dosage forms of the present invention, for use in a method of weight management in combination with an anti-diabetes agent; wherein the anti-diabetes agent is metformin.

5 COMBINATION THERAPIES

The salts of the present invention can be used in combination with suitable pharmaceutical agents.

In some embodiments the salts of the present invention can be used in combination with a second anti-obesity agent. Anti-obesity agents include, for example, adrenergic reuptake 10 inhibitors, apolipoprotein-B secretion/microsomal triglyceride transfer protein inhibitors, β_3 adrenergic receptor agonists, bombesin agonists, cannabinoid 1 receptor antagonists, cholecystokinin-A agonists, ciliary neurotrophic factors, dopamine agonists, galanin antagonists, ghrelin receptor antagonists, glucagon-like peptide-1 receptor agonists, 15 glucocorticoid receptor agonists or antagonists, histamine-3 receptor antagonists or reverse agonists, human agouti-related proteins, leptin receptor agonists, lipase inhibitors, MCR-4 agonists, melanin concentrating hormone antagonists, melanocyte-stimulating hormone receptor analogs, monoamine reuptake inhibitors, neuromedin U receptor agonists, neuropeptide-Y antagonists, orexin receptor antagonists, stimulants, sympathomimetic agents, thyromimetic agents, and urocortin binding protein antagonists.

20 In some embodiments, the second anti-obesity agent is selected from: 4-methylamphetamine, 5-HTP, amfecloral, amfepentorex, amfepramone, aminorex, amphetamine, amphetaminil, atomoxetine, benfluorex, benzphetamine, bromocriptine, bupropion, cathine, cathinone, ceticlistat, chlorphentermine, ciclazindol, clobenzorex, cloforex, clominorex, clortermine, dapiclermin, dehydroepiandrosterone, dehydroepiandrosterone analogues, 25 dexmethylphenidate, dextroamphetamine, dextromethamphetamine, difemetorex, dimethylcathinone, dinitrophenol, diphenmethoxidine, ephedra, ephedrine, ethylamphetamine, etolorex, fenbutrazate, fencamfamine, fenethylline, fenproporex, fludorex, fluminorex, furfenorex, galactomannan, glucomannan, ibipinabant, indanorex, khat, L-dopa, leptin, a leptin analog, levopropylhexedrine, lisdexamphetamine, L-phenylalanine, L-tryptophan, L-tyrosine, N-30 [[trans-4-[(4,5-dihydro[1]benzothiepino[5,4-d]thiazol-2-yl)amino]cyclohexyl]methyl]methanesulfonamide, manifaxine, mazindol, mefenorex, metformin, methamphetamine, methylphenidate, naloxone, naltrexone, oleoyl-estrone, orlistat, otenabant, oxyntomodulin, P57, pemoline, peptide YY, phendimetrazine, phenethylamine, phenmetrazine, phenpentermine, phentermine, phenylpropanolamine, pipradrol, prolintane, 35 propylhexedrine, pseudoephedrine, pyrovalerone, radafaxine, reboxetine, rimonabant, setazindol, sibutramine, simmondsin, sterculia, surinabant, synephrine, taranabant, tesofensine,

topiramate, viloxazine, xylopropamine, yohimbine, zonisamide, and zylofuramine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

In some embodiments, the second anti-obesity agent is selected from: 4-methylamphetamine, amfecloral, amfepentorex, amfepramone, aminorex, amphetamine, 5 amphetaminil, atomoxetine, benfluorex, benzphetamine, bupropion, cathine, cathinone, chlorphentermine, ciclazindol, clobenzorex, cloforex, clominorex, clortermine, dexmethylphenidate, dextroamphetamine, dextromethamphetamine, difemethorex, dimethylcathinone, diphenmetoxidine, ephedra, ephedrine, ethylamphetamine, etolorex, fenbutrazate, fencamfamine, fenethylline, fenproporex, fludorex, fluminorex, furfenorex, 10 indanorex, khat, levopropylhexedrine, lisdexamphetamine, manifaxine, mazindol, mefenorex, methamphetamine, methylphenidate, pemoline, phendimetrazine, phenethylamine, phenmetrazine, phenpentermine, phentermine, phenylpropanolamine, pipradrol, prolintane, propylhexedrine, pseudoephedrine, pyrovalerone, radafaxine, reboxetine, setazindol, sibutramine, synephrine, taranabant, tesofensine, viloxazine, xylopropamine, and zylofuramine, 15 and pharmaceutically acceptable salts, solvates, and hydrates thereof.

In some embodiments, the second anti-obesity agent is selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

In some embodiments the salts of the present invention can be used in combination with 20 an anti-diabetes agent. Anti-diabetes agents include, for example, DPP-IV inhibitors, biguanides, alpha-glucosidase inhibitors, insulin analogues, sulfonylureas, SGLT2 inhibitors, meglitinides, thiazolidinediones, anti-diabetic peptide analogues, and GPR119 agonists.

In some embodiments, the anti-diabetes agent is selected from: sitagliptin, vildagliptin, saxagliptin, alogliptin, linagliptin, phenformin, metformin, buformin, proguanil, acarbose, 25 miglitol, voglibose, tolbutamide, acetohexamide, tolazamide, chlorpropamide, glipizide, glibenclamide, glimepiride, gliclazide, dapagliflozin, remigliflozin, sergliflozin, and 4-[6-(6-methanesulfonyl-2-methyl-pyridin-3-ylamino)-5-methoxy-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester.

In some embodiments, the anti-diabetes agent is a DPP-IV inhibitor selected from the 30 following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: 3(R)-amino-1-[3-(trifluoromethyl)-5,6,7,8-tetrahydro[1,2,4]triazolo[4,3-a]pyrazin-7-yl]-4-(2,4,5-trifluorophenyl)butan-1-one; 1-[2-(3-hydroxyadamant-1-ylamino)acetyl]pyrrolidine-2(S)-carbonitrile; (1S,3S,5S)-2-[2(S)-amino-2-(3-hydroxyadamantan-1-yl)acetyl]-2-azabicyclo[3.1.0]hexane-3-carbonitrile; 2-[6-[3(R)-aminopiperidin-1-yl]-3-methyl-2,4-dioxo-35 1,2,3,4-tetrahydropyrimidin-1-ylmethyl]benzonitrile; 8-[3(R)-aminopiperidin-1-yl]-7-(2-butynyl)-3-methyl-1-(4-methylquinazolin-2-ylmethyl)xanthine; 1-[N-[3(R)-pyrrolidinyl]glycyl]pyrrolidin-2(R)-yl boronic acid; 4(S)-fluoro-1-[2-[(1R,3S)-3-(1H-1,2,4-

triazol-1-ylmethyl)cyclopentylamino]acetyl]pyrrolidine-2(S)-carbonitrile; 1-[(2S,3S,11bS)-2-amino-9,10-dimethoxy-2,3,4,6,7,11b-hexahydro-1*H*-pyrido[2,1-a]isoquinolin-3-yl]-4(S)-fluoromethyl)pyrrolidin-2-one; (2S,4S)-2-cyano-4-fluoro-1-[(2-hydroxy-1,1-dimethyl)ethylamino]acetylpyrrolidine; 8-(*cis*-hexahydro-pyrrolo[3,2-b]pyrrol-1-yl)-3-methyl-7-(3-methyl-but-2-enyl)-1-(2-oxo-2-phenylethyl)-3,7-dihydro-purine-2,6-dione; 1-((3S,4S)-4-amino-1-(4-(3,3-difluoropyrrolidin-1-yl)-1,3,5-triazin-2-yl)pyrrolidin-3-yl)-5,5difluoropiperidin-2-one; (R)-2-((6-(3-aminopiperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)methyl)-4-fluorobenzonitrile; 5-{(S)-2-[2-((S)-2-cyano-pyrrolidin-1-yl)-2-oxo-ethylamino]-propyl}-5-(1*H*-tetrazol-5-yl)10,11-dihydro-5*H*-dibenzo[a,d]cycloheptene-2,8-dicarboxylic acid bis-dimethylamide; ((2S,4S)-4-(4-(3-methyl-1-phenyl-1*H*-pyrazol-5-yl)piperazin-1-yl)pyrrolidin-2-yl)(thiazolidin-3-yl)methanone; (2S,4S)-1-[2-[(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile; 6-[(3*R*)-3-amino-piperidin-1-yl]-5-(2-chloro-5-fluoro-benzyl)-1,3-dimethyl-1,5dihydro-pyrrolo[3,2-d]pyrimidine-2,4-dione; 2-({6-[(3*R*)-3-amino-3-methylpiperidin-1-yl]-1,3-dimethyl-2,4-dioxo-1,2,3,4-tetrahydro-5*H*-pyrrolo[3,2-d]pyrimidin-5-yl}methyl)-4-fluorobenzonitrile; (2*S*)-1-{{2-(5-methyl-2-phenyloxazol-4-yl)-ethylamino]-acetyl}-pyrrolidine-2-carbonitrile; (2*S*)-1-{{1,1-dimethyl-3-(4-pyridin-3-yl-imidazol-1-yl)-propylamino]-acetyl}-pyrrolidine-2-carbonitrile; (3,3-difluoropyrrolidin-1-yl)-((2S,4S)-4-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrrolidin-2-yl)methanone; (2S,4S)-1-[(2*S*)-2-amino-3,3-bis(4-fluorophenyl)propanoyl]-4-fluoropyrrolidine-2-carbonitrile; (2*S,5R*)-5-ethynyl-1-{*N*-(4-methyl-1-(4-carboxy-pyridin-2-yl)piperidin-4-yl)glycyl}pyrrolidine-2-carbonitrile; and (1*S,6R*)-3-{[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8*H*)-yl]carbonyl}-6-(2,4,5-trifluorophenyl)cyclohex-3-en-1-amine.

In some embodiments, the anti-diabetes agent is an alpha-glucosidase inhibitor selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: (2*R,3R,4R,5R*)-4-((2*R,3R,4R,5S,6R*)-5-((2*R,3R,4S,5S,6R*)-3,4-dihydroxy-6-methyl-5-((1*S,4R,5S,6S*)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-enylamino)tetrahydro-2*H*-pyran-2-yloxy)-3,4-dihydroxy-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-yloxy)-2,3,5,6-tetrahydroxyhexanal; (2*R,3R,4R,5S*)-1-(2-hydroxyethyl)-2-(hydroxymethyl)piperidine-3,4,5-triol; and (1*S,2S,3R,4S,5S*)-5-(1,3-dihydroxypropan-2-ylamino)-1-(hydroxymethyl)cyclohexane-1,2,3,4-tetraol.

In some embodiments, the anti-diabetes agent is a sulfonylurea selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: *N*-(4-(*N*-(cyclohexylcarbamoyl)sulfamoyl)phenethyl)-5-methylpyrazine-2-carboxamide); 5-chloro-*N*-(4-(*N*-(cyclohexylcarbamoyl)sulfamoyl)phenethyl)-2-methoxybenzamide; and 3-ethyl-4-methyl-*N*-(4-(*N*-(1*r,4r*)-4-methylcyclohexylcarbamoyl)sulfamoyl)phenethyl)-2-oxo-2,5-dihydro-1*H*-pyrrole-1-carboxamide.

In some embodiments, the anti-diabetes agent is an SGLT2 inhibitor selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: (2S,3R,4R,5S,6R)-2-(4-chloro-3-(4-ethoxybenzyl)phenyl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol; ethyl ((2R,3S,4S,5R,6S)-3,4,5-trihydroxy-6-(4-(4-isopropoxybenzyl)-1-isopropyl-5-methyl-1*H*-pyrazol-3-yl)oxy)tetrahydro-2*H*-pyran-2-yl)methyl carbonate; and ethyl ((2R,3S,4S,5R,6S)-3,4,5-trihydroxy-6-(2-(4-methoxybenzyl)phenoxy)tetrahydro-2*H*-pyran-2-yl)methyl carbonate.

10 In some embodiments, the anti-diabetes agent is a meglitinide selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: (S)-2-ethoxy-4-(2-(3-methyl-1-(2-(piperidin-1-yl)phenyl)butylamino)-2-oxoethyl)benzoic acid; (R)-2-((1*r*,4*R*)-4-isopropylcyclohexanecarboxamido)-3-phenylpropanoic acid; and (S)-2-benzyl-4-((3a*R*,7a*S*)-1*H*-isoindol-2(3*H*,3a*H*,4*H*,5*H*,6*H*,7*H*,7a*H*)-yl)-4-oxobutanoic acid.

15 In some embodiments, the anti-diabetes agent is a biguanide selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof: metformin, phenformin, buformin, and proguanil.

In some embodiments, the anti-diabetes agent is metformin.

In some embodiments, the anti-diabetes agent is a GPR119 agonist selected from the GPR119 agonists disclosed in the following PCT applications: WO2006083491, WO2008081204, WO2009123992, WO2010008739, WO2010029089, and WO2010149684.

20 In some embodiments, the anti-diabetes agent is 4-[6-(6-methanesulfonyl-2-methylpyridin-3-ylamino)-5-methoxy-pyrimidin-4-yl]piperidine-1-carboxylic acid isopropyl ester.

In some embodiments, the anti-diabetes agent is 5-(4-(4-(3-fluoro-4-(methylsulfonyl)phenoxy)butan-2-yl)piperidin-1-yl)-3-isopropyl-1,2,4-oxadiazole.

25 Other anti-obesity agents, and anti-diabetes agents including the agents set forth *infra*, are well known, or will be readily apparent in light of the instant disclosure, to one of ordinary skill in the art. It will be understood that the scope of combination therapy of the salts of the present invention with other anti-obesity agents and with anti-diabetes agents is not limited to those listed above, but includes in principle any combination with any pharmaceutical agent or pharmaceutical composition useful for the treatment of overweight, obese, and diabetic

30 individuals.

One aspect of the present invention pertains to salts of the present invention, characterized in that the salts is administered in conjunction with a second anti-obesity agent as described herein.

35 One aspect of the present invention pertains to salts of the present invention, characterized in that the salt is administered in conjunction with an anti-diabetes agent as described herein.

One aspect of the present invention pertains to salts of the present invention for use in combination with a second anti-obesity agent for use in weight management.

One aspect of the present invention pertains to salts of the present invention for use in combination with an anti-diabetes agent for use in weight management and the treatment of diabetes.

One aspect of the present invention pertains to methods of weight management in an individual in need thereof, comprising administering to the individual a salt of the present invention and a second anti-obesity agent wherein the salt and the second anti-obesity agent are administered to the individual simultaneously, separately, or sequentially.

One aspect of the present invention pertains to methods of weight management and treating diabetes in an individual in need thereof, comprising administering to the individual a salt of the present invention and an anti-diabetes agent wherein the salt and the anti-diabetes agent are administered to the individual simultaneously, separately, or sequentially.

One aspect of the present invention pertains to methods of weight management in an individual in need thereof, wherein the individual has been or is being treated with a second anti-obesity agent, the method comprising administering to the individual a therapeutically effective amount of a salt of the present invention.

One aspect of the present invention pertains to methods of weight management and treatment of diabetes in an individual in need thereof, wherein the individual has been or is being treated with an anti-diabetes agent, the method comprising administering to the individual a therapeutically effective amount of a salt of the present invention.

One aspect of the present invention pertains to anti-obesity agents, characterized in that the anti-obesity agent is administered in conjunction with a salt of the present invention.

One aspect of the present invention pertains to anti-diabetes agents, characterized in that the anti-diabetes agent is administered in conjunction with a salt of the present invention.

One aspect of the present invention pertains to anti-obesity agents for use in combination with a salt of the present invention for use in weight management.

One aspect of the present invention pertains to anti-diabetes agents for use in combination with a salt of the present invention for use in weight management and the treatment of diabetes.

One aspect of the present invention pertains to methods of weight management in an individual in need thereof, comprising administering to the individual an anti-obesity agent and a salt of the present invention wherein the anti-obesity agent and the salt are administered to the individual simultaneously, separately, or sequentially.

One aspect of the present invention pertains to methods of weight management and treating diabetes in an individual in need thereof, comprising administering to the individual an

anti-diabetes agent and a salt of the present invention wherein the anti-diabetes agent and the salt are administered to the individual simultaneously, separately, or sequentially.

5 One aspect of the present invention pertains to methods of weight management in an individual in need thereof, wherein the individual has been or is being treated with a salt of the present invention, the method comprising administering to the individual a therapeutically effective amount of a second anti-obesity agent.

10 One aspect of the present invention pertains to methods of weight management and treatment of diabetes in an individual in need thereof, wherein the individual has been or is being treated with a salt of the present invention, the method comprising administering to the individual a therapeutically effective amount of an anti-diabetes agent.

15 The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

EXAMPLES

20 The following examples are provided to further define the invention without, however, limiting the invention to the particulars of these examples. The compounds and salts thereof described herein, *supra* and *infra*, are named according to the CS ChemDraw Ultra Version 7.0.1, AutoNom version 2.2, or CS ChemDraw Ultra Version 9.0.7. In certain instances common names are used and it is understood that these common names would be recognized by those skilled in the art.

25 Powder X-ray Diffraction (PXRD) studies were conducted using an X'Pert PRO MPD powder diffractometer (PANalytical, Inc.; EQ0233) with a Cu source set at 45 kV and 40 mA, Cu(K α) radiation and an X'Celerator detector. Samples were placed on a PXRD sample plate either as-is or ground slightly to reduce the size of large particles or crystals. Data were collected with the samples spinning from 5 ° to 40 °2 θ . Data were analyzed by X'Pert Data Viewer software, version 1.0a, to determine crystallinity and/or crystal form, and by X'Pert 30 HighScore software, version 1.0b, to generate the tables of PXRD peaks.

Differential scanning calorimetry (DSC) studies were conducted using a TA Instruments, Q2000 (EQ1980) at heating rate 10 °C/min. The instruments were calibrated by the vendor for temperature and energy using the melting point and enthalpy of fusion of an indium standard.

35 Thermogravimetric analyses (TGA) were conducted using a TA Instruments TGA Q5000 (EQ1982) at heating rate 10 °C/min. The instrument was calibrated by the vendor using

Alumel and Nickel Curie points for the furnace temperature and a standard weight for the balance.

Dynamic moisture-sorption (DMS) studies were conducted using a dynamic moisture-sorption analyzer, VTI Corporation, SGA-100, equipment # 0228. Samples were prepared for 5 DMS analysis by placing 5 mg to 20 mg of a sample in a tared sample holder. The sample was placed on the hang-down wire of the VTI balance. A drying step was run, typically at 40 °C and 0.5-1% RH for 1-2 h. The isotherm temperature is 25 °C. Defined % RH holds typically ranged from 10% RH to 90% RH or 95% RH, with intervals of 10 to 20% RH. A % weight change smaller than 0.010% over a specified number of minutes (typically 10-20), or up to 2 h, 10 whichever occurs first, is required before continuing to the next % RH hold. The water content of the sample equilibrated as described above was determined at each % RH hold.

If saturated in water with excess solid, a deliquescent compound or salt thereof equilibrated in a closed system at a given temperature produces a % RH in that closed system that is equal to its deliquescent %RH (DRH) at that temperature. Fractional relative humidity is 15 equal to water activity (a_w) in the vapor phase and at equilibrium in a closed system, the a_w in an aqueous solution is equal to the a_w in the vapor phase above the solution (see Equation 1).

Equation 1

$$\frac{DRH}{100\%} = \frac{\%RH}{100\%} \text{ (above enclosed sat aq sol'n at equil)} = a_w(\text{vapor}) = a_w(\text{liquid})$$

A water activity meter was used to measure DRH for selected salts described herein. 20 The instrument used for this study is a Decagon Devices AquaLab 4TE water activity meter, equipment # 2169. This instrument is designed with temperature control and a small headspace above the enclosed sample to establish equilibrium between solution and vapor phases quickly. Measured a_w values at 25 °C for samples of aqueous-saturated Compound 1 salts with excess solid were multiplied by 100% to get DRH values in % RH.

25 Acquity ultra performance liquid chromatography (UPLC) from Waters was used for solubility and stoichiometry determination. Instrument number is SY-EQ 1889. UPLC was equipped with Acquity PDA detector. UPLC mobile phase solvent A was 0.1% TFA in DI-water, solvent B was 0.1% TFA in acetonitrile. The mobile phase gradient as shown in the table below:

Time (min)	Flow (mL/min)	%A	%B	Curve
	0.600	95.0	5.0	
2.00	0.600	5.0	95.0	6
2.50	0.600	5.0	95.0	6
2.75	0.600	95.0	5.0	1
5.00	0.000	95.0	5.0	11

30 Column temperature was 40 ± 5 °C. Acquity UPLC® HSS T3 1.8µm, 2.1 × 50 mm column was used.

A known amount of sample was dissolved in water and analyzed by UPLC. The weight percent of Compound 1 in the salt samples was determined by comparing the UV signal to that of a standard, Compound 1 hydrochloride salt hemihydrate, or Compound 1 free base. The percentage of Compound 1 or the percentage of the counterion determined was compared to the theoretical values to establish the stoichiometry.

Example 1: Preparation of Form I of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine Hemi-edisylate Salt (Compound 1 Hemi-edisylate Salt, Form I).

The title salt was prepared by the dropwise addition of 0.5 equivalents of aqueous 1,2-ethanedisulfonic acid dihydrate (~3.7 M) to a solution of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine free base in either acetonitrile or isopropyl acetate with vigorous stirring. Immediate precipitation was observed. The solid obtained was washed with isopropyl alcohol and allowed to dry on the filter.

The title salt was an anhydrous crystalline material with a melting onset of ~298 °C. It was non-hygroscopic by DMS analysis, picking up just 0.14% weight out to and including the 95% RH hold at 25 °C. The DRH was determined by water activity measurement of saturated aqueous solution with excess solid to be 99.7% RH at 25 °C, which indicated that the title salt was not deliquescent.

A known amount of the title salt was dissolved in water and analyzed by UPLC. The amount of Compound 1 in sample was determined to be 68.2%. This is in good agreement with the theoretical value, 67.3%. The solubility of the title salt in water was determined by UPLC to be 61 mg/mL, with a final pH of 6.

The powder X-ray diffraction pattern of the title salt is shown in Figure 1. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 2. DMS analysis of the title salt is shown in Figure 3.

Example 2: Preparation of Form I of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine Phosphate Salt (Compound 1 Phosphate Salt, Form I).

The title salt was prepared by dropwise addition of ortho-phosphoric acid (85%) (0.5-1 mole equivalent) to a solution of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine free base in isopropyl acetate or acetonitrile with vigorous stirring. Immediate precipitation was observed in all experiments. Initially amorphous material was slurried in acetone; initially crystalline material was slurried/ripened in *n*-propanol for 3 days.

The title salt was a 1:1 salt based on stoichiometry determination. The melting onset by DSC was ~208 °C. The TGA result for a crystalline sample prior to the *n*-propanol slurry is consistent with an anhydrous salt. It was non-hygroscopic, picking up 0.14% weight out to and including the 90% RH hold at 25 °C during DMS analysis. The title salt was non-deliquescent;

the DRH by water activity measurement of a saturated solution in water with excess solid was 100% RH at 25 °C.

5 A known amount of the title salt was dissolved in water and analyzed by UPLC. The amount of Compound 1 in the sample was 64.9%, slightly lower but in good agreement with the theoretical amount, 66.6%. The solubility of the title salt, was 29.4 mg/mL, with a final pH of 4.6.

The powder X-ray diffraction pattern of the title salt is shown in Figure 4. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 5. DMS analysis of the title salt is shown in Figure 6.

10

Example 3: Preparation of Form I of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine Citrate Salt Hemihydrate (Compound 1 Citrate Salt Hemihydrate, Form I).

15 The title salt was prepared by dropwise addition of 1 mole equivalent of citric acid in hot MeOH to a solution of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine in isopropyl acetate. Precipitation occurred spontaneously. Attempts to prepare a hemicitrate salt resulted in oily products.

TGA data for the title salt, showed that it was solvated. The mass loss matches closely with a hemihydrate (observed 2.6%, theoretical 2.3%). The onset of dehydration is near 80 °C for the scan rate, 10 °C/min.

20

The title salt lost only a small amount of its water of hydration during the drying step at 40 °C and ~1% RH for 1 h. It was not hygroscopic, picking up just 0.50% out to and including the 90% RH hold at 25 °C, and not deliquescent. The DRH determined by water activity measurement of a saturated aqueous solution with excess solid was 100% RH at 25 °C.

25

A known amount of the title salt was dissolved in water and analyzed by UPLC. The amount of Compound 1 in the sample was 53.2%. This is slightly higher than, but in fair agreement with the theoretical amount for a 1:1 citrate hemihydrate salt, 49.3%. The solubility in water was determined to be 33.9 mg/mL of the salt at a pH of 3.75.

30

The powder X-ray diffraction pattern of the title salt is shown in Figure 7. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 8. DMS analysis of the title salt is shown in Figure 9.

Example 4: Preparation of Form I of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine Hemi-oxalate Salt (Compound 1 Hemi-oxalate Salt, Form I).

35

The title salt was prepared by dropwise addition of 1 mole equivalent of oxalic acid as a solid or as a solution in MeOH (~2.5 M) to a solution of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine in isopropyl acetate. In each case the same crystal form was produced.

The title salt was anhydrous and displayed a melting onset temperature about 212 °C with weight loss by TGA starting just prior to the melting onset. The title salt was determined to be non-hygroscopic and non-deliquescent at 25 °C. The DRH determined by water activity measurement of a saturated aqueous sample with excess solid was 100% RH (non-deliquescent) 5 at 25 °C.

A known amount of the title salt was dissolved in water and analyzed by UPLC. The amount of Compound 1 in the sample was 74.9%. The solid was slurried in cyclohexane and then analyzed a second time. The amount of Compound 1 in the second sample was 82.5%. The theoretical value for a hemi-oxalate salt of Compound 1 is 81.2%.

10 The solubility of the title salt, in water was determined by UPLC to be 23.3 mg/mL with a final pH 4.6.

The powder X-ray diffraction pattern of the title salt is shown in Figure 10. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 11. DMS analysis of the title salt is shown in Figure 12.

15 **Example 5: Preparation of Form I of (R)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine Succinate Salt (Compound 1 Succinate Salt, Form I).**

The title salt was prepared by the addition of succinic acid (0.5-1 eq.) in hot EtOH to a solution of (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine in isopropyl acetate. After overnight stirring, a solid was recovered by suction filtration and washed in isopropyl 20 acetate.

The title salt showed a melting onset by DSC of 179.1 °C. TGA showed no residual solvent, but did show apparent loss of the salt or component thereof prior to the melting onset.

25 The title salt was found to be non-hygroscopic by DMS analysis, picking up 0.07% weight out to and including the 90% RH hold at 25 °C. It was non-deliquescent; DRH was measured by water activity determination for a saturated aqueous solution with excess solid to be 100.0% RH at 25 °C.

A known amount of the title salt was dissolved in water and analyzed by UPLC. The amount of Compound 1 in the sample was 65-69%. This is slightly higher than theoretical, 62.4%, for a 1:1 salt, but much lower than the theoretical value for a hemisuccinate salt, 76.8%. 30 Aqueous solubility of the title salt was 27.9 mg/mL, with a final pH 4.7.

The powder X-ray diffraction pattern of the title salt is shown in Figure 13. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 14. DMS analysis of the title salt is shown in Figure 15.

35 **Example 6: Preparation of Form I of (R)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine Oxoglutarate Salt (Compound 1 Oxoglutarate Salt, Form I).**

The title salt was prepared by addition of one equivalent of α -oxo-glutaric acid to a solution of (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine in ethyl acetate at 60

°C. α -Oxo-glutaric acid in ethyl acetate at 60 °C was added dropwise with vigorous stirring. Precipitation occurred immediately and the suspension was allowed to cool and stir overnight. The resulting solid was recovered by filtration and air-dried in a fume hood overnight.

5 A known amount of Compound 1 oxoglutarate salt was dissolved in methanol and analyzed by UPLC. The percentage of Compound 1 in the salt sample was determined to be 59.7%. This is slightly higher than but in good agreement with the theoretical percentage of Compound 1 in an anhydrous Compound 1 oxoglutarate salt (57.3%).

The aqueous solubility of the title salt was determined by UPLC to be > 65.1 mg/mL, with a final pH of 3.19.

10 The powder X-ray diffraction pattern of the title salt is shown in Figure 16. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 17. DMS analysis of the title salt is shown in Figure 18.

15 **Example 7: Preparation of Form I of (*R*)-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine Oxoglutarate Salt Solvate (Compound 1 Oxoglutarate Salt Solvate, Form I).**

The title salt was prepared by addition of a molar equivalent of α -oxo-glutaric acid to a solution of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine in acetonitrile at 60 °C. α -Oxo-glutaric acid in acetonitrile at 60 °C was added dropwise with vigorous stirring. Precipitation occurred immediately and the suspension was allowed to cool and stir overnight.

20 The resulting solid was recovered by filtration and air-dried in a fume hood overnight.

A known amount of Compound 1 oxoglutarate salt solvate was dissolved in methanol and analyzed by UPLC. The percentage of Compound 1 in the salt sample was determined to be 60.1%. This is slightly higher than the theoretical percentage Compound 1 in a solvate of Compound 1 oxoglutarate salt (54.%).

25 The aqueous solubility of Compound 1 oxoglutarate solvate was determined by UPLC to be >68.9 mg/mL, with a final pH of 3.21.

The powder X-ray diffraction pattern of the title salt is shown in Figure 19. Thermal analysis (TGA and DSC) of the title salt is shown in Figure 20. DMS analysis of the title salt is shown in Figure 21.

30 Those skilled in the art will recognize that various modifications, additions, substitutions, and variations to the illustrative examples set forth herein can be made without departing from the spirit of the invention and are, therefore, considered within the scope of the invention.

What is claimed is:

1. A salt selected from:
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt;
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate salt;
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt;
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate salt;
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine succinate salt; and
 - (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt; andpharmaceutically acceptable salts, solvates and hydrates thereof.
2. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-edisylate salt.
3. The salt according to claim 2, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 23.12° , about 6.00° , and about 19.70° .
4. The salt according to claim 2, having an X-ray powder diffraction pattern substantially as shown in Figure 1.
5. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine phosphate salt.
6. The salt according to claim 5, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 19.27° , about 25.06° , and about 25.77° .
7. The salt according to claim 5, having an X-ray powder diffraction pattern substantially as shown in Figure 4.
8. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine citrate salt hemihydrate.
9. The salt according to claim 8, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 11.93° , about 18.64° , and about 24.52° .
10. The salt according to claim 8, having an X-ray powder diffraction pattern substantially as shown in Figure 7.

11. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hemi-oxalate salt.
12. The salt according to claim 11, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 6.34° , about 21.69° , and about 31.85° .
13. The salt according to claim 11, having an X-ray powder diffraction pattern substantially as shown in Figure 10.
14. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine succinate salt.
15. The salt according to claim 14, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 27.62° , about 20.65° , and about 15.64° .
16. The salt according to claim 14, having an X-ray powder diffraction pattern substantially as shown in Figure 13.
17. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt.
18. The salt according to claim 17, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 13.39° , about 23.57° , and about 21.22° .
19. The salt according to claim 17, having an X-ray powder diffraction pattern substantially as shown in Figure 16.
20. The salt according to claim 1, that is (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine oxoglutarate salt solvate.
21. The salt according to claim 20, having an X-ray powder diffraction pattern comprising peaks, in terms of 2θ , at about 26.02° , about 11.53° , and about 22.65° .
22. The salt according to claim 20, having an X-ray powder diffraction pattern substantially as shown in Figure 19.

23. A pharmaceutical composition comprising a salt according to any one of claims 1 to 22, and a pharmaceutically acceptable carrier.
24. A process for preparing a pharmaceutical composition comprising admixing a salt according to any one of claims 1 to 22, and a pharmaceutically acceptable carrier.
25. A dosage form comprising a therapeutically effective amount of a salt selected from: a pharmaceutically acceptable salt of (*R*)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine and pharmaceutically acceptable solvates and hydrates thereof, wherein said dosage form is a non-hygroscopic dosage form.
26. The dosage form according to claim 25, wherein said salt absorbs:
 - less than about 2% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 1% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 0.6% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 0.5% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 0.4% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 0.3% water by weight after about 2 h at about 90% RH and about 25 °C;
 - less than about 0.2% water by weight after about 2 h at about 90% RH and about 25 °C; or
 - less than about 0.1% water by weight after about 2 h at about 90% RH and about 25 °C.
27. A dosage form comprising a therapeutically effective amount of a salt according to any one of claims 1 to 22.
28. A method for weight management, comprising administering to an individual in need thereof, a therapeutically effective amount of a salt according to any one of claims 1 to 22, a pharmaceutical composition according to claim 23, or a dosage form according to any one of claims 25 to 27.

29. The method according to claim 28, wherein said weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.
30. The method according to claim 28 or 29, as an adjunct to diet and exercise.
31. The method according to any one of claims 28 to 30, wherein said individual in need of weight management is selected from:
 - an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$;
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; and
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein said weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.
32. The method according to any one of claims 28 to 31, further comprising administering a second anti-obesity agent to said individual.
33. The method according to claim 32, wherein said second anti-obesity agent is selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.
34. The method according to any one of claims 28 to 33, further comprising administering an anti-diabetes agent to said individual.
35. The method according to claim 34, wherein said anti-diabetes agent is metformin.
36. Use of a salt according to any one of claims 1 to 22, in the manufacture of a medicament for weight management in an individual.
37. The use according to claim 36, wherein said weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.
38. The use according to claim 36 or 37, wherein said medicament is used as an adjunct to diet and exercise.

39. The use according to any one of claims 36 to 38, wherein said individual in need of weight management is selected from:
 - an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$;
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; and
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein said weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.
40. The use according to any one of claims 36 to 39, wherein said medicament is used in combination with a second anti-obesity agent.
41. The use according to claim 40, wherein said second anti-obesity agent is selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.
42. The use according to any one of claims 36 to 41, wherein said medicament is used in combination with an anti-diabetes agent; wherein said anti-diabetes agent is metformin.
43. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of treatment of the human or animal body by therapy.
44. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of weight management.
45. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, for use in a method of weight management; wherein said weight management comprises one or more of: weight loss, maintenance of weight loss, decreased food consumption, increasing meal-related satiety, reducing pre-meal hunger, and reducing intra-meal food intake.

46. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use as an adjunct to diet and exercise for weight management.
47. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of weight management; wherein said individual in need of weight management is selected from:
 - an obese patient with an initial body mass index $\geq 30 \text{ kg/m}^2$;
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; and
 - an overweight patient with an initial body mass index $\geq 27 \text{ kg/m}^2$ in the presence of at least one weight related comorbid condition; wherein said weight related co-morbid condition is selected from: hypertension, dyslipidemia, cardiovascular disease, glucose intolerance, and sleep apnea.
48. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of weight management in combination with a second anti-obesity agent.
49. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of weight management in combination with a second anti-obesity agent selected from: chlorphentermine, clortermine, phenpentermine, and phentermine, and pharmaceutically acceptable salts, solvates, and hydrates thereof.
50. The salt according to any one of claims 1 to 22, or the pharmaceutical composition according to claim 23, or the dosage form according to any one of claims 25 to 27, for use in a method of weight management in combination with an anti-diabetes agent; wherein said anti-diabetes agent is metformin.