

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2020218908 B2**

(54) Title
Benzothiadiazepine compounds and their use as bile acid modulators

(51) International Patent Classification(s)
C07D 285/36 (2006.01) **A61P 3/06** (2006.01)
A61K 31/554 (2006.01) **A61P 3/08** (2006.01)
A61P 1/16 (2006.01) **A61P 9/00** (2006.01)

(21) Application No: **2020218908** (22) Date of Filing: **2020.02.06**

(87) WIPO No: **WO20/161217**

(30) Priority Data

(31) Number	(32) Date	(33) Country
201911004690	2019.02.06	IN
1950464-6	2019.04.12	SE
201911049981	2019.12.04	IN

(43) Publication Date: **2020.08.13**

(44) Accepted Journal Date: **2025.01.23**

(71) Applicant(s)
Albireo AB

(72) Inventor(s)
GILLBERG, Per-Göran;MATTSSON, Jan;STARKE, Ingemar;KULKARNI, Santosh S.

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 03/022286 A1
WO 03/091232 A2
WO 98/38182 A1



(51) International Patent Classification:

A61P 1/16 (2006.01) A61P 9/00 (2006.01)
 A61P 3/06 (2006.01) C07D 285/36 (2006.01)
 A61P 3/08 (2006.01) A61K 31/554 (2006.01)

(21) International Application Number:

PCT/EP2020/052942

(22) International Filing Date:

06 February 2020 (06.02.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

201911004690 06 February 2019 (06.02.2019) IN
 1950464-6 12 April 2019 (12.04.2019) SE
 201911049981 04 December 2019 (04.12.2019) IN

(71) Applicant: **ALBIREO AB** [SE/SE]; Arvid Wallgrens
 backe 20, 413 46 Göteborg (SE).

(72) Inventors: **GILLBERG, Per-Göran**; Södermalmsgatan
 22, 431 69 Mölndal (SE). **MATTSSON, Jan**; Kullaviksvä-
 gen 137, 429 33 Kullavik (SE). **STARKE, Ingemar**;
 Frigångsgatan 8C, 413 01 Göteborg (SE). **KULKARNI,**
Santosh S.; A206, Temple Trees Apartment, J P Nagar 6th
 Phase, Kanakapura Main Road, Bangalore 560078 (IN).

(74) Agent: **NOVITAS PATENT AB**; P.O.Box 55557, 102 04
 Stockholm (SE).


(81) Designated States (unless otherwise indicated, for every
 kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
 DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
 HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
 KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR,
 TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
 kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
 UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
 MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
 TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
 KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: BENZOTHIADIAZEPINE COMPOUNDS AND THEIR USE AS BILE ACID MODULATORS



(57) Abstract: The invention relates to 1,2,5-benzothiadiazepine derivatives of formula (I). These compounds are bile acid modulators having apical sodium-dependent bile acid transporter (ASBT) and/or liver bile acid transport (LBAT) inhibitory activity. The invention also relates to pharmaceutical compositions comprising these compounds and to the use of these compounds in the treatment of cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

(I)

BENZOTHIADIAZEPINE COMPOUNDS AND THEIR USE AS BILE ACID MODULATORS

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 This application claims priority to Indian Application No. 201911004690, filed February 6, 2019; Swedish Application No. 1950464-6, filed April 12, 2019; and Indian Application No. 201911049981, filed December 4, 2019, the disclosures of which are incorporated herein by reference in their entireties.

10 TECHNICAL FIELD

The invention relates to 1,2,5-benzothiadiazepine derivatives of formula (I). These compounds are bile acid modulators having apical sodium-dependent bile acid transporter (ASBT) and/or liver bile acid transport (LBAT) inhibitory activity. The invention also relates to pharmaceutical compositions
15 comprising these compounds and to the use of these compounds in the treatment of cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

BACKGROUND

20 Bile acids are physiological detergents that play an important role in the intestinal absorption and transport of lipids, nutrients and vitamins. They are also signaling molecules that activate nuclear receptors and cell signaling pathways that regulate lipid, glucose and energy metabolism. Bile acids are steroid acids that are synthesized from cholesterol in the liver and stored in the gallbladder as
25 mixed micelles. During digestion, the duodenum triggers the release of hormones that cause the gallbladder to contract, thereby releasing bile acids in the small intestine where they enable absorption of fat-soluble vitamins and cholesterol. When they reach the ileum, bile acids are reabsorbed from the intestine and secreted into portal blood to return to the liver via the portal venous circulation. Over 90% of the bile acids are thus recycled and returned to the liver. These bile
30 acids are then transported across the sinusoidal membrane of hepatocytes and re-secreted across the canalicular membrane into bile. In this first pass, 75-90% of bile acids are taken up by hepatocytes, completing one round of enterohepatic circulation. The fraction of bile acids that escapes being cleared in the liver enters the systemic circulation where the free bile acids are filtered
35 by the renal glomerulus, efficiently reclaimed in the proximal tubules and exported back into the systemic circulation. Interestingly, most of the bile acids secreted across the canalicular membrane

into bile are derived from the recirculating pool with less than 10% coming from new *de novo* hepatic synthesis. The small fraction of bile acids that is not reabsorbed in the ileum reaches the colon.

Within the intestinal lumen, the primary bile acids are transformed into secondary bile acids under the action of intestinal bacteria, mainly by single or dual dehydroxylation reactions of the steroid nucleus. The bile acids that escape intestinal absorption are thereafter excreted into the faeces.

Overall, the efficient transport system helps maintain a constant bile acid pool, ensuring sufficiently high levels of conjugated bile acids in the intestine to promote lipid absorption as well as reduce the small intestinal bacterial load. The system also minimizes fecal and urinary bile acid loss and protects the intestinal and hepatobiliary compartments by eliminating potentially cytotoxic detergents (as reviewed by Kusters and Karpen (*Xenobiotica* 2008, vol. 38, p. 1043-1071); by Chiang (*J. Lipid Res.* 2009, vol. 50, p. 1955-1966); and by Dawson (*Handb. Exp. Pharmacol.* 2011, vol. 201, p. 169-203)).

The regulation of the bile acid pool size has been found to play a key role in cholesterol homeostasis by hepatic conversion of cholesterol to bile acid, which represents a major route for elimination of cholesterol from the body. The liver plays an essential role in removing endogenous and xenobiotic compounds from the body. The normal hepatobiliary secretion and enterohepatic circulation are required for the elimination of endogenous compounds such as cholesterol and bilirubin and their metabolites from the body, thereby maintaining lipid and bile acid homeostasis. (Kusters and Karpen, *Xenobiotica* 2008, vol. 38, p. 1043-1071).

The reabsorption of bile acids in the ileum may be inhibited by apical sodium-dependent bile acid transporter (ASBT) inhibitor compounds. Inhibition of bile acid reabsorption has been reported useful in the treatment of several diseases, including dyslipidemia, diabetes, obesity, constipation, cholestatic liver diseases, non-alcoholic steatohepatitis and other hepatic diseases. A number of ASBT inhibitor compounds has been disclosed over the past decades, see e.g. WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/03818, WO 98/07449, WO 98/40375, WO 99/35135, WO 99/64409, WO 99/64410, WO 00/47568, WO 00/61568, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 01/66533, WO 01/68096, WO 02/32428, WO 02/50051, WO 03/020710, WO 03/022286, WO 03/022825, WO 03/022830, WO 03/061663, WO 03/091232, WO 03/106482, WO 2004/006899, WO 2004/076430, WO 2007/009655, WO 2007/009656, WO 2011/137135, DE 19825804, EP 864582, EP 489423, EP 549967, EP 573848, EP 624593, EP 624594, EP 624595, EP 624596, EP 0864582, EP 1173205 and EP 1535913.


Despite the number of ASBT inhibitor compounds that have been previously reported, there is a need for additional bile acid modulating compounds that have an optimized profile with respect to potency, selectivity and bioavailability.

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

DETAILED DESCRIPTION OF THE INVENTION

It has been discovered that certain 1,2,5-benzothiadiazepine derivatives are potent inhibitors of the apical sodium-dependent bile acid transporter (ASBT) and/or the liver bile acid transporter (LBAT), and may be useful for treating diseases wherein inhibition of the bile acid circulation is desirable.

In a first aspect, the invention relates to a compound of formula (I)



wherein

R^1 and R^2 are each independently C_{1-4} alkyl;

R^3 is independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, cyano, nitro, amino, N -(C_{1-4} alkyl)amino, N,N -di(C_{1-4} alkyl)amino and N -(aryl- C_{1-4} alkyl)amino;

n is an integer 1, 2 or 3;

R^4 is selected from the group consisting of hydrogen, halogen, hydroxy, cyano, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyloxy, C_{1-4} alkylthio, C_{3-6} cycloalkylthio, amino, N -(C_{1-4} alkyl)-amino and N,N -di(C_{1-4} alkyl)amino;

R^{5A} , R^{5B} , R^{5C} and R^{5D} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, amino and C_{1-4} alkyl; and

R^6 is selected from the group consisting of hydrogen and C_{1-4} alkyl;
or a pharmaceutically acceptable salt thereof.

In some embodiments, R^1 is *n*-butyl.

In some embodiments, R^2 is C_{2-4} alkyl. In a preferred embodiment, R^2 is methyl. In another preferred embodiment, R^2 is ethyl. In another preferred embodiment, R^2 is *n*-propyl. In yet another preferred embodiment, R^2 is *n*-butyl.

5

In some embodiments, R^3 is independently selected from the group consisting of hydrogen, halogen, hydroxy, amino, cyano, C_{1-4} haloalkyl, C_{1-4} alkoxy and C_{1-4} haloalkoxy. In another embodiment, R^3 is hydrogen. In a preferred embodiment, R^3 is independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, cyano, trifluoromethyl, methoxy and trifluoromethoxy.

10

In a preferred embodiment, *n* is 1, i.e. the phenyl-ring is substituted with only one substituent R^3 . In another preferred embodiment, R^3 is in the para-position.

In some embodiments, R^4 is selected from the group consisting of halogen, hydroxy, cyano, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, amino, *N*-(C_{1-4} alkyl)amino and *N,N*-di(C_{1-4} alkyl)amino. In a preferred embodiment, R^4 is selected from the group consisting of fluoro, chloro, bromo, hydroxy, cyano, methyl, methoxy, ethoxy, methylthio, ethylthio, amino, methylamino and dimethylamino. In another embodiment, R^4 is selected from the group consisting of fluoro, chloro, bromo, methoxy, ethoxy, methylthio, ethylthio and dimethylamino. In another preferred embodiment, R^4 is selected from the group consisting of chloro, bromo, methylthio and dimethylamino.


20

In some embodiments, R^{5A} and R^{5B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, amino and methyl. In some embodiments, R^{5A} and R^{5B} are each independently hydrogen or hydroxy. In another embodiment, R^{5A} is hydrogen or hydroxy and R^{5B} is hydrogen. In some embodiments, R^{5C} and R^{5D} are each independently hydrogen or methyl. In another embodiment, R^{5C} is hydrogen or methyl and R^{5D} is hydrogen.

25

In one embodiment, R^6 is hydrogen. In another embodiment, R^6 is methyl.

In a preferred embodiment, the compound of formula (I) is a compound of formula (I-a)



wherein

R^2 is C_{1-4} alkyl;

5 R^3 is independently selected from the group consisting of hydrogen, halogen, hydroxy, cyano, C_{1-4} haloalkyl, C_{1-4} alkoxy and C_{1-4} haloalkoxy;

n is an integer 1 or 2;


R^4 is selected from the group consisting of halogen, hydroxy, cyano, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, amino, N -(C_{1-4} alkyl)amino and N,N -di(C_{1-4} alkyl)amino;

10 R^{5A} is selected from the group consisting of hydrogen, halogen, hydroxy, amino and C_{1-4} alkyl; and

R^{5C} is hydrogen or C_{1-4} alkyl;

or a pharmaceutically acceptable salt thereof.

15 In another preferred embodiment, the compound of formula (I) is a compound of formula (I-b)



wherein

R^2 is methyl, ethyl, n -propyl or n -butyl;

20 R^3 is selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, cyano, trifluoromethyl, methoxy, and trifluoromethoxy;

R^4 is selected from the group consisting of selected from the group consisting of fluoro, chloro, bromo, hydroxy, cyano, methoxy, ethoxy, methylthio, ethylthio and dimethylamino; and

R^{5A} is selected from the group consisting of hydrogen, halogen, hydroxy, amino and methyl;

or a pharmaceutically acceptable salt thereof.

In another preferred embodiment, the compound of formula (I) is a compound of formula (I-b) as defined above, yet wherein

- 5 R^3 is selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy and methoxy; and

R^4 is selected from the group consisting of fluoro, chloro, bromo, methoxy, ethoxy, methylthio, ethylthio and dimethylamino.

- 10 Preferred compounds of the invention are compounds of formula (I-b), as defined above, wherein R^2 to R^{5A} are as indicated in Table 1 below, or a pharmaceutically acceptable salt thereof:

Table 1

R^2	R^3	R^4	R^{5A}
CH ₃	H	SCH ₃	H
CH ₃	F	SCH ₃	H
CH ₃	OCH ₃	SCH ₃	H
CH ₃	OH	SCH ₃	H
CH ₃	Cl	SCH ₃	H
CH ₃	H	SCH ₃	OH
CH ₃	F	SCH ₃	OH
CH ₃	OCH ₃	SCH ₃	OH
CH ₃	OH	SCH ₃	OH
CH ₃	Cl	SCH ₃	OH
CH ₃	H	N(CH ₃) ₂	H
CH ₃	F	N(CH ₃) ₂	H
CH ₃	OCH ₃	N(CH ₃) ₂	H
CH ₃	OH	N(CH ₃) ₂	H
CH ₃	Cl	N(CH ₃) ₂	H
CH ₃	H	N(CH ₃) ₂	OH
CH ₃	F	N(CH ₃) ₂	OH
CH ₃	OCH ₃	N(CH ₃) ₂	OH
CH ₃	OH	N(CH ₃) ₂	OH
CH ₃	Cl	N(CH ₃) ₂	OH

R ²	R ³	R ⁴	R ^{5A}
CH ₂ CH ₃	H	SCH ₃	H
CH ₂ CH ₃	F	SCH ₃	H
CH ₂ CH ₃	OCH ₃	SCH ₃	H
CH ₂ CH ₃	OH	SCH ₃	H
CH ₂ CH ₃	Cl	SCH ₃	H
CH ₂ CH ₃	H	SCH ₃	OH
CH ₂ CH ₃	F	SCH ₃	OH
CH ₂ CH ₃	OCH ₃	SCH ₃	OH
CH ₂ CH ₃	OH	SCH ₃	OH
CH ₂ CH ₃	Cl	SCH ₃	OH
CH ₂ CH ₃	H	N(CH ₃) ₂	H
CH ₂ CH ₃	F	N(CH ₃) ₂	H
CH ₂ CH ₃	OCH ₃	N(CH ₃) ₂	H
CH ₂ CH ₃	OH	N(CH ₃) ₂	H
CH ₂ CH ₃	Cl	N(CH ₃) ₂	H
CH ₂ CH ₃	H	N(CH ₃) ₂	OH
CH ₂ CH ₃	F	N(CH ₃) ₂	OH
CH ₂ CH ₃	OCH ₃	N(CH ₃) ₂	OH
CH ₂ CH ₃	OH	N(CH ₃) ₂	OH
CH ₂ CH ₃	Cl	N(CH ₃) ₂	OH
CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	H
CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	H
CH ₂ CH ₂ CH ₂ CH ₃	OCH ₃	SCH ₃	H
CH ₂ CH ₂ CH ₂ CH ₃	OH	SCH ₃	H
CH ₂ CH ₂ CH ₂ CH ₃	Cl	SCH ₃	H
CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	OH
CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	OH
CH ₂ CH ₂ CH ₂ CH ₃	OCH ₃	SCH ₃	OH
CH ₂ CH ₂ CH ₂ CH ₃	OH	SCH ₃	OH
CH ₂ CH ₂ CH ₂ CH ₃	Cl	SCH ₃	OH
CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	H
CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	H

R ²	R ³	R ⁴	R ^{5A}
CH ₂ CH ₂ CH ₂ CH ₃	OCH ₃	N(CH ₃) ₂	H
CH ₂ CH ₂ CH ₂ CH ₃	OH	N(CH ₃) ₂	H
CH ₂ CH ₂ CH ₂ CH ₃	Cl	N(CH ₃) ₂	H
CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	OH
CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	OH
CH ₂ CH ₂ CH ₂ CH ₃	OCH ₃	N(CH ₃) ₂	OH
CH ₂ CH ₂ CH ₂ CH ₃	OH	N(CH ₃) ₂	OH
CH ₂ CH ₂ CH ₂ CH ₃	Cl	N(CH ₃) ₂	OH

In a particular embodiment, the compound of formula (I) is selected from the group consisting of:

3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydrobenzo-1,2,5-thiadiazepin-8-yl)oxy)propanoic acid;

5 3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

(S)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

10 (R)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

(S)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

15 (R)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;

20 (S)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;

(R)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;

3-((3,3-dibutyl-7-chloro-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

3-((3,3-dibutyl-5-(4-hydroxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

3-((3-Butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

5 (S)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

(R)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

10 O-(3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

(S)-O-((R)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

(R)-O-((R)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

15 (S)-O-((S)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine; and

(R)-O-((S)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

or a pharmaceutically acceptable salt thereof.

20

As used herein, the term “halo” refers to fluoro, chloro, bromo and iodo.

As used herein, the term “C₁₋₆ alkyl” refers to a straight or branched alkyl group having from 1 to 6 carbon atoms, and the term “C₁₋₄ alkyl” refers to a straight or branched alkyl group having from 1 to 4
25 carbon atoms. Examples of C₁₋₄ alkyl include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

As used herein, the term “C₁₋₄ haloalkyl” refers to a straight or branched C₁₋₄ alkyl group, as defined herein, wherein one or more hydrogen atoms have been replaced with halogen. Examples of C₁₋₄
30 haloalkyl include chloromethyl, fluoroethyl and trifluoromethyl.

As used herein, the terms “C₁₋₄ alkoxy” and “C₁₋₄ alkylthio” refer to a straight or branched C₁₋₄ alkyl group attached to the remainder of the molecule through an oxygen or sulphur atom, respectively.

As used herein, the term "C₃₋₆ cycloalkyl" refers to a monocyclic saturated hydrocarbon ring having from 3 to 6 carbon atoms. Examples of C₃₋₆ cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "aryl" denotes an aromatic monocyclic ring composed of 6 carbon atoms or an aromatic bicyclic ring system composed of 10 carbon atoms. Examples of aryl include phenyl, naphthyl and azulenyl.

The term "amino" refers to an -NH₂ group. As used herein, the terms "*N*-(C₁₋₄ alkyl)amino" and "*N,N*-di(C₁₋₄ alkyl)amino" refer to an amino group wherein one or both hydrogen atom(s), respectively, are replaced with a straight or branched C₁₋₄ alkyl group. Examples of *N*-(C₁₋₄ alkyl)amino include methylamino, ethylamino and tert-butylamino, and examples of *N,N*-di-(C₁₋₄ alkyl)amino include dimethylamino and diethylamino.

As used herein, the term "*N*-(aryl-C₁₋₄ alkyl)amino" refers to an amino group wherein a hydrogen atom is replaced with an aryl-C₁₋₄ alkyl group. Examples of *N*-(aryl-C₁₋₄ alkyl)amino include benzylamino and phenylethylamino.

In a second aspect of the invention, there is provided a pharmaceutical composition comprising a therapeutically effective amount of a compound according to the first aspect, and one or more pharmaceutically acceptable excipients.

As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions and/or dosage forms that are suitable for human pharmaceutical use and that are generally safe, non-toxic and neither biologically nor otherwise undesirable.

As used herein, the term "about" refers to a value or parameter herein that includes (and describes) embodiments that are directed to that value or parameter per se. For example, description referring to "about 20" includes description of "20." Numeric ranges are inclusive of the numbers defining the range. Generally speaking, the term "about" refers to the indicated value of the variable and to all values of the variable that are within the experimental error of the indicated value (e.g., within the 95% confidence interval for the mean) or within 10 percent of the indicated value, whichever is greater.

The 1,2,5-benzothiadiazepine compounds of formula (I), or pharmaceutically acceptable salts thereof, are inhibitors of the apical sodium-dependent bile acid transporter (ASBT inhibitors), of the liver bile acid transporter (LBAT inhibitors), or of both the apical sodium-dependent bile acid and liver bile acid transporters (dual ASBT/LBAT inhibitors). They are therefore useful in the treatment or prevention of conditions, disorders and diseases wherein inhibition of bile acid circulation is desirable, such as cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

In a third aspect of the invention, there is therefore provided a method of treatment or prevention of a condition, disease or disorder mediated by inhibition of bile acid circulation in a subject in need thereof comprising administering to the subject an effective amount of a compound according to the first aspect or the pharmaceutical composition of the second aspect.

In a fourth aspect of the invention, there is therefore provided a use of a compound according to the first aspect or the pharmaceutical composition of the second aspect in the preparation of a medicament for treatment or prevention of a condition, disease or disorder mediated by inhibition of bile acid circulation.

Cardiovascular diseases and disorders of fatty acid metabolism and glucose utilization include, but are not limited to, hypercholesterolemia; disorders of fatty acid metabolism; type 1 and type 2 diabetes mellitus; complications of diabetes, including cataracts, micro- and macrovascular diseases, retinopathy, neuropathy, nephropathy and delayed wound healing, tissue ischaemia, diabetic foot, arteriosclerosis, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, stroke, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders and vascular restenosis; diabetes-related diseases such as insulin resistance (impaired glucose homeostasis), hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, dyslipidemia, hyperlipidemia including hypertriglyceridemia, metabolic syndrome (syndrome X), atherosclerosis and hypertension; and for increasing high density lipoprotein levels.

Gastrointestinal diseases and disorders include constipation (including chronic constipation, functional constipation, chronic idiopathic constipation (CIC), intermittent/sporadic constipation, constipation secondary to diabetes mellitus, constipation secondary to stroke, constipation secondary to chronic kidney disease, constipation secondary to multiple sclerosis, constipation secondary to Parkinson's disease, constipation secondary to systemic sclerosis, drug induced

constipation, irritable bowel syndrome with constipation (IBS-C), irritable bowel syndrome mixed (IBS-M), pediatric functional constipation and opioid induced constipation); Crohn's disease; primary bile acid malabsorption; irritable bowel syndrome (IBS); inflammatory bowel disease (IBD); ileal inflammation; and reflux disease and complications thereof, such as Barrett's esophagus, bile reflux esophagitis and bile reflux gastritis.

A liver disease as defined herein is any disease in the liver and in organs connected therewith, such as the pancreas, portal vein, the liver parenchyma, the intrahepatic biliary tree, the extrahepatic biliary tree, and the gall bladder. In some cases, a liver disease a bile acid-dependent liver disease. Liver diseases and disorders include, but are not limited to an inherited metabolic disorder of the liver; inborn errors of bile acid synthesis; congenital bile duct anomalies; biliary atresia; post-Kasai biliary atresia; post-liver transplantation biliary atresia; neonatal hepatitis; neonatal cholestasis; hereditary forms of cholestasis; cerebrotendinous xanthomatosis; a secondary defect of BA synthesis; Zellweger's syndrome; cystic fibrosis-associated liver disease; alpha1-antitrypsin deficiency; Alagilles syndrome (ALGS); Byler syndrome; a primary defect of bile acid (BA) synthesis;

progressive familial intrahepatic cholestasis (PFIC) including PFIC-1, PFIC-2, PFIC-3 and non-specified PFIC, post-biliary diversion PFIC and post-liver transplant PFIC; benign recurrent intrahepatic cholestasis (BRIC) including BRIC1, BRIC2 and non-specified BRIC, post-biliary diversion BRIC and post-liver transplant BRIC; autoimmune hepatitis; primary biliary cirrhosis (PBC); liver fibrosis; non-alcoholic fatty liver disease (NAFLD); non-alcoholic steatohepatitis (NASH); portal hypertension; cholestasis; Down syndrome cholestasis; drug-induced cholestasis; intrahepatic cholestasis of pregnancy (jaundice during pregnancy); intrahepatic cholestasis; extrahepatic cholestasis; parenteral nutrition associated cholestasis (PNAC); low phospholipid-associated cholestasis; lymphedema cholestasis syndrome 1 (LSC1); primary sclerosing cholangitis (PSC); immunoglobulin G4 associated cholangitis; primary biliary cholangitis; cholelithiasis (gall stones); biliary lithiasis; choledocholithiasis; gallstone pancreatitis; Caroli disease; malignancy of bile ducts; malignancy causing obstruction of the biliary tree; biliary strictures; AIDS cholangiopathy; ischemic cholangiopathy; pruritus due to cholestasis or jaundice; pancreatitis; chronic autoimmune liver disease leading to progressive cholestasis; hepatic steatosis; alcoholic hepatitis; acute fatty liver; fatty liver of pregnancy; drug-induced hepatitis; iron overload disorders; congenital bile acid synthesis defect type 1 (BAS type 1); drug-induced liver injury (DILI); hepatic fibrosis; congenital hepatic fibrosis; hepatic cirrhosis; Langerhans cell histiocytosis (LCH); neonatal ichthyosis sclerosing cholangitis (NISCH); erythropoietic protoporphyria (EPP); idiopathic adulthood ductopenia (IAD); idiopathic neonatal hepatitis (INH); non syndromic paucity of interlobular bile ducts (NS PILBD); North American Indian childhood cirrhosis (NAIC); hepatic sarcoidosis; amyloidosis; necrotizing enterocolitis; serum bile acid-caused toxicities, including cardiac rhythm disturbances (e.g., atrial fibrillation) in setting of abnormal serum bile acid profile, cardiomyopathy associated with liver cirrhosis ("cholecardia"), and skeletal muscle wasting associated with cholestatic liver disease; viral hepatitis (including hepatitis A, hepatitis B, hepatitis C, hepatitis D and hepatitis E); hepatocellular carcinoma (hepatoma); cholangiocarcinoma; bile acid-related gastrointestinal cancers; and cholestasis caused by tumours and neoplasms of the liver, of the biliary tract and of the pancreas. Compounds of formula (I), or pharmaceutically acceptable salts thereof, are also useful in the enhancement of corticosteroid therapy in liver disease.

Other diseases that may be treated or prevented by the compounds of formula (I), or pharmaceutically acceptable salts thereof, include hyperabsorption syndromes (including abetalipoproteinemia, familial hypobetalipoproteinemia (FHBL), chylomicron retention disease (CRD) and sitosterolemia); hypervitaminosis and osteopetrosis; hypertension; glomerular hyperfiltration; pruritus of renal failure; The compounds are also useful in the protection against liver- or metabolic disease-associated kidney injury.

The transport of bile acids in the human body is controlled by the action of the members of the SLC10 family of solute carrier proteins, in particular by the Na⁺-taurocholate cotransporting polypeptide (NTCP, also called liver bile acid transporter (LBAT); gene symbol *SLC10A1*), which is expressed in the sinusoidal membrane of hepatocytes, and by the apical sodium dependent bile acid transporter (ASBT, also called ileal bile acid transporter (IBAT), ISBT, ABAT or NTCP2; gene symbol *SLC10A2*), which is expressed in the apical membrane of ileal enterocytes, proximal renal tubule cells, biliary epithelium, large cholangiocytes and gallbladder epithelial cells. In the liver, bile acids are efficiently extracted from portal blood by the liver bile acid transporter (LBAT) and re-secreted across the canalicular membrane by the bile salt export pump (BSEP; gene symbol *ABCB11*).

10 The reabsorption of bile acids in the ileum is handled by the apical sodium-dependent bile acid transporter (ASBT), where it is commonly referred to as ileal bile acid transporter (IBAT). Both LBAT and ASBT function as electrogenic sodium-solute cotransporters that move two or more Na⁺ ions per molecule of solute.

15 Xenobiotics and endobiotics, including bile acids, are taken up by the liver from portal blood and secreted into bile by distinct transport proteins with individualized substrate specificities. Glycine- and taurine-conjugated bile acids exist in anionic form and are unable to cross membranes by diffusion, and thus, are completely dependent on membrane transport proteins to enter or exit the hepatocyte (Kosters and Karpen, *Xenobiotica* 2008, vol. 38, p. 1043-1071). ASBT and LBAT prefer

20 glycine- and taurine-conjugated bile salts over their unconjugated counterparts and demonstrate a higher affinity for dihydroxy bile salts than for trihydroxy bile salts. No non-bile acid substrates have been identified for ASBT yet, however, LBAT was also found to transport a variety of steroid sulfates, hormones and xenobiotics.

25 LBAT is not as thoroughly characterized as ASBT in terms of drug inhibition requirements. Dong et al. have identified FDA approved drugs that inhibit human LBAT and compared LBAT and ASBT inhibition requirements. A series of LBAT inhibition studies were performed using FDA approved drugs, in concert with iterative computational model development. Screening studies identified 27 drugs as novel LBAT inhibitors, including irbesartan (K_i = 11.9 μM) and ezetimibe (K_i = 25.0 μM). The common

30 feature pharmacophore indicated that two hydrophobes and one hydrogen bond acceptor were important for inhibition of LBAT. From 72 drugs screened in vitro, a total of 31 drugs inhibited LBAT, while 51 drugs (i.e. more than half) inhibited ASBT. Hence, while there was inhibitor overlap, ASBT unexpectedly was more permissive to drug inhibition than was LBAT, and this may be related to LBAT's possessing fewer pharmacophore features (Dong et al., *Mol. Pharm.* 2013, vol. 10, p. 1008–

35 1019).

Vaz et al. describe the identification of LBAT deficiency as a new inborn error of metabolism with a relatively mild clinical phenotype. The identification of LBAT deficiency confirms that this transporter is the main import system for conjugated bile salts into the liver, but also indicates that auxiliary
5 transporters are able to sustain the enterohepatic cycle in its absence (Vaz et al., *Hepatology* 2015, vol. 61, p. 260-267). These findings support the hypothesis that LBAT inhibition is a safe mechanism of action, as the hepatocytes still have the possibility to take up the necessary amount of bile acids.

Liu et al. describe the identification of a new type of hypercholanemia that is associated with
10 homozygosity for the p.Ser267Phe mutation in *SLC10A1* (LBAT). The allele frequency of this mutation in gene *SLC10A1* varies in different populations, with the highest incidence occurring in Southern China (8% and 12% in Chinese Han and Dai respectively) and in Vietnam (11%). This “hidden” hypercholanemia was believed to affect 0.64% of the Southern Han, 1.44% of the Dai Chinese population, and 1.21% of the Vietnamese population. An increase in conjugated and unconjugated
15 serum BA levels in the homozygous individuals was also observed. Liu et al. suggest that this finding is most likely due to reduced BA transport from the portal circulation into hepatocytes. This supports the hypothesis that the physiological function of the enterohepatic circulation is not only to recycle bile acids but also to clear bile acids from the circulation to achieve homeostasis (Karpen and Dawson, *Hepatology* 2015, vol. 61, p. 24-27). Alternatively, the liver may synthesize increased levels
20 of bile acids to compensate for the reduced enterohepatic recirculation in the homozygous carriers. As LBAT also transports unconjugated bile acids, the increase of the unconjugated bile acids in this study was not surprising (Liu et al., *Scientific Reports* 2017, 7: 9214, p. 1-7).

LBAT has been found to be downregulated in several forms of cholestatic liver injury and cholestasis,
25 whereas ASBT has been found to be downregulated in a variety of gastrointestinal disorders such as Crohn’s disease, primary bile acid malabsorption, inflammatory bowel disease, and ileal inflammation but upregulated in cholestasis. LBAT also functions as a cellular receptor for viral entry of the hepatitis B virus (HBV) and hepatitis D virus (HDV), which in turn is the major cause of liver disease and hepatocellular carcinoma.

30 ASBT inhibition has been investigated for decreasing plasma cholesterol levels and improving insulin resistance, as well as to relieving the hepatic bile acid burden in cholestatic liver disease. In addition, ASBT inhibition has been found to restore insulin levels and normoglycemia, thus establishing ASBT inhibition as a promising treatment for type 2 diabetes mellitus. ASBT inhibitors are also used for
35 treatment of functional constipation.

As ASBT is predominantly expressed in the ileum (where it is often referred to as IBAT), ASBT inhibitors need not be systemically available. On the other hand, ASBT is also expressed in the proximal tubule cells of the kidneys. ASBT inhibitors that are systemically available may therefore also inhibit the reuptake of bile acids in the kidneys. It is believed that this would lead to increased levels of bile acids in urine, and to an increased removal of bile acids from the body via the urine. Systemically available ASBT inhibitors that exert their effect not only in the ileum but also in the kidneys are therefore expected to lead to a greater reduction of bile acid levels than non-systemically available ASBT inhibitors that only exert their effect in the ileum.

Compounds having a high ASBT inhibiting potency are particularly suitable for the treatment of liver diseases that cause cholestasis, such as progressive familial intrahepatic cholestasis (PFIC), Alagilles syndrome, biliary atresia and non-alcoholic steatohepatitis (NASH).

Biliary atresia is a rare pediatric liver disease that involves a partial or total blockage (or even absence) of large bile ducts. This blockage or absence causes cholestasis that leads to the accumulation of bile acids that damages the liver. In some embodiments, the accumulation of bile acids occurs in the extrahepatic biliary tree. In some embodiments, the accumulation of bile acids occurs in the intrahepatic biliary tree. The current standard of care is the Kasai procedure, which is a surgery that removes the blocked bile ducts and directly connects a portion of the small intestine to the liver. There are currently no approved drug therapies for this disorder.

Provided herein are methods for treating biliary atresia in a subject in need thereof, the methods comprising administration of a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the subject has undergone the Kasai procedure prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the subject is administered a compound of formula (I), or a pharmaceutically acceptable salt thereof, prior to undergoing the Kasai procedure. In some embodiments, the treatment of biliary atresia decreases the level of serum bile acids in the subject.

In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described in Danese et al., PLoS One. 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids

prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the treatment of biliary atresia includes treatment of pruritus.

PFIC is a rare genetic disorder that is estimated to affect between one in every 50,000 to 100,000 children born worldwide and causes progressive, life-threatening liver disease.

One manifestation of PFIC is pruritus, which often results in a severely diminished quality of life. In some cases, PFIC leads to cirrhosis and liver failure. Current therapies include Partial External Biliary Diversion (PEBD) and liver transplantation, however, these options can carry substantial risk of post-surgical complications, as well as psychological and social issues.

Three alternative gene defects have been identified that correlate to three separate PFIC subtypes known as types 1, 2 and 3.

- PFIC, type 1, which is sometimes referred to as “Byler disease,” is caused by impaired bile secretion due to mutations in the ATP8B1 gene, which codes for a protein that helps to maintain an appropriate balance of fats known as phospholipids in cell membranes in the bile ducts. An imbalance in these phospholipids is associated with cholestasis and elevated bile acids in the liver. Subjects affected by PFIC, type 1 usually develop cholestasis in the first months of life and, in the absence of surgical treatment, progress to cirrhosis and end-stage liver disease before the end of the first decade of life.
- PFIC, type 2, which is sometimes referred to as “Byler syndrome,” is caused by impaired bile salt secretion due to mutations in the ABCB11 gene, which codes for a protein, known as the bile salt export pump, that moves bile acids out of the liver. Subjects with PFIC, type 2 often develop liver failure within the first few years of life and are at increased risk of developing a type of liver cancer known as hepatocellular carcinoma.
- PFIC, type 3, which typically presents in the first years of childhood with progressive cholestasis, is caused by mutations in the ABCB4 gene, which codes for a transporter that moves phospholipids across cell membranes.

In addition, TJP2 gene, NR1H4 gene or Myo5b gene mutations have been proposed to be causes of PFIC. In addition, some subjects with PFIC do not have a mutation in any of the ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b genes. In these cases, the cause of the condition is unknown.

Exemplary mutations of the ATP8B1 gene or the resulting protein are listed in Tables 2 and 3, with numbering based on the human wild type ATP8B1 protein (e.g., SEQ ID NO: 1) or gene (e.g., SEQ ID NO: 2). Exemplary mutations of the ABCB11 gene or the resulting protein are listed in Tables 4 and 5, with numbering based on the human wild type ABCB11 protein (e.g., SEQ ID NO: 3) or gene (e.g., SEQ ID NO: 4).

As can be appreciated by those skilled in the art, an amino acid position in a reference protein sequence that corresponds to a specific amino acid position in SEQ ID NO: 1 or 3 can be determined by aligning the reference protein sequence with SEQ ID NO: 1 or 3 (e.g., using a software program, such as ClustalW2). Changes to these residues (referred to herein as “mutations”) may include single or multiple amino acid substitutions, insertions within or flanking the sequences, and deletions within or flanking the sequences. As can be appreciated by those skilled in the art, a nucleotide position in a reference gene sequence that corresponds to a specific nucleotide position in SEQ ID NO: 2 or 4 can be determined by aligning the reference gene sequence with SEQ ID NO: 2 or 4 (e.g., using a software program, such as ClustalW2). Changes to these residues (referred to herein as “mutations”) may include single or multiple nucleotide substitutions, insertions within or flanking the sequences, and deletions within or flanking the sequences. See also Kooistra, et al., “KLIFS: A structural kinase-ligand interaction database,” *Nucleic Acids Res.* 2016, vol. 44, no. D1, pp. D365-D371, which is incorporated by reference in its entirety herein.

Canonical protein sequence of ATP8B1 (SEQ ID NO: 1) – Uniprot ID O43520

5 MSTERDSETT FDEDSQPNDE VVPYSDDETE DELDDQGS AV EPEQNRVNRE AEENREPFRK
 ECTWQVKAND RKYHEQPHFM NTKFLCIKES KYANNAIKTY KYNAFTFIPM NLFEQFKRAA
 NLYFLALLIL QAVPQISTLA WYTTLVPLLV VLGVTAIKDL VDDVARHKMD KEINNRTCEV
 IKDGRFKVAK WKEIQVG DVI RLKKND FVPA DILLSSSEP NSLCYVETAE LDGETNLKFK
 MSLEITDQYL QREDTLATFD GFIECEEPNN RLDKFTGT LF WRNTSFPLDA DKILLRGCVI
 RNTDFCHGLV IFAGADTKIM KNSGKTRFKR TKIDYLMNYM VYTIFVVLIL LSAGLAIGHA
 YWEAQVGNSS WYLYDGEDDT PSYRGFLIFW GYIIIVLNTMV PISLYVSVEV IRLGQSHFIN
 10 WDLQMYAEK DTPAKARTTT LNEQLGQIHY IFSDKTGT LT QNIMTFKKCC INGQIYGDHR
 DASQHNHNKI EQVDFSWNTY ADGKLAFYDH YLIEQIQSGK EPEVRQFFFL LAVCHTMVMD
 RTDQQLNYQA ASPDEGALVN AARNFGFAFL ARTQNTITIS ELGTERTYNV LAILDENSDR
 KRMSIIVRTP EGNIKLYCKG ADTVIYERLH RMNPTKQETQ DALDIFANET LRTLCLCYKE
 IEEKEFTAWN KKFMAASVAS TNRDEALDKV YEEIEKDLIL LGATAIEDKL QDGV PETISK
 LAKADIKI WV LTGDKKETAE NIGFACELLT EDTTICYGED INSL LHARME NQRNRGGVYA
 15 KFAPPVQESF FPPGGNRALI ITGSWLNEIL LEKKT KRNKI LKLKFP RTEE ERRMRTQSKR
 RLEAKKEQRQ KNFVDLACEC SAVICCRVTP KQKAMVVDLV KRYKKAITLA IGDGANDVNM
 IKTAHIGVGI SGQEGMQAVM SSDYSFAQFR YLQRLLLVHG RWSYIRMCKF LRYFFYKNFA
 FTLVHFWYSF FNGYSAQTAY EDWFITLYNV LYTSLPVLLM GLLDQDVSDK LSLRFPGLYI
 VGQRDLLFNY KRFFVSLLHG VLTSMILFFI PLGAYLQTVG QDGEAPSDYQ SFAVTIASAL
 20 VITVNFQIGL DTSYWTFVNA FSIFGSIALY FGIMFDFHSA GIHVLFP SAF QFTGTASNAL
 RQPYIWL TII LAVAVCLLPV VAIRFLSMTI WPSES DKI QK HRKRLKAE EQ WQRRQQVFRR
 GVSTRRSAYA FSHQRGYADL ISSGRSIRKK RSPLDAIVAD GTAEYRRTGD S

Canonical DNA Sequence for ATP8B1 (SEQ ID NO: 2)

25 ATG AGT ACA GAA AGA GAC TCA GAA ACG ACA TTT GAC GAG GAT TCT CAG CCT
 AAT GAC GAA GTG GTT CCC TAC AGT GAT GAT GAA ACA GAA GAT GAA CTT GAT
 GAC CAG GGG TCT GCT GTT GAA CCA GAA CAA AAC CGA GTC AAC AGG GAA GCA
 GAG GAG AAC CGG GAG CCA TTC AGA AAA GAA TGT ACA TGG CAA GTC AAA GCA
 AAC GAT CGC AAG TAC CAC GAA CAA CCT CAC TTT ATG AAC ACA AAA TTC TTG
 30 TGT ATT AAG GAG AGT AAA TAT GCG AAT AAT GCA ATT AAA ACA TAC AAG TAC
 AAC GCA TTT ACC TTT ATA CCA ATG AAT CTG TTT GAG CAG TTT AAG AGA GCA
 GCC AAT TTA TAT TTC CTG GCT CTT CTT ATC TTA CAG GCA GTT CCT CAA ATC
 TCT ACC CTG GCT TGG TAC ACC ACA CTA GTG CCC CTG CTT GTG GTG CTG GGC
 GTC ACT GCA ATC AAA GAC CTG GTG GAC GAT GTG GCT CGC CAT AAA ATG GAT
 35 AAG GAA ATC AAC AAT AGG ACG TGT GAA GTC ATT AAG GAT GGC AGG TTC AAA
 GTT GCT AAG TGG AAA GAA ATT CAA GTT GGA GAC GTC ATT CGT CTG AAA AAA
 AAT GAT TTT GTT CCA GCT GAC ATT CTC CTG CTG TCT AGC TCT GAG CCT AAC
 AGC CTC TGC TAT GTG GAA ACA GCA GAA CTG GAT GGA GAA ACC AAT TTA AAA
 TTT AAG ATG TCA CTT GAA ATC ACA GAC CAG TAC CTC CAA AGA GAA GAT ACA
 40 TTG GCT ACA TTT GAT GGT TTT ATT GAA TGT GAA GAA CCC AAT AAC AGA CTA
 GAT AAG TTT ACA GGA ACA CTA TTT TGG AGA AAC ACA AGT TTT CCT TTG GAT
 GCT GAT AAA ATT TTG TTA CGT GGC TGT GTA ATT AGG AAC ACC GAT TTC TGC
 CAC GGC TTA GTC ATT TTT GCA GGT GCT GAC ACT AAA ATA ATG AAG AAT AGT
 GGG AAA ACC AGA TTT AAA AGA ACT AAA ATT GAT TAC TTG ATG AAC TAC ATG
 45 GTT TAC ACG ATC TTT GTT GTT CTT ATT CTG CTT TCT GCT GGT CTT GCC ATC
 GGC CAT GCT TAT TGG GAA GCA CAG GTG GGC AAT TCC TCT TGG TAC CTC TAT
 GAT GGA GAA GAC GAT ACA CCC TCC TAC CGT GGA TTC CTC ATT TTC TGG GGC
 TAT ATC ATT GTT CTC AAC ACC ATG GTA CCC ATC TCT CTC TAT GTC AGC GTG
 GAA GTG ATT CGT CTT GGA CAG AGT CAC TTC ATC AAC TGG GAC CTG CAA ATG

	TAC	TAT	GCT	GAG	AAG	GAC	ACA	CCC	GCA	AAA	GCT	AGA	ACC	ACC	ACA	CTC	AAT
	GAA	CAG	CTC	GGG	CAG	ATC	CAT	TAT	ATC	TTC	TCT	GAT	AAG	ACG	GGG	ACA	CTC
	ACA	CAA	AAT	ATC	ATG	ACC	TTT	AAA	AAG	TGC	TGT	ATC	AAC	GGG	CAG	ATA	TAT
	GGG	GAC	CAT	CGG	GAT	GCC	TCT	CAA	CAC	AAC	CAC	AAC	AAA	ATA	GAG	CAA	GTT
5	GAT	TTT	AGC	TGG	AAT	ACA	TAT	GCT	GAT	GGG	AAG	CTT	GCA	TTT	TAT	GAC	CAC
	TAT	CTT	ATT	GAG	CAA	ATC	CAG	TCA	GGG	AAA	GAG	CCA	GAA	GTA	CGA	CAG	TTC
	TTC	TTC	TTG	CTC	GCA	GTT	TGC	CAC	ACA	GTC	ATG	GTG	GAT	AGG	ACT	GAT	GGT
	CAG	CTC	AAC	TAC	CAG	GCA	GCC	TCT	CCC	GAT	GAA	GGT	GCC	CTG	GTA	AAC	GCT
	GCC	AGG	AAC	TTT	GGC	TTT	GCC	TTC	CTC	GCC	AGG	ACC	CAG	AAC	ACC	ATC	ACC
10	ATC	AGT	GAA	CTG	GGC	ACT	GAA	AGG	ACT	TAC	AAT	GTT	CTT	GCC	ATT	TTG	GAC
	TTC	AAC	AGT	GAC	CGG	AAG	CGA	ATG	TCT	ATC	ATT	GTA	AGA	ACC	CCA	GAA	GGC
	AAT	ATC	AAG	CTT	TAC	TGT	AAA	GGT	GCT	GAC	ACT	GTT	ATT	TAT	GAA	CGG	TTA
	CAT	CGA	ATG	AAT	CCT	ACT	AAG	CAA	GAA	ACA	CAG	GAT	GCC	CTG	GAT	ATC	TTT
	GCA	AAT	GAA	ACT	CTT	AGA	ACC	CTA	TGC	CTT	TGC	TAC	AAG	GAA	ATT	GAA	GAA
15	AAA	GAA	TTT	ACA	GAA	TGG	AAT	AAA	AAG	TTT	ATG	GCT	GCC	AGT	GTG	GCC	TCC
	ACC	AAC	CGG	GAC	GAA	GCT	CTG	GAT	AAA	GTA	TAT	GAG	GAG	ATT	GAA	AAA	GAC
	TTA	ATT	CTC	CTG	GGA	GCT	ACA	GCT	ATT	GAA	GAC	AAG	CTA	CAG	GAT	GGA	GTT
	CCA	GAA	ACC	ATT	TCA	AAA	CTT	GCA	AAA	GCT	GAC	ATT	AAG	ATC	TGG	GTG	CTT
	ACT	GGA	GAC	AAA	AAG	GAA	ACT	GCT	GAA	AAT	ATA	GGA	TTT	GCT	TGT	GAA	CTT
20	CTG	ACT	GAA	GAC	ACC	ACC	ATC	TGC	TAT	GGG	GAG	GAT	ATT	AAT	TCT	CTT	CTT
	CAT	GCA	AGG	ATG	GAA	AAC	CAG	AGG	AAT	AGA	GGT	GGC	GTC	TAC	GCA	AAG	TTT
	GCA	CCT	CCT	GTG	CAG	GAA	TCT	TTT	TTT	CCA	CCC	GGT	GGA	AAC	CGT	GCC	TTA
	ATC	ATC	ACT	GGT	TCT	TGG	TTG	AAT	GAA	ATT	CTT	CTC	GAG	AAA	AAG	ACC	AAG
	AGA	AAT	AAG	ATT	CTG	AAG	CTG	AAG	TTC	CCA	AGA	ACA	GAA	GAA	GAA	AGA	CGG
25	ATG	CGG	ACC	CAA	AGT	AAA	AGG	AGG	CTA	GAA	GCT	AAG	AAA	GAG	CAG	CGG	CAG
	AAA	AAC	TTT	GTG	GAC	CTG	GCC	TGC	GAG	TGC	AGC	GCA	GTC	ATC	TGC	TGC	CGC
	GTC	ACC	CCC	AAG	CAG	AAG	GCC	ATG	GTG	GTG	GAC	CTG	GTG	AAG	AGG	TAC	AAG
	AAA	GCC	ATC	ACG	CTG	GCC	ATC	GGA	GAT	GGG	GCC	AAT	GAC	GTG	AAC	ATG	ATC
	AAA	ACT	GCC	CAC	ATT	GGC	GTT	GGA	ATA	AGT	GGA	CAA	GAA	GGA	ATG	CAA	GCT
30	GTC	ATG	TCG	AGT	GAC	TAT	TCC	TTT	GCT	CAG	TTC	CGA	TAT	CTG	CAG	AGG	CTA
	CTG	CTG	GTG	CAT	GGC	CGA	TGG	TCT	TAC	ATA	AGG	ATG	TGC	AAG	TTC	CTA	CGA
	TAC	TTC	TTT	TAC	AAA	AAC	TTT	GCC	TTT	ACT	TTG	GTT	CAT	TTC	TGG	TAC	TCC
	TTC	TTC	AAT	GGC	TAC	TCT	GCG	CAG	ACT	GCA	TAC	GAG	GAT	TGG	TTC	ATC	ACC
	CTC	TAC	AAC	GTG	CTG	TAC	ACC	AGC	CTG	CCC	GTG	CTC	CTC	ATG	GGG	CTG	CTC
35	GAC	CAG	GAT	GTG	AGT	GAC	AAA	CTG	AGC	CTC	CGA	TTC	CCT	GGG	TTA	TAC	ATA
	GTG	GGA	CAA	AGA	GAC	TTA	CTA	TTC	AAC	TAT	AAG	AGA	TTC	TTT	GTA	AGC	TTG
	TTG	CAT	GGG	GTC	CTA	ACA	TCG	ATG	ATC	CTC	TTC	TTC	ATA	CCT	CTT	GGA	GCT
	TAT	CTG	CAA	ACC	GTA	GGG	CAG	GAT	GGA	GAG	GCA	CCT	TCC	GAC	TAC	CAG	TCT
	TTT	GCC	GTC	ACC	ATT	GCC	TCT	GCT	CTT	GTA	ATA	ACA	GTC	AAT	TTC	CAG	ATT
40	GGC	TTG	GAT	ACT	TCT	TAT	TGG	ACT	TTT	GTG	AAT	GCT	TTT	TCA	ATT	TTT	GGA
	AGC	ATT	GCA	CTT	TAT	TTT	GGC	ATC	ATG	TTT	GAC	TTT	CAT	AGT	GCT	GGA	ATA
	CAT	GTT	CTC	TTT	CCA	TCT	GCA	TTT	CAA	TTT	ACA	GGC	ACA	GCT	TCA	AAC	GCT
	CTG	AGA	CAG	CCA	TAC	ATT	TGG	TTA	ACT	ATC	ATC	CTG	GCT	GTT	GCT	GTG	TGC
	TTA	CTA	CCC	GTC	GTT	GCC	ATT	CGA	TTC	CTG	TCA	ATG	ACC	ATC	TGG	CCA	TCA
45	GAA	AGT	GAT	AAG	ATC	CAG	AAG	CAT	CGC	AAG	CGG	TTG	AAG	GCG	GAG	GAG	CAG
	TGG	CAG	CGA	CGG	CAG	CAG	GTG	TTC	CGC	CGG	GGC	GTG	TCA	ACG	CGG	CGC	TCG
	GCC	TAC	GCC	TTC	TCG	CAC	CAG	CGG	GGC	TAC	GCG	GAC	CTC	ATC	TCC	TCC	GGG
	CGC	AGC	ATC	CGC	AAG	AAG	CGC	TCG	CCG	CTT	GAT	GCC	ATC	GTG	GCG	GAT	GGC
	ACC	GCG	GAG	TAC	AGG	CGC	ACC	GGG	GAC	AGC	TGA						

Table 2. Exemplary ATP8B1 Mutations

Amino acid position 3 (e.g., T3K) ²⁷
Amino acid position 23 (e.g., P23L) ⁵
Amino acid position 45 (e.g., N45T) ^{5,8,9}
Amino acid position 46 (e.g., R46X) ^{A,25}
Amino acid position 62 (e.g., C62R) ²⁸
Amino acid position 63 (e.g., T63T) ⁴¹
Amino acid position 70 (e.g., D70N) ^{1,6}
Amino acid position 71 (e.g., R71H) ⁴³
Amino acid position 78 (e.g., H78Q) ¹⁹
Amino acid position 82 (e.g., T82T) ⁴¹
Amino acid position 92 (e.g., Y92Y) ⁴¹
Amino acid position 93 (e.g., A93A) ⁶
Amino acid position 96 (e.g., A96G) ²⁷
Amino acid position 114 (e.g., E114Q) ⁸
Amino acid position 127 (e.g., L127P ⁶ , L127V ³⁶)
Amino acid position 177 (e.g., T177T) ⁶
Amino acid position 179 (e.g., E179X) ²⁹
Δ Amino acid positions 185-282 ⁴⁴
Amino acid position 197 (e.g., G197Lfs*10) ²²
Amino acid position 201 (e.g., R201S ²⁷ , R201H ³⁵)
Amino acid position 203 (e.g., K203E ^{5,8} , K203R ⁹ , K203fs ²⁵)
Amino acid position 205 (e.g., N205fs ⁶ , N205Kfs*2 ³⁵)
Amino acid position 209 (e.g., P209T) ⁴
Amino acid position 217 (e.g., S217N) ⁴³
Amino acid position 232 (e.g., D232D) ³⁰
Amino acid position 233 (e.g., G233R) ³⁸
Amino acid position 243 (e.g., L243fs*28) ³³
Amino acid position 265 (e.g., C265R) ²⁵
Amino acid position 271 (e.g., R271X ¹³ , R271R ³⁰)
Amino acid position 288 (e.g., L288S) ⁶
Amino acid position 294 (e.g., L294S) ⁴³

Amino acid position 296 (e.g., R296C) ¹¹
Amino acid position 305 (e.g., F305I) ²⁸
Amino acid position 306 (e.g., C306R) ²³
Amino acid position 307 (e.g., H307L) ³⁵
Amino acid position 308 (e.g., G308V ¹ , G308D ⁶ , G308S ³⁵)
Amino acid position 314 (e.g., G314S) ¹³
Amino acid position 320 (e.g., M320Vfs*13) ¹¹
Amino acid position 337 (e.g., M337R) ¹⁸
Amino acid position 338 (e.g., N338K) ¹⁸
Amino acid position 340 (e.g., M340V) ¹⁸
Amino acid position 344 (e.g., I344F) ^{6,20}
Amino acid position 349 (e.g., I349T) ⁴¹
Amino acid position 358 (e.g., G358R) ²⁸
Amino acid position 367 (e.g., G367G) ⁴¹
Amino acid position 368 (e.g., N368D) ⁴¹
Amino acid position 393 (e.g., I393V) ²⁷
Amino acid position 403 (e.g., S403Y) ⁶
Amino acid position 407 (e.g., S407N) ⁴⁰
Amino acid position 412 (e.g., R412P) ⁶
Amino acid position 415 (e.g., Q415R) ²⁷
Amino acid position 422 (e.g., D422H) ³⁵
Amino acid position 429 (e.g., E429A) ⁶
Amino acid position 446 (e.g., G446R) ^{4,11}
Amino acid position 453 (e.g., S453Y) ⁶
Amino acid position 454 (e.g., D454G) ⁶
Amino acid position 455 (e.g., K455N) ⁴³
Amino acid position 456 (e.g., T456M ^{3,6} , T456K ³⁵)
Amino acid position 457 (e.g., G457G ⁶ , G457fs*6 ³³)
Amino acid position 469 (e.g., C469G) ⁴¹
Amino acid position 478 (e.g., H478H) ⁴¹
Amino acid position 500 (e.g., Y500H) ⁶
Amino acid position 525 (e.g., R525X) ⁴
Δ Amino acid position 529 ⁶

Amino acid position 535 (e.g., H535L ⁶ , H535N ⁴¹)
Amino acid position 553 (e.g., P553P) ⁴³
Amino acid position 554 (e.g., D554N ^{1,6} , D554A ³⁵)
Δ Amino acid positions 556-628 ⁴⁴
Δ Amino acid positions 559-563 ³⁵
Amino acid position 570 (e.g., L570L) ⁴¹
Amino acid position 577 (e.g., I577V) ¹⁹
Amino acid position 581 (e.g., E581K) ³⁵
Amino acid positions 554 and 581 (e.g., D554A+E581K) ³⁵
Amino acid position 585 (e.g., E585X) ²¹
Amino acid position 600 (e.g., R600W ^{2,4} , R600Q ⁶)
Amino acid position 602 (e.g., R602X) ^{3,6}
Amino acid position 628 (e.g., R628W) ⁶
Amino acid position 631 (e.g., R631Q) ²⁸
Δ Amino acid positions 645-699 ⁴
Amino acid position 661 (e.g., I661T) ^{1,4,6}
Amino acid position 665 (e.g., E665X) ^{4,6}
Amino acid position 672 (e.g., K672fs ⁶ , K672Vfs*1 ³⁵)
Amino acid position 674 (e.g., M674T) ¹⁹
Amino acid positions 78 and 674 (e.g., H78Q/M674T) ¹⁹
Amino acid position 684 (e.g., D684D) ⁴¹
Amino acid position 688 (e.g., D688G) ⁶
Amino acid position 694 (e.g., I694T ⁶ , I694N ¹⁷)
Amino acid position 695 (e.g., E695K) ²⁷
Amino acid position 709 (e.g., K709fs ⁶ , K709Qfs*41 ¹³)
Amino acid position 717 (e.g., T717N) ⁴
Amino acid position 733 (e.g., G733R) ⁶
Amino acid position 757 (e.g., Y757X) ⁴
Amino acid position 749 (e.g., L749P) ²¹
Amino acid position 792 (e.g., P792fs) ⁶
Δ Amino acid position 795-797 ⁶
Amino acid position 809 (e.g., I809L) ²⁷
Amino acid position 814 (e.g., K814N) ²⁸

Amino acid position 833 (e.g., R833Q ²⁷ , R833W ⁴¹)
Amino acid position 835 (e.g., K835Rfs*36) ³⁵
Amino acid position 845 (e.g., K845fs) ²⁵
Amino acid position 849 (e.g., R849Q) ²⁴
Amino acid position 853 (e.g., F853S, F853fs) ⁶
Amino acid position 867 (e.g., R867C ¹ , R867fs ⁶ , R867H ²³)
Amino acid position 885 (e.g., K885T) ⁴¹
Amino acid position 888 (e.g., T888T) ⁴¹
Amino acid position 892 (e.g., G892R) ⁶
Amino acid position 912 (e.g., G912R) ³⁵
Amino acid position 921 (e.g., S921S) ⁴¹
Amino acid position 924 (e.g., Y924C) ²⁸
Amino acid position 930 (e.g., R930X ⁶ , R930Q ²⁸)
Amino acid position 941 (e.g., R941X) ³⁵
Amino acid position 946 (e.g., R946T) ⁴¹
Amino acid position 952 (e.g., R952Q ^{5,9,15} , R952X ⁶)
Amino acid position 958 (e.g., N958fs) ⁶
Amino acid position 960 (e.g., A960A) ⁴¹
Δ Amino acid position 971 ⁴³
Amino acid position 976 (e.g., A976E ⁴¹ , A976A ⁴³)
Amino acid position 981 (e.g., E981K) ²⁰
Amino acid position 994 (e.g., S994R) ⁴
Amino acid position 1011 (e.g., L1011fs*18) ³³
Amino acid position 1012 (e.g., S1012I) ¹⁰
Amino acid position 1014 (e.g., R1014X) ^{6,11}
Amino acid position 1015 (e.g., F1015L) ²⁷
Amino acid position 1023 (e.g., Q1023fs) ⁶
Amino acid position 1040 (e.g., G1040R) ^{1,6}
Amino acid position 1044 (e.g., S0144L) ³⁴
Amino acid position 1047 (e.g., L1047fs) ⁶
Amino acid position 1050 (e.g., I1050K) ³¹
Amino acid position 1052 (e.g., L1052R) ²⁸
Amino acid position 1095 (e.g., W1095X) ¹¹

Amino acid position 1098 (e.g., V1098X) ³⁵
Amino acid position 1131 (e.g., Q1131X) ⁴⁴
Amino acid position 1142 (e.g., A1142Tfs*35) ⁴³
Amino acid position 1144 (e.g., Y1144Y) ⁴³
Amino acid position 1150 (e.g., I1150T) ⁴¹
Amino acid position 1152 (e.g., A1152T) ³⁰
Amino acid position 1159 (e.g., P1159P) ^{25,43}
Amino acid position 1164 (e.g., R1164X) ⁶
Amino acid position 1193 (e.g., R1193fs*39) ³³
Amino acid position 1197 (e.g., V1197L) ⁴¹
Amino acid position 1208 (e.g., A1208fs) ⁶
Amino acid position 1209 (e.g., Y1209Lfs*28) ⁴
Amino acid position 1211 (e.g., F1211L) ²⁷
Amino acid position 1219 (e.g., D1219H ⁵ , D1219G ²⁷)
Amino acid position 1223 (e.g., S1223S) ⁴¹
Amino acid position 1233 (e.g., P1233P) ⁴¹
Amino acid position 1241 (e.g., G1241fs) ⁶
Amino acid position 1248 (e.g., T1248T) ⁴³
Splice site mutation IVS3+1_+3delGTG ⁶
Splice site mutation IVS3-2A>G ⁶
IVS6+5T>G ^{17,25}
Splice site mutation IVS8+1G>T ⁶
IVS9-G>A ²⁶
IVS12+1G>A ²⁵
Splice site mutation IVS17-1G>A ⁶
Splice site mutation IVS18+2T>C ⁶
Splice site mutation IVS20-4CT>AA
Splice site mutation IVS21+5G>A ⁶
Splice site mutation IVS23-3C>A ⁶
Splice site mutation IVS26+2T>A ⁶
g.24774-42062del ⁴
c.-4C>G ⁴¹
c.145C>T ¹²

c.181-72G>A ⁹
c.182-5T>A ⁴¹
c.182-72G>A ⁴¹
c.246A>G ⁹
c.239G>A ³⁹
c.279+1_279+3delGTG ⁴⁶
c.280-2A>G ⁴⁶
c.625_62715delinsACAGTAAT ⁴⁶
c.554+122C>T ⁹
c.555-3T>C ²⁷
c.625+5 G>T ⁴
Amino acid position 209 (e.g., P209T) and c.625+5 G>T ⁴
c.628-30G>A ⁴¹
c.628-31C>T ⁴¹
c.698+1G>T ⁴⁶
c.698+20C>T ⁴¹
c.782-1G>A ⁴⁶
c.782-34G>A ⁴¹
Δ795-797 ¹⁴
c.782 -1G>A ⁴
c.852A>C ²⁷
c.941-1G>A ⁴⁶
c.1014C>T ⁹
c.1029+35G>A ⁹
c.1221-8C.G ⁴¹
1226delA ¹⁶
c.1429+1G>A ⁴⁶
c.1429+2T>G ¹³
c.1429+49G>A ⁴¹
c.1430-42A>G ⁴¹
c.1493T>C ¹²
c.1587_1589delCTT ⁴⁶
c.1630+2T>G ²⁷

c.1631-10T>A ⁴¹
c.1637-37T>C ⁴¹
1660 G>A ¹⁴
1798 C>T ¹⁴
1799 G>A ¹⁴
c.1819-39_41delAA ⁹
c.1819+1G>A ³¹
c.1820-27G>A ⁴¹
c.1918+8C>T ²⁷
c.1933-1G>AK46
c.2097+2T>C ³²
c.2097+60T>G ⁴¹
c.2097+89T>C ⁴¹
c.2097+97T>G ⁴¹
c.2210-114T>C ⁹
2210delA ¹⁶
c.2210-45_50dupATAAAA ⁹
c.2285+29C.T ⁴¹
c.2285+32A>G ⁴¹
c.2286-4_2286-3delinsAA ⁴⁶
c.2418+5G>A ⁴⁶
c.2707+3G>C ²⁷
c.2707+9T>G ⁴¹
c.2707+43A>G ⁴¹
c.2709-59T>C ⁴¹
c.2931+9A>G ⁴¹
c.2931+59T>A ⁴¹
c.2932-3C>A ⁴⁶
c.2932+59T>A ⁹
c.2937A>C ²⁷
c.3016-9C>A ³¹
c.3033-3034del ¹⁹
3122delTCCTA/

insACATCGATGTTGATGTTAGG ⁴⁵
3318 G>A ¹⁴
c.3400+2T>A ⁴⁶
c.3401-175C>T ⁹
c.3401-167C>T ⁹
c.3401-108C>T ⁹
c.3531+8G>T ^{9,15}
c.3532-15C>T ⁹
Δ Phe ex 15 ⁴
Ex1_Ex13del ⁶
Ex2_Ex6del ³³
Ex12_Ex14del ²⁷
Skipped Exon 24 ⁴⁵
del5'UTR-ex18 ¹¹
c.*11C>T ⁴¹
c.*1101 + 366G > A ⁷
g.92918del565 ³¹
GC preceding exon 16 (e.g., resulting in a 4 bp deletion) ⁴²
Frameshift from the 5' end of exon 16 ⁴²
5' 1.4 kb deletion ⁴⁶

Table 3. Selected ATP8B1 Mutations Associated with PFIC-1

Amino acid position 23 (e.g., P23L) ⁵
Amino acid position 78 (e.g., H78Q) ¹⁹
Amino acid position 93 (e.g., A93A) ⁶
Amino acid position 96 (e.g., A96G) ²⁷
Amino acid position 127 (e.g., L127P) ⁶
Amino acid position 197 (e.g., G197Lfs*10) ²²
Amino acid position 205 (e.g., N205fs) ⁶
Amino acid position 209 (e.g., P209T) ⁴
Amino acid position 233 (e.g., G233R) ³⁸
Amino acid position 243 (e.g., L243fs*28) ³³
Amino acid position 288 (e.g., L288S) ⁶

Amino acid position 296 (e.g., R296C) ¹¹
Amino acid position 308 (e.g., G308V ^{1,6})
Amino acid position 320 (e.g., M320Vfs*13) ¹¹
Amino acid position 403 (e.g., S403Y) ⁶
Amino acid position 407 (e.g., S407N) ⁴⁰
Amino acid position 412 (e.g., R412P) ⁶
Amino acid position 415 (e.g., Q415R) ²⁷
Amino acid position 429 (e.g., E429A) ⁶
Amino acid position 446 (e.g., G446R) ⁴
Amino acid position 456 (e.g., T456M) ^{3,6}
Amino acid position 457 (e.g., G457G ⁶ , G457fs*6 ³³)
Amino acid position 500 (e.g., Y500H) ⁶
Amino acid position 525 (e.g., R525X) ⁴
Δ Amino acid position 529 ⁶
Amino acid position 535 (e.g., H535L) ⁶
Amino acid position 554 (e.g., D554N) ^{1,6}
Amino acid position 577 (e.g., I577V) ¹⁹
Amino acid position 585 (e.g., E585X) ²¹
Amino acid position 600 (e.g., R600W) ⁴
Amino acid position 602 (e.g., R602X) ^{3,6}
Amino acid position 661 (e.g., I661T) ^{4,6}
Amino acid position 665 (e.g., E665X) ^{4,6}
Δ Amino acid positions 645-699 ⁴
Amino acid position 672 (e.g., K672fs) ⁶
Amino acid position 674 (e.g., M674T) ¹⁹
Amino acid positions 78 and 674 (e.g., H78Q/M674T) ¹⁹
Amino acid position 688 (e.g., D688G) ⁶
Amino acid position 694 (e.g., I694N) ¹⁷
Amino acid position 695 (e.g., E695K) ²⁷
Amino acid position 709 (e.g., K709fs) ⁶
Amino acid position 717 (e.g., T717N) ⁴
Amino acid position 733 (e.g., G733R) ⁶
Amino acid position 749 (e.g., L749P) ²¹

Amino acid position 757 (e.g., Y757X) ⁴
Amino acid position 792 (e.g., P792fs) ⁶
Amino acid position 809 (e.g., I809L) ²⁷
Amino acid position 853 (e.g., F853S, F853fs) ⁶
Amino acid position 867 (e.g., R867fs) ⁶
Amino acid position 892 (e.g., G892R) ⁶
Amino acid position 930 (e.g., R930X ⁶ , R952Q ¹⁵)
Amino acid position 952 (e.g., R952X) ⁶
Amino acid position 958 (e.g., N958fs) ⁶
Amino acid position 981 (e.g., E981K) ²⁰
Amino acid position 994 (e.g., S994R) ⁴
Amino acid position 1014 (e.g., R1014X) ^{6,11}
Amino acid position 1015 (e.g., F1015L) ²⁷
Amino acid position 1023 (e.g., Q1023fs) ⁶
Amino acid position 1040 (e.g., G1040R) ^{1,6}
Amino acid position 1047 (e.g., L1047fs) ⁶
Amino acid position 1095 (e.g., W1095X) ¹¹
Amino acid position 1208 (e.g., A1208fs) ⁶
Amino acid position 1209 (e.g., Y1209Lfs*28) ⁴
Amino acid position 1211 (e.g., F1211L) ²⁷
Amino acid position 1219 (e.g., D1219H ⁵ , D1219G ²⁷)
Splice site mutation IVS3+1_+3delGTG ⁶
Splice site mutation IVS3-2A>G ⁶
IVS6+5T>G ¹⁷
Splice site mutation IVS8+1G>T ⁶
IVS9-G>A ²⁶
Splice site mutation IVS17-1G>A ⁶
Splice site mutation IVS18+2T>C ⁶
Splice site mutation IVS21+5G>A ⁶
g.24774-42062del ⁴
c.145C>T ¹²
c.239G>A ³⁹
c.625+5 G>T ⁴

Amino acid position 209 (e.g., P209T) and c.625+5 G>T ⁴
c.782 -1G>A ⁴
c.1493T>C ¹²
c.1630+2T>G ²⁷
1660 G>A ¹⁴
c.2707+3G>C ²⁷
c.2097+2T>C ³²
c.3033-3034del ¹⁹
3318 G>A ¹⁴
c.3158+8G>T ¹⁵
Δ Phe ex 15 ⁴
Ex1_ Ex13del ⁶
Ex2_ Ex6del ³³
Ex12_ Ex14del ²⁷
del5'UTR-ex18 ¹¹
c.*1101 + 366G > A ⁷
GC preceding exon 16 (e.g., resulting in a 4 bp deletion) ⁴²
Frameshift from the 5' end of exon 16 ⁴²

^A A mutation to 'X' denotes an early stop codon

References for Tables 2 and 3

- ¹ Folmer et al., Hepatology. 2009, vol. 50(5), p. 1597-1605.
- 5 ² Hsu et al., Hepatol Res. 2009, vol. 39(6), p. 625-631.
- ³ Alvarez et al., Hum Mol Genet. 2004, vol. 13(20), p. 2451-2460.
- ⁴ Davit-Spraul et al., Hepatology 2010, vol. 51(5), p. 1645-1655.
- ⁵ Vitale et al., J Gastroenterol. 2018, vol. 53(8), p. 945-958.
- ⁶ Klomp et al., Hepatology 2004, vol. 40(1), p. 27-38.
- 10 ⁷ Zarenezhad et al., Hepatitis Monthly: 2017, vol. 17(2); e43500.
- ⁸ Dixon et al., Scientific Reports 2017, vol. 7, 11823.
- ⁹ Painter et al., Eur J Hum Genet. 2005, vol. 13(4), p. 435-439.
- ¹⁰ Deng et al., World J Gastroenterol. 2012, vol. 18(44), p. 6504-6509.
- ¹¹ Giovannoni et al., PLoS One. 2015, vol. 10(12): e0145021.
- 15 ¹² Li et al., Hepatology International 2017, vol. 11, No. 1, Supp. Supplement 1, pp. S180. Abstract Number: OP284.

- ¹³ Togawa et al., Journal of Pediatric Gastroenterology and Nutrition 2018, vol. 67, Supp. Supplement 1, pp. S363. Abstract Number: 615.
- ¹⁴ Miloh et al., Gastroenterology 2006, vol. 130, No. 4, Suppl. 2, pp. A759-A760. Meeting Info.: Digestive Disease Week Meeting/107th Annual Meeting of the American-Gastroenterological-
5 Association. Los Angeles, CA, USA. May 19.
- ¹⁵ Dröge et al., Zeitschrift für Gastroenterologie 2015, vol. 53, No. 12. Abstract Number: A3-27. Meeting Info: 32. Jahrestagung der Deutschen Arbeitsgemeinschaft zum Studium der Leber. Dusseldorf, Germany. 22 Jan 2016-23 Jan 2016
- ¹⁶ Mizuochi et al., Clin Chim Acta. 2012, vol. 413(15-16), p. 1301-1304.
- ¹⁷ Liu et al., Hepatology International 2009, vol. 3, No. 1, p. 184-185. Abstract Number: PE405. Meeting Info: 19th Conference of the Asian Pacific Association for the Study of the Liver. Hong Kong, China. 13 Feb 2009-16 Feb 2009
- ¹⁸ McKay et al., Version 2. F1000Res. 2013; 2: 32. DOI: 10.12688/f1000research.2-32.v2
- ¹⁹ Hasegawa et al., Orphanet J Rare Dis. 2014, vol. 9:89.
- ²⁰ Stone et al., J Biol Chem. 2012, vol. 287(49), p. 41139-51.
- ²¹ Kang et al., J Pathol Transl Med. 2019 May 16. doi: 10.4132/jptm.2019.05.03. [Epub ahead of print]
- ²² Sharma et al., BMC Gastroenterol. 2018, vol. 18(1), p. 107.
- ²³ Uegaki et al., Intern Med. 2008, vol. 47(7), p. 599-602.
- ²⁴ Goldschmidt et al., Hepatol Res. 2016, vol. 46(4), p. 306-311.
- ²⁵ Liu et al., J Pediatr Gastroenterol Nutr. 2010, vol. 50(2), p. 179-183.
- ²⁶ Jung et al., J Pediatr Gastroenterol Nutr. 2007, vol. 44(4), p. 453-458.
- ²⁷ Bounford. University of Birmingham. Dissertation Abstracts International, (2016) Vol. 75, No. 1C. Order No.: AAI10588329. ProQuest Dissertations & Theses.
- ²⁸ Stolz et al., Aliment Pharmacol Ther. 2019, vol. 49(9), p. 1195-1204.
- ²⁹ Ivashkin et al., Hepatology International 2016, vol. 10, No. 1, Suppl. SUPPL. 1, pp. S461. Abstract Number: LBO-38. Meeting Info: 25th Annual Conference of the Asian Pacific Association for the Study of the Liver, APASL 2016. Tokyo, Japan. 20 Feb 2016-24 Feb 2016
- ³⁰ Blackmore et al., J Clin Exp Hepatol. 2013, vol. 3(2), p. 159-161.
- ³¹ Matte et al., J Pediatr Gastroenterol Nutr. 2010, vol. 51(4), p. 488-493.
- ³² Squires et al., J Pediatr Gastroenterol Nutr. 2017, vol. 64(3), p. 425-430.
- ³³ Hayshi et al., EBioMedicine. 2018, vol. 27, p. 187-199.
- ³⁴ Nagasaka et al., J Pediatr Gastroenterol Nutr. 2007, vol. 45(1), p. 96-105.
- ³⁵ Wang et al., PLoS One. 2016; vol. 11(4): e0153114.
- ³⁶ Narchi et al., Saudi J Gastroenterol. 2017, vol. 23(5), p. 303-305.

- ³⁷ Alashkar et al., Blood 2015, vol. 126, No. 23. Meeting Info.: 57th Annual Meeting of the American-Society-of-Hematology. Orlando, FL, USA. December 05 -08, 2015. Amer Soc Hematol.
- ³⁸ Ferreira et al., Pediatric Transplantation 2013, vol. 17, Suppl. SUPPL. 1, pp. 99. Abstract Number: 239. Meeting Info: IPTA 7th Congress on Pediatric Transplantation. Warsaw, Poland. 13 Jul 2013-16 Jul 2013.
- ³⁹ Pauli-Magnus et al., J Hepatol. 2005, vol. 43(2), p. 342-357.
- ⁴⁰ Jericho et al., Journal of Pediatric Gastroenterology and Nutrition 2015, vol. 60(3), p. 368-374.
- ⁴¹ van der Woerd et al., PLoS One. 2013, vol. 8(11): e80553.
- ⁴² Copeland et al., J Gastroenterol Hepatol. 2013, vol. 28(3), p. 560-564.
- ⁴³ Dröge et al., J Hepatol. 2017, vol. 67(6), p. 1253-1264.
- ⁴⁴ Chen et al., Journal of Pediatrics 2002, vol. 140(1), p. 119-124.
- ⁴⁵ Jirsa et al., Hepatol Res. 2004, vol. 30(1), p. 1-3.
- ⁴⁶ van der Woerd et al., Hepatology 2015, vol. 61(4), p. 1382-1391.
- In some embodiments, the mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N, F529del, I661T, E665X, R930X, R952X, R1014X, and G1040R.

Canonical Protein Sequence of ABCB11 (SEQ ID NO: 3) – Uniprot ID O95342

```

MSDSVILRSI KKFGEENDGF ESDKSYNNNDK KSRLQDEKKG DGVRVGGFFQL FRFSSSTDIW
20 LMFVGSGLCAF LHGIAQPGVL LIFGTMTDVF IDYDVELQEL QIPGKACVNN TIVWTNSSLN
QNMTNGTRCG LLNIESEMIK FASYIAGIAV AVLITGYIQI CFWVIAAARQ IQKMRKFYFR
RIMRMEIGWF DCNSVGELNT RFSDDINKIN DAIADQMALF IQRMTSTICG FLLGGFFRGWK
LTLVIISVSP LIGIGAATIG LSVSKFTDYE LKAYAKAGVV ADEVISSMRT VAAFGGEKRE
VERYEKNLVF AQRWGIRKGI VMGFFTGFVW CLIFLCYALA FWYGSTLVLD EGEYTPGTLV
25 QIFLSVIVGA LNLGNASPC L EAFATGAAA TSIFETIDRK PIIDCMSEDG YKLDRIKGEI
EFHNVTFHYP SRPEVKILND LNMVIKPGEM TALVGPSGAG KSTALQLIQR FYDPCEGMVT
VDGHDIRSLN IQWLRDQIGI VEQEPVLFST TIAENIRYGR EDATMEDIVQ AAKEANAYNF
IMDLPQQFDT LVGEGGGQMS GGQKQ RVAIA RALIRNPKIL LLDMATSLD NESEAMVQEV
LSKIQHGHTI ISVAHRLSTV RAADTIIGFE HGTAVERGTH EELLERKGVY FTLVTLQSQG
30 NQALNEEDIK DATEDDMLAR TFSRGSYQDS LRASIRQSK SQLSYLVHEP PLAVVDHKST
YEEDRKDKDI PVQEEVEPAP VRRILKFSAP EWPYMLVGSV GAAVNGTVTP LYAFLFSQIL
GTFSIPDKEE QRSQINGVCL LFMVAMGCVSL FTQFLQGYAF AKSGELLTKR LRKFGFRAML
GQDIAWFDDL RNSPGALTTR LATDASQVQG AAGSQIGMIV NSFTNVTVM IIAFSFSWKL
SLVILCFFPF LALSGATQTR MLTGFA SRDK QALEMVGQIT NEALSNI RTV AGIGKERRFI
35 EALETELEKP FKTAIQKANI YGFCFAFAQC IMFIANSASY RGGYLISNE GLHFSYVFRV
ISAVVLSATA LGRAFSYTPS YAKAKISAAR FFQLLDRQPP ISVYNTAGEK WDNFQ GKIDF
VDCKFTYPSR PDSQVLNGLS VSISPGQTLA FVGSSGCGKS TSIQLLERFY DPDQ GKVMID
GHDSKKVNVQ FLRSNIGIVS QEPVLFACSI MDNIKYGDNT KEIPMERVIA AAKQAQLHDF
VMSLPKEYET NVGSQGSQLS RGEKQRIATA RAIVRDPKIL LLDEATSALD TESEKTVQVA
40 LDKAREGR TC IVIAHRLSTI QNADIIAVMA QGVVIEKGTH EELMAQKGAY YKLVT TGSPI S

```

Canonical DNA Sequence of ABCB11 (SEQ ID NO: 4)

5 ATG TCT GAC TCA GTA ATT CTT CGA AGT ATA AAG AAA TTT GGA GAG GAG AAT
 GAT GGT TTT GAG TCA GAT AAA TCA TAT AAT AAT GAT AAG AAA TCA AGG TTA
 CAA GAT GAG AAG AAA GGT GAT GGC GTT AGA GTT GGC TTC TTT CAA TTG TTT
 CGG TTT TCT TCA TCA ACT GAC ATT TGG CTG ATG TTT GTG GGA AGT TTG TGT
 GCA TTT CTC CAT GGA ATA GCC CAG CCA GGC GTG CTA CTC ATT TTT GGC ACA
 ATG ACA GAT GTT TTT ATT GAC TAC GAC GTT GAG TTA CAA GAA CTC CAG ATT
 CCA GGA AAA GCA TGT GTG AAT AAC ACC ATT GTA TGG ACT AAC AGT TCC CTC
 10 AAC CAG AAC ATG ACA AAT GGA ACA CGT TGT GGG TTG CTG AAC ATC GAG AGC
 GAA ATG ATC AAA TTT GCC AGT TAC TAT GCT GGA ATT GCT GTC GCA GTA CTT
 ATC ACA GGA TAT ATT CAA ATA TGC TTT TGG GTC ATT GCC GCA GCT CGT CAG
 ATA CAG AAA ATG AGA AAA TTT TAC TTT AGG AGA ATA ATG AGA ATG GAA ATA
 GGG TGG TTT GAC TGC AAT TCA GTG GGG GAG CTG AAT ACA AGA TTC TCT GAT
 GAT ATT AAT AAA ATC AAT GAT GCC ATA GCT GAC CAA ATG GCC CTT TTC ATT
 15 CAG CGC ATG ACC TCG ACC ATC TGT GGT TTC CTG TTG GGA TTT TTC AGG GGT
 TGG AAA CTG ACC TTG GTT ATT ATT TCT GTC AGC CCT CTC ATT GGG ATT GGA
 GCA GCC ACC ATT GGT CTG AGT GTG TCC AAG TTT ACG GAC TAT GAG CTG AAG
 GCC TAT GCC AAA GCA GGG GTG GTG GCT GAT GAA GTC ATT TCA TCA ATG AGA
 ACA GTG GCT GCT TTT GGT GGT GAG AAA AGA GAG GTT GAA AGG TAT GAG AAA
 20 AAT CTT GTG TTC GCC CAG CGT TGG GGA ATT AGA AAA GGA ATA GTG ATG GGA
 TTC TTT ACT GGA TTC GTG TGG TGT CTC ATC TTT TTG TGT TAT GCA CTG GCC
 TTC TGG TAC GGC TCC ACA CTT GTC CTG GAT GAA GGA GAA TAT ACA CCA GGA
 ACC CTT GTC CAG ATT TTC CTC AGT GTC ATA GTA GGA GCT TTA AAT CTT GGC
 AAT GCC TCT CCT TGT TTG GAA GCC TTT GCA ACT GGA CGT GCA GCA GCC ACC
 25 AGC ATT TTT GAG ACA ATA GAC AGG AAA CCC ATC ATT GAC TGC ATG TCA GAA
 GAT GGT TAC AAG TTG GAT CGA ATC AAG GGT GAA ATT GAA TTC CAT AAT GTG
 ACC TTC CAT TAT CCT TCC AGA CCA GAG GTG AAG ATT CTA AAT GAC CTC AAC
 ATG GTC ATT AAA CCA GGG GAA ATG ACA GCT CTG GTA GGA CCC AGT GGA GCT
 GGA AAA AGT ACA GCA CTG CAA CTC ATT CAG CGA TTC TAT GAC CCC TGT GAA
 30 GGA ATG GTG ACC GTG GAT GGC CAT GAC ATT CGC TCT CTT AAC ATT CAG TGG
 CTT AGA GAT CAG ATT GGG ATA GTG GAG CAA GAG CCA GTT CTG TTC TCT ACC
 ACC ATT GCA GAA AAT ATT CGC TAT GGC AGA GAA GAT GCA ACA ATG GAA GAC
 ATA GTC CAA GCT GCC AAG GAG GCC AAT GCC TAC AAC TTC ATC ATG GAC CTG
 CCA CAG CAA TTT GAC ACC CTT GTT GGA GAA GGA GGA GGC CAG ATG AGT GGT
 35 GGC CAG AAA CAA AGG GTA GCT ATC GCC AGA GCC CTC ATC CGA AAT CCC AAG
 ATT CTG CTT TTG GAC ATG GCC ACC TCA GCT CTG GAC AAT GAG AGT GAA GCC
 ATG GTG CAA GAA GTG CTG AGT AAG ATT CAG CAT GGG CAC ACA ATC ATT TCA
 GTT GCT CAT CGC TTG TCT ACG GTC AGA GCT GCA GAT ACC ATC ATT GGT TTT
 GAA CAT GGC ACT GCA GTG GAA AGA GGG ACC CAT GAA GAA TTA CTG GAA AGG
 40 AAA GGT GTT TAC TTC ACT CTA GTG ACT TTG CAA AGC CAG GGA AAT CAA GCT
 CTT AAT GAA GAG GAC ATA AAG GAT GCA ACT GAA GAT GAC ATG CTT GCG AGG
 ACC TTT AGC AGA GGG AGC TAC CAG GAT AGT TTA AGG GCT TCC ATC CGG CAA
 CGC TCC AAG TCT CAG CTT TCT TAC CTG GTG CAC GAA CCT CCA TTA GCT GTT
 GTA GAT CAT AAG TCT ACC TAT GAA GAA GAT AGA AAG GAC AAG GAC ATT CCT
 45 GTG CAG GAA GAA GTT GAA CCT GCC CCA GTT AGG AGG ATT CTG AAA TTC AGT
 GCT CCA GAA TGG CCC TAC ATG CTG GTA GGG TCT GTG GGT GCA GCT GTG AAC
 GGG ACA GTC ACA CCC TTG TAT GCC TTT TTA TTC AGC CAG ATT CTT GGG ACT
 TTT TCA ATT CCT GAT AAA GAG GAA CAA AGG TCA CAG ATC AAT GGT GTG TGC
 CTA CTT TTT GTA GCA ATG GGC TGT GTA TCT CTT TTC ACC CAA TTT CTA CAG
 50 GGA TAT GCC TTT GCT AAA TCT GGG GAG CTC CTA ACA AAA AGG CTA CGT AAA
 TTT GGT TTC AGG GCA ATG CTG GGG CAA GAT ATT GCC TGG TTT GAT GAC CTC

AGA AAT AGC CCT GGA GCA TTG ACA ACA AGA CTT GCT ACA GAT GCT TCC CAA
 GTT CAA GGG GCT GCC GGC TCT CAG ATC GGG ATG ATA GTC AAT TCC TTC ACT
 AAC GTC ACT GTG GCC ATG ATC ATT GCC TTC TCC TTT AGC TGG AAG CTG AGC
 CTG GTC ATC TTG TGC TTC TTC CCC TTC TTG GCT TTA TCA GGA GCC ACA CAG
 5 ACC AGG ATG TTG ACA GGA TTT GCC TCT CGA GAT AAG CAG GCC CTG GAG ATG
 GTG GGA CAG ATT ACA AAT GAA GCC CTC AGT AAC ATC CGC ACT GTT GCT GGA
 ATT GGA AAG GAG AGG CGG TTC ATT GAA GCA CTT GAG ACT GAG CTG GAG AAG
 CCC TTC AAG ACA GCC ATT CAG AAA GCC AAT ATT TAC GGA TTC TGC TTT GCC
 TTT GCC CAG TGC ATC ATG TTT ATT GCG AAT TCT GCT TCC TAC AGA TAT GGA
 10 GGT TAC TTA ATC TCC AAT GAG GGG CTC CAT TTC AGC TAT GTG TTC AGG GTG
 ATC TCT GCA GTT GTA CTG AGT GCA ACA GCT CTT GGA AGA GCC TTC TCT TAC
 ACC CCA AGT TAT GCA AAA GCT AAA ATA TCA GCT GCA CGC TTT TTT CAA CTG
 CTG GAC CGA CAA CCC CCA ATC AGT GTA TAC AAT ACT GCA GGT GAA AAA TGG
 GAC AAC TTC CAG GGG AAG ATT GAT TTT GTT GAT TGT AAA TTT ACA TAT CCT
 15 TCT CGA CCT GAC TCG CAA GTT CTG AAT GGT CTC TCA GTG TCG ATT AGT CCA
 GGG CAG ACA CTG GCG TTT GTT GGG AGC AGT GGA TGT GGC AAA AGC ACT AGC
 ATT CAG CTG TTG GAA CGT TTC TAT GAT CCT GAT CAA GGG AAG GTG ATG ATA
 GAT GGT CAT GAC AGC AAA AAA GTA AAT GTC CAG TTC CTC CGC TCA AAC ATT
 GGA ATT GTT TCC CAG GAA CCA GTG TTG TTT GCC TGT AGC ATA ATG GAC AAT
 20 ATC AAG TAT GGA GAC AAC ACC AAA GAA ATT CCC ATG GAA AGA GTC ATA GCA
 GCT GCA AAA CAG GCT CAG CTG CAT GAT TTT GTC ATG TCA CTC CCA GAG AAA
 TAT GAA ACT AAC GTT GGG TCC CAG GGG TCT CAA CTC TCT AGA GGG GAG AAA
 CAA CGC ATT GCT ATT GCT CGG GCC ATT GTA CGA GAT CCT AAA ATC TTG CTA
 CTA GAT GAA GCC ACT TCT GCC TTA GAC ACA GAA AGT GAA AAG ACG GTG CAG
 25 GTT GCT CTA GAC AAA GCC AGA GAG GGT CGG ACC TGC ATT GTC ATT GCC CAT
 CGC TTG TCC ACC ATC CAG AAC GCG GAT ATC ATT GCT GTC ATG GCA CAG GGG
 GTG GTG ATT GAA AAG GGG ACC CAT GAA GAA CTG ATG GCC CAA AAA GGA GCC
 TAC TAC AAA CTA GTC ACC ACT GGA TCC CCC ATC AGT TGA

30 **Table 4. Exemplary ABCB11 Mutations**

Amino acid position 1 (e.g., M1V) ⁹
Amino acid position 4 (e.g., S4X) ^{A,64}
Amino acid position 8 (e.g., R8X) ⁸⁸
Amino acid position 19 (e.g., G19R) ⁵⁶
Amino acid position 24 (e.g., K24X) ³⁵
Amino acid position 25 (e.g., S25X) ^{5,14}
Amino acid position 26 (e.g., Y26Ifs*7) ³⁸
Amino acid position 36 (e.g., D36D) ²⁷
Amino acid position 38 (e.g., K38Rfs*24) ⁷³
Amino acid position 43 (e.g., V43I) ⁵⁷
Amino acid position 49 (e.g., Q49X) ⁷³
Amino acid position 50 (e.g., L50S, L50W) ⁵⁷
Amino acid position 52 (e.g., R52W ²⁶ , R52R ²⁸)

Amino acid position 56 (e.g., S56L) ⁵⁸
Amino acid position 58 (e.g., D58N) ⁶²
Amino acid position 62 (e.g., M62K) ⁹
Amino acid position 66 (e.g., S66N) ¹⁷
Amino acid position 68 (e.g., C68Y) ⁴¹
Amino acid position 50 (e.g., L50S) ^{5,7}
Amino acid position 71 (e.g., L71H) ⁷³
Amino acid position 74 (e.g., I74R) ⁷¹
Amino acid position 77 (e.g., P77A) ⁷³
Amino acid position 87 (e.g., T87R) ⁶⁷
Amino acid position 90 (e.g., F90F) ^{7,27}
Amino acid position 93 (e.g., Y93S ¹³ , Y93X ⁸⁸)
Amino acid position 96 (e.g., E96X) ⁸⁸
Amino acid position 97 (e.g., L97X) ³⁹
Amino acid position 101 (e.g., Q101Dfs*8) ⁹
Amino acid position 107 (e.g., C107R) ³⁶
Amino acid position 112 (e.g., I112T) ⁹
Amino acid position 114 (e.g., W114R) ^{2,9}
Amino acid position 123 (e.g. M123T) ⁶⁷
Amino acid position 127 (e.g., T127Hfs*6) ⁵
Amino acid position 129 (e.g., C129Y) ²⁵
Amino acid position 130 (e.g., G130G) ⁷⁷
Amino acid position 134 (e.g., I134I) ²⁸
Amino acid position 135 (e.g., E135K ^{7,13} , E135L ¹⁷)
Amino acid position 137 (e.g., E137K) ⁷
Amino acid position 157 (e.g., Y157C) ⁵
Amino acid position 161 (e.g., C161X) ³⁹
Amino acid position 164 (e.g., V164Gfs*7 ³⁰ , V164I ⁸⁵)
Amino acid position 167 (e.g., A167S ⁴ , A167V ⁷ , A167T ^{9,17})
Amino acid position 181 (e.g., R181I) ³⁵
Amino acid position 182 (e.g., I182K) ⁹
Amino acid position 183 (e.g., M183V ⁸ , M183T ⁹)
Amino acid position 185 (e.g., M185I) ⁷³

Amino acid position 186 (e.g., E186G) ^{2,7,22}
Amino acid position 188 (e.g., G188W) ⁷³
Amino acid position 194 (e.g., S194P) ⁷
Amino acid position 198 (e.g., L198P) ⁷
Amino acid position 199 (e.g., N199Ifs*15X) ⁸⁸
Amino acid position 206 (e.g., I206V) ²⁸
Amino acid position 212 (e.g., A212T) ⁷³
Amino acid position 217 (e.g., M217R) ⁸⁸
Amino acid position 225 (e.g., T225P) ⁵⁷
Amino acid position 226 (e.g., S226L) ⁹
Amino acid position 232 (e.g., L232Cfs*9) ⁹
Amino acid position 233 (e.g., L233S) ⁸⁶
Amino acid position 238 (e.g., G238V) ^{2,7}
Amino acid position 242 (e.g., T242I) ^{5,7}
Amino acid position 245 (e.g., I245Tfs*26) ⁵⁷
Amino acid position 256 (e.g., A256G) ⁹
Amino acid position 260 (e.g., G260D) ⁷
Amino acid position 269 (e.g., Y269Y) ²⁷
Amino acid position 277 (e.g., A277E) ⁷⁷
Amino acid position 283 (e.g., E283D) ⁷³
Amino acid positions 212 and 283 (e.g., A212T+E283D) ⁷³
Amino acid position 284 (e.g., V284L ^{7,39} , V284A ⁷ , V284D ²³)
Amino acid position 297 (e.g., E297G ^{1,2,5,7} , E297K ⁷)
Amino acid position 299 (e.g., R299K) ²⁸
Amino acid position 303 (e.g., R303K ⁸ , R303M ⁶³ R303fsX321 ⁸³)
Amino acid position 304 (e.g., Y304X) ²⁶
Amino acid position 312 (e.g., Q312H) ⁷
Amino acid position 313 (e.g., R313S) ^{5,7}
Amino acid position 314 (e.g., W314X) ⁵⁷
Amino acid position 318 (e.g., K318Rfs*26) ²⁹
Amino acid position 319 (e.g., G319G) ⁷
Amino acid position 327 (e.g., G327E) ^{5,7}
Amino acid position 330 (e.g., W330X) ²⁴

Amino acid position 336 (e.g., C336S) ^{2,7}
Amino acid position 337 (e.g., Y337H) ^{21,27}
Amino acid position 342 (e.g., W342G) ⁵⁰
Amino acid position 354 (e.g., R354X) ⁹
Amino acid position 361 (e.g., Q361X ⁵⁷ , Q361R ⁷⁴)
Amino acid position 366 (e.g., V366V ²⁸ , V366D ⁵⁷)
Amino acid position 368 (e.g., V368Rfs*27) ⁵
Amino acid position 374 (e.g., G374S) ³
Amino acid position 380 (e.g., L380Wfs*18) ⁵
Amino acid position 382 (e.g., A382G) ⁸⁸
Δ Amino acid positions 382-388 ⁵
Δ Amino acid positions 383-389 ⁵⁷
Amino acid position 387 (e.g., R387H) ⁹
Amino acid position 390 (e.g., A390P) ^{5,7}
Amino acid position 395 (e.g., E395E) ²⁸
Amino acid position 404 (e.g., D404G) ⁹
Amino acid position 410 (e.g., G410D) ^{5,7}
Amino acid position 413 (e.g., L413W) ^{5,7}
Amino acid position 415 (e.g., R415X) ⁴²
Amino acid position 416 (e.g., I416I) ²⁷
Amino acid position 420 (e.g., I420T) ⁹
Amino acid position 423 (e.g., H423R) ¹³
Amino acid position 432 (e.g., R432T) ^{1,2,7}
Amino acid position 436 (e.g., K436N) ⁴⁰
Amino acid position 440 (e.g., D440E) ⁸⁸
Amino acid position 444 (e.g., V444A) ²
Amino acid position 454 (e.g., V454X) ⁴⁹
Amino acid position 455 (e.g., G455E) ⁹
Amino acid position 457 (e.g., S457Vfs*23) ⁸⁸
Amino acid position 461 (e.g., K461E) ^{2,7}
Amino acid position 462 (e.g., S462R) ⁸⁸
Amino acid position 463 (e.g., T463I) ^{5,7}
Amino acid position 466 (e.g., Q466K) ^{5,7}

Amino acid position 470 (e.g., R470Q ^{5,7} , R470X ⁹)
Amino acid position 471 (e.g., Y472X) ⁵
Amino acid position 472 (e.g., Y472C ^{5,27} , Y472X ¹⁴)
Amino acid position 473 (e.g., D473Q ³⁵ , D473V ⁸⁸)
Amino acid position 475 (e.g., C475X) ²⁹
Amino acid position 481 (e.g., V481E) ^{5,7}
Amino acid position 482 (e.g., D482G) ^{2,5,7}
Amino acid position 484 (e.g., H484Rfs*5) ⁹
Amino acid position 487 (e.g., R487H ² , R487P ⁵)
Amino acid position 490 (e.g., N490D) ^{5,7}
Amino acid position 493 (e.g., W493X) ⁸
Amino acid position 496 (e.g., D496V) ⁸⁸
Amino acid position 498 (e.g., I498T) ^{2,7}
Amino acid position 499 (e.g., G499E) ⁷³
Amino acid position 501 (e.g., V501G) ⁶⁸
Amino acid position 504 (e.g., E504K) ⁷⁹
Amino acid position 510 (e.g., T510T) ⁷
Amino acid position 512 (e.g., I512T) ^{5,7}
Amino acid position 515 (e.g., N515T ^{5,7} , N515D ⁶⁴)
Amino acid position 516 (e.g., I516M) ¹⁷
Amino acid position 517 (e.g., R517H) ^{5,7}
Amino acid position 520 (e.g., R520X) ⁵
Amino acid position 523 (e.g., A523G) ¹³
Amino acid position 528 (e.g., I528Sfs*21 ⁵ , I528X ⁹ , I528T ⁷³)
Amino acid position 535 (e.g., A535A ⁷ , A535X ⁸⁹)
Amino acid position 540 (e.g., F540L) ⁴⁶
Amino acid position 541 (e.g., I541L ^{5,7} , I541T ^{5,17})
Amino acid position 546 (e.g., Q546K ³⁹ , Q546H ⁷³)
Amino acid position 548 (e.g., F548Y) ^{5,7}
Amino acid position 549 (e.g., D549V) ⁹
Amino acid position 554 (e.g., E554K) ²¹
Amino acid position 556 (e.g., G556R) ⁶⁷
Amino acid position 558 (e.g., Q558H) ²³

Amino acid position 559 (e.g., M559T) ⁵⁷
Amino acid position 562 (e.g., G562D ^{5,7} , G562S ⁷³)
Amino acid position 570 (e.g., A570T ^{2,5,7} , A570V ²⁶)
Amino acid position 575 (e.g., R575X ^{2,5} , R575Q ²¹)
Amino acid position 580 (e.g., L580P) ⁵⁷
Amino acid position 586 (e.g., T586I) ⁷
Amino acid position 587 (e.g., S587X) ⁷³
Amino acid position 588 (e.g., A588V ^{5,7} , A588P ⁷³)
Amino acid position 591 (e.g., N591S) ^{2,7}
Amino acid position 593 (e.g., S593R) ^{2,7}
Amino acid position 597 (e.g., V597V ⁹ , V597L ¹³)
Amino acid position 603 (e.g., K603K) ⁵⁵
Amino acid position 609 (e.g., H609Hfs*46) ²⁶
Amino acid position 610 (e.g., I610Gfs*45 ⁹ , I610T ⁵⁷) ⁹
Amino acid position 615 (e.g., H615R) ²⁶
Amino acid position 616 (e.g., R616G ²⁸ , R616H ⁷³)
Amino acid position 619 (e.g., T619A) ²⁸
Amino acid position 623 (e.g., A623A) ²⁸
Amino acid position 625 (e.g., T625Nfs*5) ²⁶
Amino acid position 627 (e.g., I627T) ⁷
Amino acid position 628 (e.g., G628Wfs*3) ⁷⁰
Amino acid position 636 (e.g., E636G) ²
Amino acid position 648 (e.g., G648Vfs*6 ⁵ , G648V ⁵⁰)
Amino acid position 655 (e.g., T655I) ⁷
Amino acid position 669 (e.g., I669V) ²⁶
Amino acid position 676 (e.g., D676Y) ¹¹
Amino acid position 677 (e.g., M677V) ^{7,13}
Amino acid position 679 (e.g., A679V) ⁵⁸
Amino acid position 685 (e.g., G685W) ⁶⁰
Amino acid position 696 (e.g., R696W ²⁷ , R696Q ⁵⁸)
Amino acid position 698 (e.g., R698H ^{7,9} , R698K ⁶¹ , R698C ⁸⁸)
Amino acid position 699 (e.g., S699P) ⁹
Amino acid position 701 (e.g., S701P) ⁵⁸

Amino acid position 702 (e.g., Q702X) ⁸⁹
Amino acid position 709 (e.g., E709K) ⁷
Amino acid position 710 (e.g., P710P) ⁷
Amino acid position 712 (e.g., L712L) ²⁸
Amino acid position 721 (e.g., Y721C) ⁸⁸
Amino acid position 729 (e.g., D724N) ³⁹
Amino acid position 731 (e.g., P731S) ²³
Amino acid position 740 (e.g., P740Qfs*6) ⁷³
Amino acid position 758 (e.g., G758R) ⁵
Amino acid position 766 (e.g., G766R) ^{5,24}
Amino acid position 772 (e.g., Y772X) ⁵
Amino acid position 804 (e.g., A804A) ⁷
Amino acid position 806 (e.g., G806D ⁴⁴ , G806G ⁵⁵)
Amino acid position 809 (e.g., S809F) ⁸¹
Amino acid position 817 (e.g., G817G) ⁸⁸
Amino acid position 818 (e.g., Y818F) ⁷
Amino acid position 824 (e.g., G824E) ⁴²
Amino acid position 825 (e.g., G825G) ⁷³
Amino acid position 830 (e.g., R830Gfs*28) ⁷³
Amino acid position 832 (e.g., R832C ^{7,26} , R832H ⁴¹)
Amino acid position 842 (e.g., D842G) ²
Amino acid position 848 (e.g., D848N) ⁷³
Amino acid position 855 (e.g., G855R) ¹¹
Amino acid position 859 (e.g., T859R) ^{5,7}
Amino acid position 865 (e.g., A865V) ²⁷
Amino acid position 866 (e.g., S866A) ⁵⁷
Amino acid position 868 (e.g., V868D) ⁷³
Amino acid position 869 (e.g., Q869P) ⁷³
Amino acid position 875 (e.g., Q875X) ⁷³
Amino acid position 877 (e.g., G877R) ⁵⁶
Amino acid position 879 (e.g., I879R) ⁸⁸
Amino acid position 893 (e.g., A893V) ⁵⁷
Amino acid position 901 (e.g., S901R ¹⁷ , S901I ⁷³)

Amino acid position 903 (e.g., V903G) ⁵⁷
Δ Amino acid position 919 ¹²
Amino acid position 923 (e.g., T923P) ^{2,7}
Amino acid position 926 (e.g., A926P) ^{2,7}
Amino acid position 928 (e.g., R928X ¹⁵ , R928Q ⁴⁰)
Amino acid position 930 (e.g., K930X ⁵ , K930Efs*79 ^{5,10} , K930Efs*49 ²⁶)
Amino acid position 931 (e.g., Q931P) ²⁷
Amino acid position 945 (e.g., S945N) ⁵⁷
Amino acid position 948 (e.g., R948C) ^{5,7,26}
Amino acid position 958 (e.g., R958Q) ²⁸
Amino acid position 969 (e.g., K969K) ⁸⁸
Δ Amino acid positions 969-972 ⁵
Amino acid position 973 (e.g., T973I) ⁵⁷
Amino acid position 976 (e.g., Q976R ⁵⁸ , Q976X ⁸⁸)
Amino acid position 979 (e.g., N979D) ^{5,7}
Amino acid position 981 (e.g., Y981Y) ²⁸
Amino acid position 982 (e.g., G982R) ^{2,5,7}
Amino acid positions 444 and 982 (e.g., V444A+G982R) ³⁸
Amino acid position 995 (e.g., A995A) ²⁸
Amino acid position 1001 (e.g., R1001R) ⁹
Amino acid position 1003 (e.g., G1003R) ²⁴
Amino acid position 1004 (e.g., G1004D) ^{2,7}
Amino acid position 1027 (e.g., S1027R) ²⁶
Amino acid position 1028 (e.g., A1028A ^{7,10,88} , A1028E ⁸⁸)
Amino acid position 1029 (e.g., T1029K) ⁵
Amino acid position 1032 (e.g., G1032R) ¹²
Amino acid position 1041 (e.g., Y1041X) ⁹
Amino acid position 1044 (e.g., A1044P) ⁸⁸
Amino acid position 1050 (e.g., R1050C) ^{2,7,57}
Amino acid position 1053 (e.g., Q1053X) ⁵⁷
Amino acid position 1055 (e.g., L1055P) ³⁶
Amino acid position 1057 (e.g., R1057X ² , R1057Q ⁵⁸)
Amino acid position 1058 (e.g., Q1058Hfs*38 ⁹ , Q1058fs*38 ¹⁷ , Q1058X ⁷³)

Amino acid position 1061 (e.g., I1061Vfs*34) ⁹
Amino acid position 1083 (e.g., C1083Y) ⁴⁷
Amino acid position 1086 (e.g., T1086T) ²⁸
Amino acid position 1090 (e.g., R1090X) ^{2,5}
Amino acid position 1099 (e.g., L1099Lfs*38) ²⁶
Amino acid position 1100 (e.g., S1100Qfs*38) ¹³
Amino acid position 1110 (e.g., A1110E) ^{5,7}
Amino acid position 1112 (e.g., V1112F) ⁷⁰
Amino acid position 1116 (e.g., G1116R ⁷ , G1116F ^{9,17} , G1116E ³⁶)
Amino acid position 1120 (e.g., S1120N) ⁸⁸
Amino acid position 1128 (e.g., R1128H ^{2,7} , R1128C ^{5,7,13})
Amino acid position 1131 (e.g., D1131V) ²⁷
Amino acid position 1144 (e.g., S1144R) ⁷
Amino acid position 1147 (e.g., V1147X) ⁵
Amino acid position 1153 (e.g., R1153C ^{2,5,7} , R1153H ⁵)
Amino acid position 1154 (e.g., S1154P) ^{5,7}
Amino acid position 1162 (e.g., E1162X) ³⁹
Δ Amino acid position 1165 ⁸⁸
Amino acid position 1164 (e.g., V1164Gfs*7)
Amino acid position 1173 (e.g., N1173D) ⁵⁷
Amino acid position 1175 (e.g., K1175T) ⁵⁸
Amino acid position 1186 (e.g., E1186K) ⁷
Amino acid position 1192 (e.g., A1192Efs*50) ⁹
Amino acid position 1196 (e.g., Q1196X) ⁸⁸
Amino acid position 1197 (e.g., L1197G) ⁷
Amino acid position 1198 (e.g., H1198R) ²⁷
Amino acid position 1204 (e.g., L1204P) ⁸⁸
Amino acid position 1208 (e.g., Y1208C) ⁷³
Amino acid position 1210 (e.g., T1210P ^{5,7} , T1210F ⁵⁷)
Amino acid position 1211 (e.g., N1211D) ⁷
Amino acid position 1212 (e.g., V1212F) ³⁶
Amino acid position 1215 (e.g., Q1215X) ⁵
Amino acid position 1221 (e.g., R1221K) ⁵³

Amino acid position 1223 (e.g., E1223D) ⁷
Amino acid position 1226 (e.g., R1226P) ⁷³
Amino acid position 1228 (e.g., A1228V) ⁷
Amino acid position 1231 (e.g., R1231W ^{5,7} , R1231Q ^{5,7})
Amino acid position 1232 (e.g., A1232D) ¹⁷
Amino acid position 1235 (e.g., R1235X) ^{5,12}
Amino acid position 1242 (e.g., L1242I) ^{5,7}
Amino acid position 1243 (e.g., D1243G) ⁶⁷
Amino acid position 1249 (e.g., L1249X) ⁷³
Amino acid position 1256 (e.g., T1256fs*1296) ⁸³
Amino acid position 1268 (e.g., R1268Q) ^{2,7}
Amino acid position 1276 (e.g., R1276H) ³⁰
Amino acid position 1283 (e.g., A1283A ²⁸ , A1283V ⁸⁸)
Amino acid position 1292 (e.g., G1292V) ⁷³
Amino acid position 1298 (e.g., G1298R) ⁵
Amino acid position 1302 (e.g., E1302X) ⁵
Amino acid position 1311 (e.g., Y1311X) ⁵⁷
Amino acid position 1316 (e.g., T1316Lfs*64) ¹⁵
Amino acid position 1321 (e.g., S1321N) ⁵⁷
Intron 4 ((+3)A>C) ¹
IVS4-74A>T ⁸⁹
Splice site mutation 3' Intron 5 c.3901G>A ⁵
Splice site mutation 5; Intron 7 c.6111G>A ⁵
Splice site mutation IVS7+1G>A ¹⁴
IVS7+5G>A ⁴⁰
IVS8+1G>C ⁷⁶
Splice site mutation 5' Intron 9 c.9081delG ⁵
Splice site mutation 5' Intron 9 c.9081G>T ⁵
Splice site mutation 5' Intron 9 c.9081G>A ⁵
Splice site mutation IVS9+1G>T ¹⁴
Splice site mutation 3' Intron 13 c.143513_1435-8del ⁵
Splice site mutation IVS13del-13^8 ¹⁴
Splice site mutation 3' Intron 16 c.20128T>G ⁵

Splice site mutation IVS16-8T>G ¹⁴
Splice site mutation 5' Intron 18 c.21781G>T ⁵
Splice site mutation 5' Intron 18 c.21781G>A ⁵
Splice site mutation 5' Intron 18 c.21781G>C ⁵
Splice site mutation 3' Intron 18 c.21792A>G ⁵
Splice site mutation IVS18+1G>A ¹⁴
Splice site mutation 5' Intron 19 c.2343+1G>T ⁵
Splice site mutation 5' Intron 19 c.2343+2T>C ⁵
Splice site mutation IVS19+2T>C ¹⁴
Splice site mutation IVS19+1G>A ²²
Splice site mutation 3' Intron 21 c.26112A>T ⁵
IVS22+3A>G ⁸⁹
IVS 23-8 G-A ³⁶
IVS24+5G>A ⁵¹
Splice site mutation 5' Intron 24 c.32131delG ⁵
IVS35-6C>G ⁸⁹
Putative splice mutation 1198-1G>C ¹⁷
Putative splice mutation 1810-3C>G ¹⁷
Putative splice mutation 2178+1G>A ¹⁷
Putative splice mutation 2344-1G>T ¹⁷
Putative splice mutation c.2611-2A>T ³⁹
Putative splice mutation 3213+1_3213+2delinsA ¹⁷
c.-24C>A ^{44,78}
c.76 13 G>T ⁹
c.77-19T>A ⁵²
c.90_93delGAAA ¹⁸
c.124G>A ⁶⁹
c.150 +3 A>C ¹⁰
174C>T ⁵⁴
c.245T>C ⁸⁷
c.249_250insT ¹⁸
270T>C ⁵⁴
402C>T ⁵⁴

585G>C ⁵⁴
c.611+1G>A ⁷⁰
c.611+4A>G ³⁶
c.612-15_-6del10bp ⁵⁵
c.625A>C ³¹
c.627+5G>T ³¹
c.625A>C/ c.627+5G>T ³¹
696G>T ⁵⁴
c. 784+1G>C ⁴⁹
807T>C ⁵⁴
c.886C>T ³¹
c.890A>G ⁵⁹
c.908+1G>A ⁵⁷
c.908+5G>A ⁵⁵
c.908delG ⁵⁹
c.909-15A>G ⁶⁶
957A>G ⁵⁴
c.1084-2A>G ⁵⁷
1145 1bp deletion ⁹⁰
1281C>T ^{54,57}
c.1309-165C > T ¹⁹
c.1434 + 174G > A ¹⁹
c.1434 + 70C > T ¹⁹
c.1530C>A ⁵⁷
c.1587-1589delCTT ³¹
c.1621A>C ^{33,59}
c.1638+32T>C ⁶⁶
c.1638+80C>T ⁶⁶
1671C>T ⁵⁴
1791G>T ⁵⁴
1939delA ¹⁴
c.2075+3A>G ⁵³
c.2081T>A ³¹

c.2093G>A ⁶⁵
2098delA ¹⁶
c.2138-8T>G ⁶⁷
2142A>G ⁵⁴
c.2178+1G>T ^{36,39}
c.2179-17C>A ⁶⁶
c.2344-157T>G ⁶⁶
c.2344-17T>C ⁶⁶
c.2417G>A ⁷⁸
c.2541delG ⁸⁷
c.2620C>T ^{32,33}
c.2815-8A>G ⁵⁵
c.3003A>G ³⁷
c.3084A>G ^{48,54}
c.3213 +4 A>G ^{9,37}
c.3213 +5 G>A ⁹
c.3268C>T ⁷⁵
3285A>G ⁵⁴
c.3382C>T ⁷⁵
3435A>G ⁵⁴
c.3491delT ⁷²
c.3589C>T ⁵⁷
c.3765(+1 +5)del5 ⁴²
c.3766-34A>G ⁶⁶
c.3767-3768insC ⁶
c.3770delA ⁶⁷
c.3826C>T ⁷²
c.3846C>T ⁵⁷
c.3929delG ⁶⁷
c.*236A>G ⁶⁶
1145delC ⁸
Ex13_Ex17del ⁸²

Table 5. Selected ABCB11 Mutations Associated with PFIC-2

Amino acid position 1 (e.g., M1V) ⁹
Amino acid position 4 (e.g., S4X) ⁶⁴
Amino acid position 19 (e.g., G19R) ⁵⁶
Amino acid position 25 (e.g., S25X) ¹⁴
Amino acid position 26 (e.g., Y26Ifs*7) ³⁸
Amino acid position 50 (e.g., L50S) ^{7,57}
Amino acid position 52 (e.g., R52W) ²⁶
Amino acid position 58 (e.g., D58N) ⁶²
Amino acid position 62 (e.g., M62K) ⁹
Amino acid position 66 (e.g., S66N) ¹⁷
Amino acid position 68 (e.g., C68Y) ⁴¹
Amino acid position 93 (e.g., Y93S) ¹³
Amino acid position 101 (e.g., Q101Dfs*8) ⁹
Amino acid position 107 (e.g., C107R) ³⁶
Amino acid position 112 (e.g., I112T) ⁹
Amino acid position 114 (e.g., W114R) ^{2,9}
Amino acid position 129 (e.g., C129Y) ²⁵
Amino acid position 135 (e.g., E135K ¹³ , E135L ¹⁷)
Amino acid position 167 (e.g., A167V ⁷ , A167T ^{9,17})
Amino acid position 182 (e.g., I182K) ⁹
Amino acid position 183 (e.g., M183V ⁸ , M183T ⁹)
Amino acid position 225 (e.g., T225P) ⁵⁷
Amino acid position 226 (e.g., S226L) ⁹
Amino acid position 232 (e.g., L232Cfs*9) ⁹
Amino acid position 233 (e.g., L233S) ⁸⁶
Amino acid position 238 (e.g., G238V) ^{2,7}
Amino acid position 242 (e.g., T242I) ⁷
Amino acid position 245 (e.g., I245Tfs*26) ⁵⁷
Amino acid position 256 (e.g., A256G) ⁹
Amino acid position 260 (e.g., G260D) ⁵⁷
Amino acid position 284 (e.g., V284L) ⁷
Amino acid position 297 (e.g., E297G) ^{2,7}

Amino acid position 303 (e.g., R303K ⁸ , R303M ⁶³ , R303fsX321 ⁸³)
Amino acid position 304 (e.g., Y304X) ²⁶
Amino acid position 312 (e.g., Q312H) ⁷
Amino acid position 313 (e.g., R313S) ⁷
Amino acid position 314 (e.g., W314X) ⁵⁷
Amino acid position 318 (e.g., K318Rfs*26) ²⁹
Amino acid position 327 (e.g., G327E) ⁷
Amino acid position 330 (e.g., V330X) ²⁴
Amino acid position 336 (e.g., C336S) ^{2,7}
Amino acid position 337 (e.g., Y337H) ²¹
Amino acid position 342 (e.g., W342G) ⁵⁰
Amino acid position 354 (e.g., R354X) ⁹
Amino acid position 361 (e.g., Q361X) ⁵⁷
Amino acid position 366 (e.g., V366D) ⁵⁷
Amino acid position 386 (e.g., G386X) ³⁴
Δ Amino acid positions 383-389 ⁵⁷
Amino acid position 387 (e.g., R387H) ⁹
Amino acid position 390 (e.g., A390P) ⁷
Amino acid position 410 (e.g., G410D) ⁷
Amino acid position 413 (e.g., L413W) ⁷
Amino acid position 415 (e.g., R415X) ⁴²
Amino acid position 420 (e.g., I420T) ⁹
Amino acid position 454 (e.g., V454X) ⁴⁹
Amino acid position 455 (e.g., G455E) ⁹
Amino acid position 461 (e.g., K461E) ^{2,7}
Amino acid position 463 (e.g., T463I) ⁷
Amino acid position 466 (e.g., Q466K) ⁷
Amino acid position 470 (e.g., R470Q ⁷ , R470X ⁹)
Amino acid position 472 (e.g., Y472X ¹⁴ , Y472C ²⁷)
Amino acid position 475 (e.g., C475X) ²⁹
Amino acid position 481 (e.g., V481E) ⁷
Amino acid position 482 (e.g., D482G) ^{2,7}
Amino acid position 484 (e.g., H484Rfs*5) ⁹

Amino acid position 487 (e.g., R487H ² , R487P ⁸⁴)
Amino acid position 490 (e.g., N490D) ⁷
Amino acid position 493 (e.g., W493X) ⁸
Amino acid position 498 (e.g., I498T) ⁷
Amino acid position 501 (e.g., V501G) ⁶⁸
Amino acid position 512 (e.g., I512T) ⁷
Amino acid position 515 (e.g., N515T ⁷ , N515D ⁶⁴)
Amino acid position 516 (e.g., I516M) ¹⁷
Amino acid position 517 (e.g., R517H) ⁷
Amino acid position 520 (e.g., R520X) ⁵⁷
Amino acid position 523 (e.g., A523G) ¹³
Amino acid position 528 (e.g., I528X) ⁹
Amino acid position 540 (e.g., F540L) ⁴⁶
Amino acid position 541 (e.g., I541L ⁷ , I541T ¹⁷)
Amino acid position 548 (e.g., F548Y) ⁷
Amino acid position 549 (e.g., D549V) ⁹
Amino acid position 554 (e.g., E554K) ²¹
Amino acid position 559 (e.g., M559T) ⁵⁷
Amino acid position 562 (e.g., G562D) ⁷
Amino acid position 570 (e.g., A570T ⁷ , A570V ²⁶)
Amino acid position 575 (e.g., R575X ² , R575Q ²¹)
Amino acid position 588 (e.g., A588V) ⁷
Amino acid position 591 (e.g., N591S) ^{9,17}
Amino acid position 593 (e.g., S593R) ^{2,7}
Amino acid position 597 (e.g., V597V ⁹ , V597L ¹³)
Amino acid positions 591 and 597 (e.g., N591S+V597V) ⁹
Amino acid position 603 (e.g., K603K) ⁵⁵
Amino acid position 609 (e.g., H609Hfs*46) ²⁶
Amino acid position 610 (e.g., I610Gfs*45) ⁹
Amino acid position 615 (e.g., H615R) ²⁶
Amino acid position 625 (e.g., T625Nfs*5) ²⁶
Amino acid position 627 (e.g., I627T) ⁷
Amino acid position 636 (e.g., E636G) ²

Amino acid position 669 (e.g., I669V) ²⁶
Amino acid position 698 (e.g., R609H) ⁹
Amino acid positions 112 and 698 (e.g., I112T+R698H) ⁹
Amino acid position 699 (e.g., S699P) ⁹
Amino acid position 766 (e.g., G766R) ²⁴
Amino acid position 806 (e.g., G806G) ⁵⁵
Amino acid position 824 (e.g., G824E) ⁴²
Amino acid position 832 (e.g., R832C ^{7,26} , R832H ⁴¹)
Amino acid position 842 (e.g., D842G) ²
Amino acid position 859 (e.g., T859R) ⁷
Amino acid position 865 (e.g., A865V) ⁴⁵
Amino acid position 877 (e.g., G877R) ⁵⁶
Amino acid position 893 (e.g., A893V) ⁵⁷
Amino acid position 901 (e.g., S901R) ¹⁷
Amino acid position 903 (e.g., V903G) ⁵⁷
Δ Amino acid position 919 ¹²
Amino acid position 928 (e.g., R928X) ^{15,21}
Amino acid position 930 (e.g., K930Efs*79 ¹⁰ , K930Efs*49 ²⁶)
Amino acid position 948 (e.g., R948C) ^{7,26}
Amino acid position 979 (e.g., N979D) ⁷
Amino acid position 982 (e.g., G982R) ^{2,7}
Amino acid positions 444 and 982 (e.g., V444A+G982R) ³⁸
Amino acid position 1001 (e.g., R1001R) ⁹
Amino acid position 1003 (e.g., G1003R) ²⁴
Amino acid position 1004 (e.g., G1004D) ^{2,7}
Amino acid position 1027 (e.g., S1027R) ²⁶
Amino acid position 1028 (e.g., A1028A) ¹⁰
Amino acid position 1032 (e.g., G1032R) ¹²
Amino acid position 1041 (e.g., Y1041X) ⁹
Amino acid position 1050 (e.g., R1050C) ⁵⁷
Amino acid position 1053 (e.g., Q1053X) ⁵⁷
Amino acid position 1055 (e.g., L1055P) ³⁶
Amino acid position 1057 (e.g., R1057X) ²

Amino acid position 1058 (e.g., Q1058Hfs*38 ⁹ , Q1058fs*38 ¹⁷)
Amino acid position 1061 (e.g., I1061Vfs*34) ⁹
Amino acid position 1083 (e.g., C1083Y) ⁴⁷
Amino acid position 1090 (e.g., R1090X) ²
Amino acid position 1099 (e.g., L1099Lfs*38) ²⁶
Amino acid position 1100 (e.g., S1100Qfs*38) ¹³
Amino acid position 1110 (e.g., A1110E) ⁷
Amino acid position 1116 (e.g., G1116R ⁷ , G1116F ^{9,17} , G1116E ³⁶)
Amino acid position 1128 (e.g., R1128C) ^{7,13}
Amino acid position 1131 (e.g., D1131V) ²⁷
Amino acid position 1144 (e.g., S1144R) ⁷
Amino acid position 1153 (e.g., R1153C ^{2,7} , R1153H ^{7,26})
Amino acid position 1154 (e.g., S1154P) ⁷
Amino acid position 1173 (e.g., N1173D) ⁵⁷
Amino acid position 1192 (e.g., A1192Efs*50) ⁹
Amino acid position 1198 (e.g., H1198R) ²⁷
Amino acid position 1210 (e.g., T1210P ⁷ , T1210F ⁵⁷)
Amino acid position 1211 (e.g., N1211D) ⁷
Amino acid position 1212 (e.g., V1212F) ³⁶
Amino acid position 1231 (e.g., R1231W ⁷ , R1223Q ⁷)
Amino acid position 1232 (e.g., A1232D) ¹⁷
Amino acid position 1235 (e.g., R1235X) ¹²
Amino acid position 1242 (e.g., L1242I) ⁷
Amino acid position 1256 (e.g., T1256fs*1296) ⁸³
Amino acid position 1268 (e.g., R1268Q) ^{2,7}
Amino acid position 1302 (e.g., E1302X) ⁵⁷
Amino acid position 1311 (e.g., Y1311X) ⁵⁷
Amino acid position 1316 (e.g., T1316Lfs*64) ¹⁵
Intron 4 ((+3)A>C) ¹
Splice site mutation IVS7+1G>A ¹⁴
IVS8+1G>C ⁷⁶
Splice site mutation IVS9+1G>T ¹⁴
Splice site mutation IVS13del-13 ⁸ -8 ¹⁴

Splice site mutation IVS16-8T>G ¹⁴
Splice site mutation IVS18+1G>A ¹⁴
Splice site mutation IVS19+2T>C ¹⁴
IVS 23-8 G-A ³⁶
IVS24+5G>A ⁵¹
Putative splice mutation 1198-1G>C ¹⁷
Putative splice mutation 1810-3C>G ¹⁷
Putative splice mutation 2178+1G>A ¹⁷
Putative splice mutation 2344-1G>T ¹⁷
Putative splice mutation 3213+1_3213+2delinsA ¹⁷
c.-24C>A ⁷⁸
c.76 13 G>T ⁹
c.77-19T>A ⁵²
c.90_93delGAAA ¹⁸
c.124G>A ⁶⁹
c.150 +3 A>C ¹⁰
c.249_250insT ¹⁸
c.611+1G>A ⁸⁴
c.611+4A>G ³⁶
c.612-15_-6del10bp ⁵⁵
c.625A>C ³¹
c.627+5G>T ³¹
c.625A>C/ c.627+5G>T ³¹
c.886C>T ³¹
c.890A>G ⁵⁹
c.908+1G>A ⁵⁷
c.908+5G>A ⁵⁵
c.908delG ⁵⁹
1273 1bp deletion ⁹¹
c.1084-2A>G ⁵⁷
c.1445A>G ⁵⁹
c.1587-1589delCTT ³¹
c.1621A>C ⁵⁹

1939delA ¹⁴
c.2081T>A ³¹
2098delA ¹⁶
c.2343+1 G>T ⁸⁰
c.2178+1G>T ³⁶
c.2417G>A ⁷⁸
c.2620C>T ³²
c.2815-8A>G ⁵⁵
c.3003A>G ³⁷
c.3213 +4 A>G ^{9,37}
c.3213 +5 G>A ⁹
c.3268C>T ⁷⁵
c.3382C>T ⁷⁵
c.3765(+1 +5)del5 ⁴²
c.3767-3768insC ⁶
1145delC ⁸
Ex13_Ex17del ⁸²

^A A mutation to 'X' denotes an early stop codon

References for Tables 4 and 5

- ¹ Noe et al., J Hepatol. 2005, vol. 43(3), p. 536-543.
- 5 ² Lam et al., Am J Physiol Cell Physiol. 2007, vol. 293(5), p. C1709-16.
- ³ Stindt et al., Liver Int. 2013, vol. 33(10), p. 1527-1735.
- ⁴ Gao et al., Shandong Yiyao 2012, vol. 52(10), p. 14-16.
- ⁵ Strautnieks et al., Gastroenterology. 2008, vol. 134(4), p. 1203-1214.
- ⁶ Kagawa et al., Am J Physiol Gastrointest Liver Physiol. 2008, vol. 294(1), p. G58-67.
- 10 ⁷ Byrne et al., Hepatology. 2009, vol. 49(2), p. 553-567.
- ⁸ Chen et al., J Pediatr. 2008, vol. 153(6), p. 825-832.
- ⁹ Davit-Spraul et al., Hepatology 2010, vol. 51(5), p. 1645-1655.
- ¹⁰ Dröge et al., Sci Rep. 2016, vol. 6: 24827.
- ¹¹ Lang et al., Pharmacogenet Genomics. 2007, vol. 17(1), p. 47-60.
- 15 ¹² Ellinger et al., World J Gastroenterol. 2017, vol. 23(29), p. :5295-5303.
- ¹³ Vitale et al., J Gastroenterol. 2018, vol. 53(8), p. 945-958.
- ¹⁴ Knisely et al., Hepatology. 2006, vol. 44(2), p. 478-86.

- ¹⁵ Ellis et al., *Hepatology*. 2018, vol. 67(4), p. 1531-1545.
- ¹⁶ Lam et al., *J Hepatol*. 2006, vol. 44(1), p. 240-242.
- ¹⁷ Varma et al., *Hepatology* 2015, vol. 62(1), p. 198-206.
- ¹⁸ Treepongkaruna et al., *World J Gastroenterol*. 2009, vol. 15(34), p. 4339-4342.
- 5 ¹⁹ Zarenezhad et al., *Hepatitis Monthly*: 2017, vol. 17(2); e43500.
- ²⁰ Hayashi et al., *Hepatol Res*. 2016, vol. 46(2), p. 192-200.
- ²¹ Guorui et al., *Linchuang Erke Zazhi* 2013, vol. 31(10), 905-909.
- ²² van Mil et al., *Gastroenterology*. 2004, vol. 127(2), p. 379-384.
- ²³ Anzivino et al., *Dig Liver Dis*. 2013, vol. 45(3), p. 226-232.
- 10 ²⁴ Park et al., *World J Gastroenterol*. 2016, vol. 22(20), p. 4901-4907.
- ²⁵ Imagawa et al., *J Hum Genet*. 2018, vol. 63(5), p. 569-577.
- ²⁶ Giovannoni et al., *PLoS One*. 2015, vol. 10(12): e0145021.
- ²⁷ Hu et al., *Mol Med Rep*. 2014, vol. 10(3), p. 1264–1274.
- ²⁸ Lang et al., *Drug Metab Dispos*. 2006, vol. 34(9), p. 1582-1599.
- 15 ²⁹ Masahata et al., *Transplant Proc*. 2016, vol. 48(9), p. 3156-3162.
- ³⁰ Holz et al., *Hepatol Commun*. 2018, vol. 2(2), p. 152-154.
- ³¹ Li et al., *Hepatology International* 2017, vol. 11, No. 1, Supp. Supplement 1, pp. S180. Abstract Number: OP284.
- ³² Francalanci et al., *Laboratory Investigation* 2011, vol. 91, Supp. SUPPL. 1, pp. 360A. Abstract
- 20 Number: 1526.
- ³³ Francalanci et al., *Digestive and Liver Disease* 2010, vol. 42, Supp. SUPPL. 1, pp. S16. Abstract Number: T.N.5.
- ³⁴ Shah et al., *J Pediatr Genet*. 2017, vol. 6(2), p. 126-127.
- ³⁵ Gao et al., *Hepatitis Monthly* 2017, vol. 17(10), e55087/1-e55087/6.
- 25 ³⁶ Evason et al., *Am J Surg Pathol*. 2011, vol. 35(5), p. 687-696.
- ³⁷ Davit-Spraul et al., *Mol Genet Metab*. 2014, vol. 113(3), p. 225-229.
- ³⁸ Maggiore et al., *J Hepatol*. 2010, vol. 53(5), p. 981-6.
- ³⁹ McKay et al., *Version 2. F1000Res*. 2013; 2: 32. DOI: 10.12688/f1000research.2-32.v2
- ⁴⁰ Liu et al., *Pediatr Int*. 2013, vol. 55(2), p. 138-144.
- 30 ⁴¹ Waisbourd-Zinman et al., *Ann Hepatol*. 2017, vol. 16(3), p. 465-468.
- ⁴² Griffin, et al., *Canadian Journal of Gastroenterology and Hepatology* 2016, vol. 2016. Abstract Number: A200. Meeting Info: 2016 Canadian Digestive Diseases Week, CDDW 2016. Montreal, QC, United States. 26 Feb 2016-29 Feb 2016
- ⁴³ Qiu et al., *Hepatology* 2017, vol. 65(5), p. 1655-1669.
- 35 ⁴⁴ Imagawa et al., *Sci Rep*. 2017, 7:41806.

- ⁴⁵ Kang et al., J Pathol Transl Med. 2019 May 16. doi: 10.4132/jptm.2019.05.03. [Epub ahead of print]
- ⁴⁶ Takahashi et al., Eur J Gastroenterol Hepatol. 2007, vol. 19(11), p. 942-6.
- ⁴⁷ Shimizu et al., Am J Transplant. 2011, vol. 11(2), p. 394-398.
- ⁴⁸ Krawczyk et al., Ann Hepatol. 2012, vol. 11(5), p. 710-744.
- 5 ⁴⁹ Sharma et al., BMC Gastroenterol. 2018, vol. 18(1), p. 107.
- ⁵⁰ Sattler et al., Journal of Hepatology 2017, vol. 66, No. 1, Suppl. S, pp. S177. Meeting Info.:
International Liver Congress / 52nd Annual Meeting of the European-Association-for-the-Study-of-
the-Liver. Amsterdam, NETHERLANDS. April 19 -23, 2017. European Assoc Study Liver.
- ⁵¹ Jung et al., J Pediatr Gastroenterol Nutr. 2007, vol. 44(4), p. 453-458.
- 10 ⁵² Sciveres. Digestive and Liver Disease 2010, vol. 42, Suppl. SUPPL. 5, pp. S329. Abstract Number:
CO18. Meeting Info: 17th National Congress SIGENP. Pescara, Italy. 07 Oct 2010-09 Oct 2010
- ⁵³ Sohn et al., Pediatr Gastroenterol Hepatol Nutr. 2019, vol. 22(2), p. 201-206.
- ⁵⁴ Ho et al., Pharmacogenet Genomics. 2010, vol. 20(1), p. 45-57.
- ⁵⁵ Wang et al., Hepatol Res. 2018, vol. 48(7), p. 574-584.
- 15 ⁵⁶ Shaprio et al., J Hum Genet. 2010, vol. 55(5), p. 308-313.
- ⁵⁷ Bounford. University of Birmingham. Dissertation Abstracts International, (2016) Vol. 75, No. 1C.
Order No.: AAI10588329. ProQuest Dissertations & Theses.
- ⁵⁸ Stolz et al., Aliment Pharmacol Ther. 2019, vol. 49(9), p. 1195-1204.
- ⁵⁹ Jankowska et al., J Pediatr Gastroenterol Nutr. 2014, vol. 58(1), p. 92-95.
- 20 ⁶⁰ Kim. Journal of Pediatric Gastroenterology and Nutrition 2016, vol. 62, Suppl. SUPPL. 1, pp. 620.
Abstract Number: H-P-045. Meeting Info: 49th Annual Meeting of the European Society for
Paediatric Gastroenterology, Hepatology and Nutrition, ESPGHAN 2016. Athens, Greece. 25 May
2016-28 May 2016.
- ⁶¹ Pauli-Magnus et al., Hepatology 2003, vol. 38, No. 4 Suppl. 1, pp. 518A. print. Meeting Info.: 54th
25 Annual Meeting of the American Association for the Study of Liver Diseases. Boston, MA, USA.
October 24-28, 2003. American Association for the Study of Liver Diseases.
- ⁶² Li et al., Hepatology International 2017, vol. 11, No. 1, Suppl. Supplement 1, pp. S362. Abstract
Number: PP0347. Meeting Info: 26th Annual Conference of the Asian Pacific Association for the
Study of the Liver, APASL 2017. Shanghai, China. 15 Feb 2017-19 Feb 2017.
- 30 ⁶³ Rumbo et al., Transplantation 2018, vol. 102, No. 7, Suppl. Supplement 1, pp. S848. Abstract
Number: P.752. Meeting Info: 27th International Congress of The Transplantation Society, TTS
2018. Madrid, Spain. 30 Jun 2018-05 Jul 2018.
- ⁶⁴ Lee et al., Pediatr Gastroenterol Hepatol Nutr. 2017, vol. 20(2), p. 114-123.
- ⁶⁵ Sherrif et al., Liver international: official journal of the International Association for the Study of
35 the Liver 2013, vol. 33, No. 8, pp. 1266-1270.

- ⁶⁶ Blackmore et al., J Clin Exp Hepatol. 2013, vol. 3(2), p. 159-161.
- ⁶⁷ Matte et al., J Pediatr Gastroenterol Nutr. 2010, vol. 51(4), p. 488-493.
- ⁶⁸ Lin et al., Zhongguo Dang Dai Er Ke Za Zhi. 2018, vol. 20(9), p. 758-764.
- ⁶⁹ Harmanci et al., Experimental and Clinical Transplantation 2015, vol. 13, Suppl. SUPPL. 2, pp. 76.
- 5 Abstract Number: P62. Meeting Info: 1st Congress of the Turkic World Transplantation Society. Astana, Kazakhstan. 20 May 2015-22 May 2015.
- ⁷⁰ Herbst et al., Mol Cell Probes. 2015, vol. 29(5), p. 291-298.
- ⁷¹ Moghadamrad et al., Hepatology. 2013, vol. 57(6), p. 2539-2541.
- ⁷² Holz et al., Zeitschrift fur Gastroenterologie 2016, vol. 54, No. 8. Abstract Number: KV275. Meeting
- 10 Info: Viszeralmedizin 2016, 71. Jahrestagung der Deutschen Gesellschaft fur Gastroenterologie, Verdauungs- und Stoffwechselkrankheiten mit Sektion Endoskopie - 10. Herbsttagung der Deutschen Gesellschaft fur Allgemein- und Viszeralchirurgie. Hamburg, Germany. 21 Sep 2016-24 Sep 2016.
- ⁷³ Wang et al., PLoS One. 2016; vol. 11(4): e0153114.
- 15 ⁷⁴ Hao et al., International Journal of Clinical and Experimental Pathology 2017, vol. 10(3), p. 3480-3487.
- ⁷⁵ Arnell et al., J Pediatr Gastroenterol Nutr. 2010, vol. 51(4), p. 494-499.
- ⁷⁶ Sharma et al., Indian Journal of Gastroenterology 2017, vol. 36, No. 1, Suppl. Supplement 1, pp. A99. Abstract Number: M-20. Meeting Info: 58th Annual Conference of the Indian Society of
- 20 Gastroenterology, ISGCON 2017. Bhubaneswar, India. 14 Dec 2017-17 Dec 2017.
- ⁷⁷ Beauséjour et al., Can J Gastroenterol. 2011, vol. 25(6), p. 311-314.
- ⁷⁸ Imagawa et al., Journal of Pediatric Gastroenterology and Nutrition 2016, vol. 63, Suppl. Supplement 2, pp. S51. Abstract Number: 166. Meeting Info: World Congress of Pediatric
- Gastroenterology, Hepatology and Nutrition 2016. Montreal, QC, Canada. 05 Oct 2016-08 Oct
- 25 2016.
- ⁷⁹ Peng et al., Zhonghua er ke za zhi (Chinese journal of pediatrics) 2018, vol. 56, No. 6, pp. 440-444.
- ⁸⁰ Tibesar et al., Case Rep Pediatr. 2014, vol. 2014: 185923.
- ⁸¹ Ng et al., Journal of Pediatric Gastroenterology and Nutrition 2018, vol. 66, Suppl. Supplement 2, pp. 860. Abstract Number: H-P-127. Meeting Info: 51st Annual Meeting European Society for
- 30 Paediatric Gastroenterology, Hepatology and Nutrition, ESPGHAN 2018. Geneva, Switzerland. 09 May 2018-12 May 2018.
- ⁸² Wong et al., Clin Chem. 2008, vol. 54(7), p. 1141-1148.
- ⁸³ Pauli-Magnus et al., J Hepatol. 2005, vol. 43(2), p. 342-357.
- ⁸⁴ Jericho et al., Journal of Pediatric Gastroenterology and Nutrition. 60, vol. 3, p. 368-374.

⁸⁵ Scheimann et al., Gastroenterology 2007, vol. 132, No. 4, Suppl. 2, pp. A452. Meeting Info.:

Digestive Disease Week Meeting/108th Annual Meeting of the American-Gastroenterological-Association. Washington, DC, USA. May 19 -24, 2007. Amer Gastroenterol Assoc; Amer Assoc Study Liver Dis; Amer Soc Gastrointestinal Endoscopy; Soc Surg Alimentary Tract.

5 ⁸⁶ Jaquotot-Haerranz et al., Rev Esp Enferm Dig. 2013, vol. 105(1), p. 52-54.

⁸⁷ Khosla et al., American Journal of Gastroenterology 2015, vol. 110, No. Suppl. 1, pp. S397. Meeting Info.: 80th Annual Scientific Meeting of the American-College-of-Gastroenterology. Honolulu, HI, USA. October 16 -21, 2015.

⁸⁸ Dröge et al., J Hepatol. 2017, vol. 67(6), p. 1253-1264.

10 ⁸⁹ Liu et al., Liver International 2010, vol. 30(6), p. 809-815.

⁹⁰ Chen et al., Journal of Pediatrics 2002, vol. 140(1), p. 119-124.

⁹¹ U.S. Patent No. 9,295,677

In some embodiments, the mutation in ABCB11 is selected from A167T, G238V, V284L, E297G,
15 R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and R1268Q.

Provided are methods of treating PFIC (e.g., PFIC-1 and PFIC-2) in a subject that includes performing an assay on a sample obtained from the subject to determine whether the subject has a mutation associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b mutation), and
20 administering (e.g., specifically or selectively administering) a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, to the subject determined to have a mutation associated with PFIC. In some embodiments, the mutation is a ATP8B1 or ABCB11 mutation. For example, a mutation as provided in any one of Tables 1-4. In some embodiments, the mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N, F529del, I661T, E665X, R930X,
25 R952X, R1014X, and G1040R. In some embodiments, the mutation in ABCB11 is selected from A167T, G238V, V284L, E297G, R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and R1268Q.

Also provided are methods for treating PFIC (e.g., PFIC-1 and PFIC-2) in a subject in need thereof, the
30 method comprising: (a) detecting a mutation associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b mutation) in the subject; and (b) administering to the subject a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, methods for treating PFIC can include administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, to a
35 subject having a mutation associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or

Myo5b mutation). In some embodiments, the mutation is a ATP8B1 or ABCB11 mutation. For example, a mutation as provided in any one of Tables 1-4. In some embodiments, the mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N, F529del, I661T, E665X, R930X, R952X, R1014X, and G1040R. In some embodiments, the mutation in ABCB11 is selected from A167T, G238V, V284L, E297G, R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and R1268Q.

In some embodiments, the subject is determined to have a mutation associated with PFIC in a subject or a biopsy sample from the subject through the use of any art recognized tests, including next generation sequencing (NGS). In some embodiments, the subject is determined to have a mutation associated with PFIC using a regulatory agency-approved, e.g., FDA-approved test or assay for identifying a mutation associated with PFIC in a subject or a biopsy sample from the subject or by performing any of the non-limiting examples of assays described herein. Additional methods of diagnosing PFIC are described in Gunaydin, M. et al., Hepat Med. 2018, vol. 10, p. 95-104, incorporated by reference in its entirety herein.

In some embodiments, the treatment of PFIC (e.g., PFIC-1 or PFIC-2) decreases the level of serum bile acids in the subject. In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described in Danese et al., PLoS One. 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the treatment of PFIC includes treatment of pruritus.

Since LBAT is expressed on hepatocytes, LBAT and dual ASBT/LBAT inhibitor substances need to have at least some bioavailability and free fraction in blood. Because LBAT inhibitor compounds only need to survive from the intestine to the liver, it is expected that a relatively low systemic exposure of such compounds will be sufficient, thereby minimizing the potential risk for any side effects in the rest of the body. It is expected that inhibition of LBAT and ASBT will have at least additive effects in decreasing the intrahepatic bile acid concentration. It is also expected that a dual ASBT/LBAT inhibitor may be able to reduce bile acid levels without inducing diarrhoea, as is sometimes observed with ASBT inhibitors.

Compounds having a high LBAT inhibiting potency and sufficient bioavailability are expected to be particularly suitable for the treatment of hepatitis. Compounds having a dual ASBT/LBAT inhibiting potency and sufficient bioavailability are expected to be particularly suitable for the treatment of non-alcoholic steatohepatitis (NASH).

5

NASH is a common and serious chronic liver disease that resembles alcoholic liver disease, but that occurs in people who drink little or no alcohol. In NASH patients, fat accumulation in the liver, known as nonalcoholic fatty liver disease (NAFLD) or steatosis, and other factors such as high LDL cholesterol and insulin resistance induce chronic inflammation in the liver and may lead to progressive scarring of tissue, known as fibrosis, and cirrhosis, followed eventually by liver failure and death. Patients with NASH have been found to have significantly higher total serum bile acid concentrations than healthy subjects under fasting conditions (2.2- to 2.4-fold increase in NASH) and at all post-prandial time points (1.7- to 2.2-fold increase in NASH). These are driven by increased taurine- and glycine-conjugated primary and secondary bile acids. Patients with NASH exhibited greater variability in their fasting and post-prandial bile acid profile. These results indicate that patients with NASH have higher fasting and post-prandial exposure to bile acids, including the more hydrophobic and cytotoxic secondary species. Increased bile acid exposure may be involved in liver injury and the pathogenesis of NAFLD and NASH (Ferslew et al., *Dig Dis Sci.* 2015, vol. 60, p. 3318–3328). It is therefore likely that ASBT and/or LBAT inhibition will be beneficial for the treatment of NASH.

20

NAFLD is characterized by hepatic steatosis with no secondary causes of hepatic steatosis including excessive alcohol consumption, other known liver diseases, or long-term use of a steatogenic medication (Chalasani et al., *Hepatology* 2018, vol. 67(1), p. 328-357). NAFLD can be categorized into non-alcoholic fatty liver (NAFL) and non-alcoholic steatohepatitis (NASH). According to Chalasani et al., NAFL is defined as the presence of $\geq 5\%$ hepatic steatosis without evidence of hepatocellular injury in the form of hepatocyte ballooning. NASH is defined as the presence of $\geq 5\%$ hepatic steatosis and inflammation with hepatocyte injury (e.g., ballooning), with or without any liver fibrosis. NASH is also commonly associated with hepatic inflammation and liver fibrosis, which can progress to cirrhosis, end-stage liver disease, and hepatocellular carcinoma. While liver fibrosis is not always present in NASH, the severity of the fibrosis, when present, can be linked to long-term outcomes.

30

There are many approaches used to assess and evaluate whether a subject has NAFLD and if so, the severity of the disease, including differentiating whether the NAFLD is NAFL or NASH. In some embodiments, the severity of NAFLD can be assessed using the NAS. In some embodiments, treatment of NAFLD can be assessed using the NAS. In some embodiments, the NAS can be

35

determined as described in Kleiner et al., *Hepatology*. 2005, 41(6):1313-1321, which is hereby incorporated by reference in its entirety. See, for example, Table 6 for a simplified NAS scheme adapted from Kleiner.

5 **Table 6. Example of the NAFLD Activity Score (NAS) with Fibrosis Stage**

Feature	Degree	Score
Steatosis	<5%	0
	5-33%	1
	>33-66%	2
	>66%	3
Lobular Inflammation	No foci	0
	<2 foci/200x	1
	2-4 foci/200x	2
	>4 foci/200x	3
Ballooning degeneration	None	0
	Few	1
	Many cells/Prominent ballooning	2
Fibrosis	None	0
	Perisinusoidal or periportal	1
	Perisinusoidal & portal/periportal	2
	Bridging fibrosis	3
	Cirrhosis	4

In some embodiments, the NAS is determined non-invasively, for example, as described in U.S. Application Publication No. 2018/0140219, which is incorporated by reference herein in its entirety. In some embodiments, the NAS is determined for a sample from the subject prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the NAS is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a lower NAS score during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof compared to prior to

10

administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof indicates treatment of NAFLD (e.g., NASH). For example, a decrease in the NAS by 1, by 2, by 3, by 4, by 5, by 6, or by 7 indicates treatment of NAFLD (e.g., NASH). In some embodiments, the NAS following administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the NAS after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less.

Additional approaches of assessing and evaluating NASH in a subject include determining one or more of hepatic steatosis (e.g., accumulation of fat in the liver); hepatic inflammation; biomarkers indicative of one or more of liver damage, hepatic inflammation, liver fibrosis, and/or liver cirrhosis (e.g., serum markers and panels). Further examples of physiological indicators of NASH can include liver morphology, liver stiffness, and the size or weight of the subject's liver.

In some embodiments, NASH in the subject is evidenced by an accumulation of hepatic fat and detection of a biomarker indicative of liver damage. For example, elevated serum ferritin and low titers of serum autoantibodies can be common features of NASH.

In some embodiments, methods to assess NASH include magnetic resonance imaging, either by spectroscopy or by proton density fat fraction (MRI-PDFF) to quantify steatosis, transient elastography (FIBROSCAN®), hepatic venous pressure gradient (HPVG), hepatic stiffness measurement with MRE for diagnosing significant liver fibrosis and/or cirrhosis, and assessing histological features of liver biopsy. In some embodiments, magnetic resonance imaging is used to detect one or more of steatohepatitis (NASH-MRI), liver fibrosis (Fibro-MRI), and steatosis. See, for example, U.S. Application Publication Nos. 2016/146715 and 2005/0215882, each of which are incorporated herein by reference in their entireties.

In some embodiments, treatment of NASH can include a decrease of one or more symptoms associated with NASH; reduction in the amount of hepatic steatosis; a decrease in the NAS; a

decrease in hepatic inflammation; a decrease in the level of biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis; and a reduction in fibrosis and/or cirrhosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis in the subject following administration of one or more doses of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

In some embodiments, treatment of NASH comprises a decrease of one or more symptoms associated with NASH in the subject. Exemplary symptoms can include one or more of an enlarged liver, fatigue, pain in the upper right abdomen, abdominal swelling, enlarged blood vessels just beneath the skin's surface, enlarged breasts in men, enlarged spleen, red palms, jaundice, and pruritus. In some embodiments, the subject is asymptomatic. In some embodiments, the total body weight of the subject does not increase. In some embodiments, the total body weight of the subject decreases. In some embodiments, the body mass index (BMI) of the subject does not increase. In some embodiments, the body mass index (BMI) of the subject decreases. In some embodiments, the waist and hip (WTH) ratio of the subject does not increase. In some embodiments, the waist and hip (WTH) ratio of the subject decreases.

In some embodiments, treatment of NASH can be assessed by measuring hepatic steatosis. In some embodiments, treatment of NASH comprises a reduction in hepatic steatosis following administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as described herein. In some embodiments, hepatic steatosis is determined by one or more methods selected from the group consisting of ultrasonography, computed tomography (CT), magnetic resonance imaging, magnetic resonance spectroscopy (MRS), magnetic resonance elastography (MRE), transient elastography (TE) (e.g., FIBROSCAN®), measurement of liver size or weight, or by liver biopsy (see, e.g., Di Lascio et al., *Ultrasound Med Biol.* 2018, vol. 44(8), p. 1585-1596; Lv et al., *J Clin Transl Hepatol.* 2018, vol. 6(2), p. 217-221; Reeder et al., *J Magn Reson Imaging.* 2011, vol. 34(4), spcone; and de Lédinghen V, et al., *J Gastroenterol Hepatol.* 2016, vol. 31(4), p. 848-855, each of which are incorporated herein by reference in their entireties). A subject diagnosed with NASH can have greater than about 5% hepatic steatosis, for example, greater than about 5% to about 25%, about 25% to about 45%, about 45% to about 65%, or greater than about 65% hepatic steatosis. In some embodiments, a subject with greater than about 5% to about 33% hepatic steatosis has stage 1 hepatic steatosis, a subject with about 33% to about 66% hepatic steatosis has stage 2 hepatic steatosis, and a subject with greater than about 66% hepatic steatosis has stage 3 hepatic steatosis.

In some embodiments, the amount of hepatic steatosis is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the amount of hepatic steatosis is determined during the period of time or after the period of time of administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some
5 embodiments, a reduction in the amount of hepatic steatosis during the period of time or after the period of time of administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a reduction in the amount of hepatic steatosis by about 1% to about 50%, about 25% to about 75%, or about 50% to
10 about 100% indicates treatment of NASH. In some embodiments, a reduction in the amount of hepatic steatosis by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% indicates treatment of NASH.

15 In some embodiments, the presence of hepatic inflammation is determined by one or more methods selected from the group consisting of biomarkers indicative of hepatic inflammation and a liver biopsy sample(s) from the subject. In some embodiments, the severity of hepatic inflammation is determined from a liver biopsy sample(s) from the subject. For example, hepatic inflammation in a liver biopsy sample can be assessed as described in Kleiner et al., *Hepatology* 2005, vol. 41(6), p.
20 1313-1321 and Brunt et al., *Am J Gastroenterol* 1999, vol. 94, p. 2467-2474, each of which are hereby incorporated by reference in their entireties. In some embodiments, the severity of hepatic inflammation is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the severity of hepatic inflammation is determined during the period of time or after the period of time of administration of a compound
25 of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the severity of hepatic inflammation during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a decrease in the severity of hepatic inflammation
30 by about 1% to about 50%, about 25% to about 75%, or about 50% to about 100% indicates treatment of NASH. In some embodiments, a decrease in the severity of hepatic inflammation by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% indicates treatment of NASH.

In some embodiments, treatment of NASH comprises treatment of fibrosis and/or cirrhosis, e.g., a decrease in the severity of fibrosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis. In some embodiments, the presence of fibrosis and/or cirrhosis is determined by one or more methods selected from the group consisting of transient elastography (e.g., FIBROSCAN®), non-invasive markers of hepatic fibrosis, and histological features of a liver biopsy. In some embodiments, the severity (e.g., stage) of fibrosis is determined by one or more methods selected from the group consisting of transient elastography (e.g., FIBROSCAN®), a fibrosis-scoring system, biomarkers of hepatic fibrosis (e.g., non-invasive biomarkers), and hepatic venous pressure gradient (HVPG). Non-limiting examples of fibrosis scoring systems include the NAFLD fibrosis scoring system (see, e.g., Angulo et al., *Hepatology* 2007, vol. 45(4), p. 846-54), the fibrosis scoring system in Brunt et al., *Am. J. Gastroenterol.* 1999, vol. 94, p. 2467-2474, the fibrosis scoring system in Kleiner et al., *Hepatology* 2005, vol. 41(6), p. 1313-1321, and the ISHAK fibrosis scoring system (see Ishak et al., *J. Hepatol.* 1995, vol. 22, p. 696-699), the contents of each of which are incorporated by reference herein in their entireties.

In some embodiments, the severity of fibrosis is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the severity of fibrosis is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the severity of fibrosis during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. In some embodiments, a decrease in the severity of fibrosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis indicates treatment of NASH. In some embodiments, the severity of fibrosis is determined using a scoring system such as any of the fibrosis scoring systems described herein, for example, the score can indicate the stage of fibrosis, e.g., stage 0 (no fibrosis), stage 1, stage 2, stage 3, and stage 4 (cirrhosis) (see, e.g., Kleiner et al.). In some embodiments, a decrease in the stage of the fibrosis is a decrease in the severity of the fibrosis. For example, a decrease by 1, 2, 3, or 4 stages is a decrease in the severity of the fibrosis. In some embodiments, a decrease in the stage, e.g., from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 indicates treatment of NASH. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to

stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 following administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof.

In some embodiments, the presence of NASH is determined by one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis or scoring systems thereof. In some embodiments, the severity of NASH is determined by one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis or scoring systems thereof. The level of the biomarker can be determined by, for example, measuring, quantifying, and monitoring the expression level of the gene or mRNA encoding the biomarker and/or the peptide or protein of the biomarker. Non-limiting examples of biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis and/or scoring systems thereof include the aspartate aminotransferase (AST) to platelet ratio index (APRI); the aspartate aminotransferase (AST) and alanine aminotransferase (ALT) ratio (AAR); the FIB-4 score, which is based on the APRI, alanine aminotransferase (ALT) levels, and age of the subject (see, e.g., McPherson et al., Gut 2010, vol. 59(9), p. 1265-9, which is incorporated by reference herein in its entirety); hyaluronic acid; pro-inflammatory cytokines; a panel of biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST®, FIBROSURE®), a panel of biomarkers consisting of bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPAScore®; see, e.g., Adams et al., Clin. Chem. 2005, vol. 51(10), p. 1867-1873), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); a panel of biomarkers consisting of tissue inhibitor of

metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score, see, e.g., Lichtinghagen R, et al., J Hepatol. 2013 Aug;59(2):236-42, which is incorporated by reference herein in its entirety). In some embodiments, the presence of fibrosis is determined by one or more of the FIB-4 score, a panel of

5 biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST®, FIBROSURE®), a panel of biomarkers consisting of bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPAScore®; see, e.g., Adams et al., Clin. Chem. 2005,

10 vol. 51(10), p. 1867-1873), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); and a panel of biomarkers consisting of tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score). In some embodiments, the level of aspartate aminotransferase (AST) does not increase. In some

15 embodiments, the level of aspartate aminotransferase (AST) decreases. In some embodiments, the level of alanine aminotransferase (ALT) does not increase. In some embodiments, the level of alanine aminotransferase (ALT) decreases. In some embodiments, the "level" of an enzyme refers to the concentration of the enzyme, e.g., within blood. For example, the level of AST or ALT can be expressed as Units/L.

20 In some embodiments, the severity of fibrosis is determined by one or more of the FIB-4 score, a panel of biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST®, FIBROSURE®), a

25 panel of biomarkers consisting of bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPAScore®; see, e.g., Adams et al., Clin. Chem. 2005, vol. 51(10), p. 1867-1873, which is incorporated by reference herein in its entirety), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); and a panel of biomarkers consisting of tissue inhibitor of

30 metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score).

In some embodiments, hepatic inflammation is determined by the level of liver inflammation biomarkers, e.g., pro-inflammatory cytokines. Non-limiting examples of biomarkers indicative of liver

35 inflammation include interleukin-(IL) 6, interleukin-(IL) 1 β , tumor necrosis factor (TNF)- α ,

transforming growth factor (TGF)- β , monocyte chemotactic protein (MCP)-1, C-reactive protein (CRP), PAI-1, and collagen isoforms such as Col1a1, Col1a2, and Col4a1 (see, e.g., Neuman, et al., Can. J. Gastroenterol. Hepatol. 2014, vol. 28(11), p. 607-618 and U.S. Patent No. 9,872,844, each of which are incorporated by reference herein in their entireties). Liver inflammation can also be assessed by change of macrophage infiltration, e.g., measuring a change of CD68 expression level. In some embodiments, liver inflammation can be determined by measuring or monitoring serum levels or circulating levels of one or more of interleukin-(IL) 6, interleukin-(IL) 1 β , tumor necrosis factor (TNF)- α , transforming growth factor (TGF)- β , monocyte chemotactic protein (MCP)-1, and C-reactive protein (CRP).

In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis is determined for a sample from the subject prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% indicates treatment of NASH. In some embodiments, the decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis following administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis,

and/or liver cirrhosis during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%.

In some embodiments, the treatment of NASH decreases the level of serum bile acids in the subject. In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described in Danese et al., PLoS One. 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the NASH is NASH with attendant cholestasis. In cholestasis, the release of bile, including bile acids, from the liver is blocked. Bile acids can cause hepatocyte damage (see, e.g., Perez MJ, Briz O. World J. Gastroenterol. 2009, vol. 15(14), p. 1677-1689) likely leading to or increasing the progression of fibrosis (e.g., cirrhosis) and increasing the risk of hepatocellular carcinoma (see, e.g., Sorrentino P et al., *Dig. Dis. Sci.* 2005, vol. 50(6), p. 1130-1135 and Satapathy SK and Sanyal AJ. *Semin. Liver Dis.* 2015, vol. 35(3), p. 221-235, each of which are incorporated by reference herein in their entireties). In some embodiments, the treatment of NASH includes treatment of pruritus. In some embodiments, the treatment of NASH with attendant cholestasis includes treatment of pruritus. In some embodiments, a subject with NASH with attendant cholestasis has pruritus.

Exemplary biomarkers for NASH are provided in Table 7.

Table 7. Exemplary NASH biomarkers**Liver Fibrosis Biomarkers**

Aspartate aminotransferase (AST) to platelet ratio index (APRI)

Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) ratio (AAR)

FIB-4 score¹

Hyaluronic acid

Pro-inflammatory cytokines

A panel including α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST[®], FIBROSURE[®])

A panel including bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPAScore^{®2})

A panel including tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT[®])

A panel including tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score³)

Liver inflammation biomarkers^{4,5}

Interleukin-(IL) 6

Interleukin-(IL) 1 β

Tumor necrosis factor (TNF)- α

Transforming growth factor (TGF)- β

Monocyte chemotactic protein (MCP)-1

C-reactive protein (CRP)

PAI-1

Collagen isoforms (e.g., Col1a1, Col1a2, and Col4a1)

Change of macrophage infiltration (e.g., a change of CD68 expression level)

References for Table 7

¹ McPherson et al., Gut. 2010, vol. 59(9), p. 1265-1269.

² Adams, et al. Clin Chem. 2005, vol. 51(10), p. 1867-1873.

5 ³ Lichtinghagen, et al. J Hepatol. 2013, vol. 59(2), p. 236-242.

⁴ Neuman, et al. Can J Gastroenterol Hepatol. 2014, vol. 28(11), p. 607-618.

⁵ U.S. Patent No. 9,872,844

10 Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show a higher free fraction in plasma. In some embodiments, the free fraction is greater than about 0.2%, such as greater than about 0.4%, such as greater than about 0.6%, such as greater than about 0.8%, such as greater than about 1.0%, such as greater than about 1.25%, such as greater than about 1.5%, such as greater than about 1.75%, such as greater than about 2.0%, such as greater than about 2.5%, such as greater than about 3%, such as greater than about 4%, such as greater than about 5%, such as
15 greater than about 7.5%, such as greater than about 10%, or such as greater than about 20%.

20 Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may be excreted in urine. In some embodiments, the fraction of the compound that is excreted in urine is greater than about 0.2%, such as greater than about 0.4%, such as greater than about 0.6%, such as greater than about 0.8%, such as greater than about 1.0%, such as greater than about 2%, such as greater than about 3%, such as greater than about 5%, such as greater than about 7.5%, such as greater than about 10%, such as greater than about 15%, such as greater than about 20%, such as greater than about 30%, or such as greater than about 50%.

25 Following absorption from the intestine, some compounds of formula (I), or pharmaceutically acceptable salts thereof, may be circulated via the enterohepatic circulation. In some embodiments, the fraction of the compound that is circulated via the enterohepatic circulation is greater than about 0.1%, such as greater than about 0.2%, such as greater than about 0.3%, such as greater than about 0.5%, such as greater than about 1.0%, such as greater than about 1.5%, such as greater than about 2%, such as greater than about 3%, such as greater than about 5%, such as greater than about 7%,
30 such as greater than about 10%, such as greater than about 15%, such as greater than about 20%, such as greater than about 30% or such as greater than about 50%.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may cause renal excretion of bile salts. In some embodiments, the fraction of circulating bile acids that is excreted by the renal route is greater than about 1 %, such as greater than about 2%, such as greater than about 5%, such as greater than about 7%, such as greater than about 10%, such as greater than about 15%,
5 such as greater than about 20%, or such as greater than about 25%.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show improved or optimal permeability. The permeability may be measured in Caco2 cells, and values are given as Papp (apparent permeability) values in cm/s. In some embodiments, the permeability is greater than at
10 least about 0.1×10^{-6} cm/s, such as greater than about 0.2×10^{-6} cm/s, such as greater than about 0.4×10^{-6} cm/s, such as greater than about 0.7×10^{-6} cm/s, such as greater than about 1.0×10^{-6} cm/s, such as greater than about 2×10^{-6} cm/s, such as greater than about 3×10^{-6} cm/s, such as greater than about 5×10^{-6} cm/s, such as greater than about 7×10^{-6} cm/s, such as greater than about 10×10^{-6} cm/s, such as greater than about 15×10^{-6} cm/s.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show an improved or optimal bioavailability. In some embodiments, the oral bioavailability is greater than about 5%, such as greater than about 7%, such as greater than about 10%, such as greater than about 15%, such as greater than about 20%, such as greater than about 30%, such as greater than
20 about 40%, such as greater than about 50 %, such as greater than about 60 %, such as greater than about 70% or such as greater than about 80%. In other embodiments, the oral bioavailability is between about 10 and about 90%, such as between about 20 and about 80%, such as between about 30 and about 70% or such as between about 40 and about 60%.

25 Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may be a substrate to relevant transporters in the kidney.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may give rise to concentrations of bile acids in the intestine, the liver and in serum that do not cause adverse
30 gastrointestinal effects.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may decrease the concentration of bile acids in the liver without causing gastrointestinal disorders such as diarrhoea.

As used herein, the terms "treatment", "treat" and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, a base-addition salt of a compound of the invention which is sufficiently acidic, such as an alkali metal salt (e.g., a sodium or potassium salt), an alkaline earth metal salt (e.g., a calcium or magnesium salt), an ammonium salt, or a salt with an organic base which affords a physiologically acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may have chiral centres and/or geometric isomeric centres (E- and Z-isomers). It is to be understood that the invention encompasses all such optical isomers, diastereoisomers and geometric isomers that possess ASBT and/or LBAT inhibitory activity. The invention also encompasses any and all tautomeric forms of compounds of formula (I), or pharmaceutically acceptable salts thereof, that possess ASBT and/or LBAT inhibitory activity. Certain compounds of formula (I), or pharmaceutically acceptable salts thereof, may exist in unsolvated as well as solvated forms, such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms that possess ASBT and/or LBAT inhibitory activity.

In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients. The excipients may e.g. include fillers, binders, disintegrants, glidants and lubricants. In general, pharmaceutical compositions may be prepared in a conventional manner using conventional excipients.

Examples of suitable fillers include, but are not limited to, dicalcium phosphate dihydrate, calcium sulfate, lactose (such as lactose monohydrate), sucrose, mannitol, sorbitol, cellulose, microcrystalline

cellulose, dry starch, hydrolyzed starches and pregelatinized starch. In certain embodiments, the filler is mannitol and/or microcrystalline cellulose.

Examples of suitable binders include, but are not limited to, starch, pregelatinized starch, gelatin, sugars (such as sucrose, glucose, dextrose, lactose and sorbitol), polyethylene glycol, waxes, natural and synthetic gums (such as acacia gum and tragacanth gum), sodium alginate, cellulose derivatives (such as hydroxypropylmethylcellulose (or hypromellose), hydroxypropylcellulose and ethylcellulose) and synthetic polymers (such as acrylic acid and methacrylic acid copolymers, methacrylic acid copolymers, methyl methacrylate copolymers, aminoalkyl methacrylate copolymers, polyacrylic acid/polymethacrylic acid copolymers and polyvinylpyrrolidone (povidone)). In certain embodiments, the binder is hydroxypropylmethylcellulose (hypromellose).

Examples of suitable disintegrants include, but are not limited to, dry starch, modified starch (such as (partially) pregelatinized starch, sodium starch glycolate and sodium carboxymethyl starch), alginic acid, cellulose derivatives (such as sodium carboxymethylcellulose, hydroxypropyl cellulose, and low substituted hydroxypropyl cellulose (L-HPC)) and cross-linked polymers (such as carmellose, croscarmellose sodium, carmellose calcium and cross-linked PVP (crospovidone)). In certain embodiments, the disintegrant is croscarmellose sodium.

Examples of suitable glidants and lubricants include, but are not limited to, talc, magnesium stearate, calcium stearate, stearic acid, glyceryl behenate, colloidal silica, aqueous silicon dioxide, synthetic magnesium silicate, fine granulated silicon oxide, starch, sodium lauryl sulfate, boric acid, magnesium oxide, waxes (such as carnauba wax), hydrogenated oil, polyethylene glycol, sodium benzoate, polyethylene glycol, and mineral oil. In certain embodiments, the glidant or lubricant is magnesium stearate or colloidal silica.

The pharmaceutical composition may be conventionally coated with one or more coating layers. Enteric coating layers or coating layers for delayed or targeted release of the compound of formula (I), or pharmaceutically acceptable salts thereof, are also contemplated. The coating layers may comprise one or more coating agents, and may optionally comprise plasticizers and/or pigments (or colorants).

Example of suitable coating agents include, but are not limited to, cellulose-based polymers (such as ethylcellulose, hydroxypropylmethylcellulose (or hypromellose), hydroxypropylcellulose, cellulose acetate phthalate, cellulose acetate succinate, hydroxypropyl methylcellulose acetate succinate and

hydroxypropyl methylcellulose phthalate), vinyl-based polymers (such as polyvinyl alcohol) and polymers based on acrylic acid and derivatives thereof (such as acrylic acid and methacrylic acid copolymers, methacrylic acid copolymers, methyl methacrylate copolymers, aminoalkyl methacrylate copolymers, polyacrylic acid/polymethacrylic acid copolymers). In certain embodiments, the coating agent is hydroxypropylmethylcellulose. In other embodiments, the coating agent is polyvinyl alcohol.

Examples of suitable plasticizers include, but are not limited to, triethyl citrate, glyceryl triacetate, tributyl citrate, diethyl phthalate, acetyl tributyl citrate, dibutyl phthalate, dibutyl sebacate and polyethylene glycol. In certain embodiments, the plasticizer is polyethylene glycol.

Examples of suitable pigments include, but are not limited to, titanium dioxide, iron oxides (such as yellow, brown, red or black iron oxides) and barium sulfate.

The pharmaceutical composition may be in a form that is suitable for oral administration, for parenteral injection (including intravenous, subcutaneous, intramuscular and intravascular injection), for topical administration or for rectal administration. In a preferred embodiment, the pharmaceutical composition is in a form that is suitable for oral administration, such as a tablet or a capsule.

The dosage required for the therapeutic or prophylactic treatment will depend on the route of administration, the severity of the disease, the age and weight of the patient and other factors normally considered by the attending physician, when determining the appropriate regimen and dosage level for a particular patient.

The amount of the compound to be administered will vary for the patient being treated, and may vary from about 1 µg/kg of body weight to about 50 mg/kg of body weight per day. A unit dose form, such as a tablet or capsule, will usually contain about 1 to about 250 mg of active ingredient, such as about 1 to about 100 mg, or such as about 1 to about 50 mg, or such as about 1 to about 20 mg, e.g. about 2.5 mg, or about 5 mg, or about 10 mg, or about 15 mg. The daily dose can be administered as a single dose or divided into one, two, three or more unit doses. An orally administered daily dose of a bile acid modulator is preferably within about 0.1 to about 250 mg, more preferably within about 1 to about 100 mg, such as within about 1 to about 5 mg, such as within about 1 to about 10 mg, such as within about 1 to about 15 mg, or such as within about 1 to about 20 mg.

In another aspect, the invention relates to a compound of formula (I), or a pharmaceutically acceptable salt thereof, for use as a medicament. The invention also relates to the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as a medicament.

5 In another aspect, the invention relates to a compound of formula (I), or a pharmaceutically acceptable salt thereof, for use in the treatment or prevention of any of the diseases recited herein. The invention also relates to the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment or prevention of any of the diseases recited herein. The invention also relates to a method of treating or preventing any of the
10 diseases recited herein in a subject, such as man, comprising administering to the subject in need of such treatment or prevention a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

Combination therapy

15 In one aspect of the invention, the compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with at least one other therapeutically active agent, such as with one, two, three or more other therapeutically active agents. The compound of formula (I), or a pharmaceutically acceptable salt thereof, and the at least one other therapeutically active agent may
20 be administered simultaneously, sequentially or separately. Therapeutically active agents that are suitable for combination with the compounds of formula (I) include, but are not limited to, known active agents that are useful in the treatment of any of the aforementioned conditions, disorders and diseases.

25 In one embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with another ASBT inhibitor. Suitable ASBT inhibitors are disclosed in WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/03818, WO 98/07449, WO 98/40375, WO 99/35135, WO 99/64409, WO 99/64410, WO 00/47568, WO 00/61568, WO 00/38725, WO 00/38726, WO 00/38727,
30 WO 00/38728, WO 00/38729, WO 01/66533, WO 01/68096, WO 02/32428, WO 02/50051, WO 03/020710, WO 03/022286, WO 03/022825, WO 03/022830, WO 03/061663, WO 03/091232, WO 03/106482, WO 2004/006899, WO 2004/076430, WO 2007/009655, WO 2007/009656, WO 2011/137135, DE 19825804, EP 864582, EP 489423, EP 549967, EP 573848, EP 624593, EP 624594, EP 624595, EP 624596, EP 0864582, EP 1173205 and EP 1535913, all of which are
35 incorporated herein by reference in their entireties.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a bile acid binder (also referred to as a bile acid sequestrant, or a resin), such as colesevelam, cholestyramine or cholestipol. In a preferred embodiment of such a combination, the bile acid binder is formulated for colon release. Examples of such formulations are disclosed in e.g. WO 2017/138877, WO 2017/138878, WO 2019/032026 and WO 2019/032027, all of which are incorporated herein by reference in their entireties.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a DPP-IV inhibitor, including gliptins such as sitagliptin, vildagliptin, saxagliptin, linagliptin, gemigliptin, anagliptin, teneligliptin, alogliptin, trelagliptin, omarigliptin, evogliptin, gosogliptin and dutogliptin, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an HMG CoA reductase inhibitor, such as fluvastatin, lovastatin, pravastatin, simvastatin, atorvastatin, pitavastatin cerivastatin, mevastatin, rosuvastatin, bervastatin or dalvastatin, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a cholesterol absorption inhibitor such as ezetimibe, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PPAR alpha agonist, including fibrates such as clofibrate, bezafibrate, ciprofibrate, clinofibrate, clofibrade, fenofibrate, gemfibrozil, ronifibrate and simfibrate, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PPAR gamma agonist, including thiazolidinediones such as pioglitazone, rosiglitazone and lobeglitazone, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual PPAR alpha/gamma agonist, including glitazars such as saroglitazar, aleglitazar, muraglitazar or tesaglitazar, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual PPAR alpha/delta agonist, such as elafibranor.

5 In yet another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a pan PPAR agonist (i.e. a PPAR agonist that has activity across all subtypes: α , γ and δ), such as IVA337.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a farnesoid X receptor (FXR) modulators, including FXR agonists such as cafestol, chenodeoxycholic acid, 6 α -ethyl-chenodeoxycholic acid (obeticholic acid; INT-747), fexaramine, tropifexor, cilofexor and MET409.

15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a TGR5 receptor modulator, including TGR5 agonists such as 6 α -ethyl-23(S)-methylcholic acid (INT-777).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual FXR/TGR5 agonist such as INT-767.

20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with ursodeoxycholic acid (UDCA). In yet another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with nor-ursodeoxycholic acid (nor-UDCA).

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an FGF19 modulator, such as NGM282.

30 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an FGF21 agonist, such as BMS-986036.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an integrin inhibitor, such as PLN-74809 and PLN-1474.

35 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a CCR2/CCR5 inhibitor, such as cenicriviroc.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a caspase protease inhibitor, such as emricasan.

- 5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a galectin-3 inhibitor, such as GR-MD-02.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a stearyl-CoA desaturase (SCD) Inhibitor, such as aramchol

- 10 (arachidyl amido cholanoic acid).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an apoptosis signal-regulating kinase 1 (ASK1) inhibitor, such as selonsertib.

15

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an LOXL2 inhibitor, such as simtuzumab.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an ACC inhibitor, such as GS-0976.

20

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a thyroid hormone receptor- β agonist, such as MGL3196.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a GLP-1 agonist such as liraglutide.

25

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual glucagon-like peptide and glucagon receptor agonists, such as SAR425899.

30

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a mitochondrial pyruvate carrier inhibitor, such as MSDC-0602K.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an anti-oxidant agent, such as vitamin E.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an SGLT1 inhibitor, an SGLT2 inhibitor or a dual SGLT1 and SGLT2 inhibitor. Examples of such compounds are dapagliflozin, sotagliflozin, canagliflozin, empagliflozin, LIK066 and SGL5213.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a diacylglycerol O-Acyltransferase 2 (DGAT2) inhibitor, such as DGAT2RX and PF-06865571.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a fatty acid synthase (FASN) Inhibitor, such as TVB-2640.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an AMP-activated protein kinase (AMPK) activator, such as PXL-770.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glucocorticoid receptor antagonist (GR), a mineralocorticoid receptor antagonist (MR), or a dual GR/MR antagonist. Examples of such compounds are MT-3995 and CORT-118335.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a cannabinoid receptor 1 (CB1) antagonist, such as IM102.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a Klotho β (KLB) and fibroblast growth factor receptor (FGFR) activator, such as MK-3655 (previously known as NGM-313).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chemokine (c-c motif) ligand 24 (CCL24) inhibitor, such as CM101.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an A3 antagonist, such as PBF-1650.

5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a P2x7 receptor antagonist, such as SGM 1019.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with P2Y13 receptor agonists, such as CER-209.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sulfated oxysterol, such as Dur-928.

15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a leukotriene D4 (LTD4) receptor antagonist, such as MN-001.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a type 1 natural killer T cell (NKT1) inhibitor, such as GRI-0621.

20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an anti-lipopolysaccharide (LPS) compound, such as IMM-124E.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a VAP1 inhibitor, such as BI1467335.

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an A3 adenosine receptor agonist, such as CF-102.

30 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a SIRT-1 activator, such as NS-20.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a nicotinic acid receptor 1 agonist, such as ARI-3037MO.

35 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a TLR4 antagonist, such as JKB-121.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a ketohexokinase inhibitor, such as PF-06835919.

- 5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an adiponectin receptor agonist, such as ADP-335.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an autotaxin inhibitor, such as PAT-505 and PF8380.

10

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chemokine (c-c motif) receptor 3 (CCR3) antagonist, such as bertilimumab.

- 15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chloride channel stimulator, such as cobiprostone and lubiprostone.

- 20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a heat shock protein 47 (HSP47) inhibitor, such as ND-L02-s0201.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sterol regulatory element-binding protein (SREBP) transcription factor inhibitor, such as CAT-2003 and MDV-4463.

25

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a biguanidine, such as metformin.

- 30 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with insulin.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glycogen phosphorylase inhibitor and/or a glucose-6-phosphatase inhibitor.

35

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sulfonylurea, such as glipizid, glibenklamid and glimepirid.

5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a meglitinide, such as repaglinide, nateglinide and ormgilitinide.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glucosidase inhibitor, such as acarbose or miglitol.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a squalene synthase inhibitor, such as TAK-475.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PTPB1 inhibitor, such as trodusquemine, ertiprotafib, JTT-551
15 and claramine.

Preparation of compounds

The compounds of the invention can be prepared as a free acid or a pharmaceutically acceptable salt
20 thereof by the processes described below. Throughout the following description of such processes it is understood that, where appropriate, suitable protecting groups will be added to, and subsequently removed from the various reactants and intermediates in a manner that will be readily understood by one skilled in the art of organic synthesis. Conventional procedures for using such protecting groups as well as examples of suitable protecting groups are for example described in *Greene's*
25 *Protective Groups in Organic Synthesis* by P.G.M Wutz and T.W. Greene, 4th Edition, John Wiley & Sons, Hoboken, 2006.

General methods

30 All solvents used were of analytical grade. Commercially available anhydrous solvents were routinely used for reactions. Starting materials were available from commercial sources or prepared according to literature procedures. 3,3-Dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide may be prepared as described in WO 03/022286 (method 24). Room temperature refers to 20 - 25 °C. Solvent mixture compositions
35 are given as volume percentages or volume ratios.

LCMS:

Instrument name: Agilent 1290 infinity II.

Method A: Mobile phase: A: 0.1% HCOOH in H₂O: ACN (95:5), B: ACN; flow rate: 1.5 mL/min; column:
5 ZORBAX XDB C-18 (50 x 4.6 mm) 3.5 µM.

Method B: Mobile phase: A: 10 mM NH₄HCO₃ in water, B: ACN; flow rate: 1.2 mL/min; column:
XBridge C8 (50 x 4.6 mm), 3.5 µM.

Method C: Mobile phase: A: 0.1% HCOOH in water: ACN (95:5), B: ACN; flow rate: 1.5 mL/min;
column: ATLANTIS dC18 (50 x 4.6 mm), 5 µM.

10 **Method D:** Mobile phase: A: 10 mM NH₄OAc in water, B: ACN; flow rate: 1.2 mL/min; column: Zorbax
Extend C18 (50 x 4.6mm) 5 µM.

Method E: Mobile phase: A: 0.1% TFA in water: ACN (95:5), B: 0.1% TFA in ACN; flow rate: 1.5
mL/min; Column: XBridge C8 (50 x 4.6 mm), 3.5 µM.

Method F: Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow Rate: 0.8 mL/min; column:
15 ZORBAX ECLIPSE PLUS C18 (50 x 2.1 mm), 1.8 µM.

Method G: Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow Rate: 0.8 mL/min; column:
Acquity UPLC BEH C18 (2.1 x 50 mm), 1.7 µm.

UPLC:

20 **Instrument name:** waters Acquity I Class

Method A: Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate: 0.8 mL/min;
Column: Acquity UPLC HSS T3 (2.1 x 50) mm; 1.8 µm.

HPLC:

25 **Instrument name:** Agilent 1260 Infinity II series instruments as followed using % with UV detection
(maxplot).

Method A: Mobile phase: A: 10 mM NH₄HCO₃ in water, B: ACN; flow rate: 1.0 mL/min; column:
XBridge C8 (50 x 4.6 mm, 3.5 µm).

Method B: Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow rate: 2.0 mL/min; column:
30 XBridge C8 (50 x 4.6 mm, 3.5 µm).

Method C: Mobile phase: A: 10 mM NH₄OAc in milli-q water, B: ACN; flow rate: 1.0 mL/min; column:
Phenomenex Gemini C18 (150 x 4.6 mm, 3.0 µm).

Method D: Mobile phase: A: 0.1% TFA in water, B: ACN; flow rate: 1.0 mL/min; column: ATLANTIS
dC18 (250 x 4.6 mm, 5.0 µm).

35

Chiral HPLC:

Instrument name: Agilent 1260 Infinity II

Method A: Mobile phase: A: 0.1% TFA in *n*-hexane; B: ethanol, flow: 1.0 mL/min; Column: CHIRALPAK IA (250 x 4.6 mm, 5.0 µm).

5

Chiral SFC:

Instrument name: PIC SFC 10 (analytical)

Ratio between CO₂ and co-solvent is ranging between 60:40 and 80:20

10 **Method A:** Mobile phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: YMC Amylose-SA (250 x 4.6 mm, 5 µm).

Method B: Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralpak AD-H (250 x 4.6 mm, 5 µm).

Method C: Mobile Phase: 20 mM ammonia in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC (250 x 4.6 mm, 5 µm).

15 **Method D:** Mobile Phase: methanol; flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

Method E: Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 5 mL/min; column: Lux C4.

Method F: Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC.

Method G: Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Lux A1.

20 **Method H:** Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

Method I: Mobile phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 4.6 mm, 5 µm).

25 **Method J:** Mobile phase: 0.5% isopropylamine in IPA; flow rate: 5 mL/min; column: YMC Cellulose-SC AD-H (250 x 4.6 mm, 5 µm).

Method K: Mobile phase: 0.5% Isopropylamine in methanol; flow rate: 4 mL/min; column: (R,R)-Whelk-01 (250 x 4.6 mm, 5 µm).

Method L: Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralcel OX-H (250 x 4.6 mm, 5 µm).

30 **Method M:** Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 5 mL/min; column: YMC Cellulose-SC (250 x 4.6 mm, 5 µm).

Method N: Mobile phase: methanol, flow rate: 5 mL/min; column: Chiralcel OX-H (250 x 4.6 mm, 5 µm).

Prep-HPLC:**Instrument name:** Agilent 1290 Infinity II**Method A:** Mobile phase: A: 0.1% TFA in water; Mobile phase; B: 0.1% TFA in CAN; flow rate: 2.0 mL/min; Column: X-Bridge C8 (50 X 4.6 mm, 3.5 µm).

- 5 **Method B:** Mobile phase: A: 10 mM NH₄OAc in water; B: ACN; flow rate: 35 mL/min; column: X select C18 (30 x 150 mm, 5 µm).

Method C: Mobile phase: A: 10 mM NH₄HCO₃ in water; B: ACN; flow rate: 1.0 mL/min; column: XBridge C8 (50 x 4.6 mm, 3.5 µm).

- 10 **Method D:** Mobile phase: A: 0.1% HCOOH in water; B: ACN; flow rate: 1.0 mL/min; column: X-select C18 (30 x 150 mm, 5 µm).

Chiral Preparative SFC:**Instrument name:** PIC SFC 100 and PSC SFC 400Ratio between CO₂ and co-solvent is ranging between 60:40 and 80:20

- 15 **Method A:** Mobile phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: YMC Amylose-SA (250 x 30 mm, 5 µm).

Method B: Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralpak AD-H (250 x 30 mm, 5 µm).

- 20 **Method C:** Mobile phase: 20 mM ammonia in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC (250 x 30 mm, 5 µm).

Method D: Mobile phase: methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 30 mm, 5 µm).**Method E:** Mobile phase: methanol; flow rate: 3 mL/min; column: Lux A1 (250 x 30 mm, 5 µm).**Method F:** Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Lux A1 (250 x 30 mm, 5 µm).

- 25 **Method G:** Mobile phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 30 mm, 5 µm).

Method H: Mobile Phase: 0.5% isopropylamine in IPA, flow rate: 5 mL/min; column: YMC Amylose-SC (250 x 30 mm, 5 µm).

- 30 **Method J:** Mobile phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralcel OX-H (250 x 30 mm, 5 µm).

Method K: Mobile phase: 0.5% isopropylamine in methanol; flow rate: 5 mL/min; column: YMC Cellulose-SC (250 x 30 mm, 5 µm).**Method L:** Mobile phase: methanol; flow rate: 5 mL/min; column: Chiralcel OX-H (250 x 30 mm, 5 µm).

35

Chiral Preparative HPLC:**Instrument name:** Agilent 1260 Infinity II**Method A:** Mobile phase: A: 0.1% TFA in *n*-hexane; B: ethanol; flow rate: 15 mL/min; Column: CHIRALPAK IA (250 x 19 mm, 5.0 µm).

5

Abbreviations


	ACN	acetonitrile
	DCM	dichloromethane
10	DMAP	4-dimethylaminopyridine
	DMF	dimethylformamide
	IPA	isopropyl alcohol
	LCMS	liquid chromatography - mass spectrometry
	HPLC	high-performance liquid chromatography
15	PE	petroleum ether
	SFC	supercritical fluid chromatography
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	TLC	thin layer chromatography
20	UPLC	ultra performance liquid chromatography

The invention will now be described by the following examples which do not limit the invention in any respect. All cited documents and references are incorporated by reference.

EXAMPLES

Intermediate 1

Ethyl 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoate




To a stirred suspension of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (12 g, 21.1 mmol) in ethyl acrylate (80 mL) at room temperature, DMAP (257 mg, 2.11 mmol) was added. The reaction mixture was heated in a sealed tube for 72 hours at 100 °C. The progress of the reaction was monitored TLC, which indicated the incomplete conversion (~40%) of the starting material. The reaction mixture was evaporated and dried under vacuum to afford the crude title compound, which was forwarded to the next step without any further purification. **Yield:** 15 g (crude, brown gum).

LCMS: (Method A) 669.3 ($M^+ + H$), Rt. 3.66 min, 36.45% (Max).

Intermediate 2

3-((3,3-Dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid



Preparation 1:

To a stirred solution of ethyl 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoate (Intermediate 1; 40 g, 59.79 mmol) in 1,4-dioxane (200 mL) at room temperature, HCl (6N, 200 ml) was added dropwise and the reaction mixture was heated for 16 hours at 80 °C. After completion of the reaction (monitored by LCMS), the reaction mixture was diluted with ice cold water (500 mL) and the aqueous layer was extracted with EtOAc (2 x 200 mL). The organic part was dried over anhydrous Na₂SO₄ and evaporated under vacuum. The resulting crude material was purified by flash column chromatography (eluent: 20-70 % EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


Yield: 39% (15 g, brown gum).

LCMS: (Method E) 641.3 (M⁺+H), Rt. 2.93 min, 75.38% (Max).

Preparation 2:

To a stirred suspension of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (500 mg, 0.88 mmol) in THF (5 mL) at 0 °C, potassium *tert*-butoxide (99 mg, 0.88 mmol) was added and the reaction mixture was stirred for 10 minutes at room temperature. Then β-propiolactone (76 mg, 1.05 mmol) was added to the reaction mixture at 0° C and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with 1.5N HCl at 0 °C and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was dried over anhydrous Na₂SO₄ and evaporated under vacuum. The resulting crude product was purified by flash column chromatography (eluent: 10-50% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 53% (300 mg, white solid).

LCMS: (Method D) 641.3 (M⁺+H), Rt. 3.48 min, 96.63% (Max).

Intermediate 3**Ethyl 2-aminobutanoate hydrochloride**


To a stirred solution of 2-aminobutanoic acid (100 g, 0.97 mol) in ethanol (750 mL), thionyl chloride (78 mL, 1.07 mol) was added at 0 °C. The reaction mixture was then heated for 16 hours at 80 °C. After completion of the reaction, the reaction mixture was concentrated under vacuum to afford the crude title compound which was used as such for the next step without any further purification.

Yield: 93% (152 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 8.66 (bs, 3H), 4.25-4.16 (m, 2H), 3.98-3.85 (m, 1H), 1.84 (t, *J* = 7.2 Hz, 2H), 1.23 (t, *J* = 6.8 Hz, 3H), 0.92 (t, *J* = 7.6 Hz, 3H).

Intermediate 4

5 Ethyl (*E*)-2-(benzylideneamino) butanoate




To a stirred solution of ethyl 2-aminobutanoate hydrochloride (Intermediate 3; 152 g, 0.91 mol) in DCM (900 mL), triethyl amine (152 mL, 1.09 mol) was added at 0 °C over a period of 30 minutes. Magnesium sulfate (98 g, 0.82 mol) was added portionwise to the reaction mixture at 0 °C.

- 10 Benzaldehyde (84 mL, 0.82 mol) was then added to the reaction mixture at 0 °C over a period of 20 minutes and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was filtered through celite and the filtrate was concentrated under vacuum. The resulting crude was dissolved in petroleum ether (1000 mL) and again filtered through celite. The filtrate was then concentrated under vacuum to afford the title
- 15 compound. This crude material was forwarded as such to the next step without any further purification. **Yield:** 90% (180 g, pale brown liquid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 8.40 (s, 1H), 7.79-7.76 (m, 2H), 7.49-7.47 (m, 3H), 4.16-4.10 (m, 2H), 3.98-3.95 (m, 1H), 1.92-1.89 (m, 1H), 1.79-1.74 (m, 1H), 1.19 (t, *J* = 7.2 Hz, 3H), 0.85 (t, *J* = 7.2 Hz, 3H).

20 Intermediate 5

Ethyl (*E*)-2-(benzylideneamino)-2-ethylhexanoate




- To a stirred solution of NaH (60%; 32.8 g, 0.82 mol) in DMF (100 mL) at 0 °C, ethyl (*E*)-2-(benzylideneamino)butanoate (Intermediate 4; 180 g, 0.82 mol) in DMF (800 mL) was slowly added
- 25 over a period of 30 minutes. The reaction mixture was then stirred for 1.5 hours at room temperature. *n*-Butyl iodide (93 mL, 0.82 mol) was added to the reaction mixture at 0 °C and the mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with 2-propanol (100 mL) at 0 °C and then diluted with water (1000 mL). The aqueous layer was extracted with petroleum ether (1000 mL). The organic

layer was washed with brine (200 mL) and dried over anhydrous Na_2SO_4 . The organic part was concentrated under vacuum and the resulting crude material was forwarded as such to the next step without any further purification. **Yield:** 88% (200 g, yellow liquid).

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.34 (s, 1H), 7.80 - 7.77 (m, 2H), 7.47-7.44 (m, 3H), 4.16 (q, J = 7.0 Hz, 2H), 2.51-1.79 (m, 4H), 1.31-1.18 (m, 7H), 0.88 - 0.84 (m, 6H).

Intermediate 6

Ethyl 2-amino-2-ethylhexanoate




To a stirred solution of ethyl (*E*)-2-(benzylideneamino)-2-ethylhexanoate (Intermediate 5; 200 g, 0.73 mol) in petroleum ether (500 mL), dilute HCl (1000 mL, 1.5 N) was added at 0 °C and the reaction mixture was stirred vigorously for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the organic layer was separated and the aqueous layer was washed with EtOAc (2 x 100 mL). The aqueous layer was then basified (pH ~8.5) by using solid sodium bicarbonate (200 g) and extracted with EtOAc (2 x 200 mL). The organic layer was washed with water (2 x 15 mL). The combined organic part was dried over anhydrous Na_2SO_4 and concentrated under vacuum to afford the title compound. The crude material was forwarded as such to the next step without any further purification. **Yield:** 80% (110 g, pale yellow liquid).

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 4.08 (q, J = 7.1 Hz, 2H), 1.68 - 1.00 (m, 13H), 0.85 (t, J = 7.2 Hz, 3H), 0.77 (t, J = 7.4 Hz, 3H).

Intermediate 7

2-Amino-2-ethyl-N-phenylhexanamide




To a stirred solution of aniline (48.3 mL, 534 mmol) in THF (250 mL) at -78 °C, *n*-BuLi (2.6M in hexanes; 205 mL, 534 mmol) was added dropwise over a period of 30 minutes, and the reaction mixture was stirred for 45 minutes at -25 °C to -30 °C. Then ethyl 2-amino-2-ethylhexanoate (Intermediate 6; 50 g, 267 mmol) in THF (250 mL) was added to the reaction mixture at -78 °C and

the reaction mixture was stirred for 2 hours at -78 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (500 mL) at -78 °C. The reaction mixture was extracted with EtOAc (2 x 250 mL) and the organic layer was washed with water (2 x 15 mL). The organic part was dried over anhydrous Na₂SO₄ and concentrated under vacuum to afford the title compound as crude. The crude product was dissolved in petroleum ether (1000 mL). The organic part was washed with 30% methanol in water (2 x 250 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude was forwarded as such to the next step without any further purification. **Yield:** 66 g (crude, brown liquid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.64 (d, *J* = 8.4 Hz, 2H), 7.30 (t, *J* = 7.4 Hz, 2H), 7.05 (t, *J* = 7.4 Hz, 1H), 6.55 (d, *J* = 8.5 Hz, 1H), 1.76-1.07 (m, 10H), 0.86-0.77 (m, 6H).

Intermediate 8

2-Ethyl-N1-phenylhexane-1,2-diamine




To a stirred solution of 2-amino-2-ethyl-N-phenylhexanamide (Intermediate 7; 66 g, 0.28 mol) in THF (600 mL), borane dimethylsulfide (2M in THF, 253 mL, 0.51 mol) was added at 0 °C and the reaction mixture was heated for 16 hours at 70 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with methanol (300 mL) at 0 °C. The reaction mixture was then heated for 2 hours at 70 °C. The reaction mixture was concentrated under vacuum and the obtained residue was dissolved in EtOAc (1000 mL). The organic layer was washed with water (2 x 150 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 40% EtOAc in hexane; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 82% (50 g, brown liquid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.04 (t, *J* = 7.2 Hz, 2H), 6.61 (d, *J* = 8.4 Hz, 2H), 6.49 (t, *J* = 7.2 Hz, 1H), 5.15 (t, *J* = 4.8 Hz, 1H), 2.79 (d, *J* = 5.6 Hz, 2H), 1.39 - 1.17 (m, 10H), 0.88-0.79 (m, 6H).

Intermediate 9


1,2-Bis(2,4-dibromo-5-methoxyphenyl)disulfane



To a stirred solution of 3-methoxybenzenethiol (100 g, 0.7 mol) in methanol (1000 mL), bromine (73 mL, 1.4 mol) was added dropwise at 0 °C and the reaction mixture was stirred for 24 hours at room temperature. The reaction mixture was evaporated under vacuum and the obtained crude was diluted with EtOAc (2000 mL) and washed with water (2 x 500 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was dissolved in glacial acetic acid (600 mL), bromine (20 mL) was added dropwise at room temperature and the reaction mixture was stirred for 2 hours at room temperature. The obtained solid was filtered off, triturated with DCM and dried under vacuum to afford the pure title compound. **Yield:** 37% (78 g, white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.69 (s, 2H), 7.17 (s, 2H), 3.84 (s, 6H).

Intermediate 10

2,4-Dibromo-5-methoxybenzenesulfonyl chloride




To a stirred suspension of 1,2-bis(2,4-dibromo-5-methoxyphenyl)disulfane (Intermediate 9; 20.0 g, 33.67 mmol) and potassium nitrate (17.02 g, 168.35 mmol) in acetonitrile (200 mL) was dropwise added sulfonyl chloride (13.6 mL, 168.35 mmol) at 0 °C. The reaction mixture was stirred for 24 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was poured into crushed ice and the solid obtained was filtered off. The solid was washed with water and dried under vacuum to afford the pure title compound. **Yield:** 91% (22.5 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 8.05 (s, 1H), 7.66 (s, 1H), 4.01 (s, 3H).

Intermediate 11

2,4-Dibromo-5-methoxy-N-(3-((phenylamino)methyl)heptan-3-yl)benzenesulfonamide




To a stirred solution of 2-ethyl-N1-phenylhexane-1,2-diamine (Intermediate 8; 4.9 g, 22.34 mmol) in THF (10 mL) were added 2,4-dibromo-5-methoxybenzenesulfonyl chloride (Intermediate 10; 10.5 g, 28.91 mmol) and triethylamine (9.3 mL, 67.02 mmol) at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction

mixture was diluted with EtOAc (50 mL). The organic layer was washed with water (2 x 15 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 59% (7.2 g, white solid).


- 5 ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.01 (s, 1H), 7.60 (s, 1H), 7.50 (s, 1H), 7.03 (t, *J* = 8.1 Hz, 2H), 6.54 - 6.46 (m, 3H), 4.80 (t, *J* = 5.1 Hz, 1H), 3.86 (s, 3H), 3.07-2.96 (m, 2H), 1.66-1.41 (m, 4H), 1.15-0.95 (m, 4H), 0.78-0.69 (m, 6H).

Intermediate 12

- 10 **7-Bromo-3-butyl-3-ethyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**




- To a stirred solution of 2,4-dibromo-5-methoxy-*N*-(3-((phenylamino)methyl)heptan-3-yl)benzenesulfonamide (Intermediate 11; 7.2 g, 13.1 mmol) in DMF (50 mL) were added potassium carbonate (3.62 g, 26.2 mmol) and copper powder (834 mg, 13.1 mmol) and the reaction mixture was heated for 24 hours at 150 °C. After completion of the reaction (monitored by TLC), the reaction mixture was filtered through celite and washed with EtOAc (25 mL). The filtrate part was concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 83% (5.1 g, white solid).
- 15 ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.43-7.30 (m, 4H), 7.15-7.13 (m, 2H), 7.03-7.01 (m, 2H), 4.00-3.60 (m, 5H), 1.62-1.34 (m, 4H), 1.08-0.95 (m, 4H), 0.74-0.71 (m, 6H). **LCMS:** (Method A) 467.0 (M⁺), Rt. 3.06 min, 95.31% (max).
- 20

Intermediate 13**7-Bromo-3-butyl-3-ethyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

- 5 To a stirred solution of 7-bromo-3-butyl-3-ethyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 12; 20.0 g, 42.7 mmol) in *N*-methyl-2-pyrrolidone (100 mL) were added Cs₂CO₃ (27.8 g, 85.5 mmol) and *p*-methoxybenzyl bromide (7.98 mL, 39.5 mmol) at 0 °C and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with EtOAc (200 mL) and the organic
- 10 layer was washed with water (2 x 50 mL). The organic part was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 64% (16 g, white solid).

LCMS: (Method A) 587.2 (M⁺), Rt. 3.51 min, 92.94% (max).

15

Intermediate 14**3-Butyl-3-ethyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**


- 20 To a stirred solution of 7-bromo-3-butyl-3-ethyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 13; 16.0 g, 27.2 mmol) in DMF (120

mL), sodium thiomethoxide (9.5 g, 136.1 mmol) was added and the reaction mixture was heated for 16 hours at 60 °C. After completion of reaction (monitored by LCMS), the reaction mixture was diluted with EtOAc (200 mL) and the organic layer was washed with water (2 x 50 mL). The organic part was dried over anhydrous Na₂SO₄, then concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 65% (9.2 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 10.37 (bs, 1H), 7.31-7.22 (m, 5H), 7.01-6.65 (m, 6H), 4.32-4.13 (m, 2H), 4.10-3.90 (m, 2H), 3.74 (s, 3H), 2.15 (s, 3H), 1.62-1.34 (m, 4H), 1.08-0.98 (m, 4H), 0.74-0.65 (m, 6H). **LCMS:** (Method E) 541.2 (M⁺+H), Rt. 2.86 min, 93.67% (max).

Intermediate 15


3-((3-butyl-3-ethyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid



To a stirred solution of 3-butyl-3-ethyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 14; 1 g, 1.85 mmol) in THF (3 mL) at 0 °C, potassium *tert*-butoxide (208 mg, 1.85 mmol) was added and the reaction mixture was stirred for 15 minutes. A solution of β-propiolactone (148 mg, 2.03 mmol) in THF (2 mL) was then added dropwise and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL) and then diluted with water (5 mL). The aqueous layer was extracted with EtOAc (2 x 20 mL), and the combined organic layer was washed with water (20 mL) and brine (20 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 35% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 48% (550 mg, white solid). **LCMS:** (Method A) 613.3 (M⁺+H), Rt. 3.04 min, 91.99% (Max).

Intermediate 16

Methyl 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate




- 5 To a stirred solution of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (500 mg, 0.87 mmol) in DMF (10 mL), Cs_2CO_3 (0.57 g, 1.76 mmol) and methyl-2,3-epoxypropanoate (0.18 g, 1.76 mmol) were added and the reaction mixture was heated for 12 hours at 50 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (10 mL) and the aqueous layer was extracted with EtOAc
- 10 (2 x 15 mL). The combined organic layer was washed with water (15 mL) and brine (15 mL) and dried over anhydrous Na_2SO_4 . The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 33% (200 mg, colourless gum).
- LCMS:** (Method A) 671.2 ($\text{M}^+\text{+H}$), Rt. 3.32 min, 44.28% (Max).

15

Intermediate 17

3-((3,3-Dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid




- 20 To a solution of methyl 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 16; 200

mg, 0.31 mmol) in 1,4-dioxane (3 mL), dilute HCl (6 N, 3 mL) was added and the reaction mixture was heated for 16 hours at 80 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with ice-cold water (5 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was forwarded to the next step as such without any further purification. **Yield:** 200 mg (crude, colourless gum).

LCMS: (Method A) 657.2 (M⁺+H), Rt. 3.0 min, 36.70% (Max).

10 Intermediate 18

Methyl 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate




To a stirred solution of 3,3-dibutyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (0.63 g, 0.002 mmol) in DMF (9 mL) was added Cs₂CO₃ (0.92 g, 0.003 mmol) and methyl-2,3-epoxypropanoate (1.28 g, 0.0126 mmol; added portionwise in 3 equal amounts in 72 hours), and the reaction mixture was stirred for 72 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 20 mL) and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic layer was washed with water (20 mL) and brine (20 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 28% (220 mg, white solid).

LCMS: (Method E) 550.8 (M⁺+H), Rt. 3.22 min, 92.74% (Max). **HPLC:** (Method B) Rt. 6.15 min, 94.48% (Max).

Intermediate 19

Methyl (S)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate and methyl (R)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate



The two enantiomers of racemic methyl 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 18; 216 mg, 0.39 mmol) were separated by chiral SFC (method A). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

Each of the two fractions was then individually treated for further purification. The obtained residue was acidified with dilute HCl (1.5 N, pH~4) and the aqueous layer extracted with EtOAc (3 X 5 mL).


The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum at 40 °C to afford a purified enantiomer of the title compound.

Enantiomer 1: **Yield:** 46% (100 mg, colourless gum). **LCMS:** (Method E) 551.2 (M⁺+H), Rt. 2.77 min, 98.09% (Max). **Chiral SFC:** (Method A) Rt. 3.58 min, 99.10% (Max).

Enantiomer 2: **Yield:** 41% (90 mg, colourless gum). **LCMS:** (Method E) 551.1 (M⁺+H), Rt. 2.77 min, 91.29% (Max). **Chiral SFC:** (Method A) Rt. 4.59 min, 99.24% (Max)

Intermediate 20

3-((3,3-Dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid




To a stirred solution of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (300 mg, 0.52 mmol) in THF (10 mL), potassium *tert*-

butoxide (65 mg, 0.58 mmol) and β -butyrolactone (68 mg, 0.79 mmol) were added and the reaction mixture was heated for 16 hours at 60 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na_2SO_4 . The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 40% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 72% (250 mg, colourless gum). **LCMS:** (Method E) 655.3 ($\text{M}^+\text{+H}$), Rt. 3.03 min, 89.11% (Max).

10 Intermediate 21

Ethyl 2-aminohexanoate hydrochloride




To a stirred solution of DL-Norleucine (100 g, 0.76 mol) in ethanol (1L), thionyl chloride (60.8 mL, 0.84 mol) was added at 0 °C and the reaction mixture was heated for 16 hours at 80 °C. After completion of the reaction, the reaction mixture was concentrated under vacuum to afford the crude title compound which was used as such for next step without any further purification. **Yield:** 97% (145 g, brown gummy solid).

^1H NMR (400 MHz, CDCl_3): δ 8.80 (s, 3H), 4.33-4.23 (m, 2H), 4.09-4.07 (m, 1H), 2.09-2.04 (m, 2H), 1.61-1.56 (m, 1H), 1.48-1.33 (m, 6H), 1.03-0.88 (m, 3H).

20

Intermediate 22

Ethyl (*E*)-2-(benzylideneamino)hexanoate




To a stirred solution of ethyl 2-aminohexanoate hydrochloride (Intermediate 21; 145 g, 0.74 mol) in DCM (1.5 L), triethylamine (124 mL, 1.2 mol) was added at 0 °C over a period of 30 minutes. Magnesium sulfate (89.2 g, 0.74 mol) was then added portionwise to the reaction mixture at 0 °C. Benzaldehyde (75.6 mL, 0.74 mol) was then added to the reaction mixture at 0 °C over a period of 20 minutes and the reaction mixture was then stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was filtered through celite and

the filtrate was concentrated under vacuum. The resulting crude was dissolved in petroleum ether (1000 mL), again filtered through celite and the filtrate was concentrated under vacuum to afford the title compound. This crude material was forwarded as such to the next step without any further purification. **Yield:** 98% (180 g, brown liquid).

- 5 ¹H NMR (400 MHz, CDCl₃): δ 8.30 (s, 1H), 7.82-7.80 (m, 2H), 7.47-7.42 (m, 3H), 4.26-4.20 (m, 2H), 3.99-3.95 (m, 1H), 2.06-2.00 (m, 1H), 1.96-1.89 (m, 1H), 1.40-1.24 (m, 7H), 0.95-0.91 (m, 3H).

Intermediate 23

Ethyl (*E*)-2-(benzylideneamino)-2-butylhexanoate



10


To a stirred solution of NaH (60%, 29.1 g, 0.73 mol) in DMF (250 mL) at 0 °C was slowly added a solution of ethyl (*E*)-2-(benzylideneamino) hexanoate (Intermediate 22; 180 g, 0.73 mol) in DMF (250 mL) over a period of 30 minutes. The reaction mixture was stirred 1.5 hours at room temperature. A solution of *n*-butyl iodide (82.7 mL, 0.73 mol) in DMF (250 mL) was then added at 0 °C and the

15 reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (1000 mL) and the aqueous layer was extracted with petroleum ether (1000 mL). The organic layer was washed with brine (200 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude material was forwarded as such to the next step without any further purification.

- 20 **Yield:** 95% (210 g, brown liquid).

Intermediate 24

Ethyl 2-amino-2-butylhexanoate



- 25 To a stirred solution of ethyl (*E*)-2-(benzylideneamino)-2-butylhexanoate (Intermediate 23; 210 g, 0.76 mol) in petroleum ether (1000 mL), dilute HCl (1000 mL, 1.5 N) was added at 0 °C and the reaction mixture was stirred vigorously for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the organic layer was separated and washed with EtOAc (2 x 100 mL).


The aqueous layer was then basified (pH ~10) using sodium hydroxide solution and extracted with DCM (2 x 1000 mL). The combined organic part was washed with water (2 x 1000 mL), brine (1000 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum to afford the title compound which was forwarded as such to the next step without any further purification.

5 **Yield:** 52% (85 g, brown liquid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 4.11-4.04 (m, 2H), 1.78-1.61 (m, 4H), 1.60-1.51 (m, 3H), 1.50-1.22 (m, 7H), 1.20-0.99 (m, 3H), 0.95-0.75 (m, 6H).


Intermediate 25

10 **2-Amino-2-butyl-N-phenylhexanamide**




To a stirred solution of aniline (19.1 mL, 209 mmol) in THF (250 mL) at -78 °C, *n*-BuLi (2.6 M in hexane, 250.7 mL, 627 mmol) was added dropwise over a period of 30 minutes and the reaction mixture was stirred for 45 minutes at -25 °C to -30 °C. A solution of ethyl 2-amino-2-butylhexanoate (Intermediate 24; 45 g, 209 mmol) in THF (200 mL) was then added to the reaction mixture at -78 °C and the reaction mixture was stirred for 2 hours at that temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched at -78 °C with isopropanol (100 mL) and then ice-cold water (500 mL), and the reaction mixture was allowed to stir for 1 hour at room temperature. The aqueous part was extracted with EtOAc (2 x 250 mL) and the combined organic layer was washed with water (2 x 50 mL). The organic part was dried over anhydrous Na₂SO₄ and concentrated under vacuum to afford the title compound as crude. The obtained crude material was dissolved in petroleum ether (1000 mL), washed with 30% methanol in water (2 x 250 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude was forwarded as such to the next step without any further purification. **Yield:** 60 g (crude, brown liquid).

LCMS: (Method A) 263.3 (M⁺), Rt. 1.29 min, 95.84% (max).


Intermediate 26**2-Butyl-N1-phenylhexane-1,2-diamine**

To a stirred solution of 2-amino-2-butyl-N-phenylhexanamide (Intermediate 25; 100 g, 0.38 mol) in THF (1L), borane dimethyl sulphide complex (2 M in THF, 476 mL, 0.95 mol) was added at 0 °C and the reaction mixture was heated for 16 hours at 70 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with methanol (300 mL) at 0 °C and the reaction mixture was heated for 2 hours at 70 °C. The reaction mixture was then concentrated under vacuum. The obtained residue was dissolved in EtOAc (1000 mL), and the organic layer was washed with water (2 x 300 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 0-25% EtOAc in hexane; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 39% (37 g, brown liquid).

Intermediate 27**2,4-Dichloro-5-methoxybenzenesulfonic acid**


To a stirred solution of chlorosulphonic acid (60 mL) at 0 °C, 2,4-dichloroanisole (20 g, 113 mmol) was added portionwise and the reaction mixture was stirred for 3 hours between 0 and 10 °C. After completion of the reaction (monitored by TLC), the reaction mixture was poured onto crushed ice with vigorous stirring and the obtained solid was filtered off. The solid was washed several times with ice-cold water (50 mL), petroleum ether (50 mL) and then dried under vacuum to afford the title compound. **Yield:** 70% (19 g, off-white solid).

LCMS: (Method A) 255.0 (M⁺-H), Rt. 2.66 min, 96.06% (Max).

Intermediate 28**2,4-Dichloro-5-methoxy-N-(5-((phenylamino)methyl)nonan-5-yl)benzenesulfonamide**

A solution of 2,4-dichloro-5-methoxybenzenesulfonic acid (Intermediate 27; 10 g, 41.5 mmol) in
 5 SOCl_2 (20 mL) was heated for 12 hours at 70 °C. After complete consumption of starting material, the
 reaction mixture was concentrated under vacuum and the obtained residue was dissolved in THF (50
 mL). A solution of triethylamine (17 mL, 124.7 mmol) in THF (100 mL) and then 2-butyl-N1-phenyl-
 hexane-1,2-diamine (Intermediate 26; 12.4 g, 49.9 mmol) were added at 0 °C, and the reaction
 mixture was stirred for 4 hours at room temperature. After completion of the reaction (monitored by
 10 TLC), the reaction mixture was quenched with ice-cold water (30 mL) and the aqueous layer was
 extracted with EtOAc (2 x 100 mL). The combined organic layer was washed with water (100 mL) and
 brine (100 mL) and dried over anhydrous Na_2SO_4 . The organic part was filtered and concentrated
 under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent:
 25-35% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 40% (8 g, off-white
 15 solid).

LCMS: (Method E) 487.1 ($\text{M}^+\text{+H}$), Rt. 2.93 min, 43.66% (Max).

Intermediate 29**3,3-Dibutyl-7-chloro-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

To a stirred solution of 2,4-dichloro-5-methoxy-N-(5-((phenylamino)methyl)nonan-5-yl)benzene-
 sulfonamide (Intermediate 28; 4 g, 8.21 mmol) in DMF (20 mL), anhydrous K_2CO_3 (2.26 g, 16.42
 mmol) and copper powder (0.52 g, 8.21 mmol) were added and the reaction mixture was heated for
 16 hours at 120 °C. After completion of the reaction (monitored by TLC), the reaction mixture was
 25 filtered through celite and the celite pad was washed with DCM (10 mL). The filtrate was
 concentrated under vacuum and the resulting crude material was purified by Isolera column


chromatography (eluent: 2-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 24% (900 mg, off-white solid).

LCMS: (Method E) 451.1 ($M^+ + H$), Rt. 2.96 min, 34.15% (Max).

5 Intermediate 30

3,3-Dibutyl-7-chloro-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide




To a solution of 3,3-dibutyl-7-chloro-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothia-
diazepine 1,1-dioxide (Intermediate 29; 0.9 g, 1.2 mmol) in DCM (4 mL) at -78 °C, BBr₃ (1 M in DCM; 4
10 mL, 4 mmol) was added and the reaction mixture was stirred for 3 h between -10 °C and 0 °C. After
completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold
water (10 mL). The organic layer was washed with water (10 mL) and brine (10 mL). The organic part
was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude
material was purified by Isolera column chromatography (eluent: 0-20% EtOAc/PE; silica gel: 230-400
15 mesh) to afford the title compound. **Yield:** 22% (195 mg, yellow gum).

LCMS: (Method E) 437.1 ($M^+ + H$), Rt. 2.80 min, 53.01% (Max).

Intermediate 31

5-(4-Bromophenyl)-3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-2,3,4,5-tetrahydro- 20 1,2,5-benzothiadiazepine 1,1-dioxide




To a stirred solution of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-
tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (3 g, 5.27 mmol) in DMF (18 mL) at 0 °C was added

a solution of N-bromosuccinimide (1.03 g, 5.80 mmol) in DMF (12 mL), and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was poured into ice-cold water (20 mL), stirred and filtered. The obtained solid was purified by Isolera column chromatography (eluent: 25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 82% (2.8 g, white solid).

LCMS: (Method B) 648.2 ($M^+ + 2$), Rt. 4.65 min, 67.27% (Max).

Intermediate 32

3,3-Dibutyl-8-hydroxy-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide




To a solution of 5-(4-bromophenyl)-3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 31; 2.0 g, 3.09 mmol) in DMF (20 mL) at 0 °C, cuprous bromide (44 mg, 0.31 mmol) was added, followed by a freshly prepared solution of sodium methoxide (prepared *in situ* by adding sodium (0.35 g, 15.4 mmol) to dry methanol (10 mL)). The reaction mixture was then heated for 24 hours at 100 °C. After completion of the reaction (monitored by UPLC), the reaction mixture was quenched with dilute HCl (1.5 N, 10 mL) and the aqueous layer was extracted with 1:1 mixture of EtOAc and PE (2 x 30 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 81% (1.5 g, pale yellow solid).

LCMS: (Method E) 599.3 ($M^+ + H$), Rt. 3.38 min, 99.50% (Max).

Intermediate 33

3-((3,3-Dibutyl-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid




- 5 To a solution of 3,3-dibutyl-8-hydroxy-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 32; 1.5 g, 2.64 mmol) in THF (20 mL), potassium *tert*-butoxide (326 mg, 2.9 mmol) was added and the reaction mixture was stirred for 10 minutes at 0 °C. Then β -propiolactone (209 mg, 2.9 mmol) was added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC),
 10 the reaction mixture was quenched with dilute HCl (1.5 N, 20 mL) and the aqueous layer was extracted with EtOAc (2 x 30 mL). The combined organic layer was washed with water (30 mL) and brine (30 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 0-70% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 48% (850 mg, white
 15 solid).

LCMS: (Method E) 670.8 (M⁺+H), Rt. 3.37 min, 94.99% (Max).

Intermediate 34

3-((3,3-Dibutyl-5-(4-methoxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid




To a solution of 3-((3,3-dibutyl-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (Intermediate 33; 850 mg, 1.27 mmol) in toluene (10 mL), triphenylamine (621 mg, 2.54 mmol) was added and the reaction mixture was stirred for 10 minutes at 0 °C. TFA (1.9 mL, 25.4 mmol) was then added and the reaction mixture was allowed to stir for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 0-5% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 92% (650 mg, off-white solid).

LCMS: (Method E) 550.8 ($M^+ + H$), Rt. 3.24 min, 94.98% (Max).

Intermediate 35


Isopropyl 3-((3,3-dibutyl-5-(4-hydroxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoate




To a solution of 3-((3,3-dibutyl-5-(4-methoxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (Intermediate 34; 300 mg, 0.54 mmol) in DCM (10 mL) at -78 °C, BBr_3 (1.1 mL, 1.09 mmol) was added and the reaction mixture was stirred for 2 hours at -30 °C. After completion of the reaction (monitored by UPLC), the reaction mixture was quenched with isopropyl alcohol and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 0-5% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 150 mg (off-white solid).

LCMS: (Method A) 578.8 ($M^+ + H$), Rt. 2.88 min, 83.38% (Max).

(Note: during quenching with isopropyl alcohol, the esterification product of the desired compound observed which was indicated by LCMS)

Intermediate 36**3-Butyl-7-(dimethylamino)-3-ethyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

- 5 To a degassed solution of 7-bromo-3-butyl-3-ethyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 13; 3.5 g, 6.34 mmol) in 1,4-dioxane (35 mL) were added sodium *tert*-butoxide (1.2 g, 12.6 mmol), xantphos (0.073g, 0.13 mmol) and Pd₂dba₃ (0.058 g, 0.06 mmol) and the solution was degassed for 10 minutes under N₂ atmosphere. *N,N*-dimethylamine (3.5 mL, 31.7 mmol) was added and the reaction mixture was heated for 48
- 10 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated and the obtained residue was diluted with EtOAc (75 mL). The organic layer was washed with water (2 x 75 mL) and brine (75 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 30% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 40% (1.4 g, brown gum).
- 15 ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.34-7.28 (m, 5H), 7.11-7.02 (m, 3H), 6.84 (d, *J* = 8.8 Hz, 2H), 6.28 (s, 1H), 4.50 (bs, 2H), 4.12 (bs, 2H), 3.92 (s, 3H), 3.81 (s, 3H), 2.68 (s, 6H), 1.42-1.25 (m, 2H), 1.19-1.05 (m, 2H), 0.95-0.81 (m, 4H), 0.73-0.58 (m, 6H). **LCMS:** (Method E) 552.1 (M⁺+H), Rt. 2.80 min, 81.8% (max).


Intermediate 37**3-Butyl-7-(dimethylamino)-3-ethyl-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

To a stirred solution of 3-butyl-7-(dimethylamino)-3-ethyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 36; 0.7 g, 1.22 mmol) in DMF (10 mL), sodium methoxide (0.5 g, 6.32 mmol) was added and the reaction mixture was heated for 16 hours at 100 °C. The completion of the reaction (monitored by TLC and LCMS), the reaction mixture was concentrated under vacuum and the obtained residue was diluted with EtOAc (20 mL). The organic layer was washed with water (2 x 20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 70% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 60% (0.4 g, off-white solid).

LCMS: (Method E) 418.2 (M⁺+H), Rt. 2.14 min, 88.9% (max).

Intermediate 38

Methyl tritylserinate




To a stirred solution of methyl serinate hydrochloride (0.65 g, 0.5 mmol) in DCM (10 mL), triethylamine (2 mL, 1.50 mmol) was added and the reaction mixture was cooled to 0 °C. Then trityl chloride (1.67 g, 0.6 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (10 mL) and the aqueous layer was extracted with DCM (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 7% EtOAc /PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 33% (0.7 g, white solid).

¹H NMR (400 MHz, CDCl₃): δ 7.54-7.50 (m, 6H), 7.32-7.27 (m, 6H), 7.24-7.20 (m, 3H), 3.74-3.72 (m, 1H), 3.62-3.51 (m, 2H), 3.33 (s, 3H), 3.00 (bs, 1H), 2.33 (bs, 1H).

Intermediate 39

Methyl *O*-(3-butyl-3-ethyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)-*N*-tritylserinate.




- 5 To a stirred solution of 3-butyl-3-ethyl-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 14; 400 mg, 0.74 mmol) in toluene (10 mL) at 0 °C, triphenylphosphine (388 mg, 1.48 mmol) and methyl tritylserinate (Intermediate 38; 333 mg, 0.88 mmol) were added. The reaction mixture was stirred for 5 minutes, then DIAD (225 mg, 1.11 mmol) was added dropwise at 0 °C and the reaction mixture was heated for
- 10 3 hours at 120 °C. After the completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum. The obtained residue was diluted with water (10 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column
- 15 chromatography (eluent: 10-25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.
- Yield:** 99% (660 mg, off-white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ. 7.45 - 7.44 (m, 7H), 7.32 - 7.26 (m, 11H), 7.20 - 7.17 (m, 3H), 7.06 (s, 1H), 6.91 - 6.89 (m, 4H), 4.25 (s, 2H), 3.75 (s, 3H), 3.74 (d, *J* = 0.4 Hz, 2H), 3.19 (t, *J* = 0.4 Hz, 4H), 3.15 (s, 2H), 2.11 (s, 3H), 1.24 - 1.23 (m, 3H), 1.21 - 1.13 (m, 2H), 0.99 (s, 3H), 0.66 (s, 6H).

20

Intermediate 40

Methyl *O*-(3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serinate




To a stirred solution of methyl O-(3-butyl-3-ethyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)-N-tritylserinate (Intermediate 39; 300 mg, 0.44 mmol) in toluene (5 mL) at 0 °C, triphenylamine (166 mg, 0.67 mmol) and TFA (774 mg, 6.78 mmol) were added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture poured into ice-cold water (10 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL). The organic part was then dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 25% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 79% (140 mg, green solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.36-7.29 (m, 4H), 7.20-7.14 (m, 2H), 7.02-6.98 (m, 1H), 6.52 (s, 1H), 4.42 (s, 1H), 4.33 (s, 2H), 3.78-3.72 (m, 5H), 2.08 (s, 3H), 1.40-1.35 (m, 2H), 1.28-1.24 (m, 2H), 1.20-1.15 (m, 3H), 1.13-1.08 (m, 3H), 1.04-0.97 (m, 6H). **LCMS:** (Method E) 522.3 (M⁺+H), Rt. 2.51 min, 92.42% (Max).

Example 1

3-((3,3-Dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid




To a stirred solution of 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (Intermediate 2; 15 g, 23.41 mmol) in dry DCM (150 mL) at 0 °C, TFA (45 mL) and triethylsilane (45 mL) were added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by LCMS), the reaction mixture was diluted with ice water (25 mL). The aqueous layer was extracted with EtOAc (2 x 100 mL). The combined organic layer was dried over anhydrous Na₂SO₄ and evaporated under vacuum. The resulting crude material was purified by Prep-HPLC (Method D) to afford title compound. **Yield:** 47% (5.8 g, off-white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.44 (bs, 1H), 7.29 - 7.19 (m, 4H), 7.18 - 7.02 (m, 2H), 6.98 - 6.89 (m, 1H), 6.58 - 6.54 (m, 1H), 4.22 (t, *J* = 5.6 Hz, 2H), 3.95 - 3.72 (m, 2H), 2.70 (t, *J* = 6.0 Hz, 2H), 2.08 (s,

3H), 1.53 - 1.24 (m, 6H), 1.08 - 1.01 (m, 6H), 0.77 - 0.73 (m, 6H). **LCMS:** (Method D) 521.3 ($M^+ + H$), Rt. 2.92 min, 99.24% (Max). **HPLC:** (Method B) Rt. 6.22 min, 98.91% (Max).

Example 2

5 3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid




To a stirred solution of 3-((3-butyl-3-ethyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (Intermediate 15; 550 mg, 0.9 mmol) in DCM (6 mL), TFA (2 mL) and triethylsilane (2 mL) were added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-60% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 67% (300 mg, white solid).

15 **^1H NMR** (400 MHz, CD_3OD): δ 7.34-7.30 (m, 3H), 7.17 (d, $J = 7.6$ Hz, 2H), 7.02 (t, $J = 7.6$ Hz, 1H), 6.62 (s, 1H), 4.30 (t, $J = 6.4$ Hz, 2H), 3.89 (s, 2H), 2.81 (t, $J = 6.4$ Hz, 2H), 2.10 (s, 3H), 1.85-1.58 (m, 2H), 1.56-1.41 (m, 2H), 1.40-1.25 (m, 1H), 1.23-0.95 (m, 3H), 1.07-0.79 (m, 6H). **LCMS:** (Method A) 493.2 ($M^+ + H$), Rt. 2.65 min, 94.21% (Max). **HPLC:** (Method B) Rt. 5.62 min, 94.00% (Max).

20 Examples 3 and 4

(S)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid and (R)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid



25 The two enantiomers of racemic 3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (300 mg, 0.61 mmol) were separated by

chiral SFC (method F). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.


Each of the two fractions was then individually treated for further purification. The obtained residue was acidified with dilute HCl (1.5 N, pH~4) and the aqueous layer was extracted with EtOAc (3 x 15 mL). The combined organic layer was washed with water (15 mL) and brine (15 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum at 40 °C to afford a purified enantiomer of the title compound.

Enantiomer 1: **Yield:** 25% (75 mg, pale brown solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.42 (s, 1H), 7.28-7.20 (m, 4H), 7.09-7.07 (m, 1H), 6.93 (t, *J* = 6.8 Hz, 1H), 6.59 (s, 1H), 4.23 (t, *J* = 5.6 Hz, 2H), 3.78 (bs, 2H), 2.77-2.69 (m, 2H), 2.09 (s, 3H), 1.73-1.59 (m, 1H), 1.56-1.47 (m, 1H), 1.41-1.32 (m, 2H), 1.12-1.28 (m, 2H), 1.10-0.97 (m, 3H), 0.76 (t, *J* = 7.2 Hz, 6H). **LCMS:** (Method E) 493.2 (M⁺+H), Rt. 2.54 min, 95.95% (Max). **HPLC:** (Method B) Rt. 5.62 min, 95.16% (Max). **Chiral SFC:** (Method H) Rt. 5.11 min, 98.47% (Max).

Enantiomer 2: **Yield:** 33% (100 mg, pale brown solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.39 (s, 1H), 7.27-7.19 (m, 4H), 7.06 (d, *J* = 6.0 Hz, 2H), 6.92 (t, *J* = 6.8 Hz, 1H), 6.58 (s, 1H), 4.22 (t, *J* = 5.6 Hz, 2H), 3.78 (bs, 2H), 2.70 (t, *J* = 6.0 Hz, 2H), 2.08 (s, 3H), 1.71-1.57 (m, 1H), 1.56-1.45 (m, 1H), 1.44-1.31 (m, 2H), 1.29-1.17 (m, 1H), 1.14-0.88 (m, 3H), 0.75-0.71 (m, 6H). **LCMS:** (Method E) 493.2 (M⁺+H), Rt. 2.54 min, 93.60% (Max). **HPLC:** (Method B) Rt. 5.62 min, 92.78% (Max). **Chiral SFC:** (Method H) Rt. 5.91 min, 97.35% (Max).

Example 5

3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid




To a stirred solution of 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid (Intermediate 17; 200 mg, 0.3 mmol) in DCM (10 mL), TFA (0.6 mL, 3 vol) and triethylsilane (0.6 mL, 3 vol) were added and

the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (5 mL) and the aqueous layer was extracted with DCM (2 x 5 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by prep HPLC (method A) to afford the title compound. **Yield:** 11% (18 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.29-7.21 (m, 4H), 7.19-7.01 (m, 2H), 6.93 (t, *J* = 7.2 Hz, 1H), 6.58 (s, 1H), 4.35-4.17 (m, 1H), 4.03-3.91 (m, 2H), 3.51-3.35 (m, 2H), 2.10 (s, 3H), 1.56-0.97 (m, 12H), 0.75 (t, *J* = 6.4 Hz, 6H). **LCMS:** (Method A) 537.3 (M⁺+H), Rt. 2.73 min, 94.57% (Max). **HPLC:** (Method B) Rt. 5.79 min, 96.65% (Max).

Examples 6 and 7

(S)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid and (R)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid



To a stirred solution of enantiomer 1 of methyl 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 19; 90 mg, 0.16 mmol) in 1,4-dioxane (2 mL), dilute HCl (6 N, 3 mL) was added and the reaction mixture was heated for 16 h at 75 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with ice-cold water (2 mL) and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was triturated with petroleum ether to afford enantiomer 1 of the title compound.

Enantiomer 2 of the title compound was obtained following the same procedure, starting from 80 mg of enantiomer 2 of Intermediate 19. The absolute configuration of the two enantiomers is not known.


Enantiomer 1: **Yield:** 79% (70 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.74 (bs, 1H), 7.26-7.22 (m, 4H), 7.13-7.11 (m, 2H), 6.95 (t, *J* = 6.8 Hz, 1H), 6.56 (s, 1H), 4.35-4.33 (m, 1H), 4.24-4.15 (m, 2H), 3.78 (bs, 2H), 2.09 (s, 3H), 1.53-1.51 (m, 2H), 1.41-1.33 (m, 2H), 1.31-1.20 (m, 2H), 1.19-0.95 (m, 6H), 0.75 (t, *J* = 6.4 Hz, 6H). **LCMS:** (Method D) 537.2 (M⁺+H), Rt. 2.86 min, 93.83% (Max). **HPLC:**

(Method A) Rt. 5.43 min, 93.28% (Max). **Chiral SFC:** (Method H) Rt. 3.34 min, 100% (Max).

Enantiomer 2: **Yield:** 64% (50 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.30-7.22 (m, 4H), 7.13-7.11 (m, 2H), 6.97-6.95 (m, 1H), 6.56 (s, 1H), 4.27-4.22 (m, 2H), 4.16-4.12 (m, 1H), 3.83 (bs, 2H), 2.09 (s, 3H), 1.53-1.51 (m, 2H), 1.41-1.33 (m, 2H), 1.31-1.20 (m, 2H), 1.19-0.95 (m, 6H), 0.75 (t, *J* = 6.8 Hz, 6H). **LCMS:** (Method D) 537.2 (M⁺+H), Rt. 2.86 min, 96.22% (Max). **HPLC:** (Method A) Rt. 5.45 min, 95.09% (Max). **Chiral SFC:** (Method H) Rt. 1.85 min, 100% (Max).


Example 8

3-((3,3-Dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid




To a stirred solution of 3-((3,3-dibutyl-2-(4-methoxybenzyl)-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid (Intermediate 20; 250 mg, 0.38 mmol) in DCM (10 mL), TFA (0.75 mL, 3 vol) and triethylsilane (0.75 mL, 3 vol) were added and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (5 mL) and the aqueous layer was extracted with DCM (2 x 5 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by prep HPLC (Method A) to afford the title compound. **Yield:** 24% (50 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.34-7.18 (m, 4H), 7.15-7.05 (m, 2H), 6.97-6.88 (m, 1H), 6.53 (s, 1H), 4.75-4.72 (m, 1H), 3.80 (s, 2H), 2.55-2.50 (m, 2H), 2.06 (s, 3H), 1.62-1.45 (m, 2H), 1.44-1.34 (m, 2H), 1.32-1.28 (m, 5H), 1.25-0.90 (m, 6H), 0.75-0.72 (m, 6H). **LCMS:** (Method A) 535.3 (M⁺+H), Rt. 2.95 min, 98.47% (Max). **HPLC:** (Method B) Rt. 6.29 min, 96.34% (Max).

Example 9**3-((3,3-dibutyl-7-chloro-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid**

- 5 To a stirred solution of 3,3-dibutyl-7-chloro-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 30; 200 mg, 0.45 mmol) in THF (3 mL) at 0 °C, potassium *tert*-butoxide (56 mg, 0.5 mmol) was added and the reaction mixture was stirred for 15 minutes. A solution of β -propiolactone (32 mg, 0.45 mmol) in THF (1 mL) was then added dropwise and the reaction mixture was stirred for 6 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 2 mL) and then diluted with water (2 mL). The aqueous layer was extracted with EtOAc (2 x 5 mL) and the combined organic layer was washed with water (5 mL) and brine (5 mL). The organic part was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 3% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 20% (45 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.30 (bs, 1H), 7.62-7.43 (m, 1H), 7.40-7.27 (m, 3H), 7.26-7.12 (m, 2H), 7.09-6.98 (m, 1H), 6.80 (s, 1H), 4.25 (t, *J* = 5.6 Hz, 2H), 3.87 (bs, 2H), 2.72 (t, *J* = 6.0 Hz, 2H), 1.56-1.45 (m, 2H), 1.44-1.31 (m, 2H), 1.29-1.19 (m, 2H), 1.15-0.91 (m, 6H), 0.89-0.72 (m, 6H). **LCMS:** (Method E) 509.1 (M⁺+H), Rt. 2.78 min, 96.38% (Max). **HPLC:** (Method B) Rt. 6.18 min, 96.38% (Max).


Example 10**3-((3,3-dibutyl-5-(4-hydroxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid**

To a solution of isopropyl 3-((3,3-dibutyl-5-(4-hydroxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoate (Intermediate 35; 120 mg, 0.20 mmol) in 1,4-dioxane (2 mL), dilute HCl (6 N, 4 mL) was added and the reaction mixture was heated for 12 hours at 70 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with ice-cold water (5 mL) and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 40-50% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 54% (60 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.37 (s, 1H), 9.30 (s, 1H), 7.39 (s, 1H), 7.09-7.06 (m, 3H), 6.76 (d, *J* = 8.8 Hz, 2H), 6.26 (s, 1H), 4.16 (t, *J* = 6.0 Hz, 2H), 3.87 (s, 2H), 2.67 (t, *J* = 6.0 Hz, 2H), 2.00 (s, 3H), 1.59-1.35 (m, 4H), 1.33-1.07 (m, 4H), 1.06-0.81 (m, 4H), 0.74 (t, *J* = 6.8 Hz, 6H). **LCMS:** (Method E) 537.2 (M⁺+H), Rt. 2.37 min, 98.62% (Max). **HPLC:** (Method B) Rt. 5.22 min, 98.80% (Max).

Example 11

3-((3-Butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid



To a stirred solution of 3-butyl-7-(dimethylamino)-3-ethyl-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 37; 600 mg, 1.43 mmol) in THF (3 mL) at 0 °C, potassium *tert*-butoxide (177 mg, 1.58 mmol) was added and the reaction mixture was stirred for 15 minutes. A solution of β-propiolactone (103 mg, 1.43 mmol) in THF (2 mL) was then added dropwise and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL) and diluted with water (5 mL). The aqueous layer was extracted with EtOAc (2 x 20 mL), and the combined organic layer was washed with water (20 mL) and brine (20 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 3-5% MeOH/DCM; silica gel: 230-400 mesh) and the obtained material was re-purified by prep HPLC (Method A) to afford the title compound.


Yield: 6% (42 mg, off-white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.41 (s, 1H), 7.25 (t, *J* = 8.0 Hz, 2H), 7.14-6.97 (m, 4H), 6.90 (t, *J* = 6.8 Hz, 1H), 6.23 (s, 1H), 4.16 (t, *J* = 5.6 Hz, 2H), 3.79 (s, 2H), 2.79-2.72 (m, 2H), 2.52 (s, 6H), 1.71-1.47 (m, 2H), 1.46-1.33 (m, 2H), 1.32-1.17 (m, 2H), 1.11-0.97 (m, 2H), 0.76-0.67 (m, 6H). **LCMS:** (Method E) 490.2 (M⁺+H), Rt. 2.41 min, 98.92% (Max). **HPLC:** (Method B) Rt. 4.28 min, 99.05% (Max).

5

Examples 12 and 13

(S)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid and (R)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid



10

The two enantiomers of racemic 3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid (Example 11; 65 mg, 0.13 mmol) were separated by chiral SFC (method I). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

15

Each of the two fractions was then individually treated for further purification. The obtained residue was acidified with dilute HCl (1.5 N, pH~4) and the aqueous layer extracted with EtOAc (3 x 5 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum at 40 °C to afford a purified enantiomer of the title compound.

20

Enantiomer 1: **Yield:** 6% (5 mg, off-white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.40 (s, 1H), 7.25 (t, *J* = 7.6 Hz, 2H), 7.15 (s, 1H), 7.08-7.02 (m, 3H), 6.91 (t, *J* = 7.2 Hz, 1H), 6.23 (s, 1H), 4.16 (t, *J* = 5.6 Hz, 2H), 3.80 (bs, 2H), 2.79-2.71 (m, 2H), 2.58 (s, 6H), 1.71-1.59 (m, 1H), 1.58-1.45 (m, 1H), 1.44-1.32 (m, 2H), 1.29-1.15 (m, 1H), 1.14-0.91 (m, 3H), 0.79-0.69 (m, 6H). **LCMS:** (Method E) 490.1 (M⁺+H), Rt. 2.39 min, 98.52% (Max). **HPLC:** (Method B) Rt. 4.32 min, 98.57% (Max). **Chiral SFC:** (Method M) Rt. 3.09 min, 98.77% (Max).


25

Enantiomer 2: **AS0649:Yield:** 5% (7 mg, off-white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.40 (s, 1H), 7.25 (t, *J* = 7.6 Hz, 2H), 7.15 (s, 1H), 7.08-7.03 (m, 3H), 6.91 (t, *J* = 7.2 Hz, 1H), 6.24 (s, 1H), 4.16 (t,

30

$J = 5.6$ Hz, 2H), 3.80 (bs, 2H), 2.79-2.71 (m, 2H), 2.58 (s, 6H), 1.71-1.59 (m, 1H), 1.58-1.45 (m, 1H), 1.44-1.31 (m, 2H), 1.29-1.14 (m, 1H), 1.14-0.91 (m, 3H), 0.79-0.69 (m, 6H). **LCMS:** (Method E) 489.9 ($M^+ + H$), Rt. 2.38 min, 99.66% (Max). **HPLC:** (Method B) Rt. 4.28 min, 98.43% (Max). **Chiral SFC:** (Method M) Rt. 4.25 min, 97.61% (Max).

5

Example 14***O*-(3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine**

- 10 To a stirred solution of methyl *O*-(3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serinate (Intermediate 40; 140 mg, 0.27 mmol) in 1,4-dioxane (2 mL), lithium hydroxide (23 mg, 0.53 mmol) was added and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 2 mL) and the aqueous layer was extracted
- 15 with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na_2SO_4 . The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 28% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 31% (42 mg, off-white solid).
- ^1H NMR** (400 MHz, $DMSO-d_6$): δ 7.85 (s, 2H), 7.29 (t, $J = 6.4$ Hz, 3H), 7.11 (d, $J = 6.0$ Hz, 2H), 6.97 (t, $J = 6.8$ Hz, 1H), 6.56 (s, 1H), 4.37 - 4.34 (m, 1H), 4.20 (t, $J = 8.0$ Hz, 1H), 3.89 (s, 1H), 3.63 (s, 1H), 3.10 - 3.05 (m, 1H), 2.10 (s, 3H), 1.63 - 1.63 (m, 8H), 1.05 - 0.72 (m, 6H). **LCMS:** (Method E) 508.3 ($M^+ + H$), Rt. 2.39 min, 98.34% (Max). **HPLC:** (Method B) Rt. 4.58 min, 97.26s% (Max).
- 20

BIOLOGICAL ASSAYS

25

IBAT (h/m) assay protocol

- 10,000 cells (Human or Mouse IBAT-overexpressing cells) were seeded in 96-wells plate (Corning CLS3809) in 200 μ L MEM-alpha medium (Gibco 12571-063) supplemented with 10% FBS (Gibco 10438026) containing Puromycin (Gibco A1113803) (10 μ g/mL) and incubated at 37 $^{\circ}$ C in 5% CO_2 for
- 30

48 hours. After incubation, media was decanted from the wells and cells were washed two times with 300 µL of basal MEM-alpha medium (FBS-free). After decanting basal MEM-alpha medium each time, plates were tapped against paper towel to ensure maximum removal of residual media.

Test inhibitor dilutions (highest test concentration being 10 µM, 3-fold serial dilution, 10 points)

5 prepared in DMSO (Sigma D2650) were added in incubation mix (maintaining 0.2% final DMSO concentration) containing 0.25 µM 3H-taurocholic acid (ARC ART-1368) and 5 µM of cold taurocholic acid (Sigma T4009). 50 µL of incubation mix containing test inhibitors was then added to the wells (in duplicate) and the plates were incubated for 20 minutes in a CO₂ incubator at 37 °C. After incubation, the reaction was stopped by keeping the plates on ice water mix for 2-3 minutes and then the
10 incubation mix was aspirated completely from the wells. The wells were washed two times with 250 µL of chilled unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10 mM) HBSS (Gibco 14175079) (pH 7.4). The plates were tapped against a paper towel after every wash to ensure maximum removal of blocking buffer.

100 µL of MicroScint-20 (PerkinElmer 6013621) was added to the wells and kept overnight at room
15 temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well).

LBAT (h/m) assay protocol

20 20,000 cells (Human or Mouse LBAT-overexpressing cells) were seeded in 96-wells plate (Corning CLS3809) in 100 µL MEM-alpha medium (Gibco 12571-063) supplemented with 10% FBS (Gibco 10438026) containing Geneticin (Gibco 10131-027) (1 mg/mL) and incubated at 37 °C in 5% CO₂ for 24 hours. After incubation, media was decanted from the wells and cells were washed two times with 300 µL of basal MEM-alpha medium (FBS-free). After decanting basal MEM-alpha medium each
25 time, plates were tapped against paper towel to ensure maximum removal of residual media.

For human LBAT, incubation mix was prepared by adding test inhibitor dilutions (3-fold serial dilution in DMSO (Sigma D2650), 10 points) in MEM-alpha (without FBS) containing 0.3 µM 3H-taurocholic acid (ARC ART-1368) and 7.5 µM cold taurocholic acid (Sigma T4009) (maintaining 0.2% final DMSO concentration). For mouse LBAT, incubation mix was prepared by adding test inhibitor dilutions (3-
30 fold serial dilution in DMSO, 10 points) in MEM-alpha (without FBS) containing 0.3 µM 3H-taurocholic acid and 25 µM cold taurocholic acid maintaining 0.2% final DMSO concentration).

50 µL of incubation mix containing test inhibitors was then added to the wells (in duplicate) and the plates were incubated for 20 minutes in a CO₂ incubator at 37 °C. After incubation, the reaction was stopped by keeping the plates on ice water mix for 2-3 minutes and then the incubation mix was
35 aspirated completely from the wells. The wells were washed two times with 250 µL of chilled

unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10 mM) HBSS (Gibco 14175079) (pH 7.4). The plates were tapped against a paper towel after every wash to ensure maximum removal of blocking buffer.

100 µL of MicroScint-20 (PerkinElmer 6013621) was added to the wells and kept overnight at room temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well, with normal plate orientation).

Bidirectional permeability assay (Caco-2 cells)

Caco-2 cells (Evotec) were seeded at a density of 70,000 cells/well in Millicell® 24-well cell culture insert plates and maintained in an incubator (37 °C, 5% CO₂, 95% RH) for 21 days with media change on alternate days.

Stock solutions (10 mM) of the test compounds, atenolol (low permeability marker), propranolol (high permeability marker) and digoxin (substrate for P-gp transport pathway) were prepared in dimethylsulfoxide (DMSO). An intermediate stock solution (1 mM) was prepared by diluting 10 µL of 10 mM master stock solution with 90 µL of neat DMSO. A working stock solution (10 µM) was prepared by diluting 50 µL of 1 mM with 4950 µL of FaSSIF buffer. Post addition of compounds to the FaSSIF, samples were subjected to sonication for 2 hours, and centrifuged at 4000 RPM for 30 minutes at 37 °C. The 4 mL of resultant supernatant was directly used in the assay. The final DMSO concentration in the transport experiments was 1%.

On the day of assay, Caco-2 monolayers were washed twice with transport buffer (HBSS, pH 7.4) and pre-incubated for 30 min (37 °C, 5% CO₂, 95% RH) in an incubator. The electrical resistance of the monolayers was measured with a Millicell® - ERS system. Monolayers with trans-epithelial electrical resistance (TEER) values greater than 350 ohm.cm² were selected for the assay.

The assay was conducted in absorptive direction (A2B) and secretory (B2A) directions. Transport experiments were initiated by addition of transport assay buffer (FaSSIF buffer prepared in HBSS) consisting of compounds to the donor compartment (apical chamber A-B; basolateral chamber B-A) in duplicate (n=2) wells. Drug free HBSS buffer (pH 7.4) containing 1% bovine serum albumin (BSA) was introduced to the receiver (A-B-basolateral; B-A- Apical) compartments. The volumes of apical and basolateral compartments were 0.4 and 0.8 mL, respectively. After adding dosing solution, plates were incubated in an incubator for 120 minutes at 37 °C. After 120 minutes, donor and receiver samples were collected and matrix matched (1:1, 30 µL study sample + 30 µL blank buffer) with the opposite buffer. Dosing samples matrix matched (1:1, 30 µL study sample + 30 µL blank buffer) with the opposite buffer. Samples were processed by adding acetonitrile containing internal standard (60

μL study sample + 200 μL acetonitrile containing internal standard -Tolbutamide, 500 ng/mL).

Samples were vortexed and centrifuged at 4000 rpm for 10 minutes. The obtained supernatant (100 μL) was diluted with 100 μL of water and transferred to fresh 96 well plates. The concentration of compounds in the samples was analyzed by liquid chromatography tandem mass spectrometry (LC-MS/MS) method using discovery grade bio-analytical method, as applicable.

The mean apparent permeability (P_{app} , $\times 10^{-6}$ cm/sec) of the test compounds, atenolol, propranolol and digoxin were calculated as follows:

$$P_{app} = \frac{dq}{dt} \times \frac{1}{C_0} \times \frac{1}{A}$$

- 10 where dq/dt = rate of transport (rate of transport of compound in the receiver compartment), C_0 = initial concentration in the donor compartment, A = surface area of the effective filter membrane.

HepaRG-based assay protocol

- 15 A cryopreserved vial of differentiated HepaRG cells (Biopredic International HPR116080) is thawed in HepaRG Thawing/Plating/General Purpose Medium (Biopredic International ADD670C) supplemented with 200 mM Glutamax (Gibco 35050061) following the protocol provided by Biopredic International. 70,000 cells per well are seeded in 96-wells plate (Corning CLS3809) in 100 μL of HepaRG Thawing/Plating/General Purpose Medium supplemented with 200 mM Glutamax and
- 20 incubated at 37 °C in 5% CO₂ for 24 hours. Post incubation, the seeding media is replaced by HepaRG Maintenance/Metabolism Medium (Biopredic International ADD620C) and incubated for 6 days, with fresh HepaRG Maintenance/Metabolism Medium replenishment every 48 hours. After 7 days incubation post seeding, incubation media is decanted from the wells and cells are washed two times with 250 μL of William's E Basal Media (Gibco 12551032). After decanting William's E Basal Media
- 25 each time, plates are tapped against paper towel to ensure maximum removal of residual media. Incubation mix is prepared by adding test inhibitor dilutions (3-fold serial dilution in DMSO (Sigma D2650)) in William's E media (basal) containing 0.3 μM 3H-taurocholic acid (ARC ART-1368) and 7.5 μM cold taurocholic acid (Sigma T4009) (maintaining 0.2% final DMSO concentration). 50 μL of incubation mix containing test inhibitors is then added to the wells (in duplicate) and the plates are
- 30 incubated for 30 minutes in 5% CO₂ incubator at 37 °C. After incubation, the reaction is stopped by keeping the plates on ice water mix for 2-3 minutes and the incubation mix is then aspirated completely from the wells. The wells are washed two times with 250 μL of chilled unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10mM) HBSS (Gibco 14175079) (pH

7.4). The plates are tapped against a paper towel after every wash to ensure maximum removal of blocking buffer.

100 µL of MicroScint-20 (PerkinElmer 6013621) is added to the wells and kept overnight at room temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well, with normal plate orientation).

Preparation of test compound dilutions

All test compounds were provided in powder form at room temperature. 10 mM DMSO stocks of the test compounds were prepared, aliquoted and stored at -20 °C. From the 10 mM DMSO stock of the compounds, a 3-fold serial dilution in DMSO was prepared to get a total of 10 dilutions of the test compounds. 0.5 µL of this dilution in DMSO was added to 250 µL of FBS-free basal media containing 3H-taurocholic acid and cold taurocholic acid to prepare the incubation mixture.

Bioavailability studies

Male mice (C57BL/6 or CD1) or Wistar rats of 8-9 weeks old were used. For each test compound, two groups of 3 animals each were used. One group was administered a single intravenous dose of 1 mg/kg (vehicle 100% DMSO) through the tail vein and the other group was administered a single oral dose of 10 mg/kg through gavage needle. The group that was administered an oral dose was fasted overnight. Blood samples were collected after 0.083, 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours following intravenous administration, and after 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours following oral administration. Blood samples were taken from saphenous vein. 0.2% EDTA was used as the anticoagulant. The samples were analyzed by a discovery grade bioanalytical method developed for the estimation of test compound in plasma, using an LC-MS/MS system.

Results

Biological data for the compounds of the examples is shown in Table 8 below.

Table 8

Example	hLBAT IC ₅₀ (nM)	hIBAT IC ₅₀ (nM)	Permeability (Caco-2)		Bioavailability (%)
			P _{app} A2B (x 10 ⁻⁶ cm/sec)	ER	
1	2908	7	9.2	0.9	34 (C57BL/6) 59 (CD1) 98 (Na ⁺ salt; rat)
2	736	4			
3	283	23			
4	3654	2	7.6	2.4	45 (CD1) 90 (rat)
5	1882	5	1.0	4.2	
6	3658	14			
7	2225	2	0.4	1.9	
8	2633	12	9.5	0.5	
9	1617	21	10.0	0.5	
10	7205	2	2.1	3.1	
11	1175	3	13.1	1.2	
12	2222	3	14.2	1.1	
13	2222	15			
14	342	6	0.0		

PD model: Evaluation of test compound on total bile acids levels in male C57BL/6 mice.

- 5 C57BL/6N Tac mice of 8-9 weeks old are used to study the effect of bile acid modulators on bile acid levels. After completion of quarantine and acclimatization period, animals are randomized based on bodyweight into x experimental groups: (i) vehicle control, and (ii) test compound y mg/kg po once daily. Animals are treated with test compound for 7 days. On day 5 of the study, animals are individually housed in fresh cages. On day 7, feces are collected from each cage, followed by blood withdrawal from each animal through retro-orbital route. Animals are euthanized to collect liver and terminal ileum from each animal for further analysis. Bodyweight and food consumption are measured twice weekly. Serum lipid profiles are analyzed in serum samples of day 7. Total bile acids in serum is measured in the serum samples of day 7. Fecal bile excretion is measured in the fecal
- 10

sample of day 7. Hepatic expression of CYP7A1 and SHP are quantified in the liver samples of day 7. Liver triglycerides and total cholesterol are analyzed in the liver samples of day 7.

Urine bile acid model: Evaluation of test compounds on urine bile acid levels in male C57BL/6 mice.


5

C57BL/6N Tac mice of 8-9 weeks old are used to study the effect of bile acid modulators on bile acid levels. After completion of quarantine and acclimatization period, animals are randomized based on bodyweight into x experimental groups: (i) vehicle control, and (ii) test compound y mg/kg po once daily. Animals are treated with test compound for 7 days. On day 6 of the study, animals are

10 transferred to a metabolic cage. On day 7, feces and urine are collected from each metabolic cage, followed by blood withdrawal from each animal through retro-orbital route. Animals are euthanized to collect kidney from each animal for further analysis. Bodyweight is measured twice weekly. Total bile acids in serum is measured in serum samples of day 7. Fecal bile acid excretion is measured in the fecal sample of day 7. Urine excretion of bile acids is measured in the sample of day 7. Kidney
15 expression of ASBT, OSTa, OSTAb and MRP2 is quantified in the samples of day 7.

The claims defining the invention are as follows:

1. A compound of formula (I)



wherein

R^1 and R^2 are each independently C_{1-4} alkyl;

R^3 is independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, cyano, nitro, amino, N -(C_{1-4} alkyl)amino, N,N -di(C_{1-4} alkyl)amino and N -(aryl- C_{1-4} alkyl)amino;

n is an integer 1, 2 or 3;

R^4 is selected from the group consisting of hydrogen, halogen, hydroxy, cyano, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyloxy, C_{1-4} alkylthio, C_{3-6} cycloalkylthio, amino, N -(C_{1-4} alkyl)amino and N,N -di(C_{1-4} alkyl)amino;

R^{5A} , R^{5B} , R^{5C} and R^{5D} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, amino and C_{1-4} alkyl; and

R^6 is selected from the group consisting of hydrogen and C_{1-4} alkyl;
or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein R^1 is n-butyl.
3. A compound according to claim 1 or 2, wherein R^2 is n-butyl.
4. A compound according to claim 1 or 2, wherein R^2 is ethyl.

5. A compound according to any one of claims 1 to 4, wherein R³ is independently selected from the group consisting of hydrogen, halogen, hydroxy, amino, cyano, C₁₋₄ haloalkyl, C₁₋₄ alkoxy and C₁₋₄ haloalkoxy.
6. A compound according to any one of claims 1 to 5, wherein R⁴ is selected from the group consisting of halogen, hydroxy, cyano, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, amino, *N*-(C₁₋₄ alkyl)amino and *N,N*-di(C₁₋₄ alkyl)amino.
7. A compound according to any one of claims 1 to 6, wherein R^{5A} and R^{5B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, amino and methyl.
8. A compound according to any one of claims 1 to 7, wherein R⁶ is hydrogen or methyl.
9. A compound according to claim 1, selected from the group consisting of:
 - 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydrobenzo-1,2,5-thiadiazepin-8-yl)oxy)propanoic acid;
 - 3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;
 - (*S*)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;
 - (*R*)-3-((3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;
 - 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;
 - (*S*)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;
 - (*R*)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;
 - 3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;
 - (*S*)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;
 - (*R*)-3-((3,3-dibutyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)butanoic acid;

3-((3,3-dibutyl-7-chloro-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

3-((3,3-dibutyl-5-(4-hydroxyphenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

3-((3-Butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

(S)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

(R)-3-((3-butyl-7-(dimethylamino)-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)propanoic acid;

O-(3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

(S)-O-((R)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

(R)-O-((R)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

(S)-O-((S)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine; and

(R)-O-((S)-3-butyl-3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)serine;

or a pharmaceutically acceptable salt thereof.

10. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of claims 1 to 9, and one or more pharmaceutically acceptable excipients.
11. A method of treatment or prevention of a condition, disease or disorder mediated by inhibition of bile acid circulation in a subject in need thereof comprising administering to the subject an effective amount of a compound according to any one of claims 1 to 9 or the pharmaceutical composition of claim 10.
12. Use of a compound according to any one of claims 1 to 9 or the pharmaceutical composition of claim 10 in the preparation of a medicament for treatment or prevention of a condition, disease or disorder mediated by inhibition of bile acid circulation.

13. The method according to claim 11 or the use according to claim 12, wherein the condition, disease or disorder is a cardiovascular disease or a disorder of fatty acid metabolism or a glucose utilization disorder selected from hypercholesterolemia; disorders of fatty acid metabolism; type 1 and type 2 diabetes mellitus; complications of diabetes, including cataracts, micro- and macrovascular diseases, retinopathy, neuropathy, nephropathy and delayed wound healing, tissue ischaemia, diabetic foot, arteriosclerosis, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, stroke, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders and vascular restenosis; diabetes-related diseases selected from insulin resistance (impaired glucose homeostasis), hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, dyslipidemia, hyperlipidemia including hypertriglyceridemia, metabolic syndrome (syndrome X), atherosclerosis and hypertension; and low high density lipoprotein levels.
14. The method according to claim 11 or the use according to claim 12, wherein the condition, disease or disorder is a gastrointestinal disease or disorder, selected from constipation (including chronic constipation, functional constipation, chronic idiopathic constipation (CIC), intermittent/sporadic constipation, constipation secondary to diabetes mellitus, constipation secondary to stroke, constipation secondary to chronic kidney disease, constipation secondary to multiple sclerosis, constipation secondary to Parkinson's disease, constipation secondary to systemic sclerosis, drug induced constipation, irritable bowel syndrome with constipation (IBS-C), irritable bowel syndrome mixed (IBS-M), pediatric functional constipation and opioid induced constipation); Crohn's disease; primary bile acid malabsorption; irritable bowel syndrome (IBS); inflammatory bowel disease (IBD); ileal inflammation; and reflux disease and complications thereof, selected from Barrett's esophagus, bile reflux esophagitis and bile reflux gastritis.
15. The method according to claim 11 or the use according to claim 12, wherein the condition, disease or disorder is a liver disease or disorder, selected from inherited metabolic disorder of the liver; inborn errors of bile acid synthesis; congenital bile duct anomalies; biliary atresia; post-Kasai biliary atresia; post-liver transplantation biliary atresia; neonatal hepatitis; neonatal cholestasis; hereditary forms of cholestasis; cerebrotendinous xanthomatosis; a secondary defect of BA synthesis; Zellweger's syndrome; cystic fibrosis-associated liver disease; alpha1-antitrypsin deficiency; Alagilles syndrome (ALGS); Byler syndrome; a primary defect of bile acid (BA) synthesis; progressive familial intrahepatic cholestasis (PFIC) including PFIC-1, PFIC-2, PFIC-3 and non-specified PFIC, post-biliary diversion PFIC and post-liver transplant PFIC; benign

recurrent intrahepatic cholestasis (BRIC) including BRIC1, BRIC2 and non-specified BRIC, post-biliary diversion BRIC and post-liver transplant BRIC; autoimmune hepatitis; primary biliary cirrhosis (PBC); liver fibrosis; non-alcoholic fatty liver disease (NAFLD); non-alcoholic steatohepatitis (NASH); portal hypertension; cholestasis; Down syndrome cholestasis; drug-induced cholestasis; intrahepatic cholestasis of pregnancy (jaundice during pregnancy); intrahepatic cholestasis; extrahepatic cholestasis; parenteral nutrition associated cholestasis (PNAC); low phospholipid-associated cholestasis; lymphedema cholestasis syndrome 1 (LSC1); primary sclerosing cholangitis (PSC); immunoglobulin G4 associated cholangitis; primary biliary cholangitis; cholelithiasis (gall stones); biliary lithiasis; choledocholithiasis; gallstone pancreatitis; Caroli disease; malignancy of bile ducts; malignancy causing obstruction of the biliary tree; biliary strictures; AIDS cholangiopathy; ischemic cholangiopathy; pruritus due to cholestasis or jaundice; pancreatitis; chronic autoimmune liver disease leading to progressive cholestasis; hepatic steatosis; alcoholic hepatitis; acute fatty liver; fatty liver of pregnancy; drug-induced hepatitis; iron overload disorders; congenital bile acid synthesis defect type 1 (BAS type 1); drug-induced liver injury (DILI); hepatic fibrosis; congenital hepatic fibrosis; hepatic cirrhosis; Langerhans cell histiocytosis (LCH); neonatal ichthyosis sclerosing cholangitis (NISCH); erythropoietic protoporphyria (EPP); idiopathic adulthood ductopenia (IAD); idiopathic neonatal hepatitis (INH); non syndromic paucity of interlobular bile ducts (NS PILBD); North American Indian childhood cirrhosis (NAIC); hepatic sarcoidosis; amyloidosis; necrotizing enterocolitis; serum bile acid-caused toxicities, including cardiac rhythm disturbances in setting of abnormal serum bile acid profile, cardiomyopathy associated with liver cirrhosis ("cholecardia"), and skeletal muscle wasting associated with cholestatic liver disease; viral hepatitis (including hepatitis A, hepatitis B, hepatitis C, hepatitis D and hepatitis E); hepatocellular carcinoma (hepatoma); cholangiocarcinoma; bile acid-related gastrointestinal cancers; and cholestasis caused by tumours and neoplasms of the liver, of the biliary tract and of the pancreas; or enhancement of corticosteroid therapy in liver disease.

16. The method according to claim 11 or the use according to claim 12, wherein the condition, disease or disorder is hyperabsorption syndrome (including abetalipoproteinemia, familial hypobetalipoproteinemia (FHBL), chylomicron retention disease (CRD) and sitosterolemia); hypervitaminosis and osteopetrosis; hypertension; glomerular hyperfiltration; and pruritus of renal failure; or protection against liver- or metabolic disease-associated kidney injury.

SEQUENCE LISTING

<110> ALBIREO AB

<120> BENZOTHIADIAZEPINE COMPOUNDS AND THEIR USE AS BILE ACID MODULATORS

<130> NP0461WO

<150> IN 201911004690

<151> 2019-02-06

<150> SE 1950464-6

<151> 2019-04-12

<150> IN 201911049981

<151> 2019-12-04

<160> 4

<170> PatentIn version 3.5

<210> 1

<211> 1251

<212> PRT

<213> Homo sapiens

<400> 1

Met	Ser	Thr	Glu	Arg	Asp	Ser	Glu	Thr	Thr	Phe	Asp	Glu	Asp	Ser	Gln
1				5					10					15	

Pro	Asn	Asp	Glu	Val	Val	Pro	Tyr	Ser	Asp	Asp	Glu	Thr	Glu	Asp	Glu
			20					25					30		

Leu	Asp	Asp	Gln	Gly	Ser	Ala	Val	Glu	Pro	Glu	Gln	Asn	Arg	Val	Asn
		35					40					45			

Arg	Glu	Ala	Glu	Glu	Asn	Arg	Glu	Pro	Phe	Arg	Lys	Glu	Cys	Thr	Trp
	50					55					60				

Gln	Val	Lys	Ala	Asn	Asp	Arg	Lys	Tyr	His	Glu	Gln	Pro	His	Phe	Met
65					70					75					80

Asn	Thr	Lys	Phe	Leu	Cys	Ile	Lys	Glu	Ser	Lys	Tyr	Ala	Asn	Asn	Ala
				85					90					95	

Ile	Lys	Thr	Tyr	Lys	Tyr	Asn	Ala	Phe	Thr	Phe	Ile	Pro	Met	Asn	Leu
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

100

105

110

Phe Glu Gln Phe Lys Arg Ala Ala Asn Leu Tyr Phe Leu Ala Leu Leu
 115 120 125

Ile Leu Gln Ala Val Pro Gln Ile Ser Thr Leu Ala Trp Tyr Thr Thr
 130 135 140

Leu Val Pro Leu Leu Val Val Leu Gly Val Thr Ala Ile Lys Asp Leu
 145 150 155 160

Val Asp Asp Val Ala Arg His Lys Met Asp Lys Glu Ile Asn Asn Arg
 165 170 175

Thr Cys Glu Val Ile Lys Asp Gly Arg Phe Lys Val Ala Lys Trp Lys
 180 185 190

Glu Ile Gln Val Gly Asp Val Ile Arg Leu Lys Lys Asn Asp Phe Val
 195 200 205

Pro Ala Asp Ile Leu Leu Leu Ser Ser Ser Glu Pro Asn Ser Leu Cys
 210 215 220

Tyr Val Glu Thr Ala Glu Leu Asp Gly Glu Thr Asn Leu Lys Phe Lys
 225 230 235 240

Met Ser Leu Glu Ile Thr Asp Gln Tyr Leu Gln Arg Glu Asp Thr Leu
 245 250 255

Ala Thr Phe Asp Gly Phe Ile Glu Cys Glu Glu Pro Asn Asn Arg Leu
 260 265 270

Asp Lys Phe Thr Gly Thr Leu Phe Trp Arg Asn Thr Ser Phe Pro Leu
 275 280 285

Asp Ala Asp Lys Ile Leu Leu Arg Gly Cys Val Ile Arg Asn Thr Asp
 290 295 300

Phe Cys His Gly Leu Val Ile Phe Ala Gly Ala Asp Thr Lys Ile Met
 305 310 315 320

Lys Asn Ser Gly Lys Thr Arg Phe Lys Arg Thr Lys Ile Asp Tyr Leu
325 330 335

Met Asn Tyr Met Val Tyr Thr Ile Phe Val Val Leu Ile Leu Leu Ser
340 345 350

Ala Gly Leu Ala Ile Gly His Ala Tyr Trp Glu Ala Gln Val Gly Asn
355 360 365

Ser Ser Trp Tyr Leu Tyr Asp Gly Glu Asp Asp Thr Pro Ser Tyr Arg
370 375 380

Gly Phe Leu Ile Phe Trp Gly Tyr Ile Ile Val Leu Asn Thr Met Val
385 390 395 400

Pro Ile Ser Leu Tyr Val Ser Val Glu Val Ile Arg Leu Gly Gln Ser
405 410 415

His Phe Ile Asn Trp Asp Leu Gln Met Tyr Tyr Ala Glu Lys Asp Thr
420 425 430

Pro Ala Lys Ala Arg Thr Thr Thr Leu Asn Glu Gln Leu Gly Gln Ile
435 440 445

His Tyr Ile Phe Ser Asp Lys Thr Gly Thr Leu Thr Gln Asn Ile Met
450 455 460

Thr Phe Lys Lys Cys Cys Ile Asn Gly Gln Ile Tyr Gly Asp His Arg
465 470 475 480

Asp Ala Ser Gln His Asn His Asn Lys Ile Glu Gln Val Asp Phe Ser
485 490 495

Trp Asn Thr Tyr Ala Asp Gly Lys Leu Ala Phe Tyr Asp His Tyr Leu
500 505 510

Ile Glu Gln Ile Gln Ser Gly Lys Glu Pro Glu Val Arg Gln Phe Phe
515 520 525

Phe Leu Leu Ala Val Cys His Thr Val Met Val Asp Arg Thr Asp Gly

530

535

540

Gln Leu Asn Tyr Gln Ala Ala Ser Pro Asp Glu Gly Ala Leu Val Asn
 545 550 555 560

Ala Ala Arg Asn Phe Gly Phe Ala Phe Leu Ala Arg Thr Gln Asn Thr
 565 570 575

Ile Thr Ile Ser Glu Leu Gly Thr Glu Arg Thr Tyr Asn Val Leu Ala
 580 585 590

Ile Leu Asp Phe Asn Ser Asp Arg Lys Arg Met Ser Ile Ile Val Arg
 595 600 605

Thr Pro Glu Gly Asn Ile Lys Leu Tyr Cys Lys Gly Ala Asp Thr Val
 610 615 620

Ile Tyr Glu Arg Leu His Arg Met Asn Pro Thr Lys Gln Glu Thr Gln
 625 630 635 640

Asp Ala Leu Asp Ile Phe Ala Asn Glu Thr Leu Arg Thr Leu Cys Leu
 645 650 655

Cys Tyr Lys Glu Ile Glu Glu Lys Glu Phe Thr Glu Trp Asn Lys Lys
 660 665 670

Phe Met Ala Ala Ser Val Ala Ser Thr Asn Arg Asp Glu Ala Leu Asp
 675 680 685

Lys Val Tyr Glu Glu Ile Glu Lys Asp Leu Ile Leu Leu Gly Ala Thr
 690 695 700

Ala Ile Glu Asp Lys Leu Gln Asp Gly Val Pro Glu Thr Ile Ser Lys
 705 710 715 720

Leu Ala Lys Ala Asp Ile Lys Ile Trp Val Leu Thr Gly Asp Lys Lys
 725 730 735

Glu Thr Ala Glu Asn Ile Gly Phe Ala Cys Glu Leu Leu Thr Glu Asp
 740 745 750

Thr Thr Ile Cys Tyr Gly Glu Asp Ile Asn Ser Leu Leu His Ala Arg
755 760 765

Met Glu Asn Gln Arg Asn Arg Gly Gly Val Tyr Ala Lys Phe Ala Pro
770 775 780

Pro Val Gln Glu Ser Phe Phe Pro Pro Gly Gly Asn Arg Ala Leu Ile
785 790 795 800

Ile Thr Gly Ser Trp Leu Asn Glu Ile Leu Leu Glu Lys Lys Thr Lys
805 810 815

Arg Asn Lys Ile Leu Lys Leu Lys Phe Pro Arg Thr Glu Glu Glu Arg
820 825 830

Arg Met Arg Thr Gln Ser Lys Arg Arg Leu Glu Ala Lys Lys Glu Gln
835 840 845

Arg Gln Lys Asn Phe Val Asp Leu Ala Cys Glu Cys Ser Ala Val Ile
850 855 860

Cys Cys Arg Val Thr Pro Lys Gln Lys Ala Met Val Val Asp Leu Val
865 870 875 880

Lys Arg Tyr Lys Lys Ala Ile Thr Leu Ala Ile Gly Asp Gly Ala Asn
885 890 895

Asp Val Asn Met Ile Lys Thr Ala His Ile Gly Val Gly Ile Ser Gly
900 905 910

Gln Glu Gly Met Gln Ala Val Met Ser Ser Asp Tyr Ser Phe Ala Gln
915 920 925

Phe Arg Tyr Leu Gln Arg Leu Leu Leu Val His Gly Arg Trp Ser Tyr
930 935 940

Ile Arg Met Cys Lys Phe Leu Arg Tyr Phe Phe Tyr Lys Asn Phe Ala
945 950 955 960

Phe Thr Leu Val His Phe Trp Tyr Ser Phe Phe Asn Gly Tyr Ser Ala

965

970

975

Gln Thr Ala Tyr Glu Asp Trp Phe Ile Thr Leu Tyr Asn Val Leu Tyr
 980 985 990

Thr Ser Leu Pro Val Leu Leu Met Gly Leu Leu Asp Gln Asp Val Ser
 995 1000 1005

Asp Lys Leu Ser Leu Arg Phe Pro Gly Leu Tyr Ile Val Gly Gln
 1010 1015 1020

Arg Asp Leu Leu Phe Asn Tyr Lys Arg Phe Phe Val Ser Leu Leu
 1025 1030 1035

His Gly Val Leu Thr Ser Met Ile Leu Phe Phe Ile Pro Leu Gly
 1040 1045 1050

Ala Tyr Leu Gln Thr Val Gly Gln Asp Gly Glu Ala Pro Ser Asp
 1055 1060 1065

Tyr Gln Ser Phe Ala Val Thr Ile Ala Ser Ala Leu Val Ile Thr
 1070 1075 1080

Val Asn Phe Gln Ile Gly Leu Asp Thr Ser Tyr Trp Thr Phe Val
 1085 1090 1095

Asn Ala Phe Ser Ile Phe Gly Ser Ile Ala Leu Tyr Phe Gly Ile
 1100 1105 1110

Met Phe Asp Phe His Ser Ala Gly Ile His Val Leu Phe Pro Ser
 1115 1120 1125

Ala Phe Gln Phe Thr Gly Thr Ala Ser Asn Ala Leu Arg Gln Pro
 1130 1135 1140

Tyr Ile Trp Leu Thr Ile Ile Leu Ala Val Ala Val Cys Leu Leu
 1145 1150 1155

Pro Val Val Ala Ile Arg Phe Leu Ser Met Thr Ile Trp Pro Ser
 1160 1165 1170

Glu Ser Asp Lys Ile Gln Lys His Arg Lys Arg Leu Lys Ala Glu
1175 1180 1185

Glu Gln Trp Gln Arg Arg Gln Gln Val Phe Arg Arg Gly Val Ser
1190 1195 1200

Thr Arg Arg Ser Ala Tyr Ala Phe Ser His Gln Arg Gly Tyr Ala
1205 1210 1215

Asp Leu Ile Ser Ser Gly Arg Ser Ile Arg Lys Lys Arg Ser Pro
1220 1225 1230

Leu Asp Ala Ile Val Ala Asp Gly Thr Ala Glu Tyr Arg Arg Thr
1235 1240 1245

Gly Asp Ser
1250

<210> 2
<211> 3756
<212> DNA
<213> Homo sapiens

<400> 2
atgagtacag aaagagactc agaaacgaca ttgacgagg attctcagcc taatgacgaa 60
gtggttccct acagtgatga tgaaacagaa gatgaacttg atgaccaggg gtctgctgtt 120
gaaccagaac aaaaccgagt caacagggaa gcagaggaga accgggagcc attcagaaaa 180
gaatgtacat ggcaagtcaa agcaaacgat cgcaagtacc acgaacaacc tcactttatg 240
aacacaaaat tcttgtgtat taaggagagt aaatatgcga ataatgcaat taaaacatac 300
aagtacaacg catttacctt tataccaatg aatctgtttg agcagtttaa gagagcagcc 360
aatttatatt tcctggctct tcttatctta caggcagttc ctcaaattct taccctggct 420
tggtacacca cactagtgcc cctgcttggt gtgctgggctg tcaactgcaat caaagacctg 480
gtggacgatg tggctcgcca taaaatggat aaggaaatca acaataggac gtgtgaagtc 540
attaaggatg gcaggttcaa agttgctaag tggaaagaaa ttcaagttgg agacgtcatt 600
cgtctgaaaa aaaatgattt tgttccagct gacattctcc tgctgtctag ctctgagcct 660
aacagcctct gctatgtgga aacagcagaa ctggatggag aaaccaattt aaaatttaag 720

atgtcacttg	aatcacaga	ccagtacctc	caaagagaag	atacattggc	tacatttgat	780
ggttttattg	aatgtgaaga	acccaataac	agactagata	agtttacagg	aacactattt	840
tggagaaaca	caagttttcc	tttggatgct	gataaaatth	tgttacgtgg	ctgtgtaatt	900
aggaacaccg	atttctgcca	cggcttagtc	atthtttgag	gtgctgacac	taaaataatg	960
aagaatagt	ggaaaaccag	atttaaaaga	actaaaattg	attacttgat	gaactacatg	1020
gtttacacga	tctttgttgt	tcttattctg	ctttctgctg	gtcttgccat	cggccatgct	1080
tattgggaag	cacaggtggg	caattcctct	tggtacctct	atgatggaga	agacgataca	1140
ccctcctacc	gtggattcct	cattttcttg	ggctatatca	ttgttctcaa	caccatggta	1200
cccattctct	tctatgtcag	cgtggaagt	attcgtcttg	gacagagtca	cttcatcaac	1260
tgggacctgc	aatgtacta	tgctgagaag	gacacacccg	caaaagctag	aaccaccaca	1320
ctcaatgaac	agctcgggca	gatccattat	atcttctctg	ataagacggg	gacactcaca	1380
caaaatatca	tgacctttaa	aaagtgtgt	atcaacgggc	agatatatgg	ggaccatcgg	1440
gatgcctctc	aacacaacca	caacaaaata	gagcaagttg	atthtagctg	gaatacatat	1500
gctgatggga	agcttgcat	ttatgaccac	tatcttattg	agcaaatcca	gtcagggaaa	1560
gagccagaag	tacgacagtt	cttcttcttg	ctcgcagttt	gccacacagt	catggtggat	1620
aggactgatg	gtcagctcaa	ctaccaggca	gcctctcccg	atgaagggtg	cctggtaaac	1680
gctgccagga	actttggctt	tgcttctctc	gccaggagcc	agaacaccat	caccatcagt	1740
gaactgggca	ctgaaaggac	ttacaatgtt	cttgccattt	tggacttcaa	cagtgaccgg	1800
aagcgaatgt	ctatcattgt	aagaacccca	gaaggcaata	tcaagcttta	ctgtaaagg	1860
gctgacactg	ttatthtatga	acggttacat	cgaatgaatc	ctactaagca	agaaacacag	1920
gatgccctgg	atatctttgc	aatgaaact	cttagaacc	tatgcctttg	ctacaaggaa	1980
attgaagaaa	aagaatthac	agaatggaat	aaaaagttta	tggctgccag	tgtggcctcc	2040
accaaccggg	acgaagctct	ggataaagta	tatgaggaga	ttgaaaaaga	cttaattctc	2100
ctgggagcta	cagctattga	agacaagcta	caggatggag	ttccagaaac	catttcaaaa	2160
cttgcaaaag	ctgacattaa	gatctgggtg	cttactggag	acaaaaagga	aactgctgaa	2220
aatataggat	ttgcttgatga	acttctgact	gaagacacca	ccatctgcta	tggggaggat	2280
attaattctc	ttcttcatgc	aaggatggaa	aaccagagga	atagagggtg	cgtctacgca	2340

aagtttgcac ctctgtgca ggaatctttt tttccacccg gtggaaaccg tgccttaatc	2400
atcactgggtt cttggttgaa tgaaattctt ctcgagaaaa agaccaagag aaataagatt	2460
ctgaagctga agttcccaag aacagaagaa gaaagacgga tgcggaccca aagtaaaagg	2520
aggctagaag ctaagaaaga gcagcggcag aaaaactttg tggacctggc ctgcgagtgc	2580
agcgcagtca tctgctgccg cgtcaccccc aagcagaagg ccatgggtgg ggacctgggt	2640
aagaggtaca agaaagccat cacgctggcc atcggagatg gggccaatga cgtgaacatg	2700
atcaaaactg cccacattgg cgttggaaata agtggacaag aaggaatgca agctgtcatg	2760
tcgagtgact attcctttgc tcagttccga tatctgcaga ggctactgct ggtgcatggc	2820
cgatgggtctt acataaggat gtgcaagttc ctacgatact tcttttacia aaactttgcc	2880
tttactttgg ttcatctctg gtactccttc ttcaatggct actctgcgca gactgcatac	2940
gaggattgggt tcatcacctt ctacaacgtg ctgtacacca gcctgcccgt gctcctcatg	3000
gggctgctcg accaggatgt gagtgcacaa ctgagcctcc gattccctgg gttatacata	3060
gtgggacaaa gagacttact attcaactat aagagattct ttgtaagctt gttgcatggg	3120
gtcctaacat cgatgatcct cttcttcata cctcttggag cttatctgca aaccgtaggg	3180
caggatggag aggcaccttc cgactaccag tcttttgccg tcaccattgc ctctgctctt	3240
gtaataacag tcaattttcca gattggcttg gatacttctt attggacttt tgtgaatgct	3300
ttttcaattt ttggaagcat tgcactttat tttggcatca tgtttgactt tcatagtgtc	3360
ggaatacatg ttctctttcc atctgcattt caatttacag gcacagcttc aaacgctctg	3420
agacagccat acatttggtt aactatcatc ctggctgttg ctgtgtgctt actaccgctc	3480
gttgccattc gattcctgtc aatgaccatc tggccatcag aaagtgataa gatccagaag	3540
catcgcaagc ggttgaaggc ggaggagcag tggcagcgac ggcagcaggt gttccgccgg	3600
ggcgtgtcaa cgcggcgctc ggcctacgcc ttctcgcacc agcggggcta cgcggaacctc	3660
atctcctccg ggcgcagcat ccgcaagaag cgctcgccgc ttgatgccat cgtggcggat	3720
ggcaccgcgg agtacaggcg caccggggac agctga	3756

<210> 3
 <211> 1321
 <212> PRT
 <213> Homo sapiens

<400> 3

Met Ser Asp Ser Val Ile Leu Arg Ser Ile Lys Lys Phe Gly Glu Glu
1 5 10 15

Asn Asp Gly Phe Glu Ser Asp Lys Ser Tyr Asn Asn Asp Lys Lys Ser
20 25 30

Arg Leu Gln Asp Glu Lys Lys Gly Asp Gly Val Arg Val Gly Phe Phe
35 40 45

Gln Leu Phe Arg Phe Ser Ser Ser Thr Asp Ile Trp Leu Met Phe Val
50 55 60

Gly Ser Leu Cys Ala Phe Leu His Gly Ile Ala Gln Pro Gly Val Leu
65 70 75 80

Leu Ile Phe Gly Thr Met Thr Asp Val Phe Ile Asp Tyr Asp Val Glu
85 90 95

Leu Gln Glu Leu Gln Ile Pro Gly Lys Ala Cys Val Asn Asn Thr Ile
100 105 110

Val Trp Thr Asn Ser Ser Leu Asn Gln Asn Met Thr Asn Gly Thr Arg
115 120 125

Cys Gly Leu Leu Asn Ile Glu Ser Glu Met Ile Lys Phe Ala Ser Tyr
130 135 140

Tyr Ala Gly Ile Ala Val Ala Val Leu Ile Thr Gly Tyr Ile Gln Ile
145 150 155 160

Cys Phe Trp Val Ile Ala Ala Ala Arg Gln Ile Gln Lys Met Arg Lys
165 170 175

Phe Tyr Phe Arg Arg Ile Met Arg Met Glu Ile Gly Trp Phe Asp Cys
180 185 190

Asn Ser Val Gly Glu Leu Asn Thr Arg Phe Ser Asp Asp Ile Asn Lys
195 200 205

Ile Asn Asp Ala Ile Ala Asp Gln Met Ala Leu Phe Ile Gln Arg Met
210 215 220

Thr Ser Thr Ile Cys Gly Phe Leu Leu Gly Phe Phe Arg Gly Trp Lys
225 230 235 240

Leu Thr Leu Val Ile Ile Ser Val Ser Pro Leu Ile Gly Ile Gly Ala
245 250 255

Ala Thr Ile Gly Leu Ser Val Ser Lys Phe Thr Asp Tyr Glu Leu Lys
260 265 270

Ala Tyr Ala Lys Ala Gly Val Val Ala Asp Glu Val Ile Ser Ser Met
275 280 285

Arg Thr Val Ala Ala Phe Gly Gly Glu Lys Arg Glu Val Glu Arg Tyr
290 295 300

Glu Lys Asn Leu Val Phe Ala Gln Arg Trp Gly Ile Arg Lys Gly Ile
305 310 315 320

Val Met Gly Phe Phe Thr Gly Phe Val Trp Cys Leu Ile Phe Leu Cys
325 330 335

Tyr Ala Leu Ala Phe Trp Tyr Gly Ser Thr Leu Val Leu Asp Glu Gly
340 345 350

Glu Tyr Thr Pro Gly Thr Leu Val Gln Ile Phe Leu Ser Val Ile Val
355 360 365

Gly Ala Leu Asn Leu Gly Asn Ala Ser Pro Cys Leu Glu Ala Phe Ala
370 375 380

Thr Gly Arg Ala Ala Ala Thr Ser Ile Phe Glu Thr Ile Asp Arg Lys
385 390 395 400

Pro Ile Ile Asp Cys Met Ser Glu Asp Gly Tyr Lys Leu Asp Arg Ile
405 410 415

Lys Gly Glu Ile Glu Phe His Asn Val Thr Phe His Tyr Pro Ser Arg

420

425

430

Pro Glu Val Lys Ile Leu Asn Asp Leu Asn Met Val Ile Lys Pro Gly
 435 440 445

Glu Met Thr Ala Leu Val Gly Pro Ser Gly Ala Gly Lys Ser Thr Ala
 450 455 460

Leu Gln Leu Ile Gln Arg Phe Tyr Asp Pro Cys Glu Gly Met Val Thr
 465 470 475 480

Val Asp Gly His Asp Ile Arg Ser Leu Asn Ile Gln Trp Leu Arg Asp
 485 490 495

Gln Ile Gly Ile Val Glu Gln Glu Pro Val Leu Phe Ser Thr Thr Ile
 500 505 510

Ala Glu Asn Ile Arg Tyr Gly Arg Glu Asp Ala Thr Met Glu Asp Ile
 515 520 525

Val Gln Ala Ala Lys Glu Ala Asn Ala Tyr Asn Phe Ile Met Asp Leu
 530 535 540

Pro Gln Gln Phe Asp Thr Leu Val Gly Glu Gly Gly Gly Gln Met Ser
 545 550 555 560

Gly Gly Gln Lys Gln Arg Val Ala Ile Ala Arg Ala Leu Ile Arg Asn
 565 570 575

Pro Lys Ile Leu Leu Leu Asp Met Ala Thr Ser Ala Leu Asp Asn Glu
 580 585 590

Ser Glu Ala Met Val Gln Glu Val Leu Ser Lys Ile Gln His Gly His
 595 600 605

Thr Ile Ile Ser Val Ala His Arg Leu Ser Thr Val Arg Ala Ala Asp
 610 615 620

Thr Ile Ile Gly Phe Glu His Gly Thr Ala Val Glu Arg Gly Thr His
 625 630 635 640

Glu Glu Leu Leu Glu Arg Lys Gly Val Tyr Phe Thr Leu Val Thr Leu
645 650 655

Gln Ser Gln Gly Asn Gln Ala Leu Asn Glu Glu Asp Ile Lys Asp Ala
660 665 670

Thr Glu Asp Asp Met Leu Ala Arg Thr Phe Ser Arg Gly Ser Tyr Gln
675 680 685

Asp Ser Leu Arg Ala Ser Ile Arg Gln Arg Ser Lys Ser Gln Leu Ser
690 695 700

Tyr Leu Val His Glu Pro Pro Leu Ala Val Val Asp His Lys Ser Thr
705 710 715 720

Tyr Glu Glu Asp Arg Lys Asp Lys Asp Ile Pro Val Gln Glu Glu Val
725 730 735

Glu Pro Ala Pro Val Arg Arg Ile Leu Lys Phe Ser Ala Pro Glu Trp
740 745 750

Pro Tyr Met Leu Val Gly Ser Val Gly Ala Ala Val Asn Gly Thr Val
755 760 765

Thr Pro Leu Tyr Ala Phe Leu Phe Ser Gln Ile Leu Gly Thr Phe Ser
770 775 780

Ile Pro Asp Lys Glu Glu Gln Arg Ser Gln Ile Asn Gly Val Cys Leu
785 790 795 800

Leu Phe Val Ala Met Gly Cys Val Ser Leu Phe Thr Gln Phe Leu Gln
805 810 815

Gly Tyr Ala Phe Ala Lys Ser Gly Glu Leu Leu Thr Lys Arg Leu Arg
820 825 830

Lys Phe Gly Phe Arg Ala Met Leu Gly Gln Asp Ile Ala Trp Phe Asp
835 840 845

Asp Leu Arg Asn Ser Pro Gly Ala Leu Thr Thr Arg Leu Ala Thr Asp

850

855

860

Ala Ser Gln Val Gln Gly Ala Ala Gly Ser Gln Ile Gly Met Ile Val
 865 870 875 880

Asn Ser Phe Thr Asn Val Thr Val Ala Met Ile Ile Ala Phe Ser Phe
 885 890 895

Ser Trp Lys Leu Ser Leu Val Ile Leu Cys Phe Phe Pro Phe Leu Ala
 900 905 910

Leu Ser Gly Ala Thr Gln Thr Arg Met Leu Thr Gly Phe Ala Ser Arg
 915 920 925

Asp Lys Gln Ala Leu Glu Met Val Gly Gln Ile Thr Asn Glu Ala Leu
 930 935 940

Ser Asn Ile Arg Thr Val Ala Gly Ile Gly Lys Glu Arg Arg Phe Ile
 945 950 955 960

Glu Ala Leu Glu Thr Glu Leu Glu Lys Pro Phe Lys Thr Ala Ile Gln
 965 970 975

Lys Ala Asn Ile Tyr Gly Phe Cys Phe Ala Phe Ala Gln Cys Ile Met
 980 985 990

Phe Ile Ala Asn Ser Ala Ser Tyr Arg Tyr Gly Gly Tyr Leu Ile Ser
 995 1000 1005

Asn Glu Gly Leu His Phe Ser Tyr Val Phe Arg Val Ile Ser Ala
 1010 1015 1020

Val Val Leu Ser Ala Thr Ala Leu Gly Arg Ala Phe Ser Tyr Thr
 1025 1030 1035

Pro Ser Tyr Ala Lys Ala Lys Ile Ser Ala Ala Arg Phe Phe Gln
 1040 1045 1050

Leu Leu Asp Arg Gln Pro Pro Ile Ser Val Tyr Asn Thr Ala Gly
 1055 1060 1065

Glu Lys Trp Asp Asn Phe Gln Gly Lys Ile Asp Phe Val Asp Cys
1070 1075 1080

Lys Phe Thr Tyr Pro Ser Arg Pro Asp Ser Gln Val Leu Asn Gly
1085 1090 1095

Leu Ser Val Ser Ile Ser Pro Gly Gln Thr Leu Ala Phe Val Gly
1100 1105 1110

Ser Ser Gly Cys Gly Lys Ser Thr Ser Ile Gln Leu Leu Glu Arg
1115 1120 1125

Phe Tyr Asp Pro Asp Gln Gly Lys Val Met Ile Asp Gly His Asp
1130 1135 1140

Ser Lys Lys Val Asn Val Gln Phe Leu Arg Ser Asn Ile Gly Ile
1145 1150 1155

Val Ser Gln Glu Pro Val Leu Phe Ala Cys Ser Ile Met Asp Asn
1160 1165 1170

Ile Lys Tyr Gly Asp Asn Thr Lys Glu Ile Pro Met Glu Arg Val
1175 1180 1185

Ile Ala Ala Ala Lys Gln Ala Gln Leu His Asp Phe Val Met Ser
1190 1195 1200

Leu Pro Glu Lys Tyr Glu Thr Asn Val Gly Ser Gln Gly Ser Gln
1205 1210 1215

Leu Ser Arg Gly Glu Lys Gln Arg Ile Ala Ile Ala Arg Ala Ile
1220 1225 1230

Val Arg Asp Pro Lys Ile Leu Leu Leu Asp Glu Ala Thr Ser Ala
1235 1240 1245

Leu Asp Thr Glu Ser Glu Lys Thr Val Gln Val Ala Leu Asp Lys
1250 1255 1260

Ala Arg Glu Gly Arg Thr Cys Ile Val Ile Ala His Arg Leu Ser

1265

1270

1275

Thr Ile Gln Asn Ala Asp Ile Ile Ala Val Met Ala Gln Gly Val
 1280 1285 1290

Val Ile Glu Lys Gly Thr His Glu Glu Leu Met Ala Gln Lys Gly
 1295 1300 1305

Ala Tyr Tyr Lys Leu Val Thr Thr Gly Ser Pro Ile Ser
 1310 1315 1320

<210> 4

<211> 3966

<212> DNA

<213> Homo sapiens

<400> 4

atgtctgact cagtaattct tcgaagtata aagaaatttg gagaggagaa tgatggtttt 60
 gagtcagata aatcatataa taatgataag aaatcaaggt tacaagatga gaagaaaggt 120
 gatggcgtaa gagttggctt ctttcaattg tttcggtttt cttcatcaac tgacatttgg 180
 ctgatgtttg tgggaagttt gtgtgcattt ctccatggaa tagcccagcc aggcgtgcta 240
 ctcatTTTTG gcacaatgac agatgttttt attgactacg acgttgagtt acaagaactc 300
 cagattccag gaaaagcatg tgtgaataac accattgtat ggactaacag ttccctcaac 360
 cagaacatga caaatggaac acgttgtggg ttgctgaaca tcgagagcga aatgatcaaa 420
 tttgccagtt actatgctgg aattgctgtc gcagtactta tcacaggata tattcaaata 480
 tgcttttggg tcattgccgc agctcgtcag atacagaaaa tgagaaaatt ttactttagg 540
 agaataatga gaatggaaat aggggtggtt gactgcaatt cagtggggga gctgaataca 600
 agatttcttg atgatattaa taaaatcaat gatgccatag ctgaccaaatt ggcccttttc 660
 attcagcgca tgacctcgac catctgtggt ttctgttgg gatttttcag gggttggaaa 720
 ctgaccttgg ttattatttc tgtcagccct ctcatggga ttggagcagc caccattggt 780
 ctgagtgtgt ccaagtttac ggactatgag ctgaaggcct atgccaaaagc aggggtggtg 840
 gctgatgaag tcatttcac aatgagaaca gtggctgctt ttggtggtga gaaaagagag 900
 gttgaaaggt atgagaaaaa tcttgtgttc gccagcggtt ggggaattag aaaaggaata 960
 gtgatgggat tctttactgg attcgtgtgg tgtctcatct ttttgtgtta tgcactggcc 1020

ttctggtacg gctccacact tgtcctggat gaaggagaat atacaccagg aacccttgtc	1080
cagattttcc tcagtgtcat agtaggagct ttaaattcttg gcaatgcctc tccttgtttg	1140
gaagcctttg caactggacg tgcagcagcc accagcattt ttgagacaat agacaggaaa	1200
cccatcattg actgcatgtc agaagatggg tacaagttgg atcgaatcaa ggggtgaaatt	1260
gaattccata atgtgacctt ccattatcct tccagaccag aggtgaagat tctaaatgac	1320
ctcaacatgg tcattaaacc aggggaaatg acagctctgg taggaccagcagg tggagctgga	1380
aaaagtacag cactgcaact cattcagcga ttctatgacc cctgtgaagg aatgggtgacc	1440
gtggatggcc atgacattcg ctctcttaac attcagtggc ttagagatca gattgggata	1500
gtggagcaag agccagttct gttctctacc accattgcag aaaatatctg ctatggcaga	1560
gaagatgcaa caatggaaga catagtccaa gctgccaaagg aggccaatgc ctacaacttc	1620
atcatggacc tgccacagca atttgacacc cttgttggag aaggaggagg ccagatgagt	1680
ggtagggcaga aacaaagggg agctatcgcc agagccctca tccgaaatcc caagattctg	1740
cttttggaca tggccacctc agctctggac aatgagagtg aagccatggg gcaagaagtg	1800
ctgagtaaga ttcagcatgg gcacacaatc atttcagttg ctcatcgctt gtctacggtc	1860
agagctgcag ataccatcat tggttttgaa catggcactg cagtggaaag agggacccat	1920
gaagaattac tggaaaggaa aggtgtttac ttcactctag tgactttgca aagccaggga	1980
aatcaagctc ttaatgaaga ggacataaag gatgcaactg aagatgacat gcttgcgagg	2040
acctttagca gagggagcta ccaggatagt ttaagggctt ccatccggca acgctccaag	2100
tctcagcttt cttacctggg gcacgaacct ccattagctg ttgtagatca taagtctacc	2160
tatgaagaag atagaaagga caaggacatt cctgtgcagg aagaagttga acctgcccc	2220
gtaggagga ttctgaaatt cagtgtcca gaatggccct acatgctggg aggggtctgtg	2280
ggtagcagctg tgaacgggac agtcacaccc ttgtatgcct ttttattcag ccagattctt	2340
gggacttttt caattcctga taaagaggaa caaagggtcac agatcaatgg tgtgtgccta	2400
ctttttgtag caatgggctg tgtatctctt ttcacccaat ttctacaggg atatgccttt	2460
gctaaatctg gggagctcct aacaaaaagg ctacgtaaatt ttggtttcag ggcaatgctg	2520
gggcaagata ttgcctgggt tgatgacctc agaaatagcc ctggagcatt gacaacaaga	2580
cttgctacag atgcttccca agttcaaggg gctgccggct ctcagatcgg gatgatagtc	2640

aattccttca ctaacgtcac tgtggccatg atcattgcct tctccttttag ctggaagctg	2700
agcctgggtca tcttgtgctt cttccccttc ttggctttat caggagccac acagaccagg	2760
atgttgacag gatttgcctc tcgagataag caggccctgg agatgggtggg acagattaca	2820
aatgaagccc tcagtaacat ccgcactgtt gctggaattg gaaaggagag gcggttcatt	2880
gaagcacttg agactgagct ggagaagccc ttcaagacag ccattcagaa agccaatatt	2940
tacggattct gctttgcctt tgcccagtgc atcatgttta ttgcgaattc tgcttcctac	3000
agatatggag gttacttaat ctccaatgag gggctccatt tcagctatgt gttcagggtg	3060
atctctgcag ttgtactgag tgcaacagct cttggaagag ccttctctta caccccaagt	3120
tatgcaaaag ctaaaatata agctgcacgc ttttttcaac tgctggaccg acaaccccca	3180
atcagtgtat acaatactgc aggtgaaaaa tgggacaact tccaggggaa gattgatttt	3240
gttgattgta aatttacata tccttctcga cctgactcgc aagttctgaa tggctcttca	3300
gtgtcgatta gtccagggca gacactggcg tttgttggga gcagtggatg tggcaaaagc	3360
actagcattc agctgttgga acgtttctat gatcctgatc aagggaaggt gatgatagat	3420
ggtcatgaca gcaaaaaagt aaatgtccag ttcctccgct caaacattgg aattgtttcc	3480
caggaaccag tgttgtttgc ctgtagcata atggacaata tcaagtatgg agacaacacc	3540
aaagaaattc ccatggaaag agtcatagca gctgcaaaac aggtcagct gcatgatttt	3600
gtcatgtcac tcccagagaa atatgaaact aacgttgggt cccaggggtc tcaactctct	3660
agagggggaga aacaacgcat tgctattgct cgggccattg tacgagatcc taaaatcttg	3720
ctactagatg aagccacttc tgccttagac acagaaagtg aaaagacggt gcaggttgct	3780
ctagacaaag ccagagaggg tcggacctgc attgtcattg cccatcgctt gtccaccatc	3840
cagaacgcgg atatcattgc tgtcatggca caggggggtg tgattgaaaa ggggacccat	3900
gaagaactga tggcccaaaa aggagcctac tacaactag tcaccactgg atcccccatc	3960
agttga	3966