





(86) Date de dépôt PCT/PCT Filing Date: 2009/07/22
 (87) Date publication PCT/PCT Publication Date: 2010/01/28
 (85) Entrée phase nationale/National Entry: 2011/01/13
 (86) N° demande PCT/PCT Application No.: EP 2009/059440
 (87) N° publication PCT/PCT Publication No.: 2010/010127
 (30) Priorité/Priority: 2008/07/23 (EP08161005.7)

(51) Cl.Int./Int.Cl. *A61K 31/397* (2006.01),
A61P 21/00 (2006.01)
 (71) Demandeur/Applicant:
NOVARTIS AG, CH
 (72) Inventeurs/Inventors:
AHMED, SYED SOHAIL, IT;
LONDEI, MARCO, US;
WRIGHT, TIMOTHY, US;
GERGELY, PETER, CH
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : MODULATEURS DU RECEPTEUR DE SPHINGOSINE 1 PHOSPHATE ET LEUR UTILISATION POUR LE
TRAITEMENT DE L'INFLAMMATION MUSCULAIRE
 (54) Title: SPHINGOSINE 1 PHOSPHATE RECEPTOR MODULATORS AND THEIR USE TO TREAT MUSCLE
INFLAMMATION





(57) **Abrégé/Abstract:**

The use of an S1P receptor modulator of the formula (Ia) or (Ib) wherein the meaning of the different residues is that indicated in claim 1, in the preparation of a medicament for preventing, inhibiting or treating an inflammatory condition selected from polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 January 2010 (28.01.2010)(10) International Publication Number
WO 2010/010127 A1

- (51) **International Patent Classification:**
A61K 31/397 (2006.01) A61P 21/00 (2006.01)
- (21) **International Application Number:**
PCT/EP2009/059440
- (22) **International Filing Date:**
22 July 2009 (22.07.2009)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
08161005.7 23 July 2008 (23.07.2008) EP
- (71) **Applicant (for all designated States except US):** NOVARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** AHMED, Syed Sohail [US/IT]; Via Socino, 56, I-53100 Siena (IT). LONDEI, Marco [IT/US]; 410 Pearl Street, La Jolla, California 92037 (US). WRIGHT, Timothy [US/US]; 471 Puritan Road, Swampscott, Massachusetts 01907 (US). GERGELY, Peter [HU/CH]; Auf der Wacht 14, CH-4104 Oberwil (CH).
- (74) **Agent:** HUTCHISON, John; Novartis Pharma AG, Patent Department, CH-4002 Basel (CH).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report (Art. 21(3))
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** SPHINGOSINE 1 PHOSPHATE RECEPTOR MODULATORS AND THEIR USE TO TREAT MUSCLE INFLAMMATION(57) **Abstract:** The use of an S1P receptor modulator of the formula (Ia) or (Ib) wherein the meaning of the different residues is that indicated in claim 1, in the preparation of a medicament for preventing, inhibiting or treating an inflammatory condition selected from polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis.

WO 2010/010127 A1

SPHINGOSINE 1 PHOSPHATE RECEPTOR MODULATORS AND THEIR USE TO TREAT MUSCLE INFLAMMATION

Field of the Invention

The present invention relates generally to specific sphingosine 1 phosphate (S1P) receptor modulators, and more specifically to their use to treat muscle inflammation such as polymyositis and other inflammatory conditions such as dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis.


Background


Polymyositis is an immune-mediated disorder leading to an inflammation of the muscles. It results in weakness that can be severe. Polymyositis is a chronic illness with periods of increased symptoms, called flares or relapses, and minimal or no symptoms, known as remissions.

The therapy of such muscle inflammation diseases, and in particular polymyositis, is only partially effective, and in most cases only offers a short delay in disease progression despite anti-inflammatory and immunosuppressive treatment. Accordingly, there is a need for agents which are effective in the inhibition or treatment of muscle diseases, e.g. polymyositis, including reduction of, alleviation of, stabilization of or relief from the symptoms which affect the muscles.

Summary of the invention

In accordance with a **first** aspect of the invention, there is provided the use of an S1P receptor modulator of the formula Ia or Ib:






wherein

A is $-\text{C}(\text{O})\text{OR}_5$, $-\text{OP}(\text{O})(\text{OR}_5)_2$, $-\text{P}(\text{O})(\text{OR}_5)_2$, $-\text{S}(\text{O})_2\text{OR}_5$, $-\text{P}(\text{O})(\text{R}_5)\text{OR}_5$ or 1*H*-tetrazol-5-yl, R_5 being H or C_{1-6} alkyl;

5 W is a bond, C_{1-3} alkylene or C_{2-3} alkenylene;

Y is C_{6-10} aryl or C_{2-9} heteroaryl e.g. C_{3-9} heteroaryl, optionally substituted by 1 to 3 radicals selected from halogen, $-\text{OH}$, $-\text{NO}_2$, C_{1-6} alkyl, C_{1-6} alkoxy; halo-substituted C_{1-6} alkyl and halo-substituted C_{1-6} alkoxy;

Z is chosen from:



10

wherein the left and right asterisks of Z indicate the point of attachment between $-\text{C}(\text{R}_3)(\text{R}_4)-$ and A of Formula Ia or Ib, respectively; R_6 is chosen from hydrogen and C_{1-6} alkyl; and J_1 and J_2 are independently methylene or a heteroatom chosen from S, O and NR_5 ; wherein R_5 is chosen from hydrogen and C_{1-6} alkyl; and any alkylene of Z can be

15 further substituted by one to three radicals chosen from halo, hydroxy, C_{1-6} alkyl; or R_6 can

be attached to a carbon atom of Y to form a 5-7 member ring e.g. a heterocyclic group as indicated in WO 04/103306A, e.g. azetidine;

R₁ is C₆₋₁₀aryl or C₂₋₉heteroaryl e.g. C₃₋₉heteroaryl, optionally substituted by C₁₋₆alkyl, C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₃₋₉heteroaryl, C₃₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl or C₃₋₈heterocycloalkylC₁₋₄alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R₁ may be substituted by 1 to 5 groups selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy and halo substituted-C₁₋₆alkyl or -C₁₋₆alkoxy; and any alkyl group of R₁ can optionally have a methylene replaced by an atom or group chosen from -S-, -S(O)-, -S(O)₂-, -NR₅- and -O-; wherein R₅ is chosen from hydrogen or C₁₋₆alkyl;

R₂ is H, C₁₋₆alkyl, halo substituted C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl; and each of R₃ or R₄, independently, is H, halogen, OH, C₁₋₆alkyl, C₁₋₆alkoxy or halo substituted C₁₋₆alkyl or C₁₋₆alkoxy;

and the N-oxide derivatives thereof or prodrugs thereof, individual isomers and mixtures of isomers thereof; and the pharmacologically acceptable salts, solvates or hydrates thereof,

in the preparation of a medicament for preventing, inhibiting or treating an inflammatory condition selected from polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis.

In accordance with a **second** aspect of the invention, there is provided an S1P receptor modulator as defined in the first aspect for use in preventing, inhibiting or treating an inflammatory condition selected from polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis.

In accordance with a **third** aspect of the invention, there is provided a method for preventing, inhibiting or treating an inflammatory condition such as polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis, in a subject in need thereof, comprising administering to said subject a therapeutically effective amount of an S1P receptor modulator as defined in the first aspect.

In accordance with a **fourth** aspect of the invention, there is provided a pharmaceutical composition comprising an S1P receptor modulator as defined in the first aspect for use in preventing, inhibiting or treating an inflammatory condition selected from polymyositis, dermatomyositis and nerve-muscle diseases e.g. muscular dystrophies and inclusion body myositis together with one or more pharmaceutically acceptable diluents or carriers therefor.

Brief description of the Figures



Figures 1(a) and 1(b) show the beneficial effects of the compounds of the invention on cytokine induced myotube atrophy.

- 5 Figure 1(a) shows myotube diameter following exposure to DMSO (control); varying concentrations of compound A; and cytokine with varying concentrations of compound A. Figure 1(b) shows myotube diameter as a percentage of the control myotube diameter (exposed to the DMSO control) following exposure to varying concentrations of compound A; and cytokine with varying concentrations of compound A.

10 **Detailed description of the invention**

S1P receptor modulators which can be used according to the invention are compounds of formula Ia or Ib, e.g. as disclosed in WO 04/103306A, WO 05/000833, WO 05/103309 or WO 05/113330, the contents of which are hereby incorporated by reference to the extent permitted by national law.

- 15 For example, the S1P receptor modulator may be a compound of formula Ia or Ib as defined below:



20


wherein

A is $-\text{C}(\text{O})\text{OR}_5$, $-\text{OP}(\text{O})(\text{OR}_5)_2$, $-\text{P}(\text{O})(\text{OR}_5)_2$, $-\text{S}(\text{O})_2\text{OR}_5$, $-\text{P}(\text{O})(\text{R}_5)\text{OR}_5$ or 1*H*-tetrazol-5-yl, R_5 being H or C_{1-6} alkyl;

W is a bond, C_{1-3} alkylene or C_{2-3} alkenylene;

Y is C₆₋₁₀aryl or C₂₋₉heteroaryl e.g. C₃₋₉heteroaryl, optionally substituted by 1 to 3 radicals selected from halogen, -OH, -NO₂, C₁₋₆alkyl, C₁₋₆alkoxy; halo-substituted C₁₋₆alkyl and halo-substituted C₁₋₆alkoxy;

Z is chosen from:



5

wherein the left and right asterisks of Z indicate the point of attachment between – C(R₃)(R₄)– and A of Formula Ia or Ib, respectively; R₆ is chosen from hydrogen and C₁₋₆alkyl; and J₁ and J₂ are independently methylene or a heteroatom chosen from S, O and NR₅; wherein R₅ is chosen from hydrogen and C₁₋₆alkyl; and any alkylene of Z can be further substituted by one to three radicals chosen from halo, hydroxy, C₁₋₆alkyl; or R₆ can be attached to a carbon atom of Y to form a 5-7 member ring e.g. a heterocyclic group as indicated in WO 04/103306A, e.g. azetidine;

R₁ is C₆₋₁₀aryl or C₂₋₉heteroaryl e.g. C₃₋₉heteroaryl, optionally substituted by C₁₋₆alkyl, C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₃₋₉heteroaryl, C₃₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl or C₃₋₈heterocycloalkylC₁₋₄alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R₁ may be substituted by 1 to 5 groups selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy and halo substituted-C₁₋₆alkyl or -C₁₋₆alkoxy; and any alkyl group of R₁ can optionally have a methylene replaced by an atom or group chosen

from $-S-$, $-S(O)-$, $-S(O)_2-$, $-NR_5-$ and $-O-$; wherein R_5 is chosen from hydrogen or C_{1-6} alkyl;

R_2 is H, C_{1-6} alkyl, halo substituted C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl; and

each of R_3 or R_4 , independently, is H, halogen, OH, C_{1-6} alkyl, C_{1-6} alkoxy or halo substituted

5 C_{1-6} alkyl or C_{1-6} alkoxy;

and the N-oxide derivatives thereof or prodrugs thereof, individual isomers and mixtures of isomers thereof; and the pharmacologically acceptable salts, solvates or hydrates thereof.

For these compounds of Formula Ia or Ib, in one embodiment R_1 is phenyl, naphthyl or thienyl optionally substituted by C_{6-10} aryl, C_{6-10} aryl C_{1-4} alkyl, C_{2-9} heteroaryl, C_{2-9} heteroaryl C_{1-4} alkyl,


10 C_{3-8} cycloalkyl, C_{3-8} cycloalkyl C_{1-4} alkyl, C_{3-8} heterocycloalkyl, C_{3-8} heterocycloalkyl C_{1-4} alkyl or C_{1-6} alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl group of R_1

can be optionally substituted by one to five radicals chosen from halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted- C_{1-6} alkyl and halo-substituted- C_{1-6} alkoxy; and any alkyl group of R_1 can

optionally have a methylene replaced by an atom or group chosen from $-S-$, $-S(O)-$, $-S(O)_2-$, $-NR_5-$ and $-O-$; wherein R_5 is hydrogen or C_{1-6} alkyl.

15


In another embodiment, Y is chosen from:



20 wherein R_7 is hydrogen or C_{1-6} alkyl; and the left and right asterisks of Y indicate the point of attachment a) either between $-C(R_2)=NOR_1$ and the $-CR_3R_4-$, or between $-CR_3R_4-$ and $-C(R_2)=NOR_1$ of Formula Ia, respectively, or b) either between $-CR_3R_4-$ and W or between W and $-CR_3R_4-$ of Formula Ib, respectively; wherein any aryl or heteroaryl of Y


can be optionally substituted with 1 to 3 radicals chosen from halo, hydroxy, nitro, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted C₁₋₆alkyl and halo-substituted C₁₋₆alkoxy.

In a further embodiment, R₁ is chosen from:



wherein the asterisk is the point of attachment of R₁ with W; R₈ is C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₂₋₉heteroaryl, C₂₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl, C₃₋₈heterocycloalkylC₁₋₄alkyl or C₁₋₆alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl group of R₈ can be optionally substituted by one to three radicals chosen from halo, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted-C₁₋₆alkyl and halo-substituted-C₁₋₆alkoxy; and any alkyl group of R₈ can optionally have a methylene replaced by an atom or group chosen from -S-, -S(O)-, -S(O)₂-, -NR₅- and -O-; wherein R₅ is hydrogen or C₁₋₆alkyl; and R₉ is chosen from halo, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted-C₁₋₆alkyl and halo-substituted-C₁₋₆alkoxy.


In another embodiment, A is -C(O)OH; and Z is chosen from:



wherein the left and right asterisks of Z indicate the point of attachment between -C(R₃)(R₄)- and A of Formula Ia or Ib, respectively; R₆ is chosen from hydrogen and C₁₋₆alkyl; and R₃ and R₄ are both hydrogen.

In a further embodiment, Y is chosen from phenyl, pyridinyl, thienyl and furanyl; wherein any phenyl, pyridinyl, thienyl or furanyl of Y is optionally substituted with 1 to 3 radicals chosen from methyl, ethyl, cyclopropyl, chloro, bromo, fluoro and methoxy; or where Y is phenyl, R₆ can be attached to a carbon atom of Y to form 3,4-dihydro-1H-isoquinolin-2-yl.

In another embodiment, W is a bond or methylene; R₁ is chosen from:



wherein R_8 is chosen from phenyl, cyclohexyl, thienyl, 3,3-dimethyl-butyl, pyridinyl, cyclopentyl and piperidinyl; wherein R_8 can be optionally substituted by 1 to 3 radicals
 5 chosen from trifluoromethyl, methoxy, fluoro, trifluoromethoxy and methyl; and R_9 is chosen from trifluoromethyl, fluoro, methyl, chloro, methoxy and ethyl.

Preferred compounds of the invention include:

3-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-
 {4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic
 10 acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-
 azetidine-3-carboxylic acid, 3-({2-Chloro-6-[1-(4-cyclohexyl-3-trifluoromethyl-
 benzyloxyimino)-ethyl]-pyridin-3-ylmethyl}-amino)-propionic acid, 3-({6-[1-(4-Cyclohexyl-3-
 trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-pyridin-3-ylmethyl}-amino)-propionic acid, 3-{4-
 [1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 4-{4-[1-(Biphenyl-4-
 15 ylmethoxyimino)-ethyl]-benzylamino}-butyric acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-
 ethyl]-benzyl}-azetidine-3-carboxylic acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-
 benzyl}-piperidine-3-carboxylic acid, {4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-
 acetic acid, 3-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-cyclopentanecarboxylic
 acid, 3-{4-[1-(4'-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic
 20 acid, 3-{4-[1-(5-Phenyl-furan-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-
 (3'-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-
 (3-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-
 (4'-Methoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-
 (Biphenyl-3-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-2-yl-
 25 benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-2-yl-3-
 trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Fluoro-
 biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Trifluoromethoxy-
 biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3'-Trifluoromethoxy-
 biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(2-Trifluoromethyl-
 30 biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-azetidine-3-carboxylic acid, 1-{4-[1-(2-
 Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-pyrrolidine-3-carboxylic acid, 1-{4-
 [1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-piperidine-3-carboxylic acid,

3-{4-[1-(3'-Methoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 2-Hydroxy-3-{4-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Methyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Phenyl-thiophen-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-pyrrolidine-3-carboxylic acid, 3-{4-[1-(4-Furan-3-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-3-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-3-yl-2-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 2-Fluoro-3-{4-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-butyric acid, 3-{4-[1-(5-Phenyl-thiophen-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3-Fluoro-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Fluoro-2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Methyl-2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Furan-2-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2'-Fluoro-2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-(4-{1-[4-(3,3-Dimethyl-butyl)-3-trifluoromethyl-benzyloxyimino]-ethyl}-benzylamino)-propionic acid, 3-{4-[1-(4-Furan-3-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Pyridin-3-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Pyridin-4-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2-Fluoro-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-({2-Methoxy-6-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-pyridin-3-ylmethyl}-amino)-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{2-Bromo-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclopentyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-({6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-3-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-thiophen-2-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-2-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-furan-2-ylmethyl}-amino)-propionic acid, 3-({2-[1-(4-

Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-4-ylmethyl}-amino)-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-fluoro-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 1-{6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-pyridin-3-ylmethyl}-azetidine-3-carboxylic acid, 3-{2-Ethyl-4-[1-(4-piperidin-1-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-methyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(3-Chloro-4-cyclohexyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-methoxy-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-methoxy-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-methyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-methyl-benzyl}-azetidine-3-carboxylic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-cyclopropyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-cyclopropyl-benzyl}-azetidine-3-carboxylic acid, 3-{2-Ethyl-4-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 1-{4-[1-(4-Cyclohexyl-3-methyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 1-{2-Chloro-4-[1-(4-cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-benzyl}-azetidine-3-carboxylic acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-fluoro-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-fluoro-benzyl}-azetidine-3-carboxylic acid, 3-{6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-3,4-dihydro-1H-isoquinolin-2-yl}-propionic acid, 3-{6-[1-(4-Cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-3,4-dihydro-1H-isoquinolin-2-yl}-propionic acid, 3-{4-[1-(2-Trifluoromethyl-biphenyl-4-yl)-ethylideneaminooxymethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid and 1-{4-[1-(4-Cyclohexyl-3-ethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid.

The invention provides forms of the compound that have the hydroxyl or amine group present in a protected form; these function as prodrugs. Prodrugs are compounds that are converted into an active drug form after administration, through one or more chemical or biochemical transformations. Forms of the compounds of the present invention that are readily converted into the claimed compound under physiological conditions are prodrugs of the claimed compounds and are within the scope of the present invention. Examples of prodrugs include forms where a hydroxyl group is acylated to form a relatively labile ester such as an acetate ester, and forms where an amine group is acylated with the carboxylate group of glycine or an L-amino acid such as serine, forming an amide bond that is particularly susceptible to hydrolysis by common metabolic enzymes.

When the compounds of Formula Ia or Ib have asymmetric centers in the molecule, various optical isomers are obtained. The present invention also encompasses enantiomers, racemates, diastereoisomers and mixtures thereof. Moreover, when the compounds of Formula Ia or Ib include geometric isomers, the present invention embraces cis-compounds, trans-compounds and mixtures thereof. Similar considerations apply in relation to starting materials exhibiting asymmetric carbon atoms or unsaturated bonds as mentioned above.

Compounds of Formula Ia or Ib can exist in free form or in salt form, e.g. addition salts with inorganic or organic acids. Where hydroxyl groups are present, these groups can also be present in salt form, e.g. an ammonium salt or salts with metals such as lithium, sodium, potassium, calcium, zinc or magnesium, or a mixture thereof. Compounds of Formula Ia or Ib and their salts in hydrate or solvate form are also part of the invention.

A preferred compound of formula Ia is e.g. 1-{4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid (Compound A), or a pharmaceutically acceptable salt or prodrug thereof.

Unless otherwise stated, alkyl, alkoxy, alkenyl or alkynyl may be straight or branched.

Halo or halogen means F, Cl, Br or I, preferably F or Cl. Halo-substituted alkyl groups and compounds can be partially halogenated or perhalogenated, whereby in the case of multiple halogenation, the halogen substituents can be identical or different. A preferred perhalogenated alkyl group is for example trifluoromethyl or trifluoromethoxy.

Any double bonds can be in the cis- or trans- configuration. "Alkynyl" as a group and as structural element of other groups and compounds contains at least one C \equiv C triple bond and can also contain one or more C=C double bonds, and can, so far as possible, be either

straight-chain or branched. Any cycloalkyl group, alone or as a structural element of other groups can contain from 3 to 8 carbon atoms, preferably from 3 to 6 carbon atoms. "Alkylene" and "alkenylene" are divalent radicals derived from "alkyl" and "alkenyl" groups, respectively. In this application, any alkyl group of R₁ can be optionally interrupted by a member of the group selected from -S-, -S(O)-, -S(O)₂-, -NR²⁰- and -O- (wherein R²⁰ is hydrogen or C₁₋₆alkyl). These groups include -CH₂-O-CH₂-, -CH₂-S(O)₂-CH₂-, -(CH₂)₂-NR²⁰-CH₂-, -CH₂-O-(CH₂)₂-, and the like.

"Aryl" means a monocyclic or fused bicyclic aromatic ring assembly containing six to ten ring carbon atoms. For example aryl, e.g. C₆₋₁₀aryl, can be phenyl, biphenyl or naphthyl, such as phenyl or naphthyl, preferably phenyl. A fused bicyclic ring can be partially saturated, for example, 1,2,3,4-tetrahydro-naphthalene, and the like.

"Heteroaryl" means aryl, as defined in this application, with the addition of at least one heteroatom moiety selected from N, O or S, and each ring is comprised of 5 to 6 ring atoms, unless otherwise stated. For example, C₂heteroaryl includes oxadiazole, triazole, and the like. C₉heteroaryl includes quinoline, 1,2,3,4-tetrahydro-quinoline, and the like. C₂₋₉heteroaryl as used in this application includes thienyl, pyridinyl, furanyl, isoxazolyl, benzoxazolyl or benzo[1,3]dioxolyl, preferably thienyl, furanyl or pyridinyl. A fused bicyclic heteroaryl ring system can be partially saturated, for example, 2,3-dihydro-1H-isoindole, 1,2,3,4-tetrahydro-quinoline, and the like.

According to the invention, inflammatory myopathies include, but are not limited to muscle inflammations, polymyositis, dermatomyositis, nerve-muscle diseases such as muscular dystrophies, and inclusion body myositis (IBM).

It has now been found that the S1P receptor modulators of the invention have an inhibitory effect on inflammatory myopathies, e.g. polymyositis.

In a series of further specific or alternative embodiments, the present invention provides:

1.1 A method for preventing, inhibiting or treating muscle inflammation or inflammatory myopathies, e.g. polymyositis, in a subject in need thereof, comprising administering to said subject a therapeutically effective amount of a S1P receptor modulator of the invention modulator, e.g. a compound of formulae Ia or Ib, e.g. Compound A.

1.2 A method for alleviating or delaying progression of the symptoms of an inflammatory myopathy, e.g. polymyositis, in a subject in need thereof, in which method muscle inflammation associated with said disease is prevented or inhibited, comprising

administering to said subject a therapeutically effective amount of a S1P receptor modulator of the invention, e.g. a compound of formulae Ia or Ib, e.g. Compound A.

1.3 A method for reducing or preventing or alleviating relapses in an inflammatory myopathy, e.g. polymyositis, in a subject in need thereof, in which method muscle inflammation associated with said disease is prevented or inhibited, comprising administering to said subject a therapeutically effective amount of a S1P receptor modulator of the invention, e.g. a compound of formulae Ia or Ib, e.g. Compound A.

1.4 A method for slowing progression of an inflammatory myopathy, e.g. polymyositis, in a subject being in a relapsing-remitting phase of the disease, in which method muscle inflammation associated with said disease is prevented or inhibited, comprising administering to said subject a therapeutically effective amount of a S1P receptor modulator of the invention modulator, e.g. a compound of formulae I or Ib, e.g. Compound A.

2. A pharmaceutical composition for use in any one of the methods 1.1 to 1.5, comprising a S1P receptor modulator of the invention, e.g. a compound of formulae Ia or Ib, e.g. Compound A, as defined hereinabove, together with one or more pharmaceutically acceptable diluents or carriers therefor.

3. A S1P receptor modulator of the invention, e.g. a compound of formula Ia or Ib, e.g. Compound A, as defined herein above, for use in any one of the methods 1.1 to 1.5.

4. A S1P receptor modulator of the invention, e.g. a compound of formula Ia or Ib, e.g. Compound A, as defined herein above, for use in the preparation of a medicament for use in any one of the methods 1.1 to 1.5.

5. Use of a S1P receptor modulator of the invention, e.g. a compound of formula Ia or Ib, e.g. Compound A as defined herein above, in any one of the methods 1.1 to 1.5 e.g. 1.1.

6. Use of a S1P receptor modulator of the invention, e.g. a compound of formulae Ia or Ib, e.g. Compound A as defined herein above, in the preparation of a medicament for use in a method according to one of the methods 1.1 to 1.5 e.g. 1.1.

7. A method, use, or pharmaceutical composition according to any one of the preceding paragraphs 1.1-1.5 and 2-6 wherein the S1P receptor modulator of the invention is 1-{4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-

benzyl}-azetidine-3-carboxylic acid in free form or in a pharmaceutically acceptable salt form.

Administration

5 Daily dosages required in practicing the method of the present invention when a S1P receptor modulator of the invention alone is used will vary depending upon, for example, the compound used, the host, the mode of administration and the severity of the condition to be treated. A preferred daily dosage range is about from 0.01 to 100 mg as a single dose or in divided doses. Suitable daily dosages for patients are on the order of from e.g. 0.03 to 2.5
10 mg/kg per body weight. The S1P receptor modulators of the invention may be administered by any conventional route, in particular enterally, e.g. orally, e.g. in the form of tablets, capsules, parenterally, e.g. in the form of injectable solutions or suspensions, or topically, e.g. in the form of lotions, gels, ointments or creams, or in a nasal or a suppository form. Suitable unit dosage forms for oral administration comprise from ca. 0.5 to about 100 mg,
15 e.g. 1 to 50 mg S1P receptor modulator of the invention, together with one or more pharmaceutically acceptable diluents or carriers therefore.

The compounds of Formula Ia or Ib may be administered as the sole active ingredient or in conjunction with, e.g. as an adjuvant to, other drugs e.g. immunosuppressive or immunomodulating agents or other anti-inflammatory agents. For example the compounds
20 of Formula Ia or Ib may be used in combination with a calcineurin inhibitor, e.g. cyclosporin A or FK 506; an ascomycin having immunosuppressive properties, e.g. ABT-281, ASM981, etc.; corticosteroids; cyclophosphamide; azathioprene; methotrexate; leflunomide; mizoribine; mycophenolic acid; mycophenolate mofetil; 15-deoxyspergualine or an immunosuppressive homologue, analogue or derivative thereof; immunosuppressive
25 monoclonal antibodies, e.g. monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD8, CD25, CD28, CD40, CD45, CD58, CD80, CD86 or their ligands; other immunomodulatory compounds, e.g. a recombinant binding molecule having at least a portion of the extracellular domain of CTLA4 or a mutant thereof, e.g. an at least extracellular portion of CTLA4 or a mutant thereof joined to a non-CTLA4 protein sequence,
30 e.g. CTLA4Ig (for ex. designated ATCC 68629) or a mutant thereof, e.g. LEA29Y ; adhesion molecule inhibitors, e.g. LFA-1 antagonists, ICAM-1 or -3 antagonists, VCAM-4 antagonists or VLA-4 antagonists.

Methods of preparing the compounds for use in the invention

The compounds for use in the aspects of the invention may, for example, be prepared by the methods specified in WO 2004/103306.

Examples

- 5 Utility of the S1P receptor modulators of the invention, e.g. 1-{4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid (Compound A), in preventing or treating an inflammatory myopathy, e.g. polymyositis, as hereinabove specified, may be demonstrated in vitro, in animal test methods as well as in clinic, for example in accordance with the methods hereinafter described.

In vitro: effect on cytokine-induced atrophy of primary human myotubes.

Human skeletal muscle (skMC) cells are obtained from Cambrex (#CC-2561). For experiments skMC stocks are thawed and maintained in SkBM (Lonza CC-3161) containing 20% FCS and 0.1% gentamycin at 37°C, 5% CO₂. After 4-5 days cells are seeded for experiments onto six-well plates coated with matrigel (450,000/well) and grown at 37°C, 5% CO₂ for one day. Cells are then washed 3x with SkBM and differentiated into myotubes with SkBM containing 1µM SB431542 (ALK4/5 inhibitor; Sigma #S4317) for 4 days (SB431542 is removed 24h prior cell treatment). Myotubes are then treated with test compound either in the absence or in the presence of a cytokine cocktail (TNFα 10 ng/ml, IL-1β 2 ng/ml, IFNγ 10 ng/ml) for 24h in SkBM plus 0.1% gentamycin, washed once with CSB buffer: 80mM pipes, 5nM EGTA, 1mM MgCl 6H₂O and 4% PEG35000 (Fluka #94646) and fixed with 4% paraformaldehyde (Electron Microscopy Sciences #15714) in CSB for 15 min at room temperature. Cells are then rinsed with CSB, permeabilized with 0.2% Triton X-100 (Merk #1.12298.0100) for 20 min at room temperature, rinsed with CSB and blocked with 10% normal goat serum blocking solution (Zymed Laboratories #50-062Z) for 20 min at room temperature. Primary antibody (anti-myosin heavy chain antibody; Upstate #05-716) diluted 1:500 in PBS containing 1.5% goat serum is added for 1h at room temperature. Myotubes are then washed 2x with CSB (5 min/wash) then add secondary antibody (Alexa Fluor 488 F(ab'); Invitrogen #A11017) diluted 1:750 in PBS is added for 1h at room temperature. Myotubes are washed once with CSB (10 min) then with PBS (Invitrogen # 14190) and double distilled water. Finally, ProLong Gold antifade reagent with DAPI (Invitrogen #P36931) is added and myotubes are photographed. Average diameters of at least 40 myotubes are measured for each condition at three points separated by 50 µm along the length of the myotube.

Human primary myoblasts, differentiated for 4 days and treated with cytokine cocktail for 24 h, fixed, and assayed for changes in myotube diameters show distinct atrophic phenotype with an approx. 20% decrease in myotube diameter as compared to vehicle control. Addition of Compound A at a concentration of 1nM is sufficient to block almost completely the cytokine-induced atrophy. Higher concentrations of Compound A exert the same effect. The results are shown in Figures 1(a) and 1(b).

In vivo: controllable muscle-specific promoter system to up-regulate MHC class I in the skeletal muscle of mice, as described e.g. in K. Nagaraju et al (PNAS, August 1, 2000, Vol 97, No. 16, p 9209-9214), the content thereof being included by reference.

Briefly, mice transgenic for both a transactivator (tTA) under the control of a muscle creatine kinase promoter (T^+) and the tetracycline-responsive element (TRE)-H-2K^b (H^+) are used. In H^+T^+ mice, the binding of a tetracycline analog to tTA prevents transactivator from binding to the TRE region, thereby preventing target gene expression. Thus, the transgenic H-2K^b expression can be induced by removing the tetracycline analog and suppressed by administering it. Mice having only the tetracycline-responsive element (TRE)-H-2Kb (H^+),
10 serve as control. H^+T^+ mice develop clinical, biochemical, histological, and immunological features similar to human myositis. First signs of disease are muscle weakness (about 3 months of age in females, i.e. 2 months after transgene induction).


Behavioral assays are used to measure developing muscle weakness (e.g. treadmill, RotaRod, open field locomotor activity, running wheel, grip test). Other readouts of the
15 model include blood markers indicative for muscle damage (CK, GOT etc.), and histopathology and immunohistochemistry of skeletal muscle (incl. mononuclear cell infiltrate, CD3⁺ T cell infiltrate, ICAM, etc.).

Application of compound in the therapeutic mode may start around 3 months of age in female mice (2 month after transgene induction), when first clinical signs of muscle
20 weakness occur and may last e.g. for 2 months. Compounds are applied i.p., s.c. or via sirup drops into the mouth. Oral gavage is not recommended due to dystrophy of pharyngeal muscles. Groups of n=8 mice are used. Compounds are applied to mice having both transgenes, H^+T^+ and mice having only the tetracycline-responsive element (TRE)-H-2Kb (H^+ , control). Vehicle-treated animals are compared to compound-treated animals.

25 Application of compound in the prophylactic mode starts from day of transgene induction. Both, therapeutic and prophylactic treatment mode can be combined with other therapeutics, like e.g. prednisone or comparable steroids.

CLAIMS

1. The use of an S1P receptor modulator of the formula Ia:



5


wherein

A is $-C(O)OR_5$, $-OP(O)(OR_5)_2$, $-P(O)(OR_5)_2$, $-S(O)_2OR_5$, $-P(O)(R_5)OR_5$ or 1H-tetrazol-5-yl, R_5 being H or C_{1-6} alkyl;

W is a bond, C_{1-3} alkylene or C_{2-3} alkenylene;

10 Y is C_{6-10} aryl or C_{2-9} heteroaryl e.g. C_{3-9} heteroaryl, optionally substituted by 1 to 3 radicals selected from halogen, $-OH$, $-NO_2$, C_{1-6} alkyl, C_{1-6} alkoxy; halo-substituted C_{1-6} alkyl and halo-substituted C_{1-6} alkoxy;

Z is chosen from:



15

wherein the left and right asterisks of Z indicate the point of attachment between $-C(R_3)(R_4)-$ and A of Formula Ia, respectively; R_6 is chosen from hydrogen and C_{1-6} alkyl; and

J₁ and J₂ are independently methylene or a heteroatom chosen from S, O and NR₅; wherein R₅ is chosen from hydrogen and C₁₋₆alkyl; and any alkylene of Z can be further substituted by one to three radicals chosen from halo, hydroxy, C₁₋₆alkyl; or R₆ can be attached to a carbon atom of Y to form a 5-7 member ring e.g. a heterocyclic group as indicated in WO
5 04/103306A, e.g. azetidine;

R₁ is C₆₋₁₀aryl or C₂₋₉heteroaryl e.g C₃₋₉heteroaryl, optionally substituted by C₁₋₆alkyl, C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₃₋₉heteroaryl, C₃₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl or C₃₋₈heterocycloalkylC₁₋₄alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R₁ may be substituted by 1 to 5 groups selected from
10 halogen, C₁₋₆alkyl, C₁₋₆alkoxy and halo substituted-C₁₋₆alkyl or -C₁₋₆alkoxy; and any alkyl group of R₁ can optionally have a methylene replaced by an atom or group chosen from -S-, -S(O)-, -S(O)₂-, -NR₅- and -O-; wherein R₅ is chosen from hydrogen or C₁₋₆alkyl;


R₂ is H, C₁₋₆alkyl, halo substituted C₁₋₆alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl; and each of R₃ or R₄, independently, is H, halogen, OH, C₁₋₆alkyl, C₁₋₆alkoxy or halo substituted
15 C₁₋₆alkyl or C₁₋₆alkoxy;

and the N-oxide derivatives thereof or prodrugs thereof, individual isomers and mixtures of isomers thereof; and the pharmacologically acceptable salts, solvates or hydrates thereof, in the preparation of a medicament for preventing, inhibiting or treating an inflammatory condition selected from polymyositis or dermatomyositis.

20

2. The use of claim 1, wherein, in the compound of formula Ia, R₁ is phenyl, naphthyl or thienyl optionally substituted by C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₂₋₉heteroaryl, C₂₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl, C₃₋₈heterocycloalkylC₁₋₄alkyl or C₁₋₆alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl group of R₁ can be
25 optionally substituted by one to five radicals chosen from halo, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted-C₁₋₆alkyl and halo-substituted-C₁₋₆alkoxy; and any alkyl group of R¹ can optionally have a methylene replaced by an atom or group chosen from -S-, -S(O)-, -S(O)₂-, -NR₅- and -O-; wherein R₅ is hydrogen or C₁₋₆alkyl.


30 3. The use of claim 1, wherein, in the compound of formula Ia, Y is chosen from:



wherein R₇ is hydrogen or C₁₋₆alkyl; and the left and right asterisks of Y indicate the point of attachment a) either between -C(R₂)=NOWR₁ and the -CR₃R₄-, or between -CR₃R₄- and -C(R₂)=NOWR₁ of Formula Ia, respectively; wherein any aryl or heteroaryl of Y can be optionally substituted with 1 to 3 radicals chosen from halo, hydroxy, nitro, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted C₁₋₆alkyl and halo-substituted C₁₋₆alkoxy.

4. The use of claim 1, wherein, in the compound of formula Ia, R₁ is chosen from:

10




wherein the asterisk is the point of attachment of R₁ with W; R₈ is C₆₋₁₀aryl, C₆₋₁₀arylC₁₋₄alkyl, C₂₋₉heteroaryl, C₂₋₉heteroarylC₁₋₄alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₄alkyl, C₃₋₈heterocycloalkyl, C₃₋₈heterocycloalkylC₁₋₄alkyl or C₁₋₆alkyl; wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl group of R₈ can be optionally substituted by one to three radicals chosen from halo, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted-C₁₋₆alkyl and halo-substituted-C₁₋₆alkoxy; and any alkyl group of R₈ can optionally have a methylene replaced by an atom or group chosen from -S-, -S(O)-, -S(O)₂-, -NR₅- and -O-; wherein R₅ is hydrogen or C₁₋₆alkyl; and R₉ is chosen from halo, C₁₋₆alkyl, C₁₋₆alkoxy, halo-substituted-C₁₋₆alkyl and halo-substituted-C₁₋₆alkoxy.

15

20

5. The use of claim 1, wherein, in the compound of formula Ia, A is $-C(O)OH$; Z is chosen from:



5


wherein the left and right asterisks of Z indicate the point of attachment between $-C(R_3)(R_4)-$ and A of Formula Ia, respectively; R_6 is chosen from hydrogen and C_{1-6} alkyl; and R_3 and R_4 are both hydrogen.

10

6. The use of claim 5, wherein, in the compound of formula Ia, Y is chosen from phenyl, pyridinyl, thienyl and furanyl; wherein any phenyl, pyridinyl, thienyl or furanyl of Y is optionally substituted with 1 to 3 radicals chosen from methyl, ethyl, cyclopropyl, chloro, bromo, fluoro and methoxy; or where Y is phenyl, R_6 can be attached to a carbon atom of Y to form 3,4-dihydro-1H-isoquinolin-2-yl.

15

7. The use of claim 6, wherein, in the compound of formula Ia, W is a bond or methylene; R_1 is chosen from:



20

wherein R_8 is chosen from phenyl, cyclohexyl, thienyl, 3,3-dimethyl-butyl, pyridinyl, cyclopentyl and piperidinyl; wherein R_8 can be optionally substituted by 1 to 3 radicals chosen from trifluoromethyl, methoxy, fluoro, trifluoromethoxy and methyl; and R_9 is chosen from trifluoromethyl, fluoro, methyl, chloro, methoxy and ethyl.

25

8. The use of claim 6, wherein the compound of formula Ia is selected from: 3-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 3-({2-Chloro-6-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-

- pyridin-3-ylmethyl)-amino)-propionic acid, 3-({6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-pyridin-3-ylmethyl)-amino)-propionic acid, 3-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 4-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-butyric acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-
- 5 azetidine-3-carboxylic acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-piperidine-3-carboxylic acid, {4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-acetic acid, 3-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-cyclopentanecarboxylic acid, 3-{4-[1-(4'-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(5-Phenyl-furan-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3'-
- 10 Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Methoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(Biphenyl-3-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-2-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-2-yl-3-
- 15 trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Fluoro-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Trifluoromethoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3'-Trifluoromethoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-azetidine-3-carboxylic acid, 1-{4-[1-(2-
- 20 Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-pyrrolidine-3-carboxylic acid, 1-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-piperidine-3-carboxylic acid, 3-{4-[1-(3'-Methoxy-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 2-Hydroxy-3-{4-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Methyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-
- 25 [1-(4-Phenyl-thiophen-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(Biphenyl-4-ylmethoxyimino)-ethyl]-benzyl}-pyrrolidine-3-carboxylic acid, 3-{4-[1-(4-Furan-3-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-3-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Thiophen-3-yl-2-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 2-Fluoro-3-{4-[1-(2-
- 30 trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2-Trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-butyric acid, 3-{4-[1-(5-Phenyl-thiophen-2-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(3-Fluoro-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Fluoro-2-trifluoromethyl-

biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4'-Methyl-2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Furan-2-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2'-Fluoro-2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-(4-
5 {1-[4-(3,3-Dimethyl-butyl)-3-trifluoromethyl-benzyloxyimino]-ethyl]-benzylamino)-propionic acid, 3-{4-[1-(4-Furan-3-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Pyridin-3-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Pyridin-4-yl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(2-Fluoro-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 3-({2-Methoxy-6-[1-(2-trifluoromethyl-
10 biphenyl-4-ylmethoxyimino)-ethyl]-pyridin-3-ylmethyl}-amino)-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{2-Bromo-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclopentyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic
15 acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-({6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-3-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-thiophen-2-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-2-ylmethyl}-amino)-propionic acid, 3-({5-[1-(4-Cyclohexyl-3-
20 trifluoromethyl-benzyloxyimino)-ethyl]-furan-2-ylmethyl}-amino)-propionic acid, 3-({2-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-pyridin-4-ylmethyl}-amino)-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-fluoro-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 1-{6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-pyridin-3-
25 ylmethyl}-azetidine-3-carboxylic acid, 3-{2-Ethyl-4-[1-(4-piperidin-1-yl-3-trifluoromethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-methyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(3-Chloro-4-cyclohexyl-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-methoxy-benzyloxyimino)-ethyl]-2-ethyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-methoxy-
30 benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-methyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-methyl-benzyl}-azetidine-3-carboxylic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-cyclopropyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-

2-cyclopropyl-benzyl)-azetidine-3-carboxylic acid, 3-{2-Ethyl-4-[1-(2-trifluoromethyl-biphenyl-4-ylmethoxyimino)-ethyl]-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 1-{4-[1-(4-Cyclohexyl-3-methyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid, 1-{2-Chloro-4-[1-(4-cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-benzyl}-azetidine-3-carboxylic acid, 3-{2-Chloro-4-[1-(4-cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-fluoro-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-fluoro-benzyl}-azetidine-3-carboxylic acid, 3-{6-[1-(4-Cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-3,4-dihydro-1H-isoquinolin-2-yl}-propionic acid, 3-{6-[1-(4-Cyclohexyl-3-ethyl-benzyloxyimino)-ethyl]-3,4-dihydro-1H-isoquinolin-2-yl}-propionic acid, 3-{4-[1-(2-Trifluoromethyl-biphenyl-4-yl)-ethylideneaminooxymethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-benzylamino}-propionic acid, 3-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzylamino}-propionic acid, 1-{4-[1-(4-Cyclohexyl-3-trifluoromethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid and 1-{4-[1-(4-Cyclohexyl-3-ethyl-phenyl)-ethylideneaminooxymethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid.

9. The use of claim 6, wherein the compound of formula Ia is 1-{4-[1-(4-cyclohexyl-3-trifluoromethyl-benzyloxyimino)-ethyl]-2-ethyl-benzyl}-azetidine-3-carboxylic acid (Compound A), or a pharmaceutically acceptable salt or prodrug thereof.

10. An S1P receptor modulator as defined in any one of claims 1 to 9 for use in preventing, inhibiting or treating an inflammatory condition selected from polymyositis or dermatomyositis.

11. A method for preventing, inhibiting or treating an inflammatory condition selected from polymyositis or dermatomyositis, in a subject in need thereof, comprising administering to said subject a therapeutically effective amount of an S1P receptor modulator as defined in any one of claims 1 to 9.

12. A pharmaceutical composition comprising an S1P receptor modulator as defined in any one of claims 1 to 9 for use in preventing, inhibiting or treating an inflammatory condition selected from polymyositis or dermatomyositis together with one or more pharmaceutically acceptable diluents or carriers therefor.

5

13. A method, use, pharmaceutical composition or S1P receptor modulator for use substantially as hereinbefore defined and described.

Figures




Figure 1(a)

5






Figure 1(b)

10



(1a)



(1b)