

(19) **DANMARK**

(10) **DK/EP 3473670 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 08 K 5/34 (2006.01)** **C 08 J 5/18 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-07-13**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-04-15**
- (86) Europæisk ansøgning nr.: **18201146.0**
- (86) Europæisk indleveringsdag: **2016-02-16**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-04-24**
- (30) Prioritet: **2015-02-20 EP 15155952**
- (62) Stamansøgningsnr: **16704632.5**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **BASF SE, Carl-Bosch-Strasse 38, 67056 Ludwigshafen am Rhein, Tyskland**
- (72) Opfinder: **Mueller, Daniel, Hardmattstr. 434, 5082 Kaisten, Schweiz**
Weyland, Tania, Hardmattstr. 434, 5082 Kaisten, Schweiz
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **LYSSTABILISEREDE POLYOLEFINFILM, -BÅND OG -MONOFILAMENTER**
- (56) Fremdragne publikationer:
US-A- 5 006 587

DESCRIPTION

[0001] The present invention relates to a process for the reduction of water carry-over of a light stabilized polyolefin film, tape or monofilament which is passed through a water bath during production, to the polyolefin film, tape or monofilament made in accordance to that process and to the use of specific light stabilizers to reduce the water carry-over of a light stabilized polyolefin film, tape or monofilament.

[0002] A widespread method for producing polyolefin films, tapes or monofilaments consists in extruding the polymer melt through a suitable apparatus and, in the form of a film, tape or monofilament, into a water bath, where the film, tape or monofilament is cooled and solidified. The film, tape or monofilament may then in turn be passed out of the water bath and subjected to further steps of processing. On emerging from the water bath the film, tape or monofilament can entrain water, which interferes with subsequent steps of processing. This effect, the entrainment of water from the water bath, is often termed "water carry-over" in the technical literature, and this can be abbreviated to WCO.

[0003] Stretching of the film at a suitable temperature results in orientation and further crystallization of the polymer, resulting in the specific properties. This second step of processing may take place directly on the film, but the primary film is often split into tapes prior to the stretching process. Since the desired properties are obtained during the stretching process, precise adherence to all process parameters is essential here. Even traces of moisture on the film or on the tapes prior to the stretching process alter the subsequent stretching and orientation so as to produce very severe variations in the quality of the product obtained. The effects of this go beyond merely the corresponding major variations in the quality of the resultant final product. Even during production or processing, the poor quality can cause break-offs of the tapes, for example, and thus stop production. A source of water which can lead to the problems mentioned is the cooling bath into which the primary film is extruded. Although most plastics, in particular the polyolefins, such as polyethylene (PE) or polypropylene (PP) are very hydrophobic and therefore have little tendency to absorb water, it is frequently found that at relatively high production speeds water droplets or a thin film of water adhere to the film when this is drawn over the water bath. In addition, various additives such as light stabilizers have to be added to the polymer to ensure that the final product has good functionality. Some of these additives contribute to increased entrainment of water from the water bath. The additives, and also a maximum processing speed, are essential if suitable products are to be produced at low cost. A reduction in processing speed leads to uneconomic production. The addition of necessary additives, some of which lead to a worsening of WCO (i.e. more WCO and therefore poorer product) is also impossible to avoid, because otherwise the product properties brought about by the additives cannot be achieved.


[0004] The reduction of WCO is currently promoted via simple design measures during machinery manufacture. For example, the draw-off of the film from the water bath is usually vertically upward, so that gravity alone maximizes the amount of water dropping away. In

addition, use is often made of squeeze rollers and/or air knives to remove the maximum amount of water from the film. There are other technical methods, depending on the specific configuration of the machine. US 5 006 587 A discloses a polyolefin comprising a hindered amine light stabilizer, which in the examples is Chimasorb 944FL and as such corresponding to component (A) of the present invention, together with fluoborate salt in order to reduce water carry-over by the polyolefin film.

[0005] There continues to be a requirement for a process for improving WCO during the production of polyolefin films, tapes and monofilaments. Ideally, no undesirable side-effects should arise, and the process should also be capable of problem-free use for existing formulations.

[0006] Surprisingly, it has now been found that the addition of a specific second sterically hindered amine light stabilizer to a light stabilized polyolefin film, tape or monofilament which contain already a sterically hindered amine light stabilizer can markedly reduce WCO during the production. This is unexpected particularly because the two sterically hindered amines belong to the same class of compounds. It was completely surprising and in no way predictable that particular combinations of two sterically hindered amine light stabilizers show a synergistic improvement in WCO properties.


[0007] The invention therefore provides a process for reducing water carry-over of a light stabilized polyolefin film (preferably monolayer), tape (preferably monolayer) or monofilament which contains component (A) as light stabilizer and which is passed through a water bath during production, characterized in that the polyolefin film, tape or monofilament additionally comprises component (B) as light stabilizer, component (A) is at least one compound selected from the group consisting of compounds of the formula (A-I), compounds of the formula (A-II) and compounds of the formula (A-III),




wherein A_1 is hydrogen or C_1 - C_4 alkyl,

A_2 is a direct bond or C_1 - C_{10} alkylene, and

a_1 is a number from 2 to 20;






wherein A_3 , A_4 and A_5 independently of one another are C_2 - C_{18} alkylene,

A_6 , A_7 , A_8 , A_9 and A_{10} independently of one another are hydrogen, C_1 - C_{12} alkyl, C_5 - C_{12} cycloalkyl or a group of the formula (a-1),



the radicals A_{11} independently of one another are hydrogen, C_1 - C_{12} alkyl or C_5 - C_{12} cycloalkyl, and

a_2 is a number from 1 to 20;



wherein A_{12} is C_2 - C_{18} alkylene, C_5 - C_7 cycloalkylene or C_1 - C_4 alkylenedi(C_5 - C_7 cycloalkylene),

A_{13} and A_{14} independently of one another are hydrogen, C_1 - C_{12} alkyl, C_5 - C_{12} cycloalkyl or a group of the formula (a-2),




or the radicals A_{13} and A_{14} , together with the nitrogen atom to which they are bonded, form a 5- to 10-membered heterocyclic ring,

the radicals A_{15} independently of one another are hydrogen, C_1 - C_{12} alkyl or C_5 - C_{12} cycloalkyl, and


a_3 is a number from 2 to 20;

component (B) is at least one compound selected from the group consisting of compounds of


the formula (B-I), compounds of the formula (B-II), compounds of the formula (B-III) and compounds of the formula (B-IV),



wherein R₁ is C₁-C₁₈alkyl, C₁-C₁₈hydroxyalkyl, cyclohexyl or hydroxycyclohexyl or R₁ is a group -C(C₆H₅)(H)CH₂-OH and R₂ is C₁-C₂₅alkyl or a group of the formula (b-1);




wherein x is an integer from 2 to 8 and y is zero or 1;



wherein R₃ is C₂-C₁₈alkylene, C₅-C₇cycloalkylene or C₁-C₄alkylenedi(C₅-C₇cycloalkylene),


the radicals R₄ independently of one another are C₁-C₁₂alkyl or C₅-C₁₂cycloalkyl,

R₅ and R₆ independently of one another are hydrogen, C₁-C₁₂alkyl, C₅-C₁₂cycloalkyl or a group of the formula (b-2),





or the radicals R₅ and R₆, together with the nitrogen atom to which they are bonded, form a 5- to 10-membered heterocyclic ring and

b₁ is a number from 1 to 20;




wherein the radicals R₈, R₉ and R₁₀ independently of one another are C₂-C₁₈alkylene and the radicals R₇ independently of one another are a group of the formula (b-3)






wherein R_{11} and R_{12} independently of one another are hydrogen, C_1 - C_{12} alkyl, C_5 - C_{12} cycloalkyl or a group of the formula (b-4)



and the radicals R_{13} independently of one another are C_1 - C_{12} alkyl or C_5 - C_{12} cycloalkyl;



wherein

R_5 is C_1 - C_{25} alkyl,

n is a number from 1 to 10, and

W is a wax residue comprising between 50 and 1000 carbon atoms, preferably a residue of a polyethylene or polypropylene wax.

[0008] Components (A) and (B) are preferably in one layer, if the film is a multilayer film.

[0009] The compounds of components (A) and (B) are known and largely commercially available and can be prepared according to known methods.

[0010] The compounds of component (A) can be prepared for example in analogy to the methods described in US-A-4,233,412, US-A-4,477,615 (CAS 136,504-96-6) and US-A-4,108,829, US-A-4,086,204, US-A-4,331,586, US-A-6,046,304 and US-A-6,297,299.

[0011] The compounds of component (B) can be prepared for example in analogy to the methods described in US-A-6,388,072, US-A-6,117,995, US-A-6,420,462 and US-A-6,677,451, and US-A-5,844,026.

[0012] Preferred compounds of component (A) are the commercial products Tinuvin®622, Sabostab® UV119, Uvasorb®HA88, Uvasorb®HA10, Chimassorb®944, Chimassorb®2020, Cyasorb®UV 3346, Cyasorb®UV 3529 and Dastib®1082.


[0013] Preferred compounds of component (B) are the commercial products Tinuvin®NOR 371, Tinuvin®123, ADK STAB®LA81 or Flamestab®NOR 116.

[0014] The polyolefin is preferably a polyethylene or a polypropylene and contains for example 0.01 to 10%, in particular 0.1 to 1%, relative to the weight of the polyolefin, of the sum of components (A) and (B).


[0015] The weight ratio of component (A) to component (B) is for example 1:20 to 20:1, preferably 1:15 to 15:1.

[0016] The meanings of the terminal groups which saturate the free valences in the compounds of the formulae (A-I), (A-II), (A-III) and (B-II) depend on the processes used for their preparation. The terminal groups can also be modified after the preparation of the compounds.


[0017] If the compounds of the formula (A-I) are prepared, for example, by reacting a compound of the formula




in which A_1 is hydrogen or methyl, with a dicarboxylic acid diester of the formula $Y_0\text{-OOC-A}_2\text{-COO-Y}_0$, in which Y_0 is, for example, methyl, ethyl or propyl, and A_2 is as defined above, the terminal group bonded to the 2,2,6,6-tetramethyl-4-oxypiperidin-1-yl radical is hydrogen or -CO-A₂-COO-Y₀, and the terminal group bonded to the diacyl radical is -O-Y₀ or




[0018] In the compounds of the formula (A-II) the terminal groups bonded to the triazine radical is, for example, Cl or a group



and the terminal group bonded to the amino radical is, for example, hydrogen or a group







and a_3 is a number from 2 to 10.

[0022] The preparation of this compound is described in Example 10 of US-A-6,046,304.

[0023] In the compounds of the formula (B-II), the end group bonded to the diamino radical is for example hydrogen or



and the terminal group bonded to the triazine radical is for example



[0024] Examples of alkyl having up to 25 carbon atoms are methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 2-ethylbutyl, n-pentyl, isopentyl, 1-methyl-pentyl, 1,3-dimethyl-butyl, n-hexyl, 1-methyl-hexyl, n-heptyl, isoheptyl, 1,1,3,3-tetra-methyl-butyl, 1-methyl-heptyl, 3-methyl-heptyl, n-octyl, 2-ethyl-hexyl, 1,1,3-tri-methyl-hexyl, 1,1,3,3-tetra-methyl-pentyl, nonyl, decyl, undecyl, 1-methyl-undecyl, dodecyl, 1,1,3,3,5,5-hexa-methyl-hexyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl and octadecyl.

[0025] Examples of C_5 - C_{12} cycloalkyl are cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and cyclododecyl. Cyclohexyl is preferred.

[0026] Examples of alkylene having up to 18 carbon atoms are methylene, ethylene, propylene, trimethylene, tetramethylene, pentamethylene, 2,2-dimethyltrimethylene, hexamethylene, decamethylenen and octadecamethylene.

[0027] A preferred example of C_5 - C_7 cycloalkylene is cyclohexylene.

[0028] A preferred example of C_1 - C_4 alkylenedi(C_5 - C_7 cycloalkylene) is cyclohexylene-methylene-cyclohexylene.

[0029] A preferred example of a 5-to 7-membered heterocyclic ring is a morpholine group.

[0030] In a preferred process A_1 is hydrogen,

A_2 is C_2 - C_6 alkylene, and

a_1 is a number from 2 to 10;

A_3 , A_4 and A_5 independently of one another are C_2 - C_6 alkylene,

A_6 , A_7 , A_8 , A_9 and A_{10} independently of one another are C_1 - C_4 alkyl,

the radicals A_{11} independently of one another are hydrogen or C_1 - C_4 alkyl, and

a_2 is a number from 1 to 10;

A_{12} is C_2 - C_6 alkylene,

A_{13} and A_{14} independently of one another are hydrogen, C_1 - C_4 alkyl, cyclohexyl or a group of the formula (a-2),

or the radicals A_{13} and A_{14} , together with the nitrogen atom to which they are bonded, form a morpholino group,

the radicals A_{15} independently of one another are hydrogen or C_1 - C_4 alkyl, and a_3 is a number from 2 to 10,

R_1 is C_1 - C_{11} alkyl or C_2 - C_6 hydroxyalkyl, R_2 is C_{10} - C_{20} alkyl or a group of the formula (b-1); R_3 is C_2 - C_6 alkylene,

the radicals R_4 independently of one another are C_1 - C_4 alkyl or cyclohexyl;

R_5 and R_6 independently of one another are hydrogen, C_1 - C_4 alkyl, cyclohexyl or a group of the formula (b-2),


or the radicals R_5 and R_6 , together with the nitrogen atom to which they are bonded, form a morpholino group and

b_1 is a number from 1 to 10;

R_8 , R_9 and R_{10} independently of one another are C_2 - C_6 alkylene and the radicals R_7 independently of one another are a group of the formula (b-3),


R_{11} and R_{12} independently of one another are C_1 - C_4 alkyl, and the radicals R_{13} independently of one another are C_1 - C_4 alkyl or cyclohexyl.

[0031] According to a preferred embodiment the compound of the formula (A-I) corresponds to the formula (A-I-1)




wherein a_1 is a number from 2 to 10,

the compound of the formula (A-II) corresponds to the formula (A-II-1) or (A-II-2),




wherein a_2 is a number from 1 to 10,




wherein a_2 is a number from 1 to 10,


the compound of the formula (A-III) corresponds to the formula (A-III-1), (A-III-2), (A-III-3) or (A-III-4),




wherein a_3 is a number from 2 to 10,



wherein a_3 is a number from 2 to 10,






wherein A_{15} is hydrogen or methyl and a_3 is a number from 2 to 10,






wherein a_3 is a number from 2 to 10.

the compound of the formula (B-I) corresponds to the formula (B-I-1), (B-I-2) or (B-I-3),





the compound of the formula (B-II) corresponds to the formula (B-II-1),







wherein a_3 is a number from 2 to 10,



wherein b_1 is a number from 1 to 10,



wherein R_7 is a group of the formula (b-3-1).



[0033] The polyolefin may additionally contain one or more conventional additives. Suitable examples are listed below.

1. Antioxidants

[0034] 1.1. Alkylated monophenols, for example 2,6-di-tert-butyl-4-methylphenol, 2-tert-butyl-4,6-dimethyl phenol, 2,6-di-tert-butyl-4-ethylphenol, 2,6-di-tert-butyl-4-n-butylphenol, 2,6-di-tert-butyl-4-isobutylphenol, 2,6-dicyclopentyl-4-methylphenol, 2-(α -methylcyclohexyl)-4,6-dimethylphenol, 2,6-dioctadecyl-4-methylphenol, 2,4,6-tricyclohexylphenol, 2,6-di-tert-butyl-4-

methoxymethylphenol, nonylphenols which are linear or branched in the side chains, for example, 2,6-di-nonyl-4-methylphenol, 2,4-dimethyl-6-(1'-methylundec-1'-yl)phenol, 2,4-dimethyl-6-(1'-methylheptadec-1'-yl)phenol, 2,4-dimethyl-6-(1'-methyltridec-1'-yl)phenol and mixtures thereof.

[0035] 1.2. Alkylthiomethylphenols, for example 2,4-dioctylthiomethyl-6-tert-butylphenol, 2,4-dioctylthiomethyl-6-methylphenol, 2,4-dioctylthiomethyl-6-ethylphenol, 2,6-didodecylthiomethyl-4-nonylphenol.

[0036] 1.3. Hydroquinones and alkylated hydroquinones, for example 2,6-di-tert-butyl-4-methoxyphenol, 2,5-di-tert-butylhydroquinone, 2,5-di-tert-amylhydroquinone, 2,6-diphenyl-4-octadecyloxyphenol, 2,6-di-tert-butylhydroquinone, 2,5-di-tert-butyl-4-hydroxyanisole, 3,5-di-tert-butyl-4-hydroxyanisole, 3,5-di-tert-butyl-4-hydroxyphenyl stearate, bis(3,5-di-tert-butyl-4-hydroxyphenyl) adipate.

[0037] 1.4. Tocopherols, for example α -tocopherol, β -tocopherol, γ -tocopherol, δ -tocopherol and mixtures thereof (vitamin E).

[0038] 1.5. Hydroxylated thiodiphenyl ethers, for example 2,2'-thiobis(6-tert-butyl-4-methylphenol), 2,2'-thiobis(4-octylphenol), 4,4'-thiobis(6-tert-butyl-3-methylphenol), 4,4'-thiobis(6-tert-butyl-2-methylphenol), 4,4'-thiobis(3,6-di-sec-amylphenol), 4,4'-bis(2,6-dimethyl-4-hydroxyphenyl)disulfide.

[0039] 1.6. Alkylidenebisphenols, for example 2,2'-methylenebis(6-tert-butyl-4-methylphenol), 2,2'-methylenebis(6-tert-butyl-4-ethylphenol), 2,2'-methylenebis[4-methyl-6-(α -methylcyclohexyl)phenol], 2,2'-methylenebis(4-methyl-6-cyclohexylphenol), 2,2'-methylenebis(6-nonyl-4-methylphenol), 2,2'-methylenebis(4,6-di-tert-butyl phenol), 2,2'-ethylidenebis(4,6-di-tert-butylphenol), 2,2'-ethylidenebis(6-tert-butyl-4-isobutylphenol), 2,2'-methylenebis[6-(α -methylbenzyl)-4-nonyl phenol], 2,2'-methylenebis[6-(α , α -dimethylbenzyl)-4-nonylphenol], 4,4'-methylenebis(2,6-di-tert-butylphenol), 4,4'-methylenebis(6-tert-butyl-2-methylphenol), 1,1-bis(5-tert-butyl-4-hydroxy-2-methylphenyl)butane, 2,6-bis(3-tert-butyl-5-methyl-2-hydroxybenzyl)-4-methylphenol, 1,1,3-tris(5-tert-butyl-4-hydroxy-2-methylphenyl)butane, 1,1-bis(5-tert-butyl-4-hydroxy-2-methyl-phenyl)-3-n-dodecylmercaptobutane, ethylene glycol bis[3,3-bis(3'-tert-butyl-4'-hydroxyphenyl)butyrate], bis(3-tert-butyl-4-hydroxy-5-methyl-phenyl)dicyclopentadiene, bis[2-(3'-tert-butyl-2'-hydroxy-5'-methylbenzyl)-6-tert-butyl-4-methylphenyl]terephthalate, 1,1-bis-(3,5-dimethyl-2-hydroxyphenyl)butane, 2,2-bis(3,5-di-tert-butyl-4-hydroxyphenyl)propane, 2,2-bis(5-tert-butyl-4-hydroxy-2-methylphenyl)-4-n-dodecylmercaptobutane, 1,1,5,5-tetra-(5-tert-butyl-4-hydroxy-2-methylphenyl)pentane.

[0040] 1.7. O-, N- and S-benzyl compounds, for example 3,5,3',5'-tetra-tert-butyl-4,4'-dihydroxydibenzyl ether, octadecyl-4-hydroxy-3,5-dimethylbenzylmercaptoacetate, tridecyl-4-hydroxy-3,5-di-tert-butylbenzylmercaptoacetate, tris(3,5-di-tert-butyl-4-hydroxybenzyl)amine, bis(4-tert-butyl-3-hydroxy-2,6-dimethylbenzyl)dithioterephthalate, bis(3,5-di-tert-butyl-4-

hydroxybenzyl)sulfide, isooctyl-3,5-di-tert-butyl-4-hydroxybenzylmercaptoacetate.

[0041] 1.8. Hydroxybenzylated malonates, for example dioctadecyl-2,2-bis(3,5-di-tert-butyl-4-hydroxybenzyl)malonate, di-octadecyl-2-(3-tert-butyl-4-hydroxy-5-methylbenzyl)malonate, di-dodecylmercaptoethyl-2,2-bis (3,5-di-tert-butyl-4-hydroxybenzyl)malonate, bis[4-(1,1,3,3-tetramethylbutyl)phenyl]-2,2-bis(3,5-di-tert-butyl-4-hydroxybenzyl)malonate.

[0042] 1.9. Aromatic hydroxybenzyl compounds, for example 1,3,5-tris(3,5-di-tert-butyl-4-hydroxybenzyl)-2,4,6-trimethylbenzene, 1,4-bis(3,5-di-tert-butyl-4-hydroxybenzyl)-2,3,5,6-tetramethylbenzene, 2,4,6-tris(3,5-di-tert-butyl-4-hydroxybenzyl)phenol.

[0043] 1.10. Triazine compounds, for example 2,4-bis(octylmercapto)-6-(3,5-di-tert-butyl-4-hydroxyanilino)-1,3,5-triazine, 2-octylmercapto-4,6-bis(3,5-di-tert-butyl-4-hydroxyanilino)-1,3,5-triazine, 2-octylmercapto-4,6-bis(3,5-di-tert-butyl-4-hydroxyphenoxy)-1,3,5-triazine, 2,4,6-tris(3,5-di-tert-butyl-4-hydroxyphenoxy)-1,2,3-triazine, 1,3,5-tris(3,5-di-tert-butyl-4-hydroxybenzyl)isocyanurate, 1,3,5-tris(4-tert-butyl-3-hydroxy-2,6-dimethylbenzyl)isocyanurate, 2,4,6-tris(3,5-di-tert-butyl-4-hydroxyphenylethyl)-1,3,5-triazine, 1,3,5-tris(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)-hexahydro-1,3,5-triazine, 1,3,5-tris(3,5-dicyclohexyl-4-hydroxybenzyl)isocyanurate.

[0044] 1.11. Benzylphosphonates, for example dimethyl-2,5-di-tert-butyl-4-hydroxybenzylphosphonate, diethyl-3,5-di-tert-butyl-4-hydroxybenzyl phosphonate, dioctadecyl-3,5-di-tert-butyl-4-hydroxybenzylphosphonate, dioctadecyl-5-tert-butyl-4-hydroxy-3-methylbenzylphosphonate, the calcium salt of the monoethyl ester of 3,5-di-tert-butyl-4-hydroxybenzylphosphonic acid.

[0045] 1.12. Acylaminophenols, for example 4-hydroxylauranilide, 4-hydroxystearanilide, octyl N-(3,5-di-tert-butyl-4-hydroxyphenyl)carbamate.

[0046] 1.13. Esters of β -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid with mono- or polyhydric alcohols, e.g. with methanol, ethanol, n-octanol, i-octanol, octadecanol, 1,6-hexanediol, 1,9-nonanediol, ethylene glycol, 1,2-propanediol, neopentyl glycol, thiodiethylene glycol, diethylene glycol, triethylene glycol, pentaerythritol, tris(hydroxyethyl)isocyanurate, N,N'-bis(hydroxyethyl)oxamide, 3-thiaundecanol, 3-thiapentadecanol, trimethylhexanediol, trimethylolpropane, 4-hydroxymethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane.

[0047] 1.14. Esters of β -(5-tert-butyl-4-hydroxy-3-methylphenyl)propionic acid with mono- or polyhydric alcohols, e.g. with methanol, ethanol, n-octanol, i-octanol, octadecanol, 1,6-hexanediol, 1,9-nonanediol, ethylene glycol, 1,2-propanediol, neopentyl glycol, thiodiethylene glycol, diethylene glycol, triethylene glycol, pentaerythritol, tris(hydroxyethyl)isocyanurate, N,N'-bis(hydroxyethyl)oxamide, 3-thiaundecanol, 3-thiapentadecanol, trimethylhexanediol, trimethylolpropane, 4-hydroxymethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane; 3,9-bis[2-{3-(3-tert-butyl-4-hydroxy-5-methylphenyl)propionyloxy}-1,1-dimethylethyl]-2,4,8,10-tetraoxaspiro[5.5]undecane.

[0048] 1.15. Esters of β -(3,5-dicyclohexyl-4-hydroxyphenyl)propionic acid with mono- or polyhydric alcohols, e.g. with methanol, ethanol, octanol, octadecanol, 1,6-hexanediol, 1,9-nonanediol, ethylene glycol, 1,2-propanediol, neopentyl glycol, thiodiethylene glycol, diethylene glycol, triethylene glycol, pentaerythritol, tris(hydroxyethyl)isocyanurate, N,N'-bis(hydroxyethyl)oxamide, 3-thiaundecanol, 3-thiapentadecanol, trimethylhexanediol, trimethylolpropane, 4-hydroxymethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane.

[0049] 1.16. Esters of 3,5-di-tert-butyl-4-hydroxyphenyl acetic acid with mono- or polyhydric alcohols, e.g. with methanol, ethanol, octanol, octadecanol, 1,6-hexanediol, 1,9-nonanediol, ethylene glycol, 1,2-propanediol, neopentyl glycol, thiodiethylene glycol, diethylene glycol, triethylene glycol, pentaerythritol, tris(hydroxyethyl)isocyanurate, N,N'-bis(hydroxyethyl)oxamide, 3-thiaundecanol, 3-thiapentadecanol, trimethylhexanediol, trimethylolpropane, 4-hydroxymethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane.

[0050] 1.17. Amides of β -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid e.g. N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hexamethylenediamide, N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)trimethylenediamide, N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hydrazide, N,N'-bis[2-(3-[3,5-di-tert-butyl-4-hydroxyphenyl]propionyloxy)ethyl]oxamide (Naugard®XL-1, supplied by Uniroyal).


1.18. Ascorbic acid (vitamin C)

[0051] 1.19. Aminic antioxidants, for example N,N'-di-isopropyl-p-phenylenediamine, N,N'-di-sec-butyl-p-phenylenediamine, N,N'-bis(1,4-dimethylpentyl)-p-phenylenediamine, N,N'-bis(1-ethyl-3-methylpentyl)-p-phenylenediamine, N,N'-bis(1-methylheptyl)-p-phenylenediamine, N,N'-dicyclohexyl-p-phenylenediamine, N,N'-diphenyl-p-phenylenediamine, N,N'-bis(2-naphthyl)-p-phenylenediamine, N-isopropyl-N'-phenyl-p-phenylenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-p-phenylenediamine, N-(1-methylheptyl)-N'-phenyl-p-phenylenediamine, N-cyclohexyl-N'-phenyl-p-phenylenediamine, 4-(p-toluenesulfamoyl)diphenylamine, N,N'-dimethyl-N,N'-di-sec-butyl-p-phenylenediamine, diphenylamine, N-allyldiphenylamine, 4-isopropoxydiphenylamine, N-phenyl-1-naphthylamine, N-(4-tert-octylphenyl)-1-naphthylamine, N-phenyl-2-naphthylamine, octylated diphenylamine, for example p,p'-di-tert-octyldiphenylamine, 4-n-butylaminophenol, 4-butyrylamino-phenol, 4-nonanoylamino-phenol, 4-dodecanoylamino-phenol, 4-octadecanoylamino-phenol, bis(4-methoxyphenyl)amine, 2,6-di-tert-butyl-4-dimethylaminomethylphenol, 2,4'-diaminodiphenylmethane, 4,4'-diaminodiphenylmethane, N,N,N',N'-tetramethyl-4,4'-diaminodiphenylmethane, 1,2-bis[(2-methylphenyl)amino]ethane, 1,2-bis(phenylamino)propane, (o-tolyl)biguanide, bis[4-(1',3'-dimethylbutyl)phenyl]amine, tert-octylated N-phenyl-1-naphthylamine, a mixture of mono- and dialkylated tert-butyl/tert-octyldiphenylamines, a mixture of mono- and dialkylated nonyldiphenylamines, a mixture of mono- and dialkylated dodecyldiphenylamines, a mixture of mono- and dialkylated isopropyl/isohexyldiphenylamines, a mixture of mono- and dialkylated tert-butyldiphenylamines,

2,3-dihydro-3,3-dimethyl-4H-1,4-benzothiazine, phenothiazine, a mixture of mono- and dialkylated tert-butyl/tert-octylphenothiazines, a mixture of mono- and dialkylated tert-octylphenothiazines, N-allylphenothiazine, N,N,N',N'-tetraphenyl-1,4-diaminobut-2-ene.

2. UV absorbers and light stabilizers

[0052] 2.1. 2-(2'-Hydroxyphenyl)benzotriazoles, for example 2-(2'-hydroxy-5'-methylphenyl)-benzotriazole, 2-(3',5'-di-tert-butyl-2'-hydroxyphenyl)benzotriazole, 2-(5'-tert-butyl-2'-hydroxyphenyl)benzotriazole, 2-(2'-hydroxy-5'-(1,1,3,3-tetramethylbutyl)phenyl)benzotriazole, 2-(3',5'-di-tert-butyl-2'-hydroxyphenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-methylphenyl)-5-chloro-benzotriazole, 2-(3'-sec-butyl-5'-tert-butyl-2'-hydroxyphenyl)benzotriazole, 2-(2'-hydroxy-4'-octyloxyphenyl)benzotriazole, 2-(3',5'-di-tert-amyl-2'-hydroxyphenyl)benzotriazole, 2-(3',5'-bis-(α,α -dimethylbenzyl)-2'-hydroxyphenyl)benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)-carbonyl]ethyl)-2'-hydroxyphenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)carbonyl]ethyl)-2'-hydroxyphenyl)benzotriazole, 2-(3'-dodecyl-2'-hydroxy-5'-methylphenyl)benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-iso-octyloxycarbonyl)ethyl)phenyl)benzotriazole, 2,2'-methylene-bis[4-(1,1,3,3-tetramethylbutyl)-6-benzotriazole-2-yl]phenol]; the transesterification product of 2-[3'-tert-butyl-5'-(2-methoxycarbonyl)ethyl)-2'-hydroxyphenyl]-2H-benzotriazole with polyethylene glycol 300;



where R = 3'-tert-butyl-4'-hydroxy-5'-2H-benzotriazol-2-ylphenyl, 2-[2'-hydroxy-3'-(α,α -dimethylbenzyl)-5'-(1,1,3,3-tetramethylbutyl)-phenyl]benzotriazole; 2-[2'-hydroxy-3'-(1,1,3,3-tetramethylbutyl)-5'-(α,α -dimethylbenzyl)-phenyl]benzotriazole.

[0053] 2.2. 2-Hydroxybenzophenones, for example the 4-hydroxy, 4-methoxy, 4-octyloxy, 4-decyloxy, 4-dodecyloxy, 4-benzyloxy, 4,2',4'-trihydroxy and 2'-hydroxy-4,4'-dimethoxy derivatives.

[0054] 2.3. Esters of substituted and unsubstituted benzoic acids, for example 4-tert-butylphenyl salicylate, phenyl salicylate, octylphenyl salicylate, dibenzoyl resorcinol, bis(4-tert-butylbenzoyl)resorcinol, benzoyl resorcinol, 2,4-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate, hexadecyl 3,5-di-tert-butyl-4-hydroxybenzoate, octadecyl 3,5-di-tert-butyl-4-hydroxybenzoate, 2-methyl-4,6-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate.

[0055] 2.4. Acrylates, for example ethyl α -cyano- β,β -diphenylacrylate, isooctyl α -cyano- β,β -diphenylacrylate, methyl α -carbomethoxycinnamate, methyl α -cyano- β -methyl-p-methoxycinnamate, butyl α -cyano- β -methyl-p-methoxy-cinnamate, methyl α -carbomethoxy-p-

methoxycinnamate, N-(β -carbomethoxy- β -cyanovinyl)-2-methylindoline, neopentyl tetra(α -cyano- β , β -diphenylacrylate).

[0056] 2.5. Nickel compounds, for example nickel complexes of 2,2'-thio-bis[4-(1,1,3,3-tetramethylbutyl)phenol], such as the 1:1 or 1:2 complex, with or without additional ligands such as n-butylamine, triethanolamine or N-cyclohexyldiethanolamine, nickel dibutyldithiocarbamate, nickel salts of the monoalkyl esters, e.g. the methyl or ethyl ester, of 4-hydroxy-3,5-di-tert-butylbenzylphosphonic acid, nickel complexes of ketoximes, e.g. of 2-hydroxy-4-methylphenylundecylketoxime, nickel complexes of 1-phenyl-4-lauroyl-5-hydroxypyrazole, with or without additional ligands.

[0057] 2.6. Sterically hindered amines, for example carbonic acid bis(1-undecyloxy-2,2,6,6-tetramethyl-4-piperidyl)ester, bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate, bis(2,2,6,6-tetramethyl-4-piperidyl)succinate, bis(1,2,2,6,6-pentamethyl-4-piperidyl)sebacate, bis(1-octyloxy-2,2,6,6-tetramethyl-4-piperidyl)sebacate, bis(1,2,2,6,6-pentamethyl-4-piperidyl) n-butyl-3,5-di-tert-butyl-4-hydroxybenzylmalonate, the condensate of 1-(2-hydroxyethyl)-2,2,6,6-tetramethyl-4-hydroxypiperidine and succinic acid, linear or cyclic condensates of N,N'-bis(2,2,6,6-tetramethyl-4-piperidyl)hexamethylenediamine and 4-tert-octylamino-2,6-dichloro-1,3,5-triazine, tris(2,2,6,6-tetramethyl-4-piperidyl)nitrilotriacetate, tetrakis(2,2,6,6-tetramethyl-4-piperidyl)-1,2,3,4-butanetetracarboxylate, 1,1'-(1,2-ethanediyl)-bis(3,3,5,5-tetramethylpiperazinone), 4-benzoyl-2,2,6,6-tetramethylpiperidine, 4-stearyloxy-2,2,6,6-tetramethylpiperidine, bis(1,2,2,6,6-pentamethylpiperidyl)-2-n-butyl-2-(2-hydroxy-3,5-di-tert-butylbenzyl)malonate, 3-n-octyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decane-2,4-dione, bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)sebacate, bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)succinate, linear or cyclic condensates of N,N'-bis(2,2,6,6-tetramethyl-4-piperidyl)hexamethylenediamine and 4-morpholino-2,6-dichloro-1,3,5-triazine, the condensate of 2-chloro-4,6-bis(4-n-butylamino-2,2,6,6-tetramethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane, the condensate of 2-chloro-4,6-di-(4-n-butylamino-1,2,2,6,6-pentamethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane, 8-acetyl-3-dodecyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decane-2,4-dione, 3-dodecyl-1-(2,2,6,6-tetramethyl-4-piperidyl)pyrrolidine-2,5-dione, 3-dodecyl-1-(1,2,2,6,6-pentamethyl-4-piperidyl)pyrrolidine-2,5-dione, a mixture of 4-hexadecyloxy- and 4-stearyloxy-2,2,6,6-tetramethylpiperidine, a condensate of N,N'-bis(2,2,6,6-tetramethyl-4-piperidyl)hexamethylenediamine and 4-cyclohexylamino-2,6-dichloro-1,3,5-triazine, a condensate of 1,2-bis(3-aminopropylamino)ethane and 2,4,6-trichloro-1,3,5-triazine as well as 4-butylamino-2,2,6,6-tetramethylpiperidine (CAS Reg. No. [136504-96-6]); a condensate of 1,6-hexanediamine and 2,4,6-trichloro-1,3,5-triazine as well as N,N-dibutylamine and 4-butylamino-2,2,6,6-tetramethylpiperidine (CAS Reg. No. [192268-64-7]); N-(2,2,6,6-tetramethyl-4-piperidyl)-n-dodecylsuccinimide, N-(1,2,2,6,6-pentamethyl-4-piperidyl)-n-dodecylsuccinimide, 2-undecyl-7,7,9,9-tetramethyl-1-oxa-3,8-diaza-4-oxo-spiro[4,5]decane, a reaction product of 7,7,9,9-tetramethyl-2-cycloundecyl-1-oxa-3,8-diaza-4-oxospiro-[4,5]decane and epichlorohydrin, 1,1-bis(1,2,2,6,6-pentamethyl-4-piperidyloxycarbonyl)-2-(4-methoxyphenyl)ethene, N,N'-bis-formyl-N,N'-bis(2,2,6,6-tetramethyl-4-piperidyl)hexamethylenediamine, a diester of 4-methoxymethylenemalonic acid with 1,2,2,6,6-

pentamethyl-4-hydroxypiperidine, poly[methylpropyl-3-oxy-4-(2,2,6,6-tetramethyl-4-piperidyl)]siloxane, a reaction product of maleic acid anhydride- α -olefin copolymer with 2,2,6,6-tetramethyl-4-aminopiperidine or 1,2,2,6,6-pentamethyl-4-aminopiperidine, 2,4-bis[N-(1-cyclohexyloxy-2,2,6,6-tetramethylpiperidine-4-yl)-N-butylamino]-6-(2-hydroxyethyl)amino-1,3,5-triazine, 1-(2-hydroxy-2-methylpropoxy)-4-octadecanoyloxy-2,2,6,6-tetramethylpiperidine, 5-(2-ethylhexanoyl)oxymethyl-3,3,5-trimethyl-2-morpholinone, Sanduvor (Clariant; CAS Reg. No. 106917-31-1], 5-(2-ethylhexanoyl)oxymethyl-3,3,5-trimethyl-2-morpholinone, the reaction product of 2,4-bis[(1-cyclohexyloxy-2,2,6,6-piperidine-4-yl)butylamino]-6-chloro-s-triazine with N,N'-bis(3-aminopropyl)ethylenediamine), 1,3,5-tris(N-cyclohexyl-N-(2,2,6,6-tetramethylpiperazine-3-one-4-yl)amino)-s-triazine, 1,3,5-tris(N-cyclohexyl-N-(1,2,2,6,6-pentamethylpiperazine-3-one-4-yl)amino)-s-triazine.

[0058] 2.7. Oxamides, for example 4,4'-dioctyloxyoxanilide, 2,2'-diethoxyoxanilide, 2,2'-dioctyloxy-5,5'-di-tert-butoxanilide, 2,2'-didodecyloxy-5,5'-di-tert-butoxanilide, 2-ethoxy-2'-ethyloxanilide, N,N'-bis(3-dimethylaminopropyl)oxamide, 2-ethoxy-5-tert-butyl-2'-ethoxanilide and its mixture with 2-ethoxy-2'-ethyl-5,4'-di-tert-butoxanilide, mixtures of o- and p-methoxy-disubstituted oxanilides and mixtures of o- and p-ethoxy-disubstituted oxanilides.

[0059] 2.8. 2-(2-Hydroxyphenyl)-1,3,5-triazines, for example 2,4,6-tris(2-hydroxy-4-octyloxyphenyl)-1,3,5-triazine, 2-(2-hydroxy-4-octyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2-(2,4-dihydroxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2,4-bis(2-hydroxy-4-propyloxyphenyl)-6-(2,4-dimethylphenyl)-1,3,5-triazine, 2-(2-hydroxy-4-octyloxyphenyl)-4,6-bis(4-methylphenyl)-1,3,5-triazine, 2-(2-hydroxy-4-dodecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2-(2-hydroxy-4-tridecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2-[2-hydroxy-4-(2-hydroxy-3-butyloxypropoxy)phenyl]-4,6-bis(2,4-dimethyl)-1,3,5-triazine, 2-[2-hydroxy-4-(2-hydroxy-3-octyloxypropyloxy)phenyl]-4,6-bis(2,4-dimethyl)-1,3,5-triazine, 2-[4-(dodecyloxy/tridecyloxy-2-hydroxypropoxy)-2-hydroxyphenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2-[2-hydroxy-4-(2-hydroxy-3-dodecyloxypropoxy)phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2-(2-hydroxy-4-hexyloxy)phenyl-4,6-diphenyl-1,3,5-triazine, 2-(2-hydroxy-4-methoxyphenyl)-4,6-diphenyl-1,3,5-triazine, 2,4,6-tris[2-hydroxy-4-(3-butoxy-2-hydroxypropoxy)phenyl]-1,3,5-triazine, 2-(2-hydroxyphenyl)-4-(4-methoxyphenyl)-6-phenyl-1,3,5-triazine, 2-[2-hydroxy-4-[3-(2-ethylhexyl-1-oxy)-2-hydroxypropyloxy]phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine, 2,4-bis(4-[2-ethylhexyloxy]-2-hydroxyphenyl)-6-(4-methoxyphenyl)-1,3,5-triazine, 2-(4,6-bis-biphenyl-4-yl-1,3,5-triazin-2-yl)-5-(2-ethyl-(n)-hexyloxy)phenol.





[0060] 3. Metal deactivators, for example N,N'-diphenyloxamide, N-salicylal-N'-salicyloyl hydrazine, N,N'-bis(salicyloyl)hydrazine, N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hydrazine, 3-salicyloylamino-1,2,4-triazole, bis(benzylidene)oxalyl dihydrazide, oxanilide, isophthaloyl dihydrazide, sebacoyl bisphenylhydrazide, N,N'-diacetyladiopoyl dihydrazide, N,N'-bis(salicyloyl)oxalyl dihydrazide, N,N'-bis(salicyloyl)thiopropionyl dihydrazide.




[0061] 4. Phosphites and phosphonites, for example triphenyl phosphite, diphenylalkyl

phosphites, phenyldialkyl phosphites, tris(nonylphenyl) phosphite, trilauryl phosphite, trioctadecyl phosphite, distearyl pentaerythritol diphosphite, tris(2,4-di-tert-butylphenyl) phosphite, diisodecyl pentaerythritol diphosphite, bis(2,4-di-tert-butylphenyl) pentaerythritol diphosphite, bis(2,4-di-cumylphenyl) pentaerythritol diphosphite, bis(2,6-di-tert-butyl-4-methylphenyl) pentaerythritol diphosphite, diisodecyloxypentaerythritol diphosphite, bis(2,4-di-tert-butyl-6-methylphenyl) pentaerythritol diphosphite, bis(2,4,6-tris(tert-butylphenyl) pentaerythritol diphosphite, tristearyl sorbitol triphosphite, tetrakis(2,4-di-tert-butylphenyl) 4,4'-biphenylene diphosphonite, 6-isooctyloxy-2,4,8,10-tetra-tert-butyl-12H-dibenz[d,g]-1,3,2-dioxaphosphocin, bis(2,4-di-tert-butyl-6-methylphenyl) methyl phosphite, bis(2,4-di-tert-butyl-6-methylphenyl) ethyl phosphite, 6-fluoro-2,4,8,10-tetra-tert-butyl-12-methyl-dibenz[d,g]-1,3,2-dioxaphosphocin, 2,2',2''-nitrilo[triethyltris(3,3',5,5'-tetra-tert-butyl-1,1'-biphenyl-2,2'-diyl) phosphite], 2-ethylhexyl (3,3',5,5'-tetra-tert-butyl-1,1'-biphenyl-2,2'-diyl) phosphite, 5-butyl-5-ethyl-2-(2,4,6-tri-tert-butylphenoxy)-1,3,2-dioxaphosphirane.

[0062] The following phosphites are especially preferred:

Tris(2,4-di-tert-butylphenyl) phosphite (Irgafos®168, Ciba Specialty Chemicals Inc.), tris(nonylphenyl) phosphite,





[0063] 5. Hydroxylamines. for example N,N-dibenzylhydroxylamine, N,N-diethylhydroxylamine, N,N-dioctylhydroxylamine, N,N-dilaurylhydroxylamine, N,N-ditetradecylhydroxylamine, N,N-dihexadecylhydroxylamine, N,N-dioctadecylhydroxylamine, N-hexadecyl-N-octadecylhydroxylamine, N-heptadecyl-N-octadecylhydroxylamine, N,N-dialkylhydroxylamine derived from hydrogenated tallow amine.

[0064] 6. Nitrones. for example, N-benzyl-alpha-phenylnitron, N-ethyl-alpha-methylnitron, N-octyl-alpha-heptylnitron, N-lauryl-alpha-undecylnitron, N-tetradecyl-alpha-tridecylinnitron, N-hexadecyl-alpha-pentadecylnitron, N-octadecyl-alpha-heptadecylnitron, N-hexadecyl-alpha-heptadecylnitron, N-octadecyl-alpha-pentadecylnitron, N-heptadecyl-alpha-heptadecylnitron, N-octadecyl-alpha-hexadecylnitron, nitron derived from N,N-dialkylhydroxylamine derived from hydrogenated tallow amine.

[0065] 7. Thiosynergists. for example dilauryl thiodipropionate, dimistyl thiodipropionate, distearyl thiodipropionate, pentaerythritol tetrakis[3-(dodecylthio)propionate] or distearyl disulfide.

[0066] 8. Peroxide scavengers. for example esters of β -thiodipropionic acid, for example the lauryl, stearyl, myristyl or tridecyl esters, mercaptobenzimidazole or the zinc salt of 2-mercaptobenzimidazole, zinc dibutyldithiocarbamate, dioctadecyl disulfide, pentaerythritol tetrakis(β -dodecylmercapto)propionate.

[0067] 9. Polyamide stabilizers. for example copper salts in combination with iodides and/or phosphorus compounds and salts of divalent manganese.

[0068] 10. Basic co-stabilizers. for example melamine, polyvinylpyrrolidone, dicyandiamide, triallyl cyanurate, urea derivatives, hydrazine derivatives, amines, polyamides, polyurethanes,

alkali metal salts and alkaline earth metal salts of higher fatty acids, for example calcium stearate, zinc stearate, magnesium behenate, magnesium stearate, sodium ricinoleate and potassium palmitate, antimony pyrocatecholate or zinc pyrocatecholate.

[0069] 11. Nucleating agents, for example inorganic substances, such as talcum, metal oxides, such as titanium dioxide or magnesium oxide, phosphates, carbonates or sulfates of, preferably, alkaline earth metals; organic compounds, such as mono- or polycarboxylic acids and the salts thereof, e.g. 4-tert-butylbenzoic acid, adipic acid, diphenylacetic acid, sodium succinate or sodium benzoate; polymeric compounds, such as ionic copolymers (ionomers). Especially preferred are 1,3:2,4-bis(3',4'-dimethylbenzylidene)sorbitol, 1,3:2,4-di(paramethyldibenzylidene)sorbitol, and 1,3:2,4-di(benzylidene)sorbitol.

[0070] 12. Fillers and reinforcing agents, for example calcium carbonate, silicates, glass fibres, glass beads, asbestos, talc, kaolin, mica, barium sulfate, metal oxides and hydroxides, carbon black, graphite, wood flour and flours or fibers of other natural products, synthetic fibers.

[0071] 13. Other additives, for example plasticisers, lubricants, emulsifiers, pigments, rheology additives, catalysts, flow-control agents, optical brighteners, flameproofing agents, antistatic agents and blowing agents.

[0072] 14. Benzofuranones and indolinones, for example those disclosed in U.S. 4,325,863; U.S. 4,338,244; U.S. 5,175,312; U.S. 5,216,052; U.S. 5,252,643; DE-A-4316611; DE-A-4316622; DE-A-4316876; EP-A-0589839, EP-A-0591102; EP-A-1291384 or 3-[4-(2-acetoxyethoxy)phenyl]-5,7-di-tert-butylbenzofuran-2-one, 5,7-di-tert-butyl-3-[4-(2-stearoyloxyethoxy)phenyl]benzofuran-2-one, 3,3'-bis[5,7-di-tert-butyl-3-(4-[2-hydroxyethoxy]phenyl)benzofuran-2-one], 5,7-di-tert-butyl-3-(4-ethoxyphenyl)benzofuran-2-one, 3-(4-acetoxy-3,5-dimethylphenyl)-5,7-di-tert-butylbenzofuran-2-one, 3-(3,5-dimethyl-4-pivaloyloxyphenyl)-5,7-di-tert-butylbenzofuran-2-one, 3-(3,4-dimethylphenyl)-5,7-di-tert-butylbenzofuran-2-one, 3-(2,3-dimethylphenyl)-5,7-di-tert-butylbenzofuran-2-one, 3-(2-acetyl-5-isooctylphenyl)-5-isooctylbenzofuran-2-one.

[0073] The weight ratio of component (A) to the conventional additive is for example 1:100 to 100:1, preferably 1:100 to 10:1, in particular 1:10 to 10:1.

[0074] According to a preferred embodiment the polyolefin additionally contains as component (C) a phenolic antioxidant, preferably octadecyl 3-(3,5-di-tert-butyl-4-hydroxyphenyl)propionate.

[0075] Component (C) may be present in a concentration of for example 0.01 to 10%, relative to the weight of the polyolefin.

[0076] According to another preferred embodiment component (A) is a compound of the formula (A-I-1) and a compound of the formula (A-III-2), and component (B) is a compound of the formula (B-I-1) or (B-II-1), and the polyolefin optionally comprises octadecyl 3-(3,5-di-tert-

butyl-4-hydroxyphenyl)propionate.

[0077] A further embodiment of the present invention is a polyolefin film, tape or monofilament made in accordance with the process described above.

[0078] Another embodiment of the present invention is the use of component (B) as light stabilizer to reduce the water carry-over of a light stabilized polyolefin film, tape or monofilament which contains component (A) as light stabilizer and which is passed through a water bath during production.

[0079] Still another embodiment of the present invention is a process for preparing a light stabilized polyolefin film, tape or monofilament, comprising the step of passing said light stabilized polyolefin film, tape or monofilament through a water bath at a haul off speed of 5 to 100 m/min, preferably 4 to 80 m/min or 10 to 50 m/min, in particular 20 to 40 m/min, characterized in that said polyolefin, film, tape or monofilament contains components (A) and (B) as defined above. All preferences described above also relate to this process which is useful for reducing the water carry-over from the water bath.


[0080] The examples below illustrate the invention in greater detail. All percentages and parts are by weight, unless stated otherwise.

Light stabilizers used in the following examples:

Compound (A-I-1):

(Tinuvin®622)

[0081]




wherein a_1 is a number from 2 to 10.


Compound (A-II-1-a):

(Sabostab® UV119)


[0082]



wherein X is the group




and Y is the group




Compound (A-III-2):

(Chimassorb®2020)


[0083]



wherein E₁ is




, E₂ is



and a_3 is a number from 2 to 10.


Compound (B-I-1):

[0084]




Compound (B-II-1):


[0085]



wherein E_3 is




, E_4 is




and wherein b_1 is a number from 2 to 10.

Compound (B-III-1):

(Flamestab®NOR 116)

[0086]

wherein R₇ is a group of the formula



Determination of synergism for the experiments indicated below:

[0087] The synergistic WCO (water carry-over) effect of two light stabilizers (α) and (β) is determined by a comparison of the expected values of the WCO with the actually measured values of the WCO activity. The expected values are calculated on the basis of the additivity law (B. Ranby and J.F. Rabek; Photodegradation, Photo-oxidation and Photostabilization of Polymers, Principles and Applications, John Wiley & Sons, London, New York, Sydney, Toronto, 1975, pages 418 and 419) according to the following equation for a 1:1 ratio of the two light stabilizers:

$$\text{Expected WCO activity} = \frac{\text{WCO activity of 100 \% } (\alpha) + \text{WCO activity of 100 \% } (\beta)}{2}$$

[0088] There is a synergistic WCO effect of the two light stabilizers in the samples below, when the "measured WCO" < the "calculated WCO" (Low WCO values are desired).

[0089] The above equation is adapted in appropriate manner, when the ratio of the two light stabilizers is different from 1:1; for example if the ratio is 2:1, the expected WCO effect can be calculated according to the following equation:

$$\text{Expected WCO activity} = \frac{2 \times [\text{WCO activity of 100 \% } (\alpha)] + 1 \times [\text{WCO activity of 100 \% } (\beta)]}{3}$$

Example 1:

[0090] The light stabilizers indicated in Tables 1a and 1b are mixed in a high speed mixer (MTI, Mischtechnik International, Germany) to polyethylene (Dowlex®SC 2108). The mixture is extruded at 200°C in a double screw extruder (Krauss Maffei Berstorff, Germany) to give granules which are subsequently converted in a semi industrial tape line (Leonard, Italy) to tapes at 220°C. Water carry-over is expressed in length (mm) to which water from the water cooling bath is carried along with the extrudate (200 micron) before entering the drawing part. The lower the water carry-over, the faster and more economically a film, drawn tape or monofilament can be processed. The results are shown in Tables 1a and 1b.

Table 1a:

Haul off speed (m/min)	30	35	40
Light stabilizer	Water carry-over in mm *)		
0.8% of Compound (A-I-1)	120	130	140
0.8% of Compound (B-II-1)	20	40	50
0.7% of Compound (A-I-1)	80	100	120
plus 0.1% of Compound (B-II-1)			
	Calculated: 108	Calculated: 119	Calculated: 129
*) Low values are desired.			

Table 1b:

Haul off speed (m/min)	30	35	40
Light stabilizer	Water carry-over in mm *)		
0.8% of Compound (A-II-1-a)	380	450	450
0.8% of Compound (B-III-1)	30	50	60
0.5% of Compound (A-II-1-a)	70	120	130
plus 0.3% of Compound (B-III-1)			
	Calculated: 249	Calculated: 300	Calculated: 304
*) Low values are desired.			

Example 2:

[0091] The light stabilizers indicated in Tables 2a and 2b are mixed in a high speed mixer (MTI, Mischtechnik International, Germany) to polyethylene (Dowlex®SC 2108) which contains 0.5 % by weight, relative to the weight of the polyethylene, of Compound (A-I-1). The mixture is extruded at 200°C in a double screw extruder (Krauss Maffei Berstorff, Germany) to give granules which are subsequently converted in a semi industrial tape line (Leonard, Italy) to tapes at 220°C. Water carry-over is expressed in length (mm) to which water from the water cooling bath is carried along with the extrudate (200 micron) before entering the drawing part.

The lower the water carry-over, the faster and more economically a film, drawn tape or monofilament can be processed. The results are shown in Tables 2a and 2b.

Table 2a:

Light stabilizer	Water carry-over in mm *) at a haul off speed of 27 m/min
0.3% of Compound (A-III-2)	350
0.3% of Compound (B-II-1)	60
0.2% of Compound (A-III-2) plus 0.1% of Compound (B-II-1)	180
	Calculated: 253
*) Low values are desired.	

Table 2b:

Light stabilizer	Water carry-over in mm *) at a haul off speed of 27 m/min
0.3% of Compound (A-III-2)	350
0.3% of Compound (B-I-1)	70
0.25% of Compound (A-III-2)	70
plus	
0.05% of Compound (B-I-1)	
	Calculated: 303
*) Low values are desired.	

Example 3:

[0092] The light stabilizers indicated in Tables 3a and 3b are mixed in a high speed mixer (MTI, Mischtechnik International, Germany) to polypropylene (Polychim®A 10 TB). The mixture is extruded at 230°C in a double screw extruder (Krauss Maffei Berstorff, Germany) to give granules which are subsequently converted in a semi industrial tape line (Leonard, Italy) to tapes at 240°C. Water carry-over is expressed in length (mm) to which water from the water cooling bath is carried along with the extrudate (240 micron) before entering the drawing part. The lower the water carry-over, the faster and more economically a film, drawn tape or monofilament can be processed. The results are shown in Tables 3a and 3b.

Table 3a:

Light stabilizer	Water carry-over in mm *) at a haul off speed of 20 m/min
0.6% of Compound (A-III-2)	> 600
0.6% of Compound (B-I-1)	130
0.3% of Compound (A-III-2)	210

Light stabilizer	Water carry-over in mm *) at a haul off speed of 20 m/min
plus	
0.3% of Compound (B-I-1)	
	Calculated: >365
*) Low values are desired.	

Table 3b:

Light stabilizer	Water carry-over in mm *) at a haul off speed of 20 m/min
0.6% of Compound (A-III-2)	> 600
0.6% of Compound (B-I-1)	130
0.4% of Compound (A-III-2)	350
plus	
0.2% of Compound (B-I-1)	
	Calculated: >443
*) Low values are desired.	

Example 4:

[0093] The light stabilizers indicated in Table 4 are mixed in a high speed mixer (MTI, Mischtechnik International, Germany) to polyethylene (Dowlex®SC 2108) which contains 0.5 % by weight, relative to the weight of the polyethylene, of Compound (A-I-1). The mixture is extruded at 200°C in a double screw extruder (Krauss Maffei Berstorff, Germany) to give granules which are subsequently converted in a semi industrial tape line (Leonard, Italy) to tapes at 220°C. Water carry-over is expressed in length (mm) to which water from the water cooling bath is carried along with the extrudate (100 micron) before entering the drawing part. The lower the water carry-over, the faster and more economically a film, drawn tape or monofilament can be processed. The results are indicated in Table 4.

Table 4:

Light stabilizer	Water carry-over in mm *) at a haul off speed of 21 m/min
0.3% of Compound (A-III-2)	140
0.25% of Compound (A-III-2)	100
plus	
0.05% of Compound (B-II-1)	
plus	
0.025% of octadecyl 3-(3,5-di-tert-butyl-4-	

Light stabilizer	Water carry-over in mm *) at a haul off speed of 21 m/min
hydroxyphenyl)propionate	
*) Low values are desired.	

[0094] In the above examples "%" means "% by weight relative to the weight of the polyolefin".

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US5006587A](#) [0004]
- [US4233412A](#) [0010]
- [US4477615A](#) [0010]
- [US4108829A](#) [0010]
- [US4086204A](#) [0010]
- [US4331586A](#) [0010]
- [US6046304A](#) [0010] [0022]
- [US6297299A](#) [0010]
- [US6388072A](#) [0011]
- [US6117995A](#) [0011]
- [US6420462A](#) [0011]
- [US6677451A](#) [0011]
- [US5844026A](#) [0011]
- [US4325863A](#) [0072]
- [US4338244A](#) [0072]
- [US5175312A](#) [0072]
- [US5216052A](#) [0072]
- [US5252643A](#) [0072]
- [DE4316611A](#) [0072]
- [DE4316622A](#) [0072]
- [DE4316876A](#) [0072]
- [EP0589839A](#) [0072]
- [EP0591102A](#) [0072]

- [EP1291384A](#) [\[0072\]](#)


Non-patent literature cited in the description

- CHEMICAL ABSTRACTS, 136,504-96-6 [\[0010\]](#)
- CHEMICAL ABSTRACTS, 136504-96-6 [\[0057\]](#)
- CHEMICAL ABSTRACTS, 192268-64-7 [\[0057\]](#)
- CHEMICAL ABSTRACTS, 106917-31-1 [\[0057\]](#)
- Photodegradation, Photo-oxidation and Photostabilization of Polymers **B. RANBY** **J.F. RABEK** Principles and Applications John Wiley & Sons 1975 0000418-419- [\[0087\]](#)

1. Blanding, der omfatter forbindelserne med formel (A-I-1) og (A-III-2) sammen med en forbindelse med formel (B-I-1) eller en forbindelse med formel (B-II-1)


$$\left[\begin{array}{c} \text{N} \text{---} (\text{CH}_2)_6 \text{---} \text{N} \text{---} \text{C}_6\text{H}_3\text{N}_2 \text{---} \text{N} \text{---} \text{C}_6\text{H}_3\text{N}_2 \text{---} \text{N} \\ \text{H}_3\text{C} \quad \text{CH}_3 \quad \text{H}_3\text{C} \quad \text{CH}_3 \quad \text{H}_3\text{C} \quad \text{CH}_3 \\ \text{H}_3\text{C} \quad \text{H} \quad \text{H}_3\text{C} \quad \text{H} \quad \text{H}_3\text{C} \quad \text{CH}_3 \\ \text{H} \quad \text{H} \quad \text{C}_4\text{H}_9 \quad \text{H}_3\text{C} \quad \text{CH}_3 \end{array} \right]_{a_3} \quad (\text{A-III-2})$$
CC(C)(O)CO[N+]1(C)CC(C)(C)CC1OC(=O)CCCCCCCCCCCCCCC

(B-I-1)



2. Blanding ifølge krav 1, der omfatter forbindelserne med formel (A-I-1) og (A-III-2) sammen med en forbindelse med formel (B-I-1).

3. Blanding ifølge krav 1, der omfatter forbindelserne med formel (A-I-1) og (A-III-2) sammen med en forbindelse med formel (B-II-1).