



US 20110135613A1

(19) **United States**

(12) **Patent Application Publication**
HUANG et al.

(10) **Pub. No.: US 2011/0135613 A1**

(43) **Pub. Date: Jun. 9, 2011**

(54) **METHODS FOR TREATING
APOLIPOPROTEIN E4-ASSOCIATED
DISORDERS**

Publication Classification

(75) Inventors: **Yadong HUANG**, San Francisco,
CA (US); **Yaisa Andrews-Zwilling**,
San Francisco, CA (US)

(51) **Int. Cl.**
A61K 35/12 (2006.01)
A61K 31/7088 (2006.01)
A61K 35/30 (2006.01)
A61P 25/00 (2006.01)
C12N 5/10 (2006.01)

(73) Assignee: **The J. David Gladstone Institutes**,
San Francisco, CA (US)

(52) **U.S. Cl.** **424/93.21**; 514/44 A; 424/93.7;
435/325

(21) Appl. No.: **12/958,057**

(22) Filed: **Dec. 1, 2010**

(57) **ABSTRACT**

Related U.S. Application Data

(60) Provisional application No. 61/356,977, filed on Jun.
21, 2010, provisional application No. 61/266,449,
filed on Dec. 3, 2009.

The present disclosure provides a method of increasing the
functionality of a GABAergic interneuron in the hilus of the
hippocampus of an individual having at least one apolipoprotein
E4 (apoE4) allele. The method generally involves reducing
tau levels in the interneuron.




FIG. 1




FIG. 2







FIG. 3







FIG. 3 (Cont.)




FIG. 4




FIG. 5




FIG. 6






FIG. 6 (Cont.)





FIG. 7




FIG. 8







FIG. 8 (Cont.)




FIG. 9




FIG. 9 (Cont.)




FIG. 10




FIG. 10 (Cont.)

catg agtttgccat gttgagcagg actatttctg gcacttgcaa gtcccatgat
ttcttcggta attctgaggg tggggggagg gacatgaaat catcttagct tagctttctg
tctgtgaatg tctatatagt gtattgtgtg ttttaacaaa tgatttacac tgactgttgc
tgtaaaagtg aatttggaat taaagttatt actctgat

(nucleotides 6547-6758 of tau1 mRNA; SEQ ID NO:7)

FIG. 11A

acggccgagc ggcagggcgc tcgcgcgcgc ccactagtgg ccggaggaga aggctcccgc
ggaggccgcg ctgcccgcgc cctcccctgg ggaggctcgc gttcccctg ctgcgcctg
cgccgcccgc cggcctcagg aacgcgcctt ctgcgcgcgc gcgcgcctc gcagtcaccg
ccaccacca gctccggcac caacagcagc gccgctgcca ccgccaccct tctgcccgcg

(nucleotides 1-240 of tau1 mRNA; SEQ ID NO:8)

FIG. 11B

METHODS FOR TREATING APOLIPOPROTEIN E4-ASSOCIATED DISORDERS

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 61/266,449, filed Dec. 3, 2009, and U.S. Provisional Patent Application No. 61/356,977, filed Jun. 21, 2010, which applications are incorporated herein by reference in their entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with government support under Grant No. P01 AG022074 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND

[0003] Apolipoprotein (apo) E, a polymorphic protein with three isoforms (apoE2, apoE3, and apoE4), is essential for lipid homeostasis. Carriers of apoE4 are at higher risk for developing Alzheimer's disease (AD). The hippocampus is one of the first regions of the brain damaged in AD, and memory deficits and disorientation are among the early symptoms.

[0004] Tau protein is expressed in central nervous system and plays a critical role in the neuronal architecture by stabilizing intracellular microtubule network. Impairment of the physiological role of the tau protein either by truncation, hyperphosphorylation or by disturbing the balance between the six naturally occurring tau isoforms leads to the formation of neurofibrillary tangles (NFT), dystrophic neurites and neuropil threads. These structures represent ultrastructural hallmarks of Alzheimer's disease (AD). The major protein subunit of these structures is microtubule associated protein Tau. The amount of NFT found in autopsies of AD patients correlates with clinical symptoms including intellectual decline. Therefore, Tau protein plays a critical role in AD pathology.

LITERATURE

[0005] Roberson et al. (2007) *Science* 316:750; Brunden et al. (Oct. 1, 2009) *Nature Reviews Drug Discovery* 8:783.

SUMMARY

[0006] The present disclosure provides a method of increasing the functionality of a GABAergic interneuron in the hilus of the hippocampus of an individual having at least one apolipoprotein E4 (apoE4) allele. The method generally involves reducing tau levels in the interneuron.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] FIGS. 1A-I depict age-dependent decrease in numbers of GABAergic interneurons in the hilus of dentate gyrus of female ApoE4-KI mice.

[0008] FIGS. 2A-M illustrate that presynaptic GABAergic input onto granule cells is reduced in female ApoE4-KI mice.

[0009] FIGS. 3A-H depict the correlation of hilar GABAergic interneuron impairment with spatial learning deficits in ApoE4-KI mice.

[0010] FIGS. 4A-D illustrate that GABAA receptor potentiator pentobarbital rescues spatial learning and memory deficits in ApoE4-KI mice.

[0011] FIGS. 5A-P depict localization of apoE4(Δ 272-299) in the hippocampus and its effects on neurodegeneration and tau pathology in the presence and absence of Tau.

[0012] FIGS. 6A-R depict loss of GABAergic interneurons in the hilus of the dentate gyrus of ApoE4(Δ 272-299)mE^{-/-}Tau^{+/+} mice and rescue by tau removal.

[0013] FIGS. 7A-E illustrate that eliminating Tau prevents the neurotoxic effect of ApoE4 fragments on primary hippocampal GABAergic neurons

[0014] FIGS. 8A-H depict spatial learning and memory deficits in ApoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice and rescue by Tau removal.

[0015] FIGS. 9A-F illustrate the performance in the cued platform trial does not correlate with the number of hilar GABAergic interneurons in apoE3-KI and apoE4-KI mice.

[0016] FIGS. 10A-H depict: (A) the effect of eliminating tau on apoE4 fragment-caused abnormal anxiety in apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice; (B-D) performance in the cued platform trial and the number of hilar GABAergic interneurons in apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice; (E) the effect of treatment with the GABAA receptor antagonist picrotoxin (Picro) on the number of hilar GABAergic interneurons in ApoE4(Δ 272-299) mE^{-/-}Tau^{-/-} mice; (F and G) the effect of treatment with a low dose of picrotoxin on the learning and memory performance in wildtype and mE^{-/-}Tau^{+/+} mice; and (H) the effect of the GABAA receptor potentiator pentobarbital on the learning deficit in apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice.

[0017] FIGS. 11A and 11B depict exemplary tau target sequences (SEQ ID NOs:7 and 8).

DEFINITIONS

[0018] As used herein, an "apoE4-associated disorder" is any disorder that is caused by the presence of apoE4 in a cell, in the serum, in the interstitial fluid, in the cerebrospinal fluid, or in any other bodily fluid of an individual; any physiological process or metabolic event that is influenced by apoE4 domain interaction; any disorder that is characterized by the presence of apoE4; a symptom of a disorder that is caused by the presence of apoE4 in a cell or in a bodily fluid; a phenomenon associated with a disorder caused by the presence in a cell or in a bodily fluid of apoE4; and the sequelae of any disorder that is caused by the presence of apoE4. ApoE4-associated disorders include apoE4-associated neurological disorders and disorders related to high serum lipid levels. ApoE4-associated neurological disorders include, but are not limited to, sporadic Alzheimer's disease; familial Alzheimer's disease; poor outcome following a stroke; poor outcome following traumatic head injury; and cerebral ischemia. Phenomena associated with apoE4-associated neurological disorders include, but are not limited to, neurofibrillary tangles; amyloid deposits; memory loss; and a reduction in cognitive function. ApoE4-related disorders associated with high serum lipid levels include, but are not limited to, atherosclerosis, and coronary artery disease. Phenomena associated with such apoE4-associated disorders include high serum cholesterol levels.

[0019] The term "Alzheimer's disease" (abbreviated herein as "AD") as used herein refers to a condition associated with formation of neuritic plaques comprising amyloid β protein primarily in the hippocampus and cerebral cortex, as well as

impairment in both learning and memory. "AD" as used herein is meant to encompass both AD as well as AD-type pathologies.

[0020] The term "phenomenon associated with Alzheimer's disease" as used herein refers to a structural, molecular, or functional event associated with AD, particularly such an event that is readily assessable in an animal model. Such events include, but are not limited to, amyloid deposition, neuropathological developments, learning and memory deficits, and other AD-associated characteristics.

[0021] As used herein, the term "neural stem cell" (NSC) refers to an undifferentiated neural cell that can proliferate, self-renew, and differentiate into the main adult neural cells of the brain. NSCs are capable of self-maintenance (self-renewal), meaning that with each cell division, one daughter cell will also be a stem cell. The non-stem cell progeny of NSCs are termed neural progenitor cells. Neural progenitor cells generated from a single multipotent NSC are capable of differentiating into neurons, astrocytes (type I and type II), and oligodendrocytes. Hence, NSCs are "multipotent" because their progeny have multiple neural cell fates. Thus, NSCs can be functionally defined as a cell with the ability to: 1) proliferate, 2) self-renew, and 3) produce functional progeny that can differentiate into the three main cell types found in the central nervous system: neurons, astrocytes and oligodendrocytes.

[0022] As used herein, the terms "neural progenitor cell" or "neural precursor cell" refer to a cell that can generate progeny such as neuronal cells (e.g., neuronal precursors or mature neurons), glial precursors, mature astrocytes, or mature oligodendrocytes. Typically, the cells express some of the phenotypic markers that are characteristic of the neural lineage. A "neuronal progenitor cell" or "neuronal precursor cell" is a cell that can generate progeny that are mature neurons. These cells may or may not also have the capability to generate glial cells.

[0023] A "neurosphere" is a group of cells derived from a single neural stem cell as the result of clonal expansion. A method for culturing neural stem cells to form neurospheres has been described in, for example, U.S. Pat. No. 5,750,376.

[0024] The terms "polynucleotide" and "nucleic acid," used interchangeably herein, refer to a polymeric form of nucleotides of any length, either ribonucleotides or deoxynucleotides. Thus, this term includes, but is not limited to, single-, double-, or multi-stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, or a polymer comprising purine and pyrimidine bases or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases.

[0025] The nucleic acid may be double stranded, single stranded, or contain portions of both double stranded or single stranded sequence. As will be appreciated by those in the art, the depiction of a single strand ("Watson") also defines the sequence of the other strand ("Crick"). By the term "recombinant nucleic acid" herein is meant nucleic acid, originally formed in vitro, in general, by the manipulation of nucleic acid by endonucleases, in a form not normally found in nature. Thus an isolated nucleic acid, in a linear form, or an expression vector formed in vitro by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell or organism, it will replicate non-recombinantly, i.e. using the in vivo cellular machinery of the host cell rather than

in vitro manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention.

[0026] Nucleic acid sequence identity (as well as amino acid sequence identity) is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 residues long, more usually at least about 30 residues long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul et al. (1990), *J. Mol. Biol.* 215:403-10 (using default settings, i.e. parameters $w=4$ and $T=17$).

[0027] The terms "polypeptide," "peptide," and "protein," used interchangeably herein, refer to a polymeric form of amino acids of any length, which can include coded and non-coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified peptide backbones. The term includes fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence, fusions with heterologous and homologous leader sequences, with or without N-terminal methionine residues; immunologically tagged proteins; and the like. NH_2 refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxyl group present at the carboxyl terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J. Biol. Chem.*, 243 (1969), 3552-59 is used.

[0028] As used herein, the terms "treatment," "treating," and the like, refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse affect attributable to the disease. "Treatment," as used herein, covers any treatment of a disease in a mammal, particularly in a human, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; and (c) relieving the disease, i.e., causing regression of the disease.

[0029] The terms "individual," "subject," "host," and "patient," used interchangeably herein, refer to a mammal, including, but not limited to, murines (rats, mice), non-human primates, humans, canines, felines, ungulates (e.g., equines, bovines, ovines, porcines, caprines), etc.

[0030] A "therapeutically effective amount" or "efficacious amount" refers to the amount of a compound, the amount of a recombinant expression vector, or a number of cells that, when administered to a mammal or other subject for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the expression vector, or the cell, the disease and its severity and the age, weight, etc., of the subject to be treated.

[0031] Before the present invention is further described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0032] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0033] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

[0034] It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a tau polypeptide” includes a plurality of such tau polypeptides and reference to “the GABAergic interneuron” includes reference to one or more GABAergic interneurons and equivalents thereof known to those skilled in the art, and so forth. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as “solely,” “only” and the like in connection with the recitation of claim elements, or use of a “negative” limitation.

[0035] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

DETAILED DESCRIPTION

[0036] The present disclosure provides a method of increasing the functionality of a GABAergic interneuron (an interneuron that produces γ -aminobutyric acid (GABA)) in the hilus of the hippocampus of an individual having at least one apolipoprotein E4 (apoE4) allele. The method generally involves reducing tau levels in the interneuron. The present disclosure provides a method of increasing cognitive function in an individual having at least one apoE4 allele.

[0037] In some embodiments, a subject method involves administering to an individual in need thereof an effective amount of an interfering nucleic acid that specifically reduces the level of tau polypeptide in a GABAergic interneuron. In other embodiments, a subject method involves administering to an individual in need thereof an effective number of stem cells that have been genetically modified to reduce the level of tau polypeptide produced by the stem cell, or by a neuron (e.g., a GABAergic neuron) generated from the stem cell.

[0038] In some embodiments, a subject method is effective to increase the number of GAD67-positive interneurons in the hippocampus (e.g., in the hilus of the dentate gyrus of the

hippocampus) of an individual by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, or more than 50%, compared with the number of GAD67-positive interneurons in the absence of treatment, or before treatment, with the method. In some embodiments, a GAD67-positive interneuron is also somatostatin-positive. In other embodiments, a GAD67-positive interneuron is also neuropeptide Y-positive.

[0039] Thus, for example, an effective amount of an interfering nucleic acid that reduces the level of tau polypeptide in a GABAergic interneuron is an amount that is effective, when administered in one or more doses, in monotherapy or in combination therapy, to increase the number of GAD67-positive interneurons in the hippocampus of an individual by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, or more than 50%, compared with the number of GAD67-positive interneurons in the absence of treatment, or before treatment, with the interfering nucleic acid.

[0040] As another example, an effective number of genetically modified stem cells (e.g., stem cells that have been genetically modified to reduce the level of tau polypeptide produced by the stem cell, or by a neuron (e.g., a GABAergic neuron) generated from the stem cell of genetically modified stem cell) is a number that is effective, when administered in one or more doses, in monotherapy or in combination therapy, to increase the number of GAD67-positive interneurons in the hippocampus of an individual by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, or more than 50%, compared with the number of GAD67-positive interneurons in the absence of treatment, or before treatment, with the genetically modified stem cells.

[0041] GAD67 (glutamic acid decarboxylase, 67 kDa isoform) has been described in the literature; see, e.g., Lariviere et al. (2002) *Mol. Biol. Evol.* 19:2325; GenBank Accession No. AAB26937; and Yamashita et al. (1993) *Biochem. Biophys. Res. Comm.* 192:1347.

[0042] In some embodiments, a subject method is effective to increase the functionality of a GABAergic interneuron in the hippocampus of an individual by at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 75%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 10-fold, or more than 10-fold, compared with the functionality of the GABAergic interneuron in the hippocampus of the individual in the absence of treatment, or before treatment, with the subject method. The functionality of a GABAergic interneuron includes basal GABA release, KCl-evoked GABA release, and neuregulin-evoked GABA release.

[0043] For example, in some embodiments, an effective amount of an interfering nucleic acid that reduces the level of tau polypeptide in a GABAergic interneuron is an amount that is effective, when administered in one or more doses, in monotherapy or in combination therapy (e.g., in combination with stem cell therapy or in combination therapy with at least one additional therapeutic agent), to increase the functionality of a GABAergic interneuron in the hippocampus of an individual by at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 75%, at least

about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 10-fold, or more than 10-fold, compared with the functionality of the GABAergic interneuron in the hippocampus of the individual in the absence of treatment, or before treatment, with the interfering nucleic acid.

[0044] As another example, an effective number of genetically modified stem cells (e.g., stem cells that have been genetically modified to reduce the level of tau polypeptide produced by the stem cell, or by a neuron (e.g., a GABAergic neuron) generated from the stem cell of genetically modified stem cell) is a number that is effective, when administered in one or more doses, in monotherapy or in combination therapy, to increase the functionality of a GABAergic interneuron in the hippocampus of an individual by at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 75%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 10-fold, or more than 10-fold, compared with the functionality of the GABAergic interneuron in the hippocampus of the individual in the absence of treatment, or before treatment, with the genetically modified stem cells.

[0045] In some embodiments, a subject method is effective to ameliorate at least one phenomenon associated with an apoE4-associated neurological disorder, where such phenomena include, e.g., neurofibrillary tangles; amyloid deposits; memory loss; and a reduction in cognitive function. Thus, for example, in some embodiments, a subject method is effective to reduce memory loss and at least slow the reduction in cognitive function. For example, in some embodiments, a subject method is effective to increase memory function and/or to increase cognitive function. Thus, for example, an effective amount of an interfering nucleic acid that reduces the level of tau polypeptide in a GABAergic interneuron is an amount that is effective, when administered in one or more doses, in monotherapy or in combination therapy, to reduce memory loss, to increase memory functions, to reduce loss of cognitive function, or to increase cognitive function. As another example, an effective number of genetically modified stem cells (e.g., stem cells that have been genetically modified to reduce the level of tau polypeptide produced by the stem cell, or by a neuron (e.g., a GABAergic neuron) generated from the stem cell of genetically modified stem cell) is a number that is effective, when administered in one or more doses, in monotherapy or in combination therapy, to reduce memory loss, to increase memory functions, to reduce loss of cognitive function, or to increase cognitive function.

Interfering Nucleic Acid

[0046] As noted above, in some embodiments, an interfering nucleic acid is used to interfere with production of tau transcripts and production of tau polypeptide. Interfering nucleic acids include small nucleic acid molecules, such as a short interfering nucleic acid (siNA), a short interfering RNA (siRNA), a double-stranded RNA (dsRNA), a micro-RNA (miRNA), and a short hairpin RNA (shRNA).

[0047] The terms “short interfering nucleic acid,” “siNA,” “short interfering RNA,” “siRNA,” “short interfering nucleic acid molecule,” “short interfering oligonucleotide molecule,” and “chemically-modified short interfering nucleic acid molecule” as used herein refer to any nucleic acid molecule capable of inhibiting or down regulating gene expression, for example by mediating RNA interference “RNAi” or gene silencing in a sequence-specific manner. Design of RNAi

molecules, given a target gene, is routine in the art. See also US 2005/0282188 (which is incorporated herein by reference) as well as references cited therein. See, e.g., Pushparaj et al. *Clin Exp Pharmacol Physiol*. 2006 May-June; 33(5-6): 504-10; Lutzelberger et al. *Handb Exp Pharmacol*. 2006; (173):243-59; Aronin et al. *Gene Ther*. 2006 March; 13(6): 509-16; Xie et al. *Drug Discov Today*. 2006 January; 11(1-2):67-73; Grunweller et al. *Curr Med. Chem*. 2005; 12(26): 3143-61; and Pekaraik et al. *Brain Res Bull*. 2005 Dec. 15; 68(1-2):115-20. Epub 2005 Sep. 9.

[0048] Methods for design and production of siRNAs to a desired target are known in the art, and their application to tau genes for the purposes disclosed herein will be readily apparent to the ordinarily skilled artisan, as are methods of production of siRNAs having modifications (e.g., chemical modifications) to provide for, e.g., enhanced stability, bioavailability, and other properties to enhance use as therapeutics. In addition, methods for formulation and delivery of siRNAs to a subject are also well known in the art. See, e.g., US 2005/0282188; US 2005/0239731; US 2005/0234232; US 2005/0176018; US 2005/0059817; US 2005/0020525; US 2004/0192626; US 2003/0073640; US 2002/0150936; US 2002/0142980; and US2002/0120129, each of which are incorporated herein by reference.

[0049] Publicly available tools to facilitate design of siRNAs are available in the art. See, e.g., DEQOR: Design and Quality Control of RNAi (available on the internet at cluster1.mpi-cbg.de/Deqor/deqor.html). See also, Henschel et al. *Nucleic Acids Res*. 2004 Jul. 1; 32 (Web Server issue):W113-20. DEQOR is a web-based program which uses a scoring system based on state-of-the-art parameters for siRNA design to evaluate the inhibitory potency of siRNAs. DEQOR, therefore, can help to predict (i) regions in a gene that show high silencing capacity based on the base pair composition and (ii) siRNAs with high silencing potential for chemical synthesis. In addition, each siRNA arising from the input query is evaluated for possible cross-silencing activities by performing BLAST searches against the transcriptome or genome of a selected organism. DEQOR can therefore predict the probability that an mRNA fragment will cross-react with other genes in the cell and helps researchers to design experiments to test the specificity of siRNAs or chemically designed siRNAs.

[0050] Suitable tau gene targets include, e.g., a contiguous stretch of from about 10 nucleotides (nt) to about 15 nt, from about 15 nt to about 20 nt, from about 20 nt to about 25 nt, from about 25 nt to about 30 nt, from about 30 nt to about 35 nt, from about 35 nt to about 40 nt, from about 40 nt to about 50 nt, from about 50 nt to about 60 nt, from about 60 nt to about 70 nt, from about 70 nt to about 80 nt, from about 80 nt to about 90 nt, or from about 90 nt to about 100 nt, of nucleotides 6547-6758 of SEQ ID NO:1 (tau isoform 1 mRNA; GenBank NM_016835), nucleotides 5596-5807 of SEQ ID NO:2 (tau isoform 2 mRNA; GenBank NM_005910); nucleotides 5596-5807 of SEQ ID NO:3 (tau isoform 3 mRNA; GenBank NM_016834), nucleotides 5329-5540 of SEQ ID NO:4 (tau isoform 4 mRNA; GenBank NM_016841), nucleotides 5509-5720 of SEQ ID NO:5 (tau isoform 5 mRNA; GenBank NM_001123067), or nucleotides 6601-6812 of SEQ ID NO:6 (tau isoform 6 mRNA; GenBank NM_001123066). SEQ ID NOs:1-6 provide nucleotide sequences of tau isoform 1-6 mRNA.

[0051] Suitable tau gene targets include, e.g., a contiguous stretch of from about 10 nucleotides (nt) to about 15 nt, from about 15 nt to about 20 nt, from about 20 nt to about 25 nt, from about 25 nt to about 30 nt, from about 30 nt to about 35 nt, from about 35 nt to about 40 nt, from about 40 nt to about 50 nt, from about 50 nt to about 60 nt, from about 60 nt to about 70 nt, from about 70 nt to about 80 nt, from about 80 nt to about 90 nt, or from about 90 nt to about 100 nt, of nucleotides 1-240 of tau isoform 1 mRNA (GenBank NM_016835). Nucleotides 1-240 of tau isoform 1 mRNA (GenBank NM_016835) are shown in FIG. 11B (SEQ ID NO:8).

[0052] Suitable tau gene targets include, e.g., nucleotides 6547-6758 of SEQ ID NO:1 (tau isoform 1 mRNA; GenBank NM_016835), nucleotides 5596-5807 of SEQ ID NO:2 (tau isoform 2 mRNA; GenBank NM_005910); nucleotides 5596-5807 of SEQ ID NO:3 (tau isoform 3 mRNA; GenBank NM_016834), nucleotides 5329-5540 of SEQ ID NO:4 (tau isoform 4 mRNA; GenBank NM_016841), nucleotides 5509-5720 of SEQ ID NO:5 (tau isoform 5 mRNA; GenBank NM_001123067), and nucleotides 6601-6812 of SEQ ID NO:6 (tau isoform 6 mRNA; GenBank NM_001123066). Suitable tau gene targets include, e.g., SEQ ID NO:7 (nucleotides 6547-6758 of tau1 mRNA); and SEQ ID NO:8 (nucleotides 1-240 of tau1 mRNA).

[0053] Other suitable target sequences will be readily apparent upon inspection of a sequence alignment of, e.g., SEQ ID NO:1 (tau isoform 1 mRNA; GenBank NM_016835), SEQ ID NO:2 (tau isoform 2 mRNA; GenBank NM_005910); SEQ ID NO:3 (tau isoform 3 mRNA; GenBank NM_016834), SEQ ID NO:4 (tau isoform 4 mRNA; GenBank NM_016841), SEQ ID NO:5 (tau isoform 5 mRNA; GenBank NM_001123067), and SEQ ID NO:6 (tau isoform 6 mRNA; GenBank NM_001123066).

[0054] It should be understood that the sequences provided above are the target sequences of the mRNAs encoding the target gene, and that the siRNA oligonucleotides used would comprise a sequence complementary to the target.

[0055] siNA molecules can be of any of a variety of forms. For example the siNA can be a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. siNA can also be assembled from two separate oligonucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary. In this embodiment, each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double stranded structure, for example wherein the double stranded region is about 15 to about 30, e.g., about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 base pairs; the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof (e.g., about 15 to about 25 or more nucleotides of the siNA molecule are complementary to the target nucleic acid or a portion thereof).

[0056] Alternatively, the siNA can be assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the siNA are linked by a nucleic acid-based or non-nucleic acid-based linker(s). The siNA can be a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof.

[0057] The siNA can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either in vivo or in vitro to generate an active siNA molecule capable of mediating RNAi. The siNA can also comprise a single stranded polynucleotide having nucleotide sequence complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof (e.g., where such siNA molecule does not require the presence within the siNA molecule of nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide can further comprise a terminal phosphate group, such as a 5'-phosphate (see for example Martinez et al., 2002, Cell., 110, 563-574 and Schwarz et al., 2002, Molecular Cell, 10, 537-568), or 5',3'-diphosphate.

[0058] In certain embodiments, the siNA molecule contains separate sense and antisense sequences or regions, wherein the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linker molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. In certain embodiments, the siNA molecules comprise nucleotide sequence that is complementary to nucleotide sequence of a target gene. In another embodiment, the siNA molecule interacts with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene.

[0059] As used herein, siNA molecules need not be limited to those molecules containing only RNA, but further encompass chemically-modified nucleotides and non-nucleotides. In certain embodiments, the short interfering nucleic acid molecules of the invention lack 2'-hydroxy (2'-OH) containing nucleotides. siNAs do not necessarily require the presence of nucleotides having a 2'-hydroxy group for mediating RNAi and as such, siNA molecules of the invention optionally do not include any ribonucleotides (e.g., nucleotides having a 2'-OH group). Such siNA molecules that do not require the presence of ribonucleotides within the siNA molecule to support RNAi can however have an attached linker or linkers or other attached or associated groups, moieties, or chains containing one or more nucleotides with 2'-OH groups. Optionally, siNA molecules can comprise ribonucleotides at about 5, 10, 20, 30, 40, or 50% of the nucleotide positions. The modified short interfering nucleic acid molecules of the invention can also be referred to as short interfering modified oligonucleotides "siMON."

[0060] As used herein, the term siNA is meant to be equivalent to other terms used to describe nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically-modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term RNAi is meant to be equivalent to other terms used to describe sequence specific RNA interference, such as post transcriptional gene silencing, translational inhibition, or epigenetics. For example, siNA molecules of the invention can be used to epigenetically silence a target gene at the post-transcriptional level or the pre-transcriptional level. In a non-limiting example, epigenetic regulation of gene expression by siNA molecules of the invention can result from siNA mediated modification of chromatin structure or methylation pattern to alter gene expression (see, for example, Verdel et al., 2004, *Science*, 303, 672-676; Pal-Bhadra et al., 2004, *Science*, 303, 669-672; Allshire, 2002, *Science*, 297, 1818-1819; Volpe et al., 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall et al., 2002, *Science*, 297, 2232-2237).

[0061] siNA molecules contemplated herein can comprise a duplex forming oligonucleotide (DFO) see, e.g., WO 05/019453; and US 2005/0233329, which are incorporated herein by reference). siNA molecules also contemplated herein include multifunctional siNA, (see, e.g., WO 05/019453 and US 2004/0249178). The multifunctional siNA can comprise sequence targeting, for example, two regions of tau.

[0062] siNA molecules contemplated herein can comprise an asymmetric hairpin or asymmetric duplex. By "asymmetric hairpin" as used herein is meant a linear siNA molecule comprising an antisense region, a loop portion that can comprise nucleotides or non-nucleotides, and a sense region that comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complementary nucleotides to base pair with the antisense region and form a duplex with loop. For example, an asymmetric hairpin siNA molecule can comprise an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g. about 15 to about 30, or about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides) and a loop region comprising about 4 to about 12 (e.g., about 4, 5, 6, 7, 8, 9, 10, 11, or 12) nucleotides, and a sense region having about 3 to about 25 (e.g., about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25) nucleotides that are complementary to the antisense region. The asymmetric hairpin siNA molecule can also comprise a 5'-terminal phosphate group that can be chemically modified. The loop portion of the asymmetric hairpin siNA molecule can comprise nucleotides, non-nucleotides, linker molecules, or conjugate molecules as described herein.

[0063] By "asymmetric duplex" as used herein is meant a siNA molecule having two separate strands comprising a sense region and an antisense region, wherein the sense region comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complementary nucleotides to base pair with the antisense region and form a duplex. For example, an asymmetric duplex siNA molecule of the invention can comprise an antisense region having length sufficient to mediate RNAi in a cell or in vitro system

(e.g. about 15 to about 30, or about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides) and a sense region having about 3 to about 25 (e.g., about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25) nucleotides that are complementary to the antisense region.

[0064] Stability and/or half-life of siRNAs can be improved through chemically synthesizing nucleic acid molecules with modifications (base, sugar and/or phosphate) can prevent their degradation by serum ribonucleases, which can increase their potency (see e.g., Eckstein et al., International Publication No. WO 92/07065; Perrault et al., 1990 *Nature* 344, 565; Pieken et al., 1991, *Science* 253, 314; Usman and Cedergren, 1992, *Trends in Biochem. Sci.* 17, 334; Usman et al., International Publication No. WO 93/15187; and Rossi et al., International Publication No. WO 91/03162; Sproat, U.S. Pat. No. 5,334,711; Gold et al., U.S. Pat. No. 6,300,074; and Burgin et al., supra; all of which are incorporated by reference herein, describing various chemical modifications that can be made to the base, phosphate and/or sugar moieties of the nucleic acid molecules described herein. Modifications that enhance their efficacy in cells, and removal of bases from nucleic acid molecules to shorten oligonucleotide synthesis times and reduce chemical requirements are desired.

[0065] For example, oligonucleotides are modified to enhance stability and/or enhance biological activity by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-O-allyl, 2'-H, nucleotide base modifications (for a review see Usman and Cedergren, 1992, *TIBS*. 17, 34; Usman et al., 1994, *Nucleic Acids Symp. Ser.* 31, 163; Burgin et al., 1996, *Biochemistry*, 35, 14090). Sugar modification of nucleic acid molecules have been extensively described in the art (see Eckstein et al., International Publication PCT No. WO 92/07065; Perrault et al. *Nature*, 1990, 344, 565-568; Pieken et al. *Science*, 1991, 253, 314-317; Usman and Cedergren, *Trends in Biochem. Sci.*, 1992, 17, 334-339; Usman et al. International Publication PCT No. WO 93/15187; Sproat, U.S. Pat. No. 5,334,711 and Beigelman et al., 1995, *J. Biol. Chem.*, 270, 25702; Beigelman et al., International PCT publication No. WO 97/26270; Beigelman et al., U.S. Pat. No. 5,716,824; Usman et al., U.S. Pat. No. 5,627,053; Woolf et al., International PCT Publication No. WO 98/13526; Thompson et al., U.S. Ser. No. 60/082,404 which was filed on Apr. 20, 1998; Karpeisky et al., 1998, *Tetrahedron Lett.*, 39, 1131; Eamshaw and Gait, 1998, *Biopolymers (Nucleic Acid Sciences)*, 48, 39-55; Verma and Eckstein, 1998, *Annu. Rev. Biochem.*, 67, 99-134; and Burlina et al., 1997, *Bioorg. Med. Chem.*, 5, 1999-2010; each of which are hereby incorporated in their totality by reference herein). In view of such teachings, similar modifications can be used as described herein to modify the siNA nucleic acid molecules of disclosed herein so long as the ability of siNA to promote RNAi in cells is not significantly inhibited.

[0066] Short interfering nucleic acid (siNA) molecules having chemical modifications that maintain or enhance activity are contemplated herein. Such a nucleic acid is also generally more resistant to nucleases than an unmodified nucleic acid. Accordingly, the in vitro and/or in vivo activity should not be significantly lowered. Nucleic acid molecules delivered exogenously are generally selected to be stable within cells at least for a period sufficient for transcription and/or translation of the target RNA to occur and to provide

for modulation of production of the encoded mRNA and/or polypeptide so as to facilitate reduction of the level of the target gene product.

[0067] Production of RNA and DNA molecules can be accomplished synthetically and can provide for introduction of nucleotide modifications to provide for enhanced nuclease stability. (see, e.g., Wincott et al., 1995, *Nucleic Acids Res.* 23, 2677; Caruthers et al., 1992, *Methods in Enzymology* 211, 3-19, incorporated by reference herein. In one embodiment, nucleic acid molecules of the invention include one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) G-clamp nucleotides, which are modified cytosine analogs which confer the ability to hydrogen bond both Watson-Crick and Hoogsteen faces of a complementary guanine within a duplex, and can provide for enhanced affinity and specificity to nucleic acid targets (see, e.g., Lin et al. 1998, *J. Am. Chem. Soc.*, 120, 8531-8532). In another example, nucleic acid molecules can include one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) LNA "locked nucleic acid" nucleotides such as a 2',4'-C methylene bicyclo nucleotide (see, e.g., Wengel et al., WO 00/66604 and WO 99/14226).

[0068] siNA molecules can be provided as conjugates and/or complexes, e.g., to facilitate delivery of siNA molecules into a cell. Exemplary conjugates and/or complexes include those composed of an siNA and a small molecule, lipid, cholesterol, phospholipid, nucleoside, antibody, toxin, negatively charged polymer (e.g., protein, peptide, hormone, carbohydrate, polyethylene glycol, or polyamine). In general, the transporters described are designed to be used either individually or as part of a multi-component system, with or without degradable linkers. These compounds can improve delivery and/or localization of nucleic acid molecules into cells in the presence or absence of serum (see, e.g., U.S. Pat. No. 5,854,038). Conjugates of the molecules described herein can be attached to biologically active molecules via linkers that are biodegradable, such as biodegradable nucleic acid linker molecules.

[0069] Interfering RNAs may be generated exogenously by chemical synthesis, by *in vitro* transcription, or by cleavage of longer double-stranded RNA with dicer or another appropriate nuclease with similar activity. Chemically synthesized interfering RNAs, produced from protected ribonucleoside phosphoramidites using a conventional DNA/RNA synthesizer, may be obtained from commercial suppliers such as Ambion Inc. (Austin, Tex.), Invitrogen (Carlsbad, Calif.), or Dharmacon (Lafayette, Colo.).

[0070] Interfering RNAs are purified by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof, for example. Alternatively, interfering RNA may be used with little if any purification to avoid losses due to sample processing.

[0071] Interfering RNAs can also be expressed endogenously from plasmid or viral expression vectors or from minimal expression cassettes, for example, polymerase chain reaction (PCR)-generated fragments comprising one or more promoters and an appropriate template or templates for the interfering RNA. Examples of commercially available plasmid-based expression vectors for shRNA include members of the pSilencer series (Ambion, Austin, Tex.) and pCpG-siRNA (InvivoGen, San Diego, Calif.). Viral vectors for expression of interfering RNA may be derived from a variety of viruses including adenovirus, adeno-associated virus, lentivirus (e.g., HIV, FIV, and EIAV), and herpes virus. Examples of commercially available viral vectors for shRNA expression

include pSilencer adeno (Ambion, Austin, Tex.) and pLenti6/BLOCK-iTTM-DEST (Invitrogen, Carlsbad, Calif.). Selection of viral vectors, methods for expressing the interfering RNA from the vector and methods of delivering the viral vector are within the ordinary skill of one in the art. Examples of kits for production of PCR-generated shRNA expression cassettes include Silencer Express (Ambion, Austin, Tex.) and siXpress (Minus, Madison, Wis.).

[0072] An interfering RNA can be delivered in a delivery system that provides tissue targetable delivery. In addition, a suitable formulation for an interfering nucleic acid can include one or more additional properties: 1) nucleic acid binding into a core that can release the siRNA into the cytoplasm; 2) protection from non-specific interactions; 3) and tissue targeting that provides cell uptake. In some embodiments, the composition comprises a modular polymer conjugate targeting hippocampal neurons (e.g., interneurons) by coupling a peptide ligand specific for those cells to one end of a protective polymer, coupled at its other end to a cationic carrier for nucleic acids. For example, a suitable polymer conjugate can have three functional domains: peptide ligand specific for a target cell; protective polymer; and cationic carrier for nucleic acids. Another suitable formulation includes surface coatings attached to a preformed nanoparticle.

[0073] Suitable formulations for delivery of an interfering nucleic acid include polymers, polymer conjugates, lipids, micelles, self-assembly colloids, nanoparticles, sterically stabilized nanoparticles, and ligand-directed nanoparticles.

Recombinant Expression Vector

[0074] In some embodiments, a subject method involves administering to an individual in need thereof an effective amount of a recombinant expression vector that provides for production of a nucleic acid that reduces the level of tau polypeptide in a GABAergic interneuron, e.g., a recombinant expression vector comprising a nucleotide sequence that encodes an interfering nucleic acid that selectively reduces the level of a tau polypeptide in a cell that produces tau. Thus, in some embodiments, a recombinant expression vector is administered to an individual in need thereof, where the recombinant expression vector comprises a nucleotide sequence encoding an interfering RNA that specifically reduces a tau transcript and/or polypeptide in a cell (e.g., in a GABAergic interneuron). In some embodiments, the nucleotide sequence encoding an interfering RNA that specifically reduces a tau transcript and/or polypeptide in a cell is operably linked to a transcriptional control element (e.g., a promoter) that is active in a GABAergic interneuron.

[0075] Expression vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences encoding heterologous proteins. A selectable marker operative in the expression host may be present. Suitable expression vectors include, but are not limited to, viral vectors (e.g. viral vectors based on vaccinia virus; poliovirus; adenovirus (see, e.g., Li et al., *Invest Ophthalmol Vis Sci* 35:2543 2549, 1994; Borrás et al., *Gene Ther* 6:515 524, 1999; Li and Davidson, *PNAS* 92:7700 7704, 1995; Sakamoto et al., *H Gene Ther* 5:1088 1097, 1999; WO 94/12649, WO 93/03769; WO 93/19191; WO 94/28938; WO 95/11984 and WO 95/00655); adeno-associated virus (see, e.g., Ali et al., *Hum Gene Ther* 9:8186, 1998, Flannery et al., *PNAS* 94:6916 6921, 1997; Bennett et al., *Invest Ophthalmol Vis Sci* 38:2857 2863, 1997; Jomary et

al., *Gene Ther* 4:683-690, 1997; Rolling et al., *Hum Gene Ther* 10:641-648, 1999; Ali et al., *Hum Mol Genet.* 5:591-594, 1996; Srivastava in WO 93/09239, Samulski et al., *J. Vir.* (1989) 63:3822-3828; Mendelson et al., *Virol.* (1988) 166:154-165; and Flotte et al., *PNAS* (1993) 90:10613-10617); SV40; herpes simplex virus; human immunodeficiency virus (see, e.g., Miyoshi et al., *PNAS* 94:10319-23, 1997; Takahashi et al., *J Virol* 73:7812-7816, 1999); a retroviral vector (e.g., Murine Leukemia Virus, spleen necrosis virus, and vectors derived from retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, human immunodeficiency virus, myeloproliferative sarcoma virus, and mammary tumor virus); and the like.

[0076] Depending on the host/vector system utilized, any of a number of suitable transcription and translation control elements, including constitutive and inducible promoters, transcription enhancer elements, transcription terminators, etc. may be used in the expression vector (see e.g., Bitter et al. (1987) *Methods in Enzymology*, 153:516-544).

[0077] Non-limiting examples of suitable eukaryotic promoters (promoters functional in a eukaryotic cell) include cytomegalovirus (CMV) immediate early, herpes simplex virus (HSV) thymidine kinase, early and late SV40, long terminal repeats (LTRs) from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art. The expression vector may also contain a ribosome binding site for translation initiation and a transcription terminator. The expression vector may also include appropriate sequences for amplifying expression.

[0078] A recombinant vector will in some embodiments include one or more selectable markers. In addition, the expression vectors will in many embodiments contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture.

[0079] Other gene delivery vehicles and methods may be employed, including polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example Curiel (1992) *Hum. Gene Ther.* 3:147-154; ligand linked DNA, for example see Wu (1989) *J. Biol. Chem.* 264:16985-16987; eukaryotic cell delivery vehicles cells; deposition of photopolymerized hydrogel materials; hand-held gene transfer particle gun, as described in U.S. Pat. No. 5,149,655; ionizing radiation as described in U.S. Pat. No. 5,206,152 and in WO 92/11033; nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip (1994) *Mol. Cell. Biol.* 14:2411-2418, and in Woffendin (1994) *Proc. Natl. Acad. Sci.* 91:1581-1585.

[0080] Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and U.S. Pat. No. 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm. Liposomes that can act as gene delivery vehicles are described in U.S. Pat. No. 5,422,120, PCT Nos. WO 95/13796, WO 94/23697, and WO 91/14445, and EP No. 524 968.

[0081] Liposome or lipid nucleic acid delivery vehicles can also be used. Liposome complexes for gene delivery are described in, e.g., U.S. Pat. No. 7,001,614. For example,

liposomes comprising DOTAP and at least one cholesterol and/or cholesterol-derivative, present in a molar ratio range of 2.0 mM to 10 mM provide an effective delivery system, e.g., where the molar ratio of DOTAP to cholesterol is 1:1.3:1. The cationic lipid N-[(2,3-dioleoyloxy)propyl]-L-lysineamide (LADOP) can be used in a composition for delivering a polynucleotide, where LADOP-containing liposomes are described in, e.g., U.S. Pat. No. 7,067,697. Liposome formulations comprising amphipathic lipids having a polar head-group and aliphatic components capable of promoting transfection are suitable for use and are described in, e.g., U.S. Pat. No. 6,433,017. Lipid-conjugated polyamide compounds can be used to deliver nucleic acid; see, e.g., U.S. Pat. No. 7,214,384.

[0082] Suitable synthetic polymer-based carrier vehicles are described in, e.g., U.S. Pat. No. 6,312,727. Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:11581-11585. Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of hand-held gene transfer particle gun, as described in U.S. Pat. No. 5,149,655; use of ionizing radiation for activating transferred gene, as described in U.S. Pat. No. 5,206,152 and PCT No. WO 92/11033.

Genetically Modified Stem Cells

[0083] A stem cell used in a subject method is genetically modified in such a way as to reduce the level of tau polypeptide produced by the stem cell or a progeny of the stem cell. A parent (or "host") stem cell is genetically modified with an exogenous nucleic acid. The exogenous nucleic acid will in some embodiments comprise a nucleotide sequence that has sufficient homology to an endogenous tau polypeptide-encoding nucleic acid such that the exogenous nucleic acid will undergo homologous recombination with the endogenous tau polypeptide-encoding nucleic acid and will functionally disable the endogenous tau polypeptide-encoding nucleic acid. The term "functionally disabled," as used herein, refers to a genetic modification of a nucleic acid, which modification results in production of a gene product encoded by the nucleic acid that is produced at below normal levels, and/or is non-functional.

[0084] In some embodiments, the endogenous tau gene of a genetically modified stem cell is deleted. Any method for deleting a gene can be used. For example, a recombinase-mediated knockout method can be used, e.g., using a Cre/Lox system (the Cre/lox site-specific recombination system known in the art employs the bacteriophage P1 protein Cre recombinase and its recognition sequence loxP; see Rajewsky et al., *J. Clin. Invest.*, 98:600-603 (1996); Sauer, *Methods*, 14:381-392 (1998); Gu et al., *Cell*, 73:1155-1164 (1993); Araki et al., *Proc. Natl. Acad. Sci. USA*, 92:160-164 (1995); Lakso et al., *Proc. Natl. Acad. Sci. USA*, 89:6232-6236 (1992); and Orban et al., *Proc. Natl. Acad. Sci. USA*, 89:6861-6865 (1992)); a FLP/FRT recombination system (see, e.g., Brand and Perrimon, 1993, *Development* 118:401-415); and the like. As another example, a deletion-based conditional knockout method can be used. As another example, e.g., as described in U.S. Pat. No. 7,625,755, an inducible gene silencer comprising: a splice acceptor

sequence; an internal ribosomal entry site (IRES) sequence; a nucleotide sequence coding for a reporter protein; a polyadenylation sequence; and a pair of oppositely oriented recombination site sequences, which cause single cycle inversions in the presence of a suitable recombinase enzyme, flanking the aforementioned elements, can be used.

[0085] Alternatively, mutations that can cause reduced expression level (e.g., reduced transcription and/or translation efficiency, and decreased mRNA stability) of an endogenous tau-encoding nucleic acid may also be introduced into an endogenous tau gene by homologous recombination.

[0086] In addition, cells that have been genetically altered with recombinant genes or by antisense technology, to provide a gain or loss of genetic function, may be utilized with the invention. Methods for generating genetically modified cells are known in the art, see for example “Current Protocols in Molecular Biology”, Ausubel et al., eds, John Wiley & Sons, New York, N.Y., 2000. The genetic alteration may be a knock-out, usually where homologous recombination results in a deletion that knocks out expression of a targeted gene; or a knock-in, where a genetic sequence not normally present in the cell is stably introduced.

[0087] A variety of methods can be used to achieve a knock-out, including site-specific recombination, expression of anti-sense or dominant negative mutations, and the like. Knockouts have a partial or complete loss of function in one or both alleles of the endogenous gene in the case of gene targeting. In some embodiments, expression of the targeted gene product is undetectable or insignificant in the cells being analyzed; this may be achieved by introduction of a disruption of the coding sequence, e.g. insertion of one or more stop codons, insertion of a DNA fragment, etc., deletion of coding sequence, substitution of stop codons for coding sequence, etc. In some cases the introduced sequences are ultimately deleted from the genome, leaving a net change to the native sequence.

[0088] Different approaches may be used to achieve the “knock-out”. A chromosomal deletion of all or part of the native gene may be induced, including deletions of the non-coding regions, particularly the promoter region, 3' regulatory sequences, enhancers, or deletions of gene that activate expression of the targeted genes. A functional knock-out may also be achieved by the introduction of an anti-sense construct that blocks expression of the native genes (for example, see Li and Cohen (1996) *Cell* 85:319-329). “Knock-outs” also include conditional knock-outs, for example where alteration of the target gene occurs upon exposure of the animal to a substance that promotes target gene alteration, introduction of an enzyme that promotes recombination at the target gene site (e.g. Cre in the Cre-lox system), or other method for directing the target gene alteration.

[0089] The genetic construct may be introduced into tissues or host cells by any number of routes, including calcium phosphate transfection, viral infection, microinjection, electroporation or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth et al. (1992), *Anal. Biochem.* 205:365-368. The DNA may be coated onto gold microparticles, and delivered intradermally by a particle bombardment device, or “gene gun” as described in the literature (see, for example, Tang et al. (1992), *Nature* 356:152-154), where gold microprojectiles are coated with the DNA, then bombarded into cells.

[0090] A number of selection systems can be used for introducing the genetic changes, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, *Cell* 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, *Proc. Natl. Acad. Sci. USA* 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, *Cell* 22:817) genes can be employed in tk⁻; hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, *Natl. Acad. Sci. USA* 77:3567; O'Hare, et al., 1981, *Proc. Natl. Acad. Sci. USA* 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, *Proc. Natl. Acad. Sci. USA* 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, *J. Mol. Biol.* 150:1); and hygro, which confers resistance to hygromycin (Santerre, et al., 1984, *Gene* 30:147).

[0091] The stem cells used for transplantation can be allogeneic, autologous, or xenogeneic, relative to the individual being treated (e.g., the recipient individual into whom the stem cells are being transplanted). For example, in some cases, the stem cells (e.g., NSC or iNSC) are obtained from a human donor individual who is the same as the human individual being treated (the recipient). As another example, in some cases the stem cells (e.g., NSC or iNSC) are obtained from a human donor individual who is other than the human individual being treated (the recipient).

[0092] Neural stem cells of various species have been described. See, e.g., WO 93/01275, WO 94/09119, WO 94/10292, WO 94/16718, and Cattaneo et al., *Mol. Brain Res.*, 42, pp. 161-66 (1996). In some embodiments, NSCs, when maintained in certain culture conditions (e.g., a mitogen-containing (e.g., epidermal growth factor or epidermal growth factor plus basic fibroblast growth factor), serum-free culture medium), grow in suspension culture to form aggregates of cells known as “neuro spheres.”

[0093] NSCs can be generated from somatic cells (where the NSCs are referred to as “induced NSCs”); pluripotent stem cells; induced pluripotent stem cells (iPS); or fetal or adult tissue that contains NSCs. Suitable tissue sources of NSCs include, but are not limited to, hippocampus, septal nuclei, cortex, cerebellum, ventral mesencephalon, and spinal cord.

[0094] A suitable NSC exhibits one or more of the following properties: 1) expression of Nestin; 2) expression of Sox2; 3) expression of Musashil; 4) ability to undergo self-renewal, either as a monolayer or in suspension cultures as neurospheres; 5) ability to differentiate into neurons, specific subtypes of neurons, astrocytes, and oligodendrocytes; and 6) morphological characteristics typical for NSCs. A suitable iNSC can also express CD133 and Vimentin. Nestin, Sox2, and Musashil are well described in the literature as hallmark genes expressed in NSCs. See, e.g., GenBank Accession Nos. NP_006608, CAA46780, and CAI16338 for Nestin. For Musashil, see, e.g., GenBank Accession No. BAB69769; and Shu et al. (2002) *Biochem. Biophys. Res. Comm.* 293:150.

[0095] A suitable NSC is generally negative for markers that identify mature neurons, astrocytes, and oligodendrocytes. Thus, e.g., a suitable NSC is generally microtubule-associated protein-2 (MAP2) negative, neuron-specific nuclear protein (NeuN) negative, Tau negative, S10013 negative, oligodendrocyte marker O4 negative, and oligodendrocyte lineage transcription factor Olig2 negative. These mark-

ers of mature neural markers are well described in the literature. For MAP2, see, e.g., GenBank Accession Nos. AAA59552, AAB48098, AAI43246, and AAH38857. For NeuN, see, e.g., Wolf et al. (1996) *J. Histochem. & Cytochem.* 44:1167. For S10013, see, e.g., GenBank Accession Nos. NP_006263.1 (*H. sapiens* S10013); NP_033141 (*Mus musculus* S100β); CAG46920.1 (*Homo sapiens* S100β); and see also, Allore et al. (1990) *J. Biol. Chem.* 265:15537. For O4, see, e.g., Schachner et al. (1981) *Dev. Biol.* 83:328; Bansal et al. (1989) *J. Neurosci. Res.* 24:548; and Bansal and Pfeiffer (1989) *Proc. Natl. Acad. Sci. USA* 86:6181. For Olig2, see, e.g., Lu et al. (2001) *Proc. Natl. Acad. Sci. USA* 98:10851; Ligon et al. (2004) *J. Neuropathol. Exp. Neurol.* 63:499.

Tissue Sources

[0096] Suitable tissue sources of neural stem cells include the CNS, including the cerebral cortex, cerebellum, midbrain, brainstem, spinal cord and ventricular tissue; and areas of the peripheral nervous system (PNS) including the carotid body and the adrenal medulla. Exemplary areas include regions in the basal ganglia, e.g., the striatum which consists of the caudate and putamen, or various cell groups, such as the globus pallidus, the subthalamic nucleus, the nucleus basalis, or the substantia nigra pars compacta. In some embodiments, the neural tissue is obtained from ventricular tissue that is found lining CNS ventricles (e.g., lateral ventricles, third ventricle, fourth ventricle, central canal, cerebral aqueduct, etc.) and includes the subependyma.

[0097] Non-autologous human neural stem cells can be derived from fetal tissue following elective abortion, or from a post-natal, juvenile or adult organ donor. Autologous neural tissue can be obtained by biopsy, or from patients undergoing neurosurgery in which neural tissue is removed, for example, during epilepsy surgery, temporal lobectomies and hippocampalectomies. Neural stem cells have been isolated from a variety of adult CNS ventricular regions, including the frontal lobe, conus medullaris, thoracic spinal cord, brain stem, and hypothalamus. In each of these cases, the neural stem cell exhibits self-maintenance and generates a large number of progeny which include neurons, astrocytes and oligodendrocytes.

Induced NSCs

[0098] Suitable NSCs include induced NSCs (iNSCs). An iNSC can be generated by introducing into a somatic cell one or more of: an exogenous Sox2 polypeptide, an Oct-3/4 polypeptide, an exogenous c-Myc polypeptide, an exogenous Klf4 polypeptide, an exogenous Nanog polypeptide, and an exogenous Lin28 polypeptide.

[0099] Sox2 polypeptides, Oct-3/4 polypeptides, c-Myc polypeptides, and Klf4 polypeptides, are known in the art and are described in, e.g., U.S. Patent Publication No. 2009/0191159. Nanog polypeptides and Lin28 polypeptides are known in the art and are described in, e.g., U.S. Patent Publication No. 2009/0047263. See also the following GenBank Accession Nos.: 1) GenBank Accession Nos. NP_002692, NP_001108427; NP_001093427; NP_001009178; and NP_038661 for Oct-3/4; 2) GenBank Accession Nos. NP_004226, NP_001017280, NP_057354, AAP36222, NP_034767, and NP_446165 for Klf4 and Klf4 family members; 3) GenBank Accession Nos. NP_002458, NP_001005154, NP_036735, NP_034979, P0C0N9, and

NP_001026123 for c-Myc; 4) GenBank Accession Nos. AAP49529 and BAC76999, for Nanog; 5) GenBank Accession Nos. AAH28566 and NP_078950, for Lin28; and 6) GenBank Accession Nos. NP_003097, NP_001098933, NP_035573, ACA58281, BAA09168, NP_001032751, and NP_648694 for Sox2 amino acid sequences.

[0100] A multipotent iNSC can be induced from a wide variety of mammalian somatic cells. Examples of suitable mammalian cells include, but are not limited to: fibroblasts (including dermal fibroblasts, human foreskin fibroblasts, etc.), bone marrow-derived mononuclear cells, skeletal muscle cells, adipose cells, peripheral blood mononuclear cells, macrophages, hepatocytes, keratinocytes, oral keratinocytes, hair follicle dermal cells, gastric epithelial cells, lung epithelial cells, synovial cells, kidney cells, skin epithelial cells, and osteoblasts.

[0101] A somatic cell can also originate from many different types of tissue, e.g., bone marrow, skin (e.g., dermis, epidermis), muscle, adipose tissue, peripheral blood, foreskin, skeletal muscle, or smooth muscle. The cells can also be derived from neonatal tissue, including, but not limited to: umbilical cord tissues (e.g., the umbilical cord, cord blood, cord blood vessels), the amnion, the placenta, or other various neonatal tissues (e.g., bone marrow fluid, muscle, adipose tissue, peripheral blood, skin, skeletal muscle etc).

[0102] A somatic cell can be obtained from any of a variety of mammals, including, e.g., humans, non-human primates, murines (e.g., mice, rats), ungulates (e.g., bovines, equines, ovines, caprines, etc.), felines, canines, etc.

[0103] A somatic cell can be derived from neonatal or post-natal tissue collected from a subject within the period from birth, including cesarean birth, to death. For example, the tissue may be from a subject who is >10 minutes old, >1 hour old, >1 day old, >1 month old, >2 months old, >6 months old, >1 year old, >2 years old, >5 years old, >10 years old, >15 years old, >18 years old, >25 years old, >35 years old, >45 years old, >55 years old, >65 years old, >80 years old, <80 years old, <70 years old, <60 years old, <50 years old, <40 years old, <30 years old, <20 years old or <10 years old. The subject may be a neonatal infant. In some cases, the subject is a child or an adult. In some examples, the tissue is from a human of age 2, 5, 10 or 20 hours. In other examples, the tissue is from a human of age 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 9 months or 12 months. In some cases, the tissue is from a human of age 1 year, 2 years, 3 years, 4 years, 5 years, 18 years, 20 years, 21 years, 23 years, 24 years, 25 years, 28 years, 29 years, 31 years, 33 years, 34 years, 35 years, 37 years, 38 years, 40 years, 41 years, 42 years, 43 years, 44 years, 47 years, 51 years, 55 years, 61 years, 63 years, 65 years, 70 years, 77 years, or 85 years old.

[0104] The cells can be from non-embryonic tissue, e.g., at a stage of development later than the embryonic stage. In other cases, the cells may be derived from an embryo. In some cases, the cells may be from tissue at a stage of development later than the fetal stage. In other cases, the cells may be derived from a fetus.

[0105] The cells to be induced or reprogrammed can be obtained from a single cell or a population of cells. The population may be homogeneous or heterogeneous. The cells can be a population of cells found in a human cellular sample, e.g., a biopsy or blood sample.

[0106] Methods for obtaining human somatic cells are well established, as described in, e.g., Schantz and Ng (2004), *A Manual for Primary Human Cell Culture*, World Scientific

Publishing Co., Pte, Ltd. In some cases, the methods include obtaining a cellular sample, e.g., by a biopsy (e.g., a skin sample), blood draw, or alveolar or other pulmonary lavage. It is to be understood that initial plating densities of cells prepared from a tissue can vary, due to a variety of factors, e.g., expected viability or adherence of cells from that particular tissue.

[0107] An exogenous polypeptide can be introduced into a somatic cell by contacting the somatic cell with the exogenous polypeptide (e.g., a Sox2 polypeptide, as described above) wherein the exogenous polypeptide is taken up into the cell.

[0108] In some embodiments, an exogenous polypeptide (e.g., a Sox2 polypeptide) comprises a protein transduction domain, e.g., an exogenous polypeptide is linked, covalently or non-covalently, to a protein transduction domain.

[0109] "Protein Transduction Domain" or PTD refers to a polypeptide, polynucleotide, carbohydrate, or organic or inorganic compound that facilitates traversing a lipid bilayer, micelle, cell membrane, organelle membrane, or vesicle membrane. A PTD attached to another molecule facilitates the molecule traversing a membrane, for example going from extracellular space to intracellular space, or cytosol to within an organelle. In some embodiments, a PTD is covalently linked to the amino terminus of an exogenous polypeptide (e.g., a Sox2 polypeptide). In some embodiments, a PTD is covalently linked to the carboxyl terminus of an exogenous polypeptide (e.g., a Sox2 polypeptide). Exemplary protein transduction domains include but are not limited to a minimal undecapeptide protein transduction domain (corresponding to residues 47-57 of HIV-1 TAT comprising YGRKKRRQRRR; SEQ ID NO:9); a polyarginine sequence comprising a number of arginines sufficient to direct entry into a cell (e.g., 3, 4, 5, 6, 7, 8, 9, 10, or 10-50 arginines); a VP22 domain (Zender et al., Cancer Gene Ther. 2002 June; 9(6):489-96); an *Drosophila* Antennapedia protein transduction domain (Noguchi et al., Diabetes 2003; 52(7):1732-1737); a truncated human calcitonin peptide (Trehin et al. Pharm. Research, 21:1248-1256, 2004); polylysine (Wender et al., PNAS, Vol. 97:13003-13008); RRQRRTSKLMKR (SEQ ID NO:10); Transportan GWTLNSAGYLLGKINL-KALAALAKKIL (SEQ ID NO:11); KALAWREAK-LAKALAKALAKHLAKALAKALKCEA (SEQ ID NO:12); and RQIKIWFQNRRMKWKK (SEQ ID NO:13). Exemplary PTDs include but are not limited to, YGRKKRRQRRR (SEQ ID NO:9), RKKRRQRRR (SEQ ID NO:14); an arginine homopolymer of from 3 arginine residues to 50 arginine residues; Exemplary PTD domain amino acid sequences include, but are not limited to, any of the following:

YGRKKRRQRRR;	(SEQ ID NO: 9)
RKKRRQRRR;	(SEQ ID NO: 14)
YARAAARQARA;	(SEQ ID NO: 15)
THRLPRRRRRR;	(SEQ ID NO: 16)
and	
GGRRARRRRRR.	(SEQ ID NO: 17)

[0110] In some embodiments, introduction of an exogenous polypeptide (e.g., an exogenous Sox2 polypeptide) into a somatic cell is achieved by genetic modification of the somatic cell with an exogenous nucleic acid comprising a

nucleotide sequence encoding the polypeptide. Exogenous nucleic acids include a recombinant expression vector comprising a nucleotide sequence encoding an exogenous polypeptide (e.g., an exogenous Sox2 polypeptide). Suitable recombinant expression vectors include plasmids, as well as viral-based expression vectors, e.g., lentivirus vectors, adenovirus vectors, adeno-associated virus vectors, etc., which are well known in the art.

iPS Cells

[0111] In some embodiments, NSCs are generated from induced pluripotent stem (iPS) cells. iPS cells are generated from somatic cells, including skin fibroblasts, using, e.g., known methods. iPS cells produce and express on their cell surface one or more of the following cell surface antigens: SSEA-3, SSEA-4, TRA-1-60, TRA-1-81, TRA-2-49/6E, and Nanog. In some embodiments, iPS cells produce and express on their cell surface SSEA-3, SSEA-4, TRA-1-60, TRA-1-81, TRA-2-49/6E, and Nanog. iPS cells express one or more of the following genes: Oct-3/4, Sox2, Nanog, GDF3, REX1, FGF4, ESG1, DPPA2, DPPA4, and hTERT. In some embodiments, an iPS cell expresses Oct-3/4, Sox2, Nanog, GDF3, REX1, FGF4, ESG1, DPPA2, DPPA4, and hTERT. iPS can be induced to differentiate into neural cells that express one or more of: β III-tubulin, tyrosine hydroxylase, AADC, DAT, ChAT, LMX1B, and MAP2. Methods of generating iPS are known in the art, and any such method can be used to generate iPS. See, e.g., Takahashi and Yamanaka (2006) Cell 126:663-676; Yamanaka et al. (2007) Nature 448:313-7; Wernig et al. (2007) Nature 448:318-24; Maherali (2007) Cell Stem Cell 1:55-70.

[0112] iPS cells can be generated from somatic cells (e.g., skin fibroblasts) by genetically modifying the somatic cells with one or more expression constructs encoding Oct-3/4 and Sox2. In some embodiments, somatic cells are genetically modified with one or more expression constructs comprising nucleotide sequences encoding Oct-3/4, Sox2, c-myc, and Klf4. In some embodiments, somatic cells are genetically modified with one or more expression constructs comprising nucleotide sequences encoding Oct-4, Sox2, Nanog, and LIN28.

[0113] iPS cells can be induced to differentiate into neural cells using any of a variety of published protocols (see, e.g., Muotri et al., 2005, Proc. Natl. Acad. Sci. USA. 102:18644; Takahashi et al, 2007, Cell 131:861). For example, in some embodiments, iPS cells are cultured on mitotically inactivated (e.g., mitomycin C-treated or irradiated) mouse embryonic fibroblasts (Specialty Media, Lavellette, N.J.) in DMEM/F12 Glutamax (GIBCO), 20% knockout serum replacement (GIBCO), 0.1 mM nonessential amino acids (GIBCO), 0.1 mM 2-mercaptoethanol (GIBCO), and 4 ng/ml bFGF-2 (R & D Systems). iPS cell neuronal differentiation can be induced by coculturing the iPS cells with PA6 cells for 3-5 weeks under the following differentiation conditions: DMEM/F12 Glutamax (GIBCO), 10% knockout serum replacement (GIBCO), 0.1 mM nonessential amino acids (GIBCO), and 0.1 mM 2-mercaptoethanol (GIBCO). Alkaline phosphatase activity can be measured using the Vector Red Alkaline Phosphatase substrate kit I from Vector Laboratories. Neuronal differentiation can be monitored by immunostaining with various neuronal cell markers.

Combination Therapies

[0114] In some embodiments, a subject method further includes administering at least one additional therapeutic agent. Suitable additional therapeutic agents include, but are not limited to, acetylcholinesterase inhibitors, including, but not limited to, Aricept (donepezil), Exelon (rivastigmine), metrifonate, and tacrine (Cognex); non-steroidal anti-inflammatory agents, including, but not limited to, ibuprofen and indomethacin; cyclooxygenase-2 (Cox2) inhibitors such as Celebrex; and monoamine oxidase inhibitors, such as Selegiline (Eldepryl or Deprenyl). Dosages for each of the above agents are known in the art. For example, Aricept is generally administered at 50 mg orally per day for 6 weeks, and, if well tolerated by the individual, at 10 mg per day thereafter.

[0115] Another suitable additional therapeutic agent is an apoE4 “structure corrector” that reduces apoE4 domain interaction. Agents that reduce apoE4 domain interaction include, e.g., an agent as described in U.S. Patent Publication No. 2006/0073104); and in Ye et al. (2005) *Proc. Natl. Acad. Sci. USA* 102:18700.

[0116] Another suitable additional therapeutic agent is an agent that inhibits tau aggregation, e.g., a naphthoquinone derivative that inhibits tau aggregation, as described in U.S. Pat. No. 7,605,179. Another suitable additional therapeutic agent is an agent that inhibits phosphorylation of tau, e.g., a 3-substituted-4-pyrimidone derivative that inhibits tau protein kinase 1, as described in U.S. Pat. No. 7,572,793.

Formulations, Dosages, and Routes of Delivery

[0117] An agent active agent (e.g., an interfering nucleic acid; a recombinant expression vector; a population of genetically modified stem cells; at least a second therapeutic agent) can be provided together with a pharmaceutically acceptable excipient. Pharmaceutically acceptable excipients are known to those skilled in the art, and have been amply described in a variety of publications, including, for example, A. Gennaro (1995) “Remington: The Science and Practice of Pharmacy”, 19th edition, Lippincott, Williams, & Wilkins. In the discussion, below, of formulations, dosages, and routes of delivery, an “active agent” will refer to an agent discussed herein, e.g., a recombinant expression vector, a population of genetically modified stem cells, or at least a second therapeutic agent, unless otherwise specified.

Nucleic Acids

[0118] Nucleic acids can be formulated in a variety of ways in order to facilitate delivery to the surface of the intestinal cells. The form (e.g., liquid, solid, pill, capsule) and composition of the formulation will vary according to the method of administration used. For example, where the formulation is administered orally, the nucleic acid can be formulated as a tablet, pill, capsule, solution (e.g., gel, syrup, slurry, or suspension), or other suitable form.

[0119] The formulation can contain components in addition to nucleic acid, where the additional components aid in the delivery of the nucleic acid to the target intestinal cell. The nucleic acid can be present in a pharmaceutical composition of the invention with additional components such as, but not limited to, stabilizing compounds and/or biocompatible pharmaceutical-carriers, e.g., saline, buffered saline, dextrose, or water. The nucleic acid can also be administered alone or in combination with other agents, including other therapeutic agents. The formulation can also contain organic and inor-

ganic compounds to, for example, facilitate nucleic acid delivery to and uptake by the target cell (e.g., detergents, salts, chelating agents, etc.).

[0120] Where the nucleic acid formulation is administered orally, the formulation can contain buffering agents or comprise a coating to protect the nucleic acid from stomach acidity and/or facilitate swallowing. In addition or alternatively, the oral formulation can be administered during an interdigestive period (between meals or at bedtime) when stomach pH is less acidic or with the administration of inhibitors of acid secretion such as H2 blockers (e.g., cimetidine) or proton pump inhibitors (e.g., PROLISECT™). The formulation can also comprise a time-release capsule designed to release the nucleic acid upon reaching the surface of the target intestinal cells.

[0121] A nucleic acid can be formulated in a complex with a liposome. Such complexes comprise a mixture of lipids which bind to nucleic acid, providing a hydrophobic core and hydrophilic coat which allows the genetic material to be delivered into cells. Suitable liposomes include DOPE (dioleoyl phosphatidyl ethanol amine), CUDMEDA (N-(5-cholestrum-3-β-ol 3-urethanyl)-N',N'-dimethylethylene diamine).

[0122] Other formulations can also be used for nucleic acids. Such formulations include RNA coupled to a carrier molecule (e.g., an antibody or a receptor ligand) which facilitates delivery to a target cell. An RNA can be chemically modified. By the term “chemical modification” is meant modifications of nucleic acids to allow, for example, coupling of the nucleic acid compounds to a carrier molecule such as a protein or lipid, or derivative thereof. Exemplary protein carrier molecules include antibodies specific to target cells.

[0123] A nucleic acid can be formulated with any of a variety of natural polymers, synthetic polymers, synthetic co-polymers, and the like. Generally, the polymers are biodegradable, or can be readily eliminated from the subject. Naturally occurring polymers include polypeptides and polysaccharides. Suitable synthetic polymers include, but are not limited to, polylysines, and polyethyleneimines (PEI; Boussif et al., *PNAS* 92:7297-7301, 1995) which molecules can also serve as condensing agents. These carriers may be dissolved, dispersed or suspended in a dispersion liquid such as water, ethanol, saline solutions and mixtures thereof. A wide variety of synthetic polymers are known in the art and can be used.

[0124] A nucleic acid can be formulated in a lipid-based vehicle. Lipid-based vehicles include cationic liposomes such as disclosed by Felgner et al (U.S. Pat. Nos. 5,264,618 and 5,459,127; *PNAS* 84:7413-7417, 1987; *Annals N.Y. Acad. Sci.* 772:126-139, 1995); they may also consist of neutral or negatively charged phospholipids or mixtures thereof including artificial viral envelopes as disclosed by Schreier et al. (U.S. Pat. Nos. 5,252,348 and 5,766,625). Nucleic acid/liposome complexes are suitable, and can comprise a mixture of lipids which bind to nucleic acid by means of cationic charge (electrostatic interaction). Cationic liposomes that are suitable for use include 3β-[N-(N',N'-dimethyl-aminoethane)-carbamoyl]-cholesterol (DC-Chol), 1,2-bis(oleoyloxy-3-trimethylammonio-propane (DOTAP) (see, for example, WO 98/07408), lysinyphosphatidylethanolamine (L-PE), lipopolyamines such as lipospermine, N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propanaminium bromide, dimethyl dioctadecyl ammonium bromide (DDAB), dioleoylphosphatidyl ethanolamine (DOPE), dio-

leoylphosphatidyl choline (DOPC), N(1,2,3-dioleoyloxy)propyl-N,N,N-triethylammonium (DOTMA), DOSPA, DMRIE, GL-67, GL-89, Lipofectin, and Lipofectamine (Thiery et al. (1997) *Gene Ther.* 4:226-237; Felgner et al., *Annals N.Y. Acad. Sci.* 772:126-139, 1995; Eastman et al., *Hum. Gene Ther.* 8:765-773, 1997). Polynucleotide/lipid formulations described in U.S. Pat. No. 5,858,784 can also be used in the methods described herein. Many of these lipids are commercially available from, for example, Boehringer-Mannheim, and Avanti Polar Lipids (Birmingham, Ala.). Also suitable are the cationic phospholipids found in U.S. Pat. Nos. 5,264,618, 5,223,263 and 5,459,127. Other suitable phospholipids which may be used include phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingomyelin, phosphatidylinositol, and the like. Cholesterol may also be included.

Stem Cells

[0125] For administration to a mammalian host, a genetically modified stem cell population (e.g., a genetically modified NSC population) can be formulated as a pharmaceutical composition. A pharmaceutical composition can be a sterile aqueous or non-aqueous solution, suspension or emulsion, which additionally comprises a physiologically acceptable carrier (i.e., a non-toxic material that does not interfere with the activity of the active ingredient). Any suitable carrier known to those of ordinary skill in the art may be employed in a subject pharmaceutical composition. Representative carriers include physiological saline solutions, gelatin, water, alcohols, natural or synthetic oils, saccharide solutions, glycols, injectable organic esters such as ethyl oleate or a combination of such materials. Optionally, a pharmaceutical composition may additionally contain preservatives and/or other additives such as, for example, antimicrobial agents, antioxidants, chelating agents and/or inert gases, and/or other active ingredients.

[0126] For example, a genetically modified stem cell population (e.g., a genetically modified NSC population) can be supplied in the form of a pharmaceutical composition comprising an isotonic excipient prepared under sufficiently sterile conditions for human administration. For general principles in medicinal formulation, see, e.g., *Cell Therapy: Stem Cell Transplantation, Gene Therapy, and Cellular Immunotherapy*, by G. Morstyn & W. Sheridan eds, Cambridge University Press, 1996; and *Hematopoietic Stem Cell Therapy*, E. D. Ball, J. Lister & P. Law, Churchill Livingstone, 2000.

[0127] In some embodiments, a genetically modified stem cell population (e.g., a genetically modified NSC population) is encapsulated, according to known encapsulation technologies, including microencapsulation (see, e.g., U.S. Pat. Nos. 4,352,883; 4,353,888; and 5,084,350). Where the NSCs are encapsulated, in some embodiments the NSCs are encapsulated by macroencapsulation, as described in U.S. Pat. Nos. 5,284,761; 5,158,881; 4,976,859; 4,968,733; 5,800,828 and published PCT patent application WO 95/05452.

[0128] A unit dosage form of a genetically modified stem cell population (e.g., a genetically modified NSC population) can contain from about 10^3 cells to about 10^9 cells, e.g., from about 10^3 cells to about 10^4 cells, from about 10^4 cells to about 10^5 cells, from about 10^5 cells to about 10^6 cells, from about 10^6 cells to about 10^7 cells, from about 10^7 cells to about 10^8 cells, or from about 10^8 cells to about 10^9 cells.

[0129] A genetically modified stem cell population (e.g., a genetically modified NSC population) will in some embodiments be transplanted into a patient according to conventional techniques, into the CNS, as described for example, in U.S. Pat. Nos. 5,082,670 and 5,618,531, or into any other suitable site in the body. In one embodiment, a population of NSCs is transplanted directly into the CNS. Parenchymal and intrathecal sites are also suitable. It will be appreciated that the exact location in the CNS will vary according to the disease state. Cells may be introduced by, for example, stereotaxic implantation or intracerebral grafting into the CNS of a patient.

[0130] In some embodiments, a genetically modified NSC population is administered as a cell suspension. In other embodiments, a genetically modified NSC population is administered as neurospheres. In other embodiments, a genetically modified NSC population is administered in an encapsulated form. In other embodiments, a genetically modified NSC population is contained with a reservoir, and the reservoir is implanted into the individual.

[0131] A single dose of a genetically modified NSC population can contain from about 10^3 cells to about 10^9 cells, e.g., from about 10^3 cells to about 10^4 cells, from about 10^4 cells to about 10^5 cells, from about 10^5 cells to about 10^6 cells, from about 10^6 cells to about 10^7 cells, from about 10^7 cells to about 10^8 cells, or from about 10^8 cells to about 10^9 cells. In some embodiments, multiple doses of a genetically modified NSC population are administered to an individual in need of such treatment. Doses can be administered at regular intervals (e.g., once a week, once a month, once every 6 weeks, once every 8 weeks, once every 6 months, etc.). Alternatively doses beyond an initial dose can be administered according to need, as determined by a medical professional, e.g., based on reappearance of symptoms associated with an apoE4-associated neurodegenerative disorder.

Genetically Modified Stem Cells

[0132] The present disclosure provides genetically modified stem cells and progeny thereof, where the stem cells (and/or progeny of such stem cells, such as NSCs, etc.) are genetically modified with one or more nucleic acids, and where the genetic modification results in a reduced level of tau polypeptide produced by the genetically modified cell, compared to a parent host cell or compared to a cell (e.g., a neuron) that normally produces tau.

[0133] Tau amino acid sequences are known in the art, as are nucleotide sequences encoding tau polypeptides. See, e.g., the nucleotide and amino acid sequences found under the GenBank accession numbers in parentheses in the following: Human Tau transcript variant 1 mRNA (NM_016835.3) and isoform 1 protein (NP_058519.2); human Tau transcript variant 2 mRNA (NM_005910.4) and isoform 2 protein (NP_005901.2); human Tau transcript variant 3 mRNA (NM_016834.3) and isoform 3 protein (NP_058518.1); human Tau transcript variant 4 mRNA (NM_016841.3) and isoform 4 protein (NP_058525.1); human Tau transcript variant 5 mRNA (NM_001123067.2) and isoform 5 protein (NP_001116539.1); and human Tau transcript variant 6 mRNA (NM_001123066.2) and isoform 6 protein (NP_001116539.1).

[0134] A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence set forth in

any one of GenBank Accession Nos. NP_058519, NP_005901, NP_058518, NP_058525, NP_001116539, and NP_001116539.

[0135] A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with an amino acid sequence encoded by any one of SEQ ID NOs:1-6. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:1. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:2. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:3. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:4. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:5. A tau polypeptide can comprise an amino acid sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity with the amino acid sequence encoded by SEQ ID NO:6.

[0136] A subject genetically modified stem cell and/or a progeny thereof, can synthesize less than about 80%, less than about 70%, less than about 50%, less than about 25%, less than about 10%, less than about 5%, or less than about 1%, of the amount of tau polypeptides synthesized by a parent (control) stem cell or progeny thereof that has not been genetically modified so as to reduce the level of tau transcript and/or polypeptide.

[0137] Methods of generating a subject genetically modified stem cell are described above.

[0138] The present disclosure provides a composition comprising a subject genetically modified stem cell, or progeny thereof.

[0139] A subject composition various components in addition to the genetically modified stem cells. For example, a subject composition can include a subject genetically modified stem cell and a culture medium. In some cases, the culture medium comprises one or more growth factors. In some embodiments, the culture medium is a serum-free culture medium. In some cases, the composition comprises genetically modified stem cells and a cryopreservative agent, e.g., a cryopreservation medium.

[0140] A subject composition can include a subject genetically modified stem cell and a matrix, e.g., a matrix component. Suitable matrix components include, e.g., collagen;

gelatin; fibrin; fibrinogen; laminin; a glycosaminoglycan; elastin; hyaluronic acid; a proteoglycan; a glycan; poly(lactic acid); poly(vinyl alcohol); poly(vinyl pyrrolidone); poly(ethylene oxide); cellulose; a cellulose derivative; starch; a starch derivative; poly(caprolactone); poly(hydroxy butyric acid); mucin; and the like. In some embodiments, the matrix comprises one or more of collagen, gelatin, fibrin, fibrinogen, laminin, and elastin; and can further comprise a non-proteinaceous polymer, e.g., can further comprise one or more of poly(lactic acid), poly(vinyl alcohol), poly(vinyl pyrrolidone), poly(ethylene oxide), poly(caprolactone), poly(hydroxy butyric acid), cellulose, a cellulose derivative, starch, and a starch derivative. In some embodiments, the matrix comprises one or more of collagen, gelatin, fibrin, fibrinogen, laminin, and elastin; and can further comprise hyaluronic acid, a proteoglycan, a glycosaminoglycan, or a glycan. Where the matrix comprises collagen, the collagen can comprise type I collagen, type II collagen, type III collagen, type V collagen, type XI collagen, and combinations thereof.

[0141] The matrix can be a hydrogel. A suitable hydrogel is a polymer of two or more monomers, e.g., a homopolymer or a heteropolymer comprising multiple monomers. Suitable hydrogel monomers include the following: lactic acid, glycolic acid, acrylic acid, 1-hydroxyethyl methacrylate (HEMA), ethyl methacrylate (EMA), propylene glycol methacrylate (PEMA), acrylamide (AAM), N-vinylpyrrolidone, methyl methacrylate (MMA), glycidyl methacrylate (GDMA), glycol methacrylate (GMA), ethylene glycol, fumaric acid, and the like. Common cross linking agents include tetraethylene glycol dimethacrylate (TEGDMA) and N,N'-methylenebisacrylamide. The hydrogel can be homopolymeric, or can comprise co-polymers of two or more of the aforementioned polymers. Exemplary hydrogels include, but are not limited to, a copolymer of poly(ethylene oxide) (PEO) and poly(propylene oxide) (PPO); Pluronic™ F-127 (a difunctional block copolymer of PEO and PPO of the nominal formula EO₁₀₀-PO₆₅-EO₁₀₀, where EO is ethylene oxide and PO is propylene oxide); poloxamer 407 (a tri-block copolymer consisting of a central block of poly(propylene glycol) flanked by two hydrophilic blocks of poly(ethylene glycol)); a poly(ethylene oxide)-poly(propylene oxide)-poly(ethylene oxide) co-polymer with a nominal molecular weight of 12,500 Daltons and a PEO:PPO ratio of 2:1); a poly(N-isopropylacrylamide)-base hydrogel (a PNIPAAm-based hydrogel); a PNIPAAm-acrylic acid copolymer (PNIPAAm-co-AAc); poly(2-hydroxyethyl methacrylate); poly(vinyl pyrrolidone); and the like.

[0142] The cell density in a subject iNSC/matrix composition can range from about 10² cells/mm³ to about 10⁹ cells/mm³, e.g., from about 10² cells/mm³ to about 10⁴ cells/mm³, from about 10⁴ cells/mm³ to about 10⁶ cells/mm³, from about 10⁶ cells/mm³ to about 10⁷ cells/mm³, from about 10⁷ cells/mm³ to about 10⁸ cells/mm³, or from about 10⁸ cells/mm³ to about 10⁹ cells/mm³.

[0143] The matrix can take any of a variety of forms, or can be relatively amorphous. For example, the matrix can be in the form of a sheet, a cylinder, a sphere, etc.

Subjects Suitable for Treatment

[0144] A variety of subjects are suitable for treatment with a subject method. Suitable subjects include any individual, particularly a human, who has an apoE4-associated disorder, who is at risk for developing an apoE-associated disorder,

who has had an apoE-associated disorder and is at risk for recurrence of the apoE4-associated disorder, or who is recovering from an apoE4-associated disorder.

[0145] Subjects suitable for treatment with a subject method include individuals who have one apoE4 allele; and individuals who have two apoE4 alleles. In other words, suitable subjects include those who are homozygous for apoE4 and those who are heterozygous for apoE4. For example, an individual can have an apoE3/apoE4 genotype, or an apoE4/apoE4 genotype. In some embodiments, the subject has been diagnosed as having Alzheimer's disease.

EXAMPLES

[0146] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Celsius, and pressure is at or near atmospheric. Standard abbreviations may be used, e.g., bp, base pair(s); kb, kilobase(s); pl, picoliter(s); s or sec, second(s); min, minute(s); h or hr, hour(s); aa, amino acid(s); kb, kilobase(s); bp, base pair(s); nt, nucleotide(s); i.m., intramuscular(ly); i.p., intraperitoneal(ly); s.c., subcutaneous(ly); and the like.

Example 1

GABAergic Interneuron Dysfunction Impairs Hippocampal Neurogenesis in Adult Apolipoprotein E4 Knockin Mice

[0147] It is shown that apoE4-KI mice have a significant age-dependent decrease in hilar GABAergic interneurons that correlates with the extent of apoE4-induced learning and memory deficits in aged mice. In neurotoxic apoE4 fragment transgenic mice, the interneuron loss was even more pronounced and correlated with the extent of learning and memory deficits. The interneuron loss and learning and memory deficits in these mice were prevented by tau removal, and the prevention was abolished by blocking GABA signaling with picrotoxin. In both groups of mice, the GABAA receptor potentiator pentobarbital rescued the learning and memory deficits. Thus, apoE4 causes age- and tau-dependent hilar GABAergic interneuron impairment, leading to learning and memory deficits in mice.

Materials and Methods

[0148] Reagents and Cell Culture. Minimal essential medium (MEM), Opti-MEM, and fetal bovine serum were from Invitrogen. Primary hippocampal neuronal cultures were prepared from P0 pups of apoE4(Δ 272-299)mE^{-/-}Tau^{+/+}, apoE4(Δ 272-299)mE^{-/-}Tau^{-/-}, mE^{-/-}Tau^{+/+}, mE^{-/-}Tau^{-/-} and wildtype mice, as reported (Li et al., 2009). Hippocampi were isolated on postnatal day 0, and dissociated cells were plated at 125,000 cells/ml in Neurobasal medium supplemented with B27, 100 U/ml penicillin G, and 100 μ g/ml streptomycin. The genotypes of cultures were deter-

mined by polymerase chain reaction (PCR) analysis of the tails of the pups from which the cells were obtained. After 14 days in vitro, the cultures were fixed in 4% paraformaldehyde in phosphate-buffered saline (135 mM NaCl, 2.7 mM KCl, 43 mM Na₂HPO₄, 14 mM KH₂PO₄, pH 7.4) for 30 min at room temperature. After permeabilization in phosphate-buffered saline with 0.1% Triton for 10 min, cells were placed in blocking buffer (phosphate-buffered saline with 10% normal serum from the same species that produced the secondary antibody and 0.01% Triton) for 30 min. Primary anti-GAD67 (1:250; Chemicon) were applied overnight at 4° C. and visualized with anti-mouse (GAD67) IgG conjugated with Alexa Fluor 488. Cells were counter-stained with 4',6-diamidino-2-phenylindole (DAPI). To measure GABAergic neuronal survival in hippocampal neuron cultures, GAD67-positive neurons were counted in 15-30 random fields under a fluorescence microscope at 200 \times magnification (Li et al., 2009).

[0149] Mice and Treatments. Human apoE3-KI and apoE4-KI mice (Sullivan et al., 2004) were from Taconic. Tau^{-/-} mice (Dawson et al., 2001) were backcrossed onto the mE^{-/-} background, and mE^{-/-}Tau^{-/-} mice were crossed with apoE4(Δ 272-299)mE^{-/-}Tau^{+/+} mice. Studies were conducted on female mice at 1, 3, 6, 12, 16, or 21 months of age. All mice were on a C57BL/6 genetic background. Some apoE4-KI and apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice received daily intraperitoneal injections of pentobarbital (20 mg/kg) or saline in their home cages for 21 days before the first day of Morris Water Maze (MWM) training and 1 h after daily training. Some apoE4(Δ 272-299) mE^{-/-}Tau^{-/-} mice were given daily injections of picrotoxin (1 mg/kg) or saline intraperitoneally in their home cages for 3 days before MWM training and 30 min before daily training. Brain tissues were collected after a 1-min transcardial perfusion with 0.9% NaCl. One hemibrain from each mouse was fixed in 4% paraformaldehyde, sectioned (30 μ m) with a microtome, and immunostained as described below. All experiments were performed in accordance with NIH and institutional guidelines.

[0150] Immunohistochemistry and Image Collection. Sliding microtome sections (30 μ m) were immunostained with the following primary antibodies: polyclonal goat anti-human apoE (1:8000 for fluorescence; Calbiochem), rabbit anti-neuropeptide Y (1:8000 for gamma-diamino butyric acid (DAB); ImmunoStar), rat anti-somatostatin (1:100 for DAB; Chemicon), mouse anti-GAD67 (1:1000 for DAB; Chemicon), mouse anti-MAP2 (1:500 for fluorescence; Sigma), mouse anti-synaptophysin (1:500 for fluorescence; DakoCytomation), and phosphorylation-dependent monoclonal antibody AT8 (p-Ser202; 1:100 for DAB; Endogen). Primary antibodies were detected with biotinylated goat anti-rabbit or goat anti-rat IgG (both 1:200; Vector Laboratories), Alexa Fluor 488-labeled goat anti-rabbit IgG (1:2000; Invitrogen), or Alexa Fluor 594-labeled donkey anti-mouse IgG (1:2000; Invitrogen). Stained sections were examined with a Radiance 2000 laser-scanning confocal system (Bio-Rad) mounted on a Nikon Optiphot-2 microscope. Images were processed with Photoshop CS (Adobe Systems).

[0151] Quantitative Analyses of Immunostained Brain Sections. GABAergic interneurons in the hilus of the dentate gyrus were quantified by counting GAD67-, neuropeptide Y (NPY)-, and somatostatin-positive cells in every tenth serial coronal section throughout the rostrocaudal extent of the hippocampus by an investigator blinded to genotype and

treatment (Li et al., 2009). Results are presented as the total number of positive cells counted per hemisphere, multiplied by two (for both hemibrains), and then by 10 (for every tenth serial section).

[0152] Morris Water Maze. The water maze pool (diameter, 122 cm) contained opaque water (22-23° C.) with a platform 10 cm in diameter. The platform was submerged 1.5 cm during hidden platform sessions (Harris et al., 2003; Raber et al., 1998; Roberson et al., 2007) and marked with black-and-white-striped mast (15 cm high) during cued training sessions. Mice were trained to locate the hidden platform (hidden days 1-5) and the cued platform (visible days 1-3) in two daily sessions (3.5 h apart), each consisting of two 60-s trials (hidden and cued training) with a 15-min intertrial interval. The platform location remained constant in the hidden platform sessions but was changed for each cued platform session. Entry points were changed semirandomly between trials. Twenty-four, 72, and 96 hours after the last hidden platform training, a 60-s probe trial (platform removed) was performed. Entry points for the probe trial were in the west quadrant, and the target quadrant was in the southeast quadrant. Performance was monitored with an EthoVision video-tracking system (Noldus Information Technology).

[0153] Elevated Plus Maze. The elevated plus maze tests "emotionality" and unconditioned anxiety-related behaviors that involve a conflict between the rodent's desire to explore a novel environment and anxiogenic elements such as elevation and an unfamiliar, brightly illuminated area (Roberson et al., 2007). The maze consists of two open arms and two closed arms equipped with rows of infrared photo-cells interfaced with a computer (Hamilton). Mice were placed individually into the center of the maze and allowed to explore for 10 min. The number of beam breaks was recorded to calculate the amount of time spent and distance moved in each arm and the number of entries into the open and closed arms. After each mouse was tested, the maze was cleaned with 70% ethanol to standardize odors.

[0154] Electrophysiology. ApoE3-KI and apoE4-KI mice were sacrificed and processed for slice preparation as described (Li et al., 2009). Brains were quickly removed into an ice-cold solution containing (in mM) 110 choline chloride, 2.5 KCl, 1.3 KH₂PO₄, 25 NaHCO₃, 0.5 CaCl₂, 7 MgCl₂, 10 dextrose, 1.3 sodium ascorbate, and 0.6 sodium pyruvate (300-305 mOsm). Horizontal slices (350 μm thick) were cut with a Vibratome, maintained in continuously oxygenated external solution (in mM: 125 NaCl, 2.5 KCl, 1.3 KH₂PO₄, 25 NaHCO₃, 2 CaCl₂, 1.3 MgCl₂, 1.3 sodium ascorbate, 0.6 sodium pyruvate, 10 dextrose, pH 7.4) at 30° C. for at least 40 min, and incubated at room temperature for at least 60 min before recording. Whole-cell voltage-clamp recordings from dentate gyrus granule cells were obtained with an infrared differential interference contrast video microscopy system. Patch electrodes (3-6 MΩ) were pulled from borosilicate glass capillary tubing (World Precision Instruments) on a horizontal Flaming-Brown microelectrode puller (model P-97, Sutter Instruments). Intracellular patch pipette solution contained (in mM) 120 Cs-gluconate, 10 HEPES, 0.1 EGTA, 15 CsCl₂, 4 MgCl₂, 4 Mg-ATP, and 0.3 Na₂-GTP, pH 7.25 (285-290 mOsm). To measure mIPSCs, slices were perfused with artificial cerebrospinal fluid containing 20 μM DNQX, 50 μM D-AP5, and 1 μM TTX. Whole-cell voltage-clamp data were low-pass filtered at 6 kHz (-3 dB, eight-pole Bessel), digitally sampled at 10 kHz with a Multiclamp 700A amplifier (Axon Instruments), and acquired with a Digidata-

1322 digitizer and pClamp 9.2 software (Axon Instruments). Whole-cell access resistance was monitored throughout the recording, and cells were rejected if values changed by >25%. mIPSCs were analyzed with a program provided by Dr. John Huguenard (Stanford University).

[0155] Statistical Analyses. Values are expressed as mean±SEM or mean±SD. Statistical analyses were performed with GraphPad Prism, Statview 5.0 (SAS Institute) or SPSS-10 (SPSS). Differences between means were assessed by t test, Mann-Whitney U test, one-factor ANOVA, or two-factor ANOVA, followed by Bonferroni, Tukey-Kramer or Fisher's PLSD post hoc tests. P<0.05 was considered statistically significant.

Results

[0156] Age-Dependent Decrease in GABAergic Interneurons in the Hilus of Dentate Gyrus of Female ApoE4-KI Mice. To assess the effect of aging and apoE4 on GABAergic interneurons in the hippocampus, their numbers were quantified in the hilus of dentate gyrus of female apoE3-KI and apoE4-KI mice at 1, 3, 6, 12, 16, and 21 months of age. Female mice were studied because they are susceptible to apoE4-induced learning and memory deficits (Raber et al., 1998; Raber et al., 2000). Anti-GAD67 and anti-somatostatin immunostaining, as shown representatively in 16-month-old apoE3-KI and apoE4-KI mice (FIGS. 1A-1D), revealed a significantly greater age-dependent decrease in GABAergic interneurons in the hilus of the dentate gyrus of female apoE4-KI mice than in age- and sex-matched apoE3-KI mice (FIGS. 1E and 1F). The significant difference between apoE4-KI and apoE3-KI mice was first observed at 6 months and was most pronounced at 16 and 21 months (FIGS. 1E and 1F). ApoE3-KI mice had a milder age-dependent decrease in hilar GABAergic interneurons (FIGS. 1E and 1F). Interestingly, the number of GABAergic interneurons in the hippocampal CA1 area did not differ in apoE3-KI and apoE4-KI mice at 16 months (FIGS. 1G-1I), suggesting a region-specific detrimental effect of apoE4 on GABAergic interneurons.

[0157] Presynaptic GABAergic Input onto Granule Cells is Reduced in Female ApoE4-KI Mice. The axonal termini of GABAergic interneurons on granule cells in the dentate gyrus of female apoE3-KI and apoE4-KI mice at 16 months of age were assessed by anti-GAD67 and anti-synaptophysin double immunofluorescence staining and confocal imaging analysis. The GABAergic axonal termini on granule cells were significantly decreased at the absolute level (GAD67 fluorescence intensity) and relative to the presynaptic marker synaptophysin (GAD67/synaptophysin ratio) in apoE4-KI mice (FIGS. 2A-2H). To assess the functional consequence of this finding, whole-cell patch-clamp recordings from granule cells were performed; glutamate currents were blocked with 6,7-dinitroquinoxaline-2,3-dione (DNQX) (20 μM) and D(-)-2-amino-5-phosphonovaleric acid (D-AP5) (50 μM), and action potential-mediated GABA release was blocked with tetrodotoxin (TTX) (1 μM). Consistent with the above findings, the frequency of miniature inhibitory postsynaptic currents (mIPSCs) was ~40% lower in apoE4-KI mice than in apoE3-KI mice (FIGS. 2I-2K). The mIPSC amplitude and membrane resistance were not altered significantly (FIGS. 2L and 2M). These results suggest that apoE4-KI mice have fewer functional GABAergic synapses onto granule cells.

[0158] Hilar GABAergic Interneuron Impairment Precedes Learning and Memory Deficits in Female ApoE4-KI Mice. Next, the spatial learning and memory of female apoE3-KI

and apoE4-KI mice at 12, 16, and 21 months of age was tested in the Morris water maze (MWM). At 12 months, apoE3-KI and apoE4-KI mice performed equally well in the hidden platform and probe trials, suggesting normal learning and memory in both groups. At 16 (FIG. 3A) and 21 (FIG. 4D) months, apoE3-KI mice quickly learned to find the hidden platform, which requires spatial learning, but apoE4-KI mice showed deficits. Swim speeds did not differ, indicating that the impairment was not due to motor deficits. ApoE3-KI and apoE4-KI mice performed equally well in visible platform trials, which test general performance deficits (FIG. 3A and FIG. 4D). In the probe trial 96 h after the last hidden platform trial, 16-month-old apoE4-KI mice had a deficit in memory retention (FIG. 3B, probe 3), although they performed as well as apoE3-KI mice in probe trials at 24 and 72 h (FIG. 3B, probe 2). Thus, hilar GABAergic interneuron impairment, first observed at 6 months of age, precedes the learning and memory deficits, which were first observed at 16 months of age, in apoE4-KI mice.

[0159] Hilar GABAergic Interneuron Impairment Correlates with Spatial Learning Deficits in Female ApoE4-KI Mice. In days 1-5 of the hidden platform trials, the number of hilar GABAergic interneurons correlated inversely with escape latency of apoE4-KI, but not apoE3-KI, mice at 16 months of age (FIGS. 3C-3F); no correlation was observed in visible platform trials (FIGS. 9A-9D). Similar results were obtained at 21 months of age (FIGS. 3G and 3H and FIGS. 9E and 9F). Interestingly, at both ages, all apoE3-KI mice had more than 2500 hilar GABAergic interneurons (FIGS. 3D, 3F, and 3H), whereas ~50% of the apoE4-KI mice had fewer than 2500 (FIGS. 3C, 3E, and 3G) and had greater learning deficits in the hidden platform trials (FIGS. 3C, 3E, and 3G).

[0160] The individual numbers of hilar GABAergic interneurons was looked at in female apoE4-KI mice at 6 or 12 months of age, when they also had, on average, significantly fewer hilar GABAergic interneurons than apoE3-KI mice at similar ages (FIGS. 1E and 1F). Interestingly, none of those mice had fewer than 2500 hilar GABAergic interneurons, and none of those mice had learning and memory deficits at 12 months, as mentioned above. Thus, 2500 might be the threshold number of hilar GABAergic interneurons that determines normal versus impaired learning performance of female mice in the MWM.

[0161] Pentobarbital Rescues Spatial Learning and Memory Deficits in Female ApoE4-KI Mice. To determine whether the loss of GABAergic interneurons contributes directly to the learning and memory deficits, 16-month-old female apoE4-KI mice were treated with the GABAA receptor potentiator pentobarbital for 4 weeks. This treatment rescued the learning and memory deficits (FIGS. 4A and 4B) but did not alter the number of hilar GABAergic interneurons (FIG. 4C). The learning deficit was also rescued in 21-month-old female apoE4-KI mice (FIG. 4D).

[0162] AD-Like Neurodegeneration Occurs in Transgenic Mice Expressing Low Levels of ApoE4(Δ 272-299). It was reported that neurons under stress, including neurons cultured in vitro (Harris et al., 2004b; Xu et al., 2008), express apoE and that neuronal apoE undergoes proteolytic cleavage to generate neurotoxic fragments, with apoE4 being more susceptible to the cleavage than apoE3 (Brecht et al., 2004; Harris et al., 2003; Huang et al., 2001). In primary hippocampal neuronal cultures, apoE4 impairs the survival of

GABAergic interneurons by generating more neurotoxic apoE fragments and increasing the levels of phosphorylated tau (p-tau) (Li et al., 2009).

[0163] To assess the contributions of apoE4 fragments and p-tau to hilar GABAergic interneuron impairment and behavioral deficits in vivo, transgenic mice were studied expressing low levels of apoE4(Δ 272-299), a major neurotoxic fragment in mouse and AD brains (Brecht et al., 2004; Harris et al., 2003), under the control of the neuron-specific Thy-1 promoter. These mice develop AD-like neurodegeneration and spatial learning and memory deficits (Harris et al., 2003). To eliminate confounding effects of mouse apoE, the original apoE4(Δ 272-299) transgenic line was crossed with apoE knockout (mE^{-/-}) mice to generate apoE4(Δ 272-299)mE^{-/-}Tau^{+/+} mice. To assess the effect of tau removal on AD-like neuronal and behavioral deficits caused by apoE4 fragments, apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice were crossed with mE^{-/-}Tau^{+/-} mice to generate littermates of apoE4(Δ 272-299) mE^{-/-}Tau^{+/+}, apoE4(Δ 272-299)mE^{-/-}Tau^{-/-}, and mE^{-/-}Tau^{+/-} and mE^{-/-}Tau^{-/-} mice. Eliminating endogenous tau did not alter the expression levels of apoE4(Δ 272-299). Age- and sex-matched wildtype mice were included as controls.

[0164] Morphological studies revealed neuronal deficits in the hippocampus of 12-month-old apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice, including presynaptic accumulation of apoE4 fragments as determined by anti-apoE and anti-synaptophysin (a presynaptic marker) or anti-MAP2 (a dendritic marker) double immunostaining (FIGS. 5A-5F), neurodegeneration as determined by hematoxylin/eosin and anti-MAP2 immunostaining (FIGS. 5G-5J), and tau pathology as determined by anti-p-tau (AT8 monoclonal antibody) immunostaining (FIGS. 5L, 5M, 5O, and 5P) in the hilus of the dentate gyrus, the hippocampal CA3 area, and the subiculum. Neurodegeneration and tau pathology occurred earliest in the hilus (FIGS. 5G-5J and 5L).

[0165] Tau Removal Prevents Loss of Hilar GABAergic Interneurons in Female ApoE4(Δ 272-299) Mice. Immunostaining for GAD67 (FIGS. 6A-6E), neuropeptide Y (NPY) (FIGS. 6F-6J), and somatostatin (FIGS. 6K-6O) revealed 40-50% fewer GABAergic interneurons in the hilus of apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice than in mE^{-/-}Tau^{+/+} or wildtype controls (FIGS. 6P-6R). Eliminating tau prevented neuronal deficits in apoE4 fragment transgenic mice, including loss of GABAergic interneurons in the hilus (FIG. 6), neurodegeneration (compare FIGS. 5K to 5I and 5J), and tau pathology in hilar interneurons (compare FIGS. 5N to 5L and 5M).

[0166] In 14-day primary hippocampal neuronal cultures, immunostaining for GAD67 (FIGS. 7A and 7B) revealed ~50% fewer GABAergic neurons in cultures from apoE4(Δ 272-299) mE^{-/-}Tau^{+/+} mice than from mE^{-/-}Tau^{+/+} controls (FIG. 7E) and markedly lower GAD67 immunoreactivity in neurites of surviving GABAergic neurons (compare FIG. 7B to 7A). Tau removal increased the survival of GABAergic neurons from apoE4(Δ 272-299)mE^{-/-}Tau^{+/+} mice to levels higher than in mE^{-/-}Tau^{+/+} mice (FIGS. 7A-7C and 7E). Removing tau also increased the survival of GABAergic neurons from mE^{-/-}Tau^{-/-} mice to levels higher than those of neurons from mE^{-/-}Tau^{+/+} mice (FIGS. 7A, 7D, and 7E). Thus, eliminating endogenous tau rescues apoE4 fragment-caused GABAergic interneuron impairment both in mice and in primary hippocampal neuronal cultures.

[0167] Tau Removal Prevents Spatial Learning and Memory Deficits in Female ApoE4(Δ 272-299) Mice. To assess effects of tau removal on learning and memory deficits induced by apoE4 fragments, 12-month-old female mice were tested in the MWM. In the hidden platform trial, mE^{-/-}-Tau^{+/+} and wildtype mice quickly learned the task, but apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice showed a deficit (FIG. 8A). Swim speeds of the mice did not differ. ApoE4(Δ 272-299)mE^{-/-}-Tau^{-/-} mice performed as well as mE^{-/-}-Tau^{+/+} and wildtype mice in the hidden platform trial (FIG. 8A). Thus, tau removal prevented the apoE4 fragment-induced learning deficit. In subsequent visible platform trials, all groups of mice performed equally well (FIG. 8A). In the probe trial 24 h after the last hidden platform trial, apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice had deficits in the target crossing and target quadrant tests that were eliminated by tau removal (FIGS. 8B and 8C). Interestingly, in the elevated plus maze, which assesses hippocampus-independent anxiety, apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice had increased anxiety that was unaffected by tau removal (FIG. 10A), suggesting that elimination of tau specifically affects hippocampus-dependent learning and memory performance.

[0168] Hilar GABAergic Interneuron Impairment Correlates with Spatial Learning Deficits in Female ApoE4(Δ 272-299) Mice with Tau. In apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice, the number of GABAergic interneurons in the hilus correlated inversely with escape latency on days 1-5 of the hidden platform test (FIGS. 8D-8F). Importantly, as in apoE4-KI mice (FIGS. 3C, 3E, and 3G), apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice with fewer than 2500 hilar GABAergic interneurons had greater learning deficits in the hidden platform trials (FIGS. 8D-8F), consistent with a threshold of \approx 2500 hilar GABAergic interneurons for normal versus impaired learning performance in the MWM. The number of hilar GABAergic interneurons did not correlate with performance in visible platform trials in apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice (FIGS. 10B-10D).

[0169] Tau Removal Prevents ApoE4-Induced Learning and Memory Deficits by Protecting Against Hilar GABAergic Interneuron Impairment. Finally, whether the rescue of learning and memory deficits by tau removal reflects protection against GABAergic interneuron impairment was determined. ApoE4(Δ 272-299)mE^{-/-}-Tau^{-/-} mice were treated with a subthreshold dose (1 mg/kg) of picrotoxin, a GABAA receptor antagonist, to block GABA signaling. The rescue was abolished (FIGS. 8G and 8H), but the number of hilar GABAergic interneurons was unaltered (FIG. 10E). Picrotoxin at this dose did not alter learning and memory in wildtype or mE^{-/-}-Tau^{+/+} mice (FIGS. 10F and 10G). In contrast, treatment of apoE4(Δ 272-299)mE^{-/-}-Tau^{+/+} mice with pentobarbital, a GABAA receptor potentiator, rescued the learning deficit (FIG. 10H). Evidently, tau removal rescues apoE4 fragment-induced learning and memory deficits by preventing the loss of GABAergic interneurons.

[0170] FIG. 1. Age-Dependent Significant Decrease in Numbers of GABAergic Interneurons in the Hilus of Dentate Gyrus of Female ApoE4-KI Mice. (A-D) Representative photomicrographs (200 \times) from female apoE3-KI and apoE4-KI mice at 16 months of age show GABAergic interneurons in the hilus after staining with anti-GAD67 (A and B) and anti-somatostatin (C and D). (E and F) Hilar GABAergic interneurons positive for GAD67 (E) or somatostatin (F) in female apoE3-KI and apoE4-KI mice at 1, 3, 6, 12, 16, and 21 months of age (n=4-12 mice per group) were quantified as described

in Experimental Procedures. Values are mean \pm SEM. *p<0.05; **p<0.01 (t test). (G and H) Representative photomicrographs (200 \times) show GABAergic interneurons in CA1 region of the hippocampus after staining with anti-GAD67. (I) Quantification of GAD67-positive GABAergic interneurons in CA1 region of the hippocampus in 16-month-old female apoE3-KI (n=10) and apoE4-KI (n=12) mice. Values are mean \pm SEM.

[0171] FIG. 2. Presynaptic GABAergic Input onto Granule Cells is Reduced in Female ApoE4-KI Mice. (A-F) Representative confocal images of the granule cell layer of the dentate gyrus of female apoE3-KI (A-C) and apoE4-KI (D-F) mice at 16 months of age stained with anti-GAD67 (A and D) and anti-synaptophysin (B and E). Merged images are shown in panels C and F. (G) GAD67 immunoreactivity (IR) of sections from different mice was quantified and normalized by area. Values are mean \pm SEM (four images per mouse and 4-5 mice per genotype). ***p<0.005 versus apoE3-KI mice (t test). (H) The ratio of GAD67-IR to synaptophysin-IR (a general presynaptic marker) in sections from different mice. Values are mean \pm SEM (four images per mouse and 4-5 mice per genotype). ***p<0.005 vs. apoE3-KI mice (t test). (I and J) Traces of miniature inhibitory postsynaptic currents (mIPSCs) in granule cells from apoE3-KI (I) or apoE4-KI (J) mice during whole-cell voltage clamp recording in the presence of DNQX (20 μ M), D-AP5 (50 μ M), and TTX (1 μ M). Scale bars, 20 pA and 700 ms. (K) Average mIPSC frequency in granule cells was lower in apoE4-KI mice than in apoE3-KI mice. Values are mean \pm SEM (n=9-10 cells per genotype). **p<0.01 versus apoE3-KI mice (t test). (L) Average mIPSC amplitude in granule cells was similar in apoE3-KI and apoE4-KI mice. Values are mean \pm SEM (n=9-10 cells per genotype). (M) Average membrane resistance of granule cells was similar in apoE3-KI and apoE4-KI mice. Values are mean \pm SEM (n=8-11 cells per genotype).

[0172] FIG. 3. Correlation of Hilar GABAergic Interneuron Impairment with Spatial Learning Deficits in ApoE4-KI Mice. (A) Ten apoE3-KI and 12 apoE4-KI female mice were tested at 16 months of age in the MWM. Points represent averages of daily trials. HD, hidden platform day (two trials/session, two sessions/day); HD0, first trial on HD1; VD, visible platform day (two trials/session, two sessions/day). Y-axis indicates time to reach the target platform (escape latency, mean \pm SEM). In the hidden platform days, learning curves differed significantly by genotypes (p<0.01, repeated-measures ANOVA). (B) The probe trials of female apoE3-KI and apoE4-KI mice at 16 months of age were performed 72 h (probe 2) and 96 h (probe 3) after the last hidden platform training. Percent time spent in the target quadrant versus the other quadrants differed by genotype in probe 3 (p<0.05). Values are mean \pm SEM. *p<0.05, ***p<0.005 (t test). (C and D) Escape latency in hidden platform days 1-5 correlated inversely with the number of GAD67-positive hilar GABAergic interneurons in apoE4-KI mice (C, n=12) but not apoE3-KI mice (D, n=10) at 16 months of age. (E and F) Escape latency in hidden platform days 1-5 correlated inversely with the number of somatostatin-positive hilar GABAergic interneurons in apoE4-KI mice (E, n=12) but not apoE3-KI mice (F, n=10) at 16 months of age. (G and H) Eight apoE3-KI and eight apoE4-KI female mice were tested at 21 months of age in the MWM. Escape latency in hidden platform days 1-5 correlated inversely with the number of somatostatin-positive hilar GABAergic interneurons in apoE4-KI mice (G, n=8) but not apoE3-KI mice (H, n=8) at 21 months of age.

[0173] FIG. 4. GABAA Receptor Potentiator Pentobarbital Rescues Spatial Learning and Memory Deficits in ApoE4-KI Mice. (A) Female 16-month-old apoE4-KI mice were treated with pentobarbital (PB, 20 mg/kg i.p.) or saline (n=6-13 per group) for 21 days before and daily during the MWM test. Untreated apoE3-KI mice (n=10) served as controls. The learning curve of pentobarbital-treated apoE4-KI mice differed from that of saline-treated apoE4-KI mice ($p < 0.05$, repeated-measures ANOVA) but resembled that of untreated apoE3-KI mice. Values are mean \pm SEM. HD, hidden day; VD, visible day. (B) In the probe trial 96 h after the last hidden session, pentobarbital treatment rescued memory deficits in 16-month-old apoE4-KI mice in the target quadrant and target cross tests (n=6-13 mice/group). Values are mean \pm SEM. *** $p < 0.005$ (t test). (C) Total number of GAD67-positive GABAergic interneurons in the hilus of apoE3-KI mice, apoE4-KI mice, and apoE4-KI mice treated with pentobarbital. Values are mean \pm SEM. ** $p < 0.01$, *** $p < 0.005$ vs. apoE3-KI mice (t test). (D) Female 21-month-old apoE4-KI mice were treated with pentobarbital (PB, 20 mg/kg) or saline (n=8 per group) for 21 days before and daily during the MWM test. Saline-treated apoE3-KI mice (n=8) served as controls. The learning curve of pentobarbital-treated apoE4-KI mice differed from that of saline-treated apoE4-KI mice ($p < 0.05$, repeated-measures ANOVA) but resembled that of saline-treated apoE3-KI mice. Values are mean \pm SEM. HD, hidden day; VD, visible day.

[0174] FIG. 5. Localization of apoE4(Δ 272-299) in the hippocampus and its effects on neurodegeneration and tau pathology in the presence and absence of tau. (A-D) Double immunofluorescence staining for apoE (green) and NeuN (red) in the hippocampus of apoE4(Δ 272-299)mE-/-Tau+/+ mice (magnification: A, 100 \times ; B, C, and D 400 \times). (E) Double immunofluorescence staining for apoE (green) and synaptophysin (Syn, red) in the CA3 region of apoE4(Δ 272-299)mE-/-Tau+/+ mice (600 \times). (F) Double immunofluorescence staining for apoE (green) and MAP2 (red) in the CA3 region of apoE4(Δ 272-299)mE-/-Tau+/+ mice (magnification, 600 \times). (G and H) Hematoxylin-eosin (HE) staining of the dentate gyrus of apoE4(Δ 272-299)mE-/-Tau+/+(G) and wildtype (H) mice (magnification, 200 \times). (I-K) Immunofluorescence staining for MAP2 in the hilus of the dentate gyrus of apoE4(Δ 272-299)mE-/-Tau+/+ (I), wildtype (J), and apoE4(Δ 272-299)mE-/-Tau-/- (K) mice (magnification, 200 \times). (L-N) Anti-p-tau (AT8 monoclonal antibody) immunostaining of the hilus of apoE4(Δ 272-299)mE-/-Tau+/+ (L), mE-/-Tau+/+ (M), and apoE4(Δ 272-299)mE-/-Tau-/- (N) mice (magnification, 400 \times). (O and P) Anti-p-tau (AT8 monoclonal antibody) immunostaining of the CA3 region (O) of the hippocampus and the subiculum (P) of apoE4(Δ 272-299)mE-/-Tau+/+ mice (magnification, 400 \times). All mice were 11-13 months of age.

[0175] FIG. 6. Loss of GABAergic Interneurons in the Hilus of the Dentate Gyrus of ApoE4(Δ 272-299)mE-/-Tau+/+Mice and Rescue by Tau Removal. The brains of 14 mE-/-Tau+/+, 10 apoE4(Δ 272-299)mE-/-Tau+/+, 12 apoE4(Δ 272-299)mE-/-Tau-/-, eight mE-/-Tau-/-, and 16 wildtype mice (all females) were collected at 12 months of age after behavioral assessment, sectioned, and immunostained with antibodies against GAD67, neuropeptide Y (NPY), or somatostatin. (A-O) Photomicrographs (200 \times) of GABAergic interneurons in the hilus after staining with anti-GAD67 (A-E), anti-NPY (F-J), or anti-somatostatin (K-O). (P-R) Total number of GAD67-positive (P), NPY-positive (q), and somatostatin-positive (R) GABAergic interneurons in the hilus. Values are mean \pm SEM. *** $p < 0.005$ (t test).

[0176] FIG. 7. Eliminating Tau Prevents the Neurotoxic Effect of ApoE4Fragments on Primary Hippocampal GABAergic Neurons. (A-D) Primary hippocampal neurons from individual P0 pups (mE-/-Tau+/+, apoE4(Δ 272-299)mE-/-Tau+/+, apoE4(Δ 272-299)mE-/-Tau-/-, and mE-/-Tau-/-) were cultured for 14 days in vitro (DIV 14) and double stained with anti-GAD67 (green) and DAPI (blue). Shown are representative images collected from three mice of each genotype and five fields per coverslip (magnification, 200 \times). (E) Numbers of GAD67-positive neurons were quantified as described in Experimental Procedures. Values are mean \pm SEM. ** $p < 0.01$, *** $p < 0.005$ (t test).

[0177] FIG. 8. Spatial Learning and Memory Deficits in ApoE4(Δ 272-299)mE-/-Tau+/+Mice and Rescue by Tau Removal. (A) Fourteen mE-/-Tau+/+, 10 apoE4(Δ 272-299)mE-/-Tau+/+, 12 apoE4(Δ 272-299)mE-/-Tau-/-, eight mE-/-Tau-/-, and 16 wildtype mice (all females) were tested at 12 months of age in the MWM. Values are mean \pm SEM. In the hidden platform days, learning curves differed significantly by genotype ($p < 0.001$, repeated-measures ANOVA). In post-hoc comparisons, apoE4(Δ 272-299)mE-/-Tau+/+ mice learned poorly ($p < 0.01$ vs. other groups). ApoE4(Δ 272-299)mE-/-Tau-/-, mE-/-Tau+/+, and wildtype mice performed at a similar level. (B) In the probe trial 24 h after the last hidden platform training, the number of target platform crossings versus crossings of the equivalent area in the other quadrants differed by genotype ($p < 0.05$). In post-hoc comparisons, apoE4(Δ 272-299)mE-/-Tau-/- mice performed better than apoE4(Δ 272-299)mE-/-Tau+/+ mice ($p < 0.01$) in the target crossing test. Only apoE4(Δ 272-299)mE-/-Tau+/+ mice showed impaired memory in the probe trail, and the deficit was rescued by tau removal. Values are mean \pm SEM. *** $p < 0.005$. (C) In the probe trial 24 h after the last hidden platform training, the time spent in the target quadrant versus the other quadrants differed by genotypes ($p < 0.01$). In post-hoc comparisons, only apoE4(Δ 272-299)mE-/-Tau+/+ mice showed impaired memory in the probe test, and the deficit was rescued by tau removal. Values are mean \pm SEM. *** $p < 0.005$. (D-F) Latency on hidden days 1-5 correlated inversely with the number of GAD67-positive (D), somatostatin-positive (E), and NPY-positive (F) GABAergic interneurons in the hilus in apoE4(Δ 272-299)mE-/-Tau+/+ mice. n=10 per analysis. (G) ApoE4(Δ 272-299)mE-/-Tau-/- mice were treated with picrotoxin (Picro, 1 mg/kg i.p.) or saline (n=6-8 per group) for 3 days before and daily during the MWM test. Saline-treated apoE4(Δ 272-299)mE-/-Tau+/+ and mE-/-Tau+/+ mice (n=6-8 per group) served as controls. The learning curve of picrotoxin-treated apoE4(Δ 272-299)mE-/-Tau-/- mice resembled that of saline-treated apoE4(Δ 272-299)mE-/-Tau+/+ mice, which differed significantly from those of saline-treated controls ($p < 0.01$). Values are mean \pm SEM. (H) In the probe trial 24 h after the last hidden session, picrotoxin-treated apoE4(Δ 272-299)mE-/-Tau-/- mice performed significantly worse than saline-treated apoE4(Δ 272-299)mE-/-Tau-/- ($p < 0.05$) or mE-/-Tau+/+ mice ($p < 0.01$) in the target crossing test. n=6-8 mice per group. Values are mean \pm SEM. * $p < 0.05$, ** $p < 0.01$.

[0178] FIG. 9. Performance in the cued platform trial does not correlate with the number of hilar GABAergic interneurons in apoE3-KI and apoE4-KI mice at 16 and 21 months of age. Ten apoE3-KI and 12 apoE4-KI mice at 16 months of age and eight apoE3-KI and eight apoE4-KI mice at 21 months of age (all females) were tested in the MWM. (A and B) Performance in the cued platform trial did not correlate with the

number of GAD67-positive GABAergic interneurons in apoE4-KI mice (A, n=12) or apoE3-KI mice (B, n=10) at 16 months of age. (C and D) Performance in the cued platform trial did not correlate with the number of somatostatin-positive GABAergic interneurons in apoE4-KI mice (C, n=12) or apoE3-KI mice (D, n=10) at 16 months of age. (E and F) Performance in the cued platform trial did not correlate with the number of somatostatin-positive GABAergic interneurons in apoE4-KI mice (E, n=8) or apoE3-KI mice (F, n=8) at 21 months of age.

[0179] FIG. 10. (A) Eliminating tau does not rescue apoE4 fragment-caused abnormal anxiety in apoE4(Δ 272-299) mE-/-Tau+/+ mice. Nine mE-/-Tau+/+, 10 apoE4(Δ 272-299)mE-/-Tau+/+, 12 apoE4(Δ 272-299)mE-/-Tau-/-, six mE-/-Tau-/-, and eight wildtype mice (all females). were tested in an elevated plus maze at 12 months of age. Values are mean \pm SEM. ***p<0.001 (t test). (B-D) Ten female apoE4 (Δ 272-299)mE-/-Tau+/+ were tested at 12 months of age in the MWM. Performance in the cued platform trial did not correlate with the number of GAD67-positive (B), somatostatin-positive (C), or NPY-positive (D) hilar GABAergic interneurons in apoE4(Δ 272-299)mE-/-Tau+/+ mice. (E) Treatment with the GABAA receptor antagonist picrotoxin (Picro) does not alter the number of hilar GABAergic interneurons in ApoE4(Δ 272-299)mE-/-Tau-/- mice. Female apoE4(Δ 272-299)mE-/-Tau-/- mice at 12 months of age were treated with picrotoxin (Picro, 1 mg/kg i.p.) or saline (n=6-8 per group) for 3 days before the MWM test and every day during the test. Age-matched, saline-treated apoE4(Δ 272-299) mE-/-Tau+/+ and mE-/-Tau+/+ mice (n=6-8 per group) served as controls. Total number of GAD67-positive interneurons in the hilus was quantified after the behavioral test. Values are mean \pm SEM. ***p<0.005 (t test). (F and G) Treatment with a low dose of picrotoxin does not alter the learning and memory performance in wildtype and mE-/-Tau+/+ mice. Female wildtype and mE-/-Tau+/+ mice at 12 months of age were treated with intraperitoneal injections of picrotoxin (Picro, 1 mg/kg) or saline (n=8 per group) for 3 days before the MWM test and every day during the test. Age-matched, saline-treated wildtype and mE-/-Tau+/+ mice (n=8 per group) served as controls. There was no significant difference among the learning curves (F). In the probe trial performed 24 h after the last hidden platform training, the time spent in the target quadrant versus the other quadrants does not differ by genotypes or treatment (G). Values are mean \pm SEM. ***p<0.005 (t test). (H) Treatment with GABAA receptor potentiator pentobarbital rescues the learning deficit in apoE4(Δ 272-299)mE-/-Tau+/+ mice. Female mE-/-Tau+/+ and apoE4(Δ 272-299)mE-/-Tau+/+ mice at 12 months of age were treated with intraperitoneal injections of pentobarbital (PB, 20 mg/kg) or saline (n=7-9 per group) for 21 days before the MWM test and every day during the test. In the hidden platform sessions, learning curves differed significantly by genotype and treatment (p<0.01, repeated-measures ANOVA). In post-hoc comparisons, apoE4(Δ 272-299)mE-/-Tau+/+ mice learned poorly versus mE-/-Tau+/+ mice (p<0.005). ApoE4(Δ 272-299)mE-/-Tau+/+ mice treated with pentobarbital learned better than saline-treated apoE4(Δ 272-299)mE-/-Tau+/+ mice (p<0.01). Values are mean \pm SEM.

REFERENCES

- [0180]** Bareggi, S. R., Franceschi, M., Bonini, L., Zecca, L., and Smirne, S. (1982). Decreased CSF concentrations of homovanillic acid and γ -aminobutyric acid in Alzheimer's disease. Age- or disease-related modifications? *Arch Neurol* 39, 709-712.
- [0181]** Bell, R. D., and Zlokovic, B. V. (2009). Neurovascular mechanisms and blood-brain barrier disorder in Alzheimer's disease. *Acta Neuropathol* 118, 103-113.
- [0182]** Bour, A., Grootendorst, J., Vogel, E., Kelche, C., Dodart, J.-C., Bales, K., Moreau, P.-H., Sullivan, P. M., and Mathis, C. (2008). Middle-aged human apoE4 targeted-replacement mice show retention deficits on a wide range of spatial memory tasks. *Behavioral Brain Res* 193, 174-182.
- [0183]** Brecht, W. J., Harris, F. M., Chang, S., Tesseur, I., Yu, G.-Q., Xu, Q., Fish, J. D., Wyss-Coray, T., Buttini, M., Mucke, L., et al. (2004). Neuron-specific apolipoprotein E4 proteolysis is associated with increased tau phosphorylation in brains of transgenic mice. *J Neurosci* 24, 2527-2534.
- [0184]** Brosh, I., and Barkai, E. (2009). Learning-induced enhancement of feedback inhibitory synaptic transmission. *Learn Mem* 16, 413-416.
- [0185]** Brunden, K. R., Trojanowski, J. Q., and Lee, V. M. Y. (2009). Advances in tau-focused drug discovery for Alzheimer's disease and related tauopathies. *Nat Rev Drug Discov* 8, 783-793.
- [0186]** Bu, G. (2009). Apolipoprotein E and its receptors in Alzheimer's disease: pathways, pathogenesis and therapy. *Nat Rev Neurosci* 10, 333-344.
- [0187]** Caselli, R. J., Dueck, A. C., Osborne, D., Sabbagh, M. N., Connor, D. J., Ahern, G. L., Baxter, L. C., Rapcsak, S. Z., Shi, J., Woodruff, B. K., et al. (2009). Longitudinal modeling of age-related memory decline and the APOE ϵ 4 effect. *N Engl J Med* 361, 255-263.
- [0188]** Corder, E. H., Saunders, A. M., Strittmatter, W. J., Schmechel, D. E., Gaskell, P. C., Small, G. W., Roses, A. D., Haines, J. L., and Pericak-Vance, M. A. (1993). Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261, 921-923.
- [0189]** Crowther, R. A. (1993). Tau protein and paired helical filaments of Alzheimer's disease. *Curr Opin Struct Biol* 3, 202-206.
- [0190]** Cui, Y., Costa, R. M., Murphy, G. G., Elgersma, Y., Zhu, Y., Gutmann, D. H., Parada, L. F., Mody, I., and Silva, A. J. (2008). Neurofibromin regulation of ERK signaling modulates GABA release and learning. *Cell* 135, 549-560.
- [0191]** Davies, P., Katzman, R., and Terry, R. D. (1980). Reduced somatostatin-like immunoreactivity in cerebral cortex from cases of Alzheimer disease and Alzheimer senile dementia. *Nature* 288, 279-280.
- [0192]** Dawson, H. N., Ferreira, A., Eyster, M. V., Ghoshal, N., Binder, L. I., and Vitek, M. P. (2001). Inhibition of neuronal maturation in primary hippocampal neurons from tau deficient mice. *J Cell Sci* 114, 1179-1187.
- [0193]** Dennis, N. A., Browndyke, J. N., Stokes, J., Need, A., Burke, J. R., Welsh-Bohmer, K. A., and Cabeza, R. (2009). Temporal lobe functional activity and connectivity in young adult APOE ϵ 4 carriers. *Alzheimers Dement* 5, 1-9.
- [0194]** Farrer, L. A., Cupples, L. A., Haines, J. L., Hyman, B., Kukull, W. A., Mayeux, R., Myers, R. H., Pericak-Vance, M. A., Risch, N., and Van Duijn, C. M. (1997).

- Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. *J Am Med Assoc* 278, 1349-1356.
- [0195] Filippini, N., MacIntosh, B. J., Hough, M. G., Goodwin, G. M., Frisoni, G. B., Smith, S. M., Matthews, P. M., Beckmann, C. F., and Mackay, C. E. (2009). Distinct patterns of brain activity in young carriers of the APOE- ϵ 4 allele. *Proc Natl Acad Sci USA* 106, 7209-7214.
- [0196] Götz, J., Chen, F., van Dorpe, J., and Nitsch, R. M. (2001). Formation of neurofibrillary tangles in P301L tau transgenic mice induced by Ab42 fibrils. *Science* 293, 1491-1495.
- [0197] Grouselle, D., Winsky-Sommerer, R., David, J. P., Delacourte, A., Dournaud, P., and Epelbaum, J. (1998). Loss of somatostatin-like immunoreactivity in the frontal cortex of Alzheimer patients carrying the apolipoprotein epsilon 4 allele. *Neurosci Lett* 255, 21-24.
- [0198] Hardy, J., Cowburn, R., Barton, A., Reynolds, G., Dodd, P., Wester, P., O'Carroll, A. M., Lofdahl, E., and Winblad, B. (1987). A disorder of cortical CABAergic innervation in Alzheimer's disease. *Neurosci Lett* 73, 192-196.
- [0199] Hardy, J., and Selkoe, D. J. (2002). The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* 297, 353-356.
- [0200] Harris, F. M., Brecht, W. J., Xu, Q., Mahley, R. W., and Huang, Y. (2004a). Increased tau phosphorylation in apolipoprotein E4 transgenic mice is associated with activation of extracellular signal-regulated kinase: Modulation by zinc. *J Biol Chem* 279, 44795-44801.
- [0201] Harris, F. M., Brecht, W. J., Xu, Q., Tesseur, I., Kekoni, L., Wyss-Coray, T., Fish, J. D., Masliah, E., Hopkins, P. C., Scearce-Levie, K., et al. (2003). Carboxyl-terminal-truncated apolipoprotein E4 causes Alzheimer's disease-like neurodegeneration and behavioral deficits in transgenic mice. *Proc Natl Acad Sci USA* 100, 10966-10971.
- [0202] Harris, F. M., Tesseur, I., Brecht, W. J., Xu, Q., Mullendorff, K., Chang, S., Wyss-Coray, T., Mahley, R. W., and Huang, Y. (2004b). Astroglial regulation of apolipoprotein E expression in neuronal cells. Implications for Alzheimer's disease. *J Biol Chem* 279, 3862-3868.
- [0203] Hartman, R. E., Wozniak, D. F., Nardi, A., Olney, J. W., Sartorius, L., and Holtzman, D. M. (2001). Behavioral phenotyping of GFAP-*apoE3* and -*apoE4* transgenic mice: *ApoE4* mice show profound working memory impairments in the absence of Alzheimer's-like neuropathology. *Exp Neurol* 170, 326-344.
- [0204] Herz, J., and Beffert, U. (2000). Apolipoprotein E receptors: Linking brain development and Alzheimer's disease. *Nat Rev Neurosci* 1, 51-58.
- [0205] Hoe, H. S., and Rebeck, G. W. (2008). Functional interactions of APP with the apoE receptor family. *J Neurochem* 106, 2263-2271.
- [0206] Hu, J. H., Ma, Y. H., Jiang, J., Yang, N., Duan, S. H., Jiang, Z. H., Mei, Z. T., Fei, J., and Guo, L. H. (2004). Cognitive impairment in mice over-expressing gamma-aminobutyric acid transporter 1 (GAT1) *Neuroreport* 15, 9-12.
- [0207] Huang, Y. (2006a). Apolipoprotein E and Alzheimer disease. *Neurology* 66 (Suppl. 1), S79-S85.
- [0208] Huang, Y. (2006b). Molecular and cellular mechanisms of apolipoprotein E4 neurotoxicity and potential therapeutic strategies. *Curr Opin Drug Discov Dev* 9, 627-641.
- [0209] Huang, Y., Liu, X. Q., Wyss-Coray, T., Brecht, W. J., Sanan, D. A., and Mahley, R. W. (2001). Apolipoprotein E fragments present in Alzheimer's disease brains induce neurofibrillary tangle-like intracellular inclusions in neurons. *Proc Natl Acad Sci USA* 98, 8838-8843.
- [0210] Irizarry, M. C., Rebeck, G. W., Cheung, B., Bales, K., Paul, S. M., Holtzman, D. M., and Hyman, B. T. (2000). Modulation of A β deposition in APP transgenic mice by an apolipoprotein E null background. *Ann NY Acad Sci* 920, 171-178.
- [0211] Jasinska, M., Siucinska, E., Cybulska-Klosowicz, A., Pyza, E., Furness, D. N., Kossut, M., and Glazewski, S. (2010). Rapid, learning-induced inhibitory synaptogenesis in murin barrel field. *J Neurosci* 30, 1176-1184.
- [0212] Kim, J., Basak, J. M., and Holtzman, D. M. (2009). The role of apolipoprotein E in Alzheimer's disease. *Neuron* 63, 287-303.
- [0213] Lewis, J., Dickson, D. W., Lin, W.-L., Chisholm, L., Corral, A., Jones, G., Yen, S.-H., Sahara, N., Skipper, L., Yager, D., et al. (2001). Enhanced neurofibrillary degeneration in transgenic mice expressing mutant tau and APP. *Science* 293, 1487-1491.
- [0214] Li, G., Bien-Ly, N., Andrews-Zwilling, Y., Xu, Q., Bernardo, A., Ring, K., Halabisky, B., Deng, C., Mahley, R. W., and Huang, Y. (2009). GABAergic interneuron dysfunction impairs hippocampal neurogenesis in adult apolipoprotein E4 knockin mice. *Cell Stem Cell* 5, 634-645.
- [0215] Mahley, R. W. (1988). Apolipoprotein E: Cholesterol transport protein with expanding role in cell biology. *Science* 240, 622-630.
- [0216] Mahley, R. W., Weisgraber, K. H., and Huang, Y. (2006). Apolipoprotein E4: A causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci USA* 103, 5644-5651.
- [0217] Namba, Y., Tomonaga, M., Kawasaki, H., Otomo, E., and Ikeda, K. (1991). Apolipoprotein E immunoreactivity in cerebral amyloid deposits and neurofibrillary tangles in Alzheimer's disease and kuru plaque amyloid in Creutzfeldt-Jakob disease. *Brain Res* 541, 163-166.
- [0218] Nitz, D., and McNaughton, B. (2004). Differential modulation of CA1 and dentate gyrus interneurons during exploration of novel environments. *Am Physiol Soc* 91, 863-872.
- [0219] Palop, J. J., and Mucke, L. (2009). Epilepsy and cognitive impairments in Alzheimer disease. *Neurol Res* 66, 435-440.
- [0220] Perrin, R. J., Fagan, A. M., and Holtzman, D. M. (2009). Multimodal techniques for diagnosis and prognosis of Alzheimer's disease. *Nature* 461, 917-922.
- [0221] Raber, J., Wong, D., Buttini, M., Orth, M., Bellosta, S., Pitas, R. E., Mahley, R. W., and Mucke, L. (1998). Isoform-specific effects of human apolipoprotein E on brain function revealed in ApoE knockout mice: Increased susceptibility of females. *Proc Natl Acad Sci USA* 95, 10914-10919.
- [0222] Raber, J., Wong, D., Yu, G.-Q., Buttini, M., Mahley, R. W., Pitas, R. E., and Mucke, L. (2000). Apolipoprotein E and cognitive performance. *Nature* 404, 352-354.

- [0223] Rapoport, M., Dawson, H. N., Binder, L. I., Vitek, M. P., and Ferreira, A. (2002). Tau is essential to b-amyloid-induced neurotoxicity. *Proc Natl Acad Sci USA* 99, 6364-6369.
- [0224] Roberson, E. D., Scarce-Levie, K., Palop, J. J., Yan, F., Cheng, I. H., Wu, T., Gerstein, H., Yu, G.-Q., and Mucke, L. (2007). Reducing endogenous tau ameliorates amyloid β -induced deficits in an Alzheimer's disease mouse model. *Science* 316, 750-754.
- [0225] Seidl, R., Cairns, N., Singewald, N., Kaehler, S. T., and Lubec, G. (2001). Differences between GABA levels in Alzheimer's disease and Down syndrome with Alzheimer-like neuropathology. *Naunyn Schmiedeberg Arch Pharmacol* 363, 139-145.
- [0226] Selkoe, D. J. (1991). The molecular pathology of Alzheimer's disease. *Neuron* 6, 487-498.
- [0227] Small, S. A., and Duff, K. (2008). Linking A β and tau in late-onset Alzheimer's disease: A dual pathway hypothesis. *Neuron* 60, 534-542.
- [0228] Strittmatter, W. J., Saunders, A. M., Goedert, M., Weisgraber, K. H., Dong, L.-M., Jakes, R., Huang, D. Y., Pericak-Vance, M., Schmechel, D., and Roses, A. D. (1994). Isoform-specific interactions of apolipoprotein E with microtubule-associated protein tau: Implications for Alzheimer disease. *Proc Natl Acad Sci USA* 91, 11183-11186.
- [0229] Strittmatter, W. J., Saunders, A. M., Schmechel, D., Pericak-Vance, M., Enghild, J., Salvesen, G. S., and Roses, A. D. (1993). Apolipoprotein E: High-avidity binding to b-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci USA* 90, 1977-1981.
- [0230] Sullivan, P. M., Mace, B. E., Maeda, N., and Schmechel, D. E. (2004). Marked regional differences of brain human apolipoprotein E expression in targeted replacement mice. *Neuroscience* 124, 725-733.
- [0231] Tanzi, R. E., and Bertram, L. (2001). New frontiers in Alzheimer's disease genetics. *Neuron* 32, 181-184.
- [0232] Tesseur, I., Van Dorpe, J., Bruynseels, K., Bronfman, F., Sciot, R., Van Lommel, A., and Van Leuven, F. (2000a). Prominent axonopathy and disruption of axonal transport in transgenic mice expressing human apolipoprotein E4 in neurons of brain and spinal cord. *Am J Pathol* 157, 1495-1510.
- [0233] Tesseur, I., Van Dorpe, J., Spittaels, K., Van den Haute, C., Moechars, D., and Van Leuven, F. (2000b). Expression of human apolipoprotein E4 in neurons causes hyperphosphorylation of protein tau in the brains of transgenic mice. *Am J Pathol* 156, 951-964.
- [0234] Vepsäläinen, S., Helisalmi, S., Koivisto, A. M., Tapaninen, T., Hiltunen, M., and Soininen, H. (2007). Somatostatin genetic variants modify the risk for Alzheimer's disease among Finnish patients. *J Neurol* 254, 1504-1508.
- [0235] Weisgraber, K. H. (1994). Apolipoprotein E: Structure-function relationships. *Adv Protein Chem* 45, 249-302.
- [0236] Wisniewski, T., and Frangione, B. (1992). Apolipoprotein E: A pathological chaperone protein in patients with cerebral and systemic amyloid. *Neurosci Lett* 135, 235-238.
- [0237] Xu, Q., Walker, D., Bernardo, A., Brodbeck, J., Bal-estra, M. E., and Huang, Y. (2008). Intron-3 retention/splicing controls neuronal expression of apolipoprotein E in the CNS. *J Neurosci* 28, 1452-1459.
- [0238] Xue, S., Jia, L., and Jia, J. (2009). Association between somatostatin gene polymorphisms and sporadic Alzheimer's disease in Chinese population. *Neurosci Lett* 465, 181-183.
- [0239] Zhao, C., Deng, W., and Gage, F. H. (2008). Mechanisms and functional implications of adult neurogenesis. *Cell* 132, 645-660.
- [0240] Zhong, N., and Weisgraber, K. H. (2009). Understanding the association of apolipoprotein E4 with Alzheimer's disease: Clues from its structure. *J Biol Chem* 284, 6027-6031.
- [0241] Zimmer, R., Teelken, A. W., Trieling, W. B., Weber, W., Weihmayr, T., and Lauter, H. (1984). γ -Aminobutyric acid and homovanillic acid concentration in the CSF of patients with senile dementia of Alzheimer's type. *Arch Neurol* 41, 602-604.
- [0242] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 6762

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

acggccgagc ggcagggcgc tcgcgcgcgc ccactagtgg ccggaggaga aggtccccgc 60

ggaggccgcg ctgcccgccc cctcccctgg ggaggetcgc gtccccctg ctccgcctg 120

cgccgcccgc cggcctcagg aacgcgcct cttcgcgggc ggcgcctc gcagtcacgc 180

-continued

ccaccacca gctccggcac caacagcagc gccgctgcca ccgcccacct tctgccgceg	240
ccaccacagc caccttctcc tctctcegtg tctctccccg tctctgcctc tgtcgactat	300
caggtagaact ttgaaccagg atggctgagc cccgccagga gttcgaagtg atggaagatc	360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag	420
accaagaggg tgacacggac gctggcctga aagaatctcc cctgcagacc cccactgagg	480
acggatctga ggaaccgggc tctgaaacct ctgatgctaa gagcactcca acagcgggaag	540
atgtgacagc acccttagtg gatgagggag ctccccggcaa gcaggctgcc gcgcagcccc	600
acacggagat cccagaagga accacagctg aagaagcagg cattggagac acccccagcc	660
tggaagacga agctgctggt cacgtgaccc aagagcctga aagtggtaag gtgggtccagg	720
aaggcttctc ccgagagcca gggccccag gtctgagcca ccagctcatg tccggcatgc	780
ctggggctcc cctcctgcct gagggcccca gagaggccac acgccaacct tcggggacag	840
gacctgagga cacagagggc ggcgcaccg cccctgagct gctcaagcac cagcttctag	900
gagacctga ccaggagggg ccgccctga agggggcagg gggcaaagag aggccgggga	960
gcaaggagga ggtggatgaa gaccgcgacg tcgatgagtc ctcccccaa gactccccctc	1020
cctccaaggc ctccccagcc caagatgggc ggcctcccca gacagccgcc agagaagcca	1080
ccagcatccc aggttccca gcggagggtg ccattccccct cctgtggat ttctctcca	1140
aagtttccac agagatccca gcctcagagc ccgacgggcc cagtgtaggg cgggcccagg	1200
ggcaggatgc cccccggag ttcaagtttc acgtggaaat cacaccaac gtgcagaagg	1260
agcaggcgca ctcgaggag catttgggaa gggctgcatt tccaggggcc cctggagagg	1320
ggccagaggc ccggggcccc tctttgggag aggcacacaa agaggctgac cttccagagc	1380
cctctgaaaa gcagcctgct gctgctccgc gggggaagcc cgtcagccgg gtcctcaac	1440
tcaaagctcg catggtcagt aaaagcaaag acgggactgg aagcgtgac aaaaaagcca	1500
agacatccac acgttctct gctaaaacct tgaaaaatag gccttgctt agccccaac	1560
acccactcc tggtagctca gaccctctga tccaacctc cagccctgct gtgtgcccag	1620
agccacttc ctctcctaaa cacgtctct ctgtcactc ccgaactggc agttctggag	1680
caaaggagat gaaactcaag ggggctgatg gtaaaacgaa gatcgccaca ccgcggggag	1740
cagccccctc aggccagaag ggccaggcca acgccaccag gattccagca aaaaccccgc	1800
ccgctccaaa gacaccacc agctctggtg aacctccaaa atcaggggat cgcagcggt	1860
acagcagccc cggctcccca ggcactcccc gcagccgctc ccgcaccccg tcccttccaa	1920
ccccaccac ccgggagccc aagaaggtgg cagtggtecg tactccacc aagtcgect	1980
cttcgccaa gagccgctg cagacagccc ccgtgccat gccagactg aagaatgtca	2040
agtccaagat cggctccact gagaacctga agcaccagcc gggaggcggg aaggtgcaga	2100
taattaataa gaagctggat cttagcaacg tccagtccaa gtgtggtca aaggataata	2160
tcaaacagc cccgggaggc ggcagtgtgc aaatagtcta caaacagtt gacctgagca	2220
aggtgacctc caagtgtggc tcattaggca acatccatca taaaccagga ggtggccagg	2280
tggaagtaaa atctgagaag ctgacttca aggcagagat ccagtcgaag attgggtccc	2340
tggacaatat caccacgctc cctggoggag gaaataaaaa gattgaaacc cacaagctga	2400
ccttccgcga gaacgcaaaa gccaaagacg accacggggc ggagatcgtg tacaagtcgc	2460

-continued

cagtgggtgc	tggggacacg	tctccacggc	atctcagcaa	tgtctcctcc	accggcagca	2520
tcgacatggt	agactcgccc	cagctcgcca	cgctagctga	cgaggtgtct	gcctccctgg	2580
ccaagcaggg	tttgtgatca	ggccctggg	gcggtcaata	attgtggaga	ggagagaatg	2640
agagagtgtg	aaaaaaaa	gaataatgac	ccggcccccg	ccctctgccc	ccagctgtct	2700
ctcgcagttc	ggtaattgg	ttaatcaact	aacctgcttt	tgtcactcgg	ctttggctcg	2760
ggacttcaaa	atcagtgatg	ggagtaagag	caaatttcat	ctttccaaat	tgatgggtgg	2820
gctagtaata	aaatatttaa	aaaaaaaaacat	tcaaaaacat	ggccacatcc	aacatttctt	2880
caggcaattc	cttttgattc	ttttttcttc	cccctccatg	tagaagaggg	agaaggagag	2940
gctctgaaag	ctgctctcgg	gggatttcaa	gggactgggg	gtgccaaacca	cctctggccc	3000
tgttgtgggg	gtgtcacaga	ggcagtggca	gcaacaaagg	atltgaaact	tggtgtgttc	3060
gtggagccac	aggcagacga	tgtcaacett	gtgtgagtgt	gacgggggtt	gggggggggc	3120
gggaggccac	gggggaggcc	gaggcagggg	ctgggcagag	gggagaggaa	gcacaagaag	3180
tgggagtggg	agaggaagcc	acgtgctgga	gagtagacat	ccccctcctt	gccgctggga	3240
gagccaaggc	ctatgccacc	tgcagcgtct	gagcggccgc	ctgtccttgg	tgccggggg	3300
tgggggcctg	ctgtgggtca	gtgtgccacc	ctctgcaggg	cagcctgtgg	gagaagggac	3360
agcgggtaaa	aagagaaggc	aagctggcag	gaggtggca	cttcgtggat	gacctcctta	3420
gaaaagactg	accttgatgt	cttgagagcg	ctggcctctt	cctccctccc	tgcagggtag	3480
ggggcctgag	ttgaggggct	tccctctgct	ccacagaaac	cctgttttat	tgagttctga	3540
aggttggaac	tgctgccatg	atlttggcca	ctttgcagac	ctgggacttt	agggctaacc	3600
agttctcttt	gtaaggactt	gtgcctcttg	ggagacgtcc	accctgttcc	aagcctgggc	3660
cactggcacc	tctggagtgt	gtgggggtct	gggaggcagg	tcccagagccc	cctgtccttc	3720
ccacggccac	tgcagtcacc	cctgtctcgg	ccgctgtgct	gttgtctgcc	gtgagagccc	3780
aatcactgcc	tatacccctc	atcacacgtc	acaatgtccc	gaattcccag	cctcaccacc	3840
ccttctcagt	aatgaccctg	gttggttgca	ggaggtacct	actccatact	gaggggtgaaa	3900
ttaaggggag	gcaaagtcca	ggcacaagag	tgggacccca	gcctctcact	ctcagttcca	3960
ctcatccaac	tgggaccctc	accacgaatc	tcgatgatctg	atccgggtcc	ctgtctcttc	4020
ctcccgctac	agatgtgagc	cagggcactg	ctcagctgtg	accctaggtg	tttctgctt	4080
gttgacatgg	agagagccct	ttcccctgag	aaggcctggc	cccttctctg	gctgagccca	4140
cagcagcagg	ctgggtgtct	tggttgtcag	tgggtggcacc	aggatggaag	ggcaaggcac	4200
ccagggcagg	cccacagtcc	cgctgtcccc	cacttgcacc	ctagcttgta	gctgccaacc	4260
tcccagacag	cccagcccgc	tgtcagctc	cacatgcata	gtatcagccc	tccacaccgc	4320
acaaagggga	acacaccccc	ttggaatgg	ttcttttccc	ccagtcccag	ctggaagcca	4380
tgtgtctgt	tctgtggag	cagctgaaca	tatacataga	tgttgccctg	ccctccccat	4440
ctgcaccctg	ttgagttgta	gttgatttg	tctgtttatg	cttgattca	ccagagtgc	4500
tatgatagtg	aaaagaaaa	aaaaaaaaaa	aaaggacgca	tgtatcttga	aatgcttgta	4560
aagaggttct	taacccacc	tcacgaggtg	tctctcacc	ccacactggg	actcgtgtgg	4620
cctgtgtggt	gccaccctgc	tggggcctcc	caagtttga	aaggcttcc	tcagcactg	4680
ggaccaaca	gagaccagct	tctagcagct	aaggaggccg	ttcagctgtg	acgaaggcct	4740

-continued

```

gaagcacagg attaggactg aagcgatgat gtccccctcc ctaactcccc ttggggctcc 4800
ctgtgtcagg gcacagacta ggtcttgtgg ctggtctggc ttgcggcgcg aggatggttc 4860
tctctgggtca tagcccgaag tctcatggca gtcccaaagg aggcttacia ctectgcatc 4920
acaagaaaaa ggaagccact gccagctggg gggatctgca gctcccagaa gctccgtgag 4980
cctcagccac ccctcagact gggttcctct ccaagctcgc cctctggagg ggcagcgag 5040
cctcccacca agggccctgc gaccacagca gggattggga tgaattgcct gtectggatc 5100
tgctctagag gcccagctg cctgcctgag gaaggatgac ttgacaagtc aggagacact 5160
gttcccaag ccttgaccag agcacctcag cccgctgacc ttgcacaaac tccatctgct 5220
gccatgagaa aagggagacc gcctttgcaa aacattgctg cctaaagaaa ctacagagcc 5280
tcaggcccaa ttctgccact tctggtttgg gtacagttaa aggcaacct gagggacttg 5340
gcagtagaaa tccagggcct cccctggggc tggcagcttc gtgtgcagct agagctttac 5400
ctgaaaggaa gtctctgggc ccagaactct ccaccaagag cctccctgcc gttcgtgag 5460
tcccagcaat tctcctaagt tgaagggatc tgagaaggag aaggaaatgt ggggtagatt 5520
tgggtgtggt tagagatatg cccccctcat tactgccaac agtttcggct gcatttcttc 5580
acgcaccteg gttcctcttc ctgaagttct tgtgccctgc tcttcagcac catgggctct 5640
cttatacgga aggctctggg atctccccct tgtgggggca ggctcttggg gccagcctaa 5700
gatcatggtt tagggtgatc agtgctggca gataaattga aaaggcacgc tggcttgtga 5760
tcttaaatga ggacaatccc cccagggctg ggcactctc ccctcccctc acttctccca 5820
cctcgagagc cagtgtcctt ggggtgggta gataggatat actgtatgcc ggctccttca 5880
agctgctgac tcactttatc aatagttcca tttaaattga cttcagtggt gagactgtat 5940
cctgtttgct attgcttgtt gtgctatggg gggagggggg aggaatgtgt aagatagtta 6000
acatgggcaa agggagatct tggggtgcag cacttaaact gcctcgtaac ccttttcatg 6060
atltcaacca catttgctag agggagggag cagccacgga gttagaggcc cttggggttt 6120
ctcttttcca ctgacaggct ttcccaggca gctggctagt tcattccctc cccagccagg 6180
tgcaggcgta ggaatatgga catctggttg ctttgccctg ctgccctctt tcaggggtcc 6240
taagcccaca atcatgcctc cctaagacct tggcatcctt ccctctaagc cgttggcacc 6300
tctgtgccac ctctcacact ggctccagac acacagctg tgcttttggg gctgagatca 6360
ctcgttccac cctcctcacc tttgttctcc aagtaaagcc acgaggtcgg ggcgagggca 6420
gaggtgatca cctgcgtgtc ccactctacag acctgcccgt tcataaaact tctgatttct 6480
cttcagcttt gaaaagggtt accctgggca ctggcctaga gcctcacctc ctaatagact 6540
tagcccatg agtttgccat gttgagcagg actatttctg gcacttgcaa gtcccatgat 6600
ttcttcggta attctgaggg tggggggagg gacatgaaat catcttagct tagctttctg 6660
tctgtgaatg tctatatagt gtattgtgtg ttttaacaaa tgatttacac tgactgttgc 6720
tgtaaaagtg aatttggaat taaagttatt actctgatta aa 6762

```

```

<210> SEQ ID NO 2
<211> LENGTH: 5811
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 2

```

-continued

acggccgagc ggcagggcgc tcgcgcgcgc ccactagtgg ccggaggaga aggctcccgc	60
ggaggccgcg ctgcccgcct cctcccctgg ggaggctcgc gttcccctg ctgcgctc	120
cgccgcccgc cggcctcagg aacgcgcctt cttcgccggc gcgcgccctc gcagtcaccg	180
ccaccacca gctccggcac caacagcagc gccgctgcca ccgcccacct tctgcccgcg	240
ccaccacagc caccttctcc tcctcogctg tcctctcccg tcctcgcctc tgtcgactat	300
cagggtgaact ttgaaccagg atggctgagc ccgcccagga gttcgaagtg atggaagatc	360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag	420
accaagaggg tgacacggac gctggcctga aagaatctcc cctgcagacc cccactgagg	480
acggatctga ggaaccgggc tctgaaacct ctgatgctaa gagcaactca acagcgggaag	540
atgtgacagc acccttagtg gatgagggag ctcccggcaa gcaggctgcc gcgagcccc	600
acacggagat cccagaagga accacagctg aagaagcagg cattggagac acccccagcc	660
tggaagacga agctgctggt cacgtgaccc aagctcgcct ggtcagtaaa agcaaagacg	720
ggactggaag cgatgacaaa aaagccaagg gggctgatgg taaaaagaa atcgccacac	780
cgcggggagc agcccctcca ggcagaagg gccaggccaa cgccaccagg attccagcaa	840
aaaccccgcc cgtccaaaag acaccacca gctctggtga acctccaaa tcaggggatc	900
gcagcgggta cagcagcccc ggctccccag gcaactccgg cagccgctcc cgcaccccg	960
cccttccaac cccaccacc cgggagccca agaaggtggc agtggtcctg actccacca	1020
agtgcgcgtc ttccgccaag agccgcctgc agacagcccc cgtgcccctg ccagacctga	1080
agaatgtcaa gtccaagatc ggctccactg agaacctgaa gcaccagccg ggaggcggga	1140
aggtgcagat aattaataag aagctggatc ttagcaactg ccagtccaag tgtggctcaa	1200
aggataatat caaacacgtc ccgggaggcg gcagtggtca aatagtctac aaaccagttg	1260
acctgagcaa ggtgacctcc aagtgtggct cattaggcaa catccatcat aaaccaggag	1320
gtggccaggt ggaagtaaaa tctgagaagc ttgacttcaa ggacagagtc cagtcgaaga	1380
ttgggtccct ggacaatatc acccagctcc ctggcggagg aaataaaaag attgaaacct	1440
acaagctgac cttccgagc aacgccaag ccaagacaga ccacggggcg gagatcgtgt	1500
acaagtgcgc agtgggtgct ggggacacgt ctccacggca tctcagcaat gtctcctcca	1560
ccggcagcat cgacatggta gactcgcccc agctcgccac gctagctgac gaggtgtctg	1620
cctccctggc caagcaggtt ttgtgatcag gccctgggg cggtcaataa ttgtggagag	1680
gagagaatga gagagtgtgg aaaaaaaaa aataatgacc cggccccgc cctctgcccc	1740
cagctgctcc tcgcatgttg gttaatgtgt taatcaacta acctgctttt gtcactcggc	1800
tttggctcgg gacttcaaaa tcagtgatgg gagtaagagc aaatttcac tttccaaatt	1860
gatgggtggg ctagtataa aatatttaa aaaaaacatt caaaaacatg gccacatcca	1920
acatttctc aggcaattcc ttttgattct tttttctcc cctccatgt agaagagggga	1980
gaaggagagg ctctgaaagc tgctctggg ggatttcaag ggactggggg tgccaaccac	2040
ctctggccct gttgtggggg tgtcacagag gcagtggcag caacaaagga tttgaaactt	2100
ggtgtgttcg tggagccaca gccagacgat gtcaaccttg tgtgagtgtg acgggggttg	2160
gggtggggcg ggaggccacg ggggagggcg aggcaggggc tgggcagagg ggagaggaag	2220
cacaagaagt gggagtggga gaggaagcca cgtgctggag agtagacatc cccctccttg	2280

-continued

ccgctgggag agccaaggcc tatgccacct gcagcgtctg agcggccgcc tgtccttgg	2340
ggccgggggt gggggcctgc tgtgggtcag tgtgccacc tctgcagggc agcctgtggg	2400
agaagggaca gcggttaaaa agagaaggca agctggcagg aggggtggcac ttcgtggatg	2460
acctccttag aaaagactga ccttgatgtc ttgagagcgc tggcctcttc ctccctccct	2520
gcagggtagg gggcctgagt tgaggggctt ccctctgctc cacagaaacc ctgttttatt	2580
gagttctgaa ggttgaact gctgccatga ttttggccac tttgcagacc tgggacttta	2640
gggctaacca gttctcttgg taaggacttg tgcctcttgg gagacgtcca cccgtttcca	2700
agcctgggcc actggcatct ctggagtgtg tgggggtctg ggaggcaggc cccgagcccc	2760
ctgtccttcc cacggccact gcagtcaccc ctgtctgctc cgctgtgctg ttgtctgccg	2820
tgagagccca atcactgcct ataccctca tcacacgtca caatgtcccg aattcccage	2880
ctcaccacc cttctcagta atgacctgg ttggttcag gaggtaccta ctccatactg	2940
agggtgaaat taaggaagg caaagtccag gcacaagagt gggaccccag cctctcactc	3000
tcagttccac tcatccaact gggacctca ccacgaatct catgatctga ttcggttccc	3060
tgtctctcc tcccgcaca gatgtgagcc agggcactgc tcagctgtga ccctaggtgt	3120
ttctgccttg ttgacatgga gagagccctt tcccctgaga aggcctggcc ccttctctgtg	3180
ctgagccca agcagcaggc tgggtgtctt ggttgcagc ggtggacca ggatggaagg	3240
gcaaggcacc cagggcaggc ccacagtccc gctgtcccc acttgcaacc tagctttag	3300
ctgccaacct cccagacagc ccagcccgt gctcagctcc acatgcatag taccagccct	3360
ccacaccga caaaggggaa cacaccccct tggaaatggt tcttttcccc cagtcccage	3420
tggaagccat gctgtctgtt ctgctggagc agctgaacat atacatagat gttgccctgc	3480
ctccccatc tgcacctgt tgagtttag ttggattgt ctgtttatgc ttggattcac	3540
cagagtgact atgatagtga aaagaaaaa aaaaaaaaa aaggacgcat gatatctgaa	3600
atgcttgtaa agaggtttct aaccaccct cacgaggtgt ctctacccc cacactggga	3660
ctcgtgtggc ctgtgtggg ccacctgct ggggcctccc aagttttgaa aggccttct	3720
cagcacctgg gacccaacag agaccagctt ctagcagcta aggaggccgt tcagctgtga	3780
cgaaggcctg aagcacagga ttaggactga agcgatgatg tccccttccc tacttcccct	3840
tggggctccc tgtgtcaggg cacagactag gtcttgtggc tggctcggct tgcggcgcga	3900
ggatggttct ctctggtcat agcccgaagt ctcatggcag tccc aaagga ggcttacaac	3960
tcctgcatca caagaaaaag gaagccactg ccagctgggg ggatctgcag ctcccagaag	4020
ctcgtgagc ctccagccacc cctcagactg ggttctctc caagctcgcc ctctggaggg	4080
gcagcgcagc ctcccacaa gggccctgcg accacagcag ggattgggat gaattgcctg	4140
tcctggatct gctctagagg cccaagctgc ctgcctgagg aaggatgact tgacaagtca	4200
ggagacactg tccc aaagc cttgaccaga gcacctcagc ccgctgacct tgcacaaact	4260
ccatctgctg ccattgagaa agggaaagcc ctttgcaaa acattgctgc ctaagaaac	4320
tcagcagcct caggcccaat tctgccactt ctggtttggg tacagttaaa ggcaacctg	4380
agggacttgg cagtagaat ccagggcctc ccctggggct ggcagctctg tgtgcagcta	4440
gagctttacc tgaaggaag tctctgggcc cagaactctc caccaagagc ctccctgccg	4500
ttcgtgagt cccagcaatt ctccaaagt gaaggatct gagaaggaga aggaaatgtg	4560

-continued

```

gggtagattt ggtggtggtt agagatatgc cccctcatt actgccaaca gtttcggctg 4620
catttcttca cgcacctcgg ttcctcttcc tgaagttctt gtgccctgct cttcagcacc 4680
atgggccttc ttatacggaa ggctctggga tctccccctt gtgggggag gctcttgggg 4740
ccagcctaag atcatggttt aggggatca gtgctggcag ataaattgaa aaggcacgct 4800
ggcttgatgat cttaaagtag gacaatcccc ccagggtggt gcaactcctcc cctccccctca 4860
cttctccccac ctgcagagcc agtgtccttg ggtgggctag ataggatata ctgtatgccg 4920
gctccttcaa gctgctgact cactttatca atagttccat ttaaattgac ttcagtgggtg 4980
agactgtatc ctgtttgcta ttgcttggtg tgctatgggg ggagggggga ggaatgtgta 5040
agatagttaa catgggcaaa gggagatctt ggggtgcagc acttaaactg cctcgtaac 5100
cttttcatga tttcaaccac atttgctaga gggagggagc agccacggag ttagaggccc 5160
ttggggtttc tcttttccac tgacaggctt tcccaggcag ctggctagt ttcctccctcc 5220
ccagccaggt gcaggcgtag gaatatggac atctggttgc tttggcctgc tgcctcttt 5280
caggggtcct aagcccacaa tcatgcctcc ctaagacctt ggcaccttc cctctaagcc 5340
gttggcacct ctgtgccacc tctcacactg gctccagaca cacagcctgt gcttttgag 5400
ctgagatcac tcgcttcacc ctccctcatct ttgttctcca agtaaagcca cgaggtcggg 5460
gcgagggcag aggtgatcac ctgcgtgtcc catctacaga cctgcccgtt cataaaactt 5520
ctgatttctc ttcagcttg aaaagggtta ccctgggac tggcctagag cctcacctcc 5580
taatagactt agccccatga gtttgccatg ttgagcagga ctatttctgg cacttgcaag 5640
tcccattgatt tcttcggtaa ttctgagggg ggggggaggg acatgaaatc atcttagctt 5700
agctttctgt ctgtgaatgt ctatatatgt tattgtgtgt ttaacaaat gatttacact 5760
gactgttctg gtaaaagtga atttgaaat aaagttatta ctctgattaa a 5811

```

<210> SEQ ID NO 3

<211> LENGTH: 5637

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

acggccgagc ggcagggcgc tcgcgcgcgc ccactagtgg ccgaggaga aggctcccgc 60
ggaggcgcgc ctgcccgccc cctcccctgg ggaggctcgc gttcccctg ctgcgcctg 120
cgcccccgc cggcctcagg aacgcgcctt cttcgcggc gcgcgcctc gcagtcaccg 180
ccaccacca gctccggcac caacagcagc gccgctgcc cgcgccacct tctgccgccc 240
ccaccacagc caccttctcc tctctcctg tcctctccc tcctcgcctc tgtcgactat 300
caggtgaact ttgaaccagg atggctgagc cccgccagga gttcgaagt atggaagatc 360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag 420
accaagaggg tgacacggac gctggcctga aagctgaaga agcaggcatt ggagacacc 480
ccagcctgga agacgaagct gctggtcacg tgacccaagc tcgcatggtc agtaaaagca 540
aagacgggac tggaagcgat gacaaaaaag ccaagggggc tgatggtaaa acgaagatcg 600
ccacaccgcg gggagcagcc cctccaggcc agaagggcca ggccaacgcc accaggatc 660
cagcaaaaac cccgcccctt ccaaagacac caccagctc tggtgaaact ccaaaatcag 720
gggatcgcag cggctacagc agccccgctt ccccaggcac tcccggcagc cgtcccgc 780

```

-continued

ccccgtccct tccaacccca cccacccggg agcccaagaa ggtggcagtg gtcctactc	840
cacccaagtc gccgtcttcc gccaaagacc gcctgcagac agccccctg cccatgccag	900
acctgaagaa tgtcaagtcc aagatcggct ccaactgagaa cctgaagcac cagccgggag	960
gcgggaaggt gcagataatt aataagaagc tggatcttag caactgccag tccaagtgtg	1020
gctcaaagga taatatcaaa cacgtcccgg gaggcggcag tgtgcaaata gtctacaaac	1080
cagttgacct gagcaagggt acctccaagt gtggctcatt aggcaacatc catcataaac	1140
caggagggtg ccagggtgaa gtaaaatctg agaagcttga cttcaaggac agagtccagt	1200
cgaagattgg gtccctggac aatatcacc acgtccctgg cggaggaaat aaaagattg	1260
aaaccacaaa gctgacctc cgcgagaac ccaaagccaa gacagaccac ggggcccaga	1320
tcgtgtacaa gtcgccagtg gtgtctgggg acactctcc acggcatctc agcaatgtct	1380
cctccaccgg cagcatcgac atggtagact cgcaccagct cgcacgcta gctgacgagg	1440
tgctgcctc cctggccaag cagggtttgt gatcaggccc ctggggcggg caataattgt	1500
ggagaggaga gaatgagaga gtgtggaaaa aaaaagaata atgaccggc ccccgcctc	1560
tgccccagc tgctcctgc agttcggta attggtaat cacttaacct gctttgtca	1620
ctcggctttg gctcgggact tcaaaatcag tgatgggagt aagagcaaat ttcactttc	1680
caaatgatg ggtgggctag taataaata ttaaaaaaa aacattcaaa aacatggcca	1740
catccaacat ttctcagcc aattcctttt gattcttttt tcttccccct ccatgtagaa	1800
gagggagaag gagagctct gaaagctgct tctgggggat ttcaaggac tgggggtgcc	1860
aaccacctct gcccctgtg tgggggtgtc acagaggcag tggcagcaac aaaggattg	1920
aaacttgggt gtctcgtgga gccacaggca gacgatgtca accttgtgtg agtgtgacgg	1980
gggttggggg ggggcccggg gccacggggg aggccgaggc aggggctggg cagaggggag	2040
aggaagcaca agaagtggga gtgggagagg aagccactg ctggagagta gacatcccc	2100
tccttgcccg tgggagagcc aaggcctatg ccacctgcag cgtctgagcg gccgcctgtc	2160
cttgggtggc gggggtggg gcctgctgtg ggtcagtggt ccaccctctg cagggcagcc	2220
tgtgggagaa gggacagcgg gtaaaaagag aaggcaagct ggcaggaggg tggcacttcg	2280
tggatgacct ccttagaaaa gactgacctt gatgtcttga gagcgtggc ctcttcctc	2340
ctccctgcag ggtagggggc ctgagttgag gggcttccct ctgctccaca gaaacctgt	2400
tttattgagt tctgaaggtt ggaactgctg ccatgatttt ggccactttg cagacctggg	2460
actttagggc taaccagttc tctttgtaag gacttgtgcc tcttgggaga cgtccaccg	2520
ttccaagcc tgggccaactg gcactctctg agtgtgtggg ggtctgggag gcaggctccg	2580
agccccctgt ccttcccacg gccactgcag tcaccctgt ctgcgccgt gtgctgtgt	2640
ctgccgtgag agcccaatca ctgcctatac cctcatcac acgtcacaat gtcccgaatt	2700
cccagcctca ccacccttc tcagtaatga cctgggttg ttgcaggagg tacctactc	2760
atactgaggg tgaatataag ggaaggcaaa gtccaggcac aagagtggga cccagcctc	2820
tcactctcag ttccactcat ccaactggga cctcaccac gaatctcatg atctgattcg	2880
gttccctgtc tctctctccc gtcacagatg tgagccaggg caactgctcag ctgtgacct	2940
aggtgtttct gccttgttga catggagaga gccctttccc ctgagaaggc ctggccctt	3000
cctgtgctga gccacagca gcaggctggg tgtcttggtt gtcagtgggt gcaccaggat	3060

-continued

ggaagggcaa ggcacccagg gcaggcccac agtcccgtg tccccactt gcaccctage	3120
ttgtagctgc caacctccca gacagcccag cccgctgctc agtcccacat gcatagtatc	3180
agccctccac acccgacaaa ggggaacaca ccccttggga aatggttctt tcccccaagt	3240
cccagctgga agccatgctg tctgttctgc tggagcagct gaacatatac atagatgttg	3300
ccctgcctc cccatctgca ccctgttgag ttgtagttgg atttgtctgt ttatgcttgg	3360
attcaccaga gtgactatga tagtgaaaag aaaaaaaaa aaaaaaagg acgcatgtat	3420
cttgaatgc ttgtaaagag gtttctaacc caccctcacg aggtgtctct cccccca	3480
ctgggactcg tgtggcctgt gtggtgccac cctgctggg cctccaagt ttgaaaggc	3540
ttctctcagc acctgggacc caacagagac cagcttctag cagctaagga ggccgttcag	3600
ctgtgacgaa ggctgaagc acaggattag gactgaagcg atgatgtccc ctccctact	3660
tcccctggg gctccctgtg tcagggcaca gactaggtct tgtggctggg ctggcttgcg	3720
gcgcgaggat ggttctctct ggtcatagcc cgaagtctca tggcagtcac aaaggagget	3780
tacaactcct gcatcacaag aaaaaggaag ccaactgccag ctggggggat ctgcagctcc	3840
cagaagctcc gtgagcctca gccaccctc agactgggtt cctctccaag ctgcccctc	3900
ggaggggagc cgcagcctcc caccaagggc cctgcgacca cagcagggat tgggatgaat	3960
tgcctgtcct ggatctgctc tagaggccca agctgcctgc ctgaggaagg atgactgac	4020
aagtcaaggag aactgttcc caaagcctg accagagcac ctacgccgc tgacctgca	4080
caaaactccat ctgctgccat gagaaaagg aagccgctt tgcaaacat tgctgcctaa	4140
agaaactcag cagcctcagg cccaattctg ccacttctgg tttgggtaca gttaaaggca	4200
acctgaggg acttgccagt agaaatccag ggcctccct ggggctggca gcttctgtg	4260
cagctagagc tttacctgaa aggaagtctc tgggccaga actctccacc aagagcctc	4320
ctgccgttcg ctgagctcca gcaattctcc taagttgaag ggatctgaga aggagaagga	4380
aatgtggggt agatttgggt gtggttagag atatgcccc ctcattactg ccaacagttt	4440
cggctgcatt tcttcacgca cctcggttcc tcttctgaa gttcttctgc cctgctctc	4500
agcaccatgg gccttcttat acggaaggct ctgggatctc ccccttgtgg gggcaggtc	4560
ttggggccag cctaagatca tggtttaggg tgatcagtc tggcagataa attgaaaagg	4620
cacgtggct tgtgatctta aatgaggaca atccccccag ggctgggac tctccctc	4680
ccctcacttc tcccactgc agagccagt tccttgggtg ggctagatag gatatactgt	4740
atgccggctc cttcaagctg ctgactcact ttatcaatag ttccattta attgacttca	4800
gtggtgagac tgtatctctg ttgctattgc ttgttctgct atggggggag gggggaggaa	4860
tgtgtaagat agttaacatg ggcaaaggga gatcttgggg tgcagcactt aaactgcctc	4920
gtaacccttt tcatgatctc aaccacatt gctagaggga gggagcagcc acggagttag	4980
aggcccttg ggtttctct tccactgac aggcttctcc aggcagctgg ctagtctatt	5040
ccctccccag ccaggtgcag gcgtaggaat atggacatct ggttcttgg cctgctgcc	5100
ctcttctcag ggtcctaagc ccacaatcat gcctccctaa gaccttggca tcttccctc	5160
taagccgttg gcacctctgt gccacctctc aactgctc cagacacaca gctgtgctt	5220
ttggagctga gatcactcgc ttcacctcc tcatcttctg tctccaagta aagccacgag	5280
gtcggggcga gggcagagg gatcacctgc gtgtccatc tacagacctg cggcttcata	5340

-continued

```

aaacttctga tttctcttca gctttgaaaa gggttaccct gggcactggc ctagagcctc 5400
acctccta at agacttagcc ccatgagttt gccatgttga gcaggactat ttctggcact 5460
tgcaagtccc atgatttctt cggtaattct gaggggtggg ggaggacat gaaatcatct 5520
tagcttagct ttctgtctgt gaatgtctat atagtgtatt gtgtgtttta acaaatgatt 5580
tactactgact gttgctgtaa aagtgaattt ggaaataaag ttattactct gattaaa 5637

```

<210> SEQ ID NO 4

<211> LENGTH: 5544

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```

acggccgagc ggcagggcgc tcgcgcgcgcc cactagtgg ccgaggaga aggctcccgc 60
ggaggccgcg ctgcccgcct cctcccctgg ggaggctcgc gttcccctg ctgcgccctg 120
cgccgcccgc cggcctcagg aacgcgcctt cttcgccggc gcgcccctc gcagtcaccg 180
ccaccacca gctccggcac caacagcagc gccgctgcc accccacct tctgcccgcg 240
ccaccacagc caccttctcc tctctcgctg tctctcccg tctcgcctc tctcgactat 300
cagggtgaact ttgaaccagg atggctgagc cccgccagga gttcgaagt atggaagatc 360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag 420
accaagaggg tgacacggac gctggcctga aagctgaaga agcaggcatt ggagacacc 480
ccagcctgga agacgaagct gctggtcacg tgaccaagc tcgcatggtc agtaaaagca 540
aagacgggac tgggaagcat gacaaaaaag ccaagggggc tgatggtaaa acgaagatcg 600
ccacaccgcg gggagcagcc cctccaggcc agaagggcca ggccaacgcc accaggattc 660
cagcaaaaac cccgcccctt ccaaagacac caccagctc tggtgaaact ccaaaatcag 720
gggatcgcag cggctacagc agccccggct ccccaggcac tcccggcagc cgtcccgcga 780
ccccgtccct tccaacccca cccaccggg agcccaagaa ggtggcagtg gtcctgactc 840
cacccaagtc gccgtcttcc gccaaagacc gcctgcagac agccccctg cccatgccag 900
acctgaagaa tgtcaagtcc aagatcggct cactgagaa cctgaagcac cagccgggag 960
gcgggaaggt gaaatagtc tacaaccag ttgacctgag caaggtgacc tccaagtgtg 1020
gctcattagc caatccat cataaaccag gaggtggcca ggtggaagta aaatctgaga 1080
agcttgactt caaggacaga gtccagtcga agattgggtc cctggacaat atcaccacg 1140
tccctggcgg aggaaataaa aagattgaaa cccacaagct gaccttcgc gagaacgcca 1200
aagccaagac agaccacggg gcggagatcg tgtacaagtc gccagtggtg tctggggaca 1260
cgtctccacg gatctcagc aatgtctct caccggcag catcgacatg gtagactcgc 1320
cccagctcgc cacgctagct gacgaggtgt ctgcctcct ggccaagcag ggtttgtgat 1380
caggcccctg gggcgtcaa taattgtgga gaggagagaa tgagagagtg tggaaaaaaa 1440
aagaataatg acccgcccc cgcctctgc cccagctgc tctcgcagt tcggttaatt 1500
ggttaatcac ttaacctgct tttgtcactc ggctttggct cgggacttca aaatcagtga 1560
tgggagtaag agcaaatctc atctttccaa attgatgggt gggctagtaa taaaatattt 1620
aaaaaaaaac attcaaaaac atggccacat ccaacatttc ctcaggcaat tctttttgat 1680
tcttttttct tccccctcca tgtagaagag ggagaaggag aggctctgaa agctgcttct 1740

```


-continued

gggggatttc aagggactgg ggggtgccaac cacctctggc cctgttgtgg ggggtgcaca	1800
gaggcagtgg cagcaacaaa ggatttgaaa cttgggtgtgt tcgtggagcc acaggcagac	1860
gatgtcaacc ttgtgtgagt gtgacggggg ttgggggtgg gggggaggcc acgggggagg	1920
ccgaggcagg ggctgggcag aggggagagg aagcacaaga agtgggagtg ggagaggaag	1980
ccacgtgctg gagagtagac atccccctcc ttgccgctgg gagagccaag gcctatgcca	2040
cctgcagcgt ctgagcggcc gcctgtcctt ggtggcggg ggtgggggccc tgctgtgggt	2100
cagtgtgcca ccctctgcag ggcagcctgt gggagaaggg acagcgggta aaaagagaag	2160
gcaagctggc aggagggtgg cacttcgtgg atgacctcct tagaaaagac tgacctgat	2220
gtcttgagag cgctggcctc ttccctccctc cctgcagggt agggggcctg agttgagggg	2280
cttccctctg ctccacagaa accctgtttt attgagttct gaaggttga actgctgcca	2340
tgatthtggc cactttgcag acctgggact ttagggctaa ccagttctct ttgtaaggac	2400
ttgtgctct tgggagacgt ccaccgctt ccaagcctgg gccactggca tctctggagt	2460
gtgtgggggt ctgggagga ggtcccagc ccctgtcct tcccacggcc actgcagtca	2520
ccccgtctg cgcctgtgtg ctggtgtctg ccgtgagagc ccaatcactg cctatacccc	2580
tcatacaacg tcacaatgac cgaattccc agcctacca ccccttctca gtaatgacct	2640
tggttggttg caggaggtac ctactccata ctgaggggtga aattaaggga aggcaaagtc	2700
caggcacaag agtgggacct cagcctctca ctctcagttc cactcatcca actgggacct	2760
tcaccacgaa tctcatgatc tgattcgggt ccctgtctcc tcctcccgtc acagatgtga	2820
gccagggcac tgctcagctg tgacctaggt tgtttctgccc ttgttgacat ggagagagcc	2880
ctttcccctg agaaggcctg gcccttctc gtgctgagcc cacagcagca ggtgggtgt	2940
cttgggtgtc agtgggtgga ccaggatgga agggcaaggc acccagggca ggcccacagt	3000
cccgtgtgccc cccactgca ccctagcttg tagctgcca cctcccagac agcccagccc	3060
gctgctcagc tccacatgca tagtatcagc cctccacacc cgacaaaagg gaacacacct	3120
ccttggaat ggttcttttc cccagctccc agctggaagc catgctgtct gttctgctgg	3180
agcagctgaa catatacata gatgtgccc tgcctcccc atctgcaccc tgttgagttg	3240
tagttggatt tgtctgttta tgcttgatt caccagagtg actatgatag tgaagaagaa	3300
aaaaaaaaa aaaaaggagc catgtatctt gaaatgcttg taaagaggtt tctaaccac	3360
cctcacgagg tgtctctcac ccccaactg ggactcgtgt ggctgtgtg gtgccacct	3420
gctggggcct cccaagtttt gaaaggctt cctcagcacc tgggacccaa cagagaccag	3480
cttctagcag ctaaggaggc cgttcagctg tgacgaaggc ctgaagcaca ggattaggac	3540
tgaagcagat atgtcccct ccctaactcc ccttggggct ccctgtgtca gggcacagac	3600
taggtcttgt ggtggtctg gcttgcggcg cgaggatggt tctctctggt catagcccga	3660
agtctcatgg cagtcccaaa ggaggcttac aactcctgca tcacaagaaa aaggaagcca	3720
ctgccagctg ggggatctg cagctcccag aagctccgtg agcctcagcc acccctcaga	3780
ctgggtcct ctccaagctc gccctctgga ggggcagcgc agcctcccac caagggcct	3840
gagaccacag cagggttggt gatgaattgc ctgtcctgga tctgctctag aggcccagc	3900
tgctgctgct aggaaggatg acttgacaag tcaggagaca ctgttccca agccttgacc	3960
agagcacctc agcccctga ccttgacaaa actccatctg ctgccatgag aaaagggaag	4020

-continued

```

ccgcctttgc aaaacattgc tgcctaaaga aactcagcag cctcaggccc aattctgcca 4080
cttctggttt gggtagcatt aaagcaacc ctgagggact tggcagtaga aatccagggc 4140
ctcccctggg gctggcagct tcgtgtgcag ctgagccttt acctgaaagg aagtctctgg 4200
gcccagaact ctccaccaag agcctccctg cgttcgctg agtcccagca attctcctaa 4260
gttgaagga tctgagaagg agaagaaat gtgggtaga tttggtggtg gttagagata 4320
tgccccctc attactgcca acagtttcgg ctgcatttct tcacgcacct cggttcctct 4380
tctgaagtt cttgtgcctt gctcttcagc accatgggccc ttcttatacg gaaggctctg 4440
ggatctcccc cttgtggggg caggctcttg gggccagcct aagatcatgg ttaggggtga 4500
tcagtctgag cagataaatt gaaaaggcac gctggcttgt gatcttaaat gaggacaatc 4560
ccccagggc tgggcaactc tccccctccc tcaactctcc cacctgcaga gccagtgctc 4620
ttgggtgggc tagataggat atactgtatg ccggctcctt caagctgctg actcaactta 4680
tcaatagttc cattaataat gacttcagtg gtgagactgt atcctgtttg ctattgcttg 4740
ttgtgctatg gggggagggg ggaggaatgt gtaagatagt taacatgggc aaaggagat 4800
cttgggtgac agcaactaaa ctgcctcgta acccttttca tgatttcaac cacatttgct 4860
agagggaggg agcagccacg gagttagagg cccttggggg ttctcttttc cactgacagg 4920
ctttcccagg cagctgggta gttcattccc tcccagcca ggtgcaggcg taggaatatg 4980
gacatctggt tgctttggcc tgctgccctc ttccagggg cctaagccca caatcatgcc 5040
tcctaagac cttggcatcc ttccctctaa gccgttgcca cctctgtgcc acctctcaca 5100
ctggctccag acacacagcc tgtgcttttg gagctgagat cactcgcttc accctcctca 5160
tctttgttct ccaagtaaag ccacgaggtc ggggcgaggg cagaggtgat cacctgcgtg 5220
tcccatctac agacctgcgg cttcataaaa cttctgattt ctcttcagct ttgaaaaggg 5280
ttaccctggg cactggccta gagcctcacc tcctaataga cttagcccca tgagtttgcc 5340
atgttgagca ggactatttc tggcacttgc aagtcccatg atttcttcgg taattctgag 5400
ggtgggggga gggacatgaa atcatcttag cttagcttcc tgtctgtgaa tgtctatata 5460
gtgtattgtg tgttttaaca aatgatttac actgactgtt gctgtaaaag tgaatttgga 5520
aataaagtta ttactctgat taaa 5544

```

<210> SEQ ID NO 5

<211> LENGTH: 5724

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

acggccgagc ggcagggcgc tcgcgcgcgc ccactagtgg ccgagagaga aggctcccgc 60
ggagggccgc ctgccccccc cctcccctgg ggaggctcgc gttcccctg ctgcgcctg 120
cgccgccgc cggcctcagg aacgcgcctt cttcgccggc gcgcgcctc gcagtcaccg 180
ccaccacca gctccggcac caacagcagc gccgctgcca ccgcccaact tctgccgccg 240
ccaccacagc caccttctcc tctctcgtg tcctctcccg tcctcgcctc tgtcgactat 300
caggtgaact ttgaaccagg atggtgagc cccgccagga gttcgaagtg atggaagatc 360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag 420
accaagaggg tgacacggac gctggcctga aagaatctcc cctgcagacc cccactgagg 480

```

-continued

acggatctga ggaaccgggc tctgaaacct ctgatgctaa gagcactcca acagcggaag	540
ctgaagaagc aggcattgga gacaccccca gcctggaaga cgaagctgct ggtcacgtga	600
cccaagctcg catggtcagt aaaagcaaag acgggactgg aagcgtgac aaaaaagcca	660
agggggctga tggtaaaacg aagatcgcca caccgcgggg agcagcccct ccaggccaga	720
agggccaggc caacgccacc aggattccag caaaaacccc gcccgctcca aagacaccac	780
ccagctctgg tgaacctcca aaatcagggg atcgcagcgg ctacagcagc cccggctccc	840
caggcactcc cggcagccgc tcccgcaccc cgtcccttcc aaccccaccc acccgggagc	900
ccaagaaggt ggcagtggtc cgtactccac ccaagtcgcc gtcttccgcc aagagccgcc	960
tgacagcagc ccccgtgccc atgccagacc tgaagaatgt caagtccaag atcggctcca	1020
ctgagaacct gaagcaccag cggggaggcg ggaaggtgca gataattaat aagaagctgg	1080
atcttagcaa cgtccagtcc aagtgtggct caaaggataa tatcaaacac gtcccgggag	1140
gcggcagtgt gcaaatagtc tacaaccag ttgacctgag caaggtgacc tccaagtgtg	1200
gctcattagc caacatccat cataaaccag gaggtggcca ggtggaagta aaatctgaga	1260
agcttgactt caaggacaga gtccagtcga agattgggtc cctggacaat atcaccacg	1320
tccctggcgg aggaaataaa aagattgaaa cccacaagct gaccttcgc gagaacgcca	1380
aagccaagac agaccacggg gcggagatcg tgtacaagtc gccagtggtg tctggggaca	1440
cgtctccacg gcatctcagc aatgtctcct ccaccggcag catcgacatg gtagactcgc	1500
cccagctcgc cacgctagct gacgaggtgt ctgcctcct ggccaagcag ggtttgtgat	1560
caggccctg gggcggtcaa taattgtgga gaggagagaa tgagagagtg tggaaaaaaa	1620
aagaataatg acccgccccc cgccctctgc ccccagctgc tcctcgcagt tcggttaatt	1680
ggttaatcac ttaacctgct tttgtcactc ggctttggct cgggacttca aaatcagtga	1740
tgggagtaag agcaaatctc atctttccaa attgatgggt gggctagtaa taaaaatatt	1800
aaaaaaaaac attcaaaaac atggccacat ccaacatttc ctcaggcaat tccttttgat	1860
tcttttttct tccccctcca tgtagaagag ggagaaggag aggctctgaa agctgcttct	1920
gggggatttc aagggactgg ggggtccaac cacctctggc cctgttgtgg ggggtgcaca	1980
gaggcagtgg cagcaacaaa ggattgaaa cttggtgtgt tcgtggagcc acaggcagac	2040
gatgtcaacc ttgtgtgagt gtgacggggg ttgggggtgg gcgggaggcc acgggggagg	2100
ccgaggcagc ggctgggcag aggggagagg aagcacaaga agtgggagtg ggagaggaag	2160
ccacgtgctg gagagtagac atccccctcc ttgccgctgg gagagccaag gcctatgcca	2220
cctgcagcgt ctgagcggcc gcctgtcctt ggtggccggg ggtgggggcc tgcgtgggt	2280
cagtgtgcca ccctctgcag ggcagcctgt gggagaaggg acagcgggta aaaagagaag	2340
gcaagctggc aggaggggtg cacttcgtgg atgacctcct tagaaaagac tgacctgat	2400
gtcttgagag cgtcggcctc ttcctccctc cctgcagggt agggggcctg agttgagggg	2460
cttccctctg ctccacagaa accctgtttt attgagttct gaaggttggg actgctgcca	2520
tgatthtggc cactthtgcag acctgggact ttagggctaa ccagttctct ttgtaaggac	2580
ttgtgcctct tgggagacgt ccaccgttt ccaagcctgg gccactggca tctctggagt	2640
gtgtgggggt ctgggaggca ggtcccgagc cccctgtcct tcccacggcc actgcagtca	2700
cccctgtctg cgcctgtgtg ctgthtctg ccgtgagagc ccaatcactg cctatacccc	2760

-continued

tcatacacag tcacaatgtc ccgaattccc agcctcacca ccccttctca gtaatgacce	2820
tggttggttg caggaggtac ctactccata ctgaggggtga aattaaggga aggcaaagtc	2880
caggcacaag agtgggaccc cagcctctca ctctcagttc cactcatcca actgggaccc	2940
tcaccacgaa tctcatgatc tgattcgggt cctctgtctcc tcctcccgtc acagatgtga	3000
gccagggcac tgctcagctg tgaccctagg tgtttctgcc ttgttgacat ggagagagcc	3060
ctttcccctg agaaggcctg gccccttctc gtgctgagcc cacagcagca ggctgggtgt	3120
cttgggtgtc agtgggtgga ccaggatgga agggcaaggc acccagggca ggcccacagt	3180
cccgtgtccc cccacttgca ccctagcttg tagctgcca cctcccagac agcccagccc	3240
gctgctcagc tccacatgca tagtatcagc cctccacacc cgacaaaagg gaacacaccc	3300
ccttggaat ggttcttttc ccccagtcct agctggaagc catgctgtct gttctgctgg	3360
agcagctgaa catatacata gatgttgccc tgccctcccc atctgcaacc tgttgagttg	3420
tagttggatt tgtctgttta tgcttgatt caccagagtg actatgatag tgaagaaaa	3480
aaaaaaaaa aaaaaggagc catgtatctt gaaatgcttg taaagaggtt tctaaccac	3540
cctcacgagg tgtctctcac cccacactg ggactcgtgt ggctgtgtg gtgccacct	3600
gctggggcct cccaagtttt gaaaggcttt cctcagcacc tgggaccaa cagagaccag	3660
cttctagcag ctaaggaggc cgttcagctg tgacgaaggc ctgaagcaca ggattaggac	3720
tgaagcagat atgtcccctt ccctacttcc ccttggggct ccctgtgtca gggcacagac	3780
taggtcttgt ggctggctg gcttgcggcg cgaggatggt tctctctggt catagcccga	3840
agtctcatgg cagtcccaaa ggaggcttac aactcctgca tcacaagaaa aaggaagcca	3900
ctgccagctg ggggatctg cagctcccag aagctccgtg agcctcagcc acccctcaga	3960
ctgggttctc ctccaagctc gccctctgga ggggcagcgc agcctccac caagggcct	4020
gcgaccacag cagggatggt gatgaattgc ctgtcctgga tctgctctag aggcccaagc	4080
tgctgctctg aggaaggatg acttgacaag tcaggagaca ctgttccca agccttgacc	4140
agagcacctc agcccctga ccttgcaaaa actccatctg ctgccatgag aaaagggaag	4200
ccgcctttgc aaaacattgc tgccaaaga aactcagcag cctcaggccc aattctgcca	4260
cttctggttt ggttacagtt aaaggcaacc ctgagggact tggcagtaga aatccagggc	4320
ctcccctggg gctggcagct tegtgtgcag cttagagctt acctgaaagg aagtctctgg	4380
gcccagaact ctccaccaag agcctccctg ccgttcgctg agtcccagca attctcctaa	4440
gttgaagga tctgagaagg agaaggaaat gtgggtaga tttggtggtg gttagagata	4500
tgccccctc attactgcca acagttctgg ctgcattctc tcaegcact cggttctct	4560
tctgaagtt cttgtgacct gctctcagc accatgggccc ttcttatacg gaaggctctg	4620
ggatctcccc cttgtggggg caggctcttg gggccagcct aagatcatgg tttagggtga	4680
tcagtgtctg cagataaatt gaaaaggcac gctggcttgt gatcttaaat gaggacaatc	4740
ccccagggc tgggcactcc tcccctccc tcaactctcc cacctgcaga gccagtgtcc	4800
ttgggtgggc tagataggat atactgtatg ccggtcctt caagctgctg actcacttta	4860
tcaatagttc catttaaat gacttcagtg gtgagactgt atcctgtttg ctattgcttg	4920
ttgtgctatg gggggagggg ggaggaatgt gtaagatagt taacatgggc aaaggagat	4980
cttgggtgac agcaactaaa ctgcctcgta acccttttca tgatttcaac cacatttgct	5040

-continued

```

agagggaggg agcagccacg gagttagagg cccttggggg ttctcttttc cactgacagg 5100
ctttcccagg cagctggcta gttcattccc tccccagcca ggtgcaggcg taggaatatg 5160
gacatctggt tgctttggcc tgctgcccct tttcaggggt cctaagccca caatcatgcc 5220
tccctaagac cttggcatcc ttccctctaa gccgttgga cctctgtgcc acctctcaca 5280
ctggctccag acacacagcc tgtgcttttg gagctgagat cactcgcttc acctcctca 5340
tctttgttct ccaagtaaag ccacgaggtc ggggcgaggg cagaggatgat cacctgcgtg 5400
tcccactcac agacctgcgg cttcataaaa cttctgattt ctcttcagct ttgaaaaggg 5460
ttacctggg cactggccta gaggctcacc tccataataga cttagcccca tgagtttgcc 5520
atgttgagca ggactatttc tggcacttgc aagtcccag atttcttcgg taattctgag 5580
gggtggggga gggacatgaa atcatcttag cttagctttc tgtctgtgaa tgtctatata 5640
gtgtattgtg tgttttaaca aatgatttac actgactggt gctgtaaaag tgaatttga 5700
aataaagtta ttactctgat taaa 5724

```

<210> SEQ ID NO 6

<211> LENGTH: 6816

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

acggccgagc ggcagggcgc tcgcgcgccg ccactagtgg ccggaggaga aggctcccgc 60
ggaggccgcg ctgcccgccc cctcccctgg ggaggctcgc gttcccctg ctgcgcctg 120
cgccgcccgc cggcctcagg aacgcgcctt cttcgccggc gcgcgcctc gcagtcaccg 180
ccaccacca gctccggcac caacagcagc gccgctgcca ccgccacct tctgcccgg 240
ccaccacagc caccttctcc tctctcgtg tctctcccg tcctcgctc tgtcgactat 300
caggtgaact ttgaaccagg atggctgagc cccgccagga gttcgaagt atggaagatc 360
acgctgggac gtacgggttg ggggacagga aagatcaggg gggctacacc atgcaccaag 420
accaagaggg tgacacggac gctggcctga aagaatctcc cctgcagacc cccactgagg 480
acggatctga ggaaccgggc tctgaaacct ctgatgctaa gagcactcca acagcggag 540
atgtgacagc accttagtg gatgagggag ctcccggcaa gcaggctgcc gcgcagcccc 600
acacggagat cccagaagga accacagctg aagaagcagg cattggagac acccccagcc 660
tggaagacga agctgctggt cacgtgaccc aagagcctga aagtggtaag gtggtccagg 720
aaggcttctt ccgagagcca ggccccccag gtctgagcca ccagctcatg tccggcatgc 780
ctggggctcc cctcctgctt gaggggccca gagaggccac acgccaacct tcggggacag 840
gacctgagga cacagagggc ggccgcccag ccctgagct gctcaagcac cagcttctag 900
gagacctgca ccaggagggg ccgcccgtga agggggcagg gggcaaagag aggccgggga 960
gcaaggagga ggtggatgaa gaccgagcgc tcgatgagtc ctcccccaa gaetcccctc 1020
cctccaagc ctcccagcc caagatgggc ggctcccga gacagccgcc agagaagcca 1080
ccagcatccc aggttccca gcggaggggt ccatcccct ccctgtggat ttctctcca 1140
aagtttccac agagatccca gcctcagagc ccgacgggcc cagtgtaggg cgggccaaag 1200
ggcaggatgc ccccctggag ttcacgtttc acgtgaaat cacaccaac gtgcagaagg 1260
agcaggcgca ctcgaggag catttgggaa gggctgcatt tccaggggcc cctggagagg 1320

```

-continued

ggccagagge	ccggggcccc	tctttgggag	aggacacaaa	agaggctgac	cttccagage	1380
cctctgaaaa	gcagcctgct	gctgctccgc	gggggaagcc	cgtcagccgg	gtccctcaac	1440
tcaaagctcg	catggctcagt	aaaagcaaa	acgggactgg	aagcgatgac	aaaaagcca	1500
agacatccac	acgttctct	gctaaaacct	tgaaaaatag	gccttgctt	agcccaaac	1560
accccaactcc	tggtagctca	gaccctctga	tccaaccctc	cagccctgct	gtgtgcccag	1620
agccaccttc	ctctcctaaa	cacgtctctt	ctgtcacttc	ccgaactggc	agttctggag	1680
caaaggagat	gaaactcaag	ggggctgatg	gtaaaacgaa	gatcgccaca	ccgcccgggag	1740
cagccctcc	aggccagaag	ggccaggcca	acgccaccag	gattccagca	aaaaccccgc	1800
ccgctccaaa	gacaccacc	agctctcgca	ctaagcaagt	ccagagaaga	ccaccccctg	1860
cagggcccg	atctgagaga	ggtgaacctc	caaatcagg	ggatcgccgc	ggctacagca	1920
gccccggctc	cccaggcact	cccggcagcc	gctcccgcac	cccgtccctt	ccaaccccac	1980
ccacccggga	gcccagaag	gtggcagtgg	tccgtactcc	acccaagtcc	ccgtcttccg	2040
ccaagagccg	cctgcagaca	gccccctg	ccatgccaga	cctgaagaat	gtcaagtcca	2100
agatcggtc	cactgagaac	ctgaagcacc	agccgggagg	cggaaggtg	cagataatta	2160
ataagaagct	ggatcttagc	aacgtccagt	ccaagtgtgg	ctcaaaggat	aatatcaaac	2220
acgtcccggg	aggcggcagt	gtgcaaatag	tctacaaacc	agttgacctg	agcaaggtga	2280
cctccaagt	tggtcatta	ggcaacatcc	atcataaac	aggaggtggc	caggtggaag	2340
taaaatctga	gaagcttgac	ttcaaggaca	gagtccagtc	gaagattggg	tccttgaca	2400
atatcaccca	cgctccctgg	ggaggaaata	aaaagattga	aaccacaag	ctgaccttc	2460
gcgagaacgc	caaagccaag	acagaccacg	ggcgggagat	cggtgacaag	tcgccagtgg	2520
tgtctgggga	cacgtctcca	cggcatctca	gcaatgtctc	ctccaccggc	agcatcgaca	2580
tggtagactc	gccccagctc	gccacgctag	ctgacgaggt	gtctgctcc	ctggccaagc	2640
agggtttg	atcagcccc	tggggcggtc	aataattgtg	gagaggagag	aatgagagag	2700
tgtgaaaaa	aaaagaata	tgaccggcc	cccgccctct	gccccagct	gctcctcgca	2760
gttcggtaa	ttggttaac	acttaacctg	ctttgtcac	tcggcttgg	ctcgggactt	2820
caaatcagt	gatgggagta	agagcaaat	tcctcttcc	aaattgatgg	gtgggctagt	2880
ataaaaat	ttaaaaaaa	acattcaaaa	acatggccac	atccaacatt	tcctcaggca	2940
attcctttt	attcctttt	cttccccctc	catgtagaag	agggagaagg	agaggctctg	3000
aaagctgctt	ctgggggatt	tcaagggact	gggggtgcca	accacctctg	gccctgttgt	3060
gggggtgca	cagaggcagt	ggcagcaaca	aaggatttga	aacttggtgt	gttcgtggag	3120
ccacaggcag	acgatgtcaa	ccttgtgtga	gtgtgacggg	ggttgggggtg	gggcccggag	3180
ccacggggga	ggccgaggca	ggggctgggc	agaggggaga	ggaagcaca	gaagtgggag	3240
tgggagagga	agccacgtgc	tggagagtag	acatccccct	ccttgccgct	gggagagcca	3300
aggcctatgc	caactgcagc	gtctgagcgg	ccgcctgtcc	ttggtggccg	ggggtggggg	3360
cctgctgtgg	gtcagtgctc	caccctctgc	agggcagcct	gtgggagaag	ggacagcggg	3420
taaaaagaga	aggcaagctg	gcaggagggt	ggcacttcgt	ggatgacctc	cttagaaaag	3480
actgaccttg	atgtcttgag	agcgtggcc	tcttctccc	tcctgcagg	gtagggggcc	3540
tgagttgagg	ggcttccctc	tgctccacag	aaaccctggt	ttattgagtt	ctgaaggttg	3600

-continued

gaactgctgc	catgattttg	gccactttgc	agacctggga	ctttagggct	aaccagttct	3660
ctttgtaagg	acttgtgcct	cttgggagac	gtccaccctg	ttccaagcct	gggccactgg	3720
catctctgga	gtgtgtgggg	gtctgggagg	caggtcccga	gccccctgtc	cttcccacgg	3780
ccactgcagt	caccctgtgc	tgccgctgtg	tgctgtttgc	tgccgtgaga	gcccacatcac	3840
tgccataacc	cctcatcaca	cgtcacaatg	tcccgaattc	ccagcctcac	cacccttctt	3900
cagtaatgac	cctggttggt	tgccaggagg	acctactcca	tactgagggt	gaaattaagg	3960
gaaggcaaa	tccaggcaca	agagtgggac	cccagcctct	cactctcagt	tccactcacc	4020
caactgggac	cctcaccacg	aatctcatga	tctgattcgg	ttccctgtct	cctcctcccg	4080
tcacagatgt	gagccagggc	actgtctcag	tgtagcccta	ggtgtttctg	ccttgttgac	4140
atggagagag	ccctttcccc	tgagaaggcc	tggcccttc	ctgtgctgag	cccacagcag	4200
caggctgggt	gtcttggttg	tcagtgggtg	caccaggatg	gaaggcaag	gcaccagggg	4260
caggcccaca	gtcccctgtg	ccccacttg	cacctagct	tgtagctgcc	aaactcccag	4320
acagcccagc	ccgctgtctc	gctccacatg	catagtatca	gccctccaca	cccgacaaag	4380
gggaacacac	ccccttgaa	atggttcttt	tccccagtc	ccagctgga	gccatgctgt	4440
ctgttctgct	ggagcagctg	aacatataca	tagatgttgc	cctgccctcc	ccatctgcac	4500
cctgttgagt	tgtagttgga	tttgtctggt	tatgcttggg	ttcaccagag	tgactatgat	4560
agtgaaaaga	aaaaaaaa	aaaaaaagga	cgcagtatc	ttgaaatgct	tgtaaagagg	4620
tttctaacc	accctcagca	ggtgtctctc	acccccacac	tgggactcgt	gtggcctgtg	4680
tggtgccacc	ctgctggggc	ctcccagtt	tgaaaggct	ttcctcagca	cctgggaccc	4740
aacagagacc	agcttctagc	agctaaggag	gccgttcagc	tgtagcgaag	gcctgaagca	4800
caggattag	actgaagcga	tgatgtcccc	ttccctactt	ccccttgggg	ctccctgtgt	4860
cagggcacag	actaggtctt	gtggtgggtc	tggcttgccg	cgcgaggatg	gttctctctg	4920
gtcatagccc	gaagtctcat	ggcagtccca	aaggaggctt	acaactcctg	catcacaaga	4980
aaaaggaagc	cactgccagc	tggggggatc	tgtagctccc	agaagctccg	tgagcctcag	5040
ccaccctca	gactgggttc	ctctccaagc	tcgccctctg	gaggggcagc	gcagcctccc	5100
accaagggcc	ctgagccacc	agcagggatt	gggatgaatt	gcctgtcctg	gatctgetct	5160
agaggcccaa	gctgctgccc	tgaggaagga	tgacttgaca	agtcaggaga	cactgttccc	5220
aaagccttga	ccagagcacc	tcagcccgtc	gaccttgcac	aaactccacc	tgctgccatg	5280
agaaaagga	agccgccttt	gcaaaacatt	gctgcctaaa	gaaactcagc	agcctcaggc	5340
ccaattctgc	cacttctggt	ttgggtacag	ttaaaggcaa	ccctgaggga	cttggcagta	5400
gaaatccagg	gcctcccctg	gggctggcag	cttctgtgtc	agctagagct	ttaoctgaaa	5460
ggaagtctct	gggccagaaa	ctctccacca	agagcctccc	tgccgttcgc	tgagtcaccag	5520
caattctcct	aagttgaagg	gatctgagaa	ggagaaggaa	atgtggggta	gatttgggtg	5580
tggttagaga	tatgcccccc	tcattactgc	caacagtttc	ggctgcattt	cttcacgcac	5640
ctcggttcct	cttccgtgaa	ttcttgtgcc	ctgctcttca	gcaccatggg	ccttcttata	5700
cggaggctc	tgggatctcc	cccttgtggg	ggcaggetct	tggggccagc	ctaagatcat	5760
ggtttagggt	gatcagtgct	ggcagataaa	tgaaaaggc	acgctggctt	gtgatcttaa	5820
atgaggacaa	tccccccagg	gctgggcact	cctcccctcc	cctcacttct	cccacctgca	5880

-continued

```

gagccagtgt ccttgggtgg gctagatagg atatactgta tgccggctcc tccaagctgc 5940
tgactcactt tatcaatagt tccatttaaa ttgacttcag tggtagact gtatcctgtt 6000
tgctattgct tgttgtgcta tggggggagg ggggaggaat gtgtaagata gttaacatgg 6060
gcaaagggag atcttggggt gcagcactta aactgcctcg taaccctttt catgatttca 6120
accacatttg ctagagggag ggagcagcca cggagttaga ggcccttggg gtttctcttt 6180
tccactgaca ggctttccca ggcagctggc tagttcattc cctcccagc cagggtgcagg 6240
cgtaggaata tggacatctg gttgctttgg cctgctgccc tctttcaggg gtccctaagcc 6300
cacaatcatg cctccctaag accttggcat ccttcctct aagccgttgg cacctctgtg 6360
ccacctctca cactggctcc agacacacag cctgtgcttt tggagctgag atcactcgct 6420
tcaccctcct catctttggt ctccaagtaa agccacgagg tcggggcgag ggcagaggtg 6480
atcacctgcg tgtcccactc acagacctgc ggcttcataa aacttctgat ttctctcag 6540
ctttgaaaag ggttaccctg ggcactggcc tagagcctca cctcctaata gacttagccc 6600
catgagtttg ccatgttgag caggactatt tctggcactt gcaagtccca tgatttcttc 6660
ggtaattctg aggggtgggg gagggacatg aatcatctt agcttagctt tctgtctgtg 6720
aatgtctata tagtgtattg tgtgttttaa caaatgattt aactgactg ttgctgtaaa 6780
agtgaatttg gaaataaagt tattactctg attaaa 6816

```

```

<210> SEQ ID NO 7
<211> LENGTH: 212
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 7

```

```

catgagtttg ccatgttgag caggactatt tctggcactt gcaagtccca tgatttcttc 60
ggtaattctg aggggtgggg gagggacatg aatcatctt agcttagctt tctgtctgtg 120
aatgtctata tagtgtattg tgtgttttaa caaatgattt aactgactg ttgctgtaaa 180
agtgaatttg gaaataaagt tattactctg at 212

```

```

<210> SEQ ID NO 8
<211> LENGTH: 240
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 8

```

```

acggccgagc ggccggggcgc tcgcgcgccg ccactagtgg ccggaggaga aggctcccgc 60
ggaggccgcg ctgcccgccc cctcccctgg ggaggctcgc gtteccgctg ctgcccctg 120
cgccgcccgc cggcctcagg aacgcgcctt cttcgccggc gcgcccctc gcagtcaccg 180
ccaccacca gctccggcac caacagcagc gccgctgcca ccgccacct tctgccgccc 240

```

```

<210> SEQ ID NO 9
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 9

```

```

Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
1           5           10

```


-continued

<210> SEQ ID NO 10
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 10

Arg Arg Gln Arg Arg Thr Ser Lys Leu Met Lys Arg
 1 5 10

<210> SEQ ID NO 11
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 11

Gly Trp Thr Leu Asn Ser Ala Gly Tyr Leu Leu Gly Lys Ile Asn Leu
 1 5 10 15

Lys Ala Leu Ala Ala Leu Ala Lys Lys Ile Leu
 20 25

<210> SEQ ID NO 12
 <211> LENGTH: 33
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 12

Lys Ala Leu Ala Trp Glu Ala Lys Leu Ala Lys Ala Leu Ala Lys Ala
 1 5 10 15

Leu Ala Lys His Leu Ala Lys Ala Leu Ala Lys Ala Leu Lys Cys Glu
 20 25 30

Ala

<210> SEQ ID NO 13
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 13

Arg Gln Ile Lys Ile Trp Phe Gln Asn Arg Arg Met Lys Trp Lys Lys
 1 5 10 15

<210> SEQ ID NO 14
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 14

Arg Lys Lys Arg Arg Gln Arg Arg Arg
 1 5

<210> SEQ ID NO 15

-continued

```

<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 15

```

```

Tyr Ala Arg Ala Ala Ala Arg Gln Ala Arg Ala
1           5           10

```

```

<210> SEQ ID NO 16
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 16

```

```

Thr His Arg Leu Pro Arg Arg Arg Arg Arg Arg
1           5           10

```

```

<210> SEQ ID NO 17
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 17

```

```

Gly Gly Arg Arg Ala Arg Arg Arg Arg Arg Arg
1           5           10

```

What is claimed is:

1. A method of increasing the functionality of a GABAergic interneuron in the hilus of the hippocampus of an individual having at least one apolipoprotein E4 (apoE4) allele, the method comprising reducing the level of tau in the interneuron.

2. The method of claim 1, wherein said reducing comprises administering to the individual a tau-specific interfering nucleic acid that reduces the level of a tau polypeptide in the GABAergic interneuron.

3. The method of claim 1, wherein said tau-specific interfering nucleic acid is encoded by a nucleotide sequence operably linked to a neuron-specific transcriptional control element, wherein the nucleotide sequence is present in a recombinant expression vector.

4. The method of claim 2, wherein said administering is via a local route of administration.

5. The method of claim 4, wherein said administering is via intracranial administration.

6. The method of claim 1, wherein said reducing comprises administering to the individual a genetically modified stem cell that has a reduced level of tau.

7. The method of claim 6, wherein the genetically modified stem cell has a knockout of an endogenous tau-encoding nucleic acid.

8. The method of claim 6, wherein the genetically modified stem cell is generated from a host cell from a donor individual who is the same as the individual being treated.

9. The method of claim 6, wherein the genetically modified stem cell is generated from a host cell from a donor individual who is other than the individual being treated.

10. The method of claim 6, wherein said genetically modified stem cell is a genetically modified induced pluripotent stem cell.

11. The method of claim 6, wherein said genetically modified stem cell is a genetically modified neural stem cell (NSC).

12. The method of claim 11, wherein the genetically modified NSC is an induced NSC (iNSC).

13. The method of claim 12, wherein the iNSC is induced from a somatic cell obtained from the individual being treated.

14. The method of claim 1, wherein said increasing the functionality of a GABAergic interneuron results in an increase in cognitive function in the individual.

15. The method of claim 14, wherein said cognitive function is learning or memory.

16. A genetically modified stem cell that exhibits reduced production of a tau polypeptide compared to a parent stem cell.

17. The genetically modified stem cell of claim 16, wherein said genetically modified stem cell is a neural stem cell.

18. The genetically modified stem cell of claim 17, wherein said genetically modified stem cell is an induced neural stem cell.

* * * * *