



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2032907 C 2002/05/14

(11)(21) **2 032 907**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) **Date de dépôt/Filing Date:** 1990/12/21

(41) **Mise à la disp. pub./Open to Public Insp.:** 1991/06/23

(45) **Date de délivrance/Issue Date:** 2002/05/14

(30) **Priorités/Priorities:** 1989/12/22 (456,162) US;
1990/11/07 (610,180) US

(51) **Cl.Int.⁶/Int.Cl.⁶** C07K 14/18, G01N 33/576, G01N 33/543

(72) **Inventeurs/Inventors:**

Lesniewski, Richard R., US;
Mehta, Smriti U., US;
Desai, Suresh M., US;
Casey, James M., US;
Devare, Sushil G., US;
Sarin, Virender K., US;
Leung, Tat, US

(73) **Propriétaire/Owner:**

ABBOTT LABORATORIES, US

(74) **Agent:** GOUDREAU GAGE DUBUC

(54) **Titre :** TEST POUR LA RECHERCHE DU VIRUS DE L'HEPATITE C

(54) **Title:** HEPATITIS C ASSAY

(57) **Abrégé/Abstract:**

The present invention provides an improved assay for detecting the presence of an antibody to an HCV antigen in a sample by contacting the sample with polypeptide containing at least one epitope of an HCV antigen. Preferred assay formats include a confirmatory assay, a combination assay, a synthetic polypeptide-based assay, an immunodot assay, and a competition assay.



-44-

ABSTRACT

HEPATITIS C ASSAY

5

The present invention provides an improved assay for detecting the presence of an antibody to an HCV antigen in a sample by contacting the sample with polypeptide containing at least one epitope of an HCV antigen. Preferred assay formats include a confirmatory assay, a combination assay, a synthetic polypeptide-based assay, an immunodot assay, and a competition assay.

15

20

25

30

35

-1-

HEPATITIS C ASSAY

This invention relates generally to an assay for identifying the presence in a sample of an antibody which is immunologically reactive with a hepatitis C virus antigen and specifically to an assay for detecting a complex of an antibody and a polypeptide having at least one epitope of a hepatitis C virus antigen.

BACKGROUND

Acute viral hepatitis is clinically diagnosed by a well-defined set of patient symptoms, including jaundice, hepatic tenderness, and an increase in the serum levels of alanine aminotransferase and aspartate aminotransferase. Additional serologic immunoassays are generally performed to diagnose the specific type of viral causative agent. Historically, patients presenting clinical hepatitis symptoms and not otherwise infected by hepatitis A, hepatitis B, Epstein-Barr or cytomegalovirus were clinically diagnosed as having non-A non-B hepatitis (NANBH) by default. The disease may result in chronic liver damage.

Each of the well-known, immunologically characterized hepatitis-inducing viruses, hepatitis A virus (HAV), hepatitis B virus (HBV), and hepatitis D virus (HDV) belongs to a separate family of viruses and has a distinctive viral organization, protein structure, and mode of replication.

Attempts to identify the NANBH virus by virtue of genomic similarity to one of the known hepatitis viruses have failed, suggesting that NANBH has a distinct organization and structure. [Fowler, et al., J. Med. Virol., 12:205-213 (1983) and Weiner, et al., Med. Virol., 21:239-247 (1987)].

Progress in developing assays to detect antibodies specific for NANBH has been particularly hampered by difficulties in correctly identifying antigens associated with NANBH. See, for example, Wands, J., et al., U.S. Patent 4,870,076, Wands, et al., Proc. Nat'l. Acad. Sci., 83:6608-6612 (1986), Ohori, et al., J. Med. Virol., 12:161-178 (1983), Bradley, et al., Proc. Nat'l. Acad. Sci., 84:6277-6281, (1987), Akatsuka, T., et al., J. Med.

Virol, 20:43-56 (1986), Seto, B., et al., WO90-02206

Takahashi, K., et al., European Patent Application No. 0 293 274, published November 30, 1988, and Seelig, R., et al., in PCT Application PCT/EP88/00123.

Recently, another hepatitis-inducing virus has been unequivocally identified as hepatitis C virus (HCV) by Houghton, M., et al., European Patent Application publication number 0 318 216, May 31, 1989. Related papers describing this virus include Kuo, G., et al., Science, 244:359-361 (1989) and Choo, Q., et al., Science, 244:362-364 (1989). Houghton, M., et al. reported isolating cDNA sequences from HCV which encode antigens which react immunologically with antibodies present in patients infected with NANBH, thus establishing that HCV is the viral agent causing NANBH.

The cDNA sequences associated with HCV were isolated from a cDNA library prepared from the RNA obtained from pooled serum from a chimpanzee with chronic HCV infection. The cDNA library contained cDNA sequences of approximate mean size of about 200 base pairs. The cDNA library was screened for encoded epitopes expressed in clones that could bind to antibodies in sera from patients who had previously experienced NANBH.

In the European Patent Application, Houghton, M., et al. also described the preparation of several superoxide dismutase fusion polypeptides (SOD) and the use of these SOD fusion polypeptides to develop an HCV screening assay. The most complex SOD fusion polypeptide described in the European Patent Application, designated C100-3, was described as containing 154 amino acids of human SOD at the aminoterminal, 5 amino acid residues derived from the expression of a synthetic DNA adapter containing a restriction site, EcoRI, 363 amino acids derived from the expression of a cloned HCV cDNA fragment, and 5 carboxy terminal amino acids derived from an MS2 cloning vector nucleotide sequence. The DNA sequence encoding this polypeptide was transformed into yeast cells using a plasmid. The

transformed cells were cultured and expressed a 54,000 molecular weight polypeptide which was purified to about 80% purity by differential extraction.

Other SOD fusion polypeptides designated SOD NANB₅₋₁₋₁, and SOD-NANB₈₁ were expressed in recombinant bacteria. The E.coli fusion polypeptides were purified by differential extraction and by chromatography using anion and cation exchange columns. The purification procedures were able to produce SOD-NANB₅₋₁₋₁ as about 80% pure and SOD-NANB₈₁ as about 50% pure.

The recombinant SOD fusion polypeptides described by Houghton, M., et al. were coated on microtiter wells or polystyrene beads and used to assay serum samples. Briefly, coated microtiter wells were incubated with a sample in a diluent. After incubation, the microtiter wells were washed and then developed using either a radioactively labelled sheep anti-human antibody or a mouse antihuman IgG-HRP (horseradish peroxidase) conjugate. These assays were used to detect both post acute phase and chronic phase HCV infection. Due to the preparative methods, assay specificity required adding yeast or E. coli extracts to the samples in order to prevent undesired immunological reactions with any yeast or E.coli antibodies present in samples.

Ortho Diagnostic Systems Inc. have developed a research immunoenzyme assay to detect antibodies to HCV antigens. The Ortho assay procedure is a three-stage test for serum/plasma carried out in a microwell coated with the recombinant yeast/hepatitis C virus SOD fusion polypeptide C100-3.

In the first stage, a test specimen is diluted directly in the test well and incubated for a specified length of time. If antibodies to HCV antigens are present in the specimen, antigen-antibody complexes will be formed on the microwell surface. If no antibodies are present, complexes will not be formed and the unbound serum or plasma proteins will be removed in a washing step.

In the second stage, anti-human IgG murine monoclonal antibody horseradish peroxidase conjugate is added to the

-4-

microwell. The conjugate binds specifically to the antibody portion of the antigen-antibody complexes. If antigen-antibody complexes are not present, the unbound conjugate will also be removed by a washing step.

In the third stage, an enzyme detection system composed of o-phenylenediamine 2HCl (OPD) and hydrogen peroxide is added to the test well. If bound conjugate is present, the OPD will be oxidized, resulting in a colored end product. After formation of the colored end product, dilute sulfuric acid is added to the microwell to stop the color-forming detection reaction.

The intensity of the colored end product is measured with a microwell reader. The assay may be used to screen patient serum and plasma.

It is established that HCV may be transmitted by contaminated blood and blood products. In transfused patients, as many as 10% will suffer from post-transfusion hepatitis. Of these, approximately 90% are the result of infections diagnosed as HCV. The prevention of transmission of HCV by blood and blood products requires reliable, sensitive and specific diagnosis and prognostic tools to identify HCV carriers as well as contaminated blood and blood products. Thus, there exists a need for an HCV assay which uses reliable and efficient reagents and methods to accurately detect the presence of HCV antibodies in samples.

BRIEF SUMMARY

The present invention provides an improved assay for detecting the presence of an antibody to an HCV antigen in a sample by contacting the sample with polypeptide containing at least one epitope of an HCV antigen.

One assay format according to the invention provides a confirmatory assay for unequivocally identifying the presence of an antibody that is immunologically reactive with an HCV antigen. Briefly, a fluid sample is used to prepare first and second aliquots. The aliquots are then contacted with at least two polypeptides duplicative of a continuous amino acid sequence putatively contained in proteins expressed by clones containing

HCV cDNA sequences containing at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide. Finally, the antibody-antigen complex is detected. The improvement comprises contacting the first aliquot with recombinant polypeptide C100-3, and contacting the second aliquot with one or more polypeptides selected from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866, pl899, p30, p380.LG, p447, p607, p607a, p643b, p666, p691 and p2302. Preferred polypeptides are selected from the group consisting of pl684, pl689, pl866, p380, p643b, p666, p2302 and p380.LG.

Another assay format provides a combination assay for detecting the presence of an antibody that is immunologically reactive with an HCV antigen in a fluid sample by contacting the sample with a polypeptide containing at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide and detecting the antibody-polypeptide complex. The improvement comprises contacting the sample with a solid support containing commonly bound recombinant polypeptide C100-3 and a polypeptide selected from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866, pl899, p380, p380.LG, p447, p607, p607a, p643b, p666, p691 and p2303.

Another assay format provides an assay for identifying the presence of an antibody that is immunologically reactive with an HCV antigen in a fluid sample comprising contacting the sample with a polypeptide containing at least one epitope of an HCV antigen selected from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866, pl899, p380, p380.LG, p447, p607, p607a, p643b, p666, p691 and p2303 under conditions suitable for complexing the antibody with the polypeptide and detecting the antibody-polypeptide complex.

Another assay format provides an immunodot assay for identifying the presence of an antibody that is immunologically reactive with an HCV antigen by concurrently contacting a sample with at least two polypeptides each containing distinct epitopes of an HCV antigen under conditions suitable for complexing the

antibody with at least one of the polypeptides and detecting the antibody-polypeptide complex by reacting the complex with color-producing reagents. The improvement comprises employing polypeptides selected from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866, pl899, p380, p380.LG, p447, p607, p607a, p643b, p666, p691, p2302 and C100-3. Preferred polypeptides are selected from the group consisting of pl684, pl694, pl684, pl866, p380, p643b, p666, p2302, p380.LG and C100-3.

Another assay format provides a competition assay directed to the confirmation that positive results are not false by identifying the presence of an antibody that is immunologically reactive with an HCV antigen in a fluid sample where the sample is used to prepare first and second immunologically equivalent aliquots. The first aliquot is contacted with solid support containing a bound polypeptide which contains at least one epitope of an HCV antigen under conditions suitable for complexing with the antibody to form a detectable antibody-polypeptide complex and the second aliquot is first contacted with unbound polypeptide and then contacted with the same, solid support containing bound polypeptide. The improvement comprises selecting the polypeptide from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866, pl899, p380, p380.LG, p447, p607, p607a, p643b, p666, p691 and p2302.

In all of the assays, the sample is diluted before contacting the polypeptide absorbed on a solid support. Samples may be obtained from different biological samples such as whole blood, serum, plasma, cerebral or spinal fluid, and lymphocyte or cell culture supernatants. Solid support materials may include cellulose materials, such as paper and nitrocellulose, natural and synthetic polymeric materials, such as polyacrylamide, polystyrene, and cotton, porous gels such as silica gel, agarose, dextran and gelatin, and inorganic materials such as deactivated alumina, magnesium sulfate and glass. Suitable solid support materials may be used in assays in a variety of well known

physical configurations, including microtiter wells, test tubes, beads, strips, membranes, and microparticles. A preferred solid support for a non-immunodot assay is a polystyrene bead. A preferred solid support for an immunodot assay is nitrocellulose.

Suitable methods and reagents for detecting an antibody-antigen complex in an assay of the present invention are commercially available or known in the relevant art. Representative methods may employ detection reagents such as enzymatic, radioisotopic, fluorescent, luminescent, or chemiluminescent reagents. These reagents may be used to prepare hapten-labelled antihapten detection systems according to known procedures, for example, a biotin-labelled antibiotin system may be used to detect an antibody-antigen complex.

The present invention also encompasses assay kits including polypeptides which contain at least one epitope of an HCV antigen bound to a solid support as well as needed sample preparation reagents, wash reagents, detection reagents and signal producing reagents.

Other aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following detailed description which provides illustrations of the invention in its presently preferred embodiments.

DESCRIPTION OF THE DRAWINGS

FIGURES 1a and 1b illustrate the HCV genome.

FIGURE 2 illustrates the use of antigenic polypeptides to identify the presence of antibodies in a chimpanzee inoculated with HCV.

FIGURES 3a and 3b illustrate the sensitivity increase using a combination assay format.

FIGURE 4 illustrates a test cartridge for an immunodot assay.

FIGURE 5 illustrates a seroconversion graph wherein the amount of anti-NS-5 S/N antibody, shown as the solid line between closed circles, and the amount of anti-HCV 2.0 S/CO antibody

shown as the solid lines between open squares, is plotted against days post presentation.

DETAILED DESCRIPTION

The present invention is directed to an assay to detect an antibody to an HCV antigen in a sample. Human serum or plasma is diluted in a sample diluent and incubated with a polystyrene bead coated with a polypeptide that includes an HCV antigenic epitope. If antibodies are present in the sample they will form a complex with the antigenic polypeptide and become affixed to the polystyrene bead. After the complex has formed, unbound materials and reagents are removed by washing the bead and the bead-antigen-antibody complex is reacted with a solution containing horseradish peroxidase labeled goat antibodies directed against human antibodies. This peroxidase enzyme then binds to the antigen-antibody complex already fixed to the bead. In a final reaction the horseradish peroxidase is contacted with o-phenylenediamine and hydrogen peroxide which results in a yellow-orange color. The intensity of the color is proportional to the amount of antibody which initially binds to the antigen fixed to the bead. The preferred polypeptides having HCV antigenic epitopes were selected from portions of the HCV genome which encoded polypeptides which possessed amino acid sequences similar to other known immunologically reactive agents and which were identified as having some immunological reactivity. (The immunological reactivity of a polypeptide was initially identified by reacting the cellular extract of E. coli clones which had been transformed with cDNA fragments of the HCV genome with HCV infected serum. The clones presumably expressed polypeptides encoded by the incorporated cDNA which were immunologically reactive with serum known to contain antibody to HCV antigens.) An analysis of a given amino acid sequence, however, only provides rough guides to predicting immunological reactivity. There is no invariably predictable way to ensure immunological activity short of preparing a given amino acid sequence and testing the suspected sequence in an assay. As

-9-

Biosystems' small scale rapid cycle protocol. Protected amino acids were coupled using preformed symmetric anhydride chemistry except for asparagine, glutamine, arginine and histidine which were double coupled using N-N'-dicyclohexylcarbodiimide (DCC)/1-hydroxybenzotriazole (HOBT) chemistry. In the first coupling, protected amino acids were coupled using preformed symmetric anhydrides dissolved in dimethyl-formamide (DMF). The symmetric anhydride of an individual amino acid was formed in methylene chloride followed by solvent exchange to DMF before transferring to the reaction vessel of the peptide synthesizer. The second coupling of symmetric anhydride was also conducted in DMF. The N-amino group of all amino acids used was protected by a t-butyloxycarbonyl (t-BOC) linkage. The side chain functional groups of various amino acids were protected by the following groups:

Arg-Tos	(Tosyl)
Lys-2ClZ	(2-chlorobenzyloxycarbonyl)
Thr,Ser-Bzl	(Benzyl)
Tyr-2BrZ	(2-Bromobenzyloxycarbonyl)
Cys-4MeBzl	(4-Methylbenzyl)
Asp,Glu-OBzl	(O-Benzyl)
His-DNP	(Dinitrophenyl)

The fully protected peptide-resin (0.28g) was allowed to swell in methylene chloride (CH_2Cl_2) for 5 minutes. The peptide-resin was transferred to a manual reaction vessel, treated twice with 5% thiophenol in DMF for twenty minutes each followed by six CH_2Cl_2 washes for one minute each, and then transferred to the reaction vessel of the synthesizer. The t-BOC protecting group was then removed using 60% TFA/ CH_2Cl_2 according to the manufacturer's protocol and the partially deprotected peptide-resin was then dried overnight under house vacuum at room temperature.

Partially deprotected peptide-resin was then treated with dimethyl sulfide (DMS (1 ml), p-cresol (1 ml), p-thiocresol (0.2 g) and HF (10 mL) at 0°C for one hour to cleave the peptide from the resin support. The HF/DMS and other volatiles were distilled

-10-

off in vacuo at 0°C. The cleaved peptide and resin were washed three times with 15 ml aliquots of diethyl ether, and the cleaved peptide was extracted by washing three times each with 10 ml aliquots of 40% aqueous acetic acid and 15% aqueous acetic acid, respectively. The aqueous extracts were combined and washed three times with 15 ml aliquots of diethyl ether and then lyophilized to yield a crude peptide.

The crude peptide was analyzed for purity using reversed-phase high performance liquid chromatography on a C₄, 4.6 x 30mm column (Brownlee, Applied Biosystems, Inc., Foster City, California), flow rate one ml/minute employing 0.1% aqueous TFA (A) and 100% acetonitrile (B) as the solvent system. The preferred solvent gradient employed for this peptide analysis started with 30% B solvent. The column was maintained at 30% B for one minute followed by an increase over 20 minutes using a linear gradient to 55% B and maintained for one minute. Finally, the column was brought back to 30% B over a two minute period. The presence of peptide in the effluent was monitored simultaneously at 225 nm and 280 nm. The composition of the purified peptide was determined by acid hydrolysis. After removal of the acid, the hydrolysate was analyzed on a Beckmar 6300 amino acid analyzer.

If increased quantities of purified polypeptide were desired, semi-preparative reversed phase high performance liquid chromatography was performed in a similar manner using a C₄, 10 x 100 mm column (Brownlee, Applied Biosystems Inc., Foster City, California) using the same aqueous 0.1% TFA (A) and 100% acetonitrile (B) solvent system described above. The preferred solvent gradient for a semi-preparative run started with 27% B at 3 ml/minute for two minutes followed by an increase over 20 minutes using a linear gradient to 50% B. The concentration was maintained at 50% B for one minute and then reduced to 27% B within one minute.

Other peptides described herein were assembled on solid support in a manner analogous to the synthesis described above. The amino acids tryptophan and methionine, if present, were used

-11-

without any side chain protection. Usually, after incorporating methionine during the chain assembly, ethanedithiol (0.1% v/v) was added to TFA for all subsequent removal of t-BOC groups. However, if histidine protected by DNP was present in the sequence, ethanedithiol was not added to TFA; instead, indole (1% w/v) was used. Also, after incorporating tryptophan, indole (1% w/v) was added to the TFA solution.

HF cleavage from the resin and purification of the peptides were achieved essentially as described above.

The peptides synthesized as described above were evaluated for their antigenic/immunogenic properties. A summary of the amino acid sequences, beginning with the amino terminus and ending with the carboxy terminus, of immunologically reactive peptides is presented in Tables 2 and 3.

TABLE 1

CHIRON UPDATE	GENOMIC REGION	SAMPLE A00642 DILUTION	RESULT	SAMPLE #401 DILUTION	RESULT	SAMPLE #423 DILUTION	RESULT
1694-1735	C-100	1:800	POS.	1:40	POS.	1:20	POS.
1866-1930	C-100	1:100	POS.	-	-	-	-
1689-1805	C-100	1:500	POS.	1:40	POS.	1:20	POS.
1684-1750	C-100	1:500	POS.	1:40	POS.	1:20	POS.
1899-1930	C-100	1:50	POS.	1:40	POS.	1:20	NEG.
1192-1240	33C	1:800	POS.	1:40	POS.	1:20	POS.
1223-1240	33C	1:200	POS.	1:40	POS.	1:20	POS.
1-75	Putative Core	1:100	POS.	1:40	POS.	1:20	POS.
35-75	Putative Core	1:50	NEG.	1:40	NEG.	1:20	POS.
99-126	Putative Core	1:50	NEG.	1:40	NEG.	1:20	POS.
1569-1593	C-100	1:25	NEG.	1:40	NEG.	1:20	NEG.
1357-1407	33C	1:25	NEG.	1:40	NEG.	1:20	NEG.
1418-1457	33C	1:25	NEG.	1:40	NEG.	1:20	NEG.
195-262	Putative Core/ Envelope	1:50	NEG.	1:40	NEG.	1:20	NEG.
230-262	Putative Core/ Envelope	1:25	NEG.	1:40	NEG.	1:20	NEG.

A sample is considered positive if the absorbance at 492 nm $\geq$ 4X absorbance value of the negative control (S/N $\geq$ 4.0).

A00642 Human plasma sample convalescent from NANB (HCV) Hepatitis. Patient was clinically diagnosed with NANB and was negative for HBV and HAV markers.

#401 Human paid plasma donor positive by screening assays based on C100-3. No known clinical history.

#423 Human paid plasma donor positive by screening assays based on C100-3. No known clinical history.

-13-

TABLE 2

(Note: H signifies the amino terminus;
OH signifies the carboxyl terminus.)

5	p1 (1-75)	H-M-S-T-N-P-K-P-Q-K-K-N-K-R-N-T- N-R-R-P-Q-D-V-K-F-P-G-G-G-Q-I- V-G-G-V-Y-L-L-P-R-R-G-P-R-L- G-V-R-A-T-R-K-T-S-E-R-S-Q- P-R-G-R-R-Q-P-I-P-K-A-R-R-P- E-G-R-T-OH
10	p35 (35-75)	H-Y-L-L-P-R-R-G-P-R-L-G-V-R-A-T-R-K-T- S-E-R-S-Q-P-R-G-R-R-Q-P-I-P-K-A-R-R- P-E-G-R-T-OH
15	p99 (99-126)	H-S-P-R-G-S-R-P-S-W-G-P-T-D-P-R-R-R-S- R-N-L-G-K-V-I-D-T-L-OH
20	p195 (195-262)	H-R-N-S-T-G-L-Y-H-V-T-N-D-C-P-N-S-S-I-V-Y- E-A-A-D-A-I-L-H-T-P-G-C-V-P-C-V-R-E-G-N-A- S-R-C-W-V-A-M-T-P-T-V-A-T-R-D-G-K-L-P-A-T- Q-L-R-R-H-I-OH
25	p230 (230-262)	H-V-R-E-G-N-A-S-R-C-W-V-A-M-T-P-T-V-A-T- R-D-G-K-L-P-A-T-Q-L-R-R-H-I-OH
30	p1192 (1192-1240)	H-A-V-D-F-I-P-V-E-N-L-E-T-T-M-R-S-P-V- F-T-D-N-S-S-P-P-V-V-P-Q-S-F-Q-V-A-H- L-H-A-P-T-G-S-G-K-S-T-K-V-OH
	p1223 (1223-1240)	H-F-Q-V-A-H-L-H-A-P-T-G-S-G-K-S-T-K-V-OH
35	p1357 (1357-1407)	H-Y-V-P-H-P-N-I-E-E-V-A-L-S-T-T-G-E-I-P-F- Y-G-K-A-I-P-L-E-V-I-K-G-G-R-H-L-I-F-C- H-S-K-K-K-C-D-E-L-A-A-K-L-OH

-14-

TABLE 2 (CONT'D)

(Note: H signifies the amino terminus;
OH signifies the carboxyl terminus.)

5	p1418 (1418-1457)	H-R-G-L-D-V-S-V-I-P-T-S-G-D-V-V-V- V-A-T-D-A-L-M-T-G-Y-T-G-D-F-D-S-V- I-D-C-N-T-C-OH
10	p1569 (1569-1593)	H-D-A-H-F-L-S-Q-T-K-Q-S-G-E-N-L-P-Y-L-V- A-Y-Q-A-T-V-OH
15	p1684 (1684-1750)	H-G-R-V-V-L-S-G-K-P-A-I-I-P-D-R-E-V-L-Y R-E-F-D-E-M-E-E-C-S-Q-H-L-P-Y-I-E-Q-G-M M-L-A-E-Q-F-K-Q-K-A-L-G-L-L-Q-T-A-S-R- Q-A-E-V-I-A-P-A-V-OH
20	p1689 (1689-1805)	H-S-G-K-P-A-I-I-P-D-R-E-V-L-Y-R-E-F D-E-M-E-E-C-S-Q-H-L-P-Y-I-E-Q-G-M- M-L-A-E-Q-F-K-Q-K-A-L-G-L-L-Q-T- A-S-R-Q-A-E-V-I-A-P-A-V-Q-T-N-W- Q-K-L-E-T-F-W-A-K-H-M-W-N-F-I-S- G-I-Q-Y-L-A-G-L-S-T-L-P-G-N-P-A- I-A-S-L-M-A-F-T-A-A-V-T-S-P-L-T-T-S-Q-OH
25	p1694 (1694-1735)	H-I-I-P-D-R-E-V-L-Y-R-E-F-D-E- M-E-E-C-S-Q-H-L-P-Y-I-E-Q-G- M-M-L-A-E-Q-F-K-Q-K-A-L-G-L-OH
30	p1866 (1866-1930)	H-F-K-I-M-S-G-E-V-P-S-T-E-D-L-V-N- L-L-P-A-I-L-S-P-G-A-L-V-V-G-V-V- C-A-A-I-L-R-R-H-V-G-P-G-E-G-A-V- Q-W-M-N-R-L-I-A-F-A-S-R-G-N-H-V-S-OH
35	p1899 (1899-1930)	H-A-A-I-L-R-R-H-V-G-P-G-E-G-A-V- Q-W-M-N-R-L-I-A-F-A-S-R-G-N-H-V- S-OH

-15-

TABLE 8

(NOTE: H signifies the amino terminus; OH signifies the carboxyl terminus. The two underlined Tyr residues are not part of the HCV sequence but are engineered there for ease of iodinating the peptide at a later time).

p380 (380-436)	H	-Gly-Val-Asp-Ala-Glu-Thr-His-Val-Thr-Gly-Gly-Ser-Ala-Gly-His-Thr-Val-Ser-Gly-Phe-Val-Ser-Leu-Leu-Ala-Pro-Gly-Ala-Lys-Gln-Asn-Val-Gln-Leu-Ile-Asn-Thr-Asn-Gly-Ser-Trp-His-Leu-Asn-Ser-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Leu-Asn-Thr-Gly- OH
p380.LG (380-436.LG)	H	-Gly-Val-Asp-Ala-Glu-Thr-His-Val-Thr-Gly-Gly-Ser-Ala-Gly-His-Thr-Val-Ser-Gly-Phe-Val-Ser-Leu-Leu-Ala-Pro-Gly-Ala-Lys-Gln-Asn-Val-Gln-Leu-Ile-Asn-Thr-Asn-Gly-Ser-Trp-His-Leu-Asn-Ser-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Leu-Asn-Thr-Gly- OH
p447 (447-483)	H	-Phe-Asn-Ser-Ser-Gly-Cys-Pro-Glu-Arg-Leu-Ala-Ser-Cys-Arg-Pro-Leu-Thr-Asp-Phe-Asp-Gln-Gly-Trp-GlyPro-Ile-Ser-Tyr-Ala-Asn-Gly-Ser-Gly-Pro-Asp-Gln-Arg- OH
p607 (607-627)	H	-Cys-Leu-Val-Asp-Tyr-Pro-Tyr-Arg-Leu-Trp-His-Tyr-Pro-Cys-Thr-Ile-Asn-Tyr-Thr-Ile-Phe- OH
p643a (643-663)	H	-Ala-Cys-Asn-Trp-Thr-Arg-Gly-Glu-Arg-Cys-Asp-Leu-Glu-Asp-Arg-Asp-Arg-Ser-Glu-Leu-Ser- <u>Tyr</u> -OH
p643b (643-683)	H	-Ala-Cys-Asn-Trp-Thr-Arg-Gly-Glu-Arg-Cys-Asp-Leu-Glu-Asp-Arg-Asp-Arg-Ser-Glu-Leu-Ser-Pro-Leu-Leu-Leu-Thr-Thr-Gln-Trp-Gln-Val-Leu-Pro-Cys-Ser-Phe-Thr-Thr-Leu-Pro- OH
p666 (666-683)	H	-Leu-Leu-Thr-Thr-Thr-Gln-Trp-Gln-Val-Leu-Pro-Cys-Ser-Phe-Thr-Thr-Leu-Pro- <u>Tyr</u> -OH
p691 (691-714)	H	-His-Leu-His-Gln-Asn-Ile-Val-Asp-Val-Gln-Tyr-Leu-Tyr-Gly-Val-Gly-Ser-Ser-Ile-Ala-Ser-Trp-Ala-Ile- OH
p2302 (2302-2352)	H	-Lys-Lys-Pro-Asp-Tyr-Gln-Pro-Pro-Val-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Pro-Lys-Ser-Pro-Pro-Val-Pro-Pro-Pro-Lys-Lys-Lys-Arg-Thr-Val-Val-Leu-Thr-Glu-Ser-Thr-Leu-Ser-Thr-Ala-Leu-Ala-Glu-Leu-Ala-Thr-Arg-Ser-Phe- OH

-16-

The polypeptides illustrated in Tables 2 and 8 may also be prepared in a stepwise fashion or in a fragment coupling protocol using various side chain protection methodologies known to those skilled in the art. The polypeptides may also be prepared using enzymatic methodology.

Further, the polypeptides useful in the practice of this invention may be prepared using recombinant technologies. Briefly, DNA sequences which encode the desired polypeptides are preferably assembled from fragments of the total desired sequence. The fragments are generally prepared using well known automated processes and apparatus. After the complete sequence has been prepared the desired sequence is incorporated into an expression vector which is transformed into a host cell. The DNA sequence is then expressed by the host cell to give the desired polypeptide which is harvested from the host cell or from the medium in which the host cell is cultured. In most cases, the manufactured DNA sequence is assembled using codons which are known to be best expressed in the host cell. When smaller peptides are to be made using recombinant technologies it may be advantageous to prepare a single DNA sequence which encodes several copies of the desired polypeptide in a connected chain. The long chain is then isolated and the chain is cleaved into the shorter, desired sequences.

The amino acid sequence for pl684 is reverse translated to give the codons listed in Table 3 which are optimized (where not inconsistent with assembly and synthesis of fragments) to facilitate high level expression in E. coli. Individual oligonucleotides are synthesized on Applied Biosystem 380A DNA synthesizer using methods and reagents recommended by the manufacturer. These purified oligonucleotides are annealed and ligated together to assemble the entire DNA sequence for digestion with BamH1 and Sall, allowing ligation into pUC18. The resulting plasmid is suitably transformed into E. coli JM103 cells. Table 3 also lists preferred codons to express pl and pl223.

p1

p1223

p1634

				5					10					15					20	
Gly	Arg	Val	Val	Leu	Ser	Gly	Lys	Pro	Ala	Ile	Ile	Pro	Asp	Arg	Glu	Val	Leu	Tyr	Arg	
GGT	CGT	GTT	GTT	CTG	TCT	GGT	AAA	CCG	GCT	ATC	ATC	CCG	GAC	CGT	GAA	GTT	CTG	TAC	CGT	
				25					30					35					40	
Glu	Phe	Asp	Glu	Met	Glu	Glu	Cys	Ser	Gln	His	Leu	Pro	Tyr	Ile	Glu	Gln	Gly	Met	Met	
GAA	TTC	GAC	GAA	ATG	GAA	GAA	TGC	TCT	CAG	CAC	CTG	CCG	TAC	ATC	GAA	CAG	GGT	ATG	ATG	
				45					50					55					60	
Leu	Ala	Glu	Gln	Phe	Lys	Gln	Lys	Ala	Leu	Gly	Leu	Leu	Gln	Thr	Ala	Ser	Arg	Gln	Ala	
CTG	GCT	GAA	CAG	TTC	AAA	CAG	AAA	GCT	CTG	GGT	CTG	CTG	CAG	ACC	GCT	TCT	CGT	CAG	GCT	
				65																
Glu	Val	Ile	Ala	Pro	Ala	Val														
GAA	GTT	ATC	GCT	CCG	GCT	GTT														

-18-

The polypeptides illustrated in Tables 2 and 8 may also be prepared in a stepwise fashion or in a fragment coupling protocol using various side chain protection methodologies known to those skilled in the art. The polypeptides may also be prepared using enzymatic methodology.

Further, the polypeptides useful in the practice of this invention may be prepared using recombinant technologies. Briefly, DNA sequences which encode the desired polypeptides are preferably assembled from fragments of the total desired sequence. The fragments are generally prepared using well known automated processes and apparatus. After the complete sequence has been prepared the desired sequence is incorporated into an expression vector which is transformed into a host cell. The DNA sequence is then expressed by the host cell to give the desired polypeptide which is harvested from the host cell or from the medium in which the host cell is cultured. In most cases, the manufactured DNA sequence is assembled using codons which are known to be best expressed in the host cell. When smaller peptides are to be made using recombinant technologies it may be advantageous to prepare a single DNA sequence which encodes several copies of the desired polypeptide in a connected chain. The long chain is then isolated and the chain is cleaved into the shorter, desired sequences.

The amino acid sequence for pl684 is reverse translated to give the codons listed in Table 3 which are optimized (where not inconsistent with assembly and synthesis of fragments) to facilitate high level expression in E. coli. Individual oligonucleotides are synthesized on Applied Biosystem 380A DNA synthesizer using methods and reagents recommended by the manufacturer. These purified oligonucleotides are annealed and ligated together to assemble the entire DNA sequence for digestion with BamHI and Sall, allowing ligation into pUC18. The resulting plasmid is suitably transformed into E. coli JM103 cells. Table 3 also lists preferred codons to express pl and pl223.

-19-

In order to establish that a clone expresses the DNA sequence, it is grown at 37 °C in 50 ml Luria Broth, in a 250 ml Erlenmeyer flask. When the culture reaches an OD600 of 0.3-0.5, IPTG is added to a final concentration of 1 mM to induce expression. Samples (1.5 ml) are removed at one hour intervals, and the cells are pelleted and resuspended to an OD600 of 10.0 in 2X SDS/PAGE loading buffer. Aliquots (15 ul) of the prepared samples are loaded on a 15% SDS/PAGE gel, the expressed polypeptides separated, and then electrophoretically transferred to nitrocellulose for immunoblotting. The nitrocellulose sheet containing the transferred proteins is incubated with a blocking solution for one hour and incubated overnight at 4 °C with HCV patients' sera diluted in TBS containing 5% E. coli JM103 lysate. The nitrocellulose sheet is washed three times in TBS, then incubated with HRP0-labeled goat anti-human IgG, diluted in TBS containing 10% fetal calf sera. The nitrocellulose is washed three times with TBS and the color is developed in TBS containing 2 mg/ml 4-chloro-1-naphthol, 0.02% hydrogen peroxide and 17% methanol. Strong immunoreactive band formation with HCV patients' sera indicates that the synthetic polypeptide is expressed in E. coli in immunologically reactive form.

Preferred formats for assays using the polypeptides described above are provided in the following examples. Example 1 describes a confirmatory assay. Example 2 describes a combination assay. Example 3 describes a synthetic polypeptide-based assay. Example 4 describes an immunodot assay. Example 5 describes a competition assay. Example 5 describes an EIA assay in which peptides 380-436 and 447-483, 643-683 and 2302-2352 are used. Example 6 describes an EIA utilizing peptide p380.LG. Example 7 describes an EIA utilizing peptide 2302 (NS-5) compared to an EIA utilizing antigens NS3 (CKS-33C), NS4 (C-100) or CORE (CRS-CORE). Example 8 describes the PEPSCAN^{*} protocol followed.

* Trademark

Example 1. CONFIRMATORY ASSAY

The confirmatory assay uses at least two polypeptides containing HCV antigenic epitopes which are preferably prepared and isolated from different sources. One polypeptide is used to screen serum or plasma samples. The other polypeptide is used to confirm the presence of a HCV antibody in a sample initially identified as containing a HCV antibody by the screening procedure.

In the presently preferred confirmatory assay, the screening procedure uses a recombinant C100-3 polypeptide. The C100-3 recombinant polypeptide is believed to contain multiple epitopes as well as an immunodominant region defined by the 1689-1806 amino acid sequence. The C100-3 polypeptide is expressed in recombinant yeast cells and isolated from the cell extract as described in EPA Publication Number 0 318 216. Other recombinant polypeptides containing amino acid sequences essentially duplicative of C100-3 may also be used.

The other peptide used in the confirmatory assay is a synthetic peptide selected from the group consisting of pl, p35, p99, pl192, pl223, pl684, pl689, pl694, pl866 and pl899. Preferably the peptide is pl684 or pl866. These peptides were prepared following procedures described above. In the confirmatory assay, both C100-3 and the synthetic peptides, pl684, pl694 or pl866, were separately coated onto polystyrene beads. A combination of synthetic peptides coated on a polystyrene bead may also be used if desired. The polystyrene beads are first washed with distilled water and propanol then incubated with crude or purified HCV synthetic peptides diluted to 0.1-20.0 ug/ml in a 0.1 M solution of an appropriate buffer containing about 0.4-0.5 M NaCl, about 0.0022% Triton X-100 and adjusted to about pH 6.5-10.0. The following buffers, tris, $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, boric acid, and citrate buffers are preferred and are optimized for each peptide; preferred buffers, pH and coating concentration for the synthetic peptides are listed in Table 4. Successful coatings have also been accomplished with lower or higher pH. The beads are incubated in the antigen solution for

-21-

about two hours at 38-42 °C, washed in phosphate buffer solution (PBS) and soaked in 0.1% Triton X-100 in PBS for sixty minutes at 38-42 °C. The beads are then washed two times in PBS, overcoated with a solution of 5% (w/v) bovine serum albumin in PBS for sixty minutes and washed three times with PBS. Finally, the beads are overcoated with 5% (w/v) sucrose in PBS and dried under nitrogen or air.

The peptides are each individually coated onto polystyrene beads and used in an antibody capture format. Ten microliters of sample are added to the wells of a reaction tray along with 400 ul of a sample diluent and a peptide coated bead. The sample diluent consists of about 10% (v/v), or less, bovine serum and about 20% (v/v), or less, goat serum in 20 mM Tris phosphate buffer containing 0.20%, or less, (v/v) Triton X-100, 3% (w/v), or less, bovine serum albumin. When the recombinant yeast C100-3 polypeptide is used, antibodies to yeast antigens which may be present in a sample are reacted with yeast extracts which are added to the sample diluent (typically about 200 ug/ml). The addition of yeast extracts to the sample diluent is used to prevent false positive results. The final material is sterile filtered and filled in plastic bottles, and preserved with 0.1% sodium azide.

After one hour of incubation at 40 °C, the beads are washed and 200 ul of conjugate is added to the wells of the reaction tray.

The preferred conjugate is goat anti-human IgG horseradish peroxidase conjugate. Concentrated conjugate is purchased from Kirkegaard and Perry Laboratories, Inc., Gaithersburg, Maryland, and titered to determine a working concentration. A twenty-fold concentrate of the working conjugate solution is then prepared by diluting the concentrate in diluent. The conjugate diluent includes 10% (v/v) bovine serum, 10% (v/v) goat serum and 0.15% Triton-X100 in 20 mM Tris buffer, pH 7.5 with 0.01% gentamicin sulfate, pink dye and antifungal agents as preservatives. The conjugate is sterile filtered and filled in plastic bottles.

TABLE 4

PEPTIDE	COATING CONCEN. ug/ml	COATING BUFFER	OTHER COMPONENTS IN COATING SOLUTION
1-75	2.0	0.1M NaPHOSPHATE pH 6.5	0.4M NaCl, .0022%TRITON X-100
35-75	2.0	0.1M NaPHOSPHATE pH 6.5	0.4M NaCl, .0022%TRITON X-100
99-126	2.0	0.1M NaPHOSPHATE pH 6.5	0.4M NaCl, .0022%TRITON X-100
195-262	2.0	0.1M NaPHOSPHATE pH 6.5	0.4M NaCl, .0022%TRITON X-100
230-262	2.0	0.1M NaPHOSPHATE pH 6.5	0.4M NaCl, .0022%TRITON X-100
1357-1407	2.0	0.1M BORIC ACID pH 9.0	0.4M NaCl, .0022%TRITON X-100
1418-1457	2.0	0.1M BORIC ACID pH 9.0	0.4M NaCl, .0022%TRITON X-100
1569-1593	3.0	0.1M TRIS/HCl pH 8.5	0.5M NaCl, .0022%TRITON X-100
1899-1930	2.0	0.1M TRIS/HCl pH 8.5	0.5M NaCl, .0022%TRITON X-100
1192-1240	2.0	0.1M BORIC ACID pH 9.0	0.4M NaCl, .0022%TRITON X-100

TABLE 4 (CONT'D)

1223-1240	5.0	0.1M BORIC ACID pH 9.0	0.4M NaCl, .0022% TRITON X-100
1684-1750	1.0	0.1M BORIC ACID pH 10.0	0.4M NaCl, .0022% TRITON X-100
1689-1805	1.0	0.1M BORIC ACID pH 10.0	0.4M NaCl, .0022% TRITON X-100
1694-1735	3.0	0.1M TRIS/HCL pH 8.5	0.5M NaCl, .0022% TRITON X-100
1866-1930	0.75	0.1M TRIS/HCL pH 8.5	0.5M NaCl, .0022% TRITON X-100
380-436	3.0	0.1 M TRIS/HCL pH 8.5	0.9% NaCl
447-483	3.0		
643-683	3.0	0.1 M TRIS/HCL pH 8.5	0.5 M NaCl 0.0022% TRITON X-100
2302-2352	3.0	0.1 M Borate	0.4 M NaCl 0.0022% TRITON X-100

-24-

After one hour of incubation with the conjugate at 40° C, the beads are washed, exposed to the OPD substrate for thirty minutes at room temperature and the reaction terminated by the addition of 1 N H₂SO₄. The absorbance is read at 492 nm.

Samples found to be repeatably reactive by a screening assay using the polypeptide C100-3 are tested in duplicate using pl684 or pl689 coated beads. Reactive specimens are considered confirmed samples. Samples not reacting with pl684 or pl689 are tested in duplicate with pl694 and pl866 beads. Samples reacting with one or both of these peptides are considered confirmed. Those specimens not reacting with any of these peptides are considered nonconfirmed.

In order to maintain acceptable specificity, the cutoff for the assay should be at least 5-15 standard deviations above the absorbance value of a normal population mean. Consistent with these criteria, a cutoff for the assay may be selected which clearly separated most of the presumed "true negatives" from "true positive" specimens. A general cutoff value may be calculated as about 2.1 to 8 times the negative control mean absorbance value.

Confirmatory Assay Performance

1. Intravenous Drug User Samples

Samples were collected from a population of intravenous drug users enrolled in an NIH-funded study. The population consisted of individuals who were acknowledged users of intravenous drugs selected over a two-year period from patients at the Edward Hines Jr. Veteran's Administration Hospital in Maywood, Illinois by Dr. Connie Pachucki and members of the Infectious Diseases staff.

As illustrated in Table 5, a total of 296 specimens, each obtained from a single donor, were screened using recombinant yeast C100-3 polypeptide. A total of 271 of 296 (91.6%) specimens initially tested positive; upon retesting, 269 of 271 (99.3%) were repeat positives.

Confirmatory testing indicated that 263 of 269 (97.8%) of the repeat positives were reactive with pl689, five specimens

-25-

were non-reactive with pl689, and one specimen was not tested with any of the confirmatory polypeptides. Four of the five specimens which were non-reactive with pl689 were reactive with pl866 only; one specimen which was non-reactive with pl689 was reactive with pl694 only.

All specimens which were repeatably reactive were confirmed reactive in assays using the HCV synthetic peptides.

TABLE 5

<u>INTRAVENOUS DRUG USERS SAMPLES</u>					
<u>Confirmatory Testing</u>					
C100-3	C100-3				
Initial	Repeat				
Positive	Positive	pl689	pl866	pl694	No. of Repeat Positives Confirmed
271/296	269/271	263/269	4/5	1/5	268/269

2. Chimpanzees Samples

Confirmatory assays were used to evaluate 92 samples from six chimpanzees. All were initially reactive with recombinant C100-3. (Duplicate, repeat testing of chimp sera was not done because of the rare nature of these specimens and their utility for serological studies with other HCV antigens). Eighty-three of 92 (90.2%) specimens were confirmed reactive using pl689. Confirmation of initial reactives improved to 96.7% (89 of 92) when repeat testing with pl694 and pl866 was done.

3. Chiron Corporation Non-A, Non-B Hepatitis Virus Proficiency Panel #2

A proficiency panel comprised of neat and diluted human plasma including specimens containing antibodies to HCV C100-3 was provided by scientists at the Chiron Corporation (12 specimens). This panel contains specimens ranging from low to high reactivity in other assays, non-reactive presumed "true

-26-

negative" specimens, and reactive specimens diluted to give low-level or negative results.

Results using the confirmatory assay on the Chiron Corporation Non-A, Non-B Hepatitis Virus Proficiency Panel #2 indicated 9 of 9 (100%) of the specimens reactive by the preliminary screening assay are confirmed by pl689. All negative specimens were non-reactive.

4. Confirmatory Testing on NANB Panel II

A panel of highly pedigreed human sera from Dr. H. Alter, NIH, Bethesda, MD, containing infectious HCV sera, negative sera and other disease controls were tested. A total of 44 specimens were present in the panel.

All specimens (16/16, 100%) reactive in the assay using C100-3 were confirmed by pl689, as shown in Table 6. Again, there were no nonspecific reactives as all pedigreed negative or "other disease" controls were non-reactive in the peptide assay.

Data presented demonstrate the efficacy of the confirmatory assay for detection of antibodies to HCV antigens. The current assay is both sensitive and specific for detection of antibodies to HCV antigens.

The data further support the utility of the confirmatory strategy using synthetic peptides. The synthetic peptides serve as an independent source of antigen for use in immunoassays. The ability to confirm an average of 99% of repeatably active specimens in high risk or pedigreed positive HCV panels, establishes the utility of this strategy.

Example 2. COMBINATION ASSAY

The combination assay uses more than one polypeptide antigen coated on the same bead. To prepare multiple polypeptide-containing beads, the polystyrene beads described in Example 1 are contacted simultaneously with the polypeptide in appropriate buffer solutions. After the beads have been contacted with the polypeptides, the bead is treated further as described above.

-27-

TABLE 6

COMPARISON OF P1689 RESULTS WITH HCV SCREENING
ASSAY RESULTS ON NANB PANEL II (H. ALTER, NIH)

SAMPLE	C100-3		ORTHO C100-3	P1689 EIA
	MANUAL S/CO	MACHINE S/CO	S/CO	S/CO
1	>5.88	>6.47	>6.38	>8.33
2	0.63	0.93	0.27	0.45
3	>5.88	>6.47	>6.38	>8.33
4	>5.88	>6.47	>6.38	>8.33
5	0.43	0.35	0.16	0.43
6	>5.88	>6.47	>6.38	>8.33
7	0.46	0.73	0.36	0.32
8	0.41	0.50	0.32	0.38
9	1.87	2.21	0.91	2.84
10	0.35	0.41	0.32	0.30
11	0.48	0.45	0.27	0.46
12	0.32	0.41	0.17	0.39
13	0.48	0.69	0.32	0.51
14	0.37	0.40	0.19	0.32
15	>5.88	>6.47	>6.38	>8.33
16	>5.88	>6.47	>6.38	>8.33
17	0.34	0.40	0.20	0.44
18	3.01	3.68	0.68	6.80
19	0.74	0.61	0.53	0.72
20	0.53	0.59	0.28	0.33
21	>5.88	>6.47	>6.38	>8.33
22	0.24	0.26	0.20	0.23
23	>5.88	>6.47	>6.38	>8.33
24	0.69	0.64	0.53	0.70
25	0.50	0.60	0.49	0.40
26	3.41	4.11	0.77	6.61
27	0.62	0.74	0.30	0.65
28	0.61	0.77	0.08	0.47
29	0.34	0.42	0.13	0.33
30	1.58	2.40	1.26	2.65
31	0.32	0.35	0.22	0.37
32	>5.88	>6.47	>6.38	>8.33
33	0.45	0.48	0.24	0.45
34	>5.88	>6.47	>6.38	>8.33
35	>5.88	>6.47	>6.38	>8.33
36	0.37	0.36	0.21	0.40
37	0.40	0.46	0.24	0.52
38	>5.88	>6.47	>6.38	>8.33
39	0.40	0.49	0.80	0.46
40	0.53	0.59	0.30	0.56
41	0.41	0.28	0.15	0.32
42	0.52	0.56	0.33	0.60
43	0.28	0.30	0.38	0.33
44	0.44	0.57	0.35	0.53

-28-

For a polystyrene bead containing both C100-3 and pl694 the sensitivity of the assay increases. As graphically illustrated in Figure 3a, adding about 0.3, 0.95, and 3 micrograms of pl694 to the coating solution, respectively, shows a significant increase in the signal when the detection procedures of Example 1 are utilized. Figure 3b graphically illustrates the data which show no corresponding increase in the signals (such as may attend non-specific binding) generated from negative human plasma.

Example 3. SYNTHETIC POLYPEPTIDE-BASED ASSAY

The use of synthetic polypeptides which contain epitopes of HCV antigens provide immunological assays which have increased sensitivity and may be more specific than HCV immunological assays using the SOD fusion polypeptide C100-3. The use of shorter amino acid sequences on polystyrene bead provides an increase in sensitivity.

The increased sensitivity of an assay employing synthetic polypeptide compared to recombinant C100-3 polypeptide was demonstrated in a serial dilution study. The serial dilution study employed fifteen samples which were identified as having antibodies to HCV antigens using a recombinant C100-3 screening assay. Each positive sample was assayed using recombinant C100-3 polypeptide in one assay and *pl689 polypeptide in a second assay, and the samples were then diluted twofold until the S/CO value was less than one. In twelve samples the pl689 polypeptide gave increased sensitivity (larger S/CO values) at all dilutions. In two samples, the pl689 polypeptide and the recombinant yeast C100-3 polypeptides were essentially equivalent. In one sample, the pl689 polypeptide gave a negative response to a positive sample at all dilutions.

Additional studies on samples from serial bleeds of three chimps which developed an acute resolved case of HCV infections and three chimps which developed chronic HCV infections showed different immunological responses believed to be due to both the type of infection and the polypeptide used in the assay. This study assayed serum from serial bleeds of six chimps inoculated

-29-

with HCV. The assay protocols were similar to those described in Example 1 above with the following differences.

The antibodies, IgG, IgM and IgA were detected using affinity purified goat antibodies to human IgG, IgM and IgA coupled to horseradish peroxidase (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Maryland) which were used at working concentrations of 0.2 ug/ml Anti-IgG, 0.5 ug/ml Anti-IgM and 0.2 ug/ml Anti-IgA. Serum dilutions for each assay were 1:41 for IgG, 1:101 for IgM, and 1:41 for IgA.

The polypeptides that were used in the study include C100-3, pl694, pl684, pl689, and pl866.

Briefly, beads containing the polypeptides were incubated with diluted serum for one hour at 40 C, the beads were washed and incubated with the appropriate goat antibody for one hour at 40 C. The beads were washed again and the assay was developed by incubating the beads with OPD for thirty minutes at room temperature. The color development was quenched with 1 N sulfuric acid and the results read at 492 nm.

All chimps developed antibodies that were detected by C100-3, pl684, and pl866 within 7 to 17 week post-inoculation (WPI). Within each chimp, IgG antibody reacting with C100-3, pl684, and pl866 appeared at approximately the same time. The response to pl694 and pl866 was variable within that time period with indications that antibody to these two peptides can be either undetectable or significantly delayed following HCV infection. These data suggest that, out of the five peptides tested, antibody to C100-3, pl684, or pl689 would be the earliest and most consistent serologic indicator of HCV infection.

IgM antibody was detected in only three of the six chimps studied. The response of each of the three animals to C100-3, pl684, pl689 and pl694 was detected in 7 to 10 WPI whereas IgM antibody to pl866 was undetectable in two chimps and delayed in the third. All IgM responses were short lived with levels falling below positive (S/N less than 3.0) within 2 to 22 weeks.

The explanation and significance of finding IgM antibodies in 3 chimps with acute resolved disease while not detecting IgM

-30-

antibodies in 3 chimps with chronic infection is unexpected. Preliminary experimental results indicated that false negative IgM results due to preferential IgM binding is an unlikely explanation. If the pattern observed in these 6 chimps with the five peptides holds true, antibody assays will provide important HCV prognostic information.

A positive IgA response (S/N greater than 3.0) was detected in only 2 of the 6 chimps and proved to be either biphasic or significantly later than the IgG or IgM response. Although these 2 chimps had chronic disease no conclusions regarding the significance of IgA antibodies can be made since sera from the three resolved chimps is available only through 30 to 40 WPI.

The results show the polypeptides when used to assay for antibodies to HCV antigens are useful to follow the progression of *HCV infection and that the polypeptides exhibit unexpected sensitivity to different antibodies generated during the clinical progression of HCV infection.

Example 4. IMMUNODOT ASSAY.

The immunodot assay system uses a panel of purified synthetic polypeptides placed in an array on a nitrocellulose solid support. The prepared solid support is contacted with a sample and captures specific antibodies to HCV antigens. The captured antibodies are detected by a conjugate-specific reaction. Preferably, the conjugate-specific reaction is quantified using a reflectance optics assembly.

In an immunodot assay, a nitrocellulose-base test cartridge is treated with multiple antigenic polypeptides. A test cartridge which may be used in an automated process for performing an immunodot assay described above is illustrated in Figure 4. Each polypeptide is contained within a specific reaction zone on the test cartridge. After all the antigenic

-31-

polypeptides have been placed on the nitrocellulose, excess binding sites on the nitrocellulose are blocked. The test cartridge is then contacted with a sample such that each antigenic polypeptide in each reaction zone will react if the sample contains the appropriate antibody. After reaction, the test cartridge is washed and any antigen-antibody reactions are identified using suitable well known reagents.

As described in the patent applications listed above, the entire process is amenable to automation.

In a preferred immunodot assay, the synthetic polypeptides pl223, pl684, pl689 and pl866 were diluted into an aqueous buffered solution (polypeptide diluent: 0.03% Triton X-100 and 0.1% sodium azide in 50 mM Hepes buffer, pH 7.6) and applied to a preassembled nitrocellulose test cartridge at about 40 ng in each reaction zone. After drying the cartridge overnight at room temperature, the nonspecific binding capacity of the nitro-cellulose phase was blocked. The blocking solution contained 1% porcine gelatin, 1% casein enzymatic hydrolysate, 5% Tween-20*, 0.1% sodium azide, 0.5 M sodium chloride and 20 mM Tris, pH 7.5.

Test cartridges were incubated with samples 00642 and 423 (see Table 1) and ALT 27. The sample ALT 27 was obtained from a volunteer donor having elevated alanine aminotransferase levels. After sample incubation, sequential incubations with a biotin-conjugated goat anti-human immunoglobulin-specific antibody, an alkaline phosphatase-conjugated rabbit anti-biotin specific antibody, and 5-bromo-4-chloro-3-indolyl phosphate produced a colored product at the site of the reaction.

A detectable reaction is defined by the formation of a visually discernable product at the antigen site on the array; when quantified by the instrument, a reflectance density (Dr) value of greater than or equal to approximately 0.0150 above background is obtained. None of the tested polypeptides elicited

* Trademark

-32-

a detectable reaction with a negative control serum that was previously demonstrated negative for antibodies to HCV antigens using a recombinant C100-3 polypeptide.

A reaction with each of the synthetic polypeptides pl684, pl689, pl694 and pl866 occurred when the prepared test cells were incubated with either sample 00642 (1:100 dilution in negative serum) or sample 423 (1:40 dilution in negative serum).

Polypeptide pl223, in addition to polypeptides pl684, pl689, pl694 and pl866 demonstrated a significant reaction with the elevated ALT 27 specimen. In all specimens, highest reactivity was obtained with pl689. Enhanced reactivity of polypeptide pl684 with sample 00642 was achieved through subtle modification of the antigen dilution (the modified polypeptide diluent was 0.5 M sodium chloride, 0.0022% Triton X-100 and 0.1 M Tris/HCl, pH 8.5).

The net reflectance (Dr) for a test cartridge containing the polypeptides pl223, pl684, pl689, pl694, and pl866 which indicate a positive or negative response is set out in Table 7.

Example 5. COMPETITION ASSAY.

The synthetic peptides containing antigenic HCV epitopes are useful for competition assays. To perform a neutralization assay, peptides representing epitopes within the C100-3 region such as pl694, pl684 or pl689 are solubilized and mixed with a specimen

diluent to a final concentration of 0.5-50 ug/ml. Ten microliters of specimen or diluted specimen is added to a reaction well followed by 400 ul of the specimen diluent containing peptide and if desired, the mixture may be pre-incubated for about fifteen minutes to two hours. A bead coated with C100-3 antigen of HCV is then added to the reaction well and incubated for one hour at 40 C. After washing, 200 ul of a peroxidase labeled goat anti-human IgG in conjugate diluent is added and incubated for one hour at 40 C. After washing, OPD substrate is added and incubated at room temperature for thirty minutes. The reaction is

TABLE 7

Sample	NET REFLECTANCE DENSITY (D)						p1223	p1684	p1689	p1694	p1866	p1223	p1684	p1689	p1694	p1866
	p1223	p1684	p1689	p1694	p1866											
Neg Ctrl	-.0051	-.0042	.0047	.0012	.0134		-	-	-	-	-	-	-	-	-	-
	-.004	.0004	.0068	.0000	.0119		-	-	-	-	-	-	-	-	-	-
Sample 00642 (1:100)	.0011	.0281	.4401	.0748	.1444		-	+	+	+	+	-	+	+	+	+
	.0002	.0368	.4552	.0853	.1564		-	+	+	+	+	-	+	+	+	+
Sample 423 (1:40)	.0073	.1501	1.5389	.1922	.0325		-	+	++	+	+	-	+	+	+	+
	.3040	10.40	29.64	13.00	7.450		+	++	+++	++	++	+	++	++	++	++
Sample ALT27	.1469	9.619	30.20	13.26	7.887		+	++	+++	++	++	+	++	++	++	++
							+	++	+++	++	++	+	++	++	++	++

+/- Values Based on S/N

- Dr/Bkg < 2.5
+ 2.5 < Dr/Bkg < 100
++ 100 < Dr/Bkg < 1000
+++ Dr/Bkg > 1000

-34-

terminated by the addition of 1 N sulfuric acid and the absorbance read at 492 nm.

Samples containing antibodies to the C100-3 antigen generate a reduced signal caused by the competitive binding of the peptides to these antibodies in solution. The percentage of competitive binding may be calculated by comparing the absorbance value of the sample in the presence of a synthetic peptide to the absorbance value of the sample assayed in the absence of a synthetic peptide at the same dilution.

-35-

EXAMPLE 6 . EIA ASSAY

Beads were coated with either peptides 380-436 and 447-483, 643-683 and 2302-2353 according to the method described in Example 1, except that peptides 380-436 and 447-483 were coated simultaneously on the same solid phase, both sequences being from the putative envelope region of HCV. Either peptide alone had activity in this type of assay. EIA was performed using each bead configuration described herein. The EIA method performed was as is described in Example 1, with the cutoff set at four times the negative control value. Table 9 presents data obtained from these assays in which serum specimens from patients diagnosed with chronic NANBH were assayed.

TABLE 9
ANTIGEN
POS./NO. TESTED

p380	p643b	p2302
70/165 (42%)	62/165 (38%)	102/165 (62%)

EXAMPLE 7. EIA UTILIZING p380.LG AND p380.

Beads were coated either with p380.LG or p380 according to Example 1. An EIA following the procedure of Example 1 was used to assay samples. As can be seen by the data presented in Table 10, the p380.LG peptide detected antigen in specimens that were negative to p380. The p380.LG sequence is highly variable in this region. Therefore, there is reasonable probability that differentiation between HCV "serotypes" based on reactivity of human specimens to one or the other of these envelope region peptide sequences is possible. The data of Table 10 suggest that p380.LG can detect chronically infected HCV patients who are negative to p380.

TABLE 10

SAMPLE	p380		p380LG	
	OD	S/N	OD	S/N
#8	.155	2.12	.399	5.87
#28	.246	3.37	.950	13.97
#23	.114	1.56	.458	6.74

-36-

EXAMPLE 8. EIA UTILIZING p2302.

Beads were coated with either p2303 or "HCV2.0S/CO" according to the method of Example 1. The bead coated with "HCV 2.0/SCO" comprised antigen from the NS3 (CKS-33C), NS4 (C-100) and Core (CKS-CORE) regions of the HCV genome. A patient sample which exhibited seroconversion to p2302, but not to HCV.20 S/CO, is shown in Figure 5. Thus, this peptide improves the ability to detect HCV infected individuals.

EXAMPLE 9. PEPSCAN PROTOCOL

NS1 region of HCV genome from a.a. 600-720 was mapped with PEPSCAN analysis, which is serological analyses of series of overlapping peptides spanning the protein sequence to identify immunogenic domains. A total of 106 overlapping hexamer peptides were synthesized on polypropylene pins following the manufacturer's instructions (Cambridge Research Bioscience, Valley Stream, N.Y.). Fab dimers of IgG purified from sera of individuals seropositive for HCV were tested with these peptides. Based on the reactivity in EIA (performed as described by the manufacturer) four peptide sequences were selected as illustrated in Table 11.

Each of these peptides were synthesized by a stepwise solid phase synthesis, starting with the Carboxy terminus residue. A panel of sera positive for antibodies to C-100 protein of CHV was tested for their reactivity to NS1 peptide by microtiter EIA as described below.

EIA PROTOCOL

Wells of microtiter plates were coated with 100 ul of the peptide at 10 ug/ml in 0.02M bicarbonate buffer, pH 9.5 at ambient temperatures for 12-16 hrs. After washing with Phosphate buffered saline which also contained 0.01% Sodium Dodecyl sulphate (SDS) and 0.05% Tween-20 (available from BioRad Laboratories, Richmond, CA.), free sites were overcoated with 1% BSA in bicarbonate buffer pH 9.5. Plates were stored at 4 C following a final wash.

Sera from individuals seropositive for antibodies to HCV C-100 were serially diluted in 100 ul of a buffer containing 20MM sodium phosphate, pH 7.4, 0.15M NaCl, 20% normal goat serum, 10% fetal calf serum, 5 MM EDTA, 10MM EGTA, 50MM Tris, 0.2% Tween-20 with sodium azide as preservative, pH 6.8. The diluted sera were reacted with peptides in microtiter wells for 3 hours at

-37-

TABLE 11

AMINO ACID SEQUENCE OF PEPTIDES SELECTED
FROM THE NS1 REGION OF HCV GENOME
(A.A. 600-720) BASED ON PEPSCAN ANALYSIS

A.A. NO. OF HCV GENOME	PEPTIDE SEQUENCE
607-627	CLVDYPYRLWHYPCTINYTIF
643-663	ACNWTRGERCDLEDRDRSELSY
666-683	LLTTTQWQVLPCSFTTLPY
691-714	HLHQNIVDVQYLYGVGSSIASWAI

-38-

TABLE 12

REACTIVITY OF A PANEL OF HCV SEROPOSITIVE
SAMPLES WITH NS1 PEPTIDES
BY MICROTITER EIA

SAMPLE ID*	PEPTIDES SELECTED FROM NS1 REGION (A.A. NUMBERS)			
	607-627	643-663	666-683	691-714
15	-	-	+	-
22	-	-	-	-
23	-	-	-	-
24	+/-	-	-	-
25	+/-	-	-	-
32	+/-	+	+	-
36	-	-	-	-
46	-	-	-	-
50	-	-	-	-
65	-	-	-	-
70	-	+	+	-
71	-	-	+	-
75	-	-	-	-
89	-	++	-	+
95	-	++	++	+
100	-	++	-	-
102	-	-	-	-
108	-	-	-	-
130	-	-	+	-
137	-	-	-	-
LG	+	++	+++	+
301060	-	++	+++	+
PB3178	++	+	+++	+
PB3180	++	+	+++	++
300423	++	-	+++	+
% POSITIVE	16	36	44	27

* ALL SAMPLES SHOWED THE PRESENCE OF ANTIBODIES TO HEPATITIS C
VIRUS BY EIA AS WELL AS WESTERN BLOT ANALYSIS.

-39-

37°C or overnight at ambient temperatures. The plates were washed and 100 ul of appropriately diluted goat anti-mouse (HH) Horseradish Peroxidase (HRPO)-conjugated antibody (Jackson immunochemicals, West Grove, PA) was added. The plates were incubated at 37°C for 2 hours. After a final wash, 100 ul of 0-phenylenediamine 2HCl (OPD) color reagent was added. The reaction was carried out at room temperature in the dark for 20-25 minutes, and stopped by the addition of 100 ul of IN H SO₄. The absorbance of the reaction mixture recorded at 492 NM. A negative control which was previously confirmed to be negative for HCV infection was included with each plate in triplicate. The sample was considered reactive if the absorbance of the sample at a 1:2000 dilution was three times the absorbance of the negative control at the same dilution. Table 12 illustrates the reactivity of these samples with each of the peptides.

The legend for Table 12 is as follows:

+ = Sample showing A₄₉₂ 3 X neg. control

++ = Sample titering to 1:5000 dilution

+++ = Strong reactivity with sample titering to 1:10,000 dilution.

It is envisioned that these peptides may be used for the development of unique polyclonal and monoclonal antibodies. Other variations of applications and modifications of the specific embodiments of the invention as set forth herein will be apparent to those skilled in the art. Accordingly, the invention is intended to be limited only in accordance with the appended claims.

-40-

What is claimed is:

1. An assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample comprising:

5 contacting the sample with a polypeptide containing at least one epitope of an HCV antigen selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, p1899, p380, p380.LG, p447, 10 p607, p643a, p643b, p666, p691 and p2302 under conditions suitable for complexing the antibody with the polypeptide; and detecting the antibody-polypeptide complex.

15 2. The assay of claim 1 wherein the antigen is p1866, p380, p380.LG, p643b, p666, or p2302.

3. In a combination assay for detecting the presence of an antibody immunologically reactive with 20 an HCV antigen in a fluid sample wherein the sample is contacted with a polypeptide containing at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide and wherein the antibody-polypeptide complex is detected, 25 the improvement comprising:

 contacting the sample with a solid support having commonly bound recombinant polypeptide C100-3 and a polypeptide selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, 30 p1899, p380, p380.LG, p447, p607, p643a, p643b, p666, p691 and p2302.

4. In a confirmatory assay for identifying the presence of an antibody in a fluid sample 35 immunologically reactive with an HCV antigen wherein

-41-

the sample is used to prepare first and second aliquots and the first aliquot is contacted with a first recombinant polypeptide C100-3 which contains at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide and wherein the first antibody-antigen complex is detected, the improvement comprising:

contacting the second aliquot with a second polypeptide selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, p1899, p380, p380.LG, p447, p607, p643a, p643b, p666, p691 and p2302 under conditions suitable to form a second antibody-antigen complex; and detecting the second antibody-antigen complex.

5. The assay of claim 4 wherein the second antigen is p1684, p380, p380.LG, p643b, p666 or p2302.

6. In an immunodot assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample wherein the sample is concurrently contacted with at least two polypeptides each containing distinct epitopes of an HCV antigen under conditions suitable for complexing the antibody with the polypeptides and wherein the antibody-polypeptide is detected by reacting the complex with color-producing reagents, the improvement comprising:

employing polypeptides selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, p1899, p380, p380.LG, p447, p607, p643a, p643b, p666, p691, p2302 and C100-3 bound to a solid support.

-42-

7. In a competition assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample wherein the sample is used to prepare first and second immunologically equivalent aliquots wherein the first aliquot is contacted with solid support containing a bound polypeptide which contains at least one epitope of an HCV antigen under conditions suitable for complexing with the antibody to form a detectable antibody-polypeptide complex and wherein the second aliquot is first contacted with unbound polypeptide and then contacted with the solid support containing the bound polypeptide, the improvement comprising:

selecting the polypeptide from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, p1899, p380, p380.LG, p447, p607, p643a, p643b, p666, p691, and p2302.

8. An immunoassay kit comprising:

a polypeptide containing at least one epitope of an HCV antigen selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1866, p1899, p380, p380.LG, p447, p607, p643a, p643b, p666, p691, and p2302.

one or more sample preparation reagents; and

one or more detection and signal producing reagents.

9. The polypeptide p1866.


-43-


10. The polypeptides p1, p35, and p99.


11. The polypeptide p380.LG.


5

12. The polypeptide p2302.










Test - 3b



Test - 3a

