



(86) Date de dépôt PCT/PCT Filing Date: 2001/07/04

(87) Date publication PCT/PCT Publication Date: 2002/01/17

(85) Entrée phase nationale/National Entry: 2002/12/16

(86) N° demande PCT/PCT Application No.: SE 2001/001544

(87) N° publication PCT/PCT Publication No.: 2002/004446

(30) Priorités/Priorities: 2000/07/07 (0002603-9) SE;  
2000/07/27 (0002788-8) SE

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C07D 471/08, A61K 31/435, A61P 9/06,  
C07D 471/20


(71) Demandeur/Applicant:  
ASTRAZENECA AB, SE

(72) Inventeurs/Inventors:  
ANDERSSON, KJELL, SE;  
BJORE, ANNIKA, SE;  
BJORSNE, MAGNUS, SE;  
PONTEN, FRITIOF, SE;  
STRANDLUND, GERT, SE;  
SVENSSON, PEDER, SE;  
TOTTIE, LOUISE, SE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : NOUVEAUX COMPOSES DE BISPIDINE ET LEUR UTILISATION DANS LE TRAITEMENT DES ARYTHMIES  
CARDIAQUES

(54) Title: NEW BISPIDINE COMPOUNDS AND THEIR USE IN THE TREATMENT OF CARDIAC ARRHYTHMIAS



(57) Abrégé/Abstract:

There is provided compounds of formula (I), wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3a</sup>, R<sup>3b</sup> and R<sup>41</sup> to R<sup>46</sup> have meanings given in the description, which are useful in the prophylaxis and in the treatment of arrhythmias, in particular atrial and ventricular arrhythmias.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number  
**WO 02/04446 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 471/08**,  
471/20, A61K 31/435, A61P 9/06

(21) International Application Number: PCT/SE01/01544

(22) International Filing Date: 4 July 2001 (04.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0002603-9 7 July 2000 (07.07.2000) SE  
0002788-8 27 July 2000 (27.07.2000) SE

(71) Applicant (for all designated States except US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ANDERSSON, Kjell** [SE/SE]; AstraZeneca AB, S-151 85 Södertälje (SE). **BJÖRE, Annika** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **BJÖRSNE, Magnus** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **PONTÉN, Fritiof** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **STRANDLUND, Gert** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **SVENSSON, Peder** [SE/SE]; Nordenskiöldsgatan 12, S-413 09 Göteborg (SE). **TOTTIE, Louise** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE).

(74) Agent: **GLOBAL INTELLECTUAL PROPERTY**; AstraZeneca AB, S-151 85 Södertälje (SE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG)


— of inventorship (Rule 4.17(iv)) for US only

#### Published:

— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEW BISPIDINE COMPOUNDS AND THEIR USE IN THE TREATMENT OF CARDIAC ARRHYTHMIAS



(57) Abstract: There is provided compounds of formula (I), wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3a</sup>, R<sup>3b</sup> and R<sup>41</sup> to R<sup>46</sup> have meanings given in the description, which are useful in the prophylaxis and in the treatment of arrhythmias, in particular atrial and ventricular arrhythmias.

## NEW BISPIDINE COMPOUNDS AND THEIR USE IN THE TREATMENT OF CARDIAC ARRHYTHMIAS

### Field of the Invention

5

This invention relates to novel pharmaceutically useful compounds, in particular compounds which are useful in the treatment of cardiac arrhythmias.

### 10 Background and Prior Art

Cardiac arrhythmias may be defined as abnormalities in the rate, regularity, or site of origin of the cardiac impulse or as disturbances in conduction which causes an abnormal sequence of activation. Arrhythmias may be  
15 classified clinically by means of the presumed site of origin (i.e. as supraventricular, including atrial and atrioventricular, arrhythmias and ventricular arrhythmias) and/or by means of rate (i.e. bradyarrhythmias (slow) and tachyarrhythmias (fast)).

20 In the treatment of cardiac arrhythmias, the negative outcome in clinical trials (see, for example, the outcome of the Cardiac Arrhythmia Suppression Trial (CAST) reported in New England Journal of Medicine, 321, 406 (1989)) with "traditional" antiarrhythmic drugs, which act primarily by slowing the conduction velocity (class I antiarrhythmic drugs), has  
25 prompted drug development towards compounds which selectively delay cardiac repolarization, thus prolonging the QT interval. Class III antiarrhythmic drugs may be defined as drugs which prolong the transmembrane action potential duration (which can be caused by a block of outward  $K^+$  currents or from an increase of inward ion currents) and  
30 refractoriness, without affecting cardiac conduction.

One of the key disadvantages of hitherto known drugs which act by delaying repolarization (class III or otherwise) is that they all are known to exhibit a unique form of proarrhythmia known as *torsades de pointes* (turning of points), which may, on occasion be fatal. From the point of view of safety, the minimisation of this phenomenon (which has also been shown to be exhibited as a result of administration of non-cardiac drugs such as phenothiazines, tricyclic antidepressants, antihistamines and antibiotics) is a key problem to be solved in the provision of effective antiarrhythmic drugs.

10


Antiarrhythmic drugs based on bispidines (3,7-diazabicyclo[3.3.1]nonanes), are known from *inter alia* international patent applications WO 91/07405 and WO 99/31100, European patent applications 000 074, 301 245, 306 871, 308 843, 461 574 and 665 228, German patent applications DE 24 28 792, DE 26 58 558 and DE 27 44 248 and US patents 3,962,449, 4,556,662, 4,550,112, 4,459,301, 5,468,858 and 5,786,481, as well as journal articles including *inter alia*: *J. Med. Chem.* 39, 2559 (1996); *Pharmacol. Res.* 24, 149 (1991); *Circulation*, 90, 2032 (1994); *Anal. Sci.* 9, 429, (1993); and *J. Med Chem.* 20, 1668 (1977). Known bispidine-based antiarrhythmic compounds include bisamil (syn-9-(4-chlorobenzoyloxy)-3-methyl-7-ethyl-3,7-diazabicyclo[3.3.1]nonane), tedisamil (3,7-di-(cyclopropylmethyl)-9,9-tetramethylene-3,7-diazabicyclo[3.3.1]nonane), SAZ-VII-22 (3-(4-chlorobenzoyl)-7-*iso*-propyl-3,7-diazabicyclo[3.3.1]nonane), SAZ-VII-23 (3-benzoyl-7-*iso*-propyl-3,7-diazabicyclo[3.3.1]nonane), GLG-V-13 (3-[4-(1*H*-imidazol-1-yl)benzoyl]-7-*iso*-propyl-3,7-diazabicyclo[3.3.1]nonane), KMC-IV-84 (7-[4'-(1*H*-imidazol-1-yl)-benzenesulfonyl]-3-*iso*-propyl-3,7-diazabicyclo[3.3.1]nonane dihydroperchlorate and ambasilide (3-(4-aminobenzoyl)-7-benzyl-3,7-diazabicyclo[3.3.1]nonane).

Further bispidine compounds are known from *inter alia*: *Eur. J. Med. Chem.* 25, 1 (1990); *Bull. Polish Acad. Sci. Chem.* 34(5-6), 205 (1986); *J. Org. Chem.* 60, 8148 (1995); *Eur. J. Med. Chem. - Chimica Therapeutica* 12(4), 301 (1977); *Phosphorous, Sulfur and Silicon* 123, 385 (1997); *J. Org. Chem.* 42(6), 937 (1977); and *J. Molecular Structure* 127, 185 (1985).

We have surprisingly found that a novel group of 3,7-diazabicyclo[3.3.1]-nonane-based compounds exhibit electrophysiological activity, preferably class III electrophysiological activity, and are therefore expected to be useful in the treatment of cardiac arrhythmias.


### Disclosure of the Invention

According to the invention there is provided compounds of formula I,



wherein

R<sup>1</sup> represents a structural fragment of formula Ia,



20

R<sup>4</sup> represents H, halo, C<sub>1-4</sub> alkyl, -D-OR<sup>7</sup>, -D-N(R<sup>8</sup>)R<sup>9</sup>, or R<sup>4</sup>, together with R<sup>5</sup>, represents =O;

$R^5$  represents H,  $C_{1-4}$  alkyl, or  $R^5$ , together with  $R^4$ , represents  $=O$ ;

D represents a direct bond or  $C_{1-4}$  alkylene;

$R^7$  represents H,  $C_{1-6}$  alkyl, -E-aryl, -E-Het<sup>1</sup>,  $-C(O)R^{10a}$ ,  $-C(O)OR^{10b}$  or  $-C(O)N(R^{11a})R^{11b}$ ;

5  $R^8$  represents H,  $C_{1-6}$  alkyl, -E-aryl, -E-Het<sup>1</sup>,  $-C(O)R^{10a}$ ,  $-C(O)OR^{10b}$ ,  $-S(O)_2R^{10c}$ ,  $-[C(O)]_nN(R^{11a})R^{11b}$  or  $-C(NH)NH_2$ ;

$R^9$  represents H,  $C_{1-6}$  alkyl, -E-aryl, or  $-C(O)R^{10d}$ ;

E represents, at each occurrence when used herein, a direct bond or  $C_{1-4}$  alkylene;

10  $R^{10a}$  to  $R^{10d}$  independently represent, at each occurrence when used herein,  $C_{1-6}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo, aryl and Het<sup>2</sup>), aryl, Het<sup>3</sup>, or  $R^{10a}$  and  $R^{10d}$  independently represent H;

$R^{11a}$  and  $R^{11b}$  independently represent, at each occurrence when used herein,

15 H,  $C_{1-6}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo, aryl and Het<sup>4</sup>), aryl, Het<sup>5</sup>, or  $R^{11a}$  and  $R^{11b}$  together represent  $C_{3-7}$  alkylene, which alkylene group is optionally interrupted by an oxygen atom;

n represents 1 or 2;

20

A represents -G-, -J-N( $R^{12}$ )- or -J-O- (in which latter two groups, J is attached to the bispidine nitrogen atom);

B represents -L-, -L-N( $R^{13}$ )-, -N( $R^{13}$ )-L-, -L-S(O)<sub>p</sub>- or -L-O- (in which latter two groups, L is attached to the carbon atom bearing  $R^4$  and  $R^5$ );

25 G represents a direct bond or  $C_{1-6}$  alkylene;

J represents  $C_{2-6}$  alkylene;

L represents a direct bond or  $C_{1-4}$  alkylene;

p represents 0, 1 or 2;

$R^{12}$  and  $R^{13}$  independently represent H or  $C_{1-4}$  alkyl;

30

$R^6$  represents aryl, Het<sup>6</sup> (both of which groups are optionally substituted and/or terminated (as appropriate) by one or more substituents selected from -OH, cyano, halo, nitro, C<sub>1-6</sub> alkyl (optionally terminated by -N(H)C(O)OR<sup>14a</sup>), C<sub>1-6</sub> alkoxy, aryl, Het<sup>7</sup>, -N(R<sup>15a</sup>)R<sup>15b</sup>, -C(O)R<sup>15c</sup>,  
 5 -C(O)OR<sup>15d</sup>, -C(O)N(R<sup>15e</sup>)R<sup>15f</sup>, -N(R<sup>15g</sup>)C(O)R<sup>15h</sup>, -N(R<sup>15i</sup>)C(O)N(R<sup>15j</sup>)R<sup>15k</sup>,  
 -N(R<sup>15m</sup>)S(O)<sub>2</sub>R<sup>14b</sup>, -S(O)<sub>q</sub>R<sup>14c</sup>, -OS(O)<sub>2</sub>R<sup>14d</sup> and -S(O)<sub>2</sub>N(R<sup>15n</sup>)R<sup>15p</sup>) or, when R<sup>4</sup> and R<sup>5</sup> together represent =O, R<sup>6</sup> may represent C<sub>1-6</sub> alkyl;  
 q represents 0, 1 or 2;

10 R<sup>2</sup> represents -CN, Het<sup>8</sup>, -C(O)R<sup>16</sup>, -C(S)OR<sup>17</sup>, -C(S)N(R<sup>18</sup>)R<sup>19</sup>,  
 -[C(O)]<sub>2</sub>N(R<sup>20a</sup>)R<sup>20b</sup>, -[C(O)]<sub>2</sub>OR<sup>21</sup>, -S(O)<sub>2</sub>R<sup>22</sup>, -S(O)<sub>2</sub>N(R<sup>23</sup>)R<sup>24</sup>,  
 -C(=N-CN)N(R<sup>25</sup>)R<sup>26</sup>, -C(=N-CN)OR<sup>27</sup> or C<sub>1-12</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from -C(O)R<sup>28</sup>, -C(O)N(R<sup>29a</sup>)R<sup>29b</sup>, -N(R<sup>30</sup>)R<sup>31</sup>, -OR<sup>32</sup>, -S(O)<sub>r</sub>R<sup>33</sup>,  
 15 halo, -CN, nitro, aryl and Het<sup>9</sup>);

R<sup>16</sup> represents H, aryl, Het<sup>10</sup> or C<sub>1-6</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN, -N(R<sup>34</sup>)R<sup>35</sup>, aryl and Het<sup>11</sup>);

20 R<sup>34</sup> represents, H, C<sub>1-6</sub> alkyl, aryl, Het<sup>12</sup>, -C(O)R<sup>36a</sup> or -C(O)OR<sup>36b</sup>;

R<sup>18</sup> represents H, aryl, Het<sup>13</sup>, -C(O)R<sup>36a</sup>, -C(O)OR<sup>36b</sup> or C<sub>1-6</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN, -C(O)R<sup>36a</sup> and -C(O)OR<sup>36b</sup>);

25

R<sup>22</sup> represents Het<sup>14</sup>, aryl, or C<sub>1-6</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN, Het<sup>15</sup> and aryl);

$R^{23}$  represents H,  $C_{1-6}$  alkyl, aryl,  $Het^{16}$ ,  $-C(O)R^{36a}$ ,  $-C(O)OR^{36b}$  or  $-C(O)SR^{36b}$ ;

$R^{25}$  represents H or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN,  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from  $C_{1-4}$  alkyl and -OH),  $C_{1-6}$  alkoxy and aryl);

$R^{27}$  represents  $C_{1-6}$  alkyl or aryl;

$R^{28}$  represents H,  $C_{1-6}$  alkyl, aryl or  $Het^{17}$ ;

$R^{29a}$  and  $R^{29b}$  independently represent H,  $C_{1-6}$  alkyl, aryl or  $Het^{18}$ ;

$R^{30}$  represents H,  $C_{1-6}$  alkyl, aryl,  $Het^{19}$ ,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ ;

$R^{31}$  represents H,  $C_{1-6}$  alkyl, aryl or  $Het^{20}$ ;

$R^{32}$  represents H,  $C_{1-6}$  alkyl, aryl,  $Het^{21}$ ,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ ;

$R^{33}$  represents  $C_{1-6}$  alkyl, aryl or  $Het^{22}$ ;

r represents 0, 1 or 2;

$R^{36a}$  and  $R^{36b}$  independently represent, at each occurrence when used herein,  $C_{1-6}$  alkyl, or  $R^{36a}$  represents H;

$R^{37a}$  to  $R^{37d}$  independently represent, at each occurrence when used herein,  $C_{1-6}$  alkyl, aryl or  $Het^{23}$ , or  $R^{37a}$ ,  $R^{37c}$  and  $R^{37d}$  independently represent H;

Het<sup>1</sup> to Het<sup>23</sup> independently represent, at each occurrence when used herein, five- to twelve-membered heterocyclic groups containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur;

- 5 R<sup>3a</sup> and R<sup>3b</sup> independently represent H, C<sub>1-4</sub> alkyl, -OR<sup>38a</sup>, -SR<sup>38b</sup>, -N(R<sup>39</sup>)R<sup>38c</sup>, or R<sup>3a</sup> and R<sup>3b</sup> together represent C<sub>3-5</sub> alkylene, -O-Z-O-, -O-Z-S- or -S-Z-S-;

R<sup>39</sup> represents H, C<sub>1-6</sub> alkyl or a structural fragment of formula Ia as defined above;

- 10 Z represents C<sub>2-3</sub> alkylene optionally substituted by one or more C<sub>1-4</sub> alkyl groups;

R<sup>41</sup> to R<sup>46</sup> independently represent H or C<sub>1-3</sub> alkyl;

- 15 R<sup>14a</sup> to R<sup>14d</sup>, R<sup>17</sup> and R<sup>21</sup> independently represent C<sub>1-6</sub> alkyl;  
R<sup>15a</sup> to R<sup>15p</sup>, R<sup>19</sup>, R<sup>20a</sup>, R<sup>20b</sup>, R<sup>24</sup>, R<sup>26</sup>, R<sup>35</sup> and R<sup>38a</sup> to R<sup>38c</sup> independently represent H or C<sub>1-6</sub> alkyl;

- wherein each aryl and Het (Het<sup>1</sup> to Het<sup>23</sup>) group, unless otherwise specified,  
20 is optionally substituted;

or a pharmaceutically acceptable derivative thereof;

provided that:

- 25 (a) when R<sup>1</sup> represents a structural fragment of formula Ia in which:  
R<sup>4</sup> and R<sup>5</sup> together represent =O;  
A represents a direct bond;  
then B does not represent a direct bond, -N(R<sup>13</sup>)-L- (in which group  
-N(R<sup>13</sup>)- is attached to the carbon atom bearing R<sup>4</sup> and R<sup>5</sup>), -N(R<sup>13</sup>)-,  
30 -S(O)<sub>p</sub>- or -O-;

- (b) when  $R^5$  represents H or  $C_{1-4}$  alkyl; and  
 A represents  $-J-N(R^{12})-$  or  $-J-O-$ ;  
 then B does not represent  $-N(R^{13})-L-$ ,  $-N(R^{13})-$ ,  $-S(O)_p-$  or  $-O-$ ;
- (c) when  $R^4$  represents  $-D-OR^7$ ,  $-D-N(R^8)R^9$  in which D represents a  
 5 direct bond, then:
- (i) A does not represent  $-J-N(R^{12})-$  or  $-J-O-$ ; and  
 (ii) B does not represent  $-N(R^{13})-L-$ ,  $-N(R^{13})-$ ,  $-S(O)_p-$  or  $-O-$ ;
- (d) when  $R^{3a}$  and  $R^{3b}$  and both represent H;  
 and  $R^1$  represents unsubstituted benzyl;  
 10 then  $R^2$  does not represent unsubstituted benzyl or optionally  
 substituted benzoyl; and
- (e) the compound is not:
- (i)  $N^1$ -phenyl-3-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-  
 propanamide;
- 15 (ii) 3-benzyl-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-6,8-  
 dimethyl-3,7-diazabicyclo[3.3.1]nonane;
- (iii) 3-benzyl-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-6-methyl-  
 3,7-diazabicyclo[3.3.1]nonane;
- (iv)  $N$ -{2-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-1-[(4-cyano-  
 20 phenoxy)methyl]ethyl}methanesulfonamide;
- (v) 3-benzyl-7-[3-(2-propyl-1,3-dioxolan-2-yl)propyl]-3,7-diaza-  
 bicyclo[3.3.1]nonane; or
- (vi) 7-benzyl-3,7-diazabicyclo[3.3.1]nonane-3-ethanol;
- 25 which compounds are referred to hereinafter as "the compounds of the  
 invention".

Unless otherwise specified, alkyl groups and alkoxy groups as defined  
 herein may be straight-chain or, when there is a sufficient number (i.e. a  
 30 minimum of three) of carbon atoms, be branched-chain and/or cyclic.

Further, when there is a sufficient number of carbon atoms (i.e. a minimum of four), such alkyl and alkoxy groups may also be part cyclic/acyclic. Such alkyl and alkoxy groups may also be saturated or, when there is a sufficient number of carbon atoms (i.e. a minimum of two), be unsaturated  
 5 and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkyl and alkoxy groups may also be substituted by one or more halo, and especially fluoro, atoms.

Unless otherwise specified, alkylene groups as defined herein may be  
 10 straight-chain or, when there is a sufficient number of carbon atoms (i.e. a minimum of two), be branched-chain. Such alkylene chains may also be saturated or, when there is a sufficient number of carbon atoms (i.e. a minimum of two), be unsaturated and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkylene groups may also  
 15 be substituted by one or more halo atoms.

The term "aryl", when used herein, includes C<sub>6-10</sub> aryl groups such as phenyl, naphthyl and the like. Unless otherwise specified, aryl groups may be substituted by one or more substituents including -OH, cyano, halo,  
 20 nitro, C<sub>1-6</sub> alkyl (optionally terminated by -N(H)C(O)OR<sup>14a</sup>), C<sub>1-6</sub> alkoxy, Het<sup>1</sup>, aryl (which aryl group may not be substituted with any further aryl groups), -N(R<sup>15a</sup>)R<sup>15b</sup>, -C(O)R<sup>15c</sup>, -C(O)OR<sup>15d</sup>, -C(O)N(R<sup>15e</sup>)R<sup>15f</sup>,  
 -N(R<sup>15g</sup>)C(O)R<sup>15h</sup>, -N(R<sup>15i</sup>)C(O)N(R<sup>15j</sup>)R<sup>15k</sup>, -N(R<sup>15m</sup>)S(O)<sub>2</sub>R<sup>14b</sup>, -S(O)<sub>q</sub>R<sup>14c</sup>,  
 -OS(O)<sub>2</sub>R<sup>14d</sup> and -S(O)<sub>2</sub>N(R<sup>15n</sup>)R<sup>15p</sup> (wherein Het<sup>1</sup>, R<sup>14a</sup> to R<sup>14d</sup>, R<sup>15a</sup> to R<sup>15p</sup>  
 25 and q are as hereinbefore defined). When substituted, aryl groups are preferably substituted by between one and three substituents.

The term "halo", when used herein, includes fluoro, chloro, bromo and iodo.

Het (Het<sup>1</sup> to Het<sup>23</sup>) groups that may be mentioned include those containing 1 to 4 heteroatoms (selected from the group oxygen, nitrogen and/or sulfur) and in which the total number of atoms in the ring system are between five and twelve. Het (Het<sup>1</sup> to Het<sup>23</sup>) groups may be fully saturated, partly  
5 unsaturated, wholly aromatic, partly aromatic and/or bicyclic in character. Heterocyclic groups that may be mentioned include benzodioxanyl, benzodioxepanyl, benzodioxolyl, benzofuranyl, benzo-furazanyl, benzimidazolyl, benzomorpholinyl, benzothiophenyl, chromanyl, cinnolinyl, dioxanyl, furanyl, hydantoinyl, imidazolyl, imidazo[1,2-  
10 *a*]pyridinyl, indolyl, isoquinolinyl, isoxazolyl, maleimido, morpholinyl, 2-oxazolidonyl, oxazolyl, phthalazinyl, piperazinyl, piperidinyl, purinyl, pyranyl, pyrazinyl, pyrazolyl, pyridinyl, pyrimidyl, pyrrolidinonyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, quinazolinyl, quinolinyl, sulfolanyl, 3-sulfolenyl, tetrahydropyranyl, tetrahydrofuranyl, tetrazolyl, thiadiazolyl, thiazolyl, thienyl, thiochromanyl, triazolyl and the like. Values of Het<sup>1</sup> that  
15 may be mentioned include pyridyl. Values of Het<sup>6</sup> that may be mentioned include benzodioxanyl, benzomorpholinyl, furanyl, isoquinolinyl, isoxazolyl, 2-oxazolidonyl, piperazinyl, pyrazolyl, pyrrolidinonyl and 1,2,3-thiadiazolyl. Values of Het<sup>8</sup> that may be mentioned include pyrimidyl, quinazolinyl, tetrazolyl, thiazolyl and 1,2,4-triazolyl. Values of Het<sup>9</sup> that  
20 may be mentioned include benzomorpholinyl, 2-oxazolidonyl and piperazinyl. Values of Het<sup>10</sup> that may be mentioned include furanyl, isoxazolyl, pyrazolyl, pyrrolidinonyl and 1,2,3-thiadiazolyl. Values of Het<sup>14</sup> that may be mentioned include imidazolyl, sulfolanyl, thienyl and quinolinyl. Values of Het<sup>15</sup> that may be mentioned include morpholinyl.  
25 Values of Het<sup>17</sup> that may be mentioned include benzomorpholinyl. Values of Het<sup>21</sup> that may be mentioned include isoquinolinyl.

Unless otherwise specified, Het (Het<sup>1</sup> to Het<sup>23</sup>) groups may be substituted  
30 by one or more substituents including =O, -OH, cyano, halo, nitro, C<sub>1-6</sub>

alkyl (optionally terminated by  $-N(H)C(O)OR^{14a}$ ),  $C_{1-6}$  alkoxy,  $Het^1$ , aryl,  $-N(R^{15a})R^{15b}$ ,  $-C(O)R^{15c}$ ,  $-C(O)OR^{15d}$ ,  $-C(O)N(R^{15e})R^{15f}$ ,  $-N(R^{15g})C(O)R^{15h}$ ,  $-N(R^{15i})C(O)N(R^{15j})R^{15k}$ ,  $-N(R^{15m})S(O)_2R^{14b}$ ,  $-S(O)_qR^{14c}$ ,  $-OS(O)_2R^{14d}$  and  $-S(O)_2N(R^{15n})R^{15p}$  (wherein  $Het^1$ , aryl,  $R^{14a}$  to  $R^{14d}$ ,  $R^{15a}$  to  $R^{15p}$  and  $q$  are as  
 5 hereinbefore defined). When a  $Het$  ( $Het^1$  to  $Het^{23}$ ) group is substituted by one or more  $Het^1$  and/or aryl group(s), that (those) said  $Het^1$  and/or aryl substituent(s) may not itself (themselves) be substituted by any aryl and/or  $Het^1$  group(s). Substituents on  $Het$  ( $Het^1$  to  $Het^{23}$ ) groups may, where appropriate, be located on any atom in the ring system including a  
 10 heteroatom. The point of attachment of  $Het$  ( $Het^1$  to  $Het^{23}$ ) groups may be *via* any atom in the ring system including (where appropriate) a heteroatom.  $Het$  ( $Het^1$  to  $Het^{23}$ ) groups may also be in the *N*- or *S*-oxidised form.

Pharmaceutically acceptable derivatives include salts and solvates. Salts  
 15 which may be mentioned include acid addition salts. Pharmaceutically acceptable derivatives also include, at the 3,7-diazabicyclo[3.3.1]nonane nitrogen,  $C_{1-4}$  alkyl quaternary ammonium salts and *N*-oxides, provided that when a *N*-oxide is present:

- (a) no  $Het$  ( $Het^1$  to  $Het^{23}$ ) group contains an unoxidised S-atom;
- 20 (b)  $p$  does not represent 0 when B represents  $-L-S(O)_p-$ ;
- (c)  $q$  does not represent 0 when the group  $-S(O)_qR^{14c}$  is present as a substituent on aryl,  $Het$  ( $Het^1$  to  $Het^{23}$ ) or  $R^6$ ; and/or
- (d)  $r$  does not represent 0 when the group  $-S(O)_rR^{33}$  is present as a substituent on an alkyl group that  $R^2$  represents.

25

The compounds of the invention may exhibit tautomerism. All tautomeric forms and mixtures thereof are included within the scope of the invention.

The compounds of the invention may also contain one or more asymmetric  
 30 carbon atoms and may therefore exhibit optical and/or diastereoisomerism.

Diastereoisomers may be separated using conventional techniques, e.g. chromatography or fractional crystallisation. The various stereoisomers may be isolated by separation of a racemic or other mixture of the compounds using conventional, e.g. fractional crystallisation or HPLC, techniques. Alternatively the desired optical isomers may be made by reaction of the appropriate optically active starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation, for example with a homochiral acid followed by separation of the diastereomeric esters by conventional means (e.g. HPLC, chromatography over silica). All stereoisomers are included within the scope of the invention.

Abbreviations are listed at the end of this specification.

Compounds of the invention that may be mentioned include compounds of formula I, as defined above, with the additional provisos that:

(i) when  $R^2$  represents  $C_{1-6}$  alkyl (optionally substituted by one or two aryl groups), then:

(I) when  $R^4$  represents H,  $C_{1-4}$  alkyl,  $-OR^7$ , or  $R^4$ , together with  $R^5$  represents  $=O$ ;

$R^7$  represents H,  $C_{1-6}$  alkyl or  $-C(O)R^{10a}$ ; and

$R^6$  represents aryl;

then B does not represent  $-L-$ ; and/or

(II) when A represents a single bond; and

$R^4$  and  $R^5$  together represent  $=O$ ;

then  $R^{3a}$  and  $R^{3b}$  do not both represent  $C_{1-4}$  alkyl or they do not together represent  $C_{3-5}$  alkylene;

(ii) when  $R^2$  represents  $-C(O)R^{16}$  and the group  $-A-C(R^4)(R^5)-B-$  represents  $C_{1-6}$  alkylene, then:

(I)  $R^{16}$  does not represent aryl; and/or

(II)  $R^{3a}$  and  $R^{3b}$  do not both represent  $C_{1-4}$  alkyl or they do not together represent  $C_{3-5}$  alkylene; and

(iii) when  $R^2$  represents  $-S(O)_2R^{22}$ ;  
 $R^{3a}$  and  $R^{3b}$  independently represent H or  $C_{1-4}$  alkyl; and  
 5  $R^6$  represents aryl;  
 then A and B do not simultaneously represent direct bonds,  
 in which above provisos aryl groups, unless otherwise specified, are optionally substituted as described hereinbefore.

10 Compounds of the invention that may also be mentioned include compounds of formula I, as defined above, with the additional provisos that: when  $R^{3a}$  and  $R^{3b}$  independently represent H,  $C_{1-4}$  alkyl, OH or  $N(R^{39})R^{38c}$ ; and  
 $R^{39}$  represents H or  $C_{1-6}$  alkyl, then:

15 (a) when  $R^2$  represents CN or  $C_{1-6}$  alkyl optionally substituted by OH,  $N(R^{30})R^{31}$  or Het<sup>9</sup>;  
 $R^{30}$  and  $R^{31}$  independently represent H,  $C_{1-6}$  alkyl or  $C_{3-8}$  cycloalkyl;  
 Het<sup>9</sup> represents an unsubstituted, saturated 3- to 8-membered heterocycle containing one nitrogen atom (*via* which atom the  
 20 heterocyclic group is attached to the rest of the molecule);  
 $R^4$  and  $R^5$  both represent H; and  
 $R^6$  represents phenyl substituted in the *meta*- or *para*-position (relative to the group B) by  $CO_2H$  or  $NH_2$ , then:

(i) when A represents a direct bond,  $C_{1-6}$  *n*-alkylene or -J-O-; and  
 25 J represents  $C_{2-3}$  *n*-alkylene;  
 then B does not represent -L- $N(R^{13})$ - or -L-O- (in which latter two groups, L represents a direct bond or  $C_{1-4}$  *n*-alkylene and is attached to the carbon atom bearing  $R^4$  and  $R^5$ ); and  
 (ii) when A represents -J-O-; and  
 30 J represents  $C_{2-3}$  *n*-alkylene;

then B does not represent a direct bond; and

(b) when  $R^6$  represents  $Het^6$ ;

$Het^6$  represents an unsubstituted, saturated 3- to 8-membered heterocycle containing one nitrogen atom (*via* which atom the heterocyclic group is attached to the rest of the molecule); and

the group  $-A-C(R^4)(R^5)-B-$  represents  $C_{1-6}$  *n*-alkylene;

then  $R^2$  does not represent:

(i)  $C_{1-6}$  *n*-alkyl, which alkyl group is optionally interrupted by O and is terminated by  $N(R^{30})R^{31}$  or  $OR^{32}$ ; wherein

one of  $R^{30}$  and  $R^{31}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$  and the other represents H or  $C_{1-6}$  alkyl; and

$R^{32}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ;

(ii)  $-C(O)R^{16}$ , wherein  $R^{16}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ;

(iii)  $-S(O)_2R^{22}$ , wherein  $R^{22}$  represents phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ;

(iv)  $-S(O)_2N(R^{23})R^{24}$ ; wherein

$R^{23}$  represents phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ; and

$R^{24}$  represents H or  $C_{1-6}$  alkyl; and

(v)  $C_{3-4}$  *n*-alkyl, which alkyl group is terminated by phenyl, which phenyl group is substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ , and which alkyl group is interrupted at the  $\beta$ -position (relative to the point of attachment of the phenyl group) by O.

Further compounds of the invention that may be mentioned include compounds of formula I, as defined above, with the additional proviso that:

$R^6$  does not represent:

- (i) an unsubstituted, saturated 3- to 8-membered heterocycle containing one nitrogen atom (*via* which atom the heterocyclic group is attached to the rest of the molecule); or
- (ii) phenyl substituted in the *meta*- or *para*-position (relative to the group B) by  $\text{CO}_2\text{H}$  or  $\text{NH}_2$ .

Further compounds of the invention that may also be mentioned include those in which  $R^6$  represents aryl, which group is optionally substituted and/or terminated (as appropriate) by one or more substituents selected from -OH, cyano, halo, nitro,  $\text{C}_{1-6}$  alkyl (optionally terminated by -N(H)C(O)OR<sup>14a</sup>),  $\text{C}_{1-6}$  alkoxy, aryl, Het<sup>7</sup>, -C(O)R<sup>15c</sup>, -C(O)N(R<sup>15e</sup>)R<sup>15f</sup>, -N(R<sup>15g</sup>)C(O)R<sup>15h</sup>, -N(R<sup>15i</sup>)C(O)N(R<sup>15j</sup>)R<sup>15k</sup>, -N(R<sup>15m</sup>)S(O)<sub>2</sub>R<sup>14b</sup>, -S(O)<sub>q</sub>R<sup>14c</sup>, -OS(O)<sub>2</sub>R<sup>14d</sup> and -S(O)<sub>2</sub>N(R<sup>15n</sup>)R<sup>15p</sup>, or, when  $R^4$  and  $R^5$  together represent =O,  $R^6$  may represent  $\text{C}_{1-6}$  alkyl.

Preferred compounds of the invention also include those in which:

- $R^4$  represents H,  $\text{C}_{1-2}$  alkyl, -OR<sup>7</sup> or N(H)R<sup>8</sup>, or  $R^4$ , together with  $R^5$ , represents =O;
- $R^5$  represents H, or  $R^5$ , together with  $R^4$ , represents =O;
- $R^7$  represents H,  $\text{C}_{1-4}$  alkyl, optionally substituted phenyl, -C(O)R<sup>10a</sup>, or -C(O)N(R<sup>11a</sup>)R<sup>11b</sup>;
- $R^8$  represents H,  $\text{C}_{1-4}$  alkyl, -C(O)R<sup>10a</sup>, -C(O)OR<sup>10b</sup> or -C(O)N(R<sup>11a</sup>)R<sup>11b</sup>;
- $R^{10a}$  and  $R^{10b}$  independently represent, at each occurrence when used herein,  $\text{C}_{1-5}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo and phenyl), optionally substituted phenyl, or  $R^{10a}$  represents H;

$R^{11a}$  and  $R^{11b}$  independently represent, at each occurrence when used herein, H or  $C_{1-5}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo and phenyl);

A represents -G- or -J-N( $R^{12}$ )-;

5 B represents a direct bond,  $C_{1-4}$  alkylene, -L-N(H)-, -L-S(O)<sub>2</sub>- or -L-O- (in which latter three groups, L is attached to the carbon atom bearing  $R^4$  and  $R^5$ );

G represents a direct bond or  $C_{1-4}$  alkylene;

J represents  $C_{2-4}$  alkylene;

10 L represents  $C_{1-4}$  alkylene;

$R^6$  represents phenyl, Het<sup>6</sup> (both of which groups are optionally substituted by one or more substituents selected from cyano, halo, nitro,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, optionally substituted phenyl, -N(H) $R^{15b}$ , -C(O) $R^{15c}$ , -C(O)N(H) $R^{15f}$ , -N(H)C(O) $R^{15h}$ , -N(H)C(O)N(H) $R^{15k}$ , -N(H)S(O)<sub>2</sub> $R^{14b}$ ,  
15 -S(O)<sub>2</sub> $R^{14c}$  and -S(O)<sub>2</sub>N( $R^{15n}$ ) $R^{15p}$ ), or, when  $R^4$  and  $R^5$  together represent =O,  $R^6$  may represent  $C_{1-5}$  alkyl;

$R^2$  represents -CN, Het<sup>8</sup>, -C(O) $R^{16}$ , -C(S)OR<sup>17</sup>, -C(S)N(H) $R^{18}$ ,  
-[C(O)]<sub>2</sub>N(H) $R^{20b}$ , -[C(O)]<sub>2</sub>OR<sup>21</sup>, -S(O)<sub>2</sub> $R^{22}$ , -S(O)<sub>2</sub>N( $R^{23}$ ) $R^{24}$ ,  
-C(=N-CN)N( $R^{25}$ ) $R^{26}$ , -C(=N-CN)OR<sup>27</sup> or  $C_{1-6}$  alkyl (which alkyl group is  
20 optionally substituted and/or terminated by one or more substituents selected from -C(O) $R^{28}$ , -C(O)N(H) $R^{29b}$ , -N( $R^{30}$ ) $R^{31}$ , -OR<sup>32</sup>, -S(O)<sub>2</sub> $R^{33}$ , halo, -CN, optionally substituted phenyl and Het<sup>9</sup>);

$R^{16}$  represents optionally substituted phenyl, Het<sup>10</sup> or  $C_{1-6}$  alkyl (which alkyl group is optionally unsaturated and/or optionally substituted and/or  
25 terminated by one or more substituents selected from halo, -CN, -N(H) $R^{34}$  and optionally substituted phenyl);

$R^{34}$  represents, H,  $C_{1-4}$  alkyl, -C(O) $R^{36a}$  or -C(O)OR<sup>36b</sup>;

$R^{18}$  represents H, -C(O)OR<sup>36b</sup> or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo  
30 and -C(O)OR<sup>36b</sup>);

- $R^{22}$  represents Het<sup>14</sup>, optionally substituted phenyl or C<sub>1-4</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, Het<sup>15</sup> and optionally substituted phenyl);
- $R^{23}$  represents H, C<sub>1-4</sub> alkyl, -C(O)OR<sup>36b</sup> or -C(O)SR<sup>36b</sup>;
- 5  $R^{25}$  represents H or C<sub>1-6</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, C<sub>1-6</sub> alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from C<sub>1-4</sub> alkyl and -OH), C<sub>1-4</sub> alkoxy, naphthyl and optionally substituted phenyl);
- 10  $R^{27}$  represents optionally substituted phenyl;
- $R^{28}$  represents C<sub>1-5</sub> alkyl, optionally substituted phenyl or Het<sup>17</sup>;
- $R^{29b}$  represents H, C<sub>1-4</sub> alkyl or optionally substituted phenyl;
- $R^{30}$  represents H, optionally substituted phenyl, -C(O)R<sup>37a</sup> or -C(O)OR<sup>37b</sup>;
- $R^{31}$  represents H, C<sub>1-2</sub> alkyl or optionally substituted phenyl;
- 15  $R^{32}$  represents H, C<sub>1-4</sub> alkyl (which alkyl group is optionally interrupted by oxygen), optionally substituted phenyl or Het<sup>21</sup>;
- $R^{33}$  represents C<sub>1-6</sub> alkyl or optionally substituted phenyl;
- $R^{37a}$  and  $R^{37b}$  independently represent, at each occurrence when used herein, C<sub>1-5</sub> alkyl, optionally substituted phenyl, or  $R^{37a}$  represents H;
- 20  $R^{3a}$  and  $R^{3b}$  independently represent H, C<sub>1-2</sub> alkyl, -SR<sup>38b</sup>, -N(R<sup>39</sup>)R<sup>38c</sup>, or  $R^{3a}$  and  $R^{3b}$  together represent C<sub>3-4</sub> alkylene or -O-Z-O-;
- $R^{39}$  represents H, C<sub>1-2</sub> alkyl or a structural fragment of formula Ia;
- Z represents C<sub>2-3</sub> alkylene;
- $R^{41}$  to  $R^{46}$  independently represent H or C<sub>1-2</sub> alkyl;
- 25  $R^{14b}$ ,  $R^{14c}$ ,  $R^{17}$  and  $R^{21}$  independently represent C<sub>1-4</sub> alkyl;
- $R^{15b}$  to  $R^{15p}$ ,  $R^{20b}$ ,  $R^{24}$ ,  $R^{26}$ ,  $R^{38b}$  and  $R^{38c}$  independently represent H or C<sub>1-5</sub> alkyl;
- optional substituents on phenyl groups are one or more substituents selected from cyano, halo, nitro, C<sub>1-2</sub> alkyl, C<sub>1-2</sub> alkoxy, Het<sup>1</sup>, -NH<sub>2</sub>, -C(O)R<sup>15c</sup>,

$-\text{C}(\text{O})\text{N}(\text{H})\text{R}^{15\text{f}}$ ,  $-\text{N}(\text{H})\text{C}(\text{O})\text{R}^{15\text{h}}$ ,  $-\text{N}(\text{H})\text{C}(\text{O})\text{N}(\text{H})\text{R}^{15\text{k}}$ ,  $-\text{N}(\text{H})\text{S}(\text{O})_2\text{R}^{14\text{b}}$  and  $-\text{S}(\text{O})_2\text{N}(\text{R}^{15\text{n}})\text{R}^{15\text{p}}$ .

When  $\text{R}^{3\text{a}}$  and/or  $\text{R}^{3\text{b}}$  represent(s)  $-\text{N}(\text{R}^{39})\text{R}^{38\text{c}}$  in which  $\text{R}^{39}$  represents a structural fragment of formula Ia, preferred compounds of formula I include those in which, in that  $\text{R}^{39}$  group:

$\text{R}^4$  represents H,  $-\text{OR}^7$  or  $\text{N}(\text{H})\text{R}^8$ , or  $\text{R}^4$ , together with  $\text{R}^5$ , represents  $=\text{O}$ ;

$\text{R}^5$  represents H, or  $\text{R}^5$ , together with  $\text{R}^4$ , represents  $=\text{O}$ ;

$\text{R}^7$  represents H, phenyl (which group is optionally substituted by one to three methoxy groups),  $-\text{C}(\text{O})\text{CH}_3$ , or  $-\text{C}(\text{O})\text{N}(\text{H})-\text{C}_{1-4}$  alkyl;

$\text{R}^8$  represents H,  $-\text{C}(\text{O})\text{O}-\text{C}_{1-4}$  alkyl or  $-\text{C}(\text{O})\text{N}(\text{H})\text{CH}_3$ ;

A represents  $\text{C}_{1-3}$  alkylene or  $-\text{C}_{2-3}$  alkylene- $\text{N}(\text{H})-$ ;

B represents a direct bond,  $-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{H})-$ ,  $-\text{CH}_2-\text{S}(\text{O})_2-$ ,  $-\text{CH}_2-\text{O}-$  (in which latter three groups,  $\text{CH}_2$  is attached to the carbon atom bearing  $\text{R}^4$  and  $\text{R}^5$ ) or  $-\text{O}-$ ;

$\text{R}^6$  represents phenyl optionally substituted by up to three substituents (in the *para*- and/or *ortho*- positions) selected from cyano,  $-\text{N}(\text{H})\text{C}(\text{O})\text{N}(\text{H})\text{CH}_3$ ,  $-\text{N}(\text{H})\text{S}(\text{O})_2\text{CH}_3$  and  $-\text{S}(\text{O})_2\text{N}(\text{CH}_3)_2$ .

More preferred compounds of the invention include those in which:

$\text{R}^4$  represents H,  $-\text{OR}^7$  or  $\text{N}(\text{H})\text{R}^8$ , or  $\text{R}^4$ , together with  $\text{R}^5$ , represents  $=\text{O}$ ;

$\text{R}^5$  represents H, or  $\text{R}^5$ , together with  $\text{R}^4$ , represents  $=\text{O}$ ;

$\text{R}^7$  represents H,  $\text{C}_{1-2}$  alkyl, optionally substituted phenyl,  $-\text{C}(\text{O})\text{R}^{10\text{a}}$ , or  $-\text{C}(\text{O})\text{N}(\text{R}^{11\text{a}})\text{R}^{11\text{b}}$ ;

$\text{R}^8$  represents H,  $\text{C}_{1-2}$  alkyl,  $-\text{C}(\text{O})\text{OR}^{10\text{b}}$  or  $-\text{C}(\text{O})\text{N}(\text{R}^{11\text{a}})\text{R}^{11\text{b}}$ ;

$\text{R}^{10\text{a}}$  and  $\text{R}^{10\text{b}}$  independently represent, at each occurrence when used herein,  $\text{C}_{1-5}$  alkyl (optionally substituted or terminated by phenyl), optionally substituted phenyl, or  $\text{R}^{10\text{a}}$  represents H;

$\text{R}^{11\text{a}}$  and  $\text{R}^{11\text{b}}$  independently represent, at each occurrence when used herein,

H or  $\text{C}_{1-5}$  alkyl (optionally substituted or terminated by phenyl);

A represents -G- or -J-N(R<sup>12</sup>)-;

B represents a direct bond, C<sub>1-4</sub> alkylene, -L-N(H)-, -L-S(O)<sub>2</sub>- or -L-O- (in which latter three groups, L is attached to the carbon atom bearing R<sup>4</sup> and R<sup>5</sup>);

5 G represents a direct bond or C<sub>1-4</sub> alkylene;

J represents C<sub>2-4</sub> alkylene;

L represents C<sub>1-4</sub> alkylene;

R<sup>6</sup> represents phenyl or Het<sup>6</sup> (which two groups are optionally substituted by one or more substituents selected from cyano, halo, C<sub>1-2</sub> alkyl, C<sub>1-2</sub> alkoxy, -C(O)R<sup>15c</sup>, -N(H)C(O)R<sup>15h</sup>, -N(H)C(O)N(H)R<sup>15k</sup>, -N(H)S(O)<sub>2</sub>R<sup>14b</sup>,  
10 -S(O)<sub>2</sub>R<sup>14c</sup> and -S(O)<sub>2</sub>N(R<sup>15n</sup>)R<sup>15p</sup>).

When R<sup>2</sup> represents -S(O)<sub>2</sub>R<sup>22</sup>, more preferred compounds of the invention also include those in which:

R<sup>4</sup>, together with R<sup>5</sup>, represents =O;

15 A represents C<sub>1-4</sub> alkylene;

B represents a direct bond or C<sub>1-4</sub> alkylene;

R<sup>6</sup> represents C<sub>1-5</sub> alkyl.

Further preferred compounds of the invention include those in which:

20 A represents -G-, -J-N(R<sup>12</sup>)- or -J-O- (in which latter two groups, J is attached to the bispidine nitrogen atom);

G represents C<sub>1-6</sub> alkylene;

R<sup>4</sup> represents -D-OR<sup>7</sup>, -D-N(R<sup>8</sup>)R<sup>9</sup>, or R<sup>4</sup>, together with R<sup>5</sup>, represents =O;

R<sup>2</sup> represents -CN, Het<sup>8</sup>, -C(O)R<sup>16</sup>, -C(S)OR<sup>17</sup>, -C(S)N(R<sup>18</sup>)R<sup>19</sup>,  
25 -[C(O)]<sub>2</sub>N(R<sup>20a</sup>)R<sup>20b</sup>, -[C(O)]<sub>2</sub>OR<sup>21</sup>, -S(O)<sub>2</sub>R<sup>22</sup>, -S(O)<sub>2</sub>N(R<sup>23</sup>)R<sup>24</sup>,  
-C(=N-CN)N(R<sup>25</sup>)R<sup>26</sup>, -C(=N-CN)OR<sup>27</sup> or C<sub>1-12</sub> alkyl (which alkyl group is substituted and/or terminated by one or more substituents selected from -C(O)R<sup>28</sup>, -C(O)N(R<sup>29a</sup>)R<sup>29b</sup>, -N(R<sup>30</sup>)R<sup>31</sup>, -OR<sup>32</sup>, -S(O)<sub>r</sub>R<sup>33</sup>, halo, -CN, nitro and Het<sup>9</sup>);

$R^{16}$  represents H,  $Het^{10}$  or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN,  $-N(R^{34})R^{35}$ , aryl and  $Het^{11}$ );


$R^6$  represents aryl,  $Het^6$  (both of which groups are substituted and/or terminated (as appropriate) by one or more substituents selected from -OH, cyano, halo, nitro,  $C_{1-6}$  alkyl (optionally terminated by  $-N(H)C(O)OR^{14a}$ ),  $C_{1-6}$  alkoxy, aryl,  $Het^7$ ,  $-N(R^{15a})R^{15b}$ ,  $-C(O)R^{15c}$ ,  $-C(O)OR^{15d}$ ,  $-C(O)N(R^{15e})R^{15f}$ ,  $-N(R^{15g})C(O)R^{15h}$ ,  $-N(R^{15i})C(O)N(R^{15j})R^{15k}$ ,  $-N(R^{15m})S(O)_2R^{14b}$ ,  $-S(O)_qR^{14c}$ ,  $-OS(O)_2R^{14d}$  and  $-S(O)_2N(R^{15n})R^{15p}$ ) or, when  $R^4$  and  $R^5$  together represent  $=O$ ,  $R^6$  may represent  $C_{1-6}$  alkyl.

Preferred compounds of the invention include the compounds of the Examples disclosed hereinafter.

## 15 Preparation


According to the invention there is also provided a process for the preparation of compounds of formula I which comprises:

20 (a) reaction of a corresponding compound of formula II,




wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined, with a compound of formula III,

21



wherein  $L^1$  represents a leaving group (e.g. mesylate, tosylate or halo) and  $R^4$ ,  $R^5$ ,  $R^6$ , A and B are as hereinbefore defined, for example at between  $-10^\circ\text{C}$  and reflux temperature in the presence of a suitable base (e.g. triethylamine or  $\text{K}_2\text{CO}_3$ ) and an appropriate organic solvent (e.g. dichloromethane, acetonitrile or DMSO);


(b) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which A represents  $\text{C}_2$  alkylene and  $R^4$  and  $R^5$  together represent  $=\text{O}$ , reaction of a corresponding compound of formula II, as hereinbefore defined, with a compound of formula IV,



wherein  $R^6$  and B are as hereinbefore defined, for example at room temperature in the presence of a suitable organic solvent (e.g. ethanol);

(c) for compounds of formula I in which  $R^{3a}$  or  $R^{3b}$  represents  $-\text{N}(\text{R}^{39})\text{R}^{38c}$  and  $R^{39}$  represents a structural fragment of formula Ia, reaction of a corresponding compound of formula I in which  $R^{3a}$  or  $R^{3b}$  (as appropriate) represents  $-\text{N}(\text{H})\text{R}^{38c}$ , wherein  $R^{38c}$  is as hereinbefore defined, with a compound of formula III as hereinbefore defined, for example under conditions described hereinbefore (see process step (a));

(d) for compounds of formula I in which  $R^1$  represents a fragment of formula Ia in which A represents  $\text{CH}_2$  and  $R^4$  represents  $-\text{OH}$  or  $-\text{N}(\text{H})\text{R}^8$ , reaction of a corresponding compound of formula II, as hereinbefore defined, with a compound of formula V,



wherein X represents O or N(R<sup>8</sup>) and R<sup>5</sup>, R<sup>6</sup>, R<sup>8</sup> and B are as hereinbefore defined, for example at elevated temperature (e.g. 60°C to reflux) in the presence of a suitable solvent (e.g. a lower alkyl alcohol (e.g. IPA), acetonitrile, or a mixture of a lower alkyl alcohol and water);

(e) for compounds of formula I in which R<sup>3a</sup> or R<sup>3b</sup> represents -N(R<sup>39</sup>)R<sup>38c</sup> and R<sup>39</sup> represents a structural fragment of formula Ia in which A represents CH<sub>2</sub> and R<sup>4</sup> represents -OH or -N(H)R<sup>8</sup>, reaction of a corresponding compound of formula I in which R<sup>3a</sup> or R<sup>3b</sup> (as appropriate) represents -N(H)R<sup>38c</sup>, wherein R<sup>38c</sup> is as hereinbefore defined, with a compound of formula V as hereinbefore defined, for example under conditions described hereinbefore (see process step (d));

(f) for compounds of formula I in which A represents C<sub>1-6</sub> alkylene, B represents C<sub>1-4</sub> alkylene and R<sup>4</sup> and R<sup>5</sup> both represent H, reduction of a corresponding compound of formula I in which R<sup>4</sup> and R<sup>5</sup> together represent =O, in the presence of a suitable reducing agent and under appropriate reaction conditions, for example by activating the relevant C=O group using an appropriate agent (such as tosylhydrazine) in the presence of a suitable reducing agent (e.g. sodium borohydride or sodium cyanoborohydride) and an appropriate organic solvent (e.g. a lower (e.g. C<sub>1-6</sub>) alkyl alcohol);


(g) for compounds of formula I in which R<sup>4</sup> and R<sup>5</sup> both represent H and (1) A represents a single bond or -J-N(R<sup>12</sup>) and B represents C<sub>1-4</sub> alkylene, or (2) A represents C<sub>1-6</sub> alkylene and B represents N(R<sup>13</sup>) or -N(R<sup>13</sup>)-L-, reduction of a corresponding compound of formula I in which R<sup>4</sup> and R<sup>5</sup>

together represent =O, in the presence of a suitable reducing agent (e.g.  $\text{LiAlH}_4$ ) and an appropriate solvent (e.g. THF);

(h) for compounds of formula I in which A represents  $\text{C}_{1-6}$  alkylene, B represents a direct bond,  $\text{C}_{1-4}$  alkylene,  $-\text{L}-\text{N}(\text{R}^{13})-$ ,  $-\text{L}-\text{S}(\text{O})_p-$  or  $-\text{L}-\text{O}-$  (in which latter three groups L represents  $\text{C}_{1-4}$  alkylene),  $\text{R}^4$  represents OH and  $\text{R}^5$  represents H, reduction of a corresponding compound of formula I in which  $\text{R}^4$  and  $\text{R}^5$  together represent =O, in the presence of a suitable reducing agent (e.g.  $\text{NaBH}_4$ ) and an appropriate organic solvent (e.g. THF);

10

(i) for compounds of formula I in which  $\text{R}^{3a}$  and  $\text{R}^{3b}$  both represent H, reduction of a corresponding compound of formula VI,




wherein  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^{41}$  to  $\text{R}^{46}$  are as hereinbefore defined, and in which the bridgehead  $\text{C}=\text{O}$  group may be activated using an appropriate agent, such as tosylhydrazine, in the presence of a suitable reducing agent (e.g. sodium borohydride, sodium cyanoborohydride) and an appropriate organic solvent (e.g. a lower alkyl alcohol), or under standard Wolff-Kischner conditions known to those skilled in the art; when the  $\text{C}=\text{O}$  group is activated, the activation step may be carried out at between room and reflux temperature in the presence of an appropriate organic solvent (e.g. a lower alkyl alcohol such as methanol, ethanol or IPA), whereafter the reducing agent may be added to the reaction mixture and the reduction carried out at between  $60^\circ\text{C}$

and reflux, advantageously in the presence of a suitable organic acid (e.g. acetic acid);

(j) for compounds of formula I in which one of  $R^{3a}$  and  $R^{3b}$  represents H, and the other represents -OH, reduction of a corresponding compound of formula VI, as hereinbefore defined, in the presence of a mild reducing agent, e.g. sodium borohydride, and an appropriate organic solvent (e.g. a lower alcohol such as methanol or ethanol);

(k) for compounds of formula I in which  $R^{3a}$  and  $R^{3b}$  both represent  $-OR^{38a}$  or  $-SR^{38b}$ , or in which  $R^{3a}$  and  $R^{3b}$  together represent  $-O-Z-O-$ ,  $-O-Z-S-$  or  $-S-Z-S-$ , reaction of a corresponding compound of formula VI, as hereinbefore defined, with a compound of formula  $HOR^{38a}$ ,  $HSR^{38b}$ ,  $HO-Z-OH$ ,  $HO-Z-SH$  or  $HS-Z-SH$  (as appropriate), wherein  $R^{38a}$ ,  $R^{38b}$  and Z are as hereinbefore defined, under appropriate reaction conditions, for example by refluxing in the presence of a suitable protic or Lewis acid catalyst (e.g. *p*TSA, trimethylsilyl chloride or boron trifluoride) and an appropriate organic solvent (e.g. toluene or diethyl ether);

(l) for compounds of formula I in which one of  $R^{3a}$  and  $R^{3b}$  represents  $-NH_2$  and the other represents H, reduction of a compound of formula VII,



VII

wherein  $R^1$ ,  $R^2$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined, in the presence of a suitable reducing agent (e.g.  $\text{LiAlH}_4$ ), for example under conditions that are well known to those skilled in the art;

- 5 (m) for compounds of formula I in which one or both of  $R^{3a}$  and  $R^{3b}$  represent  $-\text{N}(\text{R}^{39})\text{R}^{38c}$  in which one or both of  $R^{39}$  and  $R^{38c}$  represents  $\text{C}_{1-6}$  alkyl, alkylation of a corresponding compound of formula I in which  $R^{3a}$  and/or  $R^{3b}$  represent  $-\text{N}(\text{R}^{39})\text{R}^{38c}$  (as appropriate) in which  $R^{39}$  and/or  $R^{38c}$  (as appropriate) represent H, using a compound of formula VIII,


10



VIII

wherein  $R^a$  represents  $\text{C}_{1-6}$  alkyl and  $L^1$  is as hereinbefore defined, for example under conditions that are well known to those skilled in the art;

- 15 (n) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which B represents  $-\text{L}-\text{O}-$ , reaction of a compound of formula IX,



wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^4$ ,  $R^5$ ,  $R^{41}$  to  $R^{46}$ , A and L are as hereinbefore defined, with a compound of formula X,

20




X

in which  $R^6$  is as hereinbefore defined, for example under Mitsunobu-type conditions e.g. at between ambient (e.g.  $25^\circ\text{C}$ ) and reflux temperature in the presence of a tertiary phosphine (e.g. tributylphosphine or triphenyl-

phosphine), an azodicarboxylate derivative (e.g. diethylazodicarboxylate or 1,1'-(azodicarbonyl)dipiperidine) and an appropriate organic solvent (e.g. dichloromethane or toluene);

- 5 (o) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which A represents  $C_{1-6}$  alkylene and B represents  $-N(R^{13})-L-$  (wherein the group  $-N(R^{13})-$  is attached to the carbon atom bearing  $R^4$  and  $R^5$ ), reaction of a compound of formula XI,




- 10 wherein  $A^a$  represents  $C_{1-6}$  alkylene and  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^4$ ,  $R^5$ ,  $R^{13}$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined with a compound of formula XII,



- wherein  $L^2$  represents a leaving group such as halo, alkane sulfonate, perfluoroalkane sulfonate or arenesulfonate, and  $R^6$  and L are as  
15 hereinbefore defined, for example at 40°C in the presence of a suitable organic solvent (e.g. acetonitrile);

- (p) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which  $R^4$  represents  $-D-NH_2$ , reduction of a corresponding  
20 compound of formula XIII,

27



wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^5$ ,  $R^6$ ,  $R^{41}$  to  $R^{46}$ , A, B and D are as hereinbefore defined, for example by hydrogenation at a suitable pressure in the presence of a suitable catalyst (e.g. palladium on carbon) and an appropriate solvent  
 5 (e.g. a water-ethanol mixture);

(q) for compounds of formula I in which  $R^4$  represents  $-D-N(R^9)C(O)NH(R^{11b})$ , reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-N(R^9)H$  with a compound of formula XIV,



wherein  $R^{11b}$  is as hereinbefore defined, for example at ambient temperature (e.g. 25°C) in the presence of a suitable solvent (e.g. benzene);

(r) for compounds of formula I in which  $R^4$  represents  $-D-N(H)[C(O)]_2NH_2$ ,  
 15 reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-NH_2$  with oxalic acid diamide, for example at between -10 and 25°C in the presence of a suitable coupling agent (e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide), an appropriate activating agent (e.g. 1-hydroxybenzotriazole), a suitable base (e.g. triethylamine) and a reaction-inert organic  
 20 solvent (e.g. DMF);

(s) for compounds of formula I in which  $R^4$  represents  $-D-N(R^8)R^9$ , wherein  $R^8$  and  $R^9$  are as hereinbefore defined, provided that  $R^8$  does not represent H, reaction of a corresponding compound of formula I, in which  $R^4$  represents  $-D-N(H)R^9$  with a compound of formula XV,



wherein  $R^{8a}$  represents  $R^8$  as hereinbefore defined except that it does not represent H, and  $L^3$  represents a leaving group such as halo (e.g. chloro or bromo), *p*-nitrophenolate,  $C_{1-4}$  alkoxide,  $C_{1-4}$  alkylthiolate,  $-OC(O)R^{10a}$ ,  $-OC(O)OR^{10b}$ , or  $-OS(O)_2R^{10c}$ , wherein  $R^{10a}$  to  $R^{10c}$  are as hereinbefore defined, for example under conditions that are well known to those skilled in the art;


(t) for compounds of formula I in which  $R^4$  represents  $-D-OR^7$  in which  $R^7$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>, reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-OH$  with a compound of formula XVI,



wherein  $R^{7a}$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>, wherein Het<sup>1</sup> is as hereinbefore defined, for example at between ambient (e.g. 25°C) and reflux temperature, under Mitsunobu-type conditions (i.e. in the presence of e.g. triphenylphosphine, an azodicarboxylate derivative (e.g. 1,1'-(azodicarbonyl)dipiperidine) and a suitable organic solvent (e.g. dichloromethane));

(u) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which  $R^4$  represents  $-D-OR^7$  (in which  $R^7$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>), reaction of a corresponding compound of formula XVII,

29



wherein  $L^2$ ,  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^5$ ,  $R^6$ ,  $R^{41}$  to  $R^{46}$ , A, B and D are as hereinbefore defined with a compound of formula XVI as hereinbefore defined, for example at between ambient (e.g. 25°C) and reflux temperature, under  
 5 Williamson-type conditions (i.e. in the presence of an appropriate base (e.g. KOH or NaH) and a suitable organic solvent (e.g. dimethylsulfoxide or DMF));


(v) for compounds of formula I in which  $R^4$  represents  $-D-OR^7$ , wherein  $R^7$   
 10 is as hereinbefore defined, provided that it does not represent H, reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-OH$  with a compound of formula XVIII,



wherein  $R^{7b}$  represents  $R^7$  as hereinbefore defined, except that it does not  
 15 represent H, and  $L^4$  represents a leaving group such as OH, halo, alkane sulfonate, arene sulfonate or  $-OC(O)R^{10a}$ , wherein  $R^{10a}$  is as hereinbefore defined, for example at between room and reflux temperature, optionally in the presence of a reaction-inert organic solvent (e.g. THF or  $CH_2Cl_2$ ), a suitable base (e.g. triethylamine or  $K_2CO_3$ ) and/or an appropriate coupling  
 20 agent (e.g. 1,3-dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, optionally combined with a suitable catalyst such as 4-dimethylaminopyridine) (for example, when  $R^{7b}$  represents  $-C(O)R^{10a}$  and  $L^4$  represents OH, this reaction may be performed at ambient temperature

(e.g. 25°C) in the presence of a coupling agent such as 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide, an appropriate catalyst such as 4-(dimethyl-amino)pyridine and a solvent such as THF);

- 5 (w) for compounds of formula I in which  $R^4$  represents halo, substitution of a corresponding compound of formula I in which  $R^4$  represents -OH, using an appropriate halogenating agent (e.g., for compounds in which  $R^4$  represents fluoro, reaction with diethylaminosulfurtrifluoride);
- 10 (x) reaction of a corresponding compound of formula XIX,




wherein  $R^1$ ,  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined, with a compound of formula XX,



- 15 wherein  $L^5$  represents a leaving group such as halo, OH, alkane sulfonate, arene sulfonate,  $C_{1-4}$  alkoxy, phenoxy,  $-OC(O)R^{16}$ ,  $-OC(O)OR^{21}$  or  $-OS(O)_2R^{22}$ , and  $R^2$ ,  $R^{16}$ ,  $R^{21}$  and  $R^{22}$  are as hereinbefore defined, for example at between -10°C and reflux temperature, optionally in the presence of a suitable solvent (e.g.  $CHCl_3$ ,  $CH_3CN$ , 2-propanol, diethyl
- 20 ether, or mixtures thereof) and/or an appropriate base (e.g.  $K_2CO_3$ , pyridine or triethylamine);

(y) for compounds of formula I in which  $R^2$  represent  $C_{1-12}$  alkyl, which alkyl group is substituted at the C-2 carbon (relative to the bispidine

nitrogen) with OH or N(H)R<sup>30</sup>, and is otherwise optionally substituted with one or more further substituents as specified hereinbefore for R<sup>2</sup>, reaction of a compound of formula XIX as hereinbefore defined with a compound of formula XXA



wherein X<sub>a</sub> represents O or N(R<sup>30</sup>) and R<sup>2a</sup> represents C<sub>1-10</sub> alkyl, optionally substituted with one or more substituents as specified hereinbefore for R<sup>2</sup>, for example as described hereinbefore for preparation of compounds of formula I (process step (d));

(z) for compounds of formula I in which R<sup>2</sup> represents tetrazol-5-yl, reaction of a corresponding compound of formula I in which R<sup>2</sup> represents -CN with a suitable source of the azide ion (e.g. sodium azide), for example at elevated temperature (e.g. 100°C) in the presence of an appropriate solvent (e.g. DMF) and optionally in the presence of a suitable proton source (e.g. NH<sub>4</sub>Cl);

(aa) for compounds of formula I which are bispidine-nitrogen *N*-oxide derivatives, oxidation of the corresponding bispidine nitrogen of a corresponding compound of formula I, in the presence of a suitable oxidising agent (e.g. *m*CPBA), for example at 0°C in the presence of a suitable organic solvent (e.g. DCM);

(ab) for compounds of formula I which are C<sub>1-4</sub> alkyl quaternary ammonium salt derivatives, in which the alkyl group is attached to a bispidine nitrogen, reaction, at the bispidine nitrogen, of a corresponding compound of formula I with a compound of formula XXI,



wherein  $R^b$  represents  $C_{1-4}$  alkyl and  $L^2$  is as hereinbefore defined, for example at room temperature in the presence of an appropriate organic solvent (e.g. DMF), followed by purification (using e.g. HPLC) in the presence of a suitable counter-ion provider (e.g.  $NH_4OAc$ );


5

(ac) conversion of one substituent on  $R^6$  to another using techniques well known to those skilled in the art; or

(ad) conversion of one  $R^2$  group to another using techniques well known to those skilled in the art.

10


Compounds of formula II may be prepared by reaction of a corresponding compound of formula XXII,



15 wherein  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined, with a compound of formula XX as hereinbefore defined, for example as described hereinbefore for synthesis of compounds of formula I (process step (x)).

20 Compounds of formula II in which  $R^{3a}$  and  $R^{3b}$  both represent H may be prepared by reduction of a corresponding compound of formula XXIII,

33



wherein  $R^2$  and  $R^{41}$  to  $R^{46}$  are as hereinbefore defined, and in which the  $C=O$  group may be activated using an appropriate agent, such as tosylhydrazine, for example as described hereinbefore for the synthesis of  
 5 compounds of formula I (process step (i)).

Compounds of formula II in which one of  $R^{3a}$  and  $R^{3b}$  represents  $-OH$  and the other represents  $C_{1-4}$  alkyl may be prepared by reaction of a compound of formula XXIII, or a protected derivative thereof, with a compound of  
 10 formula XXIV,



wherein  $R^{alk}$  represents  $C_{1-4}$  alkyl and Hal represents chloro, bromo or iodo, for example at between  $-25^\circ C$  and ambient temperature in the presence of a suitable solvent (e.g. diethyl ether).

15

Compounds of formula III may be prepared by standard techniques. For example compounds of formula III in which:

(1) B represents  $-L-O-$  may be prepared by coupling a compound of formula  
 20 X, as hereinbefore defined, to a compound of formula XXV,



wherein  $L^6$  represents a suitable leaving group (e.g. halo) and  $R^4$ ,  $R^5$ , A, L and  $L^1$  are as hereinbefore defined; or

(2) B represents  $-N(R^{13})-L-$  and  $R^4$  and  $R^5$  together represent  $=O$  may be prepared by coupling a compound of formula XXVI,




wherein  $R^6$ ,  $R^{13}$  and L are as hereinbefore defined, to a compound of formula XXVII,



wherein  $L^6$ , A and  $L^1$  are as hereinbefore defined;


in both cases, under conditions which are well known to those skilled in the art.

Compounds of formula III in which A represents  $C_2$ -alkylene and  $R^4$  represents  $-OR^7$ , in which  $R^7$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup> may alternatively be prepared by reaction of a compound of formula XVI as hereinbefore defined with a compound of formula XXVIII,



wherein  $R^y$  represents  $C_{1-4}$  alkyl or aryl (which two groups are optionally substituted with one or more substituents selected from  $C_{1-4}$  alkyl or halo) and  $R^5$ ,  $R^6$  and B are as hereinbefore defined, for example at between ambient temperature (e.g.  $25^\circ\text{C}$ ) and reflux temperature in the presence of a suitable base (e.g.  $K_2CO_3$ ) and an appropriate organic solvent (e.g. acetonitrile), followed by conversion of the ester functionality to an  $L^1$  group (in which  $L^1$  is as hereinbefore defined), under conditions that are well known to those skilled in the art.

Compounds of formula III in which A represents C<sub>2-6</sub> alkylene may be prepared by reduction of a corresponding compound of formula XXIX,




wherein A<sup>b</sup> represents a direct bond or C<sub>1-4</sub> alkylene, and R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and B are as hereinbefore defined, with a suitable borane or borane-Lewis base complex (e.g. borane-dimethyl sulfide) in the presence of an appropriate solvent (e.g. diethyl ether, THF, or a mixture thereof), followed by oxidation of the resulting borane adduct with a suitable oxidising agent (e.g. sodium perborate) and then conversion of the resulting OH group to an L<sup>1</sup> group under conditions known to those skilled in the art.

Compounds of formula III in which A represents C<sub>2-6</sub> alkylene and B represents -L-N(R<sup>13</sup>)- (wherein L represents C<sub>1-4</sub> alkylene) may be prepared by coupling a compound of formula XXX,



wherein R<sup>6</sup> and L<sup>6</sup> are as hereinbefore defined, with a compound of formula XXXI,




wherein L<sup>a</sup> represents C<sub>1-4</sub> alkylene, A<sup>c</sup> represents C<sub>2-6</sub> alkylene, and R<sup>4</sup>, R<sup>5</sup> and R<sup>13</sup> are as hereinbefore defined, for example at between room and reflux temperature, optionally in the presence of a suitable solvent and/or an appropriate base, followed by conversion of the OH group to an L<sup>1</sup> group under conditions known to those skilled in the art.

Compounds of formula III in which B represents -L-S(O)- or -L-S(O)<sub>2</sub>- may be prepared by oxidation of corresponding compounds of formula III in

which B represents -L-S-, wherein L is as hereinbefore defined, in the presence of an appropriate amount of a suitable oxidising agent (e.g. *m*CPBA) and an appropriate organic solvent.

- 5 Compounds of formula V may be prepared in accordance with techniques which are known to those skilled in the art. For example, compounds of formula V in which:


- (1) B represents -CH<sub>2</sub>O- and X represents O may be prepared by reaction of  
 10 a compound of formula X, as hereinbefore defined, with a compound of formula XXXII,



XXXII

- wherein R<sup>5</sup> and L<sup>2</sup> are as hereinbefore defined, for example at elevated temperature (e.g. between 60°C and reflux temperature) in the presence of a  
 15 suitable base (e.g. K<sub>2</sub>CO<sub>3</sub> or NaOH) and an appropriate organic solvent (e.g. acetonitrile or toluene/water), or as otherwise described in the prior art;

- (2) R<sup>5</sup> represents H and X represents O may be prepared by reduction of a compound of formula XXXIII,




XXXIII

- 20 wherein R<sup>6</sup> and B are as hereinbefore defined, for example at between -15°C and room temperature in the presence of a suitable reducing agent (e.g. NaBH<sub>4</sub>) and an appropriate organic solvent (e.g. THF), followed by an internal displacement reaction in the resultant intermediate, for example at


room temperature in the presence of a suitable base (e.g.  $K_2CO_3$ ) and an appropriate organic solvent (e.g. acetonitrile);

(3) B represents  $-L-$ ,  $-L-N(R^{13})-$ ,  $-L-S(O)_2-$  or  $-L-O-$  (in all four of which groups L represents  $C_{1-4}$  alkylene) and X represents O may be prepared by oxidation of a compound of formula XXXIV,



wherein  $B^a$  represents  $-L-$ ,  $-L-N(R^{13})-$ ,  $-L-S(O)_2-$  or  $-L-O-$  (in all four of which groups L represents a single bond or  $C_{1-3}$  alkylene), and  $R^5$ ,  $R^6$  and  $R^{13}$  are as hereinbefore defined, in the presence of a suitable oxidising agent (e.g. *m*CPBA), for example by refluxing in the presence of a suitable organic solvent (e.g. DCM); or

(4) B represents  $-L-O-$  (in which group L represents  $C_{1-4}$  alkylene) and X represents  $N(R^8)$  (wherein  $R^8$  represents  $-C(O)OR^{10b}$  or  $-S(O)_2R^{10c}$ ) may be prepared by cyclisation of a compound of formula XXXV,



wherein  $R^{8b}$  represents  $-C(O)OR^{10b}$  or  $-S(O)_2R^{10c}$  and  $R^5$ ,  $R^6$ ,  $R^{10b}$ ,  $R^{10c}$ ,  $L^a$  and  $L^2$  are as hereinbefore defined, for example at between  $0^\circ C$  and reflux temperature in the presence of a suitable base (e.g. sodium hydroxide), an appropriate solvent (e.g. dichloromethane, water, or a mixture thereof) and, if necessary a phase transfer catalyst (such as tetrabutylammonium hydrogensulfate).

Compounds of formula VII may be prepared by reaction of a corresponding compound of formula VI with hydroxylamine, for example at elevated temperature (e.g. at reflux) in the presence of a suitable organic solvent (e.g. methanol).

5

Compounds of formulae IX, XI, XIII and XVII may be prepared in a similar fashion to compounds of formula I (see, for example, process steps (a), (b) and (x)).

- 10 Compounds of formula XIII may alternatively be prepared by reaction of a corresponding compound of formula I in which  $R^4$  represents -D-OH, with a compound of formula XXXVI,



XXXVI

- 15 wherein  $R^y$  is as hereinbefore defined, for example at between -10 and 25°C in the presence of a suitable solvent (e.g. dichloromethane), followed by reaction with a suitable source of the azide ion (e.g. sodium azide) for example at between ambient and reflux temperature in the presence of an appropriate solvent (e.g. DMF) and a suitable base (e.g.  $NaHCO_3$ ).

- 20 Compounds of formula XVII may alternatively be prepared by replacement of the OH group of a compound of formula I in which  $R^4$  represents -D-OH with an  $L^2$  group under conditions that are well known to those skilled in the art.

- 25 Compounds of formula XIX may be prepared by reaction of a corresponding compound of formula XXII, as hereinbefore defined, with a compound of formula III, as hereinbefore defined.


- 30 Compounds of formula XIX in which A represents  $C_2$  alkylene and  $R^4$  and  $R^5$  together represent =O may be prepared by reaction of a corresponding

compound of formula XXII, as hereinbefore defined, with a compound of formula IV, as hereinbefore defined, for example as described hereinbefore for synthesis of compounds of formula I (process step (b)).

- 5 Compounds of formula XIX in which A represents  $\text{CH}_2$  and  $\text{R}^4$  represents  $-\text{OH}$  or  $-\text{N}(\text{H})\text{R}^8$  may be prepared by reaction of a corresponding compound of formula XXII, as hereinbefore defined, with a compound of formula V as hereinbefore defined, for example as described hereinbefore for synthesis of compounds of formula I (process step (d)).

10

Compounds of formula XIX in which  $\text{R}^{3a}$  and  $\text{R}^{3b}$  both represent H may alternatively be prepared by reduction of a corresponding compound of formula XXXVII,



XXXVII

- 15 wherein  $\text{R}^1$  and  $\text{R}^{41}$  to  $\text{R}^{46}$  are as hereinbefore defined, and in which the  $\text{C}=\text{O}$  group may be activated using an appropriate agent, such as tosylhydrazine, for example as described hereinbefore for the synthesis of compounds of formula I (process step (i)).

- 20 Compounds of formulae II and XIX in which one or more of  $\text{R}^{41}$ ,  $\text{R}^{42}$ ,  $\text{R}^{45}$  and/or  $\text{R}^{46}$  represent  $\text{C}_{1-3}$  alkyl may alternatively be prepared by reaction of a compound of formula II or XIX (as appropriate) in which  $\text{R}^{41}$ ,  $\text{R}^{42}$ ,  $\text{R}^{45}$  and/or  $\text{R}^{46}$  (as appropriate) represent H, with an appropriate alkylating agent (e.g. dimethyl sulfate), for example in the presence of a suitable strong base (e.g. *s*-


BuLi), N,N,N',N'-tetramethylethylenediamine and a reaction-inert solvent (e.g. THF).

Compounds of formula XX in which  $R^2$  represents  $-C(=N-CN)N(R^{25})R^{26}$  and  $L^5$  represents phenoxy may be prepared by reaction of a corresponding compound of formula XXXVIII,



wherein  $R^{25}$  and  $R^{26}$  are as hereinbefore defined, with diphenyl cyanocarbonimidoate, for example at between  $-10^\circ\text{C}$  and room temperature in the presence of an appropriate solvent (e.g. isopropanol).


Compounds of formula XXII are known in the literature or are readily available using known techniques. For example, compounds of formula XXII in which  $R^{3a}$  and  $R^{3b}$  together represent  $\text{C}_{3-5}$  alkylene,  $-\text{O}-\text{Z}-\text{O}-$ ,  $-\text{O}-\text{Z}-\text{S}-$  or  $-\text{S}-\text{Z}-\text{S}-$  and  $R^{41}$  to  $R^{46}$  all represent H, may be prepared by reduction of a compound of formula XXXIX,



wherein  $R^c$  and  $R^d$  together represent  $\text{C}_{3-5}$  alkylene,  $-\text{O}-\text{Z}-\text{O}-$ ,  $-\text{O}-\text{Z}-\text{S}-$  or  $-\text{S}-\text{Z}-\text{S}-$ , wherein Z is as hereinbefore defined, in the presence of a suitable reducing agent (e.g.  $\text{LiAlH}_4$ ) under conditions that are well known to those skilled in the art.


Compounds of formula XXIX in which B represents  $\text{C}_{1-4}$  alkylene may be prepared by coupling a compound of formula XL,

41




wherein B<sup>b</sup> represents C<sub>1-4</sub> alkylene and Hal, A<sup>b</sup>, R<sup>4</sup> and R<sup>5</sup> are as hereinbefore defined, with a compound of formula XXX, as hereinbefore defined, for example at between -25°C and room temperature in the presence of a suitable zinc(II) salt (e.g. anhydrous ZnBr<sub>2</sub>), an appropriate catalyst (e.g. Pd(PPh<sub>3</sub>)<sub>4</sub>) and a reaction-inert organic solvent (e.g. THF, toluene or diethyl ether).

Compounds of formulae VI, XXIII and XXXVII (in which, in all cases, R<sup>41</sup> and R<sup>42</sup> both represent H), may be prepared, advantageously, by reaction of (as appropriate) either (i) a compound of formula XLI,



wherein R<sup>z</sup> represents C<sub>1-10</sub> alkyl or C<sub>1-3</sub> alkylaryl (e.g. alkylphenyl, such as benzyl) and R<sup>43</sup> to R<sup>46</sup> are as hereinbefore defined, or (ii) 4-piperidone (or a protected derivative thereof), with (as appropriate) either (1) a compound of formula XLII,



wherein R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, A and B are as hereinbefore defined, or (2) NH<sub>3</sub> (or a protected (e.g. benzyl) derivative thereof), in all cases in the presence of a formaldehyde (i.e. an appropriate source of formaldehyde, such as paraformaldehyde or formalin solution) and, in the case of compounds of formulae VI and XXIII, conversion of the C(O)OR<sup>z</sup> group in the resultant

intermediate to an  $R^2$  group using techniques such as those described herein (e.g. removal of the  $C(O)OR^z$  group followed by carrying out a coupling, e.g. according to process step (x) above).

- 5 The formation of compounds of formulae VI, XXIII and XXXVII may be carried out in this way for example at between room temperature and reflux (depending upon the concentration of the reactants) in the presence of an appropriate solvent (e.g. ethanol or methanol) and, preferably, in the presence of an organic acid (e.g. a  $C_{1-6}$  carboxylic acid, especially acetic  
10 acid).

It will be also appreciated by those skilled in the art that compounds of formula XXII in which  $R^{3a}$  and  $R^{3b}$  both represent H may also be prepared *via* this method (i.e. by reaction of a 4-piperidone (or a protected derivative  
15 thereof) with  $NH_3$  (or a protected derivative thereof) in the presence of a formaldehyde), provided that the intermediate so formed is subsequently reduced under appropriate reaction conditions.


The skilled person will also appreciate that this process may also be used to  
20 prepare compounds of formula I in which  $R^{45}$  and  $R^{46}$  are H, and  $R^{41}$  and/or  $R^{42}$  are other than H, for example by:

- (i) reacting a compound of formula XLI in which  $R^{45}$  and/or  $R^{46}$  is/are other than H with, for example, benzylamine or a derivative thereof;
- (ii) removal of the  $-C(O)OR^z$  unit;
- 25 (iii) reaction at the free bispidine nitrogen of the resultant compound with a compound of formula III, IV or V (as appropriate), as hereinbefore defined;
- (iv) removal of the benzyl protecting group; and
- (v) reaction at the free bispidine nitrogen of the resultant compound with a  
30 compound of formula XX as hereinbefore defined,

under conditions well known to those skilled in the art including those described hereinbefore. This reaction will be accompanied by, at some point, conversion of the bridgehead carbonyl functionality to the desired  $R^{3a}/R^{3b}$  groups.

5

Compounds of formula XXXIX may be prepared in accordance with techniques which are well known to those skilled in the art. For example, compounds of formula XXXIX in which  $R^c$  and  $R^d$  together represent  $C_{3-5}$  alkylene may be prepared by reaction of a compound of formula XLIII,



XLIII

10

wherein  $R^e$  and  $R^f$  together represent  $C_{3-5}$  alkylene, with a mixture of phosphoric acid and sulfuric acid, for example at  $120^\circ\text{C}$ .

15

Compounds of formula XLII are well known in the literature or are readily available using known techniques. For example, compounds of formula XLII in which  $R^4$  represents OH,  $R^5$  represents H and A represents  $\text{CH}_2$  may be prepared by reaction of a corresponding compound of formula V wherein  $R^5$  represents H and X represents O with ammonium hydroxide under conditions which are well known to those skilled in the art.

20

Compounds of formulae IV, VIII, X, XII, XIV, XV, XVI, XVIII, XX (in which  $R^2$  represents other than  $-\text{C}(=\text{N}-\text{CN})\text{N}(\text{R}^{25})\text{R}^{26}$  and/or  $\text{L}^5$  represents other than phenoxy), XXA, XXI, XXIV, XXV, XXVI, XXVII, XXVIII, XXX, XXXI, XXXII, XXXIII, XXXIV, XXXV, XXXVI, XXXVIII, XL, XLI, XLIII and derivatives thereof, are either commercially available, are

25

known in the literature, or may be obtained either by analogy with the processes described herein, or by conventional synthetic procedures, in accordance with standard techniques, from readily available starting materials using appropriate reagents and reaction conditions.

5

Substituents on the aryl (e.g. phenyl), and (if appropriate) heterocyclic, group(s) in compounds defined herein may be converted to other claimed substituents using techniques well known to those skilled in the art. For example, hydroxy may be converted to alkoxy, phenyl may be halogenated  
10 to give halophenyl, nitro may be reduced to give amino, amino may be acetylated to give acetylamino, etc.

The skilled person will also appreciate that various standard substituent or functional group interconversions and transformations within certain  
15 compounds of formula I will provide other compounds of formula I. For example, carbonyl may be reduced to hydroxy or alkylene, hydroxy may be acylated to give alkylcarbonyloxy, nitro may be reduced to amino, amido may be reduced to amino, amino may be sulfonated or acylated to give sulfonylamino or acylamino (respectively), and certain acyclic groups may  
20 be converted to certain heterocyclic groups under conditions known to those skilled in the art, for example as described in Comprehensive Heterocyclic Chemistry II, edited by AR Katritzky, CW Rees and EFV Scriven, 1<sup>st</sup> Edition, Elsevier Science Ltd., Volumes 1-11 (1996).

25 The compounds of the invention may be isolated from their reaction mixtures using conventional techniques.

It will be appreciated by those skilled in the art that, in the process described above, the functional groups of intermediate compounds may be,  
30 or may need to be, protected by protecting groups.

Functional groups which it is desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl and diarylalkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl and alkylcarbonyl groups (e.g. methyl- and ethylcarbonyl groups). Suitable protecting groups for amino include benzyl, *tert*-butyloxycarbonyl, 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for amidino and guanidino include benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C<sub>1-6</sub> alkyl or benzyl esters.

The protection and deprotection of functional groups may take place before or after any of the reaction steps described hereinbefore.

Protecting groups may be removed in accordance with techniques which are well known to those skilled in the art and as described hereinafter.

The use of protecting groups is fully described in "Protective Groups in Organic Chemistry", edited by J.W.F. McOmie, Plenum Press (1973), and "Protective Groups in Organic Synthesis", 3<sup>rd</sup> edition, T.W. Greene & P.G.M. Wutz, Wiley-Interscience (1999).

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative, and, on some occasions, more convenient, manner, the individual process steps mentioned herein may be performed in a different order, and/or the individual reactions may be performed at a different stage in the overall route (i.e. substituents may be added to and/or chemical transformations performed upon, different intermediates to those associated hereinbefore with a particular reaction).

This will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the availability of key intermediates and the protecting group strategy (if any) to be adopted. Clearly, the type of chemistry involved will influence the choice of reagent that is used in the said synthetic steps, the need, and type, of protecting groups that are employed, and the sequence for accomplishing the synthesis.

It will also be appreciated by those skilled in the art that, although certain protected derivatives of compounds of formula I, which may be made prior to a final deprotection stage, may not possess pharmacological activity as such, they may be administered parenterally or orally and thereafter metabolised in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". Moreover, certain compounds of formula I may act as prodrugs of other compounds of formula I.

All prodrugs of compounds of formula I are included within the scope of the invention.

Some of the intermediates referred to hereinbefore are novel. According to a further aspect of the invention there is thus provided: (a) a compound of formula II, as hereinbefore defined, or a protected derivative thereof; (b) a compound of formula VI, as hereinbefore defined, or a protected derivative thereof; (c) a compound of formula VII, as hereinbefore defined, or a protected derivative thereof; (d) a compound of formula IX, as hereinbefore defined, or a protected derivative thereof; (e) a compound of formula XI, as hereinbefore defined, or a protected derivative thereof; (f) a compound of formula XIII, as hereinbefore defined, or a protected derivative thereof; (g) a compound of formula XVII, as hereinbefore defined, or a protected derivative thereof; (h) a compound of formula XIX, as hereinbefore defined

(provided that at least one of  $R^{3a}$  and  $R^{3b}$  represents  $-N(R^{39})R^{38c}$ , wherein  $R^{39}$  represents a structural fragment of formula Ia, as hereinbefore defined), or a protected derivative thereof; (i) a compound of formula XXII, as hereinbefore defined (provided that at least one of  $R^{3a}$  and  $R^{3b}$  represents  $-N(R^{39})R^{38c}$ , wherein  $R^{39}$  represents a structural fragment of formula Ia, as  
 5 hereinbefore defined), or a protected derivative thereof; and (j) a compound of formula XXIII, as hereinbefore defined, or a protected derivative thereof.

Compounds of formula II that may be mentioned include those in which:

10 when  $R^{3a}$  and  $R^{3b}$  independently represent H,  $C_{1-4}$  alkyl, OH or  $N(R^{39})R^{38c}$ ; and  
 $R^{39}$  represents H or  $C_{1-6}$  alkyl;  
 then  $R^2$  does not represent:

- (i) CN;
- 15 (ii)  $C_{1-6}$  alkyl optionally substituted by OH,  $N(R^{30})R^{31}$  or Het<sup>9</sup>; wherein  $R^{30}$  and  $R^{31}$  independently represent H,  $C_{1-6}$  alkyl or  $C_{3-8}$  cycloalkyl); and  
 Het<sup>9</sup> represents an unsubstituted, saturated 3- to 8-membered heterocycle containing one nitrogen atom (*via* which atom the  
 20 heterocyclic group is attached to the rest of the molecule);
- (iii)  $C_{1-6}$  *n*-alkyl, which alkyl group is optionally interrupted by O and is terminated by  $N(R^{30})R^{31}$  or  $OR^{32}$ ; wherein  
 one of  $R^{30}$  and  $R^{31}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$   
 25 and the other represents H or  $C_{1-6}$  alkyl; and  
 $R^{32}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ;
- (iv)  $-C(O)R^{16}$ , wherein  $R^{16}$  represents H or phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $CO_2H$  or  $NH_2$ ;

- (v)  $-\text{S}(\text{O})_2\text{R}^{22}$ , wherein  $\text{R}^{22}$  represents phenyl substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $\text{CO}_2\text{H}$  or  $\text{NH}_2$ ;
- (vi)  $-\text{S}(\text{O})_2\text{N}(\text{R}^{23})\text{R}^{24}$ ; wherein  
 $\text{R}^{23}$  represents phenyl substituted in the *meta*- or *para*-position  
 5 (relative to the point of attachment) by  $\text{CO}_2\text{H}$  or  $\text{NH}_2$ ; and  
 $\text{R}^{24}$  represents H or  $\text{C}_{1-6}$  alkyl; and
- (vii)  $\text{C}_{3-4}$  *n*-alkyl, which alkyl group is terminated by phenyl, which phenyl group is substituted in the *meta*- or *para*-position (relative to the point of attachment) by  $\text{CO}_2\text{H}$  or  $\text{NH}_2$ , and which alkyl group is interrupted  
 10 at the  $\beta$ -position (relative to the point of attachment of the phenyl group) by O.

Compounds of formula II, IX and XI that may be mentioned include those in which:

- 15 when  $\text{R}^{3a}$  and  $\text{R}^{3b}$  independently represent H,  $\text{C}_{1-4}$  alkyl, OH or  $\text{N}(\text{R}^{39})\text{R}^{38c}$ ;  
 $\text{R}^{39}$  represents H or  $\text{C}_{1-6}$  alkyl; and  
 $\text{R}^4$  and  $\text{R}^5$  both represent H;  
 then  $\text{R}^2$  does not represent CN,  $-\text{C}(\text{O})\text{R}^{16}$ ,  $-\text{S}(\text{O})_2\text{R}^{22}$ ,  $-\text{S}(\text{O})_2\text{N}(\text{R}^{23})\text{R}^{24}$  or  
 $\text{C}_{1-6}$  alkyl optionally substituted as defined hereinbefore in respect of  $\text{R}^2$ .

20

Further compounds of formulae II, IX and XI that may be mentioned include those in which:

- $\text{R}^2$  represents  $\text{Het}^8$ ,  $-\text{C}(\text{O})\text{R}^{16}$ ,  $-\text{C}(\text{S})\text{OR}^{17}$ ,  $-\text{C}(\text{S})\text{N}(\text{R}^{18})\text{R}^{19}$ ,  
 $-\text{[C}(\text{O})\text{]}_2\text{N}(\text{R}^{20a})\text{R}^{20b}$ ,  $-\text{[C}(\text{O})\text{]}_2\text{OR}^{21}$ ,  $-\text{S}(\text{O})_2\text{R}^{22}$ ,  $-\text{S}(\text{O})_2\text{N}(\text{R}^{23})\text{R}^{24}$ ,  
 25  $-\text{C}(=\text{N}-\text{CN})\text{N}(\text{R}^{25})\text{R}^{26}$ ,  $-\text{C}(=\text{N}-\text{CN})\text{OR}^{27}$  or  $\text{C}_{1-12}$  alkyl (which alkyl group is substituted and/or terminated by one or more substituents selected from  
 $-\text{C}(\text{O})\text{R}^{28}$ ,  $-\text{C}(\text{O})\text{N}(\text{R}^{29a})\text{R}^{29b}$ ,  $-\text{N}(\text{R}^{30})\text{R}^{31}$ ,  $-\text{OR}^{32}$ ,  $-\text{S}(\text{O})\text{R}^{33}$ , halo, -CN and nitro);

$R^{16}$  represents  $Het^{10}$  or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN,  $-N(R^{34})R^{35}$ , aryl and  $Het^{11}$ );

$R^{22}$  represents  $Het^{14}$  or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH, -CN,  $Het^{15}$  and aryl);

$R^{23}$  represents H,  $C_{1-6}$  alkyl,  $Het^{16}$ ,  $-C(O)R^{36a}$ ,  $-C(O)OR^{36b}$  or  $-C(O)SR^{36b}$ ;

$R^{30}$  represents  $Het^{19}$ ,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ ;

$R^{32}$  represents  $C_{1-6}$  alkyl,  $Het^{21}$ ,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ .

### Medical and pharmaceutical use

The compounds of the invention are useful because they possess pharmacological activity. They are therefore indicated as pharmaceuticals.

Thus, according to a further aspect of the invention there is provided the compounds of the invention for use as pharmaceuticals.

In particular, the compounds of the invention exhibit myocardial electrophysiological activity, for example as demonstrated in the test described below.

The compounds of the invention are thus expected to be useful in both the prophylaxis and the treatment of arrhythmias, and in particular atrial and ventricular arrhythmias.

The compounds of the invention are thus indicated in the treatment or prophylaxis of cardiac diseases, or in indications related to cardiac diseases, in which arrhythmias are believed to play a major role, including ischaemic

heart disease, sudden heart attack, myocardial infarction, heart failure, cardiac surgery and thromboembolic events.

In the treatment of arrhythmias, compounds of the invention have been found to selectively delay cardiac repolarization, thus prolonging the QT interval, and, in particular, to exhibit class III activity. Although compounds of the invention have been found to exhibit class III activity in particular, in the treatment of arrhythmias, their mode(s) of activity is/are not necessarily restricted to this class.

According to a further aspect of the invention, there is provided a method of treatment of an arrhythmia which method comprises administration of a therapeutically effective amount of a compound of the invention to a person suffering from, or susceptible to, such a condition.

### **Pharmaceutical preparations**

The compounds of the invention will normally be administered orally, subcutaneously, intravenously, intraarterially, transdermally, intranasally, by inhalation, or by any other parenteral route, in the form of pharmaceutical preparations comprising the active ingredient either as a free base, a pharmaceutically acceptable ion exchanger or a non-toxic organic or inorganic acid addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated, as well as the route of administration, the compositions may be administered at varying doses.

The compounds of the invention may also be combined with any other drugs useful in the treatment of arrhythmias and/or other cardiovascular disorders.

According to a further aspect of the invention there is thus provided a pharmaceutical formulation including a compound of the invention in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

Suitable daily doses of the compounds of the invention in therapeutic  
5 treatment of humans are about 0.005 to 10.0 mg/kg body weight at oral administration and about 0.005 to 5.0 mg/kg body weight at parenteral administration.

The compounds of the invention have the advantage that they are effective  
10 against cardiac arrhythmias.

Compounds of the invention may also have the advantage that they may be more efficacious than, be less toxic than, have a broader range of activity (including exhibiting any combination of class I, class II, class III and/or  
15 class IV activity (especially class I and/or class IV activity in addition to class III activity)) than, be more potent than, be longer acting than, produce fewer side effects (including a lower incidence of proarrhythmias such as *torsades de pointes*) than, be more easily absorbed than, or that they may have other useful pharmacological properties over, compounds known in  
20 the prior art.

## Biological Tests

### Test A

#### 25 Primary Electrophysiological Effects In Anaesthetised Guinea Pigs

Guinea pigs weighing between 660 and 1100 g were used. The animals were housed for at least one week before the experiment and had free access to food and tap water during that period.

Anaesthesia was induced by an intraperitoneal injection of pentobarbital (40 to 50 mg/kg) and catheters were introduced into one carotid artery (for blood pressure recording and blood sampling) and into one jugular vein (for drug infusions). Needle electrodes were placed on the limbs for recording of ECGs (lead II). A thermistor was placed in the rectum and the animal was placed on a heating pad, set to a rectal temperature of between 37.5 and 38.5°C.

A tracheotomy was performed and the animal was artificially ventilated with room air by use of a small animal ventilator, set to keep blood gases within the normal range for the species. In order to reduce autonomic influences both vagi were cut in the neck, and 0.5 mg/kg of propranolol was given intravenously, 15 minutes before the start of the experiment.

The left ventricular epicardium was exposed by a left-sided thoracotomy, and a custom-designed suction electrode for recording of the monophasic action potential (MAP) was applied to the left ventricular free wall. The electrode was kept in position as long as an acceptable signal could be recorded, otherwise it was moved to a new position. A bipolar electrode for pacing was clipped to the left atrium. Pacing (2 ms duration, twice the diastolic threshold) was performed with a custom-made constant current stimulator. The heart was paced at a frequency just above the normal sinus rate during 1 minute every fifth minute throughout the study.

The blood pressure, the MAP signal and the lead II ECG were recorded on a Mingograph ink-jet recorder (Siemens-Elema, Sweden). All signals were collected (sampling frequency 1000 Hz) on a PC during the last 10 seconds of each pacing sequence and the last 10 seconds of the following minute of sinus rhythm. The signals were processed using a custom-made program developed for acquisition and analysis of physiological signals measured in

experimental animals (see Axenborg and Hirsch, *Comput. Methods Programs Biomed.* 41, 55 (1993)).

The test procedure consisted of taking two basal control recordings, 5 minutes apart, during both pacing and sinus rhythm. After the second control recording, the first dose of the test substance was infused in a volume of 0.2 mL into the jugular vein catheter for 30 seconds. Three minutes later, pacing was started and a new recording was made. Five minutes after the previous dose, the next dose of test substance was administered. Six to ten  
10 consecutive doses were given during each experiment.

#### Data analysis

Of the numerous variables measured in this analysis, three were selected as the most important for comparison and selection of active compounds. The  
15 three variables selected were the MAP duration at 75 percent repolarization during pacing, the atrio-ventricular (AV) conduction time (defined as the interval between the atrial pace pulse and the start of the ventricular MAP) during pacing, and the heart rate (defined as the RR interval during sinus rhythm). Systolic and diastolic blood pressure were measured in order to  
20 judge the haemodynamic status of the anaesthetised animal. Further, the ECG was checked for arrhythmias and/or morphological changes.

The mean of the two control recordings was set to zero and the effects recorded after consecutive doses of test substance were expressed as  
25 percentage changes from this value. By plotting these percentage values against the cumulative dose administered before each recording, it was possible to construct dose-response curves. In this way, each experiment generated three dose-response curves, one for MAP duration, one for AV-conduction time and one for the sinus frequency (RR interval). A mean  
30 curve of all experiments performed with a test substance was calculated, and

potency values were derived from the mean curve. All dose-response curves in these experiments were constructed by linear connection of the data points obtained. The cumulative dose prolonging the MAP duration by 10% from the baseline was used as an index to assess the class III electrophysiological  
5 potency of the agent under investigation ( $D_{10}$ ).

### Test B

#### Glucocorticoid-treated mouse fibroblasts as a model to detect blockers of the delayed rectifier K current

10 IC<sub>50</sub> for K channel blockade was determined using a microtitre plate based screen method, based on membrane potential changes of glucocorticoid-treated mouse fibroblasts. The membrane potential of glucocorticoid-treated mouse fibroblasts was measured using fluorescence of the bisoxonol dye DiBac<sub>4(3)</sub>, which could be reliably detected using a fluorescence laser  
15 imaging plate reader (FLIPR). Expression of a delayed rectifier potassium channel was induced in mouse fibroblasts by 24 hours exposure to the glucocorticoid dexamethasone (5  $\mu$ M). Blockade of these potassium channels depolarised the fibroblasts, resulting in increased fluorescence of DiBac<sub>4(3)</sub>.

20

Mouse ltk fibroblasts (L-cells) were purchased from American Type Culture Collection (ATCC, Manassa, VA), and were cultured in Dulbeccos modified eagle medium supplemented with fetal calf serum (5% vol/vol), penicillin (500 units/mL), streptomycin (500  $\mu$ g/mL) and L-alanine-L-  
25 glutamine (0.862 mg/mL). The cells were passaged every 3-4 days using trypsin (0.5 mg/mL in calcium-free phosphate buffered saline, Gibco BRL). Three days prior to experiments, cell-suspension was pipetted out into clear-bottom, black plastic, 96-well plates (Costar) at 25 000 cells/well.

The fluorescence probe DiBac<sub>4(3)</sub> (DiBac Molecular probes) was used to measure membrane potential. DiBac<sub>4(3)</sub> maximally absorbs at 488 nM and emits at 513 nM. DiBac<sub>4(3)</sub> is a bisoxonol, and thus is negatively charged at pH 7. Due to its negative charge, the distribution of DiBac<sub>4(3)</sub> across the membrane is dependent upon the transmembrane potential: if the cell depolarizes (i.e. the cell interior becomes less negative relative to cell exterior), the DiBac<sub>4(3)</sub> concentration inside the cell increases, due to electrostatic forces. Once inside the cell, DiBac<sub>4(3)</sub> molecules can bind to lipids and proteins, which causes an increase in fluorescence emission. Thus, a depolarization will be reflected by an increase in DiBac<sub>4(3)</sub> fluorescence. The change in DiBac<sub>4(3)</sub> fluorescence was detected by a FLIPR.

Prior to each experiment, the cells were washed 4 times in phosphate-buffered saline (PBS) to remove all culture media. The cells were then treated with 5  $\mu$ M DiBac<sub>4(3)</sub> (in 180  $\mu$ L of PBS) at 35°C. Once a stable fluorescence was reached (usually after 10 min), 20  $\mu$ L of the test substance was added, using FLIPR's internal 96 well pipetting system. Fluorescence measurements were then taken every 20 sec for a further 10 min. All experiments were carried out at 35°C, due to the high temperature sensitivity of both delayed rectifier potassium channel conductance and DiBac<sub>4(3)</sub> fluorescence. Test substances were prepared in a second 96 well plate, in PBS containing 5  $\mu$ M DiBac<sub>4(3)</sub>. The concentration of substance prepared was 10 times that of the desired concentration in the experiment as an additional 1:10 dilution occurred during addition of substance during the experiment. Dofetilide (10  $\mu$ M) was used as a positive control, i.e. to determine the maximum increase in fluorescence.

Curve-fitting, used to determine the IC<sub>50</sub> values, was performed with the Graphpad Prism program (Graphpad Software Inc., San Diego, CA).

### Test C

#### 5 Metabolic Stability of Test Compounds

An *in vitro* screen was set up to determine the metabolic stability of the compounds of the invention.

The hepatic S-9 fraction from dog, man, rabbit and rat with NADPH as co-  
10 factor was used. The assay conditions were as follows: S-9 (3 mg/mL), NADPH (0.83 mM), Tris-HCl buffer (50 mM) at pH 7.4 and 10  $\mu$ M of test compound.

The reaction was started by addition of test compound and terminated after 0,  
15 1, 5, 15 and 30 minutes by raising the pH in the sample to above 10 (NaOH; 1 mM). After solvent extraction, the concentration of test compound was measured against an internal standard by LC (fluorescence/UV detection).

The percentage of test compound remaining after 30 minutes (and thus  $t_{1/2}$ )  
20 was calculated and used as a measure for metabolic stability.

The invention is illustrated by way of the following examples.

### **Examples**

25

#### General Experimental Procedures

Mass spectra were recorded on one of the following instruments: a Finnigan  
MAT TSQ 700 triple quadrupole mass spectrometer equipped with an  
electrospray interface (FAB-MS); a Perkin-Elmer SciX API 150ex  
30 spectrometer; a VG Quattro II triple quadrupole; a VG Platform II single

quadrupole; or a Micromass Platform LCZ single quadrupole mass spectrometer (the latter three instruments were equipped with a pneumatically assisted electrospray interface (LC-MS)).  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR measurements were performed on a BRUKER ACP 300 and Varian  
5 300, 400 and 500 spectrometers, operating at  $^1\text{H}$  frequencies of 300, 400 and 500 MHz respectively, and at  $^{13}\text{C}$  frequencies of 75.5, 100.6 and 125.7 MHz respectively. Alternatively,  $^{13}\text{C}$  NMR measurements were performed on a BRUKER ACE 200 spectrometer at a frequency of 50.3 MHz.

10

Rotamers may or may not be denoted in spectra depending upon ease of interpretation of spectra. Unless otherwise stated, chemical shifts are given in ppm with the solvent as internal standard.

15 Example 1

*tert*-Butyl 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo-[3.3.1]non-3-yl}ethylcarbamate

(i) *tert*-Butyl 2-bromoethylcarbamate

20 A mixture of 2-bromoethylamine hydrobromide (10.0 g, 0.049 mol), NaOH (1.84 g, 0.046 mol), water (50 mL) and THF (200 mL) was cooled to 0°C. Di-*tert*-butyl dicarbonate (10.1 g, 0.046 mol) was added slowly to the mixture, which was then stirred at rt overnight. The mixture was concentrated *in vacuo* and the residue dissolved in DCM. This organic  
25 solution was washed with water and purified by chromatography on silica, eluting with DCM, to give 5.6 g (50%) of the title compound.

(ii) 4-[3-(3,7-Diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile

HCl-saturated EtOAc (600 mL) was added to a solution of *tert*-butyl 7-[3-  
30 (4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]nonane-3-

carboxylate (62 g; see Example 2 of international patent application No. PCT/SE98/02276) in EtOAc (600 mL) and the mixture was stirred at rt. for 4 h. The solvent was removed under reduced pressure, the residue was dissolved in MeCN (1.3 L) and K<sub>2</sub>CO<sub>3</sub> (100 g) was added. The suspension  
5 was stirred for 12 h and filtered. Concentration of the filtrate gave the title compound in a 90% yield.

<sup>13</sup>C NMR (CDCl<sub>3</sub>) : δ 28.9, 29.2, 32.3, 50.9, 57.7, 60.8, 62.1, 66.0, 71.2, 104.0, 115.3, 119.1, 133.9, 162.1.

10 (iii) tert-Butyl 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diaza-bicyclo[3.3.1]non-3-yl}ethylcarbamate

4-[3-(3,7-Diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile (see step (ii) above; 7.7 g, 25.6 mmol) and *tert*-butyl 2-bromoethylcarbamate (see step (i) above; 5.7 g, 2.56 mmol) and K<sub>2</sub>CO<sub>3</sub> (3.5 g, 2.56 mmol) were  
15 mixed in CH<sub>3</sub>CN (50 mL) and stirred at 60°C for 60 h. The reaction mixture was filtered and evaporated. The residue was purified using column chromatography (DCM: 10-20% MeOH saturated with NH<sub>3</sub> (g)), to yield 8 g (71%) of the title compound.

<sup>13</sup>C NMR (CDCl<sub>3</sub>) : δ 28.4, 29.6, 30.3, 32.0, 36.9, 54.8, 58.4, 58.6, 59.5,  
20 64.8, 71.1, 78.8, 104.1, 115.3, 119.2, 133.9, 156.4, 162.2.

FAB-MS (M+1)<sup>+</sup> = 445 (m/z)

Example 2

25 4-{3-[7-(3,3-Dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile

4-[3-(3,7-Diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile (see Example 1(ii) above; 0.6 g, 2.0 mmol) and 1-chloropinacolone (0.27 g, 2.0 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.27 g, 20 mmol) were mixed in CH<sub>3</sub>CN

and stirred at 60°C for 1 h, and then at r.t. overnight. The reaction mixture was filtered and evaporated, giving 0.7 g (90%) of the title compound.

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) :  $\delta$  26.2, 30.0, 30.6, 31.9, 43.5, 55.1, 57.3, 57.6, 59.2, 59.8, 61.7, 64.8, 71.1, 103.9, 115.3, 119.2, 133.9, 162.3, 212.2.

5 FAB-MS  $(\text{M}+1)^+ = 400$  (m/z)

### Example 3

4-{3-[7-(2-Ethyl-2H-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile

10

(i) 3,7-Dibenzyl-3,7-diazabicyclo[3.3.1]nonane-9-one

The sub-title compound was prepared according to the procedure described in *J. Org. Chem.*, 41(9), 1976, pp. 1593-1597.

15 (ii) 3,7-Dibenzyl-3,7-diazabicyclo[3.3.1]nonane

The sub-title compound was prepared according to the procedure described in *J. Org. Chem.*, 41(9), 1976, pp. 1593-1597, using 3,7-dibenzyl-3,7-diazabicyclo[3.3.1]nonane-9-one (see step (i) above) in place of *N*-benzyl-*N'*-methylbispidone.

20

(iii) 3-Benzyl-3,7-diazabicyclo[3.3.1]nonane

A solution of 3,7-dibenzyl-3,7-diazabicyclo[3.3.1]nonane (see step (ii) above; 97 g, 6.4 mmol) in aqueous ethanol (95%) was hydrogenated over 5% Pd/C at 1 atm. until tlc indicated that the reaction was complete. The catalyst was removed by filtration through a pad of Celite®, and the filtrate concentrated under reduced pressure to give the sub-title compound in quantitative yield.

25

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) :  $\delta$  30.1, 33.4, 36.0, 52.5, 59.6, 64.3, 126.9, 128.3, 128.7, 138.8

(iv) 3-Benzyl-7-cyano-3,7-diazabicyclo[3.3.1]nonane

3-Benzyl-3,7-diazabicyclo[3.3.1]nonane (see step (iii) above; 20 g, 92 mmol) was dissolved in ether (120 mL). Cyanogen bromide (9.8 g, 92 mmol) dissolved in ether (80 mL) was added dropwise at 0°C. The mixture was stirred at 0°C for 15 minutes, and then at r.t. overnight, after which a white precipitate formed. The ether was evaporated. Water and a saturated Na<sub>2</sub>CO<sub>3</sub> (aq) solution were added. The mixture was extracted with ether. The ether layer was separated and dried (MgSO<sub>4</sub>), giving 20.3 g (91%) of the sub-title compound.

10

(v) 3-Benzyl-7-(2H-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]nonane, ammonium salt

A mixture of 3-benzyl-7-cyano-3,7-diazabicyclo[3.3.1]nonane (see step (iv) above; 10.2 g, 42 mmol), NaN<sub>3</sub> (2.92 g, 45 mmol), NH<sub>4</sub>Cl (2.41 g, 45 mmol) and DMF (50 mL) was stirred at 100°C for 22 h. DMF was evaporated, toluene was added and evaporated, which resulted in 12 g of an orange-coloured powder. The product was purified by preparative reversed phase HPLC, giving 5.8 g (46%) of the sub-title compound.

20 (vi) 3-Benzyl-7-(2-ethyl-2H-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]nonane

A mixture of 3-benzyl-7-(2H-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]nonane, ammonium salt (see step (v) above; 4 g, 13 mmol), ethyl iodide (2.20 mL, 26 mmol) and NaOH (0.62 g, 15.6 mmol) was refluxed for 2 h. The solvent was evaporated and the residue purified by flash chromatography (hexane:ethyl acetate (1:1), MeOH (NH<sub>3</sub>) 0-32%), giving 1.5 g (37%) of the sub-title compound.

25

(vii) 3-(2-Ethyl-2*H*-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]nonane  
3-Benzyl-7-(2-ethyl-2*H*-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]-  
nonane (see step (vi) above; 0.5 g, 1.6 mmol) dissolved in ethanol (5 mL of  
95%) was hydrogenated over 5% Pd/C at 1 atm overnight. The catalyst was  
5 filtered over a pad of Celite®, and the residue was evaporated to give 0.2 g  
(56%) of the sub-title compound.

(viii) 4-{3-[7-(2-Ethyl-2*H*-1,2,3,4-tetrazol-5-yl)-3,7-diazabicyclo[3.3.1]-  
non-3-yl]-2-hydroxypropoxy} benzonitrile  
10 4-(2-Oxiranylmethoxy)benzonitrile (see international patent application  
WO 99/31100; 0.18 g, 1 mmol) and 3-(2-ethyl-2*H*-1,2,3,4-tetrazol-5-yl)-  
3,7-diazabicyclo[3.3.1]nonane (see step (vii) above; 0.2 g, 0.9 mmol) were  
mixed in isopropanol:H<sub>2</sub>O (1.1 mL of 10:1) and the mixture was stirred at  
60°C overnight. The solvent was evaporated and the residue was purified  
15 by chromatography (hexane:ethyl acetate (1:1), MeOH (NH<sub>3</sub>) 0-32%),  
giving 0.28 g (77%) of the title compound.

<sup>13</sup>C NMR (CD<sub>3</sub>CN) : δ 14.4, 29.7, 30.0, 31.2, 48.6, 51.2, 51.3, 58.6, 60.8,  
61.0, 66.3, 71.9, 104.3, 116.2, 118.2, 119.9, 134.9, 163.3, 170.1.

#### 20 Example 4

4-{2-Hydroxy-3-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]-  
propoxy} benzonitrile  
4-[3-(3,7-Diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile  
(see Example 1(ii) above; 1 g, 3.3 mmol) 2-bromothiazole (0.54 g,  
25 3.3 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.91 g, 6.5 mmol) were mixed in DMF (15 mL).  
The mixture was stirred overnight at 60°C. The solvent was evaporated,  
toluene was added and then evaporated. The residue was dissolved in ethyl  
acetate and washed with NaOH solution (2 M). The organic layer was  
separated and dried (Na<sub>2</sub>SO<sub>4</sub>). The residue was purified by chromatography  
30 (hexane:ethyl acetate (1:1)). Yield: 0.57 g (45%).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) : δ 29.1, 29.4, 30.5, 52.8, 53.0, 57.9, 60.7, 65.3, 70.4, 104.1, 105.6, 115.3, 119.2, 133.9, 139.7, 162.0, 171.2.

### Example 5

5 *N*<sup>7</sup>-Cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(3,4,5-trimethoxybenzyl)-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide

#### (i) Phenyl *N*<sup>7</sup>-cyano-*N*-(3,4,5-trimethoxybenzyl)carbamimidoate

3,4,5-Trimethoxybenzylamine (1.08 mL, 6.3 mmol) was dissolved in  
10 isopropanol (10 mL). The mixture was cooled to 0°C before diphenyl cyanocarbonimidoate (1.5 g, 6.3 mmol) was added in portions. The reaction mixture was allowed to reach r.t. and then stirred overnight at that temperature. The precipitate that formed was filtered off and was then purified by chromatography DCM:MeOH (gradient 100:0 to 99:1). Yield:  
15 1.37 g (63.5%)

#### (ii) *N*<sup>7</sup>-Cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(3,4,5-trimethoxybenzyl)-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide

4-[3-(3,7-Diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile  
20 (see Example 1(ii) above; 1 g, 3.3 mmol) and phenyl *N*<sup>7</sup>-cyano-*N*-(3,4,5-trimethoxybenzyl)carbamimidoate (see step (i) above; 1.13 g, 3.3 mmol) were mixed with isopropanol (15 mL) and then stirred at reflux for 3 h. The mixture was cooled to r.t. and the product that formed was filtered off. The product was purified by chromatography (DCM:MeOH (gradient 0 to 1%),  
25 giving 1.67 g (92%) of the title compound.

<sup>13</sup>C NMR (CDCl<sub>3</sub>) : δ 29.0, 29.3, 31.2, 47.9, 50.9, 51.3, 56.1, 57.1, 60.0, 60.7, 61.3, 65.6, 70.6, 103.9, 105.1, 115.2, 118.3, 119.1, 133.2, 133.9, 137.4, 153.3, 160.2, 161.9, 176.3, 176.4.

Example 64-{3-Amino-4-[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butoxy}-benzonitrile5 (i) 4-(3-Butenyloxy)benzonitrile

4-Cyanophenol (30 g, 250 mmol) was mixed with K<sub>2</sub>CO<sub>3</sub> (72.5 g, 525 mmol) and stirred for 60 min. 4-Bromo-1-butene (50 g, 370 mmol) was added dropwise, and then the reaction mixture was stirred at 60°C overnight. The solids were filtered off and then the solvents were  
10 evaporated. The residue was dissolved in DCM and washed with 1 N NaOH. The organic layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated, giving 37 g (58 %) of the sub-title compound.

(ii) 4-[2-(2-Oxiranyl)ethoxy]benzonitrile

15 4-(3-Butenyloxy)benzonitrile (see step (i) above; 37 g, 0.21 mol) was mixed with *m*CPBA (61.6 g, 0.25 mol) and DCM (700 mL) and stirred at r.t. for 4 h. The reaction mixture was filtered and 2 mL of DMSO was added to destroy the excess *m*CPBA. The mixture was washed with NaHCO<sub>3</sub>, then separated, dried and evaporated to give 38.7 g (97%) of the sub-title  
20 compound.

(iii) 4-(4-Amino-3-hydroxybutoxy)benzonitrile

4-[2-(2-Oxiranyl)ethoxy]benzonitrile (see step (ii) above; 38.5 g, 204 mmol) was mixed with aqueous NH<sub>3</sub> (1200 mL, conc.) and isopropanol  
25 (450 mL). The mixture was stirred at r.t. for 24 h. The solid (by-product) was filtered off, and the solvents were evaporated, giving 39.1 g (93%) of the sub-title compound.

(iv) tert-Butyl 4-(4-cyanophenoxy)-2-hydroxybutylcarbamate

4-(4-Amino-3-hydroxybutoxy)benzonitrile (see step (iii) above; 34.3 g, 166 mmol) was dissolved in THF:H<sub>2</sub>O (600 mL of 8:2). Di-*tert*-butyl dicarbonate (36.3 g, 166 mmol) was added at 0°C. The mixture was then stirred at r.t. overnight before being evaporated to give 50 g (100%) of the sub-title compound (which was used in the next step without further purification).

10 (v) 1-{[(*tert*-Butoxycarbonyl)amino]methyl}-3-(4-cyanophenoxy)propyl methanesulfonate

*tert*-Butyl 4-(4-cyanophenoxy)-2-hydroxybutylcarbamate (see step (iv) above; 38.1 g, 120 mmol) and 4-(dimethylamino)pyridine (10 mol%) were dissolved in pyridine (200 mL). The mixture was cooled to 0°C. Methanesulfonyl chloride (10.7 mL, 0.136 mol) was then added dropwise at 15 0°C. The mixture was allowed to reach r.t. before the pyridine was evaporated. DCM was added and the solution was washed with 2 M HCl and water before being dried and evaporated. The compound was purified by chromatography on silica, eluting with DCM (5% ethyl acetate), to give 27 g of the sub-title compound.

20

(vi) tert-Butyl 2-[2-(4-cyanophenoxy)ethyl]-1-aziridinecarboxylate

1-{[(*tert*-Butoxycarbonyl)amino]methyl}-3-(4-cyanophenoxy)propyl methanesulfonate (see step (v) above; 25.3 g, 0.066 mol) was mixed with tetrabutylammonium hydrogen sulphate (2.7 g; 7.8 mmol) and DCM (170 mL). The mixture was cooled to 0°C and NaOH (50% (aq)) was added slowly. The mixture was then allowed to reach r.t. before water and DCM were added. The organic layer was separated, washed with water, dried and then evaporated to give the sub-title compound. Yield: 19 g (99%). The product was used in the next step without further purification.

30

(vii) 3-Benzyl-7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]nonane

Butanesulfonyl chloride was added dropwise at 0°C to a mixture of 3-benzyl-3,7-diazabicyclo[3.3.1]nonane (see Example 3(iii) above; 13 g, 60 mmol), K<sub>2</sub>CO<sub>3</sub> (60 mmol) and MeCN (100 mL). The mixture was  
5 allowed to reach r.t. and was then stirred overnight at r.t. The reaction mixture was filtered through a plug of silica, which was then eluted with ethyl acetate to give 15 g (75%) of the sub-title compound.

(viii) 3-(Butylsulfonyl)-3,7-diazabicyclo[3.3.1]nonane

10 3-Benzyl-7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]nonane (see step (vii) above; 12.7 g, 38 mmol) was dissolved in ethanol (150 mL of 95%) and hydrogenated over 5% Pd/C at 1 atm. overnight. TLC analysis showed that no reaction had occurred. The catalyst was filtered off and new catalyst (5% Pd/C) was added, together with H<sub>2</sub>O (10 mL) and acetic acid (2 mL).  
15 The mixture was then hydrogenated at 1 atm. overnight. The catalyst was filtered off, 2 N NaOH was added and the mixture was extracted with toluene. Evaporation of the toluene solution gave 8 g (85%) of the sub-title compound.

20 (ix) *tert*-Butyl 1-{[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-methyl}-3-(4-cyanophenoxy)propylcarbamate

*tert*-Butyl 2-[2-(4-cyanophenoxy)ethyl]-1-aziridinecarboxylate (see step (vi) above; 1 g, 3.5 mmol) was mixed with 3-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]nonane (see step (viii) above; 0.85 g, 3.5 mmol) in  
25 isopropanol (25 mL). The mixture was stirred at 60°C overnight. Evaporation and purification of the residue on silica (DCM, 2% MeOH) gave 1.5 g (81%) of the sub-title compound.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) : δ 1.44 (3H, t), 1.91 (9H, s), 2.08 (2H, m), 2.1 (2H, m), 2.2 (2H, m), 2.4 (2H, s (broad)), 2.6 (2H, m), 2.8 (2H, d), 2.9 (2H, dd), 3.3-

3.6 (6H, m), 4.3 (3H, m), 4.6 (2H, m), 5.7 (1H, s (broad)), 7.4 (2H, dd), 8.0 (2H, dd).

(x) 4-{3-Amino-4-[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-  
5 butoxy} benzonitrile

*tert*-Butyl 1-{[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}-3-(4-cyanophenoxy)propylcarbamate (see step (ix) above; 0.9 g, 1.7 mmol) was dissolved in ethyl acetate (20 mL). Ethyl acetate saturated with HCl (g) (50 mL) was added at 0°C. The resulting mixture was allowed  
10 to reach r.t. for 2 h. The solvent was evaporated and the resulting residue was dissolved in water to give a solution that was freeze-dried. This gave the dihydrochloride salt of the title compound.

ES-MS (M + H)<sup>+</sup> = 435 (m/z)

15 Example 7

The compounds listed below were prepared either in accordance with, or analogously to, methods described herein, or were otherwise prepared according to the following procedure: The appropriate 3- or 7-unsubstituted bispidine (dissolved in CHCl<sub>3</sub>) was reacted with 2 eq. of the appropriate  
20 electrophile (dissolved in CH<sub>3</sub>CN) in the presence of a base (2.5 eq. of K<sub>2</sub>CO<sub>3</sub>). The reaction mixtures were warmed to 50-100°C. When the reaction was ready (as determined by mass spectral analysis), the inorganic salts were filtered off and the reaction mixture was added to an ion exchange solid phase extraction plug (CBA). The plug was washed with  
25 CHCl<sub>3</sub> and the product was finally eluted with CHCl<sub>3</sub>:MeOH:Et<sub>3</sub>N (8:1:1). The products were analysed by HPLC and MS. Compounds with a purity less than 90% were purified by preparative HPLC.

Mass spectra of the compounds, where recorded, are in brackets:

- 4-(3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]-non-3-yl}-2-hydroxypropoxy)benzonitrile;
- 4-(3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-9-(1,2-ethylenedioxy)-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;
- 5 4-{3-[7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-9,9-bis(propylsulfanyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile;
- 3,7-bis(4-nitrophenethyl)-3,7-diazabicyclo[3.3.1]nonane;
- 4-(3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-9,9-tetramethylene-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;
- 10 4-{2-[7-(4-cyanophenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl} benzonitrile;
- N*-{4-[(7-{4-[(methylsulfonyl)amino]benzyl}-3,7-diazabicyclo[3.3.1]non-3-yl)methyl]phenyl}methanesulfonamide;
- 4-cyano-*N*-[2-(7-{2-[(4-cyanobenzoyl)amino]ethyl}-3,7-diazabicyclo-
- 15 [3.3.1]non-3-yl)ethyl]benzamide;
- 4-(2-{7-[2-(4-cyanophenyl)-2-hydroxyethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-1-hydroxyethyl)benzonitrile;
- 4-{3-[(3,7-dibenzyl-3,7-diazabicyclo[3.3.1]non-9-yl)(methyl)amino]-2-hydroxypropoxy} benzonitrile;
- 20 2-[(7-{[6-cyano-4-(methylsulfonyl)-3,4-dihydro-2*H*-1,4-benzoxazin-2-yl]-methyl}-3,7-diazabicyclo[3.3.1]non-3-yl)methyl]-4-(methylsulfonyl)-3,4-dihydro-2*H*-1,4-benzoxazine-6-carbonitrile;
- 4-[2-hydroxy-3-(7-methyl-3,7-diazabicyclo[3.3.1]non-3-yl)propoxy]benzonitrile;
- 25 2-[(7-methyl-3,7-diazabicyclo[3.3.1]non-3-yl)methyl]-4-(methylsulfonyl)-3,4-dihydro-2*H*-1,4-benzoxazine-6-carbonitrile;
- 4-[3-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile;
- 4-{2-hydroxy-3-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-
- 30 propoxy} benzonitrile;

- 4-(((2*R*)-3-{7-[(2*R*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy]benzonitrile;
- 4-(((2*S*)-3-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy]benzonitrile;
- 5 4-[3-(7-butyryl-3,7-diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]-benzonitrile;
- 4-{3-[7-(ethylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile;
- 4-(((2*S*)-3-{7-[(2*R*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy]benzonitrile;
- 10 4-{[(2*S*)-2-hydroxy-3-(7-propionyl-3,7-diazabicyclo[3.3.1]non-3-yl)-propyl]oxy}benzonitrile;
- 4-{[(2*R*)-2-hydroxy-3-(7-propionyl-3,7-diazabicyclo[3.3.1]non-3-yl)-propyl]oxy}benzonitrile;
- 15 2-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-ethyl-2-oxoacetamide;
- 2-{7-[(2*R*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-ethyl-2-oxoacetamide;
- tert*-butyl (1*S*)-2-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-1-[(4-cyanophenoxy)methyl]ethylcarbamate;
- 20 *tert*-butyl (1*S*)-2-(4-cyanophenoxy)-1-({7-[(ethylamino)carbothioyl]-3,7-diazabicyclo[3.3.1]non-3-yl}methyl)ethylcarbamate;
- 7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-ethyl-3,7-diazabicyclo[3.3.1]nonane-3-carbothioamide;
- 25 *tert*-butyl (1*S*)-1-({7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}carbonyl)-2-methylpropylcarbamate;
- 2-(4-cyanophenoxy)-1-{[7-(ethylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}ethyl *tert*-butylcarbamate;
- 4-(((2*S*)-3-{7-[(2*S*)-2-amino-3-methylbutanoyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy]benzonitrile;
- 30

- 4- {[(2*S*)-2-hydroxy-3-(7-{[(2*S*)-5-oxopyrrolidinyl]carbonyl}-3,7-diaza-  
bicyclo[3.3.1]non-3-yl)propyl]oxy} benzonitrile;
- N*-(5-cyano-2-{3-[7-(ethylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-  
hydroxypropoxy} phenyl)-*N'*-ethylurea;
- 5 *N*-(2-{2-amino-3-[7-(ethylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-  
propoxy}-5-cyanophenyl)-*N'*-ethylurea;
- 4-{3-[7-(3,3-dimethylbutanoyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-  
hydroxypropoxy} benzonitrile;
- N*-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]-  
10 non-3-yl}-2-oxoethyl)acetamide;
- tert*-butyl (1*S*)-2-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-  
diazabicyclo[3.3.1]non-3-yl}-1-(4-methoxybenzyl)-2-oxoethylcarbamate;
- 4-[(2*S*)-3-{7-[(2*S*)-2-amino-3-(4-methoxyphenyl)propanoyl]-3,7-diaza-  
bicyclo[3.3.1]non-3-yl}-2-hydroxypropyl]oxy] benzonitrile;
- 15 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-  
yl}-*N*-(2,6-dimethylphenyl)acetamide (*m/z* = 463);
- tert*-butyl 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo-  
[3.3.1]non-3-yl}-2-oxoethylcarbamate (*m/z* = 344);
- 4-{3-[7-(2-aminoethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxy-  
20 propoxy} benzonitrile (*m/z* = 345);
- N*-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]-  
non-3-yl} ethyl)-4-nitrobenzamide (*m/z* = 494);
- 4-amino-*N*-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo-  
[3.3.1]non-3-yl} ethyl)benzamide (*m/z* = 464);
- 25 *N*-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]-  
non-3-yl} ethyl)-4-[(methylsulfonyl)amino]benzamide (*m/z* = 542);
- 4-(acetylamino)-*N*-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-  
diazabicyclo[3.3.1]non-3-yl} ethyl)benzamide (*m/z* = 506);
- 2-[7-(2-{[4-(acetylamino)benzoyl]amino} ethyl)-3,7-diazabicyclo[3.3.1]-  
30 non-3-yl]-1-[(4-cyanophenoxy)methyl]ethyl acetate (*m/z* = 548);

- 4-(3-{7-[(3,5-dimethyl-4-isoxazolyl)carbonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 425$ );
- 4-{2-hydroxy-3-[7-(2-isopropyl-2*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}benzonitrile ( $m/z = 412$ );
- 5 4-(2-hydroxy-3-{7-[(5-methyl-3-isoxazolyl)carbonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 411$ );
- 4-[3-(7-{[3-(*tert*-butyl)-1-methyl-1*H*-pyrazol-5-yl]carbonyl}-3,7-diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile ( $m/z = 466$ );
- 4-(2-hydroxy-3-{7-[(4-methyl-1,2,3-thiadiazol-5-yl)carbonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 428$ );
- 10 4-[3-(7-cyano-3,7-diazabicyclo[3.3.1]non-3-yl)-2-hydroxypropoxy]benzonitrile ( $m/z = 327$ );
- 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-oxoacetamide ( $m/z = 373$ );
- 15 *N*-(3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-*N*-(3,4-dimethoxyphenyl)-4-nitrobenzamide ( $m/z = 644$ );
- 4-{[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]sulfonyl}benzonitrile ( $m/z = 404$ );
- 4-{2-hydroxy-3-[7-(1-phenyl-1*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}benzonitrile ( $m/z = 446$ );
- 20 4-{2-hydroxy-3-[7-(1-methyl-1*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}benzonitrile ( $m/z = 384$ );
- 4-{2-hydroxy-3-[7-(2-methyl-2*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}benzonitrile ( $m/z = 384$ );
- 25 *N*-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-cyclopropyl-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide;
- S*-propyl {7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}sulfonylcarbamothioate ( $m/z = 483$ );
- 4-((*E*)-3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-3-oxo-1-propenyl)benzonitrile ( $m/z = 457$ );
- 30

- ethyl {7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]-non-3-yl} carbothioylcarbamate;
- 4-(2-hydroxy-3-{7-[(2-oxo-1,3-oxazolidin-4-yl)methyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile;
- 5 *tert*-butyl 2-{7-[2-amino-3-(4-cyanophenoxy)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl} ethylcarbamate ( $m/z = 444$ );
- tert*-butyl 3-[(7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl)carbothioylamino]propanoate;
- tert*-butyl 2-{7-[3-(4-cyanoanilino)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl} ethylcarbamate ( $m/z = 428$ );
- 10 4-(3-{7-[(*E*)-3-(3,4-dimethoxyphenyl)-2-propenoyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;
- 4-(3-{7-[(*E*)-3-(4-fluorophenyl)-2-propenoyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 450$ );
- 15 2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-isopropylacetamide ( $m/z = 401$ );
- 4-[(2*S*)-3-[7-(cyclopropylmethyl)-9,9-tetramethylen-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropyl]oxy)benzonitrile ( $m/z = 410$ );
- phenyl *N*-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]nonane-3-carboximidoate ( $m/z = 446$ );
- 20 4-{[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-carbonyl}benzonitrile ( $m/z = 420$ );
- 4-{3-[7-(3-amino-1*H*-1,2,4-triazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile ( $m/z = 384$ );
- 25 *N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(2-hydroxyethyl)-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide ( $m/z = 413$ );
- tert*-butyl 3-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-1-methyl-3-oxopropylcarbamate;

- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-[(5-hydroxy-1,3,3-trimethylcyclohexyl)methyl]-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide ( $m/z = 523$ );
- 4-[2-hydroxy-3-(7-{2-hydroxy-3-[(2-methyl-1-oxo-1,2-dihydro-4-iso-quinolinyloxy]propyl}-3,7-diazabicyclo[3.3.1]non-3-yl)propoxy]benzonitrile;
- 4-(3-{7-[(3,4-dimethoxyphenyl)sulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 502$ );
- 4-{3-[7-(benzylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile ( $m/z = 456$ );
- 4-{3-[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile ( $m/z = 422$ );
- 7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N,N*-dimethyl-3,7-diazabicyclo[3.3.1]nonane-3-sulfonamide;
- 4-{3-[7-(3-aminobutanoyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile;
- 4-{3-[7-(1,3-dimethyl-2,6-dioxo-1,2,3,6-tetrahydro-4-pyrimidinyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile ( $m/z = 440$ );
- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(2-methoxyethyl)-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide ( $m/z = 427$ );
- 4-{2-hydroxy-3-[7-(2,2,2-trifluoroacetyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy} benzonitrile ( $m/z = 398$ );
- 4-{2-hydroxy-3-[7-(3,3,3-trifluoropropanoyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy} benzonitrile;
- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-[(1*S*)-1-(1-naphthyl)ethyl]-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide ( $m/z = 523$ );
- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-isopropyl-*N*-methyl-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide ( $m/z = 425$ );

- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide;
- N*'-cyano-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-(3,4-dimethoxyphenethyl)-*N*-methyl-3,7-diazabicyclo[3.3.1]nonane-3-carboximidamide
- 5 ( $m/z = 547$ );
- 4-(2-hydroxy-3-{7-[(3-methyl-8-quinoliny)lsulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 507$ );
- 4-(2-hydroxy-3-{7-[(1-methyl-1*H*-imidazol-4-yl)sulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 446$ );
- 10 4-(2-hydroxy-3-{7-[(trifluoromethyl)sulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 434$ );
- 4-{2-hydroxy-3-[7-(2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-propoxy}benzonitrile ( $m/z = 372$ );
- 4-{3-[7-(2-furoyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-
- 15 benzonitrile ( $m/z = 396$ );
- 4-[2-hydroxy-3-(7-{[5-(2-pyridinyl)-2-thienyl]sulfonyl}-3,7-diazabicyclo[3.3.1]non-3-yl)propoxy]benzonitrile ( $m/z = 525$ );
- N*-[4-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}acetyl)phenyl]methanesulfonamide ( $m/z = 513$ );
- 20 4-[2-hydroxy-3-(7-{[2-(4-morpholinyl)ethyl]sulfonyl}-3,7-diazabicyclo[3.3.1]non-3-yl)propoxy]benzonitrile;
- 4-{3-[7-(4-amino-6,7-dimethoxy-2-quinazolinyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile;
- 3,7-bis[3-(4-cyanophenoxy)-2-hydroxypropyl]-7-aza-3-azoniabicyclo-
- 25 [3.3.1]nonan-3-olate ( $m/z = 493$ );
- 4-(3-{7-[(3-fluorophenyl)sulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 460$ );
- 4-{3-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile ( $m/z = 466$ );

- 4-[4-(7-butyryl-3,7-diazabicyclo[3.3.1]non-3-yl)-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z = 506$ );
- 4-[4-[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z = 556$ );
- 5 4-{1-(3,4-dimethoxyphenoxy)-4-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile ( $m/z = 534$ );
- 4-[4-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z = 600$ );
- 10 4-[4-(7-butyryl-3,7-diazabicyclo[3.3.1]non-3-yl)butyl]benzonitrile ( $m/z = 354$ );
- 4-{2-[7-(butylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile ( $m/z = 404$ );
- 4-{4-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile ( $m/z = 382$ );
- 15 4-{4-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile ( $m/z = 448$ );
- 4-[2-(7-butyryl-3,7-diazabicyclo[3.3.1]non-3-yl)ethoxy]benzonitrile ( $m/z = 342$ );
- 4-{2-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile ( $m/z = 370$ );
- 20 4-{2-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile ( $m/z = 436$ );
- O*-ethyl 7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]nonane-3-carbothioate;
- 25 benzonitrile, 4-[2-hydroxy-3-[7-[(tetrahydro-3-thienyl)sulfonyl]-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy]-, *S,S*-dioxide;
- 4-({(2*S*)-2-amino-3-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}oxy)benzonitrile;
- 4-{2-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile ( $m/z = 355$ );
- 30

- 4-{1-(3,4-dimethoxyphenoxy)-4-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl} benzonitrile ( $m/z = 519$ );
- 4-({3-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-sulfonyl)benzonitrile ( $m/z = 417$ );
- 5 4-cyano-*N*-{3-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzamide ( $m/z = 396$ );
- 4-{3-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile ( $m/z = 356$ );
- 4-{2-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}-
- 10 benzonitrile ( $m/z = 326$ );
- 4-[4-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z = 490$ );
- 4-({3-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-amino)benzonitrile ( $m/z = 339$ );
- 15 4-{3-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z = 438$ );
- 4-({3-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile ( $m/z = 383$ );
- 4-(4-(4-cyanophenyl)-1-{2-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-oxoethyl}-1*H*-pyrazol-5-yl)benzonitrile ( $m/z = 535$ );
- 20 4-{3-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z = 482$ );
- 4-(2-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-ethoxy)benzonitrile ( $m/z = 374$ );
- 25 *N*-[2-(4-cyanophenoxy)-1-({7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}methyl)ethyl]-*N'*-methylurea ( $m/z = 460$ );
- 4-[(3-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-propyl)amino]benzonitrile ( $m/z = 387$ );

- 4-[4-(4-cyanophenyl)-1-(2-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-oxoethyl)-1*H*-pyrazol-5-yl]benzonitrile ( $m/z$  = 539);
- 4-{3-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile ( $m/z$  = 410);
- 4-{2-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy} benzonitrile ( $m/z$  = 380);
- 4-{1-(3,4-dimethoxyphenoxy)-4-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl} benzonitrile ( $m/z$  = 544);
- 4-({3-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl} amino)benzonitrile ( $m/z$  = 393);
- 4-({3-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-sulfonyl)benzonitrile ( $m/z$  = 442);
- 4-cyano-*N*-{3-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl} benzamide ( $m/z$  = 421);
- 4-{3-[7-(4-fluorobenzyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z$  = 492);
- 4-{2-hydroxy-3-[7-(isopropylsulfonyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy} benzonitrile;
- 4-{3-[7-(1-cyanoethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy} benzonitrile;
- 4-{2-[7-(1-cyanoethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy} benzonitrile;
- 4-[4-[7-(1-cyanoethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z$  = 489);
- 4-{3-[7-(2-cyanopropyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z$  = 451);
- 4-({3-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl} amino)benzonitrile ( $m/z$  = 449);

- 4-({3-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-propyl}sulfonyl)benzonitrile ( $m/z = 498$ );
- 4-(4-(4-cyanophenyl)-1-{2-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-oxoethyl}-1*H*-pyrazol-5-yl)benzonitrile ( $m/z = 601$ );
- 5 4-cyano-*N*-{3-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzamide ( $m/z = 477$ );
- 4-{1-(3,4-dimethoxyphenoxy)-4-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile ( $m/z = 548$ );
- 4-(4-(4-cyanophenyl)-1-{2-oxo-2-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-1*H*-pyrazol-5-yl)benzonitrile ( $m/z = 549$ );
- 10 4-cyano-*N*-{3-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzamide ( $m/z = 425$ );
- 4-(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 478$ );
- 15 4-[(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile ( $m/z = 461$ );
- 4-(3-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile ( $m/z = 436$ );
- 4-(1-(3,4-dimethoxyphenoxy)-4-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}butyl)benzonitrile ( $m/z = 570$ );
- 20 *N*-{2-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-1-[(4-cyanophenoxy)methyl]ethyl}-*N'*-methylurea ( $m/z = 526$ );
- 4-[1-(2-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-oxoethyl)-4-(4-cyanophenyl)-1*H*-pyrazol-5-yl]benzonitrile ( $m/z =$
- 25 605);
- 4-({3-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile ( $m/z = 368$ );
- 4-{2-hydroxy-3-[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z = 467$ );

- N*-(2-(4-cyanophenoxy)-1-{[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}ethyl)-*N'*-methylurea ( $m/z = 412$ );
- 4-({3-[7-(cyclopropylmethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-sulfonyl)benzonitrile ( $m/z = 388$ );
- 5 4-(2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-3,7-diazabicyclo[3.3.1]non-3-yl}acetyl)benzonitrile;
- N*-(2-(4-cyanophenoxy)-1-{[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}ethyl)-*N'*-methylurea ( $m/z = 456$ );
- 4-({3-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}sulfonyl)benzonitrile ( $m/z = 432$ );
- 10 4-cyano-*N*-{3-[7-(3,3-dimethyl-2-oxobutyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzamide ( $m/z = 411$ );
- 4-(2-hydroxy-3-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile ( $m/z = 404$ );
- 15 4-(1-(3,4-dimethoxyphenoxy)-4-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}butyl)benzonitrile ( $m/z = 538$ );
- 4-[(3-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile ( $m/z = 436$ );
- 4-cyano-*N*-(3-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzamide ( $m/z = 415$ );
- 20 4-(2-hydroxy-3-{7-[2-(2-methoxyethoxy)ethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)-*N,N*-dimethylbenzenesulfonamide ( $m/z = 486$ );
- N*-{2-[7-(1-cyanoethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-1-[(4-cyanophenoxy)methyl]ethyl}-*N'*-methylurea;
- 25 4-cyano-*N*-{3-[7-(2-cyanopropyl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzamide ( $m/z = 380$ );
- N*-(2-(4-cyanophenoxy)-1-{[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}ethyl)-*N'*-methylurea ( $m/z = 522$ );
- 4-{3-[7-(3,4-dimethoxyphenethyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z = 548$ );
- 30

- 4-{2-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy} benzonitrile ( $m/z = 384$ );
- 4-{2-hydroxy-3-[7-(4-oxoheptyl)-3,7-diazabicyclo[3.3.1]non-3-yl]-propoxy}-*N,N*-dimethylbenzenesulfonamide ( $m/z = 496$ );
- 5 *N*-[2-(4-cyanophenoxy)-1-({7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl} methyl)ethyl]-*N'*-methylurea ( $m/z = 496$ );
- 4-[(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl} propyl)sulfonyl] benzonitrile ( $m/z = 510$ );
- 10 4-[4-(4-cyanophenyl)-1-(2-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-oxoethyl)-1*H*-pyrazol-5-yl]- benzonitrile ( $m/z = 510$ );
- 4-cyano-*N*-(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl} propyl) benzamide ( $m/z = 489$ );
- 15 4-(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)-*N,N*-dimethylbenzenesulfonamide ( $m/z = 560$ );
- 4-(2-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl} ethoxy)- benzonitrile ( $m/z = 406$ );
- 20 *N*-[2-(4-cyanophenoxy)-1-({7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl} methyl)ethyl]-*N'*-methylurea ( $m/z = 492$ );
- 4-[(3-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-propyl)amino] benzonitrile ( $m/z = 419$ );
- 4-[(3-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-propyl)sulfonyl] benzonitrile ( $m/z = 468$ );
- 25 4-[4-(4-cyanophenyl)-1-(2-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-oxoethyl)-1*H*-pyrazol-5-yl] benzonitrile ( $m/z = 571$ );
- 4-cyano-*N*-(3-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl} propyl) benzamide ( $m/z = 447$ );

- 4-(3-{7-[3-(ethylsulfonyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)-*N,N*-dimethylbenzenesulfonamide ( $m/z = 518$ );
- 4-(2-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile ( $m/z = 440$ );
- 5 4-[4-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile ( $m/z = 604$ );
- 4-[(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile ( $m/z = 453$ );
- 4-[(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile ( $m/z = 502$ );
- 10 4-(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)-*N,N*-dimethylbenzenesulfonamide ( $m/z = 552$ );
- 4-(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;
- 15 *N*-(2-(4-cyanophenoxy)-1-{[7-(1,3-thiazol-2-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}ethyl)-*N'*-methylurea ( $m/z = 441$ );
- 4-{1-(3,4-dimethoxyphenoxy)-4-[7-(2-ethyl-2*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile; and
- 4-({3-[7-(2-ethyl-2*H*-1,2,3,4-tetraazol-5-yl)-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile.
- 20

#### Example 8

Title compounds of the above Examples were tested in Test A above and were found to exhibit  $D_{10}$  values of at least 6.0.

25

#### Example 9

Title compounds of the above Examples were tested in Test B above and were found to exhibit  $pIC_{50}$  values of at least 5.5.

**Abbreviations**


|    |                |   |                                                     |
|----|----------------|---|-----------------------------------------------------|
|    | Ac             | = | acetyl                                              |
|    | API            | = | atmospheric pressure ionisation (in relation to MS) |
| 5  | aq.            | = | aqueous                                             |
|    | br             | = | broad (in relation to NMR)                          |
|    | Bt             | = | benzotriazole                                       |
|    | <i>t</i> -BuOH | = | <i>tert</i> -butanol                                |
|    | CI             | = | chemical ionisation (in relation to MS)             |
| 10 | <i>m</i> CPBA  | = | <i>meta</i> -chloroperoxybenzoic acid               |
|    | d              | = | doublet (in relation to NMR)                        |
|    | DBU            | = | diazabicyclo[5.4.0]undec-7-ene                      |
|    | DCM            | = | dichloromethane                                     |
|    | dd             | = | doublet of doublets (in relation to NMR)            |
| 15 | DMAP           | = | 4-dimethylaminopyridine                             |
|    | DMF            | = | <i>N,N</i> -dimethylformamide                       |
|    | DMSO           | = | dimethylsulfoxide                                   |
|    | EDC            | = | 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide     |
|    | Et             | = | ethyl                                               |
| 20 | EtOAc          | = | ethyl acetate                                       |
|    | eq.            | = | equivalents                                         |
|    | ES             | = | electrospray (in relation to MS)                    |
|    | FAB            | = | fast atom bombardment (in relation to MS)           |
|    | h              | = | hour(s)                                             |
| 25 | HCl            | = | hydrochloric acid                                   |
|    | HEPES          | = | 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid  |
|    | HPLC           | = | high performance liquid chromatography              |
|    | IPA            | = | <i>iso</i> -propyl alcohol (propan-2-ol)            |
|    | m              | = | multiplet (in relation to NMR)                      |
| 30 | Me             | = | methyl                                              |

|    |       |   |                                                           |
|----|-------|---|-----------------------------------------------------------|
|    | MeCN  | = | acetonitrile                                              |
|    | MeOH  | = | methanol                                                  |
|    | min.  | = | minute(s)                                                 |
|    | m.p.  | = | melting point                                             |
| 5  | MS    | = | mass spectroscopy                                         |
|    | NADPH | = | nicotinamide adenine dinucleotide phosphate, reduced form |
|    | OAc   | = | acetate                                                   |
|    | Pd/C  | = | palladium on carbon                                       |
|    | q     | = | quartet (in relation to NMR)                              |
| 10 | rt    | = | room temperature                                          |
|    | s     | = | singlet (in relation to NMR)                              |
|    | t     | = | triplet (in relation to NMR)                              |
|    | TEA   | = | triethylamine                                             |
|    | THF   | = | tetrahydrofuran                                           |
| 15 | tlc   | = | thin layer chromatography                                 |

Prefixes *n*-, *s*-, *i*-, *t*- and *tert*- have their usual meanings: normal, secondary, *iso*, and tertiary.

# Claims

1. A compound of formula I,



wherein

R<sup>1</sup> represents a structural fragment of formula Ia,



- 10 R<sup>4</sup> represents H, halo, C<sub>1-4</sub> alkyl, -D-OR<sup>7</sup>, -D-N(R<sup>8</sup>)R<sup>9</sup>, or R<sup>4</sup>, together with R<sup>5</sup>, represents =O;  
 R<sup>5</sup> represents H, C<sub>1-4</sub> alkyl, or R<sup>5</sup>, together with R<sup>4</sup>, represents =O;  
 D represents a direct bond or C<sub>1-4</sub> alkylene;  
 R<sup>7</sup> represents H, C<sub>1-6</sub> alkyl, -E-aryl, -E-Het<sup>1</sup>, -C(O)R<sup>10a</sup>, -C(O)OR<sup>10b</sup> or  
 15 -C(O)N(R<sup>11a</sup>)R<sup>11b</sup>;  
 R<sup>8</sup> represents H, C<sub>1-6</sub> alkyl, -E-aryl, -E-Het<sup>1</sup>, -C(O)R<sup>10a</sup>, -C(O)OR<sup>10b</sup>,  
 -S(O)<sub>2</sub>R<sup>10c</sup>, -[C(O)]<sub>n</sub>N(R<sup>11a</sup>)R<sup>11b</sup> or -C(NH)NH<sub>2</sub>;  
 R<sup>9</sup> represents H, C<sub>1-6</sub> alkyl, -E-aryl, or -C(O)R<sup>10d</sup>;  
 E represents, at each occurrence when used herein, a direct bond or C<sub>1-4</sub>  
 20 alkylene;  
 R<sup>10a</sup> to R<sup>10d</sup> independently represent, at each occurrence when used herein,  
 C<sub>1-6</sub> alkyl (optionally substituted and/or terminated by one or more

substituents selected from halo, aryl and Het<sup>2</sup>), aryl, Het<sup>3</sup>, or R<sup>10a</sup> and R<sup>10d</sup> independently represent H;

R<sup>11a</sup> and R<sup>11b</sup> independently represent, at each occurrence when used herein, H, C<sub>1-6</sub> alkyl (optionally substituted and/or terminated by one or more substituents selected from halo, aryl and Het<sup>4</sup>), aryl, Het<sup>5</sup>, or R<sup>11a</sup> and R<sup>11b</sup> together represent C<sub>3-7</sub> alkylene, which alkylene group is optionally interrupted by an oxygen atom;

n represents 1 or 2;

10 A represents -G-, -J-N(R<sup>12</sup>)- or -J-O- (in which latter two groups, J is attached to the bispidine nitrogen atom);

B represents -L-, -L-N(R<sup>13</sup>)-, -N(R<sup>13</sup>)-L-, -L-S(O)<sub>p</sub>- or -L-O- (in which latter two groups, L is attached to the carbon atom bearing R<sup>4</sup> and R<sup>5</sup>);

G represents a direct bond or C<sub>1-6</sub> alkylene;

15 J represents C<sub>2-6</sub> alkylene;

L represents a direct bond or C<sub>1-4</sub> alkylene;

p represents 0, 1 or 2;

R<sup>12</sup> and R<sup>13</sup> independently represent H or C<sub>1-4</sub> alkyl;

20 R<sup>6</sup> represents aryl, Het<sup>6</sup> (both of which groups are optionally substituted and/or terminated (as appropriate) by one or more substituents selected from -OH, cyano, halo, nitro, C<sub>1-6</sub> alkyl (optionally terminated by -N(H)C(O)OR<sup>14a</sup>), C<sub>1-6</sub> alkoxy, aryl, Het<sup>7</sup>, -N(R<sup>15a</sup>)R<sup>15b</sup>, -C(O)R<sup>15c</sup>, -C(O)OR<sup>15d</sup>, -C(O)N(R<sup>15e</sup>)R<sup>15f</sup>, -N(R<sup>15g</sup>)C(O)R<sup>15h</sup>, -N(R<sup>15i</sup>)C(O)N(R<sup>15j</sup>)R<sup>15k</sup>,  
25 -N(R<sup>15m</sup>)S(O)<sub>2</sub>R<sup>14b</sup>, -S(O)<sub>q</sub>R<sup>14c</sup>, -OS(O)<sub>2</sub>R<sup>14d</sup> and -S(O)<sub>2</sub>N(R<sup>15n</sup>)R<sup>15p</sup>) or, when R<sup>4</sup> and R<sup>5</sup> together represent =O, R<sup>6</sup> may represent C<sub>1-6</sub> alkyl;

q represents 0, 1 or 2;

R<sup>2</sup> represents -CN, Het<sup>8</sup>, -C(O)R<sup>16</sup>, -C(S)OR<sup>17</sup>, -C(S)N(R<sup>18</sup>)R<sup>19</sup>,  
30 -[C(O)]<sub>2</sub>N(R<sup>20a</sup>)R<sup>20b</sup>, -[C(O)]<sub>2</sub>OR<sup>21</sup>, -S(O)<sub>2</sub>R<sup>22</sup>, -S(O)<sub>2</sub>N(R<sup>23</sup>)R<sup>24</sup>,

$-C(=N-CN)N(R^{25})R^{26}$ ,  $-C(=N-CN)OR^{27}$  or  $C_{1-12}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from  $-C(O)R^{28}$ ,  $-C(O)N(R^{29a})R^{29b}$ ,  $-N(R^{30})R^{31}$ ,  $-OR^{32}$ ,  $-S(O)_rR^{33}$ , halo,  $-CN$ , nitro, aryl and  $Het^9$ );

5

$R^{16}$  represents H, aryl,  $Het^{10}$  or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo,  $-OH$ ,  $-CN$ ,  $-N(R^{34})R^{35}$ , aryl and  $Het^{11}$ );

$R^{34}$  represents, H,  $C_{1-6}$  alkyl, aryl,  $Het^{12}$ ,  $-C(O)R^{36a}$  or  $-C(O)OR^{36b}$ ;

10

$R^{18}$  represents H, aryl,  $Het^{13}$ ,  $-C(O)R^{36a}$ ,  $-C(O)OR^{36b}$  or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo,  $-OH$ ,  $-CN$ ,  $-C(O)R^{36a}$  and  $-C(O)OR^{36b}$ );

15  $R^{22}$  represents  $Het^{14}$ , aryl, or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo,  $-OH$ ,  $-CN$ ,  $Het^{15}$  and aryl);

$R^{23}$  represents H,  $C_{1-6}$  alkyl, aryl,  $Het^{16}$ ,  $-C(O)R^{36a}$ ,  $-C(O)OR^{36b}$  or  
20  $-C(O)SR^{36b}$ ;

$R^{25}$  represents H or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo,  $-OH$ ,  $-CN$ ,  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or  
25 terminated by one or more substituents selected from  $C_{1-4}$  alkyl and  $-OH$ ),  $C_{1-6}$  alkoxy and aryl);

$R^{27}$  represents  $C_{1-6}$  alkyl or aryl;

30  $R^{28}$  represents H,  $C_{1-6}$  alkyl, aryl or  $Het^{17}$ ;

$R^{29a}$  and  $R^{29b}$  independently represent H,  $C_{1-6}$  alkyl, aryl or Het<sup>18</sup>;

$R^{30}$  represents H,  $C_{1-6}$  alkyl, aryl, Het<sup>19</sup>,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ ;

5  $R^{31}$  represents H,  $C_{1-6}$  alkyl, aryl or Het<sup>20</sup>;

$R^{32}$  represents H,  $C_{1-6}$  alkyl, aryl, Het<sup>21</sup>,  $-C(O)R^{37a}$ ,  $-C(O)OR^{37b}$  or  $-C(O)N(R^{37c})R^{37d}$ ;

10  $R^{33}$  represents  $C_{1-6}$  alkyl, aryl or Het<sup>22</sup>;

r represents 0, 1 or 2;

$R^{36a}$  and  $R^{36b}$  independently represent, at each occurrence when used herein,  $C_{1-6}$  alkyl, or  $R^{36a}$  represents H;

15  $R^{37a}$  to  $R^{37d}$  independently represent, at each occurrence when used herein,  $C_{1-6}$  alkyl, aryl or Het<sup>23</sup>, or  $R^{37a}$ ,  $R^{37c}$  and  $R^{37d}$  independently represent H;

Het<sup>1</sup> to Het<sup>23</sup> independently represent, at each occurrence when used herein, five- to twelve-membered heterocyclic groups containing one or more  
20 heteroatoms selected from oxygen, nitrogen and/or sulfur;

$R^{3a}$  and  $R^{3b}$  independently represent H,  $C_{1-4}$  alkyl,  $-OR^{38a}$ ,  $-SR^{38b}$ ,  $-N(R^{39})R^{38c}$ , or  $R^{3a}$  and  $R^{3b}$  together represent  $C_{3-5}$  alkylene,  $-O-Z-O-$ ,  $-O-Z-S-$  or  $-S-Z-S-$ ;

25  $R^{39}$  represents H,  $C_{1-6}$  alkyl or a structural fragment of formula Ia as defined above;

Z represents  $C_{2-3}$  alkylene optionally substituted by one or more  $C_{1-4}$  alkyl groups;

30  $R^{41}$  to  $R^{46}$  independently represent H or  $C_{1-3}$  alkyl;

$R^{14a}$  to  $R^{14d}$ ,  $R^{17}$  and  $R^{21}$  independently represent  $C_{1-6}$  alkyl;  
 $R^{15a}$  to  $R^{15p}$ ,  $R^{19}$ ,  $R^{20a}$ ,  $R^{20b}$ ,  $R^{24}$ ,  $R^{26}$ ,  $R^{35}$  and  $R^{38a}$  to  $R^{38c}$  independently represent H or  $C_{1-6}$  alkyl;

- 5 wherein each aryl and Het ( $Het^1$  to  $Het^{23}$ ) group, unless otherwise specified, is optionally substituted;

or a pharmaceutically acceptable derivative thereof;

- 10 provided that:

- (a) when  $R^1$  represents a structural fragment of formula Ia in which:  
 $R^4$  and  $R^5$  together represent  $=O$ ;

A represents a direct bond;

- 15 then B does not represent a direct bond,  $-N(R^{13})-L-$  (in which group  $-N(R^{13})-$  is attached to the carbon atom bearing  $R^4$  and  $R^5$ ),  $-N(R^{13})-$ ,  $-S(O)_p-$  or  $-O-$ ;

- (b) when  $R^5$  represents H or  $C_{1-4}$  alkyl; and  
A represents  $-J-N(R^{12})-$  or  $-J-O-$ ;

then B does not represent  $-N(R^{13})-L-$ ,  $-N(R^{13})-$ ,  $-S(O)_p-$  or  $-O-$ ;

- 20 (c) when  $R^4$  represents  $-D-OR^7$ ,  $-D-N(R^8)R^9$  in which D represents a direct bond, then:

(i) A does not represent  $-J-N(R^{12})-$  or  $-J-O-$ ; and

(ii) B does not represent  $-N(R^{13})-L-$ ,  $-N(R^{13})-$ ,  $-S(O)_p-$  or  $-O-$ ;

- (d) when  $R^{3a}$  and  $R^{3b}$  and both represent H;

- 25 and  $R^1$  represents unsubstituted benzyl;

then  $R^2$  does not represent unsubstituted benzyl or optionally substituted benzoyl; and

- (e) the compound is not:

- (i)  $N^1$ -phenyl-3-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-  
30 propanamide;

- (ii) 3-benzyl-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-6,8-dimethyl-3,7-diazabicyclo[3.3.1]nonane;
- (iii) 3-benzyl-7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-6-methyl-3,7-diazabicyclo[3.3.1]nonane;
- 5 (iv) *N*-{2-(7-benzyl-3,7-diazabicyclo[3.3.1]non-3-yl)-1-[(4-cyanophenoxy)methyl]ethyl}methanesulfonamide;
- (v) 3-benzyl-7-[3-(2-propyl-1,3-dioxolan-2-yl)propyl]-3,7-diazabicyclo[3.3.1]nonane; or
- (vi) 7-benzyl-3,7-diazabicyclo[3.3.1]nonane-3-ethanol.
- 10
2. A compound as claimed in Claim 1, wherein  $R^4$  represents H,  $C_{1-2}$  alkyl,  $-OR^7$  or  $N(H)R^8$ , or  $R^4$ , together with  $R^5$ , represents  $=O$ .
3. A compound as claimed in Claim 1 or Claim 2, wherein  $R^5$  represents
- 15 H, or  $R^5$ , together with  $R^4$ , represents  $=O$ .
4. A compound as claimed in any one of Claims 1 to 3, wherein  $R^7$  represents H,  $C_{1-4}$  alkyl, optionally substituted phenyl,  $-C(O)R^{10a}$ , or  $-C(O)N(R^{11a})R^{11b}$ .
- 20
5. A compound as claimed in any one of Claims 1 to 3, wherein  $R^8$  represents H,  $C_{1-4}$  alkyl,  $-C(O)R^{10a}$ ,  $-C(O)OR^{10b}$  or  $-C(O)N(R^{11a})R^{11b}$ .
6. A compound as claimed in any one of Claims 1 to 5, wherein  $R^{10a}$  and
- 25  $R^{10b}$  independently represent, at each occurrence when used herein,  $C_{1-5}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo and phenyl), optionally substituted phenyl, or  $R^{10a}$  represents H.

7. A compound as claimed in Claim 4 or Claim 5, wherein  $R^{11a}$  and  $R^{11b}$  independently represent, at each occurrence when used herein, H or  $C_{1-5}$  alkyl (optionally substituted and/or terminated by one or more substituents selected from halo and phenyl).
8. A compound as claimed in any one of Claims 1 to 7, wherein A represents -G- or -J-N( $R^{12}$ )-.
9. A compound as claimed in Claim 8, wherein G represents a direct bond or  $C_{1-4}$  alkylene.
10. A compound as claimed in any one of Claims 1 to 8, wherein J represents  $C_{2-4}$  alkylene.
11. A compound as claimed in any one of Claims 1 to 10, wherein B represents a direct bond,  $C_{1-4}$  alkylene, -L-N(H)-, -L-S(O)<sub>2</sub>- or -L-O- (in which latter three groups, L is attached to the carbon atom bearing  $R^4$  and  $R^5$ ).
12. A compound as claimed in any one of Claims 1 to 11, wherein L represents  $C_{1-4}$  alkylene.
13. A compound as claimed in any one of Claims 1 to 12, wherein  $R^6$  represents phenyl, Het<sup>6</sup> (both of which groups are optionally substituted by one or more substituents selected from cyano, halo, nitro,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, optionally substituted phenyl, -N(H) $R^{15b}$ , -C(O) $R^{15c}$ , -C(O)N(H) $R^{15f}$ , -N(H)C(O) $R^{15h}$ , -N(H)C(O)N(H) $R^{15k}$ , -N(H)S(O)<sub>2</sub> $R^{14b}$ , -S(O)<sub>2</sub> $R^{14c}$  and -S(O)<sub>2</sub>N( $R^{15n}$ ) $R^{15p}$ ), or, when  $R^4$  and  $R^5$  together represent =O,  $R^6$  may represent  $C_{1-5}$  alkyl.

14. A compound as claimed in any one of Claims 1 to 13, wherein  $R^2$  represents  $-\text{CN}$ ,  $\text{Het}^8$ ,  $-\text{C}(\text{O})\text{R}^{16}$ ,  $-\text{C}(\text{S})\text{OR}^{17}$ ,  $-\text{C}(\text{S})\text{N}(\text{H})\text{R}^{18}$ ,  $-\text{[C}(\text{O})\text{]}_2\text{N}(\text{H})\text{R}^{20\text{b}}$ ,  $-\text{[C}(\text{O})\text{]}_2\text{OR}^{21}$ ,  $-\text{S}(\text{O})_2\text{R}^{22}$ ,  $-\text{S}(\text{O})_2\text{N}(\text{R}^{23})\text{R}^{24}$ ,  $-\text{C}(=\text{N}-\text{CN})\text{N}(\text{R}^{25})\text{R}^{26}$ ,  $-\text{C}(=\text{N}-\text{CN})\text{OR}^{27}$  or  $\text{C}_{1-6}$  alkyl (which alkyl group is  
 5 optionally substituted and/or terminated by one or more substituents selected from  $-\text{C}(\text{O})\text{R}^{28}$ ,  $-\text{C}(\text{O})\text{N}(\text{H})\text{R}^{29\text{b}}$ ,  $-\text{N}(\text{R}^{30})\text{R}^{31}$ ,  $-\text{OR}^{32}$ ,  $-\text{S}(\text{O})_2\text{R}^{33}$ , halo,  $-\text{CN}$ , optionally substituted phenyl and  $\text{Het}^9$ ).

15. A compound as claimed in Claim 14, wherein  $R^{16}$  represents  
 10 optionally substituted phenyl,  $\text{Het}^{10}$  or  $\text{C}_{1-6}$  alkyl (which alkyl group is optionally unsaturated and/or optionally substituted and/or terminated by one or more substituents selected from halo,  $-\text{CN}$ ,  $-\text{N}(\text{H})\text{R}^{34}$  and optionally substituted phenyl).

15 16. A compound as claimed in Claim 15, wherein  $R^{34}$  represents, H,  $\text{C}_{1-4}$  alkyl,  $-\text{C}(\text{O})\text{R}^{36\text{a}}$  or  $-\text{C}(\text{O})\text{OR}^{36\text{b}}$ .

17. A compound as claimed in Claim 14, wherein  $R^{18}$  represents H,  $-\text{C}(\text{O})\text{OR}^{36\text{b}}$  or  $\text{C}_{1-6}$  alkyl (which alkyl group is optionally substituted and/or  
 20 terminated by one or more substituents selected from halo and  $-\text{C}(\text{O})\text{OR}^{36\text{b}}$ ).

18. A compound as claimed in Claim 14, wherein  $R^{22}$  represents  $\text{Het}^{14}$ , optionally substituted phenyl or  $\text{C}_{1-4}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from  
 25 halo,  $\text{Het}^{15}$  and optionally substituted phenyl).

19. A compound as claimed in Claim 14, wherein  $R^{23}$  represents H,  $\text{C}_{1-4}$  alkyl,  $-\text{C}(\text{O})\text{OR}^{36\text{b}}$  or  $-\text{C}(\text{O})\text{SR}^{36\text{b}}$ .

20. A compound as claimed in Claim 14, wherein  $R^{25}$  represents H or  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents selected from halo, -OH,  $C_{1-6}$  alkyl (which alkyl group is optionally substituted and/or terminated by one or more substituents  
5 selected from  $C_{1-4}$  alkyl and -OH),  $C_{1-4}$  alkoxy, naphthyl and optionally substituted phenyl).
21. A compound as claimed in Claim 14, wherein  $R^{27}$  represents optionally substituted phenyl.
- 10 22. A compound as claimed in Claim 14, wherein  $R^{28}$  represents  $C_{1-5}$  alkyl, optionally substituted phenyl or Het<sup>17</sup>.
- 15 23. A compound as claimed in any one of Claims 1 to 14 or 22, wherein  $R^{29b}$  represents H,  $C_{1-4}$  alkyl or optionally substituted phenyl.
24. A compound as claimed in any one of Claims 14, 22 and 23, wherein  $R^{30}$  represents H, optionally substituted phenyl,  $-C(O)R^{37a}$  or  $-C(O)OR^{37b}$ .
- 20 25. A compound as claimed in any one of Claims 14 or 22 to 24, wherein  $R^{31}$  represents H,  $C_{1-2}$  alkyl or optionally substituted phenyl.
26. A compound as claimed in any one of Claims 14 or 22 to 25, wherein  $R^{32}$  represents H,  $C_{1-4}$  alkyl (which alkyl group is optionally interrupted by  
25 oxygen), optionally substituted phenyl or Het<sup>21</sup>.
27. A compound as claimed in any one of Claims 14 or 22 to 26, wherein  $R^{33}$  represents  $C_{1-6}$  alkyl or optionally substituted phenyl.

28. A compound as claimed in any one of Claims 1 to 14 or 22 to 27, wherein  $R^{37a}$  and  $R^{37b}$  independently represent, at each occurrence when used herein,  $C_{1-5}$  alkyl, optionally substituted phenyl, or  $R^{37a}$  represents H.

5 29. A compound as claimed in any one of Claims 1 to 28, wherein  $R^{3a}$  and  $R^{3b}$  independently represent H,  $C_{1-2}$  alkyl,  $-SR^{38b}$ ,  $-N(R^{39})R^{38c}$ , or  $R^{3a}$  and  $R^{3b}$  together represent  $C_{3-4}$  alkylene or  $-O-Z-O-$ .

30. A compound as claimed in Claim 29, wherein  $R^{39}$  represents H,  $C_{1-2}$  alkyl or a structural fragment of formula Ia.

31. A compound as claimed in any one of Claims 1 to 30, wherein Z represents  $C_{2-3}$  alkylene.

15 32. A compound as claimed in any one of Claims 1 to 31, wherein  $R^{41}$  to  $R^{46}$  independently represent H or  $C_{1-2}$  alkyl.


33. A compound as claimed in Claim 13 or Claim 14, wherein  $R^{14b}$ ,  $R^{14c}$ ,  $R^{17}$  and  $R^{21}$  independently represent  $C_{1-4}$  alkyl.

20 34. A compound as claimed in any one of Claims 13, 14 and 29, wherein  $R^{15b}$  to  $R^{15p}$ ,  $R^{20b}$ ,  $R^{24}$ ,  $R^{26}$ ,  $R^{38b}$  and  $R^{38c}$  independently represent H or  $C_{1-5}$  alkyl.

25 35. A compound as claimed in any one of Claims 1 to 34, wherein optional substituents on phenyl groups are one or more substituents selected from cyano, halo, nitro,  $C_{1-2}$  alkyl,  $C_{1-2}$  alkoxy,  $Het^1$ ,  $-NH_2$ ,  $-C(O)R^{15c}$ ,  $-C(O)N(H)R^{15f}$ ,  $-N(H)C(O)R^{15h}$ ,  $-N(H)C(O)N(H)R^{15k}$ ,  $-N(H)S(O)_2R^{14b}$  and  $-S(O)_2N(R^{15n})R^{15p}$ .


36. A pharmaceutical formulation including a compound as defined in any one of Claims 1 to 35 in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.
- 5 37. A pharmaceutical formulation for use in the prophylaxis or the treatment of an arrhythmia, comprising a compound as defined in any one of Claims 1 to 35.
38. A compound as defined in any one of Claims 1 to 35, but without  
10 proviso (e), for use as a pharmaceutical.
39. A compound as defined in any one of Claims 1 to 35, but without proviso (e), for use in the prophylaxis or the treatment of an arrhythmia.
- 15 40. The use of a compound as defined in any of one Claims 1 to 35, but without proviso (e), as active ingredient for the manufacture of a medicament for use in the prophylaxis or the treatment of an arrhythmia.
41. The use as claimed in Claim 40, wherein the arrhythmia is an atrial or  
20 a ventricular arrhythmia.
42. A method of prophylaxis or treatment of an arrhythmia which method comprises administration of a therapeutically effective amount of a compound as defined in any one of Claims 1 to 35, but without proviso (e),  
25 to a person suffering from, or susceptible to, such a condition.
43. A process for the preparation of a compound of formula I as defined in Claim 1 which comprises:
- (a) reaction of a corresponding compound of formula II,

94



II

wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1, with a compound of formula III,




III

5 wherein  $L^1$  represents a leaving group and  $R^4$ ,  $R^5$ ,  $R^6$ , A and B are as defined in Claim 1;

(b) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which A represents  $C_2$  alkylene and  $R^4$  and  $R^5$  together represent  $=O$ , reaction of a corresponding compound of formula II, as

10 defined above, with a compound of formula IV,




IV

wherein  $R^6$  and B are as defined in Claim 1;

(c) for compounds of formula I in which  $R^{3a}$  or  $R^{3b}$  represents  $-N(R^{39})R^{38c}$  and  $R^{39}$  represents a structural fragment of formula Ia, reaction of a  
15 corresponding compound of formula I in which  $R^{3a}$  or  $R^{3b}$  (as appropriate) represents  $-N(H)R^{38c}$ , wherein  $R^{38c}$  is as defined in Claim 1, with a compound of formula III as defined above;


(d) for compounds of formula I in which  $R^1$  represents a fragment of formula Ia in which A represents  $CH_2$  and  $R^4$  represents  $-OH$  or  $-N(H)R^8$ ,

20 reaction of a corresponding compound of formula II, as defined above, with a compound of formula V,



wherein X represents O or N(R<sup>8</sup>) and R<sup>5</sup>, R<sup>6</sup>, R<sup>8</sup> and B are as defined in Claim 1;

- (e) for compounds of formula I in which R<sup>3a</sup> or R<sup>3b</sup> represents -N(R<sup>39</sup>)R<sup>38c</sup> and R<sup>39</sup> represents a structural fragment of formula Ia in which A represents CH<sub>2</sub> and R<sup>4</sup> represents -OH or -N(H)R<sup>8</sup>, reaction of a corresponding compound of formula I in which R<sup>3a</sup> or R<sup>3b</sup> (as appropriate) represents -N(H)R<sup>38c</sup>, wherein R<sup>38c</sup> is as defined in Claim 1, with a compound of formula V as defined above;
- (f) for compounds of formula I in which A represents C<sub>1-6</sub> alkylene, B represents C<sub>1-4</sub> alkylene and R<sup>4</sup> and R<sup>5</sup> both represent H, reduction of a corresponding compound of formula I in which R<sup>4</sup> and R<sup>5</sup> together represent =O;
- (g) for compounds of formula I in which R<sup>4</sup> and R<sup>5</sup> both represent H and (1) A represents a single bond or -J-N(R<sup>12</sup>) and B represents C<sub>1-4</sub> alkylene, or (2) A represents C<sub>1-6</sub> alkylene and B represents N(R<sup>13</sup>) or -N(R<sup>13</sup>)-L-, reduction of a corresponding compound of formula I in which R<sup>4</sup> and R<sup>5</sup> together represent =O;
- (h) for compounds of formula I in which A represents C<sub>1-6</sub> alkylene, B represents a direct bond, C<sub>1-4</sub> alkylene, -L-N(R<sup>13</sup>)-, -L-S(O)<sub>p</sub>- or -L-O- (in which latter three groups L represents C<sub>1-4</sub> alkylene), R<sup>4</sup> represents OH and R<sup>5</sup> represents H, reduction of a corresponding compound of formula I in which R<sup>4</sup> and R<sup>5</sup> together represent =O;
- (i) for compounds of formula I in which R<sup>3a</sup> and R<sup>3b</sup> both represent H, reduction of a corresponding compound of formula VI,




wherein  $R^1$ ,  $R^2$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1, and in which the bridgehead C=O group may be activated;

(j) for compounds of formula I in which one of  $R^{3a}$  and  $R^{3b}$  represents H, and the other represents -OH, reduction of a corresponding compound of formula VI, as defined above;

(k) for compounds of formula I in which  $R^{3a}$  and  $R^{3b}$  both represent  $-OR^{38a}$  or  $-SR^{38b}$ , or in which  $R^{3a}$  and  $R^{3b}$  together represent -O-Z-O-, -O-Z-S- or -S-Z-S-, reaction of a corresponding compound of formula VI, as defined above, with a compound of formula  $HOR^{38a}$ ,  $HSR^{38b}$ , HO-Z-OH, HO-Z-SH or HS-Z-SH (as appropriate), wherein  $R^{38a}$ ,  $R^{38b}$  and Z are as defined in Claim 1;

(l) for compounds of formula I in which one of  $R^{3a}$  and  $R^{3b}$  represents  $-NH_2$  and the other represents H, reduction of a compound of formula VII,



wherein  $R^1$ ,  $R^2$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1;


- (m) for compounds of formula I in which one or both of  $R^{3a}$  and  $R^{3b}$  represent  $-N(R^{39})R^{38c}$  in which one or both of  $R^{39}$  and  $R^{38c}$  represents  $C_{1-6}$  alkyl, alkylation of a corresponding compound of formula I in which  $R^{3a}$  and/or  $R^{3b}$  represent  $-N(R^{39})R^{38c}$  (as appropriate) in which  $R^{39}$  and/or  $R^{38c}$
- 5 (as appropriate) represent H, using a compound of formula VIII,



VIII

wherein  $R^a$  represents  $C_{1-6}$  alkyl and  $L^1$  is as defined above;

- (n) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which B represents  $-L-O-$ , reaction of a compound of
- 10 formula IX,



wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^4$ ,  $R^5$ ,  $R^{41}$  to  $R^{46}$ , A and L are as defined in Claim 1, with a compound of formula X,




X

- 15 in which  $R^6$  is as defined in Claim 1;

(o) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which A represents  $C_{1-6}$  alkylene and B represents  $-N(R^{13})-L-$  (wherein the group  $-N(R^{13})-$  is attached to the carbon atom bearing  $R^4$  and  $R^5$ ), reaction of a compound of formula XI,

98




wherein  $A^a$  represents  $C_{1-6}$  alkylene and  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^4$ ,  $R^5$ ,  $R^{13}$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1, with a compound of formula XII,



5 wherein  $L^2$  represents a leaving group and  $R^6$  and  $L$  are as defined in Claim 1;

(p) for compounds of formula I in which  $R^1$  represents a structural fragment of formula Ia in which  $R^4$  represents  $-D-NH_2$ , reduction of a corresponding compound of formula XIII,



10 wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^5$ ,  $R^6$ ,  $R^{41}$  to  $R^{46}$ ,  $A$ ,  $B$  and  $D$  are as defined in Claim 1;

(q) for compounds of formula I in which  $R^4$  represents  $-D-N(R^9)C(O)NH(R^{11b})$ , reaction of a corresponding compound of formula I


15 in which  $R^4$  represents  $-D-N(R^9)H$  with a compound of formula XIV,



wherein  $R^{11b}$  is as defined in Claim 1;

(r) for compounds of formula I in which  $R^4$  represents  $-D-N(H)[C(O)]_2NH_2$ , reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-NH_2$  with oxalic acid diamide;

- 5 (s) for compounds of formula I in which  $R^4$  represents  $-D-N(R^8)R^9$ , wherein  $R^8$  and  $R^9$  are as defined in Claim 1, provided that  $R^8$  does not represent H, reaction of a corresponding compound of formula I, in which  $R^4$  represents  $-D-N(H)R^9$  with a compound of formula XV,




XV

- 10 wherein  $R^{8a}$  represents  $R^8$  as defined in Claim 1 except that it does not represent H, and  $L^3$  represents a leaving group;

(t) for compounds of formula I in which  $R^4$  represents  $-D-OR^7$  in which  $R^7$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>, reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-OH$  with a compound of


- 15 formula XVI,



XVI

wherein  $R^{7a}$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>, wherein Het<sup>1</sup> is as defined in Claim 1;

- (u) for compounds of formula I in which  $R^1$  represents a structural fragment  
20 of formula Ia in which  $R^4$  represents  $-D-OR^7$  (in which  $R^7$  represents  $C_{1-6}$  alkyl,  $-E$ -aryl or  $-E$ -Het<sup>1</sup>), reaction of a corresponding compound of formula XVII,



XVII

wherein  $R^2$ ,  $R^{3a}$ ,  $R^{3b}$ ,  $R^5$ ,  $R^6$ ,  $R^{41}$  to  $R^{46}$ , A, B and D are as defined in Claim 1 and  $L^2$  is as defined above, with a compound of formula XVI as defined above;


- (v) for compounds of formula I in which  $R^4$  represents  $-D-OR^7$ , wherein  $R^7$  is as defined in Claim 1, provided that it does not represent H, reaction of a corresponding compound of formula I in which  $R^4$  represents  $-D-OH$  with a compound of formula XVIII,



- wherein  $R^{7b}$  represents  $R^7$  as defined in Claim 1, except that it does not represent H, and  $L^4$  represents a leaving group;

(w) for compounds of formula I in which  $R^4$  represents halo, substitution of a corresponding compound of formula I in which  $R^4$  represents  $-OH$ , using an appropriate halogenating agent;

(x) reaction of a corresponding compound of formula XIX,




wherein  $R^1$ ,  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1, with a compound of formula XX,



wherein  $L^5$  represents a leaving group and  $R^2$  is as defined in Claim 1;

- (y) for compounds of formula I in which  $R^2$  represent  $C_{1-12}$  alkyl, which alkyl group is substituted at the C-2 carbon (relative to the bispidine nitrogen) with OH or  $N(H)R^{30}$ , and is otherwise optionally substituted with one or more further substituents as specified in Claim 1 for  $R^2$ , reaction of a

compound of formula XIX as defined above with a compound of formula  
XXA



- wherein  $X_a$  represents O or  $N(R^{30})$  and  $R^{2a}$  represents  $C_{1-10}$  alkyl, optionally  
 5 substituted with one or more substituents as specified in Claim 1 for  $R^2$ ;  
 (z) for compounds of formula I in which  $R^2$  represents tetrazol-5-yl,  
 reaction of a corresponding compound of formula I in which  $R^2$  represents  
 -CN with a source of the azide ion;  
 (aa) for compounds of formula I which are bispidine-nitrogen *N*-oxide  
 10 derivatives, oxidation of the corresponding bispidine nitrogen of a  
 corresponding compound of formula I, in the presence of a suitable  
 oxidising agent;  
 (ab) for compounds of formula I which are  $C_{1-4}$  alkyl quaternary ammonium  
 salt derivatives, in which the alkyl group is attached to a bispidine nitrogen,  
 15 reaction, at the bispidine nitrogen, of a corresponding compound of formula  
 I with a compound of formula XXI,



- wherein  $R^b$  represents  $C_{1-4}$  alkyl and  $L^2$  is as defined above;  
 (ac) conversion of one substituent on  $R^6$  to another;  
 20 (ad) conversion of one  $R^2$  group to another; or  
 (ae) deprotection of a protected derivative of a compound of formula I as  
 defined in Claim 1.

44. A compound of formula II, as defined in Claim 43, or a protected  
 25 derivative thereof.

45. A compound of formula VI, as defined in Claim 43, or a protected  
 derivative thereof.

46. A compound of formula VII, as defined in Claim 43, or a protected derivative thereof.

47. A compound of formula IX, as defined in Claim 43, or a protected derivative thereof.


48. A compound of formula XI, as defined in Claim 43, or a protected derivative thereof.

49. A compound of formula XIII, as defined in Claim 43, or a protected derivative thereof.

50. A compound of formula XVII, as defined in Claim 43, or a protected derivative thereof.

51. A compound of formula XIX, as defined in Claim 43 (provided that at least one of  $R^{3a}$  and  $R^{3b}$  represents  $-N(R^{39})R^{38c}$ , wherein  $R^{39}$  represents a structural fragment of formula Ia, as defined in Claim 1), or a protected derivative thereof.

52. A compound of formula XXII,




XXII


wherein  $R^{3a}$ ,  $R^{3b}$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1 (provided that at least one of  $R^{3a}$  and  $R^{3b}$  represents  $-N(R^{39})R^{38c}$ , wherein  $R^{39}$  represents a

structural fragment of formula Ia, as defined in Claim 1), or a protected derivative thereof.

53. A compound of formula XXIII,



wherein  $R^2$  and  $R^{41}$  to  $R^{46}$  are as defined in Claim 1, or a protected derivative thereof.



(I)