

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number
WO 2006/068796 A2

(51) International Patent Classification:
A61K 31/397 (2006.01)

(21) International Application Number:
PCT/US2005/043361

(22) International Filing Date:
28 November 2005 (28.11.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/632,490 2 December 2004 (02.12.2004) US

(71) Applicant (for all designated States except US): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BARNETT, Stanley, F.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **BOGUSKY, Michael, J.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **LEISTER, William, H.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **LINDSLEY, Craig, W.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF AKT ACTIVITY

(57) Abstract: The instant invention provides for canthine analogs that inhibit Akt activity. In particular, the compounds disclosed selectively inhibit one or two of the Akt isoforms. The invention also provides for compositions comprising such inhibitory compounds and methods of inhibiting Akt activity by administering the compound to a patient in need of treatment of cancer.

WO 2006/068796 A2

TITLE OF THE INVENTION

INHIBITORS OF AKT ACTIVITY

BACKGROUND OF THE INVENTION

5 The present invention relates to canthine analogs which are inhibitors of the activity of one or more of the isoforms of the serine/threonine kinase, Akt (also known as PKB; hereinafter referred to as "Akt"). The present invention also relates to pharmaceutical compositions comprising such compounds and methods of using the instant compounds in the treatment of cancer.

10 Apoptosis (programmed cell death) plays essential roles in embryonic development and pathogenesis of various diseases, such as degenerative neuronal diseases, cardiovascular diseases and cancer. Recent work has led to the identification of various pro- and anti-apoptotic gene products that are involved in the regulation or execution of programmed cell death. Expression of anti-apoptotic genes, such as Bcl2 or Bcl-x_L, inhibits apoptotic cell death induced by various stimuli. On the other hand, expression of pro-apoptotic genes, such as Bax or Bad, leads to programmed cell death (Adams et al.
15 *Science*, 281:1322-1326 (1998)). The execution of programmed cell death is mediated by caspase-1 related proteinases, including caspase-3, caspase-7, caspase-8 and caspase-9 etc (Thornberry et al. *Science*, 281:1312-1316 (1998)).

20 The phosphatidylinositol 3'-OH kinase (PI3K)/Akt pathway appears important for regulating cell survival/cell death (Kulik et al. *Mol. Cell. Biol.* 17:1595-1606 (1997); Franke et al, *Cell*, 88:435-437 (1997); Kauffmann-Zeh et al. *Nature* 385:544-548 (1997) Hemmings *Science*, 275:628-630 (1997); Dudek et al., *Science*, 275:661-665 (1997)). Survival factors, such as platelet derived growth factor (PDGF), nerve growth factor (NGF) and insulin-like growth factor-1 (IGF-1), promote cell survival under various conditions by inducing the activity of PI3K (Kulik et al. 1997, Hemmings 1997). Activated PI3K leads to the production of phosphatidylinositol (3,4,5)-triphosphate (PtdIns(3,4,5)-P3),
25 which in turn binds to, and promotes the activation of, the serine/threonine kinase Akt, which contains a pleckstrin homology (PH)-domain (Franke et al *Cell*, 81:727-736 (1995); Hemmings *Science*, 277:534 (1997); Downward, *Curr. Opin. Cell Biol.* 10:262-267 (1998), Alessi et al., *EMBO J.* 15: 6541-6551 (1996)). Specific inhibitors of PI3K or dominant negative Akt mutants abolish survival-promoting activities of these growth factors or cytokines. It has been previously disclosed that inhibitors of PI3K
30 (LY294002 or wortmannin) blocked the activation of Akt by upstream kinases. In addition, introduction of constitutively active PI3K or Akt mutants promotes cell survival under conditions in which cells normally undergo apoptotic cell death (Kulik et al. 1997, Dudek et al. 1997).

35 Three members of the Akt subfamily of second-messenger regulated serine/threonine protein kinases have been identified and termed Akt1/ PKB α , Akt2/PKB β , and Akt3/PKB γ (hereinafter referred to as "Akt1", "Akt2" and "Akt3"), respectively. The isoforms are homologous, particularly in regions encoding the catalytic domains. Akts are activated by phosphorylation events occurring in

response to PI3K signaling. PI3K phosphorylates membrane inositol phospholipids, generating the second messengers phosphatidyl-inositol 3,4,5-trisphosphate and phosphatidylinositol 3,4-bisphosphate, which have been shown to bind to the PH domain of Akt. The current model of Akt activation proposes recruitment of the enzyme to the membrane by 3'-phosphorylated phosphoinositides, where
5 phosphorylation of the regulatory sites of Akt by the upstream kinases occurs (B.A. Hemmings, *Science* 275:628-630 (1997); B.A. Hemmings, *Science* 276:534 (1997); J. Downward, *Science* 279:673-674 (1998).

Phosphorylation of Akt1 occurs on two regulatory sites, Thr³⁰⁸ in the catalytic domain activation loop and on Ser⁴⁷³ near the carboxy terminus (D. R. Alessi *et al. EMBO J.* 15:6541-6551
10 (1996) and R. Meier *et al. J. Biol. Chem.* 272:30491-30497 (1997)). Equivalent regulatory phosphorylation sites occur in Akt2 and Akt3. The upstream kinase, which phosphorylates Akt at the activation loop site has been cloned and termed 3'-phosphoinositide dependent protein kinase 1 (PDK1). PDK1 phosphorylates not only Akt, but also p70 ribosomal S6 kinase, p90RSK, serum and glucocorticoid-regulated kinase (SGK), and protein kinase C. The upstream kinase phosphorylating the
15 regulatory site of Akt near the carboxy terminus has not been identified yet, but recent reports imply a role for the integrin-linked kinase (ILK-1), a serine/threonine protein kinase, or autophosphorylation.

Analysis of Akt levels in human tumors showed that Akt2 is overexpressed in a significant number of ovarian (J. Q. Cheng *et al. Proc. Natl. Acad. Sci. U.S.A.* 89:9267-9271(1992)) and pancreatic cancers (J. Q. Cheng *et al. Proc. Natl. Acad. Sci. U.S.A.* 93:3636-3641 (1996)). Similarly,
20 Akt3 was found to be overexpressed in breast and prostate cancer cell lines (Nakatani *et al. J. Biol. Chem.* 274:21528-21532 (1999)).

The tumor suppressor PTEN, a protein and lipid phosphatase that specifically removes the 3' phosphate of PtdIns(3,4,5)-P3, is a negative regulator of the PI3K/Akt pathway (Li *et al. Science* 275:1943-1947 (1997), Stambolic *et al. Cell* 95:29-39 (1998), Sun *et al. Proc. Natl. Acad. Sci. U.S.A.*
25 96:6199-6204 (1999)). Germline mutations of PTEN are responsible for human cancer syndromes such as Cowden disease (Liaw *et al. Nature Genetics* 16:64-67 (1997)). PTEN is deleted in a large percentage of human tumors and tumor cell lines without functional PTEN show elevated levels of activated Akt (Li *et al. supra*, Guldberg *et al. Cancer Research* 57:3660-3663 (1997), Risinger *et al. Cancer Research* 57:4736-4738 (1997)).

30 These observations demonstrate that the PI3K/Akt pathway plays important roles for regulating cell survival or apoptosis in tumorigenesis.

Inhibition of Akt activation and activity can be achieved by inhibiting PI3K with inhibitors such as LY294002 and wortmannin. However, PI3K inhibition has the potential to indiscriminately affect not just all three Akt isozymes but also other PH domain-containing signaling
35 molecules that are dependent on PtdIns(3,4,5)-P3, such as the Tec family of tyrosine kinases.

Furthermore, it has been disclosed that Akt can be activated by growth signals that are independent of PI3K.

Alternatively, Akt activity can be inhibited by blocking the activity of the upstream kinase PDK1. No specific PDK1 inhibitors have been disclosed. Again, inhibition of PDK1 would result in inhibition of multiple protein kinases whose activities depend on PDK1, such as atypical PKC isoforms, SGK, and S6 kinases (Williams et al. *Curr. Biol.* 10:439-448 (2000)).

It is an object of the instant invention to provide novel compounds that are inhibitors of Akt.

It is also an object of the present invention to provide pharmaceutical compositions that comprise the novel compounds that are inhibitors of Akt.


It is also an object of the present invention to provide a method for treating cancer that comprises administering such inhibitors of Akt activity.

SUMMARY OF THE INVENTION

The instant invention provides for canthine analogs that inhibit Akt activity. In particular, the compounds disclosed selectively inhibit one or two of the Akt isoforms. The invention also provides for compositions comprising such inhibitory compounds and methods of inhibiting Akt activity by administering the compound to a patient in need of treatment of cancer.


DETAILED DESCRIPTION OF THE INVENTION

The compounds of the instant invention are useful in the inhibition of the activity of the serine/threonine kinase Akt. In a first embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A:



wherein:

a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is independently 0, 1, 2, 3 or 4; p is independently 0, 1, 2, 3, 4 or 5; r is 0 or 1; s is 0 or 1; and t is 2, 3, 4, 5 or 6;



is selected from: C₃-C₈ cycloalkyl, aryl, heteroaryl and heterocyclyl;

R¹ is independently selected from: (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR^c(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR^cS(O)_mR^a,
 5 oxo, CHO, NO₂, NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R^Z;

R² is independently selected from: (C₁-C₆)alkyl-heterocyclyl, (C₁-C₆)alkyl-NR⁵R⁶, (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl,
 10 (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR^c(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR^cS(O)_mR^a, oxo, CHO, NO₂, NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^Z;

R⁵ and R⁶ are independently selected from: H, (C=O)O_bR^a, C₁-C₁₀ alkyl, aryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, heterocyclyl, C₃-C₈ cycloalkyl, SO₂R^a, and (C=O)NR^b₂, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R^Z, or R⁵ and R⁶ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in
 20 addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R^Z;

R^Z is selected from: (C=O)_rO_s(C₁-C₁₀)alkyl, O_r(C₁-C₃)perfluoroalkyl, (C₀-C₆)alkylene-S(O)_mR^a, oxo, OH, halo, CN, (C=O)_rO_s(C₂-C₁₀)alkenyl, (C=O)_rO_s(C₂-C₁₀)alkynyl, (C=O)_rO_s(C₃-C₆)cycloalkyl, (C=O)_rO_s(C₀-C₆)alkylene-aryl, (C=O)_rO_s(C₀-C₆)alkylene-heterocyclyl,
 25 (C=O)_rO_s(C₀-C₆)alkylene-N(R^b)₂, C(O)R^a, (C₀-C₆)alkylene-CO₂R^a, C(O)H, (C₀-C₆)alkylene-CO₂H, C(O)N(R^b)₂, S(O)_mR^a, S(O)₂N(R^b)₂NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from R^b, OH, (C₁-C₆)alkoxy, halogen, aryl, heterocyclyl, CO₂H, CN, O(C=O)C₁-C₆ alkyl, oxo, and N(R^b)₂, wherein said
 30 heterocyclyl is optionally substituted with from one to three substituents selected from oxo, OH, N(R^d)₂, and -O(C₁-C₆)alkyl;

R^a is: substituted or unsubstituted (C₁-C₁₀)alkyl, substituted or unsubstituted (C₂-C₁₀)alkenyl, substituted or unsubstituted (C₂-C₁₀)alkynyl, substituted or unsubstituted (C₃-

C₁₀)cycloalkyl, substituted or unsubstituted aryl, (C₁-C₆)perfluoroalkyl, 2,2,2-trifluoroethyl, or substituted or unsubstituted heterocyclyl;


R^b is: H, (C₁-C₁₀)alkyl, substituted or unsubstituted aryl, substituted or unsubstituted benzyl, substituted or unsubstituted heterocyclyl, (C₃-C₁₀)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl or S(O)₂R^a;

R^C is selected from: H, C₁-C₁₀ alkyl, aryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, heterocyclyl, C₃-C₁₀ cycloalkyl, C₁-C₆ perfluoroalkyl, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R^Z, and

R^d is independently selected from: H and (C₁-C₆)alkyl;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a second embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A1:




wherein:

R' and R" are independently selected from: H, (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR⁵(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR⁵S(O)_mR^a, oxo, CHO, NO₂, O(C=O)O_bC₁-C₁₀ alkyl, and O(C=O)O_bC₃-C₈ cycloalkyl, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one to three substituents selected from R^Z; or R' and R" can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R^Z;

R⁸ and R⁹ are independently selected from: H, (C₁-C₆)alkyl and (C₁-C₆)perfluoroalkyl, or R⁸ and R⁹ are combined to form -(CH₂)_t- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)_m, -N(R^b)C(O)-, and -N(COR^a)-;

and all other substituents and variables are as defined in the first embodiment;
or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a third embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A3:



5

wherein:


q is 0, 1, 2, 3 or 4;

R⁷ is independently selected from: (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR⁵(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR⁵S(O)_mR^a, oxo, CHO, NO₂, O(C=O)O_bC₁-C₁₀ alkyl, and O(C=O)O_bC₃-C₈ cycloalkyl, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R^z;

15

and all other substituents and variables are as defined in the second embodiment;
or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a fourth embodiment the inhibitors of the instant invention are illustrated by the Formula A4:




wherein:

all other substituents and variables are as defined in the third embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a fifth embodiment the inhibitors of the instant invention are illustrated by the

5 Formula A5:



wherein:


Q is selected from: heterocyclyl, said heterocyclyl optionally substituted with one to three substituents selected from R^Z;

10

all other substituents and variables are as defined in the fourth embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a sixth embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A6:




15

wherein:

all other substituents and variables are as defined in the second embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a seventh embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A7:




wherein:

all other substituents and variables are as defined in the third embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In an eighth embodiment the inhibitors of the instant invention are illustrated by the

Formula A8:




wherein:

all other substituents and variables are as defined in the fourth embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a ninth embodiment the inhibitors of the instant invention are illustrated by the

Formula A9:



wherein:

all other substituents and variables are as defined in the fifth embodiment;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

5 Specific compounds of the instant invention include:

1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-5); and

1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-6);

10 or a pharmaceutically acceptable salt or stereoisomer thereof.

Examples of the compounds of the instant invention include TFA salts of the following compounds:


1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-5) and

15 1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-6):


or a stereoisomer thereof

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, all such stereoisomers being included in the present invention.

In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the invention, even though only one tautomeric structure is depicted. For example, any claim to compound A below is understood to include tautomeric structure B, and vice versa, as well as mixtures thereof.



Tetrazoles exist as a mixture of 1H/2H tautomers. The tautomeric forms of the tetrazol moiety are also within the scope of the instant invention.



When any variable (e.g. R¹, R², R^z, etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds. Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms. If the ring system is polycyclic, it is intended that the bond be attached to any of the suitable carbon or nitrogen atoms on the proximal ring(s) only.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase "optionally substituted with one or more substituents" should be taken to be equivalent to the phrase "optionally substituted with at least one substituent" and in such cases the preferred embodiment will have from zero to four substituents, and the more preferred embodiment will have from zero to three substituents.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C₁-C₁₀, as in "C₁-C₁₀ alkyl" is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, "C₁-C₁₀ alkyl" specifically includes methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *i*-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and so on. The term "cycloalkyl" means a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms. For example, "cycloalkyl" includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, and so on.

"Alkoxy" represents either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge. "Alkoxy" therefore encompasses the definitions of alkyl and cycloalkyl above.

If no number of carbon atoms is specified, the term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Thus, "C₂-C₆ alkenyl" means an alkenyl radical having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl, 2-methylbutenyl and cyclohexenyl. The straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Thus, "C₂-C₆ alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. The straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.

In certain instances, substituents may be defined with a range of carbons that includes zero, such as (C₀-C₆)alkylene-aryl. If aryl is taken to be phenyl, this definition would include phenyl itself as well as -CH₂Ph, -CH₂CH₂Ph, CH(CH₃)CH₂CH(CH₃)Ph, and so on.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydro-naphthyl, indanyl and biphenyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring.

The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, carbazolyl, cinnolinyl, quinoxalinyl, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, indolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heteroaryl. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively. Such heteroaryl moieties for substituent Q include but are not limited to: 2-benzimidazolyl, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 1-isoquinolinyl, 3-isoquinolinyl and 4-isoquinolinyl.

The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 3- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrathydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzoimidazolyl, benzoimidazolonyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolyl, furanyl, imidazolyl, indolyl, indolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolyl, quinolyl, quinoxalyl, tetrahydropyranyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidyl, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholyl, thiomorpholyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolyl, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidyl, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof. Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.

As used herein, the terms "substituted C₁-C₁₀ alkyl", "substituted C₂-C₁₀ alkenyl", and "substituted C₂-C₁₀ alkynyl" are intended to include the branch or straight-chain alkyl group of the specified number of carbon atoms, wherein the carbon atoms may be substituted with 1 to 3 substituents selected from the group which includes, but is not limited to, halo, C₁-C₂₀ alkyl, CF₃, NH₂, N(C₁-C₆ alkyl)₂, NO₂, oxo, CN, N₃, -OH, -O(C₁-C₆ alkyl), C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₀-C₆ alkyl) S(O)₀₋₂, (C₀-C₆ alkyl)S(O)₀₋₂(C₀-C₆ alkyl), (C₀-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, -O(C₁-C₆ alkyl)CF₃, (C₀-C₆ alkyl)C(O)-, (C₀-C₆ alkyl)OC(O)-, (C₀-C₆ alkyl) O(C₁-C₆ alkyl)-, (C₀-C₆ alkyl)C(O)₁₋₂(C₀-C₆ alkyl)-, (C₀-C₆ alkyl)OC(O)NH-, aryl, aralkyl, heterocycle, heterocyclylalkyl, halo-aryl, halo-aralkyl, halo-heterocycle, halo-heterocyclylalkyl, cyano-aryl, cyano-aralkyl, cyano-heterocycle and cyano-heterocyclylalkyl.

As used herein, the terms "substituted C₃-C₁₀ cycloalkyl", "substituted aryl", and "substituted heterocycle", are intended to include the cyclic group containing from 1 to 3 substituents in addition to the point of attachment to the rest of the compound. Preferably, the substituents are selected from the group which includes, but is not limited to, halo, C₁-C₂₀ alkyl, CF₃, NH₂, N(C₁-C₆ alkyl)₂, NO₂, oxo, CN, N₃, -OH, -O(C₁-C₆ alkyl), C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₀-C₆ alkyl) S(O)₀₋₂, (C₀-C₆ alkyl)S(O)₀₋₂(C₀-C₆ alkyl)-, (C₀-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, -O(C₁-C₆ alkyl)CF₃, (C₀-C₆ alkyl)C(O)-, (C₀-C₆ alkyl)OC(O)-, (C₀-C₆alkyl)O(C₁-C₆ alkyl)-, (C₀-C₆

alkyl)C(O)₁₋₂(C₀-C₆ alkyl)-, (C₀-C₆ alkyl)OC(O)NH-, aryl, aralkyl, heteroaryl, heterocyclalkyl, halo-aryl, halo-aralkyl, halo-heterocycle, halo-heterocyclalkyl, cyano-aryl, cyano-aralkyl, cyano-heterocycle and cyano-heterocyclalkyl.

As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro (Ch), fluoro (F), bromo (Br) and iodo (I).

In an embodiment, n is independently 0, 1 or 2.

In another embodiment, n is 0.

In an embodiment, p is independently 0, 1 or 2.


In another embodiment, p is 0.




In an embodiment, A is selected from: phenyl.

In an embodiment, Q is selected from: 2-azepinone, benzimidazolyl, benzimidazolonyl, 2-diazapinone, imidazolyl, 2-imidazolidinone, indolyl, isoquinoliny, morpholiny, piperidyl, piperaziny, pyridyl, pyrrolidiny, 2-piperidinone, 2-pyrimidinone, 2-pyrrolidinone, quinoliny, tetrazolyl, tetrahydrofuryl, tetrahydroisoquinoliny, thienyl, pyrazolopyrimidinyl, pyrazolyl, thiazolyl, oxadiazolyl and triazolyl, optionally substituted with 1-3 R^Z.

In another embodiment, Q is selected from:



which are optionally substituted with one to three substituents selected from R^Z.



In another embodiment, Q is selected from:

In an embodiment, R¹ is selected from: H, (C=O)aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl and heterocyclyl, said alkyl, aryl cycloalkyl

and heterocyclyl optionally substituted with one to three substituents selected from: (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl.

In another embodiment, R¹ is selected from: H, (C=O)_aOb(C₁-C₆)alkyl, NH₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, phenyl and (C₃-C₈)cycloalkyl, said alkyl, phenyl and cycloalkyl

5 optionally substituted with one to three substituents selected from: (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl.

In an embodiment, R² is independently selected from: (C₁-C₆)alkyl-heterocyclyl, (C₁-C₆)alkyl-NR⁵R⁶, (C=O)_aObC₁-C₁₀ alkyl, (C=O)_aObaryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aOb heterocyclyl, (C=O)_aObC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, ObC₁-C₆ perfluoroalkyl,

10 O_a(C=O)_bNR⁵R⁶, NR^c(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR^cS(O)_mR^a, CHO, NO₂, NR^c(C=O)ObR^a, O(C=O)ObC₁-C₁₀ alkyl, O(C=O)ObC₃-C₈ cycloalkyl, O(C=O)Obaryl, and O(C=O)Ob-heterocycle, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^Z.

In another embodiment, R² is independently selected from: (C₁-C₆)alkyl-heterocyclyl, (C₁-C₆)alkyl-NR⁵R⁶, said heterocyclyl optionally substituted with one, two or three substituents selected from R^Z.

In another embodiment, R² is selected from: CH₃-NR⁵R⁶.

In yet another embodiment, R² is selected from: halo, OH, N(R^d)₂, oxo and O_a(C₁-C₆)alkyl.

20 In an embodiment, R⁵ and R⁶ are selected from: H, (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl and heterocyclyl, said alkyl, aryl cycloalkyl and heterocyclyl optionally substituted with one to three substituents selected from: (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl.

In an embodiment, R⁷ is selected from: H, (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl and heterocyclyl, said alkyl, aryl cycloalkyl and heterocyclyl optionally substituted with one to three substituents selected from: R^Z.

In an embodiment, R⁸ and R⁹ are selected from H and (C₁-C₆)alkyl.

In another embodiment, R⁸ and R⁹ are H.

30 In an embodiment, R^Z is selected from: H, (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl and heterocyclyl, said alkyl, aryl cycloalkyl and heterocyclyl optionally substituted with one to three substituents selected from: (C=O)_aOb(C₁-C₆)alkyl, NR^b₂, OH, oxo, halo, NO₂, C(O)H, C(O)OH, CF₃, aryl, (C₃-C₈)cycloalkyl and heterocyclyl, wherein said heterocyclyl is optionally substituted with one to three substituents selected from oxo, OH, N(R^d)₂, and O_a(C₁-C₆)alkyl.

Included in the instant invention is the free form of compounds of Formula A, as well as the pharmaceutically acceptable salts and stereoisomers thereof. Some of the isolated specific compounds exemplified herein are the protonated salts of amine compounds. The term "free form" refers to the amine compounds in non-salt form. The encompassed pharmaceutically acceptable salts not only include the isolated salts exemplified for the specific compounds described herein, but also all the typical pharmaceutically acceptable salts of the free form of compounds of Formula A. The free form of the specific salt compounds described may be isolated using techniques known in the art. For example, the free form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free forms may differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise pharmaceutically equivalent to their respective free forms for purposes of the invention.

The pharmaceutically acceptable salts of the instant compounds can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

Thus, pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed by reacting a basic instant compound with an inorganic or organic acid. For example, conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like, as well as salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic (TFA) and the like.

When the compound of the present invention is acidic, suitable "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine,

ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine tripropylamine, tromethamine and the like.

5 The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al.*, "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977:66:1-19.

10 It will also be noted that the compounds of the present invention are potentially internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary nitrogen atom.

UTILITY

15 The compounds of the instant invention are inhibitors of the activity of Akt and are thus useful in the treatment of cancer, in particular cancers associated with irregularities in the activity of Akt and downstream cellular targets of Akt. Such cancers include, but are not limited to, ovarian, pancreatic, breast and prostate cancer, as well as cancers (including glioblastoma) where the tumor suppressor PTEN is mutated (Cheng *et al.*, *Proc. Natl. Acad. Sci.* (1992) 89:9267-9271; Cheng *et al.*, *Proc. Natl. Acad. Sci.* (1996) 93:3636-3641; Bellacosa *et al.*, *Int. J. Cancer* (1995) 64:280-285; Nakatani *et al.*, *J. Biol. Chem.* (1999) 274:21528-21532; Graff, *Expert. Opin. Ther. Targets* (2002) 6(1):103-113; and Yamada and Araki, *J. Cell Science.* (2001) 114:2375-2382; Mischel and Cloughesy, *Brain Pathol.* (2003) 13(1):52-61).

25 The compounds, compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, 30 sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, 35 Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma,

transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone:
 5 osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma,
 10 gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell
 15 tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic
 20 syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

25 Akt signaling regulates multiple critical steps in angiogenesis. Shiojima and Walsh, *Circ. Res.* (2002) 90:1243-1250. The utility of angiogenesis inhibitors in the treatment of cancer is known in the literature, see J. Rak et al. *Cancer Research*, 55:4575-4580, 1995 and Dredge et al., *Expert Opin. Biol. Ther.* (2002) 2(8):953-966, for example. The role of angiogenesis in cancer has been shown in numerous types of cancer and tissues: breast carcinoma (G. Gasparini and A.L. Harris, *J. Clin.*
 30 *Oncol.*, 1995, 13:765-782; M. Toi et al., *Japan. J. Cancer Res.*, 1994, 85:1045-1049); bladder carcinomas (A.J. Dickinson et al., *Br. J. Urol.*, 1994, 74:762-766); colon carcinomas (L.M. Ellis et al., *Surgery*, 1996, 120(5):871-878); and oral cavity tumors (J.K. Williams et al., *Am. J. Surg.*, 1994, 168:373-380). Other cancers include, advanced tumors, hairy cell leukemia, melanoma, advanced head and neck, metastatic renal cell, non-Hodgkin's lymphoma, metastatic breast, breast adenocarcinoma,
 35 advanced melanoma, pancreatic, gastric, glioblastoma, lung, ovarian, non-small cell lung, prostate, small cell lung, renal cell carcinoma, various solid tumors, multiple myeloma, metastatic prostate, malignant

glioma, renal cancer, lymphoma, refractory metastatic disease, refractory multiple myeloma, cervical cancer, Kaposi's sarcoma, recurrent anaplastic glioma, and metastatic colon cancer (Dredge et al., *Expert Opin. Biol. Ther.* (2002) 2(8):953-966). Thus, the Akt inhibitors disclosed in the instant application are also useful in the treatment of these angiogenesis related cancers.

5 Tumors which have undergone neovascularization show an increased potential for metastasis. In fact, angiogenesis is essential for tumor growth and metastasis. (S.P. Cunningham, et al., *Can. Research*, 61: 3206-3211 (2001)). The Akt inhibitors disclosed in the present application are therefore also useful to prevent or decrease tumor cell metastasis.

10 Further included within the scope of the invention is a method of treating or preventing a disease in which angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of the present invention. Ocular neovascular diseases are an example of conditions where much of the resulting tissue damage can be attributed to aberrant infiltration of blood vessels in the eye (see WO 00/30651, published 2 June 2000). The undesirable infiltration can be triggered by ischemic retinopathy, such as that resulting from
15 diabetic retinopathy, retinopathy of prematurity, retinal vein occlusions, etc., or by degenerative diseases, such as the choroidal neovascularization observed in age-related macular degeneration. Inhibiting the growth of blood vessels by administration of the present compounds would therefore prevent the infiltration of blood vessels and prevent or treat diseases where angiogenesis is implicated, such as ocular diseases like retinal vascularization, diabetic retinopathy, age-related macular degeneration, and
20 the like.

Further included within the scope of the invention is a method of treating or preventing a non-malignant disease in which angiogenesis is implicated, including but not limited to: ocular diseases (such as, retinal vascularization, diabetic retinopathy and age-related macular degeneration), atherosclerosis, arthritis, psoriasis, obesity and Alzheimer's disease (Dredge et al., *Expert Opin. Biol.*
25 *Ther.* (2002) 2(8):953-966). In another embodiment, a method of treating or preventing a disease in which angiogenesis is implicated includes: ocular diseases (such as, retinal vascularization, diabetic retinopathy and age-related macular degeneration), atherosclerosis, arthritis and psoriasis.

Further included within the scope of the invention is a method of treating hyperproliferative disorders such as restenosis, inflammation, autoimmune diseases and allergy/asthma.

30 Further included within the scope of the instant invention is the use of the instant compounds to coat stents and therefore the use of the instant compounds on coated stents for the treatment and/or prevention of restenosis (WO03/032809).

Further included within the scope of the instant invention is the use of the instant compounds for the treatment and/or prevention of osteoarthritis (WO03/035048).

35 Further included within the scope of the invention is a method of treating hyperinsulinism.

The compounds of the invention are also useful in preparing a medicament that is useful in treating the diseases described above, in particular cancer.

In an embodiment of the invention, the instant compound is a selective inhibitor whose inhibitory efficacy is dependent on the PH domain. In this embodiment, the compound exhibits a
5 decrease in *in vitro* inhibitory activity or no *in vitro* inhibitory activity against truncated Akt proteins lacking the PH domain.

In a further embodiment, the instant compound is selected from the group of a selective inhibitor of Akt1, a selective inhibitor of Akt2 and a selective inhibitor of both Akt1 and Akt2.

In another embodiment, the instant compound is selected from the group of a selective
10 inhibitor of Akt1, a selective inhibitor of Akt2, a selective inhibitor of Akt3 and a selective inhibitor of two of the three Akt isoforms.

In another embodiment, the instant compound is a selective inhibitor of all three Akt isoforms, but is not an inhibitor of one, two or all of such Akt isoforms that have been modified to delete the PH domain, the hinge region or both the PH domain and the hinge region.

The present invention is further directed to a method of inhibiting Akt activity which
15 comprises administering to a mammal in need thereof a pharmaceutically effective amount of the instant compound.

The compounds of this invention may be administered to mammals, including humans, either alone or, in combination with pharmaceutically acceptable carriers, excipients or diluents, in a
20 pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible
25 powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in
30 admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example,
35 magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the

gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropylmethyl-cellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty

acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring agents, preservatives and antioxidants.

5 Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of sterile injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

10 The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

15 The injectable solutions or microemulsions may be introduced into a patient's blood-stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

20 The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as
25 a solution in 1,3-butane diol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

30 Compounds of Formula A may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

35 For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula A are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

When a composition according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In an embodiment, a suitable amount of an inhibitor of Akt is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount of inhibitor of between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, or between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day. Another therapeutic dosage that comprises the instant composition includes from about 0.01 mg to about 1000 mg of inhibitor of Akt. In another embodiment, the dosage comprises from about 1 mg to about 1000 mg of inhibitor of Akt.

The instant compounds are also useful in combination with known therapeutic agents and anti-cancer agents. For example, instant compounds are useful in combination with known anti-cancer agents. Combinations of the presently disclosed compounds with other anti-cancer or chemotherapeutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V.T. Devita and S. Hellman (editors), 6th edition (February 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and other angiogenesis inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, inhibitors of cell proliferation and survival signaling, and agents that interfere with cell cycle checkpoints. The instant compounds are particularly useful when co-administered with radiation therapy.

"Estrogen receptor modulators" refers to compounds that interfere with or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)ethoxy]phenyl)]-2H-1-

benzopyran-3-yl]-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

"Androgen receptor modulators" refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism. Examples of androgen receptor modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

"Retinoid receptor modulators" refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl) retinamide, and N-4-carboxyphenyl retinamide.

"Cytotoxic/cytostatic agents" refer to compounds which cause cell death or inhibit cell proliferation primarily by interfering directly with the cell's functioning or inhibit or interfere with cell myosis, including alkylating agents, tumor necrosis factors, intercalators, hypoxia activatable compounds, microtubule inhibitors/microtubule-stabilizing agents, inhibitors of mitotic kinesins, histone deacetylase inhibitors, inhibitors of kinases involved in mitotic progression, inhibitors of kinases involved in growth factor and cytokine signal transduction pathways, antimetabolites, biological response modifiers, hormonal/anti-hormonal therapeutic agents, haematopoietic growth factors, monoclonal antibody targeted therapeutic agents, topoisomerase inhibitors, proteosome inhibitors, ubiquitin ligase inhibitors, and aurora kinase inhibitors.

Examples of cytotoxic/cytostatic agents include, but are not limited to, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methyl-pyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum(II)]tetrachloride, diarizidynylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, 4-demethoxy-3-deamino-3-aziridynyl-4-methylsulphonyl-daunorubicin (see WO 00/50032), Raf kinase inhibitors (such as Bay43-9006) and mTOR inhibitors (such as Wyeth's CCI-779).

An example of a hypoxia activatable compound is tirapazamine.

Examples of proteosome inhibitors include but are not limited to lactacystin and MLN-341 (Velcade).

Examples of microtubule inhibitors/microtubule-stabilising agents include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl) benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, the epothilones (see for example U.S. Pat. Nos. 6,284,781 and 6,288,237) and BMS188797. In an embodiment the epothilones are not included in the microtubule inhibitors/microtubule-stabilising agents.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrzolo[3,4,5-kl]acridine-2-(6H) propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]-indolizino[1,2b]quinoline-10,13(9H,15H)dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2-(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c] quinolin-7-one, and dimesna.

Examples of inhibitors of mitotic kinesins, and in particular the human mitotic kinesin KSP, are described in PCT Publications WO 01/30768, WO 01/98278, WO 03/049527, WO 03/049679, WO 03/050064, WO 03/050122, WO 03/049678 and WO 03/039460. In an embodiment inhibitors of mitotic kinesins include, but are not limited to inhibitors of KSP, inhibitors of MKLP1, inhibitors of CENP-E, inhibitors of MCAK and inhibitors of Rab6-KIFL.

Examples of "histone deacetylase inhibitors" include, but are not limited to, SAHA, TSA, oxamflatin, PXD101, MG98 and scriptaid. Further reference to other histone deacetylase inhibitors may be found in the following manuscript; Miller, T.A. et al. *J. Med. Chem.* 46(24):5097-5116 (2003).

"Inhibitors of kinases involved in mitotic progression" include, but are not limited to, inhibitors of aurora kinase, inhibitors of Polo-like kinases (PLK; in particular inhibitors of PLK-1), inhibitors of bub-1 and inhibitors of bub-R1. An example of an "aurora kinase inhibitor" is VX-680.

"Antiproliferative agents" includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine,

carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-
 5 [2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine, 3-aminopyridine-2-carboxaldehyde thiosemicarbazone and
 10 trastuzumab.

Examples of monoclonal antibody targeted therapeutic agents include those therapeutic agents which have cytotoxic agents or radioisotopes attached to a cancer cell specific or target cell specific monoclonal antibody. Examples include Bexxar.

15 "HMG-CoA reductase inhibitors" refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see U.S. Patent Nos. 4,231,938, 4,294,926 and 4,319,039), simvastatin (ZOCOR®; see U.S. Patent Nos. 4,444,784, 4,820,850 and 4,916,239), pravastatin (PRAVACHOL®; see U.S. Patent Nos. 4,346,227, 4,537,859, 4,410,629, 5,030,447 and 5,180,589),
 20 fluvastatin (LESCOL®; see U.S. Patent Nos. 5,354,772, 4,911,165, 4,929,437, 5,189,164, 5,118,853, 5,290,946 and 5,356,896), atorvastatin (LIPITOR®; see U.S. Patent Nos. 5,273,995, 4,681,893, 5,489,691 and 5,342,952) and cerivastatin (also known as rivastatin and BAYCHOL®; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs",
 25 *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention.

30 "Prenyl-protein transferase inhibitor" refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase).

Examples of prenyl-protein transferase inhibitors can be found in the following
 35 publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Patent No. 5,420,245, U.S. Patent No. 5,523,430,

U.S. Patent No. 5,532,359, U.S. Patent No. 5,510,510, U.S. Patent No. 5,589,485, U.S. Patent No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Patent No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Patent No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Patent No. 5,532,359. For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see *European J. of Cancer*, Vol. 35, No. 9, pp.1394-1401 (1999).

"Angiogenesis inhibitors" refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- α , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (*PNAS*, Vol. 89, p. 7384 (1992); *JNCI*, Vol. 69, p. 475 (1982); *Arch. Ophthalmol.*, Vol. 108, p.573 (1990); *Anat. Rec.*, Vol. 238, p. 68 (1994); *FEBS Letters*, Vol. 372, p. 83 (1995); *Clin. Orthop.* Vol. 313, p. 76 (1995); *J. Mol. Endocrinol.*, Vol. 16, p.107 (1996); *Jpn. J. Pharmacol.*, Vol. 75, p. 105 (1997); *Cancer Res.*, Vol. 57, p. 1625 (1997); *Cell*, Vol. 93, p. 705 (1998); *Intl. J. Mol. Med.*, Vol. 2, p. 715 (1998); *J. Biol. Chem.*, Vol. 274, p. 9116 (1999)), steroidal anti-inflammatories (such as corticosteroids, mineralocorticoids, dexamethasone, prednisone, prednisolone, methylpred, betamethasone), carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez et al., *J. Lab. Clin. Med.* 105:141-145 (1985)), and antibodies to VEGF (see, *Nature Biotechnology*, Vol. 17, pp.963-968 (October 1999); Kim et al., *Nature*, 362, 841-844 (1993); WO 00/44777; and WO 00/61186).

Other therapeutic agents that modulate or inhibit angiogenesis and may also be used in combination with the compounds of the instant invention include agents that modulate or inhibit the coagulation and fibrinolysis systems (see review in *Clin. Chem. La. Med.* 38:679-692 (2000)). Examples of such agents that modulate or inhibit the coagulation and fibrinolysis pathways include, but are not limited to, heparin (see *Thromb. Haemost.* 80:10-23 (1998)), low molecular weight heparins and carboxypeptidase U inhibitors (also known as inhibitors of active thrombin activatable fibrinolysis

inhibitor [TAFIa]) (see *Thrombosis Res.* 101:329-354 (2001)). TAFIa inhibitors have been described in WO 03/13526.

“Agents that interfere with cell cycle checkpoints” refer to compounds that inhibit protein kinases that transduce cell cycle checkpoint signals, thereby sensitizing the cancer cell to DNA
5 damaging agents. Such agents include inhibitors of ATR, ATM, the Chk1 and Chk2 kinases and cdk and cdc kinase inhibitors and are specifically exemplified by 7-hydroxystaurosporin, flavopiridol, CYC202 (Cyclacel) and BMS-387032.

“Inhibitors of cell proliferation and survival signalling pathway” refer to compounds that inhibit signal transduction cascades downstream of cell surface receptors. Such agents include inhibitors
10 of serine/threonine kinases (including but not limited to inhibitors of Akt such as described in WO 02/083064, WO 02/083139, WO 02/083140, WO 02/083138, WO 03/086279, WO 03/086394, WO 03/086403, WO 03/086404 and WO 04/041162), inhibitors of Raf kinase (for example BAY-43-9006), inhibitors of MEK (for example CI-1040 and PD-098059), inhibitors of mTOR (for example Wyeth CCI-779), and inhibitors of PI3K (for example LY294002).

As described above, the combinations with NSAID's are directed to the use of NSAID's which are potent COX-2 inhibiting agents. For purposes of this specification an NSAID is potent if it possesses an IC₅₀ for the inhibition of COX-2 of 1μM or less as measured by cell or microsomal assays.

The invention also encompasses combinations with NSAID's which are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are
20 defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of IC₅₀ for COX-2 over IC₅₀ for COX-1 evaluated by cell or microsomal assays. Such compounds include, but are not limited to those disclosed in U.S. Patent 5,474,995, U.S. Patent 5,861,419, U.S. Patent 6,001,843, U.S. Patent 6,020,343, U.S. Patent 5,409,944, U.S. Patent 5,436,265, U.S. Patent 5,536,752, U.S. Patent 5,550,142, U.S. Patent 5,604,260, U.S. 5,698,584, U.S. Patent
25 5,710,140, WO 94/15932, U.S. Patent 5,344,991, U.S. Patent 5,134,142, U.S. Patent 5,380,738, U.S. Patent 5,393,790, U.S. Patent 5,466,823, U.S. Patent 5,633,272 and U.S. Patent 5,932,598, all of which are hereby incorporated by reference.

Inhibitors of COX-2 that are particularly useful in the instant method of treatment are: 3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone; and
30 5-chloro-3-(4-methylsulfonyl)phenyl-2-(2-methyl-5-pyridinyl)pyridine; or a pharmaceutically acceptable salt thereof.

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to, the following: parecoxib, BEXTRA® and CELEBREX® or a pharmaceutically acceptable salt thereof.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin,
35 ukrain, ranpirnase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-

6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1H-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-pyrrolocarbonylimino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-
 5 [(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_3$ integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_5$ integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the $\alpha_v\beta_3$
 10 integrin and the $\alpha_v\beta_5$ integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins. The term also refers to antagonists of any combination of $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins.

Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-
 15 5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571,
 20 CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

Combinations with compounds other than anti-cancer compounds are also encompassed
 25 in the instant methods. For example, combinations of the instantly claimed compounds with PPAR- γ (i.e., PPAR-gamma) agonists and PPAR- δ (i.e., PPAR-delta) agonists are useful in the treatment of certain malignancies. PPAR- γ and PPAR- δ are the nuclear peroxisome proliferator-activated receptors γ and δ . The expression of PPAR- γ on endothelial cells and its involvement in angiogenesis has been reported in the literature (see *J. Cardiovasc. Pharmacol.* 1998; 31:909-913; *J. Biol. Chem.*
 30 1999;274:9116-9121; *Invest. Ophthalmol Vis. Sci.* 2000; 41:2309-2317). More recently, PPAR- γ agonists have been shown to inhibit the angiogenic response to VEGF in vitro; both troglitazone and rosiglitazone maleate inhibit the development of retinal neovascularization in mice. (*Arch. Ophthalmol.* 2001; 119:709-717). Examples of PPAR- γ agonists and PPAR- γ/α agonists include, but are not limited to, thiazolidinediones (such as DRF2725, CS-011, troglitazone, rosiglitazone, and pioglitazone),
 35 fenofibrate, gemfibrozil, clofibrate, GW2570, SB219994, AR-H039242, JTT-501, MCC-555, GW2331, GW409544, NN2344, KRP297, NP0110, DRF4158, NN622, GI262570, PNU182716, DRF552926, 2-

[(5,7-dipropyl-3-trifluoromethyl-1,2-benzisoxazol-6-yl)oxy]-2-methylpropionic acid WO 01/60807, and 2(R)-7-(3-(2-chloro-4-(4-fluorophenoxy) phenoxy)propoxy)-2-ethylchromane-2-carboxylic acid WO 02/026729.

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with gene therapy for the treatment of cancer. For an overview of genetic strategies to treating cancer see Hall et al (*Am. J. Hum. Genet.* 61:785-789, 1997) and Kufe et al (*Cancer Medicine*, 5th Ed, pp 876-889, BC Decker, Hamilton 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Patent No. 6,069,134, for example), a uPA/uPAR antagonist ("Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and Dissemination in Mice," *Gene Therapy*, August 1998;5(8):1105-13), and interferon gamma (*J. Immunol.* 2000;164:217-222).

The compounds of the instant invention may also be administered in combination with an inhibitor of inherent multidrug resistance (MDR), in particular MDR associated with high levels of expression of transporter proteins. Such MDR inhibitors include inhibitors of p-glycoprotein (P-gp), such as LY335979, XR9576, OC144-093, R101922, VX853 and PSC833 (valsopodar).

A compound of the present invention may be employed in conjunction with anti-emetic agents to treat nausea or emesis, including acute, delayed, late-phase, and anticipatory emesis, which may result from the use of a compound of the present invention, alone or with radiation therapy. For the prevention or treatment of emesis, a compound of the present invention may be used in conjunction with other anti-emetic agents, especially neurokinin-1 receptor antagonists, 5HT3 receptor antagonists, such as ondansetron, granisetron, tropisetron, and zatisetron, GABAB receptor agonists, such as baclofen, a corticosteroid such as Decadron (dexamethasone), Kenalog, Aristocort, Nasalide, Preferid, Benecorten or others such as disclosed in U.S. Patent Nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712, an antidopaminergic, such as the phenothiazines (for example prochlorperazine, fluphenazine, thioridazine and mesoridazine), metoclopramide or dronabinol. In another embodiment, conjunctive therapy with an anti-emesis agent selected from a neurokinin-1 receptor antagonist, a 5HT3 receptor antagonist and a corticosteroid is disclosed for the treatment or prevention of emesis that may result upon administration of the instant compounds.

Neurokinin-1 receptor antagonists of use in conjunction with the compounds of the present invention are fully described, for example, in U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699, 5,719,147; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0 436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275, 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535,

0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and
0 776 893; PCT International Patent Publication Nos. WO 90/05525, 90/05729, 91/09844, 91/18899,
92/01688, 92/06079, 92/12151, 92/15585, 92/17449, 92/20661, 92/20676, 92/21677, 92/22569,
93/00330, 93/00331, 93/01159, 93/01165, 93/01169, 93/01170, 93/06099, 93/09116, 93/10073,
5 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440,
94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843,
94/08997, 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767,
94/15903, 94/19320, 94/19323, 94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040,
95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679,
10 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338,
95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 96/05193, 96/05203, 96/06094, 96/07649,
96/10562, 96/16939, 96/18643, 96/20197, 96/21661, 96/29304, 96/29317, 96/29326, 96/29328,
96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362,
97/18206, 97/19084, 97/19942 and 97/21702; and in British Patent Publication Nos. 2 266 529, 2 268
15 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689. The preparation
of such compounds is fully described in the aforementioned patents and publications, which are
incorporated herein by reference.

In an embodiment, the neurokinin-1 receptor antagonist for use in conjunction with the
compounds of the present invention is selected from: 2-(R)-(1-(R)-(3,5-
20 bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-
triazolo)methyl)morpholine, or a pharmaceutically acceptable salt thereof, which is described in U.S.
Patent No. 5,719,147.

A compound of the instant invention may also be administered with an agent useful in
the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis
25 receptor activator (such as epoetin alfa).

A compound of the instant invention may also be administered with an agent useful in
the treatment of neutropenia. Such a neutropenia treatment agent is, for example, a hematopoietic
growth factor which regulates the production and function of neutrophils such as a human granulocyte
colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

30 A compound of the instant invention may also be administered with an immunologic-
enhancing drug, such as levamisole, isoprinosine and Zadaxin.

A compound of the instant invention may also be useful for treating or preventing
cancer, including bone cancer, in combination with bisphosphonates (understood to include
bisphosphonates, diphosphonates, bisphosphonic acids and diphosphonic acids). Examples of
35 bisphosphonates include but are not limited to: etidronate (Didronel), pamidronate (Aredia), alendronate
(Fosamax), risedronate (Actonel), zoledronate (Zometa), ibandronate (Boniva), incadronate or

cimadronate, clodronate, EB-1053, minodronate, neridronate, piridronate and tiludronate including any and all pharmaceutically acceptable salts, derivatives, hydrates and mixtures thereof.

Thus, the scope of the instant invention encompasses the use of the instantly claimed compounds in combination with a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, PPAR- γ agonists, PPAR- δ agonists, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia, an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, a bisphosphonate and an agent that interferes with a cell cycle checkpoint.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

In an embodiment, the angiogenesis inhibitor to be used as the second compound is selected from a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP (matrix metalloprotease) inhibitor, an integrin blocker, interferon- α , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, or an antibody to VEGF. In an embodiment, the estrogen receptor modulator is tamoxifen or raloxifene.

Also included in the scope of the claims is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula A in combination with

radiation therapy and/or in combination with a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, PPAR- γ agonists, PPAR- δ agonists, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia, an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, a bisphosphonate and an agent that interferes with a cell cycle checkpoint.

And yet another embodiment of the invention is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula A in combination with paclitaxel or trastuzumab.

The invention further encompasses a method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Formula A in combination with a COX-2 inhibitor.

The instant invention also includes a pharmaceutical composition useful for treating or preventing cancer that comprises a therapeutically effective amount of a compound of Formula A and a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, a PPAR- γ agonist, a PPAR- δ agonist, an inhibitor of cell proliferation and survival signaling, a bisphosphonate, and an agent that interferes with a cell cycle checkpoint.

All patents, publications and pending patent applications identified are hereby incorporated by reference.

Abbreviations used in the description of the chemistry and in the Examples that follow are: AEBSF (p-aminoethylbenzenesulfonyl fluoride); BSA (bovine serum albumin); BuLi (n-Butyl lithium); CDCl₃ (chloroform-d); CuI (copper iodide); CuSO₄ (copper sulfate); DCE (dichloroethane); DCM (dichloromethane); DEAD (diethyl azodicarboxylate); DMF (N,N-dimethylformamide); DMSO (dimethyl sulfoxide); DTT (dithiothreitol); EDTA (ethylene-diamine-tetra-acetic acid); EGTA (ethylene-glycol-tetra-acetic acid); EtOAc (ethyl acetate); EtOH (ethanol); HOAc (acetic acid); HPLC (high-performance liquid chromatography); HRMS (high resolution mass spectrum); LCMS (liquid chromatograph-mass spectrometer); LHMDs (lithium bis(trimethylsilyl)amide); LRMS (low resolution mass spectrum); MeOH (methanol); MP-B(CN)H₃ (Macroporous cyanoborohydride); NaHCO₃ (sodium bicarbonate); Na₂SO₄ (sodium sulfate); Na(OAc)₃BH (sodium triacetoxymborohydride); NH₄OAc (ammonium acetate); NBS (N-bromosuccinamide); NMR (nuclear magnetic resonance); PBS (phosphate buffered saline); PCR (polymerase chain reaction); Pd(dppf) ([1,1'-bis(diphenylphosphino)ferrocene]


palladium); Pd(Ph₃)₄ (palladium(0) tetrakis-triphenylphosphine); POCl₃ (phosphorous oxychloride); PS-DIEA (polystyrene diisopropylethylamine); PS-PPh₃ (polystyrene-triphenyl phosphine); TBAF (tetrabutylammonium fluoride); THF (tetrahydrofuran); TFA (trifluoroacetic acid); and TMSCH₂N₂ (trimethylsilyldiazomethane); CH₂Cl₂/DCM (dichloromethane); DIEA (diisopropylethylamine).

5 Several methods for preparing the compounds of this invention are illustrated in the following General Reaction Scheme and Scheme 1. Starting materials and the requisite intermediates are in some cases commercially available, or can be prepared according to literature procedures or as illustrated herein.

10 The compounds of this invention may be prepared by employing reactions as shown in the General Reaction Scheme, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. Substituent numbering as shown in the General Reaction Scheme does not necessarily correlate to that used in the claims and often, for clarity, a single substituent is shown attached to the compound where multiple substituents are allowed under the definitions hereinabove. Reactions used to generate the compounds of this invention are prepared by employing
15 reactions as shown in the General Reaction Scheme and Scheme 1 herein, in addition to other standard manipulations such as ester hydrolysis, cleavage of protecting groups, etc., as may be known in the literature or exemplified in the experimental procedures.

20 In some cases the final product may be further modified, for example, by manipulation of substituents. These manipulations may include, but are not limited to, reduction, oxidation, alkylation, acylation, and hydrolysis reactions which are commonly known to those skilled in the art. In some cases the order of carrying out the foregoing reaction schemes may be varied to facilitate the reaction or to avoid unwanted reaction products. The following General Reaction Scheme and Scheme 1 are provided so that the invention might be more fully understood. These examples are illustrative only and should not be constructed as limiting the invention in any way.


GENERAL REACTION SCHEME



5 As illustrated in the General Reaction Scheme, for the compounds of the instant invention, commercially available *p*-bromobenzil **1-1** is treated with a primary or secondary amine to generate **1-2**. (An alternate synthesis of **1-2** can also be found in PCT publication WO2003/086394). Treatment of this material with the known indole-tethered acyl hydrazide **1-3** and excess ammonium acetate in HOAc at 220°C under microwave irradiation delivers, via 1,24-triazine formation and subsequent Diels-Alder reaction, the two regioisomeric unnatural canthine alkaloids, **1-4** and **1-5**, in a 1:1 ratio.

10

SCHEME 1



1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-5) and 1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-6)

1-(4-{[4-(2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl)piperidin-1-yl]methyl}phenyl)-2-phenylethane-1,2-dione (1-3)

To a 100 mL round-bottom flask, equipped with stirrer bar and purged with nitrogen was added *p*-bromomethyl benzil (1-1) (1.0 g, 3.3 mmol) in dry DCM (30 mL). To this was added DIEA (2 mL) and 1-piperidin-4-yl-1,3-dihydro-2*H*-benzimidazol-2-one (1-2) (716 mg, 3.3 mmol) at room temperature and allowed to stir overnight. In the morning, this solution was diluted with DCM and washed with water and 1 N HCl. Concentration of the organics afforded 1.3 g of a bright yellow solid. Analytical LCMS afforded a single peak. *m/z* 440.1.

1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-5) and 1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one (1-6)

To a 5 mL microwave vial was added (1-3) (131 mg, 0.3 mmol), 4-(1*H*-indol-1-yl)butanohydrazide (1-4) [reference for the preparation of 1-4: Lindsley, C.W.; Wisnoski, D.D.; Leister, W.H., Wang, Y.; Zhao, Z. 'A 'One Pot' Microwave-Mediated Synthesis of the Basic Canthine Skeleton: Expedient Access to Unnatural β -Carboline Alkaloids' *Tetrahedron Lett.* 2003, 44, 4495.] (65 mg, 0.3 mmol), ammonium acetate (231 mg, 3.0 mmol, 10 equiv.) in 2.5 mL of glacial acetic acid. The reaction vessel was heated in a single-mode microwave synthesizer at 220 °C for 40 minutes (with almost 16 bar of pressure). After 40 minutes, the instrument rapidly cooled the reaction vessel to 40 °C. The crude sample was immediately injected onto a mass-guided prep HPLC system and the pure regioisomeric canthines 1-5 and 1-6 were separated and purified. Analytical data for (1-5): Analytical LCMS indicates a single peak at 2.176 min., (MeCN/H₂O/0.05%TFA), 4 min. gradient, >99% pure; ¹H NMR (CDCl₃, 600 MHz), δ 9.09 (br s, 1H), 7.72 (t, J=7.78 Hz, 7.73Hz, 1H), 7.62 (d, J=8.48 Hz, 1H), 7.45 (m, 3H), 7.42 (d, J=5.89 Hz, 2H), 7.41 (d, J=5.89 Hz, 2H), 7.39 (obs d, 1H), 7.33 (d, J=8.16 Hz, 1H), 7.31 (t, J=7.61 Hz, 6.98 Hz, 2H), 7.18 (t, J=8.84 Hz, 7.79 Hz, 1H), 7.01 (br m, 3H), 4.60 (br s, 1H), 4.44 (t, J=7.00 Hz, 5.62 Hz, 2H), 4.24 (br s, 2H), 3.71 (br t, 2H), 3.59 (br d, 2H), 2.88 (m, 4H), 2.62 (t, J=5.46 Hz, 6.79 Hz, 2H), 1.94 (br s, 2H); ¹³C NMR (125 MHz, CDCl₃, 25°C) δ 154.5, 143.8, 139.6, 138.8, 134.7, 134.1, 132.5, 131.8, 131.5, 131.2, 131.1, 130.1, 129.7, 129.4, 129.2, 128.4, 127.8, 125.1, 122.2, 122.0, 121.9, 120.9, 110.4, 110.3, 109.9, 60.0, 51.7, 47.3, 41.2, 26.0, 23.7, 21.8; HRMS calc'd for C₃₉H₃₆N₅O (M+H), 590.2914; found 590.2926; Analytical data for (1-6): Analytical LCMS indicates a single peak at 2.389 min., (MeCN/H₂O/0.05%TFA), 4 min. gradient, >99% pure; ¹H NMR (CDCl₃, 600 MHz), δ 8.40 (br s, 1H), 7.72 (t, J=7.95 Hz, 8.1 Hz, 1H), 7.60 (d, 8.79 Hz, 1H), 7.57 (d, 7.35 Hz, 2H), 7.47 (d, J=7.95 Hz, 1H), 7.43 (d, J=7.35 Hz, 2H), 7.36 (d, J=8.21 Hz, 1H), 7.31 (d, J=7.91 Hz, 2H), 7.28 (obs, 1H), 7.23 (t, J=7.60 Hz, 8.97 Hz, 2H), 7.21 (obs, 1H), 7.12 (t, J=7.30 Hz, 1H), 7.09 (t, J=7.90 Hz, 1H), 7.07 (d, J=8.21 Hz, 1H), 4.71 (m, 1H), 4.44 (t, J=5.43 Hz, 2H), 4.33 (s, 2H), 3.73 (t, J=5.56 Hz, 2H), 3.69 (br d, J=11.4 Hz, 2H), 2.98 (q, J=13.36 Hz, 12.65 Hz, 2H), 2.85 (t, J=12.4 Hz, 2H), 2.62 (m, 2H), 2.03, (d, J=13.17 Hz, 2H); ¹³C NMR (125 MHz, DMSO-d₆, 25°C) δ 153.2, 142.0, 140.7, 140.5, 137.0, 131.5, 131.3, 130.2, 129.4, 128.4, 128.0, 127.9, 127.8, 127.5, 125.6, 122.9, 120.6, 120.2, 119.9, 119.5, 110.8, 108.8, 108.2, 58.5, 50.8, 46.5, 40.3, 25.2, 24.7, 21.1; HRMS calc'd for C₃₉H₃₆N₅O (M+H), 590.2914; found 590.2921. The two regioisomers were further confirmed by ROE correlations.

EXAMPLE 1

Cloning of the human Akt isoforms and Δ PH-Akt1

The pS2neo vector (deposited in the ATCC on April 3, 2001 as ATCC PTA-3253) was prepared as follows: The pRmHA3 vector (prepared as described in *Nucl. Acid Res.* 16:1043-1061 (1988)) was cut with BglII and a 2734 bp fragment was isolated. The pUCHsneo vector (prepared as described in *EMBO J.* 4:167-171 (1985)) was also cut with BglII and a 4029 bp band was isolated. These two isolated fragments were ligated together to generate a vector termed pS2neo-1. This plasmid contains a polylinker between a metallothionine promoter and an alcohol dehydrogenase poly A addition

site. It also has a neo resistance gene driven by a heat shock promoter. The pS2neo-1 vector was cut with Psp5II and BsiWI. Two complementary oligonucleotides were synthesized and then annealed (CTGCGGCCGC (SEQ.ID.NO.: 1) and GTACGCGGCCGCAG (SEQ.ID.NO.: 2)). The cut pS2neo-1 and the annealed oligonucleotides were ligated together to generate a second vector, pS2neo. Added in this conversion was a NotI site to aid in the linearization prior to transfection into S2 cells.

Human Akt1 gene was amplified by PCR (Clontech) out of a human spleen cDNA (Clontech) using the 5' primer:

5'CGCGAATTCAGATCTACCATGAGCGACGTGGCTATTGTG 3' (SEQ.ID.NO.: 3), and the 3' primer: 5'CGCTCTAGAGGATCCTCAGGCCGTGCTGCTGGC3' (SEQ.ID.NO.: 4). The 5' primer included an EcoRI and BglII site. The 3' primer included an XbaI and BamHI site for cloning purposes. The resultant PCR product was subcloned into pGEM3Z (Promega) as an EcoRI/Xba I fragment. For expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt1 gene using the PCR primer: 5'GTACGATGCTGAACGATATCTTCG 3' (SEQ.ID.NO.: 5). The resulting PCR product encompassed a 5' KpnI site and a 3' BamHI site which were used to subclone the fragment in frame with a biotin tag containing insect cell expression vector, pS2neo.

For the expression of a pleckstrin homology domain (PH) deleted (Δ aa 4-129, which includes deletion of a portion of the Akt1 hinge region) version of Akt1, PCR deletion mutagenesis was done using the full length Akt1 gene in the pS2neo vector as template. The PCR was carried out in 2 steps using overlapping internal primers

(5'GAATACATGCCGATGGAAAGCGACGGGGCTGAAGAGATGGAGGTG 3' (SEQ.ID.NO.: 6), and 5'CCCCTCCATCTCTTCAGCCCCGTCGCTTTCATCGGCATG

TATTC 3' (SEQ.ID.NO.: 7)) which encompassed the deletion and 5' and 3' flanking primers which encompassed the KpnI site and middle T tag on the 5' end. The final PCR product was digested with KpnI and SmaI and ligated into the pS2neo full length Akt1 KpnI/SmaI cut vector, effectively replacing the 5' end of the clone with the deleted version.

Human Akt3 gene was amplified by PCR of adult brain cDNA (Clontech) using the amino terminal oligo primer:

5' GAATTCAGATCTACCATGAGCGATGTTACCATTTGTG 3' (SEQ.ID.NO.: 8); and the carboxy terminal oligo primer :

5' TCTAGATCTTATTCTCGTCCACTTGCAGAG 3'(SEQ.ID.NO.: 9).

These primers included a 5' EcoRI/BglII site and a 3' XbaI/BglII site for cloning purposes. The resultant PCR product was cloned into the EcoRI and XbaI sites of pGEM4Z (Promega). For expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt3 clone using the PCR primer:

5'GGTACCATGGAATACATGCCGATGGAAAGCGATGTTACCATTTGTGAAG 3'(SEQ.ID.NO.:

10). The resultant PCR product encompassed a 5' KpnI site which allowed in frame cloning with the biotin tag containing insect cell expression vector, pS2neo.

Human Akt2 gene was amplified by PCR from human thymus cDNA (Clontech) using the amino terminal oligo primer:

5' AAGCTTAGATCTACCATGAATGAGGTGTCTGTC 3' (SEQ.ID.NO.: 11); and the carboxy terminal oligo primer: 5'GAATTCGGATCCTCACTCGCGGATGCTGGC 3' (SEQ.ID.NO.: 12).

These primers included a 5' HindIII/BglII site and a 3' EcoRI/BamHI site for cloning purposes. The resultant PCR product was subcloned into the HindIII/EcoRI sites of pGem3Z (Promega). For

expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt2 using the

PCR primer: 5'GGTACCATGGAATACATGCCGATGGAAAATGAGGTGTCTGTTCATCAAAG 3'

(SEQ.ID.NO.: 13). The resultant PCR product was subcloned into the pS2neo vector as described above.

EXAMPLE 2

Expression of human Akt isoforms and ΔPH-Akt1

The DNA containing the cloned Akt1, Akt2, Akt3 and ΔPH-Akt1 genes in the pS2neo

expression vector was purified and used to transfect *Drosophila* S2 cells (ATCC) by the calcium

phosphate method. Pools of antibiotic (G418, 500 μg/ml) resistant cells were selected. Cell were

expanded to a 1.0 L volume (~7.0 x 10⁶ / ml), biotin and CuSO₄ were added to a final concentration of 50 μM and 50 mM respectively. Cells were grown for 72 h at 27°C and harvested by centrifugation. The cell paste was frozen at -70°C until needed.

EXAMPLE 3

Purification of human Akt isoforms and ΔPH-Akt1

Cell paste from one liter of S2 cells, described in Example 2, was lysed by sonication with 50 mls 1% CHAPS in buffer A: (50mM Tris pH 7.4, 1mM EDTA, 1mM EGTA, 0.2mM AEBSF, 10μg/ml benzamidine, 5μg/ml of leupeptin, aprotinin and pepstatin each, 10% glycerol and 1mM DTT).

The soluble fraction was purified on a Protein G Sepharose fast flow (Pharmacia) column loaded with 9mg/ml anti-middle T monoclonal antibody and eluted with 75 μM EYMPME (SEQ.ID.NO.: 14) peptide in buffer A containing 25% glycerol. Akt/PKB containing fractions were pooled and the protein purity evaluated by SDS-PAGE. The purified protein was quantitated using a standard Bradford protocol.

Purified protein was flash frozen on liquid nitrogen and stored at -70°C.

Akt and Akt pleckstrin homology domain deletions purified from S2 cells required activation. Akt and Akt pleckstrin homology domain deletions were activated (Alessi et al. *Current Biology* 7:261-269) in a reaction containing 10 nM PDK1 (Upstate Biotechnology, Inc.), lipid vesicles (10 μM phosphatidylinositol-3,4,5-trisphosphate – Metreya, Inc, 100 μM phosphatidylcholine and 100 μM phosphatidylserine – Avanti Polar lipids, Inc.) and activation buffer (50 mM Tris pH7.4, 1.0 mM DTT, 0.1 mM EGTA, 1.0 μM Microcystin-LR, 0.1 mM ATP, 10 mM MgCl₂, 333 μg/ml BSA and

0.1mM EDTA). The reaction was incubated at 22°C for 4 hours. Aliquots were flash frozen in liquid nitrogen.

EXAMPLE 4

Akt Kinase Assays

Activated Akt isoforms and pleckstrin homology domain deletion constructs were assayed utilizing a GSK-derived biotinylated peptide substrate. The extent of peptide phosphorylation was determined by Homogeneous Time Resolved Fluorescence (HTRF) using a lanthanide chelate(Lance)-coupled monoclonal antibody specific for the phosphopeptide in combination with a streptavidin-linked allophycocyanin (SA-APC) fluorophore which will bind to the biotin moiety on the peptide. When the Lance and APC are in proximity (i.e. bound to the same phosphopeptide molecule), a non-radiative energy transfer takes place from the Lance to the APC, followed by emission of light from APC at 665 nm.

Materials required for the assay:

- A. Activated Akt isozyme or pleckstrin homology domain deleted construct
- 15 B. Akt peptide substrate: GSK3 α (S21) Peptide #3928 biotin-GGRARTSSFAEPG (SEQ.ID.NO.:15), Macromolecular Resources.
- C. Lance labeled anti-phospho GSK3 α monoclonal antibody (Cell Signaling Technology, clone # 27).
- D. SA-APC (Prozyme catalog no. PJ25S lot # 896067).
- 20 E. Microfluor[®]B U Bottom Microtiter Plates (Dynex Technologies, Catalog no. 7205).
- F. Discovery[®] HTRF Microplate Analyzer, Packard Instrument Company.
- G. 100 X Protease Inhibitor Cocktail (PIC): 1 mg/ml benzamidine, 0.5 mg/ml pepstatin, 0.5 mg/ml leupeptin, 0.5 mg/ml aprotinin.
- H. 10X Assay Buffer: 500 mM HEPES, pH 7.5, 1% PEG, mM EDTA, 1 mM EGTA, 1% BSA, 20 mM θ -Glycerol phosphate.
- 25 I. Quench Buffer: 50 mM HEPES pH 7.3, 16.6 mM EDTA, 0.1% BSA, 0.1% Triton X-100, 0.17 nM Lance labeled monoclonal antibody clone # 27, 0.0067 mg/ml SA-APC
- J. ATP/MgCl₂ working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 125 mM KCl, 5% Glycerol, 25 mM MgCl₂, 375 TM ATP
- 30 K. Enzyme working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 5% Glycerol, active Akt. The final enzyme concentrations were selected so that the assay was in a linear response range.
- L. Peptide working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 5% Glycerol, 2 TM GSK3 biotinylated peptide # 3928

The reaction is assembled by adding 16 TL of the ATP/MgCl₂ working solution to the appropriate wells of a 96-well microtiter plate. Inhibitor or vehicle (1.0 Tl) is added followed by 10 Tl of

peptide working solution. The reaction is started by adding 13 μ l of the enzyme working solution and mixing. The reaction is allowed to proceed for 50 min and then stopped by the addition of 60 μ l HTRF quench buffer. The stopped reactions were incubated at room temperature for at least 30 min and then read on the Discovery instrument.

5 Procedure for Streptavidin Flash Plate Assay:

Step 1:

A 1 μ l solution of the test compound in 100% DMSO was added to 20 μ l of 2X substrate solution (20 μ M GSK3 Peptide, 300 μ M ATP, 20 mM MgCl_2 , 20 μ Ci / ml [γ - ^{33}P] ATP, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC, 0.1% BSA and 100 mM KCl). Phosphorylation reactions were
10 initiated by adding 19 μ l of 2X Enzyme solution (6.4 nM active Akt/PKB, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC and 0.1% BSA). The reactions were then incubated at room temperature for 45 minutes.

Step 2:

The reaction was stopped by adding 170 μ l of 125 mM EDTA. 200 μ l of stopped
15 reaction was transferred to a Streptavidin Flashplate[®] PLUS (NEN Life Sciences, catalog no. SMP103). The plate was incubated for ≥ 10 minutes at room temperature on a plate shaker. The contents of each well was aspirated, and the wells rinsed 2 times with 200 μ l TBS per well. The wells were then washed 3 times for 5 minutes with 200 μ l TBS per well with the plates incubated at room temperature on a platform shaker during wash steps.

20 The plates were covered with sealing tape and counted using the Packard TopCount with the appropriate settings for counting [^{33}P] in Flashplates.

Procedure for Streptavidin Filter Plate Assay:

Step 1:

25 The enzymatic reactions as described in Step 1 of the Streptavidin Flash Plate Assay above were performed.

Step 2:

The reaction was stopped by adding 20 μ l of 7.5M Guanidine Hydrochloride. 50 μ l of the stopped reaction was transferred to the Streptavidin filter plate (SAM^{2™} Biotin Capture Plate, Promega, catalog no. V7542) and the reaction was incubated on the filter for 1-2 minutes before applying
30 vacuum.

The plate was then washed using a vacuum manifold as follows: 1) 4 x 200 μ l/well of 2M NaCl; 2) 6 x 200 μ l/well of 2M NaCl with 1% H_3PO_4 ; 3) 2 x 200 μ l/well of diH_2O ; and 4) 2 x 100 μ l/well of 95% Ethanol. The membranes were then allowed to air dry completely before adding scintillant.

The bottom of the plate was sealed with white backing tape, 30 µl/well of Microscint 20 (Packard Instruments, catalog no. 6013621) was added. The top of the plate was sealed with clear sealing tape, and the plate then counted using the Packard TopCount with the appropriate settings for [³³P] with liquid scintillant.

5 Procedure for Phosphocellulose Filter Plate Assay:

Step 1:

The enzymatic reactions were performed as described in Step 1 of the Streptavidin Flash Plate Assay (above) utilizing KKGGRARTSSFAEPG (SEQ.ID.NO.: 16) as the substrate in place of biotin-GGRARTSSFAEPG.

10 Step 2:

The reaction was stopped by adding 20 µl of 0.75% H₃PO₄. 50 µl of stopped reaction was transferred to the filter plate (UNIFILTER™, Whatman P81 Strong Cation Exchanger, White Polystyrene 96 Well Plates, Polyfiltronics, catalog no. 7700-3312) and the reaction incubated on the filter for 1-2 minutes before applying vacuum.

15 The plate was then washed using a vacuum manifold as follows: 1) 9 x 200 µl/well of 0.75% H₃PO₄; and 2) 2 x 200 µl/well of diH₂O. The bottom of the plate was sealed with white backing tape, then 30 µl/well of Microscint 20 was added. The top of the plate was sealed with clear sealing tape, and the plate counted using the Packard TopCount with the appropriate settings for [³³P] and liquid scintillant.

20 PKA assay:

Each individual PKA assay consists of the following components:

- A. 5X PKA assay buffer (200 mM Tris pH7.5, 100 mM MgCl₂, 5mM θ -mercaptoethanol, 0.5 mM EDTA)
- B. 50 µM stock of Kemptide (Sigma) diluted in water
- 25 C. ³³P-ATP prepared by diluting 1.0 µl ³³P-ATP [10 mCi/ml] into 200 Tl of a 50 µM stock of unlabeled ATP
- D. 10 µl of a 70 nM stock of PKA catalytic subunit (UBI catalog # 14-114) diluted in 0.5 mg/ml BSA
- E. PKA/Kemptide working solution: equal volumes of 5X PKA assay buffer, Kemptide
- 30 solution and PKA catalytic subunit.

The reaction is assembled in a 96 deep-well assay plate. The inhibitor or vehicle (10 Tl) is added to 10 Tl of the ³³P-ATP solution. The reaction is initiated by adding 30 Tl of the PKA/Kemptide working solution to each well. The reactions were mixed and incubated at room temperature for 20 min. The reactions were stopped by adding 50 Tl of 100 mM EDTA and 100 mM sodium pyrophosphate and

35 mixing.

The enzyme reaction product (phosphorylated Kemptide) was collected on p81 phosphocellulose 96 well filter plates (Millipore). To prepare the plate, each well of a p81 filter plate was filled with 75 mM phosphoric acid. The wells were emptied through the filter by applying a vacuum to the bottom of the plate. Phosphoric acid (75 mM, 170 μ l) was added to each well. A 30 μ l aliquot from each stopped PKA reaction was added to corresponding wells on the filter plate containing the phosphoric acid. The peptide was trapped on the filter following the application of a vacuum and the filters were washed 5 times with 75 mM phosphoric acid. After the final wash, the filters were allowed to air dry. Scintillation fluid (30 μ l) was added to each well and the filters counted on a TopCount (Packard).

10 PKC assay:

Each PKC assay consists of the following components:

- A. 10X PKC co-activation buffer: 2.5 mM EGTA, 4mM CaCl₂
- B. 5X PKC activation buffer: 1.6 mg/ml phosphatidylserine, 0.16 mg/ml diacylglycerol, 100 mM Tris pH 7.5, 50 mM MgCl₂, 5 mM θ -mercaptoethanol
- 15 C. ³³P-ATP prepared by diluting 1.0 μ l ³³P-ATP [10 mCi/ml] into 100 μ l of a 100 μ M stock of unlabeled ATP
- D. Myelin basic protein (350 μ g/ml, UBI) diluted in water
- E. PKC (50ng/ml, UBI catalog # 14-115) diluted into 0.5 mg/ml BSA
- F. PKC/Myelin Basic Protein working solution: Prepared by mixing 5 volumes each of
- 20 PKC co-activation buffer and Myelin Basic protein with 10 volumes each of PKC activation buffer and PKC.

The assays were assembled in 96 deep-well assay plates. Inhibitor or vehicle (10 TI) was added to 5.0 μ l of ³³P-ATP. Reactions were initiated with the addition of the PKC/Myelin Basic Protein working solution and mixing. Reactions were incubated at 30°C for 20 min. The reactions were stopped by adding 50 TI of 100 mM EDTA and 100 mM sodium pyrophosphate and mixing. Phosphorylated Myelin Basic Protein was collected on PVDF membranes in 96 well filter plates and quantitated by scintillation counting.

Specific compounds of the instant invention were tested in the assay described above and were found to have IC₅₀ of $\leq$ 50 μ M against one or more of Akt1, Akt2 and Akt3.

30 EXAMPLE 5

Cell based Assays to Determine Inhibition of Akt/PKB

Cells (for example LnCaP or a PTEN^(-/-) tumor cell line with activated Akt/PKB) were plated in 100 mM dishes. When the cells were approximately 70 to 80% confluent, the cells were refed with 5 mls of fresh media and the test compound added in solution. Controls included untreated cells, vehicle treated cells and cells treated with either LY294002 (Sigma) or wortmanin (Sigma) at 20 μ M or 200 nM, respectively. The cells were incubated for 2, 4 or 6 hrs, and the media removed. The cells were

washed with PBS, scraped and transferred to a centrifuge tube. They were pelleted and washed again with PBS. Finally, the cell pellet was resuspended in lysis buffer (20 mM Tris pH8, 140 mM NaCl, 2 mM EDTA, 1% Triton, 1 mM Na Pyrophosphate, 10 mM β -Glycerol Phosphate, 10 mM NaF, 0.5 mM NaVO₄, 1 μ M Microcystine, and 1x Protease Inhibitor Cocktail), placed on ice for 15 minutes and gently vortexed to lyse the cells. The lysate was spun in a Beckman tabletop ultra centrifuge at 100,000 x g at 4°C for 20min. The supernatant protein was quantitated by a standard Bradford protocol (BioRad) and stored at -70° C until needed.

Proteins were immunoprecipitated (IP) from cleared lysates as follows: For Akt1/PKB1, lysates are mixed with Santa Cruz sc-7126 (D-17) in NETN (100mM NaCl, 20mM Tris pH 8.0, 1mM EDTA, 0.5% NP-40) and Protein A/G Agarose (Santa Cruz sc-2003) was added. For Akt2/PKB β , lysates were mixed in NETN with anti-Akt2 agarose (Upstate Biotechnology #16-174) and for Akt3/PKBK, lysates were mixed in NETN with anti-Akt3 agarose (Upstate Biotechnology #16-175). The IPs were incubated overnight at 4° C, washed and separated by SDS-PAGE.

Western blots were used to analyze total Akt, pThr308 Akt1, pSer473 Akt1, and corresponding phosphorylation sites on Akt2 and Akt3, and downstream targets of Akt using specific antibodies (Cell Signaling Technology): Anti-Total Akt (cat. no. 9272), Anti-Phospho Akt Serine 473 (cat. no. 9271), and Anti-Phospho Akt Threonine 308 (cat. no. 9275). After incubating with the appropriate primary antibody diluted in PBS + 0.5% non-fat dry milk (NFDM) at 4 °C overnight, blots were washed, incubated with Horseradish peroxidase (HRP)-tagged secondary antibody in PBS + 0.5% NFDM for 1 hour at room temperature. Proteins were detected with ECL Reagents (Amersham/Pharmacia Biotech RPN2134).

EXAMPLE 6

Heregulin Stimulated Akt Activation

MCF7 cells (a human breast cancer line that is PTEN^{+/+}) were plated at 1x10⁶ cells per 100mM plate. When the cells were 70 – 80% confluent, they were refed with 5 ml of serum free media and incubated overnight. The following morning, compound was added and the cells were incubated for 1 – 2 hrs, after which time heregulin was added (to induce the activation of Akt) for 30 minutes and the cells were analyzed as described above.

EXAMPLE 7

Inhibition Of Tumor Growth


In vivo efficacy of an inhibitor of the growth of cancer cells may be confirmed by several protocols well known in the art.

Human tumor cell lines which exhibit a deregulation of the PI3K pathway (such as LnCaP, PC3, C33a, OVCAR-3, MDA-MB-468 or the like) are injected subcutaneously into the left flank of 6-10 week old female nude mice (Harlan) on day 0. The mice are randomly assigned to a vehicle, compound or combination treatment group. Daily subcutaneous

administration begins on day 1 and continues for the duration of the experiment. Alternatively, the inhibitor test compound may be administered by a continuous infusion pump. Compound, compound combination or vehicle is delivered in a total volume of 0.2 ml. Tumors are excised and weighed when all of the vehicle-treated animals exhibited lesions of 0.5 - 1.0 cm in diameter, typically 4 to 5.5 weeks after the cells were injected. The average weight of the tumors in each treatment group for each cell line is calculated.

WHAT IS CLAIMED IS:

1. A compound of the Formula A:



5 wherein:

a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is independently 0, 1, 2, 3 or 4; p is independently 0, 1, 2, 3, 4 or 5; r is 0 or 1; s is 0 or 1; and t is 2, 3, 4, 5 or 6;

10  is selected from: C₃-C₈ cycloalkyl, aryl, heteroaryl and heterocyclyl;

R¹ is independently selected from: (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR^c(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR^cS(O)_mR^a, oxo,
15 CHO, NO₂, NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R^z;

R² is independently selected from: (C₁-C₆)alkyl-heterocyclyl, (C₁-C₆)alkyl-NR⁵R⁶, (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, (C=O)_aO_bC₃-C₈ cycloalkyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁵R⁶, NR^c(C=O)NR⁵R⁶, S(O)_mR^a, S(O)₂NR⁵R⁶, NR^cS(O)_mR^a, oxo, CHO, NO₂, NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected
25 from R^z;

R⁵ and R⁶ are independently selected from: H, (C=O)O_bR^a, C₁-C₁₀ alkyl, aryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, heterocyclyl, C₃-C₈ cycloalkyl, SO₂R^a, and (C=O)NR^b₂, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R^Z, or R⁵ and R⁶ can be taken together with the nitrogen to which they are attached to form a
 5 monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R^Z;

R^Z is selected from: (C=O)_rO_s(C₁-C₁₀)alkyl, O_r(C₁-C₃)perfluoroalkyl, (C₀-C₆)alkylene-S(O)_mR^a, oxo,
 10 OH, halo, CN, (C=O)_rO_s(C₂-C₁₀)alkenyl, (C=O)_rO_s(C₂-C₁₀)alkynyl, (C=O)_rO_s(C₃-C₆)cycloalkyl, (C=O)_rO_s(C₀-C₆)alkylene-aryl, (C=O)_rO_s(C₀-C₆)alkylene-heterocyclyl, (C=O)_rO_s(C₀-C₆)alkylene-N(R^b)₂, C(O)R^a, (C₀-C₆)alkylene-CO₂R^a, C(O)H, (C₀-C₆)alkylene-CO₂H, C(O)N(R^b)₂, S(O)_mR^a, S(O)₂N(R^b)₂NR^c(C=O)O_bR^a, O(C=O)O_bC₁-C₁₀ alkyl, O(C=O)O_bC₃-C₈ cycloalkyl, O(C=O)O_baryl, and O(C=O)O_b-heterocycle, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally
 15 substituted with up to three substituents selected from R^b, OH, (C₁-C₆)alkoxy, halogen, aryl, heterocyclyl, CO₂H, CN, O(C=O)C₁-C₆ alkyl, oxo, and N(R^b)₂, wherein said heterocyclyl is optionally substituted with from one to three substituents selected from oxo, OH, N(R^d)₂, and -O(C₁-C₆)alkyl;

R^a is: substituted or unsubstituted (C₁-C₁₀)alkyl, substituted or unsubstituted (C₂-C₁₀)alkenyl,
 20 substituted or unsubstituted (C₂-C₁₀)alkynyl, substituted or unsubstituted (C₃-C₁₀)cycloalkyl, substituted or unsubstituted aryl, (C₁-C₆)perfluoroalkyl, 2,2,2-trifluoroethyl, or substituted or unsubstituted heterocyclyl;


R^b is: H, (C₁-C₁₀)alkyl, substituted or unsubstituted aryl, substituted or unsubstituted benzyl,
 25 substituted or unsubstituted heterocyclyl, (C₃-C₁₀)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl or S(O)₂R^a;

R^c is selected from: H, C₁-C₁₀ alkyl, aryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, heterocyclyl, C₃-C₁₀ cycloalkyl, C₁-C₆ perfluoroalkyl, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is
 30 optionally substituted with one or more substituents selected from R^Z, and

R^d is independently selected from: H and (C₁-C₆)alkyl;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

2. A compound according to Claim 1 of the Formula A5:




wherein:

- 5 Q is selected from: heterocyclyl, said heterocyclyl optionally substituted with one to three substituents selected from R^Z;

and all other substituents and variables are as defined in Claim 1;

- 10 or a pharmaceutically acceptable salt or a stereoisomer thereof.

3. A compound according to Claim 1 of the Formula A9:



15 wherein:

Q is selected from: heterocyclyl, said heterocyclyl optionally substituted with one to three substituents selected from R^Z;

20 and all other substituents and variables are as defined in Claim 1;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

4. A compound which is selected from:

1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one; and

1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

5. A TFA salt of a compound according to Claim 1 which is selected from:

1-{1-[4-(1-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-2-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one; and

1-{1-[4-(2-phenyl-5,6-dihydro-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-1-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2*H*-benzimidazol-2-one;

or a stereoisomer thereof.

6. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

7. The use of the compound according to Claim 1 for the preparation of a medicament useful in the treatment or prevention of cancer in a mammal in need of such treatment.

SEQUENCE LISTING

<110> Merck & Co., Inc
Barnett, Stanley F.
Bogusky, Michael J.
Leister, William H.
Lindsley, Craig W.

<120> INHIBITORS OF AKT ACTIVITY

<130> 21821

<150> 60/632,490

<151> 2004-12-02

<160> 16

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely synthetic DNA Sequence

<400> 1

ctgcggccgc

10

<210> 2

<211> 14

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely synthetic DNA Sequence

<400> 2

gtacgcggcc gcag

14

<210> 3

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely synthetic DNA Sequence

<400> 3

cgcgaattca gatctaccat gagcgacgtg gctattgtg

39

<210> 4

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely synthetic DNA Sequence

<400> 4
cgctctagag gatcctcagg ccgtgctgct ggc 33

<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 5
gtacgatgct gaacgatatc ttcg 24

<210> 6
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 6
gaatacatgc cgatggaaag cgacggggct gaagagatgg aggtg 45

<210> 7
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 7
cccctccatc tcttcagccc cgtcgctttc catcggcattg tattc 45

<210> 8
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 8
gaattcagat ctacatgag cgatggtacc attgtg 36

<210> 9
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 9
tctagatctt attctcgtcc acttgacagag 30

<210> 10
<211> 48

<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 10
ggtaccatgg aatacatgcc gatggaaagc gatgttacca ttgtgaag 48

<210> 11
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 11
aagcttagat ctaccatgaa tgaggtgtct gtc 33

<210> 12
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 12
gaattcggat cctcactcgc ggatgctggc 30

<210> 13
<211> 49
<212> DNA
<213> Artificial Sequence

<220>
<223> Completely synthetic DNA Sequence

<400> 13
ggtaccatgg aatacatgcc gatggaaaat gaggtgtctg tcatcaaag 49

<210> 14
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Completely synthetic Amino Acid Sequence

<400> 14
Glu Tyr Met Pro Met Glu
1 5

<210> 15
<211> 13
<212> PRT
<213> Artificial Sequence

<220>

<223> Completely synthetic Amino Acid Sequence

<400> 15

Gly Gly Arg Ala Arg Thr Ser Ser Phe Ala Glu Pro Gly
1 5 10

<210> 16

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely synthetic Amino Acid Sequence

<400> 16

Lys Lys Gly Gly Arg Ala Arg Thr Ser Ser Phe Ala Glu Pro Gly
1 5 10 15