



US 20170319711A1

(19) **United States**(12) **Patent Application Publication**  
**Gerber et al.**(10) **Pub. No.: US 2017/0319711 A1**(43) **Pub. Date: Nov. 9, 2017**(54) **ANTIBODY-DRUG CONJUGATES**(71) Applicant: **WYETH LLC**, Madison, NJ (US)(72) Inventors: **Hans-Peter Gerber**, Montclair, NJ (US); **John Francis DiJoseph**, Woodbridge, NJ (US); **Kiran Manohar Khandke**, Nanuet, NY (US); **Kimberly Ann Marquette**, Somerville, MA (US); **Puja Sapra**, River Edge, NJ (US); **Lioudmila Gennadievna Tchistiakova**, Stoneham, MA (US)(73) Assignee: **WYETH LLC**, Madison, NJ (US)(21) Appl. No.: **15/161,972**(22) Filed: **May 23, 2016****Related U.S. Application Data**

(63) Continuation of application No. 14/052,942, filed on Oct. 14, 2013, now abandoned, which is a continuation of application No. 13/616,030, filed on Sep. 14, 2012, now Pat. No. 8,586,049, which is a continu-

ation of application No. 13/435,731, filed on Mar. 30, 2012, now Pat. No. 8,309,094.

(60) Provisional application No. 61/602,349, filed on Feb. 23, 2012, provisional application No. 61/593,549, filed on Feb. 1, 2012, provisional application No. 61/470,576, filed on Apr. 1, 2011.

**Publication Classification**(51) **Int. Cl.****A61K 47/68** (2006.01)**C07K 16/30** (2006.01)**A61K 47/68** (2006.01)**A61K 39/00** (2006.01)**A61K 39/00** (2006.01)(52) **U.S. Cl.**CPC ..... **A61K 47/6851** (2017.08); **C07K 16/30**(2013.01); **A61K 47/6803** (2017.08); **A61K****2039/505** (2013.01); **A61K 2039/545**(2013.01); **C07K 2317/732** (2013.01); **C07K****2317/92** (2013.01)

(57)

**ABSTRACT**

Disclosed are anti-5T4 antibody drug conjugates and methods for preparing and using the same.

## ANTIBODY-DRUG CONJUGATES

[0001] This application claims priority from copending provisional application Nos. 61/602,349 filed Feb. 23, 2012, 61/593,549 filed Feb. 1, 2012 and 61/470,576 filed Apr. 1, 2011 the entire disclosure of which is hereby incorporated by reference.

## FIELD

[0002] The present invention generally relates to anti-5T4 antibody-drug conjugates for the treatment of cancer.

## BACKGROUND

[0003] Antibody-drug conjugates (ADCs) combine the binding specificity of monoclonal antibodies with the potency of chemotherapeutic agents. The technology associated with the development of monoclonal antibodies to tumor associated target molecules, the use of more effective cytotoxic agents, and the design of chemical linkers to covalently bind these components, has progressed rapidly in recent years (Ducry L., et al. Bioconjugate Chemistry, 21:5-13, 2010).

[0004] Promising ADCs such as SGN-75 (U.S.2009/148942) and trastuzumab-DM1 (U.S.2009/0226465) are currently in clinical trials. However, as other tumor associated antigens are considered for targets, numerous challenges remain. Each monoclonal antibody must be characterized separately, an appropriate linker designed, and a suitable cytotoxic agent identified that retains its potency upon delivery to tumor cells. One must consider the antigen density on the cancer target and whether normal tissues express the target antigen. Other considerations include whether the entire ADC is internalized upon binding the target; whether a cytostatic or cytotoxic drug is preferable when considering possible normal tissue exposure and/or the type and stage of the cancer being treated; and, whether the linker connecting the antibody to the drug payload is a cleavable or a non-cleavable linkage. Furthermore, the antibody to drug moiety conjugation ratio must be sufficient without compromising the binding activity of the antibody and/or the potency of the drug. It is evident that ADCs are complex biologics and the challenges to develop an effective ADC remain significant.

[0005] The human 5T4 tumor associated antigen is the target antigen of the present invention. It has recently been shown that the 5T4 antigen is expressed in high levels on certain highly tumorigenic cells, also called tumor-initiating cells (WO2010/111659). Tumor-initiating cells show resistance to standard therapies and are believed to be responsible for tumor recurrence and metastasis and therefore present yet another obstacle for ADC development.

[0006] The novel anti-5T4 ADCs of the present invention overcome the challenges associated with ADC technology and provide highly specific and potent ADCs that bind to tumor cells expressing the 5T4 antigen and deliver sufficient cytotoxic drug to the cells, thus providing an innovative and effective treatment for cancer.

## SUMMARY

[0007] In one embodiment, an antibody-drug conjugate of the present invention has the formula: Ab-(LU-D)<sub>p</sub> or a pharmaceutically acceptable salt thereof wherein, Ab is an anti-5T4 antibody or antigen binding portion thereof, com-

prising a heavy chain variable region having a VH CDR1 region as shown in SEQ ID NO: 5, a VH CDR2 region as shown in SEQ ID NO: 6, and a VH CDR3 region as shown in SEQ ID NO: 7; LU is a linker unit selected from the group consisting of maleimidocaproyl and maleimidocaproyl-Val-Cit-PABA; p is an integer from about 1 to about 8; and D is a Drug unit selected from the group consisting of MMAE, MMAF, and MMAD.

[0008] The present invention further provides anti-5T4 antibody-drug conjugates wherein said anti-5T4 antibody or antigen binding portion thereof, comprises a heavy chain variable region having (a) a VH CDR1 region as shown in SEQ ID NO: 5, (b) a VH CDR2 region as shown in SEQ ID NO: 6, and (c) a VH CDR3 region as shown in SEQ ID NO: 7.

[0009] The present invention further provides an anti-5T4 antibody-drug conjugate wherein said anti-5T4 antibody or antigen binding portion thereof, comprises a light chain variable region having (a) a VL CDR1 region as shown in SEQ ID NO: 8, (b) a VL CDR2 region as shown in SEQ ID NO: 9, and (c) a VL CDR3 region as shown in SEQ ID NO: 10.

[0010] The present invention further provides an anti-5T4 antibody-drug conjugate wherein said anti-5T4 antibody or antigen binding portion thereof, further comprises a heavy chain variable region having (a) a VH CDR1 region as shown in SEQ ID NO: 5, (b) a VH CDR2 region as shown in SEQ ID NO: 6, and (c) a VH CDR3 region as shown in SEQ ID NO: 7 and a light chain variable region having (a) a VL CDR1 region as shown in SEQ ID NO: 8, (b) a VL CDR2 region as shown in SEQ ID NO: 9, and (c) a VL CDR3 region as shown in SEQ ID NO: 10.

[0011] The present invention further provides an anti-5T4 antibody-drug conjugate wherein said anti-5T4 antibody or antigen binding portion thereof, comprises the VH region of SEQ ID NO: 3 and the VL region of SEQ ID NO: 4.

[0012] The present invention further provides an anti-5T4 antibody-drug conjugate wherein said anti-5T4 antibody consists of a heavy chain having SEQ ID NO: 1 and a light chain having SEQ ID NO: 2.

[0013] The present invention further provides an anti-5T4 antibody-drug conjugate wherein:

[0014] (a) said anti-5T4 antibody consists of a heavy chain having SEQ ID NO:1 and a light chain having SEQ ID NO: 2, (b) said LU is maleimidocaproyl, (c) said Drug is MMAF, and (d) p is an integer of about 4.

[0015] The present invention further provides an anti-5T4 antibody-drug conjugate wherein:

[0016] (a) said anti-5T4 antibody consists of a heavy chain having SEQ ID NO:1 and a light chain having SEQ ID NO: 2, (b) said LU is maleimidocaproyl-Val-Cit-PABA, (c) said Drug is MMAE, and (d) p is an integer of about 4.

[0017] The present invention further provides an anti-5T4 antibody-drug conjugate wherein:

[0018] (a) said anti-5T4 antibody consists of a heavy chain having SEQ ID NO:1 and a light chain having SEQ ID NO: 2, (b) said LU is maleimidocaproyl-Val-Cit-PABA, (c) said Drug is MMAD, and (d) p is an integer from about 1 to about 8.

[0019] The present invention further provides an anti-5T4 antibody-drug conjugate wherein: (a) said anti-5T4 antibody consists of a heavy chain having SEQ ID NO:15 and a light chain having SEQ ID NO: 2, (b) said LU is maleimido-

caproyl-Val-Cit-PABA, (c) said Drug is MMAE, and (d) p is an integer of about 1 to about 8.

**[0020]** The present invention provides an anti-5T4 antibody-drug conjugate wherein said antibody recognizes an epitope on human 5T4 antigen wherein said epitope comprises amino acid residues 173-258 and 282-361 of the amino acid sequence of SEQ ID NO: 11.

**[0021]** The present invention provides a pharmaceutical composition comprising an antibody-drug conjugate indicated above and a pharmaceutically acceptable carrier.

**[0022]** The present invention further provides a method of treating a 5T4-positive cancer in a patient in need thereof, comprising administering to said patient an antibody-drug conjugate indicated above.

**[0023]** The present invention further provides a method of treating a 5T4-positive cancer wherein said cancer is selected from the group consisting of carcinomas of the bladder, breast, cervix, colon, endometrium, kidney, lung, esophagus, ovary, prostate, pancreas, liver, skin, stomach, and testes.

**[0024]** More preferably, the present invention provides a method of treating a 5T4-positive cancer wherein said cancer is selected from the group consisting of colorectal, breast, pancreatic, and non-small cell lung carcinomas.

**[0025]** The invention further provides an antibody-drug conjugate indicated above for use in therapy. The invention further provides the use of an antibody-drug conjugate indicated above for the manufacture of a medicament.

**[0026]** The invention further provides the use indicated above, wherein said use is for the treatment of a 5T4-positive cancer and wherein said cancer is selected from the group consisting of carcinomas of the bladder, breast, cervix, endometrium, kidney, lung, esophagus, ovary, prostate, pancreas, skin, stomach, and testes.

**[0027]** More preferably, the invention further provides the use indicated above, wherein said use is for the treatment of a 5T4-positive cancer wherein said cancer is selected from the group consisting of colorectal, breast, pancreatic, and non-small cell lung carcinomas.

**[0028]** The invention further provides a nucleic acid that encodes an anti-5T4 antibody, a vector comprising said nucleic acid, and a host cell comprising said vector.

**[0029]** The invention further provides a process for producing an anti-5T4 antibody comprising cultivating the host cell comprising the above mentioned vector and recovering the antibody from the cell culture.

**[0030]** The invention further provides a process for producing an anti-5T4 antibody-drug conjugate comprising: (a) taking the antibody recovered from the cell culture, (b) chemically linking said antibody via a linker unit selected from the group consisting of maleimidocaproyl or maleimidocaproyl-Val-Cit to a Drug unit selected from the group consisting of MMAE, MMAD, or MMAF, and (c) purifying the antibody-drug conjugate.

#### DETAILED DESCRIPTION

**[0031]** The present invention provides anti-5T4 antibody-drug conjugates for the treatment of cancer. In order that the present invention is more readily understood, certain terms are first defined.

**[0032]** All amino acid abbreviations used in this disclosure are those accepted by the United States Patent and Trademark Office as set forth in 37 C.F.R. §1.822 (B)(I).

**[0033]** 5T4 refers to the 5T4 oncofetal antigen, a 72 kDa highly glycosylated transmembrane glycoprotein comprising a 42 kDa non-glycosylated core (see U.S. Pat. No. 5,869,053). Human 5T4 is expressed in numerous cancer types, including carcinomas of the bladder, breast, cervix, colon, endometrium, kidney, lung, esophagus, ovary, prostate, pancreas, liver, skin, stomach, and testes. Highly tumorigenic cells, also called cancer stem cells or tumor-initiating cells have been shown to have high levels of 5T4 expression (W02010/111659). Anti-5T4 antibodies of the invention include antibodies that specifically bind the human 5T4 antigen (see U.S. 2007/0231333).

**[0034]** An “antibody” is an immunoglobulin molecule capable of specific binding to a target, such as a carbohydrate, polynucleotide, lipid, polypeptide, etc., through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule. As used herein, the term “antibody” encompasses not only intact polyclonal or monoclonal antibodies, but also any antigen binding fragment (i.e., “antigen-binding portion”) or single chain thereof, fusion proteins comprising an antibody, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site including, for example without limitation, Fab, Fab', F(ab')<sub>2</sub>, an Fd fragment consisting of the VH and CH1 domains, an Fv fragment consisting of the VL and VH domains of a single arm of an antibody, an isolated complementarity determining region (CDR), scFv, single domain antibodies (e.g., shark and camelid antibodies), maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv.

**[0035]** An antibody includes an antibody of any class, such as IgG, IgA, or IgM (or sub-class thereof), and the antibody need not be of any particular class. Depending on the antibody amino acid sequence of the constant region of its heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2. The heavy-chain constant regions that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

**[0036]** A “variable region” of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. As known in the art, the variable regions of the heavy and light chain each consist of four framework regions (FRs) connected by three complementarity determining regions (CDRs) also known as hypervariable regions, contribute to the formation of the antigen binding site of antibodies. If variants of a subject variable region are desired, particularly with substitution in amino acid residues outside of a CDR region (i.e., in the framework region), appropriate amino acid substitution, preferably, conservative amino acid substitution, can be identified by comparing the subject variable region to the variable regions of other antibodies which contain CDR1 and CDR2 sequences in the same canonical class as the subject variable region (Chothia and Lesk, J Mol Biol 196 (4): 901-917, 1987). When choosing FR to flank subject CDRs, e.g., when

humanizing or optimizing an antibody, FRs from antibodies which contain CDR1 and CDR2 sequences in the same canonical class are preferred.

**[0037]** A “CDR” of a variable domain are amino acid residues within the variable region that are identified in accordance with the definitions of the Kabat, Chothia, the cumulation of both Kabat and Chothia, AbM, contact, and/or conformational definitions or any method of CDR determination well known in the art. Antibody CDRs may be identified as the hypervariable regions originally defined by Kabat et al. See, e.g., Kabat et al., 1992, Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, NIH, Washington, D.C. The positions of the CDRs may also be identified as the structural loop structures originally described by Chothia and others. See, e.g., Chothia et al., 1989, Nature 342:877-883. Other approaches to CDR identification include the “AbM definition,” which is a compromise between Kabat and Chothia and is derived using Oxford Molecular’s AbM antibody modeling software (now Accelrys®), or the “contact definition” of CDRs based on observed antigen contacts, set forth in MacCallum et al., 1996, J. Mol. Biol., 262:732-745. In another approach, referred to herein as the “conformational definition” of CDRs, the positions of the CDRs may be identified as the residues that make enthalpic contributions to antigen binding. See, e.g., Makabe et al., 2008, Journal of Biological Chemistry, 283:1156-1166. Still other CDR boundary definitions may not strictly follow one of the above approaches, but will nonetheless overlap with at least a portion of the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. As used herein, a CDR may refer to CDRs defined by any approach known in the art, including combinations of approaches. The methods used herein may utilize CDRs defined according to any of

derived from non-human immunoglobulin. Preferably, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity.

**[0040]** The term “chimeric antibody” is intended to refer to antibodies in which the variable region sequences are derived from one species and the constant region sequences are derived from another species, such as an antibody in which the variable region sequences are derived from a mouse antibody and the constant region sequences are derived from a human antibody.

**[0041]** Antibodies of the invention can be produced using techniques well known in the art, e.g., recombinant technologies, phage display technologies, synthetic technologies or combinations of such technologies or other technologies readily known in the art (see, for example, Jayasena, S. D., Clin. Chem., 45: 1628-50 (1999) and Fellouse, F. A., et al, J. Mol. Biol., 373 (4):924-40 (2007)).

**[0042]** Tables 1 and 2 below depict preferred CDRs for the antibodies of the present invention.

TABLE 1

Anti-body	LCDR1	LCDR2	LCDR3
A1	KASQSVSNDVA SEQ ID NO: 8	FATNRYT SEQ ID NO: 9	QQDYSSPWT SEQ ID NO: 10
A3	KASQDVDTAVA SEQ ID NO: 17	WASTRLT SEQ ID NO: 18	QQYSSYPYT SEQ ID NO: 19

TABLE 2

Anti-body	HCDR1	HCDR2	HCDR3
A1	NFGMN SEQ ID NO: 5	WINTNTGEPYAEFPKG SEQ ID NO: 6	DWDGAYFPDY SEQ ID NO: 7
A1-IgG4	GYTFTNFGMN SEQ ID NO: 14	WINTNTGEPYAEFPKG SEQ ID NO: 6	DWDGAYFPDY SEQ ID NO: 7
A3	TYAMN SEQ ID NO: 22	RIRSKSNYYATYYADSVKD SEQ ID NO: 23	QWDYDVRAMNY SEQ ID NO: 24

these approaches. For any given embodiment containing more than one CDR, the CDRs may be defined in accordance with any of Kabat, Chothia, extended, AbM, contact, and/or conformational definitions.

**[0038]** The term “monoclonal antibody” (Mab) refers to an antibody that is derived from a single copy or clone, including e.g., any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Preferably, a monoclonal antibody of the invention exists in a homogeneous or substantially homogeneous population.

**[0039]** “Humanized” antibody refers to forms of non-human (e.g. murine) antibodies that are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) that contain minimal sequence

**[0043]** The present invention includes an antibody or antigen binding portion thereof, that comprises:

**[0044]** a) a light chain variable region comprising:

**[0045]** i) a LCDR1 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 8 and 17;

**[0046]** ii) a LCDR2 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 9 and 18; and

**[0047]** iii) a LCDR3 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 10 and 19; and

**[0048]** b) a heavy chain variable region comprising:

**[0049]** i) a HCDR1 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5 and 22;

**[0050]** ii) a HCDR2 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 6 and 23; and p1 iii) a LCDR1 having an amino acid sequence selected from the group consisting of SEQ ID NOs: 7 and 24.

**[0051]** A preferred antibody or antigen binding portion thereof, of the invention comprises:

**[0052]** a) a LCVR comprising: a LCDR1 of SEQ ID NO: 8, a LCDR2 of SEQ ID NO: 9, and a LCDR3 of SEQ ID NO: 10; and

**[0053]** b) a HCVR comprising: a HCDR1 of SEQ ID NO: 5, a HCDR2 of SEQ ID NO: 6, and a HCDR3 of SEQ ID NO: 7.

**[0054]** Preferred monoclonal antibodies of the invention are referred to herein as A1 (a humanized anti-5T4 IgG1 antibody); A1-IgG4 (a humanized anti-5T4 IgG4 antibody); A3 (a mouse/human chimeric antibody); and A3hu (a humanized anti-5T4 IgG1 antibody). The SEQ ID NOs of the amino acid sequences encoding Mabs A1, A1-IgG4 and A3 are provided in Table 3 below:

TABLE 3

Mab	LC	HC	LCVR	LCDR1	LCDR2	LCDR3	HCVR	HCDR1	HCDR2	HCDR3
A1	2	1	4	8	9	10	3	5	6	7
A1-IgG4	2	12	4	8	9	10	13	5	6	7
A3	2	15	21	22	23	24	16	17	18	19
A3hu	30	25	31	32	33	34	26	27	28	29

**[0055]** The phrases “an antibody recognizing an antigen” and “an antibody specific for an antigen” are used interchangeably herein with the term “an antibody which binds specifically to an antigen.”

**[0056]** Anti-5T4 Antibody-Drug conjugate refers to an anti-5T4 antibody or antigen binding portion thereof, as described herein linked to a cytotoxic drug moiety (D) via a linker unit molecule (LU).

**[0057]** Linker Unit (LU): LU describes the direct or indirect linkage of the antibody to the drug. Attachment of a linker to a mAb can be accomplished in a variety of ways, such as through surface lysines, reductive-coupling to oxidized carbohydrates, and through cysteine residues liberated by reducing interchain disulfide linkages. A variety of ADC linkage systems are known in the art, including hydrazone-, disulfide- and peptide-based linkages.

**[0058]** Drug (D): A drug is any substance having biological or detectable activity, for example, therapeutic agents, detectable labels, binding agents, etc., and prodrugs, which are metabolized to an active agent in vivo. The terms drug and payload are used interchangeably. In some embodiments, the Drug is an auristatin, such as auristatin E (also known in the art as a derivative of dolastatin-10) or a derivative thereof. The auristatin can be, for example, an ester formed between auristatin E and a keto acid. For example, auristatin E can be reacted with paraacetyl benzoic acid or benzoylvaleric acid to produce AEB and AEVB, respectively. Other typical auristatins include AFP, MMAF, and MMAE. The synthesis and structure of exemplary auristatins are described in U.S. Pat. Nos. 6,884,869, 7,098,308, 7,256,257, 7,423,116, 7,498,298 and 7,745,394, each of which is incorporated by reference herein in its entirety and for all purposes.

**[0059]** Auristatins have been shown to interfere with microtubule dynamics and nuclear and cellular division and have anticancer activity. Auristatins of the present invention bind tubulin and can exert a cytotoxic or cytostatic effect on a 5T4 expressing cell or cell line. There are a number of different assays, known in the art, that can be used for determining whether an auristatin or resultant antibody-drug conjugate exerts a cytostatic or cytotoxic effect on a desired cell or cell line. Methods for determining whether a compound binds tubulin are known in the art. See, for example, Muller et al., *Anal. Chem* 2006, 78, 4390-4397; Hamel et al., *Molecular Pharmacology*, 1995 47: 965-976; and Hamel et al., *The Journal of Biological Chemistry*, 1990 265:28, 17141-17149.

**[0060]** Examples of drugs or payloads are selected from the group consisting of DM1 (maytansine, N2'-deacetyl-N2'-(3-mercapto-1-oxopropyl)- or N2'-deacetyl-N2'-(3-mercapto-1-oxopropyl)-maytansine), mc-MMAD (6-maleimidocaproyl-monomethylauristatin-D or N-methyl-L-valyl-N-[(1S,2R)-2-methoxy-4-[(2S)-2-[(1R,2R)-1-methoxy-2-

methyl-3-oxo-3-[[[(1S)-2-phenyl-1-(2-thiazolyl)ethyl]amino]propyl]-1-pyrrolidinyl]-1-[(1S)-1-methylpropyl]-4-oxobutyl]-N-methyl-(9CI)-L-valinamide), mc-MMAF (maleimidocaproyl-monomethylauristatin F or N-[6-(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-1-oxohexyl]-N-methyl-L-valyl-L-valyl-(3R,4S,5S)-3-methoxy-5-methyl-4-(methylamino)heptanoyl-( $\alpha$ R, $\beta$ R,2S)- $\beta$ -(3-methoxy- $\alpha$ -methyl-2-pyrrolidinepropanoyl-L-phenylalanine) and mc-Val-Cit-PABA-MMAE (6-maleimidocaproyl-ValCit-(p-aminobenzyloxycarbonyl)-monomethylauristatin E or N-[[[4-[[N-[6-(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-1-oxohexyl]-L-valyl-N5-(aminocarbonyl)-L-ornithyl]amino]phenyl]methoxy]carbonyl]-N-methyl-L-valyl-N-[(1S,2R)-4-[(2S)-2-[(1R,2R)-3-[(1R,2S)-2-hydroxy-1-methyl-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]-1-pyrrolidinyl]-2-methoxy-1-[(1S)-1-methylpropyl]-4-oxobutyl]-N-methyl-L-valinamide). DM1 is a derivative of the tubulin inhibitor maytansine while MMAD, MMAE, and MMAF are auristatin derivatives. The preferred payloads of the present invention are selected from the group consisting of mc-MMAF and mc-Val-Cit-PABA-MMAE.

**[0061]** The term “epitope” refers to that portion of a molecule capable of being recognized by and bound by an antibody at one or more of the antibody's antigen-binding regions. Epitopes often consist of a chemically active surface grouping of molecules such as amino acids or sugar side chains and have specific three-dimensional structural characteristics as well as specific charge characteristics. The term “antigenic epitope” as used herein, is defined as a portion of a polypeptide to which an antibody can specifically bind as determined by any method well known in the art, for example, by conventional immunoassays. A “non-linear epitope” or “conformational epitope” comprises non-

contiguous polypeptides (or amino acids) within the antigenic protein to which an antibody specific to the epitope binds.

**[0062]** The term “binding affinity ( $K_D$ )” as used herein, is intended to refer to the dissociation rate of a particular antigen-antibody interaction. The  $K_D$  is the ratio of the rate of dissociation, also called the “off-rate ( $k_{off}$ )”, to the association rate, or “on-rate ( $k_{on}$ )”. Thus,  $K_D$  equals  $k_{off}/k_{on}$  and is expressed as a molar concentration (M). It follows that the smaller the  $K_D$ , the stronger the affinity of binding. Therefore, a  $K_D$  of 1  $\mu$ M indicates weak binding affinity compared to a  $K_D$  of nM.  $K_D$  values for antibodies can be determined using methods well established in the art. One method for determining the  $K_D$  of an antibody is by using surface plasmon resonance (SPR), typically using a biosensor system such as a Biacore® system.

**[0063]** The term “specifically binds” as used herein in reference to the binding between an antibody and a 5T4 antigen and the antibody binds the 5T4 antigen with a  $K_D$  less than about 30 nM as determined by SPR at 25° C.

**[0064]** Pharmaceutically acceptable salt as used herein refers to pharmaceutically acceptable organic or inorganic salts of a molecule or macromolecule.

**[0065]** The term “potency” is a measurement of biological activity and may be designated as  $I_{50}$ , or effective concentration of antibody needed to inhibit 50% of growth of a 5T4 positive cell line as described in Example 3. Alternatively, potency may refer to anti-tumor activity as determined in an in vivo tumor xenograph model as shown in Example 4.

**[0066]** The terms “polynucleotide” or “nucleic acid molecule”, as used herein, are intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

**[0067]** The polynucleotides that encode the antibodies of the present invention may include the following: only the coding sequence for the variant, the coding sequence for the variant and additional coding sequences such as a functional polypeptide, or a signal or secretory sequence or a pro-protein sequence; the coding sequence for the antibody and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the antibody. The term “polynucleotide encoding an antibody” encompasses a polynucleotide which includes additional coding sequence for the variant but also a polynucleotide which includes additional coding and/or non-coding sequence. It is known in the art that a polynucleotide sequence that is optimized for a specific host cell/expression system can readily be obtained from the amino acid sequence of the desired protein (see GENEART® AG, Regensburg, Germany).

**[0068]** The polynucleotides encoding the antibodies of the present invention will typically include an expression control polynucleotide sequence operably linked to the antibody coding sequences, including naturally-associated or heterologous promoter regions known in the art. Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host cell line, the host cell is propagated under conditions suitable for expressing the nucleotide sequences, and, as desired, for the collection and purification of the antibodies. Preferred eukaryotic cell lines

include the CHO cell lines, various COS cell lines, HeLa cells, myeloma cell lines, transformed B-cells, or human embryonic kidney cell lines. The most preferred host cell is a CHO cell line.

**[0069]** The present invention encompasses antibodies or antigen-binding portions thereof that bind to a specific epitope on the 5T4 antigen. The epitope identified is a nonlinear or conformational epitope comprising a first contact with the human 5T4 antigen (SEQ ID NO: 11) between amino acid residues 173 and 252 and comprising a second contact between amino acid residues 276 and 355 (see Example 7). Thus, the CDRs and heavy and light chain variable regions described herein are used to make full-length antibodies as well as functional fragments and analogs that maintain the binding affinity of the protein employing the CDRs specific for the above mentioned epitope of the 5T4 antigen.

**[0070]** The binding affinity of antibodies of the present invention is determined using SPR (Example 6). In these experiments the 5T4 antigens are immobilized at low densities onto a BIAcore® chip and antibodies are flowed past. Build up of mass at the surface of the chip is measured. This analytical method allows the determination in real time of both on and off rates to obtain affinity ( $K_D$ ) for binding. The humanized antibodies of the present invention have a  $K_D$  of between about 0.30 and about 30 nM; about 0.30 and about 20 nM; about 0.30 and about 10 nM; about 0.5 and about 7 nM; about 1.0 and about 5 nM; and about 1.0 and about 3 nM.

#### Conjugation of Drugs to an Antibody

**[0071]** The drug has, or is modified to include, a group reactive with a conjugation point on the antibody. For example, a drug can be attached by alkylation (e.g., at the epsilon-amino group lysines or the N-terminus of antibodies), reductive amination of oxidized carbohydrate, transesterification between hydroxyl and carboxyl groups, amidation at amino groups or carboxyl groups, and conjugation to thiols. In some embodiments, the number of drug moieties, p, conjugated per antibody molecule ranges from an average of 1 to 8; 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, or 1 to 2. In some embodiments, p ranges from an average of 2 to 8, 2 to 7, 2 to 6, 2 to 5, 2 to 4 or 2 to 3. In other embodiments, p is an average of 1, 2, 3, 4, 5, 6, 7 or 8. In some embodiments, p ranges from an average of about 1 to about 8; about 1 to about 7, about 1 to about 6, about 1 to about 5, about 1 to about 4, about 1 to about 3, or about 1 to about 2. In some embodiments, p ranges from about 2 to about 8, about 2 to about 7, about 2 to about 6, about 2 to about 5, about 2 to about 4 or about 2 to about 3. For examples of chemistries that can be used for conjugation, see, e.g., Current Protocols in Protein Science (John Wiley & Sons, Inc.), Chapter 15 (Chemical Modifications of Proteins) (the disclosure of which is incorporated by reference herein in its entirety.)

**[0072]** For example, when chemical activation of the protein results in formation of free thiol groups, the protein may be conjugated with a sulfhydryl reactive agent. In one aspect, the agent is one which is substantially specific for free thiol groups. Such agents include, for example, maleimide, haloacetamides (e.g., iodo, bromo or chloro), haloesters (e.g., iodo, bromo or chloro), halomethyl ketones (e.g., iodo, bromo or chloro), benzylic halides (e.g., iodide, bromide or chloride), vinyl sulfone and pyridylthio.

### Linkers

**[0073]** The drug can be linked to an antibody by a linker. Suitable linkers include, for example, cleavable and non-cleavable linkers. A cleavable linker is typically susceptible to cleavage under intracellular conditions. Suitable cleavable linkers include, for example, a peptide linker cleavable by an intracellular protease, such as lysosomal protease or an endosomal protease. In exemplary embodiments, the linker can be a dipeptide linker, such as a valine-citrulline (val-cit), a phenylalanine-lysine (phe-lys) linker, or maleimido-capronic-valine-citrulline-p-aminobenzyloxycarbonyl (mc-Val-Cit-PABA) linker. Another linker is Sulfosuccinimidyl-4-[N-maleimidomethyl]cyclohexane-1-carboxylate (smcc). Sulfo-smcc conjugation occurs via a maleimide group which reacts with sulfhydryls (thiols, -SH), while its Sulfo-NHS ester is reactive toward primary amines (as found in Lysine and the protein or peptide N-terminus). Yet another linker is maleimidocaproyl (mc). Other suitable linkers include linkers hydrolyzable at a specific pH or a pH range, such as a hydrazone linker. Additional suitable cleavable linkers include disulfide linkers. The linker may be covalently bound to the antibody to such an extent that the antibody must be degraded intracellularly in order for the drug to be released e.g. the mc linker and the like.

**[0074]** A linker can include a group for linkage to the antibody. For example, linker can include an amino, hydroxyl, carboxyl or sulfhydryl reactive groups (e.g., maleimide, haloacetamides (e.g., iodo, bromo or chloro), haloesters (e.g., iodo, bromo or chloro), halomethyl ketones (e.g., iodo, bromo or chloro), benzylic halides (e.g., iodide, bromide or chloride), vinyl sulfone and pyridylthio). See generally Wong, *Chemistry of Protein Conjugation and Cross-linking*; CRC Press, Inc., Boca Raton, 1991.

### Immunotherapy

**[0075]** For immunotherapy, an antibody can be conjugated to a suitable drug, such as a cytotoxic or cytostatic agent, an immunosuppressive agent, a radioisotope, a toxin, or the like. The conjugate can be used for inhibiting the multiplication of a tumor cell or cancer cell, causing apoptosis in a tumor or cancer cell, or for treating cancer in a patient. The conjugate can be used accordingly in a variety of settings for the treatment of animal cancers. The conjugate can be used to deliver a drug to a tumor cell or cancer cell. Without being bound by theory, in some embodiments, the conjugate binds to or associates with a cancer-cell or a tumor-associated antigen, and the conjugate and/or drug can be taken up inside a tumor cell or cancer cell through receptor-mediated endocytosis. The antigen can be attached to a tumor cell or cancer cell or can be an extracellular matrix protein associated with the tumor cell or cancer cell. Once inside the cell, one or more specific peptide sequences within the conjugate (e.g., in a linker) are hydrolytically cleaved by one or more tumor-cell or cancer-cell-associated proteases, resulting in release of the drug. The released drug is then free to migrate within the cell and induce cytotoxic or cytostatic or other activities. In some embodiments, the drug is cleaved from the antibody outside the tumor cell or cancer cell, and the drug subsequently penetrates the cell, or acts at the cell surface.

### Therapy for Cancer

**[0076]** As discussed above, cancers, including, but not limited to, a tumor, metastasis, or other disease or disorder

characterized by uncontrolled cell growth, can be treated or prevented by administration of a protein-drug conjugate.

**[0077]** In other embodiments, methods for treating or preventing cancer are provided, including administering to a patient in need thereof an effective amount of a conjugate and a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is that with which treatment of the cancer has not been found to be refractory. In some embodiments, the chemotherapeutic agent is that with which the treatment of cancer has been found to be refractory. The conjugate can be administered to a patient that has also undergone a treatment, such as surgery for treatment for the cancer. In another embodiment, the additional method of treatment is radiation therapy.

### Multi-Drug Therapy for Cancer

**[0078]** Methods for treating cancer include administering to a patient in need thereof an effective amount of an antibody-drug conjugate and another therapeutic agent that is an anti-cancer agent. Suitable anticancer agents include, but are not limited to, methotrexate, taxol, L-asparaginase, mercaptopurine, thioguanine, hydroxyurea, cytarabine, cyclophosphamide, ifosfamide, nitrosoureas, cisplatin, carboplatin, mitomycin, dacarbazine, procarbazine, topotecan, nitrogen mustards, cytoxan, etoposide, 5-fluorouracil, BCNU, irinotecan, camptothecins, bleomycin, doxorubicin, idarubicin, daunorubicin, dactinomycin, plicamycin, mitoxantrone, asparaginase, vinblastine, vincristine, vinorelbine, paclitaxel, calicheamicin, and docetaxel.

**[0079]** The ADCs of the present invention can be in the form of a pharmaceutical composition for administration that are formulated to be appropriate for the selected mode of administration, and pharmaceutically acceptable diluent or excipients, such as buffers, surfactants, preservatives, solubilizing agents, isotonicity agents, stabilizing agents, carriers, and the like. Remington's *Pharmaceutical Sciences*, Mack Publishing Co., Easton Pa., 18<sup>th</sup> ed., 1995, incorporated herein by reference, provides a compendium of formulation techniques as are generally known to practitioners.

**[0080]** These pharmaceutical compositions may be administered by any means known in the art that achieve the generally intended purpose to treat cancer. The preferred route of administration is parenteral, defined herein as referring to modes of administration that include but not limited to intravenous, intramuscular, intraperitoneal, subcutaneous, and intraarticular injection and infusion. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

**[0081]** Compositions within the scope of the invention include all compositions wherein an ADC is present in an amount that is effective to achieve the desired medical effect for treating cancer. While individual needs may vary from one patient to another, the determination of the optimal ranges of effective amounts of all of the components is within the ability of the clinician of ordinary skill.

### EXAMPLE 1

#### Preparation of an Anti-5T4 ADC

**[0082]** 5T4-A1 antibody drug conjugate (ADC) is prepared via partial reduction of the mAb with tris(2-carboxy-

ethyl)phosphine (TCEP) followed by reaction of reduced Cys residues with the desired maleimide terminated linker-payload. In particular, 5T4-A1 mAb is partially reduced via addition of 2.8 molar excess of tris(2-carboxyethyl)phosphine (TCEP) in 100 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid buffer), pH 7.0 and 1 mM diethylenetriaminepentaacetic acid (DTPA) for 2 h at 37° C. The desired linker-payload is then added to the reaction mixture at a linker-payload/mAb-thiol molar ratio of 5.5 (maleimidocapronic-monomethylauristatin F [mc-MMAF]) or 8 (maleimidocapronic-valine-citruline-p-aminobenzyloxycarbonyl-monomethylauristatin E [mc-Val-Cit-PABA-MMAE]) and reacted for an additional 1 h at 25° C. in the presence of 15% v/v of dimethylacetamide (DMA). After the 1 h incubation period, N-ethylmaleimide (4.5 fold excess for mc-MMAF and 2 fold excess for mc-Val-Cit-PABA-MMAE) is added to cap the unreacted thiols and is allowed to react for 15 minutes followed by addition of 6 fold excess L-Cys to quench any unreacted linker-payload. The reaction mixture is dialyzed overnight at 4° C. in phosphate buffered saline (PBS), pH 7.4, and purified via SEC (AKTA explorer, Superdex200 10/30 GL column). The ADC is further characterized via size exclusion chromatography (SEC) for purity, hydrophobic interaction chromatography (HIC), and liquid chromatography electrospray ionisation tandem mass spectrometry (LC-ESI MS) to calculate loading, and the concentration is determined via UV spectrophotometer.

#### EXAMPLE 2

##### Binding Studies

**[0083]** Cells expressing the 5T4 antigen, and the negative control Raji cells, are plated at a density of 500,000 cells/well on non-tissue culture treated 96 well plates and kept on ice. Dilutions of the A1 and A1-IgG4 antibodies or A1-mcMMAF ADC are made in 3% bovine serum albumin BSA in Dulbecco's phosphate buffered saline (DPBS) and added to the plate at a final concentration of 10 µg/mL. The plates are then incubated on ice for 1 hour followed by 2 washes. The secondary antibody, PE (phycoerythrin) conjugated Goat Anti-Human IgG Fc is added to the wells. After 30 minutes of incubation at 4 C, the mean fluorescence intensity is then measured using a flow cytometer.

**[0084]** The data in Table 4 indicates that the A1 antibody binds a diverse panel of 5T4 positive cell lines. The data in Table 5 indicates that similar binding on several different cell lines is observed with the A1 and A1-IgG4 antibodies as well as the A1-mcMMAF ADC.

TABLE 4

Human Cell Lines	A1 antibody Mean Fluorescent Intensity
MDAMB435/5T4 (melanoma)	15000
MDAMB468 (breast)	3000
MDAMD361-DYT2 (breast)	4500
NCI-H157 (lung)	4100
A431 (epithelial)	2000
Caki (kidney)	2500
PC3mm2 (prostate)	4500
PC14PE6 (lung)	3200
Panc1 (pancreatic)	3500
BxPC3 (pancreatic)	1000
Su8686 (pancreatic)	3700
H1975	1600

TABLE 4-continued

Human Cell Lines	A1 antibody Mean Fluorescent Intensity
37622A-(Primary Lung cancer cells)	10600
Raji (negative control)	<100

TABLE 5

Cell Lines	Mean Fluorescent Intensity				
	A1	A1-mcMMAF	A1-IgG4	A1-IgG4-CM	IgG control
MDAMD361-DYT2	7000	6900	5500	4800	<100
A431	3900	3400	2400	2000	<100
MDAMB468	3500	2800	2500	1800	<100
PC3mm2	3400	2700	2100	1500	<100
Raji	<200	<200	<200	<100	<100

#### EXAMPLE 3

##### Cytotoxicity Assay

**[0085]** Cell lines expressing 5T4, and the negative control Raji cell line, are cultured with increasing concentrations of ADC. After four days, viability of each culture is assessed. IC<sub>50</sub> values are calculated by logistic non-linear regression and are presented as ng Ab/mL. A1-mcMMAF, A1-vcMMAE, A3-mcMMAF and A3-vcMMAE are shown to inhibit the growth of 5T4 expressing cell lines (MDAMB435/5T4, MDAMB468, and MDAMB361DYT2), while being inactive on 5T4 negative cells (Raji), Table 6.

TABLE 6

ADC	IC <sub>50</sub> (ng Ab/ml)			
	MDAMB435/5T4	MDAMB361DYT2	MDAMB468	Raji (5T4-)
A1-mcMMAF	1.3	104.2	534.2	>45,000
A1-vcMMAE	6.8	157.7	7667	>45,000
A3-mcMMAF	0.3	31.0	27.7	>45,000
A3-vcMMAE	3.5	86.8	160.6	>45,000
Non-binding Ab-mcMMAF	21258	~50,000	73059	>45,000
Non-binding Ab-vcMMAE	7979	27650	23819	>45,000

**[0086]** Additionally, 5T4+primary lung tumor 37622a cells are isolated and grown in culture. Cells are cultured with increasing concentrations of ADC. Ten days later, viability of each culture is assessed using the MTS method. IC<sub>50</sub> values were calculated by logistic non-linear regression and are presented as ng Ab/ml. A1-mcMMAF, A1-vcMMAE, A3-mcMMAF, and A3-vcMMAE inhibit the growth of the primary lung tumor cells, Table 7.



TABLE 7

ADC	37622a primary lung IC <sub>50</sub> (ng Ab/ml)
A1-mcMMAF	504.1
A1-vcMMAE	443.1
A3-mcMMAF	77.2
A3-vcMMAE	78.1
Nonbinding Ab- mcMMAF	>45,000

## EXAMPLE 4

## Subcutaneous Xenograft Model

**[0087]** Female, athymic (nude) mice (or another strain of immunosuppressed mice) are injected s.c. with MDAMB435/5T4, MDAMB361DYT2, or H1975 tumor cells. Mice with staged tumors, approximately 0.1 to 0.3 g (n=6 to 10 mice/treatment group) are administered intravenously Q4Dx4 with normal saline (vehicle), A1-mcMMAF, A1-vcMMAE, A1-mcMMAD, A1-smccDM1, A3-mcMMAF, A3-vcMMAE, or a nonbinding control antibody conjugated to either mcMMAF or vcMMAE, at the dose of 3 mg Ab/kg. All ADCs are dosed based on Ab content. Tumors are measured at least once a week and their size (mm<sup>2</sup>±SEM) is calculated as mm<sup>2</sup>=0.5×(tumor width<sup>2</sup>)×(tumor length).

**[0088]** The data in Table 8 indicates that A1-mcMMAF, A1-vcMMAE, A1-vcMMAD, A3-mcMMAF, and A3-vcMMAE inhibit the growth of MDAMB435/5T4 xenografts while A1-mcMMAD and A1-smccDM1 were not active in this model.

**[0089]** The data in Table 9 indicates that A1-mcMMAF, A1-vcMMAE, A1-vcMMAD, A1-smccDM1, A3-mcMMAF, and A3-vcMMAE inhibit the growth of MDAMB361DYT2 xenografts while A1-mcMMAD was not active in this model.

**[0090]** The data in Table 10 indicates that A1-mcMMAF, A1-vcMMAE, A1-vcMMAD, A3-mcMMAF, and A3-vcMMAE inhibit the growth of H1975 xenografts while A1-mcMMAD and A1-smccDM1 were not active in this model.

TABLE 8

Compound (3 mg/kg)	MDAMB435/5T4 xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
Q4dx4)	Day 0	Day 17	Day 42	Day 65	Day 85
Vehicle	169 ± 8	531 ± 73	1255 ± 190	GT	GT
A1 mcMMAF	168 ± 15	53 ± 12	67 ± 56	174 ± 119	364 ± 278

TABLE 8-continued

Compound (3 mg/kg)	MDAMB435/5T4 xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
Q4dx4)	Day 0	Day 17	Day 42	Day 65	Day 85
A1 vcMMAE	168 ± 8	4 ± 4	10 ± 10	91 ± 91	200 ± 200
A1 mcMMAD	168 ± 12	390 ± 112	GT	GT	GT
A1 smccDM1	174 ± 10	429 ± 62	1255 ± 227	1781 ± 388	GT
A1 vcMMAD	169 ± 12	17 ± 7	0 ± 0	0 ± 0	0 ± 0
A3 mcMMAF	174 ± 12	105 ± 27	216 ± 143	448 ± 220	GT
A3 vcMMAE	172 ± 13	0 ± 0	0 ± 0	0 ± 0	0 ± 0
Nonbinding Ab mcMMAF	170 ± 11	100 ± 15	314 ± 121	838 ± 381	GT
Nonbinding Ab vcMMAE	172 ± 11	168 ± 53	461 ± 178	GT	GT

GT = group terminated due to large tumor size

TABLE 9

Compound (3 mg/kg)	MDAMB361DYT2 xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
Q4dx4)	Day 0	Day 19	Day 47	Day 90	Day 131
Vehicle	353 ± 10	363 ± 58	558 ± 149	1117 ± 348	GT
A1 mcMMAF	348 ± 14	76 ± 32	0 ± 0	7 ± 7	11 ± 11
A1 vcMMAE	356 ± 11	86 ± 8	0 ± 0	9 ± 9	34 ± 27
A1 vcMMAD	352 ± 26	130 ± 15	0 ± 0	0 ± 0	0 ± 0
A3 mcMMAF	342 ± 23	128 ± 10	79 ± 30	105 ± 49	353 ± 234
A3 vcMMAE	354 ± 21	111 ± 20	21 ± 21	72 ± 72	155 ± 155
A1 mcMMAD	347 ± 15	380 ± 66	775 ± 199	GT	GT
A1 vcMMAD	352 ± 26	130 ± 15	0 ± 0	0 ± 0	0 ± 0
A1 smccDM1	353 ± 25	123 ± 9	51 ± 25	98 ± 41	330 ± 146
Nonbinding Ab mcMMAF	342 ± 38	407 ± 93	869 ± 198	GT	GT
Nonbinding Ab vcMMAE	344 ± 20	303 ± 78	346 ± 185	595 ± 362	GT

GT = group terminated due to large tumor size

TABLE 10

		H1975 Xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
Compound	Q4dx4	Day 0	Day 8	Day 15	Day 22	Day 40
Vehicle		423 ± 14	1154 ± 136	2229 ± 240	GT	GT
A1 mcMMAF	3	425 ± 14	619 ± 46	519 ± 45	581 ± 79	2840 ± 207
A1 vcMMAE	3	425 ± 12	702 ± 45	929 ± 90	926 ± 116	GT
A1 vcMMAD	3	427 ± 18	739 ± 59	467 ± 19	240 ± 14	625 ± 317

TABLE 10-continued

Compound	Dose (mg/kg)	H1975 Xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
		Q4dx4	Day 0	Day 8	Day 15	Day 22
A3 mcMMAF	3		426 ± 10	980 ± 79	1343 ± 140	1261 ± 203
A3 vcMMAE	3		431 ± 14	944 ± 52	993 ± 71	GT
A1 mcMMAD	3		427 ± 16	837 ± 69	1468 ± 139	GT
A1 smccDM1	3		423 ± 18	901 ± 83	1852 ± 167	GT
Nonbinding Ab- mcMMAF	3		423 ± 16	1026 ± 68	1861 ± 224	GT
Nonbinding Ab-vcMMAE	3		427 ± 13	1213 ± 67	1959 ± 139	GT

GT = group terminated due to large tumor size

**[0091]** Alternatively, nude mice with 37622a primary tumor cell xenografts established subcutaneously are treated iv Q4D×4 with A1-mcMMAF, A1-mcMMAD, A1-vcMMAD, or A3-mcMMAF at the dose of 3 mg Ab/kg and the tumor growth is monitored over the period of 96 days. Table 11 demonstrates that A1-mcMMAF, A1-vcMMAD and A3-mcMMAF inhibit the growth of 37622a primary tumor xenografts compared to vehicle control treated animals while A1-mcMMAD was not active in this model.

effective against the MDAMB361DYT2 xenograft but had essentially no effect against the MDAMB435/5T4 and the H1975 xenografts even though all the xenografts have a high expression of the 5T4 target antigen. This data illustrates that the effectiveness of the linker-payload could not be predicted even when the same high affinity antibody is utilized or even when the same ADC is used.

TABLE 11

Compound	Dose (mg/kg)	37622a Primary Tumor Xenografts Tumor volume (mm <sup>3</sup> , x ± sem)				
		Q4dx4	Day 1	Day 22	Day 46	Day 68
Vehicle			111 ± 18	503 ± 155	1174 ± 247	GT
A1- mcMMAF	3		111 ± 18	67 ± 11	124 ± 47	233 ± 105
A1- mcMMAD	3		127 ± 28	376 ± 119	862 ± 377	GT
A1- vcMMAD	3		108 ± 14	52 ± 14	13 ± 5	50 ± 37
A3- mcMMAF	3		131 ± 28	99 ± 26	211 ± 128	463 ± 210

GT = group terminated due to large tumor size

**[0092]** Unexpectedly, the data in Tables 8-11 show that ADCs with the same antibody and drug payload but with different linkers had a dissimilar efficacy profile i.e. A1-mcMMAD vs A1-vcMMAD in all four xenograft models. In addition, the data show that ADCs with the same antibody and linker but with different drug payloads also had a different efficacy profile i.e. A1-mcMMAF vs A1-mcMMAD, in all four xenograft models. Thus, the drug MMAD is effective in all four xenograft models when linked to the A1 antibody by the vc linker but has no activity in any of the xenograft models tested when linked by the mc linker. In contrast, the drug MMAF is highly effective in all 4 xenograft models when linked to the A1 antibody with the mc linker while the chemically related drug MMAD has no activity in all 4 xenograft models when linked to the same antibody by the same linker.

**[0093]** Yet another unexpected observation is seen with the ADC A1-smccDM1 (Tables 8-10). This ADC was very

## EXAMPLE 5

## Antibody Dependent Cell-Mediated Cytotoxicity (ADCC)

## ADCC Assay

**[0094]** Blood from a healthy volunteer is collected into a BD Vacutainer CPT cell preparation tube with sodium heparin. Human peripheral blood mononucleocytes (PBMC) are harvested and resuspended in assay buffer (RPMI 1640 supplemented with 10 mM HEPES) at 2.5×10<sup>7</sup> cells/ml. Target cells (MDAMB435/5T4 or MDAMB435/neo) are seeded at a density of 1×10<sup>4</sup> cells/well in a 96 well assay plate. A1 antibody or A1-mcMMAF are added, then human PBMC effector cells (5×10<sup>5</sup>) are dispensed into the wells for an effector:target cell ratio (E:T) of 50:1. The assay plate is incubated at 37° C. for 4 hours for ADCC activity. The plate is harvested by adding equal volume of CytoTox-One

reagent (Promega). Stop solution (Promega; 50 ul) is added to each well and lactate dehydrogenase release was quantified by measuring fluorescence intensity. As a positive control, 2  $\mu$ l of lysis buffer per well is added to generate a maximum LDH release (100% cytotoxicity) in control wells. Percent cytotoxicity is calculated using the following equation:

$$\% \text{ Specific Cytotoxicity} = \frac{\text{experimental} - \text{effector spontaneous} - \text{target spontaneous}}{\text{target maximum} - \text{target spontaneous}} \times 100$$

**[0095]** Where “experimental” corresponds to the signal measured in one of the experimental conditions, “effector spontaneous” corresponds to the signal measured in the presence of PBMC alone, “target spontaneous” corresponds to the signal measured in the presence of target cells alone, and “target maximum” corresponds to the signal measured in the presence of detergent-lysed target cells alone.

**[0096]** The ADCC activity of A1-IgG1 Ab and A1-mcMMAF compared to A1-IgG4 Ab is shown in Table 12. Both the A1 antibody and A1-mcMMAF demonstrate comparable ADCC activity indicating that the ADCC activity of A1-mcMMAF may contribute to its anti-tumor activity.

TABLE 12

Compound	% Cytotoxicity
A1-IgG1	37 $\pm$ 8
A1-mcMMAF	34 $\pm$ 1
A1-IgG4	9 $\pm$ 5

## EXAMPLE 6

## Binding Affinity

**[0097]** Surface plasmon resonance (SPR) analysis is performed utilizing the BIAcore® to determine the affinity constants for A1-IgG1 and A1-IgG4 binding to either human or cynomolgus 5T4 at pH 6.0 and pH 7.4. BIAcore® technology utilizes changes in the refractive index at the surface layer upon binding of the huA1 antibody variants to the human 5T4 protein immobilized on the surface layer. Binding is detected by SPR of laser light refracting from the surface. Analysis of the signal kinetics on-rate and off-rate allows the discrimination between non-specific and specific interactions. The 5T4 proteins used for this analysis consisted of the human or cynomolgus 5T4 ectodomain fused to the human IgG1-Fc domain and low densities (45.1 and 45.4 RU for human and cynomolgus respectively) are immobilized onto a CM5 chip to accurately measure affinity constants.

**[0098]** The measurement of specific binding to the 5T4 ectodomain is attained by subtracting binding to a reference surface that had only human IgG1-Fc protein immobilized onto the CM5 chip at the same density to that on the 5T4-Fc surfaces. Next, various concentrations of A1, A1-IgG4, or A3 antibodies in either HBS-EP pH 7.4 or MES-EP pH 6.0 buffer are injected over the surface. The surface is regenerated two times with Glycine pH 1.7+0.05% Surfactant P20 (GE Healthcare, BR-1000-54) between injection cycles.

**[0099]** Results show that the A1 has a slightly higher affinity for human 5T4 using the low-density 5T4 surface at

both pH 6.0 and pH 7.4 relative to A1-IgG4 (1.5-fold and 1.2-fold respectively, Table 13). Additionally, A1 exhibited slightly better binding to cynomolgus 5T4 at both pH 6.0 and pH 7.4 compared to A1-IgG4 (1.7-fold and 1.2-fold respectively) and both A1 and A1-IgG4 bound human 5T4, 3-4 fold better than cynomolgus 5T4 (Table 12).

TABLE 13

Antibody	Antigen	pH	ka (1/Ms)	kd (1/s)	KD (nM)
A1-IgG1	hu5T4	6.0	4.31E+05	4.59E-04	1.06
A1-IgG4	hu5T4	6.0	6.26E+05	8.93E-04	1.43
A1-IgG1	cyno5T4	6.0	2.33E+05	6.41E-04	2.76
A1-IgG4	cyno5T4	6.0	2.02E+05	9.50E-04	4.70
A1-IgG1	hu5T4	7.4	2.75E+05	1.32E-04	0.48
A1-IgG4	hu5T4	7.4	3.28E+05	1.72E-04	0.52
A1-IgG1	cyno5T4	7.4	1.51E+05	2.73E-04	1.80
A1-IgG4	cyno5T4	7.4	1.81E+05	3.82E-04	2.11

**[0100]** Comparing the A1 and A3 antibodies, it is apparent that the A1 antibody binds human and cynomolgus 5T4 better at pH 7.4 relative to pH 6.0 while the A3 antibody exhibits enhanced binding at pH 6.0 compared to pH 7.4, Table 14.

TABLE 14

Antibody	Antigen	pH	ka (1/Ms)on	kd (1/s)off	KD (nM)
A1	hu5T4	6.0	4.31E+05	4.59E-04	1.06
A3	hu5T4	6.0	3.51E+05	4.17E-05	0.12
A1	cyno5T4	6.0	2.33E+05	6.41E-04	2.76
A3	cyno5T4	6.0	4.58E+05	1.87E-04	0.41
A1	hu5T4	7.4	2.75E+05	1.32E-04	0.48
A3	hu5T4	7.4	1.79E+05	3.06E-05	0.17
A1	cyno5T4	7.4	1.51E+05	2.73E-04	1.80
A3	cyno5T4	1.98E+05	1.62E-04	0.82	1.98E+05

## EXAMPLE 7

## Epitope Mapping Using 5T4 Chimeras

**[0101]** To identify the epitopes to which each of the A1 and A3 antibodies bind, an enzyme linked immunosorbent assay (ELISA) is performed using (1) 5T4 ectodomain Fc construct and (2) human/mouse 5T4 chimera constructs transiently expressed in COS-1 cells. The ectodomain includes the amino-terminal region, two leucine-rich repeats, and the intervening hydrophilic region. Mouse and rat 5T4 ectodomains contain a 6 amino acid direct repeat within their hydrophilic region.

**[0102]** Fusion proteins containing a 5T4 ectodomain and a Fc constant region from human IgG1 are prepared using human 5T4 (amino acids 1-355), mouse 5T4 (amino acids 1-361), rat 5T4 (amino acids 1-361), cynomolgus monkey 5T4 (amino acids 1-355), chimpanzee 5T4 (amino acids 1-355), and black-tailed marmoset (amino acids 1-355). The binding results with human/mouse 5T4 chimera constructs are summarized in Table 14, which indicates specific binding, partial binding, or lack of binding, by the A1 and A3 antibodies.

[0103] Table 15 refers to binding ability of the antibodies to the various human/mouse chimeras and the nomenclature is designated by mouse 5T4 content. When no binding is observed, this indicates where the antibody binds human 5T4 since these antibodies do not bind mouse 5T4. For example, the A3 antibody has the most N-terminal binding epitope (between 83-163) and this is shown by lack of binding to the 5T4 chimera that has residues 83-163 replaced by mouse 5T4, hence A3 can no longer bind. Based upon these results, it is determined that humanized A1 antibody has a first contact with human 5T4 between amino acid residues 173 and 252 and a second contact with human 5T4 between amino acid residues 276 and 355. The A3 antibody binds the first leucine-rich repeat region of human 5T4 between amino acid residues 83 through 163. The number of amino acid residues corresponds to the human 5T4 antigen amino acid sequence of SEQ ID NO: 11.

TABLE 15

5T4 Antigen Construct	Antibody	
	A1	A3
Human/mouse 83-163	+	-
Human/mouse 173-361	-	+
Human/mouse 173-258	+/-	+
Human/mouse 282-361	+/-	+

## EXAMPLE 8

Comparison of A1-mcMMAF ADC with  
A1-IgG4-CM ADC

[0104] A1-mcMMAF is compared to A1-IgG4-AcBut calicheamicin (A1-IgG4-CM) for both safety and efficacy. A1-4-CM is comprised of the A1-IgG4 antibody conjugated with the linker, AcBut[-(4' acetylphenoxy)butanoic acid], to a calicheamicin payload. The calicheamicins are potent antitumor agents of a class of enediyne antibiotics derived from the bacterium *Micromonospora echinospora*.

[0105] The cell binding activity of A1 Ab, A1-IgG4 Ab, A1-mcMMAF ADC and A1-IgG4-CM ADC are compared using several 5T4 positive cell lines (see Example 2, Table 5). The data indicates that similar binding is observed with the A1 and A1-IgG4 antibodies as well as the A1-mcMMAF ADC, all of which have a higher mean fluorescent intensity than A1-IgG4-CM for all the 5T4 positive cell lines tested.

[0106] A1-mcMMAF and A1-IgG4-CM are tested side-by-side in the MDAMB435/5T4 subcutaneous xenograft model. Both ADCs are given iv (Q4dx2) when the tumors reach approximately 200 mm<sup>2</sup> in size. The anti-tumor activity of A1-IgG4-CM at a dose of 3 mg/kg is similar to the anti-tumor activity of A1-mcMMAF administered at dose of 10 mg/kg (Table 16). Based upon these results, the anti-tumor activity of A1-IgG4-CM is approximately 3.3 fold more potent than A1-mcMMAF.

TABLE 16

Compound	Dose (mg/kg) Q4dx4	Tumor volume (mm <sup>3</sup> , x ± sem)				
		Day 0	Day 7	Day 21	Day 31	Day 45
Vehicle	0	123 ± 8	195 ± 36	402 ± 56	635 ± 111	1309 ± 332
A1-mcMMAF	3	124 ± 11	121 ± 8	166 ± 29	227 ± 42	361 ± 89

TABLE 16-continued

Compound	Dose (mg/kg) Q4dx4	Tumor volume (mm <sup>3</sup> , x ± sem)				
		Day 0	Day 7	Day 21	Day 31	Day 45
A1-mcMMAF	10	123 ± 14	76 ± 11	0 ± 0	3 ± 3	2 ± 1
A1-IgG4-CM	3	121 ± 12	140 ± 15	32 ± 10	24 ± 10	26 ± 15

[0107] It could be expected that the 3.3 fold enhanced potency of A1-IgG4-CM over that of A1-mcMMAF would translate into a 3.3 fold enhanced safety margin of A1-mcMMAF over that of A1-IgG4-CM in an animal toxicity study. However, when the safety profile of A1-IgG4-CM in cynomolgus macques is reviewed, it is determined that A1-IgG4-CM is at least 100 fold more toxic than A1-mcMMAF in the cynomolgus macque. When A1-IgG4-CM is administered at 0.032, 0.095 and 0.32 mg Ab/kg/cycle (2, 6, 20 µg calicheamicin/kg/cycle) to male (n=3) and female (n=3) cynomolgus macques, toxicity is observed at each dose level. After 2 cycles (2 doses), 4 out of 6 animals in the 0.095 treatment group are either euthanized or found dead. On the other hand, no deaths are observed at dosages up to 10 mg/kg with A1-mcMMAF (247 pg mcMMAF/kg/cycle), after 2 cycles (2 doses), over the same 4 week time period. In summary, the 10 mg/kg dosage group of A1-mcMMAF is safe while the 0.096 mg/kg dosage group of A1-IgG4-CM is deemed toxic when both are administered twice to cynomolgus macques in a 4 week observation period.

[0108] Unexpectedly, these results demonstrate a 105 fold (10/0.095=105) safety margin of A1-mcMMAF over that of A1-IgG4-CM, rather than the expected 3.3 fold safety margin based on the relative anti-tumor potency of each ADC. This data reveals the unpredictable nature of antibody-drug conjugates that utilize antibodies to the same antigen target but are conjugated to a different drug payload.

## EXAMPLE 9

A1-mcMMAF Mouse PK/PD Modeling and  
Clinical Dose Predictions

[0109] PK/PD modeling has been used to quantify the tumor response of A1-mcMMAF in mouse xenograft studies, in order to determine efficacious concentration across cell lines. The transit compartment tumor kill PK/PD model used was previously described by Simeoni et al. (Simeoni et al, *Cancer Res*, 64:1094, (2004). The model has been modified to account for linear, exponential and logistic growth of tumor, and saturative killing by the drug. PK/PD model parameters include:

[0110]  $k_{g\text{ ex}}$  exponential growth

[0111]  $k_g$  logistic growth

[0112]  $w_0$  initial tumor volume

[0113]  $\tau$  transduction rate

[0114]  $k_{max}$  maximum kill rate

[0115]  $kC_{50}$  concentration at half max kill rate

[0116] The PK/PD modeling results are used to calculate the Tumor Static Concentration (TSC, Equation 1). This is the drug concentration where tumor growth is equal to tumor death rates and tumor volume remains unchanged. TSC can be defined as the minimal concentration required for effi-

cacy. TSC is used to give guidance on clinical dose selection, with concentrations of >TSC required for efficacy in the clinic.

[0117] For A1-mcMMAF, mouse PK was determined in a separate study (3mg/kg IV, female athymic nu/nu mice). Mouse xenograft studies were completed using 3 different 5T4 cell lines with A1-mcMMAF administered at dose levels between 1 and 30 mg/kg every 4 days: cell line MDAMB435/5T4 (dosed at 1, 3, 10, and 30 mg/kg), cell line H1975 (dosed at 1, 3, and 10 mg/kg) and cell line 37622A (dosed at 1 and 10 mg/kg). PK/PD modeling was performed as described and TSCs are reported in Table 17.

[0118] Mouse PK/PD parameters for each xenograft cell line were combined with predicted human PK of A1-mcMMAF to simulate doses required for efficacy in the clinic. Using this methodology, A1-mcMMAF has a predicted minimally efficacious clinical dose of about 0.22 to about 2.3 mg/kg Q3 weeks [every three weeks] (Table 17).

[0119] In an embodiment of the present invention, dose ranges can be in the range from about 0.18 mg/kg to about 2.7 mg/kg, from about 0.22 mg/kg to about 2.6 mg/kg, from about 0.27 mg/kg to about 2.5 mg/kg, from about 0.32 mg/kg to about 2.3 mg/kg, from about 0.37 mg/kg to about 2.15 mg/kg, from about 0.42 mg/kg to about 2.10 mg/kg, from about 0.47 mg/kg to about 2.05 mg/kg, from about 0.52 mg/kg to about 2.00 mg/kg, from about 0.57 mg/kg to about 1.95 mg/kg, from about 0.62 mg/kg to about 1.90 mg/kg, from about 0.67 mg/kg to about 1.85 mg/kg, from about 0.72 mg/kg to about 1.80 mg/kg, from about 0.82 mg/kg to about 1.70 mg/kg, from about 0.92 mg/kg to about 1.60 mg/kg, from about 1.02 mg/kg to about 1.50 mg/kg, from about 1.12 mg/kg to about 1.40 mg/kg, or from about 1.20 mg/kg to about 1.30 mg/kg, with dosing at Q3 weeks. Preferably, dose ranges can be in the range from about 0.22 mg/kg to about 2.3 mg/kg.

Equation 1

$$\text{If } \frac{k_{gEx}}{k_g} w_0 \leq 1, \quad TSC = \frac{k_{gEx} \cdot k_{C50}}{k_{max} - k_{gEx}} \quad 1.1$$
$$\text{If } \frac{k_{gEx}}{k_g} w_0 > 1, \quad TSC = \frac{k_g \cdot k_{C50}}{w_0 \cdot k_{max} - k_g} \quad 1.2$$

TABLE 17

Cell Line	TSC [80% confidence] (ug/ml)	Predicted Stasis Dose [80% confidence] (mg/kg Q3 weeks)
MDAMB435/5T4	1.1 [0.9, 1.4]	0.22 [0.18, 0.28]
37622A	5.1 [2.1, 9.9]	1.1 [0.6, 2.0]
H1975	11.6 [9.6, 14.1]	2.3 [2.0, 2.7]

SEQUENCE LISTING

[0120]

Humanized A1 human IgG1 heavy chain  
SEQ ID NO: 1  
EVQLVESGGGLVQPGGSLRLSCAASGYTFTNFGMNWVRQAPGKGLEWVAW  
INTNTGEPRYAEEFKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDW

-continued

DGAYFFDYWGQGLTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKD  
YFPEPVTWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTY  
ICNVNHKPSNTKVDKKVEPKSCDKTHTCTPPCPAPELLGGPSVFLFPPKPK  
DTLMISRTPETCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS  
TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV  
YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVVL  
DSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK  
Humanized A1 human Kappa light chain:  
SEQ ID NO: 2  
DIQMTQSPSSLSASVGRVTITCKASQSVSNDAVAYQQKPGKAPKLLIYF  
ATNRYTGVPSTRFSGSGYGTDFLTITISLQPEDFATYYCQQDYSSPWTFGQ  
GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV  
DNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQG  
LSSPVTKSFNRGEC  
A1-VH  
SEQ ID NO: 3  
EVQLVESGGGLVQPGGSLRLSCAASGYTFTNFGMNWVRQAPGKGLEWVAW  
INTNTGEPRYAEEFKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDW  
DGAYFFDYWGQGLTLVTVSS  
A1-VL  
SEQ ID NO: 4  
DIQMTQSPSSLSASVGRVTITCKASQSVSNDAVAYQQKPGKAPKLLIYF  
ATNRYTGVPSTRFSGSGYGTDFLTITISLQPEDFATYYCQQDYSSPWTFGQ  
GTKVEIK  
A1-HC CDR1  
SEQ ID NO: 5  
NFGMN  
A1-HC CDR2  
SEQ ID NO: 6  
WINTNTGEPRYAEEFKG  
A1-HC CDR3  
SEQ ID NO: 7  
DWDGAYFFDY  
A1-LC-CDR1  
SEQ ID NO: 8  
KASQSVSNDA  
A1-LC-CDR2  
SEQ ID NO: 9  
FATNRYT  
A1-LC-CDR3  
SEQ ID NO: 10  
QQDYSSPWT  
Human 5T4 antigen  
SEQ ID NO: 11  
MPGGCSRGAAGDGLRLRLALVLLGNVSSSSPTSSASS  
FSSSAPFLASAVSAQPPLPDQCPALCECSEAARTVKCVNR  
NLTEVPTDLPAYVRNFLTGNQLAVLPAGAFARRPPLAEL  
AALNLSGSRLDEVRAFAFEHLPSLRQLDLSHNPLADLSPF

-continued

AFSGSNASVSAPSPLVELILNHI VPPEDERQNRSEFGMVV

AALLAGRALQGLRRLELASNHFLYLPRDVLAQLPSLRHLD

LSNNSLVSLTYVSFRNLTHLESLHLEDNALKVLHNGTLAE

LQGLPHIRVFLDNNPWVCDCHMADMVTLKETEVVQGKDR

LTCAYPEKMRNRVLELNSADLDCDPIPPSLQTSYVFLG

IVLALIGAIFLLVLYLNRKGIKKWMHNIIRDACRDHMEGYH

YRYEINADPRLTNLSSNSDV

Humanized A1 human IgG4m heavy chain:

SEQ ID NO: 12

EVQLVESGGGLVQPGGSLRLSCAASGYTFTNFGMNWVRQAPGKGLEWVAW

INTNTGEPRYAEFEKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDWDGAYFFDYWGQGLTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKD

YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTTY

TCNVDHKPSNTKVDKRVESKYGPPCPPCAPEFLGGPSVFLFPPKPKDTL

MISRTPEVTCVVVDVSDQEDPEVFQFNWYVDGVEVHNAKTKPREEQFNSTYR

VVSVLTVLHQDLNGKEYCKVSNKGLPSSIEKTIKAKGQPREPQVYTL

PPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD

GSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKSLSLSLGK

A1 human IgG4m VH (A1-IGG4-VH)

SEQ ID NO: 13

EVQLVESGGGLVQPGGSLRLSCAASGYTFTNFGMNWVRQAPGKGLEWVAW

INTNTGEPRYAEFEKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDW

DGAYFFDYWGQGLTLTVSS

A1-IgG4-VH-CDR1

SEQ ID NO: 14

NFGMN

Chimeric A3 heavy chain (muA3-huIgG1)

SEQ ID NO: 15

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVAR

IRSKSNNYATYYADSVKDRFTISRDDSQSMYLYLQMNNLKTEDTAMYYCVR

QWDYDVRAMNYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL

VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGT

QTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP

KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQ

YNSTYRVVSVLTVLHQDLNGKEYCKVSNKALPAPIEKTISKAKGQPRE

PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTP

PVLDSDGSFFLYSLKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSP

GK

Chimeric A3 VH

SEQ ID NO: 16

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVAR

IRSKSNNYATYYADSVKDRFTISRDDSQSMYLYLQMNNLKTEDTAMYYCVR

QWDYDVRAMNYWGQGLTLTVSS

-continued

Chimeric A3 VH-CDR1

SEQ ID NO: 17

TYAMN

Chimeric A3 VH-CDR2

SEQ ID NO: 18

RIRSKSNNYATYYADSVKD

Chimeric A3 VH-CDR3

SEQ ID NO: 19

QWDYDVRAMNY

Chimeric A3 light chain (muA3-huKappa)

SEQ ID NO: 20

DIVMTQSHIFMSTSVGDRVSITCKASQDQDVTAVAWYQQKPGQSPKLLIYW

ASTRLTGVPDRFTGSGSGTDFTLTISNVQSEDLADYFCQQYSSYPYTFGQ

GTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV

DNALQSGNSQESVTEQDSKDYSTYLSSTLTLSKADYEKHKVYACEVTHQG

LSSPVTKSFNRGEC

Chimeric A3 VL

SEQ ID NO: 21

DIVMTQSHIFMSTSVGDRVSITCKASQDQDVTAVAWYQQKPGQSPKLLIYW

ASTRLTGVPDRFTGSGSGTDFTLTISNVQSEDLADYFCQQYSSYPYTFGQ

GTKLEIK

Chimeric A3 VL-CDR1

SEQ ID NO: 22

KASQDQDVA

Chimeric A3 VL-CDR2

SEQ ID NO: 23

WASTRLT

Chimeric A3 VL-CDR3

SEQ ID NO: 24

QQYSSYPYT

Humanized A3 human IgG1 heavy chain:

SEQ ID NO: 25

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVAR

IRSKSNNYATYYADSVKDRFTISRDDAKNSLYLQMNSLRAEDTAVYYCVR

QWDYDVRAMNYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL

VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGT

QTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP

KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQ

YNSTYRVVSVLTVLHQDLNGKEYCKVSNKALPAPIEKTISKAKGQPRE

PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTP

PVLDSDGSFFLYSLKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSP

GK

Humanized A3 VH:

SEQ ID NO: 26

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVAR

IRSKSNNYATYYADSVKDRFTISRDDAKNSLYLQMNSLRAEDTAVYYCVR

QWDYDVRAMNYWGQGLTLTVSS

-continued

Humanized A3 VH-CDR1:  
 TYAMN  
 SEQ ID NO: 27

Humanized A3 VH-CDR2:  
 RIRSKSNYYATYYADSVKD  
 SEQ ID NO: 28

Humanized A3 VH-CDR3:  
 QWDYDVRAMNY  
 SEQ ID NO: 29

Humanized A3 human Kappa light chain:  
 DIQMTQSPSSLSASVGDRTITCKASQDVTAVAWYQQKPGKAPKLLIYW  
 SEQ ID NO: 30

ASTRLTGVPSTRFSGSGSGTDFTLTISLQPEDFATYYCQQYSSYPYTFGQ  
 GTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV

-continued

DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQG  
 LSSPVTKSFNRGEC

Humanized A3 VL:  
 DIQMTQSPSSLSASVGDRTITCKASQDVTAVAWYQQKPGKAPKLLIYW  
 SEQ ID NO: 31

ASTRLTGVPSTRFSGSGSGTDFTLTISLQPEDFATYYCQQYSSYPYTFGQ  
 GTKLEIK

Humanized A3 VL-CDR1:  
 KASQDVTAVA  
 SEQ ID NO: 32

Humanized A3 VL-CDR2:  
 WASTRLT  
 SEQ ID NO: 33

Humanized A3 VL-CDR3:  
 QQYSSYPYT  
 SEQ ID NO: 34

---

 SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 34

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 449

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: "Synthetic Sequence Humanized A1 human IgG1 heavy chain

&lt;400&gt; SEQUENCE: 1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Phe  
 20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ala Trp Ile Asn Thr Asn Thr Gly Glu Pro Arg Tyr Ala Glu Glu Phe  
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr  
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Ala Arg Asp Trp Asp Gly Ala Tyr Phe Phe Asp Tyr Trp Gly Gln Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe  
 115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu  
 130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp  
 145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu  
 165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser  
 180 185 190

-continued

---

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro  
 195 200 205  
 Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys  
 210 215 220  
 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro  
 225 230 235 240  
 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser  
 245 250 255  
 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp  
 260 265 270  
 Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn  
 275 280 285  
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val  
 290 295 300  
 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu  
 305 310 315 320  
 Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys  
 325 330 335  
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr  
 340 345 350  
 Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr  
 355 360 365  
 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu  
 370 375 380  
 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu  
 385 390 395 400  
 Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys  
 405 410 415  
 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu  
 420 425 430  
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly  
 435 440 445

Lys

<210> SEQ ID NO 2  
 <211> LENGTH: 214  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Humanized A1 human Kappa  
 light chain

&lt;400&gt; SEQUENCE: 2

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
 1 5 10 15  
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Ser Asn Asp  
 20 25 30  
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Phe Ala Thr Asn Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60  
 Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
 65 70 75 80



-continued

---

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Asp Tyr Ser Ser Pro Trp  
                     85                    90                    95  
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala  
                     100                    105                    110  
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly  
                     115                    120                    125  
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala  
                     130                    135                    140  
 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln  
                     145                    150                    155                    160  
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser  
                     165                    170                    175  
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
                     180                    185                    190  
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
                     195                    200                    205  
 Phe Asn Arg Gly Glu Cys  
                     210

<210> SEQ ID NO 3  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-VH

<400> SEQUENCE: 3

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1                    5                    10                    15  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Phe  
                     20                    25                    30  
 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                     35                    40                    45  
 Ala Trp Ile Asn Thr Asn Thr Gly Glu Pro Arg Tyr Ala Glu Glu Phe  
                     50                    55                    60  
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr  
                     65                    70                    75                    80  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                     85                    90                    95  
 Ala Arg Asp Trp Asp Gly Ala Tyr Phe Phe Asp Tyr Trp Gly Gln Gly  
                     100                    105                    110  
 Thr Leu Val Thr Val Ser Ser  
                     115

<210> SEQ ID NO 4  
 <211> LENGTH: 107  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-VL

<400> SEQUENCE: 4

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
 1                    5                    10                    15  
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Ser Asn Asp

-continued

---

20	25	30
Val Ala Trp Tyr Gln Gln Lys	Pro Gly Lys Ala Pro Lys Leu Leu Ile	
35	40	45
Tyr Phe Ala Thr Asn Arg Tyr	Thr Gly Val Pro Ser Arg Phe Ser Gly	
50	55	60
Ser Gly Tyr Gly Thr Asp Phe	Thr Leu Thr Ile Ser Ser Leu Gln Pro	
65	70	75
Glu Asp Phe Ala Thr Tyr Tyr	Cys Gln Gln Asp Tyr Ser Ser Pro Trp	
85	90	95
Thr Phe Gly Gln Gly Thr Lys	Val Glu Ile Lys	
100	105	

<210> SEQ ID NO 5  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-HC CDR1

<400> SEQUENCE: 5

Asn Phe Gly Met Asn  
 1 5

<210> SEQ ID NO 6  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-HC CDR2

<400> SEQUENCE: 6

Trp Ile Asn Thr Asn Thr Gly Glu Pro Arg Tyr Ala Glu Glu Phe Lys  
 1 5 10 15

Gly

<210> SEQ ID NO 7  
 <211> LENGTH: 10  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-HC CDR3

<400> SEQUENCE: 7

Asp Trp Asp Gly Ala Tyr Phe Phe Asp Tyr  
 1 5 10

<210> SEQ ID NO 8  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-LC-CDR1

<400> SEQUENCE: 8

Lys Ala Ser Gln Ser Val Ser Asn Asp Val Ala  
 1 5 10

<210> SEQ ID NO 9  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence

---

-continued

---

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence A1-LC-CDR2

&lt;400&gt; SEQUENCE: 9

Phe Ala Thr Asn Arg Tyr Thr  
1 5

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 9

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence A1-LC-CDR3

&lt;400&gt; SEQUENCE: 10

Gln Gln Asp Tyr Ser Ser Pro Trp Thr  
1 5

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 420

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence Human 5T4 antigen

&lt;400&gt; SEQUENCE: 11

Met Pro Gly Gly Cys Ser Arg Gly Pro Ala Ala Gly Asp Gly Arg Leu  
1 5 10 15  
Arg Leu Ala Arg Leu Ala Leu Val Leu Leu Gly Trp Val Ser Ser Ser  
20 25 30  
Ser Pro Thr Ser Ser Ala Ser Ser Phe Ser Ser Ser Ala Pro Phe Leu  
35 40 45  
Ala Ser Ala Val Ser Ala Gln Pro Pro Leu Pro Asp Gln Cys Pro Ala  
50 55 60  
Leu Cys Glu Cys Ser Glu Ala Ala Arg Thr Val Lys Cys Val Asn Arg  
65 70 75 80  
Asn Leu Thr Glu Val Pro Thr Asp Leu Pro Ala Tyr Val Arg Asn Leu  
85 90 95  
Phe Leu Thr Gly Asn Gln Leu Ala Val Leu Pro Ala Gly Ala Phe Ala  
100 105 110  
Arg Arg Pro Pro Leu Ala Glu Leu Ala Ala Leu Asn Leu Ser Gly Ser  
115 120 125  
Arg Leu Asp Glu Val Arg Ala Gly Ala Phe Glu His Leu Pro Ser Leu  
130 135 140  
Arg Gln Leu Asp Leu Ser His Asn Pro Leu Ala Asp Leu Ser Pro Phe  
145 150 155 160  
Ala Phe Ser Gly Ser Asn Ala Ser Val Ser Ala Pro Ser Pro Leu Val  
165 170 175  
Glu Leu Ile Leu Asn His Ile Val Pro Pro Glu Asp Glu Arg Gln Asn  
180 185 190  
Arg Ser Phe Glu Gly Met Val Val Ala Ala Leu Leu Ala Gly Arg Ala  
195 200 205  
Leu Gln Gly Leu Arg Arg Leu Glu Leu Ala Ser Asn His Phe Leu Tyr  
210 215 220  
Leu Pro Arg Asp Val Leu Ala Gln Leu Pro Ser Leu Arg His Leu Asp  
225 230 235 240

Leu	Ser	Asn	Asn	Ser	Leu	Val	Ser	Leu	Thr	Tyr	Val	Ser	Phe	Arg	Asn	
				245					250							255
Leu	Thr	His	Leu	Glu	Ser	Leu	His	Leu	Glu	Asp	Asn	Ala	Leu	Lys	Val	
			260				265								270	
Leu	His	Asn	Gly	Thr	Leu	Ala	Glu	Leu	Gln	Gly	Leu	Pro	His	Ile	Arg	
			275				280								285	
Val	Phe	Leu	Asp	Asn	Asn	Pro	Trp	Val	Cys	Asp	Cys	His	Met	Ala	Asp	
			290				295								300	
Met	Val	Thr	Trp	Leu	Lys	Glu	Thr	Glu	Val	Val	Gln	Gly	Lys	Asp	Arg	
305						310								315	320	
Leu	Thr	Cys	Ala	Tyr	Pro	Glu	Lys	Met	Arg	Asn	Arg	Val	Leu	Leu	Glu	
			325				330								335	
Leu	Asn	Ser	Ala	Asp	Leu	Asp	Cys	Asp	Pro	Ile	Leu	Pro	Pro	Ser	Leu	
			340				345								350	
Gln	Thr	Ser	Tyr	Val	Phe	Leu	Gly	Ile	Val	Leu	Ala	Leu	Ile	Gly	Ala	
			355				360								365	
Ile	Phe	Leu	Leu	Val	Leu	Tyr	Leu	Asn	Arg	Lys	Gly	Ile	Lys	Lys	Trp	
370						375								380		
Met	His	Asn	Ile	Arg	Asp	Ala	Cys	Arg	Asp	His	Met	Glu	Gly	Tyr	His	
385						390								395	400	
Tyr	Arg	Tyr	Glu	Ile	Asn	Ala	Asp	Pro	Arg	Leu	Thr	Asn	Leu	Ser	Ser	
			405				410								415	
Asn	Ser	Asp	Val													
				420												
<210> SEQ ID NO 12																
<211> LENGTH: 446																
<212> TYPE: PRT																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic Sequence Humanized A1 human IgG4m heavy chain																
<400> SEQUENCE: 12																
Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	
1				5				10							15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asn	Phe	
			20				25								30	
Gly	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35				40								45	
Ala	Trp	Ile	Asn	Thr	Asn	Thr	Gly	Glu	Pro	Arg	Tyr	Ala	Glu	Glu	Phe	
			50				55								60	
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	
65						70								75	80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85				90								95	
Ala	Arg	Asp	Trp	Asp	Gly	Ala	Tyr	Phe	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	
			100				105								110	
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	
			115				120								125	
Pro	Leu	Ala	Pro	Cys	Ser	Arg	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	
130						135								140		
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	
145						150								155	160	

&lt;400&gt; SEQUENCE: 13

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asn	Phe
			20					25					30		
Gly	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35					40					45			
Ala	Trp	Ile	Asn	Thr	Asn	Thr	Gly	Glu	Pro	Arg	Tyr	Ala	Glu	Glu	Phe
	50					55					60				

-continued

---

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr  
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Ala Arg Asp Trp Asp Gly Ala Tyr Phe Phe Asp Tyr Trp Gly Gln Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 14  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence A1-IgG4-VH-CDR1

<400> SEQUENCE: 14

Asn Phe Gly Met Asn  
 1 5

<210> SEQ ID NO 15  
 <211> LENGTH: 452  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 heavy chain  
 (muA3-huIgG1)

<400> SEQUENCE: 15

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Lys Gly  
 1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Thr Tyr  
 20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ala Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp  
 50 55 60

Ser Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Gln Ser Met  
 65 70 75 80

Leu Tyr Leu Gln Met Asn Asn Leu Lys Thr Glu Asp Thr Ala Met Tyr  
 85 90 95

Tyr Cys Val Arg Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr Trp  
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro  
 115 120 125

Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr  
 130 135 140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr  
 145 150 155 160

Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro  
 165 170 175

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr  
 180 185 190

Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn  
 195 200 205

-continued

---

His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser
210						215					220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
225					230					235					240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
				245					250					255	
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
			260					265					270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
		275					280					285			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
		290				295					300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
305					310					315					320
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro
				325					330					335	
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
			340					345					350		
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val
		355					360					365			
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
	370					375					380				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
385					390					395					400
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr
				405					410					415	
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val
			420					425					430		
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu
		435					440					445			
Ser	Pro	Gly	Lys												
		450													

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 122

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence Chimeric A3 VH

&lt;400&gt; SEQUENCE: 16

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Lys	Gly
1				5					10					15	
Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Asn	Thr	Tyr
		20						25					30		
Ala	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35					40					45			
Ala	Arg	Ile	Arg	Ser	Lys	Ser	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp
		50				55					60				
Ser	Val	Lys	Asp	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Gln	Ser	Met
65					70					75					80
Leu	Tyr	Leu	Gln	Met	Asn	Asn	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Met	Tyr
			85						90						95

-continued

---

Tyr Cys Val Arg Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr Trp  
                   100                  105                  110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
           115                  120

<210> SEQ ID NO 17  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VH-CDR1

<400> SEQUENCE: 17

Thr Tyr Ala Met Asn  
 1                  5

<210> SEQ ID NO 18  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VH-CDR2

<400> SEQUENCE: 18

Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser  
 1                  5                  10                  15

Val Lys Asp

<210> SEQ ID NO 19  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VH-CDR3

<400> SEQUENCE: 19

Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr  
 1                  5                  10

<210> SEQ ID NO 20  
 <211> LENGTH: 214  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 light chain  
 (muA3-huKappa)

<400> SEQUENCE: 20

Asp Ile Val Met Thr Gln Ser His Ile Phe Met Ser Thr Ser Val Gly  
 1                  5                  10                  15

Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asp Thr Ala  
                   20                  25                  30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile  
           35                  40                  45

Tyr Trp Ala Ser Thr Arg Leu Thr Gly Val Pro Asp Arg Phe Thr Gly  
           50                  55                  60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Gln Ser  
 65                  70                  75                  80

Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Tyr Pro Tyr  
           85                  90                  95



-continued

---

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala  
 100 105 110  
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly  
 115 120 125  
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala  
 130 135 140  
 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln  
 145 150 155 160  
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser  
 165 170 175  
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
 180 185 190  
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
 195 200 205  
 Phe Asn Arg Gly Glu Cys  
 210

<210> SEQ ID NO 21  
 <211> LENGTH: 107  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VL

<400> SEQUENCE: 21

Asp Ile Val Met Thr Gln Ser His Ile Phe Met Ser Thr Ser Val Gly  
 1 5 10 15  
 Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asp Thr Ala  
 20 25 30  
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Trp Ala Ser Thr Arg Leu Thr Gly Val Pro Asp Arg Phe Thr Gly  
 50 55 60  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Gln Ser  
 65 70 75 80  
 Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Tyr Pro Tyr  
 85 90 95  
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys  
 100 105

<210> SEQ ID NO 22  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VL-CDR1

<400> SEQUENCE: 22

Lys Ala Ser Gln Asp Val Asp Thr Ala Val Ala  
 1 5 10

<210> SEQ ID NO 23  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Chimeric A3 VL-CDR2

-continued

&lt;400&gt; SEQUENCE: 23

Trp Ala Ser Thr Arg Leu Thr  
1 5

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 9

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence Chimeric A3 VL-CDR3

&lt;400&gt; SEQUENCE: 24

Gln Gln Tyr Ser Ser Tyr Pro Tyr Thr  
1 5

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 452

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Sequence Humanized A3 human IgG1 heavy chain

&lt;400&gt; SEQUENCE: 25

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15  
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Thr Tyr  
20 25 30  
Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45  
Ala Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp  
50 55 60  
Ser Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ala Lys Asn Ser  
65 70 75 80  
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr  
85 90 95  
Tyr Cys Val Arg Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr Trp  
100 105 110  
Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro  
115 120 125  
Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr  
130 135 140  
Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr  
145 150 155 160  
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro  
165 170 175  
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr  
180 185 190  
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn  
195 200 205  
His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser  
210 215 220  
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu  
225 230 235 240  
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu  
245 250 255

-continued

---

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser  
                   260                  265                  270

His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu  
                   275                  280                  285

Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr  
                   290                  295                  300

Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn  
                   305                  310                  315                  320

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro  
                   325                  330                  335

Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln  
                   340                  345                  350

Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val  
                   355                  360                  365

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val  
                   370                  375                  380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro  
                   385                  390                  395                  400

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr  
                   405                  410                  415

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val  
                   420                  425                  430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu  
                   435                  440                  445

Ser Pro Gly Lys  
                   450

<210> SEQ ID NO 26  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VH

<400> SEQUENCE: 26

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1                  5                  10                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Thr Tyr  
                   20                  25                  30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                  40                  45

Ala Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp  
                   50                  55                  60

Ser Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ala Lys Asn Ser  
                   65                  70                  75                  80

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr  
                   85                  90                  95

Tyr Cys Val Arg Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr Trp  
                   100                  105                  110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
                   115                  120

<210> SEQ ID NO 27

---

-continued

---

<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VH-CDR1

<400> SEQUENCE: 27

Thr Tyr Ala Met Asn  
1 5

<210> SEQ ID NO 28  
<211> LENGTH: 19  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VH-CDR2

<400> SEQUENCE: 28

Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser  
1 5 10 15

Val Lys Asp

<210> SEQ ID NO 29  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VH-CDR3

<400> SEQUENCE: 29

Gln Trp Asp Tyr Asp Val Arg Ala Met Asn Tyr  
1 5 10

<210> SEQ ID NO 30  
<211> LENGTH: 214  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 human Kappa  
light chain

<400> SEQUENCE: 30

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Asp Thr Ala  
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
35 40 45

Tyr Trp Ala Ser Thr Arg Leu Thr Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Ser Tyr Pro Tyr  
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala  
100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly  
115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala  
130 135 140

---

-continued

---

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln  
145 150 155 160  
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser  
165 170 175  
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
180 185 190  
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
195 200 205  
Phe Asn Arg Gly Glu Cys  
210

<210> SEQ ID NO 31  
<211> LENGTH: 107  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VL

<400> SEQUENCE: 31

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15  
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Asp Thr Ala  
20 25 30  
Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
35 40 45  
Tyr Trp Ala Ser Thr Arg Leu Thr Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60  
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80  
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Ser Tyr Pro Tyr  
85 90 95  
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys  
100 105

<210> SEQ ID NO 32  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VL-CDR1

<400> SEQUENCE: 32

Lys Ala Ser Gln Asp Val Asp Thr Ala Val Ala  
1 5 10

<210> SEQ ID NO 33  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VL-CDR2

<400> SEQUENCE: 33

Trp Ala Ser Thr Arg Leu Thr  
1 5

<210> SEQ ID NO 34  
<211> LENGTH: 9

-continued

---

<212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Sequence Humanized A3 VL-CDR3

<400> SEQUENCE: 34

Gln Gln Tyr Ser Ser Tyr Pro Tyr Thr  
 1 5

---

We Claim:

1. An antibody-drug conjugate of the formula:

Ab-(LU-D)<sub>p</sub>

or a pharmaceutically acceptable salt thereof wherein;

(a) Ab is an anti-5T4 antibody or antigen binding portion thereof, comprising:

- (i) a VH CDR1 region as shown in SEQ ID NO: 5,
- (ii) a VH CDR2 region as shown in SEQ ID NO: 6,
- (iii) a VH CDR3 region as shown in SEQ ID NO: 7;
- (iv) a VL CDR1 region as shown in SEQ ID NO: 8,
- (v) a VL CDR2 region as shown in SEQ ID NO: 9, and
- (vi) a VL CDR3 region as shown in SEQ ID NO: 10,

(b) LU is a linker unit selected from the group consisting of maleimidocaproyl and maleim idocaproyl-Val-Cit,

(c) p is an integer from about 1 to about 8, and

(d) D is a Drug unit selected from the group consisting of MMAE, MMAD, and MMAF.

2. The antibody-drug conjugate of claim 1 wherein said anti-5T4 antibody or antigen binding portion thereof, comprises the VH region of SEQ ID NO: 3 and the VL region of SEQ ID NO: 4.

3. The antibody-drug conjugate of claim 2 wherein said anti-5T4 antibody or antigen binding portion thereof, consists of a heavy chain having SEQ ID NO:1 and a light chain having SEQ ID NO: 2.

4. The antibody-drug conjugate of claim 3, wherein said antibody or antigen binding portion thereof, recognizes an epitope on human 5T4 antigen, wherein said epitope comprises amino acid residues 173-258 and 282-361 of the amino acid sequence of SEQ ID NO:11.

5. The antibody-drug conjugate of claim 1 wherein:

(a) said anti-5T4 antibody consists of a heavy chain having SEQ ID NO:1 and a light chain having SEQ ID NO: 2,

(b) said LU is maleimidocaproyl,

(c) said Drug is MMAF, and

(d) p is 4.

6. A pharmaceutical composition comprising the antibody-drug conjugate of claim 1 and a pharmaceutically acceptable carrier.

7. A method of treating 5T4-positive cancer in a patient in need thereof, comprising administering to said patient the antibody-drug conjugate according to claim 1.

8. The method of claim 7 wherein said cancer is selected from the group consisting of colorectal, breast, pancreatic, and non-small cell lung carcinomas.

\* \* \* \* \*