

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2001222619 C1**

(54) Title
Processes for preparing clarithromycin polymorphs

(51) International Patent Classification(s)
A61K 31/70 (2006.01) **C07H 17/08** (2006.01)
C07H 1/00 (2006.01)

(21) Application No: **2001222619** (22) Date of Filing: **2000.12.15**

(87) WIPO No: **WO01/51059**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/175,477	2000.01.11	US

(43) Publication Date: **2001.07.24**

(43) Publication Journal Date: **2001.10.04**

(44) Accepted Journal Date: **2005.04.28**

(44) Amended Journal Date: **2006.01.12**

(71) Applicant(s)
Teva Pharmaceutical Industries Ltd.

(72) Inventor(s)
Masarwa, Basem; Lifshitz, Igor; Avrutov, Ilya; Schwartz, Edi

(74) Agent / Attorney
Shelston IP, Level 21 60 Margaret Street, Sydney, NSW, 2000

(56) Related Art
US 5,844,105

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2001 (19.07.2001)

PCT

(10) International Publication Number
WO 01/51059 A1

(51) International Patent Classification⁷: **A61K 31/70,**
C07H 1/00, 17/08

SCHWARTZ, Edi [IL/IL]; Remez 70, Rechovot (IL).
MASARWA, Basem [IL/IL]; P.O. Box 3857, 40400 Taibe (IL).

(21) International Application Number: **PCT/US00/33844**

(22) International Filing Date:
15 December 2000 (15.12.2000)

(74) Agents: **BRAINARD, Charles, R.** et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/175,477 11 January 2000 (11.01.2000) **US**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except BB, US*):
TEVA PHARMACEUTICAL INDUSTRIES LTD. [IL/IL]; Basel Street 5, P.O. Box 3190, 49131 Petah Tikva (IL).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for BB only*): **TEVA PHARMACEUTICALS USA, INC.** [US/US]; 1090 Horsham Road, P.O. Box 1090, North Wales, PA 19454-1090 (US).

Published:
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LIFSHITZ, Igor** [IL/IL]; Shpinoza 9/11, Petach Tikva (IL). **AVRUTOV, Ilya** [IL/IL]; Beery Street 50, 42842 Bat Hefer (IL).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/51059 A1

(54) Title: **PROCESSES FOR PREPARING CLARITHROMYCIN POLYMORPHS**

(57) Abstract: The present invention relates to processes for converting clarithromycin Form I to clarithromycin Form II, which include slurrying clarithromycin the Form I in water. The present invention also relates to processes for the preparation of clarithromycin Form II by converting erythromycin A to clarithromycin and thereafter converting clarithromycin Form I to clarithromycin Form II by slurrying.

PROCESSES FOR PREPARING CLARITHROMYCIN POLYMORPHS

CROSS REFERENCE TO RELATED APPLICATION

The present application claims the benefit of U.S. Provisional Application No. 60/175,477 filed on January 11, 2000.

5 FIELD OF THE INVENTION

The invention relates to methods for making polymorphic Form II of clarithromycin via slurring polymorphic Form I of clarithromycin in water.


By slurrying in water, clarithromycin Form I can be converted to clarithromycin Form II. The invention also relates to clarithromycin Form II prepared by slurrying 10 polymorphic Form I in water and pharmaceutical compositions containing the same.

BACKGROUND OF THE INVENTION

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

6-0-methyl erythromycin A (clarithromycin) is a semisynthetic macrolide antibiotic related to erythromycin A. Clarithromycin exhibits excellent antibacterial activity against gram-positive bacteria, some gram-negative bacteria, anaerobic bacteria, *Mycoplasma*, and *Chlamydia*. It is stable under acidic conditions and is efficacious when administered orally. Clarithromycin is useful therapy for infections of the upper respiratory tract in children and adults. Clarithromycin is stable under acidic conditions and is efficacious when administered orally.

The chemical structure of clarithromycin is:



Several crystal forms of clarithromycin and/or solvates of clarithromycin, "Form 0," "Form I", and "Form II" have been identified as indicated in U.S. Patent No.

5,945,405. The crystal form of clarithromycin is typically identified by the powder x-ray diffraction patterns. Different crystalline forms of clarithromycin may have different thermal stability, cost of preparation, dissolution characteristics and bioavailability.

Various methods of preparing clarithromycin described in the patent literature have been reported to result in different forms of clarithromycin. Purification of crude clarithromycin has been reported to convert one form of clarithromycin to another form. For example, methods in which the compound is purified by recrystallization from ethanol, have been reported to result in the initial formation of Form 0 solvate (see U.S. Patent No. 5,945,405) or in the initial formation of Form I (See U.S. Patent No. 5,858,986). The Form 0 solvate may be used as a therapeutic agent, as described in U.S. Patent No. 5,945,405. The '405 patent discloses that the Form 0 solvate may be converted to the non-solvated Form I by removing the solvent by drying at a temperature of from about 0°C to about 50°C. Another form, Form II, is reported to be relatively thermodynamically stable compared to Form I. The clarithromycin currently marketed in the United States under the trademark Biaxin® is formulated using Form II.

Several methods of converting clarithromycin Form 0 or Form I to Form II have been described. One such method, as described in U.S. Patent No. 4,945,405 and U.S. Patent No. 5,858,986, is to heat Form 0 under vacuum at a temperature of between about 70°C and 110°C. According to this patent, the Form 0 solvate first converts to Form I clarithromycin and then to Form II. This method is described in U.S. Patent No. 4,945,405 and U.S. Patent No. 5,858,986. However, this vacuum drying step of converting Form 0 to Form II is expensive in both energy and material handling. Clarithromycin Form II has been reported to be formed when Form I is crystallized from various solvents, including ethanol and water (US 5,844,105).

SUMMARY OF THE INVENTION

The invention relates to methods for making polymorphic forms of clarithromycin via slurring clarithromycin Form I in water. By slurring in water, clarithromycin Form I

can be converted to clarithromycin Form II. The invention also relates clarithromycin form II formed by this slurring process and pharmaceutical compositions containing the same.

The present invention relates to a process for converting clarithromycin Form I to clarithromycin Form II, which includes slurring clarithromycin Form I in water. The present invention also relates to processes for the preparation of clarithromycin Form II by converting erythromycin A to clarithromycin and thereafter, converting clarithromycin Form I to clarithromycin Form II by slurring.

According to a first aspect of the invention there is provided a process for converting polymorphic Form I of clarithromycin into polymorphic Form II of clarithromycin, which comprises:

- (a) slurring a solid comprising clarithromycin Form I in water to form clarithromycin Form II; and
- (b) isolating the clarithromycin Form II.

According to a second aspect of the invention there is provided a process for converting clarithromycin Form I to clarithromycin Form II comprising the steps of:

- (a) slurring clarithromycin Form I in water to form clarithromycin Form II; and
- (b) isolating the clarithromycin Form II.

According to a third aspect of the invention there is provided a process for preparing clarithromycin Form II comprising the steps of:

- (a) converting erythromycin A to clarithromycin Form I ;
- (b) slurring the clarithromycin Form I in water to form clarithromycin Form II; and
- (c) isolating the clarithromycin Form II.

According to a fourth aspect of the invention there is provided clarithromycin Form II, formed by slurring clarithromycin Form I in water.

According to a fifth aspect of the invention there is provided a pharmaceutical composition comprising

- a therapeutically effective amount of clarithromycin Form II formed by slurring clarithromycin Form I in water, and
- at least one pharmaceutically acceptable carrier.

Unless the context clearly requires otherwise, throughout the description and the claims, the words 'comprise', 'comprising', and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".


5 DETAILED DESCRIPTION OF THE INVENTION

We have found that clarithromycin Form I undergoes transformation to Form II by simply slurring the Form I clarithromycin with water. The present process for converting clarithromycin Form I to Form II is simpler and less expensive than previously described methods of converting Form I to Form II, which require for
10 example, the use of organic solvents such as ethanol.

Form II prepared by this process can be formulated into pharmaceutical compositions that include a therapeutically effective amount of clarithromycin Form II and one or more pharmaceutically acceptable carriers. Those pharmaceutical compositions can be administered as a therapeutically effective antibiotic. Each of these
15 aspects of the present invention is described in further detail below.

The terms "6-O-methylerythromycin A" and "clarithromycin" are used interchangeably herein and are meant to include clarithromycin in any crystalline form or mixtures thereof, as well as amorphous solids, syrups, or semisolids comprising 6-O-methylerythromycin A in any state of purity, unless specified otherwise, or as the
20 context requires.

The term "Form I" is used herein to mean the crystal Form I of clarithromycin. Form I has been characterized by 2-theta angle positions in the powder x-ray diffraction pattern of 5.2 ± 0.2 , 6.7 ± 0.2 , 10.2 ± 0.2 , 12.3 ± 0.2 , 14.2 ± 0.2 , 15.4 ± 0.2 , 15.7 ± 0.2 , and 16.4 ± 0.2 . Form I has also been characterized by an exothermic transition at 132.2°C by
25 differential scanning calorimetry, and by endothermic peaks at 223.4°C , and 283.3°C ,



followed by an exothermic peak at 306.9°C.

The term "Form II" is used herein to mean the crystal Form II of clarithromycin. Form II has been characterized by 2-theta angle positions in the powder x-ray diffraction pattern of 8.5°±0.2, 9.5°±0.2, 10.8°±0.2, 11.5°±0.2, 11.9°±0.2, 12.4°±0.2, 13.7°±0.2, 14.1°±0.2, 15.2°±0.2, 16.5°±0.2, 16.9°±0.2, 17.3°±0.2, 18.1°±0.2, 18.4°±0.2, 19.0°±0.2, 19.9°±0.2, and 20.5°±0.2. Form II has also been characterized by melting at 223.4°C, and by an endothermic peak at 283.3°C by differential scanning calorimetry.

The term "Form 0" is used herein to mean the crystal Form 0 or Form 0 solvate of 6-O-methylerythromycin A. Form 0 solvate is characterized by 2-theta angle positions in the powder x-ray diffraction pattern of 4.6°±0.2, 6.5°±0.2, 7.6°±0.2, 9.2°±0.2, 10.2°±0.2, 11.0°±0.2, 11.6°±0.2, 12.5°±0.2, 13.8°±0.2, 14.8°±0.2, 17.0°±0.2, 18.2°±0.2, 18.9°±0.2 and 19.5°±0.2. This pattern may be somewhat varied depending on the solvent. For example a Form 0 ethanolate may be characterized by 2-theta angle positions of 4.7°±0.2, 6.6°±0.2, 7.7°±0.2, 9.3°±0.2, 10.4°±0.2, 11.1°±0.2, 11.9°±0.2, 12.7°±0.2, 13.9°±0.2, 15.0°±0.2, 17.2°±0.2, 18.5°±0.2, 19.1°±0.2, 19.7°±0.2, 23.1°±0.2 and 24.0°±0.2.

Formation of clarithromycin Form II

The present invention relates to processes for converting clarithromycin Form I to clarithromycin Form II, which include slurrying clarithromycin Form I with water. The present invention also relates to a process of preparing clarithromycin Form II from erythromycin A, which includes slurrying clarithromycin Form I in water. In one embodiment, this method includes first converting erythromycin A to clarithromycin and thereafter slurrying clarithromycin Form I with water to form clarithromycin Form II. There are several methods by which clarithromycin may be formed from erythromycin A, as described further below. In certain of these methods, the initial form of clarithromycin that is formed is Form I. In other methods, the initial form of clarithromycin that is formed is not Form I, but rather another form that is subsequently converted to Form I.

As used herein, the terms "slurrying" and "slurried" are intended to include stirring particles in a liquid, preferably water.

In the present invention, a wet solid of clarithromycin is obtained, for example by treating clarithromycin with ethanol, filtering and washing the precipitate with ethanol. The wet solid is slurried with water, preferably at ambient temperature, for about 1 hour. The slurrying step may include stirring the clarithromycin and water.

5 After clarithromycin Form I is converted to clarithromycin Form II by slurrying, the resulting crystalline clarithromycin Form II may then be isolated, preferably by filtration and dried in an oven, preferably a vacuum oven, at a temperature of between about ambient temperature and about 70°C, preferably from about 40°C to about 60°C, even more preferably, about 50°C. Preferably, the drying takes place under vacuum.

10 In the slurrying step of any of the embodiments that involve converting Form I to Form II described herein, clarithromycin Form I is slurried in about 2 to about 25 ml of water per gram of clarithromycin Form I. More preferably clarithromycin Form I is slurried in about 3 to about 10 ml of water per gram of clarithromycin Form I. Most preferably clarithromycin Form I is slurried in about 5 ml of water per gram of
15 clarithromycin Form I.

For purposes of this specification, ambient temperature is from about 20°C to about 25°C.

20 The term "treating" as used herein refers to suspending, dissolving, washing, mixing, crystallizing or recrystallizing clarithromycin in any of the solvents described above.

As indicated above, processes for preparing clarithromycin Form II according to the present invention may include first converting erythromycin A to clarithromycin prior to slurrying the clarithromycin Form I with water to form clarithromycin Form II. This
25 conversion of erythromycin A to clarithromycin may be performed by any method used by those in the art. In general, clarithromycin is prepared by methylation of the 6-hydroxy group of erythromycin A. However, in the conversion process it is necessary to protect various groups, especially hydroxy groups at positions of erythromycin A, which are potentially reactive with alkylating agents, prior to alkylation of the 6-hydroxy group.
30 Examples of methods of preparing clarithromycin are described for example, in U.S.

Patent Nos. 4,331,803, 4,670,549, 4,672,109, 4,990,602, 5,858,986, 5,844,105 and 5,945,405, which are each herein incorporated by reference in their entirety. By way of example, U.S. Patent Nos. 4,990,602 and 5,858,986 describe a method of preparing clarithromycin from erythromycin A by protecting the 9-oxo group with a substituted oxime group, protection of the C-2' and C-4'' hydroxy groups, methylation of the C-6 hydroxy group, and deoxygenation and removal of the protecting groups. Removal of the protecting groups and deoxygenation of 6-O-methyl-2', 4''-bis(trimethylsilyl)-erythromycin A 9-O-(2-methoxyprop-2-yl)oxime (or other protected oxime) are then accomplished by treatment with acid and deoxygenating agent to give clarithromycin. Examples of suitable deoxygenating agents include inorganic sulfur oxide compounds such as sodium hydrogen sulfite, sodium pyrosulfate, sodium thiosulfate, sodium sulfite, sodium hydrosulfite, sodium metabisulfite, sodium bisulfite, sodium dithionate, potassium hydrogen sulfite, potassium thiosulfate, potassium metabisulfite and the like. Particularly preferred deoxygenating agents are sodium metabisulfite and sodium bisulfite and the like. Deprotection may occur prior to deoxygenation or deprotection and deoxygenation may be accomplished in a single step ("one-pot" synthesis) by treatment with acid, such as formic acid, and deoxygenating agent.

Various other methods of converting erythromycin A to clarithromycin that presently exist or methods that may be contemplated in the future may also be used.

Clarithromycin Form II formed by slurrying Form I

The present invention also relates to clarithromycin Form II prepared by slurrying clarithromycin Form I in water.

Pharmaceutical compositions including clarithromycin Form II formed by slurrying Form I

Another aspect of the present invention provides pharmaceutical compositions including a therapeutically effective amount of clarithromycin Form II prepared by slurrying Form I in water or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable carriers.

The pharmaceutical compositions of this invention are administered to humans and other animals by any acceptable route within sound medical judgment, such as orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray. Preferably, the pharmaceutical compositions of this invention are formulated for oral administration in solid or liquid form, for parenteral injection, or for rectal administration.

The term "parenteral" administration as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of this invention for parenteral injection include for example pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

The pharmaceutical compositions of the present invention may also contain one or more adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include within the pharmaceutical composition one or more isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the clarithromycin Form II, it is desirable to slow the absorption of the clarithromycin Form II from subcutaneous or

intramuscular injection. This may be accomplished for example by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the clarithromycin Form II then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed

- 5 absorption of a parenterally administered clarithromycin Form II form may be accomplished by dissolving or suspending the clarithromycin Form II in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the clarithromycin Form II in biodegradable polymers such as polylactide-polyglycolide.

- Depending upon the ratio of drug to polymer and the nature of the particular polymer
10 employed, the rate of clarithromycin Form II release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the clarithromycin Form II in liposomes or microemulsions that are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions, which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

- 5 Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the clarithromycin Form II is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such pharmaceutically acceptable excipients or carriers include sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example,
- 10 carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such
- 15 as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

- Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular
- 20 weight polyethylene glycols and the like.

- Solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition such that they release the clarithromycin Form II only, or preferentially, in a certain part of the
- 25 intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

 The active compounds can also be in micro-encapsulated form, if appropriate, with one or more excipients, such as the above-mentioned excipients.

Liquid dosage form for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the clarithromycin Form II, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Oral compositions according to the present invention may also include one or more adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the clarithromycin Form II.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods of forming liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976),

p. 33 et seq.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The clarithromycin Form II is mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers, or propellants, which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of clarithromycin Form II in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the clarithromycin Form II that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend on various factors such as the activity of the particular compound, the route of administration, the severity of the condition being treated, the size of the patient and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the clarithromycin Form II at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. Thus, an appropriate therapeutically effective amount may be determined by a doctor, veterinarian or other healthcare professional having experience in determining appropriate dosages for treatment.

The following examples are provided to enable one skilled in the art to practice the invention and are merely illustrative of the invention. The examples should not be read as limiting the scope of the invention as defined in the claims.

Example 1

Clarithromycin (5g, assay ~ 95%) was dissolved in 60 milliliters of ethanol at reflux. The solution was then cooled to about 10°C and the precipitate was filtered and washed with 5 milliliters of cold ethanol. 6 grams of a wet solid were obtained and then dried at 50°C under vacuum to obtain dry product, which powder x-ray diffraction analysis (PXRD) indicated was clarithromycin crystalline Form I. The product so obtained was slurried with stirring in 25 milliliters of water at ambient temperature for about 1 hour and the solid substance was then filtered. Powder x-ray diffraction analysis was performed on the wet product and the product

was determined to be polymorph II. The product was dried at temperature of about 50°C under vacuum. The final product was 4.5 grams of pure clarithromycin (assay above 98%) polymorphic Form II.

5 Example 1 shows a particular embodiment of the present invention, which includes converting clarithromycin Form I to Form II by slurring. It would be evident to those in the art based on the present disclosure that similar processes may be employed in which the process parameters are varied from those in the examples, for example if different amounts, temperatures, isolation techniques, etc. are employed, as long as the yielded product is
10 clarithromycin Form II.

 The present invention provides methods for converting clarithromycin Form I to clarithromycin Form II by slurring in water, clarithromycin Form II prepared by such methods, and pharmaceutical compositions including clarithromycin Form II prepared by slurring Form I. Although the present invention has been described with respect to certain exemplary
15 embodiments, there are many other variations of the above-described embodiments, which will be apparent to those skilled in the art, even where elements or steps have not explicitly been designated as exemplary. It is understood that these modifications are within the teaching of the present invention.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A process for converting polymorphic Form I of clarithromycin into polymorphic Form II of clarithromycin, which comprises:
 - (a) slurring a solid comprising clarithromycin Form I in water to form clarithromycin Form II; and
 - (b) isolating the clarithromycin Form II.
2. A process for converting clarithromycin Form I to clarithromycin Form II comprising the steps of:
 - (a) slurring clarithromycin Form I in water to form clarithromycin Form II; and
 - (b) isolating the clarithromycin Form II.
3. The process of claim 2, wherein the clarithromycin Form I is slurried in about 2 to about 25 ml of water per gram of clarithromycin Form I.
4. The process of claim 2, wherein the clarithromycin Form I is slurried in about 3 to about 10 ml of water per gram of clarithromycin Form I.
5. The process of claim 2, wherein the slurring takes place at about ambient temperature.
6. The process of claim 2, further comprising drying the clarithromycin Form II that results from slurring the clarithromycin Form I in water.
7. The process of claim 6, wherein the drying takes place under vacuum.
8. A process for preparing clarithromycin Form II comprising the steps of:
 - (a) converting erythromycin A to clarithromycin Form I;
 - (b) slurring the clarithromycin Form I in water to form clarithromycin Form II; and
 - (c) isolating the clarithromycin Form II.
9. The process of claim 8, wherein the clarithromycin Form I is slurried in about 2 to about 25 ml of water per gram of clarithromycin Form I.

10. The process of claim 8, wherein the clarithromycin Form I is slurried in about 3 to about 10 ml of water per gram of clarithromycin Form I.
11. The process of claim 8, wherein the slurrying takes place at about ambient temperature.
- 5 12. The process of claim 8, further comprising drying the clarithromycin Form II that results from slurrying the clarithromycin Form I in water.
13. The process of claim 12, wherein the drying takes place under vacuum.
14. Clarithromycin Form II, formed by slurrying clarithromycin Form I in water.
15. A pharmaceutical composition comprising
 - 10 a therapeutically effective amount of clarithromycin Form II formed by slurrying clarithromycin Form I in water, and
 - at least one pharmaceutically acceptable carrier.
16. The process of claim 1, wherein the solid in step a) is obtained by a process comprising the steps of:
 - 15 (i) converting erythromycin A to clarithromycin Form I to form a solid comprising clarithromycin Form I; and
 - (ii) washing the solid in ethanol.
17. The process of claim 16, wherein the solid is wet.
18. The process of claim 1, wherein the solid is slurried in water for about one hour in
 - 20 step a).
19. The process of claim 18, wherein the solid is slurried in water for about one hour at a temperature of about 20°C to about 25°C in step a).
20. The process of claim 1, wherein the solid is slurried in about 2 ml to about 25 ml of water per gram of the clarithromycin Form I.

21. A process for converting polymorphic Form I of clarithromycin into polymorphic Form II of clarithromycin, substantially as herein described with reference to any one of the examples but excluding comparative examples.
22. A process for converting clarithromycin Form I to clarithromycin Form II,
5 substantially as herein described with reference to any one of the examples but excluding comparative examples.
23. A process for preparing clarithromycin Form II, substantially as herein described.
24. A pharmaceutical composition comprising a therapeutically effective amount of clarithromycin Form II, substantially as herein described with reference to any one of the
10 examples but excluding comparative examples.

DATED this 27th day of July, 2005

Shelston IP

Attorneys for: TEVA PHARMACEUTICAL INDUSTRIES LTD