




- (51) International Patent Classification:
C07K 16/28 (2006.01) *G01N 33/53* (2006.01)
C07K 16/30 (2006.01) *A61P 35/00* (2006.01)
A61K 39/395 (2006.01)
- (21) International Application Number: PCT/IL2012/050402
- (22) International Filing Date: 10 October 2012 (10.10.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
PCT/IL2011/000808
11 October 2011 (11.10.2011) IL
- (71) Applicant: TEL HASHOMER MEDICAL RESEARCH INFRASTRUCTURE AND SERVICES LTD. [IL/IL]; The Chaim Sheba Medical Center Tel Hashomer, 52621 Ramat Gan (IL).
- (72) Inventor; and
(71) Applicant : CCAM BIOTHERAPEUTICS LTD. [IL/IL]; Tel-Hai Industrial Park, P.O. Box 408, 11013 Kiryat-Shmona (IL).
- (72) Inventors: MARKEL, Gal; 50/6 Bnei Dan Street, 62264 Tel Aviv (IL). BEN MOSHE, Tehila; Koranit, 20181 Misgav (IL). SAPIR, Yair; Hadekel 16, 42860 Burgata (IL). MANDEL, Ilana; Hate'ena 3, 29066 Kiryat Yam (IL). SCHACHTER, Jacob; 258 Derech Ben Gurion, 53326 Givatayim (IL).
- (74) Agents: WEBB & CO et al.; P.O. Box 2189, 76121 Rehovot (IL).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

[Continued on next page]

(54) Title: ANTIBODIES TO CARCINOEMBRYONIC ANTIGEN-RELATED CELL ADHESION MOLECULE (CEACAM)

Figure 9



(57) Abstract: The present invention provides antibodies, as well as molecules having at least the antigen-binding portion of an antibody, recognizing a specific epitope of the protein CEACAM1 and optionally binds also other subtypes of the CEACAM protein family. Disclosed antibodies and antibody fragments are characterized by specific CDR sequences. Methods of production and use in therapy and diagnosis, of such antibodies and antibody fragments are also provided.



EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

— with international search report (Art. 21(3))

— with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

— with sequence listing part of description (Rule 5.2(a))

**ANTIBODIES TO CARCINOEMBRYONIC ANTIGEN-RELATED CELL
ADHESION MOLECULE (CEACAM)**

FIELD OF THE INVENTION

5 The present invention relates to therapeutic and diagnostic antibodies, useful in diseases involving Carcinoembryonic Antigen-Related Cell Adhesion Molecule (CEACAM), expression, activation or function. In particular, the present invention provides antibodies having specific complementarity determining regions (CDRs) and improved properties over other antibodies which recognize CEACAM1.

10

BACKGROUND OF THE INVENTION

 The transmembrane protein Carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM1, also known as biliary glycoprotein (BGP), CD66a and C-CAM1), is a member of the carcinoembryonic antigen family (CEA) that also belongs to the immunoglobulin superfamily. CEACAM1 interacts with other known CEACAM proteins, including CD66a (CEACAM1), CD66e (CEACAM6) and CD66e (CEACAM5, CEA) proteins. It is expressed on a wide spectrum of cells, ranging from epithelial cells to those of hemopoietic origin (e.g. immune cells).

 Many different functions have been attributed to the CEACAM1 protein. It was shown that the CEACAM1 protein is over expressed in some carcinomas of colon, prostate, as well as other types of cancer. Additional data support the central involvement of CEACAM1 in angiogenesis and metastasis. CEACAM1 also plays a role in the modulation of innate and adaptive immune responses. For example, CEACAM1 was shown to be an inhibitory receptor for activated T cells contained within the human intestinal epithelium (WO99/52552 and Morales et al. J. Immunol. 1999, 163, 1363-1370). Additional reports have indicated that CEACAM1 engagement either by T Cell Receptor cross-linking with Monoclonal antibodies (mAbs) or by Neisseria gonorrhoeae Opa proteins inhibits T cell activation and proliferation.

 Melanoma is a malignancy of pigment-producing cells (melanocytes), responsible for 75% of skin cancer-related mortality worldwide, mainly due to extensive metastasis. Metastatic melanoma (MM) responds feebly to most anticancer regimens, and mean overall survival mean for patients with MM is 8.5 months. There is evidence that overexpression of CEACAM1 can be correlated with poor prognosis and is detected in the majority of metastatic melanoma cases. CEACAM1 is rarely expressed by normal melanocytes, but

frequently found on melanoma cells. CEACAM1 expression on primary cutaneous melanoma lesions strongly predicts the development of metastatic disease with poor prognosis. Moreover, increased CEACAM1 expression was observed on NK cells derived from some patients with metastatic melanoma compared with healthy donors.

5 Evidence indicates that CEACAM1 may have an important role in virus infections. For example, Markel et al. (J. Clinical Investigation 2002, 110, 943-953) demonstrated that lymphocytes isolated from the deciduae of CMV-infected patients express the CEACAM1 protein in increased levels. The increased CEACAM1 expression on the decidual lymphocytes might diminish the local immune response and serve as another mechanism
10 developed by the virus to avoid recognition and clearance primarily by activated decidual lymphocytes. Albarran-Somoza et al. (Journal of Histochemistry & Cytochemistry 2006, 54, 1393), who studied the protein expression pattern of CEACAM1 in cervical cancer and precursor lesions in the context of human papillomavirus (HPV) infection, showed that CEACAM1 immunostaining is significantly increased in high-grade squamous intraepithelial
15 lesions (SIL) in comparison with low-grade SIL and normal cervical tissues. The authors suggested that CEACAM1 upregulation may be related to integration of HPV DNA in high-grade SIL and that CEACAM1 may be an important biological marker in SIL and cervical cancer progression. Altogether this evidence indicates that CEACAM1 plays an important role in various viral infections. In addition, CEACAM1 over expression may serve as marker
20 of various viral infections.

WO2007/063424 and U.S. Patent Application No. 20070110668 disclose methods for regulating the immune system, and in particular methods for the regulation of a specific immune response, including the regulation of lymphocyte activity. These methods comprise both the negative and positive modulation of CEACAM1 protein function.

25 U.S. Patent Application No. 20070071758 teaches methods and compositions for enhancing the efficacy of tumor-infiltrating lymphocyte (TIL) therapy in the treatment of cancer by negatively modulating the activity of the CEACAM1 protein, such as for example, by using an immunoglobulin specific for CEACAM1.

U.S. Patent Application No. 20080108140 discloses methods of modulating specific
30 immune responses to create a protective immunity in the treatment of autoimmune diseases and diseases requiring the transplantation of tissue. In particular, it relates to the suppression of immune responses in a targeted fashion, by increasing the functional concentration of the CEACAM1 protein in the target tissue.

U.S. Patent Application No. 20040047858 discloses specific antibodies which are capable of modulating T cell activity via CEACAM1 and uses thereof in treating immune response related diseases (e.g. graft versus host disease, autoimmune diseases, cancers etc.).

U.S. Patent Application Nos. 20020028203, 20050169922 and 20080102071 disclose compositions which bind T cell inhibitory receptor molecules and modulate (i.e. enhance or suppress) T cell activity (e.g. cytotoxicity and proliferation), such as biliary glycoprotein binding agents, and methods of using such compositions such as for treatment of diseases (e.g. an autoimmune disease, immunodeficiency, cancer etc.).

WO 2010/125571 to the present inventor discloses a murine monoclonal antibody produced by a specific hybridoma cell. The mAb is highly selective to CEACAM1 and does not cross-react with other members of the CEACAM family.

None of the known antibodies which recognize CEACAM1 have the spectrum of binding specificity of the monoclonal antibodies of the present invention. Thus, there is an unmet need to provide antibodies recognizing specific subsets of CEACAM proteins which can be used diagnostically and therapeutically in diseases involving CEACAM expression or activation.

SUMMARY OF THE INVENTION

The present invention discloses monoclonal antibodies which recognize a specific set of CEACAM subtypes. Advantageously, the antibodies of the invention show binding to CEACAM1 and at least one additional subtype selected from CEACAM5 and CEACAM3. The antibodies of the invention are characterized by having unique CDR sequence and framework combinations and by binding to newly identified epitopes within the CEACAM1 molecule. The unique specificity of the monoclonal antibodies of the present invention, broaden their therapeutic utility for treatment and diagnosis of additional types of malignancies and viral infections. The present invention also provides methods for identifying and isolating such antibodies, methods for their production, and therapeutic and diagnostic uses thereof.

The monoclonal antibodies according to the present invention have specific combinations of CDRs and possess unique properties and improved specificity and potency over known anti CEACAM1 antibodies.

According to one aspect, the present invention provides a monoclonal antibody which recognizes CEACAM1, or an antibody fragment thereof comprising at least an antigen-binding portion thereof, having heavy-chain CDRs comprising sequences set forth in SEQ ID

NOs: 1, 2 and 3, and light-chain CDRs comprising sequences set forth in SEQ ID NOs: 4, 5 and 6, and analogs and derivatives thereof.

5 According to some embodiments a monoclonal antibody or antibody fragment which recognizes CEACAM1 is provided having a heavy-chain CDR1 comprising a sequence set forth in SEQ ID NO: 1, a heavy-chain CDR2 comprising a sequence set forth in SEQ ID NO: 2 a heavy-chain CDR3 comprising a sequence set forth in SEQ ID NO: 3, a light-chain CDR1 comprising a sequence set forth in SEQ ID NO: 4, a light-chain CDR2 comprising a sequence set forth in SEQ ID NO: 5 and a light-chain CDR3 comprising a sequence set forth in SEQ ID NO:6, and analogs and derivatives thereof.

10 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, comprising heavy chain CDRs having the sequences set forth in SEQ ID NOs: 7, 8 and 9.

15 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, comprising heavy chain CDRs having the sequences set forth in SEQ ID NOs: 13, 14 and 15.

According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, comprising light chain CDRs having the sequences set forth in SEQ ID NOs: 10, 11 and 12.

20 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, wherein the light chain CDRs having the sequences set forth in SEQ ID NOs: 16, 17 and 18.

According to other embodiments a monoclonal antibody is provided having CDR sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17, and 18.

25 According to yet other embodiments, a monoclonal antibody is provided having CDR sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12.

Analogous and derivatives of the monoclonal antibody or fragment thereof, having at least 90% sequence identity with the antigen-binding portion of the reference sequence are also within the scope of the present invention.

30 According to some embodiments, analogous and derivatives of the monoclonal antibody or fragment thereof having at least 95% sequence identity with the antigen-binding portion of the reference sequence are provided. According to a specific embodiment the antibody comprises a heavy chain variable domain sequence having a sequence set forth in SEQ ID NO: 26:

QVQLQQSGAELVRPGTSVKVSKASGYAFTNNLIEWVKQRPGQGLEWIGVINPGSG
DTNYNEKFKGKATLTADKSSNTAYMQLSSLTSDDSA VYFCARGDY YGGFAVDYW
GQGTSVTVSS, or an analog or derivative thereof having at least 97% sequence identity
with the heavy chain sequence.

5 According to yet another embodiment the antibody comprises a light chain variable
domain sequence having a sequence set forth in SEQ ID NO: 28:

DIQMTQTTSSLSASLGDRVTISCRTSQDIGNYLNWYQQKPDGTVKLLIYYTSRLHSG
VPSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGKSLPRTEFGGGTKLEIK, or an analog
or derivative thereof having at least 97% sequence identity with the light chain sequence.

10 According to a specific embodiment the antibody or fragment thereof comprises a
heavy chain variable domain having a sequence set forth in SEQ ID NO: 26 and a light chain
variable domain having a sequence set forth in SEQ ID NO: 28, or an analog or derivative
thereof having at least 97% sequence identity with the antibody or fragment sequence.

The present invention encompasses monoclonal antibodies isolated from hybridoma
15 cells or other biological systems, as well as monoclonal antibodies produced recombinantly
or synthetically. A monoclonal antibody according to the present invention may contain a
constant region from any mammalian species, including but not limited to mouse, rat and
human. A monoclonal antibody according to the present invention includes a chimeric
antibody, a humanized antibody, a fully human antibody, a xenogeneic antibody, and an
20 antibody fragment comprising at least the antigen-binding portion of an antibody. According
to a specific embodiment the antibody fragment is selected from the group consisting of: Fab,
Fab', F(ab')₂, Fd, Fd', Fv, dAb, isolated CDR region, single chain antibody, "diabodies", and
"linear antibodies".

25 According to some particular embodiments the present invention provides a
monoclonal antibody, or an antibody fragment comprising:

- i. a framework sequence selected from the group consisting of: mouse IgG2a, mouse
IgG2b, mouse IgG3, human IgG1, human IgG2, human IgG3; and
- ii. a set of six CDRs having sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17,
and 18; or a set of six CDRs having sequences set forth in SEQ ID NOs: 7, 8, 9, 10,
30 11, and 12; and analogs and derivatives thereof having at least 97% sequence identity
with said CDR sequences, wherein the monoclonal antibody or fragment binds with
an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.

According to some embodiments, the monoclonal antibody or fragment thereof binds
with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.

According to other embodiments, the monoclonal antibody or fragment thereof binds with an affinity of at least about 10^{-8} M to CEACAM1.

According to some specific embodiments, the monoclonal antibody is a chimeric monoclonal antibody.

5 According to some embodiments, the chimeric antibody comprises human-derived constant regions.

According to some embodiments the human constant regions of the chimeric antibody are selected from the group consisting of: human IgG1, human IgG2, and human IgG3

10 According to a particular embodiment, a chimeric or humanized monoclonal antibody which recognizes CEACAM1 is provided comprising the six CDRs having sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17, and 18; or the six CDRs having sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12; and analogs and derivatives thereof having at least 95% sequence identity with said CDR sequences, and a constant region subclass selected from human IgG1, human IgG2 and human IgG3, wherein the monoclonal antibody binds
15 with an affinity of at least about 5×10^{-7} M to at least two CEACAM subtypes.

According to a specific embodiment the chimeric or humanized monoclonal antibody or fragment thereof, comprises a constant region subclass of human IgG1 subtype.

20 According to another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided comprising a heavy chain sequence set forth in SEQ ID NO: 30.

According to yet another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided comprising a light chain sequence set forth in SEQ ID NO: 31.

25 According to yet another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided having a heavy chain sequence set forth in SEQ ID NO: 30, and light chain sequence set forth in SEQ ID NO: 31.

30 According to a particular embodiment, a monoclonal antibody which recognizes CEACAM1 is provided produced from DNA sequences of the heavy and light chains contained in a plasmid deposited on September 28, 2011 under ATCC Accession Number PTA-12130.

Monoclonal antibodies of the present invention exhibit according to some embodiments specific binding to more than one CEACAM subtype. According to some embodiments, the monoclonal antibody binds at least two different CEACAM subtypes.

According to some specific embodiments the monoclonal antibody binds to CEACAM1 and at least one of CEACAM3 and CEACAM5. According to a particular embodiment the monoclonal antibody binds to CEACAM1 and CEACAM5. According to another particular embodiment the monoclonal antibody binds to CEACAM1 and CEACAM3. According to yet other embodiments, a monoclonal antibody according to the present invention binds to CEACAM subtypes 1, 3 and 5.

According to particular embodiments, a monoclonal antibody according to the present invention does not bind to CEACAM4 and CEACAM6.

According to yet another aspect the present invention provides a monoclonal antibody which recognizes CEACAM1, or a fragment thereof comprising at least the antigen-binding portion, which is capable of binding the same epitope on the CEACAM1 molecule to which a monoclonal antibody to CEACAM1 having a heavy chain sequence set forth in SEQ ID NO: 26 or SEQ ID NO: 30 and a light chain sequence set forth as SEQ ID NO: 28 or SEQ ID NO: 31, binds.

According to some embodiments, the monoclonal antibody is reactive with an epitope within residues 17-29 and 68-79 of human CEACAM1 having the sequences VLLLVHNLPPQLF (SEQ ID NO:32) and YPNASLLIQNVT (SEQ ID NO:33) respectively.

According to some embodiments, the epitope on the CEACAM1 molecule to which the monoclonal antibody binds is an epitope comprising amino acid residues within the sequences VLLLVHNLPPQLF (SEQ ID NO: 32) and YPNASLLIQNVT (SEQ ID NO: 33).

According to other embodiments, the monoclonal antibody according to the invention binds an epitope comprising at least four amino acids of the sequence VLLLVHNLPPQLF (SEQ ID NO: 32).

According to yet other embodiments, the monoclonal antibody according to the invention binds an epitope within sequences VLLLVHNLPPQLF (SEQ ID NO: 32) and PNASLLI (SEQ ID NO: 34).

According to some embodiments, the monoclonal antibody or fragment thereof binds to the same epitope of which an antibody having the six CDR sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11 and 12 binds.

According to yet other embodiments, the monoclonal antibody or fragment thereof binds to the same epitope of which an antibody having the CDR sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17 and 18 binds.

According to a particular embodiment, the monoclonal antibody or fragment thereof binds to the same epitope which is bound by an antibody produced from DNA sequences deposited on September 28, 2011 under ATCC Accession Number PTA-12130.

5 The present invention provides, according to yet another aspect, an isolated peptide sequence of 6-20 amino acids comprising at least three amino acids from the sequence VLLLVHNLPPQLF (SEQ ID NO: 32) and at least three amino acids from the sequence YPNASLLIQNVT (SEQ ID NO: 33). Analogs and derivatives of said peptide, having at least 85%, 90%, 95% or 98% homology with the parent sequence are also within the scope of the present invention.

10 According to some embodiments, the isolated peptide comprises at least six amino acids from the sequence VLLLVHNLPPQLF (SEQ ID NO: 32).

According to yet other embodiments, the isolated peptide comprises the amino acids of the sequences VLLLVHNLPPQLF (SEQ ID NO: 32) and PNASLLI (SEQ ID NO: 34).

15 Use of the isolated peptides for production of monoclonal or polyclonal antibodies is also within the scope of the present invention as well as their use in diagnosis or treatment.

Within the scope of the present invention are also nucleic acid molecules encoding an antibody or antibody fragment according to the invention, having affinity and specificity for CEACAM1.

20 According to this aspect, an isolated polynucleotide sequence encoding an antibody which recognizes CEACAM1 or an antibody fragment thereof is disclosed.

25 According to some embodiments, the isolated polynucleotide sequence comprises a DNA sequence set forth in SEQ ID NO: 25 or analog thereof having at least 90% sequence identity with said DNA sequence. According to other embodiments, the isolated polynucleotide sequence comprises a DNA sequence set forth in SEQ ID NO: 27 or analog thereof having at least 90% sequence identity with said DNA sequence.

Plasmids comprising at least one polynucleotide sequence encoding a monoclonal antibody or fragment thereof according to the invention are also disclosed, as well as host cells comprising these plasmids.

30 According to a particular embodiment, a plasmid comprising polynucleotide sequences set forth in SEQ ID NOs: 25 and 27, deposited on September 28, 2011 under ATCC Accession Number PTA-12130, is disclosed.

In another aspect the present invention is related to a pharmaceutical composition useful for preventing, attenuating or treating a disease or disorder associated with CEACAM1, CEACAM3 or CEACAM5 expression, activation or function. A pharmaceutical

composition according to the invention comprises a therapeutically effective amount of a monoclonal antibody which recognizes CEACAM1, CEACAM3 or CEACAM5 or an antibody fragment thereof comprising at least an antigen-binding portion; and a pharmaceutically acceptable carrier.

5 According to some embodiments, the pharmaceutical composition comprises a monoclonal antibody capable of binding to CEACAM1 with a binding affinity of at least 10^{-8} kD.

10 According to additional embodiments, the pharmaceutical composition comprises a monoclonal antibody capable of binding with an affinity of at least about 10^{-8} kD to CEACAM1 and with affinity of at least about 5×10^{-7} M to at least one of CEACAM3 and CEACAM5.

According to a particular embodiment, the pharmaceutical composition comprises a monoclonal antibody capable of binding with an affinity of at least about 5×10^{-7} kD to CEACAM1, CEACAM3 and CEACAM5.

15 According to certain embodiments the disease or disorder associated with CEACAM1, CEACAM3 and/or CEACAM5 expression, activation or function is a cell proliferative disease or disorder. According to some embodiments the cell proliferative disease or disorder is cancer.

20 According to some embodiments, the cancer associated with over-expression of CEACAM5 is selected from the group consisting of: gastrointestinal, colorectal (CRC) pancreatic non-small cell lung cancer (NSCL), breast, thyroid, stomach, ovarian and uterine.

According to a specific embodiment the cancers associated with over expression of CEACAM1 are melanoma, pancreatic cancer, all types of lung cancers and myeloma.

25 The pharmaceutical composition according to the present invention may be administered as a stand alone treatment or in addition to a treatment with any other therapeutic agent. According to a specific embodiment, antibodies according to the present invention are administered to a subject in need thereof as part of a treatment regimen in conjunction with at least one anti-cancer agent. The pharmaceutical composition according to the present invention may be administered together with the other agent or separately.

30 In another aspect the present invention provides diagnostic compositions useful for detecting at least one CEACAM subtype selected from the group consisting if: CEACAM1, CEACAM3 and CEACAM5, in a subject. A diagnostic composition according to the invention comprises a therapeutically effective amount of a monoclonal antibody having affinity of at least about 5×10^{-7} M to CEACAM1, CEACAM3 or CEACAM5 or an antibody

fragment thereof comprising at least an antigen-binding portion; and an optional carrier or excipient.

In yet another aspect the present invention is related to a method of preventing, attenuating or treating a disease or disorder associated with expression, activation or function of CEACAM, comprising administering to a subject in need thereof a pharmaceutical composition comprising a therapeutically effective amount of an antibody to CEACAM; and a pharmaceutically acceptable carrier.

According to some embodiments the disease or disorder is a cell proliferative disease or disorder. According to certain embodiments the cell proliferative disease or disorder is cancer. According to a specific embodiment the monoclonal antibody, or fragment thereof, has an affinity of at least about 10^{-8} M to CEACAM1 and the cancer is melanoma.

According to other embodiments, the monoclonal antibody, or fragment thereof, has an affinity of at least about 5×10^{-7} M to CEACAM5 and the cancer is selected from the group consisting of: gastrointestinal, colorectal (CRC) pancreatic, non-small cell lung cancer (NSCL), breast, thyroid, stomach, ovarian, myeloma and uterine.

According to an additional embodiment, the disease or disorder associated with over expression of CEACAM1 is a viral infection.

According to some embodiments, the viral infection is caused by a virus selected from the group consisting of: DNA viruses, such as but not limited to cytomegalovirus (CMV), adenovirus, hepatitis virus and human papillomavirus (HPV); and RNA viruses such as but not limited to influenza virus and human immuno-deficiency virus (HIV).

According to an aspect of the present invention there is provided a method of immunomodulation, the method comprising contacting a CEACAM-expressing lymphocyte with the antibody or antibody fragment.

According to an aspect of the present invention there is provided a method of inhibiting migration of a CEACAM expressing tumor cell, the method comprising contacting the CEACAM expressing tumor cell with the antibody or antibody fragment, thereby inhibiting migration of a CEACAM expressing tumor cell.

According to some embodiments, the tumor cell comprises a melanoma tumor cell.

According to an aspect of the present invention there is provided a method of treating cancer, the method comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antibody fragment, thereby treating the cancer in the subject.

According to an aspect of the present invention there is provided a method of inhibiting CEACAM homotypic or heterotypic protein-protein interaction, the method comprising contacting a CEACAM1-expressing lymphocyte with the antibody or antibody fragment, thereby inhibiting CEACAM1 homotypic or heterotypic protein-protein interaction.

According to some embodiments, the isolated antibody or antibody fragment is attached to a cytotoxic moiety.

According to some embodiments, the cytotoxic moiety comprises a cytotoxin, a chemokine, a chemotherapeutic composition, a pro-apoptotic, an interferon, a radioactive moiety, or combinations thereof.

According to some embodiments, the antibody or antibody fragment is attached to an identifiable moiety.

According to some embodiments, cells of the cancer are characterized by over expression of CEACAM1 as compared to unaffected cells.

According to some embodiments, the method of treating cancer further comprises administering to the subject lymphocytes.

According to some embodiments, the lymphocytes comprise T cells or NK cells. According to some embodiments, the lymphocytes express CEACAM1. According to other embodiments, the CEACAM1-expressing lymphocyte is a Tumor Infiltrating Lymphocyte (TIL). According to other embodiments, the CEACAM1-expressing lymphocyte is a cytotoxic T cell.

The antibody of the present invention can be used to block CEACAM on either or both immune effector cells (CEACAM expressing lymphocytes e.g., tumor infiltrating cells, T cells or NK cells) and target cells (e.g., CEACAM expressing pathological cells such as cancer cells). Examples of cancer cells which are candidates for this therapy include, but are not limited to, melanoma, lung, thyroid, breast, colon, prostate, hepatic, bladder, renal, cervical, pancreatic, leukemia, lymphoma, myeloid, ovarian, uterus, sarcoma, biliary, or endometrial cells.

According to a further aspect of the invention there is provided a method of rendering a CEACAM expressing tumor cell susceptible to immunomodulation. The method comprising contacting the CEACAM expressing tumor cell (e.g., melanoma, lung, thyroid, breast, colon, prostate, hepatic, bladder, renal, cervical, pancreatic, leukemia, lymphoma, myeloid, ovarian, uterus, sarcoma, biliary or endometrial cell) with the antibody or antibody

fragment described above, thereby rendering the CEACAM expressing tumor cell susceptible to immunomodulation.

5 Additionally or alternatively, the present invention also envisages a method of immunomodulation (e.g., inhibiting CEACAM1 homotypic or heterotypic protein-protein interaction), by contacting a CEACAM1-expressing lymphocyte with the antibody or antibody fragment described herein.

The therapeutic or preventive methods of the present teachings can be effected ex-vivo (e.g., used in T cell based adoptive immunotherapy) or in-vivo.

10 Antibodies of some embodiments of the invention can have anti cancer activity which is independent from its immunomodulatory activity described above.

In another aspect, the present invention provides a method for increasing the duration or progression of response or survival of a subject having cancer, comprising administering to the subject effective amounts of a composition comprising an antibody which recognizes CEACAM and an anti-neoplastic composition, wherein said anti-neoplastic composition
15 comprises at least one chemotherapeutic agent, whereby the co-administration of the antibody and the anti-neoplastic composition effectively increases the duration or progression of response or survival.

Furthermore, the present invention provides a method for treating a subject having cancer, comprising administering to the subject effective amounts of a composition
20 comprising an antibody to CEACAM and an anti-neoplastic composition whereby co-administration of the antibody to CEACAM and the anti-neoplastic composition effectively increases the response incidence in the group of subjects.

Aside from therapeutic applications, antibodies of the present invention can also be used in diagnostic applications.

25 Thus, according to a further aspect there is provided a method for diagnosing a cancer in a subject in need thereof, the method comprising contacting a biological sample derived from the subject (in-vivo, in vitro or ex-vivo) with the antibody or antibody fragment described herein, wherein a complex formation beyond a predetermined threshold is indicative of the cancer in the subject. According to some embodiments, cells of the cancer
30 are characterized by over expression of CEACAM as compared to unaffected cells.

According to a particular embodiment the diagnosed cancer is selected from the group consisting of: melanoma, pancreatic cancer, lung cancer and myeloma.

According to another particular embodiment the measured protein is CEACAM5 and the diagnosed cancer is selected from the group consisting of: gastrointestinal, colorectal

(CRC) pancreatic non-small cell lung cancer (NSCL), breast, thyroid, stomach, ovarian and uterine.

As mentioned, the method of the invention is affected under conditions sufficient to form an immunocomplex; such conditions (e.g., appropriate concentrations, buffers, temperatures, reaction times) as well as methods to optimize such conditions are known to those skilled in the art, and examples are disclosed herein. As used herein the phrase “immunocomplex” refers to a complex which comprises the antibody of the invention and the CEACAM. Determining a presence or level of the immunocomplex of the invention may be direct or by detecting an identifiable (detectable) moiety which may be attached to the antibody.

The level of the immunocomplex in the tested cell (e.g., a cell of a subject in need thereof) is compared to a predetermined threshold. It will be appreciated that the antibody of the present invention can also be used to measure the amount of serum soluble CEACAM. Regardless, the threshold may be determined based on a known reference level and/or a level in a control cell or serum. The control cell can be obtained from a control, healthy subject (e.g., a subject not suffering from the cancer) or from the same subject prior to disease initiation or following treatment. According to some embodiments of the invention, the control subject is of the same species e.g. human, preferably matched with the same age, weight, sex etc. as the subject in need thereof.

To facilitate diagnosis, the above teachings can be combined with other methods of diagnosing cancer which are well known in the art include but are not limited to imaging, molecular tests and surgical biopsies.

According to another aspect of present invention a method for detecting or quantifying the presence of CEACAM in is provided. Thus, the present invention also provides methods for diagnosing conditions associated with CEACAM expression using antibodies which recognizes CEACAM. Diagnostic methods according to the invention may be performed according to specific embodiments, *in-vitro* or *ex-vivo*. The antibodies according to the present invention may be also used to configure screening methods. For example, an ELISA assay can be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art.

According to one embodiment a method is provided for detecting or quantifying the presence of CEACAM, comprising the steps of:

- i. incubating a biological sample with an antibody to CEACAM or an antibody fragment thereof comprising at least an antigen-binding portion;
- ii. detecting the bound CEACAM using a detectable probe;
- iii. comparing the amount of (ii) to a standard curve obtained from reference samples containing known amounts of CEACAM; and
- iv. calculating the amount of the CEACAM in the sample from the standard curve.

According to another embodiment a method for diagnosing a disease or disorder associated with CEACAM expression is provided comprising the steps of:

- i. incubating a biological sample with an antibody to CEACAM or an antibody fragment thereof comprising at least an antigen-binding portion;
- ii. detecting the bound CEACAM using a detectable probe;
- iii. comparing the amount of (ii) to a standard curve obtained from reference samples containing known amounts of CEACAM;
- iv. calculating the amount of the CEACAM in the biological sample from the standard curve; and
- v. comparing the amount of (iv) to a normal CEACAM amount.

According to some embodiments the biological sample is a body fluid of a mammalian subject. According to particular embodiments, the mammalian subject is human.

The antibodies of the present invention may be also used in screening assays for assessing the CEACAM levels in patients and for prediction of the effectiveness of treatment. The screening assays with the antibodies of the present invention may allow determination of the levels of CEACAM and therefore prediction of treatment outcome and planning of an appropriate treatment regimen.

According to other embodiments, the level of at least one of CEACAM1, CEACAM3 and CEACAM5 is assessed. According to a particular embodiment the level of CEACAM1 is assessed.

According to some embodiments of the invention, the antibody or antibody fragment is attached to an identifiable moiety.

It will be appreciated that such attachment of antibodies, or fragments thereof, and identifiable moiety can be effected using chemical conjugation or by recombinant DNA technology according to methods well known in the art.

The identifiable moiety can be a member of a binding pair, which is identifiable via its interaction with an additional member of the binding pair and a label which is directly

visualized. In one example, the member of the binding pair is an antigen which is identified by a corresponding labeled antibody. In one example, the label is a fluorescent protein or an enzyme producing a colorimetric reaction.

Another aspect of the present invention relates to the use of an antibody to CEACAM or an antibody fragment thereof, for diagnosis or treatment of a cell proliferative or angiogenesis-related disease or disorder or a viral infection.

According one embodiment the cell proliferative disease is melanoma.

According to other embodiments the cell proliferative disease or disorder is a cancer selected from the group consisting of: gastrointestinal, colorectal (CRC) pancreatic non-small cell lung cancer (NSCL), breast, thyroid, stomach, ovarian, uterine, and myeloma.

According to one embodiment, the present invention provides use of an antibody to CEACAM or an antibody fragment thereof comprising at least an antigen-binding portion, for preparation of a medicament for treatment of a disorder or disease associated with expression or activation of, including but not limited to cancer and viral infection.

The invention also relates to use of an antibody to CEACAM or an antibody fragment thereof, for the manufacture of a diagnostic composition for the diagnosis of a cell proliferative or angiogenesis-related disease or disorder or a viral infection.

Essentially all of the uses known or envisioned in the prior art for CEACAM1, CEACAM3 and CEACAM5 antibodies can be accomplished with the antibodies of the present invention which are shown to possess improved affinity toward these proteins and superior inhibitory and in indirect immunomodulatory effects on CEACAM1 bearing cells. These uses include diagnostic, prophylactic and therapeutic techniques.

Further embodiments and the full scope of applicability of the present invention will become apparent from the detailed description given hereinafter.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is an SDS-PAGE image showing light and heavy chains of the chimeric antibody CM10.

Figure 2 shows specific binding curve of CM10 to purified hCEACAM1.

Figure 3 demonstrates specific binding of CM10 to CEACAM1 as detected by Flow Cytometry analysis.

Figure 4 confirms that CM10 blocks CEACAM1-CEACAM1 interaction between cells. Mouse IL-2 secretion of effectors cells (BW/221 cells expressing CEACAM1) incubated in the presence of various CM10 concentrations, was measured by ELISA.

Figure 5 shows CM10 enhancement of the specific killing activity of CEACAM1-positive melanoma cells.

Figure 6 demonstrates that CM10 stimulates the killing activity of tumor-infiltrating lymphocyte (TILs).

5 **Figure 7** demonstrates that CM10 enhances the killing activity of NK cells on CEACAM1 positive melanoma cell lines.

Figure 8 CM10 immunomodulatory effect inhibits tumor growth in-vivo. Arrows indicate time of administration (CM10 cycles, TIL triangles, CM10 and TIL open squares).

Figure 9 is a schematic presentation of CM10 immunomodulatory mode of action.

10 **Figure 10** represents CEACAM1 binding intensity level in tumors as determined by anti CEACAM1 antibody.

Figure 11 shows quantification of CM10 molecules bound per cell.

Figure 12 confirms that CM10 has no effect on PBMC Proliferation. Results represent average proliferation rates from three donors for each treatment.

15 **Figure 13** presents FACS analysis of binding between CM10 to CEACAM family proteins. CEACAM1, 5, 6 and 8 were expressed by 721.221 cells, and CEACAM3 and 4 by HEK293T cells.

Figure 14 represents results of complement-dependent cytotoxicity (CDC) assay in melanoma cell lines.

20 **Fig 15** demonstrates that CM10 enhances granzyme B secretion of TIL in the presence of CEACAM1 and HLA-A2 positive melanoma cells.

Fig 16 shows that CM10 blocks CEACAM1-CEACAM5 interactions.

Fig 17 presents CM10 enhancement of HLA restricted T cell killing.

Fig 18 demonstrates the immunomodulatory activity of CM10 inhibits tumor growth in-vivo.

25 **Fig 19:** indicates that CM10 enhances the killing activity of NK cells on CEACAM1-positive pancreatic cancer cell lines COLO-357 and BXPC3.

Fig 20: demonstrates that CM10 enhances granzyme B secretion of NK cells in the presence of CEACAM1-positive pancreatic cancer cell lines.

30 **DETAILED DESCRIPTION OF THE INVENTION**

The present invention provides antibodies which recognize CEACAM1 comprising specific sets of CDR sequences which possess improved and unique specificity, selectivity, affinity and/or activity.

Antibodies according to the present invention bind CEACAM1 with higher affinity than other anti-CEACAM1 antibodies, they blocks the function of CEACAM1, while not all anti CEACAM antibodies do, and more efficiently than polyclonal anti CEACAM antibodies. Furthermore, antibodies according to the present invention are effective against cancer cells, in particular melanoma cells: the antibodies render melanoma cells more susceptible to lymphocytes, inhibit melanoma growth rate in vivo, an effect which is enhanced when the antibody is combined with adoptive T cell transfer in vivo.

It is shown here for the first time that the in vivo anti-melanoma effect of anti-CEACAM1 antibodies according to the invention is a combined direct anti-tumor effect as well as immunomodulatory effect rendering the cells more susceptible to reactive lymphocytes.

An antibody according to the present invention, fragments and derivatives can be used as an effective tool for diagnosis, immunomodulation and cancer treatment.

The antibody inhibits CEACAM1 homophilic interactions, as determined by co-incubation of immune effector cells and target cells expressing CEACAM1 and assaying IL-2 secretion and by the in vitro killing assays.

The antibody of the present invention is shown to enhances CM10 enhances HLA restricted T cell killing and to enhance granzyme B (a serine protease involves in mediation apoptosis of the target cells) secretion from effector NK and T cells in the presence of specific target cells, thus enabling the observed enhanced killing of the target cells by the antibody. It is also inhibits the binding between CEACAM1 to CEACAM5 in a dose-dependent manner therefore can be used to treat malignancies that express high level of CEACAM5 and exploit the CEACAM1-CEACAM5 axis in order to suppress the immune cells.

In addition it is herein shown that, an antibody according to the invention is effective in inhibiting melanoma cells invasion. Furthermore, *in vivo* administration of an antibody according to the invention, either alone or in combination with reactive lymphocytes was shown effective in inhibiting growth of melanoma tumors. The combination of adoptive human T cell transfer with monoclonal antibody injections exhibited significant synergism and strongly inhibited xenograft growth compared to the isotype control group.

According to a further aspect of the invention there is provided an isolated antibody or antibody fragment having the same binding specificity and selectivity to an antibody defined herein comprising an antigen recognition domain having specific CDR segments described above. According to this aspect, isolated antibody or antibody fragment is capable

of binding the same epitope determinant of the CEACAM1 protein as does the antibody described above by its specific CDR segments sequences.

The proposed sequence of an epitope to which a monoclonal antibody according to the invention binds is disclosed herein for the same time, together with proposed isolated peptides derived from this epitope which can be used for raising additional monoclonal antibodies.

Monoclonal antibodies (mAbs) can be designed to selectively target tumor cells and elicit a variety of responses once bound. These agents can destruct tumor cells in different ways such as blocking tumor cell proliferation or activating the immune system. Chimeric monoclonal antibodies according to the present invention were designed to specifically bind and neutralize various functions of the CEACAM1 protein and other CEACAM subtype proteins, and to induce the specific death of tumor cells. Without wishing to be bound to any theory, it is suggested that monoclonal antibodies according to the present invention act also via activation of the immune system against cancerous cells.

Both the clinical and biological evidence highlight CEACAM1 as a promising target for the development of targeted-immunotherapy. CEACAM1 is not found on normal melanocytes, but undergoes neo-expression and is widely expressed on the vast majority of metastatic melanoma specimens. It has been previously demonstrated mechanistically that CEACAM1 protects melanoma cells by inhibiting effector functions of NK cells and T cells.

It is herein demonstrated for the first time that CM10 is a chimeric monoclonal antibody which binds with high affinity to human CEACAM1. *In-vitro*, CM10 efficiently blocked CEACAM1-homophilic interactions in a dose dependent manner and improves CEACAM1 positive melanoma cells killing by T cells and NK cells. Moreover, CM10 significantly inhibited the *in- vivo* growth of melanoma xenografts when administered systemically along with melanoma-reactive human T lymphocytes (tumor-infiltrating lymphocytes, TILs). Without wishing to be bound to any theory, this is in line with the suggested mechanism of action; abrogation of immune-protective interactions of the tumor cells with the activated lymphocytes.

Several evidences reported that CEACAM1 is expressed by a wide variety of epithelial cells, including colon, prostate, breast, kidney etc. Extensive examination of CEACAM1 expression profile on normal and malignant tissues by IHC have been performed. The expression analysis showed a strong staining of melanoma cells, as compared to no staining of the vast majority of the tissues tested in a normal human tissue. Nevertheless, some selective staining was observed in restricted sites of several organs. When more

quantitative method was used to quantify the number of CM10 mAb molecules bound to malignant and normal primary cells, very low CM10 molecules could be detected in normal cells, which may indicate that CM10 binds mostly to patient's tumor cells. Furthermore it is shown that CM10 has no effect on primary cells proliferation and is completely or almost completely, unable to induce CDC or ADCC indicating the potential safety of the monoclonal antibody in human subjects.

Since CM10 has an immunomodulation activity, possible immune-related side effects, are evaluated. Following PBMC activation, CEACAM1 is upregulated on the activated lymphocytes (Gray-Owen and Blumberg 2006, Nat Rev Immunol 6, 433-46). *Ex-vivo* human PBMC proliferation assay revealed that CM10 has no effect on naïve and activated PBMC proliferative response.

The main advantage of CEACAM1 blockade over abrogation of generalized inhibitory mechanisms is the expected selectivity to the vicinity of the tumor and therefore fewer adverse events compare to other general immune toxicity agents.

As demonstrated in the present invention, CM10 shows encouraging activity and safety profile and is a promising candidate for cancer immunotherapy and can be used as a strategy to selectively enhance the anti-tumor properties of the endogenous immune response in several malignancies, such as melanoma and non-small cell lung cancer.

Binding to additional CEACAM subtypes increases the therapeutic profile of the antibody, thus it can be used for diagnosis and treatment of other types of malignancies which do not extensively express CEACAM1 but express CEACAM5, for example.

CEACAM5 has been found to be over-expressed in a high percentage of many human tumors, including 90% of gastrointestinal, colorectal (CRC) and pancreatic cancers, 70% of non-small cell lung cancer cells and 50% of breast cancers. It is also over-expressed in thyroid, stomach, ovarian and uterine cancers (Thompson, Grunert et al. 1991, J Clin Lab Anal 5, 344-66). CEACAM5 even serves as a clinical marker for liver metastasis in CRC and post-surgical surveillance of colon cancer (Duffy 2001, Clin Chem 47, 624-30). The evidence that CM10 is capable to bind CEACA5 is very important and can expand the possible indications that can be treated by CM10 from 4-5 types of malignancies to above 10. The anti-CEACAM5 agents that have entered clinical trials include anti-CEACAM5 antibodies conjugated to toxic substances such as radioactive substances for both diagnostic purposes and for the treatment of various malignancies. It seems that even these toxic conjugated forms don't show safety problems, which can indicate that CEACAM5 is a safe target.

The human counterparts of murine IgG subclasses are based on similarities in biological and functional activities. Murine IgG2a and IgG2b and human IgG1 and IgG3 share the ability to fix complement and bind to protein antigens (Hussain et al., 1995, Clinical and Diagnostic Laboratory Immunology 726-732). Murine IgG1 and human IgG4 are considered to be similar because of their property of binding to mast cells. Human IgG4 is the only human IgG subclass which does not activate complement and the subclasses IgG1 and 3 are the most effective in activating complements. For mouse it is the subclasses IgG2a and IgG2b which are active with IgG1 and possibly IgG3 being inactive (Clark MR., Chem Immunol. 1997;65:88-110).

Several known monoclonal antibodies which recognize CEACAM1 are of subtype mouse IgG1. As the human equivalent of mouse IgG1 is IgG4 it would be expected to create a chimeric antibody comprising the human IgG4 constant framework. Unexpectedly, according to some embodiments of the present invention chimeric monoclonal antibodies comprise a human IgG1 constant framework.

According to one aspect, the present invention provides a monoclonal antibody which recognizes CEACAM1, or an antibody fragment comprising at least an antigen-binding portion thereof, comprising at least one heavy-chain CDR comprising a sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and at least one light-chain CDR comprising a sequence selected from the group consisting of: SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6, and analogs and derivatives thereof.

According to some embodiments, analogs and derivatives of the monoclonal antibody or fragment thereof, having at least 90% sequence identity with the sequence of the reference sequence are disclosed.

According to other embodiments analogs and derivatives of the monoclonal antibody or fragment thereof having at least 95% sequence identity with the reference sequence are disclosed.

According to yet other embodiments, analogs and derivatives of the monoclonal antibody or fragment thereof having at least 98% sequence identity with the CDR sequence of the reference antibody are disclosed.

According to one embodiment the antibody or antibody fragment comprises at least two heavy-chain CDRs comprising a sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and at least one light-chain CDRs comprising a sequence selected from the group consisting of: SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID

NO: 6, and analogs and derivatives thereof having at least 97% sequence identity with the sequence of the monoclonal antibody or fragment thereof.

According to other embodiments the antibody or antibody fragment comprises at least one heavy-chain CDR comprising a sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and at least two light-chain CDRs comprising a sequence selected from the group consisting of: SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6, and analogs and derivatives thereof having at least 97% sequence identity with the sequence of the monoclonal antibody or fragment thereof.

According to yet other embodiments the antibody or antibody fragment comprises at least two heavy-chain CDRs comprising a sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and at least two light-chain CDRs comprising a sequence selected from the group consisting of: SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6, and analogs and derivatives thereof having at least 97% sequence identity with the sequence of the monoclonal antibody or fragment thereof.

According to some embodiments the antibody or antibody fragment comprises at least one heavy-chain CDR sequence of at least five amino acids derived from a sequence selected from the group consisting of: SEQ ID NO: 19, SEQ ID NO: 20 and SEQ ID NO: 21, and at least one light-chain CDR sequence of at least five amino acids derived from a sequence selected from the group consisting of: SEQ ID NO: 22, SEQ ID NO: 23 and SEQ ID NO: 24, and analogs and derivatives thereof having at least 97% sequence identity with the sequence of the monoclonal antibody or fragment thereof.

According to other embodiments, the antibody binding site of the antibody or fragment thereof consists of three heavy chain CDRs selected from the group consisting of SEQ ID NOs: 7, 8, 9, 13, 14 and 15 and three light chain CDRs selected from the group consisting of SEQ ID NOs: 10, 11, 12, 16, 17, 18, and analogs and derivatives thereof having at least 97% sequence identity with the antibody binding site.

According to yet other embodiments, the antibody binding site consists of the six CDRs of SEQ ID NOs: 13, 14, 15, 16, 17, and 18.

According to other embodiments, the antibody binding site consists of the six CDRs of SEQ ID NOs: 7, 8, 9, 10, 11, and 12.

The CDR sequences according to the invention were identified using two different algorithm methods: IMGT algorithm (Lefranc et al., 1999, Nucleic Acids Research, 27, 209-212); and KABAT algorithm (Wu TT and Kabat E.A., 1970, J. Exp. Med. 132, 211-250). The sequences revealed by both methods are disclosed.

According to some embodiments, the heavy chain CDR1 of the antibody according to the invention or a fragment thereof is selected from>NNLIE (SEQ ID NO: 7) and GYAFTNNL (SEQ ID NO: 13).

5 According to some embodiments, the heavy chain CDR2 of the antibody according to the invention or a fragment thereof is selected from>VINPGSGDTNYNEKFKG (SEQ ID NO: 8) and INPGSGDT (SEQ ID NO: 14).

According to some embodiments, the heavy chain CDR3 of the antibody according to the invention or a fragment thereof is selected from>GDYYGGFAVDY (SEQ ID NO: 9) and ARGDYYGGFAVDY (SEQ ID NO: 15).

10 According to some embodiments, the light chain CDR1 of the antibody according to the invention or a fragment thereof is selected from>RTSQDIGNYLN (SEQ ID NO: 10) and QDIGNY (SEQ ID NO: 16).

15 According to some embodiments, the light chain CDR2 of the antibody according to the invention or a fragment thereof is selected from>YTSRLHS (SEQ ID NO: 11) and YTS (SEQ ID NO: 17).

According to some embodiments, the light chain CDR3 of the antibody according to the invention or a fragment thereof is selected from>QQGKSLP (SEQ ID NO: 12) and QQGKSLPRT (SEQ ID NO: 18).

20 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, wherein the heavy chain CDRs consist of the sequences of SEQ ID NOs: 7, 8 and 9.

According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, wherein the heavy chain CDRs consist of the sequences of SEQ ID NOs: 13, 14 and 15.

25 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, wherein the light chain CDRs consist of the sequences of SEQ ID NOs: 10, 11 and 12.

30 According to some embodiments a monoclonal antibody which recognizes CEACAM1 or a fragment thereof comprising at least an antigen binding portion is provided, wherein the light chain CDRs consist of the sequences of SEQ ID NOs: 16, 17 and 18.

According to a specific embodiment the antibody comprises the heavy chain variable domain sequence:

According to a specific embodiment the antibody or fragment thereof comprises a heavy chain variable domain sequence consisting of the of SEQ ID NO: 26 and a light chain

variable domain sequence consisting of SEQ ID NO: 28, or an analog or derivative thereof having at least 90% sequence identity with the antibody or fragment sequence.

According to some particular embodiments the present invention provides a monoclonal antibody, or an antibody fragment comprising a set of six CDRs selected from i. SEQ ID NOs: 13, 14, 15, 16, 17, and 18 and ii. SEQ ID NOs: 7, 8, 9, 10, 11, and 12; and analogs and derivatives thereof having at least 97% sequence identity with said CDR sequences, and a framework sequence selected from mouse IgG2a, mouse IgG2b, mouse IgG3, human IgG1, human IgG2, human IgG3, wherein the monoclonal antibody binds with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.

According to a particular embodiment, a chimeric monoclonal antibody which recognizes CEACAM1 is provided, comprising at least one CDR sequence selected from the group consisting of: SEQ ID NOs: 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, and 18; and analogs and derivatives thereof having at least 97% sequence identity with said CDR sequences, and a constant region sequence selected from human IgG1, human IgG2 and human IgG3, wherein the monoclonal antibody binds with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.

According to a particular embodiment, a chimeric or humanized monoclonal antibody which recognizes CEACAM1 is provided comprising a set of six CDRs selected from i. SEQ ID NOs: 13, 14, 15, 16, 17, and 18 and ii. SEQ ID NOs: 7, 8, 9, 10, 11, and 12; and analogs and derivatives thereof having at least 97% sequence identity with said CDR sequences, and a constant region subclass selected from human IgG1, human IgG2 and human IgG3, wherein the monoclonal antibody binds with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.

According to yet another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided comprising a heavy chain sequence according to SEQ ID NO: 30.

According to yet another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided comprising a light chain sequence according to SEQ ID NO: 31.

According to yet another particular embodiment a chimeric monoclonal antibody or a fragment thereof comprising at least the antigen-binding portion, is provided comprising a human IgG1 heavy chain sequence according to SEQ ID NO: 30, and a human IgG1 light chain sequence according to SEQ ID NO: 31.

Definitions

The term “CEACAM1” is used to refer to the protein product of the CEACAM1 gene e.g., NP_001020083.1, NP_001703.2. In humans, 11 different CEACAM1 splice variants have been detected so far. Individual CEACAM1 isoforms differ with respect to the number of extracellular immunoglobulin-like domains (for example, CEACAM1 with four extracellular immunoglobulin-like domains is known as CEACAM1-4), membrane anchorage and/or the length of their cytoplasmic tail (for example, CEACAM1-4 with a long cytoplasmic tail is known as CEACAM1-4L and CEACAM1-4 with a short cytoplasmic tail is known as CEACAM1-4S). The N-terminal domain of CEACAM1 starts immediately after the signal peptide and its structure is regarded as IgV-type. For example, in CEACAM1 annotation P13688, the N-terminal IgV-type domain is comprised of 108 amino acids, from amino acid 35 to 142. This domain was identified as responsible for the homophilic binding activity (Watt et al., 2001, Blood. 98, 1469-79). All variants, including these splice variants are included within the term “CEACAM1”.

An “anti-CEACAM1 antibody”, “an antibody which recognizes CEACAM1”, “an antibody against CEACAM1”, or “an antibody to CEACAM1” is an antibody that binds to the CEACAM1 protein with sufficient affinity and specificity. Typically, an antibody according to the present teachings is capable of binding CEACAM1 with a minimal affinity of about 10^{-8} or 10^{-9} M. Some of the monoclonal antibodies of the present invention are capable of binding CEACAM3, 5 and/or 8 with a minimal affinity of about 5×10^{-7} M.

Preferably, the anti-CEACAM1 antibody of the invention can be used as a diagnostic or therapeutic agent in targeting and interfering with diseases or conditions wherein the CEACAM1 expression or activity is involved.

An “antigen” is a molecule or a portion of a molecule capable of eliciting antibody formation and being bound by an antibody. An antigen may have one or more than one epitope. The specific reaction referred to above is meant to indicate that the antigen will react, in a highly selective manner, with its corresponding antibody and not with the multitude of other antibodies which may be evoked by other antigens. An antigen according to the present invention is a CEACAM1 protein or a fragment thereof.

The term “antigenic determinant” or “epitope” according to the invention refers to the region of an antigen molecule that specifically reacts with particular antibody. Peptide sequences derived from an epitope can be used, alone or in conjunction with a carrier moiety, applying methods known in the art, to immunize animals and to produce additional polyclonal or monoclonal antibodies. Isolated peptides derived from an epitope may be used

in diagnostic methods to detect antibodies and as therapeutic agents when inhibition of said antibodies is required.

Antibodies, or immunoglobulins, comprise two heavy chains linked together by disulfide bonds and two light chains, each light chain being linked to a respective heavy chain by disulfide bonds in a "Y" shaped configuration. Proteolytic digestion of an antibody yields Fv (Fragment variable) and Fc (fragment crystalline) domains. The antigen binding domains, Fab, include regions where the polypeptide sequence varies. The term F(ab')₂ represents two Fab' arms linked together by disulfide bonds. The central axis of the antibody is termed the Fc fragment. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains (C_H). Each light chain has a variable domain (V_L) at one end and a constant domain (C_L) at its other end, the light chain variable domain being aligned with the variable domain of the heavy chain and the light chain constant domain being aligned with the first constant domain of the heavy chain (CH1). The variable domains of each pair of light and heavy chains form the antigen-binding site. The domains on the light and heavy chains have the same general structure and each domain comprises four framework regions, whose sequences are relatively conserved, joined by three hypervariable domains known as complementarity determining regions (CDR1-3). These domains contribute specificity and affinity of the antigen-binding site. The isotype of the heavy chain (gamma, alpha, delta, epsilon or mu) determines immunoglobulin class (IgG, IgA, IgD, IgE or IgM, respectively). The light chain is either of two isotypes (kappa, κ or lambda, λ) found in all antibody classes.

The term "antibody" is used in the broadest sense and includes monoclonal antibodies (including full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired biological activity.

The antibody according to the present invention is a molecule comprising at least the antigen-binding portion of an antibody. Antibody or antibodies according to the invention include intact antibodies, such as polyclonal antibodies or monoclonal antibodies (mAbs), as well as proteolytic fragments thereof such as the Fab or F(ab')₂ fragments. Further included within the scope of the invention are chimeric antibodies; human and humanized antibodies; recombinant and engineered antibodies, and fragments thereof. Furthermore, the DNA encoding the variable region of the antibody can be inserted into the DNA encoding other antibodies to produce chimeric antibodies. Single chain antibodies also fall within the scope of the present invention.

"Antibody fragments" comprise only a portion of an intact antibody, generally including an antigen binding site of the intact antibody and thus retaining the ability to bind antigen. Examples of antibody fragments encompassed by the present definition include: (i) the Fab fragment, having VL, CL, VH and CH1 domains; (ii) the Fab' fragment, which is a Fab fragment having one or more cysteine residues at the C-terminus of the CH1 domain; (iii) the Fd fragment having VH and CH1 domains; (iv) the Fd' fragment having VH and CH1 domains and one or more cysteine residues at the C-terminus of the CH1 domain; (v) the Fv fragment having the VL and VH domains of a single arm of an antibody; (vi) the dAb fragment (Ward et al., Nature 1989, 341, 544-546) which consists of a VH domain; (vii) isolated CDR regions; (viii) F(ab')₂ fragments, a bivalent fragment including two Fab' fragments linked by a disulphide bridge at the hinge region; (ix) single chain antibody molecules (e.g. single chain Fv; scFv) (Bird et al., Science 1988, 242, 423-426; and Huston et al., PNAS (USA) 1988, 85,5879-5883); (x) "diabodies" with two antigen binding sites, comprising a heavy chain variable domain (VH) connected to a light chain variable domain (VL) in the same polypeptide chain (see, e.g., EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 1993, 90, 6444-6448); (xi) "linear antibodies" comprising a pair of tandem Fd segments (VH-CH1-VH-CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions (Zapata et al. Protein Eng., 1995, 8, 1057-1062; and U.S. Pat. No. 5,641,870).

Single chain antibodies can be single chain composite polypeptides having antigen binding capabilities and comprising amino acid sequences homologous or analogous to the variable regions of an immunoglobulin light and heavy chain i.e. linked V_H-V_L or single chain Fv (scFv).

A "neutralizing antibody" as used herein refers to a molecule having an antigen-binding site to a specific receptor or ligand target capable of reducing or inhibiting (blocking) activity or signaling through a receptor, as determined by in vivo or in vitro assays, as per the specification.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier

"monoclonal" is not to be construed as requiring production of the antibody by any particular method. mAbs may be obtained by methods known to those skilled in the art. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al., Nature 1975, 256, 495, or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson et al., Nature 1991, 352, 624-628 or Marks et al., J. Mol. Biol., 1991, 222:581-597, for example.

The mAbs of the present invention may be of any immunoglobulin class including IgG, IgM, IgE, IgA. A hybridoma producing a mAb may be cultivated *in vitro* or *in vivo*. High titers of mAbs can be obtained *in vivo* production where cells from the individual hybridomas are injected intraperitoneally into pristine-primed Balb/c mice to produce ascites fluid containing high concentrations of the desired mAbs. mAbs of isotype IgM or IgG may be purified from such ascites fluids, or from culture supernatants, using column chromatography methods well known to those of skill in the art.

The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; and Morrison et al., Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984)). In addition, complementarity determining region (CDR) grafting may be performed to alter certain properties of the antibody molecule including affinity or specificity. A non-limiting example of CDR grafting is disclosed in US patent 5,225,539.

Chimeric antibodies are molecules, the different portions of which are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Antibodies which have variable region framework residues substantially from human antibody (termed an acceptor antibody) and complementarity determining regions substantially from a mouse antibody (termed a donor antibody) are also referred to as humanized antibodies. Chimeric antibodies are primarily used to reduce immunogenicity in application and to increase yields in production, for example, where murine mAbs have higher yields from hybridomas but higher

immunogenicity in humans, such that human/murine chimeric mAbs are used. Chimeric antibodies and methods for their production are known in the art (for example PCT patent applications WO 86/01533, WO 97/02671, WO 90/07861, WO 92/22653 and US patents 5,693,762, 5,693,761, 5,585,089, 5,530,101 and 5,225,539).

5 "Humanized" forms of non-human (e.g., murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman
10 primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise
15 substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further
20 details, see Jones et al., *Nature* 1986, 321, 522-525; Riechmann et al., *Nature* 1988, 332, 323-329; and Presta, *Curr. Op. Struct. Biol.*, 1992 2, 593-596.

A "human antibody" is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human
25 antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art. In one embodiment, the human antibody is selected from a phage library, where that phage library expresses human antibodies (Vaughan et al. *Nature Biotechnology* 1996 14,309-314; Sheets et al. *PNAS (USA)*, 1998, 95, 6157-6162); Hoogenboom and Winter, *J. Mol. Biol.*,
30 1991, 227, 381; Marks et al., *J. Mol. Biol.*, 1991, 222, 581). Human antibodies can also be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire.

This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al, *Bio/Technology 10*: 779-783 (1992); Lonberg et al., *Nature 368*: 856-859 (1994); Morrison, *Nature 368*:812-13 (1994); Fishwild et al., *Nature Biotechnology 14*: 845-51 (1996);
5 Neuberger, *Nature Biotechnology 14*: 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol. 13*:65-93 (1995). Alternatively, the human antibody may be prepared via immortalization of human B lymphocytes producing an antibody directed against a target antigen (such B lymphocytes may be recovered from an individual or may have been immunized in vitro). See, e.g., Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan
10 R. Liss, p. 77 (1985); Boerner et al., *J. Immunol., 147* (1):86-95 (1991); and U.S. Pat No. 5,750,373.

By the term "single chain variable fragment (scFv)" is meant a fusion of the variable regions of the heavy and light chains of immunoglobulin, linked together with a short (usually serine, glycine) linker. Single chain antibodies can be single chain composite
15 polypeptides having antigen binding capabilities and comprising amino acid sequences homologous or analogous to the variable regions of an immunoglobulin light and heavy chain (linked V_H - V_L or single chain Fv (scFv)). Both V_H and V_L may copy natural monoclonal antibody sequences or one or both of the chains may comprise a CDR-FR construct of the type described in US patent 5,091,513, the entire contents of which are incorporated herein
20 by reference. The separate polypeptides analogous to the variable regions of the light and heavy chains are held together by a polypeptide linker. Methods of production of such single chain antibodies, particularly where the DNA encoding the polypeptide structures of the V_H and V_L chains are known, may be accomplished in accordance with the methods described, for example, in US patents 4,946,778, 5,091,513 and 5,096,815, the entire contents of each of
25 which are incorporated herein by reference.

A "molecule having the antigen-binding portion of an antibody" as used herein is intended to include not only intact immunoglobulin molecules of any isotype and generated by any animal cell line or microorganism, but also the antigen-binding reactive fraction thereof, including, but not limited to, the Fab fragment, the Fab' fragment, the $F(ab')_2$
30 fragment, the variable portion of the heavy and/or light chains thereof, Fab mini-antibodies (see WO 93/15210, US patent application 08/256,790, WO 96/13583, US patent application 08/817,788, WO 96/37621, US patent application 08/999,554, the entire contents of which are incorporated herein by reference), dimeric bispecific mini-antibodies (see Muller et al., 1998) and chimeric or single-chain antibodies incorporating such reactive fraction, as well as

any other type of molecule or cell in which such antibody reactive fraction has been physically inserted, such as a chimeric T-cell receptor or a T-cell having such a receptor, or molecules developed to deliver therapeutic moieties by means of a portion of the molecule containing such a reactive fraction. Such molecules may be provided by any known
5 technique, including, but not limited to, enzymatic cleavage, peptide synthesis or recombinant techniques.

Antibodies according to the invention can be obtained by administering CEACAM1, or epitope-bearing fragments, analogs, or cells expressing, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in
10 the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., Nature 256: 495-497 (1975); Kozbor et al., Immunology Today 4: 72 (1983); Cole et al., pg. 77-96 in MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc. (1985).

Besides the conventional method of raising antibodies *in vivo*, antibodies can be
15 generated *in vitro* using phage display technology. Such a production of recombinant antibodies is much faster compared to conventional antibody production and they can be generated against an enormous number of antigens. Furthermore, when using the conventional method, many antigens prove to be non-immunogenic or extremely toxic, and therefore cannot be used to generate antibodies in animals. Moreover, affinity maturation
20 (i.e., increasing the affinity and specificity) of recombinant antibodies is very simple and relatively fast. Finally, large numbers of different antibodies against a specific antigen can be generated in one selection procedure. To generate recombinant monoclonal antibodies one can use various methods all based on display libraries to generate a large pool of antibodies with different antigen recognition sites. Such a library can be made in several ways: One
25 can generate a synthetic repertoire by cloning synthetic CDR3 regions in a pool of heavy chain germline genes and thus generating a large antibody repertoire, from which recombinant antibody fragments with various specificities can be selected. One can use the lymphocyte pool of humans as starting material for the construction of an antibody library. It is possible to construct naive repertoires of human IgM antibodies and thus create a human
30 library of large diversity. This method has been widely used successfully to select a large number of antibodies against different antigens. Protocols for bacteriophage library construction and selection of recombinant antibodies are provided in the well-known reference text Current Protocols in Immunology, Colligan et al (Eds.), John Wiley & Sons, Inc. (1992-2000), Chapter 17, Section 17.1.

Non-human antibodies may be humanized by any methods known in the art. In one method, the non-human complementarity determining regions (CDRs) are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

5 For example, US Patent 5,585,089 of Queen et al. discloses a humanized immunoglobulin and methods of preparing same, wherein the humanized immunoglobulin comprises complementarity determining regions (CDRs) from a donor immunoglobulin and heavy and light chain variable region frameworks from human acceptor immunoglobulin heavy and light chains, wherein said humanized immunoglobulin comprises amino acids
10 from the donor immunoglobulin framework outside the Kabat and Chothia CDRs, wherein the donor amino acids replace corresponding amino acids in the acceptor immunoglobulin heavy or light chain frameworks.

US Patent 5,225,539, of Winter, also discloses an altered antibody or antigen-binding fragment thereof and methods of preparing same, wherein a variable domain of the antibody
15 or antigen-binding fragment has the framework regions of a first immunoglobulin heavy or light chain variable domain and the complementarity determining regions of a second immunoglobulin heavy or light chain variable domain, wherein said second immunoglobulin heavy or light chain variable domain is different from said first immunoglobulin heavy or light chain variable domain in antigen binding specificity, antigen binding affinity, species,
20 class or subclass.

Anti-idiotypic antibodies specifically immunoreactive with an antibody of the invention are also comprehended.

Techniques for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this
25 invention. Also, transgenic mice, or other organisms such as other mammals, can be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

Alternatively, phage display technology can be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR
30 amplified v-genes of lymphocytes from humans screened for possessing anti-CEACAM1 or from libraries (McCafferty, et al., 1990, Nature 348, 552-554; Marks, et al., 1992, Biotechnology 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson et al., 1991, Nature 352:628).

The above-described antibodies can be employed to isolate or to identify clones

expressing the polypeptides to purify the polypeptides by, for example, affinity chromatography.

The invention also provides conservative amino acid variants of the antibody molecules according to the invention. Variants according to the invention also may be made that conserve the overall molecular structure of the encoded proteins. Given the properties of the individual amino acids comprising the disclosed protein products, some rational substitutions will be recognized by the skilled worker. Amino acid substitutions, *i.e.* "conservative substitutions," may be made, for instance, on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved.

A "disorder" is any condition that would benefit from treatment with the antibody. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include benign and malignant tumors; leukemias and lymphoid malignancies; neuronal, glial, astrocytal, hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, angiogenic, immunologic disorders or hyperpermeability states.

The term "therapeutically effective amount" refers to an amount of a drug effective to treat a disease or disorder in a mammal. In the case of cancer, the therapeutically effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the disorder. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. For cancer therapy, efficacy *in vivo* can, for example, be measured by assessing the duration of survival, time to disease progression (TTP), the response rates (RR), duration of response, and/or quality of life.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include melanoma, lung, thyroid, breast, colon, prostate,

hepatic, bladder, renal, cervical, pancreatic, leukemia, lymphoma, myeloid, ovarian, uterus, sarcoma, biliary, or endometrial cancer.

According to some embodiments, the antibody of the present invention is attached to a cytotoxic or therapeutic moiety. The cytotoxic or therapeutic moiety can be, for example, a
5 cytotoxic moiety, a toxic moiety, a cytokine moiety, a bi-specific antibody moiety, a cytotoxin, a chemokine, a chemotherapy, a pro-apoptotic, interferon, a radioactive moiety, or combinations thereof, examples of which are provided infra.

The term "anti-neoplastic composition" refers to a composition useful in treating cancer comprising at least one active therapeutic agent capable of inhibiting or preventing
10 tumor growth or function, and/or causing destruction of tumor cells. Therapeutic agents suitable in an anti-neoplastic composition for treating cancer include, but not limited to, chemotherapeutic agents, radioactive isotopes, toxins, cytokines such as interferons, and antagonistic agents targeting cytokines, cytokine receptors or antigens associated with tumor cells. Preferably the therapeutic agent is a chemotherapeutic agent.

As used herein the term "diagnosing" refers to determining presence or absence of a
15 pathology, classifying a pathology or a symptom, determining a severity of the pathology, monitoring pathology progression, forecasting an outcome of a pathology and/or prospects of recovery.

Pharmacology

The present invention also contemplates pharmaceutical formulations for human
20 medical use, which comprise as the active agent at least one antibody which recognizes CEACAM1, for the manufacture of a therapeutic or diagnostic composition for the treatment, diagnosis or prophylaxis of the conditions variously described herein.

In such pharmaceutical and medicament formulations, the active agent is preferably
25 utilized together with one or more pharmaceutically acceptable carrier(s) and optionally any other therapeutic ingredients. The carrier(s) must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the formulation and not unduly deleterious to the recipient thereof. The active agent is provided in an amount effective to achieve the desired pharmacological effect, as described above, and in a quantity appropriate to achieve
30 the desired daily dose.

Typically, the molecules of the present invention comprising the antigen binding portion of an antibody or comprising another polypeptide including a peptidomimetic will be suspended in a sterile saline solution for therapeutic uses. The pharmaceutical compositions may alternatively be formulated to control release of active ingredient (molecule comprising

the antigen binding portion of an antibody) or to prolong its presence in a patient's system. Numerous suitable drug delivery systems are known and include, e.g., implantable drug release systems, hydrogels, hydroxymethylcellulose, microcapsules, liposomes, microemulsions, microspheres, and the like. Controlled release preparations can be prepared through the use of polymers to complex or adsorb the molecule according to the present invention. For example, biocompatible polymers include matrices of poly(ethylene-co-vinyl acetate) and matrices of a polyanhydride copolymer of a stearic acid dimer and sebaric acid. The rate of release of the molecule according to the present invention, i.e., of an antibody or antibody fragment, from such a matrix depends upon the molecular weight of the molecule, the amount of the molecule within the matrix, and the size of dispersed particles.

The pharmaceutical composition of this invention may be administered by any suitable means, such as orally, topically, intranasally, subcutaneously, intramuscularly, intravenously, intra-arterially, intraarticular, intralesionally or parenterally. Ordinarily, intravenous (i.v.), intraarticular, topical or parenteral administration will be preferred.

It will be apparent to those of ordinary skill in the art that the therapeutically effective amount of the molecule according to the present invention will depend, *inter alia* upon the administration schedule, the unit dose of molecule administered, whether the molecule is administered in combination with other therapeutic agents, the immune status and health of the patient, the therapeutic activity of the molecule administered and the judgment of the treating physician. As used herein, a "therapeutically effective amount" refers to the amount of a molecule required to alleviate one or more symptoms associated with a disorder being treated over a period of time.

Although an appropriate dosage of a molecule of the invention varies depending on the administration route, type of molecule (polypeptide, polynucleotide, organic molecule etc.) age, body weight, sex, or conditions of the patient, and should be determined by the physician in the end, in the case of oral administration, the daily dosage can generally be between about 0.01mg to about 500 mg, preferably about 0.01mg to about 50 mg, more preferably about 0.1mg to about 10 mg, per kg body weight. In the case of parenteral administration, the daily dosage can generally be between about 0.001mg to about 100 mg, preferably about 0.001mg to about 10 mg, more preferably about 0.01mg to about 1 mg, per kg body weight. The daily dosage can be administered, for example in regimens typical of 1-4 individual administration daily. Other preferred methods of administration include intraarticular administration of about 0.01mg to about 100 mg per kg body weight. Various considerations in arriving at an effective amount are described, e.g., in Goodman and

Gilman's: The Pharmacological Bases of Therapeutics, 8th ed., Pergamon Press, 1990; and Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Co., Easton, Pa., 1990.

Suitable dosing regimens of combination chemotherapies are known in the art and described in, for example, Saltz et al. Proc ASCO 1999, 18, 233a and Douillard et al., Lancet
5 2000, 355, 1041-7.

The molecules of the present invention as active ingredients are dissolved, dispersed or admixed in an excipient that is pharmaceutically acceptable and compatible with the active ingredient as is well known. Suitable excipients are, for example, water, saline, phosphate buffered saline (PBS), dextrose, glycerol, ethanol, or the like and combinations thereof.
10 Other suitable carriers are well known to those skilled in the art. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents.

The pharmaceutical composition according to the present invention may be administered together with an anti-neoplastic composition. According to a specific
15 embodiment the anti-neoplastic composition comprises at least one chemotherapeutic agent. The chemotherapy agent, which could be administered together with the antibody according to the present invention, or separately, may comprise any such agent known in the art exhibiting anticancer activity, including but not limited to: mitoxantrone, topoisomerase inhibitors, spindle poison vincas: vinblastine, vincristine, vinorelbine (taxol), paclitaxel,
20 docetaxel; alkylating agents: mechlorethamine, chlorambucil, cyclophosphamide, melphalan, ifosfamide; methotrexate; 6-mercaptopurine; 5-fluorouracil, cytarabine, gemcitabin; podophyllotoxins: etoposide, irinotecan, topotecan, dacarbazine; antibiotics: doxorubicin (adriamycin), bleomycin, mitomycin; nitrosoureas: carmustine (BCNU), lomustine, epirubicin, idarubicin, daunorubicin; inorganic ions: cisplatin, carboplatin; interferon,
25 asparaginase; hormones: tamoxifen, leuprolide, flutamide, and megestrol acetate.

According to a specific embodiment, the chemotherapeutic agent is selected from the group consisting of alkylating agents, antimetabolites, folic acid analogs, pyrimidine analogs, purine analogs and related inhibitors, vinca alkaloids, epipodophyllotoxins, antibiotics, L-asparaginase, topoisomerase inhibitor, interferons, platinum coordination complexes,
30 anthracenedione substituted urea, methyl hydrazine derivatives, adrenocortical suppressant, adrenocorticosteroides, progestins, estrogens, antiestrogen, androgens, antiandrogen, and gonadotropin-releasing hormone analog. According to another embodiment, the chemotherapeutic agent is selected from the group consisting of 5-fluorouracil (5-FU), leucovorin (LV), irinotecan, oxaliplatin, capecitabine, paclitaxel and doxorubicin. Two or more

chemotherapeutic agents can be used in a cocktail to be administered in combination with administration of the anti-CEACAM1 antibody.

The following examples are intended to illustrate how to make and use the compounds and methods of this invention and are in no way to be construed as a limitation.

5 Although the invention will now be described in conjunction with specific embodiments thereof, it is evident that many modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such modifications and variations that fall within the spirit and broad scope of the appended claims.

10 **EXAMPLES**

Means for preparing and characterizing antibodies are well known in the art. A description follows as to exemplify techniques for the production, characterization and use of anti-CEACAM1 antibodies in accordance with the present invention.

15 Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., 1989; "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. 1994; Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland 1989; Perbal, 20 "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York 1988; Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York 1998; methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III 25 Cellis, J. E., ed. 1994; "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. 1994; Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT 1994; Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York 1980; available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 30 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. 1985; "Transcription and Translation" Hames, B. D., and Higgins S. J., Eds. 1984; "Animal Cell Culture" Freshney, R. I., ed. 1986; "Immobilized Cells and Enzymes" IRL

Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., 1984 and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA 1990; Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press 1996. The
5 procedures therein are believed to be well known in the art.

Example 1: Generation and characterization of monoclonal antibodies which recognized CEACAM

Monoclonal antibodies that effectively block the CEACAM1 homophilic interactions
10 in vitro at nanomolar concentrations were generated by immunizing mice with recombinant human CEACAM1 protein. Hybridomas producing the CEACAM1-blocking antibodies were produced and re-cloned several times to yield a stable clone.

The DNA and amino acid sequence of one exemplary monoclonal antibody which recognizes CEACAM1 was determined by Fusion Antibodies Ltd. mRNA was extracted
15 from the hybridoma cell pellets and total RNA was extracted from the pellets using RNA extraction protocol. RT-PCR-cDNA was created from the RNA by reverse-transcription with an oligo(dT) primer. PCR reactions using variable domain primers were used to amplify both the VH and VL regions of the monoclonal antibody DNA.

The VH and VL products were cloned into the Invitrogen sequencing vector pCR2.1
20 and transformed into TOP10 for positive transformants. Selected colonies were picked and analyzed through sequencing. The resulted DNA and amino acid sequences determined are:

Variable heavy chain (VH)

DNA sequence of the VH domain:

ATGGGATGGACCTTGGTCTTTCTCTTTCTCCTGTCAGTAACTGCAGGTGTTCACTC
25 CCAGGTCCAGCTGCAGCAGTCTGGAGCTGAGCTGGTAAGGCCTGGGACTTCAGT
GAAGGTGTCCTGCAAGGCTTCTGGATACGCCTTCACTAATAACTTGATAGAGTGG
GTAAAACAGAGGCCTGGACAGGGCCTTGAGTGGATTGGAGTGATTAATCCTGGA
AGTGGTGATACTAACTACAATGAGAAGTTCAAGGGCAAGGCAACACTGACTGCA
GACAAATCCTCCAACACTGCCTACATGCAGCTCAGCAGCCTGACATCTGATGACT
30 CTGCGGTCTATTTCTGTGCAAGAGGGGATTACTACGGTGGCTTTGCTGTGGACTA
CTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCAGCCAAAACGACACCCCCATCC
GTTTATCCCTTGGCCCCTGGAAGCTTGGG (SEQ ID NO: 25).

Amino acid sequence of the VH domain:

QVQLQQSGAELVRPGTSVKVSKASGYAFTNNLIEWVKQRPGQGLEWIGVINPGSG
 DTNYNEKFKGKATLTADKSSNTAYMQLSSLTSDDSA VYFCARGDYGGFAVDYWG
 QGTSVTVSS (SEQ ID NO: 26).

5 Variable light chain (VL)**DNA sequence of the VL domain:**

ATGGTGTCTCAGCTCAGTTCCTTGGTCTCCTGTTGCTCTGTTTTCAAGGAACCAG
 ATGTGATATCCAGATGACACAGACTACATCCTCCCTGTCTGCCTCTCTGGGAGAC
 AGAGTCACCATCAGTTGCAGGACAAGTCAGGACATTGGCAATTATTTAAACTGG
 10 TATCAGCAGAAACCAGATGGAAGTGTAAACTCCTGATCTACTACACATCAAGAT
 TACTCAGGAGTCCCATCAAGGTTTCAAGTGGCAGTGGGTCTGGAACAGATTATTC
 TCTCACCATTAGCAACCTGGAGCAAGAAGATATTGCCACTTACTTTTGCCAACAG
 GGTAAGCCTTCCTCGGACGTTCCGGTGGAGGCACCAAGTTGGAAATCAAACGG
 GCTGATGCTGCACCAACTGTATCCATCTTCCCACCATCCAGTGAGCAGTTAACAT
 15 CTGGAGGTGCCTCAGTCGTGTGCTTCTTGAACA ACTTCTACCC CAGAGA (SEQ ID
 NO: 27).

Amino acid sequence of the VL domain:

DIQMTQTTSSLSASLGDRVTISCRTSQDIGNYLNWYQQKPDGTVKLLIYYTSRLHSGV
 PSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGKSLPRTFGGGTKLEIK (SEQ ID NO:
 20 28).

N-terminal amino-acid sequencing and Mass-Spectra analysis were used to confirm the VL and VH identities.

Example 2: Verification of N-terminal amino acid sequence

25 Amino acid sequence analysis of the light chain was performed by the Edman degradation method to verify the N-terminus sequence of the light chain of one of the monoclonal antibodies. The obtained N-terminal sequence was: DIQMTQTTSS (SEQ ID NO: 29), which is in accordance with the N-terminal expected sequence based on the DNA sequence.

30

Example 3: Complementary determining region (CDR) sequences

The CDR segments were identified using two different algorithm methods:

1. IMGT algorithm (Lefranc et al., 1999, Nucleic Acids Research, 27, 209-212);
2. KABAT algorithm (Wu TT and Kabat E.A., 1970, J. Exp. Med. 132, 211-250).

Table 1 summarizes the determined CDR sequences using the two methods as well as the minimal consensus sequence and combined sequence of sequences identified using both methods.

Table 1. CDR sequences

	VH1	VH2	VH3	VL1	VL2	VL3
IMGT	GYAFTNNL (SEQ ID NO: 13)	INPGSGDT (SEQ ID NO: 14)	ARGDYYGG FAVDY (SEQ ID NO: 15)	QDIGNY (SEQ ID NO: 16)	YTSR (SEQ ID NO: 17)	QQGKSLPR T (SEQ ID NO: 18)
KABAT	NNLIE (SEQ ID NO: 7)	VINPGSGDT NYNEKFKG (SEQ ID NO: 8)	GDYYGGFA VDY (SEQ ID NO: 9)	RTSQDIGNY LN (SEQ ID NO: 10)	YTSRLHS (SEQ ID NO: 11)	QQGKSLP (SEQ ID NO: 12)
Combined sequence	GYAFTNNLI E (SEQ ID NO: 19)	VINPGSGDT NYNEKFKG (SEQ ID NO: 20)	ARGDYYGG FAVDY (SEQ ID NO: 21)	RTSQDIGNY LN (SEQ ID NO: 22)	YTSRLHS (SEQ ID NO: 23)	QQGKSLPR T (SEQ ID NO: 24)
Consensus sequence	X ₁ NNLX ₂ * (SEQ ID NO: 1)	INPGSGDT (SEQ ID NO: 2)	GDYYGGFA VDY (SEQ ID NO: 3)	QDIGNY (SEQ ID NO: 4)	YTSR (SEQ ID NO: 5)	QQGKSLP (SEQ ID NO: 6)

5 * wherein X₁ is absent or is Thr (T) and X₂ is absent or is Ile (I)

Example 4: Design and production of a chimeric monoclonal antibody

The DNA sequence of the variable heavy and light chains (SEQ ID NOs 25 and 27) were used to construct a chimeric antibody, comprising the human IgG1 isotype constant domains and constant light (CL) human IgKappa domain. Although the parent monoclonal antibody is mouse IgG1 and its human equivalent is IgG4, a human IgG1 framework was used to construct some of the chimeric antibodies of the present invention. The DNA sequences for the light chain and heavy chain were synthesized and cloned into the expression vector pFUSION-DHFR1 under separate promoters.

15 Transient Transfection of CHO cells

Suspension CHO cells (Invitrogen, UK) were cultivated at 130rpm, 8% CO₂, 37°C in Pro CHO 5 serum free medium (Lonza, UK) in 250 and 500ml vented Erlenmeyer flasks (Corning, Netherlands). On the day of transfection, cells were seeded at a density of 2.0 X 10⁶ cells/ml, 2.5g/ml of plasmid DNA (Geneart, Germany) was transfected into the cells using Polyethylenimine (Polysciences Inc, PA, US). Transfected cultures were incubated at 20 130 rpm, 8% CO₂, 37°C for 9-10 days. Prior to harvest of the culture supernatants were spinned at 4,000 rpm for 40 minutes.

Media was harvested and purified in two separate batches. The media was filtered through a 0.8µm gyrodisc filter and purified using a 1ml Protein A column. The antibody was

purified by FPLC. 320ml sample was loaded at 0.2mls/min overnight and increased to 0.5mls/min after 17 hours. Column was washed/equilibrated with PBS at 0.5mls/min before elution with pH 3.0 Gly/HCL elution buffer. A good peak was observed and fractions 1 to 5 were quantified by Bradford Assay. The Bradford assay showed protein present in fractions 1-4 which were pooled and dialyzed for buffer exchange overnight in 1 liter of PBS (4°C, 120 RPM). A concentration of 1.823 mg/ml was observed for the 4 ml sample, therefore a total yield of approximately 7.32 mg was purified. In the second batch, the results of the Bradford assay showed protein present in fractions 1-3 which were pooled and dialyzed. From the 320 ml conditioned medium a total yield of approximately 7.32 mg was purified (batch A). From a 910 ml culture a total yield of approximately 15.74 mg was purified (batch B). Total transient expression yielded about 23 mg of purified protein. After concentration determinations and SDS/PAGE analysis, 19.65 mg of the chimeric antibody were yielded. The purified antibody samples were analyzed by SDS-PAGE to assess purity. Figure 1 depicts the SDS-PAGE gel image showing light and heavy chains of the chimeric antibody. MS analysis revealed that the molecular weight of CM10 heavy chain is 48.6 KDa and of light chain is 23.3 KDa.

The resulted antibody, denoted CM10, has the following amino acid sequence of the heavy and light chains:

Heavy chain amino acid sequence (without signal peptide):

20 **QVQLQQSGAELVRPGTSVKVSCKASGYAFTNNLIEWVKQRPGQGLEWIGVINPGSG**
DTNYNEKFKGKATLTADKSSNTAYMQLSSLTSDDSAVYFCARGDYGGFAVDYWG
QGTSVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN
25 AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPRE
PQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGDSFF
LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 30).

Variable domain is in bold, CDRs according to IMGT are underlined.

Light chain amino acid sequence (without signal peptide):

30 **DIQMTQTTSSLSASLGDRVTISCRTSQDIGNYLNWYQQKPDGTVKLLIYYTSRLHSG**
VPSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGKSLPRTFGGGTKLEIKRTVAAPSV
FIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL
SSTLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:31)

Variable domain is in bold, CDRs according to IMGT are underlined.

A plasmid containing the DNA sequences of the heavy and light chains of an exemplary chimeric monoclonal antibody denoted CM10 was deposited on September 28, 2011 under ATCC Accession Number PTA-12130.

5 **Example 5: Affinity characterization of the chimeric monoclonal antibody CM10**

Binding of CM10 to purified human CEACAM1

The binding specificity of CM10 to human CEACAM1 was tested in ELISA assay using purified human CEACAM1. Indirect ELISA using 23 double dilutions of CM10 were used to generate specific binding curve. The results shown in Figure 2 represent average O.D. from triplicate \pm SE. Similar results were obtain from other ten independent experiments.

In order to test whether the chimerization process affect the binding affinity of the antibody, the chimeric antibody CM10 was evaluated for CEACAM1 binding by competitive ELISA and by BIAcore analysis.

For the ELISA, recombinant purified human CEACAM1 was bound to the plate. The chemically biotinylated CM10 was used as tracer at a constant concentration and was competed with increasing concentrations of unlabelled CM10. Following incubation and washing, the plate was developed with a StrepAvidin- HRP conjugate and the color reaction was developed with TMB as an HRP substrate.

Using 50 ng/ml of tracer, the apparent affinity values detected for CM10 were 1.2-1.6 nM.

BIAcore analysis

Each antibody was immobilized onto a single channel of a CM5 sensor chip of Biacore3000 instrument by NHS-EDC coupling chemistry.

Recombinant CEACAM1 was flowed at 50 μ l/min over the chip in various concentrations (0.19, 0.39, 0.78, 1.56, 3.12, 6.25, 12.5 and 25nM). The running buffer was PBS-ET(10mM p-buffer pH7.4, 150mM NaCl, 3.4mM EDTA and 0.005% tween 20). The data were analyzed using BIAEvaluation software 3.0 and the KD values of CM10-CEACAM1 affinity, calculated from three independent experiments were (KD): 4.07- 5.05 nM (average 4.56 nM).

Binding specificity of CM10 to membrane-bound endogenous CEACAM1

In order to test the binding of CM10 to membrane-bound endogenous CEACAM1, a FACS analysis was performed. Several human melanoma cell lines were screened for hCEACAM1 expression while 526mel cell line was used as positive control and 003mel as

negative control line. 526mel, 003mel, Malme 3M, Skmel 5 and A375 cell lines were stained with CM10. Empty histograms represent mAb staining while darker histograms represent background staining. At least 5000 cells were used to analyze CEACAM1 expression in each histogram.

5 As can be appreciated from Figure 3, CM10 detects membrane-bound endogenous CEACAM1. Malme3M and Skmel5 cell lines showed high expression of CEACAM1 while no expression could be detected in A375 melanoma cell-line.

Conclusion

10 The chimerization process was carried out successfully. The chimeric antibody bind CEACAM1 in an affinity of 1.4 nM as validated by two different approaches The FACS analysis testing the binding specificity of CM10 to various melanoma cell lines, demonstrate that the antibody retained its biological binding ability.

Example 6. Assessing the activity of CM10

15 Blocking cell-cell interaction assay

The assay which determines the ability of anti CEACAM1 mAb to blocks CEACAM1 cell-cell interaction uses murine T cells (BW cells) that are stably transfected with a chimeric molecule composed of the extracellular portion of human CEACMA1 fused to mouse z-chain (BW/CEACAM1). Engagement of CEACAM1 by co-incubation of
20 BW/CEACAM1 cells with B cells stably transfected with CEACAM1 (221/CEACAM1), lead to the secretion of mouse IL-2, mediated by the z-chain.

Effectors cells (BW, expressing CEACAM1) were incubated in the presence of CM10 or PBS for 30 minutes on ice. Following the incubation the effectors cells were co-culture over night with target cell expressing CEACAM1 (221+) or negative to CEACAM1 (221-). Mouse
25 IL-2 secretion was measured by commercial ELISA. The results shown in Figure 4 represent average IL-2 secretion from duplicate wells. As shown in the figure, upon addition of CM10, CEACAM1-mediated cell-cell interactions between T and B cells were abolished in a dose-dependent manner as indicated by blockage of IL-2 secretion.

In-vitro immunomodulatory killing assays

30 T cells Killing assay: Melanoma-reactive T cells **TILs** (Tumor Infiltrating Lymphocytes, derived from melanoma patients) can destroy melanoma cells with matched HLA. TILs were purchased from ELLA Institute at Shiba medical center and were growth according to the clinical lab protocols. CFSE-labeled melanoma cells (SKmel5) were pre-incubated with CM10 (10µg/ml) for 30 minutes on ice. TIL were added for additional 10

hours incubation at 37°C. Percentage of killing was determined by PI-staining of the CFSE labeled melanoma cells. Effector-to-target ratio was 5:1. In another assay, CFSE-labeled melanoma cells were pre-incubated with CM10 for 30 minutes on ice. TIL were added for additional 5 hours incubation at 37°C. Percentage of killing was determined by PI-staining of the CFSE labeled melanoma cells. Results represent average of % specific killing from triplicate wells \pm SE per treatment. Effector-to-target ratio was 5:1.

The results describes in Figure 5 show that the killing activity of T cells is enhanced in the presence of the anti CEACAM1 mAb, CM10. Furthermore, in assay conditions where TILs where unable to kill any melanoma cells (short incubation or low TIL ratio), addition of CM10 stimulated the killing activity of TILs while no killing could be detected with IgG1 isotype control (Figure 6).

NK cells Killing assay

Natural killer cells (NK cells) are a type of cytotoxic lymphocyte that can destroy malignant cells by releasing small proteins called perforin and granzyme that cause the target cell to die by apoptosis. NK cells can be activated through several different pathways among them cytokines, FC receptor, and MHC class 1 absence at the target cells. Several activation and inhibition receptors to various ligands on target cells regulate the final cytotoxicity activity of NK cells. NK 92MI cells are IL-2 independent NK cell line that were purchased from the ATCC.

NK 92MI were incubated with CM10 (0.2 μ g/ml, 1 μ g/ml or 5 μ g/ml) or isotype match control Ab (5 μ g/ml) for 30 minutes at 37°C, target cells expressing CEACAM1 were added for additional 5 hours. Percentage of killing was determined by classical LDH release assay. Results represent average of % cytotoxicity from triplicate wells \pm SE per treatment. Effector-to-target ratio was 2.5:1.

The assay describes in Figure 7 shows that CM10 strongly enhanced the killing activity of NK cells on two melanoma cell lines expressing CEACAM1 (SKMel 5 and G361), compare to PBS or isotype match IgG. Similar results have been demonstrated in two other CEACAM1+ melanoma cell lines and in various effector-to-target (E:T) ratios.

Antibody-Dependent Cell-mediated Cytotoxicity

Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC) is a mechanism of cell-mediated immunity whereby an effector cell of the immune system (mostly NK cells) actively lyses target cells that has been bound by a specific antibody. In order to assess the safety profile of CM10 we have conducted preliminary ADCC assay where the ability of

CM10 to induce ADCC was examined in three melanoma cell lines (CEACAM1 positive cell lines: G361 and SKmel5 and CEACAM1 negative cell line: SKmel28).

The ability of CM10 to induce ADCC was examined in comparison to a positive control antibody (Polyclonal Ab anti CEACAM1 that showed ADCC activity in preliminary experiment). Isotype matched antibody served as negative control (hIgG1 K). The results indicate that CM10 do not trigger ADCC in the setting tested.

Complement-Dependent Cytotoxicity

Complement proteins are found in the blood, and their action "complements" the work of antibodies. Complement-dependent cytotoxicity (CDC) is a mechanism of killing cells in which antibody bound to the target cell surface fixes complement, results in assembly of the membrane attack complex that create pores in the target cell membrane and finally lead to cell lyses.

In order to access the safety profile of CM10, the ability of CM10 to induce CDC was examined in two melanoma cell lines (SKmel28 and SKmel5) expressing CEACAM1. Commercial pooled human serum was used as complement proteins source. The commercial monoclonal antibody Rituximab incubated with Daudi cells was used as the assay positive control and commercial IgG1K as isotype match control to CM10. Melanoma cell lines – SKMEL5 and SKMEL28 or positive control Daudi cells were incubated with CM10 or Rituxiamab respectively for 1 hour in room temperature followed by the addition of normal human serum at a final concentration of 50% for 2 additional hours in a humidified incubator (37°C, 5% CO₂). The percent of lysed cells was determined by Propidium iodine (PI) staining. The results (Figure 14) represent the average + S.E of 2 individual experiments preformed in duplicates, indicating that CM10 did not induce CDC lysis in the setting tested.

In-vivo efficacy experiment

The purpose of this experiment is to test the direct effect of CM10 on melanoma cells in-vivo, as well as to evaluate the immunomodulatory effect, which is missing in the xenograft setting.

Calibration of xenograft experiments

CEACAM1 positive human melanoma cell line purchased from "ATCC" (SKMel5) and NOD-SCID, age matched mice from "Harlan laboratories" were used. The calibration assay was conducted in order to monitor the growth of the tumors and to find the optimal TILs regime. SKMel5 melanoma cells were injected SC (subcutaneous) to SCID-NOD mice and tumor volume was monitored by physical measurements. When the tumor volume reached 100mm³, the mice where divided into 5 randomized groups. The TILs were

injected either IT (Intra Tumoral) at two different concentrations or IV (Intra Venus) at one concentration (20×10^6 per mice) while one group received only one injection and the second group received 2 TIL injections. Each TIL injection was followed by 5 days of hIL-2 administration. The calibration experiment demonstrated that TIL IV injection has higher effect on tumor size than IT administration. In addition repetitive TIL regime provides better tumor growth inhibition over single injection, as could be predicted from T cells half life. Based on this data, TIL will be administrated every 10 days by IV injections, in future xenograft experiments.

In-vivo immunomodulatory, anti-cancer activity of CM10

Human CEACAM1 positive SKmel5 melanoma cells were injected SC to SCID-NOD mice. When the tumors reached a volume of approximately 100mm^3 , the mice were randomized to one of the following treatment groups: a) Weekly IV injections of PBS; b) Weekly IV injections of 0.45mg CM10; c) Three IV injection of 20×10^6 anti-tumor reactive human T cells (TIL) and weekly IV injections of PBS; d) Three IV injections of 20×10^6 anti-tumor reactive human T cells and weekly IV injections of 0.45mg CM10. A person blind to the experimental setting measured the tumors volume 2-3 times per week. The results of Figure 8 represent average tumor volume \pm SE from 6-10 mice per group. Arrows indicate time of administration (CM10 cycle, TIL triangles, CM10 and TIL open squares). As shown, a moderate inhibition of tumor growth was observed either with CM10 alone or with TIL only, but the differences did not reach statistical significance when compared to the control treatment. Strikingly, the combination of adoptive human T cell transfer with CM10 injections exhibited significant synergism and strongly inhibited xenograft growth. This observation concurs with the *in-vitro* data showing the potentiating effect of CM10 on T cell killing (Figures 5 and 6).

Significant growth inhibition was observed in the group treated with CM10 in the presence of TIL. These results reinforce the immunostimulatory effect of anti CEACAM1 mAb in different cell lines, and indicate that CM10 can be used as promising immunomodulatory antibody. This observation concurs with the *in-vitro* data showing the stimulatory effect of CM10 on melanoma cells killing by T cell.

Conclusions

CEACAM1 is known as a regulator of lymphocyte activation. CM10 is an antibody that blocks the interactions between two CEACAM1 molecules (Figure 4) and therefore eliminates the inhibitory signals mediated by CEACAM1, results in stronger cytotoxic lymphocytes activation against tumor cells (Figures 6 and 7). The *in-vivo* xenograft result

(Figure 8) reinforce the immunostimulatory nature of CM10 and demonstrate significant growth inhibition of tumors in mice treated with CM10 in the presence of TIL. The scheme presented in Figure 9 demonstrates a non-limitative theory of the mode of action of CM10 that prevents CEACAM1-CEACAM1 interaction enabling activation of killing signals by immune system cells.

Example 7: CM10 in-vitro safety assessment

The effect of CM10 on normal human cells

In order to evade from the immune system cancer cells alter the expression of many molecules. Several evidences have showed that CEACAM1 expression is increasing during the malignance transformation of melanoma cells. According to the literature CEACAM1 is also expressed on normal cells, therefore it is important to map the possible binding sites of CM10 in the body and to identify if binding of CM10 to normal cells may lead to any undesired outcome.

Normal human tissues cross reactivity

In this study, the binding intensity of anti-CEACAM1 mAb in a human tissue microarray containing normal and malignant melanoma samples was examined. The binding intensity was assessed using a standard pathological scoring system. Tissue micro array (TMA) containing 100 cases of malignant melanoma (primary, metastasis) and of benign nevi were analyzed for anti CEACAM1 binding intensity by standard IHC procedure. Each core of tumor was graded from 0 to +3. As shown in Figure 10, binding intensity of anti-CEACAM1 mAb was seen in more than 50% of the melanoma samples and in 65% of metastatic melanoma samples.

The multi normal human organ tissue microarray (TMA) included 33 types of normal organs, each type taken from 3 normal human individuals. The age ranged from 2 - 67 years, 43 specimens were derived females and 57 specimens from males. The following tissues were negative for anti-CEACAM1 mAb binding: Cerebrum, cerebellum, ovary, pancreas, parathyroid gland, hypophysis, thyroid gland, tonsil, bone marrow, spleen, thymus, lung, cardiac muscle, stomach, skeletal muscle, skin, peripheral nerves, mesothelium and retina. A cell-specific staining was detected in some organs, mainly on the luminal side of epithelial cells forming ducts or glands in hollow visceral organs such as: brush border of small intestine; some apical colonic glands; Breast ductal epithelium; Liver bile canaliculi; inner surface of renal tubules; few Endometrial glands ;luminal part of Salivary gland. In addition, some low cellular staining was observed in adrenal gland cortex, apical surface of prostatic

glands, Leydig cells of testis and single scattered cells in the pancreas. The only cells of the immune system that found positive were neutrophils within capillaries. No staining of lymphocytes was found in tissues and lymphatic organs. Finally, weak to moderate positive staining was found in endothelial cells of small blood vessels at selective sites, including:
5 ovary, adrenal gland, kidney, and rarely in pancreas, prostate, hypophysis and endometrium.

The IHC analysis showed a strong anti CEACAM1 staining of melanoma cells, as compared to no staining of the vast majority of the tissues tested in a normal human tissue. Nevertheless, some selective staining was observed in the luminal aspect of epithelial cells of ducts or glands in hollow viscera. This cellular aspect is generally less accessible to an
10 antibody administered via the peripherally blood.

Quantification of CM10 molecules bound per cell

In order to quantify the exact number of CM10 molecules bound to each cell types the QuantiBRITE kit was used. Using the kit the MFI (mean fluorescence intensity) was directly translated to the number of molecules bound per cell. Three human primary cells of tissues,
15 which were found to be positive for anti-CEACAM1 binding, were purchased from ATCC. HUVEC cells to represent the positive staining found in endothelial cells; primary prostate epithelial cells, since the apical surface of prostate glands showed positive staining, and primary renal proximal tubule epithelial cells since the inner surface of tubules stained positive. In more details, SkMel 5, G361, Malme 3M, NK 92MI, HUVEC, Renal primary
20 cells and prostate primary cells were grown according to ATCC protocols. CM10 was conjugated to a PE molecule (RPE LYNX Rapid Conjugation Kits Serotec) according to the manufacturer's protocol and was used (1µg/ml) with the QuantiBRITE PE beads kit (BD) to determine the ABC (antibodies bound per cell) by Flow cytometry. The number of CM10 molecules bound per cell was analyzed using flow cytometry in the indicated primary cells in
25 comparison to melanoma cell lines. At least 10000 cells were counted for each cell line.

Quantitative analysis (Figure 11, the results represent the average of 2-3 independent experiments \pm SE) showed that the CEACAM1-positive melanoma cells bind between 20,000-50,000 CM10 molecules, while normal endothelial and epithelial cells (e.g. HUVEC, Kidney and Prostate) which have been reported to exert some CEACAM1 expression, bind
30 up to 2,000 CM10 molecules only. Furthermore, high numbers of CM10 antibodies were bound to the NK cells (\sim 20000) which correlates with published data showing high CEACAM1 expression on activated lymphocytes. These results reinforce the safety profile of CM10 (low expression on primary cells) and its activity (NK results) as a player in activated lymphocyte-mediated cell lysis.

Proliferation of human primary cells in the presence of CM10

Since some positive staining was found in normal tissues the effect of CM10 on primary cell growth was examined. HUVEC and primary prostate cells were grown according to the ATCC protocols and were monitored for cell proliferation using XTT standard assay. No effect on cell-proliferation could be detected.

Summary

The binding profile of anti CEACAM1 mAb to normal tissues and melanomas cells was identified. CEACAM1 is absent on normal melanocytes but undergoes neo-expression and is widely expressed on the vast majority of metastatic melanoma specimens (Figure 10), in other normal organs there is restricted expression of CEACAM1 in specific cells within several tissues. A quantitative analysis, measuring the number of CM10 molecules that are bound to cells, revealed a very low number of bound-CM10 on normal human primary cells (Figure 11). These results imply that the majority of CM10 molecules injected to patients will mainly target cancer cells and not normal tissues due to expression differences. Furthermore, CM10 has no effect on cell proliferation, CDC activity, and negligible or low ADCC activity which suggests that the binding of CM10 to non-target cells, would not result in unwanted cell outcome.

The effect of CM10 on the immune system

CM10 is an immunostimulatory antibody that blocks the interaction between two CEACAM1 molecules and by doing so mediates stimulation of lymphocytes against malignant cells. It is important to verify that the antibody will not cause unleashed stimulation of the immune system which can cause severe adverse events. In normal lymphocytes there is a neglect expression of CEACAM1 on the cell membrane, only following cell activation CEACAM1 is mobilized to the membrane, where it rapidly and strongly up-regulated on activated lymphocytes (Gray-Owen and Blumberg 2006, Nat Rev Immunol 6, 433-46). As demonstrated above, no cross reactivity to normal lymphatic tissues (Spleen, Thymus, Bone Marrow) or to lymphocytes was observed; nevertheless various *in-vitro* and *ex-vivo* immuno-toxicity analyses were performed to help in predicting potential side effects.

The effect of CM10 on human lymphocytes proliferation

One of the most common and acceptable ways to evaluate safety of immunomodulatory antibodies *in-vitro* is by examining its effects on proliferation and cytokine secretion of normal human PBMC (peripheral blood mononuclear cells). Human PBMC from 3 unrelated donors were isolated and incubated with CM10 using 3 different

concentrations (2 μ g/ml, 20 μ g/ml or 200 μ g/ml) or with a control mAb IgG1K (200 μ g/ml) for 60 minutes and with eight replicate wells. PHA (1 μ g/ml) was added to 4 of the assay replicate wells and cells were incubated for 96 hours. 3H-Thymidine incorporation used to assay cell proliferation. Mock stimulated cells and PHA only stimulated cells used as assay's
5 negative and positive controls respectively. The proliferation was assessed in resting lymphocytes as well as in activated lymphocytes (PHA treated). Mock stimulated cells and PHA only stimulated cells used as assay's negative and positive controls respectively. The results (Figure 12) clearly show that CM10 has no effect on the proliferation of naïve or activated human lymphocytes. In addition to the PBMC proliferation study, the effect of
10 CM10 on cytokine secretion from human PBMC is assessed. Cytokine secretion studies define the immunomodulatory effect of CM10 and assist in predicting potential side effects.

Example 8. CM10 selectivity panel.

Characterization of the binding profile was performed using Cell lines over-
15 expressing the different CEACAM family proteins and flow cytometry analysis. .

In humans, the CEA family is encoded by 18 genes and 11 pseudogenes on chromosome 19q13.2. Several closely related members belong to the CEACAM family (CEACAM1,3,4,5,6,7,8) and are differentially expressed by various human cell types. The CEACAM proteins have been implicated in various adhesion mediated effects that govern
20 the growth and differentiation of normal and cancerous cells (Gray-Owen and Blumberg 2006, Nat Rev Immunol 6, 433-46). The closely related proteins in the family share a high amino acid similarity that varies from 45% up to 90% similarity between certain members.

A standard FACS protocol was used with CM10 conjugated to a Biotin molecule and Strep-Avidin APC as secondary agent. 721.221 cells expressing CEACAM 1,5,6,8 or HEK
25 293 T transient expressing CEACAM 3, 4, were stained with biotinylated CM10 (1 μ g/ml) and Strep-Avidin APC as secondary agent. Empty histograms represent mAb staining while red histograms represent background staining. At least 10000 cells were used to analyze CM10 binding in each histogram. FAB was calculated by dividing the MFI of the stained cells in the MFI of the background staining. The results demonstrated in Figure 13, clearly
30 indicate that CM10 bind strongly to cells expressing CEACAM1. Moderate staining was observed in cells expressing CEACAM3 and 5. Weak or neglect binding was demonstrated in cells expressing CEACAM 4, 6, 8.

Conclusions

CM10 is a mAb developed to recognize human CEACAM1, a protein that was found to be associated with cancer, in general, and with Melanoma in particular. The over-expression of CEACAM1 has been identified in a few malignancies among them melanoma, NSCLC, Thyroid cancer and gastric cancer. The evidence indicates that over-expression of CEACAM1 can be correlated with poor prognosis in melanoma and NSCLC patients.

CEACAM5 has been found to be over-expressed in a high percentage of many human tumors, including 90% of gastrointestinal, colorectal (CRC) and pancreatic cancers, 70% of non-small cell lung cancer cells and 50% of breast cancers. It is also over-expressed in thyroid, stomach, ovarian and uterine cancers (Thompson, Grunert et al. 1991, J Clin Lab Anal 5, 344-66). CEACAM5 even serves as a clinical marker for liver metastasis in CRC and post-surgical surveillance of colon cancer (Duffy 2001, Clin Chem 47, 624-30). The evidence that CM10 is capable to bind CEACA5 is very important and can expand the possible indications that can be treated by CM10 from 4-5 types of malignancies to above 10. The anti-CEACAM5 agents that have entered clinical trials include anti-CEACAM5 antibodies conjugated to toxic substances such as radioactive substances for both diagnostic purposes and for the treatment of various malignancies. It seems that even these toxic conjugated forms don't show safety problems, which can indicate that CEACAM5 is a safe target (Liersch, et al. 2005, J Clin Oncol 23(27), 6763-70; Ychou, et al. 2008, Clin Cancer Res 14(11), 3487-93). On the other hand, none of these agents target the immunological regulation of tumors, which can be targeted by an antibody which can bind both CEACAM1 and CEACAM5, such as CM10.

Example 9: Epitope mapping of CM10

To reconstruct discontinuous epitopes of the target molecule, a library of structured peptides was synthesized. The target molecule was the N-domain of the human CEACAM1 molecule having the sequence:

```

1      QLTTESMPFN VAEGKEVLLL VHNLPQQLFG YSWYKGERVD GNRQIVGYAI50
51     GTQQATPGPA NSGRETIYPN ASLLIQNVTQ NDTGFYTLQV IKSDLVNEEA100
30    101     TGQFHVYP108 (SEQ ID NO:35).
```

The library was synthesized using Pepscan's Chemically Linked Peptides on Scaffolds (CLIPS) technology which allows to structure peptides into single loops, double-loops, triple loops, sheet-like folds, helix-like folds and combinations thereof (Timmerman *et al.* 2007, J. Mol. Recognit. **20**:283-99 and Slootstra et al. 1996, Molecular

Diversity 1: 87-96). CLIPS templates are coupled to cysteine residues. The side-chains of multiple cysteines in the peptides are coupled to one or two CLIPS templates. For example, a 0.5 mM solution of the T2 CLIPS template 1,3-bis (bromomethyl) benzene is dissolved in ammonium bicarbonate (20 mM, pH 7.9)/acetonitrile (1:1(v/v)). This solution is added onto the peptide arrays. The CLIPS template will bind to side-chains of two cysteines as present in the solid-phase bound peptides of the peptide-arrays (455 wells plate with 3 μ l wells). The peptide arrays are gently shaken in the solution for 30 to 60 minutes while completely covered in solution. Finally, the peptide arrays are washed extensively with excess of H₂O and sonicated in disrupt-buffer containing 1 percent SDS/0.1 percent beta-mercaptoethanol in PBS (pH 7.2) at 70°C for 30 minutes, followed by sonication in H₂O for another 45 minutes. The T3 CLIPS carrying peptides were made in a similar way but with three cysteines.

A total number of 3002 peptides was synthesized that includes: CLIPS matrix set combining two different areas of the entire protein wherein native cysteines are replaced by alanines; Linear peptide sets of length 6,10,15,20,25 and 33; Linear peptide sets of length 15, where the middle residue is replaced by alanine; CLIPS conformational loop set of length 15 wherein native cysteines are replaced by alanines; and CLIPS conformational loop set of length 15 with the middle residue replace by alanine wherein native cysteines are replaced by alanines.

The binding of antibody to each of the synthesized peptides was tested in a PEPSCAN-based ELISA. The peptide arrays were incubated with primary antibody (CM10) solution (overnight at 4°C). After washing, the peptide arrays were incubated with a 1/1000 dilution of an antibody peroxidase conjugate for one hour at 25°C. After washing, the peroxidase substrate 2,2'-azino-di-3-ethylbenzthiazoline sulfonate (ABTS) and 2 microliters/milliliter of 3 percent H₂O₂ were added. After one hour, the color development was measured. The color development was quantified with a charge coupled device (CCD) - camera and an image processing system.

The values obtained from the CCD camera range from 0 to 3000 mAU, similar to a standard 96-well plate ELISA-reader. The results are quantified and stored into the Peplab database. The binding values are extracted for analysis using graphical representation of data where the values taken by a variable in a two-dimensional map are represented as colors.

Based on the observed binding patterns, the core of the epitope of CM10.S91 consists of residues ₁₇VLLLVHNLPPQLF₂₉. A secondary binding region was observed on sequence around ₆₈YPNASLLIQNVT₇₉. When the observed binding patterns are compared to the

available crystal structure of the target protein, it is proposed that the discontinuous epitope for CM10.S91 consists of ₁₇VLLLVHNLPPQLF₂₉ (core) together with ₆₈PNASLLI₇₅ (support).

5 **Example 10: Additional experiments assessing the activity of CM10**

Granzyme B secretion by effector cells as a potential PD marker

The development of robust pharmacodynamic (PD) markers is critical for improving the success of drugs in clinical trials and assists in selection of an optimal drug dose to balance efficacy and toxicity. PD markers are often proximal in a molecular pathway to the drug target and are used to measure the effect of a drug regardless of therapeutic effect. Granzyme B is a serine protease that is released through cytoplasmic granules of cytotoxic T cells (CTLs) and NK cells upon recognition of specific targets. This in turn mediates the apoptosis of the target cells. We sought to determine if CM10 enhances granzyme B secretion from effector NK and T cells in the presence of specific target cells, thus enabling the observed enhanced killing of the target cells by the antibody.

TIL were incubated with CM10 (0.2µg/ml, 1µg/ml, 5µg/ml or 10 µg/ml) or an isotype matched control Ab (IPI-Ipilimumab 10 µg/ml) for 30 minutes at 37°C; Target positive melanoma cells expressing CEACAM1 and HLA-A2 - SKMEL5, were added for an overnight incubation at effector-to-target ratio of 2.5:1. Granzyme B secretion was measured by ELISA. Results represent the mean ± S.E of granzyme B release values from 6 repeats per treatment. Similar results were seen in another independent experiment.

A represented in Fig 15 TIL secretes low initial levels of granzyme B that remained low in the presence of CM10 alone. Only when the target cells were added to the TIL, granzyme B elevated dramatically. Yet, CM10 was able to significantly increase granzyme B secretion above the high initial levels. Similar results were seen when NK92MI cells were used as effector cells demonstrating that CM10 can enhance granzyme B secretion from NK cells in a similar manner. The use of Granzyme B as PD marker to CM10 treatment is also examined in complementary set of experiments including in vivo models and patients specimens.

30 CM10 blocks CEACAM1-CEACAM5 interactions

Carcinoembryonic antigen (CEA, CEACAM5, and CD66e) has been found to be associated with various types of cancers, particularly colorectal carcinoma, as well as gastrointestinal, pancreatic cancers, non-small cell lung cancer ,breast cancers , thyroid,

stomach and ovarian cancer. The molecule has been developed to be a target for cancer diagnosis and therapy.

Homophilic CEACAM1 interactions between a MHC-I-deficient melanoma cell line and NK cells significantly inhibited killing of the cancer cells by the NK cells (Markel et al J Immunol. 2002;168:2803–2810). Similarly, forced expression of CEACAM5 in HLA class I-defective 721.221 cells also suppressed NK-mediated cytotoxicity (Stern et al J Immunol. 2005;174:6692–6701), indicating CEACAM5-CEACAM1 heterophilic interactions are also able to trigger inhibitory CEACAM1 signaling in immune cells.

CEACAM1 positive cell line (SKMel 5) was incubated with an isotype control or CM10 in various concentrations, unbound CM10 was washed twice, followed by incubation with or without soluble biotinylated CEACAM5. The binding of the biotinylated CEACAM5 to the cells was detected by a secondary streptavidin-APC and FACS analysis. The results represent the mean of duplicate wells \pm SE. Similar results were obtained in two other independent experiments.

The results of Fig 16 demonstrate that CM10 inhibits the binding between CEACAM1 to CEACAM5 in a dose-dependent manner. Therefore CM10 can be used to treat malignancies that express high level of CEACAM5 and exploit the CEACAM1-CEACAM5 axis in order to suppress the immune cells.

CM10 enhances HLA restricted T cell killing

Cytotoxic T cells destroy virally infected cells and tumor cells, and are also implicated in transplant rejection. These cells recognize their targets by binding to antigens associated with HLA molecules, which are present on the surface of nearly every cell of the body. The killing of infected or malignant cells requires the recruitment of several pathways in order to generate proper response.

This set of assays was sought to determine if CM10 enhances the “natural” CTL activity which is HLA restricted or alternatively causes a general stimulation of the immune system.

TIL were incubated with CM10 (0.2 μ g/ml, 1 μ g/ml, 5 μ g/ml or 10 μ g/ml) or an isotype matched control antibody for 30 minutes at 37°C. Target melanoma cells SKMEL5, expressing CEACAM1 and HLA-A2 were pre-incubated with or without HLA-A2 blocking monoclonal antibody for 1 hour and were added for an overnight incubation at effector-to-target ratio of 2.5:1. Killing was determined by standard LDH release assay. Results represent the mean \pm S.E of LDH release values from 3 repeats per treatment.

The results in Fig 17 demonstrate that in the presence of HLA-A2 blocking antibodies CM10 was unable to augment melanoma cell killing by the TIL and that CM10's activity is HLA restricted. Similar result was seen in another independent experiment. It is therefore suggested that in accordance with these results CM10 will improve the activity of CTLs against malignant cells in vivo and will not facilitate non-specific activation of the immune system.

The immunomodulatory activity of CM10 inhibits tumor growth in-vivo

One of the most widely used models to assess the in vivo action of a novel anti cancer agents is the human tumor xenograft model. In this model, human tumor cells are transplanted under the skin into immunocompromised mice that do not reject human cells. Depending upon the number of cells injected and the cells' growth rate, the tumor develops over several weeks, and the response to appropriate therapeutic regimes can be studied in vivo. The advantage of a model that involves treatment in early stages of tumor formation is the opportunity to examine the agent's influence in a setting that can be more similar to metastasis spread in patients and to identify the effect of the treatment on initial tumor implantation.

The immunomodulatory activity of CM10 in vivo was therefore examined in SCID-NOD mice engrafted with human melanoma cells under prophylactic treatment regime. SCID-NOD mice were randomized according to the five treatment groups indicated in figure 18. 6×10^6 SKmel5 cells (melanoma cell line) were injected SC in Day 0. Mice were treated with antibodies/TIL as indicated in the figure. Arrows indicate time of administration; Thin arrows represent antibodies-0.25mg/mouse/injection (CM10, IgG control and Ipilimumab control) and wide arrows for TIL (20×10^6 cell per mouse per injection). The tumors' volume was blindly measured 2-3 times per week. The results represent average tumor volumes \pm SE from 8-9 mice per group.

Fig 18 shows that the combination of adoptive human T cell transfer with CM10 injections exhibited significant synergism and strongly inhibited xenograft growth compared to the isotype control group (69% TGI). The results demonstrate that CM10 also has an effect as a single agent (45% TGI at day 32) and probably worked directly on the tumor cells or the tumors' surroundings by inhibiting proliferation, angiogenesis or through other biological pathways.

In the same set of experiments the tumors' total weight was determined at the experiment termination day. The results obtained support the volume measurements, and

show a dramatic decrease in the tumor burden of the TIL+ CM10 treated group and also statistically significant reduction in the tumors' weight in the CM10 treated group.

CM10 enhances the immunomodulatory killing activity of NK cells on CEACAM1-positive pancreatic cancer cell lines

5 Preliminary screening showed that pancreatic cancer cell lines and patient specimens express high levels of CEACAM1 (94% from pancreatic cancer specimens were CEACAM1 positive). Therefore, an experiment was conducted to examine if CM10 can enhanced the activity of immune cells against pancreatic cancer cell lines. Killing assays were conducted by co-culture of NK92MI cells that were pre-treated with various concentrations of CM10
10 with different CEACAM1 positive target pancreatic cancer cell lines. NK 92MI were incubated with CM10 (0.2µg/ml, 1µg/ml or 10 µg/ml) or an isotype matched control Ab (10 µg/ml) for 30 minutes at 37°C; **A.** Target cells expressing CEACAM1 were added for additional 5 hours at effector-to-target ratio of 2.5:1 (COLO-357) or overnight at an effector-to-target ratio of 5:1 (BXPC3). Results represent an average of % cytotoxicity ±SE as
15 determined by classical LDH release assay from triplicate wells per treatment.

The results represented in figure 19 show that although the killing of the pancreatic cancer cells by the NK cells was relatively low, CM10 enhanced this activity by up to 2 folds from the baseline cytotoxicity (PBS or control Ab). Similar results were seen in additional two lines.

20 CM10 enhances granzyme B secretion of NK cells in the presence of CEACAM1-positive pancreatic cancer cell lines

The experiment was designed to determine whether similarly to the melanoma cells, CM10 can enhance the secretion of granzyme B from NK cells in the presence of pancreatic cancer cells. In a similar setting to the killing assay, NK cells were pre-treated with various
25 concentrations of CM10 and then co-cultured with four different target pancreatic cancer cell lines. The supernatants were analyzed for granzyme B content by a specific ELISA. NK 92MI were incubated with CM10 (0.2µg/ml, 1µg/ml or 10 µg/ml) or an isotype matched control antibody (10 µg/ml) for 30 minutes at 37°C. Target pancreatic cells expressing CEACAM1 were added for additional 5 hours at effector-to-target ratio of 2.5:1 **A.** (COLO-
30 357 and ASPC-1) and **B.** (SU8686) or E:T of 10:1 **C.** (BXPC3). Granzyme B secretion was measured by ELISA. Results represent the mean ±S.E of granzyme B release values from 2-6 repeats per treatment.

Figure 20 clearly demonstrates that CM10 increases granzyme B secretion from NK cells in the presence of pancreatic cancer cells.

Example 11: Humanized and Human Antibodies

A humanized antibody typically has a human framework grafted with non human CDRs. Thus, a humanized antibody has one or more amino acid sequence introduced into it
5 from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones et al., Nature, 321:522-525, 1986; Riechmann et al., Nature, 332:323-327, 1988; Verhoeyen et al., Science, 239:1534-1536, 1988), by substituting rodent CDRs or CDR
10 sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are
15 substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened
20 against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims et al., J. Immunol., 151:2296, 1993; Chothia et al., J. Mol. Biol., 196:901, 1987). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same
25 framework may be used for several different humanized antibodies (Carter et al., Proc. Natl. Acad. Sci. USA, 89:4285, 1992; Presta et al., J. Immunol., 151:2623, 1993).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the
30 parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin

sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

Alternatively, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge (e.g., Jakobovits et al., Proc. Natl. Acad. Sci. USA, 90:2551, 1993; Jakobovits et al., Nature, 362:255-258, 1993; Bruggermann et al., Year in Immuno., 7:33, 1993; and Duchosal et al. Nature 355:258, 1992. Human antibodies can also be derived from phage-display libraries (Hoogenboom et al., J. Mol. Biol., 227:381, 1991; Marks et al., J. Mol. Biol., 222:581-597, 1991; Vaughan et al. Nature Biotech 14:309, 1996).

Example 12. Antibody Fragments

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (e.g., Morimoto et al., Journal of Biochemical and Biophysical Methods 24:107-117, 1992 and Brennan et al., Science, 229:81, 1985). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from E. coli and chemically coupled to form F(ab')₂ fragments (Carter et al., Bio/Technology 10:163-167, 1992). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv).

Example 13. Potency of anti CEACAM antibody against viral infections

The following experiments are used to determine the potential of CM10 against viral infection. The experiments include different target cells, various virus and several in vivo and in vitro models.

5 Detection/diagnosis: Examination of CEACAM expression level in cell lines and primary cells infected with different virus types, by FACS analysis, RT PCR and Immunohistochemistry.

Prevention: pre-incubation of target cells with anti CEACAM antibody and determination of the viral load or viral replication post viral infection.

10 Treatment: after viral infection the infected cell are incubated with immune system cell and the killing ability of the effectors cells and the viral load is examined. In addition, the viral load and replication and overall survival are determined in vivo after virus infection and treatment with anti CEACAM antibodies.

15 The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without undue experimentation and without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of
20 equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation. The means, materials, and steps for carrying out various disclosed functions may take a variety of alternative forms without departing from the invention.

CLAIMS

1. A monoclonal antibody or an antibody fragment thereof which recognizes human CEACAM1 having a heavy-chain CDR1 comprising a sequence set forth in SEQ ID NO: 1, a heavy-chain CDR2 comprising a sequence set forth in SEQ ID NO: 2 a heavy-chain CDR3 comprising a sequence set forth in SEQ ID NO: 3, a light-chain CDR1 comprising a sequence set forth in SEQ ID NO: 4, a light-chain CDR2 comprising a sequence set forth in SEQ ID NO: 5 and a light-chain CDR3 comprising a sequence set forth in SEQ ID NO: 6, and analogs and derivatives thereof.
2. The monoclonal antibody or fragment thereof according to claim 1, comprising heavy chain CDR1 having the sequence set forth in SEQ ID NO: 7, heavy chain CDR2 having the sequence set forth in SEQ ID NO: 8 and heavy chain CDR3 having the sequence set forth in SEQ ID NO: 9.
3. The monoclonal antibody or fragment thereof according to claim 1, comprising heavy chain CDR1 having the sequence set forth in SEQ ID NO: 13, heavy chain CDR2 having the sequence set forth in SEQ ID NO: 14 and heavy chain CDR3 having the sequence set forth in SEQ ID NO: 15.
4. The monoclonal antibody or fragment thereof according to claim 1, comprising light chain CDR1 having the sequence set forth in SEQ ID NO: 10, light chain CDR2 having the sequence set forth in SEQ ID NO: 11 and light chain CDR3 having the sequence set forth in SEQ ID NO: 12.
5. The monoclonal antibody or fragment thereof according to claim 1, comprising light chain CDR1 having the sequence set forth in SEQ ID NO: 16, light chain CDR2 having the sequence set forth in SEQ ID NO: 17, and light chain CDR3 having the sequence set forth in SEQ ID NO: 18.
6. The monoclonal antibody or fragment thereof according to claim 1, having CDR sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17, and 18.
7. The monoclonal antibody or fragment thereof according to claim 1, having CDR sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12.
8. An analog or derivative of a monoclonal antibody or fragment thereof according to any one of claims 1-7, having at least 90% sequence identity with the antigen-binding portion of the said monoclonal antibody.
9. The monoclonal antibody or fragment thereof, according to claim 1, comprising a heavy chain variable domain sequence having a sequence set forth in SEQ ID NO: 26, or an

analog or derivative thereof having at least 97% sequence identity with said heavy chain sequence.

10. The monoclonal antibody or fragment thereof according to claim 1, comprising a light chain variable domain sequence having a sequence set forth in SEQ ID NO: 28, or an analog or derivative thereof having at least 97% sequence identity with said light chain sequence.
11. The monoclonal antibody or fragment thereof according to claim 1, comprising a heavy chain variable domain having a sequence set forth in SEQ ID NO: 26 and a light chain variable domain having a sequence set forth in SEQ ID NO: 28, or an analog or derivative thereof having at least 97% sequence identity with the antibody or fragment sequence.
12. The monoclonal antibody or fragment thereof according to claim 1, comprising:
 - i. a framework sequence selected from the group consisting of: mouse IgG2a, mouse IgG2b, mouse IgG3, human IgG1, human IgG2, human IgG3; and
 - ii. six CDRs having sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17, and 18; or six CDRs having sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12; and analogs and derivatives thereof having at least 97% sequence identity with said CDR sequences, wherein the monoclonal antibody or fragment binds with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least two CEACAM subtypes.
13. The monoclonal antibody or fragment thereof according to claim 12 capable of binding with an affinity of at least about 10^{-8}M to CEACAM1.
14. The monoclonal antibody or fragment thereof according to claim 13 capable of binding with an affinity of at least about $5 \times 10^{-7} \text{M}$ to at least one of CEACAM3 and CEACAM5.
15. The monoclonal antibody or fragment thereof, according to claim 12 wherein the monoclonal antibody or fragment is a chimeric monoclonal antibody.
16. The chimeric monoclonal antibody or fragment thereof according to claim 15, comprising human derived constant regions selected from the group consisting of: human IgG1, human IgG2, and human IgG3
17. The chimeric monoclonal antibody or fragment thereof according to claim 16, comprising the six CDRs having sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17, and 18; or the six CDRs having sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12; and analogs and derivatives thereof having at least 95% sequence identity with said CDR sequences, wherein the monoclonal antibody binds with an affinity of at least about 10^{-8}M to CEACAM1.

18. The chimeric monoclonal antibody or fragment thereof, according to claim 17, comprising a constant region subclass of human IgG1 subtype.
19. The chimeric monoclonal antibody or fragment thereof according to claim 17, comprising a heavy chain sequence set forth in SEQ ID NO: 30.
- 5 20. The chimeric monoclonal antibody or fragment thereof according to claim 17, comprising a light chain sequence set forth in SEQ ID NO: 31.
21. The chimeric monoclonal antibody or fragment thereof according to claim 17, comprising a heavy chain sequence set forth in SEQ ID NO: 30, and light chain sequence set forth in SEQ ID NO: 31.
- 10 22. A monoclonal antibody which recognizes CEACAM1 produced from DNA sequences of the heavy and light chains contained in a plasmid deposited on September 28, 2011 under ATCC Accession Number PTA-12130.
23. A monoclonal antibody which recognizes CEACAM1, or a fragment thereof comprising at least the antigen-binding portion, which is capable of binding the same epitope on the
15 CEACAM1 molecule to which a monoclonal antibody having the six CDR sequences set forth in SEQ ID NOs: 7, 8, 9, 10, 11 and 12, or the six CDR sequences set forth in SEQ ID NOs: 13, 14, 15, 16, 17 and 18, binds.
24. The monoclonal antibody or fragment thereof of claim 23 wherein the antibody is reactive with an epitope within residues 17-29 and 68-79 of human CEACAM1 having
20 the sequences VLLLVHNLPPQLF (SEQ ID NO:32) and YPNASLLIQNVT (SEQ ID NO:33) respectively.
25. The monoclonal antibody or fragment thereof according to claim 23, wherein the epitope on the CEACAM1 molecule to which the monoclonal antibody binds is an epitope comprising amino acid residues within the sequences VLLLVHNLPPQLF (SEQ ID NO:
25 32) and YPNASLLIQNVT (SEQ ID NO: 33).
26. The monoclonal antibody or fragment thereof according to claim 23, wherein the epitope on the CEACAM1 molecule to which the monoclonal antibody binds comprises at least four amino acids of the sequence VLLLVHNLPPQLF (SEQ ID NO: 32).
27. The monoclonal antibody or fragment thereof according to claim 23, wherein the epitope
30 on the CEACAM1 molecule to which the monoclonal antibody binds is na epitope within sequences VLLLVHNLPPQLF (SEQ ID NO: 32) and PNASLLI (SEQ ID NO: 34).
28. An isolated peptide sequence of 6-20 amino acids comprising at least three amino acids from the sequence VLLLVHNLPPQLF (SEQ ID NO: 32) and at least three amino acids

from the sequence YPNASLLIQNVT (SEQ ID NO: 33), or an analog or derivative having at least 85% homology with the parent sequence.

29. The isolated peptide of claim 28 comprising at least six amino acids from the sequence VLLLVHNLPPQLF (SEQ ID NO: 32).
- 5 30. The isolated peptide of claim 28 comprising the amino acids of the sequences VLLLVHNLPPQLF (SEQ ID NO: 32) and PNASLLI (SEQ ID NO: 34).
31. Use of an isolated peptide according to claim 28 for production of antibodies which recognizes CEACAM1.
32. An isolated polynucleotide encoding a monoclonal antibody or antibody fragment
10 according to any one of claims 1-27.
33. The isolated polynucleotide sequence according to claim 32, comprising a DNA sequences set forth in SEQ ID NO: 25 or SEQ ID NO: 27 or analogs thereof, having at least 90% sequence identity with said sequences.
34. A plasmid comprising at least one isolated polynucleotide sequence according to claim
15 32.
35. A plasmid according to claim 34 deposited on September 28, 2011 under ATCC Accession Number PTA-12130.
36. A pharmaceutical composition comprising a therapeutically effective amount of a
20 monoclonal antibody or fragment according to any one of claims 1-27; and a pharmaceutically acceptable carrier.
37. The pharmaceutical composition of claim 36 for treatment of a disease or disorder associated with CEACAM1, CEACAM3 or CEACAM5 expression, activation or function.
38. The pharmaceutical composition of claim 37 wherein the disease or disorder is a cell
25 proliferative disease or disorder.
39. The pharmaceutical composition of claim 38 wherein the cell proliferative disease or disorder is a cancer selected from the group consisting of: gastrointestinal, colorectal (CRC), pancreatic, non-small cell lung (NSCL), breast, thyroid, stomach, ovarian and uterine.
- 30 40. The pharmaceutical composition of claim 38 wherein the cell proliferative disease or disorder is a cancer selected from the group consisting of: melanoma, pancreatic cancer, lung cancer and myeloma.
41. A diagnostic composition comprising at least one monoclonal antibody or antibody fragment according to any one of claims 1-27; and an optional carrier or excipient.

42. A method of preventing, attenuating or treating a disease or disorder associated with expression, activation or function of CEACAM, comprising administering to a subject in need thereof a pharmaceutical composition comprising a therapeutically effective amount of a pharmaceutical composition according to claim 36.
- 5 43. The method of claim 42 wherein the disease or disorder is a cancer.
44. The method of claim 42 wherein the disease or disorder is a viral infection.
45. The method according to claim 42, wherein the isolated antibody or antibody fragment contained in the pharmaceutical composition is attached to a cytotoxic moiety and the disease is cancer.
- 10 46. The method of claim 46, further comprising administering to the subject CEACAM1-expressing lymphocytes.
47. The method of claim 46 wherein the lymphocytes comprise T cells, NK cells or Tumor Infiltrating Lymphocyte.
48. A method of immunomodulation, the method comprising contacting a CEACAM-expressing lymphocyte with the antibody or antibody fragment according to any one of
15 claims 1-27.
49. A method of inhibiting migration of a CEACAM expressing tumor cell, the method comprising contacting the CEACAM expressing tumor cell with the antibody or antibody fragment, according to any one of claims 1-27, thereby inhibiting migration of a
20 CEACAM expressing tumor cell.
50. A method of inhibiting CEACAM homotypic or heterotypic protein-protein interaction, the method comprising contacting a CEACAM1-expressing lymphocyte with the antibody or antibody fragment, according to any one of claims 1-27, thereby inhibiting CEACAM1 homotypic or heterotypic protein-protein interaction.
- 25 51. A method for increasing the duration or progression of response or survival of a subject having cancer, comprising administering to the subject effective amounts of a composition comprising a monoclonal antibody or antibody fragment, according to any one of claims 1-27, and an anti-neoplastic composition, wherein said anti-neoplastic composition comprises at least one chemotherapeutic agent, whereby the co-
30 administration of the antibody and the anti-neoplastic composition effectively increases the duration or progression of survival.
52. A method for diagnosing a cancer in a subject in need thereof, the method comprising contacting a biological sample derived from the subject with the diagnostic composition

of claim 41, wherein a complex formation beyond a predetermined threshold is indicative of the cancer in the subject.

53. A method for diagnosing a disease or disorder associated with CEACAM expression is provided comprising the steps of:
- 5 i. incubating a biological sample with a monoclonal antibody or antibody fragment according to any one of claims 1-27;
 - ii. detecting the bound CEACAM using a detectable probe;
 - iii. comparing the amount of (ii) to a standard curve obtained from reference samples containing known amounts of CEACAM;
 - 10 iv. calculating the amount of the CEACAM in the biological sample from the standard curve; and
 - v. comparing the amount of (iv) to a normal CEACAM amount.
54. Use of a monoclonal antibody or an antibody fragment according to any one of claims 1-27, for diagnosis, prevention or treatment of a cell proliferative or angiogenesis-related
- 15 disease or disorder or a viral infection.
55. Use of a monoclonal antibody or an antibody fragment according to any one of claims 1-27, for preparation of a medicament for treatment of a disorder or disease associated with expression or activation of CEACAM.
56. Use of a monoclonal antibody or an antibody fragment according to any one of claims 1-
- 20 27, for preparation of a diagnostic composition for the diagnosis of a cell proliferative or angiogenesis-related disease or disorder or a viral infection.

Figure 1




Figure 2




Figure 3






Figure 4



5/22

Figure 5



6/22

Figure 6





Figure 7







Figure 8

Figure 9



9/22




Figure 10




Figure 11

11/22




Figure 12

12/22




Figure 13





Figure 13 continued

Figure 13 continued

HEK 293T Transient expression







Figure 14

Figure 15








Figure 16

Figure 17



19/22

Figure 18







Figure 19

Figure 20






Figure 20 continued

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2012/050402

A. CLASSIFICATION OF SUBJECT MATTER IPC (2013.01) C07K 16/28, C07K 16/30, A61K 39/395, G01N 33/53, A61P 35/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC (2013.01) C07K, A61K, G01N Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Databases consulted: NCBI, THOMSON INNOVATION, REGISTRY, Google Scholar Search terms used: CEACAM1, BGP, CD66a, C-CAM 1, CEACAM5, CD66e		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Pavoni E, et al., Selection, affinity maturation, and characterization of a human scFv antibody against CEA protein (2006). BMC Cancer, 6:41. Retrieved from the Internet <URL: http://www.biomedcentral.com/1471-2407/6/41 > <doi:10.1186/1471-2407-6-41>. 24 Feb 2006 (2006/02/24) The whole document	1-14,23-34,36-56
Y		15-22,35
X	Zheng C, et al., A novel anti-CEACAM5 monoclonal antibody, CC4, suppresses colorectal tumor growth and enhances NK cells-mediated tumor immunity (2011). PLoS One. 6(6): e21146. <URL: http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0021146 > < doi: 10.1371/journal.pone.0021146>. 22 Jun 2011 (2011/06/22) The whole document	1-14,23-34,36-56
Y		15-22,35
X	WO 0113937 A1 SKUBITZ KEITH M [US]; SKUBITZ AMY P N [US] 01 Mar 2001 (2001/03/01) SEQ ID NOs: 17, 119, 178, 187- Tables I, II, VII, VIII, page 3, paragraphs 1-2	28,29
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: “A” document defining the general state of the art which is not considered to be of particular relevance “E” earlier application or patent but published on or after the international filing date “L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) “O” document referring to an oral disclosure, use, exhibition or other means “P” document published prior to the international filing date but later than the priority date claimed “T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention “X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone “Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art “&” document member of the same patent family		
Date of the actual completion of the international search 20 Jan 2013		Date of mailing of the international search report 20 Jan 2013
Name and mailing address of the ISA: Israel Patent Office The Technology Park, Bldg.5, Malcha, Jerusalem, 96951, Israel Facsimile No. 972-2-5651616		Authorized officer RON-COHEN Yael yaelrc@justice.gov.il Telephone No. 972-2-5651737

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2012/050402

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Presta LG, et al., Humanization of an anti-vascular endothelial growth factor monoclonal antibody for the therapy of solid tumors and other disorders (1997). Cancer Res.57(20):4593-9. <URL: http://cancerres.aacrjournals.org/content/57/20/4593.long > 15 Oct 1997 (1997/10/15) abstract, p 4594, right column second paragraph, page 4595 right column third paragraph	15-22,35
Y	Kim SJ, et al., Antibody engineering for the development of therapeutic antibodies (2005). Mol Cells. 20(1):17-29..<URL: http://cms.daegu.ac.kr/sgpark/molecular%20biology/antibody%20engineering.pdf > 31 Aug 2005 (2005/08/31) page 19 left column, paragraph 1	15-22,35

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/IL2012/050402

Patent document cited search report	Publication date	Patent family member(s)	Publication Date
WO 0113937 A1	01 Mar 2001	AU 6940700 A	19 Mar 2001
		CA 2383562 A1	01 Mar 2001
		EP 1212075 A1	12 Jun 2002
		EP 1212075 A4	23 Nov 2005
		WO 0113937 A1	01 Mar 2001
		WO 0113937 A9	06 Sep 2002