



(22) Date de dépôt/Filing Date: 2004/03/05

(41) Mise à la disp. pub./Open to Public Insp.: 2004/09/16

(62) Demande originale/Original Application: 2 517 899

(30) Priorité/Priority: 2003/03/06 (US60/453,134)


(51) Cl.Int./Int.Cl. *C12N 15/31* (2006.01),
A61K 38/16 (2006.01), *A61K 39/102* (2006.01),
A61K 39/40 (2006.01), *A61P 27/16* (2006.01),
A61P 31/04 (2006.01), *A61P 37/04* (2006.01),
C07K 14/285 (2006.01), *C07K 16/12* (2006.01),
C12Q 1/68 (2006.01), *G01N 33/569* (2006.01),
A61K 39/395 (2006.01), *A61K 31/7088* (2006.01)

(71) Demandeurs/Applicants:
CHILDREN'S HOSPITAL, INC., US;
THE BOARD OF REGENTS OF THE UNIVERSITY OF
OKLAHOMA, US

(72) Inventeurs/Inventors:
BAKALETZ, LAUREN O., US; ...

(54) Titre : GENES D'UN ISOLAT DE L'OTITE MOYENNE DE HAEMOPHILUS INFLUENZAE NON TYPABLE

(54) Title: GENES OF AN OTITIS MEDIA ISOLATE OF NONTYPEABLE HAEMOPHILUS INFLUENZAE



(57) Abrégé/Abstract:

The invention relates to the polynucleotide sequence of a nontypeable strain of Haemophilus influenzae (NTHi) and polypeptides encoded by the polynucleotides and uses thereof. The invention also relates to NTHi genes which are upregulated during or in response to NTHi infection of the middle ear and/or the nasopharynx.



(21) **2 796 381**

(13) **A1**

(72) Inventeurs(suite)/Inventors(continued): MUNSON, ROBERT S., JR., US; DYER, DAVID W., US

(74) Agent: SMART & BIGGAR

64267-1392D

Abstract

The invention relates to the polynucleotide sequence of a nontypeable strain of Haemophilus influenzae (NTHi) and polypeptides encoded by the polynucleotides and uses thereof. The invention also relates to NTHi genes which are upregulated during or in response
5 to NTHi infection of the middle ear and/or the nasopharynx.

64267-1392D

1

**GENES OF AN OTITIS MEDIA ISOLATE OF NONTYPEABLE
HAEMOPHILUS INFLUENZAE**

This is a division of Canadian Patent Application Serial No. 2,517,899 filed on March 5, 2004.

5 It should be understood that the expression "the present invention" or the like used in this specification encompasses not only the subject matter of this divisional application, but also that of the parent application.

Related Applications

The present application claims priority benefit from U.S. Provisional
10 Application 60/453,134 filed March 6, 2003.

Field of Invention

The invention relates to the polynucleotide sequence of a nontypeable strain of *Haemophilus influenzae* (NTHi) genome, NTHi genes contained within the genome and polypeptides encoded by the polynucleotides. The invention also relates
15 to uses of these NTHi polynucleotides and NTHi polypeptides including vaccines and methods of treating and preventing NTHi related disorders. The invention also relates to NTHi genes which are upregulated during or in response to NTHi infection of the middle ear or nasopharynx.

Background

20 Otitis media (OM) is a highly prevalent pediatric disease worldwide and is the primary cause for emergency room visits by children (Infante-Rivand and Fernandez, *Epidemiol. Rev.*, 15: 444-465, 1993). Recent statistic indicate that 24.5 million physician office visits were made for OM in 1990, representing a greater than 200% increase over those reported in the 1980's. While rarely associated with
25 mortality any longer, the morbidity associated with OM is significant. Hearing loss is a common problem associated with this disease, often times affecting a child's behavior, education and development of language skills (Baldwin, *Am. J. Otol.*, 14: 601-604, 1993; Hunter *et al.*, *Ann. Otol. Rhinol. Laryngol. Suppl.*, 163: 59-61, 1994; Teele *et al.*, *J. Infect. Dis.*, 162: 685-694, 1990). The socioeconomic impact of OM
30 is also great, with direct and indirect costs of diagnosing and managing OM exceeding \$5 billion annually in the U.S. alone (Kaplan *et al.*, *Pediatr. Infect. Dis. J.*, 16: S9-11, 1997).

64267-1392D

1a

Whereas antibiotic therapy is common and the surgical placement of tympanostomy tubes has been successful in terms of draining effusions, clearing infection and relieving pain associated with the accumulation of fluids in the middle ear, the emergence of multiple antibiotic-resistant bacteria and the invasive nature
5 associated with tube placement, has illuminated the need for more effective and

accepted approaches to the management and preferably, the prevention of OM.

Surgical management of chronic OM involves the insertion of tympanostomy tubes through the tympanic membrane while a child is under general anesthesia. While this procedure is commonplace (prevalence rates are ~ 13%; Bright *et al.*, *Am. J. Public Health*, 83(7): 1026-8, 1993) and is highly effective in terms of relieving painful symptoms by draining the middle ear of accumulated fluids, it too has met with criticism due to the invasive nature of the procedure and its incumbent risks (Berman *et al.*, *Pediatrics*, 93(3):353-63, 1994; Bright *et al.*, *supra.*; Cimon, *ASM News*, 60: 527-528; Paap, *Ann. Pharmacother.*, 30(11): 1291-7, 1996).

Progress in vaccine development is most advanced for *Streptococcus pneumoniae*, the primary causative agent of acute OM (AOM), as evidenced by the recent approval and release of a seven-valent capsular-conjugate vaccine, PREVNAR[®] (Eskola and Kilpi, *Pediatr. Infect. Dis. J.* 16: S72-78, 2000). While PREVNAR[®] has been highly efficacious for invasive pneumococcal disease, coverage for OM has been disappointing (6-8%) with reports of an increased number of OM cases due to serotypes not included in the vaccine (Black *et al.*, *Pediatr. Infect. Dis. J.*, 19: 187-195; Eskola *et al.*, *Pediatr. Infect. Dis. J.*, 19: S72-78, 2000; Eskola *et al.*, *N. Engl. J. Med.* 344: 403-409, 2001; Snow *et al.*, *Otol. Neurotol.*, 23: 1-2, 2002). Less progress has been made for non-typeable *Haemophilus influenzae* (NTHi), the gram-negative pathogen that predominates in chronic OM with effusion (Klein, *Pediatr. Infect. Dis. J.*, 16: S5-8, 1997; Spinola *et al.*, *J. Infect. Dis.*, 154: 100-109, 1986). Hampering development of effective vaccines against NTHi, is the currently incomplete understanding of the pathogenesis of NTHi-induced middle ear disease. Contributing to this delay is a lack of understanding of the dynamic interplay between microbe-expressed virulence factors and the host's immune response as the disease progresses from one of host immunological tolerance of a benign nasopharyngeal commensal, to that of an active defensive reaction to an opportunistic invader of the normally sterile middle ear space.

Currently there is a poor understanding of how NTHi causes OM in children. The identification of putative virulence factors necessary for induction of OM will contribute significantly to the understanding of the host-pathogen interaction and ultimately, the identification of potential vaccine candidates and targets of

chemotherapy. There is a tremendous need to develop more effective and accepted approaches to the management and preferably, the prevention of otitis media.

Vaccine development is a very promising and cost effective method to accomplish this goal (Giebank, *Pediatr. Infect. Dis. J.*, 13(11): 1064-8, 1994; Karma *et al.*, *Int. J. Pediatr. Otorhinolaryngol.*, 32(Suppl.): S127-34, 1995).

Summary of Invention

The present invention provides for the identification and characterization of the genomic sequence of NTHi *H. influenzae* strain 86-028NP and the polypeptide sequences encoded thereby. The 3-fold analysis of the NTHi genomic sequence is set out in a series of contig sequences denoted as SEQ ID NO: 1-576, and the subsequent 8-fold analysis of the genomic sequence is set out in a series of 11 contig sequences denoted as SEQ ID NOS: 675-685. These contigs are raw data and one of skill in the art may assemble these contigs by comparing overlapping sequences to construct the complete genome of the NTHi stain 86-028NP using routine methods.

The present invention also provides for antibodies specific for the NTHi polypeptides of the invention. Methods of detecting NTHi bacteria in a human or in sample, such as serum, sputum, ear fluid, blood, urine, lymphatic fluid and cerebrospinal fluid are contemplated. These methods include detecting NTHi polynucleotides with specific polynucleotide probes or detecting NTHi polypeptides with specific antibodies. The invention also contemplates diagnostic kits which utilize these methods of detecting NTHi bacteria.

The present invention also contemplates methods of eliciting an immune response by administering a NTHi polypeptide of the invention or a NTHi peptide thereof. These methods include administering the NTHi polypeptide or NTHi peptide as a vaccine for treatment and/or prevention of diseases caused by NTHi infection, such as OM. The following NTHi genes are upregulated during or in response to middle ear and/or nasopharynx infections; and the polypeptides encoded by these genes and peptides thereof are contemplated as possible OM vaccine candidates and/or target of chemotherapy: *hisB*, *lppB*, *sapA*, *lolA*, *rbsC*, *purE*, *ribB*, *arcB*, *uxuA*, *dsbB*, *ureH*, *licC*, HI1647, *ispZ*, *radC*, *mukF*, *glpR*, *ihfB*, *argR*, *cspD*, HI0094, HI1163, HI1063, HI0665, HI1292, HI1064. NTHi *hisB* gene is set out as

nucleotide sequence SEQ ID NO: 615 and encodes the amino acid sequence set out as SEQ ID NO: 616. NTHi *sapA* gene is set out as nucleotide sequence SEQ ID NO: 617 and encodes the amino acid sequence set out as SEQ ID NO: 618. NTHi *rhsC* gene is set out as nucleotide sequence SEQ ID NO: 619 and encodes the amino acid sequence set out as SEQ ID NO: 620. NTHi *purE* gene is set out as nucleotide sequence SEQ ID NO: 621 and encodes the amino acid sequence set out as SEQ ID NO: 622. NTHi *ribB* gene is set out as nucleotide sequence SEQ ID NO: 623 and encodes the amino acid sequence set out as SEQ ID NO: 624. NTHi *arcB* gene is set out as nucleotide sequence SEQ ID NO: 625 and encodes the amino acid sequence set out as SEQ ID NO: 626. NTHi *uxuA* gene is set out as nucleotide sequence SEQ ID NO: 627 and encodes the amino acid sequence set out as SEQ ID NO: 628. NTHi *dsbB* gene is set out as nucleotide sequence SEQ ID NO: 629 and encodes the amino acid sequence set out as SEQ ID NO: 630. NTHi *ureH* gene is set out as nucleotide sequence SEQ ID NO: 631 and encodes the amino acid sequence set out as SEQ ID NO: 632. NTHi *licC* gene is set out as nucleotide sequence SEQ ID NO: 633 and encodes the amino acid sequence set out as SEQ ID NO: 634. NTHi HI1647 gene is set out as nucleotide sequence SEQ ID NO: 635 and encodes the amino acid sequence set out as SEQ ID NO: 636. NTHi *ispZ* gene is set out as nucleotide sequence SEQ ID NO: 637 and encodes the amino acid sequence set out as SEQ ID NO: 638. NTHi *radC* gene is set out as nucleotide sequence SEQ ID NO: 639 and encodes the amino acid sequence set out as SEQ ID NO: 640. NTHi *mukF* gene is set out as nucleotide sequence SEQ ID NO: 641 and encodes the amino acid sequence set out as SEQ ID NO: 642. NTHi *glpR* gene is set out as nucleotide sequence SEQ ID NO: 643 and encodes the amino acid sequence set out as SEQ ID NO: 644. NTHi *ihfB* gene is set out as nucleotide sequence SEQ ID NO: 645 and encodes the amino acid sequence set out as SEQ ID NO: 646. NTHi *argR* gene is set out as nucleotide sequence SEQ ID NO: 647 and encodes the amino acid sequence set out as SEQ ID NO: 648. NTHi *cspD* gene is set out as nucleotide sequence SEQ ID NO: 649 and encodes the amino acid sequence set out as SEQ ID NO: 650. NTHi HI1163 gene is set out as nucleotide sequence SEQ ID NO: 651 and encodes the amino acid sequence set out as SEQ ID NO: 652. NTHi HI1063 gene is set out as nucleotide sequence SEQ ID NO: 653 and encodes the amino acid sequence set out as SEQ ID NO: 654. NTHi HI0665 gene is set out as nucleotide sequence SEQ ID NO: 655 and encodes the amino acid sequence

set out as SEQ ID NO: 656. NTHi HI1292 gene is set out as nucleotide sequence SEQ ID NO: 657 and encodes the amino acid sequence set out as SEQ ID NO: 658.

The novel NTHi genes included in the polynucleotide sequences presented as SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Tables 4 and 4B are also up-regulated during infection of the middle ear and/or the nasopharynx, and therefore are contemplated to encode OM vaccine candidates and/or targets of chemotherapy. In addition, the following NTHi genes are contemplated to be virulence-associated genes and therefore are contemplated to encode possible OM vaccine candidates and/or targets of chemotherapy: HI1386, HI1462, HI1369, *lav*, HI1598. NTHi HI1386 gene sequence is set out as SEQ ID NO: 659 and encodes the amino acid sequence set out as SEQ ID NO: 660. NTHi HI1462 gene sequence is set out as SEQ ID NO: 661 and encodes the amino acid sequence set out as SEQ ID NO: 662. NTHi HI1369 gene sequence is set out as SEQ ID NO: 665 and encodes the amino acid sequence set out as SEQ ID NO: 666. NTHi *lav* gene sequence is set out as SEQ ID NO: 663 and encodes the amino acid sequence set out as SEQ ID NO: 664. NTHi HI1598 gene sequence is set out as SEQ ID NO: 669 and SEQ ID NO: 671 and encodes the amino acid sequence set out as SEQ ID NO: 670 and SEQ ID NO: 672. Additional NTHi genes associated with virulence include the polynucleotide sequences presented as SEQ ID NO: 667 and SEQ ID NO: 673.

As a method of treating or preventing NTHi infection, the present invention contemplates administering a molecule that inhibits expression or the activity of the NTHi polypeptides, which are upregulated or active during infection. In particular, the invention contemplates methods of treating or preventing NTHi infection comprising modulating NTHi protein expression by administering an antisense oligonucleotide that specifically binds to NTHi genes that are upregulated during NTHi infections, such genes include *hisB*, *lppB*, *sapA*, *lola*, *rbsC*, *purE*, *ribB*, *arcB*, *uxuA*, *dsbB*, *ureH*, *licC*, HI1647, *ispZ*, *radC*, *mukF*, *glpR*, *ihfB*, *argR*, *cspD*, HI0094, HI1163, HI1063, HI0665, HI1292, HI1064. The invention also contemplates methods of treating or preventing NTHi infection comprising administering antibodies or small molecules that modulate the activity of the proteins encoded by these genes. The novel NTHi genes included in the polynucleotide sequences presented as SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide

sequences set out in Tables 4 and 4B are also up-regulated during infection of the middle ear and/or the nasopharynx and therefore antisense oligonucleotides that specifically bind these polynucleotide sequences are also contemplated.

Polynucleotides and Polypeptides of the Invention

5 The present invention provides for the sequences of the NTHi strain 86-028NP genome. This genomic sequence is presented as a series of contig sequences denoted herein as "contigs 1-576". Each contig is assigned a sequence identification number that correlates with its "contig number". Therefore, the contigs of the present invention as set out as SEQ ID NOS: 1-576. These contig
10 polynucleotide sequences may be assembled into the complete genome sequence of the NTHi strain 86-028NP using routine methods. Upon completion of 8-fold sequence analysis of the NTHi strain 82-028NP genome, the genomic sequence was assembled into 11 contigs which are denoted herein as SEQ ID NOS: 675-685.

 The present invention provides for the NTHi polynucleotide sequences
15 and open reading frames contained within the contigs of SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5. The present invention also provides for the polypeptide sequences encoded by the NTHi polynucleotides of the present invention such as the amino acid sequences set out in Table 3B, Table 4B and Table 5. The invention provides for
20 polynucleotides that hybridize under stringent conditions to (a) the complement of the nucleotides sequence of SEQ ID NOS: 1-576; SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5 herein (b) a polynucleotide which is an allelic variant of any polynucleotides recited above; (c) a polynucleotide which encodes a species homolog of any of the proteins recited above;
25 or (d) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the NTHi polypeptides of the present invention.

 The NTHi polynucleotides of the invention also include nucleotide sequences that are substantially equivalent to the polynucleotides recited above. Polynucleotides according to the invention can have, *e.g.*, at least 65%, at least 70%,
30 at least 75%, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more typically at least 90%, 91%, 92%, 93%, or 94% and even more typically at least 95%, 96%, 97%, 98% or 99% sequence identity to the NTHi polynucleotides recited above.

Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to the NTHi nucleotide sequences of SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5
5 herein, or compliments thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, *e.g.*, 15, 17, or 20 nucleotides or more that are selective for (*i.e.*, specifically hybridize to any one of the polynucleotides of the invention) are contemplated. Probes capable of specifically
10 hybridizing to a polynucleotide can differentiate NTHi polynucleotide sequences of the invention from other polynucleotide sequences in the same family of genes or can differentiate NTHi genes from other bacterial genes, and are preferably based on unique nucleotide sequences.

The term "stringent" is used to refer to conditions that are commonly
15 understood in the art as stringent. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of stringent conditions for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride, 0.0015M sodium citrate, and 50% formamide at 42°C. See Sambrook *et al.*,
20 *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, (Cold Spring Harbor, N.Y. 1989). More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used, however, the rate of hybridization will be affected. In instances wherein hybridization of deoxyoligonucleotides is concerned, additional exemplary stringent
25 hybridization conditions include washing in 6x SSC 0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos).

Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples
30 are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO₄, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or other non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration

and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4, however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson *et al.*, *Nucleic Acid*
5 *Hybridisation: A Practical Approach*, Ch. 4, IRL Press Limited (Oxford, England). Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of different sequence relatedness to form hybrids.

The sequences falling within the scope of the present invention are not
10 limited to these specific sequences, but also include allelic and species variations thereof. Allelic and species variations can be routinely determined by comparing the sequence provided in SEQ ID NOS: 1-576, SEQ ID NOS: 675-685, and nucleotide sequences out in Table 3B, Table 4B and Table 5 herein, preferably the open reading frames therein, a representative fragment thereof, or a nucleotide sequence at least
15 90% identical, preferably 95% identical, to the open reading frames within SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5 with a sequence from another isolate of the same species. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including
20 GAP (Devereux *et al.*, *Nucl. Acid. Res.*, 12: 387, 1984; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Altschul *et al.*, *J. Mol. Biol.*, 215: 403-410, 1990). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (*BLAST Manual*, Altschul *et al.* NCB/NLM/NIH Bethesda, MD 20894; Altschul *et*
25 *al.*, *supra*). The well known Smith Waterman algorithm may also be used to determine identity.

Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific open reading frames (ORF) disclosed herein. In other words, in the coding region of
30 an ORF, substitution of one codon for another codon that encodes the same amino acid is expressly contemplated.

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising: the amino acid sequences encoded by the nucleotide

sequences included within the polynucleotide sequences set out as SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5, or the corresponding full length or mature protein. The polypeptides of the invention include the amino acid sequences of SEQ ID NO: 616, SEQ ID NO: 618, SEQ ID NO: 620, SEQ ID NO: 622, SEQ ID NO: 624, SEQ ID NO: 626, SEQ ID NO: 628, SEQ ID NO: 628, SEQ ID NO: 630, SEQ ID NO: 632, SEQ ID NO: 634, SEQ ID NO: 636, SEQ ID NO: 638, SEQ ID NO: 640, SEQ ID NO: 642, SEQ ID NO: 644, SEQ ID NO: 646, SEQ ID NO: 648, SEQ ID NO: 650, SEQ ID NO: 652, SEQ ID NO: 654, SEQ ID NO: 656, SEQ ID NO: 658, SEQ ID NO: 660, SEQ ID NO: 662, SEQ ID NO: 664, SEQ ID NO: 666, SEQ ID NO: 668, SEQ ID NO: 670, SEQ ID NO: 672, SEQ ID NO: 674, SEQ ID NO: 687, SEQ ID NO: 689, SEQ ID NO: 691, SEQ ID NO: 693, SEQ ID NO: 695, SEQ ID NO: 697, SEQ ID NO: 699, SEQ ID NO: 701, SEQ ID NO: 703, SEQ ID NO: 705, SEQ ID NO: 707, SEQ ID NO: 709, SEQ ID NO: 711, SEQ ID NO: 713, SEQ ID NO: 715, SEQ ID NO: 717, SEQ ID NO: 719, SEQ ID NO: 721, SEQ ID NO: 723, SEQ ID NO: 725, SEQ ID NO: 727, SEQ ID NO: 729, SEQ ID NO: 731, SEQ ID NO: 733, SEQ ID NO: 735, SEQ ID NO: 737, SEQ ID NO: 739, SEQ ID NO: 741, SEQ ID NO: 743, SEQ ID NO: 745, SEQ ID NO: 747, SEQ ID NO: 749, SEQ ID NO: 751, SEQ ID NO: 753, SEQ ID NO: 755, SEQ ID NO: 757, SEQ ID NO: 759, SEQ ID NO: 761, 763, SEQ ID NO: 765, SEQ ID NO: 767, SEQ ID NO: 769 or SEQ ID NO: 771 which are set out in Table 3B, Table 4B and Table 5 herein.

Polypeptides of the invention also include polypeptides preferably with biological or immunogenic activity that are encoded by: (a) an open reading frame contained within the nucleotide sequences set forth as SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5, or (b) polynucleotides that hybridize to the complement of the polynucleotides of (a) under stringent hybridization conditions.

The invention also provides biologically active or immunologically active variants of the amino acid sequences of the present invention; and "substantial equivalents" thereof (*e.g.*, with at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, 86%, 87%, 88%, 89%, at least about 90%, 91%, 92%, 93%, 94%, typically at least about 95%, 96%, 97%, more typically at least about 98%, or most typically at least about 99% amino acid identity) that

retain biological and/or immunogenic activity. Polypeptides encoded by allelic variants may have a similar, increased, or decreased activity compared to polypeptides encoded by the polynucleotides included within the nucleotide sequences presented in SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the
5 nucleotide sequences set out in Table 3B, Table 4B and Table 5 herein, and the polypeptides having an amino acid sequence set out in Table 3B, Table 4B and Table 5 herein

NTHi peptides refer to fragments of the NTHi polypeptides encoded by the nucleotide sequences presented in SEQ ID NOS: 1-576, SEQ ID NOS: 675-
10 685 or the nucleotide sequences set out in Table 3B, Table 4B and Table 5 herein, and the polypeptides having the amino acid sequences set out in Table 3B, Table 4B and Table 5 herein. The preferred NTHi peptides are biologically and/or immuniologically active.

The present invention further provides isolated NTHi polypeptides or
15 NTHi peptides encoded by the NTHi nucleic acid fragments of the present invention or by degenerate variants of the nucleic acid fragments of the present invention. The term "degenerate variant" refers to nucleotide fragments which differ from a nucleic acid fragment of the present invention (*e.g.*, an ORF) by nucleotide sequence but, due to the degeneracy of the genetic code, encode an identical NTHi polypeptide
20 sequence. Preferred nucleic acid fragments of the present invention are the ORFs that encode proteins.

The invention also provides for NTHi polypeptides with one or more conservative amino acid substitutions that do not affect the biological and/or immunogenic activity of the polypeptide. Alternatively, the NTHi polypeptides of the
25 invention are contemplated to have conservative amino acids substitutions which may or may not alter biological activity. The term "conservative amino acid substitution" refers to a substitution of a native amino acid residue with a nonnative residue, including naturally occurring and nonnaturally occurring amino acids, such that there is little or no effect on the polarity or charge of the amino acid residue at that position.
30 For example, a conservative substitution results from the replacement of a non-polar residue in a polypeptide with any other non-polar residue. Further, any native residue in the polypeptide may also be substituted with alanine, according to the methods of "alanine scanning mutagenesis". Naturally occurring amino acids are characterized

based on their side chains as follows: basic: arginine, lysine, histidine; acidic: glutamic acid, aspartic acid; uncharged polar: glutamine, asparagine, serine, threonine, tyrosine; and non-polar: phenylalanine, tryptophan, cysteine, glycine, alanine, valine, proline, methionine, leucine, norleucine, isoleucine. General rules for amino acid substitutions are set forth in Table 1 below.

Table 1
Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asn
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe,	Leu
Leu	Norleucine, Ile, Val, Met,	Leu
Lys	Arg, 1,4 Diaminobutyric	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Arg
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala,	Leu

Antisense polynucleotides complementary to the polynucleotides encoding the NTHi polypeptides are also provided.

The invention contemplates that polynucleotides of the invention may be inserted in a vector for amplification or expression. For expression, the polynucleotides are operatively linked to appropriate expression control sequence

such as a promoter and polyadenylation signal sequences. Further provided are cells comprising polynucleotides of the invention. Exemplary prokaryotic hosts include bacteria such as *E. coli*, *Bacillus*, *Streptomyces*, *Pseudomonas*, *Salmonella* and *Serratia*.

5 The term "isolated" refers to a substance removed from, and essentially free of, the other components of the environment in which it naturally exists. For example, a polypeptide is separated from other cellular proteins or a DNA is separated from other DNA flanking it in a genome in which it naturally occurs.

Antibodies and Methods for Eliciting an Immune Response

10 The invention provides antibodies which bind to antigenic epitopes unique to (*i.e.*, are specific for) NTHi polypeptides. Also provided are antibodies which bind to antigenic epitopes common among multiple *H. influenzae* subtypes but unique with respect to any other antigenic epitopes. The antibodies may be polyclonal antibodies, monoclonal antibodies, antibody fragments which retain their
15 ability to bind their unique epitope (*e.g.*, Fv, Fab and F(ab)2 fragments), single chain antibodies and human or humanized antibodies. Antibodies may be generated by techniques standard in the art.

 It is known in the art that antibodies to the capsular polysaccharide of *H. influenzae* exhibit the ability to kill bacteria *in vitro* assays. These antibodies are
20 also known to protect against challenge with *H. influenzae* in animal model systems. These studies indicate antibody to the capsular polysaccharides are likely to elicit a protective immune response in humans. The present invention provides for antibodies specific for the NTHi polypeptides of the present invention and fragments thereof, which exhibit the ability to kill both *H. influenzae* bacteria and to protect humans
25 from NTHi infection. The present invention also provides for antibodies specific for the NTHi polypeptides of the invention which reduce the virulence, inhibit adherence, inhibit cell division, and/or inhibit penetration into the epithelium of *H. influenzae* bacteria or enhance phagocytosis of the *H. influenzae* bacteria.

In vitro complement mediated bactericidal assay systems (Musher *et al.*, *Infect. Immun.* 39: 297-304, 1983; Anderson *et al.*, *J. Clin. Invest.* 51: 31-38, 1972) may be used to measure the bactericidal activity of anti-NTHi antibodies.
30 Further data on the ability of NTHi polypeptides and NTHi peptides to elicit a

protective antibody response may be generated by using animal models of infection such as the chinchilla model system described herein.

It is also possible to confer short-term protection to a host by passive immunotherapy via the administration of pre-formed antibody against an epitope of NTHi, such as antibodies against NTHi OMP, LOS or noncapsular proteins. Thus, the contemplated vaccine formulations can be used to produce antibodies for use in passive immunotherapy. Human immunoglobulin is preferred in human medicine because a heterologous immunoglobulin may provoke an immune response to its foreign immunogenic components. Such passive immunization could be used on an emergency basis for immediate protection of unimmunized individuals exposed to special risks. Alternatively, these antibodies can be used in the production of anti-idiotypic antibody, which in turn can be used as an antigen to stimulate an immune response against NTHi epitopes.

The invention contemplates methods of eliciting an immune response to NTHi in an individual. These methods include immune responses which kill the NTHi bacteria and immune responses which block *H. influenzae* attachment to cells. In one embodiment, the methods comprise a step of administering an immunogenic dose of a composition comprising a NTHi protein or NTHi peptide of the invention. In another embodiment, the methods comprise administering an immunogenic dose of a composition comprising a cell expressing a NTHi protein or NTHi peptide of the invention. In yet another embodiment, the methods comprise administering an immunogenic dose of a composition comprising a polynucleotide encoding a NTHi protein or NTHi peptide of the invention. The polynucleotide may be a naked polynucleotide not associated with any other nucleic acid or may be in a vector such as a plasmid or viral vector (*e.g.*, adeno-associated virus vector or adenovirus vector). Administration of the compositions may be by routes standard in the art, for example, parenteral, intravenous, oral, buccal, nasal, pulmonary, rectal, or vaginal. The methods may be used in combination in a single individual. The methods may be used prior or subsequent to NTHi infection of an individual.

An "immunological dose" is a dose which is adequate to produce antibody and/or T cell immune response to protect said individual from NTHi infection, particularly NTHi infection of the middle ear and/or the nasopharynx or lower airway. Also provided are methods whereby such immunological response

slows bacterial replication. A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells. A NTHi protein or an antigenic peptide thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein. Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the NTHi polypeptides encoded by the polynucleotide of the invention or antigenic peptides thereof.

The invention correspondingly provides compositions suitable for eliciting an immune response to NTHi infection, wherein the antibodies elicited block binding of NTHi bacterium to the host's cells. The compositions comprise NTHi proteins or NTHi peptides of the invention, cells expressing the NTHi polypeptide, or polynucleotides encoding the polypeptides. The compositions may also comprise other ingredients such as carriers and adjuvants.

Genes that are up-regulated in NTHi infection of the middle ear and/or the nasopharynx and genes that are associated with NTHi virulence are described herein. The polypeptides and peptides thereof which are encoded by these NTHi genes are contemplated to be useful for eliciting an immune response for treating or preventing disorders associated with NTHi infection, such as OM. Some of the polypeptides encoded by these genes include: histidine biosynthesis protein, lipoprotein B, peptide ABC transporter, periplasmic SapA precursor, outer membrane lipoproteins carrier protein precursor, ribose transport system permease protein, phosphoribosylaminoimidazole carboxylase catalytic subunit, PurE, Phosphoribosylaminoimidazole carboxylase catalytic subunit, ornithine

carbamolytransferase, mannonate dehydratase, disulfide oxidoreductase, urease.
accessory protein, phosphocholine cytidylyltransferase, putative pyridoxine
biosynthesis protein, singlet oxygen resistance protein, intracellular septation protein,
DNA repair protein, MukF protein, glycerol-3-phosphate regulon repressor,
5 integration host factor beta subunit, arginine repressor, cold shock like protein, stress
response protein, LicA, MukF, RadA and those hypothetical proteins encoded by
HI0094, HI1163, HI0665, HI1292, HI1064 HI186, HI0352 genes. NTHi OMPs, LOS
and noncapsular proteins are also contemplated to elicit an immune response for
prevention and treatment of disorders associated with NTHi infection.

10 An "immunogenic dose" of a composition of the invention is one that
generates, after administration, a detectable humoral and/or cellular immune response
in comparison to the immune response detectable before administration or in
comparison to a standard immune response before administration. The invention
contemplates that the immune response resulting from the methods may be protective
15 and/or therapeutic.

The invention includes methods of blocking binding of NTHi bacteria
to host cells in an individual. The methods comprise administering antibodies or
polypeptides of the invention that block binding of NTHi cellular attachment.
Alternatively, administration of one or more small molecules that block binding of
20 NTHi cell attachment is contemplated. *In vitro* assays may be used to demonstrate
the ability of an antibody, polypeptide or small molecule of the invention to block
NTHi cell attachment.

Pharmaceutical compositions comprising antibodies of the invention,
polypeptides of the invention and/or small molecules of the invention that block NTHi
25 cellular attachment are provided. The pharmaceutical compositions may consist of
one of the foregoing active ingredients alone, may comprise combinations of the
foregoing active ingredients or may comprise additional active ingredients used to
treat bacterial infections. The pharmaceutical compositions may comprise one or
more additional ingredients such as pharmaceutically effective carriers. Dosage and
30 frequency of the administration of the pharmaceutical compositions are determined by
standard techniques and depend, for example, on the weight and age of the individual,
the route of administration, and the severity of symptoms. Administration of the

pharmaceutical compositions may be by routes standard in the art, for example, parenteral, intravenous, oral, buccal, nasal, pulmonary, rectal, or vaginal.

Also provided by the invention are methods for detecting NTHi infection in an individual. In one embodiment, the methods comprise detecting NTHi polynucleotides of the invention in a sample using primers or probes that specifically
5 bind to the polynucleotides. Detection of the polynucleotide may be accomplished by numerous techniques routine in the art involving, for example, hybridization and PCR.

The antibodies of the present invention may also be used to provide
10 reagents for use in diagnostic assays for the detection of NTHi antigens (NTHi polypeptides and peptides thereof) in various body fluids of individuals suspected of *H. influenzae* infection. In another embodiment, the NTHi proteins and peptides of the present invention may be used as antigens in immunoassays for the detection of NTHi in various patient tissues and body fluids including, but not limited to: blood, serum,
15 ear fluid, spinal fluid, sputum, urine, lymphatic fluid and cerebrospinal fluid. The antigens of the present invention may be used in any immunoassay system known in the art including, but not limited to: radioimmunoassays, ELISA assays, sandwich assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, fluorescent immunoassays, protein A immunoassays and
20 immunoelectrophoresis assays.

Vaccines and Chemotherapeutic Targets

An aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with a NTHi antigen protein or an antigenic peptide thereof.

25 The present invention also provides for vaccine formulations which comprise an immunogenic recombinant NTHi protein or NTHi peptide of the invention together with a suitable carrier. The NTHi polypeptides and peptides thereof contemplated as vaccine candidates and/or targets of chemotherapy include, but are not limited to, histidine biosynthesis protein, lipoprotein B, peptide ABC
30 transporter, periplasmic SapA precursor, outer membrane lipoproteins carrier protein precursor, ribose transport system permease protein, phosphoribosylaminoimidazole carboxylase catalytic subunit, PurE, 3,4-dihydroxy-2-butanone 4-phosphate synthase,

ornithine carbamolytransferase, mannonate dehydratase, disulfide oxidoreductase, urease accessory protein, phosphocholine cytidylyltransferase, putative pyridoxine biosynthesis protein, singlet oxygen resistance protein, intracellular septation protein, DNA repair protein, MUKF protein, glycerol-3-phosphate regulon repressor, integration host factor beta subunit, arginine repressor, cold shock like protein, stress response protein, LicA, RadA and those hypothetical proteins encoded by HI0094, HI1163, HI0665, HI1292, HI1064 HI1386, HI0352 genes, NTHi OMPs, NTHi LOS and NTHi noncapsular proteins and polypeptides encoded by the novel NTHi polynucleotide sequences present in the nucleotide sequences set out as SEQ ID NOS: 1-576, SEQ ID NOS: 675-685 and the nucleotide sequences set out in Table 3B, Table 4B and Table 5 herein, and the polypeptides having the amino acid sequences set out in Table 3B, Table 4B and Table 5 herein.

Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

A. Peptide Vaccines

Peptide therapeutic agents, such as peptide vaccines, are well known in the art and are of increasing use in the pharmaceutical arts. Consistent drawbacks to the parenteral administration of such peptide compounds have been the rapidity of breakdown or denaturation. Infusion pumps, as well as wax or oil implants, have been employed for chronic administration of therapeutic agents in an effort to both prolong the presence of peptide-like therapeutic agents and preserve the integrity of such

agents. Furthermore, the peptide-like agent should (with particular reference to each epitope of the peptide-like agent) ideally maintain native state configuration for an extended period of time and additionally be presented in a fashion suitable for triggering an immunogenic response in the challenged animal or immunized human.

5 The NTHi antigenic peptides of the invention can be prepared in a number of conventional ways. The short peptides sequences can be prepared by chemical synthesis using standard means. Particularly convenient are solid phase techniques (see, *e.g.*, Erikson *et al.*, *The Proteins* (1976) v. 2, Academic Press, New York, p. 255). Automated solid phase synthesizers are commercially available. In
10 addition, modifications in the sequence are easily made by substitution, addition or omission of appropriate residues. For example, a cysteine residue may be added at the carboxy terminus to provide a sulfhydryl group for convenient linkage to a carrier protein, or spacer elements, such as an additional glycine residue, may be
15 incorporated into the sequence between the linking amino acid at the C-terminus and the remainder of the peptide. The short NTHi peptides can also be produced by recombinant techniques. The coding sequence for peptides of this length can easily be synthesized by chemical techniques, *e.g.*, the phosphotriester method described in Matteucci *et al.*, *J Am Chem Soc.*, 103: 3185 (1981).

 Some of the NTHi peptide sequences contemplated herein may be
20 considered too small to be immunogenic, they may be linked to carrier substances in order to confer this property upon them. Any method of creating such linkages known in the art may be used. Linkages can be formed with heterobifunctional agents that generate a disulfide link at one functional group end and a peptide link at the other, such as a disulfide amide forming agent, *e.g.*, N-succidimidyl-3-(2-pyridyldithio)
25 propionate (SPDP) (See, *e.g.*, Jansen *et al.*, *Immun. Rev.* 62:185, 1982) and bifunctional coupling agents that form a thioether rather than a disulfide linkage such as reactive esters of 6-maleimidocaproic acid, 2-bromoacetic acid, 2-iodoacetic acid, 4-(N-maleimido-methyl) cyclohexane-1-carboxylic acid and the like, and coupling agent which activate carboxyl groups by combining them with succinimide or 1-
30 hydroxy-2-nitro-4-sulfonic acid, for sodium salt such as succinimmidyl 4-(N-maleimido-methyl) cyclohexane-1-carboxylate (SMCC).

B. Vaccine Compositions and Administration

A priming dose of the immunogen that is followed by one or more booster exposures to the immunogen may be necessary to be an effective vaccine (Kramp *et al.*, *Infect. Immun.*, 25: 771-773, 1979; Davis *et al.*, *Immunology Letters*, 14: 341-8 1986 1987). Examples of proteins or polypeptides that could beneficially enhance the immune response if co-administered include cytokines (*e.g.*, IL-2, IL-12, GM-CSF), cytokine-inducing molecules (*e.g.* Leaf) or costimulatory molecules. Helper (HTL) epitopes could be joined to intracellular targeting signals and expressed separately from the CTL epitopes. This would allow direction of the HTL epitopes to a cell compartment different than the CTL epitopes. If required, this could facilitate more efficient entry of HTL epitopes into the MHC class II pathway, thereby improving CTL induction. In contrast to CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (*e.g.* TGF- β) may be beneficial in certain diseases.

Ideally, an immunogen will exhibit two properties; the capacity to stimulate the formation of the corresponding antibodies and the propensity to react specifically with these antibodies. Immunogens bear one or more epitopes which are the smallest part of an immunogen recognizable by the combining site of an antibody. In particular instances, immunogen, fractions of immunogens or conditions under which the immunogen is presented are inadequate to precipitate the desired immunological response resulting in insufficient immunity. This is often the case with peptides or other small molecules used as immunogens. Other substances such as immunomodulators (*e.g.*, cytokines such as the interleukins) may be combined in vaccines as well.

The vaccine art recognizes the use of certain substances called adjuvants to potentate an immune response when used in conjunction with an immunogen. Adjuvants are further used to elicit an immune response that is faster or greater than would be elicited without the use of the adjuvant. In addition, adjuvants may be used to create an immunological response using less immunogen than would be needed without the inclusion of adjuvant, to increase production of certain antibody subclasses that afford immunological protection or to enhance components of the immune response (*e.g.*, humoral, cellular). Known adjuvants include emulsions such as Freund's Adjuvants and other oil emulsions, *Bordetella pertussis*, MF59,

purified saponin from *Quillaja saponaria* (QS21), aluminum salts such as hydroxide, phosphate and alum, calcium phosphate, (and other metal salts), gels such as aluminum hydroxide salts, mycobacterial products including muramyl dipeptides, solid materials, particles such as liposomes and virosomes. Examples of natural and bacterial products known to be used as adjuvants include monophosphoryl lipid A (MPL), RC-529 (synthetic MPL-like acylated monosaccharide), OM-174 which is a lipid A derivative from *E. coli*, holotoxins such as cholera toxin (CT) or one of its derivatives, pertussis toxin (PT) and heat-labile toxin (LT) of *E. coli* or one of its derivatives, and CpG oligonucleotides. Adjuvant activity can be affected by a number of factors, such as carrier effect, depot formation, altered lymphocyte recirculation, stimulation of T-lymphocytes, direct stimulation of B-lymphocytes and stimulation of macrophages.

Vaccines are typically prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients, which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, *e.g.*, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants, which enhance the effectiveness of the vaccine. The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25-70%.

Vaccines may also be administered through transdermal routes utilizing jet injectors, microneedles, electroporation, sonoporation, microencapsulation, polymers or liposomes, transmucosal routes and intranasal routes using nebulizers, aerosols and nasal sprays. Microencapsulation using natural or synthetic polymers such as starch, alginate and chitosan, D-poly L-lactate (PLA), D-poly DL-lactic-coglycolic microspheres, polycaprolactones, polyorthoesters, polyanhydrides and polyphosphazenes polyphosphatazanes are useful for both transdermal and transmucosal administration. Polymeric complexes comprising synthetic poly-ornithate, poly-lysine and poly-arginine or amphipathic peptides are useful for transdermal delivery systems. In addition, due to their amphipathic nature, liposomes are contemplated for transdermal, transmucosal and intranasal vaccine delivery systems. Common lipids used for vaccine delivery include N-(1,2,3-(dioleoyl-dihydroxypropyl)-N,N,N, - trimethylammonium-methyl sulfate (DOTAP), dioleoyloxy-propyl - trimethylammonium chloride DOTMA, dimystyloxypropyl-3-dimethyl-hydroxyethyl ammonium (DMRIE), dimethyldioctadecyl ammonium bromide (DDAB) and 9N(N',N-dimethylaminoethane) carbamoyl) cholesterol (DC-Chol). The combination of helper lipids and liposomes will enhance up-take of the liposomes through the skin. These helper lipids include, dioleoyl phosphatidylethanolamine (DOPE), dilauroylphosphatidylethanolamine (DLPE), dimyristoyl phosphatidylethanolamine (DMPE), dipalmitoylphosphatidylethanolamine (DPPE). In addition, triterpenoid glycosides or saponins derived from the Chilean soap tree bark (*Quillaja saponaria*) and chitosan (deacetylated chitan) have been contemplated as useful adjuvants for intranasal and transmucosal vaccine delivery.

The proteins may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, *e.g.*, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, *e.g.*, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, and procaine.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, capacity of the subject's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual. However, suitable dosage ranges are of the order of several hundred micrograms active ingredient per individual. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed in one or three month intervals by a subsequent injection or other administration.

Upon immunization with a vaccine composition as described herein, the immune system of the host responds to the vaccine by producing large amounts of CTLs specific for the desired antigen, and the host becomes at least partially immune to later infection, or resistant to developing chronic infection. Vaccine compositions containing the NTHi polypeptide or NTHi peptides of the invention are administered to a patient susceptible to or otherwise at risk of bacterial infection to elicit an immune response against the antigen and thus enhance the patient's own immune response capabilities. Such an amount is defined to be an "immunogenically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally range from about 1.0 µg to about 5000 per 70 kilogram patient, more commonly from about 10 to about 500 mg per 70 kg of body weight. For therapeutic or immunization purposes, the NTHi polypeptide or NTHi peptides of the invention can also be expressed by attenuated viral hosts, such as vaccinia or fowlpox. This approach involves the use of vaccinia virus as a vector to express nucleotide sequences that encode the peptides of the invention. Upon introduction into an acutely or chronically infected host or into a noninfected host, the recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits a host CTL response.

Humoral immune response may be measured by many well known methods, such as Single Radial Immunodiffusion Assay (SRID), Enzyme Immunoassay (EIA) and Hemagglutination Inhibition Assay (HAI). In particular, SRID utilizes a layer of a gel, such as agarose, containing the immunogen being tested. A well is cut in the gel and the serum being tested is placed in the well.

Diffusion of the antibody out into the gel leads to the formation of a precipitation ring whose area is proportional to the concentration of the antibody in the serum being tested. EIA, also known as ELISA (Enzyme Linked Immunoassay), is used to determine total antibodies in the sample. The immunogen is adsorbed to the surface of a microtiter plate. The test serum is exposed to the plate followed by an enzyme linked immunoglobulin, such as IgG. The enzyme activity adherent to the plate is quantified by any convenient means such as spectrophotometry and is proportional to the concentration of antibody directed against the immunogen present in the test sample. HAI utilizes the capability of an immunogen such as viral proteins to agglutinate chicken red blood cells (or the like). The assay detects neutralizing antibodies, *i.e.*, those antibodies able to inhibit hemagglutination. Dilution of the test serum are incubated with a standard concentration of immunogen, followed by the addition of the red blood cells. The presence of neutralizing antibodies will inhibit the agglutination of the red blood cells by the immunogen. Tests to measure cellular immune response include determination of delayed-type hypersensitivity or measuring the proliferative response of lymphocytes to target immunogen.

Nontypeable *Haemophilus influenzae* (NTHi)

H. influenzae is a small, nonmotile gram negative bacterium. Unlike other *H. influenzae* strains, the nontypeable *H. influenzae* (NTHi) strains lack a polysaccharide capsule and are sometimes denoted as "nonencapsulated." NTHi strains are genetically distinct from encapsulated strains and are more heterogenous than the type b *H. influenzae* isolates. NTHi presents a complex array of antigens to the human host. Possible antigens that may elicit protection include OMPs, lipopolysaccharides, lipoproteins, adhesion proteins and noncapsular proteins.

Humans are the only host for *H. influenzae*. NTHi strains commonly reside in the upper respiratory tract including the nasopharynx and the posterior oropharynx, the lower respiratory tract and the female genital tract. NTHi causes a broad spectrum of diseases in humans, including but not limited to, otitis media, pneumonia, sinusitis, septicemia, endocarditis, epiglottitis, septic arthritis, meningitis, postpartum and neonatal infections, postpartum and neonatal sepsis, acute and chronic salpingitis, epiglottis, pericarditis, cellulitis, osteomyelitis, endocarditis, cholecystitis, intraabdominal infections, urinary tract infection, mastoiditis, aortic graft infection, conjunctivitis, Brazilian purpuric fever, occult bacteremia and

exacerbation of underlying lung diseases such as chronic bronchitis, bronchiectasis and cystic fibrosis.

Epidemiologic studies of NTHi have indicated that the strains are heterogeneous with respect to outer membrane protein profiles (Barenkamp *et al.*, *Infect. Immun.*, 36: 535-40, 1982), enzyme allotypes (Musser *et al.*, *Infect. Immun.*, 52: 183-191, 1986), and other commonly used epidemiologic tools. There have been several attempts to subtype NTHi, but none of the methodologies have been totally satisfactory. The outer-membrane protein composition of NTHi consists of approximately 20 proteins. All NTHi strains contains two common OMP's with molecular weights of 30,000 and 16,600 daltons. NTHi strains may be subtyped based on two OMP's within the 32,000-42,000 dalton range. The NTHi liposaccharide profile is fundamentally different than the enteric gram negative bacteria and separates into 1-4 distinct bands ranging from less than 20,000 daltons.

A prototype NTHi isolate is the low passage isolate 86-028NP which was recovered from a child with chronic otitis media. This strain has been well characterized *in vitro* (Bakaletz *et al.*, *Infect. Immun.*, 53: 331-5, 1988; Holmes *et al.*, *Microb. Pathog.*, 23: 157-66, 1997) as well as in the chinchilla OM model (described herein) (Bakaletz *et al.*, *Vaccine*, 15: 955-61, 1997; Suzuki *et al.*, *Infect. Immun.*, 62: 1710-8, 1994; DeMaria *et al.*, *Infect. Immun.*, 64: 5187-92, 1996). The 86-028NP strain was used, as described herein, to identify genes that are up-regulated in expression in the chinchilla model of otitis media and genes that are necessary for NTHi survival in the chinchilla middle ear.

DFI Strategy

A differential fluorescence induction (DFI) strategy was used herein to identify NTHi genes induced during OM in a chinchilla animal model. Several methods have been developed to identify bacterial genes that contribute to the virulence of an organism during infection. Such methods include *in vivo* expression technology (IVET) in which bacterial promoters regulate the expression of gene(s) required for synthesis of essential nutrients required for survival in the host; signature-tagged mutagenesis (STM) enabling tag-specific identification of genes that alter the virulence properties of a microorganism when mutated; DNA microarray technology to globally screen for transcriptionally active genes, and DFI which uses FACS analysis to select for transcriptionally active promoters (Chiang *et al.*, *Annu. Rev.*

Microbiol., 53: 129-154, 1999). DFI is a high-throughput method that allows for the identification of differentially regulated genes regardless of the basal level of expression and does not exclude those that are essential for growth *in vitro*.

DFI has been successfully utilized in many microorganisms. For example, a GFP reporter system and flow cytometry was used to study mycobacterial gene expression upon interaction with macrophages (Dhandayuthapani *et al.*, *Mol. Microbiol.*, 17: 901-912, 1995). A promoter trap system was used to identify genes whose transcription was increased when *Salmonellae* were subjected to environments simulating *in vivo* growth and when internalized by cultured macrophage-like cells (Valdivia and Falkow, *Mol. Microbiol.*, 22: 367-378, 1996; Valdivia and Falkow, *Science*, 277: 2007-2011, 1997; Valdivia and Falkow, *Curr. Opin. Microbiol.*, 1: 359-363, 1998). In addition, DFI has been used to identify promoters expressed in *S. pneumoniae* and *S. aureus* when grown under varied *in vitro* conditions simulating infection (Marra *et al.*, *Infect. Immun.*, 148: 1483-1491, 2002; Schneider *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 97: 1671-1676, 2000). In addition, DFI has been utilized to study gene regulation in *Bacillus cereus* in response to environmental stimuli (Dunn and Handelsman, *Gene*, 226: 297-305, 1999), in *S. pneumoniae* in response to a competence stimulatory peptide (Bartilson *et al.*, *Mol. Microbiol.*, 39: 126-135, 2001), and upon interaction with and invasion of host cells in *Bartonella henselae* Lee and Falkow, *Infect. Immun.*, 66: 3964-3967, 1998), *Listeria monocytogenes* Wilson *et al.*, *Infect. Immun.*, 69: 5016-5024, 2001), *Brucella abortus* (Eskra *et al.*, *Infect. Immun.*, 69: 7736-7742, 2001), and *Escherichia coli* (Badger *et al.*, *Mol. Microbiol.*, 36: 174-182, 2000).

Whereas DFI has been successfully used to identify promoters active in cell culture models of infection or *in vitro* conditions designed to simulate an *in vivo* environment, few have applied DFI to identify promoters regulated in a specific biological niche within the whole animal. This is likely due to the numerous challenges associated with sorting from an *in vivo* environment. The host inflammatory response, dissemination and/or clearance of bacterial cells from the site of infection, as well as adherence of bacteria to epithelial cells, possibly via biofilm formation, can make bacteria inaccessible for retrieval from the living animal. These factors, among others, contribute to the complexity of the microenvironment and the heterogeneity of gene expression as the bacteria sense and respond to these changes.

Recently, DFI has been used to identify promoters expressed in *S. pneumoniae* when the bacteria were screened in a mouse model of respiratory tract infection and a gerbil infection model of OM (Marra *et al.*, *Infect. Immun.* 70: 1422-33, 2002; Marra *et al.*, *Microbiol.*, 148: 1483-91, 2002).

5 Animal Model

The chinchilla model is a widely accepted experimental model for OM. In particular, a chinchilla model of NTHi-induced OM has been well characterized (Bakaletz *et al.*, *J. Infect. Dis.*, 168: 865-872, 1993; Bakaletz and Holmes, *Clin. Diagn. Lab. Immunol.*, 4: 223-225, 1997; Suzuki and Bakaletz, *Infect. Immun.*, 62: 1710-1718, 1994), and has been used to determine the protective efficacy of several NTHi outer membrane proteins, combinations of outer membrane proteins, chimeric synthetic peptide vaccine components, and adjuvant formulations as vaccinogens against OM (Bakaletz *et al.*, *Vaccine*, 15: 955-961, 1997; Bakaletz *et al.*, *Infect. Immun.*, 67: 2746-2762, 1999; Kennedy *et al.*, *Infect. Immun.*, 68: 2756-2765, 2000).

15 In particular, there is an unique *in vivo* model wherein adenovirus predisposes chinchillas to *H. influenzae*-induced otitis media, which allowed for the establishment of relevant cell, tissue and organ culture systems for the biological assessment of NTHi (Bakaletz *et al.*, *J. Infect. Dis.*, 168: 865-72, 1993; Suzuki *et al.*, *Infect. Immunity* 62: 1710-8, 1994). Adenovirus infection alone has been used to
20 assess for the transudation of induced serum antibodies into the tympanum (Bakaletz *et al.*, *Clin. Diagnostic Lab Immunol.*, 4(2): 223-5, 1997) and has been used as a co-pathogen with NTHi, to determine the protective efficacy of several active and passive immunization regimens targeting various NTHi outer membrane proteins, combinations of OMPs, chimeric synthetic peptide vaccine components, and adjuvant
25 formulations as vaccinogens against otitis media (Bakaletz *et al.*, *Infect Immunity*, 67(6): 2746-62, 1999; Kennedy *et al.*, *Infect Immun.*, 68(5): 2756-65, 2000; Novotny *et al.*, *Infect Immunity* 68(4): 2119-28, 2000; Poolman *et al.*, *Vaccine* 19 (Suppl. 1): S108-15, 2000).

Genes Upregulated *In vivo* in Response to NTHi Infection of the Middle Ear

30 In order to identify differentially regulated promoters in response to NTHi infection of the middle ear, a promoter trap library was constructed and sorting parameters defined. A portion of the promoter trap library was inoculated directly into the chinchilla middle ear and OM development was monitored by video otoscopy

and tympanometry at 24 and 48 hours. In addition, the middle ear fluids were recovered 24 and 48 hours after infection. Two-color FACS analysis was used to isolated bacteria that were expressing GFP from other cells and debris associated with the effusion. Following isolation, DNA sequence of the *Haemophilus* inserts 5' of the *gfpmut3* gene were determined and analyzed. In this manner, we identified genes that are up-regulated as NTHi sense and respond to the environment of the chinchilla middle ear during AOM. The following genes were identified and due to their up-regulation during NTHi infection, they may play a role in NTHi infection and virulence.

As described below in Example 7, following the DFI procedure described above and subsequent FACS analysis of *gfp*-expressing clones, 52 candidate clones containing potential *in vivo*-regulated promoters were isolated. The genes these clones control were categorized based upon general description and function within the cell and include general metabolic processes, environmental informational processing and membrane transport, membrane proteins and hypothetical proteins. Eight of these 52 clones contain sequences that are unique to NTHi strain 86-028NP. Importantly, 3 clones were isolated from independent screens in more than one animal thereby verifying the method of isolation.

In order to independently confirm the FACS data, we determined the relative expression of candidate genes by quantitative RT-PCR. The parent strain 86-028NP, was used for these studies. Thus, wild-type gene expression without the influence of plasmid copy number on gene regulation was analyzed, allowing for the indication of false-positive clone identification by FACS. Of the 44 candidate clones containing sequence similar to that identified in *H. influenzae* strain Rd, quantitative comparison of gene expression *in vitro* and *in vivo* confirmed up-regulated gene expression for twenty-six genes (60%) when NTHi respond to environmental cues present in the chinchilla middle ear. This analysis identified *in vivo*-regulated promoters which drive expression of genes involved in membrane transport, environmental informational processing, cellular metabolism, gene regulation, as well as hypothetical proteins with unknown function. (See Table 4 in Example 6).

Quantitative RT-PCR demonstrated a two-fold increase in *lola* expression, enabling lipoprotein transport from the inner membrane to the outer membrane. Bacteria grow rapidly in the middle ear environment reaching 5.0×10^8

CFU NTHi ml middle ear fluid within 48 hours. The bacteria sense and respond to the environment, acquiring or synthesizing the necessary nutrients for growth and survival. The gene encoding the membrane component in ribose sugar transport, *rbsC* (SEQ ID NO: 619), showed a 5-fold increase in expression *in vivo* compared to cells
 5 growing *in vitro*. In addition, many genes involved in metabolic processes show a dramatic increase in gene expression *in vivo* compared to cells growing *in vitro*. These include a riboflavin synthesis gene, *ribB* (SEQ ID NO: 623), a purine nucleotide biosynthetic gene *purE* (SEQ ID NO: 621), ornithine carbamoyltransferase, *arcB* (SEQ ID NO: 625), involved in arginine degradation via
 10 the urea cycle and *uxuA* (SEQ ID NO: 627), encoding mannonate hydrolase, required for the uptake of D-glucuronate and transformation into glyceraldehyde 3-phosphate. In addition, but to a lesser degree, genes for histidine biosynthesis (*hisB*; SEQ ID NO: 615), DNA repair (*radC*; SEQ ID NO: 639) and a putative intracellular septation transmembrane protein (*ispZ*; SEQ ID NO: 637) were up-regulated.

15 Disulfide bond formation is important for folding and assembly of many secreted proteins in bacteria. In prokaryotes, DsbA and DsbB make up the oxidative pathway responsible for the formation of disulfides. DsbB reoxidizes DsbA, which donates disulfide bonds directly to unfolded polypeptides, and DsbB has been demonstrated to generate disulfides *de novo* from oxidized quinones (Collet and
 20 Bardwell, *Mol. Microbiol.*, 44: 1-8, 2002). In *H. influenzae* strain Rd, DsbA is required for competence for transformation (Tomb, *Proc. Natl. Acad. Sci. U.S.A.*, 89: 10252-10256, 1992). Herein, an approximate 3-fold increase in *dsbB* gene (SEQ ID NO: 629) transcription was demonstrated, illuminating an important role for disulfide interchange for NTHi growing in the middle ear environment.

25 Bacteria colonization of the middle ear, a normally sterile environment, results in a host inflammatory response and subsequent neutrophil infiltration. Bacteria have evolved numerous strategies to combat this host response. NTHi increase gene expression (4-fold) of *ureH* (SEQ ID NO: 631), a homologue of a gene required for expression of active urease in *Helicobacter*, shown to be involved in
 30 acid tolerance (Young *et al.*, *J. Bacteriol.*, 178: 6487-6495, 1996). Recently, it has been reported that urease activity may play a role in chronic *Actinobacillus pleuropneumoniae* infection by counteracting the decrease in pH occurring upon infection (Baltes *et al.*, *Infect. Immun.*, 69: 472-478, 2000; Baltes *et al.*, *Infect.*

Immun., 69: 472-478, 2001; Bosse and MacInnes, *Can. J. Vet. Res.*, 64: 145-150). A biotype analysis on NTHi isolates from middle ear effusions demonstrated that 87% are urease positive (DeMaria *et al.*, *J. Clin. Microbiol.*, 20: 1102-1104, 1984).

However, the role of urease in NTHi virulence is unknown. Similarly, an increase in
 5 expression of a gene whose product demonstrates 88% sequence identity to a pyridoxine biosynthesis protein in *S. pneumoniae* and 60% homology to a putative singlet oxygen resistance protein that may function as an antioxidant.

Phosphorylcholine (ChoP) has been implicated in the pathogenesis of NTHi (Weiser
et al., *Infect. Immun.*, 65: 943-950, 1997). NTHi modulates ChoP expression by
 10 phase variation, decorating the LOS on the cell surface. ChoP may contribute to NTHi persistence in the respiratory tract via decreased susceptibility to antimicrobial peptides (Lysecko *et al.*, *Infect. Immun.*, 68: 1664-1671, 2000) and alter the sensitivity to serum killing mediated by C-reactive protein (CRP) (Weiser *et al.*, *J. Exp. Med.*, 187: 631-640, 1998). The microenvironment of the nasopharynx and
 15 middle ear cavity may select for the ChoP⁺ phenotype, as ChoP⁺ strains show greater colonization of the chinchilla nasopharynx (Tong *et al.*, *Infect. Immun.*, 68: 4593-4597, 2000). Expression of the *licC* gene (SEQ ID NO: 633) was also increased. The *licC* gene encodes a phosphorylcholine cytidyltransferase that plays a role in the biosynthesis of phosphorylcholine-derivatized LOS (Rock *et al.*, *J. Bacteriol.*, 183:
 20 4927-4931, 2001).

Also included among the *in vivo*-induced genes is a set whose products subsequently regulate gene expression or DNA replication. These genes include transcriptional regulation of glycerol metabolism by the *glp* repressor, *glpR* (SEQ ID NO: 643), the arginine repressor gene, *argR* (SEQ ID NO: 647), and the integration
 25 host factor (IHF) beta subunit, *ihfB* (SEQ ID NO: 645). IHF is a histone-like protein that binds DNA at specific sequences, an accessory factor involved in replication, site-specific recombination and transcription, altering the activity of a large number of operons (Goosen and van de Putte, *Mol. Microbiol.* 16: 1-7, 1995). In addition, CspD inhibits DNA replication during stationary phase-induced stress response in *E.*
 30 *coli* (Yamanaka *et al.*, *Mol. Microbiol.*, 39: 1572-1584, 2001) and the *mukF* (SEQ ID NO: 641) gene protein homologue contributes to a remodeling of the nucleoid structure into a more compact form prior to cell segregation (Sawitzke and Austin, *Proc. Natl. Acad. Sci. U.S.A.*, 62: 1710-1718, 2000). The DFI strategy described

herein also identified promoters induced *in vivo* for genes of unknown function. The hypothetical protein, HI0094, demonstrated an 8-fold increase in gene expression during early OM but its role remains unknown. HI1163 (SEQ ID NO: 651) showed 58% amino acid identity with the hypothetical YdiJ proteins, a putative oxidase, of *E. coli*.

A high-density transposon mutagenesis strategy was used to identify *H. influenzae* genes essential for growth on rich medium (Akerley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 99: 966-971, 2002). Six genes were identified in the screen described herein that are included in essential gene set described in Akerley' *et al.*, *supra.* (*hisB*, *lppB*, *lola*, *ispZ*, *mukF* and unknown HI0665). Recently genes of non-typeable *H. influenzae* that are expressed upon interaction with two human respiratory tract-derived epithelial cell lines have been identified. These genes included those involved in metabolic processes, stress responses, gene expression, cell envelope biosynthesis, DNA-related processes, cell division and ORF's encoding proteins of unknown function. (Ulsen *et al.*, *Mol. Microbiol.*, 45: 485-500, 2002). Similarly the stress response gene, *cspD* (SEQ ID NO: 649), genes involved in purine and riboflavin biosynthesis, and a protein of unknown function, *vapA* was identified in the screen described herein. Expression of *vapA* was detected *in vitro*, yet *vapA* gene expression increased two-fold *in vivo*. These unique approaches identified known genes that are upregulated in NTHi-induced OM and therefore are likely to play a role in NTHi infection and virulence; and may be potential candidates for vaccines and antisense therapies and other therapeutic methods of treatment of NTHi related disorders.

The DFI strategy resulted in the identification of promoters induced *in vivo* for genes of unknown function as well. The hypothetical protein, HI0094, demonstrated a 8-fold increase in gene expression during early OM but its role remains unknown. HI1163 (SEQ ID NO: 651) showed 58% amino acid identity with the hypothetical YdiJ proteins, a putative oxidase, of *E. coli*. Therefore, these hypothetical genes are likely to play a role in OM induced by NTHi infection.

Brief Description of Figures

Figure 1 depicts the LKP gene region in a panel of *Haemophilus* isolates. The strain 86-028NP sequence is identical in this region to the sequence in

NTHi strain R3001. Both of these NTHi lack the *hif* gene cluster encoding the hemagglutinating pilus.

Figure 2 depicts the *rfaD* region in a panel of *Haemophilus* isolates. The gene arrangement in the *rfaD* region of the strain 86-028NP genome is similar to that of the strain Rd genome but different than the arrangement of these genes seen in the genome of most NTHi examined.

Figures 3A-3M set out the nucleotide sequences (SEQ ID NOS: 589-614) described in Table 4, which were identified to be upregulated during OM infection (see Example 6). The nucleotides (nt.) which correspond to known genes and those nt. which correspond to the contig sequences set out as SEQ ID NO: 1-576 are also presented.

Detailed Description

The following examples illustrate the invention wherein Example 1 describes the sequence of the NTHi genome, Example 2 describes the identified contigs and initial gene discovery, Example 3 describes construction of the NTHi promoter trap library, Example 4 describes the analyses of 86-028NP derivatives expressing GFP, Example 5 demonstrates direct labelling of bacteria from middle ear fluids, Example 6 describes identification of promoters induced *in vivo* in acute otitis media, Example 7 describes identification of virulence-associated genes, and Example 8 describes identification of unique NTHi gene sequences.

Example 1

Sequence of a Non-Typeable *Haemophilus influenzae* Genome

NTHi strain 86-028NP is a minimally passaged clinical isolate obtained from a pediatric patient who underwent tympanostomy and tube insertion for chronic OM at Columbus Children's Hospital. (Bakaletz *et al. Infection and Immunity*, 56(2): 331-335, 1988) The 86-028NP strain was deposited with the American Type Tissue Collection (Manassas, VA 20108 USA) on October 16, 2002 and assigned accession no. PTA-4764.

In an effort to more broadly approach the identification of the virulence determinants in NTHi, the genome of the NTHi 86-028NP strain was sequenced to 3-fold coverage. Chromosomal DNA was prepared from strain 86-028NP using the Puregene protocol and sheared to 2-4 kb in size with a Hydroshear

instrument (Gene Machines). The sheared DNA was ethanol-precipitated, end-repaired using a mixture of Klenow enzyme and T4 DNA polymerase, and size-selected by agarose gel electrophoresis to obtain 2-4 kb fragments as described in Chissoe *et al.* (*Methods: a Companion to Methods of Enzymology* 3: 55-65, 1991) and
 5 Sambrook *et al.* (*Molecular Cloning: a Laboratory Manual*, 2nd Ed. Cold Springs Harbor, NY, 1989). These fragments were cloned into vector pUC18 using the *Sma*I restriction site (phosphatase-treated) and transformed into *E. coli* XL-1 Blue, selecting for ampicillin resistance. Colonies that contain inserts were identified by blue/white screening on LB-Amp plates containing X-gal, and transferred into 96-deep well
 -10—plates containing 1.5 ml of TB-Amp (TB = Terrific Broth) broth. The deep-well plate cultures were grown overnight (18-22 hours) at 37°C. Template preparation, sequencing and contig assembly were performed.

Automated template preparation was performed on the Beckman Biomek 2000 automated robotics workstation as described in Chissoe *et al.*, (*supra.*)
 15 Briefly, each 96-deep well plate, containing the clones prepared above, was centrifuged to pellet the cells, the supernatant decanted, and the cells frozen (if necessary) at -20°C. Four 96-deep well blocks were placed on the Biomek table, and the liquid handling robot was used to prepare the template using an automated version of a typical SDS-NaOH lysis protocol as described in Chissoe *et al.*, (*supra.*). The
 20 final ethanol-precipitated templates were each dissolved in 50 µl ddH₂O, and used for DNA sequencing.

Sequencing reactions were run by re-arraying the templates (from 96-well plates) into 384-well plates, using the Robbins Hydra 96 robot. Cycle-sequencing reactions were run using PE Big-Dye™ terminators and universal primers
 25 (M13 forward and reverse), cleaned up over Sephadex G50 columns, and analyzed on a PE Biosystems 3700 capillary electrophoresis DNA sequencer according to the manufacturer's instructions. Sequencing reads (8219) were assembled into 576 contigs (SEQ ID NOS: 1- 576 herein). The statistics for the 3-fold sequencing are shown in Table 2A. The total unique sequence in assembly 17 is 1.74Mb.

30

Table 2A

WO 2004/078949

PCT/US2004/007001

33

Contig Size	Total Number	Total Length	% of Cumulative
0 – 1 kb	65	55961	3.2%
1 – 2 kb	228	333665	19.2%
2 – 3 kb	101	243059	14.0%
3 – 4 kb	49	172385	9.9%
4 – 5 kb	45	196699	11.3%
5 – 10 kb	74	515152	29.6%
10 – 20 kb	11	144591	8.3%
20 – 30 kb	3	77352	4.4%

Subsequently, 8-fold sequencing analysis of the NTHi genome was carried out. The 8-fold sequencing assembled the NTHi genome into 11 contigs. Contigs 5, 8, 9, 10, 12-18 are denoted as SEQ ID NOS: 675-685 herein. The statistics for the 8-fold sequencing are shown in Table 2B.

5

TABLE 2B.

Contig Size	Total Number	Total Length	% of Cumulative
0 – 1 kb	5	3950	0.2%
1 – 2 kb	3	4316	0.2%
2 – 3 kb	0	0	0.0%
3 – 4 kb	1	3964	0.2%
4 – 5 kb	0	0	0.0%
5 – 10 kb	0	0	0.0%
10 – 20 kb	1	15147	0.8%
20 – 30 kb	2	51888	2.7%
30 - 40 kb	0	0	0.0%
40 - 50 kb	0	0	0.0%
50 - 100 kb	1	85814	4.5%
>100 kb	5	1760339	91.4%

Example 2

Contig Description and Initial Gene Discovery

Seventy-five of the 88 contigs with length ≥ 5000 bp, identified with the 3-fold sequence analysis, show significant similarity via BLASTN to genes in *H.*

10 *influenzae* strain Rd. To visualize the potential relationship between the gene order in *H. influenzae* strain 86-028NP and *H. influenzae* strain Rd, the 86-028NP three-fold

contig set and the Rd gene set were bidirectionally compared using BLASTN. The results were plotted in gene-order verses contig space by sorting the contigs based on gene coordinates of the Rd genes hit, anchoring each contig at the smallest coordinate found as described in Ray *et al.*, (*Bioinformatics* 17: 1105-12, 2001). Compared in
 5 this fashion, an incomplete assembly of a genome with identical gene order to a completely known genome would display a monotonically increasing stair-stepped form.

BLASTX was used to identify hits to sequences with homology to genes in the strain Rd genome as well as genes not found in *H. influenzae* strain Rd.
 10 Hits to strain Rd sequences were removed from the data set and the other hits summarized in Table 3A. The data are presented as follows: contig # (= SEQ ID NO: #), column 1; E score for each hit, column 2; the name of the protein that had homology to a portion of the amino acid translation of the cited contig, column 3; the organism producing the homologue, column 4; and the Genbank protein identifier for
 15 each of the proteins cited in column 3, column 5; the corresponding nucleotides within the contig (referenced by SEQ ID NO:). In most instances, several homologues were identified but for clarity, the protein of greatest homology is cited in Table 3A.

The sequences for some of the genes listed in Table 3A were identified
 20 within the 8-fold sequencing of the NTHi genome. Table 3B lists the location of these genes within the 11 contigs, the full length open reading frame sequence (identified by SEQ ID NO:), the derived amino acid sequence encoded by the open reading frame and the gene with high homology identified by BLASTX (as listed in Table 3A).

25 To examine the relative short range gene arrangements in NTHi and the Rd strain, the gene order in two gene clusters that have been well-described were compared. First, the genes present in the hemagglutinating pilus (LKP) gene region were examined. (Mhlanga-Mutangadura *et al.*, *J Bacteriol.* 180(17): 4693-703, 1998). The pilus gene cluster is located between the *purE* and *pepN* genes, only
 30 fragments of which are depicted in Figure 1. The serotype b strain, Eagan, contains the *hifABCDE* gene cluster and produces hemagglutinating pili. Strain Rd lacks the *hicAB* genes as well as the *hifABCDE* gene cluster. In general, the nontypeable strains previously examined contained the *hicAB* genes but not the *hif* genes that

encode the hemagglutinating pilus. The strain 86-028NP sequence (described herein) is identical in this region to the sequence in NTHi strain R3001 (Figure 1). The *rfaD* gene encodes an enzyme involved in the biosynthesis of endotoxin. In addition, the *rfaD* gene from NTHi strain 2019 has been characterized by Nichols *et al.* (*Infect*
 5 *Immunity* 65(4): 1377-86, 1997). In strain 2019, the *rfaD* gene is immediately upstream of the *rfaF* gene that encodes another enzyme involved in endotoxin biosynthesis. The gene arrangement in strain Rd is different; the *rfaD* and *rfaF* genes are separated by approximately 11 kb of sequence. Most nontypeable strains examined contained the gene arrangement seen in strain 2019. In contrast, strain 86-
 10 028NP has a gene arrangement identical to that seen in strain Rd (Figure 2).

A global analysis of the current assembly indicates that the gene content and order are similar to that in strain Rd. A more detailed analysis revealed that there are a substantial number of NTHi genes not previously seen in the *Pasteurellaceae* and some regions where the NTHi gene content and order is different
 15 than that seen in strain Rd. Thus, the current data suggest that the strain 86-028NP genome will contain a complex mosaic of Rd and non-Rd like features.

The DFI strategy also identified novel NTHi sequences that had increased gene expression. A list of these novel contig sequences that contain genes or gene fragments that have homology to ORFs in other organisms (primarily gram-
 20 negative bacteria) is set out in Table 3A. For example, the nucleotide sequence of contig 442 (SEQ ID NO: 442), nucleotides 1498-1845 are highly homologous to the sequences encoding amino acids 1-116 of H. influenzae strain Rd lipoprotein B (LppB). The gene is positioned between the stationary phase survival gene, *surE*, and a gene encoding a 43 kD antigenic outer membrane lipoprotein that is highly
 25 homologous to the recently identified bacterial lipoprotein, LppB/NlpD, which has been associated with virulence (Padmalayam *et al.*, *Infect. Immun.*, 68: 4972-4979, 2000). Recently, Zhang and coworkers demonstrated that *nlpD* and *surE* gene expression was induced during stationary phase of bacterial growth in *Thermotoga maritima* (Zhang *et al.*, *Structure (Camb)*, 9: 1095-1106, 2001). Therefore, under
 30 stress-induced conditions in the middle ear, this NTHi lipoprotein may be expressed.

Table 3A

Contig	E score	Hit Identity	Organism	Genbank Protein	SEQ ID NO:
104	4.00E-59	CpdB	Pasteurella multocida	NP_246953.1	nt. 204-659 of SEQ ID NO: 104
106	9.00E-10	hypothetical protein PH0217 -	Pyrococcus horikoshii	G71244	nt. 40-309 of SEQ ID NO: 106
106	1.00E-08	unknown	Pasteurella multocida	NP_246871.1	nt. 605-694 of SEQ ID NO: 106
106	2.00E-20	Orf122	Chlorobium tepidum	AAG12204.1	nt. 7-210 of SEQ ID NO: 106
110	3.00E-05	ArdC antirestriction protein	IncW plasmid pSa	AAD52160.1	compliment of nt. 959-1162 of SEQ ID NO: 110
110	1.00E-33	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458676.1	compliment of nt. nt. 181-825 of SEQ ID NO: 110
111	5.00E-12	putative membrane protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458664.1	compliment of nt. 45-287 of SEQ ID NO: 111
111	6.00E-41	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458658.1	compliment of nt. 1091-1480 of SEQ ID NO: 111
114	7.00E-80	unknown	Pasteurella multocida	NP_245828.1	compliment of nt. 118-696 of SEQ ID NO: 114
115	2.00E-09	A111R	Paramecium bursaria Chlorella virus 1	NP_048459.1	nt. 555-869 of SEQ ID NO: 115
118	5.00E-45	DNA methylase HsdM, putative	Vibrio cholerae	NP_231404.1	nt. 44-439 of SEQ ID NO: 118
122	2.00E-18	unknown	Pasteurella multocida	NP_245314.1	nt. 865-1302 of SEQ ID NO: 122
123	4.00E-99	RNA POLYMERASE SIGMA-32 FACTOR	Proteus mirabilis	P50509	nt. 351-782 of SEQ ID NO: 123
124	9.00E-58	ACETOLACTATE SYNTHASE (ACETOHYDROXY-ACID SYNTHASE) (ALS)	Spirulina platensis	P27868	nt. 603-1025 of SEQ ID NO: 124
130	0	restriction modification system-R protein	Neisseria meningitidis	CAA09003.1	nt. 495-1559 of SEQ ID NO: 130
131	6.00E-91	uronate isomerase (glucuronate isomerase)	Salmonella enterica subsp. enterica serovar Typhi	NP_457532.1	compliment of nt. 661-1380 of SEQ ID NO: 131
133	3.00E-30	GyrA	Pasteurella multocida	NP_245778.1	compliment of nt. 1447-1626 of SEQ ID NO: 133
133	1.00E-27	DNA GYRASE SUBUNIT A	Pectobacterium carotovorum	P41513	compliment of nt. 1302-1442 of SEQ ID NO: 133
138	7.00E-06	KicA	Pasteurella multocida	NP_245545.1	compliment of nt. 92-157 of SEQ ID NO: 138

138	1.00E-148	TYPE II RESTRICTION ENZYME HAEII (ENDONUCLEASE HAEII) (R.HAEII)	Haemophilus aegyptius	O30869	compliment of nt. 164-1045 of SEQ ID NO: 138
143	4.00E-06	Gifsy-1 prophage protein	Salmonella typhimurium LT2	NP_461555.1	compliment of nt. 228-632 of SEQ ID NO: 143
143	1.00E-14	hypothetical protein	Bacteriophage VT2-Sa	NP_050531.1	compliment of nt. 778-1248 of SEQ ID NO: 143
143	5.00E-09	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	CAD09979.1	compliment of nt. 715-1026 of SEQ ID NO: 143
143	6.00E-10	hypothetical 14.9kd protein	Escherichia coli	NP_065324.1	nt. 3-173 of SEQ ID NO: 143
147	1.00E-38	GTP-binding elongation factor, may be inner membrane protein	Escherichia coli O157:H7 EDL933	NP_289127.1	compliment of nt. 172-342 of SEQ ID NO: 147
147	2.00E-14	GTP-binding membrane protein (lepA)	Borrelia burgdorferi	NP_212222.1	compliment of nt. 17-181 of SEQ ID NO: 147
148	6.00E-17	galactokinase	Homo sapiens	AAC35849.1	compliment of nt. 746-1246 of SEQ ID NO: 148
148	7.00E-96	GALACTOKINASE (GALACTOSE KINASE)	Actinobacillus pleuropneumoniae	P94169	compliment of nt. 232-741 of SEQ ID NO: 148
149	1.00E-92	GTP-binding protein TypA/BipA	Buchnera sp. APS	NP_240245.1	compliment of nt. 265-1077 of SEQ ID NO: 149
15	2.00E-21	ORF 1	Escherichia coli	CAA39631.1	nt. 665-850 of SEQ ID NO: 15
150	6.00E-17	unknown	Pasteurella multocida	NP_245919.1	nt. 171-665 of SEQ ID NO: 150
153	7.00E-07	outer membrane protein A	Rickettsia conorii	T30852	nt. 51-623 of SEQ ID NO: 153
155	7.00E-40	cytochrome d ubiquinol oxidase, subunit II	Vibrio cholerae	NP_233259.1	nt. 583-1002 of SEQ ID NO: 155
157	7.00E-13	unknown	Pasteurella multocida	NP_245490.1	compliment of nt. 1170-1367 of SEQ ID NO: 157
157	2.00E-05	glycosyl transferase	Neisseria gonorrhoeae	AAA68012.1	nt. 85-189 of SEQ ID NO: 157
158	1.00E-152	MltC	Pasteurella multocida	NP_246259.1	compliment of nt. 36-530 of SEQ ID NO: 158
161	3.00E-25	lipoprotein, putative	Vibrio cholerae	NP_230232.1	nt. 870-1439 of SEQ ID NO: 161
163	9.00E-53	chorismate synthase	Caulobacter crescentus	NP_421948.1	nt. 1283-2029 of SEQ ID NO: 163
168	3.00E-13	COPPER-TRANSPORTING ATPASE 1 (COPPER PUMP 1)	Mus musculus	Q64430	nt. 66-995 of SEQ ID NO: 168
168	2.00E-22	Cu transporting ATPase P	Homo sapiens	2001422A	nt. 135-989 of SEQ ID NO: 168
174	8.00E-48	magnesium/cobalt transport protein	Mesorhizobium loti	NP_103977.1	nt. 918-1205 of SEQ ID NO: 174
175	5.00E-26	vacB protein	Buchnera sp. APS	NP_240369.1	compliment of nt. 1-1587 of SEQ ID NO: 175

176	3.00E-21	putative ABC transport system permease protein [	Campylobacter jejuni	NP_282774.1	compliment of nt. 259-1089 of SEQ ID NO: 176
183	5.00E-29	PROBABLE ATP SYNTHASE A CHAIN TRANSMEMBRAN E PROTEIN	Ralstonia solanacearum	NP_521442.1	compliment of nt. 42-677 of SEQ ID NO: 183
185	6.00E-85	putative exported protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458655.1	compliment of nt. 162-1529 of SEQ ID NO: 185
187	2.00E-05	transketolase	Homo sapiens	AAA61222.1	nt. 709-819 of SEQ ID NO: 187
188	1.00E-116	ribonuclease E	Xylella fastidiosa 9a5c	NP_299884.1	compliment of nt. 280-1704 of SEQ ID NO: 188
192	1.00E-38	ImpA	Pasteurella multocida	NP_245829.1	nt. 35-448 of SEQ ID NO: 192
193	3.00E-08	Orf80	Enterobacteria phage 186	NP_052285.1	nt. 1612-1818 of SEQ ID NO: 193
193	1.00E-06	holin	Haemophilus somnus	AAC45168.1	nt. 370-576 of SEQ ID NO: 193
193	0.007	unknown	Enterobacteria phage 186	NP_052260.1	nt. 1376-1609 of SEQ ID NO: 193
193	2.00E-48	lysozyme	Haemophilus somnus	AAC45169.1	nt. 608-1093 of SEQ ID NO: 193
199	4.00E-21	unknown protein	Escherichia coli O157:H7 EDL933, prophage CP-933V	NP_288675.1	nt. 398-778 of SEQ ID NO: 199
199	2.00E-49	hypothetical protein	Bacteriophage 933W	NP_049495.1	compliment of nt. 1907-2392 of SEQ ID NO: 199
20	1.00E-62	RpL14	Pasteurella multocida	NP_246344.1	compliment of nt. 233-601 of SEQ ID NO: 20
200	2.00E-62	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458658.1	compliment of nt. 431-997 of SEQ ID NO: 200
200	3.00E-16	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458657.1	compliment of nt. 1028-1264 of SEQ ID NO: 200
201	2.00E-26	TsaA	Pasteurella multocida	NP_245732.1	compliment of nt. 1618-1809 of SEQ ID NO: 201
209	6.00E-16	TsaA	Pasteurella multocida	NP_245732.1	compliment of nt. 2-136 of SEQ ID NO: 209
211	2.00E-15	unknown	Pasteurella multocida	NP_245535.1	compliment of nt. 23-211 of SEQ ID NO: 211
211	1.00E-70	PUTATIVE ATPASE PROTEIN	Ralstonia solanacearum	NP_520082.1	compliment of nt. 475-915 of SEQ ID NO: 211
212	3.00E-18	hypothetical protein	Escherichia coli O157:H7	NP_309775.1	compliment of nt. 895-1035 of SEQ ID NO: 212
216	1.00E-173	unknown	Pasteurella multocida	NP_245069.1	nt. 35-1543 of SEQ ID NO: 216
217	9.00E-18	diacylglycerol kinase	Vibrio cholerae	NP_233101.1	nt. 2083-2208 of SEQ ID NO: 217

221	4.00E-34	Tail-Specific Protease	Chlamydia trachomatis	NP_219953.1	nt. 849-1421 of SEQ ID NO: 221
222	4.00E-23	AGR_C_3689p	Agrobacterium tumefaciens str. C58 (Cereon)	NP_355005.1	compliment of nt. 940-1305 of SEQ ID NO:222
224	9.00E-19	unknown	Pasteurella multocida	NP_245536.1	nt. 15-308 of SEQ ID NO: 224
225	1.00E-89	portal vector -like protein, in phage P2 [Salmonella typhimurium LT2]	Salmonella typhimurium LT2Fels-2 prophage	NP_461651.1	nt. 31-750 of of SEQ ID NO: 225
229	2.00E-25	anaerobic ribonucleotide reductase	Salmonella typhimurium	CAB62266.1	nt. 1806-2108 of SEQ ID NO: 229
234	3.00E-08	conserved hypothetical protein	Xylella fastidiosa 9a5c	NP_299850.1	nt. 1680-2048 of SEQ ID NO: 234
234	1.00E-42	Methionine sulfoxide reductase C-terminal domain related protein, YPPQ ortholog	Clostridium acetobutylicum	NP_348177.1	compliment of nt. 415-654 of SEQ ID NO: 234
235	4.00E-16	phage-related tail protein	Wolbachia endosymbiont of Drosophila melanogaster	AAK85310.1	compliment of nt. 931-1929 of SEQ ID NO: 235
235	6.00E-56	similar to orfG protein in phage 186, Salmonella typhimurium LT2	Salmonella typhimurium LT2, Fels-2 prophage	NP_461625.1	compliment of nt. 313-1863 of SEQ ID NO: 235
236	6.00E-20	conserved hypothetical protein	Pseudomonas aeruginosa	NP_252693.1	nt. 1572-1916 of SEQ ID NO: 236
240	5.00E-27	MODIFICATION METHYLASE BEPI	Brevibacterium epidermidis	P10283	compliment of nt. 922-1305 of SEQ ID NO: 240
241	2.00E-15	phage-related protein	Xylella fastidiosa 9a5c	NP_299573.1	compliment of nt. 865-1305 of SEQ ID NO: 241
241	4.00E-08	hypothetical protein	phage SPP1	T42296	nt. 73-636 of SEQ ID NO: 241
241	4.00E-07	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458686.1	nt. 10-468 of SEQ ID NO: 241
242	2.00E-29	translation elongation factor EF-G	chloroplast - soybean	S35701	compliment of nt. 18-1085 of SEQ ID NO: 242
247	3.00E-23	GTP CYCLOHYDROLA SE I (GTP-CH-I)	Synechococcus sp. PCC 7942	Q54769	compliment of nt. 1009-1257c of SEQ ID NO: 247
248	6.00E-05	phospho-N- acetylmuramoyl- pentapeptide- transferase	Aquifex aeolicus	NP_213025.1	nt. 830-1747 of SEQ ID NO: 248
25	2.00E-86	PROBABLE TRANSPORT TRANSMEMBRAN E PROTEIN	Ralstonia solanacearum	NP_522358.1	compliment of nt. 309-854 of SEQ ID NO: 25
25	7.00E-06	major facilitator family transporter	Caulobacter crescentus	NP_419155.1	compliment of nt. 134-283 of SEQ ID NO: 25
250	1.00E-150	CpdB	Pasteurella multocida	NP_246953.1	compliment of nt. 36-1016 of SEQ ID NO: 250

252	3.00E-57	alanyl-tRNA synthetase	Vibrio cholerae	AAA99922.1	compliment of nt. 1418-1951 of SEQ ID NO: 252
253	1.00E-108	similar to glutathione Reductase	Listeria monocytogenes EGD-e	NP_464432.1	compliment of nt. 411-1358 of SEQ ID NO: 253
259	3.00E-39	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458654.1	compliment of nt. 342-1037 of SEQ ID NO: 259
259	3.00E-17	possible exported protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458653.1	compliment of nt. 1251-1607 of SEQ ID NO: 259
261	5.00E-74	hypothetical protein 6 - Haemophilus influenzae	Haemophilus influenzae	S27582	compliment of nt. 3-422 of SEQ ID NO: 261
263	1.00E-94	putative transposase	Haemophilus paragallinarum	AAD01406.1	nt. 2142-2672 of SEQ ID NO: 263
264	1.00E-126	unknown	Actinobacillus actinomycetemcomitans	NP_067554.1	nt. 40-714 of SEQ ID NO: 264
264	1.00E-103	unknown	Actinobacillus actinomycetemcomitans	NP_067555.1	nt. 695-1309 of SEQ ID NO: 264
264	2.00E-21	unknown	Actinobacillus actinomycetemcomitans	NP_067556.1	nt. 1302-1448 of SEQ ID NO: 264
265	6.00E-27	Aminopeptidase 2	chloroplast	Q42876	nt. 556-1539 of SEQ ID NO: 265
268	1.00E-116	MutY	Pasteurella multocida	NP_246257.1	nt. 1003-1581 of SEQ ID NO: 268
272	1.00E-07	hypothetical protein	Bacteriophage 933W	NP_049495.1	compliment of nt. 77-232 of SEQ ID NO: 272
274	3.00E-13	unknown	Pasteurella multocida	NP_246952.1	compliment of nt. 1658-1975 of SEQ ID NO: 274
275	3.00E-20	CafA	Neisseria gonorrhoeae	AAG24267.1	nt. 1299-1571 of SEQ ID NO: 275
276	1.00E-45	mukE protein	Vibrio cholerae	NP_231351.1	compliment of nt. 650-1390 of SEQ ID NO: 276
276	1.00E-69	KicA	Pasteurella multocida	NP_245545.1	compliment of nt. 647-1321 of SEQ ID NO: 276
278	2.00E-56	3-oxoacyl-[acyl-carrier-protein] synthase III	Salmonella enterica subsp. enterica serovar Typhi	NP_455686.1	nt. 1366-1944 of SEQ ID NO: 278
281	5.00E-56	unknown	Pasteurella multocida	NP_246261.1	compliment of nt. 31-678 of SEQ ID NO: 281
282	3.00E-09	orf25; similar to T gene of P2	bacteriophage phi CTX	NP_490625.1	compliment of nt. 511-1032 of SEQ ID NO: 282
282	1.00E-08	orf11; similar to phage P2 gene S-like product, which is involved in tail synthesis,	Haemophilus somnus	AAC45165.1	compliment of nt. 1450-1584 of SEQ ID NO: 282
282	9.00E-27	putative bacteriophage tail protein	Salmonella enterica subsp. enterica serovar Typhi	NP_457167.1	compliment of nt. 3-509 of SEQ ID NO: 282

286	5.00E-18	plasmid-related protein	Listeria innocua plasmid	NP_471066.1	compliment of nt. 887-1501 of SEQ ID NO: 286
287	8.00E-20	GTP cyclohydrolase II	Escherichia coli O157:H7 EDL933	NP_287920.1	nt. 2-145 of SEQ ID NO: 287
289	1.00E-168	MODIFICATION METHYLASE HAEII	Haemophilus aegyptius	O30868	compliment of nt. 138-1091 of SEQ ID NO: 289
289	5.00E-11	TYPE II RESTRICTION ENZYME HAEII	Haemophilus aegyptius	O30869	compliment of nt. 22-132 of SEQ ID NO: 289
289	6.00E-47	mukF homolog	Haemophilus influenzae biotype aegyptius	AAB70828.1	compliment of nt. 1107-1385 of SEQ ID NO: 289
294	1.00E-171	LICA PROTEIN	Haemophilus influenzae RM7004	P14181	compliment of nt. 677-1564 of SEQ ID NO: 294
297	1.00E-158	DNA methylase HsdM, putative	Vibrio cholerae	NP_231404.1	compliment of nt. 12-1136 of SEQ ID NO: 297
302	0	HEME-BINDING PROTEIN A	Haemophilus influenzae DL42	P33950	nt.3-1316 of SEQ ID NO: 302
304	6.00E-19	hypothetical protein 6	Haemophilus influenzae	S27582	nt. 121-267 of SEQ ID NO: 304
305	6.00E-40	putative recombinase - phage associated	Streptococcus pyogenes M1 GAS	NP_269557.1	nt. 65-805 of SEQ ID NO: 305
305	7.00E-22	single stranded DNA-binding protein	Shewanella sp. F1A	AAB57886.1	nt. 1607-2014 of SEQ ID NO: 305
305	1.00E-43	phage-related protein	Bacillus halodurans	NP_244410.1	nt. 92-751 of SEQ ID NO: 305
312	1.00E-28	PUTATIVE BACTERIOPHAGE -RELATED TRANSMEMBRANE PROTEIN	Ralstonia solanacearum	NP_518994.1	nt. 1819-2673 of SEQ ID NO:312
312	9.00E-25	similar to BASEMENT MEMBRANE-SPECIFIC HEPARAN SULFATE PROTEOGLYCAN CORE PROTEIN PRECURSOR (HSPG)	Homo sapiens	XP_068727.1	nt. 27-1001 of SEQ ID NO: 312
315	2.00E-45	uracil permease	Deinococcus radiodurans	NP_296001.1	compliment of nt. 525-1592 of SEQ ID NO: 315
318	7.00E-15	CzcD	Pasteurella multocida	NP_246276.1	compliment of nt. 3-227 of SEQ ID NO: 318
320	2.00E-60	orf3; similar to endonuclease subunit of the phage P2 terminase (gene M)	Haemophilus somnus	AAC45159.1	compliment of nt. 606-1241 of SEQ ID NO: 320
320	2.00E-09	orf4; similar to head completion/stabilization protein (gene L) of phage P2	Haemophilus somnus	AAC45160.1	compliment of nt. 52-285 of SEQ ID NO: 320

320	3.00E-35	orf2; similar to major capsid protein precursor of phage P2 (gene N)	Haemophilus somnus	AAC45158.1	compliment of nt. 1271-1624 of SEQ ID NO: 320
323	4.00E-37	dedC protein	Escherichia coli	AAA23966.1	compliment of nt. 74-463 of SEQ ID NO: 323
324	1.00E-153	conserved hypothetical protein	Neisseria meningitidis MC58	NP_274972.1	compliment of nt. 930-1943 of SEQ ID NO: 324
326	5.00E-52	selenophosphate synthetase	Eubacterium acidaminophilum	CAB53511.1	compliment of nt. 1186-2292 of SEQ ID NO: 326
328	1.00E-129	secretion protein SecD	Pseudomonas aeruginosa	NP_252510.1	compliment of nt. 8-625 of SEQ ID NO: 328
333	3.00E-08	unknown	Pasteurella multocida	NP_245489.1	compliment of nt. 5-418 of SEQ ID NO: 333
336	6.00E-38	probable methyl transferase	Pseudomonas aeruginosa	NP_253353.1	compliment of nt. 2547-2819 of SEQ ID NO: 336
338	2.00E-98	Pmi	Pasteurella multocida	NP_245766.1	nt. 144-842 of SEQ ID NO: 338
339	2.00E-07	tRNA nucleotidyltransferase	Escherichia coli	QQECPE	nt. 2331-2540 of SEQ ID NO: 339
340	0	DNA gyrase, subunit A, type II topoisomerase	Salmonella typhimurium LT2	NP_461214.1	compliment of nt. 93-1799 of SEQ ID NO: 340
342	4.00E-12	tolA protein	Haemophilus influenzae	JC5212	nt. 980-1318 of SEQ ID NO: 342
344	1.00E-07	MODIFICATION METHYLASE HPHIA	Haemophilus parahaemolyticus	P50192	compliment of nt. 849-1034 of SEQ ID NO: 344
344	8.00E-05	ABC transporter protein 1	Leishmania major	AAF31030.1	compliment of nt. 17-205 of SEQ ID NO: 344
349	3.00E-44	conserved hypothetical protein	Neisseria meningitidis MC58	NP_273467.1	compliment of nt. 1397-1903 of SEQ ID NO: 349
349	8.00E-09	hypothetical protein	Pseudomonas aeruginosa	NP_252667.1	compliment of nt. 795-1121 of SEQ ID NO: 349
349	9.00E-10	conserved hypothetical secreted protein	Helicobacter pylori 26695	NP_207009.1	compliment of nt. 1319-1816 of SEQ ID NO: 349
349	2.00E-06	putative TPR repeat protein	Salmonella typhimurium LT2	NP_463149.1	compliment of nt. 2244-2558 of SEQ ID NO: 349
35	1.00E-23	type I restriction-modification system specificity determinant	Xylella fastidiosa 9a5c	NP_300003.1	compliment of nt. 29-388 of SEQ ID NO: 35
352	1.00E-116	putative peptidase	Escherichia coli K12	NP_416827.1	compliment of nt. 951-1640 of SEQ ID NO: 352
352	0	unknown	Pasteurella multocida	NP_245275.1	compliment of nt. 86-946 of SEQ ID NO: 352
354	5.00E-86	putative uronate isomerase	Salmonella typhimurium LT2	NP_462052.1	compliment of nt. 168-914 of SEQ ID NO: 354
356	1.00E-07	isomerase-like protein (DsbD) -	Escherichia coli	S57220	nt. 5-73 of SEQ ID NO: 356

358	1.00E-07	USG protein	Pediococcus pentosaceus	CAC16793.1	nt.534-1307 of SEQ ID NO: 358
358	0.005	HsdS protein	Escherichia coli	CAA10700.1	nt. 26-205 of SEQ ID NO: 358
361	1.00E-152	maltodextrin phosphorylase	Escherichia coli O157:H7 EDL933	NP_289957.1	compliment of nt. 77-922 of SEQ ID NO: 361
363	6.00E-06	BH2505~unknown conserved protein	Bacillus halodurans	NP_243371.1	nt. 554-844 of SEQ ID NO: 363
368	1.00E-12	H02F09.3.p	Caenorhabditis elegans	NP_508295.1	compliment of nt. 1069-1977 of SEQ ID NO: 368
368	6.00E-27	hypothetical glycine-rich protein	Mesorhizobium loti	NP_102360.1	compliment of nt. 1201-1986 of SEQ ID NO: 368
37	6.00E-09	putative ATP-binding component of a transport system	Escherichia coli K12	NP_415469.1	compliment of nt. 455-691 of SEQ ID NO: 37
372	7.00E-18	conserved hypothetical protein	Clostridium perfringens	BAB80319.1	compliment of nt. 1763-1924 of SEQ ID NO: 372
376	7.00E-24	putative bacteriophage protein	Salmonella enterica subsp. enterica serovar Typhi	NP_456379.1	compliment of nt. 158-808 of SEQ ID NO: 376
376	8.00E-10	hypothetical protein	Xylella fastidiosa 9a5c	NP_298882.1	compliment of nt. 1129-1671 of SEQ ID NO: 376
376	9.00E-06	lin1713	Listeria innocua	NP_471049.1	compliment of nt. 913-1557 of SEQ ID NO: 376
377	6.00E-05	Vng1732c	Halobacterium sp. NRC-1	NP_280487.1	nt. 2378-2587 of SEQ ID NO: 377
377	1.00E-11	INVASIN PRECURSOR (OUTER MEMBRANE ADHESIN)	Yersinia enterocolitica	P31489	compliment of nt. 127-345 of SEQ ID NO: 377
382	4.00E-16	unknown	Pasteurella multocida	NP_246871.1	compliment of nt. 967-1068 of SEQ ID NO: 382
383	4.00E-36	putative transposase	Streptomyces avermitilis	BAB69302.1	nt. 488-1162 of SEQ ID NO: 383
383	3.00E-58	recombinase	IncN plasmid R46	NP_511241.1	compliment of nt. 1-393 of SEQ ID NO: 383
383	4.00E-24	transposase	Escherichia coli	I69674	nt. 1294-1740 of SEQ ID NO: 383
383	0	tnpA	Yersinia enterocolitica	CAA73750.1	nt. 1782-2834 of SEQ ID NO: 383
385	2.00E-31	unknown	Pasteurella multocida	NP_246065.1	nt. 1515-1772 of SEQ ID NO: 385
386	5.00E-65	cydC [	Escherichia coli	AAA66172.1	compliment of nt. 3438-4115 of SEQ ID NO: 386
386	4.00E-33	ABC transporter, ATP-binding protein	Mesorhizobium loti	NP_105463.1	compliment of nt. 2569-3390 of SEQ ID NO: 386
388	3.00E-45	60 KDA INNER-MEMBRANE PROTEIN HOMOLOG	Coxiella burnetii	P45650	compliment of nt. 3211-3759 of SEQ ID NO: 388

WO 2004/078949

PCT/US2004/007001

44

390	4.00E-25	putative DNA-binding protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458175.1	nt. 1051-1416 of SEQ ID NO: 390
390	3.00E-13	transcriptional regulator	Bacillus halodurans	NP_241773.1	compliment of nt. 84-578 of SEQ ID NO: 390
390	3.00E-06	DNA translocase stage III sporulation prot homolog	Staphylococcus aureus subsp. aureus Mu50	NP_372265.1	compliment of nt. 620-871 of SEQ ID NO: 390
395	7.00E-31	ATPase, Cu ⁺⁺ transporting, beta polypeptide	Homo sapiens	NP_000044.1	compliment of nt. 615-1406 of SEQ ID NO: 395
397	3.00E-23	terminase large subunit	Bacteriophage HK620	NP_112076.1	compliment of nt. 2363-2725 of SEQ ID NO: 397
397	3.00E-16	hypothetical protein	Xylella fastidiosa 9a5c	NP_297824.1	compliment of nt. 1517-1744 of SEQ ID NO: 397
398	4.00E-67	orf32	Haemophilus phage HP2	NP_536839.1	compliment of nt. 1288-1866 of SEQ ID NO: 398
398	8.00E-24	putative cytoplasmic protein	Salmonella typhimurium LT2	NP_463063.1	compliment of nt. 798-1220 of SEQ ID NO: 398
398	2.00E-83	orf31	Haemophilus phage HP1	NP_043502.1	compliment of nt. 1881-2510 of SEQ ID NO: 398
399	1.00E-94	HEME/HEMOPEXI N-BINDING PROTEIN	Haemophilus influenzae N182	P45355	nt. 88-774 of SEQ ID NO: 399
401	3.00E-63	Sty SBLI	Salmonella enterica	CAA68058.1	nt. 1690-2742 of SEQ ID NO: 401
401	3.00E-06	RESTRICTION-MODIFICATION ENZYME SUBUNIT M3	Mycoplasma pulmonis	NP_325912.1	nt. 79-489 of SEQ ID NO: 401
402	2.00E-13	OPACITY PROTEIN OPA66 PRECURSOR	Neisseria gonorrhoeae	Q05033	compliment of nt. 2634-2915 of SEQ ID NO: 402
406	8.00E-13	type I restriction enzyme EcoR124II R	Neisseria meningitidis MC58	NP_273876.1	nt. 281-520 of SEQ ID NO: 406
407	6.00E-65	unknown	Pasteurella multocida	NP_246237.1	nt. 938-2450 of SEQ ID NO: 407
407	5.00E-99	PepE	Pasteurella multocida	NP_245391.1	nt. 1216-1917 of SEQ ID NO: 407
407	1.00E-16	Hemoglobin-haptoglobin binding protein A	Haemophilus influenzae Tn106	Q48153	nt. 1-141 of SEQ ID NO: 407
409	1.00E-106	hypothetical protein 1	Haemophilus influenzae	S27577	compliment of nt. 2524-3159 of SEQ ID NO: 409
411	4.00E-29	heme-repressible hemoglobin-binding protein	Haemophilus influenzae, type b, strain HI689	AAB46794.1	nt. 391-615 of SEQ ID NO: 411
411	0	Hemoglobin-haptoglobin binding protein A	Haemophilus influenzae Tn106	Q48153	nt. 651-3263 of SEQ ID NO: 411
412	2.00E-07	REGULATORY PROTEIN CRO (ANTIREPRESSOR)	bacteriophage 434	P03036	compliment of nt. 59-259 of SEQ ID NO: 412
412	4.00E-06	hypothetical protein	Bacteriophage P27	CAC83535.1	nt. 1436-1654 of SEQ ID NO: 412

413	8.00E-07	hypothetical protein	Deinococcus radiodurans	NP_294301.1	compliment of nt. 791-1012 of SEQ ID NO: 413
414	9.00E-65	conserved hypothetical protein	Vibrio cholerae	NP_230092.1	nt. 1696-2103 of SEQ ID NO: 414
414	3.00E-93	unknown	Pasteurella multocida	NP_246834.1	nt. 1777-2109 of SEQ ID NO: 414
416	2.00E-17	unknown	Pasteurella multocida	NP_246629.1	compliment of nt. 2565-2831 of SEQ ID NO: 416
416	4.00E-26	hypothetical protein o154	Escherichia coli	S30728	compliment of nt. 1928-2254 of SEQ ID NO: 416
416	3.00E-37	transport protein TatC	Pseudomonas aeruginosa	NP_253757.1	compliment of nt. 1494-2018 of of SEQ ID NO: 416
417	1.00E-66	weakly similar to methyltransferases	Listeria innocua	NP_471073.1	compliment of nt. 999-1928 of SEQ ID NO: 417
417	5.00E-05	DNA-BINDING PROTEIN RDGA	Pectobacterium carotovorum	Q47587	compliment of nt. 3526-4212 of SEQ ID NO: 417
417	2.00E-29	putative phage-related protein	Yersinia pestis	NP_407132.1	compliment of nt. 2546-2938 of SEQ ID NO: 417
417	3.00E-06	Adenine-specific DNA methylase	Thermoplasma acidophilum	NP_393798.1	compliment of nt. 826-1020 of SEQ ID NO: 417
43	9.00E-16	PcnB	Pasteurella multocida	NP_245801.1	nt. 511-870 of SEQ ID NO: 43
434	2.00E-97	beta' subunit of RNA polymerase	Nephroselmis olivacea	NP_050840.1	compliment of nt. 32-1534 of SEQ ID NO: 434
435	4.00E-52	MODIFICATION METHYLASE BEPI	Brevibacterium epidermidis	P10283	compliment of nt. 11-565 of SEQ ID NO: 435
435	4.00E-57	pentafunctional arom polypeptide (contains: 3-dehydroquinase synthase, 3-dehydroquinase dehydratase (3-dehydroquinase), shikimate 5-dehydrogenase, shikimate kinase, and eps synthase)	Saccharomyces cerevisiae	NP_010412.1	compliment of nt. 757-2064 of SEQ ID NO: 435
437	5.00E-70	dihydrofolate reductase	Haemophilus influenzae (clinical isolate R1042)	S52336	nt. 2393-2767 of SEQ ID NO: 437
438	1.00E-106	polyA polymerase	Vibrio cholerae	NP_230244.1	nt. 3-1124 of SEQ ID NO: 438
439	6.00E-60	Porphyrin biosynthetic protein	Salmonella enterica subsp. enterica serovar Typhi	NP_457816.1	nt. 2343-2783 of SEQ ID NO: 439
441	5.00E-73	RimM	Pasteurella multocida	NP_246234.1	compliment of nt. 151-441 of SEQ ID NO: 441
442	9.00E-31	LIPOPROTEIN NLPD	Salmonella typhimurium	P40827	compliment of nt. 3362-3520 of SEQ ID NO: 442

444	6.00E-24	glycine betaine transporter	Staphylococcus aureus subsp. aureus Mu50	NP_371872.1	compliment of nt. 2242-2514 of SEQ ID NO: 444
452	6.00E-28	unknown	Pasteurella multocida	NP_245616.1	compliment of nt. 533-883 of SEQ ID NO: 452
452	0	Type I restriction enzyme EcoPrrI M protein	Escherichia coli	Q47163	nt. 3291-4154 of SEQ ID NO: 452
452	2.00E-75	type I restriction enzyme M protein	Ureaplasma urealyticum	NP_077929.1	nt. 4156-4662 of SEQ ID NO: 452
455	9.00E-56	PROBABLE BACTERIOPHAGE PROTEIN	Ralstonia solanacearum	NP_520059.1	nt. 2028-2774 of SEQ ID NO: 455
455	2.00E-55	orf2; similar to major capsid protein precursor of phage P2 (gene N),	Haemophilus somnus	AAC45158.1	nt. 2864-3490 of SEQ ID NO: 455
455	1.00E-175	gpP	Enterobacteria phage P2	NP_046758.1	compliment of nt. 127-1812 of SEQ ID NO: 455
456	1.00E-38	hypothetical protein	Pseudomonas putida	NP_542872.1	compliment of nt. 1010-1282 of SEQ ID NO: 456
456	1.00E-172	hypothetical protein	Pseudomonas putida	NP_542873.1	compliment of nt. 1443-2006 of SEQ ID NO: 456
457	1.00E-116	hypothetical protein (galE 5' region) - Haemophilus influenzae	Haemophilus influenzae	S15287	compliment of nt. 62-961 of SEQ ID NO: 457
457	1.00E-134	dTDPglucose 4,6-dehydratase	Actinobacillus actinomycetemcomitans	T00102	nt. 2637-3656 of SEQ ID NO: 457
459	2.00E-10	RNA polymerase gamma-subunit	Synechocystis sp. PCC 6803	NP_441586.1	nt. 25-117 of SEQ ID NO: 459
461	9.00E-51	conserved hypothetical protein	Staphylococcus aureus subsp. aureus Mu50	NP_370593.1	nt. 4124-4624 of SEQ ID NO: 461
462	9.00E-06	NADH dehydrogenase	Burkholderia pseudomallei	AAG01016.1	nt. 703-828 of SEQ ID NO: 462
465	3.00E-41	GTP-binding protein Era	Synechocystis sp. PCC 6803	NP_441951.1	compliment of nt. 2470-2787 of SEQ ID NO: 465
466	1.00E-15	putative bacteriophage protein	Salmonella enterica subsp. enterica serovar Typhi	NP_455548.1	nt. 837-1478 of SEQ ID NO: 466
466	1.00E-90	orf31	Haemophilus phage HP1	NP_043502.1	nt. 2396-3199 of SEQ ID NO: 466
469	0	Hemoglobin and hemoglobin-haptoglobin binding protein C precursor	Haemophilus influenzae HI689	Q9X442	compliment of nt. 427-3459 of SEQ ID NO: 469
471	8.00E-05	transposase, putative	Neisseria meningitidis MC58	NP_274608.1	nt. 2957-3217 of SEQ ID NO: 471
472	6.00E-08	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458660.1	compliment of nt. 2881-3270 of SEQ ID NO: 472
472	5.00E-23	antirestriction protein	Mesorhizobium loti	NP_106707.1	nt. 4908-5324 of SEQ ID NO: 472

472	1.00E-75	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458661.1	compliment of nt. 1931-2776 of SEQ ID NO: 472
472	9.00E-72	hypothetical protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458662.1	compliment of nt. 544-1689 of SEQ ID NO: 472
475	3.00E-25	unknown	Pasteurella multocida	NP_244952.1	nt. 3207-3626 of SEQ ID NO: 475
476	8.00E-73	putative DNA-binding protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458175.1	compliment of nt. 3339-4310 of SEQ ID NO: 476
476	6.00E-47	anticodon nuclease	Neisseria meningitidis MC58	NP_273873.1	compliment of nt. 4397-4885 of SEQ ID NO: 476
478	3.00E-06	methionin synthase-like enzyme	Arabidopsis thaliana	CAB38313.1	compliment of nt. 3554-3679 of SEQ ID NO: 478
478	3.00E-05	unknown	Pasteurella multocida	NP_245444.1	compliment of nt. 164-250 of SEQ ID NO: 478
479	1.00E-18	conserved hypothetical protein	Xylella fastidiosa 9a5c	NP_298841.1	nt. 2302-2658 of SEQ ID NO: 479
48	3.00E-19	Dca	Neisseria gonorrhoeae	AAF12796.1	compliment of nt. 225-746 of SEQ ID NO: 48
482	1.00E-06	hypothetical protein	Neisseria meningitidis MC58	NP_275122.1	nt. 2055-2189 of SEQ ID NO: 482
482	9.00E-28	conserved hypothetical protein	Neisseria meningitidis MC58	NP_274383.1	nt. 1689-1898 of SEQ ID NO: 482
487	5.00E-75	conserved hypothetical protein	Neisseria meningitidis Z2491	NP_284304.1	nt. 2541-2978 of SEQ ID NO: 487
488	2.00E-64	unknown	Pasteurella multocida	NP_246617.1	nt. 2983-3540 of SEQ ID NO: 488
488	8.00E-93	1-deoxy-D-xylulose 5-phosphate reductoisomerase	Zymomonas mobilis	AAD29659.1	nt. 1344-1880 of SEQ ID NO: 488
491	5.00E-51	rubredoxin oxidoreductase homolog	Clostridium acetobutylicum	AAB50346.1	compliment of nt. 1690-2439 of SEQ ID NO: 491
492	1.00E-27	phosphotransferase system enzyme IIA-like protein	Staphylococcus aureus	AAK83253.1	compliment of nt. 755-970 of SEQ ID NO: 492
493	2.00E-84	unknown	Actinobacillus actinomycetemcomitans	AAC70895.1	nt. 3333-3935 of SEQ ID NO: 493
493	4.00E-49	unknown	Helicobacter pylori J99	NP_223898.1	nt. 3345-4010 of SEQ ID NO: 493
493	9.00E-31	transcriptional factor MdcH	Acinetobacter calcoaceticus	AAF20290.1	nt. 1885-2793 of SEQ ID NO: 493
493	6.00E-30	HimA	Pasteurella multocida	NP_245565.1	nt. 1129-1260 of SEQ ID NO: 493
494	4.00E-85	putative prophage integrase	Yersinia pestis	NP_404712.1	nt. 900-2099 of SEQ ID NO: 494
494	4.00E-63	DNA methyltransferase	Xylella fastidiosa 9a5c	NP_299063.1	compliment of nt. 5544-6170 of SEQ ID NO: 494
494	6.00E-19	MODIFICATION METHYLASE SCRFIA	Lactococcus lactis subsp. cremoris	P34877	compliment of nt. 5019-6113 of SEQ ID NO: 494

WO 2004/078949

PCT/US2004/007001

48

497	0	transferrin-binding protein 1	Haemophilus influenzae (strain PAK 12085)	S70906	nt. 3251-4999 of SEQ ID NO: 497
50	5.00E-07	AcpP	Pasteurella multocida	NP_246856.1	nt. 2-136 of SEQ ID NO: 50
501	7.00E-50	conserved hypothetical protein	Vibrio cholerae	NP_231403.1	compliment of nt. 3649-4872 of SEQ ID NO: 501
501	0	type I restriction enzyme HsdR, putative	Vibrio cholerae	NP_231400.1	compliment of nt. 1551-3440 of SEQ ID NO: 501
501	4.00E-13	ATP-dependent DNA helicase RecG-related protein	Deinococcus radiodurans	NP_295921.1	compliment of nt. 5317-5844 of SEQ ID NO: 501
501	5.00E-11	conserved hypothetical	Ureaplasma urealyticum	NP_077868.1	compliment of nt. 5098-5769 of SEQ ID NO: 501
504	2.00E-44	OUTER MEMBRANE PROTEIN P2 PRECURSOR (OMP P2)	Haemophilus influenzae AG30010	Q48218	compliment of nt. 4681-5019 of SEQ ID NO: 504
507	0	SpoT	Pasteurella multocida	NP_245857.1	compliment of nt. 3685-5316 of SEQ ID NO: 507
51	6.00E-87	glucosamine--fructose-6-phosphate aminotransferase (isomerizing)	Vibrio cholerae	NP_230141.1	nt. 30-470 of SEQ ID NO: 51
512	2.00E-28	dipeptide transport system permease protein	Yersinia pestis	NP_407439.1	compliment of nt. 1095-1580 of SEQ ID NO: 512
512	3.00E-82	SapC	Pasteurella multocida	NP_245850.1	compliment of nt. 730-1095 of SEQ ID NO: 512
514	9.00E-06	putative integral membrane protein	Campylobacter jejuni	NP_281236.1	compliment of nt. 577-684 of SEQ ID NO: 514
514	3.00E-11	orf, hypothetical protein	Escherichia coli O157:H7 EDL933	NP_286004.1	compliment of nt. 449-568 of SEQ ID NO: 514
518	0	putative inner membrane trans-acylase protein	Neisseria meningitidis Z2491	NP_284893.1	nt. 92-1927 of SEQ ID NO: 518
519	4.00E-30	hypothetical protein	Mesorhizobium loti	NP_108196.1	compliment of nt. 2221-3159 of SEQ ID NO: 519
519	2.00E-12	conserved hypothetical protein	Listeria innocua	NP_471067.1	compliment of nt. 3994-5241 of SEQ ID NO: 519
519	6.00E-20	hypothetical protein	Mesorhizobium loti	NP_108198.1	compliment of nt. 707-1552 of SEQ ID NO: 519
519	4.00E-26	putative bacteriophage protein	Salmonella enterica subsp. enterica serovar Typhi	NP_455526.1	compliment of nt. 3982-5163 of SEQ ID NO: 519
52	3.00E-94	OUTER MEMBRANE PROTEIN P2 PRECURSOR (OMP P2)	Haemophilus influenzae	Q48218	nt. 45-788 of SEQ ID NO: 52

520	0	excision nuclease subunit A	Escherichia coli K12	NP_418482.1	compliment of nt. 6309-7745 of SEQ ID NO: 520
521	5.00E-08	zinc/manganese ABC transporter substrate binding protein	Rickettsia conorii	NP_359651.1	nt. 2236-2652 of SEQ ID NO: 521
521	1.00E-140	unknown	Pasteurella multocida	NP_245865.1	nt. 338-1390 of SEQ ID NO: 521
521	1.00E-86	ORF_f432	Escherichia coli	AAB40463.1	nt. 203-1390 of SEQ ID NO: 521
522	3.00E-22	unknown	Pasteurella multocida	NP_246093.1	nt. 670-885 of SEQ ID NO: 522
526	5.00E-33	exodeoxyribonuclease V alpha chain	Yersinia pestis	NP_404635.1	nt. 5582-6202 of SEQ ID NO: 526
526	1.00E-62	exodeoxyribonuclease V, 67 kDa subunit	Vibrio cholerae	NP_231950.1	nt. 5675-6193 of SEQ ID NO: 526
527	1.00E-147	unknown	Pasteurella multocida	NP_245980.1	nt. 4283-5203 of SEQ ID NO: 527
527	0	Mfd	Pasteurella multocida	NP_245978.1	nt. 7545-8759 of SEQ ID NO: 527
527	0	transcription-repair coupling factor (TrcF)	Salmonella enterica subsp. enterica serovar Typhi	NP_455708.1	nt. 7611-8762 of SEQ ID NO: 527
527	0	PROBABLE TRANSCRIPTION-REPAIR COUPLING FACTOR PROTEIN	Ralstonia solanacearum	NP_519763.1	nt. 7611-8870 of SEQ ID NO: 527
528	1.00E-48	undecaprenyl pyrophosphate synthetase	Chlamydia muridarum	NP_297109.1	nt. 2918-3712 of SEQ ID NO: 528
528	0	leucyl-tRNA synthetase	Vibrio cholerae	NP_230603.1	compliment of nt. 180-2822 of SEQ ID NO: 528
529	1.00E-104	DNA PRIMASE	Legionella pneumophila	P71481	compliment of nt. 3316-3960 of SEQ ID NO: 529
534	9.00E-29	putative integrase	Salmonella typhimurium LT2	NP_461690.1	nt. 4668-5009 of SEQ ID NO: 534
534	6.00E-18	hypothetical protein NMA0153	Neisseria meningitidis Z2491	NP_283002.1	compliment of nt. 5933-6337 of SEQ ID NO: 534
534	2.00E-23	hypothetical protein	Deinococcus radiodurans	NP_294868.1	nt. 6908-7654 of SEQ ID NO: 534
534	1.00E-88	prophage CP4-57 integrase	Escherichia coli K12	NP_417111.1	nt. 5057-5875 of SEQ ID NO: 534
535	1.00E-115	phosphate acetyltransferase	Buchnera sp. APS	NP_240007.1	nt. 3385-4596 of SEQ ID NO: 535
536	3.00E-35	cobalt membrane transport protein CbiQ	Actinobacillus pleuropneumoniae	AAD49727.1	compliment of nt. 3531-4136 of SEQ ID NO: 536
536	6.00E-37	unknown	Pasteurella multocida	NP_245305.1	compliment of nt. 6478-6921 of SEQ ID NO: 536
539	2.00E-26	Orf122	Chlorobium tepidum	AAG12204.1	compliment of nt. 1778-2008 of SEQ ID NO: 539
540	1.00E-77	heat shock protein HtpX	Neisseria meningitidis MC58	NP_273864.1	compliment of nt. 2567-3481 of SEQ ID NO: 540
541	0	IleS	Pasteurella multocida	NP_246601.1	nt. 3167-4549 of SEQ ID NO: 541

545	2.00E-09	DNA-BINDING PROTEIN RDGB	Pectobacterium carotovorum	Q47588	nt. 3816-3977 of SEQ ID NO: 545
545	2.00E-11	putative transposase	Sinorhizobium meliloti	NP_437741.1	compliment of nt. 2786-3019 of SEQ ID NO: 544
545	2.00E-07	Hypothetical 42.5 kd protein in thrW- argF intergenic region	Escherichia coli	BAA77933.1	compliment of nt. 2614-2811 of SEQ ID NO: 545
545	4.00E-18	putative IS element transposase	Salmonella enterica subsp. enterica serovar Typhi	NP_454711.1	nt. 1955-2230 of SEQ ID NO: 545
546	0	HEME/HEMOPEXI N-BINDING PROTEIN	Haemophilus influenzae	P45354	nt. 5551-7809 of SEQ ID NO: 546
546	0	HEME/HEMOPEXI N UTILIZATION PROTEIN B	Haemophilus influenzae	P45356	nt. 3842-5536 of SEQ ID NO: 546
546	0	HEME/HEMOPEXI N UTILIZATION PROTEIN C	Haemophilus influenzae	P45357	nt. 1638-3176 of SEQ ID NO: 546
546	2.00E-12	HasR	Pasteurella multocida	NP_246561.1	nt. 3149-3763 of SEQ ID NO: 546
549	0	unknown	Pasteurella multocida	NP_246821.1	nt. 2526-3512 of SEQ ID NO: 549
549	1.00E-121	putative membrane protein	Yersinia pestis	NP_404859.1	nt. 605-1108 of SEQ ID NO: 549
549	0	unknown	Pasteurella multocida	NP_246822.1	nt. 1122-1664 of SEQ ID NO: 549
551	1.00E-157	type I restriction- modification system endonuclease	Xylella fastidiosa 9a5c	NP_300016.1	compliment of nt. 7396-8322 of SEQ ID NO: 551
552	1.00E-100	valyl-tRNA synthetase	Deinococcus radiodurans	NP_293872.1	compliment of nt. 6691-8688 of SEQ ID NO: 552
552	0	VALYL-TRNA SYNTHETASE	Haemophilus parainfluenzae	P36432	compliment of nt. 5850-6647 of SEQ ID NO: 552
553	0	DNA-directed RNA polymerase, beta subunit	Vibrio cholerae	NP_229982.1	nt. 2668-6699 of SEQ ID NO: 553
554	0	iron utilization protein B	Haemophilus influenzae	T10887	nt. 991-2508 of SEQ ID NO: 554
559	1.00E-100	PREPROTEIN TRANSLOCASE SECA SUBUNIT	Bacillus firmus	P96313	nt. 3420-4472 of SEQ ID NO: 559
56	2.00E-23	RpL30	Pasteurella multocida	NP_246336.1	compliment of nt. 656-832 of SEQ ID NO: 56
56	9.00E-13	RpS5	Pasteurella multocida	NP_246337.1	compliment of nt. 843-1064 of SEQ ID NO: 56
560	1.00E-157	Na ⁺ /H ⁺ antiporter	Vibrio cholerae	NP_231535.1	2 compliment of nt. 279-2989 of SEQ ID NO: 560
562	1.00E-72	putative biotin sulfoxide reductase 2	Yersinia pestis	NP_404419.1	nt. 7862-8878 of SEQ ID NO: 562
562	1.00E-125	restriction modification system-R protein	Neisseria meningitidis	CAA09003.1	nt. 2-985 of SEQ ID NO: 562
563	0	IMMUNOGLOBULI N A1 PROTEASE	Haemophilus influenzae HK715	P45384	compliment of nt. 4127-9508 of SEQ ID NO: 563

563	0	3-ISOPROPYLMALATE DEHYDRATASE (IPMI)	Schizosaccharomyces pombe	O14289	nt. 1980-3983 of SEQ ID NO: 563
564	2.00E-79	orf32	Haemophilus phage HP2	NP_536839.1	nt. 6241-6831 of SEQ ID NO: 564
564	7.00E-33	probable variable tail fibre protein	Salmonella enterica subsp. enterica serovar Typhi	NP_457882.1	nt. 3707-4177 of SEQ ID NO: 564
564	2.00E-14	M protein	Enterobacteria phage 186	NP_052264.1	nt. 1905-2213 of SEQ ID NO: 564
564	4.00E-44	similar to tail fiber protein (gpH) in phage P2	Salmonella typhimurium LT2, Fels-2 prophage	NP_461635.1	nt. 3171-3692 of SEQ ID NO: 564
564	2.00E-85	gpJ	Enterobacteria phage P2	NP_046773.1	nt. 2267-3166 of SEQ ID NO: 564
564	1.00E-24	hypothetical protein	Neisseria meningitidis Z2491	NP_284534.1	nt. 6852-7334 of SEQ ID NO: 564
564	4.00E-26	gpV	Enterobacteria phage P2	NP_046771.1	nt. 1337-1912 of SEQ ID NO: 564
564	2.00E-47	similar to [SwissProt P44255	Escherichia coli	BAA16182.1	nt. 11383-11961 of SEQ ID NO: 564
564	2.00E-51	hypothetical protein NMA1315	Neisseria meningitidis Z2491	NP_284066.1	nt. 10452-11180 of SEQ ID NO: 564
564	0	orf31	Haemophilus phage HP1	NP_043502.1	nt. 4160-6226 of SEQ ID NO: 564
564	2.00E-09	rep	Haemophilus phage HP2	NP_536816.1	compliment of nt. 9986-10234 of SEQ ID NO: 564
565	2.00E-57	resolvase/integrase-like protein	Haemophilus influenzae biotype aegyptius	AAL47097.1	nt. 11885-12445 of SEQ ID NO: 565
565	1.00E-93	integrase	Actinobacillus actinomycetemcomitans	AAC70901.1	compliment of nt. 4118-4900 of SEQ ID NO: 565
565	6.00E-35	probable phage integrase	Salmonella enterica subsp. enterica serovar Typhi	NP_458745.1	compliment of nt. 4148-4990 of SEQ ID NO: 565
565	1.00E-107	hypothetical protein	Xylella fastidiosa 9a5c	NP_299042.1	compliment of nt. 5066-6817 of SEQ ID NO: 565
566	1.00E-126	hypothetical protein (galE 5' region) -	Haemophilus influenzae	S15287	compliment of nt. 10726-11607 of SEQ ID NO: 566
567	0	unknown	Pasteurella multocida	NP_246387.1	nt. 5343-7688 of SEQ ID NO: 567
568	1.00E-151	multidrug resistance membrane translocase	Escherichia coli O157:H7	NP_311575.1	nt. 6-1403 of SEQ ID NO: 568
568	1.00E-141	YhbX/YhjW/YijP/YjdB family protein	Neisseria meningitidis MC58	NP_275002.1	compliment of nt. 11213-12634 of SEQ ID NO: 568

570	1.00E-180	hypothetical protein 3 (ksgA-lic2B intergenic region)	Haemophilus influenzae (strain RM7004)	S71024	compliment of nt. 12845-13720 of SEQ ID NO: 570
571	0	glycerophosphodie ster phosphodiesterase	Haemophilus influenzae (isolate 772)	A43576	nt. 1656-2693 of SEQ ID NO: 571
571	1.00E-137	outer membrane protein P4 precursor - Haemophilus influenzae	Haemophilus influenzae	A43604	nt. 6145-6909 of SEQ ID NO: 571
571	2.00E-72	CG8298 gene product [alt 1]	Drosophila melanogaster	AAF58597.1	nt. 3813-5339 of SEQ ID NO: 571
572	1.00E-40	hypothetical protein TC0130	Chlamydia muridarum (strain Nigg)	G81737	nt. 3734-4099 of SEQ ID NO: 572
572	5.00E-10	hypothetical protein	Pyrococcus horikoshii	NP_142215.1	nt. 4472-4888 of SEQ ID NO: 572
572	3.00E-11	109aa long hypothetical protein	Sulfolobus tokodaii	NP_377117.1	nt. 7303-7470 of SEQ ID NO: 572
572	8.00E-43	hypothetical protein	Chlamydophila pneumoniae AR39	NP_445524.1	nt. 4289-4618 of SEQ ID NO: 572
572	9.00E-08	CDH1-D	Gallus gallus	AAL31950.1	nt. 7183-7521 of SEQ ID NO: 572
575	1.00E-173	topoisomerase B	Salmonella enterica subsp. enterica serovar Typhi	NP_458624.1	nt. 18980-20923 of SEQ ID NO: 575
575	1.00E-100	DNA helicase	Salmonella enterica subsp. enterica serovar Typhi	NP_458617.1	nt. 10399-11706 of SEQ ID NO: 575
65	2.00E-53	Sufl	Pasteurella multocida	NP_245041.1	nt. 3-821 of SEQ ID NO: 65
67	4.00E-39	putative MFS family transport protein (1st mdule)	Salmonella typhimurium LT2	NP_462786.1	compliment of nt. 125-1033 of SEQ ID NO: 67
7	4.00E-29	putative membrane protein	Salmonella enterica subsp. enterica serovar Typhi	NP_458664.1	compliment of nt. 2-559 of SEQ ID NO: 7
72	2.00E-51	serine transporter	Vibrio cholerae	NP_230946.1	nt. 18-803 of SEQ ID NO: 72
74	3.00E-90	hypothetical 21.8K protein (in locus involved in transformation) -	Haemophilus influenzae	JH0436	compliment of nt. 248-766 of SEQ ID NO: 74
77	2.00E-18	RecX protein	Legionella pneumophila	CAC33485.1	nt. 480-920 of SEQ ID NO: 77
82	4.00E-95	unknown	Pasteurella multocida	NP_246414.1	nt. 128-955 of SEQ ID NO: 82
83	2.00E-66	unknown	Pasteurella multocida	NP_246777.1	nt. 5-556 of SEQ ID NO: 83
83	6.00E-33	CTP SYNTHASE	Helicobacter pylori J99	NP_223042.1	compliment of nt. 1027-1338 of SEQ ID NO: 83.
83	4.00E-34	CTP synthase	Campylobacter jejuni	NP_281249.1	compliment of nt. 1024-1275 of SEQ ID NO: 83
84	1.00E-16	REPRESSOR PROTEIN CI	Bacteriophage phi-80	P14819	nt. 823-1233 of SEQ ID NO: 84
84	2.00E-05	orf, hypothetical protein	Escherichia coli K12	NP_415875.1	compliment of nt. 533-700 of SEQ ID NO: 84

WO 2004/078949

PCT/US2004/007001

53

84	4.00E-11	orf33	bacteriophage phi CTX	NP_490633.1	compliment of nt. 32-466 of SEQ ID NO: 84
85	3.00E-42	SpoT	Pasteurella multocida	NP_245857.1	nt. 899-1261 of SEQ ID NO: 85
90	1.00E-103	putative methylase	Bacteriophage Tuc2009	NP_108695.1	compliment of nt. 478-1206 of SEQ ID NO: 90
90	4.00E-11	probable adenine specific DNA methyltransferase	Thermoplasma acidophilum	NP_394624.1	compliment of nt. 397-1140 of SEQ ID NO: 90

TABLE 3B

Hit Identity	Full Length Nucleotide Sequence	Amino Acid Sequence	Location in Contig	Homology to Genbank Protein
CpdB	SEQ ID NO: 686	SEQ ID NO: 687	nt. 38041-36068 of SEQ ID NO: 681 (contig 14)	NP_246953.1
putative membrane protein	SEQ ID NO: 688	SEQ ID NO: 689	nt. 906601-908094 of SEQ ID NO: 685 (contig 18)	NP_458664.1
GTP-binding protein TypA/BipA	SEQ ID NO: 690	SEQ ID NO: 691	nt. 42557-40995 of SEQ ID NO: 683 (contig 16)	NP_240245.1
outer membrane protein A	SEQ ID NO: 692	SEQ ID NO: 693	nt. 7000420-704187 of SEQ ID NO: 685 (contig 18)	T30852
vacB protein	SEQ ID NO: 694	SEQ ID NO: 695	nt. 39184-36836 of SEQ ID NO: 683 (contig 16)	NP_240369.1
putative ABC transport system permease protein [	SEQ ID NO: 696	SEQ ID NO: 697	nt. 59155-58370 of SEQ ID NO: 685 (contig 18)	NP_282774.1
putative exported protein	SEQ ID NO: 698	SEQ ID NO: 699	nt. 901142-902542 of SEQ ID NO: 685 (contig 18)	NP_458655.1
ImpA	SEQ ID NO: 700	SEQ ID NO: 701	nt. 348187-347747 of SEQ ID NO: 685 (contig 18)	NP_245829.1
TsaA	SEQ ID NO: 702	SEQ ID NO: 703	nt. 74941-75548 of SEQ ID NO: 684 (contig 17)	NP_245732.1
PROBABLE TRANSPORT TRANSMEMBRANE PROTEIN	SEQ ID NO: 704 SEQ ID NO: 706	SEQ ID NO: 705 SEQ ID NO: 707	nt. 74436-75176 of SEQ ID NO: 685 (contig 18) nt. 75160-75660 of SEQ ID NO: 685 (contig 18)	NP_522358.1
possible exported protein	SEQ ID NO: 708	SEQ ID NO: 709	nt. 899618-900262 of SEQ ID NO: 685 (contig 18)	NP_458653.1
LICA PROTEIN	SEQ ID NO: 710	SEQ ID NO: 711	nt. 356917-355958 of SEQ ID NO: 685 (contig 18)	P14181
HEME-BINDING PROTEIN A	SEQ ID NO: 712	SEQ ID NO: 713	NT. 26114-27739 of SEQ ID NO: 683 (contig 16)	P33950

similar to BASEMENT MEMBRANE- SPECIFIC HEPARAN SULFATE PROTEOGLYCAN CORE PROTEIN PRECURSOR (HSPG)	SEQ ID NO: 714	SEQ ID NO: 715	nt. 311610-312683 of SEQ ID NO: 685 (contig 18)	XP_068727.1
CzcD	SEQ ID NO: 716	SEQ ID NO: 717	nt. 34865-35542 of SEQ ID NO: 681 (contig 14)	NP_246276.1
conserved hypothetical protein	SEQ ID NO: 718	SEQ ID NO: 719	nt. 194993-193977 of SEQ ID NO: 685 (contig 18)	NP_274972.1
secretion protein SecD	SEQ ID NO: 720	SEQ ID NO: 721	nt. 203707-201857 of SEQ ID NO: 683 (contig 17)	NP_252510.1
ABC transporter protein 1	SEQ ID NO: 722	SEQ ID NO: 723	nt. 3943-5859 of SEQ ID NO: 681 (contig 14)	AAF31030.1
conserved hypothetical protein	SEQ ID NO: 724 SEQ ID NO: 726 SEQ ID NO: 728	SEQ ID NO: 725 SEQ ID NO: 727 SEQ ID NO: 729	nt. 331090-331749 of SEQ ID NO: 685 (contig 18) nt. 331938-332492 of SEQ ID NO: 685 (contig 18) nt. 332681-33232 of SEQ ID NO: 685 (contig 18)	NP_273467.1
INVASIN PRECURSOR (OUTER MEMBRANE ADHESIN)	SEQ ID NO: 730	SEQ ID NO: 731	nt. 416757-417020 of SEQ ID NO: 685 (contig 18)	P31489
HEME/HEMOPEXI N-BINDING PROTEIN	SEQ ID NO: 732	SEQ ID NO: 733	nt. 229430-232195 of SEQ ID NO: 384 (contig 17)	P45355
OPACITY PROTEIN OPA66 PRECURSOR	SEQ ID NO: 734	SEQ ID NO: 735	nt. 375592-375879 of SEQ ID NO: 384 (contig 17)	Q05033
Hemoglobin- haptoglobin binding protein A	SEQ ID NO: 736	SEQ ID NO: 737	nt. 45709-42566 of SEQ ID NO: 681 (contig 14)	Q48153
transport protein TatC	SEQ ID NO: 738	SEQ ID NO: 739	nt. 134452-135222 of SEQ ID NO: 384 (contig 17)	NP_253757.1
LIPOPROTEIN NLPD	SEQ ID NO: 740	SEQ ID NO: 741	nt. 18895-20112 of SEQ ID NO: 682 (contig 15)	P40827
Hemoglobin and hemoglobin- haptoglobin binding protein C precursor	SEQ ID NO: 742	SEQ ID NO: 743	nt. 34181-31041 of SEQ ID NO: 682 (contig 15)	Q9X442
HimA	SEQ ID NO: 744	SEQ ID NO: 745	nt. 382795-383085 of SEQ ID NO: 685 (contig 18)	NP_245565.1
transferrin-binding protein 1	SEQ ID NO: 746	SEQ ID NO: 747	nt. 178537-175799 of SEQ ID NO: 683 (contig 16)	S70906
SapC	SEQ ID NO: 748	SEQ ID NO: 749	nt. 197754-196867 of SEQ ID NO: 685 (contig 18)	NP_245850.1

heat shock protein HtpX	SEQ ID NO: 750	SEQ ID NO: 751	nt. 40414-41265 of SEQ ID NO: 682 (contig 15)	NP_273864.1
HEME/HEMOPEXIN-BINDING PROTEIN	SEQ ID NO: 752	SEQ ID NO: 753	nt. 229430-232195 of SEQ ID NO: 684 (contig 17)	P45354
HEME/HEMOPEXIN UTILIZATION PROTEIN B	SEQ ID NO: 754	SEQ ID NO: 755	nt. 227721-229418 of SEQ ID NO: 684 (contig 17)	P45356
HEME/HEMOPEXIN UTILIZATION PROTEIN C	SEQ ID NO: 756	SEQ ID NO: 757	nt. 225516-227645 of SEQ ID NO: 684 (contig 17)	P45357 NP_246561.1
iron utilization protein B	SEQ ID NO: 758	SEQ ID NO: 759	nt. 32076-33611 of SEQ ID NO: 684 (contig 17)	T10887
PREPROTEIN TRANSLOCASE SECA SUBUNIT	SEQ ID NO: 760	SEQ ID NO: 761	nt. 82314-84785 of SEQ ID NO: 683 (contig 16)	P96313
IMMUNOGLOBULIN A1 PROTEASE	SEQ ID NO: 762	SEQ ID NO: 763	nt. 171647-166263 of SEQ ID NO: 683 (contig 16)	P45384
multidrug resistance membrane translocase	SEQ ID NO: 764	SEQ ID NO: 765	nt. 74524-72992 of SEQ ID NO: 683 (contig 16)	NP_311575.1
YhbX/YhjW/YijP/YjdB family protein	SEQ ID NO: 766	SEQ ID NO: 767	nt. 61734-63200 of SEQ ID NO: 683 (contig 16)	NP_275002.1
putative membrane protein	SEQ ID NO: 768	SEQ ID NO: 769	nt. 906601-908094 of SEQ ID NO: 685 (contig 18)	NP_458664.1
putative membrane protein	SEQ ID NO: 770	SEQ ID NO: 771	nt. 16185-17942 of SEQ ID NO: 683 (contig)	NP_404859.1

Example 3

Construction of the NTHi Promoter Trap Library

- 5 To identify potential virulence determinants of NTHi, bacterial gene expression was monitored by differential fluorescence induction (DFI) during early disease progression in one specific anatomical niche of a chinchilla model of NTHi-induced otitis media (OM). Genomic DNA fragments from NTHi strain 86-028NP were cloned upstream of the promoterless *gfpmut3* gene using a promoter trap library.
- 10 Plasmid pGZRS39A, a derivative of pGZRS-1 isolated from *Actinobacillus pleuropneumoniae*, is an *A. pleuropneumoniae*-*Escherichia coli* shuttle vector. This plasmid contains the origin of replication from *A. pleuropneumoniae*, the *lacZa* gene from pUC19 and the kanamycin resistance gene from Tn903. (West *et al.*, *Genes*, 160: 81-86, 1995).
- 15 The promoter trap vector was constructed by cloning the GTP mutant *gfpmut3* gene, as a *Bam*HI to *Eco*RI fragment into pGZRS-39A to form pRSM2167. This mutant GTP gene contains two amino acid changes, S65G and S72A, that

enhance fluorescence emission when excited at 488 nm. This mutant also has high solubility and fast kinetics of chromophore formation (Cormack *et al.*, *Gene*, 173: 33-38, 1996). This plasmid was transformed by electroporation into NTHi strain 86-028NP, generating the parent-plasmid strain 86-028NP/pRSM2169.

5 Random genomic DNA fragments (described in Example 1) were prepared for ligation into the promoter probe vector. Genomic DNA was isolated from strain 86-028NP using the Puregene DNA isolation kit (Gentra Systems, Minneapolis, MN) according to the manufacturer's protocol. Due to restriction barriers, it was necessary to isolate the plasmid DNA and use this for the library
10 generation. The isolated DNA was partially digested with *Sau3AI* (NEB, Beverly, MA; 0.25 units/ μ g DNA) for 1 hour at 37°C, separated by gel electrophoresis and DNA fragments 0.5-1.5 kb in size were recovered using the Qiagen gel extraction kit. For vector preparation, pRSM2167 was isolated from an overnight culture using the Wizard Plus Maxiprep DNA purification system (Promega, Madison WI) according
15 to the manufacturer's protocol.

Plasmid DNA was linearized by *Bam*HI digestion and 5' phosphate groups removed by treatment with calf intestinal alkaline phosphatase (CIAP; GibcoBRL Life Technologies). Genomic DNA fragments were ligated with the linearized, phosphatase-treated vector and electroporated into competent NTHi strain
20 86-028NP prepared for electroporation according to a modified protocol (Mitchell *et al.*, *Nucleic Acids Res.*, 19: 3625-3628, 1991). When plasmid DNA was electroporated back into NTHi strain 86-028NP, transformation efficiency was improved by one-thousand fold. Briefly, cells were grown to an $OD_{600}=0.3$ in sBHI (brain heart infusion) broth at 37°C, 220 rpm. Cells were chilled on ice for 30
25 minutes and subsequently washed with an equal volume of 0.5xSG (1xSG: 15% glycerol, 272 mM sucrose) at 4°C. Washes were repeated a total of three times. Subsequently, the cells were diluted in 1xSG to a 100x concentrated volume. The cells were electroporated using the BioRad Gene Pulser II set at 200 ohms, 2.5 kV and 25 μ F and then diluted in 1 ml prewarmed sBHI, incubated for 2 hours at 37°C,
30 5% CO₂ and plated on chocolate agar for overnight growth of transformants.

Transformants were selected and frozen in pools of 1000 clones in skim milk containing 20% glycerol (vol/vol). A 68,000 member *gfp* promoter probe library was generated. Using the probability calculation of Clarke and Carbon (*Cell*,

9: 91-99, 1976), to achieve a 99% probability of having a given DNA sequence represented in a library of 300 bp fragments of strain 86-028NP DNA (1.8×10^6 bp/genome), a library of 27,629 clones was needed. Therefore the present library represents 2.5 fold coverage of the 86-028NP genome.

5 In order to assess the quality of the library, fifty clones were selected at random, grown overnight on chocolate agar and the plasmids were isolated and insert DNA sequenced. A majority (64%) of the selected clones had insert sizes ranging between 200 and 500 bp while 32% exceeded 500 bp. The majority of inserts showed homology to unique *H. influenzae* strain Rd open reading frames (ORFs), and 15
10 clones had sequence unique to strain 86-028NP DNA. Of those clones with homology to strain Rd, 60% were in the correct orientation, 36% of which contained sequence upstream an ORF. Although a majority of clones had an insert size less than 500 bp, no correlation was found between small insert size and increased GFP expression. In fact four clones exhibited slight to moderate fluorescence *in vitro*, 3 of
15 which had insert sizes between 200-500 base pairs and one had an insert that was greater than 700 base pairs.

A fraction of the library (approximately 1000 clones) was grown on chocolate agar, harvested in PBS and analyzed by flow cytometry for GFP fluorescence. Compared to strain 86-028NP/pRSM2169 that contains the promoter
20 trap vector without insert DNA, the pool of library clones displays an increased fluorescence intensity. Thus, the library contains clones with promoters at varying levels of activity.

Example 4

Analysis of 86-028NP Derivatives Expressing GFP

25 In order to establish the FACS parameters necessary to identify and sort *gfp*-expressing bacteria, a panel of isolates demonstrating varying levels of *gfp* expression was utilized. Background fluorescence was assessed using strain 86-028NP/pRSM2169 (negative control), therefore any observed fluorescence would be due to the *lacZ* promoter driving *gfp* expression. However, this strain does not
30 produce detectable levels of GFP and in fact, does not demonstrate increased fluorescence when compared to the parent strain 86-028NP. A high-level *gfp*-expressing isolate was generated by cloning a 500 bp fragment containing the strong promoter for outer membrane protein P2 expression into *SalI*-*BamHI* digested

pRSM2167. This plasmid was transformed into 86-028NP by electroporation, generating the high-level *gfp* expressing strain 86-028NP/pRSM2211 (highly fluorescent control). This strain demonstrated an approximate 100 fold increase in GFP fluorescence compared to strain 86-028NP/pRSM2169. An intermediate
5 fluorescent derivative clone, 86-028NP/pKMM4B5 (intermediate fluorescent control), was isolated by FACS analysis and used both in preliminary experiments and as a control for cell sorting. The DNA fragment containing a promoter driving *gfp* expression *in vitro* is unique to strain 86-028NP, having no known homology to DNA of other organisms. This clone exhibits an approximate 10 fold increase in
10 fluorescence compared to strain 86-028NP/pRSM2169.

The control strains were resuspended from growth on chocolate agar and labeled with cross-reactive Phycoprobe R-PE anti-human IgG (H+L) antibody (10 µg/ml in 100 µl PBS; Biomeda Corp) for 30 minutes at 4° C. Following three successive washes to remove unbound antibody, bacteria were resuspended in 300 µl
15 DPBS for FACS analysis. These control preparations were used to set the appropriate size and fluorescence gates using a Coulter Epics Elite flow cytometer (Coulter Corp.) equipped with an argon laser emitting at 488 nm. Bacteria were gated for size based on log forward angle and side scatter detection and for sorting by FITC/PE labeling of bacteria. Sorted cells were collected into cold sBHI and plated on chocolate agar.
20 After overnight growth, cells were collected for a secondary round of infection or were individually selected and grown overnight, screened by individual clone for fluorescence when grown *in vitro*, and frozen in skim milk containing 20% (vol/vol) glycerol prior to plasmid isolation and sequencing of insert DNA. Sorting efficiency of control strains was confirmed using a Coulter EPICS flow cytometer (Coulter
25 Corp.).

Many plasmids were segregated rapidly *in vitro* in the absence of antibiotic selection. Thus, in order to assess whether the promoter trap vector used here was prone to this event, a single colony of strain 86-028NP/pRSM2211 (highly fluorescent control) was isolated on chocolate agar and passaged 20 times in the
30 absence of antibiotic selection. No significant decrease in fluorescence intensity was observed when compared to bacteria grown in the presence of antibiotic. In addition, the plasmid is maintained in the absence of antibiotic selection *in vivo*. Similar bacterial counts were observed when bacteria-containing middle ear fluids collected

from a chinchilla were plated on chocolate agar with or without kanamycin. These data demonstrate that the promoter trap vector was stably maintained in the absence of antibiotic selection.

In addition to problems with plasmid stability, early studies on the use of GFP as a reporter to study host-pathogen interactions demonstrated that GFP could be continuously synthesized as a cytoplasmic protein with low toxicity, having minimal effects on the bacterial cell-surface dynamics (Chalfie *et al.*, *Science*, 263: 802-805, 1994). The construction of a high level *gfp*-expressing derivative allowed the assessment of the GFP toxicity on NTHi. Growth curves of both the wild-type strain (86-028NP) and the high GFP producing strain 86-028NP/pRSM2211 were compared when grown under similar conditions. The growth rates were similar, indicating that GFP expression was not toxic to the cells.

The 86-028NP *gfp*-expressing derivatives were used to define the parameters for efficient cell sorting. Strain 86-028NP/pRSM2169 was mixed with the intermediate *gfp*-expressing derivative, strain 86-028NP/pKMM4B5, at a 100:1 ratio, simulating the *in vivo* environment that is expected to contain a small percentage of *gfp*-expressing clones relative to the total bacterial population. This mixture was subjected to FACS analysis, collecting the 1.8% most fluorescent population and the 52% least fluorescent population. Flow cytometric analysis of the sorted populations revealed an enrichment of strain 86-028NP/pKMM4B5 to 65% of the bacterial population, a phenomenon that was not observed when sorting on the negative population. Subsequent rounds of sorting would be expected to further enrich for this intermediate fluorescent population. The inability to decrease the amount of fluorescent bacteria in the negative sort was attributed to the size of the gate set for negative sorting. GFP-negative cells were enriched by gating on the 10% least fluorescent population.

Example 5

Direct Labeling of Bacteria from Middle Ear Fluids

A similar strategy (as described in Example 5) was applied to sort fluorescent clones from effusions obtained from the chinchilla middle ear during AOM. Our ability to use differential fluorescence induction (DFI) *in vivo* was dependent upon our ability to sort *gfp*-expressing bacteria from non-fluorescent bacteria, fluorescent and non-fluorescent cellular debris, and eukaryotic cells.

Healthy adult chinchillas (*Chinchilla lanigera*) with no evidence of middle ear infection by either otoscopy or tympanometry were used to screen the library for promoter activity *in vivo*. Two pools of the NTHi/pRSM2169 library (1000 clones each) were grown overnight on chocolate agar containing kanamycin.

5 The library was combined and diluted in cold 10 mM sterile PBS to 3.3×10^6 CFU/ml and 300 μ l (1.0×10^6 CFU; 500 CFU/clone) was used to inoculate the left and the right chinchilla transbullar cavity (2000 clones/ear). OM development was monitored by video otoscopy and tympanometry at 24 and 48 hours. The bacteria multiplied in the middle ear cavity, reaching a concentration 500 times the inoculum dose by 48

10 hours as expected (Bakaletz *et al.*, *Infect. Immunity* 67: 2746-62, 1999). This bacterial adaptation to the host environment results in an inflammatory response, indicated by erythema, vessel dilation and bulging of the tympanic membrane, infiltration of polymorphonuclear cells (PMN's), and accumulation of fluid in the middle ear cavity as observed by otoscopy and microscopic examination of recovered

15 effusions. Twenty-four and 48 hours later, middle ear fluids were retrieved by epitympanic tap, and prepared for FACS.

It is important to note that this analysis was limited to those bacteria recoverable in the middle ear fluid. In some cases it was necessary to lavage the middle ear cavity to collect the bacteria for FACS analysis. Thus, this analysis

20 includes genes up-regulated when NTHi are loosely adherent to mucosae. NTHi has been observed to form a biofilm in the middle ear cavity in a chinchilla model of OM (Erhlich *et al.*, *JAMA*, 287: 1710-5, 2002). Since the protocols described herein select for clones recovered from the planktonic population, it is not expected to recover those clones in which genes are up-regulated when the bacteria are associated

25 with mucosal biofilms. Homogenization of middle ear mucosae and subsequent bacterial cell isolation however, would enable us to recover these clones. It is also possible that some GFP-expressing clones were recovered in the effusion, yet were adherent to eukaryotic cells present in the effusion as exfoliated cells, or in aggregates. These bacteria are difficult to recover from the effusion without

30 compromising the sorting efficiency. Therefore the middle ear fluids were treated with a mucolytic agent, then centrifuged to remove large aggregates and eukaryotic cells and prior to labeling.

Chinchilla middle ear fluids were diluted, if necessary, to 250 μ l with sterile saline. An equal volume of N-acetyl-L-cysteine (0.5%; w/v) in DPBS (pH 7.4) was added for 5 minutes at room temperature as a mucolytic agent (Miyamoto and Bakaletz, *Microb. Pathog.*, 21: 343-356 1996). Fluids were centrifuged (300 x g, 5 min) to remove cellular debris, red blood cells and inflammatory cells, and supernatants containing bacteria were transferred to a fresh tube. Bacteria were incubated with chinchilla antiserum (1:50 dilution) directed against a whole OMP preparation, derived from NTHi strain 86-028NP, for 45 minutes at 4°C, pelleted by centrifugation (2000 x g, 5 min) and washed twice with cold DPBS containing 0.05% bovine serum albumin. Bacteria were subsequently labeled with cross-reactive phycoprobe R-PE anti-human IgG (H+L) antibody (10 μ g/ml in 100 μ l PBS; Biomed Corp) for 30 minutes at 4°C. Following three successive washes to remove unbound antibody, cells were resuspended in 300 μ l DPBS for FACS analysis.

Example 6

15 Identification of Promoters Induced *In Vivo* in Acute Otitis Media

H. influenzae 86-028NP transformed with the promoter trap library was grown overnight on chocolate agar. To select against those clones containing promoters that expressed *gfp in vitro*, the library was subjected to one round of FACS analysis (as described in Example 6), collecting only those clones expressing low-level amounts of GFP. These clones were pooled and used to inoculate the chinchilla middle ear transbullarily. Following 24 and 48 hours of infection, bacteria-containing effusions were removed by epitympanic tap. Bacteria were indirectly labeled with R-PE-labeled antibody and subjected to FACS analysis by gating on fluorescently tagged bacteria but sorting for those that were also expressing. These clones were used to reinfect animals for further enrichment. Following the final round of sorting, single colony isolates were screened *in vitro* for lack of fluorescence.

Those clones isolated by FACS analysis (positive for GFP fluorescence *in vivo*), which did not emit fluorescence *in vitro* were prepared for plasmid isolation and identification of insert DNA sequence. These clones were grown overnight on chocolate agar plates containing kanamycin and prepared for plasmid isolation using the Qiaprep Miniprep Kit (Qiagen) according to the manufacturer's protocol. Plasmid insert DNA was sequenced using the primer 5'-TGCCCATTAACATCACCATCTA-3' (SEQ ID NO: 588) that is complementary to

the *gfpmut3* gene and downstream of the insert DNA. Sequencing reactions were performed using the ABI prism BigDye[®] terminator cycle sequencing ready reaction kit (Applied Biosystems) according to manufacturer's protocol using a GeneAmp PCR System 9700 (Applied Biosystems). The sequences were then purified by
5 passage through sephadex G-50 in a 96-well multiscreen HV plate (Millipore) and subsequently analyzed on an ABI Prism 3100 DNA analyzer (Applied Biosystems).

Insert sequences were compared to the complete annotated sequence of *H. influenzae* strain Rd. Those inserts with no nucleotide homology to strain Rd were subsequently analyzed using the BLASTN and BLASTX algorithms. Further
10 sequence analysis was performed with DNASTAR (Madison, Wisc). Inserts in the correct orientation and containing sequence 5' to a predicted ORF contained a putative promoter that was preferentially active when the NTHi bacteria were in the chinchilla middle ear.

Fifty-two clones with putative promoters that were regulated *in vivo*
15 were isolated. Of the 44 candidate clones containing sequence similar to that identified in *H. influenzae* strain Rd, quantitative comparison of gene expression *in vitro* and *in vivo* confirmed up-regulated gene expression for twenty-six genes (60%) when NTHi respond to environmental cues present in the chinchilla middle ear and these genes are summarized in Table 4A below. The *in vivo*-regulated promoters
20 driving expression of genes are predicted to be involved in membrane transport, environmental informational processing, cellular metabolism, gene regulation, as well as hypothetical proteins with unknown function.

In order to confirm the induction of putative promoter candidates *in vivo*, the relative amount of messenger RNA expression was compared when NTHi
25 strain 86-028NP was grown *in vitro* to mid-log phase or *in vivo* for 48 hours. The RNA was isolated using TRIzol LS reagent (Gibco Life Technologies) according to the manufacturer's protocol. DNA was removed from the RNA preparation using DNA-free kit (Ambion) according to the manufacturer's protocol. DNase I treated RNA samples were purified by passage through a Qiagen RNeasy column. RNA
30 purity and integrity was assessed by 260/280 nm spectrophotometer readings and on the Agilent 2100 Bioanalyzer (Agilent Technologies), respectively.

In order to independently confirm the FACS data, we determined the relative expression of candidate genes by quantitative RT-PCR. The parent strain 86-028NP, was used for these studies. Real-time quantitative RT-PCR using the one-step QuantiTect SYBR Green RT-PCR kit (Qiagen) assessed transcription levels

5 according to the manufacture's instructions. Briefly, using primers generated to an open reading frame downstream of the putative *in vivo*-induced promoters identified by FACS analysis, gene-specific mRNA was reverse transcribed and amplified by RT-PCR on the ABI Prism 7700 sequence detection system (Applied Biosystems). The amount of product was calculated using a standard curve generated to known

10 amounts of bacterial genomic DNA (10^2 - 10^7 genomic copies DNA) by amplifying a fragment of the gyrase (*gyr*) gene. Controls were analyzed in parallel to verify the absence of DNA in the RNA preparation (-RT control) as well as the absence of primer dimers in control samples lacking template RNA. In addition, RT-PCR products were analyzed by gel electrophoresis and, in all cases, a single product was

15 observed at the appropriate base pair size. Amounts of bacterial RNA between samples were normalized relative to *gyr* expression, shown to be constitutively expressed under various growth conditions that we tested *in vitro*. Known amounts of bacterial genomic DNA (10^2 - 10^7 genomic copies DNA) were used to generate a standard curve for RT-PCR quantitation by amplifying a fragment of the gyrase (*gyr*)

20 gene. Gyrase is constitutively expressed *in vitro* under various growth conditions and was therefore used to normalize total bacterial RNA levels between samples. Relative gene expression *in vivo* was compared to that of gene expression *in vitro* and data expressed as fold-increase are summarized in Table 4A.

The 8-fold sequencing of the NTHi genome identified the full length

25 open reading frames for the majority of genes listed in Table 4A. Table 4B provides the full length nucleotide sequence within the NTHi genome and the corresponding amino acid sequence. The fold induction of the gene due to environmental cues present in the chinichilla middle ear and the product or function of the gene are repeated in Table 4B for convenience.

Table 4A

Category	Gene or ORF	SEQ ID NO:	GenBank Protein ID	Fold Induction	Product or Function
Amino acid metabolism	<i>hisB</i>	589	NP_438632	2.9	Histidine biosynthesis bifunctional protein
Lipoprotein	<i>lppB</i>	590	NP_438862.1	2.6	Lipoprotein B homologue
Membrane transport	<i>sapA</i>	591	NP_439780.1	2.8	Peptide ABC transporter; periplasmic SapA precursor
	<i>lolA</i>	592	NP_439736.1	2.4	Outer membrane lipoproteins carrier protein precursor
	<i>rbsC</i>	593	NP_438661.1	5.1	Ribose transport system permease protein
Purine synthesis	<i>purE</i>	594	NP_439757.1	51.7	Phosphoribosylaminoimidazole carboxylase catalytic subunit; PurE
Biosynthetic and metabolic functions	<i>ribB</i>	595	NP_438923.1	8.3	3,4-dihydroxy-2-butanone 4-phosphate synthase; riboflavin biosynthesis
	<i>arcB</i>	596	NP_438753.1	10	Ornithine carbamoyltransferase; arginine degradation
	<i>uxuA</i>	597	NP_438228.1	3.1	Mannonate dehydratase; production of glyceraldehyde 3-phosphate
	<i>dsbB</i>	598	NP_438589.1	2.6	Disulfide oxidoreductase; disulfide bond formation protein B
	<i>ureH</i>	599	NP_438693.1	3.9	Urease accessory protein
	<i>licC</i>	600	NP_439688.1	2.3	Phosphocholine (ChoP) cytidyltransferase
	HI1647	601	NP_439789.1	2.0	Putative pyridoxin biosynthesis protein; singlet oxygen resistance protein
DNA replication, repair	<i>ispZ</i>	602	P43810	2.5	Probable intracellular septation protein
	<i>radC</i>	603	NP_439113.1	2.1	DNA repair protein
	<i>mukF</i>	604	P45185	2.0	MukF protein homologue; remodeling of nucleoid structure
Gene regulation	<i>glpR</i>	605	NP_438777.1, NP_439170.1	2.8	Glycerol-3-phosphate regulon repressor
	<i>ihfB</i>	606	P43724	2.5	Integration host factor beta subunit
	<i>argR</i>	607	NP_439365.1	2.7	Arginine repressor
	<i>cspD</i>	608	NP_439584.1	2.1	Cold shock like protein; stress response protein
Hypothetical or unknown proteins	HI0094	609	NP_438267.1	8.3	Hypothetical protein
	HI1163	610	NP_439321.1	2.3	Conserved hypothetical protein; putative oxidase
	HI1063	611	NP_439221.1	2.7	Hypothetical protein
	HI0665	612	NP_438824.1	2.8	Hypothetical protein
	HI1292	613	NP_439444.1	2.6	Hypothetical protein
	HI1064	614	NP_439222.1	2.6	Hypothetical protein

Table 4B

Category	Gene or ORF	Full Length Nucleotide Sequence	Amino Acid Sequence	Location in Contig	Fold Induction	Product or Function
Amino acid metabolism	<i>hisB</i>	SEQ ID NO: 615	SEQ ID NO: 616	nt. 68378-67290 of SEQ ID NO: 680 (contig 13)	2.9	Histidine biosynthesis bifunctional protein
Membrane transport	<i>sapA</i>	SEQ ID NO: 617	SEQ ID NO: 618	nt. 200403-198709 of SEQ ID NO: 685 (contig 18)	2.8	Peptide ABC transporter; periplasmic SapA precursor
	<i>rbsC</i>	SEQ ID NO: 619	SEQ ID NO: 620	nt. 42773-41802 of SEQ ID NO: 680 (contig 13)	5.1	Ribose transport system permease protein
Purine synthesis	<i>purE</i>	SEQ ID NO: 621	SEQ ID NO: 622	nt. 219625-219131 of SEQ ID NO: 685 (contig 18)	51.7	Phosphoribosylaminimidazole carboxylase catalytic subunit; PurE
Biosynthetic and metabolic functions	<i>ribB</i>	SEQ ID NO: 623	SEQ ID NO: 624	nt. 131537-132184 of SEQ ID NO: 682 (contig 15)	8.3	3,4-dihydroxy-2-butanone 4-phosphate synthase; riboflavin biosynthesis
	<i>arcB</i>	SEQ ID NO: 625	SEQ ID NO: 626	nt. 49710-48706 of SEQ ID NO: 681 (contig 14)	10	Ornithine carbamoyltransferase; arginine degradation
	<i>uxuA</i>	SEQ ID NO: 627	SEQ ID NO: 628	nt. 840671-841855 of SEQ ID NO: 685 (contig 18)	3.1	Mannonate dehydratase; production of glyceraldehyde 3-phosphate
	<i>dsbB</i>	SEQ ID NO: 629	SEQ ID NO: 630	nt. 388050-388583 of SEQ ID NO: 384 (contig 17)	2.6	Disulfide oxidoreductase; disulfide bond formation protein B
	<i>ureH</i>	SEQ ID NO: 631	SEQ ID NO: 632	nt. 4452-5267 of SEQ ID NO: 680 (contig 13)	3.9	Urease accessory protein
	<i>licC</i>	SEQ ID NO: 633	SEQ ID NO: 634	nt. 355083-354382 of SEQ ID NO: 385 (contig 18)	2.3	Phosphocholine (ChoP) cytidyltransferase
	HI1647	SEQ ID NO: 635	SEQ ID NO: 636	nt. 664017-664892 of SEQ ID NO: 685 (contig 18)	2.0	Putative pyridoxin biosynthesis protein; singlet oxygen resistance protein
DNA replication, repair	<i>ispZ</i>	SEQ ID NO: 637	SEQ ID NO: 638	nt. 4512-5069 of SEQ ID NO: 683 (contig 16)	2.5	Probable intracellular septation protein
	<i>radC</i>	SEQ ID NO: 639	SEQ ID NO: 640	nt. 132695-132030 of SEQ ID NO: 683 (contig 16)	2.1	DNA repair protein

	<i>mukF</i>	SEQ ID NO: 641	SEQ ID NO: 642	nt. 504549- 503215 of SEQ ID NO: 685 (contig 18)	2.0	MukF protein homologue; remodeling of nucleoid structure
Gene regulation	<i>glpR</i>	SEQ ID NO: 643	SEQ ID NO: 644	nt. 72716-73483 of SEQ ID NO: 682 (contig 15)	2.8	Glycerol-3- phosphate regulon repressor
	<i>ihfB</i>	SEQ ID NO: 645	SEQ ID NO: 646	nt. 661004- 660720 of SEQ ID NO: 685 (contig 18)	2.5	Integration host factor beta subunit
	<i>argR</i>	SEQ ID NO: 647	SEQ ID NO: 648	nt. 178540- 178085 of SEQ ID NO: 685 (contig 18)	2.7	Arginine repressor
	<i>cspD</i>	SEQ ID NO: 649	SEQ ID NO: 650	nt. 435310- 435528 of SEQ ID NO: 685 (contig 18)	2.1	Cold shock like protein; stress response protein
Hypotheti- cal or unknown proteins	HI1163	SEQ ID NO: 651	SEQ ID NO: 652	nt. 137202- 134119 of SEQ ID NO: 685 (contig 18)	2.3	Conserved hypothetical protein; putative oxidase
	HI1063	SEQ ID NO: 653	SEQ ID NO: 654	nt. 35158-34937 of SEQ ID NO: 685 (contig 18)	2.7	Hypothetical protein
	HI0665	SEQ ID NO: 655	SEQ ID NO: 656	nt. 17949-18980 of SEQ ID NO: 679 (contig 12)	2.8	Hypothetical protein
	HI1292	SEQ ID NO: 657	SEQ ID NO: 658	nt. 555002- 555799 of SEQ ID NO: 685 (contig 18)	2.6	Hypothetical protein

Example 7

Identification of Virulence-Associated Genes

In many bacterial species, a subset of virulence-associated genes is regulated by errors in replication of short repeats. These repeats may be 5' to a gene or in the coding sequence, and their presence is an indication of controlled expression of the gene, which indicates association with virulence. Addition or deletion of a repeat results in the expression or of lack of expression of the particular virulence determinant.

The NTHi *H. influenzae* strain 86-028NP contig set was queried for short oligonucleotide repeats. The region surrounding the repeats was analyzed to identify the gene(s) associated with the repeat. Table 5 lists the identified repeats and the ORF (identified by BLAST) associated with each repeat.

Further sequence analysis has identified the full length nucleotide sequence of the virulence-associated genes and the corresponding amino acid

sequences encoded by the ORF. The derived amino acid sequences are highly homologous to the listed Genbank sequence.

Table 5

Repeat	Location in 3-fold Contigs	Location in 8-fold Contigs	Full Length Nucleotide Sequence	Amino Acid Sequence	Genebank Accession No.
SEQ ID NO: 581	115 nt. 473-540 of SEQ ID NO: 115	nt. 484533-483643 of SEQ ID NO: 685 (contig 18)	SEQ ID NO: 659	SEQ ID NO: 660	NP_439538.1
SEQ ID NO: 582	377 nt. 546-597 of SEQ ID NO: 337	nt. 416274-414910 of SEQ ID NO: 685 (contig 18)	SEQ ID NO: 661	SEQ ID NO: 662	P45217
SEQ ID NO: 583	505 nt. 310-393 of SEQ ID NO: 505	nt. 414500-416614 of SEQ ID NO: 684 (contig 17)	SEQ ID NO: 663	SEQ ID NO: 664	AAK76425
SEQ ID NO: 584	508 nt. 2079-2120 of SEQ ID NO: 508	nt. 506516-507913 of SEQ ID NO: 685 (contig 18)	SEQ ID NO: 665	SEQ ID NO: 666	NP_439520
SEQ ID NO: 585	518 nt. 758-789 of SEQ ID NO: 518	nt. 354274-352406 of SEQ ID NO: 684 (contig 17)	SEQ ID NO: 667	SEQ ID NO: 668	NP_284893
SEQ ID NO: 586	543 nt. 1814-196 of SEQ ID NO: 543	nt. 347864-243236 of SEQ ID NO: 685 (contig 18)	SEQ ID NO: 669	SEQ ID NO: 670	AAA20524

64267-1392D

68

Repeat	Location in 3-fold Contigs	Location in 8-fold Contigs	Full Length Nucleotide Sequence	Amino Acid Sequence	Genebank Accession No.
SEQ ID NO: 586	543 nt. 1814-196 of SEQ ID NO: 543	nt. 699709-704187 of SEQ ID NO: 685 (contig 18)	SEQ ID NO: 671	SEQ ID NO: 672	AAD56660
SEQ ID NO: 587	567 nt. 13309-13320 of SEQ ID NO: 567	nt. 85546-84689 of SEQ ID NO: 681 (contig 14)	SEQ ID NO: 673	SEQ ID NO: 674	ZP_00053190

Example 8**Identification of Unique NTHi Gene Sequences**

Genes associated with NTHi virulence were also identified by

- 5 comparing the level of expression of the gene when the NTHi bacterium was infecting a tissue verses the level of expression of the same gene when the NTHi was grown on artificial laboratory media. These novel genes were identified using the promoter trap techniques described above in Examples 4-6, and subsequently comparisons with the known Rd genome demonstrated these genes are unique to
- 10 NTHi strain 86-028NP.

The DNA sequence identified using this screening procedure are set forth as SEQ ID NOS: 577-580. These sequences did not contain genes or gene fragments that have homologues in the *H. influenzae* Rd. genome sequence. Even though these are completely novel sequences, due to their expression level during

15 NTHi infection in the chinchilla middle ear, it is likely that expression of these genes are involved in NTHi virulence.

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 64267-1392D Seq 06-11-12 v1.txt)

- 20 A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

64267-1392D

69

CLAIMS:

1. An isolated polynucleotide comprising the nucleotide sequence of any one of SEQ ID NOS: 577-579, SEQ ID NOS: 589-614, SEQ ID NOS: 675-685, SEQ ID NO: 615, SEQ ID NO: 617, SEQ ID NO: 619, SEQ ID NO: 621, SEQ ID NO: 623, SEQ ID NO: 625,
 - 5 SEQ ID NO: 627, SEQ ID NO: 629, SEQ ID NO: 631, SEQ ID NO: 633, SEQ ID NO: 635, SEQ ID NO: 637, SEQ ID NO: 639, SEQ ID NO: 641, SEQ ID NO: 643, SEQ ID NO: 645, SEQ ID NO: 647, SEQ ID NO: 649, SEQ ID NO: 651, SEQ ID NO: 653, SEQ ID NO: 655, SEQ ID NO: 657, SEQ ID NO: 659, SEQ ID NO: 661, SEQ ID NO: 665, SEQ ID NO: 667, SEQ ID NO: 669,671, SEQ ID NO: 673, SEQ ID NO: 686, SEQ ID NO: 688, SEQ ID
 - 10 NO: 692, SEQ ID NO: 694, SEQ ID NO: 696, SEQ ID NO: 698, SEQ ID NO: 700, SEQ ID NO: 702, SEQ ID NO: 704, SEQ ID NO: 706, SEQ ID NO: 708, SEQ ID NO: 710, SEQ ID NO: 712, SEQ ID NO: 714, SEQ ID NO: 716, SEQ ID NO: 718, SEQ ID NO: 720, SEQ ID NO: 722, SEQ ID NO: 724, SEQ ID NO: 726, SEQ ID NO: 728, SEQ ID NO: 730, SEQ ID NO: 732, SEQ ID NO: 734, SEQ ID NO: 736, SEQ ID NO: 738, SEQ ID NO: 740, SEQ ID
 - 15 NO: 742, SEQ ID NO: 744, SEQ ID NO: 746, SEQ ID NO: 748, SEQ ID NO: 750, SEQ ID NO: 752, SEQ ID NO: 754, SEQ ID NO: 756, SEQ ID NO: 758, SEQ ID NO: 760, SEQ ID NO: 762, SEQ ID NO: 764, SEQ ID NO: 766, SEQ ID NO: 768, and SEQ ID NO: 770.
2. An isolated polypeptide comprising an amino acid sequence encoded by the polynucleotide of claim 1 or a fragment thereof.
- 20 3. An isolated polypeptide comprising an amino acid sequence of any one of SEQ ID NO: 616, SEQ ID NO: 618, SEQ ID NO: 620, SEQ ID NO: 622, SEQ ID NO: 624, SEQ ID NO: 626, SEQ ID NO: 628, SEQ ID NO: 628, SEQ ID NO: 630, SEQ ID NO: 632, SEQ ID NO: 634, SEQ ID NO: 636, SEQ ID NO: 638, SEQ ID NO: 640, SEQ ID NO: 642, SEQ ID NO: 644, SEQ ID NO: 646, SEQ ID NO: 648, SEQ ID NO: 650, SEQ ID NO: 652,
 - 25 SEQ ID NO: 654, SEQ ID NO: 656, SEQ ID NO: 658, SEQ ID NO: 660, SEQ ID NO: 662, SEQ ID NO: 666, SEQ ID NO: 668, SEQ ID NO: 670, SEQ ID NO: 672, SEQ ID NO: 674, SEQ ID NO: 687, SEQ ID NO: 689, SEQ ID NO: 691, SEQ ID NO: 693, SEQ ID NO: 695, SEQ ID NO: 697, SEQ ID NO: 699, SEQ ID NO: 701, SEQ ID NO: 703, SEQ ID NO: 705,

64267-1392D

70

SEQ ID NO: 707, SEQ ID NO: 709, SEQ ID NO: 711, SEQ ID NO: 713, SEQ ID NO: 715, SEQ ID NO: 717, SEQ ID NO: 719, SEQ ID NO: 721, SEQ ID NO: 723, SEQ ID NO: 725, SEQ ID NO: 727, SEQ ID NO: 729, SEQ ID NO: 731, SEQ ID NO: 733, SEQ ID NO: 735, SEQ ID NO: 737, SEQ ID NO: 739, SEQ ID NO: 741, SEQ ID NO: 743, SEQ ID NO: 745, 5 SEQ ID NO: 747, SEQ ID NO: 749, SEQ ID NO: 751, SEQ ID NO: 753, SEQ ID NO: 755, SEQ ID NO: 757, SEQ ID NO: 759, SEQ ID NO: 761, 763, SEQ ID NO: 765, SEQ ID NO: 767, SEQ ID NO: 769, and SEQ ID NO: 771.

4. A composition comprising the polypeptide of claim 2 or 3 and a pharmaceutically acceptable carrier.

10 5. An antibody that specifically binds to the polypeptide of claim 2 or 3 or fragment thereof.

6. A composition comprising the antibody of claim 5 and a pharmaceutically acceptable carrier.

7. A method for detecting NTHi bacteria in a biological sample comprising:

15 (a) contacting the polynucleotide of claim 1 or a fragment thereof with a biological sample, and

(b) detecting hybridization of the polynucleotide within the sample.

8. A method for detecting NTHi bacteria in a biological sample comprising:

(a) contacting the antibody of claim 5 with a biological sample, and

20 (b) detecting binding of the antibody within the sample.

9. The method of claim 7 or 8 wherein the biological sample is selected from the group consisting of serum, sputum, ear fluid, blood, urine, lymphatic fluid, and cerebrospinal fluid.

64267-1392D

71

10. Use, for eliciting an immune response to NTHi bacteria in a patient at risk of NTHi bacterial infection, of an immunogenically effective dose of the polypeptide of claim 2 or 3 or a fragment thereof.
11. A vaccine comprising the polypeptide of claim 2 or 3 or a fragment thereof and
5 a pharmaceutically suitable carrier.
12. Use for preventing NTHi bacterial infection in a patient at risk of NTHi bacterial infection, of the polypeptide of claim 2 or 3 that blocks cellular attachment of NTHi bacteria.
13. Use for preventing NTHi bacterial infection in a patient at risk of NTHi
10 bacterial infection, of the antibody of claim 5 that blocks cellular attachment of NTHi bacteria.
14. Use, in the manufacture of a medicament, for preventing NTHi bacterial infection in a patient at risk of NTHi bacterial infection, of the polypeptide of claim 2 or 3 that blocks cellular attachment of NTHi bacteria.
- 15 15. Use, in the manufacture of a medicament, for preventing NTHi bacterial infection in a patient at risk of NTHi bacterial infection, of the antibody of claim 5 that blocks cellular attachment of NTHi bacteria.
16. The use of any one of claims 13 to 15 wherein the NTHi infection is in the middle ear.
- 20 17. Use, for treatment or prevention of NTHi bacterial infection, of a molecule that inhibits expression or activity of the polypeptide of claim 2 or 3.
18. Use, in the manufacture of a medicament, for treatment or prevention of NTHi bacterial infection, of a molecule that inhibits expression or activity of the polypeptide of claim 2 or 3.

64267-1392D

72

19. The use of claim 17 or 18 wherein the polypeptide is encoded by an NTHi gene selected from the group consisting of hisB, lppB, sapA, rbsC, pure, rib, arcB, uxuA, licC, ispZ, ritukF, glpR, il, cspD, lav, HI1647, HI0094, HI1163, HI0665, HI1292, HI1064, HI1386, HI1462, HI1369, and HI1598.

5 20. The use of claim 17 or 18 wherein the polypeptide comprises the amino acid sequence selected from the group consisting of SEQ ID NO: 616, SEQ ID NO: 618, SEQ ID NO: 620, SEQ ID NO: 622, SEQ ID NO: 624, SEQ ID NO: 626, SEQ ID NO: 628, SEQ ID NO: 634, SEQ ID NO: 638, SEQ ID NO: 642, SEQ ID NO: 644, SEQ ID NO: 646, SEQ ID NO: 650, SEQ ID NO: 652, SEQ ID NO: 656, SEQ ID NO: 658, SEQ ID NO: 660, SEQ ID
10 NO: 662, SEQ ID NO: 666, SEQ ID NO: 668, SEQ ID NO: 670, SEQ ID NO: 672, and SEQ ID NO: 674.

21. The use of claim 17 or 18 wherein the molecule is an antisense oligonucleotide.

22. The use of claim 17 or 18 wherein the molecule is an antibody.

15 23. The use of claim 17 or 18 wherein the molecule is a small molecule.

24. The use of any one of claims 17 to 23 wherein the NTHi infection is in the middle ear.

Figure 1




Figure 2




Figure 3A

SEQ ID NO: 589

TTNAACAAGANATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCA
TATGTATATCTCCTTCTTAAATCTAGAGGATCAAGCAACTTAATTTTTGGT
TTGCGACATAGACAATTTTCTTCGGGCTTATGCGGGCAAATGAGTACTTCA
TCAAAGGTAATGCCCTGAGATTCAAATAATGCCATCATTACATTGTGGGG
CTTATCAAAATTTGTTTGCGGAAAAGCATCCGTACCAAGCCCATCTTGATT
AGAACTATCACAAAACGGTATTTTCGCTTTCAGTCTGAGGAGTGCGGGAA
TCACTTTCGGTTCTAATTTTAGTTTCTCTAAACTGTCAATTTGGAAATCAGT
TTTTGGTTCATCAATTAATGTGCCGTCTCGGTCGATAAAAAGAGTAGGTTG
CATATTTTCTCCGTGTATGTTATATGAATGGTATGAGGTTTATCATTTGCT
TTATACAAACAGTATAAACTAAATAATCTCGAATCGCTCTGTGACTTAA
ATAGGAGGTGGTTCATTTTACCTCCTTAATTGCTTCAACAACCTTCTCACA
CTCATTCCTTGTTCCACAGTAATGCGAATACAATTTTGT

Nt 80-410 of SEQ ID NO: 589 corresponds to compliment of nt 1-331 of the *hisB* gene.

Nt 80-600 of SEQ ID NO: 589 corresponds to nt 4655-5175 of the contig 532 (SEQ ID NO: 532).

SEQ ID NO: 590

TAACAAGAACTTGGGACAAAGNCCAGTGAAAAGTTCTTCTCCTTTACTCA
TATGTATATCTCCTTCTTAAATCTAGAGGATCAGCATCACATCGGGCGGAA
TAGGAAAAAATATCGCCTCGATAAAACTTACAAAAGAGAGCCAAAAAAC
AGCAAAACGATGTTTTGACCATTCCATCGTTTTATCGTACATCGTGCCGAA
AATTTTCATATTTTTCTCTATTTGGTAAATTCCTGCTCTAACCAATTTTGCA
AAGAAAGCAATGAATGATATGCGGTAAAGATCCTCTAGAGTCGACCTGCAG
GCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTG
TTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
AAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGC
TCACTGCCCCGCTTTCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAATGA
ATCGGCCAACGCGCGGGGAGAGGCGGTTTGCGTATTGGGCGCCAGATACC
GGCTCCATCGGAGAACTGTCCGAGGTTATGTTGACCTGCAGGG

Nt 78-210 of SEQ ID NO: 590 corresponds to compliment of nt 1-133 of the *lppB* gene.

Nt 78-285 of SEQ ID NO: 590 corresponds to compliment of nt 1423-1630c of the contig 442 (SEQ ID NO: 442).

Figure 3B

SEQ ID NO: 591

NACAGATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGTA
TATCTCCTTCTTAAATCTAGAGGATCCACAATAATATGTTCTATCTTGGCTT
CTTTTTTCCAATAGTTTTTCGTTACGCACTAAGCGAACATATTGGTTATATA
CATAATCTTTTACTTGATAAGGCCCTGTGCCTACTGGGTGGGTATCTAATT.
GAGCAAGGTTGTCATCTGCGCTTAATTGATAGGCATATTCTTGTGAAAAA
ATAATGGCATACTGGCTGGCAAGATGCGACAAAATGGAGGAATCTGGTGC
AAATAATTCAATTTTTTACTTGATAAGGCGAAAGTGCGGTCACAGATTTGAT
TTTTTCGTTAAGTTTAATGCTATCAAAATAAGGAAAACGCACTTTTCTTGC
TTGTTTCGTGAAACACTCTATATTGTGGATTACTATAGGNAACCATNCGCCT
CTGCTAAGGTTGGTAAATAAGTATTATGCCCTAATACACGATTAATCGAA
AATACTTCNGTNTTCAGNGNTAAAAACACCNTGNNGGGGNAANNCNANG
GGGTTTGGGGGAAATTTTACCCCCGGGACGTAAATTNAATAAA

Nt 74-514 of SEQ ID NO: 591 corresponds to compliment of nt 369-811 of the *sapA* gene.

Nt 74-514 of SEQ ID NO: 591 corresponds to nt 3458-3897 of the contig 512 (SEQ ID NO: 512).

SEQ ID NO: 592

TTAACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCAT
ATGTATATCTCCTTCTTAAATCTAGAGGATCGTAGAACCATAAGGTTTTAC
CATCAGAAATAATCTGGGTTTCTTGAGGGGTTTTAGTCTCCATACGGAATA
AATTTGGGCGTTTAATTTGAAGTTTGCCACTTCCTTGTTGAACATTTTTTCC
ACTTCCAGAAGTCACTGTTTGCACAAATTCTGCACTTAATACATCGACTTT
AGCTAAACGCATTTGTAATTCATTGCGCATCAGCCAATGCCAAATTACT
CAAACCAAGTAAGGTAAGTGCGGCAAATTTTAAGGTTGTTTTTTTCATTTT
ACTTTCCTTTTAAATTAGTATTCTGGACGATGCGATAAAATTCACGCTTA
CCATTTTGCATTGGGCTGACAATCCCTTGCTCTTCCATTTGATCCTCTAGAG
TCGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCT
GTGTGAAATTGNTATCCGNTCACAATTCCCACAACATACGAGCCNGAAGC
ATAAAGTGTAAGCCTGGGGGGNCTNATGAGGGAGCTA

Nt 78-345 of SEQ ID NO: 592 corresponds to the compliment of nt 11-278 of the *lola* gene:

Nt 78-454 of SEQ ID NO: 592 corresponds to compliment of nt 1039-1415 of the contig 360 (SEQ ID NO: 360).

Figure 3C

SEQ ID NO: 593

TTAACAGAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATAT
GTATATCTCCTTCTTAAATCTAGAGGATCTATACCTGCAATTAAAATAACG
AAAGTCATACCCACTGCAATAATCGCATTAAACCGAAGTTTGGCGTAAAAT
ATTCAAAATATTATCTACGCTAAAAAAATCAGGATTAATCATTGATACAA
TCGCGATAAGAATAATCAACGCAATAAAAGAACGCTGTTCAATCAAAAAT
CTTCCTATTTGAAAATTAGATGTTTTCATTTTTTCATCATACTTACCTACTCTT
ATTTACCAATAGCTGCTGCTAATAATTTTTCTTGAGTTGCGTCTTTGCGAG
AAAATTCTGCACTAATTTTACCTTCACGCATTACCAACACTCTATCGCTCA
TACCAAGCACTTCTGGCATATCAGATGAGATCCTCTAGAGTCGACCTGCA
GGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATT
GNTATCCGNTCACAATTNCACACAACATACGAGCCNGNAGCATAAAGGG
GAANNCCGGGGGGGCCTAANGGGGNGCTACNNNATTNATNGGGTG

Nt 77-290 of SEQ ID NO: 593 corresponds to compliment of nt 1-214 of the *rbsC* gene.

Nt 77-438 of SEQ ID NO: 593 corresponds to compliment of nt 1813-2174 of the ,
contig 445 (SEQ ID NO: 445).

SEQ ID NO: 594

AAGTACTCTGTNTNTCTCCTTCTTAAATCTAGAGGATCGGGATTATCAAGC
ACATTACTCGTTTGGCTTTCACGGAAAGATTGCAAGCGTGAAAGCAACTT
CTGCATCCCAACTTGCTAGAATTTGTGCGGCTAACAAATCCAGCATTTGCCG
CACCCGCAGGGCCAATAGCCAATGTTCCGACTGGGATTCCTTTTGGCATT
GCACAATTGAATAAAGGCTATCTACGCCACTTAACATAGAACTTTTACTG
GCACACCCAGCACAGGCACAAGGGTTTTGGCTGCGATCATGCCGGGTAAA
TGTGCCGCACCGCCAGCCCCAGCAATAATTACTTTATAGCCATTTTTTTGT
GCATTTTCAGCAAATTCAAAAAGTTTATCAGGCGTACGATGAGCAGAGAC
GATTTCCACATGATAAGGTACTTTTAATTCATCTAAAATCTGAGTTGCCTC
TTGCATGGTAGCCCAATCGCTTTTGGAAACCCATCACAACAGCAATTTGTGC
AGTTTTTGACATGCTATTTTCTCAATTTTCTAATTAAAAAGGTGGCGTAGA
ATAGCATAGATTACATATATTGAGCAAACGTTTGCTATNTAT

Nt 105-519 of SEQ ID NO: 594 corresponds to compliment of nt 1-415 of the *purE* gene.

Nt 34-281 of SEQ ID NO: 594 corresponds to compliment of nt 286-40 of the contig
536 (SEQ ID NO: 536).

Figure 3D

SEQ ID NO: 595

TTATGNGACCNTGGGACAAGTCCAGTGAAAAGTTCTTCTCCTTTANTCATA
TGTATATCTCCTTCTTAAATCTAGAGGATCTTCATCATCTAAAACTAATAC
GCCAGTTCCATGTTTGAAAGCATTAAATTGCATTAAGTACGCGTTCTTCAGC
GGTGTTGCCGAATGGAGATAAAATTGACTGATTCATAGTAATTTCTTAATT
TGAGATTTCAATTAAATACCAGAATCAGGGCTGAGAAATGCAAATAAAAT
GAACGATTTGGGAACCCAACCTCGTTTTATTCTCTTTCATCCAGACTTTACT
GTCGGCTTTGGAATTTACCAAATCTGCTGACTTTAAAAGTGAAAATCAG
GACAATTTTCGCAGGAAAAATTTGCCAAAAATTGACCGCACTTTTAAACGC
TCGTGGGCTTTACCACCGGTAGGGAATTTACCCTGCCCTGAGAATGCGA
GCTAAGTATAACGCAAAATACCTAGGCACTAAATCAGAACGTGAAAAATA
TTTTCATTTAGCCTACAATAAACACACTCAATTCTTACGCTATCAAGAGCA
GATTATGTCAAAAACAAAAGAGAAAAAAGTCGGTGTCATTTTCG

Nt 78-189 of SEQ ID NO: 595 corresponds to complement of nt 1-112 of the *ribB* gene.

Nt 237-565 of SEQ ID NO: 595 corresponds to complement of nt 2216-2544 of the contig 290 (SEQ ID NO: 290).

SEQ ID NO: 596

TNACAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATAT
GTATATCTCCTTCTTAAATCTAGAGGATCTAAGAGATATTTAATTTACAGT
TCTGTGTGATGAACAAGACTAAGTAAATGTCTATTTTTTCATATTGAAAGCC
ATAATAATCTCCTTGTGTTTTGACTATACATTGGGGAATAAACATACATCA
TTCGAGATACTGCTTGTTGTTGCGAACAGTTGCATTATAAGAGAGGGTAAAAT
GAAATTTGTTGTCGTTTTTCAACTGGTAATGCCAGTATGTAGAATTTTATT
GCGAGATTTTATAGTATTATTAGAATTTAATAAAAAATAATCTAACTCTTTAT
TGTTATTTGTTTGATTTCTAATAGTAGATGAATTTAAACGAAAATATGTTT
TAGTTTATTTTTGATGAAGAGGAATGNGTCTATCTATNCTTAATAAAAGNG
CGGNTATTTTTATTGNAGNTTTAATATATTGATCCTNTAGAGAGCGACCC
NCACGCNTGCGCNCTTGGGGCGANCATGGGGCATATAGTTGTTCCCTGTG
NNAANGTGTTATCCGTTACANTTCCCCACAAAANACGAN

Nt 77-155 of SEQ ID NO: 596 corresponds to complement of nt 1-79 of the *arcB* gene.

Nt 77-494 of SEQ ID NO: 596 corresponds to complement of nt 772-1189 of contig 461 (SEQ ID NO: 461).

Figure 3E

SEQ ID NO: 597

AACAGNATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGT
ATATCTCCTTCTTAAATCTAGAGGATCAGTACGAGTCCAGTCTAAAACAG
GCATAAAATTGTAACAAACAGTGTCAATTCCGCATTGAGCAAGCATTGCG
AGAGAGTTTGTGTTATAGTTATTAATCCATTTTGTATAATTCCAGTACTGA
GTTTTAATCTCTTCATGAACCTGGTACACTTTCCACTACTGACCAGCTTAAA
CCAGCATTCTCTATTTTCAGTTTTTGCAGTTTTTTAATTTCTTCGATACTCCAC
ACTTCACCGTTAGGAATATGGTGAAGTGCAGTGACTATAACCAGTTGCGCC
TGCTTGACGAATATCAGATAAAGAAACAGGATCCTCTANAGTCGACCTGC
AGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGNTTCCTGTGNGAAA
CTGGTATCCGCTCACAATTCACCNAACATACGAGCCNGAAGNCATAAAG
NGNTAANCCNTGGGGNGCCTAATGAGTGAGCTAACTCACATTAATTGNGT
TGGGCTTACTGCCCCGTTTTTCAGNNGGGAAACCTGGCGCGCCAGCT

Nt 83-388 of SEQ ID NO: 597 corresponds to complement of nt 34-335 of the *uxuA* gene.

Nt 156-388 of SEQ ID NO: 597 corresponds to complement of nt 779-1008 of the contig 5 (SEQ ID NO: 5).

SEQ ID NO: 598

TNACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATA
TGTATATCTCCTTCTTAAATCTAGAGGATCAACCTCGTAACTTAGCAAAAT
CAGTTACCGTTGAATAAAGTGCGGTTGATTTTAATGTAATAAAAAAAACG
GTGAAATTTACCGTATTTTTTTTAGTTAAATAATCTTCTTTGTTTCCGTGTT
TTAATCACTTGTGCGATTAACAATAATGTGAGAATAACCACATATAGTGA
AAAAATAACCACTAGCCATTGCACCATCGTTAAACCAAATAATGACCATT
GGCTTTCATTACAAGAGCCAGTTGGGTAAATATAAATGGAAACCATTGA
TGAAATGGTAAAGTCTCGGGAAAATTTGGAATAAACTCACATTGTTTCCA
AGGAGCTGGATTCATTTGCAAATCAAGATGACGAAATGAAATCAATAAGC
CCTTAATACTGCTGAATAAACCTAAAGCCAAGGCGATTAATCGAAGTATG
AAAGCAAGAGGTTGAAGTAATGCAATAATGCCCCGCTACAAATAAGCCAAT
CATTGCTAAACGTTTCATAAACACAGAGCACACAAGGTTGTAAGCCC

Nt 176-600 of SEQ ID NO: 598 corresponds to complement of nt 108-532 of the *dsbB* gene.

Nt 78-600 of SEQ ID 598 corresponds to nt 819-1341 of the contig 560 (SEQ ID NO: 560).

Figure 3F

SEQ ID NO: 599

NACAAGATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGT
ATATCTCCTTCTTAAATCTAGAGGATCGCCCGCCAGCACCCCAGGGGAGG
AGGACATCTGCATCGCATTCAAGCCATTCGCCCACGCATCATCGTAACTTG
GCAACGTTATCACTTTAAACGGTGGCGTCGCAAAGTATTAGCAAGCTGG
GTTTTACCGCTGGAGGAAAGTTTGGTGGAAAGGGATAATTTGCTGTTTCAT
AGATTTTGTAGGGGCTAATTACATTAGCCCCGATATTTACCAATAATTGG
GCAAATGTAATTTGCCCCTACGATTAACGAACCAAAGATGCAGGTTCTTC
CACATTCTTCAACAACGCATATTTTTCGATCCTCTAGAGTCGACCTGCAGG
CATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTGT
TATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
AGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTC
ACTGCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATT

Nt 75-252 of SEQ ID NO: 599 corresponds to compliment of nt 1-178 of the *ureH* gene.

Nt 75-385 of SEQ ID NO: 599 corresponds to nt 825-1135 of the contig 258 (SEQ ID NO: 258).

SEQ ID NO: 600

TANACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCAT
ATGTATATCTCCTTCTTAAATCTAGAGGATCCAATCTCAATACGTATTACT
TGCCCATTACTATTTAAAATCGGTAGCCATTCATTGTGCGTTTTAGAACGG
ATAACCGTGAAATATTTGCTGTGTGATGGTTTCGTTAGAAAAATATTACGA
TTAAGCACTACATCCGCATCAATAACATAACAATCACTAAAGAAGTCTTG
TGCCAGAGAAAAGGAATAAATGCTGTTATATTCGCGATATTTTTCGTTGTA
AATCAAGGTGCAGTCATATTTTTTCTTTAGATACTCGAATTGTTTCGTGTAA
ATACCCTGTAACAATGACAATATTGTCGATATTTGCTTGACGTAGAAAAGT
GAGTGTCGGTTCTAAGTTAGGTGTGCCGTGAATATCTAACAATGATTTATG
TGTGCTTTGTGTGATGTCTTTAAACGACCGGCTTANTCNTGCAGCTAAAA
TAATCGCATTCATTTTTGTTCCTTTGNAAAAAAGTGTACCGATCCTCTA
GAGTCGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGN

Nt 82-482 of SEQ ID NO: 600 corresponds to compliment of nt 40-440 of the *licC* gene.

Corresponding sequence not in contig set.

Figure 3G

SEQ ID NO: 601

TNACAAGANTTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATAT
GTATATCTCCTTCTTAAATCTAGAGGATCGCTCATTCGAGAGACACCTCCA
ACAGCACGAATATCTGCAGGAATTCGTTCCAAAGCCATCACTGCAGCTGC
ACCAGCAGCTTCTGCAATGCGAGCTTGTTTCAGGGTTTTGAACATCCATGAT
AACACCACCTTTGAGCATCTGTGCTAAGTTTTTATTTAGTTCATAACGATT
TTCAGCCATTTCTTTGTCCTCATTATTTGGTTATTTGTATTGATAGCTACGT
TTTTTATTGTAATGTAGAAAAGTCTTTAGAACAAGTTTGAAAAAATGCGTA
TATATTAAGGCATAATGTCTAGTGTGCTTAATTTTCAGTTTTTTGTATAA
ACGAAGTGTAGAACGTCAGAAGGATTGTCCAATTTAATAATAGAAAAAAG
AGCTACTTTAGCTGAAAGAATTTTCGTCCTTAAGTGCCACAATAGTTATTAA
AAAAACCTTGAAATGTTTTTAAAAACCTTTTTCTATGTGGTTAGATTTTCT
GAAATTTAAGAAAAAGCTTGAAAGGGAGTTTACGCTTTCC

Nt 75-263 of SEQ ID NO: 601 corresponds to complement of nt 1-189 of the HI1647 gene.

Nt 75-600 of SEQ ID NO: 601 corresponds to complement of nt 1035-1560 of the contig 471 (SEQ ID NO: 471).

SEQ ID NO: 602

TTNACAAGGAATCGGGACAACCTAGCAGTGAAAAGTTCTTCTCCTTTACTA
CATATGTATATCTCCTTCTTAAATCTAGAGGATCTGCAAATTGTTGCAAC
CACTAATACGATGGCTGCTTCACGTACACCACCGAGTTTGTAAGTGATAA
AAAATAAAATTAAAGGGATAAAGTCAAGGAGTTGTTTCATATTGATTAAT
CCTTCATAAATAAGCTGTAGAAACGATAAGTCACTACTAGCATGAAAATA
TTAAGTAATGCGGTGAAAATGCCTATTACCATATCAAATATAGCATTATTG
GAAAATGCGCTTAATTGGAAAATAAGAATTGGCACGAGAAAATAAACCA
GTAGGGTGTAATAAACAATACGCCTTTTCGAGTATTTCTCGCATCCAAA
TTTTTCGGATAGTTTGCGAAAGTGCCTCTTGAGTAGAAAGATAATGTACTA
CCGTAAATTTTAAACGGACAAAGAACCAACGCCAACAAGCATCGCGAT
AAGTGACATAATAGAGGGCGATTTTTTTGTGAGTAATGCANCAAAGGCTT
CACCAAGNACCAGTAACATTGGTGCAACCATTAATAGATCCTCTTAG

Nt 81-191 of SEQ ID NO: 602 corresponds to complement of nt 1-111 of the *ispZ* gene.

Nt 81-593 of SEQ ID NO: 602 corresponds to complement of nt 850-1361 of the contig 418 (SEQ ID NO: 418).

Figure 3H

SEQ ID NO: 603

NACAAGANTTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATG
TATATCTCCTTCTTAAATCTAGAGGATCCTTGAAACATCACCTCAGATTTA
ATCAATCTGTGGATTCTATCCAAGAAAAGTACCACGAAATTTTCTCTTTCC
TCATTACCTATAATGGTTTGAAAATAGAGTTTAGCTATTGATGGCTCAGTA
AATTCCTGTGCACCTTGGTAAAAGGGCTTTTCTTCCATCACTTTAGTTAAA
ATAACCTGTGCCTGTTCTAAAATAGCTTGTTGTTGCTCGTTTAATTTAAAC
ATTGATATTCCTTCTAAAATAGGGAATATCGCCCATAAGGGATATATCCC
CTATGGGTAAGGTGCGAAAAGAATGCTTATCGCACATTAAAGTTAATGTT
ATTAATCAGAAGTTTTTAAACTGTCGATAATCTGTTTAATTTTTGCTGCAA
ATTCTAAATCGTGCTCAGATTCATCATCTATAACAATATTCCAAATAATTAC
TCATATCAGCAATTAGTGCTTCTTTTGTGGCGAAGGTTGGAACATAATTG
ATCCTCTAGAGTCGACCTGCAGGCATGCAAGCTTGGCGTA

Nt 65-525 of SEQ ID NO: 603 corresponds to compliment of nt 18-477 of the *radC* gene.

Nt 76-564 of SEQ ID NO: 603 corresponds to compliment of nt 23606-23869 of the contig 575 (SEQ ID NO: 576)

SEQ ID NO:604

TTAACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCAT
ATGTATATCTCCTTCTTAAATCTAGAGGATCGGCTTCTAACATTTCGCCGT
CTAAACGTTTCGTTGTTATAAATTGCAATCGCCAGCAGGAAAACCAAGCGT
TCGGTAGGTAAATTCAAAGAAAACCTCGCGATCTTTTGCCCAAGAGACGAG
TTCAGGAATAGTTTGGGATGTTTCAATCATTGAATTGCAAAGCGTTAAAAT
TGAACAAAAAATTGGAAAGATTATACAGGATTTTTTTGCGGATCCTCTAGA
GTCGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCC
TGTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAA
GCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTA
ATTGCGTTGCGCTCACTGCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAG
CTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCGTATTGG
GCGCCAGATACCGGNCTCCATNGGANAAACNGTCCCAGGGTNTNT

Nt 79-296 of SEQ ID NO: 604 corresponds to compliment of nt 30-247 of the *mukF* gene.

Nt 79-296 of SEQ ID NO: 604 corresponds to compliment of nt 4026-4243 of the contig 508 (SEQ ID NO: 508).

Figure 3I

SEQ ID NO: 605

TAACAGNATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATG
TATATCTCCTTCTTAAATCTAGAGGATCCACGTAATCAGAATTTTCTGCAG
AAGATGGCGATGCCGCACCACCGTGATGGCGGCGAATTAAATCTAACTCC
GCCAAAATATTCAAATCACGACGAATCGTTTGAGGGCTAACATCTAAGGC
AGCAACCAATTCTTCCGTGCTTAAATAGCCAGATTGTTCCACCAGTTTAAT
AATTTTTTGATGGCGTAACGATTGTTTCATAGCGAATCCTTTATCCAAATT
AAACGGACTTACTCTAGCGAATTTTTGCGCAATTTGCTATCAATAAATCCC
CAAATCAAGCCGACAATCAAGCCTGAAATATGTGCAGCATTTCCCATTTTC
AACACCAAATAAGGGGGCTAATAAAACCTAATGCGATCCTCTAGAGTCGAC
CTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTG
AAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAA
AGTGNAAGCCTGGGGNGCCTAAGGGTGAGCTAACTCACATTATT

Nt 76-283 of SEQ ID NO: 605 corresponds to compliment of nt 1-208 of the *glpR* gene.

Corresponding sequence not in contig set.

SEQ ID NO:606

TAACAAGAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATAT
GTATATCTCCTTCTTAAATCTAGAGGATCACCATTTTCAAGAGATTGAGAA
ATAAACTCTAAAATATCTTTTACCATATTTTCAATTTCTTTTGCAGATAAA
GTTGGCTGTTTTGCTGACAATTTTCCATAAGTTCTGACTTAGTCATCCTCT
ATACTTCCTTAATATTATTAAATTAAGATCCTCTAGAGTCGACCTGCAGG
CATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTGT
TATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
AGCCTGGGGTGCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTC
ACTGCCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAAT
CGGCCAACGCGCGGGGAGAGGCGGTTTGCGTATTGGGCGCCAGATACCGG
CCTCCATCGGAGAACTGTCCGAGGTTATGTTGACCTGCAGGGGGGGGGGG
GGCCCCTGAGGTCTGCCTCGNGAAGAAGGGGTTGCTGACTC

Nt 77-200 of SEQ ID NO: 606 corresponds to compliment of nt 1-124 of the *ihfB* gene.

Nt 77-236 of SEQ ID NO: 606 corresponds to compliment of nt 1455-1614 of the contig 408 (SEQ ID NO: 408).

Figure 3J

SEQ ID NO: 607

NACAGNNTTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGT
ATATCTCCTTCTTAAATCTAGAGGATCAACATCTAAGACAAGATTTTTTAA
TGGACTACTGGTATTGGGAACACTTAATTCATTTGGTAAGCAATAGACCAT
TTCCATTTTTGTATTACGAGTGCGAACC GCACCAAATTTGCTGAGCATTCT.
CGATATTTTTGATTGATTAATAACCAGTAAAGCCTTGTTTTTTTAAGGCATC
GACAATTTTCATTTTGAGAACCAAAGCGTTCTTGATTGAGTAATTCTTTAAA
AGCACGAGTTAAATTGTCAGTCATCTTTATTTTTTAATAGTCAAAATTTGC
ATAAGAATTGCATAAAAATTNNNCTTTCATCAACTGGAATATGGNTTNGN
ACCNNGGGGAAGAAATTTTTTTNNNNAAAATNGGAACNAGATCCNTTNA
GNCAACCTGCAGGCATGCAAGCTNGGCGNAATCANGGCATAGCTGNTTCC
TGGGNAAATTGNTATCCGTTANAATCCCNCNACATNCNANCCGNAGCAT
AAGNGAAAGCCTGGGGGCNTANGANGNGGCTACCCCATNATT

Nt 75-330 of SEQ ID NO: 607 corresponds to compliment of nt 1-256 of the *argR* gene.

Corresponding sequence not found in contig set.

SEQ ID NO: 608

TTTAACAAGGAATCCGGGCACAACCTACCAGTGAAAAGTTCTCTCTCCTTTA
CTCATATGTATATCTCCTTCTTAAATCTAGAGGATCCTTTATCGCTGTGGA
GAACTTCAAATTGTACTTTTTGACCTGCTTTTAATGAACGATAGCCGTCCA
TTTCAATCACTGAATAATGTGCAAAAATATCGGCATCTACACCTTCTGCGG
AAATGAAACCAAATCCTTTTGCATTATTGAACCATTTTACAATACCAATTT
CCATAAAAGACCTCTCTAGGCTTAGCCTATTAAAACAATAAATCAACAAG
ACTCTCGCCTTATCACCTACTAATTAGGTCCGTTATGTTTAAATATTTTGAA
CGGTTATGCAACGAGCAATATTCAAAAATGATAAGTAGATCCTCTAGAGT
CGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTNCCT
GNGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG
CATAAAGTGNTAAGCNTGGGGTGCCTAATGAGTGNNCTAACTNACATAAA
TTGAGTTANGCTCACTGCNCCGTTTTCCAGNNGGNGANACCNG

Nt 84-259 of SEQ ID NO: 608 corresponds to compliment of nt 1-176 of the *ihfB* gene.

Corresponding sequence not found in contig set.

Figure 3K

SEQ ID NO: 609

TAACAGCATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATG
TATATCTCCTTCTTAAATCTAGAGGATCTATTACAGAACCATCATTTCCCC
AAATCAAGAATAGCTTTGATGAAAATAAGTAAAATCGGTAAGACAATAG
GAATACTCACGACATAGCCCGTAATGGCTAATGCCCATTCCTTCTTTTCT
TACCTAGAAATTTGATAAAACTGAACGCCATTTTTTTCAGCAACTGGGCAG
ATTGAAGTTTCACCATGTAAAGGTTCTCGAATAATGCTACTTACAAAGGA
AGAAGTCGCTTTATTTGGTTTTCTGGACGGATACAAATTGTGAGCAGGCG
AACACAGAGACCATCAACAAAACCTTTTCTTGTATAGTCATTAATAAGTA
ATTCATTCATTGCTTTTTGTGCACCATAAGTCGATTGTGGCGTTACAGCAG
TAGAATCCTGAATAATTTCTGGTAAATCACCACCAAATACAGCAAGTGAA
CTAGAAAAAATAAAACGAATTTTAGTATCAGATTCATCAAAATATGAGAA
AGTGGCGACATTCCATATTTGTCCACTAACAAATTTTGATCCTCTA

Nt 93-413 of SEQ ID NO: 609 corresponds to compliment of nt 1-321 of the HI0094 gene.

Nt 76-595 of SEQ ID NO: 609 corresponds to nt 1467-1985 of the contig 379 (SEQ ID NO: 379).

SEQ ID NO: 610

TTNNACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCA
TATGTATATCTCCTTCTTAAATCTAGAGGATCATCAAGATAATCAAGTACA
TATTGTTCAATTTGAGGAATGCGATTTAGACTTGGCAACATAGTATTTAAC
TCATTTAAAGGTAAAAATGGCAGGTTATTGATAATATCTTAAGGCGTTAAT
GATGTCGAATTAGATTTTGTGACATTTTAAAGGAGTGTTTTATGGAGTAAAT
GAGTCAAGAAAAGTGTGTTGTTTGGATGAAAAATAAACNAAAAATTCAAA
AGAATTTGATCCTCTAGAGTCGACCTGCAGGCATGCAAGCTTGGCGTAAT
CATGGTCATAGCTGNTTCCTGTGTGAAATTGNTATCCGCTCACAATTCCAC
ACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATG
AGTGAGCTAACTCACATTAATTGCGNNGCGCTCNCTNNCCCCCTTTNCNCA
GGGGAAAACCCTGNCGCNCCAGCTGCNTTTAATGAATCNGCCCNCGCG
GGGAGGGGCGNNTTGCGTNTTGGGCGCCANANCCCGNCTCCATCG

Nt 80-143 of SEQ ID NO: 610 corresponds to compliment of nt 1-64 of the HI1163 gene.

Nt 80-316 of SEQ ID NO: 610 corresponds to compliment of nt 117-353 of the contig 513 (SEQ ID NO: 513).

Figure 3L

SEQ ID NO: 611

NACAAGNAATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATAT
GTATATCTCCTTCTTAAATCTAGAGGATCGTCATCTTTACGATATTTTTCAT
CAATTCGTTTTTGGCAATCCAAAGTCATTTTTATCCGATTCGTTAGAAAATA
AATCTTCTGTTTGCCTTAATTGAGGATTTTGAATTCCTATCCAATTTGCAAT
CCCTTCTAAGAAATTTAAACCTGATTTAAACACCTTATATTCTTTGCGTTCC
ATATCATCTGATGAAATTTTGAATAGCGGAATATTATGGTGCTCTCGGCTG
AAACAGTTTTGATTGAACAAAAGTATATTGTTTTTTTCATCTTGTTGGTGG
CACAAACCGTGATCCTCTAGAGTCGACCTGCAGGCATGCAAGCTTGGCGT
AATCATGGTCATAGCTGTTTCCTGTGTGAAATTGTTATCCGCTCACAATTC
CACACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTA
ATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCCGNTTCCA
GTCGGGAAANCCTGTCGTGCCAGCTGCATTAATGAATCG

Nt 76-260 of SEQ ID NO: 611 corresponds to compliment of nt 1-185 of the gene
HI1063.

Nt 76-374 of SEQ ID NO: 611 corresponds to nt 229-527 of the contig 48 (SEQ ID
NO: 48).

SEQ ID NO: 612

ATTNNACAAGNAATTGGGCACAACCTCCAGTGAAAAGTTCTTCTCCTTTACT
CATATGTATATCTCCTTCTTAAATCTAGAGGATCATCTTTGATAATAAGCT
GTAATTTTGGTTGAAAACCTGAAATGCTCATCCCTTGCTGTTTTTGGGGTT
TCACTGTAATAAATTCTTGGCGAGAAAAGGGCAATTCAGGATTAATAATGT
TTATTGCCTGTAAGATAGTGTAACCTTTTGTGCTGTAACCAGAATGAAC
TCATTTTTTTTCAATGGCTTTAATAAAATACGACAAAATTCATTGTTTTTC
TTCCTGAATTTGCACTGCGCCTAACATATTTTCCCCGTTATTAAGTAAAAA
GCGAAACATATCGCTTTCATCAATTCGTTGATGAAGTGCATATTTGTGTTT
TAGCCAGCCTTCTGGGACGAGGGATGCAAAATAAGGGAATAAGGTGTNA
GAATGAAAAGGACTTTGCTCAATAGGAAGCTGAGAGAGAGCGGTATTCC
TTGATAATCGGGATTGTAAGCAAAATGAAAACCTCGATGATCCTCTAGAG
TCGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGC

Nt 84-298 of SEQ ID NO: 612 corresponds to compliment of nt 1-215 of the HI0665
gene.

Nt 82-549 of SEQ ID NO: 612 corresponds to compliment of nt 4726-5193 of the
contig 557 (SEQ ID NO: 557).

Figure 3M

SEQ ID NO: 613

NACAGNATTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGT
ATATCTCCTTCTTAAATCTAGAGGATCACTGCATTGAGTAAACGACTTTGT
GATTCAATATTATTTTCGAGTGCAGTCAGGCGAAAATAGGTTTCGATTA
TACATAAATTTTTTAATTGCTGTCTGATTGGTTTAAGCCGTTTAAACGCAT
TTCACTTAAACGTTGTGAAATGCCGAGTTCGCTAAAAAAGGCAATAATAT
CAAATGTATTCAATTCATCCGCTGTTTTGCCATTGATTTGATTGAATAAAA
TCCAAAGCAAGCCATTCATAATTCTGGCTTCGCTAAATCCGCTGAATTGAA
AAGTGCGGTCATGTTTAGGCATAATTTGAAACCACATTTGTGCTTCGCAGC
CAGNNAATAGGTTGCATTTGAGCGAGTTCATTATCGCTTGGGCGAGGNN
ANNTTTTGNCTNCNCNGGAAAAATCAGGNNATNGCGATCCTTTAGAGTCG
ACCTGCAGGCATGCAAGCTTGGCNGTAATCATGGNCATAGCTGNTTCCTG
GGGGAAATNGTTNTCCGNTNCAATCCCCACAACATACGAGC

Nt 75-158 of SEQ ID NO: 613 corresponds to complement of nt 1-84 of the HI1292 gene.

Nt 142-455 of SEQ ID NO: 613 corresponds to nt 12-324 of the contig 313 (SEQ ID NO: 313).

SEQ ID NO: 614

CAAGACTTGGGACAACCTCCAGTGAAAAGTTCTTCTCCTTTACTCATATGTA
TATCTCCTTCTTAAATCTAGAGGATCTACAGTTGGTTCTGTAAACATCCCA
GAACCGATTAAAATAAAATAACCCGCGAATGCGGCACAAATCACCGCAA
AAAGTGCGGGCAAAATTTGGCTTGTTTTTTTCGTGTTTCATCATTGTCTTCCC
TGAAAAGAAAATATCGCTATTATCCACCTTTCACTCAGAATTTCCATATTT
AACTTCTTCACTCTTGAGCGGTTATTTTATAGAATAAGCCAATTTTTTTTAA
GCCAGAAGGAAAAAACAATGAGCCAACCAATTTATAAACGTATTTTATTG
AAATTAAGCGGTGAAGCATTACAAGGAGAAGATGGTCTTGGTATCGATCC
TCTAGAGTCGACCTGCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGC
TGTTTCCTGTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAG
CCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTC
ACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCAGTCGGGAAA

Nt 83-265 of SEQ ID NO: 614 corresponds to complement of nt 1-181 of the HI1064 gene.

Nt 74-406 of SEQ ID NO: 614 corresponds to nt 440-772 of the contig 178 (SEQ ID NO: 178).

