




**ΚΥΠΡΙΑΚΟ ΓΡΑΦΕΙΟ ΔΙΠΛΩΜΑΤΩΝ
ΕΥΡΕΣΙΤΕΧΝΙΑΣ
THE PATENT OFFICE OF CYPRUS**

**ΑΡΙΘΜΟΣ ΔΗΜΟΣΙΕΥΣΗΣ CY2012
PUBLICATION NUMBER**

ΑΡΙΘΜΟΣ ΔΗΜΟΣΙΕΥΣΗΣ
ΓΡΑΦΕΙΟΥ ΔΙΠΛΩΜΑΤΩΝ ΕΥΡΕΣΙΤΕΧΝΙΑΣ
ΗΝΩΜΕΝΟΥ ΒΑΣΙΛΕΙΟΥ
UK PATENT OFFICE
PUBLICATION NUMBER GB2259647

Το έγγραφο που παρουσιάζεται πιο κάτω καταχωρήθηκε στο «Γραφείο Διπλωμάτων Ευρεσιτεχνίας» στην Αγγλία σύμφωνα με το Νόμο Κεφ. 266 πριν την 1^η Απριλίου 1998. Δημοσίευση έγινε μετέπειτα από το Γραφείο Διπλωμάτων Ευρεσιτεχνίας του Ηνωμένου Βασιλείου μόνο στην Αγγλική γλώσσα.

**The document provided hereafter was filed at "The Patent Office"
in England under the law CAP.266 before the 1st of April 1998.
It was published afterwards by the UK patent office only in English.**

(12) **UK Patent Application** (19) **GB** (11) **2 259 647** (13) **A**
(43) Date of A publication **24.03.1993**

(21) Application No **9219749.0**

(22) Date of filing **18.09.1992**

(30) Priority data

(31) **9120131**

(32) **20.09.1991**

(33) **GB**

(71) Applicant

Glaxo Group Limited

(Incorporated in the United Kingdom)

**Glaxo House, Berkeley Avenue, Greenford, Middlesex,
UB6 0NN, United Kingdom**

(72) Inventor

Andrew Alexander McColm

(74) Agent and/or Address for Service

C L Brewer

**Glaxo Holdings plc, Glaxo House, Berkeley Avenue,
Greenford, Middlesex, UB6 0NN, United Kingdom**

(51) INT CL⁵

A61K 31/29

(52) UK CL (Edition L)

**A5B BKC B180 B21Y B213 B215 B216 B28Y B285
B44Y B444 B446 B45Y B451 B452 B453 B54Y
B540 B56Y B565
U1S S1318**

(56) Documents cited

GB 2220937 A WO 92/11849 A1

(58) Field of search

UK CL (Edition K) A5B BKC

INT CL⁵ A61K

Online databases: WPI; CLAIMS,CAS ONLINE

(54) **Ranitidine salt compositions**

(57) Compositions useful in treating gastro intestinal disorders contain (i) ranitidine bismuth citrate and (ii) one or more *Helicobacter pylori*-inhibiting antibiotics.

Suitable antibiotics are metronidazole, tetracyclin or amoxycillin. Packs containing the compounds in (i) and (ii) are also disclosed. The active compounds can be given simultaneously or sequentially.

GB 2 259 647 A

MEDICAMENTS
FOR TREATING GASTROINTESTINAL DISORDERS

5 The present invention relates to improvements in the treatment of gastrointestinal disorders. More particularly it relates to the co-administration of a salt formed between ranitidine and a complex of bismuth with a carboxylic acid, with antibiotics.

 In our published UK Patent Specification No. 2220937A we describe and claim salts formed between ranitidine and a complex of bismuth with a carboxylic acid,
10 particularly tartaric acid and, more especially, citric acid. One such salt is N-[2-[[[5-[(dimethylamino)methyl]-2-furanyl]methyl]thio]ethyl]-N'-methyl-2-nitro-1,1-ethenediamine 2-hydroxy-1,2,3-propanetricarboxylate bismuth (3⁺) complex, also known as ranitidine bismuth citrate.

 The salts disclosed in UK Patent Specification No. 2220937A possess the
15 H₂-antagonist antisecretory properties associated with ranitidine, together with antibacterial activity against Helicobacter pylori (formerly Campylobacter pylori). In addition, such salts possess cytoprotective properties, and display activity against the human gastric pepsins, with preferential inhibition of pepsin 1, a pepsin isozyme associated with peptic ulcer. The salts disclosed in UK Patent Specification No.
20 2220937A thus possess a particularly advantageous combination of properties for the treatment of gastrointestinal disorders, especially peptic ulcer disease and other gastroduodenal conditions, for example gastritis and non-ulcer dyspepsia.

 We have now shown that the antibacterial activity of ranitidine bismuth citrate against Helicobacter organisms may be significantly enhanced by co-administering the
25 compound with one or more antibiotics.

 The present invention thus provides, in one aspect, the use of (i) ranitidine bismuth citrate and (ii) one or more Helicobacter pylori-inhibiting antibiotics in the manufacture of medicaments for simultaneous, separate or sequential use in treating or preventing gastrointestinal disorders.

30 The term "gastrointestinal disorder" as used herein encompasses a disease or other disorder of the gastrointestinal tract, including for example a disorder not manifested by the presence of ulcerations in the gastric mucosa (eg gastritis, non-ulcer dyspepsia,

esophageal reflux disease and gastric motility disorders), and peptic ulcer disease (eg gastric and duodenal ulcer disease).

The ranitidine bismuth citrate is preferably co-administered with one or two antibiotics but, in particularly difficult cases, three antibiotics may be required.

- 5 A wide variety of antibiotics may be used according to the invention, including for example nitroimidazole antibiotics (e.g. tinidazole or metronidazole), tetracyclines (e.g. tetracyclin, doxycyclin and minocyclin), pencillins (e.g. amoxycillin, ampicillin and mezlocillin), cephalosporins (e.g. cefachlor, cefadroxil, cephradine, cefuroxime, cefuroxime axetil, cephalixin, cefpodoxime proxetil, ceftazidime and ceftriaxone),
10 carbopenems (e.g. imipenem and meropenem), amino-glycosides (e.g. paromomycin), macrolide antibiotics (e.g. erythromycin, clarithromycin and azithromycin), lincosamide antibiotics (e.g. clindamycin), 4-quinolones (e.g. ofloxacin, ciprofloxacin, pefloxacin and norfloxacin), rifamycins (e.g. rifampicin), nitrofurantoin and derivatives of 10-(1-hydroxyethyl)-11-oxo-1-azatricyclo[7.2.0.0.3.8]undec-2-ene-2-carboxylic acid described
15 in published European Patent Specification No. 0416953 and published International Patent Specification No. WO92/03437.

Antibiotics which may be administered orally are generally preferred. Examples include metronidazole, tetracyclin (especially as tetracyclin hydrochloride), amoxycillin, cefuroxime axetil and clarithromycin.

- 20 Preferred combinations include ranitidine bismuth citrate and metronidazole; ranitidine bismuth citrate and tetracyclin; ranitidine bismuth citrate and cefuroxime axetil; ranitidine bismuth citrate and amoxycillin; ranitidine bismuth citrate and clarithromycin; ranitidine bismuth citrate, metronidazole and amoxycillin; ranitidine bismuth citrate, metronidazole and tetracyclin; and ranitidine bismuth citrate, tetracyclin
25 and clarithromycin.

Combinations of ranitidine bismuth citrate and cefuroxime and ranitidine bismuth citrate, cefuroxime and metronidazole may also be preferred.

- Particularly preferred combinations comprise ranitidine bismuth citrate co-administered with either metronidazole, tetracyclin, metronidazole and amoxycillin, or
30 metronidazole and tetracyclin. Such combinations have shown a synergistic antibacterial effect against Helicobacter pylori in vitro and against Helicobacter mustelae in vivo in ferrets. Thus, for example, in combination studies in vivo, co-administration of

ranitidine bismuth citrate with one or two antibiotics hereinabove has produced a level of antibacterial activity against Helicobacter mustelae which is better than that shown by the active ingredients individually or, in the case of co-administration with two antibiotics, that shown by the two antibiotics together.

5 The ranitidine bismuth citrate and the one or more antibiotics are preferably co-administered in the form of separate pharmaceutical compositions for simultaneous and/or sequential use. Alternatively the ranitidine bismuth citrate and the antibiotic(s) may be administered as a single pharmaceutical composition for oral use comprising effective amounts of the active ingredients.

10 Thus, according to a further aspect, the invention provides a product containing (i) ranitidine bismuth citrate and (ii) one or more Helicobacter pylori-inhibiting antibiotics as a combined preparation for simultaneous, separate or sequential use in treating or preventing gastrointestinal disorders.

When the ranitidine bismuth citrate and the one or more antibiotics are administered
15 as separate preparations, each of the antibiotics may conveniently be provided in the manner known in the art and/or commercially for the compounds concerned. Administration of both the ranitidine bismuth citrate and the antibiotic(s) by the oral route is preferred, although the antibiotic(s), where appropriate, may also be given by another route, for example parenterally (e.g. intravenously, intramuscularly or
20 subcutaneously).

The ranitidine bismuth citrate may conveniently be formulated as tablets (including chewable tablets), capsules (of either the hard or soft type), or as a liquid preparation, as described for example in UK Patent Specification Nos. 2220937A and 2248185A. Tablets are generally preferred.

25 As stated hereinabove, ranitidine bismuth citrate and the antibiotic(s) may be administered as a single pharmaceutical composition for oral use. Thus according to a further aspect the invention provides a pharmaceutical composition, for oral use in human or veterinary medicine, comprising ranitidine bismuth citrate and one or more Helicobacter pylori-inhibiting antibiotics. Such compositions may be formulated in a
30 conventional manner using additional pharmaceutically acceptable carriers or excipients as appropriate. Thus, the composition may be prepared by conventional means with additional carriers or excipients such as binding agents (e.g. pregelatinised maize starch,

polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrates (e.g. starch or sodium starch glycollate); or wetting agents (e.g. sodium lauryl sulphate). An alkaline salt of the type described in UK Patent
5 Specification No. 2248185A may be added to improve the rate of disintegration and/or dissolution of the composition. Tablets may be coated by methods well known in the art. The preparations may also contain flavouring, colouring and/or sweetening agents as appropriate.

The compositions may be prepared according to conventional techniques well
10 known in the pharmaceutical industry. Thus, for example, the ranitidine bismuth citrate and the antibiotic(s) may be admixed together, if desired, with suitable excipients. Tablets may be prepared, for example, by direct compression of such a mixture. Capsules may be prepared by filling the blend along with suitable excipients into gelatin capsules, using a suitable filling machine.

15 When ranitidine bismuth citrate and the antibiotic(s) are administered as a single pharmaceutical composition for oral use the composition is preferably in the form of a capsule or, more particularly, a tablet.

Low packing density oral pharmaceutical compositions comprising a soluble bismuth-containing pharmaceutical agent, including ranitidine bismuth citrate, optionally
20 also comprising a Helicobacter pylori-inhibiting antibiotic were recently described in International Patent Specification No. WO92/11849. Thus, according to a further aspect, the present invention provides a pharmaceutical composition for oral use comprising ranitidine bismuth citrate and one or more Helicobacter pylori-inhibiting antibiotics wherein the packing density of the composition is not less than 1g/ml.

25 The compositions for use according to the invention may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredients. The pack may for example comprise metal or plastic foil, such as a blister pack. Where the ranitidine bismuth citrate and the one or more antibiotics are intended for administration as separate compositions these may be presented in the form
30 of, for example, a multiple (e.g. twin) pack.

Thus, according to a further aspect the present invention provides a multiple-container pack for use in treating or preventing gastrointestinal disorders, one of

the containers containing ranitidine bismuth citrate and the other(s) containing a Helicobacter pylori-inhibiting antibiotic.

The doses at which the ranitidine bismuth citrate and the one or more antibiotics may be administered to man (of approximately 70kg body weight) will depend on the
5 route of administration of the antibiotic and on the nature and severity of the condition being treated. It will also be appreciated that it may be necessary to make routine variations to the dosage depending on the age and weight of the patient.

A proposed dosage of ranitidine bismuth citrate for use according to the invention is 150mg to 1.5g, preferably 200 - 800mg per unit dose. The unit dose may be
10 administered, for example, 1 to 4 times per day.

The one or more antibiotics may conveniently be administered at doses within the normal dosage range at which the compound(s) are therapeutically effective, or at higher doses if required. The antibiotic(s) may be taken one or more times daily as appropriate.

In a further aspect, the present invention provides a method for treating or
15 preventing gastrointestinal disorders in a human or animal subject, which comprises administering to said subject effective amounts of ranitidine bismuth citrate and one or more Helicobacter pylori-inhibiting antibiotics.

The methods of the present invention comprise administering the Helicobacter pylori-inhibiting antibiotic(s) and ranitidine bismuth citrate either concurrently or
20 non-concurrently. As used herein, concurrent administration means that the agents are given within 24 hours of each other, whereas non-concurrent administration means that the agents are given more than 24 hours apart. When the agents are administered concurrently, it may be preferable to administer the agents within about 1 hour of each other or, more preferably, within about 5 minutes of each other.

25 For the methods of the present invention, the duration of administration of the agents during either concurrent or non-concurrent dosing will vary according to the specific gastrointestinal disorder being treated. However, a typical regime would be to administer ranitidine bismuth citrate for 4 to 8 weeks and during this period to administer one or more antibiotics for 1 to 2 weeks.

30 As stated hereinabove, certain combinations of ranitidine bismuth citrate with antibiotics have shown a synergistic effect in vitro against Helicobacter pylori and in vivo against Helicobacter mustelae.

In Vitro Synergy Methodology

Synergy was measured dependent on the rate of kill observed in minimal bactericidal concentrations (MBCs) using a 2-dimensional microtitre checkerboard method¹. The checkerboard is produced by serially diluting agent A prior to addition to the plates. Agent B was then diluted in the wells containing A to a final volume of 50µl. Each well is then inoculated with 50µl of H.pylori cultured in broth and the plates incubated at 37°C. Plates are then sampled at 24, 48 and 72 hours for H.pylori growth using a 0 - 6 quantitative scale. Mean fractional inhibitory concentrations (FICs) were then determined from 2-dimensional isobolograms. A mean FIC index of less than 1 = synergy².

To confirm synergy, time kill studies using the apparently effective combinations were subsequently performed in 3 ml broth cultures.

$$\text{FIC index} = \text{FIC}_A + \text{FIC}_B$$

$$\text{where } \text{FIC}_A = \frac{\text{MBC}_A \text{ in presence of B}}{\text{MBC}_A}$$

$$\text{and } \text{FIC}_B = \frac{\text{MBC}_B \text{ in presence of A}}{\text{MBC}_B}$$

In Vivo Synergy Methodology

Ferrets naturally colonised with H.mustelae were treated with either ranitidine bismuth citrate (24mg/kg), amoxycillin and metronidazole (10mg/kg and 20mg/kg respectively) or the combination of all three agents. Compounds were given orally three times daily for 28 days with the exception of metronidazole (day 0 - 9 only). The apparent colonisation frequency was assessed from gastric antral biopsies obtained by endoscopy before, during and after the therapy phase. A culture - positive biopsy means that the compound is still colonised.

Results

In vitro, the combination of ranitidine bismuth citrate with tetracyclin, metronidazole, amoxycillin or cefuroxime demonstrated a synergistic effect. Indeed, the

combination of ranitidine bismuth citrate with tetracyclin or metronidazole gave a mean FIC index < 0.5.

In vivo, the combination of ranitidine bismuth citrate with metronidazole and amoxycillin proved superior to either ranitidine bismuth citrate alone or the antibiotics alone and lead to 2/6 infections being eradicated. The ranitidine bismuth citrate or antibiotic alone groups failed to eradicate but showed good suppression. The removal of metronidazole from the regime after ten days dosing resulted in a subsequent relapse of infections by day 28, particularly in the metronidazole/amoxycillin group.

10 References

1. Berenbaum, M.C., Applied Microbiology 16: 890-895, 1968.
2. McLaren, A and McDowell, S.R., Irish Journal of Medical Science, 5th Workshop on Gastroduodenal Pathology and Helicobacter pylori, p98 (1992).

15

20

25

30

CLAIMS

1. The use of (i) ranitidine bismuth citrate and (ii) one or more Helicobacter pylori-inhibiting antibiotics in the manufacture of medicaments for simultaneous,
5 separate or sequential use in treating or preventing gastrointestinal disorders.
2. The use according to Claim 1 in which the compounds (i) and (ii) are presented as separate compositions for said use.
- 10 3. A product containing compounds (i) and (ii) as defined in Claim 1 as a combined preparation for simultaneous, separate or sequential use in treating or preventing gastrointestinal disorders.
4. A pharmaceutical composition, for oral use, which comprises both a compound (i)
15 and a compound (ii) as defined in Claim 1, together with a pharmaceutical carrier or excipient.
5. A use, product or composition according to any preceding claim in which (ii) is one or more antibiotics selected from metronidazole, tetracyclin, amoxycillin, cefuroxime
20 axetil and clarithromycin.
6. A use, product or composition according to Claim 5 in which (ii) is selected from metronidazole, tetracyclin, cefuroxime axetil, amoxycillin, clarithromycin, metronidazole and amoxycillin, metronidazole and tetracyclin, and tetracyclin and clarithromycin.
25
7. A use, product or composition according to Claim 6 in which (ii) is selected from metronidazole, tetracyclin, metronidazole and amoxycillin, and metronidazole and tetracyclin.
- 30 8. A use or product according to any of Claims 1 - 3 in which (ii) is selected from cefuroxime, and cefuroxime and metronidazole.

9. A use or product according to any of Claims 1 - 7 in which compounds (i) and (ii) are in forms suitable for oral administration.

10. A use or product according to any preceding claim in which compound (i) is
5 formulated as a tablet.

11. A multiple-container pack for use in treating or preventing gastrointestinal disorders, one of the containers containing (i) and the other(s) containing (ii) as defined in the preceding claims.

10

12. A composition according to any of Claims 4 - 7, or a pack according to Claim 11, in associated with instructions for the use of both (i) and (ii) in treating or preventing gastrointestinal disorders.

15 13. A method for the preparation of a composition according to any of Claims 4 - 7 which comprises admixing (i) and (ii) together, if desired, with suitable excipients.

14. A method for treating or preventing gastrointestinal disorders in a human or animal subject, which comprises administering to said subject effective amounts of compounds
20 (i) and (ii) as defined in the preceding claims.

25

30

- 10 -

Patents Act 1977
Examiner's report to the Comptroller under
Section 17 (The Search Report)

Application number

GB 9219749.0

Relevant Technical fields

(i) UK Cl (Edition K) A5B (BKC)

(ii) Int Cl (Edition 5) A61K

Search Examiner

M R WENDT

Databases (see over)

(i) UK Patent Office

(ii) ONLINE DATABASES: WPI, CLAIMS, CAS ONLINE

Date of Search

24 NOVEMBER 1992

Documents considered relevant following a search in respect of claims 1 TO 13

Category (see over)	Identity of document and relevant passages	Relevant to claim(s)
A	GB 2220937 A (GLAXO) see Example 4. Abstract (referred to in Application)	1
P,X	WO 92/11849 A1 (P & G) see Claims 1, 2, 6, 8, 9, 11 (referred to in Applica- tion)	1-13

SF2(p)

SJJ - doc99\fil001058

Category	Identity of document and relevant passages	Relevant to claim(s).

Categories of documents

X: Document indicating lack of novelty or of inventive step.

Y: Document indicating lack of inventive step if combined with one or more other documents of the same category.

A: Document indicating technological background and/or state of the art.

P: Document published on or after the declared priority date but before the filing date of the present application.

E: Patent document published on or after, but with priority date earlier than, the filing date of the present application.

&: Member of the same patent family, corresponding document.

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).