



US 20040137071A1

(19) **United States**

(12) **Patent Application Publication**  
**Unger**

(10) **Pub. No.: US 2004/0137071 A1**

(43) **Pub. Date: Jul. 15, 2004**

(54) **NANOCAPSULE ENCAPSULATION SYSTEM  
AND METHOD**

(60) Provisional application No. 60/185,282, filed on Feb.  
28, 2000.

(75) Inventor: **Gretchen M. Unger**, Chaska, MN (US)

**Publication Classification**

Correspondence Address:

**PATTERSON, THUENTE, SKAAR &  
CHRISTENSEN, P.A.**

**4800 IDS CENTER  
80 SOUTH 8TH STREET  
MINNEAPOLIS, MN 55402-2100 (US)**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 33/24**; A61K 9/14

(52) **U.S. Cl.** ..... **424/489**; 424/649

(57) **ABSTRACT**

The present invention generally relates to nanocapsules and methods of preparing these nanocapsules. The present invention includes a method of forming a surfactant micelle and dispersing the surfactant micelle into an aqueous composition having a hydrophilic polymer to form a stabilized dispersion of surfactant micelles. The method further includes mechanically forming droplets of the stabilized dispersion of surfactant micelles, precipitating the hydrophilic polymer to form precipitated nanocapsules, incubating the nanocapsules to reduce a diameter of the nanocapsules, and filtering or centrifuging the nanocapsules.

(73) Assignee: **Genesegues, Inc.**

(21) Appl. No.: **10/652,814**

(22) Filed: **Aug. 29, 2003**

**Related U.S. Application Data**

(63) Continuation of application No. 09/796,575, filed on Feb. 28, 2001, now Pat. No. 6,632,671.






Figure 1

**Figure 2A**




Nanocapsules prepared under different dispersion conditions.

**Figure 2B**




Cumulative release studies for nanocapsule formulations.

**Figure 2C**




Quantitative recovery of DNA from receiver solutions.

**Figure 3**




Nanocapsule modulation of cellular uptake.

**Figure 4**




Dose response for a nanocapsule formula.

**Figure 5A**




Nanocapsule-delivered transgene production in porcine dermis.

**Figure 5B**




Macromolecule delivery across keratinized barrier epithelia.

**Figure 6**




Incorporation of nanocapsules into a suture coating.

**Figure 7A**




PVP nanocapsules are taken up by fibroblasts but not keratinocytes.

Figure 7B




Nanocapsule design for tumor-targeting.

Figure 7C




Nanocapsule coating design for increased drug safety.

**Figure 8A**



Cellular uptake and lysosomal sequestration of RNA oligomers complexed with polyethyleneimine.

**Figure 8B**



Nanocapsules avoid lysosomal sequestration at 18 hours postaddition.



## NANOCAPSULE ENCAPSULATION SYSTEM AND METHOD

### CROSS-REFERENCE TO RELATED APPLICATION(S)

[0001] This application is a continuation of application Ser. No. 09/796,575 filed Feb. 28, 2001, which claims the benefit of U.S. Provisional Application No. 60/185,282 filed Feb. 28, 2000.

### BACKGROUND OF THE INVENTION

[0002] The present invention generally relates to a field of controlled-release delivery systems for macromolecules, particularly those for nucleic acids and gene therapy. More specifically, the present invention relates to nanocapsules having a diameter of less than about 50 nanometers, in which a bioactive component is located in a core of the nanocapsule, and to methods of forming these nanocapsules.

[0003] Over the past several decades, active and extensive research into the use of nanoparticles in the delivery of bioactive agents has generated a number of approaches in the preparation of nanoparticles. These approaches typically include the use of heat, high pressure homogenization, or high intensity ultrasound sonication to prepare nanoparticles having a diameter of more than 100 nanometers, or high amounts of solvents or oils, cytotoxic chemicals, such as cross-linking agents, adjuvants, catalysts or any combination of any of these, to prepare nanoparticles having a diameter of less than 100 nanometers. Furthermore, these approaches are challenging due to a number of variables.

[0004] For example, when organic solvents are included in the manufacturing process for nanoparticles, the organic solvent may denature the bioactive agent which reduces most, if not all, efficacy of the bioactive agent. In fact, denaturation of the bioactive agent may promote a toxic response upon administration of the nanoparticle, to a human subject, for example.

[0005] In addition, when an organic solvent is used to prepare nanoparticles, the organic solvent may undergo degradation to form a low pH environment that destroys the efficacy of the bioactive agent. Therefore, organic

[0006] As a result, organic solvents are typically removed during the manufacturing process of nanoparticles. However, inclusion of one or more organic solvent removal techniques generally increases the costs and complexity of forming nanoparticles.

[0007] The incorporation of high pressure homogenization or high intensity ultrasound sonication to prepare nanoparticles typically results in entangling or embedding the bioactive agent in a polymeric matrix of the nanoparticle. Entangling or embedding the bioactive agent in the polymeric matrix may also denature the bioactive agent to thereby reduce the efficacy of the bioactive agent.

[0008] Entangling or embedding the bioactive agent in the polymeric matrix of the nanoparticle may also reduce the efficacy of the bioactive agent by permitting premature release of the bioactive agent prior to reaching a target cell. Premature release of the bioactive agent typically promotes cytotoxicity or cell death during administration of the nanoparticle.

[0009] Furthermore, nanoparticles that reach the target cell are typically transported into the target cell via endosomal regulated pathways that results in lysosomal degradation of the bioactive agent and the nanoparticle. Therefore, functional activity of the bioactive agent inside the target cell may not occur since the bioactive agent and the nanoparticle undergoes degradation. As used herein, the term "functional activity" refers to an ability of a bioactive agent to function within a target cell for purposes of providing a therapeutic effect on the target cell.

[0010] Additionally, high pressure homogenization or high intensity ultrasound sonication techniques often require complex and expensive equipment that generally increases costs in preparing nanoparticles. Therefore, an urgent need exists to prepare nanoparticles without the use of cytotoxic chemicals like organic solvents or the use of complex and expensive equipment. Furthermore, an urgent need exists to prepare nanoparticles that do not entangle nor embed the bioactive agent in the nanoparticle so that cytotoxic responses are minimized. Additionally, an urgent need exists to develop a nanoparticle that maybe transported into a target cell where the bioactive agent is released to accomplish therapeutic delivery of the bioactive agent.

### BRIEF SUMMARY OF THE INVENTION

[0011] The present invention generally relates to nanocapsules and methods of preparing these nanocapsules. The present invention includes a method of forming a surfactant micelle and dispersing the surfactant micelle into an aqueous composition having a hydrophilic polymer to form a stabilized dispersion of surfactant micelles. The method further includes mechanically forming droplets of the stabilized dispersion of surfactant micelles, precipitating the hydrophilic polymer to form precipitated nanocapsules, incubating the nanocapsules to reduce a diameter of the nanocapsules, and filtering or centrifuging the nanocapsules.

### BRIEF DESCRIPTION OF THE DRAWINGS

[0012] FIG. 1 is a schematic of a method of the present invention for preparing nanocapsules.

[0013] FIG. 1A illustrates atomic force microscopy of nanocapsule formulations prepared under different dispersion conditions.

[0014] FIG. 1B illustrates results from an experiment documenting quantitative recovery of small amounts of DNA from releasing solutions.

[0015] FIG. 1C illustrates cumulative release over 72 hours for nanocapsules prepared under different dispersion conditions.

[0016] FIG. 2 illustrates relative pinocytotic activity of HacaT keratinocyte cultures treated with DNA complexes, nanocapsules containing DNA or no treatment.

[0017] FIG. 3 illustrates the results of western blotting of total protein from rat fibroblast cultures.

[0018] FIG. 4A illustrates

[0019] FIG. 4B illustrates immunofluorescence microscopy of porcine dermal tissue sections from the experiment of FIG. 4.

[0020] FIG. 5 shows incorporation of nanocapsules into a solid dosage form.

[0021] FIG. 6A illustrates polyvinylpyrrolidone nanocapsule uptake and Green Fluorescent Protein (GFP) expression in 35 mm human dermal fibroblast and immortalized keratinocyte cultures.

[0022] FIG. 6B illustrates tumor targeting of GFP plasmid DNA by Tenascin nanocapsules.

[0023] FIG. 6C illustrates an effect of nanocapsules that are coated with Tenascin and nanocapsules that are not coated with Tenascin on growth inhibition of squamous cell carcinoma and human dermal fibroblast (HDF) cultures.

[0024] FIG. 7 shows uptake of HDF cultures treated with nanocapsules containing 20 mer Fitec-labeled O-methyl RNA oligonucleotides.

#### DETAILED DESCRIPTION

[0025] The present invention generally relates to nanocapsules having a diameter of less than about 50 nanometers (nm). The present invention also relates to a method of preparing these nanocapsules. According to the method of the present invention, a nanocapsule is formed by partitioning a bioactive component within a core of surfactant molecules, and surrounding the surfactant molecules with a biocompatible polymer shell.

[0026] A method for producing the nanocapsule is generally depicted at 10 in FIG. 1. In the method 10, a bioactive component 12 is homogeneously dispersed into a first aqueous composition 14 to form a hydrophilic composition (not shown). Next, a surfactant composition 16, including a surfactant component (not shown) that contains a plurality of surfactant molecules, and an optional biocompatible oil component 18, are introduced into a first dispersing apparatus 20 along with the hydrophilic composition. The surfactant composition 16 is subjected to conditions in the first dispersing apparatus 20 that initiate at least partial adsorption of the surfactant molecules onto a surface of the bioactive component 12.

[0027] Partial adsorption of surfactant molecules onto the surface of the bioactive component 12 initiates partitioning of the bioactive component 12 into a core of a shell formed from the surfactant molecules in the first aqueous composition 14. Adsorption of the surfactant molecules onto the surface of the bioactive component 12 may proceed until an entire surface of the bioactive component 12 is covered by the surfactant molecules to complete partitioning of the bioactive component 12 into the core of surfactant molecules and form a surfactant micelle 22.

[0028] Next, a biocompatible polymer component 24 is added to the surfactant micelle 22 to stabilize the surfactant micelle 22 located in the first aqueous composition 14. Preferably, the biocompatible polymer component 24 surrounds the surfactant micelle 22 in a stabilizing apparatus 26 to form a stabilized surfactant micelle 28.

[0029] After stabilization, the stabilized surfactant micelle 28 is transferred from the stabilizing apparatus 26 into a second aqueous composition 30 located in a second dispersing apparatus 32. Preferably, the second aqueous composition 30 includes a solute (not shown) that is capable of precipitating the biocompatible polymer component 24 that

coats the stabilized surfactant micelle 28. After precipitating the biocompatible polymer component 24 of the stabilized surfactant micelle 28, dispersed, optionally atomized precipitated nanocapsules 36, hereinafter referred to as nanocapsules 36, are formed.

[0030] It has been discovered that dispersing a surfactant composition, that includes a surfactant component having a hydrophile-lipophile-balance (HLB) value of less than about 6.0 units, into an aqueous composition that contains a bioactive component forms surfactant micelles that surround the bioactive component. It has further been discovered that stabilizing the surfactant micelles by adding a biocompatible polymer coats the surfactant micelles to form nanocapsules having a diameter of less than about 50 nm.

[0031] As used herein, the term “nanoparticle” refers to a particle having a matrix-type structure with a size of less than about 1,000 nanometers. When the nanoparticle includes a bioactive component, the bioactive component is entangled or embedded in the matrix-type structure of the nanoparticle.

[0032] The term “nanosphere”, as used herein, refers to a particle having a solid spherical-type structure with a size of less than about 1,000 nanometers. When the nanosphere includes a bioactive component, the bioactive component is adsorbed onto the surface of the nanosphere or embedded in the nanosphere.

[0033] Similarly, the term “nanocore”, as used herein, refers to a particle having a solid core with a size of less than about 1,000 nanometers. When the nanocore includes a bioactive component, the bioactive component is entangled in the nanocore.

[0034] As used herein, the term “nanocapsule” refers to a particle having a hollow core that is surrounded by a shell, such that the particle has a size of less than about 1,000 nanometers. When a nanocapsule includes a bioactive component, the bioactive component is located in the core that is surrounded by the shell of the nanocapsule. The term “nanocapsule” is not meant to encompass, and generally does not include, a particle having a size of less than about 1,000 nanometers, in which a bioactive component is entangled or embedded in the matrix of the nanocapsule or adsorbed onto the surrounding shell of the nanocapsule.

[0035] The bioactive component 12 may be included into the first aqueous composition 14 as a liquid, vapor or in granular form. The form of the bioactive component 12 that is selected preferably permits the bioactive component 12 to (1) remain stable prior to dissolving or dispersing into the first aqueous composition 14, (2) be homogeneously dispersed into the first aqueous composition 14, (3) be optionally condensed to reduce a size of the bioactive component 12, (4) be partitioned into the core of the surfactant micelles 22, (5) be released upon degradation of the biocompatible polymer shell 24 of the nanocapsule 36, and (6) be functionally active upon release from the nanocapsule 36.

[0036] The bioactive component 12 may be characterized as “hydrophilic” or “hydrophobic”. As used herein, the term “hydrophilic” and “hydrophilicity” refers to an ability of a molecule to adsorb water or form one or more hydrogen-bond(s) with water. All references to “hydrophilic” is also understood as encompassing any portion of the molecule that is capable of adsorbing water or forming one or more

hydrogen-bond(s) with water. As used herein, the term “hydrophobic” and “hydrophobicity” refers to an ability of a molecule to not adsorb water nor form one or more hydrogen-bond(s) with water. All references to “hydrophobic” is also understood as encompassing any portion of the molecule that is not capable of adsorbing water nor forming one or more hydrogen-bond(s) with water.

[0037] When the bioactive component **12** is a hydrophilic bioactive component, the hydrophilic bioactive component may be directly added to the first aqueous composition **14**. As an alternative, the hydrophilic bioactive component **12** may be optionally dissolved or dispersed in one or more solvents, such as water, a nonpolar solvent, a polar solvent, or any combination of any of these.

[0038] As used herein, the term “nonpolar solvent” refers to a solvent that does not have a permanent electric dipole moment, and therefore has no ability for an intramolecular association with a polar solvent. Additionally, a nonpolar solvent may be characterized as a solvent that includes molecules having a dielectric constant of less than about 20 units. Similarly, the term “immiscible”, as used herein, refers to an inability of two or more substances, such as two or more liquids, solids, vapors, or any combination of any of these, to form an intramolecular association with another substance. Some non-exhaustive examples of nonpolar solvents may be found in Perry’s Chemical Engineer’s Handbook, Sixth Edition, which is incorporated herein by reference.

[0039] As used herein, the term “polar solvent” refers to a solvent that has a permanent electrical dipole moment, and therefore has an ability to form an intramolecular association with another polar substance, such as a liquid, a solid, a vapor or any combination of any of these. Additionally, a polar solvent may be characterized as a solvent that includes molecules having a dielectric constant of more than about 20 units. Likewise, the term “miscible”, as used herein, refers to an ability of two or more substances to form an intramolecular association with each other. Some non-exhaustive examples of polar solvents may be found in Perry’s Chemical Engineer’s Handbook, Sixth Edition, which has been incorporated herein by reference.

[0040] When the bioactive component **12** is a hydrophobic bioactive component, the hydrophobic bioactive component may be dispersed or dissolved in a solvent that is capable of dispersing or dissolving the hydrophobic molecule, such as the above-mentioned water, a nonpolar solvent, a polar solvent, or any combination of any of these. Preferably, when the bioactive component **12** is a hydrophobic bioactive component **12**, the hydrophobic bioactive component **12** is dissolved or dispersed in a water-miscible solvent, such as, acetone, acetonitrile, ethanol, dimethyl acetamide (DMA), tetrahydrofuran (THF), dioxane, dimethylsulfoxide (DMSO), and dimethylformamide (DMF). Other suitable non-exhaustive examples of water-miscible solvents may be found in Perry’s Chemical Engineer’s Handbook, Sixth Edition, which has been incorporated herein by reference.

[0041] As noted, the bioactive component **12** may be optionally condensed in the first aqueous composition **14** prior to forming the surfactant micelle **16**. For example, when the bioactive component is a polynucleotide, the polynucleotide may be condensed using a DNA-condensing

agent. As used herein, a “DNA-Condensing Agent” is a molecule that facilitates condensation or a size reduction of DNA.

[0042] While condensation of the bioactive component **12** is not critical to the present invention, condensation of the bioactive component **12** maybe practiced to reduce the size of the bioactive component **12**. Condensation of the bioactive component **12** generally reduces the size of the bioactive component **12** prior to partitioning into the core of the surfactant micelle **16**. Reducing the size of the bioactive component **12** may permit maximum incorporation of the bioactive component **12** into the surfactant micelle **22** or may assist a reduction in the overall size of the nanocapsule **36**. Increasing the amount of the bioactive component **12** that may be included as part of the nanocapsule **36** permits incorporation of macromolecules having a large number of monomers, such as a large number of base pairs or amino acids, for example. Some non-exhaustive examples of condensing agents have been reviewed in Rolland, A. P. (1998). *Crit. Rev. Therapeutic Drug. Carr. Syst.* 15:143-198, and is incorporated herein by reference.

[0043] The bioactive component **12** may further include additional components that are compatible with, and that do not interfere with solvation or dispersion of the bioactive component **12**. Some non-exhaustive examples of additional components that may be added to the bioactive component **12** include a DNA-associating moiety, which refers to a molecule, or portions thereof, that interact in a non-covalent fashion with nucleic acids. DNA-associating moieties may include, but are not limited to, a major-and minor-groove binder, a DNA intercalator, a polycation, a DNA-masking component, a membrane-permeabilizing component, a sub-cellular-localization component, or the like. Major- and minor-groove binders, as used herein, are molecules thought to interact with DNA by associating with the major or minor groove of double-stranded DNA.

[0044] Similarly, the term “DNA intercalator”, as used herein, refer to a planar molecule or planar portion of a molecule thought to intercalate into DNA by inserting themselves between, and parallel to, a nucleotide base pair. As used herein, a “polycation” is thought to associate with the negative charges on the DNA backbone. The DNA-associating moiety may be covalently linked through a “reactive group” to a functional component of this invention. The reactive group is easily reacted with a nucleophile on the functional component. Some non-exhaustive examples of reactive groups (with their corresponding reactive nucleophiles) include, but are not limited to N-hydroxysuccinimide (e.g., amine), maleimide and maleimidophenyl (e.g., sulfhydryl), pyridyl disulfide (e.g., sulfhydryl), hydrazide (e.g., carbohydrate), and phenylglyoxal (e.g., arginine).

[0045] The term “DNA-masking component”, as used herein, refers to a molecule capable of masking all or part of a polynucleotide following release from a nanocapsule to increase its circulatory half-life by inhibiting attack by degrading reagents, such as nucleases, present in the circulation and/or interfering with uptake by the reticuloendothelial system. Similarly, the term “membrane-permeabilizing component”, as used herein, refers to any component that aids in the passage of a polynucleotide or encapsulated polynucleotide across a membrane. Therefore, “membrane

permeabilizing component” encompasses in part a charge-neutralizing component, usually a polycation, that neutralizes the large negative charge on a polynucleotide, and enables the polynucleotide to traverse the hydrophobic interior of a membrane.

**[0046]** Many charge-neutralizing components can act as membrane-permeabilizers. Membrane-permeabilization may also arise from amphipathic molecules. A “membrane permeabilizer”, as used herein, is a molecule that can assist a normally impermeable molecule to traverse a cellular membrane and gain entrance to the cytoplasm of the cell. The membrane permeabilizer may be a peptide, bile salt, glycolipid, phospholipid or detergent molecule. Membrane permeabilizers often have amphipathic properties such that one portion is hydrophobic and another is hydrophilic, permitting them to interact with membranes.

**[0047]** The term “subcellular-localization component”, as used herein, refers to a molecule capable of recognizing a subcellular component in a targeted cell. Recognized subcellular components include the nucleus, ribosomes, mitochondria, and chloroplasts. Particular subcellular-localization components include the “nuclear-localization components” that aid in carrying molecules into the nucleus and are known to include the nuclear localization peptides and amino acid sequences.

**[0048]** The bioactive component **12** may be included at an amount that is therapeutically effective to transform a plurality of cells, such as in vitro, in vivo or ex vivo cells. As used herein, “transform” refers to a presence and/or functional activity of the bioactive component in the plurality of cells after exposing the nanocapsules to the plurality of cells.

**[0049]** Furthermore, those of ordinary skill in the art will recognize that the amount of the bioactive component **12** may vary depending upon the bioactive component **12**, the temperature, pH, osmolarity, any solutes, any additional component or optional solvents present in the second aqueous component **30**, the surfactant composition **16**, a type or an amount of the surfactant micelle **22**, the biocompatible polymer component **24**, any desired characteristics of the stabilized surfactant micelle **28**, any desired characteristics of the nanocapsules **36**, or any combination of any of these.

**[0050]** The bioactive component **12** of the nanocapsule **36** maybe supplied as an individual macromolecule or supplied in various prepared mixtures of two or more macromolecules that are subsequently combined to form the bioactive component **12**. Some non-exhaustive examples of hydrophilic macromolecules that may be suitable for inclusion as part of the bioactive component **12** include, but are not limited to polynucleotides, polypeptides, genetic material, peptide nucleic acids, aptamers, carbohydrates, mini-chromosomes, molecular polymers, aggregates or associations of an inorganic or organic nature, genes, any other hydrophilic macromolecule or any combination of any of these.

**[0051]** Some non-exhaustive examples of hydrophobic macromolecules that may be included part of the bioactive component **12** include, but are not limited to, adrenergic, adrenocortical steroid, adrenocortical suppressant, aldosterone antagonist, and anabolic agents; analeptic, analgesic, anesthetic, anorectic, and anti-acne agents; anti-adrenergic, anti-allergic, anti-amebic, anti-anemic, and anti-anginal agents; anti-arthritis, anti-asthmatic, anti-atherosclerotic,

antibacterial, and anticholinergic agents; anticoagulant, anti-convulsant, antidepressant, antidiabetic, and antidiarrheal agents; antidiuretic, anti-emetic, anti-epileptic, antifibrinolytic, and antifungal agent; antihemorrhagic, inflammatory, antimicrobial, antimigraine, and antimiotic agents; antimycotic, antinauseant, antineoplastic, antineutropenic, and antiparasitic agents; antiproliferative, antipsychotic, antirheumatic, antiseborrheic, and antisecretory agents; antispasmodic, antithrombotic, antiulcerative, antiviral, and appetite suppressant agents; blood glucose regulator, bone resorption inhibitor, bronchodilator, cardiovascular, and cholinergic agents; fluorescent, free oxygen radical scavenger, gastrointestinal motility effector, glucocorticoid, and hair growth stimulant agent; hemostatic, histamine H2 receptor antagonists; hormone; hypocholesterolemic, and hypoglycemic agents; hypolipidemic, hypotensive, and imaging agents, immunizing and agonist agents; mood regulators, mucolytic, mydriatic, or nasal decongestant; neuromuscular blocking agents; neuroprotective, NMDA antagonist, non-hormonal sterol derivative, plasminogen activator, and platelet activating factor antagonist agent; platelet aggregation inhibitor, psychotropic, radioactive, scabicide, and sclerosing agents; sedative, sedative-hypnotic, selective adenosine A1 antagonist, serotonin antagonist, and serotonin inhibitor agent; serotonin receptor antagonist, steroid, thyroid hormone, thyroid hormone, and thyroid inhibitor agent; thyromimetic, tranquilizer, amyotrophic lateral sclerosis, cerebral ischemia, and Paget’s disease agent; unstable angina, vasoconstrictor, vasodilator, wound healing, and xanthine oxidase inhibitor agent; immunological agents, antigens from pathogens, such as viruses, bacteria, fungi and parasites, optionally in the form of whole inactivated organisms, peptides, proteins, glycoproteins, carbohydrates, or combinations thereof, any examples of pharmacological or immunological agents that fall within the above-mentioned categories and that have been approved for human use that may be found in the published literature, any other hydrophobic bioactive component, or any combination of any of these.

**[0052]** As used herein, the term “polypeptide” refers to a polymer of amino acids not limited by the number of amino acids. It is also to be understood that the term “polypeptide” is meant to encompass an oligopeptide, a peptide, or a protein, for example.

**[0053]** As used herein, the term “polynucleotide” refers to RNA or DNA sequences of more than 1 nucleotide in either single chain, duplex or multiple chain form. The term “polynucleotide” is also meant to encompass polydeoxyribonucleotides containing 2'-deoxy-D-ribose or modified forms thereof, RNA and any other type of polynucleotide which is an N-glycoside or C-glycoside of a purine or pyrimidine base, or modified purine or pyrimidine base or basic nucleotide. The polynucleotide may encode promoter regions, operator regions, structural regions, termination regions, combinations thereof or any other genetically relevant material. Similarly, the term “genetic” as used herein, refers to any material capable of modifying gene expression.

**[0054]** The first aqueous composition **14** may be included in the method of the present invention as a gel, liquid, or in vapor form. The form of the first aqueous composition **14** that is selected preferably permits the first aqueous composition **14** to (1) remain stable prior to dissolving or dispersing the bioactive component, the surfactant composition **16**,

the surfactant micelle 22, or optionally the stabilizer surfactant micelle 28, (2) homogeneously disperse the bioactive component 12, the surfactant composition 16, the surfactant micelle 22, or optionally the stabilizer surfactant 28, (3) function as a continuous phase in an oil-in-water emulsion, (4) not interfere with, or mask the functional activity of the bioactive component 12, and (5) not modify or degrade the bioactive component 12, the surfactant composition 16, the surfactant micelle 22, or optionally the stabilized surfactant micelle 28.

[0055] The first aqueous composition 14 may include only water, or may optionally include additional solutes or solvents that do not interfere with the method of forming the nanocapsules 36 nor mask the functional activity of the bioactive component 12. Furthermore, those of ordinary skill in the art will recognize that an amount of the first aqueous composition 14 used to prepare the nanocapsules 36 may vary depending upon the bioactive component 12, the surfactant composition 16, the temperature, pH, osmolarity, optional solutes or optional solvents, the surfactant micelle 22, the biocompatible polymer component 24, any desired characteristics of the stabilized surfactant micelle 28 or the nanocapsules 36.

[0056] The bioactive component 12 may be added to the first aqueous composition 14 or the first aqueous composition 14 may be added to the bioactive component 12. While the order of addition of the bioactive component 12 and the first aqueous composition 14 is not critical to the present invention, the hydrophilic composition (not shown) that is formed when the bioactive component 12 is dissolved or dispersed in the first aqueous composition 14 is preferably capable of maintaining a homogeneous solution or dispersion in the hydrophilic composition.

[0057] The first aqueous composition 14 may be supplied as an individual component or supplied in various prepared mixtures of two or more components that are subsequently combined to form the first aqueous composition 14. Some non-exhaustive examples of the first aqueous composition 14 include, but are not limited to, the above-mentioned water, nonpolar solvents, polar solvents, or any combination of any of these. Preferably, water is the first aqueous composition 14.

[0058] The surfactant composition 16 maybe introduced into the bioactive component 12, the first aqueous composition 14, the hydrophilic composition as a liquid, vapor or in granular form. The form of the surfactant composition 16 that is selected preferably permits the surfactant composition 16 to (1) remain stable prior to introducing into the bioactive component 12, the first aqueous composition 14, or the hydrophilic composition, (2) be homogeneously dispersed into the bioactive component 12, the first aqueous composition 14, or the hydrophilic composition, (3) form a micellar structure, (4) be adsorbed onto a surface of the bioactive component 12, the first aqueous composition 14, the hydrophilic composition (5) displace the first aqueous composition that is located on the surface of the bioactive component 12, (6) partition the bioactive component 12 or the hydrophilic composition into a core of the micellar structure to form the surfactant micelle 22, and (7) provide a thermodynamic driving force that is effective to reduce a size of the bioactive component 12, surfactant micelle 22, the stabilized surfactant 28 or the nanocapsule 36.

[0059] As used herein, a "surfactant" refers to any molecule containing a polar portion that thermodynamically prefers to be solvated by a polar solvent, and a hydrocarbon portion that thermodynamically prefers to be solvated by a non-polar solvent. The term "surfactant" is also meant to encompass anionic, cationic, or non-ionic surfactants. As used herein, the term "anionic surfactant" refers to a surfactant with a polar portion that ionizes to form an anion in aqueous solution. Similarly, a "cationic surfactant" refers to a surfactant having a cationic polar portion that ionizes to form a cation in aqueous solution. Likewise, a "non-ionic" surfactant refers to a surfactant having a polar portion that does not ionize in aqueous solution.

[0060] While not wanting to be bound to theory, it is generally believed that a surfactant refers to a molecule that is effective to reduce a surface or an interfacial tension between a first substance dispersed in a second substance such that the first substance is solvated and any molecular groups of the first substance are dispersed. Typically, a hydrodynamic diameter of the first substance increases after addition of the surfactant. Nonetheless, the surfactant composition 16 is believed to be effective to reduce the size or diameter of the surfactant micelles 22 in the first aqueous composition 14, to thereby reduce the size of the nanocapsule 36 when practicing the present invention.

[0061] The surfactant composition 16 may include the surfactant component only (not shown), or may optionally include the biocompatible oil component 18. The surfactant component may be characterized on the HLB scale that ranges from less than about 1 to more than about 13 units.

[0062] A surfactant component having an HLB value of less than about 6.0 units may be described as being poorly, or not dispersable in an aqueous or water-based composition. In addition, a surfactant component having an HLB value of less than about 6.0 units may be characterized as a hydrophobic or non-ionic surfactant. A surfactant component having an HLB value of more than about 7.0 units may be described as being capable of forming a milky to translucent to clear dispersion when the surfactant having an HLB value of more than about 7.0 units is dispersed in an aqueous or water-based composition.

[0063] Preferably, the surfactant component of the surfactant composition 16 has an HLB value of less than about 6.0 units when practicing the method of the present invention. Still more preferably the surfactant component of the surfactant composition 16 has an HLB value of less than about 5.0 units to facilitate preparation of nanocapsules having a diameter of less than about 50 nm.

[0064] The surfactant component may also be characterized in terms of a critical micelle concentration (CMC) value. Preferably, the surfactant component of the surfactant composition 16 has a CMC value of less than about 300 micromolars ( $\mu\text{M}$ ). Still more preferably, the surfactant component has a CMC value of less than about 200  $\mu\text{M}$ .

[0065] While not wanting to be bound to theory, it is believed that the surfactant component of the surfactant composition 16 adsorbs onto the surface of the bioactive component 12 when introduced into the first aqueous composition 14 to minimize exposure of a surface of the hydrophobic surfactant component to a thermodynamically unfavorable environment created by the first aqueous

composition **14**. Therefore, the surfactant component adsorbs onto the surface of the bioactive component to reduce the surface area of the surfactant component that may be exposed to the first aqueous composition **14**. Adsorption of the surfactant component onto the bioactive component **12** is believed to facilitate the size reduction of the bioactive component **12** and/or the surfactant micelle **22**.

[0066] The surfactant component of the surfactant composition **16** may be supplied as individual surfactants or supplied in various prepared mixtures of two or more surfactants that are subsequently combined to form the surfactant composition **16**. Some non-exhaustive examples of suitable surfactants having an HLB value of less than about 6.0 units or a CMC value of less than about 200  $\mu\text{m}$  be listed in *Dermatological Formulations* (Barry, B., Marcel Dekker, (1983)), or in *Percutaneous absorption: drug, cosmetics, mechanisms, methodology*, 3<sup>rd</sup> ed., Bronough, R. ed., 1999, or the *Handbook of Industrial Surfactants* (Ash, M., Ed., Gower Pub. (1993), which are incorporated herein by reference. As an example, the surfactant component may be 2, 4, 7, 9-tetramethyl-5-decyn-4, 7-diol(TM-diol), blends of 2, 4, 7, 9-tetramethyl-5-decyn-4, 7-diol(TM-diol), molecules having one or more acetylenic diol groups, cetyl alcohol or any combination of any of these.

[0067] The optional biocompatible oil component **18** of the surfactant composition **16** may be combined with the surfactant component as a liquid, vapor or in granular form. The form of the optional biocompatible oil component **18** that is selected preferably permits the optional biocompatible oil component **18** to (1) remain stable prior to introduction into the surfactant composition **16**, (2) be homogeneously blended into the surfactant composition **16**, (3) dissolve or disperse the surfactant component, and (4) increase the hydrophobicity of the surfactant composition **16**, and therefore, the degree to which the size of the bioactive component **12**, the surfactant micelle **22**, the stabilizer surfactant micelle **28**, or the nanocapsule **36** may be reduced when practicing the present invention.

[0068] Preferably, the concentration of the optional biocompatible oil component **18** in the surfactant composition **16** ranges from about  $10^{-7}$  weight percent to about 10 weight percent, based upon a total volume of the stabilized surfactant micelles **28** in the first aqueous composition **14**. Concentrations of the optional biocompatible oil component **18** higher than about 10 weight percent, based upon the total volume of the surfactant composition **18**, may be less desirable because such higher concentrations increase a phase volume of the biocompatible oil, and consequently may cause difficulties in preparing, dispersing and/or handling the surfactant micelles **22**, the stabilized surfactant micelles **28** or the nanocapsules **36**. Concentrations of the optional biocompatible oil component lower than about  $10^{-7}$  weight percent in the surfactant composition **16** may be less preferred, because such lower concentrations would not be effective to solvate the surfactant component, or increase the hydrophobicity of the surfactant composition **16**, and may ultimately increase the diameter of the nanocapsules **36**.

[0069] The optional biocompatible oil component **18** of the surfactant composition **16** may be supplied as an individual biocompatible oil or supplied in various prepared mixtures of two or more biocompatible oils that are subsequently combined to form the optional biocompatible oil

component **18**. Some non-exhaustive examples of suitable biocompatible oils that may be included as part of the biocompatible oil component **18** may be found in *Dermatological Formulations* (Barry, B., Marcel Dekker, (1983)), or in *Percutaneous absorption: drug, cosmetics, mechanisms, methodology*, 3<sup>rd</sup> ed., Bronough, R. ed., 1999, or in the *Handbook of Industrial Surfactants* (Ash, M., Ed., Gower Pub. (1993), which have been incorporated herein by reference. Preferably, food or USP grade oils, such as DMSO, DMF, castor oil, or any combination thereof, are used to practice the present method.

[0070] The surfactant composition **16** may be included at an amount that is effective to form the micellar structure that partitions the bioactive component **12**, the first aqueous composition **14** or the hydrophilic composition into the core of the micellar structure when forming the surfactant micelle **22**. Still more preferably, the surfactant composition **16** is included at an amount that is effective to provide a maximum thermodynamic driving force that minimizes the size of the bioactive component **12**, the surfactant micelle **22**, and ultimately, the size of the nanocapsule **36** when practicing the present invention.

[0071] Furthermore, those of ordinary skill in the art will recognize that the amount of the surfactant composition **16** may be varied based upon the bioactive component **12**, the first aqueous composition **14**, a ratio of the surfactant component to the optional biocompatible oil **18**, any desired characteristics of the surfactant micelles **22**, the stabilized surfactant micelles **28** or the nanocapsules **36**. For example, a surfactant composition containing a surfactant component having an HLB value of about 6.0 units mixed with a nonpolar biocompatible oil like castor oil, may provide the same degree of a thermodynamic driving force as a second surfactant composition containing a surfactant component of about 4.0 units mixed with DMSO.

[0072] The amount of the surfactant composition **16** may range up to about 0.5 weight percent, based upon a total volume of the stabilized surfactant micelles dispersed in the first aqueous composition **14**. Still more preferably, the amount of the surfactant composition **16** is less than about 0.25 weight percent, based upon the total volume of the stabilized surfactant micelles **28** dispersed in the first aqueous composition **14**. Most preferably, the surfactant composition **16** is present at an amount of less than about 0.05 weight percent, based upon the total volume of the stabilized surfactant micelles **28** dispersed in the first aqueous composition **14**. As one non-exhaustive example, the surfactant composition **16** may be added to the total volume of the hydrophilic composition at a concentration of about 500 ppm, based on the total volume of the stabilized surfactant micelles **28** in the first aqueous composition **14**.

[0073] The first dispersing apparatus **20** initiates and promotes formation of the micellar structures that are based on the bioactive component **12**, the first aqueous composition **14** and the surfactant composition **16**. Adsorption of surfactant component onto the surface of the bioactive component **12**, or hydrophilic composition continues until all of the surfactant molecules cover, and therefore, entrap the bioactive component **12** or hydrophilic composition in the core of the micellar structure to form surfactant micelles **22**. Formation of a plurality of surfactant micelles **22** in the first aqueous composition **14** forms a dispersion of surfactant micelles **22**.

[0074] In general, any conventional dispersing apparatus 20 that is capable of homogeneously blending or dispersing may be suitable for use in forming the dispersion of surfactant micelles in accordance with the present invention. Furthermore, those of ordinary skill in the art will recognize that the first dispersing apparatus 20 may vary depending upon the desired characteristics of the nanocapsules 36. For example, the first dispersing apparatus 20 may include any device, such as a sonicating or a vortexing apparatus (not shown), or the like to disperse the bioactive component 12 in the hydrophilic composition, and the formation of the surfactant micelles 22 after addition of the surfactant composition 16. Nonetheless, while the first dispersing apparatus 20 may include a sonicating or a vortexing apparatus, the sonicating or the vortexing apparatus is not critical when practicing the method of the present invention.

[0075] As used herein, a “surfactant micelle” may be characterized as a close packed mono-molecular barrier of surfactant molecules at an interface between the bioactive composition 12 and the surfactant composition 16, such that the barrier encapsulates the bioactive component 12, the first aqueous composition 14 or the hydrophilic composition. It is also to be understood that the term “surfactant micelle” encompasses partial or hemi-surfactant micelles that partially enclose the bioactive component 12, the first aqueous composition 14 or the hydrophilic composition.

[0076] When the bioactive component 12 is a hydrophilic bioactive component, the polar portion of the surfactant molecule associates with a surface of the hydrophilic bioactive component. When the bioactive component 12 is a hydrophobic bioactive component, the hydrocarbon portion of the surfactant micelle associates with a surface of the hydrophobic bioactive component.

[0077] The formation of a surfactant micelle typically occurs at a well defined concentration known as the critical micelle concentration. As noted, surfactant components having a CMC value of less than about 200 micromolars are preferred when practicing the present invention.

[0078] After forming the dispersion of surfactant micelles 22, the dispersion of surfactant micelles 22 is transferred into the stabilizing apparatus 26 where a biocompatible polymer component 24 is added to stabilize the dispersion of surfactant micelles 22. Alternatively, the biocompatible polymer component 24 may be added to the dispersion of surfactant micelles 22 in the first dispersing apparatus 20 which obviates the need for the stabilizing apparatus 26.

[0079] The biocompatible polymer component 24 stabilizes the dispersion of surfactant micelles 22 to form stabilized surfactant micelles 28 within the first aqueous composition 14. Therefore, a dispersion of stabilized surfactant micelles 28 are present within the first aqueous composition 14 after addition of the biocompatible polymer component 24.

[0080] As used herein, the term “biocompatible” refers to a material that is capable of interacting with a biological system without causing cytotoxicity, undesired protein or nucleic acid modification or activation of an immune response.

[0081] The biocompatible polymer component 24 may be introduced into the dispersion of surfactant micelles 22 as a liquid, vapor or in granular form. The form of the biocom-

patible polymer component 24 that is selected preferably permits the biocompatible polymer component 24 to (1) remain stable prior to addition into the dispersion of surfactant micelles 22, (2) be homogeneously dispersed into the dispersion of surfactant micelles 22, (3) increase a viscosity of the first aqueous composition 14, (4) form a boundary layer at an interface of the surfactant micelle 22 and the first aqueous composition 14, (5) be absorbed onto a surface of the surfactant micelles 22, (6) be capable of iontophoretic exchange, (7) be capable of being precipitated upon addition of a solute, (8) be capable of enzymatic degradation, surface and/or bulk erosion, (9) not interfere with or mask the functional activity of the bioactive component 12, (10) prevent aggregation and/or agglomeration of the dispersion of surfactant micelles 22, and (11) be capable of obtaining a particular dissolution profile.

[0082] The biocompatible polymer component 24 may be included at an amount that is effective to coat and therefore stabilize the surfactant micelle 22. Furthermore, those of ordinary skill in the art will recognize that the amount of the biocompatible polymer component 24 used to stabilize the surfactant micelles 22 may vary depending upon the bioactive component 12, the first aqueous composition 14, the surfactant composition 16, the temperature, pH, osmolarity, presence of any optional solutes or optional solvents, the surfactant micelle 22, any desired characteristics of the stabilized surfactant micelle 28, the nanocapsules 36, or a desired dissolution profile.

[0083] While the concentration of the biocompatible polymer component 24 is not critical to the present invention, the concentration of the biocompatible polymer component 24 is preferably based upon the bioactive component and on the desired dissolution profile. When the concentration of the biocompatible polymer component 24 is too high, the shell of the nanocapsule 36 may not dissolve. If the concentration of the biocompatible polymer component 24 is too low, the shell of the nanocapsule 36 may dissolve rapidly in a manner that promotes cytotoxicity, for example. In addition, too low a concentration of the biopolymer component 24 may not provide an effective degree of mechanical force to stabilize the surfactant micelles 28.

[0084] Concentrations of the biocompatible polymer component 24 that are too high may also be less desirable because such higher concentrations may increase the viscosity of the first aqueous composition 14, and consequently may cause difficulties in preparing, mixing and/or transferring the stabilizer surfactant micelles 28. Concentrations of the biocompatible polymer component 24 that are too low may be less preferred, because lower concentrations may not provide the needed viscosity to stabilize the surfactant micelles, nor be capable of effectively coating the surfactant micelles 22 to prevent aggregation of the surfactant micelles 22 in the first aqueous composition 14.

[0085] The biocompatible polymer component 24 may be supplied as individual biocompatible polymers or supplied in various prepared mixtures of two or more biocompatible polymers that are subsequently combined to form the biocompatible polymer component 18. Some non-exhaustive examples of biocompatible polymers include polyamides, polycarbonates, polyalkylenes, polyalkylene glycols, polyalkylene oxides, polyalkylene terephthalates, polyvinyl alcohols, polyvinyl ethers, polyvinyl esters, polyvinyl halides,

polyvinylpyrrolidone, polyglycolides, polysiloxanes, polyurethanes and copolymers thereof, alkyl cellulose, hydroxy-alkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, polymers of acrylic and methacrylic esters, methyl cellulose, ethyl cellulose, hydroxypropyl cellulose, hydroxy-propyl methyl cellulose, hydroxybutyl methyl cellulose, cellulose acetate, cellulose propionate, cellulose acetate butyrate, cellulose acetate phthalate, carboxylethyl cellulose, cellulose triacetate, cellulose sulphate sodium salt, poly(methyl methacrylate), poly(ethylmethacrylate), poly(butylmethacrylate), poly(isobutylmethacrylate), poly(hexylmethacrylate), poly(isodecylmethacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), polyethylene, polypropylene poly(ethylene glycol), poly(ethylene oxide), poly(ethylene terephthalate), poly(vinyl alcohols), poly(vinyl acetate, poly vinyl chloride polystyrene, polyvinylpyrrolidone, polyhyaluronic acids, casein, gelatin, glutin, polyanhydrides, polyacrylic acid, alginate, chitosan, poly(methyl methacrylates), poly(ethyl methacrylates), poly(butylmethacrylate), poly(isobutylmethacrylate), poly(hexylmethacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate) and combination of any of these.

[0086] Additionally, biocompatible polymers that have been modified for enzymatic degradation, or change upon application of light, ultrasonic energy, radiation, a change in temperature, pH, osmolarity, solute or solvent concentration may also be included as part of the biocompatible polymer component 24. Preferably, the biocompatible polymer component 24 is a hydrophilic polymer that is capable of substantially coating, and preferably continuously coating the surfactant micelle 22. Still more preferably, the hydrophilic biocompatible polymer component 24 is capable of iontophoretic exchange.

[0087] Though descriptions of the present invention are primarily made in terms of a hydrophilic biocompatible polymer component 24, it is to be understood that any other biocompatible polymer, such as hydrophobic biocompatible polymers may be substituted in place of the hydrophilic biocompatible polymer, in accordance with the present invention, while still realizing benefits of the present invention. Likewise, it is to be understood that any combination of any biocompatible polymer may be included in accordance with the present invention, while still realizing benefits of the present invention.

[0088] In general, any conventional apparatus and technique that is suitable for permitting the biocompatible polymer component 24 to stabilize the surfactant micelles 22 may be used as the stabilizing apparatus 26 in accordance with the present invention. Furthermore, any other device, such as high pressure homogenization or high ultrasound sonication is preferably not included during stabilization.

[0089] After stabilizing the surfactant micelles 22, the stabilized surfactant micelles 28 may be transferred into a second aqueous composition 30 located in a second dispersing apparatus 32. The stabilized surfactant micelles 28 may be transferred by mechanically forming droplets of the stabilized surfactant micelle 28 that are subsequently introduced into the second aqueous composition 30.

[0090] The second aqueous composition 30 may include water only, or may optionally include a solute to precipitate the biocompatible polymer component 24 surrounding the stabilized surfactant micelle 28. Some non-exhaustive examples of solutes that may be used to precipitate the biocompatible polymer 24 include ionic species derived from elements listed in the periodic table.

[0091] Preferably, the second aqueous composition 30 includes a solute in an amount that is effective to precipitate the biocompatible polymer component 24 and form the dispersed, and optionally atomized nanocapsules 36 of the present invention. As used herein, the term "precipitate" refers to a solidifying or a hardening of the biocompatible polymer component 24 that surrounds the stabilized surfactant micelles 28. It is also to be understood that the term "precipitation" is also meant to encompass any crystallization of the biocompatible polymer 24 that may occur when the biocompatible polymer component 24 is exposed to the solute.

[0092] Additionally, any other component that is capable of modulating the efficacy the nanocapsules 36 may be included as part of the second aqueous composition to thereby modulate the functional activity of the nanocapsule 36. For example, the second aqueous composition may include additional coating excipients, such as a cell recognition component or various ionic species, such as  $Mn^{2+}$ ,  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Al^{3+}$ ,  $Be^{2+}$ ,  $Li^{30}$ ,  $Ba^{2+}$ ,  $Gd^{3+}$ , or any other ionic species that is capable of interacting with the biocompatible polymer component 24.

[0093] The term "cell recognition component", as used herein, refers to a molecule capable of recognizing a component on a surface of a targeted cell. Cell recognition components may include an antibody to a cell surface antigen, a ligand for a cell surface receptor, such as cell surface receptors involved in receptor-mediated endocytosis, peptide hormones, and the like.

[0094] It has been observed that when the stabilized surfactant micelles 28 are allowed to incubate in the second aqueous composition 30 that includes the solute to precipitate the biocompatible polymer component 24, the nanocapsules 36 undergo a reduction in size. Furthermore, the formation of a flocculated suspension of the nanocapsules 36 has also been observed after incubating the stabilized surfactant micelles 28 in the second aqueous composition.

[0095] As used herein, "a flocculated suspension" refers to the formation of a loose aggregation of discrete particles held together in a network-like structure either by physical absorption of bioactive components, bridging during chemical interaction (precipitation), or when longer range van der Waals forces of attraction exceed shorter range forces of repulsion. The flocculated suspension of nanocapsules 36 may entrap varying amounts of the first aqueous composition 14 or the second aqueous composition 30 within the network-like structure. Additionally, the flocculated suspension of nanocapsules may be gently tapped to disperse the nanocapsules 36.

[0096] The stabilized surfactant micelles 28 may be transferred into the second aqueous composition 30 via atomization through a nozzle (not shown) having a particular orifice size or through an aerosolizing apparatus (not shown). Atomizing or aerosolizing the stabilized surfactant



micelles **28** typically includes the application of a shear force that may be capable of further dispersing the stabilized surfactant micelles **28**. Furthermore, the application of the shear force during transfer may also be effective to (1) reduce the size of the nanocapsules **36**, or (2) break up any agglomerates or associations between stabilized surfactant micelles **28** that may have formed in the stabilizing apparatus **26**. Nozzle pressures of less than about 100 psi, for example, may be used to atomize the stabilized surfactant micelles **28**.

[0097] The diameter of the nanocapsules **36** may also be varied depending upon the orifice size of the nozzle that may be used to transfer the stabilized surfactant micelles **28** into the second aqueous composition. Alternatively, the stabilized surfactant micelles **28** may be added to the second aqueous composition **30** containing the solute that precipitates the biocompatible polymer **24** to form a dispersion of nanocapsules **36** for purposes of providing the dispersion for sub-cutaneous delivery of the nanocapsules, for example.

[0098] After precipitating and/or optionally incubating the nanocapsules **36** in the second aqueous composition **30**, the nanocapsules **36** may be filtered, centrifuged or dried to obtain separate and discrete nanocapsules **36**. The nanocapsules **36** may be frozen or reconstituted for later use or may be delivered to a target cell or tissue by such routes of administration as oral, intravenous, subcutaneous, intraperitoneal, intrathecal, intramuscular, inhalational, topical, transdermal, suppository (rectal), pessary (vaginal), intra-urethral, intraportal, intrahepatic, intra-arterial, intra-ocular, transtympanic, intraocular, intrathecal, or any combination of any of these.

[0099] The nanocapsules **36** having a diameter of less than about 50 nm are advantageous in the delivery of bioactive components to target cells for several reasons. First, nanocapsules **36** having a diameter of less than about 50 nm enhances delivery of bioactive components by protecting the bioactive components against degradation during transport to the target cell.

[0100] Second, nanocapsules **36** having a diameter of less than about 50 nm promotes efficient cellular uptake. Efficient cellular uptake into the target cell typically occurs when a particle has a diameter of less than about 50 nm, as opposed to when a particle has a diameter of more than about 50 nm.

[0101] Third, it is believed that uptake of the nanocapsules **36** by the target cell occurs via transport systems, such as a non-endosomal pathway, that prevents lysosomal degradation of the nanocapsules **36**. Indeed, it is believed that the nanocapsules **36** of the present invention are efficiently exported into a cell via a caveolin-regulated pathway that circumvents most, if not all, endosomal-regulated pathways that typically degrade nanocapsules **36**.

[0102] Fourth, the nanocapsules **36** have a biocompatible polymer shell that is separate from the bioactive component. In fact, the bioactive component is not entangled in, embedded in, or adsorbed onto the biocompatible polymer shell of the nanocapsules **36**. When the bioactive component is not entangled in, embedded in, or adsorbed onto the biocompatible polymer shell, the cell that incorporate the nanocapsules **36** avoid apoptosis or cell death.

[0103] Fifth, enclosing the bioactive component within a core surrounded by the biocompatible polymer shell when

preparing the nanocapsules **36** in accordance with the present method is advantageous in avoiding premature degradation of the nanocapsules **36**, or a cytotoxic response during in vivo transport of the nanocapsule. Enclosing the bioactive component within the core results in a linear release rate of the bioactive component without any zero burst phenomenon during release from the nanocapsules **36**.

[0104] The linear release rate of the bioactive component from the nanocapsule without any zero burst phenomenon is also an advantageous feature as the linear release rate allows rational design of coating dissolution profiles to minimize cytotoxicity. As used herein, the term "dissolution profile" refers to a rate at which the biocompatible polymer shell is dissolved or degraded to release a bioactive agent from a core of a nanocapsule.

[0105] Another benefit of the nanocapsules **36** prepared by the method of the present invention is that little, if any, addition of an organic solvent is required to form the nanocapsules **36**. Eliminating the use of most, if not all, organic solvents from the method of the present invention is beneficial since organic solvents may damage the bioactive component **12**, destroy the target cells, or be toxic during preparation of the nanocapsule **36**. The elimination of most, if not all, use of organic solvents eliminates the need for complex solvent removal techniques, such as solvent dilution, vacuum evaporation, or the like, and obviates any associated costs or complex process strategies during preparation of the nanocapsules **36**.

[0106] The nanocapsules **36** of the present invention further permits stable encapsulation of a bioactive component, and in particular, hydrophilic bioactive components, such as polynucleotides and polypeptides. "Stable encapsulation", as used herein, refers to maintenance of the encapsulated bioactive component's structure. For nucleic acids, the appearance of low molecular weight nucleic acid breakdown products, which may be assayed for by electrophoresis, is substantially eliminated. The nanocapsules **36** may also be used to encapsulate any bioactive component regardless of water solubility or charge density.

## APPLICATIONS

[0107] The nanocapsules **36** may be combined with additional polymeric binders, surfactants, fillers, and other excipients to incorporate the nanocapsules **36** into solid dosage forms such as granules, tablets, pellets, films or coatings for use in enhanced bioactive component **12** delivery. In this way, design of the dissolution profile, control of the particle size, and cellular uptake remains at the level of the nanocapsule. Such applications include, but are not limited to, creation of rapidly dissolving nanocapsule pellets for pulmonary delivery or nanocapsule films for device-mediated delivery.

[0108] In another application, the nanocapsules **36** may be designed for specific cellular or tissue uptake by polymer selection and/or inclusion of cell-recognition components in the nanocapsule biocompatible polymer shell or coating. Such coatings will have utility for specific or increased delivery of the bioactive agent to the target cell. Such applications include, but are not limited to tumor-targeting of chemotherapeutic agents or anti-sense DNA, antigen delivery to antigen-presenting cells, ocular delivery of

ribozymes to retinal cells, transdermal delivery of protein antibodies, or transtympanic membrane delivery of peptide nucleic acids.

**[0109]** Property Determination and Characterization Techniques

**[0110]** Various analytical techniques are employed herein. An explanation of these techniques follows:

**[0111] FIG. 1A:** Samples were prepared on freshly cleaved mica as dispensed, dried in air and imaged using a Nanoscope II multimode AFM (Digital Instruments) with a J type scanner and ambient tapping mode holder. 125  $\mu\text{m}$  long silicon cantilevers type IBMSC were from IBM and have resonant frequencies of 250-450 kHz. All imaging was in tapping mode, images were 512x512 pixels and scanning frequency was 1 kHz. Height, amplitude and phase images were collected. Images were processed in DI software and analyzed in NIH Image SXM. A: Formula Q from 2-phase system, low HLB surfactant, B: Formula S from 2-phase system, high HLB surfactant, C: Formula T from 1-phase system, high HLB surfactant, D: Formula V from 2-phase system, surfactant below CMC.

**[0112] FIG. 1B:** Nanocapsules were released into a solution of 10% isobutanol in Phosphate-buffered Saline (PBS), pH=7.2. Samples were run in duplicate. **FIG. 1C:** Nominal 300 ng samples of DNA were aliquoted from a master batch containing surfactant and processed through commercial miniprep columns. Eluate was recycled through Qiaquick columns and collected either 3 times (4, 5) or twice (6,7) or recycled through Zymoclean columns and collected twice (8,9). Samples were alcohol precipitated using a commercial coprecipitant, electrophoresed on 1.5% agarose gels modified with Synergel, stained with SybrGold dye, digitized on a Storm 860 and compared to unmodified but reprecipitated samples from the same master batch (10,11). Lanes 1-3: 100, 50 and 5 ng of ?\_DNA.

**[0113] FIG. 2:** Endocytic activity was assessed by immunosignal levels of clathrin (Chemicon). Potocytotic activity was assessed by immunosignal for caveolin-1 as described in the literature (Transduction Laboratories). Lysosomal activity was detected by a monoclonal antibody to Lamp-1 (Transduction Laboratories). Nanocapsule localization was detected by streptavidin-biotin immunocomplexes directed against sheep IgG (Jackson Laboratories). Nanocapsule coatings were spiked with ovine IgG to enable this detection strategy.

**[0114] FIG. 3:** immortalized Rt-1 fibroblast cultures at 70% confluence were treated for 4 days with increasing amounts of nanocapsule formula K and transiently treated (3 hours) with an optimized liposomal formula (dose, 500 ng) Results are expressed as a percentage of cellular actin integrated intensity and compared to the liposomal formula. Expression vector was code 448: pEF/myc-his/GFP (Invitrogen).

**[0115] FIG. 4A:** Radiated porcine biopsies were snapfrozen 7 days after treatment with saline or 6  $\mu\text{g}$  of controlled release nanocapsules, then homogenized in RIPA. 100  $\mu\text{g}$  lysate samples were electrophoresed on SDS-Page gradient gels, transferred to nitocellulose membranes and detected for either  $\beta$ -galactosidase (121 Kd) or involucrin (~100 kD) using chemiluminescence. Results were normalized to the post-transfer gel stained with Coomassie due to interference

at 100 kD from a gel defect. Involucrin, a component of the cornified membrane, manufactured by suprabasal cells can be detected in radiated porcine skin and will be used for future normalization purposes. Lane A: N, topical, biopsy oc-2; B: N, topical, biopsy oc-3; C: 0, topical, biopsy 1-1; D: PBS only, biopsy 1-5; E: N, subcutaneous injection, biopsy 1-6.

**[0116] FIG. 4B:** The  $\beta$ -galactosidase reporter protein was detected by a monoclonal antibody directed at an incorporated fusion protein tag. A: N, topical, biopsy oc-1, detection with anti-Xpress<sup>®</sup>; B: Matching view to A with detection for anti-von Willenbrand factor (Sigma); C: untreated biopsy, detection with anti-Xpress<sup>®</sup>(Invitrogen).

**[0117] FIG. 5:** Nanocapsules were incorporated into an aqueous suture coating and sutures were applied to pigskin biopsies in organ culture. Nanocapsules were detected with Cy3 conjugated-streptavidin-biotin complexes to incorporate ovine IgG and GFP transgene expression was detected by rabbit polyclonal antibodies to GFP (Abcom) in combination with Fitec-conjugated polyclonal antibodies to rabbit IgG and Alexa 488-conjugated polyclonal antibodies to Fitec (Molecular Probes). Controls omitting primary antibodies were included for signal-to-background level estimation.

**[0118] FIG. 6A:** Nanocapsules were detected as previously described and GFP transgene expression was detected by rabbit polyclonal antibodies to GFP in combination with Cy3-conjugated antibodies to rabbit IgG (Jackson Laboratories).

**[0119] FIG. 6B:** GFP expression was detected as described in **FIG. 5** and cell nuclei were counterstained with 10  $\mu\text{g}/\text{ml}$  bisbenzamide.

**[0120] FIG. 6C:** Carcinoma cells and HDF's were seeded overnight into 96 well plates at 2000 and 6000 cells per well respectively. Cisplatin preparations were added to wells for 18 hours as noted on the graph than washed out. After 72 hours cell viability of assessed by a commercial MTT assay (WST assay, Boehringer Mannheim). Wells were executed in duplicate.

**[0121] FIG. 7:** Colocalization with lysosomes was detected using a monoclonal antibody to Lamp-1 (Transduction Laboratories). AFM images are included of O-methyl RNA formulated by nanoencapsulation or complexation with 27 KD polyethyleneimine.

## EXAMPLES

**[0122]** The present invention is more particularly described in the following Examples which are intended as illustrations only since numerous modifications and variations within the scope of the present invention will be apparent to those skilled in the art.

**[0123]** Reagents:

**[0124]** A. Nucleic acid condensing agents

**[0125]** Poly(ethylenimine) (PEI) at 27 KiloDalton (kD). PEI was used at optimized conditions (90% charge neutralization)

**[0126]** Polylysine (PLL) at 70-150,000 molecular weight. PLL condensing materials were conjugated with nuclear signal localization peptides, either SV-40 T antigen or cys-

gly-tyr-gly-pro-lys-lys-lys-arg-lys-val-gly-gly using carboximide chemistry available from Pierce Chemical (Rockford, Ill.).

[0127] Preparations of nuclear matrix proteins (NMP). NMP were collected from a rat fibroblast cell line, and a human keratinocyte cell line using a procedure described in Gerner et al. *J. Cell. Biochem.* 71 (1998):363-374 which is incorporated herein by reference. Protein preparations were conjugated with nuclear signal localization peptides as described.

#### [0128] B. Surfactants

[0129] 2, 4, 7, 9 - tetramethyl-5-decyn-4, 7 - diol (TM-diol): HLB=4-5, CMC is not determined

[0130] Poly(oxy-1, 2-ethanediol),  $\alpha$ -(4-nonylphenol)- $\omega$ -hydroxy, Tergitol NP-40 (NP40): HLB=17.8, CMC 180  $\mu$ M,

[0131] Polyoxyethylene 20 sorbitan monooleate (Tween 80): HLB=10, CMC 920  $\mu$ M, Cetyl Alcohol: HLB=4, CMC is not determined.

#### [0132] C. Polymers

[0133] Hyaluronan, bacterially-derived, 1 million kiloDalton (MM kD) and conjugated with nuclear localization signal peptides as described in U.S. Pat. No. 5,846,561, which is incorporated herein by reference.

[0134] Hyaluronan, derived from human umbilical cord, about 4MMKD and not conjugated.

[0135] Povidone (polyvinylpyrrolidone, PVP) 10,000 kD MW and not bioconjugated Povidone (polyvinylpyrrolidone, PVP) 40,000 Kd MW and not bioconjugated Povidone (polyvinylpyrrolidone, PVP) 360,000 kD MW and not bioconjugated Tenascin, 220 kD and not bioconjugated.

#### [0136] D. Expression Vectors

[0137] 334: pcDNA/His/LacZ, produces galactosidase, incorporates CMV promoter, based on pcDNA 3.1. (Invitrogen), 8.6 kB

[0138] 425: pEGFP-c/farn, enhanced GFP (green fluorescent protein) expression vector modified with a farnasyl moiety to improve microscopy, CMV promoter, 4.6 kB

[0139] 423: pEGFP-c3/p57(Kpn/Sma) Clontech enhanced GFP (green fluorescent protein) expression vector modified with a nuclear localization tag from a cyclin dependent kinase to improve microscopy, 4.6 kB

#### [0140] E. Cells

[0141] CCRL 1764: Immortalized rat neonatal fibroblast cell line

[0142] HaCaT: immortalized human keratinocyte cell line

[0143] Ca9: human tumor cells derived from a squamous cell carcinoma of tongue origin.

#### Example 1A—Effect of changing dispersion conditions on hydrophillic nanocapsules.

[0144] The importance of appropriate dispersion conditions was investigated in the following series of formulations. Formulae were produced by i) predispersing 25  $\mu$ g of DNA (425) on ice using a bath sonicator, ii) condensing DNA in a small amount of water by vortexing then incubating on ice for 20 minutes, iii) adding surfactant then oil

followed by 30 seconds of probe sonication at 10 Watts, iv) diffusion dilution to 3 milliliters (mL) by first adding saline then 1 MM kD hyaluronan polymer (1%) as a protective colloid, v) mechanically shearing emulsion into droplets by pumping through a 250 micrometer ( $\mu$ m) orifice into 22 mL of PBS, 10 millimolar (mM)  $\text{Ca}^{2+}$ , 200 mM  $\text{Li}^{+}$ , vi) incubating overnight end over end and vii) centrifuging to recover nanoparticles for resuspension and filter sterilization. The condenser-to-DNA weight ratio was determined by dye exclusion at 90% charge neutralization. TM-diols were used in this experiment to represent water-immiscible surfactants, while Tergitol NP40 and Tween 80 were used to represent water-soluble and even more water-soluble emulsifiers/dispersing aids.

[0145] Dispersion conditions were systematically varied to discourage micelle formation in aqueous media by i) choosing water-soluble surfactants (Formulae S,T,U, and V), ii) removing the dispersed phase (Formula T) and iii) decreasing surfactant loading below that required for micelle formation. Formula U featured use of a water-miscible oil (silicone oil).

[0146] Formulae were characterized physically and tested for functionality in in vitro gene transfer. Quantitative results are summarized in Table 1A:

TABLE 1A

	Effect of changing dispersion conditions on hydrophillic nanocapsules.		
	Formula		
	Q	R	S
Experimental	surf > CMC	surf > CMC	surf > CMC
Modification:			
Critical Micelle Concentration (CMC)	~0	~0	360 ppm
Pre-aerosol surfactant Concentration (3 ml basis)	500 ppm	500 ppm	600 ppm
HLB number	4-5	4	17.8
Phases	Water/ misc. oil	Water/ misc. oil	Water/ misc. oil
Formula Characteristics:			
Nucleic Acid Incorporation (%)	86 $\pm$ 8	67 $\pm$ 1.4	50.3 $\pm$ 12
Low MW DNA Appearance (% above bkground, Post nanocapsule digest by electrophoresis)	15.00	76	93.00
Supercoil retention (post 100 hrs release) (area %, initial distribution = 76% supercoiled)	87%	65%	66%
Particle Size (mean $\pm$ SE)	42 $\pm$ 2	45 $\pm$ 3	73 $\pm$ 4
Secondary Structure(s)	25%	30%	70%
Flocculation Status	100-200 nm stringy flocs	500 nm stringy flocs	300 nm spheroid aggregates
Comments:			
Performance:			
Transduced GFP Protein Generation	420	340	0

TABLE 1A-continued

Effect of changing dispersion conditions on hydrophillic nanocapsules.	Formula		
	Q	R	S
(pixel units, % of control liposome formula, 100 $\mu$ g total protein, Day 11)			

[0147]

[0148] Nanocapsule sizing was determined by tapping mode AFM and images are illustrated in FIG. 1A. The data indicate average nanocapsule sizes less than 50 nm are achievable only with multi-phase systems in combination with low water solubility surfactants (Table 1A: Formulae Q,R vs. S,T,U,V, and W). Furthermore, only nanocapsules of less than 50 nm resulted in detectable transgene production in CRL-1764 rat fibroblast cells (Table 1A). Effective dispersion also corresponded with decreased aggregation and enhanced DNA stability (as indicated by decreased electrophoretic breakdown products). The starting DNA was partially relaxed (76% supercoiled by electrophoresis). Using this value as a basis, supercoil retention in DNA still

TABLE 1A

Effect of changing dispersion conditions on hydrophillic nanocapsules.	Formula			
	T	U	W	V
Experimental Modification:	surf > CMC	surf > CMC	surf > CMC	surf < CMC
Critical Micelle Concentration (CMC)	360 ppm	360 ppm	1200 ppm	360 ppm
Pre-aerosol surfactant Concentration (3 ml basis)	600 ppm	600 ppm	4000 ppm	90 ppm
HLB number	17.8	17.8	10	17.8
Phases	Water only	Water/immisc. oil	Water/misc. oil	Water/misc. oil
Formula Characteristics:				
Nucleic Acid Incorporation (%)	39 $\pm$ 1.7	32.8 $\pm$ 6	37 $\pm$ 1.41	57.6 $\pm$ 16
Low MW DNA Appearance (% above bkgground, Post nanocapsule digest by electrophoresis)	53.00	66	28	41.00
Supercoil retention (post 100 hrs release) (area %, initial distribution = 76% supercoiled)	59%	43%	65%	80%
Particle Size (mean $\pm$ SE)	226 $\pm$ 11 S < 10%	291 $\pm$ 25 S < 10%	150 $\pm$ 7 S > 40%	199 $\pm$ 11 S > 80% 400 nm
Secondary Structure(s) Flocculation Status			yeast-like aggregates	aggregates
Comments:	ppt. during aerosolization		ppt. during aerosolization	ppt. during aerosolization
Performance: Transduced GFP Protein Generation (pixel units, % of control liposome formula, 100 $\mu$ g total protein, Day 11)	0	0	0	0

encapsulated following 100 hrs of release testing, was excellent in multi-phase systems.

**[0149]** Release profiles for hydrophilic dispersed atomized nanocapsules were linear, showed no zero burst and resulted in about 60% release after 72 hours (See **FIG. 1B**). Formula W, manufactured with the most water-soluble surfactant in the series (Tween 80) failed to completely release loaded DNA. **FIG. 1C** illustrates that small amounts of DNA (in this case 300 nanograms of DNA) can be recovered accurately in a procedure comprising butanol extraction of 10% butanol/saline releasing fluid followed by isolation on a miniprep column and measurement of absorbance at 260 nm excitation. Results obtained from UV spectroscopy are confirmed by electrophoresis of recovered DNA following alcohol coprecipitation with a commercial coprecipitant aid. Experiment 1A demonstrates the importance of a multi-phase system in creating coated particles from the micellar solution, defines surfactant requirements and validates method for measuring in vitro release profiles.

#### Example 1B—Effect of process parameters on particle functionality

**[0150]** To investigate the effect of modulating process parameters on nanocapsule functionality for DNA delivery, a series of formulas (designed to release in 3 days) were prepared and measured transduction efficiency of these formulas for delivering a nuclear Green Fluorescent Protein (GFP) reporter transgene in rat fibroblast cultures 5 days later. Charge neutralization of the DNA molecule, the surfactant/oil system, total surfactant phase volume, the inclusion of probe sonication, the absolute requirement for atomization and receiving bath osmolality were modulated. Results for this experiment are summarized in the Table 1B:

**[0151]** Aqueous dispersion of DNA condensates with poorly soluble surfactants in the inventive method produced average nanocapsule diameters under 50 nm. A 30 number of successful operating regimes were feasible with varying effects on encapsulation yield. In a cetyl alcohol/castor oil system, under condensation resulted in an average particle diameter increase from 20 to 12 nm (Table 1B: F1 vs. F2). The same decrease in condenser weight ratio induced a particle size increase from 24 to 36 nm, while still maintaining nanocapsule functionality for transgene delivery, when using a TM-diol/DMSO surfactant system for initial micelle formation (Table 1B: F4 vs. F5). This finding teaches surfactant selection impacts final average nanocapsule diameters.

**[0152]** Moderate energy input was removed (dropped probe sonication, atomization but kept bath sonication) during nanocapsule formation and resulted in functional particles with decreased yield (Table 1B: F3 vs. F4). This finding indicates that optimal nanocapsule production is not dependent on any spontaneous micro-emulsification process. Cosolvent phase volume was reduced from 4 weight percent to 500 ppm without any negative effect on particle functionality (Table 1B: F4 vs. F6). This finding indicates that essentially solvent-free nanocapsules can be made by the inventive method. Finally, salt was removed from the atomization receiving bath without any negative effects on nanocapsule functionality (Table 1B: F6 vs. F7).

#### Example 2—Effect of nanocapsule sizing on a nanocapsule uptake in human keratinocytes

**[0153]** The effect of nanocapsule sizing on intracellular trafficking in immortalized HacaT human keratinocyte cultures (HacaT's) was investigated in this example. In this

TABLE 1B

Effective of process parameters on particle functionality								
Nano capsule Design	Formula Name	charge neutralization by condenser	Surfactant	Biocompatible Oil	Oil Phase Volume (%; 4.5 ml basis)	Emulsify by sonication	Atomize Diameter (μm)	Receiving bath Osmolality (mOs)
1	q.co.2	+	Cetyl OH	Castor oil/Etoh	4	+	250	220
2	q.co	—	Cetyl OH	Castor oil/Etoh	4	+	250	220
3	o.35	+	TM-diol	DMSO	4	+	1.4	220
4	ea0.2	+	TM-diol	DMSO	4	—	—	220
5	ea0.1	—	TM-diol	DMSO	4	—	—	220
6	ed0.2	+	TM-diol	DMSO	0.05	—	250	220
7	ed0a.12.di	+	TM-diol	DMSO	0.05	—	250	0

Nanocapsule Design	Formula name	Nanocapsule diameter (nm)* n = 20	Encapsulation yield (%; mean ± SE)	Transduction Efficiency, (5 days, rat fibroblasts)
1	q.co.2	20 ± 3, rods	48.6 ± 11	87 ± 7%
2	q.co	12 ± 0.7, irregular	48.6 ± 2	71 ± 28%
3	o.35	17 ± 1.2, spheres	82.3 ± 7 (4)	86 ± 2%
4	ea0.2	24 ± 2, s/r	32 ± 10	72 ± 2%
5	ea0.1	36 ± 3, irregular	57 ± 2	85 ± 1%
6	ed0.2	39 ± 3, r/e	39 ± 5	96%
7	ed0a12.di	39 ± 3, ellipse	69 ± 2	100%

\*Nanocapsule diameter is reported as average of the minor and major particle axis using digital image analysis, while nanocapsule morphology is reported as irregular, rods (r), ellipse (e) or spheres (s).

series of formulae, these micellar dispersion were sheared by syringes of different orifice diameter. The coating weight was also lowered from 1:1 DNA:Polymer (w/w) to 1:40 to shorten the dissolution profile from 5 to 3 days. In these experiments, nanocapsule formulae were compared to standard polyplexes of DNA and PEI, and lipoplexed plasmid DNA. Table 2 summarizes the experimental design and results:

TABLE 2

Effect of particle size on nanocapsule functionality for gene transfer					
Formula Name	Particle Size (mean, nm; morphology)	4 hr. colocalization with caveolin-1*	4 hr. colocalization with cathrin	10 hr. colocalization with lysosomes	Transduction Efficiency. (5 days, human keratinocytes)
o.22 (64)	47 ± 3, rods	0	++	+	16 ± 13
o.27 (57)	21 ± 2, rods	+	+	ND	81 ± 8
o.35 (85)	17 ± 1.2, spheres	+++	0	0	78 ± 9
pei-	67 ± 4,	0	+++	+++	40 ± 15
pDNA	spheres, irreg				
Lipoplex	48 ± 2	+	+	+++	41 ± 27
pDNA	200 nm aggregates				

Key: 0 = no change from unstimulated condition,

+ greater than 25% increase,

++ greater than 50% increase,

+++ greater than 75% increase in number of cells stimulated.

ND = not determined.

[0154] It was observed that compared to the unstimulated state, nanocapsules increased cellular pinocytotic activity compared to standard formulations, and smaller nanocapsules shifted pinocytotic activity to caveolae from clathrin-coated pits (Table 2: Formula O vs. pei-dna and lipoplex pDNA). It was further observed that nanocapsules avoided lysosome co-localization at 10 hours post-addition with smaller nanocapsules being particularly effective (see Table 2: Formula vs. pei-dna and lipoplex pDNA). These results are illustrated further in FIG. 2. This improvement is further emphasized by comparison with published uptake studies for HacaT keratinocytes. Compared to primary keratinocytes, uptake of naked DNA oligonucleotides (20  $\mu$ m) were very poor in HacaT's and showed accumulation of oligonucleotides in punctate vesicles consistent lysosomes at 2 hours. Using hydrophilic dispersed atomized nanocapsules of the inventive method, complete avoidance of lysosomes at 10 hours post-addition was demonstrated. These results indicate that products of the inventive process will have increased and prolonged effectiveness.

#### Example 3—Effect of nanoparticle delivery on DNA and reagent-induced cytotoxicity

[0155] To test whether soluble exogenous DNA released from liposomes or dendrimers induces apoptosis, Rt-1's were treated with loaded/unloaded liposome complexes, dendrimer complexes, nanoparticles and 1  $\mu$ g/ml etoposide, a DNA intercalating agent as a positive control. Cultures

were treated with standard formulas for 3 hours then assayed for gene product expression 30 hours later. Cultures were treated with nanocapsules for 4 days to ensure full DNA release during the experiment. Controls included as a positive control for apoptotic cell death, 1  $\mu$ g/ml etoposide, a DNA intercalating agent was applied to cultures overnight before experiment termination. Other controls included standard PEI-DNA complexes, empty nanocapsules and nanocapsules containing empty vector plasmid DNA. Hydrophilic nanoparticles were produced for this experiment as described earlier using a 35-gage syringe.

[0156] One of the later steps in apoptosis is DNA fragmentation mediated by activation of endonucleases during the apoptic program. Therefore, DNA fragmentation was assayed by end-labeling of fragments using an exogenous enzyme and a substituted nucleotide (TUNEL: tdt-mediated uridine nucleotide and labeling. Results are expressed as a Fragmentation Index, or the percent of cells in the total culture exhibiting BRDU end-labeled DNA. Cultures were run in duplicate. The experimental design and results are summarized in Table 3:

TABLE 3

Effect of nanocapsule coating weight on nonspecific reagent and plasmid DNA-associated cytotoxicity.				
	Formula			
	$\kappa$ .35	$\zeta$	o(Omicron)	b.35
<u>Particle Design:</u>				
DNA Condensing Agent	Denatured h. keratinocyte nuclear protein	100 Kd MW Polylysine	27 kD PEI	27 kD PEI
Coating Ratio (DNA/polymer)	0.1	0.25	0.25	0.01
<u>Performance:</u>				
dose: (30 hrs for Std. Formulas, 100 hrs for nanocapsules)	4.6	4.1	4	5
Cytotoxicity: (Fragmentation Index, %)	ND	0.26 $\pm$ 0.15	2 $\pm$ 0.7	1.9 $\pm$ 0.6
cytotoxicity controls: (1 $\mu$ g etoposide (8 hr): 25 $\pm$ 10%) (Pei-DNA polyplexes (100 hr): 24 $\pm$ 7%) (Empty vector nanocapsules: 1.25 $\pm$ 1.25%) (Empty vector nanocapsules: 0.9 $\pm$ 0.7%)				
Transduction Efficiency: (% cells 120 hrs, dose as listed)	31 $\pm$ 2	ND	85 $\pm$ 7	32 $\pm$ 3
<u>Formula Characteristics:</u>				
Nucleic: Acid Incorporation: (%)	55 $\pm$ 10	27 $\pm$ 7	54 $\pm$ 5	65 $\pm$ 4
Cumulative Release: (% 48 hr)	70	75 $\pm$ 8	83 $\pm$ 12	ND
Particle Size (mean $\pm$ SE, nm)	26 $\pm$ 2	22 $\pm$ 2	20 $\pm$ 1	35 $\pm$ 2
Agglomerates (as dispensed)	few	50% 80 $\pm$ 6	200 nm	200 nm
	Formula			
	Y.35	Lipoplex GP	Lipoplex L+	Polyplex
<u>Particle Design:</u>				
DNA Condensing Agent	27 kD PEI	cationic lipid	cationic lipid	dendrimer
Coating Ratio (DNA/polymer)	0.0025			
<u>Performance:</u>				
dose: (30 hrs for Std. Formulas, 100 hrs for nanocapsules)	5	1 $\mu$ g 500 ng 0 ng	500 ng 250 ng 0 ng	2 $\mu$ g 1 $\mu$ g 0 $\mu$ g
Cytotoxicity: (Fragmentation Index, %)	9 $\pm$ 8	27 $\pm$ 8	9.3 $\pm$ 0.2	6.63 $\pm$ 1.4
cytotoxicity controls: (1 $\mu$ g etoposide (8 hr): 25 $\pm$ 10%) (Pei-DNA polyplexes (100 hr): 24 $\pm$ 7%) (Empty vector nanocapsules:		6 $\pm$ 3 4 $\pm$ 2.5	12.8 $\pm$ 1.5 7.8 $\pm$ 0.1	5.7 $\pm$ 1.8 3.1 $\pm$ 0.3

TABLE 3-continued

Effect of nanocapsule coating weight on nonspecific reagent and plasmid DNA-associated cytotoxicity.					
1.25 ± 1.25%) (Empty vector nanocapsules: 0.9 ± 0.7%) Transduction Efficiency: (%) cells) 120 hrs, dose as listed) Formula Characteristics:	24	± 4	17 ± 2	dead	dead
Nucleic Acid Incorporation: (%)	667	±0.2	ND	ND	ND
Cumulative Release: (% 48 hr)	ND		ND	ND	ND
Particle Size (mean ± SE, nm)	57	± 5	48 ± 2	ND	22.4 ± 2
Agglomerates (as dispensed)	g.t.50% 300 nm		300 nm		25% 300 nm hard-fused

[0157]

TABLE 3B

Dose response of nanoencapsulated pDNA		
Formula	Dose (100 hr.)	GFP/Actin Production (density ratio, %)
K.35	9 µg	94.8
K.35	4.5 µg	83.5
K.35	1.5 µg	83.3
Lipoplex GP	0.5 µg	94.9

[0158] It was observed that use of controlled-release nanocapsules reduced the fraction of apoptotic cells in fibroblast cultures 3 to 100 fold. Conventional reagents without DNA showed a 4-fold increase in FI over empty nanoparticles, but increased another 50-100% without additional reagents in the presence of additional DNA. Decreasing the coating weight from 1:40 to 1:400 resulted in an increase in average nanocapsule diameter from 20 to 57 nm and the appearance of regions of apoptotic induction in cultures (Table 3: F omicron vs. F upsilon 35). Decreasing the coating weight from 1:40 to an intermediate 1:100 reduced transduction efficiency without increasing particle size and the appearance of cytotoxicity. These findings indicate that nanocapsule design plays a role in maintaining nanocapsule integrity and that size effects and dissolution profiles can contribute to observed cytotoxicity and functionality. We concluded that application of nanocapsule formulations increased dosing to useful efficiency levels without induction of an apoptotic program.

[0159] Table 3B exemplifies this improvement with a dose response of Formula K.35 measured in fibroblast lysates. GFP production was measured in fibroblast lysates after 4 days of treatment with increasing doses of nanocapsules. A 9.5 µg dose of nanocapsules equaled the production of a liposomal formulation without any evidence of cytotoxicity.

Example 4—Nanocapsule delivery of macromolecules to porcine tissue across keratinized barrier epithelia by transdermal and subcutaneous means

[0160] The utility of nanocapsules for nonviral nucleic acid delivery to tissue in a pig biopsy organ culture system was investigated. 6 and 8 mm circular biopsies were collected under sterile conditions from sedated research animals and cultured on meshes in partial contact with media containing 20% Fetal Calf Serum. Biopsies were either injected with 90 µl (6.3 µg) or treated topically with 3×30 µl aliquots. Biopsies were snapfrozen 7 days later and sectioned/homogenized for β-galactosidase production measurement. Formulation information and results from this experiment are summarized in Table 4:

TABLE 4:

Table 4: Functionality of dispersed atomized nanocapsules for macromolecule delivery across keratinized barrier membranes.		
Exp. Modification (from Formula Q)	Formula	
	N	O
	coating wt. is 2.5x Polymer MW is 1x	coating wt. is 2.5x Polymer MW is 4x
Formula Characteristics:		
Nucleic Acid Incorporation (%)	70.00	70.50
Cumulative Release (%, 169 hr 2.5 µg sample)	83	83. ± 1.5
Low MW DNA in postdigested Electrophoresis Samples	0	0
Supercoil retention (237 hr release, initial = 69.7% sc/released)	100%	100%
Particle Size (mean, SE, major species)	18.2 ± 0.2 nm	ND
Particle Description	spherical	
Secondary structure:	20% 100 nm flocs	



TABLE 4:-continued

Table 4: Functionality of dispersed atomized nanocapsules for macromolecule delivery across keratinized barrier membranes.

## Performance:

Transduced Protein Production (pixel units, % of neg control, 100 $\mu$ g total protein, normalized by protein)	312 $\pm$ 74 (topical) 142 (s.c.)	
Reporter (gene Product Distribution (6.3 $\mu$ g dose, 6 mm (N), 8 mm (O) porcine biopsy, 1 wk) keratinocytes (% cells), n = 2 fields/200 cells, neg ctrl: 6% endothelial cells, (% vwf+area) papillary and/or reticular, n = 2-4 fields, neg ctrl: 1.07 $\pm$ 0.72 dermis (% area); negative ctrl: 0.24 $\pm$ 0.03, n = 4/20x fields)	100%  73 $\pm$ 20 (pap)  32 $\pm$ 15 (ret)  2.74 $\pm$ 0.96*	100%  13.8 $\pm$ 0.5 (pap)  8 $\pm$ 2 (ret)  1.77 $\pm$ 0.49*

\*= p < 0.05

[0161] Western blotting of radiated tissue lysates showed a 3-fold increase in  $\beta$ -galactosidase in duplicate biopsies treated topically with Formula N over an identically cultured 6mm biopsy treated with saline. Only a 2-fold increase was measured in a 8 mm biopsy treated topically with formula O nanoparticles (see FIG. 4B). Formula O was produced with a higher molecular weight analog of the N polymer suggesting a difference in particle morphology, a dose effect or differing in situ release profiles between the two formulations related to this difference. To identify initial cell type-specific differences in nanocapsule delivery effectiveness, tissue sections were analyzed for  $\beta$ -galactosidase expression in double-label experiments using antibodies to cell-specific epitopes (see FIG. 4B). Digital image analysis of these sections indicated that radiated keratinocytes and endothelial cells are readily transduced in organ culture 7 days after treatment with a 10 day releasing formula. Specific quantitation of fibroblastic cells was not possible without inclusion of a cell-specific marker, however, an 11-fold increase in area of expression was measured in N biopsy dermis (see FIG. 4B). Interestingly, for both the formulae N and O topically-treated biopsies examined, the area percentage of blood vessels transduced decreased about 50% in nearby fields between 100  $\mu$ m and 300  $\mu$ m of depth (Table 4: papillary vs. reticular endothelial cells). These data suggest that nanocapsules are penetrating the epidermis to enter the dermis.

Example 5—Incorporation of inventive nanocapsules into a solid dosage form for additional utility in physical targeting

[0162] Nanocapsules containing a nuclear GFP transgene or empty vector were incorporated into a suture coating by vortexing the following components: i) 50  $\mu$ g of nanocapsules containing plasmid DNA, ii) 200  $\mu$ g of bovine mucin, and iii) 75  $\mu$ g of sucrose (60% w/w) in a 1000  $\mu$ l volume. Sutures were aseptically coated by drawing sutures 5x through punctured microcentrifuge tubes. Coating functionality for gene transfer was tested by applying sutures in

cultured porcine skin biopsies. Biopsies were cultured on a mesh such that the biopsy bottom was in contact with cell culture media. Biopsies were treated for 7 days, then snap-frozen and sectioned for immunofluorescence microscopy to assess nanocapsule penetration and transgene delivery.

[0163] Nanocapsule penetration was detected by streptavidin-biotin immunocomplexes directed at sheep IgG. Nanocapsule coatings are spiked with ovine IgG to enable this detection strategy. FIG. 5A shows distribution of sheep IgG signal throughout porcine dermal tissue with accumulation on capillaries. In FIG. 5A', primary antibody is omitted during slide processing to determine level of background fluorescence. A suture is visible in this view. Sutures were identifiable as smooth objects without positive nuclear counterstain. GFP expression was confirmed using a polyclonal GFP antibody to obviate the effect of nonspecific tissue green fluorescence. FIG. 5B shows GFP expression throughout the suture-treated dermis using a GFP polyclonal antibody. A suture was visible 750 microns away. FIG. 5C shows the lack of GFP expression in a biopsy treated with empty vector coating. This example demonstrates the usefulness of nanocapsules for use in physically targeted macromolecule delivery.

Example 6—Utility of nanocapsules for local targeting by design of nanocapsule coating

[0164] Fibroblast targeting

[0165] GFP nanocapsules were prepared by dispersion atomization as described in Example 1. Polyvinylpyrrolidone (PVP, MW 10,000) was used as the coating basis. A coating weight ratio of 1:40 was used and rod-shaped nanocapsules of 23 $\pm$ 2 nm were produced. 1  $\mu$ g of PVP nanocapsules were applied to both human dermal fibroblasts (HDF) and HacaT keratinocyte cultures for 4 hours then fixed for detection for nanocapsule uptake by streptavidin-biotin immunocomplexes to sheep IgG. Nanocapsule coatings are spiked with ovine IgG to enable this detection strategy. FIG. 6 illustrates positive nuclear localization of PVP nanocapsules in HDF's and negative colocalization of PVP nanocapsules in keratinocytes (FIG. 6: 6a vs. 6b). Views of untreated cultures are included for comparison (6a', 6b'). Cultures were also treated with 5  $\mu$ g of PVP nanocapsules for 5 days then tested for GFP transgene production. Consistent with uptake studies results, only the fibroblast cultures showed production of GFP transgene (FIG. 6: 6a" vs. 6b").

[0166] Tumor-targeting

[0167] GFP nanocapsules were prepared by dispersion atomization as described in example 1. Tenascin (TN, MW 200,000) was used as the coating basis. A coating weight ratio of 1:20 was used and spherical nanocapsules of 19 $\pm$ 0.9 nm were produced. 500 ng of TN nanocapsules were applied topically in successive small aliquots to pig biopsies maintained in organ culture. Biopsies were rinsed in media after 3 minutes of topical application and culture media was changed to preclude any delivery other than topical.

[0168] To simulate tumor nests of epithelial-derived origin, biopsies had been seeded 12 hours previously with 50,000 human squamous carcinoma cells. 7 days later biopsies were snapfrozen and sectioned for immunological detection of GFP production. In FIG. 6B, view "a" shows

intense GFP fluorescence in the tumor center, view “b” confirms this GFP expression with polyclonal antibodies to GFP, view “c” shows cell positioning in the section using a counterstain for cell nuclei and view “d” shows the level of background fluorescence by omission of GFP antibodies. Tumor origin was confirmed by positive detection with antibody to keratin 10/1, an epithelial marker. Comparison of view “b” and view “c” indicates that GFP expression is limited to cells within the tumor. As already demonstrated in example 5, expression throughout a tissue is also feasible and can be modulated by coating design. This example demonstrates that nanocapsule delivery can be productively targeted.

**[0169]** Cell-specific delivery for enhanced drug therapeutic window

**[0170]** Nanocapsules were prepared as described in Example 1 to encapsulate cisplatin, a hydrophobic molecule and a common cancer chemotherapeutic with serious side effects. A coating weight ratio of 1:100 was used and

Example 7—Utility of nanoencapsulation for improved cellular uptake of other species used as pharmaceutical, nutraceutical, research or cosmetic agents

**[0171]** Nanocapsules containing either 500 kD FITC-labelled dextran, 20 mer FITC-labelled mer O-methylated RNA oligonucleotide and 16 mer phosphodiester DNA oligonucleotide were prepared as described in Example 1. A 1:40 coating weight ratio was used and 1 MM kD hyaluronan was used as a coating basis. PEI was used to condense the phosphodiester DNA oligonucleotide, but no PEI was included in the dextran or RNA oligonucleotide formulas. Nanocapsule functionality for drug delivery was tested by evaluating changes in relative pinocytotic activity and cellular uptake in 35 mm cultures of human dermal fibroblast. Nanocapsule formulas were compared to naked species or species formulated as complexes. Quantitative results are summarized in Table 7.

TABLE 6

Nanocapsulation improves cellular uptake of other species used as pharmaceutical, nutraceutical, research or cosmetic agents. At 18 hours post-addition, lysosomes are only evident in conventionally formulated species.								
		Particle size (mean, SE, nm, morphology)	4.5 hours post-addition		18 hours post-addition			
Bioactive			Increase in cellular uptake activity, (% cells above baseline, <u>caveolin-1/clathrin</u> )	Nuclear Uptake Efficiency (% cells,	Bioactive component Colocalization with lysosomes, (% cells, human fibroblasts)	Detection persistence (% cell, human fibroblast)		
Component	Formulation		dose	fibroblast)		fibroblasts)	fibroblast)	
500 kd	nanocapsule	22 ± 2, s/r	89/20	25 µg*	95 ± 2	2 ± 2	5 µg	88 ± 11
fluc-dextran	naked, FITclabelled	—	75/18	100 µg	10	100 ± 10	100 µg	61 ± 20
20 mer o-	nanocapsule	13 ± 0.7, r	78/90	2 µg	74 ± 5	0 ± 0	5 µg	80 ± 6
methylated	naked, FITclabelled	—	—/73	5 µg	14 ± 7	—	—	—
RNA oligo	PEI/FITclabelled	236 ± 26, r	—/—	—	—	100 ± 0	5 µg	94 ± 10
16 mer PO	nanocapsule	17 ± 1, r	70/94	1 µg	34 ± 25	0 ± 0	5 µg	91 ± 8
DNA oligo	PEI/FITclabelled	67 ± 4, s/r	72%	2 µg	95 ± 2	80 ± 7	5 µg	66
			lysosomes					
Nominal n		20 particles	70 cells		140 cells	50 cells		50 cells

\*Dose was estimated for encapsulation dextran assuming 100% encapsulation.

s = sphere

r = rod

irregular nanocapsules of 29±3 nm were produced. Targeting efficacy was demonstrated by changes in the dose response for cell growth inhibition in fibroblast vs. squamous cell carcinoma cultures. Cells were seeded overnight into 96 well plates, treated for 18 hours with increasing amounts of encapsulated or unencapsulated drug, then assessed for cell growth inhibition using an MTT assay 48 hours later for total growth time of 72 hours. Results are illustrated in FIG. 6C. The data shows that tenascin nanocapsules protected nontarget cells from cell death (zero death) at drug levels that killed using unencapsulated drug (FIG. 6Aa: open vs. closed circles). In carcinoma cultures, TN nanocapsules productively decreased the inhibition concentration (IC<sub>50</sub>) an estimated 300% from 525 to 160 µg/ml. Example 6 demonstrates the usefulness of nanocapsules for use in coating-targeted macromolecule delivery.

**[0172]** Table 7 shows that average diameters for all nanocapsules were below 50 nm by AFM. PEI complexes of DNA oligonucleotides were measured at 67 nm and PEI complexes of uncharged RNA O-methyl oligonucleotides were measured at 236 nm. As discussed in Example 2 using keratinocyte cultures and plasmid DNA, nanocapsules stimulate pinocytotic activity as indicated by increased signal levels of clathrin and caveolin-1. In the 500 kD dextran case, pinocytotic activity shifts productively towards caveolae with nanoencapsulation (Table 7, 500 kD Dextran). At 4.5 hours post-addition, nuclear uptake is enhanced for encapsulated dextran and RNA relative to naked species. For the nanocapsules of DNA oligonucleotides, cellular uptake is decreased relative to complexed oligonucleotide, however, a majority of that DNA oligonucleotides is already nonproductively sequestered in lyso-

somes by that 4.5 hours (Table 7). At 18 hours post-addition, nanocapsules species show continued exclusion from lysosomes, while naked species show high levels of sequestration. These results are illustrated in **FIG. 7A and 7B**. Views “a” and “b” show Fite detection in cultures at 18 hours. That distribution is exclusively nuclear for the nanocapsules of RNA oligonucleotides (**FIG. 7B: a vs. a'**). Punctate inclusion are visible in the cultures treated with the complexed RNA oligonucleotides that co-localize with an immunological marker for lysosomes (**FIG. 7A: a vs. a'**). Particle sizing results from AFM microscopy are included to demonstrate dramatic difference in sizing following encapsulation. (**FIG. 7A, 7B: b vs. b, b'**). Formulas encapsulating lower molecular

weight dextrans and unstabilized RNA were also prepared with analogous uptake, nanocapsule size and yield to demonstrate that encapsulation can provide not only a targeting function but aid in stabilizing molecules sensitive to chemical or enzymatic degradation. These examples demonstrate the usefulness of nanocapsules **36** for use in delivery of a broad range of macromolecules.

[0173] Although the present invention has been described with reference to preferred embodiments, workers skilled in the art will recognize that changes may be made in form and detail without departing from the spirit and scope of the invention.

---

SEQUENCE LISTING

---

<160> NUMBER OF SEQ ID NOS: 5

<210> SEQ ID NO 1

<211> LENGTH: 6232

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Supplied by Invitrogen of Carlsbad, California

<400> SEQUENCE: 1

```

gtaccgaatt caagcttcgt gaggtccgg tgcccgtcag tgggcagagc gcacatcgcc      60
cacagtcccc gagaagttag ggggaggggt cggaattga accggtgcct agagaagggtg    120
gcgcggggta aactgggaaa gtgatgtcgt gtactggctc cgcctttttc ccgaggggtgg    180
gggagaaccg tatataagtg cagtagtcgc cgtgaacgtt ctttttcgca acgggtttgc     240
cgccagaaca caggtaatgt ccgtgtgttg ttcccgcggg cctggcctct ttacgggtta     300
tggcccttgc gtgccttgaa ttacttccac ctggctccag tacgtgatto ttgatccga     360
gctggagcca gggcggggcc ttgcgcttta ggagccctt cgcctcgtgc ttgagttgag     420
gcctggcctg ggcgtgggg ccgccgcgtg cgaatctggt ggcacctcg cgcctgtctc     480
gctgctttcg ataagtctct agccatttaa aatttttgat gacctgctgc gacgcttttt     540
ttctggcaag atagtcttgt aaatgcgggc caggatctgc aactgggtat ttcggttttt     600
gggcccgcgg ccggcgacgg gggccgtgcg tcccagcgca catgttcggc gaggcggggc     660
ctgcgagcgc ggccaccgag aatcggaagg gggtagtctc aagctggccg gcctgctctg     720
gtgcctggcc tcgcgccgcc gtgtatcgcc ccgccctggg cggcaaggct ggcccggtcg     780
gcaccagttg cgtgagcgga aagatggccg cttcccggcc ctgctccagg gggctcaaaa     840
tggaggacgc ggcgctcggg agagcggggc ggtgagtcac ccacacaaag gaaaagggcc     900
tttccgtcct cagccgtcgc ttcattgtgac tccacggagt accgggcgcc gtccaggcac     960
ctcgattagt tctggagctt ttggagtacg tcgtcttttag gttgggggga ggggttttat    1020
gcgatggagt tttccacac tgagtgggtg gagactgaag ttaggccagc ttggcacttg    1080
atgtaattct ccttggaatt tggccttttt gagtttggt cttggttcat tctcaagcct    1140
cagacagtgg ttcaaagttt ttttcttcca ttccaggtgt cgtgaacacg tggccaccat    1200
ggcccagtg cagctgcaga tggctagcaa aggagaagaa cttttcactg gagttgtccc    1260
aattcttggt gaattagatg gtgatgttaa tgggcacaaa ttttctgtca gtggagaggg    1320
tgaaggtgat gctacatacg gaaagcttac ccttaaattt atttgacta ctggaaact     1380

```

-continued

---

acctgttcca	tggccaacac	ttgtcactac	tttctcttat	ggtgttcaat	gcttttcccg	1440
ttatccggat	catatgaaac	ggcatgactt	tttcaagagt	gccatgcccg	aagggtatgt	1500
acaggaacgc	actatatctt	tcaaagatga	cgggaactac	aagacgcgtg	ctgaagtcaa	1560
gtttgaaggt	gatacccttg	ttaatcgtat	cgagttaaaa	ggtattgatt	ttaaagaaga	1620
tggaacatt	ctcggacaca	aactcgagta	caactataac	tcacacaatg	tatacatcac	1680
ggcagacaaa	caaaagaatg	gaatcaaagc	taacttcaa	attcgccaca	acattgaaga	1740
tggaatccgtt	caactagcag	accattatca	acaaaatact	ccaattggcg	atggccctgt	1800
ccttttacca	gacaaccatt	acctgtcgac	acaatctgcc	ctttcgaaag	atcccaacga	1860
aaagcgtgac	cacatggtcc	ttcttgagtt	tgtaactgct	gctgggatta	cacatggcat	1920
ggatgagctc	tacaaagcgg	ccgcagatcc	aaaaaagaag	agaaggtag	atccaaaaaa	1980
gaagagaaa	gtagatccaa	aaaagaagag	aaaggtagat	acggccgcag	aacaaaaact	2040
catctcagaa	gaggatctga	atggggccgc	atagtctaga	agctcgctga	tcagcctcga	2100
ctgtgccttc	tagttgccag	ccatctgttg	tttgcccctc	cccgtgcct	tccttgacct	2160
tggaaggtgc	cactcccact	gtcctttcct	aataaaatga	ggaaattgca	tcgcattgtc	2220
tgagtaggtg	tcattctatt	ctggggggtg	gggtggggca	ggacagcaag	ggggaggatt	2280
gggaagacaa	tagcaggcat	gctggggatg	gcccgggctc	tatggcttct	gaggcgaaa	2340
gaaccagctg	gggtcttagg	gggtatcccc	acgcgccctg	tagcggcgca	ttaagcgcg	2400
cgggtgtggt	ggttacgcgc	agcgtgaccg	ctacacttgc	cagcgcccta	gcgcccgctc	2460
ctttcgcttt	cttcccttcc	tttctcgcca	cgttcgccgg	ctttcccctg	caagctctaa	2520
atcggggcat	cccttttagg	ttccgattta	gtgctttacg	gcacctcgac	cccaaaaaac	2580
ttgattaggg	tgatggttca	cgtagtgggc	catcgccctg	atagacggtt	tttcgcccct	2640
tgacgttgg	gtccacgttc	tttaatatgt	gactcttggt	ccaaactgga	acaacactca	2700
accctatctc	ggtctattct	tttgatttat	aagggatttt	ggggatttcg	gcctattggt	2760
taaaaaatga	gctgatttaa	caaaaattta	acgcgaatta	attctgtgga	atgtgtgtca	2820
gttagggtgt	ggaaagtccc	caggctcccc	aggcaggcag	aagtatgcaa	agcatgcac	2880
tcaattatgc	agcaaccagg	tgtggaaa	ccccaggctc	cccagcaggc	agaagtatgc	2940
aaagcatgca	tctcaattag	tcagcaacca	tagtcccgcc	cctaactccg	cccatcccgc	3000
ccctaactcc	gcccagttcc	gcccattctc	cgcccctagg	ctgactaatt	ttttttat	3060
atgcagaggc	cgaggccgcc	tctgcctctg	agctattcca	gaagtagtga	ggaggctttt	3120
ttggaggcct	aggcttttgc	aaaaagctcc	cgggagggtcc	acaatgattg	aacaagatgg	3180
attgcacgca	ggttctccgg	ccgcttgggt	ggagaggcta	ttcggctatg	actgggcaca	3240
acagacaatc	ggctgctctg	atgccgcgtg	gttccggctg	tcagcgcagg	ggcgcccggt	3300
tctttttgtc	aagaccgacc	tgtccggtgc	cctgaatgaa	ctccaggacg	aggcagcgcg	3360
gctatcgtgg	ctggccacga	cgggcgttcc	ttgcgcagct	gtgctcgacg	ttgtcactga	3420
agcgggaagg	gactgggtgc	tattgggcga	agtgccgggg	caggatctcc	tgatcatctca	3480
ccttgctcct	gccgagaaa	tatccatcat	ggctgatgca	atgcggcggc	tgcatacgct	3540
tgatccggct	acctgccc	tcgaccacca	agcgaaacat	cgcacgcagc	gagcacgtac	3600
tcggatggaa	gccggctctg	tcgatcagga	tgatctggac	gaagagcatc	aggggctcgc	3660

-continued

---

gccagccgaa	ctgttcgcca	ggctcaaggc	gcgtatgcc	gacggcgagg	atctcgtcgt	3720
gactcatggc	gatgcctgct	tgccgaatat	catggtggaa	aatggccgct	tttctggatt	3780
catcgactgt	ggccggctgg	gtgtggcgga	ccgctatcag	gacatagcgt	tggtaccctg	3840
tgatattgct	gaagagcttg	gcggcgaaatg	ggctgaccgc	ttcctcgtgc	tttacggtat	3900
cgcgcgtccc	gattcgcagc	gcctcgcctt	ctatcgcctt	cttgacgagt	tcttctgagc	3960
gggactctgg	ggttcgaaat	gaccgaccaa	gcgacgcca	acctgccatc	acgagatttc	4020
gattccaccg	ccgccttcta	tgaaagggtg	ggcttcggaa	tcgttttccg	ggacgccggc	4080
tggtgatcc	tccagcgcgg	ggatctcatg	ctggagttct	tcgcccaccc	caacttgttt	4140
attgcagctt	ataatggtta	caaataaagc	aatagcatca	caaatttcac	aaataaagca	4200
tttttttcac	tgcatcttag	ttgtggtttg	tccaaactca	tcaatgtatc	ttatcatgtc	4260
tgtataccgg	atctttccgc	ttcctcgtc	actgactcgc	tcgcctcggg	cgttcggctg	4320
cggcgagcgg	tatcagctca	ctcaaaggcg	gtaatacggg	tatccacaga	atcaggggat	4380
aacgcaggaa	agaacatgtg	agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	4440
gcgttgctgg	cgtttttcca	taggctccgc	ccccctgacg	agcatcacia	aaatcgacgc	4500
tcaagtcaga	ggtggcgaaa	cccgcagga	ctataaagat	accaggcggt	tccccctgga	4560
agctccctcg	tcgcgtctcc	tgttccgacc	ctgcgcgtta	ccgataacct	gtccgccttt	4620
ctcccttcgg	gaagcgtggc	gctttctcaa	tgctcacgct	gtaggatatc	cagttcgggtg	4680
taggtcgttc	gtccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	4740
gccttatccg	gtaactatcg	tcttgagtcc	aaccgcgtaa	gacacgactt	atcgccactg	4800
gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtg	tacagagttc	4860
ttgaagtggg	ggcctaacta	cggctacact	agaaggacag	tatttggtat	ctgcgctctg	4920
ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	4980
gctggttagcg	gtggtttttt	tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	5040
caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	aaactcacgt	5100
taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	tttaaatata	5160
aaatgaagtt	ttaaatacat	ctaaagtata	tatgagtaaa	cttggtctga	cagttaccaa	5220
tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcac	catagttgcc	5280
tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	ccccagtgct	5340
gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	aaaccagcca	5400
gccggaaggg	ccgagcgcag	aagtggctct	gcaactttat	ccgcctccat	ccagtctatt	5460
aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	5520
gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	atcagctcc	5580
ggttcccaac	gatcaaggcg	agttacatga	tccccatgt	tgtgcaaaaa	agcgggttagc	5640
tccttcggtc	ctccgacgt	gtcagaagt	aagttggcgg	cagtgttatc	actcatgggt	5700
atggcagcac	tcataaatc	tcttactgtc	atgccatccg	taagatgctt	tctgtgact	5760
ggtgagtact	caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	5820
ccggcgtaaa	tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	gtcatcatt	5880
ggaaaacggt	cttcggggcg	aaaactctca	aggatcttac	cgctgttgag	atccagttcg	5940

## -continued

---

```

atgtaaccca ctctgtgcacc caactgatct tcagcatctt ttactttcac cagcgtttct 6000
gggtgagcaa aaacaggaag gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa 6060
tggtgaatac tcatactctt cctttttcaa tattattgaa gcatttatca gggttattgt 6120
ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaatagg ggttccgcgc 6180
acatttcccc gaaaagtgcc acctgacgtc agatcgacgg atcgggagat cg 6232

```

```

<210> SEQ ID NO 2
<211> LENGTH: 2200
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Gly Ala Met Thr Gln Leu Leu Ala Gly Val Phe Leu Ala Phe Leu
1           5           10           15
Ala Leu Ala Thr Glu Gly Gly Val Leu Lys Lys Val Ile Arg His Lys
20          25          30
Arg Gln Ser Gly Val Asn Ala Thr Leu Pro Glu Glu Asn Gln Pro Val
35          40          45
Val Phe Asn His Val Tyr Asn Ile Lys Leu Pro Val Gly Ser Gln Cys
50          55          60
Ser Val Asp Leu Glu Ser Ala Ser Gly Glu Lys Asp Leu Ala Pro Pro
65          70          75          80
Ser Glu Pro Ser Glu Ser Phe Gln Glu His Thr Val Asp Gly Glu Asn
85          90          95
Gln Ile Val Phe Thr His Arg Ile Asn Ile Pro Arg Arg Ala Cys Gly
100         105         110
Cys Ala Ala Ala Pro Asp Val Lys Glu Leu Leu Ser Arg Leu Glu Glu
115         120         125
Leu Glu Asn Leu Val Ser Ser Leu Arg Glu Gln Cys Thr Ala Gly Ala
130         135         140
Gly Cys Cys Leu Gln Pro Ala Thr Gly Arg Leu Asp Thr Arg Pro Phe
145         150         155         160
Cys Ser Gly Arg Gly Asn Phe Ser Thr Glu Gly Cys Gly Cys Val Cys
165         170         175
Glu Pro Gly Trp Lys Gly Pro Asn Cys Ser Glu Pro Glu Cys Pro Gly
180         185         190
Asn Cys His Leu Arg Gly Arg Cys Ile Asp Gly Gln Cys Ile Cys Asp
195         200         205
Asp Gly Phe Thr Gly Glu Asp Cys Ser Gln Leu Ala Cys Pro Ser Asp
210         215         220
Cys Asn Asp Gln Gly Lys Cys Val Asn Gly Val Cys Ile Cys Phe Glu
225         230         235         240
Gly Tyr Ala Gly Ala Asp Cys Ser Arg Glu Ile Cys Pro Val Pro Cys
245         250         255
Ser Glu Glu His Gly Thr Cys Val Asp Gly Leu Cys Val Cys His Asp
260         265         270
Gly Phe Ala Gly Asp Asp Cys Asn Lys Pro Leu Cys Leu Asn Asn Cys
275         280         285
Tyr Asn Arg Gly Arg Cys Val Glu Asn Glu Cys Val Cys Asp Glu Gly
290         295         300

```

## -continued

---

Phe	Thr	Gly	Glu	Asp	Cys	Ser	Glu	Leu	Ile	Cys	Pro	Asn	Asp	Cys	Phe	305	310	315	320
Asp	Arg	Gly	Arg	Cys	Ile	Asn	Gly	Thr	Cys	Tyr	Cys	Glu	Glu	Gly	Phe	325	330	335	
Thr	Gly	Glu	Asp	Cys	Gly	Lys	Pro	Thr	Cys	Pro	His	Ala	Cys	His	Thr	340	345	350	
Gln	Gly	Arg	Cys	Glu	Glu	Gly	Gln	Cys	Val	Cys	Asp	Glu	Gly	Phe	Ala	355	360	365	
Gly	Leu	Asp	Cys	Ser	Glu	Lys	Arg	Cys	Pro	Ala	Asp	Cys	His	Asn	Arg	370	375	380	
Gly	Arg	Cys	Val	Asp	Gly	Arg	Cys	Glu	Cys	Asp	Asp	Gly	Phe	Thr	Gly	385	390	395	400
Ala	Asp	Cys	Gly	Glu	Leu	Lys	Cys	Pro	Asn	Gly	Cys	Ser	Gly	His	Gly	405	410	415	
Arg	Cys	Val	Asn	Gly	Gln	Cys	Val	Cys	Asp	Glu	Gly	Tyr	Thr	Gly	Glu	420	425	430	
Asp	Cys	Ser	Gln	Leu	Arg	Cys	Pro	Asn	Asp	Cys	His	Ser	Arg	Gly	Arg	435	440	445	
Cys	Val	Glu	Gly	Lys	Cys	Val	Cys	Glu	Gln	Gly	Phe	Lys	Gly	Tyr	Asp	450	455	460	
Cys	Ser	Asp	Met	Ser	Cys	Pro	Asn	Asp	Cys	His	Gln	His	Gly	Arg	Cys	465	470	475	480
Val	Asn	Gly	Met	Cys	Val	Cys	Asp	Asp	Gly	Tyr	Thr	Gly	Glu	Asp	Cys	485	490	495	
Arg	Asp	Arg	Gln	Cys	Pro	Arg	Asp	Cys	Ser	Asn	Arg	Gly	Leu	Cys	Val	500	505	510	
Asp	Gly	Gln	Cys	Val	Cys	Glu	Asp	Gly	Phe	Thr	Gly	Pro	Asp	Cys	Ala	515	520	525	
Glu	Leu	Ser	Cys	Pro	Asn	Asp	Cys	His	Gly	Gln	Gly	Arg	Cys	Val	Asn	530	535	540	
Gly	Gln	Cys	Val	Cys	His	Glu	Gly	Phe	Met	Gly	Lys	Asp	Cys	Lys	Glu	545	550	555	560
Gln	Arg	Cys	Pro	Ser	Asp	Cys	His	Gly	Gln	Gly	Arg	Cys	Val	Asp	Gly	565	570	575	
Gln	Cys	Ile	Cys	His	Glu	Gly	Phe	Thr	Gly	Leu	Asp	Cys	Gly	Gln	His	580	585	590	
Ser	Cys	Pro	Ser	Asp	Cys	Asn	Asn	Leu	Gly	Gln	Cys	Val	Ser	Gly	Arg	595	600	605	
Cys	Ile	Cys	Asn	Glu	Gly	Tyr	Ser	Gly	Glu	Asp	Cys	Ser	Glu	Val	Ser	610	615	620	
Pro	Pro	Lys	Asp	Leu	Val	Val	Thr	Glu	Val	Thr	Glu	Glu	Thr	Val	Asn	625	630	635	640
Leu	Ala	Trp	Asp	Asn	Glu	Met	Arg	Val	Thr	Glu	Tyr	Leu	Val	Val	Tyr	645	650	655	
Thr	Pro	Thr	His	Glu	Gly	Gly	Leu	Glu	Met	Gln	Phe	Arg	Val	Pro	Gly	660	665	670	
Asp	Gln	Thr	Ser	Thr	Ile	Ile	Gln	Glu	Leu	Glu	Pro	Gly	Val	Glu	Tyr	675	680	685	
Phe	Ile	Arg	Val	Phe	Ala	Ile	Leu	Glu	Asn	Lys	Lys	Ser	Ile	Pro	Val	690	695	700	
Ser	Ala	Arg	Val	Ala	Thr	Tyr	Leu	Pro	Ala	Pro	Glu	Gly	Leu	Lys	Phe				

-continued

705	710	715	720
Lys Ser Ile Lys Glu Thr Ser Val Glu Val Glu Trp Asp Pro Leu Asp	725	730	735
Ile Ala Phe Glu Thr Trp Glu Ile Ile Phe Arg Asn Met Asn Lys Glu	740	745	750
Asp Glu Gly Glu Ile Thr Lys Ser Leu Arg Arg Pro Glu Thr Ser Tyr	755	760	765
Arg Gln Thr Gly Leu Ala Pro Gly Gln Glu Tyr Glu Ile Ser Leu His	770	775	780
Ile Val Lys Asn Asn Thr Arg Gly Pro Gly Leu Lys Arg Val Thr Thr	785	790	795
Thr Arg Leu Asp Ala Pro Ser Gln Ile Glu Val Lys Asp Val Thr Asp	805	810	815
Thr Thr Ala Leu Ile Thr Trp Phe Lys Pro Leu Ala Glu Ile Asp Gly	820	825	830
Ile Glu Leu Thr Tyr Gly Ile Lys Asp Val Pro Gly Asp Arg Thr Thr	835	840	845
Ile Asp Leu Thr Glu Asp Glu Asn Gln Tyr Ser Ile Gly Asn Leu Lys	850	855	860
Pro Asp Thr Glu Tyr Glu Val Ser Leu Ile Ser Arg Arg Gly Asp Met	865	870	875
Ser Ser Asn Pro Ala Lys Glu Thr Phe Thr Thr Gly Leu Asp Ala Pro	885	890	895
Arg Asn Leu Arg Arg Val Ser Gln Thr Asp Asn Ser Ile Thr Leu Glu	900	905	910
Trp Arg Asn Gly Lys Ala Ala Ile Asp Ser Tyr Arg Ile Lys Tyr Ala	915	920	925
Pro Ile Ser Gly Gly Asp His Ala Glu Val Asp Val Pro Lys Ser Gln	930	935	940
Gln Ala Thr Thr Lys Thr Thr Leu Thr Gly Leu Arg Pro Gly Thr Glu	945	950	955
Tyr Gly Ile Gly Val Ser Ala Val Lys Glu Asp Lys Glu Ser Asn Pro	965	970	975
Ala Thr Ile Asn Ala Ala Thr Glu Leu Asp Thr Pro Lys Asp Leu Gln	980	985	990
Val Ser Glu Thr Ala Glu Thr Ser Leu Thr Leu Leu Trp Lys Thr Pro	995	1000	1005
Leu Ala Lys Phe Asp Arg Tyr Arg Leu Asn Tyr Ser Leu Pro Thr	1010	1015	1020
Gly Gln Trp Val Gly Val Gln Leu Pro Arg Asn Thr Thr Ser Tyr	1025	1030	1035
Val Leu Arg Gly Leu Glu Pro Gly Gln Glu Tyr Asn Val Leu Leu	1040	1045	1050
Thr Ala Glu Lys Gly Arg His Lys Ser Lys Pro Ala Arg Val Lys	1055	1060	1065
Ala Ser Thr Glu Gln Ala Pro Glu Leu Glu Asn Leu Thr Val Thr	1070	1075	1080
Glu Val Gly Trp Asp Gly Leu Arg Leu Asn Trp Thr Ala Ala Asp	1085	1090	1095
Gln Ala Tyr Glu His Phe Ile Ile Gln Val Gln Glu Ala Asn Lys	1100	1105	1110



-continued

---

Val	Glu	Ala	Ala	Arg	Asn	Leu	Thr	Val	Pro	Gly	Ser	Leu	Arg	Ala
1115						1120					1125			
Val	Asp	Ile	Pro	Gly	Leu	Lys	Ala	Ala	Thr	Pro	Tyr	Thr	Val	Ser
1130						1135					1140			
Ile	Tyr	Gly	Val	Ile	Gln	Gly	Tyr	Arg	Thr	Pro	Val	Leu	Ser	Ala
1145						1150					1155			
Glu	Ala	Ser	Thr	Gly	Glu	Thr	Pro	Asn	Leu	Gly	Glu	Val	Val	Val
1160						1165					1170			
Ala	Glu	Val	Gly	Trp	Asp	Ala	Leu	Lys	Leu	Asn	Trp	Thr	Ala	Pro
1175						1180					1185			
Glu	Gly	Ala	Tyr	Glu	Tyr	Phe	Phe	Ile	Gln	Val	Gln	Glu	Ala	Asp
1190						1195					1200			
Thr	Val	Glu	Ala	Ala	Gln	Asn	Leu	Thr	Val	Pro	Gly	Gly	Leu	Arg
1205						1210					1215			
Ser	Thr	Asp	Leu	Pro	Gly	Leu	Lys	Ala	Ala	Thr	His	Tyr	Thr	Ile
1220						1225					1230			
Thr	Ile	Arg	Gly	Val	Thr	Gln	Asp	Phe	Ser	Thr	Thr	Pro	Leu	Ser
1235						1240					1245			
Val	Glu	Val	Leu	Thr	Glu	Glu	Val	Pro	Asp	Met	Gly	Asn	Leu	Thr
1250						1255					1260			
Val	Thr	Glu	Val	Ser	Trp	Asp	Ala	Leu	Arg	Leu	Asn	Trp	Thr	Thr
1265						1270					1275			
Pro	Asp	Gly	Thr	Tyr	Asp	Gln	Phe	Thr	Ile	Gln	Val	Gln	Glu	Ala
1280						1285					1290			
Asp	Gln	Val	Glu	Glu	Ala	His	Asn	Leu	Thr	Val	Pro	Gly	Ser	Leu
1295						1300					1305			
Arg	Ser	Met	Glu	Ile	Pro	Gly	Leu	Arg	Ala	Gly	Thr	Pro	Tyr	Thr
1310						1315					1320			
Val	Thr	Leu	His	Gly	Glu	Val	Arg	Gly	His	Ser	Thr	Arg	Pro	Leu
1325						1330					1335			
Ala	Val	Glu	Val	Val	Thr	Glu	Asp	Leu	Pro	Gln	Leu	Gly	Asp	Leu
1340						1345					1350			
Ala	Val	Ser	Glu	Val	Gly	Trp	Asp	Gly	Leu	Arg	Leu	Asn	Trp	Thr
1355						1360					1365			
Ala	Ala	Asp	Asn	Ala	Tyr	Glu	His	Phe	Val	Gln	Val	Gln	Glu	Val
1370						1375					1380			
Asn	Lys	Val	Glu	Ala	Ala	Gln	Asn	Leu	Thr	Leu	Pro	Gly	Ser	Leu
1385						1390					1395			
Arg	Ala	Val	Asp	Ile	Pro	Gly	Leu	Glu	Ala	Ala	Thr	Pro	Tyr	Arg
1400						1405					1410			
Val	Ser	Ile	Tyr	Gly	Val	Ile	Arg	Gly	Tyr	Arg	Thr	Pro	Val	Leu
1415						1420					1425			
Ser	Ala	Glu	Ala	Ser	Thr	Ala	Lys	Glu	Pro	Glu	Ile	Gly	Asn	Leu
1430						1435					1440			
Asn	Val	Ser	Asp	Ile	Thr	Pro	Glu	Ser	Phe	Asn	Leu	Ser	Trp	Met
1445						1450					1455			
Ala	Thr	Asp	Gly	Ile	Phe	Glu	Thr	Phe	Thr	Ile	Glu	Ile	Ile	Asp
1460						1465					1470			
Ser	Asn	Arg	Leu	Leu	Glu	Thr	Val	Glu	Tyr	Asn	Ile	Ser	Gly	Ala
1475						1480					1485			

-continued

Glu Arg	Thr Ala His Ile Ser	Gly Leu Pro Pro Ser	Thr Asp Phe
1490	1495	1500	
Ile Val	Tyr Leu Ser Gly Leu	Ala Pro Ser Ile Arg	Thr Lys Thr
1505	1510	1515	
Ile Ser	Ala Thr Ala Thr Thr	Glu Ala Leu Pro Leu	Leu Glu Asn
1520	1525	1530	
Leu Thr	Ile Ser Asp Ile Asn	Pro Tyr Gly Phe Thr	Val Ser Trp
1535	1540	1545	
Met Ala	Ser Glu Asn Ala Phe	Asp Ser Phe Leu Val	Thr Val Val
1550	1555	1560	
Asp Ser	Gly Lys Leu Leu Asp	Pro Gln Glu Phe Thr	Leu Ser Gly
1565	1570	1575	
Thr Gln	Arg Lys Leu Glu Leu	Arg Gly Leu Ile Thr	Gly Ile Gly
1580	1585	1590	
Tyr Glu	Val Met Val Ser Gly	Phe Thr Gln Gly His	Gln Thr Lys
1595	1600	1605	
Pro Leu	Arg Ala Glu Ile Val	Thr Glu Ala Glu Pro	Glu Val Asp
1610	1615	1620	
Asn Leu	Leu Val Ser Asp Ala	Thr Pro Asp Gly Phe	Arg Leu Ser
1625	1630	1635	
Trp Thr	Ala Asp Glu Gly Val	Phe Asp Asn Phe Val	Leu Lys Ile
1640	1645	1650	
Arg Asp	Thr Lys Lys Gln Ser	Glu Pro Leu Glu Ile	Thr Leu Leu
1655	1660	1665	
Ala Pro	Glu Arg Thr Arg Asp	Leu Thr Gly Leu Arg	Glu Ala Thr
1670	1675	1680	
Glu Tyr	Glu Ile Glu Leu Tyr	Gly Ile Ser Lys Gly	Arg Arg Ser
1685	1690	1695	
Gln Thr	Val Ser Ala Ile Ala	Thr Thr Ala Met Gly	Ser Pro Lys
1700	1705	1710	
Glu Val	Ile Phe Ser Asp Ile	Thr Glu Asn Ser Ala	Thr Val Ser
1715	1720	1725	
Trp Arg	Ala Pro Thr Ala Gln	Val Glu Ser Phe Arg	Ile Thr Tyr
1730	1735	1740	
Val Pro	Ile Thr Gly Gly Thr	Pro Ser Met Val Thr	Val Asp Gly
1745	1750	1755	
Thr Lys	Thr Gln Thr Arg Leu	Val Lys Leu Ile Pro	Gly Val Glu
1760	1765	1770	
Tyr Leu	Val Ser Ile Ile Ala	Met Lys Gly Phe Glu	Glu Ser Glu
1775	1780	1785	
Pro Val	Ser Gly Ser Phe Thr	Thr Ala Leu Asp Gly	Pro Ser Gly
1790	1795	1800	
Leu Val	Thr Ala Asn Ile Thr	Asp Ser Glu Ala Leu	Ala Arg Trp
1805	1810	1815	
Gln Pro	Ala Ile Ala Thr Val	Asp Ser Tyr Val Ile	Ser Tyr Thr
1820	1825	1830	
Gly Glu	Lys Val Pro Glu Ile	Thr Arg Thr Val Ser	Gly Asn Thr
1835	1840	1845	
Val Glu	Tyr Ala Leu Thr Asp	Leu Glu Pro Ala Thr	Glu Tyr Thr
1850	1855	1860	
Leu Arg	Ile Phe Ala Glu Lys	Gly Pro Gln Lys Ser	Ser Thr Ile

-continued

1865	1870	1875
Thr Ala Lys Phe Thr Thr	Asp Leu Asp Ser Pro	Arg Asp Leu Thr
1880	1885	1890
Ala Thr Glu Val Gln Ser	Glu Thr Ala Leu Leu	Thr Trp Arg Pro
1895	1900	1905
Pro Arg Ala Ser Val Thr	Gly Tyr Leu Leu Val	Tyr Glu Ser Val
1910	1915	1920
Asp Gly Thr Val Lys Glu	Val Ile Val Gly Pro	Asp Thr Thr Ser
1925	1930	1935
Tyr Ser Leu Ala Asp Leu	Ser Pro Ser Thr His	Tyr Thr Ala Lys
1940	1945	1950
Ile Gln Ala Leu Asn Gly	Pro Leu Arg Ser Asn	Met Ile Gln Thr
1955	1960	1965
Ile Phe Thr Thr Ile Gly	Leu Leu Tyr Pro Phe	Pro Lys Asp Cys
1970	1975	1980
Ser Gln Ala Met Leu Asn	Gly Asp Thr Thr Ser	Gly Leu Tyr Thr
1985	1990	1995
Ile Tyr Leu Asn Gly Asp	Lys Ala Gln Ala Leu	Glu Val Phe Cys
2000	2005	2010
Asp Met Thr Ser Asp Gly	Gly Gly Trp Ile Val	Phe Leu Arg Arg
2015	2020	2025
Lys Asn Gly Arg Glu Asn	Phe Tyr Gln Asn Trp	Lys Ala Tyr Ala
2030	2035	2040
Ala Gly Phe Gly Asp Arg	Arg Glu Glu Phe Trp	Leu Gly Leu Asp
2045	2050	2055
Asn Leu Asn Lys Ile Thr	Ala Gln Gly Gln Tyr	Glu Leu Arg Val
2060	2065	2070
Asp Leu Arg Asp His Gly	Glu Thr Ala Phe Ala	Val Tyr Asp Lys
2075	2080	2085
Phe Ser Val Gly Asp Ala	Lys Thr Arg Tyr Lys	Leu Lys Val Glu
2090	2095	2100
Gly Tyr Ser Gly Thr Ala	Gly Asp Ser Met Ala	Tyr His Asn Gly
2105	2110	2115
Arg Ser Phe Ser Thr Phe	Asp Lys Asp Thr Asp	Ser Ala Ile Thr
2120	2125	2130
Asn Cys Ala Leu Ser Tyr	Lys Gly Ala Phe Trp	Tyr Arg Asn Cys
2135	2140	2145
His Arg Val Asn Leu Met	Gly Arg Tyr Gly Asp	Asn Asn His Ser
2150	2155	2160
Gln Gly Val Asn Trp Phe	His Trp Lys Gly His	Glu His Ser Ile
2165	2170	2175
Gln Phe Ala Glu Met Lys	Leu Arg Pro Ser Asn	Phe Arg Asn Leu
2180	2185	2190
Glu Gly Arg Arg Lys Arg	Ala	
2195	2200	

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 8578

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Supplied by Invitrogen of Carlsbad, California

## -continued

&lt;400&gt; SEQUENCE: 3

gacggatcgg gagatctccc gatccctat ggtcgactct cagtacaatc tgctctgatg	60
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggctcgt gagtagtgcg	120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc	180
ttagggttag gcgttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt	240
gattattgac tagttattaa tagtaataca ttacggggtc attagtcat agcccatata	300
tggagttccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc	360
cccgccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc	420
attgacgtca atgggtggac tatttacggt aaactgcca cttggcagta catcaagtgt	480
atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt	540
atgccagta catgacctta tgggactttc ctacttgga gtacatctac gtattagtca	600
tcgctattac catggtgatg cggttttggc agtacatcaa tggcggtgga tagcggtttg	660
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc	720
aaaatcaacg ggactttcca aaatgtcgtg acaactccgc ccattgacg caaatgggcg	780
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaacca	840
ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa gctggctagc	900
gtttaaactt aagcttacca tgggggggtc tcatcatcat catcatcatg gtatggctag	960
catgactggt ggacagcaaa tgggtcggga tctgtacgac gatgacgata aggtacctaa	1020
ggatcagctt ggagttgatc ccgtcgtttt acaacgtcgt gactgggaaa accctggcgt	1080
tacccaactt aatcgccttg cagcacatcc ccctttcgcc agctggcgta atagcgaaga	1140
ggccgcacc gatcgccctt cccaacagtt gcgcagcctg aatggcgaat ggcgctttgc	1200
ctggtttccg gcaccagaag cgggtccgga aagctggctg gagtgcgac ttcctgaggc	1260
cgatactgtc gtcgtccctt caaactggca gatgcacggt tacgatgcgc ccatctacac	1320
caacgtaacc tatccatta cgggtcaatcc gccgtttgtt cccacggaga atccgacggg	1380
ttgttactcg ctacattta atgttgatga aagctggcta caggaaggcc agacgcgaat	1440
tatttttgat ggcgtaact cggcgtttca tctgtggtgc aacgggcgct gggtcggtta	1500
cggccaggac agtcgtttgc cgtctgaatt tgacctgagc gcatttttac gcgccggaga	1560
aaaccgcctc gcggtgatgg tgctgcgttg gagtgcggc agttatcttg aagatcagga	1620
tatgtggcgg atgagcggca ttttcgtga cgtctcgttg ctgcataaac cgactacaca	1680
aatcagcgat ttccatgttg ccaactcgctt taatgatgat ttcagccgag ctgtactgga	1740
ggctgaagtt cagatgtgag gcgagttgcg tgactaccta cgggtaacag tttctttatg	1800
gcagggtgaa acgcaggtcg ccagcggcac cgcgcctttc ggcggtgaaa ttatcgatga	1860
gcgtggtggt tatgccgacg gcgtcacact acgtctgaac gtcgaaaacc cgaactgtg	1920
gagcgccgaa atcccgaaac tctatcgtgc ggtggttgaa ctgcacaccg ccgacggcac	1980
gctgattgaa gcagaagcct gcgatgtcgg tttccgcgag gtgcggattg aaaatggtct	2040
gctgctgctg aacggcaagc cgttgctgat tcgaggcgtt aacgctcacg agcatcatcc	2100
tctgcatggt caggctcatg atgagcagac gatggtgcag gatattctgc tgatgaagca	2160
gaacaacttt aacgcctgtc gctgttcgca ttatccgaac catccgctgt ggtacacgct	2220

## -continued

gtgcgaccgc tacggcctgt atgtgggtgga tgaagccaat attgaaaccc acggcatggt	2280
gccaatgaat cgtctgaccg atgatccgcg ctggctaccg gcgatgagcg aacgcgtaac	2340
gcgaatgggt cagcgcgcat gtaatcaccg gagtgtgac atctggctgc tggggaatga	2400
atcaggccac ggcgctaate acgacgcgct gtatcgctgg atcaaactctg tcgacccctc	2460
ccgcccgggt cagtatgaag gcggcggagc cgacaccacg gccaccgata ttatttgccc	2520
gatgtacgcg cgcgtggatg aagaccagcc ctccccggct gtgccgaaat ggtccatcaa	2580
aaaaatggctt tcgctacctg gagagacgcg cccgctgac ctttgcgaaat acgcccacgc	2640
gatgggtaac agtcttggcg gtttcgctaa atactggcag gcgtttcgtc agtatccccg	2700
ttacagggc ggcttcgtct gggactgggt ggatcagtcg ctgattaaat atgatgaaaa	2760
cggcaacccc tggtcggctt acggcgggtga ttttggcgat acgccgaacg atcgccagtt	2820
ctgtatgaac ggtctggtct ttgccgaccg caccgcgcat ccagcgctga cggaagcaaa	2880
acaccagcag cagtttttcc agttccgttt atccgggcaa accatcgaag tgaccagcga	2940
atacctgttc cgtcatagcg ataacgagct cctgcactgg atggtggcgc tggatggtaa	3000
gccgctggca agcggtgaa gtcctctgga tgcgctcca caaggtaaac agttgattga	3060
actgcctgaa ctaccgcagc cggagagcgc cgggcaactc tggctcacag tacgcgtagt	3120
gcaaccgaac gcgaccgcat ggtcagaagc cgggcacatc agcgccctggc agcagtggcg	3180
tctggcggaa aaactcagtg tgacgctccc cgcgcgtcc caccgatcc cgcactgac	3240
caccagcgaa atggattttt gcatcgagct gggtaataag cgttggcaat ttaaccgcca	3300
gtcaggcttt ctttcacaga tgtggattgg cgataaaaa caactgctga cgcgcgtgcg	3360
cgatcagttc acccgtgcac cgctggataa cgacattggc gtaagtgaag cgaccgcgat	3420
tgaccctaac gcctgggtcg aacgtggaa ggcggcgggc cattaccagg ccgaagcagc	3480
gttgttcgag tgcacggcag atacacttgc tgatgcggtg ctgattacga ccgctcacgc	3540
gtggcagcat caggggaaaa ccttatttat cagccggaaa acctaccgga ttgatggtag	3600
tggtaaatg gcgattaccg ttgatgttga agtggcgcgc gatacaccgc atccgcgcg	3660
gattggcctg aactgccagc tggcgcaggt agcagagcgg gtaaaactggc tcggattagg	3720
gccgcaagaa aactatccc accgccttac tgcgcctgt tttgaccgct gggatctgcc	3780
attgtcagac atgtataccc cgtacgtctt cccgagcgaa aacggtctgc gctgcgggac	3840
gcgcgaattg aattatggcc cacaccagtg gcgcggcgac ttccagttca acatcagccg	3900
ctacagtcaa cagcaactga tggaaaccag ccacgcgcat ctgctgcacg cggaagaagg	3960
cacatggctg aatatcgacg gtttccatat ggggattggt ggcgacgact cctggagccc	4020
gtcagtatcg gcggagtacc agctgagcgc cggctcgtac cattaccagt tggctcgtg	4080
tcaaaaataa taaagccgaa ttctgcagat atccagcaca gtggcggccg ctcgagtcta	4140
gagggcccgt ttaaaccgcg tgatcagcct cgactgtgcc ttctagtgtc cagccactctg	4200
ttgtttgccc ctccccctgt ccttccttga ccttggaaag tgccactccc actgtccttt	4260
cctaataaaa tgaggaaatt gcatcgcatt gtctgagtag gtgtcattct attctggggg	4320
gtgggtggg gcaggacagc aagggggagg attgggaaga caatagcagg catgctgggg	4380
atgcggtggg ctctatggct tctgagcgcg aaagaaccag ctggggctct aggggggtatc	4440
cccacgcgcc ctgtagcggc gcattaagcg cggcgggtgt ggtggttacg cgcagcgtga	4500

## -continued

ccgctacact	tgccagcgcc	ctagcgccc	ctcctttcgc	tttcttcctt	tcctttctcg	4560
ccacgttcgc	cggctttccc	cgtcaagctc	taaatcgggg	catcccttta	gggttcgat	4620
ttagtgtctt	acggcacctc	gacccccaaa	aacttgatta	gggtgatggg	tcacgtagt	4680
ggccatcgcc	ctgatagacg	gtttttcgcc	ctttgacgtt	ggagtccacg	ttctttaata	4740
gtggactctt	gttccaaact	ggaacaacac	tcaaccctat	ctcggcttat	tccttttgatt	4800
tataagggat	tttggggatt	tcggcctatt	ggttaaaaaa	tgagctgatt	taacaaaaat	4860
ttaacgcgaa	ttaatctgt	ggaatgtgtg	tcagttagg	tgtggaaagt	ccccaggctc	4920
cccaggcagg	cagaagtatg	caaagcatgc	atctcaatta	gtcagcaacc	aggtgtggaa	4980
agtccccagg	ctccccagca	ggcagaagta	tgcaaagcat	gcctctcaat	tagtcagcaa	5040
ccatagtccc	gcccctaact	ccgcccctcc	cggcccctaac	tccgcccagt	tccgcccatt	5100
ctccgcccc	tggtgacta	atttttttta	tttatgcaga	ggcggaggcc	gcctctgcct	5160
ctgagctatt	ccagaagtag	tgaggaggct	tttttgagg	cctaggcttt	tgcaaaaagc	5220
tcccgggagc	ttgtatatcc	attttcggt	ctgatcaaga	gacaggatga	ggatcgtttc	5280
gcatgattga	acaagatgga	ttgcacgcag	gttctccggc	cgttggggtg	gagaggctat	5340
tcggctatga	ctgggcacaa	cagacaatcg	gctgctctga	tgccgccgtg	ttccggctgt	5400
cagcgcagg	gcgcccgtt	ctttttgtca	agaccgacct	gtccgggtgc	ctgaatgaac	5460
tgaggagcga	ggcagcgcg	ctatcggtgc	tgccacgac	ggcggttct	tgccagctg	5520
tgctcgagct	gtcactgaa	gcgggaagg	actggctgct	attgggcgaa	gtgccggggc	5580
aggatctcct	gtcatctcac	cttgctcctg	ccgagaaagt	atccatcatg	gctgatgcaa	5640
tgccggcggt	gcatacgctt	gatccggcta	cctgcccatt	cgaccaccaa	gcgaaacatc	5700
gcatcgagcg	agcacgtact	cggatggaag	cgggtcttgt	cgatcaggat	gatctggacg	5760
aagagcatca	ggggctcgcg	ccagccgaac	tggtcgccag	gctcaaggcg	cgcatgccc	5820
acggcgagga	tctcgtcggt	acctatggcg	atgcctgctt	gccgaatata	atggtggaaa	5880
atggccgctt	ttctggattc	atcgactgtg	gccggctggg	tgtggcgga	cgctatcagg	5940
acatagcggt	ggctaccggt	gatattgctg	aagagcttgg	cggcgaaatg	gctgaccgct	6000
tcctcgtgct	ttacggtatc	gccgctccc	attcgacg	catcgcttc	tatcgcttc	6060
ttgacgagtt	cttctgagcg	ggactctggg	gttcgaaatg	accgaccaag	cgacgccc	6120
cctgccatca	cgagatttct	attccaccgc	cgccttctat	gaaagggttg	gcttcggaat	6180
cgttttccgg	gacgcgggct	ggatgatcct	ccagcgcg	gatctcatgc	tgaggttctt	6240
cgcaccccc	aacttgctta	ttgcagctta	taatgggtac	aaataaagca	atagcatcac	6300
aaatttcaca	aataaagcat	ttttttcact	gcattctagt	tgtggttgt	ccaaactcat	6360
caatgtatct	tatcatgtct	gtataccgtc	gacctctagc	tagagcttgg	cgtaatcatg	6420
gtcatagctg	ttctcgtgtg	gaaattgtta	tccgctcaca	attccacaca	acatacgagc	6480
cggagcata	aagtgtaaa	cctgggggtg	ctaagtagtg	agctaactca	cattaattgc	6540
gttgcgctca	ctgcccgtt	tccagtcggg	aaacctgtcg	tgccagctgc	attaatgaat	6600
cggccaacgc	gcggggagag	gcggtttcg	tattgggcgc	tcttcgctt	cctcgctcac	6660
tgactcgctg	cgctcggtcg	ttcggtcg	gcgagcggt	tcagctcact	caaaggcggt	6720
aatacggtta	tccacagaat	caggggataa	cgcaggaaag	aacatgtgag	caaaaggcca	6780

## -continued

gcaaaaggcc aggaaccgta aaaaggccgc gttgctggcg tttttccata ggtcccgccc	6840
ccctgacgag catcacaaaa atcgacgctc aagtcagagg tggcgaaacc cgacaggact	6900
ataaagatac caggcgcttc cccctggaag ctccctcgtg cgctctcctg ttccgaccct	6960
gccgcttacc ggatacctgt ccgcctttct cccttcggga agcgtggcgc tttctcaatg	7020
ctcacgctgt aggtatctca gttcgggtga ggtcgttcgc tccaagctgg gctgtgtgca	7080
cgaaccccc gttcagcccc accgctgcgc cttatccggt aactatcgtc ttgagtccaa	7140
cccggtaaga cagcacttat cgccactggc agcagccact ggtaacagga ttagcagagc	7200
gaggtatgta gccggtgcta cagagtctct gaagtggtag cctaactacg gctacactag	7260
aaggacagta ttggtatct gcgctctgct gaagccagtt accttcggaa aaagagttgg	7320
tagctcttga tccggcaaac aaaccaccgc tggtagcggg ggtttttttg tttgcaagca	7380
gcagattacg cgcaaaaaa aaggatctca agaagatcct ttgatctttt ctacggggtc	7440
tgacgctcag tggaacgaaa actcagctta agggattttg gtcagtatat tatcaaaaag	7500
gatcttcacc tagatccttt taaattaaaa atgaagtttt aaatcaatct aaagtatata	7560
tgagtaaaact tggctctgaca gttaccaatg cttaatcagt gaggcaccta tctcagcgat	7620
ctgtctatct cgttcaccca tagttgcctg actccccgtc gtgtagataa ctacgatacg	7680
ggagggttta ccatctggcc ccagtgcgc aatgataccg cgagaccac gctcaccggc	7740
tccagattta tcagcaataa accagccagc cggaagggcc gagcgagaa gtggtcctgc	7800
aactttatcc gcctccatcc agtctattaa ttggtgcgg gaagctagag taagtagttc	7860
gccagttaat agtttgcgca acggtgttgc cattgctaca ggcatcgtgg tgcacgctc	7920
gtcgtttggt atggcttcat tcagctccgg tccccacga tcaaggcgag ttacatgatc	7980
ccccatggtg tgcaaaaaa cggttagctc cttcgggtcct ccgacgttg tcagaagtaa	8040
gttgcccgca gtgttatcac tcagtgttat ggcagcactg cataattctc ttactgtcat	8100
gccatccgta agatgccttt ctgtgactgg tgagtactca accaagtcac tctgagaata	8160
gtgtatgcgg cgaccgagtt gctcttgccc ggcgtcaata cgggataata ccgcgccaca	8220
tagcagaact ttaaaagtgc tcattcattg aaaacgttct tcggggcgaa aactctcaag	8280
gatcttaccg ctgttgagat ccagttcgat gtaaccact cgtgcaccca actgatcttc	8340
agcatctttt actttcacca gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc	8400
aaaaaaggga ataaggcgca caccgaaatg ttgaatactc atactcttcc tttttcaata	8460
ttattgaagc atttatcagg gttattgtct catgagcgga tacatatattg aatgtattta	8520
gaaaaataaa caaatagggg ttccgcgcac atttcccga aaagtgccac ctgacgtc	8578

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 4748

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Provided by Dr. Brett Levay-Young of the  
University of Minnesota

&lt;400&gt; SEQUENCE: 4

tagttattaa tagtaataca ttacggggtc attagttcat agcccatata tggagttccg	60
cgttacataa cttacggtta atggcccgcc tggctgaccg cccaacgacc ccgcccatt	120
gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca	180

-continued

---

atgggtggag tatttacggt aaactgccca cttggcagta catcaagtgt atcatatgcc	240
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta	300
catgacctta tgggactttc ctacttgga gtacatctac gtattagtca tcgctattac	360
catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggttg actcacggg	420
atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg	480
ggactttcca aaatgtcgta acaactccgc cccattgacg caaatgggag gtaggcgtgt	540
acggtgggag gtctatataa gcagagctgg tttagtgaac cgtcagatcc gctagcgcta	600
ccggtcgcca ccatggtgag caagggcgag gagctgttca ccggggtggt gcccatcctg	660
gtcagctgg acggcgacgt aaacggccac aagttcagcg tgtccggcga gggcgagggc	720
gatgccacct acggcaagct gaccctgaag ttcactctga ccaccggcaa gctgcccggtg	780
ccctggccca ccctcgtgac caccctgacc tacggcgtgc agtgcctcag ccgctacccc	840
gaccacatga agcagcacga cttcttcaag tccgccatgc ccgaaggcta cgtccaggag	900
cgcaccatct tcttcaagga cgacggcaac tacaagacc gcgccgaggt gaagttcgag	960
ggcgacaccc tgggtaaccg catcgagctg aagggcatcg acttcaagga ggacggcaac	1020
atcctggggc acaagctgga gtacaactac aacagccaca acgtctatat catggccgac	1080
aagcagaaga acggcatcaa ggtgaacttc aagatccgcc acaacatcga ggacggcagc	1140
gtgcagctcg ccgaccacta ccagcagaac acccccatcg gcgacggccc cgtgctgctg	1200
cccgacaacc actacctgag caccagtc ccctgagca aagacccaa cgagaagcgc	1260
gatcacatgg tcctgctgga gttcgtgacc gccgccggga tcaactctcg catggacgag	1320
ctgtacaagt actcagatct cgagctcaag cttaaccctc cggacgagag cggccctggc	1380
tgtatgtcct gcaagtgcgt gctgtcctga tcaccggatc tagataactg atcataatca	1440
gccataccac attttagtag gttttacttg ctttaaaaaa cctccacac ctccccctga	1500
acctgaaaca taaaatgaat gcaattgttg ttgttaactt gtttattgca gcttataatg	1560
gttacaataa aagcaatagc atcacaatt tcacaaataa agcatttttt tcaactgcatt	1620
ctagttgtgg tttgtccaaa ctcatcaatg tatcttaacg cgtaaattgt aagcggtaat	1680
attttgtaa aattcgcgtt aaatttttgt taaatcagct cattttttta ccaataggcc	1740
gaaatcggca aaatccctta taaatcaaaa gaatagaccg agatagggtt gagtgttgtt	1800
ccagtttgga acaagagtcc actattaaag aacgtggact ccaacgtcaa agggcgaaaa	1860
accgtctatc agggcgatgg ccactacgt gaaccatcac cctaatacag ttttttggg	1920
tcgaggtgcc gtaaagcact aaatcggaac cctaaaggga gccccgatt tagagcttga	1980
cggggaagc cggcgaaagt ggcgagaaag gaagggaaga aagcgaaag agcgggcgct	2040
agggcgctgg caagtgtagc ggtcacgctg cgcgtaacca ccacaccgc cgcgcttaat	2100
gcgcgctac agggcgctgc aggtggcact tttcggggaa atgtgcgcgg aaccctatt	2160
tgtttatatt tctaaatata ttcaaatatg tatccgctca tgagacaata accctgataa	2220
atgcttcaat aatattgaaa aaggaagagt cctgaggcgg aaagaaccag ctgtggaatg	2280
tgtgtcagtt aggggtgtga aagtccccag gctccccagc aggcagaagt atgcaagca	2340
tgcatctcaa ttagtcagca accaggtgtg gaaagtcccc aggtccccca gcaggcagaa	2400
gtatgcaaa catgcatctc aattagtcag caaccatagt cccgcccta actccgcca	2460



-continued

tccccccct aactccgcc agttccgcc attctccgcc ccatggctga ctaatttttt	2520
ttatttatgc agaggccgag gccgcctcgg cctctgagct attccagaag tagtgaggag	2580
gcttttttgg aggcctaggc ttttgcaaa atcgatcaag agacaggatg aggatcgttt	2640
cgcatgattg aacaagatgg attgcacgca ggttctccgg ccgcttgggt ggagaggcta	2700
ttcggtatg actgggcaca acagacaatc ggctgctctg atgccgccgt gttccggctg	2760
tcagcgcagg ggcgcccggt tctttttgtc aagaccgacc tgtccggtgc cctgaatgaa	2820
ctgcaagacg aggcagcgcg gctatcgtag ctggccacga cggcggttcc ttgcgcagct	2880
gtgctcgacg ttgtcactga agcgggaagg gactggctgc tattgggcga agtgccgggg	2940
caggatctcc tgtcatctca ccttgctcct gccgagaaag tatccatcat ggctgatgca	3000
atcgcgcggc tgcatacgct tgatccggct acctgcccac tcgaccacca agcgaaacat	3060
cgcatcgagc gagcagctac tcggatggaa gccggtcttg tcgatcagga tgatctggac	3120
gaagagcatc aggggctcgc gccagccgaa ctgttcgcca ggctcaaggc gagcatgccc	3180
gacggcgagg atctcgtcgt gaccatggc gatgcctgct tgccgaatat catggtggaa	3240
aatggccgct tttctggatt catcgactgt ggccggctgg gtgtggcgga ccgctatcag	3300
gacatagcgt tggctacccg tgatattgct gaagagcttg gcggcgaaat ggctgaccgc	3360
ttcctcgtgc tttacggtat cggcgctccc gattcgagc gcacgcctt ctatcgctt	3420
cttgacgagt tcttctgagc gggactctgg ggttcgaaat gaccgaccaa gcgacgccc	3480
acctgccatc acgagatttc gattccaccg ccgccttcta tgaaggttg ggcttcggaa	3540
tcgttttccg ggacgcggc tggatgatcc tccagcgcg ggatctcatg ctggagtct	3600
tcgccacccc tagggggagg ctaactgaaa cacggaagga gacaataccg gaaggaaccc	3660
gcgctatgac ggcaataaaa agacagaata aaacgcacgg tgttgggtcg tttgttcata	3720
aacgcggggt tcggtcccag ggctggcact ctgtcgatc cccaccgaga cccattggg	3780
gccaatcacg ccgcttttct tccttttccc caccacccc cccaagtctg ggtgaaggcc	3840
cagggtcgc agccaacgct gggcgggcag gccctgccat agcctcaggt tactcatata	3900
tactttagat tgatttaaaa ctctattttt aatttaaaag gatctagggt aagatccttt	3960
ttgataatct catgacaaa atcccttaac gtgagttttc gttccactga gcgtcagacc	4020
ccgtagaaaa gatcaaagga tcttcttgag atcctttttt tctgcgcgta atctgctgct	4080
tgcaaaaaa aaaaccaccg ctaccagcgg tggtttggtt gccggatcaa gagctaccaa	4140
ctctttttcc gaaggttaact ggcttcagca gagcgagat accaaatact gtccttctag	4200
tgtagccgta gttaggccac cacttcaaga actctgtagc accgcctaca tacctcgctc	4260
tgtaaatcct gttaccagtg gctgctgcca gtggcgataa gtcgtgtctt accgggttg	4320
actcaagacg atagttaccg gataaggcgc agcggtcggg ctgaacgggg ggttcgtgca	4380
cacagcccag cttggagcga acgacctaca ccgaactgag atacctacag cgtgagctat	4440
gagaaagcgc cagcgttccc gaaggagaa aggcggacag gtatccggtg agcggcaggg	4500
tcggaacagg agagcgcacg agggagcttc cagggggaaa cgctgtgtat ctttatagtc	4560
ctgtcggggt tcgccacctc tgacttgagc gtcgattttt gtgatgctcg tcaggggggc	4620
ggagcctatg gaaaaacgcc agcaacgcgg cctttttacg gttcctggcc ttttgctggc	4680
cttttgctca catgttcttt cctgcgttat cccctgattc tgtggataac cgtattaccg	4740

---

-continued

---

ccatgcat

4748

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 4992

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Supplied by BD Biosciences Clonetechn of Palo Alto, California

&lt;400&gt; SEQUENCE: 5

tagttattaa tagtaatacaa ttacggggtc attagttcat agcccatata tggagttccg	60
cgttacataa cttacggtaa atggcccgcc tggctgaccg ccaacgacc cccgccatt	120
gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca	180
atgggtggag tatttacggt aaactgcca cttggcagta catcaagtgt atcatatgcc	240
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta	300
catgacctta tgggactttc ctacttgga gtacatctac gtattagtca tcgctattac	360
catggtgatg cggttttgcc agtacatcaa tgggcgtgga tagcggtttg actcacggg	420
atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg	480
ggactttcca aaatgtcgta acaactccgc cccattgacg caaatgggcg gtaggcgtgt	540
acggtgggag gtctatataa gcagagctgg tttagtgaac cgtcagatcc gctagcgcta	600
ccggtcgcca ccatggtgag caaggcgag gagctgttca ccggggtggt gccatcctg	660
gtcagctgg acggcgacgt aaacggccac aagttcagcg tgtccggcga gggcgaggcg	720
gatgccacct acggcaagct gacctgaag ttcactctga ccaccggcaa gctgcccgtg	780
ccctggccca ccctcgtgac caccctgacc tacggcgtgc agtgcttcag ccgctacccc	840
gaccacatga agcagcacga cttcttcaag tccgccatgc ccgaaggcta cgtccaggag	900
cgacccatct tcttcaagga cgacggcaac tacaagacc gcgccgaggt gaagttcgag	960
ggcgacaccc tgggtgaacc catcgagctg aaggcgatcg acttcaagga ggacggcaac	1020
atcctggggc acaagctgga gtacaactac aacagccaca acgtctatat catggccgac	1080
aagcagaaga acggcatcaa ggtgaacttc aagatccgcc acaacatga ggacggcagc	1140
gtgcagctcg ccgaccacta ccagcagaac acccccatcg gcgacggccc cgtgctgctg	1200
cccgaacaac actacctgag caccagtcg gccctgagca aagaccccaa cgagaagcgc	1260
gatcacatgg tcctgctgga gttcgtgacc gccgccggga tcaactctcg catggacgag	1320
ctgtacaagt actcagatct cgagctcaag cttaccatgg ggggttttca tcatcatcat	1380
catcatggta tggctagcat gactggtgga cagcaaatgg gtcgggatct gtaacagcat	1440
gacgataagg ggactgctgc ggccaatgag aacgacttct tcgccaagcg caagagaact	1500
gcgcaggaga acaaggcgtc gaacgacgtc cctccagggt gtccctctcc aaacgtggct	1560
cctgggggtg gcgcggtgga gcagaccccg cgcaaacgtc tgagatgagg atccagtgtg	1620
gtggaattct gcagatatcc agcacagtgg cggccgctcg agtctagata actgatcata	1680
atcagccata ccacatttgt agaggtttta cttgctttta aaaacctccc acacctccc	1740
ctgaacctga aacataaaat gaatgcaatt gttgtgttta acttgtttat tgcagcttat	1800
aatggttaca aataaagcaa tagcatcaca aatttcacaa ataaagcatt tttttcactg	1860

## -continued

cattctagtt gtggtttgtc caaactcatc aatgtatctt aacgcgtaaa ttgtaagcgt	1920
taatatatttg ttaaaattcg cgttaaattt ttgttaaadc agctcatttt ttaaccaata	1980
ggccgaaatc ggcaaaatcc cttataaadc aaaagaatag accgagatag ggttgagtgt	2040
tggtccagtt tggaacaaga gtccactatt aaagaacgtg gactccaacg tcaaagggcg	2100
aaaaaccgtc tatcagggcg atggccact acgtgaacca tcaccctaata caagtttttt	2160
ggggtcgagg tgccgtaaag cactaaatcg gaaccctaaa gggagccccc gatttagagc	2220
ttgacgggga aagccggcga acgtggcgag aaaggaaggg aagaaagcga aaggagcggg	2280
cgctagggcg ctggcaagtg tagcggtcac gctgcgcgta accaccacac ccgccgcgct	2340
taatgcgccg ctacagggcg cgtcaggtgg cacttttcgg ggaatgtgc gcggaacccc	2400
tatttgttta tttttctaaa tacattcaaa tatgtatccg ctcatgagac aataacctg	2460
ataaatgctt caataatatt gaaaagga gagtcctgag gcgaaagaa ccagctgtgg	2520
aatgtgtgtc agttagggtg tggaaagtcc ccaggctccc cagcaggcag aagtatgcaa	2580
agcatgcac tcaattagtc agcaaccagg tgtggaaagt cccagggctc cccagcaggc	2640
agaagtatgc aaagcatgca tctcaattag tcagcaacca tagtcccgcc cctaactccg	2700
cccatcccg ccctaactcc gccagttcc gccattctc cgcgccatgg ctgactaatt	2760
ttttttattt atgcagagcg cgaggccgcc tcggcctctg agctattcca gaagtagtga	2820
ggaggctttt ttggaggcct aggccttttc aaagatcgat caagagacag gatgaggatc	2880
gtttcgcagt attgaacaag atggattgca cgcaggttct ccggccgctt gggtgagag	2940
gctattcggc tatgactggg cacaacagac aatcggctgc tctgatgccg ccgtgttccg	3000
gctgtcagcg caggggcgcc cggttctttt tgtcaagacc gacctgtccg gtgccctgaa	3060
tgaactgcaa gacgaggcag cgcggctatc gtggctggcc acgacggcg ttctctgcgc	3120
agctgtgtct gacgttgtca ctgaagcggg aagggactgg ctgctattgg gcgaagtgcc	3180
ggggcaggat ctctgtcat ctacactgc tcctgccgag aaagtatcca tcatggetga	3240
tgcaatgcgg cggctgcata cgcttgatcc ggctacctgc ccattcgacc accaagcgaa	3300
acatcgcac gagcgagcac gtactcggat ggaagccggt cttgtcgatc aggatgatct	3360
ggacgaagag catcaggggc tcgcgccagc cgaactgttc gccaggctca aggcgagcat	3420
gcccagcggc gaggatctcg tcgtgaccca tggcgatgcc tgcttgccga atatcatggt	3480
ggaaaatggc cgcttttctg gattcatcga ctgtggccgg ctgggtgtgg cggaccgcta	3540
tcaggacata gcgttggtta ccctgatata tgctgaagag cttggcgcg aatgggctga	3600
ccgcttctc gtgctttacg gtatcgccgc tcccgattcg cagcgcatcg cttctatcg	3660
ccttcttgac gaggttctct gagcgggact ctggggttcg aaatgaccga ccaagcgacg	3720
cccaacctgc catcacgaga ttctgattcc accgccgcct tctatgaaag gttgggcttc	3780
ggaatcgttt tccggagcgc cggctggatg atcctccagc gcggggatct catgctggag	3840
ttcttcgccc accctagggg gaggctaact gaaacacgga aggagacaat accggaagga	3900
acccgcgcta tgacggcaat aaaaagacag aataaaacgc acggtgttgg gtcgtttgtt	3960
cataaacgcg gggttcggtc ccagggttgg cactctgtcg atacccacc gagacccat	4020
tggggccaat acgcccgct ttcttcttt tccccacccc accccccaag ttcgggtgaa	4080
ggcccagggc tcgcagccaa cgtcggggcg gcaggccctg ccatagcctc aggttactca	4140

## -continued

tatatacttt agattgattt aaaacttcat ttttaattta aaaggatcta ggtgaagatc	4200
ctttttgata atctcatgac caaaatccct taacgtgagt tttcgttcca ctgagcgtca	4260
gaccccgtag aaaagatcaa aggatcttct tgagatcctt tttttctgcg cgtaatctgc	4320
tgcttgcaaa caaaaaaacc accgctacca gcggtggttt gtttgccgga tcaagagcta	4380
ccaactcttt ttccgaaggt aactggcttc agcagagcgc agataccaaa tactgtcctt	4440
ctagtgtagc cgtagttagg ccaccacttc aagaactctg tagcaccgcc tacatacctc	4500
gctctgctaa tcctgttacc agtggtgct gccagtgccg ataagtcgtg tcttaccggg	4560
ttggactcaa gacgatagtt accggataag gcgcagcggc cggtgctgaac ggggggttcg	4620
tgacacacgc ccagcttgga gcgaacgacc tacaccgaac tgagatacct acagcgtgag	4680
ctatgagaaa gcgccacgct tcccgaaggg agaaaggcgg acaggtatcc ggtaagcggc	4740
agggtcggaa caggagagcg cacgagggag cttccagggg gaaacgcctg gtatctttat	4800
agtctgtcgc ggtttcgcca cctctgactt gagcgtcgat ttttgtgatg ctgcgtcagg	4860
ggggcgagcc tatggaaaaa cgccagcaac gcggcctttt tacggttcct ggcttttgc	4920
tggccttttg ctacatggtt ctttcctgcg ttatcccctg attctgtgga taaccgtatt	4980
accgccatgc at	4992

## 1. A composition comprising:

a plurality of particles comprising a surfactant having an HLB value of less than about 6.0 units associated with a functional composition and a polymer, with the particles having an average diameter of less than about 100 nanometers as measured by atomic force microscopy of the particles following drying of the particles, wherein the functional composition is a member of the group consisting of a bioactive component and a diagnostic agent.

2. The composition of claim 1 wherein the surfactant is a non-ionic surfactant.

3. The composition of claim 1 wherein the surfactant has an HLB value of less than about 5.0 units.

4. The composition of claim 3 wherein the average diameter of the particles is less than about 50 nm.

5. The composition of claim 1 wherein the functional composition is a member of the group consisting of peptides, proteins, and carbohydrates.

6. The composition of claim 5 wherein the average diameter is less than about 50 nm.

7. The composition of claim 1 wherein the functional composition is cisplatin.

8. The composition of claim 1 wherein the functional composition is inorganic.

9. The composition of claim 1 wherein the functional composition is a diagnostic agent that is a fluorescent molecule.

10. The composition of claim 1 wherein the functional composition is a bioactive component that comprises a hydrophilic component.

11. The composition of claim 10 wherein the bioactive component is condensed.

12. The composition of claim 1 wherein the functional composition is a bioactive component that is a member of

the group consisting of aptamers, mini-chromosomes, steroids, adrenergic, adrenocortical steroid, adrenocortical suppressant, aldosterone antagonist, and anabolic agents; analgesic, anesthetic, anorectic, anti-acne agents; anti-adrenergic, anti-allergic, anti-amebic, anti-anemic, and anti-anginal agents; anti-arthritis, anti-asthmatic, anti-atherosclerotic, antibacterial, and anticholinergic agents; anticoagulant, anticonvulsant, antidepressant, antidiabetic, and antidiarrheal agents; antidiuretic, anti-emetic, anti-epileptic, antifibrinolytic, and antifungal agent; antihemorrhagic, inflammatory, antimicrobial, antimigraine, and antimiotic agents; antimycotic, antinauseant, antineoplastic, antineutropenic, and antiparasitic agents; antiproliferative, antipsychotic, antirheumatic, antiseborrheic, and antisecretory agents; antipasmodic, antihrombotic, anti-ulcerative, antiviral and appetite suppressant agents.

13. The composition of claim 1 wherein the functional composition is a bioactive component that is a member of the group consisting of blood glucose regulator, bone resorption inhibitor, bronchodilator, cardiovascular, and cholinergic agents; fluorescent, free oxygen radical scavenger, gastrointestinal motility effector, glucocorticoid, and hair growth stimulant agent; hemostatic, histamine H<sub>2</sub> receptor antagonists; hormone; hypocholesterolemic, and hypoglycemic agents; hypolipidemic, hypotensive, and imaging agents, immunizing and agonist agents; mood regulators, mucolytic, mydriatic, nasal decongestant; neuromuscular blocking agents; neuroprotective, NMDA antagonist, non-hormonal sterol derivative, plasminogen activator, and platelet activating factor antagonist agent.

14. The composition of claim 1 wherein the functional composition is a bioactive component that is a member of the group consisting of platelet aggregation inhibitor, psychotropic, radioactive, scabicide, and sclerosing agents; sedative, sedative-hypnotic, selective adenosine A1 antagonist, serotonin antagonist, and serotonin inhibitor agent;

serotonin receptor antagonist, steroid, thyroid hormone, thyroid hormone, thyroid inhibitor agent; thyromimetic, tranquilizer, amyotrophic lateral sclerosis, cerebral ischemia, Paget's disease agent; unstable angina, vasoconstrictor, vasodilator, wound healing, and xanthine oxidase inhibitor agent; and immunological agents.

15. The composition of claim 1 wherein the functional composition is a bioactive component that comprises a member of the group consisting of antigens isolated from pathogens, viral antigens, fungal antigens, parasitic antigens, and inactivated pathogenic organisms.

16. The composition of claim 1 wherein the surfactant has a critical micelle concentration of less than about 200 micromolar.

17. The composition of claim 1 further comprising a biocompatible oil.

18. The composition of claim 1 wherein the functional composition is a bioactive component that comprises a polynucleic acid, oligonucleotide, antisense molecule, or a polypeptide.

19. The composition of claim 18 wherein the bioactive component is condensed.

20. The composition of claim 19 wherein the composition further comprises a water-miscible solvent.

21. The composition of claim 1 wherein the surfactant is selected from the group consisting of 2, 4, 7, 9-tetramethyl-5-decyn-4, 7-diol, molecules containing an acetylenic diol portion, and blends of 2, 4, 7, 9-tetramethyl-5-decyn-4, 7-diol.

22. The composition of claim 1 wherein the particles further comprise a cell recognition component.

23. The composition of claim 22 wherein the cell recognition component is a ligand, peptide hormone, or an antibody.

24. The composition of claim 23 wherein the cell recognition component comprises tenascin, hyaluronan, or polyvinylpyrrolidone, or a fragment thereof.

25. The composition of claim 24 wherein the average diameter of the particles is less than about 50 nm.

26. The composition of claim 24 wherein the cell recognition component comprises tenascin or a fragment thereof.

27. The composition of claim 1 wherein the average diameter of the particles is less than about 50 nm.

28. The composition of claim 1 wherein the polymer is an iontophoretic polymer.

29. The composition of claim 1 wherein the polymer is a hydrophobic polymer.

30. The composition of claim 1 wherein the polymer is a hydrophilic polymer.

31. The composition of claim 1 wherein the polymer is chosen from the group consisting of polyamides, polycarbonates, polyalkylenes, polyalkylene glycols, polyalkylene oxides, polyalkylene terephthalates, polyvinyl alcohols, polyvinyl ethers, polyvinyl esters, polyvinyl halides, polyvinylpyrrolidone, polyglycolides, polysiloxanes, polyurethanes and copolymers thereof, alkyl cellulose, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, polymers of acrylic and methacrylic esters, methyl cellulose, ethyl cellulose, hydroxypropyl cellulose, hydroxy-propyl methyl cellulose, hydroxybutyl methyl cellulose, cellulose acetate, cellulose propionate, cellulose acetate butyrate, cellulose acetate phthalate, carboxylethyl cellulose, cellulose triacetate, and cellulose sulphate sodium salt.

32. The composition of claim 1 wherein the polymer is chosen from the group consisting of poly(methyl methacrylate), poly(ethylmethacrylate), poly(butylmethacrylate), poly(isobutylmethacrylate), poly(hexylmethacrylate), poly(isodecylmethacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), poly(ethylene glycol), poly(ethylene oxide), and poly(ethylene terephthalate).

33. The composition of claim 1 wherein the polymer is chosen from the group consisting of poly(vinyl alcohols), poly(vinyl acetate), poly(vinyl chloride polystyrene), polyvinylpyrrolidone, polyhyaluronic acids, casein, gelatin, gluten, polyanhydrides, polyacrylic acid, alginate, chitosan, poly(methyl methacrylates), poly(ethyl methacrylates), poly(butylmethacrylate), poly(isobutylmethacrylate), poly(hexylmethacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), and poly(octadecyl acrylate).

34. The composition of claim 1 wherein the hydrophilic polymer is a member of the group consisting of proteinaceous materials, peptides, carbohydrates.

35. A method of delivering a functional composition across keratinized barrier epithelia to a cell, the method comprising introducing the composition of claim 1 at a position that is separated from the cell by a keratinized barrier epithelium, wherein at least a portion of the plurality of particles passes through the keratinized barrier epithelium to the cell.

36. The method of claim 35 wherein the functional composition is delivered transcutaneously.

37. The method of claim 35 wherein the composition of claim 1 is prepared as a medicament, and the medicament is administered to a patient.

38. A solution comprising the composition of claim 1, the solution comprising a concentration of cations between 20 and 2000 millimolar.

39. The composition of claim 1, further comprising a cation chosen from the group consisting of  $\text{Ni}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Al}^{3+}$ ,  $\text{Be}^{2+}$ ,  $\text{Li}^+$ ,  $\text{Ba}^{2+}$ , and  $\text{Gd}^{3+}$ .

40. A medicament comprising the composition of claim 1.

41. The medicament of claim 39 further comprising a form selected from the group consisting of granules, tablets, pellets, films, oral, intravenous, subcutaneous, intraperitoneal, intrathecal, intramuscular, inhalation, topical, transdermal, suppository, pessary, intra urethral, intraportal, intraocular, transtympanic, intrahepatic, intra-arterial, intrathecal, transmucosal, coatings, buccal, and combinations thereof.

42. A method of delivering a functional composition to a patient, the method comprising administering a medicament to the patient that comprises the composition of claim 1.

43. A cell comprising the composition of claim 1, wherein the plurality of particles is associated with the cell.

44. A cell comprising the composition of claim 18, wherein the plurality of particles is associated with the cell.

45. A method of transfecting a cell, the method comprising exposing the cell to the composition of claim 18.

46. A matrix for binding bioactive or diagnostic particles, the matrix comprising the particles of claim 1 and a binder.

47. A method of delivering a medical agent to a cell having caveolae, the method comprising: associating the medical agent with an organic functional composition in

vitro to make an association of the medical agent and the organic functional composition, wherein the association is passable through cellular caveolae for delivery of the medical agent.

**48.** The method of claim 47 wherein the association of the agent and the functional composition comprises a particle.

**49.** The method of claim 48 wherein the particle has a diameter of less than about 100 nanometers as measured by atomic force microscopy of the particles following drying of the particles.

**50.** The method of claim 48 wherein the particle has a diameter of less than about 50 nm.

**51.** The method of claim 48 wherein the particle further comprises a surfactant having an HLB value of less than about 6.0 units.

**52.** The method of claim 51 wherein the particle has an average diameter of less than about 50 nanometers as measured by atomic force microscopy of the particles following drying of the particles.

**53.** The method of claim 51 further comprising exposing the particle to the cell.

**54.** The method of claim 47 wherein the association of the agent and the functional composition has a maximum dimension of no more than about 50 nm nanometers as measured by atomic force microscopy following drying of the association of the agent and the functional composition.

**55.** The method of claim 47 wherein the functional composition comprises a surfactant.

**56.** The method of claim 47 further comprising exposing the association of the agent and the functional composition to the cell.

**57.** The method of claim 47 further comprising administering a medicament to a patient, the medicament comprising the association of the medical agent and the organic functional composition.

**58.** The method of claim 47 wherein the functional composition comprises a surfactant and a hydrophilic polymer.

**59.** The method of claim 47 wherein the agent comprises a bioactive component that a member of the group consisting of peptides, proteins, and carbohydrates.

**60.** The method of claim 47 wherein the agent comprises a member of the group consisting of a bioactive component and a diagnostic agent.

**61.** The method of claim 47 wherein the agent comprises a fragment of a nucleic acid that comprises a nucleic acid sequence.

**62.** The method of claim 61 wherein the functional composition comprises a surfactant.

**63.** The method of claim 47 wherein the association is introduced at a position that is separated from the cell by keratinized barrier epithelia, and the association passes through the keratinized barrier epithelia to the cell.

**64.** The method of claim 63 further comprising exposing the cell to the association of the agent and the functional composition

**65.** The method of claim 47 wherein the functional composition comprises carbon and hydrogen.

\* \* \* \* \*