



(51) International Patent Classification:

A61K 31/437 (2006.01) C07D 471/04 (2006.01)
A61P 35/02 (2006.01)

(21) International Application Number:

PCT/US2018/044862

(22) International Filing Date:

01 August 2018 (01.08.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/540,051 01 August 2017 (01.08.2017) US
62/617,180 13 January 2018 (13.01.2018) US

(71) Applicant: PTC THERAPEUTICS, INC. [US/US]; 100 Corporate Court, Middlesex Business Center, South Plainfield, NJ 07080 (US).

(72) Inventors: CAO, Liangxian; 17 Winesap Drive, East Brunswick, NJ 08816 (US). WEETALL, Maria; 70 Burnham Parkway, Morristown, NJ 07960 (US).

(74) Agent: SCOLA, Daniel, A., Jr. et al.; Hoffmann & Baron, LLP, 6 Campus Drive, Parsippany, NJ 07054 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,


CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— with international search report (Art. 21(3))

(54) Title: DHODH INHIBITOR FOR USE IN TREATING HEMATOLOGIC CANCERS



(57) Abstract: A method of use for Cpd 1 as an inhibitor of dihydroorotate dehydrogenase (DHODH) function in treating or ameliorating a hematological cancer in a subject in need thereof comprising, administering an effective amount of Cpd 1 to the subject, having structure (I); or a form or pharmaceutical composition thereof.



DHODH INHIBITOR FOR USE IN TREATING HEMATOLOGIC CANCERS

REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Patent application 62/540,051, filed August 1, 2017; and also claims priority to United States Provisional Patent application 62/617,180, filed January 13, 2018, the entire contents of each of which are incorporated herein in their entirety and for all purposes.

INTRODUCTION

Cancer cells drive uncontrolled proliferation by regulating and integrating multiple biosynthetic processes, consequently affecting nucleotide and protein expression [16]. One of the affected biosynthetic processes is the expression of dihydroorotate dehydrogenase (DHODH), a rate-limiting enzyme whose function regulating de novo biosynthesis of pyrimidine nucleotides [1] is affected during uncontrolled cell cycle division. Consistent with such an effect, small molecule inhibition of DHODH function has recently been shown to prevent growth of several hematological cancers [2-4] where the hematological cancer is characterized by reduced expression of uridine salvage enzymes and a higher dependence on de novo pyrimidine nucleotide synthesis.


Another affected biosynthetic process in certain hematological cancers (~13%) is the dysregulated expression of a mutant isocitrate dehydrogenase (mIDH) protein. Some inhibitors functioning as antiproliferative agents demonstrate selectivity for mutant IDH1 or mutant IDH2 over wildtype enzyme [29, 30], while other small molecules such as enasidenib have recently been shown to selectively inhibit mutant IDH2 enzyme activity [31].

Accordingly, there remains a great need for useful chemotherapeutic agents having potent inhibition of DHODH functional activity, favorable pharmaceutical properties and extensive clinical experience for use in treating such hematological cancers. In other instances, useful combinations of chemotherapeutic agents having synergistic antiproliferative activity, inhibiting both DHODH functional activity and IDH enzyme

activity are desired for treating hematological cancers in subjects having high unmet medical need.

SUMMARY

4-chlorophenyl (*S*)-6-chloro-1-(4-methoxyphenyl)-1,3,4,9-tetrahydro-2H-pyrido[3,4-*b*]indole-2-carboxylate is a pharmacologically active enantiomer (hereinafter referred to as “Cpd 1” or “Cpd 1 *S*-enantiomer”), having the structure:



Cpd 1 has demonstrated potent activity as an inhibitor of dihydroorotate dehydrogenase (DHODH) function, showing broad and potent activity against hematological cancers in both in vitro and in vivo preclinical and clinical studies. Moreover, a combination therapy of an inhibitor of mutant isocitrate dehydrogenase (mIDH) function with Cpd 1, an inhibitor of DHODH function, has recently been discovered as a potentiator of the antiproliferative effect of both mutant and wildtype IDH1 and IDH2 cells.

Provided herein is a method of use for Cpd 1 or a form or pharmaceutical composition thereof as an inhibitor of dihydroorotate dehydrogenase (DHODH) function in treating or ameliorating a hematological cancer in a subject in need thereof comprising, administering to the subject an effective amount of Cpd 1. More particularly, the hematological cancer is a leukemia or myelodysplastic syndrome (MDS) characterized by reduced expression of uridine salvage enzymes and dependence on de novo pyrimidine nucleotide synthesis. Also provided herein is a method of use for Cpd 1 or a form or pharmaceutical composition thereof in combination with other chemotherapeutic agents having additive and synergistic activity for inhibiting functional activity of both DHODH and IDH.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1a shows Cpd 1 inhibition of total intracellular pyrimidine nucleotide production in HT1080 cells; Figure 1b shows the results of ¹⁵N-glutamine labeling studies comparing inhibition of synthesis of de novo pyrimidine nucleotide production (UMP, CTP and UTP) in HT1080 cells treated with the Cpd 1 active *S*-enantiomer and Cpd 1 (*R*), the inactive *R*-enantiomer, relative to vehicle control; Figure 1c shows that Cpd 1 acts upstream of UMPS, blocking UMPS inhibitor pyrazofurin induced increase of orotic acid in HT1080 cells; Figure 1d shows Cpd 1 inhibited production of VEGFA protein (a DHODH surrogate for inhibition of DHODH function) in HT1080 cells was also blocked by addition of exogenous uridine rather than other nucleosides, including adenosine, cytidine and guanosine; Figure 1e shows that S-phase cell cycle arrest induced by Cpd 1 was blocked by addition of exogenous uridine or cytidine.

Figure 2a shows free Cpd 1 and brequinar selectively “Competing On” with DHODH binding to the Cpd 1-beads by Western blot; Figure 2b shows elution of DHODH from selective “Competing Off” by free Cpd 1 and brequinar; Figure 2c compares the effect of Cpd 1 on the activity of isolated mitochondrial DHODH with the known DHODH inhibitors brequinar and teriflunomide using Cpd 1 (*R*), the inactive *R*-enantiomer as a negative control.

Figure 3a shows serum levels of DHO correlated with Cpd 1 concentration in a NF2 clinical study subject; Figure 3b shows Cpd 1 increased serum DHO in three additional NF2 subjects.

Figure 4a compares Cpd 1 overall activity in hematopoietic cell lines with solid cancer cell lines relative to Figure 5a; Figure 4b provides box-plot microarray log₂ expression levels for CDA and UPP1, two pyrimidine salvage enzymes for producing uridine from cytidine and uracil, respectively; Figure 4c compares relative UMP production from the salvage pathway in Cpd 1 sensitive and insensitive cell lines; Figure 4d compares other known DHODH inhibitors with Cpd 1 for inhibiting proliferation of MOLM-13 AML cells; Figure 4e shows inhibition of in vitro proliferation in various subject derived AML cell lines treated with Cpd 1.

Figure 5a shows sensitivity to Cpd 1 for a panel of 240 cell lines having various tissue origins; Figure 5b shows effect of Cpd 1 treatment on the number of human leukemia cells in blood from mice in a systemic leukemia lethality model; Figure 5c shows survival of mice in the systemic leukemia lethality model.

Figure 6 shows effect on tumor growth progress in a MOLM-13 acute myeloid leukemia (AML) model after treatment with vehicle, Cpd 1, cytarabine (AraC), or a combination of Cpd 1 and AraC over the study period.

Figure 7 compares bioavailability of Cpd 1 and GSK983 orally administered to mice.

Figure 8 MOLM-13 AML cells were treated with Cpd 1 (100 nM) or DMSO control for 72 hours, then stained with anti-CD14 and CD11b antibodies.

Figure 9 compares the percent reduction in proliferation by treatment with the combination of Cpd 1 (inhibitor of DHODH function) and enasidenib (mutant IDH2 inhibitor) alone or in combination in HL-60 cells (human promyelocytic leukemia cells) treated for 72 hours.


Figure 10a compares the percent reduction in proliferation of total blasts and immature blasts with the combination of Cpd 1 (inhibitor of DHODH function) and enasidenib (mutant IDH2 inhibitor) alone or in combination in assays utilizing blood samples obtained from AML subjects and treated ex vivo for 72 hours. Enasidenib enhanced the activity of Cpd 1 in both mutant and wildtype IDH1 and IDH2 cells.

Figure 10b compares the percent reduction in proliferation of total blasts and immature blasts with the combination of Cpd 1 (inhibitor of DHODH function) and enasidenib (mutant IDH2 inhibitor) alone or in combination in assays utilizing blood samples obtained from ALL subjects and treated ex vivo for 72 hours. Enasidenib enhanced the activity of Cpd 1 in both mutant and wildtype IDH1 and IDH2 cells.

Figure 11 compares the mean tumor volume in IDH2 wild-type cells after treatment with Cpd 1 alone, Agios Compound AG-221 alone and in combination. The use of the combination of a Cpd 1 DHODH functional inhibitor and a AG-221 IDH2 functional inhibitor demonstrates synergistic activity.

DETAILED DESCRIPTION

Provided herein is a method of use for Cpd 1 or a form or pharmaceutical composition thereof as an inhibitor of dihydroorotate dehydrogenase (DHODH) function in treating or ameliorating a hematological cancer in a subject in need thereof comprising, administering an effective amount of Cpd 1 to the subject, having the structure:



or a form or pharmaceutical composition thereof.

Comparative Gene Expression Analysis

The identification of Cpd 1 as an inhibitor of DHODH function for selective use in treating or ameliorating a hematological cancer was obtained from a gene expression analysis comparing Cpd 1-resistant HT1080 cells with Cpd 1-sensitive HT1080 cells, as shown in Table 1.

Table 1

Gene	Genbank#	Expression	Fold Change
STOM	M81635	Ubiquitous membrane protein deficient in high Na ⁺ , low K ⁺ stomatocytic erythrocytes	12.30
CDA	NM_001785	Cytidine deaminase	11.15
F8A	NM_012151	Coagulation factor VIII-associated (intronic transcript)	6.35
STOM	AI537887	tp32g06.x1 NCI_CGAP_Ut4 homo sapiens cDNA clone image: 2189530 (having a 3' amino acid terminus similar to the human gb:m81635 erythrocyte band 7 integral membrane protein human mRNA sequence)	6.24
KLHL1	AF126749	SCA8 mRNA, repeat region	5.13

Gene	Genbank#	Expression	Fold Change
EIF1AY	BC005248	Eukaryotic translation initiation factor 1A, Y-linked	0.07
LOC81569	BF594459	Actin like protein	0.12
SORT1	BE622952	Sortilin 1	0.16
FLJ21986	BF724137	Hypothetical protein FLJ21986	0.16
MTMR1	AI167164	Myotubularin related protein 1	0.17

The gene expression analysis was performed with a cDNA microarray (provided by Affymatrix) using Cpd 1-resistant cells generated by culturing HT1080 cells in escalating concentrations of Cpd 1, then doubling Cpd 1 concentration weekly from 1 nM to 1000 nM and isolating total mRNA from the Cpd 1-resistant and wild-type HT1080 cells. Table 1 lists the genes with the greatest increase or decrease in expression in the Cpd 1-resistant HT1080 cell line, when compared to the parent HT1080 cells.

For example, cytidine deaminase (CDA) expression was increased 11-fold in Cpd 1-resistant HT1080 cells when compared with levels in the wild type HT1080 cells.

¹⁵N-Glutamine Tracing Study

Since CDA is an enzyme in the pyrimidine nucleotide salvage pathway [5], the effect of Cpd 1 treatment on intracellular nucleotide levels was also evaluated using an ¹⁵N-glutamine tracing study. The study results demonstrate that, compared to ¹⁵N-glutamine (1 μ M), Cpd 1 selectively decreased the total levels of deoxy-pyrimidine nucleotides (dCTP and dTTP) but not deoxy-purine nucleotides dATP and dGTP in HT1080 cells treated for 8 hours with Cpd 1 (100 nM) relative to vehicle control (0.5% DMSO), as determined by LC/MS (Figure 1a). Therefore, Cpd 1 inhibits de novo pyrimidine synthesis, while its inactive *R*-enantiomer had no effect as determined by LCMS against ¹⁵N-labelled CTP, UTP and UMP (Figure 1b). Further studies demonstrated that inhibition was dose-dependent and was measurable after only 30 minutes of treatment (data not shown).

Table 2 shows metabolism of ¹⁵N-glutamine in cells treated with Cpd 1 enantiomers (100 nm), brequinar (100 nm) (a known DHODH inhibitor) and pyrazofurin-

monophosphate (MP) (100 nm) (a known UMPS inhibitor) or DMSO control for 8 hours in the presence of ^{15}N -glutamine. The resulting metabolic activity is shown (μM), where BLDL indicates activity Below Detection Limits. The metabolism of ^{15}N -glutamine in cells treated with Cpd 1 was similar to that in cells treated with brequinar [6], in contrast to the UMP synthase (UMPS) inhibitor pyrazofurin-monophosphate [7], which increased ^{15}N -orotate levels by almost 160 fold.

Table 2

Treatment	^{15}N -DHO (μM)	^{15}N -Orotate (μM)	^{15}N -UMP (μM)
0.5% DMSO	BLDL	6.7 ± 0.3	1.4 ± 0.4
Cpd 1 (<i>R</i>)	BLDL	4.4 ± 0.4	0.7 ± 0.2
Cpd 1	18.1 ± 5.5	44.8 ± 6.1	BLDL
Brequinar	11.9 ± 3.7	50.1 ± 3.1	BLDL
Pyrazofurin-MP	1.3 ± 1.1	974 ± 175	BLDL

Comparative Inhibition of De Novo Pyrimidine Nucleotide Synthesis

In another study, treatment with pyrazofurin-MP resulted in a greater than 40-fold increase of ^{15}N -orotate, the substrate of UMPS, compared with vehicle (0.5% DMSO) (Figure 1c). However, co-treatment with Cpd 1 blocked the pyrazofurin-MP-induced increase of ^{15}N -orotate, consistent with inhibition of DHODH function, indicating that Cpd 1 acts upstream of UMPS. Having thus shown that Cpd 1 inhibits de novo pyrimidine nucleotide synthesis, subsequent nucleoside rescue studies using inhibition of VEGF protein production as a surrogate for inhibition of DHODH function demonstrated that the addition of uridine to the culture medium blocked Cpd 1 inhibition of VEGFA protein production (Figure 1d). In contrast, adenosine, cytidine and guanosine did not have a similar effect. Moreover, Cpd 1 induced S phase cell cycle delay in HT1080 cells was prevented by the addition of the pyrimidine nucleosides uridine or cytidine (Figure 1e), whereas addition of the purine nucleosides adenosine or guanosine had no effect (data not shown).

Comparative DHODH Binding

The binding partner of Cpd 1 was identified by pull down studies comparing the active Cpd 1 *S*-enantiomer and the inactive *R*-enantiomer linked to epoxy-activated Sepharose beads. Western blot analysis demonstrated that binding of DHODH to the Cpd 1-linked beads was inhibited by preincubation with either free Cpd 1 or brequinar (Figure 2a). In contrast, binding to prohibitin, another mitochondrial protein, was not blocked by either Cpd 1 or brequinar. Subsequent competitive elution studies demonstrated that DHODH was selectively eluted from the Cpd 1 beads by the free Cpd 1 *S*-enantiomer or brequinar, but not by the inactive *R*-enantiomer or the DMSO control (Figure 2b). These findings demonstrate that Cpd 1 specifically binds DHODH. Computer modelling using the available crystal structure of the human DHODH protein [8] shows that Cpd 1 and brequinar analog C44 bind to the same pocket with similar docking scores [9,10] of 9.55 for C44 compared to 9.27 for Cpd 1, consistent with pull down studies demonstrating DHODH binding competition between brequinar and Cpd 1. When purified mitochondria were used as the source of DHODH, treatment with Cpd 1 inhibited DHODH activity more potently than did teriflunomide and with similar potency as brequinar (orotate production was measured by LC-MS/MS after 30 minutes of treatment, Figure 2c). These data indicate that Cpd 1 inhibits de novo pyrimidine nucleotide synthesis via directly targeting DHODH.

Tumor Cell Line Panel Analysis

Since pyrimidine nucleotides can be synthesized by either or both the de novo or the salvage pathway [11], to determine whether some cancer cell lines were insensitive to Cpd 1 due to increased pyrimidine nucleotide salvage activity, thus affecting the selective use of Cpd 1 to certain cancer types, the IC₅₀ (concentration that inhibits cell proliferation by 50%) of Cpd 1 was determined against a panel of 240 tumor cell lines. After a 3-day incubation period, the results shown in Figure 5a indicate that Cpd 1 is an active inhibitor of DHODH function in certain hematologic cancers; where, ALL: acute lymphoblastic leukemia; AML: acute myeloid leukemia; ANLL: acute nonlymphocytic leukemia; CLL: chronic lymphocytic leukemia; CML: chronic myeloid leukemia; H&N: head and neck, HCC: hepatocellular carcinoma; NSCLC: non-small cell lung cancer; SCLC: small cell lung cancer. In Figure 5a, tumor cells having an IC₅₀ ≤ 2 μM were

classified as a sensitive responder (68 lines) compared with those having an $IC_{50} > 2 \mu M$, considered a resistant non-responder (172 lines). When grouped by hematopoietic cancer vs solid tumor, 57% (32/56) of hematopoietic lines were sensitive to Cpd 1, whereas only 18% (34/184) of the solid tumor lines were sensitive to Cpd 1 (Figure 4a).

Gene expression of pyrimidine nucleotide synthesis enzymes in these 240 lines was also analyzed using published microarray data in the Cancer Cell Line Encyclopedia (CCLE) database [12]. Of the 240 cell lines, 232 cell lines were found in the CCLE database, including 54 hematopoietic and 178 solid tumor cell lines.

Table 3 compares average expression of certain genes: dihydroorotate dehydrogenase (DHODH), uridine monophosphate synthetase (UMPS), uridine phosphorylase 1 (UPP1), cytidine deaminase (CDA), uridine-cytidine kinase 1 (UCK1), uridine -cytidine kinase 1-like 1 (UCKL1), uridine phosphorylase 2 (UPP2), uridine-cytidine kinase 2 (UCK2), cytosolic cytidine monophosphate (UMP-CMP) kinase 1 (CMPK1), providing results for average expression in hematopoietic cancer lines ($Hemat^{Avg}$), solid tumor lines ($Solid^{Avg}$), and standard deviation in hematopoietic cancer lines ($Hemat^{SD}$), solid tumor lines ($Solid^{SD}$) and differential expression in hematopoietic tumor vs. solid tumor ($\Delta H-S$) with *t*-test p-values comparing hematopoietic cancer ($n = 54$) with solid tumors ($n = 178$).

As shown in Table 3, two salvage enzymes, CDA and UPP1 for converting cytidine and uracil to uridine, respectively, had lower expression (3.2-fold for CDA and 4.6-fold for UPP1) (where fold reduction is based on the $\log_2$ value of $\Delta H-S$) in the hematopoietic cancer lines than in the solid tumor lines (see also Figure 4b).

Table 3

Gene	$Hemat^{Avg}$	$Solid^{Avg}$	$Hemat^{SD}$	$Solid^{SD}$	$\Delta H-S$	p-value
DHODH ¹	6.5	6.1	0.5	0.7	0.5	1
UPMS ¹	6.6	6.5	0.3	0.4	0.0	1
UPP1 ²	6.3	8.5	1.8	2.0	- 2.2	8.9 E-12
CDA ²	4.5	6.2	1.0	2.4	- 1.7	2.9 E-12
UCK1 ²	6.8	7.0	0.4	0.5	- 0.2	0.0006
UCKL1 ²	7.7	7.7	0.4	0.6	0.0	0.39
UPP2 ²	3.8	3.7	0.2	0.1	0.1	1
UCK2 ²	9.0	8.8	0.6	0.6	0.3	1

CMPK1²	10.9	10.4	0.5	0.6	0.6	1
--------------------------	------	------	-----	-----	-----	---

Salvage and De Novo Pathway Panel

LC-MS/MS analysis of UMP pools from salvage and de novo pathways in a panel of tumor cell lines was performed, where four sensitive cell lines (HeLa, HT1080, A549 and K562) and four insensitive lines (U87MG, MCF7, PC3 and Huh7) were cultured in the presence of ¹⁵N-glutamine for 8 hours and unlabeled and ¹⁵N-labeled UMP in the cell lysates were measured by LC-MS/MS. In general, Cpd 1 sensitive cells have less salvage UMP production relative to de novo UMP production when compared to the insensitive tumor cells (Figure 4c). These cumulative data suggest that a hematologic cancer having lower uridine salvage activity enhances the therapeutic effect of Cpd 1.

DHODH Inhibitor Leukemia Cell Line Comparison

After confirming in vivo DHODH inhibition activity in solid tumor clinical studies and in vitro activity in leukemia cell lines, the potency of Cpd 1 was also compared with that of several well-known DHODH inhibitors [6;13;14] in leukemia cells. Cpd 1 was the most potent inhibitor with an IC₅₀ of about 1 nM, having 10-1000-fold greater potency than well-known inhibitors of DHODH function, such as brequinar, vidofludimus or teriflunomide (Figure 4d). Acute myeloid leukemia (AML) subject-derived cells were treated for 72 hours with Cpd 1, teriflunomide and brequinar, where proliferation was measured with Celltiter Glo (Promega), demonstrating inhibition of proliferation in all five AML subject derived cell lines tested (Figure 4e). Unlike teriflunomide and brequinar, Cpd 1 did not show myeloid suppression in subjects with solid tumors [24].

MOLT-4 Human Acute Lymphoblastic Leukemia Lethality Model

After confirming in vitro activity in leukemia cells, the efficacy of Cpd 1 in two mouse models of leukemia was assessed. In the MOLT-4 human acute lymphoblastic leukemia (ALL) lethality model, NOD/SCID mice were injected intravenously (IV) with MOLT-4 cells (CD45+) then treated with Cpd 1 (10 mg/kg, qd) or doxorubicin (0.5 mg/kg or 1 mg/kg IP once per week) initiated 7 days after tumor inoculation. Cpd 1 treatment significantly reduced the number of circulating MOLT-4 cells in mice compared to doxorubicin (Figure 5b). Doxorubicin was tested at the highest dose tolerated by NOD/SCID mice. Vehicle-dosed mice had a median survival time (MST) of 46 days and

did not survive beyond 71 days. Treatment with doxorubicin did not prolong survival time (Figure 6a). In contrast, the MST for the Cpd 1 treated mice was 136 days ($p < 0.05$, one way ANOVA, multiple comparisons vs vehicle).

MOLM-13 Human Acute Myeloid Leukemia Model

In another xenograft study, MOLM-13 AML cells were injected into the flank of nude mice to generate a solid tumor then mice were treated with vehicle, Cpd 1, cytarabine (AraC), a deoxycytidine analog used as standard chemotherapy to treat AML [15], or a combination of Cpd 1 and AraC (Figure 6). Cpd 1 treatment alone resulted in significant tumor growth delay when compared with vehicle or AraC treatment; the median time to reach a tumor volume of 1000 mm³ was 23 days for Cpd 1 vs 7 and 13 days for vehicle and AraC, respectively. The combination of Cpd 1 with AraC further delayed tumor growth for 39 days. The increased activity of the combination of AraC with Cpd 1 may reflect a more efficient incorporation of AraC into DNA due to a reduction in the levels of competing endogenous pyrimidine nucleotides.

These results suggest that Cpd 1 potently inhibits leukemia cell proliferation dependent on de novo pyrimidine nucleotide synthesis by selectively affecting DHODH function.

Definitions

As used herein, the term “about” means a range around a given value wherein the resulting value is substantially the same as the expressly recited value. In one aspect, “about” means within 25% of a given value or range. For example, the phrase “about 70% by weight” comprises at least all values from 52% to 88% by weight. In another aspect, the term “about” means within 10% of a given value or range. For example, the phrase “about 70% by weight” comprises at least all values from 63% to 77% by weight. In another aspect, the term “about” means within 7% of a given value or range. For example, the phrase “about 70% by weight” comprises at least all values from 65% to 75% by weight.

Concentrations, amounts, cell counts, percentages and other numerical values may be presented herein in a range format. It is to be understood that such range format is used merely for convenience and brevity and should be interpreted flexibly to include not only the numerical values explicitly recited as the limits of the range but also to

include all the individual numerical values or sub-ranges encompassed within that range as if each numerical value and sub-range is explicitly recited.

As used herein, the terms “therapies” and “therapy” can refer to any protocol(s), method(s), compositions, formulations, and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a condition or disorder or one or more symptoms thereof (*e.g.*, hematologic cancer or one or more symptoms or one or more conditions associated therewith).

In certain aspects, the terms “therapies” and “therapy” refer to drug therapy such as chemotherapy, adjuvant therapy, radiation, surgery, biological therapy, supportive therapy, antiviral therapy and/or other therapies useful in treatment, management, prevention, or amelioration of a condition or disorder or one or more symptoms thereof (*e.g.*, hematologic cancer or one or more symptoms or one or more conditions associated therewith). In certain aspects, the term “therapy” refers to a therapy other than Cpd 1 or a pharmaceutical composition thereof. In specific aspects, an “additional therapy” and “additional therapies” refer to a therapy other than a treatment using Cpd 1 or a pharmaceutical composition thereof. In a specific aspect, a therapy includes the use of Cpd 1 as an adjuvant therapy. For example, using Cpd 1 in conjunction with a drug therapy such as chemotherapy, biological therapy, surgery, supportive therapy, antiviral therapy and/or other therapies useful in treatment, management, prevention, or amelioration of a condition or disorder or one or more symptoms thereof (*e.g.*, hematologic cancer or one or more symptoms or one or more conditions associated therewith).

As used herein, the term “human infant” refers to a newborn to 1 year old year human.

As used herein, the term “human toddler” refers to a human that is 1 year to 3 years old.

As used herein, the term “human child” refers to a human that is 1 year to 18 years old.

As used herein, the term “human adult” refers to a human that is 18 years or older.

As used herein, the term “middle-aged human” refers to a human between the ages of 30 and 64.

As used herein, the term “elderly human” refers to a human 65 years or older.

As used herein, the term “subject” refers to an individual being administered a therapy as described herein. In a specific aspect, the individual is a human.

As used herein, the term “effective amount” in the context of administering Cpd 1 to a subject having a hematologic cancer refers to the dose of Cpd 1 that results in a beneficial or therapeutic effect. In specific aspects, an “effective amount” of Cpd 1 refers to an amount of Cpd 1 which is sufficient to achieve at least one, two, three, four or more of the following beneficial or therapeutic effects: (i) inhibition of a hematologic cancer; (ii) regression of the hematologic cancer; (iii) eradication, removal, or complete remission of the hematologic cancer; (iv) prevention of the development or onset of one or more symptoms associated with the hematologic cancer; (v) reduction or amelioration of the severity of one or more symptoms associated with the hematologic cancer; (vi) the reduction in the number of one or more symptoms associated with cancer; (vii) amelioration of the severity of one or more symptoms associated with the hematologic cancer; (viii) reduction in the duration of one or more symptoms associated with the hematologic cancer; (ix) prevention in the recurrence of proliferation or one or more symptoms associated with the hematologic cancer; (x) a reduction in mortality; (xi) an increase in survival rate of subjects; (xii) an increase in relapse free survival; (xiii) an increase in the number of hematologic cancer subjects in remission; (xiv) reduction in hospitalization of a subject; (xv) reduction in hospitalization length; (xvi) a decrease in hospitalization rate; (xvii) an increase in the survival of a subject; (xviii) an increase in symptom-free survival of cancer subjects; (xix) an increase in the length of a period of remission of a hematologic cancer in a subject; (xx) improvement in quality of life (QOL) as assessed by methods well known in the art, *e.g.*, QOL questionnaires and the like; (xxi) a reduction in proliferation before treatment with another chemotherapeutic agent; (xxii) a reduction in proliferation before treatment with radiation; (xxiii) a reduction in proliferation before treatment with surgery; (xxiv) enhancement of or improvement of the therapeutic effect of another therapy; (xxv) an additive antiproliferative effect in combination with another therapy; (xxvi) a synergistic antiproliferative effect in combination with another therapy; (xxvii) a decrease in the concentration of plasma DHODH of a subject having a hematologic cancer; (xxviii) a decrease in circulating proliferative cells in the plasma of a subject having a

hematologic cancer; (xxix) an alteration (*e.g.*, a decrease) in a biomarker for a hematologic cancer; (xxx) reduction of the concentration of DHODH in a biological specimen (*e.g.*, plasma, serum, cerebral spinal fluid, urine, or any other biofluids) from a subject having a hematologic cancer; (xxxi) a decrease in the concentration of DHODH in a biological specimen (*e.g.*, the plasma, serum, urine or cerebrospinal fluid (CSF)) from a subject having a hematologic cancer; (xxxii) reduction of the concentration of one or more biomarkers in a biological specimen (*e.g.*, plasma, serum, cerebral spinal fluid, urine, or any other biofluids) from a subject having a hematologic cancer; (xxxiii) proliferative cell count is maintained after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan; (xxxiv) white cell count is maintained after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan; (xxxv) proliferative cell count does not increase or increases by less than expected after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan; (xxxvi) white cell count does not increase or increases by less than expected after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan; (xxxvii) proliferative cell count is decreased after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan; or, (xxxviii) white cell count is decreased after administration of a therapy as described herein as measured by conventional methods available to one skilled in the art, such as magnetic

resonance imaging (MRI), dynamic contrast-enhanced MRI (DCE-MRI), X-ray, computed tomography (CT) scan, or a positron emission tomography scan.

As used herein, the term “in a 24 hour period” refers to a period of time over which a condition is maintained; for example, the effective amount of Cpd 1 is identified when the mean plasma concentration of Cpd 1 is achieved and maintained for a plurality of 24 hour periods. In other words, the mean plasma concentration of Cpd 1 may be reached in a suitable time, which may be more or less than 24 hours.

As used herein, the term “a therapy as described herein” refers to a method of use for Cpd 1 or a form or pharmaceutical composition thereof as an inhibitor of DHODH function in treating or ameliorating a hematological cancer in a subject in need thereof comprising, administering to the subject an effective amount of Cpd 1. In one aspect, the hematological cancer is a leukemia. In another aspect, the leukemia is an acute or chronic form of leukemia. In another aspect, the leukemia is characterized by reduced expression of uridine salvage enzymes and dependence on de novo pyrimidine nucleotide synthesis. In another aspect of the therapy described herein, the method of use for Cpd 1 or a form or pharmaceutical composition comprises a combination with other chemotherapeutic agents having synergistic antiproliferative activity. In one aspect, the other chemotherapeutic agent inhibits DHODH functional activity. In another aspect, the other chemotherapeutic agent inhibits IDH expression.

As used herein, the term “pharmaceutically acceptable salt(s)” refers to a salt prepared from a pharmaceutically acceptable non-toxic acid or base including an inorganic acid and base and an organic acid and base; see, for example, *Remington's Pharmaceutical Sciences*, 18th eds., Mack Publishing, Easton PA (1990) or *Remington: The Science and Practice of Pharmacy*, 19th eds., Mack Publishing, Easton PA (1995).

As used herein, the term “Cpd 1” generally refers to a 4-chlorophenyl (S)-6-chloro-1-(4-methoxyphenyl)-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indole-2-carboxylate compound and pharmaceutically acceptable salts thereof. “Cpd 1” may be a substantially pure (e.g., about 90%, about 95%, about 98%, about 99%, or about 99.9% pure) single stereoisomer or a mixture of two or more stereoisomers. In various aspects, the term “Cpd 1” refers to Compound #10 disclosed in International Publication Nos.

WO2005/089764, WO2006/113703, WO2008/127715 and WO2008/127714, all of which are incorporated by reference herein in their entirety.

Method of Use

As demonstrated herein, Cpd 1 inhibits de novo pyrimidine synthesis by targeting inhibition of DHODH function, a rate-limiting enzyme catalyzing oxidation of DHO to orotate [1]. An inhibitor of DHODH function and other therapeutics affecting DHODH alter the nucleotide balance to induce cell-cycle arrest [17]. Exogenously added uridine prevents both the inhibition of DHODH enzyme function and cell cycle arrest mediated by Cpd 1, while exogenously added cytidine only rescues the cell cycle arrest, suggesting that inhibition of DHODH function and cell cycle arrest may be coincidental.

Low pyrimidine pool levels may affect certain translation factor(s) or mechanisms required for the non-canonical protein synthesis of stress-regulated mRNAs. Treatment of Cpd 1-resistant HT1080 cells with Cpd 1 showed that eIF1AY gene expression decreased over 90% when compared with wild type HT1080 cells (Table 1). EIF1A is an isoform that controls translation start codon recognition [18]. Low uridine levels may also affect the mRNA substrate directly: depletion of pyrimidine nucleotides in cancer cells can cause a starvation-like stress response leading to changes in methylation and pseudouridylation of mRNA modifications [19]. Although Cpd 1 demonstrates less inhibition of DHODH functional activity in purified rhDHODH enzymatic assays, Cpd 1 is a more potent inhibitor of DHODH function in hematologic cancer cells than other well-known DHODH inhibitors. Without being limited by theory, this apparent discrepancy is likely due to the presence of the cell membrane lipid bilayer which may facilitate the entry of the hydrophobic Cpd 1 into the binding pocket of the DHODH enzyme [25].

Another potent hDHODH inhibitor brequinar was tested in the 1990s in clinical trials for solid tumors, but was associated with myeloid suppression and has a limited therapeutic window [20;21]. The side-effects associated with these DHODH inhibitors are believed to be due to off-target activities such as kinase inhibition [21;22]. Cpd 1 avoids off-target activities having demonstrated lack of inhibition of kinase enzyme activity in a panel of biologically important human kinases. The antiviral compound GSK983 was recently reported as a DHODH inhibitor [23]. When compared with Cpd 1 at a dose of

10 mg/kg, GSK983 demonstrated a lower plasma concentration as determined by HPLC-MS/MS (Figure 7). In contrast, Cpd 1 (100 nM) induces in vitro differentiation of MOML-13 acute myeloid leukemia (AML) cells relative to DMSO control after 72 hours treatment when stained with anti-CD14 and CD11b antibodies (BD Bioscience). FACS analysis was done on FACSCalibur (Figure 8).

There are no DHODH inhibitors known or approved for use in treating hematologic cancers. The broad activity of Cpd 1 across various leukemia/lymphoma cells demonstrated herein suggest that de novo pyrimidine synthesis is required in proliferation of leukemia cells. These results are consistent with a recent report on DHODH inhibitors [26]. Accordingly, potent and selective activity, favorable pharmaceutical properties and extensive clinical experience suggest that Cpd 1 is a useful agent for treatment of leukemia.

In one aspect, methods for inhibiting or reducing DHODH function in a proliferating cell or cell line are described herein.

In another aspect, a method for inhibiting or reducing DHODH function in a proliferating cell or cell line comprises, contacting Cpd 1 or a composition thereof with a proliferating cell or cell line, which proliferating cell or cell line may be naïve or has been shown to be affected by the inhibition or a reduction in DHODH function.

In another aspect, non-limiting examples of such cells or cell lines are selected from HL-60, HeLa, HT1080, HCT116, HEK293, NCI H460, U-87MG, ASPC-1, PL-45, HPAF-2, PC-3, MDA-MB-231, MDA-MB-468, A431, SNU-1, AGS, Kato III, A549, Calu-6, A375, SY5Y, SKOV3, Capan-1, sNF96.2, TIVE-L1, TIVE-L2, LNCaP cells and the like. In a more specific aspect, the cell or cell line may be a hematologic cancer cell.

In one aspect, a method for inhibiting or reducing DHODH function in a subject having a hematologic cancer comprises, administering Cpd 1 or a composition thereof to the subject as described herein.

In a specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing DHODH function.

In a specific aspect, the hematologic cancer capable of being treated by inhibiting or reducing DHODH function demonstrates, prior to administration of Cpd 1, the reduced expression of uridine salvage enzymes, wherein the low expression of uridine salvage enzymes and dependence on de novo pyrimidine nucleotide synthesis in the subject or in a biological sample taken from the subject indicates that the hematological cancer is amendable to treatment with Cpd 1.

In a specific aspect, a method for inhibiting or reducing DHODH function as described herein inhibits or reduces DHODH function by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95 %, or 100% relative to DHODH function prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for inhibiting or reducing DHODH function as described herein inhibits or reduces DHODH function in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to DHODH function prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for inhibiting or reducing DHODH function as described herein inhibits proliferation or reduces an in vitro or in vivo proliferating cell or cell line population by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95 %, or 100%, relative to the in vitro or in vivo proliferating cell or cell line population prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for inhibiting or reducing DHODH function as described herein inhibits proliferation or reduces an in vitro or in vivo proliferating cell or cell line population in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the in vitro or in vivo proliferating cell or cell line population prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In various aspects, a method for inhibiting or reducing DHODH function as described herein reduces the concentration of DHODH in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 effective to inhibit or reduce DHODH function in the subject is described herein.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces DHODH function by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95 %, or 100% relative to DHODH function of Cpd 1 to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces DHODH function in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to DHODH function prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein reduces the concentration of DHODH in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 effective to inhibit proliferation or reduce an in vitro or in vivo proliferating cell or cell line population in the subject is described herein.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits proliferation or reduces an in vitro or in vivo proliferating cell or cell line population in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%,

95 %, or 100% relative to proliferation or in vitro or in vivo proliferating cell or cell line population in the subject prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits proliferation or reduces an in vitro or in vivo proliferating cell or cell line population in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to proliferation or in vitro or in vivo proliferating cell or cell line population in the subject prior to administration of Cpd 1 to the subject, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits proliferation or reduces an in vitro or in vivo proliferating cell or cell line population in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 effective to inhibit proliferation or reduce an in vitro or in vivo proliferating cell or cell line population in the subject in combination with another therapy (*e.g.*, one or more additional therapies that do not comprise Cpd 1, or that comprise a different anti-proliferative agent) to a subject in need thereof is described herein.

Such methods may involve administering Cpd 1 prior to, concurrent with, or subsequent to administration of the additional therapy. In certain aspects, such methods have an additive or synergistic effect.

In a specific aspect, presented herein is a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprising, administering to a subject in need thereof an effective amount of Cpd 1 and an effective amount of another therapy.

Specific examples of cancers that can be prevented, treated or ameliorated in accordance with the methods provided herein include, but are not limited to, the following: leukemias such as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; plasma cell leukemia and MDS.

In certain aspects, cancers that can be prevented, treated or ameliorated in accordance with the methods provided herein are selected from acute myeloid leukemia, acute lymphocytic leukemia, and MDS.

In one aspect, presented herein is a method for preventing, treating or ameliorating a hematologic cancer, comprising: (a) administering to a subject in need thereof one or more doses of Cpd 1 or a pharmaceutical composition thereof; and (b) monitoring the concentration of certain biomarkers, before and/or after step (a).

In a specific aspect, the monitoring step (b) is carried out before and/or after a certain number of doses (*e.g.*, 1, 2, 4, 6, 8, 10, 12, 14, 15, or 29 doses, or more doses; 2 to 4, 2 to 8, 2 to 20 or 2 to 30 doses) or a certain time period (*e.g.*, 1, 2, 3, 4, 5, 6, or 7 days; or 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 45, 48, or 50 weeks) of administering Cpd 1 or a pharmaceutical composition thereof.

In a specific aspect, one or more of these monitoring parameters are detected prior to administration of Cpd 1 or pharmaceutical composition thereof to the subject.

In a specific aspect, a decrease in the proliferation of an *in vitro* or *in vivo* proliferating cell or cell line population following administration of Cpd 1 or a pharmaceutical composition thereof indicates that the course of treatment is effective for preventing, treating or ameliorating the hematologic cancer.

In a specific aspect, a change in the proliferation of an *in vitro* or *in vivo* proliferating cell or cell line population following administration of Cpd 1 or a pharmaceutical composition thereof may indicate that the dosage, frequency and/or length of administration of Cpd 1

or a pharmaceutical composition thereof may be adjusted (*e.g.*, increased, reduced or maintained).

In a specific aspect, the concentration of certain biomarkers in biological specimens of a subject is monitored before, during and/or after a course of treatment for a hematologic cancer involving the administration of Cpd 1 or a pharmaceutical composition thereof to the subject.

The dosage, frequency and/or length of administration of Cpd 1 or a pharmaceutical composition thereof to a subject might be modified as a result of the proliferation of an *in vitro* or *in vivo* proliferating cell or cell line population. Alternatively, the changes in these monitoring parameters (*e.g.*, concentration of certain biomarkers) might indicate that the course of treatment involving the administration of the Cpd 1 or a pharmaceutical composition thereof is effective in preventing, treating or ameliorating the hematologic cancer.

The concentration of certain biomarkers in a subject may be detected by any technique known to one of skill in the art. In certain aspects, the method for detecting the concentration of certain biomarkers of a subject comprises obtaining a biological sample (*e.g.*, tissue or fluid sample) from the subject and detecting the concentration of the biomarkers in the biological sample (*e.g.*, from plasma, serum, cerebral spinal fluid, urine, or any other biofluids), that has been subjected to certain types of treatment (*e.g.*, centrifugation), and detection by use of immunological techniques, such as ELISA.

In a specific aspect, an ELISA assay, as described herein, may be used to detect the concentration of the biomarkers in a biological sample (*e.g.*, from plasma, serum, cerebral spinal fluid, urine, or any other biofluids) that has been subjected to certain types of treatment (*e.g.*, centrifugation). Other techniques known in the art that may be used to detect the concentration of the biomarkers in a biological sample include multiplex or proteomic assays.

In specific aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein alleviate or manage one, two or more symptoms associated with the hematologic cancer. Alleviating or managing one, two or more symptoms of the hematologic cancer may be used as a clinical endpoint for efficacy of Cpd 1 or a

pharmaceutical composition thereof for preventing, treating or ameliorating the hematologic cancer. In some aspects, the methods for preventing, treating or ameliorating the hematologic cancer provided herein reduce the duration and/or severity of one or more symptoms associated with the hematologic cancer.

In some aspects, the methods for preventing, treating or ameliorating the hematologic cancer provided herein inhibit the onset, progression and/or recurrence of one or more symptoms associated with the hematologic cancer. In some aspects, the methods for treating the hematologic cancer provided herein reduce the number of symptoms associated with the hematologic cancer.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 or a pharmaceutical composition thereof effective to inhibit or reduce the function of DHODH and IDH in the subject is described herein.

In a specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and IDH in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and IDH in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and IDH prior to administration of Cpd 1 or a pharmaceutical composition thereof to the subject, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and IDH in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and IDH prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and IDH in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 or a pharmaceutical composition thereof effective to inhibit or reduce the function of DHODH and wild-type IDH or mutant IDH in the subject is described herein.

In another specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and wild-type IDH or mutant IDH in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH or mutant IDH in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH or mutant IDH in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function

of DHODH and wild-type IDH or mutant IDH in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering an amount of Cpd 1 or a pharmaceutical composition thereof effective to inhibit or reduce the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject is described herein.

In a more specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering a combination product having an amount of Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof effective to inhibit or reduce the function of DHODH and IDH in the subject is described herein.

In a specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and IDH in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and IDH in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and IDH prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and IDH in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and IDH prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof by administering the combination product as described herein inhibits or reduces the function of DHODH and IDH in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering a combination product having an

amount of Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof effective to inhibit or reduce the function of DHODH and wild-type IDH or mutant IDH in the subject is described herein.

In another specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and wild-type IDH or mutant IDH in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH or mutant IDH in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and wild-type IDH or mutant IDH prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH or mutant IDH in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and wild-type IDH or mutant IDH prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering a combination product having an amount of Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof effective

to inhibit or reduce the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject is described herein.

In a more specific aspect, the subject is diagnosed with a hematologic cancer capable of being treated by inhibiting or reducing the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof as described herein inhibits or reduces the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 in the subject in a range of from about 5% to about 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, or from about 40% to about 100%, or any range in between, relative to the function of DHODH and wild-type IDH1, wild-type IDH2, mutant IDH1 or mutant IDH2 prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In various aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof by administering the combination product as described herein decreases the concentration of DHODH and IDH or certain biomarkers in a subject as assessed by methods well known in the art, *e.g.*, ELISA.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer as described herein decreases the concentrations of one or more biomarkers in the subject by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 80%, 85%, 90%, 95%, or 100% relative to the respective concentration observed prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art, *e.g.*, ELISA.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer as described herein decreases the concentrations of one or more biomarkers in the blood of a subject in the range of about 5% to 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, 30% to 100%, or any range in between, relative to the respective concentration observed prior to administration of the combination product having Cpd 1 or a pharmaceutical composition thereof in combination with an amount of an inhibitor of IDH function or a pharmaceutical composition thereof, as assessed by methods well known in the art, *e.g.*, ELISA.

In certain aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein prolong or delay the G1/S or late G1/S phase of the cell cycle (*i.e.*, the period between the late resting or pre-DNA synthesis phase, and the early DNA synthesis phase).

In some aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein reduce, ameliorate, or alleviate the severity of the hematologic cancer and/or one or more symptoms thereof.

In other aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein reduce hospitalization (*e.g.*, the frequency or duration of hospitalization) of a subject diagnosed with the hematologic cancer.

In some aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein reduce hospitalization length of a subject diagnosed with the hematologic cancer.

In certain aspects, the methods provided herein increase the survival of a subject diagnosed with a hematologic cancer. In specific aspects, the methods provided herein increase the survival of a subject diagnosed with a hematologic cancer by about 6 months or more, about 7 months or more, about 8 months or more, about 9 months or more, or about 12 months or more.

In particular aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein inhibit or reduce the progression of the hematologic cancer, or one or more symptoms associated therewith. In specific aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein enhance or improve the therapeutic effect of another therapy (*e.g.*, an anti-cancer agent, radiation, drug therapy, such as chemotherapy, anti-androgen therapy, or surgery). In certain aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein involve the use of Cpd 1 or a pharmaceutical composition thereof as an adjuvant therapy.

In particular aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein reduce the mortality of subjects diagnosed with the hematologic cancer. In certain aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase the number of subjects in remission or decrease the hospitalization rate. In other aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein prevent the development, onset or progression of one or more symptoms associated with the hematologic cancer.

In particular aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase symptom-free survival of cancer subjects. In some aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein do not cure the cancer in subjects, but prevent the progression or worsening of the disease. In some aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein improve the subject's quality of life.

In specific aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein decrease the number of CTCs in the blood of a subject by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%,

80%, 85%, 90%, 95%, 99%, or 100%, relative to the number of CTCs observed prior to administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art.

In particular aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein decrease the number of CTCs in the blood of a subject in the range of about 5% to 20%, 10% to 20%, 10% to 30%, 15% to 40%, 15% to 50%, 20% to 30%, 20% to 40%, 20% to 50%, 30% to 60%, 30% to 70%, 30% to 80%, 30% to 90%, 30% to 95%, 30% to 99%, 30% to 100%, or any range in between, relative to the number of CTCs in the blood observed prior to the administration of Cpd 1 or a pharmaceutical composition thereof, as assessed by methods well known in the art, such as CellSearch immunomagnetic-capture.

In certain aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase the cancer-free survival rate of subjects diagnosed with the cancer. In some aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase relapse-free survival. In certain aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase the number of subjects in remission. In other aspects, the methods for preventing, treating or ameliorating a hematologic cancer provided herein increase the length of remission in subjects.

Treatment Population

In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human who has or is diagnosed with a hematologic cancer. In other aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human predisposed or susceptible to a hematologic cancer. In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human at risk of developing a hematologic cancer.

In one aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human infant. In another aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human toddler. In another aspect, a subject treated for a hematologic cancer in accordance with the

methods provided herein is a human child. In another aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human adult. In another aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a middle-aged human. In another aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is an elderly human.

In certain aspects, a subject treated for cancer in accordance with the methods provided herein has a cancer that metastasized to other areas of the body, such as the bones, lung and liver. In certain aspects, a subject treated for cancer in accordance with the methods provided herein is in remission from the cancer. In some aspects, a subject treated for cancer in accordance with the methods provided herein that has a recurrence of the cancer. In certain aspects, a subject treated in accordance with the methods provided herein is experiencing recurrence of one or more leukemias associated with the hematologic cancer.

In certain aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that is about 1 to about 5 years old, about 5 to 10 years old, about 10 to about 18 years old, about 18 to about 30 years old, about 25 to about 35 years old, about 35 to about 45 years old, about 40 to about 55 years old, about 50 to about 65 years old, about 60 to about 75 years old, about 70 to about 85 years old, about 80 to about 90 years old, about 90 to about 95 years old or about 95 to about 100 years old, or any age in between.

In a specific aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that is 18 years old or older. In a particular aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human child that is between the age of 1 year old to 18 years old. In a certain aspect, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that is between the age of 12 years old and 18 years old. In a certain aspect, the subject is a male human. In another aspect, the subject is a female human. In one aspect, the subject is a female human that is not

pregnant or is not breastfeeding. In one aspect, the subject is a female that is pregnant or will/might become pregnant, or is breast feeding.

In particular aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that is in an immunocompromised state or immunosuppressed state. In certain aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human receiving or recovering from immunosuppressive therapy. In certain aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that has or is at risk of getting a hematologic cancer. In certain aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human who is, will or has undergone surgery, drug therapy, such as chemotherapy, hormonal therapy and/or radiation therapy.

In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is administered Cpd 1 or a pharmaceutical composition thereof or a pharmaceutical composition thereof, or a combination therapy before any adverse effects or intolerance to therapies other than Cpd 1 develops. In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a refractory subject. In certain aspects, a refractory subject is a subject refractory to a standard therapy (*e.g.*, surgery, radiation, anti-androgen therapy and/or drug therapy such as chemotherapy). In certain aspects, a subject with a hematologic cancer is refractory to a therapy when the hematologic cancer has not significantly been eradicated and/or the one or more symptoms have not been significantly alleviated. The determination of whether a patient is refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of a treatment of a hematologic cancer, using art-accepted meanings of “refractory” in such a context.

In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that has proven refractory to therapies other than treatment with Cpd 1 or a pharmaceutical composition thereof, but is no longer on these therapies. In certain aspects, a subject treated for a hematologic cancer in accordance

with the methods provided herein is a human already receiving one or more conventional anti-cancer therapies, such as surgery, drug therapy such as chemotherapy, anti-androgen therapy or radiation. Among these subjects are refractory subjects, subjects who are too young for conventional therapies, and subjects with recurring hematologic cancers despite treatment with existing therapies.

In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human susceptible to adverse reactions to conventional therapies. In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that has not received a therapy, *e.g.*, drug therapy such as chemotherapy, surgery, anti-androgen therapy or radiation therapy, prior to the administration of Cpd 1 or a pharmaceutical composition thereof. In other aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that has received a therapy prior to administration of Cpd 1 or a pharmaceutical composition thereof. In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is a human that has experienced adverse side effects to the prior therapy or the prior therapy was discontinued due to unacceptable levels of toxicity to the human.

In some aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is not, has not and/or will not receive a drug that is primarily metabolized by CYP2D6. In particular aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein has not and will not receive a drug that is primarily metabolized by CYP2D6 1, 2, 3 or 4 weeks before receiving Cpd 1 or a pharmaceutical composition thereof and 1, 2, 3 or 4 weeks after receiving Cpd 1 or a pharmaceutical composition thereof. Examples of such drugs include, without limitation, some antidepressants (*e.g.*, tricyclic antidepressants and selective serotonin uptake inhibitors), some antipsychotics, some beta-adrenergic receptor blockers, certain antiviral agents and certain anti-arrhythmic agents. In specific aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein is not, has not and/or will not receive tamoxifen. In particular aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein has not and will not receive tamoxifen 1, 2, 3 or 4 weeks before receiving Cpd 1 or a pharmaceutical

composition thereof and 1, 2, 3 or 4 weeks after receiving Cpd 1 or a pharmaceutical composition thereof. In specific aspects, a subject treated for a hematologic cancer in accordance with the methods provided herein has received tamoxifen, *e.g.*, for 1, 2, 3 or 4 weeks before receiving Cpd 1 or a pharmaceutical composition thereof.

Dosage and Administration

In accordance with the methods for preventing, treating or ameliorating a hematologic cancer provided herein, Cpd 1 or a pharmaceutical composition thereof can be administered to a subject in need thereof by a variety of routes in amounts which result in a beneficial or therapeutic effect. Cpd 1 or a pharmaceutical composition thereof may be orally administered to a subject in need thereof in accordance with the methods for preventing, treating or ameliorating a hematologic cancer provided herein. The oral administration of Cpd 1 or a pharmaceutical composition thereof may facilitate subjects in need of such treatment complying with a regimen for taking Cpd 1 or a pharmaceutical composition thereof. Thus, in a specific aspect, Cpd 1 or a pharmaceutical composition thereof is administered orally to a subject in need thereof. In another aspect, Cpd 1 or a pharmaceutical composition thereof provided herein can be administered orally, with or without food or water.

Other routes of administration include, but are not limited to, intravenous, intradermal, intrathecal, intramuscular, subcutaneous, intranasal, inhalation, transdermal, topical, transmucosal, intracranial, epidural and intra-synovial. In one aspect, Cpd 1 or a pharmaceutical composition thereof is administered systemically (*e.g.*, parenterally) to a subject in need thereof. In one aspect, Cpd 1 or a pharmaceutical composition thereof is administered via a route that permits Cpd 1 or a pharmaceutical composition thereof to cross the blood-brain barrier (*e.g.*, orally).

Evaluation has indicated that Cpd 1 penetrates the blood-brain barrier. Table 4 provides brain tissue plasma concentration ratios determined by whole-body autoradiography at specified times after a single oral administration of ¹⁴C-Cpd 1 to rats (50 mg/kg).

Table 4. Blood-Brain Barrier Penetration

Tissue	6 Hours		12 Hours		24 Hours		48 Hours		72 Hours	
	M	F	M	F	M	F	M	F	M	F
Cerebellum	1.55	1.23	1.85	2.85	1.74	1.59	1.21	1.17	NA	2.04
Cerebrum	1.52	1.22	1.75	2.79	1.89	1.57	1.35	1.68	NA	1.56
Medulla	1.60	1.42	1.98	3.82	1.83	1.69	1.20	2.01	NA	1.88
Olfactory lobe	1.42	1.38	1.35	2.45	1.23	1.13	0.97	NA	NA	3.33
Pituitary gland	4.06	4.27	3.22	5.48	2.72	2.33	0.89	3.68	NA	1.58
Spinal cord	1.14	0.89	1.24	1.92	1.75	1.60	1.43	1.60	1.84	2.75

In accordance with the methods for preventing, treating or ameliorating a hematologic cancer provided herein that involve administration of Cpd 1 or a pharmaceutical composition thereof in combination with one or more additional therapies, Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies may be administered by the same route or a different route of administration.

The dosage and frequency of administration of Cpd 1 or a pharmaceutical composition thereof is administered to a subject in need thereof in accordance with the methods for preventing, treating or ameliorating a hematologic cancer provided herein will be efficacious while minimizing any side effects. The exact dosage and frequency of administration of Cpd 1 or a pharmaceutical composition thereof can be determined by a practitioner, in light of factors related to the subject that requires treatment.

Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. The dosage and frequency of administration of Cpd 1 or a pharmaceutical composition thereof may be adjusted over time to provide an effective amount of Cpd 1 or a pharmaceutical composition thereof or to maintain the desired effect.

As described herein, the methods for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof presented herein comprises, administering to the

subject an effective amount of Cpd 1 or a pharmaceutical composition thereof, wherein the effective amount is an initial dose, then a loading dose followed by a maintenance dose thereafter.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises, administering to the subject an effective amount of Cpd 1 or a pharmaceutical composition thereof, wherein the effective amount is a loading dose followed by a maintenance dose thereafter.

In a specific aspect, the effective amount is an initial dose administered to the subject for one day once per day.

In a specific aspect, the effective amount is a loading dose administered to the subject for seven days once or twice per day.

In a specific aspect, the effective amount is a loading dose administered to the subject for seven days once per day.

In a specific aspect, the effective amount is a maintenance dose administered to the subject once per day.

In another aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject wherein the effective amount is an initial dose, then a loading dose followed by a maintenance dose, wherein the dose is selected from a dose in a range of from about 1 mg to about 200 mg, from about 1 mg to about 210 mg, from about 1 mg to about 190 mg, from about 1 mg to about 180 mg, from about 1 mg to about 170 mg, from about 1 mg to about 160 mg, from about 1 mg to about 150 mg, from about 1 mg to about 140 mg, from about 1 mg to about 130 mg, from about 1 mg to about 120 mg, from about 1 mg to about 110 mg, from about 1 mg to about 100 mg, from about 1 mg to about 90 mg, from about 1 mg to about 80 mg, from about 1 mg to about 70 mg, from about 1 mg to about 60 mg, from about 1 mg to about 50 mg, from about 1 mg to about 40 mg, from about 1 mg to about 30 mg, from about 1 mg to about 20 mg, from about 1 mg to about 10 mg, from about 10 mg to about 200 mg, from about 20 mg to about 200 mg, from about 30 mg to about

200 mg, from about 40 mg to about 200 mg, from about 50 mg to about 200 mg, from about 60 mg to about 200 mg, from about 70 mg to about 200 mg, from about 80 mg to about 200 mg, from about 90 mg to about 200 mg, from about 100 mg to about 200 mg, from about 110 mg to about 200 mg, from about 120 mg to about 200 mg, from about 130 mg to about 200 mg, from about 140 mg to about 200 mg, from about 150 mg to about 200 mg, from about 160 mg to about 200 mg, from about 170 mg to about 200 mg, from about 180 mg to about 200 mg, from about 190 mg to about 200 mg, from about 200 mg to about 210 mg, from about 10 mg to about 200 mg, from about 10 mg to about 190 mg, from about 10 mg to about 180 mg, from about 10 mg to about 170 mg, from about 10 mg to about 160 mg, from about 10 mg to about 150 mg, from about 10 mg to about 140 mg, from about 10 mg to about 130 mg, from about 10 mg to about 120 mg, from about 10 mg to about 110 mg, from about 10 mg to about 100 mg, from about 10 mg to about 90 mg, from about 10 mg to about 80 mg, from about 10 mg to about 70 mg, from about 10 mg to about 60 mg, from about 10 mg to about 50 mg, from about 10 mg to about 40 mg, from about 10 mg to about 30 mg, from about 10 mg to about 20 mg, from about 20 mg to about 200 mg, from about 20 mg to about 190 mg, from about 20 mg to about 180 mg, from about 20 mg to about 170 mg, from about 20 mg to about 160 mg, from about 20 mg to about 150 mg, from about 20 mg to about 140 mg, from about 20 mg to about 130 mg, from about 20 mg to about 120 mg, from about 20 mg to about 100 mg, from about 20 mg to about 90 mg, from about 20 mg to about 80 mg, from about 20 mg to about 70 mg, from about 20 mg to about 60 mg, from about 20 mg to about 50 mg, from about 20 mg to about 40 mg, from about 20 mg to about 30 mg, from about 30 mg to about 40 mg, from about 30 mg to about 50 mg, from about 30 mg to about 60 mg, from about 30 mg to about 70 mg, from about 40 mg to about 50 mg, from about 40 mg to about 60 mg, from about 40 mg to about 70 mg, from about 50 mg to about 60 mg, from about 50 mg to about 70 mg, from about 60 mg to about 70 mg, from about 90 mg to about 210 mg, from about 90 mg to about 200 mg, from about 90 mg to about 190 mg, from about 90 mg to about 180 mg, from about 90 mg to about 170 mg, from about 90 mg to about 160 mg, from about 90 mg to about 150 mg, from about 90 mg to about 140 mg, from about 90 mg to about 130 mg, from about 90 mg to about 120 mg, from about 90 mg to

about 110 mg, from about 100 mg to about 210 mg, from about 100 mg to about 200 mg, from about 100 mg to about 190 mg, from about 100 mg to about 180 mg, from about 100 mg to about 170 mg, from about 100 mg to about 160 mg, from about 100 mg to about 150 mg, from about 100 mg to about 140 mg, from about 100 mg to about 130 mg, from about 100 mg to about 120 mg, from about 100 mg to about 110 mg, from about 110 mg to about 210 mg, from about 120 mg to about 210 mg, from about 130 mg to about 210 mg, from about 140 mg to about 210 mg, from about 150 mg to about 210 mg, from about 160 mg to about 210 mg, from about 170 mg to about 210 mg, from about 180 mg to about 210 mg, from about 190 mg to about 210 mg, from about 1 mg to about 110 mg, from about 1 mg to about 100 mg, from about 1 mg to about 90 mg, from about 1 mg to about 80 mg, from about 1 mg to about 70 mg, from about 1 mg to about 60 mg, from about 1 mg to about 50 mg, from about 1 mg to about 40 mg, from about 1 mg to about 30 mg, from about 1 mg to about 20 mg, from about 1 mg to about 10 mg, from about 10 mg to about 110 mg, from about 10 mg to about 100 mg, from about 10 mg to about 90 mg, from about 10 mg to about 80 mg, from about 10 mg to about 70 mg, from about 10 mg to about 60 mg, from about 10 mg to about 50 mg, from about 10 mg to about 40 mg, from about 10 mg to about 30 mg, from about 10 mg to about 20 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject wherein the effective amount is a dose selected from a dose in a range of from about 1 mg to about 200 mg, from about 1 mg to about 210 mg, from about 1 mg to about 190 mg, from about 1 mg to about 180 mg, from about 1 mg to about 170 mg, from about 1 mg to about 160 mg, from about 1 mg to about 150 mg, from about 1 mg to about 140 mg, from about 1 mg to about 130 mg, from about 1 mg to about 120 mg, from about 1 mg to about 110 mg, from about 1 mg to about 100 mg, from about 1 mg to about 90 mg, from about 1 mg to about 80 mg, from about 1 mg to about 70 mg, from about 1 mg to about 60 mg, from about 1 mg to about 50 mg, from about 1 mg to about 40 mg, from about 1 mg to about 30 mg, from about 1 mg to about 20 mg, from about 1 mg to about 10 mg, from about

10 mg to about 200 mg, from about 20 mg to about 200 mg, from about 30 mg to about 200 mg, from about 40 mg to about 200 mg, from about 50 mg to about 200 mg, from about 60 mg to about 200 mg, from about 70 mg to about 200 mg, from about 80 mg to about 200 mg, from about 90 mg to about 200 mg, from about 100 mg to about 200 mg, from about 110 mg to about 200 mg, from about 120 mg to about 200 mg, from about 130 mg to about 200 mg, from about 140 mg to about 200 mg, from about 150 mg to about 200 mg, from about 160 mg to about 200 mg, from about 170 mg to about 200 mg, from about 180 mg to about 200 mg, from about 190 mg to about 200 mg, from about 200 mg to about 210 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dose selected from a dose in a range of from about 10 mg to about 200 mg, from about 10 mg to about 190 mg, from about 10 mg to about 180 mg, from about 10 mg to about 170 mg, from about 10 mg to about 160 mg, from about 10 mg to about 150 mg, from about 10 mg to about 140 mg, from about 10 mg to about 130 mg, from about 10 mg to about 120 mg, from about 10 mg to about 110 mg, from about 10 mg to about 100 mg, from about 10 mg to about 90 mg, from about 10 mg to about 80 mg, from about 10 mg to about 70 mg, from about 10 mg to about 60 mg, from about 10 mg to about 50 mg, from about 10 mg to about 40 mg, from about 10 mg to about 30 mg, from about 10 mg to about 20 mg, from about 20 mg to about 200 mg, from about 20 mg to about 190 mg, from about 20 mg to about 180 mg, from about 20 mg to about 170 mg, from about 20 mg to about 160 mg, from about 20 mg to about 150 mg, from about 20 mg to about 140 mg, from about 20 mg to about 130 mg, from about 20 mg to about 120 mg, from about 20 mg to about 120 mg, from about 20 mg to about 100 mg, from about 20 mg to about 90 mg, from about 20 mg to about 80 mg, from about 20 mg to about 70 mg, from about 20 mg to about 60 mg, from about 20 mg to about 50 mg, from about 20 mg to about 40 mg, from about 20 mg to about 30 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose selected from a dose in a range of from about 30 mg to about 40 mg, from about 30 mg to about 50 mg, from about 30 mg to about 60 mg, from about 30 mg to about 70 mg, from about 40 mg to about 50 mg, from about 40 mg to about 60 mg, from about 40 mg to about 70 mg, from about 50 mg to about 60 mg, from about 50 mg to about 70 mg, from about 60 mg to about 70 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose selected from a dose in a range of from about 40 mg to about 50 mg, from about 40 mg to about 60 mg, from about 50 mg to about 60 mg, from about 50 mg to about 70 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose selected from a dose in a range of from about 40 mg to about 60 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose of about 50 mg administered orally once per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose selected from a dose in a range of from about 90 mg to about 210 mg, from about 90 mg to about 200 mg, from about 90 mg to about 190 mg, from

about 90 mg to about 180 mg, from about 90 mg to about 170 mg, from about 90 mg to about 160 mg, from about 90 mg to about 150 mg, from about 90 mg to about 140 mg, from about 90 mg to about 130 mg, from about 90 mg to about 120 mg, from about 90 mg to about 110 mg, from about 100 mg to about 210 mg, from about 100 mg to about 200 mg, from about 100 mg to about 190 mg, from about 100 mg to about 180 mg, from about 100 mg to about 170 mg, from about 100 mg to about 160 mg, from about 100 mg to about 150 mg, from about 100 mg to about 140 mg, from about 100 mg to about 130 mg, from about 100 mg to about 120 mg, from about 100 mg to about 110 mg, from about 110 mg to about 210 mg, from about 120 mg to about 210 mg, from about 130 mg to about 210 mg, from about 140 mg to about 210 mg, from about 150 mg to about 210 mg, from about 160 mg to about 210 mg, from about 170 mg to about 210 mg, from about 180 mg to about 210 mg, from about 190 mg to about 210 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose selected from a dose in a range of from about 90 mg to about 210 mg, from about 90 mg to about 200 mg, from about 90 mg to about 190 mg, from about 100 mg to about 210 mg, from about 100 mg to about 200 mg, from about 100 mg to about 190 mg, from about 110 mg to about 210 mg, from about 120 mg to about 210 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose in a range of from about 100 mg to about 200 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount

of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose of about 100 mg administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 1 mg to about 110 mg, from about 1 mg to about 100 mg, from about 1 mg to about 90 mg, from about 1 mg to about 80 mg, from about 1 mg to about 70 mg, from about 1 mg to about 60 mg, from about 1 mg to about 50 mg, from about 1 mg to about 40 mg, from about 1 mg to about 30 mg, from about 1 mg to about 20 mg, from about 1 mg to about 10 mg, from about 10 mg to about 110 mg, from about 10 mg to about 100 mg, from about 10 mg to about 90 mg, from about 10 mg to about 80 mg, from about 10 mg to about 70 mg, from about 10 mg to about 60 mg, from about 10 mg to about 50 mg, from about 10 mg to about 40 mg, from about 10 mg to about 30 mg, from about 10 mg to about 20 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 1 mg to about 110 mg, from about 1 mg to about 100 mg, from about 1 mg to about 90 mg, from about 10 mg to about 110 mg, from about 10 mg to about 100 mg, from about 10 mg to about 90 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 1 mg to about 100 mg, from about 10 mg to about 100 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 1 mg to about 100 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 10 mg to about 100 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose of about 50 mg administered orally once or twice per day.

In another aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose, then a loading dose followed by a maintenance dose, wherein the dose is selected from a dose in a range of from about 4 mg to about 84 mg, from about 4 mg to about 80 mg, from about 4 mg to about 76 mg, from about 4 mg to about 72 mg, from about 4 mg to about 68 mg, from about 4 mg to about 64 mg, from about 4 mg to about 60 mg, from about 4 mg to about 56 mg, from about 4 mg to about 52 mg, from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 4 mg to about 28 mg, from about 4 mg to about 24 mg, from about 4 mg to about 20 mg, from about 4 mg to about 16 mg, from about 4 mg to about 12 mg, from about 4 mg to about 8 mg, from about 8 mg to about 80 mg, from about 12 mg to about 80 mg,

from about 16 mg to about 80 mg, from about 20 mg to about 80 mg, from about 24 mg to about 80 mg, from about 28 mg to about 80 mg, from about 32 mg to about 80 mg, from about 36 mg to about 80 mg, from about 40 mg to about 80 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, from about 52 mg to about 80 mg, from about 56 mg to about 80 mg, from about 60 mg to about 80 mg, from about 64 mg to about 80 mg, from about 68 mg to about 80 mg, from about 72 mg to about 80 mg, from about 76 mg to about 80 mg, from about 8 mg to about 84 mg, from about 12 mg to about 84 mg, from about 16 mg to about 84 mg, from about 20 mg to about 84 mg, from about 24 mg to about 84 mg, from about 28 mg to about 84 mg, from about 32 mg to about 84 mg, from about 36 mg to about 84 mg, from about 40 mg to about 84 mg, from about 44 mg to about 84 mg, from about 48 mg to about 84 mg, from about 52 mg to about 84 mg, from about 56 mg to about 84 mg, from about 60 mg to about 84 mg, from about 64 mg to about 84 mg, from about 68 mg to about 84 mg, from about 72 mg to about 84 mg, from about 76 mg to about 84 mg, from about 12 mg to about 16 mg, from about 12 mg to about 20 mg, from about 12 mg to about 24 mg, from about 12 mg to about 28 mg, from about 12 mg to about 28 mg, from about 16 mg to about 20 mg, from about 16 mg to about 24 mg, from about 20 mg to about 24 mg, from about 20 mg to about 28 mg, from about 24 mg to about 28 mg, from about 32 mg to about 88 mg, 32 mg to about 84 mg, from about 32 mg to about 80 mg, from about 32 mg to about 76 mg, from about 32 mg to about 72 mg, from about 32 mg to about 68 mg, from about 32 mg to about 64 mg, from about 32 mg to about 60 mg, from about 32 mg to about 56 mg, from about 32 mg to about 52 mg, from about 32 mg to about 48 mg, from about 32 mg to about 44 mg, from about 32 mg to about 40 mg, from about 32 mg to about 36 mg, from about 36 mg to about 88 mg, from about 36 mg to about 84 mg, from about 36 mg to about 80 mg, from about 36 mg to about 76 mg, from about 36 mg to about 72 mg, from about 36 mg to about 68 mg, from about 36 mg to about 64 mg, from about 36 mg to about 60 mg, from about 36 mg to about 56 mg, from about 36 mg to about 52 mg, from about 36 mg to about 48 mg, from about 36 mg to about 44 mg, from about 36 mg to about 40 mg, from about 40 mg to about 88 mg, 40 mg to about 84 mg, from about 40 mg to about 80 mg, from about 40 mg to about 76 mg, from about 40 mg to about 72 mg, from about 40 mg to about 68 mg, from about 40 mg to about 64 mg, from

about 40 mg to about 60 mg, from about 40 mg to about 56 mg, from about 40 mg to about 52 mg, from about 40 mg to about 48 mg, from about 40 mg to about 44 mg, from about 44 mg to about 88 mg, from about 48 mg to about 88 mg, from about 52 mg to about 88 mg, from about 56 mg to about 88 mg, from about 60 mg to about 88 mg, from about 64 mg to about 88 mg, from about 68 mg to about 88 mg, from about 72 mg to about 88 mg, from about 76 mg to about 88 mg, from about 80 mg to about 88 mg, from about 84 mg to about 88 mg, from about 44 mg to about 84 mg, from about 48 mg to about 84 mg, from about 52 mg to about 84 mg, from about 56 mg to about 84 mg, from about 60 mg to about 84 mg, from about 64 mg to about 84 mg, from about 68 mg to about 84 mg, from about 72 mg to about 84 mg, from about 76 mg to about 84 mg, from about 80 mg to about 84 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, from about 52 mg to about 80 mg, from about 56 mg to about 80 mg, from about 60 mg to about 80 mg, from about 64 mg to about 80 mg, from about 68 mg to about 80 mg, from about 72 mg to about 80 mg, from about 76 mg to about 80 mg, from about 84 mg to about 80 mg, from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 4 mg to about 28 mg, from about 4 mg to about 24 mg, from about 4 mg to about 20 mg, from about 4 mg to about 16 mg, from about 4 mg to about 12 mg, from about 4 mg to about 8 mg, from about 8 mg to about 48 mg, from about 8 mg to about 44 mg, from about 8 mg to about 40 mg, from about 8 mg to about 36 mg, from about 8 mg to about 32 mg, from about 8 mg to about 28 mg, from about 8 mg to about 24 mg, from about 8 mg to about 20 mg, from about 8 mg to about 16 mg, from about 8 mg to about 12 mg, from about 12 mg to about 48 mg, from about 12 mg to about 44 mg, from about 12 mg to about 40 mg, from about 12 mg to about 36 mg, from about 12 mg to about 32 mg, from about 12 mg to about 28 mg, from about 12 mg to about 24 mg, from about 12 mg to about 20 mg, from about 12 mg to about 16 mg, from about 16 mg to about 48 mg, from about 20 mg to about 48 mg, from about 24 mg to about 48 mg, from about 26 mg to about 48 mg, from about 30 mg to about 48 mg, from about 34 mg to about 48 mg, from about 38 mg to about 48 mg, from about 40 mg to about 48 mg, from about 44 mg to about 48 mg, from about 16 mg to about 44 mg, from about 20 mg to about 44 mg, from about 24 mg to about 44 mg, from about 26

mg to about 44 mg, from about 30 mg to about 44 mg, from about 34 mg to about 44 mg, from about 38 mg to about 44 mg, from about 40 mg to about 44 mg, from about 16 mg to about 40 mg, from about 20 mg to about 40 mg, from about 24 mg to about 40 mg, from about 26 mg to about 40 mg, from about 30 mg to about 40 mg, from about 34 mg to about 40 mg, from about 38 mg to about 40 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dose selected from a dose in a range of from about 4 mg to about 84 mg, from about 4 mg to about 80 mg, from about 4 mg to about 76 mg, from about 4 mg to about 72 mg, from about 4 mg to about 68 mg, from about 4 mg to about 64 mg, from about 4 mg to about 60 mg, from about 4 mg to about 56 mg, from about 4 mg to about 52 mg, from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 4 mg to about 28 mg, from about 4 mg to about 24 mg, from about 4 mg to about 20 mg, from about 4 mg to about 16 mg, from about 4 mg to about 12 mg, from about 4 mg to about 8 mg, from about 8 mg to about 80 mg, from about 12 mg to about 80 mg, from about 16 mg to about 80 mg, from about 20 mg to about 80 mg, from about 24 mg to about 80 mg, from about 28 mg to about 80 mg, from about 32 mg to about 80 mg, from about 36 mg to about 80 mg, from about 40 mg to about 80 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, from about 52 mg to about 80 mg, from about 56 mg to about 80 mg, from about 60 mg to about 80 mg, from about 64 mg to about 80 mg, from about 68 mg to about 80 mg, from about 72 mg to about 80 mg, from about 76 mg to about 80 mg, from about 8 mg to about 84 mg, from about 12 mg to about 84 mg, from about 16 mg to about 84 mg, from about 20 mg to about 84 mg, from about 24 mg to about 84 mg, from about 28 mg to about 84 mg, from about 32 mg to about 84 mg, from about 36 mg to about 84 mg, from about 40 mg to about 84 mg, from about 44 mg to about 84 mg, from about 48 mg to about 84 mg, from about 52 mg to about 84 mg, from about 56 mg to about 84 mg, from about 60 mg to about 84 mg, from about 64 mg to about 84 mg, from about 68 mg to about 84 mg, from about 72

mg to about 84 mg, from about 76 mg to about 84 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose selected from a dose in a range of from about 12 mg to about 16 mg, from about 12 mg to about 20 mg, from about 12 mg to about 24 mg, from about 12 mg to about 28 mg, from about 12 mg to about 28 mg, from about 16 mg to about 20 mg, from about 16 mg to about 24 mg, from about 20 mg to about 24 mg, from about 20 mg to about 28 mg, from about 24 mg to about 28 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose selected from a dose in a range of from about 16 mg to about 20 mg, from about 16 mg to about 24 mg, from about 20 mg to about 24 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an initial dose of about 20 mg administered orally once per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose selected from a dose in a range of from about 32 mg to about 88 mg, 32 mg to about 84 mg, from about 32 mg to about 80 mg, from about 32 mg to about 76 mg, from about 32 mg to about 72 mg, from about 32 mg to about 68 mg, from about 32 mg to about 64 mg, from about 32 mg to about 60 mg, from about 32 mg to about 56 mg, from about 32 mg to about 52 mg, from about 32 mg to about 48 mg, from about 32 mg to about 44 mg, from about 32 mg to about 40 mg, from about 32 mg to

about 36 mg, from about 36 mg to about 88 mg, from about 36 mg to about 84 mg, from about 36 mg to about 80 mg, from about 36 mg to about 76 mg, from about 36 mg to about 72 mg, from about 36 mg to about 68 mg, from about 36 mg to about 64 mg, from about 36 mg to about 60 mg, from about 36 mg to about 56 mg, from about 36 mg to about 52 mg, from about 36 mg to about 48 mg, from about 36 mg to about 44 mg, from about 36 mg to about 40 mg, from about 40 mg to about 88 mg, 40 mg to about 84 mg, from about 40 mg to about 80 mg, from about 40 mg to about 76 mg, from about 40 mg to about 72 mg, from about 40 mg to about 68 mg, from about 40 mg to about 64 mg, from about 40 mg to about 60 mg, from about 40 mg to about 56 mg, from about 40 mg to about 52 mg, from about 40 mg to about 48 mg, from about 40 mg to about 44 mg, from about 44 mg to about 88 mg, from about 48 mg to about 88 mg, from about 52 mg to about 88 mg, from about 56 mg to about 88 mg, from about 60 mg to about 88 mg, from about 64 mg to about 88 mg, from about 68 mg to about 88 mg, from about 72 mg to about 88 mg, from about 76 mg to about 88 mg, from about 80 mg to about 88 mg, from about 84 mg to about 88 mg, from about 44 mg to about 84 mg, from about 48 mg to about 84 mg, from about 52 mg to about 84 mg, from about 56 mg to about 84 mg, from about 60 mg to about 84 mg, from about 64 mg to about 84 mg, from about 68 mg to about 84 mg, from about 72 mg to about 84 mg, from about 76 mg to about 84 mg, from about 80 mg to about 84 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, from about 52 mg to about 80 mg, from about 56 mg to about 80 mg, from about 60 mg to about 80 mg, from about 64 mg to about 80 mg, from about 68 mg to about 80 mg, from about 72 mg to about 80 mg, from about 76 mg to about 80 mg, from about 84 mg to about 80 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose selected from a dose in a range of from about 36 mg to about 88 mg, from about 36 mg to about 84 mg, from about 36 mg to about 80 mg, from about 36 mg to about 76 mg, from about 36 mg to about 72 mg, from about 40 mg to about 88 mg, 40 mg to about 84 mg, from about 40 mg to about 80 mg, from about 40 mg to

about 76 mg, from about 40 mg to about 72 mg, from about 44 mg to about 88 mg, from about 48 mg to about 88 mg, from about 44 mg to about 84 mg, from about 48 mg to about 84 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose selected from a dose in a range of from about 36 mg to about 84 mg, from about 36 mg to about 80 mg, from about 36 mg to about 76 mg, 40 mg to about 84 mg, from about 40 mg to about 80 mg, from about 40 mg to about 76 mg, from about 44 mg to about 80 mg, from about 48 mg to about 80 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose in a range of from about 40 mg to about 80 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose of about 40 mg administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a loading dose of about 40 mg administered orally once per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from

about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 4 mg to about 28 mg, from about 4 mg to about 24 mg, from about 4 mg to about 20 mg, from about 4 mg to about 16 mg, from about 4 mg to about 12 mg, from about 4 mg to about 8 mg, from about 8 mg to about 48 mg, from about 8 mg to about 44 mg, from about 8 mg to about 40 mg, from about 8 mg to about 36 mg, from about 8 mg to about 32 mg, from about 8 mg to about 28 mg, from about 8 mg to about 24 mg, from about 8 mg to about 20 mg, from about 8 mg to about 16 mg, from about 8 mg to about 12 mg, from about 12 mg to about 48 mg, from about 12 mg to about 44 mg, from about 12 mg to about 40 mg, from about 12 mg to about 36 mg, from about 12 mg to about 32 mg, from about 12 mg to about 28 mg, from about 12 mg to about 24 mg, from about 12 mg to about 20 mg, from about 12 mg to about 16 mg, from about 16 mg to about 48 mg, from about 20 mg to about 48 mg, from about 24 mg to about 48 mg, from about 26 mg to about 48 mg, from about 30 mg to about 48 mg, from about 34 mg to about 48 mg, from about 38 mg to about 48 mg, from about 40 mg to about 48 mg, from about 44 mg to about 48 mg, from about 16 mg to about 44 mg, from about 20 mg to about 44 mg, from about 24 mg to about 44 mg, from about 26 mg to about 44 mg, from about 30 mg to about 44 mg, from about 34 mg to about 44 mg, from about 38 mg to about 44 mg, from about 40 mg to about 44 mg, from about 16 mg to about 40 mg, from about 20 mg to about 40 mg, from about 24 mg to about 40 mg, from about 26 mg to about 40 mg, from about 30 mg to about 40 mg, from about 34 mg to about 40 mg, from about 38 mg to about 40 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 8 mg to about 48 mg, from about 8 mg to about 44 mg, from about 8 mg to about 40 mg, from about 8 mg to about 36 mg, from about 8 mg to about 32 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose selected from a dose in a range of from about 4 mg to about 48 mg, from about 4 mg to about 44 mg, from about 4 mg to about 40 mg, from about 4 mg to about 36 mg, from about 4 mg to about 32 mg, from about 8 mg to about 48 mg, from about 8 mg to about 44 mg, from about 8 mg to about 40 mg, from about 8 mg to about 36 mg, from about 8 mg to about 32 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose in a range of from about 4 mg to about 40 mg, and the like, or any range in between, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose of about 20 mg, administered orally once or twice per day.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a maintenance dose of about 20 mg, administered orally once per day.

In some aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that is expressed as mg per meter squared (mg/m^2). The mg/m^2 for Cpd 1 may be determined, for example, by multiplying a conversion factor for an animal by an animal dose in mg per kilogram (mg/kg) to obtain the dose in mg/m^2 for human dose equivalent. For regulatory purposes, the following conversion factors may be

used: Mouse = 3, Hamster = 4.1, Rat = 6, Guinea Pig = 7.7. (based on Freireich *et al.*, Cancer Chemother. Rep. 50(4):219-244 (1966)). The height and weight of a human may be used to calculate a human body surface area applying Boyd's Formula of Body Surface Area. In specific aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is an amount in the range of from about 0.1 mg/m² to about 1000 mg/m², or any range in between.

In one aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a target mean plasma concentration of Cpd 1 in a subject with a hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In another aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 4 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 8 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 12 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 16 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 20 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 24 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 28 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 32 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 36 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 40 µg·hr/mL to approximately 44 µg·hr/mL, from approximately 4 µg·hr/mL to approximately 40 µg·hr/mL, from approximately 8 µg·hr/mL to approximately 40 µg·hr/mL, from approximately 12 µg·hr/mL to approximately 40 µg·hr/mL, from approximately 16 µg·hr/mL to approximately 40 µg·hr/mL, from approximately 20 µg·hr/mL to approximately 40 µg·hr/mL, from approximately 24 µg·hr/mL to approximately 40 µg·hr/mL, from

approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 40 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 40 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 40 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 44 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 44 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12

$\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $16 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $20 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $24 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $28 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $32 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $44 \mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately $4 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $8 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $12 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $16 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $20 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $24 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $28 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $32 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $40 \mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately $4 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $8 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $12 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $16 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately $20 \mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately $36 \mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

$\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 36 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount

of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 32 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 28 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 24 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, from approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between, in a subject with the hematologic cancer or an animal model (*e.g.*, an animal model with a pre-established hematologic cancer).

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 6 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective

amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 10 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 14 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 18 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount

of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

To achieve such plasma concentrations, an initial dose, then a loading dose followed by a maintenance dose of Cpd 1 or a pharmaceutical composition thereof may be administered. In certain aspects, subsequent doses of Cpd 1 or a pharmaceutical composition thereof may be adjusted accordingly based on the mean plasma concentrations of Cpd 1 achieved with initial doses of Cpd 1 or a pharmaceutical composition thereof administered to the subject.

In a specific aspect, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, the effective amount may be adjusted based on the mean Area Under the Curve (AUC) in a 24 hour period (AUC_{0-24}), wherein a mean $\text{AUC}_{0-24} < 8 \mu\text{g}\cdot\text{hr}/\text{mL}$ will allow a dose increase of up to about 100% of the initial dose administered; wherein a mean AUC_{0-24} between ≥ 8 and $< 14 \mu\text{g}\cdot\text{hr}/\text{mL}$ will allow a dose increase of up to about 50% of the initial dose administered, wherein a mean $\text{AUC}_{0-24} \geq 14$ and $< 16 \mu\text{g}\cdot\text{hr}/\text{mL}$ will allow a dose increase of up to about 33% of the initial dose administered, wherein a mean $\text{AUC}_{0-24} \geq 16$ and $< 18 \mu\text{g}\cdot\text{hr}/\text{mL}$ will allow a dose increase of up to about 25% of the initial dose administered.

In specific aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves a reduced target mean plasma concentration of one or more biomarkers in a subject with the hematologic cancer or an animal model (*e.g.*, a hematologic cancer animal model).

In particular aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount is a dosage that achieves the desired tissue to mean plasma concentration

ratios of Cpd 1 or a pharmaceutical composition thereof as determined, *e.g.*, by any imaging techniques known in the art such as whole-body autoradiography, in a subject with the hematologic cancer or an animal model (such as an animal model with a pre-established hematologic cancer).

In some aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of an effective amount of Cpd 1 or a pharmaceutical composition thereof to the subject, wherein the effective amount may or may not be the same for each dose. In particular aspects, a first (*i.e.*, initial) dose of Cpd 1 or a pharmaceutical composition thereof is administered to a subject in need thereof for a first period of time, followed by a second (*i.e.*, loading) dose of Cpd 1 or a pharmaceutical composition thereof is administered to the subject for a second period of time and, subsequently, a third (*i.e.*, maintenance) dose of Cpd 1 or a pharmaceutical composition thereof is administered to the subject for a second period of time. The first dose may be more than the second dose, or the first dose may be less than the second dose. In similar fashion, the third dose of Cpd 1 or a pharmaceutical composition thereof may be more or less than the second dose and more or less than the first dose.

In some aspects, the dosage amounts described herein refer to total amounts administered; that is, if more than one Compound is administered, then, in some aspects, the dosages correspond to the total amount administered. In a specific aspect, oral compositions contain about 5% to about 95% of Cpd 1 by weight.

The length of time that a subject in need thereof is administered Cpd 1 or a pharmaceutical composition thereof in accordance with a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof will be the time period that is determined by cancer free survival or freedom from symptoms. In certain aspects, a method for treating a hematologic cancer presented herein comprises the administration of Cpd 1 or a pharmaceutical composition thereof for a period of time until the severity and/or number of one or more symptoms associated with the hematologic cancer decreases.

In some aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of Cpd 1 or a pharmaceutical composition thereof for up to 48 weeks. In other aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of Cpd 1 or a pharmaceutical composition thereof for up to 4 weeks, 8 weeks, 12 weeks, 16 weeks, 20 weeks, 24 weeks, 26 weeks (0.5 year), 52 weeks (1 year), 78 weeks (1.5 years), 104 weeks (2 years), or 130 weeks (2.5 years) or more.

In certain aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of Cpd 1 or a pharmaceutical composition thereof for an indefinite period of time. In some aspects, a method for treating a hematologic cancer presented herein comprises the administration of Cpd 1 or a pharmaceutical composition thereof for a period of time followed by a period of rest (*i.e.*, a period wherein Cpd 1 or a pharmaceutical composition thereof is not administered) before the administration of Cpd 1 or a pharmaceutical composition thereof is resumed.

In specific aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises the administration of Cpd 1 or a pharmaceutical composition thereof in cycles, *e.g.*, 1 week cycles, 2 week cycles, 3 week cycles, 4 week cycles, 5 week cycles, 6 week cycles, 8 week cycles, 9 week cycles, 10 week cycles, 11 week cycles, or 12 week cycles. In such cycles, Cpd 1 or a pharmaceutical composition thereof may be administered once, twice, three times, or four times daily.

In specific aspects, the period of time of administration of Cpd 1 or a pharmaceutical composition thereof may be dictated by one or more monitoring parameters, *e.g.*, concentration of certain biomarkers.

In particular aspects, the period of time of administration of Cpd 1 or a pharmaceutical composition thereof may be adjusted based on one or more monitoring parameters, *e.g.*, concentration of biomarkers.

In certain aspects, in accordance with a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof, Cpd 1 or a pharmaceutical

composition thereof is administered to a subject in need thereof prior to, concurrently with, or after a meal (*e.g.*, breakfast, lunch, or dinner). In specific aspects, in accordance with the methods for treating a hematologic cancer presented herein, Cpd 1 or a pharmaceutical composition thereof is administered to a subject in need thereof in the morning (*e.g.*, between 5 am and 12 pm).

In certain aspects, in accordance with a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof, Cpd 1 or a pharmaceutical composition thereof is administered to a subject in need thereof at noon (*i.e.*, 12 pm). In particular aspects, in accordance with the methods for treating a hematologic cancer presented herein, Cpd 1 or a pharmaceutical composition thereof is administered to a subject in need thereof in the afternoon (*e.g.*, between 12 pm and 5 pm), evening (*e.g.*, between 5 pm and bedtime), and/or before bedtime.

In specific aspects, a dose of Cpd 1 or a pharmaceutical composition thereof is administered to a subject once per day, twice per day, three times per day; once, twice or three times every other day (*i.e.*, on alternate days); once, twice or three times every two days; once, twice or three times every three days; once, twice or three times every four days; once, twice or three times every five days; once, twice, or three times once a week, biweekly or monthly.

In certain aspects, a dose of Cpd 1 or a pharmaceutical composition thereof is administered to a subject once or twice per day.

In certain aspects, a dose of Cpd 1 or a pharmaceutical composition thereof is administered to a subject once per day.

Combination Therapies

Presented herein are combination therapies for the treatment of a hematologic cancer which involve the administration of Cpd 1 or a pharmaceutical composition thereof in combination with one or more additional therapies to a subject in need thereof. In a specific aspect, presented herein are combination therapies for the treatment of a hematologic cancer which involve the administration of an effective amount of Cpd 1 or

a pharmaceutical composition thereof in combination with an effective amount of another therapy to a subject in need thereof.

As used herein, the term “in combination,” refers, in the context of the administration of Cpd 1 or a pharmaceutical composition thereof, to the administration of Cpd 1 or a pharmaceutical composition thereof prior to, concurrently with, or subsequent to the administration of one or more additional therapies (e.g., agents, surgery, or radiation) for use in treating a hematologic cancer. The use of the term “in combination” does not restrict the order in which one or more therapeutic agents and one or more additional therapies are administered to a subject. In specific aspects, the interval of time between the administration of Cpd 1 or a pharmaceutical composition thereof and the administration of one or more additional therapies may be about 1-5 minutes, 1-30 minutes, 30 minutes to 60 minutes, 1 hour, 1-2 hours, 2-6 hours, 2-12 hours, 12-24 hours, 1-2 days, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 15 weeks, 20 weeks, 26 weeks, 52 weeks, 11-15 weeks, 15-20 weeks, 20-30 weeks, 30-40 weeks, 40-50 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 1 year, 2 years, or any period of time in between. In certain aspects, Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies are administered less than 1 day, 1 week, 2 weeks, 3 weeks, 4 weeks, one month, 2 months, 3 months, 6 months, 1 year, 2 years, or 5 years apart.

In some aspects, the combination therapies provided herein involve administering Cpd 1 or a pharmaceutical composition thereof daily, and administering one or more additional therapies once a week, once every 2 weeks, once every 3 weeks, once every 4 weeks, once every month, once every 2 months (e.g., approximately 8 weeks), once every 3 months (e.g., approximately 12 weeks), or once every 4 months (e.g., approximately 16 weeks). In certain aspects, Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies are cyclically administered to a subject. Cycling therapy comprises the administration of Cpd 1 or a pharmaceutical composition thereof for a period of time, followed by the administration of one or more additional therapies for a period of time, and repeating this sequential administration. In certain

aspects, cycling therapy may also include a period of rest where Cpd 1 or a pharmaceutical composition thereof or the additional therapy is not administered for a period of time (e.g., 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 10 weeks, 20 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 2 years, or 3 years). In an aspect, the number of cycles administered is from 1 to 12 cycles, from 2 to 10 cycles, or from 2 to 8 cycles.

In some aspects, a method for preventing, treating or ameliorating a hematologic cancer in a subject in need thereof comprises administering Cpd 1 or a pharmaceutical composition thereof as a single agent for a period of time prior to administering Cpd 1 or a pharmaceutical composition thereof in combination with an additional therapy. In certain aspects, the methods for treating a hematologic cancer provided herein comprise administering an additional therapy alone for a period of time prior to administering Cpd 1 or a pharmaceutical composition thereof in combination with the additional therapy.

In some aspects, the administration of Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies in accordance with the methods presented herein have an additive effect relative the administration of Cpd 1 or a pharmaceutical composition thereof or said one or more additional therapies alone. In some aspects, the administration of Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies in accordance with the methods presented herein have a synergistic effect relative to the administration of Cpd 1 or a pharmaceutical composition thereof or said one or more additional therapies alone.

As used herein, the term “synergistic,” refers to the effect of the administration of Cpd 1 or a pharmaceutical composition thereof in combination with one or more additional therapies (e.g., agents), which combination is more effective than the additive effects of any two or more single therapies (e.g., agents).

In a specific aspect, a synergistic effect of a combination therapy permits the use of lower dosages (*i.e.*, sub-optimal doses) of Cpd 1 or a pharmaceutical composition

thereof or an additional therapy and/or less frequent administration of Cpd 1 or a pharmaceutical composition thereof or an additional therapy to a subject.

In certain aspects, the ability to utilize lower dosages of Cpd 1 or a pharmaceutical composition thereof or of an additional therapy and/or to administer Cpd 1 or a pharmaceutical composition thereof or said additional therapy less frequently reduces the toxicity associated with the administration of Cpd 1 or a pharmaceutical composition thereof or of said additional therapy, respectively, to a subject without reducing the efficacy of Cpd 1 or a pharmaceutical composition thereof or of said additional therapy, respectively, in the treatment of a hematologic cancer.

In some aspects, a synergistic effect results in improved efficacy of Cpd 1 or a pharmaceutical composition thereof and each of said additional therapies in treating a hematologic cancer. In some aspects, a synergistic effect of a combination of Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies avoids or reduces adverse or unwanted side effects associated with the use of any single therapy.

The combination of Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies can be administered to a subject in the same pharmaceutical composition. Alternatively, Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies can be administered concurrently to a subject in separate pharmaceutical compositions. Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies can be administered sequentially to a subject in separate pharmaceutical compositions. Cpd 1 or a pharmaceutical composition thereof and one or more additional therapies may also be administered to a subject by the same or different routes of administration.

The combination therapies provided herein involve administering to a subject in need thereof Cpd 1 or a pharmaceutical composition thereof in combination with conventional, or known, therapies for treating a hematologic cancer. Other therapies for a hematologic cancer or a condition associated therewith are aimed at controlling or relieving one or more symptoms. Accordingly, in some aspects, the combination therapies provided herein involve administering to a subject in need thereof a pain

reliever, or other therapies aimed at alleviating or controlling one or more symptoms associated with a hematologic cancer or a condition associated therewith.

Specific examples of anti-cancer agents that may be used in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include: a hormonal agent (*e.g.*, aromatase inhibitor, selective estrogen receptor modulator (SERM), and estrogen receptor antagonist), chemotherapeutic agent (*e.g.*, microtubule disassembly blocker, antimetabolite, topoisomerase inhibitor, and DNA crosslinker or damaging agent), anti-angiogenic agent (*e.g.*, VEGF antagonist, receptor antagonist, integrin antagonist, vascular targeting agent (VTA)/vascular disrupting agent (VDA)), radiation therapy, and conventional surgery.

Non-limiting examples of hormonal agents that may be used in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include aromatase inhibitors, SERMs, and estrogen receptor antagonists. Hormonal agents that are aromatase inhibitors may be steroidal or nonsteroidal. Non-limiting examples of nonsteroidal hormonal agents include letrozole, anastrozole, aminoglutethimide, fadrozole, and vorozole. Non-limiting examples of steroidal hormonal agents include aromasin (exemestane), formestane, and testolactone. Non-limiting examples of hormonal agents that are SERMs include tamoxifen (branded/marketed as Nolvadex®), afimoxifene, arzoxifene, bazedoxifene, clomifene, femarelle, lasofoxifene, ormeloxifene, raloxifene, and toremifene. Non-limiting examples of hormonal agents that are estrogen receptor antagonists include fulvestrant. Other hormonal agents include but are not limited to abiraterone and lonaprisan.

Non-limiting examples of chemotherapeutic agents that may be used in combination with Cpd 1 or a pharmaceutical composition thereof for treating cancer include microtubule disassembly blocker, antimetabolite, topoisomerase inhibitor, and DNA crosslinker or damaging agent. Chemotherapeutic agents that are microtubule disassembly blockers include, but are not limited to, taxenes (*e.g.*, paclitaxel (branded/marketed as TAXOL®), docetaxel, abraxane, larotaxel, ortataxel, and tesetaxel); epothilones (*e.g.*, ixabepilone); and vinca alkaloids (*e.g.*, vinorelbine, vinblastine, vindesine, and vincristine (branded/marketed as ONCOVIN®)).

Chemotherapeutic agents that are antimetabolites include, but are not limited to, folate antimetabolites (*e.g.*, methotrexate, aminopterin, pemetrexed, raltitrexed); purine antimetabolites (*e.g.*, cladribine, clofarabine, fludarabine, mercaptopurine, pentostatin, thioguanine); pyrimidine antimetabolites (*e.g.*, 5-fluorouracil, capecitabine, gemcitabine (GEMZAR®), cytarabine, decitabine, floxuridine, tegafur); and deoxyribonucleotide antimetabolites (*e.g.*, hydroxyurea).

Chemotherapeutic agents that are topoisomerase inhibitors include, but are not limited to, class I (camptotheca) topoisomerase inhibitors (*e.g.*, topotecan (branded/ marketed as HYCAMTIN®) irinotecan, rubitecan, and belotecan); class II (podophyllum) topoisomerase inhibitors (*e.g.*, etoposide or VP-16, and teniposide); anthracyclines (*e.g.*, doxorubicin, epirubicin, Doxil, aclarubicin, amrubicin, daunorubicin, idarubicin, pirarubicin, valrubicin, and zorubicin); and anthracenediones (*e.g.*, mitoxantrone, and pixantrone).

Chemotherapeutic agents that are DNA crosslinkers (or DNA damaging agents) include, but are not limited to, alkylating agents (*e.g.*, cyclophosphamide, mechlorethamine, ifosfamide (branded/ marketed as IFEX®), trofosfamide, chlorambucil, melphalan, prednimustine, bendamustine, uramustine, estramustine, carmustine (branded/ marketed as BiCNU®), lomustine, semustine, fotemustine, nimustine, ranimustine, streptozocin, busulfan, mannosulfan, treosulfan, carboquone, N,N,N'-triethylenethiophosphoramidate, triaziquone, triethylenemelamine); alkylating-like agents (*e.g.*, carboplatin (branded/ marketed as PARAPLATIN®), cisplatin, oxaliplatin, nedaplatin, triplatin tetranitrate, satraplatin, picoplatin); nonclassical DNA crosslinkers (*e.g.*, procarbazine, dacarbazine, temozolomide (branded/ marketed as TEMODAR®), altretamine, mitobronitol); and intercalating agents (*e.g.*, actinomycin, bleomycin, mitomycin, and plicamycin).

Non-limiting examples of anti-angiogenic agents that may be used in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include VEGF antagonists, receptor antagonists, integrin antagonists (*e.g.*, vitaxin, cilengitide, and S247), and VTAs/VDAs (*e.g.*, fosbretabulin). VEGF antagonists include, but are not to, anti-VEGF antibodies (*e.g.*, bevacizumab (branded/ marketed as

AVASTIN[®]) and ranibizumab (branded/marketed as LUCENTIS[®]), VEGF traps (*e.g.*, aflibercept), VEGF antisense or siRNA or miRNA, and aptamers (*e.g.*, pegaptanib (branded/marketed as MACUGEN[®])). Anti-angiogenic agents that are receptor antagonists include, but are not limited to, antibodies (*e.g.*, ramucirumab) and kinase inhibitors (*e.g.*, sunitinib, sorafenib, cediranib, pazopanib, vandetanib, axitinib, and AG-013958) such as tyrosine kinase inhibitors. Other non-limiting examples of anti-angiogenic agents include ATN-224, anecortave acetate (branded/marketed as RETAANE[®]), microtubule depolymerization inhibitor such as combretastatin A4 prodrug, and protein or protein fragment such as collagen 18 (endostatin).

Non-limiting examples of other therapies that may be administered to a subject in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include:

- (1) a statin such as lovastatin (*e.g.*, branded/marketed as MEVACOR[®]);
- (2) an mTOR inhibitor such as sirolimus which is also known as Rapamycin (*e.g.*, branded/marketed as RAPAMUNE[®]), temsirolimus (*e.g.*, branded/marketed as TORISEL[®]), everolimus (*e.g.*, branded/marketed as AFINITOR[®]), and deforolimus;
- (3) a farnesyltransferase inhibitor agent such as tipifarnib (*e.g.*, branded/marketed as ZARNESTRA[®]);
- (4) an antifibrotic agent such as pirfenidone;
- (5) a pegylated interferon such as PEG-interferon alfa-2b;
- (6) a CNS stimulant such as methylphenidate (branded/marketed as RITALIN[®]);
- (7) a HER-2 antagonist such as anti-HER-2 antibody (*e.g.*, trastuzumab) and kinase inhibitor (*e.g.*, lapatinib);
- (8) an IGF-1 antagonist such as an anti-IGF-1 antibody (*e.g.*, AVE1642 and IMC-A11) or an IGF-1 kinase inhibitor;
- (9) EGFR/HER-1 antagonist such as an anti-EGFR antibody (*e.g.*, cetuximab, panitumumab) or EGFR kinase inhibitor (*e.g.*, erlotinib (*e.g.*, branded/marketed as TARCEVA[®]), gefitinib);
- (10) SRC antagonist such as bosutinib;
- (11) cyclin dependent kinase (CDK) inhibitor such as seliciclib;

- (12) Janus kinase 2 inhibitor such as lestaurtinib;
- (13) proteasome inhibitor such as bortezomib;
- (14) phosphodiesterase inhibitor such as anagrelide;
- (15) inosine monophosphate dehydrogenase inhibitor such as tiazofurine;
- (16) lipoxygenase inhibitor such as masoprocol;
- (17) endothelin antagonist;
- (18) retinoid receptor antagonist such as tretinoin or alitretinoin;
- (19) immune modulator such as lenalidomide, pomalidomide, or thalidomide (*e.g.*, branded/ marketed as THALIDOMID[®]);
- (20) kinase (eg, tyrosine kinase) inhibitor such as imatinib (*e.g.*, branded/ marketed as GLEEVEC[®]), dasatinib, erlotinib, nilotinib, gefitinib, sorafenib, sunitinib (*e.g.*, branded/ marketed as SUTENT[®]), lapatinib, AEE788, or TG100801;
- (21) non-steroidal anti-inflammatory agent such as celecoxib (branded/ marketed as CELEBREX[®]);
- (22) human granulocyte colony-stimulating factor (G-CSF) such as filgrastim (branded/ marketed as NEUPOGEN[®]);
- (23) folinic acid or leucovorin calcium;
- (24) integrin antagonist such as an integrin $\alpha 5 \beta 1$ -antagonist (*e.g.*, JSM6427);
- (25) nuclear factor kappa beta (NF- $\kappa\beta$) antagonist such as OT-551, which is also an anti-oxidant;
- (26) hedgehog inhibitor such as CUR61414, cyclopamine, GDC-0449, or anti-hedgehog antibody;
- (27) histone deacetylase (HDAC) inhibitor such as SAHA (also known as vorinostat (branded/ marketed as ZOLINZA[®])), PCI-24781, SB939, CHR-3996, CRA-024781, ITF2357, JNJ-26481585, or PCI-24781;
- (28) retinoid such as isotretinoin (*e.g.*, branded/ marketed as ACCUTANE[®]);
- (29) hepatocyte growth factor/scatter factor (HGF/SF) antagonist such as HGF/SF monoclonal antibody (*e.g.*, AMG 102);
- (30) synthetic chemical such as antineoplaston;
- (31) anti-diabetic such as rosiglitazone maleate (*e.g.*, branded/ marketed as AVANDIA[®]);

- (32) antimalarial and amebicidal drug such as chloroquine (*e.g.*, branded/marketed as ARALEN[®]);
- (33) synthetic bradykinin such as RMP-7;
- (34) platelet-derived growth factor receptor inhibitor such as SU-101;
- (35) receptor tyrosine kinase inhibitors of Flk-1/KDR/VEGFR2, FGFR1 and PDGFR beta such as SU5416 and SU6668;
- (36) anti-inflammatory agent such as sulfasalazine (*e.g.*, branded/marketed as AZULFIDINE[®]); and
- (37) TGF-beta antisense therapy.

Non-limiting examples of other therapies that may be administered to a subject in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include: a synthetic nonapeptide analog of naturally occurring gonadotropin releasing hormone such as leuprolide acetate (branded/marketed as LUPRON[®]); a nonsteroidal, anti-androgen such as flutamide (branded/marketed as EULEXIN[®]) or nilutamide (branded/marketed as NILANDRON[®]); a non-steroidal androgen receptor inhibitor such as bicalutamide (branded/marketed as CASODEX[®]); steroid hormone such as progesterone; anti-fungal agent such as Ketoconazole (branded/marketed as NIZORAL[®]); glucocorticoid such as prednisone; estramustine phosphate sodium (branded/marketed as EMCYT[®]); and bisphosphonate such as pamidronate, alendronate, and risedronate.

Additional specific examples of therapies that may be used in combination with Cpd 1 or a pharmaceutical composition thereof for treating a hematologic cancer include, but are not limited to, agents associated with cancer immunotherapy (*e.g.*, cytokines, interleukins, and cancer vaccines).

Specific examples of agents alleviating side-effects associated with a hematologic cancer that can be used as therapies in combination with Cpd 1 or a pharmaceutical composition thereof, include, but are not limited to: antiemetics, *e.g.*, Ondansetron hydrochloride (branded/marketed as Zofran[®]), Granisetron hydrochloride (branded/marketed as Kytril[®]), Lorazepam (branded/marketed as Ativan[®]) and Dexamethasone (branded/marketed as Decadron[®]).

In certain aspects, combination therapies provided herein for treating a hematologic cancer comprise administering Cpd 1 or a pharmaceutical composition thereof in combination with one or more agents used to treat and/or manage a side effect, such as, bleeding (usually transient, low-grade epistaxis), arterial and venous thrombosis, hypertension, delayed wound healing, asymptomatic proteinuria, nasal septal perforation, reversible posterior leukoencephalopathy syndrome in association with hypertension, light-headedness, ataxia, headache, hoarseness, nausea, vomiting, diarrhea, rash, subungual hemorrhage, myelodysplastic syndromes, myelosuppression, fatigue, hypothyroidism, QT interval prolongation, or heart failure.

In certain aspects, Cpd 1 or a pharmaceutical composition thereof is not used in combination with a drug that is primarily metabolized by CYP2D6 (such as an antidepressant (*e.g.*, a tricyclic antidepressant, a selective serotonin reuptake inhibitor, and the like), an antipsychotic, a beta-adrenergic receptor blocker, or certain types of anti-arrhythmics) to treat a hematologic cancer.

Kits

Provided herein is a pharmaceutical pack or kit comprising one or more containers filled with Cpd 1 or a pharmaceutical composition thereof. Additionally, one or more other therapies useful for the treatment of a hematologic cancer, or other relevant agents can also be included in the pharmaceutical pack or kit. Also provided herein is a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions described herein. Optionally associated with such kits can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Examples

General Methods. All culture media and supplement agents were purchased from (Gibco BRL, Invitrogen). Tumor cell lines purchased from ATCC were maintained in DMEM (adherent cells) or RPMI-1640 (suspension cells) containing 1g/L glucose, supplemented with 10% fetal bovine serum (FBS), penicillin (50 IU/mL), and

streptomycin (50 µg/mL). Cells were cultured at 37°C in a humidified atmosphere containing 5% CO₂ and 21% O₂ in air (normoxia). For hypoxic exposure, cells were placed in a hypoxia incubator (NAPCO 8000 WJ) with a mixture of gas consisting of 1% O₂, 5% CO₂, and 94% N₂. All primary cells were purchased from AllCells, Inc and cultured in RPMI-1640 (for peripheral blood mononuclear cells, PBMC) or DMEM (for adherent cells) supplemented with 10% FBS, penicillin (50 IU/mL), and streptomycin (50 µg/mL).

Western Blot Analysis. Cells were treated with test compounds for 24 to 48 hours as indicated in the Results section. Subsequently, the conditioned medium was harvested by mixing with equal volume of 2x SDS-PAGE sample buffer (Bio-Rad). Whole-cell extracts were obtained by lysing cells either directly in 1x SDS-PAGE sample buffer (Bio-Rad) or in M-PER lysis buffer containing 150 mmol/L NaCl, 2 mmol/L EDTA and 1x Halt protease inhibitors (Pierce, Rockford, IL). Proteins in the samples were resolved on a Tris-glycine Criterion gel (Bio-Rad) and transferred onto a 0.45 µm nitrocellulose membrane. The membranes were then immunoblotted with specific antibodies indicated in the Results section. Antibody (C1) specific for VEGFA was purchased from Santa Cruz Biotechnologies, CA (1:200 dilution); Antibody specific for human β-actin was purchased from Abcam (1:10000 dilution); Antibody specific for V5-tag was purchased from Invitrogen (1:5000 dilution). Antibody specific for DHODH was purchased from Proteintech (Cat#: 14877-1-AP, 1:500 dilution); and antibody specific for human prohibitin was purchased from Thermo Scientific (Cat#: PA5-12274 and PA5-14133, 1:1000 dilution). Immunodetection was done using the corresponding secondary antibodies conjugated with infrared dyes or horseradish peroxidase. The expression levels of proteins were detected with Odyssey (LI-COR) or using enhanced chemiluminescence (Pierce, Rockford, IL).

Determination of ELISA EC₅₀ and Cytotoxicity CC₅₀: All ELISAs were performed using commercially available ELISA kits (R&D Systems) according to the manufacturer's instructions. The screen of 240 cell lines were performed in Crown Bioscience (Shanghai, China). Cells were treated with Cpd 1 in a serial of doses for 72 hours and the inhibition of cell proliferation was determined using a standard assay, CellTiter-Glo® Luminescent Cell Viability Assay (Promega, Madison, WI) that measures

total cellular adenosine triphosphate (ATP) concentrations as an indicator of cell viability. Data generated from ELISA or cytotoxicity studies were plotted with Prism software, with the Y-axis typically representing the % inhibition and the X-axis representing log compound concentration. A sigmoid dose response with a variable slope regression curve was generated for each compound. Maximal inhibition was set at 100% and minimal at 0%. EC₅₀ and CC₅₀ values were automatically calculated after curve fitting using the Prism software.

Human Tumor Xenograft Studies. All studies involving animals were performed in strict accordance with guidelines promulgated by the American Association for Accreditation of Laboratory Animal Care with the oversight of animal use and care committees.

Human Xenograft Model: HT1080 or MOLM-13 tumor cells in log phase growth (5-10 x 10⁶ cells/mouse, depending on cell lines) were implanted subcutaneously in male athymic nude mice. When the average tumor size reached the indicated size, mice were randomly divided into groups (10-15 mice/group) and administered vehicle or test compounds orally, once a day or twice per day as indicated. Tumor size was measured using calipers at periodic intervals during the study and tumor volume was calculated as $L \times W^2/2$, in which L represents the longest and W is the shortest diameter of the tumors. At the end of the experiments, tumors were excised and homogenized on ice in Tris-HCl buffer containing a cocktail of proteinase inhibitors using a Powergen homogenizer fitted with Omni-Tip disposable/reusable probes (both from Fisher Scientific). Intratumoral levels of growth factors and proteins were measured using commercially available ELISA kits (R&D Systems). Protein concentrations of the homogenates were measured (Bio-Rad protein assay kit) and intratumoral growth factor levels normalized to the total protein concentration.

Systemic Leukemia Lethality Model: Male NOD-SCID mice were inoculated with MOLT-4 human ALL tumor cells (1 x 10⁷ cells in 200 µL PBS) by intravenous (IV) injection. Eight days after tumor inoculation, mice were randomized into two groups (10 mice per groups) and treated with either vehicle or Cpd 1 (10 mg/kg qd). The mice were dosed and observed until moribund, at which time they were euthanized. At 2- and

4-weeks post-inoculation, whole blood was obtained from 5 mice per group by retro-orbital bleeding, stained with an antibody against human CD45, and the number of CD45-positive cells was determined by FACs. Data in Figure 5b represents % of human CD45+ from blood of each mouse; where * indicates $p < 0.05$, one way ANOVA, with multiple comparisons vs vehicle.

Quantification of Pyrimidine Nucleotide Synthesis Metabolites in Cpd 1 Treated

Cells. HT1080 cells in log phase growth were seeded in 10 cm dishes (4×10^6 cells/dish) and cultured in regular DMEM supplemented with 10% fetal bovine serum (FBS), penicillin (50 IU/mL), and streptomycin (50 μ g/mL). After overnight incubation, the culture medium was replenished with glutamine free DMEM containing 1g/L glucose, 10% fetal bovine serum (FBS), penicillin (50 IU/mL), streptomycin (50 μ g/mL) and 1 mM 15 N-glutamine (Sigma, Cat#: 490024). After culturing for the indicated time in the presence of compounds or vehicle control (0.5% DMSO), the cells were washed once with cold PBS (5 mL) and harvested and lysed in cold distilled H₂O (0.5 mL) with -80 °C methanol (1 mL) using a plastic cell scraper. The cell lysates were then centrifuged at 10,000 x g for 15 minutes at 4°C, and supernatant collected for LC-MS/MS analysis of the 15 N-labelled de novo pyrimidine synthesis metabolites.

LC-MS/MS Detection of Metabolites. Quantification of 15 N-labelled de novo pyrimidine synthesis metabolites was carried out on an Accela pump and a PAL auto-sampler coupled to a TSQ Quantum Ultra mass spectrometer. The mass spectrometer was equipped with a heated electrospray ionization source operated in negative-ion mode (Thermo Fisher Scientific, Waltham, MA). The ion spray voltage was set at 2500 V, capillary temperature at 350 °C, vaporizer temperature at 300 °C; sheath gas pressure at 50 units and auxiliary gas pressure at 5 units. A Thermo Fisher Scientific Hypercab column (3 μ , 50 x 2.1 mm) was used and maintained at 50 °C for metabolite separation. Mobile phase A was 10 mM NH₄HCO₃ in water, pH 9.4, and mobile phase B was 10 mM NH₄HCO₃ in ACN-water (9:1). The flow rate was set at 0.50 mL/min. Ion transitions monitored were at m/z 156.0 $\rightarrow$ m/z 112.0 for 15 N-orotic acid, m/z 158.0 $\rightarrow$ m/z 114.0 for 15 N-dihydrorotic acid, m/z 324.0 $\rightarrow$ m/z 79.0 for 15 N-ump.

Conjugation of Cpd 1 *S*- and *R*-Enantiomer to Sepharose-6B beads. Sepharose 6B-Cpd 1: Epoxy-activated Sepharose 6B (GE healthcare cat. # 17-0480-01, 1g) was stirred in water (20 mL) for 2h at RT, then filtered. The wet Sepharose 6B was diluted with pH 11 phosphate/NaOH buffer (5 mL). To a solution of 4-chlorophenyl (*S*)-6-chloro-1-(4-methoxyphenyl)-1,3,4,9-tetrahydro-2H-pyrido[3,4-*b*]indole-2-carboxylate (1 mmole) in DMF (12 mL) at 40 °C was added Sepharose-6B in phosphate/NaOH buffer. The reaction mixture was stirred further 24 hours at 40 °C. The Sepharose 6B-Cpd 1 product was filtered and washed with DMF (100 mL), water (100 mL), pH 4 buffer (100 mL), pH 11 buffer (100 mL), water (100 mL), and 0.5M NaCl/water. The Sepharose-6B-Cpd 1 *R*-enantiomer was prepared using 4-chlorophenyl (*R*)-6-chloro-1-(4-methoxyphenyl)-1,3,4,9-tetrahydro-2H-pyrido[3,4-*b*]indole-2-carboxylate and the preceding reaction procedures.

Pull-down of Proteins Bound to Cpd 1-beads. Cells in log phase growth were washed with cold PBS once and then collected by trypsin digestion. After centrifugation at 1200 rpm for 5 minutes, the cells were re-suspended in lysis buffer (PBS + 0.5% Triton X100 + 1x proteinase inhibitors (Halt, Roche, Cat#: 78440) and incubated on ice for 10 minutes; supernatants were collected after passing cells through a 22-gauge needle 10 times and centrifuging at 13,000 rpm for 15 min to remove cell debris. Subsequently, 1 mL of cell lysate solution (2 mg/mL protein) was added to each reaction tube containing 25 μ L (packed volume) Cpd 1 *S*-enantiomer or control *R*-enantiomer beads and incubated for 2 hours at 4°C with gentle agitation on a rotator; after three times wash with washing Buffer (1 mL) for 5 minutes each, the protein bound with beads were eluted with indicated buffer as shown in the result section. Proteins in the eluted samples were identified via LC-MS/MS or western blot analysis.

Mitochondria Isolated from Cultured K562 Cells as Source of DHODH for In Vitro Enzyme Inhibition Study. Mitochondria were isolated from K562 cells in log phase growth with a Dounce homogenizer as reported [27]. Briefly, about 2×10^9 K562 cells were collected by centrifugation, re-suspended in 11 mL of ice-cold RSB hypo buffer (10 mM NaCl, 1.5 mM MgCl₂, 10 mM Tris-HCl, pH 7.5) in a 15-mL Dounce homogenizer. After swelling for 5–10 min on ice, the swollen cells were lysed with 10 strokes of the B

pestle. A volume of 8 mL of 2.5× MS homogenization buffer (525 mM mannitol, 175 mM sucrose, 12.5 mM Tris-HCl, 2.5 mM EDTA, pH 7.5) was immediately added to the cell lysate and mixed to give a final concentration of 1× MS homogenization buffer. The homogenate was centrifuged at 1300g for 5 min to remove nuclei, unlysed cells, and large membrane fragments. The mitochondria in the supernatant were pelleted at 12,000g for 15 min and re-suspended in 1 mL 1× MS homogenization buffer for analysis of downstream DHODH activity as described below.

In Vitro DHODH Activity Assays. The chromogen reduction assay was carried out as previously reported [28]. DHODH activity was determined in the presence of compounds or vehicle control using the standard colorimetric DHODH continuous assay in which the oxidation of dihydroorotic acid (DHO) and subsequent reduction of ubiquinone is coupled with and measured by monitoring the reduction of 2,6-dichlorophenolindophenol (DCPIP). Briefly, full length purified recombinant human DHODH or mitochondria isolated from K562 cells were used as enzyme sources to test compound inhibition of DHODH activity. Enzymatic assays were conducted at 25 °C in a reaction buffer containing HEPES buffer (100 mM, pH 8.0), 150 mM NaCl, 5% glycerol and 0.05% Triton X-100, 200 μM L-dihydroorotate, 20 μM QD, 100 μM DCIP. The total volume of each reaction mixture was 100 μl containing 10 nM DHODH or an amount of isolated mitochondria yielding a similar level of DHODH activity. Compounds were prepared as 200 x stock solutions in DMSO and the final concentration of DMSO was 0.5%. The DCIP reduction was monitored by measuring the absorbance at OD₆₁₀ nm on a BioTek Power Wave XS2 and the enzyme reaction velocity was derived for each reaction using the GEN5 software.

Statistical Analysis. Data are given as the mean ± SD or SEM as indicated for quantitative experiments. For statistical analysis, p-values were derived using unpaired Student's *t*-tests for any study with only two groups presented. Otherwise, comparisons of groups were performed on log-transformed data using a one way ANOVA test. All analyses were made using GraphPad Prism Software.

In vitro Combination Assay with HL-60 cells. HL-60 cells (human promyelocytic leukemia cells) were treated for 72 hours in the presence of Cpd 1 (DHODH inhibitor)

and enasidenib (mutant IDH2 inhibitor IDHIFA[®]) alone and in combination. HL-60 cells are reported to express the wild-type IDH2 gene/protein. Concentrations of Cpd 1 tested included: 0, 1.4, 4.1, 12.3, 37, 111, 333 and 1000 nM. Concentrations of enasidenib were tested at: 0, 0.1, 0.3, 1.3, 6.4, 32, 160, 800, 4,000 and 20,000 nM. The combinations were evaluated in a full checkerboard titration. The extent of proliferation at 72 hours was determined by addition of Cell Titer Glo, and the percent inhibition calculated as referenced to vehicle/DMSO treated HL-60 cells.

Results. At all concentrations tested, enasidenib had no significant antiproliferative effect as single agent (see Figure 9 with Cpd 1 nil concentration). Cpd 1 at a single agent test concentration of 111 nM had approximately 20% proliferative inhibition. However, the combination of enasidenib with Cpd 1, each at the two highest test concentrations of 4000:333 and 20000:1000 nM, respectively, demonstrated significant synergistic activity.

Conclusion. An inhibitor of DHODH function such as Cpd 1 in combination with an inhibitor of mutant IDH2 enzyme activity such as enasidenib may have utility in treating mutant and wild-type IDH2 leukemias.

In Vitro Combination Assays with Primary Leukemia Blood Samples. Blood samples from 11 leukemia subjects (10 AML and one ALL) were shipped overnight from clinical investigation sites. Upon arrival, red blood cells were lysed and the remaining cells counted and resuspended at the appropriate concentration in serum free media supplemented with cytokines. The samples were plated in 384-well microtiter plates and treated with the Cpd 1 *S*-enantiomer or *R*-enantiomer in triplicate. After incubation at 37 °C for 72 hours, the samples were stained with antibodies (antibody panel details below) and evaluated using an Intellicyt iQue Plus flow cytometer. A single concentration of Cpd 1 was tested (10 nM for cells from five AML subjects, or 20 nM for cells from the other five AML subjects and one ALL subject). A single concentration of enasidenib was tested (4000 nM) across cells from all 11 subjects.

Leukemia cells were quantified using the following FACS staining cell panel: DAPI, CD33, CD34, c-kit, CD3/CD19, CD66b, CD14, CD38. Live cells were gated using

FSC/SSC and DAPI exclusion, and then further defined by cell surface marker expression.

Blood cells from subject 332AML harbored an oncogenic mutant IDH2 allele. Subjects 328AML, 333AML, 342AML, 347AML, and 351AML did not express an oncogenic mutant IDH2 allele. Genotype data was not available for the other 6 subjects.

For each subject an interaction score was calculated as: Measured minus Expected.

Measured = fraction of blasts after treatment with Cpd 1 and enasidenib.

Expected = fraction of blasts after treatment with Cpd 1 multiplied by the fraction of blasts after treatment with enasidenib as monotherapy.

Results. Table 5 summarizes the effect of Cpd 1, enasidenib, and the combination against 11 leukemia subject samples. At 4000 nM, enasidenib did not have activity as a monotherapy against any of the subject samples. Cpd 1 significantly reduced proliferation in 10/11 samples (AML347 did not respond to Cpd 1). Further, the addition of enasidenib enhanced the activity of Cpd 1 in 10/11 samples (no further increase in inhibition was seen with the addition of enasidenib in sample 356). In some samples, the additional of enasidenib only marginally increased the anti-proliferative effect of Cpd 1 but in other samples a more pronounced effect was seen (for example, in subject sample 369). All samples were from AML subjects except sample 340 from an ALL subject. Abbreviations: ND= no data; mut = mutant; wt = wildtype

Figure 10 shows data from two representative samples, demonstrating that enasidenib enhanced the anti-proliferative effect of Cpd 1 in an AML subject (Figure 10a) and an ALL subject (Figure 10a).

Table 5. % Inhibition of Total Blasts; Effect of Cpd 1, enasidenib, or the Combination on Primary Leukemia Samples

Sample ID	328	332	333	340	342	347	351	356	359	369	371
IDH2 mutational status	wt	mut	wt	ND	wt	wt	Wt	ND	ND	ND	ND
Cpd 1	76.6	41.5	70	24	91.9	-11.6	28.9	47	75.7	49.7	52.9
Enasidenib	1.7	-10.9	-36.6	-6.5	6.4	-8.9	8.8	26.1	-24.1	-0.4	-7.2

Combination	94.4	58	82.9	39.7	93.8	11.3	41.1	44.1	81.6	70.7	64.4
Interaction Coefficient	-0.17	-0.23	-0.24	-0.21	-0.01	-0.33	-0.06	0.17	-0.12	-0.21	-0.15

Conclusion. An inhibitor of DHODH function such as Cpd 1 in combination with an inhibitor of mutant IDH2 function such as enasidenib has demonstrated synergistic activity which may have utility in treating mutant and wild-type IDH2 leukemia.

Reference List

- [1] Rawls J, Knecht W, Diekert K, Lill R, Löffler M. Requirements for the mitochondrial import and localization of dihydroorotate dehydrogenase. *Eur J Biochem* 2000 Apr;267(7):2079-87.
- [2] Zhang X, Yang J, Chen M, Li L, Huan F, Li A, et al. Metabolomics profiles delineate uridine deficiency contributes to mitochondria-mediated apoptosis induced by celastrol in human acute promyelocytic leukemia cells. *Oncotarget* 2016 Jul 19;7(29):46557-72.
- [3] Baumann P, Mandl-Weber S, Volkl A, Adam C, Bumeder I, Oduncu F, et al. Dihydroorotate dehydrogenase inhibitor A771726 (leflunomide) induces apoptosis and diminishes proliferation of multiple myeloma cells. *Mol Cancer Ther* 2009 Feb;8(2):366-75.
- [4] Sykes DB, Kfoury YS, Mercier FE, Wawer MJ, Law JM, Haynes MK, et al. Inhibition of Dihydroorotate Dehydrogenase Overcomes Differentiation Blockade in Acute Myeloid Leukemia. *Cell* 2016 Sep 22;167(1):171-86.
- [5] Cacciamani T, Vita A, Cristalli G, Vincenzetti S, Natalini P, Ruggieri S, et al. Purification of human cytidine deaminase: molecular and enzymatic characterization and inhibition by synthetic pyrimidine analogs. *Arch Biochem Biophys* 1991 Nov 1;290(2):285-92.
- [6] Peters GJ, Schwartzmann G, Nadal JC, Laurensse EJ, van Groenigen CJ, van der Vijgh WJ, et al. In vivo inhibition of the pyrimidine de novo enzyme dihydroorotic acid dehydrogenase by brequinar sodium (DUP-785; NSC 368390) in mice and patients. *Cancer Res* 1990 Aug 1;50(15):4644-9.
- [7] Ohnuma T, Roboz J, Shapiro ML, Holland JF. Pharmacological and biochemical effects of pyrazofurin in humans. *Cancer Res* 1977 Jul;37(7 Pt 1):2043-9.
- [8] McLean LR, Zhang Y, Degnen W, Peppard J, Cabel D, Zou C, et al. Discovery of novel inhibitors for DHODH via virtual screening and X-ray crystallographic structures. *Bioorg Med Chem Lett* 2010 Mar 15;20(6):1981-4.
- [9] Jain AN. Effects of protein conformation in docking: improved pose prediction through protein pocket adaptation. *J Comput Aided Mol Des* 2009 Jun;23(6):355-74.
- [10] Jain AN. Surflex: fully automatic flexible molecular docking using a molecular similarity-based search engine. *J Med Chem* 2003 Feb 13;46(4):499-511.

- [11] Lane AN, Fan TW. Regulation of mammalian nucleotide metabolism and biosynthesis. *Nucleic Acids Res* 2015 Feb 27;43(4):2466-85.
- [12] Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature* 2012 Mar 28;483(7391):603-7.
- [13] Baumann P, Mandl-Weber S, Volkl A, Adam C, Bumeder I, Oduncu F, et al. Dihydroorotate dehydrogenase inhibitor A771726 (leflunomide) induces apoptosis and diminishes proliferation of multiple myeloma cells. *Mol Cancer Ther* 2009 Feb;8(2):366-75.
- [14] Fitzpatrick LR, Small JS, Doblhofer R, Ammendola A. Vidofludimus inhibits colonic interleukin-17 and improves hapten-induced colitis in rats by a unique dual mode of action. *J Pharmacol Exp Ther* 2012 Sep;342(3):850-60.
- [15] Wang X, Harrison JS, Studzinski GP. Enhancement of arabinocytosine (AraC) toxicity to AML cells by a differentiation agent combination. *J Steroid Biochem Mol Biol* 2016 Nov;164:72-8.
- [16] Cunningham JT, Moreno MV, Lodi A, Ronen SM, Ruggero D. Protein and nucleotide biosynthesis are coupled by a single rate-limiting enzyme, PRPS2, to drive cancer. *Cell* 2014 May 22;157(5):1088-103.
- [17] Mohamad Fairus AK, Choudhary B, Hosahalli S, Kavitha N, Shatrah O. Dihydroorotate dehydrogenase (DHODH) inhibitors affect ATP depletion, endogenous ROS and mediate S-phase arrest in breast cancer cells. *Biochimie* 2017 Apr;135:154-63.
- [18] Saini AK, Nanda JS, Lorsch JR, Hinnebusch AG. Regulatory elements in eIF1A control the fidelity of start codon selection by modulating tRNA(i)(Met) binding to the ribosome. *Genes Dev* 2010 Jan 1;24(1):97-110.
- [19] Carlile TM, Rojas-Duran MF, Zinshteyn B, Shin H, Bartoli KM, Gilbert WV. Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. *Nature* 2014 Nov 6;515(7525):143-6.
- [20] Cody R, Stewart D, DeForni M, Moore M, Dallaire B, Azarnia N, et al. Multicenter phase II study of brequinar sodium in patients with advanced breast cancer. *Am J Clin Oncol* 1993 Dec;16(6):526-8.
- [21] Xu X, Williams JW, Shen J, Gong H, Yin DP, Blinder L, et al. In vitro and in vivo mechanisms of action of the antiproliferative and immunosuppressive agent, brequinar sodium. *J Immunol* 1998 Jan 15;160(2):846-53.
- [22] Xu X, Williams JW, Gong H, Finnegan A, Chong AS. Two activities of the immunosuppressive metabolite of leflunomide, A77 1726. Inhibition of pyrimidine nucleotide synthesis and protein tyrosine phosphorylation. *Biochem Pharmacol* 1996 Aug 23;52(4):527-34.
- [23] Deans RM, Morgens DW, Okesli A, Pillay S, Horlbeck MA, Kampmann M, et al. Parallel shRNA and CRISPR-Cas9 screens enable antiviral drug target identification. *Nat Chem Biol* 2016 May;12(5):361-6.


- [24] Packer RJ, Rood BR, Turner DC, Stewart CF, Fisher M, Smith C, et al. Phase I and pharmacokinetic trial of PTC299 in pediatric patients with refractory or recurrent central nervous system tumors: a PBTC study. *J Neurooncol* 2015 Jan;121(1):217-24.
- [25] Basso LGM, Mendes LFS, Costa-Filho AJ. The two sides of a lipid-protein story. *Biophys Rev* 2016 Jun;8(2):179-91.
- [26] Lewis TA, Sykes DB, Law JM, Munoz B, Rustiguel JK, Nonato MC, et al. Development of ML390: A Human DHODH Inhibitor That Induces Differentiation in Acute Myeloid Leukemia. *ACS Med Chem Lett* 2016 Dec 8;7(12):1112-7.
- [27] Clayton DA, Shadel GS. Isolation of mitochondria from cells and tissues. *Cold Spring Harb Protoc* 2014 Oct 1;2014(10):db.
- [28] Chen SF, Perrella FW, Behrens DL, Papp LM. Inhibition of dihydroorotate dehydrogenase activity by brequinar sodium. *Cancer Res* 1992 Jul 1;52(13):3521-7.
- [29] Yen K, Travins J, Wang F, David MD, Artin E, Straley K, Padyana A, Gross S, DeLaBarre B, Tobin E, Chen Y, Nagaraja R, Choe S, Jin L, Konteatis Z, Cianchetta G, Saunders JO, Salituro FG, Quivoron C, Opolon P, Bawa O, Saada V, Paci A, Broutin S, Bernard OA, de Botton S, Marteyn BS, Pilichowska M, Xu Y, Fang C, Jiang F, Wei W, Jin S, Silverman L, Liu W, Yang H, Dang L, Dorsch M, Penard-Lacronique V, Biller SA, Su SM. AG-221, a First-in-Class Therapy Targeting Acute Myeloid Leukemia Harboring Oncogenic IDH2 Mutations. *Cancer Discov*. 2017 May;7(5):478-493.
- [30] Nassereddine S, Lap CJ, Haroun F, Tabbara I. The role of mutant IDH1 and IDH2 inhibitors in the treatment of acute myeloid leukemia. *Ann Hematol*. 2017 Dec;96(12):1983-1991.
- [31] Amatangelo MD, Quek L, Shih A, Stein EM, Roshal M, David MD, Marteyn B, Famoud NR, de Botton S, Bernard OA, Wu B, Yen KE, Tallman MS, Papaemmanuil E, Penard-Lacronique V, Thakurta A, Vyas P, Levine RL; Enasidenib induces acute myeloid leukemia cell differentiation to promote clinical response; *Blood*, 10Aug2017, 130:732-741.

Without regard to whether a document cited herein was specifically and individually indicated as being incorporated by reference, all documents referred to herein are incorporated by reference into the present application for any and all purposes to the same extent as if each individual reference was fully set forth herein.

Having now fully described the subject matter of the claims, it will be understood by those having ordinary skill in the art that the same can be performed within a wide range of equivalents without affecting the scope of the subject matter or aspects described herein. It is intended that the appended claims be interpreted to include all such equivalents.

What is claimed is:

1. A method of use for a 4-chlorophenyl (S)-6-chloro-1-(4-methoxyphenyl)-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indole-2-carboxylate compound or a form or pharmaceutical composition thereof for preventing, treating or ameliorating a hematologic cancer characterized by reduced expression of uridine salvage enzymes and dependence on de novo pyrimidine nucleotide synthesis in a subject in need thereof comprises, administering to the subject an effective amount of the compound, having the structure:



or a form or pharmaceutical composition thereof.

2. The method of claim 1, wherein the hematological cancer is a leukemia.
3. The method of claim 2, wherein the leukemia is selected from acute lymphoblastic leukemia, acute myeloid leukemia, acute nonlymphocytic leukemia, chronic lymphocytic leukemia or chronic myeloid leukemia.
4. The method of claim 1, wherein the effective amount of the compound is an initial dose of about 50 mg administered orally once per day, then a loading dose of about 100 mg administered orally once or twice per day, followed by a maintenance dose of about 50 mg administered orally once or twice per day.
5. The method of claim 4, wherein the initial dose is administered for one day, the loading dose is administered once or twice per day, and the maintenance dose is once per day.
6. The method of claim 5, wherein the loading dose is administered once per day.
7. The method of claim 1, wherein the effective amount of the compound is an initial dose of about 20 mg administered orally once per day, then a loading dose of about 40

mg administered orally once or twice per day, followed by a maintenance dose of about 20 mg administered orally once or twice per day.

8. The method of claim 7, wherein the initial dose is administered for one day, the loading dose is administered once or twice per day, and the maintenance dose is once per day.

9. The method of claim 8, wherein the loading dose is administered once per day.

10. The method of claim 1, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 in a 24 hour period in a range of from approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ to approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$, and the like, or any range in between.

11. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 4 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

12. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 6 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

13. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 8 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

14. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 10 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

15. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 12 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

16. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 14 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

17. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 16 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

18. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 18 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

19. The method of claim 10, wherein the effective amount is a dosage that achieves a mean plasma concentration of Cpd 1 of approximately 20 $\mu\text{g}\cdot\text{hr}/\text{mL}$ in a 24 hour period.

Figure 1.






Figure 2.

2a.



2b.



2c.





Figure 3.

3a.



3b.






Figure 4.


4a.




4b.



4c.



4d.



4e.






Figure 5

5a.



5b.



5c.




Figure 6




Figure 7




Figure 8




Figure 9





Figure 10

10a.



10b.





Figure 11



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/44862

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 31/437; A61P 35/02; C07D 471/04 (2018.01)

CPC - A61K 31/437; C07D 471/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2012/0202763 A1 (ALMSTEAD, N et al.) 09 August 2012; figure 1; paragraphs [0059], [0068], [0225] - table 1, compound #10, page 199, [0253], [0264], [0324], [0336], [0344], [0347], [0353]	1-3, 10-19
Y	~ FIEDLER, W et al., An open-label, Phase I study of cediranib (RECENTIN) in patients with acute myeloid leukemia, Leukemia Research 34(2), pages 196-202, 2010; abstract only	1-3, 10-19
A	US 2005/0276808 A1 (CEDARBAUM, JM) 15 December 2005; paragraphs [0004], [0024]; claims 1-2 & 5	4-9
A	US 2017/0143712 A1 (ACERTA PHARMA B.V.) 25 May 2017; paragraphs [0030], [0061], [0073], [0111]	4-9

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 September 2018 (10.09.2018)

Date of mailing of the international search report

25 SEP 2018

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450
Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774