



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2106277 C 2005/08/09

(11)(21) **2 106 277**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 1993/09/15

(41) Mise à la disp. pub./Open to Public Insp.: 1995/03/16

(45) Date de délivrance/Issue Date: 2005/08/09

(51) Cl.Int.⁵/Int.Cl.⁵ C07C 281/14, A01N 47/34

(72) Inventeurs/Inventors:

ANDERSON, RICHARD JAMES, US;
CLOUDSDALE, IAN STUART, US;
HARR, JOST, CH;
LAMOREAUX, ROBERT JAMES, US;
SCHAEFER, KRISTINE JOYCE PETERSON, US

(73) Propriétaire/Owner:

BASF AKTIENGESELLSCHAFT, DE

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : NOUVELLES COMPOSITIONS CONTENANT UN INHIBITEUR DE TRANSPORT D'AUXINE ET D'UN AUTRE
HERBICIDE

(54) Title: NOVEL COMPOSITIONS CONTAINING AN AUXIN TRANSPORT INHIBITOR AND ANOTHER HERBICIDE

(57) Abrégé/Abstract:

The present invention is directed to a herbicidal composition comprising a herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor and at least one other herbicide, wherein the auxin transport inhibitor is present in an amount producing a potentiating effect and wherein the weight ratio of the auxin transport inhibitor to the other herbicide is 1:2000 to 100:1.



Abstract of the Disclosure


The present invention is directed to a herbicidal composition comprising a herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor and at least one other herbicide, wherein the auxin transport inhibitor is present in an amount producing a potentiating effect and wherein the weight ratio of the auxin transport inhibitor to the other herbicide is 1:2000 to 100:1.

-2-

NOVEL COMPOSITIONS CONTAINING AN AUXIN TRANSPORT INHIBITOR
AND ANOTHER HERBICIDE

The present invention concerns the use of auxin transport inhibitors as potentiators or enhancers of herbicides, as well as co-application of such auxin transport inhibitors and herbicides, compositions containing at least one auxin transport inhibitor in combination with at least one herbicide and use of these in combatting or controlling undesired plant growth and in plant growth regulation.



Auxin transport inhibitors are compounds which themselves are herbicides and act by inhibiting transmembrane movement of auxin which accumulates in the cells and affects plant growth. Examples of auxin transport inhibitors are e.g. naptalam, TIBA (2,3,5-triiodobenzoic acid) and DPX 1840 [3,3a-dihydro-2-(p-methoxyphenyl)-8H-pyrazolo [5,1-a]isoindol-8-one] (cf E.M. Beyer, Jr., Plant Physiol., 50, 322 (1972), E.M. Beyer, Jr. et al., Plant Physiol., 57, 839 (1976)) and semicarbazones such as described in US Patents 5,098,462 and 5,098,466 and EP-A-219451. Especially preferred auxin transport inhibitors in the practice of the invention are compounds of formula A




wherein,

X and Y represent independently, hydrogen, fluorine or chlorine, and

R is the group



, or



wherein Z is hydrogen, fluorine or chlorine and M is hydrogen, or a salt forming moiety e.g. an alkali metal cation or an optionally substituted ammonium cation.

Compounds of formula A are generally disclosed, e.g. in US Patents 5,098,462 and 5,098,466 and in European Patent Application 219451, as well as processes for their production, their use as herbicides and plant growth regulators and herbicidal and plant growth regulating compositions containing them. These patents make no reference to the specific compound group of formula A or its potentiating activity.

The term herbicides, as used herein, refers to compounds which combat or control undesired plant growth. This class of compounds may be divided into sub-classes according to the primary type or mode of action the herbicide has on the plant. For example according to G.F. Warren of Purdue University, Indiana, USA, herbicides can be classified as auxin transport inhibitors, growth regulator herbicides, photosynthesis inhibitors, pigment inhibitors, growth inhibitors, amino acid synthesis inhibitors, lipid biosynthesis inhibitors, cell wall biosynthesis inhibitors, rapid cell membrane disruptors as well as "miscellaneous" herbicides which do not come under one of the preceding categories. (Growth regulator herbicides include, e.g. auxin agonists.)

In accordance with the present invention it has now surprisingly been found that auxin transport inhibitors, which are usually highly active herbicides in their own right, potentiate the activity of other herbicides on co-application therewith. In the context of this invention herbicides are to be understood as including desiccants and defoliants.

In one preferred embodiment there is provided a herbicidal composition comprising a herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor which is 2-acetyl nicotinic acid 4-(3,5-difluorophenyl) semi-carbazone in free acid or salt form, and at least one other herbicide which is selected from the group consisting of another auxin transport inhibitor, a growth regulator herbicide, a photosynthesis inhibitor, a pigment inhibitor, a cell wall biosynthesis inhibitor, a cell wall biosynthesis inhibitor and a rapid cell membrane disrupter wherein the auxin transport inhibitor is present

-3a-

in an amount producing a potentiating effect, and wherein the weight ratio of the auxin transport inhibitor to the other herbicide is 1:2000 to 100:1.

Potentiating, as herein used, refers to the interaction of the auxin transport inhibitor with the herbicide such that the activity is greater than the predicted activity, based upon the activity observed for the auxin transport inhibitor and the herbicide separately. Thus, co-application results in herbicidal activity which is significantly superior to the additive effectiveness of the individual active substances.

This potentiation manifests itself in various forms. Thus, co-application enables application rates to be employed for the auxin transport inhibitor(s) and/or herbicide(s) which would be insufficiently effective if employed alone, or enables various types of weeds to be controlled which would not be controlled by application of each individual active ingredient alone at the same rates as in the mixture.

Furthermore, co-application results in herbicidal activity which is significantly superior to the additive effectiveness of the individual active substances. Moreover, the auxin transport inhibitors of this application are able to increase the efficacy of a herbicide such that the maximum level of control or growth regulation for a given application rate of a herbicide is increased, or alternatively, the application rate of a herbicide giving optimum control or growth regulation can be reduced.

Under co-application is to be understood concurrent, or immediately sequential application (e.g. within 24 hours), application as a tank mix or application of fixed combination premixes.

Non-limiting examples of herbicides which may be potentiated by use of auxin transport inhibitors, especially compounds of formula A in accordance with the invention include

1. other auxin transport inhibitors, e.g. naptalam;
2. growth regulators, including a) benzoic acids, e.g. dicamba; b) phenoxy acids i) acetic acid type, e.g. 2,4-D, MCPA, ii) propionic acid type, e.g. 2,4-DP, MCPP, iii) butyric acid type, e.g. 2,4-DB, MCPB; c) picolinic acids and related compounds, e.g. picloram, triclopyr, fluroxypyr, clopyralid;
3. photosynthesis inhibitors, including a) s-triazines i) chloro substituted, e.g. atrazine, simazine, cyanazine, ii) methoxy substituted, e.g. prometon, iii) methylthio substituted, e.g. ametryn, prometryn; b) other triazines, e.g. hexazinone, metribuzin; c) substituted ureas, e.g. diuron, fluometruon, linuron, tebuthiuron, thidiazuron, forchlorfenuron; d)

-5-

uracils, e.g. bromacil, terbacil; e) others, e.g. bentazon, desmidepham, methazole, phenmedipham, propanil, pyrazon, pyridate;

4. pigment inhibitors, including a) pyridazinones, e.g. norflurazon; b) isoxazolones, e.g. clomazone; c) triketones and cyclic diones of the type described in US Patents 4,695,673; 4,921,526; 5,006,150; 5,089,046, EP-A-338992; EP-A-394889 and EP-A-506907 including for example 2-(2-chloro-4-methylsulfonylbenzoyl)-1,3-cyclohexanedione (sulcotrione); 2-(4-methylsulfonyloxy-2-nitrobenzoyl)-4,4,6,6-tetramethyl-1,3-cyclohexanedione; 3-(4-methylsulfonyloxy-2-nitrobenzoyl)-bicyclo-[3,2,1]octane-2,4-dione; 3-(4-methylsulfonyl-2-nitrobenzoyl)-bicyclo-[3,2,1]octane-2,4-dione; 4-(4-chloro-2-nitrobenzoyl)-2,6,6-trimethyl-2H-1,2-oxazine-3,5(4H,6H)dione; 4-(4-methylthio-2-nitrobenzoyl)-2,6,6-trimethyl-2H-1,2-oxazine-3,5(4H,6H)-dione; 3-(4-methylthio-2-nitrobenzoyl)-bicyclo[3,2,1]octane-2,4-dione; 4-(2-nitro-4-trifluoromethoxybenzoyl)-2,6,6-trimethyl-2H-1,2-oxazine-3,5(4H,6H)-dione; d) others, e.g. amitrole, fluridone;
5. growth inhibitors, including a) mitotic disruptors i) dinitroanilines, e.g. trifluralin, prodiamine, benefin, ethalfluralin, isopropalin, oryzalin, pendimethalin; ii) others, e.g. DCPA, dithiopyr, thiazapyr, pronamide; b) inhibitors of shoots of emerging seedlings i) thiocarbamates, e.g. EPTC, butylate, cycloate, molinate, pebulate, thiobencarb, triallate, vernolate; c) inhibitors of roots only of seedlings, e.g. bensulide, apropanide, siduron; d) inhibitors of roots and shoots of seedlings, including chloroacetamides e.g. alachlor, acetochlor, metolachlor, diethatyl, propachlor, and thiophenamines such as dimethenamid (2-chloro-N-(1-methyl-2-ethoxy-ethyl)-N-(2,4-dimethyl-thien-3-yl)acetamide; cf US Patent 4,666,502), and others e.g. cinmethylin;
6. amino acid synthesis inhibitors, including a) glyphosate, glufosinate; b) sulfonylureas, e.g. metsulfuron, metsulfuron-methyl, ethametsulfuron, nicosulfuron, triasulfuron, primisulfuron, bensulfuron, chlorimuron, chlorimuron-ethyl, chlorsulfuron, sulfometuron, thifensulfuron, tribenuron, triflusulfuron, clopyrasulfuron and pyrazasulfuron; c) sulfonamides, e.g. flumetsulam (DE498); d) imidazolinones, e.g.

-6-

imazaquin, imazamethabenz, imazapyr, imazethapyr;

7. lipid biosynthesis inhibitors, including a) cyclohexanediones, e.g. sethoxydim, clethodim; b) aryloxyphenoxys, e.g. fluazifop-P-butyl, diclofop-methyl, haloxyfop-methyl, quizalofop; c) others, e.g. fenoxaprop-ethyl;
8. cell wall biosynthesis inhibitors, e.g. dichlobenil, isoxaben;
9. rapid cell membrane disruptors, including a) bipyridiliums, e.g. paraquat, diquat; b) diphenyl ethers, e.g. acifluorfen, fomesafen, lactofen, oxyfluorfen; c) glutamine synthetase inhibitors, e.g. glufosinate; d) others, e.g. oxadiazon;
10. miscellaneous, including a) carbamates, e.g. asulam; b) nitriles, e.g. bromoxynil, ioxynil; c) hydantocidin and derivatives; d) various, e.g. paclobutrazol, ethofumesate, quinclorac (BAS 514), difenzoquat, endothall, fosamine, DSMA, MSMA;
11. Others

Compounds of the type described in EP-A-315889; EP-A-461,079 and EP-A-549524; and PCT Application WO 91/10653 including for example

3-[(4,6-dimethoxy-2-pyrimidinyl)hydroxymethyl]- N-methyl-2-pyridine carboxamide;
 4,7-dichloro-3-(4,6-dimethoxy-2-pyrimidinyl)- 3-hexanoyl-oxyphthalide;
 3-[(4,6-dimethoxy-2-pyrimidinyl)carbonyl]-N,N-dimethyl- 2-pyridine carboxamide;
 3,6-dichloro-2-[(4,6-dimethoxy-2-pyrimidinyl)-carbonyl] benzoic acid;
 6-chloro-2-[(4,6-dimethoxy-2-pyrimidinyl)thio]-benzoic acid (DPX-PE350 or pyriothiobac) and salts thereof.

It will be noted that in some cases one auxin transport inhibitor may potentiate the effect of another. The nature of the effect of the auxin transport inhibitor is such that it has the potential to enhance the activity of different classes of herbicides. The present invention therefore also concerns a method of combatting or controlling undesired plant growth or otherwise regulating plant growth which comprises co-applying to a locus where such

combatting or control is desired an herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor and at least one other herbicide, wherein the auxin transport inhibitor is applied at a potentiating rate.

Application rates for co-application will of course vary depending upon climatic conditions, season, soil ecology, weeds to be combatted and the like, however, successful results can be obtained e.g. with rates of auxin transport inhibitor of 0.00011 kg to 1.1 kg/ha (0.0001 lb to 1.0 lb/A), preferably 0.0011 to 0.55 kg/ha (0.001 to 0.5 lb/A), especially 0.011 to 0.11 kg/ha (0.01 to 0.1 lb/A) in co-application with rates for partner herbicides which correspond to or are significantly lower than recommended for use thereof individually (application rates hereinafter set forth are calculated from measurements originally made in lb/A using the conversion factor 1 lb/A = 1.1 kg/ha).

The suitability of specific co-applications for pre- or post-emergent uses and selectivity will of course depend on the partners chosen.

The activity of compounds of formula A is described in the above mentioned patents and that of other known auxin transport inhibitors and of suitable herbicidal partners is described in the literature or on commercially available forms thereof (cf also CROP PROTECTION CHEMICALS REFERENCE, Chemical & Pharmaceutical Press, NY, NY).

The invention also provides herbicidal or plant growth regulating compositions comprising at least one auxin transport inhibitor and at least one other herbicide, wherein the auxin transport inhibitor is present in a potentiating amount. Especially preferred compositions contain a compound of formula A.

Such compositions contain the active substances in association with agriculturally acceptable diluents. They may be employed in either solid or liquid forms e.g. in the form of a wettable powder or an emulsifiable concentrate, incorporating conventional diluents.

Such compositions may be produced in conventional manner, e.g. by mixing the active ingredient with a diluent and optionally other formulating ingredients such as surfactants and oils.

The term diluents as used herein means any liquid or solid agriculturally acceptable material which may be added to the active constituent to provide a more easily or improved applicable form, or to achieve a usable or desirable strength of activity. Examples of diluents are talc, kaolin, diatomaceous earth, xylene, non-phytotoxic oils, or water.

Particular formulations, to be applied in spraying forms such as water dispersible concentrates, water dispersible granules, or wettable powders, may contain surfactants such as wetting and dispersing agents, e.g. the condensation product of formaldehyde with naphthalene sulphonate, an alkylarylsulphonate, a lignin sulphonate, a fatty alkyl sulphate, an ethoxylated alkylphenol or an ethoxylated fatty alcohol.

In general, the formulations include from 0.01 to 90% by weight of active agent(s) and from 0 to 20% by weight of agriculturally acceptable surfactant, the active agent consisting either of at least one auxin transport inhibitor and at least one other herbicide. Concentrate forms of compositions generally contain between about 2 and 90%, preferably between about 5 and 80% by weight of active agent. Application forms of formulation may for example contain from 0.01 to 20% by weight of active agent.

When employing concurrent, immediately sequential or tank mix applications the non-auxin transport inhibitor partner(s) can be employed in commercially available form if appropriate and at rates equivalent to or preferably below those recommended by the manufacturer. The auxin transport inhibitor can be formulated as described in the above mentioned EP-A-219451, US Patents 5,098,462 or 5,098,466.

On co-application according to the present invention other compounds having biological activity, e.g. compounds having insecticidal or fungicidal activity, may also be included.

The preferred modes of application include tank mix prepared e.g. by adding an auxin transport inhibitor to a tank containing the other herbicide partner and an appropriate surfactant and fixed combination premixes.

Depending on the choice of co-application partners both pre- and post-emergent activity on a large range of broadleaf and grassy weeds may be achieved. Non-limiting examples of such weeds are

Setaria sp. - foxtail

Brachiaria platyphylla - broadleaf signalgrass

Ipomoea sp. - morningglories

Abutilon theophrasti - velvetleaf

Hibiscus trionum - Venice mallow

Solanum sp. - nightshades e.g. silverleaf nightshade

Avena fatua - wild oats

Sinapis alba - white mustard

Amaranthus sp. - pigweeds, spiny amaranth

Xanthium strumarium - common cocklebur

Sorghum halepense - johnsongrass

Echinochloa crus-galli - barnyardgrass

Polygonum sp. - smartweeds, wild buckwheat, prostrate knotweed

Cassia obtusifolia - sicklepod

Digitaria sp. - e.g. crabgrasses

Bromus tectorum - downy brome

Apera spica-venti - windgrass

Chenopodium album - common lambsquarter

Sorghum bicolor - shattercane

Portulaca oleracea - common purslane

Sida spinosa - prickly sida

Campsis radicans - trumpet creeper

Rottboellia exaltata - itchgrass

Cynodon dactylon - bermudagrass

Agropyron repens - quackgrass
Cyperus sp. - nutsedges
Panicum sp. e.g. - prosomillet
Lespedeza sp - lespedezas
Trifolium sp - clovers
Hippuris vulgaris - marestail
Asclepias sp - milkweeds
Salvia sp - e.g. lanceleaf sage
Salsola iberica - russian thistle
Convolvulus arvensis - field bindweed
Cirsium arvense - Canada thistle
Proboscidea louisianica - devilsclaw
Senecio sp. - common groundsel
Chorispora tennela - blue mustard
Alopecurus myosuroides - blackgrass
Sisymbrium altissimum - tumble mustard
Caperionia palustris - texasweed

Crop selectivity will also usually depend upon choice of partners.

Compounds of formula A for example exhibit excellent selectivity in corn and small grain crops.

It will be appreciated that mixtures of an auxin transport inhibitor with more than one other herbicide, e.g. 3-way mixes, are also contemplated.

Preferred auxin transport inhibitors are those of formula A especially those wherein M is hydrogen or a sodium, potassium, isopropylammonium or 2-(2-hydroxyethoxy)ethylammonium cation (Compounds A1).

Other compound groups comprise compound of formula A wherein Z represents hydrogen (compounds A2); Z represents fluorine (compounds A3); Z represents chlorine (compounds A4).

Particularly preferred individual auxin transport inhibitors are 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone in free acid or in salt form especially its sodium salt form, 2-acetylnicotinic acid 4-(3-fluorophenyl)semicarbazone in free acid form or in salt form especially in sodium salt form, and 2-acetylnicotinic acid-4-(3-chlorophenyl)-semicarbazone in free acid form or in sodium salt form.

Preferred classes of herbicidal mix partners are growth regulator herbicides, such as benzoic acids, phenoxy acetic acids, picolinic acids and related compounds, growth inhibitors such as inhibitors of roots and shoots of seedlings, rapid cell membrane disruptors such as bipyridiliums, amino acid synthesis inhibitors such as sulfonylureas and sulfonamides.

Examples of specific, preferred herbicidal partners for co-application are dicamba, thidiazuron, 2,4-D, dimethenamid, atrazine, cyanazine, norflurazon, fluroxypyr, primisulfuron, nicosulfuron, pendimethalin, chlorpyralid, paraquat, ethofumesate, flumetsulam (DE498) and for certain situations, glyphosate.

Non-limiting examples of specific combinations are those containing e.g. 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone 2-(2-hydroxyethoxy)ethyl-ammonium salt (a); or 2-acetylnicotinic acid 4-(3-fluorophenyl)semicarbazone sodium salt (b); or 2-acetylnicotinic acid 4-(3-chlorophenyl)semicarbazone sodium salt (c); 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone (d); 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone sodium salt (e); each with e.g. dicamba (z); dimethenamid (y); 2,4-D (x); or thidiazuron (w).

As stated above, application rates may depend on a variety of factors. In general, satisfactory results are obtained when applying the co-application partners at the rates given below:

Compound (a), (b), (c) or (d) 0.0011 to 1.1 kg/ha preferably 0.011 to 0.55 kg/ha, especially 0.011 to 0.11 kg/ha.

Compound (z) 0.011 to 2.2 kg/ha, preferably 0.05 to 0.55 kg/ha, especially 0.11 to 0.55 kg/ha.

Compound (y) 0.11 to 4.4 kg/ha, preferably 0.275 to 1.0 kg/ha, especially 0.55 to 1.0 kg/ha.

Compound (x) 0.011 to 2.2 kg/ha, preferably 0.11 to 1.1 kg/ha, especially 0.275 to 0.825 kg/ha.

Compound (w) 0.011 to 1.1 kg/ha, preferably 0.055 to 0.55 kg/ha, especially 0.088 to 0.44 kg/ha.


The weight ratio of individual components in fixed premixes will vary according to the intended application rate thus for example, the ratio of compound (a) to compound (z) in a premix may vary e.g. from 1:2000 to 100:1, preferably 1:50 to 5:1, especially 1:50 to 1:1 e.g. 1:50 to 1:2.5.

Examples of a 3-way mixtures are compound (e) with compound (z) and nicosulfuron or dimethenamid.

Also for example a mixture of compound (d) or a salt thereof with compound (z) may have added thereto a grass active herbicide such as from classes 5 and 7 listed above.

The compounds of formula A wherein Z is chlorine or fluorine are new and also form part of the invention. The invention therefore further concerns compounds of formula XA

-13-



wherein X, Y and M are as defined for formula A and Z' represents chlorine or fluorine.

The invention also concerns use of compounds of formula XA alone or in combination with other active compounds in combatting weeds, herbicidal compositions containing compounds of formula XA either alone or in combination with other active compounds and processes for preparing compounds of formula XA.

Particular compounds of formula XA are for example those wherein X is fluorine or chlorine, Y is hydrogen or fluorine and Z is 6-fluorine or 6-chlorine (ortho to carboxylic) in salt or free acid form.

Compounds of formula XA wherein Z' is 6-fluorine or 6-chlorine are particularly preferred.

The use of compounds of formula XA and their formulation as herbicidal compositions can be carried out as described herein or as described in US Patents 5,098,462; 5,098,466 and in EP-A-219451.

The following examples are intended to illustrate without in any way limiting the invention.

EXAMPLE 1Field Test

A field trial was conducted against velvetleaf and pigweed in field corn. Application was post-emergent 35 days after seeding with weeds at a height of 41 to 89 cm. Application was of tank mixed formulation by foliar applied broadcast treatment. Compound (a) above was tank mixed as is. Compound (z) above was tank mixed in the form of a 0.5 kg/l s.c. commercially available under the trademark BANVEL®. Aquagene is a commercially available surfactant, (Universal Coop Incorporated, Minneapolis, MN). Values set forth are calculated from measurements originally made in acres, lbs, inches and gallons according to the conversion factors: 1 hectare = 2.47 acres; 1 kg = 2.2 lbs; 1 m = 3.28 ft; and 1 gallon = 3.78 l.

The results may be summarized as follows.

<u>Treatment</u>	Rate <u>kg/ha</u>	% of control	
		<u>Velvetleaf</u>	<u>Pigweed</u>
Aquagene ¹	0	0	0
Compound (z)	0.275	18	30
Compound (a)	0.011	13	18
Compound (z) +	0.275	27	40
Aquagene ¹			
Compound (a) +	0.011	13	18
Aquagene ¹			
Compound (a) +	0.011	57	90
Compound (z)	0.275		
Compound (a) +	0.011	65	93
Compound (z) +	0.275		
Aquagene ¹			
(¹ 0.9 l/ha)			

The tank mix combination of (a) plus (z) was significantly better than either treatment alone. The adjuvant provided some increase in control, but was not responsible for the surprising increase in control observed for the herbicide combination. The combination of (a) plus (z) yielded a response markedly superior to the additive effect of either herbicide which when applied alone at the stated rate showed unsatisfactory weed control.

There was no significant effect on the field corn.

EXAMPLE 2

Greenhouse Test

A greenhouse test was conducted against velvetleaf, pigweed, morningglory and cocklebur. Treatment was at 10 days post-emergence and evaluation at 18 days after treatment. Compound (e) was formulated as technical a.i. in a mixture of equal parts acetone and water with 1/2% surfactant. Compound (z) was used in the commercially available form BANVEL[®] herbicide (=480g/L a.i. equivalent) in water with 1/2% surfactant. The tank mixes were applied in a linear spray chamber with 3 repetitions per concentration.

<u>Treatment</u>	Rate kg/ha	% control			
		<u>Velvetleaf</u>	<u>Pigweed</u>	<u>Morningglory</u>	<u>Cocklebur</u>
Compound (e)	0.01	55	50	55	25
Compound (z)*	0.02	35	35	35	75
Compound (e)+	0.01	98	100	98	100
Compound (z)	0.02				

The results indicate synergism utilizing Limpel's formula and are statistically significant utilizing Duncan's multiple range test.

*as the commercially available BANVEL[®] herbicide.

EXAMPLE 3

Preparation of 2-acetyl-6-fluorobenzoic acid 4-(3,5-difluorophenyl)semicarbazone (Table A cpd 1)

a) Preparation of 3-fluorophthalic anhydride

15 g of 3-fluorophthalic acid are mixed with 16.6 g of acetic anhydride and refluxed for 3 hrs. After removal of unreacted acetic anhydride the remaining white solid is recrystallized from toluene.

b) Preparation of 2-acetyl-6-fluorobenzoic acid

9 g of 3-fluorophthalic anhydride and 6.8 g of malonic acid are mixed in 80 ml of triethylamine and heated in an oil bath at 71-72° until evolution of gas ceases. The reaction mixture is mixed with 50 ml of 10% HCl/H₂O and extracted with ether. The ether is evaporated off and the resulting black oil chromatographed on a column using 1 l of 20% of ethylacetate/hexane followed by 1 l of 30% ethylacetate/hexane to yield first the 3-fluoro-isomer followed by the desired 6-fluoro isomer; m.p. 76-81.5°.

c) Preparation of title compound

3 g of 6-fluoro-2-acetyl benzoic acid and 3 g of 4-(3,5-difluorophenyl)semicarbazide are mixed in 20 ml of methanol and heated until clear. The solution is then stirred at R.T. for 24 hr. A white solid forms which is filtered and dried in vacuum at 60° to yield the title product m.p. 227° (decomp.). The corresponding sodium salt is made by reaction of the free acid with 25% sodium methoxylate/methanol.

The following compounds of formula XA may be prepared analogously.

TABLE A

<u>Cpd no</u>	<u>X</u>	<u>Y</u>	<u>Z'</u>	<u>m.p.</u>
1	F	F	6-F	acid 227° (decomp), Na + salt
2	F	H	6-F	acid 174° (decomp)
3	Cl	H	6-F	acid 157° (decomp)
4	F	F	6-Cl	acid 174° (decomp)
5	F	H	6-Cl	acid 176° (decomp)
6	Cl	H	6-Cl	acid 204° (decomp)

CLAIMS:

1. A herbicidal composition comprising a herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor which is 2-acetyl nicotinic acid 4-(3,5-difluorophenyl) semi-carbazone in free acid or salt form, and at least one other herbicide which is selected from the group consisting of another auxin transport inhibitor, a growth regulator herbicide, a photosynthesis inhibitor, a pigment inhibitor, a growth inhibitor, an amino acid synthesis inhibitor, a lipid biosynthesis inhibitor, a cell wall biosynthesis inhibitor and a rapid cell membrane disrupter wherein the auxin transport inhibitor is present in an amount producing a potentiating effect, and wherein the weight ratio of the auxin transport inhibitor to the other herbicide is 1:2000 to 100:1.
2. A herbicidal composition according to Claim 1 wherein the herbicide is selected from the group consisting of a growth regulator herbicide, a growth inhibitor, a rapid cell membrane disrupter and an amino acid synthesis inhibitor.
3. A herbicidal composition according to Claim 1 in which the herbicide is a benzoic acid, a phenoxy acetic acid, a phenoxy propionic acid, a phenoxy butyric acid, a picolinic acid, a chloro-substituted s-triazine, a methoxy-substituted s-triazine, a methylthio-substituted s-triazine, a triazine, a substituted urea, a uracil, a pyridazinone, an isoxazolone, a dinitroaniline, a thiocarbamate, a chloroacetamide, a thiophenamine, a sulfonylurea, a sulfonamide, an imidazolinone, a cyclohexanedione, an aryloxyphenoxy, a bipyridilium, a diphenyl ether, a carbamate or a nitrile.
4. A herbicidal composition according to Claim 3 wherein the herbicide is a benzoic acid, a phenoxy acetic acid, a picolinic acid or a bipyridilium.
5. A herbicidal composition according to Claim 1 wherein the herbicide is dicamba, thidiazuron, 2,4-D, dimethenamid, norflurazon, fluroxypyr, chloropyralid, paraquat, glyphosate, glufosinate, ethofumesate, nicosulfuron, quinclorac or flumetsulam.

6. A herbicidal composition according to Claim 1 or 3 wherein the auxin transport inhibitor is 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone 2-(2-hydroxyethoxy)ethylammonium salt; or 2-acetylnicotinic acid 4-(3,5-difluorophenyl)semicarbazone and the other herbicide is dicamba dimethenamid; 2,4-D; or thidiazuron.
7. A herbicidal composition according to any one of Claims 1 to 4 wherein the other auxin transport inhibitor is selected from the group consisting of naptalam, TIBA, and DPX 1840.
8. A herbicidal composition according to any one of Claims 1 to 7 wherein the weight ratio of auxin transport inhibitor to the other herbicide is 1:50 to 5:1.
9. A herbicidal composition according to any one of Claims 1 to 7 wherein the weight ratio of auxin transport inhibitor to the other herbicide is 1:50 to 1:1.
10. A herbicidal composition according to any one of Claims 1 to 7 wherein the weight ratio of auxin transport inhibitor to the other herbicide is 1:50 to 1:2.5.
11. A method for combating or controlling undesired plant growth or otherwise regulating plant growth comprising co-applying to a locus where such combatting or control is desired a herbicidally or plant growth regulating effective aggregate amount of at least one auxin transport inhibitor and at least one other herbicide as defined in any one of Claims 1 to 10 wherein the auxin transport inhibitor is applied in a potentiating rate.
12. A method according to Claim 11 wherein the auxin transport inhibitor is applied at a rate of 0.0011 to 1.1 kg/ha.