

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number
WO 2004/048594 A2

- (51) International Patent Classification⁷: **C12Q**
- (21) International Application Number:
PCT/US2003/037356
- (22) International Filing Date:
21 November 2003 (21.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/428,013 21 November 2002 (21.11.2002) US
- (71) Applicant: **EPICENTRE TECHNOLOGIES** [US/US];
726 Post Road, Madison, WI 53713 (US).
- (72) Inventors: **DAHL, Gary, A.**; 2131 Kendall Avenue,
Madison, WI 53726 (US). **JENDRISAK, Jerome, J.**;
2721 Mason Street, Madison, WI 53705 (US). **DAVY-
DOVA, Elena**; 5623 S. Cottage Grove, #1, Chicago, IL
60637 (US). **ROTHMAN-DENES, Lucia**; 5830 South
Stony Island Ave, Apt. 9A, Chicago, IL 60637 (US).
GERDES, Svetlana; 4 Mesa Verde Court, Madison, WI
53705 (US).
- (74) Agent: **VINAROV, Sara, D.**; Quarles & Brady LLP, P.O.
Box 2113, Madison, WI 53701-2113 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

[Continued on next page]

(54) Title: PREPARATION AND USE OF SINGLE-STRANDED TRANSCRIPTION SUBSTRATES FOR SYNTHESIS OF
TRANSCRIPTION PRODUCTS CORRESPONDING TO TARGET SEQUENCES

Bacteriophage N4 vRNAP promoters

-11
A/G
A G
G:C
C:G
X:X'
X:X' +1
3' C:G A A A/T A C C 5'
-17

(57) Abstract: The present invention provides methods, compositions and kits for using an RNA polymerase for making transcription products corresponding to a target sequence by obtaining single-stranded DNA transcription substrates that comprise a single-stranded promoter that is operably joined to a single-stranded target sequence. The invention has broad applicability for research, diagnostic and therapeutic applications, such as preparing cDNA corresponding to full-length mRNA, making sense or anti-sense probes, detecting gene- or organism-specific sequences, cloning, cell signaling or making RNA for use in RNAi.

WO 2004/048594 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

PREPARATION AND USE OF SINGLE-STRANDED TRANSCRIPTION
SUBSTRATES FOR SYNTHESIS OF TRANSCRIPTION PRODUCTS
CORRESPONDING TO TARGET SEQUENCES

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application is a Continuation in Part of U.S. Patent Application No. 10/53,219, which claims priority to U.S. Provisional Patent Application Serial No. 60/292,845, filed May 22, 2001. This application also claims priority to U.S. Provisional Patent Application Serial No. 60/428,013, filed November 21, 2002. The entire disclosure of all referenced priority applications is specifically incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR
DEVELOPMENT

[0002] The government may own rights in the present invention pursuant to grant number R01 A1 12575 from the National Institute of Health.

FIELD OF THE INVENTION

[0003] The present invention relates generally to methods, compositions and kits for using an RNA polymerase for making transcription products corresponding to a target sequence by obtaining single-stranded DNA transcription substrates that comprise a single-stranded promoter that is operably joined to a single-stranded target sequence. The invention has broad applicability for research, diagnostic and therapeutic applications, such as preparing cDNA corresponding to full-length mRNA, making sense or anti-sense probes, detecting gene- or organism-specific sequences, cloning, cell signaling or making RNA for use in RNAi.

BACKGROUND OF THE INVENTION

Description of Related Art

[0004] Transcription of DNA into mRNA is regulated by the promoter region of the DNA. The promoter region contains a sequence of bases that signals RNA polymerase to associate with the DNA, and to initiate the transcription of mRNA using one of the DNA strands as a template to make a corresponding complementary strand of RNA. RNA polymerases from different species typically recognize promoter regions comprised of different sequences. In order to express a protein-encoding gene in a host cell, either the promoter driving transcription of the protein-encoding gene must be recognized by a host RNA polymerase, or an RNA

polymerase which recognizes the promoter driving transcription of the protein-encoding gene must be provided to the host cell (U.S. Pat. No. 6,218,145).

[0005] There are many reasons for which it is beneficial to obtain cDNA and transcribe one or more target nucleic acid sequences. For example, *in vitro* transcription is frequently used in methods to make probes corresponding to mRNA sequences in samples in order to profile the expression of genes in cells of a particular type versus another type or versus a similar type in response to different conditions or stimuli. Contemporary gene expression profiling is typically performed by simultaneously hybridizing labeled probes prepared from one or more samples to arrays or microarrays having sequences for up to hundreds or thousands of different genes attached to a surface. In some cases, the expression or lack of expression of particular genes may correlate with or be indicative of the presence or status of a disease state, such as, but not limited to, a cancer.

[0006] Examples of methods that involve *in vitro* transcription for making probes for gene expression profiling are described in: Murakawa *et al.*, DNA 7:287-295, 1988; Phillips and Eberwine, Methods in Enzymol. Suppl. 10:283-288, 1996; Ginsberg *et al.*, Ann. Neurol. 45:174-181, 1999; Ginsberg *et al.*, Ann. Neurol. 48:77-87, 2000; VanGelder *et al.*, Proc. Natl. Acad. Sci. USA 87:1663-1667, 1990; Eberwine *et al.*, Proc. Natl. Acad. Sci. USA 89:3010-3014, 1992; U.S. Patent Nos. 5,021,335; 5,168,038; 5,545,522; 5,514,545; 5,716,785; 5,891,636; 5,958,688; 6,291,170; and PCT Patent Applications WO 00/75356 and WO 02/065093.

[0007] Still other methods use *in vitro* transcription as part of a process for amplifying and detecting one or more target nucleic acid sequences in order to detect the presence of a pathogen, such as a viral or microbial pathogen, that is a causative agent for a disease or to detect a gene sequence that is related to a disease or the status of a disease for medical purposes. Examples of methods that use *in vitro* transcription for this purpose include U.S. Patent Nos. 5,130,238; 5,194,370; 5,399,491; 5,409,818; 5,437,990; 5,466,586; 5,554,517; 5,665,545; 6,063,603; 6,090,591; 6,100,024; 6,410,276; Kwok *et al.*, Proc. Natl. Acad. Sci. USA 86:1173, 1989; Fahy *et al.*, In: PCR Methods and Applications, pp. 25-33, 1991; PCT Patent Application Nos. WO 89/06700 and WO 91/18155; and European Patent Application Nos. 0427073 A2 and 0427074 A2.

[0008] Most DNA-dependent RNA polymerases read double-stranded DNA, limiting RNA synthesis to systems in which a double-stranded DNA template is available. The synthesis of RNA using single-stranded DNA is not as common. All of the methods referenced above for making probes for gene expression profiling or for amplifying and detecting one or more target

nucleic acid sequences require, at a minimum, the use of a double-stranded transcription promoter, and in most cases, also require the use of a double-stranded DNA as a template. Synthesizing RNA using a single-stranded DNA template immobilized on a solid support is described in U.S. Pat. No. 5,700,667, but transcription of the single-stranded template still required formation of a double-stranded promoter region for binding of the RNA polymerase.

[0009] In contrast to the methods in the art, the present invention provides methods, compositions and kits for transcription of target nucleic acid sequences using RNA polymerases that bind single-stranded DNA promoters and read single-stranded DNA templates.

BRIEF SUMMARY OF THE INVENTION

[0010] The invention provides methods of use for a various novel deletion mutants of bacteriophage N4-coded, virion RNA polymerases (mini-vRNAPs) and compositions and kits therefore. The novel polymerases are described by an isolated nucleic acid comprising a region encoding a polypeptide having the amino sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15. The nucleic acid may comprise the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14. The vRNAP and mini-vRNA polymerase transcribe nucleic acid operatively linked to an N4 promoter such as a P2 promoter of SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29. The promoter of SEQ ID NO:16 or SEQ ID NO:28 is preferred.

[0011] The current invention can use a polypeptide encoded by an isolated polynucleotide comprising a sequence identical or complementary to at least 14 contiguous nucleotides of SEQ ID NO:1. The polynucleotide may comprise at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more contiguous nucleotides of SEQ ID NO:1. The polynucleotide may comprise all contiguous nucleotides of SEQ ID NO:3 or all contiguous nucleotides of SEQ ID NO:1. Similarly, the polynucleotide may comprise at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more nucleotides complementary to at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more contiguous nucleotides of SEQ ID NO:1.

[0012] A purified N4 virion RNA polymerase of the current invention can comprise at least 20 contiguous amino acids of SEQ ID NO:2. It is preferred that the polymerase contain at least 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000 or more contiguous amino acids of SEQ ID NO:2.

[0013] In another aspect, the current invention can use a polypeptide encoded by an isolated nucleic acid comprising a region encoding at least 6 contiguous amino acids of SEQ ID NO:2, wherein the polypeptide has RNA polymerase activity under appropriate reaction conditions. It is preferred that this polypeptide comprises at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000 or more contiguous amino acids of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, or SEQ ID NO:15. The encoded polypeptide may have at least one hexahistidine tag or other tag, or the encoded polypeptide may lack a tag. The polypeptide may be a mutant of the peptide found in SEQ ID NO:2 or SEQ ID NO:4, such as an enzyme possessing an amino acid substitution at position Y678.

[0014] An embodiment of the current invention comprises a method of making RNA. This method comprises: (a) obtaining a N4 virion RNA polymerase (i.e. the polypeptide); (b) obtaining DNA wherein the DNA preferably contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the DNA; and (d) culturing the RNA polymerase and the DNA under conditions effective to allow RNA synthesis. Optionally, the method may comprise synthesizing polynucleotides containing modified ribonucleotides or deoxyribonucleotides. The DNA is preferably single-stranded DNA or denatured double-stranded DNA. Step (c) may occur in a host cell such as an *E. coli* host cell.

[0015] The amino acid sequence of the RNA polymerase is preferably the sequence essentially as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:15, or a mutant form of the polymerase of SEQ ID NO:4 or SEQ ID NO:6. The mutation may be, for example, at position number Y678. The RNA transcript may contain derivatized nucleotides.

[0016] An aspect of the current invention comprises using an N4 vRNAP promoter to direct transcription. The promoter is preferentially an N4 promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29. The P2 promoter of SEQ ID NO:16 or SEQ ID NO:28 is preferred. The promoter sequence may be upstream of the transcription initiation site. The promoter may comprise a set of inverted repeats forming a hairpin with a 2-7 base pair long stem and 3-5 base loop having purines in the central and/or next to the central position of the loop.

[0017] The preferred conditions of the transcription method claimed herein includes a pH in step (c) of between 6 and 9, with a pH of between 7.5 and 8.5 more preferred. Mg^{+2} or Mn^{+2} , preferably Mg^{+2} may be admixed. Preferred temperatures for the reaction are 25° C to

50° C with the range of 30° C to 45° C being more preferred and the range of 32° C to 42° C being most preferred. The admixing may occur *in vivo* or *in vitro*.

[0018] An aspect of the current invention also includes translation of the RNA after transcription. A reporter gene such as an α -peptide of β -galactosidase may be used. It is preferred the transcription comprises admixing an *E. coli* single-stranded binding protein (EcoSSB), a SSB protein homologous to EcoSSB or another naturally occurring or chimeric SSB protein homologous to EcoSSB with the polymerase and DNA. Yet another aspect of the current invention is the transcription method in which no EcoSSB is admixed with the RNA polymerase and DNA; the product of this method is a DNA/RNA hybrid.

[0019] The DNA admixed with the RNA polymerase of the current invention may be single-stranded linear DNA or single-stranded circular DNA such as bacteriophage M13 DNA. The DNA may be denatured DNA, such as single-stranded, double-stranded linear or double-stranded circular denatured DNA. The DNA may also be double-stranded DNA under certain conditions. The RNA may be pure RNA or may contain modified nucleotides. Mixed RNA-DNA oligonucleotides may also be synthesized with the Y678F mutant mini-vRNAP (SEQ ID NO:8) of the current invention.

[0020] The synthesized RNA may comprise a detectable label such as a fluorescent tag, biotin, digoxigenin, 2'-fluoro nucleoside triphosphate, or a radiolabel such as a ³⁵S- or ³²P-label. The synthesized RNA may be adapted for use as a probe for blotting experiments or in-situ hybridization. Nucleoside triphosphates (NTPs) or derivatized NTPs may be incorporated into the RNA, and may optionally have a detectable label. Deoxynucleoside triphosphates may be incorporated into the RNA.

[0021] The RNA may be adapted for use for NMR structural determination. Short RNAs such as those between 10 and 1000 bases or between 10 and 300 bases may be used. The RNA may be adapted for use in spliceosome assembly, splicing reactions or for antisense or RNA interference experiments. Also, the RNA may be adapted for use in probing for a complementary nucleotide sequence or for use as a probe in RNase protection studies.

[0022] Yet another aspect of the current invention comprises delivering RNA into a cell after transcription of the RNA. The delivery may be by microinjection, transfection, electroporation, or other methods in the art. Another aspect of the invention comprises amplifying the RNA after transcription.

[0023] Another embodiment of the current invention comprises a method of making RNA comprising: (a) obtaining a N4 virion RNA polymerase; (b) obtaining a single-stranded

DNA oligonucleotide wherein the oligonucleotide contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the oligonucleotide; and (d) culturing the RNA polymerase and the oligonucleotide under conditions effective to allow RNA synthesis. The polymerase preferentially has the amino sequence set forth in SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. In this embodiment, it is preferred that the DNA has between 20 and 200 bases.

[0024] Yet another embodiment of the invention comprises a method of making RNA comprising: (a) obtaining a N4 virion RNA polymerase; (b) obtaining a single-stranded DNA wherein the DNA contains a N4 virion RNA polymerase promoter sequence; (c) obtaining a ribonucleoside triphosphate (XTP) or a derivatized ribonucleoside triphosphate; (d) admixing the RNA polymerase, the DNA and the XTP; and (e) culturing the RNA polymerase and the oligonucleotide under conditions effective to allow RNA synthesis wherein the RNA is a derivatized RNA. The RNA polymerase preferentially has the amino sequence set forth in SEQ ID NO:4 or SEQ ID NO:6 or a mutant of the polymerase of SEQ ID NO:4 or SEQ ID NO:6, such as a mutant with a mutation at position number Y678 or the polymerase of SEQ ID NO:8.

[0025] Another embodiment of the invention comprises a method for making transcription product corresponding to a target nucleic acid sequence, the method comprising: (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter; (b) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic acid sequence that is present in or complementary to at least a portion of a target nucleic acid in a sample; (c) obtaining a ssDNA transcription substrate by operably joining to the single-stranded DNA a single-stranded promoter sequence that binds the RNA polymerase; (d) obtaining nucleoside triphosphates (NTPs) that are substrates for the RNA polymerase and that are complementary to the canonical nucleic acid bases; (e) admixing the RNA polymerase, the ssDNA transcription substrate and the NTPs; and (f) incubating the RNA polymerase and the ssDNA transcription substrate under conditions effective to allow synthesis of transcription product. The RNA polymerase preferentially has the amino sequence set forth in SEQ ID NO:4 or SEQ ID NO:6 or a mutant of the polymerase of SEQ ID NO:4 or SEQ ID NO:6, such as a mutant with a mutation at position number Y678 or the polymerase of SEQ ID NO:8.

[0026] Yet another embodiment of the invention comprises a method for obtaining additional rounds of synthesis of transcription product corresponding to a target nucleic acid sequence, the method comprising: (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter; (b) obtaining a first transcription product by transcription of a

first ssDNA transcription substrate corresponding to a target nucleic acid sequence; (c) obtaining a reverse transcriptase; (d) reverse transcribing the first transcription product; (e) obtaining first-strand cDNA complementary to the first transcription product; (f) obtaining a second ssDNA transcription substrate by operably joining to the first-strand cDNA a single-stranded promoter sequence that binds the RNA polymerase; (g) admixing the RNA polymerase and the second ssDNA transcription substrate; and (e) incubating the RNA polymerase and the second ssDNA transcription substrate under conditions effective to allow synthesis of a second transcription product. This method for obtaining additional rounds of synthesis of transcription product can be repeated to obtain synthesis of still more transcription products. The RNA polymerase preferentially has the amino sequence set forth in SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

[0027] Another embodiment of the invention comprises a method for *in vivo* or *in vitro* protein synthesis comprising: (a) obtaining an RNA polymerase having the amino sequence set forth in SEQ ID NO:4, SEQ ID NO:6 or a mutant thereof; (b) obtaining DNA wherein the DNA contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the DNA; (d) culturing the RNA polymerase and the DNA under conditions effective to allow RNA synthesis; and (e) culturing the RNA *in vivo* or *in vitro* under conditions effective to allow protein synthesis. Step (e) may comprise using a two plasmid system or a one plasmid system in which a reporter gene and the RNA polymerase gene are located on the same plasmid.

[0028] Another embodiment of the invention provides for a method that comprises a target nucleic acid sequence comprising a 3'-portion that encodes a first sequence, a 5'-portion that encodes a second sequence that is complementary to the first sequence, and a middle portion that joins the 3'portion and the 5'portion, wherein the middle portion comprises a sequence that is not complementary to either the 3'-portion or the 5'portion and wherein the transcription product comprises a hairpin RNA corresponding to a target nucleic acid sequence in a target nucleic acid, wherein the hairpin RNA is siRNA has RNA interference activity in a cell.

[0029] Another embodiment of the invention provides for a method of cloning a target nucleic acid, the method comprising:

- a) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic acid sequence that is present in or complementary to the target nucleic acid;
- b) joining to the single-stranded DNA a single-stranded polynucleotide comprising a single-stranded origin of replication and a marker gene;

- c) making a circular ssDNA molecule by covalently joining the 3'-end and the 5'-end of the product of step (b);
- d) transforming the circular ssDNA molecule into a host cell, in which the marker gene is expressible, wherein the host cell is capable of replicating the circular ssDNA molecule; and
- e) growing the host cell under conditions that support the expression of the marker gene.

[0030] Still another embodiment of the invention provides for a method for detecting an analyte in a sample, the method comprising:

- a) obtaining a transcription signaling system comprising a ssDNA comprising:
 - i) a promoter sequence that binds an RNA polymerase that can transcribe RNA using a single-stranded promoter, and
 - ii) a signal sequence that is operably joined to the promoter sequence;
- b) joining the transcription signaling system to an analyte-binding substance;
- c) contacting the analyte-binding substance to which the transcription signaling system is joined with a sample under conditions effective to allow binding of an analyte to the analyte-binding substance and forming a specific binding pair;
- d) removing the specific binding pair from the sample;
- e) incubating the specific-binding pair with an RNA polymerase that can transcribe RNA using a single-stranded promoter under conditions effective to allow synthesis of a transcription product; and
- f) detecting the transcription product.

[0031] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although suitable methods and materials for the practice or testing of the present invention are described below, other methods and materials similar or equivalent to those described herein, which are well known in the art, can also be used.

[0032] Other objects, advantages and features of the present invention will become apparent from the following specification taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0033] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0034] FIG. 1--Bacteriophage N4 vRNAP promoters on single-stranded templates. These promoters are characterized by conserved sequences and a 5 bp stem, 3 base loop hairpin structure.

[0035] FIG. 2A and FIG. 2B--N4 vRNAP and generation of mini-vRNAP. FIG. 2A shows a schematic of the N4 vRNAP protein with three motifs: the T/DxxGR motif found in DNA-dependent polymerases, the P-loop, an ATP/GTP-binding motif present in some nucleotide-binding proteins, and motif B (Rx₃Kx₆₋₇YG), one of three motifs common to the Pol I and Pol α DNA polymerases and the T7-like RNA polymerases. FIG. 2B shows the mini-vRNAP.

[0036] FIG. 3A and FIG. 3B--Identification of the minimal transcriptionally active domain of N4 vRNAP by proteolytic cleavage. FIG. 3A, SDS-PAGE analysis of the products of vRNAP digestion with trypsin. FIG. 3B N-terminal sequencing of the three initial proteolytic fragments indicated that the stable active polypeptide (mini-vRNAP) corresponds to the middle 1/3 of vRNAP, the region containing the three motifs described in FIG. 2A.

[0037] FIG. 4--ORFs for full length polymerase, mini-vRNAP and mutants thereof were cloned under pBAD control with an N-terminal hexahistidine tag. In other experiments, ORFs for mini-vRNAP and the Y678F mutant thereof, both of which lacked an N-terminal hexahistidine tag, were cloned in *E. coli* under pT7 control in cells which inducibly express T7 RNAP; the polypeptides obtained showed similar activities to the polypeptides with hexahistidine tags shown in this figure.

[0038] FIG. 5--Purification of cloned vRNAP and mini-vRNAP. The left hand side shows the relative amounts of full size and mini-vRNAP proteins purified on TALON columns from the same volume of induced cells. Further concentration on a monoQ column reveals that, in contrast to full size vRNAP, mini-vRNAP is stable after induction (right).

[0039] FIG. 6--Activation of N4 vRNAP transcription by EcoSSB at three different ssDNA concentrations. The extent of EcoSSB activation is template-concentration dependent, with highest activation at low DNA template concentration.

[0040] FIG. 7A, FIG. 7B, FIG. 7C, and FIG. 7D--Effect of EcoSSB on ssDNA template recycling. In the absence of EcoSSB, no increase in transcription was observed beyond 10 min of incubation (FIG. 7A). Addition of template at 20 min to the reaction carried out in the absence of EcoSSB led to a dramatic increase in RNA synthesis (FIG. 7B). RNA synthesis increased linearly throughout the period of incubation (FIG. 7C). Addition of EcoSSB at 20 min led to a slow rate of transcriptional recovery (FIG. 7D).

[0041] FIG. 8--Effect of EcoSSB on the state of template DNA and product RNA in vRNAP transcription. Native gel electrophoresis was carried out in the absence and in the presence of EcoSSB. Transcription was performed at an intermediate (5 nM) DNA concentration, at which only a 2-fold effect of EcoSSB is observed. Either ^{32}P -labeled template (right panel) or labeled NTPs (left panel) were used to analyze the state of the template (right panel) or RNA product (left panel) in the absence or presence of EcoSSB.

[0042] FIG. 9A, FIG. 9B, and FIG. 9C--Transcription initiation by vRNAP and mini-vRNAP. The initiation properties of the full length and mini-vRNA polymerases were compared at similar molar concentrations (FIG. 9A) using the catalytic autolabeling assay and two reaction conditions: using a template containing +1C, the benzaldehyde derivative of GTP and $\alpha\text{-}^{32}\text{P}$ -ATP, or a template containing +1T, the benzaldehyde derivative of ATP and $\alpha\text{-}^{32}\text{P}$ -GTP. Comparison of the results in FIGS. 9B and 9C demonstrates that mini-vRNAP exhibits initiation properties similar to full size vRNAP.

[0043] FIG. 10--Effect of EcoSSB on transcription of vRNAP and mini-vRNAP. The elongation and termination properties of vRNAP and mini-vRNAP are compared.

[0044] FIG. 11A and FIG. 11B--Determination of mini-vRNAP promoter contacts. A 20-base oligonucleotide containing wild type promoter P2 sequence binds with a 1 nM Kd (FIG. 11A). Most oligonucleotides substituted with 5-Iodo-dU at specific positions showed close to wild type affinity except for the oligonucleotides substituted at positions -11 (at the center of the loop) and -8, indicating that these positions are essential for promoter recognition (FIG. 11B). UV crosslinking indicates that mini-vRNAP primarily contacts the -11 position.

[0045] FIG. 12--Binding affinities of stem-length promoter mutants. Wild type promoter P2 with a 5 bp stem has a Kd of 1 nM (top). The stem was shortened by removal of 3' bases (left). The stem can be shortened by two base pairs without change in the binding affinity. The effect of lengthening the stem by addition of 3' bases is shown (right). The stem can be lengthened by two base pairs without change in the binding affinity.

[0046] FIG. 13A and FIG. 13B--Identification of the transcription start site by catalytic autolabeling. A series of templates were constructed with a single C placed at different distances from the center of the hairpin (position -11) by addition or deletion of the tract of As present at promoter P2 (FIG. 13A). The affinity of mini-vRNAP for these promoters was measured by filter binding, and transcription initiation was measured by catalytic autolabeling of mini-vRNAP. All templates showed similar binding affinities. However, only the template with a C positioned 12 bases downstream from the center of the hairpin was able to support transcription initiation (FIG. 13B).

[0047] FIG. 14--UV crosslinking of mutant mini-vRNAPases to promoter oligonucleotides. Two mutants (K670A and Y678F) were tested for their ability to bind to wild type promoters. Both mutant RNA polymerases bound to promoter DNA with wild type affinities and crosslinked to 5-Iodo-dU substituted P2 DNA templates at positions -11 and +3 as well as the wild type enzyme, indicating that these polymerase mutations do not affect promoter binding.

[0048] FIG. 15--Run-off transcription by mutant mini-vRNAPases. The wild type and Y678F (SEQ ID NO:8) enzymes displayed similar activities at both template excess and template-limiting conditions, while the K670A enzyme exhibited decreased activity under both conditions. Under limiting template conditions, all three enzymes were activated by EcoSSB (right panel). However, the Y678F enzyme showed reduced discrimination between incorporation of ribo- and deoxyribonucleoside triphosphates.

[0049] FIG. 16--Mutant mini-vRNAPases in transcription initiation. The initiation properties of the three enzymes were compared using catalytic autolabeling. The K670A enzyme displays significantly reduced activity with the GTP derivative. The Y678F enzyme, in contrast to wild type polymerase, incorporates dATP as efficiently as rATP in a single round of phosphodiester bond formation.

[0050] FIG. 17A, FIG. 17B, and FIG. 17C--Detection of *in vivo* activities of N4 vRNAP and mini-vRNAP. Transcription of β -galactosidase α -peptide by full size and mini-vRNAP was assayed on inducing-Xgal media (FIG. 17A). Plasmid (pACYC) templates were constructed with a reporter gene (α -peptide of β -galactosidase) under the control of vRNAP promoter P2 cloned in either of two orientations (FIG. 17B). Induction of mini-vRNAP led to production and accumulation of detectable levels of the protein, whereas full-length vRNAP was degraded (FIG. 17C).

- [0051]** FIG. 18--Schematic of an embodiment of the invention for obtaining a ssDNA transcription substrate using a promoter splice template oligo.
- [0052]** FIG. 19--Schematic of an embodiment of a continuous transcription assay of the invention that utilizes a promoter splice template oligo and a blocker oligo.
- [0053]** FIG. 20--Schematic of an embodiment of the invention for obtaining a ssDNA transcription substrate using a promoter ligation oligo.
- [0054]** FIG. 21--Schematic of an embodiment of a continuous transcription assay that uses a promoter ligation oligo and a blocker oligo.
- [0055]** FIG. 22--Embodiments of the invention for obtaining both circular and linear ssDNA transcription substrates using a promoter primer.
- [0056]** FIG. 23--Schematic of an embodiment of a continuous transcription assay that uses a promoter primer that also comprises transcription termination sequences to generate a circular ssDNA transcription substrate.
- [0057]** FIG. 24--Embodiment that uses a promoter primer comprising an anti-sense sequence of an N4 vRNAP promoter to make a circular anti-sense-promoter-containing first-strand cDNA that serves as a template for rolling circle replication, thereby synthesizing a linear second-strand ssDNA transcription substrate which can be used to synthesize more RNA.

DETAILED DESCRIPTION OF THE INVENTION

[0058] Methods are provided for using various deletion mutants of bacteriophage N4-coded, virion RNA polymerases (mini-vRNAPs), which have been developed, cloned and expressed in *E. coli*, and purified in active form. In the presence of *E. coli* SSB protein, mini-vRNAPs efficiently transcribe single-stranded DNA (ssDNA) transcription substrates that comprise a single-stranded DNA target sequence as a template if the template is operably joined to a sequence comprising a single-stranded 17-base promoter. The enzymes efficiently incorporate derivatized nucleoside triphosphates, and a specific amino acid mutant of mini-vRNAP (Y678F) also incorporates other nucleotides, such as 2'-deoxynucleoside triphosphates, with greater efficiency. Methods, compositions and kits are provided for making ssDNA transcription substrates for synthesis of transcription products corresponding to a target sequence and for obtaining additional rounds of transcription by using transcription products synthesized in earlier rounds for obtaining additional ssDNA transcription substrates. Methods are provided for obtaining replicable clones, including expression clones, of target sequences by using oligos comprising a single-stranded origin of replication and a sequence encoding a selectable or

screenable marker for obtaining circular ssDNA, which is used to transform a host cell. Other methods are provided for obtaining signaling systems for analytes by using mini-vRNAP to transcribe ssDNA comprising a promoter and a signal sequence, which ssDNA is joined to an analyte-binding substance.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0059] The present invention relates to novel methods, compositions, and kits for synthesizing RNA corresponding to one or multiple target nucleic acid sequences in or from a sample. The target sequence can comprise at least a portion of one or more target nucleic acids comprising either RNA or DNA from any source. By way of example, but not of limitation, the target nucleic acid sequences that are transcribed using a method of the invention can comprise first-strand cDNA corresponding to from one to substantially all mRNA molecules in a sample. Alternatively, the target nucleic acid sequence that is transcribed using a method of the invention can comprise a specific sequence in genomic DNA of a particular organism. The present invention overcomes deficiencies in the art by providing a stable RNA polymerase that uses single-stranded DNA transcription substrate that comprises both a sense promoter and a template for transcription and provides a unique system to synthesize RNAs of a desired sequence. Thus, in contrast to other methods currently used in the art, the methods of the present invention do not require synthesis of double-stranded DNA, use of a proto-promoter, or annealing of a promoter-complementary oligonucleotide for transcription. The RNA polymerase and mini-vRNA polymerase can be used to synthesize RNA for use as probes for expression studies using arrays or microarrays or in RNase protection studies of DNAs or RNAs, in situ hybridization studies, and in Southern and Northern blot analysis, for the synthesis of defined RNA:DNA hybrids, for NMR structure determination of RNAs, for *in vitro* studies of spliceosome assembly, splicing reactions and antisense or RNA interference experiments, for *in vitro* translation or microinjection, and for nucleic acid amplification. The present invention allows for the synthesis of derivatized RNA and can use ssDNA in the form of single-stranded oligonucleotides, denatured DNA or DNA cloned into M13 templates.

I. RNA Polymerases

a. Structure and Promoter Recognition of DNA-Dependent RNA Polymerases

[0060] Inspection of the sequences of phage, archaeobacterial, eubacterial, eukaryotic and viral DNA-dependent RNA polymerases has revealed the existence of two enzyme families.

The eubacterial, eukaryotic, archaeobacterial, chloroplast and the vaccinia virus RNA polymerases are complex multisubunit enzymes (5-14 subunits) composed of two large subunits, one to several subunits of intermediate molecular weight (30-50-kDa) and none to several subunits of small molecular weight (<30-kDa) (Archambault and Friesen, Microbiol. Rev. 57:703-724, 1993; Record et al., Cell and Molecular Biology 1:792-821, 1995. Eubacterial RNA polymerases are the simplest with an $\alpha_2\beta\beta'$ core structure. Sequence comparison of the genes coding for the different subunits of these enzymes has revealed: 1-sequence homology in eight segments (A to H) between β' and the largest subunit of other RNA polymerases, 2-sequence homology in nine segments (A to I) between β and the next largest subunit of other RNA polymerases, 3-sequence homology in 3 segments (1.1, 1.2 and 2) between a and a subunit in RNA polymerases I, II and III (Puhler, et al., Proc. Natl. Acad. Sci. USA 86:4569-4573, 1989; Sweetser, et al., Proc. Natl. Acad. Sci. USA 84:1192-1196, 1987). Not surprisingly, the crystal structures of yeast RNAP II and *E. coli* RNAP core revealed remarkable similarities (Zhang, et al., Cell 98:811-824, 1999; Cramer, et al., Scienceexpress, www.sciencexpress.org. 19 April, 2001).

[0061] In contrast, members of the phage T7-like (T3, SP6) family of RNA polymerases consist of a single (~100 kDa) polypeptide which catalyzes all functions required for accurate transcription (Cheetham, et al., Curr. Op. In Struc. Biol. 10:117-123, 2000). The heterodimeric bacteriophage N4 RNAP II, nuclear-coded mitochondrial, and Arabidopsis chloroplast RNA polymerases show sequence similarity to the phage RNA polymerases (Cermakian, et al., Nuc. Acids Res. 24:648-654, 1996; Hedtke, et al., Science 277:809-811, 1997; Zehring, et al., J. Biol. Chem. 258:8074-8080, 1983). Three sequence motifs -A and C, which contain the two aspartic acids required for catalysis, and motif B- are conserved in polymerases that use DNA as a template (Delarue, et al., Protein Engineering 3:461-467, 1990). The crystal structure of T7 RNAP resembles a "cupped right hand" with "palm," "fingers" and "thumb" subdomains (Sousa, et al., Nature 364:593-599, 1993). The two catalytic aspartates are present in the "palm" of the structure. This structure is shared by the polymerase domains of *E. coli* DNA polymerase I and HIV reverse transcriptase (Sousa, Trends in Biochem. Sci. 21:186-190, 1996). Genetic, biochemical and structural information indicates that T7 RNA polymerase contains additional structures dedicated to nascent RNA binding, promoter recognition, dsDNA unwinding and RNA:DNA hybrid unwinding (Cheetham, et al., Curr. Op. In Struc. Biol. 10:117-123, 2000; Sousa, Trends in Biochem. Sci. 21:186-190, 1996).

[0062] Both Class I and Class II RNA polymerases recognize specific sequences, called promoters, on B form double-stranded DNA. Eubacterial promoters (except those recognized by σ^{54}) are characterized by two regions of sequence homology: the -10 and the -35 hexamers (Gross, *et al.*, Cold Spring Harbor Symp. Quant. Biol. 63:141-156, 1998). Specificity of promoter recognition is conferred to the core enzyme by the σ subunit, which makes specific interactions with the -10 and -35 sequences through two distinct DNA binding domains (Gross, *et al.*, Cold Spring Harbor Symp. Quant. Biol. 63:141-156, 1998). This modular promoter structure is also present at the promoters for eukaryotic RNA polymerases I, II and III. Transcription factors TFIIB and TFIIC direct recognition of RNAP III to two separate sequences (boxes A and C, separated by defined spacing) at the 5S gene promoter, while transcription factors TFIIB and TFIIC direct recognition of this enzyme to blocks A and B, separated by variable distance (31-74 bp) at the tRNA promoters (Paule, *et al.*, Nuc. Acids Res. 28:1283-1298, 2000). Sequences important for RNAP I transcription initiation at the human rRNA promoters are also restricted to two regions: the "core" region located at -40 to +1 and the "upstream" region present at -160 to -107 (Paule, *et al.*, Nuc. Acids Res. 28:1283-1298, 2000). Assembly of the initiation complex at RNAP II promoters requires several general transcription factors (TFIIA, TFIIB, TFIID, TFIIIE, TFIIF and TFIIF). Recognition involves three core elements: the TATA box located at position -30 and recognized by TBP, the initiator element located near -1, and the downstream promoter element near +30 (Roeder, Trends Biochem. Sci. 21:327-335, 1996).

[0063] Promoters for the T7-like and mitochondrial RNAPases are simpler. The T7-type RNAP promoters span a continuous highly conserved 23 bp region extending from position -17 to +6 relative to the start site of transcription (+1) (Rong, *et al.*, Proc. Natl. Acad. Sci. USA 95:515-519, 1998). The yeast mitochondrial RNAP promoters are even smaller, extending from -8 to +1 (Shadel, *et al.*, J. Biol. Chem. 268:16083-16086, 1993). One exception are the promoters for N4 RNAP II, which are restricted to two blocks of conserved sequence: a/tTTTA at +1 and AAGACCTG present 18-26 bp upstream of +1 (Abravaya, *et al.*, J. Mol. Biol. 211:359-372, 1990).

[0064] The activity of the multisubunit class of RNA polymerases is enhanced by activators at weak promoters. Transcription activators generally bind at specific sites on double-stranded DNA upstream of the -35 region (with the exception of the T4 sliding clamp activator), or at large distances in the cases of enhancers (Sanders, *et al.*, EMBO Journal 16:3124-3132, 1997). Activators modulate transcription by increasing the binding (formation of closed

complex) or isomerization (formation of open complex) steps of transcription through interactions with the α or σ subunits of RNAP (Hochschild, et al., Cell 92:597-600, 1998). An exception is N4SSB, the activator of *E. coli* RNAP σ^{70} at the bacteriophage N4 late promoters, which activates transcription through direct interactions with the β' subunit of RNAP in the absence of DNA binding (Miller, et al., Science 275:1655-1657, 1997).

[0065] Proteins that bind to ssDNAs with high affinity but without sequence specificity have been purified and characterized from several prokaryotes, eukaryotes, and their viruses (Chase, et al., Ann. Rev. Biochem. 55:130-136, 1986). These proteins (SSBs), which are required for replication, recombination and repair, bind stoichiometrically and, in many cases, cooperatively to ssDNA to cover the transient single-stranded regions of DNA that normally arise *in vivo* as a result of replication, repair and recombination. Binding to DNA results in the removal of hairpin structures found on ssDNA, providing an extended conformation for proteins involved in DNA metabolism. Several lines of evidence suggest that single-stranded DNA binding proteins play a more dynamic role in cellular processes. Genetic and biochemical evidence indicates that these proteins are involved in a multitude of protein-protein interactions including transcription activation (Rothman-Denes, et al., Genes Develpmnt. 12:2782-2790, 1999).

b. The Bacteriophage N4 Virion RNA Polymerase

[0066] Bacteriophage N4 virion RNA polymerase (N4 vRNAP) is present in N4 virions and is injected into the *E. coli* cell at the beginning of infection, where it is responsible for transcription of the N4 early genes (Falco, et al., Proc. Natl. Acad. Sci. (USA) 74:520-523, 1977; Falco, et al., Virology 95:454-465, 1979; Malone, et al., Virology 162:328-336, 1988). The N4 vRNAP gene maps to the late region of the N4 genome (Zivin, et al., J. Mol. Biol. 152:335-356, 1981). N4 vRNAP purified from virions is composed of a single polypeptide with an apparent molecular mass of approximately 320,000 kDa (Falco, et al., Biol. Chem. 255:4339-4347, 1980). In contrast to other DNA-dependent RNAPases, N4 vRNAP recognizes promoters on single-stranded templates (Falco, et al., Proc. Natl. Acad. Sci. (USA) 75:3220-3224, 1978). These promoters are characterized by conserved sequences and a 5 bp stem, 3 base loop hairpin structure (FIG. 1) (Haynes, et al., Cell 41:597-605, 1985; Glucksmann, et al., Cell 70:491-500, 1992). *In vivo*, *E. coli* gyrase and single-stranded binding protein are required for transcription by N4 vRNAP (Falco, et al., J. Biol. Chem. 255:4339-4347, 1980; Markiewicz, et al., Genes and Dev. 6:2010-2019, 1992).

[0067] Sequencing of the N4 vRNAP gene revealed an ORF coding for a protein 3,500 amino acids in length (SEQ ID NO:1-2). Inspection of the sequence revealed no extensive homology to either the multisubunit or the T7-like families of RNA polymerases. However, three motifs are present (FIG. 2A): the T/DxxGR motif found in DNA-dependent polymerases, and Motif B (Rx₃Kx₆₋₇YG), one of three motifs common to the Pol I and Pol α DNA polymerases and the T7-like RNA polymerases.

c. Transcription Using N4 vRNAP

[0068] RNA synthesis requires RNA polymerase, a DNA template, an activated precursor (the ribonucleoside triphosphates ATP, GTP, UTP and CTP (XTP)), and divalent metal ions such as Mg⁺² or Mn⁺². The metal ion Mg⁺² is strongly preferred. Synthesis of RNA begins at the promoter site on the DNA. This site contains a sequence which the RNA polymerase recognizes and binds. The RNA synthesis proceeds until a termination site is reached. N4 vRNAP termination signals comprise a hairpin loop that forms in the newly synthesized RNA which is followed by a string of uracils (poly U). The sequence of the terminator signals for vRNAP present in the N4 genome include SEQ ID NOS:21-26. These N4 vRNAP termination signals possess all of the characteristics of eubacterial sequence-dependent terminators. The ribonucleoside triphosphate may be derivatized with, for example, biotin. Derivatized XTPs can be used for the preparation of derivatized RNA. Exemplary methods for making derivatized XTPs are disclosed in detail in Rashtchian *et al.*, "Nonradioactive Labeling and Detection of Biomolecules," C. Kessler, Ed., Springer-Verlag, New York, pp. 70-84, 1992, herein incorporated by reference.

[0069] Single-stranded DNA of varying lengths can be used as a template for RNA synthesis using the N4 vRNAP or mini-vRNAP. Oligonucleotides and polynucleotides of intermediate length may be used. One particular single-stranded DNA that may be used is M13 DNA. M13 genomic DNA exists temporarily inside infected *E. coli* cells as a double-stranded DNA plasmid and is packaged as a small, single-stranded circular DNA into phage particles. M13 phage particles are secreted by an infected cell and single-stranded DNA can be purified from these particles for use as a transcription template. Initially M13 phage vectors required a working knowledge of phage biology and were primarily used for creating single-strand DNA molecules for DNA sequencing. M13-derived cloning vectors called "phagemids" take advantage of M13 replication to produce single-strand molecules, but can be propagated as conventional ColE1-based replicating double-strand plasmids.

[0070] EcoSSB is essential for N4 vRNAP transcription *in vivo* (Falco *et al.*, Proc. Natl. Acad. Sci. (USA) 75:3220-3224, 1978; Glucksmann, *et al.*, Cell 70:491-500, 1992, herein incorporated by reference). EcoSSB is a specific activator of N4 vRNAP on single-stranded and supercoiled double-stranded DNA templates. EcoSSB, unlike other SSBs, does not melt the N4 vRNAP promoter hairpin structure (Glucksmann-Kuis, *et al.*, Cell 84:147-154, 1996). EcoSSB has a high specificity for N4 vRNAP and mini-vRNAP resulting from EcoSSB's ability to stabilize the template-strand hairpin, whereas the nontemplate strand hairpin is destabilized. Other single-stranded DNA binding proteins destabilize the template-strand hairpin (Glucksmann-Kuis, *et al.*, Cell 84:147-154, 1996; Dai *et al.*, Genes Develpmnt. 12:2782-2790, 1998). EcoSSB mediates template recycling during transcription by N4 vRNAP (Davidova, EK and Rothman-Denes, LB, Proc. Natl. Acad. Sci. USA 100:9250-9255, 2003, incorporated herein by reference). When EcoSSB is not used in N4 vRNAP transcription *in vitro*, a DNA:RNA hybrid is formed, preventing template reutilization.

II. Genes and DNA Segments

[0071] Important aspects of the present invention concern isolated DNA segments and recombinant vectors encoding N4 vRNAP or more particularly mini-vRNAP or a mutant of mini-vRNAP and the creation and use of recombinant host cells through the application of DNA technology, that express a wild type, polymorphic or mutant vRNAP. Other aspects of the present invention concern isolated nucleic acid segments and recombinant vectors encoding vRNAP. Sequences of SEQ ID NO:1, 3, 5, 7, 14 and biologically functional equivalents thereof are used in the current invention. Single-stranded DNA oligonucleotides and polynucleotides can be used as DNA templates.

[0072] The present invention concerns isolated nucleic acid segments that are capable of expressing a protein, polypeptide or peptide that has RNA polymerase activity. As used herein, the term "nucleic acid segment" refers to a nucleic acid molecule that has been isolated free of total genomic DNA of a particular species. Therefore, a nucleic acid segment encoding vRNAP refers to a nucleic acid segment that contains wild-type, polymorphic or mutant vRNAP coding sequences yet is isolated away from, or purified free from, total bacterial or N4 phage genomic DNA. Included within the term "nucleic acid segment," are nucleic acid segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like.

[0073] Similarly, a nucleic acid segment comprising an isolated or purified vRNAP gene refers to a nucleic acid segment including vRNAP protein, polypeptide or peptide coding sequences and, in certain aspects, regulatory sequences, isolated substantially away from other naturally occurring genes or protein encoding sequences. In this respect, the term "gene" is used for simplicity to refer to a functional protein, polypeptide or peptide encoding unit. As will be understood by those of skill in the art, this functional term includes both genomic sequences, cDNA sequences and engineered segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, vRNAPs and mutants of vRNAP encoding sequences.

[0074] "Isolated substantially away from other coding sequences" means that the gene of interest, in this case the vRNAP, or more particularly mini-vRNAP genes, forms the significant part of the coding region of the nucleic acid segment, and that the nucleic acid segment does not contain large portions of naturally-occurring coding DNA, such as large chromosomal fragments or other functional genes or cDNA coding regions. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

[0075] The term "a sequence essentially as set forth in SEQ ID NO:2 means, for example, that the sequence substantially corresponds to a portion of SEQ ID NO:2 and has relatively few amino acids that are not identical to, or a biologically functional equivalent of, the amino acids of SEQ ID NO:2. This applies with respect to all peptide and protein sequences herein, such as those of SEQ ID NO:4, 6, 8 and 15.

[0076] The term "biologically functional equivalent" is well understood in the art and is further defined in detail herein. Accordingly, sequences that have about 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or about 99%, and any range derivable therein, such as, for example, about 70% to about 80%, and more preferably about 81% and about 90%; or even more preferably, between about 91% and about 99%; of amino acids that are identical or functionally equivalent to the amino acids of SEQ ID NO:2 will be sequences that are "essentially as set forth in SEQ ID NO:2," provided the biological activity of the protein is maintained. In particular embodiments, the biological activity of a vRNAP protein, polypeptide or peptide, or a biologically functional equivalent, comprises transcription. A preferred transcriptional activity that may be possessed

by a vRNAP protein, polypeptide or peptide, or a biologically functional equivalent, is RNA synthesis using single-stranded N4 vRNAP promoter-containing DNA as a template.

[0077] In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that include within their sequence a nucleic acid sequence essentially as set forth in SEQ ID NO:1. The term "essentially as set forth in SEQ ID NO:1" is used in the same sense as described above and means that the nucleic acid sequence substantially corresponds to a portion of SEQ ID NO:1 and has relatively few codons that are not identical, or functionally equivalent, to the codons of SEQ ID NO:1. Again, nucleic acid segments that encode proteins, polypeptide or peptides exhibiting RNAP activity will be most preferred.

[0078] The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine and serine, and also refers to codons that encode biologically equivalent amino acids. For optimization of expression of vRNAP in human cells, the following codons are used, with preference of use from left to right: Alanine Ala A GCC GCT GCA GCG; Cysteine Cys C TGC TGT; Aspartic acid Asp D GAG GAT; Glutamic acid Glu E GAG GAA; Phenylalanine Phe F TTC TTT; Glycine Gly G GGC GGG GGA GGT; Histidine His H CAC CAT; Isoleucine Ile I ATC ATT ATA; Lysine Lys K AAG AAA; Leucine Leu L CTG CTC TTG CTT CTA TTA; Methionine Met M ATG; Asparagine Asn N AAC AAT; Proline Pro P CCC CCT CCA CCG; Glutamine Gln Q CAG CAA; Arginine Arg R CGC AGG CGG AGA CGA CGT; Serine Ser S AGC TCC TCT AGT TCA TCG; Threonine Thr T ACC ACA ACT ACG; Valine Val V GTG GTC GTT GTA; Tryptophan Trp W TGG; Tyrosine Tyr Y TAC TAT. Thus, the most preferred codon for alanine is "GCC," and the least is "GCG." Codon usage for various organisms and organelles can be found at the website <http://www.kazusa.or.jp/codon/>, incorporated herein by reference, allowing one of skill in the art to optimize codon usage for expression in various organisms using the disclosures herein. Thus, it is contemplated that codon usage may be optimized for other animals, as well as other organisms such as a prokaryote (e.g., an eubacteria), an archaea, an eukaryote (e.g., a protist, a plant, a fungus, an animal), a virus and the like, as well as organelles that contain nucleic acids, such as mitochondria or chloroplasts, based on the preferred codon usage as would be known to those of ordinary skill in the art.

[0079] It will also be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids or 5' or 3' sequences, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the

sequence meets the criteria set forth above, including the maintenance of biological protein, polypeptide or peptide activity. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region or may include various internal sequences, i.e., introns, which are known to occur within genes.

[0080] Excepting intronic or flanking regions, and allowing for the degeneracy of the genetic code, sequences that have about 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or about 99%, and any range derivable therein, such as, for example, about 50% to about 80%, and more preferably about 81% and about 90%; or even more preferably, between about 91% and about 99%; of nucleotides that are identical to the nucleotides of SEQ ID NO:1 will be sequences that are "essentially as set forth in SEQ ID NO:1".

a. Nucleic Acid Hybridization

[0081] The nucleic acid sequences disclosed herein also have a variety of uses. Contiguous sequences from vRNAP nucleic acid sequences can be used, for example, as templates to synthesize vRNAP.

[0082] Naturally, the present invention also encompasses DNA segments that are complementary, or essentially complementary, to the sequence set forth in SEQ ID NO:1, 3, 5, 7 and 14. Nucleic acid sequences that are "complementary" are those that are capable of base-pairing according to the standard Watson-Crick complementary rules. As used herein, the term "complementary sequences" means nucleic acid sequences that are complementary, as may be assessed by the same nucleotide comparison set forth above, or as defined as being capable of hybridizing to the nucleic acid segment of SEQ ID NO:1 under stringent conditions such as those described herein.

[0083] As used herein, a "DNA/RNA hybrid" is understood to mean that a single strand of RNA is hybridized to a single strand of DNA.

[0084] The term "appropriate reaction conditions" as described herein mean that temperature, pH, buffer, and other parameters are adjusted to optimize the reaction rate and yield.

[0085] As used herein, "hybridization," "hybridizes" or "capable of hybridizing" is understood to mean the forming of a double or triple stranded molecule or a molecule with partial double or triple stranded nature. The term "hybridization," "hybridize(s)" or "capable of hybridizing" encompasses the terms "stringent condition(s)" or "high stringency" and the terms "low stringency" or "low stringency condition(s)."

[0086] As used herein "stringent condition(s)" or "high stringency" are those conditions that allow hybridization between or within one or more nucleic acid strand(s) containing complementary sequence(s), but precludes hybridization of random sequences. Stringent conditions tolerate little, if any, mismatch between a nucleic acid and a target strand. Such conditions are well known to those of ordinary skill in the art, and are preferred for applications requiring high selectivity. Non-limiting applications include isolating a nucleic acid, such as a gene or a nucleic acid segment thereof, or detecting at least one specific mRNA transcript or a nucleic acid segment thereof, and the like.

[0087] Stringent conditions may comprise low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50° C to about 70° C. It is understood that the temperature and ionic strength of a desired stringency are determined in part by the length of the particular nucleic acid(s), the length and nucleobase content of the target sequence(s), the charge composition of the nucleic acid(s), and to the presence or concentration of formamide, tetramethylammonium chloride or other solvent(s) in a hybridization mixture.

[0088] It is also understood that these ranges, compositions and conditions for hybridization are mentioned by way of non-limiting examples only, and that the desired stringency for a particular hybridization reaction is often determined empirically by comparison to one or more positive or negative controls. Depending on the application envisioned it is preferred to employ varying conditions of hybridization to achieve varying degrees of selectivity of a nucleic acid towards a target sequence. In a non-limiting example, identification or isolation of a related target nucleic acid that does not hybridize to a nucleic acid under stringent conditions may be achieved by hybridization at low temperature and/or high ionic strength. For example, a medium stringency condition could be provided by about 0.1 to 0.25 M NaCl at temperatures of about 37° C to about 55° C. Under these conditions, hybridization may occur even though the sequences of probe and target strand are not perfectly complementary, but are mismatched at one or more positions. In another example, a low stringency condition could be provided by about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20° C to

about 55° C. Of course, it is within the skill of one in the art to further modify the low or high stringency conditions to suit a particular application. For example, in other embodiments, hybridization may be achieved under conditions of 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl₂, 1.0 mM dithiothreitol, at temperatures between approximately 20° C to about 37° C. Other hybridization conditions utilized could include approximately 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, at temperatures ranging from approximately 40° C to about 72° C.

[0089] Accordingly, the nucleotide sequences of the disclosure may be used for their ability to selectively form duplex molecules with complementary stretches of genes or RNAs or to provide primers for amplification of DNA or RNA from tissues. Depending on the application envisioned, it is preferred to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence.

[0090] The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, enhancers, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

[0091] For example, nucleic acid fragments may be prepared that include a contiguous stretch of nucleotides identical to or complementary to SEQ ID NO:1, 3, 5, 7 or 14. Nucleic acid fragments for use as a DNA transcription template may also be prepared. These fragments may be short or of intermediate lengths, such as, for example, about 8, about 10 to about 14, or about 15 to about 20 nucleotides, and that are chromosome-sized pieces, up to about 35,000, about 30,000, about 25,000, about 20,000, about 15,000, about 10,000, or about 5,000 base pairs in length, as well as DNA segments with total lengths of about 1,000, about 500, about 200, about 100 and about 50 base pairs in length (including all intermediate lengths of these lengths listed above, i.e., any range derivable therein and any integer derivable therein such a range) are also contemplated to be useful.

[0092] For example, it will be readily understood that "intermediate lengths," in these contexts, means any length between the quoted ranges, such as 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 70, 75, 80, 85, 90, 95,

100, 105, 110, 115, 120, 130, 140, 150, 160, 170, 180, 190, including all integers through the 200-500; 500-1,000; 1,000-2,000; 2,000-3,000; 3,000-5,000; 5,000-10,000 ranges, up to and including sequences of about 12,001, 12,002, 13,001, 13,002, 15,000, 20,000 and the like.

[0093] Various nucleic acid segments may be designed based on a particular nucleic acid sequence, and may be of any length. By assigning numeric values to a sequence, for example, the first residue is 1, the second residue is 2, etc., an algorithm defining all nucleic acid segments can be created:

[0094] n to $n+y$

[0095] where n is an integer from 1 to the last number of the sequence and y is the length of the nucleic acid segment minus one, where $n+y$ does not exceed the last number of the sequence. Thus, for a 10-mer, the nucleic acid segments correspond to bases 1 to 10, 2 to 11, 3 to 12 . . . and/or so on. For a 15-mer, the nucleic acid segments correspond to bases 1 to 15, 2 to 16, 3 to 17 . . . and/or so on. For a 20-mer, the nucleic segments correspond to bases 1 to 20, 2 to 21, 3 to 22 . . . and/or so on. In certain embodiments, the nucleic acid segment may be a probe or primer. As used herein, a "probe" generally refers to a nucleic acid used in a detection method or composition. As used herein, a "primer" generally refers to a nucleic acid used in an extension or amplification method or composition.

[0096] The use of a hybridization probe of between 17 and 100 nucleotides in length, or in some aspect of the invention even up to 1-2 Kb or more in length, allows the formation of a duplex molecule that is both stable and selective. Molecules having complementary sequences over stretches greater than 20 bases in length are generally preferred, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of particular hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having complementary sequences over stretches of 20 to 30 nucleotides, or even longer where desired. Such fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means or by introducing selected sequences into recombinant vectors for recombinant production.

[0097] In general, it is envisioned that the hybridization probes described herein will be useful both as reagents in solution hybridization, as in PCR, for detection of expression of corresponding genes, as well as in embodiments employing a solid phase. In embodiments involving a solid phase, the test DNA (or RNA) is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to hybridization with selected probes under desired conditions. The selected conditions will depend on the

particular circumstances based on the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, etc.). Following washing of the hybridized surface to remove non-specifically bound probe molecules, hybridization is detected, or even quantified, by means of the label.

b. Nucleic Acid Amplification

[0098] Nucleic acid used as a template for amplification is isolated from cells contained in the biological sample, according to standard methodologies (Sambrook *et al.*, In: Molecular Cloning: A Laboratory Manual 2 rev.ed., Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 1989). The nucleic acid may be genomic DNA or fractionated or whole cell RNA. Where RNA is used, it may be desired to convert the RNA to a complementary DNA. In one embodiment, the RNA is whole cell RNA and is used directly as the template for amplification.

[0099] Pairs of primers that selectively hybridize to nucleic acids are contacted with the isolated nucleic acid under conditions that permit selective hybridization. The term "primer," as defined herein, is meant to encompass any nucleic acid that is capable of priming the synthesis of a nascent nucleic acid in a template-dependent process. Typically, primers are oligonucleotides from ten to twenty or thirty base pairs in length, but longer sequences can be employed. Primers may be provided in double-stranded or single-stranded form, although the single-stranded form is preferred.

[00100] Once hybridized, the nucleic acid:primer complex is contacted with one or more enzymes that facilitate template-dependent nucleic acid synthesis. Multiple rounds of amplification, also referred to as "cycles," are conducted until a sufficient amount of amplification product is produced.

[00101] Next, the amplification product is detected. In certain applications, the detection may be performed by visual means. Alternatively, the detection may involve indirect identification of the product via chemiluminescence, radioactive scintigraphy of incorporated radiolabel or fluorescent label, or even via a system using electrical or thermal impulse signals (Affymax technology).

[00102] A number of template dependent processes are available to amplify the marker sequences present in a given template sample. One of the best known amplification methods is the polymerase chain reaction (referred to as PCR) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, each incorporated herein by reference in its entirety.

[00103] Briefly, in PCR, two primer sequences are prepared that are complementary to regions on opposite complementary strands of the marker sequence. An excess of deoxynucleoside triphosphates are added to a reaction mixture along with a DNA polymerase, e.g., Taq polymerase. If the marker sequence is present in a sample, the primers will bind to the marker and the polymerase will cause the primers to be extended along the marker sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the marker to form reaction products, excess primers will bind to the marker and to the reaction products, and the process is repeated.

[00104] A reverse transcriptase PCR amplification procedure may be performed in order to quantify the amount of mRNA amplified. Alternative methods for reverse transcription utilize thermostable, RNA-dependent DNA polymerases. These methods are described in WO 90/07641, filed Dec. 21, 1990, incorporated herein by reference. Polymerase chain reaction methodologies are well known in the art.

[00105] Another method for amplification is the ligase chain reaction ("LCR"), disclosed in EPA No. 320 308, incorporated herein by reference in its entirety. In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR, bound ligated units dissociate from the target and then serve as "target sequences" for ligation of excess probe pairs. U.S. Pat. No. 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence.

[00106] Q-beta Replicase, described in PCT Application No. PCT/US87/00880, incorporated herein by reference, may also be used as still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence, which can then be detected.

[00107] An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[alpha-thio]-triphosphates in one strand of a restriction site may also be useful in the amplification of nucleic acids in the present invention.

[00108] Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement

and synthesis, i.e., nick translation. A similar method, called Repair Chain Reaction (RCR), involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA. Target specific sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having 3' and 5' sequences of non-specific DNA and a middle sequence of specific RNA is hybridized to DNA that is present in a sample. Upon hybridization, the reaction is treated with RNase H, and the products of the probe identified as distinctive products that are released after digestion. The original template is annealed to another cycling probe and the reaction is repeated.

[00109] Still another amplification method described in GB Application No. 2 202 328, and in PCT Application No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template- and enzyme-dependent synthesis. The primers may be modified by labeling with a capture moiety (e.g., biotin) and/or a detector moiety (e.g., enzyme). In the latter application, an excess of labeled probes are added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

[00110] Other nucleic acid amplification procedures include transcription-based amplification systems (TAS), including nucleic acid sequence based amplification (NASBA) and 3SR (Gingeras *et al.*, PCT Application WO 88/10315, incorporated herein by reference). In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a clinical sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer which has target specific sequences. Following polymerization, DNA/RNA hybrids are digested with RNase H while double-stranded DNA molecules are heat denatured again. In either case, the single-stranded DNA is made fully double-stranded by addition of second target specific primer, followed by polymerization. The double-stranded DNA molecules are then multiply transcribed by an RNA polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into single-stranded DNA, which is then converted to double-stranded DNA, and

then transcribed once again with an RNA polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target specific sequences.

[00111] Davey *et al.*, EPA No. 329 822 (incorporated herein by reference in its entirety) disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a template for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from the resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in duplex with either DNA or RNA). The resultant ssDNA is a template for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to the template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting in a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be chosen to be in the form of either DNA or RNA.

[00112] Miller *et al.*, PCT Application WO 89/06700 (incorporated herein by reference in its entirety) disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This scheme is not cyclic, i.e., new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" and "one-sided PCR" (Frohman, In: PCR Protocols: A Guide To Methods And Applications, Academic Press, N.Y., 1990, incorporated herein by reference).

[00113] Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide," thereby amplifying the di-oligonucleotide, may also be used in the amplification step of the present invention.

c. Nucleic Acid Detection

[00114] In certain embodiments, it will be advantageous to employ nucleic acid sequences of the present invention such as all or part of SEQ ID NO:1, 3, 5, 7, 14 or a mutant

thereof in combination with an appropriate means, such as a label, for hybridization assays, RNase protection and Northern hybridization. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of being detected. In preferred embodiments, one may desire to employ a fluorescent label or an enzyme tag such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmentally undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a detection means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

[00115] In embodiments wherein nucleic acids are amplified, it may be desirable to separate the amplification product from the template and the excess primer for the purpose of determining whether specific amplification has occurred. In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods (Sambrook *et al.*, In: Molecular Cloning: A Laboratory Manual 2 rev.ed., Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 1989).

[00116] Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography which may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for using them including column, paper, thin-layer and gas chromatography.

[00117] Amplification products must be visualized in order to confirm amplification of the marker sequences. One typical visualization method involves staining of a gel with ethidium bromide and visualization under UV light. Alternatively, if the amplification products are integrally labeled with radio- or fluorometrically-labeled nucleotides, the amplification products can then be exposed to x-ray film or visualized under the appropriate stimulating spectra, following separation.

[00118] In one embodiment, visualization is achieved indirectly. Following separation of amplification products, a labeled, nucleic acid probe is brought into contact with the amplified marker sequence. The probe preferably is conjugated to a chromophore but may be radiolabeled. In another embodiment, the probe is conjugated to a binding partner, such as an antibody or biotin, and the other member of the binding pair carries a detectable moiety.

[00119] In one embodiment, detection is by Southern blotting and hybridization with a labeled probe. The techniques involved in Southern blotting are well known to those of skill in the art. Briefly, amplification products are separated by gel electrophoresis. The gel is then

contacted with a membrane, such as nitrocellulose, permitting transfer of the nucleic acid and non-covalent binding. Subsequently, the membrane is incubated with a chromophore-conjugated probe that is capable of hybridizing with a target amplification product. Detection is by exposure of the membrane to x-ray film or ion-emitting detection devices.

[00120] One example of the foregoing is described in U.S. Pat. No. 5,279,721, incorporated by reference herein, which discloses an apparatus and method for the automated electrophoresis and transfer of nucleic acids. The apparatus permits electrophoresis and blotting without external manipulation of the gel and is ideally suited to carrying out methods according to the present invention.

[00121] Other methods for genetic screening to accurately detect mutations in genomic DNA, cDNA or RNA samples may be employed, depending on the specific situation.

[00122] Historically, a number of different methods have been used to detect point mutations, including denaturing gradient gel electrophoresis ("DGGE"), restriction enzyme polymorphism analysis, chemical and enzymatic cleavage methods, and others. The more common procedures currently in use include direct sequencing of target regions amplified by PCR (see above) and single-strand conformation polymorphism analysis ("SSCP").

[00123] Another method of screening for point mutations is based on RNase cleavage of base pair mismatches in RNA/DNA and RNA/RNA heteroduplexes. As used herein, the term "mismatch" is defined as a region of one or more unpaired or mispaired nucleotides in a double-stranded RNA/RNA, RNA/DNA or DNA/DNA molecule. This definition thus includes mismatches due to insertion/deletion mutations, as well as single and multiple base point mutations.

[00124] U.S. Pat. No. 4,946,773 describes an RNase A mismatch cleavage assay that involves annealing single-stranded DNA or RNA test samples to an RNA probe, and subsequent treatment of the nucleic acid duplexes with RNase A. After the RNase cleavage reaction, the RNase is inactivated by proteolytic digestion and organic extraction, and the cleavage products are denatured by heating and analyzed by electrophoresis on denaturing polyacrylamide gels. For the detection of mismatches, the single-stranded products of the RNase A treatment, electrophoretically separated according to size, are compared to similarly treated control duplexes. Samples containing smaller fragments (cleavage products) not seen in the control duplex are scored as positive.

[00125] Currently available RNase mismatch cleavage assays, including those performed according to U.S. Pat. No. 4,946,773, require the use of radiolabeled RNA probes. Myers and

Maniatis in U.S. Pat. No. 4,946,773 describe the detection of base pair mismatches using RNase A. Other investigators have described the use of an *E. coli* enzyme, RNase I, in mismatch assays. Because it has broader cleavage specificity than RNase A, RNase I would be a desirable enzyme to employ in the detection of base pair mismatches if components can be found to decrease the extent of non-specific cleavage and increase the frequency of cleavage of mismatches. The use of RNase I for mismatch detection is described in literature from Promega Biotech. Promega markets a kit containing RNase I that is shown in their literature to cleave three out of four known mismatches, provided the enzyme level is sufficiently high.

[00126] The RNase Protection assay was first used to detect and map the ends of specific mRNA targets in solution. The assay relies on being able to easily generate high specific activity radiolabeled RNA probes complementary to the mRNA of interest by *in vitro* transcription. Originally, the templates for *in vitro* transcription were recombinant plasmids containing bacteriophage promoters. The probes are mixed with total cellular RNA samples to permit hybridization to their complementary targets, then the mixture is treated with RNase to degrade excess unhybridized probe. Also, as originally intended, the RNase used is specific for single-stranded RNA, so that hybridized double-stranded probe is protected from degradation. After inactivation and removal of the RNase, the protected probe (which is proportional in amount to the amount of target mRNA that was present) is recovered and analyzed on a polyacrylamide gel.

[00127] The RNase Protection assay was adapted for detection of single base mutations. In this type of RNase A mismatch cleavage assay, radiolabeled RNA probes transcribed *in vitro* from wild-type sequences are hybridized to complementary target regions derived from test samples. The test target generally comprises DNA (either genomic DNA or DNA amplified by cloning in plasmids or by PCR), although RNA targets (endogenous mRNA) have occasionally been used. If single nucleotide (or greater) sequence differences occur between the hybridized probe and target, the resulting disruption in Watson-Crick hydrogen bonding at that position ("mismatch") can be recognized and cleaved in some cases by single-strand specific ribonuclease. To date, RNase A has been used almost exclusively for cleavage of single-base mismatches, although RNase I has recently been shown as useful also for mismatch cleavage. There are recent descriptions of using the MutS protein and other DNA-repair enzymes for detection of single-base mismatches.

[00128] Nuclease S1 analysis of reaction products can be used to measure RNA. An exemplary procedure for S1 analysis involves hybridization reaction with the RNA of interest

(0.005-0.1 mg) and an excess of S1 probe which comprises a labeled oligonucleotide complementary to 20-80 or more sequential nucleotides of a specific RNA in S1 hybridization buffer (80% formamide, 0.4 M NaCl, 1 mM EDTA, 40 mM Pipes, pH 6.4). After denaturation for 4 min at 94° C, overnight hybridization at 30° C and precipitation with ethanol, the S1 probe/RNA mixture is resuspended in S1 buffer (0.26 M NaCl, 0.05 M sodium acetate, pH 4.6, and 4.5 mM zinc sulfate). The sample is divided into two volumes and 100 units of S1 nuclease (Sigma Chemical Company) is added to one tube. The samples are incubated for 60 minutes at 37° C; then EDTA (10 mM final concentration) and 15 g polyI-polyC RNA are added and the sample is extracted with phenol/chloroform and precipitated in ethanol. The samples are then subjected to polyacrylamide gel electrophoresis.

[00129] One method to produce a radiolabeled RNA probe with high specific activity includes admixing a radiolabeled NTP during transcription. Suitable isotopes for radiolabeling include ³⁵S- and ³²P-labeled UTP, GTP, CTP or ATP. For optimal results, a gel-purified radiolabeled RNA probe which is preferentially 300-500 bases in length, with a specific activity of 1-3 X10⁸ cpm/μg should be generated using the RNA polymerase of the current invention. In order to produce this *in vitro* transcript, it is often advisable to use a high specific activity (e.g., [α-³²P]CTP at 3,000Ci/mmol) NTP. To prevent background hybridization, it is important to remove plasmid template DNA by digestion which can be done with, for example, RQ1 RNase-Free DNase followed by phenol:chloroform:isoamyl alcohol extraction and ethanol precipitation.

[00130] Another method for producing radiolabeled probes includes using a riboprobe system which can produce high specific activity, radiolabeled RNA probes or microgram quantities of *in vitro* transcript. Riboprobes are useful with radiolabeled RNA probes in many applications including RNase protection, Northern hybridization, S1 analysis and *in situ* hybridization assays. The principle components of an *in vitro* transcription are the riboprobe, an RNA polymerase, a DNA template which includes a phage RNA polymerase promoter and ribonucleotide triphosphates.

d. Cloning vRNAP Genes

[00131] The present invention contemplates cloning vRNAP, or more particularly mini-vRNAP genes. A technique often employed by those skilled in the art of protein production today is to obtain a so-called "recombinant" version of the protein, to express it in a recombinant cell and to obtain the protein, polypeptide or peptide from such cells. These techniques are

based upon the "cloning" of a nucleic acid molecule encoding the protein from a DNA library, i.e., on obtaining a specific DNA molecule distinct from other portions of DNA. This can be achieved by, for example, cloning a cDNA molecule, or cloning a genomic-like DNA molecule.

[00132] The first step in such cloning procedures is the screening of an appropriate DNA library, such as, for example, from a phage, bacteria, yeast, fungus, mouse, rat, monkey or human. The screening protocol may utilize nucleotide segments or probes that are designed to hybridize to cDNA or genomic sequences of vRNAPs from protists. Additionally, antibodies designed to bind to the expressed vRNAP proteins, polypeptides, or peptides may be used as probes to screen an appropriate viral, eubacterial, archaeobacterial or eukaryotic DNA expression library. Alternatively, activity assays may be employed. The operation of such screening protocols are well known to those of skill in the art and are described in detail in the scientific literature, for example, in Sambrook *et al.*, In: *Molecular Cloning: A Laboratory Manual* 2 rev.ed., Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 1989, incorporated herein by reference. Moreover, as the present invention encompasses the cloning of genomic segments as well as cDNA molecules, it is contemplated that suitable genomic cloning methods, as known to those in the art, may also be used.

[00133] Encompassed by the invention are DNA segments encoding relatively small peptides, such as, for example, peptides of from about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 35, about 40, about 45, to about 50 amino acids in length, and more preferably, of from about 15 to about 30 amino acids in length; as set forth in SEQ ID NO:2, 4, 6, 8 or 15 and also larger polypeptides up to and including proteins corresponding to the full-length sequences set forth in SEQ ID NO:2 and SEQ ID NO:15, and any range derivable therein and any integer derivable in such a range. In addition to the "standard" DNA and RNA nucleotide bases, modified bases are also contemplated for use in particular applications of the present invention.

III. Recombinant Vectors, Promoters, Host Cells and Expression

[00134] Recombinant vectors form an important further aspect of the present invention. The term "expression vector or construct" means any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a proteinaceous molecule,

but it need not be, such as in the case of mini-vRNAP transcribing an RNA using a single-stranded DNA template. Thus, in certain embodiments, expression includes both transcription of a single-stranded DNA and translation of an RNA into the protein product. In other embodiments, expression only includes transcription of the nucleic acid. A recombinant vector can also be used for delivery of the RNA of the current invention.

[00135] Particularly useful vectors are contemplated to be those vectors in which the coding portion of the DNA segment, whether encoding a full length protein or smaller polypeptide or peptide, is positioned under the transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrases "operatively positioned," "under control" or "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

[00136] One particularly useful vector is pBAD. The pBAD expression vectors allow for greater control of bacterial expression of recombinant proteins and allow tight regulation for turning expression on or off. pBAD vectors allow for dose dependent induction for modulation of expression levels. The pBAD expression system helps overcome two of the most common problems of heterologous protein expression in bacteria: toxicity of the recombinant protein to the host and insolubility of the recombinant protein when it is expressed at high, uncontrolled levels. In both cases, a tightly-regulated expression system is critical for maximizing recombinant protein yields. The pBAD expression system is based on the araBAD operon which controls the arabinose metabolic pathway in *E. coli* and allows for precise modulation of heterologous expression to levels that are optimal for recovering high yields of the protein of interest (Guzman et al., J. Bact. 177:4121-4130, 1995).

a. Promoters

[00137] Any promoters normally found in a host cell in the native state can be used in the present invention to drive expression of N4 vRNA or mini-vRNA polymerase. Also, promoters not normally found in the host cell in the native state that are recognized by a native, normally native host cell RNA polymerase, or non-native RNA polymerase expressed in the cell can be used in the present invention to drive expression of the RNA polymerase. Other promoters may be selected from a nucleic acid sequence database accessible to those of skill in the art, e.g., GenBank, or the promoter can be isolated by a screening method. A promoter recognized by the

host cell can be operably linked to the gene or genes encoding the N4 RNA polymerase. The operable linkage can be constructed using any known techniques for DNA manipulation, as referred to herein.

[00138] Promoters are described as either constitutive or inducible. Constitutive promoters actively drive expression of genes under their control. Inducible promoters, in contrast, are activated in response to specific environmental stimuli. Both constitutive and inducible promoters can be used in the present invention for expressing non-host genes in a host cell.

[00139] Inducible promoters include, but are not limited to, trp, tac, lac, ara, reca, λPr, and λP1. These promoters and others that can be used in the present invention for expression of the N4 vRNA or mini-vRNA polymerase, in embodiments in which the host cell is *E. coli*, are described by Makrides, Microbiological Reviews 60, 512-538, 1996, herein incorporated by reference. Further, in embodiments of the present invention wherein the host cell is a microbe other than *E. coli*, such as *Saccharomyces*, *Bacillus*, and *Pseudomonas*, any inducible promoter known to those skilled in the art to be active in the host cell can be used to drive expression of the heterologous RNA polymerase. (U.S. Pat. No. 6,218,145).

[00140] The promoter may be in the form of the promoter that is naturally associated with N4 vRNA or mini-vRNA polymerase, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment or exon, for example, using recombinant cloning and/or PCR technology, in connection with the compositions disclosed herein (PCR technology is disclosed in U.S. Pat. No. 4,683,202 and U.S. Pat. No. 4,682,195, each incorporated herein by reference).

[00141] In other embodiments, it is contemplated that certain advantages will be gained by positioning the coding DNA segment under the control of a recombinant, or heterologous, promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with N4 vRNA or mini-vRNA polymerase in its natural environment. Such promoters may include promoters normally associated with other genes, and/or promoters isolated from any other bacterial, viral, eukaryotic, protist, or mammalian cell, and/or promoters made by the hand of man that are not "naturally occurring," i.e., containing different elements from different promoters, or mutations that increase, decrease, or alter expression.

[00142] Naturally, it will be important to employ a promoter that effectively directs the expression of the DNA segment in the cell type, organism, or even animal, chosen for

expression. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook *et al.* In: Molecular Cloning: A Laboratory Manual 2 rev.ed., Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 1989, incorporated herein by reference. The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins, polypeptides or peptides.

[00143] At least one module in a promoter generally functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

[00144] Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the thymidine kinase promoter, the spacing between promoter elements can be increased to 50 base pairs apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

[00145] The particular promoter that is employed to control the expression of a nucleic acid is not believed to be critical, so long as it is capable of expressing the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a human or viral promoter.

[00146] In various other embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter and the Rous sarcoma virus long terminal repeat can be used to obtain high-level expression of the instant nucleic acids. The use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression are contemplated as well, provided that the levels of expression are sufficient for a given purpose.

[00147] In certain embodiments of the invention, promoter sequences may be used that are recognized specifically by a DNA-dependent RNA polymerase, such as, but not limited to, those described by Chamberlin and Ryan, In: *The Enzymes*. San Diego, Calif., Academic Press, 15:87-108, 1982, and by Jorgensen *et al.*, *J. Biol. Chem.* 266:645-655, 1991. These promoters can be used to express a wild-type or mutant form of a miniV RNA polymerase of the invention. Several RNA polymerase promoter sequences are especially useful, including, but not limited to, promoters derived from SP6 (e.g., Zhou and Doetsch, *Proc. Nat. Acad. Sci. USA* 90:6601-6605, 1993), T7 (e.g., Martin, and Coleman, *Biochemistry* 26:2690-2696, 1987) and T3 (e.g., McGraw *et al.*, *Nucl. Acid. Res.* 13:6753-6766, 1985). An RNA polymerase promoter sequence derived from *Thermus thermophilus* can also be used (see, e.g., Wendt *et al.*, *Eur. J. Biochem.* 191:467-472, 1990; Faraldo *et al.*, *J. Bact.* 174:7458-7462, 1992; Hartmann *et al.*, *Biochem.* 69:1097-1104, 1987; Hartmann *et al.*, *Nucl. Acids Res.* 19:5957-5964, 1991). The length of the promoter sequence will vary depending upon the promoter chosen. For example, the T7 RNA polymerase promoter can be only about 25 bases in length and act as a functional promoter, while other promoter sequences require 50 or more bases to provide a functional promoter.

[00148] In other embodiments of the invention, a promoter is used that is recognized by an RNA polymerase from a T7-like bacteriophage. The genetic organization of all T7-like phages that have been examined has been found to be essentially the same as that of T7. Examples of T7-like phages according to the invention include, but are not limited to *Escherichia coli* phages T3, ϕ I, ϕ II, W31, H, Y, A1, 122, cro, C21, C22, and C23; *Pseudomonas putida* phage gh-1; *Salmonella typhimurium* phage SP6; *Serratia marcescens* phages IV; *Citrobacter* phage ViIII; and *Klebsiella* phage No. 11 (Hausmann, *Current Topics in Microbiology and Immunology* 75:77-109, 1976; Korsten *et al.*, *J. Gen. Virol.* 43:57-73, 1975; Dunn, *et al.*, *Nature New Biology* 230:94-96, 1971; Towle, *et al.*, *J. Biol. Chem.* 250:1723-1733, 1975; Butler and Chamberlin, *J. Biol. Chem.* 257:5772-5778, 1982).

[00149] When a T7 RNA polymerase promoter, or another T7-like RNA polymerase promoter is used to express a wild-type or mutant form of a gene for a miniV RNA polymerase of the invention, the gene can be expressed in a host cell which expresses the T7 RNA polymerase, or the corresponding T7-like RNA polymerase for the promoter used, wherein the RNA polymerase for the promoter is expressed either constitutively, or more preferably, from an inducible promoter. By way of example, a T7 RNA polymerase expression system, such as, but

not limited to, the expression systems disclosed in, for example, U.S. Pat. Nos. 5,693,489 and 5,869,320, the disclosures of which are incorporated herein by reference in their entirety.

b. Enhancers

[00150] Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of DNA with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

[00151] The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

[00152] Additionally any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base EPDB, <http://www.epd.isb-sib.ch/>) could also be used to drive expression.

[00153] Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[00154] Turning to the expression of the proteinaceous molecules after transcription using the vRNAP, mini-vRNAP, or mutants thereof of the present invention, once a suitable clone or clones have been obtained, whether they be cDNA based or genomic, one may proceed to prepare an expression system.

[00155] The engineering of DNA segment(s) for expression in a prokaryotic or eukaryotic system may be performed by techniques generally known to those of skill in recombinant expression. It is believed that virtually any expression system may be employed in the expression of the proteinaceous molecules of the present invention.

[00156] Both cDNA and genomic sequences are suitable for eukaryotic expression, as the host cell will generally process the genomic transcripts to yield functional mRNA for translation into proteinaceous molecules. Generally speaking, it may be more convenient to employ as the recombinant gene a cDNA version of the gene. It is believed that the use of a cDNA version

will provide advantages in that the size of the gene will generally be much smaller and more readily employed to transfect the targeted cell than will a genomic gene, which will typically be up to an order of magnitude or more larger than the cDNA gene. However, it is contemplated that a genomic version of a particular gene may be employed where desired.

[00157] In expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal and the bovine growth hormone polyadenylation signal, convenient and known to function well in various target cells. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

c. Antisense, RNAi and Ribozymes

[00158] In some embodiments of the invention the vRNA polymerase can be used to synthesize antisense RNA, RNAi or interference RNA or ribozymes.

[00159] The term "antisense nucleic acid" is intended to refer to the oligonucleotides complementary to the base sequences of DNA and RNA. Antisense oligonucleotides, when introduced into a target cell, specifically bind to their target nucleic acid and interfere with transcription, RNA processing, transport, translation, and/or stability. Targeting double-stranded (ds) DNA with oligonucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. An antisense nucleic acid may be complementary to SEQ ID NO:1, 3, 5, 7 or 14, complementary to a mini-vRNAP encoding sequence or to mini-vRNAP non-coding sequences. Antisense RNA constructs, or DNA encoding such antisense RNAs, may be employed to inhibit gene transcription or translation or both within a host cell, either *in vitro* or *in vivo*, such as within a host animal, including a human subject.

[00160] Antisense constructs may be designed to bind to the promoter and other control regions, exons, introns or even exon-intron boundaries (splice junctions) of a gene. It is contemplated that the most effective antisense constructs may include regions complementary to intron/exon splice junctions. Thus, antisense constructs with complementary regions within 50-200 bases of an intron-exon splice junction may be used. It has been observed that some exon sequences can be included in the construct without seriously affecting the target selectivity thereof. The amount of exonic material included will vary depending on the particular exon and

intron sequences used. One can readily test whether too much exon DNA is included simply by testing the constructs *in vitro* to determine whether normal cellular function is affected or whether the expression of related genes having complementary sequences is affected.

[00161] As stated above, "complementary" or "antisense" means polynucleotide sequences that are substantially complementary over their entire length and have very few base mismatches. For example, sequences of fifteen bases in length may be termed complementary when they have complementary nucleotides at thirteen or fourteen positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire length and have no base mismatches. Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology, but also contains a non-homologous region (e.g., ribozyme) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

[00162] It may be advantageous to combine portions of genomic DNA with cDNA or synthetic sequences to generate specific constructs. For example, where an intron is desired in the ultimate construct, a genomic clone will need to be used. The cDNA or a synthesized polynucleotide may provide more convenient restriction sites for the remaining portion of the construct and, therefore, would be used for the rest of the sequence.

[00163] While all or part of the gene sequence may be employed in the context of antisense construction, statistically, any sequence 17 bases long should occur only once in the human genome and, therefore, suffice to specify a unique target sequence. Although shorter oligomers are easier to make and increase *in vivo* accessibility, numerous other factors are involved in determining the specificity of hybridization. Both binding affinity and sequence specificity of an oligonucleotide to its complementary target increases with increasing length. It is contemplated that oligonucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more base pairs will be used. One can readily determine whether a given antisense nucleic acid is effective at targeting of the corresponding host cell gene simply by testing the constructs *in vivo* to determine whether the endogenous gene's function is affected or whether the expression of related genes having complementary sequences is affected.

[00164] In certain embodiments, one may wish to employ antisense constructs which include other elements, for example, those which include C-5 propyne pyrimidines. Oligonucleotides which contain C-5 propyne analogues of uridine and cytidine have been shown

to bind RNA with high affinity and to be potent antisense inhibitors of gene expression (Wagner *et al.*, Science 260:1510-1513, 1993).

[00165] As an alternative to targeted antisense delivery, targeted ribozymes may be used. The term "ribozyme" refers to an RNA-based enzyme capable of targeting and cleaving particular base sequences in oncogene DNA and RNA. Ribozymes either can be targeted directly to cells, in the form of RNA oligonucleotides incorporating ribozyme sequences, or introduced into the cell as an expression construct encoding the desired ribozymal RNA. Ribozymes may be used and applied in much the same way as described for antisense nucleic acids. Sequences for ribozymes may be included in the DNA template to eliminate undesired 5' end sequences in RNAs generated through T7 RNA polymerase transcription.

[00166] Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, Proc. Natl. Acad. Sci. USA 84:8788-8792, 1987; Gerlach *et al.*, Nature 328:802-805, 1987; Forster and Symons, Cell 49:211-220, 1987). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, Cell 27:487-496, 1981; Michel and Westhof, J. Mol. Biol. 216:585-610, 1990; Reinhold-Hurek and Shub, Nature 357:173-176, 1992). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

[00167] Ribozyme catalysis has primarily been observed as part of sequence specific cleavage/ligation reactions involving nucleic acids (Joyce, Nature 338:217-244, 1989; Cech *et al.*, Cell 27:487-496, 1981). For example, U.S. Pat. No. 5,354,855 reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that of known ribonucleases and approaching that of the DNA restriction enzymes. Thus, sequence-specific ribozyme-mediated inhibition of gene expression may be particularly suited to therapeutic applications (Scanlon *et al.*, Proc. Natl. Acad. Sci. USA 88:10591-10595, 1991; Sarver *et al.*, Science 247:1222-1225, 1990; Sioud *et al.*, J. Mol. Biol. 223:831-835, 1992). Recently, it was reported that ribozymes elicited genetic changes in some cell lines to which they were applied; the altered genes included the oncogenes H-ras, c-fos and genes of HIV. Most of this work involved the modification of a target mRNA, based on a specific mutant codon that is cleaved by a specific ribozyme. In light of the information included herein and the knowledge of one of ordinary skill

in the art, the preparation and use of additional ribozymes that are specifically targeted to a given gene will now be straightforward.

[00168] Several different ribozyme motifs have been described with RNA cleavage activity (reviewed in Symons, Annu. Rev. Biochem. 61:641-671, 1992). Examples of ribozymes include sequences from the Group I self-splicing introns including tobacco ringspot virus (Prody, et al., Science 231:1577-1580, 1986), avocado sunblotch viroid (Palukaitis, et al., Virology 99:145-151, 1979; Symons, Nucl. Acids Res. 9:6527-6537, 1981), and Lucerne transient streak virus (Forster and Symons, Cell 49:211-220, 1987). Sequences from these and related viruses are referred to as hammerhead ribozymes based on a predicted folded secondary structure.

[00169] Other suitable ribozymes include sequences from RNase P with RNA cleavage activity (Yuan, et al., Proc. Natl. Acad. Sci. USA 89:8006-8010, 1992; Yuan and Altman, Science, 263:1269-1273, 1994), hairpin ribozyme structures (Berzal-Herranz, et al., Genes and Devel. 6:129-134, 1992; Chowrira et al., Biochemistry 32:1088-1095, 1993) and hepatitis δ virus based ribozymes (Perrotta and Been, Biochemistry 31:16-21, 1992). The general design and optimization of ribozyme directed RNA cleavage activity has been discussed in detail (Haseloff and Gerlach, Nature 334:585-591, 1988; Symons, Annu. Rev. Biochem. 61:641-671, 1992; Chowrira, et al., J. Biol. Chem. 269:25856-25864, 1994; and Thompson, et al., Nature Medicine 1:277-278, 1995).

[00170] The other variable on ribozyme design is the selection of a cleavage site on a given target RNA. Ribozymes are targeted to a given sequence by virtue of annealing to a site by complementary base pair interactions. Two stretches of homology are required for this targeting. These stretches of homologous sequences flank the catalytic ribozyme structure defined above. Each stretch of homologous sequence can vary in length from 7 to 15 nucleotides. The only requirement for defining the homologous sequences is that, on the target RNA, they are separated by a specific sequence which is the cleavage site. For hammerhead ribozymes, the cleavage site is a dinucleotide sequence on the target RNA, uracil (U) followed by either an adenine, cytosine or uracil (A, C or U; Perriman, et al., Gene 113:157-163, 1992; Thompson, et al., Nature Medicine 1:277-278, 1995). The frequency of this dinucleotide occurring in any given RNA is statistically 3 out of 16. Therefore, for a given target messenger RNA of 1000 bases, 187 dinucleotide cleavage sites are statistically possible.

[00171] Designing and testing ribozymes for efficient cleavage of a target RNA is a process well known to those skilled in the art. Examples of scientific methods for designing and

testing ribozymes are described by Chowrira *et al.*, *J. Biol. Chem.* 269:25856-25864 (1994) and Lieber and Strauss, *Mol. Cell. Biol.* 15: 540-551 (1995), each incorporated by reference. The identification of operative and preferred sequences for use in ribozymes is simply a matter of preparing and testing a given sequence, and is a routinely practiced "screening" method known to those of skill in the art.

[00172] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements.

d. Host Cells

[00173] Host cells may be derived from prokaryotes or eukaryotes, including yeast cells, insect cells, and mammalian cells, depending upon whether the desired result is replication of the vector or expression of part or all of the vector-encoded nucleic acid sequences. Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials (www.atcc.org). An appropriate host can be determined by one of skill in the art based on the vector backbone and the desired result. A plasmid or cosmid, for example, can be introduced into a prokaryotic host cell for replication of many vector copies. Bacterial cells used as host cells for vector replication and/or expression include DH5 α , BL 21, JM109, and KC8, as well as a number of commercially available bacterial hosts such as SURE[®] Competent Cells and SOLOPACK Gold Cells (STRATAGENE[®], La Jolla, Calif.). Alternatively, bacterial cells such as *E. coli* LE392 could be used as host cells. Appropriate yeast cells include *Saccharomyces cerevisiae*, *Saccharomyces pombe*, and *Pichia pastoris*.

[00174] Examples of eukaryotic host cells for replication and/or expression of a vector include HeLa, NIH3T3, Jurrat, 293, Cos, CHO, Saos, BHK, C127 and PC12. Many host cells from various cell types and organisms are available and would be known to one of skill in the

art. Similarly, a viral vector may be used in conjunction with either a eukaryotic or prokaryotic host cell, particularly one that is permissive for replication or expression of the vector.

[00175] Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and/or their polypeptides, proteins, or peptides.

[00176] It is proposed that vRNAP, or more particularly mini-vRNAP may be co-expressed with other selected proteinaceous molecules such as EcoSSB and other proteins of interest, wherein the proteinaceous molecules may be co-expressed in the same cell or vRNAP gene may be provided to a cell that already has another selected proteinaceous molecule. Co-expression may be achieved by co-transfecting the cell with two distinct recombinant vectors, each bearing a copy of either of the respective DNAs. Alternatively, a single recombinant vector may be constructed to include the coding regions for both of the proteinaceous molecules, which could then be expressed in cells transfected with the single vector. In either event, the term "co-expression" herein refers to the expression of both the vRNAP gene and the other selected proteinaceous molecules in the same recombinant cell.

[00177] As used herein, the terms "engineered" and "recombinant" cells or host cells are intended to refer to a cell into which an exogenous DNA segment or gene, such as a cDNA or gene encoding vRNAP, mini-vRNAP or a mutant thereof, has been introduced. Therefore, engineered cells are distinguishable from naturally occurring cells which do not contain a recombinantly introduced exogenous DNA segment or gene. Engineered cells are thus cells having a gene or genes introduced through the hand of man. Recombinant cells include those having an introduced cDNA or genomic gene, and also include genes positioned adjacent to a promoter not naturally associated with the particular introduced gene.

[00178] To express a recombinant vRNAP, whether mutant or wild-type, in accordance with the present invention one would prepare an expression vector that comprises a wild-type, or mutant vRNAP proteinaceous molecule-encoding nucleic acid under the control of one or more promoters. To bring a coding sequence "under the control of" a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame generally between about 1 and about 50 nucleotides "downstream" of the chosen promoter. The "upstream"

promoter directs transcription of the DNA and promotes expression of the encoded recombinant protein, polypeptide or peptide. This is the meaning of "recombinant expression" in this context.

[00179] Many standard techniques are available to construct expression vectors containing the appropriate nucleic acids and transcriptional/translational control sequences in order to achieve protein, polypeptide or peptide expression in a variety of host expression systems. Cell types available for expression include, but are not limited to, bacteria, such as *E. coli* and *B. subtilis*, transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors.

[00180] Certain examples of prokaryotic hosts are *E. coli* strain RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776 (ATCC No. 31537) as well as *E. coli* W3110 (F-, lambda-, prototrophic, ATCC No. 273325); bacilli such as *Bacillus subtilis*; and other enterobacteriaceae such as *Salmonella typhimurium*, *Serratia marcescens*, and various *Pseudomonas* species.

[00181] In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication origin, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of its own proteins.

[00182] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM™-11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as *E. coli* LE392.

[00183] Further useful vectors include pIN vectors; and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble proteins for later purification and separation or cleavage.

[00184] The following details concerning recombinant protein production in bacterial cells, such as *E. coli*, are provided by way of exemplary information on recombinant protein production in general, the adaptation of which to a particular recombinant expression system will be known to those of skill in the art.

[00185] Bacterial cells, for example, *E. coli*, containing the expression vector are grown in any of a number of suitable media, for example, LB. The expression of the recombinant proteinaceous molecule may be induced, e.g., by adding IPTG or any appropriate inducer to the media or by switching incubation to a higher temperature, depending on the regulated promoter used. After culturing the bacteria for a further period, generally of between 2 and 24 hours, the cells are collected by centrifugation and washed to remove residual media.

[00186] The bacterial cells are then lysed, for example, by disruption in a cell homogenizer, by sonication or cell press and centrifuged to separate the dense inclusion bodies and cell membranes from the soluble cell components. This centrifugation can be performed under conditions whereby the dense inclusion bodies are selectively enriched by incorporation of sugars, such as sucrose, into the buffer and centrifugation at a selective speed.

[00187] If the recombinant proteinaceous molecule is expressed in the inclusion bodies, as is the case in many instances, these can be washed in any of several solutions to remove some of the contaminating host proteins, then solubilized in solutions containing high concentrations of urea (e.g., 8M) or chaotropic agents such as guanidine hydrochloride in the presence of reducing agents, such as β -mercaptoethanol or DTT (dithiothreitol).

[00188] Under some circumstances, it may be advantageous to incubate the proteinaceous molecule for several hours under conditions suitable for the proteinaceous molecule to undergo a refolding process into a conformation which more closely resembles that of the native proteinaceous molecule. Such conditions generally include low proteinaceous molecule concentrations, less than 500 mg/ml, low levels of reducing agent, concentrations of urea less than 2 M and often the presence of reagents such as a mixture of reduced and oxidized glutathione which facilitate the interchange of disulfide bonds within the proteinaceous molecule.

[00189] The refolding process can be monitored, for example, by SDS-PAGE, or with antibodies specific for the native molecule (which can be obtained from animals vaccinated with the native molecule or smaller quantities of recombinant proteinaceous molecule). Following refolding, the proteinaceous molecule can then be purified further and separated from the refolding mixture by chromatography on any of several supports including ion exchange resins, gel permeation resins or on a variety of affinity columns.

[00190] For expression in *Saccharomyces*, the plasmid YRp7, for example, is commonly used. This plasmid already contains the *trp1* gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076

or PEP4-1. The presence of the *trp1* lesion as a characteristic of the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

[00191] Suitable promoter sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate protein, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector downstream of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination.

[00192] In addition to micro-organisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from vertebrate or invertebrate culture. In addition to mammalian cells, these include insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus); and plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing one or more RNAP coding sequences.

[00193] Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteinaceous molecules. Appropriate cells lines or host systems can be chosen to ensure the correct modification and processing of the foreign proteinaceous molecule expressed.

[00194] A number of viral-based expression systems may be utilized, for example, commonly used promoters are derived from polyoma, Adenovirus 2, and most frequently Simian Virus 40 (SV40). The early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication. Smaller or larger SV40 fragments may also be used, provided there is included the approximately 250 bp sequence extending from the HindIII site toward the BglI site located in the viral origin of replication.

[00195] In cases where an adenovirus is used as an expression vector, the coding sequences may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of

the viral genome (e.g., region E1, E3, or E4) will result in a recombinant virus that is viable and capable of expressing an RNA in infected hosts.

[00196] Specific initiation signals may also be used for more efficient translation using the vRNAP of the current invention. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may additionally need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be in-frame (or in-phase) with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements and transcription terminators.

[00197] In eukaryotic expression, one will also typically desire to incorporate into the transcriptional unit an appropriate polyadenylation site (e.g., 5'-AATAAA-3') if one was not contained within the original cloned segment. Typically, the poly A addition site is placed about 30 to 2000 nucleotides "downstream" of the termination site of the proteinaceous molecule at a position prior to transcription termination.

[00198] For long-term, high-yield production of a recombinant vRNAP protein, polypeptide or peptide, stable expression is preferred. For example, cell lines that stably express constructs encoding a vRNAP protein, polypeptide or peptide may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with vectors controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

[00199] A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (tk), hypoxanthine-guanine phosphoribosyltransferase (hgprt) and adenine phosphoribosyltransferase (aprt) genes, in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dihydrofolate reductase (dhfr), that confers resistance to methotrexate; gpt, that confers

resistance to mycophenolic acid; neomycin (neo), that confers resistance to the aminoglycoside G-418; and hygromycin (hygro), that confers resistance to hygromycin.

[00200] Large scale suspension culture of bacterial cells in stirred tanks is a common method for production of recombinant proteinaceous molecules. Two suspension culture reactor designs are in wide use--the stirred reactor and the airlift reactor. The stirred design has successfully been used on an 8000 liter capacity for the production of interferon. Cells are grown in a stainless steel tank with a height-to-diameter ratio of 1:1 to 3:1. The culture is usually mixed with one or more agitators, based on bladed disks or marine propeller patterns. Agitator systems offering less shear forces than blades have been described. Agitation may be driven either directly or indirectly by magnetically coupled drives. Indirect drives reduce the risk of microbial contamination through seals on stirrer shafts.

[00201] The airlift reactor for microbial fermentation relies on a gas stream to both mix and oxygenate the culture. The gas stream enters a riser section of the reactor and drives circulation. Gas disengages at the culture surface, causing denser liquid free of gas bubbles to travel downward in the downcomer section of the reactor. The main advantage of this design is the simplicity and lack of need for mechanical mixing. Typically, the height-to-diameter ratio is 10:1. The airlift reactor scales up relatively easily, has good mass transfer of gases and generates relatively low shear forces.

[00202] It is contemplated that the vRNAP proteins, polypeptides or peptides of the invention may be "overexpressed," i.e., expressed in increased levels relative to its natural expression in cells. Such overexpression may be assessed by a variety of methods, including radio-labeling and/or proteinaceous molecule purification. However, simple and direct methods are preferred, for example, those involving SDS/PAGE and proteinaceous composition staining or western blotting, followed by quantitative analyses, such as densitometric scanning of the resultant gel or blot. A specific increase in the level of the recombinant protein, polypeptide or peptide in comparison to the level in natural cells is indicative of overexpression, as is a relative abundance of the specific proteinaceous molecule in relation to the other proteins produced by the host cell and, e.g., visible on a gel.

IV. Methods of Gene Transfer

[00203] In order to mediate the effect of transgene expression in a cell, it will be necessary to transfer the expression constructs (e.g., a therapeutic construct) of the present invention into a cell. Such transfer may employ viral or non-viral methods of gene transfer.

This section provides a discussion of methods and compositions of gene or nucleic acid transfer, including transfer of antisense sequences.

[00204] The vRNAP genes are incorporated into a viral vector to mediate gene transfer to a cell. Additional expression constructs encoding EcoSSB and other therapeutic agents as described herein may also be transferred via viral transduction using infectious viral particles, for example, by transformation with an adenovirus vector of the present invention.

Alternatively, a retrovirus, bovine papilloma virus, an adeno-associated virus (AAV), a lentiviral vector, a vaccinia virus, a polyoma virus, or an infective virus that has been engineered to express a specific binding ligand may be used. Similarly, nonviral methods which include, but are not limited to, direct delivery of DNA such as by injection, electroporation, calcium phosphate precipitation, liposome mediated transfection, and microprojectile bombardment may be employed. Thus, in one example, viral infection of cells is used in order to deliver therapeutically significant genes to a cell. Typically, the virus simply will be exposed to the appropriate host cell under physiologic conditions, permitting uptake of the virus.

[00205] Microinjection can be used for delivery into a cell. Microinjection involves the insertion of a substance such as RNA into a cell through a microelectrode. Typical applications include the injection of drugs, histochemical markers (such as horseradish peroxidase or lucifer yellow) and RNA or DNA in molecular biological studies. To extrude the substances through the very fine electrode tips, either hydrostatic pressure (pressure injection) or electric currents (iontophoresis) is employed.

V. Proteinaceous Compositions

[00206] In certain embodiments, the present invention concerns novel compositions or methods comprising at least one proteinaceous molecule. The proteinaceous molecule may have a sequence essentially as set forth in SEQ ID NO:2, 4, 6, 8 or 15. The proteinaceous molecule may be a vRNAP or more preferably a mini-vRNAP, or a delivery agent. The proteinaceous molecule may also be a mutated mini-vRNAP.

[00207] As used herein, a "proteinaceous molecule," "proteinaceous composition," "proteinaceous compound," "proteinaceous chain" or "proteinaceous material" generally refers to, but is not limited to, a protein of greater than about 200 amino acids or the full length endogenous sequence translated from a gene; a polypeptide of greater than about 100 amino acids; and/or a peptide of from about 3 to about 100 amino acids. All the "proteinaceous" terms described above may be used interchangeably herein.

[00208] In certain embodiments the size of the at least one proteinaceous molecule may comprise, but is not limited to, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, about 60, about 61, about 62, about 63, about 64, about 65, about 66, about 67, about 68, about 69, about 70, about 71, about 72, about 73, about 74, about 75, about 76, about 77, about 78, about 79, about 80, about 81, about 82, about 83, about 84, about 85, about 86, about 87, about 88, about 89, about 90, about 91, about 92, about 93, about 94, about 95, about 96, about 97, about 98, about 99, about 100, about 110, about 120, about 130, about 140, about 150, about 160, about 170, about 180, about 190, about 200, about 210, about 220, about 230, about 240, about 250, about 275, about 300, about 325, about 350, about 375, about 400, about 425, about 450, about 475, about 500, about 525, about 550, about 575, about 600, about 625, about 650, about 675, about 700, about 725, about 750, about 775, about 800, about 825, about 850, about 875, about 900, about 925, about 950, about 975, about 1000, about 1100, about 1200, about 1300, about 1400, about 1500, about 1750, about 2000, about 2250, about 2500 or greater amino molecule residues, and any range derivable therein.

[00209] As used herein, an "amino molecule" refers to any amino acid, amino acid derivative or amino acid mimic as would be known to one of ordinary skill in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

[00210] Accordingly, the term "proteinaceous composition" encompasses amino molecule sequences comprising at least one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid, including but not limited to the following, beginning with the corresponding abbreviation: Aad 2-Aminoadipic acid; EtAsn N-Ethylasparagine; Baad 3-Aminoadipic acid; Hyl Hydroxylysine; Bala β -alanine, β -Amino-propionic acid; AHyl allo-Hydroxylysine; Abu 2-Aminobutyric acid; 3Hyp 3-Hydroxyproline; 4Abu 4-Aminobutyric acid, piperidinic; 4Hyp 4-Hydroxy-proline acid; Acp 6-Aminocaproic

acid; Ide Isodesmosine; Ahe 2-Amino-heptanoic acid; Alle allo-Isoleucine; Aib 2-Aminoisobutyric acid; MeGly N-Methylglycine, sarcosine; Baib 3-Aminoisobutyric acid; MeIle N-Methylisoleucine; Apm 2-Aminopimelic acid; MeLys 6-N-Methyllysine; Dbu 2,4-Diaminobutyric acid; MeVal N-Methylvaline; Des Desmosine; Nva Norvaline; Dpm 2,2'-Diaminopimelic acid; Nle Norleucine; Dpr 2,3-Diaminopropionic acid; Orn Ornithine EtGly N-Ethylglycine.

[00211] In certain embodiments the proteinaceous composition comprises at least one protein, polypeptide or peptide, such as vRNAP or mini-vRNAP. In further embodiments the proteinaceous composition comprises a biocompatible protein, polypeptide or peptide. As used herein, the term "biocompatible" refers to a substance which produces no significant untoward effects when applied to, or administered to, a given organism according to the methods and amounts described herein. Such untoward or undesirable effects are those such as significant toxicity or adverse immunological reactions. In preferred embodiments, biocompatible protein, polypeptide or peptide containing compositions will generally be mammalian proteins or peptides or synthetic proteins or peptides each essentially free from toxins, pathogens and harmful immunogens.

[00212] Proteinaceous compositions may be made by any technique known to those of skill in the art, including the expression of proteins, polypeptides or peptides through standard molecular biological techniques, the isolation of proteinaceous compounds from natural sources, or the chemical synthesis of proteinaceous materials. The nucleotide and protein, polypeptide and peptide sequences for various genes have been previously disclosed, and may be found at computerized databases known to those of ordinary skill in the art. One such database is the National Center for Biotechnology Information's Genbank and GenPept databases (<http://www.ncbi.nlm.nih.gov/>). The coding regions for these known genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art. Alternatively, various commercial preparations of proteins, polypeptides and peptides are known to those of skill in the art.

[00213] In certain embodiments, a proteinaceous compound may be purified. Generally, "purified" will refer to a specific or desired protein, polypeptide, or peptide composition that has been subjected to fractionation to remove various other proteins, polypeptides, or peptides, and which composition substantially retains its activity, as may be assessed, for example, by the protein assays, as would be known to one of ordinary skill in the art for the specific or desired protein, polypeptide or peptide.

[00214] In certain embodiments, the proteinaceous composition may comprise at least one antibody. A mini-vRNAP antibody may comprise all or part of an antibody that specifically recognizes mini-vRNAP. As used herein, the term "antibody" is intended to refer broadly to any immunologic binding agent such as IgG, IgM, IgA, IgD and IgE. Generally, IgG and/or IgM are preferred because they are the most common antibodies in the physiological situation and because they are most easily made in a laboratory setting.

[00215] The term "antibody" is used to refer to any antibody-like molecule that has an antigen binding region, and includes antibody fragments such as Fab', Fab, F(ab')₂, single domain antibodies (DABs), Fv, scFv (single chain Fv), and the like. The techniques for preparing and using various antibody-based constructs and fragments are well known in the art. Means for preparing and characterizing antibodies are also well known in the art (See, e.g., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988; incorporated herein by reference).

[00216] It is contemplated that virtually any protein, polypeptide or peptide containing component may be used in the compositions and methods disclosed herein. However, it is preferred that the proteinaceous material is biocompatible. In certain embodiments, it is envisioned that the formation of a more viscous composition will be advantageous in that the high viscosity will allow the composition to be more precisely or easily applied to the tissue and to be maintained in contact with the tissue throughout the procedure. In such cases, the use of a peptide composition, or more preferably, a polypeptide or protein composition, is contemplated. Ranges of viscosity include, but are not limited to, about 40 to about 100 poise. In certain aspects, a viscosity of about 80 to about 100 poise is preferred.

[00217] Proteins and peptides suitable for use in this invention may be autologous proteins or peptides, although the invention is clearly not limited to the use of such autologous proteins. As used herein, the term "autologous protein, polypeptide or peptide" refers to a protein, polypeptide or peptide which is derived or obtained from an organism. Organisms that may be used include, but are not limited to, a bovine, a reptilian, an amphibian, a piscine, a rodent, an avian, a canine, a feline, a fungal, a plant, or a prokaryotic organism, with a selected animal or human subject being preferred. The "autologous protein, polypeptide or peptide" may then be used as a component of a composition intended for application to the selected animal or human subject. In certain aspects, the autologous proteins or peptides are prepared, for example from whole plasma of the selected donor. The plasma is placed in tubes and placed in a freezer at about -80° C for at least about 12 hours and then centrifuged at about 12,000 times g for about

15 minutes to obtain the precipitate. The precipitate, such as fibrinogen may be stored for up to about one year.

VI. Protein Purification

[00218] To prepare a composition comprising a vRNAP or mini-vRNAP, it is desirable to purify the components or variants thereof. Purification of the mini-vRNAP (SEQ ID NO:4) can be done in two step using affinity columns. The mini-vRNAP of SEQ ID NO:6 has been modified to comprise a His tag such that purification can be done in a single step when using metal affinity columns such as those which employ nickel, cobalt or zinc. The full length vRNAP of SEQ ID NO:15 is also His tagged for purification.

[00219] According to one embodiment of the present invention, purification of a peptide comprising vRNAP can be utilized ultimately to operatively link this domain with a selective agent. Protein purification techniques are well known to those of skill in the art. These techniques involve, at one level, the crude fractionation of the cellular milieu to polypeptide and non-polypeptide fractions. Having separated the polypeptide from other proteins, the polypeptide of interest may be further purified using chromatographic and electrophoretic techniques to achieve partial or complete purification (or purification to homogeneity). Analytical methods particularly suited to the preparation of a pure peptide are ion-exchange chromatography, exclusion chromatography; polyacrylamide gel electrophoresis; isoelectric focusing. A particularly efficient method of purifying peptides is affinity chromatography.

[00220] A tag may be used for protein or peptide purification and detection such as hexahistidine (6-His, HHHHHH), FLAG (DYKDDDDK), hemagglutinin (HA, YPYDVPDYA) and c-myc (EQKLISEEDL). Other tags also have been generated, most of which are very small, comprising only a few amino acids, and are therefore likely to have little to no effect on the conformation of the mature protein or peptide. These small tags do not require any special conformation to be recognized by antibodies. Systems for protein purification using these tags include NTA resin (6-His) or the FLAG fusion system marketed by IBI (FLAG) where the fusion protein is affinity-purified on an antibody column.

[00221] Certain aspects of the present invention concern the purification, and in particular embodiments, the substantial purification, of an encoded protein or peptide, such as a vRNAP. The term "purified protein or peptide" as used herein, is intended to refer to a composition, isolatable from other components, wherein the protein or peptide is purified to any degree

relative to its naturally-obtainable state. A purified protein or peptide therefore also refers to a protein or peptide, free from the environment in which it may naturally occur.

[00222] Generally, "purified" will refer to a protein or peptide composition, such as the vRNAP, that has been subjected to fractionation to remove various other components, and which composition substantially retains its expressed biological activity. Where the term "substantially purified" is used, this designation will refer to a composition in which the protein or peptide forms the major component of the composition, such as constituting about 50%, about 60%, about 70%, about 80%, about 90%, about 95% or more of the proteins in the composition.

[00223] Various methods for quantifying the degree of purification of the protein or peptide will be known to those of skill in the art in light of the present disclosure. These include, for example, determining the specific activity of an active fraction, or assessing the amount of polypeptides within a fraction by SDS/PAGE analysis. A preferred method for assessing the purity of a fraction is to calculate the specific activity of the fraction, to compare it to the specific activity of the initial extract, and to thus calculate the degree of purity, herein assessed by a "-fold purification" number. The actual units used to represent the amount of activity will, of course, be dependent upon the particular assay technique chosen to follow the purification and whether or not the expressed protein or peptide exhibits a detectable activity.

[00224] Various techniques suitable for use in protein purification will be well known to those of skill in the art. These include, for example, precipitation with ammonium sulphate, PEG, antibodies and the like or by heat denaturation, followed by centrifugation; chromatography steps such as ion exchange, gel filtration, reverse phase, hydroxylapatite and affinity chromatography; isoelectric focusing; gel electrophoresis; and combinations of such and other techniques. As is generally known in the art, it is believed that the order of conducting the various purification steps may be changed, or that certain steps may be omitted, and still result in a suitable method for the preparation of a substantially purified protein or peptide.

[00225] There is no general requirement that the protein or peptide always be provided in their most purified state. Indeed, it is contemplated that less substantially purified products will have utility in certain embodiments. Partial purification may be accomplished by using fewer purification steps in combination, or by utilizing different forms of the same general purification scheme. For example, it is appreciated that a cation-exchange column chromatography performed utilizing an HPLC apparatus will generally result in a greater "-fold" purification than the same technique utilizing a low pressure chromatography system. Methods exhibiting a

lower degree of relative purification may have advantages in total recovery of protein product, or in maintaining the activity of an expressed protein.

[00226] It is known that the migration of a polypeptide can vary, sometimes significantly, with different conditions of SDS/PAGE. It will therefore be appreciated that under differing electrophoresis conditions, the apparent molecular weights of purified or partially purified expression products may vary.

[00227] Ion exchange chromatography is a preferred method of separation. Using columns resins such as the metal affinity chromatography resin TALON are also preferred. TALON resin has an enhanced resolving power for polyhistidine-tagged proteins. This results in greater purity with less effort. TALON employs cobalt, an electropositive metal with a remarkably high affinity for polyhistidine-tagged proteins and a low affinity for other proteins. Often, no discernible binding of host proteins occurs and a separate wash step is not required. The binding properties of cobalt allow protein elution under mild pH conditions that protect protein integrity.

[00228] Further concentration of the proteins can be done on an anion exchange column, such as the MonoQ column, a high resolution, anion exchange column. This column works at pressures less than 5 MPa, has a high capacity and gives very high chromatographic resolution.

[00229] High Performance Liquid Chromatography (HPLC) is characterized by a very rapid separation with extraordinary resolution of peaks. This is achieved by the use of very fine particles and high pressure to maintain an adequate flow rate. Separation can be accomplished in a matter of minutes, or at most an hour. Moreover, only a very small volume of the sample is needed because the particles are so small and close-packed that the void volume is a very small fraction of the bed volume. Also, the concentration of the sample need not be very great because the bands are so narrow that there is very little dilution of the sample.

[00230] Gel chromatography, or molecular sieve chromatography, is a special type of partition chromatography that is based on molecular size. The theory behind gel chromatography is that the column, which is prepared with tiny particles of an inert substance that contain small pores, separates larger molecules from smaller molecules as they pass through or around the pores, depending on their size. As long as the material of which the particles are made does not adsorb the molecules, the sole factor determining rate of flow is the size. Hence, molecules are eluted from the column in decreasing size, so long as the shape is relatively constant. Gel chromatography is unsurpassed for separating molecules of different size because separation is independent of all other factors such as pH, ionic strength, temperature, etc. There

also is virtually no adsorption, less zone spreading and the elution volume is related in a simple matter to molecular weight.

[00231] Affinity chromatography, a particularly efficient method of purifying peptides, is a chromatographic procedure that relies on the specific affinity between a substance to be isolated and a molecule that it can specifically bind to. This is a receptor-ligand type interaction. The column material is synthesized by covalently coupling one of the binding partners to an insoluble matrix. The column material is then able to specifically adsorb the substance from the solution. Elution occurs by changing the conditions to those in which binding will not occur (e.g., alter pH, ionic strength, and temperature). Tags, as described herein above, can be used in affinity chromatography.

[00232] The matrix should be a substance that itself does not adsorb molecules to any significant extent and that has a broad range of chemical, physical and thermal stability. The ligand should be coupled in such a way as to not affect its binding properties. The ligand also should provide relatively tight binding, and it should be possible to elute the substance without destroying the sample or the ligand. One of the most common forms of affinity chromatography is immunoaffinity chromatography. The generation of antibodies that would be suitable for use in accordance with the present invention is discussed below.

[00233] An affinity column may have an N4 promoter which the vRNAP or mini-vRNAP proteins recognize attached to a matrix. This column would be suitable for use for the purification of polymerases with no additional tags such as histidine tags.

VII. Separation, Quantitation, and Identification Methods

[00234] Following synthesis of the RNA, it may be desirable to separate the amplification products of several different lengths from each other and from the template and the excess primer.

a. Gel Electrophoresis

[00235] In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods.

b. Chromatographic Techniques

[00236] Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography which may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for

using them including column, paper, thin-layer and gas chromatography. In yet another alternative, labeled cDNA products, such as biotin-labeled or antigen-labeled, can be captured with beads bearing avidin or antibody, respectively.

c. Microfluidic Techniques

[00237] Microfluidic techniques include separation on a platform such as microcapillaries, designed by ACLARA BioSciences Inc., or the LabChip™ "liquid integrated circuits" made by Caliper Technologies Inc. These microfluidic platforms require only nanoliter volumes of sample, in contrast to the microliter volumes required by other separation technologies. Miniaturizing some of the processes involved in genetic analysis has been achieved using microfluidic devices. For example, published PCT Application No. WO 94/05414, to Northrup and White, incorporated herein by reference, reports an integrated micro-PCR apparatus for collection and amplification of nucleic acids from a specimen. U.S. Pat. Nos. 5,304,487 to Wilding *et al.*, and 5,296,375 to Kricka *et al.*, discuss devices for collection and analysis of cell containing samples and are incorporated herein by reference. U.S. Pat. No. 5,856,174 describes an apparatus which combines the various processing and analytical operations involved in nucleic acid analysis and is incorporated herein by reference.

d. Capillary Electrophoresis

[00238] In some embodiments, it may be desirable to provide an additional, or alternative means for analyzing the amplified genes. In these embodiments, micro capillary arrays are contemplated to be used for the analysis.

[00239] Microcapillary array electrophoresis generally involves the use of a thin capillary or channel which may or may not be filled with a particular separation medium. Electrophoresis of a sample through the capillary provides a size based separation profile for the sample. Microcapillary array electrophoresis generally provides a rapid method for size-based sequencing, PCR product analysis and restriction fragment sizing. The high surface to volume ratio of these capillaries allows for the application of higher electric fields across the capillary without substantial thermal variation across the capillary, consequently allowing for more rapid separations. Furthermore, when combined with confocal imaging methods, these methods provide sensitivity in the range of attomoles, which is comparable to the sensitivity of radioactive sequencing methods. Typically, these methods comprise photolithographic etching of micron scale channels on a silica, silicon or other crystalline substrate or chip, and can be

readily adapted for use in the present invention. In some embodiments, the capillary arrays may be fabricated from the same polymeric materials described for the fabrication of the body of the device, using the injection molding techniques described herein.

[00240] Rectangular capillaries are known as an alternative to the cylindrical capillary glass tubes. Some advantages of these systems are their efficient heat dissipation due to the large height-to-width ratio and, hence, their high surface-to-volume ratio and their high detection sensitivity for optical on-column detection modes. These flat separation channels have the ability to perform two-dimensional separations, with one force being applied across the separation channel, and with the sample zones detected by the use of a multi-channel array detector.

[00241] In many capillary electrophoresis methods, the capillaries, e.g., fused silica capillaries or channels etched, machined or molded into planar substrates, are filled with an appropriate separation/sieving matrix. Typically, a variety of sieving matrices are known in the art may be used in the microcapillary arrays. Examples of such matrices include, e.g., hydroxyethyl cellulose, polyacrylamide, agarose and the like. Generally, the specific gel matrix, running buffers and running conditions are selected to maximize the separation characteristics of the particular application, e.g., the size of the nucleic acid fragments, the required resolution, and the presence of native or undenatured nucleic acid molecules. For example, running buffers may include denaturants, chaotropic agents such as urea or the like, to denature nucleic acids in the sample.

e. Mass Spectroscopy

[00242] Mass spectrometry provides a means of "weighing" individual molecules by ionizing the molecules in vacuo and making them "fly" by volatilization. Under the influence of combinations of electric and magnetic fields, the ions follow trajectories depending on their individual mass (m) and charge (z). For low molecular weight molecules, mass spectrometry has been part of the routine physical-organic repertoire for analysis and characterization of organic molecules by the determination of the mass of the parent molecular ion. In addition, by arranging collisions of this parent molecular ion with other particles (e.g., argon atoms), the molecular ion is fragmented forming secondary ions by the so-called collision induced dissociation (CID). The fragmentation pattern/pathway very often allows the derivation of detailed structural information. Other applications of mass spectrometric methods known in the

art can be found summarized in Methods in Enzymology, Vol. 193: "Mass Spectrometry" (J. A. McCloskey, editor), 1990, Academic Press, New York.

[00243] Due to the apparent analytical advantages of mass spectrometry in providing high detection sensitivity, accuracy of mass measurements, detailed structural information by CID in conjunction with an MS/MS configuration and speed, as well as on-line data transfer to a computer, there has been considerable interest in the use of mass spectrometry for the structural analysis of nucleic acids. The biggest hurdle to applying mass spectrometry to nucleic acids is the difficulty of volatilizing these very polar biopolymers. Therefore, "sequencing" had been limited to low molecular weight synthetic oligonucleotides by determining the mass of the parent molecular ion and through this, confirming the already known sequence, or alternatively, confirming the known sequence through the generation of secondary ions (fragment ions) via CID in an MS/MS configuration utilizing, in particular, for the ionization and volatilization, the method of fast atomic bombardment (FAB mass spectrometry) or plasma desorption (PD mass spectrometry).

[00244] Two ionization/desorption techniques are electrospray/ion spray (ES) and matrix-assisted laser desorption/ionization (MALDI). As a mass analyzer, a quadrupole is most frequently used. The determination of molecular weights in femtomole amounts of sample is very accurate due to the presence of multiple ion peaks, which all could be used for the mass calculation.

[00245] MALDI mass spectrometry, in contrast, can be particularly attractive when a time-of-flight (TOF) configuration is used as a mass analyzer. Since, in most cases, no multiple molecular ion peaks are produced with this technique, the mass spectra, in principle, look simpler compared to ES mass spectrometry. DNA molecules up to a molecular weight of 410,000 Daltons could be desorbed and volatilized. More recently, the use of infra red lasers (IR) in this technique (as opposed to UV-lasers) has been shown to provide mass spectra of larger nucleic acids such as synthetic DNA, restriction enzyme fragments of plasmid DNA, and RNA transcripts up to a size of 2180 nucleotides.

[00246] In Japanese Patent No. 59-131909, an instrument is described which detects nucleic acid fragments separated either by electrophoresis, liquid chromatography or high speed gel filtration. Mass spectrometric detection is achieved by incorporating into the nucleic acids atoms which normally do not occur in DNA such as S, Br, I or Ag, Au, Pt, Os, Hg.

f. Energy Transfer

[00247] Labeling hybridization oligonucleotide probes with fluorescent labels is a well known technique in the art and is a sensitive, nonradioactive method for facilitating detection of probe hybridization. More recently developed detection methods employ the process of fluorescence energy transfer (FET) rather than direct detection of fluorescence intensity for detection of probe hybridization. FET occurs between a donor fluorophore and an acceptor dye (which may or may not be a fluorophore) when the absorption spectrum of one (the acceptor) overlaps the emission spectrum of the other (the donor) and the two dyes are in close proximity. Dyes with these properties are referred to as donor/acceptor dye pairs or energy transfer dye pairs. The excited-state energy of the donor fluorophore is transferred by a resonance dipole-induced dipole interaction to the neighboring acceptor. This results in quenching of donor fluorescence. In some cases, if the acceptor is also a fluorophore, the intensity of its fluorescence may be enhanced. The efficiency of energy transfer is highly dependent on the distance between the donor and acceptor, and equations predicting these relationships have been developed. The distance between donor and acceptor dyes at which energy transfer efficiency is 50% is referred to as the Forster distance (R_0). Other mechanisms of fluorescence quenching are also known including, for example, charge transfer and collisional quenching.

[00248] Energy transfer and other mechanisms which rely on the interaction of two dyes in close proximity to produce quenching are an attractive means for detecting or identifying nucleotide sequences, as such assays may be conducted in homogeneous formats. Homogeneous assay formats are simpler than conventional probe hybridization assays which rely on detection of the fluorescence of a single fluorophore label, as heterogeneous assays generally require additional steps to separate hybridized label from free label.

[00249] Homogeneous methods employing energy transfer or other mechanisms of fluorescence quenching for detection of nucleic acid amplification have also been described. Higuchi *et al.*, *Biotechnology* 10:413-417 (1992) disclose methods for detecting DNA amplification in real-time by monitoring increased fluorescence of ethidium bromide as it binds to double-stranded DNA. The sensitivity of this method is limited because binding of the ethidium bromide is not target specific and background amplification products are also detected. WO 96/21144 discloses continuous fluorometric assays in which enzyme-mediated cleavage of nucleic acids results in increased fluorescence. Fluorescence energy transfer is suggested for use in the methods, but only in the context of a method employing a single fluorescent label which is quenched by hybridization to the target.

[00250] Signal primers or detector probes which hybridize to the target sequence downstream of the hybridization site of the amplification primers have been described for use in detection of nucleic acid amplification (U.S. Pat. No. 5,547,861). The signal primer is extended by the polymerase in a manner similar to extension of the amplification primers. Extension of the amplification primer displaces the extension product of the signal primer in a target amplification-dependent manner, producing a double-stranded secondary amplification product which may be detected as an indication of target amplification. The secondary amplification products generated from signal primers may be detected by means of a variety of labels and reporter groups, restriction sites in the signal primer which are cleaved to produce fragments of a characteristic size, capture groups, and structural features such as triple helices and recognition sites for double-stranded DNA binding proteins.

[00251] Many donor/acceptor dye pairs known in the art and may be used in the present invention. These include, for example, fluorescein isothiocyanate (FITC)/tetramethylrhodamine isothiocyanate (TRITC), FITC/Texas Red (Molecular Probes), FITC/N-hydroxysuccinimidyl 1-pyrenebutyrate (PYB), FITC/eosin isothiocyanate (EITC), N-hydroxysuccinimidyl 1-pyrenesulfonate (PYS)/FITC, FITC/Rhodamine X, FITC/tetramethylrhodamine (TAMRA), and others. The selection of a particular donor/acceptor fluorophore pair is not critical. For energy transfer quenching mechanisms, it is only necessary that the emission wavelengths of the donor fluorophore overlap the excitation wavelengths of the acceptor, i.e., there must be sufficient spectral overlap between the two dyes to allow efficient energy transfer, charge transfer or fluorescence quenching. P-(dimethyl aminophenylazo) benzoic acid (DABCYL) is a non-fluorescent acceptor dye which effectively quenches fluorescence from an adjacent fluorophore, e.g., fluorescein or 5-(2'-aminoethyl) aminonaphthalene (EDANS). Any dye pair which produces fluorescence quenching in the detector nucleic acids of the invention are suitable for use in the methods of the invention, regardless of the mechanism by which quenching occurs. Terminal and internal labeling methods are both known in the art and may be routinely used to link the donor and acceptor dyes at their respective sites in the detector nucleic acid.

g. In Vitro Studies

[00252] The synthesized RNA of the current invention may be used for *in vitro* studies of spliceosome assembly, splicing reactions, or antisense experiments.

[00253] The spliceosome is a large, multisubunit complex consisting of small, nuclear ribonucleoprotein particles (snRNPs). There are a total of 5 snRNAs: U1, U2, U4, U5, and U6

which are small and uridine rich. Each snRNP has 1 or 2 of these RNAs. In addition to catalyzing the splicing reaction, the spliceosome retains intermediate products, positions splice sites for precise joining of the exons, and prevents exons from diffusing away after cleavage and before ligation. Spliceosome catalysis involves concerted cleavage/ligation reactions in which the 2'-OH of branch site A attacks the 5' splice site to form a 2'-5' phosphodiester bond with the first nucleotide of the intron. The resulting 3'-OH at the end of the 5' exon attacks the 3' splice site to release the lariat form of the intron and join the two exons together with a normal 3'-5' phosphodiester bond. At least 50 different proteins are involved in spliceosome assembly and function. In the group I and group II introns, splicing is improved (in velocity and accuracy) by protein factors.

VIII. Methods for Making Transcription Products Corresponding to a Target Nucleic Acid Sequence

[00254] In one aspect, the invention comprises a method for making a transcription product corresponding to a target nucleic acid sequence, the method comprising: (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter; (b) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic sequence that is present in or complementary to at least a portion of a target nucleic acid in a sample; (c) obtaining a ssDNA transcription substrate by operably joining to the single-stranded DNA a single-stranded polynucleotide comprising a promoter sequence that binds the RNA polymerase; (d) obtaining nucleoside triphosphates (NTPs) that are substrates for the RNA polymerase and that are complementary to the canonical nucleic acid bases; (e) admixing the RNA polymerase, the ssDNA transcription substrate and the NTPs; and (f) incubating the RNA polymerase and the ssDNA transcription substrate under conditions effective to allow synthesis of transcription product.

[00255] In yet another embodiment, the invention comprises a method for obtaining additional rounds of synthesis of a transcription product corresponding to a target nucleic acid sequence, the method comprising: (a) obtaining a first transcription product by transcription of a first ssDNA transcription substrate corresponding to a target nucleic acid sequence; (b) obtaining a reverse transcriptase; (c) reverse transcribing the first transcription product; (d) obtaining first-strand cDNA complementary to the first transcription product; (e) obtaining a second ssDNA transcription substrate by operably joining to the first-strand cDNA a single-stranded polynucleotide comprising a promoter sequence that binds an RNA polymerase that can transcribe RNA using a single-stranded promoter; (f) obtaining an RNA polymerase that can

transcribe RNA using a single-stranded promoter; (g) obtaining nucleoside triphosphates (NTPs) that are substrates for the RNA polymerase; (h) admixing the RNA polymerase, the second ssDNA transcription substrate and the NTPs; and (i) incubating the RNA polymerase and the second ssDNA transcription substrate under conditions effective to allow synthesis of a second transcription product. This method for obtaining additional rounds of synthesis of transcription product can be repeated in a stepwise manner to obtain synthesis of still more transcription products. Alternatively, in other embodiments, the invention comprises continuous transcription, wherein additional ssDNA transcription substrates are obtained continuously by reverse transcription of transcription products to obtain first-strand cDNA and operable joining of a promoter to the first-strand cDNA, and wherein, the additional ssDNA transcriptional substrates are used to make more transcription products.

[00256] The invention comprises use of any polymerase that synthesizes a transcription product using a single-stranded transcription promoter and a single-stranded DNA template to which the promoter is operably or functionally joined or linked. An RNA polymerase that synthesizes a transcription product using a ssDNA transcription substrate comprising a ssDNA transcription promoter that is operably joined to a target sequence has not previously been known in the art for making a transcription product corresponding to a target sequence. A preferred RNA polymerase of the invention is an N4 mini-vRNAP enzyme or a mutant form of an N4 mini-vRNAP enzyme. The RNA polymerase preferentially has the amino sequence set forth in SEQ ID NO:4 or SEQ ID NO:6 or a mutant of the polymerase of SEQ ID NO:4 or SEQ ID NO:6, such as a mutant with a mutation at position number Y678 or the polymerase of SEQ ID NO:8. The vRNAP and mini-vRNA polymerase transcribe nucleic acid operatively linked to an N4 promoter such as a P2 promoter of SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29. The promoter of SEQ ID NO:16 or SEQ ID NO:28 is preferred.

[00257] Other preferred promoters of the invention comprise single-stranded pseudopromoters or synthetic promoters that are recognized by an RNA polymerase (RNAP) so as to function in a method of the invention. A "pseudopromoter" or "synthetic promoter" of the present invention can be any single-stranded sequence that is identified and/or selected to be functional as a promoter for *in vitro* transcription by an RNA polymerase that binds the promoter with specificity and functions as a promoter for the RNA polymerase in a transcription reaction. A promoter comprising a pseudopromoter or synthetic promoter of the invention can be made as described by Ohmichi *et al.* (*Proc. Natl. Acad. Sci. USA* 99:54-59, 2002), which is incorporated herein by reference. If a pseudopromoter or synthetic promoter is used as a

promoter in a method or assay of the invention, then the corresponding RNA polymerase for which the pseudopromoter or synthetic promoter was identified and/or selected is used in the method. By way of example, but not of limitation, a target probe with a promoter comprising a ssDNA pseudopromoter can be obtained and used in a method or assay of the invention that uses *E. coli* RNAP or a T7-type phage RNAP, such as, but not limited to, T7 RNAP, T3 RNAP, or SP6 RNAP, as described by Ohmichi *et al.* (*Proc. Natl. Acad. Sci. USA* 99:54-59, 2002) and incorporated herein by reference. By "T7-type RNAPs" we mean T7, T3, ϕ , ϕ III, W31, gh1, Y, A1122, SP6 and mitochondrial RNAPs. Mutant RNAPs, such as, but not limited to, T7 RNAP Y639F mutant enzyme, T3 RNAP Y573F mutant enzyme, SP6 RNAP Y631F mutant enzyme, T7 RNAP having altered amino acids at both positions 639 and 784, T3 RNAP having altered amino acids at both positions 573 and 785, or SP6 RNAP having altered amino acids at both positions 631 and 779, or other mutant forms of an RNAP that functions in a method of the invention can also be used in embodiments of methods or assays of the invention with single-stranded pseudopromoters obtained as described by Ohmichi *et al.* (*Proc. Natl. Acad. Sci. USA* 99:54-59, 2002).

A. Supplemental Definitions and Methods

1. Transcription Product

[00258] The term "transcription product" as used herein can mean RNA or, in view of the ability of certain polymerases of the invention, such as the Y678F mutant polymerase having SEQ ID NO:8, to use deoxynucleotide substrates, a transcription product can also comprise DNA or a mixture of both RNA and DNA. A transcription product does not necessarily have perfect sequence complementarity or identity to the target sequence. For example, a transcription product can include nucleotide analogs such as deoxyinosine or deoxyuridine, intentional sequence alterations, such as sequence alterations introduced through a primer comprising a sequence that is hybridizable, but not complementary, to the target sequence, and/or sequence errors that occur during transcription.

[00259] Also, for a variety of reasons, a nucleic acid or polynucleotide of the invention may comprise one or more modified nucleic acid bases, sugar moieties, or internucleoside linkages. By way of example, some reasons for using nucleic acids or polynucleotides that contain modified bases, sugar moieties, or internucleoside linkages include, but are not limited to: (1) modification of the T_m; (2) changing the susceptibility of the polynucleotide to one or more nucleases; (3) providing a moiety for attachment of a label; (4) providing a label or a

quencher for a label; or (5) providing a moiety, such as biotin, for attaching to another molecule which is in solution or bound to a surface.

2. Sample/Target/Target Nucleic Acid/Target Sequence

[00260] A “sample” or a “biological sample” according to the present invention is used in its broadest sense. A sample is any specimen that is collected from or is associated with a biological or environmental source, or which comprises or contains biological material, whether in whole or in part, and whether living or dead.

[00261] Biological samples may be plant or animal, including human, fluid (e.g., blood or blood fractions, urine, saliva, sputum, cerebral spinal fluid, pleural fluid, milk, lymph, or semen), swabs (e.g., buccal or cervical swabs), solid (e.g., stool), microbial cultures (e.g., plate or liquid cultures of bacteria, fungi, parasites, protozoans, or viruses), or cells or tissue (e.g., fresh or paraffin-embedded tissue sections, hair follicles, mouse tail snips, leaves, or parts of human, animal, plant, microbial, viral, or other cells, tissues, organs or whole organisms, including subcellular fractions or cell extracts), as well as liquid and solid food and feed products and ingredients such as dairy items, vegetables, meat and meat by-products, and waste. Biological samples may be obtained from all of the various families of domestic plants or animals, as well as wild animals or plants.

[00262] Environmental samples include environmental material such as surface matter, soil, water, air, or industrial samples, as well as samples obtained from food and dairy processing instruments, apparatus, equipment, utensils, disposable and non-disposable items. These examples are not to be construed as limiting the sample types applicable to the present invention.

[00263] In short, a sample comprises a specimen from any source that contains or may contain a naturally occurring target nucleic acid.

[00264] A sample on which the assay method of the invention is carried out can be a raw specimen of biological material, such as serum or other body fluid, tissue culture medium or food material. More typically, the method is carried out on a sample that is a processed specimen, derived from a raw specimen by various treatments to remove materials that would interfere with detection of a target nucleic acid or an amplification product thereof. Methods for processing raw samples to obtain a sample more suitable for the assay methods of the invention are well known in the art.

[00265] An “analyte” means a substance whose presence, concentration or amount in a sample is being determined in an assay. An analyte is sometimes referred to as a “target substance” or a “target molecule” or a “target analyte” of an assay. An analyte may also be referred to more specifically. Embodiments of the present invention pertain to analytes that are naturally occurring nucleic acids, and the analyte may be referred to as a “target nucleic acid” or a “target polynucleotide” or a “target oligonucleotide” or a “target sequence,” depending on the particular case. A composition, kit, or method of the invention can be used for an “analyte-specific reagent” to detect a target nucleic acid analyte or another analyte or analyte-binding substance in a sample.

[00266] With respect to the present invention, an analyte is often associated with a biological entity that is present in a sample if and only if the analyte is present. Such biological entities include viroids (analyte is, e.g., a nucleic acid or a segment thereof); viruses (analyte is, e.g., a viral genome, or a segment of viral genome); other microorganisms (analyte is, e.g., a segment of the genome or the RNA of the microorganism); abnormal cells, such as cancer cells (analyte is, e.g., an oncogene); or an abnormal gene (analyte is, e.g., a gene segment which includes the altered bases which render the gene abnormal, or a messenger RNA segment which includes altered bases as a result of having been transcribed from the abnormal gene).

[00267] From the description of analyte, it is apparent that the present invention has widespread applicability, including in applications in which nucleic acid probe hybridization assays are often employed. Thus, among other applications, the invention is useful in diagnosing diseases in plants and animals, including humans; and in testing products, such as food, blood, and tissue cultures, for contaminants.

[00268] A “target” of the present invention is a biological organism or material that is the reason or basis for which a biological assay or a diagnostic assay is performed. By way of example, but not of limitation, an assay of the present invention may be performed to detect a target that is a virus which is indicative of a present disease or a risk of future disease (e.g., HIV which is believed to result in AIDS), or a target that is a gene which is indicative of antibiotic resistance (e.g., an antibiotic resistance gene in an infectious pathogenic bacterium), or a target that is a gene which, if absent, may be indicative of disease (e.g., a deletion in an essential gene). In developing assays according to the present invention, it is important to identify target analytes that yield assay results that are sufficiently specific, accurate, and sensitive to be meaningful related to the presence or condition of the target.

[00269] A target analyte that is a “target polynucleotide” or a “target nucleic acid” comprises at least one nucleic acid molecule or portion of at least one nucleic acid molecule, whether the molecule or molecules is or are DNA, RNA, or both DNA and RNA, and wherein each the molecule has, at least in part, a defined nucleotide sequence. The target polynucleotide may also have at least partial complementarity with other molecules used in an assay, such as, but not limited to, primers, splice template oligos, ligation splint oligos, capture probes or detection probes. The target polynucleotide may be single- or double- stranded. A target polynucleotide of the present invention may be of any length. However, it must comprise a polynucleotide sequence of sufficient sequence specificity and length so as to be useful for its intended purpose. By way of example, but not of limitation, a target nucleic acid that is to be detected using a sequence-complementary detection probe must have a sequence of sufficient sequence specificity and length so as remain hybridized by the detection probe under assay hybridization conditions wherein sequences that are not target polynucleotides are not hybridized. A target polynucleotide having sufficient sequence specificity and length for an assay of the present invention may be identified, using methods known to those skilled in the art, by comparison and analysis of nucleic acid sequences known for a target and for other sequences which may be present in the sample. For example, sequences for nucleic acids of many viruses, bacteria, humans (e.g., for genes and messenger RNA), and many other biological organisms can be searched using public or private databases, and sequence comparisons, folded structures, and hybridization melting temperatures (i.e., T_m 's) may be obtained using computer software known to those knowledgeable in the art.

[00270] The terms “source of target nucleic acid” or “source of target polynucleotide” refers to any sample that contains a naturally occurring target nucleic acid, RNA or DNA.

[00271] Thus, a method of the present invention can be carried out on nucleic acid from a variety of sources, including unpurified nucleic acids, or nucleic acids purified using any appropriate method in the art, such as, but not limited to, various “spin” columns, cationic membranes and filters, or salt precipitation techniques, for which a wide variety of products are commercially available (e.g., MasterPure™ DNA & RNA Purification Kits from Epicentre Technologies, Madison, WI, USA). Methods of the present invention can also be carried out on nucleic acids isolated from viroids, viruses or cells of a specimen and deposited onto solid supports as described by Gillespie and Spiegelman (*J. Mol. Biol.* 12:829-842, 1965), including solid supports on dipsticks and the inside walls of microtiter plate wells. The method can also be carried out with nucleic acid isolated from specimens and deposited on solid support by “dot”

blotting (Kafatos, *et al.*, Nucl. Acids Res. 7:1541-1552, 1979); White, and Bancroft, *J. Biol. Chem.* 257:8569-8572, 1982); Southern blotting (Southern, E., *J. Mol. Biol.* 98:503-517, 1975); “northern” blotting (Thomas, *Proc. Natl. Acad. Sci. USA* 77:5201-5205, 1980); and electroblotting (Stellwag, and Dahlberg, Nucl. Acids Res. 8:299-317, 1980). The method can also be carried out for nucleic acids spotted on membranes, on slides, or on chips as arrays or microarrays, or the method can be carried out to prepare probes for detecting or quantifying nucleic acids present in a sample based on hybridization to nucleic acids spotted or synthesized on one of these surfaces. Nucleic acid of specimens can also be assayed by the method of the present invention applied to water phase hybridization (Britten, and Kohne, *Science* 161:527-540, 1968) and water/organic interphase hybridizations (Kohne, *et al.*, *Biochemistry* 16:5329-5341, 1977). Water/organic interphase hybridizations have the advantage of proceeding with very rapid kinetics but are not suitable when an organic phase-soluble linking moiety, such as biotin, is joined to the nucleic acid affinity molecule.

[00272] The methods of the present invention can also be carried out on amplification products obtained by amplification of a naturally occurring target nucleic acid, provided that the target sequence in the target nucleic acid is amplified by the method used only if the target nucleic acid is present in the sample. Suitable amplification methods include, but are not limited to, PCR, RT-PCR, NASBA, TMA, 3SR, LCR, LLA, SDA (e.g., Walker *et al.*, Nucleic Acids Res. 20:1691-1696, 1992), RCA, Multiple Displacement Amplification (Molecular Staging), ICAN™ or UCAN™ (TAKARA), Loop-AMP (EIKEN), and SPIA™ or Ribo-SPIA™ (NuGEN Technologies). There are various reasons for using a nucleic acid that is a product of another amplification method as a target nucleic acid for an assay of the present invention, such as, but not limited to, for obtaining more sensitive detection of targets, greater specificity, or to decrease the time required to obtain an assay result.

[00273] The methods of the invention can also be carried out on nucleic acids isolated from specimens and deposited onto solid supports by dot-blotting, or by adsorption onto walls of microtiter plate wells or solid support materials on dipsticks, on membranes, on slides, or on chips as arrays or microarrays.

[00274] Still further, the methods of the invention are applicable to detecting cellular nucleic acids in whole cells from a specimen, such as a fixed or paraffin-embedded section, or from microorganisms immobilized on a solid support, such as replica-plated bacteria or yeast. In some embodiments, the methods of the invention can be used to make a transcription product corresponding to a target sequence to detect target nucleic acids in living cells.

[00275] A target nucleic acid can be a nucleic acid from any source in purified or unpurified form. For example, a target nucleic acid comprising DNA, can be dsDNA or ssDNA such as mitochondrial DNA, chloroplast DNA, chromosomes, plasmids or other episomes, the genomes of bacteria, yeasts, viruses, viroids, mycoplasma, molds, or other microorganisms, or genomes of fungi, plants, animals, or humans. Target nucleic acids comprising RNA can be tRNA, mRNA, rRNA, mitochondrial RNA, chloroplast RNA, micro RNA, or other RNA molecules, without limit. Target nucleic acids can also be mixtures of DNA and RNA, including, but not limited to, mixtures of the above nucleic acids or fragments thereof, or DNA-RNA hybrids. The target nucleic acid can be only a minor fraction of a complex mixture such as a biological sample and can be obtained from various biological materials by procedures known in the art. Numerous methods for purification of a particular target nucleic are known in the art, if further purification is necessary.

[00276] The term "target nucleic acid sequence" or "target sequence" refers to the particular nucleotide sequence of the target nucleic acid(s) that is/are to be transcribed to make a transcription product. A "target sequence" comprises one or more sequences within one or more target nucleic acids. A target sequence can also have "complexing sequences" which are added during processes of methods of the invention for the purpose of facilitating joining of a target sequence to a polynucleotide comprising a single-stranded transcription promoter in order to obtain a ssDNA transcription substrate of the invention. A complexing sequence can provide a complementary sequence to which an oligonucleotide (e.g., a primer and/or splice template) used in a method of the invention can anneal or complex. A complexing sequence usually comprises a "tail" sequence that is added by means such as those discussed herein, including, but not limited to, non-templated addition of dCMP residues to first-strand cDNA by reverse transcriptase pausing at cap structures of mRNA (in the presence or absence of manganese cations) and/or controlled ribonucleotide tailing using TdT. If a complexing sequence is added to a target sequence during a process of a method of the invention, it is desirable that a complexing sequence is chosen that does not to affect the specificity of the transcription products made using the resulting transcription substrate comprising the target sequence.

[00277] A target nucleic acid can be either single-stranded or double-stranded RNA, DNA, or mixed RNA and DNA. A target nucleic acid is sometimes referred to more specifically by the type of nucleic acid. By way of example, but not of limitation, a target nucleic acid can be a "target RNA" or an "RNA target," or a "target mRNA," or a "target DNA" or a "DNA target." Similarly, the target sequence can be referred to as "a target RNA sequence"

or a "RNA target sequence", or as a "target mRNA sequence" or a "target DNA sequence," or the like. In some embodiments, the target sequence comprises one or more entire target nucleic acids, such as, one or all full-length mRNA molecules in a particular sample. In other embodiments, the target sequence comprises only a portion of one or more target nucleic acid molecules. When the target nucleic acid is originally single-stranded, the term "target sequence" is also meant to refer to the sequence complementary to the "target sequence." When the "target nucleic acid" is originally double-stranded, the term "target sequence" may refer to both the sense strand of the sequence or its complement, or both, depending on the intended purpose of the method. The target sequence may be known or not known, in terms of its actual sequence. In some instances, the terms "target sequence," "target nucleic acid," "target polynucleotide," and variations thereof are used interchangeably.

3. cDNA/First-Strand cDNA/Second-Strand cDNA/Reverse Transcriptase/RnaseH

[00278] In some important embodiments of the invention, the target sequence of a transcription substrate comprises cDNA. In general, "cDNA" refers to "complementary DNA" that is synthesized by primer extension using a DNA polymerase, including, but not limited to, an RNA-dependent DNA polymerase or reverse transcriptase, using at least a portion of a target nucleic acid as a template, and which cDNA is "homologous to" or "base pairs with" at least a portion of the target nucleic acid template. In some embodiments of the invention, which are preferred embodiments, cDNA is obtained by reverse transcription primer extension using a reverse transcriptase and a target nucleic acid comprising messenger RNA (mRNA) obtained from a biological sample as a template, and which cDNA is homologous to the mRNA. Methods in the art related to making cDNA from mRNA involve synthesis of double-stranded cDNA comprising first-strand cDNA and second-strand cDNA, which usually are synthesized sequentially using different methods. However herein, we often refer to "first-strand cDNA" even when a method of the invention results in synthesis of only one strand of DNA that is complementary to the mRNA (i.e., the term "first-strand cDNA" is not intended to imply that there is also a second-strand cDNA). In some embodiments, the terms "first-strand cDNA" or "cDNA" refer to a single-stranded DNA molecule obtained by reverse transcription of any RNA molecule, even if it is not mRNA. In other embodiments, the terms "first-strand cDNA" or "cDNA" refer to a single-stranded DNA molecule obtained by primer extension using a target nucleic acid comprising either a single-stranded DNA or one strand of a double-stranded DNA as a template for a DNA polymerization reaction.

[00279] An “RNA-dependent DNA polymerase” or “reverse transcriptase” is an enzyme that can synthesize a complementary DNA copy (“cDNA”) from an RNA template. All known reverse transcriptases also have the ability to make a complementary DNA copy from a DNA template; thus, they are both RNA- and DNA-dependent DNA polymerases. A “template” is the nucleic acid molecule that is copied by a nucleic acid polymerase. The synthesized copy is complementary to the template. Both RNA and DNA are always synthesized in the 5’-to-3’ direction and the two strands of a nucleic acid duplex always are aligned so that the 5’ ends of the two strands are at opposite ends of the duplex (and, by necessity, so then are the 3’ ends). A primer is required for both RNA and DNA templates to initiate synthesis by a DNA polymerase. Examples of reverse transcriptases that can be used in methods of the present invention include, but are not limited to, AMV reverse transcriptase, MMLV reverse transcriptase, Tth DNA polymerase, rBst DNA polymerase large fragment, also called IsoTherm™ DNA Polymerase (Epicentre Technologies, Madison, WI, USA), and BcaBEST™ DNA polymerase (Takara Shuzo Co, Kyoto, Japan). In some cases, a mutant form of a reverse transcriptase, such as, an MMLV reverse transcriptase that lacks RNase H activity is used. In still other embodiments, IsoTherm™ DNA polymerase is most suitable. In other embodiments, a wild-type enzyme is preferred.

[00280] In embodiments of the invention that obtain a transcription substrate in which a ssDNA target sequence comprising first-strand cDNA is synthesized using a reverse transcriptase and a primer that is complementary to and anneals to the 3’-end of a target nucleic acid comprising mRNA, the primer can comprise oligo(dT) or modified oligo(dT), or it can comprise an oligo d(T) anchor primer, wherein one (or a small number) of the 3’-nucleotides of the anchor primer comprise either a specific base for a specific mRNA or a randomized nucleotide (i.e., synthesized with a mixture of all four nucleotides) for priming all mRNA molecules in a sample, or the primer can comprise an oligonucleotide having a specific sequence that is complementary to the sequence of a specific mRNA molecule, or in some cases, it can comprise an oligonucleotide having a random sequence (i.e., synthesized using a mixture of all four nucleotides for every position of the primer).

[00281] If a target nucleic acid is RNA, such as mRNA, and it is desirable to remove the RNA that is annealed to first-strand cDNA following reverse transcription using a promoter primer, this can be accomplished by one of several means. By way of example, but not of limitation, the RNA can be removed by treatment with RNase H, by treatment with a base, such as, but not limited to sodium or potassium hydroxide, or the RNA can be removed from the

hybrid by heat denaturation. In preferred embodiments for some applications, the RNA is removed by an RNase H activity of a reverse transcriptase that is used for reverse transcription (or primer extension), such as, but not limited to, MMLV reverse transcriptase. Alternatively, in some embodiments, the RNA is dissociated from the first-strand cDNA by incubating the hybrid or performing the reverse transcription in the presence of a single-strand binding (SSB) protein of the invention, such as, but not limited to *E. coli* SSB (*EcoSSB*).

[00282] In some embodiments of the invention, especially in embodiments for obtaining additional rounds of transcription products, a separate RNase H enzyme is also used, whether or not the reverse transcriptase has RNase H activity. If RNase H activity is desirable in an embodiment for obtaining multiple rounds of transcription, but a separate RNase H enzyme is not added, MMLV reverse transcriptase (wild-type RNase H-positive) can be used. AMV reverse transcriptase can be used in some embodiments for obtaining multiple rounds of transcription in which a separate RNase H enzyme is also added. Kacian *et al.* (U.S. Patent No. 5,399,491) disclose information related to the effects of adding different amounts of a separate RNase H enzyme to transcription-mediated amplification assays that use either MMLV or AMV reverse transcriptase, which reference and information is incorporated herein by reference and made a part of the present disclosure.

[00283] “Ribonuclease H” or “RNase H” is an enzyme that degrades the RNA portion of an RNA:DNA duplex. An RNase H can be an endonuclease or an exonuclease. Most wild-type reverse transcriptase enzymes have an RNase H activity in addition to their polymerase activity. However, other sources of the RNase H are available without an associated polymerase activity. The degradation may result in separation of RNA from an RNA:DNA complex. Alternatively, the RNase H may simply cut the RNA at various locations such that portions of the RNA melt off or permit enzymes to unwind portions of the RNA. When used in an embodiment of the invention, RNaseH enzymes that can be used include, but are not limited to, *E. coli* RNase H, *Thermus thermophilus* RNase H, and *Thermus flavus* RNase H (U.S. Patents Nos. 5,268,289; 5,459,055; and 5,500,370, incorporated herein by reference). The latter two enzymes, which are thermostable and, therefore, maintain more consistent activity in reactions and are more easily stored and shipped, are preferred in most embodiments in which a separate RNase H enzyme is used. Other RNase H enzymes that can be used are those that are described by Sagawa *et al.* in PCT Patent Publication No. WO 02/16639; and in PCT Patent Publications Nos. WO 00/56877 and AU 00/29742, all of which are incorporated herein by reference. In other embodiments, it is

desirable to use a less thermally stable enzyme, such as *E. coli* RNase H, because it is easier to inactivate the enzyme in a reaction mixture.

[00284] Kacian *et al.* disclosed in U.S. Patent No. 5,399,491, incorporated herein by reference, that the number, distribution, and position of putative RNase H cut sites determine, in part, the usefulness of a given primer and that amplification can be improved by inclusion of intentional mismatches or insertion of sequences in order to affect the number, distribution, and position of putative RNase H cut sites. Thus, in preferred processes of the invention for removing RNA from RNA:DNA hybrids following reverse transcription to make first-strand cDNA, the RNA target sequence is determined and then analyzed to determine where RNase H degradation will cause cuts or removal of sections of RNA from the duplex upon synthesis of first-strand cDNA. The processes of the invention include conducting experiments to determine the effect on amplification of the target sequence of the degradation of the RNA target sequence by RNase H present in the reverse transcriptase and/or separate RNase H enzyme(s) used, including, but not limited to, AMV reverse transcriptase, and both RNase H-plus and RNase H-minus MMLV reverse transcriptase, and *E. coli* RNase H or thermostable RNase H enzymes that are stable for more than 10 minutes at 70° C (U.S. Patents Nos. 5,268,289; 5,459,055; and 5,500,370, incorporated herein by reference), such as, but not limited to, Hybridase™ thermostable RNase H (Epicentre Technologies, Madison, WI, USA), Tth RNase H, and Tfl RNase H, or by different combinations of a reverse transcriptase and a separate RNase H.

[00285] In selecting a primer, including a promoter primer of the invention, for use in reverse transcription of an RNA target sequence to make first-strand cDNA, it is preferable that the primer be selected so that it will hybridize to a section of RNA which is substantially nondegraded by the RNase H present in the reaction mixture. If there is substantial degradation, the cuts in the RNA strand in the region of the primer may stop or inhibit DNA synthesis and prevent extension of the primer. Thus, it is desirable to select a primer that will hybridize with a sequence of the RNA target, located so that when the RNA is subjected to RNase H, there is no substantial degradation that would prevent formation of the primer extension product.

4. Transcription Substrate

[00286] As used herein, a “transcription substrate” or a “ssDNA transcription substrate” according to the present invention comprises a ssDNA comprising a target sequence that is operably joined to a transcription promoter, wherein an RNA polymerase can bind to the transcription promoter with specificity and synthesize a transcription product corresponding to

the target sequence under suitable transcription conditions. In order to be operably joined to the target sequence, the transcription promoter is 3'-of the target sequence. A transcription substrate of the invention can also have additional nucleic acid sequences that are 5'-of and/or 3'-of the transcription promoter sequence, but a transcription substrate is not required to have such additional other sequences. By way of example, but not of limitation, a transcription substrate can have a transcription initiation site 5'-of the promoter sequence. Also, in some embodiments of the invention, a transcription substrate can have one or more transcription termination sequences, one or more sites for DNA cleavage to permit controlled linearization of a circular first-strand cDNA that is a transcription substrate, one or more origins ("ori's") of replication (preferably an *ori* for a single-stranded replicon, such as, but not limited to, a phage M13 replicon), a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene or a beta-galactosidase gene, respectively, or one or more transposon recognition sequences, such as outer end ("OE") or mosaic end ("ME") sequences for a Tn5-type transposon,) that can be recognized and used by a transposase for *in vitro* or *in vivo* transposition, or one or more sites that are recognized by a recombinase (such as, but not limited to, the cre-lox system), and/or other sequences and genetic elements for a particular purpose, including, but not limited to, sequences that are transcribed by the RNA polymerase so as to provide additional regions of complementarity in the RNA transcription products for annealing of primers for reverse transcription in order to make cDNA for additional rounds of transcription. As described herein above, a "transcription substrate" of the present invention is single-stranded, whereas double-stranded DNA molecules, having at a minimum double-stranded promoters and usually also double-stranded templates, are used in other methods in the art.

[00287] Since a transcription substrate of the present invention is single-stranded, the terms "3'-of" and "5'-of" are used herein with respect to the present invention to refer to the position or orientation of a particular nucleic acid sequence or genetic element, such as, but not limited to, a transcription promoter, relative to other sequences or genetic elements within the DNA strand comprising the ssDNA transcription substrate. Thus, although the synthesis of RNA in a 5'-to-3' direction during transcription is thought of as proceeding in a "downstream" direction, the transcription promoter on the ssDNA transcription substrate is referred to herein as being 3'-of the target sequence. Those with knowledge in the art will understand these terms in the context of nucleic acid chemistry and structure, particularly related to the 3'- and 5'-positions of sugar moieties of canonical nucleic acid nucleotides. By way of example, a

transcription promoter that is "3'-of the target sequence" on a linear transcription substrate refers a promoter sequence that is at or closer to the 3'-end of the transcription substrate relative to the target sequence on the same strand. If a first nucleic acid sequence is 3'-of a second sequence on one strand, the complement of the first sequence will be 5'-of the complement of the second sequence on the complementary strand. The description of the invention will be understood with respect to the relative 5' or 3' position and orientation of a sequence or genetic element within a particular nucleic acid strand, unless explicitly stated to the contrary.

5. Nucleic Acids, Polynucleotides and Analogs Thereof

[00288] A "nucleic acid" or "polynucleotide" of the invention is a polymer molecule comprising a series of "mononucleosides," also referred to as "nucleosides," in which the 3'-position of the pentose sugar of one nucleoside is linked by an internucleoside linkage, such as, but not limited to, a phosphodiester bond, to the 5'-position of the pentose sugar of the next nucleoside. A nucleoside linked to a phosphate group is referred to as a "nucleotide." The nucleotide that is linked to the 5'-position of the next nucleotide in the series is referred to as "5' of" or the "5' nucleotide" and the nucleotide that is linked to the 3'-position of the 5' nucleotide is referred to as "3' of" or the "3' nucleotide." The pentose sugar of the nucleic acid can be ribose, in which case, the nucleic acid or polynucleotide is referred to as "RNA," or it can be 2'-deoxyribose, in which case, the nucleic acid or polynucleotide is referred to as "DNA." Alternatively, especially if the nucleic acid is synthesized chemically, the nucleic acid can be composed of both DNA and RNA mononucleotides. In both RNA and DNA, each pentose sugar is covalently linked to one of four common "nucleic acid bases" (each also referred to as a "base"). Three of the predominant naturally-occurring bases that are linked to the sugars (adenine, cytosine and guanine) are common for both DNA and RNA, while one base is different; DNA has the additional base thymine, while RNA has the additional base uridine. Those in the art commonly think of a small polynucleotide as an "oligonucleotide." The term "oligonucleotide" as used herein is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, preferably about 10 to 200 nucleotides, but there is no defined limit to the length of an oligonucleotide. The exact size will depend on many factors, which in turn depends on the ultimate function or use of the oligonucleotide.

[00289] In order to accomplish the goals of the invention, there is no limit to the composition of the nucleic acids or polynucleotides of the invention including any splice template oligos, primers, including promoter primers, promoter ligation oligos, ligation splint

oligos, detection probes, such as, but not limited to molecular beacons (U.S. Patents Nos. 5,925,517 and 6,103,476 of Tyagi *et al.* and 6,461,817 of Alland *et al.*, which are incorporated herein by reference), capture probes, oligonucleotides, or other nucleic acids used or detected in the assays or methods, so long as each of the nucleic acid functions for its intended use. By way of example, but not of limitation, the nucleic acid bases in the mononucleotides may comprise guanine, adenine, uracil, thymine, or cytosine, or alternatively, one or more of the nucleic acid bases may comprise xanthine, allyamino-uracil, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl adenines, 2-propyl and other alkyl adenines, 5-halouracil, 5-halo cytosine, 5-propynyl uracil, 5-propynyl cytosine, 7-deazaadenine, 7-deazaguanine, 7-deaza-7-methyl-adenine, 7-deaza-7-methyl-guanine, 7-deaza-7-propynyl-adenine, 7-deaza-7-propynyl-guanine and other 7-deaza-7-alkyl or 7-aryl purines, N2-alkyl-guanine, N2-alkyl-2-amino-adenine, purine 6-aza uracil, 6-aza cytosine and 6-aza thymine, 5-uracil (pseudo uracil), 4-thiouracil, 8-halo adenine, 8-amino-adenine, 8-thiol adenine, 8-thiolalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines and 8-halo guanines, 8-amino-guanine, 8-thiol guanine, 8-thiolalkyl guanines, 8-hydroxyl guanine and other 8-substituted guanines, other aza and deaza uracils, other aza and deaza thymidines, other aza and deaza cytosine, aza and deaza adenines, aza and deaza guanines or 5-trifluoromethyl uracil and 5-trifluorocytosine. Still further, they may comprise a nucleic acid base that is derivatized with a biotin moiety, a digoxigenin moiety, a fluorescent or chemiluminescent moiety, a quenching moiety or some other moiety. The invention is not limited to the nucleic acid bases listed; this list is given to show the broad range of bases which may be used for a particular purpose in a method.

[00290] In some embodiments of the invention, a molecule comprising a "peptide nucleic acid" (PNA) or a molecule comprising both a nucleic acid and a PNA, as described in U.S. Pat. Nos. 5,539,082; 5,641,625; 5,700,922; 5,705,333; 5,714,331; 5,719,262; 5,736,336; 5,773,571; 5,786,461; 5,817,811; 5,977,296; 5,986,053; 6,015,887; and 6,020,126 (and references therein), can also be used. In general, a PNA molecule is a nucleic acid analog consisting of a backbone comprising, for example, N-(2-aminoethyl)glycine units, to each of which a nucleic acid base is linked through a suitable linker, such as, but not limited to an aza, amido, ureido, or methylene carbonyl linker. The nucleic acid bases in PNA molecules bind complementary single-stranded DNA or RNA according to Watson-Crick base-pairing rules. However, the T_m 's for PNA/DNA or PNA/RNA duplexes or hybrids are higher than the T_m 's for DNA/DNA, DNA/RNA, or RNA/RNA duplexes. PNA provides tighter binding and greater binding stability than a nucleic acid of similar base sequence (e.g., see U.S. Pat. No. 5,985,563). Also, since PNA is not

naturally occurring, PNA molecules are highly resistant to protease and nuclease activity. PNA can be prepared according to methods known in the art, such as, but not limited to, methods described in the above-mentioned patents, and references therein.

[00291] When a molecule comprising both a nucleic acid and a peptide nucleic acid (PNA) is used in the invention, modified bases can be used in one or both parts. For example, binding affinity can be increased by the use of certain modified bases in both the nucleotide subunits that make up the 2'-deoxyoligonucleotides of the invention and in the peptide nucleic acid subunits. Such modified bases may include 5-propynylpyrimidines, 6-azapyrimidines, and N-2, N-6 and O-6 substituted purines including 2-aminopropyl-adenine. Other modified pyrimidine and purine base are also expected to increase the binding affinity of macromolecules to a complementary strand of nucleic acid.

[00292] With respect to nucleic acids or polynucleotides of the invention, one or more of the sugar moieties can comprise ribose or 2'-deoxyribose, or alternatively, one or more of the sugar moieties can be some other sugar moiety, such as, but not limited to, 2'-fluoro-2'-deoxyribose or 2'-O-methyl-ribose, which provide resistance to some nucleases, or 2'-amino-2'-deoxyribose or 2'-azido-2'-deoxyribose, which can be used to label transcription products by reacting them with visible, fluorescent, infrared fluorescent or other detectable dyes or chemicals having an electrophilic, photoreactive or other reactive chemical moiety.

[00293] The internucleoside linkages of nucleic acids or polynucleotides of the invention can be phosphodiester linkages, or alternatively, one or more of the internucleoside linkages can comprise modified linkages, such as, but not limited to, phosphorothioate, phosphorodithioate, phosphoroselenate, or phosphorodiselenate linkages, which are resistant to some nucleases.

[00294] A variety of methods are known in the art for making nucleic acids having a particular sequence or that contain particular nucleic acid bases, sugars, internucleoside linkages, chemical moieties, and other compositions and characteristics. Any one or any combination of these methods can be used to make a nucleic acid, polynucleotide, or oligonucleotide for the present invention. The methods include, but are not limited to: (1) chemical synthesis (usually, but not always, using a nucleic acid synthesizer instrument); (2) post-synthesis chemical modification or derivatization; (3) cloning of a naturally occurring or synthetic nucleic acid in a nucleic acid cloning vector (e.g., see Sambrook, *et al.*, Molecular Cloning: A Laboratory Approach Second Edition, 1989, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY and Sambrook and Russell, Molecular Cloning, A Laboratory Manual, Third Edition, 2001, Cold Spring Harbor Laboratory Press,) such as, but not limited to a plasmid, bacteriophage (e.g.,

M13 or lambda), phagemid, cosmid, fosmid, YAC, or BAC cloning vector, including vectors for producing single-stranded DNA; (4) primer extension using an enzyme with DNA template-dependent DNA polymerase activity, such as, but not limited to, Klenow, T4, T7, *rBst*, *Taq*, *Tfl*, or *Tth* DNA polymerases, including mutated, truncated (e.g., *exo-minus*), or chemically-modified forms of such enzymes; (5) PCR (e.g., see Dieffenbach, C.W., and Dveksler, eds., PCR Primer: A Laboratory Manual, 1995, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY); (6) reverse transcription (including both isothermal synthesis and RT-PCR) using an enzyme with reverse transcriptase activity, such as, but not limited to, reverse transcriptases derived from avian myeloblastosis virus (AMV), Moloney murine leukemia virus (MMLV), *Bacillus stearothermophilus* (*rBst*), or *Thermus thermophilus* (*Tth*); (7) *in vitro* transcription using an enzyme with RNA polymerase activity, such as, but not limited to, SP6, T3, or T7 RNA polymerase, *Tth* RNA polymerase, *E. coli* RNA polymerase, or SP6 or T7 R&DNA™ Polymerase (Epicentre Technologies, Madison, WI, USA), or another enzyme; (8) use of restriction enzymes and/or modifying enzymes, including, but not limited to *exo-* or *endonucleases*, kinases, ligases, phosphatases, methylases, glycosylases, terminal transferases, including kits containing such modifying enzymes and other reagents for making particular modifications in nucleic acids; (9) use of polynucleotide phosphorylases to make new randomized nucleic acids; (10) other compositions, such as, but not limited to, a ribozyme ligase to join RNA molecules; and/or (11) any combination of any of the above or other techniques known in the art. Oligonucleotides and polynucleotides, including chimeric (i.e., composite) molecules and oligonucleotides with modified bases, sugars, or internucleoside linkages are commercially available (e.g., TriLink Biotechnologies, San Diego, CA, USA or Integrated DNA Technologies, Coralville, IA).

[00295] A "portion" or "region," used interchangeably herein, of a polynucleotide or oligonucleotide is a contiguous sequence of 2 or more bases. In other embodiments, a region or portion is at least about any of 3, 5, 10, 15, 20, 25 contiguous nucleotides.

6. Ligation / Ligase

[00296] "Ligation" refers to the joining of a 5'-phosphorylated end of one nucleic acid molecule to a 3'-hydroxyl end of the same or another nucleic acid molecule by an enzyme called a "ligase." Alternatively, in some embodiments of the invention, ligation is effected by a type I topoisomerase moiety attached to one end of a nucleic acid (see U.S. Patent No. 5,766,891, incorporated herein by reference). The terms "ligating," "ligation," and "ligase" are often used

in a general sense herein and are meant to comprise any suitable method and composition for joining a 5'-end of one nucleic acid to a 3'-end of the same or another nucleic acid. Different ligases are preferred in different embodiments of the invention, as discussed elsewhere herein.

[00297] In general, if a nucleic acid to be ligated comprises RNA, a ligase such as, but not limited to, T4 RNA ligase, a ribozyme ligase, Tsc RNA Ligase (Prokaria Ltd., Reykjavik, Iceland), or another ligase can be used for non-homologous joining of the ends. T4 DNA ligase can also be used to ligate RNA molecules when a 5'-phosphoryl end is adjacent to a 3'-hydroxyl end annealed to a complementary sequence (e.g., see U.S. Patent No. 5,807,674 of Tyagi, incorporated herein by reference).

[00298] If the nucleic acids to be joined comprise DNA and the 5'-phosphorylated and the 3'-hydroxyl ends are ligated when the ends are annealed to a complementary DNA so that the ends are adjacent (such as, when a "ligation splint" is used), then enzymes such as, but not limited to, T4 DNA ligase, Ampligase® DNA Ligase (Epicentre Technologies, Madison, WI, USA), Tth DNA ligase, Tfl DNA ligase, or Tsc DNA Ligase (Prokaria Ltd., Reykjavik, Iceland) can be used. However, the invention is not limited to the use of a particular ligase and any suitable ligase can be used. Still further, Faruqi discloses in U.S. Patent No. 6,368,801 that T4 RNA ligase can efficiently ligate DNA ends of nucleic acids that are adjacent to each other when hybridized to an RNA strand. Thus, T4 RNA ligase is a suitable ligase of the invention in embodiments in which DNA ends are ligated on a ligation splint oligo comprising RNA or modified RNA, such as, but not limited to modified RNA that contains 2'-F-dCTP and 2'-F-dUTP made using the DuraScribe™ T7 Transcription Kit (Epicentre Technologies, Madison, WI, USA) or the N4 mini-vRNAP Y678F mutant enzyme described herein. With respect to ligation on an homologous ligation template, especially ligation using a "ligation splint" or a "ligation splint oligo" (as discussed elsewhere herein), a region, portion, or sequence that is "adjacent" to another sequence directly abuts that region, portion, or sequence.

[00299] In other embodiments comprising intramolecular ligation of linear ssDNA, ligation can be effected in the absence of a ligation splint using a ligase that can catalyze non-homologous ligation of ssDNA, such as, but not limited to, ThermoPhage™ RNA Ligase II (Prokaria Ltd., Reykjavik, Iceland), which is derived from phage TS2126 that infects *Thermus scotoductus*.

7. DNA Polymerases/Strand-Displacing DNA Polymerases/Strand Displacement/Rolling Circle Replication

[00300] A “DNA-dependent DNA polymerase” is an enzyme that synthesizes a complementary DNA (“cDNA”) copy from a DNA template. Examples are DNA polymerase I from *E. coli* and bacteriophage T7 DNA polymerase. All known DNA-dependent DNA polymerases require a complementary primer to initiate synthesis. It is known that under suitable conditions a DNA-dependent DNA polymerase may synthesize (i.e., “reverse transcribe”) a complementary DNA copy from an RNA template, a process that is also referred to as “reverse transcription.”

[00301] Some DNA polymerases are able to displace the strand complementary to the template strand as a new DNA strand is synthesized by the polymerase. This process is called “strand displacement” and the DNA polymerases that have this activity are referred to herein as “strand-displacing DNA polymerases.” The template for strand displacement DNA synthesis using a method of the invention can be a linear or circular ssDNA. If the DNA template is a single-stranded circle, primed DNA synthesis proceeds around and around the circle, with continual displacement of the strand ahead of the replicating strand, a process called “rolling circle replication.” Rolling circle replication results in synthesis of tandem copies of the circular template. The suitability of a DNA polymerase for use in an embodiment of the invention that comprises strand displacement on linear templates or rolling circle replication can be readily determined by assessing its ability to carry out rolling circle replication. By way of example, but not of limitation, the ability of a polymerase to carry out rolling circle replication can be determined by using the polymerase in a rolling circle replication assay such as those described by Fire and Xu (Proc. Natl. Acad. Sci. USA 92:4641-4645, 1995), incorporated herein by reference. It is preferred that a DNA polymerase be a strand displacing DNA polymerase and lack a 5'-to-3' exonuclease activity for strand displacement polymerization reactions using both linear or circular templates since a 5'-to-3' exonuclease activity, if present, might result in the destruction of the synthesized strand. It is also preferred that DNA polymerases for use in the disclosed strand displacement synthesis methods are highly processive. The ability of a DNA polymerase to strand-displace can vary with reaction conditions, in addition to the particular enzyme used. Strand displacement and DNA polymerase processivity can also be assayed using methods described in Kong *et al.* (J. Biol. Chem. 268:1965-1975, 1993 and references cited therein, all of which are incorporated herein by reference).

[00302] Preferred strand displacing DNA polymerases of the invention are rBst DNA polymerase large fragment (also called IsoTherm™ DNA Polymerase (Epicentre Technologies, Madison, WI, USA), BcaBEST™ DNA polymerase (Takara Shuzo Co., Kyoto, Japan), ϕ 29 DNA polymerase (U.S. Pat. Nos. 5,576,204 and 5,001,050 to Blanco *et al.*, incorporated herein by reference), SequiTherm™ DNA polymerase (Epicentre Technologies, Madison, WI, USA), and MMLV reverse transcriptase. Other strand-displacing DNA polymerases which can be used include, but are not limited to, phage M2 DNA polymerase (Matsumoto *et al.*, *Gene* 84:247, 1989), phage ϕ PRD1 DNA polymerase (Jung *et al.*, *Proc. Natl. Acad. Sci. USA* 84: 8287, 1987), VENT® DNA polymerase (Kong *et al.*, *J. Biol. Chem.* 268:1965-1975, 1993), Klenow fragment of DNA polymerase I (Jacobsen *et al.*, *Eur. J. Biochem.* 45:623-627, 1974), T5 DNA polymerase (Chatterjee *et al.*, *Gene* 97:13-19, 1991), PRD1 DNA polymerase (Zhu and Ito, *Biochim. Biophys. Acta* 1219: 267-276, 1994), modified T7 DNA polymerase (Tabor and Richardson, *J. Biol. Chem.* 262:15,330-15,333, 1987); Tabor and Richardson, *J. Biol. Chem.* 264:6447-6458, 1989); Sequenase™ (U.S. Biochemicals, Cleveland, OH, USA), and T4 DNA polymerase holoenzyme (Kaboord and Benkovic, *Curr. Biol.* 5:149-157, 1995), all of which references, are incorporated herein by reference. Strand displacing DNA polymerases are also useful in some embodiments of the invention for strand displacement replication of linear first-strand cDNA, and in other embodiments, for rolling circle replication of circular first-strand cDNA. IsoTherm™ DNA polymerase (rBst DNA polymerase large fragment; Epicentre) is most preferred because, in addition to having strand-displacing DNA polymerase activity, it can also be used as a reverse transcriptase for synthesis of first-strand cDNA from RNA target nucleic acids (e.g., U.S. Patent No. 6,030,814 of Jendrisak *et al.*, incorporated herein by reference). BcaBEST™ DNA polymerase (Takara Shuzo Co., Kyoto, Japan) can also be used as a reverse transcriptase as well as a strand-displacing DNA polymerase.

[00303] Strand displacement can be facilitated through the use of a strand displacement factor, such as helicase. It is considered that any DNA polymerase that can perform rolling circle replication in the presence of a strand displacement factor is suitable for use in embodiments of the invention that comprise strand displacement or rolling circle replication, even if the DNA polymerase does not perform rolling circle replication in the absence of such a factor. Strand displacement factors useful in rolling circle replication include, but are not limited to, BMRF1 polymerase accessory subunit (Tsurumi *et al.*, *J. Virology* 67:7648-7653, 1993), adenovirus DNA-binding protein (Zijderveld and van der Vliet, *J. Virology* 68:1158-1164, 1994), herpes simplex viral protein ICP8 (Boehmer and Lehman, *J. Virology* 67:711-715,

1993); Skaliter and Lehman, Proc. Natl. Acad. Sci. USA 91:10,665-10,669, 1994), single-stranded DNA binding proteins (SSB; Rigler and Romano, J. Biol. Chem. 270:8910-8919, 1995), and calf thymus helicase (Siegel et al., J. Biol. Chem. 267:13,629-13,635, 1992), all of which are incorporated herein by reference.

8. Hybridize/Hybridization

[00304] The terms "hybridize" and "hybridization" refer to the formation of complexes between nucleotide sequences that are sufficiently complementary to form complexes via Watson-Crick base pairing. With respect to the present invention, nucleic acid sequences that "hybridize" or "anneal" with each other should form "hybrids" or "complexes" that are sufficiently stable to serve the intended purpose. By way of example, but not of limitation, where a primer or splice template oligo hybridizes or anneals with a target nucleic acid in a sample or with a "tailed" target sequence, respectively, each respective complex or hybrid should be sufficiently stable to serve the respective priming functions required for a DNA polymerase to copy the target sequence by primer extension of the annealed primer or to extend the 3'-end of the target sequence using the annealed splice template oligo as a template, respectively.

9. Reaction Conditions

[00305] Appropriate reaction media and conditions for carrying out the methods of the present invention are those that permit nucleic acid transcription and other reactions according to the methods of the present invention. For example, the conditions below can be used for *in vitro* transcription with N4 vRNAP, mini-vRNAP, and mini-vRNAP Y678F enzymes of the invention. An *in vitro* transcription reaction is prepared by setting up a reaction mixture containing the following final concentrations of components, added in the order given: 0.1 micromolar of a N4 vRNAP promoter-containing DNA oligo; 1.0 micromolar *Eco*SSB Protein; 1X transcription buffer comprising 40 mM Tris-HCl (pH 7.5), 6 mM MgCl₂, 2 mM spermidine, and 10 mM NaCl; 1 mM DTT; 0.5 mM of each NTP (ATP, CTP, GTP and UTP); deionized RNase-free water so the final volume will be 50 microliters after addition of an RNAP; and 0.1 micromolar of N4 vRNAP, mini-vRNAP or mini-vRNAP Y678F enzyme. In some embodiments of the invention, 2'-F-dUTP and 2'-F-dCTP are used at a final concentration of 0.5 mM each in place of UTP and CTP in order to obtain synthesis of modified RNA which is resistant to ribonuclease A-type enzymes. Other modified nucleoside triphosphates can also be

used in place of or in addition to the canonical NTPs for specific applications. The reaction mixture is then incubated at 37° C to permit synthesis of RNA from the template. The reaction can be followed by gel electrophoresis on a PAGE gel.

[00306] The invention is not limited to these reaction conditions or concentrations of reactants. Those with skill in the art will know that other suitable reaction conditions under which an RNA polymerase of the invention can be used can be found by simple experimentation, and any of these reaction conditions are also included within the scope of the invention. Such media and conditions are known to persons of skill in the art, and are described in various publications, such as U.S. Pat. No. 5,679,512 and PCT Pub. No. WO99/42618, incorporated herein by reference. For example, a buffer maybe Tris buffer, although other buffers can also be used as long as the buffer components are non-inhibitory to enzyme components of the methods of the invention. The pH is preferably from about 5 to about 11, more preferably from about 6 to about 10, even more preferably from about 7 to about 9, and most preferably from about 7.5 to about 8.5. The reaction medium can also include bivalent metal ions such as Mg^{+2} or Mn^{+2} , at a final concentration of free ions that is within the range of from about 0.01 to about 10 mM, and most preferably from about 1 to 6 mM. The reaction medium can also include other salts, such as KCl, that contribute to the total ionic strength of the medium. For example, the range of a salt such as KCl is preferably from about 0 to about 100 mM, more preferably from about 0 to about 75 mM, and most preferably from about 0 to about 50 mM. The reaction medium can further include additives that could affect performance of the reactions, but that are not integral to the activity of the enzyme components of the methods. Such additives include proteins such as BSA, and non-ionic detergents such as NP40 or Triton. Reagents, such as DTT, that are capable of maintaining activities enzyme with sulfhydryl groups can also be included. Such reagents are known in the art. Where appropriate, an RNase inhibitor, such as, but not limited to a placental ribonuclease inhibitor (e.g., RNasin®, Promega Corporation, Madison, WI, USA) or an antibody RNase inhibitor, that does not inhibit the activity of an RNase employed in the method can also be included. Any aspect of the methods of the present invention can occur at the same or varying temperatures. Preferably, the reactions are performed isothermally, which avoids the cumbersome thermocycling process. The reactions are carried out at a temperature that permits hybridization of the oligonucleotides of the present invention to the target sequence and/or first-strand cDNA of a method of the invention and that does not substantially inhibit the activity of the enzymes employed. The temperature can be in the range of preferably about 25° C to about 85° C, more preferably about

30° C to about 75° C, and most preferably about 37° C to about 70° C. In the processes that include RNA transcription, the temperature for the transcription steps is lower than the temperature(s) for the preceding steps. In these processes, the temperature of the transcription steps can be in the range of preferably about 25° C to about 85° C, more preferably about 30° C to about 75° C, and most preferably about 37° C to about 55° C.

[00307] As disclosed in U.S. Patent Nos. 6,048,696 and 6,030,814, as well as in German Patent No. DE4411588C1, all of which are incorporated herein by reference and made part of the present invention, it is preferred in many embodiments to use a final concentration of about 0.25 M, about 0.5 M, about 1.0 M, about 1.5 M, about 2.0 M, about 2.5 M or between about 0.25 M and 2.5 M betaine (trimethylglycine) in DNA polymerase or reverse transcriptase reactions in order to decrease DNA polymerase stops and increase the specificity of reactions which use a DNA polymerase.

[00308] Nucleotide and/or nucleotide analogs, such as deoxyribonucleoside triphosphates, that can be employed for synthesis of reverse transcription or primer extension products in the methods of the invention are provided in an amount that is determined to be optimal or useful for a particular intended use.

[00309] The oligonucleotide components of reactions of the invention are generally in excess of the number of target nucleic acid sequence to be amplified. They can be provided at about or at least about any of the following: 10 , 10^2 , 10^4 , 10^6 , 10^8 , 10^{10} , 10^{12} times the amount of target nucleic acid. Promoter splice template oligos, splice templates, promoter ligation oligos, ligation splint oligos, promoter primers, blocker sequence oligos, and strand-displacement primers, and the like, can each be provided at about or at least about any of the following concentrations: 50 nM, 100 nM, 500 nM, 1000 nM, 2500 nM, 5000 nM, or 10,000 nM, but higher or lower concentrations can also be used. By way of example, but not of limitation, a concentration of one or more oligonucleotides may be desirable for production of one or more target nucleic acid sequences that are used in another application or process. The invention is not limited to a particular concentration of an oligonucleotide, so long as the concentration is effective in a particular method of the invention.

[00310] In some embodiments, the foregoing components are added simultaneously at the initiation of the process. In other embodiments, components are added in any order prior to or after appropriate time points during the process, as required and/or permitted by the reaction. Such time points can readily be identified by a person of skill in the art. The enzymes used for nucleic acid reactions according to the methods of the present invention are generally added to

the reaction mixture following a step for denaturation of a double-stranded target nucleic acid in or from a sample, and/or following hybridization of primers and/or oligos of a reaction to a denatured double-stranded or single-stranded target nucleic acid, as determined by their thermal stability and/or other considerations known to the person of skill in the art.

[00311] The reactions can be stopped at various time points, and resumed at a later time. The time points can readily be identified by a person of skill in the art. Methods for stopping the reactions are known in the art, including, for example, cooling the reaction mixture to a temperature that inhibits enzyme activity. Methods for resuming the reactions are also known in the art, including, for example, raising the temperature of the reaction mixture to a temperature that permits enzyme activity. In some embodiments, one or more of the components of the reactions is replenished prior to, at, or following the resumption of the reactions. Alternatively, the reaction can be allowed to proceed (i.e., from start to finish) without interruption.

10. Detection and Identification of Reaction Products

[00312] In some embodiments, the detection of the product is indicative of the presence of the target sequence. Quantitative analysis, including analysis in real time, can also be performed in some embodiments. Direct and indirect detection methods (including quantification) are well known in the art. For example, by comparing the amount of product amplified from a test sample containing an unknown amount of a polynucleotide containing a target sequence to the product of a reference sample that has a known quantity of a polynucleotide that contains the target sequence, the amount of target sequence in the test sample can be determined. The methods of the present invention can also be extended to analysis of sequence alterations and sequencing of the target nucleic acid. The amplified nucleic acid can be sequenced using any suitable procedure. Many such procedures are known. Preferred forms of sequencing for use with amplified sequences produced from some embodiments are nanosequencing methods described by Jalanko *et al.*, Clinical Chemistry 38:39-43, 1992; Nikiforov *et al.*, Nucleic Acids Research 22:4167-4175, 1994; and Kobayashi *et al.*, Molecular and Cellular Probes 9:175-182, 1995, and primer extension sequencing, as described in PCT Application WO 97/20948, all of which references are included herein by reference. Further, detection could be effected by, for example, examination of translation products from RNA products.

B. Methods for Obtaining a ssDNA Comprising a Target Sequence

1. General Aspects and Methods for Obtaining a Target Sequence

[00313] An initial step in obtaining a target sequence is rendering the target nucleic acid single-stranded. If the target nucleic acid is a double-stranded DNA (dsDNA), the initial step is target denaturation. The denaturation step may be thermal denaturation or any other method known in the art, such as alkali treatment.

[00314] In some embodiments of the invention in which the target nucleic acid in a sample is DNA, the ssDNA target sequence comprises either ssDNA that is present in a biological sample or ssDNA that is obtained by denaturation of dsDNA in the sample.

[00315] In other embodiments, the ssDNA target sequence comprises ssDNA that is obtained as a result of a "primer extension reaction," meaning an *in vitro* or *in vivo* DNA polymerization reaction using either ssDNA or denatured dsDNA that is present in the sample as a template and an oligonucleotide as a primer under DNA polymerization reaction conditions. A "primer" is an oligonucleotide (oligo), generally with a free 3'-OH group, for which at least the 3'-portion of the oligo is complementary to a portion of the template and which oligo "binds" (or "complexes" or "anneals" or "hybridizes"), by hydrogen bonding and other molecular forces, to the template to give a primer/template complex for initiation of synthesis by a DNA polymerase, and which is extended (i.e., "primer extended") by the addition of covalently bonded bases linked at its 3'-end which are complementary to the template in the process of DNA synthesis. The result is a primer extension product. Virtually all DNA polymerases (including reverse transcriptases) that are known require complexing of an oligonucleotide to a single-stranded template ("priming") to initiate DNA synthesis, whereas RNA replication and transcription (copying of RNA from DNA) generally do not require a primer.

[00316] In some embodiments, the target nucleic acid in the sample or the primer extension product, or both, are made into smaller DNA fragments by methods known in the art in order to generate a DNA target sequence. In some embodiments using samples containing DNA target nucleic acids, a ssDNA target sequence and/or a transcription substrate is obtained using a strand displacement method, such as but not limited to, a methods described in PCT Patent Publication Nos. WO 02/16639; WO 00/56877; and AU 00/29742; of Takara Shuzo Company, Kyoto, Japan; U.S. Patent Nos. 5,523,204; 5,536,649; 5,624,825; 5,631,147; 5,648,211; 5,733,752; 5,744,311; 5,756,702; and 5,916,779 of Becton Dickinson and Company; U.S. Patent Nos. 6,238,868; 6,309,833; and 6,326,173 of Nanogen/ Becton Dickinson Partnership; U.S. Patent Nos. 5,849,547; 5,874,260; and 6,218,151 of Bio Merieux; U.S. Patent

Nos. 5,786,183; 6,087,133; and 6,214,587 of Gen-Probe, Inc.; U.S. Patent No. 6,063,604 of Wick *et al.*; U.S. Patent No. 6,251,639 of Kurn; U.S. Patent No. 6,410,278; and PCT Publication No. WO 00/28082 of Eiken Kagaku Kabushiki Kaishi, Tokyo, Japan; U.S. Patent Nos. 5,591,609; 5,614,389; 5,773,733; 5,834,202; and 6,448,017 of Auerbach; and U.S. Patent Nos. 6,124,120; and 6,280,949 of Lizardi, all of which are incorporated herein by reference. In still other embodiments, the ssDNA target sequence is obtained from a rolling circle replication reaction. The 3'-end of the DNA target sequence can be defined, if it need be defined, by using any suitable method known in the art, such as, but not limited to, a method discussed in the section herein entitled "Methods for Defining the 5'- and 3'-Ends of Target Sequences That Comprise Only a Portion of a Larger RNA or DNA Target Nucleic Acid."

[00317] If the target nucleic acid is RNA, the initial step for obtaining a target sequence comprises synthesis of a single-stranded first-strand cDNA by reverse transcription of the RNA target, meaning an *in vitro* reaction that utilizes an RNA present in a sample as a template and a nucleic acid oligonucleotide that is complementary to at least a portion of a sequence of the RNA template as a primer in order to synthesize ssDNA using an RNA-dependent DNA polymerase (i.e., reverse transcriptase) under reaction conditions. Techniques for the synthesis of cDNA from RNA are known in the art. In some embodiments, a first-strand cDNA for use in methods of the invention is synthesized *in situ* in cells or tissue in a tissue section using methods similar to those described in U.S. Patent Nos. 5,168,038; 5,021,335; and 5,514,545, which are incorporated herein by reference. Thus, the first-strand cDNA is synthesized by contacting the cells or tissue in the tissue section under hybridizing conditions with a primer, wherein the primer hybridizes to one or more target sequences in the cell or tissue. The present invention comprises a method for making a transcription substrate comprising first-strand cDNA that is complementary to a target sequence in cells or tissue in a tissue section, the method comprising: (a) contacting the cells or tissue in the tissue section under hybridizing conditions with a promoter primer, the promoter primer comprising (i) a 5'-portion comprising a ssDNA transcription promoter for an RNA polymerase that synthesizes RNA using a ssDNA transcription substrate, and (ii) a 3'-end that is complementary to a target sequence in a target nucleic acid; and (b) contacting the cells or tissue containing the promoter primer in the tissue section with a reverse transcriptase under reverse transcription conditions so as to obtain a first-strand cDNA that is complementary to the target sequence; and (c) obtaining first-strand cDNA, wherein the first-strand cDNA comprises a ssDNA transcription substrate for the an RNA polymerase that binds to the promoter.

[00318] An oligonucleotide primer can be complementary to a specific known sequence in the RNA target in a sample, or an oligonucleotide primer can comprise a mixture of all possible or many possible sequences, such as, but not limited to, a primer comprising a random hexamer priming sequence. Random primers can be made by using an oligonucleotide synthesizer by including nucleotide reagents that are complementary to each of the four canonical bases (i.e., all four nucleotides) during the chemical synthesis of each nucleotide position of the oligonucleotide that is complementary to the target sequence. In embodiments of the invention using samples containing mRNA targets, the ssDNA target sequence comprises first-strand cDNA that is made by reverse transcription of the mRNA using an oligonucleotide primer comprising either a specific sequence which is complementary to a known sequence of a specific mRNA or, if the mRNA has a poly(A) tail at its 3'-end, an oligo(dT) primer or an oligo(dT) anchor primer. In other embodiments of the invention, a promoter primer is used, which serves both to prime synthesis of the first-strand cDNA target sequence and to join a transcription promoter to the target sequence.

2. Methods for Defining the 5'- and 3'-Ends of Target Sequences That Comprise Only a Portion of a Larger RNA or DNA Target Nucleic Acid

[00319] When a method of the invention is used to obtain a transcription product corresponding to the complete sequence(s) of one or a multitude of nucleic acid molecules, such as, but not limited to the complete sequences (excluding the cap structure) of substantially all polyadenylated mRNA molecules in a sample, it is not necessary to devise methods to define the 5'- and 3'-ends of the sequences. However, if a method of the invention is used to obtain a transcription product corresponding to a target sequence that comprises only a portion of a larger RNA or DNA nucleic acid in a sample, then methods are needed to delimit the target sequence that becomes the transcription template.

[00320] There are two general approaches for delimiting the ends of the target sequence that becomes the transcription template sequence. In the first direct approach, methods are used to determine the size and end sequences of a target nucleic acid molecule or molecules present in the sample itself. In the second indirect approach, instead of changing the size and end sequences of the target nucleic acid molecules present in a sample, methods are used to determine the size and end sequences of one or more first-strand cDNA molecules that is/are synthesized by reverse transcription or primer extension, respectively, of RNA or of at least one strand of DNA in a sample.

[00321] With respect to the direct approach, a number of methods are known in the art for cleaving a nucleic acid molecule at or near a specific sequence, and any of the methods which delimit the size and end sequences of a target nucleic acid for an application of the present invention can be used. By way of example, but not of limitation, a DNA in a sample comprising a dsDNA molecule or a ssDNA molecule to which an appropriate complementary DNA oligo is annealed can be digested with a restriction endonuclease, provided a restriction site that would provide a suitable 5'-end and/or 3'-end sequence is present. Alternatively, one or more DNA oligonucleotides having a double-stranded segment that contains a *FokI* restriction enzyme site and a single-stranded segment that binds to the desired cleavage site on a first-strand cDNA can be used. As is well known in the art, this type of oligonucleotide can be used with the restriction enzyme *FokI* to cut a single-stranded DNA at almost any desired sequence (Szybalski, W., Gene 40:169-173, 1985; Podhajski A. J. and Szybalski W., Gene 40:175, 1985, incorporated herein by reference).

[00322] By way of further example, but not of limitation, a ssRNA target nucleic acid present in a sample can be cleaved using a ribonuclease H in regions to which complementary oligonucleotides comprising at least three-to-four deoxynucleotides are annealed. Alternatively, a linear DNA oligonucleotide can be annealed to an RNA in a sample at a location that encodes a recognition site of a restriction enzyme that can cut RNA:DNA heteroduplexes. Cutting the target RNA:DNA oligo with the enzyme will then generate a defined end. Alternatively, an RNA or DNA oligo or polynucleotide with a sequence complementary to the region of an RNA target sequence that is intended to become a transcription substrate can be annealed to the RNA and the sequences of the RNA to which the oligo or polynucleotide is not annealed can be digested using a single-strand-specific ribonuclease, such as RNase A or RNase T1. Still further, either RNA or DNA nucleic acids of known sequence can be cleaved at specific sites using a 5'-nuclease or Cleavase™ enzyme and specific oligonucleotides, as described by Kwiatkowski, et al., (Molecular Diagnosis 4:353-364, 1999) and in U.S. Patent No. 6,001,567 and related patents assigned to Third Wave Technologies (Madison, WI, USA), which are incorporated herein by reference.

[00323] In general, with respect to the second indirect approach, the 5'-end of the primer that is used for reverse transcription of RNA in a sample or for primer extension of at least one strand of DNA in a sample defines the 5'-end of the first-strand cDNA target sequence. Thus, a sample target nucleic acid that is reverse transcribed or primer extended to make a first-strand cDNA target sequence need not have a defined 3'-end.

[00324] In order to generate a defined 3'-end on a first-strand cDNA (i.e., corresponding to the 5'-end of the target sequence), a number of methods can be used to obtain a target sequence of the present invention. By way of example, but not of limitation, if a specific sequence is present in the first-strand cDNA that corresponds to a restriction endonuclease site that would provide a suitable 3'-end sequence, a complementary DNA oligo can be annealed to this sequence and the site can be cleaved with the restriction enzyme. The complementary DNA oligo used to provide the double-stranded restriction site can optionally have a 2', 3'-dideoxynucleotide or another terminator nucleotide at its 3'-end so that it cannot be extended by a DNA polymerase. Alternatively, the 3'-end of the target sequence can be defined using a DNA oligonucleotide having a double-stranded segment that contains a *FokI* restriction enzyme site and a single-stranded segment that binds to the desired cleavage site on a first-strand cDNA (Szybalski, W., Gene 40:169-173, 1985; Podhajski A. J. and Szybalski W., Gene 40:175, 1985). Still further, a 5'-nuclease can be used to cleave a first-strand cDNA at a defined 3'-end as discussed above.

[00325] In addition to the above methods, the 3'-end of a first-strand cDNA can also be limited by other methods. A preferred method of the invention is to use a "blocking oligo" or a "blocker sequence," as disclosed by Laney, et al. in U.S. Patent No. 5,679,512, and by Kurn in U.S. Patent No. 6,251,639, both of which are incorporated herein by reference. The "blocker sequence" or "blocker oligo" is a polynucleotide, which is usually a synthetic polynucleotide that is single-stranded and comprises a sequence that is hybridizable, and preferably complementary, to a segment of target nucleic acid, wherein the blocking oligo anneals to the target nucleic acid so as to block further primer extension of the 3'-end of first-strand cDNA at a desired position. Some embodiments of strand displacement methods of the present invention for obtaining a ssDNA transcription substrate comprise use of a blocking oligo. The blocking oligo comprises nucleotides that bind to the target nucleic acid with an affinity, preferably a high affinity, such that the blocker sequence resists displacement by DNA polymerase in the course of primer extension, in preferably more than about 30%, more preferably more than about 50%, even more preferably more than about 75%, and most preferably more than about 90%, of primer extension events. The length and composition of the blocker polynucleotide should be such that excessive random non-specific hybridization is avoided under the conditions of the methods of the present invention. The length of the blocker polynucleotide is preferably from about 3 to about 30 nucleotides, more preferably from about 5 to about 25 nucleotides, even more preferably from about 8 to about 20 nucleotides, and most preferably from about 10 to

about 15 nucleotides. In other embodiments, the blocker polynucleotide is at least about any of the following: 3, 5, 8, 10, 15; and less than about any of the following: 20, 25, 30, 35. It is understood that the length can be greater or less as appropriate under the reaction conditions of the methods of this invention. The complementarity of the blocker polynucleotide is preferably at least about 25%, more preferably at least about 50%, even more preferably at least about 75%, and most preferably at least about 90%, to its intended binding sequence on the target nucleic acid. In some embodiments, the blocker sequence that hybridizes to a DNA target nucleic acid is attached to the DNA such that displacement of the blocker sequence by the polymerase that affects primer extension is substantially, or at least sufficiently, inhibited. Suitable methods for achieving such attachment include techniques known in the art, such as using a cytosine analog that contains a G-clamp heterocycle modification as described by Flanagan *et al.*, (*Proc. Natl. Acad. Sci. USA* 96:3513-3518, 1999); and locked nucleic acids as described, e.g., by Kumar *et al.*, (*Bioorg. Med. Chem. Lett.* 8:2219-2222, 1998; and by Wahlestedt *et al.* (*Proc. Natl. Acad. Sci. USA* 97:5633-5638, 2000), all of which are incorporated herein by reference. Other suitable methods include using, where appropriate, sequences with a high GC content and/or cross-linking. Any of these methods for obtaining enhanced attachment may be used alone or in combination. Alternatively, a molecule comprising a peptide nucleic acid ("PNA") can be used.

[00326] Still further, another method that can be used to limit the 3'-end of a first-strand cDNA is to use a thermocycler with short DNA synthesis elongation cycles during reverse transcription or primer extension to synthesize first-strand cDNA. The length of the primer extension product can be somewhat controlled by the length of the DNA synthesis cycle. Conditions can be determined to define an approximate chain length of first-strand cDNA by controlling the temperature and time interval of DNA synthesis before denaturing the growing first-strand cDNA from the template by raising the temperature.

[00327] Further, the 3'-end of a first-strand cDNA that is to become the template sequence for a transcription reaction can be defined by first amplifying the target nucleic acid sequence using any suitable amplification method that delimits the end sequence. By way of example, but not of limitation, it can be prepared using PCR, RT-PCR, NASBA, TMA, 3SR, Ligation Chain Reaction (LCR), Linked Linear Amplification (BioRad), SDA, RCA, ICANTM (Takara: Sagawa *et al.* in PCT Patent Publication No. WO 02/16639; and in PCT Patent Publications Nos. WO 00/56877 and AU 00/29742; or a strand-displacement method of Kurn (U.S. Patent No. 6,251,639), all of which are incorporated herein by reference.

[00328] If a 3'-end of a target sequence need not be at an exact location, and can be random or imprecise, which is the case in some embodiments of the invention, there are a number of other methods that can be used for making smaller fragments of a DNA molecule, whether for a target nucleic acid, a target sequence, or otherwise. By way of example, but not of limitation, a target nucleic acid can be fragmented by physical means, such as by movement in and out of a syringe needle or other orifice or by sonication, preferably with subsequent end repair, such as using a T4 DNA polymerase or a kit, such as the End-It™ DNA End Repair Kit (Epicentre Technologies, Madison, WI, USA). Still another method that can be used is to incorporate dUMP randomly into the first-strand cDNA during reverse transcription or primer extension by using dUTP in place of a portion of the TTP in the reaction. The dUMP will be incorporated randomly in place of TMP at a frequency based on the ratio of dUTP to TTP. Then, the first-strand cDNA can be cleaved at sites of dUMP incorporation by treatment (e.g., see U.S. Patent No. 6,048,696, incorporated herein by reference) with uracil-N-glycosylase (UNG) and endonuclease IV (endo IV), which are available from Epicentre Technologies (Madison, WI, USA). UNG hydrolyzes the N-glycosidic bond between the deoxyribose sugar and uracil in single- and double-stranded DNA that contains uracil in place of thymidine. It has no activity on uracil residues in RNA or on dUTP. Endo IV cleaves the phosphodiester linkage at the abasic site. It may be useful to use a thermolabile UNG (e.g., HK™-UNG from Epicentre Technologies, Madison, Wisconsin, USA) for some applications. (Also, incorporation of dUMP at specific sites within a synthetic oligonucleotide or, for example, within a promoter primer of the invention between the 3'-target-sequence-complementary portion and the promoter sequence, introduces specific cleavage sites which can be used at any time to cleave a resulting nucleic acid which contains the site by treatment with UNG and endo IV.) Still further, in some cases, the 3'-end of a first-strand cDNA can be defined by treatment with exonuclease III (Henikoff, S., *Gene* 28:351, 1984). In still other cases, the 3'-end of a first-strand cDNA that is annealed to a DNA target nucleic acid can be incubated with T4 DNA polymerase or unmodified T7 DNA polymerase (Epicentre Technologies, Madison, WI) in the absence or the presence of dNTPs in the reaction; these enzymes have the 3'-to-5' exonuclease activity in the absence of dNTPs, but the polymerase activity predominates in the presence of dNTPs. These are only some of the methods that can be used to define the 3'-ends of a first-strand cDNA, and the invention is not limited to these methods, which are presented only as examples.

C. Methods for Obtaining a Transcription Substrate

1. Introduction

[00329] The invention includes a number of embodiments for obtaining a transcription substrate. In a first general embodiment, a ssDNA transcription substrate is obtained by using a promoter splice template oligo, comprising an anti-sense transcription promoter in its 5'-portion, as a template for DNA polymerase- or reverse transcriptase-catalyzed extension of the 3'-end of first-strand cDNA that is complementary to a target sequence. This first general embodiment results in a transcription substrate with a sense transcription promoter 3'-of a target sequence. In a second general embodiment, a ssDNA transcription substrate is obtained by ligating a promoter ligation oligo, comprising a sense transcription promoter, to the 3'-end of a target sequence. In this embodiment, the target sequence can comprise first-strand cDNA that is complementary to a DNA or RNA target sequence or, if the target nucleic acid is DNA and the 3'-end of the target sequence is suitably delimited, the promoter ligation oligo can be ligated directly onto the 3'-end of the target sequence. In a third general embodiment, a ssDNA transcription substrate is obtained by synthesizing first-strand cDNA by a reverse transcription or primer extension of a promoter primer comprising a sense transcription promoter in its 5'-portion and a sequence complementary to the target sequence at its 3'-end and then ligating the resulting linear first-strand cDNA to obtain a circular ssDNA transcription substrate comprising a promoter-containing circular first-strand cDNA. A linear ssDNA transcription substrate resulting from the first or second general embodiments can also be ligated to make a circular ssDNA transcription substrate. In still other embodiments, any of the above circular ssDNA transcription substrates is linearized 3'-of a sense transcription promoter sequence to obtain a linear ssDNA transcription substrate having the promoter 3'-of the target sequence.

[00330] In still another embodiment, a ssDNA transcription substrate comprising second-strand cDNA is obtained by (a) synthesizing first-strand cDNA using a reverse transcriptase or a DNA polymerase and a primer comprising an anti-sense transcription promoter in its 5'-portion and a sequence complementary to the target sequence at its 3'-end; (b) ligating the resulting linear first-strand cDNA to obtain a circular first-strand cDNA having an anti-sense transcription promoter; and (c) obtaining a linear second-strand cDNA comprising a ssDNA transcription substrate by DNA synthesis using the circular first-strand cDNA as a template, a primer with a 3'-end complementary to a sequence on first-strand cDNA and a DNA polymerase, preferably a strand-displacing DNA polymerase that results in strand-displacement DNA synthesis of the

linear transcription substrate. In this embodiment, the transcription product of the transcription substrate comprises an anti-sense transcription product.

[00331] In yet another embodiment, a ssDNA transcription substrate comprising second-strand cDNA is obtained by (a) synthesizing linear first-strand cDNA having an anti-sense promoter 3'-of a target sequence using a primer, a reverse transcriptase or DNA polymerase, and a splice template oligo having sense promoter in its 5'-portion and a sequence complementary to the target sequence at its 3'-end; (b) ligating the resulting linear first-strand cDNA to obtain a circular first-strand cDNA having an anti-sense transcription promoter; and (c) obtaining a linear second-strand cDNA comprising a ssDNA transcription substrate by DNA synthesis using the circular first-strand cDNA as a template, a primer with a 3'-end complementary to a sequence on the first-strand cDNA and a DNA polymerase, preferably a strand-displacing DNA polymerase that results in strand-displacement DNA synthesis of the linear transcription substrate. In this embodiment, the transcription product of the transcription substrate comprises an anti-sense transcription product.

[00332] In still another embodiment, a ssDNA transcription substrate comprising second-strand cDNA is obtained by (a) synthesizing linear first-strand cDNA using a primer and a reverse transcriptase or DNA polymerase; (b) ligating an oligonucleotide comprising an anti-sense promoter to the 3'-end of the first-strand cDNA target sequence; (c) ligating the resulting linear first-strand cDNA having an anti-sense promoter to obtain a circular first-strand cDNA having an anti-sense transcription promoter; and (c) synthesizing a linear second-strand cDNA, comprising a ssDNA transcription substrate, by DNA synthesis using the circular first-strand cDNA as a template, a primer with a 3'-end complementary to a sequence on first-strand cDNA and a DNA polymerase, preferably a strand-displacing DNA polymerase that results in strand-displacement DNA synthesis of the linear transcription substrate. In this embodiment, the transcription product of the transcription substrate comprises an anti-sense transcription product.

[00333] In all of the above embodiments for obtaining a transcription substrate comprising second-strand cDNA, the transcription substrate, including the promoter sequence, is nevertheless single-stranded, rather than double-stranded as used in the prior art.

[00334] In addition, the invention also comprises embodiments in which a linear ssDNA transcription substrate comprising second-strand cDNA is obtained without attaching either a sense or an anti-sense promoter sequence to the target sequence of the first-strand cDNA. By way of example, but not of limitation, a splice template oligo or a ligation oligo can be used to attach a primer-binding sequence to the 3'-end of a target sequence comprising first-strand

cDNA. The splice template oligo or ligation oligo does not comprise a complete sense or anti-sense promoter sequence, although the splice template oligo or ligation oligo can (but need not) comprise part of a sequence that is complementary to a sense or anti-sense promoter, respectively. If the splice template oligo or ligation oligo comprises part of a promoter sequence, the first-strand cDNA will have a portion of an anti-sense promoter sequence on its 3'-end, providing an annealing site for a primer comprising a sense promoter sequence that can be used for synthesis of a ssDNA transcription substrate comprising second-strand cDNA. If the splice template oligo or ligation oligo does not comprise part of a promoter sequence, then the respective oligo will have another sequence that provides an annealing site for another primer that has a complete sense promoter of the invention, thereby permitting synthesis of ssDNA transcription substrate comprising second-strand cDNA by primer extension.

[00335] The embodiments of processes for obtaining a transcription substrate for methods of the invention described above are provided only as examples, and are not intended to limit the present invention. The description above and herein will reveal and make evident to those with knowledge in the art numerous other embodiments of methods and processes for obtaining a transcription substrate for use in making a transcription product corresponding to a target nucleic acid sequence in a method of the invention, and the invention includes all of those methods and processes for obtaining a transcription substrate for use in the methods.

[00336] In embodiments in which the target sequences comprise mRNA, whether of a single species of mRNA or all of the mRNA in a particular sample, the transcription products can subsequently be used for a variety of applications. By way of example, but not of limitation, transcription products can be used for *in vitro* or *in vivo* translation, for use as RNAi to silence one or more genes *in vivo*, for spotting on a surface to make expression arrays or microarrays, or for making hybridization probes for arrays or microarrays for gene expression profiling or other uses. In still other embodiments, methods of the invention can be used to make first-strand cDNA from mRNA, which in turn can be used for techniques such as random amplification of cDNA ends (RACE) or to make hybridization probes.

2. Obtaining a ssDNA Transcription Substrate Using a Splice Template Oligo or Promoter Splice Template Oligo

a. Definitions and Methods for Using Splice Template Oligos

[00337] A "splice template" or "splice template oligo" is an oligonucleotide that complexes with a single-stranded target nucleic acid and is used as a template to extend the 3'-

terminus of the target sequence in order to add a specific sequence. The 3'-portion of a splice template is sufficiently complementary to the 3'-terminus of the target sequence that is to be extended to anneal thereto. A DNA- or RNA-dependent DNA polymerase is then used to extend the target nucleic acid molecule using the sequence in the 5'-portion of the splice template oligo as a template. The extension product of the primer-extended molecule has the specific sequence at its 3'-terminus that is complementary to the sequence in the 5'-portion of the splice template oligo.

[00338] A preferred embodiment of a splice template of the present invention is a "promoter splice template" or a "promoter splice template oligo." A promoter splice template oligo comprises a sequence in its 3'-portion that is sufficiently complementary to the 3'-end of the target sequence to anneal thereto and a sequence in its 5'-portion that is complementary to a sequence comprising a single-stranded transcription promoter of the invention. Thus, a promoter splice template oligo can provide a template for synthesis of a sequence comprising a transcription promoter at the 3'-end of first-strand cDNA obtained either by reverse transcriptase primer extension of a target sequence comprising an RNA target nucleic acid, such as, but not limited to, an mRNA target, or by DNA polymerase primer extension of a target sequence comprising DNA. If the promoter splice template is annealed to the end of first-strand cDNA made by primer extension of a target nucleic acid, and the promoter splice template comprises a sequence that is complementary to a functional transcription promoter (i.e., the promoter splice template has an anti-sense promoter sequence), then the first-strand cDNA will have a sense promoter and will therefore comprise a ssDNA transcription substrate of the invention. In one embodiment, which is a preferred embodiment, transcription of a ssDNA transcription substrate results in synthesis of transcription products having a sequence identical to that of the target sequence.

[00339] However, the invention also comprises other embodiments. Thus, in one embodiment, if the 3'-end of a DNA target nucleic acid in a sample has a defined 3'-end and is itself primer-extended on a promoter splice template oligo comprising an anti-sense promoter sequence (i.e., the target sequence is used directly without being copied using a DNA polymerase to obtain first-strand cDNA), then the transcription products made using the ssDNA transcription substrate will be complementary to the target nucleic acid (i.e., the transcription products can comprise anti-sense RNA or "aRNA"). In another embodiment, if the splice template comprises a sense promoter sequence, primer extension of first-strand cDNA annealed to the promoter splice template results in first-strand cDNA having an anti-sense (i.e., non-

functional) promoter. However, if this linear anti-sense-promoter-containing first-strand cDNA is circularized using a ligase under ligation conditions, and replicated by rolling circle replication using a strand displacement primer and a strand-displacing DNA polymerase, the resulting second-strand cDNA will comprise a ssDNA transcription substrate with functional sense promoters. Transcription of this transcription substrate results in synthesis of transcription products that are complementary to the target sequence (i.e., antisense transcription products or aRNA).

[00340] If the 3' terminus of a splice template is not blocked and is complementary to the target nucleic acid, the splice template itself may also act as a primer and be primer-extended by the DNA polymerase using the target nucleic acid molecule as a template. Since an RNA polymerase of the present invention can bind to a promoter in a ssDNA transcription substrate and initiate and synthesize RNA therefrom, in most embodiments of the present invention, the 3'-terminus of a splice template, including a promoter splice template oligo, is blocked in any of a variety of ways, including, but not limited to, using a terminal 2',3'-dideoxynucleotide or a 3'-terminal sequence that is non-complementary to the target, or in other ways well known to those skilled in the art. It is also possible to block the 3'-end using a label, or a small molecule that is a member of a specific binding pair, such as biotin.

[00341] Some embodiments of the invention comprise methods that use a promoter splice template oligo for obtaining a ssDNA transcription substrate for making a transcription product corresponding to a target nucleic acid sequence, wherein the ssDNA transcription substrate comprises a promoter-containing first-strand cDNA and wherein the promoter-containing first-strand cDNA comprises (i) a transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions and (ii) a target nucleic acid sequence, at least a portion of which target sequence either comprises, or, is complementary to, at least a portion of a nucleic acid sequence in a target nucleic acid present in a sample, and wherein the transcription promoter is 3'-of the target sequence in the ssDNA transcription substrate.

[00342] With respect to embodiments that use a promoter splice template oligo, one embodiment of the invention comprises a method for obtaining a ssDNA transcription substrate of the invention for making a transcription product corresponding to a target sequence in a target nucleic acid comprising single-stranded DNA or RNA, the method comprising:

[00343] a. obtaining a primer for synthesis of a first-strand cDNA, the primer comprising a sequence that is complementary to the 3'-end of the target sequence that is to be transcribed;

- [00344]** b. optionally, obtaining a blocking oligo, the blocking oligo comprising a sequence that anneals tightly to a sequence on the target nucleic acid so as to delimit the 3'-end of a primer extension product of the primer using the target nucleic acid as a template, wherein the blocking oligo is not displaced by the primer extension product, and wherein the blocking oligo is not itself capable of being primer-extended by a DNA polymerase;
- [00345]** c. annealing the primer and the blocking oligo, if used, to the target nucleic acid;
- [00346]** d. primer-extending the primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the target sequence to which the primer is annealed;
- [00347]** e. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA;
- [00348]** f. optionally, adding one or more tail sequences to the 3'-end of the first-strand cDNA obtained by primer extension of the primer;
- [00349]** g. obtaining a promoter splice template oligo, wherein the promoter splice template oligo comprises (i) a 3'-end that is sufficiently homologous to the 3'-end of the linear first-strand cDNA to hybridize therewith, and (ii) a 5'-portion that comprises a sequence that is complementary to transcription promoter for an RNA polymerase that can synthesize RNA using a ssDNA transcription substrate;
- [00350]** h. annealing the promoter splice template oligo to the linear first-strand cDNA, including the tail, if present;
- [00351]** i. primer-extending the first-strand cDNA including the tail, if present, with a DNA polymerase under DNA polymerization conditions so as to obtain a promoter-containing first-strand cDNA that has a 3'-portion that is complementary to the portion of the promoter splice template oligo that is not hybridizable to the first-strand cDNA including the tail, if present;
- [00352]** j. removing or dissociating the promoter splice template oligo from the promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate; and
- [00353]** k. obtaining the ssDNA transcription substrate.
- [00354]** As discussed elsewhere herein, a target nucleic acid of the invention can be DNA or RNA of any type. By way of example, but not of limitation, a target sequence can comprise a target nucleic acid comprising a single species of mRNA or a target sequence can comprise a target nucleic acid comprising all of the mRNA in a sample.

[00355] In synthesizing first-strand cDNA from mRNA, when the reverse transcriptase reaches the 5'-end of the mRNA, it can add a non-templated dCMP residue to the 3'-end of full-length first-strand cDNA. The two dCMP residues, including the dG-templated dCMP resulting from an mRNA G[5']ppp[5']G cap structure, provide an annealing site for a splice template oligo or a promoter splice template oligo that comprises a sequence that has at least two G nucleotide residues at or near its 3'-end. Thus, when annealed to the 3'-end of "tailed" first-strand cDNA, the splice template oligo or promoter splice template oligo serves as a template for primer extension of first-strand cDNA. If a promoter splice template oligo is used, a sequence comprising a transcription promoter sequence is added to the 3'-end. The present invention differs from the method of Chenchik *et al.* (U.S. Patent No. 5,962,272, incorporated herein by reference) in a number of ways. By way of example, a template switch oligonucleotide of Chenchik *et al.* has at least one ribonucleotide at its 3'-end portion, and uses the 3'-end of the template switch oligonucleotide to prime synthesis of second-strand cDNA strand. In contrast, since a transcription substrate of the present invention comprises ssDNA, a promoter splice template oligo of the present invention is not used to prime synthesis of second-strand cDNA and the 3'-terminal nucleotide of the promoter splice template oligo is preferably a terminator nucleotide such as, but not limited to, a dideoxynucleotide.

[00356] In some embodiments of methods of the invention for making a transcription substrate using a splice template oligo or promoter splice template oligo, three to four non-templated dCMP residues are added to the 3'-end of the target sequence in first-strand cDNA by using a reverse transcriptase in the presence manganese cations and/or two to four rNMP residues are added to the 3'-end by controlled ribonucleotide tailing using terminal deoxynucleotidyl transferase (Schmidt, W.M. and Mueller, M.W., *Nucleic Acids Res.* 24:1789-91, 1996; *Ibid*, *Nucleic Acids Res.* 27:e31 (i-iv), 1999, both incorporated herein by reference). These additional three to four dCMP residues and/or two to four rNMP residues on the 3'-end of the target sequence in first-strand cDNA provide a site with an even greater degree of homology (compared to Chenchik's method) for annealing to the 3'-end of the splice template oligo or promoter splice template oligo. By way of example, but not of limitation, the splice template oligo or promoter splice template oligo can have two to four nucleotides that are complementary to corresponding rNMP residues at the 3'-end of first-strand cDNA from controlled ribonucleotide tailing and the splice template oligo or promoter splice template oligo can have three to four G nucleotide residues that are complementary to the dCMP residues at or near the 3'-end of first-strand cDNA, providing additional nucleotides for annealing to the splice

template oligo. A promoter splice template oligo annealed to the 3'-end of first-strand cDNA serves as a template for extension of first-strand cDNA by an enzyme with DNA polymerase activity, with concomitant synthesis of a sequence comprising a transcription promoter sequence at its 3'-end. The DNA polymerase used for primer extension of the first-strand cDNA can be a DNA-dependent DNA polymerase or an RNA-dependent DNA polymerase (i.e., reverse transcriptase), depending on whether the composition of the splice template oligo or promoter splice template oligo is DNA or RNA, respectively. Preferably, a DNA polymerase with both DNA-dependent and RNA-dependent DNA polymerase activity is used. By way of example, but not of limitation, preferred DNA polymerases of the invention include MMLV reverse transcriptase (including both RNase H-containing enzyme and mutant forms of the enzyme which lack substantial RNase H activity, depending on the particular application and method of the invention used), AMV reverse transcriptase, and rBst DNA polymerase large fragment (Epicentre Technologies, Madison, WI, USA). Since no synthesis of a second cDNA strand is needed or performed to prepare transcription substrates for the RNA polymerases of the invention, the 3'-terminal nucleotide of the splice template oligo or promoter splice template oligo is preferably a terminator nucleotide such as, but not limited to, a dideoxynucleotide. In some embodiments, the target sequence used comprises full-length first-strand cDNA prepared from mRNA in a sample.

[00357] Thus, one embodiment of the invention that uses a promoter splice template oligo is a method for making a transcription product corresponding to a target sequence comprising a target nucleic acid comprising mRNA in a sample (FIG. 18), the method comprising:

[00358] a. obtaining target nucleic acid comprising mRNA;

[00359] b. obtaining a primer for synthesis of a first-strand cDNA that is complementary to the mRNA, the primer chosen from among: (i) an oligo(dT) primer, and (ii) an oligo(dT) anchor primer; (iii) a primer that is complementary to a specific sequence at the 3'-end of an mRNA, and (iv) a primer in a mixture of primers, the primer comprising a sequence of nucleotides, each of which nucleotides comprises a random nucleotide base that is complementary to any of the four canonical nucleotide bases;

[00360] c. annealing the primer to the target mRNA;

[00361] d. primer-extending (or reverse transcribing) the primer annealed to the target mRNA with a DNA polymerase (reverse transcriptase) under DNA synthesis (reverse transcription) conditions so as to obtain a linear first-strand cDNA that is complementary to the target mRNA to which the primer was annealed;

- [00362]** e. optionally, adding a tail to the first-strand cDNA using one or more methods;
- [00363]** f. optionally, removing the target mRNA that is annealed to the linear first-strand cDNA;
- [00364]** g. obtaining a promoter splice template oligo, wherein the promoter splice template oligo comprises (i) a 3'-end that is sufficiently homologous to the 3'-end of the linear first-strand cDNA, including the tail, if present, to hybridize therewith, and (ii) a 5'-portion that comprises a sequence that is complementary to transcription promoter for an RNA polymerase that can synthesize RNA using a ssDNA transcription substrate;
- [00365]** h. annealing the promoter splice template oligo to the linear first-strand cDNA, including the tail, if present;
- [00366]** i. primer-extending the first-strand cDNA with a DNA polymerase under DNA polymerization conditions so as to obtain a promoter-containing first-strand cDNA that has a 3'-portion that is complementary to the portion of the promoter splice template oligo that is not hybridizable to the first-strand cDNA including the tail, if present;
- [00367]** j. removing or dissociating the promoter splice template oligo from the promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate;
- [00368]** k. obtaining the ssDNA transcription substrate;
- [00369]** l. contacting the ssDNA transcription substrate from step j with an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions so as to obtain transcription products; and
- [00370]** m. obtaining the transcription products;
- [00371]** Thus a general embodiment of the invention comprises a method for using a promoter splice template oligo to make a transcription product corresponding to a target sequence in a target nucleic acid, the method comprising:
- [00372]** a. obtaining a ssDNA transcription substrate by primer extension of the 3'-end of a target sequence using a promoter splice template oligo as a template;
- [00373]** b. contacting the ssDNA transcription substrate with an RNA polymerase that transcribes the ssDNA transcription substrate using the transcription promoter under transcription conditions so as to obtain transcription products; and
- [00374]** c. obtaining the transcription products.
- [00375]** The methods described above for obtaining a ssDNA transcription substrate using a promoter splice template oligo and for making a transcription product corresponding to a

target sequence can be performed in a stepwise manner, or, under suitable reaction conditions, they can be performed continuously in a single reaction mixture.

[00376] Thus, one embodiment of the invention that uses a promoter splice template oligo comprises a method for obtaining additional rounds of transcription of a target sequence comprising a target nucleic acid (FIG. 19), the method comprising:

[00377] a. obtaining transcription products from step c of the method above;

[00378] b. obtaining a primer for synthesis of a first-strand cDNA, the primer comprising a sequence that is complementary to the 3'-end of the transcription products comprising the target sequence that is to be transcribed;

[00379] c. annealing the primer to the transcription products;

[00380] d. primer-extending the primer annealed to the transcription products with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the transcription products to which the primer was annealed;

[00381] e. optionally, removing the transcription products that are annealed to the linear first-strand cDNA;

[00382] f. obtaining a promoter splice template, wherein the promoter splice template comprises a 3'-end that is sufficiently homologous to the 3'-end of the linear first-strand cDNA to hybridize therewith, and a 5'-portion that comprises a sequence that is complementary to a transcription promoter for an RNA polymerase that can synthesize RNA using a ssDNA transcription substrate;

[00383] g. annealing the promoter splice template to the linear first-strand cDNA;

[00384] h. primer-extending the first-strand cDNA with a DNA polymerase under DNA polymerization conditions so as to obtain a promoter-containing first-strand cDNA that has a 3'-portion that is complementary to the portion of the promoter splice template oligo that is not hybridizable to the first-strand cDNA including the tail, if present;

[00385] i. removing or dissociating the promoter splice template oligo from the promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate;

[00386] j. obtaining the ssDNA transcription substrate;

[00387] k. contacting the ssDNA transcription substrate from step j with an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions so as to obtain additional transcription products; and

[00388] l. obtaining the additional transcription products; and

[00389] m. optionally, repeating steps a through m to obtain additional rounds of transcription to obtain transcription products.

[00390] Another embodiment of the invention comprises a method for obtaining a ssDNA transcription substrate of the invention for making a transcription product corresponding to a target nucleic acid sequence; wherein the ssDNA transcription substrate comprises a promoter-containing second-strand cDNA and wherein the promoter-containing second-strand cDNA comprises (i) a transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions and (ii) a target nucleic acid sequence, at least a portion of which target sequence either comprises or is complementary to at least a portion of a nucleic acid sequence in a target nucleic acid present in a sample, and wherein the transcription promoter is 3'-of the target sequence in the ssDNA transcription substrate; the method comprising:

[00391] (a) annealing to a single-stranded target nucleic acid (i) a primer, the primer being complementary to the 3'-end of the single-stranded target nucleic acid sequence, and, optionally, (ii) a blocking oligo that anneals to the target nucleic acid at a sequence that delimits the 5'-end of the target sequence in the target nucleic acid; and

[00392] (b) extending the primer by reverse transcription or primer extension with a DNA polymerase so as to obtain a first-strand cDNA that is complementary to the target nucleic acid sequence; and

[00393] (c) optionally, tailing the first-strand cDNA by adding one or more tail sequences to the 3'-end of the first-strand cDNA obtained by reverse transcription or primer extension using the primer; and

[00394] (d) annealing to the 3'-end of the first-strand cDNA a promoter splice template oligo, wherein the promoter splice template oligo comprises: (i) a 3'-portion that is hybridizable to the 3'-end of the first-strand cDNA including the tail, if present, and (ii) a 5'-portion comprising a sequence that encodes a transcription promoter for an RNA polymerase that can synthesize RNA using a ssDNA transcription substrate; and

[00395] (e) extending the first-strand cDNA including the tail, if present, by reverse transcription or primer extension with a DNA polymerase so as to obtain an anti-sense-promoter-containing first-strand cDNA that has a 3'-portion that is complementary to the portion of the promoter splice template oligo that is not hybridizable to the target nucleic acid sequence including the tail, if present; and

[00396] (f) removing or dissociating the target nucleic acid from the anti-sense-promoter-containing first-strand cDNA; and

[00397] (g) circularizing the promoter-containing first-strand cDNA with a ligase under ligation reaction conditions; and

[00398] (h) annealing to the circular anti-sense-promoter-containing first-strand cDNA a strand-displacement primer, wherein the strand-displacement primer is complementary to a sequence on the anti-sense-promoter-containing first-strand cDNA; and

[00399] (i) incubating the circular anti-sense-promoter-containing first-strand cDNA to which the strand-displacement primer is annealed with a strand-displacing DNA polymerase under strand-displacing DNA polymerization conditions so as to obtain linear promoter-containing second-strand cDNA, wherein the linear promoter-containing second-strand cDNA comprises a ssDNA transcription substrate; and

[00400] (j) obtaining the ssDNA transcription substrate.

[00401] Those embodiments of the invention in which a promoter splice template is used to generate a ssDNA transcription substrate comprising promoter-containing second-strand cDNA will be better understood by reading of the invention specification pertaining to DNA polymerases, particularly, pertaining to strand-displacing DNA polymerases and rolling circle replication. Circularization of anti-sense-promoter-containing first-strand cDNA permits generation of a linear ssDNA comprising sense-promoter-containing second-strand cDNA and positions the transcription promoter 3'-of the target sequence, thereby permitting transcription of target sequence using an RNA polymerase of the present invention. Transcription of a ssDNA transcription substrate comprising second-strand cDNA by an RNA polymerase of the invention results in synthesis of anti-sense RNA with respect to the target nucleic acid sequence, whereas transcription of a ssDNA transcription substrate comprising first-strand cDNA by an RNA polymerase of the invention results in synthesis of sense RNA with respect to the target nucleic

[00402] In still another embodiment of the invention, a splice template oligo is used to add a sense promoter sequence for a T7-type RNAP to the 3'-end of a first-strand cDNA primer extension product comprising a target sequence. This embodiment is similar in most respects to embodiments for using a promoter splice template oligo to add a single-stranded sense promoter for an N4 mini-vRNAP to the 3'-end of a first-strand cDNA comprising a target sequence. However, since T7-type RNAPs, including, without limitation, T7, T3 and SP6 RNAP, generally use double-stranded promoter sequences for transcription, in this embodiment an operable promoter comprises both a sense promoter sequence and an anti-sense promoter

sequence, which sequences are annealed to form a complex. As defined herein, a "sense promoter sequence" for a double-stranded promoter means the promoter sequence of an operable double-stranded promoter that is joined to the 3'-end of the template strand that is transcribed, and an "anti-sense promoter sequence" is a sequence that is complementary to the sense promoter sequence. Promoter sequences for many RNA polymerases are well known in the art and suitable sense and anti-sense promoter sequences can be identified. By way of example, but not of limitation, suitable sense and anti-sense promoter sequences for T7 RNAP can be found in writings by JF Milligan et al. (Nucleic Acids Res., 15: 8783-8798, 1987) and by FW Studier et al. (In: Methods in Enzymology, Vol. 185, pp. 60-89, 1990, edited by DV Goeddel, Academic Press, Inc., San Diego, CA), both incorporated herein by reference.

[00403] Thus, this embodiment of the invention comprises a method for obtaining a substrate for transcription by a T7-type RNAP that uses a double-stranded promoter, the method comprising: (a) obtaining a single-stranded target nucleic acid comprising a target sequence; (b) obtaining a primer that anneals to the 3'-end of the target sequence; (c) annealing the primer to the target sequence under annealing conditions; (d) synthesizing first-strand cDNA by primer extension of the primer annealed to the 3'-end of the target sequence using a DNA polymerase or reverse transcriptase under polymerization conditions; (e) optionally, tailing the first-strand cDNA; (f) obtaining a splice template oligo, the splice template oligo comprising an anti-sense sequence of a double promoter and wherein the 3'-end portion of the splice template oligo anneals to the 3'-end of the first-strand cDNA including the tail sequence, if present; (g) annealing the splice template oligo first-strand cDNA; (h) primer extending the 3'-end of the first-strand cDNA using the annealed splice template oligo as a template using a DNA polymerase or reverse transcriptase under polymerization conditions; (i) obtaining a ssDNA "pro-transcription substrate," the pro-transcription substrate comprising the primer-extended first-strand cDNA having at its 3'-end a sense strand promoter sequence; (j) annealing to the pro-transcription substrate an anti-sense promoter oligo, the anti-sense promoter oligo comprising an the anti-sense promoter sequence that is complementary to the sense strand promoter sequence of the pro-transcription substrate; (k) obtaining a "transcription substrate complex," the transcription substrate complex comprising the complex between the pro-transcription substrate and the anti-sense promoter oligo. The transcription substrate complex can be used to make a transcription product using a T7-type RNAP that binds to the double-stranded promoter in the transcription substrate complex and that uses the single-stranded template attached thereto as a template for transcription under transcription conditions. In another embodiment, the ssDNA

pro-transcription substrate is circularized using a ligase under ligation conditions prior to annealing an anti-sense promoter oligo, thereby obtaining a "circular transcription substrate complex." Transcription of a circular transcription substrate complex using a T7-type RNAP under transcription conditions comprises "rolling circle transcription," which, unless a transcription termination sequence is present in the circular transcription substrate, makes concatameric transcription products. The methods of the present invention differ from the methods disclosed in PCT Patent Application No. WO 02/065093 which synthesize double-stranded templates for use as transcription substrates.

b. Composition of Promoter Splice Template Oligos and Splice Template Oligos

[00404] In most embodiments of the present invention, a splice template oligo or a promoter splice template oligo comprises DNA nucleotides, but a promoter splice template oligo or a splice template oligo can comprise DNA and/or RNA nucleotides and/or modified DNA and/or RNA nucleotides. In some embodiments, a promoter splice template oligo or a splice template oligo can comprise modified RNA having 2'-F-dCMP and 2'-F-dUMP nucleotides in place of CMP and UMP (e.g., made using Epicentre's DuraScribe™ T7 Transcription Kit or N4 mini-vRNAP Y678F enzyme, as described elsewhere herein). The presence of 2'-fluoropyrimidines in the oligo makes it resistant to RNaseA-type ribonucleases. As discussed above, since no synthesis of a second cDNA strand is needed or performed to prepare transcription substrates for the RNA polymerases of the invention, the 3'-terminal nucleotide of the promoter splice template oligo or a splice template oligo is preferably a terminator nucleotide such as, but not limited to, a dideoxynucleotide. In some embodiments, 3 or 4 of the internucleotide linkages at the 5'-end of a promoter splice template oligo or a splice template oligo can comprise phosphorothioate linkages or other modified linkages, as discussed elsewhere herein, in order to protect the oligos from degradation by some exonucleases that can be present in a sample (e.g., Nikiforov *et al.*, PCR Methods and Applications 3:285-291, 1994, incorporated herein by reference).

[00405] In some embodiments, such as in those embodiments in which the ssDNA transcription substrate comprises a second-strand cDNA that is generated by strand displacement during rolling circle replication using anti-sense-promoter-containing first-strand cDNA as a template, a promoter splice template oligo can also comprise a sequence that encodes a primer-binding site for a strand-displacement primer, which primer-binding site is generally 5'-of the sense promoter sequence in the promoter splice template oligo, thus resulting in a primer

binding site 3'-of the anti-sense transcription promoter sequence in anti-sense-promoter-containing first-strand cDNA. However, a sequence in the splice template oligo that encodes a primer-binding-site is not required. Alternatively, one or more rolling circle strand displacement primers that prime at one or more other sites on the anti-sense-promoter-containing first-strand cDNA rolling circle replication template can be used. In some embodiments, a sequence that encodes a primer-binding site in the promoter splice template oligo is not desirable because this sequence will be replicated in the rolling circle replication product in addition to the anti-sense sequence of the target nucleic acid sequence.

[00406] In general, a splice template oligo that does not comprise a sense or anti-sense promoter sequence is used in conjunction with alternative methods or processes for incorporating a transcription promoter into a ssDNA transcription template, as discussed elsewhere herein. By way of example, but not of limitation, a splice template oligo that does not comprise a sense or anti-sense promoter sequence can be used to add a sequence for a primer-binding site or to add other sequences or genetic elements for other applications in conjunction with use of a promoter primer to incorporate a transcription promoter into a ssDNA transcription template.

3. Obtaining ssDNA Transcription Substrates by Ligation of a Promoter Ligation Oligo to a Target Sequence

a. Definitions and Methods for Using Promoter Ligation Oligos

[00407] A "promoter ligation oligo" is a ssDNA comprising all or substantially all of a sequence for a single-stranded transcription promoter of the invention that is joined to a ssDNA polynucleotide comprising a target sequence in order to make a transcription substrate of the invention. A promoter ligation oligo comprising a sense transcription promoter can be phosphorylated at its 5'-end or it can be phosphorylated during the processes of the methods of the invention using a polynucleotide kinase (PNK), such as, but not limited to T4 PNK, and ATP under suitable reaction conditions. In other embodiments, the promoter ligation oligo can have a bound topoisomerase moiety, or it can be ligated using a topoisomerase during the processes of the methods of the invention.

[00408] In some embodiments, the 5'-phosphorylated promoter ligation oligo comprising a sense promoter is ligated to the 3'-hydroxyl end of a ssDNA polynucleotide comprising a target sequence. In other embodiments, the 3'-hydroxyl end of a promoter ligation oligo comprising a sense promoter sequence is ligated to the 5'-end of a ssDNA polynucleotide

comprising a target sequence, and then the 5'-end of the promoter ligation oligo is phosphorylated and ligated to the 3'-hydroxyl end of the target sequence to make a circular ssDNA transcription substrate. In still other embodiments, a DNA oligo that encodes all or substantially all a sequence that is complementary to a transcription promoter (i.e., it comprises an anti-sense promoter) is ligated to either the 3'-end or the 5'-end of a target sequence, with the nucleic acid having a 5'-end that is ligated being phosphorylated. In some embodiments in which a promoter ligation oligo comprising an anti-sense promoter is used, the molecule to which the promoter ligation oligo is ligated to make a circular ssDNA template for rolling circle replication, which in turn, makes a single-stranded second-strand DNA that comprises a ssDNA transcription substrate for an RNA polymerase of the invention. In still other embodiments, a linear ssDNA comprising an anti-sense promoter 3'-of the target sequence can be used to make a linear ssDNA transcription substrate by strand displacement amplification.

[00409] A "ligation splint" or a "ligation splint oligo" is an oligo that is used to provide an annealing site or a "ligation template" for joining two ends of one nucleic acid (i.e., "intramolecular joining") or two ends of two nucleic acids (i.e., "intermolecular joining") using a ligase or another enzyme with ligase activity. The ligation splint holds the ends adjacent to each other and "creates a ligation junction" between the 5'-phosphorylated and a 3'-hydroxylated ends that are to be ligated. For example, when a ligation splint oligo is used to join a sense promoter ligation oligo to the 3'-end of a first-strand cDNA comprising a target sequence, the ligation splint oligo has a sequence complementary to the 3'-end of the target sequence, including a "tailed" target sequence, if any, and a second adjacent sequence that is complementary to the 5'-end of a the 5'-phosphorylated promoter ligation oligo. Ligases that can be used to ligate suitable ends that are annealed to a ligation splint comprising DNA include, but are not limited to, Ampligase® DNA Ligase (EPICENTRE Technologies, Madison, WI), Tth DNA ligase, Tfl DNA ligase, Tsc DNA ligase (Prokaria, Ltd., Reykjavik, Iceland), or T4 DNA ligase. These ligases can be used for both intermolecular and intramolecular ligations when a ligation splint comprising DNA is used to bring the respect ends adjacent. If a ligation splint comprising RNA is used, T4 DNA ligase can be used to join the ends that are annealed to the ligation splint. In some embodiments, a ligase that catalyzes non-homologous intramolecular ligation of a 5'-phosphorylated end with a 3'-hydroxyl end can be used in methods of the invention for circularization of single-stranded DNA without a ligation splint. By way of example, but not of limitation, a ligation splint is not required for ligation of a ssDNA using ThermoPhage™ RNA Ligase II (Prokaria, Ltd., Reykjavik, Iceland).

[00410] Some embodiments of the invention comprise methods that use a promoter ligation oligo for obtaining a ssDNA transcription substrate for making a transcription product corresponding to a target nucleic acid sequence, wherein the ssDNA transcription substrate comprises a promoter-containing first-strand cDNA and wherein the promoter-containing first-strand cDNA comprises a transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions and a target nucleic acid sequence, at least a portion of which target sequence either comprises, or, is complementary to, at least a portion of a nucleic acid sequence in a target nucleic acid present in a sample, and wherein the transcription promoter is 3'-of the target sequence in the ssDNA transcription substrate.

[00411] With respect to embodiments that use a promoter ligation oligo, one embodiment of the invention comprises a method for obtaining a ssDNA transcription substrate of the invention for making a transcription product corresponding to a target sequence in a target nucleic acid comprising single-stranded DNA or RNA, the method comprising:

[00412] a. obtaining a primer for synthesis of a first-strand cDNA, the primer comprising a sequence that is complementary to the 3'-end of the target sequence that is to be transcribed;

[00413] b. optionally, obtaining a blocking oligo, the blocking oligo comprising a sequence that anneals tightly to a sequence on the target nucleic acid so as to delimit the 3'-end of a primer extension product of the primer using the target nucleic acid as a template, wherein the blocking oligo is not displaced by the primer extension product, and wherein the blocking oligo is not itself capable of being primer-extended by a DNA polymerase;

[00414] c. annealing the primer and the blocking oligo, if used, to the target nucleic acid;

[00415] d. primer-extending the primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the target sequence to which the primer is annealed;

[00416] e. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA;

[00417] f. optionally, adding one or more tail sequences to the 3'-end of the first-strand cDNA obtained by primer extension of the primer;

[00418] g. obtaining a promoter ligation oligo, wherein the promoter ligation oligo comprises a sense transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions, wherein the promoter

ligation oligo is either phosphorylated at its 5'-terminus, or is subsequently phosphorylated using a polynucleotide kinase and ATP;

[00419] h. obtaining a ligation splint oligo, wherein the ligation splint oligo has a sequence that is complementary to the 5'-end of the promoter ligation oligo and the 3'-end of the first-strand cDNA, wherein the 5'-end of the promoter ligation oligo is adjacent to the 3'-end of the first-strand cDNA when annealed to the ligation splint oligo;

[00420] i. incubating under annealing conditions: (i) the first-strand cDNA; (ii) the promoter ligation oligo; and (iii) the ligation splint oligo;

[00421] j. ligating using a ligase or a topoisomerase under ligation conditions, the 5'-end of the promoter ligation oligo to the 3'-end of the first-strand cDNA, which ends are annealed to the ligation splint oligo, so as to obtain a promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate;

[00422] k. optionally, removing or dissociating the ligation splint oligo; and

[00423] l. obtaining the ssDNA transcription substrate.

[00424] For methods of the invention that use a promoter ligation oligo, a target nucleic acid can be DNA or RNA. By way of example, but not of limitation, a target sequence can comprise a target nucleic acid comprising a single species of mRNA or it can comprise all of the mRNA in a sample. If the target nucleic acid comprises mRNA, an initial step of the method is reverse transcription of mRNA to obtain a target sequence comprising first-strand cDNA. In these embodiments, the transcription promoter is added to the 3'-end of a target sequence comprising first-strand cDNA by ligation using a ligase under ligation conditions. By way of example, but not of limitation, a ssDNA transcription substrate of the invention can be prepared by ligating a sense-strand promoter ligation oligo, which has been phosphorylated at its 5'-end using T4 polynucleotide kinase or another suitable polynucleotide kinase and ATP under suitable reaction conditions, to the 3'-end of a target sequence comprising first-strand cDNA that has additional nucleotides which have been added (e.g., see U.S. Patent No. 5,962,272 and Schmidt and Mueller, Nucleic Acids Res. 27:e31 [i-iv], 1999, both incorporated herein by reference). The tail at the 3'-end of the first-strand cDNA is not required, but in those embodiments in which it is used, a tail provides a region of homology for a ligation splint oligo. In some embodiments of this aspect of the invention, a ligation splint oligo, which has a 5'-portion that is complementary to the 5'-end of a phosphorylated sense-strand promoter oligo and a second 3'-portion that is complementary to the 3'-end of a target sequence, such as, but not limited to, "tailed" target sequence, is used to provide an annealing site that will keep the two

ends adjacent to one another for ligation. In most embodiments of this method, the promoter ligation oligo comprises ssDNA or modified ssDNA, wherein the modified nucleotides, if present, do not affect the ability of the transcription promoter to function fully as a promoter for *in vitro* transcription. In some embodiments, the 3'-terminal nucleotide of a ligation splint is a dideoxynucleotide or another termination nucleotide, so that the ligation splint cannot serve as a primer for polymerases in the reaction.

[00425] Thus, one embodiment of the invention for using a promoter ligation oligo is a method for making a transcription product corresponding to a target sequence comprising a target nucleic acid comprising mRNA (FIG. 20), the method comprising

[00426] a. obtaining a target nucleic acid comprising mRNA;

[00427] b. obtaining a primer for synthesis of a first-strand cDNA that is complementary to the mRNA, the primer chosen from among: (i) an oligo(dT) primer, and (ii) an oligo(dT) anchor primer; (iii) a primer that is complementary to a specific sequence at the 3'-end of an mRNA, and (iv) a primer in a mixture of primers, the primer comprising a sequence of nucleotides, each of which nucleotides comprises a random nucleotide base that is complementary to any of the four canonical nucleotide bases;

[00428] c. annealing the primer to the target mRNA;

[00429] d. primer-extending (or reverse transcribing) the primer annealed to the target mRNA with a DNA polymerase (reverse transcriptase) under DNA synthesis (reverse transcription) conditions so as to obtain a linear first-strand cDNA that is complementary to the target mRNA to which the primer was annealed;

[00430] e. optionally, adding a tail to the first-strand cDNA using one or more methods;

[00431] f. optionally, removing the target mRNA that is annealed to the linear first-strand cDNA;

[00432] g. obtaining a promoter ligation oligo, wherein the promoter ligation oligo comprises a sense transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions, wherein the promoter ligation oligo is either phosphorylated at its 5'-terminus, or is subsequently phosphorylated using a polynucleotide kinase and ATP;

[00433] h. obtaining a ligation splint oligo, wherein the ligation splint oligo has a sequence that is complementary to the 5'-end of the promoter ligation oligo and the 3'-end of the first-strand cDNA, wherein the 5'-end of the promoter ligation oligo is adjacent to the 3'-end of the first-strand cDNA when annealed to the ligation splint oligo;

[00434] i. incubating under annealing conditions the first-strand cDNA, the promoter ligation oligo, and the ligation splint oligo;

[00435] j. ligating using a ligase or a topoisomerase moiety under ligation conditions, the 5'-end of the promoter ligation oligo to the 3'-end of the first-strand cDNA, which ends are annealed to the ligation splint oligo, so as to obtain a promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate;

[00436] k. optionally, removing or dissociating the ligation splint oligo;

[00437] l. obtaining the ssDNA transcription substrate.

[00438] Thus, a general embodiment of the invention comprises a method for using a promoter ligation oligo for making a transcription product corresponding to a target sequence in a target nucleic acid, the method comprising:

[00439] a. obtaining a ssDNA transcription substrate by carrying out steps a through l of the method above;

[00440] b. contacting the ssDNA transcription substrate with an RNA polymerase that transcribes the ssDNA transcription substrate using the transcription promoter under transcription conditions so as to obtain transcription products; and

[00441] c. obtaining the transcription products.

[00442] The methods described above for obtaining a ssDNA transcription substrate using a promoter ligation oligo and for making a transcription product corresponding to a target sequence can be performed in a stepwise manner, or, under suitable reaction conditions, they can be performed continuously in a single reaction mixture.

[00443] Thus, one embodiment of the invention that uses a promoter ligation oligo comprises a method for obtaining additional rounds of transcription of a target sequence in a target nucleic acid (FIG. 21), the method comprising:

[00444] a. obtaining the transcription products from step c of the method above;

[00445] b. obtaining a primer for synthesis of a first-strand cDNA, the primer comprising a sequence that is complementary to the 3'-end of the transcription products comprising the target sequence that is to be transcribed;

[00446] c. annealing the primer to the target nucleic acid;

[00447] d. primer-extending the primer annealed to the transcription products with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the transcription products to which the primer was annealed;

- [00448]** e. optionally, removing the transcription products that are annealed to the linear first-strand cDNA;
- [00449]** f. optionally, adding one or more tail sequences to the 3'-end of the first-strand cDNA obtained by primer extension of the primer;
- [00450]** g. obtaining a promoter ligation oligo, wherein the promoter ligation oligo comprises a sense transcription promoter for an RNA polymerase that synthesizes RNA from a ssDNA promoter-containing template under transcription conditions, wherein the promoter ligation oligo is either phosphorylated at its 5'-terminus, or is subsequently phosphorylated using a polynucleotide kinase and ATP;
- [00451]** h. obtaining a ligation splint oligo, wherein the ligation splint oligo has a sequence that is complementary to the 5'-end of the promoter ligation oligo and the 3'-end of the first-strand cDNA, wherein the 5'-end of the promoter ligation oligo is adjacent to the 3'-end of the first-strand cDNA when annealed to the ligation splint oligo;
- [00452]** i. incubating under annealing conditions the first-strand cDNA, the promoter ligation oligo and the ligation splint oligo;
- [00453]** j. ligating using a ligase or a topoisomerase moiety under ligation conditions, the 5'-end of the promoter ligation oligo to the 3'-end of the first-strand cDNA, which ends are annealed to the ligation splint oligo, so as to obtain a promoter-containing first-strand cDNA, the promoter-containing first-strand cDNA comprising a ssDNA transcription substrate;
- [00454]** k. optionally, removing or dissociating the ligation splint oligo;
- [00455]** l. obtaining the ssDNA transcription substrate;
- [00456]** m. contacting the ssDNA transcription substrate from step l with an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions so as to obtain transcription products; and
- [00457]** n. obtaining the additional transcription products; and
- [00458]** o. optionally, repeating steps a through o to obtain additional rounds of transcription to obtain transcription products.
- [00459]** In other embodiments for obtaining a ssDNA transcription substrate of the invention, a promoter ligation oligo comprising a sense transcription promoter is ligated to a 5'-phosphorylated end of a target nucleic acid sequence comprising DNA or to a first-strand cDNA prepared by reverse transcription or primer extension of a target sequence in a target nucleic acid. In these embodiments of the invention, the resulting first-strand cDNA having a sense transcription promoter at its 5'-end is then circularized by ligation with a ligase under ligation

reaction conditions in order to bring the promoter in proper juxtaposition with the 3'-end of the target sequence for transcription of the target sequence by an RNA polymerase of the invention, thereby obtaining a circular ssDNA transcription substrate of the invention. In this embodiment, the ligation splint oligo comprises a 5'-portion and a 3'-portion, wherein the 5'-portion is complementary to the 5'-end of the DNA or first-strand cDNA target sequence and the 3'-portion is complementary to the 3'-end of the promoter ligation oligo. In yet another embodiment, a circular ssDNA transcription substrate can be linearized 3'-of the transcription promoter using methods discussed elsewhere herein in order to obtain a linear ssDNA transcription substrate of the invention.

[00460] In other embodiments of the invention for obtaining a ssDNA transcription substrate, a promoter ligation oligo that comprises an anti-sense transcription promoter is ligated using a ligase under ligation reaction conditions to either the 3'-end or the 5'-end of a ssDNA target sequence or a first-strand cDNA made by primed reverse transcription or primer extension of a target sequence in a target nucleic acid, wherein either the promoter ligation oligo or the first-strand cDNA target sequence, respectively, is 5'-phosphorylated. By way of example, but not of limitation, in one embodiment, a 5'-phosphorylated promoter ligation oligo, which comprises an anti-sense transcription promoter, is ligated in a first ligation reaction using a ligation splint oligo, to the 3'-end of the first-strand cDNA target sequence, generating a linear first-strand cDNA with an anti-sense promoter (i.e., a non-functional complement of a functional promoter). The anti-sense-promoter-containing first-strand cDNA is then phosphorylated at its 5'-end and ligated using a ligase in a second ligation reaction to generate a circular anti-sense-promoter-containing first-strand cDNA molecule. A similar circular DNA molecule can be made following ligation of an anti-sense-promoter ligation oligo to the 5'-end of first-strand cDNA, followed by 5'-phosphorylation, and a second ligation reaction, thereby generating the similar circular first-strand cDNA with an anti-sense transcription promoter. These circular molecules can be used as templates for rolling circle replication using a strand-displacing DNA polymerase of the invention and one or more strand displacement primers that is/are complementary to one or more sequences in the circular molecules. Strand displacement replication of these circular anti-sense-promoter-containing first-strand cDNA molecules using one or more strand displacement primers generates linear second-strand cDNA that is released by strand displacement into the reaction mixture as ssDNA that has a sense (i.e., functional) transcription promoter for an RNA polymerase of the invention. Thus, these linear single-stranded promoter-containing second-strand cDNA molecules comprise ssDNA transcription

substrates of the invention. Transcription of these ssDNA transcription substrates by an RNA polymerase of the invention results in synthesis of anti-sense transcription products corresponding to the target sequence. In some embodiments of this method for obtaining a ssDNA transcription substrate of the invention, a ligation splint oligo is used in the second ligation reaction to ligate linear anti-sense-promoter-containing first-strand cDNA to generate circular anti-sense-promoter-containing first-strand cDNA, whereas in other embodiments, the linear single-stranded cDNA is ligated in the second ligation reaction without a ligation splint oligo using a ligase, such as, but not limited to, ThermoPhage™ RNA Ligase II (Prokaria, Reykjavik, Iceland). In still other embodiments ligation splint oligos are used in both the first and second ligation reactions, and both ligation reactions are carried out simultaneously to form circular first-strand cDNA in one step. Primer extension with a second-strand cDNA synthesis primer using the circular first-strand cDNA as a template for a strand displacing DNA polymerase of the invention generates linear ssDNA having tandem repeats of sense transcription promoters and the target sequence, which can be used as a transcription substrate of the invention. In other embodiments, circular first-strand cDNA can be linearized to form a linear first-strand cDNA that can be replicated by strand displacement replication methods in order to generate a linear ssDNA second-strand cDNA transcription substrate of the invention. In some embodiments of this aspect of the method, an anti-sense promoter ligation oligo also has, in addition to an anti-sense transcription promoter, at least one additional sequence that comprises a primer binding site that is complementary to a strand displacement primer in order to provide a priming site outside of the target sequence for making second-strand cDNA. This primer-binding site can be used for primed rolling circle replication of the circular transcription substrate or, following linearization of the circular first-strand cDNA strand displacement template as described elsewhere herein, the primer binding site can be used for strand displacement replication of the linear ssDNA transcription substrate. However, in some embodiments it is not desirable to add another sequence (i.e., the primer-binding site) that is replicated along with the target sequence.

[00461] In some embodiments, remaining linear nucleic acids, such as, but not limited to, ligation splint oligos, are removed during the reaction using the gene 6 exonuclease of phage T7. This exonuclease digests DNA starting from the 5'-end of a double-stranded structure. It has been used successfully for the generation of single-stranded DNA after PCR amplification (Holloway *et al.*, Nucleic Acids Res. 21:3905-3906, 1993; Nikiforov *et al.*, PCR Methods and Applications 3:285-291, 1994, incorporated herein by reference). The gene 6 exonuclease of

phage T7 can be added after ligation, together with the rolling circle DNA polymerase to remove unligated oligos. To protect the ssDNA transcription substrate from degradation, a strand displacement primer for rolling circle replication can contain 3 or 4 phosphorothioate linkages at the 5'-end, to make this molecule resistant to the exonuclease (Nikiforov *et al.*, PCR Methods and Applications 3:285-291, 1994). The exonuclease degrades unprotected linear molecules as they become associated with the rolling circle DNA product. Based on this description, those with knowledge in the art will understand and know other embodiments of the invention in which this process of the invention for removing single-stranded DNA oligos can be used to advantage, and the invention comprises all such embodiments.

[00462] From the above description, those with knowledge in the art will understand other embodiments of methods and processes of the invention for making linear or circular ssDNA transcription substrates, all of which are included herein.

b. Composition of Promoter Ligation Oligos and Ligation Splint Oligos

[00463] In most embodiments, a promoter ligation oligo of the invention comprises DNA or modified DNA nucleotides. In addition to the sequence of the transcription promoter, the promoter ligation oligo can also comprise additional nucleic acid sequences that are 3'-of or 5'-of the transcription promoter, which may be included for a particular purpose. By way of example, but not of limitation, a promoter ligation oligo can have an additional sequence that is 3'-of or 5'-of the transcription promoter that serves as a priming site for synthesis of a second-strand cDNA. In some embodiments, the promoter ligation oligo can contain phosphorothioate linkages between the nucleotides at its 5'-end (e.g., between 3 or 4 of the nucleotides) in order to protect it from degradation by some exonucleases. The 5'-terminus of the promoter ligation oligo either has a 5'-phosphoryl group or it is phosphorylated using a polynucleotide kinase, such as, but not limited to, T4 PNK and ATP during the processes of a method of the invention. Promoter ligation oligos can be synthesized on an oligo synthesizer, which is usually preferred, or enzymatically, using methods discussed elsewhere herein.

[00464] A "ligation splint" or a "ligation splint oligo" is an oligonucleotide that is used to provide an annealing site or a "ligation template" on which a 5'-phosphorylated end and a 3'-hydroxyl end of one or two different nucleic acids, such as, but not limited to, a promoter ligation oligo and a first-strand cDNA target nucleic acid sequence, can hybridize so as to bring the two ends adjacent to one another in a ligation reaction for joining by a ligase (or a topoisomerase or other enzyme with ligase activity). A ligation splint comprises a sufficient

number and composition of nucleotides, and is present in sufficient concentration, so that the complementary sequences of both the 5'- and 3'-ends remain annealed to the ligation splint under ligation conditions so to permit ligation of the ends to occur. Thus, in most embodiments, the ligation splint comprises at least about 4 nucleotides up to about 20 nucleotides, but the invention is not limited to a specific number of nucleotides. An appropriate sequence (and T_m), size, and concentration for a ligation splint can be determined empirically by those with knowledge in the art. In most embodiments of the present invention, it is preferable that the 3'-terminal nucleotide of a ligation splint is a dideoxynucleotide or another termination nucleotide, so that the ligation splint oligo cannot serve as a primer for polymerases in the reaction. In most embodiments in which there are multiple rounds of transcription of a target nucleic acid sequence, the other nucleotides in a ligation splint oligo comprise deoxynucleotides. However, in some embodiments, the other nucleotides can comprise ribonucleotides and/or purine ribonucleotides and 2'-fluoro-pyrimidine nucleotides, which confer resistance to RNase A-type nucleases. The composition of a ligation splint oligo will also depend on the ligase and ligation conditions used. By way of example, Ampligase® Thermostable Ligase (Epicentre Technologies, Madison, WI, USA) will only ligate the 5'-phosphoryl and 3'-hydroxyl ends of DNA that is annealed to a DNA ligation splint and will not ligate DNA ends annealed to RNA. However, T4 RNA ligase has been reported to efficiently ligate DNA ends that are annealed to an RNA ligation splint (Faruqi *et al.*, U.S. Patent No. 6,368,801, incorporated herein by reference). Ligation splint oligos can be synthesized on an oligo synthesizer, which is usually preferred, or enzymatically, using methods discussed elsewhere herein.

c. Ligases and Ligation Methods for Attaching a Promoter Ligation Oligo to a Target Sequence

[00465] The invention is not limited to a specific ligase. However, when a promoter ligation oligo is used, preferably the ligase is not active in ligating blunt ends and is highly selective for ligation of a deoxynucleotide having a 5'-phosphate and a deoxynucleotide having 3'-hydroxyl group when these respective 5'- and 3'- nucleotides are adjacent to each other when annealed to a ligation splint. Ampligase® Thermostable DNA Ligase (Epicentre Technologies, Madison, Wisconsin, USA), Tth DNA ligase, Tfl DNA ligase, and Tsc DNA Ligase (Prokaria Ltd., Reykjavik, Iceland) are NAD-dependent DNA ligases that are not active on blunt ends and that ligate the 5'-phosphate and 3'-hydroxyl termini of DNA ends that are adjacent to one another when annealed to a complementary DNA molecule, and are suitable ligases in embodiments of the invention that use a ligation splint oligo comprising DNA. However, the

invention is not limited to the use of a particular ligase and any suitable ligase can be used. For example, T4 DNA ligase can be used in embodiments of the invention that use a ligation splint. Still further, Faruqi discloses in U.S. Patent No. 6,368,801 that T4 RNA ligase can efficiently ligate DNA ends of nucleic acids that are adjacent to each other when hybridized to an RNA strand. Thus, T4 RNA ligase is a suitable ligase of the invention in embodiments in which DNA ends are ligated on a ligation splint oligo comprising RNA or modified RNA, such as, but not limited to modified RNA that contains 2'-F-dCTP and 2'-F-dUTP made using the DuraScribe™ T7 Transcription Kit (Epicentre Technologies, Madison, WI, USA) or the N4 mini-vRNAP Y678F mutant enzyme described herein. If a promoter ligation oligo is initially ligated to the 5'-end of a target sequence, the promoter sequence can be operably joined to the 3'-end of the target sequence by first phosphorylating the 5'-end using a polynucleotide kinase, such as, but not limited to, T4 polynucleotide kinase, and then ligating the linear ssDNA to form a circular ssDNA using a ligase that catalyzes non-homologous intramolecular ligation, such as, but not limited to ThermoPhage™ RNA Ligase II (Prokaria, Ltd., Reykjavik, Iceland). The invention is also not limited to the use of a ligase for covalently joining the 5'-end to the 3'-end of the same or different nucleic acid molecules in the various embodiments of the invention. By way of example, other ligation methods such as, but not limited to, topoisomerase-mediated ligation (e.g., U.S. Patent No. 5,766,891, incorporated herein by reference) can be used.

4. Obtaining ssDNA Transcription Substrates Using a Promoter Primer

a. Methods for Using Promoter Primers

[00466] A "promoter primer" is a primer, generally with a free 3'-OH group, that comprises a sequence that is complementary to a target sequence at its 3'-end and which encodes a transcription promoter in its 5'-portion. A promoter primer can have a sequence at its 3'-end that is complementary to a specific known sequence in a target nucleic acid, in which case it is referred to as a "specific-sequence promoter primer." However, other embodiments of promoter primers can also be used in methods of the invention. An "oligo(dT) promoter primer" has an oligo(dT) sequence at its 3'-end, and is used mainly in embodiments of the invention which pertain to mRNA molecules having polyadenylated [i.e., poly(A) tails], although an oligo(dT) promoter primer can also be used in embodiments in which another target nucleic acid is tailed with poly(A) or poly(dA). An "anchored oligo d(T) promoter primer, in addition to having an oligo(dT) sequence in its 3'-portion, also has one (or a small number) of nucleotides 3'-of the oligo(dT) sequence, called "anchor nucleotides," which anneal to the 3'-portion of the

mRNA target sequence just prior to the poly(A) sequence. Thus, the anchor nucleotides serve to "anchor" the mRNA-complementary portion of the anchored oligo(dT) promoter primer to the beginning of the protein-coding sequence of the mRNA target sequence. The anchor nucleotides can comprise either a specific base for a specific mRNA or a randomized nucleotide (i.e., synthesized with a mixture of all four nucleotides) for priming all mRNA molecules in a sample. A "random-sequence promoter primer" has a random sequence, such as, but not limited to a random hexamer sequence, at its 3'-end. In most cases, a random-sequence promoter primer comprises a mixture of primers with all possible sequences (e.g., all possible hexamers) in its target sequence-complementary portion. Random-sequence promoter primers can be made by including all four canonical nucleotide reagents during the chemical synthesis of each of the nucleotide positions of the random sequence (e.g., the hexamer sequence) of the target sequence-complementary portion of the primer. A random-sequence promoter primer can be used in those embodiments of the invention in which it is desired to amplify all target nucleic acid sequences in a sample, or to amplify all target nucleic acid sequences in a random manner, such as for making a library of all target nucleic acid sequences. However, although all sequences may be amplified, use of a random-sequence promoter primer does not necessarily generate only full-length copies of target nucleic acids (e.g., full-length cDNA copies of mRNA molecules from a cell). Thus, embodiments of the invention which use random-sequence promoter primers are usually used when full-length copies of a target nucleic acid sequence are not required, such as, for obtaining hybridization probes for some applications.

[00467] In embodiments of the invention in which a transcription substrate of the invention comprises first-strand cDNA obtained by reverse transcription or primer extension of a promoter primer using a target nucleic acid as a template, the transcription promoter in the promoter primer comprises a sense promoter sequence that is located in the 5'-portion of the promoter primer. Thus, the transcription promoter in the linear first-strand cDNA obtained by reverse transcriptase- or DNA polymerase-catalyzed extension of the promoter primer using the target nucleic acid as a template is not operable as a promoter for transcription of the target sequence since the promoter is not operably joined to the 3'-end of the target sequence. A method of the present invention solves this problem by operably joining the single-stranded sense transcription promoter in the 5'-portion of the linear first-strand cDNA to the 3'-end of the target sequence using a ligase or another joining means, thus forming a circular ssDNA transcription substrate for an RNA polymerase that can bind the single-stranded promoter and transcribe the target sequence joined thereto.

[00468] Thus, in one embodiment of the invention, a promoter primer having a sequence complementary to a target sequence at its 3'-end and a transcription promoter in its 5'-portion is used to obtain a ssDNA transcription substrate of the invention (FIG. 22). After annealing to a target nucleic acid, the promoter primer is used to prime first-strand cDNA synthesis using a DNA polymerase or reverse transcriptase under suitable reaction conditions in order to obtain linear first-strand cDNA.

[00469] The linear promoter-containing first-strand cDNA is then ligated using a ligase under suitable ligation conditions, or using another joining method, such as, but not limited to a topoisomerase (e.g., see U.S. Patent No. 5,766,891, incorporated herein by reference), under suitable joining conditions, so as to obtain a circular promoter-containing first-strand cDNA. By ligation of the phosphorylated 5'-end of linear first-strand cDNA to its 3'-end, the transcription promoter is joined to the target sequence so that *in vitro* transcription using an RNA polymerase of the invention under transcription conditions will synthesize transcription products corresponding to the target sequence.

[00470] In the example in FIG. 22, the target sequence comprises target nucleic acid comprising all mRNA molecules in a sample and the transcription products that are made by transcription of the transcription substrate comprise RNA that is essentially the same as the sense mRNA. Thus, in this embodiment, the circular first-strand cDNA comprises a ssDNA transcription substrate of the invention, which embodiment is useful for many applications. If there is no sequence in the circular ssDNA transcription substrate that results in termination of transcription, transcription continues around and around the circular ssDNA transcription substrate multiple times and generates concatemers of sense transcription products (i.e., comprising tandem copies of the same nucleic acid sequence as an mRNA target nucleic acid sequence in the sample), which concatemers are useful for certain applications of the invention.

[00471] In some embodiments of the invention, one or more transcription termination sequences is/are incorporated into the promoter primer between the target sequence-complementary 3'-region and the transcription promoter 5'-portion in order to permit synthesis of single-copy rather than concatemeric sense transcription products. For example, if a transcription termination sequence is present in the promoter primer, the transcription product can correspond in length to a single copy of an mRNA target nucleic acid sequence following *in vitro* transcription of the circular ssDNA transcription substrate using an RNA polymerase of the invention. Transcription termination sequences are known in the art and those with knowledge in the art will know how to find information about the sequences, as well as experimental

methods for identifying additional termination sequences that can be used. By way of example, but not of limitation, information about transcription termination sequences can be found in a book entitled "RNA Polymerases and the Regulation of Transcription," edited by Reznikoff, W.S., et al., (Elsevier Science Publishing Co., Inc., New York, 1987) and in Section 17 of a book by Miller, J.H. entitled "A Short Course in Bacterial Genetics. A Laboratory Handbook for Escherichia coli and Related Bacteria" (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1992), both incorporated herein by reference.

[00472] In still other embodiments, the circular ssDNA transcription substrate is linearized prior to use for *in vitro* transcription in order to form a "linear ssDNA transcription substrate." By way of example, but not of limitation, the circular transcription substrate can be linearized by treatment with uracil-N-glycosylase ("UNG") and endonuclease IV ("endo IV") (e.g., see methods in U.S. Patent No. 6,048,696, which is incorporated herein by reference) if the promoter primer is synthesized to have a dUMP nucleotide between the target sequence-complementary 3'-region and the transcription promoter in its 5'-portion. However, the use of a promoter primer comprising a dUMP nucleotide can only be used in embodiments of the invention which are performed in a stepwise manner, because the presence of UNG in a continuous reaction would cleave the transcription promoter portion of a promoter primer from the target-complementary portion of the promoter primer, thus destroying the ability of the promoter primer to generate more ssDNA transcription substrates. There are a number of other methods known in the art for linearizing a circular DNA molecule, which can be used in embodiments of the invention, and those with knowledge in the art will know or know how to find suitable methods for use in the invention. By way of example, but not of limitation, a number of such methods which can be used are described herein in the section entitled "Methods for Defining the 5'- and 3'-Ends of Target Sequences That Comprise Only a Portion of a Larger RNA or DNA Target Nucleic Acid."

[00473] Other embodiments of the invention comprise use of a promoter primer for synthesis of a second-strand cDNA as a transcription substrate of the invention. In those embodiments, a transcription promoter can be incorporated into the second-strand cDNA by, either (a) synthesizing first-strand cDNA using a promoter primer that has a sequence in its 5'-portion comprising a sequence that is complementary to a sense transcription promoter (i.e., it comprises an anti-sense promoter primer) or, (b) using a promoter primer comprising a sense transcription promoter for synthesis of second-strand cDNA. The promoter sequence used in a promoter primer can be determined based on knowledge of sequences of sense promoters for

RNA polymerases of the invention and of the rules of nucleic acid base complementarity and the directionality of DNA and RNA synthesis by DNA and RNA polymerases. By way of example, if a sense transcription promoter is desired in a second-strand cDNA that is made by primer extension of first-strand cDNA, which is in turn made by reverse transcription of mRNA, one can work backwards from having a sense transcription promoter of known sequence in a transcription substrate at the 3'-end of a target nucleic acid sequence that is to be amplified by an RNA polymerase of the invention, in order to determine the appropriate sense or anti-sense sequence (and position of the sequence with respect to the target sequence) that is needed, respectively, for promoter primers that are used for first-strand cDNA synthesis, or for second-strand cDNA synthesis.

[00474] In some embodiments of the invention, more than one promoter can be present on the promoter primer. By way of example, but not of limitation, a promoter primer can encode two promoter sequences, both of which encode sense promoters on first-strand cDNA (e.g., for two different RNA polymerases of the invention). Alternatively, a first promoter sequence of the promoter primer and the resulting first-strand cDNA can comprise a sense promoter and a second promoter sequence can comprise an anti-sense promoter, in which case, the first promoter sequence will be anti-sense and the second promoter sequence will be sense in second-strand cDNA. In embodiments that include additional rounds of transcription by using RNA from the first round to obtain a second transcription substrate for transcription, it is necessary to take into account the fate of the promoter sequences through the subsequent rounds of transcription when designing the reaction.

[00475] In addition to the transcription promoter sequence and the target-complementary sequence, a promoter primer of the invention can also have additional nucleic acid sequences that are 5'-of and/or 3'-of the transcription promoter sequence, but a promoter primer is not required to have such additional other sequences. By way of example, but not of limitation, a promoter primer can have a transcription initiation site 5'-of the promoter sequence. In some embodiments of the invention, a promoter primer can have one or more transcription termination sequences, one or more sites for DNA cleavage, (such as, but not limited to, a dUMP residue that can be cleaved using uracil-N-glycosylase and endonuclease IV, or other cleavage methods discussed elsewhere herein) to permit controlled linearization of a circular first-strand cDNA that is a transcription substrate, one or more origins ("ori's") of replication (preferably an *ori* for a single-stranded replicon, such as, but not limited to, a phage M13 replicon), a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene or a beta-

galactosidase gene, respectively, or one or more transposon recognition sequences (e.g., OE or ME sequences) that can be recognized and used by a transposase for *in vitro* or *in vivo* transposition, or one or more sites that are recognized by a recombinase (such as, but not limited to, the cre-lox system), and/or other sequences or genetic elements for a particular purpose. After reading the specification of the present invention, those with knowledge in the art will know that a sequence that is 5'-of a functional promoter will be transcribed by an RNA polymerase of the invention, and will therefore know where to position particular additional sequences or genetic elements relative to the promoter sequence in a promoter primer. In some embodiments of the invention, a promoter primer has a 5'-phosphate or is phosphorylated at its 5'-end using an enzyme, such as, but not limited to, a polynucleotide kinase (e.g., T4 PNK), during the processes of a method of the invention. A primary reason for providing a 5'-phosphate group on a promoter primer is to permit ligation of linear first-strand cDNA following reverse transcription or primer extension of a promoter primer on a target sequence in order to make a ssDNA transcription substrate comprising circular first-strand cDNA.

[00476] A number of examples of embodiments that use promoter primers of the invention are described below. The invention comprises all methods for using a promoter primer wherein the transcription promoter can be used by an RNA polymerase that can synthesize RNA using a ssDNA transcription substrate and is not limited to only the example embodiments presented.

[00477] With respect to methods that use a promoter primer, one embodiment of the invention comprises a method for obtaining a circular ssDNA transcription substrate of the invention for making a transcription product corresponding to a target sequence in a target nucleic acid, the method comprising:

[00478] a. obtaining a single-stranded promoter primer for synthesis of a first-strand cDNA, the promoter primer comprising a sequence at its 3'-end that is complementary to the 3'-end of the target sequence that is to be transcribed, and a 5'-portion comprising a single-stranded sense transcription promoter, wherein the promoter functions to direct transcription by an RNA polymerase that can use a ssDNA transcription substrate for synthesis of RNA or modified RNA that is complementary to the target sequence, and optionally, a phosphate group or a topoisomerase moiety on its 5'-end;

[00479] b. annealing the promoter primer to the target nucleic acid;

[00480] c. primer-extending or the promoter primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand

cDNA that is complementary to the target sequence or sequences to which the promoter primer was annealed, and wherein, the linear first-strand cDNA comprises the transcription promoter at a position which is 5'-of the target sequence in the first-strand cDNA which is complementary to the target sequence;

[00481] d. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA;

[00482] e. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently attached to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA, wherein the circular first-strand cDNA comprises a circular ssDNA transcription substrate; and

[00483] f. obtaining the circular ssDNA transcription substrate.

[00484] Another embodiment of the invention comprises a method for obtaining a linear ssDNA transcription substrate for making a transcription product corresponding to a target sequence in a target nucleic acid, the method comprising:

[00485] a. obtaining a circular ssDNA transcription substrate by carrying out steps (a) through (f) of the method above; then

[00486] b. linearizing the circular ssDNA transcription substrate at a site 3'-of the transcription promoter and 5'-of the cDNA target sequence that is complementary to the target nucleic acid sequence, wherein a linear ssDNA transcription substrate is obtained; and

[00487] c. obtaining the linear ssDNA transcription substrate.

[00488] A general embodiment of the invention comprises a method for making a transcription product corresponding to a target sequence in a target nucleic acid, the method comprising:

[00489] a. obtaining a ssDNA transcription substrate, chosen from among a circular ssDNA transcription substrate and a linear ssDNA transcription substrate;

[00490] b. contacting the ssDNA transcription substrate with an RNA polymerase that transcribes the ssDNA transcription substrate using the single-stranded transcription promoter under transcription conditions so as to obtain transcription product; and

[00491] c. obtaining the transcription product.

[00492] The target nucleic acid can be DNA or RNA. By way of example, but not of limitation, a target sequence can comprise a target nucleic acid comprising a single species of mRNA or a target sequence can comprise a target nucleic acid, which can comprise all of the mRNA in a sample.

[00493] Thus, one embodiment of the invention is a method for using a promoter primer for making a transcription product corresponding to a target sequence comprising a target nucleic acid comprising mRNA (FIG. 22), the method comprising

[00494] a. obtaining a target nucleic acid comprising mRNA;

[00495] b. obtaining a single-stranded promoter primer for synthesis of a first-strand cDNA, the promoter primer comprising: (i) a sequence at its 3'-end that is complementary to the 3'-end of the target sequence that is to be transcribed, wherein the complementary sequence on the promoter primer is chosen from among (a) and oligo(dT) sequence, (b) an anchored oligo(dT) sequence, (c) a sequence that is complementary to a specific sequence at the 3'-end of an mRNA, and (d) a sequence of nucleotides, each of which nucleotides comprises a random nucleotide base that is complementary to any of the four canonical nucleotide bases in RNA; and (ii) a 5'-portion comprising a single-stranded sense transcription promoter, wherein the promoter functions to direct transcription by an RNA polymerase that can use a ssDNA transcription substrate for synthesis of RNA or modified RNA that is complementary to the target sequence; and (iii) optionally, a phosphate group or a topoisomerase moiety on its 5'-end;

[00496] c. annealing the promoter primer to the target nucleic acid;

[00497] d. primer-extending the promoter primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the target sequence or sequences to which the promoter primer was annealed, and wherein, the linear first-strand cDNA comprises the transcription promoter at a position which is 5'-of the target sequence in the first-strand cDNA which is complementary to the target sequence;

[00498] e. optionally, removing the RNA that is annealed to the linear first-strand cDNA;

[00499] f. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently attached to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA, wherein the circular first-strand cDNA comprises a circular ssDNA transcription substrate;

[00500] g. optionally, linearizing the circular ssDNA transcription substrate at a site that is 3'-of the promoter sequence and 5'-of the cDNA target sequence that is complementary to the target sequence, wherein a linear ssDNA transcription substrate is obtained;

[00501] h. contacting the circular ssDNA transcription substrate from step f, or the linear ssDNA transcription substrate from step g with an RNA polymerase that transcribes the ssDNA

transcription substrate using the promoter under transcription conditions so as to obtain transcription product; and

[00502] i. obtaining the transcription product.

[00503] The methods for obtaining a ssDNA transcription substrate and for making a transcription product corresponding to a target sequence can be performed in a stepwise manner, or, under suitable reaction conditions, they can be performed continuously in a single reaction mixture.

[00504] Thus, one embodiment of the invention comprises a method for obtaining additional rounds of transcription of a target sequence in a target nucleic acid (FIG. 23), the method comprising:

[00505] a. obtaining the transcription products from step c of the methods above (such as for example, a linear ssDNA transcription substrate for making a transcription product corresponding to a target sequence in a target nucleic acid);

[00506] b. obtaining a single-stranded promoter primer for synthesis of a first-strand cDNA, the promoter primer comprising: (i) a sequence at its 3'-end that is complementary to the 3'-end of the target sequence that is to be transcribed, and (ii) a 5'-portion comprising a single-stranded sense transcription promoter, wherein the promoter functions to direct transcription by an RNA polymerase that can use a ssDNA transcription substrate for synthesis of RNA or modified RNA that is complementary to the target sequence, and (iii) optionally, a phosphate group or a topoisomerase moiety on its 5-end;

[00507] c. annealing the promoter primer to the target nucleic acid;

[00508] d. primer-extending the promoter primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the target sequence or sequences to which the promoter primer was annealed, and wherein, the linear first-strand cDNA comprises the transcription promoter at a position that is 5'-of the target sequence in the first-strand cDNA which is complementary to the target sequence;

[00509] e. optionally, removing the RNA that is annealed to the linear first-strand cDNA;

[00510] f. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently attached to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA, wherein the circular first-strand cDNA comprises a circular ssDNA transcription substrate;

[00511] g. optionally, linearizing the circular ssDNA transcription substrate at a site that is 3'-of the promoter sequence and 5'-of the cDNA target sequence that is complementary to the target sequence, wherein a linear ssDNA transcription substrate is obtained;

[00512] h. contacting the circular ssDNA transcription substrate from step f, or the linear ssDNA transcription substrate from step g with an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions so as to obtain transcription products; and

[00513] i. obtaining the additional transcription products.

[00514] Different embodiments of methods of the invention can also be used to make a transcription product corresponding to target sequences that are internal to a target nucleic acid sequence. By way of example, some embodiments can be used to make a transcription product corresponding to sequences, such as, but not limited to, wild-type or mutated sequence in genomic DNA. In these embodiments, one or more processes, such as, but not limited to, annealing a blocking oligo to the target nucleic acid, are required to limit the 3'-end of the target sequence that is transcribed, as is discussed elsewhere herein. In general, the target nucleic acid must be single-stranded for use in a method of the invention. Thus, a double-stranded nucleic acid must be denatured.

[00515] Thus, one embodiment of the invention for using a promoter primer is a method for making a transcription product corresponding to a target sequence that comprises only a portion of a target nucleic acid comprising single-stranded DNA or RNA, the method comprising

[00516] a. obtaining a target nucleic acid comprising single-stranded DNA or RNA; then

[00517] b. obtaining a single-stranded promoter primer for synthesis of a first-strand cDNA, the promoter primer comprising: (i) a sequence at its 3'-end that is complementary to the 3'-end of the target sequence that is to be transcribed, and (ii) a 5'-portion comprising a single-stranded sense transcription promoter, wherein the promoter functions to direct transcription by an RNA polymerase that can use a ssDNA transcription substrate for synthesis of RNA or modified RNA that is complementary to the target sequence, and (iii) optionally, a phosphate group or a topoisomerase moiety on its 5-end;

[00518] c. obtaining a blocking oligo, the blocking oligo comprising a sequence that anneals tightly to a sequence on the target nucleic acid so as to delimit the 3'-end of a primer extension product of the promoter primer using the target nucleic acid as a template, wherein the

blocking oligo is not displaced by the primer extension product, and wherein the blocking oligo is not itself capable of being primer extended by a DNA polymerase;

[00519] d. annealing the promoter primer and the blocking oligo to the target nucleic acid;

[00520] e. primer-extending the promoter primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand cDNA that is complementary to the target sequence to which the promoter primer was annealed and that is not complementary to the sequence where the blocking oligo is annealed, and wherein, the linear first-strand cDNA comprises the transcription promoter at a position which is 5'-of the target sequence in the first-strand cDNA which is complementary to the target sequence;

[00521] f. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA;

[00522] g. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently attached to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA, wherein the circular first-strand cDNA comprises a circular transcription substrate;

[00523] h. optionally, linearizing the circular ssDNA transcription substrate at a site that is 3'-of the promoter sequence and 5'-of the cDNA target sequence that is complementary to the target sequence, wherein a linear ssDNA transcription substrate is obtained;

[00524] i. contacting the circular ssDNA transcription substrate from step g, or the linear ssDNA transcription substrate from step h with an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions so as to obtain transcription products; and

[00525] j. obtaining the transcription products; and

[00526] k. optionally, repeating steps b through k to obtain additional rounds of transcription to obtain transcription products.

[00527] In still other embodiments of the invention, which are preferred embodiments, the circular first-strand cDNA does not comprise a transcription substrate of the invention. Rather, in those embodiments, the circular first-strand cDNA is used as a template for DNA synthesis using a strand-displacing DNA polymerase and at least one strand displacement primer, and in some embodiments, multiple strand displacement primers. A "strand displacement primer," as used herein, is an oligonucleotide or polynucleotide that can be primer extended by a strand displacing DNA polymerase of the invention, wherein strand displacement DNA synthesis

occurs. In general, strand displacement is more a property of the DNA polymerase and the reaction conditions used than of the primer. Thus, the composition and properties of a strand displacement primer can vary greatly. For example, in some embodiments of the invention, a strand displacement primer can be an oligonucleotide that is hybridizable to a circular DNA template, wherein the DNA synthesis product resulting from rolling circle replication by a strand displacing DNA polymerase results in displacement of the ssDNA strand displacement primer extension product, resulting in tandem complementary ssDNA copies of circular template. In other embodiments, it is preferred that the strand displacement primer has a 3'-portion that is complementary to a sequence in a circular DNA template and a 5'-portion that is non-complementary, and therefore has a "flap;" the protruding flap appears to facilitate displacement of the primer in some cases. When strand displacement is carried out on linear DNA templates, strand displacement primers can be designed to have particular nucleotide compositions and/or structures, and additional methods and reaction components can be used in order to facilitate strand displacement by liberating the strand displacement primer from the template. This is discussed further in the section of the present invention entitled "Methods of the Invention for Strand Displacement Replication on Linear Templates."

[00528] A strand-displacing DNA polymerase of the invention can be any DNA polymerase that results in strand displacement. Preferred strand-displacing DNA polymerases of the present invention lack 5'-exonuclease activity, including structure-dependent 5'-nuclease activity. Preferred strand-displacing DNA polymerases comprise rBst DNA polymerase large fragment, also called "IsoTherm™ DNA Polymerase" (Epicentre Technologies, Madison, Wisconsin, USA), Bca DNA Polymerase (TAKARA Shuzo Company, Kyoto, Japan), ϕ 29 DNA polymerase (U.S. Patent Nos. 5,198,543 and 5,001,050, incorporated herein by reference), SequiTherm™ DNA Polymerase (Epicentre Technologies, Madison, Wisconsin, USA), MMLV reverse transcriptase, and Sequenase® DNA Polymerase (USB, Cleveland, Ohio, USA). In these embodiments, a strand displacement primer is used to prime second-strand DNA synthesis using circular first-strand cDNA as a template. Once DNA synthesis has proceeded completely around the circular first-strand cDNA template, the second-strand cDNA is displaced so that the displaced single-stranded second DNA strand is released into the reaction medium. Since a transcription promoter is present at least once in every round of DNA synthesis of the circular first-strand cDNA template, this released single-stranded second-strand DNA can be used as a ssDNA transcription substrate by an RNA polymerase of the invention. However, in this case, the RNA that is synthesized using the single-stranded second-strand DNA as a ssDNA

transcription substrate comprises anti-sense RNA, rather than sense RNA as in previous embodiments that used first-strand cDNA as a template. Anti-sense RNA can be used for many applications, such as, but not limited to, for use as probes for nucleic acid arrays. As the second DNA strand continues to grow, longer and longer concatemers of the anti-sense RNA are formed.

[00529] With respect to embodiments of the invention that use a promoter primer, some embodiments obtain a ssDNA transcription template for making a transcription product corresponding to a target sequence in a target nucleic acid by first obtaining a circular first-strand cDNA comprising an anti-sense transcription promoter, wherein the circular first-strand cDNA is then used as a template for strand displacement DNA synthesis of the linear ssDNA transcription substrate by a strand-displacing DNA polymerase.

[00530] Thus, one embodiment of the invention that uses a promoter primer comprises a method for making a transcription product corresponding to a target sequence in a target nucleic acid (FIG. 24), the method comprising:

[00531] a. obtaining a target nucleic acid comprising single-stranded DNA or RNA;

[00532] b. obtaining a single-stranded anti-sense promoter primer for synthesis of a first-strand cDNA, the anti-sense promoter primer comprising: (i) a sequence at its 3'-end that is complementary to the 3'-end of the target sequence that is to be transcribed, and (ii) a 5'-portion comprising a single-stranded sequence that is complementary to a sense transcription promoter (i.e., an "anti-sense promoter"), wherein the sense promoter functions to direct transcription by an RNA polymerase that can use a ssDNA transcription substrate for synthesis of RNA or modified RNA that is complementary to the target sequence, and (iii) optionally, a phosphate group or a topoisomerase moiety on its 5-end;

[00533] c. optionally, (if the 5'-end of the target sequence in the target nucleic acid is not a 5'-terminus or otherwise delimited,) obtaining a blocking oligo, the blocking oligo comprising a sequence that anneals tightly to a sequence on the target nucleic acid so as to delimit the 3'-end of a primer extension product of the promoter primer using the target nucleic acid as a template, wherein the blocking oligo is not displaced by the primer extension product, and wherein the blocking oligo is not itself capable of being primer extended by a DNA polymerase;

[00534] d. annealing the promoter primer and, optionally, the blocking oligo to the target nucleic acid;

[00535] e. primer-extending or the promoter primer annealed to the target nucleic acid with a DNA polymerase under DNA synthesis conditions so as to obtain a linear first-strand

cDNA that is complementary to the target sequence to which the promoter primer was annealed and that is not complementary to the sequence where the blocking oligo is annealed (if used), and wherein, the linear first-strand cDNA comprises the anti-sense transcription promoter at a position which is 5'-of the target sequence in the first-strand cDNA which is complementary to the target sequence;

[00536] f. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA; then,

[00537] g. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently attached to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA;

[00538] h. obtaining a strand-displacement primer;

[00539] i. annealing the strand displacement primer to the circular first-strand cDNA;

[00540] j. obtaining a strand-displacing DNA polymerase;

[00541] k. contacting the circular first-strand cDNA to which the strand displacement primer is annealed with a strand-displacing DNA polymerase under DNA synthesis conditions so as to obtain linear second-strand cDNA, wherein the second-strand cDNA comprises a ssDNA transcription template comprising a sense transcription promoter;

[00542] l. obtaining the linear second-strand ssDNA transcription template;

[00543] m. obtaining an RNA polymerase that transcribes the ssDNA transcription substrate using the promoter under transcription conditions;

[00544] n. contacting the linear second-strand ssDNA transcription template with the RNA polymerase under transcription conditions so as to obtain anti-sense transcription products; and

[00545] o. obtaining the anti-sense transcription products; and

[00546] p. optionally, repeating steps b through p to obtain additional rounds of transcription of the anti-sense transcription products.

b. Composition of Promoter Primers of the Invention

[00547] In preferred embodiments of the invention, a promoter primer comprises DNA nucleotides. A promoter primer can also comprise one or more modified nucleotides for a particular purpose. By way of example, but not of limitation, a promoter primer can comprise one or more dUMP nucleotides 3'-of a sense transcription promoter and 5'-of the sequence that is complementary to a target nucleic acid sequence, which provides a site for linearizing a

circular ssDNA transcription substrate using UNG and endo IV, as discussed elsewhere herein. However, the invention is not limited to promoter primers comprising DNA nucleotides or modified DNA nucleotides, and in some cases, a promoter primer can comprise RNA or modified RNA nucleotides or both DNA and RNA nucleotides or modified nucleotides.

[00548] The nucleic acid target-complementary portion of a promoter primer can be complementary to a specific known sequence in the RNA target in a sample, or it can comprise a mixture of all possible or many possible sequences, such as, but not limited to, random hexamer sequences. Random primer sequences can be made by including nucleotide reagents which are complementary to all four canonical bases during the chemical synthesis of each nucleotide position of the mRNA-complementary portion of the promoter primer. In embodiments of the invention using samples containing mRNA target nucleic acids, which are preferred embodiments, the 3'-end of a promoter primer comprises either a specific sequence that is complementary to a known sequence of a specific mRNA or, if the mRNA has a poly(A) tail at its 3'-end, the 3'-end of the promoter primer can comprise an oligo(dT) sequence. In still other embodiments of the invention for mRNA target nucleic acids, the 3'-end of a promoter primer can comprise a random sequence, such as, but not limited to a random hexamer sequence.

[00549] A promoter primer of the invention comprises a transcription promoter in its 5'-portion. In most embodiments, the transcription promoter comprises a sense promoter sequence that is capable of binding an RNA polymerase of the invention. However, those with knowledge in the art will understand that, due to the fact that the transcription promoter is 5'-of the linear first-strand cDNA that is synthesized by reverse transcription of the RNA target nucleic acid using the promoter primer, an RNA polymerase of the invention cannot use the transcription promoter to synthesize RNA complementary to first-strand cDNA that is complementary to the RNA target nucleic acid. However, if the 5'-end of the linear first-strand cDNA is ligated to its 3'-end so as to form circular first-strand cDNA, then an RNA polymerase of the invention can use the transcription promoter to synthesize RNA complementary to first-strand cDNA that is complementary to the RNA target nucleic acid; thus, a circular first-strand cDNA of this embodiment comprises a transcription substrate of the invention. Thus, a preferred promoter primer of these embodiments of the invention is phosphorylated at its 5'-end in order to facilitate ligation of linear first-strand cDNA by a ligase of the invention under ligation conditions. By means of example, but not of limitation, the 5'-end of the promoter primer can be phosphorylated using T4 polynucleotide kinase and ATP under suitable reaction conditions known in the art. In other embodiments, the promoter primer has a type 1 topoisomerase moiety

at its 5'-end in order to facilitate topoisomerase-mediated ligation of linear first-strand cDNA under ligation conditions (e.g., U.S. Patent No. 5,766,891, incorporated herein by reference).

D. Composition, Design and Use of Promoter Splice Template Oligos, Splice Template Oligos, Promoter Ligation Oligos, Ligation Oligos, Promoter Primers, Primers, and Ligation Splint Oligos of the Invention

[00550] Depending on the particular method and application, a promoter splice template oligo, a splice template, a promoter ligation oligo, a ligation oligo, a promoter primer, a primer, or a ligation splint oligo, or oligos used for other purposes can comprise unmodified or modified DNA, RNA or both DNA and RNA for various applications and embodiments of the invention, some of which are discussed in greater detail in individual sections related thereto. By way of example, but not of limitation, in some embodiments, these molecules can comprise DuraScript™ RNA made using the DuraScribe™ T7 Transcription Kit (Epicentre Technologies, Madison, WI). Similar types of modified RNA oligos that contain 2'-F-dCMP and 2'-F-dUMP can also be made using the N4 mini-vRNAP Y678F mutant enzyme, described herein in the section on "RNA Polymerases of the Invention." DNA oligos can be synthesized on an oligo synthesizer, which is usually preferred, or enzymatically, using methods discussed elsewhere herein.

[00551] In addition to the transcription promoter sequence and a target-complementary sequence, if present, the various promoter-containing oligos of the invention, including, but not limited to, promoter splice template oligos, promoter ligation oligos, or promoter primers, of the invention can also have additional nucleic acid sequences that are 5'-of and/or 3'-of the transcription promoter sequence, but a promoter-containing oligo is not required to have such additional other sequences. By way of example, but not of limitation, these oligos can have a transcription initiation site 5'-of the promoter sequence. In some embodiments of the invention, a promoter-containing oligo can have one or more primer sites, one or more transcription termination sequences, one or more sites for DNA cleavage, (such as, but not limited to, a dUMP residue that can be cleaved using uracil-N-glycosylase and endonuclease IV, or other cleavage methods discussed elsewhere herein) to permit controlled linearization of a circular first-strand cDNA that is a transcription substrate, one or more origins ("ori's") of replication (preferably an *ori* for a single-stranded replicon, such as, but not limited to, a phage M13 replicon), a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene or a beta-galactosidase gene, respectively, or one or more transposon recognition sequences (e.g., OE or ME sequences) that can be recognized and used by a transposase for *in vitro* or *in*

vivo transposition, or one or more sites that are recognized by a recombinase (such as, but not limited to, the cre-lox system), and/or other sequences or genetic elements for a particular purpose. Similarly, other oligos used in a method of the invention, including those that lack a promoter sequence, such as splice template oligos, ligation oligos, or primers can have one or more of any of these additional sequences and/or genetic elements 5'-of the target-complementary or target-joining portion of each respective oligo.

[00552] The various methods of the invention, including, but not limited to the various processes for attaching one or more sense and/or anti-sense transcription promoter sequences to a target nucleic acid sequence, are not mutually exclusive of each other, and in some embodiments of the invention, more than one method can be used together. By way of example, but not of limitation, both an oligo(dT)-promoter primer and a splice template oligo that is used to primer-extend the 3'-end of a first-strand cDNA that is synthesized by reverse transcription of mRNA from a sample can be used. In some embodiments, the splice template oligo can be a promoter splice template oligo, which encodes a promoter sequence, and in other embodiments, a splice template oligo that does not encode a promoter sequence can be used. A splice template oligo that does not encode a promoter can be used, without limitation, to provide a primer binding site, a single-stranded origin of replication ("ori"), such as a phage M13 ori, a selectable or screenable marker, such as a sequence that encodes an antibiotic-resistance gene or a beta-galactosidase gene, respectively, and/or a transcription termination sequence. If the splice template oligo encodes a promoter sequence, it can encode a sequence that comprises a sense transcription promoter in the synthesized first-strand cDNA or it can encode a sequence that results in a sense promoter in a second cDNA strand that is synthesized by strand displacement DNA synthesis using first-strand cDNA as a template. In some embodiments of the invention, more than one sense transcription promoter, which can have the same DNA sequence or a different DNA sequence, can be present in first-strand cDNA and/or second-strand cDNA. The promoter sequence used in a promoter splice template oligo can be determined based on knowledge of sequences of sense promoters for RNA polymerases of the invention and of the rules of nucleic acid base complementarity and the directionality of DNA and RNA synthesis by DNA and RNA polymerases. By way of example, if a sense transcription promoter is desired in a second-strand cDNA that is made by primer extension of first-strand cDNA, which in turn is made by reverse transcription of mRNA, one can work backwards from having a sense transcription promoter of known sequence at the 3'-end of a target nucleic acid sequence that is to be transcribed by an RNA polymerase of the invention, in order to determine the anti-sense

sequence (and directionality of transcription encoded by the sequence) that is needed in the first-strand cDNA, and then, what is needed in the promoter splice template oligo to which the primer-extended first-strand cDNA must be complementary in order for the promoter splice template oligo to serve as a template for primer extension of the first cDNA strand. Based on a reading of the disclosures herein, those with knowledge in the art will understand other methods of the invention which use processes with other different combinations or permutations of promoter splice template oligos, splice template oligos, promoter ligation oligos, promoter primers, and non-promoter primers to make ssDNA transcription substrates having ssDNA promoters that are recognized by RNA polymerases of the invention and that can be used to synthesize or make a transcription product corresponding to a target nucleic acid sequence for a particular purpose. All such methods are included in the invention.

E. Circularizing Linear ssDNA

1. Methods for Cloning ssDNA and Generating Libraries of ssDNA Clones, Including Expression Clones and Expressions Libraries

[00553] In some embodiments, the methods of the invention for obtaining a circular ssDNA transcription substrate corresponding to a target sequence for a target nucleic acid comprise obtaining molecular clones of the target sequence. In still other embodiments, such as, but not limited to embodiments in which the target sequence comprises first-strand cDNA corresponding to all target mRNAs in a sample, the invention comprises methods for obtaining a library of clones comprising the target sequences of target nucleic acid. In embodiments that comprise obtaining a clone or a library of clones comprising target sequences, a promoter splice template oligo, a promoter ligation oligo, or a promoter primer that comprises a single-stranded origin of replication (*ori*), such as, but not limited to an M13 *ori*, is used, and in most embodiments, a sequence that encodes a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene or a beta-galactosidase gene, respectively, is also used, and, optionally, a transcription termination sequence or other sequences or genetic elements are also used. Methods, *E. coli* host cells, bacteriophage (including helper phage) strains, and vectors and sequences, including M13 *ori* sequences, that are needed or useful for working with M13 phages and M13 nucleic acids and vectors and sequences derived therefrom are available and well-known in the art (e.g., see Chapter 3, Volume 1 of Sambrook and Russell, Molecular Cloning, A Laboratory Manual, Third Edition, 2001, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference) and these methods and compositions can

be used for methods of the invention for obtaining replicable clones of single-stranded DNA corresponding to target sequences of target nucleic acid. Also, a number of selectable or screenable markers and selection and screening systems are known in the art, such as, but not limited to those described by Sambrook and Russell Molecular Cloning, A Laboratory Manual, Third Edition (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; e.g., see Chapter 1, Volume 1), which is incorporated herein by reference, and any suitable marker or selection or screening system can be used for the present invention.

[00554] Thus, one embodiment of the invention is a method for cloning a target sequence, the method comprising:

[00555] a. obtaining a single-stranded promoter primer comprising a 3'-end portion that has a sequence that is complementary to the 3'-end of a target sequence and a 5'-portion, the 5'-portion comprising a sequence at or near the 5'-end that encodes a single-stranded sense transcription promoter for an RNA polymerase that can make a transcription product using a ssDNA transcription substrate and additional sequences 3'-of the promoter sequence, the additional sequences comprising a single-stranded origin of replication that can be replicated in a host cell, at least one gene for a selectable or screenable marker that is expressible in the host cell, and optionally, the 5'-end of the promoter primer can have a phosphate group or a topoisomerase moiety or a phosphate or topoisomerase moiety can be added to the 5'-end of linear first-strand cDNA as an additional step of the method prior to the ligation step;

[00556] b. annealing the promoter primer to the target nucleic acid;

[00557] c. obtaining a DNA polymerase;

[00558] d. primer-extending the promoter primer annealed to the target nucleic acid with the DNA polymerase under DNA synthesis conditions;

[00559] e. obtaining a linear first-strand cDNA comprising a sequence that is complementary to the target sequence;

[00560] f. optionally, removing the target nucleic acid that is annealed to the linear first-strand cDNA;

[00561] g. ligating the linear first-strand cDNA, wherein the 5'-end of the linear first-strand cDNA is covalently joined to the 3'-end of the linear first-strand cDNA so as to obtain circular first-strand cDNA, wherein the circular first-strand cDNA comprises a circular ssDNA transcription substrate;

[00562] h. obtaining the circular ssDNA transcription substrate;

- [00563]** i. obtaining host cells that can replicate a circular ssDNA comprising the single-stranded origin of replication and in which the selectable or screenable marker is expressible;
- [00564]** j. incubating the host cells with DNA comprising a circular ssDNA transcription substrate under conditions suitable to obtain transformation;
- [00565]** k. plating the host cells that were incubated with the DNA comprising a circular ssDNA transcription substrate on medium that contains a composition that permits selection or screening for cells that contain and express the gene for the selectable or screenable marker; and
- [00566]** l. obtaining transformed host cells that contain replicable clones comprising the target sequence.

[00567] In some embodiments, the host cells that are used to obtain replicable clones comprising the target sequence are host cells that comprise an inducible gene for an RNAP, wherein the RNAP can bind to the single-strand promoter and initiate transcription of a single-strand template therefrom. In particularly suitable embodiments, the host cells comprise an inducible gene that encodes an N4 vRNAP. In other embodiments, the host cells comprise an inducible gene that encodes an RNA polymerase that can bind to a pseudopromoter that is on a ssDNA transcription substrate and initiate transcription from the pseudopromoter. Inducible promoters that can be used to express the RNAP in the host cells in the presence of an inducing substance are described elsewhere herein. When host cells comprising an inducible RNAP are used to obtain replicable clones according to this embodiment of the invention, the transformed host cells that contain replicable clones, when incubated in the presence of an inducing substance under inducing conditions, comprise expression clones. Still further, in embodiments of the invention in which the target sequence comprises target nucleic acid comprising all mRNAs in a sample, the collection of all transformed host cells that contain replicable clones comprises a library of replicable clones. In yet another embodiment, if the host cells contain an inducible gene that encodes an RNAP, wherein the RNAP can bind to a single-stranded promoter and initiate transcription of a single-strand template therefrom, the collection of all transformed host cells that contain replicable clones comprises a library of replicable expression clones.

[00568] The invention also comprises embodiments of methods for cloning a target sequence by using a promoter splice template oligo or a promoter ligation oligo that comprises sequences that encode a single-stranded origin of replication (ori), such as, but not limited to an M13 ori, that is replicable in a host cell, a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene or a beta-galactosidase gene, respectively, that is

expressible in the host cell and, optionally, a transcription termination sequence. In these embodiments, the sequences that encode the ori, the selectable or screenable marker and the transcription termination sequence, if present, are 5'-of the promoter-complementary sequence (i.e., 5'-of the anti-sense promoter sequence) in the promoter splice template oligo or 3'-of the sense promoter sequence in the promoter ligation oligo, respectively. Initial steps of the methods for cloning a target sequence using a promoter splice template oligo or a promoter ligation oligo comprise steps that are identical to those described previously for using these respective oligos for obtaining a linear ssDNA transcription substrate of the invention.

However, the next step of the methods for cloning a target sequence using a promoter splice template oligo or a promoter ligation oligo comprises obtaining a circular ssDNA transcription substrate by ligating the 5'-end of the linear ssDNA transcription substrate to the 3'-end under ligation conditions. Ligation conditions addition of a phosphate group to the 5'-end of the linear ssDNA transcription substrate, if it is not already present, using kinase, such as, but not limited to T4 polynucleotide kinase. Thereafter, the steps of the methods of the invention for using a promoter splice template oligo or a promoter ligation oligo for obtaining transformed host cells that contain replicable clones, replicable expression clones, libraries or expression libraries of a target sequence for target nucleic acid in a sample are the same as described above for methods that use a promoter primer.

[00569] Based on the above description of the invention for using promoter splice template oligos, promoter ligation oligos, and promoter primers for obtaining host cells that contain replicable clones or libraries of target sequences, those with knowledge in the art will understand that the invention also comprises similar methods for obtaining host cells that contain replicable clones or libraries of target sequences using splice template oligos, ligation oligos, and primers that lack a promoter sequence. Thus, provided an oligo encodes sequences that comprise a single-stranded origin of replication and a suitable selectable or screenable marker for the host cell used, the methods for making replicable clones using an oligo that lacks a promoter sequence, including a splice template oligo, a ligation oligo or a primer, are substantially the same as those methods used for making replicable clones using a promoter splice template oligo, a promoter ligation oligo or a promoter primer, respectively. Of course, replicable clones of a target sequence made using an oligo that lacks a promoter sequence cannot express the target sequence, and therefore, expression clones and expression libraries cannot be obtained using these oligos.

2. Ligases and Ligation Methods for Circularizing Linear ssDNA

[00570] It will be clear from the above descriptions of methods for obtaining ssDNA transcription substrates that it is useful in various embodiments to ligate linear ssDNA to obtain circular ssDNA, including, but not limited to circular ssDNA transcription substrates. The invention is not limited to a specific ligase for circularizing a linear ssDNA molecule and different ligases and ligation methods can be used in different embodiments in order to accomplish a particular purpose. In embodiments that use a ligase, the 5'-end of the linear ssDNA that is ligated to obtain a circular ssDNA must have a 5'-phosphate group or the 5'-end must be phosphorylated using a polynucleotide kinase, such as, but not limited to T4 polynucleotide kinase, during the processes of the method of the invention.

[00571] A ligase that catalyzes non-homologous intramolecular ligation, such as, but not limited to ThermoPhage™ RNA Ligase II (Prokaria, Ltd., Reykjavik, Iceland), is a suitable ligase for ligating linear ssDNA to form a circular ssDNA.

[00572] NAD-dependent DNA ligases that are not active on blunt ends, such as, but not limited to Ampligase® Thermostable DNA Ligase (Epicentre Technologies, Madison, Wisconsin, USA), Tth DNA ligase, Tfl DNA ligase, and Tsc DNA Ligase (Prokaria Ltd., Reykjavik, Iceland) can be used to ligate the 5'-phosphate and 3'-hydroxyl termini of DNA ends that are adjacent to one another when annealed to a complementary DNA molecule, and are suitable ligases in embodiments of the invention that use a ligation splint oligo comprising DNA. However, the invention is not limited to the use of a particular ligase and any suitable ligase can be used. For example, T4 DNA ligase can be used in embodiments of the invention that use a ligation splint. Still further, Faruqi discloses in U.S. Patent No. 6,368,801 that T4 RNA ligase can efficiently ligate DNA ends of nucleic acids that are adjacent to each other when hybridized to an RNA strand. Thus, T4 RNA ligase is a suitable ligase of the invention in embodiments in which DNA ends are ligated on a ligation splint oligo comprising RNA or modified RNA, such as, but not limited to modified RNA that contains 2'-F-dCTP and 2'-F-dUTP made using the DuraScribe™ T7 Transcription Kit (Epicentre Technologies, Madison, WI, USA) or the N4 mini-vRNAP Y678F mutant enzyme described herein.

[00573] The invention is also not limited to the use of a ligase for covalently joining the 5'-end to the 3'-end of the same or different nucleic acid molecules in the various embodiments of the invention. By way of example, other ligation methods such as, but not limited to, topoisomerase-mediated ligation (e.g., U.S. Patent No. 5,766,891, incorporated herein by reference) can be used.

G. Modes of Performance of the Methods of the Invention

[00574] Depending on the application and its requirements and constraints, the methods of the invention can be performed in a stepwise fashion, with one set of reactions being performed, followed by purification of a reaction product or removal of reagents or inactivation of enzymes or addition of reagents before proceeding to the next set of reactions, or, in other embodiments for other applications, the methods can be performed as a continuous set of multiple reactions in a single reaction mixture. By way of example, but not of limitation, in some embodiments, each of the separate reactions for synthesis of a cDNA library and for transcription using promoter-containing first-strand cDNA as a template for transcription can be performed separately. Still by way of example, in some embodiments in which the methods of the invention are used as part of a diagnostic assay, all of the reactions can be carried out in a single reaction mixture and the products of the transcription reaction may be detected, without ever being isolated.

[00575] The invention also comprises parts or subsets of the methods and compositions of the invention. By way of example, but not of limitation, in addition to obtaining a method for making a transcription product corresponding to the mRNA molecules in a sample, the invention also comprises improved methods for making full-length first-strand cDNA with DNA sequences corresponding to the complete 5'-end of the mRNA in the sample, and cloning the single-stranded cDNA using a single-stranded origin of replication, such as a phage M13 *ori*. This method for making and cloning full-length cDNA corresponding to mRNA in a sample can be, but need not be, part of the method for making a transcription product. Thus, the invention comprises all of the individual steps of the methods of the invention that are enabled thereby, in addition to the overall methods.

G. Examples of the Scope of Applications of the Invention

[00576] Those with knowledge in the art will understand that the present invention is novel and very broad in scope and provides improvements in methods, processes, compositions and kits related to making transcription products corresponding to a target nucleic acid sequence comprising RNA, including mRNA, or DNA in a biological sample for many applications. These methods are useful for applications such as, but not limited to, making full-length cDNA and cDNA libraries, improving gene expression analysis, and detecting a target sequence. By way of further example, but not of limitation, the present invention comprises improved methods, processes, compositions and kits that improve upon methods and applications to:

1. amplify nucleic acid molecules *in vitro*
2. amplify nucleic molecules *in vivo*
3. amplify a DNA sequence *in vitro*
4. amplify a DNA sequence *in vivo*
5. amplify a genomic DNA sequence *in vitro*
6. amplify a genomic DNA sequence *in vivo*
7. amplify an RNA sequence *in vitro*
8. amplify an RNA sequence *in vivo*
9. amplify mRNA
10. amplify rRNA
11. make cDNA complementary to an RNA
12. make full-length first-strand cDNA complementary to mRNA
13. make a cDNA library complementary to mRNA in a sample
14. synthesize RNA
15. synthesize modified RNA, such as, but not limited to, RNA containing 2'-fluoro-nucleotides for various applications, including, but not limited to, other applications for modified RNA or dsRNA herein above or below.
16. synthesize DNA
17. detect the presence of a nucleic acid sequence in a sample
18. detect the presence of a target nucleic acid sequence in a sample that is indicative of the presence of a target organism
19. detect the presence of a target nucleic acid analyte in a sample
20. detect the presence of a DNA sequence in a sample
21. detect the presence of an RNA sequence in a sample
22. detect the presence (or absence at a detectable level) of a gene in a sample
23. detect the presence (or absence at a detectable level) of a mutated gene in a sample
24. detect the presence of a target organism in a sample
25. detect the presence of a virus in a sample
26. detect the presence of a bacterium in a sample
27. detect the presence of a pathogenic organism in a sample
28. detect the presence of a beneficial organism in a sample
29. identify and quantify nucleic acids associated with RNA and DNA binding proteins using a ssDNA transcription substrate of the invention and an RNA polymerase that

can synthesize RNA using the ssDNA transcription substrate as an indirect signaling method similar to the IDAT and other methods that use a T7-type RNA polymerase, as disclosed by Zhang *et al.* (*Proc. Natl. Acad. Sci. USA* 98:5497-5502, 2001) and by Eberwine in PCT Publication No. WO 02/14476, both of which are incorporated herein by reference

30. detect an oncogene
31. detect an anti-oncogene
32. quantify the level of a virus, a microorganism, a gene, an mRNA, an rRNA, a nucleic acid analyte, or any other nucleic acid of whatever type for whatever purpose.
33. use as a probe
34. use as a probe for an array or microarray
35. make dsRNA or modified dsRNA that can be introduced into human, animal or other eukaryotic cells in serum and without use of a transfection agent.
36. synthesize dsRNA or modified dsRNA *in vitro* for use in RNAi
37. synthesize dsRNA or modified dsRNA *in vivo* for use in RNAi
38. synthesize dsRNA or modified dsRNA *in vitro* for use siRNA
39. synthesize dsRNA *in vivo* for use siRNA
40. make RNAi or modified RNAi that silences a gene encoded by a virus or other infectious agent.
41. make siRNAi or modified siRNA that silences a gene encoded by a virus or other infectious agent.
42. make RNAi or modified RNAi that silences a gene encoded by a plant, animal, human, fungal or other eukaryotic host gene that is involved with and/or interacts with a biological molecule encoded by a virus or other infectious agent.
43. make siRNAi or modified siRNA that silences a gene encoded by a plant, animal, human, fungal or other eukaryotic host gene that is involved with and/or interacts with a biological molecule encoded by a virus or other infectious agent.
44. make RNAi or modified RNAi that silences a non-essential disease-susceptibility gene encoded by a plant, animal, human, fungal or other eukaryotic host gene .
45. make siRNAi or modified siRNA that silences a non-essential disease-susceptibility gene encoded by a plant, animal, human, fungal or other eukaryotic host gene.

46. make RNAi or modified RNAi that silences a non-essential gene encoded by a plant, animal, human, fungal or other eukaryotic host gene, wherein the gene silencing results in a beneficial effect.
47. make siRNAi or modified siRNA that silences a non-essential gene encoded by a plant, animal, human, fungal or other eukaryotic host gene, wherein the gene silencing results in a beneficial effect.
48. make RNAi or modified RNAi that silences a non-essential gene encoded by a plant, animal, fungal or other eukaryotic host gene, wherein the gene silencing results in improved yield, production of a biological molecule, flavor, or other commercially beneficial effect, such as, but not limited to, cold-hardiness, salt-tolerance, shortened growing season, or increased efficiency of utilization of a nutrient.
49. make siRNAi or modified siRNA that silences a non-essential gene encoded by a plant, animal, fungal or other eukaryotic host gene, wherein the gene silencing results in improved yield, production of a biological molecule, flavor, or other commercially beneficial effect, such as, but not limited to, cold-hardiness, salt-tolerance, shortened growing season, or increased efficiency of utilization of a nutrient.
50. diagnose the presence and/or level of an infectious organism
51. diagnose a disease of any type
52. detect the presence of a nucleic acid in an environmental sample
53. differentiate, both qualitatively and quantitatively, between which mRNA molecules are present in different types of cells or in the same type of cells under different conditions or in the same or different types of cells under the same or different conditions or in response to specific stimuli or treatments
54. analyze, both qualitatively and quantitatively, gene expression profiles in cells under different defined conditions
55. analyze, both qualitatively and quantitatively, gene expression profiles in different types of cells under the same defined environmental conditions
56. analyze, both qualitatively and quantitatively, gene expression profiles in cells over time
57. analyze, both qualitatively and quantitatively, gene expression in response to specific stimuli
58. make a library or libraries of mRNA molecules and/or cDNA molecules that are present in one type of cell that are not present in another type of cell, or that are

- present in one type of cell under certain conditions but not under other conditions (i.e., a subtraction library).
59. map and clone sequences corresponding to the 5'-ends of mRNA's, including, but not limited to, those generated from a specific gene by alternative splicing and promoter usage
 60. generate improved templates for more accurate rapid amplification of cDNA ends ("RACE") techniques (e.g., see Flouriot *et al.*, Nucleic Acids Res. 27:e8 (I-iv), 1999, incorporated herein by reference).
 61. make and/or amplify mRNA for *in vitro* or *in vivo* translation, including, but not limited to coupled or step-wise transcription and translation.
 62. amplify RNA and/or DNA or modified RNA and/or DNA present in living cells, such as, but not limited to, tumor or cancer cells from a patient for introduction into dendritic cells (e.g., see U.S. Patents Nos. 5,994,126 and 6,475,483 of Steinman *et al.*, incorporated herein by reference) or other cells from the patient in order to boost or increase an *in vivo* response, such as, but not limited to, an immune response in the patient, with the goal of decreasing the size or the number of cells in the tumor or cancer in the patient.
 63. make and/or amplify RNA and/or DNA or modified RNA and/or DNA from a patient and or present in a virus or other infectious agent, wherein the RNA and/or DNA is for use as an RNA vaccine and/or a DNA vaccine, respectively.
 64. make RNA or modified RNA that can be introduced into human, animal or other eukaryotic cells in serum and without use of a transfection agent.
 65. make modified RNA containing 2'-fluoro-2'-deoxynucleotides, such as, but not limited to, 2'-F-dCMP and 2'-F-dUTP, in place of the corresponding canonical nucleotides.
 66. produce arrays or microarrays of amplified nucleic acids by attaching the amplification products onto a solid substrate.
 67. detect a mutation or a mutated form of a target nucleic acid sequence in a sample
 68. quantify the amount of a target nucleic acid or target nucleic acids in a sample.

[00577] The mini-vRNAP Y678F mutant enzyme has improved ability to incorporate 2'-fluoro-dNTPs compared to the wild-type N4 vRNAP or mini-vRNAP (Kazmierczak, K.M., *et al.*, EMBO J., 21:5815-5823, 2002, incorporated herein by reference). Modified RNA molecules that contain 2'-F-dCMP and 2'-F-dUTP are resistant to RNase A-type ribonucleases

(Sousa *et al.*, U.S. Patent No. 5,849,546), included herein by reference. Capodici *et al.*, (*J. Immunology* 169:5196-5201, 2002), included herein by reference, showed that 2'-fluoro-containing dsRNA molecules made using the DuraScribe™ T7 Transcription Kit (Epicentre Technologies, Madison, WI, USA) did not require transfection reagents for delivery into cells, even in the presence of serum. Kakiuchi *et al.* (*J. Biol. Chem.* 257:1924-1928, 1982), included herein by reference, showed that use of [(2'-F-dI)_n : (2'-F-dC)_n] duplexes were 40-100 times less antigenic than [(rI).sub_n : (rC_n.)] duplexes, and did not induce an interferon response like [(rI)_n : (rC)_n.] duplexes.

H. Use of Methods of the Present Invention for Making Double-Stranded RNA for RNA Interference

1. Combining Use of a ssDNA Transcription Substrate for N4 mini-vRNAP and a Substrate for dsDNA Substrate for a T7-Type RNAP to Synthesize Both RNA Strands of dsRNA for Use as RNAi

[00578] The methods of the present invention can also be combined with methods in the art for a particular purpose. By way of example, but not of limitation, both N4 vRNAP-type promoters and T7 RNAP-type promoters can be combined for a method to make double-stranded RNA for use as RNAi or siRNA. Thus, in one embodiment, a primer comprising a T7 RNAP or other T7-type protopromoter at its 5'-end and a sequence that is complementary to the 3'-end of a target sequence can be used for reverse transcription or primer extension to synthesize first-strand cDNA that is complementary to a target sequence in a target nucleic acid. Then, the 3'-end of the first-strand cDNA can be primer-extended using a promoter splice template oligo, the promoter splice template oligo comprising a sequence at its 3'-end that is complementary to the 3'-end of the first-strand cDNA and a sequence at its 5'-end that is complementary to a sense promoter for an N4 vRNAP promoter. The resulting product has a sense N4 vRNAP promoter on its 3'-end and an anti-sense T7 RNAP protopromoter sequence on its 5'-end. Thus, an N4 mini-vRNAP can be used to synthesize is sense-strand RNA from this single-stranded first-strand cDNA template corresponding to the target sequence. Then, the first-strand cDNA can be made double stranded by primer extension using a primer complementary to the 3'-end of the first-strand cDNA (e.g., complementary to the N4 vRNAP promoter) and first-strand cDNA as a template. The resulting double-stranded cDNA can be used for transcription of RNA that is anti-sense with respect to the target sequence using T7 RNAP. The sense and anti-sense modified RNA strands can be annealed to produce double-stranded RNA that can be used for RNA interference, including, but not limited to, as siRNA.

In other embodiments of the invention, N4 mini-vRNAP Y678F and T7 RNAP Y639F (Sousa et al., U.S. Patent No. 5,849,546, incorporated herein by reference) mutant enzymes can be used in order to synthesize, respectively, sense and anti-sense modified RNA that contains 2'-fluoro-pyrimidine nucleotides in place of the canonical pyrimidine nucleotides.

2. Methods for Making Transcription Products for Use as RNAi by Obtaining a ssDNA Transcription Substrate Comprising a Hairpin Target Sequence

[00579] The present invention is not limited to this particular method for making dsRNA for use as RNAi. Another method of the invention uses a target sequence comprising a hairpin in order to synthesize a ssDNA transcription substrate that encodes both strands of a dsRNA and a non-complementary region. Transcription of the transcription substrate using an N4 vRNAP or another RNAP that recognizes a single-stranded pseudopromoter, as described herein, yields a hairpin RNA structure for use as RNAi. According to this method, a sense transcription promoter for an N4 vRNAP or a sense single-stranded pseudopromoter is attached at the 3'-end of a DNA target sequence using a promoter splice template, a promoter ligation oligo or a promoter primer, as described elsewhere herein, in order to make a ssDNA transcription substrate. The target sequence in the ssDNA transcription substrate comprises: (a) a 3'-portion, wherein the 3'-portion encodes a sequence for a first strand of a dsRNA that is to be used as RNAi; (b) a middle portion, wherein the middle portion comprises a sequence that is not complementary to either strand of the dsRNA; and (c) a 5'-portion, wherein the 5'-portion encodes the sequence for a second strand of the dsRNA that is to be used as RNAi and wherein the RNA encoded by the 5'-portion that encodes the second strand is complementary to and can anneal to the RNA encoded by the 3'-portion that encodes the first strand so as to form a hairpin structure wherein the middle portion is a loop structure of the hairpin. This ssDNA transcription substrate is used for *in vitro* transcription by an RNA polymerase that can transcribe the ssDNA transcription substrate. Suitable RNA polymerases comprise N4 vRNAPs, such as mini-vRNAP or mini-vRNAP Y678F, for transcription substrates having an N4 vRNAP promoter. In other embodiments that use a transcription substrate comprising a single-stranded pseudopromoter, an RNAP that can bind to and synthesize RNA using the pseudopromoter can be used in order to synthesize a dsRNA hairpin that can be used as RNAi. In still other embodiments, a molecule that is identical to a ssDNA transcription substrate of the invention for synthesis of hairpin RNA for use as RNAi can be obtained by chemical synthesis using an oligonucleotide synthesizer using methods known in the art. The chemically synthesized ssDNA can be used as a substrate for *in vitro* transcription using an RNA polymerase of the present invention.

[00580] In some embodiments that use a hairpin substrate for *in vitro* transcription, 2'-fluoro-pyrimidine deoxynucleotides are used in place of canonical pyrimidine nucleotides when the mini-vRNAP Y678F (Kazmierczak, K.M., *et al.*, *EMBO J.* 21:5815-5823, 2002, incorporated herein by reference) or a mutant T7-type RNAP, such as the T7 RNAP Y639F mutant enzyme (Sousa *et al.*, U.S. Patent No. 5,849,546, incorporated herein by reference), is used to synthesize dsRNA for use as RNAi.

I. Kits and Compositions of the Invention

[00581] The present invention also comprises kits and compositions for carrying out the methods of the invention. A kit of the invention comprises one or, preferably, multiple components for carrying out the various processes of a method. With respect to embodiments of the invention for making a transcription product corresponding to a target sequence of a target nucleic acid, a kit comprises an RNA polymerase of the invention that can synthesize RNA using a ssDNA transcription substrate and at least one oligonucleotide, and preferably multiple oligonucleotides, that encode a sequence for either a sense or an anti-sense transcription promoter of the invention, including a promoter splice template oligo, a promoter ligation oligo or a promoter primer for use in generating a transcription substrate of the invention. With respect to embodiments of the invention for obtaining host cells comprising replicable single-stranded clones or libraries of replicable single-stranded clones comprising target sequences of target nucleic acids in a sample, a kit comprises, at a minimum, at least one oligonucleotide, chosen from among a promoter splice template oligo, a splice template oligo, a promoter ligation oligo, a ligation oligo, or a promoter primer or primer, wherein the oligo comprises a single-stranded origin of replication (*ori*), such as, but not limited to an M13 *ori*, and a sequence that encodes a selectable or screenable marker, such as, but not limited to an antibiotic-resistance gene, or a beta-galactosidase gene, respectively, and a host cell that can replicate a circular ssDNA comprising the single-stranded origin of replication and in which the selectable or screenable marker is expressible. With respect to embodiments for obtaining host cells comprising replicable single-stranded expression clones or expression libraries comprising a target sequence of target nucleic acid, a kit of the invention comprises an oligonucleotide that comprises a promoter for an RNA polymerase of the invention that can synthesize RNA using a ssDNA transcription substrate and the host cell comprises an inducible gene for the RNA polymerase that recognizes the promoter.

[00582] Depending on the particular embodiment, a kit can also contain one or more other components. By way of example, but not of limitation, a kit can comprise one or more other oligos, including splice template oligos, ligation splint oligos, and/or primers, including, but not limited to strand-displacement primers. A kit of the invention can also contain one or more other enzymes, such as, but not limited to, a reverse transcriptase, a ligase, a DNA polymerase, which can be a strand displacing DNA polymerase, an RNaseH, and/or other enzymes for particular processes of the invention. Still further a kit can contain components or reagents for use with host cells, such as helper phages, inducing substances, or substances for making the cells competent for transformation.

[00583] A kit can be for a specific method, such as, but not limited to, a kit for the synthesis and cloning of full-length cDNA or cDNA libraries that correspond to the complete sequence of 5'-ends of mRNA molecules, a kit for generating templates for random amplification of cDNA ends (RACE), a kit for making a transcription product corresponding to mRNA from a particular type of cell or different cells, whether under the same conditions or different environmental conditions, for gene expression profiling, including generation of sense or anti-sense probes for microarrays, a kit for amplification of an RNA or DNA sequence, including a kit for performing multiple rounds of transcription of a target nucleic acid sequence in a single reaction mixture, a kit for making a transcription product corresponding to a particular target nucleic acid sequence, if present in a sample, that is diagnostic or indicative of a pathogen, a disease gene, a mutated allele, or the like, a kit for an analyte-specific assay, wherein the analyte is a nucleic acid, a kit for amplifying a target nucleic acid by strand displacement replication or rolling circle replication and/or transcription of a ssDNA transcription substrate, a kit for making RNAi, including siRNA, or modified RNAi or siRNA, including, but not limited to, 2'-F-dCMP- and 2'-F-dUMP-containing RNAi or siRNA, or any of a broad range of kits that will be understood by those with knowledge in the art by a reading of the description of the invention herein.

[00584] In general, a kit of the invention will also comprise a description of the components of the kit and instructions for their use in a particular process or method or methods of the invention. In general, a kit of the present invention will also comprise other components, such as, but not limited to, buffers, ribonucleotides and/or deoxynucleotides, including modified nucleotides in some embodiments, DNA polymerization or reverse transcriptase enhancers, such as, but not limited to betaine (trimethylglycine), and salts of monvalent or divalent cations, such as but not limited to potassium acetate or chloride and/or magnesium chloride, enzyme

substrates and/or cofactors, such as, but not limited to, ATP or NAD, and the like which are needed for optimal conditions of one or more reactions or processes of a method or a combination of methods for a particular application. A kit of the invention can comprise a set of individual reagents for a particular process or a series of sets of individual reagents for multiple processes of a method that are performed in a stepwise or serial manner, or a kit can comprise a multiple reagents combined into a single reaction mixture or a small number of mixtures of multiple reagents, each of which perform multiple reactions and/or processes in a single tube. In general, the various components of a kit for performing a particular process of a method of the invention or a complete method of the invention will be optimized so that they have appropriate amounts of reagents and conditions to work together in the process and/or method.

[00585] Components of a kit may be provided as solutions or as dried powder(s). When reagents and/or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent, in which case, the solvent may also be provided in another container.

[00586] As used herein in the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

IX. Use of RNA Polymerases of the Invention in Signaling Systems

[00587] With respect to this aspect of the invention, a "signaling system" means and comprises the multitude of substances, compositions and environmental conditions comprising a method for detecting the presence or quantity of an analyte in a sample by detecting a transcript that results from transcription of a ssDNA polynucleotide that is operably joined to an analyte-binding substance. An important composition of a method comprising a signaling system of the invention is an analyte-binding substance that is operably joined to a ssDNA polynucleotide, wherein the ssDNA polynucleotide comprises a sequence that encodes a detectable transcript, wherein the sequence is operably joined to a single-stranded promoter for an RNA polymerase that can bind the promoter of the ssDNA polynucleotide and transcribe the sequence thereof, thereby making the detectable transcript under transcription conditions.

[00588] Other methods for detecting an analyte using transcription are known in the art, including the methods described by Zhang *et al.* (*Proc. Natl. Acad. Sci. USA* 98: 5497-5502,

2001), by Eberwine in PCT Patent Application No. WO 02/14476, and by Hudson *et al.* in U.S. Patent No. 6,100,024, all of which are incorporated herein by reference. In contrast to the present invention, all of the methods of the prior art require, at a minimum, making a composition comprising a double-stranded promoter, and in most cases, the methods in the art also require preparation of double-stranded templates.

[00589] The signaling systems of the present invention differ from methods in the prior art in that they use a single-stranded DNA polynucleotide that is operably joined to an analyte-binding substance (ABS), wherein the single-stranded DNA polynucleotide comprises a single-stranded promoter that is operably joined to a single-stranded template that encodes a transcription product, and wherein the ABS is detected by making the transcription product using an RNA polymerase that binds to the single-stranded promoter comprising the single-stranded DNA polynucleotide.

[00590] The invention comprises methods, compositions and kits for using the RNA polymerases of the invention as a signaling system for an analyte of any type, including analytes such as, but not limited to, antigens, antibodies or other substances, in addition to an analyte that is a target nucleic acid.

[00591] Thus, the invention comprises a method for detecting an analyte in or from a sample, the method comprising:

[00592] 1. obtaining a transcription signaling system, the transcription signaling system comprising a ssDNA comprising: (a) a sequence for a promoter for an RNA polymerase chosen from among N4 vRNAP, N4 mini-vRNAP, and N4 mini-vRNAP Y678F enzymes or another enzyme that binds a single-stranded promoter and transcribes a single-stranded template to which it is operably joined; and (b) a signal sequence that is operably joined to the promoter sequence, wherein the signal sequence, when transcribed by the RNA polymerase, is detectable in some manner; and

[00593] 2. joining the transcription signaling system, either covalently or non-covalently, to an analyte-binding substance, wherein the joining to the analyte-binding substance is not affected by the conditions of the assay and wherein the joining to the analyte-binding substance does not affect the ability of the transcription signaling system to be transcribed using the RNA polymerase under transcription conditions;

[00594] 3. contacting the analyte-binding substance to which the transcription signaling system is joined with a sample under binding conditions, wherein the analyte, if present in the sample, binds to the analyte-binding substance so as to form a specific binding pair; and

[00595] 4. removing the specific binding pair from the sample so as to separate it from other components in the sample; and

[00596] 5. incubating the specific-binding pair under transcription conditions with an RNA polymerase, wherein the RNA polymerase can synthesize a transcription product that is complementary to the signal sequence in the ssDNA transcription signaling system under the transcription conditions;

[00597] 6. obtaining a transcription product that is complementary to the signal sequence in the ssDNA transcription signaling system; and

[00598] 7. detecting the transcription product or a substance that results from the transcription product.

[00599] This method of the invention can be used for a broad range of analytes and analyte-binding substances. By way of example, but not of limitation, the analyte can be an antigen and the analyte-binding substance can be antibody, or the analyte can be a nucleic acid and the analyte-binding substance can be another complementary nucleic acid. As discussed below, a large number of other substances exist for which a specific-binding pair can be found, all of which are within the scope of the invention.

[00600] In order to detect an analyte in a sample, an assay of this aspect of the invention uses an analyte-binding substance that "binds" to the analyte under "binding conditions." The analyte-binding substance, which can also be referred to as an "affinity molecule," an "affinity substance," a "specific binding substance," or a "binding molecule" for the analyte, is in turn detected by making a transcription product using a transcription signaling system that is joined to the analyte-binding substance. The transcription signaling system comprises a single-stranded promoter that is operably joined to a single-stranded signal sequence, wherein transcription of the signal sequence by an RNA polymerase that can use the single-stranded signaling system results in a transcription product that is detectable by some means, as discussed below.

[00601] 1. Analytes and Analyte-Binding Substances for Assays in which an RNA Polymerase that Recognizes a Single-Stranded Promoter Is Used as a Signaling System

[00602] An "analyte" or an "analyte-binding substance" of this aspect of the invention can be any of those described in U.S. Patent No. 6,562,575 of one of the present inventors, which is entitled "Analyte-Specific Assays Based on Formation of a Replicase Substrate," and which is incorporated herein in its entirety by reference so as to be made part of the present invention. An analyte can be any substance whose presence, concentration or amount in a

sample is determined in an assay. By way of example, but not of limitation, an analyte can be a biochemical molecule or a biopolymer or a segment of a biopolymer, such as a protein or peptide, including a glycoprotein or lipoprotein, an enzyme, hormone, receptor, antigen or antibody, nucleic acid (DNA or RNA), polysaccharide, or lipid.

[00603] An analyte-binding substance that is a nucleic acid, polynucleotide, oligonucleotide or a segment of a nucleic acid or polynucleotide, including nucleic acids composed of either DNA or RNA, or both DNA and RNA mononucleosides, including modified DNA or RNA mononucleosides, can also be used according to the invention to detect an analyte that does not comprise nucleic acid. For example, a method termed "SELEX," as described by Gold and Tuerk in U.S. Pat. No. 5,270,163, can be used to select a nucleic acid for use as an analyte-binding substance according to the invention. SELEX permits selection of a nucleic acid molecule that has high affinity for a specific analyte from a large population of nucleic acid molecules, at least a portion of which have a randomized sequence. For example, a population of all possible randomized 25-mer oligonucleotides (i.e., having each of four possible nucleic acid bases at every position) will contain 4^{25} (or 10^{15}) different nucleic acid molecules, each of which has a different three-dimensional structure and different analyte binding properties. SELEX can be used, according to the methods described in U.S. Pat. Nos. 5,270,163; 5,567,588; 5,580,737; 5,587,468; 5,683,867; 5,696,249; 5,723,594; 5,773,598; 5,817,785; 5,861,254; 5,958,691; 5,998,142; 6,001,577; 6,013,443; and 6,030,776, incorporated herein by reference, in order to select an analyte-binding nucleic acid with high affinity for a specific analyte that is not a nucleic acid or polynucleotide. Once selected using SELEX, nucleic acid affinity molecules can be made by any of numerous known *in vivo* or *in vitro* techniques, including, by way of example, but not of limitation, automated nucleic acid synthesis techniques, PCR, or *in vitro* transcription.

[00604] Naturally occurring nucleic acid or polynucleotide sequences that have affinity for other naturally occurring molecules such as, but not limited to, protein molecules, are also known in the art. Examples include, but are not limited to certain nucleic acid sequences such as operators, promoters, origins of replication, sequences recognized by steroid hormone-receptor complexes, restriction endonuclease recognition sequences, ribosomal nucleic acids, and so on, which are known to bind tightly to certain proteins. For example, in two well-known systems, the lac repressor and the bacteriophage lambda repressor each bind to their respective specific nucleic acid sequences called "operators" to block initiation of transcription of their corresponding mRNA molecules. Nucleic acids containing such specific sequences can be used

in the invention as analyte-binding substances for the respective proteins or other molecules for which the nucleic acid has affinity. In these cases, the nucleic acid with the specific sequence can be used according to this aspect of the invention as the analyte-binding substance for the respective specific protein, glycoprotein, lipoprotein, small molecule or other analyte that it binds. One of several techniques which are generally called "footprinting" (e.g., see Galas, D. and Schmitz, A, Nucleic Acids Res. 5:3161, 1978) can be used to identify sequences of nucleic acids which bind to a protein. Other methods are also known to those with skill in the art and can be used to identify nucleic acid sequences for use as specific analyte-binding substances for use in the invention.

[00605] A peptide nucleic acid (PNA) or a molecule comprising both a nucleic acid and a PNA can also be used according to the invention as an analyte-binding substance for an analyte that is a nucleic acid or polynucleotide. PNA as an analyte-binding substance of the invention provides tighter binding (and greater binding stability) in assays for a nucleic acid analyte (e.g., see U.S. Pat. No. 5,985,563). Also, since PNA is not naturally occurring, PNA molecules are highly resistant to protease and nuclease activity. Antibodies to PNA/DNA or PNA/RNA complexes can be used in the invention for capture, recognition, detection, identification, or quantitation of nucleic acids in biological samples, via their ability to bind specifically to the respective complexes without binding the individual molecules (U.S. Pat. No. 5,612,458).

[00606] The invention also contemplates that a combinatorial library of randomized peptide nucleic acids prepared by a method such as, but not limited to, the methods described in U.S. Pat. Nos. 5,539,083; 5,831,014; and 5,864,010, can be used to prepare analyte-binding substances for use in assays for analytes of all types, including analytes that are nucleic acids, proteins, or other analytes, without limit. As is the case for the SELEX method with nucleic acids, randomized peptide or peptide nucleic acid libraries are made to contain molecules with a very large number of different binding affinities for an analyte. After selection of an appropriate affinity molecule for an analyte from a library, the selected affinity molecule can be used in the invention as an analyte-binding substance.

[00607] An analyte-binding substance can also be an oligonucleotide or polynucleotide with a modified backbone that is not an amino acid, such as, but not limited to modified oligonucleotides described in U.S. Pat. Nos. 5,602,240; 6,610,289; 5,696,253; or 6,013,785.

[00608] The invention also contemplates that an analyte-binding substance can be prepared from a combinatorial library of randomized peptides (i.e., comprising at least four naturally-occurring amino acids). One way to prepare the randomized peptide library is to place

a randomized DNA sequence, prepared as for SELEX, downstream of a phage T7 RNA polymerase promoter, or a similar promoter, and then use a method such as, but not limited to, coupled transcription-translation, as described in U.S. Pat. Nos. 5,324,637; 5,492,817; or 5,665,563, or stepwise transcription, followed by translation. Alternatively, a randomized DNA sequence, prepared as for SELEX, can be cloned into a site in a DNA vector that, once inserted, encodes a recombinant MDV-1 RNA containing the randomized sequence that is replicatable by Q-beta replicase (e.g., between nucleotides 63 and 64 in MDV-1 (+) RNA; see U.S. Pat. No. 5,620,870). The recombinant MDV-1 DNA containing the randomized DNA sequence is downstream from a T7 RNA polymerase promoter or a similar promoter in the DNA vector. Then, following transcription, the recombinant MDV-1 RNA, containing the randomized sequence can be used to make a randomized peptide library comprising at least four naturally-occurring amino acids by coupled replication-translation as described in U.S. Pat. No. 5,556,769. An analyte-binding substance can be selected from the library by binding peptides in the library to an analyte, separating the unbound peptides, and identifying one or more peptides that is bound to analyte by means known in the art. Alternatively, high throughput screening methods can be used to screen all individual peptides in the library to identify those which can be used as analyte-binding substances. Although the identification of an analyte-binding peptide by these methods is difficult and tedious, the methods in the art are improving for doing so, and the expenditure of time and effort required may be warranted for identifying analyte-binding substances for use in assays of the invention that will be used routinely in large numbers.

[00609] A variety of other analyte-binding substances can also be used in methods for this aspect of the invention.

[00610] For an antigen analyte (which itself may be an antibody), antibodies, including monoclonal antibodies, are available as analyte-binding substances. For certain antibody analytes in samples which include only one antibody, an antibody binding protein such as *Staphylococcus aureus* Protein A can be employed as an analyte-binding substance.

[00611] For an analyte, such as a glycoprotein or class of glycoproteins, or a polysaccharide or class of polysaccharides, which is distinguished from other substances in a sample by having a carbohydrate moiety which is bound specifically by a lectin, a suitable analyte-binding substance is the lectin.

[00612] For an analyte which is a hormone, a receptor for the hormone can be employed as an analyte-binding substance. Conversely, for an analyte which is a receptor for a hormone, the hormone can be employed as the analyte-binding substance.

[00613] For an analyte which is an enzyme, an inhibitor of the enzyme can be employed as an analyte-binding substance. For an analyte which is an inhibitor of an enzyme, the enzyme can be employed as the analyte-binding substance.

[00614] Usually, an analyte molecule and an affinity molecule for the analyte molecule are related as a specific "binding pair", i.e., their interaction is only through non-covalent bonds such as hydrogen-bonding, hydrophobic interactions (including stacking of aromatic molecules), van der Waals forces, and salt bridges. Without being bound by theory, it is believed in the art that these kinds of non-covalent bonds result in binding, in part due to complementary shapes or structures of the molecules involved in the binding pair.

[00615] The term "binding" according to this aspect of the invention refers to the interaction between an analyte-binding substance or affinity molecule and an analyte as a result of non-covalent bonds, such as, but not limited to, hydrogen bonds, hydrophobic interactions, van der Waals bonds, and ionic bonds.

[00616] Based on the definition for "binding," and the wide variety of affinity molecules and analytes which can be used in the invention, it is clear that "binding conditions" vary for different specific binding pairs. Those skilled in the art can easily determine conditions whereby, in a sample, binding occurs between affinity molecule and analyte that may be present. In particular, those skilled in the art can easily determine conditions whereby binding between affinity molecule and analyte, which would be considered in the art to be "specific binding," can be made to occur. As understood in the art, such specificity is usually due to the higher affinity of affinity molecule for analyte than for other substances and components (e.g., vessel walls, solid supports) in a sample. In certain cases, the specificity might also involve, or might be due to, a significantly more rapid association of affinity molecule with analyte than with other substances and components in a sample.

[00617] "Hybridization" is the term used to refer to the process of incubating an affinity molecule comprising a nucleic acid, or a peptide nucleic acid (PNA) molecule, or a covalently linked, joined or attached nucleic acid-PNA molecule with an analyte comprising a nucleic acid under "binding conditions," which are also called "hybridization conditions." The ability of two polymers of nucleic acid containing complementary sequences to find each other and anneal through base pairing interaction is a well-recognized phenomenon. The initial observations of the "hybridization" process by Marmur and Lane (Proc. Nat. Acad. Sci. USA 46:453, 1960) and Doty, et al. (Proc. Nat. Acad. Sci. USA 46:461, 1960) have been followed by the refinement of this process into an essential tool of modern biology. "Hybridization" also refers to the

"binding" or "pairing" of complementary nucleic acid bases in a single-stranded nucleic acid, PNA, or linked nucleic acid-PNA affinity molecule with a single-stranded nucleic acid analyte, which occurs according to base pairing rules (e.g., adenine pairs with thymine or uracil and guanine pairs with cytosine). Those with skill in the art will be able to develop and make conditions which comprise binding conditions or hybridization conditions for particular nucleic acid analytes of an assay. In developing and making binding conditions for particular nucleic acid analytes analyte-binding substances, as well as in developing and making hybridization conditions for particular analytes and capture probes, certain additives can be added in the hybridization solution. By way of example, but not of limitation, dextran sulfate or polyethylene glycol can be added to accelerate the rate of hybridization (e.g., Chapter 9, Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, 1989), or betaine can be added to the hybridization solution to eliminate the dependence of T_m on basepair composition (Rees, W. A., *et al.*, *Biochemistry* 32:137-144, 1993).

[00618] The terms "degree of homology" or "degree of complementarity" are used to refer to the extent or frequency at which the nucleic acid bases on one strand (e.g., of the affinity molecule) are "complementary with" or "able to pair" with the nucleic acid bases on the other strand (e.g., the analyte). Complementarity may be "partial," meaning only some of the nucleic acid bases are matched according to base pairing rules, or complementarity may be "complete" or "total." The length (i.e., the number of nucleic acid bases comprising the nucleic acid and/or PNA affinity molecule and the nucleic acid analyte), and the degree of "homology" or "complementarity" between the affinity molecule and the analyte have significant effects on the efficiency and strength of binding or hybridization when the nucleic acid bases on the affinity molecule are maximally "bound" or "hybridized" to the nucleic acid bases on the analyte. The terms "melting temperature" or " T_m " are used as an indication of the degree of complementarity. The T_m is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands under defined conditions. Based on the assumption that a nucleic acid molecule that is used in hybridization will be approximately completely homologous or complementary to a target polynucleotide, equations have been developed for estimating the T_m for a given single-stranded sequence that is hybridized or "annealed" to a complementary sequence. For example, a common equation used in the art for oligodeoxynucleotides is: $T_m = 81.5^\circ \text{C} + 0.41 (\% \text{G+C})$ when the nucleic acid is in an aqueous solution containing 1 M NaCl (see e.g., Anderson and Young, *Quantitative Filter Hybridization*, in *Nucleic Acid Hybridization*, 1985). Other more sophisticated equations available for nucleic

acids take nearest neighbor and other structural effects into account for calculation of the T_m . Binding is generally stronger for PNA affinity molecules than for nucleic acid affinity molecules. For example the T_m of 10-mer homothymidine PNA binding to its complementary 10-mer homoadenosine DNA is 73° C, whereas the T_m for the corresponding 10-mer homothymidine DNA to the same complementary 10-mer homoadenosine DNA is only 23° C. Equations for calculating the T_m for a nucleic acid are not appropriate for PNA. Preferably, a T_m that is calculated using an equation in the art, is checked empirically and the hybridization or binding conditions are adjusted by empirically raising or lowering the stringency of hybridization as appropriate for a particular assay. As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds, under which nucleic acid hybridizations are conducted. With "high stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of complementary base sequences. Thus, conditions of "weak" or "low" stringency are often required when it is desired that nucleic acids that are not completely complementary to one another be hybridized or annealed together.

[00619] With regard to complementarity, it is important for some assays of the invention to determine whether the hybridization represents complete or partial complementarity. For example, where it is desired to detect simply the presence or absence of pathogen DNA (such as from a virus, bacterium, fungi, mycoplasma, protozoan), it is only important that the hybridization method ensures hybridization when the relevant sequence is present. In those embodiments of the invention, conditions can be selected where both partially complementary probes and completely complementary probes will hybridize.

2. Signal Sequences for Use in Signaling Systems

[00620] The methods for using RNA polymerases of the invention as a signaling system are also not limited to the signal sequences, which can vary greatly. By way of example, but not of limitation, a signal sequence can comprise a substrate for Q-beta replicase, which is detectable in the presence of the replicase under replication conditions. It can also comprise a sequence that encodes a protein, such as green fluorescent protein, that is detectable following translation of the signal sequence. Without limitation, it can also comprise a sequence that is detectable by a probe, such as, but not limited to a molecular beacon, as described by Tyagi *et al.* (U.S. Patents Nos. 5,925,517 and 6,103,476 of Tyagi *et al.* and 6,461,817 of Alland *et al.*, all of which are incorporated herein by reference).

IX. EXAMPLES

[00621] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Identification of a Transcriptionally Active Domain of N4 Virion RNA Polymerase

[00622] To determine the minimal domain possessing RNA polymerase activity, controlled proteolysis was performed followed by catalytic (transcriptional) autolabeling (Hartmann, *et al.*, 1988). Upon incubation of RNA polymerase with a benzaldehyde derivative of the initiating nucleotide, the benzaldehyde group forms a Schiff-base with the ϵ -amino group of lysines located within 12 Å of the nucleotide-binding site. The crosslinking step was performed in the presence of DNA template because it stimulates binding of the initiating nucleotide. The unstable Schiff-base is converted to a stable secondary amine by reduction under mild conditions with sodium borohydride, with concomitant reduction of any non-reacted benzaldehyde derivative. Addition of the next template-directed α - ^{32}P labeled NTP leads to phosphodiester bond formation and catalytic autolabeling of the transcriptionally active polypeptide. Controlled trypsin proteolysis of vRNAP was performed, followed by catalytic autolabeling and analysis on SDS-PAGE (FIG. 3A). Initially, three proteolytic fragments are generated, of which the smaller two are catalytically active. Upon further incubation with trypsin, a single stable, transcriptionally active product approximately 1,100 amino acids in length remains. N-terminal sequencing of the three initial proteolytic fragments (FIG. 3B) indicated that the stable active polypeptide (mini-vRNAP) corresponds to the middle 1/3 of vRNAP, the region containing the three motifs described above (FIG. 2A, SEQ ID NOS:3-4).

Example 2

Cloning and Purification of N4 mini-vRNAP

[00623] The full-size vRNAP and the mini-vRNAP (SEQ ID NOS:6 and 15) ORFs were cloned under pBAD control with an N-terminal hexahistidine tag (FIG. 4). The mini-vRNAP

domain was cloned into the pBAD B expression plasmid, which was purchased from Invitrogen. Five restriction enzyme sites within pBAD B have been altered; the SnaI site was converted to a HpaI site, and the PflMI and EcoRV sites were destroyed, all by site-directed mutagenesis. The BstBI and HindIII sites were destroyed by enzyme digestion followed by Klenow treatment and re-ligation. FIG. 5 (left) shows the relative amounts of full-length and mini-vRNAP proteins purified on TALON columns from the same volume of *E. coli* BL21 induced cells. Cloned mini-vRNAP is expressed at 100-fold higher levels than cloned full size vRNAP. Further concentration on a MonoQ column reveals that, in contrast to full size vRNAP, mini-vRNAP is stable after induction (FIG. 5, right). At least 10 mg of mini-vRNAP at a 20 mg/ml concentration are obtained from 1 L of induced cells in just two purification steps: TALON and MonoQ minicolumns. A non-histagged version of mini-vRNAP has also been cloned (SEQ ID NO:4). In this case, the enzyme is purified from a crude extract of induced cells in two steps: a promoter DNA-affinity column and MonoQ.

[00624] Mini-vRNAP possesses a high binding affinity ($K_d=1$ nM) for N4 promoter-containing DNA oligonucleotides. This property was used for purification of non-his tagged mini-vRNAP (SEQ ID NO:4) on a DNA-affinity column. The column was prepared by adsorbing a 5' biotinylated N4 promoter-containing DNA oligonucleotide onto the matrix of a 1 ml HiTrap Streptavidin column (Pharmacia/Amersham Cat.#17-5112-01) according to the manufacturer's instructions. A debris-free sonicate of bacterial cells expressing mini-vRNAP was passed through the column. To bind mini vRNAP to the DNA-affinity column, the pH in the extract and binding/washing buffer should be between 5 to 9, and the NaCl concentration should be between 50 mM and 2M. Nucleases in the extract are inhibited by addition of 2 mM EDTA. After washing the column, mini-vRNAP was eluted with warm (25° C) water; the elution temperature was raised from 4° C to 25° C to increase mini-vRNAP recovery. For complete elution, the temperature can be raised up to 43° C without significant change in the quality of the preparation. Elution under these conditions occurs due to the removal of metal ions and consequent melting of the promoter hairpin and dissociation of mini-vRNAP. Different DNA oligonucleotides containing variants of the P2 promoter (SEQ ID NOS:16-19), were used in DNA-affinity columns and tested in mini-vRNAP affinity purification. The best yield was achieved using the DNA oligonucleotide of SEQ ID NO:16. However, the DNA oligonucleotides of SEQ ID NOS:19-20 require a lower temperature than the DNA oligonucleotide of SEQ ID NO:16 for complete elution of the protein, in agreement with the lower thermal stability of the respective promoter hairpins.

[00625] Up to 1 mg of mini-vRNAP of 90% purity is obtained from a crude extract of 100 ml *E. coli* culture expressing mini-vRNAP in a single purification step using a 1 ml DNA-affinity column. The binding capacity of the DNA-affinity column was not detectably decreased by multiple use.

Example 3

Effect of EcoSSB on Transcription of Single-Stranded Templates

[00626] Inventors have previously shown that EcoSSB is required for N4 vRNAP transcription *in vivo* (Glucksmann, *et al.*, Cell 70:491-500, 1992). EcoSSB is unique in that, unlike other SSBs whose effect on vRNAP transcription was tested, it does not melt the promoter hairpin structure (Glucksmann-Kuis, *et al.*, Cell 84:147-154, 1996). Recently, inventors have reinvestigated the effect of EcoSSB on vRNAP transcription of single-stranded templates. FIG. 6 shows transcription in the absence and presence of Eco SSB at three different ssDNA template concentrations. The extent of EcoSSB activation is template-concentration dependent, with highest activation at low DNA template concentration. These results suggest that EcoSSB overcomes template limitation on ssDNA templates.

[00627] To further explore this hypothesis, the effect of addition of template or EcoSSB to transcription reactions after 20 min incubation in the absence of EcoSSB was tested. The transcription reaction mixtures (5-50 μ l) contained 20 mM Tris-HCl (pH 7.9 at 25° C), 10 mM MgCl₂, 50 mM NaCl, 1 mM dithiothreitol, 0.01-1 μ M mini-vRNAP, 1-100 nM ssDNA template (30-100 nt long, synthesized by Integrated DNA Technologies), 1 mM each of 3 non-labeled NTPs, 0.1 mM α -³²P NTP (1-2 Ci/mmol, NEN), and 1-10 μ M *E. coli* SSB. Incubation was for 1 to 80 min at 37° C at the indicated temperature. In the presence of EcoSSB, RNA synthesis increased linearly throughout the period of incubation (FIG. 7C). In the absence of EcoSSB, no increase in transcription was observed beyond 10 min of incubation (FIG. 7A). Addition of template at 20 min to the reaction carried out in the absence of EcoSSB led to a dramatic increase in RNA synthesis (FIG. 7B). Addition of EcoSSB at 20 min led to a slow rate of transcriptional recovery (FIG. 7D). These results suggest that EcoSSB converts the template from a transcriptionally inactive RNA:DNA hybrid to transcriptionally active single-stranded DNA.

[00628] To test this hypothesis, the physical states of the DNA template and the RNA product were analyzed by native gel electrophoresis in the absence and in the presence of EcoSSB. In order to have effective transcription in the absence of EcoSSB, transcription was

performed at an intermediate (5 nM) DNA concentration, at which only a 2-fold effect of EcoSSB is observed.

[00629] The results of this experiment are shown in FIG. 8. Either ^{32}P -labeled template (right panel) or labeled NTPs (left panel) were used to analyze the state of the template (right panel) or RNA product (left panel) in the absence or presence of EcoSSB. After transcription, the mixtures were split further into 3 samples: a control sample with no additions, a sample to which RNase H was added to specifically degrade RNA in RNA:DNA hybrids, and a third sample to which Nuclease S1 was added to degrade single-stranded nucleic acids. In the absence of EcoSSB, both the DNA template and the RNA product are in RNA:DNA hybrids, since the RNA product is RNase H sensitive while the DNA-containing bands show altered mobility after RNase H treatment. In the presence of EcoSSB, a significant portion of the RNA product is RNase H resistant and therefore free, although an RNase sensitive band is present that corresponds to an intermediate RNA:DNA:SSB complex. Under these conditions, the DNA is in an SSB:DNA complex. These results indicate that EcoSSB stimulates transcription through template recycling.

[00630] To define regions of EcoSSB essential for vRNAP transcription activation on single-stranded templates, the inventors have tested the effect of human mitochondrial SSB (HmtSSB), which shows extensive sequence and structural homology to EcoSSB. The N-terminus of EcoSSB contains DNA binding and tetramerization determinants while the C-terminus is involved in interaction with other replication proteins. Hmt SSB has no effect on vRNAP transcription although it does not melt the promoter hairpin. Interestingly, preliminary results using mutant EcoSSBs and EcoSSB-Hmt SSB chimeras suggest that the C-terminal region of EcoSSB is essential for vRNAP transcriptional activation.

Example 4

Characterization of mini-vRNAP Transcription Properties

[00631] The initiation properties of the full length RNA polymerase and mini-vRNAP were compared at similar molar concentrations (FIG. 9A) using the catalytic autolabeling assay and two reaction conditions: 1- using a template containing +1C, the benzaldehyde derivative of GTP and $\alpha^{32}\text{P}$ -ATP, or 2- a template containing +1T, the benzaldehyde derivative of ATP and $\alpha^{32}\text{P}$ -GTP. Comparison of the results in FIGS. 9B and 9C demonstrates that mini-vRNAP exhibits initiation properties similar to full-length vRNAP. In addition, both enzymes discriminate against dATP incorporation to the same extent. Mini-vRNAP does not synthesize

abortive products when the first four nucleotides of the transcript are comprised of 50% or more G or C nucleotides.

[00632] The elongation and termination properties of both enzymes are compared in FIG. 10. Similar run-off and terminated transcripts are synthesized. Moreover, EcoSSB activates transcription by both enzymes to the same levels. This result indicates that, if there are any sites of specific contact between vRNAP and EcoSSB, they reside in the mini-vRNAP domain.

[00633] The sequence of the terminator signals for vRNAP present in the N4 genome include SEQ ID NOS:21-26. The signals of SEQ ID NO:21 and 22 have been tested *in vitro* on single-stranded templates.

[00634] The rate of mini-vRNAP transcription has been compared to the rate of T7 RNA polymerase under the same conditions using the same DNA template. The template used was linearized pET11 containing the original T7 promoter and the N4 vRNAP P2 promoter that was introduced through cloning. The DNA template was denatured before performing transcription using N4 mini-vRNAP. The concentrations of T7 RNAP (Promega, Cat.#P2075) and mini-vRNAP were compared using SDS-PAGE. Transcription reactions contained 50 nM of polymerase, 100 nM of DNA template, 5 x transcription buffer provided with the T7 RNAP, and 1 mM of each ATP, GTP and CTP and 0.1 mM of [³²P]-UTP (1 Ci/mmol). Each reaction mixture was split in two, and *E. coli* SSB was added to one half. The mixtures were incubated at 37° C and aliquots were taken at different time points. Transcription products were electrophoresed on a 6% sequencing gel and the amount of radioactively-labeled RNA was quantitated by phosphorimaging. The results showed that: (a) transcription of T7 RNAP was not affected by the presence of *E. coli* SSB and (b) N4 mini-vRNAP synthesized 1.5 to 5 fold more RNA in the presence of EcoSSB than T7 RNAP at different time points of incubation.

[00635] The optimal temperature for mini-vRNAP transcription is 37° C It exhibits 70% activity at 30° C, 65% at 45° C, and only 20% at 50° C.

[00636] The average error frequency was estimated by determining the misincorporation frequency of each of four [³²P]-α NTPs into RNA products using template ssDNAs missing the corresponding template nucleotide in the transcribed region. The following values were obtained: $1/5 \times 10^4$ for misincorporation of G and U using "no C" (SEQ ID NO:10) and "no A" (SEQ ID NO:11) ssDNA templates, respectively; $1/4 \times 10^4$ for misincorporation of C using the "no G" (SEQ ID NO:12) template, and $1/2 \times 10^4$ for misincorporation of A using the "no T" (SEQ ID NO:13) template. For comparison, the average error frequency for T7 RNAP is $1/2 \times$

10⁴ (Huang, *et al.*, 2000). Using the method for detection of mispair formation described by Huang, *et al.* (2000), no misincorporation by mini-vRNAP was detected.

[00637] The ability of mini-vRNAP to incorporate derivatized nucleotides was measured. Transcription by mini-vRNAP in the presence of 0.1-1 mM Digoxigenin-11-UTP (cat# 1209256, Roche), Biotin-16-UTP (cat# 1388908, Roche) or underivatized UTP, yielded comparable amounts of product RNA using "control" ssDNA (SEQ ID NO:9) as a transcription template. The product RNAs synthesized in the presence of derivatized UTP have higher molecular mass than those synthesized in the presence of underivatized UTP, and the difference corresponds to the mass difference of the UTPs used. Several derivatives (i.e. 2'Fluoro-ribonucleoside triphosphates, dideoxynucleoside triphosphates) are being tested. The fluorescent analog Fluorescein-12-UTP (Roche catalog #1427857) has been tested using a template which encodes a 51 nucleotide transcript containing a run of 4 Us, and a nucleotide mix containing ATP, CTP, GTP and Fluorescein-12-UTP only. Transcription was only 3% of that achieved with UTP, biotin-6-UTP or digoxigenin-11-UTP under the same reaction conditions. However, incorporation of the fluorescent analog at higher yields is expected to occur in the presence of underivatized UTP or on templates with other sequence compositions.

Example 5

Sequence Determinants of mini-vRNAP Promoter Binding

[00638] The three N4 early promoters present in the N4 genome contain a pair of Cs separated by 4 nucleotides from the base of the 5 bp promoter stem. In the preferred promoter P2, these 4 bases are As and the Cs are followed by a T. Preferably, mini-vRNAP uses a 17 nucleotide promoter sequence located immediately upstream of the transcription initiation site. Promoters for N4 vRNA polymerase are described by Haynes *et al.*, Cell 41:597-605 (1985) and Dai *et al.*, Genes Develpmnt. 12:2782-2790 (1998), herein incorporated by reference. vRNAP-promoter recognition and activity require specific sequences and a hairpin structure on the template strand. The vRNAP promoters of SEQ ID NOS:27-29 assume a hairpin structure comprised of a 5-7 bp stem and 3 b purine-containing loop. The -11 position corresponds to the center of the loop; +1 indicates the transcription start site. Thus, promoter sequences of the invention include, but are not limited to:

[00639] -11 +1 P1 3'-CAACGAAGCGTTGAATACCT-5' SEQ ID NO:27

[00640] -11 +1 P2 3'-TTCTTCGAGGCGAAGAAAACCT-5' SEQ ID NO:28, and

[00641] -11 +1 P3 3'-CGACGAGGCGTCGAAAACCA-5' SEQ ID NO:29

[00642] Other possible vRNAP promoters of the current invention include a set of any inverted repeats forming a hairpin with a 2-7 bp long stem and 3-5 b loop having purines in the central and/or next to the central position of the loop.

[00643] To study the sequence determinants of promoter binding, 20 base-long promoter oligonucleotides, containing the wild-type vRNAP promoter P2 sequence and substituted at every position with a single 5-Iodo-dU, were used. Whenever substitutions were made in the stem, the corresponding pairing base was changed to A. These oligonucleotides were ³²P end-labeled and used to determine the enzyme's affinity for promoter DNAs by a filter binding assay and the ability to crosslink to mini-vRNAP upon UV irradiation at 320 nm. A 20-base oligonucleotide with wild type promoter P2 sequence binds with a 1 nM Kd. Most oligonucleotides showed close to wild type affinity except for the oligonucleotides substituted at positions -11 (at the center of the loop) and -8, indicating that these positions are essential for promoter recognition (FIG. 11). Surprisingly, UV crosslinking was most effective at position -11, in spite of the low binding affinity, indicating a specific contact at this position to mini-vRNAP. Crosslinking was also observed to positions +1, +2 and +3, indicating non-specific contacts with this region of the template, since 5-Iodo-dU substituted oligonucleotides at these positions showed wild-type binding affinity.

[00644] The effect of changes in the stem length of the hairpin on the ability of mini-vRNAP to bind P2 promoter DNA was analyzed. As shown above, wild type promoter P2 with a 5 bp stem has a Kd of 1 nM (FIG. 12, top). The stem was shortened by removal of 3' bases as shown in FIG. 12 (left). The stem can be shortened by two base pairs without change in the binding affinity. If two or one loop-closing base pairs remain, the binding affinity of templates is still substantial (2-10 nM). This result, although surprising, is not unexpected since it has been shown that the oligonucleotide 3'd(CGAGGCG)5' forms an unusually stable minihairpin (Yoshizawa, *et al.*, Biochemistry 36, 4761-4767, 1997). No binding is observed if one more nucleotide is removed and the loop cannot form. These results indicate that formation of a loop is essential for vRNAP-promoter recognition.

[00645] The effect of lengthening the stem by addition of 3' bases is shown in FIG. 12 (right). The stem can be lengthened by two base pairs without change in the binding affinity. On the other hand, base pairing at -2 reduces binding affinity by two orders of magnitude, with a further one order of magnitude reduction caused by base pairing at -1 and +1. These results indicate that single-strandedness of the template at positions -2, -1 and +1 is required for efficient template binding.

[00646] All three N4 early promoters present in the N4 genome contain a pair of Cs separated by 4 nucleotides from the base of the 5 bp promoter stem. In promoter P2, these 4 bases are As and the Cs are followed by a T. To identify the determinants of the site of transcription initiation, a series of templates were constructed with a single C placed at different distances from position -11 of the hairpin by addition or deletion of the tract of As present at promoter P2 (FIG. 13). The affinity of mini-vRNAP for these promoters was measured by filter binding and transcription initiation was measured by catalytic autolabeling of mini-vRNAP. All templates showed similar binding affinities. However, only the template with a C positioned 12 bases downstream from the center of the hairpin was able to support transcription initiation. This result indicates that mini-vRNAP utilizes this position as the transcription start site (+1).

Example 6

Identification of Sequence Motifs Essential for mini-vRNAP Activity

[00647] As shown in FIG. 2A, vRNAP contains the sequence Rx_3Kx_6YG , designated Motif B in the Pol I and Pol α DNA polymerases and the T7-like RNA polymerases. To determine the relevance of this motif to vRNAP activity, two mutants K670A and Y678F (SEQ ID NO:8) (position numbers in mini-vRNAP) were constructed by site-specific mutagenesis of mini-vRNAP. These two positions were chosen because, in T7-like RNA polymerases, the lysine is involved in nucleotide binding and the tyrosine in discrimination against deoxynucleoside triphosphates (Maksimova, *et al.*, Eur. J Biochem. 195:841-847, 1991; Bonner, *et al.*, EMBO J. 11:3767-3775, 1992; Osumi-Davis, *et al.*, J. Mol Biol. 226:37-45, 1992). The His-tagged Y678F mini-vRNAP gene (SEQ ID NO:7) differs from that of the mini-vRNAP domain sequence (SEQ ID NO:3) at two positions: nucleotide 2033 (A) was changed to a T, and nucleotide 2034 (T) was changed to a C.

[00648] These RNA polymerase mutants were cloned under pBAD control, purified and tested for their ability to bind to wild type promoters. Both mutant polymerases bound to promoter DNA with wild type affinities and crosslinked to 5-Iodo-dU substituted P2 DNA templates at positions -11 and +3 with wild-type affinities (FIG. 14), indicating that these mutations do not affect promoter binding.

[00649] The mutant enzymes were tested for their ability to support run-off transcription. The wild-type enzyme and Y678F enzyme (SEQ ID NO:8) displayed similar activities at both template excess and template-limiting conditions, while the K670A enzyme exhibited decreased activity under both conditions (FIG. 15). Under limiting template conditions, all three enzymes

were activated by Eco SSB (right panel). However, the Y678F enzyme showed reduced discrimination between ribo- and deoxyribonucleoside triphosphates.

[00650] The initiation properties of the three enzymes were compared using catalytic autolabeling (FIG. 16). The K670A enzyme displays significantly reduced activity with the GTP derivative. The Y678F enzyme, in contrast to wild-type polymerase, incorporates dATP as efficiently as rATP in a single round of phosphodiester bond formation.

[00651] Therefore, the behavior of the K670A and Y678F mutant enzymes indicates that Motif B is involved in catalysis, with the lysine probably required for NTP binding and the tyrosine responsible for dNTP discrimination. These results suggest that, despite its lack of extensive sequence similarity, vRNAP is a Class II T7-like RNA polymerase. Results of recent experiments revealed the location of the two carboxylates (aspartates) involved in catalysis.

Example 7

Development of an *In Vivo* System Using Mini-vRNAP and N4 vRNAP Promoters for *in vivo* Expression of RNAs and Proteins

[00652] Plasmid templates were constructed with a reporter gene (α -peptide of β -galactosidase) cloned under the control of vRNAP promoter P2 present in either of two orientations (FIG. 17B). The reporter construct was generated by cloning a cassette into plasmid pACYC177, which was obtained from New England Biolabs. The cassette contains an approximately 30 bp long fragment originating from pT7Ac (purchased from United States Biochemical), a N4 promoter, and sequence encoding the alpha fragment of lacZ (lacZ'). The N4 promoter and lacZ' were generated by oligonucleotide annealing and PCR amplification, respectively. This cassette replaces the pACY177 sequence located between the cleavage sites for restriction enzymes ApaLI and BamHI. These reporter plasmids and recombinant full-length or mini-vRNAP expressing plasmids were introduced into *E. coli* DH5 α (Δ M15), a strain that encodes the β -galactosidase θ -peptide. Expression of the reporter gene α -peptide) in this strain results in the synthesis of active β -galactosidase and consequent production of blue colonies on X-gal plates. Transcription of α -peptide by full-length and mini-vRNAP was assayed on inducing-Xgal media and shown in FIG. 17A. Induction of full-length polymerase results in small colonies with no β -galactosidase activity. This is not surprising since full-length vRNAP is degraded in these cells (FIG. 17C). In contrast, induction of mini-vRNAP led to detectable levels of the protein (FIG. 17C) and to β -galactosidase activity only from the plasmid containing

promoter P2 in the proper orientation (FIG. 17A). These results indicate that this system will be suitable for *in vivo* expression of RNAs and proteins under mini-N4 vRNAP promoter control.

Example 8

Rolling Circle Transcription of Model ssDNA Transcription Substrates

[00653] Each oligonucleotides (50 picomoles), comprising a sense P2 promoter sequence (or, in control reactions, an anti-sense sequence to the P2 promoter or no promoter) at its 5'-end, which was phosphorylated, and up to 52 additional nucleotides corresponding to a model target sequence (e.g., for the human beta actin gene) in its 3'-portion, was ligated in a reaction mixture containing 0.2 mM ATP, 1 mM DTT, and 50 micrograms per ml of BSA for 2 hours at 60° C using 200 units of ThermoPhage™ RNA Ligase II (Prokaria, Reykjavik, Iceland, #Rlig122) in 1X ThermoPhage RNA Ligase II Buffer comprising 50 mM MOPS, pH 7.5, 5 mM MgCl₂, and 10 mM KCl. Then, linear oligos were removed by digestion with Exonuclease I (EPICENTRE Technologies, Madison, WI), the Exo I was heat-inactivated, and the circular ssDNA oligos were ethanol precipitated using standard techniques.

[00654] One picomole of circular ssDNA oligonucleotide, prepared as just described, was then incubated for four hours at 37°C in a 60-microliter reaction mixture comprising one microgram of mini-vRNAP (EPICENTRE), 1 mM each NTP, 1 mM DTT, and 5 micromolar *E. coli* SSB Protein (EPICENTRE), in 1X Transcription Buffer comprising 40 mM Tris HCl, pH 7.5, 10 mM NaCl, 6 mM MgCl₂, and 1 mM spermidine. The resulting mini-vRNAP transcription products were then analyzed by electrophoresis in a 1% agarose gel containing 0.22 M formaldehyde. Transcription products, including products having a length many-fold greater than the starting oligonucleotide, were observed on the the gel, indicating efficient rolling circle transcription. No transcription products were observed if the oligo did not contain a P2 promoter, if an anti-sense sequence to the P2 promoter was used instead of the sense P2 promoter, or if an unligated linear oligo with a sense P2 promoter was used.

Example 9

Use of a Random Hexamer Promoter Primer to Obtain a Circular ssDNA Transcription Substrate for Making Transcription Product Corresponding to a Target Nucleic Acid Sequence Comprising mRNA

[00655] Messenger RNA is obtained from a sample using techniques known in the art for isolating polyadenylated RNA. A suitable amount of mRNA, such as about 1-10 micrograms of mRNA, is then used for reverse transcription using MMLV reverse transcriptase (EPICENTRE)

and a large excess of a random hexamer promoter primer comprising a P2 sense promoter in its 5'-portion, which is phosphorylated at the 5'-end, and a random hexamer sequence at its 3'-end. Following reverse transcription according to the directions supplied with the MMLV reverse transcriptase, the resulting first-strand cDNA is treated with *E. coli* RNase H (EPICENTRE) to digest the RNA, the RNaseH is inactivated with heat, and the first-strand cDNA is purified and ethanol precipitated. Then, the first-strand cDNA is circularized using ThermoPhage™ RNA Ligase II (Prokaria) as described in Example 8. Preferably, circular ssDNA molecules are obtained which comprise from 20 to about 200 nucleotides of cDNA complementary to the target mRNA, which is operably joined to the P2 promoter.

[00656] *In vitro* transcription of the resulting circular ssDNA molecules, which comprise circular ssDNA transcription substrates of the invention, as described in Example 8, yields transcription products corresponding to the mRNA in the sample. In some transcription reactions, an inorganic pyrophosphatase and high NTP concentrations are used to increase the yield of transcription product obtained. The concentrations of N4 mini-vRNAP enzyme, NTPs and pyrophosphatase can be titrated to determine the levels of each component that results in maximum yield of transcription product.

[00657] All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

[00658] All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

[00659] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes

and modifications may be made thereto without departing from the spirit or scope of the appended claims.

CLAIMS

We claim:

1. A method for making transcription product corresponding to a target nucleic acid sequence, the method comprising:

- (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter;
- (b) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic sequence that is present in or complementary to at least a portion of a target nucleic acid in a sample;
- (c) obtaining a single-stranded transcription substrate by operably joining to the single-stranded DNA a single-stranded polynucleotide comprising a promoter sequence that binds the RNA polymerase;
- (d) obtaining nucleoside triphosphates (NTPs) that are substrates for the RNA polymerase and that are complementary to canonical nucleic acid bases;
- (e) admixing the RNA polymerase, the single-stranded transcription substrate and the NTPs; and
- (f) incubating the RNA polymerase and the single-stranded transcription substrate under conditions effective to allow synthesis of transcription product.

2. A method for making transcription product corresponding to a target nucleic acid sequence, the method comprising:

- (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter;
- (b) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic acid sequence that is present in or complementary to at least a portion of a target nucleic acid in a sample;
- (c) obtaining a single-stranded transcription substrate by operably joining to the single-stranded DNA a single-stranded polynucleotide comprising a promoter sequence that binds the RNA polymerase;
- (d) admixing the RNA polymerase and the single-stranded transcription substrate; and

- (e) incubating the RNA polymerase and the single-stranded ssDNA transcription substrate under conditions effective to allow synthesis of transcription product.

3. A method for obtaining additional rounds of synthesis of transcription product corresponding to a target nucleic acid sequence, the method comprising:

- (a) obtaining an RNA polymerase that can transcribe RNA using a single-stranded promoter;
- (b) obtaining a first transcription product by transcription of a first single-stranded transcription substrate comprising a polynucleotide corresponding to a target nucleic acid sequence;
- (c) obtaining a reverse transcriptase;
- (d) reverse transcribing the first single-stranded transcription product;
- (e) obtaining first-strand cDNA complementary to the first single-stranded transcription product;
- (f) obtaining a second single-stranded transcription substrate by operably joining to the first-strand cDNA a single-stranded polynucleotide comprising a promoter sequence that binds the RNA polymerase;
- (g) admixing the RNA polymerase and the second single-stranded transcription substrate; and
- (h) incubating the RNA polymerase and the second single-stranded transcription substrate under conditions effective to allow synthesis of a second transcription product.

4. The method of claim 1 wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

5. The method of claim 1 wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

6. The method of claim 1 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

7. The method of claim 1 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

8. The method of claim 1 wherein the RNA polymerase is E. coli RNAP and the promoter is a single-stranded pseudopromoter for E.coli RNAP.

9. The method of claim 1 wherein the RNA polymerase is a T7-type RNAP and the promoter is a cognate single-stranded pseudopromoter for the T7-type RNAP.

10. The method of claim 1 wherein the RNA polymerase is T7 RNAP and the promoter is a single-stranded pseudopromoter for T7 RNAP.

11. The method of claim 1 wherein the RNA polymerase is T3 RNAP and the promoter is a single-stranded pseudopromoter for T3 RNAP.

12. The method of claim 1 wherein the RNA polymerase is SP6 RNAP and the promoter is a single-stranded pseudopromoter for SP6 RNAP.

13. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising an RNA target nucleic acid.

14. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising an mRNA target nucleic acid.

15. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising an mRNA target nucleic acid that is full-length.

16. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising mRNA target nucleic acid corresponding to substantially all mRNA in the sample.

17. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising a DNA target nucleic acid.

18. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence comprising a DNA target nucleic acid that is a product of an amplification reaction.

19. The method of claim 1 wherein the amplification reaction is selected from the group consisting of PCR, RT-PCR, NASBA, TMA, 3SR, LCR, LLA, SDA, RCA, Multiple Displacement Amplification, ICANTM, UCANTM, Loop-AMP, SPIATM and Ribo-SPIATM.

20. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that is obtained by primer extension of a larger DNA target nucleic acid.

21. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that is obtained by reverse transcriptase primer extension of at least one mRNA target nucleic acid.

22. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that is obtained by reverse transcriptase primer extension of substantially all mRNA target nucleic acid in a sample.

23. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that has a tail sequence comprising at least two nucleotides.

24. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that has a tail sequence comprising of dCMP nucleotides.

25. The method of claim 1 wherein the single-stranded DNA comprises a target nucleic acid sequence that has a tail sequence between two to ten nucleotides.

26. The method of claim 1 wherein the ssDNA transcription substrate of step (c) is obtained using a promoter splice template oligo.

27. The method of claim 26 wherein the single-stranded DNA is obtained by reverse transcription of a transcription product.

28. The method of claim 1 wherein the single-stranded DNA in step (c) is obtained by reverse transcription of a transcription product prepared using the method of claim 26.

29. The method of claim 1, wherein the ssDNA transcription substrate is obtained by using a promoter ligation oligo and a ligation splint.

30. The method of claim 1, wherein the single-stranded DNA is obtained by reverse transcription of a transcription product prepared using a promoter ligation oligo and a ligation splint.

31. The method of claim 1 wherein a single-stranded transcription substrate is obtained by DNA polymerase-catalyzed primer extension of a promoter primer using the target nucleic acid in the sample as a template, followed by ligation of the 5'-end of the primer-extended promoter primer to the 3'-end primer extension, comprising the single-stranded DNA comprising the target nucleic acid sequence, thereby operably joining the promoter to the target nucleic acid sequence to form a circular single-stranded transcription substrate.

32. The method of claim 31 wherein the target nucleic acid in the sample comprises RNA and the DNA polymerase used for primer extension is an enzyme with reverse transcriptase activity.

33. The method of claim 31 wherein the target nucleic acid in the sample comprises mRNA.

34. The method of claim 31 further comprising the DNA polymerase used for primer extension is an enzyme with reverse transcriptase activity.

35. The method of claim 31 wherein the single-stranded DNA comprising a target nucleic acid sequence is obtained by reverse transcription of a transcription product prepared using the method of claim 1.

36. The method of claim 1 wherein a linear single-stranded transcription substrate is obtained by cleaving a circular single-stranded transcription substrate obtained using the method of claim 31 at a site that is 3'-of the promoter sequence and 5'-of the target nucleic acid sequence.

37. The method of claim 1 wherein the single-stranded DNA comprising a target nucleic acid sequence is obtained by reverse transcription of a transcription product prepared using the method of claim 31.

38. The method of claim 1, wherein at least one of the NTPs comprises a 2'-amino-deoxynucleoside triphosphate.

39. The method of claim 1, wherein at least one of the NTPs comprises a 2'-amino-dCTP.

40. The method of claim 1, wherein at least one of the NTPs comprises a 2'-fluoro-deoxynucleoside triphosphate.

41. The method of claim 1, wherein at least one of the NTPs comprises a 2'-fluoro-dCTP.

42. The method of claim 1, wherein at least one of the NTPs comprises a 2'-fluoro-dUTP.

43. The method of claim 1, wherein at least one of the NTPs comprises a 2'-azido-deoxynucleoside triphosphate.

44. The method of claim 1, wherein at least one of the NTPs comprises a 2'-azido-dCTP.

45. The method of claim 1 wherein the NTPs are complementary to canonical nucleic acid bases.

46. A kit for performing the method of claim 1 wherein the kit comprises an RNA polymerase that can use a single-stranded promoter for transcription of RNA and a promoter splice template oligo.

47. The kit of claim 46 wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

48. The kit of claim 47 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

49. The kit of claim 47 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

50. The kit of claim 46 wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

51. The kit of claim 46 wherein the RNA polymerase comprises E. coli RNAP and the promoter splice template comprises a single-stranded pseudopromoter for E.coli RNAP.

52. The kit of claim 46 wherein the RNA polymerase comprises T7-type RNAP and the promoter splice template oligo comprises a single-stranded pseudopromoter for the T7-type RNAP.

53. The kit of claim 46 wherein the RNA polymerase comprises T7 RNAP and the promoter splice template oligo comprises a single-stranded pseudopromoter for T7 RNAP.

54. The kit of claim 46 wherein the RNA polymerase comprises T3 RNAP and the promoter splice template oligo comprises a single-stranded pseudopromoter for T3 RNAP.

55. The kit of claim 46 wherein the RNA polymerase comprises SP6 RNAP and the promoter splice template oligo comprises a single-stranded pseudopromoter for SP6 RNAP.

56. A kit for performing the method of claim 1 wherein the kit comprises an RNA polymerase that can use a single-stranded promoter for transcription of RNA and a promoter ligation oligo.

57. The kit of claim 56 wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

58. The kit of claim 57 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

59. The kit of claim 57 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

60. The kit of claim 56 wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

61. The kit of claim 56 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

62. The kit of claim 56 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

63. The kit of claim 56 wherein the RNA polymerase comprises E. coli RNAP and the promoter ligation oligo comprises a single-stranded pseudopromoter for E.coli RNAP.

64. The kit of claim 56 wherein the RNA polymerase comprises T7-type RNAP and the promoter ligation oligo comprises a single-stranded pseudopromoter for the T7-type RNAP.

65. The kit of claim 56 wherein the RNA polymerase comprises T7 RNAP and the promoter ligation oligo comprises a single-stranded pseudopromoter for T7 RNAP.

66. The kit of claim 56 wherein the RNA polymerase comprises T3 RNAP and the promoter ligation oligo comprises a single-stranded pseudopromoter for T3 RNAP.

67. The kit of claim 56 wherein the RNA polymerase comprises SP6 RNAP and the promoter ligation oligo comprises a single-stranded pseudopromoter for SP6 RNAP.

68. A kit for performing the method of claim 1 wherein the kit comprises an RNA polymerase that can use a single-stranded promoter for transcription of RNA and a promoter primer.

69. The kit of claim 68 wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:15.

70. The kit of claim 69 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

71. The kit of claim 69 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

72. The kit of claim 68 wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

73. The kit of claim 68 wherein the promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

74. The kit of claim 68 wherein the promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

75. The kit of claim 68 wherein the RNA polymerase comprises E. coli RNAP and the promoter primer comprises a single-stranded pseudopromoter for E.coli RNAP.

76. The kit of claim 68 wherein the RNA polymerase comprises T7-type RNAP and the promoter primer comprises a single-stranded pseudopromoter for the T7-type RNAP.

77. The kit of claim 68 wherein the RNA polymerase comprises T7 RNAP and the promoter primer comprises a single-stranded pseudopromoter for T7 RNAP.

78. The kit of claim 68 wherein the RNA polymerase comprises T3 RNAP and the promoter primer comprises a single-stranded pseudopromoter for T3 RNAP.

79. The kit of claim 68 wherein the RNA polymerase comprises SP6 RNAP and the promoter primer comprises a single-stranded pseudopromoter for SP6 RNAP

80. The method of claim 1 wherein the target nucleic acid sequence comprises a 3'-portion that encodes a first sequence, a 5'-portion that encodes a second sequence that is complementary to the first sequence, and a middle portion that joins the 3'portion and the 5'portion, wherein the middle portion comprises a sequence that is not complementary to either the 3'-portion or the 5'portion and wherein the transcription product comprises a hairpin RNA.

81. The method of claim 80, wherein the hairpin RNA corresponds to a target nucleic acid sequence in a target nucleic acid comprising an mRNA.

82. The method of claim 80 , wherein the hairpin RNA has RNA interference activity in a cell that synthesizes an mRNA target nucleic acid comprising the target nucleic acid sequence.

83. The method of claim 80 wherein the hairpin RNA comprises siRNA.

84. The method of claim 80 wherein the hairpin RNA comprises at least one modified nucleoside triphosphate.

85. The method of claim 84 wherein the modified nucleoside triphosphate is selected from the group consisting of 2'-amino-deoxynucleoside triphosphate, 2'-amino-dCTP, 2'-fluoro-deoxynucleoside triphosphate, 2'-fluoro-dCTP, 2'-fluoro-dUTP, 2'-azido-deoxynucleoside triphosphate, and 2'-azido-dCTP.

86. The method of claim 80 wherein the hairpin RNA is at least 40 nucleotides in length.

87. The method of claim 80 wherein the hairpin RNA is at least 100 nucleotides in length.

88. The method of claim 80 wherein the hairpin RNA is made with an RNA polymerase that can use a single-stranded promoter for transcription of RNA

89. The method of claim 88, wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

90. The method of claim 88, wherein the single-stranded promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

91. The method of claim 88 wherein the single-stranded promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

92. The method of claim 88, wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

93. The method of claim 88 wherein the RNA polymerase comprises E. coli RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for E. coli RNAP.

94. The method of claim 88 wherein the RNA polymerase comprises T7-type RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for the T7-type RNAP.

95. The method of claim 88 wherein the RNA polymerase comprises T7 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for T7 RNAP.

96. The method of claim 88, wherein the RNA polymerase comprises T3 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for T3 RNAP.

97. The method of claim 88 wherein the RNA polymerase comprises SP6 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for SP6 RNAP.

98. The method of claim 80 wherein the hairpin RNA is made in vitro.

99. The method of claim 80 wherein the hairpin RNA is made in vivo.

100. A method for attenuating expression of a target gene in a cell comprising introducing the hairpin RNA of claim 80 into the cell.

101. The method of claim 100 wherein the expression of a target gene in the cell is attenuated in vitro.

102. The method of claim 100 wherein the expression of the target gene in the cell is attenuated in vivo.

103. The method of claim 100, wherein the cell comprises a mammalian cell.

104. The method of claim 103, wherein the mammalian cell comprises a human cell.

105. A hairpin RNA made by the method of claim 80.
106. A cell comprising a hairpin RNA made by the method of claim 80.
107. A kit for making the hairpin of claim 80, the kit comprising an RNA polymerase that can use a single-stranded promoter for transcription of RNA and an oligonucleotide comprising a sequence corresponding to a single-stranded promoter sequence.
108. A method of cloning a target nucleic acid, the method comprising:
- a) obtaining a single-stranded DNA wherein the single-stranded DNA comprises a target nucleic acid sequence that is present in or complementary to the target nucleic acid;
 - b) joining to the single-stranded DNA a single-stranded polynucleotide comprising a single-stranded origin of replication and a marker gene;
 - c) making a circular ssDNA molecule by covalently joining the 3'-end and the 5'-end of the product of step (b);
 - d) transforming the circular ssDNA molecule into a host cell, in which the marker gene is expressible, wherein the host cell is capable of replicating the circular ssDNA molecule; and
 - e) growing the host cell under conditions that support the expression of the marker gene.
109. The method of claim 108, wherein the target nucleic acid comprises an RNA target nucleic acid.
110. The method of claim 108, wherein the target nucleic acid comprises an mRNA target nucleic acid.
111. The method of claim 108, wherein the target nucleic acid sequence comprises an mRNA target nucleic acid that is full-length.
112. The method of claim 108, wherein the target nucleic acid comprises an mRNA target nucleic acid corresponding to substantially all mRNA within a sample.

113. The method of claim 108, wherein the target nucleic acid comprises a DNA target nucleic acid.

114. The method of claim 108, wherein the target nucleic acid comprises a target nucleic acid that is a product of an amplification reaction.

115. The method of claim 114 wherein the amplification reaction is selected from the group consisting of PCR, RT-PCR, NASBA, TMA, 3SR, LCR, LLA, SDA, RCA, Multiple Displacement Amplification, ICANTM, UCANTM, Loop-AMP, SPIATM and Ribo-SPIATM.

116. The method of claim 108 wherein the target nucleic is obtained by primer extension of a larger DNA target nucleic acid.

117. The method of claim 108 wherein the target nucleic acid is obtained by reverse transcriptase primer extension of at least one mRNA target nucleic acid.

118. The method of claim 108 wherein the target nucleic acid sequence is obtained by reverse transcriptase primer extension of substantially all mRNA target nucleic acids in a sample.

119. The method of claim 108, wherein the single-stranded polynucleotide comprising a single-stranded origin of replication and a marker gene are joined to the single-stranded DNA by using a promoter splice template oligo.

120. The method of claim 108, wherein the single-stranded polynucleotide comprising a single-stranded origin of replication and a marker gene are joined to the single-stranded DNA by using a promoter ligation oligo.

121. The method of claim 108, wherein the single-stranded origin of replication comprises an M13 origin of replication.

122. The method of claim 108, wherein the marker gene comprises an antibiotic-resistance gene.

123. The method of claim 108, wherein the marker gene comprises a beta-galactosidase gene.

124. The method of claim 108, wherein the single stranded polynucleotide comprises a transposon recognition sequence.

125. The method of claim 108, wherein the single-stranded polynucleotide comprises a site that can be recognized by a recombinase.

126. The method of claim 108, wherein the circular ssDNA molecule is made by using a ligase that catalyzes non-homologous intramolecular ligation.

127. The method of claim 126, wherein the ligase is ThermoPhage™ RNA Ligase II.

128. The method of claim 108, wherein the circular DNA molecule is made by DNA polymerase-catalyzed primer extension of a primer using the target nucleic acid as a template, followed by ligation of the 3'-end of the primer extension product to the 5'-end of the primer extension product, wherein the primer comprises a single-stranded origin of replication and a marker gene.

129. A method of constructing a nucleic acid library comprising clones of substantially all nucleic acids within a sample by using the method of claim 108.

130. The method of claim 108, wherein the single-stranded polynucleotide comprises a single-stranded promoter that binds a RNA polymerase that can transcribe RNA using a promoter that is single-stranded.

131. The method of claim 130, wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

132. The method of claim 130, wherein the single-stranded promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

133. The method of claim 130 wherein the single-stranded promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

134. The method of claim 130, wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

135. The method of claim 130 wherein the RNA polymerase comprises E. coli RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for E.coli RNAP.

136. The method of claim 130 wherein the RNA polymerase comprises T7-type RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for the T7-type RNAP.

137. The method of claim 130 wherein the RNA polymerase comprises T7 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for T7 RNAP.

138. The method of claim 130, wherein the RNA polymerase comprises T3 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for T3 RNAP.

139. The method of claim 130 wherein the RNA polymerase comprises SP6 RNAP and the single-stranded promoter comprises a single-stranded pseudopromoter for SP6 RNAP.

140. The method of claim 108, wherein the single-stranded polynucleotide comprises a single-stranded promoter that binds a RNA polymerase that can transcribe RNA using a promoter that is single-stranded and wherein the host cell comprises an expressible gene encoding an RNA polymerase that can transcribe RNA using the single-stranded promoter.

141. The method of claim 140, wherein the expressible gene is operably joined to an inducible promoter.

142. The method of claim 141, wherein the inducible promoter is selected from the group consisting of a bad promoter, a lac promoter, a trp promoter, a tac promoter and a lambda promoter.

143. A method of constructing a nucleic acid library comprising clones of substantially all nucleic acids within a sample by using the method of any one of claims 130 or 140.

144. A method of constructing a nucleic acid library comprising clones of substantially all mRNAs within a sample by using the method of any one of claims 130 or 140.

145. A composition comprising a clone made by using the method of any one of claims 108 - 144.

146. A composition comprising a nucleic acid library made by using the method of claim 129.

147. A host cell comprising a circular DNA molecule made by using the method of claim 108.

148. A circular DNA molecule made by using the method of claim 108.

149. A kit for performing the method of claim 108.

150. A method for detecting an analyte in a sample, the method comprising:

- a) obtaining a transcription signaling system comprising a ssDNA comprising:
 - i) a promoter sequence that binds an RNA polymerase that can transcribe RNA using a single-stranded promoter, and
 - ii) a signal sequence that is operably joined to the promoter sequence;

- b) joining the transcription signaling system to an analyte-binding substance;
- c) contacting the analyte-binding substance to which the transcription signaling system is joined with a sample under conditions effective to allow binding of an analyte to the analyte-binding substance and forming a specific binding pair;
- d) removing the specific binding pair from the sample;
- e) incubating the specific-binding pair with an RNA polymerase that can transcribe RNA using a single-stranded promoter under conditions effective to allow synthesis of a transcription product; and
- f) detecting the transcription product.

151. The method of claim 150, wherein the analyte is selected from the group consisting of a biochemical molecule, a biopolymer a protein, a glycoprotein, a lipoprotein, an enzyme, a hormone, a receptor, an antigen, an antibody, a nucleic acid, a DNA, an RNA, a polysaccharide and a lipid.

152). The method of claim 150 wherein the promoter sequence is an N4 vRNAP promoter sequence set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

153) The method of claim 150 wherein the promoter sequence is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

154) The method of claim 150, wherein the RNA polymerase comprises a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

155) The method of claim 150, wherein the RNA polymerase comprises a polypeptide encoded by the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

156) The method of claim 150 wherein the RNA polymerase comprises E. coli RNAP and the promoter sequence comprises a single-stranded pseudopromoter for E.coli RNAP.

157) The method of claim 150 wherein the RNA polymerase comprises T7-type RNAP and the promoter sequence comprises a single-stranded pseudopromoter for the T7-type RNAP.

158) The method of claim 150 wherein the RNA polymerase comprises T7 RNAP and the promoter sequence comprises a single-stranded pseudopromoter for T7 RNAP.

159) The method of claim 150, wherein the RNA polymerase comprises T3 RNAP and the promoter sequence comprises a single-stranded pseudopromoter for T3 RNAP.

160) The method of claim 150 wherein the RNA polymerase comprises SP6 RNAP and the promoter sequence comprises a single-stranded pseudopromoter for SP6 RNAP.

161) The method of claim 150 wherein the signal sequence comprises a substrate for Q-beta replicase.

162) The method of claim 150 wherein the signal sequence comprises a sequence that encodes a detectable protein.

163) The method of claim 162 wherein the detectable protein is green fluorescent protein.

164) The method of claim 150 wherein the signal sequence comprises a sequence that is detectable by a probe.

165) The method of claim 164 wherein the sequence that is detectable by a probe comprises a molecular beacon.

166) The method of claim 150, wherein the analyte-binding substance is selected from the group consisting of a nucleic acid, a polynucleotide, an oligonucleotide, a segment of a nucleic acid or polynucleotide, a DNA, an RNA, a molecule comprising both DNA and RNA mononucleosides, modified DNA mononucleosides, a molecule obtained by a method termed "SELEX", a nucleic acid molecule having an affinity for protein molecules, a polynucleotide

molecule having an affinity for protein molecules, an operator, a promoter, an origin of replication, a restriction endonuclease recognition sequence, a ribosomal nucleic acid sequence, a sequence recognized by steroid hormone-receptor complexes, a peptide nucleic acid (PAN), a nucleic acid and a PNA, a molecule prepared by using a combinatorial library of randomized peptide nucleic acids, an oligonucleotide or polynucleotide with a modified backbone that is not an amino acid, a molecule identified by using high throughput screening methods, lectin, a receptor for a hormone, a hormone, and an enzyme inhibitor.

167) The method of claim 150 wherein the binding of step (c) comprises non-covalent bonds.

168) The method of claim 167 wherein the non-covalent bonds comprise hydrogen-bonds.

169) The method of claim 167 wherein the non-covalent bonds comprise hydrophobic interactions.

170) The method of claim 167 wherein the non-covalent bonds comprise van der Waals forces.

171) The method of claim 167 wherein the non-covalent bonds comprise salt bridges.

1/25

Bacteriophage N4 vRNAP promoters




FIG. 1

N4 vRNAP and generation of mini-vRNAP




FIG. 2

Identification of the minimal active domain of N4 vRNAP
by proteolytic cleavage.





FIG. 3

4/25




Plasmid name: pMini-vRNAP
Plasmid size: 7426 bp
Constructed by: K. M. Kazmierczak
Construction date: 2/2000
Comments: Insert cloned into Invitrogen
pBAD 8 expression plasmid

FIG 4

5/25

Purification of cloned vRNAP and mini-vRNAP

**FIG 5**

6/25

Activation of N4 vRNAP transcription by *Eco* SSB
at different ssDNA concentrations




FIG 6

7/25

Effect of *Eco* SSB on ssDNA template recycling




FIG. 7

8/25

Effect of *Eco* SSB on the state of template DNA
and product RNA in vRNAP transcription




FIG. 8

9/25

Transcription initiation by vRNAP and mini-vRNAP




FIG. 9

10/25

Effect of *Eco* SSB on transcription of vRNAP and mini-RNAP




FIG. 10

Identification of the transcription start site by catalytic autolabeling

13/25





FIG. 13

UV crosslinking of mutant mini-vRNAPases
to promoter oligonucleotides



Run-off transcription by mutant mini-vRNAPases




FIG. 15

Mutant mini-vRNAPases transcription initiation

Motif B RgiaKnplltitiY
 ↓ ↓
 A F



FIG. 16

Detection of *in vivo* activities of N4 vRNAP and mini-vRNAP




FIG. 17

18/25




FIG 18

19/25




FIG 19

20/25




FIG 20

21/25




FIG 21

22/25




FIG 22

23/25




FIG 23





FIG 24 (continued on next page)

25/25



SEQUENCE LISTING

<110> Davydova, Elena K.
 Rothman-Denes, Lucia B.
 Dahl, Gary A.
~~Meis, Judith B.~~
 Gerdes, Svetlana Y.
 Jendrisak, Jerome J.

<120> PREPARATION AND USE OF SINGLE-STRANDED TRANSCRIPTION SUBSTRATES
 FOR SYNTHESIS OF TRANSCRIPTION PRODUCTS CORRESPONDING TO TARGET
 SEQUENCES

<130> ARCD:?

<140> ?

<141> ?

<160> 29

<170> PatentIn Ver. 2.1

<210> 1

<211> 10506

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 1

```

atgtcagtat ttgatagact ggctgggttc gcagacagcg taaccaatgc aaagcaagtt 60
gacgtctcta ctgcaaccgc ccagaagaaa gctgaacaag gtgtcactac tcctcttggt 120
tctcctgatg ctgcttatca aatgcaagct gcccgactcg gtaatgttgg ggctaattgca 180
tttgaaccag ggacagtgc atcagatttc atgaatctga cccaatgca aatcatgaat 240
aagtatgggg ttgagcaagg cttacaactt atcaatgctc gtgctgatgc aggaaccag 300
gtattcaatg attcagttac tacaagaact cctggggaag aactggggga tattgctact 360
gggtgtggcc ttggttttgt taataccctt gggggcattg gtgctcttgg ggcaggctta 420
ctcaacgatg atgcagggtg tgttgttgct caacaattga gtaagtttaa tgatgctgtt 480
catgctaccc aaagccaggc attacaagat aaacgtaagc tctttgctgc tcgtaactta 540
atgaatgaag tagagagtga acgtcagtat caaacagata agaaagaagg cactaatgac 600
atagtagctt ccttatctaa atttgacgt gatgtttag gttcaattga gaatgctgct 660
caactgact ctattatttc tgatgggtta gcagaagggg taggttctct attaggtgct 720
ggtcctgtat taaggggtgc atctttactg ggtaaagcag ttgtccagc aaatactctt 780
cgtagtgctg cattgggtgg tgctattgat gcaggtagt gtactcagtc actggctcgt 840
attgctctta ctgtaggtag agctgcaccg ggtatgggtg gtgttggtgc aatggaagct 900
gggtgtgcat accaacaac tgctgatgaa attatgaaga tgagtcttaa agacttagag 960
aagtctcctg tttatcagca acatattaaa gatggtatgt cccctgaaca ggctcgtcgt 1020
cagactgcat ctgaaactgg tcttactgct gctgctattc aattacctat tgctgctgca 1080
accggtcctc tggatcccg ttttgagatg gtcctttcc gtgctggctc tttaggtgct 1140
gtaggtatga accttgccc tgaaacagt gaagaagggtg ttcagggtgc tacaggccaa 1200
ctggctcaga atattgcaca gcaacaaaac attgataaga accaagacct gcttaaaggt 1260
gtcgttacac aggtggttt aggtgctctt tatggctttg gttctgctgg tgtgtacag 1320
gctccggtg gtgctgctc tttagcagg gctgcaactg ctctgtatt gcgtaccaca 1380
atggctggtg ttaaagctgc tggtagtgta gcaggtaagg ttgtttctcc tattaagaat 1440
actttagtag ctggtggtga acgggttatg aagcagaatg aagaagcatc tcctgttgct 1500
gatgactatg ttgcacaggc agcacaagaa gctatggctc aagcaccaga agcagaagtt 1560
actattcgtg atgctgttga agcaactgat gctactccag aacagaaagt tgcagcacac 1620
cagtatgttt ctgacttaat gaatgctact cgttttaatc ctgaaaatta tcaggaagca 1680
ccagagcata ttcgtaatgc tgtagctggt tctactgacc aagtacaggt tattcagaag 1740

```

ttagcagact	tagttaacac	attagatgaa	tctaatacctc	aagcactgat	ggaagctgca	1800
tcttatatgt	atgatgctgt	ttcagagttt	gagcagttca	ttaaccgtga	ccctgctgca	1860
ctggatagca	ttcctaaaga	ttctccggct	attgagttac	tcaaccgtta	tacgaatctg	1920
acagctaata	ttcagaacac	acaaaaagta	attgggtgcac	tgaatgttat	taatcgaatg	1980
attaatgaat	ctgctcagaa	tggttctttg	aatgtgactg	aagaatccag	tccacaggaa	2040
atgcagaacg	tagcattagc	tgctgaagta	gcccctgaaa	agctcaatcc	agagtctgta	2100
aatgtttgttc	ttaaacatgc	tgctgatggg	cgtattaaac	tgaataatcg	ccagattgct	2160
gcccctccaga	atgctgctgc	aatcctgaag	ggggcacggg	aatatgatgc	agaagctgcc	2220
cgtcttggtg	tacgtcctca	agacattgtg	agtaaacaga	ttaaaacgga	tgagagcaga	2280
actcaggaag	gacaatactc	tgcgttgcaa	catgcgaata	ggattcgggc	tgcgataaac	2340
tctggttaatt	tcgagttggc	ctccgcttac	ctgaacgact	ttatgcagtt	cgcccagcac	2400
atgcagaata	aggttggagc	gttgaatgag	catcttggtta	cggggaatgc	ggataagaat	2460
aagtctgtcc	actaccaagc	tcttactgct	gacagagaat	gggttcgtag	ccgtaccgga	2520
ttgggggtga	atccctatga	cactaagtcg	gttaaatattg	cccagcaagt	tgctcttgaa	2580
gcgaaaacg	tagcggatat	tgctaattgcc	ctcgcttcgg	cttaccgga	actgaaggtc	2640
agtcataataa	aagttactcc	attggattca	cgtcttaacg	ctcctgctgc	tgagggtggtc	2700
aaggcattcc	gtcaaggcaa	tcgagacggt	cgtctctctc	aaccgaaagc	tgactccgtg	2760
aatcagggtta	aagaaactcc	tgttacaaaa	caggaaccag	ttacatctac	tgtacagact	2820
aagactcctg	ttagtgaatc	tgttaaaaaca	gaacctacta	ctaaagagtc	tagcccacag	2880
gctataaaaag	aacctgtgaa	ccagtctgaa	aaacaggatg	ttaaccttac	taatgaggac	2940
aacatcaagc	aacctactga	atctgtttaa	gaaactgaaa	cttctacaaa	agaaagtaca	3000
gttacagaag	aattaaaaga	aggtattgat	gctgtttacc	cttcattggt	aggtactgct	3060
gattctaaag	cagaggggat	taagaactat	ttcaaattgt	cctttacctt	accagaagaa	3120
cagaaatccc	gtactggttg	ttcagaagca	cctctaaaag	atgtagccca	agctctgtct	3180
tctcgtgctc	gttatgaact	ctttactgag	aaagaaactg	ctaaccctgc	ttttaatggg	3240
gaagttatta	agcgatacaa	agaactcatg	gaacatgggg	aaggatttgc	tgatattctt	3300
cgctcccgtc	tggtctaggt	ccttaacact	aaggatgttg	gtaaacgttt	tgctcaagggt	3360
acagaagcca	accgttgggg	aggtggtaag	ttacttaaca	ttgttgagca	ggatggggat	3420
acctttaagt	acaacgaaca	attgctacag	actgctgtat	tagcaggctc	tcaatggaga	3480
cttactgcta	ccagcaatac	tgctatcaaa	gatgcaaaag	atggtgctgc	tattactggt	3540
attgaccaag	ctctgctgcc	agaagggtta	gtagagcaat	ttgatactgg	tatgacactc	3600
actgaagcag	ttagttccct	ggctcagaaa	attgagtcct	actggggatt	atctcgtaat	3660
ccaaatgctc	cattgggcta	taccaaaggc	atccctacag	caatggctgc	tgaaattctg	3720
gctgcatttg	tagagtctac	tgatgttgta	gagaacatcg	tggtatgtgc	agaaattgac	3780
ccagataaca	agaagactat	tggtctgtac	accattactg	aactggattc	cttcgaccca	3840
attaatagct	tcctactgct	tattgaagaa	gctgttttag	tgaatcctac	agagaagatg	3900
ttctttgggtg	atgacattcc	tcctgtagct	aatactcagc	ttcgtaaccc	tgctgttcgt	3960
aatactccag	aacagaaggc	tgcatgaaa	gcagagcagg	ctacagagtt	ctatgtacac	4020
acccaatgg	ttcaattcta	tgagacgtta	ggtaaagacc	gtattctcga	actgatgggt	4080
gctgggtactc	tgaataaaga	gttacttaat	gataaccatg	ctaaatctct	ggaaggtaag	4140
aaccgttcag	tagaggactc	ttacaaccaa	ctgttctccg	tcattgagca	ggtaagagca	4200
cagagcggaag	acatctctac	tgtacctatt	cactatgcac	acaatatgac	ccgtgttggt	4260
cgtatgcaga	tgtttaggta	atacaatcct	caactcagca	aactggttcg	tgaggccatc	4320
ttacctacta	aagctacttt	ggatttatcg	aaccagaaca	atgaagactt	ctctgcattc	4380
cagttagggtc	tggtcagggc	attggacatt	aaagtccata	ctatgactcg	tgaggttatg	4440
tctgacgagt	tgactaaatt	actggaagggt	aatctgaaac	cagccattga	tatgatgggt	4500
gagtttaata	ccactgggtc	cttaccagaa	aacgcagttg	atgttctgaa	tacagcatta	4560
ggagatagga	agtcattcgt	agcattgatg	gctcttatgg	agtattcccg	ttacttagta	4620
gcagaggata	aatctgcatt	tgtaactcca	ctgtatgtag	aagcagatgg	tggtactaat	4680
ggtccaatca	atgccatgat	gctaattgaca	ggcgggtctgt	ttactcctga	ctggattcgt	4740
aatattgccca	aagggggcct	gttcattgggt	tctocaaata	agaccatgaa	tgagcatcgc	4800
tctactgctg	acaataatga	tttatatcaa	gcatacacta	atgctttgat	ggaatcgttg	4860
ggtaagttac	gtagtaacta	tgccctctaat	atgcctattc	agtctcagat	agacagtctt	4920
ctttctctga	tggtttgttt	tttaccggat	attaatcttg	gtgagaatgg	tgctttagaa	4980
cttaaaccgtg	gtattgctaa	gaaccactg	actattacca	tctatgggtc	tggtgctcgt	5040
ggtattgcag	gtaagctggg	tagttctggt	actgatgcca	tctatgagcg	tatgtctgat	5100
gtactgaaag	ctcgtgctaa	agacccaaat	atctctgctg	ctatggcaat	gtttggtaag	5160
caagctgctt	cagaagcaca	tgctgaagaa	cttcttgccc	gtttcctgaa	agatatggaa	5220
acactgactt	ctactgttcc	tgtaaaacgt	aaaggtgtac	tggaactaca	atccacagggt	5280
acaggagcca	aaggaaaaat	caatcctaag	acctatacca	ttaagggcga	gcaactgaag	5340
gcacttcagg	aaaatatgct	gcacttcttt	gtagaaccac	tacgtaatgg	tattactcag	5400

actgtaggtg	aaagtctggt	gtactctact	gaacaattac	agaaagctac	tcagattcaa	5460
tctgtagtgc	tggaagatat	gttcaaacag	cgagtacaag	agaagctggc	agagaaggct	5520
aaagacccaa	catggaagaa	aggtgatttc	cttactcaga	agaactgaa	tgatattcag	5580
gcttctctga	ataacttagc	ccctatgatt	gagactgggt	ctcagacttt	ctacattgct	5640
ggttcagaaa	atgcagaagt	agcaaatcag	gtattagcta	ctaaccctga	tgaccgtatg	5700
cgtgtaccaa	tgagtatcta	tgctccagca	caggccggtg	tagcagggtat	tccattttatg	5760
actattggta	ctgggtgatg	catgatgatg	caaactcctt	ccactatgaa	aggtgcacca	5820
aagaataccc	tcaaaatcct	tgatgggtatg	aacattgggt	tgaatgacat	cactgatgcc	5880
agtcgtaaa	ctaataaagc	tgtttacact	tcttggcagg	gtaaccctat	taagaatgtt	5940
tatgaatcat	atgctaagtt	catgaagaat	gtagatttca	gcaagctgtc	ccctgaagca	6000
ttggaagcaa	ttggtaaate	tgctctggaa	tatgaccaac	gtgagaatgc	tactgtagat	6060
gatattgcta	acgctgcac	tctgattgaa	cgtaacttac	gtaatatgtc	actgggtgta	6120
gatattcgct	ataaggtgct	ggataaggta	aatctgtcca	ttgaccagat	ggctgctgta	6180
ggtgctcctt	atcagaacaa	cggtaagatt	gacctcagca	atatgacccc	tgaacaacag	6240
gctgatgaac	tgaataaact	tttccgtgaa	gagttagaag	cccgtaaaca	aaaagtcgct	6300
aaggctaggg	ctgaagtcaa	agaagaaact	gtttctgaaa	aagaaccagt	gaatccagac	6360
tttggtagtg	taggccgtga	gcataaggca	tctgggtgtc	gtatcctgtc	tgctactgct	6420
attcgtaatc	tggtctaatg	tagtaatctg	ccatctactc	aggcagctac	tcttgcggag	6480
attcagaaat	cactggcagc	taaagactat	aagattatct	acggtacacc	tactcagggt	6540
gcagagtatg	ctcgtcagaa	gaatgttact	gaattgactt	ctcaggaaat	ggaagaagct	6600
caggcaggta	atattttatg	ctggactaac	ttcgtatgata	agaccattta	tctgggttagc	6660
ccatctatgg	aaaccctcat	tcatgaactg	gttcatgcct	ctaccttcga	ggaagtttat	6720
tccttctatc	agggtaatga	agtaagccct	acttctaagc	aggctattga	gaaccttgaa	6780
ggtctgatgg	aacagttccg	ttctctggat	atttccaaag	attctccaga	aatgagagaa	6840
gcataatgctg	atgctattgc	aactatcgaa	ggtcatttga	gtaatggatt	tggtgacca	6900
gctatctcta	aagctgctgc	tcttaatgag	tttatggctt	gggggttagc	taaccgtgct	6960
cttgctgcta	aacagaagag	aacatcttca	ctggttcaaa	tggtgaaaga	tggttatcag	7020
gcttaataga	aattgatttg	gggacgtaaa	caagctcctg	cattgggaga	agatagtttc	7080
tccaatctgc	tggtttaactc	tgcaattctg	atgcgtagcc	aacctacaac	tcaggcagta	7140
gctaaagatg	gcacactggt	ccatagcaaa	gcataatggt	ataatgaacg	tctgtctcag	7200
ttgaaccaga	ctttcgataa	actggtaact	gattaccttc	gtactgaccc	agttacagaa	7260
gtagaacgtc	gtggcaatgt	ggctaatagca	ttaatgagtg	ctactcgact	ggttcgtgat	7320
gttcagtctc	atggcttcaa	tatgactgct	caggaaacagt	ctgtattcca	gatggttact	7380
gctgcattag	caactgaagc	tgcgattgac	ccacatgcta	tggtcgtgct	tcaggaaactt	7440
tatacccatg	taatgaaaca	ccttacggta	gagcatttca	tggtcgaccc	tgatagtagt	7500
aaccctgctg	accgttacta	tgctcaacag	aaatatgaca	ccatctctgg	tgctaatactg	7560
gttgaagtag	atgccaaagg	tagaaccagt	ctgttaccta	cattcctggg	tctggctatg	7620
gttaatgaag	aactacgttc	aatcattaaa	gaaatgcctg	tacctaaagc	agataagaaa	7680
ttagggaaatg	atatagatac	tctgcttacc	aatgcaggta	ctcaggtaat	ggaatctctg	7740
aaccgtcgta	tggtgggtga	ccagaaagct	actaatgttc	aggacagtat	tgatgctttg	7800
tcagaaacaa	tcatggctgc	tgctttgaaa	cgagagtcct	tctatgatgc	tgtagcaacc	7860
ccataccgta	acttcattga	ccgtgcta	cagtacgtaa	cggatagcat	tgaacggttc	7920
tctgaaactg	ttattgagaa	ggcagataag	gtaattgcta	acccttctaa	tatagctgct	7980
aaaggtgttg	ctcatctggc	taaactgact	gctgctattg	catctgaaaa	acaggggtga	8040
atagtggctc	agggtgttat	gactgctatg	aaccagggtg	aagtatggca	acctttccat	8100
gacttagtta	atgacattgt	tggtcgctact	aagactaatg	ccaatgtcta	tgacttaatc	8160
aaattgggta	agagccagat	ttctcaagac	cgtcagcaat	tccgtgagca	tttacctaca	8220
gtcattgctg	gtaagttctc	tcgtaaattg	actgataccg	aatggtctgc	aatgcatact	8280
ggtttaggta	aaacagattt	agctgttcta	cgtgaaacta	tgagcatggc	tgaatttaga	8340
gatttactct	cttcatccaa	gaaagtga	gatgaaatct	ctactctgga	aaaagagatt	8400
cagaaccaag	caggtagaaa	ctggaatctg	gttcagaaga	aatctaagca	actggctcaa	8460
tacatgatta	tggtgggaag	aggtaataac	ctccttcgta	atgcccattg	tattagtcgt	8520
ttgttaggtg	aacgtattac	taatggctct	gtggcagatg	tagctgctat	tgataagctc	8580
attactttgt	actctctgga	attgatgaat	aagtctgacc	gtgacctttt	gtcagaattg	8640
gctcaatcag	aagtgggaag	tatggagttc	tccattgctt	atatgggttg	tcaacgtact	8700
gaagagtagc	gtaaaagcta	agggtgatac	cgtactctgc	tgaatcactt	taaaaggctat	8760
atccctgtag	agaaccagca	agggtggaat	ttgattattg	ctgacgataa	agagtttgct	8820
aagttaaata	gccaatcctt	tactcgtatt	ggtacttatc	aggggagcac	tggtttccgt	8880
actggttcta	aaggttatta	cttcagccca	gtagctgccc	gtgcccctta	ctctcagggt	8940
attcttcaga	acgttcgtaa	tactgctggg	ggtgtggata	ttggtactgg	ctttacgtta	9000
ggcactatgg	ttgctgggag	tattactgac	aaaccaaccg	tagagcgtat	taccaagct	9060

```

ctggctaaag gtagcgctgg gcgtgaacca ctgatgccaa tttataacag caaaggctcag 9120
gtagttgctt atgaacaatc cgttgacctt aatatgttga agcacctaaa ccaagacaat 9180
cactttgcta agatgggttg tgtatggcgt ggctcgtcagg tggaagaggc taaagcacia 9240
cgttttaatg acattctcat tgagcaatta catgctatgt atgagaaaga cattaaagac 9300
tccagtgcta ataaatctca atagttaaac ctggttaggtt aaattgatga cccagtactg 9360
gctgatgcga ttaacctgat gaacattgag actcgtcata aggccgaaga actcttcggt 9420
aaagatgagt tatgggttcg tagggatatg ctgaatgatg cacttggcta tcgtgctgca 9480
tctattggtg atgtgtggac cggtaactct cgttggtcac ctagcaccct tgatactgtt 9540
aagaagatgt tcctcggtgc attcggtaat aaggcatatc atgtagtaat gaatgctgaa 9600
aataccattc agaacttagt gaaggacgct aagacagtaa ttggtgttaa atctgttgta 9660
gtaccggcag ttaacttcct tgctaacatc taccagatga ttggacgtgg tgttcctgtt 9720
aaagatattg ctgtgaacat tcctcgtaag acgtcagaga ttaatcagta tattaatct 9780
cgtttacgtc agattgatgc ggaagcagag ctacgtgctg ctgaaggtaa ccctaactctg 9840
gttcgtaaac ttaaaactga gattcaatct attactgata gtcacgtcg tatgagtatc 9900
tggcctttga ttgaagcagg tgagttctct tctattgctg atgctggtat tagtcgtgat 9960
gacctgttag tagctgaagg taagattcat gagtacatgg aaaaacttgc taataaactt 10020
ccagaaaaag tacgtaatgc tggccgttac gctcttattg ctaaggacac tgctctgttc 10080
caggggtatcc agaaaacagt agagtattca gactttattg ctaaagccat catctatgat 10140
gatttagtga aacgtaagaa aaaatcttct tctgaagcat taggtcaggt aactgaagag 10200
tttattaact atgacagatt gcctggtcgt ttccgtggct atatggaaag tatgggtctg 10260
atgtggttct acaactttta aattcgttcc attaaagttg ctatgagcat gattagaaac 10320
aaccagtagc attctctgat tgctacagta gtacctgctc ctaccatgtt tggtaacgta 10380
ggctctaccaa ttcaggacaa catgctaacc atgctggctg aaggaagact ggattactca 10440
ttaggcttcg gacaaggatt aagagcacct accctcaatc cttggttcaa ccttactcac 10500
taataa 10506

```

<210> 2

<211> 3500

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Peptide

<400> 2

```

Met Ser Val Phe Asp Arg Leu Ala Gly Phe Ala Asp Ser Val Thr Asn
  1              5              10              15

Ala Lys Gln Val Asp Val Ser Thr Ala Thr Ala Gln Lys Lys Ala Glu
      20              25              30

Gln Gly Val Thr Thr Pro Leu Val Ser Pro Asp Ala Ala Tyr Gln Met
      35              40              45

Gln Ala Ala Arg Thr Gly Asn Val Gly Ala Asn Ala Phe Glu Pro Gly
      50              55              60

Thr Val Gln Ser Asp Phe Met Asn Leu Thr Pro Met Gln Ile Met Asn
      65              70              75              80

Lys Tyr Gly Val Glu Gln Gly Leu Gln Leu Ile Asn Ala Arg Ala Asp
      85              90              95

Ala Gly Asn Gln Val Phe Asn Asp Ser Val Thr Thr Arg Thr Pro Gly
      100             105             110

Glu Glu Leu Gly Asp Ile Ala Thr Gly Val Gly Leu Gly Phe Val Asn
      115             120             125

```

Thr Leu Gly Gly Ile Gly Ala Leu Gly Ala Gly Leu Leu Asn Asp Asp
 130 135 140
 Ala Gly Ala Val Val Ala Gln Gln Leu Ser Lys Phe Asn Asp Ala Val
 145 150 155 160
 His Ala Thr Gln Ser Gln Ala Leu Gln Asp Lys Arg Lys Leu Phe Ala
 165 170 175
 Ala Arg Asn Leu Met Asn Glu Val Glu Ser Glu Arg Gln Tyr Gln Thr
 180 185 190
 Asp Lys Lys Glu Gly Thr Asn Asp Ile Val Ala Ser Leu Ser Lys Phe
 195 200 205
 Gly Arg Asp Phe Val Gly Ser Ile Glu Asn Ala Ala Gln Thr Asp Ser
 210 215 220
 Ile Ile Ser Asp Gly Leu Ala Glu Gly Val Gly Ser Leu Leu Gly Ala
 225 230 235 240
 Gly Pro Val Leu Arg Gly Ala Ser Leu Leu Gly Lys Ala Val Val Pro
 245 250 255
 Ala Asn Thr Leu Arg Ser Ala Ala Leu Ala Gly Ala Ile Asp Ala Gly
 260 265 270
 Thr Gly Thr Gln Ser Leu Ala Arg Ile Ala Ser Thr Val Gly Arg Ala
 275 280 285
 Ala Pro Gly Met Val Gly Val Gly Ala Met Glu Ala Gly Gly Ala Tyr
 290 295 300
 Gln Gln Thr Ala Asp Glu Ile Met Lys Met Ser Leu Lys Asp Leu Glu
 305 310 315 320
 Lys Ser Pro Val Tyr Gln Gln His Ile Lys Asp Gly Met Ser Pro Glu
 325 330 335
 Gln Ala Arg Arg Gln Thr Ala Ser Glu Thr Gly Leu Thr Ala Ala Ala
 340 345 350
 Ile Gln Leu Pro Ile Ala Ala Ala Thr Gly Pro Leu Val Ser Arg Phe
 355 360 365
 Glu Met Ala Pro Phe Arg Ala Gly Ser Leu Gly Ala Val Gly Met Asn
 370 375 380
 Leu Ala Arg Glu Thr Val Glu Glu Gly Val Gln Gly Ala Thr Gly Gln
 385 390 395 400
 Leu Ala Gln Asn Ile Ala Gln Gln Gln Asn Ile Asp Lys Asn Gln Asp
 405 410 415
 Leu Leu Lys Gly Val Gly Thr Gln Ala Gly Leu Gly Ala Leu Tyr Gly
 420 425 430
 Phe Gly Ser Ala Gly Val Val Gln Ala Pro Ala Gly Ala Ala Arg Leu
 435 440 445
 Ala Gly Ala Ala Thr Ala Pro Val Leu Arg Thr Thr Met Ala Gly Val

450	455	460
Lys Ala Ala Gly Ser Val Ala Gly Lys Val Val Ser Pro Ile Lys Asn		
465	470	475 480
Thr Leu Val Ala Arg Gly Glu Arg Val Met Lys Gln Asn Glu Glu Ala		
	485 490	495
Ser Pro Val Ala Asp Asp Tyr Val Ala Gln Ala Ala Gln Glu Ala Met		
	500 505	510
Ala Gln Ala Pro Glu Ala Glu Val Thr Ile Arg Asp Ala Val Glu Ala		
	515 520	525
Thr Asp Ala Thr Pro Glu Gln Lys Val Ala Ala His Gln Tyr Val Ser		
	530 535	540
Asp Leu Met Asn Ala Thr Arg Phe Asn Pro Glu Asn Tyr Gln Glu Ala		
	545 550	555 560
Pro Glu His Ile Arg Asn Ala Val Ala Gly Ser Thr Asp Gln Val Gln		
	565 570	575
Val Ile Gln Lys Leu Ala Asp Leu Val Asn Thr Leu Asp Glu Ser Asn		
	580 585	590
Pro Gln Ala Leu Met Glu Ala Ala Ser Tyr Met Tyr Asp Ala Val Ser		
	595 600	605
Glu Phe Glu Gln Phe Ile Asn Arg Asp Pro Ala Ala Leu Asp Ser Ile		
	610 615	620
Pro Lys Asp Ser Pro Ala Ile Glu Leu Leu Asn Arg Tyr Thr Asn Leu		
	625 630	635 640
Thr Ala Asn Ile Gln Asn Thr Pro Lys Val Ile Gly Ala Leu Asn Val		
	645 650	655
Ile Asn Arg Met Ile Asn Glu Ser Ala Gln Asn Gly Ser Leu Asn Val		
	660 665	670
Thr Glu Glu Ser Ser Pro Gln Glu Met Gln Asn Val Ala Leu Ala Ala		
	675 680	685
Glu Val Ala Pro Glu Lys Leu Asn Pro Glu Ser Val Asn Val Val Leu		
	690 695	700
Lys His Ala Ala Asp Gly Arg Ile Lys Leu Asn Asn Arg Gln Ile Ala		
	705 710	715 720
Ala Leu Gln Asn Ala Ala Ala Ile Leu Lys Gly Ala Arg Glu Tyr Asp		
	725 730	735
Ala Glu Ala Ala Arg Leu Gly Leu Arg Pro Gln Asp Ile Val Ser Lys		
	740 745	750
Gln Ile Lys Thr Asp Glu Ser Arg Thr Gln Glu Gly Gln Tyr Ser Ala		
	755 760	765
Leu Gln His Ala Asn Arg Ile Arg Ser Ala Tyr Asn Ser Gly Asn Phe		
	770 775	780

Glu Leu Ala Ser Ala Tyr Leu Asn Asp Phe Met Gln Phe Ala Gln His
 785 790 795 800
 Met Gln Asn Lys Val Gly Ala Leu Asn Glu His Leu Val Thr Gly Asn
 805 810 815
 Ala Asp Lys Asn Lys Ser Val His Tyr Gln Ala Leu Thr Ala Asp Arg
 820 825 830
 Glu Trp Val Arg Ser Arg Thr Gly Leu Gly Val Asn Pro Tyr Asp Thr
 835 840 845
 Lys Ser Val Lys Phe Ala Gln Gln Val Ala Leu Glu Ala Lys Thr Val
 850 855 860
 Ala Asp Ile Ala Asn Ala Leu Ala Ser Ala Tyr Pro Glu Leu Lys Val
 865 870 875 880
 Ser His Ile Lys Val Thr Pro Leu Asp Ser Arg Leu Asn Ala Pro Ala
 885 890 895
 Ala Glu Val Val Lys Ala Phe Arg Gln Gly Asn Arg Asp Val Ala Ser
 900 905 910
 Ser Gln Pro Lys Ala Asp Ser Val Asn Gln Val Lys Glu Thr Pro Val
 915 920 925
 Thr Lys Gln Glu Pro Val Thr Ser Thr Val Gln Thr Lys Thr Pro Val
 930 935 940
 Ser Glu Ser Val Lys Thr Glu Pro Thr Thr Lys Glu Ser Ser Pro Gln
 945 950 955 960
 Ala Ile Lys Glu Pro Val Asn Gln Ser Glu Lys Gln Asp Val Asn Leu
 965 970 975
 Thr Asn Glu Asp Asn Ile Lys Gln Pro Thr Glu Ser Val Lys Glu Thr
 980 985 990
 Glu Thr Ser Thr Lys Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly
 995 1000 1005
 Ile Asp Ala Val Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala
 1010 1015 1020
 Glu Gly Ile Lys Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu
 1025 1030 1035 1040
 Gln Lys Ser Arg Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala
 1045 1050 1055
 Gln Ala Leu Ser Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu
 1060 1065 1070
 Thr Ala Asn Pro Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu
 1075 1080 1085
 Leu Met Glu His Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu
 1090 1095 1100

Ala Lys Phe Leu Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly
 1105 1110 1115 1120
 Thr Glu Ala Asn Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu
 1125 1130 1135
 Gln Asp Gly Asp Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala
 1140 1145 1150
 Val Leu Ala Gly Leu Gln Trp Arg Leu Thr Ala Thr Ser Asn Thr Ala
 1155 1160 1165
 Ile Lys Asp Ala Lys Asp Val Ala Ala Ile Thr Gly Ile Asp Gln Ala
 1170 1175 1180
 Leu Leu Pro Glu Gly Leu Val Glu Gln Phe Asp Thr Gly Met Thr Leu
 1185 1190 1195 1200
 Thr Glu Ala Val Ser Ser Leu Ala Gln Lys Ile Glu Ser Tyr Trp Gly
 1205 1210 1215
 Leu Ser Arg Asn Pro Asn Ala Pro Leu Gly Tyr Thr Lys Gly Ile Pro
 1220 1225 1230
 Thr Ala Met Ala Ala Glu Ile Leu Ala Ala Phe Val Glu Ser Thr Asp
 1235 1240 1245
 Val Val Glu Asn Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys
 1250 1255 1260
 Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro
 1265 1270 1275 1280
 Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro
 1285 1290 1295
 Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr
 1300 1305 1310
 Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala
 1315 1320 1325
 Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val
 1330 1335 1340
 Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly
 1345 1350 1355 1360
 Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser
 1365 1370 1375
 Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe
 1380 1385 1390
 Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val
 1395 1400 1405
 Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met
 1410 1415 1420
 Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile

1425	1430	1435	1440
Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp			
1445		1450	1455
Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val			
1460	1465		1470
His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu			
1475	1480		1485
Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr			
1490	1495	1500	
Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu			
1505	1510	1515	1520
Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser			
1525	1530		1535
Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr			
1540	1545		1550
Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu			
1555	1560		1565
Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys			
1570	1575	1580	
Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg			
1585	1590	1595	1600
Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu			
1605	1610		1615
Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro			
1620	1625		1630
Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu			
1635	1640		1645
Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly			
1650	1655	1660	
Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Tyr Gly Ser Gly Ala Arg			
1665	1670	1675	1680
Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu			
1685	1690		1695
Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser			
1700	1705		1710
Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala			
1715	1720		1725
Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser			
1730	1735	1740	
Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly			
1745	1750	1755	1760

Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly
 1765 1770 1775
 Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu
 1780 1785 1790
 Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr
 1795 1800 1805
 Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile Gln Ser Val Val Leu
 1810 1815 1820
 Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys Leu Ala Glu Lys Ala
 1825 1830 1835 1840
 Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu Thr Gln Lys Glu Leu
 1845 1850 1855
 Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala Pro Met Ile Glu Thr
 1860 1865 1870
 Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu Asn Ala Glu Val Ala
 1875 1880 1885
 Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg Met Arg Val Pro Met
 1890 1895 1900
 Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala Gly Ile Pro Phe Met
 1905 1910 1915 1920
 Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln Thr Leu Ser Thr Met
 1925 1930 1935
 Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe Asp Gly Met Asn Ile
 1940 1945 1950
 Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys Ala Asn Glu Ala Val
 1955 1960 1965
 Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn Val Tyr Glu Ser Tyr
 1970 1975 1980
 Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys Leu Ser Pro Glu Ala
 1985 1990 1995 2000
 Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr Asp Gln Arg Glu Asn
 2005 2010 2015
 Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser Leu Ile Glu Arg Asn
 2020 2025 2030
 Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp
 2035 2040 2045
 Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr
 2050 2055 2060
 Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln
 2065 2070 2075 2080

Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys
 2085 2090 2095
 Gln Lys Val Ala Lys Ala Arg Ala Glu Val Lys Glu Glu Thr Val Ser
 2100 2105 2110
 Glu Lys Glu Pro Val Asn Pro Asp Phe Gly Met Val Gly Arg Glu His
 2115 2120 2125
 Lys Ala Ser Gly Val Arg Ile Leu Ser Ala Thr Ala Ile Arg Asn Leu
 2130 2135 2140
 Ala Lys Ile Ser Asn Leu Pro Ser Thr Gln Ala Ala Thr Leu Ala Glu
 2145 2150 2155 2160
 Ile Gln Lys Ser Leu Ala Ala Lys Asp Tyr Lys Ile Ile Tyr Gly Thr
 2165 2170 2175
 Pro Thr Gln Val Ala Glu Tyr Ala Arg Gln Lys Asn Val Thr Glu Leu
 2180 2185 2190
 Thr Ser Gln Glu Met Glu Glu Ala Gln Ala Gly Asn Ile Tyr Gly Trp
 2195 2200 2205
 Thr Asn Phe Asp Asp Lys Thr Ile Tyr Leu Val Ser Pro Ser Met Glu
 2210 2215 2220
 Thr Leu Ile His Glu Leu Val His Ala Ser Thr Phe Glu Glu Val Tyr
 2225 2230 2235 2240
 Ser Phe Tyr Gln Gly Asn Glu Val Ser Pro Thr Ser Lys Gln Ala Ile
 2245 2250 2255
 Glu Asn Leu Glu Gly Leu Met Glu Gln Phe Arg Ser Leu Asp Ile Ser
 2260 2265 2270
 Lys Asp Ser Pro Glu Met Arg Glu Ala Tyr Ala Asp Ala Ile Ala Thr
 2275 2280 2285
 Ile Glu Gly His Leu Ser Asn Gly Phe Val Asp Pro Ala Ile Ser Lys
 2290 2295 2300
 Ala Ala Ala Leu Asn Glu Phe Met Ala Trp Gly Leu Ala Asn Arg Ala
 2305 2310 2315 2320
 Leu Ala Ala Lys Gln Lys Arg Thr Ser Ser Leu Val Gln Met Val Lys
 2325 2330 2335
 Asp Val Tyr Gln Ala Ile Lys Lys Leu Ile Trp Gly Arg Lys Gln Ala
 2340 2345 2350
 Pro Ala Leu Gly Glu Asp Met Phe Ser Asn Leu Leu Phe Asn Ser Ala
 2355 2360 2365
 Ile Leu Met Arg Ser Gln Pro Thr Thr Gln Ala Val Ala Lys Asp Gly
 2370 2375 2380
 Thr Leu Phe His Ser Lys Ala Tyr Gly Asn Asn Glu Arg Leu Ser Gln
 2385 2390 2395 2400
 Leu Asn Gln Thr Phe Asp Lys Leu Val Thr Asp Tyr Leu Arg Thr Asp

2405	2410	2415
Pro Val Thr Glu Val Glu Arg Arg Gly Asn Val Ala Asn Ala Leu Met		
2420	2425	2430
Ser Ala Thr Arg Leu Val Arg Asp Val Gln Ser His Gly Phe Asn Met		
2435	2440	2445
Thr Ala Gln Glu Gln Ser Val Phe Gln Met Val Thr Ala Ala Leu Ala		
2450	2455	2460
Thr Glu Ala Ala Ile Asp Pro His Ala Met Ala Arg Ala Gln Glu Leu		
2465	2470	2480
Tyr Thr His Val Met Lys His Leu Thr Val Glu His Phe Met Ala Asp		
2485	2490	2495
Pro Asp Ser Thr Asn Pro Ala Asp Arg Tyr Tyr Ala Gln Gln Lys Tyr		
2500	2505	2510
Asp Thr Ile Ser Gly Ala Asn Leu Val Glu Val Asp Ala Lys Gly Arg		
2515	2520	2525
Thr Ser Leu Leu Pro Thr Phe Leu Gly Leu Ala Met Val Asn Glu Glu		
2530	2535	2540
Leu Arg Ser Ile Ile Lys Glu Met Pro Val Pro Lys Ala Asp Lys Lys		
2545	2550	2560
Leu Gly Asn Asp Ile Asp Thr Leu Leu Thr Asn Ala Gly Thr Gln Val		
2565	2570	2575
Met Glu Ser Leu Asn Arg Arg Met Ala Gly Asp Gln Lys Ala Thr Asn		
2580	2585	2590
Val Gln Asp Ser Ile Asp Ala Leu Ser Glu Thr Ile Met Ala Ala Ala		
2595	2600	2605
Leu Lys Arg Glu Ser Phe Tyr Asp Ala Val Ala Thr Pro Thr Gly Asn		
2610	2615	2620
Phe Ile Asp Arg Ala Asn Gln Tyr Val Thr Asp Ser Ile Glu Arg Leu		
2625	2630	2640
Ser Glu Thr Val Ile Glu Lys Ala Asp Lys Val Ile Ala Asn Pro Ser		
2645	2650	2655
Asn Ile Ala Ala Lys Gly Val Ala His Leu Ala Lys Leu Thr Ala Ala		
2660	2665	2670
Ile Ala Ser Glu Lys Gln Gly Glu Ile Val Ala Gln Gly Val Met Thr		
2675	2680	2685
Ala Met Asn Gln Gly Lys Val Trp Gln Pro Phe His Asp Leu Val Asn		
2690	2695	2700
Asp Ile Val Gly Arg Thr Lys Thr Asn Ala Asn Val Tyr Asp Leu Ile		
2705	2710	2720
Lys Leu Val Lys Ser Gln Ile Ser Gln Asp Arg Gln Gln Phe Arg Glu		
2725	2730	2735

His Leu Pro Thr Val Ile Ala Gly Lys Phe Ser Arg Lys Leu Thr Asp
 2740 2745 2750
 Thr Glu Trp Ser Ala Met His Thr Gly Leu Gly Lys Thr Asp Leu Ala
 2755 2760 2765
 Val Leu Arg Glu Thr Met Ser Met Ala Glu Ile Arg Asp Leu Leu Ser
 2770 2775 2780
 Ser Ser Lys Lys Val Lys Asp Glu Ile Ser Thr Leu Glu Lys Glu Ile
 2785 2790 2795 2800
 Gln Asn Gln Ala Gly Arg Asn Trp Asn Leu Val Gln Lys Lys Ser Lys
 2805 2810 2815
 Gln Leu Ala Gln Tyr Met Ile Met Gly Glu Val Gly Asn Asn Leu Leu
 2820 2825 2830
 Arg Asn Ala His Ala Ile Ser Arg Leu Leu Gly Glu Arg Ile Thr Asn
 2835 2840 2845
 Gly Pro Val Ala Asp Val Ala Ala Ile Asp Lys Leu Ile Thr Leu Tyr
 2850 2855 2860
 Ser Leu Glu Leu Met Asn Lys Ser Asp Arg Asp Leu Leu Ser Glu Leu
 2865 2870 2875 2880
 Ala Gln Ser Glu Val Glu Gly Met Glu Phe Ser Ile Ala Tyr Met Val
 2885 2890 2895
 Gly Gln Arg Thr Glu Glu Met Arg Lys Ala Lys Gly Asp Asn Arg Thr
 2900 2905 2910
 Leu Leu Asn His Phe Lys Gly Tyr Ile Pro Val Glu Asn Gln Gln Gly
 2915 2920 2925
 Val Asn Leu Ile Ile Ala Asp Asp Lys Glu Phe Ala Lys Leu Asn Ser
 2930 2935 2940
 Gln Ser Phe Thr Arg Ile Gly Thr Tyr Gln Gly Ser Thr Gly Phe Arg
 2945 2950 2955 2960
 Thr Gly Ser Lys Gly Tyr Tyr Phe Ser Pro Val Ala Ala Arg Ala Pro
 2965 2970 2975
 Tyr Ser Gln Gly Ile Leu Gln Asn Val Arg Asn Thr Ala Gly Gly Val
 2980 2985 2990
 Asp Ile Gly Thr Gly Phe Thr Leu Gly Thr Met Val Ala Gly Arg Ile
 2995 3000 3005
 Thr Asp Lys Pro Thr Val Glu Arg Ile Thr Lys Ala Leu Ala Lys Gly
 3010 3015 3020
 Glu Arg Gly Arg Glu Pro Leu Met Pro Ile Tyr Asn Ser Lys Gly Gln
 3025 3030 3035 3040
 Val Val Ala Tyr Glu Gln Ser Val Asp Pro Asn Met Leu Lys His Leu
 3045 3050 3055

Asn Gln Asp Asn His Phe Ala Lys Met Val Gly Val Trp Arg Gly Arg
 3060 3065 3070
 Gln Val Glu Glu Ala Lys Ala Gln Arg Phe Asn Asp Ile Leu Ile Glu
 3075 3080 3085
 Gln Leu His Ala Met Tyr Glu Lys Asp Ile Lys Asp Ser Ser Ala Asn
 3090 3095 3100
 Lys Ser Gln Tyr Val Asn Leu Leu Gly Lys Ile Asp Asp Pro Val Leu
 3105 3110 3115 3120
 Ala Asp Ala Ile Asn Leu Met Asn Ile Glu Thr Arg His Lys Ala Glu
 3125 3130 3135
 Glu Leu Phe Gly Lys Asp Glu Leu Trp Val Arg Arg Asp Met Leu Asn
 3140 3145 3150
 Asp Ala Leu Gly Tyr Arg Ala Ala Ser Ile Gly Asp Val Trp Thr Gly
 3155 3160 3165
 Asn Ser Arg Trp Ser Pro Ser Thr Leu Asp Thr Val Lys Lys Met Phe
 3170 3175 3180
 Leu Gly Ala Phe Gly Asn Lys Ala Tyr His Val Val Met Asn Ala Glu
 3185 3190 3195 3200
 Asn Thr Ile Gln Asn Leu Val Lys Asp Ala Lys Thr Val Ile Val Val
 3205 3210 3215
 Lys Ser Val Val Val Pro Ala Val Asn Phe Leu Ala Asn Ile Tyr Gln
 3220 3225 3230
 Met Ile Gly Arg Gly Val Pro Val Lys Asp Ile Ala Val Asn Ile Pro
 3235 3240 3245
 Arg Lys Thr Ser Glu Ile Asn Gln Tyr Ile Lys Ser Arg Leu Arg Gln
 3250 3255 3260
 Ile Asp Ala Glu Ala Glu Leu Arg Ala Ala Glu Gly Asn Pro Asn Leu
 3265 3270 3275 3280
 Val Arg Lys Leu Lys Thr Glu Ile Gln Ser Ile Thr Asp Ser His Arg
 3285 3290 3295
 Arg Met Ser Ile Trp Pro Leu Ile Glu Ala Gly Glu Phe Ser Ser Ile
 3300 3305 3310
 Ala Asp Ala Gly Ile Ser Arg Asp Asp Leu Leu Val Ala Glu Gly Lys
 3315 3320 3325
 Ile His Glu Tyr Met Glu Lys Leu Ala Asn Lys Leu Pro Glu Lys Val
 3330 3335 3340
 Arg Asn Ala Gly Arg Tyr Ala Leu Ile Ala Lys Asp Thr Ala Leu Phe
 3345 3350 3355 3360
 Gln Gly Ile Gln Lys Thr Val Glu Tyr Ser Asp Phe Ile Ala Lys Ala
 3365 3370 3375
 Ile Ile Tyr Asp Asp Leu Val Lys Arg Lys Lys Lys Ser Ser Ser Glu

<400>	3						
gaaagtacag	ttacagaaga	attaaaagaa	ggtattgatg	ctgtttaccc	ttcattggta	60	
ggtactgctg	attctaaagc	agagggtatt	aagaactatt	tcaaattgtc	ctttacctta	120	
ccagaagaac	agaaatcccg	tactgttggt	tcagaagcac	ctctaaaaga	tgtagcccaa	180	
gctctgtctt	ctcgtgctcg	ttatgaactc	tttactgaga	aagaaactgc	taaccctgct	240	
tttaatgggg	aagttattaa	gcgatacaaa	gaactcatgg	aacatgggga	aggtattgct	300	
gatattcttc	gctcccgtct	ggctaagttc	cttaacacta	aggatgttgg	taaacgtttt	360	
gctcaaggta	cagaagccaa	ccgttgggta	ggtggtaagt	tacttaacat	tgttgagcag	420	
gatggggata	cctttaagta	caacgaacaa	ttgctacaga	ctgctgtatt	agcagggtctt	480	
caatggagac	ttactgctac	cagcaatact	gctatcaaag	atgcaaaaaga	tgttgctgct	540	
attactggta	ttgaccaagc	tctgctgcca	gaagggttag	tagagcaatt	tgatactggt	600	
atgacactca	ctgaagcagt	tagttccctg	gctcagaaaa	ttgagcttta	ctggggatta	660	
tctcgtaatc	caaatgcttc	attgggctat	accaaggcca	tccctacagc	aatggctgct	720	
gaaattcttg	ctgcatttgt	agagtctact	gatgttgtag	agaacatcgt	ggatatgtca	780	
gaaattgacc	cagataacaa	gaagactatt	ggtctgtaca	ccattactga	actggattcc	840	
ttcgacccaa	ttaatagctt	ccctactgct	attgaagaag	ctgttttagt	gaatcctaca	900	
gagaagatgt	tctttgggtg	tgacattcct	cctgtagcta	atactcagct	tcgtaaccct	960	
gctgttcgta	atactccaga	acagaaggct	gcattgaaag	cagagcaggc	tacagagttc	1020	
tatgtacaca	ccccaatggt	tcaattctat	gagacgttag	gtaaagaccg	tattctcgaa	1080	
ctgatgggtg	ctgggtactct	gaataaagag	ttacttaatg	ataaccatgc	taaatctctg	1140	
gaaggtaaga	accgttcagt	agaggactct	tacaaccaac	tgttctccgt	cattgagcag	1200	
gtaagagcac	agagcgaaga	catctctact	gtacctattc	actatgcata	caatatgacc	1260	
cgtgttggtc	gtatgcagat	gttaggtaaa	tacaatcctc	aatcagccaa	actggttcgt	1320	
gaggccatct	tacctactaa	agctactttg	gatttatcga	accagaacaa	tgaagacttc	1380	
tctgcattcc	agttaggctc	ggctcaggca	ttggacatta	aatgccatac	tatgactcgt	1440	
gaggttatgt	ctgacgagtt	gactaaaatta	ctggaaggta	atctgaaacc	agccattgat	1500	
atgatggttg	agtttaatac	cactgggttc	ttaccagaaa	acgcagttga	tgttctgaat	1560	
acagcattag	gagataggaa	gtcattcgta	gcattgatgg	ctcttatgga	qtattcccqt	1620	

```

tacttagtag cagaggataa atctgcattt gtaactccac tgtatgtaga agcagatggg 1680
gttactaatg gtccaatcaa tgccatgatg ctaatgacag gcggtctgtt tactcctgac 1740
tggattcgta atattgccaa agggggcctt ttcattgggt ctccaaataa gaccatgaat 1800
gagcatcgct ctactgctga caataatgat ttatatcaag catccactaa tgctttgatg 1860
gaatcgttgg gtaagttacg tagtaactat gcctctaata tgctatttca gtctcagata 1920
gacagtcctt tttctctgat ggatttggtt ttaccggata ttaatcttgg tgagaatggg 1980
gctttagaac ttaaactgtg tattgctaag aaccactga ctattaccat ctatgggtct 2040
gggtgctctg gtattgcagg taagctgggt agttctgtta ctgatgccat ctatgagcgt 2100
atgtctgatg tactgaaagc tcgtgctaaa gacccaaata tctctgctgc tatggcaatg 2160
tttggttaagc aagctgcttc agaagcacat gctgaagaac ttcttgcccg tttcctgaaa 2220
gatattgaaa cactgacttc tactgttcct gttaaacgta aaggtgtact ggaactacaa 2280
tccacaggta caggagccaa aggaaaaatc aatcctaaga cctataccat taagggcgag 2340
caactgaagg cacttcagga aaatatgctg cacttctttg tagaaccact acgtaatggg 2400
attactcaga ctgtaggtga aagtctgggt tactctactg aacaattaca gaaagctact 2460
cagattcaat ctgtagtgct ggaagatatg ttcaaacagc gagtacaaga gaagctggca 2520
gagaaggcta aagacccaac atggaagaaa ggtgatttcc ttactcagaa agaactgaat 2580
gatattcagg cttctctgaa taacttagcc cctatgattg agactgggtc tcagactttc 2640
tacattgctg gttcagaaaa tgcagaagta gcaaactcagg tattagctac taaccttgat 2700
gaccgtatgc gtgtaccaat gagtatctat gctccagcac aggcgggtgt agcagggtatt 2760
ccatttatga ctattggtac tgggtgatggc atgatgatgc aaactctttc cactatgaaa 2820
gggtgcacaa agaataccct caaaatcttt gatgggtatga acattgggtt gaatgacatc 2880
actgatgcca gtcgtaaagc taatgaagct gtttactact cttggcaggg taaccctatt 2940
aagaatgttt atgaatcata tgctaagttc atgaagaatg tagatttcag caagctgtcc 3000
cctgaagcat tggaagcaat tggtaaactc gctctggaat atgaccaacg tgagaatgct 3060
actgtagatg atattgctaa cgctgcatct ctgattgaac gtaacttacg taatattgca 3120
ctgggtgtag atattcgctc taaggtgctg gataaggtaa atctgtccat tgaccagatg 3180
gctgctgtag gtgctcctta tcagaacaac ggtaagattg acctcagcaa tatgaccctt 3240
gaacaacagg ctgatgaact gaataaactt ttccgtgaag agttagaagc ccgtaaacaa 3300
aaagtcgcta aggctagg                                     3318

```

<210> 4

<211> 1107

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic Peptide

<400> 4

```

Met Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly Ile Asp Ala Val
 1              5              10              15

Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala Glu Gly Ile Lys
 20              25              30

Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu Gln Lys Ser Arg
 35              40              45

Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala Gln Ala Leu Ser
 50              55              60

Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu Thr Ala Asn Pro
 65              70              75              80

Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu Leu Met Glu His
 85              90              95

Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu Ala Lys Phe Leu
100              105              110

```

Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly Thr Glu Ala Asn
 115 120 125
 Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu Gln Asp Gly Asp
 130 135 140
 Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala Val Leu Ala Gly
 145 150 155 160
 Leu Gln Trp Arg Leu Thr Ala Thr Ser Asn Thr Ala Ile Lys Asp Ala
 165 170 175
 Lys Asp Val Ala Ala Ile Thr Gly Ile Asp Gln Ala Leu Leu Pro Glu
 180 185 190
 Gly Leu Val Glu Gln Phe Asp Thr Gly Met Thr Leu Thr Glu Ala Val
 195 200 205
 Ser Ser Leu Ala Gln Lys Ile Glu Ser Tyr Trp Gly Leu Ser Arg Asn
 210 215 220
 Pro Asn Ala Pro Leu Gly Tyr Thr Lys Gly Ile Pro Thr Ala Met Ala
 225 230 235 240
 Ala Glu Ile Leu Ala Ala Phe Val Glu Ser Thr Asp Val Val Glu Asn
 245 250 255
 Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys Lys Thr Ile Gly
 260 265 270
 Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro Ile Asn Ser Phe
 275 280 285
 Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro Thr Glu Lys Met
 290 295 300
 Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr Gln Leu Arg Asn
 305 310 315 320
 Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala Leu Lys Ala Glu
 325 330 335
 Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val Gln Phe Tyr Glu
 340 345 350
 Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly Ala Gly Thr Leu
 355 360 365
 Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser Leu Glu Gly Lys
 370 375 380
 Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe Ser Val Ile Glu
 385 390 395 400
 Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val Pro Ile His Tyr
 405 410 415
 Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met Leu Gly Lys Tyr
 420 425 430

Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile Leu Pro Thr Lys
 435 440 445
 Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp Phe Ser Ala Phe
 450 455 460
 Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val His Thr Met Thr
 465 470 475 480
 Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu Glu Gly Asn Leu
 485 490 495
 Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr Thr Gly Ser Leu
 500 505 510
 Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu Gly Asp Arg Lys
 515 520 525
 Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser Arg Tyr Leu Val
 530 535 540
 Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr Val Glu Ala Asp
 545 550 555 560
 Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu Met Thr Gly Gly
 565 570 575
 Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys Gly Gly Leu Phe
 580 585 590
 Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg Ser Thr Ala Asp
 595 600 605
 Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu Met Glu Ser Leu
 610 615 620
 Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro Ile Gln Ser Gln
 625 630 635 640
 Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu Pro Asp Ile Asn
 645 650 655
 Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly Ile Ala Lys Asn
 660 665 670
 Pro Leu Thr Ile Thr Ile Tyr Gly Ser Gly Ala Arg Gly Ile Ala Gly
 675 680 685
 Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu Arg Met Ser Asp
 690 695 700
 Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser Ala Ala Met Ala
 705 710 715 720
 Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala Glu Glu Leu Leu
 725 730 735
 Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser Thr Val Pro Val
 740 745 750
 Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly Thr Gly Ala Lys

755					760					765					
Gly	Lys	Ile	Asn	Pro	Lys	Thr	Tyr	Thr	Ile	Lys	Gly	Glu	Gln	Leu	Lys
770						775					780				
Ala	Leu	Gln	Glu	Asn	Met	Leu	His	Phe	Phe	Val	Glu	Pro	Leu	Arg	Asn
785					790					795					800
Gly	Ile	Thr	Gln	Thr	Val	Gly	Glu	Ser	Leu	Val	Tyr	Ser	Thr	Glu	Gln
				805					810					815	
Leu	Gln	Lys	Ala	Thr	Gln	Ile	Gln	Ser	Val	Val	Leu	Glu	Asp	Met	Phe
			820					825					830		
Lys	Gln	Arg	Val	Gln	Glu	Lys	Leu	Ala	Glu	Lys	Ala	Lys	Asp	Pro	Thr
		835					840					845			
Trp	Lys	Lys	Gly	Asp	Phe	Leu	Thr	Gln	Lys	Glu	Leu	Asn	Asp	Ile	Gln
850						855					860				
Ala	Ser	Leu	Asn	Asn	Leu	Ala	Pro	Met	Ile	Glu	Thr	Gly	Ser	Gln	Thr
865					870					875					880
Phe	Tyr	Ile	Ala	Gly	Ser	Glu	Asn	Ala	Glu	Val	Ala	Asn	Gln	Val	Leu
				885					890					895	
Ala	Thr	Asn	Leu	Asp	Asp	Arg	Met	Arg	Val	Pro	Met	Ser	Ile	Tyr	Ala
			900					905					910		
Pro	Ala	Gln	Ala	Gly	Val	Ala	Gly	Ile	Pro	Phe	Met	Thr	Ile	Gly	Thr
		915					920					925			
Gly	Asp	Gly	Met	Met	Met	Gln	Thr	Leu	Ser	Thr	Met	Lys	Gly	Ala	Pro
	930					935					940				
Lys	Asn	Thr	Leu	Lys	Ile	Phe	Asp	Gly	Met	Asn	Ile	Gly	Leu	Asn	Asp
945					950					955					960
Ile	Thr	Asp	Ala	Ser	Arg	Lys	Ala	Asn	Glu	Ala	Val	Tyr	Thr	Ser	Trp
			965						970					975	
Gln	Gly	Asn	Pro	Ile	Lys	Asn	Val	Tyr	Glu	Ser	Tyr	Ala	Lys	Phe	Met
			980					985					990		
Lys	Asn	Val	Asp	Phe	Ser	Lys	Leu	Ser	Pro	Glu	Ala	Leu	Glu	Ala	Ile
		995					1000					1005			
Gly	Lys	Ser	Ala	Leu	Glu	Tyr	Asp	Gln	Arg	Glu	Asn	Ala	Thr	Val	Asp
	1010					1015					1020				
Asp	Ile	Ala	Asn	Ala	Ala	Ser	Leu	Ile	Glu	Arg	Asn	Leu	Arg	Asn	Ile
1025				1030					1035					1040	
Ala	Leu	Gly	Val	Asp	Ile	Arg	His	Lys	Val	Leu	Asp	Lys	Val	Asn	Leu
			1045						1050				1055		
Ser	Ile	Asp	Gln	Met	Ala	Ala	Val	Gly	Ala	Pro	Tyr	Gln	Asn	Asn	Gly
		1060						1065				1070			
Lys	Ile	Asp	Leu	Ser	Asn	Met	Thr	Pro	Glu	Gln	Gln	Ala	Asp	Glu	Leu
	1075					1080						1085			

Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys Gln Lys Val Ala
 1090 1095 1100

Lys Ala Arg
 1105

<210> 5

<211> 3432

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 5

```

atgggggggtt ctcacatca tcatcatcat ggtatggcta gcatgactgg tggacagcaa 60
atgggtcggg atctgtacga cgatgacgat aaggatccga gctcgagatc tgaaagtaca 120
gttacagaag aattaaaaga aggtattgat gctgtttacc cttcattggg aggtactgct 180
gattctaaag cagagggtat taagaactat ttcaaattgt cttttacctt accagaagaa 240
cagaaatccc gtactgttgg ttcagaagca cctctaaaag atgtagccca agctctgtct 300
tctcgtgctc gttatgaact ctttactgag aaagaaactg ctaaccctgc ttttaattggg 360
gaagttatta agcgatacaa agaactcatg gaacatgggg aaggatttgc tgatattctt 420
cgctcccgtc tggctaagtt ccttaacact aaggatgttg gtaaacgttt tgctcaagggt 480
acagaagcca accgttgggt aggtggtaag ttacttaaca ttggtgagca ggatggggat 540
acctttaagt acaacgaaca attgctacag actgctgtat tagcaggctc tcaatggaga 600
cttactgcta ccagcaatac tgctatcaaa gatgcaaaag atgttgctgc tattactggt 660
attgaccaag cctctgctgc agaaggttta gtagagcaat ttgatactgg tatgacactc 720
actgaagcag ttagttccct ggctcagaaa attgagtctt actggggatt atctcgtaat 780
ccaaatgctc cattgggcta taccaaaggc atccctacag caatggctgc tgaaattctg 840
gctgcatttg tagagtctac tgatgttgta gagaacatcg tggatatgtc agaaattgac 900
ccagataaca agaagactat tggctgttac accattactg aactggattc cttcgacca 960
attaatagct tccctactgc tattgaagaa gctgttttag tgaatcctac agagaagatg 1020
ttctttgggtg atgacattcc tcctgtagct aatactcagc ttcgtaacct tgctgttcgt 1080
aatactccag aacagaaggc tgcattgaaa gcagagcagg ctacagagtt ctatgtacac 1140
acccaatgg ttcaattcta tgagacgtta ggtaaagacc gtattctcga actgatgggt 1200
gctgggtactc tgaataaaga gttacttaat gataaccatg ctaaattctc ggaaggtaag 1260
aaccgttcag tagaggactc ttacaaccaa ctgttctccg tcattgagca ggtaagagca 1320
cagagcgaag acatctctac tgtacctatt cactatgcac acaatatgac ccgtgttggg 1380
cgtatgcaga gttaggttaa atacaatcct caatcagcca aactggttcg tgaggccatc 1440
ttactacta aagctacttt ggatttatcg aaccagaaca atgaagactt ctctgcattc 1500
cagttaggtc tggctcaggc attggacatt aaagtccata ctatgactcg tgaggttatg 1560
tctgacgagt tgactaaatt actggaagggt aatctgaaac cagccattga tatgatgggt 1620
gagtttaata ccaactggttc cttaccagaa aacgcagttg atgttctgaa tacagcatta 1680
ggagatagga agtcattcgt agcattgatg gctcttatgg agtattcccg ttacttagta 1740
gcagaggata aatctgcatt tgtaactcca ctgtatgtag aagcagatgg tggtactaat 1800
ggtccaatca atgccatgat gctaattgaca ggcggtctgt ttactcctga ctggattcgt 1860
aatattgcca aaggggggctt gttcattgggt tctccaaata agaccatgaa tgagcatcgc 1920
tctactgctg acaataatga tttatatcaa gcatccacta atgctttgat ggaatcgttg 1980
ggtaagttac gtagtaacta tgcctcta atgcctattc agtctcagat agacagtctt 2040
ctttctctga tggatttggt tttaccggat attaattctg gtgagaatgg tgcttagaa 2100
cttaaacgtg gtattgctaa gaaccactg actattacca tctatggttc tgggtcctgt 2160
ggatttgag gtaagctggt tagttctggt actgatgcca tctatgagcg tatgtctgat 2220
gtactgaaag ctcgtgctaa agacccaaat atctctgctg ctatggcaat gtttggttaag 2280
caagctgctt cagaagcaca tgctgaagaa cttcttgccc gtttcctgaa agatatggaa 2340
acactgactt ctactgttcc tgttaaacgt aaagggtgtac tggaaactaca atccacagg 2400
acaggagcca aaggaaaaat caatcctaag acctatacca ttaagggcga gcaactgaag 2460
gcacttcagg aaaatatgct gcacttcttt gtagaaccac tacgtaatgg tattactcag 2520
actgtagggtg aaagtctggt gtactctact gaacaattac agaaagctac tcagattcaa 2580

```

```

tctgtagtgc tggagatat gttcaaacag cgagtacaag agaagctggc agagaaggct 2640
aaagacccaa catggaagaa aggtgatttc cttactcaga aagaactgaa tgatattcag 2700
gcttctctga ataacttagc ccctatgatt gagactgggt ctcagacttt ctacattgct 2760
ggttcagaaa atgcagaagt agcaaatacag gtattagcta ctaaccttga tgaccgtatg 2820
cgtgtaccaa tgagtatcta tgctccagca caggccgggt tagcagggtat tccatttatg 2880
actattggta ctgggtgatg catgatgatg caaactcttt ccactatgaa aggtgcacca 2940
aagaataccc tcaaaatctt tgatgggatg aacattgggt tgaatgacat cactgatgcc 3000
agtcgtaaag ctaatgaagc tgtttacact tcttggcagg gtaaccctat taagaatggt 3060
tatgaatcat atgctaagtt catgaagaat gtagatttca gcaagctgtc ccctgaagca 3120
ttggaagcaa ttggtaaata tgctctggaa tatgaccaac gtgagaatgc tactgtagat 3180
gatattgcta acgctgcata tctgattgaa cgtaacttac gtaatatgtc actgggtgta 3240
gatattcgtc ataaggtgct ggataaggta aatctgtcca ttgaccagat ggctgctgta 3300
gggtgctcctt atcagaacaa cggtaagatt gacctcagca atatgacccc tgaacaacag 3360
gctgatgaac tgaataaact tttccgtgaa gagttagaag cccgtaaaca aaaagtcgct 3420
aaggctaggt aa 3432

```

<210> 6

<211> 1143

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic Peptide

<400> 6

```

Met Gly Gly Ser His His His His His His Gly Met Ala Ser Met Thr
  1             5             10             15

```

```

Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp
          20             25             30

```

```

Pro Ser Ser Arg Ser Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly
      35             40             45

```

```

Ile Asp Ala Val Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala
  50             55             60

```

```

Glu Gly Ile Lys Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu
  65             70             75             80

```

```

Gln Lys Ser Arg Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala
      85             90             95

```

```

Gln Ala Leu Ser Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu
      100            105            110

```

```

Thr Ala Asn Pro Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu
      115            120            125

```

```

Leu Met Glu His Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu
      130            135            140

```

```

Ala Lys Phe Leu Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly
      145            150            155            160

```

```

Thr Glu Ala Asn Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu
      165            170            175

```

```

Gln Asp Gly Asp Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala

```

180					185					190					
Val	Leu	Ala	Gly	Leu	Gln	Trp	Arg	Leu	Thr	Ala	Thr	Ser	Asn	Thr	Ala
	195						200					205			
Ile	Lys	Asp	Ala	Lys	Asp	Val	Ala	Ala	Ile	Thr	Gly	Ile	Asp	Gln	Ala
	210					215					220				
Leu	Leu	Pro	Glu	Gly	Leu	Val	Glu	Gln	Phe	Asp	Thr	Gly	Met	Thr	Leu
	225					230					235				240
Thr	Glu	Ala	Val	Ser	Ser	Leu	Ala	Gln	Lys	Ile	Glu	Ser	Tyr	Trp	Gly
				245					250					255	
Leu	Ser	Arg	Asn	Pro	Asn	Ala	Pro	Leu	Gly	Tyr	Thr	Lys	Gly	Ile	Pro
			260					265					270		
Thr	Ala	Met	Ala	Ala	Glu	Ile	Leu	Ala	Ala	Phe	Val	Glu	Ser	Thr	Asp
		275					280					285			
Val	Val	Glu	Asn	Ile	Val	Asp	Met	Ser	Glu	Ile	Asp	Pro	Asp	Asn	Lys
	290					295					300				
Lys	Thr	Ile	Gly	Leu	Tyr	Thr	Ile	Thr	Glu	Leu	Asp	Ser	Phe	Asp	Pro
	305					310					315				320
Ile	Asn	Ser	Phe	Pro	Thr	Ala	Ile	Glu	Glu	Ala	Val	Leu	Val	Asn	Pro
				325					330					335	
Thr	Glu	Lys	Met	Phe	Phe	Gly	Asp	Asp	Ile	Pro	Pro	Val	Ala	Asn	Thr
			340					345					350		
Gln	Leu	Arg	Asn	Pro	Ala	Val	Arg	Asn	Thr	Pro	Glu	Gln	Lys	Ala	Ala
		355					360					365			
Leu	Lys	Ala	Glu	Gln	Ala	Thr	Glu	Phe	Tyr	Val	His	Thr	Pro	Met	Val
	370					375					380				
Gln	Phe	Tyr	Glu	Thr	Leu	Gly	Lys	Asp	Arg	Ile	Leu	Glu	Leu	Met	Gly
	385					390					395				400
Ala	Gly	Thr	Leu	Asn	Lys	Glu	Leu	Leu	Asn	Asp	Asn	His	Ala	Lys	Ser
				405					410					415	
Leu	Glu	Gly	Lys	Asn	Arg	Ser	Val	Glu	Asp	Ser	Tyr	Asn	Gln	Leu	Phe
			420					425					430		
Ser	Val	Ile	Glu	Gln	Val	Arg	Ala	Gln	Ser	Glu	Asp	Ile	Ser	Thr	Val
		435					440					445			
Pro	Ile	His	Tyr	Ala	Tyr	Asn	Met	Thr	Arg	Val	Gly	Arg	Met	Gln	Met
	450					455					460				
Leu	Gly	Lys	Tyr	Asn	Pro	Gln	Ser	Ala	Lys	Leu	Val	Arg	Glu	Ala	Ile
	465					470					475				480
Leu	Pro	Thr	Lys	Ala	Thr	Leu	Asp	Leu	Ser	Asn	Gln	Asn	Asn	Glu	Asp
			485						490					495	
Phe	Ser	Ala	Phe	Gln	Leu	Gly	Leu	Ala	Gln	Ala	Leu	Asp	Ile	Lys	Val
			500				505						510		

His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu
 515 520 525
 Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr
 530 535 540
 Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu
 545 550 555 560
 Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser
 565 570 575
 Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr
 580 585 590
 Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu
 595 600 605
 Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys
 610 615 620
 Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg
 625 630 635 640
 Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu
 645 650 655
 Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro
 660 665 670
 Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu
 675 680 685
 Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly
 690 695 700
 Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Tyr Gly Ser Gly Ala Arg
 705 710 715 720
 Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu
 725 730 735
 Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser
 740 745 750
 Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala
 755 760 765
 Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser
 770 775 780
 Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly
 785 790 795 800
 Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly
 805 810 815
 Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu
 820 825 830

Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr
 835 840 845
 Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile Gln Ser Val Val Leu
 850 855 860
 Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys Leu Ala Glu Lys Ala
 865 870 875 880
 Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu Thr Gln Lys Glu Leu
 885 890 895
 Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala Pro Met Ile Glu Thr
 900 905 910
 Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu Asn Ala Glu Val Ala
 915 920 925
 Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg Met Arg Val Pro Met
 930 935 940
 Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala Gly Ile Pro Phe Met
 945 950 955 960
 Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln Thr Leu Ser Thr Met
 965 970 975
 Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe Asp Gly Met Asn Ile
 980 985 990
 Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys Ala Asn Glu Ala Val
 995 1000 1005
 Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn Val Tyr Glu Ser Tyr
 1010 1015 1020
 Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys Leu Ser Pro Glu Ala
 1025 1030 1035 1040
 Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr Asp Gln Arg Glu Asn
 1045 1050 1055
 Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser Leu Ile Glu Arg Asn
 1060 1065 1070
 Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp
 1075 1080 1085
 Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr
 1090 1095 1100
 Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln
 1105 1110 1115 1120
 Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys
 1125 1130 1135
 Gln Lys Val Ala Lys Ala Arg
 1140

<210> 7

<211> 3432

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 7

```

atgggggggtt ctcacatcatca tcacatcatcat ggtatgggcta gcatgactgg tggacagcaa 60
atggggtcggg atctgtacga cgatgacgat aaggatccga gctcgagatc tgaaagtaca 120
gttacagaag aattaaaaga aggtattgat gctgtttacc cttcattggg aggtactgct 180
gattctaaag cagaggggtat taagaactat ttcaaattgt cctttacctt accagaagaa 240
cagaaatccc gtactgttgg ttcagaagca cctctaaaag atgtagccca agctctgtct 300
tctcgtgctc gttatgaact ctttactgag aaagaaactg ctaaccctgc ttttaattggg 360
gaagttatta agcgatacaa agaactcatg gaacatgggg aaggatttgc tgatattctt 420
cgctcccgtc tggctaagtt ccttaacact aaggatgttg gtaaaccgtt tgctcaaggt 480
acagaagcca accggtgggt aggtggtaag ttacttaaca ttgttgagca ggatggggat 540
acctttaagt acaacgaaca attgctacag actgctgtat tagcaggtct tcaatggaga 600
cttactgcta ccagcaatac tgctatcaaa gatgcaaaaag atgttgctgc tattactggg 660
attgaccaag ctctgctgcc agaaggttta gtagagcaat ttgatactgg tatgacactc 720
actgaagcag ttagttccct ggctcagaaa attgagtctt actggggatt atctcgtaat 780
ccaaatgctc cattgggcta taccaaaggc atccctacag caatggctgc tgaaattctg 840
gctgcatttg tagagtctac tgatgttgta gagaacatcg tggatatgtc agaaattgac 900
ccagataaca agaagactat tggctgtgac accattactg aactggattc cttcgacca 960
attaatagct tccctactgc tattgaagaa gctgttttag tgaatcctac agagaagatg 1020
ttctttgggt atgacattcc tcctgtagct aatactcagc ttcgtaacct tgctgttcgt 1080
aatactccag aacagaaggc tgcattgaaa gcagagcagg ctacagaggt ctatgtacac 1140
acccaatgg ttcaattcta tgagacgtta ggtaaaagacc gtattctcga actgatgggt 1200
gctgggtactc tgaataaaga gttacttaat gataaccatg ctaaattctt ggaaggtaag 1260
aaccgttcag tagaggactc ttacaaccaa ctgttctccg tcattgagca ggtaagagca 1320
cagagcgaag acatctctac tgtacctatt cactatgcat acaatatgac ccgtgttggg 1380
cgtatgcaga tgttaggtaa atacaatcct caatcagcca aactggttcg tgaggccatc 1440
ttacctacta aagctacttt ggatttatcg aaccagaaca atgaagactt ctctgcattc 1500
cagttagggtc tggctcaggc attggacatt aaagtccata ctatgactcg tgaggttatg 1560
tctgacgagt tgactaaatt actggaagg taaatctgaaac cagccattga tatgatgggt 1620
gagtttaata ccaactggttc cttaccagaa aacgcagttg atgttctgaa tacagcatta 1680
ggagatagga agtcattcgt agcattgatg gctcttatgg agtattcccg ttacttagta 1740
gcagaggata aatctgcatt tgtaactcca ctgtatgtag aagcagatgg tgttactaat 1800
ggtccaatca atgccatgat gctaattgaca ggcggtctgt ttactcctga ctggattcgt 1860
aatattgcca aaggggggtt gttcattggg tctccaaata agaccatgaa tgagcatcgc 1920
tctactgctg acaataatga tttatatcaa gcatccacta atgctttgat ggaatcgttg 1980
ggtaagttac gtagtaacta tgcctcta atgcctattc agtctcagat agacagtctt 2040
ctttctctga tggatttggt tttaccggat attaatcttg gtgagaatgg tgcttttaga 2100
cttaaacgtg gtattgctaa gaaccactg actattacca tcttcggttc tgggtgctcg 2160
ggatttgtag gtaagctggg tagttctggt actgatgcca tctatgagcg tatgtctgat 2220
gtactgaaag ctctgtctaa agacccaaat atctctgctg ctatggcaat gtttggttaag 2280
caagctgctt cagaagcaca tgctgaagaa cttcttgccc gtttcctgaa agatatggaa 2340
acactgactt ctactgttcc tgttaaactg aaaggtgtac tggaaactaca atccacagg 2400
acaggagcca aaggaaaaat caatcctaag acctatacca ttaagggcga gcaactgaag 2460
gcacttcagg aaaatatgct gcacttcttt gtagaaccac tacgtaatgg tattactcag 2520
actgtagggt aaagtctggt gtactctact gaacaattac agaaagctac tcagattcaa 2580
tctgtagtgc tgggaagata gttcaaacag cgagtacaag agaagctggc agagaaggct 2640
aaagaccca catggaagaa aggtgatttc ctactcaga aagaactgaa tgatattcag 2700
gcttctctga ataacttagc ccctatgatt gagactggtt ctcagacttt ctacattgct 2760
ggttcagaaa atgcagaagt agcaaatcag gtattagcta ctaacctga tgaccgtatg 2820
cgtgtacca ttagtatcta tgctccagca caggccggtg tagcaggtat tccatttatg 2880
actattggta ctgggtgatg catgatgatg caaactcttt ccactatgaa aggtgcacca 2940
aagaataccc tcaaaatctt tgatgggtat aacattgggt tgaatgacat cactgatgcc 3000
agtcgtaaag ctaatgaagc tgtttacact tcttggcagg gtaaccctat taagaatggt 3060

```

```

tatgaatcat atgctaagtt catgaagaat gtagatttca gcaagctgtc ccctgaagca 3120
ttggaagcaa ttggtaaatac tgctctggaa tatgaccaac gtgagaatgc tactgtagat 3180
gatattgcta acgctgcatac tctgattgaa cgtaacttac gtaatattgc actgggtgta 3240
gatattcgtc ataaggtgct ggataaggta aatctgtcca ttgaccagat ggctgctgta 3300
gggtgctcctt atcagaacaa cggtaagatt gacctcagca atatgacccc tgaacaacag 3360
gctgatgaac tgaataaaact tttccgtgaa gagttagaag cccgtaaaca aaaagtcgct 3420
aaggctaggt aa 3432

```

<210> 8

<211> 1143

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic Peptide

<400> 8

```

Met Gly Gly Ser His His His His His His Gly Met Ala Ser Met Thr
  1                      5                      10                      15

Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp
          20                      25                      30

Pro Ser Ser Arg Ser Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly
          35                      40                      45

Ile Asp Ala Val Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala
  50                      55                      60

Glu Gly Ile Lys Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu
  65                      70                      75                      80

Gln Lys Ser Arg Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala
          85                      90                      95

Gln Ala Leu Ser Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu
          100                     105                     110

Thr Ala Asn Pro Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu
          115                     120                     125

Leu Met Glu His Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu
          130                     135                     140

Ala Lys Phe Leu Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly
          145                     150                     155                     160

Thr Glu Ala Asn Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu
          165                     170                     175

Gln Asp Gly Asp Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala
          180                     185                     190

Val Leu Ala Gly Leu Gln Trp Arg Leu Thr Ala Thr Ser Asn Thr Ala
          195                     200                     205

Ile Lys Asp Ala Lys Asp Val Ala Ala Ile Thr Gly Ile Asp Gln Ala
          210                     215                     220

```

Leu Leu Pro Glu Gly Leu Val Glu Gln Phe Asp Thr Gly Met Thr Leu
 225 230 235 240
 Thr Glu Ala Val Ser Ser Leu Ala Gln Lys Ile Glu Ser Tyr Trp Gly
 245 250 255
 Leu Ser Arg Asn Pro Asn Ala Pro Leu Gly Tyr Thr Lys Gly Ile Pro
 260 265 270
 Thr Ala Met Ala Ala Glu Ile Leu Ala Ala Phe Val Glu Ser Thr Asp
 275 280 285
 Val Val Glu Asn Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys
 290 295 300
 Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro
 305 310 315 320
 Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro
 325 330 335
 Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr
 340 345 350
 Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala
 355 360 365
 Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val
 370 375 380
 Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly
 385 390 395 400
 Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser
 405 410 415
 Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe
 420 425 430
 Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val
 435 440 445
 Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met
 450 455 460
 Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile
 465 470 475 480
 Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp
 485 490 495
 Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val
 500 505 510
 His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu
 515 520 525
 Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr
 530 535 540
 Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu

545		550		555		560
Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser						
	565			570		575
Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr						
	580		585		590	
Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu						
	595		600		605	
Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys						
	610		615		620	
Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg						
625		630		635		640
Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu						
	645		650		655	
Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro						
	660		665		670	
Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu						
	675		680		685	
Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly						
	690		695		700	
Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Phe Gly Ser Gly Ala Arg						
705		710		715		720
Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu						
	725		730		735	
Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser						
	740		745		750	
Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala						
	755		760		765	
Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser						
	770		775		780	
Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly						
785		790		795		800
Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly						
	805		810		815	
Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu						
	820		825		830	
Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr						
	835		840		845	
Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile Gln Ser Val Val Leu						
	850		855		860	
Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys Leu Ala Glu Lys Ala						
865		870		875		880

Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu Thr Gln Lys Glu Leu
 885 890 895
 Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala Pro Met Ile Glu Thr
 900 905 910
 Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu Asn Ala Glu Val Ala
 915 920 925
 Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg Met Arg Val Pro Met
 930 935 940
 Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala Gly Ile Pro Phe Met
 945 950 955 960
 Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln Thr Leu Ser Thr Met
 965 970 975
 Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe Asp Gly Met Asn Ile
 980 985 990
 Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys Ala Asn Glu Ala Val
 995 1000 1005
 Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn Val Tyr Glu Ser Tyr
 1010 1015 1020
 Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys Leu Ser Pro Glu Ala
 1025 1030 1035 1040
 Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr Asp Gln Arg Glu Asn
 1045 1050 1055
 Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser Leu Ile Glu Arg Asn
 1060 1065 1070
 Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp
 1075 1080 1085
 Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr
 1090 1095 1100
 Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln
 1105 1110 1115 1120
 Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys
 1125 1130 1135
 Gln Lys Val Ala Lys Ala Arg
 1140

<210> 9

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

 <223> Description of Artificial Sequence: Synthetic
 Primer

<400> 9
tcccagacaa aaggттаага ttтсатасag gattggatgc attacttcat ccaaaagaag 60
cggagcttc 69

<210> 10
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 10
tgggagagaa aaggттаага ttтgatagag gattggatgg attagttgat ggaaaagaag 60
cggagcttc 69

<210> 11
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 11
tccctgtctt ttggttttgt ttтctttctg gtttggtgc ttтctttctt ccaaaagaag 60
cggagcttc 69

<210> 12
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 12
tccacacaa aaccttaaca ttтсатасac cattccatcc attacttcat ccaaaagaag 60
cggagcttc 69

<210> 13
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 13
accagacaa aaggaaaaga aaacaaacag gaaaggaagc aaaacaacaa ccaaaagaag 60
cggagcttc 69

<210> 14
 <211> 10617
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

<400> 14
 atgggggggtt ctcacatcatca tcatcatcat ggtatggcta gcatgactgg tggacagcaa 60
 atgggtcggg atctgtacga cgatgacgat aaggatccga gctcgagatc tatgtcagta 120
 tttgatagac tggctgggtt cgcagacagc gtaaccaatg caaagcaagt tgacgtctct 180
 actgcaaccg ccagaagaa agctgaacaa ggtgtcacta ctctcttctt ttctcttgat 240
 gctgcttattc aaatgcaagc tgcccgtact ggtaatgttg gggctaattgc atttgaacca 300
 gggacagtgc aatcagattt catgaatctg accccaatgc aaatcatgaa taagtatggg 360
 gttgagcaag gcttacaact tatcaatgct cgtgctgatg cagggaacca ggtattcaat 420
 gattcagtta ctacaagaac tcctggggaa gaactggggg atattgtac tggtgttggc 480
 cttgggtttt ttaataaccct tgggggcatt ggtgctcttg gggcaggctt actcaacgat 540
 gatgcagggtg ctgttggttg tcaacaattg agtaagtta atgatgctgt tcatgctacc 600
 caaagccagg cattacaaga taaacgtaag ctctttgctg ctcgtaactt aatgaatgaa 660
 gtagagagtg aacgtcagta tcaaacagat aagaaagaag gcactaatga catagtagct 720
 tccttatcta aatttggacg tgattttgta ggttcaattg agaatgctgc tcaaactgac 780
 tctattatct ctgatgggtt agcagaaggg gtaggttctc tattaggtgc tggctcctgta 840
 ttaaggggtg catctttact gggtaaagca gttgttccag caaatactct tcgtagtgct 900
 gcattggctg gtgctattga tgcagggtact ggtactcagt cactggctcg tattgcctct 960
 actgtaggta gagctgcacc gggtaggtt ggtgttgggt caatggaagc tgggtgggtgca 1020
 taccaacaaa ctgctgatga aattatgaag atgagcttta aagacttaga gaagtctcct 1080
 gtttatcagc aacatattaa agatgggtatg tccctgaac aggcctcgctg tcagactgca 1140
 tctgaaactg gtcttactgc tgctgctatt caattaccta ttgctgctgc aaccggctcct 1200
 ctggatcccc gttttgagat ggctcctttc cgtgctggct ctttaggtgc tgtaggtatg 1260
 aaccttgccc gtgaaacagt ggaagaaggt gttcagggtg ctacaggcca actggctcag 1320
 aatattgcac agcaacaaaa cattgataag aaccaagacc tgcttaaagg tgtcggta 1380
 caggctgggt taggtgctct ttatggcttt ggttctgctg gtgttgta 1440
 ggtgctgctg gtttagcagg tgctgcaact gctcctgtat tgcgtaccac aatggctgg 1500
 gttaaagctg ctggtagtgt agcaggtaag gttgttctc ctattaagaa tacttttagta 1560
 gctcgtgggt aacgggttat gaagcagaat gaagaagcat ctctgttgc tgatgactat 1620
 gttgcacagg cagcacaaga agctatggct caagcaccag aagcagaagt tactattcgt 1680
 gatgctgttg aagcaactga tgctactcca gaacagaaag ttgcagcaca ccagtatgtt 1740
 tctgacttaa tgaatgctac tcgttttaatt cctgaaaatt atcaggaagc accagagcat 1800
 attcgtaaac ctgtagctgg ttctactgac caagtacagg ttattcagaa gttacagac 1860
 ttagttaaca cattagatga atctaactct aagcactga tggaagctgc atcttatatg 1920
 tatgatgctg tttcagagtt tgagcagttc attaaacgtg accctgctgc actggatagc 1980
 attcctaaag attctccggc tattgagtta ctcaaccgtt atacgaatct gacagcta 2040
 attcagaaca caccaaaagt aattgggtgca ctgaatgtta ttaatcgaat gattaatgaa 2100
 tctgctcaga atgggttctt gaatgtgact gaagaatcca gtccacagga aatgcagaac 2160
 gtagcattag ctgctgaagt agcccctgaa aagctcaatc cagagtctgt aaatgttgtt 2220
 cttaaacatg ctgctgatgg tcgtattaaa ctgaataatc gccagattgc tgccctccag 2280
 aatgctgctg caatcctgaa gggggcacgg gaatatgatg cagaagctgc ccgtcttgga 2340
 ttacgtcctc aagacattgt gagtaaacag attaaaacgg atgagagcag aactcaggaa 2400
 ggacaatact ctgcgttgca acatgcgaat aggattcggg ctgcgtataa ctctggta 2460
 ttcgagttgg cctccgctta cctgaacgac tttatgcagt tcgcccagca catgcagaat 2520
 aaggttgag cgttgaatga gcatcttgtt acggggaatg cggataagaa taagtctgtc 2580
 cactaccaag ctcttactgc tgacagagaa tgggttcgta gccgtaccgg attgggggtc 2640
 aatccctatg acactaagtc ggttaaattt gccagcaag ttgctcttga agcgaacacg 2700
 gtagcggata ttgctaattg cctcgcttgc gcttaccgg aactgaaggc cagtcata 2760
 aaagtactc cattggattc acgtcttaac gctcctgctg ctgagggtgt caaggcattc 2820
 cgtcaaggca atcgagacgt tgcttcttct caaccgaaag ctgactccgt gaatcagggt 2880
 aaagaaactc ctgttataaa acaggaacca gttacatcta ctgtacagac taagactcct 2940
 gtagtgtaat ctgttaaaac agaactact actaaagagt ctagccca ggctataaaa 3000

gaacctgtga accagctctga aaaacaggat gttaacctta ctaatgagga caacatcaag 3060
 caacctactg aatctgttta agaaactgaa acttctacaa aagaaagtac agttacagaa 3120
 gaattaaaag aagggtattga tgctgtttac ccttcattgg taggtactgc tgattctaaa 3180
 gcagagggtg ttaagaacta tttcaaattg tcctttacct taccagaaga acagaaatcc 3240
 cgtactgttg gttcagaagc acctctaaaa gatgtagccc aagctctgtc ttctcgtgct 3300
 cgttatgaac tctttactga gaaagaaact gctaaccctg cttttaatgg ggaagtatt 3360
 aagcgataca aagaactcat ggaacatggg gaaggatttg ctgatattct tcgctcccgt 3420
 ctggctaagt tccttaacac taaggatgtt ggtaaactgt ttgctcaagg tacagaagcc 3480
 aaccgttggg taggtggtaa gttacttaac attgttgagc aggatgggga tacctttaag 3540
 tacaacgaac aattgctaca gactgctgta ttagcaggtc ttcaatggag acttactgct 3600
 accagcaata ctgctatcaa agatgcaaaa gatgttgctg ctattactgg tattgaccaa 3660
 gctctgctgc cagaagggtt agtagagcaa tttgatactg gtatgacact cactgaagca 3720
 gttagttccc tggctcagaa aattgagctc tactggggat tatctcgtaa tccaaatgct 3780
 ccattgggct ataccaaagg catccctaca gcaatggctg ctgaaattct ggctgcattt 3840
 gtagagtcta ctgatgttgt agagaacatc gtggatatgt cagaaattga cccagataac 3900
 aagaagacta ttgggtctga caccattact gaactggatt ccttcgaccc aattaatagc 3960
 ttccctactg ctattgaaga agctgtttta gtgaatccta cagagaagat gttctttggt 4020
 gatgacattc ctctgtagc taatactcag cttcgtaacc ctgctgttcg taatactcca 4080
 gaacagaagg ctgcattgaa agcagagcag gctacagagt tctatgtaca caccccaatg 4140
 gttcaattct atgagacgtt aggtaaagac cgtattctcg aactgatggg tgctggtagt 4200
 ctgaataaag agttacttaa tgataacat gctaaatctc tggaaggtaa gaaccgttca 4260
 gtagaggact cttacaacca actgttctcc gtcattgagc aggttaagagc acagagcgaa 4320
 gacatctcta ctgtacctat tcaactatgca tacaatatga cccgtgttgg tcgtatgcag 4380
 atgttaggta aatacaatcc tcaatcagcc aaactgggtc gtgaggccat cttacctact 4440
 aaagctactt tggatttatc gaaccagaac aatgaagact tctctgcatt ccagttaggt 4500
 ctggctcagg cattggacat taaagtccat actatgactc gtgagggtat gtctgacgag 4560
 ttgactaaat tactggaagg taatctgaaa ccagccattg atatgatggt tgagtttaat 4620
 accactgggt ccttaccaga aaacgcagtt gatgttctga atacagcatt aggagatagg 4680
 aagtcattcg tagcattgat ggctcttatg gagtattccc gttacttagt agcagaggat 4740
 aaatctgcac ttgtaacctc actgtatgta gaagcagatg gtgttactaa tgggtccaatc 4800
 aatgccatga tgctaattgac aggcgggtcg tttactctcg actggattcg taatattgcc 4860
 aaagggggct tgttcattgg ttctccaaat aagaccatga atgagcatcg ctctactgct 4920
 gacaataatg atttatatca agcatccact aatgctttga tggaaatcgtt gggtaagtta 4980
 cgtagtaact atgcctctaa tatgcctatt cagtctcaga tagacagtct tctttctctg 5040
 atggatttgt ttttaccgga tattaatctt ggtgagaatg gtgctttaga acttaaacgt 5100
 ggtattgcta agaaccact gactattacc atctatggtt ctggtgctcg tggattgca 5160
 ggtaagctgg ttagttctgt tactgatgcc atctatgagc gtatgtctga tgtactgaaa 5220
 gctcgtgcta aagacccaaa tatctctgct gctatggcaa tgtttggtaa gcaagctgct 5280
 tcagaagcac atgctgaaga acttcttgcc cgtttctga aagatatgga aacactgact 5340
 tctactgttc ctgttaaacg taaagggtga ctggaactac aatccacagg tacaggagcc 5400
 aaagggaaaa tcaatcctaa gacctatacc attaaaggcg agcaactgaa ggcacttcag 5460
 gaaaatatgc tgcacttctt tgtagaacca ctacgtaatg gtattactca gactgtaggt 5520
 gaaagtctgg tgtactctac tgaacaatta cagaaagcta ctcagattca atctgtatgt 5580
 ctggaagata tgttcaaaca gcgagtacaa gagaagctgg aagaactga atgatattca ggcttctctg 5640
 acatggaaga aagggtgattt ccttactcag aaagaactga tctacattgc tgggtcagaa 5700
 aataacttag cccctatgat tgagactggg tctcagactt tctacattgc tgggtcagaa 5760
 aatgcagaag tagcaaatca ggtattagct actaaccttg atgaccgtat gcgtgtacca 5820
 atgagtatct atgctccagc acaggccggg tagcaggta ttccatttat gactattggg 5880
 actggtgatg gcatgatgat gcaaactctt tccactatga aagggtgcacc aaagaatacc 5940
 ctcaaaatct ttgatgggat gaacattggg ttgaatgaca tcaactgatgc cagtcgtaaa 6000
 gctaataaag ctgtttacac ttcttggcag ggtaacccta ttaagaatgt ttatgaatca 6060
 tatgctaagt tcatgaagaa tgtagatttc agcaagctgt cccctgaagc attggaagca 6120
 attggtaaata ctgctctgga atatgaccaa cgtgagaatg ctactgtaga tgatattgct 6180
 aacgctgcat ctctgattga acgtaactta cgtaatatgt cactgggtgt agatattcgt 6240
 cataagggtc tggataagggt aaatctgtcc attgaccaga tggctgctgt aggtgctcct 6300
 tatcagaaca acggtaagat tgacctcagc aatatgaccc ctgaacaaca ggctgatgaa 6360
 ctgaataaac ttttccgtga agagttagaa gcccgtaaac aaaaagtcgc taaggctagg 6420
 gctgaagctca aagaagaaac tgtttctgaa aaagaaccag tgaatccaga ctttggtagt 6480
 gtaggccgtg agcataaggc atctgggtgt cgatccctgt ctgctactgc tattcgtaat 6540
 ctggctaaga ttagtaatct gccatctact caggcagcta ctcttgcgga gattcagaaa 6600
 tcaactggcag ctaaaagacta taagattatc tacggtacac ctactcaggt tgcagagtat 6660

gctcgtcaga	agaatgttac	tgaattgact	tctcaggaaa	tggaagaagc	tcaggcaggt	6720
aatatattatg	gctggactaa	cttcgatgat	aagaccattt	atctgggttag	cccatctatg	6780
gaaaccctca	ttcatgaact	ggttcatgcc	tctaccttcg	aggaagttta	ttccttctat	6840
cagggtaatg	aagtaagccc	tacttctaag	caggctattg	agaaccttga	aggtctgatg	6900
gaacagttcc	gttctctgga	tatttccaaa	gattctccag	aatgagaga	agcatatgct	6960
gatgctattg	caactatcga	aggtcatttg	agtaatggat	ttgttgaccc	agctatctct	7020
aaagctgctg	ctcttaatga	gtttatggct	tgggggttag	ctaaccgtgc	tcttgctgct	7080
aaacagaaga	gaacatcttc	actggttcaa	atggtgaaag	atgtttatca	ggctattaag	7140
aaattgattt	ggggacgtaa	acaagctcct	gcattgggag	aagatatgtt	ctccaatctg	7200
ctgttttaact	ctgcaattct	gatgcgtagc	caacctacaa	ctcaggcagt	agctaaagat	7260
ggcacactgt	tccatagcaa	agcatatggg	aataatgaac	gtctgtctca	gttgaaccag	7320
actttcgata	aactggtaac	tgattacctt	cgtactgacc	cagttacaga	agtagaacgt	7380
cgtggcaatg	tggctaattg	attaatgagt	gctactcgac	tggttcgtga	tgttcagctc	7440
catggcttca	atatgactgc	tcaggaacag	tctgtattcc	agatggttac	tgctgcatta	7500
gcaactgaag	ctgcgattga	cccacatgct	atggctcggt	ctcaggaact	ttatacccat	7560
gtaattgaac	accttacggg	agagcatttc	atggctgacc	ctgatagtac	taaccctgct	7620
gaccgttact	atgctcaaca	gaaatatgac	accatctctg	gtgctaactc	ggttgaagta	7680
gatgccaaag	gtagaaccag	tctgttacct	acattcctgg	gtctggctat	ggttaatgaa	7740
gaactacgtt	caatcattaa	agaaatgcct	gtacctaaag	cagataagaa	attagggaat	7800
gatatagata	ctctgcttac	caatgcaggt	actcaggtaa	tggaaatctc	gaaccgtcgt	7860
atggctgggtg	accagaaagc	tactaatgtt	caggacagta	ttgatgcttt	gtcagaaaca	7920
atcatggctg	ctgctttgaa	acgagagtcc	ttctatgatg	ctgtagcaac	ccctaccggt	7980
aacttcattg	accgtgctaa	tcagtacgta	acggatagca	ttgaacggtt	atctgaaact	8040
gttattgaga	aggcagataa	ggtaattgct	aacccttcta	atatagctgc	taaagggtgt	8100
gctcatctgg	ctaaactgac	tgctgctatt	gcatctgaaa	aacagggtga	aatagtggtc	8160
cagggtgtta	tgactgctat	gaaccagggg	aaagtatggc	aacctttcca	tgacttagtt	8220
aatgacattg	ttggccgtac	taagactaat	gccaatgtct	atgacttaat	caaattgggt	8280
aagagccaga	tttctcaaga	ccgtcagcaa	ttccgtgagc	atttacctac	agtcattgct	8340
ggtaagtctt	ctcgtaaaat	gactgatacc	gaatggctctg	caatgcatac	tggttttaggt	8400
aaaacagatt	tagctgttct	acgtgaaact	atgacatgg	ctgaaattag	agatttactc	8460
tcttcattcca	agaaagtga	agatgaaatc	tctactctgg	aaaaagagat	tcagaaccaa	8520
gcaggtagaa	actggaatct	ggttcagaag	aaatctaagc	aactggctca	atacatgatt	8580
atgggggaag	taggtaataa	cctccttcgt	aatgcccattg	ctattagtcg	tttgtttaggt	8640
gaacgtatta	ctaattggtc	tgtggcagat	gtagctgcta	ttgataagct	cattactttg	8700
tactctctgg	aattgatgaa	taagtctgac	cgtgaccttt	tgtcagaatt	ggctcaatca	8760
gaagtgggaag	gtatggagtt	ctccattgct	tatatgggtg	gtcaacgtac	tgaagagatg	8820
cgtaaagcta	aagggtgata	ccgtactctg	ctgaatcact	ttaaaggcta	tatccctgta	8880
gagaaccagc	aagggtgtgaa	tttgattatt	gctgacgata	aagagtttgc	taagttaaat	8940
agccaatcct	ttactcgtat	tggtaacttat	caggggagca	ctggtttccg	tactggttct	9000
aaaggttatt	acttcagccc	agtagctgcc	cgtgcccctt	actctcaggg	tattcttcag	9060
aacgttcgta	atactgctgg	tgggtgtggat	attgggtactg	gctttacgtt	aggcactatg	9120
gttgctgggc	gtattactga	caaaccaacc	gtagagcgta	ttaccaaagc	tctggctaaa	9180
ggtagcgtg	ggcgtgaacc	actgatgcca	attataaca	gcaaaggcta	ggtagttgct	9240
tatgaacaat	ccgttgacct	taatatgttg	aagcacctaa	accaagacaa	tcactttgct	9300
aagatgggtg	gtgtatggcg	tggctgctcag	gtggaagagg	ctaaagcaca	acgttttaat	9360
gacattctca	ttgagcaatt	acatgctatg	tatgagaaag	acattaaaga	ctccagtgtc	9420
aataaatctc	aatatgtaaa	cctgttaggt	aaaattgatg	accagtgact	ggctgatgct	9480
attaacctga	tgaacattga	gactcgtcat	aaggccgaag	aactcttcgg	taaagatgag	9540
ttatgggttc	gtagggatat	gctgaatgat	gcacttggct	atcgtgctgc	atctattggt	9600
gatgtgtgga	ccggtaaact	tcgttggtca	cctagcacc	ttgatactgt	taagaagatg	9660
ttctcgggtg	cattcggtaa	taaggcatat	catgtagtaa	tgaatgctga	aaataccatt	9720
cagaacttag	tgaaggacgc	taagacagta	attgttggtta	aatctgttgt	agtaccggca	9780
gttaacttcc	ttgctaacat	ctaccagatg	attggacgtg	gtgttcctgt	taaagatatt	9840
gctgtgaaca	ttcctcgtaa	gacgtcagag	attaatcagt	atattaaatc	tcgtttacgt	9900
cagattgatg	cggaagcaga	gctacgtgct	gctgaaggta	accctaactc	ggttcgtaaa	9960
cttaaaactg	agattcaatc	tattactgat	agtcacgtc	gtatgagtat	ctggcctttg	10020
attgaagcag	gtgagttctc	ttctattgct	gatgtggta	ttagtcgtga	tgactgttta	10080
gtagctgaag	gtaagattca	tgagtacatg	gaaaaacttg	ctaataaact	tccagaaaaa	10140
gtacgtaatg	ctggccgtta	cgctcttatt	gctaaggaca	ctgctctgtt	ccagggtatc	10200
cagaaaacag	tagagtattc	agactttatt	gctaaagcca	tcattctatga	tgatttagtg	10260
aaacgtaaga	aaaaatcttc	ttctgaagca	ttaggtcagg	taactgaaga	gtttattaac	10320

tatgacagat tgccctggtcg tttccgtggc tatatggaaa gtatgggtct gatgtggttc 10380
 tacaacttta aaattcggtc cattaaagtt gctatgagca tgattagaaa caaccagta 10440
 cattctctga ttgctacagt agtacctgct cctaccatgt ttggtaacgt aggtctacca 10500
 attcaggaca acatgctaac catgctggct gaaggaagac tggattactc attaggttc 10560
 ggacaaggat taagagcacc taccctcaat ccttggttca accttactca ctaataa 10617

<210> 15

<211> 3537

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic Peptide

<400> 15

Met Gly Gly Ser His His His His His His Gly Met Ala Ser Met Thr
 1 5 10 15

Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp
 20 25 30

Pro Ser Ser Arg Ser Met Ser Val Phe Asp Arg Leu Ala Gly Phe Ala
 35 40 45

Asp Ser Val Thr Asn Ala Lys Gln Val Asp Val Ser Thr Ala Thr Ala
 50 55 60

Gln Lys Lys Ala Glu Gln Gly Val Thr Thr Pro Leu Val Ser Pro Asp
 65 70 75 80

Ala Ala Tyr Gln Met Gln Ala Ala Arg Thr Gly Asn Val Gly Ala Asn
 85 90 95

Ala Phe Glu Pro Gly Thr Val Gln Ser Asp Phe Met Asn Leu Thr Pro
 100 105 110

Met Gln Ile Met Asn Lys Tyr Gly Val Glu Gln Gly Leu Gln Leu Ile
 115 120 125

Asn Ala Arg Ala Asp Ala Gly Asn Gln Val Phe Asn Asp Ser Val Thr
 130 135 140

Thr Arg Thr Pro Gly Glu Glu Leu Gly Asp Ile Ala Thr Gly Val Gly
 145 150 155 160

Leu Gly Phe Val Asn Thr Leu Gly Gly Ile Gly Ala Leu Gly Ala Gly
 165 170 175

Leu Leu Asn Asp Asp Ala Gly Ala Val Val Ala Gln Gln Leu Ser Lys
 180 185 190

Phe Asn Asp Ala Val His Ala Thr Gln Ser Gln Ala Leu Gln Asp Lys
 195 200 205

Arg Lys Leu Phe Ala Ala Arg Asn Leu Met Asn Glu Val Glu Ser Glu
 210 215 220

Arg Gln Tyr Gln Thr Asp Lys Lys Glu Gly Thr Asn Asp Ile Val Ala
 225 230 235 240

Ser Leu Ser Lys Phe Gly Arg Asp Phe Val Gly Ser Ile Glu Asn Ala
 245 250 255
 Ala Gln Thr Asp Ser Ile Ile Ser Asp Gly Leu Ala Glu Gly Val Gly
 260 265 270
 Ser Leu Leu Gly Ala Gly Pro Val Leu Arg Gly Ala Ser Leu Leu Gly
 275 280 285
 Lys Ala Val Val Pro Ala Asn Thr Leu Arg Ser Ala Ala Leu Ala Gly
 290 295 300
 Ala Ile Asp Ala Gly Thr Gly Thr Gln Ser Leu Ala Arg Ile Ala Ser
 305 310 315 320
 Thr Val Gly Arg Ala Ala Pro Gly Met Val Gly Val Gly Ala Met Glu
 325 330 335
 Ala Gly Gly Ala Tyr Gln Gln Thr Ala Asp Glu Ile Met Lys Met Ser
 340 345 350
 Leu Lys Asp Leu Glu Lys Ser Pro Val Tyr Gln Gln His Ile Lys Asp
 355 360 365
 Gly Met Ser Pro Glu Gln Ala Arg Arg Gln Thr Ala Ser Glu Thr Gly
 370 375 380
 Leu Thr Ala Ala Ala Ile Gln Leu Pro Ile Ala Ala Ala Thr Gly Pro
 385 390 395 400
 Leu Val Ser Arg Phe Glu Met Ala Pro Phe Arg Ala Gly Ser Leu Gly
 405 410 415
 Ala Val Gly Met Asn Leu Ala Arg Glu Thr Val Glu Glu Gly Val Gln
 420 425 430
 Gly Ala Thr Gly Gln Leu Ala Gln Asn Ile Ala Gln Gln Gln Asn Ile
 435 440 445
 Asp Lys Asn Gln Asp Leu Leu Lys Gly Val Gly Thr Gln Ala Gly Leu
 450 455 460
 Gly Ala Leu Tyr Gly Phe Gly Ser Ala Gly Val Val Gln Ala Pro Ala
 465 470 475 480
 Gly Ala Ala Arg Leu Ala Gly Ala Ala Thr Ala Pro Val Leu Arg Thr
 485 490 495
 Thr Met Ala Gly Val Lys Ala Ala Gly Ser Val Ala Gly Lys Val Val
 500 505 510
 Ser Pro Ile Lys Asn Thr Leu Val Ala Arg Gly Glu Arg Val Met Lys
 515 520 525
 Gln Asn Glu Glu Ala Ser Pro Val Ala Asp Asp Tyr Val Ala Gln Ala
 530 535 540
 Ala Gln Glu Ala Met Ala Gln Ala Pro Glu Ala Glu Val Thr Ile Arg
 545 550 555 560

Asp Ala Val Glu Ala Thr Asp Ala Thr Pro Glu Gln Lys Val Ala Ala
 565 570 575
 His Gln Tyr Val Ser Asp Leu Met Asn Ala Thr Arg Phe Asn Pro Glu
 580 585 590
 Asn Tyr Gln Glu Ala Pro Glu His Ile Arg Asn Ala Val Ala Gly Ser
 595 600 605
 Thr Asp Gln Val Gln Val Ile Gln Lys Leu Ala Asp Leu Val Asn Thr
 610 615 620
 Leu Asp Glu Ser Asn Pro Gln Ala Leu Met Glu Ala Ala Ser Tyr Met
 625 630 635 640
 Tyr Asp Ala Val Ser Glu Phe Glu Gln Phe Ile Asn Arg Asp Pro Ala
 645 650 655
 Ala Leu Asp Ser Ile Pro Lys Asp Ser Pro Ala Ile Glu Leu Leu Asn
 660 665 670
 Arg Tyr Thr Asn Leu Thr Ala Asn Ile Gln Asn Thr Pro Lys Val Ile
 675 680 685
 Gly Ala Leu Asn Val Ile Asn Arg Met Ile Asn Glu Ser Ala Gln Asn
 690 695 700
 Gly Ser Leu Asn Val Thr Glu Glu Ser Ser Pro Gln Glu Met Gln Asn
 705 710 715 720
 Val Ala Leu Ala Ala Glu Val Ala Pro Glu Lys Leu Asn Pro Glu Ser
 725 730 735
 Val Asn Val Val Leu Lys His Ala Ala Asp Gly Arg Ile Lys Leu Asn
 740 745 750
 Asn Arg Gln Ile Ala Ala Leu Gln Asn Ala Ala Ala Ile Leu Lys Gly
 755 760 765
 Ala Arg Glu Tyr Asp Ala Glu Ala Ala Arg Leu Gly Leu Arg Pro Gln
 770 775 780
 Asp Ile Val Ser Lys Gln Ile Lys Thr Asp Glu Ser Arg Thr Gln Glu
 785 790 795 800
 Gly Gln Tyr Ser Ala Leu Gln His Ala Asn Arg Ile Arg Ser Ala Tyr
 805 810 815
 Asn Ser Gly Asn Phe Glu Leu Ala Ser Ala Tyr Leu Asn Asp Phe Met
 820 825 830
 Gln Phe Ala Gln His Met Gln Asn Lys Val Gly Ala Leu Asn Glu His
 835 840 845
 Leu Val Thr Gly Asn Ala Asp Lys Asn Lys Ser Val His Tyr Gln Ala
 850 855 860
 Leu Thr Ala Asp Arg Glu Trp Val Arg Ser Arg Thr Gly Leu Gly Val
 865 870 875 880
 Asn Pro Tyr Asp Thr Lys Ser Val Lys Phe Ala Gln Gln Val Ala Leu

37

Gly Ile Asp Gln Ala Leu Leu Pro Glu Gly Leu Val Glu Gln Phe Asp
 1220 1225 1230
 Thr Gly Met Thr Leu Thr Glu Ala Val Ser Ser Leu Ala Gln Lys Ile
 1235 1240 1245
 Glu Ser Tyr Trp Gly Leu Ser Arg Asn Pro Asn Ala Pro Leu Gly Tyr
 1250 1255 1260
 Thr Lys Gly Ile Pro Thr Ala Met Ala Ala Glu Ile Leu Ala Ala Phe
 1265 1270 1275 1280
 Val Glu Ser Thr Asp Val Val Glu Asn Ile Val Asp Met Ser Glu Ile
 1285 1290 1295
 Asp Pro Asp Asn Lys Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu
 1300 1305 1310
 Asp Ser Phe Asp Pro Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala
 1315 1320 1325
 Val Leu Val Asn Pro Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro
 1330 1335 1340
 Pro Val Ala Asn Thr Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro
 1345 1350 1355 1360
 Glu Gln Lys Ala Ala Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val
 1365 1370 1375
 His Thr Pro Met Val Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile
 1380 1385 1390
 Leu Glu Leu Met Gly Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp
 1395 1400 1405
 Asn His Ala Lys Ser Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser
 1410 1415 1420
 Tyr Asn Gln Leu Phe Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu
 1425 1430 1435 1440
 Asp Ile Ser Thr Val Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val
 1445 1450 1455
 Gly Arg Met Gln Met Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu
 1460 1465 1470
 Val Arg Glu Ala Ile Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn
 1475 1480 1485
 Gln Asn Asn Glu Asp Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala
 1490 1495 1500
 Leu Asp Ile Lys Val His Thr Met Thr Arg Glu Val Met Ser Asp Glu
 1505 1510 1515 1520
 Leu Thr Lys Leu Leu Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met
 1525 1530 1535

Val Glu Phe Asn Thr Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val
 1540 1545 1550
 Leu Asn Thr Ala Leu Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala
 1555 1560 1565
 Leu Met Glu Tyr Ser Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe
 1570 1575 1580
 Val Thr Pro Leu Tyr Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile
 1585 1590 1595 1600
 Asn Ala Met Met Leu Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile
 1605 1610 1615
 Arg Asn Ile Ala Lys Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr
 1620 1625 1630
 Met Asn Glu His Arg Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala
 1635 1640 1645
 Ser Thr Asn Ala Leu Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr
 1650 1655 1660
 Ala Ser Asn Met Pro Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu
 1665 1670 1675 1680
 Met Asp Leu Phe Leu Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu
 1685 1690 1695
 Glu Leu Lys Arg Gly Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Tyr
 1700 1705 1710
 Gly Ser Gly Ala Arg Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr
 1715 1720 1725
 Asp Ala Ile Tyr Glu Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys
 1730 1735 1740
 Asp Pro Asn Ile Ser Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala
 1745 1750 1755 1760
 Ser Glu Ala His Ala Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met
 1765 1770 1775
 Glu Thr Leu Thr Ser Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu
 1780 1785 1790
 Leu Gln Ser Thr Gly Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr
 1795 1800 1805
 Tyr Thr Ile Lys Gly Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu
 1810 1815 1820
 His Phe Phe Val Glu Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly
 1825 1830 1835 1840
 Glu Ser Leu Val Tyr Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile
 1845 1850 1855
 Gln Ser Val Val Leu Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys

1860	1865	1870
Leu Ala Glu Lys Ala Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu 1875 1880 1885		
Thr Gln Lys Glu Leu Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala 1890 1895 1900		
Pro Met Ile Glu Thr Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu 1905 1910 1915 1920		
Asn Ala Glu Val Ala Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg 1925 1930 1935		
Met Arg Val Pro Met Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala 1940 1945 1950		
Gly Ile Pro Phe Met Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln 1955 1960 1965		
Thr Leu Ser Thr Met Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe 1970 1975 1980		
Asp Gly Met Asn Ile Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys 1985 1990 1995 2000		
Ala Asn Glu Ala Val Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn 2005 2010 2015		
Val Tyr Glu Ser Tyr Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys 2020 2025 2030		
Leu Ser Pro Glu Ala Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr 2035 2040 2045		
Asp Gln Arg Glu Asn Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser 2050 2055 2060		
Leu Ile Glu Arg Asn Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg 2065 2070 2075 2080		
His Lys Val Leu Asp Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala 2085 2090 2095		
Val Gly Ala Pro Tyr Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met 2100 2105 2110		
Thr Pro Glu Gln Gln Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu 2115 2120 2125		
Leu Glu Ala Arg Lys Gln Lys Val Ala Lys Ala Arg Ala Glu Val Lys 2130 2135 2140		
Glu Glu Thr Val Ser Glu Lys Glu Pro Val Asn Pro Asp Phe Gly Met 2145 2150 2155 2160		
Val Gly Arg Glu His Lys Ala Ser Gly Val Arg Ile Leu Ser Ala Thr 2165 2170 2175		
Ala Ile Arg Asn Leu Ala Lys Ile Ser Asn Leu Pro Ser Thr Gln Ala 2180 2185 2190		

Ala Thr Leu Ala Glu Ile Gln Lys Ser Leu Ala Ala Lys Asp Tyr Lys
 2195 2200 2205
 Ile Ile Tyr Gly Thr Pro Thr Gln Val Ala Glu Tyr Ala Arg Gln Lys
 2210 2215 2220
 Asn Val Thr Glu Leu Thr Ser Gln Glu Met Glu Glu Ala Gln Ala Gly
 2225 2230 2235 2240
 Asn Ile Tyr Gly Trp Thr Asn Phe Asp Asp Lys Thr Ile Tyr Leu Val
 2245 2250 2255
 Ser Pro Ser Met Glu Thr Leu Ile His Glu Leu Val His Ala Ser Thr
 2260 2265 2270
 Phe Glu Glu Val Tyr Ser Phe Tyr Gln Gly Asn Glu Val Ser Pro Thr
 2275 2280 2285
 Ser Lys Gln Ala Ile Glu Asn Leu Glu Gly Leu Met Glu Gln Phe Arg
 2290 2295 2300
 Ser Leu Asp Ile Ser Lys Asp Ser Pro Glu Met Arg Glu Ala Tyr Ala
 2305 2310 2315 2320
 Asp Ala Ile Ala Thr Ile Glu Gly His Leu Ser Asn Gly Phe Val Asp
 2325 2330 2335
 Pro Ala Ile Ser Lys Ala Ala Ala Leu Asn Glu Phe Met Ala Trp Gly
 2340 2345 2350
 Leu Ala Asn Arg Ala Leu Ala Ala Lys Gln Lys Arg Thr Ser Ser Leu
 2355 2360 2365
 Val Gln Met Val Lys Asp Val Tyr Gln Ala Ile Lys Lys Leu Ile Trp
 2370 2375 2380
 Gly Arg Lys Gln Ala Pro Ala Leu Gly Glu Asp Met Phe Ser Asn Leu
 2385 2390 2395 2400
 Leu Phe Asn Ser Ala Ile Leu Met Arg Ser Gln Pro Thr Thr Gln Ala
 2405 2410 2415
 Val Ala Lys Asp Gly Thr Leu Phe His Ser Lys Ala Tyr Gly Asn Asn
 2420 2425 2430
 Glu Arg Leu Ser Gln Leu Asn Gln Thr Phe Asp Lys Leu Val Thr Asp
 2435 2440 2445
 Tyr Leu Arg Thr Asp Pro Val Thr Glu Val Glu Arg Arg Gly Asn Val
 2450 2455 2460
 Ala Asn Ala Leu Met Ser Ala Thr Arg Leu Val Arg Asp Val Gln Ser
 2465 2470 2475 2480
 His Gly Phe Asn Met Thr Ala Gln Glu Gln Ser Val Phe Gln Met Val
 2485 2490 2495
 Thr Ala Ala Leu Ala Thr Glu Ala Ala Ile Asp Pro His Ala Met Ala
 2500 2505 2510

Arg Ala Gln Glu Leu Tyr Thr His Val Met Lys His Leu Thr Val Glu
 2515 2520 2525
 His Phe Met Ala Asp Pro Asp Ser Thr Asn Pro Ala Asp Arg Tyr Tyr
 2530 2535 2540
 Ala Gln Gln Lys Tyr Asp Thr Ile Ser Gly Ala Asn Leu Val Glu Val
 2545 2550 2555 2560
 Asp Ala Lys Gly Arg Thr Ser Leu Leu Pro Thr Phe Leu Gly Leu Ala
 2565 2570 2575
 Met Val Asn Glu Glu Leu Arg Ser Ile Ile Lys Glu Met Pro Val Pro
 2580 2585 2590
 Lys Ala Asp Lys Lys Leu Gly Asn Asp Ile Asp Thr Leu Leu Thr Asn
 2595 2600 2605
 Ala Gly Thr Gln Val Met Glu Ser Leu Asn Arg Arg Met Ala Gly Asp
 2610 2615 2620
 Gln Lys Ala Thr Asn Val Gln Asp Ser Ile Asp Ala Leu Ser Glu Thr
 2625 2630 2635 2640
 Ile Met Ala Ala Ala Leu Lys Arg Glu Ser Phe Tyr Asp Ala Val Ala
 2645 2650 2655
 Thr Pro Thr Gly Asn Phe Ile Asp Arg Ala Asn Gln Tyr Val Thr Asp
 2660 2665 2670
 Ser Ile Glu Arg Leu Ser Glu Thr Val Ile Glu Lys Ala Asp Lys Val
 2675 2680 2685
 Ile Ala Asn Pro Ser Asn Ile Ala Ala Lys Gly Val Ala His Leu Ala
 2690 2695 2700
 Lys Leu Thr Ala Ala Ile Ala Ser Glu Lys Gln Gly Glu Ile Val Ala
 2705 2710 2715 2720
 Gln Gly Val Met Thr Ala Met Asn Gln Gly Lys Val Trp Gln Pro Phe
 2725 2730 2735
 His Asp Leu Val Asn Asp Ile Val Gly Arg Thr Lys Thr Asn Ala Asn
 2740 2745 2750
 Val Tyr Asp Leu Ile Lys Leu Val Lys Ser Gln Ile Ser Gln Asp Arg
 2755 2760 2765
 Gln Gln Phe Arg Glu His Leu Pro Thr Val Ile Ala Gly Lys Phe Ser
 2770 2775 2780
 Arg Lys Leu Thr Asp Thr Glu Trp Ser Ala Met His Thr Gly Leu Gly
 2785 2790 2795 2800
 Lys Thr Asp Leu Ala Val Leu Arg Glu Thr Met Ser Met Ala Glu Ile
 2805 2810 2815
 Arg Asp Leu Leu Ser Ser Ser Lys Lys Val Lys Asp Glu Ile Ser Thr
 2820 2825 2830
 Leu Glu Lys Glu Ile Gln Asn Gln Ala Gly Arg Asn Trp Asn Leu Val

2835	2840	2845
Gln Lys Lys Ser Lys Gln Leu Ala Gln Tyr Met Ile Met Gly Glu Val		
2850	2855	2860
Gly Asn Asn Leu Leu Arg Asn Ala His Ala Ile Ser Arg Leu Leu Gly		
2865	2870	2875 2880
Glu Arg Ile Thr Asn Gly Pro Val Ala Asp Val Ala Ala Ile Asp Lys		
	2885	2890 2895
Leu Ile Thr Leu Tyr Ser Leu Glu Leu Met Asn Lys Ser Asp Arg Asp		
	2900	2905 2910
Leu Leu Ser Glu Leu Ala Gln Ser Glu Val Glu Gly Met Glu Phe Ser		
	2915	2920 2925
Ile Ala Tyr Met Val Gly Gln Arg Thr Glu Glu Met Arg Lys Ala Lys		
	2930	2935 2940
Gly Asp Asn Arg Thr Leu Leu Asn His Phe Lys Gly Tyr Ile Pro Val		
2945	2950	2955 2960
Glu Asn Gln Gln Gly Val Asn Leu Ile Ile Ala Asp Asp Lys Glu Phe		
	2965	2970 2975
Ala Lys Leu Asn Ser Gln Ser Phe Thr Arg Ile Gly Thr Tyr Gln Gly		
	2980	2985 2990
Ser Thr Gly Phe Arg Thr Gly Ser Lys Gly Tyr Tyr Phe Ser Pro Val		
	2995	3000 3005
Ala Ala Arg Ala Pro Tyr Ser Gln Gly Ile Leu Gln Asn Val Arg Asn		
	3010	3015 3020
Thr Ala Gly Gly Val Asp Ile Gly Thr Gly Phe Thr Leu Gly Thr Met		
3025	3030	3035 3040
Val Ala Gly Arg Ile Thr Asp Lys Pro Thr Val Glu Arg Ile Thr Lys		
	3045	3050 3055
Ala Leu Ala Lys Gly Glu Arg Gly Arg Glu Pro Leu Met Pro Ile Tyr		
	3060	3065 3070
Asn Ser Lys Gly Gln Val Val Ala Tyr Glu Gln Ser Val Asp Pro Asn		
	3075	3080 3085
Met Leu Lys His Leu Asn Gln Asp Asn His Phe Ala Lys Met Val Gly		
	3090	3095 3100
Val Trp Arg Gly Arg Gln Val Glu Glu Ala Lys Ala Gln Arg Phe Asn		
3105	3110	3115 3120
Asp Ile Leu Ile Glu Gln Leu His Ala Met Tyr Glu Lys Asp Ile Lys		
	3125	3130 3135
Asp Ser Ser Ala Asn Lys Ser Gln Tyr Val Asn Leu Leu Gly Lys Ile		
	3140	3145 3150
Asp Asp Pro Val Leu Ala Asp Ala Ile Asn Leu Met Asn Ile Glu Thr		
	3155	3160 3165

Arg His Lys Ala Glu Glu Leu Phe Gly Lys Asp Glu Leu Trp Val Arg
 3170 3175 3180
 Arg Asp Met Leu Asn Asp Ala Leu Gly Tyr Arg Ala Ala Ser Ile Gly
 3185 3190 3195 3200
 Asp Val Trp Thr Gly Asn Ser Arg Trp Ser Pro Ser Thr Leu Asp Thr
 3205 3210 3215
 Val Lys Lys Met Phe Leu Gly Ala Phe Gly Asn Lys Ala Tyr His Val
 3220 3225 3230
 Val Met Asn Ala Glu Asn Thr Ile Gln Asn Leu Val Lys Asp Ala Lys
 3235 3240 3245
 Thr Val Ile Val Val Lys Ser Val Val Val Pro Ala Val Asn Phe Leu
 3250 3255 3260
 Ala Asn Ile Tyr Gln Met Ile Gly Arg Gly Val Pro Val Lys Asp Ile
 3265 3270 3275 3280
 Ala Val Asn Ile Pro Arg Lys Thr Ser Glu Ile Asn Gln Tyr Ile Lys
 3285 3290 3295
 Ser Arg Leu Arg Gln Ile Asp Ala Glu Ala Glu Leu Arg Ala Ala Glu
 3300 3305 3310
 Gly Asn Pro Asn Leu Val Arg Lys Leu Lys Thr Glu Ile Gln Ser Ile
 3315 3320 3325
 Thr Asp Ser His Arg Arg Met Ser Ile Trp Pro Leu Ile Glu Ala Gly
 3330 3335 3340
 Glu Phe Ser Ser Ile Ala Asp Ala Gly Ile Ser Arg Asp Asp Leu Leu
 3345 3350 3355 3360
 Val Ala Glu Gly Lys Ile His Glu Tyr Met Glu Lys Leu Ala Asn Lys
 3365 3370 3375
 Leu Pro Glu Lys Val Arg Asn Ala Gly Arg Tyr Ala Leu Ile Ala Lys
 3380 3385 3390
 Asp Thr Ala Leu Phe Gln Gly Ile Gln Lys Thr Val Glu Tyr Ser Asp
 3395 3400 3405
 Phe Ile Ala Lys Ala Ile Ile Tyr Asp Asp Leu Val Lys Arg Lys Lys
 3410 3415 3420
 Lys Ser Ser Ser Glu Ala Leu Gly Gln Val Thr Glu Glu Phe Ile Asn
 3425 3430 3435 3440
 Tyr Asp Arg Leu Pro Gly Arg Phe Arg Gly Tyr Met Glu Ser Met Gly
 3445 3450 3455
 Leu Met Trp Phe Tyr Asn Phe Lys Ile Arg Ser Ile Lys Val Ala Met
 3460 3465 3470
 Ser Met Ile Arg Asn Asn Pro Val His Ser Leu Ile Ala Thr Val Val
 3475 3480 3485

Pro Ala Pro Thr Met Phe Gly Asn Val Gly Leu Pro Ile Gln Asp Asn
 3490 3495 3500

Met Leu Thr Met Leu Ala Glu Gly Arg Leu Asp Tyr Ser Leu Gly Phe
 3505 3510 3515 3520

Gly Gln Gly Leu Arg Ala Pro Thr Leu Asn Pro Trp Phe Asn Leu Thr
 3525 3530 3535

His

<210> 16

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 16

ggcattactt catccaaaag aagcggagct tc

32

<210> 17

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 17

ggccatccat tacttcatcc aaaagaagcg gagcttc

37

<210> 18

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 18

ggatccaaaa gaagcggagc ttc

23

<210> 19

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 Primer

<400> 19
ggcattactt catccaaaag aagctgagct tc 32

<210> 20
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 20
ggcattactt catccaaaag aagcggagc 29

<210> 21
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 21
ggaggctcct cggagtctcc tttt 24

<210> 22
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 22
ggactacctt cgggtagtcc ttttt 25

<210> 23
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 23
agaagggggc tactaagccc ttttcttatt ttt 33

<210> 24
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 24
aagctgctcc gcagctttt 19

<210> 25
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 25
aaggctatcc ctacgggggt agcctttatt ttttt 35

<210> 26
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 26
gccctccttg tgagggtttt tt 22

<210> 27
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 27
caacgaagcg ttgaatacct 20

<210> 28
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 28
ttcttcgagg cgaagaaaac ct 22

<210> 29

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 29

cgacgaggcg tcgaaaacca