



(86) Date de dépôt PCT/PCT Filing Date: 2011/05/06
(87) Date publication PCT/PCT Publication Date: 2011/11/10
(45) Date de délivrance/Issue Date: 2020/06/02
(85) Entrée phase nationale/National Entry: 2012/10/22
(86) N° demande PCT/PCT Application No.: US 2011/035500
(87) N° publication PCT/PCT Publication No.: 2011/140429
(30) Priorité/Priority: 2010/05/06 (US61/331,909)

(51) Cl.Int./Int.Cl. *A61K 38/45* (2006.01),
A61P 1/16 (2006.01)
(72) Inventeurs/Inventors:
AUERBACH, BRUCE J., US;
HOMAN, REYNOLD, US;
KRAUSE, BRIAN, US
(73) Propriétaire/Owner:
ALPHACORE PHARMA LLC, US
(74) Agent: MACRAE & CO.

(54) Titre : ADMINISTRATION D'ESTER DE CHOLESTERYLE A DES TISSUS STEROIDOGENES
(54) Title: DELIVERY OF CHOLESTERYL ESTER TO STEROIDOGENIC TISSUES

(57) Abrégé/Abstract:

Disclosed are compositions and methods for treating conditions characterized by low HDL-CE which can lead to decreased delivery of cholesteryl ester to steroidogenic tissues, reducing the organ's ability to produce steroids especially during periods of demand, stress and or systemic inflammatory response syndrome.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 November 2011 (10.11.2011)

PCT

(10) International Publication Number
WO 2011/140429 A1(51) International Patent Classification:
A61K 38/45 (2006.01) *A61P 1/16* (2006.01)(21) International Application Number:
PCT/US2011/035500(22) International Filing Date:
6 May 2011 (06.05.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/331,909 6 May 2010 (06.05.2010) US(71) Applicant (for all designated States except US): **AL-PHACORE PHARMA LLC** [US/US]; 333 Parkland Plaza, Suite 5, Ann Arbor, Michigan 48103 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRUCE J. AUER-BACH, Bruce** [US/US]; 2425 Meadowridge Court, Ann Arbor, Michigan 48105 (US). **REYNOLD HOMAN, Reynold** [US/US]; 2607 Tessmer Road, Ann Arbor, Michigan 48103 (US). **BRIAN KRAUSE, Brian** [US/US]; 1984 Encore Ct., Ann Arbor, Michigan 48103 (US).(74) Agent: **BOTT, Cynthia, M.**; HONIGMAN MILLER SCHWARTZ AND COHN LLP, 350 East Michigan Avenue, Suite 300, Kalamazoo, Michigan 49007-3800 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: DELIVERY OF CHOLESTERYL ESTER TO STEROIDOGENIC TISSUES

(57) Abstract: Disclosed are compositions and methods for treating conditions characterized by low HDL-CE which can lead to decreased delivery of cholesteryl ester to steroidogenic tissues, reducing the organ's ability to produce steroids especially during periods of demand, stress and or systemic inflammatory response syndrome.

WO 2011/140429 A1

DELIVERY OF CHOLESTERYL ESTER TO STEROIDOGENIC TISSUES

FIELD OF INVENTION

[0001] This invention relates generally to the field of medicine, and in particular, the treatment of diseases characterized by decreased stores of cholesteryl ester in steroidogenic tissues, reducing the tissue's ability to produce steroids especially during periods of demand, stress and or systemic inflammation.

BACKGROUND

[0002] Normally, excess cholesterol is removed from tissues, such as arteries, and delivered to the liver for excretion in bile by a process known as reverse cholesterol transport (RCT). In the first step of RCT, cholesterol passes from tissue cells to high-density lipoproteins (HDL) in the circulation. In the second step, the enzyme lecithin: cholesterol acyltransferase (LCAT) enhances the cholesterol-carrying capacity of HDL by catalyzing the transesterification of a fatty acid from lecithin in HDL to cholesterol to form cholesteryl ester (CE). The cholesteryl ester product accumulates in the HDL interior until it is removed at HDL-receptors in the liver. Much of the CE entering the liver is converted to cholesterol and bile acids that are excreted in the bile. The same HDL receptor-mediated process of CE delivery occurs in steroidogenic tissues to maintain a supply of cholesterol for the production of steroids. The HDL receptor for this process is currently thought to be the scavenger receptor B1 (SR-B1).

[0003] Cholesteryl ester transfer protein (CETP) is a plasma protein responsible for the net transfer of CE from HDL to the beta-lipoproteins (VLDL and LDL). Humans deficient in CETP have greatly elevated levels of large CE-enriched HDL. Inhibitors of CETP have been shown to block the transfer of CE from HDL to LDL, thus rapidly increasing HDL-C (where C represents total cholesterol and is comprised of both cholesterol and CE) and reducing LDL-C. The resultant HDL is characteristically enriched in CE, similar to the HDL isolated from CETP deficient plasma.

[0004] The steroidogenic tissues, including adrenals, gonads (testes and ovaries) and thymus, rely on intracellular stores of cholesteryl esters as a source of cholesterol for the synthesis of steroids such as cortisol, estrogens and androgens. When needed, the ester bond is cleaved by the enzyme cholesterol esterase, liberating cholesterol for use in steroid production. In times of stress, the CE stores can be depleted quickly, and without replenishment of CE, production of steroids is greatly impaired. In vitro experiments have

demonstrated that HDL isolated from CETP deficient patients can deliver more CE through SR-B1 than HDL isolated from normal subjects.

[0005] In mice, low HDL-CE results in reduced CE content in adrenals, testes and ovaries. This decreased CE content appears to impact reproduction and the ability of the adrenals to produce cortisol during periods of stress. In humans, hypoalphalipoproteinemia (low HDL-C) is a hallmark of genetic disorders such as LCAT deficiency and Tangier's disease. Hypoalphalipoproteinemia also occurs in a variety of non-genetic disorders including infection, inflammation, auto-immune diseases, arthritic diseases, sepsis, trauma, burns, liver disease (e.g. hepatitis, fibrosis, cirrhosis, bile duct hyperplasia, bile duct atresia, organ transplant, heavy metal poisoning) and kidney disease.

[0006] Although not a primary pathology in these diseases, the decreased HDL-CE reduces the delivery of cholesterol to steroidogenic tissues, thus, decreasing the body's ability to produce critical glucocorticoids, hormones and other steroids. This reduced capacity for steroidal production can exacerbate the underlying conditions. As a compounding factor, the worsening of the underlying condition can further decrease the level of HDL-CE. In some cases this cycle can lead to a life threatening condition, even death. Following trauma or severe infection patients may have reduced HDL-C levels, or may be at risk for HDL-C levels to fall. It would be useful to normalize or increase the levels of HDL-C in order to normalize or increase steroidogenesis in these patients.

[0007] Artificial forms of HDL consisting of phospholipid and apolipoprotein A-1, or of phospholipids and apolipoprotein A-1 mimetics, are being investigated for treatment of patients having cardiovascular disorders such as atherosclerosis. Such treatments are predicated upon the ability of these artificial HDLs to remove excess cholesterol from tissues such as arteries. However, because these versions of artificial HDL lack cholesterol or cholesteryl esters, they are not immediately able to deliver CE to the adrenals as is needed to normalize or increase steroidogenesis and in fact would function to remove cholesterol from tissues. In some instances, patients at risk for adrenal insufficiency have decreased levels of LCAT and therefore have a reduced capacity to make HDL-CE (the form of cholesterol used to replenish adrenals) resulting in low levels of HDL-CE. Because these artificial forms of HDL do not correct for the deficiency of LCAT or HDL-CE they do not provide an effective treatment for critical care patients at risk for adrenal insufficiency.

[0008] Thus, there is a need for treatments to increase the availability of CE to steroidogenic tissues in patients having disorders associated with decreased function of steroidogenic tissues such as the adrenals, gonads and thymus.

SUMMARY

[0009] The compositions and methods disclosed in the present disclosure modulate the amount of steroids produced by steroidogenic tissues by normalizing or increasing HDL-CE concentrations in plasma thereby maintaining or increasing plasma levels of steroids by administering an effective dose of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a cholesterol delivery particle (CDP), or any combination thereof. The present disclosure describes methods of treating conditions characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP. The present disclosure describes methods of treating conditions characterized by adrenal insufficiency. Also disclosed are methods of treating conditions characterized by low testosterone production comprising administering to a patient in need thereof, an effective dose of a CETP inhibitor, LCAT, a compound that increases plasma LCAT activity, a CDP, or any combination thereof. Additionally described are methods of treating conditions which are characterized by low ovarian hormone production comprising administering to a patient in need thereof, an effective dose of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, or any combination thereof.

DETAILED DESCRIPTION

[0010] As used herein “adrenal insufficiency” means a condition in which the adrenal glands do not produce adequate amounts of steroid hormones. Such conditions include congenital conditions such as Addison’s disease; and non-congenital adrenal insufficiency.

[0011] As used herein non-congenital adrenal insufficiency includes “acquired conditions” for example: systemic inflammatory response syndrome; infection; inflammation; sepsis; trauma; burns; liver disease, including but not limited to hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia; kidney disease; organ transplant; heavy metal poisoning; auto-immune disease, arthritic disease; liver disease; conditions characterized by low testosterone production, including but not limited to male menopause, erectile dysfunction, or conditions characterized by low ovarian hormone production, including but not limited to menopause.

[0012] Relative adrenal insufficiency also includes conditions in which steroid levels, e.g., cortisol levels, are in the normal range, yet there is an inadequate adrenal response to suppress the inflammatory response to trauma, burns, infection, or sepsis.

[0013] “Systemic inflammatory response syndrome” or “SIRS” refers to is an inflammatory state affecting the whole body and can be caused by ischemia, inflammation, trauma, infection, or a combination of several insults, for example sepsis.

[0014] The term “treating” or other forms of the word such as “treatment”, or “treat” is used herein to mean that administration of a compound of the present invention mitigates a disease or a disorder in a host and/or reduces, inhibits, or eliminates a particular characteristic or event associated with a disorder (e.g., reduced steroidogenesis). Thus, the term "treatment" includes, preventing a disorder from occurring in a host, particularly when the host is predisposed to acquiring the disorder; inhibiting the disorder; and/or alleviating or reversing the disorder. Insofar as the methods of the present invention are directed to preventing disorders, it is understood that the term "prevent" does not require that the disease state be completely thwarted. Rather, as used herein, the term preventing refers to the ability of the skilled artisan to identify a population that is susceptible to disorders, such that administration of the compounds of the present invention may occur prior to onset of a disease. The term does not imply that the disease state be completely avoided.

[0015] Throughout the description and claims of this specification the word “comprise” and other forms of the word, such as “comprising” and “comprises,” means including but not limited to, and is not intended to exclude, for example, other additives, components, integers, or steps.

[0016] As used herein, the singular forms “a”, “an”, and “the” include plural references unless the context clearly dictates otherwise

[0017] “Between” as used herein is inclusive, e.g., “between 1 mg and 5000 mg” includes 1 mg and 5000 mg.

[0018] “From” as used herein is inclusive, e.g., “from 1 mg to 5000 mg” includes 1 mg and 5000mg.

[0019] “SC” means subcutaneous injection.

[0020] “IV” means intravenous injection or infusion.

[0021] “IM” means intramuscular injection.

[0022] “FC” is an abbreviation for free cholesterol and as used herein means non-esterified cholesterol.

[0023] “CE” is an abbreviation for cholesteryl ester

[0024] “CDP” and “cholesteryl ester delivery particle” are used interchangeably and as used herein, means a molecule that is able to provide CE to steroidogenic tissue, including but not limited to: a complex of apolipoprotein AI, phospholipid and cholesteryl ester; a

complex of apolipoprotein AI, phospholipid, cholesterol, and cholesteryl ester; a complex of one or more amphipathic peptide of 18- 40 amino acids in length, phospholipid and cholesteryl ester; a complex of one or more amphipathic peptides of 18- 40 amino acids in length, phospholipid, cholesterol and cholesteryl ester; or native HDL. Phospholipids suitable for use in producing CDP include, but are not limited to, phosphatidylcholine, sphingomyelin phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol, cardiolipin and mixtures thereof. The term CDP encompasses, but is not limited to, rHDL, mHDL, and native HDL.

[0025] “rHDL” and “reconstituted HDL” are used interchangeably and as used herein, means a complex of apolipoprotein AI, phospholipid and cholesteryl ester; or a complex of apolipoprotein AI, phospholipid, cholesterol, and cholesteryl ester.

Phospholipids suitable for use in producing rHDL include, but are not limited to, phosphatidylcholine, sphingomyelin phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol, cardiolipin and mixtures thereof.

[0026] “mHDL” and “mimetic HDL”, are used interchangeably and as used herein means a complex of one or more amphipathic peptide of 18- 40 amino acids in length, phospholipid and cholesteryl ester; or a complex of one or more amphipathic peptides of 18-40 amino acids in length, phospholipid, cholesterol and cholesteryl ester. Phospholipids suitable for use in producing mHDL include, but are not limited to, phosphatidylcholine, sphingomyelin phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol, cardiolipin and mixtures thereof. Amphipathic peptides suitable for use in producing CDP include, but are not limited to, D4F (Song et. al, (2009) Int J Biol Sci 5:637-646), (A.V. Bocharov et al.(2004) J. Biol. Chem. 279: 36072-36082), 5A and analogs (W. D'Souza et al (2010) Circ. Res. 107:217-227), A-IConA (G.M. Anantharamaiah (2007) J. Lipid Res. 48:1915-1923), trimeric apoAI variants (Graversen, et al, 2008); Ac-hE18A-NH₂ (Datta, et al, 2000; 2001), and peptides disclosed in D Busseuil et al. (2008) Br J Pharmacol. 154(4): 765-773.

[0027] “Native HDL” means HDL isolated from plasma. These particles contain phospholipid, proteins, cholesterol and cholesteryl ester.

[0028] As used herein, a “normal level of HDL-CE” means a plasma concentration of HDL-CE that is present in an average healthy untreated subject not currently on any medication which might alter HDL-CE levels.

[0029] As used herein “a normal level of a steroid” means a plasma concentration of a particular steroid that is present in an average healthy untreated subject not currently on any medication which might alter the level of that particular steroid.

[0030] “LCAT” is an abbreviation for lecithin-cholesterol acyltransferase.

[0031] “LCAT” or “LCAT polypeptide” when used herein encompass native sequence LCAT, LCAT variants, modified LCAT, LCAT derivatives and chimeric LCAT. A “native sequence LCAT” comprises a polypeptide having the same amino acid sequence as a LCAT derived from nature. Thus, a native sequence LCAT specifically encompasses naturally occurring truncated forms of LCAT, and naturally-occurring allelic variants of LCAT, naturally-occurring variant forms (e.g., alternately spliced forms). The preferred native sequence LCAT is a mature native sequence LCAT.

[0032] “Modified LCAT” means a polypeptide which has LCAT activity, wherein one or more amino acids in the native LCAT polypeptide is substituted with another amino acid, or one or more amino acids is added, or one or more amino acids is deleted from, to a portion of the native polypeptide, including, but not limited to, the N-terminal or C-terminal amino acid. Examples of substitutions include, without limitation, conservative amino acid substitutions, or substitutions with non-naturally occurring amino acids. Non-limiting exemplary conservative amino acid substitutions are provided in Table 1. For example and without limitation the modified LCAT may be a modified LCAT protein as described in United States Patent Publication No. 2009/0081182.

[0033] Derivatives of LCAT include native or modified LCAT polypeptides which have been altered to improve the activity, solubility, absorption, and/or biological half life. One of skill in the art would be familiar with methods for derivatizing polypeptides to improve their pharmacologic properties.

TABLE 1

Original Residue	Exemplary Conservative Substitutions
A	G, S
R	K
N	Q, H
D	E
C	S
Q	N

E	D
G	A, P
H	N, Q
I	L, V
L	I, V
K	R, Q, E
M	L, Y, I
F	M, L, Y
S	T
T	S
W	Y
Tyr	W, F
Val	I, L

[0034] An “agent that increases LCAT plasma activity” includes agents which increase the activity of LCAT enzyme or increase the plasma level of LCAT or both. Such agents include, but are not limited to, small molecules and biologics. For example and without limitation, compounds described in United States patent publication number US 2008/0096900.

[0035] “ApoA-I”, “ApoA-I polypeptide” and “Apolipoprotein A-I” are used interchangeably herein.

[0036] “ApoA-I” as used herein encompasses native sequence apoA-I, modified apoA-I, apoA-I derivatives, and chimeric apoA-I.

[0037] A “native sequence apoA-I” comprises a polypeptide having the same amino acid sequence as an apoA-I polypeptide derived from nature. Thus, a native sequence apoA-I specifically encompasses naturally occurring truncated forms of apoA-I, and naturally-occurring allelic variants of apoA-I, naturally-occurring variant forms (e.g., alternately spliced forms), and pre-pro or pro- apoA-I. The preferred native sequence apoA-I is a mature native sequence apoA-I.

[0038] “Modified apoA-I” means a polypeptide having an amino acid sequence which differs from a native apoA-I polypeptide sequence in that one or more amino acids is substituted with a different amino acid, or one or more amino acids is added, or one or more amino acids is deleted and wherein the polypeptide retains the ability to form a CDP.

Examples of such substitutions include, without limitation, conservative substitutions, or substitutions with non-naturally occurring amino acids.

[0039] Derivatives of apoA-I include native or modified Apo A-I polypeptides which have been altered to improve the solubility, absorption, biological half life and wherein the derivative retains the ability to form a CDP. Derivatives of polypeptides are well known in the art e.g., pegylation. One of skill in the art would be familiar with method for derivatizing polypeptides to improve their pharmacologic properties.

[0040] HDL-C refers to HDL which comprises cholesterol with or without cholesteryl ester. (where C represents total cholesterol and is comprised of both cholesterol and cholesteryl ester)

[0041] HDL-CE refers to the cholesteryl ester component of HDL.

[0042] The term “effective amount”, as used herein, means an amount of LCAT, a compound that increases plasma LCAT activity, CETP inhibitor, a CDP, or any combination thereof, effective at dosages and for periods of time necessary to achieve the desired or therapeutic result. An effective amount may vary according to factors known in the art, such as the disease state, age, sex, and weight of the subject being treated. Although particular dosage regimens may be described in examples herein, a person skilled in the art would appreciate that the dosage regimen may be altered to provide optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. In addition, the compositions of this disclosure can be administered as frequently as necessary to achieve the optimum therapeutic response.

[0043] Lipids suitable for use in producing the compositions of the present disclosure include, but are not limited to natural and synthesized (synthetic) lipids and phospholipids, phosphatidylcholine, sphingomyelin phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol, cardiolipin; phosphatidylcholine in which a fatty acid is esterified to the SN-1 position of the glycerol backbone of phosphatidylcholine in which the fatty acids include but are not limited to myristic, palmitic, palmitoleic, oleic and stearic acids; phosphatidylcholine in which a fatty acid is esterified to the SN-2 position of the glycerol backbone of phosphatidylcholine, in which the fatty acids include but are not limited to myristic, palmitic, palmitoleic, oleic and stearic acids; phosphatidylcholine in which the SN-1 or SN-2 positions of the glycerol backbone is linked by ether bonds to fatty alcohols including tetradecanol, hexadecanol and octadecanol; included also are

phospholipids as described wherein the moiety esterified to the phosphate group in place of choline is ethanolamine, serine, glycerol or inositol.

[0044] Subjects having conditions associated with adrenal insufficiency often have decreased levels of HDL-CE. Such conditions, include, but not limited to, systemic inflammatory response syndrome, infection, inflammation, auto-immune diseases, arthritic diseases, sepsis, trauma, burns, liver disease (e.g. hepatitis, fibrosis, cirrhosis, bile duct hyperplasia, bile duct atresia, organ transplant, heavy metal poisoning), and kidney disease.

[0045] Although not a primary pathology in these diseases, the decreased HDL-CE reduces the delivery of cholesterol to steroidogenic tissues, thus, decreasing the body's ability to produce critical glucocorticoids, hormones and other steroids. The adrenal gland does not store cortisol, it relies on cholesterol derived from the cholesteryl ester that is supplied by HDL-CE for synthesis of cortisol. This reduced capacity for steroidal production can exacerbate of the underlying conditions leading to serious complications. The present disclosure provides for methods of prophylactically elevating HDL-CE levels in patients who may have normal to elevated HDL-C when presenting following trauma, severe infection or prior to major surgery, in order to prevent the significant drop in HDL-C that routinely occurs in these patients. In addition, the present disclosure provides methods for treating patients currently have reduced HDL-C levels, following trauma or severe infection in order to prevent a decreased level of HDL-C levels.

[0046] LCAT is responsible for the production of HDL-CE through the esterification of the free cholesterol component of HDL-C. Thus, increasing LCAT levels or LCAT activity increases the amount of HDL-CE available for delivery to steroidogenic tissues. For example, to the adrenal glands thereby allowing for replenishment of the adrenal cholesterol stores which are used in the production of corticosteroids. The LCAT level and/or LCAT activity can be increased by any means available.

[0047] Cholesteryl ester transfer protein (CETP), transfers cholesteryl esters from HDL to very low-density lipoproteins and LDL in exchange for triglycerides. CETP activity thus reduces the level of HDL-CE. CETP inhibitors block the transfer of CE from HDL, thus increasing the amount of CE available in HDL for delivery to steroidogenic tissues. CETP inhibitors include, but are not limited to, torcetrapib, anacetrapib, dalcetrapib (JTT-705) and those described in U.S. Patent 7,470,705.

[0048] The level of cholesteryl ester in a patient can also be increased by administering to the patient an effective amount of a CDP according to the present invention.

In a particular embodiment the CDP is native HDL. In another embodiment the CDP is rHDL. In yet another embodiment the CDP is mHDL.

[0049] Accordingly, the present disclosure provides methods for modulating lipid content of steroidogenic tissues. In another embodiment the lipid content is modulated by administering an effective dose of LCAT. In a particular embodiment the lipid content is modulated by administering an effective dose of a compound that increases plasma LCAT activity. In another embodiment the lipid content is modulated by administering an effective dose of a CETP inhibitor. In another embodiment, the lipid content is modulated by administering an effective dose of a CDP. In another embodiment the lipid content is modulated by administering a effective dose of a CDP wherein the CDP is rHDL, mHDL, or native HDL. In a particular embodiment the lipid content is modulated by administering an effective dose of rHDL. In another embodiment the lipid content is modulated by administering an effective dose of mHDL. In a further embodiment the lipid content is modulated by administering an effective dose of native HDL. In another embodiment the lipid content is modulated by administering in combination, two or more agents independently selected from LCAT, a compound that increases plasma LCAT activity, native HDL, rHDL, mHDL and CETP inhibitors. The two or more agents can be administered in any order, for example, simultaneously or sequentially.

[0050] An embodiment of the present disclosure is a method of increasing steroidogenesis in a patient in need thereof comprising: administering to a patient in need thereof, an effective dose of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, or any combination thereof, whereby the steroid level is increased. In a particular embodiment the CDP is native HDL. In another embodiment the CDP is reconstituted HDL. In another embodiment the CDP is mimetic HDL. In one embodiment the steroid is a corticosteroid or sex hormone. In another embodiment the steroid is cortisol. In a further embodiment the steroid is aldosterone. In yet another embodiment the steroid is testosterone. In still another embodiment the steroid is an estrogen. An further embodiment of the present disclosure is a method of increasing steroidogenesis in a patient in need thereof comprising: administering to a patient in need thereof, an effective dose of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, or any combination thereof, whereby the steroid level is increased to about a normal level. Another embodiment of the present disclosure is a method of increasing steroidogenesis in a patient in need thereof comprising: administering to a patient in need thereof, an effective dose of an

agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, or any combination thereof, whereby the steroid level is increased to above a normal level.

[0051] One embodiment of the present invention is a method of increasing cortisol in a patient in need thereof comprising: administering to a patient in need thereof, an effective dose of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof. In a particular embodiment the CDP is native HDL. In another embodiment the CDP is reconstituted HDL. In another embodiment the CDP is mimetic HDL. In another embodiment the agent is LCAT. In another embodiment the agent is a compound that increases plasma LCAT activity.

[0052] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0053] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof.

[0054] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute

condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0055] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0056] Liver disease includes, but is not limited to, hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia. Thus, one embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of an agent selected from the group consisting of LCAT, a compound that increases plasma LCAT activity, a CETP inhibitor, a CDP, and any combination thereof. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0057] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of LCAT. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0058] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of LCAT.

[0059] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of

LCAT. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0060] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of LCAT. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0061] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of LCAT. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0062] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of a compound that increases plasma LCAT activity. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0063] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of a compound that increases plasma LCAT activity.

[0064] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of a compound that increases plasma LCAT activity. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0065] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of a compound that increases plasma LCAT activity. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0066] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of a compound that increases plasma LCAT activity. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0067] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of a CETP inhibitor. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0068] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute

condition, comprising administering to a patient in need thereof, an effective amount of a CETP inhibitor.

[0069] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of a CETP inhibitor. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0070] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of a CETP inhibitor. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0071] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of a CETP inhibitor. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0072] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of a CDP. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0073] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of a CDP.

[0074] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of a CDP. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0075] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of a CDP. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0076] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of a CDP. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0077] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of native HDL. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment

the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0078] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of native HDL.

[0079] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of native HDL. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0080] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of a native HDL. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0081] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of native HDL. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0082] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising:

administering to a patient in need thereof, an effective amount of rHDL. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0083] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of rHDL.

[0084] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of rHDL. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0085] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of rHDL. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0086] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of rHDL. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0087] Another embodiment according to the present disclosure is a method of treating a condition characterized by reduced function of steroidogenic tissues comprising: administering to a patient in need thereof, an effective amount of mHDL. In a particular embodiment, the condition is characterized by adrenal insufficiency. In another embodiment the condition is low testosterone production. In another embodiment the condition is low ovarian hormone production.

[0088] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of mHDL.

[0089] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency, wherein the condition is an acute condition, comprising administering to a patient in need thereof, an effective amount of mHDL. In one embodiment the acute condition is systemic inflammatory response syndrome. In another embodiment the acute condition is infection. In another embodiment the condition is inflammation. In yet another embodiment the acute condition is sepsis. In still another embodiment the acute condition is trauma. In another embodiment the acute condition is burn. In yet another embodiment the acute condition is liver disease. In still another embodiment the acute condition is kidney disease. In another embodiment the acute condition is organ transplant. In yet another embodiment the acute condition is heavy metal poisoning.

[0090] Another embodiment according to the present disclosure is a method of treating a condition characterized by adrenal insufficiency wherein the condition is a chronic condition, comprising administering to a patient in need thereof, an effective amount of mHDL. In one embodiment the chronic condition is an auto-immune disease. In another embodiment the chronic condition is arthritic disease. In yet another embodiment the chronic condition is liver disease. In still another embodiment the chronic condition is kidney disease.

[0091] Another embodiment of the present invention is a method of treating a liver disease wherein the liver disease is hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, or bile duct atresia, comprising administering to a patient in need thereof, an effective amount of mHDL. In a particular embodiment the liver disease is hepatitis. In another embodiment the liver disease is hepatorenal syndrome. In yet another embodiment the liver disease is fibrosis. In still another embodiment the liver disease is cirrhosis. In

another embodiment the liver disease is bile duct hyperplasia. In another embodiment the liver disease is bile duct atresia.

[0092] In certain conditions, it may be desirable to immediately increase the amount of HDL-CE to a normal or above normal level and then to maintain a normal or increased level of HDL-CE. Such conditions are generally acute conditions such as sepsis, major surgical trauma, or severe burns. One of skill in the art would understand which conditions would require an immediate increase in the amount of HDL-CE. Thus, an embodiment of the present disclosure comprises administering, to a subject in need thereof, an effective amount of a CDP as an initial dose and simultaneously or subsequently administering to the subject an amount of LCAT sufficient to maintain a desired level of HDL-CE. In preferred embodiments the desired plasma level of HDL-C is greater than 35 mg/dl.

[0093] Another embodiment is a method of treating a condition which is characterized by adrenal insufficiency or can lead to adrenal insufficiency, comprising administering, to a patient in need thereof, an effective amount of a CDP as an initial dose and then administering an amount of LCAT sufficient to maintain a desired level of HDL-CE. A particular embodiment is a method of treating sepsis comprising administering, to a patient in need thereof, an effective amount of a CDP as an initial dose and then administering an amount of LCAT sufficient to maintain a desired level of HDL-CE. In preferred embodiments the desired plasma level of HDL-C is greater than 35 mg/dl.

[0094] Another embodiment of the present disclosure comprises administering an effective amount of a CDP as an initial dose and then administering an amount of a CETP inhibitor sufficient to maintain a desired level of HDL-CE. Another embodiment is a method of treating a condition which is characterized by adrenal insufficiency or can lead to adrenal insufficiency, comprising administering, to a patient in need thereof, an effective amount of a CDP as an initial dose and then administering an amount of a CETP inhibitor sufficient to maintain a desired level of HDL-CE. A particular embodiment is a method of treating sepsis comprising administering, to a patient in need thereof, an effective amount of a CDP as an initial dose and then administering an amount of CETP inhibitor sufficient to maintain a desired level of HDL-CE.

[0095] Patients who develop systemic inflammatory response syndrome (SIRS) frequently have reduced levels of HDL-CE. Those patients with reduced levels of HDL-CE have a worse prognosis than those with normal levels of HDL-CE. Thus, an embodiment of the invention is a method of treating systemic inflammatory response syndrome comprising administering to a patient with systemic inflammatory response syndrome an effective amount

of LCAT. In a particular embodiment the effective amount of LCAT is an amount sufficient to maintain the normal level of HDL-CE. In another embodiment the effective amount of LCAT is an amount sufficient to increase the level of HDL-CE above the normal level. Conditions related to SIRS include, but are not limited to, sepsis, infection, trauma (including surgery), burns and pancreatitis. Thus, an embodiment of the present disclosure is a method of treating a patient with sepsis, infection, trauma, surgical trauma, burn, or pancreatitis comprising administering an effective amount of LCAT.

[0096] Another embodiment of the invention is a method of treating systemic inflammatory response syndrome comprising administering to a patient with systemic inflammatory response syndrome an effective amount of a CETP inhibitor. In a particular embodiment the effective amount of the CETP inhibitor is an amount sufficient to maintain the normal level of HDL-CE. In another embodiment the effective amount a CETP inhibitor is an amount sufficient to increase the level of level of HDL-CE above the normal level.

[0097] A person of ordinary skill in the art, such as a physician, would understand that, depending on the condition and severity of the condition, that different combinations of LCAT, a compound that increases plasma LCAT activity, CETP inhibitor, and CDP could be administered. The person of ordinary skill in the art would be able to determine which combination would be appropriate. For example a physician may decide to administer LCAT with a CDP, or LCAT with a CETP inhibitor, or a CDP with a CETP inhibitor, depending on the severity of crisis. Alternately, the patient may be administered LCAT (requires intravenous route) while still in the hospital critical care unit, and then discharged with instructions to take a CETP inhibitor (which are orally administered) for several days or weeks until stabilized.

[0098] Conditions such as infection, inflammation, auto-immune disease, arthritic disease, sepsis, trauma, burns, liver disease heart disease and kidney disease are sometimes associated with low testosterone levels as a result of testicular insufficiency. Low testosterone levels can lead to low testosterone syndrome (or male menopause), erectile dysfunction and other conditions. Increasing the amount of HDL-CE available for delivery to the Leydig cells in testes, allows for the replenishing of testicular cholesterol stores and normalization of testosterone production. Thus, another embodiment according to the present disclosure is a method of treating a condition characterized by low testosterone production comprising administering to a patient in need thereof, an effective dose of a CETP inhibitor, LCAT, a compound that increases plasma LCAT activity, a CDP, or any combination thereof.

[0099] Another embodiment according to the present disclosure is a method of treating a condition which is characterized by low ovarian hormone production comprising administering to a patient in need thereof, an effective dose of a CETP inhibitor, LCAT, a compound that increases plasma LCAT activity, a CDP, or any combination thereof.

[00100] LCAT, compounds that increases plasma LCAT activity, CDPs and CETP inhibitors according to the present disclosure, can be used in combination therapy with other drugs. Such therapies include, but are not limited to simultaneous or sequential administration of the drugs involved. For example, LCAT formulations can be administered with drugs that are commonly used as a standard of care for a particular condition.

Formulations

[00101] LCAT, compounds that increase plasma LCAT activity, CDPs, and CETP inhibitors, administered in the methods of the present disclosure can be formulated as pharmaceutical compositions and administered to a mammalian subject, such as a human patient in a variety of forms adapted to the chosen route of administration, i.e., orally, parenterally, by intravenous, intramuscular or subcutaneous routes.

[00102] Pharmaceutical compositions suitable for the delivery of compounds of the present disclosure and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation may be found, for example, in Remington's Pharmaceutical Sciences, 19th Edition (Mack Publishing Company, 1995).

[00103] Suitable dosage forms for oral administration include, for example, solid, semi-solid and liquid systems such as in hard or soft shell gelatin capsules, tablets, liquids, powders, lozenges (including liquid-filled), chews, gels, films, ovules, sprays, elixirs, suspensions, syrups, buccal/mucoadhesive patches and the like.

[00104] Oral dosage forms may, for example, contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active

compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and devices. The active compound may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[00105] The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Dosage

[00106] In an embodiment of the present disclosure LCAT is administered by subcutaneous injection. In another embodiment LCAT is administered by intramuscular injection. In another embodiment LCAT is administered by intravenous injection or infusion. In some embodiments the LCAT is self-administered by the patient either by subcutaneous or intramuscular injection. Self-administration of LCAT is a preferred embodiment for chronic

treatment, including, but not limited to auto-immune diseases, arthritic diseases, and chronic liver disease.

[00107] An effective daily dose of LCAT is from 1 mg to 5000 mg, or 1 mg to 2000 mg, or 10 mg to 5000 mg, or 10 mg to 1000 mg, 10 mg to 500 mg or 5 to 100 mg.

[00108] In an embodiment of the present disclosure a CDP is administered by subcutaneous injection. In another embodiment a CDP is administered by intramuscular injection. In another embodiment a CDP is administered by intravenous injection or infusion. An effective amount of a CDP is between 100 mg and 5000 mg, or between 200 mg and 20,000 mg, or between 250 mg and 10,000 mg or between 250 mg and 2000 mg, or between 500 mg and 5000 mg, between 250 mg and 1000 mg, or between 500 mg and 5000 mg or between 500 mg and 10,000 mg. An effective amount of a CETP inhibitor will vary based on the specific CETP inhibitor that is administered. An effective dose of a CETP inhibitor is between 1 and 5000 mg / day, between 10 and 1000 mg/day, between about 10 and 500 mg/day or between 1 and 250 mg/day. The specific dosage used can vary. For example, the dosage can depend on a number of factors including, but not limited to, the dosing frequency, the specific activity of the recombinant LCAT enzyme, the body weight of the patient, special requirements of the patient, special conditions of the patient (e.g., abnormal kidney or liver function), the condition being treated, etc. The dosing frequency and amount may, at the physician's discretion, fall outside of the typical range given herein. These dosages are based on an average human subject having a weight of 70kg. Determination of optimum dosages for a particular patient is well-known to those skilled in the art. The physician will readily be able to determine doses for subjects whose weight falls outside this range, such as infants and the elderly.

[00109] Depending on the disorder and the patient being treated, one skilled in the art (i.e. a physician) could determine the appropriate dose of the agent or agents depending on the condition of the patient. For example for a patient presenting with sepsis the physician might administer an initial dose of a CDP to rapidly increase HDL-C levels to a desired level and follow that treatment with an agent to maintain the desired level of HDL-C.

[00110] The efficacy of a particular dose may be assessed by reference noting the change in HDL-C or improvement in certain physiologic parameters. Suitable physiologic parameters include, but are not limited to, improved cortisol response to ACTH, improved cortisol levels, change in circulating levels of steroids, e.g., corticosteroids, glucocorticoids, cortisol, testosterone, estrogen, decreased inflammatory cytokines, and improved liver function.

[00111] For example the therapeutically effective dose of one or a combination of agents may be the amount necessary to increase levels of HDL-C and/or maintain levels of HDL-C at or above 30 mg/dl, or at or above 35 mg/dl, or at or above 40 mg/dl, or at or above 45 mg/dl, or at or above 50 mg/dl. Preferably the level of HDL-C is at or above 35 mg/dl. Or for example the therapeutically effective amount of one or a combination of agents may be the amount required to raise or maintain cortisol levels at or above 15 µg/dl, or at or above 20 µg/dl, or at or above 25 µg/dl or at or above 30 µg/dl or at or above 35 µg/dl. Preferably the cortisol level is at or above 25 µg/dl.

[00112] Measurement of biomarker levels and parameters described above may be measured using methods that are well known in the art. For example, change in HDL-CE level is easily detected with standard clinical laboratory procedures. Likewise, measuring the adrenal response to ACTH is commonly used in patients with suspected adrenal insufficiency. One of ordinary skill in the art would understand the significance of the results and may choose to adjust the dose based on assessments such as those described above.

[00113] Embodiments of compounds, compositions and methods are illustrated in the following examples. These examples are provided for illustrative purposes and are not considered limitations on the scope of compounds, compositions and methods of the present disclosure. The scope of this invention is to be defined by the claims rather than by the specifically described embodiments and examples.

Assays

[00114] Efficacy of an agent of the present invention can be determined by any suitable method. For example, one method for determining the activity of an agent or composition according to the present invention is by using an ACTH stimulation assay which measures the adrenal response to adrenocorticotrophic hormone (ACTH) by measuring cortisol production under various experimental conditions. For example, intramuscular injection of rabbits with croton oil causes an acute inflammatory response and a reduction of plasma HDL by greater than 50% within 3-days after injection. The observed inflammatory response and reduction of HDL are similar to the conditions observed in critical care patients at risk for developing adrenal insufficiency. Using this rabbit model, an efficacy of an agent of the present invention can be evaluated by measuring the recovery of cortisol response in stressed rabbits with low HDL in response to administration of the agent. The following experimental procedure is provided as a specific, non-limiting example.

[00115] Three days prior to treatment, twenty-four (24) rabbits are injected with 0.1 ml ACTH (1mg/ml), and cortisol measured at 0.5, 1, 2, and 3 hours post-stimulus. The animals are then allocated, based on cortisol response to the ACTH challenge, into groups as follow: negative control group (saline), positive control group (croton oil only), croton oil plus 10 mg/kg LCAT, croton oil plus 30 mg/kg LCAT, croton oil plus 50 mg/kg LCAT, croton oil plus 10 mg/kg rHDL, croton oil plus 30 mg/kg rHDL, croton oil plus 50 mg/kg rHDL. At time = 0 each group is administered IM injections of croton oil or saline (negative control group). At 72 and 96 hours, animals are treated with saline or LCAT or rHDL. At 98 hours (2 hours post last treatment), all animals are injected with 0.1 ml ACTH (1mg/ml), and cortisol is measured at 0.5, 1, 2, and 3 hours post ACTH challenge. The animals in the positive control group (croton oil only) are expected to show a markedly reduced production of cortisol in comparison with the negative control group. The efficacy of the agent at each dose is determined by comparison with the cortisol levels of the control groups.

[00116] Similarly, the effect of agents according to the present invention may be assessed for efficacy in treating low testosterone production or low ovarian hormone production using assays known in the art.

EXAMPLES

Example 1.

[00117] A patient is admitted into the hospital for major surgery. The patients HDL-C level is 25 mg/dl. Based on epidemiological studies, this patient is at a significantly increased risk for infection and in-hospital death following surgery. The patient is administered a dose of LCAT of 40 mg 24 hours prior to surgery. The patient's HDL-C is measured prior to surgery, and is now above 50 mg/dl, thus greatly reducing risk of infection and/or death following the procedure.

Example 2.

[00118] A burn victim is stabilized in the hospital with an HDL-C level of 12 mg/dl. Based on epidemiological studies, this patient's risk of going into shock and dying is greatly increased over the next few days due to the reduced ability of his adrenals to manufacture corticosteroids. The patient is infused with rHDL at a dose of 25 mg/kg over a period of 5 hours, raising the patient's HDL-C by 50 mg/dl. In order to maintain this level of HDL, the patient is administered a 60 mg dose of LCAT. Cortisol levels are monitored periodically to determine if the adrenal function is normalized. If adrenal function is still reduced, an additional dose of rHDL or LCAT may be administered.

Example 3.

[00119] A victim of a car accident with abdominal injuries is brought to the hospital. Her HDL is 55 mg/dl. As HDL routinely falls after trauma, the physician prescribes 150 mg/day anacetrapib and an injection of LCAT (40 mg) every 5 days until the patient is fully stabilized. This treatment maintained the patient's HDL at >75 mg/dl throughout her recovery. Adrenal function was maintained.

Example 4.

[00120] A patient is beginning to enter septic shock. Cortisol levels are <25 μ g/dl. The patient is infused with native HDL at a dose of 25 mg/kg over a period of 5 hours. The patient's HDL-C level is increased by 50 mg/dl. In order to maintain this level of HDL-C, the patient is administered a 60 mg bolus of LCAT. To aid in maintaining the HDL-CE, the patient is also given torcetrapib (120 mg b.i.d.). The patient is maintained on daily torcetrapib and a weekly LCAT injection during recovery. Cortisol levels are monitored to determine if the adrenal function is normalized. Dosages of torcetrapib and LCAT are adjusted as required.

Example 5.

[00121] A patient complains of lethargy and impotence. His testosterone levels are low, and his HDL-C is 25 mg/dl. The physician prescribes anacetrapib at a dose of 150 mg/day. For weeks later his testosterone level is rechecked, and is now within normal levels. The patient's original complaints are absent.

Example 6.

Preparation of rHDL.

[00122] An aqueous suspension of 1 mole part phosphatidylcholine, 0.2 mole part cholesterol and 0.1 mole part cholesteryl ester is sonicated in an aqueous solution containing apolipoprotein A-I as described in Song et al, Int J Biol Sci 5:637-646. The lipids are co-dissolved in a solvent or solvent mixture. Solvents are removed by evaporation and vacuum. An aqueous solution of apolipoprotein A-I is added to the dried lipids (at 1 mole part apolipoprotein A-I to every 100 mole parts phospholipid.) The protein and lipids are dispersed by ultrasonication yielding rHDL.

Example 7.

Alternate process for the preparation of rHDL.

[00123] A lyophilized cake of phosphatidylcholine, cholesterol and cholesteryl ester is rehydrated with an aqueous solution containing apolipoprotein A-I. The cake consists of phospholipid, cholesterol and cholesteryl ester in a mole ratio of 1 mole part

phosphatidylcholine, 0 to 0.5 mole parts cholesterol and 0 to 0.5 mole parts cholesteryl ester, as above. The lyophilized cake is formed by dissolving all lipids in a solvent that will form a solid at a temperature suitable for lyophilization (e.g. dioxane). The lyophilized cake is rehydrated in an aqueous solution of apolipoprotein A-I containing 1 mole part apolipoprotein A-I for every 25 to 250 mole parts phospholipid. Homogenization techniques such as heating, ultrasonication or passage through micropore filters may be necessary to obtain a stable and uniform size of rHDL.

Example 8.

Preparation of mHDL.

[00124] A lyophilized cake of 1 mole part phosphatidylcholine, 0 to 0.5 mole parts cholesterol, 0 to 0.5 mole parts cholesteryl ester and 0.1 to 1 mole part peptide are dissolved in dioxane alone or with an additional solvent such as methanol or acetic acid to aid in peptide dissolution. The solution is lyophilized. The resultant dried cake is reconstituted in an aqueous salt solution. The reconstitution process may require homogenization techniques such as heating, ultrasonication or passage through micropore filters to obtain a stable and uniform size of mHDL.

CLAIMS

1. Native sequence Lecithin:cholesterol acyltransferase (LCAT) for use in the treatment of a patient having a condition characterized by reduced function of steroidogenic tissues when compared to normal function, wherein the steroid level is increased, and wherein the condition is an acute condition selected from the group consisting of systemic inflammatory response syndrome, an autoimmune disease, arthritic disease, infection, inflammation, sepsis, trauma, burns, liver disease, kidney disease, organ transplant, and heavy metal poisoning.
2. Native sequence Lecithin:cholesterol acyltransferase (LCAT) for use to increase steroidogenesis of a steroid in a patient suffering from a condition wherein the function of steroidogenic tissues is reduced when compared to normal function, whereby the steroid level is increased, and wherein the condition is an acute condition selected from the group consisting of systemic inflammatory response syndrome, an autoimmune disease, arthritic disease, infection, inflammation, sepsis, trauma, burns, liver disease, kidney disease, organ transplant, and heavy metal poisoning.
3. The native sequence LCAT according to claim 1 or 2, wherein the liver disease is selected from the group consisting of hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, and bile duct atresia.
4. The native sequence LCAT according to claim 2, wherein the steroid is a corticosteroid or sex hormone.
5. The native sequence LCAT according to claim 4, wherein the steroid is cortisol, aldosterone, testosterone or an estrogen.
6. A composition comprising the native sequence LCAT according to any one of claims 1 to 5, and a cholesterol delivery particle (CDP).

7. The composition according to claim 6, wherein the CDP is native high density lipoproteins (HDL), reconstituted HDL or mimetic HDL, wherein:

- i) native HDL means HDL isolated from plasma;
- ii) reconstituted HDL means:
 - a) a complex of apolipoprotein A1, a phospholipid and a cholesteryl ester;
or
 - b) a complex of apolipoprotein A1, a phospholipid, cholesterol and a cholesteryl ester; and
- iii) mimetic HDL means:
 - a) a complex of one or more amphipathic peptides of 18-40 amino acids in length, a phospholipid and a cholesteryl ester; or
 - b) a complex of one or more amphipathic peptides of 18-40 amino acids in length, a phospholipid, cholesterol and a cholesteryl ester.

8. The composition according to claim 6 or 7, wherein the CDP is in the form of a bolus dose adapted for administration to the patient prior to administration of the native sequence LCAT.

9. The native sequence LCAT according to any one of claims 1-5 or the composition according to any one of claims 6-8, wherein the steroid level is increased to a normal level or is increased to above a normal level.

10. The native sequence LCAT according to any one of claims 1-5 or the composition according to any one of claims 6-8, wherein the native sequence LCAT or the composition is adapted for administration in a dose, or in multiple doses, effective to maintain a desired steroid level.

11. The native sequence LCAT or the composition of claim 10, wherein the desired steroid level is a normal level.

12. Use of native sequence Lecithin:cholesterol acyltransferase (LCAT) in the manufacture of a medicament for the treatment of a condition characterized by reduced function of

steroidogenic tissues in a patient in need thereof when compared to normal function, wherein the steroid level is increased, and wherein the condition is an acute condition selected from the group consisting of systemic inflammatory response syndrome, an autoimmune disease, arthritic disease, infection, inflammation, sepsis, trauma, burns, liver disease, kidney disease, organ transplant, and heavy metal poisoning.

13. Use of native sequence Lecithin:cholesterol acyltransferase (LCAT) in the manufacture of a medicament for increasing steroidogenesis of a steroid in a patient suffering from a condition wherein the function of steroidogenic tissues is reduced when compared to normal function, whereby the steroid level is increased, and wherein the condition is an acute condition selected from the group consisting of systemic inflammatory response syndrome, an autoimmune disease, arthritic disease, infection, inflammation, sepsis, trauma, burns, liver disease, kidney disease, organ transplant, and heavy metal poisoning.

14. The use according to claim 12 or 13, wherein the liver disease is selected from the group consisting of hepatitis, hepatorenal syndrome, fibrosis, cirrhosis, bile duct hyperplasia, and bile duct atresia.

15. The use according to claim 13, wherein the steroid is a corticosteroid or sex hormone.

16. The use according to claim 15, wherein the steroid is cortisol, aldosterone, testosterone or an estrogen.

17. The use of any one of claims 12 to 16, wherein the medicament is adapted for use with a cholesterol delivery particle (CDP).

18. The use according to claim 17, wherein the CDP is native high density lipoproteins (HDL), reconstituted HDL or mimetic HDL wherein:

- i) native HDL means HDL isolated from plasma;
 - ii) reconstituted HDL means:
 - a) a complex of apolipoprotein A1, a phospholipid and a cholesteryl ester;
- or

- b) a complex of apolipoprotein A1, a phospholipid, cholesterol and a cholesteryl ester; and
- iii) mimetic HDL means:
 - a) a complex of one or more amphipathic peptides of 18-40 amino acids in length, a phospholipid and a cholesteryl ester; or
 - b) a complex of one or more amphipathic peptides of 18-40 amino acids in length, a phospholipid, cholesterol and a cholesteryl ester.

19. The use according to any one of claims 17 or 18, wherein the CDP has already been administered in a bolus dose prior to administration of the native sequence LCAT.

20. The use according to any one of claims 13-19, wherein the steroid level is increased to a normal level or is increased to above a normal level.

21. The use according to any one of claims 13-19, wherein the native sequence LCAT is adapted for administration in a dose, or in multiple doses, effective to maintain a desired steroid level.

22. The use according to claim 21, wherein the desired steroid level is a normal level.