



HU000025409T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 025 409**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 11 176272**(51) Int. Cl.: **A61K 315/75** (2006.01)(22) A bejelentés napja: **2005. 05. 12.****A61P 35/00** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20050176272

(97) Az európai bejelentés közzétételi adatai:

EP 2392337 A2 **2011. 12. 07.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2392337 B1 **2015. 07. 29.**

(72) Feltaláló(k):

Trauner, Michael, 8010 Graz (AT)**Hofmann, Alan, La Jolla, CA California 92037 (US)****Fickert, Peter, 8010 Graz (AT)**

(73) Jogosult(ak):

Medizinische Universität Graz, 8036 Graz (AT)

(74) Képviselő:

dr. Gál Melinda, SBGK Szabadalmi Ügyvivői
Iroda, Budapest

(54)

A 24-norUDCA alkalmazása

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 392 337 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
29.07.2015 Bulletin 2015/31

(51) Int Cl.:
A61K 31/575 (2006.01) A61P 35/00 (2006.01)

(21) Application number: **11176272.0**

(22) Date of filing: **12.05.2005**

(54) Use of 24-norUDCA

Verwendung von 24-norUDCA

Utilisation de 24-norUDCA

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**
Designated Extension States:
AL BA HR LV MK YU

(43) Date of publication of application:
07.12.2011 Bulletin 2011/49

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
05749537.6 / 1 890 705

(73) Proprietor: **Medizinische Universität Graz**
8036 Graz (AT)

(72) Inventors:
• **Trauner, Michael**
8010 Graz (AT)
• **Hofmann, Alan**
La Jolla, CA California 92037 (US)
• **Fickert, Peter**
8010 Graz (AT)

(74) Representative: **Sonn & Partner Patentanwälte**
Riemergasse 14
1010 Wien (AT)

(56) References cited:

- Engstrom, PF; Sigurdson, E; Evans, AA: "Primary Neoplasms of the Liver"; "96" In: Holland; Frei: "Cancer Medicine e.5", 2000, American Cancer Society - BC Decker Inc, Hamilton, Ontario, CA, XP002664311, ISBN: 1-55009-113-1 pages 1392-1401, * page 1391, left-hand column, paragraph 1 ** page 1397; table 96.5 ** page 1398, left-hand column, paragraph 1 - right-hand column, paragraph 2 *
- TSAGARAKIS N J ET AL: "A concentration-dependent effect of ursodeoxycholate on apoptosis and caspases activities of HepG2 hepatocellular carcinoma cells", EUROPEAN JOURNAL OF PHARMACOLOGY, ELSEVIER SCIENCE, NL, vol. 640, no. 1-3, 25 August 2010 (2010-08-25), pages 1-7, XP027087786, ISSN: 0014-2999 [retrieved on 2010-06-14]
- C. Y. Ponsioen: "Novel developments in IBD-related sclerosing cholangitis", Best Practice & Research Clinical Gastroenterology, 1 April 2011 (2011-04-01), pages S15-S18, XP55013028, Retrieved from the Internet: URL: http://www.sciencedirect.com/science?_ob=MiamiImageURL&_cid=272484&_user=987766&_pii=S1521691811700051&_check=y&_origin=se arch&_zone=rslt_list_item&_coverDate=2011-04-30&wchp=dGLzVIS-zSkzk&md5=a3057a9d9f1d785d12e1654353755ea9/1-s2.0-S1521691811700051-main.pdf [retrieved on 2011-11-24]

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 392 337 B1

- HALLILBASIC E ET AL: "Side chain structure determines unique physiologic and therapeutic properties of norursodeoxycholic acid in Mdr2^{-/-} mice", HEPATOLOGY, WILLIAMS AND WILKINS, BALTIMORE, MD, US, vol. 49, no. 6, 1 June 2009 (2009-06-01), pages 1972-1981, XP002652461, ISSN: 0270-9139
- FICKERT ET AL: "24-norUrsodeoxycholic Acid Is Superior to Ursodeoxycholic Acid in the Treatment of Sclerosing Cholangitis in Mdr2 (Abcb4) Knockout Mice", GASTROENTEROLOGY, ELSEVIER, PHILADELPHIA, PA, vol. 130, no. 2, 1 February 2006 (2006-02-01), pages 465-481, XP005284548, ISSN: 0016-5085, DOI: 10.1053/J.GASTRO.2005.10.018

Description

[0001] The present invention relates to novel uses of 24-nor-ursodeoxycholic acid (*nor*UDCA) as defined in the claims.

[0002] Ursodeoxycholic acid (UDCA), the naturally occurring bile acid, which can be found in small amounts in the bile and in the blood of humans, is a widely used drug to treat liver diseases, wherein one of the most important indication areas of UDCA is the dissolution of gallstones and the treatment of primary biliary cirrhosis (PBC) and primary sclerosing cholangitis (PSC). UDCA is a naturally occurring bile acid with cytoprotective, membrane stabilizing and anti-apoptotic effects. Furthermore, UDCA lowers serum levels of bilirubin, transaminases and alkaline phosphatase as marker of cholestasis (Trauner & Graziadei 1999, Beuers & Paumgartner 2002).

[0003] Studies of UDCA in patients suffering from liver diseases, especially PBC patients, have shown that the administration of UDCA increases the rate of bile flow from the hepatocytes, thereby combating cholestasis and diluting and inhibiting toxic bile acids, which are mainly responsible for the injury to the hepatocytes. Furthermore, UDCA is also able to inhibit the immune response in the liver reducing immunological injury to the bile ducts and the liver. UDCA - as stated above - is regularly used to treat PSC and PBC.

[0004] PSC, which affects mainly males, is an inflammatory disease of the bile ducts, which may lead to cholestasis (blockage of bile transport to the gut). The blockage of bile ducts leads to the accumulation of bile acid in the liver and in bile, damages the liver and eventually causes liver failure. Most of the patients suffering from PSC show also a chronic inflammation of the colon (e.g. colitis ulcerosa). The inflammation of the bile duct can affect also the surrounding liver tissue and lead to a cicatrization of the small and large bile ducts, which will cause the constriction of the biliary tract. Consequently, such a constriction leads to a disturbed secretion of bile liquid further damaging the liver. In the course of the disease liver cirrhosis and cholangiocarcinoma may develop. Also PBC is an inflammatory disease of the bile ducts, which affects initially the smaller bile ducts and will finally result in liver cirrhosis. In contrast to PSC, PBC affects mainly female individuals and is not correlated to inflammatory diseases of the colon. The most efficient method to treat PBC and PSC is the transplantation of the liver. Until now the only promising pharmacological treatment of both diseases involves the use of UDCA. Currently, UDCA is the only drug approved for the treatment of cholestatic liver diseases (Paumgartner & Beuers 2002). UDCA is used in PBC at a dose of 12-15mg/kg/day (generally 1000-1500mg) administered orally once or twice a day. This use is approved by the U.S. Food and Drug Administration. Colchicine has been added to UDCA treatment. Colchicine is prescribed at a dose of 0.6mg twice a day because of its potential anti-inflammatory and anti-fibrotic effects. Several studies have demonstrated mild improvements in liver tests using colchicine. Nevertheless, none found a benefit for the liver histology or survival of PBC patients. Methotrexate, an immune suppressing agent, is another drug which has been tested in PBC. It is administered at a dose of 15mg per week. In small studies, methotrexate has improved symptoms, liver blood tests and progression of histology when used for several years. However, methotrexate causes severe side effects including bone marrow suppression, worsening of liver disease and potentially fatal pulmonary fibrosis.

[0005] UDCA is of limited efficacy in PSC and has not been shown to prolong survival (free of liver transplantation Trauner & Graziadei 1999, Paumgartner & Beuers 2002). Ongoing studies test whether high-dose UDCA may be more effective. Notably, UDCA reduces the risk of colon cancer in patients with PSC and ulcerative colitis. Based on the hypothesis that PSC has an immunologic cause, corticosteroids and other immunosuppressants have been tested. Oral corticosteroids yielded an initial improvement in the biochemical profile. However, lack of evidence for the long term benefit as well as bone demineralization, is an argument against the use of this regimen. Other medications such as azathioprine, cyclosporine, tested in association with corticosteroids and UDCA, have never been evaluated alone in the therapy of PSC. Methotrexate and D-penicillamine were also shown to be ineffective. Therefore, pharmaceutical therapy for PSC still needs to be optimized (Trauner & Graziadei 1999, Beuers & Paumgartner 2002).

[0006] Endoscopic treatment in PSC patients with symptomatic dominant strictures, gallstones or debris is considered to be a valuable option in addition to medical treatment. PSC patients undergoing endoscopic treatment had an increased survival, which was much higher than that predicted from survival models.

[0007] Orthotopic liver transplantation is an effective therapy for PSC and so far the only life-saving option for the end-stage disease. Following transplantation, however, PSC tends to recur in 15-30% of patients, and there is also a high recurrence rate of biliary strictures, chronic rejection, and reflux cholangitis. Unfortunately, use of immunosuppressants does not improve survival and recurrence of the disease. Thus, an urgent need exists for effective drug treatment preventing disease progression of PSC as well as recurrence after liver transplantation (Trauner & Graziadei 1999, Beuers & Paumgartner 2002).

[0008] Although UDCA, which is well tolerated with the exception of rare episodes of diarrhea and prurigo (Trauner & Graziadei 1999, Beuers & Paumgartner 2002), is predominantly used to treat cholestatic liver diseases, the efficacy of UDCA in PSC and in patients with liver diseases like progressive familial intrahepatic cholestasis type 3 is very limited (Trauner & Graziadei 1999, Jacquemin, Hermans, et al. 1997, Jacquemin 2000, Ismail, Kalicinski, et al. 1999).

[0009] In the EP 0 652 773 B1 the use of nor- and homo-bile acids derivatives, optionally conjugated with taurine, glycine or alanine, as absorption enhancers for medicaments by the enteral or other non-parenteral routes is described.

These derivatives show lipophilic and detergent properties and are not metabolized by the intestinal bacterial flora.

[0010] The EP 0 624 595 B1 discloses dimeric nor-derivatives of bile acids for the use in a medicament, especially suited for the treatment of hyperlipidemia. The substances disclosed therein consist of two single bile acid derivatives, which are covalently linked to each other.

[0011] US 4,892,868 discloses 22-Methyl-nor-ursodeoxycholic acid and 23-Methyl-ursodeoxycholic acid to be used to treat disorders of the hepatobiliary function, with particular reference to cholesterol metabolism and bile production (e.g. for the treatment of cholestasis).

[0012] It is an object of the present invention to provide alternative pharmaceuticals for the treatment of chronic liver diseases, selected from hepatocellular carcinoma and cholangiocarcinoma, which are more effective than known drugs like ursodeoxycholic acid and show less side effects than methotrexate.

[0013] Therefore, the present invention relates to the use of 24-nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof for use in the treatment and/or the prevention of hepatocellular carcinoma and cholangiocarcinoma.

[0014] It surprisingly turned out that a medicament or pharmaceutical composition comprising 24-nor-ursodeoxycholic acid, a side chain-shortened C₂₃ analogue of the naturally occurring C₂₄ bile acid ursodeoxycholic acid (UDCA), and/or pharmaceutical acceptable salts thereof can be successfully employed for the treatment of various liver diseases, because those analogues influence the physiological properties of bile acids (Hofmann 1999, Schmassmann, Hofmann, et al. 1990, Yoon, Hagey, et al. 1986, Cohen, Hofmann, et al. 1986). Although both substances are structurally very similar both substances show different characteristics when administered to mammals (see e.g. Yoon YB, Hagey LR, et al., 1986).

[0015] 24-nor-ursodeoxycholic acid and salts thereof lead to the induction of biliary bicarbonate secretion, which dilutes the toxic biliary content and protects bile duct epithelial cells against oxidative stress since bicarbonate is a potent scavenger for reactive oxygen species. This leads to the reconstitution of the cholangiocyte barrier function and will stop ongoing pericholangitis and subsequent periductal fibrosis by minimizing bile duct epithelial cell injury from the bile duct lumen. Furthermore, 24-nor-ursodeoxycholic acid and salts thereof turned out to have also anti-inflammatory and anti-fibrotic effects.

[0016] In contrast to 24-nor-deoxycholic acid and salts thereof, ursodeoxycholic acid improves only periductal fibrosis of lobular bile ducts and increases serum alanine aminotransferase (ALT) levels and induces bile infarcts. However, ursodeoxycholic acid does not improve small duct disease. The divergent effects of both bile acids in regard to liver injury may be related to the differences in the level of bile secretion i.e. that 24-nor-ursodeoxycholic acid and salts thereof stimulate primarily ductular secretion whereas ursodeoxycholic acid stimulates canalicular bile secretion upstream of the affected ducts.

[0017] The administration of 24-nor-deoxycholic acid and salts thereof induces bile acid detoxification via hydroxylation, sulfation and glucuronidation resulting in better water soluble and therefore less toxic bile acid metabolites, which dilute toxic bile acids in ductular bile and induce a ductular bicarbonate-rich choleresis reducing oxidative stress.

[0018] The generation of a bicarbonate rich choleresis by 24-nor-ursodeoxycholic acid and salts thereof has also therapeutic implications in human cholangiopathies (e.g. PSC, PBC, chronic liver graft rejection, non-suppurative destructive cholangitis), as cholehepatic shunting results in a continuing flux of molecules across the biliary ductular epithelium which helps altered bile ducts to better handle toxic/oxidative stress. For example, sulindac, a NSAID that also undergoes cholehepatic shunting in humans, has been shown to improve liver enzymes in PBC patients with incomplete response to UDCA treatment.

[0019] Methods for the preparation of 24-nor-ursodeoxycholic acid and salts thereof are known to the person skilled in the art and can preferably be prepared by a method as described in Schteingart CD and Hofmann AF (J. Lip. Res. 29(1988):1387-1395).

[0020] Of course, the drug according to the present invention can be used in humans as well as in mammals (e.g. swine, horse, primates, cattle, cat, dog).

[0021] A drug comprising 24-nor-deoxycholic acid and salts thereof can be used for the prevention/chemoprevention of a liver carcinoma, selected from hepatocellular carcinoma and cholangiocarcinoma. Also chronic liver graft rejection and non-suppurative destructive cholangitis may be treated by said drug.

[0022] Especially diseases which are a result or result in a bile duct injury can be treated by a drug comprising 24-nor-ursodeoxycholic acid and salts thereof.

[0023] Non-steroidal anti-inflammatory drug (NSAID) like ibuprofen reduce portal and lobular inflammation in the liver (which will lead to periductal fibrosis and ductular proliferation) and the formation of hepatocellular carcinoma (HCC) (Pikarsky, Porat, et al. 2004). Since 24-nor-ursodeoxycholic acid and salts thereof exhibit also anti-inflammatory properties, 24-nor-ursodeoxycholic acid may be used alone or in combination with other anti-inflammatory drugs, like NSAIDs (e.g. ibuprofen, sulindac (Bolder, Trang, et al. 1999)).

[0024] According to a preferred embodiment of the present invention 24-nor-ursodeoxycholic acid and salts thereof can be formulated for oral or intravenous administration, wherein these formulations further comprise pharmaceutically acceptable carriers, adjuvants, excipients and/or vehicles.

[0025] Solid dosage forms for oral administration can include tablets, preferably effervescent or chewable tablets, capsules, pills, powders and granules. In such solid dosage forms, 24-nor-ursodeoxycholic acid can be admixed with regularly used substances like sucrose, mannitol, sorbitol, starch and starch derivatives, lactose, lubricating agents (e.g. magnesium stearate), disintegrants and buffering agents. Tablets and pills can also be prepared with enteric coatings in order to prevent that 24-nor-ursodeoxycholic acid is affected by the stomach acids and enzymes. As immediate release tablets, these compositions may further comprise microcrystalline cellulose and/or dicalcium phosphate.

[0026] Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions and syrups containing inert diluents commonly used in the art, such as water. These dosage forms may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents. When administered by nasal aerosol or inhalation, the compositions according to the present invention may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons and/or other solubilizing or dispersing agents.

[0027] Suppositories for rectal administration of 24-nor-ursodeoxycholic acid can be prepared by mixing the compounds or compositions with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at room temperature but liquid at rectal temperature, such that they will melt in the rectum and release 24-nor-ursodeoxycholic acid and optionally other active compounds present in said suppositories.

[0028] Injectable preparations, for example sterile injectable aqueous or oleaginous suspensions, can be formulated according to the known art using suitable dispersing agents, wetting agents and/or suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that can be used are water and isotonic sodium chloride solution. Sterile fixed oils are also conventionally used as a solvent or suspending medium.

[0029] According to the present invention the dosage forms comprising 24-nor-ursodeoxycholic acid can further include conventional excipients, preferably pharmaceutically acceptable organic or inorganic carrier substances which do not react with the active compound. Suitable pharmaceutically acceptable carriers include, for instance, water, salt solutions, alcohol, oils, preferably vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, surfactants, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose and polyvinylpyrrolidone. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, like lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances which do not deleteriously react with the active compounds. For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants.

[0030] Various delivery systems are known and can be used to administer 24-nor-ursodeoxycholic acid, including, for example, encapsulation in liposomes, emulsions, microparticles, microcapsules and microgranules (e.g. EP 1 317 925). The required dosage can be administered as a single unit or in a sustained release form.

[0031] The bioavailability of 24-nor-ursodeoxycholic acid can be enhanced by micronization of the formulations using conventional techniques such as grinding, milling and spray drying in the presence of suitable excipients or agents such as phospholipids or surfactants.

[0032] According to the invention 24-nor-ursodeoxycholic acid can be formulated in a pharmaceutically acceptable salt or ester form. Pharmaceutically acceptable salts of 24-nor-ursodeoxycholic acid include preferably metal salts, in particular alkali metal salts, or other pharmaceutically acceptable salts. Suitable pharmaceutically acceptable acid addition salts may be prepared from inorganic acids, like hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid, or organic acids, like aliphatic, cycloaliphatic, aromatic, heterocyclic, carboxylic and sulfonic classes of organic acids, such as, formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, ethanesulfonic, anthranilic, mandelic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, methanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, algenic, sulfanilic, stearic, p-hydroxybutyric, cyclohexylaminosulfonic, galactaric and galacturonic acid. Pharmaceutically acceptable base addition salts include metallic salts made from lithium, aluminum, calcium, magnesium, potassium, sodium and zinc or organic salts made from primary, secondary and tertiary amines and cyclic amines. All 24-nor-ursodeoxycholic acid salts can be prepared by methods known in the state of the art (e.g. by reacting 24-nor-ursodeoxycholic acid with the appropriate acid or base). Methods for the manufacture of a drug according to the present invention comprising 24-nor-ursodeoxycholic acid and formulated for the administration as outlined herein can be found, for instance, in the "Handbook of Pharmaceutical Manufacturing Formulations" (Sarfraz K Niazi, CRC Press LLC, 2004).

[0033] The drug comprises preferably an effective amount of 24-nor-ursodeoxycholic acid and a pharmaceutically acceptable carrier and/or excipient.

[0034] According to a preferred embodiment of the present invention the medicament comprises 10 to 8000 mg, preferably 25 to 5000 mg, more preferably 50 to 1500 mg, in particular 250-500 mg, of 24-nor-ursodeoxycholic acid.

[0035] On average 24-nor-ursodeoxycholic acid and/or pharmaceutical acceptable salts thereof may preferably be

administered to a patient in an amount of 25 mg to 5 g, preferably 100 mg to 2.5 g, in particular 800 mg to 1.5 g per day. However, 1 g of 24-nor-ursodeoxycholic acid and/or pharmaceutical acceptable salts thereof is most preferably administered to a patient. It is further noted that 24-nor-ursodeoxycholic acid and/or pharmaceutical acceptable salts thereof may be administered to an individual in 1-100 mg/kg/d, preferably 5-50 mg/kg/d, more preferably 10-25 mg/kg/d, in particular 12-15 mg/kg/d. Said amounts are administered at once or preferably in more than one dose (at least 2, 3, 4, 5 or 10 doses) per day. The drug or the pharmaceutical composition according to the present invention may be administered for more than one week, preferably more than four weeks, more preferably more than six months, most preferably more than one year, in particular lifelong.

[0036] 24-nor-ursodeoxycholic acid can be administered not only in combination with pharmaceutically acceptable carriers and in dosage forms as described herein, but, of course, also in combination with one or more additional active ingredients (e.g. ursodeoxycholic acid, NSAID, like sulindac and ibuprofen) which are also known to be effective against the same or a similar disease to be treated (e.g. ursodeoxycholic acid) or against another disease, which may be preferably a result of a liver disease.

[0037] The present invention is further illustrated by the following figures and example.

Fig. 1. *norUDCA* cures sclerosing cholangitis in *Mdr2*^{-/-} mice. (A) Liver histology (H&E staining) in control diet-fed *Mdr2*^{-/-} mice (KO), UDCA-fed *Mdr2*^{-/-} mice (KO+UDCA), and *norUDCA*-fed *Mdr2*^{-/-} mice (KO+*norUDCA*) (Magnification x 10). Pronounced large bile duct disease in KO (arrow heads) which is significantly reduced in KO+UDCA (arrow heads) and absent in KO+*norUDCA*. (B) Sclerosing cholangitis in KO with periductal fibrosis, altered bile duct epithelial cells and mixed inflammatory infiltrate. This features are ameliorated in KO+ UDCA and absent in KO+*norUDCA* (Magnification x 40). (C) Sirius red staining showing significant fibrosis with periductal collagen fibers (red) in KO. Moderate reduction of fibrosis in KO+UDCA and even more pronounced reduction in KO+*norUDCA* (Magnification for b, c x 40); bd, bile duct.

Fig. 2. *norUDCA* significantly reduces hepatic hydroxyproline content in *Mdr2*^{-/-}. Hepatic hydroxyproline content in wild type (WT), control diet-fed *Mdr2*^{-/-} mice (KO), UDCA-fed *Mdr2*^{-/-} mice (KO+UDCA), and *NorUDCA*-fed *Mdr2*^{-/-} mice (KO+*norUDCA*). Hepatic hydroxyproline content is significantly increased in KO compared to WT and reduced to baseline levels in KO+*norUDCA*. Values are mean \pm SEM from n=5 per group. p<0.05, WT vs. KO; KO vs. KO+*norUDCA*.

Fig. 3. *norUDCA* significantly reduces portal neutrophils infiltration/extravasation in *Mdr2*^{-/-}. Immunohistochemistry for CD11b (red, staining neutrophils) in control diet-fed *Mdr2*^{-/-} mouse (KO), UDCA-fed *Mdr2*^{-/-} mouse (KO+UDCA), and *norUDCA*-fed *Mdr2*^{-/-} mouse (KO+*norUDCA*) of (A) interlobular and (B) lobular bile ducts. (C) Quantification of CD11b-positive cells per 20 portal fields. *norUDCA* significantly reduces the number of CD11b-positive cells in *Mdr2*^{-/-}. Values are mean \pm SEM from n=3 per group. p<0.05,

Fig. 4. *norUDCA* inhibits portal vascular cell adhesion molecule (VCAM) expression in *Mdr2*^{-/-} mice. Immunohistochemistry for VCAM (red) in interlobular (A) and lobular (B) bile ducts of control diet-fed *Mdr2*^{-/-} mouse (KO), UDCA-fed *Mdr2*^{-/-} mouse (KO+UDCA), and *norUDCA*-fed *Mdr2*^{-/-} mouse (KO+*norUDCA*). (A) No obvious difference in portal VCAM expression between KO and KO+UDCA whereas KO+*norUDCA* show significantly lower expression in bile duct proliferates. (B) At the level of lobular bile ducts both UDCA and *norUDCA* reduce cholangiocellular VCAM expression.

Fig. 5. *norUDCA* inhibits proliferation of hepatocytes and cholangiocytes in *Mdr2*^{-/-} mice. Immunohistochemistry for Ki-67 (red) in hepatocytes (A) and cholangiocytes (B) in control diet-fed *Mdr2*^{-/-} mouse (KO), UDCA-fed *Mdr2*^{-/-} mouse (KO+UDCA), and *norUDCA*-fed *Mdr2*^{-/-} mouse (KO+*norUDCA*). (A) Numerous Ki-67 positive hepatocytes in KO and KO+UDCA and scattered positive nuclei in KO+*norUDCA*. (B) Numerous Ki-67 positive cholangiocytes in KO, few positive cholangiocytes (arrow heads) in KO+UDCA and KO+*norUDCA*, respectively. (C) Number of Ki-67 positive hepatocytes per 30 HPFs and (D) number of Ki-67 positive cholangiocytes per 20 portal fields. Only *norUDCA* significantly reduces the number proliferating hepatocytes and cholangiocytes. Values are mean \pm SEM from n=3 per group. p<0.05, *WT vs. KO; t WT vs. KO+UDCA; ‡KO vs. KO+*norUDCA*.

Fig. 6. Positive correlation between biliary bicarbonate output and bile flow in *Mdr2*^{-/-}. Biliary bicarbonate output was plotted against bile flow in control diet-fed *Mdr2*^{-/-} mice (Control, open circles), UDCA-fed *Mdr2*^{-/-} mice (UDCA, open triangles), and *NorUDCA*-fed *Mdr2*^{-/-} mice (*norUDCA*, closed circles). Note the positive correlation between bicarbonate output and bile flow as well as the clustering of *NorUDCA*-treated animals showing the highest bicarbonate output in the upper right.

Fig. 7. Suggested therapeutic mechanisms of *norUDCA* in *Mdr2*^{-/-} mice. *norUDCA* is taken up by hepatocytes and secreted into canaliculi and bile ducts where it is taken up by cholangiocytes leading to ductular bicarbonate secretion. *norUDCA* is secreted back into the peribiliary plexus and shunted back to the hepatocytes (cholehepatic shunting). *NorUDCA* induces expression of Sult2a1 etc. and Mrp3 and Mrp4 which detoxifies bile salts and makes them amenable for renal elimination.

Fig. 8. Chemical structure of *norUDCA* (3- α ,7- α -dihydroxy-24-nor-5- β -cholan-23-oic acid).

EXAMPLE 1:

[0038] Mice with targeted disruption of the *Mdr2* (*Abcb4*) gene encoding a canalicular phospholipid flippase develop sclerosing cholangitis with macroscopic and microscopic features closely resembling those seen in human sclerosing cholangitis (e.g. primary sclerosing cholangitis, PSC) (Fickert, Zollner, et al. 2002, Fickert, Fuchsbichler, et al. 2004). Bile duct injury in these mice is linked to defective biliary phospholipids secretion resulting in an increased concentration of free non-micellar bile acids which subsequently cause bile duct epithelial cell (cholangiocyte) injury, pericholangitis, periductal fibrosis with ductular proliferation and finally sclerosing cholangitis (Fickert, Fuchsbichler, et al. 2004, Lammert, Wang, et al. 2004). In addition to the opportunity to study novel treatment strategies for PSC this model may be relevant for testing therapies for the wide spectrum of human liver diseases resulting from *MDR3* mutations (the human orthologue of *Mdr2*) ranging from neonatal cholestasis to adult liver disease (Jansen & Sturm 2003).

[0039] Currently, UDCA is the only drug approved for the treatment of cholestatic liver diseases (Paumgartner & Beuers 2002). However the efficacy of UDCA in PSC and in patients with liver diseases due to *MDR3* mutations (e.g. progressive familial intrahepatic cholestasis type 3) is limited (Trauner & Graziadei 1999, Jacquemin, Hermans, et al. 1997, Jacquemin 2000, Ismail, Kalicinski, et al. 1999). Side-chain shortening of UDCA could increase its therapeutic efficacy since this modification significantly influences the physiological properties of bile acids (Hofmann 1999, Schmassmann, Hofmann, et al. 1990, Yoon, Hagey, et al. 1986, Cohen, Hofmann, et al. 1986). *nor*UDCA, a side chain-shortened C₂₃ analogue of UDCA, is a potent choleric agent in different rodents (e.g. hamster, rat, guinea pig) undergoing extensive cholehepatic shunting and inducing biliary bicarbonate secretion at the bile duct level (Yoon, Hagey, et al. 1986, Cohen, Hofmann, et al. 1986). In contrast to UDCA, the effects of *nor*UDCA have never been studied in cholestasis. To test the hypothesis that cholehepatic shunting of a non-toxic bile acid may be beneficial in the treatment of cholangiopathies investigated the effects of *nor*UDCA in *Mdr2*^{-/-} mice as a model of sclerosing cholangitis (Fickert, Fuchsbichler, et al. 2004). In this example the positive effects of *nor*UDCA in the treatment for human liver diseases caused by *MDR3* mutations and human cholangiopathies such as sclerosing cholangitis (e.g. PSC) are examined.

1.1. Materials and Methods**1.1.1. Animal Experiments.**

[0040] *Mdr2*^{-/-} mice (FVB/N background) were obtained from Jackson Laboratory (The Jackson Laboratory, Bar Harbor, ME, USA). Mice were housed with a 12:12-hour light-dark cycle and permitted ad libitum consumption of water and a standard mouse diet (Sniff, Soest, Germany).

1.1.2. Bile acid feeding in *Mdr2*^{-/-} mice.

[0041] Two months-old *Mdr2*^{-/-} (a time point when sclerosing cholangitis is already fully established in these animals (Fickert, Zollner, et al. 2002) were either fed a diet supplemented with *nor*UDCA (0.5%, w/w) or UDCA as a clinical comparator (0.5%, w/w) for 4 weeks and compared to standard-diet-fed *Mdr2*^{-/-} and wild type controls.

1.1.3. Liver histology.

[0042] For conventional light microscopy, livers were fixed in 4% neutral buffered formaldehyde solution and embedded in paraffin. Sections (4 μm thick) were stained with H&E and Sirius red stain, respectively. The sections were coded and examined by a pathologist (H.D.) unaware of the animals' treatment.

1.1.4. Routine serum biochemistry.

[0043] Serum samples were stored at -70°C until analysis of alanine transaminase (ALT) and alkaline phosphatase (AP) by routine clinical chemistry performed on a Hitachi 717 analyzer (Boehringer Mannheim, Mannheim, Germany). For determination of total serum bile acid levels, a commercial 3-α-hydroxysteroid dehydrogenase assay (Ecoline S+; DiaSys, Holzheim, Germany) was used.

1.1.5. Immunohistochemistry for alpha-SMA.

[0044] Immunohistochemistry for alpha-SMA was performed on microwave treated (0.01 mmol/L citrate buffer pH 6.0) paraffin sections (4 μm thick) using the monoclonal mouse anti alpha-SMA (dilution 1:500, Sigma, St Louis, MO). Binding of the antibody was detected using the ABC system (Dako, Glostrup, Denmark) using β-amino-9-ethyl-carbazole (AEC; Dako) as substrate.

1.1.6. Immunohistochemistry for proliferation marker Ki-67.

[0045] Immunohistochemistry for Ki-67 was performed on microwave-treated (0.01 mmol/L citrate buffer pH 6.0) paraffin sections (4 µm thick) using a polyclonal rabbit anti-Ki-67 antibody (dilution 1:750, Novocastra, Newcastle upon Tyne, UK). Binding of the antibody was detected using the ABC system (Dako) using AEC (Dako) as substrate. Number of proliferating hepatocytes was calculated by counting positive nuclei in 30 high power fields in sections of 3 animals in each group. Number of proliferating bile duct epithelial cells was calculated by counting positive nuclei in 20 portal fields in sections of 3 animals in each group.

1.1.7. Immunohistochemistry for neutrophils marker CD-11b.

[0046] To quantify neutrophils CD 11b-positive cells were detected as described previously (Fickert, Fuchsbichler, et al. 2004) with the modification that binding of the antibody was detected using the ABC system (Dako) using AEC (Dako) as substrate. Number of neutrophils was calculated by counting positive cells in 20 portal fields in sections of 3 animals in each group.

1.1.8. Immunohistochemistry for vascular cell adhesion molecule (VCAM).

[0047] Immunohistochemistry for VCAM was performed on acetone fixed cryosections using the monoclonal rat anti CD106 (VCAM-1, dilution 1:30, PharMingen, San Diego, CA, USA) and binding of the antibody was detected using the ABC system (Dako) using AEC (Dako) as substrate.

1.1.9. Immunohistochemistry for 4-hydroxynonenal-protein adducts.

[0048] Liver sections were deparaffinized and then incubated with Immunopure peroxidase suppressor (Pierce, Rockford, IL) for 30 min and then protein block (DAKO, Carpinteria, CA) for 2 h. This was followed by overnight incubation with the primary anti-4-hydroxynonenal antibody (Calbiochem, San Diego, CA) at room temperature and binding of the antibody was detected using the ABC system (Dako) with AEC (Dako) as substrate.

1.1.10. Determination of hepatic hydroxyproline content.

[0049] To quantify liver fibrosis in our model hepatic hydroxyproline content was determined. The right liver lobe was homogenized in 6-normal HCl (200 mg liver tissue/4 ml HCl) and hydrolyzed at 110°C for 16 h. After filtration 50 µl were added to 450 µl 2.2% NaOH dissolved in citrate-acetate buffer (50 g citric acid x H₂O, 12 ml acetic acid, 120 g sodium acetate x 3 H₂O, 34 g NaOH ad 1 liter distilled water; pH 6.0). After adding 250 µl perchloric acid and 12 min incubation at room temperature 250 µl p-dimethylaminobenzaldehyde solution was added and incubated at 60°C for 20 min. Hydroxyproline content was measured at 565 nm using a hydroxyproline standard curve.

1.1.11. mRNA analysis and PCR of key fibrosis genes.

[0050] RNA isolation, cDNA synthesis and Taqman® real time PCR were performed as described previously (Wagner, Fickert, et al. 2003). The following primers and 5' FAM, 3' TAMRA labeled probes were used: Col1a1 fwd: caatgcaatgaagaactggactgt (Seq ID No. 1), Col1a1 rev: tctacatctctgagtttggtga (Seq ID No. 2) and Col1a1 probe: cagaagcacagcactcgccctcc (Seq ID No. 3); TIMP-1 fwd: cat-ggaaagcctctgtggtatg (Seq ID No. 4), TIMP-1 rev: aagctgcaggcat-tgatgtg (Seq ID No. 5) and TIMP-1 probe: ctcacacggcgccctaag-gaac (Seq ID No. 6); MMP-2 fwd: cttgagaaggatggcaagtatgg (Seq ID No. 7), MMP-2 rev: ttgtaggagtgccctggaa (Seq ID No. 8) and MMP-2 probe: cagatggacagccctgcaagttccc (Seq ID No. 9).

1.1.12. Bile flow measurement.

[0051] Bile flow was determined gravimetrically and normalized to liver weight as described previously (Fickert, Zollner, et al. 2001). Biliary phospholipids concentration was determined using a commercially available kit (Phospholipid B; Wako, Neuss, Germany) according to the manufacture's instructions. Biliary cholesterol concentration was determined using a commercial available kit (Cholesterol liquicolor; Human, Wiesbaden, Germany) according to the manufacture's instructions. Biliary glutathione (GSH) concentration was determined after protein precipitation in 5% metaphosphoric acid using the Glutathione Assay Kit (Calbiochem, San Diego, US) according to the manufacturer's instructions. Biliary bile acid concentration was analyzed using a 3-alpha-hydroxysteroid dehydrogenase assay (Ecoline St, DiaSys) according to the manufacturer's instructions.

1.1.13. Statistical analysis.

[0052] Data are reported as arithmetic means \pm SD. 4 to 6 animals were studied in each group. Statistical analysis was performed using Student's t-test when appropriate or ANOVA with Bonferroni post testing when three or more groups were compared. A p-value <0.05 was considered significant.

1.2. Results

1.2.1. *norUDCA significantly improves liver injury, reduces fibrosis, and cures sclerosing cholangitis in $Mdr2^{-/-}$ mice.*

[0053] Serum ALT and AP levels (as biochemical markers for liver injury and cholestasis) were significantly elevated in standard diet-fed $Mdr2^{-/-}$ compared to wild type controls (Table 1).

Table 1. Serum Biochemistry and Bile Acid Levels under various experimental conditions.

	ALT (U/L)	AP (U/L)	Bili
WT	71 \pm 19	92 \pm 13	0.09 \pm 0.03
KO	405 \pm 187	235 \pm 55	0.32 \pm 0.11
KO+UDCA	576 \pm 175*	399 \pm 73*	0.55 \pm 0.19
KO+Nor-UDCA	165 \pm 23*#	162 \pm 25*#	0.23 \pm 0.2#

NOTE. Values are expressed as mean \pm SD from n = 5 per group. ALT, alanine transaminase; AP, alkaline phosphatase; SBA, serum bile acids, WT, wild type mice; KO, *Mdr2* knock-out mice, KO+UDCA, UDCA-fed *Mdr2* knock-out mice; KO+*norUDCA*, *norUDCA*-fed *Mdr2* knock-out mice.

- *p <0.05 , KO vs. KO+UDCA and KO vs. KO+*norUDCA* (ANOVA with Bonferroni post-testing)
- #p <0.05 , KO+UDCA vs. KO+*norUDCA* (ANOVA with Bonferroni post-testing)

[0054] *norUDCA* significantly improved serum ALT and AP levels compared to standard diet-fed $Mdr2^{-/-}$ (Table 1). In parallel standard diet-fed $Mdr2^{-/-}$ displayed pronounced sclerosing cholangitis (Fig. 1) with ductular proliferation and liver fibrosis reflected by significantly elevated hepatic hydroxyproline content (Fig. 2). *norUDCA*-fed $Mdr2^{-/-}$ showed regular large- and medium-sized bile ducts with no or only modest periductal fibrosis (Fig. 1); ductular proliferation was virtually absent (not shown). In line with this histological changes, *norUDCA* significantly reduced hepatic hydroxyproline content in $Mdr2^{-/-}$ (Fig. 2) which was paralleled by a significant reduction in hepatic collagen 1 and 3 mRNA expression (Table 2).

Table 2. Real time PCR primer sequences (5'- 3')

	Forward	Reverse
Cyp2b10	CAATGGGAACGTTGGAAGA	TGATGCACTGGAAGAGGAAC
Cyp3a11	CCACCAGTAGCACACTTTCC	TTCCATCTCCATCACAGTATCA
IL-1 β	CTGGTGTGTGACGTTCCCATTA	CCGACAGCACGAGGCTTT
IL-1R	GCCAGGACCGCTCAGAGA	TGCCTCGACTGTTAGTCAAGCA
IL-6	GCCCACCAAGAACGATAGTCA	GAAGGCAACTGGATGGAAGTCT
MMP3	CCCACCAAGTCTAACTCTCTGGAA	GGGTGCTGACTGCATCAAAGA
MIP-2	CCTCAACGGAAGAACCAAAGAG	CTCAGACAGCGAGGCACATC
Mrp3	GGCAGGGCCCACTGAGT	AGTCCTCAGATGTCAGCCTAGTGA
Mrp4	TTAGATGGGCCTCTGTTCT	GCCCACAATTCCAATTCCAACCTT
i-NOS	ACATCAGGTCGGCCATCACT	CGTACCGGATGAGCTGTGAATT
Procollagen 1	GCAGGGTTCCAACGATGTTG	GCAGCCATCGACTAGGACAGA
Procollagen 3	GGTGGTTTTTCAGTTCAGCTATGG	CTGGAAAGAAGTCTGAGGAATGC

(continued)

	Forward	Reverse
TGF-beta	TCGACATGGAGCTGGTGAAA	CTGGCGAGCCTTAGTTTGGA
TNF-alpha	GACCCTCACACTCAGATCATCTTCT	CCTCCACTTGGTGGTTTGCT
TNF-R1	TGCACTAAACAGCAGAACCGAG	TTGCTCAGCCTCATGCACTG
Sult2a1	GGAAGGACCACGACTCATAAC	GATTCTTCACAAGGTTTGTGTTAC C
Ugt1a1	TCTGAGCCCTGCATCTATCTG	CCCCAGAGGCGTTGACATA

[0055] UDCA treatment in *Mdr2*^{-/-} as a clinical comparator and current standard treatment of cholestatic liver diseases (including PSC) were studied simultaneously. In contrast to *norUDCA*, UDCA significantly increased ALT and AP activity in *Mdr2*^{-/-} (Table 1) and 2 out of 5 animals showed bile infarcts in line with previous observations (Fickert, Zollner, et al. 2002). UDCA appeared to reduce periductal fibrosis of lobular and interlobular bile ducts (Fig. 1) but showed only a trend for lower hepatic hydroxyproline content without reaching statistical significance (Fig. 2). These findings clearly indicate that *norUDCA* (but not UDCA) significantly reduces liver injury, pericholangitis, and periductal fibrosis in *Mdr2*^{-/-} finally leading to healing of sclerosing cholangitis.

1.2.2. *norUDCA* reduces inflammation, oxidative stress and cell proliferation in *Mdr2*^{-/-} mice.

[0056] Since portal inflammation may represent the main trigger for periductal fibrosis in *Mdr2*^{-/-} (Pikarsky, Porat, et al. 2004) the reduction portal inflammation with *norUDCA* was analysed. Control diet-fed *Mdr2*^{-/-} had a significantly elevated number of portal neutrophils compared to wild type controls (Fig. 3). *norUDCA* also improved pericholangitis in *Mdr2*^{-/-} as reflected by a significantly reduced number of portal neutrophils compared to standard diet-fed *Mdr2*^{-/-} (Fig. 3). These apparent anti-inflammatory effects of *norUDCA* were paralleled by the significantly lower VCAM expression in the bile duct epithelial cells of interlobular and lobular bile ducts of *norUDCA*-fed *Mdr2*^{-/-} (Fig. 4). Since it was previously shown that inflammation triggers also proliferation in livers of *Mdr2*^{-/-} (Pikarsky, Porat, et al. 2004) it was tested whether this was affected by *norUDCA*. Standard diet-fed *Mdr2*^{-/-} showed a significantly elevated number of Ki-67 positive hepatocytes and cholangiocytes compared to wild type controls (Fig. 5). *norUDCA* significantly reduced the degree of hepatocyte and bile duct epithelial cell proliferation to nearly wild type control levels (Fig. 5). UDCA had no significant effects on portal inflammation (Fig. 3), VCAM expression (Fig. 4), and proliferation of hepatocytes (Fig. 5). However, UDCA reduced proliferation of bile duct epithelial cells in large bile ducts (Fig. 5). Again, these data indicate that the anti-inflammatory and anti-proliferative effects of *norUDCA* are superior to UDCA.

1.2.3. *norUDCA* induces biliary bicarbonate secretion in *Mdr2*^{-/-} mice.

[0057] To determine whether bicarbonate rich choleresis resulting from cholehepatic shunting of *norUDCA* could be responsible for the observed therapeutic effects, bile flow and composition were determined (Table 3).

Table 3. Bile flow and biliary excretion of bile acids, cholesterol, phospholipids and glutathione under various experimental conditions.

		KO	KO+UDCA	KO+ <i>norUDCA</i>
		n=4	n=6	n=5
Bile flow	μL/g /min	2.3±0.3	2.4±0.3	3.5±0.3
Bile acids	nmol/g/min	23.0±5.9	37.2±7.6*	29.5±2.3
Cholesterol	nmol/g/min	0.2±0.1	0.3±0.1	0.2±0.04
Phospholipids	nmol/g/min	0.4±0.1	0.3±0.03	0.3±0.1
Glutathione	nmol/g/min	4.6±2.0	5.6±1.1	5.8±1.2
Bicarbonate	nmol/g/min	60.9±8.0	67.1±10.0	98.8±14.6#

NOTE. Values are expressed as mean \pm SD. KO, *Mdr2* knock-out mice, KO+UDCA, UDCA-fed *Mdr2* knock-out mice; KO+*norUDCA*, *norUDCA*-fed *Mdr2* knock-out mice.

- * $p < 0.05$, KO vs. KO+UDCA (ANOVA with Bonferroni post-testing)
- # $p < 0.05$, KO vs. KO+*norUDCA* (ANOVA with Bonferroni post-testing)

[0058] Compared to control diet-fed and UDCA-fed *Mdr2*^{-/-} *norUDCA* significantly induced biliary bicarbonate secretion consistent with the concept that *norUDCA* undergoes relevant cholehepatic shunting in *Mdr2*^{-/-} (Bolder, Trang, et al. 1999). UDCA but not *norUDCA* stimulated biliary bile acid excretion. No significant effects on biliary bile acid, cholesterol, phospholipids and glutathione output were observed. These findings show that *norUDCA* leads to flushing of the injured bile ducts with a bicarbonate-enriched less toxic bile in *Mdr2*^{-/-}.

1.2.4. *norUDCA* induces phase II detoxification pathways and alternative excretory routes for bile acids in *Mdr2*^{-/-} mice.

[0059] To test the hypothesis that induction of phase I/II biotransformation pathways and alternative efflux routes for potentially toxic bile acids may also contribute to the observed beneficial effects of *norUDCA* in *Mdr2*^{-/-} the expression of key metabolic enzymes (Table 4) and serum, hepatic, and biliary bile acid composition in *norUDCA*-fed *Mdr2*^{-/-} was studied.

Table 4. Real-time PCR for quantification of relative expression levels of key metabolic enzymes and transport proteins under various experimental conditions.

	KO	KO+UDCA	KO+ <i>norUDCA</i>
Cyp2b10	100 \pm 103	751 \pm 245*	12941418*#
Cyp3a11	100 \pm 46	246 \pm 72*	241 \pm 45*
Sult2a1	n.d.	100 \pm 46	24157114948*#
Ugt1a1	100 \pm 54	137 \pm 55	304 \pm 81*#
Cyp7a1	100 \pm 78	9 \pm 4*	60 \pm 30#
Mrp3	100 \pm 33	194 \pm 55	207 \pm 73*
Mrp4	100 \pm 30	357 \pm 95	590 \pm 193*

[0060] NOTE. Values are expressed as mean \pm SD; n=5 in each group. n.d., not detectable, KO, *Mdr2* knock-out mice, KO+UDCA, UDCA-fed *Mdr2* knock-out mice; KO+*norUDCA*, *norUDCA*-fed *Mdr2* knock-out mice.

- * $p < 0.05$, KO vs. KO+UDCA and KO vs. KO+*norUDCA* (ANOVA with Bonferroni post-testing)
- # $p < 0.05$, KO+UDCA vs. KO+*norUDCA* (ANOVA with Bonferroni post-testing)

[0061] Specific attention was paid to Sult2a1 and Mrp4, since sulfation and transport of sulfated compounds are inter-related to form a coordinately regulated pathway for excretion of sulfated steroids and bile acids (Schuetz, Strom, et al. 2001). *norUDCA* had no significant effects on mRNA expression of hepatocellular uptake (Ntcp, Oatp1) and canalicular efflux (Bsep, Mrp2) systems for bile acids and organic anions (data not shown). However, *norUDCA* resulted in a robust induction of phase I and II detoxification enzymes (Table 4) with most pronounced effects on Sult2a1 expression. In addition, *norUDCA* profoundly increased the expression of alternative basolateral efflux systems such as Mrp4 and - to a lesser degree - Mrp3 (Table 4, Fig. 7). The effects of UDCA were less pronounced (Table 4, Fig. 7). This coordinated induction of biotransformation pathways and efflux systems by *norUDCA* was accompanied by the appearance of bile acid glucuronides and also sulfates reflecting the functional implications of the observed expression changes.

1.3. Discussion

[0062] It could be shown that *norUDCA* cures sclerosing cholangitis in *Mdr2*^{-/-} mice, a well characterized model system for PSC, within 4 weeks. In addition, it could be demonstrated that *norUDCA* is significantly more effective than UDCA.

[0063] The development of sclerosing cholangitis in *Mdr2*^{-/-} is directly related to defective biliary phospholipid secretion and concomitantly increased biliary levels of non-micellar-bound toxic bile acids causing bile duct injury and pericholangitis (Fickert, Zollner, et al. 2002, Fickert, Fuchsbichler, et al. 2004). Induction of biliary bicarbonate secretion in *norUDCA*-

fed *Mdr2*^{-/-} presented by this example is strongly consistent with cholehepatic shunting of norUDCA (Hofmann 1977, Yoon, Hagey, et al. 1986). Increased biliary bicarbonate secretion (i) dilutes the toxic biliary content in *Mdr2*^{-/-} and (ii) protects bile duct epithelial cells against oxidative stress since bicarbonate is a potent scavenger for reactive oxygen species. *norUDCA* stops therefore ongoing pericholangitis and subsequent periductal fibrosis in *Mdr2*^{-/-} by minimizing bile duct epithelial cell injury from the bile duct lumen. This leads to the reconstitution of the cholangiocyte barrier function which would mean that the observed anti-inflammatory and anti-fibrotic effects of *norUDCA* in *Mdr2*^{-/-} are secondary. However, it is evident that *norUDCA* has also direct anti-inflammatory and anti-fibrotic effects.

[0064] The findings of this example demonstrate that inhibition of periductal fibrosis, when accompanied with modulation of the biliary content (i.e. increasing the content of hydrophilic bile acids together with increased bicarbonate concentration within the duct), significantly improves liver injury in *Mdr2*^{-/-}.

[0065] In contrast to *norUDCA*, UDCA improved only periductal fibrosis of lobular bile ducts but increased serum ALT levels and induced bile infarcts in *Mdr2*^{-/-}. In a previous study with a shorter treatment period it was concluded that this may primarily be related to the choleric effects of UDCA in the presence of unresolved biliary obstruction comparable to findings in CBDL UDCA-fed mice (Fickert, Zollner, et al. 2002). In contrast to this previous assumption, it was found only a trend for increased bile flow in UDCA- and even more so in *norUDCA*-fed *Mdr2*^{-/-} in the current study using lower doses. However, since UDCA did not improve small duct disease in *Mdr2*^{-/-} this does not exclude the possibility of increased biliary pressure at the level of the canals of Herring in UDCA-fed *Mdr2*^{-/-} which could have led to the observed bile infarcts. The divergent effects of both bile acids in regard to liver injury are related to the differences in the level of bile secretion i.e. that *norUDCA* stimulates primarily ductular secretion whereas UDCA stimulates canalicular bile secretion upstream of the affected ducts.

[0066] A causal relationship between portal and lobular inflammation leading to periductal fibrosis and ductular proliferation as well as formation of hepatocellular carcinoma (HCC) in *Mdr2*^{-/-} has recently been demonstrated (Fickert, Fuchsichler, et al. 2004, Pikarsky, Porat, et al. 2004). In the current study *norUDCA* normalized hepatocyte and bile duct epithelial cell proliferation. Pikarsky et al. have demonstrated a reduction of inflammation and related HCC formation using the non-steroidal anti-inflammatory drug (NSAID) ibuprofen in this model (Pikarsky, Porat, et al. 2004). The therapeutic effects of NSAIDs and *norUDCA* may be combined and even amplified by sulindac, an NSAID, which undergoes cholehepatic shunting in rats (Bolder, Trang, et al. 1999).

[0067] Natural bile acids are efficiently N-acyl amidated (conjugated) in an amide linkage with glycine or taurine and then secreted into the bile canaliculus. In contrast, *nor* (C₂₃) bile acids have marked difference in their biotransformation and physiological properties when compared to their natural (C₂₄) homologues. It was shown that *norUDCA* results in a coordinated and robust induction of Sult2a1 (a transferase preferentially sulfating steroids and bile acids) and Mrp4 (a transporter of sulfated steroids and bile acids) (Schuetz, Strom, et al. 2001, Zelcer, Reid, et al. 2003). The functional implications of these findings are supported by the appearance of bile acid sulfates and glucuronides in urine of *norUDCA* fed *Mdr2*^{-/-}. Adaptive induction of bile acid detoxification by *norUDCA* via phase I (hydroxylation) and II (sulfation, glucuronidation) metabolism may result in better water soluble and therefore less toxic bile acid metabolites being eliminated by alternative hepatocellular efflux pumps (e.g. Mrp4) followed by their renal excretion as demonstrated in this example. The induction of such mechanisms by *norUDCA* was much more pronounced than that of UDCA (in this example) or agonistic ligands of CAR reported previously (Assem et al. 2004). This shows that *norUDCA* profoundly induces Sult2a1-mediated bile acid detoxification and export by adaptive overexpression of Mrp4 while *norUDCA* itself undergoes continued cholehepatic shunting. This has a dual beneficial effect by (i) displacing and diluting toxic bile acids in ductular bile and (ii) inducing a ductular bicarbonate-rich choleresis which reduces oxidative stress.

[0068] The generation of a bicarbonate rich choleresis by *norUDCA* has also therapeutic implications in human cholangiopathies (e.g. PSC, PBC, chronic liver graft rejection, non-suppurative destructive cholangitis), as cholehepatic shunting results in a continuing flux of molecules across the biliary ductular epithelium which helps altered bile ducts to better handle toxic/oxidative stress. For example, sulindac, a NSAID that also undergoes cholehepatic shunting in humans, has been shown to improve liver enzymes in PBC patients with incomplete response to UDCA treatment.

[0069] Of interest two parallels in the *norUDCA* metabolism between mice and humans exist which contrast previous observations in other rodents and experimental animals (e.g. biliary fistular rats, hamster, guinea pig). First both, mice and also men, show considerable renal excretion of *norUDCA*. In addition, it was also found that the major metabolite of *norUDCA* in mice was a glucuronide which is also in line with the findings in humans.

[0070] Second, in contrast to rats (Yoon, Hagey, et al. 1986) the estimated choleric potency of *norUDCA* in mice and humans lays about three times above that of normal bile flow in both species. However, *norUDCA* induced bile flow to a much higher degree in rats (160ml/min-kg). Nevertheless taken together these findings it could be shown that the effects of *norUDCA* in *Mdr2*^{-/-} can be directly extrapolated to human cholestatic liver diseases.

[0071] In summary it could be shown that *norUDCA* cures sclerosing cholangitis in *Mdr2*^{-/-}. *norUDCA* is an effective compound for cholestatic liver diseases particularly for human PSC and liver diseases related to *MDR3* mutations.

[0072] References:

- Bolder et al. Gastroenterology 1999; 117(4):962-971.
 Cohen et al. Gastroenterology 1986; 91(1):189-197.
 Fickert et al. Gastroenterology 2001; 121(1):170-183.
 Fickert et al. Gastroenterology 2002; 123(4):1238-1251.
 5 Fickert et al. Gastroenterology 2004; 127(1):261-274.
 Hofmann et al. Pediatr Transplant 1999; 3(3):219-224.
 Jacquemin et al. Hepatology 1997; 25(3):519-523.
 Jacquemin E. Clin Liver Dis 2000; 4(4):753-763.
 Jansen et al. Liver Int 2003; 23(5):315-322.
 10 Lammert et al. Hepatology 2004; 39(1):117-128.
 Paumgartner et al. Hepatology 2002; 36(3):525-531.
 Pikarsky et al. Nature 2004; 431(7007):461-466.
 Schmassmann et al. Hepatology 1990; 11(6):989-996.
 Trauner M et al. Aliment Pharmacol Ther 1999; 13(8):979-996.
 15 Wagner et al. Gastroenterology 2003; 125(3):825-838.
 Yoon et al. Gastroenterology 1986; 90(4):837-852.

SEQUENCE LISTING

20 [0073]

<110> Medizinische Universität Graz

<120> Use of 24-norUDCA

<130> R 59694

<140> UNKNOWN

<141> 2005-05-12

<160> 34

<170> PatentIn version 3.4

<210> 1

<211> 19

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 1

caatgggaac gttggaaga 19

<210> 2

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 2

tgatgcactg gaagaggaac 20

<210> 3

<211> 20

	<212> DNA	
	<213> Artificial	
	<220>	
5	<223> Primer	
	<400> 3	
	ccaccagtag cacacttcc	20
10	<210> 4	
	<211> 22	
	<212> DNA	
	<213> Artificial	
15	<220>	
	<223> Primer	
	<400> 4	
20	ttccatctcc atcacagtat ca	22
	<210> 5	
	<211> 22	
	<212> DNA	
	<213> Artificial	
25	<220>	
	<223> Primer	
	<400> 5	
30	ctggtgtgtg acgttcccat ta	22
	<210> 6	
	<211> 18	
	<212> DNA	
35	<213> Artificial	
	<220>	
	<223> Primer	
40	<400> 6	
	ccgacagcac gaggcctt	18
	<210> 7	
	<211> 18	
45	<212> DNA	
	<213> Artificial	
	<220>	
	<223> Primer	
50	<400> 7	
	gccaggaccg ctgagaga	18
	<210> 8	
55	<211> 22	
	<212> DNA	
	<213> Artificial	

<220>
 <223> Primer

 <400> 8
 5 tgcctcgact gttagtcaag ca 22

 <210> 9
 <211> 21
 <212> DNA
 10 <213> Artificial

 <220>
 <223> Primer

 15 <400> 9
 gccaccaag aacgatagtc a 21

 <210> 10
 <211> 22
 20 <212> DNA
 <213> Artificial

 <220>
 *<223> Primer
 25

 <400> 10
 gaaggcaact ggatggaagt ct 22

 <210> 11
 30 <211> 24
 <212> DNA
 <213> Artificial

 <220>
 35 <223> Primer

 <400> 11
 cccaccaagt ctaactctct ggaa 24

 40 <210> 12
 <211> 21
 <212> DNA
 <213> Artificial

 45 <220>
 <223> Primer

 <400> 12
 50 gggtgctgac tgcataaag a 21

 <210> 13
 <211> 22
 <212> DNA
 55 <213> Artificial

 <220>
 <223> Primer

EP 2 392 337 B1

	<400> 13	
	cctcaacgga agaaccaaag ag	22
5	<210> 14	
	<211> 20	
	<212> DNA	
	<213> Artificial	
10	<220>	
	<223> Primer	
	<400> 14	
	ctcagacagc gaggcacatc	20
15	<210> 15	
	<211> 18	
	<212> DNA	
	<213> Artificial	
20	<220>	
	<223> Primer	
	<400> 15	
25	ggcagggcca cactgagt	18
	<210> 16	
	<211> 24	
	<212> DNA	
	<213> Artificial	
30	<220>	
	<223> Primer	
	<400> 16	
35	agtcctcaga tgtcagccta gtga	24
	<210> 17	
	<211> 20	
	<212> DNA	
40	<213> Artificial	
	<220>	
	<223> Primer	
45	<400> 17	
	ttagatgggc ctctggttct	20
	<210> 18	
	<211> 24	
50	<212> DNA	
	<213> Artificial	
	<220>	
	<223> Primer	
55	<400> 18	
	gccacaatt ccaattccaa cctt	24

<210> 19
 <211> 20
 <212> DNA
 <213> Artificial
 5
 <220>
 <223> Primer
 <400> 19
 10 acatcaggtc ggccatcact 20
 <210> 20
 <211> 22
 <212> DNA
 15 <213> Artificial
 <220>
 <223> Primer
 20 <400> 20
 cgtaccggat gagctgtgaa tt 22
 <210> 21
 <211> 20
 25 <212> DNA
 <213> Artificial
 <220>
 <223> Primer
 30 <400> 21
 gcagggttcc aacgatgttg 20
 <210> 22
 35 <211> 21
 <212> DNA
 <213> Artificial
 <220>
 40 <223> Primer
 <400> 22
 gcagccatcg actaggacag a 21
 45 <210> 23
 <211> 23
 <212> DNA
 <213> Artificial
 50 <220>
 <223> Primer
 <400> 23
 55 ggtggttttc agttcagcta tgg 23
 <210> 24
 <211> 23
 <212> DNA

<213> Artificial
 <220>
 <223> Primer
 5
 <400> 24
 ctggaaagaa gtctgaggaa tgc 23
 <210> 25
 10 <211> 20
 <212> DNA
 <213> Artificial
 <220>
 15 <223> Primer
 <400> 25
 tcgacatgga gctggtgaaa 20
 20 <210> 26
 <211> 20
 <212> DNA
 <213> Artificial
 25 <220>
 <223> Primer
 <400> 26
 30 ctggcgagcc ttagtttgga 20
 <210> 27
 <211> 25
 <212> DNA
 <213> Artificial
 35 <220>
 <223> Primer
 <400> 27
 40 gaccctcaca ctcagatcat cttct 25
 <210> 28
 <211> 20
 <212> DNA
 45 <213> Artificial
 <220>
 <223> Primer
 50 <400> 28
 cctccacttg gtggttgct 20
 <210> 29
 <211> 22
 55 <212> DNA
 <213> Artificial
 <220>

<223> Primer

<400> 29
5 tgactaaac agcagaaccg ag 22

<210> 30
<211> 20
<212> DNA
<213> Artificial

10

<220>
<223> Primer

<400> 30
15 ttgctcagcc tcatgcactg 20

<210> 31
<211> 21
<212> DNA
20 <213> Artificial

<220>
<223> Primer

25 <400> 31
ggaaggacca cgactcataa c 21

<210> 32
<211> 25
30 <212> DNA
<213> Artificial

<220>
<223> Primer

35 <400> 32
gattcttcac aaggttgtg ttacc 25

<210> 33
40 <211> 21
<212> DNA
<213> Artificial

<220>
45 <223> Primer

<400> 33
tctgagccct gcacttatct g 21

50 <210> 34
<211> 19
<212> DNA
<213> Artificial

55 <220>
<223> Primer

<400> 34

ccccagaggc gttgacata

19

Claims

1. Nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof for use in the treatment and/or the prevention of hepatocellular carcinoma and cholangiocarcinoma.
2. Nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof for use in the treatment and/or the prevention of hepatocellular carcinoma and cholangiocarcinoma according to claim 1, **characterized in that** nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof is/are formulated for oral, parenteral, subcutaneous, intravenous, intramuscular, nasal, topical or rectal administration.
3. Nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof for use in the treatment and/or the prevention of hepatocellular carcinoma and cholangiocarcinoma according to claims 1 or 2, **characterized in that** nor-ursodeoxycholic acid and/or pharmaceutically acceptable salts thereof is/are to be administered to a patient in an amount of 25 mg to 5 g, preferably 100 mg to 2,5 g, in particular 800 mg to 1,5 g per day.

Patentansprüche

1. Nor-Ursodeoxycholsäure und/oder pharmazeutisch akzeptable Salze davon, zur Anwendung bei der Behandlung und/oder der Prävention von hepatozellulärem Karzinom und Cholangiokarzinom.
2. Nor-Ursodeoxycholsäure und/oder pharmazeutisch akzeptable Salze davon, zur Anwendung bei der Behandlung und/oder der Prävention von hepatozellulärem Karzinom und Cholangiokarzinom gemäß Anspruch 1, **dadurch gekennzeichnet, dass** Nor-Ursodeoxycholsäure und/oder pharmazeutisch akzeptable Salze davon für die orale, parenterale, subkutane, intravenöse, intramuskuläre, nasale, topische oder rektale Verabreichung formuliert ist/sind.
3. Nor-Ursodeoxycholsäure und/oder pharmazeutisch akzeptable Salze davon, zur Anwendung bei der Behandlung und/oder der Prävention von hepatozellulärem Karzinom und Cholangiokarzinom gemäß Ansprüchen 1 oder 2, **dadurch gekennzeichnet, dass** Nor-Ursodeoxycholsäure und/oder pharmazeutisch akzeptable Salze davon an einen Patienten in einer Menge von 25 mg bis 5 g, vorzugsweise 100 mg bis 2,5 g, insbesondere 800 mg bis 1,5 g pro Tag zu verabreichen ist/sind.

Revendications

1. Acide nor-ursodésoxycholique et/ou sels de celui-ci pharmaceutiquement acceptables destiné(s) à être utilisé(s) dans le traitement et/ou la prévention du carcinome hépatocellulaire et du cholangiocarcinome.
2. Acide nor-ursodésoxycholique et/ou sels de celui-ci pharmaceutiquement acceptables destiné(s) à être utilisé(s) dans le traitement et/ou la prévention du carcinome hépatocellulaire et du cholangiocarcinome selon la revendication 1, caractérisé(s) en ce que l'acide nor-ursodésoxycholique et/ou sels de celui-ci pharmaceutiquement acceptables est/sont formulé(s) pour une administration orale, parentérale, sous-cutanée, intraveineuse, intramusculaire, nasale, topique ou rectale.
3. Acide nor-ursodésoxycholique et/ou sels de celui-ci pharmaceutiquement acceptables destiné(s) à être utilisé(s) dans le traitement et/ou la prévention du carcinome hépatocellulaire et du cholangiocarcinome selon les revendications 1 ou 2, caractérisé(s) en ce que l'acide nor-ursodésoxycholique et/ou sels de celui-ci pharmaceutiquement acceptables doit/doivent être administré(s) à un patient en une quantité de 25 mg à 5 g, de préférence 100 mg à 2,5 g, en particulier 800 mg à 1,5 g par jour.

Fig. 1




Fig. 2




Fig. 6




Fig. 3




Fig. 4




Fig. 7




Fig. 5





Fig. 8

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0652773 B1 [0009]
- EP 0624595 B1 [0010]
- US 4892868 A [0011]
- EP 1317925 A [0030]

Non-patent literature cited in the description

- **SCHTEINGART CD ; HOFMANN AF.** *J. Lip. Res.*, 1988, vol. 29, 1387-1395 [0019]
- **SARFARAZ K NIAZI.** Handbook of Pharmaceutical Manufacturing Formulations. CRC Press LLC, 2004 [0032]
- **BOLDER et al.** *Gastroenterology*, 1999, vol. 117 (4), 962-971 [0072]
- **COHEN et al.** *Gastroenterology*, 1986, vol. 91 (1), 189-197 [0072]
- **FICKERT et al.** *Gastroenterology*, 2001, vol. 121 (1), 170-183 [0072]
- **FICKERT et al.** *Gastroenterology*, 2002, vol. 123 (4), 1238-1251 [0072]
- **FICKERT et al.** *Gastroenterology*, 2004, vol. 127 (1), 261-274 [0072]
- **HOFMANN et al.** *Pediatr Transplant*, 1999, vol. 3 (3), 219-224 [0072]
- **JACQUEMIN et al.** *Hepatology*, 1997, vol. 25 (3), 519-523 [0072]
- **JACQUEMIN E.** *Clin Liver Dis*, 2000, vol. 4 (4), 753-763 [0072]
- **JANSEN et al.** *Liver Int*, 2003, vol. 23 (5), 315-322 [0072]
- **LAMMERT et al.** *Hepatology*, 2004, vol. 39 (1), 117-128 [0072]
- **PAUMGARTNER et al.** *Hepatology*, 2002, vol. 36 (3), 525-531 [0072]
- **PIKARSKY et al.** *Nature*, 2004, vol. 431 (7007), 461-466 [0072]
- **SCHMASSMANN et al.** *Hepatology*, 1990, vol. 11 (6), 989-996 [0072]
- **TRAUNER M et al.** *Aliment Pharmacol Ther*, 1999, vol. 13 (8), 979-996 [0072]
- **WAGNER et al.** *Gastroenterology*, 2003, vol. 125 (3), 825-838 [0072]
- **YOON et al.** *Gastroenterology*, 1986, vol. 90 (4), 837-852 [0072]

Szabadalmi igénypontok

1. Nor-ursodeoxikolsav és/vagy gyógyszerészetileg elfogadhatóí sói hepatocelluláris karcinóma és kolangiokarcinóma kezelésében és/vagy megelőzésében történő alkalmazásra.

2. Nor-ursodeoxikolsav és/vagy gyógyszerészetileg elfogadhatóí sói hepatocelluláris karcinóma és kolangiokarcinóma kezelésében és/vagy megelőzésében történő 1. igénypont szerinti alkalmazásra, **azzal jellemezve, hogy** a nor-ursodeoxikolsav és/vagy gyógyszerészetileg elfogadhatóí sói orális, parenterális, szubkután, intravénás, intramuszkuláris, nazális, topikális vagy rektális adagoláshoz van(nak) formulálva.

3. Nor-ursodeoxikolsav és/vagy gyógyszerészetileg elfogadhatóí sói hepatocelluláris karcinóma és kolangiokarcinóma kezelésében és/vagy megelőzésében történő 1. vagy 2. igénypont szerinti alkalmazásra, **azzal jellemezve, hogy** a nor-ursodeoxikolsavat és/vagy gyógyszerészetileg elfogadhatóí sóit a betegnek naponta 25 mg-tól 5 g-ig terjedő mennyiségben, előnyösen 100 mg-tól 2,5 g-ig terjedő mennyiségben, különösen 800 mg-tól 1,5 g-ig terjedő mennyiségben adagoljuk.

A meghatalmazott:

Dr. Gál Mária
Szabadalmi Ügyvivő
SBBK Szabadalmi Ügyvivői Iroda
H-1062 Budapest, Andrásy ut. 113.
Telefon: 461-1000 Fax: 461-1000
Email: gal@sbgk.hu