

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2007334333 B2

**(54) Title
Activin-ActRII antagonists and uses for increasing red blood cell levels**

(51) International Patent Classification(s)
A61K 38/17 (2006.01) C12N 15/62 (2006.01)
A61P 7/06 (2006.01)

(21) Application No: 2007334333 (22) Date of Filing: 2007.12.18

(87) WIPO No: WO08/076437

(30) Priority Data

(31) Number 60/875,682 (32) Date 2006.12.18 (33) Country US

(43) Publication Date: 2008.06.26
(44) Accepted Journal Date: 2013.05.30

**(71) Applicant(s)
Acceleron Pharma Inc.**

**(72) Inventor(s)
Sherman, Matthew L.**

**(74) Agent / Attorney
Shelston IP, L 21 60 Margaret St, Sydney, NSW, 2000**

(56) Related Art
Harrisom, C. et al. TRENDS in Endocrinology and Metabolism Vol.16 No.2 March 2005, pp. 73-78
WO 2005/009460 A2
Welt, C. et al. European Journal of Endocrinology, 1998, vol.139, pp. 469-471
Coerver, K. et al. Molecular Endocrinology, vol. 10, no. 5, 1996, pp. 534-543
AU2006318449 A
WO 2006/012627 A2
WO 2004/039948 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 June 2008 (26.06.2008)

PCT

(10) International Publication Number
WO 2008/076437 A3

(51) International Patent Classification:
A61K 38/17 (2006.01) A61P 7/06 (2006.01)
C12N 15/62 (2006.01)

Matthew, L. [US/US]; 33 Janet Road, Newton, MA 02459 (US).

(21) International Application Number:
PCT/US2007/025868

(74) Agents: VINCENT, Matthew, P. et al.; Ropes & Gray LLP, One International Place, Boston, MA 02110 (US).

(22) International Filing Date:
18 December 2007 (18.12.2007)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/875,682 18 December 2006 (18.12.2006) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

(71) Applicant (for all designated States except US): ACCELERON PHARMA INC. [US/US]; 149 Sidney Street, Cambridge, MA 02139 (US).

[Continued on next page]

(54) Title: ACTIVIN-ACTRII ANTAGONISTS AND USES FOR INCREASING RED BLOOD CELL LEVELS



(57) Abstract: In certain aspects, the present invention provides compositions and methods for increasing red blood cell and/or hemoglobin levels in vertebrates, including rodents and primates, and particularly in humans.




Figure 1



FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:
18 September 2008

Published:

- *with international search report*
- *the filing date of the international application is within two months from the date of expiration of the priority period*

(48) Date of publication of this corrected version:
13 November 2008

(15) Information about Correction:
see Notice of 13 November 2008

ACTIVIN-ACTRII ANTAGONISTS AND USES FOR INCREASING RED BLOOD CELL LEVELS

RELATED APPLICATIONS

5 This application claims the benefit of U.S. Provisional Patent Application No. 60/875,682, filed December 18, 2006, which application is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

10 Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

15 The mature red blood cell, or erythrocyte, is responsible for oxygen transport in the circulatory systems of vertebrates. Red blood cells carry high concentrations of hemoglobin, a protein that binds oxygen in the lungs at relatively high partial pressure of oxygen (pO_2) and delivers oxygen to areas of the body with a relatively low pO_2 .

Mature red blood cells are produced from pluripotent hematopoietic stem cells in a process termed erythropoiesis. In post-natal individuals, erythropoiesis occurs primarily in the bone marrow and in the red pulp of the spleen. The coordinated action of various signaling pathways control the balance of cell proliferation, differentiation, 20 survival and death. Under normal conditions, red blood cells are produced at a rate that maintains a constant red cell mass in the body, and production may increase or decrease in response to various stimuli, including increased or decreased oxygen tension or tissue demand. The process of erythropoiesis begins with the formation of lineage committed precursor cells and proceeds through a series of distinct precursor cell types. The final 25 stages of erythropoiesis occur as reticulocytes are released into the bloodstream and lose their mitochondria and ribosomes while assuming the morphology of mature red blood cell. An elevated level of reticulocytes, or an elevated reticulocyte:erythrocyte ratio, in the blood is indicative of increased red blood cell production rates.

Erythropoietin (Epo) is widely recognized as the most significant positive regulator of erythropoiesis in post-natal vertebrates. Epo regulates the compensatory erythropoietic response to reduced tissue oxygen tension (hypoxia) and low red blood cell levels or low hemoglobin levels. In humans, elevated Epo levels promote red blood cell formation by _____

stimulating the generation of erythroid progenitors in the bone marrow and spleen. In the mouse, Epo enhances erythropoiesis primarily in the spleen.

Various forms of recombinant Epo are used by physicians to increase red blood cell levels in a variety of clinical settings, and particularly for the treatment of anemia. Anemia is 5 a broadly-defined condition characterized by lower than normal levels of hemoglobin or red blood cells in the blood. In some instances, anemia is caused by a primary disorder in the production or survival of red blood cells. More commonly, anemia is secondary to diseases of other systems (Weatherall & Provan (2000) Lancet 355, 1169-1175). Anemia may result from a reduced rate of production or increased rate of destruction of red blood cells or by loss 10 of red blood cells due to bleeding. Anemia may result from a variety of disorders that include, for example, chronic renal failure, myelodysplastic syndrome, rheumatoid arthritis, and bone marrow transplantation.

Treatment with Epo typically causes a rise in hemoglobins by about 1-3 g/dL in 15 healthy humans over a period of weeks. When administered to anemic individuals, this treatment regimen often provides substantial increases in hemoglobin and red blood cell levels and leads to improvements in quality of life and prolonged survival. Epo is not uniformly effective, and many individuals are refractory to even high doses (Horl et al. (2000) Nephrol Dial Transplant 15, 43-50). Over 50% of patients with cancer have an 20 inadequate response to Epo, approximately 10% with end-stage renal disease are hyporesponsive (Glaspy et al. (1997) J Clin Oncol 15, 1218-1234; Demetri et al. (1998) J Clin Oncol 16, 3412-3425), and less than 10% with myelodysplastic syndrome respond 25 favorably (Estey (2003) Curr Opin Hematol 10, 60-67). Several factors, including inflammation, iron and vitamin deficiency, inadequate dialysis, aluminum toxicity, and hyperparathyroidism may predict a poor therapeutic response, the molecular mechanisms of resistance to Epo are as yet unclear.

Thus, it is an object of the present disclosure to provide alternative compositions and methods for increasing red blood cell levels in patients.

SUMMARY OF THE INVENTION

According to a first aspect, the present invention provides a method for increasing red blood cell levels in a human patient, the method comprising administering to a human patient in need thereof an effective amount of an ActRII polypeptide selected from the group consisting of:

- 5 a) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:2;
- 10 b) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:3;
- 15 c) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:2;
- d) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:16;
- e) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:17; and
- 15 f) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:16.

According to a second aspect, the present invention provides a method for preventing or treating an anemia, the method comprising administering to a subject in need thereof an effective amount of an activin antagonist or ActRII polypeptide.

According to a third aspect, the present invention provides a method of identifying an agent that increases red blood cell levels, the method comprising:

- 25 a) identifying a test agent that binds to a ligand-binding domain of an ActRII polypeptide competitively with an ActRII ligand; and
- b) evaluating the effect of the agent on red blood cell levels in an animal.

According to a fourth aspect, the present invention provides use of an activin antagonist or ActRII polypeptide for making a medicament for increasing red blood cell levels in a human patient.

According to a fifth aspect, the present invention provides a method for increasing red blood cell levels in a human patient, the method comprising administering to a patient in need thereof an effective amount of an ActRII-Fc fusion protein, wherein the ActRII-Fc fusion protein comprises a polypeptide selected from the group consisting of:

- 5 a) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:3;
- 10 b) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:2;
- 15 c) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:7;
- d) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:17;
- 15 e) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:16;
- f) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:20;
- 20 g) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:21; and
- h) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:12.

According to a sixth aspect, the present invention provides use of an activin antagonist or ActRII polypeptide in the manufacture of a medicament for preventing or treating an anemia in a patient.

25 According to a seventh aspect, the present invention provides use of an ActRII-Fc fusion protein, wherein the ActRII-Fc fusion protein comprises an amino acid sequence selected from the group consisting of:

- a) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:3;
- 5 b) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:3;
- c) a polypeptide comprising the amino acid sequence of SEQ ID NO:3;
- d) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:2;
- 10 e) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:2;
- f) a polypeptide comprising the amino acid sequence of SEQ ID NO:2;
- g) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:7;
- h) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:7;
- 15 i) a polypeptide comprising the amino acid sequence of SEQ ID NO:7;
- j) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:17;
- k) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:17;
- 20 l) a polypeptide comprising the amino acid sequence of SEQ ID NO:17;
- m) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:16;
- n) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:16;
- 25 o) a polypeptide comprising the amino acid sequence of SEQ ID NO:16;
- p) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:20;

Unless the context clearly requires otherwise, throughout the description and the claims, the words "comprise", "comprising", and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".

5 In part, the disclosure demonstrates that activin antagonists, as well as ActRIIa and ActRIIb antagonists, can be used to increase red blood cell and hemoglobin levels. In particular, the disclosure demonstrates that a soluble form of ActRIIa acts as an inhibitor of activin and, when administered *in vivo*, increases red blood cell levels in the blood. A milder effect was observed with a soluble form of ActRIIb, which binds

10 Activin A with lesser affinity than soluble ActRIIa. While soluble ActRIIa and ActRIIb may affect red blood cell levels through a mechanism other than activin antagonism, the disclosure nonetheless demonstrates that desirable therapeutic agents may be selected on the basis of activin antagonism or ActRII antagonism or both. Such agents are referred to collectively as activin-ActRII antagonists. Therefore, in certain embodiments, the

15 disclosure provides methods for using activin-ActRII antagonists, including, for example, activin-binding ActRIIa polypeptides, activin-binding ActRIIb polypeptides, anti-activin antibodies, anti-ActRIIa antibodies, anti-ActRIIb antibodies, activin-, ActRIIb-, or ActRIIa-targeted small molecules and aptamers, and nucleic acids that decrease expression of activin, ActRIIb, or ActRIIa, to increase red blood cell and

20 hemoglobin levels in patients and to treat disorders associated with low red blood cell or hemoglobin levels in patients in need thereof. As described in U.S. Patent Application Serial No. 11/603,485, incorporated by reference herein, activin-ActRIIa antagonists can be used to promote bone growth and increase bone density. As described herein, the effects of such antagonists on red blood cell levels are more rapid and occur at lower

25 doses than the effects of such antagonists on bone. Thus, in certain embodiments, the disclosure provides methods for using an activin-ActRIIa antagonist to increase red blood cell or hemoglobin levels without causing a significant increase in bone density. For example, a method may cause less than 3%, 5%, 10% or 15% increase in bone density. This selective effect may be achieved by using, for example, lower doses of

30 activin-ActRIIa antagonist, less frequent doses, or by using an activin-ActRIIa antagonist with a shorter serum half-life at doses and frequencies calculated to provide a lower serum concentration.

- q) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:20;
- r) a polypeptide comprising the amino acid sequence of SEQ ID NO:20;
- s) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:21;
- 5 t) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:21;
- u) a polypeptide comprising the amino acid sequence of SEQ ID NO:21;
- v) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:12;
- 10 w) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:12;
- x) a polypeptide comprising the amino acid sequence of SEQ ID NO:12, in the manufacture of a medicament for increasing red blood cell levels in a human patient.

15 Unless the context clearly requires otherwise, throughout the description and the claims, the words “comprise”, “comprising”, and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of “including, but not limited to”.

20 In part, the disclosure demonstrates that activin antagonists, as well as ActRIIa and ActRIIb antagonists, can be used to increase red blood cell and hemoglobin levels. In particular, the disclosure demonstrates that a soluble form of ActRIIa acts as an inhibitor of activin and, when administered in vivo, increases red blood cell levels in the blood. A milder effect was observed with a soluble form of ActRIIb, which binds Activin A with lesser affinity than soluble ActRIIa. While soluble ActRIIa and ActRIIb 25 may affect red blood cell levels through a mechanism other than activin antagonism, the disclosure nonetheless demonstrates that desirable therapeutic agents may be selected on the basis of activin antagonism or ActRII antagonism or both. Such agents are referred to collectively as activin-ActRII antagonists. Therefore, in certain embodiments, the disclosure provides methods for using activin-ActRII antagonists, including, for


example, activin-binding ActRIIa polypeptides, activin-binding ActRIIb polypeptides, anti-activin antibodies, anti-ActRIIa antibodies, anti-ActRIIb antibodies, activin-, ActRIIb-, or ActRIIa-targeted small molecules and aptamers, and nucleic acids that decrease expression of activin, ActRIIb, or ActRIIa, to increase red blood cell and

5 hemoglobin levels in patients and to treat disorders associated with low red blood cell or hemoglobin levels in patients in need thereof. As described in U.S. Patent Application Serial No. 11/603,485, incorporated by reference herein, activin-ActRIIa antagonists can be used to promote bone growth and increase bone density. As described herein, the effects of such antagonists on red blood cell levels are more rapid and occur at lower

10 doses than the effects of such antagonists on bone. Thus, in certain embodiments, the disclosure provides methods for using an activin-ActRIIa antagonist to increase red blood cell or hemoglobin levels without causing a significant increase in bone density. For example, a method may cause less than 3%, 5%, 10% or 15% increase in bone density. This selective effect may be achieved by using, for example, lower doses of

15 activin-ActRIIa antagonist, less frequent doses, or by using an activin-ActRIIa antagonist with a shorter serum half-life at doses and frequencies calculated to provide a lower serum concentration.

In certain aspects, the disclosure provides polypeptides comprising a soluble, activin-binding ActRII polypeptide that binds to activin. The activin binding 20 polypeptide may be an ActRIIa polypeptide or an ActRIIb polypeptide. ActRII polypeptides may be formulated as a pharmaceutical preparation comprising the activin-binding ActRII polypeptide and a




pharmaceutically acceptable carrier. The activin-binding ActRII polypeptide may bind to activin with a K_D less than 1 micromolar or less than 100, 10 or 1 nanomolar. Optionally, the activin-binding ActRII polypeptide selectively binds activin versus GDF11 and/or GDF8, and optionally with a K_D that is at least 10-fold, 20-fold or 50-fold lower with respect to activin than with respect to GDF11 and/or GDF8. While not wishing to be bound to a particular mechanism of action, it is expected that this degree of selectivity for activin inhibition over GDF11/GDF8 inhibition accounts for effects on bone or erythropoiesis without a consistently measurable effect on muscle. In many embodiments, an ActRII polypeptide will be selected for causing less than 15%, less than 10% or less than 5% increase in muscle at doses that achieve desirable effects on red blood cell levels. The composition may be at least 95% pure, with respect to other polypeptide components, as assessed by size exclusion chromatography, and optionally, the composition is at least 98% pure. An activin-binding ActRIIa polypeptide for use in such a preparation may be any of those disclosed herein, such as a polypeptide having an amino acid sequence selected from SEQ ID NOs: 2, 3, 7 or 12, or having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97% or 99% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 12 or 13. An activin-binding ActRIIa polypeptide may include a functional fragment of a natural ActRIIa polypeptide, such as one comprising at least 10, 20 or 30 amino acids of a sequence selected from SEQ ID NOs: 1-3 or a sequence of SEQ ID NO: 2, lacking the C-terminal 10 to 15 amino acids (the "tail"). An activin-binding ActRIIb polypeptide for use in such a preparation may be any of those disclosed herein, such as a polypeptide having an amino acid sequence selected from SEQ ID NOs: 16, 17, 20, or 21 or having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97% or 99% identical to an amino acid sequence selected from SEQ ID NOs: 16, 17, 20, or 21. An activin-binding ActRIIb polypeptide may include a functional fragment of a natural ActRIIb polypeptide, such as one comprising at least 10, 20 or 30 amino acids of SEQ ID NOs: 15-17 or a sequence lacking the C-terminal 10 to 15 amino acids (the "tail") such as SEQ ID NO: 17.

A soluble, activin-binding ActRII polypeptide may include one or more alterations in the amino acid sequence (e.g., in the ligand-binding domain) relative to a naturally occurring ActRII polypeptide. Examples of altered ActRIIa and ActRIIb polypeptides are provided in WO 2006/012627, pp. 59-60 and pp. 55-58, respectively, which is incorporated by reference herein. The alteration in the amino acid sequence may, for example, alter glycosylation of

the polypeptide when produced in a mammalian, insect or other eukaryotic cell or alter proteolytic cleavage of the polypeptide relative to the naturally occurring ActRII polypeptide.

An activin-binding ActRII polypeptide may be a fusion protein that has, as one domain, an ActRII polypeptide, (e.g., a ligand-binding portion of an ActRIIa or ActRIIb) and one or more additional domains that provide a desirable property, such as improved pharmacokinetics, easier purification, targeting to particular tissues, etc. For example, a domain of a fusion protein may enhance one or more of in vivo stability, in vivo half life, uptake/administration, tissue localization or distribution, formation of protein complexes, multimerization of the fusion protein, and/or purification. An activin-binding ActRII fusion protein may include an immunoglobulin Fc domain (wild-type or mutant) or a serum albumin or other polypeptide portion that provides desirable properties such as improved pharmacokinetics, improved solubility or improved stability. In a preferred embodiment, an ActRII-Fc fusion comprises a relatively unstructured linker positioned between the Fc domain and the extracellular ActRII domain. This unstructured linker may correspond to the roughly 15 amino acid unstructured region at the C-terminal end of the extracellular domain of ActRII (the "tail"), or it may be an artificial sequence of 1, 2, 3, 4 or 5 amino acids or a length of between 5 and 15, 20, 30, 50 or more amino acids that are relatively free of secondary structure, or a mixture of both. A linker may be rich in glycine and proline residues and may, for example, contain a single sequence of threonine/serine and glycines or repeating sequences of threonine/serine and glycines (e.g., TG₄ (SEQ ID NO: 22) or SG₄ (SEQ ID NO: 23) singlets or repeats). A fusion protein may include a purification subsequence, such as an epitope tag, a FLAG tag, a polyhistidine sequence, and a GST fusion. Optionally, a soluble ActRII polypeptide includes one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent. A pharmaceutical preparation may also include one or more additional compounds such as a compound that is used to treat a bone disorder.

Preferably, a pharmaceutical preparation is substantially pyrogen free. In general, it is preferable that an ActRII protein be expressed in a mammalian cell line that mediates suitably natural glycosylation of the ActRII protein so as to diminish the likelihood of an unfavorable immune response in a patient. Human and CHO cell lines 5 have been used successfully, and it is expected that other common mammalian expression systems will be useful. —



As described herein, ActRIIa proteins designated ActRIIa-Fc (a form with a minimal linker between the ActRIIa portion and the Fc portion) have desirable properties, including selective binding to activin versus GDF8 and/or GDF11, high affinity ligand binding and serum half life greater than two weeks in animal models. In certain embodiments the

5 invention provides ActRII-Fc polypeptides and pharmaceutical preparations comprising such polypeptides and a pharmaceutically acceptable excipient.

In certain aspects, the disclosure provides nucleic acids encoding a soluble activin-binding ActRII polypeptide, such as an ActRIIa or ActRIIb polypeptide. An isolated polynucleotide may comprise a coding sequence for a soluble, activin-binding ActRII polypeptide, such as described above. For example, an isolated nucleic acid may include a sequence coding for an extracellular domain (e.g., ligand-binding domain) of an ActRII and a sequence that would code for part or all of the transmembrane domain and/or the cytoplasmic domain of an ActRII, but for a stop codon positioned within the transmembrane domain or the cytoplasmic domain, or positioned between the extracellular domain and the transmembrane domain or cytoplasmic domain. For example, an isolated polynucleotide may comprise a full-length ActRIIa polynucleotide sequence such as SEQ ID NO: 4 or 5 or a full-length ActRIIb polynucleotide sequence such as SEQ ID NO: 18, or a partially truncated version of ActRIIa or ActRIIb, said isolated polynucleotide further comprising a transcription termination codon at least six hundred nucleotides before the 3'-terminus or otherwise positioned such that translation of the polynucleotide gives rise to an extracellular domain optionally fused to a truncated portion of a full-length ActRII. A preferred nucleic acid sequence for ActRIIa is SEQ ID NO:14. Nucleic acids disclosed herein may be operably linked to a promoter for expression, and the disclosure provides cells transformed with such recombinant polynucleotides. Preferably the cell is a mammalian cell such as a CHO cell.

25 In certain aspects, the disclosure provides methods for making a soluble, activin-binding ActRII polypeptide. Such a method may include expressing any of the nucleic acids (e.g., SEQ ID NO: 4, 5 14, 18, or 19) disclosed herein in a suitable cell, such as a Chinese hamster ovary (CHO) cell. Such a method may comprise: a) culturing a cell under conditions suitable for expression of the soluble ActRII polypeptide, wherein said cell is transformed 30 with a soluble ActRII expression construct; and b) recovering the soluble ActRII polypeptide so expressed. Soluble ActRII polypeptides may be recovered as crude, partially purified or

highly purified fractions. Purification may be achieved by a series of purification steps, including, for example, one, two or three or more of the following, in any order: protein A chromatography, anion exchange chromatography (e.g., Q sepharose), hydrophobic interaction chromatography (e.g., phenylsepharose), size exclusion chromatography, and 5 cation exchange chromatography.

In certain aspects, an activin-ActRII antagonist disclosed herein, such as a soluble, activin-binding ActRIIa polypeptide or soluble, activin-binding ActRIIb polypeptide, may be used in a method for promoting red blood cell production or increasing red blood cell levels in a subject. In certain embodiments, the disclosure provides methods for treating a disorder 10 associated with low red blood cell counts or low hemoglobin levels (e.g., an anemia), or to promote red blood cell production, in patients in need thereof. A method may comprise administering to a subject in need thereof an effective amount of activin-ActRII antagonist. In certain aspects, the disclosure provides uses of activin-ActRII antagonists for making a medicament for the treatment of a disorder or condition as described herein.

15 In certain aspects, the disclosure provides a method for identifying an agent that stimulates production of red blood cells. The method comprises: a) identifying a test agent that binds to activin or a ligand-binding domain of an ActRII polypeptide; and b) evaluating the effect of the agent on the levels of red blood cells, hemoglobin, and/or red blood cell precursor levels (e.g., reticulocyte levels).

20

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the purification of ActRIIa-hFc expressed in CHO cells. The protein purifies as a single, well-defined peak as visualized by sizing column (left panel) and Coomassie stained SDS-PAGE (right panel) (left lane: molecular weight standards; right 25 lane: ActRIIa-hFc).

Figure 2 shows the binding of ActRIIa-hFc to activin and GDF-11, as measured by BiaCoreTM assay.

Figure 3 shows the effects of ActRIIa-hFc on red blood cell counts in female non-human primates. Female cynomolgus monkeys (four groups of five monkeys each) were 30 treated with placebo or 1mg/kg, 10 mg/kg or 30 mg/kg of ActRIIa-hFc on day 0, day 7, day

14 and day 21. Figure 3A shows red blood cell (RBC) counts. Figure 3B shows hemoglobin levels. Statistical significance is relative to baseline for each treatment group. At day 57, two monkeys remained in each group.

5 Figure 4 shows the effects of ActRIIa-hFc on red blood cell counts in male non-human primates. Male cynomolgus monkeys (four groups of five monkeys each) were treated with placebo or 1mg/kg, 10 mg/kg or 30 mg/kg of ActRIIa-hFc on day 0, day 7, day 14 and day 21. Figure 4A shows red blood cell (RBC) counts. Figure 4B shows hemoglobin levels. Statistical significance is relative to baseline for each treatment group. At day 57, two monkeys remained in each group.

10 Figure 5 shows the effects of ActRIIa-hFc on reticulocyte counts in female non-human primates. Cynomolgus monkeys (four groups of five monkeys each) were treated with placebo or 1 mg/kg, 10 mg/kg or 30 mg/kg of ActRIIa-hFc on day 0, day 7, day 14 and day 21. Figure 5A shows absolute reticulocyte counts. Figure 5B shows the percentage of reticulocytes relative to RBCs. Statistical significance is relative to baseline for each group.

15 At day 57, two monkeys remained in each group.

Figure 6 shows the effects of ActRIIa-hFc on reticulocyte counts in female non-human primates. Cynomolgus monkeys (four groups of five monkeys each) were treated with placebo or 1 mg/kg, 10 mg/kg or 30 mg/kg of ActRIIa-hFc on day 0, day 7, day 14 and day 21. Figure 6A shows absolute reticulocyte counts. Figure 6B shows the percentage of reticulocytes relative to RBCs. Statistical significance is relative to baseline for each group.

20 At day 57, two monkeys remained in each group.

Figure 7 shows results from the human clinical trial described in Example 5, where the area-under-curve (AUC) and administered dose of ActRIIa-hFc have a linear correlation, regardless of whether ActRIIa-hFc was administered intravenously (IV) or subcutaneously (SC).

25 Figure 8 shows a comparison of serum levels of ActRIIa-hFc in patients administered IV or SC.

Figure 9 shows bone alkaline phosphatase (BAP) levels in response to different dose levels of ActRIIa-hFc. BAP is a marker for anabolic bone growth.

30 Figure 10 depicts the median change from baseline of hematocrit levels from the human clinical trial described in Example 5. ActRIIa-hFc was administered intravenously (IV) at the indicated dosage.

Figure 11 depicts the median change from baseline of hemoglobin levels from the human clinical trial described in Example 5. ActRIIa-hFc was administered intravenously (IV) at the indicated dosage.

Figure 12 depicts the median change from baseline of RBC (red blood cell) count
5 from the human clinical trial described in Example 5. ActRIIa-hFc was administered intravenously (IV) at the indicated dosage.

Figure 13 depicts the median change from baseline of reticulocyte count from the human clinical trial described in Example 5. ActRIIa-hFc was administered intravenously (IV) at the indicated dosage.

10

DETAILED DESCRIPTION OF THE INVENTION

1. Overview

The transforming growth factor-beta (TGF-beta) superfamily contains a variety of growth factors that share common sequence elements and structural motifs. These proteins
15 are known to exert biological effects on a large variety of cell types in both vertebrates and invertebrates. Members of the superfamily perform important functions during embryonic development in pattern formation and tissue specification and can influence a variety of differentiation processes, including adipogenesis, myogenesis, chondrogenesis, cardiogenesis, hematopoiesis, neurogenesis, and epithelial cell differentiation. The family is
20 divided into two general branches: the BMP/GDF and the TGF-beta/Activin/BMP10 branches, whose members have diverse, often complementary effects. By manipulating the activity of a member of the TGF-beta family, it is often possible to cause significant physiological changes in an organism. For example, the Piedmontese and Belgian Blue cattle breeds carry a loss-of-function mutation in the GDF8 (also called myostatin) gene that causes
25 a marked increase in muscle mass. Grobet et al., Nat Genet. 1997, 17(1):71-4. Furthermore, in humans, inactive alleles of GDF8 are associated with increased muscle mass and, reportedly, exceptional strength. Schuelke et al., N Engl J Med 2004, 350:2682-8.

Activins are dimeric polypeptide growth factors that belong to the TGF-beta superfamily. There are three principal activin forms (A, B, and AB) that are
30 homo/heterodimers of two closely related β subunits ($\beta_A\beta_A$, $\beta_B\beta_B$, and $\beta_A\beta_B$, respectively).

The human genome also encodes an activin C and an activin E, which are primarily expressed in the liver, and heterodimeric forms containing β_C or β_E are also known. In the TGF-beta superfamily, activins are unique and multifunctional factors that can stimulate hormone production in ovarian and placental cells, support neuronal cell survival, influence 5 cell-cycle progress positively or negatively depending on cell type, and induce mesodermal differentiation at least in amphibian embryos (DePaolo et al., 1991, Proc Soc Ep Biol Med. 198:500-512; Dyson et al., 1997, Curr Biol. 7:81-84; Woodruff, 1998, Biochem Pharmacol. 55:953-963). Moreover, erythroid differentiation factor (EDF) isolated from the stimulated human monocytic leukemic cells was found to be identical to activin A (Murata et al., 1988, 10 PNAS, 85:2434). It has been suggested that activin A promotes erythropoiesis in the bone marrow. In several tissues, activin signaling is antagonized by its related heterodimer, inhibin. For example, during the release of follicle-stimulating hormone (FSH) from the pituitary, activin promotes FSH secretion and synthesis, while inhibin prevents FSH secretion and synthesis. Other proteins that may regulate activin bioactivity and/or bind to activin 15 include follistatin (FS), follistatin-related protein (FSRP) and α_2 -macroglobulin.

TGF- β signals are mediated by heteromeric complexes of type I and type II serine/threonine kinase receptors, which phosphorylate and activate downstream Smad proteins upon ligand stimulation (Massagué, 2000, Nat. Rev. Mol. Cell Biol. 1:169-178). These type I and type II receptors are transmembrane proteins, composed of a ligand-binding extracellular 20 domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II activin receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors.

25 Two related type II receptors (ActRII), ActRIIa and ActRIIb, have been identified as the type II receptors for activins (Mathews and Vale, 1991, Cell 65:973-982; Attisano et al., 1992, Cell 68: 97-108). Besides activins, ActRIIa and ActRIIb can biochemically interact with several other TGF- β family proteins, including BMP7, Nodal, GDF8, and GDF11 (Yamashita et al., 1995, J. Cell Biol. 130:217-226; Lee and McPherron, 2001, Proc. Natl. 30 Acad. Sci. 98:9306-9311; Yeo and Whitman, 2001, Mol. Cell 7: 949-957; Oh et al., 2002,

Genes Dev. 16:2749-54). ALK4 is the primary type I receptor for activins, particularly for activin A, and ALK-7 may serve as a receptor for activins as well, particularly for activin B.

As demonstrated herein, a soluble ActRIIa polypeptide (sActRIIa), which shows substantial preference in binding to activin A as opposed to other TGF-beta family members, 5 such as GDF8 or GDF11, is effective to increase red blood cell levels in vivo. While not wishing to be bound to any particular mechanism, it is expected that the effect of sActRIIa is caused primarily by an activin antagonist effect, given the very strong activin binding (picomolar dissociation constant) exhibited by the particular sActRIIa construct used in these studies. Regardless of mechanism, it is apparent from this disclosure that ActRIIa-activin 10 antagonists increase red blood cell levels in rodents, monkeys and humans. It should be noted that hematopoiesis is a complex process, regulated by a variety of factors, including erythropoietin, G-CSF and iron homeostasis. The terms "increase red blood cell levels" and "promote red blood cell formation" refer to clinically observable metrics, such as hematocrit, red blood cell counts and hemoglobin measurements, and are intended to be neutral as to the 15 mechanism by which such changes occur.

As also demonstrated herein, a soluble ActRIIb polypeptide (sActRIIb) is effective to increase reticulocyte levels in vivo, an effect which, over a longer time period is expected to cause increased hematocrit levels.

The data reported herein with respect to non-human primates are reproducible in 20 mice, rats and humans as well, and therefore, this disclosure provides methods for using ActRII polypeptides and other activin-ActRII antagonists to promote red blood cell production and increase red blood cell levels in mammals ranging from rodents to humans. Activin-ActRII antagonists include, for example, activin-binding soluble ActRIIa 25 polypeptides, activin-binding soluble ActRIIb polypeptides, antibodies that bind to activin (particularly the activin A or B subunits, also referred to as β A or β B) and disrupt ActRIIa and/or ActRIIb binding, antibodies that bind to ActRIIa and disrupt activin binding, antibodies that bind to ActRIIb and disrupt activin binding, non-antibody proteins selected for activin, ActRIIb or ActRIIa binding (see e.g., WO/2002/088171, WO/2006/055689, and WO/2002/032925 for examples of such proteins and methods for design and selection of 30 same), randomized peptides selected for activin, ActRIIb, or ActRIIa binding, often affixed to an Fc domain. Two different proteins (or other moieties) with activin, ActRIIb, or ActRIIa

binding activity, especially activin binders that block the type I (e.g., a soluble type I activin receptor) and type II (e.g., a soluble type II activin receptor) binding sites, respectively, may be linked together to create a bifunctional binding molecule. Nucleic acid aptamers, small molecules and other agents that inhibit the activin-ActRII signaling axis are included as 5 activin-ActRII antagonists. Various proteins have activin-ActRII antagonist activity, including inhibin (i.e., inhibin alpha subunit), although inhibin does not universally antagonize activin in all tissues, follistatin (e.g., follistatin-288 and follistatin-315), FSRP, activin C, alpha(2)-macroglobulin, and an M108A (methionine to alanine change at position 108) mutant activin A. Generally, alternative forms of activin, particularly those with 10 alterations in the type I receptor binding domain can bind to type II receptors and fail to form an active ternary complex, thus acting as antagonists. Additionally, nucleic acids, such as antisense molecules, siRNAs or ribozymes that inhibit activin A, B, C or E, or, particularly, ActRIIa or ActRIIb expression, can be used as activin-ActRII antagonists. The activin-ActRII antagonist to be used may exhibit selectivity for inhibiting activin-mediated signaling 15 versus other members of the TGF-beta family, and particularly with respect to GDF8 and GDF11.

The terms used in this specification generally have their ordinary meanings in the art, within the context of this invention and in the specific context where each term is used. Certain terms are discussed below or elsewhere in the specification, to provide additional 20 guidance to the practitioner in describing the compositions and methods of the invention and how to make and use them. The scope or meaning of any use of a term will be apparent from the specific context in which the term is used.

“About” and “approximately” shall generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Typically, 25 exemplary degrees of error are within 20 percent (%), preferably within 10%, and more preferably within 5% of a given value or range of values.

Alternatively, and particularly in biological systems, the terms “about” and “approximately” may mean values that are within an order of magnitude, preferably within 5-fold and more preferably within 2-fold of a given value. Numerical quantities given herein 30 are approximate unless stated otherwise, meaning that the term “about” or “approximately” can be inferred when not expressly stated.

The methods of the invention may include steps of comparing sequences to each other, including wild-type sequence to one or more mutants (sequence variants). Such comparisons typically comprise alignments of polymer sequences, e.g., using sequence alignment programs and/or algorithms that are well known in the art (for example, BLAST, 5 FASTA and MEGALIGN, to name a few). The skilled artisan can readily appreciate that, in such alignments, where a mutation contains a residue insertion or deletion, the sequence alignment will introduce a “gap” (typically represented by a dash, or “A”) in the polymer sequence not containing the inserted or deleted residue.

“Homologous,” in all its grammatical forms and spelling variations, refers to the 10 relationship between two proteins that possess a “common evolutionary origin,” including proteins from superfamilies in the same species of organism, as well as homologous proteins from different species of organism. Such proteins (and their encoding nucleic acids) have sequence homology, as reflected by their sequence similarity, whether in terms of percent identity or by the presence of specific residues or motifs and conserved positions.

15 The term “sequence similarity,” in all its grammatical forms, refers to the degree of identity or correspondence between nucleic acid or amino acid sequences that may or may not share a common evolutionary origin.

20 However, in common usage and in the instant application, the term “homologous,” when modified with an adverb such as “highly,” may refer to sequence similarity and may or may not relate to a common evolutionary origin.

2. ActRII Polypeptides

In certain aspects, the present invention relates to ActRII polypeptides. As used herein, the term "ActRII" refers to the family of type II activin receptors. This family 25 includes both the activin receptor type IIa and the activin receptor type IIb.

In certain aspects, the present invention relates to ActRIIa polypeptides. As used herein, the term “ActRIIa” refers to a family of activin receptor type IIa (ActRIIa) proteins from any species and variants derived from such ActRIIa proteins by mutagenesis or other modification. Reference to ActRIIa herein is understood to be a reference to any one of the 30 currently identified forms. Members of the ActRIIa family are generally transmembrane

proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine kinase activity.

The term “ActRIIa polypeptide” includes polypeptides comprising any naturally occurring polypeptide of an ActRIIa family member as well as any variants thereof (including mutants, fragments, fusions, and peptidomimetic forms) that retain a useful activity. See, for example, WO/2006/012627. For example, ActRIIa polypeptides include polypeptides derived from the sequence of any known ActRIIa having a sequence at least about 80% identical to the sequence of an ActRIIa polypeptide, and optionally at least 85%, 90%, 95%, 97%, 99% or greater identity. For example, an ActRIIa polypeptide of the invention may bind to and inhibit the function of an ActRIIa protein and/or activin. An ActRIIa polypeptide may be selected for activity in promoting red blood cell formation in vivo. Examples of ActRIIa polypeptides include human ActRIIa precursor polypeptide (SEQ ID NO: 1) and soluble human ActRIIa polypeptides (e.g., SEQ ID NOS: 2, 3, 7 and 12).

15 The human ActRIIa precursor protein sequence is as follows:

MGAAAKLAFAVFLISCSSA**ILGRSETQECLFFNANWEKDRTN**QTGVEP
CYGDKDKRRHCFATWKNISGSIEIVKQGCWLDDINCYDRTDCVEKKDSP
EVYFCCCEGNMCNEKFSYFPPEMEVTQOPTSNPVTPKPPYYNILLYSLVPL
MLIAGIVICAFWVYRHHKAYPPVLVPTQDPGPPPPSPLLGLKPLQLLE
20 VKARGRFGCVWKAQOLLNEYVAVKIFPIQDKQSWQNEYEVSLPGMKHEN
ILQFIGAEKRGTSDVDLWLITAFHEKGSLSDFLKANVVSNELCHIAE
TMARGLAYLHEDIPLGLDGHKPAISHRDIKSKNVLLKNNLTACIADFGL
ALKFEAGKSAGDTHGQVGTRRYMAPEVLEGAINFQRDAFLRIDMYAMGL
VILWELASRCTAADGPVDEYMLPFEETIGQHPSLEDMQEVVVHKKRGPVL
25 RDYWQKHAGMAMLCETIEECWDHDAEARLSAGCVGERITQMQLTNIIIT
TEDIVTVVTMVTNVDFPPKESSL (SEQ ID NO: 1)

The signal peptide is single underlined; the extracellular domain is in bold and the potential N-linked glycosylation sites are double underlined.

The human ActRIIa soluble (extracellular), processed polypeptide sequence is as follows:

5 ILGRSETQECLFFNANWEKDRTNQTGVEPCYGDKDKRHCFATWKNISG
 SIEIVKQGCWLDDDINCYDRTDCVEKKDSPEVYFCCCEGNMCNEKFSYFP
EMEVTQPTSNPVTPKPP (SEQ ID NO: 2)

The C-terminal “tail” of the extracellular domain is underlined. The sequence with the “tail” deleted (a Δ15 sequence) is as follows:

10 ILGRSETQECLFFNANWEKDRTNQTGVEPCYGDKDKRHCFATWKNISG
 SIEIVKQGCWLDDDINCYDRTDCVEKKDSPEVYFCCCEGNMCNEKFSYFP
 EM (SEQ ID NO:3)

The nucleic acid sequence encoding human ActRIIa precursor protein is as follows (nucleotides 164-1705 of Genbank entry NM_001616):

15 ATGGGAGCTGCTGCAAAGTTGGCGTTGCCGTCTTCTTATCTCCTGTT
 CTTCAGGTGCTATACTTGGTAGATCAGAACTCAGGAGTGTCTTTCTT
 TAATGCTAATTGGGAAAAGACAGAACCAATCAAACTGGTGTGAACCG
 TGTTATGGTGACAAAGATAAACGCCGGATTGTTGCTACCTGGAAGA
 ATATTCTGGTCCATTGAAATAGTGAACAAGGTTGTTGGCTGGATGA
 TATCAACTGCTATGACAGGACTGATTGTGTAGAAAAAAAGACAGCCCT
 GAAGTATATTTGTTGCTGTGAGGGCAATATGTGAATGAAAAGTTT
 20 CTTATTTCCAGAGATGGAAGTCACACAGCCCACTTCAAATCCAGTTAC
 ACCTAACGCCACCCTTATTACAAACATCCTGCTTATTCCCTGGTGCCACTT
 ATGTTAATTGCGGGGATTGTCATTGTCATTGTCATTGGGTGTACAGGCATC
 ACAAGATGGCCTACCCCTCTGTTACTTGTCCAACTCAAGACCCAGGACC
 ACCCCCACCTCTCCATTACTAGGGTTGAAACCACTGCAGTTATTAGAA
 25 GTGAAAGCAAGGGGAAGATTGGTGTGTCGGAAAGCCCAGTTGCTTA
 ACGAATATGTGGCTGTCAAAATATTCCAATACAGGACAAACAGTCATG
 GCAAAATGAATACGAAGTCTACAGTTGCCTGGAAATGAAGCATGAGAAC
 ATATTACAGTTCATTGGTGCAGAAAAACGAGGCACCAGTGTGATGTGG
 ATCTTGGCTGATCACAGCATTCATGAAAAAGGGTTCACTATCAGACTT
 30 TCTTAAGGCTATGTGGTCTCTGGAAATGAACTGTGTCATATTGCAGAA
 ACCATGGCTAGAGGATTGGCATATTTACATGAGGATACCTGGCCTAA
 AAGATGGCCCACAAACCTGCCATATCTCACAGGGACATCAAAAGTAAAAA

TGTGCTGTTGAAAAACAAACCTGACAGCTTGCATTGCTGACTTTGGGTTG
GCCTTAAAATTGAGGCTGGCAAGTCTGCAGGCGATAACCATGGACAGG
TTGGTACCCGGAGGTACATGGCTCCAGAGGTATTAGAGGGTGCTATAAA
CTTCCAAAGGGATGCATTTGAGGATAGATATGTATGCCATGGGATTA
5 GTCCTATGGGAACGGCTCTCGCTGTACTGCTGCAGATGGACCTGTAG
ATGAATACATGTTGCCATTGAGGAGGAAATTGGCCAGCATCCATCTCT
TGAAGACATGCAGGAAGTTGTTGTGCATAAAAAAAAGAGGCCTGTTTA
AGAGATTATTGGCAGAACATGCTGGAATGGCAATGCTCTGTGAAACCA
10 TTGAAGAATGTTGGGATCACGACGCAGAACGCCAGGTTATCAGCTGGATG
TGTAGGTGAAAGAATTACCCAGATGCAGAGACTAACAAATATTATTACC
ACAGAGGACATTGTAACAGTGGTCACAATGGTGACAAATGTTGACTTTC
CTCCCAAAGAACATCTAGTCTATGA (SEQ ID NO: 4)

The nucleic acid sequence encoding a human ActRIIa soluble (extracellular) polypeptide is as follows:

15 ATACTGGTAGATCAGAAACTCAGGAGTGTCTTTCTTAATGCTAATT
GGGAAAAAGACAGAACCAATCAAACACTGGTGTGAACCGTGTATGGTGA
CAAAGATAAACGGCGGCATTGTTTGCTACCTGGAAGAATATTCTGGT
TCCATTGAAATAGTGAACACAAGGTTGTTGGCTGGATGATATCAACTGCT
ATGACAGGACTGATTGTAGAAAAAGACAGCCCTGAAGTATATT
20 TTGTTGCTGTGAGGGCAATATGTGAATGAAAAGTTTCTTATTTCCA
GAGATGGAAGTCACACAGCCCACCTCAAATCCAGTTACACCTAACGCCAC
CC (SEQ ID NO: 5)

In certain aspects, the present invention relates to ActRIIb polypeptides. As used herein, the term "ActRIIb" refers to a family of activin receptor type IIb (ActRIIb) proteins from any species and variants derived from such ActRIIb proteins by mutagenesis or other modification. Reference to ActRIIb herein is understood to be a reference to any one of the currently identified forms. Members of the ActRIIb family are generally transmembrane proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine kinase 25 activity.

The term "ActRIIb polypeptide" includes polypeptides comprising any naturally occurring polypeptide of an ActRIIb family member as well as any variants thereof (including mutants, fragments, fusions, and peptidomimetic forms) that retain a useful activity. See, for example, WO/2006/012627. For example, ActRIIb polypeptides include 5 polypeptides derived from the sequence of any known ActRIIb having a sequence at least about 80% identical to the sequence of an ActRIIb polypeptide, and optionally at least 85%, 90%, 95%, 97%, 99% or greater identity. For example, an ActRIIb polypeptide of the invention may bind to and inhibit the function of an ActRIIb protein and/or activin. An ActRIIb polypeptide may be selected for activity in promoting red blood cell formation in 10 vivo. Examples of ActRIIb polypeptides include human ActRIIb precursor polypeptide (SEQ ID NO: 15) and soluble human ActRIIb polypeptides (e.g., SEQ ID NO: 16, 17, 20, and 21).

The human ActRIIb precursor protein sequence is as follows:

MTAPWVALALLW GSLWPG **SGRGEAETREC IYYNANWELERT** **NQSGLERC**
EGEQD KRLHCYASWAN **SSGTIELVKKGCWLDDFNCYDRQECVATEENPQ**
15 **VYFCCCEGNFCNERFTHLPEAGGPEVTYEPPPTAPTLLTVLAYSLLPIG**
GLSLIVLLAFWMYRHRKPPYGHVDIHD PGPPPSPLVGLKPLQLLEIK
ARGRFGCVWKAQLMNDFAVAKIFPLQDKQSREI FSTPGMKHENLL
QFIAAEKRGSNLEVELWLITAFHDKGSLTDYLKGNITWNELCHVAETM
SRGLSYLHEDVPWCRGEGHKPSIAHRDFKSKNVLLKSDLTAVLADFGLA
20 VRFEPGKPPGDTHGQVGTRRYMAPEVLEGAINFQRDAFLRIDMYAMGLV
LWELVSRCKAADGPVDEYMLP FEEEIGQHPSLEELQEVVVKKMRPTIK
DHWLKHPGLAQLCVTIEECWDHDAEARLSAGCVEERVSLIRRSVNGTTS
DCLVSLVTSVTNVDLPPKESSI (SEQ ID NO: 15)

The signal peptide is single underlined; the extracellular domain is in bold and the 25 potential N-linked glycosylation sites are in boxes.

The human ActRIIb soluble (extracellular), processed polypeptide sequence is as follows:

SGRGEAETREC IYYNANWELERT **NQSGLERC** EGEQD KRLHCYASWANSS
GTIELVKKGCWLDDFNCYDRQECVATEENPQVY FCCCEGNFCNERFTHL
30 **PEAGGPEVTYEPPPTAPT** (SEQ ID NO: 16)

The C-terminal “tail” of the extracellular domain is underlined. The sequence with the “tail” deleted (a Δ15 sequence) is as follows:

SGRGEAETREC~~IYYNAN~~WELRTNQSGLERCEGEQDKRLH~~CYAS~~WANSS
GTIELVKKGCWLDDFN~~CYDR~~QECVATEENPQVYFCCCEGNFCNERFTHL

5 PEA (SEQ ID NO: 17)

The nucleic acid sequence encoding a human ActRIIb precursor protein is as follows:
(nucleotides 5-1543 of Genbank entry NM_001106)

10 ATGACGGCGCCCTGGTGGCCCTGCCCTCCTGGGGATCGCTGTGGC
CCGGCTCTGGCGTGGGAGGCTGAGACACGGAGTGCATCTACTACAA
CGCCA~~ACTGGGAGCTGGAGCGCACCAACCAGAGCGGCCTGGAGCGCTGC~~
GAAGGCGAGCAGGACAAGCGGCTGCACTGCTACGCC~~CTGGCCAACA~~
GCTCTGGCACCATCGAGCTCGTGAAGAAGGGCTGCTGGCTAGATGACTT
CAACTGCTACGATAGGCAGGAGTGTGTGGCCACTGAGGAGAACCCCCAG
GTGTACTTCTGCTGTGAAGGCAACTTCTGCAACGAGCGCTCACTC
15 ATTTGCCAGAGGCTGGGGGCCCGAAGTCACGTACGAGCCACCCCCGAC
AGCCCCCACCTGCTCACGGTGTGGC~~CTACTCACTGCTGCCATCGG~~
GGCCTTCCCTCATCGT~~CCGTGGCCTTTGGATGTACCGGCATCGCA~~
AGCCCCC~~CTACGGTCATGTGGACATCCATGAGGACCCTGGCCTCCACC~~
ACCATCCC~~CTCTGGTGGGCCGAAGCCACTGCAGCTGCTGGAGATCAAG~~
20 GCTGGGGCGCTTGGCTGTGTGGAAAGGCCAGCTCATGAATGACT
TTGTAGCTGTCAAGATCTCCACTCCAGGACAAGCAGTCGTGGCAGAG
TGAACGGGAGATCTTCAGCACACCTGGCATGAAGCACGAGAACCTGCTA
CAGTTCAT~~GCTGCCGAGAAGCGAGGCTCCAACCTCGAAGTAGAGCTGT~~
GGCTCATCACGGC~~CTTCCATGACAAGGGCTCCCTACGGATTACCTCAA~~
25 GGGGAACATCATCACATGGAACGA~~CTGTGT~~CATGTAGCAGAGACGATG
TCACGAGGC~~CTCTCATACCTGCATGAGGATGTGCCCTGGTGCCTGGCG~~
AGGGCCACAAGCCGTCTATTGCC~~ACAGGGACTTTAAAGTAAGAATGT~~
ATTGCTGAAGAGCGACCTCACAGCCGTGCTGGCTGACTTGGCTGGCT
30 GTTCGATTTGAGCCAGGGAAACCTCCAGGGACACCCACGGACAGGTAG
GCACGAGACGGTACATGGCTCTGAGGTGCTCGAGGGAGCCATCAACTT
CCAGAGAGATGCCTTCC~~TGCGATTGACATGTATGCCATGGGTTGGT~~
CTGTGGAGCTTGTCTCGCTGCAAGGCTGCAGACGGACCCGTGGATG

AGTACATGCTGCCCTTGAGGAAGAGATTGCCAGCACCCCTCGTTGGA
GGAGCTGCAGGAGGTGGTGGTGCACAAGAAGATGAGGCCACCATTAAA
GATCACTGGTTGAAACACCCGGGCCTGGCCCAGCTTGTGTGACCATCG
AGGAGTGCTGGGACCATGATGCAGAGGCTCGCTGTCCGCGGGCTGTGT
5 GGAGGAGCGGGTGTCCCTGATTGGAGGTGGTCAACGGCACTACCTCG
GACTGTCTCGTTCCCTGGTGACCTCTGTACCAATGTGGACCTGCC
CTAAAGAGTCAAGCATCTAA (SEQ ID NO: 18)

The nucleic acid sequence encoding a human ActRIIa soluble (extracellular) polypeptide is as follows:

10 TCTGGCGTGGGGAGGCTGAGACACGGGAGTGCATCTACTACAACGCCA
ACTGGGAGCTGGAGCGCACCAACCAGAGCGGCCTGGAGCGCTGCGAAGG
CGAGCAGGACAAGCGGCTGCACTGCTACGCCCTCTGGCCAACAGCTCT
GGCACCATCGAGCTCGTGAAGAAGGGCTGCTGGCTAGATGACTTCAACT
GCTACGATAGGCAGGAGTGTGTGGCCACTGAGGAGAACCCCCAGGTGTA
15 CTTCTGCTGCTGTGAAGGCAACTTCTGCAACGAGCGCTTCACTCATTG
CCAGAGGCTGGGGGCCCGAAGTCACGTACGAGCCACCCCGACAGCCC
CCACC (SEQ ID NO: 19)

In a specific embodiment, the invention relates to soluble ActRII polypeptides. As 20 described herein, the term "soluble ActRII polypeptide" generally refers to polypeptides comprising an extracellular domain of an ActRIIa or ActRIIb protein. The term "soluble ActRII polypeptide," as used herein, includes any naturally occurring extracellular domain of an ActRIIa or ActRIIb protein as well as any variants thereof (including mutants, fragments and peptidomimetic forms). An activin-binding ActRII polypeptide is one that retains the 25 ability to bind to activin, including, for example, activin AA, AB, BB, or forms that include a C or E subunit. Optionally, an activin-binding ActRII polypeptide will bind to activin AA with a dissociation constant of 1 nM or less. The extracellular domain of an ActRII protein binds to activin and is generally soluble, and thus can be termed a soluble, activin-binding ActRII polypeptide. Examples of soluble, activin-binding ActRIIa polypeptides include the 30 soluble polypeptides illustrated in SEQ ID NOS: 2, 3, 7, 12 and 13. SEQ ID NO:7 is referred to as ActRIIa-hFc, and is described further in the Examples. Other examples of soluble, activin-binding ActRIIa polypeptides comprise a signal sequence in addition to the

extracellular domain of an ActRIIa protein, for example, the honey bee mellitin leader sequence (SEQ ID NO: 8), the tissue plaminogen activator (TPA) leader (SEQ ID NO: 9) or the native ActRIIa leader (SEQ ID NO: 10). The ActRIIa-hFc polypeptide illustrated in SEQ ID NO:13 uses a TPA leader. Examples of soluble, activin-binding ActRIIb polypeptides 5 include the soluble polypeptides illustrated in SEQ ID NOs: 16, 17, 20. Activin-binding ActRIIb polypeptides may also comprise a signal sequence in addition to the extracellular domain of an ActRIIb protein, for example, the honey bee mellitin leader sequence (SEQ ID NO: 8) or the tissue plaminogen activator (TPA) leader (SEQ ID NO: 9).

Functionally active fragments of ActRII polypeptides can be obtained by screening 10 polypeptides recombinantly produced from the corresponding fragment of the nucleic acid encoding an ActRII polypeptide. In addition, fragments can be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. The fragments can be produced (recombinantly or by chemical synthesis) and tested to identify those peptidyl fragments that can function as antagonists (inhibitors) of 15 ActRII protein or signaling mediated by activin.

Functionally active variants of ActRII polypeptides can be obtained by screening 20 libraries of modified polypeptides recombinantly produced from the corresponding mutagenized nucleic acids encoding an ActRII polypeptide. The variants can be produced and tested to identify those that can function as antagonists (inhibitors) of ActRII protein or signaling mediated by activin. In certain embodiments, a functional variant of the ActRIIa polypeptides comprises an amino acid sequence that is at least 75% identical to an amino acid sequence selected from SEQ ID NOs: 2 or 3. In certain cases, the functional variant has an amino acid sequence at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from SEQ ID NOs: 2 or 3. In certain embodiments, a 25 functional variant of the ActRIIb polypeptides comprises an amino acid sequence that is at least 75% identical to an amino acid sequence selected from SEQ ID NOs: 16 or 17. In certain cases, the functional variant has an amino acid sequence at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from SEQ ID NOs: 17 or 18.

30 Functional variants may be generated by modifying the structure of an ActRII polypeptide for such purposes as enhancing therapeutic efficacy, or stability (e.g., ex vivo

shelf life and resistance to proteolytic degradation *in vivo*). Such modified ActRII polypeptides when selected to retain activin binding, are considered functional equivalents of the naturally-occurring ActRII polypeptides. Modified ActRII polypeptides can also be produced, for instance, by amino acid substitution, deletion, or addition. For instance, it is 5 reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid (e.g., conservative mutations) will not have a major effect on the biological activity of the resulting molecule. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. 10 Whether a change in the amino acid sequence of an ActRII polypeptide results in a functional homolog can be readily determined by assessing the ability of the variant ActRII polypeptide to produce a response in cells in a fashion similar to the wild-type ActRII polypeptide.

In certain embodiments, the present invention contemplates specific mutations of the ActRII polypeptides so as to alter the glycosylation of the polypeptide. Such mutations may 15 be selected so as to introduce or eliminate one or more glycosylation sites, such as O-linked or N-linked glycosylation sites. Asparagine-linked glycosylation sites generally comprise a tripeptide sequence, asparagine-X-threonine or asparagine-X-serine (where "X" is any amino acid) which is specifically recognized by appropriate cellular glycosylation enzymes. The alteration may also be made by the addition of, or substitution by, one or more 20 serine or threonine residues to the sequence of the wild-type ActRII polypeptide (for O-linked glycosylation sites). A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) results in non-glycosylation at the modified tripeptide sequence. Another means of increasing the number of carbohydrate moieties on an ActRII 25 polypeptide is by chemical or enzymatic coupling of glycosides to the ActRII polypeptide. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine; (b) free carboxyl groups; (c) free sulfhydryl groups such as those of cysteine; (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline; (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan; or (f) the amide group of 30 glutamine. Removal of one or more carbohydrate moieties present on an ActRII polypeptide may be accomplished chemically and/or enzymatically. Chemical deglycosylation may involve, for example, exposure of the ActRII polypeptide to the compound

trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the amino acid sequence intact. Enzymatic cleavage of carbohydrate moieties on ActRII polypeptides can be achieved by the use of a variety of 5 endo- and exo-glycosidases as described by Thotakura et al. (1987) *Meth. Enzymol.* 138:350. The sequence of an ActRII polypeptide may be adjusted, as appropriate, depending on the type of expression system used, as mammalian, yeast, insect and plant cells may all introduce differing glycosylation patterns that can be affected by the amino acid sequence of the peptide. In general, ActRII proteins for use in humans may be expressed in a mammalian cell 10 line that provides proper glycosylation, such as HEK293 or CHO cell lines, although other mammalian expression cell lines are expected to be useful as well.

This disclosure further contemplates a method of generating mutants, particularly sets of combinatorial mutants of an ActRII polypeptide, as well as truncation mutants; pools of combinatorial mutants are especially useful for identifying functional variant sequences. The 15 purpose of screening such combinatorial libraries may be to generate, for example, ActRII polypeptide variants which bind to activin or other ligands. A variety of screening assays are provided below, and such assays may be used to evaluate variants. For example, an ActRII polypeptide variant may be screened for ability to bind to an ActRII ligand, to prevent binding of an ActRII ligand to an ActRII polypeptide or to interfere with signaling caused by 20 an ActRII ligand.

The activity of an ActRII polypeptide or its variants may also be tested in a cell-based or *in vivo* assay. For example, the effect of an ActRII polypeptide variant on the expression of genes involved in hematopoiesis may be assessed. This may, as needed, be performed in the presence of one or more recombinant ActRII ligand proteins (e.g., activin), and cells may 25 be transfected so as to produce an ActRII polypeptide and/or variants thereof, and optionally, an ActRII ligand. Likewise, an ActRII polypeptide may be administered to a mouse or other animal, and one or more blood measurements, such as an RBC count, hemoglobin, or reticulocyte count may be assessed.

Combinatorially-derived variants can be generated which have a selective or generally 30 increased potency relative to a naturally occurring ActRII polypeptide. Likewise, mutagenesis can give rise to variants which have intracellular half-lives dramatically different

than the corresponding a wild-type ActRII polypeptide. For example, the altered protein can be rendered either more stable or less stable to proteolytic degradation or other cellular processes which result in destruction of, or otherwise inactivation of a native ActRII polypeptide. Such variants, and the genes which encode them, can be utilized to alter ActRII 5 polypeptide levels by modulating the half-life of the ActRII polypeptides. For instance, a short half-life can give rise to more transient biological effects and, when part of an inducible expression system, can allow tighter control of recombinant ActRII polypeptide levels within the cell. In an Fc fusion protein, mutations may be made in the linker (if any) and/or the Fc portion to alter the half-life of the protein.

10 A combinatorial library may be produced by way of a degenerate library of genes encoding a library of polypeptides which each include at least a portion of potential ActRII polypeptide sequences. For instance, a mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential ActRII polypeptide nucleotide sequences are expressible as individual polypeptides, or alternatively, 15 as a set of larger fusion proteins (e.g., for phage display).

There are many ways by which the library of potential homologs can be generated from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be carried out in an automatic DNA synthesizer, and the synthetic genes can then be ligated into an appropriate vector for expression. The synthesis of degenerate 20 oligonucleotides is well known in the art (see for example, Narang, SA (1983) *Tetrahedron* 39:3; Itakura et al., (1981) *Recombinant DNA, Proc. 3rd Cleveland Sympos.* Macromolecules, ed. AG Walton, Amsterdam: Elsevier pp273-289; Itakura et al., (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al., (1984) *Science* 198:1056; Ike et al., (1983) *Nucleic Acid Res.* 11:477). Such techniques have been employed in the directed evolution of 25 other proteins (see, for example, Scott et al., (1990) *Science* 249:386-390; Roberts et al., (1992) *PNAS USA* 89:2429-2433; Devlin et al., (1990) *Science* 249: 404-406; Cwirla et al., (1990) *PNAS USA* 87: 6378-6382; as well as U.S. Patent Nos: 5,223,409, 5,198,346, and 5,096,815).

30 Alternatively, other forms of mutagenesis can be utilized to generate a combinatorial library. For example, ActRII polypeptide variants can be generated and isolated from a library by screening using, for example, alanine scanning mutagenesis and the like (Ruf et al.,

(1994) Biochemistry 33:1565-1572; Wang et al., (1994) J. Biol. Chem. 269:3095-3099; Balint et al., (1993) Gene 137:109-118; Grodberg et al., (1993) Eur. J. Biochem. 218:597-601; Nagashima et al., (1993) J. Biol. Chem. 268:2888-2892; Lowman et al., (1991) Biochemistry 30:10832-10838; and Cunningham et al., (1989) Science 244:1081-1085), by 5 linker scanning mutagenesis (Gustin et al., (1993) Virology 193:653-660; Brown et al., (1992) Mol. Cell Biol. 12:2644-2652; McKnight et al., (1982) Science 232:316); by saturation mutagenesis (Meyers et al., (1986) Science 232:613); by PCR mutagenesis (Leung et al., (1989) Method Cell Mol Biol 1:11-19); or by random mutagenesis, including chemical mutagenesis, etc. (Miller et al., (1992) A Short Course in Bacterial Genetics, CSHL Press, 10 Cold Spring Harbor, NY; and Greener et al., (1994) Strategies in Mol Biol 7:32-34). Linker scanning mutagenesis, particularly in a combinatorial setting, is an attractive method for identifying truncated (bioactive) forms of ActRII polypeptides.

A wide range of techniques are known in the art for screening gene products of combinatorial libraries made by point mutations and truncations, and, for that matter, for 15 screening cDNA libraries for gene products having a certain property. Such techniques will be generally adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of ActRII polypeptides. The most widely used techniques for screening large gene libraries typically comprises cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and 20 expressing the combinatorial genes under conditions in which detection of a desired activity facilitates relatively easy isolation of the vector encoding the gene whose product was detected. Preferred assays include activin binding assays and activin-mediated cell signaling assays.

In certain embodiments, the ActRII polypeptides of the invention may further 25 comprise post-translational modifications in addition to any that are naturally present in the ActRII polypeptides. Such modifications include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. As a result, the modified ActRII polypeptides may contain non-amino acid elements, such as polyethylene glycols, lipids, poly- or mono-saccharide, and phosphates. Effects of such non-amino acid 30 elements on the functionality of an ActRII polypeptide may be tested as described herein for other ActRII polypeptide variants. When an ActRII polypeptide is produced in cells by

cleaving a nascent form of the ActRII polypeptide, post-translational processing may also be important for correct folding and/or function of the protein. Different cells (such as CHO, HeLa, MDCK, 293, WI38, NIH-3T3 or HEK293) have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure
5 the correct modification and processing of the ActRII polypeptides.

In certain aspects, functional variants or modified forms of the ActRII polypeptides include fusion proteins having at least a portion of the ActRII polypeptides and one or more fusion domains. Well known examples of such fusion domains include, but are not limited to, polyhistidine, Glu-Glu, glutathione S transferase (GST), thioredoxin, protein A, protein G,
10 an immunoglobulin heavy chain constant region (Fc), maltose binding protein (MBP), or human serum albumin. A fusion domain may be selected so as to confer a desired property. For example, some fusion domains are particularly useful for isolation of the fusion proteins by affinity chromatography. For the purpose of affinity purification, relevant matrices for affinity chromatography, such as glutathione-, amylase-, and nickel- or cobalt- conjugated
15 resins are used. Many of such matrices are available in "kit" form, such as the Pharmacia GST purification system and the QIAexpress™ system (Qiagen) useful with (HIS₆) fusion partners. As another example, a fusion domain may be selected so as to facilitate detection of the ActRII polypeptides. Examples of such detection domains include the various fluorescent proteins (e.g., GFP) as well as "epitope tags," which are usually short peptide sequences for
20 which a specific antibody is available. Well known epitope tags for which specific monoclonal antibodies are readily available include FLAG, influenza virus haemagglutinin (HA), and c-myc tags. In some cases, the fusion domains have a protease cleavage site, such as for Factor Xa or Thrombin, which allows the relevant protease to partially digest the fusion proteins and thereby liberate the recombinant proteins therefrom. The liberated proteins can
25 then be isolated from the fusion domain by subsequent chromatographic separation. In certain preferred embodiments, an ActRII polypeptide is fused with a domain that stabilizes the ActRII polypeptide in vivo (a "stabilizer" domain). By "stabilizing" is meant anything that increases serum half life, regardless of whether this is because of decreased destruction, decreased clearance by the kidney, or other pharmacokinetic effect. Fusions with the Fc
30 portion of an immunoglobulin are known to confer desirable pharmacokinetic properties on a wide range of proteins. Likewise, fusions to human serum albumin can confer desirable properties. Other types of fusion domains that may be selected include multimerizing (e.g.,

dimerizing, tetramerizing) domains and functional domains (that confer an additional biological function, such as further stimulation of muscle growth).

As a specific example, the present invention provides a fusion protein comprising a soluble extracellular domain of ActRIIa fused to an Fc domain (e.g., SEQ ID NO: 6).

5 THTCPPCPAPELLGGPSVFLFPPPKD~~T~~LMISRTPEVTCVVVD (A) VS~~H~~EDPEVKF~~N~~WYV~~D~~G VEVHNAKTKP~~R~~EEQYNSTYRVVSV~~L~~T~~V~~LHQDW~~L~~NGKEYKCK (A) VSNKALPVPIEKTISKAK GQP~~R~~PQVYTL~~P~~PSREEMTKNQVSLTCLVKG~~F~~YPSDIAVEWESNGQ~~P~~ENNYK~~T~~TPV~~L~~SDG PFFLYSKLTVD~~K~~SRWQQGNVF~~S~~CSVMHEALHN (A) HYTQKSL~~S~~LS~~P~~PGK*

As an additional specific example, the present invention provides a fusion protein 10 comprising a soluble extracellular domain of ActRIIb fused to an Fc domain (e.g., SEQ ID NO: 21).

SGRGEAETREC~~I~~YYNANWELERTNQ~~S~~GLERCEGEQDKRLHCYASWANSSGTIELVKKGCWLD DFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLPEAGGPEV~~T~~YEPPPTAPTGGGHTCP PCPAP~~E~~LLGGPSVFLFPPPKD~~T~~LMISRTPEVTCVVVD~~S~~HEDPEVKF~~N~~WYV~~D~~GVEVHNAK~~T~~ K~~P~~REEQYNSTYRVVSV~~L~~T~~V~~LHQDW~~L~~NGKEYKCKVSNKALPVPIEKTISKAKGQ~~P~~REPQVYTL PPSREEMTKNQVSLTCLVKG~~F~~YPSDIAVEWESNGQ~~P~~ENNYK~~T~~TPV~~L~~SDG~~S~~FFLYSKLTVD KSRWQQGNVF~~S~~CSVMHEALHNHYTQKSL~~S~~LS~~P~~PGK

20 Optionally, the Fc domain has one or more mutations at residues such as Asp-265, lysine 322, and Asn-434. In certain cases, the mutant Fc domain having one or more of these mutations (e.g., Asp-265 mutation) has reduced ability of binding to the Fcγ receptor relative to a wildtype Fc domain. In other cases, the mutant Fc domain having one or more of these mutations (e.g., Asn-434 mutation) has increased ability of binding to the MHC class I-related Fc-receptor (FcRN) relative to a wildtype Fc domain.

25 It is understood that different elements of the fusion proteins may be arranged in any manner that is consistent with the desired functionality. For example, an ActRII polypeptide may be placed C-terminal to a heterologous domain, or, alternatively, a heterologous domain may be placed C-terminal to an ActRII polypeptide. The ActRII polypeptide domain and the heterologous domain need not be adjacent in a fusion protein, and additional domains or amino acid sequences may be included C- or N-terminal to either domain or between the 30 domains.

In certain embodiments, the ActRII polypeptides of the present invention contain one or more modifications that are capable of stabilizing the ActRII polypeptides. For example, such modifications enhance the in vitro half life of the ActRII polypeptides, enhance circulatory half life of the ActRII polypeptides or reducing proteolytic degradation of the 5 ActRII polypeptides. Such stabilizing modifications include, but are not limited to, fusion proteins (including, for example, fusion proteins comprising an ActRII polypeptide and a stabilizer domain), modifications of a glycosylation site (including, for example, addition of a glycosylation site to an ActRII polypeptide), and modifications of carbohydrate moiety (including, for example, removal of carbohydrate moieties from an ActRII polypeptide). As 10 used herein, the term "stabilizer domain" not only refers to a fusion domain (e.g., Fc) as in the case of fusion proteins, but also includes nonproteinaceous modifications such as a carbohydrate moiety, or nonproteinaceous moiety, such as polyethylene glycol.

In certain embodiments, the present invention makes available isolated and/or purified forms of the ActRII polypeptides, which are isolated from, or otherwise substantially free of, 15 other proteins. ActRII polypeptides will generally be produced by expression from recombinant nucleic acids.

3. Nucleic Acids Encoding ActRII Polypeptides

In certain aspects, the invention provides isolated and/or recombinant nucleic acids 20 encoding any of the ActRII polypeptides (e.g., soluble ActRIIa polypeptides and soluble ActRIIb polypeptides), including fragments, functional variants and fusion proteins disclosed herein. For example, SEQ ID NO: 4 encodes the naturally occurring human ActRIIa precursor polypeptide, while SEQ ID NO: 5 encodes the processed extracellular domain of 25 ActRIIa. For example, SEQ ID NO: 18 encodes the naturally occurring human ActRIIb precursor polypeptide, while SEQ ID NO: 19 encodes the processed extracellular domain of ActRIIb. The subject nucleic acids may be single-stranded or double stranded. Such nucleic acids may be DNA or RNA molecules. These nucleic acids may be used, for example, in methods for making ActRII polypeptides or as direct therapeutic agents (e.g., in a gene 30 therapy approach).

In certain aspects, the subject nucleic acids encoding ActRIIa polypeptides are further understood to include nucleic acids that are variants of SEQ ID NO: 4 or 5. In certain

aspects, the subject nucleic acids encoding ActRIIb polypeptides are further understood to include nucleic acids that are variants of SEQ ID NO: 18 or 19. Variant nucleotide sequences include sequences that differ by one or more nucleotide substitutions, additions or deletions, such as allelic variants.

5 In certain embodiments, the invention provides isolated or recombinant nucleic acid sequences that are at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to SEQ ID NOs: 4, 5, 18, or 19. One of ordinary skill in the art will appreciate that nucleic acid sequences complementary to SEQ ID NOs: 4, 5, 18, or 19, and variants of SEQ ID NOs: 4, 5, 18, or 19 are also within the scope of this invention. In further embodiments, the nucleic acid 10 sequences of the invention can be isolated, recombinant, and/or fused with a heterologous nucleotide sequence, or in a DNA library.

In other embodiments, nucleic acids of the invention also include nucleotide sequences that hybridize under highly stringent conditions to the nucleotide sequence designated in SEQ ID NOs: 4, 5, 18, or 19, complement sequence of SEQ ID NOs: 4, 5, 18, 15 or 19, or fragments thereof. As discussed above, one of ordinary skill in the art will understand readily that appropriate stringency conditions which promote DNA hybridization can be varied. One of ordinary skill in the art will understand readily that appropriate stringency conditions which promote DNA hybridization can be varied. For example, one could perform the hybridization at 6.0 x sodium chloride/sodium citrate (SSC) at about 45 °C, 20 followed by a wash of 2.0 x SSC at 50 °C. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50 °C to a high stringency of about 0.2 x SSC at 50 °C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22 °C, to high stringency conditions at about 65 °C. Both temperature and salt may be varied, or temperature or salt concentration 25 may be held constant while the other variable is changed. In one embodiment, the invention provides nucleic acids which hybridize under low stringency conditions of 6 x SSC at room temperature followed by a wash at 2 x SSC at room temperature.

Isolated nucleic acids which differ from the nucleic acids as set forth in SEQ ID NOs: 4, 5, 18, or 19 due to degeneracy in the genetic code are also within the scope of the 30 invention. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid, or synonyms (for example, CAU and CAC are

synonyms for histidine) may result in “silent” mutations which do not affect the amino acid sequence of the protein. However, it is expected that DNA sequence polymorphisms that do lead to changes in the amino acid sequences of the subject proteins will exist among 5 mammalian cells. One skilled in the art will appreciate that these variations in one or more nucleotides (up to about 3-5% of the nucleotides) of the nucleic acids encoding a particular protein may exist among individuals of a given species due to natural allelic variation. Any and all such nucleotide variations and resulting amino acid polymorphisms are within the scope of this invention.

In certain embodiments, the recombinant nucleic acids of the invention may be 10 operably linked to one or more regulatory nucleotide sequences in an expression construct. Regulatory nucleotide sequences will generally be appropriate to the host cell used for expression. Numerous types of appropriate expression vectors and suitable regulatory sequences are known in the art for a variety of host cells. Typically, said one or more regulatory nucleotide sequences may include, but are not limited to, promoter sequences, 15 leader or signal sequences, ribosomal binding sites, transcriptional start and termination sequences, translational start and termination sequences, and enhancer or activator sequences. Constitutive or inducible promoters as known in the art are contemplated by the invention. The promoters may be either naturally occurring promoters, or hybrid promoters that combine elements of more than one promoter. An expression construct may be present in a 20 cell on an episome, such as a plasmid, or the expression construct may be inserted in a chromosome. In a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selectable marker genes are well known in the art and will vary with the host cell used.

In certain aspects of the invention, the subject nucleic acid is provided in an 25 expression vector comprising a nucleotide sequence encoding an ActRII polypeptide and operably linked to at least one regulatory sequence. Regulatory sequences are art-recognized and are selected to direct expression of the ActRII polypeptide. Accordingly, the term regulatory sequence includes promoters, enhancers, and other expression control elements. Exemplary regulatory sequences are described in Goeddel; *Gene Expression Technology*: 30 *Methods in Enzymology*, Academic Press, San Diego, CA (1990). For instance, any of a wide variety of expression control sequences that control the expression of a DNA sequence when

operatively linked to it may be used in these vectors to express DNA sequences encoding an ActRII polypeptide. Such useful expression control sequences, include, for example, the early and late promoters of SV40, tet promoter, adenovirus or cytomegalovirus immediate early promoter, RSV promoters, the lac system, the *trp* system, the TAC or TRC system, T7 5 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage lambda, the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α -mating factors, the polyhedron promoter of the 10 baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. Moreover, the vector's copy number, the ability to control that copy number and the 15 expression of any other protein encoded by the vector, such as antibiotic markers, should also be considered.

A recombinant nucleic acid of the invention can be produced by ligating the cloned gene, or a portion thereof, into a vector suitable for expression in either prokaryotic cells, eukaryotic cells (yeast, avian, insect or mammalian), or both. Expression vehicles for 20 production of a recombinant ActRII polypeptide include plasmids and other vectors. For instance, suitable vectors include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

Some mammalian expression vectors contain both prokaryotic sequences to facilitate 25 the propagation of the vector in bacteria, and one or more eukaryotic transcription units that are expressed in eukaryotic cells. The pcDNA1/amp, pcDNA1/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. 30 Alternatively, derivatives of viruses such as the bovine papilloma virus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins

in eukaryotic cells. Examples of other viral (including retroviral) expression systems can be found below in the description of gene therapy delivery systems. The various methods employed in the preparation of the plasmids and in transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic 5 cells, as well as general recombinant procedures, see *Molecular Cloning A Laboratory Manual*, 3rd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press, 2001). In some instances, it may be desirable to express the recombinant polypeptides by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), 10 pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the β -gal containing pBlueBac III).

15 In a preferred embodiment, a vector will be designed for production of the subject ActRII polypeptides in CHO cells, such as a Pcmv-Script vector (Stratagene, La Jolla, Calif.), pcDNA4 vectors (Invitrogen, Carlsbad, Calif.) and pCI-neo vectors (Promega, Madison, Wisc.). As will be apparent, the subject gene constructs can be used to cause expression of the subject ActRII polypeptides in cells propagated in culture, e.g., to produce proteins, including fusion proteins or variant proteins, for purification.

20 This disclosure also pertains to a host cell transfected with a recombinant gene including a coding sequence (e.g., SEQ ID NO: 4, 5, 18, or 19) for one or more of the subject ActRII polypeptides. The host cell may be any prokaryotic or eukaryotic cell. For example, an ActRII polypeptide of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (e.g., using a baculovirus expression system), yeast, or mammalian cells. Other suitable host cells are known to those skilled in the art.

25 Accordingly, the present invention further pertains to methods of producing the subject ActRII polypeptides. For example, a host cell transfected with an expression vector encoding an ActRIIa or ActRIIb polypeptide can be cultured under appropriate conditions to allow expression of the ActRII polypeptide to occur. The ActRII polypeptide may be secreted and isolated from a mixture of cells and medium containing the ActRII polypeptide. Alternatively, the ActRII polypeptide may be retained cytoplasmically or in a membrane 30 fraction and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art.

The subject ActRII polypeptides can be isolated from cell culture medium, host cells, or both, using techniques known in the art for purifying proteins, including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, immunoaffinity purification with antibodies specific for particular epitopes of the ActRII polypeptides and affinity purification with an agent that binds to a domain fused to the ActRII polypeptide (e.g., a protein A column may be used to purify an ActRIIa-Fc or ActRIIb-Fc fusion). In a preferred embodiment, the ActRII polypeptide is a fusion protein containing a domain which facilitates its purification. In a preferred embodiment, purification is achieved by a series of column chromatography steps, including, for example, 5 three or more of the following, in any order: protein A chromatography, Q sepharose chromatography, phenylsepharose chromatography, size exclusion chromatography, and cation exchange chromatography. The purification could be completed with viral filtration and buffer exchange. As demonstrated herein, ActRIIa-hFc protein was purified to a purity of >98% as determined by size exclusion chromatography and >95% as determined by SDS 10 PAGE. This level of purity was sufficient to achieve desirable results in mice, rats and non-human primates.

15

In another embodiment, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence at the N-terminus of the desired portion of the recombinant ActRII polypeptide, can allow purification of the expressed fusion protein 20 by affinity chromatography using a Ni²⁺ metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase to provide the purified ActRII polypeptide (e.g., see Hochuli et al., (1987) *J. Chromatography* 411:177; and Janknecht et al., *PNAS USA* 88:8972).

Techniques for making fusion genes are well known. Essentially, the joining of 25 various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by 30 conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to

complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al., John Wiley & Sons: 1992).

5 4. Alternative Activin and ActRII Antagonists

The data presented herein demonstrates that antagonists of activin-ActRII signaling can be used to increase red blood cell or hemoglobin levels. Although soluble ActRIIa and ActRIIb polypeptides, and particularly ActRIIa-Fc and ActRIIb-Fc, are preferred antagonists, and although such antagonists may affect red blood cell levels through a mechanism other than activin antagonism (e.g., activin inhibition may be an indicator of the tendency of an agent to inhibit the activities of a spectrum of molecules, including, perhaps, other members of the TGF-beta superfamily, and such collective inhibition may lead to the desired effect on hematopoiesis), other types of activin-ActRII antagonists are expected to be useful, including anti-activin (e.g., activin β_A , β_B , β_C and β_E antibodies, anti-ActRIIa antibodies, anti-ActRIIb antibodies, antisense, RNAi or ribozyme nucleic acids that inhibit the production of ActRIIa and/or ActRIIb, and other inhibitors of activin, ActRIIb or ActRIIa, particularly those that disrupt activin-ActRIIa and/or activin-ActRIIb binding.

An antibody that is specifically reactive with an ActRII polypeptide (e.g., a soluble ActRIIa or ActRIIb polypeptide) and which either binds competitively to ligand with the ActRII polypeptide or otherwise inhibits ActRII-mediated signaling may be used as an antagonist of ActRII polypeptide activities. Likewise, an antibody that is specifically reactive with an activin β_A , β_B , β_C or β_E polypeptide, or any heterodimer thereof, and which disrupts ActRIIa and/or ActRIIb binding may be used as an antagonist.

By using immunogens derived from an ActRIIa polypeptide, ActRIIb polypeptide or an activin polypeptide, anti-protein/anti-peptide antisera or monoclonal antibodies can be made by standard protocols (see, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)). A mammal, such as a mouse, a hamster or rabbit can be immunized with an immunogenic form of the activin, ActRIIa or ActRIIb polypeptide, an antigenic fragment which is capable of eliciting an antibody response, or a fusion protein. Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques well known in the art. An immunogenic portion of

an ActRII or activin polypeptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies.

5 Following immunization of an animal with an antigenic preparation of an activin, ActRIIa or ActRIIb polypeptide, antisera can be obtained and, if desired, polyclonal antibodies can be isolated from the serum. To produce monoclonal antibodies, antibody-producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to 10 yield hybridoma cells. Such techniques are well known in the art, and include, for example, the hybridoma technique (originally developed by Kohler and Milstein, (1975) *Nature*, 256: 495-497), the human B cell hybridoma technique (Kozbar et al., (1983) *Immunology Today*, 4: 72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. pp. 77-96).

15 Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with an activin, ActRIIa or ActRIIb polypeptide and monoclonal antibodies isolated from a culture comprising such hybridoma cells.

The term "antibody" as used herein is intended to include whole antibodies, e.g., of any isotype (IgG, IgA, IgM, IgE, etc), and includes fragments or domains of 20 immunoglobulins which are reactive with a selected antigen. Antibodies can be fragmented using conventional techniques and the fragments screened for utility and/or interaction with a specific epitope of interest. Thus, the term includes segments of proteolytically-cleaved or recombinantly-prepared portions of an antibody molecule that are capable of selectively reacting with a certain protein. Non-limiting examples of such proteolytic and/or recombinant 25 fragments include Fab, F(ab')2, Fab', Fv, and single chain antibodies (scFv) containing a V[L] and/or V[H] domain joined by a peptide linker. The scFv's may be covalently or non-covalently linked to form antibodies having two or more binding sites. The term antibody also includes polyclonal, monoclonal, or other purified preparations of antibodies and 30 recombinant antibodies. The term "recombinant antibody", means an antibody, or antigen binding domain of an immunoglobulin, expressed from a nucleic acid that has been constructed using the techniques of molecular biology, such as a humanized antibody or a

fully human antibody developed from a single chain antibody. Single domain and single chain antibodies are also included within the term “recombinant antibody”.

In certain embodiments, an antibody of the invention is a monoclonal antibody, and in certain embodiments, the invention makes available methods for generating novel antibodies.

5 For example, a method for generating a monoclonal antibody that binds specifically to an ActRIIa polypeptide, ActRIIb polypeptide, or activin polypeptide may comprise administering to a mouse an amount of an immunogenic composition comprising the antigen polypeptide effective to stimulate a detectable immune response, obtaining antibody-producing cells (e.g., cells from the spleen) from the mouse and fusing the antibody-producing cells with myeloma cells to obtain antibody-producing hybridomas, and testing the antibody-producing hybridomas to identify a hybridoma that produces a monoclonal antibody that binds specifically to the antigen. Once obtained, a hybridoma can be propagated in a cell culture, optionally in culture conditions where the hybridoma-derived cells produce the monoclonal antibody that binds specifically to the antigen. The monoclonal antibody may be purified from the cell culture.

10

15

The adjective “specifically reactive with” as used in reference to an antibody is intended to mean, as is generally understood in the art, that the antibody is sufficiently selective between the antigen of interest (e.g., an activin, ActRIIa or ActRIIb polypeptide) and other antigens that are not of interest that the antibody is useful for, at minimum, 20 detecting the presence of the antigen of interest in a particular type of biological sample. In certain methods employing the antibody, such as therapeutic applications, a higher degree of specificity in binding may be desirable. Monoclonal antibodies generally have a greater tendency (as compared to polyclonal antibodies) to discriminate effectively between the desired antigens and cross-reacting polypeptides. One characteristic that influences the 25 specificity of an antibody:antigen interaction is the affinity of the antibody for the antigen. Although the desired specificity may be reached with a range of different affinities, generally preferred antibodies will have an affinity (a dissociation constant) of about 10^{-6} , 10^{-7} , 10^{-8} , 10^{-9} M or less.

In addition, the techniques used to screen antibodies in order to identify a desirable 30 antibody may influence the properties of the antibody obtained. For example, if an antibody is to be used for binding an antigen in solution, it may be desirable to test solution binding. A

variety of different techniques are available for testing interaction between antibodies and antigens to identify particularly desirable antibodies. Such techniques include ELISAs, surface plasmon resonance binding assays (e.g., the BiacoreTM binding assay, Biacore AB, Uppsala, Sweden), sandwich assays (e.g., the paramagnetic bead system of IGEN International, Inc., Gaithersburg, Maryland), western blots, immunoprecipitation assays, and immunohistochemistry.

5 Examples of categories of nucleic acid compounds that are activin or ActRII antagonists include antisense nucleic acids, RNAi constructs and catalytic nucleic acid constructs. A nucleic acid compound may be single or double stranded. A double stranded compound may also include regions of overhang or non-complementarity, where one or the other of the strands is single stranded. A single stranded compound may include regions of 10 self-complementarity, meaning that the compound forms a so-called “hairpin” or “stem-loop” structure, with a region of double helical structure. A nucleic acid compound may comprise a nucleotide sequence that is complementary to a region consisting of no more than 1000, no more than 500, no more than 250, no more than 100, or no more than 50, 35, 25, 22, 20, 18 or 15 nucleotides of the full-length ActRII nucleic acid sequence or activin β_A , β_B , β_C , or β_E nucleic acid sequence. The region of complementarity will preferably be at least 8 nucleotides, and optionally about 18 to 35 nucleotides. A region of complementarity may fall within an intron, a coding sequence or a noncoding sequence of the target transcript, such as 15 the coding sequence portion. Generally, a nucleic acid compound will have a length of about 20 8 to about 500 nucleotides or base pairs in length, and optionally the length will be about 14 to about 50 nucleotides. A nucleic acid may be a DNA (particularly for use as an antisense), RNA or RNA:DNA hybrid. Any one strand may include a mixture of DNA and RNA, as well as modified forms that cannot readily be classified as either DNA or RNA. Likewise, a 25 double stranded compound may be DNA:DNA, DNA:RNA or RNA:RNA, and any one strand may also include a mixture of DNA and RNA, as well as modified forms that cannot readily be classified as either DNA or RNA. A nucleic acid compound may include any of a variety of modifications, including one or modifications to the backbone (the sugar-phosphate portion in a natural nucleic acid, including internucleotide linkages) or the base portion (the 30 purine or pyrimidine portion of a natural nucleic acid). An antisense nucleic acid compound will preferably have a length of about 15 to about 30 nucleotides and will often contain one or more modifications to improve characteristics such as stability in the serum, in a cell or in

a place where the compound is likely to be delivered, such as the stomach in the case of orally delivered compounds and the lung for inhaled compounds. In the case of an RNAi construct, the strand complementary to the target transcript will generally be RNA or modifications thereof. The other strand may be RNA, DNA or any other variation. The 5 duplex portion of double stranded or single stranded "hairpin" RNAi construct will generally have a length of 18 to 40 nucleotides in length and optionally about 21 to 23 nucleotides in length, so long as it serves as a Dicer substrate. Catalytic or enzymatic nucleic acids may be ribozymes or DNA enzymes and may also contain modified forms. Nucleic acid compounds may inhibit expression of the target by about 50%, 75%, 90% or more when contacted with 10 cells under physiological conditions and at a concentration where a nonsense or sense control has little or no effect. Preferred concentrations for testing the effect of nucleic acid compounds are 1, 5 and 10 micromolar. Nucleic acid compounds may also be tested for effects on, for example, red blood cell levels.

15 5. Screening Assays

In certain aspects, the present invention relates to the use of ActRII polypeptides (e.g., soluble ActRIIa or ActRIIb polypeptides) and activin polypeptides to identify compounds (agents) which are agonist or antagonists of the activin-ActRIIa and/or activin ActRIIb signaling pathway. Compounds identified through this screening can be tested to assess their 20 ability to modulate red blood cell, hemoglobin and/or reticulocyte levels in vivo or in vitro. These compounds can be tested, for example, in animal models.

There are numerous approaches to screening for therapeutic agents for increasing red blood cell or hemoglobin levels by targeting activin and ActRII signaling. In certain embodiments, high-throughput screening of compounds can be carried out to identify agents 25 that perturb activin or ActRII-mediated effects on a selected cell line. In certain embodiments, the assay is carried out to screen and identify compounds that specifically inhibit or reduce binding of an ActRIIa or ActRIIb polypeptide to activin. Alternatively, the assay can be used to identify compounds that enhance binding of an ActRIIa or ActRIIb polypeptide to activin. In a further embodiment, the compounds can be identified by their 30 ability to interact with an activin, ActRIIb polypeptide, or ActRIIa polypeptide.

A variety of assay formats will suffice and, in light of the present disclosure, those not expressly described herein will nevertheless be comprehended by one of ordinary skill in the art. As described herein, the test compounds (agents) of the invention may be created by any combinatorial chemical method. Alternatively, the subject compounds may be naturally occurring biomolecules synthesized in vivo or in vitro. Compounds (agents) to be tested for their ability to act as modulators of tissue growth can be produced, for example, by bacteria, yeast, plants or other organisms (e.g., natural products), produced chemically (e.g., small molecules, including peptidomimetics), or produced recombinantly. Test compounds contemplated by the present invention include non-peptidyl organic molecules, peptides, polypeptides, peptidomimetics, sugars, hormones, and nucleic acid molecules. In a specific embodiment, the test agent is a small organic molecule having a molecular weight of less than about 2,000 Daltons.

The test compounds of the invention can be provided as single, discrete entities, or provided in libraries of greater complexity, such as made by combinatorial chemistry. These libraries can comprise, for example, alcohols, alkyl halides, amines, amides, esters, aldehydes, ethers and other classes of organic compounds. Presentation of test compounds to the test system can be in either an isolated form or as mixtures of compounds, especially in initial screening steps. Optionally, the compounds may be optionally derivatized with other compounds and have derivatizing groups that facilitate isolation of the compounds. Non-limiting examples of derivatizing groups include biotin, fluorescein, digoxxygenin, green fluorescent protein, isotopes, polyhistidine, magnetic beads, glutathione S transferase (GST), photoactivatable crosslinkers or any combinations thereof.

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Moreover, the effects of cellular toxicity or bioavailability of the test compound can be generally ignored in the in vitro system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding

affinity between an ActRIIa polypeptide and activin and/or between an ActRIIb polypeptide and activin.

Merely to illustrate, in an exemplary screening assay of the present invention, the compound of interest is contacted with an isolated and purified ActRIIa polypeptide which is ordinarily capable of binding to activin. To the mixture of the compound and ActRIIa polypeptide is then added a composition containing an ActRIIa ligand. Detection and quantification of ActRIIa/activin complexes provides a means for determining the compound's efficacy at inhibiting (or potentiating) complex formation between the ActRIIa polypeptide and activin. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound.

Moreover, a control assay can also be performed to provide a baseline for comparison. For example, in a control assay, isolated and purified activin is added to a composition containing the ActRIIa polypeptide, and the formation of ActRIIa/activin complex is quantitated in the absence of the test compound. It will be understood that, in general, the order in which the reactants may be admixed can be varied, and can be admixed simultaneously. Moreover, in place of purified proteins, cellular extracts and lysates may be used to render a suitable cell-free assay system. Compounds that affect ActRIIb signaling may be identified in a similar manner using an ActRIIb polypeptide and an ActRIIb ligand.

Complex formation between the ActRII polypeptide and activin may be detected by a variety of techniques. For instance, modulation of the formation of complexes can be quantitated using, for example, detectably labeled proteins such as radiolabeled (e.g., ³²P, ³⁵S, ¹⁴C or ³H), fluorescently labeled (e.g., FITC), or enzymatically labeled ActRIIa or ActRIIb polypeptide or activin, by immunoassay, or by chromatographic detection.

In certain embodiments, the present invention contemplates the use of fluorescence polarization assays and fluorescence resonance energy transfer (FRET) assays in measuring, either directly or indirectly, the degree of interaction between an ActRII polypeptide and its binding protein. Further, other modes of detection, such as those based on optical waveguides (PCT Publication WO 96/26432 and U.S. Pat. No. 5,677,196), surface plasmon resonance (SPR), surface charge sensors, and surface force sensors, are compatible with many embodiments of the invention.

Moreover, the present invention contemplates the use of an interaction trap assay, also known as the “two hybrid assay,” for identifying agents that disrupt or potentiate interaction between an ActRII polypeptide and its binding protein. See for example, U.S. Pat. No. 5,283,317; Zervos et al. (1993) Cell 72:223-232; Madura et al. (1993) J Biol Chem 268:12046-12054; Bartel et al. (1993) Biotechniques 14:920-924; and Iwabuchi et al. (1993) Oncogene 8:1693-1696). In a specific embodiment, the present invention contemplates the use of reverse two hybrid systems to identify compounds (e.g., small molecules or peptides) that dissociate interactions between an ActRII polypeptide and its binding protein. See for example, Vidal and Legrain, (1999) Nucleic Acids Res 27:919-29; Vidal and Legrain, (1999) Trends Biotechnol 17:374-81; and U.S. Pat. Nos. 5,525,490; 5,955,280; and 5,965,368.

In certain embodiments, the subject compounds are identified by their ability to interact with an ActRII or activin polypeptide of the invention. The interaction between the compound and the ActRIIa, ActRIIb, or activin polypeptide may be covalent or non-covalent. For example, such interaction can be identified at the protein level using in vitro biochemical methods, including photo-crosslinking, radiolabeled ligand binding, and affinity chromatography (Jakoby WB et al., 1974, Methods in Enzymology 46: 1). In certain cases, the compounds may be screened in a mechanism based assay, such as an assay to detect compounds which bind to an activin or ActRII polypeptide. This may include a solid phase or fluid phase binding event. Alternatively, the gene encoding an activin or ActRII polypeptide can be transfected with a reporter system (e.g., β -galactosidase, luciferase, or green fluorescent protein) into a cell and screened against the library optionally by a high throughput screening or with individual members of the library. Other mechanism based binding assays may be used, for example, binding assays which detect changes in free energy. Binding assays can be performed with the target fixed to a well, bead or chip or captured by an immobilized antibody or resolved by capillary electrophoresis. The bound compounds may be detected usually using colorimetric or fluorescence or surface plasmon resonance.

6. Exemplary Therapeutic Uses

In certain embodiments, activin-ActRII antagonists (e.g., ActRIIa or ActRIIb polypeptides) of the present invention can be used to increase red blood cell levels in

mammals such as rodents and primates, and particularly human patients. In certain embodiments, the present invention provides methods of treating or preventing anemia in an individual in need thereof by administering to the individual a therapeutically effective amount of an activin-ActRIIa antagonist, such as an ActRIIa polypeptide, or a therapeutically effective amount of an activin-ActRIIb antagonist, such as an ActRIIb polypeptide. In 5 certain embodiments, the present invention provides methods of promoting red blood cell formation in an individual by administering to the individual a therapeutically effective amount of an activin-ActRII antagonist, particularly an ActRII polypeptide. These methods may be used for therapeutic and prophylactic treatments of mammals, and particularly 10 humans.

As used herein, a therapeutic that "prevents" a disorder or condition refers to a compound that, in a statistical sample, reduces the occurrence of the disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated 15 control sample. The term "treating" as used herein includes prophylaxis of the named condition or amelioration or elimination of the condition once it has been established. In either case, prevention or treatment may be discerned in the diagnosis provided by a physician or other health care provider and the intended result of administration of the therapeutic agent.

As shown herein, activin-ActRIIa antagonists and activin-ActRIIb antagonists may be 20 used to increase red blood cell, hemoglobin or reticulocyte levels in healthy individuals, and such antagonists may be used in selected patient populations. Examples of appropriate patient populations include those with undesirably low red blood cell or hemoglobin levels, such as patients having an anemia, and those that are at risk for developing undesirably low 25 red blood cell or hemoglobin levels, such as those patients that are about to undergo major surgery or other procedures that may result in substantial blood loss. In one embodiment, a patient with adequate red blood cell levels is treated with an activin-ActRIIa antagonist to increase red blood cell levels, and then blood is drawn and stored for later use in transfusions. In one embodiment, a patient with adequate red blood cell levels is treated with an activin- 30 ActRIIb antagonist to increase red blood cell levels, and then blood is drawn and stored for later use in transfusions.

Activin-ActRII antagonists disclosed herein, and particularly ActRIIa-Fc and ActRIIb proteins, may be used to increase red blood cell levels in patients having an anemia. When observing hemoglobin levels in humans, a level of less than normal for the appropriate age and gender category may be indicative of anemia, although individual variations are taken 5 into account. For example, a hemoglobin level of 12 g/dl is generally considered the lower limit of normal in the general adult population. Potential causes include blood-loss, nutritional deficits, medication reaction, various problems with the bone marrow and many diseases. More particularly, anemia has been associated with a variety of disorders that include, for example, chronic renal failure, myelodysplastic syndrome, rheumatoid arthritis, 10 bone marrow transplantation. Anemia may also be associated with the following conditions: solid tumors (e.g. breast cancer, lung cancer, colon cancer); tumors of the lymphatic system (e.g. chronic lymphocyte leukemia, non-Hodgkins and Hodgkins lymphomas); tumors of the hematopoietic system (e.g. leukemia, myelodysplastic syndrome, multiple myeloma); radiation therapy; chemotherapy (e.g. platinum containing regimens); inflammatory and 15 autoimmune diseases, including, but not limited to, rheumatoid arthritis, other inflammatory arthritides, systemic lupus erythematosus (SLE), acute or chronic skin diseases (e.g. psoriasis), inflammatory bowel disease (e.g. Crohn's disease and ulcerative colitis); acute or chronic renal disease or failure including idiopathic or congenital conditions; acute or chronic liver disease; acute or chronic bleeding; situations where transfusion of red blood cells is not 20 possible due to patient allo- or auto-antibodies and/or for religious reasons (e.g. some Jehovah's Witnesses); infections (e.g. malaria, osteomyelitis); hemoglobinopathies, including, for example, sickle cell disease, thalassemias; drug use or abuse, e.g. alcohol misuse; pediatric patients with anemia from any cause to avoid transfusion; and elderly patients or 25 patients with underlying cardiopulmonary disease with anemia who cannot receive transfusions due to concerns about circulatory overload.

Patients may be treated with a dosing regimen intended to restore the patient to a target hemoglobin level, usually between about 10 g/dl and about 12.5 g/dl, and typically about 11.0 g/dl (see also Jacobs et al. (2000) Nephrol Dial Transplant 15, 15-19), although lower target levels may cause fewer cardiovascular side effects. Alternatively, hematocrit 30 levels (percentage of the volume of a blood sample occupied by the cells) can be used as a measure for the condition of red blood cells. Hematocrit levels for healthy individuals range from 41 to 51% for adult males and from 35 to 45% for adult females. Target hematocrit

levels are usually around 30-33%. Moreover, hemoglobin/hematocrit levels vary from person to person. Thus, optimally, the target hemoglobin/hematocrit level can be individualized for each patient.

The rapid effect on red blood cell levels of the activin-ActRIIa antagonists disclosed 5 herein indicate that these agents act by a different mechanism than Epo. Accordingly, these antagonists may be useful for increasing red blood cell and hemoglobin levels in patients that do not respond well to Epo. For example, an activin-ActRIIa antagonist may be beneficial for a patient in which administering of a normal to increased (>300 IU/kg/week) dose of Epo does not result in the increase of hemoglobin level up to the target level. Patients with an 10 inadequate Epo response are found for all types of anemia, but higher numbers of non-responders have been observed particularly frequently in patients with cancers and patients with end-stage renal disease. An inadequate response to Epo can be either constitutive (i.e. observed upon the first treatment with Epo) or acquired (e.g. observed upon repeated treatment with Epo).

15 The activin-ActRII antagonists may also be used to treat patients that are susceptible to adverse effects of Epo. The primary adverse effects of Epo are an excessive increase in the hematocrit or hemoglobin levels and polycythemia. Elevated hematocrit levels can lead to hypertension (more particularly aggravation of hypertension) and vascular thrombosis. Other adverse effects of Epo which have been reported, some of which related to hypertension, are 20 headaches, influenza-like syndrome, obstruction of shunts, myocardial infarctions and cerebral convulsions due to thrombosis, hypertensive encephalopathy, and red cell blood cell aplasia (Singibarti, (1994) J. Clin Investig 72(suppl 6), S36-S43; Horl et al. (2000) Nephrol Dial Transplant 15(suppl 4), 51-56; Delanty et al. (1997) Neurology 49, 686-689; Bunn (2002) N Engl J Med 346(7), 522-523).

25

7. Pharmaceutical Compositions

In certain embodiments, activin-ActRII antagonists (e.g., ActRIIa and ActRIIb polypeptides) of the present invention are formulated with a pharmaceutically acceptable carrier. For example, an ActRII polypeptide can be administered alone or as a component of 30 a pharmaceutical formulation (therapeutic composition). The subject compounds may be

formulated for administration in any convenient way for use in human or veterinary medicine.

In certain embodiments, the therapeutic method of the invention includes administering the composition systemically, or locally as an implant or device. When 5 administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Therapeutically useful agents other than the activin-ActRII antagonists which may also optionally be included in the composition as described above, may be administered simultaneously or sequentially with the subject compounds (e.g., ActRIIa and ActRIIb polypeptides) in the methods of the invention.

10 Typically, activin-ActRII antagonists will be administered parenterally. Pharmaceutical compositions suitable for parenteral administration may comprise one or more ActRII polypeptides in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just 15 prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents. Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, 20 vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

25 Further, the composition may be encapsulated or injected in a form for delivery to a target tissue site (e.g., bone marrow). In certain embodiments, compositions of the present invention may include a matrix capable of delivering one or more therapeutic compounds (e.g., ActRIIa or ActRIIb polypeptides) to a target tissue site (e.g., bone marrow), providing a structure for the developing tissue and optimally capable of being resorbed into the body. For example, the matrix may provide slow release of the ActRII polypeptides. Such matrices 30 may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the subject compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, 5 tricalciumphosphate, hydroxyapatite, polylactic acid and polyanhydrides. Other potential materials are biodegradable and biologically well defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are non-biodegradable and chemically defined, such as sintered 10 hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, 15 particle size, particle shape, and biodegradability.

In certain embodiments, methods of the invention can be administered for orally, e.g., 15 in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of an 20 agent as an active ingredient. An agent may also be administered as a bolus, electuary or paste.

In solid dosage forms for oral administration (capsules, tablets, pills, dragees, 25 powders, granules, and the like), one or more therapeutic compounds of the present invention may be mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose, and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, 30 potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol

monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a 5 similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and elixirs. In addition to the 10 active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid 15 esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming, and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents 20 such as ethoxylated isostearyl alcohols, polyoxyethylene sorbitol, and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

The compositions of the invention may also contain adjuvants, such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of 25 microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

30 It is understood that the dosage regimen will be determined by the attending physician considering various factors which modify the action of the subject compounds of the

invention (e.g., ActRIIa and ActRIIb polypeptides). The various factors include, but are not limited to, the patient's red blood cell count, hemoglobin level or other diagnostic assessments, the desired target red blood cell count, the patient's age, sex, and diet, the severity of any disease that may be contributing to a depressed red blood cell level, time of 5 administration, and other clinical factors. The addition of other known growth factors to the final composition may also affect the dosage. Progress can be monitored by periodic assessment of red blood cell and hemoglobin levels, as well as assessments of reticulocyte levels and other indicators of the hematopoietic process.

Experiments with primates and humans have demonstrated that effects of ActRIIa-Fc 10 on red blood cell levels are detectable when the compound is dosed at intervals and amounts sufficient to achieve serum concentrations of about 100 ng/ml or greater, for a period of at least about 20 to 30 days. Dosing to obtain serum levels of 200 ng/ml, 500 ng/ml, 1000 ng/ml or greater for a period of at least 20 to 30 days may also be used. Bone effects can be observed at serum levels of about 200 ng/ml, with substantial effects beginning at about 1000 15 ng/ml or higher, over a period of at least about 20 to 30 days. Thus, if it is desirable to achieve effects on red blood cells while having little effect on bone, a dosing scheme may be designed to deliver a serum concentration of between about 100 and 1000 ng/ml over a period of about 20 to 30 days. In humans, serum levels of 200 ng/ml may be achieved with a single dose of 0.1 mg/kg or greater and serum levels of 1000 ng/ml may be achieved with a 20 single dose of 0.3 mg/kg or greater. The observed serum half-life of the molecule is between about 20 and 30 days, substantially longer than most Fc fusion proteins, and thus a sustained effective serum level may be achieved, for example, by dosing with about 0.05 to 0.5 mg/kg on a weekly or biweekly basis, or higher doses may be used with longer intervals between dosings. For example, doses of 0.1 to 1 mg/kg might be used on a monthly or bimonthly 25 basis.

In certain embodiments, the present invention also provides gene therapy for the in vivo production of ActRII polypeptides. Such therapy would achieve its therapeutic effect by introduction of the ActRIIa or ActRIIb polynucleotide sequences into cells or tissues having the disorders as listed above. Delivery of ActRII polynucleotide sequences can be achieved 30 using a recombinant expression vector such as a chimeric virus or a colloidal dispersion

system. Preferred for therapeutic delivery of ActRII polynucleotide sequences is the use of targeted liposomes.

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or an RNA virus such as a retrovirus. The retroviral vector may be a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated.

Retroviral vectors can be made target-specific by attaching, for example, a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody. Those of skill in the art will recognize that specific polynucleotide sequences can be inserted into the retroviral genome or attached to a viral envelope to allow target specific delivery of the retroviral vector containing the ActRII polynucleotide.

Alternatively, tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

Another targeted delivery system for ActRII polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (see e.g., Fraley, et al., Trends Biochem. Sci., 6:77, 1981). Methods for efficient gene transfer using a liposome vehicle, are known in the art, see e.g., Mannino, et al., Biotechniques, 6:682, 1988. The composition of the liposome is usually a combination of phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical

characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine,

5 phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine, and distearoylphosphatidylcholine. The targeting of liposomes is also possible based on, for example, organ-specificity, cell-specificity, and organelle-specificity and is known in the art.

10 EXEMPLIFICATION

The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain embodiments and embodiments of the present invention, and are not intended to limit the invention.

15 Example 1: ActRIIa-Fc Fusion Proteins

Applicants constructed a soluble ActRIIa fusion protein that has the extracellular domain of human ActRIIa fused to a human or mouse Fc domain with a minimal linker in between. The constructs are referred to as ActRIIa-hFc and ActRIIa-mFc, respectively.

ActRIIa-hFc is shown below as purified from CHO cell lines (SEQ ID NO: 7):

20 ILGRSETQECLFFNANWEKDRTNQTGVEPCYGDKDKRRHCFATWKNISGSIEIVKQG
CWLDDINCYDRTDCVEKKDSPEVYFCCCEGNMCNEKFSYFPEMEVTQPTSNPVTPK
PPTGGGTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF
NWYVDGVEVHNAKTPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP
VPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQP
25 ENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSL
SPGK

The ActRIIa-hFc and ActRIIa-mFc proteins were expressed in CHO cell lines. Three different leader sequences were considered:

(i) Honey bee mellitin (HBML): MKFLVNVALVFMVVYISYIYA (SEQ ID NO: 8)

(ii) Tissue Plasminogen Activator (TPA): MDAMKRLCCVLLCGAVFVSP (SEQ ID NO: 9)

(iii) Native: MGAAAKLAFAVFLISCSSGA (SEQ ID NO: 10).

The selected form employs the TPA leader and has the following unprocessed amino acid sequence:

5 MDAMKRLCCVLLCGAVFVSPGAAILGRSETQECLFFNANWEKDRTNQTGVEPCY
 GDKDKRRHCFATWKNISGSIEIVKQGCWLDDINCYDRTDCVEKKDSPEVYFCCCEG
 NMCNEKFSYFPEMEVTQPTSNPVTPKPPTGGGTHTCPPCPAPELLGGPSVLFPPKPK
 DTLMisRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTPREEQYNSTYRVVS
 10 VLTVLHQDWLNGKEYKCKVSNKALPVPIEKTISKAKGQPREPQVYTLPPSREEMTKN
 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQ
 QGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:13)

This polypeptide is encoded by the following nucleic acid sequence:

15 ATGGATGCAATGAAGAGAGGGCTCTGCTGTGCTGCTGTGGAGCAGTCT
 TCGTTTCGCCCCGGCGCCGCTATACTTGGTAGATCAGAAACTCAGGAGTGTCTTT
 TTTAATGCTAATTGGGAAAAAGACAGAACCAATCAAACCTGGTGTGAACCGTGT
 ATGGTACAAAGATAAACGGCGGCATTGTTTGCTACCTGGAAGAATATTCCTGG
 TTCCATTGAATAGTGAACACAAGGTTGGCTGGATGATATCAACTGCTATGACA
 GGACTGATTGTGTAGAAAAAAAAGACAGCCCTGAAGTATATTCTGTTGCTGTGA
 20 GGGCAATATGTGTAATGAAAAGTTCTTATTTCCGGAGATGGAAGTCACACAG
 CCCACTTCAAATCCAGTTACACCTAACGCCACCCACCGGTGGAACTCACACAT
 GCCCACCGTGCCAGCACCTGAACCTCCTGGGGGACCGTCAGTCTCCTCTTCCC
 CCCAAAACCAAGGACACCCTCATGATCTCCGGACCCCTGAGGTCAAGTTCAACTGGTACGTGGAC
 GTGGTGGACGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGAC
 25 GGCCTGGAGGTGCATAATGCCAAGACAAAGCCGCGGAGGAGCAGTACAACAG
 CACGTACCGTGTGGTCAGCGTCCTCACCGTCTGCACCAAGGACTGGCTGAATGGC
 AAGGAGTACAAGTGCAGGTCTCCAACAAAGCCCTCCCAGTCCCCATCGAGAAA
 ACCATCTCCAAGCCAAGGGCAGCCCCGAGAACCAAGGTGTACACCCCTGCC
 CCATCCCGGGAGGAGATGACCAAGAACCAAGGTCAGCCTGACCTGCCTGGTCAA
 30 GGCTTCTATCCCAGCGACATGCCGTGGAGTGGAGAGCAATGGCAGCCGGAG
 AACAACTACAAGACCAAGCACGCCCTCCCGTGGACTCCGACGGCTCCTTCTCCT

ATAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGAACGTCTTCT
CATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCT
CTCCCTGTCTCCGGGTAAATGAGAATT (SEQ ID NO:14)

Both ActRIIa-hFc and ActRIIa-mFc were remarkably amenable to recombinant
5 expression. As shown in figure 1, the protein was purified as a single, well-defined peak
of protein. N-terminal sequencing revealed a single sequence of –ILGRSETQE (SEQ
ID NO: 11). Purification could be achieved by a series of column chromatography steps,
including, for example, three or more of the following, in any order: protein A
chromatography, Q sepharose chromatography, phenylsepharose chromatography, size
10 exclusion chromatography, and cation exchange chromatography. The purification
could be completed with viral filtration and buffer exchange. The ActRIIa-hFc protein
was purified to a purity of >98% as determined by size exclusion chromatography and
>95% as determined by SDS PAGE.

ActRIIa-hFc and ActRIIa-mFc showed a high affinity for ligands, particularly
15 activin A. GDF-11 or Activin A (“ActA”) were immobilized on a Biacore CM5 chip
using standard amine coupling procedure. ActRIIa-hFc and ActRIIa-mFc proteins were
loaded onto the system, and binding was measured. ActRIIa-hFc bound to activin with a
dissociation constant (K_D) of 5×10^{12} , and the protein bound to GDF11 with a K_D of
 9.96×10^{-9} . See figure 2. ActRIIa-mFc behaved similarly.

20 The ActRIIa-hFc was very stable in pharmacokinetic studies. Rats were dosed
with 1 mg/kg, 3 mg/kg or 10 mg/kg of ActRIIa-hFc protein and plasma levels of the
protein were measured at 24, 48, 72, 144 and 168 hours. In a separate study, rats were
dosed at 1 mg/kg, 10 mg/kg or 30 mg/kg. In rats, ActRIIa-hFc had an 11-14 day serum
half life and circulating levels of the drug were quite high after two weeks (11 μ g/ml,
25 110 μ g/ml or 304 μ g/ml for initial administrations of 1 mg/kg, 10 mg/kg or 30 mg/kg,
respectively.) In cynomolgus monkeys, the plasma half life was substantially greater
than 14 days and circulating levels of the drug were 25 μ g/ml, 304 μ g/ml or 1440 μ g/ml
for initial administrations of 1 mg/kg, 10 mg/kg or 30 mg/kg, respectively.

Example 2: ActRIIa-hFc Increases Red Blood Cell Levels in Non-Human Primates

The study employed four groups of five male and five female cynomolgus monkeys each, with three per sex per group scheduled for termination on Day 29, and two per sex per group scheduled for termination on Day 57. Each animal was administered the vehicle

5 (Group I) or ActRIIa-Fc at doses of 1, 10, or 30 mg/kg (Groups 2, 3 and 4, respectively) via intravenous (IV) injection on Days 1, 8, 15 and 22. The dose volume was maintained at 3 mL/kg. Various measures of red blood cell levels were assessed two days prior to the first administration and at days 15, 29 and 57 (for the remaining two animals) after the first administration.

10 The ActRIIa-hFc causes statistically significant increases in mean red blood cell parameters (red blood cell count [RBC], hemoglobin [HGB], and hematocrit [HCT]) for males and females, at all dose levels and time points throughout the study, with accompanying elevations in absolute and relative reticulocyte counts (ARTC; RTC). See Figures 3 – 6.

15 Statistical significance was calculated for each treatment group relative to the mean for the treatment group at baseline.

Notably, the increases in red blood cell counts and hemoglobin levels are roughly equivalent in magnitude to effects reported with erythropoietin. The onset of these effects is more rapid with ActRIIa-Fc than with erythropoietin.

20 Similar results were observed with rats and mice.

Example 3: ActRIIa-hFc Increases Red Blood Cell Levels in Human Patients

The ActRIIa-hFc fusion protein described in Example 1 was administered to human patients in a randomized, double-blind, placebo-controlled study that was conducted to evaluate, primarily, the safety of the protein in healthy, postmenopausal women. Forty-eight subjects were randomized in cohorts of 6 to receive either a single dose of ActRIIa-hFc or placebo (5 active:1 placebo). Dose levels ranged from 0.01 to 3.0 mg/kg intravenously (IV) and 0.03 to 0.1 mg/kg subcutaneously (SC). All subjects were followed for 120 days. In addition to pharmacokinetic (PK) analyses, the biologic activity of ActRIIa-hFc was also

assessed by measurement of biochemical markers of bone formation and resorption, and FSH levels.

To look for potential changes, hemoglobin and RBC numbers were examined in detail for all subjects over the course of the study and compared to the baseline levels. Platelet 5 counts were compared over the same time as the control. There were no clinically significant changes from the baseline values over time for the platelet counts.

PK analysis of ActRIIa-hFc displayed a linear profile with dose, and a mean half-life of approximately 25-32 days. The area-under-curve (AUC) for ActRIIa-hFc was linearly related to dose, and the absorption after SC dosing was essentially complete (see Figures 7 10 and 8). These data indicate that SC is a desirable approach to dosing because it provides equivalent bioavailability and serum-half life for the drug while avoiding the spike in serum concentrations of drug associated with the first few days of IV dosing (see Figure 8).

ActRIIa-hFc caused a rapid, sustained dose-dependent increase in serum levels of bone-specific alkaline phosphatase (BAP), which is a marker for anabolic bone growth, and a dose-dependent decrease in C-terminal type 1 collagen telopeptide and tartrate-resistant acid phosphatase 5b levels, which are markers for bone resorption. Other markers, such as P1NP 15 showed inconclusive results. BAP levels showed near saturating effects at the highest dosage of drug, indicating that half-maximal effects on this anabolic bone biomarker could be achieved at a dosage of 0.3 mg/kg, with increases ranging up to 3 mg/kg. Calculated as a 20 relationship of pharmacodynamic effect to AUC for drug, the EC50 is 51,465 (day*ng/ml). See Figure 9. These bone biomarker changes were sustained for approximately 120 days at the highest dose levels tested. There was also a dose-dependent decrease in serum FSH levels consistent with inhibition of activin.

Overall, there was a very small non-drug related reduction in hemoglobin over the 25 first week of the study probably related to study phlebotomy in the 0.01 and 0.03 mg/kg groups whether given IV or SC. The 0.1 mg/kg SC and IV hemoglobin results were stable or showed modest increases by Day 8-15. At the 0.3 mg/kg IV dose level there was a clear increase in HGB levels seen as early as Day 2 and often peaking at Day 15-29 that was not seen in the placebo subjects. At this point in the study, this change has not reached statistical 30 significance.

Overall, ActRIIa-hFc showed a dose-dependent effect on red blood cell counts and reticulocyte counts. For a summary of hematological changes, see Figures 10-13.

Example 4: Alternative ActRIIa-Fc Proteins

5 A variety of ActRIIa variants that may be used according to the methods described herein are described in the International Patent Application published as WO2006/012627 (see e.g., pp. 55-58), incorporated herein by reference in its entirety. An alternative construct may have a deletion of the C-terminal tail (the final 15 amino acids of the extracellular domain of ActRIIa. The sequence for such a construct is presented below (Fc portion 10 underlined)(SEQ ID NO: 12):

ILGRSETQECLFFNANWEKDRTNQTGVEPCYGDKDCKRRHCFATWKNISGSIEIVKQG
CWLDDINCYDRTDCVEKKDSPEVYFCCCEGNMCNEKFSYFPEMTGGGTHTCPCPA
PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVFKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPVPIEKTISKAKGQP
15 RETKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDG
SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSPGK

Example 5: ActRIIb-Fc Fusion Proteins

Applicants constructed a soluble ActRIIb fusion protein that has the extracellular 20 domain of human ActRIIb fused to a human Fc domain. A co-crystal structure of Activin and extracellular ActRIIb did not show any role for the final (C-terminal) 15 amino acids (referred to as the “tail” herein) of the extracellular domain in ligand binding. This sequence failed to resolve on the crystal structure, suggesting that these residues are present in a flexible loop that did not pack uniformly in the crystal. Thompson et al. EMBO J. 2003 Apr 25 1;22(7):1555-66. This sequence is also poorly conserved between ActRIIb and ActRIIa. Accordingly, these residues were omitted in the basic, or background, ActRIIb-Fc fusion construct. Additionally, position 64 in the background form is occupied by an alanine, which is generally considered the “wild type” form, although a A64R allele occurs naturally. Thus, the background ActRIIb-Fc fusion has the sequence (Fc portion underlined)(SEQ ID NO:20):

SGRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWANSSGTIELVK
KGCWLDDFN~~CYDRQECVATEENPQVYFCCCEGNFCNERFTHLPEAGGGTHTCPPCP~~
APELLGGPSVFLPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPVPIEKTI~~SKAKGQP~~
5 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ~~PENNYK~~TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK

Surprisingly, the C-terminal tail was found to enhance activin and GDF-11 binding, thus a preferred version of ActRIIb-Fc has a sequence (Fc portion underlined)(SEQ ID NO:21):

10 SGRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWANSSGTIELVK
KGCWLDDFN~~CYDRQECVATEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTE~~~~YEP~~
TAPTGGGTHTCPPCPAPELLGGPSVFLPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHN~~AKT~~KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA
LPVPIEKTI~~SKAKGQ~~REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG
15 Q~~PENNYK~~TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK

A variety of ActRIIb a variants that may be used according to the methods described herein are described in the International Patent Application published as WO2006/012627 (see e.g., pp. 59-60), incorporated herein by reference in its entirety.

20

Example 6: ActRIIb-hFc Stimulates Erythropoiesis in Non-Human Primates

ActRIIb-hFc (IgG1) was administered once a week for 1-month to male and female cynomolgus monkeys by subcutaneous injection. Forty-eight cynomolgus monkeys (24/sex) were assigned to one of four treatment groups (6 animals/sex/group) and were administered 25 subcutaneous injections of either vehicle or ActRIIb-hFc at 3, 10, or 30 mg/kg once weekly for 4 weeks (total of 5 doses). Parameters evaluated included general clinical pathology (hematology, clinical chemistry, coagulation, and urinalysis). ActRIIb-hFc caused statistically significant elevated mean absolute reticulocyte values by day 15 in treated animals. By day 36, ActRIIb-hFc caused several hematological changes, including elevated 30 mean absolute reticulocyte and red blood cell distribution width values and lower mean corpuscular hemoglobin concentration. All treated groups and both sexes were affected.

These effects are consistent with a positive effect of ActRIIb-hFc on the release of immature reticulocytes from the bone marrow. This effect was reversed after drug was washed out of the treated animals (by study day 56). Accordingly, we conclude that ActRIIb-hFc stimulates erythropoiesis.

5

INCORPORATION BY REFERENCE

All publications and patents mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

10 While specific embodiments of the subject matter have been discussed, the above specification is illustrative and not restrictive. Many variations will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

15

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A method for increasing red blood cell levels in a human patient, the method comprising administering to a human patient in need thereof an effective amount of an ActRII polypeptide selected from the group consisting of:
 - 5 a) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:2;
 - b) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:3;
 - c) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:2;
 - 10 d) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:16;
 - e) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:17; and
 - f) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:16.
2. The method of claim 1, wherein the polypeptide has one or more of the following characteristics:
 - i) binds to an ActRII ligand with a K_D of at least 10^{-7} M; and
 - 20 ii) inhibits ActRII signaling in a cell.
3. The method of claim 1 or 2, wherein the polypeptide binds to activin A.
4. The method of any one of claims 1 to 3, wherein said polypeptide is a fusion protein including, in addition to an ActRII polypeptide domain, one or more polypeptide portions that enhance one or more of *in vivo* stability, *in vivo* half life, uptake/administration, tissue localization or distribution, formation of protein complexes, and/or purification.

5. The method of claim 4, wherein said fusion protein includes a polypeptide portion selected from the group consisting of: an immunoglobulin Fc domain and a serum albumin.
6. The method of claim 5, wherein the fusion protein comprises a constant region from an IgG1 heavy chain.
7. The method of any one of claims 1 to 6, wherein said polypeptide includes one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.
8. The method of any one of claims 1 to 7, wherein the increase in red blood cell levels is measured as an increase in hemoglobin levels in the patient's blood.
9. A method for preventing or treating an anemia, the method comprising administering to a subject in need thereof an effective amount of an activin antagonist or ActRII polypeptide.
10. The method of claim 9, wherein the ActRII polypeptide is selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:2;
 - 20 b) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:3;
 - c) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:2;
 - d) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:16;
 - 25 e) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:17; and

- f) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO: 16.
11. The method of claims 9 or 10, wherein the ActRII polypeptide has one or more of the following characteristics:
 - 5 i) binds to an ActRII ligand with a K_D of at least 10^{-7} M; and
 - ii) inhibits ActRII signaling in a cell.
12. The method of any one of claims 9 to 11, wherein the polypeptide binds to activin A.
13. The method of any one of claims 9 to 12, wherein said ActRII polypeptide is a fusion protein including, in addition to said ActRII polypeptide, one or more polypeptide portions that enhance one or more of *in vivo* stability, *in vivo* half life, uptake/administration, tissue localization or distribution, formation of protein complexes, and/or purification.
10
14. The method of claim 13, wherein said fusion protein includes a polypeptide portion selected from the group consisting of: an immunoglobulin Fc domain and a serum albumin.
15
15. The method of claim 14, wherein the fusion protein comprises a constant region from an IgG1 heavy chain.
16. The method of any one of claims 9 to 15, wherein said ActRII polypeptide includes one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.
20
17. The method of any one of claims 9 to 16, wherein the anemia is anemia associated with multiple myeloma.
25
18. The method of any one of claims 9 to 16, wherein the anemia is anemia associated with chemotherapeutic treatment of the patient.

19. The method of any one of claims 9 to 16, wherein the anemia is anemia associated with myelodysplastic syndrome.
20. The method of any one of claims 9 to 16, wherein the anemia is anemia associated with thalassemia.
- 5 21. A method of identifying an agent that increases red blood cell levels, the method comprising:
 - a) identifying a test agent that binds to a ligand-binding domain of an ActRII polypeptide competitively with an ActRII ligand; and
 - b) evaluating the effect of the agent on red blood cell levels in an animal.
- 10 22. Use of an activin antagonist or ActRII polypeptide for making a medicament for increasing red blood cell levels in a human patient.
23. A method for increasing red blood cell levels in a human patient, the method comprising administering to a patient in need thereof an effective amount of an ActRII-Fc fusion protein, wherein the ActRII-Fc fusion protein comprises a polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:3;
 - b) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:2;
 - 15 c) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:7;
 - d) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:17;
 - e) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:16;
 - 20 f) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:20;

- g) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:21; and
- h) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:12.

5 24. The method of claim 23, wherein the patient has an anemia.

25. The method of claim 24, wherein the anemia is anemia associated with multiple myeloma.

26. The method of claim 24, wherein the anemia is anemia associated with chemotherapeutic treatment of the patient.

10 27. The method of claim 24, wherein the anemia is anemia associated with myelodysplastic syndrome.

28. The method of claim 24, wherein the anemia is anemia associated with thalassemia.

29. The method of any one of claims 23 to 28, wherein the method causes less than 15% increase in the patient's skeletal muscle mass.

15 30. The method of any one of claims 23 to 29, wherein the ActRII-Fc fusion protein is administered so as to reach a serum concentration in the patient of at least 100 ng/ml for a period of about 20 to 30 days.

31. The method of any one of claims 23 to 30, wherein the ActRII-Fc fusion protein 20 is administered so as to reach a serum concentration in the patient in the range of 100 ng/ml to 1000 ng/ml.

32. The method of any one of claims 23 to 31, wherein the ActRII-Fc fusion protein has a serum half-life of between 15 and 30 days.

33. The method of any one of claims 23 to 32, wherein the ActRII-Fc fusion protein 25 is administered to the patient no more frequently than once per week.

34. The method of claim 33, wherein the ActRII-Fc fusion protein is administered to the patient no more frequently than once per month.

35. The use of claim 22, wherein the ActRII polypeptide selected from the group

consisting of:

- a) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:2;
- 5 b) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:2;
- c) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:2;
- 10 d) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:3;
- e) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:3;
- 15 f) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:3;
- g) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:2;
- 20 h) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:16;
- i) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:16;
- j) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:16;
- 25 k) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:17;
- l) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:17;
- m) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:17; and

n) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO: 16.

36. The use of claim 35, wherein the polypeptide has one or more of the following characteristics:

5 i) binds to an ActRII ligand with a K_D of at least 10^{-7} M; and

 ii) inhibits ActRII signaling in a cell.

37. The use of claim 35 or 36, wherein said polypeptide is a fusion protein including, in addition to said activin-ActRII antagonist polypeptide, one or more polypeptide portions that enhance one or more of *in vivo* stability, *in vivo* half life, uptake/administration, tissue localization or distribution, formation of protein complexes, and/or purification.

10 38. The use of claim 37, wherein said fusion protein includes a polypeptide portion selected from the group consisting of: an immunoglobulin Fc domain and a serum albumin.

15 39. The use of any one of claims 35 to 38, wherein said polypeptide includes one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.

20 40. The use of any one of claims 35 to 39, wherein the increase in red blood cell levels is measured as an increase in hemoglobin levels in the patient's blood.

41. Use of an activin antagonist or ActRII polypeptide in the manufacture of a medicament for preventing or treating an anemia in a patient.

25 42. The use of claim 41, wherein the ActRII polypeptide is selected from the group consisting of:

 a) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:2;

- b) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:2;
- 5 c) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:2;
- d) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:3;
- 10 e) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:3;
- f) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:3;
- g) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO:2;
- h) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:16;
- 15 i) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:16;
- j) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:16;
- k) a polypeptide comprising an amino acid sequence that is at least 90% identical to SEQ ID NO:17;
- 20 l) a polypeptide comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:17;
- m) a polypeptide comprising an amino acid sequence that comprises the amino acid sequence of SEQ ID NO:17; and
- 25 n) a polypeptide comprising at least 50 consecutive amino acids selected from SEQ ID NO: 16.

43. The use of claim 41 or 42, wherein the ActRII polypeptide has one or more of the following characteristics:

- i) binds to an ActRII ligand with a K_D of at least 10^{-7} M; and
 - ii) inhibits ActRII signaling in a cell.
- 44. The use of any one of claims 41 to 43, wherein said ActRII polypeptide is a fusion protein including, in addition to said ActRII polypeptide, one or more polypeptide portions that enhance one or more of *in vivo* stability, *in vivo* half life, uptake/administration, tissue localization or distribution, formation of protein complexes, and/or purification.
- 5 45. The use of claim 44, wherein said fusion protein includes a polypeptide portion selected from the group consisting of: an immunoglobulin Fc domain and a serum albumin.
- 10 46. The use of any one of claims 41 to 45, wherein said ActRII polypeptide includes one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.
- 15 47. Use of an ActRII-Fc fusion protein, wherein the ActRII-Fc fusion protein comprises an amino acid sequence selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:3;
 - 20 b) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:3;
 - c) a polypeptide comprising the amino acid sequence of SEQ ID NO:3;
 - d) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:2;
 - 25 e) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:2;
 - f) a polypeptide comprising the amino acid sequence of SEQ ID NO:2;

- g) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:7;
- h) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:7;
- 5 i) a polypeptide comprising the amino acid sequence of SEQ ID NO:7;
- j) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:17;
- k) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:17;
- 10 l) a polypeptide comprising the amino acid sequence of SEQ ID NO:17;
- m) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:16;
- n) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:16;
- 15 o) a polypeptide comprising the amino acid sequence of SEQ ID NO:16;
- p) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:20;
- q) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:20;
- 20 r) a polypeptide comprising the amino acid sequence of SEQ ID NO:20;
- s) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:21;
- t) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:21;
- 25 u) a polypeptide comprising the amino acid sequence of SEQ ID NO:21;
- v) a polypeptide comprising an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO:12;

w) a polypeptide comprising an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO:12;

x) a polypeptide comprising the amino acid sequence of SEQ ID NO:12, in the manufacture of a medicament for increasing red blood cell levels in a human patient.

5 48. The use of claim 47, wherein the medicament causes less than a 15% increase in the patient's skeletal muscle mass.

49. The use of claim 47 or 48, wherein the medicament is for administration so as to reach a serum concentration of the fusion protein in the patient of at least 100 10 ng/ml for a period of about 20 to 30 days.

50. The use of any one of claims 47 to 48, wherein the medicament is for administration so as to reach a serum concentration of the fusion protein in the patient in the range of 100 ng/ml to 1000 ng/ml.

51. The use of any one of claims 47 to 49, wherein the ActRII-Fc fusion protein has 15 a serum half-life of between 15 and 30 days.

52. The use of any one of claims 47 to 50, wherein the medicament is for administration to the patient no more frequently than once per week.

53. The use of any one of claims 47 to 50, wherein the medicament is for administration to the patient no more frequently than once per month.

20 54. The use of any one of claims 47 to 53, wherein the patient has an anemia.

55. The use of claim 54, wherein the anemia is anemia associated with multiple myeloma.

56. The use of claim 54, wherein the anemia is anemia associated with 25 chemotherapeutic treatment of the patient.

57. The use of claim 54, wherein the anemia is anemia associated with myelodysplastic syndrome.

58. The use of claim 54 wherein the anemia is anemia associated with thalassemia.

59. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:2.
60. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:2.
- 5 61. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:2.
62. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:3.
63. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:3.
- 10 64. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:3.
65. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:16.
- 15 66. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:16.
67. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:16.
- 20 68. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:17.
69. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:17.
- 25 70. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:17.

71. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:7.
72. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:7.
- 5 73. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:7.
74. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:20.
- 10 75. The method of any one of claims 1 to 20 or 23 to 341, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:20.
76. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:20.
- 15 77. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:21.
78. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:21.
- 20 79. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:21.
80. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:12.
- 25 81. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:12.

82. The method of any one of claims 1 to 20 or 23 to 34, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:12.
83. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:2.
- 5 84. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:2.
85. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:2.
- 10 86. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:3.
87. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:3.
88. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:3.
- 15 89. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:16.
90. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:16.
19. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:16.
92. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:17.
93. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:17.
- 25 94. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:17.
95. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:7.

96. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:7.
97. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:7.
- 5 98. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:20.
99. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:20.
100. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:20.
- 10 101. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:21.
102. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:21.
- 15 103. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:21.
104. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:12.
105. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO:12.
- 20 106. The use of any one of claims 22 or 35-58, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:12.
131. A method according to any one of claims 1, 9 or 23; or use according to any one of claims 22, 41, or 47; substantially as herein described with reference to any one or more examples but excluding comparative examples.
- 25 132. The method of any one of claims 9 to 16 or 24, wherein the anemia is anemia associated with chronic renal failure.
133. The method of any one of claims 9 to 16 or 24, wherein the anemia is anemia

associated with acute renal failure.

134. The method of any one of claims 9 to 16 or 24, wherein the anemia is anemia associated with end-stage renal disease.

132. The use of claim 54, wherein the anemia is anemia associated with chronic renal failure.

133. The use of claim 54, wherein the anemia is anemia associated with acute renal failure.

134. The use of claim 54, wherein the anemia is anemia associated with end-stage renal disease.




Figure 1





Figure 2

Effect of ActRIIA-Fc on RBC Count in Female NHPs




Figure 3A

Effect of ActRIIA-Fc on Hemoglobin in Female NHPs




Figure 3B

Effect of ActRIIA-Fc on RBC Count in Male NHPs




Figure 4A

Effect of ActRIIA-Fc on Hemoglobin in Male NHPs




Figure 4B

Effect of ActRIIA-Fc on Reticulocyte Counts in Female NHPs




Figure 5A

Effect of ActRIIA-Fc on Reticulocyte Percentage in Female NHPs






Figure 5B






Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13

20073343333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt
SEQUENCE LISTING

<110> ACCELERON PHARMA INC.
<120> ACTIVIN-ACTRII ANTAGONISTS AND USES FOR INCREASING RED BLOOD CELL LEVELS
<130> PHPH-022-W01
<140> PCT/US07/025868
<141> 2007-12-18
<150> 60/875,682
<151> 2006-12-18
<160> 23
<170> PatentIn Ver. 3.3
<210> 1
<211> 513
<212> PRT
<213> Homo sapiens
<400> 1
Met Gly Ala Ala Ala Lys Leu Ala Phe Ala Val Phe Leu Ile Ser Cys
1 5 10 15
Ser Ser Gly Ala Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe
20 25 30
Phe Asn Ala Asn Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu
35 40 45
Pro Cys Tyr Gly Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp
50 55 60
Lys Asn Ile Ser Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu
65 70 75 80
Asp Asp Ile Asn Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp
85 90 95
Ser Pro Glu Val Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu
100 105 110
Lys Phe Ser Tyr Phe Pro Glu Met Glu Val Thr Gln Pro Thr Ser Asn
115 120 125
Pro Val Thr Pro Lys Pro Pro Tyr Tyr Asn Ile Leu Leu Tyr Ser Leu
130 135 140
Val Pro Leu Met Leu Ile Ala Gly Ile Val Ile Cys Ala Phe Trp Val
145 150 155 160
Tyr Arg His His Lys Met Ala Tyr Pro Pro Val Leu Val Pro Thr Gln
165 170 175
Asp Pro Gly Pro Pro Pro Ser Pro Leu Leu Gly Leu Lys Pro Leu
180 185 190
Gln Leu Leu Glu Val Lys Ala Arg Gly Arg Phe Gly Cys Val Trp Lys
195 200 205
Ala Gln Leu Leu Asn Glu Tyr Val Ala Val Lys Ile Phe Pro Ile Gln
210 215 220
Asp Lys Gln Ser Trp Gln Asn Glu Tyr Glu Val Tyr Ser Leu Pro Gly
225 230 235 240

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

Met Lys His Glu Asn Ile Leu Gln Phe Ile Gly Ala Glu Lys Arg Gly
245 250 255
Thr Ser Val Asp Val Asp Leu Trp Leu Ile Thr Ala Phe His Glu Lys
260 265 270
Gly Ser Leu Ser Asp Phe Leu Lys Ala Asn Val Val Ser Trp Asn Glu
275 280 285
Leu Cys His Ile Ala Glu Thr Met Ala Arg Gly Leu Ala Tyr Leu His
290 295 300
Glu Asp Ile Pro Gly Leu Lys Asp Gly His Lys Pro Ala Ile Ser His
305 310 315 320
Arg Asp Ile Lys Ser Lys Asn Val Leu Leu Lys Asn Asn Leu Thr Ala
325 330 335
Cys Ile Ala Asp Phe Gly Leu Ala Leu Lys Phe Glu Ala Gly Lys Ser
340 345 350
Ala Gly Asp Thr His Gly Gln Val Gly Thr Arg Arg Tyr Met Ala Pro
355 360 365
Glu Val Leu Glu Gly Ala Ile Asn Phe Gln Arg Asp Ala Phe Leu Arg
370 375 380
Ile Asp Met Tyr Ala Met Gly Leu Val Leu Trp Glu Leu Ala Ser Arg
385 390 395 400
Cys Thr Ala Ala Asp Gly Pro Val Asp Glu Tyr Met Leu Pro Phe Glu
405 410 415
Glu Glu Ile Gly Gln His Pro Ser Leu Glu Asp Met Gln Glu Val Val
420 425 430
Val His Lys Lys Arg Pro Val Leu Arg Asp Tyr Trp Gln Lys His
435 440 445
Ala Gly Met Ala Met Leu Cys Glu Thr Ile Glu Glu Cys Trp Asp His
450 455 460
Asp Ala Glu Ala Arg Leu Ser Ala Gly Cys Val Gly Glu Arg Ile Thr
465 470 475 480
Gln Met Gln Arg Leu Thr Asn Ile Ile Thr Thr Glu Asp Ile Val Thr
485 490 495
Val Val Thr Met Val Thr Asn Val Asp Phe Pro Pro Lys Glu Ser Ser
500 505 510
Leu

<210> 2
<211> 115
<212> PRT
<213> Homo sapiens

<400> 2
Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe Phe Asn Ala Asn
1 5 10 15
Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu Pro Cys Tyr Gly
20 25 30
Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp Lys Asn Ile Ser

200734333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt

35 40 45

Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu Asp Asp Ile Asn
50 55 60
Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp Ser Pro Glu Val
65 70 75 80
Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu Lys Phe Ser Tyr
85 90 95
Phe Pro Glu Met Glu Val Thr Gln Pro Thr Ser Asn Pro Val Thr Pro
100 105 110
Lys Pro Pro
115

<210> 3
<211> 100
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 3
Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe Phe Asn Ala Asn
1 5 10 15
Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu Pro Cys Tyr Gly
20 25 30
Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp Lys Asn Ile Ser
35 40 45
Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu Asp Asp Ile Asn
50 55 60
Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp Ser Pro Glu Val
65 70 75 80
Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu Lys Phe Ser Tyr
85 90 95
Phe Pro Glu Met
100

<210> 4
<211> 1542
<212> DNA
<213> Homo sapiens

<400> 4
atgggagctg ctgcaaagtt ggcgtttgcc gtctttctta tctcctgttc ttcaggtgct 60
atacttggta gatcagaaac tcagggatgt cttttcttta atgctaattg ggaaaaagac 120
agaaccaatc aaactgggtt tgaaccgtgt tatggtgaca aagataaaacg gcggcattgt 180
tttgcaccc ggaagaatat ttctggttcc attgaaatag tgaacaagg ttgttgctg 240
gatgatatca actgctatga caggactgat tttgttagaa aaaaagacag ccctgaagta 300
tattttgtt gctgtgaggg caaatatgtgt aatgaaaatgt tttcttattt tccagagatg 360
gaagtcacac agcccacttc aaatccagtt acacctaagg cacccttatta caacatcctg 420
ctctattccct tgggtgccact tatgttaatt gcggggattt tcatttgc attttgggtg 480
tacaggcatc acaagatggc ctaccctcct gtacttggc caactcaaga cccaggacca 540
cccccacctt ctccattact aggggtgaaa ccactgcagt tattagaagt gaaagcaagg 600
ggaagattt gttgtgtctg gaaagccccag ttgccttaacg aatatgtggc tgtcaaaata 660
tttccaaatac aggacaaaca gtcatggcaa aatgaatacg aagtctacag tttgccttgg 720
atgaagcatg agaacaatatt acagttcatt ggtgcagaaaa aacgaggcac cagtgttgat 780

2007343333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt

gtggatctt	ggctgatcac	agcattcat	aaaaagggtt	cactatcaga	cttcttaag	840
gctaatgtgg	tctcttgaa	tgaactgtgt	catattgcag	aaaccatggc	tagaggattg	900
gcatatttac	atgaggatat	acctggccta	aaagatggcc	acaaacctgc	catatccac	960
aggcacatca	aaagtaaaaa	tgtctgttg	aaaaacaacc	tgacagcttgc	cattgctgac	1020
tttgggttgg	ccttaaaatt	tgaggctggc	aagtctgcag	gcgataccca	tggacaggtt	1080
gttacccgga	ggtacatggc	tccagaggtt	ttagagggttgc	ctataaactt	ccaaagggtat	1140
gcattttgtt	ggatagatat	gtatgccatg	ggatttagtcc	tatgggaact	ggcttctcgc	1200
tgtactgctg	cagatggacc	tgtatgttgc	tacatgttgc	catttgagga	ggaaattggc	1260
cagcatccat	ctcttgaaga	catgcaggaa	gttgggttgc	ataaaaaaaa	gaggccgtt	1320
ttaagagatt	attggcagaa	acatgcttgc	atggcaatgc	tctgtgaaac	cattgaagaa	1380
tgtgggatc	acgacgcaga	agccaggtt	tcagctggat	gtgttagtga	aagaattacc	1440
cagatgcaga	gactaacaaa	tattattacc	acagaggaca	ttgttaacagt	ggtcacaatg	1500
gtgacaaatg	ttgactttcc	tcccaaagaa	tctagtctat	ga		1542

<210> 5
<211> 345
<212> DNA
<213> Homo sapiens

<400> 5
atacttggta gatcagaaac tcaggagtgt cttttcttta atgctaattg gaaaaaagac 60
agaaccaatc aaactgggtt tgaaccgtgt tatggtgaca aagataaacg gcggcattgt 120
tttgcacatc ggaagaatat ttctggttcc attgaaatag taaaacaagg ttgttgctg 180
gatgatatca actgctatga caggactgtat tttttgtt gctgtgaggg caatatgtgtt aatgaaaagt 300
gaagtacacac agccacttc aaatccagtt acacctaagc cacc 345

<210> 6
<211> 225
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<220>
<221> MOD_RES
<222> (43)
<223> Asp or Ala

<220>
<221> MOD_RES
<222> (100)
<223> Lys or Ala

<220>
<221> MOD_RES
<222> (212)
<223> His or Ala

<400> 6
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
1 5 10 15

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
20 25 30

Arg Thr Pro Glu Val Thr Cys Val Val Val Xaa Val Ser His Glu Asp
35 40 45

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
50 55 60

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
65 70 75 80

200734333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
85 90 95
Tyr Lys Cys Xaa Val Ser Asn Lys Ala Leu Pro Val Pro Ile Glu Lys
100 105 110
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
115 120 125
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
130 135 140
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
145 150 155 160
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
165 170 175
Asp Ser Asp Gly Pro Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
180 185 190
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
195 200 205
Ala Leu His Xaa His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
210 215 220
Lys
225

<210> 7
<211> 344
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic construct

<400> 7
Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe Phe Asn Ala Asn
1 5 10 15
Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu Pro Cys Tyr Gly
20 25 30
Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp Lys Asn Ile Ser
35 40 45
Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu Asp Asp Ile Asn
50 55 60
Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp Ser Pro Glu Val
65 70 75 80
Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu Lys Phe Ser Tyr
85 90 95
Phe Pro Glu Met Glu Val Thr Gln Pro Thr Ser Asn Pro Val Thr Pro
100 105 110
Lys Pro Pro Thr Gly Gly Thr His Thr Cys Pro Pro Cys Pro Ala
115 120 125
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
130 135 140
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt
145 150 155 160

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
165 170 175

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
180 185 190

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
195 200 205

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
210 215 220

Leu Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
225 230 235 240

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
245 250 255

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
260 265 270

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
275 280 285

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
290 295 300

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
305 310 315 320

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
325 330 335

Ser Leu Ser Leu Ser Pro Gly Lys
340

<210> 8

<211> 21

<212> PRT

<213> Apis mellifera

<400> 8

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
1 5 10 15

Ser Tyr Ile Tyr Ala
20

<210> 9

<211> 22

<212> PRT

<213> Unknown Organism

<220>

<223> Description of Unknown Organism: Tissue
Plasminogen Activator leader sequence

<400> 9

Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
1 5 10 15

Ala Val Phe Val Ser Pro
20

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

<210> 10
<211> 20
<212> PRT
<213> Unknown Organism

<220>
<223> Description of Unknown Organism: Native leader sequence

<400> 10
Met Gly Ala Ala Ala Lys Leu Ala Phe Ala Val Phe Leu Ile Ser Cys
1 5 10 15

Ser Ser Gly Ala
20

<210> 11
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 11
Ile Leu Gly Arg Ser Glu Thr Gln Glu
1 5

<210> 12
<211> 329
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic construct

<400> 12
Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe Phe Asn Ala Asn
1 5 10 15

Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu Pro Cys Tyr Gly
20 25 30

Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp Lys Asn Ile Ser
35 40 45

Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu Asp Asp Ile Asn
50 55 60

Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp Ser Pro Glu Val
65 70 75 80

Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu Lys Phe Ser Tyr
85 90 95

Phe Pro Glu Met Thr Gly Gly Thr His Thr Cys Pro Pro Cys Pro
100 105 110

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
115 120 125

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
130 135 140

Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr

200734333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt
145 150 155 160
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
165 170 175
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
180 185 190
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
195 200 205
Ala Leu Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
210 215 220
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
225 230 235 240
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
245 250 255
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
260 265 270
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
275 280 285
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
290 295 300
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
305 310 315 320
Lys Ser Leu Ser Leu Ser Pro Gly Lys
325

<210> 13
<211> 369
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
construct

<400> 13
Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
1 5 10 15
Ala Val Phe Val Ser Pro Gly Ala Ala Ile Leu Gly Arg Ser Glu Thr
20 25 30
Gln Glu Cys Leu Phe Phe Asn Ala Asn Trp Glu Lys Asp Arg Thr Asn
35 40 45
Gln Thr Gly Val Glu Pro Cys Tyr Gly Asp Lys Asp Lys Arg Arg His
50 55 60
Cys Phe Ala Thr Trp Lys Asn Ile Ser Gly Ser Ile Glu Ile Val Lys
65 70 75 80
Gln Gly Cys Trp Leu Asp Asp Ile Asn Cys Tyr Asp Arg Thr Asp Cys
85 90 95
Val Glu Lys Lys Asp Ser Pro Glu Val Tyr Phe Cys Cys Cys Glu Gly
100 105 110
Asn Met Cys Asn Glu Lys Phe Ser Tyr Phe Pro Glu Met Glu Val Thr
115 120 125

200734333 28 Mar 2011

PHPH-022-W01_Substitute_Sequence_Listing.txt

Gln Pro Thr Ser Asn Pro Val Thr Pro Lys Pro Pro Thr Gly Gly Gly
130 135 140
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
145 150 155 160
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
165 170 175
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
180 185 190
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
195 200 205
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
210 215 220
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
225 230 235 240
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Val Pro Ile Glu Lys
245 250 255
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
260 265 270
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
275 280 285
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
290 295 300
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
305 310 315 320
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
325 330 335
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
340 345 350
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
355 360 365
Lys

<210> 14
<211> 1114
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 14
atggatcaa tgaagagagg gctctgctgt gtgctgctgc tgggtggagc agtcttcgtt 60
tcgcccggcg ccgctatact tggtagatca gaaactcagg agtgtctttt tttaatgcta 120
attggaaaaa agacagaacc aatcaaactg gtgttgaacc gtgttatggg gacaaagata 180
aacggcggca ttgttttgct acctggaga atatttctgg ttccattgaa tagtggaaaca 240
agggttgtgg ctggatgata tcaactgcta tgacaggact gattgtgtag aaaaaaaaaga 300
cagccctgaa gtatatttct gttgctgtga gggcaatatg tggtaatgaaa agttttctta 360
ttttccggag atggaagtca cacagccac ttcaaattcca gttacaccta agccacccac 420
cggtgggtgg aactcacat gcccaccgtg cccagcacct gaactcctgg ggggaccgtc 480
agtcttcctc ttccccccaa aacccaagga caccctcatg atctcccgaa cccctgaggt 540

28 Mar 2011

2007334333

PHPH-022-W01_Substitute_Sequence_Listing.txt
cacatgcgtg gtgggtggacg tgagccacga agaccctgag gtcaagttca actggtagt 600
ggacggcgtg gaggtgcata atgccaagac aaagccgcgg gaggagcgt acaacacac 660
gtaccgtgtg gtcagcgtcc tcaccgtcct gcaccaggac tggctgaatg gcaaggagta 720
caagtgcag gtctccaaca aagccctccc agtccccatc gagaaaacca tctccaaagc 780
caaagggcag ccccgagaac cacaggtgtc caccctgccc ccatcccggg aggagatgac 840
caagaaccag gtcagcgtga cctgcctggt caaaggcttc tatccagcg acatcgccgt 900
ggagtgggag agcaatgggc agccggagaa caactacaag accacgcctc ccgtgctgga 960
ctccgacggc tccttcttcc tctatagcaa gtcaccgtg gacaagagca ggtggcagca 1020
ggggAACGTC ttctcatgct ccgtgatgca tgaggctctg cacaaccact acacgcagaa 1080
gagcctctcc ctgtctccgg gtaaatgaga attc 1114

<210> 15
<211> 512
<212> PRT
<213> Homo sapiens

<400> 15
Met Thr Ala Pro Trp Val Ala Leu Ala Leu Leu Trp Gly Ser Leu Trp
1 5 10 15
Pro Gly Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr
20 25 30
Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg
35 40 45
Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala
50 55 60
Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp
65 70 75 80
Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn
85 90 95
Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg
100 105 110
Phe Thr His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro
115 120 125
Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr Ser Leu Leu
130 135 140
Pro Ile Gly Gly Leu Ser Leu Ile Val Leu Leu Ala Phe Trp Met Tyr
145 150 155 160
Arg His Arg Lys Pro Pro Tyr Gly His Val Asp Ile His Glu Asp Pro
165 170 175
Gly Pro Pro Pro Pro Ser Pro Leu Val Gly Leu Lys Pro Leu Gln Leu
180 185 190
Leu Glu Ile Lys Ala Arg Gly Arg Phe Gly Cys Val Trp Lys Ala Gln
195 200 205
Leu Met Asn Asp Phe Val Ala Val Lys Ile Phe Pro Leu Gln Asp Lys
210 215 220
Gln Ser Trp Gln Ser Glu Arg Glu Ile Phe Ser Thr Pro Gly Met Lys
225 230 235 240
His Glu Asn Leu Leu Gln Phe Ile Ala Ala Glu Lys Arg Gly Ser Asn
245 250 255
Leu Glu Val Glu Leu Trp Leu Ile Thr Ala Phe His Asp Lys Gly Ser
260 265 270

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

Leu Thr Asp Tyr Leu Lys Gly Asn Ile Ile Thr Trp Asn Glu Leu Cys
275 280 285
His Val Ala Glu Thr Met Ser Arg Gly Leu Ser Tyr Leu His Glu Asp
290 295 300
Val Pro Trp Cys Arg Gly Glu Gly His Lys Pro Ser Ile Ala His Arg
305 310 315 320
Asp Phe Lys Ser Lys Asn Val Leu Leu Lys Ser Asp Leu Thr Ala Val
325 330 335
Leu Ala Asp Phe Gly Leu Ala Val Arg Phe Glu Pro Gly Lys Pro Pro
340 345 350
Gly Asp Thr His Gly Gln Val Gly Thr Arg Arg Tyr Met Ala Pro Glu
355 360 365
Val Leu Glu Gly Ala Ile Asn Phe Gln Arg Asp Ala Phe Leu Arg Ile
370 375 380
Asp Met Tyr Ala Met Gly Leu Val Leu Trp Glu Leu Val Ser Arg Cys
385 390 395 400
Lys Ala Ala Asp Gly Pro Val Asp Glu Tyr Met Leu Pro Phe Glu Glu
405 410 415
Glu Ile Gly Gln His Pro Ser Leu Glu Glu Leu Gln Glu Val Val Val
420 425 430
His Lys Lys Met Arg Pro Thr Ile Lys Asp His Trp Leu Lys His Pro
435 440 445
Gly Leu Ala Gln Leu Cys Val Thr Ile Glu Glu Cys Trp Asp His Asp
450 455 460
Ala Glu Ala Arg Leu Ser Ala Gly Cys Val Glu Glu Arg Val Ser Leu
465 470 475 480
Ile Arg Arg Ser Val Asn Gly Thr Thr Ser Asp Cys Leu Val Ser Leu
485 490 495
Val Thr Ser Val Thr Asn Val Asp Leu Pro Pro Lys Glu Ser Ser Ile
500 505 510

<210> 16
<211> 116
<212> PRT
<213> Homo sapiens

<400> 16
Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala
1 5 10 15
Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu
20 25 30
Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser
35 40 45
Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe
50 55 60
Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln
65 70 75 80
Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr
85 90 95

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro
100 105 110

Thr Ala Pro Thr
115

<210> 17

<211> 101

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 17

Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala
1 5 10 15

Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu
20 25 30

Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser
35 40 45

Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe
50 55 60

Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln
65 70 75 80

Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr
85 90 95

His Leu Pro Glu Ala
100

<210> 18

<211> 1539

<212> DNA

<213> Homo sapiens

<400> 18

atgacggcgc cctgggtggc cctcgccctc ctctggggat cgctgtggcc cggctctggg 60
cgtggggagg ctgagacacg ggagtgcac tactacaacg ccaactggga gctggagcgc 120
accaaccaga gcggcctgga gcgcgtcgaa ggcgagcagg acaagcggct gcactgctac 180
gcctcctggg ccaacagctc tggaccatc gagctcgtga agaagggctg ctggctagat 240
gacttcaact gctacgatag gcaggagtgt gtggccactg aggagaaccc ccaggtgtac 300
ttctgtctgtc gtgaaggcaa cttctgcaac gagcgcttca ctcatttgcc agaggctggg 360
ggcccgaaag tcacgtacga gcccaccccg acagccccca ccctgctcac ggtgctggcc 420
tactcactgc tgcccatcg gggcccttcc ctcatgtcc tgctggcctt ttggatgtac 480
cgccatcgca agccccctt cggcatgtg gacatccatg aggaccctgg gcctccacca 540
ccatccccctc tggtgggccc taagccactg cagctgtgg agatcaaggc tcggggcg 600
tttggctgtc tcttggaaaggc ccagctcatg aatgacttt tagctgtcaa gatcttccca 660
ctccaggaca agcagtcgtg gcagagtgaa cgggagatct tcagcacacc tggcatgaag 720
cacgagaacc tgctacagtt cattgctgcc gagaagcggag gctccaacct cgaagttagag 780
ctgtggctca tcacggccctt ccattgacaag ggctccctca cggattacct caaggggaac 840
atcatcacat ggaacgaact gtgtcatgt a c g a g a c g a c g a t g t c a c g a g a c g a t g t c a c a t a c 900
ctgcatgagg atgtgccctg gtggcgtggc gagggccaca agccgtctat tgcccacagg 960
gactttaaaa gtaagaatgt attgctgaag agcgcaccca cagccgtgct ggctgacttt 1020
ggcttggctg ttcgatttga gccaggaaa cttccagggg acaccccacgg acaggttaggc 1080
acgagacggt acatggctcc tgaggtgctc gagggagcca tcaacttcca gagagatgcc 1140
ttcctgcgca ttgacatgta tgccatgggg ttgggtgtgt gggagcttgc gtctcgctgc 1200
aaggctgcag acggaccctg ggatgagttac atgctgccct ttgaggaaga gattggccag 1260
cacccttcgt tggaggagct gcaggagggtg gtggtgacaca agaagatgag gcccaccatt 1320
aaagatcaact gttgaaaca cccggggcctg gcccagctt gtgtgaccat cgaggagtgc 1380

2007343333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

tgggaccatg atgcagaggc tcgttgtcc gcgggctgtg tggaggagcg ggtgtccctg 1440
attcggaggt cggtaacgg cactacctcg gactgtctcg ttccctggt gacctctgtc 1500
accaatgtgg acctgccccca taaagagtca agcatctaa 1539

<210> 19
<211> 348
<212> DNA
<213> Homo sapiens

<400> 19
tctgggcgtg gggaggctga gacacgggag tgcattact acaacgccaa ctgggagctg 60
gagcgcacca accagagcgg cctggagcgc tgcgaaggcg agcaggacaa gcggctgcac 120
tgctacgcct cctggccaa cagctctggc accatcgagc tcgtgaagaa gggctgctgg 180
ctagatgact tcaactgcta cgataggcag gagtgtgtgg ccactgagga gaaccccccag 240
gtgtacttct gctgctgtga aggcaacttc tgcaacgagc gcttcactca tttgccagag 300
gctgggggcc cggaaagtccac gtacgagcca ccccccacag ccccccacc 348

<210> 20
<211> 329
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 20
Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala
1 5 10 15

Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu
20 25 30

Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser
35 40 45

Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe
50 55 60

Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln
65 70 75 80

Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr
85 90 95

His Leu Pro Glu Ala Gly Gly Thr His Thr Cys Pro Pro Cys Pro
100 105 110

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
115 120 125

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
130 135 140

Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
145 150 155 160

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
165 170 175

Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
180 185 190

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
195 200 205

Ala Leu Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln

2007343333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt
210 215 220

Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
225 230 235 240
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
245 250 255
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
260 265 270
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
275 280 285
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
290 295 300
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
305 310 315 320
Lys Ser Leu Ser Leu Ser Pro Gly Lys
325

<210> 21
<211> 344
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 21
Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala
1 5 10 15
Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu
20 25 30
Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser
35 40 45
Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe
50 55 60
Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln
65 70 75 80
Val Tyr Phe Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr
85 90 95
His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro
100 105 110
Thr Ala Pro Thr Gly Gly Thr His Thr Cys Pro Pro Cys Pro Ala
115 120 125
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
130 135 140
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
145 150 155 160
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
165 170 175
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
180 185 190

200734333 28 Mar 2011

PHPH-022-w01_Substitute_Sequence_Listing.txt

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
195 200 205

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
210 215 220

Leu Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
225 230 235 240

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
245 250 255

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
260 265 270

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
275 280 285

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
290 295 300

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
305 310 315 320

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
325 330 335

Ser Leu Ser Leu Ser Pro Gly Lys
340

<210> 22
<211> 5
<212> PRT
<213> Artificial sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 22
Thr Gly Gly Gly Gly
1 5

<210> 23
<211> 5
<212> PRT
<213> Artificial sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 23
Ser Gly Gly Gly Gly
1 5