



(86) Date de dépôt PCT/PCT Filing Date: 2012/09/03
(87) Date publication PCT/PCT Publication Date: 2013/03/07
(45) Date de délivrance/Issue Date: 2021/09/07
(85) Entrée phase nationale/National Entry: 2014/02/28
(86) N° demande PCT/PCT Application No.: JP 2012/072383
(87) N° publication PCT/PCT Publication No.: 2013/032012
(30) Priorité/Priority: 2011/09/04 (JP2011-192203)

(51) Cl.Int./Int.Cl. *C07K 14/655* (2006.01),
A61K 38/00 (2006.01), *A61P 1/00* (2006.01),
A61P 1/04 (2006.01), *A61P 1/12* (2006.01),
A61P 19/02 (2006.01), *A61P 25/04* (2006.01),
A61P 25/28 (2006.01), *A61P 27/02* (2006.01),
A61P 27/06 (2006.01), *A61P 27/12* (2006.01),
A61P 29/00 (2006.01), ...

(72) Inventeurs/Inventors:
OCHIAI, HIROFUMI, JP;
SHIMODA, TAIJI, JP;
FUKAE, KAZUHIRO, JP;
MAEDA, MASATOSHI, JP;
...

(54) Titre : POLYPEPTIDE GLYCOSYLE ET COMPOSITION MEDICAMENTEUSE CONTENANT CE POLYPEPTIDE
(54) Title: GLYCOSYLATED POLYPEPTIDE AND DRUG COMPOSITION CONTAINING SAID POLYPEPTIDE

(57) Abrégé/Abstract:

[Problem] To provide a glycosylated polypeptide having affinity to somatostatin receptors and, compared to somatostatins, having improved in-blood stability. [Solution] The glycosylated polypeptide is characterized by at least two amino acids in a somatostatin or an analogue thereof being replaced by glycosylated amino acids.



(51) Cl.Int./Int.Cl. (suite/continued) *A61P 3/10* (2006.01), *A61P 35/00* (2006.01), *A61P 43/00* (2006.01),
A61P 5/02 (2006.01), *C07K 19/00* (2006.01), *C07K 2/00* (2006.01)

(72) Inventeurs(suite)/Inventors(continued): TAZURU, KEISUKE, JP; YOSHIDA, KENTA, JP

(73) Propriétaires(suite)/Owners(continued): GLYTECH, INC., JP

(74) Agent: MARKS & CLERK

Abstract

[Problem] To provide a glycosylated polypeptide having affinity to somatostatin receptors and, compared to somatostatins, having improved in-blood stability.

[Solution] The glycosylated polypeptide is characterized by at least two amino acids in a somatostatin or an analogue thereof being replaced by glycosylated amino acids.

Glycosylated Polypeptide and Drug Composition Containing said Polypeptide

Technical Field

[0001]

The present invention relates to a glycosylated polypeptide and a pharmaceutical composition comprising said polypeptide.

Background Art

[0002]

Somatostatin is a cyclic peptide present in both the central nervous system and the surrounding tissue. Somatostatin was first isolated from mammalian hypothalamus, and was identified from anterior pituitary gland as an important inhibitor of growth hormone secretion. This peptide is widely distributed in e.g. the hypothalamus, the pancreas, and the gastrointestinal tract, and its action is exerted via binding to a somatostatin receptor. In addition, somatostatin is known for its secretory suppression of growth hormone (GH) and thyroid-stimulating hormone (TSH) in the pituitary gland, as well as secretion suppression of various hormones such as gastrin, selectin, cholecystokinin (CCK), and VIP (Vasoactive Intestinal Polypeptide) in the gastrointestinal tract, and glucagon and insulin in the pancreas. It is also known to have an action to suppress gastrointestinal motility.

[0003]

Natural somatostatin having the structural formula:
Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys
(SEQ ID NO. 1) (also known as somatotropin release inhibiting factor (SRIF)) was first isolated by Guillemin and coworkers. This somatostatin exerts its effect by interacting with a family of receptors. Somatostatin receptor (SSTR) has 1 to 5 subtypes (SSTR1 - SSTR5), and among these, SSTR2 is known to be distributed in each tissues of the GH-secreting human pituitary gland adenoma, the central nervous system, the anterior pituitary gland, the retina, the adrenal medulla, the stomach, the duodenal mucosa, the small intestines, and the colon, as well as in the glucagon-secreting A-cell of the pancreatic islet. Each of these receptors is also known to be expressed in various tumors. For example, it has been reported that SSTR1 and SSTR5 are expressed in a functional pituitary adenoma, SSTR2 as well as SSTR1 and SSTR3 are expressed in a gastrointestinal tumor, SSTR3 is expressed in a pheochromocytoma, SSTR1 and SSTR5 are expressed in a prostate cancer, and SSTR5 is expressed in a colorectal

cancer (Non-Patent Literature 1). Moreover, SSTR4 is reported in regards to its function as a receptor having an antagonistically modulating action and its possibility of being essential in the treatment of a glaucoma-related disease (Non-Patent Literature 2). As such, somatostatin and analogs thereof are potentially useful therapeutic drugs for somatostatin-related diseases or various types of tumors.

[0004]

Meanwhile, because naturally-occurring somatostatin has a short half-life in blood of 2 - 3 minutes, it shows two undesirable properties of having low bioavailability and short duration of action, and thus its use or application as a therapeutic is limited. For this reason, various somatostatin analogs have been developed in order to find a somatostatin analog superior in any one of efficacy, biostability, duration of action, or selectivity considering the release suppression of growth hormone, insulin, or glucagon.

[0005]

Octreotide (Patent Literatures 1 and 2) is reported as the first approved somatostatin analog that can be clinically utilized, and this octreotide is known to have affinity towards somatostatin receptors SSTR2, SSTR3, and SSTR5.

Octreotide has been developed as a cyclic peptide consisting of eight amino acids which has a sequence of four amino acids (Phe-Trp-Lys-Thr) that is an important portion for showing the biological activity of somatostatin, Cys that forms a disulfide (S-S) bond at the two terminals of the sequence, and further D-Phe and Thr(ol) outside of the Cys at the two terminals. This octreotide can render persistence of action by improving the half-life in blood by its amino acid sequence, as well as has a higher selectivity towards growth hormone (GH) than somatostatin which enables it to have a strong action.

[0006]

Such somatostatin analogs including octreotide can be used for treating patients who have a hormone-secreting and hormone-dependent tumor. Currently, symptoms related to metastatic carcinoid tumor which is a tumor of the neuroendocrine system (flushing, diarrhea, cardiac valve disease, and abdominal pain) and symptoms related to vasoactive intestinal peptide (VIP)-secreting adenoma (watery diarrhea) are treated with octreotide.

For example, in a carcinoid and VIP-producing tumor, octreotide inhibits both secretion and action of its active factor. Accordingly, in a VIP-producing tumor

characterized in profusely-secreting diarrhea, a somatostatin analog can reduce its diarrhea by secretory inhibition of VIP as well as by directly influencing intestinal secretion.

[0007]

On the other hand, however, many neuroendocrine tumors are reported to have resistance to somatostatin analogs such as octreotide (Non-Patent Literature 3). Moreover, although octreotide is used in the treatment of acromegaly, it is reported to have no effect on approximately one third of acromegaly patients. Further, it is reported that in majority of carcinoid tumor patients, octreotide exerts its effect only during initial administration, and tachyphylaxis is caused when the duration of administration is prolonged. Further, it is reported that octreotide does not show any effect on suppression of adrenocorticotrophic hormone (ACTH) production in early Cushing's disease patients.

[0008]

In light of the problems above, development of a somatostatin analog that binds to multiple receptor subtypes with high affinity like that of a natural somatostatin is desired for a tumor expressing multiple somatostatin receptors, and it is suggested that a somatostatin analog having such affinity towards somatostatin receptors may possibly also have effect on patients who were therapeutically ineffective with or patients who have resistance to past somatostatin analogs (Non-Patent Literature 4).

Accordingly, development of a somatostatin analog having a structure similar to a naturally-occurring somatostatin, similarly having affinity towards somatostatin receptors, and having extended half-life in blood compared to somatostatin has been desired.

[0009]

Meanwhile, it has been becoming clear that sugar chains are responsible for various roles *in vivo*, and a method for adding a sugar chain to octreotide in order to extend the half-life in blood has also been proposed (such as Patent Literature 3).

However, research is delayed due to the complexity or diversity of its structure, and it cannot be said that the type of sugar chain or the position for adding a sugar chain is always optimized. A glycosylated polypeptide that has overcome the problems of past somatostatin analogs has not been reported.

Citation List

[Patent Literatures]

[0010]

[Patent Literature 1] U. S. Patent No. 4, 310, 518

[Patent Literature 2] U. S. Patent No. 4, 235, 886

[Patent Literature 3] Japanese Published Unexamined Patent Application Publication No. Hei 03 (1991)-014599

[Non-Patent Literatures]

[0011]

[Non-Patent Literature 1] Current Opinion in Pharmacology, 2004, Vol. 4, pp. 608-613

[Non-Patent Literature 2] J. Med. Chem. 2003, Vol. 46, pp. 5587-5596

[Non-Patent Literature 3] Mol. Endocrinol., 2010, Vol. 24 (1), pp. 240-249

[Non-Patent Literature 4] Molecular and Cellular Endocrinology, Vol. 286, 2008, pp. 69-74

Summary of the Invention

Problems to be Solved by the Invention

[0012]

The object of the present invention is to provide a glycosylated polypeptide having affinity towards somatostatin receptors and improved stability in blood compared to somatostatin.

Means for Solving the Problems

[0013]

As a result of repeated research to solve the above problems, the present inventors found a glycosylated polypeptide which maintains affinity towards somatostatin receptors and has improved stability in blood.

In other words, the present invention relates to a glycosylated polypeptide selected from the group consisting of: (A) SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1; (B) a polypeptide having one or a few amino acids deleted, substituted, or added from/to SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1; (C) a SRIF14 analog; (D) a polypeptide having 80% or more homology with SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1; (E) a polypeptide further comprising N amino acids (wherein N is an integer from 1 or more to 20 or less) at the N-terminal side of (A) - (D); and (F) a polypeptide further comprising M amino acids (wherein M is an integer from 1 or more to 6 or less) at the C-terminal side of (A) - (D); characterized in that at least two amino acids are substituted with glycosylated amino acids, and the polypeptide has affinity towards somatostatin receptors.

[0014]

Here, one embodiment of the glycosylated polypeptide of the present invention is characterized in that at least one of the amino acids substituted with said glycosylated amino acid is the amino acid corresponding to position 19 of SRIF14.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that at least two amino acids are substituted with glycosylated amino acids in said polypeptide (E), and at least one of the amino acids substituted with said glycosylated amino acid is present at any of said N amino acids at the N-terminal side of said polypeptide (E).

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that at least two amino acids are substituted with glycosylated amino acids in said polypeptide (F), and at least one of the amino acids substituted with said glycosylated amino acid is present at any of said M amino acids at the C-terminal side of said polypeptide (F).

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that at least two amino acids are substituted with glycosylated amino acids in said polypeptide (E), and further the sequence of said N amino acids added onto the N-terminal side is represented by X-Y-, wherein X means a sequence of any L amino acids (wherein L is an integer from 1 or more to 6 or less), and Y is a sequence selected from the group consisting of: (1) Lys, (2) Arg-Lys, (3) Glu-Arg-Lys, (4) Arg-Glu-Arg-Lys, (5) Pro-Arg-Glu-Arg-Lys, (6) Ala-Pro-Arg-Glu-Arg-Lys, (7) Met-Ala-Pro-Arg-Glu-Arg-Lys, (8) Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (9) Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (10) Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (11) Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (12) Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (13) Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, (14) Ser-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, and (15) a sequence having one or a few amino acids deleted, substituted, or added from/to the above sequences (2) - (14).

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that at least one of the amino acids substituted with said glycosylated amino acid is present in any of L amino acids which means X in X-Y- representing the sequence of said N

amino acids added onto the N-terminal side of said polypeptide (E).

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that the N amino acids added onto the N-terminal side are linked to said N-terminal side via a linker.
[0015]

Moreover, in another embodiment of the glycosylated polypeptide of the present invention, a polypeptide selected from the group consisting of: (A) SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2; (B) a polypeptide having one or a few amino acids deleted, substituted, or added from/to SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2; (C) a SRIF28 analog; (D) a polypeptide having 80% or more homology with SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2; (E) a polypeptide further comprising J amino acids (wherein J is an integer from 1 or more to 15 or less) at the N-terminal side of (A) - (D); and (F) a polypeptide further comprising K amino acids (wherein K is an integer from 1 or more to 6 or less) at the C-terminal side of (A) - (D); is characterized in that at least two amino acids are substituted with glycosylated amino acids, and the polypeptide has affinity towards somatostatin receptors.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that at least one of the amino acids substituted with said glycosylated amino acid is at least one amino acid selected from the group consisting of the amino acid corresponding to position 1, the amino acid corresponding to position 5, the amino acid corresponding to position 9, the amino acid corresponding to position 12, the amino acid corresponding to position 13, the amino acid corresponding to position 14, and the amino acid corresponding to position 19 of SRIF28.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that at least two amino acids are substituted with glycosylated amino acids in said polypeptide (E), wherein at least one of the amino acids substituted with said glycosylated amino acid is present in said J amino acids at the N-terminal side of said polypeptide (E).

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that at least two amino acids are substituted with glycosylated amino acids in said polypeptide (F), and at least one of the amino acids

substituted with said glycosylated amino acid is present in said K amino acids of the C-terminal side of said polypeptide (F).

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has increased stability in blood compared to SRIF28.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has increased half-life in blood by 10-fold or more compared to SRIF28.

[0016]

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that said affinity towards somatostatin receptors has affinity towards at least two or more receptors selected from the group consisting of SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has affinity towards any one of at least SSTR1 and SSTR4.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has affinity towards both SSTR1 and SSTR4.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has affinity towards all of SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that each of said glycosylated amino acids is glycosylated Asn or glycosylated Cys.

[0017]

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that in each of said glycosylated amino acids, the sugar chain and the amino acid are linked without a linker.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that in each of said glycosylated amino acids, the sugar chain consists of 4 or more sugars.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that in each of said glycosylated amino acids, the sugar chain is a biantennary complex-type sugar chain, a


triantennary complex-type sugar chain, or a tetraantennary complex-type sugar chain.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that in each of said glycosylated amino acids, the sugar chain is a biantennary complex-type sugar chain.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that said biantennary complex-type sugar chain is a sugar chain selected from the group consisting of a disialo sugar chain, a monosialo sugar chain, an asialo sugar chain, a diGlcNAc sugar chain, and a dimannose sugar chain.


Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that in each of said glycosylated amino acids, the sugar chain is a sugar chain represented by the following formula:

[Chemical Formula 1]



[wherein R¹ and R² are identical or different and are:

[Chemical Formula 2]



, and Ac is an acetyl group].

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that said sugar chain has at least one sialic acid at the

non-reducing terminal, and the carboxy group of said sialic acid is modified by an alkylamino group, a benzyl group, an amino group, or an aminoethylamino group, having 1 - 30 carbons.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has multiple glycosylated amino acids, wherein the sugar chains on each of said glycosylated amino acids are all identical.

Moreover, in one embodiment of the glycosylated polypeptide of the present invention, said glycosylated polypeptide is characterized in that it has Cys corresponding to Cys at position 17 and Cys at position 28 of SRIF28, and further these Cys are bound by a disulfide bond with each other.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that the C-terminal of said glycosylated polypeptide is amidated.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that an azido group is introduced at the N-terminal of said glycosylated polypeptide.

Moreover, one embodiment of the glycosylated polypeptide of the present invention is characterized in that it is labeled.

[0018]

Moreover, another aspect of the present invention relates to a pharmaceutical composition characterized in that it comprises (I) the glycosylated polypeptide described above and/or a pharmaceutically acceptable salt thereof, and (II) a pharmaceutically acceptable carrier.

Moreover, one embodiment of the pharmaceutical composition of the present invention is characterized in that the sugar chains in said glycosylated polypeptide are substantially uniform.

Moreover, one embodiment of the pharmaceutical composition of the present invention is characterized in that it is employed for treatment or prevention of a somatostatin-related disease.

Moreover, one embodiment of the pharmaceutical composition of the present invention is characterized in that said somatostatin-related disease is at least one disease selected from the group consisting of acromegaly, gigantism, Alzheimer's disease and other forms of dementia, cancer, hormone-producing tumor, endocrine tumor, carcinoid, VIPoma, insulinoma, glucagonoma, Cushing's disease, hormone secretion defect, diabetes and complications thereof, pains,

arthritis, diarrhea, gastric ulcer, inflammatory bowel disease, irritable bowel syndrome, gastrointestinal obstruction, ileus, postoperative restenosis, radiation damage, eye disease, dry eye, glaucoma, interstitial keratitis, iritis, cataract, and conjunctivitis.

[0019]

Moreover, another aspect of the present invention relates to a method for treating or preventing a somatostatin-related disease, characterized in administering an effective amount of the glycosylated polypeptide described above.

Moreover, one embodiment of the treatment or prophylactic method of the present invention is characterized in that said somatostatin-related disease is at least one disease selected from the group consisting of acromegaly, gigantism, Alzheimer's disease and other forms of dementia, cancer, hormone-producing tumor, endocrine tumor, carcinoid, VIPoma, insulinoma, glucagonoma, Cushing's disease, hormone secretion defect, diabetes and complications thereof, pains, arthritis, diarrhea, gastric ulcer, inflammatory bowel disease, irritable bowel syndrome, gastrointestinal obstruction, ileus, postoperative restenosis, radiation damage, eye disease, dry eye, glaucoma, interstitial keratitis, iritis, cataract, and conjunctivitis.

Effects of the Invention

[0020]

The glycosylated polypeptide of the present invention has affinity towards somatostatin receptors, and has significantly improved stability in blood compared to somatostatin by having at least two sugar chains in the polypeptide. Moreover, the glycosylated polypeptide of the present invention can be employed for treating a somatostatin-related disease by virtue of having the above characteristics.

[0021]

Moreover, since the sugar chain added onto the glycosylated somatostatin of the present invention is easily degraded *in vivo*, no drug-induced suffering is caused to a living body by accumulation thereof.

[0022]

Moreover, a part or all of the sugar chains added onto the glycosylated somatostatin of the present invention is a sugar chain present in the living body of e.g. mammals and birds including humans or a modified version thereof, and the possibility of showing side effects or antigenicity when administered *in vivo* is low. There is less concern

for e.g. an allergic reactions or antibody production to occur and thereby losing drug effect.

[0023]

Further, since many of the sugar chains employed herein are relatively short, those having uniform structure can be obtained without going through complex manufacturing steps. Accordingly, a pharmaceutical grade high quality glycosylated polypeptide having somatostatin activity can be stably obtained in a large scale.

Brief Description of the Drawings

[0024]

Figures 1A - 1D show each compound name of the glycosylated polypeptides of an embodiment of the present invention and the amino acid sequence information of the glycosylated polypeptide having said compound name. Moreover, Figure 1E is a graph showing the IC₅₀ value in regards to cAMP production suppressing action (agonist activity) when somatostatin receptor expression cells were treated with SRIF14, SRIF28, and the glycosylated polypeptides of an embodiment of the present invention.

Figure 2 is graphs showing the plasma concentration transition of each polypeptide when SRIF28 and S1C(disialo)-SRIF28 were administered intravenously or subcutaneously to rats. In Figure 2, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 3 is graphs showing the plasma concentration transition when SRIF28, S1C(disialo)-SRIF28, N5C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 3, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 4 is graphs showing the plasma concentration transition when SRIF28, S1C(disialo)-SRIF28, S1C(disialo)·R13C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 4, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a

graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 5 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28, 3C(disialo)-SRIF28, 5C(disialo)-SRIF28, and 10C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 5, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of the polypeptides when administered subcutaneously.

Figure 6 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, N5C(disialo)-SRIF28, A9C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, N5C(disialo)·A9C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 6, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 7 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, and S1-3C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 7, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 8 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1C(disialo(amide))-SRIF28, and S1C(disialo(aminoethylamide))-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 8, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 9 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1C(disialo)-D-Trp22-SRIF28, and S1C(disialo(Bn))-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 9, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a

graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 10 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, R13C(disialo)-SRIF28, and K14C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 10, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 11 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, E12C(disialo)-SRIF28, N19C(disialo)-SRIF28, 29C(disialo)-SRIF28, S1C(monosialo)-SRIF28, and S1C(asialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 11, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 12 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, K14C(disialo)-SRIF28, and C(disialo)-SRIF14 were administered intravenously and subcutaneously to rats. In Figure 12, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 13 is graphs showing the plasma concentration transition when C(disialo)-SRIF14, C(disialo)-C12 linker-SRIF14, and C(disialo)-PEG linker-SRIF14 were administered intravenously and subcutaneously to rats. In Figure 13, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 14 is graphs showing the plasma concentration transition when SRIF28, S1C(disialo)-SRIF28, S1C(asialo)-SRIF28, S1C(diGlcNAc)-SRIF28, S1C(diMan)-SRIF28, and S1C(GlcNAc)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 14, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma

concentration transition of each polypeptide when administered subcutaneously.

Figure 15 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1C(trisialo)-SRIF28, S1C(tetrasialo)-SRIF28, S1-2C(disialo)-SRIF28, and S1C(Asn(disialo))-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 15, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 16 is graphs showing the plasma concentration transition when SRIF28, S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, S1-3C(disialo)-SRIF28, and S1-4C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 16, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 17 is graphs showing the plasma concentration transition when SRIF14, C(disialo)-SRIF14, C(disialo)-K-SRIF14, C(disialo)-R-K-SRIF14, 2C(disialo)-R-K-SRIF14, and 3C(disialo)-R-K-SRIF14 were administered intravenously and subcutaneously to rats. In Figure 17, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 18 is graphs showing the plasma concentration transition when S1C(asialo)-SRIF28, S1-2C(asialo)-SRIF28, and S1-3C(asialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 18, the graph on the left is a graph showing the plasma concentration transition of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 19 is graphs showing the plasma concentration transition when S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, S1-2C(asialo)-SRIF28, S1-2C(disialo(amide))-SRIF28, S1-2C(disialo(Bn))-SRIF28, and C(disialo(aminoethylamide))-S1C(disialo)-SRIF28 were administered intravenously and subcutaneously to rats. In Figure 19, the graph on the left is a graph showing the plasma concentration transition

of each polypeptide when administered intravenously, and the graph on the right is a graph showing the plasma concentration transition of each polypeptide when administered subcutaneously.

Figure 20 is a graph showing the result of plasma stability test employing rat plasma for the glycosylated polypeptides of an embodiment of the present invention.

Description of Embodiments

[0025]

A "Somatostatin" herein refers to SRIF14 consisting of a sequence of 14 amino acids or SRIF28 consisting of a sequence of 28 amino acids.

In the present specification, the N-terminal Ser in the SRIF28 amino acid sequence will be set to as position 1, and the C-terminal Cys as position 28. SRIF14 is in perfect match with the amino acid sequence of positions 15 to 28 in the SRIF28 amino acid sequence. Note that position 15 of the SRIF28 amino acid sequence is Ala, and the N-terminal Ala in the SRIF14 amino acid sequence (SEQ ID NO. 1) will be set as position 15 in correspondence with position 15 of SRIF28. SRIF14 and SRIF28 have a disulfide bond at Cys at position 17 and Cys at position 28.

[0026]

SRIF14 has the amino acid sequence below (SEQ ID NO. 1). In the amino acid sequence below, 15 in "Ala₁₅" means Ala at position 15.

Ala₁₅-Gly₁₆-Cys₁₇-Lys₁₈-Asn₁₉-Phe₂₀-Phe₂₁-Trp₂₂-Lys₂₃-Thr₂₄-Phe₂₅-Thr₂₆-Ser₂₇-Cys₂₈

[0027]

SRIF28 has the amino acid sequence below (SEQ ID NO. 2).

Ser₁-Ala₂-Asn₃-Ser₄-Asn₅-Pro₆-Ala₇-Met₈-Ala₉-Pro₁₀-Arg₁₁-Glu₁₂-Arg₁₃-Lys₁₄-Ala₁₅-Gly₁₆-Cys₁₇-Lys₁₈-Asn₁₉-Phe₂₀-Phe₂₁-Trp₂₂-Lys₂₃-Thr₂₄-Phe₂₅-Thr₂₆-Ser₂₇-Cys₂₈

[0028]

An "amino acid" herein is employed in its broadest meaning, and includes not only natural amino acids but also non-natural amino acids such as amino acid variants and derivatives. Those skilled in the art will recognize in light of this broad definition that examples of amino acids herein include, e.g., natural proteinogenic L-amino acids; D-amino acids; chemically modified amino acids such as amino acid variants and derivatives; natural non-proteinogenic amino acids such as norleucine, β -alanine, and ornithine; and chemically synthesized compounds having properties well-known in the art characteristic of amino acids. Examples of non-natural amino acids include an α -

methylamino acid (such as α -methylalanine), a D-amino acid, a histidine-like amino acid (such as 2-amino-histidine, β -hydroxy-histidine, homohistidine, α -fluoromethyl-histidine, and α -methyl-histidine), an amino acid having excess methylenes on the side chain ("homo" amino acid), and an amino acid in which the carboxylic functional group amino acid in the side chain is substituted with a sulfonate group (such as cysteic acid). Some of the somatostatin analogs having affinity towards somatostatin receptors are known to comprise a non-natural amino acid. In a preferred aspect, the amino acid contained in the compound of the present invention consists only of natural amino acids.

[0029]

As used herein, when one of the amino acids or a few amino acids are said to be deleted, substituted, or added, the number of amino acids substituted etc. is not particularly limited as long as affinity towards somatostatin receptors is retained, but is 1 - 9, preferably 1 - 5, and more preferably approximately 1 - 3 amino acids, or 20% or less and preferably 10% or less of the entire length. The amino acid to be substituted or added may be a natural amino acid, a non-natural amino acid, or an amino acid analog, preferably a natural amino acid. Examples of somatostatin peptides having one of the amino acids or a few amino acids deleted, substituted, or added include, e.g., a somatostatin peptide having Trp at position 22 substituted with a D-form Trp (D-Trp), Asn at position 19 deleted (J. Med. Chem., 2001, 44, 2238-2246), Phe at position 25 substituted with Tyr, and Met at position 8 substituted with Leu (Endocrinology, 1982, 10:1049-1051).

[0030]

A "SRIF14 or SRIF28 analog" herein includes a polypeptide structurally similar to somatostatin and/or a polypeptide having an overlapping structure with somatostatin, e.g. a polypeptide having one of the amino acids or a few amino acids of somatostatin conservatively substituted, a modified somatostatin, a somatostatin fragment having affinity towards somatostatin receptors, and an elongated somatostatin having affinity towards somatostatin receptors.

[0031]

"Having one of the amino acids or a few amino acids conservatively substituted" herein refers to an amino acid substitution in which the hydrophilicity and/or hydrophobicity index are similar between the original amino acid and the amino acid to be substituted, and wherein apparent reduction or dissipation of affinity towards

somatostatin receptors before and after such substitution is not caused.

[0032]

A "modified somatostatin" herein is a modified version of somatostatin including a naturally-occurring variant of somatostatin or an artificially modified compound of somatostatin. Examples of such modifications include, e.g., alkylation, acylation (such as acetylation), amidation, carboxylation, ester formation, disulfide bond formation, glycosylation, lipidation, phosphorylation, hydroxylation, and binding of a labeling component of one or more amino acid residues of somatostatin.

[0033]

A "somatostatin fragment having affinity towards somatostatin receptors" herein is a peptide having one or more amino acids deleted from the N- and/or C-terminals of somatostatin which maintains affinity towards somatostatin receptors.

[0034]

An "elongated somatostatin having affinity towards somatostatin receptors" herein is a peptide having one or more amino acids added to the N- and/or C-terminals of SRIF28 or SRIF14 which maintains affinity towards somatostatin receptors.

[0035]

The glycosylated polypeptide of the present invention comprises a glycosylated polypeptide which is a polypeptide consisting of an amino acid sequence having 80% or more homology with the amino acid sequence represented by SEQ ID NO. 1; a polypeptide consisting of an amino acid sequence having 80% or more homology with the amino acid sequence represented by SEQ ID NO. 2; or a polypeptide having an amino acid further added to the N- or C-terminal of these polypeptides; wherein at least one amino acid is substituted with a glycosylated amino acid, and the polypeptide has affinity towards somatostatin receptors.

A polypeptide having homology with SEQ ID NO. 1 or 2 can have preferably 80% or more, 85% or more, 90% or more, and 95% or more homology, as long as it has affinity towards somatostatin receptors.

[0036]

The glycosylated polypeptide of the present invention also comprises a polypeptide having an amino acid further added to the N- and/or C-terminals of SRIF14 or SRIF28 as described above.

In the glycosylated polypeptide of the present invention, the amino acid further added to the N-terminal of SRIF14 is not particularly limited as long as it

maintains affinity towards somatostatin receptors. For example, 1 or more to 20 or less amino acids can be added.

Here, the amino acid further added to the N-terminal of SRIF14 can be represented by X-Y-. Y is an amino acid that binds directly to the N-terminal amino acid of SRIF14 polypeptide, and Y consists of any of amino acid sequences

(1) - (15) below:

- (1) Lys,
- (2) Arg-Lys,
- (3) Glu-Arg-Lys,
- (4) Arg-Glu-Arg-Lys,
- (5) Pro-Arg-Glu-Arg-Lys,
- (6) Ala-Pro-Arg-Glu-Arg-Lys,
- (7) Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (8) Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (9) Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (10) Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (11) Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (12) Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (13) Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (14) Ser-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, and
- (15) a sequence having one or a few amino acids deleted, substituted, or added from/to the above sequences (2) - (14).

Moreover, X is any amino acid from 1 or more to 6 or less, and shows any 1, 2, 3, 4, 5, or 6 amino acids. Preferably, X is a glycosylated amino acid, and more preferably glycosylated Asn or glycosylated Cys.

[0037]

In the glycosylated polypeptide of the present invention, the amino acid further added to the N-terminal of SRIF28 is not particularly limited as long as it maintains affinity towards somatostatin receptors. For example, any amino acids from 1 or more to 15 or less can be added, and any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acids can be added. The amino acid further added to the N-terminal of SRIF28 is preferably a glycosylated amino acid, and more preferably glycosylated Asn or glycosylated Cys. Moreover, all of the any 1 or more to 15 or less amino acids can be glycosylated amino acids.

[0038]

In the glycosylated polypeptide of the present invention, the amino acid further added to the C-terminal of SRIF14 or SRIF28 is not particularly limited as long as it maintains affinity towards somatostatin receptors. For

example, any amino acids from 1 or more to 6 or less can be added, and any 1, 2, 3, 4, 5, or 6 amino acids can be added. The amino acid further added to the C-terminal of SRIF14 or SRIF28 is preferably a glycosylated amino acid, and more preferably glycosylated Asn or glycosylated Cys. Moreover, all of the any 1 or more to 6 or less amino acids can be glycosylated amino acids.

[0039]

A "peptide having one or a few amino acids further added to the N-terminal side of somatostatin (position 1 of SRIF28 or position 15 of SRIF14)" herein refers to, in the case of SRIF28, those having any amino acid or glycosylated amino acid further added to the N-terminal Ser at position 1. Moreover, in the case of SRIF14, it refers to those having any amino acid or glycosylated amino acid further added to the N-terminal Ala at position 15.

Similarly, a "peptide having one or a few amino acids further added to the C-terminal side (position 28 of SRIF28 or SRIF14)" refers to those having any amino acid or glycosylated amino acid further added to Cys at position 28 of SRIF28 or SRIF14.

[0040]

An amino acid can also be further added to the glycosylated polypeptide of the present invention at its N- or C-terminal via a linker. Examples of such linkers can include, e.g., an alkyl chain or a polyethylene glycol chain having amino and carboxy groups at the two terminals so that it can form a peptide bond with the amino acid. Examples of such linkers can include, e.g., $\text{-NH-(CH}_2\text{)}_n\text{-CO-}$ (wherein n is an integer and is not limited as long as it does not inhibit the linker function of interest, but is preferably an integer from 1 - 15) or $\text{-NH-(CH}_2\text{CH}_2\text{O)}_m\text{-CH}_2\text{CH}_2\text{-CO-}$ (wherein m is an integer and is not limited as long as it does not inhibit the linker function of interest, but is preferably an integer from 1 - 7). More specifically, examples can include $\text{-NH-(CH}_2\text{)}_{11}\text{-CO-}$ (C12 linker) or $\text{-NH-(CH}_2\text{CH}_2\text{O)}_3\text{-CH}_2\text{CH}_2\text{-CO-}$ (PEG linker). Moreover, the amino acid added to the glycosylated polypeptide via a linker is not particularly limited, but is preferably a glycosylated amino acid. Examples of a glycosylated amino acid can include glycosylated Asn or glycosylated Cys.

[0041]

Moreover, in one embodiment of the present invention, an azido group can also be introduced into the glycosylated polypeptide at its N-terminal. Azidation of the N-terminal of the glycosylated polypeptide is preferred because it will allow various molecules to be selectively introduced by utilizing azide-alkyne [3+2] cycloaddition reaction or

Staudinger reaction. The method for azidating the glycosylated polypeptide is not particularly limited, but it can be obtained by e.g. condensing the N-terminal of a glycopeptide synthesized on a resin in solid phase synthesis and an azido-substituted fatty acid with a condensation agent. Azido-substituted fatty acids can include 4-azidobutanoic acid, 5-azidopentanoic acid, and 6-azidohexanoic acid.

[0042]

Moreover, in one embodiment of the present invention, a labeling compound can also be added to the glycosylated polypeptide at its N-terminal. Examples of the labeling compound used herein can include, but is not limited to, e.g. biotin, fluorescent dyes, and metal ion chelators. The labeling compound can be directly bound to the N-terminal of the glycopeptide, or it can also be bound to the N-terminal of the glycopeptide via a linker. For example, by adding biotin as the labeling compound to the N-terminal of the glycopeptide, strong binding with avidin can be utilized to enable application as research reagents, clinical test agents, or missile therapy.

The addition of a labeling compound to the glycosylated polypeptide of the present invention can be performed by a conventional method well-known to those skilled in the art. For example, the N-terminal of the glycosylated polypeptide on the resin in solid phase synthesis and a labeling compound can be condensed with a condensation agent.

[0043]

The "glycosylated polypeptide" of the present invention is characterized in that at least two amino acids are substituted by glycosylated amino acids.

A "glycosylated polypeptide" herein includes e.g. a polypeptide having at least two amino acids of somatostatin substituted with glycosylated amino acids and a polypeptide having at least two amino acids substituted with glycosylated amino acids in the above somatostatin analog, each of which is included in the glycosylated polypeptide of the present invention even if one of the amino acids or a few amino acids other than the glycosylated amino acid is further deleted, substituted, or added. A peptide having at least two amino acids substituted with glycosylated amino acids in a peptide in which the C-terminal thereof is amidated (such as SRIF14NH₂ having the amino acid sequence of Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys-NH₂ (SEQ ID NO. 3) or SRIF28NH₂ having the amino acid sequence of Ser-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-

Thr-Ser-Cys-NH₂ (SEQ ID NO. 4) is also included in the glycosylated polypeptide of the present invention. Further, a salt of these peptides is also included in glycosylated polypeptides.

[0044]

A salt as used herein may be any of acid addition salts or base addition salts. Acids generally employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, and phosphoric acid, as well as organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenylsulfonic acid, carboxylic acid, succinic acid, citric acid, benzoic acid, and acetic acid. Base addition salts include salts derived from an inorganic base such as ammonium hydroxide or alkali or alkaline earth metal hydroxides, carbonates, and bicarbonates. In particular, pharmaceutically acceptable salts are preferred.

[0045]

A "glycosylated (sugar chain attached) amino acid" herein is an amino acid having a sugar chain bound thereto, and the sugar chain and the amino acid may be bound via a linker. The binding site of the sugar chain and the amino acid is not particularly restricted, but the amino acid is preferably bound to the reducing terminal of the sugar chain.

The type of amino acid to which the sugar chain binds is not particularly limited, and any of natural amino acids and non-natural amino acids can be employed. With respect to the glycosylated amino acids having identical or similar structure as those present as glycopeptides (glycoproteins) *in vivo*, the glycosylated amino acid is preferably glycosylated Asn, like N-linked sugar chain as well as glycosylated Ser and glycosylated Thr, and glycosylated Asn, like O-linked sugar chain is particularly preferred.

[0046]

Moreover, when the sugar chain and the amino acid are bound via a linker, with respect to easy binding with the linker, the amino acid of the glycosylated amino acid is preferably an amino acid having two or more carboxy groups in a molecule such as aspartic acid and glutamic acid, an amino acid having two or more amino groups in a molecule such as lysine, arginine, histidine, and tryptophan, an amino acid having a hydroxyl group in the molecule such as serine, threonine, and tyrosine, an amino acid having a thiol group in the molecule such as cysteine, and an amino acid having an amide group in the molecule such as asparagine and glutamine. In particular, with respect to reactivity, aspartic acid, glutamic acid, lysine, arginine,

serine, threonine, cysteine, asparagine, and glutamine are preferred.

[0047]

In any glycosylated polypeptide of the present invention, it is thought that if the sugar chain structure, the structure other than the sugar chain, the number of addition sites for the sugar chain, and the number of sugar chains added are identical, there is no major difference in the half-life in blood of the glycosylated polypeptide of the present invention whether the glycosylated amino acid is glycosylated Asn (without a linker) or glycosylated Cys (with a linker).

[0048]

When the sugar chain and the amino acid are bound via a linker, linkers employed in the art can be widely used, examples of which can include, e.g., $-\text{NH}-(\text{CO})-(\text{CH}_2)_a-\text{CH}_2-$ (wherein a is an integer and is not limited as long as it does not inhibit the linker function of interest, but is preferably an integer from 0 - 4), C_{1-10} polymethylene, $-\text{CH}_2-\text{R}-$ (wherein R is a group produced by having one hydrogen atom detached from a group selected from the group consisting of an alkyl, a substituted alkyl, an alkenyl, a substituted alkenyl, an alkynyl, a substituted alkynyl, an aryl, a substituted aryl, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, and a substituted heterocyclic group), and $-(\text{CO})-(\text{CH}_2)_a-(\text{CO})-$ (wherein a is an integer and is not limited as long as it does not inhibit the linker function of interest, but is preferably an integer from 0 - 4).

[0049]

In the glycosylated amino acid, when the sugar chain and the amino acid on the somatostatin backbone are bound via a linker, it is preferred that the linker also comprises an amino acid at the sugar chain terminal. The type of amino acid is not particularly limited, but a preferred example can include Asn.

[0050]

The manufacturing method of the glycosylated polypeptide of the present invention is not to be limited in any way by a description therefor (such as the description a "glycosylated polypeptide having an amino acid substituted with a glycosylated amino acid"), and a glycosylated polypeptide manufactured with any of methods A or B described below is included in the "glycosylated polypeptide having an amino acid substituted with a glycosylated amino acid". Moreover, for example, a glycosylated polypeptide in which a sugar chain without any amino acid bound thereto is bound directly or via a linker

to an amino acid on a peptide; a glycosylated polypeptide in which a sugar or a sugar chain is further added to the sugar chain added in the glycosylated polypeptide in order to elongate the already added sugar chain; a glycosylated polypeptide in which one or a few amino acids are bound to the amino and/or carboxy group of a glycosylated amino acid, and further linked to one or more somatostatin fragments; and a glycosylated polypeptide in which a sugar chain having an amino acid bound thereto is bound via a linker to an amino acid on a peptide are also included in the glycosylated polypeptide of the present invention, as long as the final structure matches.

[0051]

In the glycosylated polypeptide of the present invention, the number of amino acids to be substituted with a glycosylated amino acid may be appropriately adjusted by e.g. bioactivity such as stability in blood or secretory suppression of growth hormone etc., the number of amino acids present in the final glycosylated polypeptide, or the molecular weight of the glycosylated polypeptide before and after glycosylation. For example, the glycosylated polypeptide of the present invention can have 2 - 15 amino acids in its amino acid sequence substituted to glycosylated amino acids. With respect to stability in blood, as long as the desired activity is obtained, it is preferred to substitute 2 or more, e.g. it is preferred to substitute 2 - 10, and more preferably to substitute 2 - 3 amino acids. In general, in a glycosylated polypeptide having one amino acid of somatostatin substituted with a glycosylated amino acid, stability in blood will increase when one or more of amino acids other than the glycosylated amino acid are further substituted by glycosylated amino acids. On the other hand, although affinity towards somatostatin receptors will tend to decrease, the stability in blood of the glycosylated polypeptide will increase, and it is therefore possible to compensate for or increase the decreased somatostatin activity.

Moreover, when multiple sugar chains are present in a glycosylated polypeptide, the each sugar chains can be added to consecutive amino acids, or can be added to amino acids with intervals in the amino acid sequence of the glycosylated polypeptide. Placing the sugar chains densely is preferred because there is no rapid increase in plasma concentration. When the sugar chains are placed densely, for example, approximately 2 - 15 consecutive glycosylated amino acids can be added to the N-terminal of SRIF14 or SRIF28. Moreover, with respect to bioavailability, the preferred glycosylation position is adding multiple sugar

chains with intervals rather than adding densely to consecutive amino acids.

[0052]

In the glycosylated polypeptide of the present invention, the site for substituting an amino acid with a glycosylated amino acid can be appropriately adjusted with respect to having affinity towards at least one somatostatin receptor, and preferably, it can be appropriately adjusted with respect to having affinity towards multiple somatostatin receptors.

[0053]

In one aspect of the present invention, the site for substituting an amino acid with a glycosylated amino acid can include, with respect to having affinity towards multiple receptors in SSTR1 - SSTR5 among affinity of the glycosylated polypeptide towards somatostatin receptors, e.g., the amino acid corresponding to position 19 of SRIF14, the amino acids added at the 1st, 2nd, 3rd, 6th, 10th, and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14, and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal side of SRIF14. Preferably, it can include the amino acids added at the 3rd, 6th, 10th, and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14.

Moreover, similarly with respect to having affinity towards multiple receptors in SSTR1 - SSTR5, it can include the amino acids corresponding to positions 1, 5, 9, 12, 13, 14, and 19 of SRIF28, and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal of SRIF28. Preferably, it can include the amino acids corresponding to positions 1, 5, 9, and 12 of SRIF28.

Moreover, particularly, examples of substitution of two or more amino acids of the glycosylated polypeptide of the present invention with glycosylated amino acids can include, with respect to the glycosylated polypeptide having affinity towards multiple somatostatin receptors, e.g., substitutions of the amino acids added at the 10th and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 6th and 10th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 2nd and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 14th and 15th positions from the

N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 14th, 15th, and 16th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; and the amino acids added at the 6th, 10th, and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14. Preferably, it can include, e.g., substitutions of the amino acids added at the 10th and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 6th and 10th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 14th and 15th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; the amino acids added at the 14th, 15th, and 16th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14; and the amino acids added at the 6th, 10th, and 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14.

Moreover, similarly, examples of substitution of two or more amino acids with glycosylated amino acids can include, with respect to the glycosylated polypeptide having affinity towards multiple somatostatin receptors, e.g., substitutions of the amino acids corresponding to positions 1 and 5; the amino acids corresponding to positions 5 and 9; the amino acids corresponding to positions 1 and 13; the amino acid corresponding to position 1 and the amino acid added at the 1st position from the N-terminal side among the amino acids further added to the N-terminal side of position 1; the amino acid corresponding to position 1 and the amino acids added at the 1st and 2nd positions from the N-terminal side among the amino acids further added to the N-terminal side of position 1; and the amino acids corresponding to positions 1, 5, and 9; preferably, substitutions of the amino acids corresponding to positions 1 and 5; substitutions of the amino acids corresponding to positions 5 and 9; substitution of the amino acid corresponding to position 1 and the amino acid added at the 1st position among the amino acids further added to the N-terminal side of position 1; the amino acid corresponding to position 1 and the amino acids added at the 1st and 2nd positions from the N-terminal side among the amino acids further added to the N-terminal side of position 1; and substitutions of the amino acids corresponding to positions 1, 5, and 9 of SRIF28.

[0054]

In one aspect of the present invention, the site for substituting an amino acid with a glycosylated amino acid can include, with respect to the stability in blood of the glycosylated polypeptide, e.g., the amino acid corresponding to position 19 of SRIF14, the amino acids added at the 1st, 2nd, 3rd, 6th, 10th, 14th positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14, and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal side of SRIF14. Preferably, it can include the amino acid corresponding to position 19 of SRIF14, the amino acids added at the 1st and 2nd positions from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14, and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal side of SRIF14. More preferably, it is the amino acid corresponding to position 19 of SRIF14, the amino acid added at the 1st position from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14, and the amino acid added at the 1st position from the C-terminal side among the amino acids further added to the C-terminal side of SRIF14.

Moreover, similarly with respect to the stability in blood of the glycosylated polypeptide, it is e.g. one or more amino acids selected from the group consisting of the amino acids corresponding to positions 1, 5, 9, 12, 13, 14, and 19 of SRIF28 and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal side of SRIF28, preferably one or more amino acids selected from the group consisting of the amino acids corresponding to positions 13, 14, and 19 of SRIF28 and the amino acids added at the 1st and 2nd positions from the C-terminal side among the amino acids further added to the C-terminal side of SRIF28, and particularly preferably one or more amino acids selected from the group consisting of the amino acids corresponding to positions 14 and 19 of SRIF28 and the amino acid added at the 1st position from the C-terminal side among the amino acids further added to the C-terminal side of SRIF28. In particular, substitution of an amino acid at a distal site from the N-terminal side of SRIF28 is also preferred.

[0055]

Here, an "amino acid corresponding to a particular amino acid" refers to an amino acid at the same position corresponding to the amino acid sequence of SRIF14 or SRIF28, as long as there is no addition or deletion etc. of

an amino acid in the glycosylated polypeptide. Moreover, if an addition or deletion of an amino acid is present in the amino acid sequence of a glycosylated polypeptide, it refers to the amino acid at the position that takes into account the shift on the amino acid sequence by the addition or deletion of an amino acid. For example, in a glycosylated polypeptide having the sequence Ser₁-Ala₂-Asn₃-Ser₄- at positions 1 to 4, when one amino acid (Trp) is added between the amino acids at positions 2 and 3 (Ser-Ala-Trp-Asn-Ser-), the amino acid corresponding to the amino acid at position 3 (Asn) refers to the amino acid (Asn) in the glycosylated polypeptide which has been shifted one to the C-terminal side by the insertion of Trp. [0056]

In one aspect of the present invention, the amino acid substituted with an glycosylated amino acid is preferably one or more amino acids selected from an amino acid other than the amino acids corresponding to positions 17, 21, 22, 23, and 28 of SRIF28, and in particular, it is more preferably one or more amino acids selected from an amino acid other than the amino acids corresponding to positions 17, 22, 23, and 28 of SRIF28. [0057]

In one aspect of the present invention, when two or more amino acids are substituted with glycosylated amino acids, any combination above can be employed for the site for substituting an amino acid with a glycosylated amino acid but is not limited thereto. For example, a combination of one site being selected from the above preferred sites and other sites being selected from any site of SRIF14 or SRIF28; or a combination of one site being selected from the above preferred sites and other sites being selected from any of one or a few amino acids further added to the N-terminal (position 1 of SRIF28 or position 15 of SRIF14) or C-terminal of somatostatin are also included in one preferred aspect of the present invention. Moreover, in a glycosylated polypeptide, it is more preferred that the two or more amino acids are substituted with glycosylated amino acids at the amino acids specified above. Moreover, it is further preferred that all the glycosylated amino acids included in the glycosylated polypeptide are substituted with glycosylated amino acids at the amino acids specified above. [0058]

Among the glycosylated polypeptides of the present invention, the combination of the two substituted amino acids in a glycosylated polypeptide having two amino acids substituted with glycosylated amino acids can be, e.g., two

amino acids selected from the group consisting of the amino acid corresponding to position 19 of SRIF14, the 1st, 2nd, 3rd, 6th, 10th, and 14th amino acids from the N-terminal side among the amino acids further added to the N-terminal side of SRIF14, and the 1st and 2nd amino acids from the C-terminal side among the amino acids further added to the C-terminal side of SRIF14. Moreover, similarly, it can include two amino acids selected from the group consisting of the amino acids corresponding to positions 1, 5, 9, 12, 13, 14, and 19 of SRIF28, as well as the 1st amino acid further added to the C-terminal of SRIF28 and the 2nd amino acid further added to the C-terminal of SRIF28.

Similarly, substitution positions for a glycosylated polypeptide having three amino acids substituted with glycosylated amino acids or a glycosylated polypeptide having four or more amino acids substituted with glycosylated amino acids can include a combination of three or four or more amino acids selected from the group consisting of the amino acids specified above. However, the glycosylated polypeptide of the present invention is not limited to the combinations listed above, and comprises other combinations as long as it has affinity towards somatostatin receptors and improved stability in blood compared to a naturally-occurring somatostatin.

[0059]

In one aspect of the present invention, the site where a deletion, substitution, or addition of an amino acid other than glycosylated amino acids occurs is preferably e.g. one or more amino acids selected from an amino acid other than the amino acids corresponding to positions 17, 22, 23, and 28 of SRIF28.

[0060]

A "sugar chain" herein refers to a compound made from a string of one or more unit sugars (monosaccharides and/or derivatives thereof). When there is a string of two or more unit sugars, each unit sugar is bound with each other by a dehydration condensation with a glycoside bond in between. Such sugar chains include, but are not limited to e.g. a wide range such as monosaccharides and polysaccharides contained *in vivo* (glucose, galactose, mannose, fucose, xylose, N-acetylglucosamine (GlcNAc), N-acetylgalactosamine (GalNAc), sialic acid, and conjugates and derivatives thereof), as well as a sugar chain degraded or derived from conjugated biomolecules such as degraded polysaccharides, glycoproteins, proteoglycans, glycosaminoglycans, and glycolipids. The sugar chain may be linear or branched.

[0061]

Moreover, a "sugar chain" herein also includes a sugar chain derivative, and examples of sugar chain derivatives include, but are not limited to, a sugar chain wherein the sugar constituting the sugar chain is e.g. a sugar having a carboxy group (such as aldonic acid in which C-position 1 is oxidized to become a carboxylic acid (such as D-gluconic acid which is oxidized D-glucose) and uronic acid in which the terminal C atom has become a carboxylic acid (D-glucuronic acid which is oxidized D-glucose)), a sugar having an amino group or an amino group derivative (such as acetylated amino group) (such as N-acetyl-D-glucosamine and N-acetyl-D-galactosamine), a sugar having both amino and carboxy groups (such as N-acetylneuraminic acid (sialic acid) and N-acetylmuramic acid), a deoxylated sugar (such as 2-deoxy-D-ribose), a sulfated sugar comprising a sulfate group, and a phosphorylated sugar comprising a phosphate group.

[0062]

A preferred sugar chain herein is a sugar chain that increases the stability in blood of somatostatin and will not dissipate affinity towards somatostatin receptors when added to somatostatin (when substituted with an amino acid of somatostatin in the form of a glycosylated amino acid). In an aspect of the present invention, a preferred sugar chain is a sugar chain that can maintain affinity towards multiple receptors of somatostatin, and further preferably a sugar chain that can maintain affinity towards all the receptors of somatostatin when added to somatostatin (when substituted with an amino acid of somatostatin in the form of a glycosylated amino acid).

[0063]

The sugar chain in the glycosylated polypeptide of the present invention is not particularly limited, and may be a sugar chain that exists as a glycoconjugate *in vivo* (such as a glycopeptide (or a glycoprotein), a proteoglycan, and a glycolipid), or it may be a sugar chain that does not exist as a glycoconjugate *in vivo*.


[0064]

A sugar chain that exists as a glycoconjugate *in vivo* is preferred with respect to the fact that the glycosylated polypeptide of the present invention is administered *in vivo*. Examples of such sugar chains include N- or O-linked sugar chains which are sugar chains bound to a peptide (or a protein) *in vivo* as a glycopeptide (or a glycoprotein). An N-linked sugar chain is preferably employed. N-linked sugar chains can include, e.g., a high-mannose form, a complex form, or a hybrid form, particularly preferably a complex form.

[0065]


Examples of preferred complex-type sugar chains used herein include, e.g., a sugar chain represented by the following general formula:

[Chemical Formula 3]



[wherein R¹ and R² are identical or different and are:

[Chemical Formula 4]



, and Ac is an acetyl group].

[0066]

In the glycosylated polypeptide of the present invention, the sugar chain may be bound to the somatostatin peptide with a method other than O- and N-linking, even if it is a sugar chain that exists as a glycoconjugate *in vivo*. For example, as described above, those in which the sugar chain is bound to Cys or Lys via a linker are also included in the glycosylated polypeptide of the present invention.

[0067]

In one aspect of the present invention, it is preferred that the sugar chain in the glycosylated polypeptide of the present invention is a sugar chain consisting of 4 or more, for example, 5 or more, 7 or more, in particular 9 or more, or 11 or more sugars.


[0068]

In one preferred aspect of the present invention, the sugar chain in the glycosylated polypeptide of the present invention is a sugar chain consisting of 5 - 11, 9 - 11, or 11 sugars.

[0069]


In one preferred aspect of the present invention, the sugar chain in the glycosylated polypeptide of the present invention is a biantennary complex-type sugar chain. A complex-type sugar chain is characterized in that it comprises two or more types of monosaccharides, and has the basic structure shown below and a lactosamine structure shown by Gal β 1-4GlcNAc.

[Chemical Formula 5]




A biantennary complex-type sugar chain refers to those having a monoantennary sugar chain consisting of 0 - 3 sugars bound to each of the two mannoses at the end of the basic structure. Examples of biantennary complex-type sugar chains are preferably e.g. a disialo sugar chain as shown below:

[Chemical Formula 6]




, a monosialo sugar chain:

[Chemical Formula 7]




[Chemical Formula 7A]




, an asialo sugar chain:

[Chemical Formula 8]




, a diGlcNAc sugar chain:

[Chemical Formula 9]



, and a dimannose sugar chain:


[Chemical Formula 10]




, and more preferably a disialo sugar chain.

Moreover, the complex-type sugar chain of the present invention includes not only the above biantennary complex-type sugar chains, but also a triantennary complex-type sugar chain (triple-branched complex-type sugar chain) and a tetraantennary complex-type sugar chain (quadruple-branched complex-type sugar chain). For example, triantennary and tetraantennary complex-type sugar chains can include a trisialo sugar chain represented by the structural formula below:

[Chemical Formula -1]




[Chemical Formula -2]



and a tetrasialo sugar chain represented by the structural formula below:


[Chemical Formula -3]




. Moreover, triantennary and tetraantennary complex-type sugar chains can also include a sugar chain having one or more sugars deleted from the non-reducing terminal of these trisialo or tetrasialo sugar chains.

Further, the complex-type sugar chain of the present invention includes those with a fucose added. Complex-type sugar chains with a fucose added can include fucose-containing complex-type sugar chains represented by the structural formula below:


[Chemical Formula -4]




[Chemical Formula -5]




[Chemical Formula -6]



[Chemical Formula -7]



[Chemical Formula -8]




. Moreover, a sugar chain having one or more sugars deleted from the non-reducing terminal of these fucose-containing complex-type sugar chains can also be included.

Moreover, a "disialo sugar chain," a "biantennary complex-type sugar chain," a "monosialo sugar chain," an "asialo sugar chain," a "diGlcNAc sugar chain," a "dimannose sugar chain," a "triantennary complex-type sugar chain," a "tetraantennary complex-type sugar chain," and a "fucose-containing complex-type sugar chain" herein include not only those shown in the above chemical formulae, but also those with a binding mode different from examples shown in the chemical formulae, and such sugar chains are also preferably employed as a sugar chain of the present invention. Examples of such sugar chains include, e.g., a disialo or monosialo sugar chain in which sialic acid and galactose are bound with an ($\alpha 2 \rightarrow 3$) bond.


Moreover, when the sugar chain is one having a sialic acid at the non-reducing terminal of the sugar chain, a sugar chain having the carboxy group of the sialic acid modified can also be employed. The modification of the carboxy group of the sialic acid is preferably a group capable of dissipating the negative charge of the carboxy group or converting it into positive charge, examples of which can include, e.g., an alkylamino group, a benzyl group, an amino group, and an aminoethylamino group, having 1 - 30 carbons. The introduction of these modifying groups will dissipate the negative charge of the carboxyl group of the sialic acid (such as benzyl or amino group) or convert it into positive charge (such as aminoethylamino group), and it is thus possible to contemplate blood clearance or control of body distribution of glycosylated polypeptide.

[0070]

Moreover, the high-mannose sugar chain employed herein is a sugar chain having 2 or more mannoses further bound to the basic structure of the complex-type sugar chain described above. Because high-mannose sugar chains are bulky, stability in blood may become higher by binding a high-mannose sugar chain to the peptide. A sugar chain comprising 5 - 9 mannoses such as a mammalian high-mannose sugar chain is preferred, but it may be a sugar chain comprising more mannoses such as a yeast high-mannose sugar chain. Examples of high-mannose sugar chains preferably employed herein can include, e.g., high-mannose-5 (M-5):
[Chemical Formula 11]



and high-mannose-9 (M-9):
[Chemical Formula 12]












[0071]

Preferred sugar chains herein can include, e.g., the sugar chains described below which are a sugar chain having an identical structure (a sugar chain in which the type of constituent sugar and the binding mode thereof are identical) with a sugar chain that exists in a human body as a glycoprotein bound to a protein (such as a sugar chain described in "FEBS LETTERS Vol. 50, No. 3, Feb. 1975"), or a sugar chain having one or more sugars deleted from the non-reducing terminal of the same.


[0072]

[Chemical Formula 13]




[0073]

[Chemical Formula 14]





[0074]

[Chemical Formula 15]



[0075]

[Chemical Formula 16]



[0076]

In one preferred aspect of the present invention, the sugar chain structure of the glycosylated amino acids in the glycosylated polypeptide of the present invention can be substantially identical, or they may have different sugar chain structures. When the sugar chain structure in the glycosylated polypeptide is substantially identical, they are preferably e.g. 90% or more identical or 99% or more identical, and it is most preferable that the sugar chain structure is completely identical. As used herein, the sugar chain structure in the glycopeptide is identical refers to the fact that in a glycosylated polypeptide having two or more sugar chains added, the type of sugar constituting the sugar chain, the binding order, and the binding mode are identical in the glycopeptide when said sugar chains are compared with each other. Moreover, in one preferred aspect of the present invention, it is preferred that the sugar chains of the glycosylated polypeptide of the present invention are uniform. As used herein, the sugar chains in the glycosylated polypeptide are uniform refers to the fact that the glycosylation sites in the peptide, the type of each sugar constituting the sugar chain, the binding order, and the binding mode between sugars are identical between glycopeptides when sugar chains are compared between glycosylated polypeptides, and that at least 90% or more, preferably 95% or more, and more preferably 99% or more of the sugar chain structure is uniform. In particular, a composition etc. comprising a glycopeptide in which the sugar chains are uniform between glycopeptides has a constant quality, and is preferred particularly in fields such as pharmaceuticals manufacture or assays. The proportion of the uniform sugar chain can be measured for example by a method employing e.g. HPLC, capillary electrophoresis, NMR, and mass spectrometry. [0077]

The preferred glycosylated polypeptides herein are the glycosylated polypeptides (SEQ ID NOs. 21 - 26, 29, 30, 33, 34, 88, 89, 125, 127, 129, 148, 149, 157, 164, and 169) manufactured in Examples 20 - 25, 28, 29, 32 - 35, 49 - 51, 63 - 64, and 66 - 68 described below. In other words, in the following SRIF28 amino acid sequence:

Ser₁-Ala₂-Asn₃-Ser₄-Asn₅-Pro₆-Ala₇-Met₈-Ala₉-Pro₁₀-Arg₁₁-Glu₁₂-Arg₁₃-Lys₁₄-Ala₁₅-Gly₁₆-Cys₁₇-Lys₁₈-Asn₁₉-Phe₂₀-Phe₂₁-Trp₂₂-Lys₂₃-Thr₂₄-Phe₂₅-Thr₂₆-Ser₂₇-Cys₂₈ (SEQ ID NO. 2),

(a1) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having one disialo sugar chain attached Cys further added at the N-terminal side (Example 20) (SEQ ID NO. 21);

(a2) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1 and Asn at position 5 (Example 21) (SEQ ID NO. 22);

(a3) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1 and Arg at position 13 (Example 22) (SEQ ID NO. 23);

(a4) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Asn at position 5 and Ala at position 9 (Example 23) (SEQ ID NO. 24);

(a5) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having two disialo sugar chain attached Cys further added at the N-terminal side (Example 24) (SEQ ID NO. 25);

(a6) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, Asn at position 5, and Ala at position 9 (Example 25) (SEQ ID NO. 26);

(a7) a glycosylated polypeptide having asialo sugar chain attached Cys substituted for Ser at position 1, and having one asialo sugar chain attached Cys further added at the N-terminal side (Example 28) (SEQ ID NO. 29);

(a8) a glycosylated polypeptide having asialo sugar chain attached Cys substituted for Ser at position 1, and having two asialo sugar chain attached Cys further added at the N-terminal side (Example 29) (SEQ ID NO. 30);

(a9) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having GlcNAc-added Cys substituted for Asn at position 19 (Example 32) (SEQ ID NO. 33);

(a10) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having dimannose glycosylated Cys substituted for Asn at position 19 (Example 33) (SEQ ID NO. 34);

(a11) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having four disialo sugar chain attached Cys further added at the N-terminal side (Example 34) (SEQ ID NO. 88);

(a12) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having nine disialo sugar chain attached Cys further added at the N-terminal side (Example 35) (SEQ ID NO. 89);

(a13) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1 and Glu at position 12 (Example 49) (SEQ ID NO. 125);

(a14) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1 in which the carboxy group of the sialic acid on the sugar chain is converted into carboxylic amide, and having one

disialo sugar chain attached Cys further added at the N-terminal side in which the carboxy group of the sialic acid on the sugar chain is converted into carboxylic amide (Example 63) (SEQ ID NO. 148);

(a15) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1 in which the carboxy group of the sialic acid on the sugar chain is protected by a benzyl group, and having one disialo sugar chain attached Cys further added at the N-terminal side in which the carboxy group of the sialic acid on the sugar chain is protected by a benzyl group (Example 64) (SEQ ID NO. 149);

(a16) a glycosylated polypeptide having disialo sugar chain attached Asn substituted for Ser at position 1, and having dimannose glycosylated Cys substituted for Asn at position 19 (Example 66) (SEQ ID NO. 157);

(a17) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having one disialo sugar chain attached Cys further added at the N-terminal side in which the carboxy group of the sialic acid on the sugar chain has formed an amide bond with one end of 1,2-ethylenediamine (Example 67) (SEQ ID NO. 164); and

(a18) a glycosylated polypeptide having disialo sugar chain attached Cys substituted for Ser at position 1, and having three disialo sugar chain attached Cys further added at the N-terminal side (Example 68) (SEQ ID NO. 169) are preferred.

Moreover, in the SRIF14 amino acid sequence:

Ala₁₅-Gly₁₆-Cys₁₇-Lys₁₈-Asn₁₉-Phe₂₀-Phe₂₁-Trp₂₂-Lys₂₃-Thr₂₄-Phe₂₅-Thr₂₆-Ser₂₇-Cys₂₈ (SEQ ID NO. 1),

(a19) a glycosylated polypeptide having disialo sugar chain attached Cys-disialo sugar chain attached Cys-Arg-Lys- further added at the N-terminal side of Ala at position 15 (Example 50) (SEQ ID NO. 127); and

(a20) a glycosylated polypeptide having disialo sugar chain attached Cys-disialo sugar chain attached Cys-disialo sugar chain attached Cys-Arg-Lys- further added at the N-terminal side of Ala at position 15 (Example 51) (SEQ ID NO. 129) are preferred.

[0078]

The glycosylated polypeptide of the present invention can be manufactured by integrating a glycosylation step into a peptide synthesis method well-known to those skilled in the art. A method utilizing an enzyme represented by transglutaminase can also be employed for glycosylation, but since there are problems in this case such as the need

for a large amount of the sugar chain to be added, the complication of purification after the final step, and the restriction for glycosylation position and the sugar chain capable of addition, it cannot be said to be a practical method for large scale manufacturing such as pharmaceuticals manufacture though it is possible to employ for synthesis of a small amount such as for assays.

[0079]

As specific examples of simple manufacturing methods of the glycosylated polypeptide of the present invention which are stable manufacturing methods of a glycosylated polypeptide having uniform sugar chain structure, a method for manufacturing a glycosylated polypeptide by using glycosylated Asn as the glycosylated amino acid and applying a well-known peptide synthesis method such as solid and liquid phase synthesis (method A), and a method for manufacturing a glycosylated polypeptide by manufacturing a peptide in which any amino acid of somatostatin is substituted with Cys according to a well-known peptide synthesis method, and then adding a sugar chain to Cys by chemical synthesis (method B) will be exemplified below. Those skilled in the art is able to manufacture various glycosylated polypeptides by referring to these manufacturing methods, and the glycosylated polypeptide obtained and the manufacturing method thereof are extremely useful especially in the field of pharmaceuticals manufacture.

[0080]

Moreover, these methods A and B can be performed in a combination of two or more. If it is a synthesis of a small amount such as for assays, it is also possible to further combine the above method employing a sugar chain elongation reaction by a transferase. Method A is described in International Publication No. 2004/005330 (US 2005222382 (A1)) and method B is described in International Publication No. 2005/010053 (US 2007060543 (A1)). Moreover, the manufacture of sugar chains having uniform sugar chain structure employed in methods A and B are described in e.g. International Publication No. 03/008431 (US 2004181054 (A1)), International Publication No. 2004/058984 (US 2006228784 (A1)), International Publication No. 2004/058824 (US 2006009421 (A1)), International Publication No. 2004/070046 (US 2006205039 (A1)), and International Publication No. 2007/011055.

[0081]

Method for Manufacturing Glycosylated Polypeptide (Method A)

The glycosylated polypeptide can be manufactured by for example a solid phase synthesis employing glycosylated Asn, the outline of which is shown below.

(1) The carboxy group of an amino acid having the amino group nitrogen protected with a lipophilic protecting group is bound to a resin. In this case, since the amino group nitrogen of the amino acid is protected with a lipophilic protecting group, self-condensation between amino acids is prevented, and the resin reacts with the amino acid to cause binding.

(2) The lipophilic protecting group of the reactant obtained is detached to form a free amino group.

(3) This free amino group and the carboxy group of any amino acid having the amino group nitrogen protected with a lipophilic protecting group are subjected to amidation reaction.

(4) The above lipophilic protecting group is detached to form a free amino group.

(5) By repeating the above steps (3) and (4) for once or more, a peptide in which any number of any amino acids are linked and having a resin bound at one end and a free amino group at the other end is obtained.

(6) Finally, by cleaving the resin with an acid, a peptide having the desired amino acid sequence can be obtained.

In (1), if glycosylated Asn having the amino group nitrogen protected with a lipophilic protecting group is employed instead of the amino acid having the amino group nitrogen protected with a lipophilic protecting group, and the carboxy group of said asparagine portion is reacted with the hydroxyl group of the resin, a peptide having a glycosylated Asn at the C-terminal can be obtained.

Moreover, after (2), or after repeating (3) and (4) for any number of times that is once or more, if glycosylated Asn having the amino group nitrogen protected with a lipophilic protecting group is employed instead of the amino acid having the amino group nitrogen protected with a lipophilic protecting group in (3), a sugar chain can be added at any position.

As such, by employing glycosylated Asn having the amino group nitrogen protected with a lipophilic protecting group instead of the amino acid having the amino group nitrogen protected with a lipophilic protecting group for twice or more times in any of steps (1) and (3), a peptide

having a sugar chain added at any two or more positions can be obtained.

After binding the glycosylated amino acid, if the lipophilic protecting group is detached to form a free amino group, and step (6) is performed immediately thereafter, a peptide having a glycosylated Asn at the N-terminal can be obtained.

[0082]

The resin may be a resin generally used in solid phase synthesis, and for example, 2-chlorotrityl chloride resin (from Merck) functionalized with chlorine, Amino-PEGA resin (from Merck) functionalized with an amino group, NovaSyn™ TGT alcohol resin (from Merck), Wang resin (from Merck), or HMPA-PEGA resin (from Merck) etc. having a hydroxyl group can be employed. Moreover, a linker may exist between the Amino-PEGA resin and the amino acid, and examples of such linkers can include, e.g., 4-hydroxymethylphenoxyacetic acid (HMPA) and 4-(4-hydroxymethyl-3-methoxyphenoxy)-butylacetic acid (HMPB). H-Cys(Trt)-Trityl NovaPEG resin (from Merck) etc. in which the C-terminal amino acid is bound to the resin in advance can also be employed.

Moreover, when the C-terminal is to be amidated, it is preferred to employ e.g. Rink-Amide-PEGA resin (from Merck) functionalized with an amino group. By cleaving this resin and the peptide with an acid, the C-terminal amino acid of the peptide can be amidated.

In the binding between the resin and the amino acid having the amino group nitrogen protected with a lipophilic protecting group, for example, in order to use a resin having a hydroxyl group or a resin functionalized with chlorine, the carboxy group of the amino acid is bound to the resin via an ester bond. Moreover, if a resin functionalized with an amino group is to be used, the carboxy group of the amino acid is bound to the resin via an amide bond.

The 2-chlorotrityl chloride resin is preferred in that it can prevent the racemization of the terminal Cys when elongating the peptide chain in solid phase synthesis.

[0083]

Any amino acid can be employed as the amino acid, and examples can include the natural amino acids serine (Ser), asparagine (Asn), valine (Val), leucine (Leu), isoleucine (Ile), alanine (Ala), tyrosine (Tyr), glycine (Gly), lysine (Lys), arginine (Arg), histidine (His), aspartic acid (Asp), glutamic acid (Glu), glutamine (Gln), threonine (Thr), cysteine (Cys), methionine (Met), phenylalanine (Phe), tryptophan (Trp), and proline (Pro).

Moreover, the D-form of the above natural amino acid can also be used.

[0084]

Examples of lipophilic protecting groups can include, e.g., carbonate- or amide-based protecting groups such as a 9-fluorenylmethoxycarbonyl (Fmoc) group, a t-butyloxycarbonyl (Boc) group, a benzyl group, an allyl group, an allyloxycarbonyl group, and an acetyl group. When introducing a lipophilic protecting group into an amino acid, e.g. when introducing an Fmoc group, introduction can be carried out by adding 9-fluorenylmethyl-N-succinimidyl carbonate and sodium hydrogen carbonate and allowing reaction. The reaction may be performed at 0 - 50°C, preferably at room temperature for approximately about 1 - 5 hours.

[0085]

As the amino acid protected with a lipophilic protecting group, those commercially available can also be used. Examples can include Fmoc-Ser-OH, Fmoc-Asn-OH, Fmoc-Val-OH, Fmoc-Leu-OH, Fmoc-Ile-OH, Fmoc-Ala-OH, Fmoc-Tyr-OH, Fmoc-Gly-OH, Fmoc-Lys-OH, Fmoc-Arg-OH, Fmoc-His-OH, Fmoc-Asp-OH, Fmoc-Glu-OH, Fmoc-Gln-OH, Fmoc-Thr-OH, Fmoc-Cys-OH, Fmoc-Met-OH, Fmoc-Phe-OH, Fmoc-Trp-OH, and Fmoc-Pro-OH.

Moreover, an amino acid protected with a lipophilic protecting group wherein the protecting group is introduced into the side chain can include, e.g., Fmoc-Arg(Pbf)-OH, Fmoc-Asn(Trt)-OH, Fmoc-Asp(OtBu)-OH, Fmoc-Cys(Acm)-OH, Fmoc-Cys(StBu)-OH, Fmoc-Cys(tBu)-OH, Fmoc-Cys(Trt)-OH, Fmoc-Glu(OtBu)-OH, Fmoc-Gln(Trt)-OH, Fmoc-His(Trt)-OH, Fmoc-Lys(Boc)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Thr(tBu)-OH, Fmoc-Trp(Boc)-OH, and Fmoc-Tyr(tBu)-OH.

[0086]

Moreover, if it is desired to add a linker in the amino acid sequence of the glycosylated polypeptide, the linker can be inserted at a preferred position by using a linker protected with a lipophilic protecting group instead of the above amino acid protected with a lipophilic protecting group in the process of solid phase synthesis.

[0087]

When employing a 2-chlorotrityl chloride resin, esterification can be performed with a base such as diisopropylethylamine (DIPEA), triethylamine, pyridine, and 2,4,6-collidine. Moreover, when employing a resin having a hydroxyl group, e.g. a well-known dehydration condensation agent such as 1-mesitylenesulfonyl-3-nitro-1,2,4-triazole (MSNT), dicyclohexylcarbodiimide (DCC), and diisopropylcarbodiimide (DIC) can be employed as the esterification catalyst. The proportion of the amino acid

and the dehydration condensation agent used is 1 part by weight of the former to generally 1 - 10 parts by weight, preferably 2 - 5 parts by weight of the latter.

[0088]

The esterification reaction is preferably performed by for example placing the resin in a solid phase column, washing this resin with a solvent, and then adding the amino acid solution. Examples of washing solvents can include, e.g., dimethylformamide (DMF), 2-propanol, and dichloromethane. Examples of solvents for dissolving the amino acid can include, e.g., dimethyl sulfoxide (DMSO), DMF, and dichloromethane. The esterification reaction may be performed at 0 - 50°C, preferably at room temperature for approximately about 10 minutes - 30 hours, preferably for approximately 15 minutes - 24 hours.

[0089]

It is also preferred to acetylate and cap the unreacted hydroxyl groups on the solid phase at this time with e.g. acetic anhydride.

[0090]

The detachment of the lipophilic protecting group can be performed by for example treatment with a base. Examples of bases can include, e.g., piperidine and morpholine. In such a case, it is preferred that this is performed in the presence of a solvent. Examples of solvents can include, e.g., DMSO, DMF, and methanol.

[0091]

The amidation reaction of the free amino group with the carboxy group of any amino acid having the amino group nitrogen protected with a lipophilic protecting group is preferably performed in the presence of an activator and a solvent.

[0092]

Examples of activators can include, e.g., dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC/HCl), diphenylphosphorylazide (DPPA), carbonyldiimidazole (CDI), diethylcyanophosphonate (DEPC), benzotriazol-1-yloxy-trispyrrolidinophosphonium (DIPCI), benzotriazol-1-yloxy-trispyrrolidinophosphonium hexafluorophosphate (PyBOP), 1-hydroxybenzotriazole (HOBt), hydroxysuccinimide (HOSu), dimethylaminopyridine (DMAP), 1-hydroxy-7-azabenzotriazole (HOAt), hydroxyphthalimide (HOPht), pentafluorophenol (Pfp-OH), 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), 1-[bis(dimethylamino)methylene]-5-chloro-1H-benzotriazolium 3-oxide hexafluorophosphate (HCTU), O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU),

O-benzotriazol-1-yl-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU), and 3,4-dihydro-3-hydrodi-4-oxa-1,2,3-benzotriazine (Dhbt).

[0093]

The amount of the activator used is preferably 1 - 20 equivalents, preferably 1 - 10 equivalents, and further preferably 1 - 5 equivalents to any amino acid having the amino group nitrogen protected with a lipophilic protecting group.

[0094]

Examples of solvents can include, e.g., DMSO, DMF, and dichloromethane. The reaction may be performed at 0 - 50°C, preferably at room temperature for approximately about 10 - 30 hours, preferably for approximately 15 minutes - 24 hours. The detachment of the lipophilic protecting group can be performed similarly to the above.

[0095]

Treatment with an acid is preferred in order to cleave the peptide chain from the resin. Examples of acids can include, e.g., trifluoroacetic acid (TFA) and hydrogen fluoride (HF).

[0096]

In this way, a glycosylated polypeptide having the desired position substituted with glycosylated Asn can be obtained. Moreover, the glycosylated polypeptide purified as such allows formation of a disulfide bond between deprotected Cys as described below.

[0097]

In one embodiment of the present invention, when sialic acid is contained in the non-reducing terminal on the sugar chain of glycosylated Asn employed in solid phase synthesis, the sialic acid is prevented from being cleaved by the acid treatment. It is therefore preferred that the carboxy group of said sialic acid is protected by a protecting group. Examples of protecting groups can include, e.g., a benzyl group, an allyl group, and a diphenylmethyl group. The method for introduction and detachment of the protecting group can be performed by a well-known method. Moreover, the detachment of the protecting group is preferably performed after the glycosylated polypeptide manufactured by solid phase synthesis is cleaved from the resin. After cleaving from the resin, if the glycosylated polypeptide is to be cyclized by allowing formation of a disulfide bond in the glycosylated polypeptide, the detachment of the protecting group may be before or after the step of forming a disulfide bond.

[0098]

Method for Manufacturing Glycosylated Polypeptide (Method B)

The glycosylated polypeptide can also be manufactured by a method of first synthesizing a peptide chain, and then adding a sugar chain to the synthesized peptide chain. Specifically, a peptide comprising Cys at the position to be glycosylated is manufactured by e.g. a solid phase synthesis method, a liquid phase synthesis method, a method of synthesis by a cell, and a method of separating and extracting those that naturally exist. Here, Cys that will not be glycosylated such as Cys at a position planned to form a disulfide bond are protected with for example an acetamidomethyl (Acm) group. Moreover, if Cys that will neither be glycosylated nor be used in the formation of a disulfide bond is to be introduced into the glycosylated polypeptide, Cys can be introduced by protecting the Cys with a protecting group during the glycosylation and disulfide bond formation steps, and then deprotecting the same. Such protecting groups can include, e.g., tert-butyl (tBu) or 4-methoxybenzyl.

Moreover, when a different sugar chain is to be added to Cys in the glycosylated polypeptide, the different sugar chain can be introduced by first unprotecting the Cys where the sugar chain is to be introduced, and then protecting the Cys where a different sugar chain is to be introduced with e.g. StBu. Specifically, when synthesizing a peptide by e.g. solid phase synthesis, Cys where the first sugar chain is to be introduced is unprotected, and Cys where the second sugar chain is to be introduced is made into Cys having a protecting group with e.g. Fmoc-Cys(StBu)-OH. Then, the sugar chain is introduced to the unprotected Cys while retaining the protecting group such as StBu. Next, the StBu group etc. is deprotected to introduce a different sugar chain to the Cys that became unprotected. Note that Cys where the first sugar chain is to be introduced and Cys where the second sugar chain is to be introduced can be one or more Cys.

In the deprotection of the StBu group, the deprotection can be performed by reaction with a reductant such as tris(2-carboxyethyl)phosphine hydrochloride (TCEP), dithiothreitol (DTT), and tributylphosphine. The above reaction may be performed generally at 0 - 80°C, preferably at 5 - 60°C, and further preferably at 10 - 35°C. The reaction time is preferably approximately 30 minutes - 5 hours in general. After completion of the reaction, it may be purified with a well-known method (such as high

performance liquid column chromatography (HPLC)) as appropriate.

When introducing different sugar chains, it is preferred that the introduction starts from sugar chains more stable against the reduction condition in the deprotection step of Cys or the acidic condition in the purification step such as HPLC. In particular, when a sialic acid-containing sugar chain is to be introduced, it is preferred that a sugar chain that does not have a sialic acid or a sugar chain having fewer sialic acid residues is introduced first.

Moreover, if it is desired to add a linker in the amino acid sequence of the glycosylated polypeptide, for example, the linker can be inserted at a preferred position of the synthesized polypeptide by using a linker protected with a lipophilic protecting group instead of an amino acid protected with a lipophilic protecting group in the process of solid phase synthesis.

Next, by reacting a haloacetylated complex-type sugar chain derivative with the peptide comprising an unprotected Cys obtained above, the sugar chain is reacted with the thiol group of the unprotected Cys and bound to the peptide. The above reaction may be performed in a phosphate buffer, a Tris-hydrochloride buffer, a citrate buffer, or a mixed solution thereof, generally at 0 - 80°C, preferably at 10 - 60°C, and further preferably at 15 - 35°C. The reaction time is generally 10 minutes - 24 hours, and preferably approximately 30 minutes - 5 hours in general. After completion of the reaction, it may be purified with a well-known method (such as HPLC) as appropriate.

[0099]

A haloacetylated complex-type sugar chain derivative is for example a compound wherein the hydroxyl group bound to the carbon at position 1 of a complex-type asparagine-linked sugar chain is substituted with $-\text{NH}-(\text{CH}_2)_a-(\text{CO})-\text{CH}_2\text{X}$ (wherein X is a halogen atom, and a is an integer and is not limited as long as it does not inhibit the linker function of interest, but is preferably an integer from 0 - 4).

[0100]

Specifically, a haloacetylated complex-type sugar chain derivative and a Cys-containing peptide are reacted in a phosphate buffer at room temperature. After completion of the reaction, a glycosylated polypeptide substituted with a glycosylated Cys can be obtained by purification with HPLC.

Moreover, the reaction can also be performed in a mixed solution of an organic solvent such as DMSO, DMF,

methanol, and acetonitrile with the above buffer. Here, the organic solvent can be added to the above buffer at a proportion in the range of 0 - 99% (v/v). For a peptide comprising an unprotected Cys with low solubility to the buffer, addition of such an organic solvent can improve the solubility to the reaction solution and is thus preferred.

Alternatively, the reaction can also be performed in an organic solvent such as DMSO, DMF, methanol, and acetonitrile or a mixed solution thereof. In this case, it is preferably performed in the presence of a base. Examples of bases can include, e.g., DIPEA, triethylamine, pyridine, and 2,4,6-collidine. Moreover, the reaction can also be performed in a mixed solution of guanidine hydrochloride or urea added to the buffer solution. Guanidine hydrochloride or urea can be added to the above buffer so that the final concentration will be 1 M - 8 M. The solubility of a peptide with low solubility in the buffer can also be improved by addition of guanidine hydrochloride or urea and is thus preferred.

Further, in order to prevent the peptide comprising an unprotected Cys from forming a dimer via a disulfide bond, tris(2-carboxyethyl)phosphine hydrochloride (TCEP) or dithiothreitol (DTT) can also be added to the buffer for reaction. TCEP or DTT can be added to the buffer so that the final concentration will be 10 μ M - 10 mM.

[0101]

Moreover, after binding the sugar chain to the Cys of interest, the protecting group of Cys protected with AcM etc. is deprotected. When the protecting group is an AcM group, it can be deprotected by subjecting to reaction with e.g. iodine, mercury acetate (II), silver nitrate (I), or silver(I) acetate in water, methanol, acetic acid, or a mixed solution thereof.

The above reaction may be performed generally at 0 - 80°C, preferably at 5 - 60°C, and further preferably at 10 - 35°C. The reaction time is preferably approximately 5 minutes - 24 hours in general. After completion of the reaction, it may be treated with DTT or hydrochloric acid etc., and then purified with a well-known method (such as HPLC) as appropriate.

In this way, a glycosylated polypeptide having the desired position substituted with glycosylated Cys can be obtained. Moreover, the glycosylated polypeptide purified as such allows formation of a disulfide bond between deprotected Cys as described below.

[0102]

Moreover, when manufacturing a glycosylated polypeptide having multiple sialic acid-containing sugar

chains such as disialo or monosialo sugar chain in the peptide sequence, a sialic acid-containing sugar chain in which the carboxy group of the sialic acid on the sugar chain to be introduced is protected by e.g. a benzyl (Bn) group, an allyl group, a diphenylmethyl group, and a phenacyl group can be employed.

When a sugar chain having the carboxy group of the sialic acid protected is introduced, a step of deprotecting the sialic acid protecting group can be carried out after a step of forming a disulfide bond in the glycosylated polypeptide described below.

Accordingly, by protecting the carboxy group of the sialic acid with a benzyl group etc., a separation and purification step by e.g. HPLC in the manufacturing step will be facilitated. Moreover, the protection of the carboxy group of the sialic acid will also enable the prevention of detachment of an acid-labile sialic acid.

The protection reaction of the carboxy group of the sialic acid on the sugar chain can be performed by a method well-known to those skilled in the art. Moreover, in a glycosylated polypeptide in which a disulfide bond was formed, the protecting group of the carboxy group of the sialic acid can be deprotected by hydrolysis under basic conditions. The above reaction may be performed generally at 0 - 50°C, preferably at 0 - 40°C, and further preferably at 0 - 30°C. The reaction time is preferably approximately 5 minutes - 5 hours in general. After completion of the reaction, it may be neutralized with a weak acid such as phosphoric or acetic acid, and then purified with a well-known method (such as HPLC) as appropriate.

[0103]

Moreover, in the glycosylated polypeptide prepared by the above methods A and B, a disulfide bond between Cys can be formed with a method well-known to those skilled in the art employing e.g. air and/or oxygen, iodine, DMSO, a mixture of oxidized and reduced glutathione, potassium ferricyanide, Ellman's reagent (5,5'-dithiobis(2-nitrobenzoic acid)), thallium(III) trifluoroacetate, and alkyltrichlorosilane sulfoxide.

When forming a disulfide bond between Cys-Cys, Cys in the glycosylated polypeptide which desirably do not form a disulfide bond is to be protected with a protecting group. As such protecting groups, a protecting group which is stable under oxidizing conditions such as AcM, tBu, 4-methoxybenzyl, and 4-methylbenzyl can be employed.

Moreover, in method B, it is also possible to perform the formation of disulfide bond before the introduction of the sugar chain. However, if a protecting group is

introduced to Cys that is to be subject to disulfide bonding, the deprotection step will come before the disulfide bond formation step.

[0104]

(Activity)

The glycosylated polypeptide of the present invention has affinity towards somatostatin receptors. Having "affinity towards somatostatin receptors" herein means having affinity towards at least one of somatostatin receptors SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5.

[0105]

The glycosylated polypeptide of the present invention preferably has affinity towards two or more somatostatin receptors, more preferably has affinity towards three or more receptors, further preferably has affinity towards four or more receptors, and most preferably has affinity towards all five receptors SSTR1 - SSTR5 similarly to a natural somatostatin (SRIF28 and SRIF14). In particular, it is preferred that it has affinity towards any one of at least SSTR1 and SSTR4 and affinity towards other SSTRs.

As such, the glycosylated polypeptide of the present invention has somatostatin activity (agonist activity) and antagonist activity towards a somatostatin receptor by having affinity towards a somatostatin receptor.

[0106]

For example, affinity towards each somatostatin receptor can be measured with e.g. competitive binding experiment *in vitro*.

In the measurement of affinity by competitive binding experiment, the affinity of the test substance (such as 50% inhibitory concentration: IC₅₀ value or binding inhibition constant: K_i value) can be measured by competitively binding a labeled ligand and the test substance to the receptor and seeking the release of the labeled ligand when the test substance is administered.

[0107]

For example, somatostatin activity (agonist activity) can be evaluated by a cAMP production suppression test *in vitro* employing somatostatin receptor expression cells as shown in Examples 69-3 and 69-4.

The cAMP production suppression test can be evaluated by treating somatostatin receptor expression cells with the glycosylated polypeptide or the control compounds SRIF14 or SRIF28, measuring the amount of cAMP accumulated in the cell after culturing for a certain amount of time, and comparing with the control compound. The amount of cAMP

can be measured by a well-known method such as the enzyme immunoassay method (EIA).

[0108]

For example, somatostatin activity (agonist activity) can be evaluated by a GH production suppression test *in vivo* as shown in Examples 89 and 90.

The GH production suppression test can be carried out for example as follows. The glycosylated polypeptide is subcutaneously administered to non-fasted rats. Rats are administered a GH release enhancer under general anesthesia, and then blood is collected in approximately 5 minutes. The collected blood is the plasma sample, and GH concentration is measured by a well-known method such as the enzyme immunoassay method (EIA). Moreover, as a control, plasma samples are obtained from rats which were administered with saline that does not contain glycosylated polypeptide, and somatostatin activity can be evaluated by comparing the GH concentrations measured.

[0109]

In the glycosylated polypeptide of the present invention, even if some extent of attenuation in affinity towards each receptor is triggered by glycosylating its structure, the half-life in blood is extended and as a result, somatostatin activity equivalent to a naturally-occurring somatostatin that is not glycosylated (hereinafter may be referred to as "non-glycosylated somatostatin") can be maintained, and it can preferably have increased somatostatin activity. In light of this extension of half-life in blood, the ratio between the K_i value of the glycosylated polypeptide of the present invention against each receptor and the K_i value of unglycosylated SRIF14 is preferably in the range of 1000:1 - 0.3:1, more preferably in the range of 100:1 - 0.3:1, and further preferably in the range of 10:1 - 0.3:1, as measured in e.g. the method described in Example 69-1.

[0110]

The stability in blood of the glycosylated polypeptide of the present invention is preferably equivalent to a naturally-occurring somatostatin (non-glycosylated somatostatin) or more. Stability in blood can be measured with a method well-known to those skilled in the art, and can be decided with e.g. stability in plasma, half-life in blood, and AUC (drug plasma concentration - area under the time curve) as indicators. Moreover, decrease in renal or hepatic clearance also contributes to an increase in stability in blood.

[0111]

The glycosylated polypeptide of the present invention has increased half-life in blood compared to a non-glycosylated SRIF28, and its half-life is increased by 4-folds or more, preferably 10-fold or more, more preferably 20-folds or more, and further preferably 30-folds or more compared to SRIF28, as measured in e.g. the experimental method shown in Example 70.

[0112]

Moreover, the glycosylated polypeptide of the present invention has preferably 4-folds or more, more preferably 10-fold or more, and further preferably 20-folds or more stability in blood compared to SRIF28, as measured in e.g. the experimental method shown in Example 88.

[0113]

(Pharmaceutical Composition)

Next, a pharmaceutical composition containing the glycosylated polypeptide of the present invention as the active ingredient will be described.

The pharmaceutical composition containing the glycosylated polypeptide of the present invention as the active ingredient is effective for the treatment or prevention of somatostatin-related diseases. As described above, various actions are known for somatostatin, and diseases related to these actions also vary. For example, examples of somatostatin-related diseases include, e.g., acromegaly, gigantism, Alzheimer's disease and other forms of dementia, cancer, hormone-producing tumor, endocrine tumor (such as carcinoid, VIPoma, insulinoma, and glucagonoma), Cushing's disease, hormone secretion defect, diabetes and complications thereof, pains, arthritis, diarrhea, gastric ulcer, inflammatory bowel disease, irritable bowel syndrome, gastrointestinal obstruction, ileus, postoperative restenosis, radiation damage, and eye disease (such as dry eye, glaucoma, interstitial keratitis, iritis, cataract, and conjunctivitis). The pharmaceutical composition containing the glycosylated polypeptide of the present invention as the active ingredient is effective for the treatment or prevention of the above diseases, in particular, acromegaly, dementia, cancer, hormone-producing tumor, endocrine tumor, Cushing's disease, hormone secretion defect, diabetes complication, diarrhea, gastrointestinal obstruction, ileus, radiation damage, eye disease, various tumors or gut-associated disease, and gastrointestinal symptoms accompanying excess hormone production.

[0114]

The above pharmaceutical composition is one formulated into the form of an ordinary pharmaceutical composition with a diluent or an excipient such as a generally used filler, expander, binder, moisturizer, disintegrant, surfactant, and lubricant.

Examples of such pharmaceutical compositions include, e.g., tablets, pills, powders, liquids, suspensions, emulsions, granules, capsules, suppositories, inhalants, ophthalmic solutions, and injections.

[0115]

The amount of the glycosylated polypeptide of the present invention contained in the pharmaceutical composition is not particularly limited and can be selected as appropriate from a broad range, but in general 1 - 70% by weight of the glycosylated polypeptide of the present invention is preferably contained in the pharmaceutical composition.

[0116]

The pharmaceutical composition containing the glycosylated polypeptide of the present invention as the active ingredient can either further contain other active ingredients, or it can be employed in combination with a pharmaceutical composition containing other active ingredients. Moreover, the pharmaceutical composition containing the glycosylated polypeptide of the present invention as the active ingredient can either comprise the glycosylated polypeptide as a pharmaceutically acceptable salt, or further contain one or more different glycosylated polypeptides of the present invention as active ingredients. Moreover, it can also be employed in combination with the pharmaceutical composition containing one or more different glycosylated polypeptides of the present invention as the active ingredient. Moreover, Examples of other ingredients that can be contained in the pharmaceutical composition can include a pharmaceutically acceptable carrier known to those skilled in the art.

[0117]

Moreover, a treatment using the glycosylated polypeptide of the present invention may include e.g. radiation therapy, and it is also useful in scintigraphy for measuring the distribution of cells and tissues expressing any of SSTR1 - SSTR5 throughout the body. The use of extracorporeal imaging by radiation scanning or magnetic resonance will enable *in vivo* semiquantitative detection.

[0118]

A radiolabeled glycosylated polypeptide is useful for therapeutic treatment of a malignant tumor expressing any

of SSTR1 - SSTR5, for example in a human body in a tissue that does not comprise a substantial amount of SSTR1 - SSTR5 in a healthy state. Moreover, the labeled glycosylated polypeptide can be administered for scintigraphy, or as a composition containing an effective amount to suppress a tumor. Examples of such labels are different isotopes such as iodine (^{123}I , ^{125}I , and ^{131}I), indium (^{111}In), carbon (^{11}C), fluorine (^{18}F), technetium ($^{99\text{m}}\text{Tc}$), and yttrium (^{90}Y), or a fluorescent label. The labeling can be carried out by a method well-known to those skilled in the art, and the label can be contained in the glycosylated polypeptide, bound directly thereto, or bound to an appropriate compound and then bound to the glycosylated polypeptide. For example, in iodization of tyrosine, the labeling can be performed by e.g. a method employing chloramine-T etc.

[0119]

The method for administering the pharmaceutical composition according to the present invention is not particularly restricted, and it is administered with a method according to various drug formulations, age, sex, and the disease condition of the patient, and other conditions. The method of administration for tablets, pills, liquids, suspensions, emulsions, granules, and capsules include e.g. oral administration. Moreover, in case of injections, it can be intravenously, intramuscularly, intradermally, subcutaneously, or intraperitoneally administered alone or mixed with an ordinary fluid replacement such as glucose or amino acid. In case of suppositories, it is intrarectally administered. In case of ophthalmic solutions, it is applied to an eye tissue such as the conjunctival sac. In case of inhalants, it is applied to the bronchial tube or the lung.

[0120]

The administration dose of the above pharmaceutical composition may be selected as appropriate according to usage, age, sex, and the disease extent of the patient, and other conditions, and for example can be an administration dose that will be 0.1 - 900 nmol, preferably 1 - 100 nmol, and more preferably 1 - 10 nmol of the glycosylated polypeptide of the present invention per 1 kg of body weight.

[0121]

The administration frequency of the above pharmaceutical composition may be selected as appropriate according to usage, age, sex, and the disease extent of the patient, and other conditions, and 3 times/day, twice/day, once/day, and further at a less frequent administration

frequency according to stability in blood thereof (such as once/week and once/month) may be selected. Preferably, the administration frequency of the above pharmaceutical composition is once or less per day.

[0122]

The sugar chain added to the glycosylated polypeptide of the present invention is easily degraded by the metabolic system in the body. Moreover, in one aspect of the present invention, said sugar chain has a structure that exists as bound to a glycopeptide (or a glycoprotein) *in vivo*. Accordingly, a pharmaceutical composition comprising the glycosylated polypeptide of the present invention and said peptide as active ingredients has advantages such as not showing side effects or antigenicity when administered *in vivo* and less concern for losing drug effect by allergic reactions or antibody production.

[0123]

Further, the glycosylated polypeptide of the present invention can be stably and easily supplied in large amounts, and it is extremely useful with respect to providing pharmaceuticals having stable and high quality.

Moreover, the present invention also provides a method for treating or preventing a somatostatin-related disease, characterized in administering an effective amount of the glycosylated polypeptide of the present invention.

[0124]

The terms used herein are to be employed to describe particular embodiments, and do not intend to limit the invention.

Moreover, the term "comprising" as used herein, unless the content clearly indicates to be understood otherwise, intends the presence of the described items (such as components, steps, elements, and numbers), and does not exclude the presence of other items (such as components, steps, elements, and numbers).

[0125]

Unless otherwise defined, all terms used herein (including technical and scientific terms) have the same meanings as those broadly recognized by those skilled in the art of the technology to which the present invention belongs. The terms used herein, unless explicitly defined otherwise, are to be construed as having consistent meanings with the meanings herein and related technical fields, and shall not be construed as having idealized or excessively formal meanings.

[0126]

The embodiments of the present invention may be described referring to schematic diagrams. In case of schematic diagrams, they may be exaggerated in presentation in order to allow clear description.

Terms such as first and second are employed to express various elements, and it should be recognized that these elements are not to be limited by these terms. These terms are employed solely for the purpose of discriminating one element from another, and it is for example possible to describe a first element as a second element, and similarly, to describe a second element as a first element without departing from the scope of the present invention.

[0127]

The present invention will now be described in further detail referring to Examples. However, the present invention can be embodied by various aspects, and shall not be construed as being limited to the Examples described herein.

Examples

[0128]

The notation system of the glycosylated polypeptides herein will be described below.

For example, S1C(disialo)·N5C(disialo)-SRIF28 shows that Ser at position 1 (S1) and Asn at position 5 of polypeptide SRIF28 are both substituted by disialo sugar chain attached Cys (C(disialo)).

Moreover, S1C(disialo)-D-Trp22-SRIF28 shows that Ser at position 1 (S1) of polypeptide SRIF28 is substituted by disialo sugar chain attached Cys (C(disialo)), and further Trp at position 22 is substituted with D-Trp.

Moreover, C(disialo)-R-K-SRIF14 shows that glycosylated Cys-Arg-Lys- is added at the N-terminal side of SRIF14.

Moreover, 29C(disialo)-SRIF28 or 30C(disialo)-SRIF28 shows peptides having one disialo sugar chain attached Cys further added to Cys at position 28 which is the C-terminal of SRIF28 (29C(disialo)-SRIF28), or having -W-disialo sugar chain attached Cys (W is any amino acid that binds to Cys at position 28) further added to Cys at position 28 which is the C-terminal of SRIF28 (30C(disialo)-SRIF28).

Moreover, S1-2C(disialo)-SRIF28 shows that Ser at position 1 present at the N-terminal of polypeptide SRIF28 is substituted with two consecutive disialo sugar chain attached Cys.

In the meantime, "disialo" means a disialo sugar chain, "monosialo" means a monosialo sugar chain, "asialo" means an asialo sugar chain, "diGlcNAc" means a diGlcNAc sugar chain, "GlcNAc" means an N-acetylglucosamine, "diMan"

means a dimannose sugar chain, "trisialo" means a trisialo sugar chain, and "tetrasialo" means a tetrasialo sugar chain. Moreover, (disialo(aminoethylamide)) means that the carboxy group of the sialic acid of the disialo sugar chain is modified by an aminoethylamino group. "Bn," "amide," or "hexadecylamide" instead of "aminoethylamide" each mean that the carboxy group of the sialic acid on the sugar chain is protected by a benzyl group, an amino group, hexadecylamino group.


[0129]

Example 1 Synthesis of S1C(disialo)-SRIF28


1-1 Glycosylation reaction of thiol

Peptide 1 (SEQ ID NO. 38) represented by the following formula (1) (from APC, Inc.) (60.6 mg, 18.3 μ mol) and compound a represented by the following formula (a) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) (85.8 mg, 36.6 μ mol, 2.0 equivalents to peptide 1) were dissolved in 33 mM phosphate buffer (pH 7.4, 5.5 mL), and reacted at room temperature for 30 minutes.

[Chemical Formula 17]



[Chemical Formula 18]



The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL™ PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous acetic acid (AcOH), B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 -> 75:25, 15 minutes, linear gradient elution] to obtain glycopeptide 2 (SEQ ID NO. 39) represented by the following formula (2) (60.5 mg, 10.9 μ mol, yield 59%).

[Chemical Formula 19]

ESI-MS: (m/z) calcd for $C_{229}H_{358}N_{50}O_{102}S_4$: $[M+3H]^{3+}$ 1858.3,
 $[M+4H]^{4+}$ 1394.0, $[M+5H]^{5+}$ 1115.4, found 1858.1, 1393.8,
 1115.2.


To glycopeptide 2 obtained in the method described in the above 1-1 (51.2 mg, 9.19 μmol) was added an aqueous solution (3.7 mL) of silver(I) acetate (18.8 mg, 113 μmol), and reacted at room temperature for 40 minutes. DTT (43.6 mg, 282 μmol) dissolved in 200 mM Tris-HCl buffer (pH 7.4, 3.7 mL) and 100 mM ascorbic acid aqueous solution (0.92 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 75:25, 15 minutes, linear gradient elution] to obtain glycopeptide 3 (SEQ ID NO. 40) represented by the following formula (3) (29.2 mg, 5.38 μmol , yield 58%).

Disialo
Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

ESI-MS: (m/z) calcd for $C_{223}H_{348}N_{48}O_{100}S_4$: $[M+3H]^{3+}$ 1810.9, $[M+4H]^{4+}$ 1358.4, $[M+5H]^{5+}$ 1086.9, $[M+6H]^{6+}$ 906.0, found 1810.7, 1358.3, 1086.6, 905.7.

Glycopeptide 3 obtained in the method described in the above 1-2 (29.2 mg, 5.38 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 10.8 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 77:23 -> 64:36, 17 minutes, linear gradient elution] to obtain a fraction containing compound (SlC(disialo)-SRIF28) represented by the following formula (4) (SEQ ID NO. 5).

65



(4)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 75:25, 15 minutes, linear gradient elution] to obtain S1C(disialo)-SRIF28 (17.2 mg, 3.17 μ mol, yield 59%).


ESI-MS: (m/z) calcd for $C_{223}H_{346}N_{48}O_{100}S_4$: $[M+3H]^{3+}$ 1810.2, $[M+4H]^{4+}$ 1357.9, $[M+5H]^{5+}$ 1086.5, found 1810.0, 1357.5, 1086.4.

[0132]

Example 2 Synthesis of N5C(disialo)-SRIF28


A compound represented by the following formula (6) (N5C(disialo)-SRIF28) (SEQ ID NO. 6) was synthesized similarly to Example 1, except that a compound represented by the following formula (5) (peptide 5) (SEQ ID NO. 41) was employed instead of peptide 1.

[Chemical Formula 22]



(5)

[Chemical Formula 23]




(6)

[0133]

Example 3 Synthesis of A9C(disialo)-SRIF28


A compound represented by the following formula (8) (A9C(disialo)-SRIF28) (SEQ ID NO. 7) was synthesized similarly to Example 1, except that a compound represented by the following formula (7) (peptide 7) (SEQ ID NO. 42) was employed instead of peptide 1.

[Chemical Formula 24]



(7)

[Chemical Formula 25]




[0134]


Example 4 Synthesis of E12C(disialo)-SRIF28

A compound represented by the following formula (10) (E12C(disialo)-SRIF28) (SEQ ID NO. 8) was synthesized similarly to Example 1, except that a compound represented by the following formula (9) (peptide 9) (SEQ ID NO. 43) was employed instead of peptide 1.

[Chemical Formula 26]



[Chemical Formula 27]




[0135]


Example 5 Synthesis of R13C(disialo)-SRIF28

A compound represented by the following formula (12) (R13C(disialo)-SRIF28) (SEQ ID NO. 9) was synthesized similarly to Example 1, except that a compound represented by the following formula (11) (peptide 11) (SEQ ID NO. 44) was employed instead of peptide 1.

[Chemical Formula 28]



[Chemical Formula 29]




[0136]

Example 6 Synthesis of K14C(disialo)-SRIF28

A compound represented by the following formula (14) (K14C(disialo)-SRIF28) (SEQ ID NO. 10) was synthesized similarly to Example 1, except that a compound represented


by the following formula (13) (peptide 13) (SEQ ID NO. 45) was employed instead of peptide 1.

[Chemical Formula 30]



(13)

[Chemical Formula 31]




(14)

[0137]

Example 7 Synthesis of A15C(disialo)-SRIF28


A compound represented by the following formula (16) (A15C(disialo)-SRIF28) (SEQ ID NO. 11) was synthesized similarly to Example 1, except that a compound represented by the following formula (15) (peptide 15) (SEQ ID NO. 46) was employed instead of peptide 1.

[Chemical Formula 32]



(15)

[Chemical Formula 33]




(16)

[0138]

Example 8 Synthesis of G16C(disialo)-SRIF28


A compound represented by the following formula (18) (G16C(disialo)-SRIF28) (SEQ ID NO. 12) was synthesized similarly to Example 1, except that a compound represented by the following formula (17) (peptide 17) (SEQ ID NO. 47) was employed instead of peptide 1.

[Chemical Formula 34]



(17)

[Chemical Formula 35]




[0139]


Example 9 Synthesis of K18C(disialo)-SRIF28

A compound represented by the following formula (20) (K18C(disialo)-SRIF28) (SEQ ID NO. 13) was synthesized similarly to Example 1, except that a compound represented by the following formula (19) (peptide 19) (SEQ ID NO. 48) was employed instead of peptide 1.

[Chemical Formula 36]



[Chemical Formula 37]




[0140]


Example 10 Synthesis of N19C(disialo)-SRIF28

A compound represented by the following formula (22) (N19C(disialo)-SRIF28) (SEQ ID NO. 14) was synthesized similarly to Example 1, except that a compound represented by the following formula (21) (peptide 21) (SEQ ID NO. 49) was employed instead of peptide 1.

[Chemical Formula 38]



[Chemical Formula 39]




[0141]


Example 11 Synthesis of F21C(disialo)-SRIF28

A compound represented by the following formula (24) (F21C(disialo)-SRIF28) (SEQ ID NO. 15) was synthesized similarly to Example 1, except that a compound represented by the following formula (23) (peptide 23) (SEQ ID NO. 50) was employed instead of peptide 1.

[Chemical Formula 40]



[Chemical Formula 41]




[0142]


Example 12 Synthesis of T26C(disialo)-SRIF28

A compound represented by the following formula (26) (T26C(disialo)-SRIF28) (SEQ ID NO. 16) was synthesized similarly to Example 1, except that a compound represented by the following formula (25) (peptide 25) (SEQ ID NO. 51) was employed instead of peptide 1.

[Chemical Formula 42]



[Chemical Formula 43]




[0143]


Example 13 Synthesis of 29C(disialo)-SRIF28

A compound represented by the following formula (28) (29C(disialo)-SRIF28) (SEQ ID NO. 17) was synthesized similarly to Example 1, except that a compound represented by the following formula (27) (peptide 27) (SEQ ID NO. 52) was employed instead of peptide 1.

[Chemical Formula 44]



[Chemical Formula 45]




[0144]


Example 14 Synthesis of 30C(disialo)-SRIF28

A compound represented by the following formula (30) (30C(disialo)-SRIF28) (SEQ ID NO. 18) was synthesized similarly to Example 1, except that a compound represented by the following formula (29) (peptide 29) (SEQ ID NO. 53) was employed instead of peptide 1.

[Chemical Formula 46]



[Chemical Formula 47]




[0145]


Example 15 Synthesis of S1C(disialo)-D-Trp22-SRIF28

A compound represented by the following formula (32) (S1C(disialo)-D-Trp22-SRIF28) (SEQ ID NO. 19) was synthesized similarly to Example 1, except that a compound represented by the following formula (31) (peptide 31) (SEQ ID NO. 54) was employed instead of peptide 1.

[Chemical Formula 48]



[Chemical Formula 49]




[0146]


Example 16 Synthesis of A9C(disialo)-D-Trp22-SRIF28

A compound represented by the following formula (34) (A9C(disialo)-D-Trp22-SRIF28) (SEQ ID NO. 20) was synthesized similarly to Example 1, except that a compound represented by the following formula (33) (peptide 33) (SEQ ID NO. 55) was employed instead of peptide 1.

[Chemical Formula 50]



[Chemical Formula 51]




[0147]


Example 17 Synthesis of C(disialo)-SRIF14

A compound represented by the following formula (36) (C(disialo)-SRIF14) (SEQ ID NO. 35) was synthesized similarly to Example 1, except that a compound represented by the following formula (35) (peptide 35) (SEQ ID NO. 56) was employed instead of peptide 1.

[Chemical Formula 52]



[Chemical Formula 53]




[0148]


Example 18 Synthesis of C(disialo)-R-K-SRIF14

A compound represented by the following formula (38) (C(disialo)-R-K-SRIF14) (SEQ ID NO. 36) was synthesized similarly to Example 1, except that a compound represented by the following formula (37) (peptide 37) (SEQ ID NO. 57) was employed instead of peptide 1.

[Chemical Formula 54]



[Chemical Formula 55]




[0149]

Example 19 Synthesis of C(disialo)-C12 linker-SRIF14

19-1 Synthesis of peptide


2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a PreludeTM peptide synthesizer, a protected peptide 39 (SEQ ID NO. 58) represented by the following formula (39) was synthesized in a state bound to the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

[Chemical Formula 56]



The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, employing HCTU as the condensation agent, Fmoc-12-aminododecanoic acid and Fmoc-Cys(Trt)-OH were condensed in sequence. After condensation, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken for 3 hours at room temperature. This leads to the detachment of the protecting group of the amino acid side chain (other than the Acm group), as well as cleaving between the peptide and the resin. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. A part of the crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 60:40 -> 36.2:63.8, 20 minutes, linear gradient elution]

to obtain a compound represented by the following formula (40) (peptide 40) (SEQ ID NO. 59) (11.2 mg).
[Chemical Formula 57]




ESI-MS: (m/z) calcd for $C_{97}H_{144}N_{22}O_{23}S_3$: $[M+2H]^{2+}$ 1042.3, $[M+3H]^{3+}$ 695.2, found 1042.0, 695.0.

[0150]

19-2 Glycosylation reaction of thiol

Peptide 40 obtained in the method described in the above 19-1 (6.8 mg, 3.3 μ mol) and compound a represented by the above formula (a) (19.1 mg, 8.15 μ mol) were dissolved in 0.2 M phosphate buffer (pH 7.4, 0.96 mL) containing 7 M guanidine hydrochloride and 330 μ M TCEP, and reacted at room temperature for 2 hours. After confirming the disappearance of raw materials with HPLC, the reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 50:50, 25 minutes, linear gradient elution] to obtain a compound represented by the following formula (41) (glycopeptide 41) (SEQ ID NO. 60) (7.1 mg, 1.6 μ mol, yield 50%).

[Chemical Formula 58]




ESI-MS: (m/z) calcd for $C_{183}H_{283}N_{29}O_{85}S_3$: $[M+3H]^{3+}$ 1449.5, $[M+4H]^{4+}$ 1087.4, $[M+5H]^{5+}$ 870.1, found 1449.3, 1087.2, 870.0.

[0151]

19-3 Deprotection of Acm group

To glycopeptide 41 obtained in the method described in the above 19-2 (10.3 mg, 2.37 μ mol) was added an aqueous solution (0.95 mL) of silver(I) acetate (9.7 mg, 58 μ mol), and reacted at room temperature for 30 minutes. DTT (22.3 mg, 145 μ mol) dissolved in 100 mM phosphate buffer (pH 7.4, 0.95 mL) and 100 mM ascorbic acid aqueous solution (0.95 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 50:50, 25 minutes, linear gradient elution] to obtain

glycopeptide 42 represented by the following formula 42 (SEQ ID NO. 61) (5.8 mg, 1.4 μ mol, yield 59%).
[Chemical Formula 59]




ESI-MS: (m/z) calcd for $C_{177}H_{273}N_{27}O_{83}S_3$: $[M+3H]^{3+}$ 1402.1, $[M+4H]^{4+}$ 1051.8, $[M+5H]^{5+}$ 841.7, found 1401.9, 1051.7, 841.5.

[0152]

19-4 Formation of disulfide bond

Glycopeptide 42 obtained in the method described in the above 19-3 (5.8 mg, 1.4 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 3.5 mL), and reacted at room temperature for 30 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 70:30 $\rightarrow$ 50:50%, 25 minutes, linear gradient elution] to obtain a fraction containing compound (C(disialo)-C12 linker-SRIF14) represented by the following formula 43 (SEQ ID NO. 37).

[Chemical Formula 60]



This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 50:50, 25 minutes, linear gradient elution] to obtain C(disialo)-C12 linker-SRIF14 (3.6 mg, 0.86 μ mol, yield 61%).
ESI-MS: (m/z) calcd for $C_{177}H_{271}N_{27}O_{83}S_3$: $[M+3H]^{3+}$ 1401.5, $[M+4H]^{4+}$ 1051.3, $[M+5H]^{5+}$ 841.3, found 1401.2, 1051.2, 841.1.

[0153]

Example 20 Synthesis of S1-2C(disialo)-SRIF28

20-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and

76


ESI-MS: (m/z) calcd for $C_{318}H_{502}N_{58}O_{165}S_5$: $[M+5H]^{5+}$ 1588.6, $[M+6H]^{6+}$ 1324.0, $[M+7H]^{7+}$ 1135.0, $[M+8H]^{8+}$ 993.3, $[M+9H]^{9+}$ 883.0, found 1588.6, 1324.0, 1135.0, 993.2, 883.0.

[0155]

20-3 Deprotection of Acn group

To glycopeptide 45 obtained in the method described in the above 20-2 (44.5 mg, 5.61 μ mol) was added an aqueous solution (2.2 mL) of silver(I) acetate (14.8 mg, 88.7 μ mol), and reacted at room temperature for 30 minutes. DTT (33.2 mg, 215 μ mol) dissolved in 200 mM phosphate buffer (pH 7.4, 2.2 mL) and 100 mM ascorbic acid aqueous solution (561 μ L) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 83:17 $\rightarrow$ 72:28, 15 minutes, linear gradient elution] to obtain glycopeptide 46 (SEQ ID NO. 64) represented by the following formula (46) (34.4 mg, 4.41 μ mol, yield 79%).

[Chemical Formula 63]




ESI-MS: (m/z) calcd for $C_{312}H_{492}N_{56}O_{163}S_5$: $[M+4H]^{4+}$ 1950.0, $[M+5H]^{5+}$ 1560.2, $[M+6H]^{6+}$ 1300.3, found 1949.8, 1560.1, 1300.2.

[0156]

20-4 Formation of disulfide bond

Glycopeptide 46 obtained in the method described in the above 20-3 (34.4 mg, 4.41 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 8.8 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 77:23 $\rightarrow$ 65:35, 16 minutes, linear gradient elution] to obtain a fraction containing compound (S1-2C(disialo)-SRIF28) represented by the following formula (47) (SEQ ID NO. 21).

[Chemical Formula 64]




This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 83:17 -> 72:28, 15 minutes, linear gradient elution] to obtain S1-2C(disialo)-SRIF28 (20.1 mg, 2.58 μ mol, yield 58%).
ESI-MS: (m/z) calcd for $C_{312}H_{490}N_{56}O_{163}S_5$: $[M+4H]^{4+}$ 1949.5, $[M+5H]^{5+}$ 1559.8, $[M+6H]^{6+}$ 1300.0, found 1949.4, 1559.7, 1299.9.

[0157]


Example 21 Synthesis of S1C(disialo)·N5C(disialo)-SRIF28

A compound represented by the following formula (49) (S1C(disialo)·N5C(disialo)-SRIF28) (SEQ ID NO. 22) was synthesized similarly to Example 20, except that a compound represented by the following formula (48) (peptide 48) (SEQ ID NO. 65) was employed instead of peptide 44.

[Chemical Formula 65]



[Chemical Formula 66]




[0158]


Example 22 Synthesis of S1C(disialo)·R13C(disialo)-SRIF28

A compound represented by the following formula (51) (S1C(disialo)·R13C(disialo)-SRIF28) (SEQ ID NO. 23) was synthesized similarly to Example 20, except that a compound represented by the following formula (50) (peptide 50) (SEQ ID NO. 66) was employed instead of peptide 44.

[Chemical Formula 67]



[Chemical Formula 68]




(51)

[0159]

Example 23 Synthesis of N5C(disialo)·A9C(disialo)-SRIF28


A compound represented by the following formula (53) (N5C(disialo)·A9C(disialo)-SRIF28) (SEQ ID NO. 24) was synthesized similarly to Example 20, except that a compound represented by the following formula (52) (peptide 52) (SEQ ID NO. 67) was employed instead of peptide 44.

[Chemical Formula 69]



(52)

[Chemical Formula 70]



(53)

[0160]

Example 24 Synthesis of S1-3C(disialo)-SRIF28


24-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a Prelude™ peptide synthesizer, a protected peptide was synthesized on the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. A part of the crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 50 x 250 mm, flow rate: 43.7 mL/min, eluent

CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 70:30, 5 minutes, then 70:30 $\rightarrow$ 55:45, 15 minutes, linear gradient elution] to obtain glycopeptide 56 (SEQ ID NO. 70) represented by the following formula (56) (7.4 mg, 0.73 μ mol, yield 64%).

[Chemical Formula 73]



(56)

ESI-MS: (m/z) calcd for $C_{401}H_{636}N_{64}O_{226}S_6$: $[M+6H]^{6+}$ 1694.7, $[M+7H]^{7+}$ 1452.7, $[M+8H]^{8+}$ 1271.3, $[M+9H]^{9+}$ 1130.1, $[M+10H]^{10+}$ 1017.2, found 1694.6, 1452.5, 1271.4, 1130.0, 1017.2.

[0163]

24-4 Formation of disulfide bond

Glycopeptide 56 obtained in the method described in the above 24-3 (7.4 mg, 0.73 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.8 mL), and reacted at room temperature for 25 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 70:30, 5 minutes, then 70:30 $\rightarrow$ 69.3:30.7, 5 minutes, linear gradient elution] to obtain a fraction containing compound (S1-3C(disialo)-SRIF28) represented by the following formula (57) (SEQ ID NO. 25).

[Chemical Formula 74]



(57)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 40:60, 20 minutes, linear gradient elution] to obtain S1-3C(disialo)-SRIF28 (4.7 mg, 0.46 μ mol, yield 63%).

ESI-MS: (m/z) calcd for $C_{401}H_{634}N_{64}O_{226}S_6$: $[M+3H]^{3+}$ 3387.7, $[M+4H]^{4+}$ 2541.0, $[M+5H]^{5+}$ 2033.0, $[M+6H]^{6+}$ 1694.3, $[M+7H]^{7+}$ 1452.4, found 3387.6, 2540.9, 2032.7, 1694.2, 1452.3.


[0164]

Example 25 Synthesis


of

S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28

A compound represented by the following formula (59) (S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28) (SEQ ID NO. 26) was synthesized similarly to Example 24, except that peptide 58 (SEQ ID NO. 71) represented by the following formula (58) was employed instead of peptide 54.
[Chemical Formula 75]




[Chemical Formula 76]




[0165]

Example 26 Synthesis of S1C(monosialo)-SRIF28

A compound represented by the following formula (60) (S1C(monosialo)-SRIF28) (SEQ ID NO. 27) was synthesized similarly to Example 1, except that compound b represented by the following formula (b) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) was employed instead of compound a.
[Chemical Formula 77]




[Chemical Formula 78]



In the final product, the ratio between the glycosylated polypeptide having the sugar chain of the following formula b1 and the glycosylated polypeptide having the sugar chain of the following formula b2 was 45:55. Note that it is possible to manufacture a glycosylated polypeptide having substantially uniform sugar chain structure by using monosialo sugar chain derivatives having identical structure.

[Chemical Formula 79]




[0166]


Example 27 Synthesis of S1C(asialo)-SRIF28

A compound represented by the following formula (61) (S1C(asialo)-SRIF28) (SEQ ID NO. 28) was synthesized similarly to Example 1, except that compound c represented by the following formula (c) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) was employed instead of compound a.

[Chemical Formula 80]



[Chemical Formula 81]




[0167]

Example 28 Synthesis of S1-2C(asialo)-SRIF28

28-1 Glycosylation reaction of thiol

Peptide 44 (21.2 mg, 6.21 μ mol) and compound c (44.5 mg, 25.3 μ mol, 4.1 equivalents to peptide 44) were dissolved in 33 mM phosphate buffer (pH 7.4, 1.9 mL), and reacted at room temperature for 1 hour. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 77:23 $\rightarrow$ 62:38, 15 minutes, linear gradient elution] to obtain glycopeptide 62 (SEQ ID NO. 72) represented by the following formula (62) (24.0 mg, 3.54 μ mol, yield 57%).

[Chemical Formula 82]



(62)


ESI-MS: (m/z) calcd for $\text{C}_{274}\text{H}_{434}\text{N}_{54}\text{O}_{133}\text{S}_5$: $[\text{M}+4\text{H}]^{4+}$ 1694.3, $[\text{M}+5\text{H}]^{5+}$ 1355.6, $[\text{M}+6\text{H}]^{6+}$ 1129.8, found 1694.3, 1355.6, 1130.0.

[0168]

28-2 Deprotection of AcM group

To glycopeptide 62 obtained in the method described in the above 28-1 (24.0 mg, 3.54 μ mol) was added an aqueous solution (1.4 mL) of silver(I) acetate (6.0 mg, 36 μ mol), and reacted at room temperature for 3 hours. DTT (14.0 mg, 90.8 μ mol) dissolved in 500 mM phosphate buffer (pH 7.4, 0.57 mL) and 100 mM ascorbic acid aqueous solution (0.35 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 65:35, 15 minutes, linear gradient elution] to obtain glycopeptide 63 (SEQ ID NO. 73) represented by the following formula (63) (20.1 mg, 3.03 μ mol, yield 86%).

[Chemical Formula 83]




(63)

ESI-MS: (m/z) calcd for $\text{C}_{268}\text{H}_{424}\text{N}_{52}\text{O}_{131}\text{S}_5$: $[\text{M}+4\text{H}]^{4+}$ 1658.7, $[\text{M}+5\text{H}]^{5+}$ 1327.2, found 1658.8, 1327.0.

[0169]

28-3 Formation of disulfide bond

Glycopeptide 63 obtained in the method described in the above 28-2 (20.1 mg, 3.03 μmol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 6.1 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 77:23 $\rightarrow$ 65:35, 16 minutes, linear gradient elution] to obtain a fraction containing compound (S1-2C(asialo)-SRIF28) represented by the following formula (64) (SEQ ID NO. 29).
[Chemical Formula 84]



(64)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 92:8 $\rightarrow$ 80:20, 16 minutes, linear gradient elution] to obtain S1-2C(asialo)-SRIF28 (11.0 mg, 1.66 μmol , yield 55%).
ESI-MS: (m/z) calcd for $\text{C}_{268}\text{H}_{422}\text{N}_{52}\text{O}_{131}\text{S}_5$: $[\text{M}+4\text{H}]^{4+}$ 1658.2, $[\text{M}+5\text{H}]^{5+}$ 1326.8, $[\text{M}+6\text{H}]^{6+}$ 1105.8, $[\text{M}+7\text{H}]^{7+}$ 948.0, $[\text{M}+8\text{H}]^{8+}$ 829.6, found 1658.1, 1326.7, 1105.6, 947.8, 829.4.


[0170]

Example 29 Synthesis of S1-3C(asialo)-SRIF28

29-1 Glycosylation reaction of thiol

Peptide 54 (21.3 mg, 6.06 μmol) and compound c (53.4 mg, 30.3 μmol , 5.0 equivalents to peptide 54) were dissolved in 33 mM phosphate buffer (pH 7.4, 1.8 mL), and reacted at room temperature for 1 hour. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 70:30, 20 minutes, linear gradient elution] to obtain glycopeptide 65 (SEQ ID NO. 74) represented by the following formula (65) (39.3 mg, 4.59 μmol , yield 76%).

[Chemical Formula 85]



(65)


ESI-MS: (m/z) calcd for $C_{341}H_{544}N_{60}O_{180}S_6$: $[M+4H]^{4+}$ 2140.2, $[M+5H]^{5+}$ 1712.3, $[M+6H]^{6+}$ 1427.1, found 2140.2, 1712.4, 1427.2.

[0171]

29-2 Deprotection of Acn group

To glycopeptide 65 obtained in the method described in the above 29-1 (39.3 mg, 4.59 μ mol) was added an aqueous solution (1.8 mL) of silver(I) acetate (18.7 mg, 112 μ mol), and reacted at room temperature for 90 minutes. DTT (43.4 mg, 28.1 μ mol) dissolved in 200 mM phosphate buffer (pH 7.4, 1.8 mL) and 100 mM ascorbic acid aqueous solution (0.46 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 68:32, 18 minutes, linear gradient elution] to obtain glycopeptide 66 (SEQ ID NO. 75) represented by the following formula (66) (27.6 mg, 3.28 μ mol, yield 71%).

[Chemical Formula 86]



(66)


ESI-MS: (m/z) calcd for $C_{335}H_{534}N_{58}O_{178}S_6$: $[M+4H]^{4+}$ 2104.6, $[M+5H]^{5+}$ 1683.9, $[M+6H]^{6+}$ 1403.4, found 2104.6, 1684.0, 1403.3.

[0172]

29-3 Formation of disulfide bond

Glycopeptide 66 obtained in the method described in the above 29-2 (27.6 mg, 3.28 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 8.2 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 72:28 $\rightarrow$ 70.5:29.5, 15 minutes, linear gradient elution] to obtain a fraction containing compound (S1-3C(asialo)-SRIF28) represented by the following formula (67) (SEQ ID NO. 30).

[Chemical Formula 87]



(67)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 96:4 $\rightarrow$ 82:18, 20 minutes, linear gradient elution] to obtain S1-3C(asialo)-SRIF28 (14.5 mg, 1.72 μ mol, yield 52%).
ESI-MS: (m/z) calcd for C₃₃₅H₅₃₂N₅₈O₁₇₈S₆: [M+4H]⁴⁺ 2104.1, [M+5H]⁵⁺ 1683.5, [M+6H]⁶⁺ 1403.1, found 2103.7, 1683.3, 1403.0.


[0173]

Example 30 Synthesis of N5N(disialo)-SRIF28

30-1 Solid phase synthesis of glycopeptide

2-chlorotriethyl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Trt)-OH (72.5 mg, 120 μ mol) and DIPEA (104.6 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a Prelude™ peptide synthesizer, a protected peptide 68 (SEQ ID NO. 76) represented by the following formula (68) was synthesized in a state bound to the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.


[Chemical Formula 88]



(68)

Next, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, compound d represented by the following formula (d) (from Otsuka Chemical Co., Ltd.) (411.9 mg, 150.4 μ mol), DMSO-DMF (1/1, v/v, 871 μ L) solution, TBTU (64.2 mg, 200 μ mol), and DIPEA (52.3 μ L, 300 μ mol) were sequentially added to the resin, and this was shaken at room temperature for 3 hours to allow condensation.


[Chemical Formula 89]



(d)

After washing with DMF, this condensation operation was repeated once. After washing the resin with DMF and dichloromethane, this was shaken with 20% piperidine in DMF for 20 minutes to deprotect the Fmoc group, and the resin was washed with DMF to synthesize a protected compound represented by the following formula (69) (peptide 69) (SEQ ID NO. 77) on the resin.


[Chemical Formula 90]



(69)

To this resin, employing HOBt/DIC as the condensation agent, Fmoc-Ser(tBu)-OH, Fmoc-Asn(Trt)-OH, Fmoc-Ala, and Fmoc-Ser(tBu)-OH were condensed in sequence. After condensation, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. Cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. This crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, A:B = 70:30] to obtain glycopeptide 70 (SEQ ID NO. 78) represented by the following formula (70) (29.1 mg, 5.26 μ mol).

[Chemical Formula 91]



(70)


ESI-MS: (m/z) calcd for $C_{235}H_{357}N_{47}O_{100}S_3$: $[M+3H]^{3+}$ 1846.6, $[M+4H]^{4+}$ 1385.2, $[M+5H]^{5+}$ 1108.4, found 1846.5, 1385.1, 1108.3.

[0174]

30-2 Formation of disulfide bond

Glycopeptide 70 obtained in the method described in the above 30-1 (12.2 mg, 2.20 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 5.5 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 66:35, 30 minutes, linear gradient elution] to obtain glycopeptide 71 (SEQ ID NO. 79) represented by the following formula (71) (8.3 mg, 1.5 μ mol, yield 68%).

[Chemical Formula 92]




ESI-MS: (m/z) calcd for $C_{235}H_{355}N_{47}O_{100}S_3$: $[M+3H]^{3+}$ 1846.5, $[M+4H]^{4+}$ 1384.7, $[M+5H]^{5+}$ 1108.0, found 1846.5, 1384.7, 1108.1.

[0175]

30-3 Deprotection of benzyl group

Glycopeptide 71 obtained in the method described in the above 30-2 (7.5 mg, 1.4 μ mol) was dissolved in 50 mM sodium hydroxide aqueous solution (20.6 mL), and reacted at 0°C for 80 minutes. 200 mM acetic acid aqueous solution (5.1 mL) was added, and the mixed solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 66.3:33.7, 20 minutes, linear gradient elution] to obtain a fraction containing compound (N5N(disialo)-SRIF28) represented by the following formula (72) (SEQ ID NO. 31).

[Chemical Formula 93]




[0176]

31-1 Solid phase synthesis of glycopeptide

[Chemical Formula 94]




[Chemical Formula 95]



After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. Cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. This crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, A:B = 71:29] to obtain glycopeptide 75 (SEQ ID NO. 82) represented by the following formula (75) (65.7 mg, 11.8 μ mol).

[Chemical Formula 96]




ESI-MS: (m/z) calcd for $C_{236}H_{358}N_{48}O_{100}S_3$: $[M+4H]^{4+}$ 1392.0, $[M+5H]^{5+}$ 1113.8, found 1391.9, 1113.8.

[0177]

31-2 Formation of disulfide bond

Glycopeptide 75 obtained in the method described in the above 31-1 (20.3 mg, 3.65 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 9.0 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 72:28 $\rightarrow$ 67:33, 30 minutes, linear gradient elution] to obtain glycopeptide 76 (SEQ ID NO. 83) represented by the following formula (76) (17.0 mg, 3.06 μ mol, yield 84%).

[Chemical Formula 97]




ESI-MS: (m/z) calcd for $C_{236}H_{356}N_{48}O_{100}S_3$: $[M+4H]^{4+}$ 1391.5, $[M+5H]^{5+}$ 1113.4, found 1391.3, 1113.2.

[0178]

31-3 Deprotection of benzyl group

Glycopeptide 76 obtained in the method described in the above 31-2 (7.0 mg, 1.3 μmol) was dissolved in 50 mM sodium hydroxide aqueous solution (19.1 mL), and reacted at 0°C for 1 hour. 200 mM acetic acid aqueous solution (9.6 mL) was added, and the mixed solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 85:15 $\rightarrow$ 77.5:22.5, 20 minutes, linear gradient elution] to obtain a compound represented by the following formula (77) (S1N(disialo)-SRIF28) (SEQ ID NO. 32) (2.7 mg, 0.50 μmol , yield 40%).

[Chemical Formula 98]



ESI-MS: (m/z) calcd for $\text{C}_{222}\text{H}_{344}\text{N}_{48}\text{O}_{100}\text{S}_3$: $[\text{M}+3\text{H}]^{3+}$ 1794.9, $[\text{M}+4\text{H}]^{4+}$ 1346.4, $[\text{M}+5\text{H}]^{5+}$ 1077.3, $[\text{M}+6\text{H}]^{6+}$ 897.9, found 1794.7, 1346.2, 1077.2, 897.7.

[0179]


Example 32 Synthesis of S1C(disialo)·N19C(GlcNAc)-SRIF28

32-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μmol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μmol) and DIPEA (104.5 μL , 600 μmol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, in a peptide solid phase synthesis method with Fmoc strategy employing a Prelude™ peptide synthesizer, a protected peptide was synthesized on a resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. A part of the crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent


A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 72:28 -> 64:36, 20 minutes, linear gradient elution] to obtain peptide 78 (SEQ ID NO. 84) represented by the following formula (78) (28.9 mg).
[Chemical Formula 99]



ESI-MS: (m/z) calcd for $C_{146}H_{226}N_{42}O_{39}S_6$: $[M+3H]^{3+}$ 1129.7, $[M+4H]^{4+}$ 847.5, found 1129.5, 847.4.

[0180]


32-2 Glycosylation of thiol and deprotection of StBu Group
Peptide 78 obtained in the method described in the above 32-1 (10.0 mg, 2.95 μ mol) and compound e represented by the following formula (e) (bromoacetamidated monosaccharide: from Otsuka Chemical Co., Ltd.) (2.0 mg, 5.90 μ mol, 2.0 equivalents to peptide 78) were dissolved in 33 mM phosphate buffer (pH 7.4, 0.89 mL) containing 20 μ M TCEP, and reacted at room temperature for 2 hours.
[Chemical Formula 100]



After the reaction, DTT (45.5 mg, 295 μ mol) dissolved in 0.1 M phosphate buffer (pH 7.4, 3.0 mL) was added, and reacted at room temperature for 3 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 65:35, 20 minutes, linear gradient elution] to obtain glycopeptide 79 (SEQ ID

NO. 85) represented by the following formula (79) (6.8 mg, 1.9 μmol , yield 64%).

[Chemical Formula 101]



(79)


ESI-MS: (m/z) calcd for $\text{C}_{152}\text{H}_{234}\text{N}_{44}\text{O}_{45}\text{S}_5$ $[\text{M}+3\text{H}]^{3+}$ 1187.0, $[\text{M}+4\text{H}]^{4+}$ 890.5, found 1187.0, 890.5.

[0181]

32-3 Glycosylation reaction of thiol

Peptide 79 obtained in the method described in the above 32-2 (6.8 mg, 1.9 μmol) and compound a represented by the above formula (a) (22.4 mg, 9.56 μmol , 5.0 equivalents to peptide 79) were dissolved in 0.1 M phosphate buffer (pH 7.4, 2.0 mL) containing 7.6 mM DTT, and reacted at room temperature for 2 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 85:15 $\rightarrow$ 65:35, 20 minutes, linear gradient elution] to obtain glycopeptide 80 (SEQ ID NO. 86) represented by the following formula (80) (3.4 mg, 0.58 μmol , yield 31%).

[Chemical Formula 102]



(80)

ESI-MS: (m/z) calcd for $\text{C}_{238}\text{H}_{373}\text{N}_{51}\text{O}_{107}\text{S}_5$: $[\text{M}+5\text{H}]^{5+}$ 1165.2, $[\text{M}+6\text{H}]^{6+}$ 971.2, $[\text{M}+7\text{H}]^{7+}$ 832.6, $[\text{M}+8\text{H}]^{8+}$ 728.6, found 1165.2, 971.1, 832.6, 728.6.


[0182]

32-4 Deprotection of Acm group

To glycopeptide 80 obtained in the method described in the above 32-3 (3.8 mg, 0.65 μmol) was added an aqueous solution (262 μL) of silver(I) acetate (2.7 mg, 16 μmol), and reacted at room temperature for 1 hour. DTT (10.0 mg, 64.8 μmol) dissolved in 100 mM phosphate buffer (pH 7.4, 426 μL) and 100 mM ascorbic acid aqueous solution (66 μL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10%

water/90% acetonitrile, gradient A:B = 90:10 -> 60:40, 30 minutes, linear gradient elution] to obtain glycopeptide 81 (SEQ ID NO. 87) represented by the following formula (81) (2.5 mg, 0.44 μ mol, yield 67%).

[Chemical Formula 103]



(81)


ESI-MS: (m/z) calcd for $C_{232}H_{363}N_{49}O_{105}S_5$: $[M+3H]^{3+}$ 1894.0, $[M+4H]^{4+}$ 1420.7, $[M+5H]^{5+}$ 1136.8, found 1893.8, 1420.6, 1136.7.

[0183]

32-4 Formation of disulfide bond

Glycopeptide 81 obtained in the method described in the above 32-3 (2.5 mg, 0.44 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.1 mL), and reacted overnight at room temperature. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 -> 70:30, 30 minutes, linear gradient elution] to obtain a compound represented by the following formula (82) (S1C(disialo)·N19C(GlcNAc)-SRIF28) (SEQ ID NO. 33) (1.5 mg, 0.26 μ mol, yield 59%).

[Chemical Formula 104]



(82)


ESI-MS: (m/z) calcd for $C_{232}H_{361}N_{49}O_{105}S_5$: $[M+3H]^{3+}$ 1893.3, $[M+4H]^{4+}$ 1420.2, $[M+5H]^{5+}$ 1136.4, found 1893.5, 1420.1, 1136.3.

[0184]

Example 33 Synthesis of S1C(disialo)·N19C(diMan)-SRIF28


A compound represented by the following formula (83) (S1C(disialo)·N19C(diMan)-SRIF28) (SEQ ID NO. 34) was synthesized similarly to Example 32, except that compound f represented by the following formula (f) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) was employed instead of compound e.

[Chemical Formula 105]



(f)

[Chemical Formula 106]



(8 3)

[0185]


Example 34 Synthesis of S1-5C(disialo)-SRIF28

34-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μmol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μmol) and DIPEA (104.5 μL , 600 μmol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a PreludeTM peptide synthesizer, a protected peptide was synthesized on the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. The crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), ϕ 50 x 250 mm, flow rate: 43.7 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 65:35, 35 minutes, linear gradient elution] to obtain peptide 84 (SEQ ID NO. 90) represented by the following formula (84).

[Chemical Formula 107]



ESI-MS: (m/z) calcd for $\text{C}_{155}\text{H}_{239}\text{N}_{47}\text{O}_{44}\text{S}_8$: $[\text{M}+3\text{H}]^{3+}$ 1241.5, $[\text{M}+4\text{H}]^{4+}$ 931.3, $[\text{M}+5\text{H}]^{5+}$ 745.3, found 1241.2, 931.2, 745.1.


[0186]

34-2 Glycosylation reaction of thiol

Peptide 84 obtained in the method described in the above 34-1 (33.7 mg, 9.06 μmol) and compound a represented by the above formula (a) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) (160.9 mg, 68.6 μmol , 7.5 equivalents to peptide 84) were dissolved in 33 mM phosphate buffer (pH 7.4, 2.7 mL) containing 10 μM TCEP, and reacted overnight at room temperature. The reaction solution was purified with HPLC [column: SHISEIDO Proteonavi (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 78.4:21.6, 20 minutes, linear gradient elution] to obtain glycopeptide 85 (SEQ ID NO. 91) represented by the following formula (85) (35.1 mg, 2.33 μmol , yield 26%).

ESI-MS: (m/z) calcd for $\text{C}_{585}\text{H}_{934}\text{N}_{82}\text{O}_{354}\text{S}_8$: $[\text{M}+6\text{H}]^{6+}$ 2507.1, $[\text{M}+7\text{H}]^{7+}$ 2149.1, $[\text{M}+8\text{H}]^{8+}$ 1880.6, found 2506.9, 2149.0, 1880.6.

[Chemical Formula 108]




[0187]

34-3 Deprotection of Acm group

To glycopeptide 85 obtained in the method described in the above 34-2 (20.6 mg, 1.37 μmol) was added an aqueous solution (0.55 mL) of silver(I) acetate (5.6 mg, 34 μmol), and reacted at room temperature for 50 minutes. Then, DTT (13.8 mg, 89 μmol) dissolved in 200 mM phosphate buffer (pH 7.4, 0.55 mL) and 100 mM ascorbic acid aqueous solution (137 μL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO Proteonavi (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 73.6:26.4,

16 minutes, linear gradient elution] to obtain glycopeptide 86 (SEQ ID NO. 92) represented by the following formula (86) (13.6 mg, 0.913 μ mol, yield 67%).

[Chemical Formula 109]




ESI-MS: (m/z) calcd for $C_{579}H_{924}N_{80}O_{352}S_8$: $[M+6H]^{6+}$ 2483.4, $[M+7H]^{7+}$ 2128.8, $[M+8H]^{8+}$ 1862.8, found 2483.2, 2128.8, 1862.9.

[0188]

34-4 Formation of disulfide bond

Glycopeptide 86 obtained in the method described in the above 34-3 (13.6 mg, 0.913 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 2.2 mL), and reacted at room temperature for 2 days. Then, the reaction solution was purified with HPLC [column: Proteonavi (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 10 mM ammonium acetate aqueous solution, B: 10 mM ammonium acetate-acetonitrile (1/9, v/v), gradient A:B = 78:22 $\rightarrow$ 70.8:29.2, 15 minutes, linear gradient elution] to obtain S1-5C(disialo)-SRIF28 (SEQ ID NO. 88) represented by the following formula (87) (10.4 mg, 0.698 μ mol, yield 76%).

[Chemical Formula 110]



ESI-MS: (m/z) calcd for $C_{579}H_{922}N_{80}O_{352}S_8$: $[M+4H]^{5+}$ 3724.1, $[M+5H]^{5+}$ 2979.5, $[M+6H]^{6+}$ 2483.1, $[M+7H]^{7+}$ 2128.5, $[M+8H]^{8+}$ 1862.6, found 3723.7, 2979.1, 2482.9, 2128.2, 1862.4.

[0189]


Example 35 Synthesis of S1-10C(disialo)-SRIF28

35-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a

Prelude™ peptide synthesizer, a protected peptide was synthesized on the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. The crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 69:31 -> 66.6:33.4, 15 minutes, linear gradient elution] to obtain peptide 88 (SEQ ID NO. 93) represented by the following formula (88) (16.8 mg).
[Chemical Formula 111]




ESI-MS: (m/z) calcd for $C_{170}H_{264}N_{52}O_{49}S_{13}$: $[M+3H]^+$ 1413.4, $[M+4H]^+$ 1060.3, $[M+5H]^+$ 848.4, found 1412.9, 1060.2, 848.1.

[0190]

35-2 Synthesis of disialo sugar chain having
bromoacetamidated benzyl protecting group

To compound a (28.9 mg, 12.3 μ mol) were sequentially added DMF (0.58 mL), lithium bromide (21.5 mg, 248 μ mol), and benzyl bromide (14.6 μ L, 122 μ mol), and reacted at 30°C for 20 hours. Benzyl bromide (14.6 μ L, 122 μ mol) was further added and reacted for 20 hours. To the reaction solution was added toluene (30 mL), and after centrifugal separation (10,000 x g, 10 minutes), the precipitate was dissolved in water (100 μ L) and purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: water:acetonitrile = 95:5 -> 70:30, 20 minutes, linear gradient elution] to obtain compound g represented by the following formula (g) (bromoacetamidated disialo sugar chain: 7.6 mg, 3.0 μ mol, yield 24%).

[Chemical Formula 112]



(g)


MALDI-MS: (m/z) calcd for $C_{100}H_{152}BrN_7O_{62}$: $[M+Na]^+$ 2544.8, found 2544.4.

[0191]

35-3 Glycosylation reaction of thiol

Peptide 88 obtained in the method described in the above 35-1 (8.1 mg, 1.9 μ mol) and compound g obtained in the method described in the above 35-2 (58.0 mg) were dissolved in 66 mM phosphate buffer (pH 6.8, 0.57 mL) containing 2.3 M guanidine hydrochloride, and reacted overnight at room temperature. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 80:20 $\rightarrow$ 74.7:25.3, 30 minutes, linear gradient elution] to obtain glycopeptide 89 (SEQ ID NO. 94) represented by the following formula (89) (12.3 mg, 0.429 μ mol, yield 23%).

[Chemical Formula 113]



(89)

ESI-MS: (m/z) calcd for $C_{1170}H_{1774}N_{122}O_{669}S_{13}$: $[M+8H]^{8+}$ 3584.7, $[M+9H]^{9+}$ 3186.5, $[M+10H]^{10+}$ 2868.0, $[M+11H]^{11+}$ 2607.4, $[M+12H]^{12+}$ 2390.2, $[M+13H]^{13+}$ 2206.4, $[M+14H]^{14+}$ 2048.9, found 3585.0, 3186.6, 2867.9, 2607.6, 2390.1, 2206.4, 2048.9.


[0192]

35-4 Deprotection of Acn group

To glycopeptide 89 obtained in the method described in the above 35-3 (49.8 mg, 1.74 μ mol) was added an aqueous

solution (0.70 mL) of silver(I) acetate (10.2 mg), and reacted at room temperature for 1 hour. A solution of DTT (16.6 mg) in 200 mM phosphate buffer (pH 7.4, 0.70 mL) and 100 mM ascorbic acid aqueous solution (0.17 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 78:22 $\rightarrow$ 73:27, 20 minutes, linear gradient elution] to obtain glycopeptide 90 (SEQ ID NO. 95) represented by the following formula (90) (33.0 mg, 1.16 μ mol, yield 67%).

[Chemical Formula 114]




ESI-MS: (m/z) calcd for $C_{1164}H_{1764}N_{120}O_{867}S_{13}$: $[M+9H]^{9+}$ 3170.8, $[M+10H]^{10+}$ 2853.8, $[M+11H]^{11+}$ 2594.4, $[M+12H]^{12+}$ 2378.3, $[M+13H]^{13+}$ 2195.4, $[M+14H]^{14+}$ 2038.7, $[M+15H]^{15+}$ 1902.9, found 3170.9, 2853.7, 2594.4, 2378.3, 2195.4, 2038.7, 1902.9.

[0193]

35-5 Formation of disulfide bond

Glycopeptide 90 obtained in the method described in the above 35-4 (2.1 mg, 0.074 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 185 μ L), and reacted at room temperature for 2 days. Then, the reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 4.6 x 250 mm, flow rate: 0.7 mL/min, eluent A: 10 mM ammonium acetate aqueous solution, B: 10 mM ammonium acetate-acetonitrile (1/9, v/v), gradient A:B = 69:31 $\rightarrow$ 65:35, 20 minutes, linear gradient elution] to obtain glycopeptide 91 (SEQ ID NO. 96) represented by the following formula (91) (1.0 mg, 0.035 μ mol, yield 47%).

[Chemical Formula 115]




ESI-MS: (m/z) calcd for $C_{1164}H_{1762}N_{120}O_{667}S_{13}$: $[M+8H]^{8+}$ 3566.7, $[M+9H]^{9+}$ 3170.5, $[M+10H]^{10+}$ 2853.6, found 3566.6, 3170.4, 2853.6.

[0194]

35-6 Deprotection of benzyl group

Glycopeptide 91 obtained in the method described in the above 35-5 (17.1 mg, 0.599 μmol) was dissolved in 50 mM sodium hydroxide aqueous solution (12 mL), and reacted at 0°C for 90 minutes. 100 mM acetic acid aqueous solution 12 mL was added, and the mixed solution was purified with HPLC [column: SHISEIDO Proteonavi, $\phi 4.6 \times 250$ mm, flow rate: 0.7 mL/min, eluent A: 10 mM ammonium acetate aqueous solution, B: 10 mM ammonium acetate-acetonitrile (1/9, v/v), gradient A:B = 72:18 $\rightarrow$ 80:20, 20 minutes, linear gradient elution] to obtain glycopeptide 92 (SEQ ID NO. 89) represented by the following formula (92) (5.8 mg, 0.22 μmol , yield 37%).

[Chemical Formula 116]




ESI-MS: (m/z) calcd for $\text{C}_{1024}\text{H}_{1642}\text{N}_{120}\text{O}_{667}\text{S}_{13}$: $[\text{M}+7\text{H}]^{7+}$ 3818.6, $[\text{M}+8\text{H}]^{8+}$ 3341.4, $[\text{M}+9\text{H}]^{9+}$ 2970.3, $[\text{M}+10\text{H}]^{10+}$ 2673.3, found 3818.3, 3341.0, 2970.1, 2673.1.

[0195]


Example 36 Synthesis of C(disialo)-SRIF28

A compound represented by the following formula (94) (C(disialo)-SRIF28) (SEQ ID NO. 98) was synthesized similarly to Example 1, except that a compound represented by the following formula (93) (peptide 93) (SEQ ID NO. 97) was employed instead of peptide 1.

[Chemical Formula 117]




[Chemical Formula 118]




[0196]

Example 37 Synthesis of R11C(disialo)-SRIF28

A compound represented by the following formula (96) (R11C(disialo)-SRIF28) (SEQ ID NO. 100) was synthesized similarly to Example 1, except that a compound represented by the following formula (95) (peptide 95) (SEQ ID NO. 99) was employed instead of peptide 1.
[Chemical Formula 119]



[Chemical Formula 120]




[0197]


Example 38 Synthesis of F20C(disialo)-SRIF28

A compound represented by the following formula (98) (F20C(disialo)-SRIF28) (SEQ ID NO. 102) was synthesized similarly to Example 1, except that a compound represented by the following formula (97) (peptide 97) (SEQ ID NO. 101) was employed instead of peptide 1.

[Chemical Formula 121]



[Chemical Formula 122]




[0198]

Example 39 Synthesis of T24C(disialo)-SRIF28


A compound represented by the following formula (100) (T24C(disialo)-SRIF28) (SEQ ID NO. 104) was synthesized similarly to Example 1, except that a compound represented by the following formula (99) (peptide 99) (SEQ ID NO. 103) was employed instead of peptide 1.

[Chemical Formula 123]



(9 9)

[Chemical Formula 124]




(1 0 0)

[0199]

Example 40 Synthesis of F25C(disialo)-SRIF28


A compound represented by the following formula (102) (F25C(disialo)-SRIF28) (SEQ ID NO. 106) was synthesized similarly to Example 1, except that a compound represented by the following formula (101) (peptide 101) (SEQ ID NO. 105) was employed instead of peptide 1.

[Chemical Formula 125]



(1 0 1)

[Chemical Formula 126]




(1 0 2)

[0200]

Example 41 Synthesis of S27C(disialo)-SRIF28


A compound represented by the following formula (104) (S27C(disialo)-SRIF28) (SEQ ID NO. 108) was synthesized similarly to Example 1, except that a compound represented by the following formula (103) (peptide 103) (SEQ ID NO. 107) was employed instead of peptide 1.

[Chemical Formula 127]



(1 0 3)

[Chemical Formula 128]




(104)

[0201]

Example 42 Synthesis of C(disialo)-K-SRIF14


A compound represented by the following formula (106) (C(disialo)-K-SRIF14) (SEQ ID NO. 110) was synthesized similarly to Example 1, except that a compound represented by the following formula (105) (peptide 105) (SEQ ID NO. 109) was employed instead of peptide 1.

[Chemical Formula 129]



(105)

[Chemical Formula 130]




(106)

[0202]

Example 43 Synthesis of S1C(disialo)-F25Y-SRIF28


A compound represented by the following formula (108) (S1C(disialo)-F25Y-SRIF28) (SEQ ID NO. 112) was synthesized similarly to Example 1, except that a compound represented by the following formula (107) (peptide 107) (SEQ ID NO. 111) was employed instead of peptide 1.

[Chemical Formula 131]




(107)

[Chemical Formula 132]



(108)




ESI-MS: (m/z) calcd for $C_{94}H_{138}N_{22}O_{26}S_3$: $[M+2H]^{2+}$ 1045.2, $[M+3H]^{3+}$ 697.1, found 1045.0, 697.0.

[0205]

45-2 Synthesis of C(disialo)-PEG linker-SRIF14

A compound represented by the following formula (112) (C(disialo)-PEG linker-SRIF14) (SEQ ID NO. 116) was synthesized similarly to Example 1, except that peptide 111 obtained in the method described in the above 45-1 was employed instead of peptide 1.

[Chemical Formula 136]




[0206]

Example 46 Synthesis of Biotin-S1C(disialo)-SRIF28

46-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a dichloromethane (3.0 mL) solution containing Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a Prelude™ peptide synthesizer, a protected peptide 113 (SEQ ID NO. 117) represented by the following formula (113) was synthesized in a state bound to the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.


[Chemical Formula 137]



The Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, employing HCTU as the condensation agent, biotin was

condensed. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. This leads to the detachment of the protecting group of the amino acid side chain (other than the Acn group), as well as cleaving between the peptide and the resin. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. The crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 -> 61:39, 18 minutes, linear gradient elution] to obtain peptide 114 (SEQ ID NO. 118) represented by the following formula (114).

[Chemical Formula 138]



(114)


ESI-MS: (m/z) calcd for $\text{C}_{153}\text{H}_{233}\text{N}_{45}\text{O}_{42}\text{S}_5$: $[\text{M}+3\text{H}]^{3+}$ 1179.4, $[\text{M}+4\text{H}]^{4+}$ 884.8, $[\text{M}+5\text{H}]^{5+}$ 708.0, found 1179.2, 884.4, 707.9.

[0207]

46-2 Synthesis of Biotin-S1C(disialo)-SRIF28

A compound represented by the following formula (115) (Biotin-S1C(disialo)-SRIF28) (SEQ ID NO. 119) was synthesized similarly to Example 1, except that peptide 114 obtained in the method described in the above 46-1 was employed instead of peptide 1.

[Chemical Formula 139]




[0208]

Example 47 Synthesis of Biotin-PEG linker-S1C(disialo)-SRIF28

47-1 Synthesis of peptide

The Fmoc protecting group of the protected peptide 113 (SEQ ID NO. 117) bound to the resin, obtained in the above 46-1, was removed by treating with 20% piperidine in DMF. After washing with DMF, employing HCTU as the condensation agent, Fmoc-NH-(PEG)₂-COOH (from Merck) and biotin were condensed in sequence. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. The crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), φ20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 -> 61:39, 22 minutes, linear gradient elution] to obtain peptide 116 (SEQ ID NO. 120) represented by the following formula (116).

[Chemical Formula 140]




ESI-MS: (m/z) calcd for C₁₆₂H₂₅₀N₄₆O₄₆S₅: [M+3H]³⁺ 1247.1, [M+4H]⁴⁺ 935.6, [M+5H]⁵⁺ 748.7, found 1246.9, 935.4, 748.6.

[0209]

47-2 Synthesis of Biotin-PEG linker-S1C(disialo)-SRIF28

A compound represented by the following formula (117) (Biotin-PEG linker-S1C(disialo)-SRIF28) (SEQ ID NO. 121) was synthesized similarly to Example 1, except that peptide 116 obtained in the method described in the above 47-1 was employed instead of peptide 1.

[Chemical Formula 141]



(117)


[0210]

Example 48 Synthesis of Azido-S1C(disialo)-SRIF28

48-1 Synthesis of peptide

The Fmoc protecting group of the protected peptide 113 (SEQ ID NO. 117) bound to the resin, obtained in the above 46-1, was removed by treating with 20% piperidine in DMF. After washing with DMF, employing HCTU as the condensation agent, 5-Azido-pentanoic acid was condensed. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. The crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 70:30 -> 60:40, 20 minutes, linear gradient elution] to obtain peptide 118 (SEQ ID NO. 122) represented by the following formula (118).

[Chemical Formula 142]



(118)


ESI-MS: (m/z) calcd for $C_{148}H_{226}N_{46}O_{41}S_4$: $[M+3H]^{3+}$ 1145.6, $[M+4H]^{4+}$ 859.5, $[M+5H]^{5+}$ 687.8, found 1145.5, 859.2, 687.5.

[0211]

48-2 Synthesis of Azido-S1C(disialo)-SRIF28

A compound represented by the following formula (119) (Azido-S1C(disialo)-SRIF28) (SEQ ID NO. 123) was synthesized similarly to Example 1, except that peptide 118 obtained in the method described in the above 48-1 was employed instead of peptide 1.

[Chemical Formula 143]




[0212]


Example 49 Synthesis of S1C(disialo)·E12C(disialo)-SRIF28

A compound represented by the following formula (121) (S1C(disialo)·E12C(disialo)-SRIF28) (SEQ ID NO. 125) was synthesized similarly to Example 20, except that a compound represented by the following formula (120) (peptide 120) (SEQ ID NO. 124) was synthesized and employed instead of peptide 44.

[Chemical Formula 144]




[Chemical Formula 145]




Example 50 Synthesis of 2C(disialo)-R-K-SRIF14

A compound represented by the following formula (123) (2C(disialo)-R-K-SRIF14) (SEQ ID NO. 127) was synthesized similarly to Example 20, except that a compound represented by the following formula (122) (peptide 122) (SEQ ID NO. 126) was synthesized and employed instead of peptide 44.

[Chemical Formula 146]



[Chemical Formula 147]




[0213]


Example 51 Synthesis of 3C(disialo)-R-K-SRIF14

A compound represented by the following formula (125) (3C(disialo)-R-K-SRIF14) (SEQ ID NO. 129) was synthesized similarly to Example 24, except that a compound represented by the following formula (124) (peptide 124) (SEQ ID NO. 128) was synthesized and employed instead of peptide 54.

[Chemical Formula 148]



[Chemical Formula 149]




Example 52 Synthesis of S1C(diGlcNAc)-SRIF28

52-1 Glycosylation reaction of thiol

Peptide 1 (SEQ ID NO. 38) (from APC, Inc.) (25.0 mg, 7.56 μ mol) and compound h represented by the following formula (h) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) (15.6 mg, 15.1 μ mol, 2.0 equivalents to peptide 1) were dissolved in 33 mM phosphate buffer (pH 7.4, 2.3 mL), and reacted at room temperature for 30 minutes.

[Chemical Formula 150]




(h)

[0214]

The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, A:B = 75:25 $\rightarrow$ 62:38, 13 minutes, linear gradient elution] to obtain glycopeptide 126 (SEQ ID NO. 130) represented by the following formula (126) (25.6 mg, 6.01 μ mol, yield 79%).

[Chemical Formula 151]



(126)

ESI-MS: (m/z) calcd for $C_{195}H_{304}N_{48}O_{76}S_4$: $[M+3H]^{3+}$ 1556.0, $[M+4H]^{4+}$ 1167.3, found 1555.7, 1167.0.


[0215]

52-2 Deprotection of AcM group

To glycopeptide 126 obtained in the method described in the above 52-1 (28.3 mg, 6.07 μ mol) was added an aqueous solution (2.4 mL) of silver(I) acetate (12.5 mg, 74.5 μ mol), and reacted at room temperature for 30 minutes. DTT (28.8 mg, 187 μ mol) dissolved in 200 mM Tris-HCl buffer (pH 7.4, 2.4 mL) and 100 mM ascorbic acid aqueous solution (0.6 mL) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 60:40, 13 minutes, linear gradient elution] to obtain glycopeptide

127 (SEQ ID NO. 131) represented by the following formula (127) (19.8 mg, 4.38 μ mol, yield 72%).

[Chemical Formula 152]




ESI-MS: (m/z) calcd for $C_{189}H_{294}N_{46}O_{74}S_4$: $[M+3H]^{3+}$ 1508.6, $[M+4H]^{4+}$ 1131.7, $[M+5H]^{5+}$ 905.6, found 1508.3, 1131.5, 905.4.

[0216]

52-3 Formation of disulfide bond

Glycopeptide 127 obtained in the method described in the above 52-2 (19.8 mg, 4.38 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 8.8 mL), and reacted at room temperature for 2 days. The reaction solution was crudely purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 60:40, 13 minutes, linear gradient elution] to obtain a fraction containing compound S1C(diGlcNAc)-SRIF28 represented by the following formula (128) (SEQ ID NO. 132).

[Chemical Formula 153]



This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 78:22, 12 minutes, linear gradient elution] to obtain S1C(diGlcNAc)-SRIF28 (11.9 mg, 2.63 μ mol, yield 60%).


ESI-MS: (m/z) calcd for $C_{189}H_{292}N_{46}O_{74}S_4$: $[M+3H]^{3+}$ 1508.0, $[M+4H]^{4+}$ 1131.2, $[M+5H]^{5+}$ 905.2, found 1507.7, 1131.0, 905.0.

[0217]

Example 53 Synthesis of S1C(diMan)-SRIF28

A compound represented by the following formula (129) (S1C(diMan)-SRIF28) (SEQ ID NO. 133) was synthesized similarly to Example 52, except that compound f was employed instead of compound h.

[Chemical Formula 154]




(129)

[0218]

Example 54 Synthesis of N19C(diMan)-SRIF28

A compound represented by the following formula (130) (N19C(diMan)-SRIF28) (SEQ ID NO. 134) was synthesized similarly to Example 52, except that peptide 21 was employed instead of peptide 1 and compound f was employed instead of compound h.

[Chemical Formula 155]




(130)

[0219]

Example 55 Synthesis of S1C(GlcNAc)-SRIF28

A compound represented by the following formula (131) (S1C(GlcNAc)-SRIF28) (SEQ ID NO. 135) was synthesized similarly to Example 52, except that compound e was employed instead of compound h.

[Chemical Formula 156]




(131)

[0220]

Example 56 Synthesis of N19C(GlcNAc)-SRIF28

A compound represented by the following formula (132) (N19C(GlcNAc)-SRIF28) (SEQ ID NO. 136) was synthesized similarly to Example 52, except that peptide 21 was employed instead of peptide 1 and compound e was employed instead of compound h.

[Chemical Formula 157]




(132)

[0221]


Example 57 Synthesis of S1C(trisialo)-SRIF28

A compound represented by the following formula (133) (S1C(trisialo)-SRIF28) (SEQ ID NO. 137) was synthesized similarly to Example 1, except that compound i represented by the following formula (i) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) was employed instead of compound a.

[Chemical Formula 158]



[Chemical Formula 159]




[0222]

Example 58 Synthesis of S1C(tetrasialo)-SRIF28


A compound represented by the following formula (134) (S1C(tetrasialo)-SRIF28) (SEQ ID NO. 138) was synthesized similarly to Example 1, except that compound j represented by the following formula (j) (bromoacetamidated oligosaccharide: from Otsuka Chemical Co., Ltd.) was employed instead of compound a.

[Chemical Formula 160]



(j)

[Chemical Formula 161]




[0223]

Example 59 Synthesis of SlC(disialo(aminoethylamide))-SRIF28


59-1 Synthesis of bromoacetamidated disialo sugar chain derivative

To compound k represented by the following formula (k) (from Otsuka Chemical Co., Ltd.) (204.1 mg, 92.1 μ mol) were added water (2 mL) and tert-butyl N-(2-aminoethyl)carbamate (0.29 mL, 0.18 mmol), and this was stirred at room temperature for 1 hour. After lyophilization, to the lyophilizate obtained were added DMF (5 mL), HATU (349 mg, 921 μ mol), and DIPEA (161 μ L, 921 μ mol), and reacted at 37°C for 18 hours. Toluene (50 mL) was added to the solution, and the deposited precipitate was collected by filtration. The precipitate was dissolved in DMF (5 mL), purified with gel filtration purification [column: Sephadex G-25, ϕ 20 x 200 mm, flow rate: 30 mL/h], and the fraction of interest was collected and lyophilized to obtain compound l represented by the following formula (l) (152.4 mg, 60.8 μ mol, yield 66%).

[Chemical Formula 162]




[Chemical Formula 163]



MALDI-MS: (m/z) calcd for $C_{98}H_{166}N_{10}O_{64}$: $[M+Na]^+$ 2530.0, found 2529.4.

Compound 1 (100 mg, 39.8 μ mol) and ammonium hydrogen carbonate (31.4 mg, 398 μ mol) were dissolved in water (1 mL), and reacted at room temperature for 7 days. After lyophilization, to the lyophilizate obtained were sequentially added water (1 mL), DCC (41.0 mg, 199 μ mol), and bromoacetic acid (27.7 mg, 199 μ mol) dissolved in DMF (1 mL). After 1 hour of reaction under ice cooling, the solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: water:acetonitrile = 84:16] to obtain compound m represented by the following formula (m) (bromoacetamidated disialo sugar chain derivative: 100 mg, 38.2 μ mol, yield 96%).

[Chemical Formula 164]



MALDI-MS: (m/z) (m/z) calcd for $C_{100}H_{168}BrN_{11}O_{64}$: $[M+Na]^+$
2648.9, found 2648.5.


[0224]

59-2 Glycosylation reaction of thiol

Compound m obtained in the method described in the above 59-1 (14.2 mg, 5.40 μ mol) and peptide 1 (15.0 mg, 4.53 μ mol) were dissolved in 33 mM phosphate buffer (pH 7.4, 1.3 mL), and reacted at room temperature for 30 minutes. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% acetic acid (AcOH) water, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 86:14 $\rightarrow$ 82:18, 20 minutes, linear gradient elution] to obtain glycopeptide 135 (SEQ ID NO. 139) represented by the following formula (135) (13.4 mg, 2.29 μ mol, yield 51%).

ESI-MS: (m/z) calcd for $C_{243}H_{386}N_{54}O_{104}S_4$: $[M+4H]^{4+}$ 1465.1, $[M+5H]^{5+}$ 1172.2, $[M+6H]^{6+}$ 977.0, found 1464.9, 1172.1, 977.1.

[Chemical Formula 165]




(1 3 5)

[0225]

59-3 Deprotection of Boc group

Glycopeptide 135 obtained in the method described in the above 59-2 (13.4 mg, 2.29 μ mol) was dissolved in 95% TFA aqueous solution (458 μ L), and this was shaken at room temperature for 5 minutes. After adding 50 mM ammonium acetate aqueous solution (pH 6.8, 33 mL), the reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 65:35, 10 minutes, linear gradient elution] to obtain glycopeptide 136 (SEQ ID NO. 140) represented by the following formula (136) (12.7 mg, 98 μ mol, yield 98%).
[Chemical Formula 166]



(136)

ESI-MS: (m/z) calcd for $C_{233}H_{370}N_{54}O_{100}S_4$: $[M+4H]^+$ 1415.1, $[M+5H]^+$ 1132.2, $[M+6H]^+$ 943.7, $[M+7H]^+$ 809.0, found 1414.9, 1132.1, 943.6, 808.9.

[0226]

59-4 Deprotection of Acm group

To glycopeptide 136 obtained in the method described in the above 59-3 (12.7 mg, 2.25 μ mol) was added an aqueous solution (0.9 mL) of silver(I) acetate (9.2 mg, 55 μ mol), and reacted at room temperature for 30 minutes. Then, DTT (21.2 mg, 137 μ mol) dissolved in 200 mM phosphate buffer (pH 7.4, 0.9 mL) and 100 mM ascorbic acid aqueous solution (225 μ L) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 73:27 $\rightarrow$ 60:40, 13 minutes, linear gradient elution] to obtain glycopeptide 137 (SEQ ID NO. 141) represented by the following formula (136) (5.2 mg, 0.94 μ mol, yield 42%).
[Chemical Formula 167]

Disialo(aminoethylamide)

Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

(137)

ESI-MS: (m/z) calcd for $C_{227}H_{360}N_{52}O_{98}S_4$: $[M+4H]^{4+}$ 1379.5, $[M+5H]^{5+}$ 1103.8, $[M+6H]^{6+}$ 920.0, $[M+7H]^{7+}$ 788.7, found 1379.4, 1103.7, 919.9, 788.6.

[0227]

59-5 Formation of disulfide bond

Glycopeptide 137 obtained in the method described in the above 59-4 (5.2 mg, 0.94 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.9 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 70:30 $\rightarrow$ 65:35, 13 minutes, linear gradient elution] to obtain a fraction containing compound (S1C(disialo(aminoethylamide))-SRIF28) represented by the following formula (138) (SEQ ID NO. 142).

[Chemical Formula 168]

Disialo(aminoethylamide)

Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

(138)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 92:8 $\rightarrow$ 85:15, 14 minutes, linear gradient elution] to obtain S1C(disialo(aminoethylamide))-SRIF28 (3.8 mg, 0.69 μ mol, yield 73%).

ESI-MS: (m/z) calcd for $C_{227}H_{358}N_{52}O_{98}S_4$: $[M+3H]^{3+}$ 1838.3, $[M+4H]^{4+}$ 1379.0, $[M+5H]^{5+}$ 1103.4, $[M+6H]^{6+}$ 919.6, $[M+7H]^{7+}$ 788.4, found 1838.0, 1378.5, 1103.2, 919.5, 788.2.


[0228]

Example 60 Synthesis of S1C(disialo(amide))-SRIF28

60-1 Synthesis of bromoacetamidated disialo sugar chain derivative

To compound k (from Otsuka Chemical Co., Ltd.) (152 mg, 68.6 μ mol) were sequentially added DMF (1.9 mL), lithium bromide (357 mg, 4.12 mmol), and phenacyl chloride (273 mg, 1.37 mmol), and reacted at 37°C. After 10 hours, water (19 mL) was added and the precipitate was removed by filtration. To the filtrate was added 25% ammonium water (5 mL), and reacted at room temperature for 18 hours, and then 100 mM phosphate buffer (pH 7.4, 80 mL) was added to allow neutralization. The solution was purified with HPLC [column: YMC Hydrosphere C18 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: 0.1% aqueous TFA:acetonitrile = 98:2 $\rightarrow$ 92:8, 30 minutes, linear gradient elution] and lyophilized to obtain compound n represented by the following formula (n) (60.9 mg, 27.4 μ mol, yield 40%). MALDI-MS: (m/z) calcd for $C_{84}H_{140}N_8O_{60}$: [M+Na]⁺ 2243.8, found 2243.6.


[Chemical Formula 169]



(n)

The obtained intermediate n (37.3 mg, 16.8 μ mol) and ammonium hydrogen carbonate (13.3 mg, 168 μ mol) were dissolved in water (0.37 mL), and reacted at room temperature for 7 days. After lyophilization, to the lyophilizate obtained were sequentially added water (0.37 mL), DCC (17.3 mg, 84 μ mol), and bromoacetic acid (11.7 mg, 84 μ mol) dissolved in DMF (0.37 mL). The solution after 1 hour of reaction under ice cooling was purified with HPLC [column: YMC Hydrosphere C18 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: 0.1% aqueous TFA:acetonitrile = 98:2 $\rightarrow$ 92:8, 30 minutes, linear gradient elution] to obtain compound o represented by the following formula (o)

(bromoacetamidated disialo sugar chain derivative: 29.0 mg, 12.4 μmol , yield 74%).
[Chemical Formula 170]



(o)


MALDI-MS: (m/z) calcd for $\text{C}_{86}\text{H}_{142}\text{BrN}_9\text{O}_{60}$: $[\text{M}+\text{Na}]^+$ 2362.7, found 2362.5.

[0229]

60-2 Synthesis of S1C(disialo(amide))-SRIF28

A compound represented by the following formula (139) (S1C(disialo(amide))-SRIF28) (SEQ ID NO. 143) was synthesized similarly to Example 1, except that compound o obtained in the method described in the above 60-1 was employed instead of compound a.

[Chemical Formula 171]



(139)

[0230]

Example 61 Synthesis of S1C(disialo(Bn))-SRIF28

A compound represented by the following formula (140) (S1C(disialo(Bn))-SRIF28) (SEQ ID NO. 144) was synthesized similarly to Example 1, except that compound g was employed instead of compound a.

[Chemical Formula 172]

Disialo(Bn)

Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Tip-Lys-Thr-Phe-Thr-Ser-Cys

(140)


[0231]

Example 62 Synthesis of S1C(disialo (hexadecylamide))-SRIF28

62-1 Synthesis of bromoacetamidated disialo sugar chain derivative

To compound k (from Otsuka Chemical Co., Ltd.) (140 mg, 63.0 μ mol) were added water (1.5 mL), methanol (1.5 mL), and hexadecylamine (300 mg, 126 μ mol), and this was stirred at room temperature for 1 hour. After lyophilization, to the lyophilizate obtained were added DMF (5 mL), HATU (239 mg, 630 μ mol), and DIPEA (110 μ L, 630 μ mol), and reacted at 37°C. After 18 hours, diethyl ether (100 mL) was added to the solution, and the deposited precipitate was collected by filtration. This precipitate was dissolved in DMF (5 mL) and purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: water:acetonitrile = 40:60 $\rightarrow$ 10:90, 30 minutes, linear gradient elution] to obtain compound p represented by the following formula (p) (71.1 mg, 26.6 μ mol, yield 42%).

[Chemical Formula 173]




(p)

MALDI-MS: (m/z) calcd for $C_{116}H_{204}N_8O_{60}$: $[M+Na]^+$ 2692.3, found 2691.9.

[0232]

The obtained compound p (71.7 mg, 26.6 μ mol) and ammonium hydrogen carbonate (21.8 mg, 266 μ mol) were dissolved in water (0.7 mL) and methanol (0.7 mL), and reacted at room temperature. After 7 days, to the lyophilizate obtained by lyophilization were sequentially added water (0.7 mL), methanol (0.7 mL), DCC (27.4 mg, 133 μ mol), and bromoacetic acid (18.5 mg, 133 μ mol) dissolved in DMF (0.7 mL). After 1 hour of reaction under ice cooling, the solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 8.0 mL/min, eluent: water:acetonitrile = 40:60 $\rightarrow$ 10:90, 30 minutes, linear gradient elution] to obtain compound q represented by the following formula (q) (bromoacetamidated disialo sugar chain: 24.9 mg, 8.9 μ mol, yield 33%).

[Chemical Formula 174]



(q)


MALDI-MS: (m/z) calcd for $C_{118}H_{206}BrN_9O_{60}$: $[M+Na]^+$ 2811.2, found 2811.0.

[0233]

62-2 Glycosylation reaction of thiol

Peptide 1 (10.4 mg, 3.14 μ mol) was dissolved in 0.5 M phosphate buffer (pH 7.4, 62 μ L) containing 30 μ M TCEP. To this solution was added a solution of compound q obtained in the method described in the above 62-1 (79.4 mg, 28.5 μ mol) in DMSO (3.7 mL), and reacted at room temperature for 20 minutes. The reaction solution was purified with HPLC [column: SHISEIDO Proteonavi (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 50:50 $\rightarrow$ 22:78, 14 minutes, linear gradient elution] to obtain

glycopeptide 141 (SEQ ID NO. 145) represented by the following formula (141) (6.4 mg, 1.1 μ mol, yield 35%).
[Chemical Formula 175]



(141)


ESI-MS: (m/z) calcd for $C_{261}H_{424}N_{52}O_{100}S_4$: $[M+3H]^{3+}$ 2007.2, $[M+4H]^{4+}$ 1505.7, $[M+5H]^{5+}$ 1204.7, $[M+6H]^{6+}$ 1004.1, found 2007.4, 1505.5, 1204.8, 1004.0.

[0234]

62-3 Deprotection of Acn group

To glycopeptide 141 obtained in the method described in the above 62-2 (6.4 mg, 1.1 μ mol) was added an aqueous solution (0.8 mL) of silver(I) acetate (3.8 mg, 23 μ mol), and reacted at room temperature for 40 minutes. Then, DTT (8.8 mg, 57 μ mol) dissolved in 200 mM phosphate buffer (pH 7.4, 377 μ L) and 100 mM ascorbic acid aqueous solution (106 μ L) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO Proteonavi (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 48:52 $\rightarrow$ 38:62, 3 minutes, then 38:62 $\rightarrow$ 30:70, 8 minutes, linear gradient elution] to obtain glycopeptide 142 (SEQ ID NO. 146) represented by the following formula (142) (3.2 mg, 0.54 μ mol, yield 49%).

[Chemical Formula 176]



(142)

ESI-MS: (m/z) calcd for $C_{255}H_{414}N_{50}O_{98}S_4$: $[M+3H]^{3+}$ 1959.9, $[M+4H]^{4+}$ 1470.1, $[M+5H]^{5+}$ 1176.3, $[M+6H]^{6+}$ 980.4, found 1959.6, 1469.9, 1176.1, 980.5.

[0235]

62-3 Formation of disulfide bond

Glycopeptide 142 obtained in the method described in the above 62-2 (3.2 mg, 0.54 μmol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.1 mL), and reacted at room temperature for 2 days. The reaction solution was purified with HPLC [column: SHISEIDO Proteonavi (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 48:52 $\rightarrow$ 38:62, 3 minutes, then 38:62 $\rightarrow$ 30:70, 8 minutes, linear gradient elution] to obtain a compound represented by the following formula (143) (S1C(disialo (hexadecylamide))-SRIF28) (SEQ ID NO. 147) (2.8 mg, 0.48 μmol , yield 89%).
[Chemical Formula 177]

Disialohexadecylamide

Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

(143)

ESI-MS: (m/z) calcd for $\text{C}_{255}\text{H}_{412}\text{N}_{50}\text{O}_{98}\text{S}_4$: $[\text{M}+3\text{H}]^{3+}$ 1959.2, $[\text{M}+4\text{H}]^{4+}$ 1469.6, $[\text{M}+5\text{H}]^{5+}$ 1175.9, $[\text{M}+6\text{H}]^{6+}$ 980.1, found 1958.9, 1469.4, 1175.7, 979.9.

[0236]

Example 63 Synthesis of S1-2C(disialo(amide))-SRIF28

A compound represented by the following formula (144) (S1-2C(disialo(amide))-SRIF28) (SEQ ID NO. 148) was synthesized similarly to Example 28, except that compound o was employed instead of compound c.

[Chemical Formula 178]

Disialo(amide)
Disialo(amide)

Cys-Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys


(144)

[0237]

Example 64 Synthesis of S1-2C(disialo(Bn))-SRIF28

A compound represented by the following formula (145) (S1-2C(disialo(Bn))-SRIF28) (SEQ ID NO. 149) was synthesized similarly to Example 28, except that compound g was employed instead of compound c.

[Chemical Formula 179]




[0238]

Example 65 Synthesis of S1C(Asn(disialo))-SRIF28


A compound represented by the following formula (146) (S1C(Asn(disialo))-SRIF28) (SEQ ID NO. 150) was synthesized similarly to Example 1, except that compound r represented by the following formula (r) (bromoacetylated glycosylated Asn: from Otsuka Chemical Co., Ltd.) was employed instead of compound a.

[Chemical Formula 180]



(r)

[Chemical Formula 181]



[0239]


Example 66 Synthesis of S1N(disialo)·N19C(diMan)-SRIF28

66-1 Solid phase synthesis of glycopeptide

2-chlorotrityl chloride resin (100 μmol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Trt)-OH (72.5 mg, 120 μmol) and DIPEA (104.6 μL, 600 μmol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20%

piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a Prelude™ peptide synthesizer, a protected peptide 147 (SEQ ID NO. 151) represented by the following formula (147) was synthesized in a state bound to the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

[Chemical Formula 182]




(147)

[0240]

Next, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, compound d (141.0 mg, 51.5 μ mol), DMSO-DMF (1/1, v/v, 1.1 mL) solution, TBTU (21.2 mg, 66.0 μ mol), and DIPEA (17.2 μ L, 98.7 μ mol) were sequentially added to the resin, and this was shaken at room temperature for 4 hours to allow condensation. After washing with DMF, this condensation operation was repeated once. After washing the resin with DMF and dichloromethane, this was shaken with 20% piperidine in DMF for 20 minutes to deprotect the Fmoc group, and the resin was washed with DMF to synthesize a protected peptide 148 (SEQ ID NO. 152) represented by the following formula (148) in a state bound to the resin.


[Chemical Formula 183]



(148)

After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold diethyl ether was added to the filtrate, and crude peptide was obtained as precipitate. This crude peptide was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, A:B = 70:30] to obtain glycopeptide 149 (SEQ ID NO. 153) represented by the following formula (149) (5.8 mg, 1.1 μ mol).

[Chemical Formula 184]



(149)


ESI-MS: (m/z) calcd for $C_{241}H_{367}N_{49}O_{101}S_4$: $[M+4H]^+$ 1424.8, $[M+5H]^+$ 1140.0, found 1424.6, 1139.9.

[0241]

66-2 Glycosylation reaction of thiol

Glycopeptide 149 obtained in the method described in the above 66-1 (10.8 mg, 1.90 μ mol) and compound f (5.3 mg, 5.1 μ mol) were dissolved in 100 mM phosphate buffer (pH 7.4, 0.8 mL) containing 141 μ M TCEP, and reacted at room temperature for 24 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, A:B = 75:25 $\rightarrow$ 60:40, 30 minutes, linear gradient elution] to obtain glycopeptide 150 (SEQ ID NO. 154) represented by the following formula (150) (4.9 mg, 0.74 μ mol, yield 39%).

[Chemical Formula 185]



(150)


ESI-MS: (m/z) calcd for $C_{277}H_{426}N_{52}O_{127}S_4$: $[M+3H]^+$ 2216.0, $[M+4H]^+$ 1662.2, $[M+5H]^+$ 1330.0, found 2215.6, 1661.9, 1330.1.

[0242]

66-3 Deprotection of AcM group

To glycopeptide 150 obtained in the method described in the above 66-2 (4.9 mg, 0.74 μ mol) was added an aqueous solution (148 μ L) of silver(I) acetate (1.5 mg, 9.0 μ mol), and reacted at room temperature for 1.5 hours reaction. DTT (3.5 mg, 23 μ mol) dissolved in 200 mM phosphate buffer (pH 7.4, 145 μ L) and 100 mM ascorbic acid aqueous solution (37 μ L) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 74:26 $\rightarrow$ 64:36, 30 minutes, linear gradient elution] to obtain glycopeptide 151 (SEQ ID NO. 155) represented by the following formula (151) (3.7 mg, 0.57 μ mol, yield 77%).

[Chemical Formula 186]



(151)


ESI-MS: (m/z) calcd for $C_{271}H_{416}N_{50}O_{125}S_4$: $[M+3H]^{3+}$ 2168.6, $[M+4H]^{4+}$ 1626.7, $[M+5H]^{5+}$ 1301.5, found 2168.6, 1626.4, 1301.3.

[0243]

66-4 Formation of disulfide bond

Glycopeptide 151 obtained in the method described in the above 66-3 (3.7 mg, 0.54 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.4 mL), and reacted at room temperature for 31 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 60:40, 30 minutes, linear gradient elution] to obtain glycopeptide 152 (SEQ ID NO. 156) represented by the following formula (152) (2.9 mg, 0.45 μ mol, yield 78%).

[Chemical Formula 187]



(152)


ESI-MS: (m/z) calcd for $C_{271}H_{414}N_{50}O_{125}S_4$: $[M+3H]^{3+}$ 2167.9, $[M+4H]^{4+}$ 1626.2, $[M+5H]^{5+}$ 1301.1, found 2167.9, 1626.0, 1301.2.

[0244]

66-5 Deprotection of benzyl group

Glycopeptide 152 obtained in the method described in the above 66-4 (2.9 mg, 0.45 μ mol) was dissolved in 50 mM sodium hydroxide aqueous solution (13.6 mL), and reacted at 0°C for 1 hour. 200 mM acetic acid aqueous solution (3.4 mL) was added, and the mixed solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 60:40, 20 minutes, linear gradient elution] to obtain a fraction containing compound

(S1N(disialo)·N19C(diMan)-SRIF28) represented by the following formula 153 (SEQ ID NO. 157).
[Chemical Formula 188]



(153)

This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 4.6 x 250 mm, flow rate: 0.7 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 95:5 $\rightarrow$ 85:15, 2 minutes, then 85:15 $\rightarrow$ 65:35, 20 minutes, linear gradient elution] to obtain S1N(disialo)·N19C(diMan)-SRIF28 (1.6 mg, 0.25 μ mol, yield 57%).


ESI-MS: calcd for $C_{257}H_{402}N_{50}O_{125}S_4$: $[M+3H]^3+$ 2107.8, $[M+4H]^4+$ 1581.1, $[M+5H]^5+$ 1265.1, found 2107.9, 1580.9, 1265.1.

[0245]

Example 67 Synthesis of
C(disialo(aminoethylamide))·S1C(disialo)-SRIF28
67-1 Synthesis of peptide

2-chlorotrityl chloride resin (100 μ mol) was taken in a column for solid phase synthesis, and after washing with DMF and dichloromethane, a solution of Fmoc-Cys(Acm)-OH (49.7 mg, 120 μ mol) and DIPEA (104.5 μ L, 600 μ mol) in dichloromethane (3.0 mL) was added, and this was shaken for 1 hour. After washing with dichloromethane and DMF, the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF, in a peptide solid phase synthesis method with Fmoc strategy employing a PreludeTM peptide synthesizer, a protected peptide 154 (SEQ ID NO. 158) represented by the following formula (154) was synthesized in a state bound to the resin. The condensation reaction was performed in DMF using HCTU as the condensation agent.

[Chemical Formula 189]



(154)


A part of the resin (50 μ mol) was taken, and the Fmoc protecting group was removed by treating with 20% piperidine in DMF. After washing with DMF and dichloromethane, TFA:water:triisopropylsilane:ethanedithiol (= 90:2.5:5:2.5) was added, and this was shaken at room temperature for 3 hours. The resin was filtered off, cold

(155)

[0246]

Compound m (50.0 mg, 19.0 μ mol) was dissolved in a TFA-H₂O (95/5, v/v, 2.5 mL) solution, and this was shaken at room temperature. After 10 minutes, diethyl ether (15 mL) was added, and the deposited precipitate was centrifuged (10,000 x g 10 minutes). The precipitate was dissolved in water and lyophilized to obtain compound s represented by the following formula (s) (bromoacetamidated disialo sugar chain: 46.0 mg, 18.9 μ mol, yield 99%).

[Chemical Formula 191]



(S)


133

[0247]

67-3 glycosylation of thiol

Peptide 155 obtained in the method described in the above 67-1 (20.6 mg, 5.89 μmol) and compound s obtained in the method described in the above 67-2 (28.6 mg, 11.8 μmol) were dissolved in 33 mM phosphate buffer (pH 7.4, 1.8 mL), and reacted at room temperature for 30 minutes. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 74:26 $\rightarrow$ 64:36, 20 minutes, linear gradient elution] to obtain glycopeptide 156 (SEQ ID NO. 160) represented by the following formula (156) (17.1 mg, 2.93 μmol , yield 50%).

[Chemical Formula 192]



(156)


ESI-MS: (m/z) calcd for $\text{C}_{240}\text{H}_{383}\text{N}_{55}\text{O}_{101}\text{S}_6$: $[\text{M}+3\text{H}]^{3+}$ 1950.1, $[\text{M}+4\text{H}]^{4+}$ 1462.8, $[\text{M}+5\text{H}]^{5+}$ 1170.5, $[\text{M}+6\text{H}]^{6+}$ 975.6, found 1949.9, 1462.6, 1170.3, 975.4.

[0248]

67-4 Deprotection of StBu Group

To glycopeptide 156 obtained in the method described in the above 67-3 (17.1 mg, 2.93 μmol) was added DTT (52.9 mg, 343 μmol) dissolved in 0.1 M phosphate buffer (pH 7.4, 3.4 mL), and reacted at room temperature for 3 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μm), $\phi 20 \times 250$ mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 95:5 $\rightarrow$ 75:25, 20 minutes, linear gradient elution] to obtain glycopeptide 157 (SEQ ID NO. 161) represented by the following formula (157) (8.6 mg, 1.5 μmol , yield 51%).

[Chemical Formula 193]



(157)


ESI-MS: (m/z) calcd for $C_{236}H_{375}N_{55}O_{101}S_5$: $[M+3H]^{3+}$ 1920.7, $[M+4H]^{4+}$ 1440.8, $[M+5H]^{5+}$ 1152.8, $[M+6H]^{6+}$ 960.9, $[M+7H]^{7+}$ 823.7, found 1920.5, 1440.6, 1152.7, 960.6, 823.6.

[0249]

67-5 Glycosylation reaction of thiol

Peptide 157 obtained in the method described in the above 67-4 (6.2 mg, 1.1 μ mol) and compound a (3.8 mg, 1.6 μ mol) were dissolved in 0.36 M phosphate buffer (pH 7.4, 339 mL) containing 1.6 mM DTT, and reacted at room temperature for 2.5 hours. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 75:25, 30 minutes, linear gradient elution] to obtain glycopeptide 158 (SEQ ID NO. 162) represented by the following formula (158) (6.4 mg, 0.80 μ mol, yield 73%).

[Chemical Formula 194]



(158)


ESI-MS: (m/z) calcd for $C_{322}H_{514}N_{62}O_{163}S_5$: $[M+4H]^{4+}$ 2006.5, $[M+5H]^{5+}$ 1605.4, $[M+6H]^{6+}$ 1338.0, found 2006.6, 1605.3, 1338.0.

[0250]

67-6 Deprotection of AcM group

To glycopeptide 158 obtained in the method described in the above 67-5 (8.2 mg, 1.0 μ mol) was added an aqueous solution (225 μ L) of silver(I) acetate (2.1 mg, 13 μ mol), and reacted at room temperature for 1 hour. DTT (4.8 mg, 31 μ mol) dissolved in 100 mM phosphate buffer (pH 7.4, 204 μ L) and 100 mM ascorbic acid aqueous solution (51 μ L) were added, and this was promptly filtered with a filter. The filtrate was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 75:25, 30 minutes, linear gradient elution] to obtain glycopeptide 159 (SEQ ID NO. 163) represented by the following formula (159) (5.5 mg, 0.70 μ mol, yield 70%).

[Chemical Formula 195]




(159)

ESI-MS: (m/z) calcd for $C_{316}H_{504}N_{60}O_{161}S_5$: $[M+4H]^{4+}$ 1971.0, $[M+5H]^{5+}$ 1577.0, $[M+6H]^{6+}$ 1314.3, found 1970.6, 1576.8, 1314.2.

[0251]

67-7 Formation of disulfide bond

Glycopeptide 159 obtained in the method described in the above 67-6 (5.4 mg, 0.69 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 1.7 mL), and reacted overnight at room temperature. The reaction solution was purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous TFA, B: 0.09% TFA/10% water/90% acetonitrile, gradient A:B = 75:25 $\rightarrow$ 65:35, 30 minutes, linear gradient elution] to obtain a fraction containing compound (C(disialo(aminoethylamide))/S1C(disialo)-SRIF28) represented by the following formula 160 (SEQ ID NO. 164). [Chemical Formula 196]



(160)


This fraction was further purified with HPLC [column: SHISEIDO CAPCELL PAK C18 UG-120 (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 0.1% aqueous AcOH, B: 0.09% AcOH/10% water/90% acetonitrile, gradient A:B = 90:10 $\rightarrow$ 75:25, 20 minutes, linear gradient elution] to obtain C(disialo(aminoethylamide))/S1C(disialo)-SRIF28 (3.1 mg, 0.40 μ mol, yield 58%).

ESI-MS: calcd for $C_{316}H_{502}N_{60}O_{161}S_5$: $[M+4H]^{4+}$ 1970.5, $[M+5H]^{5+}$ 1576.6, $[M+6H]^{6+}$ 1314.0, $[M+7H]^{7+}$ 1126.4, $[M+8H]^{8+}$ 985.8, $[M+9H]^{9+}$ 876.3, found 1970.3, 1576.4, 1313.9, 1126.4, 985.5, 876.1.

[0252]

Example 68 Synthesis of S1-4C(disialo)-SRIF28

68-1 Synthesis of peptide



Cys-Cys-Cys-Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-
-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

(164)


ESI-MS: (m/z) calcd for $C_{490}H_{780}N_{72}O_{289}S_7$: $[M+5H]^{5+}$ 2506.6,
 $[M+6H]^{6+}$ 2089.0, $[M+7H]^{7+}$ 1790.7, found 2506.5, 2088.8,
1790.4.

[0255]

68-4 Formation of disulfide bond

Glycopeptide 164 obtained in the method described in the above 68-3 (20.6 mg, 1.64 μ mol) was dissolved in 100 mM Tris-HCl buffer (pH 8.0)-DMSO (1/1, v/v, 2.1 mL), and reacted at room temperature for 2 days. Then, the reaction solution was purified with HPLC [column: SHISEIDO Proteonavi (5 μ m), ϕ 20 x 250 mm, flow rate: 7.0 mL/min, eluent A: 10 mM ammonium acetate aqueous solution, B: 10 mM ammonium acetate-acetonitrile (1/9, v/v), gradient A:B = 75:25 $\rightarrow$ 72:28, 15 minutes, linear gradient elution] to obtain S1-4C(disialo)-SRIF28 (SEQ ID NO. 169) represented by the following formula (165) (11.6 mg, 0.93 μ mol, yield 57%).

[Chemical Formula 201]



Cys-Cys-Cys-Cys-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys-
-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys

(165)

ESI-MS: (m/z) calcd for $C_{490}H_{778}N_{72}O_{289}S_7$: $[M+5H]^{5+}$ 2506.2,
 $[M+6H]^{6+}$ 2088.7, $[M+7H]^{7+}$ 1790.5, found 2506.1, 2088.6,
1790.4.

[0256]

Tables 1-1 to 1-7 show the MS spectrum data (ESI-MS) of the glycosylated SRIF peptides obtained in the methods described in Examples 1 - 68. The molecular mass was obtained by performing deconvolution of polyvalent protein mass spectrometry with MassLynx version 4.1 (from Waters).

[0257]

[Table 1]

Table 1-1

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
1	S1C(disialo)-SRIF28	5427.7	$[M+3H]^{3+}$	1810.0	5427.0
			$[M+4H]^{4+}$	1357.5	
			$[M+5H]^{5+}$	1086.4	
2	N5C(disialo)-SRIF28	5400.6	$[M+3H]^{3+}$	1801.0	5400.1
			$[M+4H]^{4+}$	1350.8	
			$[M+5H]^{5+}$	1081.2	
3	A9C(disialo)-SRIF28	5443.7	$[M+3H]^{3+}$	1815.4	5443.1
			$[M+4H]^{4+}$	1361.8	
			$[M+5H]^{5+}$	1089.6	
4	E12C(disialo)-SRIF28	5385.6	$[M+3H]^{3+}$	1796.0	5384.0
			$[M+4H]^{4+}$	1347.0	
			$[M+5H]^{5+}$	1078.0	
5	R13C(disialo)-SRIF28	5358.6	$[M+3H]^{3+}$	1787.0	5358.0
			$[M+4H]^{4+}$	1340.5	
			$[M+5H]^{5+}$	1072.6	
6	K14C(disialo)-SRIF28	5386.6	$[M+3H]^{3+}$	1796.4	5386.0
			$[M+4H]^{4+}$	1347.5	
			$[M+5H]^{5+}$	1078.2	
7	A15C(disialo)-SRIF28	5443.7	$[M+3H]^{3+}$	1815.4	5443.1
			$[M+4H]^{4+}$	1361.8	
			$[M+5H]^{5+}$	1089.6	
8	G16C(disialo)-SRIF28	5457.7	$[M+3H]^{3+}$	1820.0	5457.1
			$[M+4H]^{4+}$	1365.3	
			$[M+5H]^{5+}$	1092.2	
9	K18C(disialo)-SRIF28	5386.6	$[M+3H]^{3+}$	1796.4	5386.1
			$[M+4H]^{4+}$	1347.5	
			$[M+5H]^{5+}$	1078.2	
			$[M+6H]^{6+}$	898.7	
			$[M+7H]^{7+}$	770.5	

[Table 2]

Table 1 - 2

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
10	N19C(disialo)-SRIF28	5400.6	$[M+3H]^{3+}$	1801.0	5400.1
			$[M+4H]^{4+}$	1351.0	
			$[M+5H]^{5+}$	1080.8	
11	F21C(disialo)-SRIF28	5367.6	$[M+3H]^{3+}$	1790.0	5366.9
			$[M+4H]^{4+}$	1342.7	
			$[M+5H]^{5+}$	1074.4	
12	T26C(disialo)-SRIF28	5413.6	$[M+3H]^{3+}$	1354.3	5412.1
			$[M+4H]^{4+}$	1083.4	
			$[M+5H]^{5+}$	903.0	
			$[M+6H]^{6+}$	774.3	
13	29C(disialo)-SRIF28	5514.7	$[M+3H]^{3+}$	1839.1	5514.1
			$[M+4H]^{4+}$	1379.5	
			$[M+5H]^{5+}$	1103.8	
14	30C(disialo)-SRIF28	5615.8	$[M+3H]^{3+}$	1872.7	5615.1
			$[M+4H]^{4+}$	1404.8	
			$[M+5H]^{5+}$	1124.0	
15	S1C(disialo)-D-Trp22 -SRIF28	5427.7	$[M+3H]^{3+}$	1810.1	5427.2
			$[M+4H]^{4+}$	1357.8	
			$[M+5H]^{5+}$	1086.4	
			$[M+6H]^{6+}$	905.5	
16	A9C(disialo)-D-Trp22 -SRIF28	5443.7	$[M+3H]^{3+}$	1815.4	5442.9
			$[M+4H]^{4+}$	1361.8	
			$[M+5H]^{5+}$	1089.6	
17	C(disialo)-SRIF14	4004.1	$[M+2H]^{2+}$	2002.9	4003.3
			$[M+3H]^{3+}$	1335.4	
			$[M+4H]^{4+}$	1001.8	
18	C(disialo)-R-K-SRIF14	4288.4	$[M+3H]^{3+}$	1430.2	4287.5
			$[M+4H]^{4+}$	1072.9	
19	C(disialo)-C12linker-SRIF14	4201.4	$[M+3H]^{3+}$	1401.2	4200.5
			$[M+4H]^{4+}$	1051.2	
			$[M+5H]^{5+}$	841.1	

[Table 3]

Table 1 - 3

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
20	S1-2C(disialo)-SRIF28	7793.8	$[M+4H]^{4+}$	1949.4	7793.4
			$[M+5H]^{5+}$	1559.7	
			$[M+6H]^{6+}$	1299.9	
21	S1C(disialo) • N5C(disialo)-SRIF28	7679.7	$[M+3H]^{3+}$	1920.4	7678.7
			$[M+4H]^{4+}$	1536.8	
			$[M+5H]^{5+}$	1280.9	
22	S1C(disialo) • R13C(disialo)-SRIF28	7637.7	$[M+3H]^{3+}$	1910.2	7636.6
			$[M+4H]^{4+}$	1528.3	
			$[M+5H]^{5+}$	1273.8	
23	N5C(disialo) • A9C(disialo)-SRIF28	7695.7	$[M+5H]^{5+}$	1540.1	7695.2
			$[M+6H]^{6+}$	1283.6	
			$[M+7H]^{7+}$	1100.3	
			$[M+8H]^{8+}$	962.9	
24	S1-3C(disialo)-SRIF28	10160.0	$[M+3H]^{3+}$	3387.6	10158.2
			$[M+4H]^{4+}$	2540.9	
			$[M+5H]^{5+}$	2032.7	
			$[M+6H]^{6+}$	1694.2	
			$[M+7H]^{7+}$	1452.3	
25	S1C(disialo) • N5C(disialo) • A9C(disialo)-SRIF28	9974.8	$[M+5H]^{5+}$	1995.9	9974.3
			$[M+6H]^{6+}$	1663.2	
26	S1C(monosialo)-SRIF28	5136.4	$[M+3H]^{3+}$	1713.0	5134.9
			$[M+4H]^{4+}$	1284.7	
			$[M+5H]^{5+}$	1028.2	
			$[M+6H]^{6+}$	856.8	

[Table 4]

Table 1 — 4

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
27	S1C(asialo)-SRIF28	4845.1	$[M+3H]^{3+}$	2615.7	4843.9
			$[M+4H]^{4+}$	1212.0	
			$[M+5H]^{5+}$	969.8	
28	S1-2C(asialo)-SRIF28	6628.8	$[M+4H]^{4+}$	1658.1	6627.2
			$[M+5H]^{5+}$	1326.7	
			$[M+6H]^{6+}$	1105.6	
			$[M+7H]^{7+}$	947.8	
			$[M+8H]^{8+}$	829.4	
29	S1-3C(asialo)-SRIF28	8412.5	$[M+4H]^{4+}$	2103.7	8411.7
			$[M+5H]^{5+}$	1683.3	
			$[M+6H]^{6+}$	1403.0	
30	N5N(disialo)-SRIF28	5354.5	$[M+3H]^{3+}$	1785.7	5354.0
			$[M+4H]^{4+}$	1339.5	
			$[M+5H]^{5+}$	1071.8	
31	S1N(disialo)-SRIF28	5381.6	$[M+3H]^{3+}$	1794.7	5380.9
			$[M+4H]^{4+}$	1346.2	
			$[M+5H]^{5+}$	1077.2	
32	S1C(disialo) • N19C(GlcNAc)-SRIF28	5676.9	$[M+3H]^{3+}$	1893.5	5676.4
			$[M+4H]^{4+}$	1420.1	
			$[M+5H]^{5+}$	1136.3	
33	S1C(disialo) • N19C(diMan)-SRIF28	6366.6	$[M+3H]^{3+}$	2122.9	6366.7
			$[M+4H]^{4+}$	1592.5	
			$[M+5H]^{5+}$	1274.4	
34	S1-5C(disialo)-SRIF28	14892.4	$[M+4H]^{4+}$	3723.7	14891.3
			$[M+5H]^{5+}$	2979.1	
			$[M+6H]^{6+}$	2482.9	
			$[M+7H]^{7+}$	2128.2	
			$[M+8H]^{8+}$	1862.4	
35	S1-10C(disialo)-SRIF28	26723.2	$[M+7H]^{7+}$	3818.3	26721.8
			$[M+8H]^{8+}$	3341.0	
			$[M+9H]^{9+}$	2970.1	
			$[M+10H]^{10+}$	2673.1	

[Table 5]

Table 1 —5

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
36	C(disialo)-SRIF28	5514.7	[M+4H] ⁴⁺	1379.7	5514.6
			[M+5H] ⁵⁺	1103.9	
			[M+6H] ⁶⁺	920.1	
			[M+7H] ⁷⁺	788.8	
37	R11C(disialo)-SRIF28	5358.5	[M+3H] ³⁺	1787.4	5358.6
			[M+4H] ⁴⁺	1340.7	
			[M+5H] ⁵⁺	1072.7	
38	F20C(disialo)-SRIF28	5367.6	[M+3H] ³⁺	1790.1	5367.1
			[M+4H] ⁴⁺	1342.8	
			[M+5H] ⁵⁺	1074.4	
39	T24C(disialo)-SRIF28	5413.6	[M+3H] ³⁺	1805.7	5412.6
			[M+4H] ⁴⁺	1354.2	
			[M+5H] ⁵⁺	1083.7	
40	F25C(disialo)-SRIF28	5367.6	[M+3H] ³⁺	1790.1	5367.1
			[M+4H] ⁴⁺	1342.8	
			[M+5H] ⁵⁺	1074.4	
			[M+6H] ⁶⁺	895.4	
41	S27C(disialo)-SRIF28	5427.7	[M+3H] ³⁺	1810.1	5427.4
			[M+4H] ⁴⁺	1357.8	
42	C(disialo)-K-SRIF14	4131.2	[M+2H] ²⁺	2066.9	4131.7
			[M+3H] ³⁺	1378.2	
			[M+4H] ⁴⁺	1033.9	
43	S1C(disialo)-F25Y-SRIF28	5443.7	[M+3H] ³⁺	1815.4	5443.1
			[M+4H] ⁴⁺	1361.6	
			[M+5H] ⁵⁺	1089.6	
44	S1C(disialo)-SRIF28-amide	5426.7	[M+3H] ³⁺	1809.8	5426.2
			[M+4H] ⁴⁺	1357.6	
			[M+5H] ⁵⁺	1086.2	
			[M+6H] ⁶⁺	905.2	
45	C(disialo)-PEGlinker-SRIF14	4207.3	[M+3H] ³⁺	1403.2	4206.6
			[M+4H] ⁴⁺	1052.7	
			[M+5H] ⁵⁺	842.3	
46	Biotin-S1C(disialo)-SRIF28	5654.0	[M+3H] ³⁺	1885.4	5653.2
			[M+4H] ⁴⁺	1414.3	
			[M+5H] ⁵⁺	1131.7	
			[M+6H] ⁶⁺	943.2	
47	Biotin-PEGlinker -S1C(disialo)-SRIF28	5857.2	[M+3H] ³⁺	1953.2	5856.4
			[M+4H] ⁴⁺	1465.1	
			[M+5H] ⁵⁺	1172.1	
			[M+6H] ⁶⁺	977.1	
			[M+7H] ⁷⁺	837.6	

[Table 6]

Table 1 — 6

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
46	Azido-S1C(disialo)-SRIF28	5552.8	[M+3H] ³⁺	1851.8	5552.3
			[M+4H] ⁴⁺	1389.1	
			[M+5H] ⁵⁺	1111.7	
49	S1C(disialo)• E12C(disialo) -SRIF28	7664.7	[M+4H] ⁴⁺	1917.1	7664.1
			[M+5H] ⁵⁺	1533.8	
			[M+6H] ⁶⁺	1278.4	
50	2C(disialo)-R-K-SRIF14	6654.6	[M+3H] ³⁺	2219.2	6653.4
			[M+4H] ⁴⁺	1664.4	
			[M+5H] ⁵⁺	1331.9	
			[M+6H] ⁶⁺	1109.9	
51	3C(disialo)-R-K-SRIF14	9020.8	[M+4H] ⁴⁺	2256.1	9020.2
			[M+5H] ⁵⁺	1805.1	
			[M+6H] ⁶⁺	1504.4	
52	S1C(diGlcNAc)-SRIF28	4520.9	[M+3H] ³⁺	1507.7	4519.9
			[M+4H] ⁴⁺	1131.0	
			[M+5H] ⁵⁺	905.0	
53	S1C(diMan)-SRIF28	4114.5	[M+3H] ³⁺	1372.3	4113.8
			[M+4H] ⁴⁺	1029.5	
			[M+5H] ⁵⁺	823.8	
54	N19C(diMan)-SRIF28	4087.5	[M+3H] ³⁺	1363.3	4086.8
			[M+4H] ⁴⁺	1023.0	
55	S1C(GlcNAc)-SRIF28	3424.9	[M+2H] ²⁺	1713.3	3424.5
			[M+3H] ³⁺	1142.5	
			[M+4H] ⁴⁺	856.9	
			[M+5H] ⁵⁺	686.1	
56	N19C(GlcNAc)-SRIF28	3397.8	[M+2H] ²⁺	1699.8	3397.5
			[M+3H] ³⁺	1133.5	
			[M+4H] ⁴⁺	850.4	
57	S1C(trisialo)-SRIF28	6084.2	[M+3H] ³⁺	2028.8	6083.4
			[M+4H] ⁴⁺	1521.9	
			[M+5H] ⁵⁺	1217.7	
			[M+6H] ⁶⁺	1014.9	
58	S1C(tetrasialo)-SRIF28	6740.8	[M+3H] ³⁺	2247.7	6739.6
			[M+4H] ⁴⁺	1685.9	
			[M+5H] ⁵⁺	1349.1	
			[M+6H] ⁶⁺	1124.3	
			[M+7H] ⁷⁺	963.8	
59	S1C(disialo(aminoethylamide)) -SRIF28	5511.8	[M+3H] ³⁺	1838.0	5511.1
			[M+4H] ⁴⁺	1378.5	
			[M+5H] ⁵⁺	1103.2	
			[M+6H] ⁶⁺	919.5	
			[M+7H] ⁷⁺	788.2	

[Table 7]

Table 1 — 7

Example	Compound	Calculated molecular weight	Observed ions (m/z)		Observed molecular weight
60	S1C(disialo(amide))-SRIF28	5425.7	[M+3H] ³⁺	1809.3	5423.9
			[M+4H] ⁴⁺	1357.0	
			[M+5H] ⁵⁺	1085.8	
			[M+6H] ⁶⁺	905.2	
61	S1C(disialo(Bn))-SRIF28	5607.9	[M+3H] ³⁺	1870.0	5605.9
			[M+4H] ⁴⁺	1402.5	
			[M+5H] ⁵⁺	1122.2	
			[M+6H] ⁶⁺	935.5	
62	S1C(disialo(hexadecylamide))-SRIF28	5874.5	[M+3H] ³⁺	1958.9	5873.6
			[M+4H] ⁴⁺	1469.4	
			[M+5H] ⁵⁺	1175.7	
			[M+6H] ⁶⁺	979.9	
63	S1-2C(disialo(amide))-SRIF28	7789.9	[M+4H] ⁴⁺	1948.2	7789.9
			[M+5H] ⁵⁺	1559.0	
			[M+6H] ⁶⁺	1299.2	
64	S1-2C(disialo(Bn))-SRIF28	8154.3	[M+4H] ⁴⁺	2039.3	8153.0
			[M+5H] ⁵⁺	1631.6	
			[M+6H] ⁶⁺	1360.2	
65	S1C(Asn(disialo))-SRIF28	5542.7	[M+3H] ³⁺	1848.4	5542.2
			[M+4H] ⁴⁺	1386.6	
			[M+5H] ⁵⁺	1109.4	
			[M+6H] ⁶⁺	924.9	
66	S1N(disialo)·N19C(diMan)-SRIF28	6320.5	[M+3H] ³⁺	2107.9	6319.5
			[M+4H] ⁴⁺	1580.9	
			[M+5H] ⁵⁺	1265.1	
67	C(disialo(aminoethylamide))·S1C(disialo)-SRIF28	7878.0	[M+4H] ⁴⁺	1970.3	7878.1
			[M+5H] ⁵⁺	1576.4	
			[M+6H] ⁶⁺	1313.9	
			[M+7H] ⁷⁺	1126.4	
			[M+8H] ⁸⁺	985.5	
			[M+9H] ⁹⁺	876.1	
68	S1-4C(disialo)-SRIF28	12526.2	[M+5H] ⁵⁺	2506.1	12525.5
			[M+6H] ⁶⁺	2088.6	
			[M+7H] ⁷⁺	1790.4	

[0258]

Example 69-1 Calculation of receptor binding affinity
Competitive binding assay was performed with the method below to calculate receptor binding affinity.

Reagents employed in the competitive binding assay and their proper chemical names are as follows: HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) and BSA (bovine serum albumin).

Competitive binding assay was consigned to Ricerca Biosciences, and experiment and data analysis was performed.

The receptor subtypes and receptor membrane samples employed are shown in Table 2. Common in each binding assay, [^{125} I] Tyr¹¹-Somatostatin 14 (Tyr¹¹-SRIF14) was used as the labeled ligand, Somatostatin-14 (SRIF14) was used as the unlabeled ligand, and 25 mM HEPES containing 5 mM MgCl₂, 1 mM CaCl₂, and 0.1% BSA, pH 7.4 was used as the buffer. The test substance was used at a concentration of 0.01 nM, 0.1 nM, 1 nM, 10 nM, 100 nM, or 1000 nM, and mixed with the membrane sample into a reaction solution. Moreover, the concentrations of the labeled and unlabeled ligands added to the reaction solution are shown in Table 2. The incubation condition of the reaction solution was at 25°C for 4 hours for SSTR2 and at 25°C for 2 hours for SSTR1, SSTR3, SSTR4, and SSTR5. For each experiment run, SRIF14 was used as the positive control. For data analysis, 50% inhibitory concentration (IC₅₀ value) was determined using MathIQ™ (ID Business Solutions, UK) in a nonlinear least squares method based on the numerical data of the binding inhibition rate. The binding inhibition constant (K_i value) was calculated by the method of Cheng, Y et al. (Biochem Pharmacol, 22, 3099-3108, 1973).

[Table 8]

Table 2

	SSTR1	SSTR2	SSTR3	SSTR4	SSTR5
Labeled ligand	0.1nM	0.03nM	0.1nM	0.1nM	0.1nM
Unlabeled ligand	1μM	1μM	1μM	1μM	1μM
Source	Human recombinant CHO-K1 cells	Human recombinant CHO-K1 cells	Human recombinant CHO-K1 cells	Human recombinant Chem-1 cells	Human recombinant Chem-1 cells

[0259]

Compounds subjected to binding assay and the results of the binding experiment are shown in Table 3A. Moreover, octreotide, SRIF14 and SRIF28 were similarly evaluated as control compounds. Note that for octreotide, IC₅₀ values could not be calculated for SSTR1 and SSTR4 since the maximum concentration was 100 nM, and is thus shown as >100 nM.

Figures 1A and 1B show examples of the structures of glycosylated peptides (glycosylated forms) corresponding to the compound names in Table 3A.

[0260]

[Table 9]

Table 3A

Example		SSTR1		SSTR2		SSTR3		SSTR4		SSTR5	
		IC ₅₀ (nM)	Ki(nM)	IC ₅₀ (nM)	Ki(nM)	IC ₅₀ (nM)	Ki(nM)	IC ₅₀ (nM)	Ki(nM)	IC ₅₀ (nM)	Ki(nM)
1	S1C(disialo)-SRIF28	2.19	1.81	0.109	0.0576	0.480	0.164	2.30	2.07	0.442	0.250
2	N5C(disialo)-SRIF28	1.10	0.912	0.0945	0.0502	0.278	0.095	1.30	1.17	0.458	0.259
3	A9C(disialo)-SRIF28	2.08	1.72	0.0741	0.0394	0.593	0.203	1.72	1.54	0.471	0.266
4	E12C(disialo)-SRIF28	1.70	1.40	0.112	0.0596	0.362	0.124	2.04	1.84	0.376	0.212
5	R13C(disialo)-SRIF28	16.4	13.6	0.310	0.165	2.87	0.981	6.43	5.79	3.28	1.85
6	K14C(disialo)-SRIF28	13.8	11.4	0.481	0.256	2.59	0.887	8.02	7.22	4.01	2.27
10	N19C(disialo)-SRIF28	16.1	13.3	1.21	0.645	1.47	0.503	12.2	11.0	3.69	2.09
13	29C(disialo)-SRIF28	4.43	3.67	0.738	0.392	7.00	2.39	6.11	5.50	3.73	2.11
14	30C(disialo)-SRIF28	4.21	3.48	0.517	0.275	6.91	2.36	2.53	2.28	1.86	1.05
15	S1C(disialo)-D-Trp-SRIF28	2.23	1.85	0.0324	0.0172	0.834	0.285	5.85	5.26	0.565	0.319
16	A9C(disialo)-D-Trp-SRIF28	6.92	5.73	0.0396	0.021	0.256	0.0876	4.87	4.38	0.194	0.110
17	C(disialo)-SRIF14	32.3	26.7	0.336	0.179	4.62	1.58	8.24	7.41	5.70	3.22
18	C(disialo)-R-K-SRIF14	3.09	2.56	0.104	0.0551	0.231	0.0791	1.06	0.951	0.187	0.106
19	C(disialo)-C12linker-SRIF14	15.5	12.8	0.502	0.267	3.12	1.07	2.92	2.62	2.29	1.29
20	S1-2C(disialo)-SRIF28	18.8	15.5	0.338	0.179	1.28	0.439	5.61	5.05	0.607	0.343
21	S1C(disialo) · N5C(disialo)-SRIF28	6.89	5.70	0.224	0.119	1.58	0.539	4.00	3.60	0.728	0.411
22	S1C(disialo) · R13C(disialo)-SRIF28	177	147	6.28	3.33	11.0	3.77	45.0	40.5	14.7	8.33
23	N5C(disialo) · A9C(disialo)-SRIF28	30.0	24.9	0.793	0.421	4.09	1.40	14.2	12.8	1.81	1.02
24	S1-3C(disialo)-SRIF28	20.4	16.9	0.421	0.224	3.30	1.13	7.69	6.92	0.927	0.524
25	S1C(disialo) · N5C(disialo) · A9C(disialo)-SRIF28	69.2	57.3	0.985	0.523	7.93	2.71	25.2	22.7	4.14	2.34
26	S1C(monosialo)-SRIF28	1.43	1.19	0.0465	0.0247	0.322	0.11	1.14	1.03	0.372	0.210
27	S1C(asialo)-SRIF28	0.851	0.704	0.0224	0.0119	0.155	0.0531	0.627	0.564	0.460	0.260
28	S1-2C(asialo)-SRIF28	0.86	0.712	0.0632	0.0336	0.177	0.0604	0.78	0.702	0.19	0.107
29	S1-3C(asialo)-SRIF28	4.21	3.49	0.0978	0.0519	0.581	0.199	1.44	1.29	0.369	0.209
30	N5N(disialo)-SRIF28	3.02	2.50	0.0685	0.0364	0.579	0.198	2.86	2.57	0.465	0.263
31	S1N(disialo)-SRIF28	2.68	2.21	0.0754	0.0401	0.314	0.107	1.07	0.96	0.272	0.154
-	SRIF14	0.438	0.362	0.0142	0.00755	0.131	0.045	0.303	0.273	0.578	0.326
-	Octreotide	>100	-	0.0628	0.0334	6.41	2.19	>100	-	9.39	5.31
-	SRIF28	0.353	0.292	0.0271	0.0144	0.0868	0.0297	0.294	0.265	0.423	0.239

[0261]

The control octreotide bound to each of receptors SSTR2, SSTR3, and SSTR5, and SRIF14 and SRIF28 bound to all SSTRs. As shown in Table 3A, the compounds according to

the present invention potentially bound to all SSTRs. The binding affinity of the positive control SRIF14 against SSTR1 in terms of K_i value was 0.362 nM, and where compounds having one sugar chain was 0.704 - 26.7 nM, the compounds of the present invention having two sugar chains (the compounds of Examples 20 - 23 and 28) were 0.712 - 147 nM, showing similarly sufficient receptor binding affinity. Moreover, the compounds of the present invention having three sugar chains (the compounds of Examples 24, 25, and 29) also had sufficient receptor binding affinity, showing 3.49 nM, 16.9 nM, and 57.3 nM. Since bioavailability (BA) will be considerably increased due to the extension of its half-life in blood, they can work effectively on receptors *in vivo* even if the K_i value of the binding affinity is somewhat high. Similarly, where the binding affinity of SRIF14 against SSTR2, SSTR3, and SSTR4 in terms of K_i value were each 0.00755 nM, 0.0450 nM, and 0.273 nM, the compounds of the present invention having two or more sugar chains were each 0.0336 - 3.33 nM, 0.0604 - 3.77 nM, and 0.702 - 40.5 nM, all showing sufficient receptor binding affinity. Moreover, where the binding affinity of SRIF14 against SSTR5 was 0.326 nM, the compounds of the present invention having two or more sugar chains showed a sufficient receptor binding affinity of up to 8.33 nM.

In this way, it was found that Examples 20 - 23 and 28 are diglycosylated modified forms having affinity towards all receptors SSTR1 - SSTR5, and Examples 24, 25, and 29 are triglycosylated modified forms having affinity towards all receptors SSTR1 - SSTR5.

[0262]

Example 69-2 Calculation of receptor binding affinity
- 2

Competitive binding assay was performed with each compound shown in Table 3B with the method described in Example 69-1, and receptor binding affinity was calculated. Moreover, SRIF14 and SRIF28 were similarly evaluated as control compounds. The results of the binding assay are shown in Table 3B.

Figures 1C and 1D show examples of the structures of glycosylated peptides corresponding to the compound names in Table 3B.

[Table 10]

Table 3B

Exptl #	SSTR1			SSTR2			SSTR3			SSTR4			SSTR5		
	K _d (nM)	K _i (nM)	IC ₅₀ (nM)	K _d (nM)	K _i (nM)	IC ₅₀ (nM)	K _d (nM)	K _i (nM)	IC ₅₀ (nM)	K _d (nM)	K _i (nM)	IC ₅₀ (nM)	K _d (nM)	K _i (nM)	IC ₅₀ (nM)
1	STC (dialo)-SRIF28	7.71	6.45	0.543	0.249	1.08	0.676	1.08	0.676	6.39	1.61	1.61	0.911	0.911	1.61
7	A10C (a.salo)-SRIF28	11.4	9.43	0.938	0.408	3.40	1.19	1.19	1.19	8.18	7.36	3.84	2.17	2.17	3.84
8	G16C (dialo)-SRIF28	17.4	14.4	1.22	0.846	3.12	1.07	1.07	1.07	12.2	11.0	5.67	3.21	3.21	5.67
9	K18C (dialo)-SRIF28	45.4	37.6	12.2	8.51	14.7	5.03	5.03	5.03	62.7	54.5	160	95.0	95.0	160
12	T26C (dialo)-SRIF28	21.6	17.9	3.10	1.80	21.7	7.44	7.44	7.44	26.8	24.1	460	230	230	460
26	STC (a.salo)-SRIF28	5.38	4.46	0.213	0.112	1.18	0.405	2.35	2.12	2.35	2.12	0.816	0.400	0.400	0.816
27	STC (a.salo)-SRIF28	3.97	3.29	0.200	0.105	1.02	0.349	2.38	2.68	2.38	2.68	1.04	0.588	0.588	1.04
32	STC (dialo)-N19C (a.salo)-SRIF28	<10	<8.27	2.20	1.17	3.05	1.04	8.55	7.69	8.55	7.69	2.37	1.34	1.34	2.37
33	STC (dialo)-N19C (dialo)-SRIF28	44.0	37.0	1.73	1.08	3.63	1.24	25.0	23.0	25.0	23.0	5.69	3.22	3.22	5.69
34	STC (dialo)-N19C (dialo)-SRIF28	110	91.9	12.7	6.58	21.3	7.29	62.3	56.4	62.3	56.4	7.23	4.09	4.09	7.23
35	STC (dialo)-SRIF28	117	94.5	10.1	5.35	40.4	13.8	191	172	191	172	34.8	19.7	19.7	34.8
36	STC (dialo)-SRIF28	5.67	4.61	0.478	0.255	2.55	0.870	8.65	7.69	8.65	7.69	2.62	1.48	1.48	2.62
37	STC (dialo)-SRIF28	22.0	18.2	1.38	0.735	6.34	2.17	20.2	18.1	20.2	18.1	3.65	2.07	2.07	3.65
40	F25C (dialo)-SRIF28	>1000	>828	>1000	>531	>1000	>342	130	110	130	110	58.0	33.0	33.0	58.0
41	S27C (dialo)-SRIF28	28.4	23.5	0.326	0.170	6.77	2.31	33.4	30.1	33.4	30.1	4.94	2.80	2.80	4.94
42	STC (dialo)-K-SRIF14	20.7	17.2	1.97	1.05	3.55	1.21	11.0	9.77	11.0	9.77	3.52	1.99	1.99	3.52
43	STC (dialo)-F25C-SRIF28	1.77	1.48	0.310	0.160	0.480	0.160	1.55	1.39	1.55	1.39	1.09	0.620	0.620	1.09
44	STC (dialo)-SRIF28	4.87	4.03	1.32	0.700	5.91	2.02	8.88	6.19	8.88	6.19	3.75	2.12	2.12	3.75
45	STC (dialo)-PE-Galactose-SRIF14	38.0	30.0	4.67	2.48	8.71	2.98	21.0	19.0	21.0	19.0	>10	>5.65	>5.65	>10
46	Biotin-STC (dialo)-SRIF28	11.0	9.03	1.68	0.880	2.92	1.00	9.28	8.35	9.28	8.35	2.42	1.37	1.37	2.42
47	Biotin-PEG-Galactose-STC (dialo)-SRIF28	9.41	7.79	1.96	1.04	3.41	1.17	8.31	6.38	8.31	6.38	2.00	1.13	1.13	2.00
48	STC (dialo)-SRIF28	8.82	7.30	1.29	0.680	3.17	1.09	5.34	4.81	5.34	4.81	0.840	0.470	0.470	0.840
49	STC (dialo)-E12C (dialo)-SRIF28	22.0	19.0	2.70	1.44	4.62	1.58	20.0	18.0	20.0	18.0	2.11	1.19	1.19	2.11
50	ZC (dialo)-RK-SRIF14	48.0	40.0	5.35	2.84	9.74	3.33	49.0	44.0	49.0	44.0	6.16	3.48	3.48	6.16
51	STC (dialo)-RK-SRIF14	270	230	32.0	17.0	40.0	14.0	>100	>90.0	>100	>90.0	41.0	23.0	23.0	41.0
52	STC (dialo)-NAC-SRIF28	5.04	4.17	0.190	0.100	2.02	0.690	3.91	3.02	3.91	3.02	1.46	0.820	0.820	1.46
53	STC (dialo)-SRIF28	5.06	4.18	0.190	0.100	2.02	0.690	3.91	3.02	3.91	3.02	1.46	0.820	0.820	1.46
54	N19C (a.salo)-SRIF28	6.58	5.43	1.17	0.620	1.55	0.510	8.02	7.22	8.02	7.22	2.94	1.68	1.68	2.94
55	STC (dialo)-SRIF28	5.91	4.89	0.225	0.120	1.65	0.545	2.89	2.60	2.89	2.60	1.15	0.650	0.650	1.15
56	N19C (dialo)-SRIF28	2.60	2.15	0.940	0.500	<1	<0.342	2.38	2.15	2.38	2.15	1.80	1.07	1.07	1.80
57	STC (a.salo)-SRIF28	17.0	14.0	0.780	0.410	5.59	1.91	13.0	14.0	13.0	14.0	1.96	1.10	1.10	1.96
58	STC (a.salo)-SRIF28	17.3	14.2	1.12	0.607	5.63	1.93	12.8	12.8	12.8	12.8	2.12	1.20	1.20	2.12
59	STC (dialo)-N19C (a.salo)-SRIF28	2.35	1.94	0.101	0.0536	0.766	0.244	2.24	2.01	2.24	2.01	2.08	1.18	1.18	2.08
60	STC (dialo)-N19C (a.salo)-SRIF28	5.09	4.21	0.207	0.110	1.53	0.522	6.12	5.50	6.12	5.50	2.60	1.47	1.47	2.60
61	STC (dialo)-SRIF28	6.20	5.12	0.247	0.130	2.15	0.733	6.01	5.41	6.01	5.41	3.25	1.83	1.83	3.25
62	STC (dialo)-N19C (a.salo)-SRIF28	5.53	4.57	1.04	0.550	1.10	3.75	26.0	24.0	26.0	24.0	37.0	21.0	21.0	37.0
63	STC (dialo)-N19C (a.salo)-SRIF28	6.92	5.73	0.830	0.440	2.08	0.710	4.76	4.28	4.76	4.28	2.67	1.51	1.51	2.67
64	STC (dialo)-N19C (a.salo)-SRIF28	6.01	4.97	0.830	0.440	2.08	0.710	4.76	4.28	4.76	4.28	2.67	1.51	1.51	2.67
65	STC (dialo)-SRIF28	15.0	13.0	0.810	0.330	5.57	1.91	15.0	14.0	15.0	14.0	1.61	0.910	0.910	1.61
66	STC (dialo)-N19C (a.salo)-SRIF28	35.0	29.0	9.65	3.00	3.90	1.33	22.0	22.0	22.0	22.0	8.53	4.82	4.82	8.53
67	STC (dialo)-N19C (a.salo)-SRIF28	6.20	5.13	0.960	0.510	3.89	0.690	6.12	5.61	6.12	5.61	1.81	1.02	1.02	1.81
68	STC (dialo)-SRIF28	130	109	3.76	2.00	10.0	3.43	86.0	59.0	86.0	59.0	10.0	6.42	6.42	10.0
SRIF28	0.533	1.80	0.0480	0.0365	0.206	0.206	0.110	0.776	1.23	0.776	1.23	0.189	0.090	0.090	0.189
-	SRIF 4	1.25	2.69	0.0329	0.0933	0.0697	0.403	0.823	1.95	0.823	1.95	0.416	0.313	0.313	0.416

As shown in Table 3B, the controls SRIF14 and SRIF28 bound to all receptors SSTR1 - SSTR5. Since the number of

test runs etc. was different from Example 69-1, the K_i value of SRIF14 against each receptor was different at a high value of 2.5 to 13.2. The K_i values of the compounds of Examples 1, 7, 8, 9, 26, 27, 32, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, and 68 against SSTR1 were 1.46 - 100 nM, and similarly 0.0536 - 6.58 nM for SSTR2, 0.160 - 13.8 nM for SSTR3, 1.39 - 172 nM for SSTR4, and 0.310 - 95.0 nM for SSTR5, showing high binding affinity towards all the receptors. The K_i values of the compound of Example 12 against SSTR2 and SSTR5 were 280 - 1600-folds that of SRIF14 but the K_i values against SSTR1, SSTR3, and SSTR4 were 17.9, 7.44, and 24.1 nM, and it was thought that binding affinity was retained. The affinity of the compound of Example 40 against SSTR1, SSTR2, and SSTR3 was significantly reduced with K_i values of 310 - 5300-folds that of SRIF14 but the K_i values against SSTR4 and SSTR5 were 110 and 33.0 nM, and it was thought that binding affinity was retained. The K_i values of the compound of Example 51 against SSTR2 and SSTR4 were 170-folds and 46-folds or more that of SRIF14 but the K_i values against SSTR1, SSTR3, and SSTR5 were each 270 nM, 14.0 nM, and 23.0 nM, and it was thought that binding affinity was retained.

[0263]

Example 69-3 Agonist activity evaluation employing receptor expression cells - 1

Somatostatin receptor is a G-protein-coupled receptor (GPCR). SSTR1 - SSTR5 all suppress adenylyl cyclase activity via G-protein subfamily G_i/G_o and reduces intracellular cAMP concentration. In this experimental line, each somatostatin receptor expression cells was employed to calculate the IC_{50} value of the cAMP production suppressing action of the test substance to evaluate agonist activity. Moreover, SRIF14, SRIF28 were similarly evaluated as control compounds.

Reagents employed in the present experiment and their proper chemical names are as follows: DMEM (Dulbecco's Modified Eagle's Medium), IBMX (3-isobutyl-1-methylxanthine), and HBSS (Hank's Buffered Salt Solution).

For evaluations of SSTR2 - SSTR5, experiments were performed under the following conditions. DMEM containing 0.3% BSA and 0.5 mM IBMX was employed as the buffer, and receptor expression cells shown in Table 3C were seeded at 10^4 cells/well. The test compounds was treated by mixing at a concentration of 0.00001 nM, 0.0001 nM, 0.001 nM, 0.01 nM, 0.1 nM, 1 nM, 10 nM, 100 nM, or 1000 nM with 10 μ M

forskolin, and allowed to react at room temperature for 30 minutes. After the reaction, the cells were dissolved with 0.2 N HCl, and the amount of cAMP accumulated in the cell was measured with Cayman cyclic AMP EIA kit (Cayman, 582002).

Evaluation of SSTR1 was consigned to Cerep, and the experiment was performed. HBSS containing 20 mM HEPES (pH 7.4) and 500 μ M IBMX was used as the buffer, and SSTR1 receptor expression cells were seeded at 10^4 cells/well. The test substance was treated by mixing at a concentration of 0.001 nM, 0.01 nM, 0.1 nM, 1 nM, 10 nM, or 100 nM with 1 μ M NKH477, and allowed to react at 37°C for 20 minutes. After the reaction, the amount of cAMP accumulated in the cell was measured with Cisbio cAMP dynamic2 kit (Cisbio, 62AM4PE).

[Table 11]

Table 3C

	SSTR2	SSTR3	SSTR4	SSTR5
Distributor	Perkin Elmer			
Serial Number	ES-521-CF	ES-523-CF	ES-524-CF	ES-522-CF
Derived cell	CHO-K1 cells			

[0264]

The experimental results (IC_{50} (nM)) of the agonist activity evaluation test when employing the compounds of Examples 1, 18, 27, 28, 59, and 60 as the glycosylated form are shown in Table 3D and Figure 1E. IC_{50} of the control compounds SRIF14 and SRIF28 was 0.83 - 0.89 nM against SSTR1, and 0.0076 - 0.073 nM, 0.029 - 0.21 nM, 0.015 - 0.074 nM, and 0.066 - 0.12 nM against SSTR2, SSTR3, SSTR4, and SSTR5, respectively. IC_{50} of the agonist activity of the glycosylated compounds against SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5 were 1.0 - 2.4 nM, 0.041 - 0.10 nM, 0.12 - 0.30 nM, 0.039 - 0.085 nM, and 0.043 - 0.081 nM, respectively, and potent agonist activity was shown against all receptors SSTR1 - SSTR5. From the results shown in Examples 69-1 and 69-2, it is apparent that these compounds have receptor binding affinity, and it became clear that these compounds exert proper agonistic action by binding to a receptor.

[Table 12]

Table 30

Example	SSTR1	SSTR2	SSTR3	SSTR4	SSTR5
-	0.83 ± 0.52	0.0076 ± 0.0012	0.029 ± 0.017	0.015 ± 0.006	0.12 ± 0.03
-	0.89 ± 0.41	0.073 ± 0.014	0.21 ± 0.08	0.074 ± 0.058	0.066 ± 0.033
1	2.2 ± 1.1	0.075 ± 0.030	0.13 ± 0.06	0.065 ± 0.042	0.056 ± 0.028
18	1.2 ± 0.6	0.055 ± 0.015	0.18 ± 0.10	0.050 ± 0.025	0.052 ± 0.032
27	1.6 ± 0.6	0.041 ± 0.006	0.12 ± 0.05	0.039 ± 0.021	0.055 ± 0.031
28	1.5 ± 0.6	0.079 ± 0.033	0.15 ± 0.08	0.051 ± 0.046	0.064 ± 0.029
59	1.0 ± 0.9	0.10 ± 0.02	0.30 ± 0.15	0.085 ± 0.011	0.081 ± 0.019
60	2.4 ± 1.7	0.055 ± 0.013	0.14 ± 0.06	0.043 ± 0.023	0.043 ± 0.015

[0265]

Example 69-4 Agonist activity evaluation employing receptor expression cells - 2

Agonist activity evaluation test was carried out similarly to Example 69-3 employing each of the compounds described in Table 3E as the glycosylated form. The experimental results are shown in Table 3E. In Table 3E, fields with "-" (hyphen) shown indicates that the test was not performed.

[Table 13]

Table 3E

(-: Not performed)

Experiment		SSTR1	SSTR2	SSTR3	SSTR4	SSTR5
-	SRIF14	0.83	0.0076	0.029	0.015	0.12
-	SRIF28	0.89	0.073	0.21	0.074	0.066
4	E12C(disialo)-SRIF28	2.3	-	-	-	-
10	N19C(disialo)-SRIF28	22	-	-	-	-
15	S1C(disialo)-D-Trp22-SRIF28	3.5	-	-	-	-
17	C(disialo)-SRIF14	-	0.029	0.18	0.10	0.23
19	C(disialo)-C12linker-SRIF14	-	-	-	0.11	-
20	S1-2C(disialo)-SRIF28	-	-	0.55	0.14	0.22
21	S1C(disialo)-N5C(disialo)-SRIF28	-	0.072	0.37	0.041	0.090
23	N5C(disialo)-A8C(disialo)-SRIF28	-	0.096	0.28	0.091	0.13
24	S1-3C(disialo)-SRIF28	7.5	0.19	0.79	0.36	0.97
25	S1C(disialo)-N5C(disialo)-A9C(disialo)-SRIF28	-	1.4	2.4	0.16	1.1
26	S1C(monosialo)-SRIF28	1.5	0.059	0.15	0.046	0.062
29	S1-3C(asialo)-SRIF28	-	0.19	-	0.10	-
31	S1N(disialo)-SRIF28	2.3	-	-	-	-
32	S1C(disialo)-N19C(GlcNAc)-SRIF28	-	0.080	0.039	0.033	0.088
34	S1-5C(disialo)-SRIF28	18	0.81	2.0	0.27	-
42	C(disialo)-K-SRIF14	-	0.019	0.10	0.059	-
44	S1C(disialo)-SRIF28-amide	1.7	-	-	-	-
47	C(disialo)-PEG2linker-SRIF14	-	0.034	0.15	0.066	0.18
50	2C(disialo)-R-K-SRIF14	-	0.15	0.55	1.1	-
51	3C(disialo)-R-K-SRIF14	-	1.4	3.8	1.6	-
52	S1C(diGlcNAc)-SRIF28	1.2	0.045	0.14	0.050	0.033
53	S1C(diMan)-SRIF28	1.2	0.035	0.11	0.043	0.037
54	N19C(diMan)-SRIF28	-	-	-	0.11	-
55	S1C(GlcNAc)-SRIF28	1.4	0.040	0.14	0.049	0.031
56	N19C(GlcNAc)-SRIF28	-	-	-	0.092	-
57	S1C(trisialo)-SRIF28	4.9	0.11	0.38	0.092	0.096
58	S1C(tetrasialo)-SRIF28	2.8	0.13	0.25	0.081	0.11
61	S1C(disialo(Bn))-SRIF28	1.4	-	-	-	-
62	S1C(disialo(hexadecylamide))-SRIF28	1.3	-	-	-	-
65	S1C(Asn(disialo))-SRIF28	2.6	-	-	0.13	-
67	C(disialo(aminoethylamide))-S1C(disialo)-SRIF28	-	0.039	0.17	0.047	0.034

[0266]

In Table 3E, IC₅₀ of the agonist activity of the glycosylated compounds against SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5 were 1.2 - 22 nM, 0.019 - 1.4 nM, 0.039 - 3.8 nM, 0.033 - 1.6 nM, and 0.031 - 1.1 nM, respectively, and agonist activity was shown against receptors SSTR1 - SSTR5. From the results shown in Examples 69-1 and 69-2, it became clear that these compounds have receptor binding affinity,

and these compounds exert agonist activity by binding to a receptor.

[0267]

Example 70 Pharmacokinetics test with rats 1

In order to confirm that the glycosylated polypeptide of the present invention (glycosylated form) had improved pharmacokinetics profiles such as drug plasma concentration - area under the time curve (AUC), half-life in blood ($t_{1/2}$), mean retention time (MRT), and bioavailability compared to a non-glycosylated SRIF28, pharmacokinetics analysis by intravenous and subcutaneous administrations was performed with rats.

[0268]

70-1 Preparation of administration solution and reagent

The glycosylated form (S1C(disialo)-SRIF28) was dissolved in Japanese Pharmacopeia saline (from Otsuka Pharmaceutical factory, Inc.) to prepare a 40 μ M solution to make an administration solution. PBS solution was prepared by dissolving 1 tablet of Phosphate buffered saline (P4417 from Sigma) in 200 mL of ultrapure water. EDTA-PBS was prepared by dissolving EDTA-2Na (from Wako Pure Chemical Industries, Ltd.) in PBS to 2 mg/mL. Aprotinin-containing EDTA-PBS solution was prepared by dissolving aprotinin (010-11834 from Wako Pure Chemical Industries, Ltd.) in EDTA-PBS to 0.142 mg/mL, and was employed as an anticoagulant for collected blood.

[0269]

70-2 Preparation of plasma sample

To the tail vain or dorsal subcutaneous of male SD rats (Cr1: CD (SD), Charles River Japan, 6 weeks-old, n = 3, body weight 161.3 - 239.3 g), the administration solution prepared in the above 70-1 was administered under nonfasting condition at a dosage of 1 mL/kg with a glass syringe and a 26 G injection needle (all from Terumo Corporation) (40 nmol/kg as S1C(disialo)-SRIF28). Blood was collected from the rat cervical vein before administration, as well as at 2 minutes, 5 minutes, 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, and 8 hours after administration. 0.2 mL of the collected blood was promptly mixed with 0.2 mL of the aprotinin-containing EDTA-PBS solution prepared in the above 70-1, and left on ice for 30 minutes or more. After centrifugal separation (1,870 x g, 4°C, 10 minutes), 250 μ L of the supernatant was taken as the plasma sample. As blank plasma, plasma obtained by similarly treating the blood collected from

untreated rat cervical vein was employed. Plasma samples were frozen in storage until employed for measurement. Tips and tubes used were low absorbent products from BM Equipment Co., Ltd.

[0270]

70-3 Measurement of concentration in plasma

Measurement of plasma concentration of S1C(disialo)-SRIF28 in the plasma sample obtained in the above 70-2 was performed with Phoenix Pharmaceuticals somatostatin EIA kit (Phoenix Pharmaceuticals Inc, EK-060-03). The plasma sample was diluted with the assay buffer supplied in the EIA kit to 5, 20, 100, 400, and 1600-folds as measurement samples. The standard solution for creating a standard curve was prepared as follows. First, the blank plasma obtained in the above 70-2 was diluted with the assay buffer supplied in the EIA kit similarly to the plasma sample, and this was used as the assay buffer for standard solution preparation (for example, when diluting the plasma sample to 100-fold, 1/100 amount of blank plasma was added to the assay buffer supplied in the EIA kit, and this was used as the assay buffer for standard solution preparation). S1C(disialo)-SRIF28 was diluted with PBS solution to prepare a 100 μ M solution, and a 2 μ M solution was prepared from the 100 μ M solution. The 2 μ M S1C(disialo)-SRIF28 solution obtained was serially diluted with the assay buffer for standard solution preparation to prepare standard solutions of 20 nM, 10 nM, 2 nM, 0.4 nM, 0.08 nM, and 0.04 nM. By multiplying the results obtained and the dilution ratio, and further multiplying the dilution ratio 2 in the aprotinin-containing EDTA-PBS solution used as anticoagulant treatment, the plasma concentration was calculated. As a control, a similar operation was carried out employing non-glycosylated SRIF28 instead of the glycosylated form. The transition of plasma S1C(disialo)-SRIF28 concentration obtained is shown in Figure 2.

[0271]

70-4 Estimation of pharmacokinetics parameter

From the transition of S1C(disialo)-SRIF28 concentration obtained, AUC was calculated by the moment analysis method and the trapezoidal rule. Moreover, the predicted initial concentration (C_0) was determined by the extrapolation method for intravenous administration, $t_{1/2}$ and MRT were calculated, and the maximum plasma concentration (C_{max}) was determined from the actual value for subcutaneous administration. The pharmacokinetics parameters obtained are shown in Table 4.

[Table 14]

Table 4

	Intravenous administration				Subcutaneous administration			
	$t_{1/2}$	AUC	MRT	C_0	$t_{1/2}$	AUC	MRT	C_{max}
S1C(disialo)-SRIF28	18.1	5086	20.4	780	26.0	1273	43.8	19.9
SRIF28	0.8	358	2.3	678	1.5	15	3.2	4.4

($t_{1/2}$: min, AUC: min·nM, MRT: min, C_0 : nM, C_{max} : nM)

As is clear from the results shown in Figure 2 and Table 4, S1C(disialo)-SRIF28 has significantly extended $t_{1/2}$ and MRT compared to the non-glycosylated SRIF28, and an increase in AUC and C_{max} was recognized. These are thought to be due to the increase in resistance to degradation activity in blood by glycosylation. It is clear that the glycosylated form has improved stability *in vivo* compared to the non-glycosylated form. Moreover, as factors for the increase in AUC and C_{max} , the improvement of bioavailability according to the present invention, as well as improvement of stability thereof *in vivo* are thought to be factors.

[0272]

Example 71 Pharmacokinetics test with rats 2

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, N5C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)-SRIF28 were employed as the glycosylated forms. As a control, non-glycosylated SRIF28 was employed instead of the glycosylated form. The compound plasma concentration transition obtained is shown in Figure 3.

[0273]

Example 72 Pharmacokinetics test with rats 3

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1C(disialo)·R13C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28 were employed as the glycosylated forms. As a control, non-glycosylated SRIF28 was employed instead of the glycosylated form. The compound plasma concentration transition obtained is shown in Figure 4.

[0274]

Example 73 Pharmacokinetics test with rats 4

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28, S1-3C(disialo)-SRIF28, S1-5C(disialo)-SRIF28, and S1-10C(disialo)-SRIF28 were employed as the glycosylated forms.

The compound plasma concentration transition obtained is shown in Figure 5.

[0275]

Example 74 Pharmacokinetics test with rats 5

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, N5C(disialo)-SRIF28, A9C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, N5C(disialo)·A9C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 6.

[0276]

Example 75 Pharmacokinetics test with rats 6

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, and S1-3C(disialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 7.

[0277]

From the compound plasma concentration transition obtained in the pharmacokinetics tests of Examples 71 - 75, the pharmacokinetics parameters of each compound were calculated similarly to Example 70. Moreover, the parameters obtained were employed to calculate bioavailability from the following mathematical formula. The results are shown in Table 5A.

$$BA (\%) = (AUC_{(sc)} / Dose_{(sc)}) / (AUC_{(iv)} / Dose_{(iv)})$$

AUC_(sc): AUC for subcutaneous administration (min·nM)

Dose_(sc): Administration dose for subcutaneous administration (nmol/kg)

AUC_(iv): AUC for intravenous administration (min·nM)

Dose_(iv): Administration dose for intravenous administration (nmol/kg)

[Table 15]

Table 5A

Compound	The number of sugar chains	Value of n	Intravenous administration			Subcutaneous administration			BA		
			$t_{1/2}$	AUC	MRT	C_0	$t_{1/2}$	AUC		MRT	C_{max}
- SRIF28	None	6	1.00 ±0.20	472 ±166	2.20 ±0.11	554 ±221	2.94 ±1.59	25.5 ±13.8	4.76 ±1.80	4.47 ±1.21	5
1 S1C(disialo)-SRIF28	1	15	16.9 ±1.02	4855 ±1030	18.6 ±1.72	686 ±363	29.8 ±2.68	1773 ±281	47.9 ±2.68	28.9 ±3.23	37
2 N5C(disialo)-SRIF28	1	6	20.1 ±1.49	4421 ±1366	22.9 ±2.16	496 ±312	28.1 ±3.72	2392 ±443	51.9 ±5.92	32.4 ±4.87	54
3 A9C(disialo)-SRIF28	1	3	19.9 ±0.33	4083 ±515	25.2 ±0.38	547 ±211	30.3 ±2.45	2462 ±64.0	55.6 ±5.81	30.2 ±5.05	60
Average value			19.0	4453	22.2	576.3	29.4	2209	51.8	30.5	50
20 S1-2C(disialo)-SRIF28	2 (dense)	3	28.8 ±3.11	7306 ±625	23.3 ±2.63	1382 ±321	40.9 ±12.2	4046 ±248	67.9 ±15.0	46.3 ±9.75	55
21 S1C(disialo) · N5C(disialo)-SRIF28	2 (both intervals)	6	25.8 ±3.41	4099 ±634	24.6 ±5.97	558 ±248	38.4 ±4.93	3344 ±663	67.9 ±8.39	35.5 ±6.38	82
22 S1C(disialo) · R13C(disialo)-SRIF28	2 (both intervals)	3	23.4 ±0.66	4029 ±184	20.6 ±1.32	822 ±106	39.6 ±2.50	3407 ±436	67.7 ±5.39	38.7 ±8.73	85
23 N5C(disialo) · A9C(disialo)-SRIF28	2 (both intervals)	3	32.4 ±1.66	4465 ±833	26.6 ±0.27	664 ±104	42.7 ±9.76	3450 ±784	75.0 ±12.4	32.5 ±7.06	77
			27.2	4198	23.9	581.3	40.2	3400	70.2	35.6	81
24 S1-3C(disialo)-SRIF28	3 (dense)	6	38.5 ±2.83	5660 ±1354	24.0 ±3.92	1061 ±209	87.7 ±21.4	3884 ±543	126 ±31.7	27.5 ±10.8	69
25 S1C(disialo) · N5C(disialo) · A9C(disialo)-SRIF28	3 (both intervals)	9	39.8 ±1.87	4402 ±999	25.8 ±2.87	938 ±145	87.4 ±17.1	4015 ±944	132 ±19.8	24.8 ±9.03	91
34 S1-5C(disialo)-SRIF28	5 (dense)	3	99.5 ±14.0	6016 ±873	49.0 ±6.06	752 ±81.4	271 ±38.7	5524 ±633	389 ±50.0	13.5 ±2.85	92
35 S1-10C(disialo)-SRIF28	10 (dense)	3	219 ±6.27	100259 ±10463	272 ±14.3	821 ±264	>480	-	-	45.9 ±2.32	-

($t_{1/2}$: min; AUC: min·nM; MRT: min·nM; C_0 : nM; C_{max} : nM; BA: %)

($t_{1/2}$: min , AUC : min·nM , MRT : min·nM , C_0 : nM , C_{max} : nM , BA : %)

[0278]

As is clear from the results shown in Table 5A, the bioavailability of the glycosylated form of the present

invention increased as the number of modifying sugar chains increased. In other words, what was 5% in the non-glycosylated form was 37 - 60% in the glycosylated polypeptide having one sugar chain added, an increase of 50% on average. On the other hand, it was found that this was increased to 77 - 85%, 81% on average in those having two sugar chains added with intervals, and to 91% in those having three sugar chains added with intervals. Moreover, comparing those with dense glycosylation intervals, it was found that there was a 37% increase in those having one added, 55% in those having two added, 69% in those having three added, and 92% in those having five added (the compounds of Examples 1, 20, 24, and 34). For those having ten sugar chains added, as shown in Figure 5, AUC could not be measured because the rise in plasma concentration was maintained throughout the measurement time, and as a result bioavailability could not be calculated. From these results, it was proven that bioavailability is significantly improved by glycosylation according to the present invention compared to somatostatin without any glycosylation, and a further improved bioavailability could be obtained by adding two or more sugar chains compared to those having one sugar chain added.

As factors for the increase in bioavailability for subcutaneous administration, various pharmacokinetic factors exist. Among these factors, some are thought to be the stability of the compound in blood, or transitivity into the blood (from the administration site). As shown in Table 5A, it is recognized that the AUC for subcutaneous administration which will be an indicator for speculating the transit to blood for subcutaneous administration will increase with the increase in the number of modifying sugar chains (1: 2209 min·nM, 2 (with intervals): 3400 min·nM, and 3 (with intervals): 4015 min·nM), and it was speculated to contribute to improvement bioavailability.

Moreover, it was recognized that as the number of modifying sugar chains increased, $t_{1/2}$ and MRT extension effect for intravenous and subcutaneous administrations (for example, $t_{1/2}$ for intravenous administration was 1: 19.0 min, 2 (with intervals): 27.2 min, and 3 (with intervals): 39.8 min), and it was speculated that stability in blood was improved. On the other hand, AUC for intravenous administration was not an increase in proportion to the number of modifying sugar chains (1: 4453 min·nM, 2 (with intervals): 4198 min·nM, and 3 (with intervals): 4402 min·nM), and it is thought that the increase in the stability in blood of the compound by the

increase in the number of modifying sugar chains is not the only factor contributing to the increase in bioavailability.

C_{max} for subcutaneous administration increased compared to the non-glycosylated form up until the number of modifying sugar chains was two (0: 4.47 nM, 1: 30.5 nM, and 2 (with intervals): 35.6 nM), and three (with intervals) (24.8 nM) resulted in an adverse decrease. When considering blood transit from the administration site (subcutaneous administration in the present Example), speculated factors for the increase in AUC are two formats of rapid transit to blood, or mild but continuous transit to blood. The former has the advantage of avoiding degradation at the administration site, but if there is no problem in stability, it is speculated that the latter format has a higher transit in blood. The fact that C_{max} had decreased in those having three modifying sugar chains with high bioavailability is thought to be the result showing not a sudden but a mild and continuous migration in blood. From this, it is thought that the increase in the number of modifying sugar chains related to the present invention shows a continuous migration in blood without a rapid rise in plasma concentration (may generally be referred to as absorption delaying effect).

[0279]

As is clear from the results shown in Table 5A, there was no significant difference in $t_{1/2}$ and MRT when the modifying positions had intervals and when they were dense (for example, $t_{1/2}$ for intravenous administration was 2 (dense): 28.8 min, 2 (with intervals): 27.2 min, 3 (dense): 38.5 min, and 3 (with intervals): 39.8 min).

Moreover, with respect to AUC, for intravenous administration, AUC was increased when glycosylation was dense. In other words, for two sugar chains, this was 4029 - 4465 min·nM for compounds 21, 22, and 23 with intervals as opposed to 7306 min·nM for a dense compound 20, and for three sugar chains, compound 25 with intervals was 4402 min·nM whereas a dense compound 24 was 5660 min·nM.

On the other hand, bioavailability increased when there were intervals in the glycosylation positions compared to when they were dense. In other words, for two sugar chains, a dense compound 20 was 55% whereas compounds 21, 22, and 23 with intervals were 77 - 85% (81% on average), and for three sugar chains, a dense compound 24 was 69% whereas compound 25 with intervals was 91%. From this, it is thought that for glycosylation positions related to the present invention, multiple modifications with intervals contribute more to the improvement of bioavailability than dense multiple modifications.

[0280]

Example 76 Pharmacokinetics test with rats 7

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1C(disialo(amide))-SRIF28, and S1C(disialo(aminoethylamide))-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 8.

[0281]

Example 77 Pharmacokinetics test with rats 8

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1C(disialo(Bn))-SRIF28, and S1C(disialo)-D-Trp22-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 9.

[0282]

Example 78 Pharmacokinetics test with rats 9

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, R13C(disialo)-SRIF28, and K14C(disialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 10.

[0283]

Example 79 Pharmacokinetics test with rats 10

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, E12C(disialo)-SRIF28, N19C(disialo)-SRIF28, 29C(disialo)-SRIF28, S1C(monosialo)-SRIF28, and S1C(asialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 11.

[0284]

Example 80 Pharmacokinetics test with rats 11

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, K14C(disialo)-SRIF28, and C(disialo)-SRIF14 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 12.

[0285]

Example 81 Pharmacokinetics test with rats 12

A pharmacokinetics test was carried out similarly to Example 70, except that C(disialo)-SRIF14, C(disialo)-C12

linker-SRIF14, and C(disialo)-PEG linker-SRIF14 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 13.

[0286]

Example 82 Pharmacokinetics test with rats 13

A pharmacokinetics test was carried out similarly to Example 70, except that SRIF28, S1C(disialo)-SRIF28, S1C(asialo)-SRIF28, S1C(diGlcNAc)-SRIF28, S1C(diMan)-SRIF28, and S1C(GlcNAc)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 14.

[0287]

Example 83 Pharmacokinetics test with rats 14

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1C(tetrasialo)-SRIF28, S1C(trisialo)-SRIF28, S1C(Asn(disialo))-SRIF28, and S1-2C(disialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 15.

[0288]

Example 84 Pharmacokinetics test with rats 15

A pharmacokinetics test was carried out similarly to Example 70, except that SRIF28, S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, S1-3C(disialo)-SRIF28, and S1-4C(disialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 16.

[0289]

Example 85 Pharmacokinetics test with rats 16

A pharmacokinetics test was carried out similarly to Example 70, except that SRIF14, C(disialo)-SRIF14, C(disialo)-K-SRIF14, C(disialo)-R-K-SRIF14, 2C(disialo)-R-K-SRIF14, and 3C(disialo)-R-K-SRIF14 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 17.

[0290]

Example 86 Pharmacokinetics test with rats 17

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(asialo)-SRIF28, S1-2C(asialo)-SRIF28, and S1-3C(asialo)-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 18.

[0291]

Example 87 Pharmacokinetics test with rats 18

A pharmacokinetics test was carried out similarly to Example 70, except that S1C(disialo)-SRIF28, S1-2C(disialo)-SRIF28, S1-2C(asialo)-SRIF28, C(disialo(aminoethylamide))/S1C(disialo)-SRIF28, S1-2C(disialo(Bn))-SRIF28, and S1-2C(disialo(amide))-SRIF28 were employed as the glycosylated forms. The compound plasma concentration transition obtained is shown in Figure 19.

[0292]

From the compound plasma concentration transition obtained in pharmacokinetics tests 7 - 18 of Examples 76 - 87, the pharmacokinetics parameters of each compound were calculated similarly to Example 70. The pharmacokinetics parameters obtained are shown in Tables 5B - 5F.

[Table 16]

Table 5B

Example		Intravenous administration				Subcutaneous administration				BA
		$T_{1/2}$	AUC	MRT	C_0	$T_{1/2}$	AUC	MRT	C_{max}	
-	SRIF14	0.500	119	2.00	819	2.30	8	4.30	1.80	7
-	SRIF28	1.32	467	2.46	518	2.38	22	4.18	4.65	5
1	S1C(disialo)-SRIF28	17.7	5190	19.1	633	31.2	1946	49.2	30.1	38
4	E12C(disialo)-SRIF28	19.5	10521	27.5	1086	38.2	2891	57.5	34.7	27
5	R13C(disialo)-SRIF28	20.3	4926	22.8	481	35.4	5528	55.9	85.9	78
6	K14C(disialo)-SRIF28	20.4	6278	20.0	884	35.1	3945	55.1	57.5	62
10	N19C(disialo)-SRIF28	18.8	3854	21.5	407	37.9	2624	55.3	41.7	56
13	29C(disialo)-SRIF28	23.7	3736	25.3	359	39.3	2176	57.6	29.4	51
15	S1C(disialo)-D-Trp-SRIF28	18.3	4924	20.6	588	31.0	2220	51.9	34.0	45
18	C(disialo)-R-K-SRIF14	18.9	6571	22.5	429	28.4	1996	45.2	32.5	30
42	C(disialo)-K-SRIF14	18.8	5277	19.4	646	26.4	2446	43.7	40.5	46
17	C(disialo)-SRIF14	16.5	5614	18.3	725	27.2	3641	40.0	73.6	66
19	C(disialo)-C12linker-SRIF14	11.0	1686	7.60	500	14.4	345	22.0	13.4	21
45	C(disialo)-PEGlinker-SRIF14	16.4	3538	18.8	412	20.7	2725	39.2	47.8	77

As is clear from the results shown in Table 5B, S1C(disialo)-SRIF28, E12C(disialo)-SRIF28, R13C(disialo)-SRIF28, K14C(disialo)-SRIF28, N19C(disialo)-SRIF28, and 29C(disialo)-SRIF28, S1C(disialo)-D-Trp22-SRIF28 extended $t_{1/2}$ by 13 - 18-folds and increased AUC by 8 - 23-folds compared to the non-glycosylated form SRIF28 when administered intravenously. Moreover, C(disialo)-SRIF14, C(disialo)-K-SRIF14, and C(disialo)-R-K-SRIF14 extended $t_{1/2}$ by 33 - 38-folds and increased AUC by 44 - 55-folds compared to the non-glycosylated form SRIF14 when administered intravenously. Moreover, C(disialo)-C12 linker-SRIF14 and C(disialo)-PEG linker-SRIF14 extended $t_{1/2}$ by 22 - 33-folds and increased AUC by 14 - 30-folds compared to the non-glycosylated form SRIF14. In other words, similarly to Example 70, this was thought to be the result of improved stability in blood by glycosylation. [Table 17]

Table 50

Example		Intravenous administration				Subcutaneous administration			
		T _{1/2}	AUC	MRT	C ₀	T _{1/2}	AUC	MRT	C _{max}
-	SRIF28								
55	SIC(GlcNAc)-SRIF28	1.32	467	2.46	518	2.38	22	4.18	4.65
53	SIC(diMan)-SRIF28	1.80	690	3.00	666	5.90	63	7.90	7.10
52	SIC(diGlcNAc)-SRIF28	5.50	1445	5.20	423	9.10	124	13.9	7.30
27	SIC(Asialo)-SRIF28	10.5	3588	12.3	495	16.5	445	25.7	13.2
26	SIC(monosialo)-SRIF28	13.4	4602	12.5	749	19.7	710	29.0	17.2
1	SIC(disialo)-SRIF28	15.9	6616	15.3	1061	24.5	995	37.8	19.4
57	SIC(trisialo)-SRIF28	17.7	5190	19.1	633	31.2	1946	49.2	30.1
58	SIC(tetrasialo)-SRIF28	21.1	4428	22.0	423	36.8	1979	56.9	24.9
65	SIC(Asn(disialo))-SRIF28	20.9	4851	23.0	471	40.5	2624	59.4	32.2
59	SIC(disialo(aminoethylamide))-SRIF28	27.9	6488	35.9	365	34.0	2935	55.3	33.9
60	SIC(disialo(amide))-SRIF28	18.6	3473	12.1	746	27.1	284	33.2	9.60
61	SIC(disialo(Bn))-SRIF28	18.0	5814	12.6	1048	30.0	961	42.5	19.1
		14.4	3708	14.5	526	45.5	1047	52.6	18.7
									28

[0293]

As is clear from the results shown in Table 5C, when the size of the modifying sugar chain is altered to GlcNAc (monosaccharide), diMan (5 sugars), diGlcNAc (7 sugars), asialo (9 sugars), monosialo (10 sugars), disialo (11 sugars), trisialo (14 sugars), and tetrasialo (17 sugars), the $t_{1/2}$, AUC, and bioavailability when administered subcutaneously increased to 2 - 17-folds, 3 - 120-folds, and 2 - 11-folds, respectively, according to the size of the modifying sugar chain. From this, it became clear that altering the size of the modifying sugar chain not only improved the stability in blood, but allowed change in its increase rate. Moreover, since it is known that dimannose or asialo sugar chains etc. among these sugar chains interact with particular proteins, they are thought to be utilizable for targeting a particular protein or organs and cells having the particular protein. Moreover, as modification of the sialic acid at the non-reducing terminal, S1C(disialo(aminoethylamide))-SRIF28, S1C(disialo(amide))-SRIF28, and S1C(disialo(Bn))-SRIF28 having added a sugar chain with altered charge by e.g. the introduction of an aminoethylamide, amide, and Bn to the carboxy group extended $t_{1/2}$ by 11 - 14-folds and increased AUC by 7 - 12-folds, and improved stability in blood compared to the non-glycosylated form when administered intravenously. This shows that the carboxy group of the sialic acid has a large influence on retentivity in blood, and shows the possibility for (usage in) blood clearance or body distribution control by abstraction of the negative charge of the carboxy group (disialo(Bn) and disialo(amide)) or conversion into a positive charge (disialo(aminoethylamide)).

[Table 18]

Table 5D

Example		Intravenous administration				Subcutaneous administration				BA
		$T_{1/2}$	AUC	MRT	C_0	$T_{1/2}$	AUC	MRT	C_{max}	
-	SRIF14	0.500	119	2.00	819	2.30	8	4.30	1.80	7
-	SRIF28	1.32	467	2.46	518	2.38	22	4.18	4.65	5
18	C(disialo)-R-K-SRIF14	18.9	6571	22.5	429	28.4	1996	45.2	32.5	30
50	2C(disialo)-R-K-SRIF14	31.7	5041	24.1	704	47.7	3310	73.4	36.7	66
51	3C(disialo)-R-K-SRIF14	35.7	4268	22.2	1013	70.6	3515	112	23.5	82
1	S1C(disialo)-SRIF28	17.7	5190	19.1	633	31.2	1946	49.2	30.1	38
20	S1-2C(disialo)-SRIF28	26.2	5804	23.0	950	40.8	3091	65.3	34.6	53
24	S1-3C(disialo)-SRIF28	39.5	5834	23.8	1083	79.3	4174	115	30.7	72
68	S1-4C(disialo)-SRIF28	63.5	7070	25.2	1007	119	4554	172	22.0	64

[0294]

As is clear from the results shown in Table 5D, C(disialo)-R-K-SRIF14 modified with one disialosugar chain, 2C(disialo)-R-K-SRIF14 modified with two disialosugar

chains, and 3C(disialo)-R-K-SRIF14 modified with three disialosugar chains on SRIF14 each extended $t_{1/2}$ by 38, 63, and 71-folds and increased AUC by 55, 42, and 36-folds compared to the non-glycosylated form when administered intravenously. Similarly, S1C(disialo)-SRIF28 modified with one disialosugar chain, S1-2C(disialo)-SRIF28 modified with two disialosugar chains, S1-3C(disialo)-SRIF28 modified with three disialosugar chains, and S1-4C(disialo)-SRIF28 modified with four disialosugar chains on SRIF28 each extended $t_{1/2}$ by 13, 21, 30, and 48-folds and increased AUC by 11, 12, 12, and 15-folds compared to the non-glycosylated form when administered intravenously. In either of SRIF14 and SRIF28, it became clear that stability in blood is improved according to the number of modifying disialosugar chain.

[Table 19]

Table 5E

Example		Intravenous administration				Subcutaneous administration				BA
		$T_{1/2}$	AUC	MRT	C_0	$T_{1/2}$	AUC	MRT	C_{max}	
-	SRIF28	1.32	467	2.46	518	2.38	22	4.18	4.65	5
27	S1C(asialo)-SRIF28	13.4	4602	12.5	749	19.7	710	29.0	17.2	16
28	S1-2C(asialo)-SRIF28	8.30	2664	9.60	741	19.2	1049	36.4	19.8	39
29	S1-3C(asialo)-SRIF28	6.60	2733	6.50	1147	26.4	885	42.1	15.0	32

[0295]

As is clear from the results shown in Table 5E, S1C(asialo)-SRIF28 modified with one asialosugar chain, S1-2C(asialo)-SRIF28 modified with two asialosugar chains, and S1-3C(asialo)-SRIF28 modified with three asialosugar chains on SRIF28 each extended $t_{1/2}$ by 10, 6, and 5-folds and increased AUC by 10, 6, and 6-folds compared to the non-glycosylated form when administered intravenously. It became clear that stability in blood is also improved when the modifying sugar chain is an asialosugar chain.

[Table 20]

Table 5F

Example		Intravenous administration				Subcutaneous administration				BA
		T _{1/2}	AUC	MRT	C ₀	T _{1/2}	AUC	MRT	C _{max}	
-	SRIF28	1.32	467	2.46	518	2.38	22	4.18	4.65	5
1	SI-C(disialo)-SRIF28	17.7	5190	19.1	633	31.2	1946	49.2	30.1	38
20	SI-2C(disialo)-SRIF28	28.2	5804	23.0	950	40.8	3091	65.3	34.6	53
28	SI-2C(asialo)-SRIF28	8.30	2664	9.60	741	19.2	1049	36.4	19.8	39
63	SI-2C(disialo(amide))-SRIF28	23.8	5268	20.5	737	33.1	1297	55.8	16.3	25
64	SI-2C(disialo(Bn))-SRIF28	25.6	5525	20.7	807	33.9	1843	59.4	21.5	33
67	C(disialo(aminoethylamide))-SI-C(disialo)-SRIF28	22.5	3742	20.5	518	42.3	1289	64.7	16.2	34

[0296]

As is clear from the results shown in Table 5F, S1-2C(disialo)-SRIF28, S1-2C(asialo)-SRIF28, C(disialo(aminoethylamide))/S1C(disialo)-SRIF28, S1-2C(disialo(Bn))-SRIF28, and S1-2C(disialo(amide))-SRIF28 each extended $t_{1/2}$ by 6 - 21-folds and increased AUC by 6 - 12-folds compared to the non-glycosylated form when administered intravenously. In other words, stability in blood improved by modifying a sugar chain onto SRIF28. Moreover, $t_{1/2}$, AUC, Cmax, and BA increased in both intravenous and subcutaneous administrations according to the increase in the number of sialic acid sugar chains. On the other hand, when the negative charge of the carboxy group of the sialic acid was removed (2C(disialo(Bn))/2C(disialo(amide))), or when a positive charge was adjacently added (C(disialo(aminoethylamide))/S1C(disialo)), although $t_{1/2}$ was extended when administered subcutaneously, AUC or Cmax did not increase. Since the carboxy group of the sialic acid has a large influence on retentivity in blood or blood migration, this shows the possibility for (usage in) blood clearance or body distribution control by conversion of charge at the sugar chain terminal.

[0297]

Example 88 Plasma stability test with rat plasma

88-1 Preparation of compound solution, reagent, and rat plasma

The glycosylated form and the non-glycosylated SRIF28 were dissolved in PBS solution to prepare 2 μ M solutions as treatment solutions. 10% TFA was prepared by dissolving trifluoroacetic acid (208-02746 from Wako Pure Chemical Industries, Ltd.) in water to 10 v/v%. Rat plasma was prepared from Wistar rats (Crlj: Wistar, male, Charles River Japan, 7 weeks-old) as heparin-added plasma (heparin: Japanese Pharmacopeia heparin sodium injectable solution (Mochida Pharmaceutical Co., Ltd.)).

[0298]

88-2 Preparation of plasma-added sample

To 0.27 mL of rat plasma ($n = 3$), 0.03 mL of the glycosylated compound solution prepared in the above 88-1 was promptly mixed as plasma-added sample, and kept warm in a 37°C thermostat bath. After mixing, 0.04 mL of plasma-added sample was taken at 0 minute and over time at 1 - 24 hours, and then promptly mixed with 0.01 mL of 10% TFA. After centrifugal separation (20,000 x g, 4°C, 10 minutes), 0.04 mL of the supernatant was taken as plasma stability

measurement samples. Sampling time was 0, 1, 2, and 4 hours, or 0, 4, 8, and 24 hours. As blank plasma, plasma obtained from a similar treatment except that PBS solution was employed as the treatment solution was employed. Plasma samples were frozen in storage until employed for measurement. For each experiment run, non-glycosylated SRIF28 was employed as the positive control.

[0299]

88-3 Measurement of concentration in sample and calculation of plasma stability index

In a method similar to plasma concentration measurement of Example 70-3, the concentration of the glycosylated form remaining in the plasma stability measurement samples obtained in the above 88-2 was measured. The concentration of the glycosylated form at 0 minute after mixing was set as 100%, the residual rate over time was represented in percentage (%), and the half-life was calculated from the elimination rate constant of the following exponential formula (1) employing the calculation formula (2). Then, the half-life of the non-glycosylated SRIF28 for each experiment run was set as 1, and the plasma stability index (PS index) of the glycosylated form was calculated. The results are shown in Table 6 and Figure 20. Moreover, those having a plasma stability index of 20 or higher are shown as >20. In Figure 20, those having a plasma stability index of higher than 20 are also shown with 20 as the upper limit.

$$\text{Residual rate (\%)} = 100 \cdot e^{(k \cdot t)} \dots (1)$$

e: Base of natural logarithm

k: elimination rate constant

t: Time (hours)

$$\text{Half-life (hours)} = 0.693/k \dots (2)$$

[Table 21]

Table 6

Example		PS index
1	S1C(disialo)-SRIF28	4.5
2	N5C(disialo)-SRIF28	4.7
3	A9C(disialo)-SRIF28	6.7
4	E12C(disialo)-SRIF28	6.2
5	R13C(disialo)-SRIF28	15.9
6	K14C(disialo)-SRIF28	19.1
10	N19C(disialo)-SRIF28	>20
13	29C(disialo)-SRIF28	>20
14	30C(disialo)-SRIF28	16.8
15	S1C(disialo)-D-Trp-SRIF28	12.3
16	A9C(disialo)-D-Trp-SRIF28	17.2
21	S1C(disialo) · N5C(disialo)-SRIF28	17.8
22	S1C(disialo) · R13C(disialo)-SRIF28	>20
24	S1-3C(disialo)-SRIF28	>20
25	S1C(disialo) · N5C(disialo) · A9C(disialo)-SRIF28	>20
26	S1C(monosialo)-SRIF28	3.5
27	S1C(asialo)-SRIF28	2.6
-	Octreotide	>20
-	SRIF28	1

[0300]

As is clear from the results shown in Figure 20, the glycosylated polypeptide of the present invention was increased in plasma stability compared to SRIF28. In other words, this was 4.5 - 6.7-folds in those having one sugar chain added at positions 1, 5, 9, and 12 (the compounds of Examples 1, 2, 3, and 4), 15.9-folds for those having a sugar chain added at position 13 (the compound of Example 5), 19.1-folds for those having a sugar chain added at position 14 (the compound of Example 6), and 16.8-folds for those having a sugar chain added at position 30 (the compound of Example 14), compared to that of SRIF28. Moreover, this was 20-folds or more for those having one glycosylated amino acid further added at position 19 which is the C-terminal side (the compounds of Examples 10 and 13). It was shown that in regards to the position for glycosylation, plasma stability increases from position 1 towards the C-terminal.

[0301]

Example 89 GH production suppression test with rats

As shown in Example 69, the glycosylated polypeptide of the present invention had affinity towards each receptor of SSTR. On the other hand, some were seen to have attenuated affinity towards each SSTR, but in order to prove that even in such cases pharmacologically effective action is shown towards receptors *in vivo* by e.g. extension of half-life in blood or increase in bioavailability as shown in Examples 70 - 88, a test for evaluating the administration effect of the glycosylated polypeptide of the present invention on growth hormone (GH) production was carried out as an *in vivo* test employing rats. Increase of GH production amount into the blood is thought to cause proliferation or differentiation of cells and activation or suppression of the biosynthetic system in various organs via the paracrine effect of GH, and thereby influence biological reactions. Somatostatin released from the hypothalamus suppresses GH secretion from the anterior pituitary gland into the blood. This experimental line was carried out as a test line that can evaluate the pharmacological action of the glycosylated form on SSTR and its residence in blood after administration, with blood GH production amount as an indicator.

[0302]

89-1 Preparation of administration solution and reagent

Employing S1C(disialo)-SRIF28, N19C(disialo)-SRIF28, 29C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, and S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28 as the glycosylated forms, these were dissolved in Japanese Pharmacopeia saline (from Otsuka Pharmaceutical factory, Inc.) to prepare 1 - 100 μ M solutions as the administration solution. Moreover, GRF (GH releasing hormone, Growth hormone releasing factor) was used as the GH release enhancer. GRF was prepared by dissolving GRF injectable solution (GRF Sumitomo 50 for injection, Lot. 2006C, from Dainippon Sumitomo Pharma Co., Ltd.) in 1 mL of water for injection (Lot. 09H18C, from Fuso Pharmaceutical Industries, Ltd.), and then diluting to 25-folds with saline to obtain 2 μ g/mL. As heparin employed as the anticoagulant when collecting blood, Japanese Pharmacopeia heparin sodium injectable solution (Lot. B084, from Mochida Pharmaceutical Co., Ltd.) was used directly as stock solution.

[0303]

89-2 Preparation of plasma sample

To the dorsal subcutaneous of male SD rats (Crl: CD (SD), Charles River Japan, 6 weeks-old, n = 3, body weight 145.2 - 163.9 g), the administration solution prepared in the above 89-1 was administered under nonfasting condition at a dosage of 1 mL/kg with a glass syringe and a 26 G injection needle (all from Terumo Corporation). As the control group, saline without glycosylated polypeptide was similarly administered (vehicle group). Then, i.e. between 5 - 6 minutes after the administration of the glycosylated form, pentobarbital sodium (sompentyl, Lot. 0608101, from Kyoritsuseiyaku Corporation) was intraperitoneally administered as a general anesthetic with a glass syringe and an injection needle to give 50 mg/kg. One hour after administration of the glycosylated form, i.e. after 50 minutes or more had passed under anesthesia, GRF was administered as the GH release enhancer to the tail vein at a dosage of 1 mL/kg with a glass syringe and an injection needle. Five minutes after administration of GRF, blood was collected from the rat cervical vein with a glass syringe containing heparin and an injection needle. 0.4 mL of the collected blood was left on ice for 20 minutes or more, then centrifuged (1,870 x g, 4°C, 10 minutes), and 100 µL of the supernatant was taken as the plasma sample. As blank plasma, plasma obtained by similarly treating the blood collected from untreated rat cervical vein was employed. Plasma samples were frozen in storage until employed for measurement.

[0304]

89-3 Measurement of GH concentration in plasma

Measurement of GH concentration in the plasma sample obtained in the above 89-2 was performed with rat Growth Hormone EIA kit from SPI-Bio (SPI-Bio, A05104). The plasma sample was diluted with the assay buffer supplied in the EIA kit to 20, 100, and 500-folds as measurement samples. The standard solution for creating a standard curve followed the attached instructions by preparing a 40 ng/mL solution with distilled water, and then subjecting to serial dilution with the assay buffer to prepare 20 ng/mL, 10 ng/mL, 5 ng/mL, 2.5 ng/mL, 1.25 ng/mL, 0.63 ng/mL, and 0.31 ng/mL. By multiplying the results obtained and the dilution ratio, the GH concentration was calculated. The GH concentration in the plasma sample obtained is shown in Table 7.

[Table 22]

Table 7

Example		Dose	GH (ng/mL)
1	S1C(disialo)-SRIF28	1nmol/kg	490
1	S1C(disialo)-SRIF28	10nmol/kg	17.1
10	N19C(disialo)-SRIF28	10nmol/kg	563
10	N19C(disialo)-SRIF28	100nmol/kg	1.5
13	29C(disialo)-SRIF28	10nmol/kg	497
13	29C(disialo)-SRIF28	100nmol/kg	88.5
21	S1C(disialo)-N5C(disialo)-SRIF28	1nmol/kg	776
21	S1C(disialo)-N5C(disialo)-SRIF28	10nmol/kg	181
25	S1C(disialo)-N5C(disialo)-A9C(disialo)-SRIF28	10nmol/kg	833
25	S1C(disialo)-N5C(disialo)-A9C(disialo)-SRIF28	100nmol/kg	151
-	vehicle	-	861

[0305]

As is clear from the results shown in Table 7, the glycosylated form of the present invention suppressed GH production in rats. The ratio between the K_i value of glycosylated forms of Examples 21 and 25 against each receptor and the K_i value of unglycosylated SRIF14 is shown to be in the range of 15.7:1 - 1.2:1 (the compound of Example 21) and 158:1 - 7.1:1 (the compound of Example 25), respectively, as measured in the method of Example 69-1. Moreover, the glycosylated forms of Examples 21 and 25 are shown to have 25-folds or more (the compound of Example 21) or 30-folds or more (the compound of Example 25) increase in half-life in blood, respectively, as measured in the method of Example 70. On the other hand, from the present Example having two sugar chains, it was shown that the glycosylated form effectively exerts pharmacological action even *in vivo*.

Moreover, the glycosylated form of the present invention had GH production suppressing action even when administered 1 hour before GRF administration.

It was shown that the effective administration dose for rat GH production ability between the glycosylated forms of Examples 1 and 21 and Example 25 was different by approximately 10-folds. This is similar to the receptor affinities thereof having a difference of approximately 10-folds in terms of K_i value, as measured in the method of Example 69-1. It was shown that they have pharmacological activity even when the affinity of the glycosylated form was somewhat attenuated.

[0306]

Example 90 GH production suppression test with rats 2

Similarly to Examples 89-1 - 89-3, rat GH production suppression tests of the glycosylated compounds shown below were carried out. S1C(disialo)-SRIF28, N5C(disialo)-SRIF28, A9C(disialo)-SRIF28, E12C(disialo)-SRIF28, R13C(disialo)-SRIF28, K14C(disialo)-SRIF28, S1C(disialo)-D-Trp22-SRIF28, S1C(disialo(Bn))-SRIF28, S1C(disialo(amide))-SRIF28, S1C(disialo(aminoethylamide))-SRIF28, S1C(monosialo)-SRIF28, S1C(asialo)-SRIF28, C(disialo)-R-K-SRIF14, S1-2C(asialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28, and S1-5C(disialo)-SRIF28 were employed as the glycosylated forms. The GH concentration in the plasma sample obtained is shown in Table 8.

[Table 23]

Table 8

Example		Dose	GH (ng/mL)	The number of runs
1	S1C(disialo)-SRIF28	1nmol/kg	787	4
		3nmol/kg	182	3
		10nmol/kg	15	5
		30nmol/kg	8	1
		100nmol/kg	7	2
2	N5C(disialo)-SRIF28	1nmol/kg	969	1
		10nmol/kg	12	1
3	A9C(disialo)-SRIF28	1nmol/kg	577	1
		10nmol/kg	20	1
4	E12C(disialo)-SRIF28	10nmol/kg	10	1
5	R13C(disialo)-SRIF28	10nmol/kg	136	1
6	K14C(disialo)-SRIF28	10nmol/kg	55	1
15	S1C(disialo)-D-Trp22-SRIF28	0.3nmol/kg	986	3
		1nmol/kg	335	3
		3nmol/kg	50	1
		10nmol/kg	6	2
		30nmol/kg	2	1
18	C(disialo)-R-K-SRIF14	1nmol/kg	912	4
		3nmol/kg	291	4
		10nmol/kg	30	4
		30nmol/kg	12	1
21	S1C(disialo)-N5C(disialo)-SRIF28	1nmol/kg	776	1
		10nmol/kg	181	1
25	S1C(disialo)-N5C(disialo)-A9C(disialo)-SRIF28	100nmol/kg	151	1
26	S1C(monosialo)-SRIF28	10nmol/kg	47	1
27	S1C(asialo)-SRIF28	1nmol/kg	879	4
		3nmol/kg	515	4
		10nmol/kg	146	4
28	S1-2C(asialo)-SRIF28	3nmol/kg	997	3
		10nmol/kg	175	3
34	S1-5C(disialo)-SRIF28	10nmol/kg	842	1
59	S1C(disialo(aminoethylamide))-SRIF28	10nmol/kg	314	1
60	S1C(disialo(amide))-SRIF28	10nmol/kg	14	1
61	S1C(disialo(Bn))-SRIF28	10nmol/kg	49	1
-	vehicle	-	1012	19

[0307]

As is clear from the results shown in Table 8, the glycosylated polypeptide of the present invention suppressed GH production in rats. In the present test line, S1C(disialo)-SRIF28 suppressed GH production from 1 nmol/kg, and showed 82 - 99% of the GH production suppression effect at 3 - 10 nmol/kg. This is thought to be due to the gain in affinity towards SSTR1 - SSTR5 and improvement in the

retentivity in blood as shown in Examples 69-1, 69-2, and 70 - 87.

In Examples 69-1 and 69-2, N5C(disialo)-SRIF28, A9C(disialo)-SRIF28, E12C(disialo)-SRIF28, S1C(disialo)-D-Trp-SRIF28, S1C(disialo(Bn))-SRIF28, and C(disialo)-R-K-SRIF14 showing affinity equivalent to S1C(disialo)-SRIF28 showed 95 - 99% of the GH production suppression effect at 10 nmol/kg, showing an effect equivalent to S1C(disialo)-SRIF28.

[0308]

From the results shown in the methods of Examples 70 - 87, S1C(monosialo)-SRIF28, S1C(asialo)-SRIF28, S1-2C(asialo)-SRIF28, and S1C(disialo(amide))-SRIF28 had an AUC for subcutaneous administration which was 1/3 - 1/2 of S1C(disialo)-SRIF28, and S1C(disialo(aminoethylamide))-SRIF28 was 1/7. Meanwhile, all of these showed higher affinity than S1C(disialo)-SRIF28 by the methods shown in Examples 69-1 and 69-2. Accordingly, in this experimental line, S1C(monosialo)-SRIF28, S1C(asialo)-SRIF28, S1-2C(asialo)-SRIF28, S1C(disialo(aminoethylamide))-SRIF28, and S1C(disialo(amide))-SRIF28 suppressed 70 - 99% of the GH production at 10 nmol/kg, and is thought to have shown an effect equivalent to S1C(disialo)-SRIF28.

[0309]

R13C(disialo)-SRIF28, K14C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28, and S1-5C(disialo)-SRIF28 showed a lower affinity than S1C(disialo)-SRIF28 by the methods shown in Examples 69-1 and 69-2. Meanwhile, from the results shown in the methods of Examples 70 - 87, these compounds have AUC for subcutaneous administration that was improved to 1.8 - 3.1-folds of S1C(disialo)-SRIF28. In this experimental line, R13C(disialo)-SRIF28, K14C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)-SRIF28, S1C(disialo)·N5C(disialo)·A9C(disialo)-SRIF28, and S1-5C(disialo)-SRIF28 showed GH production suppression activity by administration of 10 or 100 nmol/kg. These results show that somatostatin activity can be compensated or increased by the increase in stability in blood even when receptor affinity is reduced.

[0310]

Example 91 Drug effect test in gastrointestinal obstruction model

As shown in Examples 89 and 90, the glycosylated polypeptides of the present invention proved to have effective action as agonists even in rats *in vivo*. Next,

in order to prove that they also show efficacy in disease models, an evaluation in rat gastrointestinal obstruction model was carried out. In gastrointestinal obstructions such as ileus, gastrointestinal symptoms such as sense of abdominal fullness, vomiting, and abdominal pain are shown by gastrointestinal tract tissue disorder and reduction in absorption ability of e.g. water or electrolyte. Their pathologies are known to be caused by obstruction of gastrointestinal content or release of digestive juice or biologically active material into the gastrointestinal tract. Somatostatin show the effect of decreasing the gastrointestinal content by secretory suppression of various digestive juices or promotion of water and electrolytes absorption via SSTTR expressed in the gastrointestinal system, and is thought to be effective for improvement of the symptoms. This experimental line was carried out as a test line to evaluate the promotion of intestinal fluid absorption or the secretory suppressing action, using the change in the intestinal fluid weight in the jejunum after bowel obstruction as an indicator. Moreover, blood parameters of deviation enzymes amylase (pancreas), lactate dehydrogenase (LDH, liver), and creatine phosphokinase (CPK, skeletal muscle, cardiac muscle etc.) were measured as indicators of tissue disorders.

[0311]

Example 91-1 Production of gastrointestinal obstruction model

The present test was consigned to Mitsubishi Chemical Medience Corporation and carried out. Male SD rats (CrI: CD (SD), Charles River Japan, 8 weeks-old, n = 5 or more, body weight 251.1 - 278.1 g) were fasted for 12 hours or more. Anesthesia was introduced with inhalation of 2% isoflurane and laughing gas:oxygen = 7:3, and this was maintained throughout the surgery. A median incision was made in the abdomen, and the jejunum at about 10cm from the musculus suspensorius duodeni was ligated with a surgical suture. Then, the incision site was promptly sutured, and the rats were fasted until compound administration. In the sham treatment group, ligation of the jejunum was not performed, but the treatment of suturing the incision site after a median incision was made in the abdomen was performed.

[0312]

Example 91-2 Preparation of compound and administration

Employing S1C(disialo)-SRIF28, C(disialo)-R-K-SRIF14, and S1-2C(asialo)-SRIF28 as the glycosylated forms, these were dissolved in Japanese Pharmacopeia saline (from Otsuka Pharmaceutical factory, Inc.) to prepare a 40 μ M solution as the administration solution. Eighteen hours after the gastrointestinal obstruction surgery, this was subcutaneously administered into the dorsal cervix at a dosage of 1 mL/kg with a glass syringe and a 25 G injection needle (all from Terumo Corporation). As the vehicle group, saline without glycosylated polypeptide was similarly administered. Moreover, octreotide was administered as the control group.

[0313]

Example 91-3 Measurement of intestinal fluid weight

One hour after compound administration, a median incision was made in the abdomen under inhalation anesthesia, and 1.5 mL of blood was collected from the abdominal vena cava. Then, the ligated jejunum on the musculus suspensorius duodeni side was resected. The fluid and blood on the jejunum surface were removed with a paper towel, nerves, blood vessels, and fat attached to the jejunum was removed, this was cut into 6 cm lengths, and wet weight was measured. This was then dried at 36 degrees for 24 hours, and dry weight was measured. The intestinal fluid weight (mg) was calculated by wet weight - dry weight. The intestinal fluid weight of the jejunum obtained is shown in Table 9.

[Table 24]

Table 9

	Intestinal fluid weight(mg)
Sham treatment	285 $\pm$ 27
vehicle	481 $\pm$ 31
octreotide	693 $\pm$ 48
S1C(disialo)-SRIF28	630 $\pm$ 83
C(disialo)-R-K-SRIF14	584 $\pm$ 95
S1-2C(asialo)-SRIF28	566 $\pm$ 34

[0314]

As apparent from Table 9, the vehicle had significantly increased intestinal fluid weight compared to

the sham treatment, and it was recognized that enhancement of intestinal fluid secretion accompanying gastrointestinal obstruction due to ligation was caused. Somatostatin or octreotide are shown to have the secretory suppression effect of the intestinal fluid into the intestinal tract and the promotional effect of intestinal fluid absorption into the bowel tissue side in such gastrointestinal obstruction models (Scand. J. Gastroenterol. 1995 May; 30 (5): 464-9), and octreotide was confirmed to have this effect also in this experimental line. S1C(disialo)-SRIF28, C(disialo)-R-K-SRIF14, and S1-2C(asialo)-SRIF28 were all recognized to have increase in the intestinal fluid weight compared to the vehicle, and it became clear that the glycosylated forms of the present invention also show efficacy such as secretory suppression of the intestinal fluid and promotion of water and electrolyte absorption. Moreover, it has become clear that in Examples 86 and 87, the AUC for subcutaneous administration of S1-2C(asialo)-SRIF28 was 1/2 compared to S1C(disialo)-SRIF28, but in Example 69-1, the affinity towards SSTR1 - SSTR5 is approximately 1.7 - 2.9-folds higher than S1C(disialo)-SRIF28. This is thought to be the reason why improvement of receptor affinity causes the drug effects in the present model to be similar even when plasma concentration is low. Similarly, this is thought to be the reason why drug effects are similar because receptor affinity is approximately 0.7 - 2.4-folds higher even though C(disialo)-R-K-SRIF14 has a slightly low AUC for subcutaneous administration compared to that of S1C(disialo)-SRIF28.

[0315]

Example 91-4 Measurement of blood parameter

Employing the blood collected in Example 91-3, amylase (IU/L), LDH (IU/L) and CPK concentrations (IU/L) were measured with an autoanalyzer 7170 (Hitachi, Ltd.). The measuring methods employed were BG5B, UV-rate, and JSCC methods, respectively. The results obtained are shown in Table 10.

[Table 25]

Table 10

	Amylase (IU/L)	LDH (IU/L)	CPK (IU/L)
Sham treatment	669 ± 164	168 ± 82	353 ± 68
vehicle	1672 ± 743	341 ± 68	377 ± 57
octreotide	1732 ± 774	269 ± 141	472 ± 215
S1C(disialo)-SRIF28	1164 ± 224	338 ± 97	455 ± 202
C(disialo)-R-K-SRIF14	985 ± 238	228 ± 155	415 ± 137
S1-2C(asialo)-SRIF28	1594 ± 573	318 ± 44	728 ± 665

[0316]

As apparent from Table 10, the vehicle had increased amylase activity and LDH activity compared to the sham treatment, and it was speculated that tissue disorder of the gastrointestinal system had developed accompanying gastrointestinal obstruction. In S1C(disialo)-SRIF28 and C(disialo)-R-K-SRIF14, the amylase activity had a low value compared to the vehicle. On the other hand, the amylase activity of octreotide was equivalent to the vehicle. As apparent from Examples 69-1 and 69-2, S1C(disialo)-SRIF28, C(disialo)-R-K-SRIF14, and S1-2C(asialo)-SRIF28 had binding affinity toward all the receptors from SSTR1 - SSTR5, whereas octreotide is a compound having specific affinity towards SSTR2, SSTR3, and SSTR5. In rat pancreas, since there is a report that all the receptors SSTR1 - SSTR5 are expressed (J. Histochem. Cytochem. 2004 Mar; 52 (3): 391-400), the possibility that the glycosylated form alleviated the tissue disorder by acting on a receptor different from that of octreotide was conceived. Moreover, in octreotide and C(disialo)-R-K-SRIF14, the LDH activity had a low value compared to the vehicle. There were no others that reduced each parameter by administration of the glycosylated form.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A glycosylated polypeptide selected from the group consisting of:

(A) SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1;

(B) a polypeptide having one or two amino acids deleted, substituted, or added from/to SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1; and

(C) a polypeptide having 80% or more identity with SRIF14 consisting of the amino acid sequence represented by SEQ ID NO. 1;

said glycosylated polypeptide further comprising (i) N amino acids at the N-terminal wherein N is an integer from 1 to 20; and/or (ii) M amino acids at the C-terminal wherein M is an integer from 1 to 6;

wherein two to ten amino acids are substituted with glycosylated amino acids;

at least one of said two to ten amino acids is present in any of said N amino acids or in any of said M amino acids;

the polypeptide binds to a somatostatin receptor; and

each of said glycosylated amino acids is glycosylated Asn or glycosylated Cys.

2. The glycosylated polypeptide according to claim 1, wherein the sequence of said N amino acids added onto the N-terminal is represented by X-Y-,

wherein X represents a sequence of any L amino acids wherein L is an integer from 1 to 6, and Y is a sequence selected from the group consisting of:

- (1) Lys,
- (2) Arg-Lys,
- (3) Glu-Arg-Lys,
- (4) Arg-Glu-Arg-Lys,
- (5) Pro-Arg-Glu-Arg-Lys,
- (6) Ala-Pro-Arg-Glu-Arg-Lys,
- (7) Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (8) Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (9) Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (10) Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (11) Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (12) Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (13) Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys,
- (14) Ser-Ala-Asn-Ser-Asn-Pro-Ala-Met-Ala-Pro-Arg-Glu-Arg-Lys, and
- (15) a sequence having one or two amino acids deleted, substituted, or added from/to the above sequences as defined in parts (2) - (14).

3. The glycosylated polypeptide according to claim 2, wherein at least one of said two to ten amino acids substituted with said glycosylated amino acids is present in said L amino acids.

4. The glycosylated polypeptide according to claim 1, wherein said N amino acids are added onto the N-terminal via a linker.

5. A glycosylated polypeptide selected from the group consisting of:

(A) SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2;

(B) a polypeptide having one to five amino acids deleted, substituted, or added from/to SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2;

(C) a polypeptide having 80% or more identity with SRIF28 consisting of the amino acid sequence represented by SEQ ID NO. 2;

(D) a polypeptide further comprising J amino acids at the N-terminal of (A) - (C) wherein J is an integer from 1 to 15; and

(E) a polypeptide further comprising K amino acids at the C-terminal of (A) - (C) wherein K is an integer from 1 to 6;

wherein two to ten amino acids are substituted with glycosylated amino acids, and

the glycosylated polypeptide binds to a somatostatin receptor.

6. The glycosylated polypeptide according to claim 5, wherein at least one of said two to ten amino acids substituted with said glycosylated amino acids is an amino acid selected from the group consisting of the amino acid corresponding to position 1, the amino acid corresponding to position 5, the amino acid corresponding to position 9, the amino acid corresponding to position 12, the amino acid corresponding to position 13, the amino acid corresponding to

position 14, and the amino acid corresponding to position 19 of SRIF28.

7. The glycosylated polypeptide according to claim 5, wherein said two to ten amino acids are substituted with glycosylated amino acids in said polypeptide as defined in part(D),

wherein at least one of the amino acids substituted with said glycosylated amino acid is present in said J amino acids at the N-terminal side of said polypeptide as defined in part (D).

8. The glycosylated polypeptide according to claim 5, wherein at least one of said two to ten amino acids substituted with said glycosylated amino acids is present in said K amino acids.

9. The glycosylated polypeptide according to any one of claims 1 to 8, wherein said polypeptide has increased stability in blood compared to SRIF28.

10. The glycosylated polypeptide according to any one of claims 1 to 9, wherein said polypeptide has increased half-life in blood by 10-fold or more compared to SRIF28.

11. The glycosylated polypeptide according to any one of claims 1 to 10, wherein said polypeptide has an ability of binding to at least two or more somatostatin receptors selected from the group consisting of SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5.

12. The glycosylated polypeptide according to any one of claims 1 to 11, wherein said polypeptide has an ability of binding to at least one of SSTR1 and SSTR4.

13. The glycosylated polypeptide according to any one of claims 1 to 11, wherein said polypeptide has an ability of binding to both SSTR1 and SSTR4.

14. The glycosylated polypeptide according to any one of claims 1 to 11, wherein said polypeptide has an ability of binding to all of SSTR1, SSTR2, SSTR3, SSTR4, and SSTR5.

15. The glycosylated polypeptide according to any one of claims 1 to 14, wherein each of said glycosylated amino acids is glycosylated Asn or glycosylated Cys.

16. The glycosylated polypeptide according to any one of claims 1 to 15, wherein a sugar chain and an amino acid present in each of said glycosylated amino acids are linked without a linker.

17. The glycosylated polypeptide according to claim 16, wherein that in each of said glycosylated amino acids, the sugar chain consists of 4 or more sugars.


18. The glycosylated polypeptide according to claim 16 or 17, wherein in each of said glycosylated amino acids, the sugar chain is a biantennary complex-type sugar chain, a

triantennary complex-type sugar chain, or a tetraantennary complex-type sugar chain.


19. The glycosylated polypeptide according to claim 16 or 17, wherein in each of said glycosylated amino acids, the sugar chain is a biantennary complex-type sugar chain.

20. The glycosylated polypeptide according to claim 18 or 19, wherein said biantennary complex-type sugar chain is a sugar chain selected from the group consisting of a disialo sugar chain, a monosialo sugar chain, an asialo sugar chain, a diGlcNAc sugar chain, and a dimannose sugar chain.

21. The glycosylated polypeptide according to claim 16, wherein in each of said glycosylated amino acids, the sugar chain is a sugar chain represented by the following Formula 1:



wherein R¹ and R² are identical or different and are represented by:



wherein Ac is an acetyl group.

22. The glycosylated polypeptide according to claim 16 or 17, wherein said sugar chain has at least one sialic acid at the non-reducing terminal, and the carboxy group of said sialic acid is modified by an alkylamino group, a benzyl group, an amino group, or an aminoethylamino group, having 1 - 30 carbons.

23. The glycosylated polypeptide according to any one of claims 1 to 22, wherein said polypeptide has two to ten glycosylated amino acids, and
the sugar chains on each of said glycosylated amino acids are all identical.

24. The glycosylated polypeptide according to any one of claims 1 to 23, wherein said polypeptide comprises Cys residues at positions 17 and 28 of SRIF28 which forms a disulfide bond therebetween.

25. The glycosylated polypeptide according to any one of claims 1 to 24, wherein the C-terminal of said glycosylated polypeptide is amidated.

26. The glycosylated polypeptide according to any one of claims 1 to 24, wherein an azido group is introduced at the N-terminal of said glycosylated polypeptide.

27. The glycosylated polypeptide according to any one of claims 1 to 26, wherein said glycosylated polypeptide is labeled.

28. A pharmaceutical composition comprising:

(I) a glycosylated polypeptide as defined in any one of claims 1 to 26 and/or a pharmaceutically acceptable salt thereof, and

(II) a pharmaceutically acceptable carrier.

29. The pharmaceutical composition according to claim 28, wherein sugar chains in said glycosylated polypeptide are 90% or more identical in respect of the type of sugar constituting the sugar chain, the binding order, and the binding mode.

30. The pharmaceutical composition according to claim 28 or 29, for use in the treatment or prevention of a somatostatin-related disease, wherein said somatostatin-related disease is at least one disease selected from the group consisting of acromegaly, gigantism, dementia, cancer, hormone-producing tumor, endocrine tumor, carcinoid, VIPoma, insulinoma, glucagonoma, diabetes, arthritis, diarrhea, gastric ulcer, inflammatory bowel disease, irritable bowel syndrome, Cushing's disease, hormone secretion defect, gastrointestinal obstruction, ileus, postoperative restenosis, radiation damage,

eye disease, dry eye, glaucoma, interstitial keratitis, iritis, cataract, and conjunctivitis.

31. The pharmaceutical composition according to claim 30, wherein said dementia is Alzheimer's disease.

32. Use of a glycosylated polypeptide as defined in any one of claims 1 to 26 for treating or preventing a somatostatin-related disease, wherein said somatostatin-related disease is at least one disease selected from the group consisting of acromegaly, gigantism, dementia, cancer, hormone-producing tumor, endocrine tumor, carcinoid, VIPoma, insulinoma, glucagonoma, diabetes, arthritis, diarrhea, gastric ulcer, inflammatory bowel disease, irritable bowel syndrome, Cushing's disease, hormone secretion defect, gastrointestinal obstruction, ileus, postoperative restenosis, radiation damage, eye disease, dry eye, glaucoma, interstitial keratitis, iritis, cataract, and conjunctivitis.

33. The use according to claim 32, wherein said dementia is Alzheimer's disease.

34. A glycosylated polypeptide selected from the group consisting of:

(A) a SRIF14 consisting of the amino acid sequence of SEQ ID NO:1;

(B) a polypeptide having one or two amino acids deleted, substituted, or added from/to SRIF14 consisting of the amino acid sequence of SEQ ID NO:1;

35. A glycosylated polypeptide selected from the group consisting of:

(A) a SRIF14 consisting of the amino acid sequence of SEQ ID NO:1;

(B) a polypeptide having one or two amino acids deleted, substituted, or added from/to SRIF14 consisting of the amino acid sequence of SEQ ID NO:1;

(C) a polypeptide having 80% or more identity with SRIF14 consisting of the amino acid sequence of SEQ ID NO:1;

(D) a polypeptide further comprising N amino acids at the N-terminal of (A) - (C), wherein N is an integer from 1 to 20; and

(E) a polypeptide further comprising M amino acids at the C-terminal of (A) - (C), wherein M is an integer from 1 to 6;

wherein two to ten amino acids are substituted with a glycosylated amino acid, and the polypeptide binds to a somatostatin receptor, and

wherein said glycosylated polypeptide comprises a sugar chain comprising at least one sialic acid at the non-reducing terminus, and said sialic acid comprises a carboxy group that is modified by an alkylamino group, a benzyl group, an amino group, or an aminoethylamino group, said alkylamino group, benzyl group, amino group, and aminoethylamino group having 1 - 30 carbons.

36. A glycosylated polypeptide selected from the group consisting of:

(A) a SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;

(B) a polypeptide having one to three amino acids deleted, substituted, or added from/to SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;


(C) a polypeptide having 80% or more identity with SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;

(D) a polypeptide further comprising J amino acids at the N-terminal of (A) - (C), wherein J is an integer from 1 to 15; and


(E) a polypeptide further comprising K amino acids at the C-terminal of (A) - (C), wherein K is an integer from 1 to 6;

wherein two to ten amino acids are substituted with glycosylated amino acids, and the polypeptide binds to a somatostatin receptor,

wherein each of said glycosylated amino acids comprises a sugar chain represented by Formula 1:



wherein R¹ and R² are identical or different and are represented by:



wherein Ac is an acetyl group.

37. A glycosylated polypeptide selected from the group consisting of:

(A) a SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;

(B) a polypeptide having one to three amino acids deleted, substituted, or added from/to SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;

(C) a polypeptide having 80% or more identity with SRIF28 consisting of the amino acid sequence of SEQ ID NO:2;

(D) a polypeptide further comprising J amino acids at the N-terminal of (A) - (C), wherein J is an integer from 1 to 15; and

(E) a polypeptide further comprising K amino acids at the C-terminal of (A) - (C), wherein K is an integer from 1 to 6;

wherein two to ten amino acids are substituted with glycosylated amino acids, and the polypeptide binds to a somatostatin receptor,

wherein said glycosylated polypeptide comprises a sugar chain comprising at least one sialic acid at the non-reducing terminus, and said sialic acid comprises a carboxy group that is modified by an alkylamino group, a benzyl group, an amino

group, or an aminoethylamino group, said alkylamino group, benzyl group, amino group, and aminoethylamino group having 1 - 30 carbons.

38. A glycosylated polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NOs: 21-26, 29, 30, 33, 34, 88, 89, 125, 127, 129, 148, 149, 157, 164 and 169.

1/14

Compound Name	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
C(dialo)-R-K-SRIF 14	C R K A G C K N F F W K T F T S C Dialo S-S																											
C(dialo)-C12inter-SRIF 14	C C12 A G C K N F F W K T F T S C Dialo S-S																											
S1-2C(dialo)-SRIF 28	C C A N S N P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1C(dialo)-NSC(dialo)-SRIF 28	C A N S C P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1C(dialo) - R13C(dialo)-SRIF 28	C A N S N P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											
NSC(dialo) - A9C(dialo)-SRIF 28	S A N S C P A M C P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1-3C(dialo)-SRIF 28	C C C A N S N P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1C(dialo) - NSC(dialo) - A9C(dialo)-SRIF 28	C A N S C P A M C P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1C(monosialo)-SRIF 28	C A N S N P A M A P R E R K A G C K N F F W K T F T S C Monosialo S-S																											
S1C(asialo)-SRIF 28	C A N S N P A M A P R E R K A G C K N F F W K T F T S C asialo S-S																											
S1-2C(asialo)-SRIF 28	C C A N S N P A M A P R E R K A G C K N F F W K T F T S C Asialo S-S																											
S1-3C(asialo)-SRIF 28	C C C A N S N P A M A P R E R K A G C K N F F W K T F T S C sub. with asialo S-S																											
NSN(dialo)-SRIF 28	S A N S N P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											
S1M(dialo)-SRIF 28	N A N S N P A M A P R E R K A G C K N F F W K T F T S C Dialo S-S																											

[Figure 1C]


Compound name	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
C(disialo)-SRIF28	C	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C	
	Disialo																													
R11C(disialo)-SRIF28	S	A	N	S	N	P	A	M	A	P	C	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	Disialo																													
F20C(disialo)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	C	F	W	K	T	F	T	S	C		
	Disialo																													
T24C(disialo)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	C	F	T	S	C		
	Disialo																													
F25C(disialo)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	C	T	S	C		
	Disialo																													
S27C(disialo)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	Disialo																													
C(disialo)-K-SRIF14	C K A G C K N F F W K T F T S C																													
	Disialo																													
S1C(disialo)-F25Y-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	Y	T	S	C		
	Disialo																													
S1C(disialo)-SRIF28- amide	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	Disialo																													
C(disialo)-PEGinker-SRIF14	C PEG A G C K N F F W K T F T S C																													
	Disialo																													
Biotin-S1C(disialo)-SRIF28	Biotin	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C	
	Disialo																													
Biotin-PEGinker-S1C(disialo)-SRIF28	Biotin	PEG	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C
	Disialo																													
Azido-S1C(disialo)-SRIF28	Azido	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C	
	Disialo																													
S1C(disialo)-E12C(disialo)-SRIF28	C	A	N	S	N	P	A	M	A	P	R	C	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	Disialo																													
2C(disialo)-R-K-SRIF14	C C R K A G C K N F F W K T F T S C																													
	Disialo																													
3C(disialo)-R-K-SRIF14	C C C R K A G C K N F F W K T F T S C																													
	Disialo																													
S1C(diglycyl)-SRIF28	C A N S N P A M A P R E R K A G C K N F F W K T F T S C																													
	Disialo																													

[Figure 1D]


Compound name	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
S1C(difMan)-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	difMan																														
N19C(difMan)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	C	F	F	W	K	T	F	T	S	C			
	S-S																														
S1C(GlcNAc)-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	GlcNAc																														
N19C(GlcNAc)-SRIF28	S	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	C	F	F	W	K	T	F	T	S	C			
	S-S																														
S1C(trisialo)-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	Trisialo																														
S1C(tetrasialo)-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	Tetrasialo																														
S1C(disialo(aminoethylamide))-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	disialo(aminoethylamide)																														
S1C(disialo(amide))-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	disialo(amide)																														
S1C(disialo(Bn))-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	disialo(Bn)																														
S1C(disialo(hexadecylamide))-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	disialo(hexadecylamide)																														
S1-2C(disialo(amide))-SRIF28	C	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	disialo(amide)																														
S1-2C(disialo(Bn))-SRIF28	C	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	disialo(Bn)																														
S1C(Asn(disialo))-SRIF28	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C			
	Asn(disialo)																														
S1N(disialo)-N19C(difMan)-SRIF28	N	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	C	F	F	W	K	T	F	T	S	C			
	Disialo																														
C(disialo(aminoethylamide))-S1C(disialo)-SRIF28	C	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C		
	disialo(aminoethylamide) disialo																														
S1-4C(disialo)-SRIF28	C	C	C	C	A	N	S	N	P	A	M	A	P	R	E	R	K	A	G	C	K	N	F	F	W	K	T	F	T	S	C
	S-S																														
	Disialo																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														
	S-S																														

Revised: 04/06/10, 04/14/10


[Figure 1E]




[Figure 2]




[Figure 3]




[Figure 4]




[Figure 5]




[Figure 6]




[Figure 7]




[Figure 8]




[Figure 9]




[Figure 10]




[Figure 11]




[Figure 12]




[Figure 13]




[Figure 14]




[Figure 15]




[Figure 16]




[Figure 17]



[Figure 18]



[Figure 19]



[Figure 20]

