



(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/GB2014/053694

(22) International Filing Date:
12 December 2014 (12.12.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1322034.8 12 December 2013 (12.12.2013) GB

(71) Applicant: **ALMAC DIAGNOSTICS LIMITED**
[GB/GB]; Almac House, 20 Seagoe Industrial Estate,
Craigavon BT63 5QD (GB).

(72) Inventors: **WALKER, Steven**; c/o Almac House, 20
Seagoe Industrial Estate, Craigavon BT63 5QD (GB). **MC-
CAVIGAN, Andrena**; c/o Almac House, 20 Seagoe In-
dustrial Estate, Craigavon BT63 5QD (GB). **DAVISON,
Timothy**; c/o Almac House, 20 Seagoe Industrial Estate,
Craigavon BT63 5QD (GB). **KENNEDY, Richard**; c/o
Almac House, 20 Seagoe Industrial Estate, Craigavon
BT63 5QD (GB). **HARKIN, Paul**; c/o Almac House, 20
Seagoe Industrial Estate, Craigavon BT63 5QD (GB).
HILL, Laura; c/o Almac House, 20 Seagoe Industrial Es-
tate, Craigavon BT63 5QD (GB).

(74) Agent: **BOULT WADE TENNANT**; Verulam Gardens,
70 Grays Inn Road, London, Greater London WC1X 8BT
(GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:
17 September 2015

(54) Title: PROSTATE CANCER CLASSIFICATION

(57) Abstract: A method is provided for characterising and/or prognosing prostate cancer in a subject comprising determining the expression level of at least one of CREM, ERFFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMT-S9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1 R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MAL-AT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in a sample from the subject. The method may be used to predict the likelihood of metastasis. Also disclosed are methods for diagnosing and selecting treatment for prostate cancer, together with corresponding meth-ods of treatment. Systems, kits and computer programs for performing the methods are also provided.



- 1 -

PROSTATE CANCER CLASSIFICATION**5 FIELD OF THE INVENTION**

The present invention relates to prostate cancer. Provided are methods for characterising and prognosing prostate cancer which rely upon biomarkers. Antibodies, kits and systems useful in the methods are also described.

10 BACKGROUND OF THE INVENTION

Prostate cancer is the most common malignancy in men with a lifetime incidence of 15.3% (Howlader 2012). Based upon data from 1999-2006 approximately 80% of prostate cancer patients present with early disease clinically confined to the prostate (Altekruse et al 2010) of which around 65% are cured by surgical resection or radiotherapy (Kattan et al 1999, Pound et al 15 1999). 35% will develop PSA recurrence of which approximately 35% will develop local or metastatic recurrence, which is non-curable. At present it is unclear which patients with early prostate cancer are likely to develop recurrence and may benefit from more intensive therapies. Current prognostic factors such as tumour grade as measured by Gleason score have prognostic value but a significant number of those considered lower grade (7 or less) still recur and a proportion of higher-grade tumours do not. Additionally there is significant heterogeneity in the prognosis of Gleason 7 tumours (Makarov et al 2002, Rasiah et al 2003). Furthermore it has become evident that the grading of Gleason score has changed leading to changes in the distribution of Gleason scores over time (Albertsen et al 2005, Smith et al 2002).

25 It is now clear that most solid tumours originating from the same anatomical site represent a number of distinct entities at a molecular level (Perou et al 2000). DNA microarray platforms allow the analysis of tens of thousands of transcripts simultaneously from archived paraffin embedded tissues and are ideally suited for the identification of molecular subgroups. This kind of approach has identified primary cancers with metastatic potential in solid tumours such as 30 breast (van 't Veer et al 2002) and colon cancer (Bertucci et al 2004).

DESCRIPTION OF THE INVENTION

The present invention is based upon the identification and verification of prostate cancer biomarkers.

35

The present inventors have identified a group of primary prostate cancers that are similar to metastatic disease at a molecular level. These tumours are defined by loss of expression of

- 2 -

several genes and defined pathways; furthermore this group is defined by activation of the proto-oncogene FOXM1 that leads to increased expression of genes involved in mitosis. A series of biomarkers that can identify tumours within this subgroup have been defined which have multivariate prognostic power and can be used to prospectively assess if a tumour is at increased likelihood of recurrence and/or metastatic development.

Thus, in a first aspect the invention provides a method for characterising and/or prognosing prostate cancer in a subject comprising:

determining the expression level of at least one of

- 10 FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36
- 15 in a sample from the subject wherein the determined expression level is used to provide a characterisation of and/or a prognosis for the prostate cancer.

According to all aspects of the invention the prostate cancer may be a primary prostate cancer.

- 20 According to a further aspect of the invention there is provided a method for diagnosing a prostate cancer with an increased metastatic potential in a subject comprising:
determining the expression level of at least one of
FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4,
25 LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36
in a sample from the subject wherein the determined expression level is used to identify whether a subject has a prostate cancer with increased metastatic potential.

- 30 In yet a further aspect, the present invention relates to a method for diagnosing a prostate cancer with an increased metastatic potential in a subject comprising:
determining the expression level of at least one of
TRPM3, PDRG1, SRSF5, PDE4D, F12 and PDK4
35 in a sample from the subject wherein the determined expression level is used to identify whether a subject has a prostate cancer with increased metastatic potential.

- 3 -

The invention also relates to a method for characterising and/or prognosing prostate cancer in a subject comprising:

determining the expression level of at least one of

FOX M1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38,
 5 CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4,
 LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2,
 RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1,
 XBP1 and ZFP36

10 in a sample from the subject in order to identify the presence or absence of cells characteristic of
 an increased likelihood of recurrence and/or metastasis wherein the determined presence or
 absence of the cells is used to provide a characterisation of and/or a prognosis for the prostate
 cancer.

In a further aspect, the present invention relates to a method for characterising and/or
 15 prognosing prostate cancer in a subject comprising:

a) obtaining a sample from the subject

b) applying an antibody specific for the protein product of at least one of

FOX M1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38,
 CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4,
 20 LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2,
 RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1,
 XBP1 and ZFP36

to the sample from the subject

c) applying a detection agent that detects the antibody-protein complex

25 d) using the detection agent to determine the level of the protein

d) wherein the determined level of the protein is used to provide a characterisation of and/or a
 prognosis for the prostate cancer.

30 The characterization, prognosis or diagnosis of the prostate cancer can also be used to guide
 treatment.

Accordingly, in a further aspect, the present invention relates to a method for selecting a
 treatment for prostate cancer in a subject comprising:

(a) determining the expression level of at least one of

35 FOX M1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38,
 CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4,
 LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2,

- 4 -

RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

in a sample from the subject wherein the determined expression level is used to provide a characterisation of and/or a prognosis for the prostate cancer and

- 5 (b) selecting a treatment appropriate to the characterisation of and/or prognosis for the prostate cancer.

In yet a further aspect, the present invention relates to a method for selecting a treatment for prostate cancer in a subject comprising:

- 10 (a) determining the expression level of at least one of
FOX M1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1,
15 XBP1 and ZFP36

in a sample from the subject wherein the determined expression level is used to provide a characterisation of and/or a prognosis for the prostate cancer

(b) selecting a treatment appropriate to the characterisation of and/or prognosis for the prostate cancer and

- 20 (c) treating the subject with the selected treatment.

The invention also relates to a method of treating prostate cancer comprising administering a chemotherapeutic agent or radiotherapy, optionally extended radiotherapy, preferably extended-field radiotherapy, to a subject or carrying out surgery on a subject wherein the subject is
25 selected for treatment on the basis of a method as described herein.

In a further aspect, the present invention relates to a chemotherapeutic agent for use in treating prostate cancer in a subject, wherein the subject is selected for treatment on the basis of a method as described herein.

30

In yet a further aspect, the present invention relates to method of treating prostate cancer comprising administering a chemotherapeutic agent or radiotherapy, optionally extended radiotherapy, preferably extended-field radiotherapy to a subject or carrying out surgery on a subject wherein the subject has an increased expression level of at least one of HJURP, PDRG1,
35 TRPM3, F12, CENPF, RNFT2, and SSTR1 and/or a decreased expression level of at least one of CREM, ERFFI1, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1,

- 5 -

MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The invention also relates to a chemotherapeutic agent for use in treating prostate cancer in a subject, wherein the subject has an increased expression level of at least one of HJURP, PDRG1, TRPM3, F12, CENPF, RNFT2, and SSTR1 and/or a decreased expression level of at least one of CREM, ERRF1, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

In certain embodiments the chemotherapeutic agent comprises, consists essentially of or consists of

- a) an anti-hormone treatment, preferably bicalutamide and/or abiraterone
- b) a cytotoxic agent
- c) a biologic, preferably an antibody and/or a vaccine, more preferably Sipuleucel-T and/or
- d) a targeted therapeutic agent

Suitable therapies and therapeutic agents are discussed in further detail herein.

The genes FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERRF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 and their protein products are described and defined in further detail in Table A below. The genes may also be referred to, interchangeably, as biomarkers.

Table A

Gene	Previous names and symbols and Synonyms (HGNC database)	Uniprot number	Protein
FOXM1	FKHL16, HFH-11, HNF-3, INS-1, "M-phase phosphoprotein 2", MPHOSPH2, MPP2, TGT3, trident	Q08050	forkhead box M1

- 6 -

TRPM3	GON-2, KIAA1616, LTRPC3, "melastatin 2"	Q9HCF6	Transient receptor potential cation channel subfamily M member 3
PDRG1	C20orf126, "chromosome 20 open reading frame 126", dJ310O13.3	Q9NUG6	p53 and DNA damage-regulated protein 1
SRSF5	SFRS5, "splicing factor, arginine/serine-rich 5", HRS, "SR splicing factor 5", SRP40	Q13243	Serine/arginine-rich splicing factor 5
PDE4D	DPDE3, "phosphodiesterase 4D, cAMP-specific (dunce (Drosophila)-homolog phosphodiesterase E3)"	Q08499	cAMP-specific 3',5'-cyclic phosphodiesterase 4D
F12	N/A	P00748	Coagulation factor XII
PK4	"pyruvate dehydrogenase kinase, isoenzyme 4"	Q16654	[Pyruvate dehydrogenase [lipoamide]] kinase isozyme 4, mitochondrial
ADAMTS1	"a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 1", C3-C5, KIAA1346, METH1	Q9UHI8	A disintegrin and metalloproteinase with thrombospondin motifs 1
ADAMTS9	KIAA1312	Q9P2N4	A disintegrin and metalloproteinase with thrombospondin motifs 9
B3GNT5	B3GN-T5, beta3Gn-T5, "lactosylceramide 1,3-N-acetyl-beta-D-glucosaminyltransferase"	Q9BYG0	Lactosylceramide 1,3-N-acetyl-beta-D-glucosaminyltransferase
CD38	"CD38 antigen (p45)", "ADP-ribosyl cyclase 1", "NAD(+) nucleosidase"	P28907	ADP-ribosyl cyclase 1
CEBPD	C/EBP-delta, CELF, CRP3, NF-IL6-beta	P49716	CCAAT/enhancer-binding protein delta
CENPF	"centromere protein F, 350/400kDa (mitosin)", hcp-1, "mitosin"	P49454	Centromere protein F

- 7 -

CREM	hCREM-2	Q03060	cAMP-responsive element modulator
DKK1	"dickkopf (Xenopus laevis) homolog 1", "dickkopf 1 homolog (Xenopus laevis)", DKK-1, SK	O94907	Dickkopf-related protein 1
EMP1	CL-20, TMP	P54849	Epithelial membrane protein 1
ERRFI1	GENE-33, MIG-6, RALT	Q9UJM3	ERBB receptor feedback inhibitor 1
F3	CD142	P13726	Tissue factor
HJURP	DKFZp762E1312, FAKTS, hFLEG1, URLC9	Q8NCD3	Holliday junction recognition protein
IL1R1	IL1R, IL1RA, CD121A, D2S1473	P14778	Interleukin-1 receptor type 1
IL8	3-10C, "alveolar macrophage chemotactic factor I", AMCF-I, b-ENAP, "beta endothelial cell-derived neutrophil activating peptide", "chemokine (C-X-C motif) ligand 8", CXCL8, GCP-1, GCP1, "granulocyte chemotactic protein 1", IL-8, K60, LECT, LUCT, "lung giant cell carcinoma-derived chemotactic protein", "lymphocyte derived neutrophil activating peptide", LYNAP, MDNCF, MONAP, "monocyte-derived neutrophil chemotactic factor", "monocyte-derived neutrophil-activating peptide", NAF, NAP-1, NAP1, "neutrophil-activating peptide 1", SCYB8, TSG-1, "tumor necrosis factor-induced gene 1"	P10145	Interleukin-8
JUNB	N/A	P17275	Transcription factor jun-B
KLF10	"TGFB inducible early growth response", TIEG, EGRA, TIEG1	Q13118	Krueppel-like factor 10
KLF4	EZF, GKLF	O43474	Krueppel-like factor 4

- 8 -

LDLR	"familial hypercholesterolemia", LDLCQ2	P01130	Low-density lipoprotein receptor
LGALS3	LGALS2, "galectin 3", GALIG, MAC-2	P17931	Galectin-3
LPAR1	EDG2, "endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2", edg-2, Gpcr26, GPR26, LPA1, Mrec1.3, rec.1.3, vzg-1	Q92633	Lysophosphatidic acid receptor 1
MALAT1	HCN, "hepcarcin", LINC00047, "long intergenic non-protein coding RNA 47", MALAT-1, "metastasis associated in lung adenocarcinoma transcript 1", NCRNA00047, NEAT2, "non- protein coding RNA 47", "nuclear enriched abundant transcript 2", "nuclear paraspeckle assembly transcript 2 (non-protein coding)", PRO1073	Q9UHZ2	Metastasis-associated lung adenocarcinoma transcript 1
MTUS1	"mitochondrial tumor suppressor 1", "AT2 receptor-interacting protein", "AT2R binding protein", ATBP, ATIP1, DKFZp586D1519, FLJ14295, ICIS, KIAA1288, "mitochondrial tumor suppressor gene 1", MP44, MTSG1	Q9ULD2	Microtubule-associated tumor suppressor 1
MYBPC1	"myosin-binding protein C, slow- type"	Q00872	Myosin-binding protein C, slow- type
NFIL3	IL3BP1, E4BP4, NF-IL3A, NFIL3A	Q16649	Nuclear factor interleukin-3- regulated protein
NR4A3	CHN, CSMF, MINOR, NOR1	Q92570	Nuclear receptor subfamily 4 group A member 3
OAT	"gyrate atrophy", HOGA, "Ornithine aminotransferase", "ornithine aminotransferase	P04181	Ornithine aminotransferase, mitochondrial

- 9 -

	precursor"		
PI15	"protease inhibitor 15", P25T1	O43692	Peptidase inhibitor 15
PTGS2	COX2	P35354	Prostaglandin G/H synthase 2
RHOBTB3	KIAA0878	O94955	Rho-related BTB domain-containing protein 3
RIN2	RASSF4	Q8WYP3	Ras and Rab interactor 2
RNFT2	TMEM118, "transmembrane protein 118", FLJ14627	Q96EX2	RING finger and transmembrane domain-containing protein 2
SELE	ELAM, ELAM1, "endothelial adhesion molecule 1", CD62E, ESEL	P16581	E-selectin
SLC15A2	"solute carrier family 15 (H ⁺ /peptide transporter), member 2", PEPT2	Q16348	Solute carrier family 15 member 2
SOCS2	CIS2, Cish2, SOCS-2, SSI-2, SSI2, "STAT-induced STAT inhibitor-2", STATI2	O14508	Suppressor of cytokine signaling 2
SOCS3	CIS3, Cish3, SOCS-3, SSI-3	O14543	Suppressor of cytokine signaling 3
SSTR1	N/A	P30872	Somatostatin receptor type 1
ST6GAL1	"sialyltransferase 1 (beta-galactoside alpha-2,6-sialyltransferase)", SIAT1, "ST6Gal I"	P15907	Beta-galactoside alpha-2,6-sialyltransferase 1
TSC22D1	TGFB114, "transforming growth factor beta 1 induced transcript 4", MGC17597, TSC22	Q15714	TSC22 domain family protein 1
XBP1	XBP2	P17861	X-box-binding protein 1
ZFP36	"zinc finger protein 36, C3H type, homolog (mouse)", G0S24, NUP475, RNF162A, TIS11, tristetraprolin, TTP	P26651	Tristetraprolin

- 10 -

In certain embodiments the expression level of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 or 46 of

TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD,
 5 CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR,
 LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3,
 RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and
 ZFP36

is determined. FOXM1 may be added to the panel in some embodiments.

10

Alternatively, the expression level of at least one of a group of 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 or 46 of

TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD,
 15 CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR,
 LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3,
 RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and
 ZFP36

is determined. FOXM1 may be included in the group in some embodiments.

20

In certain embodiments the expression level of at least one of

TRPM3, PDRG1, SRSF5, PDE4D, PDK4, F12, F3, HJURP, CENPF, MYBPC1, SELE, CEBPD,
 and XBP1

is determined.

25

In certain embodiments the expression level of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 of
 TRPM3, PDRG1, SRSF5, PDE4D, PDK4, F12, F3, HJURP, CENPF, MYBPC1, SELE, CEBPD,
 and XBP1

is determined.

30

By characterisation is meant classification and/or evaluation of the prostate cancer. Prognosis refers to predicting the likely outcome of the prostate cancer for the subject. By diagnosis is meant identifying the presence of prostate cancer.

35

According to all aspects of the invention the characterisation of and/or prognosis for the prostate cancer may comprise, consist essentially of or consist of predicting an increased likelihood of recurrence. The characterisation of and/or prognosis for the prostate cancer may comprise,

- 11 -

consist essentially of or consist of predicting a reduced time to recurrence. Recurrence may be clinical recurrence or biochemical recurrence. By biochemical recurrence is meant a rise in the level of PSA in a subject after treatment for prostate cancer. Biochemical recurrence may indicate that the prostate cancer has not been treated effectively or has recurred.

5

The characterisation of and/or prognosis for the prostate cancer may comprise, consist essentially of or consist of predicting an increased likelihood of metastasis.

Metastasis, or metastatic disease, is the spread of a cancer from one organ or part to another non-adjacent organ or part. The new occurrences of disease thus generated are referred to as metastases.

10

Characterisation of and/or prognosis for the prostate cancer may also comprise, consist essentially of or consist of determining whether the prostate cancer has a poor prognosis. A poor prognosis may be a reduced likelihood of cause-specific, i.e. cancer-specific, or long term survival. Cause- or Cancer-specific survival is a net survival measure representing cancer survival in the absence of other causes of death. Cancer survival may be for 6, 7, 8, 9, 10, 11, 12 months or 1, 2, 3, 4, 5 etc. years. Long-term survival may be survival for 1 year, 5 years, 10 years or 20 years following diagnosis. A prostate cancer with a poor prognosis may be aggressive, fast growing, and/or show resistance to treatment.

15

20 In certain embodiments an increased expression level of at least one of TRPM3, PDRG1, F12, CENPF, HJURP, RNFT2, and SSTR1 or of FOXM1 indicates an increased likelihood of recurrence and/or metastasis and/or a poor prognosis. In further embodiments a decreased expression level of at least one of SRSF5, PDE4D, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CREM, DKK1, EMP1, ERFFI1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36 indicates an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

25

In certain embodiments the methods described herein may comprise determining the expression level of at least one of TRPM3, PDRG1, F12, CENPF, HJURP, RNFT2, and SSTR1 or FOXM1 and at least one of SRSF5, PDE4D, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CREM, DKK1, EMP1, ERFFI1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36. Thus, the methods may rely upon a combination of an up-regulated marker and a down-regulated marker.

30

35

- 12 -

In certain embodiments the methods described herein comprise comparing the expression level to a reference value or to the expression level in one or more control samples or to the expression level in one or more control cells in the same sample. The control cells may be normal (i.e. cells characterised by an independent method as non-cancerous) cells. The one or
5 more control samples may consist of non-cancerous cells or may include a mixture of prostate cancer cells and non-cancerous cells. The expression level may be compared to the expression level of the same gene in one or more control samples or control cells.

The reference value may be a threshold level of expression of at least one gene set by
10 determining the level or levels in a range of samples from subjects with and without prostate cancer. The prostate cancer may be prostate cancer with or without an increased likelihood of recurrence and/or metastasis and/or a poor prognosis. Suitable methods for setting a threshold are well known to those skilled in the art. The threshold may be mathematically derived from a training set of patient data. The score threshold thus separates the test samples according to
15 presence or absence of the particular condition. The interpretation of this quantity, i.e. the cut-off threshold may be derived in a development or training phase from a set of patients with known outcome. The threshold may therefore be fixed prior to performance of the claimed methods from training data by methods known to those skilled in the art.

The reference value may also be a threshold level of expression of at least one gene set by
20 determining the level of expression of the at least one gene in a sample from a subject at a first time point. The determined levels of expression at later time points for the same subject are then compared to the threshold level. Thus, the methods of the invention may be used in order to monitor progress of disease in a subject, namely to provide an ongoing characterization and/or
25 prognosis of disease in the subject. For example, the methods may be used to identify a prostate cancer that has developed into a more aggressive or potentially metastatic form. This may be used to guide treatment decisions as discussed in further detail herein.

For genes whose expression level does not differ between normal cells and cells from a prostate
30 cancer that does not have an increased likelihood of recurrence and/or metastasis and/or a poor prognosis the expression level of the same gene in normal cells in the same sample can be used as a control.

Accordingly, in specific embodiments the expression level of at least one of

35 TRPM3, PDRG1, SRSF5, PDE4D, F12, and PDK4

in the prostate cancer cells in a sample is compared to the expression level of the same gene in the normal cells in the same sample.

- 13 -

In specific embodiments if the determined expression level of at least one of TRPM3, PDRG1, SRSF5, PDE4D, F12, and PDK4

5 is not different in the prostate cancer cells in a sample as compared to the normal cells in the same sample then the prostate cancer does not have an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

10 Different may be statistically significantly different. By statistically significant is meant unlikely to have occurred by chance alone. A suitable statistical assessment may be performed according to any suitable method.

15 In specific embodiments if the gene is TRPM3, PDRG1 or F12 and the expression level is increased in the prostate cancer cells in a sample relative to the normal cells in the same sample then the prostate cancer has an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

20 In specific embodiments if the gene is SRSF5, PDE4D or PDK4 and the expression level is decreased in the prostate cancer cells in a sample relate to the normal cells in the sample then the prostate cancer has an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

25 The methods described herein may further comprise determining the expression level of a reference gene. A reference gene may be required if the target gene expression level differs between normal cells and cells from a prostate cancer that does not have an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

30 In certain embodiments the expression level of at least one of ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 is compared to the expression level of a reference gene.

35 The reference gene may be any gene with minimal expression variance across all prostate cancer samples. Thus, the reference gene may be any gene whose expression level does not vary with likelihood of recurrence and/or metastasis and/or a poor prognosis. The skilled person is well able to identify a suitable reference gene based upon these criteria. In particular, the

- 14 -

reference gene may be TPT1, RPS14 or RPL37A. The expression level of the reference gene may be determined in the same sample as the expression level of at least one of ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERRFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The expression level of the reference gene may be determined in a different sample. The different sample may be a control sample as described above. The expression level of the reference gene may be determined in normal and/or prostate cancer cells in a sample.

The expression level of the at least one gene in the sample from the subject may be analysed using a statistical model. In specific embodiments where the expression level of at least 2 genes is measured the genes may be weighted. As used herein, the term "weight" refers to the relative importance of an item in a statistical calculation. The weight of each gene may be determined on a data set of patient samples using analytical methods known in the art. An overall score may be calculated and used to provide a characterisation of and/or prognosis for the prostate cancer.

Methods for determining the expression levels of the markers are described in greater detail herein. Typically, the methods may involve contacting a sample obtained from a subject with a detection agent, such as primers/probes/antibodies (as discussed in detail herein) specific for the marker and detecting expression products. A comparison is made against expression levels determined in a control sample to provide a characterization and/or a prognosis for the prostate cancer.

According to all aspects of the invention the expression level of the gene or genes may be measured by any suitable method. In certain embodiments the expression level is determined at the level of protein, RNA or epigenetic modification. The epigenetic modification may be DNA methylation.

The expression level may be determined by immunohistochemistry. By Immunohistochemistry is meant the detection of proteins in cells of a tissue sample by using a binding reagent such as an antibody or aptamer that binds specifically to the proteins. Thus, the expression level as determined by immunohistochemistry is a protein level. The sample may be a prostate tissue sample and may comprise prostate cancer (tumour) cells, prostatic intraepithelial neoplasia (PIN) cells, normal prostate epithelium, stroma and, optionally, infiltrating immune cells. In some embodiments the expression level of the at least one gene in the prostate cancer (tumour) cells

- 15 -

in a sample is compared to the expression level of the same gene (and/or a reference gene) in the normal cells in the same sample. In some embodiments the expression level of the at least one gene in the prostate cancer (tumour) cells in a sample is compared to the expression level of the same gene (and/or a reference gene) in the normal cells in a control sample. The normal cells may comprise, consist essentially of or consist of normal (non-cancer) prostate epithelial cells. In certain embodiments the normal cells do not comprise PIN cells and/or stroma cells. In certain embodiments the prostate cancer (tumour) cells do not comprise PIN cells and/or stroma cells. In further embodiments the expression level of the at least one gene in the prostate cancer (tumour) cells in a sample is (additionally) compared to the expression level of a reference gene in the same cells or in the prostate cancer cells in a control sample. The reference gene may be TPT1, RPS14 or RPL37A. In yet further embodiments the expression level of the at least one gene in the prostate cancer (tumour) cells in a sample is scored using a method based on intensity, proportion and/or localisation of expression in the prostate cancer (tumour) cells (without comparison to normal cells). The scoring method may be derived in a development or training phase from a set of patients with known outcome.

Accordingly, in a further aspect, the present invention relates to an antibody or aptamer that binds specifically to a protein product of at least one of

FOX M1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The antibody may be of monoclonal or polyclonal origin. Fragments and derivative antibodies may also be utilised, to include without limitation Fab fragments, ScFv, single domain antibodies, nanoantibodies, heavy chain antibodies, aptamers etc. which retain peptide-specific binding function and these are included in the definition of "antibody". Such antibodies are useful in the methods of the invention. They may be used to measure the level of a particular protein, or in some instances one or more specific isoforms of a protein. The skilled person is well able to identify epitopes that permit specific isoforms to be discriminated from one another.

Methods for generating specific antibodies are known to those skilled in the art. Antibodies may be of human or non-human origin (e.g. rodent, such as rat or mouse) and be humanized etc. according to known techniques (Jones *et al.*, Nature (1986) May 29-Jun. 4;321(6069):522-5; Roguska *et al.*, Protein Engineering, 1996, 9(10):895-904; and Studnicka *et al.*, Humanizing

- 16 -

Mouse Antibody Frameworks While Preserving 3-D Structure. Protein Engineering, 1994, Vol.7, pg 805).

In certain embodiments the expression level is determined using an antibody or aptamer conjugated to a label. By label is meant a component that permits detection, directly or indirectly. For example, the label may be an enzyme, optionally a peroxidase, or a fluorophore.

A label is an example of a detection agent. By detection agent is meant an agent that may be used to assist in the detection of the antibody-protein complex. Where the antibody is conjugated to an enzyme the detection agent may be comprise a chemical composition such that the enzyme catalyses a chemical reaction to produce a detectable product. The products of reactions catalyzed by appropriate enzymes can be, without limitation, fluorescent, luminescent, or radioactive or they may absorb visible or ultraviolet light. Examples of detectors suitable for detecting such detectable labels include, without limitation, x-ray film, radioactivity counters, scintillation counters, spectrophotometers, colorimeters, fluorometers, luminometers, and densitometers. In certain embodiments the detection agent may comprise a secondary antibody. The expression level is then determined using an unlabeled primary antibody that binds to the target protein and a secondary antibody conjugated to a label, wherein the secondary antibody binds to the primary antibody.

The invention also relates to use of an antibody as described above for characterising and/or prognosing a prostate cancer in a subject.

Additional techniques for determining expression level at the level of protein include, for example, Western blot, immunoprecipitation, immunocytochemistry, mass spectrometry, ELISA and others (see ImmunoAssay: A Practical Guide, edited by Brian Law, published by Taylor & Francis, Ltd., 2005 edition). To improve specificity and sensitivity of an assay method based on immunoreactivity, monoclonal antibodies are often used because of their specific epitope recognition. Polyclonal antibodies have also been successfully used in various immunoassays because of their increased affinity for the target as compared to monoclonal antibodies.

Suitable antibodies which may be used in the methods of the invention or included in the kits of the invention are listed in Table B below:

Table B – examples of Antibodies binding to markers of the invention

Gene ID	Antibody Supplier	Antibody Reference
ADAMTS1	Source Bioscience	LS-A1643

- 17 -

	Source Bioscience	LS-A1642
	Source Bioscience	LS-A1649
ADAMTS9	Novus	NBP1-82915
	Novus	NBP1-82916
	Sigma	HPA028577
B3GNT5	Novus	NBP1-88954
CD38	Source Bioscience	LS-A9696
	Source Bioscience	LS-A9697
	Leica Novocastra	CD38-290-L-CE
CEBPD	No suitable antibody	
CENPF	Source Bioscience	LS-B2992
	Source Bioscience	LS-B3157
	Novus	NB500-101
CREM	Sigma	HPA001818
DKK1	Source Bioscience	LS-A2867
	Source Bioscience	LS-A2865
	Source Bioscience	LS-A2868
	Novus	NBP1-95560
EMP1	Sigma	HPA056250
ERRFI1	Novus	NBP1-81835
	Sigma	HPA027206
F12	Source Bioscience	LS-B2649
	Source Bioscience	LS-B3044
	Sigma	HPA003825
F3	Novus	NBP2-15139
FOXM1	Source Bioscience	LS-B3073
	Sigma	HPA029974
	Novus	NBP1-84671
HJURP	Sigma	HPA008436
IL1R1	Source Bioscience	LS-B2859
	Novus	NBP1-30929
IL8	Source Bioscience	LS-B6427
JUNB	Source Bioscience	LS-C176087
	Novus	NBP1-89544
KLF10	Source Bioscience	LS-C119009
KLF4	Source Bioscience	LS-B5641
	Novus	NBP2-24749
LDLR	Source Bioscience	LS-B1598
	Source Bioscience	LS-B8088
	Sigma	HPA009647
LGALS3	Source Bioscience	LS-B5661

- 18 -

	Source Bioscience	LS-B1671
	Sigma	HPA003162
LPAR1	Source Bioscience	LS-A212
	Source Bioscience	LS-A211
MALAT1	NA	NA
MTUS1	Novus	NBP1-82197
MYBPC1	Novus	NBP1-86427
NR4A3	Source Bioscience	LS-A2341
	Source Bioscience	LS-A2328
	Novus	NBP1-92198
OAT	Source Bioscience	LS-B4188
	Novus	NBP1-83239
PDE4D	Source Bioscience	LS-C185640
	Source Bioscience	LS-B8230
	Novus	NBP1-31131
PDK4	Source Bioscience	LS-B3459
PDRG1	Source Bioscience	LS-C163501
	Novus	NBP2-01854
PI15	Source Bioscience	LS-C163698
PTGS2	Source Bioscience	LS-B3296
	Source Bioscience	LS-B2145
	Novus	NB110-1948
RHOBTB3	Source Bioscience	LS-C120337
	Source Bioscience	NBP1-82954
RIN2	Sigma	HPA034641
SELE	Source Bioscience	LS-B2323
SOCS2	Source Bioscience	LS-B1257
SOCS3	Source Bioscience	LS-B3373
	Source Bioscience	NBP2-00850
SRSF5	Source Bioscience	LS-B3091
	Novus	NBP1-92381
SSTR1	Source Bioscience	LS-A994
ST6GAL1	Source Bioscience	LS-B6041
	Novus	NBP1-68447
TRPM3	Novus	NBP1-46344
TSC22D1	Source Bioscience	LS-B8419
XBP1	Source Bioscience	LS-B3178
	Source Bioscience	LS-B188
	Novus	NBP1-95395
ZFP36	Source Bioscience	LS-B5606

- 19 -

Measuring mRNA in a biological sample may be used as a surrogate for detection of the level of the corresponding protein in the biological sample. Thus, the expression level of any of the genes described herein can also be detected by detecting the appropriate RNA.

5 Accordingly, in specific embodiments the expression level is determined by microarray, northern blotting, RNA-seq (RNA sequencing), in situ RNA detection or nucleic acid amplification. Nucleic acid amplification includes PCR and all variants thereof such as real-time and end point methods and qPCR. Other nucleic acid amplification techniques are well known in the art, and include methods such as NASBA, 3SR and Transcription Mediated Amplification (TMA). Other suitable
10 amplification methods include the ligase chain reaction (LCR), selective amplification of target polynucleotide sequences (US Patent No. 6,410,276), consensus sequence primed polymerase chain reaction (US Patent No 4,437,975), arbitrarily primed polymerase chain reaction (WO 90/06995), invader technology, strand displacement technology, and nick displacement amplification (WO 2004/067726). This list is not intended to be exhaustive; any nucleic acid
15 amplification technique may be used provided the appropriate nucleic acid product is specifically amplified. Design of suitable primers and/or probes is within the capability of one skilled in the art. Various primer design tools are freely available to assist in this process such as the NCBI Primer-BLAST tool. Primers and/or probes may be at least 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 (or more) nucleotides in length. mRNA expression levels may be measured by reverse
20 transcription quantitative polymerase chain reaction (RT-PCR followed with qPCR). RT-PCR is used to create a cDNA from the mRNA. The cDNA may be used in a qPCR assay to produce fluorescence as the DNA amplification process progresses. By comparison to a standard curve, qPCR can produce an absolute measurement such as number of copies of mRNA per cell. Northern blots, microarrays, Invader assays, and RT-PCR combined with capillary
25 electrophoresis have all been used to measure expression levels of mRNA in a sample. See Gene Expression Profiling: Methods and Protocols, Richard A. Shimkets, editor, Humana Press, 2004.

RNA-seq uses next-generation sequencing to measure changes in gene expression. RNA may
30 be converted into cDNA or directly sequenced. Next generation sequencing techniques include pyrosequencing, SOLiD sequencing, Ion Torrent semiconductor sequencing, Illumina dye sequencing, single-molecule real-time sequencing or DNA nanoball sequencing.

In situ RNA detection involves detecting RNA without extraction from tissues and cells. In situ
35 RNA detection includes In situ hybridization (ISH) which uses a labeled (e.g. radio labelled, antigen labelled or fluorescence labelled) probe (complementary DNA or RNA strand) to localize a specific RNA sequence in a portion or section of tissue, or in the entire tissue (whole mount

- 20 -

ISH), or in cells. The probe labeled with either radio-, fluorescent- or antigen-labeled bases (e.g., digoxigenin) may be localized and quantified in the tissue using either autoradiography, fluorescence microscopy or immunohistochemistry, respectively. ISH can also use two or more probes to simultaneously detect two or more transcripts. A branched DNA assay can also be used for RNA in situ hybridization assays with single molecule sensitivity. This approach includes ViewRNA assays. Samples (cells, tissues) are fixed, then treated to allow RNA target accessibility (RNA un-masking). Target-specific probes hybridize to each target RNA. Subsequent signal amplification is predicated on specific hybridization of adjacent probes (individual oligonucleotides that bind side by side on RNA targets). A typical target-specific probe will contain 40 oligonucleotides. Signal amplification is achieved via a series of sequential hybridization steps. A pre-amplifier molecule hybridizes to each oligo pair on the target-specific RNA, then multiple amplifier molecules hybridize to each pre-amplifier. Next, multiple label probe oligonucleotides (conjugated to an enzyme such as alkaline phosphatase or directly to fluorophores) hybridize to each amplifier molecule. Separate but compatible signal amplification systems enable multiplex assays. The signal can be visualized by measuring fluorescence or light emitted depending upon the detection system employed. Detection may involve using a high content imaging system, or a fluorescence or brightfield microscope in some embodiments.

Thus, in a further aspect the present invention relates to a kit for (in situ) characterising and/or prognosing prostate cancer in a subject comprising one or more oligonucleotide probes specific for an RNA product of at least one of FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERRFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The kit may further comprise one or more of the following components:

- a) A blocking probe
- b) A PreAmplifier
- c) An Amplifier and/or
- d) A Label molecule

The components of the kit may be suitable for conducting a viewRNA assay (<https://www.panomics.com/products/rna-in-situ-analysis/view-rna-overview>).

- 21 -

The components of the kit may be nucleic acid based molecules, optionally DNA (or RNA). The blocking probe is a molecule that acts to reduce background signal by binding to sites on the target not bound by the target specific probes (probes specific for the RNA product of the at least one gene of the invention). The PreAmplifier is a molecule capable of binding to a (a pair of) target specific probe(s) when target bound. The Amplifier is a molecule capable of binding to the PreAmplifier. Alternatively, the Amplifier may be capable of binding directly to a (a pair of) target specific probe(s) when target bound. The Amplifier has binding sites for multiple label molecules (which may be label probes).

The invention also relates to use of the kit for characterising and/or prognosing prostate cancer.

RNA expression may be determined by hybridization of RNA to a set of probes. The probes may be arranged in an array. Microarray platforms include those manufactured by companies such as Affymetrix, Illumina and Agilent. Examples of microarray platforms manufactured by Affymetrix include the U133 Plus2 array, the Almac proprietary Xcel™ array and the Almac proprietary Cancer DSAs®, including the Prostate Cancer DSA®.

In specific embodiments, expression of the at least one gene may be determined using one or more probes selected from those in Table C below:

Table C – list of probes used to measure expression levels of the genes on an array.

Gene	Probeset ID	SEQ ID No
ADAMTS1	PC3P.3828.C1_s_at	30
	PCHP.1595_s_at	75
ADAMTS9	PC3P.17014.C1_s_at	24
	PC3SNGnh.5879_at	53
	PCADA.974_s_at	62
B3GNT5	PCRS2.398_s_at	81
CD38	3Snip.8317-94a_s_at	6
	PC3P.16779.C1_s_at	23
CEBPD	>PCHP.407_s_at_1160	89
	>PC3P.4961.C1_s_at_156	90
CENPF	PC3P.14957.C1_s_at	19
	PCADA.11788_at	55
CREM	PC3P.8627.C1_at	48
	PC3P.8627.C1_s_at	49
DKK1	3Snip.431-44a_s_at	4
EMP1	PC3P.10147.C1_at	9

- 22 -

ERRFI1	PC3P.3363.C1-522a_s_at	29
	PCADA.5865_at	58
	PCADA.5865_x_at	59
	PCRS2.6810_s_at	83
F12	PCADNP.16711_at	63
F3	3Snip.8552-9072a_s_at	7
	PC3P.8519.C1_s_at	46
	PC3P.8519.C2_s_at	47
	PCHP.1022_s_at	70
FOXMI	PCHP.1211_s_at	87
HJURP	PCADA.12835_s_at	56
IL1R1	PCADA.9303_s_at	61
IL8	PCHP.274_s_at	76
KLF10	PCRS2.574_s_at	82
KLF4	PC3P.14562.C1_s_at	18
LDLR	PCHP.101_s_at	69
LGALS3	3Snip.6331-2a_s_at	5
LPAR1	PC3P.4497.C1_at	34
	PCADA.11416_s_at	54
	PCADA.5036_s_at	57
	PCRS2.2781_s_at	80
MALAT1	3Snip.3677-484a_s_at	3
	PC3P.2436.C1_s_at	25
MTUS1	PCRS2.6884_s_at	84
MYBPC1	PC3P.13654.C1_at	16
	PC3P.13654.C1_x_at	17
	PC3P.3003.C1_s_at	27
	PC3P.7685.C1_at	40
	PC3P.7685.C1_x_at	41
	PC3P.7685.C1-693a_s_at	42
NFIL3	>PC3P.9419.C1-398a_s_at_365	88
NR4A3	PC3P.11087.C1_x_at	11
	PC3P.13257.C1_at	15
OAT	PC3P.2802.C1_s_at	26
PDE4D	PC3P.11285.C1_at	12
	PCADNP.1679_s_at	64
PKD4	PC3P.16300.C1_at	20
	PC3P.16300.C1_x_at	21
	PC3P.8159.C1_s_at	43
	PC3P.8159.C1-773a_s_at	44
	PC3SNGnh.4912_at	50
	PC3SNGnh.4912_x_at	51

- 23 -

	PC3SNGnh.5369_x_at	52
	PCADNP.18913_s_at	66
	PCEM.2221_at	68
	PCPD.29484.C1_at	78
PDRG1	>PC3P.15830.C1_s_at_353	93
PI15	3Snip.2873-1277a_at	2
	PC3P.7245.C1_at	38
	PC3P.7245.C1_x_at	39
	PC3P.8311.C1-482a_s_at	45
	PCADNP.17332_s_at	65
PTGS2	3Snip.950-71a_x_at	8
	PC3P.16654.C1_s_at	22
RHOBTB3	PC3P.12138.C1_at	13
	PC3P.12138.C1_x_at	14
	PC3P.5195.C1_s_at	35
RIN2	PC3P.7127.C1_s_at	37
RNFT2	PCADNP.401_s_at	67
SELE	PCHP.1458_s_at	74
SLC15A2	3Snip.1826-385a_s_at	1
	PC3P.10260.C1_at	10
	PC3P.3316.C1_at	28
	PCRS2.7997_s_at	86
SOCS2	PC3P.5499.C1_at	36
	PCHP.128_s_at	73
SOCS3	PCHP.491_s_at	77
SRSF5	PC3P.394.CB1_s_at	32
SSTR1	>PCHP.841_s_at_4070	91
	>PC3P.12563.C1_s_at_327	92
ST6GAL1	PCRS2.699_s_at	85
TRPM3	PCADA.7751_s_at	60
TSC22D1	PC3P.41.CB2_s_at	33
	PCHP.112_s_at	71
XBP1	PC3P.3909.C1-403a_s_at	31
	PCPD.59444.C1_at	79
ZFP36	PCHP.1147_s_at	72

These probes may also be incorporated into the kits of the invention. The probe sequences may also be used in order to design primers for detection of expression, for example by RT-PCR.

5 Such primers may also be included in the kits of the invention.

- 24 -

Increased rates of DNA methylation at or near promoters have been shown to correlate with reduced gene expression levels. DNA methylation is the main epigenetic modification in humans. It is a chemical modification of DNA performed by enzymes called methyltransferases, in which a methyl group (m) is added to specific cytosine (C) residues in DNA. In mammals, methylation occurs only at cytosine residues adjacent to a guanosine residue, i.e. at the sequence CG or at the CpG dinucleotide.

Accordingly, in yet a further aspect, the present invention relates to a method for characterising and/or prognosing prostate cancer in a subject comprising:

determining the methylation status of at least one of ADAMTS9, EMP1, F3, LDLR, LGALS3, MALAT1, MTUS1, NR4A3, PTGS2, RIN2, SLC15A2, SOCS3 and TSC22D1

in a sample from the subject wherein the determined methylation status is used to provide a characterisation of and/or a prognosis for the prostate cancer.

In certain embodiments if at least one of ADAMTS9, EMP1, F3, LDLR, LGALS3, MALAT1, MTUS1, NR4A3, PTGS2, RIN2, SLC15A2, SOCS3 and TSC22D1 is (hyper)methylated the likelihood of recurrence and/or metastasis is increased.

Determination of the methylation status may be achieved through any suitable means. Suitable examples include bisulphite genomic sequencing and/or by methylation specific PCR. Various techniques for assessing methylation status are known in the art and can be used in conjunction with the present invention: sequencing, methylation-specific PCR (MS-PCR), melting curve methylation-specific PCR (McMS-PCR), MLPA with or without bisulphite treatment, QAMA (Zeschnigk et al, 2004), MSRE-PCR (Melnikov et al, 2005), MethyLight (Eads et al., 2000), ConLight-MSP (Rand et al., 2002), bisulphite conversion-specific methylation-specific PCR (BS-MSP) (Sasaki et al., 2003), COBRA (which relies upon use of restriction enzymes to reveal methylation dependent sequence differences in PCR products of sodium bisulphite - treated DNA), methylation-sensitive single-nucleotide primer extension (MS-SNuPE), methylation-sensitive single-strand conformation analysis (MS-SSCA), Melting curve combined bisulphite restriction analysis (McCOBRA) (Akey et al., 2002), PyroMethA, HeavyMethyl (Cottrell et al. 2004), MALDI-TOF, MassARRAY, Quantitative analysis of methylated alleles (QAMA), enzymatic regional methylation assay (ERMA), QBSUPT, MethylQuant, Quantitative PCR sequencing and oligonucleotide-based microarray systems, Pyrosequencing, Meth-DOP-PCR. A review of some useful techniques for DNA methylation analysis is provided in Nucleic acids

- 25 -

research, 1998, Vol. 26, No. 10, 2255-2264, Nature Reviews, 2003, Vol.3, 253-266; Oral Oncology, 2006, Vol. 42, 5-13.

Techniques for assessing methylation status are based on distinct approaches. Some include
5 use of endonucleases. Such endonucleases may either preferentially cleave methylated
recognition sites relative to non-methylated recognition sites or preferentially cleave non-
methylated relative to methylated recognition sites. Some examples of the former are Acc III,
Ban I, BstN I, Msp I, and Xma I. Examples of the latter are Acc II, Ava I, BssH II, BstU I, Hpa II,
10 and Not I. Differences in cleavage pattern are indicative for the presence or absence of a
methylated CpG dinucleotide. Cleavage patterns can be detected directly, or after a further
reaction which creates products which are easily distinguishable. Means which detect altered
size and/or charge can be used to detect modified products, including but not limited to
electrophoresis, chromatography, and mass spectrometry.

15 Alternatively, the identification of methylated CpG dinucleotides may utilize the ability of the
methyl binding domain (MBD) of the MeCP2 protein to selectively bind to methylated DNA
sequences (Cross et al, 1994; Shiraishi et al, 1999). The MBD may also be obtained from MBP,
MBP2, MBP4, poly-MBD (Jorgensen et al., 2006) or from reagents such as antibodies binding to
methylated nucleic acid. The MBD may be immobilized to a solid matrix and used for preparative
20 column chromatography to isolate highly methylated DNA sequences. Variant forms such as
expressed His-tagged methyl-CpG binding domain may be used to selectively bind to methylated
DNA sequences. Eventually, restriction endonuclease digested genomic DNA is contacted with
expressed His-tagged methyl-CpG binding domain. Other methods are well known in the art and
include amongst others methylated-CpG island recovery assay (MIRA). Another method, MB-
25 PCR, uses a recombinant, bivalent methyl-CpG-binding polypeptide immobilized on the walls of
a PCR vessel to capture methylated DNA and the subsequent detection of bound methylated
DNA by PCR.

Further approaches for detecting methylated CpG dinucleotide motifs use chemical reagents that
30 selectively modify either the methylated or non-methylated form of CpG dinucleotide motifs.
Suitable chemical reagents include hydrazine and bisulphite ions. The methods of the invention
may use bisulphite ions, in certain embodiments. The bisulphite conversion relies on treatment
of DNA samples with sodium bisulphite which converts unmethylated cytosine to uracil, while
methylated cytosines are maintained (Furuichi et al., 1970). This conversion finally results in a
35 change in the sequence of the original DNA. It is general knowledge that the resulting uracil has
the base pairing behaviour of thymidine which differs from cytosine base pairing behaviour. This
makes the discrimination between methylated and non-methylated cytosines possible. Useful

- 26 -

conventional techniques of molecular biology and nucleic acid chemistry for assessing sequence differences are well known in the art and explained in the literature. See, for example, Sambrook, J., et al., *Molecular cloning: A laboratory Manual*, (2001) 3rd edition, Cold Spring Harbor, NY; Gait, M.J.(ed.), *Oligonucleotide Synthesis, A Practical Approach*, IRL Press (1984);
5 Hames B.D., and Higgins, S.J. (eds.), *Nucleic Acid Hybridization, A Practical Approach*, IRL Press (1985); and the series, *Methods in Enzymology*, Academic Press, Inc.

Some techniques use primers for assessing the methylation status at CpG dinucleotides. Two approaches to primer design are possible. Firstly, primers may be designed that themselves do
10 not cover any potential sites of DNA methylation. Sequence variations at sites of differential methylation are located between the two primers and visualisation of the sequence variation requires further assay steps. Such primers are used in bisulphite genomic sequencing, COBRA, Ms-SnuPE and several other techniques. Secondly, primers may be designed that hybridize specifically with either the methylated or unmethylated version of the initial treated sequence.
15 After hybridization, an amplification reaction can be performed and amplification products assayed using any detection system known in the art. The presence of an amplification product indicates that a sample hybridized to the primer. The specificity of the primer indicates whether the DNA had been modified or not, which in turn indicates whether the DNA had been methylated or not. If there is a sufficient region of complementarity, e.g., 12, 15, 18, or 20
20 nucleotides, to the target, then the primer may also contain additional nucleotide residues that do not interfere with hybridization but may be useful for other manipulations. Examples of such other residues may be sites for restriction endonuclease cleavage, for ligand binding or for factor binding or linkers or repeats. The oligonucleotide primers may or may not be such that they are specific for modified methylated residues.

25 A further way to distinguish between modified and unmodified nucleic acid is to use oligonucleotide probes. Such probes may hybridize directly to modified nucleic acid or to further products of modified nucleic acid, such as products obtained by amplification. Probe-based assays exploit the oligonucleotide hybridisation to specific sequences and subsequent detection
30 of the hybrid. There may also be further purification steps before the amplification product is detected e.g. a precipitation step. Oligonucleotide probes may be labeled using any detection system known in the art. These include but are not limited to fluorescent moieties, radioisotope labeled moieties, bioluminescent moieties, luminescent moieties, chemiluminescent moieties, enzymes, substrates, receptors, or ligands.

35 In the MSP approach, DNA may be amplified using primer pairs designed to distinguish methylated from unmethylated DNA by taking advantage of sequence differences as a result of

- 27 -

sodium-bisulphite treatment (WO 97/46705). For example, bisulphite ions modify non-methylated cytosine bases, changing them to uracil bases. Uracil bases hybridize to adenine bases under hybridization conditions. Thus an oligonucleotide primer which comprises adenine bases in place of guanine bases would hybridize to the bisulphite-modified DNA, whereas an
5 oligonucleotide primer containing the guanine bases would hybridize to the non-modified (methylated) cytosine residues in the DNA. Amplification using a DNA polymerase and a second primer yield amplification products which can be readily observed, which in turn indicates whether the DNA had been methylated or not. Whereas PCR is a preferred amplification method, variants on this basic technique such as nested PCR and multiplex PCR are also included within
10 the scope of the invention.

As mentioned earlier, one embodiment for assessing the methylation status of the relevant gene requires amplification to yield amplification products. The presence of amplification products may be assessed directly using methods well known in the art. They simply may be visualized
15 on a suitable gel, such as an agarose or polyacrylamide gel. Detection may involve the binding of specific dyes, such as ethidium bromide, which intercalate into double-stranded DNA and visualisation of the DNA bands under a UV illuminator for example. Another means for detecting amplification products comprises hybridization with oligonucleotide probes. Alternatively, fluorescence or energy transfer can be measured to determine the presence of the methylated
20 DNA.

A specific example of the MSP technique is designated real-time quantitative MSP (QMSP), and permits reliable quantification of methylated DNA in real time or at end point. Real-time methods are generally based on the continuous optical monitoring of an amplification procedure and utilise
25 fluorescently labelled reagents whose incorporation in a product can be quantified and whose quantification is indicative of copy number of that sequence in the template. One such reagent is a fluorescent dye, called SYBR Green I that preferentially binds double-stranded DNA and whose fluorescence is greatly enhanced by binding of double-stranded DNA. Alternatively, labelled primers and/or labelled probes can be used for quantification. They represent a specific
30 application of the well-known and commercially available real-time amplification techniques such as TAQMAN®, MOLECULAR BEACONS®, AMPLIFLUOR® and SCORPION®, DzyNA®, Plexor™ etc. In the real-time PCR systems, it is possible to monitor the PCR reaction during the exponential phase where the first significant increase in the amount of PCR product correlates to the initial amount of target template.

35 Real-Time PCR detects the accumulation of amplicon during the reaction. Real-time methods do not need to be utilised, however. Many applications do not require quantification and Real-Time

- 28 -

PCR is used only as a tool to obtain convenient results presentation and storage, and at the same time to avoid post-PCR handling. Thus, analyses can be performed only to confirm whether the target DNA is present in the sample or not. Such end-point verification is carried out after the amplification reaction has finished.

5

According to all aspects of the invention determining the expression level of at least one of FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

10

may involve determining the level of all or a selection of the transcripts and/or proteins isoforms produced from the gene. Examples of transcripts and corresponding protein isoforms that may be detected for each gene are shown in Table D below:

15

Table D – representative transcripts and corresponding protein isoforms that may be detected in the invention

Gene ID	Ensembl Transcript IDs detected by Almac probeset	Ensembl Protein ID
TRPM3	ENST00000377111	ENSP00000366315
	ENST00000423814	ENSP00000389542
	ENST00000357533	ENSP00000350140
	ENST00000354500	Non-protein coding
	ENST00000377110	ENSP00000366314
PDRG1	ENST00000202017	ENSP00000202017
SRSF5	ENST00000553635	ENSP00000451391
	ENST00000554465	Non-protein coding
	ENST00000556184	Non-protein coding
	ENST00000557154	ENSP00000451088
	ENST00000394366	ENSP00000377892
	ENST00000557460	Non-protein coding
	ENST00000556587	Non-protein coding
	ENST00000555547	Non-protein coding
	ENST00000556330	Non-protein coding
	ENST00000554929	Non-protein coding
	ENST00000553521	ENSP00000452123
PDE4D	ENST00000507116	ENSP00000424852
	ENST00000502575	ENSP00000425917
	ENST00000502484	ENSP00000423094

- 29 -

	ENST00000340635	ENSP00000345502
	ENST00000546160	ENSP00000442734
	ENST00000505453	ENSP00000421013
	ENST00000360047	ENSP00000353152
	ENST00000405053	Non-protein coding
	ENST00000405755	ENSP00000384806
	ENST00000309641	Non-protein coding
	ENST00000514231	Non-protein coding
PDK4	ENST00000473796	Non-protein coding
	ENST00000005178	ENSP00000005178
ADAMTS1	ENST00000284984	ENSP00000284984
	ENST00000464589	Non-protein coding
ADAMTS9	ENST00000482490	Non-protein coding
	ENST00000295903	ENSP00000295903
	ENST00000481060	ENSP00000417521
	ENST00000498707	ENSP00000418735
B3GNT5	ENST00000460419	ENSP00000420778
	ENST00000326505	ENSP00000316173
CEBPD	ENST00000408965	ENSP00000386165
CENPF	ENST00000366955	ENSP00000355922
CREM	ENST00000464475	ENSP00000418450
	ENST00000488328	ENSP00000417460
	ENST00000490460	Non-protein coding
	ENST00000479070	ENSP00000420511
	ENST00000463314	ENSP00000418336
	ENST00000374734	ENSP00000363866
	ENST00000484283	ENSP00000417165
	ENST00000463960	ENSP00000419684
	ENST00000460270	ENSP00000420437
	ENST00000473940	ENSP00000420681
	ENST00000469517	Non-protein coding
	ENST00000342105	ENSP00000341875
	ENST00000461968	Non-protein coding
	ENST00000374728	ENSP00000363860
	ENST00000395887	ENSP00000379225
	ENST00000495960	Non-protein coding
	ENST00000429130	ENSP00000393538
	ENST00000348787	ENSP00000345384
	ENST00000337656	ENSP00000337138
	ENST00000333809	ENSP00000333055
	ENST00000487132	ENSP00000418798
	ENST00000374721	ENSP00000363853
	ENST00000439705	ENSP00000409220

- 30 -

	ENST00000344351	ENSP00000344365
	ENST00000345491	ENSP00000265372
	ENST00000474362	ENSP00000419018
	ENST00000361599	ENSP00000354593
	ENST00000395895	ENSP00000379232
	ENST00000354759	ENSP00000346804
	ENST00000487763	ENSP00000417807
	ENST00000356917	ENSP00000349387
DKK1	ENST00000476752	Non-protein coding
	ENST00000373970	ENSP00000363081
EMP1	ENST00000256951	ENSP00000256951
	ENST00000537612	ENSP00000445319
ERRFI1	ENST00000487559	ENSP00000467030
	ENST00000474874	ENSP00000466958
	ENST00000377482	ENSP00000366702
	ENST00000467067	ENSP00000465100
	ENST00000469499	ENSP00000466454
HJURP	ENST00000441687	ENSP00000401944
	ENST00000411486	ENSP00000414109
	ENST00000432087	ENSP00000407208
	ENST00000433484	Non-protein coding
IL1R1	ENST00000422532	Non-protein coding
	ENST00000409929	ENSP00000386776
	ENST00000233946	ENSP00000233946
	ENST00000409288	ENSP00000386478
	ENST00000413623	Non-protein coding
	ENST00000409589	ENSP00000386555
	ENST00000424272	ENSP00000415366
	ENST00000409329	ENSP00000387131
	ENST00000428279	ENSP00000410461
	ENST00000410023	ENSP00000386380
JUNB	ENST00000302754	ENSP00000303315
KLF10	ENST00000285407	ENSP00000285407
	ENST00000395884	ENSP00000379222
KLF4	ENST00000497048	Non-protein coding
	ENST00000493306	Non-protein coding
	ENST00000374672	ENSP00000363804
LDLR	ENST00000252444	ENSP00000252444
	ENST00000560628	Non-protein coding
LGALS3	ENST00000254301	ENSP00000254301
	ENST00000556438	Non-protein coding
LPAR1	ENST00000358883	ENSP00000351755
	ENST00000541779	ENSP00000445697

- 31 -

	ENST00000374430	ENSP00000363552
	ENST00000374431	ENSP00000363553
MALAT1	ENST00000534336	Non-protein coding
	ENST00000517413	Non-protein coding
	ENST00000381861	ENSP00000371285
	ENST00000520196	Non-protein coding
MTUS1	ENST00000381869	ENSP00000371293
	ENST00000262102	ENSP00000262102
	ENST00000400046	ENSP00000382921
	ENST00000544260	ENSP00000445738
	ENST00000297488	ENSP00000297488
	ENST00000541119	ENSP00000442847
	ENST00000551300	ENSP00000447116
	ENST00000361466	ENSP00000354849
	ENST00000547509	ENSP00000447362
	ENST00000552198	Non-protein coding
	ENST00000547405	ENSP00000448175
	ENST00000441232	ENSP00000388989
	ENST00000452455	ENSP00000400908
MYBPC1	ENST00000550270	ENSP00000449702
	ENST00000392934	ENSP00000376665
	ENST00000545503	ENSP00000440034
	ENST00000550514	ENSP00000447404
	ENST00000550501	Non-protein coding
	ENST00000553190	ENSP00000447900
	ENST00000360610	ENSP00000353822
	ENST00000361685	ENSP00000354845
	ENST00000549145	ENSP00000447660
	ENST00000536007	ENSP00000446128
NFIL3	ENST00000534336	ENSP00000297689
	ENST00000395097	ENSP00000378531
NR4A3	ENST00000330847	ENSP00000333122
	ENST00000260113	ENSP00000260113
PI15	ENST00000523773	ENSP00000428567
	ENST00000490885	Non-protein coding
	ENST00000559627	Non-protein coding
PTGS2	ENST00000367468	ENSP00000356438
	ENST00000466691	Non-protein coding
	ENST00000510313	ENSP00000424844
RHOBTB3	ENST00000379982	ENSP00000369318
	ENST00000504179	ENSP00000422360
	ENST00000484638	Non-protein coding
RIN2	ENST00000255006	ENSP00000255006

- 32 -

RNFT2	ENST00000319176	ENSP00000321405
	ENST00000547718	Non-protein coding
	ENST00000551251	Non-protein coding
	ENST00000407967	ENSP00000385669
	ENST00000392549	ENSP00000376332
	ENST00000257575	ENSP00000257575
SELE	ENST00000367777	ENSP00000356751
	ENST00000367775	ENSP00000356749
	ENST00000367781	ENSP00000356755
	ENST00000333360	ENSP00000331736
	ENST00000367776	ENSP00000356750
	ENST00000367779	ENSP00000356753
	ENST00000367780	ENSP00000356754
	ENST00000367782	ENSP00000356756
SLC15A2	ENST00000469422	Non-protein coding
	ENST00000295605	ENSP00000295605
	ENST00000489711	ENSP00000417085
SOCS2	ENST00000549206	ENSP00000448815
	ENST00000549122	ENSP00000447161
	ENST00000548537	ENSP00000448709
	ENST00000551883	ENSP00000474805
	ENST00000340600	ENSP00000339428
SOCS3	ENST00000330871	ENSP00000330341
SSTR1	ENST00000267377	ENSP00000267377
ST6GAL1	ENST00000470633	Non-protein coding
	ENST00000457772	ENSP00000412221
	ENST00000169298	ENSP00000169298
	ENST00000448044	ENSP00000389337
TSC22D1	ENST00000261489	ENSP00000261489
	ENST00000458659	ENSP00000397435
	ENST00000501704	ENSP00000437414
XBP1	ENST00000216037	ENSP00000216037
	ENST00000405219	ENSP00000384295
	ENST00000344347	ENSP00000343155
	ENST00000403532	ENSP00000385162
	ENST00000482720	Non-protein coding
ZFP36	ENST00000594442	ENSP00000471239
	ENST00000248673	ENSP00000248673
	ENST00000597629	ENSP00000469647
F12	ENST00000510358	Non-protein coding
	ENST00000514943	Non-protein coding
	ENST00000502854	Non-protein coding
	ENST00000504406	Non-protein coding

- 33 -

	ENST00000253496	ENSP00000253496
CD38	ENST00000226279	ENST00000226279
	ENST00000502843	Non-protein coding
F3	ENST00000370207	ENSP00000359226
	ENST00000334047	ENSP00000334145
	ENST00000480356	Non-protein coding
IL8	ENST00000307407	ENSP00000306512
OAT	ENST00000471127	Non-protein coding
	ENST00000368845	ENSP00000357838
	ENST00000539214	ENSP00000439042
FOXMI	ENST00000342628	ENSP00000342307
	ENST00000536066	Non-protein coding
	ENST00000361953	ENSP00000354492
	ENST00000359843	ENSP00000352901

The methods described herein may further comprise extracting total nucleic acid or RNA from the sample. Suitable methods are known in the art and include use of commercially available kits
5 such as Rneasy and GeneJET RNA purification kit.

In certain embodiments the methods may further comprise obtaining the sample from the subject. Typically the methods are in vitro methods performed on an isolated sample.

- 10 According to all aspects of the invention samples may be of any suitable form. The sample may comprise, consist essentially of or consist of prostate cells and often a prostate tissue sample. The prostate cells or tissue may comprise prostate cancer cells. In specific embodiments the sample comprises, consists essentially of or consists of a formalin-fixed paraffin-embedded biopsy sample. The tissue sample may be obtained by any suitable technique. Examples
15 include a biopsy procedure, optionally a fine needle aspirate biopsy procedure. Body fluid samples may also be utilised. Suitable sample types include blood, to encompass whole blood, serum and plasma samples, urine and semen.

- The methods of the invention may comprise selecting a treatment for prostate cancer in a subject
20 and optionally performing the treatment. In certain embodiments if the characterisation of and/or prognosis for the prostate cancer is an increased likelihood of recurrence and/or metastasis and/or a poor prognosis the treatment selected is one or more of
- a) an anti-hormone treatment
 - b) a cytotoxic agent
 - 25 c) a biologic

- 34 -

- d) radiotherapy
- e) targeted therapy
- f) surgery

5 By anti-hormone treatment (or hormone therapy) is meant a form of treatment which reduces the level and/or activity of selected hormones, in particular testosterone. The hormones may promote tumour growth and/or metastasis. The anti-hormone treatment may comprise a luteinizing hormone blocker, such as goserelin (also called Zoladex), buserelin, leuprorelin (also called Prostag), histrelin (Vantas) and triptorelin (also called Decapeptyl). The anti-
10 hormone treatment may comprise a gonadotrophin release hormone (GnRH) blocker such as degarelix (Firmagon) or an anti-androgen such as flutamide (also called Drogenil) and bicalutamide (also called Casodex). In specific embodiments the anti-hormone treatment may be bicalutamide and/or abiraterone.

15 The cytotoxic agent may be a platinum based agent and/or a taxane. In specific embodiments the platinum based agent is selected from cisplatin, carboplatin and oxaliplatin. The taxane may be paclitaxel, cabazitaxel or docetaxel. The cytotoxic agent may also be a vinca alkaloid, such as vinorelbine or vinblastine. The cytotoxic agent may be a topoisomerase inhibitor such as etoposide or an anthracycline (antibiotic) such as doxorubicin. The cytotoxic agent may be an
20 alkylating agent such as estramustine.

By biologic is meant a medicinal product that is created by a biological process. A biologic may be, for example, a vaccine, blood or blood component, cells, gene therapy, tissue, or a recombinant therapeutic protein. Optionally the biologic is an antibody and/or a vaccine. The
25 biologic may be Sipuleucel-T.

In certain embodiments the radiotherapy is extended radiotherapy, preferably extended-field radiotherapy.

30 Surgery may comprise radical prostatectomy. By radical prostatectomy is meant removal of the entire prostate gland, the seminal vesicles and the vas deferens. In further embodiments surgery comprises tumour resection i.e. removal of all or part of the tumour.

By targeted therapy is meant treatment using targeted therapeutic agents which are directed
35 towards a specific drug target for the treatment of prostate cancer. In specific embodiments this may mean inhibitors directed towards targets such as PARP, AKT, MET, VEGFR etc. PARP inhibitors are a group of pharmacological inhibitors of the enzyme poly ADP ribose polymerase

- 35 -

(PARP). Several forms of cancer are more dependent on PARP than regular cells, making PARP an attractive target for cancer therapy. Examples (in clinical trials) include iniparib, olaparib, rucaparib, veliparib, CEP 9722, MK 4827, BMN-673 and 3-aminobenzamide. AKT, also known as Protein Kinase B (PKB), is a serine/threonine-specific protein kinase that plays a key role in multiple cellular processes such as glucose metabolism, apoptosis, cell proliferation, transcription and cell migration. AKT is associated with tumor cell survival, proliferation, and invasiveness. Examples of AKT inhibitors include VQD-002, Perifosine, Miltefosine and AZD5363. MET is a proto-oncogene that encodes hepatocyte growth factor receptor (HGFR). The hepatocyte growth factor receptor protein possesses tyrosine-kinase activity. Examples of kinase inhibitors for inhibition of MET include K252a, SU11274, PHA-66752, ARQ197, Foretinib, SGX523 and MP470. MET activity can also be blocked by inhibiting the interaction with HGF. Many suitable antagonists including truncated HGF, anti-HGF antibodies and uncleavable HGF are known. VEGF receptors are receptors for vascular endothelial growth factor (VEGF). Various inhibitors are known such as lenvatinib, motesanib, pazopanib and regorafenib.

The methods of the present invention can guide therapy selection as well as selecting patient groups for enrichment strategies during clinical trial evaluation of novel therapeutics. For example, when evaluating a putative anti-cancer agent or treatment regime, the methods disclosed herein may be used to select individuals for clinical trials that have prostate cancer characterized as having an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

The invention also relates to a system or device for performing a method as described herein.

In a further aspect, the present invention relates to a system or test kit for characterising and/or prognosing prostate cancer in a subject, comprising:

a) one or more testing devices for determining the expression level of at least one of FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in a sample from the subject

b) a processor; and

c) storage medium comprising a computer application that, when executed by the processor, is configured to:

(i) access and/or calculate the determined expression levels of the at least one of

- 36 -

5 FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERRFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in the sample on the one or more testing devices

(ii) calculate whether there is an increased or decreased level of the at least one of

10 FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERRFI1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in the sample; and

15 (iii) output from the processor the characteristaion of and/or prognosis for the prostate cancer.

By testing device is meant a combination of components that allows the expression level of a gene to be determined. The components may include any of those described above with respect to the methods for determining expression level at the level of protein, RNA or epigenetic modification. For example the components may be antibodies, primers, detection agents and so on. Components may also include one or more of the following: microscopes, microscope slides, x-ray film, radioactivity counters, scintillation counters, spectrophotometers, colorimeters, fluorometers, luminometers, and densitometers.

25 In certain embodiments the system or test kit further comprises a display for the output from the processor.

30 The invention also relates to a computer application or storage medium comprising a computer application as defined above.

35 In certain example embodiments, provided is a computer-implemented method, system, and a computer program product for characterising and/or prognosing prostate cancer in a subject, in accordance with the methods described herein. For example, the computer program product may comprise a non-transitory computer-readable storage device having computer-readable program instructions embodied thereon that, when executed by a computer, cause the computer

- 37 -

to characterise and/or prognose prostate cancer in a subject as described herein. For example, the computer executable instructions may cause the computer to:

- (i) access and/or calculate the determined expression levels of the at least one of FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in a sample on one or more testing devices;
- (ii) calculate whether there is an increased or decreased level of the at least one of FOXM1, TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in the sample; and,
- (iii) provide an output regarding the characterization of and/or prognosis for the prostate cancer.

In certain example embodiments, the computer-implemented method, system, and computer program product may be embodied in a computer application, for example, that operates and executes on a computing machine and a module. When executed, the application may characterise and/or prognose prostate cancer in a subject, in accordance with the example embodiments described herein.

As used herein, the computing machine may correspond to any computers, servers, embedded systems, or computing systems. The module may comprise one or more hardware or software elements configured to facilitate the computing machine in performing the various methods and processing functions presented herein. The computing machine may include various internal or attached components such as a processor, system bus, system memory, storage media, input/output interface, and a network interface for communicating with a network, for example. The computing machine may be implemented as a conventional computer system, an embedded controller, a laptop, a server, a customized machine, any other hardware platform, such as a laboratory computer or device, for example, or any combination thereof. The computing machine may be a distributed system configured to function using multiple computing machines interconnected via a data network or bus system, for example.

The processor may be configured to execute code or instructions to perform the operations and functionality described herein, manage request flow and address mappings, and to perform calculations and generate commands. The processor may be configured to monitor and control

- 38 -

the operation of the components in the computing machine. The processor may be a general purpose processor, a processor core, a multiprocessor, a reconfigurable processor, a microcontroller, a digital signal processor ("DSP"), an application specific integrated circuit ("ASIC"), a graphics processing unit ("GPU"), a field programmable gate array ("FPGA"), a programmable logic device ("PLD"), a controller, a state machine, gated logic, discrete hardware components, any other processing unit, or any combination or multiplicity thereof. The processor may be a single processing unit, multiple processing units, a single processing core, multiple processing cores, special purpose processing cores, co-processors, or any combination thereof. According to certain example embodiments, the processor, along with other components of the computing machine, may be a virtualized computing machine executing within one or more other computing machines.

The system memory may include non-volatile memories such as read-only memory ("ROM"), programmable read-only memory ("PROM"), erasable programmable read-only memory ("EPROM"), flash memory, or any other device capable of storing program instructions or data with or without applied power. The system memory may also include volatile memories such as random access memory ("RAM"), static random access memory ("SRAM"), dynamic random access memory ("DRAM"), and synchronous dynamic random access memory ("SDRAM"). Other types of RAM also may be used to implement the system memory. The system memory may be implemented using a single memory module or multiple memory modules. While the system memory may be part of the computing machine, one skilled in the art will recognize that the system memory may be separate from the computing machine without departing from the scope of the subject technology. It should also be appreciated that the system memory may include, or operate in conjunction with, a non-volatile storage device such as the storage media.

The storage media may include a hard disk, a floppy disk, a compact disc read only memory ("CD-ROM"), a digital versatile disc ("DVD"), a Blu-ray disc, a magnetic tape, a flash memory, other non-volatile memory device, a solid state drive ("SSD"), any magnetic storage device, any optical storage device, any electrical storage device, any semiconductor storage device, any physical-based storage device, any other data storage device, or any combination or multiplicity thereof. The storage media may store one or more operating systems, application programs and program modules such as module, data, or any other information. The storage media may be part of, or connected to, the computing machine. The storage media may also be part of one or more other computing machines that are in communication with the computing machine, such as servers, database servers, cloud storage, network attached storage, and so forth.

The module may comprise one or more hardware or software elements configured to facilitate the computing machine with performing the various methods and processing functions presented

- 39 -

herein. The module may include one or more sequences of instructions stored as software or firmware in association with the system memory, the storage media, or both. The storage media may therefore represent examples of machine or computer readable media on which instructions or code may be stored for execution by the processor. Machine or computer readable media may
5 generally refer to any medium or media used to provide instructions to the processor. Such machine or computer readable media associated with the module may comprise a computer software product. It should be appreciated that a computer software product comprising the module may also be associated with one or more processes or methods for delivering the module to the computing machine via a network, any signal-bearing medium, or any other
10 communication or delivery technology. The module may also comprise hardware circuits or information for configuring hardware circuits such as microcode or configuration information for an FPGA or other PLD.

The input/output ("I/O") interface may be configured to couple to one or more external devices, to
15 receive data from the one or more external devices, and to send data to the one or more external devices. Such external devices along with the various internal devices may also be known as peripheral devices. The I/O interface may include both electrical and physical connections for operably coupling the various peripheral devices to the computing machine or the processor. The I/O interface may be configured to communicate data, addresses, and control signals
20 between the peripheral devices, the computing machine, or the processor. The I/O interface may be configured to implement any standard interface, such as small computer system interface ("SCSI"), serial-attached SCSI ("SAS"), fiber channel, peripheral component interconnect ("PCI"), PCI express (PCIe), serial bus, parallel bus, advanced technology attached ("ATA"), serial ATA ("SATA"), universal serial bus ("USB"), Thunderbolt, FireWire, various video buses, and the like.
25 The I/O interface may be configured to implement only one interface or bus technology.

Alternatively, the I/O interface may be configured to implement multiple interfaces or bus technologies. The I/O interface may be configured as part of, all of, or to operate in conjunction with, the system bus. The I/O interface may include one or more buffers for buffering
30 transmissions between one or more external devices, internal devices, the computing machine, or the processor.

The I/O interface may couple the computing machine to various input devices including mice, touch-screens, scanners, electronic digitizers, sensors, receivers, touchpads, trackballs,
35 cameras, microphones, keyboards, any other pointing devices, or any combinations thereof. The I/O interface may couple the computing machine to various output devices including video displays, speakers, printers, projectors, tactile feedback devices, automation control, robotic

- 40 -

components, actuators, motors, fans, solenoids, valves, pumps, transmitters, signal emitters, lights, and so forth.

The computing machine may operate in a networked environment using logical connections through the network interface to one or more other systems or computing machines across the network. The network may include wide area networks (WAN), local area networks (LAN), intranets, the Internet, wireless access networks, wired networks, mobile networks, telephone networks, optical networks, or combinations thereof. The network may be packet switched, circuit switched, of any topology, and may use any communication protocol. Communication links within the network may involve various digital or an analog communication media such as fiber optic cables, free-space optics, waveguides, electrical conductors, wireless links, antennas, radio-frequency communications, and so forth.

The processor may be connected to the other elements of the computing machine or the various peripherals discussed herein through the system bus. It should be appreciated that the system bus may be within the processor, outside the processor, or both. According to some embodiments, any of the processor, the other elements of the computing machine, or the various peripherals discussed herein may be integrated into a single device such as a system on chip ("SOC"), system on package ("SOP"), or ASIC device.

Embodiments may comprise a computer program that embodies the functions described and illustrated herein, wherein the computer program is implemented in a computer system that comprises instructions stored in a machine-readable medium and a processor that executes the instructions. However, it should be apparent that there could be many different ways of implementing embodiments in computer programming, and the embodiments should not be construed as limited to any one set of computer program instructions. Further, a skilled programmer would be able to write such a computer program to implement one or more of the disclosed embodiments described herein. Therefore, disclosure of a particular set of program code instructions is not considered necessary for an adequate understanding of how to make and use embodiments. Further, those skilled in the art will appreciate that one or more aspects of embodiments described herein may be performed by hardware, software, or a combination thereof, as may be embodied in one or more computing systems. Moreover, any reference to an act being performed by a computer should not be construed as being performed by a single computer as more than one computer may perform the act.

The example embodiments described herein can be used with computer hardware and software that perform the methods and processing functions described previously. The systems, methods, and procedures described herein can be embodied in a programmable computer,

- 41 -

computer-executable software, or digital circuitry. The software can be stored on computer-readable media. For example, computer-readable media can include a floppy disk, RAM, ROM, hard disk, removable media, flash memory, memory stick, optical media, magneto-optical media, CD-ROM, etc. Digital circuitry can include integrated circuits, gate arrays, building block logic, field programmable gate arrays (FPGA), etc.

Reagents, tools, and/or instructions for performing the methods described herein can be provided in a kit. Such a kit can include reagents for collecting a tissue sample from a patient, such as by biopsy, and reagents for processing the tissue. The kit can also include one or more reagents for performing a expression level analysis, such as reagents for performing nucleic acid amplification, including RT-PCR and qPCR, NGS, northern blot, proteomic analysis, or immunohistochemistry to determine expression levels of biomarkers in a sample of a patient. For example, primers for performing RT-PCR, probes for performing northern blot analyses, and/or antibodies or aptamers, as discussed herein, for performing proteomic analysis such as Western blot, immunohistochemistry and ELISA analyses can be included in such kits. Appropriate buffers for the assays can also be included. Detection reagents required for any of these assays can also be included. The kits may be array or PCR based kits for example and may include additional reagents, such as a polymerase and/or dNTPs for example. The kits featured herein can also include an instruction sheet describing how to perform the assays for measuring expression levels.

The kit may include one or more primer pairs complementary to at least one of TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The kit may also include one or more primer pairs complementary to a reference gene, for example primers complementary to at least one of TPT1, RPS14 or RPL37A.

Such a kit can also include primer pairs complementary to at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 or 46 of

TRPM3, PDRG1, SRSF5, PDE4D, F12, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, CREM, DKK1, EMP1, ERFF1, F3, HJURP, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3,

- 42 -

RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

The kit may include one or more primer pairs complementary to at least one of TRPM3, PDRG1, F12, CENPF, HJURP, RNFT2, and SSTR1 and one or more primer pairs complementary to at least one of SRSF5, PDE4D, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CREM, DKK1, EMP1, ERFFI1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

Kits for characterising and/or prognosing prostate cancer in a subject may permit the methylation status of at least one of ADAMTS9, EMP1, F3, LDLR, LGALS3, MALAT1, MTUS1, NR4A3, PTGS2, RIN2, SLC15A2, SOCS3 and TSC22D1 to be determined. The determined methylation status, which may be hypermethylation, is used to provide a characterisation of and/or a prognosis for the prostate cancer. Such kits may include primers and/or probes for determining the methylation status of the gene or genes directly. They may thus comprise methylation specific primers and/or probes that discriminate between methylated and unmethylated forms of DNA by hybridization. Such kits will typically also contain a reagent that selectively modifies either the methylated or non-methylated form of CpG dinucleotide motifs. Suitable chemical reagents comprise hydrazine and bisulphite ions. An example is sodium bisulphite. The kits may, however, contain other reagents as discussed hereinabove to determine methylation status such as restriction endonucleases.

Accordingly, the invention also relates to a kit for characterising and/or prognosing prostate cancer in a subject comprising one or more antibodies or aptamers as described above.

As discussed above, in certain embodiments an increased expression level of at least one of TRPM3, PDRG1, F12, CENPF, HJURP, RNFT2, and SSTR1 or of FOXM1 indicates an increased likelihood of recurrence and/or metastasis and/or a poor prognosis. In further embodiments a decreased expression level of at least one of SRSF5, PDE4D, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CREM, DKK1, EMP1, ERFFI1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36 indicates an increased likelihood of recurrence and/or metastasis and/or a poor prognosis.

- 43 -

Accordingly, the kits described herein may comprise primers, probes or antibodies/aptamers (as discussed herein) for determining the expression level of at least one of TRPM3, PDRG1, F12, CENPF, HJURP, RNFT2, and SSTR1 or FOXM1 and at least one of SRSF5, PDE4D, PDK4, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CREM, DKK1, EMP1, ERRI1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36. Thus, the kits may incorporate reagents to determine expression levels of a combination of an up-regulated marker and a down-regulated marker. Suitable antibodies and/or primers/probes can be derived from Tables B, C and D herein.

Informational material included in the kits can be descriptive, instructional, marketing or other material that relates to the methods described herein and/or the use of the reagents for the methods described herein. For example, the informational material of the kit can contain contact information, e.g., a physical address, email address, website, or telephone number, where a user of the kit can obtain substantive information about performing a gene expression analysis and interpreting the results.

The kit may further comprise a computer application or storage medium as described above.

The example systems, methods, and acts described in the embodiments presented previously are illustrative, and, in alternative embodiments, certain acts can be performed in a different order, in parallel with one another, omitted entirely, and/or combined between different example embodiments, and/or certain additional acts can be performed, without departing from the scope and spirit of various embodiments. Accordingly, such alternative embodiments are included in the examples described herein.

Although specific embodiments have been described above in detail, the description is merely for purposes of illustration. It should be appreciated, therefore, that many aspects described above are not intended as required or essential elements unless explicitly stated otherwise.

Modifications of, and equivalent components or acts corresponding to, the disclosed aspects of the example embodiments, in addition to those described above, can be made by a person of ordinary skill in the art, having the benefit of the present disclosure, without departing from the spirit and scope of embodiments defined in the following claims, the scope of which is to be accorded the broadest interpretation so as to encompass such modifications and equivalent structures.

- 44 -

DESCRIPTION OF THE FIGURES

Figure 1

Unsupervised hierarchical clustering of a FFPE prostate cancer sample set comprising 70 primary prostate cancers, 20 primary cancers with concomitant metastatic disease, 11 metastatic disease and 25 normal prostate samples FFPE.

- A. Using the most variable genes across the dataset identified a subset of primary tumours that cluster with metastatic disease application (chi squared 2.77×10^{-10})
- B. Semi-supervised hierarchical clustering using 1083 differentially expressed genes identified from the internal dataset of the dataset published by Taylor and colleagues identified a similar subcluster of primary tumours that cluster with metastatic disease application (chi squared 2.78×10^{-6}).
- C. Kaplan-Meier analysis of the probability that patients would remain disease free following surgery if they were part of the metastatic biology group or not, hazard ratios were determined by log-rank test.

Figure 2

- A. Overlap of 83 overexpressed genes with FOXM1 CHIP-Seq hits from publications, hypergeometric test of overlap p-value 9.269×10^{-5}
- B. Box plot of pearson correlation scores of 39 over-expressed targets which overlapped with FOXM1 CHIP-seq hits and remaining over-expressed targets. T-test (p-value < 0.0001).

Figure 3

- A. Great (<http://bejerano.stanford.edu/great/public/html/>) functional analysis, molecular function of genomic regions in which the hypermethylated probes are located.
- B. Venn diagram demonstrating overlap of under-expressed genes with CHIP-SEQ identified targets of EZH2 and H3K27me3, Hypergeometric test of overlap.
- C. Venn diagram demonstrating overlap of under-expressed genes with hyper-methylated and H3K27me3 modifications.

Figure 4

Venn diagram showing the overlap between the top 10,000 ranked probesets including those that are least correlated between the metastatic biology subgroup and non-metastatic biology subgroup ("Lists 1 & 2") and those that are highly correlated between the non-metastatic biology subgroup and benign groups ("List 3").

- 45 -

Figure 5

GAP analysis of sample clusters identified in internal dataset.

5 Figure 6

Functional analysis of 1182 unique genes differentially expressed genes using Toppfun (<http://toppgene.cchmc.org/>)

A. Significant molecular processes of under-expressed genes

B. Significant molecular processes of overexpressed genes.

10

Figure 7

Study outline for screening potential IHC antibodies

EXAMPLES

15 The present invention will be further understood by reference to the following experimental examples.

Results

20 **Unsupervised hierarchical clustering identifies a distinct molecular subgroup in prostate cancer defined by metastatic biology**

We hypothesized that primary prostate cancers with metastatic potential would be transcriptionally similar to metastatic disease and primary disease with known concomitant metastases. In order to identify this metastatic subgroup we took an unsupervised hierarchical clustering approach using 70 resected primary prostate cancers clinically confined to prostate, 20
25 primary prostate cancer with known concomitant metastatic disease, 11 lymph nodes with metastatic disease, and 25 normal prostate samples. Clustering was performed using the most variable probe sets across the entire dataset. GAP statistical testing (Tibshirani et al 2001) identified 2 major sample clusters with statistical significance (figure 1A, figure 5).

30 One of these molecular subgroups had significant enrichment for metastatic disease and primary tumours with known concomitant metastases (chi squared $p=2.77 \times 10^{-10}$). Importantly, 29 primary prostate samples were also found in this group, which did not present with metastatic

- 46 -

disease but shared a similar transcriptional biology. This group of tumours is hereon referred to as the “metastatic biology subgroup” and the second subgroup the “non-metastatic subgroup”.

Next we performed gene expression analysis between primary tumours in the metastatic and non-metastatic subgroups and identified 1182 differentially expressed transcripts. The majority of these transcripts were under-expressed in the metastatic subgroup (1099 under-expressed versus 83 over-expressed).

In order to identify if the 1182 differentially expressed genes were prognostic in a second dataset, we used the genes to cluster the prostate cancer dataset published by Taylor and colleagues (Taylor et al 2010), this dataset represents prostate cancers managed by surgery for which PSA follow-up is available. Consistent with our internal training set, we found 2 robust sample clusters, one of which demonstrated enrichment for metastatic samples (Chi squared $p=2.78 \times 10^{-6}$ (Figure 1B). Importantly this group also contained 63 primary tumour samples without known metastatic disease at the time of presentation. Kaplan Meier analysis demonstrated that primary tumours within the metastatic biology group had a shorter time to disease recurrence (Figure 1C) following surgery (Hazard Ratio (HR) 2.377 and p-value 0.0351). The clinical and pathological characteristics of the sample clusters are detailed in table 1. Importantly, there were no differences in other prognostic clinical factors such as stage, grade or PSA levels pre-treatment.

Molecular pathways that underlie the metastatic biology group

To establish which molecular pathways give rise to the metastatic phenotype and poor prognosis, we performed pathway analysis using 1182 differentially expressed genes between the metastatic and non-metastatic subgroups. This identified 10 significant over-expressed pathways and 20 under-expressed pathways in the metastatic subgroup (Table 2i and 2ii). Interestingly the majority of pathways overexpressed in the metastatic subgroup were related to mitotic progression (Table 2i), whereas the molecular pathways that were under-expressed were involved in cell adhesion, morphology, ATF2 and p53 transcription.

To establish which of these molecular pathways were responsible for the poor prognosis we used the genes representing each pathway to cluster the Taylor dataset and a second dataset published by Sun and colleagues (Sun et al 2009). This later dataset represents primary prostate cancers managed with surgery with PSA follow-up. A Kaplan Meier analysis of time to recurrence was used for each of the observed clusters (Tables 2i and 2ii).

- 47 -

Of the overexpressed molecular pathways, only FOXM1 transcription factor network was significantly prognostic in the Taylor dataset (HR 2.755 $p=0.0134$). Furthermore FOXM1 itself was overexpressed in the metastatic biology subgroup in our internal training dataset (FC 2.13). To establish if increased FOXM1 was responsible for the overexpression of the mitotic genes in the metastatic biology group we interrogated 2 public FoxM1 CHIP-Seq data published by Sander and colleagues and Chen and colleagues (Chen et al 2013, Sanders et al 2013). We overlapped the identified FOXM1 CHIP-Seq targets with genes overexpressed in the metastatic biology group. Significantly, 39 out of 83 overexpressed genes in the metastatic subgroup were bound by FOXM1 in either of the datasets, with 20 being common to both. This overlap was highly significant (9.269×10^{-5}). Furthermore we performed correlation analysis of all the overexpressed transcripts against FOXM1 levels in the internal dataset, (supplementary table 3). Comparison of the correlations of the 39 FOXM1 targets identified through the analysis of CHIP-Seq data to the non-CHIP targets demonstrated a highly significant increase in correlation scores for the FOXM1 targets versus those not bound by FOXM1 (t test p -value <0.0001) of the CHIP-Seq targets (Figure 2B). Taken together this data strongly suggest that FoxM1 overexpression is responsible for the transcriptional activation of a large subset of the 83 genes that were detected as over-expressed in the metastatic subgroup.

Under-expressed molecular pathways that were significantly prognostic in both the Taylor and Sun datasets were muscle contraction, adipogenesis and ATF2 transcriptional targets. The diltiazem pathway was significantly prognostic in the Taylor dataset whereas integrin signaling and transcriptional targets for p53 although lost in the Taylor dataset, only reached prognostic significance in the Sun dataset.

Epigenetic Silencing of Gene Expression Occurs in the Metastatic biology subgroup

The majority of differentially expressed genes in the metastatic biology subgroup were down-regulated. Next we asked what potential mechanisms could account for this marked loss of gene expression in the metastatic biology group. Analysis of the molecular processes identified that genes involved in chromatin binding were over-expressed (figure 6), importantly we noted that several genes known to be involved in epigenetic gene regulation were up-regulated including AR, EZH2, HELLS and UHRF1) (Table 3).

UHRF1 was overexpressed in the metastatic biology subgroup (2.375 fold). This protein has recently been shown to contribute to and to maintain epigenetic silencing in prostate cancer (Babbio et al 2012). UHRF1 can bind to hemimethylated CpGs and can recruit DNMT1 to maintain DNA methylation patterns (Bostick et al 2007, Sharif et al 2007). Increased rates of

- 48 -

DNA methylation at or near promoters has been shown to correlate to reduced gene expression, this is most probably related to accessibility of transcription factors to the gene promoters.

We therefore measured the DNA methylation levels in a subset of 22 tumours from our interim training set (11 from each subgroup) using a high content DNA methylation array (sample details in supplementary table 3). Global analysis of the 1098 under-expressed genes in the metastatic subgroup demonstrated that 418 had increased rates of DNA methylation (p-value of overlap 1.546×10^{-34}) (table 4). Furthermore, analysis of the over-expressed gene sets showed no significant hyper or hypo methylation status thereby suggesting that altered methylation status is not important in the over-expressed gene sets.

GREAT (<http://bejerano.stanford.edu/great/public/html/>) analysis of the hyper-methylated genomic regions demonstrated a number of enriched molecular processes (Figure 3A), in particular DNA binding and transcription factor functions. This suggested that methylation not only silences genes directly in the metastatic biology group, but could be responsible for the loss of genes involved in transcription causing a further loss in gene expression.

Another gene involved in epigenetic silencing, EZH2 was over 2 fold over-expressed in the metastatic biology group (Table 3). EZH2 is a component of the PRC2 (Polycomb Repressive Complex 2) one of the two classes of polycomb-group proteins or (PcG). This complex has histone methyltransferase activity and EZH2 is the catalytic subunit. Indeed, EZH2 expression is the key determinant of histone methyltransferase activity. The PRC2 complex trimethylates histone H3 on lysine 27 (i.e. H3K27me3), this site is a mark of transcriptionally silent chromatin. To establish if EZH2 function might account for at least part of the loss of gene expression in the metastatic subgroup, we used a public CHIP-Seq (Wu et al 2012) prostate cancer cell line dataset. Specifically we compared genes known to bind EZH2 and H3K27me3 to those that were suppressed in the metastatic biology subgroup (Figure 3B). A significant number of the under-expressed genes were bound by EZH2, H3K27me3 or both (p-Value 2.597×10^{-12}), thereby strongly implicating chromatin silencing via EZH2 mediated histone modification as a key mechanism for silencing of a subset of the genes within the metastatic subgroup.

Interestingly, only a proportion of the targets of epigenetic silencing (123/602) had increased rates of hyper-methylation (Figure 3C) and were predicted to have H3K27me3 related histone modification thereby suggesting that the two mechanisms may work largely independently to silence gene expression.

- 49 -

Methods for detection of the metastatic biology subgroup

Hierarchical clustering is a useful analysis method of gene expression data from a number of samples, however it cannot be used to prospectively classify individual tumours. Additionally, in a previous study we have demonstrated that tumour heterogeneity in prostate cancer causes significant discordance between tumour biopsy and resected tumour profiles from the same patient. Therefore we elected to develop markers suitable for immunohistochemistry (IHC) that would prospectively classify if a tumour were similar to the metastatic biology subgroup.

To achieve this we employed 2 methods, firstly we identified transcripts that were differentially expressed between the metastatic biology subgroup and the non-metastatic biology subgroup but had little expression difference between the non-metastatic biology subgroup and normal. This process identified 393 probesets, using the 2-sample t-test method, ~75% of these probesets were overexpressed in the non-metastatic biology subgroup in comparison to the metastatic biology subgroup. We termed this approach the targeted as the normal prostate within the test case can be used as a reference.

For the second approach we assessed the 1182 differentially expressed between the metastatic biology subgroup non-metastatic subgroup, in this instance as there maybe expression differences between the non-metastatic biology group and benign/normal there is a requirement for a reference target, to identify suitable references we identified genes with minimal expression variance within all prostate cancer samples regardless of subgroup (the top 3 genes are summarised in table 7).

Prognostic utility of IHC targets

For the first approach the 393 probesets were mapped to a gene level to assist independent evaluation in an external dataset, Taylor et al 2010. In this dataset a total of 349 of the genes were detected. We performed multivariate analysis of these 349 genes in Taylor using time-to-biochemical recurrence with Cox proportional hazard corrected for Age, Grade and Stage, this resulted in 7 genes with significant multivariate prognostic function (p-value < 0.05.), these were TRPM3, PDRG1, SRSF5, PDE4D, CNPY4, F12 and PDK4. (Table 5) Univariate survival analysis was also performed, in which 52 genes were significant with a p-value < 0.05. There was an overlap of 3 genes in these top ranked probesets; these were SRSF5, PDE4D and PDK4. The 393 probesets were also assessed using an anova test to determine if they were significantly associated to clinical factors, namely Pathology Gleason score (and Gleason score 1 and 2).

- 50 -

For the second approach the 1182 differentially expressed genes tested in the same multivariate analysis, this identified 56 genes with significant multivariate prognostic function (p-value < 0.05.), (Table 6). Univariate survival analysis was also performed, in which 304 unique genes were significant with a p-value < 0.05. There was an overlap of 41 genes in these top ranked probesets. The numbers of targets with significant multivariate prognostic function was outside the scope of validation therefore we further refined the list by cross-referencing with the prognostic pathways (Table 2i and 2ii), the FOXM1 CHIP-Seq hits and selected literature review. The top 14 genes from the focused, pathway and literature comparisons are summarised in Table 7. FOXM1 itself and the differentially expressed FOXM1 CHIP-Seq targets which demonstrated significant multivariate prognostic power are summarized in Table 9

Discussion

Since the majority of men who develop early prostate cancer will not die from the disease, there is a clear requirement to better understand the biology underlying metastatic spread. This may allow appropriate selection of high-risk patients for more aggressive primary therapies and spare low risk patients unnecessary side effects.

In this study we have identified a group of primary prostate cancers that are similar to metastatic disease at a molecular level. These tumours are defined by loss of expression of several genes and defined pathways; furthermore this group is defined by activation of the proto-oncogene FOXM1 that leads to increased expression of genes involved in mitosis.

We have define a series of markers which have multivariate prognostic power and are highly suitable for IHC development to prospectively assess if a tumour is at increased likelihood of recurrence and metastatic development.

Table 1

	Metastatic Group	Biology	Others	p-value
Mean Gleason Score	6.952		6.714	0.0730
Mean Stage	2.429		2.381	0.648
Mean PSA	9.677		7.788	0.1984

- 51 -

Patient age	57.9	57.7	0.8633
Time to relapse	40.29	56.95	0.0014**

Clinical and pathological criteria of the metastatic biology tumours and non-metastatic biology group in the Taylor dataset.

Table 2i

Pathway	Pathway p-Value	Taylor HR	Taylor p-Value	Sun HR	Sun p-Value
Genes involved in Cell Cycle, Mitotic	1.78E-10	2.118	0.0614	2.061	0.0602
PLK1 signalling events	4.42E-07	1.843	0.1306	1.54	0.2774
Genes involved in Mitotic M-M/G1 phases	1.10E-06	1.988	0.0997	0.841	0.6823
Genes involved in Mitotic Prometaphase	3.92E-06	2.062280 883	0.0779	0.9289	0.8284
Aurora B signalling	8.51E-06	1.823	0.1352	1.031	0.9268
FOXM1 transcription factor network	4.19E-04	2.755	0.0134	1.737	0.1064
Genes involved in Cyclin A1 associated events during G2/M transition	1.21E-02	1.871	0.1274	0.7773	0.4777
Genes involved in Phosphorylation of the APC/C	1.79E-02	1.995	0.0922	0.9521	0.8838
Cell cycle	3.35E-02	1.837	0.1397	0.8063	0.5324

- 52 -

Genes involved in E2F transcriptional targets at G1/S	3.98E-02	0.462	0.0699	0.4214	0.07
---	----------	-------	--------	--------	------

Significant over-expressed pathways as detected using Topptfun, pathway p-value noted, Kaplan meier survival analysis results using pathways to cluster and define class labels i Taylor and Sun datasets.

5

Table 2ii

Pathway	Pathway p-Value	Taylor HR	Taylor p-Value	Sun HR	Sun p-Value
Genes involved in Muscle contraction	2.26E-10	2.495	0.0339	2.361	0.0196
Genes involved in Smooth Muscle Contraction	1.61E-07	2.155	0.0648	2.808	0.0094
Adipogenesis	4.05E-07	2.378	0.0391	2.336	0.0117
Focal Adhesion	4.79E-07	1.726	0.1814	1.01	0.9807
Striated Muscle Contraction	4.08E-06	2.600 1	0.0189	2.24	0.0163
Genes involved in Haemostasis	1.14E-04	1.1	0.8235	1.322	0.4404
Diltiazem Pathway	5.21E-04	2.289	0.0431	1.568	0.1993
Plasma membrane estrogen receptor signalling	6.24E-04	2.179	0.0914	1.384	0.3411
Genes involved in Formation of	7.81E-04	1.872	0.1226	1.19	0.6283

- 53 -

Platelet plug					
Genes involved in Platelet degranulation	9.60E-04	1.129	0.769	1.502	0.2263
Myometrial Relaxation and Contraction Pathways	1.04E-03	2.611	0.0188	2.06	0.0342
Integrins in angiogenesis	1.74E-03	1.639	0.2234	0.8937	0.7829
ATF-2 transcription factor network	1.74E-03	4.006	0.0037	4.026	0.0004
Genes involved in Platelet Activation	1.75E-03	2.047	0.0788	1.347	0.422
Syndecan-4-mediated signaling events	2.61E-03	3.686	0.0105	1.667	0.1268
LPA receptor mediated events	5.01E-03	2.158	0.149	1.332	0.3791
Integrin Signalling Pathway	1.41E-02	1.893	0.1202	2.069	0.0306
Genes involved in Integrin cell surface interactions	2.44E-02	1.584	0.2567	1.549	0.1905
Direct p53 effectors	2.53E-02	2.151	0.0676	3.836	0.0003
Integrin-mediated cell adhesion	3.78E-02	1.941	0.1084	0.765	0.4997

Significant under-expressed pathways as detected using Toppfun, pathway p-value noted, Kaplan meier survival analysis results using pathways to cluster and define class labels i Taylor and Sun datasets.

5 Table 3

	Fold Change	FDR corrected p-	Role in transcriptional
--	-------------	------------------	-------------------------

- 54 -

		value	repression
AR	2.41796	4.86E-10	Yes
CENPA	2.87805	3.23E-09	Na
CENPF	3.00853	4.19E-09	Na
DLX1	3.22068	8.49E-08	Na
EZH2	2.7026	2.45E-12	Yes
HELLS	2.10418	6.92E-05	Yes
TOP2A	2.90041	3.84E-09	Na
UBE2T	2.36638	3.65E-07	Na
UHRF1	2.37542	2.32E-09	Yes
ZIC2	2.08528	6.52E-05	Yes

Genes annotated as chromatin binding, fold change expression of metastatic biology group versus non and FDR corrected p-Value. Published role in transcriptional repression is noted.

Table 4

Genes Set Name	Genes methylated	Hyper- test p-value
Under-expressed genes	418/1098	1.546×10^{-34}
Over-expressed genes	13/83	0.947

- 5 Over or under-expressed genes with increased hyper-methylation in the metastatic biology group, Hypergeometric test to test significance of overlap.

Table 5

	Multivariate		Univariate		Independence (pvals)			Expression
Gene	HR	pval	HR	pval	Path GGS	PathG G1	PathG G2	

- 55 -

TRPM3	6.64	0.01	5.763	0.055	0.02	0.437	0.677	Over-expressed
PDRG1	4.883	0.027	2.416	0.256	0.046	0.035	0.351	Over-expressed
	5							
SRSF5	0.378	0.029	0.398	0.031	0.11	0.382	0.165	Under-expressed
PDE4D	0.425	0.035	0.268	0	0	0.004	0	Under-expressed
F12	5.523	0.042	2.789	0.263	0.895	0.845	0.844	Over-expressed
PDK4	0.641	0.048	0.52	0.001	0.009	0.069	0.018	Under-expressed

Top ranked prognostic markers based upon multivariate survival analysis in Taylor dataset.

Table 6

	Multivariate		Univariate		Expression in Metastatic biol. Group
	HR	pval	HR	pval	
ADAMTS1	0.553383	0.0348589	0.54021	0.0194342	Under-expressed
ADAMTS9	0.4585803	0.0288481	0.371399	0.00848	Under-expressed
B3GNT5	0.5137102	0.0299695	0.3956727	0.0017219	Under-expressed
CD38	0.6091854	0.0138355	0.5143886	0.0004166	Under-expressed
CEBPD	0.3271493	0.0097009	0.1813086	8.15E-06	Under-expressed
CENPF	3.5933385	0.0416512	9.1943228	0.0001128	Over-expressed
CREM	0.2330916	0.0448842	0.1947922	0.01595	Under-expressed
DKK1	0.2871025	0.0482413	0.3428314	0.0473753	Under-expressed
EMP1	0.4347836	0.0071015	0.3768645	0.0006745	Under-expressed
ERRFI1	0.5542049	0.0300146	0.5263541	0.0078253	Under-expressed
F3	0.579862	0.0170842	0.6133141	0.0418215	Under-expressed
HJURP	13.578677	0.0428399	46.05557	0.002649	Over-expressed
IL1R1	0.4800943	0.0108723	0.3570689	0.0002425	Under-expressed
IL8	0.3158031	0.0348507	0.47006	0.0724264	Under-expressed
JUNB	0.5484282	0.0347493	0.4460346	0.0008361	Under-expressed
KLF10	0.5017168	0.030925	0.4394025	0.0066195	Under-expressed
KLF4	0.3979693	0.0274113	0.2711992	0.000817	Under-expressed
LDLR	0.4540006	0.0299326	0.2845191	0.0001791	Under-expressed
LGALS3	0.2425137	0.0163003	0.1363453	0.0005586	Under-expressed

- 56 -

LPAR1	0.4085325	0.0398731	0.2924915	0.0043957	Under-expressed
MALAT1	0.1444922	0.0264399	0.2732161	0.1174596	Under-expressed
MTUS1	0.4461261	0.0438701	0.293677	0.0001736	Under-expressed
MYBPC1	0.5829982	0.002379	0.5657745	0.0003248	Under-expressed
NFIL3	0.494893	0.0271456	0.3853505	0.0006228	Under-expressed
NR4A3	0.3498147	0.0058837	0.3287529	0.0013712	Under-expressed
OAT	0.6455529	0.0451212	0.6272926	0.0531972	Under-expressed
PDE4D	0.4404056	0.0389414	0.2744262	0.0004312	Under-expressed
PDK4	0.6173687	0.0302305	0.5004434	0.0008031	Under-expressed
PI15	0.609224	0.0068663	0.4940706	7.96E-06	Under-expressed
PTGS2	0.5919948	0.0206793	0.5621402	0.0077942	Under-expressed
RHOBTB3	0.5457431	0.0455287	0.4498483	0.0117462	Under-expressed
RIN2	0.4232609	0.0422486	0.3777279	0.0290502	Under-expressed
RNFT2	38.724825	0.0168421	52.633909	0.0070457	Over-expressed
SELE	0.4784527	0.0139667	0.4724994	0.0036146	Under-expressed
SLC15A2	0.5968128	0.03609	0.4424453	0.0005391	Under-expressed
SOCS2	0.2955096	0.003396	0.2391982	0.0005038	Under-expressed
SOCS3	0.4423332	0.042005	0.3155164	0.0014183	Under-expressed
SSTR1	9.0410923	0.0182887	15.147803	0.0033901	Over-expressed
ST6GAL1	0.6042365	0.0088117	0.5305465	0.0016829	Under-expressed
TSC22D1	0.452536	0.0301503	0.3209002	0.0011309	Under-expressed
XBP1	0.2271852	0.0008373	0.2747813	0.0006004	Under-expressed
ZFP36	0.517509	0.0108194	0.4385203	0.0001823	Under-expressed

Top ranked prognostic markers based upon multivariate survival analysis in Taylor dataset of differentially expressed genes between metastatic biology subgroup and non-metastatic bio0logy subgroup.

Table 7

Gene	Multivariate HR	pval	Univariate HR	pval	Source	Expression in Metastatic biology Group
TRPM3	6.6406	0.0105	5.7636	0.0551	Targeted	Over- expressed
PDRG1	4.8835	0.0272	2.416	0.2566	Targeted	Over-

- 57 -

						expressed
SRSF5	0.3787	0.0294	0.3986	0.031	Targeted	Under-expressed
PDE4D	0.4255	0.035	0.2685	0.0005	Targeted	Under-expressed
PDK4	0.6415	0.0484	0.52	0.0019	Targeted	Under-expressed
F12	5.5235	0.0428	2.7897	0.2638	Targeted	Over-expressed
F3	0.591462629	0.026150109	0.641392413	0.070885567	Pathway	Under-expressed
HJURP	17.72622995	0.028647576	58.21478537	0.001862129	Pathway	Over-expressed
CENPF	4.009668447	0.028698111	9.892737548	8.66E-05	Pathway	Over-expressed
MYBPC1	0.616919233	0.009107673	0.596731068	0.001740583	Pathway	Under-expressed
SELE	0.506452771	0.023940071	0.494071466	0.006723235	Pathway	Under-expressed
CEBPD	0.33366283	0.012163084	0.188671187	1.76E-05	Pathway	Under-expressed
XBP1	0.227185248	0.000837328	0.274781347	0.000600431	Literature	Under-expressed
TPT1	NA	NA	NA	NA	Reference	NA
RPS14	NA	NA	NA	NA	Reference	NA
RPL37A	NA	NA	NA	NA	Reference	NA

Summarised IHC targets with reference genes.

Table 8

	Expression in Metastatic biology Group	Hypermethylated
ADAMT S9	Under-expressed	Hypermethylated
EMP1	Under-expressed	Hypermethylated
F3	Under-expressed	Hypermethylated

- 58 -

LDLR	Under-expressed	Hypermethylated
LGALS3	Under-expressed	Hypermethylated
MALAT1	Under-expressed	Hypermethylated
MTUS1	Under-expressed	Hypermethylated
NR4A3	Under-expressed	Hypermethylated
PTGS2	Under-expressed	Hypermethylated
RIN2	Under-expressed	Hypermethylated
SLC15A2	Under-expressed	Hypermethylated
SOCS3	Under-expressed	Hypermethylated
TSC22D1	Under-expressed	Hypermethylated

Top underexpressed markers that have increased hypermethylation levels.

Table 9

Gene	Multivariate		Univariate		Expression in Metastatic Biology Group
	HR	pval	HR	pval	
FOXM1	3.6353510	0.1583446	9.4463205	0.0116584	Over-expressed
	21	19	85	25	
CENPF	3.5933384	0.0416511	9.1943227	0.0001127	Over-expressed
	84	66	68	62	
HJURP	13.578676	0.0428399	46.055570	0.0026490	Over-expressed
	93	01	07	46	
RNFT2	38.724825	0.0168420	52.633908	0.0070456	Over-expressed
	02	83	75	87	
XBP1	0.2271852	0.0008373	0.2747813	0.0006004	Under-expressed
	48	28	47	31	
SOCS2	0.2955096	0.0033959	0.2391981	0.0005037	Under-expressed
	06	53	63	79	
NR4A3	0.3498146	0.0058836	0.3287529	0.0013712	Under-expressed
	83	74	43	06	
EMP1	0.4347835	0.0071014	0.3768645	0.0006744	Under-expressed

- 59 -

	57	7	3	58	
ZFP36	0.5175090	0.0108193	0.4385202	0.0001823	Under-expressed
	16	86	53	13	
IL1R1	0.4800943	0.0108722	0.3570688	0.0002424	Under-expressed
	31	79	77	95	
SELE	0.4784527	0.0139667	0.4724993	0.0036146	Under-expressed
	26	48	6	1	
LGALS3	0.2425137	0.0163002	0.1363453	0.0005586	Under-expressed
	44	57	09	37	
MALAT1	0.1444922	0.0264399	0.2732160	0.1174596	Under-expressed
	02	02	94	35	
NFIL3	0.4948930	0.0271455	0.3853504	0.0006228	Under-expressed
	03	95	74	18	
LDLR	0.4540006	0.0299326	0.2845191	0.0001791	Under-expressed
	47	34	4	38	
ERRFI1	0.5542048	0.0300145	0.5263541	0.0078252	Under-expressed
	51	85	1	97	
KLF10	0.5017168	0.0309250	0.4394024	0.0066194	Under-expressed
	31	35	87	67	
JUNB	0.5484281	0.0347492	0.4460345	0.0008361	Under-expressed
	87	58	58	12	
MTUS1	0.4461260	0.0438700	0.2936770	0.0001736	Under-expressed
	88	83	25	24	
CREM	0.2330915	0.0448841	0.1947922	0.01595	Under-expressed
	94	66	47		
RHOBTB3	0.5457430	0.0455287	0.4498483	0.0117462	Under-expressed
	73	1	27	42	
DKK1	0.2871025	0.0482412	0.3428313	0.0473752	Under-expressed
	03	65	7	65	

FOXM1 and FOXM1 CHIP-Seq targets which were differentially expressed in the metastatic biology group.

Methods

5 Patient Samples

- 60 -

126 samples (70 resected primary prostate cancers clinically confined to prostate, 20 primary prostate cancer with known concomitant metastatic disease, 11 lymph nodes with metastatic disease, and 25 normal prostate) were provided by Addenbrookes Hospital and Karolinska Institute following local ethical approval.

- 5 The subgroup and the prognostic significance were validated and tested in dataset published by Taylor et al which contained 179 samples (131 primary tumours, 29 normal and 19 metastatic disease. Time to biochemical recurrence and recurrence status following surgery were used to test prognostic significance, 5 samples were excluded from the analysis because of (surgery type PCA0056, and neo-adjuvant treatment, PCA0050, PCA0103, PCA119 and PCA0176).
- 10 Sun et al (79 tumour samples), samples were following surgery, 79 cases, 39 of which were classified as having disease recurrence.

Gene Expression Profiling.

- 15 Total RNA was extracted from macrodissected FFPE tumor samples using the Roche High Pure RNA Paraffin Kit (Roche Diagnostics Ltd.) as described previously (Kennedy RD, Bylesjo M, Kerr P et al. Development and independent validation of a prognostic assay for stage II colon cancer using formalin-fixed paraffin-embedded tissue. J Clin Oncol 2011; 29: 4620-4626). Total RNA was amplified using the NuGEN WT-Ovation™ FFPE System (NuGEN) and hybridized to the Almac Prostate Cancer DSA™ (Affymetrix) as described previously.

20 Statistical Analysis Methods

- A one-way ANOVA analysis identified differentially expressed probesets between 29 primary metastatic biology group tumours and 41 primarynon-metastatic biology tumours group controls using a fold-change (FC) threshold of $\text{abs(FC)} > 2$ and a significance p-value threshold adjusted for False Discovery Rate (FDR) ($\text{p-valueFDR} < 0.05$). Unique genes were determined as those in
25 the sense orientation with at least 6 probes aligned.

- A combined background & variance filter was applied to the data matrix to identify the most variable genes using an in-house developed feature selection program. Firstly, a background filter was applied to remove genes with expression values too low to be distinguished from the background noise. A high threshold was used to remove a large number of probesets and ensure
30 these probesets are highly expressed (Threshold: $\leq 10^{-16}$). Secondly, an intensity dependent variance filter was applied to the data matrix to remove probesets with low variance across all samples (Threshold: $\leq 5 \cdot 10^{-16}$). Feature selection resulted in 1651 most variable probesets.

- 61 -

Hierarchical clustering (Pearson correlation distance and Ward's linkage) was applied to probesets and samples from each dataset separately. The number of sub-clusters was determined using the gap statistic.

IHC target identification

- 5 The IHC targets of interest are those that are least correlated between the metastatic and non-metastatic groups (Lists 1 & 2) and those that are highly correlated between the non-metastatic and benign groups (List 3).

Correlation p-values for probesets in each of the three lists were ranked according to these criteria. The range of p-values observed in the top 10,000 ranked probesets in each list ranged
10 from [0 - 6.62e-05] for List 1, [1.03e-19 - 6.17e-04] for List 2 and [0.99 - 0.82] for List 3.

The intersection of the top 10,000 ranked probesets in the three lists revealed 512 common probesets (Figure 4). Antisense probesets and those with less than 6 probes aligned to the probeset were removed to leave 393. Partek® Genomics Suite™ version 6.6, was used to
15 generated fold change values.

Methylation

For the 22 patients, 11 metastatic biology subgroup and 11 non-metastatic biology subgroup, DNA was extracted using Recoverall (Life technologies). Genomic DNA (800 ng) was treated with sodium bisulfite using the Zymo EZ DNA Methylation Kit™ (Zymo Research, Orange, CA,
20 USA) according to the manufacturer's procedure, with the alternative incubation conditions recommended when using the Illumina Infinium Methylation Assay. The methylation assay was performed on 4 µl bisulfite-converted genomic DNA at 50 ng/µl according to the Infinium HD Methylation Assay protocol. Samples were processed onto Illumina 450k arrays as per
25 manufacturer's procedures. Uncorrected b-values were extracted with the same software. Probesets with changes in b-values that were statistically significant were assessed using the Significance Analysis of Microarrays (SAM) (Tusher et al 2001). Using a False Discovery Rate (FDR) of 0.05, of the 235,526 probesets on the array 32,286 were hypo-methylated (corresponding to 7,222 unique genes) and 9,184 probesets (4,003 unique genes).

30 References

Albertsen PC, Hanley JA, Barrows GH, Penson DF, Kowalczyk PD, Sanders MM *et al* (2005). Prostate cancer and the Will Rogers phenomenon. *J Natl Cancer Inst* 97: 1248-1253.

- 62 -

Altekruse SF, Huang L, Cucinelli JE, McNeel TS, Wells KM, Oliver MN (2010). Spatial patterns of localized-stage prostate cancer incidence among white and black men in the southeastern United States, 1999-2001. *Cancer Epidemiol Biomarkers Prev* 19: 1460-1467.

5 Babbio F, Pistore C, Curti L, Castiglioni I, Kunderfranco P, Brino L *et al* (2012). The SRA protein UHRF1 promotes epigenetic crosstalks and is involved in prostate cancer progression. *Oncogene*.

10 Bertucci F, Salas S, Eysteries S, Nasser V, Finetti P, Ginestier C, Charafe-Jauffret E, Loriod B, Bachelart L, Montfort J, Victorero G, Viret F, Ollendorff V, Fert V, Giovaninni M, Delpero JR, Nguyen C, Viens P, Monges G, Birnbaum D, Houlgatte R. Gene expression profiling of colon cancer by DNA microarrays and correlation with histoclinical parameters. *Oncogene*. 2004 Feb 19;23(7):1377-91. PubMed PMID: 14973550.

Bostick M, Kim JK, Esteve PO, Clark A, Pradhan S, Jacobsen SE (2007). UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science* 317: 1760-1764.

15 Chen X, Muller GA, Quaas M, Fischer M, Han N, Stutchbury B *et al* (2013). The forkhead transcription factor FOXM1 controls cell cycle-dependent gene expression through an atypical chromatin binding mechanism. *Mol Cell Biol* 33: 227-236.

Howlader A (2012). SEER Cancer Statistics Review, 1978-2009.

Kattan MW, Wheeler TM, Scardino PT (1999). Postoperative nomogram for disease recurrence after radical prostatectomy for prostate cancer. *J Clin Oncol* 17: 1499-1507.

20 Makarov DV, Sanderson H, Partin AW, Epstein JI (2002). Gleason score 7 prostate cancer on needle biopsy: is the prognostic difference in Gleason scores 4 + 3 and 3 + 4 independent of the number of involved cores? *J Urol* 167: 2440-2442.

Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA *et al* (2000). Molecular portraits of human breast tumours. *Nature* 406: 747-752.

25 Pound CR, Partin AW, Eisenberger MA, Chan DW, Pearson JD, Walsh PC (1999). Natural history of progression after PSA elevation following radical prostatectomy. *JAMA* 281: 1591-1597.

30 Rasiah KK, Stricker PD, Haynes AM, Delprado W, Turner JJ, Golovsky D *et al* (2003). Prognostic significance of Gleason pattern in patients with Gleason score 7 prostate carcinoma. *Cancer* 98: 2560-2565.

- 63 -

Sanders DA, Ross-Innes CS, Beraldi D, Carroll JS, Balasubramanian S (2013). Genome-wide mapping of FOXM1 binding reveals co-binding with estrogen receptor alpha in breast cancer cells. *Genome Biol* 14: R6.

5 Sharif J, Muto M, Takebayashi S, Suetake I, Iwamatsu A, Endo TA *et al* (2007). The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature* 450: 908-912.

Smith EB, Frierson HF, Jr., Mills SE, Boyd JC, Theodorescu D (2002). Gleason scores of prostate biopsy and radical prostatectomy specimens over the past 10 years: is there evidence for systematic upgrading? *Cancer* 94: 2282-2287.

10 Sun Y, Goodison S. Optimizing molecular signatures for predicting prostate cancer recurrence. Prostate. 2009 Jul 1;69(10):1119-27. doi: 10.1002/pros.20961.

Tibshirani R, Walther G, Hastie T (2001). Estimating the number of clusters in a data set via the gap statistic. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 63: 411-423.

15 Tusher VG, Tibshirani R, Chu G (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* 98: 5116-5121.

Taylor BS, Schultz N, Hieronymus H, Gopalan A, Xiao Y, Carver BS, Arora VK, Kaushik P, Cerami E, Reva B, Antipin Y, Mitsiades N, Landers T, Dolgalev I, Major JE, Wilson M, Socci ND, Lash AE, Heguy A, Eastham JA, Scher HI, Reuter VE, Scardino PT, Sander C, Sawyers CL,
20 Gerald WL. Integrative genomic profiling of human prostate cancer. *Cancer Cell*. 2010 Jul 13;18(1):11-22. doi: 10.1016/j.ccr.2010.05.026. Epub 2010 Jun 24.

Unoki M, Kelly JD, Neal DE, Ponder BA, Nakamura Y, Hamamoto R (2009). UHRF1 is a novel molecular marker for diagnosis and the prognosis of bladder cancer. *Br J Cancer* 101: 98-105.

25 van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M *et al* (2002). Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415: 530-536.

Xu K, Wu ZJ, Groner AC, He HH, Cai C, Lis RT, Wu X, Stack EC, Loda M, Liu T, Xu H, Cato L, Thornton JE, Gregory RI, Morrissey C, Vessella RL, Montironi R, Magi-Galluzzi C, Kantoff PW, Balk SP, Liu XS, Brown M. EZH2 oncogenic activity in castration-resistant prostate cancer cells is Polycomb-independent. *Science*. 2012 Dec 14;338(6113):1465-9. doi:
30 10.1126/science.1227604. PubMed PMID: 23239736; PubMed Central PMCID: PMC3625962.

- 64 -

5

Supplementary Table 1

Characteristic	Numbers	%
Type		
Primary Tumour	70	100
Primary Tumour with Mets	21	100
Metastatic Disease	10	100
Benign	25	100
Total	126	100
Gleason Score		
<7	10	7.936507937
7	24	19.04761905
>7	56	44.44444444
NA	36	28.57142857
	126	100
Centre		

- 65 -

Cambridge	73	57.93650794
Karolinska	53	42.06349206
Total	126	100

Patient characteristics of internal sample set.

Supplementary Table 2 FOXM1 correlations

Gene Symbol	r	p-value(correlation)	Lower CI	Upper CI	N	CHIP Overlap
CENPA	0.7729 42	2.91E-26	0.69152 2	0.83496 8	126	Yes
NUSAP1	0.7702 2	5.55E-26	0.68799 6	0.83292	126	Yes
KIF11	0.7628 21	3.08E-25	0.67843 1	0.82734 3	126	Yes
BUB1	0.7607 37	4.94E-25	0.67574 2	0.82577	126	Yes
TOP2A	0.7548 79	1.81E-24	0.66819 4	0.82134 2	126	Yes
CDCA3	0.7534 81	2.46E-24	0.66639 6	0.82028 5	126	Yes
KIF11	0.7494 25	5.90E-24	0.66118 4	0.81721 3	126	Yes
ANLN	0.7491 2	6.29E-24	0.66079 3	0.81698 2	126	Yes
FAM111B	0.7460 59	1.20E-23	0.65686 6	0.81466 1	126	Yes

- 66 -

KIFC1	0.7436	2.00E-23	0.65372	0.81280	126	Yes
	11		8	4		
ASPM	0.7418	2.91E-23	0.65142	0.81143	126	Yes
	14		7	9		
CENPA	0.7366	8.26E-23	0.64485	0.80753	126	Yes
	76		7	4		
RRM2	0.7317	2.18E-22	0.63862	0.80381	126	Yes
	95		8	8		
CENPF	0.7296	3.33E-22	0.63587	0.80217	126	Yes
	36		7	3		
UBE2C	0.7253	7.65E-22	0.63039	0.79888	126	Yes
	26		2	5		
ASPM	0.7202	2.01E-21	0.62390	0.79498	126	Yes
	23		9	5		
ZWINT	0.7160	4.34E-21	0.61863	0.79180	126	Yes
	62		3	1		
NUSAP1	0.7159	4.45E-21	0.61846	0.79169	126	Yes
	26		1	8		
EZH2	0.7152	5.03E-21	0.61761	0.79118	126	Yes
	54		1	3		
PKMYT1	0.7146	5.65E-21	0.61679	0.79069	126	Yes
	12		7	1		
KIFC1	0.7136	6.79E-21	0.61552	0.78992	126	Yes
	07		5	2		
IQGAP3	0.7134	6.98E-21	0.61532	0.78980	126	Yes
	52		8	2		
SPAG5 /// -- ///	0.7057	2.75E-20	0.60562	0.78391	126	Yes
-- /// ALDOC	74		6	2		
PTTG1	0.7007	6.62E-20	0.59926	0.78003	126	Yes

- 67 -

	28		7	4		
IQGAP3	0.7007	6.65E-20	0.59923	0.78001	126	Yes
	02		4	4		
MLF1IP	0.6956	1.57E-19	0.59287	0.77612	126	Yes
	49		9	4		
CDCA8	0.6951	1.72E-19	0.59221	0.77571	126	Yes
	22		6	8		
CDCA8	0.6908	3.53E-19	0.58679	0.77238	126	Yes
	06		8	9		
AURKA	0.6901	3.94E-19	0.58595	0.77186	126	Yes
	33		4	9		
PTTG1	0.6890	4.73E-19	0.58456	0.77101	126	Yes
	26		7	6		
HJURP	0.6886	4.99E-19	0.58414	0.77075	126	Yes
	93		9	8		
KIF14	0.6863	7.34E-19	0.58119	0.76893	126	Yes
	36		6	7		
RRM2	0.6851	8.92E-19	0.57968	0.76800	126	Yes
	29		7	5		
CDK1	0.6765	3.51E-18	0.56891	0.76133	126	Yes
	07		5			
CENPF	0.6736	5.44E-18	0.56538	0.75913	126	Yes
	72		2	2		
UHRF1	0.6733	5.75E-18	0.56493	0.75885	126	Yes
	13		5	4		
KIF20A	0.6661	1.72E-17	0.55599	0.75327	126	Yes
	26		7	1		
CDK1	0.6637	2.43E-17	0.55310	0.75145	126	Yes
	94		2	7		

- 68 -

HIST1H2AJ	0.6322 12	2.03E-15	0.51415 7	0.72675 7	126	Yes
PLK1	0.6242 58	5.72E-15	0.50442 6	0.72049 8	126	Yes
RNFT2	0.6169 6	1.44E-14	0.49552 4	0.71474 1	126	Yes
CBX2	0.6042 89	6.80E-14	0.48012 7	0.70471 4	126	Yes
UBE2T	0.6028 02	8.12E-14	0.47832 4	0.70353 4	126	Yes
SGOL1	0.5922 43	2.79E-13	0.46556	0.69514 4	126	Yes
CDT1	0.5730 18	2.36E-12	0.44245 4	0.67979 5	126	Yes
RHPN1	0.5365 86	9.40E-11	0.39913 6	0.65044 9	126	Yes
AMH	0.4972 71	3.16E-09	0.35307 1	0.61839 5	126	Yes
ZNF251	0.4870 51	7.33E-09	0.34120 8	0.60999 5	126	Yes
AR	0.4869 09	7.42E-09	0.34104 4	0.60987 8	126	Yes
ZNF467	0.4847 64	8.82E-09	0.33856	0.60811 2	126	Yes
PRAME	0.4793 2	1.36E-08	0.33226 6	0.60362 3	126	Yes
MEX3A	0.4614 66	5.38E-08	0.31171 6	0.58884 7	126	Yes
ZNF467	0.4499	1.25E-07	0.29852	0.57926	126	Yes

- 69 -

	44		8	6		
APLN	0.4483	1.40E-07	0.29674	0.57796	126	Yes
	82		5	4		
GPC2	0.4455	1.72E-07	0.29348	0.57558	126	Yes
	27		8	4		
PRAME	0.4395	2.61E-07	0.28672	0.57062	126	Yes
	91		7	6		
SAC3D1	0.4355	3.44E-07	0.28218	0.56728	126	Yes
	99		9	7		
RPL11	0.4099	1.87E-06	0.25317	0.54571	126	Yes
	32		5	3		
HOXC6	0.4081	2.10E-06	0.25114	0.54419	126	No
	28		6	1		
IDUA	0.4070	2.25E-06	0.24991	0.54326	126	No
	34		7	7		
LINC00662	0.4026	2.95E-06	0.24500	0.53956	126	No
	6		6	9		
APLN	0.4013	3.21E-06	0.24348	0.53842	126	No
	04		5	1		
DLX1	0.3973	4.08E-06	0.23908	0.53509	126	No
	75		2	3		
APLN	0.3973	4.10E-06	0.23901	0.53503	126	No
	11			9		
KIFC2	0.3866	7.74E-06	0.22712	0.52600	126	No
	74		5	9		
AR	0.3836	9.22E-06	0.22377	0.52345	126	Yes
	67		4			
AR	0.3754	1.48E-05	0.21466	0.51646	126	Yes
	76		4	8		

- 70 -

--- /// ---	0.3738	1.62E-05	0.21282	0.51505	126	No
	21		7	5		
COMTD1	0.3589	3.67E-05	0.19633	0.50229	126	Yes
	2		6	7		
PTPRT	0.3571	4.04E-05	0.19435	0.50075	126	No
	22		2	3		
ECE2	0.3555	4.39E-05	0.19264	0.49942	126	No
	7		1			
CANX	0.3553	4.45E-05	0.19234	0.49919	126	Yes
	06		9	3		
MIA3	0.3409	9.36E-05	0.17655	0.48681	126	No
	39		7	8		
CPNE4	0.3388	0.000103723	0.17432	0.48505	126	No
	97			5		
ODAM	0.3287	0.000170991	0.16322	0.47627	126	Yes
	47		3	1		
AR	0.3217	0.000239105	0.15558	0.47018	126	Yes
	36		1	6		
CPNE4	0.3132	0.000355568	0.14631	0.46276	126	No
	11		7	9		
RPS2 ///	0.3111	0.000390829	0.14407	0.46096	126	No
SNORA64 ///	42		3	6		
SNORA10						
AR	0.3108	0.000395174	0.14381	0.46075	126	Yes
	99			4		
HOXC4	0.3105	0.000401081	0.14345	0.46046	126	No
	73		6	9		
PTPRT	0.3099	0.000412213	0.14280	0.45994	126	No
	7		2	3		

- 71 -

CPNE4	0.3068	0.0004738	0.13945	0.45724	126	No
	83		8	9		
DSCAM-AS1	0.3059	0.000493863	0.13845	0.45644	126	No
	57		6			
PPFIA2	0.3004	0.000629772	0.13252	0.45164	126	No
	66		1	1		
IRAK1	0.2990	0.000669126	0.13102	0.45042	126	No
	81		5	8		
---	0.2933	0.000856865	0.12485	0.44541	126	No
	55		1			
HOXC4	0.2921	0.000901978	0.12355	0.44435	126	No
	52		5	5		
FOLH1 ///	0.2890	0.00102741	0.12024	0.44165	126	No
FOLH1B	76		6	5		
---	0.2822	0.00136596	0.11288	0.43563	126	No
	26		8	1		
NCAM2	0.2790	0.00155454	0.10949	0.43284	126	No
	58		2	1		
---	0.2660	0.00259802	0.09564	0.42139	126	No
	99		08	6		
---	0.2654	0.00266239	0.09496	0.42083	126	No
	66		55	6		
MT-TG /// MT- CO3 /// MT-TR /// MT-ND4L ///	0.2628	0.00294821	0.09213	0.41848	126	No
MT-ND3	11		58	4		
NTNG2	0.2596	0.00332155	0.08879	0.41570	126	No
	71		44	1		
SLC25A16	0.2565	0.00373266	0.08548	0.41294	126	No
	63		92	2		

- 72 -

---	0.2359	0.0078127	0.06369	0.39459	126	No
	72		1	6		
EIF3K	0.2299	0.00959972	0.05731	0.38918	126	No
	2		53			
SSTR1	0.2289	0.00992512	0.05627	0.38828	126	No
	27		02	9		
CST1	0.2214	0.0127055	0.04840	0.38157	126	No
	41		62	2		
NIPAL1	0.2115	0.0174161	0.03803	0.37265	126	No
	37		39	8		
SSTR1	0.2048	0.0213703	0.03108	0.36665	126	No
	85		9	5		
---	0.1844	0.0387046	0.00983	0.34810	126	No
	26		315	7		
OXR1	0.1818	0.0415884	0.00713	0.34573	126	Yes
	21		827	6		
OPRK1	0.1615	0.0706621	-	0.32724	126	No
	88		0.01370	9		
			96			
MT-TA /// MT-TY /// MT-TC /// MT-TN	0.1399	0.118182	-	0.30729	126	No
	03		0.03588	3		
			52			
LRRN1	0.1371	0.125644	-	0.30475	126	No
	59		0.03867	8		
			93			
MT-TC /// MT-TN /// MT-TY /// MT-TA	0.1294	0.148398	-	0.29766	126	No
	98		0.04646	6		
			47			
LRRN1	0.1258	0.16021	-	0.29429	126	No
	67		0.05014	8		

- 73 -

69						
MT-ND2 ///	0.1187	0.185204	-	0.28772	126	No
MTND2P28 ///	97		0.05730	9		
MT-TW			36			
MT-TH ///	0.0828	0.356482	-	0.25406	126	No
TS2 ///	304		0.09343			
MT-ND5 ///			33			
MT-TL2						
MT-TQ	0.0693	0.440025	-	0.24137	126	No
	974		0.10680	8		
			9			
WNT5A	0.0551	0.539432	-	0.22788	126	No
	785		0.12089	9		
			8			
SNORA61 ///	0.0508	0.571383	-	0.22381	126	No
SNHG12 ///	993		0.12512	7		
SNORA44			5			
MT-TC ///	-	0.923724	-	0.16654	126	No
TN ///	0.0086		0.18324	6		
MT-TY	152		9			
MT-CO1 ///	-	0.909852	-	0.16501	126	No
TW ///	0.0101		0.18476	6		
	882		9			

Pearson correlation of over-expressed targets to FOXM1 levels across the entire internal dataset.

- 74 -

Supplementary Table 3

Samples used for methylation analysis

mRNA Class label	Sample Type	Centre	Sample Age	Surgery Type	Gleason Score	Stage
Metastatic Biology	Primary Tumour	Karolinska	1	Radical	6	T2C
Metastatic Biology	Primary Tumour	Karolinska	2	Radical	8	T3A
Metastatic Biology	Metastatic Prostate Sample	Cambridge	7	Radical	9	NA
Metastatic Biology	Metastatic Prostate Sample	Cambridge	4	TURPS	9	NA
Metastatic Biology	Primary Tumour	Karolinska	8	Radical	8	TBC
Metastatic Biology	Primary Tumour	Karolinska	1	Radical	8	T3A
Metastatic Biology	Primary Tumour	Karolinska	1	TURPS	9	T3A
Metastatic Biology	Primary Tumour	Karolinska	1	Radical	8	T2A
Metastatic Biology	Primary Tumour	Karolinska	6	Radical	7	T3B
Metastatic Biology	Primary Tumour	Karolinska	1	Radical	8	T3A
Metastatic Biology	Primary Tumour	Karolinska	1	Radical	9	T2C
Non-Met. Biology	Primary Tumour	Karolinska	2	Radical	7	T3A
Non-Met. Biology	Primary Tumour	Karolinska	2	Radical	6	T3A
Non-Met. Biology	Primary Tumour	Karolinska	2	Radical	8	T3A

- 75 -

Non-Met. Biology	Primary Tumour	Cambridge	2	Radical	7	T3B
Non-Met. Biology	Primary Tumour	Cambridge	1	Radical	9	T3B
Non-Met. Biology	Primary Tumour	Cambridge	1	Radical	7	T3A
Non-Met. Biology	Primary Tumour	Karolinska	5	Radical	8	T2C
Non-Met. Biology	Primary Tumour	Karolinska	2	Radical	9	T3A
Non-Met. Biology	Primary Tumour	Karolinska	1	Radical	9	T3A
Non-Met. Biology	Primary Tumour	Karolinska	7	Radical	8	T2C
Non-Met. Biology	Primary Tumour	Karolinska	1	Radical	8	T3B

Prostate IHC development

Approach

- 5 To identify suitable antibodies we performed analysis of 3 antibodies per target for selected targets against a 4µM full face section from biomarker positive (confirmed by microarray profiling) prostate cancer samples. Each antibody was tested using 3 antigen retrieval methods using 3 dilutions (Figure 7).

- Each full face section contained an area of tumour, prostatic intraepithelial neoplasia (PIN),
 10 normal prostate epithelium, stroma and in some sections infiltrating immune cells.

This process allowed the identification of antibodies, antigen retrieval protocols and dilutions that detect the targets of interest.

Methods

Full face FFPE sections (4µm) of prostate tumour were used.

- 15 **Test samples:**

- 76 -

•Prostate tumour (DI 20052): Age 58: Male. Pathology - Adenocarcinoma of the prostate gland.
Tumour grade: 3+4=7.

•Prostate tumour (DI 20054): Age 70: Male: Pathology - Adenocarcinoma of the prostate gland.
Tumour grade: 3+4=7.

5 **Protocol**

All incubations were carried out at room temperature, unless otherwise stated.

1. Target retrieval (FFPE):

Antigen retrieval 1- Dako PT Link and 3-in-1 pH6.1 Target Retrieval (TR) Solution.

- _97°C 20 min with automatic heating and cooling.

10 Antigen retrieval 2- Dako PT Link and 3-in-1 pH9 Target Retrieval (TR) Solution.

- _97°C 20 min with automatic heating and cooling.

Antigen retrieval 3- Microwave Vector citrate pH6.1 heat-induced epitope retrieval (HIER).

•Slides were deparaffinized and rehydrated and then boiled (3 x 5 min) with the microwave set to full power.

15 All slides were rinsed with PBS - 10 min

2. Assay steps (DAKO Envision Flex plus)

- _EnVision peroxidase block - 5min
- _Rinse
- _Dako CSAII serum-free protein block - 10min

20 • _Air-removal

- _Primary antibody - 30 min
- _Rinse x2
- _EnVision Flex/HRP - 20 min
- _Rinse x2

25 • _DAB - 10 min

- 77 -

3. Counterstain and coverslipping

- Mayer's haematoxylin counterstain
 - Dehydrated in an ascending series of ethanols
 - Clear in xylene (x3)
- 5 •Coverslip under DePeX.

Reagents – Primary antibodies**CREM - anti-cAMP responsive element modulator**

- 1) Abcam Cat No: AB64832 Tested at 4, 2 and 1 µg/ml
- 10 2) Novus biomedical Cat No: NBP1-81760 Tested at 4, 2 and 1 µg/ml
- 3) Sigma Aldrich Cat No: HPA001818-100UL Tested at 0.8, 0.4 and 0.2µg/ml (recommended concentration 0.16µg/ml)
- R-IgG – rabbit polyclonal IgG (Rabbit isotype control) Alere Cat No: X0936

15 ERRFI1 - anti-ERBB receptor feedback inhibitor 1

- 1) Abcam Cat No: ab50272 Tested at 4, 2 and 1 µg/ml
- 2) Insight biotechnology Cat No: SC-137154 Tested at 4, 2 and 1 µg/ml (Santa Cruz Biotechnology, Inc.)
- 3) Sigma Aldrich Cat No: HPA027206-100UL Tested at 4, 2 and 1 µg/ml
- 20 •M-IgG1 - mouse monoclonal IgG1 (Mouse isotype control) Alere Cat No: X0931
- R-IgG1 – rabbit polyclonal IgG (Rabbit isotype control) Alere Cat No: X0936

HJURP anti-Holliday junction recognition protein

- 1) Abcam Cat No: AB100800 Tested at 4, 2 and 1 µg/ml Rabbit polyclonal
- 25 2) Abcam Cat No: AB175577 Tested at 4, 2 and 1 µg/ml Mouse monoclonal

- 78 -

3) Biorbyt Cat No: ORB140157 Tested at 4, 2 and 1 µg/ml Rabbit polyclonal

•Rabbit isotype control Alere Cat No: X0936

•Mouse IgG1 control Alere Cat No: X0931

5 **PDK4 – anti-pyruvate dehydrogenase kinase, isozyme 4**

1) Sigma Aldrich Cat No: HPA056731-100UL Tested at 4, 2 and 1µg/ml

2) LifeSpan BioSciences Cat No: LS-B3459 Tested at 4, 2 and 1µg/ml

3) Thermo scientific Cat No: PA5-13778 Tested at 4, 2 and 1µg/ml

•R-IgG – rabbit polyclonal IgG (Rabbit isotype control) Alere Cat No: X0936

10

•**SRSF5 - anti-serine/arginine-rich splicing factor 5**

1) Novus Biomedical Cat No: H00006430-B01P Tested at 4, 2 and 1µg/ml

2) Sigma Aldrich Cat No: HPA043484-100UL Tested at 4, 2 and 1µg/ml

3) LifeSpan BioSciences Cat No: LS-B3091 Tested at 4, 2 and 1µg/ml

15 •R-IgG1 – rabbit polyclonal IgG (Rabbit isotype control) Alere Cat No: X0936

•Sigma Aldrich Cat No: F3520-1ML

•Poly mouse IgG (M-IgG1, 2a, 2b)

• M-IgG1 - Alere Cat No: X0931

• M-IgG2a - Alere Cat No: X0943

20 • M-IgG2b - Alere Cat No: X0944

PDRG1 - anti-p53 and DNA damage-regulated protein 1

1) Abcam Cat No: AB175965 Tested at 4, 2 and 1µg/ml

2) Biorbyt Cat No: ORB162334 Tested at 4, 2 and 1µg/ml

- 79 -

3) Novus Biomedical Cat No: NBP2-01854 Tested at 4, 2 and 1 µg/ml

•M-IgG1 - mouse monoclonal IgG1 (Mouse isotype control) Alere Cat No: X0931

•R-IgG1 – rabbit polyclonal IgG (Rabbit isotype control) Alere Cat No: X0936

5

Results

Following review of all the data the following targets have demonstrated IHC assays which are specific and sensitive and can be used for prostate cancer classification or prognosis.

Target	Example Suitable Antibodies	Example Retrieval Conditions	Example Concentrations
CREM	Sigma Aldrich HPA001818	pH9 PT Link	0.8 µg/ml
		pH6 Microwave	0.8 µg/ml
	Novus Biomedical NBP1-81760	pH6 PT link	4 µg/ml
ERRFI1	ABCAM AB50272	pH9 PT Link	4 µg/ml
	Sigma Aldrich HPA027206	pH9 PT Link	4 µg/ml
HJURP	Biorbyt ORB140157	pH6 PT link	4 µg/ml
PDK4	Thermo Scientific PAS-13778	pH6 PT link	4 µg/ml
		pH9 PT Link	4 µg/ml
	Sigma Aldrich HPA056731	pH9 PT Link	4 µg/ml
SRSF5	Sigma Aldrich	pH6 PT link	4 µg/ml

- 80 -

	HPA043484		
PDRG1	ABCAM AB175965	pH9 PT Link	4µg/ml

5

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims. Moreover, all embodiments described herein are considered to be broadly applicable and combinable with any and all other consistent embodiments, as appropriate.

10

15 Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

- 81 -

Claims

1. A method for characterising and/or prognosing prostate cancer in a subject comprising:
5 determining the expression level of at least one of
CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1,
ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10,
KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15,
PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1,
10 TSC22D1, XBP1 and ZFP36
in a sample from the subject wherein the determined expression level is used to provide a
characterisation of and/or a prognosis for the prostate cancer.
2. The method of claim 1 comprising determining the expression level of at least one of
15 SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, F3, CENPF, MYBPC1, SELE, CEBPD,
and XBP1.
3. The method of claim 1 or 2 wherein the characterisation of and/or prognosis for the prostate
cancer comprises, consists essentially of or consists of predicting an increased likelihood of
20 recurrence and/or predicting an increased likelihood of metastasis.
4. The method of claim 1 or 2 wherein the characterisation of and/or prognosis for the prostate
cancer comprises, consists essentially of or consists of determining whether the prostate cancer
has a poor prognosis.
25
5. The method of any preceding claim comprising comparing the expression level to a reference
value or to the expression level in one or more control samples.
6. The method of any preceding claim wherein the expression level is compared to the
30 expression level of the same gene in one or more control samples.
7. The method of any preceding claim wherein the expression level of at least one of
SRSF5, PDK4, PDRG1, TRPM3, PDE4D, and F12
in the prostate cancer cells in a sample is compared to the expression level of the same gene in
35 the normal cells in the same sample.

- 82 -

8. The method of any preceding claim further comprising determining the expression level of a reference gene.

9. The method of any preceding claim wherein the expression level of at least one of
5 CREM, ERFFI1, HJURP, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36
is compared to the expression level of a reference gene.

10. The method of claim 8 or claim 9 wherein the reference gene is TPT1, RPS14 or RPL37A.

11. The method of any preceding claim wherein an increased expression level of at least one of HJURP, PDRG1, TRPM3, F12, CENPF, RNFT2, and SSTR1 indicates an increased likelihood of
15 recurrence and/or metastasis and/or a poor prognosis or wherein a decreased expression level of at least one of CREM, ERFFI1, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36 indicates an increased
20 likelihood of recurrence and/or metastasis and/or a poor prognosis.

12. The method of any preceding claim wherein the expression level is determined at the level of protein, RNA or epigenetic modification.

13. The method of claim 12 wherein the epigenetic modification is DNA methylation.

14. The method of any preceding claim wherein the expression level is determined by immunohistochemistry.

15. The method of any preceding claim wherein the expression level is determined using an antibody conjugated to a label.

16. The method of any preceding claim wherein the expression level is determined by microarray, northern blotting, RNA-seq (RNA sequencing), in situ RNA detection or nucleic acid
35 amplification.

17. The method of any preceding claim further comprising extracting total RNA from the sample.

- 83 -

18. The method of any preceding claim further comprising obtaining the sample from the subject.

5 19. The method of any preceding claim wherein the sample comprises, consists essentially of or consists of prostate tissue.

20. The method of any preceding claim wherein the sample comprises, consists essentially of or consists of a formalin-fixed paraffin-embedded biopsy sample.

10

21. The method of any preceding claim comprising determining the expression level of at least one of HJURP, PDRG1, TRPM3, F12, CENPF, RNFT2, and SSTR1 and at least one of CREM ,
ERRFI1, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1,
EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1,
15 MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2,
SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

22. A method for selecting a treatment for prostate cancer in a subject comprising:

(a) determining the expression level of at least one of

20 CREM, ERRFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, , ADAMTS1,
ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10,
KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15,
PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1,
TSC22D1, XBP1 and ZFP36

25 in a sample from the subject wherein the determined expression level is used to provide a characterisation of and/or a prognosis for the prostate cancer and

(b) selecting a treatment appropriate to the characterisation of and/or prognosis for the prostate cancer.

30 23. A method for selecting a treatment for prostate cancer in a subject comprising:

(a) determining the expression level of at least one of

CREM, ERRFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9,
B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR,
LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3,
35 RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and
ZFP36

- 84 -

in a sample from the subject wherein the determined expression level is used to provide a characterisation of and/or a prognosis for the prostate cancer

(b) selecting a treatment appropriate to the characterisation of and/or prognosis for the prostate cancer and

5 (c) treating the subject with the selected treatment.

24. The method claim 22 or 23, wherein if the characterisation of and/or prognosis for the prostate cancer is an increased likelihood of recurrence and/or metastasis and/or a poor prognosis the treatment selected is one or more of

10 a) an anti-hormone treatment, preferably bicalutamide and/or abiraterone

b) a cytotoxic agent

c) a biologic, preferably an antibody and/or a vaccine, more preferably Sipuleucel-T

d) radiotherapy, optionally extended radiotherapy, preferably extended-field radiotherapy

e) targeted therapy

15 f) surgery.

25. A method of treating prostate cancer comprising administering a chemotherapeutic agent or radiotherapy, optionally extended radiotherapy, preferably extended-field radiotherapy, to a subject or carrying out surgery on a subject wherein the subject is selected for treatment on the basis of a method as claimed in any of claims 22 to 24.

20

26. A chemotherapeutic agent for use in treating prostate cancer in a subject, wherein the subject is selected for treatment on the basis of a method as claimed in any of claims 22 to 24.

25

27. A method of treating prostate cancer comprising administering a chemotherapeutic agent or radiotherapy, optionally extended radiotherapy, preferably extended-field radiotherapy to a subject or carrying out surgery on a subject wherein the subject has an increased expression level of at least one of HJURP, PDRG1, TRPM3, F12, CENPF, RNFT2, and SSTR1 and/or a decreased expression level of at least one of CREM, ERFFI1, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

30

35

28. A chemotherapeutic agent for use in treating prostate cancer in a subject

- 85 -

wherein the subject has an increased expression level of at least one of HJURP, PDRG1, TRPM3, F12, CENPF, RNFT2, and SSTR1 and/or a decreased expression level of at least one of CREM, ERRF11, SRSF5, PDK4, PDE4D, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, SELE, SLC15A2, SOCS2, SOCS3, ST6GAL1, TSC22D1, XBP1 and ZFP36.

29. The method of claims 25 or 27 or chemotherapeutic agent for use of claims 26 or 28 wherein the chemotherapeutic agent comprises, consists essentially of or consists of

- a) an anti-hormone treatment, preferably bicalutamide and/or abiraterone
- b) a cytotoxic agent
- c) a biologic, preferably an antibody and/or a vaccine, more preferably Sipuleucel-T and/or
- d) a targeted therapeutic agent

30. The method of claim 24 or 29, wherein the cytotoxic agent is a platinum based agent and/or a taxane.

31. The method of claim 30, wherein the platinum based agent is selected from cisplatin, carboplatin and oxaliplatin.

32. The method of claim 30, wherein the taxane is paclitaxel or docetaxel.

33. An antibody that binds specifically to a protein product of at least one of CREM, ERRF11, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36.

34. The antibody of claim 33 conjugated to a label.

35. Use of an antibody of claim 33 or 34 for characterising and/or prognosing a prostate cancer in a subject.

36. A method for diagnosing a prostate cancer with an increased metastatic potential in a subject comprising:
determining the expression level of at least one of

- 86 -

CREM, ERFFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

in a sample from the subject wherein the determined expression level is used to identify whether a subject has a prostate cancer with increased metastatic potential.

37. A method for diagnosing a prostate cancer with an increased metastatic potential in a subject comprising:

determining the expression level of at least one of SRSF5, PDK4, TRPM3, PDRG1, PDE4D, and F12

in a sample from the subject wherein the determined expression level is used to identify whether a subject has a prostate cancer with increased metastatic potential.

38. A method for characterising and/or prognosing prostate cancer in a subject comprising: determining the expression level of at least one of

CREM, ERFFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 or

in a sample from the subject in order to identify the presence or absence of cells characteristic of an increased likelihood of recurrence and/or metastasis wherein the determined presence or absence of the cells is used to provide a characterisation of and/or a prognosis for the prostate cancer.

39. A method for characterising and/or prognosing prostate cancer in a subject comprising:

a) obtaining a sample from the subject

b) applying an antibody specific for the protein product of at least one of

CREM, ERFFI1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

to the sample from the subject

c) applying a detection agent that detects the antibody-protein complex

- 87 -

d) using the detection agent to determine the level of the protein
 d) wherein the determined level of the protein is used to provide a characterisation of and/or a prognosis for the prostate cancer.

5 40. A system or device for performing the method of any previous claim.

41. A system or test kit for characterising and/or prognosing prostate cancer in a subject, comprising:

a) one or more testing devices for determining the expression level of at least one of

10 CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

15 in a sample from the subject

b) a processor; and

c) storage medium comprising a computer application that, when executed by the processor, is configured to:

(i) access and/or calculate the determined expression levels of the at least one of

20 CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

25 in the sample on the one or more testing devices

(ii) calculate whether there is an increased or decreased level of the at least one of

30 CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36

in the sample; and

(iii) output from the processor the characterisation of and/or prognosis for the prostate cancer.

35

42. The system or test kit of claim 41 further comprising a display for the output from the processor.

- 88 -

43. A computer application or storage medium comprising a computer application as defined in claim 41 or 42.

5 44. A method for characterising and/or prognosing prostate cancer in a subject comprising:
determining the methylation status of at least one of
ADAMTS9, EMP1, F3, LDLR, LGALS3, MALAT1, MTUS1, NR4A3, PTGS2, RIN2, SLC15A2,
SOCS3 and TSC22D1
in a sample from the subject wherein the determined methylation status is used to provide a
10 characterisation of and/or a prognosis for the prostate cancer.

45. The method of claim 44, wherein if at least one of
ADAMTS9, EMP1, F3, LDLR, LGALS3, MALAT1, MTUS1, NR4A3, PTGS2, RIN2, SLC15A2,
SOCS3 and TSC22D1
15 is hypermethylated the likelihood of recurrence and/or metastasis is increased.

46. A kit for characterising and/or prognosing prostate cancer in a subject comprising one or more antibodies of claim 33 or 34.

20 47. The kit of claim 46 further comprising a computer application or storage medium of claim 43.

48. A computer program product for characterising and/or prognosing prostate cancer in a subject, comprising a non-transitory computer-readable storage device having computer-readable program instructions embodied thereon that cause the computer to:

25 (i) access and/or calculate the determined expression levels of the at least one of CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and
30 ZFP36 in a sample on one or more testing devices;
(ii) calculate whether there is an increased or decreased level of the at least one of CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1, ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10, KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15, PTGS2, RHOBTB3,
35 RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1, TSC22D1, XBP1 and ZFP36 in the sample; and,
(iii) provide an output regarding the characterization of and/or prognosis for the prostate cancer.

- 89 -

49. A kit for characterising and/or prognosing prostate cancer in a subject comprising one or more oligonucleotide probes specific for an RNA product of at least one of:

CREM, ERRF1, SRSF5, PDK4, HJURP, PDRG1, TRPM3, PDE4D, F12, ADAMTS1,
ADAMTS9, B3GNT5, CD38, CEBPD, CENPF, DKK1, EMP1, F3, IL1R1, IL8, JUNB, KLF10,
5 KLF4, LDLR, LGALS3, LPAR1, MALAT1, MTUS1, MYBPC1, NFIL3, NR4A3, OAT, PI15,
PTGS2, RHOBTB3, RIN2, RNFT2, SELE, SLC15A2, SOCS2, SOCS3, SSTR1, ST6GAL1,
TSC22D1, XBP1 and ZFP36

and further comprising one or more of the following components:

a) a blocking probe

10 b) a PreAmplifier

c) an Amplifier and/or

d) a Label molecule.

15

20

25

30

35

1/8




Figure 1A




Figure 1B

2/8




Figure 1C

3/8




Figure 2A




Figure 2B




Figure 3A

5/8




Figure 3B




Figure 3C

6/8




Figure 4




Figure 5

7/8




Figure 6A




Figure 6B

8/8




Figure 7

pctgb2014053694-seq1.txt
SEQUENCE LISTING

<110> Almac Diagnostics Limited

<120> PROSTATE CANCER CLASSIFICATION

<130> MPS/NRS1/P129999W000

<160> 93

<170> PatentIn version 3.5

<210> 1

<211> 275

<212> DNA

<213> Homo sapiens

<400> 1
ctccaccttt ctccaatgac agaagccacc tttctccaat gacagaagtt acctttctcc 60
aatgacagaa gttccaggcc ccagagactc tatgtctgcc cagagactct atgtctgccc 120
gtccaagaga ctctatgtct gcccggtccat gactctatgt ctgcccggtcc atcagactgg 180
attagacaag agagatagca tggattagac aagagagata gcagcgatta gacaagagag 240
atagcagcat attagacaag agagatagca gcata 275

<210> 2

<211> 275

<212> DNA

<213> Homo sapiens

<400> 2
ggcaatgtcc aaagtcaaat atcacgtcaa atatcaccta aactggtttag actggttaga 60
ttactttctac agctaattgc aggcactggc gccctctggt cgccctctgg tggttatgaa 120
gacaaaatgg ctacttgacc tacagcaaaa ttgacctaca gcaaaagcca tttctaagcc 180
atttctgtac cataaaaatt atcatatgtt tcctacatct gacaggtttc ctacatctga 240
cagcacctaa atctgacagc acctaaaatg tttga 275

<210> 3

<211> 275

<212> DNA

<213> Homo sapiens

<400> 3
gtcttttgctt tgactactaa tctgttgact actaatctgt cttcaggact taatctgtct 60
tcaggactct ttctggggag gtaacagcac aatatctttg ctttgaacta tatacatcct 120
tgatgacatc cttgatgtat aatttgtcag gatgtataat ttgtcaggag cttgagtcag 180
gagcttgact tgattgtata attcatatct acacgagaac ctaataatat aactgccttg 240
tctttttcag ttgtcttttt caggtaatag cctgc 275

<210> 4

<211> 275

<212> DNA

<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 4
acacattgat tggtatcttg actgagattg ttatcttgac tgacaaatat ccctttaccc 60
catttaattc tagagttacc ccatttaatt ctagagtcta tctagagtct agaacgcaag 120
gatctgagtc tagaacgcaa ggatctcttg caaggatctc ttggaatgac aaatggaatg 180
acaaatgata ggtacctaaa taggtaccta aaatgtaaca tgaaattata tttcccttta 240
ggctgtgata tttcccttta ggctgtgata gtttt 275

<210> 5
<211> 275
<212> DNA
<213> Homo sapiens

<400> 5
atacaagtac tggttgaacc tgacccaagt actggttgaa cctgaccact gtactggttg 60
aacctgacca cttcaggttg aacctgacca cttcaagggtt gaacctgacc acttcaagggt 120
tgcagaacct gaccacttca aggttgacagt aatgatgctc acttggttgca gtacagatgc 180
tcacttggtg cagtacaatc atgctcactt gttgcagtac aatcatgcag tacaatcatc 240
gggttaaaaa gcagtacaat catcgggtta aaaaa 275

<210> 6
<211> 275
<212> DNA
<213> Homo sapiens

<400> 6
ataatttacc gtaagtccta cttttattha ccgtaagtcc tacatttagt agtcctacat 60
ttagtatcaa gctagtacat ttagtatcaa gctagagact aagctagaga ctgaatttga 120
actcatccaa aattcatgtg ctttttccctt aaaattcatg tgctttttcc ttctaggcct 180
ttcataccaa actaatagta atagtagttt atattctctt ccaacgtagt ttatattctc 240
ttccaacaaa tatattctct tccaacaaat gcata 275

<210> 7
<211> 275
<212> DNA
<213> Homo sapiens

<400> 7
agttttacta cttaccaggg tactgggtac tgtataaatc cttgtgctaa taaatccttg 60
tgctaaattt tgctagctat agagtatgtg gtccctgctg aatactgtat ctctatgtta 120
cataggttac atagaaagcc ctaggagact ctctgtttat ctgaacaact atttggaaca 180
actatttgct gtactgataa ggaaacagca tagtctcatt cacttaacac attttgggtca 240
ttcggaaca aacaaaatac cctctctact tttat 275

<210> 8
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 8
aactcaggag aatcttcttt acaattatta acattgatct gctgacaaaa acattgatct 60
gctgacaaaa cctgggtgatc tgctgacaaa acctgggaat acctgggaat ttgggtttgtg 120
tatgcttggg ttgtgtatgc gaatgtttca gtgtatgcga atgtttcagt gcctcgaatg 180
tttcagtgcc tcagacaaat gtttcagtg ctcagacaaa tgtgtcagtg cctcagacaa 240
atgtgtattt tgcctcagac aaatgtgtat ttaac 275

<210> 9
<211> 275
<212> DNA
<213> Homo sapiens

<400> 9
acatttcaga catatccaaa gggaaaagaa gttgaactat gactggagta ggagtaaacc 60
atgtattccc ttatcgtaaa ccatgtattc cttatcttt gtaatttgca ttactctggt 120
ggattgcatt actctggtgg attgttctag ctggtggatt gttctagtag tgtatgattg 180
ttctagtact gtattgggct tgttctagta ctgtattggg cttctgtatt gggcttcttc 240
gttaatagat gggcttcttc gttaatagat tattt 275

<210> 10
<211> 275
<212> DNA
<213> Homo sapiens

<400> 10
ctgggtctatg cccttcagtg aacagatgcc cttcagtgaa cagttgtata aaccttgtct 60
tgtcaacctt gctgatcttg tcaaccttgc tgatccagct tcttattatt acaccttgtc 120
ttgctaacaa caatgtttct ataagtcaa gtttctataa gtccaacttc ctttacaact 180
tcctttatta atgtttccat aatgtttcca tttagcctca aaaatcctga tttacttttc 240
ccataagttt atttctgcct tgaagcctag aaatt 275

<210> 11
<211> 275
<212> DNA
<213> Homo sapiens

<400> 11
tatggttcat ttgcagtgac ttttagactt ttaaggcagt actgttttagc ggcagtactg 60
tttagcactt tgatagtttt gctaaattcg aataatgttt tatatactct gtatcaagtc 120
aaaataatat ctttggccat tttgctaaga ctttggccat tttgctaaga aacaaatgtc 180
aaactgatgt cacagtagtt gtcacagtag tttttgttag ctttatgtta gctttaaatc 240
atttttgctt aaaactatgc tgtttatatt gtcatt 275

<210> 12
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 12
tattcatcct agttctgaaa acatggagtt aatcatgtta ctgaatctaa tttgcaatat 60
tagcctgcct gtatttatta gcctgcctgt atttttcatg tcatgtaaac cttttgtaca 120
ttgaaaagaa cgtttgTTTT caggctcaga aacgtttgtt ttcaggctca gacctggctc 180
agaccttaag aactgatggt accttaagaa ctgatggtct tttcttacac aaaagtctaa 240
gcagttctga aacactacag tgacattggg tcatt 275

<210> 13
<211> 275
<212> DNA
<213> Homo sapiens

<400> 13
ggaatTTTgg gttattctct aaatgttttg ggttattctc taaatgacta gtgtgcttta 60
gaagtTTaAc tgtgattaac tgtgagaagg aaggcataat gaaggcataa ttcctggctg 120
caggactggt cgcaggagga aacttaaaat attggtagta agattaatca tcctaaatca 180
tcctaaactt gggaatccac taaacttggg aatccaccac tgaaggggaa tccaccactg 240
aagcataaat ttgccttgta ataaacttgt gaaat 275

<210> 14
<211> 275
<212> DNA
<213> Homo sapiens

<400> 14
ttttgggtta ttctctaaat gactagtgtg ctttagaagt ttaactgtga gaaggcataa 60
ttcctggctg caggactggt cgcaggagga aacttaaaat ttgttacatt gctgtgtctc 120
ctacatgtgt ctctacagt ccctagaact tccctagaac tgtgacttgc ctttaaatca 180
tcctaaactt gggaatccac taaacttggg aatccaccac tgaaggggaa tccaccactg 240
aagcataaat ttgccttgta ataaacttgt gaaat 275

<210> 15
<211> 275
<212> DNA
<213> Homo sapiens

<400> 15
tacagagtgg atgaccacac tagcaggatg accacactag cacagaagag gtgattcctc 60
cccttggcgg gagaggcggg agagctctct cagtgtgaac gagctctctc agtgtgaaca 120
tgctcggaa atcaggaagc caccagctgt atggagagtg ccttgctttt atttctcaga 180
cagcagagtt ttccaaagtt tttctgctcc tctaacagca ttgctcagca ttgctcttta 240
gtgtgtgta ttagtgtgtg ttaacctgtg gtttg 275

<210> 16
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 16
tctaccatTT actaattatg tgactcaaT atttgacccc tctgagactt atatttgacc 60
cctctgagac ttggattgac ccctctgaga cttggatttc cccctctgag acttggattt 120
cacatcctct gagacttgga tttcacatct ttaagtgtaa aaagcttggt tcacaagtgt 180
aaaaagcttg gttcacagta aaatatgtac ttcctttaag gaaatgatac agtagattat 240
taagcacctt acagtagatt attaagcacc ttgct 275

<210> 17
<211> 275
<212> DNA
<213> Homo sapiens

<400> 17
tctaccatTT actaattatg tgactgtcaa atatttgacc cctctgagac atatttgacc 60
cctctgagac ttggaccctt ctgagacttg gatttcacat agtgtaaaaa gcttggttca 120
cagtaaaagc ttggttcaca gtaggtggca ataaatatgt acttccttta aggaaatatt 180
cataaggaga caacacacaa gatacagtag attattaagc accttacagt agattattaa 240
gcaccttgct taagcacctt gctttgatta aaaag 275

<210> 18
<211> 275
<212> DNA
<213> Homo sapiens

<400> 18
aagtcaggta atatacctgg tttacagtca ggtaatatat ctggtttact aataatttgt 60
acaatggttt attccataat ttgtacaatg gtttattccc gtttattccc aagtatgcct 120
taagcttatt cccaagtatg ccttaagcag tccaagtat gccttaagca gaacacccaa 180
gtatgcctta agcagaacaa tgtatTTTgc atactcaagg tgagagtatt ttgcatactc 240
aaggtgagaa atTTTgcata ctcaaggtga gaatt 275

<210> 19
<211> 275
<212> DNA
<213> Homo sapiens

<400> 19
gaagcactga tcacctgtta gcattaagca ctgatcacct gttagcattg gcactgatca 60
cctgttagca ttgccactga tcacctgtta gcattgccat tgatcacctg ttagcattgc 120
cattcatata aagctTTTTg gtaatatgtt attaaaatga caagcactat atcactgagg 180
agaaaacaat tccttagaag gaggagaaaa caattcctta gaagttccca tgtgtaggta 240
ttgaaaaagt ccatgtgtag gtattgaaaa agttt 275

<210> 20
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 20
 ataaacctgt accttagtga atattaaacc tgtaccttag tgaatatattt tttagattag 60
 ggtaatacag acactgatta gggtaataca gacactgaag ctgaagatta tttgagaact 120
 gtactagatt atttgagaac tgtacttggc actgtacttg gcataaactc atttcgtact 180
 tggcataaac tcatttcatt ggcataaact catttcattg ttatatattt taacctatgg 240
 aaaattctgt aaattctgtg ttctcctaatt attat 275

<210> 21
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 21
 ataaacctgt accttagtga atattaaacc tgtaccttag tgaatatattt gattagggta 60
 atacagacac tgaagagatt atttgagaac tgtacttggc ttgagaactg tacttggcat 120
 aaactgtact tggcataaac tcatttcatt ggcataaact catttcattg ttataacttc 180
 tattccagat gcctttgagt aacctatgga aaattctgtg ttctcaaatt ctgtgttctc 240
 ctaatattat ctgtgttctc ctaatattat tgctt 275

<210> 22
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 22
 tttttacact gtcgatgttt ccaattttac actgtcgatg tttccaatgc tacactgtcg 60
 atgtttccaa tgcatttcca atgcatcttc catgatgcat tatttaaatt agacattacc 120
 agtaaaaatt agacattacc agtaatttca aaattttctaa attcataggg tagaacatag 180
 ggtagaatca cctgtaaaag agggtagaat cacctgtaaa agcttacctg taaaagcttg 240
 tttgatttct ctgtaaaagc ttgtttgatt tctta 275

<210> 23
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 23
 taaggtaaat gccagagacg gaagcaggtc aatgccagag acggaagcct tcaatgccag 60
 agacggaagc cttttcaatg ccagagacgg aagccttttt aatgccagag acggaagcct 120
 ttttcaaata acttatatca tcagcatacc atcatcagca tacctttatt gtgatcatca 180
 gcataccttt attgtgatct cagcatacct ttattgtgat ctatcgcata cttttattgt 240
 gatctatcaa cataccttta ttgtgatcta tcaat 275

<210> 24
 <211> 275
 <212> DNA
 <213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 24
 cttccctcat ctcctattat atattgaaat tccctcttaa gctcattgtg aattccctct 60
 taagctcatt gtgttttcatt gtgttttcac ccagttcact attgtgtttt caccagttc 120
 actagacca gttcactaga ccaacagcaa ttcactagac caacagcaat accaggcaat 180
 accagtacca gagttatgtt ttagaaagtc tactagctgt ccacaagtct actagctgtc 240
 cacaaagtcg gctgtccaca aagtcgtagt tcatt 275

<210> 25
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 25
 gaggtgggag gtaacagcac aatatgggag gtaacagcac aatatctttg ctttgaacta 60
 tatacatcct tgatgacatc cttgatgtat aatttgtcag gatgtataat ttgtcaggag 120
 cttgagtcag gagcttgact tgattgtata attcatatth acacgagaac ctaattttac 180
 acgagaacct aatataactg cctaataataa ctgccttgtc tttttaatat aactgccttg 240
 tctttttcag ttgtcttttt caggtaatag cctgc 275

<210> 26
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 26
 aacttcatat tggccaacac cagggacttc atattggcca acaccagggt cttcatattg 60
 gccaacacca gggttttcat attggccaac accagggttg tcatattggc caacaccagg 120
 gttgtatatt ggccaacacc agggttgtat tattggccaa caccagggtt gtattttggc 180
 caacaccagg gttgtattct tggccaacac cagggttgta ttctaggcca acaccagggt 240
 tgtattctat gccaacacca gggttgtatt ctatg 275

<210> 27
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 27
 gcagactcct cttgcaaggc gtaccagact cctcttgcaa ggcgtacctc ctcttgcaag 60
 gcgtacctcc aaacattgca aggcgtacct ccaaataa aaggcgtacc tccaaacata 120
 attgaggcgt acctccaaac ataattgatt cactcaagca atcctgagga atactactca 180
 agcaatcctg aggaatactg tcaagcaatc ctgaggaata ctgaggagga atactgaggg 240
 agggcctggc ggaatactga gggagggcct ggcta 275

<210> 28
 <211> 275
 <212> DNA
 <213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 28
 ttcagaaata taagagcggc agccaagagc ggcagccaga ccaaccagtt agcggcagcc 60
 agaccaacca gtttttcttg atgtgaacct gaccctactc tgaacctgac cctactcctt 120
 tagaactgac cctactcctt tagaagacag taaatagctt taacccaaac tgttccaaga 180
 gacattaaca cagttctgta gggaaatgtg gcatactgta tccttgagat atcctgcagt 240
 aaagacacaa gcagtaaaga cacaagatac cttgt 275

<210> 29
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 29
 cttggcaacc atattgtcac ctgtaaccat attgtcacct gtacctgtca cctgtacctg 60
 tcacacatct tgaaagtacc tgtcacacat cttgaaaaat gactaatttt cccttacagt 120
 tcctgtttcc cttacagttc ctgcttggtc tgcttggtcc caccactga agtagaagta 180
 gctcatcgta gtgcgggccg gctcatcgta gtgcgggccg tattagggta cgttagactc 240
 agatggaaaa attagctatg tgattgagag ttatt 275

<210> 30
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 30
 gtatcctggg gaatgtctgt tcagctatcc tgggtgaatgt ctgttcagct acatgtactt 60
 tataagtatt ggtttcatgt actttataag tattggtttg tactttataa gtattggttt 120
 gggtgacttt ataagtattg gtttgggtgt ttataagta ttggtttggg tgttcttata 180
 agtattgggt tgggtgttcc tataagtatt ggtttgggtg ttcctataag tattggtttg 240
 ggtgttcctt taagtattgg tttgggtgtt ccttc 275

<210> 31
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 31
 cggatgatgc aatataggag agaatggatg atgcaatata ggagagaata gatgatgcaa 60
 tataggagag aataagatgc aatataggag agaataaagc agcaggtctt cgagctagat 120
 tgacagcagg tcttcgagct agattgacag caggtcttcg agctagattg acagaaggtc 180
 ttcgagctag attgacagaa gaggggagga gcctgaagtg tatttagggg aggagcctga 240
 agtgatattt caccagctgg taaagttttc aaata 275

<210> 32
 <211> 275
 <212> DNA
 <213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 32
attcagaatt agtttaatgc ctttaattcag aattagttta atgccttaat ttaggggtcaa 60
gggtgtcctc cactctaggg tcaaggggtgt cctccactct ggggtcaaggg tgtcctccac 120
tcttttcaag ggtgtcctcc actctttaac agctgctgga cagacacatt agagcgctgc 180
tggacagaca cattagagca cacattagag cagctgtttg ttattattag agcagctggt 240
tgttattgat ttagagcagc tgtttggtat tgata 275

<210> 33
<211> 275
<212> DNA
<213> Homo sapiens

<400> 33
tgctgaatac gctgaagtaa atcctaatac gctgaagtaa atccttggtc gctgaagtaa 60
atccttggtc actgatgaag taaatccttg ttcactgaag tgaagtcttt caattgagct 120
ggttggaagt ctttcaattg agctgggtga ttgaaaaatg ctcaagttcta actaaaacta 180
atgaaatgga tttcccagta taatgaaatg gatttcccag taggggcata tcacctgtat 240
agtagttata tcacctgtat agtagttata tgcata 275

<210> 34
<211> 275
<212> DNA
<213> Homo sapiens

<400> 34
aaagaagtcc gggagatgaa tggctggatg tcagtgcacg gttagtgttt agtgcacggt 60
tagtgtttga gcagacagag ggcagacttg taaagtacct aagtacctgt agtgaaaaga 120
atgtgggacc cgattagcag aaaggtgttt gtgtttgcac atactttata caaaagaaat 180
gaacgtggac ttttacacat gaacgtggac ttttacacat gtgcatattt ttctggaggc 240
tatatggatt aaatcatttt caatcagcgt ttggt 275

<210> 35
<211> 275
<212> DNA
<213> Homo sapiens

<400> 35
catattaagt agtaacagaa gtctgaacag aagtctgaac aattggataa gtctgaacaa 60
ttggataaat ttgacgataa atttgacttc caagacagct atttgacttc caagacagct 120
aaactaaact acactacact gttatagtta tacactacac tgttatagtt aatctgttat 180
agttaatctg acaaaaatgt gttaatctga caaaaatgtc ctcaataaag catctgttta 240
attcaacctt aaagcatctg ttttaattcaa ccttt 275

<210> 36
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 36
 tttttgctat gttagtctgc ataactaaaa ttttcctgca tagctacaat attttcctgc 60
 atagctacaa tcctgctgca tagctacaat cctgtgggtgt gctacaatcc tgtgggtgtgt 120
 caccagtgtc accataaagg tggaccctgt aagggtggacc ctgtgtgaat gagaagtga 180
 ccctgtgtga atgagaaaaat acatggccag tcttttaciaa gttgagtcctt ttacaagttg 240
 agtaggcata acaaagtttt gtggccactt atttt 275

<210> 37
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 37
 aaataaaacc agtttgcagg tgcacataaa accagtttgc aggtgcacaa gcaggtgcac 60
 aaactatgag ggtccttgcac aaactatgag ggtccttgtat tcttgtatcc acgtaacaca 120
 ggtagttgta tccacgtaac acaggtagtt gtactgtgta aagatgcata gtcacatag 180
 tcatctcatt tggttggctt tctcatttgg ttggctttgt accttccttt tttagccttg 240
 gctttttgtt tagccttggc ttttgttgaa ctaga 275

<210> 38
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 38
 aatatctaaa tatgtctcac atattattac ctaattctgt ggcaaaaatg aattctgtgg 60
 caaaaatggg gcctcctgtg gcaaaaatgg tgcctctgat tgtggcaaaa atgggtgcctc 120
 tgatggatgt tgtgatatag tattgtcagt gttgtgatat agtattgtca gtgtgggtgat 180
 atagtattgt cagtgtgtac tagtattgtc agtgtgtaca tatactttat gaaccataac 240
 aaatgtagct atgaaccata acaaatgtag ctttt 275

<210> 39
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 39
 taatcctcaa atatactgta ccattggaga agttttattc attacctaatt attacctaatt 60
 tctgtggcaa aaatgctgtg gcaaaaatgg tgcctctgat aaatgggtgcc tctgatgttg 120
 tgatagcctc tgatgtttgtg atatagtatt gtgatatagt attgtcagtg tgtacaaaac 180
 ctgtgtaaac ctctgtcctt gtaaacctct gtccttatga accatatgaa ccataacaaa 240
 ttagactttt gctttttaaa gtccattgta ttggt 275

<210> 40
 <211> 275
 <212> DNA
 <213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 40
actcttcact aggattccct atatacttca ctaggattcc ctatatatat tctagatatt 60
tactatagtg cagaagatat ttactatagt gcagaagcaa tatttactat agtgcagaag 120
caacagatta tgttttatct agcaaatttg tatgttttat ctagcaaatt tgtaagtgt 180
ttctatgggt attaaaagac tgctttctat ggttattaaa agactgctat actcttcact 240
aggattccct ctatactctt cactaggatt cccta 275

<210> 41
<211> 275
<212> DNA
<213> Homo sapiens

<400> 41
cttcactagg attccctata tatatatatg cctgttatag atatctagat gatatttact 60
atagtgcaga agcaagatta tgttttatct agcaaatttg tttcatttaa ctttccaggc 120
ttttaaatta ctttagtgct tttctatgggt gtgctttcta tggttattaa aagaccatt 180
ttataagatt tctctccata gatttctctc catatatatt tctattataa ctgctatact 240
cttcactagg ctatactctt cactaggatt cccta 275

<210> 42
<211> 275
<212> DNA
<213> Homo sapiens

<400> 42
ctcctgaaac gtaggtgaga acaaagaaac gtaggtgaga acaaagtgat acgtaggtga 60
gaacaaagtg atggatgatg gagtgagaca ttgtggcctg agtgagacat tgtggcctgg 120
aagtcgtgag acattgtggc ctggaagtct acaactgcat actaatcttt taaagactgc 180
atactaactct tttaaagcct ctgcatacta atctttttaa gcctttttaa gcctttaaca 240
gttgctttta taaagccttt aacagttgct tttaa 275

<210> 43
<211> 275
<212> DNA
<213> Homo sapiens

<400> 43
gaaatgttag tatctcaatt acaccgttag tatctcaatt acaccaactg agtatctcaa 60
ttacaccaac tgtgcatctc aattacacca actgtgcaaa tagtagatgg atactgaata 120
ctcagatgga tactgaatac tcaggccac atactcaggc ccacttaaatt tattatactc 180
aggcccactt aaattattaa gtttttgtct ttatgctatg tacaggctctt tatgctatgt 240
acagagaaat gctatgtaca gagaaatgtg ataata 275

<210> 44
<211> 275
<212> DNA
<213> Homo sapiens

pctgb2014053694-seq1.txt

<400> 44
 ataatttgaa ggttggcaga ccagtaattt gaaggttggc agaccagttg ttgaaggttg 60
 gcagaccagt tgcgcaaggt tggcagacca gttgcgctga ttggcagacc agttgcgctg 120
 attacggcag accagttgcg ctgattactc agaccagttg cgctgattac tcttaccagt 180
 tgcgctgatt actcttagag gttgcgctga ttactcttag agaaggcgct gattactctt 240
 agagaagaag cgctgattac tcttagagaa gaaga 275

<210> 45
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 45
 aatatggtta gttccagagt gcaaatttaa aattccatac acgtttgcag catttggata 60
 ctttgaaaga tgacattggt aatgaatgac tagcccaatt gaatgactag cccaattatc 120
 cttattgact agcccaatta tccttataaa gctttgatat catactactc tgcctttgat 180
 atcatactac tctgcctttg atatcatact actctgcctt tgtgggtggg cacatatgta 240
 gacactacta gggcacatat gtagacacta ctaaa 275

<210> 46
 <211> 200
 <212> DNA
 <213> Homo sapiens

<400> 46
 tgactgggaa ttgttactgt tgtacgactg ggaattgtta ctgttgact actgggaatt 60
 gttactgttg tacttctggg aattgttact gttgtactta tgggaattgt tactgttgta 120
 cttatgggaa ttgttactgt tgtacttatt ggaattgtta ctgttgact tattcgtaca 180
 atttgggtgtt tgtattagct 200

<210> 47
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 47
 tgcactttaa ctgacttaag tggcacttta actgacttaa gtggcattaa taactgactt 60
 aagtggcatt aaacataagt ggcattaaac atttgagagc acatttgaga gctaactata 120
 ttttttataa gactactata caaactacag taagactact atacaaacta cagaggacta 180
 ctatacaaac tacagagttt aactacagag tttatgattt aaggtcagag tttatgattt 240
 aaggctacta tactttaaat aaagggtgact gggaa 275

<210> 48
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 48
 aaataatgca tccagcagta caataataat gcatccagca gtacaataaa atgcatccag 60

pctgb2014053694-seq1.txt

cagtacaata aaagttgcat ccagcagtac aataaaagta atttaatgac atgcctatat	120
gagaatttaa tgacatgcct atatgagaag atgaatatat ggcatTTGca gatTTTgaat	180
atatggcatt tgcagatttt gcagattttt atattagttg ctttgcagat ttttatatta	240
gttgctttgt agatttttat attagttgct ttgtt	275

<210> 49
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 49 atgcaatatt tgtaaggctt gttccatatt tgtaaggctt gttccaatgc aaggcttgtt	60
ccaatgccac atacttgTtc caatgccaca tacttgcagc aatgccacat acttgcagct	120
cccattgcag ctcccattct atgtgtcatc agctcccatt ctatgtgtca tcaattatgt	180
gtcatcaata gtgtcctatg gtgtcatcaa tagtgtccta tgcaacatca atagtgtcct	240
atgcaataaa attatttgca ggtcttttaa tcatt	275

<210> 50
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 50 gcttggaTgt aaaacgtaat agaaaaaaag ggagagctag gaaatggact gaacttgaca	60
gctagcagaa ttggtgcaga attggTttcc attagacatt ggTttccatt agacattact	120
cccagttcca ttagacatta ctcccaggag tactcccagg agctatatta catagtccca	180
ggagctatat tacatagtaa aataggaact ttgaattacc ctctggaact ttgaattacc	240
ctctgtgcta aattaccctc tgtgctactt ttatt	275

<210> 51
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 51 tagtctttgg tgcttggaTg taaaaaaaag ggagagctag gaaatggact gaacttgaca	60
gctagcagaa ttggtgcaga attggTttcc attagacatt ggTttccatt agacattact	120
cccagtactc ccaggagcta tattacatag ttatacatcc tacatgaccc attcaacaaa	180
gctgttttac tactaactta aataggaact ttgaattacc ctctggaact ttgaattacc	240
ctctgtgcta aattaccctc tgtgctactt ttatt	275

<210> 52
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 52 gctccaactg ttggacaaaa taaagggatt gtgccagacc ttgtgttttt tgtgccagac	60
---	----

pctgb2014053694-seq1.txt

```

cttgtgtttt ttgatgtgtt ttttgattcg aactttgcc a ttttgattcg aactttgcc      120
aggccttgcc aaggcctgga acgctgaaga gcctggaacg ctgaagaata atacttcttt      180
ccaaaaagca ttaagagggt tgtttgtctg tttcctttat cctctctgtt tcctttatcc      240
tcttggtaaa aagatttaat gtttggtga taca                                275

```

```

<210> 53
<211> 275
<212> DNA
<213> Homo sapiens

```

```

<400> 53
agtgtttatc agcagactgt aaaggagact gtaaaggcaa gtaggctctg tttgtaaatt      60
ccctccaacc tgtacgtaaa ttccctccaa cctgtacaac ccctccctca aatttgagta      120
gccaaactcaa atttgagtag ccaagtatct ttgagtagcc aagtatctag ctttgatcta      180
gctttgtgta attagacagc gctttgtgta attagacagc ttcaaaatta gacagcttca      240
agtatgtatt ttgagttgga aatgatgctt agatt                                275

```

```

<210> 54
<211> 275
<212> DNA
<213> Homo sapiens

```

```

<400> 54
gttcctggac cccacaagac ttgattagct tatgtgacaa ccctcatctt atccccatcc      60
cttctgaaag taggagttgg agctcttgca atggaattca gaacagactc tggagtgtcc      120
atttatctgg agtgtccatt tagactacac tagactacac taactagact tttaagtgtg      180
gtttggtgca agtcagaata aaattctggc tagttgaatc cacaattcat ttatatacag      240
gcttcccttt aataaacacg tttatgccta tcagc                                275

```

```

<210> 55
<211> 275
<212> DNA
<213> Homo sapiens

```

```

<400> 55
caggcttcct agaggtgtgc tatacaggtg tgctatacca tgcgtctgtc ataccatgcg      60
tctgtcgttg tgcttacagt tctttggtaa gcattgtcgt ggtaagcatt gtcgtatctg      120
gtgatcattg tcgtatctgg tgatggatta tgatggatta acatatagcc tttgttctaa      180
taaaatagtc gccttcgttt taaaatagtc gccttcgttt tctgtatctt tctcttacia      240
tctgttttag gacatctttg cttatgaaac ctgta                                275

```

```

<210> 56
<211> 275
<212> DNA
<213> Homo sapiens

```

```

<400> 56
agaggagaga acacgtctta caggaagaga acacgtctta caggatggaa aagtgtgtag      60

```

pctgb2014053694-seq1.txt

ctagggttatt tcggagtagc taggttattt cggagtggtta gggtatttcg gagggttatt	120
tatctgtttt tgattcttga gactgtgagg cttgggtgac ttctctgccc ttaaagttga	180
cttctctgcc cttaaagtaa tagtgaaatt gggtccatca gagatgttcc atcagagata	240
acctcgagtt catcagagat aacctcgagt tcttg	275

<210> 57
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 57 attagatggg gatttgacgt ctgggttaat ttactgtagg actagtggca taggactagt	60
ggcagtgagt ctcagtggc agtgagtctc atattgctgt agtgagtctc atattgctgt	120
catatagtag cctaaagacc acgctttgag cacgctttga gaaactcgta aagggtgaaac	180
tcgtaaaggt ttgtcataga taattttaag ctttagttgc ctactaagta ggatgcagta	240
caagtttgtc gatgcagtac aagtttgtcc ttac	275

<210> 58
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 58 aaaaaaccct gtgccttaca caatgaaacc ctgtgcctta cacaatgtgt ccctgtgcct	60
tacacaatgt gtatacacia tgtgtatatt gagttgatac aaaaggatgat tgagacacag	120
cctctaaagg tgattgagac acagcctctg ctctgtcctg agcttagtct agtagctgag	180
cttagtctag taggaggtga cttagtctag taggaggtga cagatctagt aggaggtgac	240
agatttgtaa aataatagat ttgaagttgc tgagc	275

<210> 59
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 59 aaaaaaccct gtgccttaca caatgaaacc ctgtgcctta cacaatgtgt ggtattatac	60
aatcattcat tcagcagcaa acattcactt actgagcacc actgagcacc tactaatgtt	120
ccaggcctac taatgttcca ggtactgatt aaaggatgatt gagacacagc ctctgacaca	180
gcctctgtcc tgagcttagt ctctgtcctg agcttagtct agtagctagt aggaggtgac	240
agatttgtaa aataatagat ttgaagttgc tgagc	275

<210> 60
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 60 atactgactt gtttgcattg ggcatgcata tatactactc ttgcgagcag tactactctt	60
---	----

pctgb2014053694-seq1.txt

gcgagcagga ggcctcgagc aggaggcctg aaatctccta agcctggcaa caagggaaat	120
cccgaaaatc ccgagattgt aggcaagagg gatgctttca aaaagcctgg aggggaagtgg	180
agatgattat gtttaacgca atgtttaacg cattccttct ttggtgagct gctagcatag	240
tatggattaa aatagccaga ctatactgac ttgtt	275

<210> 61
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 61 gggaataccg agtgctcaat tttattttta tagttttctca tagttttcttc tctcatagtt	60
tcttcatggc cacagtagtt tcttcatggc cacagataaa cacagataaa cccctgaaac	120
caatcataac cctgtagttc tcactattgg ccctgtagtt ctactattg ggtcacacta	180
ttgggtcatt aacacagtca acagtcatta aatgcaagca ggtctatgca agcaggtcta	240
gtattgacta gtctagtatt gactagccct tctgg	275

<210> 62
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 62 gaggagtct gagattgact actaagacta ctaatgtctg ccatggaagt ggagcagtta	60
tggtacgagt gtcattgagt tcatatTTTT gccagttctg atttttgcca gttctgactg	120
aaggcaaagc ccaggatctt tctagcctct gatctttcta gcctctaccc agtgtcagtg	180
tcactgactt ttagtctcat tagctcattg ccctagttaa attggtacaa tgccaaatcc	240
tatcaaaggg atcaaaggg attggctgcc tgtag	275

<210> 63
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 63 atcgggctgt ggtgaccgca acaagtcggg ctgtggtgac cgcaacaagc gggagcacac	60
cgtttcctga ttgctagcac accgtttcct gattgctcag ccctccttgg tgattccgca	120
gtgagtcctt ggtgattccg cagttagaga aaggcaagat tgtgtcccat tccccgcaag	180
attgtgtccc attccccag agtccgcgc caggatggcg aggaaaggat ggcgaggaaac	240
tcaataaagt tggcgaggaa ctcaataaag tgctt	275

<210> 64
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 64 acttttgcca agcaccatac ttaaggccaa gcaccatact taagactcaa tacttaagac	60
---	----

pctgb2014053694-seq1.txt

tcaacttttt	tgcaatatca	gacaaagcac	tgtctttaag	cactagatcc	cttcttctga	120
aaatcgatcc	cttcttctga	aaatcactgt	atattttttt	agcattcact	gcatgagtta	180
agtctgcttt	cctatccctt	taagtctgct	ttcctatccc	ttatactgct	ttcctatccc	240
ttatacagaa	atcccttata	cagaagctat	ccatt			275

<210> 65
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 65						
cttgtcctcc	aagttatggg	ggatctcttg	tactgacaat	ctgtgttttc	gtactgacaa	60
tctgtgtttt	ccaggctgtg	ttttccagga	gttacgtcaa	gttttccagg	agttacgtca	120
aactaaggag	ttacgtcaaa	ctacctgtac	gttacgtcaa	actacctgta	ctgggtgtcaa	180
actacctgta	ctggtttaaa	atacatattt	tgtgctaatc	ttgttagtgt	ttgtctagca	240
tgtttgttta	gtctagcatg	tttgtttaat	ccttt			275

<210> 66
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 66						
ttcctccaaa	acaaacgaca	gcaacaaacg	acagcaacga	aaactcctta	gacagcaacg	60
aaaactcctt	aatcagaaaa	ctccttaatc	agaacactga	tcagaacact	gatccaatga	120
ggaatcagga	gaacttagtg	caagactaca	gagaacttag	tgcaagacta	caggaagact	180
acaggagtta	acagatggcc	tacaggagtt	aacagatggc	cagctgagtt	tatatcaaat	240
cctgaagaaa	atatcaaadc	ctgaagaaat	aagcc			275

<210> 67
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 67						
ggcacaagaa	attggagctg	gagaagctgg	agaagattga	tgaaagtgca	gtgcaggtgt	60
gtaaggaaat	agaacaaata	gaacagtctg	ctgggagtca	atctccaggg	tgacggaacc	120
cagtggtgac	ggaaccaggt	gtattacctg	gtgtattacc	tgctggaacc	aaggagtagg	180
ttactagtga	ataccccaat	gttactagtg	aataccccaa	tggttgaata	ccccaatggt	240
ttctccaatt	ccaatggttt	ctccaattat	gccc			275

<210> 68
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 68						
atggtgaaac	aagggatgtc	tgattaacaa	gggatgtctg	attggaagta	gatattttatt	60

pctgb2014053694-seq1.txt

taggttctag gacatctagg acattagtat cagtgaggac tcagtgagga cagtaatttc	120
ctgctgtaat ttcttgcttg tttgtatttc gtatttcagt gatcacatac acttcgtgat	180
cacatacact tctttacctg acacttcttt acctgataac gtctctacct gataacgtct	240
ctcttctcta ctacttaaag gtatcccttg aattc	275

<210> 69
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 69 ttgcaaacc ctggttgctg tatttaaacc ctggttgctg tatttgttca ctggttgctg	60
tatttgttca gtgacagatt atttgcacga actggactgt gaactggact gtgtgcaacg	120
cttttttttg gagaatgatg tccccgttgt aatgatgtcc ccgttgtagt tatgatcccc	180
gttgtagtga tgagtggctt cactgtata gaagggtttt gtagcgtagc ctgaatgtct	240
tactgtgatc cctgaatgtc ttactgtgat caatt	275

<210> 70
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 70 tgcaggagac attggtattc tgggcgcagg agacattggt attctgggca tgcactttaa	60
ctgacttaag tggcacttta actgacttaa gtggcattaa taactgactt aagtggcatt	120
aaacataagt ggcattaaac atttgagagc acatttgaga gctaactata tttttaaga	180
ctactataca aactacagag gactactata caaactacag agtttaacta cagagtttat	240
gatttaaggt cagagtttat gatttaaggt actta	275

<210> 71
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 71 aagaatttgg cacagttact cacttgaatt tggcacagtt actcactttg tgctgaatac	60
gctgaagtaa atcctaatac gctgaagtaa atccttgttc gctgaagtaa atccttgttc	120
actgatgaag tctttcaatt gagctggttg gaagtctttc aattgagctg gttgattgaa	180
aaatgctcag ttctaactaa taatgaaatg gatttccag taggggcata tcacctgtat	240
agtagttata tcacctgtat agtagttata tgcatt	275

<210> 72
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 72 tccaaaagca agtagccaaa gccgtaagca agtagccaaa gccgttgcca caagtagcca	60
---	----

pctgb2014053694-seq1.txt

aagccgttgc caaactagcc aaagccgttg ccaaacccca atgggccctt tatttatgac	120
gacttggccc tttatttatg acgactttat atttatgacg actttattta ttctatatat	180
atattgggtc gtctgcttcc atatatattg ggtcgtctgc ttccctattg ggtcgtctgc	240
ttcccttgta ttttgtaata ttgaaaacga cgata	275

<210> 73
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 73 atacaggtgc aactcaatcc ccgttgtgca actcaatccc cgtaaataaa tgtaggtata	60
ggcattctac cctttgtata ggcattctac cctttgaaat taccctttga aatagctgtg	120
tcccactttg aaatagctgt gtcccaacct agaaatatcc aagttgtcct tgaatgttgt	180
ccttgaattg tctaaccatg gtctaaccat ggacataaac agttgatgga cataaacagt	240
tgtctccctt tactgtgtag aatactttga cttaa	275

<210> 74
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 74 aagagagaac aactactgtg atcagagaac aactactgtg atcaggctat actactgtga	60
tcaggctatg tatggatgta tggaaatacag tgttattttc aattagctgt gtgaaatacc	120
agtgttagct gtgtgaaata ccagtgtggt atttatttaa gcttatgtca gaccttattt	180
aagcttatgt cagacctatt aagcttatgt cagacctatt tgacacagac ctatttgaca	240
taacactata gacataacac tataaagggtt gacaa	275

<210> 75
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 75 acatgtactt tataagtatt ggtttcatgt actttataag tattggtttg atgtacttta	60
taagtattgg tttggtgtac tttataagta ttggtttggg tactttataa gtattggttt	120
gggtgacttt ataagtattg gtttgggtgt tttataagta ttggtttggg tgttcttata	180
agtattgggt tgggtgttcc tataagtatt ggtttgggtg ttcctataag tattggtttg	240
ggtgttcctt taagtattgg tttgggtgtt ccttc	275

<210> 76
 <211> 225
 <212> DNA
 <213> Homo sapiens

<400> 76 acagccaaaa ctccacagtc aatatcagcc aaaactccac agtcaatatt agccaaaact	60
---	----

pctgb2014053694-seq1.txt

ccacagtcaa tattagccaa aactccacag tcaatattag ccaaaactcc acagtcaata	120
ttagtgatgt ttttatgtgc tctccaaatt atgtttttat gtgctctcca aattttgttt	180
ttatgtgctc tccaaatfff ttactgtttc tgattgtatg gaaat	225

<210> 77
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 77 tctgtctttt ataaagattc caccttaaag attccacctc cagtcctctc aaagattcca	60
cctccagtcc tctctcctac tcaggccctt gaggctatta tactcaggcc cttgaggcta	120
ttaggtcagg cccttgaggc tattaggaga ggcccttgag gctattagga gatgccctt	180
gaggctatta ggagatgctt agaactcaac aaaatcccaa tccaagaact caacaaaatc	240
ccaatccaag caacaaaatc ccaatccaag tcaaa	275

<210> 78
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 78 gaaatcctat gcctagaaac caattacatg actgctatft agggaaggcc gggaaggcct	60
aatctttgta acttaciaag tgattgacag agtgggctct gcctctgtca cttctagtfg	120
tggaattcta gttgtggaat ccttggaag tggcaagtgt cttaaactct ttatgactct	180
ttatgccatt gagttattat attattccca tctgttatta aagaggatta aagtatatgc	240
ctcagaggga gcctcagagg gatgtttgag tgtaa	275

<210> 79
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 79 atctacatta aagccacaag tcaccaagcc acaagtcacc ctttgctgaa ccacaagtca	60
ccctttgctg aagtcaagtc agtattagta gttggaagca agcagtgtgt tattcttgac	120
cccatgtggc acttattaag tagcttgctt aagtagcttg cttttccata attatttcca	180
taattatggc ctagcttttt agctttttaa aacactactat gaacaacacc acaagcatag	240
agttttccaa aggaaaccaa ttatactgaa tcagg	275

<210> 80
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 80 ttftaataca acaagatcaa ctagattaat acaacaagat caactagaag gatcaactag	60
aagaattcaa ctgtccaact agaagaattc aactgtctca actagaagaa ttcaactgtc	120

pctgb2014053694-seq1.txt

tcaagtcaag gtacccaat atatctcgca aaggtacccc aatatatctc gcaatggtac	180
cccaatatat ctcgcaattc tacccaata tatctcgcaa ttccacccaa tatatctcg	240
aattccaaac aatatatctc gcaattccaa acttt	275

<210> 81
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 81	
aaaaaacctt ttggtgctcc agtgtaaaac tttttggtgc tccagtgtag actttttggt	60
gctccagtgt agggcaaata aactatcagc ttggatggtc aataaactat cagcttggat	120
ggtcagcttg gatggctact tgaatagaag tggatggcca cttgaataga agatgtgaat	180
agaagatggt tatacacagt ttgtctcttg tatggcaaaa taattgctct tgtatggcaa	240
aataattagt gtatggcaaa ataattagt agttt	275

<210> 82
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 82	
cattcacctt tgtaacagt atttctcact ttgttaaca gtatttctct tatagtggat	60
gatatacaca gtggcagtgg atgatataca cagtggcaaa acacagtggc aaaacaaaag	120
tacataaaaa tgtcactata tcttccatt gtcactatat ctcccatctt aacatatatc	180
ttcccatctt acattgtttt tgtatattgg gtgtagattt ctgacattgg gtgtagattt	240
ctgacatcaa tgggtgtaga tttctgacat caaaa	275

<210> 83
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 83	
gaaggtgctc actaacagag gtacaggtgc tcactaacag aggtacatta ctaacagagg	60
tacattactg caatgtaaca gaggtacatt actgcaatgt taacagttaa acaagctggt	120
tacagcagtt aaacaagctg ttacagttt gagctattta aagcttatta tattttgcat	180
tatgtgtaca gtattggaca gcattatgtg tacagtattg gacaagtgtg cagtattgga	240
caaaggattt tacagtattg gacaaaggat tttat	275

<210> 84
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 84	
gctgtcctcc gtattacgtg accggctccg tattacgtga ccggcaaata tacgtgaccg	60
gcaaataaat ctcatgtaga atgcaaaatg ttggcacgtc gaatgcaaaa tgttggcacg	120

pctgb2014053694-seq1.txt

tcaaagttgg cacgtcaaaa atatgaatgt tgaatgtgta gacaactgta gttgtgacaa	180
ctgtagttgt gctcagtttg ggggaagtgta ttttactctg atcaaggaag tgtattttac	240
tctgatcaaa taatgctgga atactcaaga attgc	275

<210> 85
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 85 gatatggacg ttatcattgg tctggatgga cgttatcatt ggtctggtga tggctggtg	60
agatgtttca tatttattat gagttaatgc tgcctgtgtc agttaatgct gcctgtgtct	120
atgggggtcta tgggggttctg tcttctttga tgtcttcttt gatagccatc tattcgatag	180
ccatctattc atctggatca ctattcatct ggatcatggg acccttggtt tggagttgag	240
atatcagtct gttgagatat cagtctcgga aactt	275

<210> 86
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 86 tcacaggggc ctcaagaatt ggtatacagg ggcctcaaga attggtatgt ctcaagaatt	60
ggtatgtatg atgtgtgatg tgatctggtc cagccagggc atgtgatctg gtccagccag	120
ggccttgatc tgggtccagcc agggcctggc cagctctcta ggtttgatat gacttgctct	180
ctaggtttga tatgacttta ctctctaggt ttgatatgac tttagtttgt caatatagat	240
ggtaggaagc gtcaatatag atggtaggaa gcaga	275

<210> 87
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 87 gatggatgca actgaagcag agtgtggtgc ccagatgtgc gctattagat gtgcgctatt	60
agatgtttct ctgattttct ctgataatgt cccaatcat gataatgtcc ccaatcatac	120
caggggtccc caatcatacc agggagactg cagggagact ggcattgacg agaacttgac	180
gagaactcag gtggaggctt tcagctttgc aaagagccac cctagcagct gaccgcatgg	240
gtgtgagcca ctactcaata aaagcgaagg tggac	275

<210> 88
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 88 aagatgtcat ttgcaataga gcagtgtcat ttgcaataga gcagtcatt gaattttcac	60
tggacctgtg atgtcaattt tctactggacc tgtgatgtca ctttttgtgc acagattatg	120

pctgb2014053694-seq1.txt

atgaattatc actctgcctg tgtatagtca cactctgcct gtgtatagtc agatagtata	180
gtcagatagt ccatgcgaag cagatagtcc atgcgaaggc tgtattccat gcgaaggctg	240
tatatattga ttataaagtg tgtaagttac cagtt	275

<210> 89
 <211> 225
 <212> DNA
 <213> Homo sapiens

<400> 89 aaaagactga gcatgctcac tttttaaacc ctttgtattg tagataagag aaagactgag	60
catgctcact tttttaaccc tttgtattgt agataagagg aagactgagc atgctcactt	120
ttttaacaaa ccctttgtat tgtagataag agacaaaccc tttgtattgt agatacaaac	180
ccctttgtatt gtagataaga gacaaaccct ttgtattgta gataa	225

<210> 90
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 90 acttaccacc actaaactgc gagagagcag actgccggta acgcgcggcc ccgctgcagt	60
ttcttgggac ataggctaaa ctgcgagaga agctaaacgt gagcgcaaag aagctacagc	120
ctggaggaca gcagactgcc ggtaacgcgc ggacttacca ccactaaact gcgagggaga	180
gactcagcaa cgaccatac ggccgggagag actcagcaac gaccgcgttc ttgggacata	240
ggagcgcaaa tgcccgtgc agtttcttgg gacat	275

<210> 91
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 91 aaagtttaga tctgtccctt aataaaaatc ttaatcatta tccttctaag aggatacaaa	60
tttagtgctc ttaacatcaa aatcttaatc attatccttc atcagaaagt ttagatctgt	120
cccttatcat tatccttcta agaggataca ctaagaggat acaaatttag tgctccttaa	180
tcattatcct tctaagagga gaaagtttag atctgtccct taatatacaa atttagtgct	240
cttaacttgt ttgcttaaac taatgaacaa atatg	275

<210> 92
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 92 aaatcttaat cattatcctt ctaagaggat acaaatttag tgctcttaac atcagaaagt	60
ttagatctgt cccttccaag agttaagata tccctaattgt cctgaaagca gtaaccaaga	120
gttaacttaa tcattatcct tctaagagga gaaagtttag atctgtccct taatagttaa	180

pctgb2014053694-seq1.txt

gatatcccta atgttttgct taaccaagag ttaagatatc cctaataaga tatcccta	240
gttttgctta taaaaattta gtgctcttaa cttgt	275

<210> 93
 <211> 275
 <212> DNA
 <213> Homo sapiens

<400> 93 actctttact caggagggcc ttgcaagagt aggtgctgcc tagcagccct agcagccctt	60
cgtggccact ctttaagggtg ctgcctagca gcccttcgtg atgacacgtg tcaagatgcc	120
cttccatgtg ttcattaagg gctcaataaa ccagtatgaa ctgccgtgaa gtcaagaggt	180
cccagtatga actgccgtga tctttactca ggagggcctt gcagattacc atatgacacg	240
tgtcaagatg ttatgtgttc attaagggt caata	275