

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/096148 A1

(43) International Publication Date

26 June 2014 (26.06.2014)

(51) International Patent Classification:

A61K 38/26 (2006.01) C07K 14/605 (2006.01)

(21) International Application Number:

PCT/EP2013/077310

(22) International Filing Date:

19 December 2013 (19.12.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12306647.4 21 December 2012 (21.12.2012) EP

(71) Applicant: SANOFI [FR/FR]; 54 rue La Boétie, F-75008 Paris (FR).

(72) Inventors: HAACK, Torsten; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). WAGNER, Michael; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). HENKEL, Bernd; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). STENGELIN, Siegfried; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). EVERS, Andreas; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). LORENZ, Martin; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). LORENZ, Katrin; c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE).

(74) Agents: WEIB, Wolfgang et al.; Weickmann & Weickmann, Patentanwälte, Richard-Strauss-Str. 80, 81679 München (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))



WO 2014/096148 A1

(54) Title: FUNCTIONALIZED EXENDIN-4 DERIVATIVES

(57) Abstract: The present invention relates to exendin-4 derivatives and their medical use, for example in the treatment of disorders of the metabolic syndrome, including diabetes and obesity, as well as reduction of excess food intake.

Functionalized Exendin-4 Derivatives

5 Description

FIELD OF THE INVENTION

The present invention relates to exendin-4 peptide analogues which activate
10 the glucagon-like peptide 1 (GLP-1) and the glucose-dependent insulinotropic polypeptide (GIP) receptor and optionally the glucagon receptor (GCG) and their medical use, for example in the treatment of disorders of the metabolic syndrome, including diabetes and obesity, as well as reduction of excess food intake.

15

BACKGROUND OF THE INVENTION

Exendin-4 is a 39 amino acid peptide which is produced by the salivary glands of the Gila monster (*Heloderma suspectum*) (Eng J. et al., *J. Biol. 20 Chem.*, 267:7402-05, 1992). Exendin-4 is an activator of the glucagon-like peptide-1 (GLP-1) receptor, whereas it shows only very low activation of the GIP receptor and does not activate the glucagon receptor (see Table 1).

25 Table 1: Potencies of exendin-4 at human GLP-1, GIP and Glucagon receptors (indicated in pM) at increasing concentrations and measuring the formed cAMP as described in Methods.

SEQ ID NO:	peptide	EC50 hGLP-1 R [pM]	EC50 hGIP R [pM]	EC50 hGlucagon R [pM]
1	exendin-4	0.4	12500.0	>10000000

Exendin-4 shares many of the glucoregulatory actions observed with GLP-1. Clinical and non-clinical studies have shown that exendin-4 has several

- 2 -

beneficial antidiabetic properties including a glucose dependent enhancement in insulin synthesis and secretion, glucose dependent suppression of glucagon secretion, slowing down gastric emptying, reduction of food intake and body weight, and an increase in beta-cell mass and 5 markers of beta cell function (Gentilella R et al., *Diabetes Obes Metab.*, 11:544-56, 2009; Norris SL et al., *Diabet Med.*, 26:837-46, 2009; Bunck MC et al., *Diabetes Care.*, 34:2041-7, 2011).

10 These effects are beneficial not only for diabetics but also for patients suffering from obesity. Patients with obesity have a higher risk of getting diabetes, hypertension, hyperlipidemia, cardiovascular and musculoskeletal diseases.

15 Relative to GLP-1 and GIP, exendin-4 is more resistant to cleavage by dipeptidyl peptidase-4 (DPP4) resulting in a longer half-life and duration of action in vivo (Eng J., *Diabetes*, 45 (Suppl 2):152A (abstract 554), 1996; Deacon CF, *Horm Metab Res*, 36: 761-5, 2004).

20 Exendin-4 was also shown to be much more stable towards degradation by neutral endopeptidase (NEP), when compared to GLP-1, glucagon or oxyntomodulin (Druce MR et al., *Endocrinology*, 150(4), 1712-1721, 2009).

25 Nevertheless, exendin-4 is chemically labile due to methionine oxidation in position 14 (Hargrove DM et al., *Regul. Pept.*, 141: 113-9, 2007) as well as deamidation and isomerization of asparagine in position 28 (WO 2004/035623).

The amino acid sequence of exendin-4 is shown as SEQ ID NO: 1:

30 HGETFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH₂

The amino acid sequence of GLP-1(7-36)-amide is shown as SEQ ID NO: 2:

- 3 -

HAEGTFTSDVSSYLEGQAAKEFIAWLVKGR-NH₂

Liraglutide is a marketed chemically modified GLP-1 analogue in which, among other modifications, a fatty acid is linked to a lysine in position 20

5 leading to a prolonged duration of action (Drucker DJ et al, Nature Drug Disc. Rev. 9, 267-268, 2010; Buse, JB et al., Lancet, 374:39-47, 2009).

The amino acid sequence of Liraglutide is shown as SEQ ID NO: 3:

10 HAEGTFTSDVSSYLEGQAAK((S)-4-Carboxy-4-hexadecanoylamino-butyryl-
)EFLAWLVRGRG-OH

GIP (glucose-dependent insulinotropic polypeptide) is a 42 amino acid peptide that is released from intestinal K-cells following food intake. GIP and

15 GLP-1 are the two gut enteroendocrine cell-derived hormones accounting for the incretin effect, which accounts for over 70% of the insulin response to an oral glucose challenge (Baggio LL, Drucker DJ. Biology of incretins: GLP-1 and GIP. Gastroenterology 2007; 132: 2131-2157).

20 GIP's amino acid sequence is shown as SEQ ID NO: 4:

YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGKKNDWKHNITQ-OH

Glucagon is a 29-amino acid peptide which is released into the bloodstream

25 when circulating glucose is low. Glucagon's amino acid sequence is shown in SEQ ID NO: 5:

HSQGTFTSDYSKYLDSRRAQDFVQWLMNT-OH

30 During hypoglycemia, when blood glucose levels drop below normal, glucagon signals the liver to break down glycogen and release glucose, causing an increase of blood glucose levels to reach a normal level.

- 4 -

Hypoglycemia is a common side effect of insulin treated patients with hyperglycemia (elevated blood glucose levels) due to diabetes. Thus, glucagon's most predominant role in glucose regulation is to counteract insulin action and maintain blood glucose levels.

5

Holst (Holst, J. J. Physiol. Rev. 2007, 87, 1409) and Meier (Meier, J. J. Nat. Rev. Endocrinol. 2012, 8, 728) describe that GLP-1 receptor agonists, such as GLP-1, liraglutide and exendin-4, improve glycemic control in patients with T2DM by reducing fasting and postprandial glucose (FPG and PPG).

10 Peptides which bind and activate the GLP-1 receptor are described in patent applications WO1998008871, WO2008081418 and WO2008023050, the contents of which are herein incorporated by reference.

It has been described that dual activation of the GLP-1 and GIP receptors, 15 e.g. by combining the actions of GLP-1 and GIP in one preparation, leads to a therapeutic principle with significantly better reduction of blood glucose levels, increased insulin secretion and reduced body weight in mice with T2DM and obesity compared to the marketed GLP-1 agonist liraglutide (e.g. VA Gault et al., Clin Sci (Lond), 121, 107-117, 2011). Native GLP-1 and GIP 20 were proven in humans following co-infusion to interact in an additive manner with a significantly increased insulinotropic effect compared to GLP-1 alone (MA Nauck et al., J. Clin. Endocrinol. Metab., 76, 912-917, 1993).

Designing hybrid molecules which combine agonism on the GLP-1 receptor, 25 the GIP receptor and the glucagon receptor offers the therapeutic potential to achieve significantly better reduction of blood glucose levels, increased insulin secretion and an even more pronounced significant effect on body weight reduction compared to the marketed GLP-1 agonist liraglutide (e.g. VA Gault et al., Clin Sci (Lond), 121, 107-117, 2011).

30

Compounds of this invention are exendin-4 derivatives, which show agonistic activity at the GLP-1 and the GIP receptor and optionally the glucagon

- 5 -

receptor and which have – among others – preferably the following modifications: Tyr at position 1 and Ile at position 12.

Surprisingly, it was found that the modification of the selective GLP-1R

5 agonist Exendin-4 by Tyr in position 1 and Ile in position 12 results in a peptide with high dual activity at the GLP-1 and GIP receptors. This observation is surprising, since the same modification in other GLP-1 agonists, such as GLP-1 itself, does not result in high activity at the GIP receptor, as shown in Table 2.

10

Table 2: Potencies of exendin-4 and GLP-1 peptide analogues at GLP-1 and GIP receptors (indicated in pM) at increasing concentrations and measuring the formed cAMP as described in Methods.

SEQ ID NO:	peptide	EC50 [pM]	hGIP R	EC50 [pM]	hGLP-1 R
6	Tyr(1)Ile(12)-exendin-4	93.9		1.3	
7	Tyr(1)Ile(12)-GLP1	3660.0		5.0	

15 Peptides which bind and activate both the GIP and the GLP-1 receptor and optionally the glucagon receptor, and improve glycaemic control, suppress body weight gain and reduce food intake are described in patent applications WO 2011/119657 A1, WO 2012/138941 A1, WO 2010/011439 A2, WO 2010/148089 A1, WO 2011/094337 A1, WO 2012/088116 A2, the
20 contents of which are herein incorporated by reference. These applications disclose that mixed agonists of the GLP-1 receptor, the GIP receptor and optionally the glucagon receptor can be designed as analogues of the native GIP or glucagon sequences.

25 Compounds of this invention are exendin-4 peptide analogues comprising leucine in position 10 and glutamine in position 13. Krstenansky et al. (Biochemistry, 25, 3833-3839, 1986) show the importance of residues 10 to 13 of glucagon for its receptor interactions and activation of adenylate

- 6 -

5 cyclase. In the exendin-4 peptide analogues of this invention, several of the underlying residues are different from said of glucagon. In particular, residues Tyr10 and Tyr13, are replaced by leucine in position 10 and glutamine, a non-aromatic polar amino acid, in position 13. This replacement, especially in combination with isoleucine in position 23 and glutamate in position 24 leads to exendin-4 derivatives with potentially improved biophysical properties as solubility or aggregation behavior in solution. The non-conservative replacement of an aromatic amino acid with a polar amino acid in position 13 of an exendin-4 analogue surprisingly leads 10 to peptides with high activity on the GIP receptor and optionally on the glucagon receptor.

15 Furthermore, compounds of this invention are exendin-4 derivatives with fatty acid acylated residues in position 14. This fatty acid functionalization in position 14 results in an improved pharmacokinetic profile. Surprisingly, the fatty acid functionalization in position 14 also leads to peptides with a significantly higher GIPR activity, for example those shown in Example 9, Table 8.

20

BRIEF SUMMARY OF THE INVENTION

25 Provided herein are exendin-4 analogues which potently activate the GLP-1 and the GIP receptor and optionally the glucagon receptor. In these exendin-4 analogues – among other substitutions - methionine at position 14 is replaced by an amino acid carrying an –NH₂ group in the side-chain, which is further substituted with a lipophilic side-chain (e.g. a fatty acid optionally combined with a linker).

30 The invention provides a peptidic compound having the formula (I):



wherein Z is a peptide moiety having the formula (II)

5 Tyr-Aib-X3-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-X12-Gln-X14-X15-X16-
 X17-X18-X19-X20-X21-Phe-Ile-Glu-Trp-Leu-Lys-X28-X29-Gly-Pro-Ser-
 Ser-Gly-Ala-Pro-Pro-Pro-Ser-X40 (II)

X3 represents an amino acid residue selected from Gln, Glu and His,

X12 represents an amino acid residue selected from Ile and Lys,

10 X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, -C(O)O-R⁵, -C(O)NH-R⁵, -S(O)₂-R⁵ or R⁵, preferably by -C(O)-R⁵, wherein R⁵ may be a moiety comprising up to 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P,

15

X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Lys, Ile, Glu,

20 Gln, Leu, Aib, Tyr and Ala,

X18 represents an amino acid residue selected from Ala, Arg, Lys, Aib, Leu and Tyr,

X19 represents an amino acid residue selected from Ala, Val, Gln and Aib,

25 X20 represents an amino acid residue selected from Gln, Aib, Phe, Leu, Lys, His, Arg, Pip, (S)MeLys, (R)MeLys, (S)MeOrn and (R)MeOrn, X21 represents an amino acid residue selected from Asp, Glu, Leu and Tyr,

X28 represents an amino acid residue selected from Asn, Ala, Arg, Lys, Asp and Ser,

X29 represents an amino acid residue selected from Gly, Thr, Aib, D-Ala and Ala,

- 8 -

X40 is absent or represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is optionally functionalized by -C(O)-R⁵, -C(O)O-R⁵, -C(O)NH-R⁵, -S(O)₂-R⁵ or R⁵, preferably by -C(O)-R⁵, wherein R⁵ may be a moiety comprising up to 5 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P,

R¹ represents NH₂,

R² represents OH or NH₂.

10

or a salt or solvate thereof.

The compounds of the invention are GLP-1 and GIP receptor agonists and 15 optionally glucagon receptor agonists as determined by the observation that they are capable of stimulating intracellular cAMP formation. In vitro potency determination in cellular assays of agonists is quantified by determining the concentrations that cause 50% activation of maximal response (EC50) as described in Methods.

20 In certain embodiments, the invention therefore provides a peptidic compound having the formula (I):

R¹ - Z - R² (I)

25 wherein Z is a peptide moiety having the formula (II)

Tyr-Aib-X3-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-X12-Gln-X14-X15-X16-
X17-X18-X19-X20-X21-Phe-Ile-Glu-Trp-Leu-Lys-X28-X29-Gly-Pro-Ser-
Ser-Gly-Ala-Pro-Pro-Ser-X40 (II)

30

X3 represents an amino acid residue selected from Gln, Glu and His,
X12 represents an amino acid residue selected from Ile and Lys,

- 9 -

X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, -C(O)O-R⁵, -C(O)NH-R⁵, -S(O)₂-R⁵ or R⁵, preferably by -C(O)-R⁵, wherein R⁵ is a moiety comprising up to 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P,
5 X15 represents an amino acid residue selected from Asp and Glu,
X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,
X17 represents an amino acid residue selected from Arg, Lys, Ile, Glu,
10 Gln, Leu, Aib, Tyr and Ala,
X18 represents an amino acid residue selected from Ala, Arg, Lys, Aib, Leu and Tyr,
X19 represents an amino acid residue selected from Ala, Val, Gln and Aib,
15 X20 represents an amino acid residue selected from Gln, Aib, Phe, Leu, Lys, His, Arg, Pip, (S)MeLys, (R)MeLys, (S)MeOrn and (R)MeOrn,
X21 represents an amino acid residue selected from Asp, Glu, Leu and Tyr,
X28 represents an amino acid residue selected from Asn, Ala, Arg, Lys,
20 Aib and Ser,
X29 represents an amino acid residue selected from Gly, Thr, Aib, D-Ala and Ala,
X40 is absent or represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is optionally functionalized by -C(O)-R⁵, -C(O)O-R⁵, -C(O)NH-R⁵, -S(O)₂-R⁵ or R⁵,
25 preferably by -C(O)-R⁵, wherein R⁵ may be a moiety comprising up to 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P,
30 R¹ represents NH₂,
R² represents OH or NH₂.

- 10 -

or a salt or solvate thereof, wherein the peptidic compound has a relative activity of at least 0.04%, preferably at least 0.08%, more preferably at least 0.2% compared to that of natural GIP at the GIP receptor.

5

In addition, the peptidic compound, particularly with a lysine at position 14 which is further substituted with a lipophilic residue, exhibits a relative activity of at least 0.07%, preferably at least 0.1%, more preferably at least 0.14%, more preferably at least 0.35% and even more preferably at least 10 0.4% compared to that of GLP-1(7-36) at the GLP-1 receptor.

In addition, the peptidic compound, particularly with a lysine at position 14 which is further substituted with a lipophilic residue, exhibits a relative activity of at least 0.04% (i.e. $EC_{50} < 1000$ pM), more preferably 0.08% (i.e.

15 $EC_{50} < 500$ pM) and even more preferably 0.2% (i.e. $EC_{50} < 200$ pM) compared to that of natural GIP at the GIP receptor ($EC_{50} = 0.4$ pM).

20 Optionally, in some embodiments, the peptidic compound, particularly with a lysine at position 14 which is further substituted with a lipophilic residue, exhibits a relative activity of at least 0.1%, preferably at least 0.2%, more preferably at least 0.3%, more preferably at least 0.4% and even more preferably at least 0.5% compared to that of natural glucagon at the glucagon receptor.

25 The term "activity" as used herein preferably refers to the capability of a compound to activate the human GLP-1 receptor, the human GIP receptor and optionally the human glucagon receptor. More preferably the term "activity" as used herein refers to the capability of a compound to stimulate intracellular cAMP formation. The term "relative activity" as used herein is 30 understood to refer to the capability of a compound to activate a receptor in a certain ratio as compared to another receptor agonist or as compared to another receptor. The activation of the receptors by the agonists (e.g. by

- 11 -

measuring the cAMP level) is determined as described herein, e.g. as described in the examples.

According to one embodiment, the compounds of the invention have an EC₅₀ for hGLP-1 receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less.

According to one embodiment, the compounds of the invention have an EC₅₀ for hGIP receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less.

According to another embodiment, the compounds of the invention have optionally an EC₅₀ for hGlucagon receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less.

According to another embodiment, the compounds of the invention have an EC₅₀ for hGLP-1 receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more

- 12 -

preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less, and/or an EC₅₀ for hGIP receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less, and/or 5 optionally an EC₅₀ for hGlucagon receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less. 10

15

In still another embodiment, the EC₅₀ for both receptors, i.e. for the hGLP-1 receptor and for the hGIP receptor, is 500 pM or less, more preferably 200 pM or less, more preferably 150 pM or less, more preferably 100 pM or less, more preferably 90 pM or less, more preferably 80 pM or less, more 20 preferably 70 pM or less, more preferably 60 pM or less, more preferably 50 pM or less, more preferably 40 pM or less, more preferably 30 pM or less, more preferably 20 pM or less.

In still another embodiment, the EC₅₀ for all three receptors, i.e. for the 25 hGLP-1 receptor, for the hGIP receptor and for the hGlucagon receptor, is 500 pM or less, more preferably 200 pM or less, more preferably 150 pM or less, more preferably 100 pM or less, more preferably 90 pM or less, more preferably 80 pM or less, more preferably 70 pM or less, more preferably 60 pM or less, more preferably 50 pM or less, more preferably 40 pM or less, 30 more preferably 30 pM or less, more preferably 20 pM or less.

The EC₅₀ for hGLP-1 receptor, hGIP receptor and hGlucagon receptor may

- 13 -

be determined as described in the Methods herein and as used to generate the results described in Example 9.

5 The compounds of the invention have the ability to reduce the intestinal passage, to increase the gastric content and/or to reduce the food intake of a patient. These activities of the compounds of the invention can be assessed in animal models known to the skilled person and also described herein in the Methods. The results of such experiments are described in Examples 11 and 12. Preferred compounds of the invention may increase 10 the gastric content of mice, preferably of female NMRI-mice, if administered as a single dose, preferably subcutaneous dose, of 0.02 mg/kg body weight by at least 25%, more preferably by at least 30%, more preferably by at least 40%, more preferably by at least 50%, more preferably by at least 60%, more preferably by at least 70%, more preferably by at least 80%.

15 Preferably, this result is measured 1 h after administration of the respective compound and 30 mins after administration of a bolus, and/or reduces intestinal passage of mice, preferably of female NMRI-mice, if administered as a single dose, preferably subcutaneous dose, of 0.02 mg/kg body weight 20 at least by 45%; more preferably by at least 50%, more preferably by at least 55%, more preferably by at least 60%, and more preferably at least 65%; and/or reduces food intake of mice, preferably of female NMRI-mice, over a period of 22 h, if administered as a single dose, preferably subcutaneous dose of 0.01 mg/kg body weight by at least 10%, more preferably 15%, and 25 more preferably 20%.

30 The compounds of the invention have the ability to reduce blood glucose level, and/or to reduce HbA1c levels of a patient. These activities of the compounds of the invention can be assessed in animal models known to the skilled person and also described herein in the Methods. The results of such experiments are described in Examples 13, 14, 16 and 17.

- 14 -

- Preferred compounds of the invention may reduce blood glucose level of mice, preferably in female leptin-receptor deficient diabetic db/db mice over a period of 24 h, if administered as a single dose, preferably subcutaneous dose, of 0.01 mg/kg body weight by at least 4 mmol/L; more preferably by at least 6 mmol/L, more preferably by at least 8 mmol/L. If the dose is increased to 0.1 mg/kg body weight a more pronounced reduction of blood glucose levels can be observed in mice over a period of 24 h, if administered as a single dose, preferably subcutaneous dose. Preferably the compounds of the invention lead to a reduction by at least 7 mmol/L; more preferably by at least 9 mmol/L, more preferably by at least 11 mmol/L. The compounds of the invention preferably reduce the increase of HbA1c levels of mice over a period of 4 weeks, if administered at a daily dose of 0.01 mg/kg to about the ignition value.
- 15 The compounds of the invention also have the ability to reduce body weight of a patient. These activities of the compounds of the invention can be assessed in animal models known to the skilled person and also described herein in the Methods and in Examples 13 and 15.
- 20 Surprisingly, it was found that peptidic compounds of the formula (I), particularly those with a lysine (or close analogues) at position 14 which is further substituted with a lipophilic residue, showed very potent GLP-1 and GIP receptor activation; additionally in combination with amino acids like Gln in position 3 also very potent glucagon receptor activation can be provided.
- 25 It is described in the literature (Murage EN et al., Bioorg. Med. Chem. 16 (2008), 10106-10112), that a GLP-1 analogue with an acetylated Lysine at Pos.14 showed significantly reduced potency compared to natural GLP-1.
- 30 Furthermore, oxidation (in vitro or in vivo) of methionine, present in the core structure of exendin-4, is not possible anymore for peptidic compounds of the formula (I).

- 15 -

Further, compounds of the invention preferably have a high solubility at acidic and/or physiological pH values, e.g., at pH 4.5 and/or at pH 7.4 at 25°C, in another embodiment at least 0.5 mg/ml and in a particular 5 embodiment at least 1.0 mg/ml.

Furthermore, according to one embodiment, compounds of the invention preferably have a high stability when stored in solution. Preferred assay conditions for determining the stability is storage for 7 days at 25°C in 10 solution at pH 4.5 or pH 7.4. The remaining amount of peptide is determined by chromatographic analyses as described in Methods and Examples. Preferably, after 7 days at 25°C in solution at pH 4.5 or pH 7.4, the remaining peptide amount is at least 80%, more preferably at least 85%, even more preferably at least 90% and even more preferably at least 95%.

15

Preferably, the compounds of the present invention comprise a peptide moiety Z (formula II) which is a linear sequence of 39-40 amino carboxylic acids, particularly α -amino carboxylic acids linked by peptide, i.e. carboxamide, bonds.

20

In one embodiment position X14 represents an amino acid residue with a functionalized -NH_2 side chain group, such as functionalized Lys, Orn, Dab, or Dap, more preferably functionalized Lys and X40 is absent or represents Lys.

25

An amino acid residue with an -NH_2 side chain group, e.g. Lys, Orn, Dab or Dap, may be functionalized in that at least one H atom of the -NH_2 side chain group is replaced by -C(O)-R^5 , -C(O)O-R^5 , -C(O)NH-R^5 , -S(O)2-R^5 or R^5 , 30 preferably by -C(O)-R^5 , wherein R^5 is a moiety comprising up to 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P.

- 16 -

In certain embodiments, R⁵ may comprise a lipophilic moiety, e.g. an acyclic linear or branched saturated hydrocarbon group, wherein R⁵ particularly comprises an acyclic linear or branched (C₄-C₃₀) saturated or unsaturated hydrocarbon group, and/or a cyclic saturated, unsaturated or aromatic group, particularly a mono-, bi-, or tricyclic group comprising 4 to 14 carbon atoms and 0, 1, or 2 heteroatoms selected from N, O, and S, e.g. cyclohexyl, phenyl, biphenyl, chromanyl, phenanthrenyl or naphthyl, wherein the acyclic or cyclic group may be unsubstituted or substituted e.g. by halogen, -OH and/or CO₂H.

10

More preferred groups R⁵ may comprise a lipophilic moiety, e.g. an acyclic linear or branched (C₁₂-C₂₂) saturated or unsaturated hydrocarbon group. The lipophilic moiety may be attached to the -NH₂ side chain group by a linker in all stereoisomeric forms, e.g. a linker comprising one or more, e.g.

15

2, 3 or 4, amino acid linker groups such as γ -aminobutyric acid (GABA), ϵ -amino hexanoic acid (ϵ -Ahx), γ -Glu and/or β -Ala. In one embodiment the lipophilic moiety is attached to the -NH₂ side chain group by a linker. In another embodiment the lipophilic moiety is directly attached to the -NH₂ side chain group. Specific examples of amino acid linker groups are (β -Ala)₁₋₄, (γ -Glu)₁₋₄, (ϵ -Ahx)₁₋₄, or (GABA)₁₋₄. Preferred amino acid linker groups are β -Ala, γ -Glu, β -Ala- β -Ala and γ -Glu- γ -Glu.

20

Specific preferred examples for -C(O)-R⁵ groups are listed in the following Table 3, which are selected from the group consisting of (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, 4-Hexadecanoylamino-butyryl-, 4-{3-[(R)-2,5,7,8-tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy carbonyl]-propionylamino}-butyryl-, 4-octadecanoylamino-butyryl-, 4-((Z)-octadec-9-enoylamino)-butyryl-, 6-[(4,4-Diphenyl-cyclohexyloxy)-hydroxy-phosphoryloxy]-hexanoyl-, Hexadecanoyl-, (S)-4-Carboxy-4-(15-carboxy-pentadecanoylamino)-butyryl-, (S)-4-Carboxy-4-{3-[3-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoyl-amino)-propionylamino]-propionylamino}-butyryl-, (S)-4-Carboxy-4-{3-[(R)-

- 17 -

2,5,7,8-tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy-
carbonyl]-propionylamino}-butyryl-, (S)-4-Carboxy-4-((9Z,12Z)-octadeca-
9,12-dienoylamino)-butyryl-, (S)-4-Carboxy-4-[6-((2S,3R,4S,5R)-5-carboxy-
2,3,4,5-tetrahydroxy-pentanoylamino)-hexanoylamino]-butyryl-, (S)-4-
5 Carboxy-4-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-
butyryl-, (S)-4-Carboxy-4-tetradecanoylamino-butyryl-, (S)-4-(11-
Benzylloxycarbonyl-undecanoylamino)-4-carboxy-butyryl-, (S)-4-Carboxy-4-
[11-((2S,3R,4R,5R)-2,3,4,5,6-pentahydroxy-hexylcarbamoyl)-undecanoyl-
amino]-butyryl-, (S)-4-Carboxy-4-((Z)-octadec-9-enoylamino)-butyryl-, (S)-4-
10 Carboxy-4-(4-dodecloxy-benzoylamino)-butyryl-, (S)-4-Carboxy-4-henicos-
anoylamino-butyryl-, (S)-4-Carboxy-4-docosanoylamino-butyryl-, (S)-4-
Carboxy-4-((Z)-nonadec-10-enoylamino)-butyryl-, (S)-4-Carboxy-4-(4-
decyloxy-benzoylamino)-butyryl-, (S)-4-Carboxy-4-[(4'-octyloxy-biphenyl-4-
carbonyl)-amino]-butyryl-, (S)-4-Carboxy-4-(12-phenyl-dodecanoylamino)-
15 butyryl-, (S)-4-Carboxy-4-icosanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-
carboxy-4-hexadecanoylamino-butyrylamino)-butyryl-, (S)-4-Carboxy-4-((S)-
4-carboxy-4-octadecanoylamino-butyrylamino)-butyryl-, 3-(3-Octadecanoyl-
amino-propionylamino)-propionyl-, 3-(3-Hexadecanoyl-amino-propionyl-
amino)-propionyl-, 3-Hexadecanoylamino-propionyl-, (S)-4-Carboxy-4-[(R)-4-
20 ((3R,5S,7R,8R,9R,10S,12S,13R,14R,17R)-3,7,12-trihydroxy-8,10,13-
trimethyl-hexadecahydro-cyclopenta[a]-phenanthren-17-yl)-pentanoylamino]-
butyryl-, (S)-4-Carboxy-4-[(R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-
hydroxy-10,13-dimethyl-hexadecahydro-cyclopenta[a]phenanthren-17-yl)-
pentanoylamino]-butyryl-, (S)-4-Carboxy-4-((9S,10R)-9,10,16-trihydroxy-
25 hexadecanoylamino)-butyryl-, Tetradecanoyl-, 11-Carboxy-undecanoyl-, 11-
Benzylloxycarbonyl-undecanoyl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-tetra-
decanoylamino-butyrylamino)-butyryl-, 6-[Hydroxy-(naphthalene-2-yloxy)-
phosphoryloxy]-hexanoyl-, 6-[Hydroxy-(5-phenyl-pentyloxy)-phosphoryloxy]-
hexanoyl-, 4-(Naphthalene-2-sulfonylamino)-4-oxo-butyryl-, 4-(Biphenyl-4-
30 sulfonylamino)-4-oxo-butyryl-, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[
2-{2-[(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-
ethoxy}-ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetylamino]-butyrylamino}-

butyryl-, (S)-4-Carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl-, (S)-4-Carboxy-2-{(S)-4-carboxy-2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}-
5 (S)-4-Carboxy-2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl-, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}-
10 (S)-4-Carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl-, (S)-4-Carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl]-
15 (S)-4-Carboxy-2-{(S)-4-carboxy-2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}, (S)-4-Carboxy-2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl]-
20 (S)-4-Carboxy-4-[2-(2-{2-[2-(2-{(S)-4-Carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl, 2-(2-{2-[2-(2-{2-[2-(2-{(S)-4-Carboxy-4-(16-1H-tetrazol-5-yl-hexadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}, 2-(2-{2-[2-(2-{2-[2-(2-{(S)-4-Carboxy-4-(16-carboxy-hexadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-
25 (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-Carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-butyryl}, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(10-(4-carboxy-phenoxy)-decanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(7-carboxy-heptanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}-
30 (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(7-carboxy-heptanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(7-carboxy-heptanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylaminolamino]-butyryl}

- 19 -


[(S)-4-carboxy-4-(11-carboxy-undecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetylamino]-butyrylamino}-butyryl-, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[2-(2-{2-[(S)-4-carboxy-4-(13-carboxy-tridecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetylamino]-butyrylaminol-
5 (S)-4-Carboxy-4-[(S)-4-carboxy-4-[2-(2-{2-[2-(2-{2-[(S)-4-carboxy-4-(15-carboxy-pentadecanoyl-amino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetylamino]-butyrylaminol-
10 and (S)-4-Carboxy-4-[(S)-4-carboxy-4-[2-(2-{2-[2-(2-{2-[(S)-4-carboxy-4-(19-carboxy-nonadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetylamino]-butyrylaminol-
15 butyrylaminol-.

Further preferred are stereoisomers, particularly enantiomers of these groups, either S- or R-enantiomers. The term "R" in Table 3 is intended to mean the attachment site of -C(O)-R⁵ at the peptide back bone, i.e. particularly the ϵ -amino group of Lys.





- 20 -

Table 3





structure	IUPAC	name
	(S)-4-Carboxy-4-hexadecanoylamino-butyryl- γE-x53	
	(S)-4-Carboxy-4-octadecanoylamino-butyryl- γE-x70	
	4-Hexadecanoylamino-butyryl- GABA-x53	
	4-[3-[(R)-2,5,7,8-tetramethyl-2-[(4R,8R)-4,8,12-trimethyl-tridecyl]-chroman-6-yloxy carbonyl]-propionyl]amino]-butyryl- GABA-x60	
	4-octadecanoylamino-butyryl- GABA-x70	

	4-((Z)-octadec-9-enoylamino)-butyryl- GABA-x74		Hexadecanoyl- x53	(S)-4-Carboxy-4-(15-carboxy-pentadecanoylamino)-butyryl- x52
--	---	--	----------------------	---




- 22 -

 <p>(S)-4-Carboxy-4-{3-[3-((2S,3R,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoyl)amino]-propionyl}amino]butyryl</p> <p>YE-x59</p>	 <p>(S)-4-Carboxy-4-{3-[(R)-2,5,7,8-tetramethyl-2-[(4R,8R)-4,8,12-trimethyltridecyl]chroman-8-yl]oxy}butyryl</p> <p>YE-x60</p>		
		 <p>(S)-4-Carboxy-4-{(9Z,12Z)-octadeca-9,12-dienoyl}amino]butyryl</p> <p>YE-x61</p>	 <p>(S)-4-Carboxy-4-[6-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoyl)amino]-hexanoyl]butyryl</p> <p>YE-x64</p>

- 23 -

 <p>(S)-4-Carboxy-4-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-butyryl YE-x65</p>	 <p>(S)-4-Carboxy-4-tetradecanoylamino-butyryl YE-x69</p>	 <p>(S)-4-(11-Benzylxycarbonyl-undecanoyl)amino-4-carboxybutyryl YE-x72</p>	 <p>(S)-4-Carboxy-4-[11-((2S,3R,4R,5R)-2,3,4,5,6-pentahydroxy-hexylcarbamoyl)-undecanoylamino]-butyryl YE-x73</p>
--	---	--	---

- 24 -

 <p>(S)-4-Carboxy-4-((Z)-octadec-9-enoylamino)-butyryl- butyryl ester YE-^x74</p>	 <p>(S)-4-Carboxy-4-(4-dodecyloxybenzoylamino)-butyryl- butyryl ester YE-^x75</p>	 <p>(S)-4-Carboxy-4-henicosanoylamino- butyryl- butyryl ester YE-^x76</p>	 <p>(S)-4-Carboxy-4-docosanoylamino- butyryl- butyryl ester YE-^x77</p>
---	---	---	--

- 25 -

	(S)-4-Carboxy-4-((Z)-nonadec-10-enylamino)-butyryl YE-x79
	(S)-4-Carboxy-4-(4-decyloxy-benzoylamino)-butyryl YE-x80
	(S)-4-Carboxy-4-[(4'-octyloxy-biphenyl-4-carbonyl)-amino]-butyryl YE-x81





- 26 -

	(S)-4-Carboxy-4-(12-phenyl-dodecanoylamino)-butyryl-yl yE-x82
	(S)-4-Carboxy-4-icosanoylamino-butyryl-yl yE-x95
	(S)-4-Carboxy-4-((S)-4-carboxy-4-hexadecanoylamino-butyryl)amino-butyryl-yl yE-yE-x53
	(S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)amino-butyryl-yl yE-yE-x70
	3-(3-Octadecanoylamino-propionyl)amino-butyryl-yl β-Ala-β-Ala-x70
	3-(3-Hexadecanoylamino-propionyl)amino-butyryl-yl β-Ala-β-Ala-x53
	3-Hexadecanoylamino-propionyl-β-Ala-x53





- 27 -

<p>(S)-4-Carboxy-4-[<i>(R</i>)-4-((3<i>R</i>,5<i>S</i>,7<i>R</i>,8<i>R</i>,9<i>R</i>,10<i>S</i>,12<i>S</i>,13<i>R</i>,14<i>R</i>,17<i>R</i>)-3,7,12-trihydroxy-8,10,13-trimethylhexadecahydro-cyclopenta[<i>a</i>]phenanthren-17-<i>y</i>)-pentanoylamino]-butyryl-<i>y</i>E-x16</p>	<p>(S)-4-Carboxy-4-[<i>(R</i>)-4-((3<i>R</i>,5<i>R</i>,8<i>R</i>,9<i>S</i>,10<i>S</i>,13<i>R</i>,14<i>S</i>,17<i>R</i>)-3-hydroxy-10,13-dimethylhexadecahydro-cyclopenta[<i>a</i>]phenanthren-17-<i>y</i>)-pentanoylamino]-butyryl-<i>y</i>E-x19</p>	<p>(S)-4-Carboxy-4-[<i>(R</i>)-4-((9<i>S</i>,10<i>R</i>)-9,10,16-trihydroxy-hexadecanoylamino)-butyryl-<i>y</i>E-x25</p>	<p>tetradecanoyl-<i>x</i>69</p>
--	--	--	---------------------------------






- 28 -

	11-Carboxy-undecanoyl- x71		11-Benzylxycarbonyl-undecanoyl x72 (S)-4-Carboxy-4-((S)-4-carboxy-4- tetradecanoylamino-butyrylaminoo)- butyryl- YE-YE-x69		6-[Hydroxy-(naphthalen-2-yloxy)- phosphoryloxy]-hexanoyl- Phospho2
					






- 29 -






 <p>6-[Hydroxy-(5-phenyl-pentyloxy)-phosphoryloxy]-hexanoyl-</p> <p>Phospho3</p>	 <p>4-(Naphthalene-2-sulfonylamino)-4-oxo-butyryl-</p> <p>Sulfonamid 1</p>	 <p>4-(Biphenyl-4-sulfonylamino)-4-oxo-butyryl-</p> <p>Sulfonamid 2</p>	 <p>(S)-4-Carboxy-4-[(S)-4-carboxy-4-[2-(2-[2-[(S)-4-carboxy-4-(17-carboxyheptadecanoylamino)-butyrylamino]-ethoxy)-ethoxy)-acetyl]amino]-butyryl-amino]-butyryl-</p> <p>x100</p>
---	---	---	---



- 30 -

 <p>(S)-4-Carboxy-4-[2-(2-[2-{2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoyl)amino]-butyryl}amino]-ethoxy)-ethoxy]-acetylaminobutyryl</p> <p>x101</p>	 <p>(S)-4-Carboxy-2-{(S)-4-carboxy-2-[2-{2-[2-{2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoyl)amino]-ethoxy}-ethoxy]-acetylaminobutyryl}amino]-ethoxy]-acetylaminobutyryl</p> <p>x102</p>	 <p>(S)-4-Carboxy-2-[2-{2-[2-{2-[2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoyl)amino]-ethoxy}-ethoxy]-acetylaminobutyryl}amino]-ethoxy]-acetylaminobutyryl</p> <p>x103</p>	 <p>(S)-4-Carboxy-4-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoyl)amino]-butyryl}amino]-ethoxy]-acetylaminobutyryl</p> <p>x104</p>	 <p>(S)-4-Carboxy-4-[2-(2-{(S)-4-carboxy-4-(17-carboxy-heptadecanoyl)amino]-butyryl}amino]-ethoxy]-acetylaminobutyryl</p> <p>x105</p>
---	---	---	--	---

- 31 -

 <p>(S)-4-Carboxy-2-[(2-[2-(2-[2-[(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-ethoxy]-ethoxy)-acetyl]amino)-butyryl]amino]-butyryl</p> <p>x106</p>	 <p>(S)-4-Carboxy-2-[2-(2-[2-[(S)-4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyryl]amino)-ethoxy]-ethoxy]-acetyl]amino]-butyryl</p> <p>x107</p>	 <p>2-[2-[2-(2-[2-[(S)-4-Carboxy-4-(17-carboxy-heptadecanoylamino)-butyryl]amino)-ethoxy]-ethoxy]-acetyl]amino]-butyryl</p> <p>x108</p>	 <p>2-[2-[2-[(S)-4-Carboxy-4-(17-carboxy-heptadecanoylamino)-butyryl]amino)-ethoxy]-ethoxy]-acetyl]amino]-butyryl</p> <p>x109</p>	 <p>(S)-4-Carboxy-4-[(S)-4-carboxy-4-(19-carboxy-nonadecanoylamino)-butyryl]amino]-butyryl</p> <p>x110</p>
--	--	---	--	--

 <p>2-[2-{2-[2-{2-[2-[(S)-4-Carboxy-4-(16-1H-tetrazol-5-yl)hexadecanoylamino]butyrylamino]ethoxy}ethoxy]acetyl]amino]ethoxy]acetyl x111</p>	 <p>2-[2-{2-[2-{2-[2-[(S)-4-Carboxy-4-(16-carboxy-hexadecanoylamino)butyrylamino]ethoxy}ethoxy]acetyl]amino]ethoxy]acetyl x112</p>	 <p>(S)-4-Carboxy-4-[(S)-4-carboxy-4-[(17-carboxy-heptadecanoylamino)butyryl]amino]butyryl x113</p>	 <p>(S)-4-Carboxy-4-[(S)-4-carboxy-4-[(2-[2-{2-[2-[(S)-4-carboxy-4-[(4-phenoxy)decanoylamino]butyrylamino]ethoxy}ethoxy]acetyl]amino]ethoxy]acetyl]butyryl x114</p>	 <p>(S)-4-Carboxy-4-[(S)-4-carboxy-4-[2-{2-[2-{2-[2-[(S)-4-carboxy-4-(7-carboxy-heptanoylamino)butyrylamino]ethoxy}ethoxy]acetyl]amino]ethoxy]acetyl]butyryl x115</p>
---	--	---	--	---

 <p>(S)-4-Carboxy-4-((S)-4-carboxy-4-[2-(2-{2-[2-{(S)-4-carboxy-4-(11-carboxy-undecanoyl)amino}-butyryl)amino]-ethoxy)-ethoxy)-acetylaminobutyryl)amino}butyryl</p> <p>x116</p>	 <p>(S)-4-Carboxy-4-((S)-4-carboxy-4-[2-(2-{2-{(S)-4-carboxy-4-(13-carboxy-tridecanoyl)amino}-butyryl)amino]-ethoxy)-ethoxy)-acetylaminobutyryl)amino}butyryl</p> <p>x117</p>	 <p>(S)-4-Carboxy-4-((S)-4-carboxy-4-[2-(2-{2-{(S)-4-carboxy-4-(15-carboxypentadecanoyl)amino}-butyryl)amino]-ethoxy)-ethoxy)-acetylaminobutyryl)amino}butyryl</p> <p>x118</p>	 <p>(S)-4-Carboxy-4-((S)-4-carboxy-4-[2-(2-{2-{(S)-4-carboxy-4-(19-carboxynonadecanoyl)amino}-butyryl)amino]-ethoxy)-ethoxy)-acetylaminobutyryl)amino}butyryl</p> <p>x119</p>
---	---	--	---

- 34 -

In some embodiments, the invention relates to peptidic compounds of Formula (I) as defined above, wherein X14 represents an amino acid residue selected from Lys, Orn, Dab and Dap, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, X40 represents an amino acid residue selected
5 from Lys, Orn, Dab and Dap, wherein the -NH₂ side chain group can be functionalized by -C(O)-R⁵, and R⁵ is a lipophilic moiety selected from an acyclic linear or branched (C₄-C₃₀) saturated or unsaturated hydrocarbon group, and/or a cyclic saturated, unsaturated or aromatic group, wherein the lipophilic moiety may be attached to the -NH₂ side chain group by a linker
10 selected from (β-Ala)₁₋₄, (γ-Glu)₁₋₄, (ε-Ahx)₁₋₄, or (GABA)₁₋₄ in all stereoisomeric forms.

In certain embodiments, X14 represents an amino acid residue with a functionalized -NH₂ side chain group, such as functionalized Lys, Orn, Dab
15 or Dap, wherein at least one H atom of the -NH₂ side chain group is replaced by -C(O)-R⁵, which is selected from the group consisting of the substituents according to Table 3 above.

In some embodiments, X14 represents an amino acid residue selected from
20 Lys, Orn, Dab and Dap, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, X40 represents an amino acid residue selected from Lys, Orn, Dab and Dap, wherein the -NH₂ side chain group can be functionalized by -C(O)-R⁵, and -C(O)-R⁵ is selected from the group consisting of the substituents according to Table 3 above.

25

In some embodiments of the invention, position X14 and/or X40 in formula (II) represents Lysine (Lys). According to some embodiments, Lys at position 14 and optionally at position 40 is functionalized, e.g. with a group -C(O)R⁵ as described above. In other embodiments, X40 is absent and X14 is Lys
30 functionalized with -C(O)-R⁵, -C(O)O-R⁵, -C(O)NH-R⁵, -S(O)2-R⁵ or R⁵, preferably by -C(O)-R⁵, wherein R⁵ is as defined above. In particular, X14 is Lys functionalized with C(O)-R⁵, wherein R⁵ is selected from the group

- 35 -

consisting of (S)-4-carboxy-4-hexadecanoylamino-butyryl (γ E-x53), (S)-4-carboxy-4-octadecanoylamino-butyryl (γ E-x70), 4-hexadecanoylamino-butyryl (GABA-x53), 4-{3-[(R)-2,5,7,8-tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy carbonyl]-propionylamino}-butyryl-
5 (GABA-x60), 4-octadecanoylamino-butyryl (GABA-x70), 4-((Z)-octadec-9-enoylamino)-butyryl (GABA-x74), 6-[(4,4-Diphenyl-cyclohexyloxy)-hydroxy-phosphoryloxy]-hexanoyl (Phospho1), Hexadecanoyl (x53), (S)-4-Carboxy-4-(15-carboxy-pentadecanoylamino)-butyryl (x52), (S)-4-Carboxy-4-{3-[3-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-
10 propionylamino]-propionylamino}-butyryl (γ E-x59), (S)-4-Carboxy-4-{3-[(R)-2,5,7,8-tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy carbonyl]-propionylamino}-butyryl (γ E-x60), (S)-4-Carboxy-4-((9Z,12Z)-octadeca-9,12-dienoylamino)-butyryl (γ E-x61), (S)-4-Carboxy-4-[6-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-
15 hexanoylamino]-butyryl (γ E-x64), (S)-4-Carboxy-4-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-butyryl (γ E-x65), (S)-4-carboxy-4-tetradecanoylamino-butyryl (γ E-x69), (S)-4-(11-Benzyloxycarbonyl-undecanoylamino)-4-carboxy-butyryl (γ E-x72), (S)-4-carboxy-4-[11-((2S,3R,4R,5R)-2,3,4,5,6-pentahydroxy-hexylcarbamoyl)-
20 undecanoylamino]-butyryl (γ E-x73), (S)-4-Carboxy-4-((Z)-octadec-9-enoylamino)-butyryl (γ E-x74), (S)-4-Carboxy-4-(4-dodecyloxy-benzoylamino)-butyryl (γ E-x75), (S)-4-Carboxy-4-henicosanoylamino-butyryl (γ E-x76), (S)-4-Carboxy-4-docosanoylamino-butyryl (γ E-x77), (S)-4-Carboxy-4-((Z)-nonadec-10-enoylamino)-butyryl (γ E-x79), (S)-4-Carboxy-4-
25 (4-decyloxy-benzoylamino)-butyryl (γ E-x80), (S)-4-Carboxy-4-[(4'-octyloxy-biphenyl-4-carbonyl)-amino]-butyryl (γ E-x81), (S)-4-Carboxy-4-(12-phenyl-dodecanoylamino)-butyryl (γ E-x82), (S)-4-Carboxy-4-icosanoylamino-butyryl (γ E-x95), (S)-4-Carboxy-4-((S)-4-carboxy-4-hexadecanoylamino-butyryl)-butyryl (γ E- γ E-x53), (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl (γ E- γ E-x70), and 3-(3-Octadecanoylamino-propionylamino)-propionyl (β -Ala- β -Ala-x70).

- 36 -

In some embodiments, X14 is Lys functionalized with C(O)-R⁵, wherein R⁵ is selected from the group consisting of (S)-4-carboxy-4-hexadecanoylamino-butyryl (γ E-x53), (S)-4-carboxy-4-octadecanoylamino-butyryl (γ E-x70), (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl (γ E-5 γ E-x70), 4-octadecanoylamino-butyryl (GABA-x70), (S)-4-Carboxy-4-henicosanoylamino-butyryl (γ E-x76), and 3-(3-Octadecanoylamino-propionylamino)-propionyl (β -Ala- β -Ala-x70).

A further embodiment relates to a group of compounds, wherein
10 R¹ is NH₂,
R² is NH₂ or
R¹ and R² are NH₂.

A further embodiment relates to a group of compounds, wherein
15 X3 represents an amino acid residue selected from Gln, Glu and His,
X12 represents an amino acid residue selected from Ile and Lys,
X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is as described above,
20 X15 represents an amino acid residue selected from Asp and Glu,
X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,
X17 represents an amino acid residue selected from Arg, Lys, Glu, Ile, Gln, Leu, Aib, Tyr and Ala,
25 X18 represents an amino acid residue selected from Ala, Arg, Aib, Leu, Lys and Tyr,
X19 represents an amino acid residue selected from Ala, Gln, Val and Aib,
X20 represents an amino acid residue selected from Gln, Aib, Phe, Arg,
30 Leu, Lys and His,
X21 represents an amino acid residue selected from Asp, Glu, Tyr, and Leu,

- 37 -

X28 represents an amino acid residue selected from Asn, Ala, Aib, Arg and Lys,

X29 represents an amino acid residue selected from Gly, Thr, Aib, D-Ala and Ala,

5 X40 is either absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, Glu and His,

X12 represents an amino acid residue selected from Ile and Lys,

10 X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is as described above,

X15 represents an amino acid residue selected from Asp and Glu,

15 X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Lys, Glu, Gln, Leu, Aib, Tyr and Ala,

X18 represents an amino acid residue selected from Ala, Arg, Aib, Leu and Tyr,

20 X19 represents an amino acid residue selected from Ala, Val and Aib,

X20 represents an amino acid residue selected from Gln, Aib, Phe, Leu, Lys, His, Pip, (S)MeLys, (R)MeLys and (S)MeOrn,

X21 represents an amino acid residue selected from Asp, Glu and Leu,

25 X28 represents an amino acid residue selected from Asn, Ala, Aib and Ser,

X29 represents an amino acid residue selected from Gly, Thr, Aib, D-Ala and Ala,

X40 is either absent or represents Lys.

30 A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, Glu and His,

X12 represents Ile,

- 38 -

X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is as described above,

X15 represents an amino acid residue selected from Asp and Glu,

5 X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Lys, Glu, Gln, Leu, Aib, Tyr and Ala,

X18 represents an amino acid residue selected from Ala and Arg,

10 X19 represents an amino acid residue selected from Ala and Val,

X20 represents an amino acid residue selected from Gln, Aib, Lys, Pip, (S)MeLys, (R)MeLys and (S)MeOrn and His,

X21 represents an amino acid residue selected from Asp, Glu and Leu,

X28 represents an amino acid residue selected from Asn and Ala,

15 X29 represents an amino acid residue selected from Gly, Thr and D-Ala,

X40 is either absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

20 X3 represents an amino acid residue selected from Gln, Glu and His,

X12 represents an amino acid residue selected from Ile and Lys,

X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is as described above,

25 X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Lys, Glu, Gln, Leu, Aib, Tyr and Ala,

30 X18 represents an amino acid residue selected from Ala and Arg,

X19 represents an amino acid residue selected from Ala and Val,

- 39 -

X20 represents an amino acid residue selected from Gln, Aib, Lys and His,

X21 represents an amino acid residue selected from Asp, Glu and Leu,

X28 represents an amino acid residue selected from Asn and Ala,

5 X29 represents an amino acid residue selected from Gly, Thr and D-Ala,

X40 is either absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

10 X3 represents an amino acid residue selected from Gln and Glu,

X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-

15 butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyrylamo)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl- and 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-,

X15 represents an amino acid residue selected from Glu and Asp,

20 X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

25 X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

30 A further embodiment relates to a group of compounds, wherein

X3 represents Glu,

X12 represents Ile,

- 40 -

X14 represents Lys, wherein the $-\text{NH}_2$ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl- and 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-,
5 X15 represents an amino acid residue selected from Glu and Asp,
X16 represents an amino acid residue selected from Ser and Lys,
X17 represents Arg,
X18 represents Ala,
X19 represents Ala,
X20 represents an amino acid residue selected from Gln and Aib,
X21 represents an amino acid residue selected from Asp and Glu,
10 X28 represents an amino acid residue selected from Asn and Ala,
X29 represents an amino acid residue selected from Gly and Thr,
X40 is absent.

A further embodiment relates to a group of compounds, wherein
20 X3 represents Gln,
X12 represents Ile,
X14 represents Lys, wherein the $-\text{NH}_2$ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl- and 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-,
25 X15 represents an amino acid residue selected from Glu and Asp,
X16 represents an amino acid residue selected from Ser and Lys,
X17 represents Arg,
X18 represents Ala,

- 41 -

X19 represents Ala,
X20 represents an amino acid residue selected from Gln and Aib,
X21 represents an amino acid residue selected from Asp and Glu,
X28 represents an amino acid residue selected from Asn and Ala,
5 X29 represents an amino acid residue selected from Gly and Thr,
X40 is absent.

A further embodiment relates to a group of compounds, wherein

10 X14 represents Lys, wherein the -NH₂ side chain group is
functionalized by one of the groups selected from (S)-4-Carboxy-4-
hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-
butyryl-, 4-octadecanoylamino-butyryl-, Hexadecanoyl-, (S)-4-Carboxy-
4-henicosanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-
octadecanoylamino-butyryl)amino)-butyryl-, 3-(3-Octadecanoylamino-
15 propionylamino)-propionyl-.

A further embodiment relates to a group of compounds, wherein

20 X14 represents Lys, wherein the -NH₂ side chain group is
functionalized by one of the groups selected from (S)-4-Carboxy-4-
octadecanoylamino-butyryl-, 4-octadecanoylamino-butyryl-, (S)-4-
Carboxy-4-henicosanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-
carboxy-4-octadecanoylamino-butyryl)amino)-butyryl-, 3-(3-
Octadecanoylamino-propionylamino)-propionyl-.

25 A further embodiment relates to a group of compounds, wherein

X14 represents Lys, wherein the -NH₂ side chain group is
functionalized by one of the groups selected from (S)-4-Carboxy-4-
hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-
butyryl-.

30

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln and Glu,

- 42 -

X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

5

X15 represents an amino acid residue selected from Glu and Asp,

X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

10 X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

15 X40 is absent.

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, His and Glu,

X12 represents Ile,

20 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

X15 represents Glu,

25 X16 represents an amino acid residue selected from Glu and Lys,

X17 represents Glu,

X18 represents Ala,

X19 represents Val,

X20 represents Arg,

30 X21 represents Leu,

X28 represents an amino acid residue selected from Asn, Aib and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

- 43 -

X40 is absent.

A further embodiment relates to a group of compounds, wherein

X3 represents Glu,

5 X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

10 X15 represents Glu,

X16 represents an amino acid residue selected from Glu and Lys,

X17 represents Glu,

X18 represents Ala,

X19 represents Val,

15 X20 represents Arg,

X21 represents Leu,

X28 represents an amino acid residue selected from Asn, Aib and Ala,

X29 represents Gly,

X40 is absent.

20

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, His and Glu,

X12 represents an amino acid residue selected from Ile and Lys,

25 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

X15 represents an amino acid residue selected from Glu and Asp,

X16 represents Glu,

30 X17 represents an amino acid residue selected from Arg and Gln,

X18 represents an amino acid residue selected from Ala and Arg,

X19 represents Ala,

- 44 -

X20 represents an amino acid residue selected from Pip, (S)MeLys, (R)MeLys and (S)MeOrn,

X21 represents Glu,

X28 represents an amino acid residue selected from Asn, Ser and Ala,

5 X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, His and Glu,

10 X12 represents an amino acid residue selected from Ile and Lys,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, hexadecanoyl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

15 X15 represents an amino acid residue selected from Glu and Asp,

X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Leu, Aib, Tyr, Glu, Ala and Lys,

20 X18 represents an amino acid residue selected from Ala, Aib, Leu and Tyr,

X19 represents an amino acid residue selected from Ala, Val and Aib,

X20 represents Aib,

X21 represents an amino acid residue selected from Glu, Leu and Tyr,

25 X28 represents an amino acid residue selected from Asn, Arg and Ala,

X29 represents an amino acid residue selected from Gly, Ala, D-Ala and Thr,

X40 is either absent or represents Lys.

30 A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln, His and Glu,

X12 represents an amino acid residue selected from Ile and Lys,

- 45 -

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

- 5 X15 represents an amino acid residue selected from Glu and Asp,
X16 represents an amino acid residue selected from Ser, Lys and Glu,
X17 represents an amino acid residue selected from Arg, Lys, Ile, Glu and Gln,
X18 represents an amino acid residue selected from Ala, Arg and Lys,
10 X19 represents an amino acid residue selected from Ala, Val and Gln,
X20 represents an amino acid residue selected from Gln, Phe, Leu, Lys, His and Arg,
X21 represents an amino acid residue selected from Glu, Asp and Leu,
X28 represents an amino acid residue selected from Asn, Arg, Lys and
15 Ala,
X29 represents an amino acid residue selected from Gly, Aib and Thr,
X40 is either absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

- 20 X12 represents Ile.

A further embodiment relates to a group of compounds, wherein

- X19 represents Ala.

- 25 A further embodiment relates to a group of compounds, wherein
X16 represents Glu,
X20 represents an amino acid residue selected from Pip, (S)MeLys,
(R)MeLys and (S)MeOrn.

- 30 A further embodiment relates to a group of compounds, wherein
X28 represents Ala,
X29 represents Gly.

- 46 -

A further embodiment relates to a group of compounds, wherein

X28 represents Asn,

X29 represents Thr.

5

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln and Glu,

X12 represents Ile,

10 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, wherein R⁵ is selected from (S)-4-Carboxy-

4-hexadecanoylamino-butyryl- (γE-x53), (S)-4-Carboxy-4-

octadecanoylamino-butyryl- (γE-x70), (S)-4-Carboxy-4-((S)-4-carboxy-

4-octadecanoylamino-butyrylamino)-butyryl- (γE-γE-x70), 3-(3-

Octadecanoylamino-propionylamino)-propionyl- (βA-βA-x70), 4-

15 octadecanoylamino-butyryl- (GABA-x70), and (S)-4-Carboxy-4-

henicosanoylamino-butyryl- (γE-x76),

X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

20 X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

25 X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

A further embodiment relates to a group of compounds, wherein

X3 represents an amino acid residue selected from Gln and Glu,

30 X12 represents Ile,

- 47 -

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is (S)-4-Carboxy-4-hexadecanoylamino-butyryl- (γE-x53),

X15 represents an amino acid residue selected from Asp and Glu,

5 X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

10 X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

15 A further embodiment relates to a group of compounds, wherein

X3 represents Glu,

X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ is selected from (S)-4-Carboxy-

20 4-octadecanoylamino-butyryl- (γE-x70), (S)-4-Carboxy-4-((S)-4-

carboxy-4-octadecanoylamino-butyryl-amino)-butyryl- (γE-γE-x70), 3-(3-

Octadecanoylamino-propionylamino)-propionyl- (βA-βA-x70), 4-

octadecanoylamino-butyryl- (GABA-x70), and (S)-4-Carboxy-4-

henicosanoylamino-butyryl- (γE-x76),

25 X15 represents Glu,

X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

X19 represents Ala,

30 X20 represents an amino acid residue selected from Gln and Aib,

X21 represents Glu,

X28 represents an amino acid residue selected from Asn and Ala,

- 48 -

X29 represents an amino acid residue selected from Gly and Thr,
X40 is absent.

Specific examples of peptidic compounds of formula (I) are the compounds
5 of SEQ ID NO: 8-39 as well as salts and solvates thereof.

Specific examples of peptidic compounds of formula (I) are the compounds
of SEQ ID NO: 8-10 and 12-38 as well as salts and solvates thereof.

10 Specific examples of peptidic compounds of formula (I) are the compounds
of SEQ ID NO: 8-13 and 39 as well as salts and solvates thereof.

Specific examples of peptidic compounds of formula (I) are the compounds
of SEQ ID NO: 8-10 and 12-13 as well as salts and solvates thereof.

15 Specific examples of peptidic compounds of formula (I) are the compounds
of SEQ ID NO: 14-21 as well as salts and solvates thereof.

20 Specific examples of peptidic compounds of formula (I) are the compounds
of SEQ ID NO: 22-38 as well as salts and solvates thereof.

25 In certain embodiments, i.e. when the compound of formula (I) comprises
genetically encoded amino acid residues, the invention further provides a
nucleic acid (which may be DNA or RNA) encoding said compound, an
expression vector comprising such a nucleic acid, and a host cell containing
such a nucleic acid or expression vector.

30 In a further aspect, the present invention provides a composition comprising
a compound of the invention in admixture with a carrier. In preferred
embodiments, the composition is a pharmaceutically acceptable composition
and the carrier is a pharmaceutically acceptable carrier. The compound of
the invention may be in the form of a salt, e.g. a pharmaceutically acceptable

- 49 -

salt or a solvate, e.g. a hydrate. In still a further aspect, the present invention provides a composition for use in a method of medical treatment, particularly in human medicine.

- 5 In certain embodiments, the nucleic acid or the expression vector may be used as therapeutic agents, e.g. in gene therapy.

The compounds of formula (I) are suitable for therapeutic application without an additionally therapeutically effective agent. In other embodiments, 10 however, the compounds are used together with at least one additional therapeutically active agent, as described in "combination therapy".

The compounds of formula (I) are particularly suitable for the treatment or prevention of diseases or disorders caused by, associated with and/or 15 accompanied by disturbances in carbohydrate and/or lipid metabolism, e.g. for the treatment or prevention of hyperglycemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, obesity and metabolic syndrome. Further, the compounds of the invention are particularly suitable for the treatment or prevention of degenerative diseases, particularly 20 neurodegenerative diseases.

The compounds described find use, *inter alia*, in preventing weight gain or promoting weight loss. By "preventing" is meant inhibiting or reducing when compared to the absence of treatment, and is not necessarily meant to imply 25 complete cessation of a disorder.

The compounds of the invention may cause a decrease in food intake and/or increase in energy expenditure, resulting in the observed effect on body weight.

30 Independently of their effect on body weight, the compounds of the invention may have a beneficial effect on circulating cholesterol levels, being capable

- 50 -

of improving lipid levels, particularly LDL, as well as HDL levels (e.g. increasing HDL/LDL ratio).

Thus, the compounds of the invention can be used for direct or indirect
5 therapy of any condition caused or characterised by excess body weight, such as the treatment and/or prevention of obesity, morbid obesity, obesity linked inflammation, obesity linked gallbladder disease, obesity induced sleep apnea. They may also be used for treatment and prevention of the metabolic syndrome, diabetes, hypertension, atherogenic dyslipidemia,
10 atherosclerosis, arteriosclerosis, coronary heart disease, or stroke. Their effects in these conditions may be as a result of or associated with their effect on body weight, or may be independent thereof.

Preferred medical uses include delaying or preventing disease progression
15 in type 2 diabetes, treating metabolic syndrome, treating obesity or preventing overweight, for decreasing food intake, increase energy expenditure, reducing body weight, delaying the progression from impaired glucose tolerance (IGT) to type 2 diabetes; delaying the progression from type 2 diabetes to insulin-requiring diabetes; regulating appetite; inducing
20 satiety; preventing weight regain after successful weight loss; treating a disease or state related to overweight or obesity; treating bulimia; treating binge eating; treating atherosclerosis, hypertension, type 2 diabetes, IGT, dyslipidemia, coronary heart disease, hepatic steatosis, treatment of beta-blocker poisoning, use for inhibition of the motility of the gastrointestinal tract, useful in connection with investigations of the gastrointestinal tract
25 using techniques such as X-ray, CT- and NMR-scanning.

Further preferred medical uses include treatment or prevention of degenerative disorders, particularly neurodegenerative disorders such as
30 Alzheimer's disease, Parkinson's disease, Huntington's disease, ataxia, e.g spinocerebellar ataxia, Kennedy disease, myotonic dystrophy, Lewy body dementia, multi-systemic atrophy, amyotrophic lateral sclerosis, primary

- 51 -

lateral sclerosis, spinal muscular atrophy, prion-associated diseases, e.g. Creutzfeldt-Jacob disease, multiple sclerosis, telangiectasia, Batten disease, corticobasal degeneration, subacute combined degeneration of spinal cord, Tabes dorsalis, Tay-Sachs disease, toxic encephalopathy, infantile Refsum disease, Refsum disease, neuroacanthocytosis, Niemann-Pick disease, Lyme disease, Machado-Joseph disease, Sandhoff disease, Shy-Drager syndrome, wobbly hedgehog syndrome, proteopathy, cerebral β -amyloid angiopathy, retinal ganglion cell degeneration in glaucoma, synucleinopathies, tauopathies, frontotemporal lobar degeneration (FTLD), dementia, cadasil syndrome, hereditary cerebral hemorrhage with amyloidosis, Alexander disease, seipinopathies, familial amyloidotic neuropathy, senile systemic amyloidosis, serpinopathies, AL (light chain) amyloidosis (primary systemic amyloidosis), AH (heavy chain) amyloidosis, AA (secondary) amyloidosis, aortic medial amyloidosis, ApoAI amyloidosis, ApoAI amyloidosis, ApoAIV amyloidosis, familial amyloidosis of the Finnish type (FAF), Lysozyme amyloidosis, Fibrinogen amyloidosis, Dialysis amyloidosis, Inclusion body myositis/myopathy, Cataracts, Retinitis pigmentosa with rhodopsin mutations, medullary thyroid carcinoma, cardiac atrial amyloidosis, pituitary prolactinoma, Hereditary lattice corneal dystrophy, Cutaneous lichen amyloidosis, Mallory bodies, corneal lactoferrin amyloidosis, pulmonary alveolar proteinosis, odontogenic (Pindborg) tumor amyloid, cystic fibrosis, sickle cell disease or critical illness myopathy (CIM).

Further medical uses include treatment of bone related disorders, such as osteoporosis or osteoarthritis, etc., where increased bone formation and decreased bone resorption might be beneficial.

DETAILED DESCRIPTION OF THE INVENTION

30

Definitions

The amino acid sequences of the present invention contain the conventional

- 52 -

one letter and three letter codes for naturally occurring amino acids, as well as generally accepted three letter codes for other amino acids, such as Aib (α-aminoisobutyric acid), Orn (ornithine), Dab (2,4-diamino butyric acid), Dap (2,3-diamino propionic acid), Nle (norleucine), GABA (γ-aminobutyric acid) 5 or Ahx (ε-aminohexanoic acid).

Furthermore, the following codes were used for the amino acids shown in Table 4:

10

15

Table 4:

- 53 -

structure	name	code
	(S)- α -methyl-lysine	(S)MeLys
	(R)- α -methyl-lysine	(R)MeLys
	(S)- α -methyl-ornithine	(S)MeOrn
	4-amino-piperidine-4-carboxylic acid	Pip

The term „native exendin-4“ refers to native exendin-4 having the sequence HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPS-NH₂ (SEQ ID NO: 1).

The invention provides peptidic compounds as defined above.

The peptidic compounds of the present invention comprise a linear backbone of amino carboxylic acids linked by peptide, i.e. carboxamide bonds. Preferably, the amino carboxylic acids are α -amino carboxylic acids and more preferably L- α -amino carboxylic acids, unless indicated otherwise. The peptidic compounds preferably comprise a backbone sequence of 39-40 amino carboxylic acids.

The peptidic compounds of the present invention may have unmodified side-chains, but carry at least one modification at one of the side chains.

- 5 For the avoidance of doubt, in the definitions provided herein, it is generally intended that the sequence of the peptidic moiety (II) differs from native exendin-4 at least at one of those positions which are stated to allow variation. Amino acids within the peptide moiety (II) can be considered to be numbered consecutively from 0 to 40 in the conventional N-terminal to C-terminal direction. Reference to a „position“ within peptidic moiety (II) should be construed accordingly, as should reference to positions within native exendin-4 and other molecules, e.g., in exendin-4, His is at position 1, Gly at position 2, ..., Met at position 14, ... and Ser at position 39.
- 10
- 15 The amino acid residues at position 14 and optionally at position 40, having a side chain with an - NH₂ group, e.g. Lys, Orn, Dab or Dap are conjugated to a functional group, e.g. acyl groups. Thus, one or more selected amino acids of the peptides in the present invention may carry a covalent attachment at their side chains. In some cases those attachments may be
- 20 lipophilic. These lipophilic side chain attachments have the potential to reduce in vivo clearance of the peptides thus increasing their in vivo half-lives.

- 25 The lipophilic attachment may consist of a lipophilic moiety which can be a branched or unbranched, aliphatic or unsaturated acyclic moiety and/or a cyclic moiety selected from one or several aliphatic or unsaturated homocycles or heterocycles, aromatic condensed or non-condensed homocycles or heterocycles, ether linkages, unsaturated bonds and substituents, e.g. hydroxy and/or carboxy groups. The lipophilic moiety may
- 30 be attached to the peptide either by alkylation, reductive amination or by an amide bond, a carbamate or a sulfonamide bond in case of amino acids carrying an amino group at their side chain.

Nonlimiting examples of lipophilic moieties that can be attached to amino acid side chains include fatty acids, e.g. C₈₋₃₀ fatty acids such as palmitic acid, myristic acid, stearic acid and oleic acid, and/or cyclic groups as 5 described above or derivatives thereof.

There might be one or several linkers between the amino acid of the peptide and the lipophilic attachment. Nonlimiting examples of those linkers are β -alanine, γ -glutamic acid, α -glutamic acid, γ -aminobutyric acid and/or ϵ -10 amino hexanoic acid or dipeptides, such as β -Ala- β -Ala (also abbreviated β A- β A herein) and/or γ -Glu- γ -Glu (also abbreviated γ E- γ E herein) in all their stereo-isomer forms (S and R enantiomers).

Thus, one nonlimiting example of a side chain attachment is palmitic acid 15 which is covalently linked to the α -amino group of glutamic acid forming an amide bond. The γ -carboxy group of this substituted glutamic acid can form an amide bond with the side chain amino group of a lysine within the peptide.

20 In a further aspect, the present invention provides a composition comprising a compound of the invention as described herein, or a salt or solvate thereof, in admixture with a carrier.

The invention also provides the use of a compound of the present invention 25 for use as a medicament, particularly for the treatment of a condition as described below.

The invention also provides a composition wherein the composition is a pharmaceutically acceptable composition, and the carrier is a 30 pharmaceutically acceptable carrier.

The skilled person is aware of a variety of different methods to prepare the peptides that are described in this invention. These methods include but are not limited to synthetic approaches and recombinant gene expression. Thus, 5 one way of preparing these peptides is the synthesis in solution or on a solid support and subsequent isolation and purification. A different way of preparing the peptides is gene expression in a host cell in which a DNA sequence encoding the peptide has been introduced. Alternatively, the gene expression can be achieved without utilizing a cell system. The methods 10 described above may also be combined in any way.

A preferred way to prepare the peptides of the present invention is solid phase synthesis on a suitable resin. Solid phase peptide synthesis is a well established methodology (see for example: Stewart and Young, Solid Phase 15 Peptide Synthesis, Pierce Chemical Co., Rockford, Ill., 1984; E. Atherton and R. C. Sheppard, Solid Phase Peptide Synthesis. A Practical Approach, Oxford-IRL Press, New York, 1989). Solid phase synthesis is initiated by attaching an N-terminally protected amino acid with its carboxy terminus to an inert solid support carrying a cleavable linker. This solid support can be 20 any polymer that allows coupling of the initial amino acid, e.g. a trityl resin, a chlorotriyl resin, a Wang resin or a Rink resin in which the linkage of the carboxy group (or carboxamide for Rink resin) to the resin is sensitive to acid (when Fmoc strategy is used). The polymer support must be stable under the conditions used to deprotect the α -amino group during the peptide 25 synthesis.

After the first amino acid has been coupled to the solid support, the α -amino protecting group of this amino acid is removed. The remaining protected amino acids are then coupled one after the other in the order represented by 30 the peptide sequence using appropriate amide coupling reagents, for example BOP, HBTU, HATU or DIC (N,N'-diisopropylcarbodiimide) / HOEt (1-hydroxybenzotriazol), wherein BOP, HBTU and HATU are used with

- 57 -

tertiary amine bases. Alternatively, the liberated N-terminus can be functionalized with groups other than amino acids, for example carboxylic acids, etc.

- 5 Usually, reactive side-chain groups of the amino acids are protected with suitable blocking groups. These protecting groups are removed after the desired peptides have been assembled. They are removed concomitantly with the cleavage of the desired product from the resin under the same conditions. Protecting groups and the procedures to introduce protecting 10 groups can be found in Protective Groups in Organic Synthesis, 3d ed., Greene, T. W. and Wuts, P. G. M., Wiley & Sons (New York: 1999).

In some cases it might be desirable to have side-chain protecting groups that can selectively be removed while other side-chain protecting groups 15 remain intact. In this case the liberated functionality can be selectively functionalized. For example, a lysine may be protected with an ivDde ([1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-3-methylbutyl) protecting group (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603) which is labile to a very nucleophilic base, for example 4% hydrazine in DMF (dimethyl 20 formamide). Thus, if the N-terminal amino group and all side-chain functionalities are protected with acid labile protecting groups, the ivDde group can be selectively removed using 4% hydrazine in DMF and the corresponding free amino group can then be further modified, e.g. by acylation. The lysine can alternatively be coupled to a protected amino acid 25 and the amino group of this amino acid can then be deprotected resulting in another free amino group which can be acylated or attached to further amino acids.

Finally the peptide is cleaved from the resin. This can be achieved by using 30 King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The raw material can then be purified by chromatography, e.g. preparative RP-HPLC, if necessary.

Potency

As used herein, the term "potency" or "in vitro potency" is a measure for the

5 ability of a compound to activate the receptors for GLP-1, GIP or glucagon in a cell-based assay. Numerically, it is expressed as the "EC₅₀ value", which is the effective concentration of a compound that induces a half maximal increase of response (e.g. formation of intracellular cAMP) in a dose-response experiment.

10

Therapeutic uses

The compounds of the invention are agonists for the receptors for GLP-1 and for GIP as well as optionally the glucagon receptor (e.g. "dual or trigonal

15 agonists"). Such peptides that are GIP/GLP-1 co-agonists, or GIP/GLP-1/glucagon tri-agonists may provide therapeutic benefit to address a clinical need for targeting the metabolic syndrome by allowing simultaneous treatment of diabetes and obesity.

20 Metabolic syndrome is a combination of medical disorders that, when occurring together, increase the risk of developing type 2 diabetes, as well as atherosclerotic vascular disease, e.g. heart disease and stroke. Defining medical parameters for the metabolic syndrome include diabetes mellitus, impaired glucose tolerance, raised fasting glucose, insulin resistance, 25 urinary albumin secretion, central obesity, hypertension, elevated triglycerides, elevated LDL cholesterol and reduced HDL cholesterol.

Obesity is a medical condition in which excess body fat has accumulated to

the extent that it may have an adverse effect on health and life expectancy

30 and due to its increasing prevalence in adults and children it has become one of the leading preventable causes of death in modern world. It increases the likelihood of various other diseases, including heart disease, type 2

- 59 -

diabetes, obstructive sleep apnea, certain types of cancer, as well as osteoarthritis, and it is most commonly caused by a combination of excess food intake, reduced energy expenditure, as well as genetic susceptibility.

5 *Diabetes mellitus*, often simply called diabetes, is a group of metabolic diseases in which a person has high blood sugar levels, either because the body does not produce enough insulin, or because cells do not respond to the insulin that is produced. The most common types of diabetes are: (1) type 1 diabetes, where the body fails to produce insulin; (2) type 2 diabetes,
10 where the body fails to use insulin properly, combined with an increase in insulin deficiency over time, and (3) gestational diabetes, where women develop diabetes due to their pregnancy. All forms of diabetes increase the risk of long-term complications, which typically develop after many years. Most of these long-term complications are based on damage to blood
15 vessels and can be divided into the two categories "macrovascular" disease, arising from atherosclerosis of larger blood vessels and "microvascular" disease, arising from damage of small blood vessels. Examples for macrovascular disease conditions are ischemic heart disease, myocardial infarction, stroke and peripheral vascular disease. Examples for
20 microvascular diseases are diabetic retinopathy, diabetic nephropathy, as well as diabetic neuropathy.

The receptors for GLP-1 and GIP as well as glucagon are members of the family of 7-transmembrane-spanning, heterotrimeric G-protein coupled
25 receptors. They are structurally related to each other and share not only a significant level of sequence identity, but have also similar mechanisms of ligand recognition and intracellular signaling pathways.

Similarly, the peptides GLP-1, GIP and glucagon share regions of high
30 sequence identity/similarity. GLP-1 and glucagon are produced from a common precursor, preproglucagon, which is differentially processed in a tissue-specific manner to yield e.g. GLP-1 in intestinal endocrine cells and

- 60 -

glucagon in alpha cells of pancreatic islets. GIP is derived from a larger proGIP prohormone precursor and is synthesized and released from K-cells located in the small intestine.

The peptidic incretin hormones GLP-1 and GIP are secreted by intestinal 5 endocrine cells in response to food and account for up to 70% of meal-stimulated insulin secretion. Evidence suggests that GLP-1 secretion is reduced in subjects with impaired glucose tolerance or type 2 diabetes, whereas responsiveness to GLP-1 is still preserved in these patients. Thus, targeting of the GLP-1 receptor with suitable agonists offers an attractive 10 approach for treatment of metabolic disorders, including diabetes. The receptor for GLP-1 is distributed widely, being found mainly in pancreatic islets, brain, heart, kidney and the gastrointestinal tract. In the pancreas, GLP-1 acts in a strictly glucose-dependent manner by increasing secretion of insulin from beta cells. This glucose-dependency shows that activation of 15 GLP-1 receptors is unlikely to cause hypoglycemia. Also the receptor for GIP is broadly expressed in peripheral tissues including pancreatic islets, adipose tissue, stomach, small intestine, heart, bone, lung, kidney, testis, adrenal cortex, pituitary, endothelial cells, trachea, spleen, thymus, thyroid and brain. Consistent with its biological function as incretin hormone, the 20 pancreatic β -cell express the highest levels of the receptor for GIP in humans. There is some clinical evidence that the GIP-receptor mediated signaling could be impaired in patients with T2DM but GIP-action is shown to be reversible and could be restored with improvement of the diabetic status. Of note, the stimulation of insulin secretion by both incretin hormones, GIP 25 and GLP-1 is strictly glucose-dependent ensuring a fail-safe mechanism associated with a low risk for hypoglycemia.

At the beta cell level, GLP-1 and GIP have been shown to promote glucose 30 sensitivity, neogenesis, proliferation, transcription of proinsulin and hypertrophy, as well as antiapoptosis. A peptide with dual agonistic activity for the GLP-1 and the GIP receptor could be anticipated to have additive or synergistic anti-diabetic benefit. Other relevant effects of GLP-1 beyond the

- 61 -

pancreas include delayed gastric emptying, increased satiety, decreased food intake, reduction of body weight, as well as neuroprotective and cardioprotective effects. In patients with type 2 diabetes, such extrapancreatic effects could be particularly important considering the high
5 rates of comorbidities like obesity and cardiovascular disease. Further GIP actions in peripheral tissues beyond the pancreas comprise increased bone formation and decreased bone resorption as well as neuroprotective effects which might be beneficial for the treatment of osteoporosis and cognitive defects like Alzheimer's disease.

10

Glucagon is a 29 amino acid peptide hormone that is produced by pancreatic alpha cells and released into the bloodstream when circulating glucose is low. An important physiological role of glucagon is to stimulate glucose output in the liver, which is a process providing the major
15 counterregulatory mechanism for insulin in maintaining glucose homeostasis in vivo.

Glucagon receptors are however also expressed in extra-hepatic tissues such as kidney, heart, adipocytes, lymphoblasts, brain, retina, adrenal gland
20 and gastrointestinal tract, suggesting a broader physiological role beyond glucose homeostasis. Accordingly, recent studies have reported that glucagon has therapeutically positive effects on energy management, including stimulation of energy expenditure and thermogenesis, accompanied by reduction of food intake and body weight loss. Altogether,
25 stimulation of glucagon receptors might be useful in the treatment of obesity and the metabolic syndrome.

Oxyntomodulin is a peptide hormone consisting of glucagon with an eight amino acids encompassing C-terminal extension. Like GLP-1 and glucagon,
30 it is preformed in preproglucagon and cleaved and secreted in a tissue-specific manner by endocrinial cells of the small bowel. Oxyntomodulin is known to stimulate both, the receptors for GLP-1 and glucagon and is

- 62 -

therefore the prototype of a dual agonist.

As GLP-1 and GIP are known for their anti-diabetic effects, GLP-1 and glucagon are both known for their food intake-suppressing effects and

5 glucagon is also a mediator of additional energy expenditure, it is conceivable that a combination of the activities of the two or three hormones in one molecule can yield a powerful medication for treatment of the metabolic syndrome and in particular its components diabetes and obesity.

10 Accordingly, the compounds of the invention may be used for treatment of glucose intolerance, insulin resistance, pre-diabetes, increased fasting glucose, type 2 diabetes, hypertension, dyslipidemia, arteriosclerosis, coronary heart disease, peripheral artery disease, stroke or any combination of these individual disease components.

15 In addition, they may be used for control of appetite, feeding and calory intake, increase of energy expenditure, prevention of weight gain, promotion of weight loss, reduction of excess body weight and altogether treatment of obesity, including morbid obesity.

20 Further disease states and health conditions which could be treated with the compounds of the invention are obesity-linked inflammation, obesity-linked gallbladder disease and obesity-induced sleep apnea.

25 Although all these conditions could be associated directly or indirectly with obesity, the effects of the compounds of the invention may be mediated in whole or in part via an effect on body weight, or independent thereof.

30 Further, diseases to be treated are osteoporosis and neurodegenerative diseases such as Alzheimer's disease or Parkinson's disease, or other degenerative diseases as described above.

- 63 -

- Compared to GLP-1, glucagon and oxyntomodulin, exendin-4 has beneficial physicochemical properties, such as solubility and stability in solution and under physiological conditions (including enzymatic stability towards degradation by enzymes, such as DPP-4 or NEP), which results in a longer
- 5 duration of action *in vivo*. Therefore, exendin-4 might serve as good starting scaffold to obtain exendin-4 analogues with dual or even triple pharmacologies, e.g., GLP-1/GIP and optionally in addition glucagon agonism.
- 10 Nevertheless, also exendin-4 has been shown to be chemically labile due to methionine oxidation in position 14 as well as deamidation and isomerization of asparagine in position 28. Therefore, stability might be further improved by substitution of methionine at position 14 and the avoidance of sequences that are known to be prone to degradation via aspartimide formation,
- 15 especially Asp-Gly or Asn-Gly at positions 28 and 29.

Pharmaceutical compositions

- The term "pharmaceutical composition" indicates a mixture containing
- 20 ingredients that are compatible when mixed and which may be administered. A pharmaceutical composition may include one or more medicinal drugs. Additionally, the pharmaceutical composition may include carriers, buffers, acidifying agents, alkalinizing agents, solvents, adjuvants, tonicity adjusters, emollients, expanders, preservatives, physical and chemical stabilizers e.g.
- 25 surfactants, antioxidants and other components, whether these are considered active or inactive ingredients. Guidance for the skilled in preparing pharmaceutical compositions may be found, for example, in Remington: The Science and Practice of Pharmacy, (20th ed.) ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins and in R.C. Rowe et al
- 30 (Ed), Handbook of Pharmaceutical Excipients, PhP, May 2013 update.

The exendin-4 peptide derivatives of the present invention, or salts thereof,

- 64 -

are administered in conjunction with an acceptable pharmaceutical carrier, diluent, or excipient as part of a pharmaceutical composition. A "pharmaceutically acceptable carrier" is a carrier which is physiologically acceptable (e.g. physiologically acceptable pH) while retaining the
5 therapeutic properties of the substance with which it is administered. Standard acceptable pharmaceutical carriers and their formulations are known to one skilled in the art and described, for example, in Remington: The Science and Practice of Pharmacy, (20th ed.) ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins and in R.C. Rowe et al (Ed), Handbook
10 of Pharmaceutical excipients, PhP, May 2013 update. One exemplary pharmaceutically acceptable carrier is physiological saline solution.

In one embodiment carriers are selected from the group of buffers (e.g. citrate/citric acid), acidifying agents (e.g. hydrochloric acid), alkalizing agents (e.g. sodium hydroxide), preservatives (e.g. phenol), co-solvents (e.g. polyethylene glycol 400), tonicity adjusters (e.g. mannitol), stabilizers (e.g. surfactant, antioxidants, amino acids).

Concentrations used are in a range that is physiologically acceptable.

Acceptable pharmaceutical carriers or diluents include those used in
15 formulations suitable for oral, rectal, nasal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, and transdermal) administration. The compounds of the present invention will typically be administered parenterally.

20 The term "pharmaceutically acceptable salt" means salts of the compounds of the invention which are safe and effective for use in mammals. Pharmaceutically acceptable salts may include, but are not limited to, acid addition salts and basic salts. Examples of acid addition salts include chloride, sulfate, hydrogen sulfate, (hydrogen) phosphate, acetate, citrate, 25 tosylate or mesylate salts. Examples of basic salts include salts with

- 65 -

inorganic cations, e.g. alkaline or alkaline earth metal salts such as sodium, potassium, magnesium or calcium salts and salts with organic cations such as amine salts. Further examples of pharmaceutically acceptable salts are described in Remington: The Science and Practice of Pharmacy, (20th ed.)

5 ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins or in Handbook of Pharmaceutical Salts, Properties, Selection and Use, e.d. P. H. Stahl, C. G. Wermuth, 2002, jointly published by Verlag Helvetica Chimica Acta, Zurich, Switzerland, and Wiley-VCH, Weinheim, Germany.

10 The term "solvate" means complexes of the compounds of the invention or salts thereof with solvent molecules, e.g. organic solvent molecules and/or water.

15 In the pharmaceutical composition, the exendin-4 derivative can be in monomeric or oligomeric form.

The term "therapeutically effective amount" of a compound refers to a nontoxic but sufficient amount of the compound to provide the desired effect.

The amount of a compound of the formula I necessary to achieve the desired

20 biological effect depends on a number of factors, for example the specific compound chosen, the intended use, the mode of administration and the clinical condition of the patient. An appropriate "effective" amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation For example the "therapeutically effective amount" of 25 a compound of the formula (I) is about 0.01 to 50 mg/dose, preferably 0.1 to 10 mg/dose.

Pharmaceutical compositions of the invention are those suitable for parenteral (for example subcutaneous, intramuscular, intradermal or

30 intravenous), oral, rectal, topical and peroral (for example sublingual) administration, although the most suitable mode of administration depends in each individual case on the nature and severity of the condition to be treated

- 66 -

and on the nature of the compound of formula I used in each case.

Suitable pharmaceutical compositions may be in the form of separate units, for example capsules, tablets and powders in vials or ampoules, each of 5 which contains a defined amount of the compound; as powders or granules; as solution or suspension in an aqueous or nonaqueous liquid; or as an oil-in-water or water-in-oil emulsion. It may be provided in single or multiple dose injectable form, for example in the form of a pen. The compositions may, as already mentioned, be prepared by any suitable pharmaceutical 10 method which includes a step in which the active ingredient and the carrier (which may consist of one or more additional ingredients) are brought into contact.

In certain embodiments the pharmaceutical composition may be provided 15 together with a device for application, for example together with a syringe, an injection pen or an autoinjector. Such devices may be provided separate from a pharmaceutical composition or prefilled with the pharmaceutical composition.

20

Combination therapy

The compounds of the present invention, dual agonists for the GLP-1 and GIP receptors or trigonal agonists for the GLP-1, GIP and glucagon 25 receptors, can be widely combined with other pharmacologically active compounds, such as all drugs mentioned in the Rote Liste 2012 and/or the Rote Liste 2013, e.g. with all antidiabetics mentioned in the Rote Liste 2012, chapter 12, and/or the Rote Liste 2013, chapter 12, all weight-reducing agents or appetite suppressants mentioned in the Rote Liste 2012, chapter 30 1, and/or the Rote Liste 2013, chapter 1, all lipid-lowering agents mentioned in the Rote Liste 2012, chapter 58, and/or the Rote Liste 2013, chapter 58, all antihypertensives and nephroprotectives, mentioned in the Rote Liste

- 67 -

2012 and/or the Rote Liste 2013, or all diuretics mentioned in the Rote Liste 2012, chapter 36, and/or the Rote Liste 2013, chapter 36.

5 The active ingredient combinations can be used especially for a synergistic improvement in action. They can be applied either by separate administration of the active ingredients to the patient or in the form of combination products in which a plurality of active ingredients are present in one pharmaceutical preparation. When the active ingredients are administered by separate administration of the active ingredients, this can be
10 done simultaneously or successively.

Most of the active ingredients mentioned hereinafter are disclosed in the USP Dictionary of USAN and International Drug Names, US Pharmacopeia, Rockville 2011.

15 Other active substances which are suitable for such combinations include in particular those which for example potentiate the therapeutic effect of one or more active substances with respect to one of the indications mentioned and/or which allow the dosage of one or more active substances to be
20 reduced.

Therapeutic agents which are suitable for combinations include, for example, antidiabetic agents such as:

25 Insulin and Insulin derivatives, for example: Glargine / Lantus[®] , 270 - 330U/mL of insulin glargine (EP 2387989 A), 300U/mL of insulin glargine (EP 2387989 A), Glulisin / Apidra[®], Detemir / Levemir[®], Lispro / Humalog[®] / Liprolog[®], Degludec / DegludecPlus, Aspart, basal insulin and analogues (e.g. LY-2605541, LY2963016, NN1436), PEGylated insulin Lispro, Humulin[®],
30 Linjeta, SuliXen[®], NN1045, Insulin plus Symlin, PE0139, fast-acting and short-acting insulins (e.g. Linjeta, PH20, NN1218, HinsBet), (APC-002)hydrogel, oral, inhalable, transdermal and sublingual insulins (e.g.

- 68 -

Exubera[®], Nasulin[®], AfreZZa, Tregopil, TPM 02, Capsulin, Oral-lyn[®], Cobalamin[®] oral insulin, ORMD-0801, NN1953, NN1954, NN1956, VIAtab, Oshadi oral insulin). Additionally included are also those insulin derivatives which are bonded to albumin or another protein by a bifunctional linker.

5

GLP-1, GLP-1 analogues and GLP-1 receptor agonists, for example: Lixisenatide / AVE0010 / ZP10 / Lyxumia, Exenatide / Exendin-4 / Byetta / Bydureon / ITCA 650 / AC-2993, Liraglutide / Victoza, Semaglutide, Taspoglutide, Syncria / Albiglutide, Dulaglutide, rExendin-4, CJC-1134-PC, 10 PB-1023, TTP-054, Langlenatide / HM-11260C, CM-3, GLP-1 Eligen, ORMD-0901, NN-9924, NN-9926, NN-9927, Nodexen, Viador-GLP-1, CVX-096, ZYOG-1, ZYD-1, GSK-2374697, DA-3091, MAR-701, MAR709, ZP-2929, ZP-3022, TT-401, BHM-034. MOD-6030, CAM-2036, DA-15864, ARI-2651, ARI-2255, Exenatide-XTEN and Glucagon-Xten.

15

DPP-4 inhibitors, for example: Alogliptin / Nesina, Trajenta / Linagliptin / BI-1356 / Ondero / Trajenta / Tadjenta / Trayenta / Tradzenta, Saxagliptin / Onglyza, Sitagliptin / Januvia / Xelevia / Tesave / Janumet / Velmetia, Galvus / Vildagliptin, Anagliptin, Gemigliptin, Teneligliptin, Melogliptin, 20 Trelagliptin, DA-1229, Omaragliptin / MK-3102, KM-223, Evogliptin, ARI-2243, PBL-1427, Pinoxacin.

25

SGLT2 inhibitors, for example: Invokana / Canagliflozin, Forxiga / Dapagliflozin, Remoglioflozin, Sergliflozin, Empagliflozin, Ipragliflozin, Tofogliflozin, Luseogliflozin, LX-4211, Ertugliflozin / PF-04971729, RO-4998452, EGT-0001442, KGA-3235 / DSP-3235, LIK066, SBM-TFC-039,

30

Biguanides (e.g. Metformin, Buformin, Phenformin), Thiazolidinediones (e.g. Pioglitazone, Rivoglitazone, Rosiglitazone, Troglitazone), dual PPAR agonists (e.g. Aleglitazar, Muraglitazar, Tesagliptazar), Sulfonylureas (e.g. Tolbutamide, Glibenclamide, Glimepiride/Amaryl, Glipizide), Meglitinides (e.g. Nateglinide, Repaglinide, Mitiglinide), Alpha-glucosidase inhibitors (e.g.

- 69 -

Acarbose, Miglitol, Voglibose), Amylin and Amylin analogues (e.g. Pramlintide, Symlin).

5 GPR119 agonists (e.g. GSK-263A, PSN-821, MBX-2982, APD-597, ZYG-19, DS-8500), GPR40 agonists (e.g. Fasiglifam / TAK-875, TUG-424, P-1736, JTT-851, GW9508).

10 Other suitable combination partners are: Cycloset, inhibitors of 11-beta-HSD (e.g. LY2523199, BMS770767, RG-4929, BMS816336, AZD-8329, HSD-016, BI-135585), activators of glucokinase (e.g. TTP-399, AMG-151, TAK-329, GKM-001), inhibitors of DGAT (e.g. LCQ-908), inhibitors of protein tyrosinephosphatase 1 (e.g. Trodusquemine), inhibitors of glucose-6-phosphatase, inhibitors of fructose-1,6-bisphosphatase, inhibitors of glycogen phosphorylase, inhibitors of phosphoenol pyruvate carboxykinase, 15 inhibitors of glycogen synthase kinase, inhibitors of pyruvate dehydrokinase, alpha2-antagonists, CCR-2 antagonists, SGLT-1 inhibitors (e.g. LX-2761).

20 One or more lipid lowering agents are also suitable as combination partners, such as for example: HMG-CoA-reductase inhibitors (e.g. Simvastatin, Atorvastatin), fibrates (e.g. Bezafibrate, Fenofibrate), nicotinic acid and the derivatives thereof (e.g. Niacin), PPAR-(alpha, gamma or alpha/gamma) agonists or modulators (e.g. Aleglitazar), PPAR-delta agonists, ACAT inhibitors (e.g. Avasimibe), cholesterol absorption inhibitors (e.g. Ezetimibe), Bile acid-binding substances (e.g. Cholestyramine), ileal bile acid transport 25 inhibitors, MTP inhibitors, or modulators of PCSK9.

30 HDL-raising compounds such as: CETP inhibitors (e.g. Torcetrapib, Anacetrapid, Dalcetrapid, Evacetrapid, JTT-302, DRL-17822, TA-8995) or ABC1 regulators.

Other suitable combination partners are one or more active substances for the treatment of obesity, such as for example: Sibutramine, Tesofensine,

- 70 -

Orlistat, antagonists of the cannabinoid-1 receptor, MCH-1 receptor antagonists, MC4 receptor agonists, NPY5 or NPY2 antagonists (e.g. Velneperit), beta-3-agonists, leptin or leptin mimetics, agonists of the 5HT2c receptor (e.g. Lorcaserin), or the combinations of bupropione/naltrexone, 5 bupropione/zonisamide, bupropione/phentermine or pramlintide/metreleptin.

Other suitable combination partners are:

Further gastrointestinal peptides such as Peptide YY 3-36 (PYY3-36) or 10 analogues thereof, pancreatic polypeptide (PP) or analogues thereof.

Glucagon receptor agonists or antagonists, GIP receptor agonists or antagonists, ghrelin antagonists or inverse agonists, Xenin and analogues thereof.

Moreover, combinations with drugs for influencing high blood pressure, chronic heart failure or atherosclerosis, such as e.g.: Angiotensin II receptor antagonists (e.g. telmisartan, candesartan, valsartan, losartan, eprosartan, irbesartan, olmesartan, tasosartan, azilsartan), ACE inhibitors, ECE 20 inhibitors, diuretics, beta-blockers, calcium antagonists, centrally acting hypertensives, antagonists of the alpha-2-adrenergic receptor, inhibitors of neutral endopeptidase, thrombocyte aggregation inhibitors and others or combinations thereof are suitable.

25 In another aspect, this invention relates to the use of a compound according to the invention or a physiologically acceptable salt thereof combined with at least one of the active substances described above as a combination partner, for preparing a medicament which is suitable for the treatment or prevention of diseases or conditions which can be affected by binding to the 30 receptors for GLP-1 and glucagon and by modulating their activity. This is preferably a disease in the context of the metabolic syndrome, particularly one of the diseases or conditions listed above, most particularly diabetes or

- 71 -

obesity or complications thereof.

The use of the compounds according to the invention, or a physiologically acceptable salt thereof, in combination with one or more active substances

5 may take place simultaneously, separately or sequentially.

The use of the compound according to the invention, or a physiologically acceptable salt thereof, in combination with another active substance may take place simultaneously or at staggered times, but particularly within a

10 short space of time. If they are administered simultaneously, the two active substances are given to the patient together; if they are used at staggered times, the two active substances are given to the patient within a period of less than or equal to 12 hours, but particularly less than or equal to 6 hours.

15 Consequently, in another aspect, this invention relates to a medicament which comprises a compound according to the invention or a physiologically acceptable salt of such a compound and at least one of the active substances described above as combination partners, optionally together with one or more inert carriers and/or diluents.

20

The compound according to the invention, or physiologically acceptable salt or solvate thereof, and the additional active substance to be combined therewith may both be present together in one formulation, for example a tablet or capsule, or separately in two identical or different formulations, for 25 example as so-called kit-of-parts.

LEGENDS TO THE FIGURES

30 **Figure 1.** Effect of s.c. administration of compound SEQ ID NO: 13 at 10 $\mu\text{g}/\text{kg}$ on gastric emptying and intestinal passage in female NMRI-mice. Data are mean+SEM.

a) Gastric emptying

- 72 -

b) Small intestinal passage relative to small intestinal length

Figure 2. Effect of s.c. administration of compound SEQ ID NO: 9 at 1, 3 and 10 μ g/kg on gastric emptying and intestinal passage in female NMRI-mice.

5 Data are mean \pm SEM.

a) Gastric emptying

b) Small intestinal passage relative to small intestinal length

Figure 3a. Effect of s.c. administration of compound SEQ ID NO: 12, SEQ ID NO: 13 and liraglutide at 100 μ g/kg on 22-hours feed intake in female NMRI-mice. Data are mean \pm SEM.

Figure 3b. Effect of s.c. administration of compound SEQ ID NO: 9 at 3 and 10 μ g/kg on 22-hours feed intake in female NMRI-mice. Data are mean \pm SEM.

15 **Figure 4.** Effect of s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 μ g/kg on blood glucose after 6 days of treatment in female diet-induced obese C57BL/6NCrl mice (18 weeks on high-fat diet). Data are mean \pm SEM.

20 **Figure 5.** Effect of s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 μ g/kg on body weight in female diet-induced obese (DIO) C57BL/6NCrl mice (18 weeks on high-fat diet). Data are mean \pm SEM.

25 **Figure 6.** Effect of s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 μ g/kg on body weight in female diet-induced obese (DIO) C57BL/6NCrl mice calculated as relative change from baseline. Data are mean \pm SEM..

30 **Figure 7.** Effect of s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 μ g/kg on body fat content in female diet-induced obese (DIO) C57BL/6NCrl mice. Data are mean \pm SEM.

Figure 8. Effect of acute s.c. administration of compounds SEQ ID NO: 13, SEQ ID NO: 12, SEQ ID NO: 10 and SEQ ID NO: 9 at 100 µg/kg on 24h profile of blood glucose of diabetic db/db mice. Data are mean±SEM.

5

Figure 9. Effect of once-daily s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 µg/kg on blood glucose of diabetic db/db mice after 4-weeks chronic treatment. Data are mean±SEM.

10

Figure 10. Effect of once-daily s.c. administration of compound SEQ ID NO: 9 at 10, 30 and 100 µg/kg on HbA1c of diabetic db/db mice at start and at the end 4-weeks chronic treatment. Data are mean±SEM.

15

Figure 11. Effect of s.c. administration of compound SEQ ID NO: 9 and SEQ ID NO: 21 at 10 µg/kg on body weight in female diet-induced obese (DIO) C57BL/6NCrl mice following 3-weeks chronic treatment once daily. Data are mean ± SEM.

20

Figure 12. Effect of s.c. administration of compound SEQ ID NO: 9 and SEQ ID NO: 21 10 µg/kg on body weight in female diet-induced obese (DIO) C57BL/6NCrl mice following 3-weeks chronic treatment once daily. Changes in body weight were calculated as relative change from baseline. Data are mean ± SEM.

25

Figure 13. Effect of 3 weeks of treatment with SEQ ID NO: 16 at 3 and 10 µg/kg, s.c. and SEQ ID NO: 21 at 10 µg/kg, s.c. on non-fasted glucose in diabetic dbdb-mice, represented as change from baseline (0 mmol/l, day -7). Data are mean+SEM.

30

Figure 14. Effect of 3 weeks of treatment with SEQ ID NO: 16 at 3 and 10 µg/kg, s.c. and SEQ ID NO: 21 at 10 µg/kg, s.c. on HbA1c in diabetic dbdb-mice, represented as change from baseline (0 %, day -7). Data are

- 74 -

mean+SEM.

Figure 15. Effect of 3 weeks of treatment with SEQ ID NO: 16 at 3 and 10 µg/kg, s.c. and SEQ ID NO: 21 at 10 µg/kg, s.c. on oral glucose tolerance in 5 diabetic dbdb-mice, represented as change from baseline ($t = 0$ min, 0 mmol/l, immediately before glucose administration). Data are mean+SEM.

Figure 16. Effect of 3 weeks of treatment with SEQ ID NO: 16 at 3 and 10 µg/kg, s.c. and SEQ ID NO: 21 at 10 µg/kg, s.c. on oral glucose tolerance in 10 diabetic dbdb-mice, represented as area under the glucose curve (Glucose-AUC). Data are mean+SEM.

Figure 17. Effect of treatment with SEQ ID NO: 21 at 3 µg/kg, s.c. on 15 glucose lowering in non-fasted female diabetic dbdb-mice, represented as change from baseline. Data are mean+SEM.

Figure 18. Effect of treatment with SEQ ID NO: 14 at 3 µg/kg, s.c. on glucose lowering in non-fasted female diabetic dbdb-mice, represented as change from baseline. Data are mean+SEM.

20 :::::::::::

METHODS

Abbreviations employed are as follows:

5	AA	amino acid
	cAMP	cyclic adenosine monophosphate
	Boc	tert-butyloxycarbonyl
	BOP	(benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate
10	BSA	bovine serum albumin
	tBu	tertiary butyl
	Dde	1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-ethyl
	ivDde	1-(4,4-dimethyl-2,6-dioxocyclohexylidene)3-methyl-butyl
	DIC	N,N'-diisopropylcarbodiimide
15	DIPEA	N,N-diisopropylethylamine
	DMEM	Dulbecco's modified Eagle's medium
	DMF	dimethyl formamide
	EDT	ethanedithiol
	FA	formic acid
20	FBS	fetal bovine serum
	Fmoc	fluorenylmethyloxycarbonyl
	HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
	HBSS	Hanks' Balanced Salt Solution
25	HBTU	2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyl-uronium hexafluorophosphate
	HEPES	2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid
	HOEt	1-hydroxybenzotriazole
	HOSu	N-hydroxysuccinimide
30	HPLC	High Performance Liquid Chromatography
	HTRF	Homogenous Time Resolved Fluorescence
	IBMX	3-isobutyl-1-methylxanthine

- 76 -

	LC/MS	Liquid Chromatography/Mass Spectrometry
	Palm	palmitoyl
	PBS	phosphate buffered saline
	PEG	polyethylene glycole
5	PK	pharmacokinetic
	RP-HPLC	reversed-phase high performance liquid chromatography
	Stea	stearyl
	TFA	trifluoroacetic acid
	Trt	trityl
10	UV	ultraviolet

General synthesis of peptidic compounds

15 Materials:

Different Rink-Amide resins (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucylaminomethyl resin, Merck Biosciences; 4-[(2,4-Dimethoxyphenyl)(Fmoc-amino)methyl]phenoxy acetamido methyl resin, 20 Agilent Technologies) were used for the synthesis of peptide amides with loadings in the range of 0.3-0.4 mmol/g.

Fmoc protected natural amino acids were purchased from Protein Technologies Inc., Senn Chemicals, Merck Biosciences, Novabiochem, Iris Biotech or Bachem. The following standard amino acids were used throughout the syntheses: Fmoc-L-Ala-OH, Fmoc-Arg(Pbf)-OH, Fmoc-L-Asn(Trt)-OH, Fmoc-L-Asp(OtBu)-OH, Fmoc-L-Cys(Trt)-OH, Fmoc-L-Gln(Trt)-OH, Fmoc-L-Glu(OtBu)-OH, Fmoc-Gly-OH, Fmoc-L-His(Trt)-OH, Fmoc-L-Ile-OH, Fmoc-L-Leu-OH, Fmoc-L-Lys(Boc)-OH, Fmoc-L-Met-OH, Fmoc-L-Phe-OH, Fmoc-L-Pro-OH, Fmoc-L-Ser(tBu)-OH, Fmoc-L-Thr(tBu)-OH, Fmoc-L-Trp(Boc)-OH, Fmoc-L-Tyr(tBu)-OH, Fmoc-L-Val-OH. 30

- 77 -

In addition, the following special amino acids were purchased from the same suppliers as above: Fmoc-L-Lys(ivDde)-OH, Fmoc-L-Lys(Mmt)-OH, Fmoc-Aib-OH, Fmoc-D-Ser(tBu)-OH, Fmoc-D-Ala-OH, Boc-L-Tyr(tBu)-OH, Boc-L-His(Boc)-OH (available as toluene solvate) and Boc-L-His(Trt)-OH.

5

The solid phase peptide syntheses were performed for example on a Prelude Peptide Synthesizer (Protein Technologies Inc) or similar automated synthesizer using standard Fmoc chemistry and HBTU/DIPEA activation. DMF was used as the solvent. Deprotection: 20% piperidine/DMF for 2 x 2.5 10 min. Washes: 7 x DMF. Coupling: 2:5:10 200 mM AA / 500 mM HBTU / 2 M DIPEA in DMF 2 x for 20 min. Washes: 5 x DMF.

15 In cases where a Lys-side-chain was modified, Fmoc-L-Lys(ivDde)-OH or Fmoc-L-Lys(Mmt)-OH was used in the corresponding position. After completion of the synthesis, the ivDde group was removed according to a modified literature procedure (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603), using 4% hydrazine hydrate in DMF. The Mmt group was removed by repeated treatment with 1% TFA in dichloromethane. The following acylations were carried out by treating the resin with the N-hydroxy 20 succinimide esters of the desired acid or using coupling reagents like HBTU/DIPEA or HOEt/DIC.

25 All the peptides that had been synthesized were cleaved from the resin with King's cleavage cocktail consisting of 82.5% TFA, 5% phenol, 5% water, 5% thioanisole, 2.5% EDT. The crude peptides were then precipitated in diethyl or diisopropyl ether, centrifuged, and lyophilized. Peptides were analyzed by analytical HPLC and checked by ESI mass spectrometry. Crude peptides were purified by a conventional preparative HPLC purification procedure.

30 **Analytical HPLC / UPLC**

Method A: Analytical UPLC/MS was performed on a Waters UPLC system

- 78 -

with a Waters UPLC HSS 1.7 μ m C18 column (2.1 x 100 mm) at 40 °C with a gradient elution at a flow rate of 0.5 mL/min and monitored at 215 and 280 nm. The gradients were set up as 10% B to 90% B over 15 min and then 90% B for 1 min or as 15% B to 50% B over 12.5 min and then 50% B to 5 90% B over 3 min. Buffer A = 0.1 % formic acid in water and B = 0.1 % formic acid in acetonitrile. A Waters LCT Premier Time-of-Flight instrument was used as mass analyser equipped with an electrospray in the positive ion mode.

10 **Method B:** detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

column: Waters ACQUITY UPLC® CSH™ C18 1.7 μ m (150 x 2.1mm) at 50 °C

solvent: H₂O+0.5%TFA : ACN+0.35%TFA (flow 0.5 ml/min)

15 gradient: 80:20 (0 min) to 80:20 (3 min) to 25:75 (23 min) to 2:98 (23.5 min) to 2:98 (30.5 min) to 80:20 (31 min) to 80:20 (37 min)

Method C: detection at 215 nm

column: Aeris Peptide, 3.6 μ m, XB-C18 (250 x 4.6 mm) at 60 °C

20 solvent: H₂O+0.1%TFA : ACN+0.1%TFA (flow 1.5 ml/min)

gradient: 90:10 (0 min) to 90:10 (3 min) to 10:90 (43 min) to 10:90 (48 min) to 90:10 (49 min) to 90:10 (50 min)

Method D: detection at 214 nm

25 column: Waters X-Bridge C18 3.5 μ m 2.1 x 150 mm

solvent: H₂O+0.5%TFA : ACN (flow 0.55 ml/min)

gradient: 90:10 (0 min) to 40:60 (5 min) to 1:99 (15 min)

Method E: detection at 210 - 225 nm, optionally coupled to a mass analyser

30 Waters LCT Premier, electrospray positive ion mode

column: Waters ACQUITY UPLC® BEH™ C18 1.7 μ m (150 x 2.1 mm) at 50 °C

- 79 -

solvent: H₂O+1%FA : ACN+1%FA (flow 0.9 ml/min)

gradient: 95:5 (0 min) to 95:5 (2min) to 35:65 (3 min) to 65:35 (23.5 min)
to 5:95 (24 min) to 95:5 (26min) to 95:5 (30min)

5 **General Preparative HPLC Purification Procedure:**

The crude peptides were purified either on an Äkta Purifier System or on a Jasco semiprep HPLC System. Preparative RP-C18-HPLC columns of different sizes and with different flow rates were used depending on the amount of crude peptide to be purified. Acetonitrile + 0.05 to 0.1% TFA (B) and water + 0.05 to 0.1% TFA (A) were employed as eluents. Alternatively, a buffer system consisting of acetonitrile and water with minor amounts of acetic acid was used. Product-containing fractions were collected and lyophilized to obtain the purified product, typically as TFA or acetate salt.

15 **Solubility and Stability-Testing of exendin-4 derivatives**

Prior to the testing of solubility and stability of a peptide batch, its content was determined. Therefore, two parameters were investigated, its purity (HPLC-UV) and the amount of salt load of the batch (ion chromatography).

20 For solubility testing, the target concentration was 1.0 mg/mL pure compound. Therefore, solutions from solid samples were prepared in different buffer systems with a concentration of 1.0 mg/mL compound based on the previously determined content. HPLC-UV was performed after 2 h of gentle agitation from the supernatant, which was obtained by 20 min of 25 centrifugation at 4000 rpm.

The solubility was then determined by comparison with the UV peak areas obtained with a stock solution of the peptide at a concentration of 2 mg/mL in pure water or a variable amount of acetonitrile (optical control that all of the 30 compound was dissolved). This analysis also served as starting point (t0) for the stability testing.

- 80 -

For stability testing, an aliquot of the supernatant obtained for solubility was stored for 7 days at 25°C. After that time course, the sample was centrifuged for 20 min at 4000 rpm and the supernatant was analysed with HPLC-UV.

For determination of the amount of the remaining peptide, the peak areas of

5 the target compound at t0 and t7 were compared, resulting in "% remaining peptide", following the equation

$$\% \text{ remaining peptide} = [(\text{peak area peptide t7}) \times 100] / \text{peak area peptide t0}.$$

The amount of soluble degradation products was calculated from the

10 comparison of the sum of the peak areas from all observed impurities reduced by the sum of peak areas observed at t0 (i.e. to determine the amount of newly formed peptide-related species). This value was given in percentual relation to the initial amount of peptide at t0, following the equation:

15

$$\% \text{ soluble degradation products} = \{[(\text{peak area sum of impurities t7}) - (\text{peak area sum of impurities t0})] \times 100\} / \text{peak area peptide t0}$$

The potential difference from the sum of "% remaining peptide" and "%

20 soluble degradation products" to 100% reflects the amount of peptide which did not remain soluble upon stress conditions following the equation

$$\% \text{ precipitate} = 100 - [\% \text{ remaining peptide}] + [\% \text{ soluble degradation products}]$$

25

This precipitate includes non-soluble degradation products, polymers and/or fibrils, which have been removed from analysis by centrifugation.

The chemical stability is expressed as "% remaining peptide".

30

Anion Chromatography

Instrument: Dionex ICS-2000, pre/column: Ion Pac AG-18 2 x 50 mm

- 81 -

(Dionex)/AS18 2 x 250 mm (Dionex), eluent: aqueous sodium hydroxide, flow: 0.38 mL/min, gradient: 0-6 min: 22 mM KOH, 6-12 min: 22-28 mM KOH, 12-15 min: 28-50 mM KOH, 15-20min: 22mM KOH, suppressor: ASRS 300 2 mm, detection: conductivity.

5

As HPLC/UPLC method method D or E has been used.

In vitro cellular assays for GIP receptor, GLP-1 receptor and glucagon receptor efficacy

10 Agonism of compounds for the receptors was determined by functional assays measuring cAMP response of HEK-293 cell lines stably expressing human GIP, GLP-1 or glucagon receptor.

cAMP content of cells was determined using a kit from Cisbio Corp. (cat. no. 15 62AM4PEC) based on HTRF (Homogenous Time Resolved Fluorescence). For preparation, cells were split into T175 culture flasks and grown overnight to near confluence in medium (DMEM / 10% FBS). Medium was then removed and cells washed with PBS lacking calcium and magnesium, followed by proteinase treatment with accutase (Sigma-Aldrich cat. no. 20 A6964). Detached cells were washed and resuspended in assay buffer (1 x HBSS; 20 mM HEPES, 0.1% BSA, 2 mM IBMX) and cellular density determined. They were then diluted to 400000 cells/ml and 25 μ l-aliquots dispensed into the wells of 96-well plates. For measurement, 25 μ l of test compound in assay buffer was added to the wells, followed by incubation for 25 30 minutes at room temperature. After addition of HTRF reagents diluted in lysis buffer (kit components), the plates were incubated for 1 hr, followed by measurement of the fluorescence ratio at 665 / 620 nm. In vitro potency of agonists was quantified by determining the concentrations that caused 50% activation of maximal response (EC50).

30

Bioanalytical screening method for quantification of exendin-4 derivatives in mice and pigs

- 82 -

Mice were dosed 1 mg/kg subcutaneously (s.c.). The mice were sacrificed and blood samples were collected after 0.25, 0.5, 1, 2, 4, 8, 16 and 24 hours post application. Plasma samples were analyzed after protein precipitation via liquid chromatography mass spectrometry (LC/MS). PK parameters and 5 half-life were calculated using WinonLin Version 5.2.1 (non-compartment model).

Female Göttinger minipigs were dosed 0.1 mg/kg subcutaneously (s.c.). Blood samples were collected after 0.25, 0.5, 1, 2, 4, 8, 24, 32, 48, 56 and 10 72 hours post application. Plasma samples were analyzed after protein precipitation via liquid chromatography mass spectrometry (LC/MS). PK parameters and half-life were calculated using WinonLin Version 5.2.1 (non-compartment model).

Gastric emptying and intestinal passage in mice

15 Female NMRI-mice of a body weight between 20 and 30 g were used. Mice were adapted to housing conditions for at least one week.

Mice were overnight fasted, while water remained available all the time. On 20 the study day, mice were weighed, single-caged and allowed access to 500 mg of feed for 30 min, while water was removed. At the end of the 30 min feeding period, remaining feed was removed and weighed. Then, the test compound / reference compound or its vehicle in the control group was administered subcutaneously. 60 min later, to allow the compound to reach 25 relevant plasma exposure, a coloured, non-caloric bolus was instilled via gavage into the stomach. After another 30 min, the animals were sacrificed and the stomach and the small intestine prepared. The filled stomach was weighed, emptied, carefully cleaned and dried and reweighed. The stomach content, calculated as weight of filled subtracted by the weight of emptied stomach, indicated the degree of gastric emptying. The small intestine was 30 straightened without force and measured in length. Then the distance from the gastric beginning of the gut to the tip of the farthest travelled intestinal content bolus was measured. The intestinal passage was given as ratio in

- 83 -

percent of the latter distance and the total length of the small intestine. Comparable data can be obtained for both female and male mice.

Statistical analyses were performed with Everstat 6.0 by 1-way-ANOVA,
5 followed by Dunnett's as post-hoc test. Dunnett's Test was applied to compare versus vehicle control., Differences were considered statistically significant at the $p < 0.05$ level.

Automated assessment of feed intake in mice

10 Female NMRI-mice of a body weight between 20 and 30 g were used. Mice were adapted to housing conditions for at least one week and for at least one day single-caged in the assessment equipment, when basal data were recorded simultaneously. On the study day, test product was administered subcutaneously close to the lights-off phase (12 h lights off) and assessment
15 of feed consumption was directly started afterwards. Assessment included continued monitoring over 22 hours, while data are processed as mean over every 30 min. Repetition of this procedure over several days was possible. Restriction of assessment to 22 hours was for practical reasons to allow for reweighing of animals, refilling of feed and water and drug administration
20 between procedures. Results could be assessed as cumulated data over 22 hours or differentiated to 30 min intervals. Comparable data can be obtained for both female and male mice.

Statistical analyses were performed with Everstat 6.0 by two-way ANOVA on
25 repeated measures and Dunnett's post-hoc analyses. Differences were considered statistically significant at the $p < 0.05$ level.

30 Acute and subchronic effects of exendin-4 derivatives after subcutaneous treatment on blood glucose and body weight in female diet-induced obese (DIO) C57BL/6NCrl mice

18 weeks on high-fat diet (method 1)

- 84 -

Female C57BL/6NCrl mice were housed in groups in a specific pathogen-free barrier facility on a 12 h light/dark cycle with free access to water and high-fat diet. After 18 weeks on high-fat diet, mice were stratified to treatment groups (n = 8), so that each group had similar mean body weight.

5 An aged-matched group with *ad libitum* access to standard chow was included as standard control group.

Before the experiment, mice were subcutaneously (s.c.) injected with vehicle solution and weighed for 3 days to acclimate them to the procedures.

10 1) Acute effect on blood glucose in fed DIO mice: initial blood samples were taken just before first administration (s.c.) of vehicle (phosphate buffer solution) or the exendin-4 derivatives at doses of 10, 30 and 100 µg/kg (dissolved in phosphate buffer), respectively. The volume of administration was 5 mL/kg. The animals had access to water and their corresponding diet
15 during the experiment, food consumption was determined at all time points of blood sampling. Blood glucose levels were measured at t = 0.5 h, t = 1 h, t = 2 h, t = 4 h, t = 6 h, t = 8 h, and t = 24 h (method: d-glucose hexokinase, hemolysate, AU640 Beckman Coulter). Blood sampling was performed by tail incision without anaesthesia.

20 2) Subchronic effect on body weight: all animals were treated once daily s.c. in the afternoon, at the end of the light phase (12 h lights on) with either vehicle or exendin-4 derivatives at the abovementioned doses for 4 weeks. Body weight was recorded daily. On days 6 and 28, total fat mass was
25 measured by nuclear magnetic resonance (NMR) using a Bruker minispec (Ettlingen, Germany).

14 weeks of prefeeding with high-fat diet (method 2)

Female C57BL/6NCrl mice were housed in groups in a specific pathogen-free barrier facility on a 12 h light/dark cycle with free access to water and high-fat diet. After 14 weeks on high-fat diet, mice were stratified to treatment groups (n = 8), so that each group had similar mean body weight.

- 85 -

An aged-matched group with *ad libitum* access to standard chow and water was included as standard control group.

Before the experiment, mice were subcutaneously (s.c.) injected with vehicle solution and weighed for 3 days to acclimate them to the procedures.

5

Subchronic effect on body weight: all animals were treated once daily s.c. late afternoon, at the end of the light phase (LD 12:12) with either vehicle or exendin-4 derivatives at the abovementioned doses for 3 weeks. Body weight was recorded daily.

10

Statistical analyses were performed with Everstat 6.0 by repeated measures two-way ANOVA and Dunnett's post-hoc analyses (glucose profile) and 1-way-ANOVA, followed by Dunnett's post-hoc test (body weight, body fat). Differences versus vehicle-treated DIO control mice were considered 15 statistically significant at the p < 0.05 level.

Acute and subchronic effects of exendin-4 derivatives after subcutaneous treatment on blood glucose and HbA1c in female leptin-receptor deficient diabetic db/db mice (method 3)

20

Female BKS.Cg-m +/+ LepRdb/J (db/db) and BKS.Cg-m +/+ LepRdb/+ (lean control) mice were obtained from Charles River Laboratories, Germany, at an age of 9 – 10 weeks. The animals were housed in groups in a specific pathogen-free barrier facility on a 12-h light/dark cycle with free access to 25 water and rodent-standard chow. After 1 week of acclimatization, blood samples were drawn from the tail without anaesthesia and blood glucose (method: d-glucose hexokinase, hemolysate, AU640 Beckman Coulter) and HbA1c level (method: hemolysate, Cobas6000 c501, Roche Diagnostics, Germany) were determined.

30

HbA1c is a glycosylated form of haemoglobin whose level reflects the average level of glucose to which the erythrocyte has been exposed during

- 86 -

its lifetime. In mice, HbA1c is a relevant biomarker for the average blood glucose level during the preceding 4 weeks (erythrocyte life span in mouse ~ 47 days).

5 Db/db mice were stratified to treatment groups (n = 8), so that each group had similar baseline blood glucose and HbA1c levels.

10 1) Acute effect on blood glucose in fed db/db mice: initial blood samples were taken just before first administration (s.c.) of vehicle (phosphate buffer solution) or exendin-4 derivatives at doses of 3, 10, and 100 µg/kg (dissolved in phosphate buffer), respectively. The volume of administration was 5 mL/kg. The animals had access to water and chow during the experiment, food consumption was determined at all time points of blood sampling. Blood glucose levels were measured at t = 0.5 h, t = 1 h, t = 2 h, t = 4 h, t = 6 h, t = 8 h, and t = 24 h. Blood sampling was performed by tail incision without anaesthesia. Comparable data can be obtained for both female and male mice.

20 2) Subchronic effect on blood glucose and HbA1c: all animals were treated once daily s.c. in the afternoon, at the end of the light phase (12 h lights on) with either vehicle or exendin-4 derivatives at the abovementioned doses for 4 weeks. At the end of the study, blood samples (tail, no anaesthesia) were analyzed for glucose and HbA1c. Comparable data can be obtained for both female and male mice.

25

Statistical analyses were performed with Everstat 6.0 by repeated measures two-way ANOVA and Dunnett's post-hoc analyses. Differences versus vehicle-treated db/db control mice were considered statistically significant at the p < 0.05 level.

30

Effects of 4 weeks of treatment on glucose, HbA1c and oral glucose tolerance in female diabetic dbdb-mice (method 4)

8 week old, female diabetic dbdb-mice of mean non-fasted glucose value of 14.5 mmol/l and a body weight of 37-40 g were used. Mice were individually marked and were adapted to housing conditions for at least one week.

5 7 days prior to study start, baseline values for non-fasted glucose and HbA1c were determined, 5 days prior to study start, mice were assigned to groups and cages (5 mice per cage, 10 per group) according to their HbA1c values to ensure even distribution of lower and higher values between groups (stratification).

10 Mice were treated for 4 weeks, by once daily subcutaneous administration 3 hours prior to the dark phase (6 pm to 6 am). Blood samples from a tail tip incision were obtained for HbA1c on study day 21 and oral glucose tolerance was assessed in the 4th week. Oral glucose tolerance test was done in the morning without prior extra compound administration to majorly assess the

15 effect of chronic treatment and less of acute compound administration. Mice were fasted for 4 hours prior to oral glucose administration (2 g/kg, t = 0 min). Blood samples were drawn prior to glucose administration and at 15, 30, 60, 90, 120, and 180 min thereafter. Feed was returned after the last blood sampling. Results are represented as change from baseline, glucose

20 in mmol/l and HbA1c in %.

Statistical analyses are performed with Everstat Version 6.0 based on SAS by 1-way-ANOVA, followed by Dunnett's post-hoc test against vehicle-control. Differences are considered statistically significant at the p < 0.05 level.

25

Glucose lowering in non-fasted female diabetic dbdb-mice

Female diabetic dbdb-mice of mean non-fasted glucose value of 20-22 mmol/l and a body weight of 42 g +/- 0.6 g (SEM) were used. Mice were

30 individually marked and were adapted to housing conditions for at least one week.

3-5 days prior to study start mice were assigned to groups and cages (4 mice per cage, 8 per group) according to their non-fasted glucose values to

- 88 -

ensure even distribution of lower and higher values between groups (stratification). On the study day, mice were weighed and dosed (t = 0). Immediately prior to compound administration feed was removed while water remained available, and a first blood sample at a tail incision was drawn
5 (baseline). Further blood samples were drawn at the tail incision at 30, 60, 90, 120, 240, 360, and 480 min.

Statistical analyses are performed with Everstat Version 6.0 based on SAS by 2-way-ANOVA on repeated measures, followed by Dunnett's post-hoc test against vehicle-control. Differences are considered statistically
10 significant at the p < 0.05 level.

EXAMPLES

The invention is further illustrated by the following examples.

5 Example 1:

Synthesis of SEQ ID NO: 20

The solid phase synthesis was carried out on Rink-resin with a loading of 0.38 mmol/g, 75-150 µm from the company Agilent Technologies. The Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to literature (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603). Hereafter Fmoc-Glu-OtBu was coupled to the liberated amino-group employing the coupling reagents HBTU/DIPEA followed by Fmoc-deprotection with 20% piperidine in DMF. Finally heneicosanyl chloride was coupled to the amino-group of Glu in dichloromethane with DIPEA as base. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (XBridge, BEH130, Prep C18 5µM) using an acetonitrile/water gradient (both buffers with 0.05% TFA). The purified peptide was analysed by LCMS (Method C). Deconvolution of the mass signals found under the peak with retention time 31.67 min revealed the peptide mass 4647.40 which is in line with the expected value of 4647.35.

Example 2:

Synthesis of SEQ ID NO: 16

30

The solid phase synthesis was carried out on Novabiochem Rink-Amide resin (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-

- 90 -

norleucylaminomethyl resin), 100-200 mesh, loading of 0.34 mmol/g. The Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved
5 from the peptide on resin according to a modified literature procedure (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603), using 4% hydrazine hydrate in DMF. Hereafter Fmoc-Glu-OtBu was coupled to the liberated amino-group employing the coupling reagents HBTU/DIPEA followed by Fmoc-deprotection with 20% piperidine in DMF. Again Fmoc-Glu-OtBu was
10 coupled followed by Fmoc-deprotection and the final coupling of stearic acid using HBTU/DIPEA. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (Sunfire, Prep C18) using an acetonitrile/water gradient
15 (both buffers with 0.05% TFA). The purified peptide was analysed by LCMS (Method C). Deconvolution of the mass signals found under the peak with retention time 28.45 min revealed the peptide mass 4733.6 which is in line with the expected value of 4734.4.

20 Example 3:

Synthesis of SEQ ID NO: 17

The solid phase synthesis was carried out on Rink-resin with a loading of 0.38 mmol/g, 75-150 µm from the company Agilent Technologies. The Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to literature (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603). Hereafter Fmoc- γ -amino butyric acid was coupled to
25 the liberated amino-group employing the coupling reagents HBTU/DIPEA followed by Fmoc-deprotection with 20% piperidine in DMF. Finally stearic acid was coupled using HBTU/DIPEA. The peptide was cleaved from the
30

- 91 -

resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (XBridge, BEH130, Prep C18 5 μ M) using an acetonitrile/water gradient (both buffers with 0.05% TFA). The 5 purified peptide was analysed by LCMS (Method C). Deconvolution of the mass signals found under the peak with retention time 29.59 min revealed the peptide mass 4561.4 which is in line with the expected value of 4561.26.

Example 4:

10 Synthesis of SEQ ID NO: 18

The solid phase synthesis was carried out on Rink-resin with a loading of 0.38 mmol/g, 75-150 μ m from the company Agilent Technologies. The Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 15 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to literature (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603). Hereafter Fmoc- β -Ala-OH was coupled to the liberated amino-group employing the coupling reagents HBTU/DIPEA 20 followed by Fmoc-deprotection with 20% piperidine in DMF. Again Fmoc- β -Ala-OH was coupled followed by Fmoc-deprotection and the final coupling of stearic acid using HBTU/DIPEA. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative 25 HPLC on a Waters column (XBridge, BEH130, Prep C18 5 μ M) using an acetonitrile/water gradient (both buffers with 0.05% TFA). The purified peptide was analysed by LCMS (Method C). Deconvolution of the mass signals found under the peak with retention time 28.97 min revealed the peptide mass 4618.6 which is in line with the expected value of 4618.32.

30

Example 5:

Synthesis of SEQ ID NO: 9

The solid phase synthesis was carried out on Novabiochem Rink-Amide resin (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucylaminomethyl resin), 100-200 mesh, loading of 0.34 mmol/g. The 5 Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to a modified literature procedure (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603), using 4% hydrazine 10 hydrate in DMF. Hereafter Palm-Glu(γ OSu)-OtBu was coupled to the liberated amino-group. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (Sunfire, Prep C18) using an acetonitrile/water gradient 15 (both buffers with 0.1% TFA). The purified peptide was analysed by LCMS (Method B). Deconvolution of the mass signals found under the peak with retention time 12.7 min revealed the peptide mass 4577.3 which is in line with the expected value of 4577.22.

20 Example 6:

Synthesis of SEQ ID NO: 36

The solid phase synthesis was carried out on Novabiochem Rink-Amide resin (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucylaminomethyl resin), 100-200 mesh, loading of 0.34 mmol/g. The 25 Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to a modified literature procedure (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603), using 4% hydrazine 30 hydrate in DMF. Hereafter Palm-Glu(γ OSu)-OtBu was coupled to the liberated amino-group. The peptide was cleaved from the resin with King's

- 93 -

cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (Sunfire, Prep C18) using an acetonitrile/water gradient (both buffers with 0.05% TFA). The purified peptide was analysed by LCMS (Method B). Deconvolution of the mass signals found under the peak with retention time 12.53 min revealed the peptide mass 4489.57 which is in line with the expected value of 4490.13.

5 Example 7:

10 Synthesis of SEQ ID NO: 39

The solid phase synthesis was carried out on Novabiochem Rink-Amide resin (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucylaminomethyl resin), 100-200 mesh, loading of 0.34 mmol/g. The 15 Fmoc-synthesis strategy was applied with HBTU/DIPEA-activation. In position 1 Boc-Tyr(tBu)-OH and in position 14 Fmoc-Lys(ivDde)-OH was used in the solid phase synthesis protocol. The ivDde-group was cleaved from the peptide on resin according to a modified literature procedure (S.R. Chhabra et al., Tetrahedron Lett. 39, (1998), 1603), using 4% hydrazine 20 hydrate in DMF. Hereafter Palm-Glu(γ OSu)-OtBu was coupled to the liberated amino-group. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (Sunfire, Prep C18) using an acetonitrile/water gradient 25 (both buffers with 0.05% TFA). The purified peptide was analysed by LCMS (Method B). Deconvolution of the mass signals found under the peak with retention time 13.5 min revealed the peptide mass 4491.3 which is in line with the expected value of 4492.1.

30 In an analogous way, the following peptides SEQ ID NO: 8 - 41 were synthesized and characterized (Method A-E), see Table 5.

- 94 -

Table 5: list of synthesized peptides and comparison of calculated vs. found molecular weight.

SEQ ID NO:	calc. Mass	found mass
8	4576.2	4575.6
9	4577.2	4577.3
10	4478.0	4477.5
11	4462.1	4462.5
12	4548.1	4547.7
13	4506.1	4505.3
14	4561.2	4560.9
15	4605.3	4605.7
16	4734.4	4733.6
17	4561.3	4561.4
18	4618.3	4618.6
19	4648.3	4647.6
20	4647.4	4647.4
21	4520.1	4518.9
22	4464.0	4463.4
23	4565.1	4564.5
24	4522.1	4521.4
25	4579.1	4578.7
26	4620.2	4619.6
27	4563.2	4562.4
28	4504.1	4504.5
29	4477.0	4477.2
30	4420.0	4419.2
31	4505.1	4505.1
32	4477.1	4476.5
33	4519.1	4518.0
34	4533.2	4532.1

- 95 -

35	4449.0	4448.4
36	4490.1	4489.6
37	4491.1	4491.0
38	4590.3	4590.2
39	4492.1	4491.3
*40	4094.5	4092.3
*41	4194.6	4194.0

*non-acylated comparison compound

Example 8: Chemical stability and solubility

- 5 Solubility and chemical stability of peptidic compounds were assessed as described in Methods. The results are given in Table 6.

Table 6: Chemical stability and solubility

SEQ ID NO:	Stability (pH4.5) [%]	Stability (pH7.4) [%]	Solubility (pH4.5) [µg/ml]	Solubility (pH7.4) [µg/ml]	Method
8	98.0	98.0	>1000	971.7	D
9	92.5	97.7	>1000	>1000	D
12	100.0	95.3	>1000	>1000	D
13	86.8	95.9	267.7	>1000	D
14	96.0	94.0	>1000	>1000	D
15	91.0	90.0	997.0	>1000	D
17	100.0	100.0	970.0	>1000	E
31	94.0	96.0	>1000	>1000	D
35	100.0	98.0	424.5	>1000	D

10

Example 9: In vitro data on GLP-1, GIP and glucagon receptor
Potencies of peptidic compounds at the GLP-1, GIP and glucagon receptors
were determined by exposing cells expressing human glucagon receptor

- 96 -

(hGLUC R), human GIP (hGIP R) and human GLP-1 receptor (hGLP-1 R) to the listed compounds at increasing concentrations and measuring the formed cAMP as described in Methods.

- 5 The results for Exendin-4 derivatives with activity at the human GIP (hGIP R), human GLP-1 receptor (hGLP-1 R) and human glucagon receptor (hGLUC R) are shown in Table 7.
- 10

Table 7. EC₅₀ values of exendin-4 peptide analogues at GLP-1, GIP and Glucagon receptors (indicated in pM)

SEQ ID NO:	EC50 hGIP R [pM]	EC50 hGLP-1 R [pM]	EC50 hGLUC R [pM]
8	9.8	5.3	18.3
9	5.7	3.6	7710.0
10	15.1	13.2	40000.0
11	3.2	11.5	7220.0
12	8.9	12.7	1890.0
13	71.0	7.3	31.3
14	4.4	4.3	3760.0
15	8.2	8.1	5810.0
16	5.1	4.0	2890.0
17	9.6	8.7	9740.0
18	8.1	7.6	4950.0
19	13.8	4.0	707.5
20	24.5	23.2	3310.0
21	6.4	4.8	10100.0
22	16.6	32.0	11600.0
23	79.5	11.8	19100.0
24	23.5	13.5	38900.0
25	73.6	9.5	20500.0
26	19.7	4.9	8510.0
27	6.7	4.0	6390.0

- 97 -

28	10.9	3.2	9.9
29	127.0	7.0	46.8
30	22.1	12.0	226.0
31	6.5	6.0	3080.0
32	7.1	8.4	82.6
33	9.1	6.4	12900.0
34	22.2	4.6	11600.0
35	7.3	6.9	39100.0
36	6.4	3.4	5785.0
37	21.2	8.9	32.0
38	11.2	6.7	11.4
39	8.5	4.3	19300.0

Comparison Testing

A selection of inventive exendin-4 derivatives comprising a functionalized amino acid in position 14 has been tested versus corresponding compounds

5 having in this position 14 a 'non-functionalized' amino acid. The reference pair compounds and the corresponding EC50 values at GLP-1 and GIP receptors (indicated in pM) are given in Table 8. As shown, the inventive exendin-4 derivatives show a superior activity in comparison to the compounds with a 'non-functionalized' amino acid in position 14.

10

Table 8. Comparison of exendin-4 derivatives comprising a non-functionalized amino acid in position 14 vs. exendin-4 derivatives comprising a functionalized amino acid in position 14. EC50 values at GLP-1 and GIP receptors are indicated in pM. (K=lysine, Nle=norleucine, L=leucine, γ E-x53=(S)-4-Carboxy-4-hexadecanoylamino-butyryl-)

SEQ ID NO:	EC50 hGIP R [pM]	EC50 hGLP-1 R [pM]	residue in position 14
32	7.1	8.4	K(γ E-x53)
40	858	3.2	L

- 98 -

9	5.7	3.6	K(γ E-x53)
41	449	11.2	Nle

Example 10: Pharmacokinetic testing

Pharmacokinetic profiles were determined as described in Methods.

- 5 Calculated $T_{1/2}$ and c_{max} values are shown in Table 9.

Table 9. Pharmacokinetic profiles of exendin-4 derivatives.

SEQ ID NO:	Mice (1 mg/kg)		Mini pigs (0.1 mg/kg)	
	$T_{1/2}$ [h]	C_{max} [ng/ml]	$T_{1/2}$ [h]	C_{max} [ng/ml]
8	3.4	3740		
9	4.1	5470	12.2	278
10	2.7	5820		
12	2.8	3790		
13	3.1	3790		
14	2.8	5340		
15	3.5	5000		
16	5.3	3460		
18	2.1	5750		
21	4.0	5050	19.1	479
26	3.7	3120		
32	2.7	5520		
34	2.8	5130		

- 10 Example 11: Effect of SEQ ID NO: 9 and SEQ ID NO: 13 on gastric emptying and intestinal passage in female NMRI-mice

Female NMRI-mice, weighing on average 25 – 30 g, received 1, 3 and 10 μ g/kg of SEQ ID NO: 9, or 10 μ g/kg of SEQ ID NO: 13 or phosphate buffered

- 99 -

saline (vehicle control) subcutaneously, 60 min prior to the administration of the coloured bolus. 30 min later, the assessment of stomach contents and intestinal passage was done (Fig. 1 and 2).

- 5 In these studies, SEQ ID NO: 9 reduced intestinal passage by 49, 62 and 64 % (p<0.0001) and increased remaining gastric contents by 32, 79 and 111 % (p<0.0001), respectively. SEQ ID NO: 13 reduced intestinal passage by 60 % (p<0.0001) and increased remaining gastric contents by 40 % (p<0.0001), respectively. (p<0.0001 versus vehicle control, 1-W-ANOVA, followed by
10 Dunnett's post-hoc test).

Example 12:

Effect of SEQ ID NO: 12, SEQ ID NO: 13 and liraglutide on 22-hours food intake in female NMRI-mice

- 15 Fed female NMRI-mice, weighing on average 25-30 g, were administered 0.1 mg/kg of SEQ ID NO: 12, SEQ ID NO: 13, liraglutide or phosphate buffered saline (vehicle control) subcutaneously, directly prior to start of feeding monitoring. Lights-off phase (dark phase) started 4 hours later.
20 All tested compounds induced a pronounced reduction of feed intake, reaching after 22 hours for liraglutide 47% (p=0.006), for SEQ ID NO: 12 71% (p<0.0001) and SEQ ID NO: 13 93% (p=0.0003, 2-W-ANOVA-RM on ranks, post hoc Dunnett's Test) at the end of the study, respectively (Fig. 3a).

- 25 Effect of SEQ ID NO: 9 on 22-hours food intake in female NMRI-mice
Fed female NMRI-mice, weighing on average 25-30 g, were administered 3 µg/kg or 10 µg/kg of SEQ ID NO: 9 or phosphate buffered saline (vehicle control) subcutaneously, directly prior to start of feeding monitoring. Lights-off phase (dark phase) started 4 hours later.
30 SEQ ID NO: 9 induced a pronounced reduction of feed intake, reaching after 22 hours for 3 µg/kg 11% (not significant, p=0.78), and for 10 µg/kg 62% (p=

- 100 -

0.0005, 2-W-ANOVA-RM on ranks, post hoc Dunnett's Test) at the end of the study, respectively (Fig. 3b).

Example 13:

5 Subchronic effects of SEQ ID NO: 9 after subcutaneous treatment on blood glucose and body weight in female diet-induced obese (DIO) C57BL/6NCrl mice (18 weeks on high-fat diet, method 1)

10 1) Glucose profile

10 Diet-induced obese female C57BL/6NCrl mice were administered daily in the afternoon, at the end of the light phase (12 h lights on) with 10, 30 and 100 µg/kg of SEQ ID NO: 9 or phosphate buffered solution (vehicle control on standard or high-fat diet) subcutaneously. On day 6 of treatment and at predefined time points, more blood samples were taken to measure blood 15 glucose and generate the blood glucose profile over 24 h.

Already at the beginning of blood sampling on day 6 of treatment the basal blood glucose levels were dose-dependently decreased compared to DIO control mice (Fig. 4).

20 2) Body weight

Female obese C57BL/6NCrl mice were treated for 4 weeks once daily subcutaneously in the afternoon, at the end of the light phase (12 h lights on) with 10, 30 or 100 µg/kg SEQ ID NO: 9 or vehicle. Body weight was recorded daily, and body fat content was determined before the start of 25 treatment and after 4 weeks of treatment.

Comparable data can be obtained for both female and male mice.

30 Treatment with SEQ ID NO: 9 reduced body weight, whereas in the high-fat diet control group maintained body weight (Fig. 5 and Table 10). Calculating the relative body weight change from baseline values revealed a dose-dependent decrease of body weight, varying between 13.3-16.4% at 10 µg/kg, 17.6-20.9% at 30 µg/kg and 21.7-22.7% at 100 µg/kg (Fig. 6). These

- 101 -

changes resulted from a decrease in body fat, as shown by the absolute changes in body fat content (Fig. 7 and Table 10).

Table 10. Weight change in DIO mice over a 4-week treatment period (mean

5 \pm SEM)

Example (Dose)	Overall weight change (g)	Bodyfat change (g)
Control standard diet	-0.7 \pm 0.2	1.5 \pm 0.2
Control high-fat diet	-0.98 \pm 0.4	-0.8 \pm 0.4
SEQ ID NO: 9 (10 μ g/kg)	-5.0 \pm 0.5	-4.2 \pm 0.4
SEQ ID NO: 9 (30 μ g/kg)	-6.9 \pm 0.5	-6.1 \pm 0.4
SEQ ID NO: 9 (100 μ g/kg)	-7.3 \pm 0.6	-6.4 \pm 0.5
Liraglutide (100 μ g/kg)	-3.9 \pm 0.4	-3.3 \pm 0.4

Example 14: Acute and subchronic effects of SEQ ID NO: 13, SEQ ID NO: 12, SEQ ID NO: 10 and SEQ ID NO: 9 after subcutaneous treatment on blood glucose and HbA1c in female leptin-receptor deficient diabetic db/db mice (method 3)

1) Glucose profile

After blood sampling to determine the blood glucose baseline level, fed diabetic female db/db mice were administered 100 μ g/kg of SEQ ID NO:

15 13, SEQ ID NO: 12, SEQ ID NO: 10 and SEQ ID NO: 9 or phosphate buffered solution (vehicle-treated db/db control) subcutaneously in the morning, at the beginning of the light phase (12 h lights on). At predefined time points, more blood samples were taken to measure blood glucose and generate the blood glucose profile over 24 h.

20 Comparable data can be obtained for both female and male mice.

At the tested dose, of SEQ ID NO: 13, SEQ ID NO: 12, SEQ ID NO: 10 and SEQ ID NO: 9 demonstrated a significant decrease in blood glucose compared to db/db control mice, lasting longer than 24 h in the SEQ ID NO:

- 102 -

10 and SEQ ID NO: 9 dose group (Fig. 8).

2) Blood glucose & HbA1c

Female diabetic mice were treated for 4 weeks once daily subcutaneously

5 with 10, 30 or 100 µg/kg SEQ ID NO: 9 or vehicle in the morning, at the beginning of the light phase (12 h lights on). Blood glucose and HbA1c were determined before start of treatment and at the end of the study after 4 weeks of treatment. A strong and dose-dependent decrease in blood

10 glucose, superior to liraglutide in the medium and highest dose could be observed (Fig. 9). Before treatment started, no significant differences in blood glucose levels could be detected between db/db groups, only the lean control animals had significant lower glucose levels. During the 4 weeks of

15 treatment, glucose levels increased in the vehicle-treated db/db control group, indicating a worsening of the diabetic situation. All SEQ ID NO: 9– treated animals displayed a significant lower blood glucose level than the db control mice at the end of the study.

Comparable data can be obtained for both female and male mice.

Corresponding to blood glucose, at start of the study, no significant

20 differences in HbA1c levels could be detected between db/db groups, only the lean control animals had significant lower levels. During the 4 weeks of treatment, HbA1c increased in the vehicle-treated db/db control group, corresponding to the increasing blood glucose levels. Animals treated with

25 SEQ ID NO: 9 displayed a lower HbA1c level than the db/db control mice at the end of the study in all three doses (Fig. 10).

Example 15: Subchronic effects of SEQ ID NO: 9 and SEQ ID NO: 21 after subcutaneous treatment on body weight in female diet-induced obese (DIO) C57BL/6NCrl mice (14 weeks of prefeeding with high-fat diet, method 2)

30

Female obese C57BL/6NCrl mice were treated for 3 weeks once daily subcutaneously in the late afternoon, prior the end of the light phase (12 h

- 103 -

lights on) with 10 µg/kg SEQ ID NO: 9 and SEQ ID NO: 21 or vehicle. Body weight was recorded daily.

Treatment with SEQ ID NO: 9 and SEQ ID NO: 21 reduced body weight,

5 whereas the high-fat diet control group even gained body weight (Fig. 11 and Table 11). Calculating the relative body weight change from baseline values revealed a decrease of body weight up to 15.1% at 10 µg/kg SEQ ID NO: 9 and 18.0% at 10 µg/kg SEQ ID NO: 21 (Fig. 12).

10 Table 11. Weight change in DIO mice over a 3-week treatment period (mean ± SEM)

Example (Dose)	Overall weight change (g)
Control standard diet	+0.3 ± 0.2
Control high-fat diet	+2.7 ± 0.3
SEQ ID NO: 9 (10 µg/kg)	-6.2 ± 0.4
SEQ ID NO: 21 (10 µg/kg)	-7.3 ± 0.7

Example 16: Effects of 4 weeks of treatment with SEQ ID NO: 16, and SEQ ID NO: 21 on glucose, HbA1c and oral glucose tolerance in female diabetic dbdb-mice (method 4)

15 Female dbdb-mice, received 3 and 10 µg/kg of SEQ ID NO: 16 and 10 µg/kg of SEQ ID NO: 21 or phosphate buffered saline (vehicle control) once daily, subcutaneously over four weeks.

20 Both compounds reached a statistical significant reduction of non-fasted glucose compared to vehicle control at the 10 µg/kg dose (Fig. 13); p<0.05, 1-way-ANOVA, followed by Dunnett's post-hoc test.

Furthermore, both compounds prevented an increase of HbA1c in a statistical significant manner compared to vehicle control at the 10 µg/kg

25 dose (Fig. 14); (p<0.05, 1-way-ANOVA, followed by Dunnett's post-hoc test).

Treatment with SEQ ID NO: 16, and SEQ ID NO: 21 lead to improved oral glucose tolerance (Fig. 15; represented as normalized to 0 mmol/l at 0 min),

- 104 -

and reduction of AUC under the glucose curve reached statistical significance compared to vehicle control (Fig. 16); (p<0.05, 1-way-ANOVA, followed by Dunnett's post-hoc test).

- 5 Example 17: SEQ ID NO: 14 and SEQ ID NO: 21 on glucose lowering in non-fasted female diabetic dbdb-mice

Female dbdb-mice, received 3 µg/kg of SEQ ID NO: 14, SEQ ID NO: 21 or phosphate buffered saline (vehicle control) subcutaneously, at time 0 min.

10 Both compounds immediately lowered glucose values (baseline at 20-22 mmol/l), with SEQ ID NO: 14 reaching the maximal effect of ~8 mmol/l (Fig. 18) and SEQ ID NO: 21 of 10 – 12 mmol/l glucose reduction (Fig. 17), respectively, at 240 min and keeping it to the end of observation at 480 min.

15 Both compounds reached a statistical significant reduction of glucose compared to vehicle control from t = 60 min until end of observation (p<0.05, 2-way-ANOVA on repeated measures, followed by Dunnett's post-hoc test).

Table 12: Sequences

SEQ ID NO:	sequence
1	H-G-E-G-T-F-T-S-D-L-S-K-Q-M-E-E-E-A-V-R-L-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH2
2	H-A-E-G-T-F-T-S-D-V-S-S-Y-L-E-G-Q-A-A-K-E-F-I-A-W-L-V-K-G-R-NH2
3	H-A-E-G-T-F-T-S-D-V-S-S-Y-L-E-G-Q-A-A-K(γ E-x53)-E-F-I-A-W-L-V-R-G-R-G
4	Y-A-E-G-T-F-I-S-D-Y-S-I-A-M-D-K-I-H-Q-Q-D-F-V-N-W-L-L-A-Q-K-G-K-K-N-D-W-K-H-N-I-T-Q
5	H-S-Q-G-T-F-T-S-D-Y-S-K-Y-L-D-S-R-R-A-Q-D-F-V-Q-W-L-M-N-T
6	Y-G-E-G-T-F-T-S-D-L-S-I-Q-M-E-E-E-A-V-R-L-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH2
7	Y-A-E-G-T-F-T-S-D-V-S-I-Y-L-E-G-Q-A-A-K-E-F-I-A-W-L-V-K-G-R-NH2
8	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
9	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
10	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
11	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-K-R-A-A-Aib-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
12	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-K-R-A-A-Q-D-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH2
13	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-S-R-A-A-Q-D-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH2
14	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x70)-E-K-R-A-A-Q-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
15	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x70)-E-K-R-A-A-Aib-E-

	F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
16	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E- γ E-x70)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
17	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(GABA-x70)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
18	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(β A- β A-x70)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
19	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x70)-E-K-R-A-A-Q-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
20	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x76)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
21	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x70)-E-S-R-A-A-Q-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
22	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-S-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
23	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Q-D-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
24	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Aib-D-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
25	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Q-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
26	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Q-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
27	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Aib-D-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
28	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-K-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
29	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-S-R-A-A-Q-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
30	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-S-R-A-A-Aib-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
31	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-K-R-A-A-Q-D-F-

	I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
32	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
33	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
34	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Q-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
35	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Aib-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
36	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-K-R-A-A-Aib-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
37	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-S-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
38	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-D-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2
39	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-K(γ E-x53)-E-S-R-A-A-Q-E-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
40	Y-Aib-Q-G-T-F-T-S-D-L-S-I-Q-L-E-S-R-A-A-Q-D-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH2
41	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-Nle-E-K-R-A-A-Aib-E-F-I-E-W-L-K-N-T-G-P-S-S-G-A-P-P-P-S-NH2

Claims

1. A peptidic compound having the formula (I):

5



wherein Z is a peptide moiety having the formula (II)

10 Tyr-Aib-X3-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-X12-Gln-X14-X15-X16-X17-
X18-X19-X20-X21-Phe-Ile-Glu-Trp-Leu-Lys-X28-X29-Gly-Pro-Ser-Ser-Gly-
Ala-Pro-Pro-Pro-Ser-X40 (II)

X3 represents an amino acid residue selected from Gln, Glu and His,

15 X12 represents an amino acid residue selected from Ile and Lys,

X14 represents an amino acid residue having a side chain with an -NH₂ group, wherein the -NH₂ side chain group is functionalized by - C(O)-R⁵, wherein R⁵ may be a moiety comprising up to 50 or up to 100 carbon atoms and optionally heteroatoms selected from halogen, N, O, S and/or P,

20 X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser, Lys, Glu and Gln,

X17 represents an amino acid residue selected from Arg, Lys, Glu, Gln, Leu, Aib, Tyr and Ala,

X18 represents an amino acid residue selected from Ala and Arg,

25 X19 represents an amino acid residue selected from Ala and Val,

X20 represents an amino acid residue selected from Gln, Aib, Lys and His,

X21 represents an amino acid residue selected from Asp, Glu and Leu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly, Thr and D-Ala,

30 X40 is either absent or represents Lys,

R¹ represents NH₂,

R^2 represents the C-terminal group of the peptidic compound and is selected from OH and NH₂,

or a salt or solvate thereof.

5

2. A compound of claim 1, wherein

X14 represents an amino acid residue with a functionalized -NH₂ side chain group, such as functionalized Lys, Orn, Dab or Dap, wherein at least one H atom of the -NH₂ side chain group is replaced by -C(O)-R⁵, which is selected from

(S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, 4-Hexadecanoylamino-butyryl-, 4-{3-[(R)-2,5,7,8-tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy carbonyl]-propionylamino}-butyryl-, 4-octadecanoylamino-butyryl-, 4-

15 ((Z)-octadec-9-enoylamino)-butyryl-, 6-[(4,4-Diphenyl-cyclohexyloxy)-hydroxy-phosphoryloxy]-hexanoyl-, Hexadecanoyl-, (S)-4-Carboxy-4-(15-carboxy-pentadecanoylamino)-butyryl-, (S)-4-Carboxy-4-{3-[3-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-propionylamino]-propionylamino}-butyryl-, (S)-4-Carboxy-4-{3-[(R)-2,5,7,8-

20 tetramethyl-2-((4R,8R)-4,8,12-trimethyl-tridecyl)-chroman-6-yloxy carbonyl]-propionylamino}-butyryl-, (S)-4-Carboxy-4-((9Z,12Z)-octadeca-9,12-dienoylamino)-butyryl-, (S)-4-Carboxy-4-[6-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-hexanoylamino]-butyryl-, (S)-4-

25 Carboxy-4-((2S,3R,4S,5R)-5-carboxy-2,3,4,5-tetrahydroxy-pentanoylamino)-butyryl-, (S)-4-Carboxy-4-tetradecanoylamino-butyryl-, (S)-4-(11-

Benzylloxycarbonyl-undecanoylamino)-4-carboxy-butyryl-, (S)-4-Carboxy-4-[11-((2S,3R,4R,5R)-2,3,4,5,6-pentahydroxy-hexylcarbamoyl)-

undecanoylamino]-butyryl-, (S)-4-Carboxy-4-((Z)-octadec-9-enoylamino)-butyryl-, (S)-4-Carboxy-4-(4-dodecyloxy-benzoylamino)-butyryl-, (S)-4-

30 Carboxy-4-henicosanoylamino-butyryl-, (S)-4-Carboxy-4-docosanoylamino-butyryl-, (S)-4-Carboxy-4-((Z)-nonadec-10-enoylamino)-butyryl-, (S)-4-Carboxy-4-(4-decyloxy-benzoylamino)-butyryl-, (S)-4-Carboxy-4-[(4'-

- 110 -

carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-
5 acetylamino]-butyrylamino}-butyryl-, (S)-4-Carboxy-2-[2-{2-[(S)-4-carboxy-
4-(17-carboxy-heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-
acetylamino]-butyryl-, 2-(2-{2-[(S)-4-Carboxy-4-(17-carboxy-
heptadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-
ethoxy)-acetyl-, 2-(2-{2-[(S)-4-Carboxy-4-(17-carboxy-heptadecanoylamino)-
butyrylamino]-ethoxy}-ethoxy)-acetyl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-
10 {(S)-4-carboxy-4-[(S)-4-carboxy-4-(19-carboxy-nonadecanoylamino)-
butyrylamino]-butyrylamino}-butyryl-, 2-(2-{2-[(S)-4-
Carboxy-4-(16-1H-tetrazol-5-yl-hexadecanoylamino)-butyrylamino]-ethoxy}-
ethoxy)-acetylamino]-ethoxy}-ethoxy)-acetyl-, 2-(2-{2-[(S)-4-
Carboxy-4-(16-carboxy-hexadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-
acetylamino]-ethoxy)-acetyl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-[(S)-
4-carboxy-4-(17-carboxy-heptadecanoylamino)-butyrylamino]-butyrylamino}-
15 butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-{2-[2-{2-[(S)-4-
carboxy-4-[10-(4-carboxy-phenoxy)-decanoylamino]-butyrylamino]-ethoxy}-
ethoxy)-acetylamino}-ethoxy)-acetylamino]-butyryl-, (S)-4-Carboxy-4-((S)-4-
carboxy-4-[2-(2-{2-[(S)-4-carboxy-4-(7-carboxy-heptanoylamino)-
butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy)-acetylamino]-
20 butyrylamino}-butyryl-, (S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-
[(S)-4-carboxy-4-(11-carboxy-undecanoylamino)-butyrylamino]-ethoxy}-
ethoxy)-acetylamino]-ethoxy)-acetylamino]-butyrylamino}-butyryl-,
(S)-4-Carboxy-4-{(S)-4-carboxy-4-[2-(2-{2-[(S)-4-carboxy-4-(13-
carboxy-tridecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-
25 ethoxy)-acetylamino]-butyrylamino}-butyryl-, (S)-4-Carboxy-4-{(S)-4-
carboxy-4-[2-(2-{2-[(S)-4-carboxy-4-(15-carboxy-
pentadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-
ethoxy)-acetylamino]-butyrylamino}-butyryl-, and (S)-4-Carboxy-4-{(S)-4-
carboxy-4-[2-(2-{2-[(S)-4-carboxy-4-(19-carboxy-
30 nonadecanoylamino)-butyrylamino]-ethoxy}-ethoxy)-acetylamino]-ethoxy}-
ethoxy)-acetylamino]-butyrylamino}-butyryl-,
X40 is absent or represents Lys.

3. A compound of any one of claims 1 - 2, wherein

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-

5 butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, 4-octadecanoylamino-butyryl-, Hexadecanoyl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl-.

10 4. A compound according to any one of claims 1-3,

wherein X14 is Lys functionalized with C(O)-R⁵, wherein R⁵ is selected from the group consisting of (S)-4-carboxy-4-hexadecanoylamino-butyryl (γE-x53), (S)-4-carboxy-4-octadecanoylamino-butyryl (γE-x70), (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)-butyryl (γE-γE-x70), 4-

15 octadecanoylamino-butyryl (GABA-x70), (S)-4-Carboxy-4-henicosanoylamino-butyryl (γE-x76), and 3-(3-Octadecanoylamino-propionylamino)-propionyl (β-Ala-β-Ala-x70).

20 5. A compound of any one of claims 1 - 4,

wherein R² is NH₂.

6. A compound according to any one of claims 1-5,

wherein the peptidic compound has a relative activity of at least 0.04%, preferably at least 0.08%, more preferably at least 0.2% compared to that of

25 natural GIP at the GIP receptor.

7. A compound according to any one of claims 1-6, wherein the peptidic

compound exhibits a relative activity of at least 0.07%, preferably at least 0.1%, more preferably at least 0.14%, more preferably at least

30 0.35% and even more preferably at least 0.4% compared to that of GLP-1(7-36) at the GLP-1 receptor.

8. A compound according to any one of claims 6 or 7, wherein the peptidic compound further exhibits a relative activity of at least 0.1%, preferably at least 0.2%, more preferably at least 0.3%, more preferably at least 0.4% and even more preferably at least 0.5% compared to that of natural glucagon at the glucagon receptor.

5 9. A compound of any one of claims 1 - 8,

wherein

X3 represents an amino acid residue selected from Gln and Glu,

10 X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)amino)-butyryl-, 3-(3-

15 Octadecanoylamino-propionylamino)-propionyl- and 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-,

X15 represents an amino acid residue selected from Glu and Asp,

X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

20 X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

25 X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

10. A compound of any one of claims 1 – 9, wherein

X3 represents Glu,

30 X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-

butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyrylamino)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl- and 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-butyryl-,

- 5 X15 represents an amino acid residue selected from Glu and Asp,
X16 represents an amino acid residue selected from Ser and Lys,
X17 represents Arg,
X18 represents Ala,
X19 represents Ala,
10 X20 represents an amino acid residue selected from Gln and Aib,
X21 represents an amino acid residue selected from Asp and Glu,
X28 represents an amino acid residue selected from Asn and Ala,
X29 represents an amino acid residue selected from Gly and Thr,
X40 is absent.

- 15 11. A compound of any one of claims 1 – 10, wherein
X14 represents Lys, wherein the -NH₂ side chain group is functionalized by
one of the groups selected from (S)-4-Carboxy-4-octadecanoylamino-
butyryl-, 4-octadecanoylamino-butyryl-, (S)-4-Carboxy-4-henicosanoylamino-
butyryl-, (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-
butyrylamino)-butyryl-, 3-(3-Octadecanoylamino-propionylamino)-propionyl-.

12. A compound of any one of claims 1 – 9,
wherein
25 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by
one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-
butyryl-, (S)-4-Carboxy-4-octadecanoylamino-butyryl-.

- 30 13. A compound of any one of claims 1 – 9, 12,
wherein
X3 represents an amino acid residue selected from Gln and Glu,
X12 represents Ile,

- 115 -

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by one of the groups selected from (S)-4-Carboxy-4-hexadecanoylamino-butyryl- and (S)-4-Carboxy-4-octadecanoylamino-butyryl-,

X15 represents an amino acid residue selected from Glu and Asp,

5 X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

10 X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

15 14. A compound of any one of claims 1 - 13,

wherein X19 represents Ala.

15. A compound of any one of claims 1 - 14,

wherein

20 X28 represents Ala,

X29 represents Gly.

16. A compound of any one of claims 1 - 14,

wherein

25 X28 represents Asn,

X29 represents Thr.

17. A compound of any one of claims 1 - 9, wherein

X3 represents an amino acid residue selected from Gln and Glu,

30 X12 represents Ile,

X14 represents Lys, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, wherein R⁵ is selected from (S)-4-Carboxy-4-hexadecanoylamino-

- 116 -

butyryl- (γ E-x53), (S)-4-Carboxy-4-octadecanoylamino-butyryl- (γ E-x70), (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyrylamino)-butyryl- (γ E- γ E-x70), 3-(3-Octadecanoylamino-propionylamino)-propionyl- (β A- β A-x70), 4-octadecanoylamino-butyryl- (GABA-x70), and (S)-4-Carboxy-4-

5 heneicosanoylamino-butyryl- (γ E-x76),

X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser and Lys,

X17 represents Arg,

X18 represents Ala,

10 X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

X21 represents an amino acid residue selected from Asp and Glu,

X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

15 X40 is absent.

18. A compound of any one of claims 1-9,12-17, wherein

X3 represents an amino acid residue selected from Gln and Glu,

X12 represents Ile,

20 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, wherein R⁵ is (S)-4-Carboxy-4-hexadecanoylamino-butyryl- (γ E-x53),

X15 represents an amino acid residue selected from Asp and Glu,

X16 represents an amino acid residue selected from Ser and Lys,

25 X17 represents Arg,

X18 represents Ala,

X19 represents Ala,

X20 represents an amino acid residue selected from Gln and Aib,

X21 represents an amino acid residue selected from Asp and Glu,

30 X28 represents an amino acid residue selected from Asn and Ala,

X29 represents an amino acid residue selected from Gly and Thr,

X40 is absent.

19. A compound of any one of claims 1-11, 14-17, wherein
X3 represents Glu,
X12 represents Ile,
- 5 X14 represents Lys, wherein the -NH₂ side chain group is functionalized by -C(O)-R⁵, wherein R⁵ is selected from (S)-4-Carboxy-4-octadecanoylamino-butyryl- (γE-x70), (S)-4-Carboxy-4-((S)-4-carboxy-4-octadecanoylamino-butyryl)amino)-butyryl- (γE-γE-x70), 3-(3-Octadecanoylamino-propionylamino)-propionyl- (βA-βA-x70), 4-octadecanoylamino-butyryl- (GABA-x70), and (S)-4-Carboxy-4-henicosanoylamino-butyryl- (γE-x76),
- 10 X15 represents Glu,
X16 represents an amino acid residue selected from Ser and Lys,
X17 represents Arg,
X18 represents Ala,
- 15 X19 represents Ala,
X20 represents an amino acid residue selected from Gln and Aib,
X21 represents Glu,
X28 represents an amino acid residue selected from Asn and Ala,
X29 represents an amino acid residue selected from Gly and Thr,
- 20 X40 is absent.
20. The compound of any one of claims 1-19, selected from the compounds of SEQ ID NO: 8-39 or a salt or solvate thereof.
- 25 21. The compound of any one of claims 1-19, selected from the compounds of SEQ ID NO: 8-10 and 12-38 or a salt or solvate thereof.
- 30 22. The compound of claim 20, selected from the compounds of SEQ ID NO: 8-13 and 39 or a salt or solvate thereof.
23. The compound of claim 21, selected from the compounds of SEQ ID

NO: 8-10 and 12-13 or a salt or solvate thereof.

24. The compound of claim 20, selected from the compounds of SEQ ID NO: 14-21 or a salt or solvate thereof.

5

25. The compound of claim 20, selected from the compounds of SEQ ID NO: 22-38 or a salt or solvate thereof.

- 10 26. The compound of any one of claims 1-25 for use in medicine, particularly in human medicine.

- 15 27. The compound for use according to claim 26 which is present as an active agent in a pharmaceutical composition together with at least one pharmaceutically acceptable carrier.

- 20 28. The compound for use according to claim 26 or 27 together with at least one additional therapeutically active agent, wherein the additional therapeutically active agent is selected from the series of Insulin and Insulin derivatives, GLP-1, GLP-1 analogues and GLP-1 receptor agonists, polymer bound GLP-1 and GLP-1 analogues, dual GLP1/glucagon agonists, PYY3-36 or analogues thereof, pancreatic polypeptide or analogues thereof, Glucagon receptor agonists, GIP receptor agonists or antagonists, ghrelin antagonists or inverse agonists, Xenin and analogues thereof, DDP-IV inhibitors, SGLT2

- 25 inhibitors, dual SGLT2 / SGLT1 inhibitors, Biguanides Thiazolidinediones, dual PPAR agonists, Sulfonylureas, Meglitinides, alpha-glucosidase inhibitors, Amylin and Amylin analogues, GPR119 agonists, GPR40 agonists, GPR120 agonists, GPR142 agonists, systemic or low-absorbable TGR5 agonists, Cycloset, inhibitors of 11-


- 30 beta-HSD, activators of glucokinase, inhibitors of DGAT, inhibitors of protein tyrosinephosphatase 1, inhibitors of glucose-6-phosphatase, inhibitors of fructose-1,6-bisphosphatase, inhibitors of glycogen

phosphorylase, inhibitors of phosphoenol pyruvate carboxykinase, inhibitors of glycogen synthase kinase, inhibitors of pyruvate dehydrogenase kinase, alpha2-antagonists, CCR-2 antagonists, modulators of glucose transporter-4, Somatostatin receptor 3 agonists, HMG-CoA-reductase inhibitors, fibrates, nicotinic acid and the derivatives thereof, nicotinic acid receptor 1 agonists, PPAR-alpha, gamma or alpha/gamma) agonists or modulators, PPAR-delta agonists, ACAT inhibitors, cholesterol absorption inhibitors, bile acid-binding substances, IBAT inhibitors, MTP inhibitors, modulators of PCSK9, LDL receptor up-regulators by liver selective thyroid hormone receptor β agonists, HDL-raising compounds, lipid metabolism modulators, PLA2 inhibitors, ApoA-I enhancers, thyroid hormone receptor agonists, cholesterol synthesis inhibitors, omega-3 fatty acids and derivatives thereof, active substances for the treatment of obesity, such as Sibutramine, Tesofensine, Orlistat, CB-1 receptor antagonists, MCH-1 antagonists, MC4 receptor agonists and partial agonists, NPY5 or NPY2 antagonists, NPY4 agonists, beta-3-agonists, leptin or leptin mimetics, agonists of the 5HT2c receptor, or the combinations of bupropione/naltrexone (CONTRAVE), bupropione/zonisamide (EMPATIC), bupropione/phentermine or pramlintide/metreleptin, QNEXA (Phentermine+ topiramate), lipase inhibitors, angiogenesis inhibitors, H3 antagonists, AgRP inhibitors, triple monoamine uptake inhibitors (norepinephrine and acetylcholine), MetAP2 inhibitors, nasal formulation of the calcium channel blocker diltiazem, antisense against production of fibroblast growth factor receptor 4, prohibitin targeting peptide-1, drugs for influencing high blood pressure, chronic heart failure or atherosclerosis, such as angiotensin II receptor antagonists, ACE inhibitors, ECE inhibitors, diuretics, beta-blockers, calcium antagonists, centrally acting hypertensives, antagonists of the alpha-2-adrenergic receptor, inhibitors of neutral endopeptidase, thrombocyte aggregation inhibitors.

29. The compound for use according to claim 26 or 27 together with at least one additional therapeutically active agent, wherein the additional therapeutically active agent particularly is a GLP-1 agonist and/or insulin or an insulin analogue and/or a gastrointestinal peptide.
- 5
30. The compound for use according to any one of claims 26-29 for the treatment or prevention of hyperglycemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, obesity, metabolic syndrome and neurodegenerative disorders, particularly for delaying or preventing disease progression in type 2 diabetes, treating metabolic syndrome, treating obesity or preventing overweight, for decreasing food intake, increase energy expenditure, reducing body weight, delaying the progression from impaired glucose tolerance (IGT) to type 2 diabetes; delaying the progression from type 2 diabetes to insulin-requiring diabetes; regulating appetite; inducing satiety; preventing weight regain after successful weight loss; treating a disease or state related to overweight or obesity; treating bulimia; treating binge eating; 15 treating atherosclerosis, hypertension, IGT, dyslipidemia, coronary heart disease, hepatic steatosis, treatment of beta-blocker poisoning, use for inhibition of the motility of the gastro-intestinal tract, useful in connection with investigations of the gastro-intestinal tract using techniques such as X-ray, CT- and NMR-scanning.
- 20
- 25
31. The compound for use according to any one of claims 26-30 for the treatment or prevention of hyperglycemia, type 2 diabetes, obesity.

Fig. 1

a)



b)





Fig. 2


a)



■ Vehicle Control
■ SEQ ID NO: 9, 3 µg/kg

■ SEQ ID NO: 9, 1 µg/kg
■ SEQ ID NO: 9, 10 µg/kg

b)



■ Vehicle Control
■ SEQ ID NO: 9, 3 µg/kg

■ SEQ ID NO: 9, 1 µg/kg
■ SEQ ID NO: 9, 10 µg/kg

Fig. 3:
a)






Fig. 3:
b)



5/19

Fig. 4



6/19

Fig. 5:





Fig. 6:







Fig. 7:


9/19

Fig. 8:




10/19

Fig. 9:



11/19

Fig. 10:



12/19




Fig. 11:

13/19

Fig. 12:






Fig. 13:



15/19



16/19




Fig. 16:






Fig. 17:



19/19

Fig. 18:



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/077310

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K38/26 C07K14/605
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2008/081418 A1 (COVX TECHNOLOGIES IRELAND LTD [IE]; BRADSHAW CURT [US]; SAKAMURI SUKUM) 10 July 2008 (2008-07-10) paragraphs [0001], [0009], [0059], [0068], [0281], [0283] table IV ----- WO 2008/023050 A1 (NOVO NORDISK AS [DK]; LAU JESPER [DK]; HANSEN THOMAS KRUSE [DK]; CHRIS) 28 February 2008 (2008-02-28) page 1, line 28 - line 31 page 2 ----- -/-	1-31
Y		1-31

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
10 February 2014	18/02/2014
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Surdej, Patrick

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/077310

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2011/094337 A1 (UNIV INDIANA RES & TECH CORP [US]; DIMARCI RICHARD D [US]; MATAO [U] 4 August 2011 (2011-08-04) paragraph [0594] - paragraph [0596] table 8 figure 5 -----	1-31
Y	US 2011/237503 A1 (ALSINA-FERNANDEZ JORGE [US] ET AL) 29 September 2011 (2011-09-29) paragraph [0006] - paragraph [0007] tables 5-7 examples 1-5 -----	1-31

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/077310

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
WO 2008081418	A1 10-07-2008	AU 2008203641	A1 10-07-2008	BR PI0806308	A2 06-09-2011
		CA 2674112	A1 10-07-2008	EP 2118131	A1 18-11-2009
		JP 5009376	B2 22-08-2012	JP 2010514835	A 06-05-2010
		JP 2012184232	A 27-09-2012	KR 20090096498	A 10-09-2009
		KR 20120083510	A 25-07-2012	NZ 577686	A 25-11-2011
		US 2009098130	A1 16-04-2009	WO 2008081418	A1 10-07-2008
WO 2008023050	A1 28-02-2008	EP 2057189	A1 13-05-2009	US 2009318353	A1 24-12-2009
		US 2012283170	A1 08-11-2012	WO 2008023050	A1 28-02-2008
WO 2011094337	A1 04-08-2011	CA 2788304	A1 04-08-2011	CN 102834108	A 19-12-2012
		EP 2528618	A1 05-12-2012	JP 2013518115	A 20-05-2013
		KR 20120123443	A 08-11-2012	US 2012322725	A1 20-12-2012
		WO 2011094337	A1 04-08-2011		
US 2011237503	A1 29-09-2011	AR 080592	A1 18-04-2012	AU 2011232597	A1 30-08-2012
		CA 2794664	A1 29-09-2011	CN 102821779	A 12-12-2012
		CO 6630128	A2 01-03-2013	CR 20120457	A 16-10-2012
		DO P2012000244	A 15-02-2013	EA 201270723	A1 30-01-2013
		EP 2552471	A1 06-02-2013	JP 2013523647	A 17-06-2013
		KR 20120133402	A 10-12-2012	MA 34077	B1 05-03-2013
		PE 05232013	A1 25-04-2013	SG 184289	A1 29-11-2012
		TW 201201829	A 16-01-2012	US 2011237503	A1 29-09-2011
		WO 2011119657	A1 29-09-2011		