

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2013242405 B2**

(54) Title
Artificial nucleic acid molecules comprising a 5'TOP UTR

(51) International Patent Classification(s)
C12N 15/67 (2006.01)

(21) Application No: **2013242405**

(22) Date of Filing: **2013.03.27**

(87) WIPO No: **WO13/143700**

(30) Priority Data

(31) Number	(32) Date	(33) Country
PCT/EP2012/002448	2012.06.08	EP
PCT/EP2012/001334	2012.03.27	EP

(43) Publication Date: **2013.10.03**

(44) Accepted Journal Date: **2018.06.28**

(71) Applicant(s)
CureVac GmbH

(72) Inventor(s)
Thess, Andreas

(74) Agent / Attorney
Allens Patent & Trade Mark Attorneys, Deutsche Bank Place Corner Hunter and Phillip Streets, SYDNEY, NSW, 2000, AU

(56) Related Art
C. K. DAMGAARD ET AL, "Translational coregulation of 5'TOP mRNAs by TIA-1 and TIAR", GENES & DEVELOPMENT, (2011-10-01), vol. 25, no. 19, doi:10.1101/gad.17355911, ISSN 0890-9369, pages 2057 - 2068
S. LEVY ET AL, PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, (1991-04-15), vol. 88, no. 8, doi:10.1073/pnas.88.8.3319, ISSN 0027-8424, pages 3319 - 3323
ULF ANDERSSON ØROM ET AL, "MicroRNA-10a Binds the 5'UTR of Ribosomal Protein mRNAs and Enhances Their Translation", MOLECULAR CELL, (2008-05-01), vol. 30, no. 4, doi:10.1016/j.molcel.2008.05.001, ISSN 1097-2765, pages 460 - 471
LEDDA M ET AL, "Effect of 3'UTR length on the translational regulation of 5'-terminal oligopyrimidine mRNAs", GENE, ELSEVIER, AMSTERDAM, NL, (2005-01-03), vol. 344, doi:10.1016/J.GENE.2004.09.023, ISSN 0378-1119, pages 213 - 220



(51) International Patent Classification:
C12N 15/67 (2006.01)

(21) International Application Number:
PCT/EP2013/000938

(22) International Filing Date:
27 March 2013 (27.03.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/EP2012/001334 27 March 2012 (27.03.2012) EP
PCT/EP2012/002448 8 June 2012 (08.06.2012) EP

(71) Applicant: CUREVAC GMBH [DE/DE]; Paul-Ehrlich-Str. 15, 72076 Tübingen (DE).

(72) Inventor; and

(71) Applicant (for US only): THEB, Andreas [DE/DE]; Hindenburgstr. 9, 72127 Kusterdingen (DE).

(74) Agents: GRAF VON STOSCH, Andreas et al.; Graf von Stosch Patentanwalts-gesellschaft mbH, Prinzregentenstrasse 22, 80538 München (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:
27 December 2013

(54) Title: ARTIFICIAL NUCLEIC ACID MOLECULES COMPRISING A 5'TOP UTR

(57) Abstract: The invention relates to an artificial nucleic acid molecule comprising at least one 5'UTR element which is derived from a TOP gene, at least one open reading frame and optionally at least one 3'UTR element comprising a nucleic acid sequence which is preferably derived from the 3'UTR of a gene providing a stable mRNA, such as an albumin gene, or from a variant of the 3'UTR of a gene providing a stable mRNA. The invention further relates to the use of such an artificial nucleic acid molecule in gene therapy and/or genetic vaccination.



5 Artificial nucleic acid molecules comprising a 5'TOP UTR

The invention relates to artificial nucleic acid molecules comprising a 5'UTR element derived from the 5'UTR of a TOP gene, an open reading frame, and optionally a 3'UTR element, a poly(A) sequence and/or a polyadenylation signal. The invention relates further
10 to a vector comprising a 5'UTR element derived from the 5'UTR of a TOP gene, to a pharmaceutical composition comprising the artificial nucleic acid molecule or the vector, and to a kit comprising the artificial nucleic acid molecule, the vector and/or the pharmaceutical composition, preferably for use in the field of gene therapy and/or genetic vaccination.

15 Gene therapy and genetic vaccination belong to the most promising and quickly developing methods of modern medicine. They may provide highly specific and individual options for therapy of a large variety of diseases. Particularly, inherited genetic diseases but also autoimmune diseases, cancerous or tumour-related diseases as well as inflammatory diseases may be the subject of such treatment approaches. Also, it is envisaged to prevent
20 (early) onset of such diseases by these approaches.

The main conceptual rational behind gene therapy is appropriate modulation of impaired gene expression associated with pathological conditions of specific diseases. Pathologically altered gene expression may result in lack or overproduction of essential gene products, for example, signalling factors such as hormones, housekeeping factors, metabolic enzymes,
25 structural proteins or the like. Altered gene expression may not only be due to mis-regulation of transcription and/or translation, but also due to mutations within the ORF coding for a particular protein. Pathological mutations may be caused by e.g. chromosomal aberration, or by more specific mutations, such as point or frame-shift-mutations, all of them

resulting in limited functionality and, potentially, total loss of function of the gene product. However, misregulation of transcription or translation may also occur, if mutations affect genes encoding proteins which are involved in the transcriptional or translational machinery of the cell. Such mutations may lead to pathological up- or down-regulation of genes which are – as such – functional. Genes encoding gene products which exert such regulating functions, may be, e.g., transcription factors, signal receptors, messenger proteins or the like. However, loss of function of such genes encoding regulatory proteins may, under certain circumstances, be reversed by artificial introduction of other factors acting further downstream of the impaired gene product. Such gene defects may also be compensated by gene therapy via substitution of the affected gene itself.

Genetic vaccination allows evoking a desired immune response to selected antigens, such as characteristic components of bacterial surfaces, viral particles, tumour antigens or the like. Generally, vaccination is one of the pivotal achievements of modern medicine. However, effective vaccines are currently available only for a smaller number of diseases. Accordingly, infections that are not preventable by vaccination still affect millions of people every year.

Commonly, vaccines may be subdivided into “first”, “second” and “third” generation vaccines. “First generation” vaccines are, typically, whole-organism vaccines. They are based on either live and attenuated or killed pathogens, e.g. viruses, bacteria or the like. The major drawback of live and attenuated vaccines is the risk for a reversion to life-threatening variants. Thus, although attenuated, such pathogens may still intrinsically bear unpredictable risks. Killed pathogens may not be as effective as desired for generating a specific immune response. In order to minimize these risks, “second generation” vaccines were developed. These are, typically, subunit vaccines, consisting of defined antigens or recombinant protein components which are derived from pathogens.

Genetic vaccines, i.e. vaccines for genetic vaccination, are usually understood as “third generation” vaccines. They are typically composed of genetically engineered nucleic acid molecules which allow expression of peptide or protein (antigen) fragments characteristic for a pathogen or a tumor antigen *in vivo*. Genetic vaccines are expressed upon administration to a patient and uptake by competent cells. Expression of the administered

nucleic acids results in production of the encoded proteins. In the event these proteins are recognized as foreign by the patient's immune system, an immune response is triggered.

As can be seen from the above, both methods, gene therapy and genetic vaccination, are essentially based on the administration of nucleic acid molecules to a patient and subsequent transcription and/or translation of the encoded genetic information. Alternatively, genetic vaccination or gene therapy may also comprise methods which include isolation of specific body cells from a patient to be treated, subsequent *in vitro* transfection of such cells, and re-administration of the treated cells to the patient.

DNA as well as RNA may be used as nucleic acid molecules for administration in the context of gene therapy or genetic vaccination. DNA is known to be relatively stable and easy to handle. However, the use of DNA bears the risk of undesired insertion of the administered DNA-fragments into the patient's genome potentially resulting in loss of function of the impaired genes. As a further risk, the undesired generation of anti-DNA antibodies has emerged. Another drawback is the limited expression level of the encoded peptide or protein that is achievable upon DNA administration and its transcription/translation. Among other reasons, the expression level of the administered DNA will be dependent on the presence of specific transcription factors which regulate DNA transcription. In the absence of such factors, DNA transcription will not yield satisfying amounts of RNA. As a result, the level of translated peptide or protein obtained is limited.

By using RNA instead of DNA for gene therapy or genetic vaccination, the risk of undesired genomic integration and generation of anti-DNA antibodies is minimized or avoided. However, RNA is considered to be a rather unstable molecular species which may readily be degraded by ubiquitous RNAses.

In vivo, RNA-degradation contributes to the regulation of the RNA half-life time. That effect was considered and proven to fine tune the regulation of eukaryotic gene expression (Friedel *et al.*, Conserved principles of mammalian transcriptional regulation revealed by RNA half-life, Nucleic Acid Research, 2009, 1-12). Accordingly, each naturally occurring mRNA has its individual half-life depending on the gene from which the mRNA is derived. It contributes to the regulation of the expression level of this gene. Unstable RNAs are important to realize transient gene expression at distinct points in time. However, long-lived

5 RNAs may be associated with accumulation of distinct proteins or continuous expression of genes. *In vivo*, the half life of mRNAs may also be dependent on environmental factors, such as hormonal treatment, as has been shown, e.g., for insulin-like growth factor I, actin, and albumin mRNA (Johnson *et al.*, Newly synthesized RNA: Simultaneous measurement in intact cells of transcription rates and RNA stability of insulin-like growth factor I, actin, and albumin in growth hormone-stimulated hepatocytes, Proc. Natl. Acad. Sci., Vol. 88, pp. 5287-5291, 1991).

10 For gene therapy and genetic vaccination, usually stable RNA is desired. This is, on the one hand, due to the fact that the product encoded by the RNA-sequence shall accumulate *in vivo*. On the other hand, the RNA has to maintain its structural and functional integrity when prepared for a suitable dosage form, in the course of its storage, and when administered. Thus, considerable attention was dedicated to provide stable RNA molecules for gene therapy or genetic vaccination in order to prevent them from being subject to early degradation or decay.

15 It has been reported that the G/C-content of nucleic acid molecules may influence their stability. Thus, nucleic acids comprising an increased amount of guanine (G) and/or cytosine (C) residues may be functionally more stable than nucleic acids containing a large amount of adenine (A) and thymine (T) or uracil (U) nucleotides. In this context, WO02/098443 provides a pharmaceutical composition containing an mRNA that is
20 stabilised by sequence modifications in the translated region. Such a sequence modification takes advantage of the degeneracy of the genetic code. Accordingly, codons which contain a less favourable combination of nucleotides (less favourable in terms of RNA stability) may be substituted by alternative codons without altering the encoded amino acid sequence. This method of RNA stabilization is limited by the provisions of the specific nucleotide
25 sequence of each single RNA molecule which is not allowed to leave the space of the desired amino acid sequence. Also, that approach is restricted to coding regions of the RNA.

30 As an alternative option for mRNA stabilisation, it has been found that naturally occurring eukaryotic mRNA molecules contain characteristic stabilising elements. For example, they may comprise so-called untranslated regions (UTR) at their 5'-end (5'UTR) and/or at their 3'-end (3'UTR) as well as other structural features, such as a 5'-cap structure or a 3'-

poly(A) tail. Both, 5'UTR and 3'UTR are typically transcribed from the genomic DNA and are, thus, an element of the premature mRNA. Characteristic structural features of mature mRNA, such as the 5'-cap and the 3'-poly(A) tail (also called poly(A) tail or poly(A) sequence) are usually added to the transcribed (premature) mRNA during mRNA processing.

A 3'-poly(A) tail is typically a monotonous sequence stretch of adenine nucleotides added to the 3'-end of the transcribed mRNA. It may comprise up to about 400 adenine nucleotides. It was found that the length of such a 3'-poly(A) tail is a potentially critical element for the stability of the individual mRNA.

Also, it was shown that the 3'UTR of α -globin mRNA may be an important factor for the well-known stability of α -globin mRNA (Rodgers *et al.*, Regulated α -globin mRNA decay is a cytoplasmic event proceeding through 3'-to-5' exosome-dependent decapping, RNA, 8, pp. 1526-1537, 2002). The 3'UTR of α -globin mRNA is obviously involved in the formation of a specific ribonucleoprotein-complex, the α -complex, whose presence correlates with mRNA stability *in vitro* (Wang *et al.*, An mRNA stability complex functions with poly(A)-binding protein to stabilize mRNA *in vitro*, Molecular and Cellular biology, Vol 19, No. 7, July 1999, p. 4552-4560).

Irrespective of factors influencing mRNA stability, effective translation of the administered nucleic acid molecules by the target cells or tissue is crucial for any approach using nucleic acid molecules for gene therapy or genetic vaccination. Along with the regulation of stability, also translation of the majority of mRNAs is regulated by structural features like UTRs, 5'-cap and 3'-poly(A) tail. In this context, it has been reported that the length of the poly(A) tail may play an important role for translational efficiency as well. Stabilizing 3'-elements, however, may also have an attenuating effect on translation.

Further regulative elements, which may have an influence on expression levels, may be found in the 5'UTR. For example, it has been reported that synthesis of particular proteins, e.g. proteins belonging to the translational apparatus, may be regulated not only at the transcriptional but also at the translational level. For example, translation of proteins encoded by so called 'TOP-genes' may be down-regulated by translational repression. Therein, the term 'TOP-gene' relates to a gene corresponding to an mRNA that is characterized by the presence of a TOP sequence at the 5'end and in most cases by a

growth-associated translation regulation (Iadevaia *et al.*, All translation elongation factors and the e, f, and h subunits of translation initiation factor 3 are encoded by 5'-terminal oligopyrimidine (TOP) mRNAs; RNA, 2008, 14:1730-1736). In this context, a TOP sequence – also called the '5'-terminal oligopyrimidine tract' – typically consists of a C
5 residue at the cap site, followed by an uninterrupted sequence of up to 13 or even more pyrimidines (Avni *et al.*, Vertebrate mRNAs with a 5'-terminal pyrimidine tract are Candidates for translational repression in quiescent cells: characterization of the translational cis-regulatory element, Molecular and Cellular Biology, 1994, p. 3822-3833). These TOP sequences are reported to be present in many mRNAs encoding components of
10 the translational machinery and to be responsible for selective repression of the translation of these TOP containing mRNAs due to growth arrest (Meyuhas, *et al.*, Translational Control of Ribosomal Protein mRNAs in Eukaryotes, Translational Control. Cold Spring Harbor Monograph Archive. Cold Spring Harbor Laboratory Press, 1996, p. 363–388). These TOP sequences are thought to serve as a cis-regulatory element which inhibits the binding of
15 translational regulatory proteins or the translational machinery itself. As a result, the translation of these genes is inhibited at the growth arrest of cells. More specifically, when a cell is faced with starvation or treated by some chemicals such as 12-Otetradecanoyl-1-phorbol-13-acetate (TPA), mRNAs of TOP genes, which are normally associated with polysomes, change their status into the translationally inactive 'sub-polysome' while most
20 non-TOP mRNAs stay in the 'polysome' state (Yamashita *et al.*, Comprehensive detection of human terminal oligo-pyrimidine (TOP) genes and analysis of their characteristics. Nucleic Acids Res. 2008 Jun;36(11):3707-15. doi: 10.1093/nar/gkn248. Epub 2008 May 14). In this context, it was shown that the oligopyrimidine tract at the 5' end of the 5'UTR (TOP motif) was required for translational repression of TOP genes. The oligopyrimidine tract at the 5'
25 end of mammalian ribosomal protein mRNAs is required for their translational control (Levy *et al.*, Proc Natl Acad Sci U S A. 1991 Apr 15;88(8):3319-23). Furthermore, it was shown that miRNA miR-10a positively controls the translation of ribosomal proteins by binding downstream of the TOP motif present in the 5'UTRs of TOP genes. Such an enhancement of translation was dependent on the presence of the TOP motif in the 5'UTR. Furthermore this
30 translational regulation of ribosomal TOP genes was dependent on the presence of miR-10a or its human homolog miR-10b which is highly overexpressed in several tumor types and is reportedly involved in the progression of cancer (Ørom *et al.*, MicroRNA-10a binds the

5'UTR of ribosomal protein mRNAs and enhances their translation. Mol Cell. 2008 May 23;30(4):460-71).

There is provided herein nucleic acid molecules which may be suitable for application in gene therapy and/or genetic vaccination. Particularly, there is provided an mRNA species which is
5 stabilized against preterm degradation or decay without exhibiting significant functional loss in translational efficiency. Also provided are nucleic acid molecules coding for such a superior mRNA species which may be amenable for use in gene therapy and/or genetic vaccination. Further provided is a pharmaceutical composition for use in gene therapy and/or genetic vaccination. In summary, the present invention provides improved nucleic acid species which
10 overcome the above discussed disadvantages of the prior art by a cost-effective and straightforward approach.

For the sake of clarity and readability the following definitions are provided. Any technical feature mentioned for these definitions may be read on each and every embodiment of the invention. Additional definitions and explanations may be specifically provided in the context of
15 these embodiments.

Adaptive immune response: The adaptive immune response is typically understood to be an antigen-specific response of the immune system. Antigen specificity allows for the generation of responses that are tailored to specific pathogens or pathogen-infected cells. The ability to mount these tailored responses is usually maintained in the body by "memory cells". Should a
20 pathogen infect the body more than once, these specific memory cells are used to quickly eliminate it. In this context, the first step of an adaptive immune response is the activation of naïve antigen-specific T cells or different immune cells able to induce an antigen-specific immune response by antigen-presenting cells. This occurs in the lymphoid tissues and organs through which naïve T cells are constantly passing. The three cell types that may serve as antigen-
25 presenting cells are dendritic cells, macrophages, and B cells. Each of these cells has a distinct function in eliciting immune responses. Dendritic cells may

take up antigens by phagocytosis and macropinocytosis and may become stimulated by contact with e.g. a foreign antigen to migrate to the local lymphoid tissue, where they differentiate into mature dendritic cells. Macrophages ingest particulate antigens such as bacteria and are induced by infectious agents or other appropriate stimuli to express MHC molecules. The unique ability of B cells to bind and internalize soluble protein antigens via their receptors may also be important to induce T cells. MHC-molecules are, typically, responsible for presentation of an antigen to T-cells. Therein, presenting the antigen on MHC molecules leads to activation of T cells which induces their proliferation and differentiation into armed effector T cells. The most important function of effector T cells is the killing of infected cells by CD8+ cytotoxic T cells and the activation of macrophages by Th1 cells which together make up cell-mediated immunity, and the activation of B cells by both Th2 and Th1 cells to produce different classes of antibody, thus driving the humoral immune response. T cells recognize an antigen by their T cell receptors which do not recognize and bind the antigen directly, but instead recognize short peptide fragments e.g. of pathogen-derived protein antigens, e.g. so-called epitopes, which are bound to MHC molecules on the surfaces of other cells.

Adaptive immune system: The adaptive immune system is essentially dedicated to eliminate or prevent pathogenic growth. It typically regulates the adaptive immune response by providing the vertebrate immune system with the ability to recognize and remember specific pathogens (to generate immunity), and to mount stronger attacks each time the pathogen is encountered. The system is highly adaptable because of somatic hypermutation (a process of accelerated somatic mutations), and V(D)J recombination (an irreversible genetic recombination of antigen receptor gene segments). This mechanism allows a small number of genes to generate a vast number of different antigen receptors, which are then uniquely expressed on each individual lymphocyte. Because the gene rearrangement leads to an irreversible change in the DNA of each cell, all of the progeny (offspring) of such a cell will then inherit genes encoding the same receptor specificity, including the Memory B cells and Memory T cells that are the keys to long-lived specific immunity.

Adjuvant/adjuvant component: An adjuvant or an adjuvant component in the broadest sense is typically a pharmacological and/or immunological agent that may modify, e.g. enhance, the effect of other agents, such as a drug or vaccine. It is to be interpreted in a broad sense and refers to a broad spectrum of substances. Typically, these substances are

able to increase the immunogenicity of antigens. For example, adjuvants may be recognized by the innate immune systems and, e.g., may elicit an innate immune response. "Adjuvants" typically do not elicit an adaptive immune response. Insofar, "adjuvants" do not qualify as antigens. Their mode of action is distinct from the effects triggered by antigens resulting in an adaptive immune response.

Antigen: In the context of the present invention "antigen" refers typically to a substance which may be recognized by the immune system, preferably by the adaptive immune system, and is capable of triggering an antigen-specific immune response, e.g. by formation of antibodies and/or antigen-specific T cells as part of an adaptive immune response. Typically, an antigen may be or may comprise a peptide or protein which may be presented by the MHC to T-cells.

Artificial nucleic acid molecule: An artificial nucleic acid molecule may typically be understood to be a nucleic acid molecule, e.g. a DNA or an RNA, that does not occur naturally. In other words, an artificial nucleic acid molecule may be understood as a non-natural nucleic acid molecule. Such nucleic acid molecule may be non-natural due to its individual sequence (which does not occur naturally) and/or due to other modifications, e.g. structural modifications of nucleotides which do not occur naturally. An artificial nucleic acid molecule may be a DNA molecule, an RNA molecule or a hybrid-molecule comprising DNA and RNA portions. Typically, artificial nucleic acid molecules may be designed and/or generated by genetic engineering methods to correspond to a desired artificial sequence of nucleotides (heterologous sequence). In this context an artificial sequence is usually a sequence that may not occur naturally, i.e. it differs from the wild type sequence by at least one nucleotide. The term 'wild type' may be understood as a sequence occurring in nature. Further, the term 'artificial nucleic acid molecule' is not restricted to mean 'one single molecule' but is, typically, understood to comprise an ensemble of identical molecules. Accordingly, it may relate to a plurality of identical molecules contained in an aliquot.

Bicistronic RNA, multicistronic RNA: A bicistronic or multicistronic RNA is typically an RNA, preferably an mRNA, that typically may have two (bicistronic) or more (multicistronic) open reading frames (ORF). An open reading frame in this context is a sequence of codons that is translatable into a peptide or protein.

Carrier / polymeric carrier: A carrier in the context of the invention may typically be a compound that facilitates transport and/or complexation of another compound (cargo). A polymeric carrier is typically a carrier that is formed of a polymer. A carrier may be associated to its cargo by covalent or non-covalent interaction. A carrier may transport
5 nucleic acids, e.g. RNA or DNA, to the target cells. The carrier may – for some embodiments – be a cationic component.

Cationic component: The term “cationic component” typically refers to a charged molecule, which is positively charged (cation) at a pH value typically from 1 to 9, preferably at a pH value of or below 9 (e.g. from 5 to 9), of or below 8 (e.g. from 5 to 8), of
10 or below 7 (e.g. from 5 to 7), most preferably at a physiological pH, e.g. from 7.3 to 7.4. Accordingly, a cationic component may be any positively charged compound or polymer, preferably a cationic peptide or protein which is positively charged under physiological conditions, particularly under physiological conditions *in vivo*. A ‘cationic peptide or protein’ may contain at least one positively charged amino acid, or more than one
15 positively charged amino acid, e.g. selected from Arg, His, Lys or Orn. Accordingly, ‘polycationic’ components are also within the scope exhibiting more than one positive charge under the conditions given.

5'-cap: A 5'-cap is an entity, typically a modified nucleotide entity, which generally ‘caps’ the 5'-end of a mature mRNA. A 5'-cap may typically be formed by a modified
20 nucleotide, particularly by a derivative of a guanine nucleotide. Preferably, the 5'-cap is linked to the 5'-terminus via a 5'-5'-triphosphate linkage. A 5'-cap may be methylated, e.g. m7GpppN, wherein N is the terminal 5' nucleotide of the nucleic acid carrying the 5'-cap, typically the 5'-end of an RNA. Further examples of 5'cap structures include glyceryl, inverted deoxy abasic residue (moiety), 4',5' methylene nucleotide, 1-(beta-D-
25 erythrofuranosyl) nucleotide, 4'-thio nucleotide, carbocyclic nucleotide, 1,5-anhydrohexitol nucleotide, L-nucleotides, alpha-nucleotide, modified base nucleotide, threo-pentofuranosyl nucleotide, acyclic 3',4'-seco nucleotide, acyclic 3,4-dihydroxybutyl nucleotide, acyclic 3,5 dihydroxypentyl nucleotide, 3'-3'-inverted nucleotide moiety, 3'-3'-inverted abasic moiety, 3'-2'-inverted nucleotide moiety, 3'-2'-inverted abasic moiety, 1,4-butanediol
30 phosphate, 3'-phosphoramidate, hexylphosphate, aminohexyl phosphate, 3'-phosphate, 3'phosphorothioate, phosphorodithioate, or bridging or non-bridging methylphosphonate moiety.

Cellular immunity/cellular immune response: Cellular immunity relates typically to the activation of macrophages, natural killer cells (NK), antigen-specific cytotoxic T-lymphocytes, and the release of various cytokines in response to an antigen. In more general terms, cellular immunity is not based on antibodies, but on the activation of cells of the immune system. Typically, a cellular immune response may be characterized e.g. by activating antigen-specific cytotoxic T-lymphocytes that are able to induce apoptosis in cells, e.g. specific immune cells like dendritic cells or other cells, displaying epitopes of foreign antigens on their surface. Such cells may be virus-infected or infected with intracellular bacteria, or cancer cells displaying tumor antigens. Further characteristics may be activation of macrophages and natural killer cells, enabling them to destroy pathogens and stimulation of cells to secrete a variety of cytokines that influence the function of other cells involved in adaptive immune responses and innate immune responses.

DNA: DNA is the usual abbreviation for deoxy-ribonucleic-acid. It is a nucleic acid molecule, i.e. a polymer consisting of nucleotides. These nucleotides are usually deoxy-adenosine-monophosphate, deoxy-thymidine-monophosphate, deoxy-guanosine-monophosphate and deoxy-cytidine-monophosphate monomers which are – by themselves – composed of a sugar moiety (deoxyribose), a base moiety and a phosphate moiety, and polymerize by a characteristic backbone structure. The backbone structure is, typically, formed by phosphodiester bonds between the sugar moiety of the nucleotide, i.e. deoxyribose, of a first and a phosphate moiety of a second, adjacent monomer. The specific order of the monomers, i.e. the order of the bases linked to the sugar/phosphate-backbone, is called the DNA-sequence. DNA may be single stranded or double stranded. In the double stranded form, the nucleotides of the first strand typically hybridize with the nucleotides of the second strand, e.g. by A/T-base-pairing and G/C-base-pairing.

Epitope: Epitopes (also called 'antigen determinant') can be distinguished in T cell epitopes and B cell epitopes. T cell epitopes or parts of the proteins in the context of the present invention may comprise fragments preferably having a length of about 6 to about 20 or even more amino acids, e.g. fragments as processed and presented by MHC class I molecules, preferably having a length of about 8 to about 10 amino acids, e.g. 8, 9, or 10, (or even 11, or 12 amino acids), or fragments as processed and presented by MHC class II molecules, preferably having a length of about 13 or more amino acids, e.g. 13, 14, 15, 16, 17, 18, 19, 20 or even more amino acids, wherein these fragments may be selected from

any part of the amino acid sequence. These fragments are typically recognized by T cells in form of a complex consisting of the peptide fragment and an MHC molecule, i.e. the fragments are typically not recognized in their native form. B cell epitopes are typically fragments located on the outer surface of (native) protein or peptide antigens as defined herein, preferably having 5 to 15 amino acids, more preferably having 5 to 12 amino acids, even more preferably having 6 to 9 amino acids, which may be recognized by antibodies, i.e. in their native form.

Such epitopes of proteins or peptides may furthermore be selected from any of the herein mentioned variants of such proteins or peptides. In this context antigenic determinants can be conformational or discontinuous epitopes which are composed of segments of the proteins or peptides as defined herein that are discontinuous in the amino acid sequence of the proteins or peptides as defined herein but are brought together in the three-dimensional structure or continuous or linear epitopes which are composed of a single polypeptide chain.

Fragment of a sequence: A fragment of a sequence may typically be a shorter portion of a full-length sequence of e.g. a nucleic acid molecule or an amino acid sequence. Accordingly, a fragment, typically, consists of a sequence that is identical to the corresponding stretch within the full-length sequence. A preferred fragment of a sequence in the context of the present invention, consists of a continuous stretch of entities, such as nucleotides or amino acids corresponding to a continuous stretch of entities in the molecule the fragment is derived from, which represents at least 20%, preferably at least 30%, more preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, even more preferably at least 70%, and most preferably at least 80% of the total (i.e. full-length) molecule from which the fragment is derived.

G/C modified: A G/C-modified nucleic acid may typically be a nucleic acid, preferably an artificial nucleic acid molecule as defined herein, based on a modified wild-type sequence comprising a preferably increased number of guanosine and/or cytosine nucleotides as compared to the wild-type sequence. Such an increased number may be generated by substitution of codons containing adenosine or thymidine nucleotides by codons containing guanosine or cytosine nucleotides. If the enriched G/C content occurs in a coding region of DNA or RNA, it makes use of the degeneracy of the genetic code.

Accordingly, the codon substitutions preferably do not alter the encoded amino acid residues, but exclusively increase the G/C content of the nucleic acid molecule.

Gene therapy: Gene therapy may typically be understood to mean a treatment of a patient's body or isolated elements of a patient's body, for example isolated tissues/cells, by nucleic acids encoding a peptide or protein. It typically may comprise at least one of the steps of a) administration of a nucleic acid, preferably an artificial nucleic acid molecule as defined herein, directly to the patient - by whatever administration route - or *in vitro* to isolated cells/tissues of the patient, which results in transfection of the patient's cells either *in vivo/ex vivo* or *in vitro*; b) transcription and/or translation of the introduced nucleic acid molecule; and optionally c) re-administration of isolated, transfected cells to the patient, if the nucleic acid has not been administered directly to the patient.

Genetic vaccination: Genetic vaccination may typically be understood to be vaccination by administration of a nucleic acid molecule encoding an antigen or an immunogen or fragments thereof. The nucleic acid molecule may be administered to a subject's body or to isolated cells of a subject. Upon transfection of certain cells of the body or upon transfection of the isolated cells, the antigen or immunogen may be expressed by those cells and subsequently presented to the immune system, eliciting an adaptive, i.e. antigen-specific immune response. Accordingly, genetic vaccination typically comprises at least one of the steps of a) administration of a nucleic acid, preferably an artificial nucleic acid molecule as defined herein, to a subject, preferably a patient, or to isolated cells of a subject, preferably a patient, which usually results in transfection of the subject's cells either *in vivo* or *in vitro*; b) transcription and/or translation of the introduced nucleic acid molecule; and optionally c) re-administration of isolated, transfected cells to the subject, preferably the patient, if the nucleic acid has not been administered directly to the patient.

Heterologous sequence: Two sequences are typically understood to be 'heterologous' if they are not derivable from the same gene. I.e., although heterologous sequences may be derivable from the same organism, they naturally (in nature) do not occur in the same nucleic acid molecule, such as in the same mRNA.

Humoral immunity/humoral immune response: Humoral immunity refers typically to antibody production and optionally to accessory processes accompanying antibody production. A humoral immune response may be typically characterized, e.g., by Th2

activation and cytokine production, germinal center formation and isotype switching, affinity maturation and memory cell generation. Humoral immunity also typically may refer to the effector functions of antibodies, which include pathogen and toxin neutralization, classical complement activation, and opsonin promotion of phagocytosis and pathogen
5 elimination.

Immunogen: In the context of the present invention an immunogen may be typically understood to be a compound that is able to stimulate an immune response. Preferably, an immunogen is a peptide, polypeptide, or protein. In a particularly preferred embodiment, an immunogen in the sense of the present invention is the product of translation of a provided
10 nucleic acid molecule, preferably an artificial nucleic acid molecule as defined herein. Typically, an immunogen elicits at least an adaptive immune response.

Immunostimulatory composition: In the context of the invention, an immunostimulatory composition may be typically understood to be a composition containing at least one component which is able to induce an immune response or from which a component
15 which is able to induce an immune response is derivable. Such immune response may be preferably an innate immune response or a combination of an adaptive and an innate immune response. Preferably, an immunostimulatory composition in the context of the invention contains at least one artificial nucleic acid molecule, more preferably an RNA, for example an mRNA molecule. The immunostimulatory component, such as the mRNA may
20 be complexed with a suitable carrier. Thus, the immunostimulatory composition may comprise an mRNA/carrier-complex. Furthermore, the immunostimulatory composition may comprise an adjuvant and/or a suitable vehicle for the immunostimulatory component, such as the mRNA.

Immune response: An immune response may typically be a specific reaction of the adaptive
25 immune system to a particular antigen (so called specific or adaptive immune response) or an unspecific reaction of the innate immune system (so called unspecific or innate immune response), or a combination thereof.

Immune system: The immune system may protect organisms from infection. If a pathogen succeeds in passing a physical barrier of an organism and enters this organism, the innate
30 immune system provides an immediate, but non-specific response. If pathogens evade this innate response, vertebrates possess a second layer of protection, the adaptive immune

system. Here, the immune system adapts its response during an infection to improve its recognition of the pathogen. This improved response is then retained after the pathogen has been eliminated, in the form of an immunological memory, and allows the adaptive immune system to mount faster and stronger attacks each time this pathogen is encountered. According to this, the immune system comprises the innate and the adaptive immune system. Each of these two parts typically contains so called humoral and cellular components.

Immunostimulatory RNA: An immunostimulatory RNA (isRNA) in the context of the invention may typically be an RNA that is able to induce an innate immune response. It usually does not have an open reading frame and thus does not provide a peptide-antigen or immunogen but elicits an immune response e.g. by binding to a specific kind of Toll-like-receptor (TLR) or other suitable receptors. However, of course also mRNAs having an open reading frame and coding for a peptide/protein may induce an innate immune response and, thus, may be immunostimulatory RNAs.

Innate immune system: The innate immune system, also known as non-specific (or unspecific) immune system, typically comprises the cells and mechanisms that defend the host from infection by other organisms in a non-specific manner. This means that the cells of the innate-system may recognize and respond to pathogens in a generic way, but unlike the adaptive immune system, it does not confer long-lasting or protective immunity to the host. The innate immune system may be, e.g., activated by ligands of Toll-like receptors (TLRs) or other auxiliary substances such as lipopolysaccharides, TNF-alpha, CD40 ligand, or cytokines, monokines, lymphokines, interleukins or chemokines, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-21, IL-22, IL-23, IL-24, IL-25, IL-26, IL-27, IL-28, IL-29, IL-30, IL-31, IL-32, IL-33, IFN-alpha, IFN-beta, IFN-gamma, GM-CSF, G-CSF, M-CSF, LT-beta, TNF-alpha, growth factors, and hGH, a ligand of human Toll-like receptor TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, a ligand of murine Toll-like receptor TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, TLR11, TLR12 or TLR13, a ligand of a NOD-like receptor, a ligand of a RIG-I like receptor, an immunostimulatory nucleic acid, an immunostimulatory RNA (isRNA), a CpG-DNA, an antibacterial agent, or an anti-viral agent. The pharmaceutical composition according to the present invention may comprise one or more such substances. Typically, a response of the innate immune system includes recruiting

immune cells to sites of infection, through the production of chemical factors, including specialized chemical mediators, called cytokines; activation of the complement cascade; identification and removal of foreign substances present in organs, tissues, the blood and lymph, by specialized white blood cells; activation of the adaptive immune system; and/or
5 acting as a physical and chemical barrier to infectious agents.

Cloning site: A cloning site is typically understood to be a segment of a nucleic acid molecule, which is suitable for insertion of a nucleic acid sequence, e.g., a nucleic acid sequence comprising an open reading frame. Insertion may be performed by any molecular biological method known to the one skilled in the art, e.g. by restriction and ligation. A
10 cloning site typically comprises one or more restriction enzyme recognition sites (restriction sites). These one or more restrictions sites may be recognized by restriction enzymes which cleave the DNA at these sites. A cloning site which comprises more than one restriction site may also be termed a multiple cloning site (MCS) or a polylinker.

Nucleic acid molecule: A nucleic acid molecule is a molecule comprising, preferably
15 consisting of nucleic acid components. The term nucleic acid molecule preferably refers to DNA or RNA molecules. It is preferably used synonymous with the term "polynucleotide". Preferably, a nucleic acid molecule is a polymer comprising or consisting of nucleotide monomers which are covalently linked to each other by phosphodiester-bonds of a sugar/phosphate-backbone. The term "nucleic acid molecule" also encompasses modified
20 nucleic acid molecules, such as base-modified, sugar-modified or backbone-modified etc. DNA or RNA molecules.

Open reading frame: An open reading frame (ORF) in the context of the invention may typically be a sequence of several nucleotide triplets which may be translated into a peptide or protein. An open reading frame preferably contains a start codon, i.e. a combination of
25 three subsequent nucleotides coding usually for the amino acid methionine (ATG or AUG), at its 5'-end and a subsequent region which usually exhibits a length which is a multiple of 3 nucleotides. An ORF is preferably terminated by a stop-codon (e.g., TAA, TAG, TGA). Typically, this is the only stop-codon of the open reading frame. Thus, an open reading frame in the context of the present invention is preferably a nucleotide sequence, consisting
30 of a number of nucleotides that may be divided by three, which starts with a start codon (e.g. ATG or AUG) and which preferably terminates with a stop codon (e.g., TAA, TGA, or

TAG or UAA, UAG, UGA, respectively). The open reading frame may be isolated or it may be incorporated in a longer nucleic acid sequence, for example in a vector or an mRNA. An open reading frame may also be termed 'protein coding region'.

5 Peptide: A peptide or polypeptide is typically a polymer of amino acid monomers, linked by peptide bonds. It typically contains less than 50 monomer units. Nevertheless, the term peptide is not a disclaimer for molecules having more than 50 monomer units. Long peptides are also called polypeptides, typically having between 50 and 600 monomeric units.

10 Pharmaceutically effective amount: A pharmaceutically effective amount in the context of the invention is typically understood to be an amount that is sufficient to induce a pharmaceutical effect, such as an immune response, altering a pathological level of an expressed peptide or protein, or substituting a lacking gene product, e.g., in case of a pathological situation.

15 Protein A protein typically comprises one or more peptides or polypeptides. A protein is typically folded into 3-dimensional form, which may be required for to protein to exert its biological function.

20 Poly(A) sequence: A poly(A) sequence, also called poly(A) tail or 3'-poly(A) tail, is typically understood to be a sequence of adenine nucleotides, e.g., of up to about 400 adenine nucleotides, e.g. from about 20 to about 400, preferably from about 50 to about 400, more preferably from about 50 to about 300, even more preferably from about 50 to about 250, most preferably from about 60 to about 250 adenine nucleotides. A poly(A) sequence is typically located at the 3'end of an mRNA. In the context of the present invention, a poly(A) sequence may be located within an mRNA or any other nucleic acid molecule, such as, e.g., in a vector, for example, in a vector serving as template for the generation of an RNA, 25 preferably an mRNA, e.g., by transcription of the vector.

Polyadenylation: Polyadenylation is typically understood to be the addition of a poly(A) sequence to a nucleic acid molecule, such as an RNA molecule, e.g. to a premature mRNA. Polyadenylation may be induced by a so called polyadenylation signal. This signal is preferably located within a stretch of nucleotides at the 3'-end of a nucleic acid molecule, 30 such as an RNA molecule, to be polyadenylated. A polyadenylation signal typically

comprises a hexamer consisting of adenine and uracil/thymine nucleotides, preferably the hexamer sequence AAUAAA. Other sequences, preferably hexamer sequences, are also conceivable. Polyadenylation typically occurs during processing of a pre-mRNA (also called premature-mRNA). Typically, RNA maturation (from pre-mRNA to mature mRNA) comprises the step of polyadenylation.

Restriction site: A restriction site, also termed 'restriction enzyme recognition site', is a nucleotide sequence recognized by a restriction enzyme. A restriction site is typically a short, preferably palindromic nucleotide sequence, e.g. a sequence comprising 4 to 8 nucleotides. A restriction site is preferably specifically recognized by a restriction enzyme.

The restriction enzyme typically cleaves a nucleotide sequence comprising a restriction site at this site. In a double-stranded nucleotide sequence, such as a double-stranded DNA sequence, the restriction enzyme typically cuts both strands of the nucleotide sequence.

RNA, mRNA: RNA is the usual abbreviation for ribonucleic-acid. It is a nucleic acid molecule, i.e. a polymer consisting of nucleotides. These nucleotides are usually adenosine-monophosphate, uridine-monophosphate, guanosine-monophosphate and cytidine-monophosphate monomers which are connected to each other along a so-called backbone. The backbone is formed by phosphodiester bonds between the sugar, i.e. ribose, of a first and a phosphate moiety of a second, adjacent monomer. The specific succession of the monomers is called the RNA-sequence. Usually RNA may be obtainable by transcription of a DNA-sequence, e.g., inside a cell. In eukaryotic cells, transcription is typically performed inside the nucleus or the mitochondria. *In vivo*, transcription of DNA usually results in the so-called premature RNA which has to be processed into so-called messenger-RNA, usually abbreviated as mRNA. Processing of the premature RNA, e.g. in eukaryotic organisms, comprises a variety of different posttranscriptional-modifications such as splicing, 5'-capping, polyadenylation, export from the nucleus or the mitochondria and the like. The sum of these processes is also called maturation of RNA. The mature messenger RNA usually provides the nucleotide sequence that may be translated into an amino acid sequence of a particular peptide or protein. Typically, a mature mRNA comprises a 5'-cap, a 5'UTR, an open reading frame, a 3'UTR and a poly(A) sequence. Aside from messenger RNA, several non-coding types of RNA exist which may be involved in regulation of transcription and/or translation.

Sequence of a nucleic acid molecule: The sequence of a nucleic acid molecule is typically understood to be the particular and individual order, i.e. the succession of its nucleotides. The sequence of a protein or peptide is typically understood to be the order, i.e. the succession of its amino acids.

5 Sequence identity: Two or more sequences are identical if they exhibit the same length and order of nucleotides or amino acids. The percentage of identity typically describes the extent to which two sequences are identical, i.e. it typically describes the percentage of nucleotides that correspond in their sequence position with identical nucleotides of a reference-sequence. For determination of the degree of identity, the sequences to be
10 compared are considered to exhibit the same length, i.e. the length of the longest sequence of the sequences to be compared. This means that a first sequence consisting of 8 nucleotides is 80% identical to a second sequence consisting of 10 nucleotides comprising the first sequence. In other words, in the context of the present invention, identity of sequences preferably relates to the percentage of nucleotides of a sequence which have the
15 same position in two or more sequences having the same length. Gaps are usually regarded as non-identical positions, irrespective of their actual position in an alignment.

Stabilized nucleic acid molecule: A stabilized nucleic acid molecule is a nucleic acid molecule, preferably a DNA or RNA molecule that is modified such, that it is more stable to disintegration or degradation, e.g., by environmental factors or enzymatic digest, such as by
20 an exo- or endonuclease degradation, than the nucleic acid molecule without the modification. Preferably, a stabilized nucleic acid molecule in the context of the present invention is stabilized in a cell, such as a prokaryotic or eukaryotic cell, preferably in a mammalian cell, such as a human cell. The stabilization effect may also be exerted outside of cells, e.g. in a buffer solution etc., for example, in a manufacturing process for a
25 pharmaceutical composition comprising the stabilized nucleic acid molecule.

Transfection: The term 'transfection' refers to the introduction of nucleic acid molecules, such as DNA or RNA (e.g. mRNA) molecules, into cells, preferably into eukaryotic cells. In the context of the present invention, the term 'transfection' encompasses any method known to the skilled person for introducing nucleic acid molecules into cells, preferably
30 into eukaryotic cells, such as into mammalian cells. Such methods encompass, for example, electroporation, lipofection, e.g. based on cationic lipids and/or liposomes, calcium

phosphate precipitation, nanoparticle based transfection, virus based transfection, or transfection based on cationic polymers, such as DEAE-dextran or polyethylenimine etc. Preferably, the introduction is non-viral.

Vaccine: A vaccine is typically understood to be a prophylactic or therapeutic material providing at least one antigen, preferably an immunogen. The antigen or immunogen may be derived from any material that is suitable for vaccination. For example, the antigen or immunogen may be derived from a pathogen, such as from bacteria or virus particles etc., or from a tumor or cancerous tissue. The antigen or immunogen stimulates the body's adaptive immune system to provide an adaptive immune response.

Vector: The term 'vector' refers to a nucleic acid molecule, preferably to an artificial nucleic acid molecule. A vector in the context of the present invention is suitable for incorporating or harboring a desired nucleic acid sequence, such as a nucleic acid sequence comprising an open reading frame. Such vectors may be storage vectors, expression vectors, cloning vectors, transfer vectors etc. A storage vector is a vector which allows the convenient storage of a nucleic acid molecule, for example, of an mRNA molecule. Thus, the vector may comprise a sequence corresponding, e.g., to a desired mRNA sequence or a part thereof, such as a sequence corresponding to the open reading frame and the 3'UTR of an mRNA. An expression vector may be used for production of expression products such as RNA, e.g. mRNA, or peptides, polypeptides or proteins. For example, an expression vector may comprise sequences needed for transcription of a sequence stretch of the vector, such as a promoter sequence, e.g. an RNA promoter sequence. A cloning vector is typically a vector that contains a cloning site, which may be used to incorporate nucleic acid sequences into the vector. A cloning vector may be, e.g., a plasmid vector or a bacteriophage vector. A transfer vector may be a vector which is suitable for transferring nucleic acid molecules into cells or organisms, for example, viral vectors. A vector in the context of the present invention may be, e.g., an RNA vector or a DNA vector. Preferably, a vector is a DNA molecule. Preferably, a vector in the sense of the present application comprises a cloning site, a selection marker, such as an antibiotic resistance factor, and a sequence suitable for multiplication of the vector, such as an origin of replication. Preferably, a vector in the context of the present application is a plasmid vector.

Vehicle: A vehicle is typically understood to be a material that is suitable for storing, transporting, and/or administering a compound, such as a pharmaceutically active compound. For example, it may be a physiologically acceptable liquid which is suitable for storing, transporting, and/or administering a pharmaceutically active compound.

5 3'-untranslated region (3'UTR): A 3'UTR is typically the part of an mRNA which is located between the protein coding region (i.e. the open reading frame) and the poly(A) sequence of the mRNA. A 3'UTR of the mRNA is not translated into an amino acid sequence. The 3'UTR sequence is generally encoded by the gene which is transcribed into the respective mRNA during the gene expression process. The genomic sequence is first
10 transcribed into pre-mature mRNA, which comprises optional introns. The pre-mature mRNA is then further processed into mature mRNA in a maturation process. This maturation process comprises the steps of 5'capping, splicing the pre-mature mRNA to excise optional introns and modifications of the 3'-end, such as polyadenylation of the 3'-end of the pre-mature mRNA and optional endo- or exonuclease cleavages etc. In the context of the
15 present invention, a 3'UTR corresponds to the sequence of a mature mRNA which is located 3' to the stop codon of the protein coding region, preferably immediately 3' to the stop codon of the protein coding region, and which extends to the 5'-side of the poly(A) sequence, preferably to the nucleotide immediately 5' to the poly(A) sequence. The term "corresponds to" means that the 3'UTR sequence may be an RNA sequence, such as in the
20 mRNA sequence used for defining the 3'UTR sequence, or a DNA sequence which corresponds to such RNA sequence. In the context of the present invention, the term "a 3'UTR of a gene", such as "a 3'UTR of an albumin gene", is the sequence which corresponds to the 3'UTR of the mature mRNA derived from this gene, i.e. the mRNA obtained by transcription of the gene and maturation of the pre-mature mRNA. The term
25 "3'UTR of a gene" encompasses the DNA sequence and the RNA sequence of the 3'UTR.

5'-untranslated region (5'UTR): A 5'UTR is typically understood to be a particular section of messenger RNA (mRNA). It is located 5' of the open reading frame of the mRNA. Typically, the 5'UTR starts with the transcriptional start site and ends one nucleotide before
30 the start codon of the open reading frame. The 5'UTR may comprise elements for controlling gene expression, also called regulatory elements. Such regulatory elements may be, for example, ribosomal binding sites or a 5'-Terminal Oligopyrimidine Tract. The 5'UTR may be posttranscriptionally modified, for example by addition of a 5'-cap. In the context of

the present invention, a 5'UTR corresponds to the sequence of a mature mRNA which is located between the 5'cap and the start codon. Preferably, the 5'UTR corresponds to the sequence which extends from a nucleotide located 3' to the 5'-cap, preferably from the nucleotide located immediately 3' to the 5'cap, to a nucleotide located 5' to the start codon of the protein coding region, preferably to the nucleotide located immediately 5' to the start codon of the protein coding region. The nucleotide located immediately 3' to the 5'cap of a mature mRNA typically corresponds to the transcriptional start site. The term "corresponds to" means that the 5'UTR sequence may be an RNA sequence, such as in the mRNA sequence used for defining the 5'UTR sequence, or a DNA sequence which corresponds to such RNA sequence. In the context of the present invention, the term "a 5'UTR of a gene", such as "a 5'UTR of a TOP gene", is the sequence which corresponds to the 5'UTR of the mature mRNA derived from this gene, i.e. the mRNA obtained by transcription of the gene and maturation of the pre-mature mRNA. The term "5'UTR of a gene" encompasses the DNA sequence and the RNA sequence of the 5'UTR.

5'Terminal Oligopyrimidine Tract (TOP): The 5'terminal oligopyrimidine tract (TOP) is typically a stretch of pyrimidine nucleotides located at the 5' terminal region of a nucleic acid molecule, such as the 5' terminal region of certain mRNA molecules or the 5' terminal region of a functional entity, e.g. the transcribed region, of certain genes. The sequence starts with a cytidine, which usually corresponds to the transcriptional start site, and is followed by a stretch of usually about 3 to 30 pyrimidine nucleotides, more often 3 to 15 pyrimidine nucleotides. For example, the TOP may comprise 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or even more nucleotides. The pyrimidine stretch and thus the 5' TOP ends one nucleotide 5' to the first purine nucleotide located downstream of the TOP. Messenger RNA that contains a 5'-terminal oligopyrimidine tract is often referred to as TOP mRNA. Accordingly, genes that provide such messenger RNAs are referred to as TOP genes. TOP sequences have, for example, been found in genes and mRNAs encoding peptide elongation factors and ribosomal proteins.

TOP motif: In the context of the present invention, a TOP motif is a nucleic acid sequence which corresponds to a 5'TOP as defined above. Thus, a TOP motif in the context of the present invention is preferably a stretch of pyrimidine nucleotides having a length of

3-30 nucleotides. Preferably, the TOP-motif consists of at least 3 pyrimidine nucleotides, preferably at least 4 pyrimidine nucleotides, preferably at least 5 pyrimidine nucleotides, more preferably at least 6 nucleotides, more preferably at least 7 nucleotides, most preferably at least 8 pyrimidine nucleotides, wherein the stretch of pyrimidine nucleotides preferably starts at its 5' end with a cytosine nucleotide. In TOP genes and TOP mRNAs, the TOP-motif preferably starts at its 5' end with the transcriptional start site and ends one nucleotide 5' to the first purin residue in said gene or mRNA. A TOP motif in the sense of the present invention is preferably located at the 5' end of a sequence which represents a 5'UTR or at the 5' end of a sequence which codes for a 5'UTR. Thus, preferably, a stretch of 3 or more pyrimidine nucleotides is called "TOP motif" in the sense of the present invention if this stretch is located at the 5' end of a respective sequence, such as the artificial nucleic acid molecule according to the present invention, the 5'UTR element of the artificial nucleic acid molecule according to the present invention, or the nucleic acid sequence which is derived from the 5'UTR of a TOP gene as described herein. In other words, a stretch of 3 or more pyrimidine nucleotides which is not located at the 5'-end of a 5'UTR or a 5'UTR element but anywhere within a 5'UTR or a 5'UTR element is preferably not referred to as "TOP motif".

TOP gene: TOP genes are typically characterised by the presence of a 5' terminal oligopyrimidine tract. Furthermore, most TOP genes are characterized by a growth-associated translational regulation. However, also TOP genes with a tissue specific translational regulation are known. As defined above, the 5'UTR of a TOP gene corresponds to the sequence of a 5'UTR of a mature mRNA derived from a TOP gene, which preferably extends from the nucleotide located 3' to the 5' cap to the nucleotide located 5' to the start codon. A 5'UTR of a TOP gene typically does not comprise any start codons, preferably no upstream AUGs (uAUGs) or upstream open reading frames (uORFs). Therein, upstream AUGs and upstream open reading frames are typically understood to be AUGs and open reading frames that occur 5' of the start codon (AUG) of the open reading frame that should be translated. The 5'UTRs of TOP genes are generally rather short. The lengths of 5'UTRs of TOP genes may vary between 20 nucleotides up to 500 nucleotides, and are typically less than about 200 nucleotides, preferably less than about 150 nucleotides, more preferably less than about 100 nucleotides. Exemplary 5'UTRs of TOP genes in the sense of the present invention are the nucleic acid sequences extending from the nucleotide at position 5 to the nucleotide

located immediately 5' to the start codon (e.g. the ATG) in the sequences according to SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422.

In a first aspect, the present invention relates to an artificial nucleic acid molecule comprising

- 5 a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene; and
- b. at least one open reading frame (ORF).

In a first embodiment there is provided an artificial nucleic acid molecule comprising:

- 10 a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene;
- b. at least one open reading frame (ORF); and
- 15 c. at least one 3'UTR element comprising or consisting of a nucleic acid sequence derived from a 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.

In a second embodiment there is provided a vector comprising:

- a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP
- 20 gene or which is derived from a variant of the 5'UTR of a TOP gene;
- b. at least one open reading frame (ORF) and/or at least one cloning site; and
- c. at least one 3'UTR element, which comprises or consists of a nucleic acid sequence derived from a 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.

25

In a third embodiment there is provided a cell comprising the artificial nucleic acid molecule according to the first embodiment or the vector according to the second embodiment.

In a fourth embodiment there is provided a pharmaceutical composition comprising the artificial nucleic acid molecule according to the first embodiment, the vector according to the second

30 embodiment, or the cell according to the third embodiment.

In a fifth embodiment there is provided use of the artificial nucleic acid molecule according to the first embodiment, the vector according to the second embodiment, the cell according to the third embodiment, or the pharmaceutical composition according to the fourth embodiment for the manufacture of a medicament.

- 5 In a sixth embodiment there is provided use of the artificial nucleic acid molecule according to the first embodiment, the vector according to the second embodiment, the cell according to the third embodiment, or the pharmaceutical composition according to the fourth embodiment as a vaccine or in gene therapy.

- 10 In a seventh embodiment there is provided a method for treating or preventing a disorder comprising administering the artificial nucleic acid molecule according to the first embodiment, the vector according to the second embodiment, the cell according to the third embodiment, or the pharmaceutical composition according to the fourth embodiment to a subject in need thereof.

- 15 In an eighth embodiment there is provided a method of treating or preventing a disorder comprising transfection of a cell with the artificial nucleic acid molecule according to the first embodiment or the vector according to the second embodiment.

In a ninth embodiment there is provided a method for increasing protein production from an artificial nucleic acid molecule, comprising the step of providing the artificial nucleic acid molecule with:

- 20 i. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene, and
- ii. at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.

- 25 In a tenth embodiment there is provided a kit comprising an artificial nucleic acid molecule according to the first embodiment, the vector according to the second embodiment, or the cell according to the third embodiment, and/or the pharmaceutical composition according to the fourth embodiment.

Such an artificial nucleic acid molecule may be DNA or RNA. In case the artificial nucleic acid molecule is DNA it may be used for providing RNA, preferably an mRNA with a corresponding sequence as is described further below. The inventive artificial nucleic acid molecule is particularly useful in gene therapy and genetic vaccination because it may provide increased
5 and/or prolonged protein production of the protein encoded by the open reading frame. It is preferred, if the components (a) and (b) are heterologous, such that the inventive nucleic acid molecule does not occur naturally, but is an artificial chimeric recombinant nucleic acid molecule.

In this context, the term '5'UTR element' preferably refers to a nucleic acid sequence which represents a 5'UTR of an artificial nucleic acid sequence, such as an artificial mRNA, or which
10 codes for a 5'UTR of an artificial nucleic acid molecule. Thus, preferably, a 5'UTR element may be the 5'UTR of an mRNA, preferably of an artificial mRNA, or it may be the transcription template for a 5'UTR of an mRNA. Thus, a 5'UTR element preferably is a nucleic acid sequence which corresponds to the 5'UTR of an mRNA, preferably to the 5'UTR of an artificial mRNA, such as an mRNA obtained by transcription of a genetically engineered vector construct.

15 Preferably, a 5'UTR element in the sense of the present invention functions as a 5'UTR or codes for a nucleotide sequence that fulfils the function of a 5'UTR. The term '5'UTR element' may also refer to a fragment or part of a 5'UTR of an artificial nucleic acid sequence, such as an artificial mRNA, or which codes for a part or fragment of a 5'UTR of an artificial nucleic acid molecule. This means that the 5'UTR element in the sense of the present invention may be comprised in
20 the 5'UTR of an artificial

nucleic acid sequence, such as an artificial mRNA, or which codes for a 5'UTR of an artificial nucleic acid molecule.

5 According to the invention, the 5'UTR element comprises or consists of a nucleic acid sequence that is derived from the 5'UTR of a TOP gene or from a variant of the 5'UTR of a TOP gene.

The term 'a nucleic acid sequence which is derived from the 5'UTR of a TOP gene' preferably refers to a nucleic acid sequence which is based on the 5'UTR sequence of a TOP gene or on a fragment thereof. This term includes sequences corresponding to the entire 5'UTR sequence, i.e. the full length 5'UTR sequence of a TOP gene, and sequences corresponding to a fragment of the 5'UTR sequence of a TOP gene. Preferably, a fragment of a 5'UTR of a TOP gene consists of a continuous stretch of nucleotides corresponding to a continuous stretch of nucleotides in the full-length 5'UTR of a TOP gene, which represents at least 20%, preferably at least 30%, more preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, even more preferably at least 70%, even more preferably at least 80%, and most preferably at least 90% of the full-length 5'UTR of a TOP gene. Such a fragment, in the sense of the present invention, is preferably a functional fragment as described herein. A particularly preferred fragment of a 5'UTR of a TOP gene is a 5'UTR of a TOP gene lacking the 5'TOP motif, which typically corresponds to a pyrimidine stretch of 3 to 30 pyrimidine nucleotides at the 5' terminus of the 5'UTR of a TOP gene. For the above preferred embodiment of the invention employing a 5'UTR of a TOP gene, the 5'UTR (comprised by the inventive nucleic acid molecule) starts with the first nucleotide following the most 3'-terminal nucleotide of the 5'TOP motif. In case the 5'TOP motif does not correspond to the 5'terminal part of the 5'UTR of the TOP gene, the 5' UTR (of the TOP gene) employed in the inventive nucleic acid may consist of the nucleotide sequence located upstream of the 5'terminus of the 5'TOP motif and/or of the nucleotide sequence located downstream of the 3'terminus of the 5'TOP motif. In an alternative embodiment, the 5' motif of a 5'UTR of a TOP gene may be rendered dysfunctional by e.g. introducing one or more purine nucleotides, which interrupt the monotonic pyrimidine nucleotide stretch of the 5'TOP motif such that the modified (interrupted) 5'TOP motif sequence cannot exert its regulatory function any longer, in particular cannot exert its function as an element for translational control. Another way of rendering the 5' TOP motif

dysfunctional is the deletion of one or more pyrimidine nucleotides of the 5'TOP motif sequence (either at the termini and/or within the 5'TOP motif).

In one embodiment, the 5'UTR of a TOP gene will not be derived from the 5'UTR of ribosomal proteins (rp) mRNA (in particular not from mammalian 5'UTR of rp mRNA, more specifically not from rpP2 (e.g. rat rpP2), rpL32, rpL30, rpL13a (e.g. mouse transplantation antigen P198), rpS20, rpS6, rpL12 or rpS16 mRNA or not from an rpS19 mRNA (e.g. from *Xenopus*). In another embodiment, the 5'UTR of a TOP gene is not derived from the 5'UTR of a EF1alpha or (hamster) EF2 mRNA. The 5'UTRs of these afore-mentioned rp mRNAs are specifically not used, if they are linked to reporter genes in the ORF of the inventive nucleic acid. If e.g. the 5'UTR of rpS16 mRNA is used for the inventive nucleic acid, that 5'UTR will either not contain the 5'TOP motif sequence (composed of the oligonucleotide (CCTTTTCC or CCUUUUCC) or will contain a dysfunctional variant thereof by e.g. interruption of the oligopyrimidine sequence by purine nucleotides or by deletion of one or more pyrimidine nucleotides of that 5'TOP motif. Accordingly, the dysfunctional mutants may e.g. contain one or more purine nucleotides within the 5'TOP motif sequence thereby lacking the translational control function exerted by the 5'TOP motif, e.g. by abolishing its interaction with other regulatory compounds, e.g. miRNA or interaction with granule-associated proteins TIA-1 and TIAR.

The term '5'UTR of a TOP gene' preferably refers to the 5'UTR of a naturally occurring TOP gene.

The terms 'variant of the 5'UTR of a TOP gene' and 'variant thereof' in the context of a 5'UTR of a TOP gene refers to a variant of the 5'UTR of a naturally occurring TOP gene, preferably to a variant of the 5'UTR of a vertebrate TOP gene, preferably to a variant of the 5'UTR of a mammalian TOP gene, more preferably to a variant of the 5'UTR of a human TOP gene. Such variant may be a modified 5'UTR of a TOP gene. For example, a variant 5'UTR may exhibit one or more nucleotide deletions, insertions, additions and/or substitutions compared to the naturally occurring 5'UTR from which the variant is derived. Preferably, a variant of a 5'UTR of a TOP gene is at least 40%, preferably at least 50%, more preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, even more preferably at least 90%, most preferably at least 95% identical to the

naturally occurring 5'UTR the variant is derived from. Preferably, the variant is a functional variant as described herein.

5 The term "a nucleic acid sequence that is derived from a variant of the 5'UTR of a TOP gene" preferably refers to a nucleic acid sequence which is based on a variant of a 5'UTR sequence of a TOP gene or on a fragment thereof. This term includes sequences corresponding to the entire variant 5'UTR sequence, i.e. the full length variant 5'UTR sequence of a TOP gene, and sequences corresponding to a fragment of the variant 5'UTR sequence of a TOP gene. Preferably, a fragment of a variant of the 5'UTR of a TOP gene
10 consists of a continuous stretch of nucleotides corresponding to a continuous stretch of nucleotides in the full-length variant 5'UTR of a TOP gene, which represents at least 20%, preferably at least 30%, more preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, even more preferably at least 70%, even more preferably at least 80%, and most preferably at least 90% of the full-length variant 5'UTR of a TOP gene.
15 Such a fragment of a variant, in the sense of the present invention, is preferably a functional fragment as described herein.

Thus, the 5'UTR element of the artificial nucleic acid molecule may comprise or consist of a fragment of the 5'UTR of a TOP gene or of a fragment of a variant of the 5'UTR of a TOP
20 gene or may comprise or consist of the entire 5'UTR of a TOP gene or may comprise or consist of a variant of the 5'UTR of a TOP gene.

The 5'UTR element is preferably suitable for increasing protein production from the artificial nucleic acid molecule.

25 Preferably, the at least one 5'UTR element is functionally linked to the ORF. This means preferably that the 5'UTR element is associated with the ORF such that it may exert a function, such as a protein production increasing function for the protein encoded by the ORF or a stabilizing function on the artificial nucleic acid molecule. Preferably, the 5'UTR
30 element and the ORF are associated in 5'→3' direction. Thus, preferably, the artificial nucleic acid molecule comprises the structure 5'-5'UTR element-(optional)linker-ORF-3', wherein the linker may be present or absent. For example, the linker may be one or more

nucleotides, such as a stretch of 1-50 or 1-20 nucleotides, e.g., comprising or consisting of one or more restriction enzyme recognition sites (restriction sites).

Preferably, the 5'UTR element and the at least one open reading frame are heterologous.

- 5 The term 'heterologous' in this context means that the open reading frame and the 5'UTR element are not occurring naturally (in nature) in this combination. Preferably, the 5'UTR element is derived from a different gene than the open reading frame. For example, the ORF may be derived from a different gene than the 5'UTR element, e.g. encoding a different protein or the same protein but of a different species etc. For example, the ORF does not
10 encode the protein which is encoded by the gene from which the 5'UTR element is derived.

- In a preferred embodiment, the 5'UTR element, preferably the artificial nucleic acid molecule, does not comprise a complete TOP-motif or 5'TOP sequence. Thus, preferably, the 5'UTR element, preferably the artificial nucleic acid molecule, does not comprise the
15 complete TOP-motif of the TOP gene from which the nucleic acid sequence of the 5'UTR element is derived. For example, the 5'UTR element or the artificial nucleic acid molecule according to the present invention may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine residues of the TOP-motif or 5'TOP, preferably 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine residues of the TOP-motif located at the 3'side of the TOP-motif or 5'TOP.
20 For example, the 5'UTR element may comprise or consist of a nucleic acid sequence which starts at its 5'end with a pyrimidine residue that corresponds to residue 2, 3, 4, 5, 6, 7, 8, 9, 10 etc. of the TOP-motif or 5'TOP of the TOP gene from which the nucleic acid sequence of the 5'UTR element is derived.

- 25 It is particularly preferred that the 5'UTR element, preferably the artificial nucleic acid molecule according to the present invention, does not comprise a TOP-motif or a 5'TOP. For example, the nucleic acid sequence of the 5'UTR element which is derived from a 5'UTR of a TOP gene starts at its 5'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 downstream of the 5'terminal oligopyrimidine tract (TOP) of the 5'UTR of a
30 TOP gene. Position 1 downstream of the 5'terminal oligopyrimidine tract (TOP) is the first purine based nucleotide 3' of the TOP-motif or the 5'TOP. Accordingly, position 1 downstream of the 5'terminal oligopyrimidine tract is the first nucleotide following the 3'-end of the 5'terminal oligopyrimidine tract in 5'-3'-direction. Likewise, position 2

downstream of the 5'TOP is the second nucleotide following the end of the 5'terminal oligopyrimidine tract, position 3 the third nucleotide and so on.

Therefore, the 5'UTR element preferably starts 5, 10, 15, 20, 25, 30, 40 or 50 nucleotides downstream of the transcriptional start site of the 5'UTR of a TOP gene.

In some embodiments, the nucleic acid sequence of the 5'UTR element which is derived from a 5'UTR of a TOP gene terminates at its 3'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 upstream of the start codon (e.g. A(U/T)G) of the gene or mRNA it is derived from. Thus, the 5'UTR element does not comprise any part of the protein coding region. Thus, preferably, the only protein coding part of the inventive artificial nucleic acid molecule is provided by the open reading frame. However, the open reading frame is preferably derived – as said above – from a gene that is different to the gene the 5'UTR element is derived from.

It is particularly preferred that the 5'UTR element does not comprise a start codon, such as the nucleotide sequence A(U/T)G. Thus, preferably, the artificial nucleic acid molecule will not comprise any upstream AUGs (or upstream ATGs in case it is a DNA molecule). In other words, in some embodiments, it may be preferred that the AUG or ATG, respectively, of the open reading frame is the only start codon of the artificial nucleic acid molecule.

Additionally, it is preferred that the 5'UTR element does not comprise an open reading frame. Thus, preferably, the artificial nucleic acid molecule will not comprise any upstream open reading frames.

The nucleic acid sequence which is derived from the 5'UTR of a TOP gene is derived from a eukaryotic TOP gene, preferably a plant or animal TOP gene, more preferably a chordate TOP gene, even more preferably a vertebrate TOP gene, most preferably a mammalian TOP gene, such as a human or mouse TOP gene.

Preferably, the artificial nucleic acid molecule according to the present invention comprises a 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene, wherein the TOP gene is a plant or animal TOP gene, more preferably a chordate

TOP gene, even more preferably a vertebrate TOP gene, most preferably a mammalian TOP gene, such as a human or mouse TOP gene and which optionally does not comprise the nucleotide sequence A(U/T)G and optionally does not comprise an open reading frame; at least one open reading frame (ORF); wherein optionally the 5'UTR element does not
5 comprise a TOP motif and wherein optionally the 5'UTR element starts at its 5'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 downstream of the 5'terminal oligopyrimidine tract (TOP) of the 5'UTR of a TOP gene and wherein further optionally the 5'UTR element which is derived from a 5'UTR of a TOP gene terminates at its 3'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 upstream of the start codon
10 (A(U/T)G) of the gene or mRNA it is derived from.

For example, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422, from the homologs of
15 SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422, from a variant thereof, or a corresponding RNA sequence. The term "homologs of SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422" refers to sequences of other species, e.g. other species than Homo sapiens (human) or Mus musculus (mouse), which are homologous to the sequences according to SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422. For example, SEQ ID NO. 1 relates to a
20 sequence comprising the 5'UTR of Homo sapiens alpha 2 macroglobulin (A2M). A homolog of SEQ ID NO. 1 in the context of the present invention is any such sequence derived from an alpha 2 macroglobulin (A2M) gene or mRNA of another species than Homo sapiens, such as any vertebrate, preferably any mammalian alpha 2 macroglobulin (A2M) gene other than the human alpha 2 macroglobulin (A2M) gene, such as a mouse, rat, rabbit, monkey
25 etc. alpha 2 macroglobulin (A2M) gene.

In a preferred embodiment, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a nucleic acid sequence extending from nucleotide position 5 (i.e. the nucleotide that is located at position 5 in the sequence) to the nucleotide
30 position immediately 5' to the start codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence, of a nucleic acid sequence selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, from the homologs of SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or

SEQ ID NO. 1422, from a variant thereof, or a corresponding RNA sequence. It is particularly preferred that the 5' UTR element is derived from a nucleic acid sequence extending from the nucleotide position immediately 3' to the 5'TOP to the nucleotide position immediately 5' to the start codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence, of a nucleic acid sequence selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, from the homologs of SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, from a variant thereof, or a corresponding RNA sequence.

- 10 In a preferred embodiment, the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to a nucleic acid sequence
- 15 extending from nucleotide position 5 to the nucleotide position immediately 5' to the start codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence of a nucleic acid sequence selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, or a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic
- 20 acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to a nucleic acid sequence extending from nucleotide position 5 to the nucleotide position immediately 5' to the start
- 25 codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence of a nucleic acid sequence, selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, or a corresponding RNA sequence, wherein, preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 5'UTR the fragment is derived
- 30 from.

Preferably, the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least

about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to a nucleic acid sequence extending from the nucleotide position immediately 3' to the 5'TOP to the nucleotide position immediately 5' to the start codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence, of a nucleic acid sequence selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, or a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to a nucleic acid sequence extending from the nucleotide position immediately 3' to the 5'TOP to the nucleotide position immediately 5' to the start codon (located at the 3' end of the sequences), e.g. the nucleotide position immediately 5' to the ATG sequence, of a nucleic acid sequence selected from SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, or a corresponding RNA sequence, wherein, preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 5'UTR the fragment is derived from.

Preferably, the above defined fragments and variants (e.g. exhibiting at least 40% identity) of the sequences according to SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 or SEQ ID NO. 1422, are functional fragments and variants as described herein.

Furthermore, the artificial nucleic acid molecule according to the present invention may comprise more than one 5'UTR elements as described above. For example, the artificial nucleic acid molecule according to the present invention may comprise one, two, three, four or more 5'UTR elements, wherein the individual 5'UTR elements may be the same or they may be different. For example, the artificial nucleic acid molecule according to the present invention may comprise two essentially identical 5'UTR elements as described above, e.g. two 5'UTR elements comprising or consisting of a nucleic acid sequence which is derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422, from the homologs of

SEQ ID NOs. 1-1363, SEQ ID NO. 1395, SEQ ID NO. 1421 and SEQ ID NO. 1422, from a variant thereof, or a corresponding RNA sequence or from functional variants thereof, functional fragments thereof, or functional variant fragments thereof as described above.

- 5 In a particularly preferred embodiment, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a 5'UTR of a TOP gene encoding a ribosomal protein or from a variant of a 5'UTR of a TOP gene encoding a ribosomal protein. Particularly preferred 5'UTR elements comprise or consist of a nucleic acid sequence which are derived from a 5' UTR of a TOP gene coding for a ribosomal protein selected from RPSA, RPS2, RPS3, RPS3A, RPS4, RPS5, RPS6, RPS7, RPS8, RPS9,
10 RPS10, RPS11, RPS12, RPS13, RPS14, RPS15, RPS15A, RPS16, RPS17, RPS18, RPS19, RPS20, RPS21, RPS23, RPS24, RPS25, RPS26, RPS27, RPS27A, RPS28, RPS29, RPS30, RPL3, RPL4, RPL5, RPL6, RPL7, RPL7A, RPL8, RPL9, RPL10, RPL10A, RPL11, RPL12, RPL13, RPL13A, RPL14, RPL15, RPL17, RPL18, RPL18A,
15 RPL19, RPL21, RPL22, RPL23, RPL23A, RPL24, RPL26, RPL27, RPL27A, RPL28, RPL29, RPL30, RPL31, RPL32, RPL34, RPL35, RPL35A, RPL36, RPL36A, RPL37, RPL37A, RPL38, RPL39, RPL40, RPL41, RPLP0, RPLP1, RPLP2, RPLP3, UBA52. Particularly preferred are nucleic acid sequences which are derived from a 5' UTR of TOP genes vertebrate coding for ribosomal proteins, such as mammalian ribosomal
20 proteins e.g. human or mouse ribosomal proteins.

For example, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a 5'UTR of a nucleic acid sequence according to any of SEQ ID NOs: 170, 232, 244, 259, 1284, 1285, 1286, 1287, 1288, 1289, 1290, 1291, 1292, 1293, 1294, 1295, 1296, 1297, 1298, 1299, 1300, 1301, 1302, 1303, 1304, 1305, 1306, 1307, 1308,
25 1309, 1310, 1311, 1312, 1313, 1314, 1315, 1316, 1317, 1318, 1319, 1320, 1321, 1322, 1323, 1324, 1325, 1326, 1327, 1328, 1329, 1330, 1331, 1332, 1333, 1334, 1335, 1336, 1337, 1338, 1339, 1340, 1341, 1342, 1343, 1344, 1346, 1347, 1348, 1349, 1350, 1351, 1352, 1353, 1354, 1355, 1356, 1357, 1358, 1359, or 1360; a corresponding RNA sequence, a homolog thereof, or a variant thereof as described herein, preferably lacking the
30 5'TOP motif. As described above, the sequence extending from position 5 to the nucleotide immediately 5' to the ATG (which is located at the 3'end of the sequences) corresponds to the 5'UTR of said sequences.

Preferably, the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the 5'UTR of a nucleic acid sequence according to any of SEQ ID NOs.: 170, 232, 244, 259, 1284, 1285, 1286, 1287, 1288, 1289, 1290, 1291, 1292, 1293, 1294, 1295, 1296, 1297, 1298, 1299, 1300, 1301, 1302, 1303, 1304, 1305, 1306, 1307, 1308, 1309, 1310, 1311, 1312, 1313, 1314, 1315, 1316, 1317, 1318, 1319, 1320, 1321, 1322, 1323, 1324, 1325, 1326, 1327, 1328, 1329, 1330, 1331, 1332, 1333, 1334, 1335, 1336, 1337, 1338, 1339, 1340, 1341, 1342, 1343, 1344, 1346, 1347, 1348, 1349, 1350, 1351, 1352, 1353, 1354, 1355, 1356, 1357, 1358, 1359, or 1360; or a corresponding RNA sequence, preferably lacking the 5'TOP motif, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the 5'UTR of a nucleic acid sequence according to SEQ ID NOs.: 170, 232, 244, 259, 1284, 1285, 1286, 1287, 1288, 1289, 1290, 1291, 1292, 1293, 1294, 1295, 1296, 1297, 1298, 1299, 1300, 1301, 1302, 1303, 1304, 1305, 1306, 1307, 1308, 1309, 1310, 1311, 1312, 1313, 1314, 1315, 1316, 1317, 1318, 1319, 1320, 1321, 1322, 1323, 1324, 1325, 1326, 1327, 1328, 1329, 1330, 1331, 1332, 1333, 1334, 1335, 1336, 1337, 1338, 1339, 1340, 1341, 1342, 1343, 1344, 1346, 1347, 1348, 1349, 1350, 1351, 1352, 1353, 1354, 1355, 1356, 1357, 1358, 1359, or 1360; or a corresponding RNA sequence, wherein, preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 5'UTR, preferably lacking the 5'TOP motif. Preferably, the fragment exhibits a length of at least about 20 nucleotides or more, preferably of at least about 30 nucleotides or more, more preferably of at least about 40 nucleotides or more. Preferably, the fragment is a functional fragment as described herein.

Preferably, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a 5'UTR of a TOP gene encoding a ribosomal Large protein (RPL) or from a variant of a 5'UTR of a TOP gene encoding a ribosomal Large protein (RPL). For example,

the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from a 5'UTR of a nucleic acid sequence according to any of SEQ ID NOs: 67, 259, 1284-1318, 1344, 1346, 1348-1354, 1357, 1358, 1421 and 1422, a corresponding RNA sequence, a homolog thereof, or a variant thereof as described herein, preferably lacking the 5'TOP motif.

Preferably, the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the 5'UTR of a nucleic acid sequence according to any of SEQ ID NOs. 67, 259, 1284-1318, 1344, 1346, 1348-1354, 1357, 1358, 1421 and 1422 or a corresponding RNA sequence, preferably lacking the 5'TOP motif, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the 5'UTR of a nucleic acid sequence according to SEQ ID NOs: 67, 259, 1284-1318, 1344, 1346, 1348-1354, 1357, 1358, 1421 and 1422 or a corresponding RNA sequence, wherein, preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 5'UTR, preferably lacking the 5'TOP motif. Preferably, the fragment exhibits a length of at least about 20 nucleotides or more, preferably of at least about 30 nucleotides or more, more preferably of at least about 40 nucleotides or more. Preferably, the fragment is a functional fragment as described herein.

In a particularly preferred embodiment, the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), an ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, an hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), an androgen-induced 1 gene (AIG1), cytochrome c oxidase subunit VIc gene (COX6C), or a N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or

from a variant thereof, preferably from a vertebrate ribosomal protein Large 32 gene (RPL32), a vertebrate ribosomal protein Large 35 gene (RPL35), a vertebrate ribosomal protein Large 21 gene (RPL21), a vertebrate ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a vertebrate hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a vertebrate androgen-induced 1 gene (AIG1), a vertebrate cytochrome c oxidase subunit VIc gene (COX6C), or a vertebrate N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, more preferably from a mammalian ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), a mammalian ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a mammalian hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a mammalian androgen-induced 1 gene (AIG1), a mammalian cytochrome c oxidase subunit VIc gene (COX6C), or a mammalian N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, most preferably from a human ribosomal protein Large 32 gene (RPL32), a human ribosomal protein Large 35 gene (RPL35), a human ribosomal protein Large 21 gene (RPL21), a human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a human hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a human androgen-induced 1 gene (AIG1), a human cytochrome c oxidase subunit VIc gene (COX6C), or a human N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, wherein preferably the 5'UTR element does not comprise the 5'TOP of said gene.

Accordingly, in a particularly preferred embodiment, the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the nucleic acid sequence according to SEQ ID No. 1368, or SEQ ID NOs 1412-1420, or a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at

least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the nucleic acid sequence according to SEQ ID No. 1368, or SEQ ID NOs 1412-1420, wherein, preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 5'UTR. Preferably, the fragment exhibits a length of at least about 20 nucleotides or more, preferably of at least about 30 nucleotides or more, more preferably of at least about 40 nucleotides or more. Preferably, the fragment is a functional fragment as described herein.

10 Preferably, the at least one 5'UTR element exhibits a length of at least about 20 nucleotides or more, preferably of at least about 30 nucleotides or more, more preferably of at least about 40 nucleotides or more. However, it may be preferred if the 5'UTR element of the artificial nucleic acid molecule is rather short. Accordingly, it may have a length of less than about 200, preferably less than 150, more preferably less than 100 nucleotides. For example, the 5'UTR may have a length of less than about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200 nucleotides. Preferably, the 5'UTR element may have a length of about 20-25, 26-30, 31-35, 36-40, 41-45, 46-50, 51-55, 56-60, 61-65, 66-70, 71-80, 81-85, 86-90, 91-95, 96-100, 101-105, 106-110, 111-115, 116-120, 121-125, 126-130, 131-135, 136-140, 141-145, 146-150, 151-155, 156-160, 161-165, 166-170, 171-175, 176-180, 181-185, 186-190, 191-195, 196-200 or more nucleotides. For example, the 5'UTR element may have a length of about 20, 26, 31, 36, 41, 46, 51, 56, 61, 66, 71, 81, 86, 91, 96, 101, 106, 111, 116, 121, 126, 131, 136, 141, 146, 151, 156, 161, 166, 171, 176, 181, 186, 191 or 196 nucleotides. Preferably, the 5'UTR element may have a length from about 20, 30, 40 or more to less than about 200 nucleotides, more preferably from about 20, 30, 40 or more to less than about 150 nucleotides, most preferably from about 20, 30, 40 or more to less than about 100 nucleotides.

Preferred 5'UTR elements are derived from a 5' UTR of a TOP gene selected from RPSA, RPS2, RPS3, RPS3A, RPS4, RPS5, RPS6, RPS7, RPS8, RPS9, RPS10, RPS11, RPS12, RPS13, RPS14, RPS15, RPS15A, RPS16, RPS17, RPS18, RPS19, RPS20, RPS21, RPS23, RPS24, RPS25, RPS26, RPS27, RPS27A, RPS28, RPS29, RPS30, RPL3, RPL4, RPL5, RPL6, RPL7, RPL7A, RPL8, RPL9, RPL10, RPL10A, RPL11, RPL12, RPL13, RPL13A, RPL14, RPL15, RPL17, RPL18, RPL18A, RPL19, RPL21, RPL22, RPL23, RPL23A, RPL24, RPL26, RPL27,

RPL27A, RPL28, RPL29, RPL30, RPL31, RPL32, RPL34, RPL35, RPL35A, RPL36, RPL36A, RPL37, RPL37A, RPL38, RPL39, RPL40, RPL41, RPLP0, RPLP1, RPLP2, RPLP3, RPLP0, RPLP1, RPLP2, EEF1A1, EEF1B2, EEF1D, EEF1G, EEF2, EIF3E, EIF3F, EIF3H, EIF2S3, EIF3C, EIF3K, EIF3EIP, EIF4A2, PABPC1, HNRNPA1, TPT1, TUBB1, UBA52, NPM1, ATP5G2,
 5 GNB2L1, NME2, UQCRB or from a variant thereof.

In some embodiments, the artificial nucleic acid molecule comprises a 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a vertebrate TOP gene, such as a mammalian, e.g. a human TOP gene, selected from RPSA,
 10 RPS2, RPS3, RPS3A, RPS4, RPS5, RPS6, RPS7, RPS8, RPS9, RPS10, RPS11, RPS12, RPS13, RPS14, RPS15, RPS15A, RPS16, RPS17, RPS18, RPS19, RPS20, RPS21, RPS23, RPS24, RPS25, RPS26, RPS27, RPS27A, RPS28, RPS29, RPS30, RPL3, RPL4, RPL5, RPL6, RPL7, RPL7A, RPL8, RPL9, RPL10, RPL10A, RPL11, RPL12, RPL13, RPL13A, RPL14, RPL15, RPL17, RPL18, RPL18A, RPL19, RPL21, RPL22, RPL23, RPL23A, RPL24, RPL26, RPL27,
 15 RPL27A, RPL28, RPL29, RPL30, RPL31, RPL32, RPL34, RPL35, RPL35A, RPL36, RPL36A, RPL37, RPL37A, RPL38, RPL39, RPL40, RPL41, RPLP0, RPLP1, RPLP2, RPLP3, RPLP0, RPLP1, RPLP2, EEF1A1, EEF1B2, EEF1D, EEF1G, EEF2, EIF3E, EIF3F, EIF3H, EIF2S3, EIF3C, EIF3K, EIF3EIP, EIF4A2, PABPC1, HNRNPA1, TPT1, TUBB1, UBA52, NPM1, ATP5G2, GNB2L1, NME2, UQCRB, or from a variant thereof, wherein preferably the 5'UTR element
 20 does not comprise a TOP-motif or the 5'TOP of said genes, and wherein optionally the 5'UTR element starts at its 5'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 downstream of the 5'terminal oligopyrimidine tract (TOP) and wherein further optionally the 5'UTR element which is derived from a 5'UTR of a TOP gene terminates at its 3'-end with a nucleotide located at position 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 upstream of the
 25 start codon (A(U/T)G) of the gene it is derived from.

In a preferred embodiment, the artificial nucleic acid molecule according to the present invention further comprises

- 30 c. at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene, or from a variant of

the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene.

The term '3'UTR element' refers to a nucleic acid sequence which comprises or consists of
5 a nucleic acid sequence that is derived from a 3'UTR or from a variant of a 3'UTR. A 3'UTR element in the sense of the present invention may represent the 3'UTR of an mRNA, e.g., in the event that the artificial nucleic acid molecule is an mRNA, or it may represent a sequence in a nucleic acid construct, such as a vector construct, that when transcribed represents the 3'UTR of the transcription product, such as the mRNA. Thus, in the sense of
10 the present invention, preferably, a 3'UTR element may be the 3'UTR of an mRNA, preferably of an artificial mRNA, or it may be the transcription template for a 3'UTR of an mRNA. Thus, a 3'UTR element preferably is a nucleic acid sequence which corresponds to the 3'UTR of an mRNA, preferably to the 3'UTR of an artificial mRNA, such as an mRNA obtained by transcription of a genetically engineered vector construct. Preferably, the 3'UTR
15 element fulfils the function of a 3'UTR or encodes a sequence which fulfils the function of a 3'UTR. The term '3'UTR element' furthermore refers to a fragment or part of a 3'UTR of an artificial nucleic acid sequence, such as an artificial mRNA, or which codes for a part or fragment of a 3'UTR of an artificial nucleic acid molecule. This means that the 3'UTR element in the sense of the present invention may be comprised in the 3'UTR of an artificial
20 nucleic acid sequence, such as an artificial mRNA, or which codes for a 3'UTR of an artificial nucleic acid molecule.

Preferably, the 3'UTR element and the at least one open reading frame are heterologous.
25 For example, the artificial nucleic acid molecule may consist of at least two sequence parts that are derivable from two different genes, the 5'UTR element which is derivable from a TOP gene and the open reading frame and the 3'UTR which may be derivable from the gene encoding the desired protein product. More preferably, the artificial nucleic molecule consist of three sequence parts that are derivable from three different genes: the 5'UTR
30 element which is derivable from a TOP gene, the open reading frame which is derivable from the gene encoding the desired gene product and the 3'UTR element which may be derivable from a gene that relates to an mRNA with an enhanced half-life, for example a 3'UTR element as defined and described below.

Preferably, the at least one 3'UTR element is functionally linked to the ORF. This means preferably that the 3'UTR element is associated with the ORF such that it may exert a function, such as a stabilizing function on the expression of the ORF or a stabilizing
5 function on the artificial nucleic acid molecule. Preferably, the ORF and the 3'UTR element are associated in 5'→3' direction. Thus, preferably, the artificial nucleic acid molecule comprises the structure 5'-ORF-(optional)linker-3'UTR element-3', wherein the linker may be present or absent. For example, the linker may be one or more nucleotides, such as a stretch of 1-50 or 1-20 nucleotides, e.g., comprising or consisting of one or more restriction
10 enzyme recognition sites (restriction sites).

Preferably, the at least one 5'UTR element and the at least one 3'UTR element are functionally linked to the ORF. This means preferably that the 5'UTR element and the 3'UTR element are associated with the ORF such that they may exert a function, preferably
15 in an additive, more preferably in a synergistic manner, such as a stabilizing function on the expression of the ORF, a protein production increasing function for the protein encoded by the ORF, or a stabilizing function on the artificial nucleic acid molecule. Preferably, the 5'UTR element, the ORF, and the 3'UTR element are associated in 5'→3' direction. Thus, preferably, the artificial nucleic acid molecule comprises the structure 5'-5'UTR element-
20 (optional)linker-ORF-(optional)linker-3'UTR element-3', wherein the linker may be present or absent. For example, the linker may be one or more nucleotides, such as a stretch of 1-50 or 1-20 nucleotides, e.g., comprising or consisting of one or more restriction enzyme recognition sites (restriction sites).

25 In a particularly preferred embodiment, the 5'UTR element and the 3'UTR element are heterologous, e.g. preferably the 5'UTR and the 3'UTR are derived from different genes of the same or of different species. Preferably, the 3'UTR is not derived from the TOP gene the 5'UTR is derived from.

30 In a preferred embodiment, the 3'UTR element is chosen such that it exerts at least an additive, preferably a synergistic function with the 5'UTR element on the protein production from the ORF of the artificial nucleic acid molecule. Preferably, the protein production is increased in at least an additive, preferably a synergistic way by the 3'UTR element and the

5'UTR element. Thus, the protein amount of the protein encoded by the ORF, such as a reporter protein, e.g. luciferase, at a certain time point after initiation of expression of the ORF, e.g. after transfection of a test cell or cell line, is preferably at least the same, preferably higher than what would be expected if the protein production increasing effects of the 3'UTR element and the 5'UTR element were purely additive. The additive, preferably synergistic effect may, for example, be determined by the following assay. Four artificial nucleic acid molecules, e.g. mRNAs, comprising an ORF encoding, e.g. a reporter protein such as luciferase, are generated, i.e. (i) lacking UTR elements (E0), (ii) containing a 5'UTR element derived from a 5'UTR of a TOP gene or of a variant thereof (E1), (iii) containing a test 3'UTR element (E2), and (iv) containing both the 5'UTR element and the test 3'UTR element (E1E2). Expression of the ORF contained in the artificial nucleic acid molecules is initiated, for example, by transfecting a test cell line, such as a mammalian cell line, e.g. HELA cells, or primary cells, e.g. HDF cells. Samples are taken at specific time points after initiation of expression, for example, after 6 hours, 24 hours, 48 hours, and 72 hours and the amount of protein produced by expression of the ORF contained in the artificial nucleic acid molecules is measured, for example, by an ELISA assay or a luciferase test, depending on the type of protein encoded by the ORF. The predicted amount of protein at a certain time point after initiation of expression obtained by construct E1E2 if the effects of the 3'UTR element and the 5'UTR element were purely additive (PPA) may be calculated as follows:

$$PPA_x = (E1_x - E0_x) + (E2_x - E0_x) + E0_x$$

E0 is the amount of protein obtained for the construct E0 (lacking UTRs), E1 is the amount of protein obtained for the construct E1, E2 is the protein amount obtained for the construct E2, and x is the time point after initiation of expression. The effect on increasing protein production is additive if $E1E2_x = PPA_x$ and synergistic in the sense of the present invention if $E1E2_x > PPA_x$, wherein $E1E2_x$ is the amount of protein obtained from construct E1E2 at time point x. Preferably, E1E2 is at least 1.0, preferably at least 1.1, more preferably at least 1.3, more preferably at least 1.5, even more preferably at least 1.75 times PPA at a given time point post initiation of expression, such as 24 hours, 48 hours or 72 hours post initiation of expression.

Thus, in a preferred embodiment, the present invention provides an artificial nucleic acid molecule comprising (a.) at least one 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene; (b.) at least one open reading frame (ORF); and
5 (c.) at least one 3'UTR element, wherein the 3'UTR element and the 5'UTR element act at least additively, preferably synergistically to increase protein production from the ORF, preferably wherein $E1E2 \geq PPA$, preferably $E1E2$ is at least 1.0 times PPA, preferably $E1E2$ is at least 1.1 times PPA, more preferably $E1E2$ is at least 1.3 times PPA, even more preferably $E1E2$ is at least 1.5 times PPA at a given time point post initiation of expression of the ORF,
10 for example 24 hours, preferably 48 hours post initiation of expression, such as post transfection, wherein $E1E2$ and PPA are as described above.

Furthermore, it is preferred that the 3'UTR element and the 5'UTR element have at least an additive, preferably a synergistic effect on the total protein production from the artificial
15 nucleic acid molecule in a certain time span, such as within 24 hours, 48 hours, or 72 hours post initiation of expression. The additive or the synergistic effect may be determined as described above, with the difference that the area under the curve (AUC) for the amount of protein over time predicted for $E1E2$ if the effects were purely additive is compared to the actual AUC measured for $E1E2$.

20

In a preferred embodiment, the 3'UTR element comprises or consists of a nucleic acid sequence which is derived from the 3'UTR of a stable mRNA or from a variant of the 3'UTR of a stable mRNA. Thus, in a preferred embodiment, the 3'UTR element comprises or consists of a sequence which is derived from a gene providing a stable mRNA or from a
25 variant of a 3'UTR of a gene providing a stable mRNA. The term "stable mRNA", preferably refers to mRNAs which exhibit a longer half-life in mammalian cells than the average half-life of mRNA molecules in mammalian cells. Preferably, a stable mRNA in the sense of the present application refers to an mRNA which exhibits a half-life of more than 5 hours, preferably more than 8 hours, in a mammalian cell, such as in a mammalian cell line, e.g.
30 in HELA cells, or in primary cells, e.g. in HDF cells, preferably determined by using a transcription inhibitor such as actinomycin D.

For example, the half-life of an mRNA in mammalian cells, such as HELA or HDF cells, may be determined by culturing the cells in presence of a transcription inhibitor, e.g. actinomycin D, 5,6-dichloro-1- β -D-ribofuranosylbenzimidazole (DRB), or α -amanitin, harvesting the cells at different time points after inhibition of transcription, and determining
5 the amount of the mRNA present in the cell samples by methods well known to the person skilled in the art, e.g. by quantitative RT-PCR. The half-life of a particular mRNA may be calculated based on the amounts of the particular mRNA measured at the different time points post inhibition of transcription. Alternatively, pulse-chase methods, e.g. using radioactively labelled nucleotides, or constructs comprising inducible promoters may be
10 used for determining the half-life of an mRNA in mammalian cells.

It is particularly preferred that the enhanced stability of a stable mRNA in the sense of the present invention is affected by its 3'UTR. Thus, preferably, the 3'UTR element comprises or consists of a nucleic acid sequence which is derived from the 3'UTR of a stable mRNA
15 which exhibits a half-life of more than 5 hours, preferably more than 8 hours, in a mammalian cell, such as in a mammalian cell line, e.g. in HeLa cells, or in mammalian primary cells, e.g. in HDF cells, preferably determined by using a transcription inhibitor such as actinomycin D, wherein the enhanced stability of said stable mRNA is effected by its 3'UTR. The ability of a 3'UTR for enhancing stability may be tested as described herein,
20 e.g. by using a reporter open reading frame such as a luciferase encoding open reading frame. Alternatively, an artificial construct encoding the test stable mRNA may be generated, wherein the 3'UTR of the stable mRNA is replaced with a reference 3'UTR, such as a 3'UTR of a short lived mRNA, e.g. a Myc 3'UTR. The stability of the wild type stable mRNA and the 3'UTR modified mRNA may be determined as described above. In the event
25 the 3'UTR modified mRNA exhibits a shorter half-life than the wild type stable mRNA, it may be concluded that a stability enhancing effect is exerted by the 3'UTR of the stable mRNA.

In a particularly preferred embodiment, the 3'UTR element comprises or consists of a
30 nucleic acid sequence which is derived from a 3'UTR of a gene selected from the group consisting of an albumin gene, an α -globin gene, a β -globin gene, a tyrosine hydroxylase gene, a lipoxxygenase gene, and a collagen alpha gene, such as a collagen alpha 1(I) gene, or from a variant of a 3'UTR of a gene selected from the group consisting of an albumin

gene, an α -globin gene, a β -globin gene, a tyrosine hydroxylase gene, a lipoxygenase gene, and a collagen alpha gene, such as a collagen alpha 1(I) gene. In a particularly preferred embodiment, the 3'UTR element comprises or consists of a nucleic acid sequence which is derived from a 3'UTR of an albumin gene, preferably a vertebrate albumin gene, more preferably a mammalian albumin gene, most preferably a human albumin gene. In another particularly preferred embodiment, the 3'UTR element comprises or consists of a nucleic acid sequence which is derived from a 3'UTR of an α -globin gene, preferably a vertebrate α -globin gene, more preferably a mammalian α -globin gene, most preferably a human α -globin gene. For example, the 3'UTR element may comprise or consist of the center, α -complex-binding portion of the 3'UTR of an α -globin gene, such as of a human α -globin gene.

Preferably, the at least one 3'UTR element comprises or consists of a nucleic acid sequence which is derived from the 3'UTR of a vertebrate albumin gene, a vertebrate α -globin gene, a vertebrate β -globin gene, a vertebrate tyrosine hydroxylase gene, a vertebrate lipoxygenase gene, and a vertebrate collagen alpha gene, such as a vertebrate collagen alpha 1(I) gene, or from a variant thereof, preferably from the 3'UTR of a mammalian albumin gene, a mammalian α -globin gene, a mammalian β -globin gene, a mammalian tyrosine hydroxylase gene, a mammalian lipoxygenase gene, and a mammalian collagen alpha gene, such as a mammalian collagen alpha 1(I) gene, or from a variant thereof, more preferably from the 3'UTR of a human albumin gene, a human α -globin gene, a human β -globin gene, a human tyrosine hydroxylase gene, a human lipoxygenase gene, and a human collagen alpha gene, such as a human collagen alpha 1(I) gene, or from a variant thereof, even more preferably from the 3'UTR of the human albumin gene according to GenBank Accession number NM_000477.5 or from a variant thereof. In a preferred embodiment, the 3'UTR element is not derived from the 3'UTR of a *Xenopus* albumin gene. Preferably, the 3'UTR element does not comprise a poly(A) limiting element B (PLEB) of a 3'UTR from a *Xenopus* albumin gene. Preferably, the 3'UTR element does not consist of a PLEB of a 3'UTR from a *Xenopus* albumin gene.

Preferably, the 3'UTR element and the at least one open reading frame are heterologous, e.g. preferably the 3'UTR element and the ORF are derived from different genes of the same or of different species. Preferably, the ORF does not encode an α -globin protein if the

3'UTR element is derived from an α -globin gene. Preferably, the ORF does not encode a β -globin protein if the 3'UTR element is derived from a β -globin gene. Preferably, the ORF does not encode an albumin protein if the 3'UTR element is derived from an albumin gene. Preferably, the ORF does not encode a tyrosine hydroxylase protein if the 3'UTR element is derived from a tyrosine hydroxylase gene. Preferably, the ORF does not encode a lipoxygenase protein if the 3'UTR element is derived from a lipoxygenase gene. Preferably, the ORF does not encode a collagen alpha protein if the 3'UTR element is derived from a collagen alpha gene. Preferably, the ORF does not code for a protein selected from the group consisting of albumin proteins, growth hormones, e.g. human growth hormone (hGH), α -globin proteins, β -globin proteins, tyrosine hydroxylase proteins, lipoxygenase proteins, and collagen alpha proteins. Furthermore, it is preferred that the open reading frame does not code for a reporter protein, e.g., selected from the group consisting of globin proteins, in particular beta-globin,, luciferase protein, GFP proteins, e.g. EGFP, or variants thereof, for example, variants exhibiting at least 70% sequence identity to a globin protein, a luciferase protein, or a GFP protein.

The term 'a nucleic acid sequence which is derived from the 3'UTR of a [...] gene' preferably refers to a nucleic acid sequence which is based on the 3'UTR sequence of a [...] gene or on a part thereof, such as on the 3'UTR of an albumin gene, an α -globin gene, a β -globin gene, a tyrosine hydroxylase gene, a lipoxygenase gene, or a collagen alpha gene, such as a collagen alpha 1(I) gene, preferably of an albumin gene or on a part thereof. This term includes sequences corresponding to the entire 3'UTR sequence, i.e. the full length 3'UTR sequence of a gene, and sequences corresponding to a fragment of the 3'UTR sequence of a gene, such as an albumin gene, α -globin gene, β -globin gene, tyrosine hydroxylase gene, lipoxygenase gene, or collagen alpha gene, such as a collagen alpha 1(I) gene, preferably of an albumin gene. A fragment in this context preferably consists of a continuous stretch of nucleotides corresponding to a continuous stretch of nucleotides in the full-length 3'UTR, which represents at least 20%, preferably at least 30%, more preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, even more preferably at least 70%, even more preferably at least 80%, and most preferably at least 90% of the full-length 3'UTR. Such a fragment, in the sense of the present invention, is preferably a functional fragment as described herein. The term '3'UTR of a [...] gene' preferably refers to the 3'UTR of a naturally occurring gene, such as of a naturally occurring

albumin gene, α -globin gene, β -globin gene, tyrosine hydroxylase gene, lipoxygenase gene, or collagen alpha gene, such as a collagen alpha 1(I) gene, preferably of a naturally occurring albumin gene.

- 5 The terms 'variant of the 3'UTR of a [...] gene' and 'variant thereof' in the context of a 3'UTR refers to a variant of the 3'UTR of a naturally occurring gene, such as a naturally occurring albumin gene, a naturally occurring α -globin gene, a naturally occurring β -globin gene, a naturally occurring tyrosine hydroxylase gene, a naturally occurring lipoxygenase gene, or a naturally occurring collagen alpha gene, such as a naturally occurring collagen
- 10 alpha 1(I) gene, preferably to a variant of the 3'UTR of a vertebrate albumin gene, a vertebrate α -globin gene, a vertebrate β -globin gene, a vertebrate tyrosine hydroxylase gene, a vertebrate lipoxygenase gene, and a vertebrate collagen alpha gene, such as a vertebrate collagen alpha 1(I) gene, preferably to a variant of the 3'UTR of a mammalian albumin gene, a mammalian α -globin gene, a mammalian β -globin gene, a mammalian
- 15 tyrosine hydroxylase gene, a mammalian lipoxygenase gene, and a mammalian collagen alpha gene, such as a mammalian collagen alpha 1(I) gene, more preferably to a variant of the 3'UTR of a human albumin gene, a human α -globin gene, a human β -globin gene, a human tyrosine hydroxylase gene, a human lipoxygenase gene, and a human collagen alpha gene, such as a human collagen alpha 1(I) gene. Such variant may be a modified
- 20 3'UTR of a gene. For example, a variant 3'UTR may exhibit one or more nucleotide deletions, insertions, additions and/or substitutions compared to the naturally occurring 3'UTR from which the variant is derived. Preferably, a variant of a 3'UTR is at least 40%, preferably at least 50%, more preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, even more preferably at least 90%, most preferably at least
- 25 95% identical to the naturally occurring 3'UTR the variant is derived from. Preferably, the variant is a functional variant as described herein.

- The term 'a nucleic acid sequence which is derived from a variant of the 3'UTR of a [...] gene' preferably refers to a nucleic acid sequence which is based on a variant of the 3'UTR
- 30 sequence of a gene, such as on a variant of the 3'UTR of an albumin gene, an α -globin gene, a β -globin gene, a tyrosine hydroxylase gene, a lipoxygenase gene, or a collagen alpha gene, such as a collagen alpha 1(I) gene, or on a part thereof as described above. This term includes sequences corresponding to the entire sequence of the variant of the 3'UTR of

a gene, i.e. the full length variant 3'UTR sequence of a gene, and sequences corresponding to a fragment of the variant 3'UTR sequence of a gene. A fragment in this context preferably consists of a continuous stretch of nucleotides corresponding to a continuous stretch of nucleotides in the full-length variant 3'UTR, which represents at least 20%, preferably at least 30%, more preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, even more preferably at least 70%, even more preferably at least 80%, and most preferably at least 90% of the full-length variant 3'UTR. Such a fragment of a variant, in the sense of the present invention, is preferably a functional fragment of a variant as described herein.

The terms 'functional variant', 'functional fragment', and 'functional fragment of a variant' (also termed 'functional variant fragment') in the context of the present invention, mean that the fragment of the 5'UTR or the 3'UTR, the variant of the 5'UTR or the 3'UTR, or the fragment of a variant of the 5'UTR or the 3'UTR of a gene fulfils at least one, preferably more than one, function of the naturally occurring 5'UTR or 3'UTR of the gene of which the variant, the fragment, or the fragment of a variant is derived. Such function may be, for example, stabilizing mRNA and/or stabilizing and/or prolonging protein production from an mRNA and/or increasing protein production from an mRNA, preferably in a mammalian cell, such as in a human cell. It is particularly preferred that the variant, the fragment, and the variant fragment in the context of the present invention fulfil the function of stabilizing an mRNA, preferably in a mammalian cell, such as a human cell, compared to an mRNA comprising a reference 5'UTR and/or a reference 3'UTR or lacking a 5'UTR and/or a 3'UTR, and/or the function of stabilizing and/or prolonging protein production from an mRNA, preferably in a mammalian cell, such as in a human cell, compared to an mRNA comprising a reference 5'UTR and/or a reference 3'UTR or lacking a 5'UTR and/or a 3'UTR, and/or the function of increasing protein production from an mRNA, preferably in a mammalian cell, such as in a human cell, compared to an mRNA comprising a reference 5'UTR and/or a reference 3'UTR or lacking a 5'UTR and/or a 3'UTR. A reference 3'UTR may be, for example, a 3'UTR naturally occurring in combination with the ORF. Furthermore, a functional variant, a functional fragment, or a functional variant fragment of a 5'UTR or of a 3'UTR of a gene preferably does not have a substantially diminishing effect on the efficiency of translation of the mRNA which comprises such variant of a 5'UTR and/or such variant of a 3'UTR compared to the wild type 5'UTR and/or 3'UTR from which

the variant is derived. A particularly preferred function of a "functional fragment", a "functional variant" or a "functional fragment of a variant" of the 3'UTR of a gene, such as an albumin gene, α -globin gene, β -globin gene, tyrosine hydroxylase gene, lipoxygenase gene, or collagen alpha gene, such as a collagen alpha 1(I) gene, in the context of the present invention is the stabilization and/or prolongation of protein production by expression of an mRNA carrying the functional fragment, functional variant or functional fragment of a variant as described above. A particularly preferred function of a "functional fragment", a "functional variant" or a "functional fragment of a variant" of the 5'UTR in the context of the present invention is the protein production increasing function.

Preferably, the efficiency of the one or more functions exerted by the functional variant, the functional fragment, or the functional variant fragment, such as mRNA and/or protein production stabilizing efficiency and/or the protein production increasing efficiency, is at least 40%, more preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, even more preferably at least 80%, most preferably at least 90% of the mRNA and/or protein production stabilizing efficiency and/or the protein production increasing efficiency exhibited by the naturally occurring 5'UTR and/or 3'UTR of which the variant, the fragment or the variant fragment is derived.

In the context of the present invention, a fragment or part of the 3'UTR of a gene, such as an albumin gene, α -globin gene, β -globin gene, tyrosine hydroxylase gene, lipoxygenase gene, or collagen alpha gene, such as a collagen alpha 1(I) gene, or of a variant thereof preferably exhibits a length of at least about 40 nucleotides, preferably of at least about 50 nucleotides, preferably of at least about 75 nucleotides, more preferably of at least about 100 nucleotides, even more preferably of at least about 125 nucleotides, most preferably of at least about 150 nucleotides. Preferably, such fragment of the 3'UTR of a gene or of a variant of the 3'UTR of a gene is a functional fragment as described above.

In the context of the present invention, a fragment or part of the 5'UTR of a TOP gene or of a variant thereof preferably exhibits a length of at least about 20 nucleotides, preferably of at least about 30 nucleotides, more preferably of at least about 50 nucleotides. Preferably, such fragment of the 5'UTR of a TOP gene or of a variant of the 5'UTR of a TOP gene is a functional fragment as described above.

In some embodiments, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention comprises or consists of a "functional fragment", a "functional variant" or a "functional fragment of a variant" of the 3'UTR of a gene, such as of an albumin gene, α -globin gene, β -globin gene, tyrosine hydroxylase gene, lipoxygenase gene, or collagen alpha gene, such as a collagen alpha 1(I) gene, or of a variant thereof.

In some embodiments, the at least one 5'UTR element of the artificial nucleic acid molecule according to the present invention comprises or consists of a "functional fragment", a "functional variant" or a "functional fragment of a variant" of the 5'UTR of a TOP gene.

Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention increases the stability of the artificial nucleic acid molecule, e.g. increases the stability of an mRNA according to the present invention, compared to a respective mRNA (reference mRNA) lacking a 3'UTR element or comprising a reference 3'UTR element, such as a 3'UTR naturally occurring in combination with the ORF. Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention increases the stability of protein production from the artificial nucleic acid molecule according to the present invention, e.g. from an mRNA according to the present invention, compared to a respective mRNA lacking a 3'UTR element or comprising a reference 3'UTR element, such as a 3'UTR naturally occurring in combination with the ORF. Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention prolongs protein production from the artificial nucleic acid molecule according to the present invention, e.g. from an mRNA according to the present invention, compared to a respective mRNA lacking a 3'UTR element or comprising a reference 3'UTR element, such as a 3'UTR naturally occurring in combination with the ORF. Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention increases the protein production from the artificial nucleic acid molecule according to the present invention, e.g. from an mRNA according to the present invention, compared to a respective mRNA lacking a 3'UTR element or comprising a reference 3'UTR element, such as a 3'UTR naturally occurring in combination with the ORF. Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention does not negatively influence

translational efficiency of an mRNA compared to the translational efficiency of a respective mRNA lacking a 3'UTR element or comprising a reference 3'UTR element, such as a 3'UTR naturally occurring in combination with the ORF. The term 'respective mRNA' in this context means that - apart from the different 3'UTR - the reference mRNA is comparable, preferably identical, to the mRNA comprising the 3'UTR element.

Preferably, the at least one 5'UTR element of the artificial nucleic acid molecule according to the present invention increases the stability of the artificial nucleic acid molecule, e.g. increases the stability of an mRNA according to the present invention, compared to a respective mRNA (reference mRNA) lacking a 5'UTR element or comprising a reference 5'UTR element, such as a 5'UTR naturally occurring in combination with the ORF. Preferably, the at least one 5'UTR element of the artificial nucleic acid molecule according to the present invention increases protein production from the artificial nucleic acid molecule according to the present invention, e.g. from an mRNA according to the present invention, compared to a respective mRNA lacking a 5'UTR element or comprising a reference 5'UTR element, such as a 5'UTR naturally occurring in combination with the ORF. The term 'respective mRNA' in this context means that - apart from the different 5'UTR - the reference mRNA is comparable, preferably identical, to the mRNA comprising the inventive 5'UTR element.

Preferably, the at least one 5'UTR element and the at least one 3'UTR element act synergistically to increase protein production from the artificial nucleic acid molecule according to the present invention, e.g. from an mRNA according to the present invention, as described above.

The term 'stabilizing and/or prolonging protein production from an mRNA' preferably means that the protein production from the mRNA is stabilized and/or prolonged compared to the protein production from a reference mRNA, e.g. comprising a reference 3'UTR element or lacking a 3'UTR element.

'Stabilized protein expression' in this context preferably means that there is more uniform protein production from the artificial nucleic acid molecule according to the present invention over a predetermined period of time, such as over 24 hours, more preferably over

48 hours, even more preferably over 72 hours, when compared to a reference nucleic acid molecule, for example, an mRNA comprising a reference 3'UTR element or lacking a 3'UTR element. Thus, the level of protein production, e.g. in a mammalian system, from the artificial nucleic acid molecule comprising a 3'UTR element according to the present invention, e.g. from an mRNA according to the present invention, preferably does not drop to the extent observed for a reference nucleic acid molecule, such as a reference mRNA as described above. For example, the amount of a protein (encoded by the ORF) observed 6 hours after initiation of expression, e.g. 6 hours post transfection of the artificial nucleic acid molecule according to the present invention into a cell, such as a mammalian cell, may be comparable to the amount of protein observed 48 hours after initiation of expression, e.g. 48 hours post transfection. Thus, the ratio of the amount of protein encoded by the ORF, such as of a reporter protein, e.g., luciferase, observed at 48 hours post initiation of expression, e.g. 48 hours post transfection, to the amount of protein observed 6 hours after initiation of expression, e.g. 6 hours post transfection, is preferably above 0.4, preferably above 0.5, more preferably above 0.6, even more preferably above 0.7, e.g. between about 0.4 and about 4, preferably between about 0.65 and about 3, more preferably between about 0.7 and about 2 for a nucleic acid molecule according to the present invention. For a respective reference nucleic acid molecule, e.g. an mRNA comprising a reference 3'UTR element or lacking a 3'UTR element, said ratio may be, e.g. between about 0.05 and about 0.3. Thus, the present invention provides an artificial nucleic acid molecule comprising an ORF and a 3'UTR element as described above, wherein the ratio of the (reporter) protein amount observed 48 hours after initiation of expression to the (reporter) protein amount observed 6 hours after initiation of expression, preferably in a mammalian expression system, such as in mammalian cells, is preferably above 0.4, preferably above 0.5, more preferably above 0.6, even more preferably above 0.7, e.g. between about 0.4 and about 4, preferably between about 0.65 and about 3, more preferably between about 0.7 and about 2.

'Increased protein expression' in the context of the present invention may refer to increased protein expression at one time point after initiation of expression compared to a reference molecule or to an increased total protein production within a certain time period after initiation of expression. Thus, the protein level observed at a certain time point after initiation of expression, e.g. after transfection, of the artificial nucleic acid molecule

according to the present invention, e.g. after transfection of an mRNA according to the present invention, for example, 24, 48, or 72 hours post transfection, or the total protein produced in a time span of, e.g. 24, 48 or 72 hours, is preferably higher than the protein level observed at the same time point after initiation of expression, e.g. after transfection, or the total protein produced within the same time span, for a reference nucleic acid molecule, such as a reference mRNA comprising a reference 5' and/or a reference 3'UTR or lacking a 5'UTR element and/or 3'UTR element. As set forth above, it is a particularly preferred function of the 5'UTR element to affect the increase in protein production from the artificial nucleic acid molecule. Preferably, the increase in protein production effected by the 5'UTR element compared to a reference nucleic acid molecule lacking such 5'UTR element at a given time point post initiation of expression is at least 1.5-fold, more preferably at least 2-fold, more preferably at least 3-fold, even more preferably at least 4-fold, most preferably at least 5-fold of the protein production observed for a reference nucleic acid molecule lacking the 5'UTR element. The same holds preferably for the total protein production in a given time period, for example in a time period of 24, 48 or 72 hours post initiation of expression.

Said increase in stability of the artificial nucleic acid molecule, said increase in stability of protein production, said prolongation of protein production and/or said increase in protein production is preferably determined by comparison with a respective reference nucleic acid molecule lacking a 5'UTR element and/or a 3'UTR element, e.g. an mRNA lacking a 5'UTR element and/or a 3'UTR element, or a reference nucleic acid molecule comprising a reference 5'UTR element and/or a reference 3'UTR element, such as a 3'UTR and/or a 5'UTR naturally occurring with the ORF or a 5'UTR and/or a 3'UTR of a reference gene.

The mRNA and/or protein production stabilizing effect and efficiency and/or the protein production increasing effect and efficiency of the variants, fragments and/or variant fragments of the 3'UTR of an albumin gene as well as the mRNA and/or protein production stabilizing effect and efficiency and/or the protein production increasing effect and efficiency of the at least one 3'UTR element, the at least one 5'UTR element, or the at least one 3'UTR element and the at least one 5'UTR element of the artificial nucleic acid molecule according to the present invention may be determined by any method suitable for this purpose known to the skilled person. For example, artificial mRNA molecules may be

generated comprising a coding sequence for a reporter protein, such as luciferase, and no 3'UTR and/or no 5'UTR, a 5'UTR element derived from a TOP gene and/or a 3'UTR element derived from a gene as described above, a 5'UTR element derived from a reference gene and/or a 3'UTR derived from a reference gene (i.e., a reference 3'UTR element or a reference 5'UTR element, such as a 5'UTR or a 3'UTR naturally occurring with the ORF), as 3'UTR a variant of a 3'UTR of a gene as described above, as 3'UTR a fragment of a 3'UTR of a gene as described above, or as 3'UTR a fragment of a variant of a 3'UTR of a gene as described above, as 5'UTR a variant of a 5'UTR of a TOP gene, as 5'UTR a fragment of a 5'UTR of a TOP gene, or as 5'UTR a fragment of a variant of a 5'UTR of a TOP gene. Such mRNAs may be generated, for example, by *in vitro* transcription of respective vectors such as plasmid vectors, e.g. comprising a T7 promoter and a sequence encoding the respective mRNA sequences. The generated mRNA molecules may be transfected into cells by any transfection method suitable for transfecting mRNA, for example they may be electroporated into mammalian cells, such as HELA or HDF cells, and samples may be analyzed certain time points after transfection, for example, 6 hours, 24 hours, 48 hours, and 72 hours post transfection. Said samples may be analyzed for mRNA quantities and/or protein quantities by methods well known to the skilled person. For example, the quantities of reporter mRNA present in the cells at the sample time points may be determined by quantitative PCR methods. The quantities of reporter protein encoded by the respective mRNAs may be determined, e.g., by ELISA assays or reporter assays such as luciferase assays depending on the reporter protein used. The effect of stabilizing protein expression and/or prolonging protein expression may be, for example, analyzed by determining the ratio of the protein level observed 48 hours post transfection and the protein level observed 6 hours post transfection. The closer said value is to 1, the more stable the protein expression is within this time period. Said value may also be above 1 if the protein level is higher at the later time point. Such measurements may of course also be performed at 72 or more hours and the ratio of the protein level observed 72 hours post transfection and the protein level observed 6 hours post transfection may be determined to determine stability of protein expression.

Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about

60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99%, most preferably of 100% to a nucleic acid sequence selected from SEQ ID No. 1369-1377, 1391, 1392, and 1393 and wherein the variants of the sequences according to SEQ ID No. 1369-1377, 1391, 1392 and 1393 are preferably functional variants as described above.

The at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention may also comprise or consist of a fragment of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99%, most preferably of 100% to the nucleic acid sequence according to SEQ ID No. 1369-1377, 1391, 1392, or 1393 wherein the fragment is preferably a functional fragment or a functional variant fragment as described above. Preferably, the fragment is as described above, i.e. being a continuous stretch of nucleotides representing at least 20% etc. of the full-length 3'UTR the fragment is derived from. Such fragment preferably exhibits a length of at least about 40 nucleotides, preferably of at least about 50 nucleotides, preferably of at least about 75 nucleotides, more preferably of at least about 100 nucleotides, even more preferably of at least about 125 nucleotides, most preferably of at least about 150 nucleotides.

For example, such fragment may exhibit a nucleic acid sequence according to SEQ ID Nos. 1378-1390, such as

AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC
ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT
TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATT (SEQ ID No. 1378)

CATCACATTT AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA
AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG (SEQ ID No.
1379)

AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC
ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC (SEQ ID No.
1380)

5 CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT
CTTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT (SEQ ID No.
1381)

TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT
10 GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT (SEQ ID No.
1382)

AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC
CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT (SEQ ID No.
15 1383)

TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG
TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT (SEQ ID No.
1384)

20 AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA (SEQ ID No.
1385)

25 ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT
TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA AAAATGGAAA (SEQ ID No.
1386)

CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT
30 CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT
TGCCTCTTTT CTCTGTGCTT CAATTAATAA A (SEQ ID No. 1387)

TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG
TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA
A (SEQ ID No. 1388)

5 CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT
TGCCTCTTTT CTCTGTGCTT CAATTAATAA A (SEQ ID No. 1389)

AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
(SEQ ID No. 1390)

10

or the corresponding RNA sequence, or a nucleic acid sequence which is at least 40%, preferably at least about 50%, preferably at least about 60%, preferably at least about 70%, more preferably at least about 80%, more preferably at least about 90%, even more preferably at least about 95%, even more preferably at least about 99% identical to said
15 nucleic acid sequences or the corresponding RNA sequence. Thus, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention may comprise or consist of a nucleic acid fragment as described above. Obviously, the
thymidine nucleotides comprised in the fragments according to SEQ ID Nos. 1378-1390 may be replaced by uridine nucleotides.

20

Preferably, said variants, fragments or variant fragments are functional variants, functional fragments, or functional variant fragments as described above, exhibiting at least one function of the nucleic acid sequence according to SEQ ID Nos. 1369-1377, 1391, 1392, or 1393 such as stabilization of the artificial nucleic acid molecule according to the invention,
25 stabilizing and/or prolonging protein expression from the artificial nucleic acid molecule according to the invention, and/or increasing protein production, preferably with an efficiency of at least 40%, more preferably of at least 50%, more preferably of at least 60%, even more preferably of at least 70%, even more preferably of at least 80%, most preferably of at least 90% of the stabilizing efficiency and/or protein production increasing efficiency
30 exhibited by the nucleic acid sequence according to SEQ ID Nos. 1369-1377, 1391, 1392, or 1393. Preferably, variants, fragments or variant fragments are functional variants, functional fragments, or functional variant fragments exhibit the function of acting synergistically with the 5'UTR element to increase protein production from the artificial nucleic acid molecule.

Preferably, the at least one 3'UTR element of the artificial nucleic acid molecule according to the present invention exhibits a length of at least about 40 nucleotides, preferably of at least about 50 nucleotides, preferably of at least about 75 nucleotides, more preferably of at least about 100 nucleotides, even more preferably of at least about 125 nucleotides, most preferably of at least about 150 nucleotides. For example, the 3'UTR may exhibit a length of about 50 to about 300 nucleotides, preferably of about 100 to about 250 nucleotides, more preferably of about 150 to about 200 nucleotides.

Furthermore, the artificial nucleic acid molecule according to the present invention may comprise more than one 3'UTR elements as described above. For example, the artificial nucleic acid molecule according to the present invention may comprise one, two, three, four or more 3'UTR elements, wherein the individual 3'UTR elements may be the same or they may be different. For example, the artificial nucleic acid molecule according to the present invention may comprise two essentially identical 3'UTR elements as described above, e.g. two 3'UTR elements comprising or consisting of a nucleic acid sequence which is derived from the 3'UTR of an albumin gene or from a variant of the 3'UTR of an albumin gene, such as a nucleic acid sequence according to SEQ ID No. 1369 or 1376, functional variants thereof, functional fragments thereof, or functional variant fragments thereof as described above.

Surprisingly, the inventors found that an artificial nucleic acid molecule comprising a 5'UTR element comprising or consisting of a nucleic acid sequence derived from a TOP gene as described above may represent or may provide an mRNA molecule exhibiting strongly enhanced protein production from said artificial nucleic acid molecule.

The artificial nucleic acid molecule according to the present invention may be RNA, such as mRNA, DNA, such as a DNA vector, or may be a modified RNA or DNA molecule. It may be provided as a double-stranded molecule having a sense strand and an anti-sense strand, for example, as a DNA molecule having a sense strand and an anti-sense strand.

The artificial nucleic acid molecule according to the present invention may further comprise a 5'-cap. The optional 5'-cap is preferably attached to the 5'-side of the 5'UTR element.

In a preferred embodiment, the artificial nucleic acid sequence comprises a 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene encoding a ribosomal protein as described above, for example, encoding a ribosomal Large protein, or from a variant thereof, and a 3'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 3'UTR of an albumin gene or a variant thereof as described above.

In a particularly preferred embodiment, the artificial nucleic acid sequence comprises a 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), an ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, an hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), an androgen-induced 1 gene (AIG1), cytochrome c oxidase subunit VIc gene (COX6C), or a N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, preferably from a vertebrate ribosomal protein Large 32 gene (RPL32), a vertebrate ribosomal protein Large 35 gene (RPL35), a vertebrate ribosomal protein Large 21 gene (RPL21), a vertebrate ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a vertebrate hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a vertebrate androgen-induced 1 gene (AIG1), a vertebrate cytochrome c oxidase subunit VIc gene (COX6C), or a vertebrate N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, more preferably from a mammalian ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), a mammalian ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a mammalian hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a mammalian androgen-induced 1 gene (AIG1), a mammalian cytochrome c oxidase subunit VIc gene (COX6C), or a mammalian N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, most preferably from a human ribosomal protein Large 32 gene (RPL32), a human ribosomal protein Large 35 gene (RPL35), a human ribosomal protein Large 21 gene (RPL21), a human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a human hydroxysteroid (17-beta) dehydrogenase 4 gene

(HSD17B4), a human androgen-induced 1 gene (AIG1), a human cytochrome c oxidase subunit VIc gene (COX6C), or a human N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof, wherein preferably the 5'UTR element does not comprise the 5'TOP of said gene, and a 3'UTR
5 element which comprises or consists of a nucleic acid sequence which is derived from an albumin gene as described above.

In a particularly preferred embodiment, the artificial nucleic acid molecule according to the present invention comprises a 5'UTR element which comprises or consists of a nucleic acid
10 sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99% to the nucleic acid sequence according to SEQ ID No. 1368 or SEQ ID NOs 1412-1420, or a corresponding RNA
15 sequence, and a 3'UTR element which comprises or consist of a nucleic acid sequence which has an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%, even more preferably of at least about 99%, most preferably of 100% to the nucleic acid
20 sequence according to SEQ ID No. 1369, 1376, 1377, 1391, or 1392, e.g., a 5'UTR element which comprises or consists of a nucleic acid sequence which has an identity of at least about 90% to the nucleic acid sequence according to SEQ ID No. 1368 or a corresponding RNA sequence and a 3'UTR element which comprises or consist of a nucleic acid sequence which has an identity of at least about 90% to the nucleic acid sequence
25 according to SEQ ID No. 1369, 1376, 1377, 1391, or 1392.

Preferably, the artificial nucleic acid molecule according to the present invention further comprises a poly(A) sequence and/or a polyadenylation signal. Preferably, the optional poly(A) sequence is located 3' to the ORF or the at least one 3'UTR element, preferably is
30 connected to the 3'-end of the ORF or the 3'UTR element. The connection may be direct or indirect, for example, via a stretch of 2, 4, 6, 8, 10, 20 etc. nucleotides, such as via a linker of 1-50, preferably of 1-20 nucleotides, e.g. comprising or consisting of one or more restriction sites.

In one embodiment, the optional polyadenylation signal is located within the 3'UTR element. Preferably, the polyadenylation signal comprises the consensus sequence NN(U/T)ANA, with N = A or U, preferably AA(U/T)AAA or A(U/T)(U/T)AAA. Such
5 consensus sequence may be recognised by most animal and bacterial cell-systems, for example by the polyadenylation-factors, such as cleavage/polyadenylation specificity factor (CPSF) cooperating with CstF, PAP, PAB2, CFI and/or CFII. Preferably, the polyadenylation signal, preferably the consensus sequence NNUANA, is located less than about 50 nucleotides, more preferably less than about 30 nucleotides, most preferably less than about
10 25 nucleotides, for example 21 nucleotides, upstream of the 3'-end of the 3'UTR element.

Using an appropriate transcription system will then lead to attachment of a poly(A) sequence to the premature-RNA. For example, the inventive artificial nucleic acid molecule may be a DNA molecule comprising a 3'UTR element as described above and a
15 polyadenylation signal, which may result in polyadenylation of an RNA upon transcription of this DNA molecule. Accordingly, a resulting RNA may comprise a combination of the 3'UTR element followed by a poly(A) sequence.

Potential transcription systems are *in vitro* transcription systems or cellular transcription
20 systems etc. Accordingly, transcription of an artificial nucleic acid molecule according to the invention, e.g. transcription of an artificial nucleic acid molecule comprising a 5'UTR element, an open reading frame, a 3'UTR element and a polyadenylation-signal, may result in an mRNA molecule comprising a 5'UTR element, an open reading frame, a 3'UTR element and a poly(A) sequence.

25 The invention also provides an artificial nucleic acid molecule which is an mRNA molecule comprising a, 5'UTR element, an open reading frame, an optional 3'UTR element as described above and a poly(A) sequence.

30 In one embodiment, the invention provides an artificial nucleic acid molecule which is an artificial DNA molecule comprising a 5'UTR element as described above, an open reading frame and optionally a nucleic acid sequence according to any one of SEQ ID Nos. 1369-1377, 1391, and 1392 or a sequence having an identity of at least about 40% or more to a

nucleic acid sequence according to any one of SEQ ID Nos. 1369-1377, 1391, and 1392 or a fragment thereof. Furthermore, the invention provides an artificial nucleic acid molecule which is an artificial RNA molecule comprising a 5'UTR element as described above, an open reading frame and optionally an RNA sequence corresponding to a sequence
5 according to any one of SEQ ID Nos. 1369-1377, 1391, and 1392 or a sequence having an identity of at least about 40 % or more to any one of SEQ ID Nos. 1369-1377, 1391, and 1392, or a fragment thereof.

Accordingly, the invention provides an artificial nucleic acid molecule which may be a
10 template for an RNA molecule, preferably for an mRNA molecule, which is stabilised and optimized with respect to translation efficiency. In other words, the artificial nucleic acid molecule may be a DNA or RNA which may be used for production of an mRNA. The obtainable mRNA, may, in turn, be translated for production of a desired peptide or protein encoded by the open reading frame. If the artificial nucleic acid molecule is a DNA, it may,
15 for example, be used as a double-stranded storage form for continued and repetitive *in vitro* or *in vivo* production of mRNA.

In one embodiment, the artificial nucleic acid molecule according to the present invention further comprises a poly(A) sequence. The length of the poly(A) sequence may vary. For example, the poly(A) sequence may have a length of about 20 adenine nucleotides up to
20 about 300 adenine nucleotides, preferably of about 40 to about 200 adenine nucleotides, more preferably from about 50 to about 100 adenine nucleotides, such as about 60, 70, 80, 90 or 100 adenine nucleotides.

For example, the artificial nucleic acid molecule according to the present invention may
25 comprise a nucleic acid sequence corresponding to the DNA sequence

CATCACATTT AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA
AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA AAAATGGAAA
30 GAATCTAGAT CTAAAAA AAAA AAAAAA AAAAAA AAAAAA AAAAAA
AAAAA (SEQ ID No. 1377).

Transcription of such a sequence may result in an artificial nucleic acid molecule comprising a corresponding RNA sequence.

Such artificial RNA molecule may also be obtainable *in vitro* by common methods of chemical synthesis without being necessarily transcribed from a DNA progenitor.

In a particularly preferred embodiment, the artificial nucleic acid molecule according to the present invention is an RNA molecule, preferably an mRNA molecule comprising in 5'-to-3'-direction a 5'UTR element as described above, an open reading frame, a 3'UTR element as described above and a poly(A) sequence.

In a preferred embodiment, the open reading frame does not code for human albumin, provided that the 3'UTR element is identical to the 3'UTR of human albumin. In some further embodiments, it is preferred that the open reading frame does not code for human albumin according to GenBank Accession number NM_000477.5 provided that the 3'UTR element is identical to the 3'UTR of human albumin. In some further embodiments, it is preferred that the open reading frame does not code for human albumin or variants thereof provided that the 3'UTR element is a sequence which is identical to SEQ ID No. 1369. Furthermore, in some embodiments, it is preferred that the open reading frame does not code for a reporter protein, e.g., selected from the group consisting of globin proteins, luciferase proteins, GFP proteins or variants thereof, for example, variants exhibiting at least 70% sequence identity to a globin protein, a luciferase protein, or a GFP protein.

In some embodiments, it is preferred that the 3'UTR element does not consist of a histone stem-loop, preferably does not comprise a histone stem-loop. In one embodiment, the artificial nucleic acid molecule according to the present invention does not comprise a histone stem-loop. However, in some embodiments, the 3'UTR element of the artificial nucleic acid molecule or the artificial nucleic acid molecule according to the present invention may comprise a histone stem-loop in addition to the nucleic acid sequence derived from the 3'UTR of an albumin gene. Such artificial nucleic acid molecule according to the present invention, for example, may comprise in 5'-to-3'-direction a 5'UTR element, an ORF, a 3'UTR element, preferably comprising a polyadenylation signal, an optional histone stem-loop and an optional poly(A) sequence. It may also comprise in 5'-to-3'-

direction a 5'UTR element as described above, an ORF, a 3'UTR element, e.g. comprising a polyadenylation signal, a poly(A) sequence and an optional histone stem-loop.

In the context of the present invention, such a histone stem-loop is typically derived from a histone gene and comprises an intramolecular base pairing of two neighbored entirely or partially reverse complementary sequences, thereby forming a stem-loop. A stem-loop can occur in single-stranded DNA or, more commonly, in RNA. The structure is also known as a hairpin or hairpin loop and usually consists of a stem and a (terminal) loop within a consecutive sequence, wherein the stem is formed by two neighbored entirely or partially reverse complementary sequences separated by a short sequence as sort of spacer, which builds the loop of the stem-loop structure. The two neighbored entirely or partially reverse complementary sequences may be defined as e.g. stem-loop elements stem1 and stem2. The stem loop is formed when these two neighbored entirely or partially reverse complementary sequences, e.g. stem-loop elements stem1 and stem2, form base-pairs with each other, leading to a double stranded nucleic acid sequence comprising an unpaired loop at its terminal ending formed by the short sequence located between stem-loop elements stem1 and stem2 on the consecutive sequence. The unpaired loop thereby typically represents a region of the nucleic acid which is not capable of base pairing with either of these stem-loop elements. The resulting lollipop-shaped structure is a key building block of many RNA secondary structures. The formation of a stem-loop structure is thus dependent on the stability of the resulting stem and loop regions, wherein the first prerequisite is typically the presence of a sequence that can fold back on itself to form a paired double strand. The stability of paired stem-loop elements is determined by the length, the number of mismatches or bulges it contains (a small number of mismatches is typically tolerable, especially in a long double strand), and the base composition of the paired region. In the context of the present invention, optimal loop length is 3-10 bases, more preferably 3 to 8, 3 to 7, 3 to 6 or even more preferably 4 to 5 bases, and most preferably 4 bases.

An example for a histone stem-loop sequence is the sequence according to SEQ ID NO: 1394 (CAAAGGCTCTTTTCAGAGCCACCA) or the corresponding RNA sequence.

Thus, in some embodiments, the artificial nucleic acid molecule according to the present invention comprises (a.) at least one 5'UTR element as described herein, (b.) at least one open reading frame, and at least one histone-stem loop which may, for example, comprise or consist of a sequence having a sequence identity of at least about 75%, preferably of at least about 80%, preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95% to the sequence according to SEQ ID NO. 1394 or the corresponding RNA sequence, wherein preferably positions 6, 13 and 20 of the sequence having a sequence identity of at least about 75%, preferably of at least about 80%, preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95% to the sequence according to SEQ ID NO. 1394 or the corresponding RNA sequence are conserved, i.e. are identical to the nucleotides at positions 6, 13 and 20 of SEQ ID NO. 1394.

In some embodiments, the artificial nucleic acid molecule comprises further elements such as a 5'-cap, a poly(C) sequence and/or an IRES-motif. A 5'-cap may be added post-transcriptionally to the 5'end of an RNA. Further, the inventive artificial nucleic acid molecule, particularly if the nucleic acid is in the form of an mRNA or codes for an mRNA, may be modified by a sequence of at least 10 cytidines, preferably at least 20 cytidines, more preferably at least 30 cytidines (so-called "poly(C) sequence"). Particularly, the inventive nucleic acid molecule may contain, especially if the nucleic acid is in the form of an (m)RNA or codes for an mRNA, a poly(C) sequence of typically about 10 to 200 cytidine nucleotides, preferably about 10 to 100 cytidine nucleotides, more preferably about 10 to 70 cytidine nucleotides or even more preferably about 20 to 50 or even 20 to 30 cytidine nucleotides.

An internal ribosome entry site (IRES) sequence or IRES-motif may separate several open reading frames, for example if the artificial nucleic acid molecule encodes for two or more peptides or proteins. An IRES-sequence may be particularly helpful if the mRNA is a bi- or multicistronic RNA.

Furthermore, the artificial nucleic acid molecule may comprise additional 5'-elements such as a promoter containing-sequence. The promoter may drive and or regulate transcription of

the artificial nucleic acid molecule according to the present invention, for example of an artificial DNA-molecule according to the present invention.

In preferred embodiments, the invention provides artificial nucleic acid molecules, preferably mRNA molecules, comprising in 5'-to-3'-direction at least one of the following structures

5'-cap – 5'UTR element – ORF – 3'UTR element – histone stem-loop – poly(A) sequence

5'-cap – 5'UTR element – ORF – 3'UTR element – poly(A) sequence – histone stem-loop

10 5'-cap - 5'UTR element – ORF – IRES – ORF – 3'UTR element - histone stem-loop - poly(A) sequence

5'-cap – 5'UTR element – ORF – IRES – ORF – 3'UTR element – poly(A) sequence – histone stem-loop

5'-cap – 5'UTR element – ORF – 3'UTR element – poly(A) sequence – poly(C) sequence

15 5'-cap – 5'UTR element – ORF – 3'UTR element – poly(A) sequence – poly(C) sequence – histone stem-loop

5'-cap - 5'UTR element – ORF – IRES – ORF – 3'UTR element – histone stem-loop – poly(A) sequence – poly(C) sequence

20 Preferably, the artificial nucleic acid molecule, preferably the open reading frame, is at least partially G/C modified. Thus, the inventive artificial nucleic acid molecule may be thermodynamically stabilized by modifying the G (guanosine)/C (cytidine) content of the molecule. The G/C content of the open reading frame of an artificial nucleic acid molecule according to the present invention may be increased compared to the G/C content of the open reading frame of a corresponding wild type sequence, preferably by using the
 25 degeneration of the genetic code. Thus, the encoded amino acid sequence of the nucleic acid molecule is preferably not modified by the G/C modification compared to the coded amino acid sequence of the particular wild type sequence. The codons of a coding sequence or a whole nucleic acid molecule, e.g. an mRNA, may therefore be varied
 30 compared to the wild type coding sequence, such that they include an increased amount of G/C nucleotides while the translated amino acid sequence is maintained. In respect to the fact that several codons code for one and the same amino acid (so-called degeneration of

the genetic code), the most favourable codons for the stability can be determined (so-called alternative codon usage).

Depending on the amino acid to be encoded by the coding region of the inventive nucleic acid molecule as defined herein, there are various possibilities for modification of the nucleic acid sequence, e.g. the open reading frame, compared to its wild type coding region. In the case of amino acids which are encoded by codons which contain exclusively G or C nucleotides, no modification of the codon is necessary. Thus, the codons for Pro (CCC or CCG), Arg (CGC or CCG), Ala (GCC or GCG) and Gly (GGC or GGG) require no modification, since no A or U/T is present.

In contrast, codons which contain A and/or U/T nucleotides may be modified by substitution of other codons which code for the same amino acids but contain no A and/or U/T. For example

the codons for Pro can be modified from CC(U/T) or CCA to CCC or CCG;
the codons for Arg can be modified from CG(U/T) or CGA or AGA or AGG to CGC or CCG;
the codons for Ala can be modified from GC(U/T) or GCA to GCC or GCG;
the codons for Gly can be modified from GG(U/T) or GGA to GGC or GGG.

In other cases, although A or (U/T) nucleotides cannot be eliminated from the codons, it is however possible to decrease the A and (U/T) content by using codons which contain a lower content of A and/or (U/T) nucleotides. Examples of these are:

The codons for Phe can be modified from (U/T)(U/T)(U/T) to (U/T) (U/T)C;
the codons for Leu can be modified from (U/T) (U/T)A, (U/T) (U/T)G, C(U/T) (U/T) or C(U/T)A to C(U/T)C or C(U/T)G;
the codons for Ser can be modified from (U/T)C(U/T) or (U/T)CA or AG(U/T) to (U/T)CC, (U/T)CG or AGC;
the codon for Tyr can be modified from (U/T)A(U/T) to (U/T)AC;
the codon for Cys can be modified from (U/T)G(U/T) to (U/T)GC;
the codon for His can be modified from CA(U/T) to CAC;
the codon for Gln can be modified from CAA to CAG;

the codons for Ile can be modified from A(U/T)(U/T) or A(U/T)A to A(U/T)C;

the codons for Thr can be modified from AC(U/T) or ACA to ACC or ACG;

the codon for Asn can be modified from AA(U/T) to AAC;

the codon for Lys can be modified from AAA to AAG;

5 the codons for Val can be modified from G(U/T)(U/T) or G(U/T)A to G(U/T)C or G(U/T)G;

the codon for Asp can be modified from GA(U/T) to GAC;

the codon for Glu can be modified from GAA to GAG;

the stop codon (U/T)AA can be modified to (U/T)AG or (U/T)GA.

10 In the case of the codons for Met (A(U/T)G) and Trp ((U/T)GG), on the other hand, there is no possibility of sequence modification without altering the encoded amino acid sequence.

The substitutions listed above can be used either individually or in all possible combinations to increase the G/C content of the open reading frame of the inventive nucleic acid
15 sequence as defined herein, compared to its particular wild type open reading frame (i.e. the original sequence). Thus, for example, all codons for Thr occurring in the wild type sequence can be modified to ACC (or ACG).

Preferably, the G/C content of the open reading frame of the inventive artificial nucleic acid molecule as defined herein is increased by at least 7%, more preferably by at least 15%,
20 particularly preferably by at least 20%, compared to the G/C content of the wild type coding region. According to a specific embodiment at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, more preferably at least 70 %, even more preferably at least 80% and most preferably at least 90%, 95% or even 100% of the substitutable codons in the open reading frame of the inventive artificial nucleic acid molecule or a fragment, variant or derivative thereof are
25 substituted, thereby increasing the G/C content of said open reading frame.

In this context, it is particularly preferable to increase the G/C content of the open reading frame of the inventive nucleic acid sequence as defined herein, to the maximum (i.e. 100% of the substitutable codons), compared to the wild type open reading frame.

Furthermore, the open reading frame is preferably at least partially codon-optimized.
30 Codon-optimization is based on the finding that the translation efficiency may be determined by a different frequency in the occurrence of transfer RNAs (tRNAs) in cells. Thus, if

so-called "rare codons" are present in the coding region of the inventive artificial nucleic acid molecule as defined herein, to an increased extent, the translation of the corresponding modified nucleic acid sequence is less efficient than in the case where codons coding for relatively "frequent" tRNAs are present.

- 5 Thus, the open reading frame of the inventive nucleic acid sequence is preferably modified compared to the corresponding wild type coding region such that at least one codon of the wild type sequence which codes for a tRNA which is relatively rare in the cell is exchanged for a codon which codes for a tRNA which is comparably frequent in the cell and carries the same amino acid as the relatively rare tRNA. By this modification, the open reading
- 10 frame of the inventive artificial nucleic acid molecule as defined herein, is modified such that codons for which frequently occurring tRNAs are available may replace codons which correspond to rare tRNAs. In other words, according to the invention, by such a modification all codons of the wild type open reading frame which code for a rare tRNA may be exchanged for a codon which codes for a tRNA which is more frequent in the cell and
- 15 which carries the same amino acid as the rare tRNA. Which tRNAs occur relatively frequently in the cell and which, in contrast, occur relatively rarely is known to a person skilled in the art; cf. e.g. Akashi, Curr. Opin. Genet. Dev. 2001, 11(6): 660-666. Accordingly, preferably, the open reading frame is codon-optimized, preferably with respect to the system in which the nucleic acid molecule according to the present invention is to be expressed, preferably with respect to the system in which the nucleic acid molecule according
- 20 to the present invention is to be translated. Preferably, the codon usage of the open reading frame is codon-optimized according to mammalian codon usage, more preferably according to human codon usage. Preferably, the open reading frame is codon-optimized and G/C-content modified.
- 25 For further improving degradation resistance, e.g. resistance to *in vivo* degradation by an exo- or endonuclease, and/or for further improving protein production from the artificial nucleic acid molecule according to the present invention, the artificial nucleic acid molecule may further comprise modifications, such as backbone modifications, sugar modifications and/or base modifications, e.g., lipid-modifications or the like. Preferably, the tran-
- 30 scription and/or the translation of the artificial nucleic acid molecule according to the present invention is not significantly impaired by said modifications.

Nucleotide analogues/modifications that may be used in the context of the present invention may be selected, for example, from 2-amino-6-chloropurineriboside-5'-triphosphate, 2-aminoadenosine-5'-triphosphate, 2-thiocytidine-5'-triphosphate, 2-thiouridine-5'-triphosphate, 4-thiouridine-5'-triphosphate, 5-aminoallylcytidine-5'-triphosphate, 5-aminoallyluridine-5'-triphosphate, 5-bromocytidine-5'-triphosphate, 5-bromouridine-5'-triphosphate, 5-iodocytidine-5'-triphosphate, 5-iodouridine-5'-triphosphate, 5-methylcytidine-5'-triphosphate, 5-methyluridine-5'-triphosphate, 6-azacytidine-5'-triphosphate, 6-azauridine-5'-triphosphate, 6-chloropurineriboside-5'-triphosphate, 7-deazaadenosine-5'-triphosphate, 7-deazaguanosine-5'-triphosphate, 8-azaadenosine-5'-triphosphate, 8-azidoadenosine-5'-triphosphate, benzimidazole-riboside-5'-triphosphate, N1-methyladenosine-5'-triphosphate, N1-methylguanosine-5'-triphosphate, N6-methyladenosine-5'-triphosphate, O6-methylguanosine-5'-triphosphate, pseudouridine-5'-triphosphate, or puromycin-5'-triphosphate, xanthosine-5'-triphosphate. Particular preference is given to nucleotides for base modifications selected from the group of base-modified nucleotides consisting of 5-methylcytidine-5'-triphosphate, 7-deazaguanosine-5'-triphosphate, 5-bromocytidine-5'-triphosphate, and pseudouridine-5'-triphosphate.

Further, lipid-modified artificial nucleic acid molecules may typically comprise at least one linker which is covalently linked with the artificial nucleic acid molecule, and at least one lipid which is covalently linked with this linker. Alternatively, a lipid-modified artificial nucleic acid molecule may comprise at least one artificial nucleic acid molecule as defined herein and at least one, preferably bifunctional lipid which is covalently linked, preferably without a linker, with that artificial nucleic acid molecule. According to a third alternative, a lipid-modified artificial nucleic acid molecule may comprise an artificial nucleic acid molecule as defined herein, at least one linker which is covalently linked with that artificial nucleic acid molecule, at least one lipid which is covalently linked with this linker, and additionally at least one, preferably bifunctional lipid which is covalently linked, preferably without a linker, with the artificial nucleic acid molecule.

In a further aspect, the present invention provides a vector comprising

- a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene;

- b. at least one open reading frame (ORF) and/or a cloning site; and
- c. optionally, at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene, or from a
5 variant of the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene.

The at least one 5'UTR element, the optional at least one 3'UTR element and the at least one ORF are as described herein for the artificial nucleic acid molecule according to the
10 present invention. The cloning site may be any sequence that is suitable for introducing an open reading frame or a sequence comprising an open reading frame, such as one or more restriction sites. The vector comprising a cloning site is preferably suitable for inserting an open reading frame into the vector 3' to the 5'UTR element, preferably directly 3' to the 5'UTR element. Thus, the vector comprising a cloning site is preferably suitable for inserting
15 an open reading frame into the vector, preferably for inserting an open reading frame between the 5'UTR element and the optional 3'UTR element, preferably 5' to the optional 3'UTR element and 3' to the 5'UTR element. Preferably, the cloning site or the ORF is located 5' to the 3'UTR element, preferably in close proximity to the 5'-end of the 3'UTR element. For example, the cloning site or the ORF may be directly connected to the 5'-end
20 of the 3'UTR element or they may be connected via a stretch of nucleotides, such as by a stretch of 2, 4, 6, 8, 10, 20 etc. nucleotides as described above for the artificial nucleic acid molecule according to the present invention. Preferably the cloning site or the ORF is located 3' to the 5'UTR element, preferably in close proximity to the 3'-end of the 5'UTR element. For example, the cloning site or the ORF may be directly connected to the 3'-end
25 of the 5'UTR element or they may be connected via a stretch of nucleotides, such as by a stretch of 2, 4, 6, 8, 10, 20 etc. nucleotides as described above for the artificial nucleic acid molecule according to the present invention.

Preferably the vector according to the present invention is suitable for producing the
30 artificial nucleic acid molecule according to the present invention, preferably for producing an artificial mRNA according to the present invention, for example, by optionally inserting an open reading frame or a sequence comprising an open reading frame into the vector and

transcribing the vector. Thus, preferably, the vector comprises elements needed for transcription, such as a promoter, e.g. an RNA polymerase promoter. Preferably, the vector is suitable for transcription using eukaryotic, prokaryotic, viral or phage transcription systems, such as eukaryotic cells, prokaryotic cells, or eukaryotic, prokaryotic, viral or phage *in vitro* transcription systems. Thus, for example, the vector may comprise a promoter sequence, which is recognized by a polymerase, such as by an RNA polymerase, e.g. by a eukaryotic, prokaryotic, viral, or phage RNA polymerase. In a preferred embodiment, the vector comprises a phage RNA polymerase promoter such as an SP6 or T7, preferably a T7 promoter. Preferably, the vector is suitable for *in vitro* transcription using a phage based *in vitro* transcription system, such as a T7 RNA polymerase based *in vitro* transcription system.

The vector may further comprise a poly(A) sequence and/or a polyadenylation signal as described above for the artificial nucleic acid molecule according to the present invention.

The vector may be an RNA vector or a DNA vector. Preferably, the vector is a DNA vector. The vector may be any vector known to the skilled person, such as a viral vector or a plasmid vector. Preferably, the vector is a plasmid vector, preferably a DNA plasmid vector.

In a preferred embodiment, the vector according to the present invention comprises the artificial nucleic acid molecule according to the present invention.

Preferably, a vector according to the present invention comprises a sequence according to SEQ ID NOs. 1-1363, 1395, 1421, 1422, 1368, or 1412-1420, or a sequence having an identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%; even more preferably of at least about 99%; even more preferably of 100% sequence identity to a sequence according to any one of SEQ ID NOs. 1-1363, 1395, 1421, 1422, 1368, or 1412-1420, or a fragment thereof, preferably a functional fragment thereof, or a corresponding RNA sequence.

Preferably, a vector, such as a DNA vector, according to the present invention comprises a sequence according to SEQ ID NOs. 1368-1392 or 1412-1420, or a sequence having an

identity of at least about 40%, preferably of at least about 50%, preferably of at least about 60%, preferably of at least about 70%, more preferably of at least about 80%, more preferably of at least about 90%, even more preferably of at least about 95%; even more preferably of at least about 99%; even more preferably of 100% sequence identity to a
5 sequence according to any one of SEQ ID NOs. 1368-1392 or 1412-1420 or a fragment thereof, preferably a functional fragment thereof, or a corresponding RNA sequence.

Preferably, the vector is a circular molecule. Preferably, the vector is a double-stranded molecule, such as a double stranded DNA molecule. Such circular, preferably double
10 stranded DNA molecule may be used conveniently as a storage form for the inventive artificial nucleic acid molecule. Furthermore, it may be used for transfection of cells, for example, cultured cells. Also it may be used for *in vitro* transcription for obtaining an artificial RNA molecule according to the invention.

Preferably, the vector, preferably the circular vector, is linearizable, for example, by restriction enzyme digestion. In a preferred embodiment, the vector comprises a cleavage site, such as a restriction site, preferably a unique cleavage site, located immediately 3' to the ORF, or - if present - immediately 3' to the 3'UTR element, or - if present - immediately 3' to the poly(A) sequence or polyadenylation signal, or - if present - located 3' to the
20 poly(C) sequence, or - if present - located 3' to the histone stem-loop". Thus, preferably, the product obtained by linearizing the vector terminates at the 3'end with the stop codon, or - if present - the 3'-end of the 3'UTR element, or - if present - with the 3'-end of the poly(A) sequence or with the 3'-end of the polyadenylation signal, or - if present - with the 3'-end of the poly(C) sequence, or - if present - with the 3'-end of the histone stem-loop, plus
25 optionally some nucleotides remaining from the restriction site after cleavage.

In a further aspect, the present invention relates to a cell comprising the artificial nucleic acid molecule according to the present invention or the vector according to present invention. The cell may be any cell, such as a bacterial cell, insect cell, plant cell,
30 vertebrate cell, e.g. a mammalian cell. Such cell may be, e.g., used for replication of the vector of the present invention, for example, in a bacterial cell. Furthermore, the cell may be used for transcribing the artificial nucleic acid molecule or the vector according to the present invention and/or translating the open reading frame of the artificial nucleic acid

molecule or the vector according to the present invention. For example, the cell may be used for recombinant protein production.

The cells according to the present invention are, for example, obtainable by standard nucleic acid transfer methods, such as standard transfection methods. For example, the artificial nucleic acid molecule or the vector according to the present invention may be transferred into the cell by electroporation, lipofection, e.g. based on cationic lipids and/or liposomes, calcium phosphate precipitation, nanoparticle based transfection, virus based transfection, or based on cationic polymers, such as DEAE-dextran or polyethylenimine etc.

Preferably, the cell is a mammalian cell, such as a cell of human subject, a domestic animal, a laboratory animal, such as a mouse or rat cell. Preferably the cell is a human cell. The cell may be a cell of an established cell line, such as a CHO, BHK, 293T, COS-7, HELA, HEK etc. cell, or the cell may be a primary cell, e.g. a HDF cell, preferably a cell isolated from an organism. In a preferred embodiment, the cell is an isolated cell of a mammalian subject, preferably of a human subject. For example, the cell may be an immune cell, such as a dendritic cell, a cancer or tumor cell, or any somatic cell etc., preferably of a mammalian subject, preferably of a human subject.

In a further aspect, the present invention provides a pharmaceutical composition comprising the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, or the cell according to the present invention. The pharmaceutical composition according to the invention may be used, e.g., as a vaccine, for example, for genetic vaccination. Thus, the ORF may, e.g., encode an antigen to be administered to a patient for vaccination. Thus, in a preferred embodiment, the pharmaceutical composition according to the present invention is a vaccine. Furthermore, the pharmaceutical composition according to the present invention may be used, e.g., for gene therapy.

Preferably, the pharmaceutical composition further comprises one or more pharmaceutically acceptable excipients, vehicles, fillers and/or diluents. In the context of the present invention, a pharmaceutically acceptable vehicle typically includes a liquid or non-liquid basis for the inventive pharmaceutical composition. In one embodiment, the pharmaceutical composition is provided in liquid form. In this context, preferably, the vehicle is based

on water, such as pyrogen-free water, isotonic saline or buffered (aqueous) solutions, e.g. phosphate, citrate etc. buffered solutions. The buffer may be hypertonic, isotonic or hypotonic with reference to the specific reference medium, i.e. the buffer may have a higher, identical or lower salt content with reference to the specific reference medium, wherein preferably such concentrations of the afore mentioned salts may be used, which do not lead to damage of mammalian cells due to osmosis or other concentration effects. Reference media are e.g. liquids occurring in "*in vivo*" methods, such as blood, lymph, cytosolic liquids, or other body liquids, or e.g. liquids, which may be used as reference media in "*in vitro*" methods, such as common buffers or liquids. Such common buffers or liquids are known to a skilled person. Ringer-Lactate solution is particularly preferred as a liquid basis.

One or more compatible solid or liquid fillers or diluents or encapsulating compounds suitable for administration to a patient may be used as well for the inventive pharmaceutical composition. The term "compatible" as used herein preferably means that these components of the inventive pharmaceutical composition are capable of being mixed with the inventive nucleic acid, vector or cells as defined herein in such a manner that no interaction occurs which would substantially reduce the pharmaceutical effectiveness of the inventive pharmaceutical composition under typical use conditions.

The pharmaceutical composition according to the present invention may optionally further comprise one or more additional pharmaceutically active components. A pharmaceutically active component in this context is a compound that exhibits a therapeutic effect to heal, ameliorate or prevent a particular indication or disease. Such compounds include, without implying any limitation, peptides or proteins, nucleic acids, (therapeutically active) low molecular weight organic or inorganic compounds (molecular weight less than 5000, preferably less than 1000), sugars, antigens or antibodies, therapeutic agents already known in the prior art, antigenic cells, antigenic cellular fragments, cellular fractions, cell wall components (e.g. polysaccharides), modified, attenuated or de-activated (e.g. chemically or by irradiation) pathogens (virus, bacteria etc.).


Furthermore, the inventive pharmaceutical composition may comprise a carrier for the artificial nucleic acid molecule or the vector. Such a carrier may be suitable for mediating dissolution in physiological acceptable liquids, transport and cellular uptake of the pharmaceutical active artificial nucleic acid molecule or the vector. Accordingly, such a carrier

may be a component which may be suitable for depot and delivery of an artificial nucleic acid molecule or vector according to the invention. Such components may be, for example, cationic or polycationic carriers or compounds which may serve as transfection or complexation agent.

5

Particularly preferred transfection or complexation agents in this context are cationic or polycationic compounds, including protamine, nucleoline, spermine or spermidine, or other cationic peptides or proteins, such as poly-L-lysine (PLL), poly-arginine, basic poly-peptides, cell penetrating peptides (CPPs), including HIV-binding peptides, HIV-1 Tat (HIV),
 10 Tat-derived peptides, Penetratin, VP22 derived or analog peptides, HSV VP22 (Herpes simplex), MAP, KALA or protein transduction domains (PTDs), PpT620, prolin-rich peptides, arginine-rich peptides, lysine-rich peptides, MPG-peptide(s), Pep-1, L-oligomers, Calcitonin peptide(s), Antennapedia-derived peptides (particularly from *Drosophila antennapedia*), pAntp, plsl, FGF, Lactoferrin, Transportan, Buforin-2, Bac715-24, SynB, SynB(1), pVEC,
 15 hCT-derived peptides, SAP, or histones.


Furthermore, such cationic or polycationic compounds or carriers may be cationic or polycationic peptides or proteins, which preferably comprise or are additionally modified to comprise at least one –SH moiety. Preferably, a cationic or polycationic carrier is selected
 20 from cationic peptides having the following sum formula (I):




wherein $l + m + n + o + x = 3-100$, and l , m , n or o independently of each other is any
 25 number selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81-90 and 91-100 provided that the overall content of Arg (Arginine), Lys (Lysine), His (Histidine) and Orn (Ornithine) represents at least 10% of all amino acids of the oligopeptide; and Xaa is any amino acid selected from native (= naturally occurring) or non-native amino acids except of Arg, Lys, His or Orn; and x is
 30 any number selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81-90, provided, that the overall content of Xaa does not exceed 90 % of all amino acids of the oligopeptide. Any of amino acids Arg,

Lys, His, Orn and Xaa may be positioned at any place of the peptide. In this context cationic peptides or proteins in the range of 7-30 amino acids are particular preferred.

Further, the cationic or polycationic peptide or protein, when defined according to formula
 5 $\{(Arg)_i;(Lys)_m;(His)_n;(Orn)_o;(Xaa)_x\}$ (formula (I)) as shown above and which comprise or are additionally modified to comprise at least one -SH moiety, may be, without being restricted thereto, selected from subformula (Ia):



wherein $(Arg)_i;(Lys)_m;(His)_n;(Orn)_o$ and x are as defined herein, Xaa' is any amino acid selected from native (= naturally occurring) or non-native amino acids except of Arg, Lys, His, Orn or Cys and y is any number selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-30, 31-40, 41-50, 51-60, 61-70, 71-80 and 81-90, provided that
 15 the overall content of Arg (Arginine), Lys (Lysine), His (Histidine) and Orn (Ornithine) represents at least 10% of all amino acids of the oligopeptide. Further, the cationic or polycationic peptide may be selected from subformula (Ib):



wherein empirical formula $\{(Arg)_i;(Lys)_m;(His)_n;(Orn)_o;(Xaa)_x\}$ (formula (III)) is as defined herein and forms a core of an amino acid sequence according to (semiempirical) formula (III) and wherein Cys₁ and Cys₂ are Cysteines proximal to, or terminal to $(Arg)_i;(Lys)_m;(His)_n;(Orn)_o;(Xaa)_x$.

Further preferred cationic or polycationic compounds, which can be used as transfection or complexation agent may include cationic polysaccharides, for example chitosan, polybrene, cationic polymers, e.g. polyethyleneimine (PEI), cationic lipids, e.g. DOTMA: [1-(2,3-sioleyloxy)propyl]-N,N,N-trimethylammonium chloride, DMRIE, di-C14-amidine, DOTIM, SAINT, DC-Chol, BGTC, CTAP, DOPC, DODAP, DOPE: Dioleoyl phosphatidylethanol-amine, DOSPA, DODAB, DOIC, DMEPC, DOGS: Dioctadecylamidoglycylspermin, DIMRI: Dimyristo-oxypropyl dimethyl hydroxyethyl ammonium bromide, DOTAP: dioleoyloxy-3-(trimethylammonio)propane, DC-6-14: O,O-

ditetradecanoyl-N-(α -trimethylammonioacetyl)diethanolamine chloride, CLIP1: rac-[2,3-dioctadecyloxypropyl)(2-hydroxyethyl)]-dimethylammonium chloride, CLIP6: rac-[2(2,3-dihexadecyloxypropyl-oxymethyloxy)ethyl]trimethylammonium, CLIP9: rac-[2(2,3-dihexadecyloxypropyl-oxysuccinyloxy)ethyl]-trimethylammonium, oligofectamine, or cationic or polycationic polymers, e.g. modified polyaminoacids, such as β -aminoacid-polymers or reversed polyamides, etc., modified polyethylenes, such as PVP (poly(N-ethyl-4-vinylpyridinium bromide)), etc., modified acrylates, such as pDMAEMA (poly(dimethylaminoethyl methylacrylate)), etc., modified Amidoamines such as pAMAM (poly(amidoamine)), etc., modified polybetaaminoester (PBAE), such as diamine end modified 1,4 butanediol diacrylate-co-5-amino-1-pentanol polymers, etc., dendrimers, such as polypropylamine dendrimers or pAMAM based dendrimers, etc., polyimine(s), such as PEI: poly(ethyleneimine), poly(propyleneimine), etc., polyallylamine, sugar backbone based polymers, such as cyclodextrin based polymers, dextran based polymers, chitosan, etc., silan backbone based polymers, such as PMOXA-PDMS copolymers, etc., blockpolymers consisting of a combination of one or more cationic blocks (e.g. selected from a cationic polymer as mentioned above) and of one or more hydrophilic or hydrophobic blocks (e.g. polyethyleneglycole); etc.

In this context, it is particularly preferred that the inventive artificial nucleic acid molecule or the inventive vector is complexed at least partially with a cationic or polycationic compound, preferably cationic proteins or peptides. Partially means that only a part of the inventive artificial nucleic acid molecule or the inventive vector is complexed with a cationic or polycationic compound and that the rest of the inventive artificial nucleic acid molecule or the inventive vector is in uncomplexed form ("free"). Preferably the ratio of complexed nucleic acid to: free nucleic acid is selected from a range. of about 5:1 (w/w) to about 1:10 (w/w), more preferably from a range of about 4:1 (w/w) to about 1:8 (w/w), even more preferably from a range of about 3:1 (w/w) to about 1:5 (w/w) or 1:3 (w/w), and most preferably the ratio of complexed nucleic acid to free nucleic acid is selected from a ratio of about 1:1 (w/w).

30

The pharmaceutical composition according to the present invention may optionally further comprise one or more adjuvants, for example, adjuvants for stimulating the innate immune system or for enhancing cellular uptake of the artificial nucleic acid molecule or vector. In

this context, an adjuvant may be understood as any compound, which is suitable to initiate or increase an immune response of the innate immune system, i.e. a non-specific immune response. In other words, when administered, the inventive pharmaceutical composition preferably elicits an innate immune response due to the adjuvant, optionally contained
5 therein. Preferably, such an adjuvant may be an adjuvant supporting the induction of an innate immune response in a mammal. Such an adjuvant may be, for example, an immunostimulatory nucleic acid, i.e. a nucleic acid that may bind to a Toll-like-receptor or the like, preferably an immunostimulatory RNA.

10 Such adjuvants, preferably such immunostimulatory nucleic acids, may induce an innate, i.e. unspecific, immune response which may support a specific, i.e. adaptive, immune response to the peptide or protein, i.e. the antigen, encoded by the artificial nucleic acid molecule of the pharmaceutical composition, preferably the vaccine.

15 The inventive pharmaceutical composition may also additionally comprise any further compound, which is known to be immunostimulating due to its binding affinity (as ligands) to human Toll-like receptors TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, or due to its binding affinity (as ligands) to murine Toll-like receptors TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, TLR11, TLR12 or TLR13.

20

Further additives which may be included in the inventive pharmaceutical composition are, e.g., emulsifiers, such as, for example, Tween®; wetting agents, such as, for example, sodium lauryl sulfate; colouring agents; taste-imparting agents, pharmaceutical carriers; tablet-forming agents; stabilizers; antioxidants; preservatives etc.

25

The pharmaceutical composition according to the present invention preferably comprises a "safe and effective amount" of the components of the pharmaceutical composition, particularly of the inventive nucleic acid sequence, the vector and/or the cells as defined herein. As used herein, a "safe and effective amount" means an amount sufficient to significantly
30 induce a positive modification of a disease or disorder as defined herein. At the same time, however, a "safe and effective amount" preferably avoids serious side-effects and permits a sensible relationship between advantage and risk. The determination of these limits typically lies within the scope of sensible medical judgment.

In a further aspect, the present invention provides the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention for use as a medicament, for example, as vaccine (in genetic vaccination) or in gene therapy.

The artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention are particularly suitable for any medical application which makes use of the therapeutic action or effect of peptides, polypeptides or proteins, or where supplementation of a particular peptide or protein is needed. Thus, the present invention provides the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention for use in the treatment or prevention of diseases or disorders amenable to treatment by the therapeutic action or effect of peptides, polypeptides or proteins or amenable to treatment by supplementation of a particular peptide, polypeptide or protein. For example, the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention may be used for the treatment or prevention of genetic diseases, autoimmune diseases, cancerous or tumour-related diseases, infectious diseases, chronic diseases or the like, e.g., by genetic vaccination or gene therapy.

In particular, such therapeutic treatments which benefit from a stable, prolonged and/or increased presence of therapeutic peptides, polypeptides or proteins in a subject to be treated are especially suitable as medical application in the context of the present invention, since the 5'UTR element optionally in combination with the 3'UTR element provides for increased protein expression from the ORF and the 3'UTR element provides for a stable and prolonged expression of the ORF of the inventive nucleic acid molecule. Thus, a particularly suitable medical application for the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention

is vaccination, for example against infections or tumours. Thus, the present invention provides the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention for vaccination of a subject, preferably a mammalian subject, more preferably a human subject. Preferred vaccination treatments are vaccination against infectious diseases, such as bacterial, protozoal or viral infections, and anti-tumour-vaccination. Such vaccination treatments may be prophylactic or therapeutic.

Depending on the disease to be treated or prevented, the ORF may be selected. For example, the open reading frame may code for a protein that has to be supplied to a patient suffering from total lack or at least partial loss of function of a protein, such as a patient suffering from a genetic disease. Additionally, the open reading frame may be chosen from an ORF coding for a peptide or protein which beneficially influences a disease or the condition of a subject. Furthermore, the open reading frame may code for a peptide or protein which effects down-regulation of a pathological overproduction of a natural peptide or protein or elimination of cells expressing pathologically a protein or peptide. Such lack, loss of function or overproduction may, e.g., occur in the context of tumour and neoplasia, autoimmune diseases, allergies, infections, chronic diseases or the like. Furthermore, the open reading frame may code for an antigen or immunogen, e.g. for an epitope of a pathogen or for a tumour antigen. Thus, in preferred embodiments, the artificial nucleic acid molecule or the vector according to the present invention comprises an ORF encoding an amino acid sequence comprising or consisting of an antigen or immunogen, e.g. an epitope of a pathogen or a tumour-associated antigen, a 5'UTR element as described above, and optional further components, such as a 3'UTR element and/or a poly(A) sequence etc. as described above.

In the context of medical application, in particular, in the context of vaccination, it is preferred that the artificial nucleic acid molecule according to the present invention is RNA, preferably mRNA, since DNA harbours the risk of eliciting an anti-DNA immune response and tends to insert into genomic DNA. However, in some embodiments, for example, if a viral delivery vehicle, such as an adenoviral delivery vehicle is used for delivery of the artificial nucleic acid molecule or the vector according to the present invention, e.g., in the

context of gene therapeutic treatments, it may be desirable that the artificial nucleic acid molecule or the vector is a DNA molecule.

The artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional, intracranial, transdermal, intradermal, intrapulmonal, intraperitoneal, intracardial, intraarterial, and sublingual injection or infusion techniques.

Preferably, the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention is administered parenterally, e.g. by parenteral injection, more preferably by subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional, intracranial, transdermal, intradermal, intrapulmonal, intraperitoneal, intracardial, intraarterial, sublingual injection or via infusion techniques. Particularly preferred is intradermal and intramuscular injection. Sterile injectable forms of the inventive pharmaceutical composition may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents.

The artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention may also be administered orally in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions.

The artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention may also be administered topically, espe-

cially when the target of treatment includes areas or organs readily accessible by topical application, e.g. including diseases of the skin or of any other accessible epithelial tissue. Suitable topical formulations are readily prepared for each of these areas or organs. For topical applications, the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention may be formulated in a suitable ointment suspended or dissolved in one or more carriers.

In one embodiment, the use as a medicament comprises the step of transfection of mammalian cells, preferably *in vitro* transfection of mammalian cells, more preferably *in vitro* transfection of isolated cells of a subject to be treated by the medicament. If the use comprises the *in vitro* transfection of isolated cells, the use as a medicament may further comprise the (re)administration of the transfected cells to the patient. The use of the inventive artificial nucleic acid molecules or the vector as a medicament may further comprise the step of selection of successfully transfected isolated cells. Thus, it may be beneficial if the vector further comprises a selection marker. Also, the use as a medicament may comprise *in vitro* transfection of isolated cells and purification of an expression-product, i.e. the encoded peptide or protein from these cells. This purified peptide or protein may subsequently be administered to a subject in need thereof.

The present invention also provides a method for treating or preventing a disease or disorder as described above comprising administering the artificial nucleic acid molecule according to the present invention, the vector according to the present invention, the cell according to the present invention, or the pharmaceutical composition according to the present invention to a subject in need thereof.

Furthermore, the present invention provides a method for treating or preventing a disease or disorder comprising transfection of a cell with an artificial nucleic acid molecule according to the present invention or with the vector according to the present invention. Said transfection may be performed *in vitro* or *in vivo*. In a preferred embodiment, transfection of a cell is performed *in vitro* and the transfected cell is administered to a subject in need thereof, preferably to a human patient. Preferably, the cell which is to be transfected *in vitro* is an isolated cell of the subject, preferably of the human patient. Thus, the present

invention provides a method of treatment comprising the steps of isolating a cell from a subject, preferably from a human patient, transfecting the isolated cell with the artificial nucleic acid molecule according to the present invention or the vector according to the present invention, and administering the transfected cell to the subject, preferably the human patient.

The method of treating or preventing a disorder according to the present invention is preferably a vaccination method and/or a gene therapy method as described above.

As described above, the 5'UTR element and the optional 3'UTR element are capable of increasing protein production from an artificial nucleic acid molecule, such as an mRNA or vector, comprising the 5'UTR element and an ORF. Thus, in a further aspect, the present invention relates to a method for increasing protein production from an artificial nucleic acid molecule comprising the step of associating the artificial nucleic acid molecule, preferably the ORF contained within the artificial nucleic acid molecule, with (i) at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene as described above and (ii) optionally at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene, or from a variant of the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene as described above.

The term "associating the artificial nucleic acid molecule or the vector with a 5'UTR element and an optional 3'UTR element" in the context of the present invention preferably means functionally associating or functionally combining the artificial nucleic acid molecule, such as the mRNA or the vector, with the 5'UTR element and the optional 3'UTR element. This means that the artificial nucleic acid molecule, preferably the ORF contained within the artificial nucleic acid molecule, the 5'UTR element and the optional 3'UTR element as described above are associated or coupled such that the function of the 5'UTR element and the optional 3'UTR element, e.g., protein production increasing function, is exerted. Typically, this means that the 5'UTR element and the optional 3'UTR element are

integrated into the artificial nucleic acid molecule, preferably into the mRNA molecule or the vector, such that the open reading frame is positioned 3' to the 5'UTR element, preferably between the 5'UTR element and the optional 3'UTR element.

- 5 In a further aspect, the present invention provides the use of at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene as described above and optionally at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of a chordate
10 gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene, or from a variant of the 3'UTR of a chordate gene, preferably a vertebrate gene, more preferably a mammalian gene, most preferably a human gene as described above for increasing protein production from an artificial nucleic acid molecule, such as an mRNA or a vector.

15

The use according to the present invention preferably comprises associating the artificial nucleic acid molecule with the 5'UTR element and the optional 3'UTR element as described above.

- 20 The method for increasing protein production from an artificial nucleic acid molecule and the above use may also comprise associating the artificial nucleic acid molecules with one or more further elements, such as a polyadenylation signal, a poly(A) sequence, a poly(C) sequence and/or a histone stem loop as described above.

- 25 The compounds and ingredients of the inventive pharmaceutical composition may also be manufactured and traded separately of each other. Thus, the invention relates further to a kit or kit of parts comprising an artificial nucleic acid molecule according to the invention, a vector according to the invention, a cell according to the invention, and/or a pharmaceutical composition according to the invention. Preferably, such kit or kit of parts may,
30 additionally, comprise instructions for use, cells for transfection, an adjuvant, a means for administration of the pharmaceutical composition, a pharmaceutically acceptable carrier and/or an pharmaceutically acceptable solution for dissolution or dilution of the artificial nucleic acid molecule, the vector, the cells or the pharmaceutical composition.

It is to be noted that, throughout the description and claims of this specification, the word 'comprise' and variations of the word, such as 'comprising' and 'comprises', is not intended to exclude other variants or additional components, integers or steps. Modifications and improvements to the invention will be readily apparent to those skilled in the art. Such
5 modifications and improvements are intended to be within the scope of this invention.

Any reference to or discussion of any document, act or item of knowledge in this specification is included solely for the purpose of providing a context for the present invention. It is not suggested or represented that any of these matters or any combination thereof formed at the
10 priority date part of the common general knowledge, or was known to be relevant to an attempt to solve any problem with which this specification is concerned.

The following Figures, Sequences and Examples are intended to illustrate the invention further. They are not intended to limit the subject-matter of the invention thereto.

- 5 Fig. 1 shows the nucleotide sequence of a *Photinus pyralis* luciferase encoding nucleic acid molecule PpLuc(GC) – A64N64. This artificial construct does not comprise a 5'UTR element or a 3'UTR element in the sense of the present invention. The coding region for PpLuc(GC) is depicted in italics.
- 10 Fig. 2 shows the nucleotide sequence of PpLuc(GC) – albumin7 – A64N64. The 3'UTR of human albumin, with a T7 termination signal as well as a HindIII and XbaI restriction site removed by three single point mutations, was inserted between the ORF and poly(A) of the construct shown in Figure 1. The coding region for PpLuc(GC) is depicted in italics. The albumin 3'UTR is underlined.
- 15 Fig. 3 shows the nucleotide sequence of RPL32 – PpLuc(GC) – A64N64. The 5'UTR of human ribosomal protein Large 32 gene lacking the 5' terminal oligopyrimidine tract (RPL32) according to SEQ ID NO. 1368 was inserted 5' of the ORF in the construct shown in Figure 1. The coding region for PpLuc(GC) is depicted in italics. The RPL32 5'UTR is underlined.
- 20 Fig. 4 shows the nucleotide sequence of RPL32 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human ribosomal protein Large 32 gene lacking the 5' terminal oligopyrimidine tract (RPL32) according to SEQ ID NO. 1368 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.
- 25 Fig. 5 is a graphical representation of the effect of the TOP 5'UTR element which is derived from the 5'UTR of the TOP gene RPL23 according to SEQ ID NO. 1368, the albumin 3' UTR element according to SEQ ID NO. 1376 and the combination of the TOP 5'UTR element and the albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the TOP 5'UTR element increases luciferase levels
- 30

compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SD (relative light units $\pm$ standard deviation) for triplicate transfections. RLU are summarized in Example 5.1.

Fig. 6 shows the nucleotide sequence of RPL35 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human ribosomal protein Large 35 gene lacking the 5' terminal oligopyrimidine tract (RPL35) according to SEQ ID NO. 1412 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.

Fig. 7 shows the nucleotide sequence of RPL21 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human ribosomal protein Large 21 gene lacking the 5' terminal oligopyrimidine tract (RPL21) according to SEQ ID NO. 1413 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.

Fig. 8 shows the nucleotide sequence of atp5a1 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1 gene lacking the 5' terminal oligopyrimidine tract (atp5a1) according to SEQ ID NO. 1414 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.

Fig. 9 shows the nucleotide sequence of HSD17B4 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human hydroxysteroid (17-beta) dehydrogenase 4 gene lacking the 5' terminal oligopyrimidine tract (HSD17B4) according to SEQ ID NO. 1415 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.

Fig. 10 shows the nucleotide sequence of AIG1 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human androgen-induced 1 gene lacking the 5' terminal oligopyrimidine tract (AIG1) according to SEQ ID NO. 1416 and the albumin7

3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.

- Fig. 11 shows the nucleotide sequence of COX6C – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human cytochrome c oxidase subunit VIc gene lacking the 5' terminal oligopyrimidine tract (COX6C) according to SEQ ID NO. 1417 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.
- Fig. 12 shows the nucleotide sequence of ASAH1 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of human N-acylsphingosine amidohydrolase (acid ceramidase) 1 lacking the 5' terminal oligopyrimidine tract (ASAH1) according to SEQ ID NO. 1418 and the albumin7 3'UTR according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.
- Fig. 13 shows the nucleotide sequence of mRPL21 – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of murine ribosomal protein Large 21 gene lacking the 5' terminal oligopyrimidine tract (mRPL21) according to SEQ ID NO. 1419 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.
- Fig. 14 shows the nucleotide sequence of mRPL35A – PpLuc(GC) – albumin7 – A64N64. The 5'UTR of murine ribosomal protein Large 35a gene lacking the 5' terminal oligopyrimidine tract (mRPL35A) according to SEQ ID NO. 1420 and the albumin7 3'UTR element according to SEQ ID NO. 1376 were inserted 5' and 3' of the ORF in the construct shown in Figure 1, respectively.
- Fig. 15 shows the nucleotide sequence of RPL35 – PpLuc(GC) – A64N64. The 5'UTR of human ribosomal protein Large 35 gene lacking the 5' terminal oligopyrimidine tract (RPL35) according to SEQ ID NO. 1412 was inserted 5' of the ORF in the construct shown in Figure 1.
- Fig. 16 shows the nucleotide sequence of RPL21 – PpLuc(GC) – A64N64. The 5'UTR of human ribosomal protein Large 21 gene lacking the 5' terminal oli-

gopyrimidine tract (RPL21) according to SEQ ID NO. 1413 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 17 shows the nucleotide sequence of atp5a1 – PpLuc(GC) – A64N64. The 5'UTR of human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1 gene lacking the 5' terminal oligopyrimidine tract (atp5a1) according to SEQ ID NO. 1414 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 18 shows the nucleotide sequence of HSD17B4 – PpLuc(GC) – A64N64. The 5'UTR of human hydroxysteroid (17-beta) dehydrogenase 4 gene lacking the 5' terminal oligopyrimidine tract (HSD17B4) according to SEQ ID NO. 1415 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 19 shows the nucleotide sequence of AIG1 – PpLuc(GC) – A64N64. The 5'UTR of human androgen-induced 1 gene lacking the 5' terminal oligopyrimidine tract (AIG1) according to SEQ ID NO. 1416 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 20 shows the nucleotide sequence of COX6C – PpLuc(GC) – A64N64. The 5'UTR of human cytochrome c oxidase subunit VIc gene lacking the 5' terminal oligopyrimidine tract (COX6C) according to SEQ ID NO. 1417 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 21 shows the nucleotide sequence of ASAH1 – PpLuc(GC) – A64N64. The 5'UTR of human N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene lacking the 5' terminal oligopyrimidine tract (ASAH1) according to SEQ ID NO. 1418 was inserted 5' of the ORF in the construct shown in Figure 1.

Fig. 22 is a graphical representation of the effect of different TOP 5'UTR elements on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. TOP 5'UTR elements strongly increase luciferase levels compared to mRNA lacking a 5'UTR element. mRNAs comprising 5'UTR elements derived from the 5'UTRs of the TOP genes ASAH1, COX6C, AIG1, HSD17B4, atp5a1, RPL21, RPL35 and RPL32 were

transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The TOP 5'UTR elements increases luciferase levels compared to mRNA lacking a 5'UTR element. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. RLU are summarized in Example 5.2.

Fig. 23 is a graphical representation of the effect of the RPL35 TOP 5'UTR element, the albumin 3'UTR element and the combination of RPL35 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the RPL35 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of RPL35 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 24 is a graphical representation of the effect of the RPL21 TOP 5'UTR element, the albumin 3'UTR element and the combination of RPL21 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the RPL21 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of RPL21 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 25 is a graphical representation of the effect of the atp5a1 TOP 5'UTR element, the albumin 3'UTR element and the combination of atp5a1 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the atp5a1 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of atp5a1 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 26 is a graphical representation of the effect of the HSD17B4 TOP 5'UTR element, the albumin 3'UTR element and the combination of HSD17B4 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the HSD17B4 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of HSD17B4 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 27 is a graphical representation of the effect of the AIG1 TOP 5'UTR element, the albumin 3'UTR element and the combination of AIG1 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression,

while the AIG1 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of AIG1 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 28 is a graphical representation of the effect of the COX6C TOP 5'UTR element, the albumin 3'UTR element and the combination of COX6C TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the COX6C TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of COX6C TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 29 is a graphical representation of the effect of the ASAH1 TOP 5'UTR element, the albumin 3'UTR element and the combination of ASAH1 TOP 5'UTR element and albumin 3'UTR element on luciferase expression from mRNA. A variety of mRNAs were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. The albumin 3'UTR element extends luciferase expression, while the ASAH1 TOP 5'UTR element increases luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly, the combination of ASAH1 TOP 5'UTR element and albumin 3'UTR element further strongly increases the luciferase level, much above the level observed with either of the individual elements, thus acting synergistically. Data are

graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. Synergy is summarized in Example 5.3.

Fig. 30 is a graphical representation of the effect of the TOP 5'UTR element from mouse genes on luciferase expression from mRNA. mRNAs containing either a mouse or a human TOP 5'UTR element were transfected into human dermal fibroblasts (HDF) by lipofection. Luciferase levels were measured at 24, 48, and 72 hours after transfection. Mouse TOP 5'UTR elements strongly increase luciferase levels compared to mRNA lacking a 5'-element, similarly as the human TOP 5'UTR element. Data are graphed as mean RLU $\pm$ SEM (relative light units $\pm$ standard error) for triplicate transfections. RLU are summarized in Example 5.4.

SEQ ID No. 1-1363, 1395, 1421, and 1422 Sequences comprising 5'UTRs of TOP genes

	SEQ ID No. 1364	PpLuc(GC) – A64N64
15	SEQ ID No. 1365	PpLuc(GC) – albumin7 – A64N64
	SEQ ID No. 1366	RPL32 – PpLuc(GC) – A64N64
	SEQ ID No. 1367	RPL32 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID No. 1368	5'UTR of human ribosomal protein Large 32 lacking the 5' terminal oligopyrimidine tract
20	SEQ ID No. 1369	Human albumin 3'UTR
	SEQ ID No. 1370	3'UTR of Homo sapiens hemoglobin, alpha 1 (HBA1)
	SEQ ID No. 1371	3'UTR of Homo sapiens hemoglobin, alpha 2 (HBA2)
	SEQ ID No. 1372	3'UTR of Homo sapiens hemoglobin, beta (HBB)
	SEQ ID No. 1373	3'UTR of Homo sapiens tyrosine hydroxylase (TH)
25	SEQ ID No. 1374	3'UTR of Homo sapiens arachidonate 15-lipoxygenase (ALOX15)
	SEQ ID No. 1375	3'UTR of Homo sapiens collagen, type I, alpha 1 (COL1A1)
	SEQ ID No. 1376	albumin7 3'UTR
	SEQ ID No. 1377	Human albumin 3'UTR + poly(A) sequence
	SEQ ID No. 1378	Human albumin 3'UTR fragment 1
30	SEQ ID No. 1379	Human albumin 3'UTR fragment 2
	SEQ ID No. 1380	Human albumin 3'UTR fragment 3
	SEQ ID No. 1381	Human albumin 3'UTR fragment 4

	SEQ ID No. 1382	Human albumin 3'UTR fragment 5
	SEQ ID No. 1383	Human albumin 3'UTR fragment 6
	SEQ ID No. 1384	Human albumin 3'UTR fragment 7
	SEQ ID No. 1385	Human albumin 3'UTR fragment 8
5	SEQ ID No. 1386	Human albumin 3'UTR fragment 9
	SEQ ID No. 1387	Human albumin 3'UTR fragment 10
	SEQ ID No. 1388	Human albumin 3'UTR fragment 11
	SEQ ID No. 1389	Human albumin 3'UTR fragment 12
	SEQ ID No. 1390	Human albumin 3'UTR fragment 13
10	SEQ ID No. 1391	Albumin7 3'UTR - poly(A) sequence - poly(C) sequence – HL
	SEQ ID No. 1392	Albumin7 3'UTR - poly(A) sequence - poly(C) sequence
	SEQ ID No. 1393	Center, α -complex-binding portion of the 3'UTR of an α -globin gene
	SEQ ID No. 1394	Histone stem-loop
	SEQ ID NO. 1396	RPL35 – PpLuc(GC) – albumin7 – A64N64
15	SEQ ID NO. 1397	RPL21 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1398	ATP5A1 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1399	HSD17B4 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1400	AIG1 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1401	COX6C – PpLuc(GC) – albumin7 – A64N64
20	SEQ ID NO. 1402	ASA1 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1403	mRPL21 – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1404	mRPL35A – PpLuc(GC) – albumin7 – A64N64
	SEQ ID NO. 1405	RPL35 – PpLuc(GC) – A64N64
	SEQ ID NO. 1406	RPL21 – PpLuc(GC) – A64N64
25	SEQ ID NO. 1407	ATP5A1 – PpLuc(GC) – A64N64
	SEQ ID NO. 1408	HSD17B4 – PpLuc(GC) – A64N64
	SEQ ID NO. 1409	AIG1 – PpLuc(GC) – A64N64
	SEQ ID NO. 1410	COX6C – PpLuc(GC) – A64N64
	SEQ ID NO. 1411	ASA1 – PpLuc(GC) – A64N64
30	SEQ ID NO. 1412	5'UTR of human ribosomal protein Large 35 (RPL35) lacking the 5' terminal oligopyrimidine tract
	SEQ ID NO. 1413	5'UTR of human ribosomal protein Large 21 (RPL21) lacking the 5' terminal oligopyrimidine tract

	SEQ ID NO. 1414	5'UTR of human ATP synthase, H ⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) lacking the 5' terminal oligopyrimidine tract
5	SEQ ID NO. 1415	5'UTR of human hydroxysteroid (17-beta) dehydrogenase 4 (HSD17B4) lacking the 5' terminal oligopyrimidine tract
	SEQ ID NO. 1416	5'UTR of human androgen-induced 1 (AIG1) lacking the 5' terminal oligopyrimidine tract
	SEQ ID NO. 1417	5'UTR of human cytochrome c oxidase subunit VIc (COX6C) lacking the 5' terminal oligopyrimidine tract
10	SEQ ID NO. 1418	5'UTR of human N-acylsphingosine amidohydrolase (acid ceramidase) 1 (ASAH1) lacking the 5' terminal oligopyrimidine tract
	SEQ ID NO. 1419	5'UTR of mouse ribosomal protein Large 21 (mRPL21) lacking the 5' terminal oligopyrimidine tract
15	SEQ ID NO. 1420	5'UTR of mouse ribosomal protein large 35A (mRPL35A) lacking the 5' terminal oligopyrimidine tract

Examples

1. Preparation of DNA-templates

A vector for *in vitro* transcription was constructed containing a T7 promoter followed by a GC-enriched sequence coding for *Photinus pyralis* luciferase (PpLuc(GC)) and an A64 poly(A) sequence. The poly(A) sequence was followed by a restriction site used for linearization of the vector before *in vitro* transcription. mRNA obtained from this vector accordingly by *in vitro* transcription is designated as „PpLuc(GC) – A64N64“.

This vector was modified to include untranslated sequences 5' or 3' of the open reading frame (5'UTR or 3'UTR, respectively). In summary, vectors comprising the following mRNA encoding sequences have been generated (the mRNA coding sequences are depicted in Figures 1 to 4 and 6 to 21):

- SEQ ID No. 1364 (Fig. 1): PpLuc(GC) – A64N64
- SEQ ID No. 1365 (Fig. 2): PpLuc(GC) – albumin7 – A64N64
- SEQ ID No. 1366 (Fig. 3): RPL32 – PpLuc(GC) – A64N64

- SEQ ID No. 1367 (Fig. 4): RPL32 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1396 (Fig. 6): RPL35 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1397 (Fig. 7): RPL21 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1398 (Fig. 8): ATP5A1 – PpLuc(GC) – albumin7 – A64N64
 5 SEQ ID NO. 1399 (Fig. 9): HSD17B4 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1400 (Fig. 10): AIG1 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1401 (Fig. 11): COX6C – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1402 (Fig. 12): ASAH1 – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1403 (Fig. 13): mRPL21 – PpLuc(GC) – albumin7 – A64N64
 10 SEQ ID NO. 1404 (Fig. 14): mRPL35A – PpLuc(GC) – albumin7 – A64N64
 SEQ ID NO. 1405 (Fig. 15): RPL35 – PpLuc(GC) – A64N64
 SEQ ID NO. 1406 (Fig. 16): RPL21 – PpLuc(GC) – A64N64
 SEQ ID NO. 1407 (Fig. 17): ATP5A1 – PpLuc(GC) – A64N64
 SEQ ID NO. 1408 (Fig. 18): HSD17B4 – PpLuc(GC) – A64N64
 15 SEQ ID NO. 1409 (Fig. 19): AIG1 – PpLuc(GC) – A64N64
 SEQ ID NO. 1410 (Fig. 20): COX6C – PpLuc(GC) – A64N64
 SEQ ID NO. 1411 (Fig. 21): ASAH1 – PpLuc(GC) – A64N64

20 2. *In vitro* transcription

The DNA-template according to Example 1 was linearized and transcribed *in vitro* using T7-Polymerase. The DNA-template was then digested by DNase-treatment. mRNA transcripts contained a 5'-CAP structure obtained by adding an excess of N7-Methyl-Guanosine-5'-Triphosphate-5'-Guanosine to the transcription reaction. mRNA thus obtained was purified
 25 and resuspended in water.

3. Luciferase expression by mRNA lipofection

Human dermal fibroblasts (HDF) were seeded in 24 well plates at a density of 5×10^4 cells per well. The following day, cells were washed in opti-MEM and then transfected with
 30 50 ng per well of Lipofectamine2000-complexed PpLuc-encoding mRNA in opti-MEM. As a control, mRNA not coding for PpLuc was lipofected separately. mRNA coding for *Renilla reniformis* luciferase (RrLuc) was transfected together with PpLuc mRNA to control for transfection efficiency (20 ng of RrLuc mRNA per well). 90 minutes after start of transfection,

opti-MEM was exchanged for medium. 24, 48, 72 hours after transfection, medium was aspirated and cells were lysed in 200 µl of lysis buffer (25 mM Tris, pH 7.5 (HCl), 2 mM EDTA, 10% glycerol, 1% Triton X-100, 2 mM DTT, 1 mM PMSF). Lysates were stored at -20°C until luciferase activity was measured.

5

Alternatively, HDF were seeded in 96 well plates three days before transfection at a density of 10^4 cells per well. Immediately before lipofection, cells were washed in opti-MEM. Cells were lipofected with 25 ng of PpLuc-encoding mRNA per well complexed with Lipofectamine2000. mRNA coding for Renilla reniformis luciferase (RrLuc) was transfected together with PpLuc mRNA to control for transfection efficiency (2.5 ng of RrLuc mRNA per well). 90 minutes after start of transfection, opti-MEM was exchanged for medium. 24, 48, 72 hours after transfection, medium was aspirated and cells were lysed in 100 µl of lysis buffer (Passive Lysis Buffer, Promega). Lysates were stored at -80°C until luciferase activity was measured.

10

15

4. Luciferase measurement

Luciferase activity was measured as relative light units (RLU) in a BioTek SynergyHT plate reader. PpLuc activity was measured at 15 seconds measuring time using 50 µl of lysate and 200 µl of luciferin buffer (75 µM luciferin, 25 mM Glycylglycin, pH 7.8 (NaOH), 15 mM MgSO₄, 2 mM ATP). RrLuc activity was measured at 15 seconds measuring time using 50 µl of lysate and 200 µl of coelenterazin buffer (40 µM coelenterazin in phosphate buffered saline adjusted to 500 mM NaCl).

20

Alternatively, luciferase activity was measured as relative light units (RLU) in a Hidex Chameleon plate reader. PpLuc activity was measured at 2 seconds measuring time using 20 µl of lysate and 50 µl of luciferin buffer (Beetle-Juice, PJK GmbH). RrLuc activity was measured at 2 seconds measuring time using 20 µl of lysate and 50 µl of coelenterazin buffer (Renilla-Juice, PJK GmbH).

25

30

Results

5.1 The combination of TOP 5'UTR element and albumin 3'UTR element increases protein expression from mRNA in a synergistic manner.

To investigate the effect of the combination of a TOP 5'UTR element and an albumin 3'UTR element on protein expression from mRNA, mRNAs with different UTRs were synthesized: mRNAs either lacked both TOP 5'UTR element and albumin 3'UTR element, or contained either a TOP 5'UTR element (RPL32) or an albumin 3'UTR element (albumin7), or both TOP 5'UTR element and albumin 3'UTR element. Luciferase encoding mRNAs or control mRNA were transfected into human dermal fibroblasts (HDF). Luciferase levels were measured at 24, 48, and 72 hours after transfection. The PpLuc signal was corrected for transfection efficiency by the signal of cotransfected RrLuc (see following Table 1 and Figure 5).

Table 1:

mRNA	RLU at 24 hours	RLU at 48 hours	RLU at 72 hours
PpLuc(GC)-A64N64	115147	28973	8371
PpLuc(GC)-albumin7-A64N64	120234	48546	38138
RPL32-PpLuc(GC)-A64N64	671815	168741	21709
RPL32-PpLuc(GC)-albumin7-A64N64	913310	381288	100890

Luciferase was clearly expressed from mRNA having neither TOP 5'UTR nor albumin 3'UTR (PpLuc(GC)-A64N64). The albumin 3'UTR element extended luciferase expression, while the TOP 5'UTR element increased luciferase levels compared to mRNA lacking 5'- and 3'UTR elements. Strikingly however, the combination of TOP 5'UTR element and albumin 3'UTR element further strongly increased the luciferase level, much above the level observed with either of the individual elements. The magnitude of the rise in luciferase level due to combining TOP 5'UTR element and albumin 3'UTR element in the same mRNA demonstrates that they are acting synergistically.

The synergy between TOP 5'UTR element and albumin 3'UTR element was quantified by dividing the signal from mRNA combining both elements by the sum of the signal from mRNA lacking both elements plus the rise in signal effected by the TOP 5'UTR element plus the rise in signal effected by the albumin 3'UTR element. This calculation was performed for the three time points individually and for total protein expressed from 0 to 72 hours calculated from the area under the curve (AUC) (see following Table 2).

Table 2:

24 h					
RPL32	albumin	RLU	Δ RLU	RLU predicted (additive)	synergy
—	—	115147			
—	+	120234	5088		
+	—	671815	556668		
+	+	913310		676903	1,35
48 h					
RPL32	albumin	RLU	Δ RLU	RLU predicted (additive)	synergy
—	—	28973			
—	+	48546	19573		
+	—	168741	139768		
+	+	381288		188313	2,02
72 h					
RPL32	albumin	RLU	Δ RLU	RLU predicted (additive)	synergy
—	—	8371			
—	+	38138	29767		
+	—	21709	13338		
+	+	100890		51476	1,96
AUC 0 - 72 hours					
RPL32	albumin	RLU	Δ RLU	RLU predicted (additive)	synergy
—	—	3559000			
—	+	4508000	949000		
+	—	20430000	16871000		

+	+	32280000		21379000	1,51
---	---	----------	--	----------	------

The synergy thus calculated specifies how much higher the luciferase level from mRNA combining TOP 5'UTR element and albumin 3'UTR element is than would be expected if the effects of TOP 5'UTR element and albumin 3'UTR element were purely additive. The

5 luciferase level from mRNA combining TOP 5'UTR element and albumin 3'UTR element was up to two times higher than if their effects were purely additive. This result confirms that the combination of TOP 5'UTR element and albumin 3'UTR element effects a markedly synergistic increase in protein expression.

10 5.2 TOP 5'UTR elements increase protein expression from mRNA.

To investigate the effect of TOP 5'UTR elements on protein expression from mRNA, mRNAs comprising different TOP 5'UTR elements were synthesized. In addition, mRNAs contained the albumin7 3'UTR element. Luciferase encoding mRNAs were transfected into human dermal fibroblasts (HDF). Luciferase levels were measured at 24, 48, and

15 72 hours after transfection (see following Table 3 and Figure 22).

Table 3:

5'UTR	RLU at 24 hours	RLU at 48 hours	RLU at 72 hours
none	114277	121852	68235
RPL32	332236	286792	114148
RPL35	495917	234070	96993
RPL21	563314	352241	156605
atp5a1	1000253	538287	187159
HSD17B4	1179847	636877	299337
AIG1	620315	446621	167846

COX6C	592190	806065	173743
ASAH1	820413	529901	198429

Luciferase was clearly expressed from mRNA lacking a 5'UTR element. Strikingly however, all TOP 5'UTR elements strongly increased the luciferase level.

5 5.3 The combination of TOP 5'UTR elements and albumin 3'UTR element increases protein expression from mRNA in a synergistic manner.

To investigate the effect of the combination of TOP 5'UTR elements and an albumin 3'UTR element on protein expression from mRNA, mRNAs comprising different UTR elements were synthesized: mRNAs either lacked both TOP 5'UTR element and albumin 3'UTR element, or contained an albumin 3'UTR element, or contained one of different TOP 5'UTR elements, or contained both one of different TOP 5'UTR elements and an albumin 3'UTR element. Luciferase encoding mRNAs were transfected into human dermal fibroblasts (HDF). Luciferase levels were measured at 24, 48, and 72 hours after transfection (see Figures 23 to 30). Luciferase was clearly expressed from mRNA having neither a TOP 5'UTR element nor an albumin 3'UTR element. The albumin 3'UTR element extended luciferase expression, while TOP 5'UTR elements increased luciferase levels compared to mRNA lacking 5' and 3'UTRs. Strikingly however, the combinations of TOP 5'UTR elements and albumin 3'UTR element further strongly increased the luciferase level, much above the level observed with either of the individual elements. The magnitude of the rise in luciferase level due to combining TOP 5'UTR element and albumin 3'UTR element in the same mRNA demonstrates that they are acting synergistically.

The synergy between TOP 5'UTR element and albumin 3'UTR element was quantified by dividing the signal from mRNA combining both elements by the sum of the signal from mRNA lacking both elements plus the rise in signal effected by the TOP 5'UTR element plus the rise in signal effected by the albumin 3'UTR element. This calculation was performed for total protein expressed from 0 to 72 hours calculated from the area under the curve (AUC) (see following Table 4).

Table 4:

TOP 5'UTR	Synergy with albumin 3'UTR
RPL35	2,25
RPL21	1,30
atp5a1	3,19
HSD17B4	2,18
AIG1	2,03
COX6C	1,56
ASAH1	1,84

The synergy thus calculated specifies how much higher the luciferase level from mRNA combining TOP 5'UTR elements and albumin 3'UTR element is than would be expected if the effects of TOP 5'UTR element and albumin 3'UTR element were purely additive. The luciferase level from mRNA combining TOP 5'UTR element and albumin 3'UTR element was up to three times higher than if their effects were purely additive. This result confirms that the combination of TOP 5'UTR element and albumin 3'UTR element effects a markedly synergistic increase in protein expression.

5.4 TOP 5'UTR elements from mouse genes increase protein expression from mRNA.

To investigate the effect of TOP 5'UTR elements from mouse genes on protein expression from mRNA, mRNAs with two different mouse TOP 5'UTR elements were synthesized. In addition, mRNAs contained the albumin7 3'UTR element. Luciferase encoding mRNAs were transfected into human dermal fibroblasts (HDF). For comparison, mRNA containing the human RPL32 TOP 5'UTR element was transfected. Luciferase levels were measured at 24, 48, and 72 hours after transfection (see following Table 5 and Figure 30).

Table 5:

5'UTR	RLU at 24 hours	RLU at 48 hours	RLU at 72 hours
none	114277	121852	68235
32L	332236	286792	114148
m21L	798233	351894	139249
m35AL	838609	466236	174949

- Luciferase was clearly expressed from mRNA lacking a 5'UTR element. Both mouse
 5 TOP 5'UTR elements strongly increased the luciferase level, similarly as the human TOP
 5'UTR element.

SEQUENCES:

- 10 Homo sapiens alpha-2-macroglobulin (A2M): gctccttcttttctgcaacatg
 (Seq ID No: 1)
- Homo sapiens acyl-CoA dehydrogenase, C-4 to C-12 straight chain
 (ACADM):
 ggctctctttccgcgctgcggtcagcctcggcgtccacagagagggccagaggtggaacgcaga
 aaaccaaaccaggactatcagagattgcccgagaggggatg (Seq ID No: 2)
- 15 Homo sapiens arylsulfatase E (chondrodysplasia punctata 1) (ARSE):
 cttcctcttcttgatcggggattcaggaaggagcccaggagcagaggaagtagagagagagacaac
 atg (Seq ID No: 3)
- Homo sapiens Bruton agammaglobulinemia tyrosine kinase (BTK):
 20 tgtccttctctcttgactgtaagaatatgtctccagggccagtgtctgctgcatcgagtccac
 cttccaagtctggcatctcaatgcattctgggaagctacctgcattaagtcaggactgagcacaca
 ggtgaactccagaaagaagaagctatg (Seq ID No: 4)
- Homo sapiens complement component 2 (C2):
 tgaccttttccctccgcggctctctacctctcgcccccctagggaggacaccatg
 (Seq ID No: 5)
- 25 Homo sapiens cyclin-dependent kinase 4 (CDK4):
 gggcctctctagcttgcggcctgtgtctatggtcgggccctctgctccagctgctccggaccgag

ctcgggtgtatggggccgtaggaaccgggtccggggcccgataacgggcccggcccccacagcacc
cgggctggcgtgaggggtctcccttgatctgagaatg (Seq ID No: 6)

5 Homo sapiens cytochrome P450, family 17, subfamily A, polypeptide
1 (CYP17A1): agctcttctactccactgctgtctatcttgccctgccggcaccagccaccatg
(Seq ID No: 7)

Homo sapiens endoglin (ENG):
cttcctctacccggttggcaggcggcctggcccagcccccttctctaaggaagcgcatcttctgcct
ccctgggcccggcgggctggatg (Seq ID No: 8)

10 Homo sapiens excision repair cross-complementing rodent repair def
iciency, complementation group 3 (ERCC3):
tcttctctctgctgctgtagctgccatg (Seq ID No: 9)

Homo sapiens excision repair cross-complementing rodent repair def
iciency, complementation group 5 (ERCC5):
15 ctgtcttttcttccgggaggcgggtgacagctgctgagacgtgttgacagccagagtctctccgcttta
atgcgctccattagtgccgtcccccactggaaaaccgtggcttctgtattatttgccatctttgt
tgtgtaggagcagggagggttctctcccggggtcctaggcggcgggtgcagtcgcgtagaagaat
tagagtagaagttgtcggggccgctcttaggacgcagccgcctcatg (Seq ID No: 10)

Homo sapiens ferritin, light polypeptide (FTL):
20 cgtcccctcgcagttcggcgggtcccgcgggtctgtctcttgcttcaacagtgtttggacggaacag
atccggggactctcttccagcctccgaccgccctccgatttctctccgcttgcaacctccgggac
catcttctcggccatctcctgcttctgggacctgccagcaccgtttttgtggttagctccttcttg
ccaaccaaccatg (Seq ID No: 11)

Homo sapiens galactosylceramidase (GALC):
ccgcctccctggggcgccggagtcagtgtagccacacaatg (Seq ID No: 12)

25 Homo sapiens gap junction protein, alpha 1, 43kDa (GJA1):
ttttctttcattagggggaaggcgtgaggaaagtacaaacagcagcggagttttaaacttttaaat
agacaggtctgagtgccctgaacttgcccttttcttttacttcatcctccaaggagttcaatcactt
ggcgtgacttcactacttttaagcaaaagagtgggtgccaggcaacatg (Seq ID No: 13)

30 Homo sapiens gap junction protein, beta 1, 32kDa (GJB1):
cattctctgggaaaggcagcagcagccagggtgtggcagtgacagggaggtgtgaatgaggcagga
tg (Seq ID No: 14)

Homo sapiens glucose-6-phosphate isomerase (GPI):
cgctccttccctcctcggtcgcgtctcactcagtgtagcttctagtccccgccatg
(Seq ID No: 15)

35 Homo sapiens hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase
/enoyl-CoA hydratase (trifunctional protein), alpha subunit
(HADHA): ctgtcctcttcagctcaagatg (Seq ID No: 16)

Homo sapiens hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase
/enoyl-CoA hydratase (trifunctional protein), beta subunit
40 (HADHB):
gggcccttttctgggcaggacccgccccttgggtcccgagagccttgggtacttggacctgaaccttg
ctccgagagggagtcctcgcggacgtcagccaagattccagaatg (Seq ID No: 17)

Homo sapiens complement factor H (CFH):

cttccttttgcagcaagttctttcctgcactaatcacaattcttggaagaggagaactggacgttg
tgaacagagtttagctggtaaattgtcctcttaaaagatccaaaaaatg (Seq ID No: 18)

Homo sapiens sarcoglycan, gamma

5 (35kDa dystrophin-associated glycoprotein) (SGCG):

agcccttttctccagggacagttgctgaagcttcattcctttgctctcattctgtaagtcataaaaa
gtttgaaacattctgtctgtggttagagctcgggccagctgtagttcattcgccagtggtgttttct
taatatctaagatg (Seq ID No: 19)

Homo sapiens lipase A, lysosomal acid, cholesterol esterase

10 (LIPA):

gggtcccttatccgcaccccgccctgagagctggcactgcgactcgagacagcggcccggcagga
cagctccagaatg (Seq ID No: 20)

Homo sapiens lipoprotein lipase (LPL):

15 cccctcttctcctcctcaagggaaagctgccacttctagctgccctgccatcccctttaaagg
gcgacttgctcagcgccaaaccgcggtccagccctctccagcctccggctcagccggctcatcag
tcggtccgcgccttgagctcctccagagggacgcgccccgagatg (Seq ID No: 21)

Homo sapiens mutL homolog 1, colon cancer, nonpolyposis type 2

(E. coli) (MLH1): ggctcttctggcgccaaaatg (Seq ID No: 22)

Homo sapiens Niemann-Pick disease, type C1 (NPC1):

20 cttccttctcctgaccggcgcgcgagcctgctgccgcggtcagcgccctgctcctcctccgctcc
tcttgcgcggggtgctgaaacagccggggaagtagagccgctccggggagcccaaccagccgaa
cgccgcccggcgtcagcagccttgcgcgccacagcatg (Seq ID No: 23)

Homo sapiens peroxisomal biogenesis factor 12 (PEX12):

25 ggcctctcttccgcccagcatcccagaggtcctgggtggtttcatttccgggtgcggcttctgtca
taaagcggagacctcccttcaaacgtggcgctgctgggttgtttgcgccctgcctgggggtcagcgag
caaggacgggcgcgggcggggatactcaaagccaacagctggagtcagcccttggtgtcccggtc
acagtggcacgactgaatcctcagagtcgggtggttttgagctctcacgattggggaggaggggg
cgtttctggttcgcagctccagaggattgcgttcttccccatacctgtccccacagtcacgct
ctgccctgacgtgcagcatttgacaagttacccctcgccacatactacttccacccacgtccgag
30 ttaactttgttcttaaccttcttgagactacctcggcctccaggtctttttttccagttcattt
ttgcccataagattgagtttgcagtttccagatatcatgcagaaagtttacctttaagactgagcac
ccatctgatactcttctcccgaaaaagttcatgctcacgagagagtttgtgggaaaagtgaagc
cagtacacgcaggaaactatg (Seq ID No: 24)

Homo sapiens peroxisomal biogenesis factor 6 (PEX6):

35 cgctccttcacctcctcgttggtgtcctgtcaccatg (Seq ID No: 25)

Homo sapiens phosphofructokinase, muscle (PFKM):

gagccttcttgtcagcatctgttagtgagggttggaagcctctcctccttccccctcctctttg
cctccacctggctcctccccatgttcgtccatccccctccccctttcccaaggacaatctgcaa
gaaagcagcggcgaggagagctaagactaaaagagtggatcatg (Seq ID No: 26)

40 Homo sapiens serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 (SERPINA1):

ctgtctcctcagcttcaggcaccaccactgacctgggacagtgaatcgacaatg
(Seq ID No: 27)

Homo sapiens phosphatase and tensin homolog (PTEN):

agttctctcctctcggaaagctgcagccatgatggaagtttgagagttgagccgctgtgagggcaggg
ccgggctcaggcgagggagatgagagacggcgggcgccgagccctctcagcgctgt
gagcagccgcgggggcagcgccctcggggagccggcgccgctgcggcgggcagcgggcggttt
5 ctcgcctcctctctcgtcttttctaaccgtgcagcctcttctcctcggttctcctgaaagggaaaggtg
gaagccgtgggctcggggcgagccggtgaggcgggcgggcgggcgggcgccacctcccgtcctgg
agcggggggggagaagcgggcgggcgggcgggcgggcggtgcagctccaggagggggtctgagt
cgctgtcaccatttccagggtgggaacgcggagagttggtctctcccttctactgcctccaa
cacggcgggcgggcgggcgggcgccacatccagggacccggggcggttttaaacctcccgtccgcccgcg
10 ccgaccccccggtggcccggtcctcgaggccgcccggcgaggcagccgttcggaggattattcgt
cttctccccattccgctgccgcccgtgccaggcctctgggtgctgaggagaagcaggcccagtcgc
tgcaaccatccagcagccgcccagcagccattaccggctgcggtccagagccaagcggcgggcag
agcgaggggcatcagctaccgccaagtccagagccatttccatcctgcagaagaagccccgccacc
agcagcttctgccatctctctcctcttcttcttcagccacaggctcccagacatg
15 (Seq ID No: 28)

Homo sapiens solute carrier family 3

(cystine, dibasic and neutral amino acid transporters, activator of
cystine, dibasic and neutral amino acid transport), member 1
(SLC3A1): cctcccttactgcaggaaggcactccgaagacataagtgcggtgagacatg
20 (Seq ID No: 29)

Homo sapiens aldehyde dehydrogenase 3 family, member A2 (ALDH3A2):

ccgcctcccactcccagcgcccccgaccgtgcagttctctgcaggaccaggccatg
(Seq ID No: 30)

Homo sapiens bleomycin hydrolase (BLMH):

gtttctcccagcctcagcctccccgcgcccgcgcccgcgcccgcgagccggtttcctttttc
25 cggcgctccgggtgcgagagacaggtcggggccccctaggcagcgagccgcagcgcaatcccggcgc
tcgccaaggaccctggaagctaccgttacccccgcgggcagcggtgggcgcatg
(Seq ID No: 31)

Homo sapiens cathepsin K (CTSK):

cctcctcctcttaccctaaatttccagccgatcactggagctgacttccgcaatcccgatggaata
30 aatctagcaccctgatggtgtgccacactttgctgccgaaacgaagccagacaacagatttcca
tcagcaggatg (Seq ID No: 32)

Homo sapiens GM2 ganglioside activator (GM2A):

gcttctttgcgtaaccaataactggaaggcattttaaaggcacctctgccgccacagaccttgagtt
35 aactccgcccagaccacccttcccgatg (Seq ID No: 33)

Homo sapiens hydroxysteroid (17-beta) dehydrogenase 4 (HSD17B4):

ccgcctcctcctgtcccgcagtcggcggtccagcggtctgcttggttcgtgtgtgtcgttgagg
ccttattcatg (Seq ID No: 34)

Homo sapiens neutrophil cytosolic factor 2 (NCF2):

ctctctctgcttcttttcttttctctctcatggtagggttatgagtcagttgccaaaagggtgggga
40 catttctgatgcatttgcaacactgagaagttatcttaaggagggtgggccccatttctactcat
ctggcccagaaagtgaacaccttgggggcccactaaggcagccctgctaggggagacgctccaacct
gtcttctctctgtctcctggcagctctcttggcctcctagtttctacctaattcatg
(Seq ID No: 35)

Homo sapiens 3-oxoacid CoA transferase 1 (OXCT1):

cagcctcctcctgcctcaccgcccgaagatg (Seq ID No: 36)

Homo sapiens sulfite oxidase (SUOX):

ccgcccccttctcgagaactcgagagctgggctggtaaaattgcagtgctgaagacactggacccg
caaaaggctgtccctcccaaactgggattctgggctcactgagttcacctgcgagtcagccctac
ctgcactgctctggtctagtacaaacaggctgctggcattgagggacggagtcctccaactcctggc
5 ctctagcagtcctcctgtgtaggtctcccaaagtgtagtgtgtccggaattggtgggttcttggg
ctcactgacttcaagaatgaagccgcggaccctcgagctctgtacaatg (Seq ID No: 37)

Homo sapiens albumin (ALB): ttttctcttctgtcaacccacacgcctttggca-
caatg (Seq ID No: 38)

Homo sapiens arylsulfatase A (ARSA):

10 ctccctctagcgcttccccccggcccgactccgctggtcagcgccaagtgacttacgcccccgac
cctgagccccggaccgctagggcgaggaggatcagatctccgctcgagaatctgaagggtgccttggc
ctggaggagtccgctcccagccccgggtctcccggtactgtcgggccccggccctctggagcttca
ggaggcgccggtcaggggtcggggagtatttgggtccgggtctcaggggaaggcgccctgggtc
15 tgcggtatcggaagagcctgctggagccaagtagccctccctctcttgggacagacccctcggtc
ccatg (Seq ID No: 39)

Homo sapiens elastin (ELN):

ctccctccctctttccctcacagccgacgaggcaacaattaggctttggggataaaaacgaggtgcg
gagagcgggctggggcatttctccccgagatg (Seq ID No: 40)

20 Homo sapiens hemoglobin, alpha 2 (HBA2): cactcttctggtccccaca-
gactcagagagaacccaccatg (Seq ID No: 41)

Homo sapiens hexosaminidase B (beta polypeptide) (HEXB):

cttcctctgatccgggcccggcggaagtcgggtcccgaggctccggctcggcagaccggcgga
agcagccgagcggccatg (Seq ID No: 42)

Homo sapiens mannosidase, alpha, class 2B, member 1 (MAN2B1):

25 cggcctttccagggccggggaacccaggaggaagctgctgagccatg (Seq ID No: 43)

Homo sapiens recombination activating gene 2 (RAG2):

cactctctttacagtcagccttctgcttgccacagtcatagtgggcagtcagtgaatcttcccaa
gtgctgacaattaatacctgggttagcggaagattcagagaggcgtgagcagccctctggcct
tcagacaaaaatctacgtaccatcagaaactatg (Seq ID No: 44)

30 Homo sapiens CD53 molecule (CD53):

tctcctttttacacaaatagccccgatatctgtgttaccagccttgtctcggccacctcaaggata
atcactaaattctgccgaaaggactgaggaacggtgcctggaaaagggaagaatatcacggcatg
(Seq ID No: 45)

Homo sapiens Fc fragment of IgG, low affinity IIIa, receptor

35 (CD16a) (FCGR3A): tggctccctttagggtccggatatctttgggtgacttgtccactccag-
tgtggcatcatg (Seq ID No: 46)

Homo sapiens interleukin 1, beta (IL1B):

aaacctcttcgaggcacaaggcacaacaggctgctctgggattctcttcagccaatcttcattgct
caagtgtctgaagcagccatg (Seq ID No: 47)

40 Homo sapiens CD4 molecule (CD4):

ctgtctctcttcatttaagcacgactctgcagaaggaacaaagcaccctccccactgggctcctgg
ttgcagagctccaagtcctcacacagatacgctgtttgagaagcagcgggcaagaaagacgcaag

cccagaggccctgccatttctgtgggctcaggtccctactggctcaggcccctgcctccctcggca
aggccacaatg (Seq ID No: 48)

Homo sapiens serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 5 (SERPINA5):

- 5 agccctctgcccttcttgagcccgaggactgccacctccactgtgtgcacactcagctacgggac
acatttcaggtatccaaggcagcagaggtgagtgggtcccccagctctgtgaccttatgtccac
actaactctggcagagcctccgtttctcatagaacaaagaacagccaccatg
(Seq ID No: 49)

Homo sapiens vitronectin (VTN):

- 10 tgccctccttccctgtctctgcctctccctcccttcctcaggcatcagagcggagacttcagggag
accagagcccagcttgccaggcactgagctagaagccctgccatg (Seq ID No: 50)

Homo sapiens aldehyde dehydrogenase 9 family, member A1 (ALDH9A1):
ccgcccctcccgcgccccgccccctcccgcgccccgtcagcctctgccgaggagctgcgtccgcc
ctcatg (Seq ID No: 51)

- 15 Homo sapiens annexin A1 (ANXA1):

cttcctttaaaatcctataaaatcagaagcccaagtctccactgccagtgtgaaatcttcagagaa
gaatttctcttttagttctttgcaagaaggttagagataaagacactttttcaaaaatg
(Seq ID No: 52)

- 20 Homo sapiens ATPase, Na⁺/K⁺ transporting, alpha 1 polypeptide
(ATP1A1): ttttctctctgattctccagcgacaggaccggcgccgggactgagcaccgc-
caccatg (Seq ID No: 53)

Homo sapiens ATPase, Na⁺/K⁺ transporting, alpha 2 polypeptide
(ATP1A2):

- 25 ctttctctgtctgccagggtctccgactgtcccagacgggctgggtgtgggcttgggatcctcctgg
tgacctctcccgttaaggtccctcagccactctgccccaaagatg (Seq ID No: 54)

Homo sapiens calcium channel, voltage-dependent, beta 3 subunit
(CACNB3):

- 30 ccctccttcgcgctctctcgctccctgccgcccgcagggctgcggggctcggtggcatctccc
gggcgcgccccgcagtccttgcccctgcctccgggcccgtcccgcccccgcgccgctcgctcccc
cgaccgggactccccatg (Seq ID No: 55)

Homo sapiens cholinergic receptor, nicotinic, alpha 7 (neuronal)
(CHRNA7):

gtgcctctgtggcgcaggcgcaggccccggcgacagccgagacgtggagcgcgcccggctcgctgc
agctccgggactcaacatg (Seq ID No: 56)

- 35 Homo sapiens cytochrome P450, family 51, subfamily A, polypeptide
1 (CYP51A1):

gcttctctcgttccgtcgattgggaggagcgggtggcgacctcggccttcagtgtttccgacggagt
gaatg (Seq ID No: 57)

Homo sapiens glutamate decarboxylase 1 (brain, 67kDa) (GAD1):

- 40 atctctctcttctcctggcgctcgctgagaggggaactagcgagaacgaggaagcagctggagg
tgacgccgggagattacgcctgtcagggccgagccgagcggatcgctgggctgtgagaggaa
aggcgggagtgcccggctcgctgtcgagagccgagcctgtttctgcgcccggaccagtcgaggact
ctggacagtagaggccccgggacgaccgagctgatg (Seq ID No: 58)

Homo sapiens gamma-glutamyl carboxylase (GGCX):

aattctcctggcggcctccgttcagacgcggcagctgtgacccacctgcctcctccgcagagcaat
g (Seq ID No: 59)

Homo sapiens glutamate receptor, metabotropic 3 (GRM3):

5 tccccctctttccccaacctcctcctctcttctactccacccctccgttttccactccccactga
ctcggatgcctggatgttctgccaccgggcagtggtccagcgtgcagccgggagggggcaggggca
gggggcaactgtgacaggaagctgcgcgcacaagttggccatttcgagggcaaaataagttctcct
tggatttggaaaggacaaagccagtaagctacctcttttgtgtcggatgaggaggaccaaccatga
gccagagcccggtgcaggctcacgcgcgcgtgccaccgcggtcagctccagttcctgccagga
10 gttgtcgggtgcgaggaattttgtgacaggctctgttagtctgttcctcccttatttgaaggacagg
ccaaagatccagtttggaaatgagagaggactagcatgacacattggctccaccattgatatctcc
cagaggtacagaaacaggattcatgaagatg (Seq ID No: 60)

Homo sapiens guanylate cyclase 1, soluble, alpha 3 (GUCY1A3):

15 ggttcctttggggtgatcaaagagggagacacagacacagagagacaaaggcaaggaggactgtct
gggagccacgcgggcgatacagtttccgaggcacgcgcgtcccgctagcctgttgaacaggtag
acatgagcgacccaagctgcggatttgcgaggcgcgcctggagctgctagagatccggaagcaca
gccccgaggtgtgcaagccaccaagtcaagttcctaacgagtcttcagaggaggcagcaggaagc
tcagagagctgcaaagcaaccgtgcccattctgtcaagacattcctgagaagaacatacaagaaagt
cttcctcaaagaaaaaccagtcggagccgagtcctatcttcacactttggcagagagtatttgcaaa
20 ctgattttcccagagtttgaacggctgaatgttgcaacttcagagaaacattggcaagcacaata
aaagaaagcaggaaatctttggaaagagaagactttgaaaaacaattgcagagcaagcagttgca
gcaggagttccagtgagggttatcaaagaatctcttggtgaagagggtttttaaataatgttacgag
gaagatgaaaacatccttgggggtggttgaggcaccccttaaagattttttaaagcgttcagtacc
cttctgaaacagagcagccattgccaaagcaggaaaaaggggcaggcttgaggacgcctccatt
25 ctatgcctggataaggaggatgattttctacatgtttactacttcttccttaagagaaccacctcc
ctgattcttcccgcatcataaaggcagctgctcacgtattatatgaaacggaagtggaagtgtcg
ttaatg (Seq ID No: 61)

Homo sapiens 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR):

30 ggctccttccgctccgcgactgcgttaactggagccaggctgagcgtcggcgccggggttcgggtg
cctctagttagatctggaggatccaaggattctgtagctacaatg (Seq ID No: 62)

Homo sapiens IMP (inosine 5'-monophosphate) dehydrogenase 2

(IMPDH2): aggtctctgcggcgcggtcctcgagacacgcggcggtgtcctgtgttgccatg
(Seq ID No: 63)

Homo sapiens leukotriene A4 hydrolase (LTA4H):

35 acttcctttccggcgtgcaccgcgaatccctcctcctcttctttacctctctccctcctcctcag
gttctctatcgacgagctctggtagctgagcgttgggctgtaggtcgctgtgtgtgatcccca
gagccatg (Seq ID No: 64)

Homo sapiens neuropeptide Y receptor Y1 (NPY1R):

40 ccttctttaataagcaggagcgaaaaagacaaattccaaaggaggattgttcagttcaagggaatga
agaattcagaataattttggtaaatggattccaatatggggaataagaataagctgaacagttgac
ctgctttgaagaaacatactgtccatttgtctaaaataatctataacaaccaaaccaatcaaaatg
(Seq ID No: 65)

Homo sapiens pyruvate dehydrogenase (lipoamide) beta (PDHB):

cggccctctgttgcgtttggcagcggatagaggacacgaccaagatg (Seq ID No: 66)

5

10

15

20

25

30 Homo sapiens wntless homolog (Drosophila) (WLS):

35

ttttctctttc aaactgcccagacggttg gacaggagctagacacacagaagaaaagaagacaaag
aacgggttaggaaaattaaaaagggttaccatg (Seq ID No: 73)

40

Homo sapiens structural maintenance of chromosomes 4 (SMC4):

45

ggcgggtgataatcgccgcactttttttcaaattagtggtatcccagaaatcattgcgcgcatttgta
acgaattttccggttcgagtttgatatttttaggcgccattttcgagtgaaggacccggagccgaaacac
cggtaggagcgagggtgggtactacacaaccgtctccagccttggtctgagtggactgtcctgc
agcgacctg (Seq ID No: 75)

- 5 Homo sapiens GLE1 RNA export mediator homolog (yeast) (GLE1):
tggccttcccggcggtgattcgagggcttggttggtcagaagggggcgtcagagaagctgcccc
ttagccaacctg (Seq ID No: 76)

- Homo sapiens tripartite motif containing 6 (TRIM6):
gagtctttcggcctgggtggaggacgcggctgcttcaagtccttggtctgatccaggccacagat
10 tccaggattctacaggcaggaaacatcttagaaatcagggttgggcaggcaggagccaggagagta
gctacaatg (Seq ID No: 77)

- Homo sapiens ecotropic viral integration site 2A (EVI2A):
tattcttttttactgcagattttactttaaggctcatattctccaagtctattctgctttaaaaaga
agacaagaaaagaagtggtttatcaaaatcacgttataatcagattttgaccaagcattttgtaag
15 tatacaaatgtcagccaatgacatataacaaccattttcttataaaaccttgatgttcaaaagcctg
actagcagtggcatccatg (Seq ID No: 78)

- Homo sapiens heterogeneous nuclear ribonucleoprotein L (HNRNPL):
tgctcttttcgatccgggacggccggctcaggctcgccgcccagctggagaactacgatgaccgcga
caaaacccctgcctccccagttgtccacatcaggggctgattgacgggtgtggtggaagcagacct
20 tgtggaggccttgaggagtttggaacctatcagctatgtggtggtaatg (Seq ID No: 79)

- Homo sapiens mitochondrial translational initiation factor 2
(MTIF2):
cattcttccgggtccagaaggatctccgcccgtgctcagaatccagggggccggggctgtagat
tccttgacaaggatatcctagcggcgaaacaacaccgtactgggagtcagaacgtctgggttctag
25 tcttgactgccattaactagcgggtatgacattggagaagcttttttgacccttctggatttccgtt
tccttttctgtaaaatgaggagcttggaagatccggaaaatgaggcccataggaaacaagtgactt
gctgagtccagataacactgactgtcagagagaaacatg (Seq ID No: 80)

- Homo sapiens nuclear factor of kappa light polypeptide gene enhanc
er in B-cells inhibitor, zeta (NFKBIZ):
30 tggcctcctcttgccacgaggtcagacggcgagttcttagagaaaaaggctgcttagctgctgctt
atcatgtaacctcaaaaggaaactgatcgcttttctcatgctgtcacgtacttgggttattatcgc
tgattacagctggaacaattgatttgccttactgatttgtgtgacttgactcttcaaacacaaa
ggttaacaggaagatctcgagggccctggctgaacttcaccttttggtttcttggcctgatgctg
aactctcgaggttgagccccatag (Seq ID No: 81)

- 35 Homo sapiens v-erb-b2 erythroblastic leukemia viral oncogene homol
og 3 (avian) (ERBB3):
atccctccccggactccggctccggctccgattgcaatttgcaacctccgctgccgtcgccgcagc
agccaccaattcgccagcgggtcaggtggctcttgccctcgatgtcctagcctaggggcccccgggc
cggacttggctgggctcccttcaccctctgcggagtcatg (Seq ID No: 82)

- 40 Homo sapiens podoplanin (PDPN): ccgcctcctcgggagagataaatg
(Seq ID No: 83)

Homo sapiens ribonucleotide reductase M1 (RRM1):
gcgccccctttgtgcgtcacgggtggcgggcgcggaaggggatttgattgttgcgcctctgctct
gaagaaagtgtgtctggtccaactccagttctttccctgagcagcgccctggaacctaacctt

cccactctgtcaccttctcgatcccgcgcggcgctttagagccgcagtcagtccttgatccttcag
agcctcagccactagctgcgatg (Seq ID No: 84)

Homo sapiens solute carrier family 2 (facili-
tated glucose transporter), member 4 (SLC2A4):

5 gcgctttttccccagccccgctccaccagatccgcgggagccccactgctctccgggtccttggc
ttgtggctgtgggtcccatcgggcccgcctcgcacgtcactccgggacccccgcggcctccgcag
gttctgcgctccaggccggagtcagagactccaggatcggttctttcatcttcgccgcccctgcgc
gtccagctcttctaagacgagatg (Seq ID No: 85)

10 Homo sapiens steroid-5-alpha-reductase, alpha polypeptide 1
(3-oxo-5 alpha-steroid delta 4-dehydrogenase alpha 1) (SRD5A1):
aaccctttctgcagagtcgggcagtgccggactccggtagccgccctccggtagccgcccctcc
tgccccgcgcgcgcgcctatatgttgccgcgcgcgcctctggggcatggagcacgctgccag
ccctggcgatg (Seq ID No: 86)

15 Homo sapiens thromboxane A synthase 1 (platelet) (TBXAS1):
gttcccttttctacctgcagagcacggttcccataagggcggcgagatcagcctcctgtctcatct
ggaagaccaccactctgggtctcagaggaatg (Seq ID No: 87)

Homo sapiens transketolase (TKT):

ctatctctgtgtgtccgcgtgtgcgcccgggtccccgcctgccgcaccatg (Seq ID No: 88)

20 Homo sapiens tumor necrosis factor receptor superfamily, member 1A
(TNFRSF1A):

cctcctcctccagctcttcctgtcccgctgttgcaaacactgcctcactcttcccctcccaccttct
ctcccctcctctctgctttaattttctcagaattctctggactgaggctccagttctggcctttgg
ggttcaagatcactgggaccaggccgtgatctctatgcccgagtctcaaccctcaactgtcacccc
aaggcacttgggacgtcctggacagaccgagtcggggaagccccagcactgccgctgccacactg
25 ccctgagcccaaatgggggagtgagaggccatagctgtctggcatg (Seq ID No: 89)

Homo sapiens tubulin, beta 2A class IIa (TUBB2A):

aggtctctgcgcagcccagcccgcgggtccacgcgcgcaccgctccgagggccagcgccaccgcg
tccgcagccggcaccatg (Seq ID No: 90)

30 Homo sapiens actin, beta (ACTB): tcgcctttgccgatccgcgcggcggtcca-
caccgcgcgcagctcaccatg (Seq ID No: 91)

Homo sapiens adenylosuccinate synthase (ADSS):

ggctccttcttctctctgcatgtggctggcgccgcagagcagttcagttcgctcactcctcgccgg
ccgcctctccttcgggctctcctcgcgtcactggagccatg (Seq ID No: 92)

Homo sapiens alanyl (membrane) aminopeptidase (ANPEP):

35 cgttctctgcctggcctgaggctccctgagccgcctccccaccatcaccatg
(Seq ID No: 93)

Homo sapiens beaded filament structural protein 1, filensin
(BFSP1):

40 gcctcctttctttctcagcccagacctggccctctggagagggttttggagtcctgggtaggcagg
gtacctcaggcagcaggcagcacaccttggatgtgagctgaatggattttcaaatttcacagaagg
agcctccatgctggagaaagtatgtatg (Seq ID No: 94)

Homo sapiens basic transcription factor 3 (BTF3):

cggcctcccttttagctgccatcttgcgctccccgcgtgtgtgcgcctaattctcaggtgggtccaccg

agaccccttgagcaccaaccctagtagtccccgcgcggcccccttattcgctccgacaagatg
(Seq ID No: 95)

Homo sapiens complement component 1, q subcomponent binding protein (C1QBP): ttgtcctttgcatctgcacgtgttcgcagtcgtttccgcgatg
5 (Seq ID No: 96)

Homo sapiens calsequestrin 1 (fast-twitch, skeletal muscle) (CASQ1):

10 tttccttttcttaatatggcgatgagctcttaggccagtggtggggaccggggctgaggtgccctgga
cactggaggagggggaggggaaggagccccctgggagcctggggtagaagtgtaggaggtgggaggat
tccgccccgcgatggagctgtcctggcctcagaaggttatccgtctctcctgccaaccatggagaca
tatttagacaggaccaggtggggactgaggggtgccaatcagggggcagctccggttccctccc
cgccccctgtcctatttctccacctgacctttttcccttggtctctgtcggcagtttctccagga
cccagcagtgccctctgtccactgctctgggccattccccaatccccctcccacttgagccccta
actcagaatctgggacccaggggccccctccctacccagctaacctcttctggaccaggagagcca
15 acccagatcccactacctccatg (Seq ID No: 97)

Homo sapiens caveolin 3 (CAV3):

gtctctctgccccctctctgcccccaagtattttcagccccagccggccacacagctcggatctcctc
ctgtggatccccccagctctgcgatg (Seq ID No: 98)

20 Homo sapiens serpin peptidase inhibitor, clade H
(heat shock protein 47), member 1, (collagen binding protein 1)
(SERPINH1):

aggtcttttggttttttttggcgagctggggcgccctccggaagcgtttccaactttccagaagtt
tctcgggacgggcaggaggggtggggactgccatatatagatcccgggagcaggggagcgggcta
agagtagaatcgtgtcgcggctcgagagcgagagtcacgtcccggcgctagcccagcccagccag
25 gccaccgtggtgcacgcaaaccacttcttgccatg (Seq ID No: 99)

Homo sapiens CD68 molecule (CD68):

tttcctcctttccaagagagggctgagggagcaggggtgagcaactggtgcagacagcctagctgg
actttgggtgaggcgggttcagccatg (Seq ID No: 100)

30 Homo sapiens cell division cycle 20 homolog (S. cerevisiae)
(CDC20):

gggtcccttttctgtccctgagcacctgcctcctttcctccagggctccgtaggcaccaactgc
aaggacccctccccctgcgggcgtcccatg (Seq ID No: 101)

Homo sapiens cadherin 13, H-cadherin (heart) (CDH13):

35 gagcctctcctcaaagcctggctccacggaaaatatgctcagtcagccgcgtgcatgaatgaaa
acgccgcggggcgcttctagtcggacaaaatg (Seq ID No: 102)

Homo sapiens regulator of chromosome condensation (RCC1) and BTB
(POZ) domain containing protein 2 (RCBTB2):

40 cgctcccttcgtttccgtctcggccgggcacccgagcgcacccgcccaggccgggcccgtttcagg
gggaggcgccaactcatcgcggcgccgggccccctgaccgtgcagtaaccgctaccagggaggcgga
gcggaacaaggctccggcctgcgaggagtcacattaactttgctctagaagacaactttacaaggat
ctaaaaggaacaggattaaagatgactgaatactgggttccagaaatttaaaacaatcagcttagc
aaatcatatatattcttctgtggagctgagaattgatgtccgctcttccccgtgatttggaactttcc
aatcccagagaaaagttgacaaagggactgccaggactgagtcacatg
(Seq ID No: 103)

Homo sapiens cold inducible RNA binding protein (CIRBP):

ccccccctcactcgcgcgtaggaggtcgggtcgttggtgcgctgtcttcccgcttgcgctcag
ggacctgcccgactcagtggccgcatg (Seq ID No: 104)

Homo sapiens LIM domain binding 2 (LDB2):

5 cctcctctcctctccctctcctctcctgctatagagggctccgacagcagttcccagccagcgtgt
tcagcctgcctgcctgcctgcctctgtgtgtgtgtgagcgtgtgtgcgtgcgtctactttgtactg
ggaagaacacagcccatgtgctctgcatggacgttactgatactctgtttagcttgattttcgaaa
agcaggcaagatg (Seq ID No: 105)

Homo sapiens chloride channel, nucleotide-sensitive, 1A (CLNS1A):

10 ctgcctcttccagggcgggcggtgtggtgcacgcattgctgtgctccaactccctcagggcctgtg
ttgccgcactctgtgctatg (Seq ID No: 106)

Homo sapiens collapsin response mediator protein 1 (CRMP1):

15 cctcctccttctcccgccctcctcgcgcgacccgggcggtgctggcagccggagcggcgggcgggcg
gccgagcagccggggcagccgcgcgtgggcatccacgggcgcgagcctccgtccgtgtctctatc
cctcccgggcctttgtcagcgcgcccgtgggagcggggccgagagcgcgggttccagtcagacag
ccccgcaggtcagcggccgggcccagggcgccagagggggccatg (Seq ID No: 107)

Homo sapiens catenin (cadherin-associated protein), delta 1
(CTNND1):

20 ttgcctttggctgggtgcaacttccatttttaggtgttggatctgagggggaaaaaaaagagagagg
gagagagagagaaagaagagcaggaaagatcccgaaaggaggaagaggtggcgaaaaatcaactgc
cctgctggatttgtctttctcagcaccttggcgaagccttgggtttctttcttaaaggactgattt
ttagaactccacatttgaggtgtgtggcttttgaagaaaatgtatgtactgacgggaaaaggagga
taagcaagtcgaatttttgtcttacgctctctccttccctgcttccctccttgcgtgtggtggctggga
tgcttcttccatgattttttgaatctagactgggctgttctctgtgttaaaccaatcagttgcgac
25 cttctcttaacagtgtgaagtgggggtctctctccctccttctccttctctgtgattcacctt
cctttttaccctgcccgtgcggcggtccgccccttaccttcatg (Seq ID No: 108)

Homo sapiens diacylglycerol kinase, alpha 80kDa (DGKA):

30 ccgtcccctccagcccagctcgggctccagctccagcgcggcgcttcagctgcgaccgcgagccc
tctcaagcaagatataacttccccaaagtcacacagtggtatcagagctaagaatgggaccagata
tgactgatctagtctgtttccaaaaccgtgctgtattatattaacgcctaccctctgaagaggtcc
aagcaacggaagtactactacgaagctgcctttctggccatccttgagaaaaatagacagatgagt
tcttgccagtgccttaggcctccatctctctcccttgcgtgtaccaccttcaccaccatccatg
cgaccccaagagccttaagtactctagaagagactccaggcaggggaagctgaaaggacctttcac
tccctacttttggccagggccttctgtgccacctgccaagaccagcaggcctaccctctgaagagg
35 tccaagcaacggaagtactactacgaagctgcctttctggccatccttgagaaaaatagacagatg
(Seq ID No: 109)

Homo sapiens aspartyl-tRNA synthetase (DARS):

40 cgatctttctggagccgcacctccacgcggagtcgagcgcgtgtgctgagaccccagggtcggga
gggcggagactgggagggagggaagccctttggcctgccttacggaagcctgcgagggagggt
ggtgtccactgccagttccgtgtcccgatg (Seq ID No: 110)

Homo sapiens dynein, cytoplasmic 1, intermediate chain 2
(DYNC1I2):

45 agttcttctcgatcgtgtcagtttgtaaggcagggcggaagttggattcctggcctgagaatatt
aggcgtagttttccagtttttggcaaagcggaataacttaaggccctgggttgactgggttcttt
gttttatctaccggccttctgctttacgacaggtcacaaacatg (Seq ID No: 111)

Homo sapiens dedicator of cytokinesis 1 (DOCK1):

tttcctccccatcctgtcgcggctcgaaaggaatggaaaatggcggcctagacgcggagtttcctg
cccgaaccgcggcggctccggcggcgccatg (Seq ID No: 112)

Homo sapiens dihydropyrimidinase-like 2 (DPYSL2):

5 ctctctcttttttttccgccctagctggggctgtgttgaggagaggaagaaagagagacagagga
ttgcattcatccggttacgttcttgaaatttcctaataagcaagaccagcgaagcgggttgacccttt
tcaatcttgcaaaggaaaaaaacaaaacaaaaaaacccaagtccccttcccggcagtttt
tgccttaaagctgccctcttgaaattaatttttccaggagagagatg (Seq ID No: 113)

Homo sapiens developmentally regulated GTP binding protein 2
(DRG2):

10 tgttctcttttggttccggggcgacgctactctgtcgcggccgtcagaccggaattgccggtgccg
ccgccaccgctgtctgtgcgccacctctgtctgtaccatg (Seq ID No: 114)

Homo sapiens eukaryotic translation elongation factor 1 alpha 1
(EEF1A1):

15 cgttctttttcgcaacgggtttgccgccagaacacaggtgtcgtgaaaactaccctaaaagccaa
aatg (Seq ID No: 115)

Homo sapiens eukaryotic translation elongation factor 1 gamma
(EEF1G):

20 tctcctctttccccctcccttctctccggggcggttactttgcggcagcgccgagaaccccaccc
cctttctttgcggaatcaccatg (Seq ID No: 116)

Homo sapiens eukaryotic translation initiation factor 2, subunit 3
gamma, 52kDa (EIF2S3): atttccttctcttttggcaacatggcgggc
(Seq ID No: 117)

Homo sapiens eukaryotic translation initiation factor 4B (EIF4B):

25 gggctcttttgcgttctctttccctctcccaacatg (Seq ID No: 118)

Homo sapiens eukaryotic translation initiation factor 4 gamma, 2
(EIF4G2): tattcttttgaagattcttcggtgtcaagccgccaagt
(Seq ID No: 119)

Homo sapiens epithelial membrane protein 1 (EMP1):

30 cttcccctcagtgcggtcacatacttccagaagagcggaccagggtgctgccagcacctgccact
cagagcgctctgtcgtggtggacccttcagaactctctttgctcacaagttaccaaaaaaaaaaaga
gccaacatg (Seq ID No: 120)

Homo sapiens fibrillarin (FBL):

35 cgctcttttccacgtgcgaaagccccggactcgtggagttgtgaacgccgcggactccggagccgc
acaaaccagggtcgccatg (Seq ID No: 121)

Homo sapiens exostoses (multiple)-like 2 (EXTL2):

40 ctgtcccttgctccaggcgctcacttttgcgggcggcactttttccaggttgtaataccagctaagt
gagaaggatagatgcacgctacttggtttagaaaaaaaaaacaataatgacaaacgagacgccccct
tccgttttatgataactaagctgcagggaaataaatcggtggccctactgcaatctactgcactc
gagaacatcacagaaaattctttgatttatcttaatagtgacaagtgagcctgcttctgtcaatt
actgaagctataaggagattttttaaaattaaacttcaacacaatg (Seq ID No: 122)

Homo sapiens solute carrier family 37 (glu-
cose-6-phosphate transporter), member 4 (SLC37A4):

ccgcctctgttcaggacactgggtccccttggagcctccccaggcttaatgattgtccagaaggcg
gctataaagggagcctgggaggctgggtggaggagggagcagaaaaacccaactcagcagatctg
ggaactgtgagagcggcaagcaggaactgtggtcagaggctgtgcgtcttggctggtaggcctgc
tcttttctaccatg (Seq ID No: 123)

5 Homo sapiens GDP dissociation inhibitor 2 (GDI2):
agccctcccctcctcgctccctcccctcctcctcccgcagttcttctcttcccgtctgaggtgg
cggtcggtctcgccttgtcgccagctccattttcctctcttctcttcccctttccttcgcgcccc
agagcgctcccagcctcgtagggtgggtcacggagcccctgcgccttttcttctgctcgggtcctgc
gtccgcgcctgccccgcatg (Seq ID No: 124)

10 Homo sapiens UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, p
olypeptide 1 (B4GALT1): cacccttcttaaagcggcgggcgggaagatg
(Seq ID No: 125)

Homo sapiens GDP-mannose 4,6-dehydratase (GMD5):
15 ggccctcccctgcacggcctcccgtgcgcccctgtcagactgtggcgggccggtcgcgcggtgcgctc
tccctcccctgcccgcagcctggagaggcgcttcgtgctgcacacccccgcgttccctgcgggcaccg
cgctgcctctgcgcgctccgcctgcccgcgaccgcacgcccgcgcgggacatg
(Seq ID No: 126)

Homo sapiens histone deacetylase 2 (HDAC2):
20 ggccccctcctcgcgagttggtgccgctgccacctccgattccgagctttcggcacctctgccggg
tggtaccgagccttcccggcgccccctcctctcctcccaccggcctgcccttcccgcgggactat
cgccccacgtttccctcagcccttttctctccggccgagccgcggcggcagcagcagcagcagc
agcagcaggaggaggagcccgggtggcgcggtggcgggggagcccatg (Seq ID No: 127)

Homo sapiens protein arginine methyltransferase 2 (PRMT2):
25 gggccttcccggctgacggcctgcgtgcactgcgcttgcgcggttgagggcggtggctcaggctc
ctggaaaggaccgtccacccctccgcgctggcggtgtggacgcggaactcagcggagaaaacgcgat
tgagagcagtggtggtgattacactatcactggaaaaatacgaattgagaagaaggaaaagactgga
agatgcagaccttggttccctgtagtggaacactgtaagggtcccagaaatggaaaagaaaatgaa
ataaatcagcagttatgaggcagagcctaagagaactatg (Seq ID No: 128)

30 Homo sapiens immunoglobulin (CD79A) binding protein 1 (IGBP1):
gttctctctccccaagatg (Seq ID No: 129)

Homo sapiens eukaryotic translation initiation factor 3, subunit E
(EIF3E): actcccttttctttggcaagatg (Seq ID No: 130)

Homo sapiens activated leukocyte cell adhesion molecule (ALCAM):
35 gtccctctactcagagcagcccggagaccgctgccgcccgtgccgctgctaccaccgctgccacct
gaggagaccgcgcgcccccccgctgcgcgcctcctgcgagtccttcttagcacctggcgtttcatgc
acattgccactgccattattattatcattccaatacaaggaaaataaaagaagataccagcgaaaa
gaaccgcttacacctttccgaattactcaagtgtctcctggaaacagagggtcggtgtccccggag
gagcagccgaaggggcccggtgggtggtggtgacggggaggaggaggagttgggggacattgcgtgg
tggaagttgcgtgcggcagagaaccgaagggtgcagcgccacagcccaggggacggtgtgtctggg
40 agaagacgtgccctgcgtcgggacccgccagcgcgcgggcacccgcggggcccgggacgacgccc
cctcctgcggcggtggactccgtcagtgggccaccaagaaggaggaggaatatg
(Seq ID No: 131)

Homo sapiens acyloxyacyl hydrolase (neutrophil) (AOAH):
45 ttttctttatcctgcagtcctttacctcagcagaaccgcacaccacagactccctccagctctttgt
gtgtggctctctcagggtccaacaagagcaagctgtgggtctgtgagtgtttatgtgtgcttttat

tcacttcacacttattgaaaagtgtgtatgtgagaggggtgggggtgtgtgtgtcaaagagagtgagg
aagagaaggagagagagatcaattgattctgcagcctcagctccagcatccctcagttgggagctt
ccaaagccgggtgatcacttgggggtgcatagctcggagatg (Seq ID No: 132)

Homo sapiens ADP-ribosylation factor 1 (ARF1):

5 ccgcccccttaccggtgccccgcgcccggaggcgctgacgtggccgcccgtcagagccgcatct
tgtgggagcaaaaccaacgcctggctcggagcagcagcctctgaggtgtccctggccagtgtcctt
ccacctgtccacaagcatg (Seq ID No: 133)

Homo sapiens ADP-ribosylation factor 6 (ARF6):

10 gcgccttttccggcagcgccgcccgcagaactgggaggaggagttggaggccggaggaggagcccgcg
ctcggggcgccggtggaggcagcgaccgagttcccgcgaggatccatgacctgacgggggccccg
gagccgcgctgcctctcgggtgtcctgggtcgggtggggagcccagtgtcgcaggccggcgggcg
gcccggagggtgcagtctccctcgcggtgagaggaaggcgaggagcggaaccgcggcgccgctc
gcgcggcgccctgcggggggaagggcagttccggggccggccgcgcctcagcaggcgccggtccc
15 agcgcagtctcaggggccgggtggcgccgactggagaaatcaagttgtcgggtcgggtgatgcc
cgagtgcggggggcctgggcctctgcccttaggaggcaactcccacgcaggccgcaaaggcgct
ctcgcggccgagaggttcgtttcgtttcgcggcgccggttgttggtgaggggacccgg
gacacctgaatgcccccgccccggctcctccgacgcgatg (Seq ID No: 134)

Homo sapiens ras homolog family member A (RHOA):

20 cgccctcccgccgcccgcgcctcgcctctcgcgctaccctcccgccgcccgcggctcctccgtc
ggttctctcgttagtccacggctctggtcttcagctaccgccttcgtctccgagtttgcgactcg
ggaccggcgctccccggcggaagaggctggactcggattcgttgctgagcaatg
(Seq ID No: 135)

Homo sapiens ras homolog family member G (RHOG):

25 cggcctcccgctctcaattccttctcgagcccggagccgctgcgcgccccccagctccccgcct
cggggaggggcaccaggtcactgcagccagaggggtccagaagagagaggaggcactgcctccacta
cagcaactgcacccacgatg (Seq ID No: 136)

Homo sapiens ATP synthase, H⁺ transporting, mitochondrial F1 compl
ex, O subunit (ATP5O): ctctcttcccactcgggtttgacctacagccgcccgggagaa-
gatg (Seq ID No: 137)

30 Homo sapiens B lymphoid tyrosine kinase (BLK):

ccacctctgtctgctgccggcagaaagccacaagccatgaaaactgattgagatgagaagaattca
tctgggactggcttttgccttaggatggtgttggaagttgctcgttgctagggagcctgctcca
ctgtaagggtgtcaggatctgaagagctatggtgaaacaccactgaagcattgccaaggatg
(Seq ID No: 138)

35 Homo sapiens B-cell translocation gene 1, anti-proliferative
(BTG1):

gcactctcttcgcctctcggagctggaaatgcagctattgagatcttcgaatgctgcggagctggag
gcggaggcagctggggagggtccgagcgatgtgaccaggccgcatcgctcgtctcttctctctcc
tgccgcctcctgtctcgaaaataacttttttagtctaaagaaagaaagacaaaagtagtcgtccgc
40 ccctcacgccctctcttctctcagccttccgcccgggtgaggaagccgggggtggtgctccgccg
tcggggccgcgcccggagccccagccgcccgggcccgcacgcgcccccatg
(Seq ID No: 139)

Homo sapiens calcium modulating ligand (CAMLG):

45 cggcctctagtcacgcctcgcagcgccgccaacatcaccgccactgccacccctcccagactg
tggacgggaggatg (Seq ID No: 140)

Homo sapiens calnexin (CANX):

aggcctcttggttctgcggcacgtgacggtcggggccgcctccgcctctctctttactgcggcgcg
ggcaaggtgtgcggggcggaaggggcacgggcacccccgcggtccccgggaggctagagatcatg
(Seq ID No: 141)

- 5 Homo sapiens calpain 2, (m/II) large subunit (CAPN2):
cgacctttctctgcgcagtagcgccgcccgggaccgcagcatg (Seq ID No: 142)

Homo sapiens caveolin 1, caveolae protein, 22kDa (CAV1):

gcgccctttttttccccccatacaatacaagatcttccttcctcagttcccttaaagcacagcccag
ggaaacctcctcacagttttcatccagccacggggccagcatg (Seq ID No: 143)

- 10 Homo sapiens CD1d molecule (CD1D):
cgacctctttgcagctcgcacagctaagggcgagggcgcccttcggcagaagcagcaaaccgcccgg
caagcccagcgaggagggctgccggggtctgggcttgggaattggctggcaccagcggaagggga
cgtgagctgagcggcgggggagaagagtgcgcaggtcagagggcggcgcagcggcgctccgcga
ggtccccacgcccggcgatatg (Seq ID No: 144)

- 15 Homo sapiens CD22 molecule (CD22):
tctccttttctctcagatgctgccagggtcctgaagaggggaagacacgcggaaacaggcttgca
cccagacacgacaccatg (Seq ID No: 145)

Homo sapiens CD37 molecule (CD37):

- 20 cttcctcttttggggttcttcctttctctctcagctctccgtctctctttctctctcagcctcttt
ctttctccctgtctccccactgtcagcacctcttctgtgtggtgagtgagcgttaccacca
ggtgaagatg (Seq ID No: 146)

Homo sapiens CD38 molecule (CD38):

gcctctctcttctgctgcctagcctcctgccggcctcatcttcgccagccaaccccgccctggagccc
tatg (Seq ID No: 147)

- 25 Homo sapiens CD48 molecule (CD48):
cggcctttttctagccaggctctcaactgtctcctgcggttgctgggaagtctggaaggaagcatg
(Seq ID No: 148)

Homo sapiens chromogranin B (secretogranin 1) (CHGB):

cttcctttccgcacaggggcccgcgagcggggccatg (Seq ID No: 149)

- 30 Homo sapiens chloride channel, voltage-sensitive 3 (CLCN3):

ttccccttccgtgggtcagggccgggtccgggtccggaacctgcagcccctttcccagtggtctagtt
cgcccgtagcccggaataatgagcaaggagggtgtggtgggttgaaagccatcctactttactccc
gagttagagcatggattcagtttttagtcttaagggggaagtgcagattggagatttttattttta
tttgggcagaagcaggttgactctagggatctccagagcgagaggatttaacttcatgttgctccc
35 gtgtttgaaggaggacaataaaaagtcccaccgggcaaaaattttcgtaacctctgcggtagaaaacg
tcaggatatcttttaaatcgcgatagttttcgctgtgtcaggcctttcttcggtggagctccgaggg
agctaggttctaggtttgaaacagatgcagaatccaaaggcagcgcaaaaaacagccaccgatttt
gctatgtctctgagctgcgagataatcagacagctaaatg (Seq ID No: 150)

Homo sapiens colipase, pancreatic (CLPS):

- 40 ttccccttccgtgggtcagggccgggtccgggtccggaacctgcagcccctttcccagtggtctagtt
cgcccgtagcccggaataatgagcaaggagggtgtggtgggttgaaagccatcctactttactccc
gagttagagcatggattcagtttttagtcttaagggggaagtgcagattggagatttttattttta
tttgggcagaagcaggttgactctagggatctccagagcgagaggatttaacttcatgttgctccc
gtgtttgaaggaggacaataaaaagtcccaccgggcaaaaattttcgtaacctctgcggtagaaaacg

tcaggtatcttttaaatcgcgatagttttcgctgtgtcaggctttcttcggtggagctccgagggg
agctaggttctaggtttgaaacagatgcagaatccaaaggcagcgcaaaaaacagccaccgatttt
gctatgtctctgagctgcgagataatcagacagctaaatg (Seq ID No: 151)

Homo sapiens cytochrome c oxidase subunit IV isoform 1 (COX4I1):
5 ctacccttttccgctccacgggtgacctccgtgcggccgggtgcgggaggagtcttcctcgatcccg
tggtgctccgcggcgccgcttgcctctcttcgggtgcgggacaccgggtgtagagggcggtcgcg
gcgggcagtgggcggcagaatg (Seq ID No: 152)

Homo sapiens cytochrome c oxidase subunit VIIc (COX7C):
10 ctttcttttcagtccttgcgacccggggaacaaggctcgtgaaaaaaaaaggctcttggtgaggtgccg
ccatttcattctgtcctcattctctgcgccttttcgcagagcttccagcagcggtatg
(Seq ID No: 153)

Homo sapiens activating transcription factor 2 (ATF2):
cagccttttctccaggggtgctttgtaaacacggctgtgctcagggctcgcgggtgaccgaaagg
atcatgaactagtgacctggaaagggtagtagatggaaacttgagaaaggactgcttattgataac
15 agctaaggtattcctggaagcagagtaaataaagctcatggcccaccagctagaaagtattcttgc
catgagaaaaagaatgtgataagttattcaacttatg (Seq ID No: 154)

Homo sapiens casein kinase 1, alpha 1 (CSNK1A1):
agatccctttccagagtgtctctgcgcctgagaagaagcggtcccggggactgggggcattttgtg
20 ttggctggagctggagtaacaagatggcgctcgtccgcggagtgcaggggtccctctgggcccggag
ccggcggcagtggtggcagcggtatcgccgccctagctcaccgcgccccctttccagcccgcgacg
tcgccgcgcaagcagggcagcggcggccgagagaaacaagtggcccagcctggttaaccgcccgaga
agcccttcacaaactgcggcctggcaaaaagaaacctgactgagcggcggtgatcaggttcccctc
tgctgattctgggccccgaacccccggtaaaaggcctccgtgttccgtttcctgccgccctcctccgt
25 agccttgccctagtgtaggagccccgaggcctccgtcctcttcccagaggtgtcggggccttgcccc
agcctccatcttctgtctctcaggatg (Seq ID No: 155)

Homo sapiens catenin (cadherin-associated protein), beta 1, 88kDa
(CTNNB1):
aagcctctcgggtctgtggcagcagcggttgccccggccccgggagcggagagcagggggaggcggag
acggaggaaggctctgaggagcagcttcagtcccccgcgagccgccaccgcaggtcgaggacggtcg
30 gactcccgcggcgaggagcctgttccccctgagggtatttgaagtataccatacaactgttttga
aatccagcgtggacaatg (Seq ID No: 156)

Homo sapiens dCMP deaminase (DCTD):
ccgcctcctccccgacttccttccctgagcacggcgggcgggggacgagcaccggcctgcgcgc
ggagccggcaccggatgaccaacatg (Seq ID No: 157)

Homo sapiens damage-specific DNA binding protein 1, 127kDa (DDB1):
35 ctgtcttttctgcttgtgtccctctttctagtgctcgcgctcgagtcccgcagggccgctccaagcct
cgacatg (Seq ID No: 158)

Homo sapiens desmin (DES):
ctgtctccccctcgccgcattccactctccggccggccgctgcccgccgctcctccgtgcgccccg
40 cagcctcgccccgcgcccgcacatg (Seq ID No: 159)

Homo sapiens deoxyhypusine synthase (DHPS):
cgttccctacttctgtgtcttgcggagacgcgcgcgtcggggtttaacgcgtttctgggcccgc
gtaagcccggcctaggggcagctttgactcgagagccggctataggcgcatg
(Seq ID No: 160)

Homo sapiens dihydrolipoamide S-acetyltransferase (DLAT):

caccctttcggatgcctcccctagaaccctaccactttccacccctttccgctctgttatttctccc
aaacttgcgcccgacagggccctctggaacactcctgccccgtagtgccctcgtccccgctccg
tagagaaagagcgtgcggtgcccgcattttctggcctggggagcgggtggagtaaacctgcgggaac
5 cattttacgacaacgtgcggtgtgcggtgtggctgacggcaacgccgctgctcttgagaggtca
ctccggagacggcggttggttttgggggtgtggggggttggtggcactatg (Seq ID No: 161)

Homo sapiens down-regulator of transcription 1, TBP-binding (nega-
tive cofactor 2) (DR1):

ccttccctggcatctggagggaccacgggttgccgcgtcttcggcttccacgatctgcgttcgggct
10 acgcggccacggcgccagccactgcgaactcccactgtgcctggctctgtccatattagttcccagg
cgccgctcgccgttccagcagcggcagcggcagcggcagcggcgacatgttgtagggcgccggcg
cgggtgtctgaaggatggtttggccgaggcgccggaacggctgctggcgccggcgccagcggcag
cggggcctcgggctctatagagccgagcccgctgggtaccgcgggtaccgcggcgaggccagtg
cccctggatcttgctctgctccgacgcggttggggaccagttaggcgacagcggccgcccctctg
15 aggagacacgaaggtggttccccagccgctcaaatttccggaccaccgcgctttccctcctcagc
ctgggctgtgctctctctagaatcctcgggcccccactttcttccaaactcatcctaaatctctc
acacacgcgagtggttccagccctcaagccagctgctcctccgttcattttctgcaccctcttcgc
aaagcaccggcgatcactctccgagggcgactttttgagaaatctcggtggagtagtggaacca
gagctggggagtttttaaaagccggggcgcgagaaacaggaaggtactatg
20 (Seq ID No: 162)

Homo sapiens endothelin receptor type A (EDNRA):

ttttctttttcgtgcgagccctcgcgcgcgctacagtcacccgctggtctgacgattgtggaga
ggcgggtggagaggcttcatccatcccaccggctcgctcgccggggattgggggtcccagcgagacctc
cccgggagaagcagtgcccaggaggttttctgaagccggggaagctgtgcagccgaagccgcggcc
25 ggcgggagcccgggacaccggccaccctccgcgccaccaccctcgccggtccgggttctctg
gcccaggcgccgcgggacccggcagctgtctgcgcacgccgagctccacggtgaaaaaaaagtga
aggtgtaaaagcagcacaagtgcataagagatatcttctcaaatttgccctcaagatg
(Seq ID No: 163)

Homo sapiens eukaryotic translation elongation factor 1 alpha 2
(EEF1A2):

cagtccctctggctgagacctcggctccggaatcactgcagccccctcgccctgagccagagcac
30 cccgggtcccgcagccctcacactcccagcaaatg (Seq ID No: 164)

Homo sapiens eukaryotic translation elongation factor 2 (EEF2):

cgttctcttccgcgctcgctcgcccatcctcggcgcgactcgcttctttcggttctacctgggag
35 aatccaccgccatccgccaccatg (Seq ID No: 165)

Homo sapiens eukaryotic translation initiation factor 4A2

(EIF4A2): ctgtcttttcagtcgggcgctgagtggtttttcggatcatg

(Seq ID No: 166)

Homo sapiens egf-like module containing, mucin-like, hormone recep-

40 tor-like 1 (EMR1): gtttcttttctttgaatgacagaactacagcataatg
(Seq ID No: 167)

Homo sapiens enolase 2 (gamma, neuronal) (ENO2):

gcgctcctccgcccgcggccgggagccgcagccgcggccgactgccactcccgtctctcag
cgccgcgctcgccaccgccaccgccaccgccactaccaccgtctgagctctgagctcccagatccc
45 agccatcatg (Seq ID No: 168)

Homo sapiens esterase D (ESD):

ccgccttttacttcggcccgcttcttctggtcactccgccaccgtagaatcgctaccatttggtg
caagcaaaaagcaatcagcaattggacaggaaaagaatg (Seq ID No: 169)

Homo sapiens Finkel-Biskis-Reilly murine sarcoma virus

5 (FBR-MuSV) ubiquitously expressed (FAU):

cttcctcttttctcgactccatcttcgcggtagctgggaccgccgttcagtcgccaatatg
(Seq ID No: 170)

Homo sapiens Friend leukemia virus integration 1 (FLI1):

10 ctgtctcttttcgctccgctacaacaacaaacgtgcacaggggagtgagggcagggcgctcgaggg
ggcagcgagggagggcccagggcgccagggaggccgcgcccgggctaatacgaaggggctgcgaggt
caggctgtaaccgggtcaatgtgtggaatattggggggctcggtcgagacttggccaaatg
(Seq ID No: 171)

Homo sapiens fibromodulin (FMOD):

15 gccccttttcacaatatattgattaggaatttggggcgggaccctgggtctggcacaggcacgcacac
tctcagtagactctttcactcctctctctcttctctctcacacgttctccaacccaaggaggcca
gacagagggacgtgggtcactctctgaaaagttcaacttgagagacaaaatg
(Seq ID No: 172)

Homo sapiens ferritin, heavy polypeptide 1 (FTH1):

20 cgttcttcgccgagagtcgtcggggtttcctgcttcaacagtgttgagcgaacccggcgctcgt
tccccaccccgccggccgcccatagccagccctccgtcacctcttcaccgcacccctcgactgcc
ccaagggcccccgcgcgctccagcgccgcagccaccgcgcgcgcgcctctccttagtcg
ccgccatg (Seq ID No: 173)

Homo sapiens glyceraldehyde-3-phosphate dehydrogenase (GAPDH):

25 cgctctctgctcctcctgttcgacagtcagccgcacatcttcttttgctcgccagccgagccacatc
gctcagacaccatg (Seq ID No: 174)

Homo sapiens glycyl-tRNA synthetase (GARS):

caccctctctggacagcccagggccgcagggtcatg (Seq ID No: 175)

Homo sapiens glutamic-oxaloacetic transaminase 2, mitochondrial
(aspartate aminotransferase 2) (GOT2):

30 ctgtccttaccttcagcaggagccggttccctgtgtgtgtgtccgctcgccctctgctccgtcctg
cggctgcccactgcctcctacggtccaccatg (Seq ID No: 176)

Homo sapiens general transcription factor IIF, polypeptide 1, 74kD
a (GTF2F1):

35 gcgcctcttccggttaccttttcccagcgccagaggcgccctaggggttggggctctcgctcaggcac
agagaccgcagaccgagcgggcgttccccgggatcgagggacgcgcacgccagaggagacgaaag
gaaccggggtcggaccagatcggaaccactgaccattgccatg (Seq ID No: 177)

Homo sapiens glycogen synthase 1 (muscle) (GYS1):

40 cggcctccttctgcctaggtcccaacgcttcggggcaggggtgcggtcttgcaataggaagccgag
cgtcttgcaagcttcccgctgggaccagctactcgccccgcaccctacctgggtgcattccctag
acacctccggggtccctacctggagatccccggagcccccttctcgccagccatg
(Seq ID No: 178)

Homo sapiens major histocompatibility complex, class I, C (HLA-C):

cattctccccagaggccgagatg (Seq ID No: 179)

ctctccccctcctccccctcccgctccaagattcgccgcccgcggccgcgcaggagtagc
cgccgccggagccgcgcgcagccatg (Seq ID No: 190)

Homo sapiens keratin 19 (KRT19):

gctcctcccgcgaatcgcagcttctgagaccagggttgctccgtccgtgctccgcctcgccatg
(Seq ID No: 191)

Homo sapiens laminin, beta 1 (LAMB1):

5 attcccttctttgggctcgggggctcccggagcagggcgagagctcgcgctcgccggaaggaagac
gggaagaaagggcagggcggtcggcgggcggtcttctccactcctctgccgcgtccccgtggctgca
gggagccggcatg (Seq ID No: 192)

Homo sapiens ribosomal protein SA (RPSA):

10 ctgtcttttccgtgctacctgcagaggggtccatacggcggttggtctggattcccgctcgtaactta
aagggaattttcacatg (Seq ID No: 193)

Homo sapiens lymphocyte cytosolic protein 1 (L-plastin) (LCP1):

ttttctttcctggctgatgattgtcattctagtcacttccctgccttgtagccacacaccagggt
tgacaaagctgttctgcagatcagaaagaaggggttcctgggtcatacaccagtactaccaaggaca
gcttttttctgcaagatctgttacctaagcaataaaaaatg (Seq ID No: 194)

15 Homo sapiens lectin, galactoside-binding, soluble, 1 (LGALS1):

ccatctctctcgggtggagtcttctgacagctggtgcgcctgcccgggaacatcctcctggactca
atcatg (Seq ID No: 195)

Homo sapiens SH2 domain containing 1A (SH2D1A):

20 ttctctcttttttgcacatctggctgaactgggagtcagggtggttgacttgtgcctggctgcagta
gcagcggcatctcccttgacagttctcctcctcggcctgcccgaagagtcaccaggccatg
(Seq ID No: 196)

Homo sapiens mannosidase, alpha, class 2A, member 1 (MAN2A1):

25 tgttccttttccctccgcttctctgacctagctgcgcggccccggccccgggagctgccgaacccgc
gcctcccctgggtgaggaggacacgcctgcctcgtcgagaaaacttttctgcccactcagttgg
ggcggcggtggcaggaagtgcggggcagcgacctctcctccgctgccccgcgcgcctgcccggagg
tcggcgctgagcttgcatcaagtttggtggggcccccttcccagttgccggcgagctctgcctc
gagaggggcgcccagccccggggagggcggcaggccagggcgaaggccaagggcggtgtggtggcgc
cggagactaggtgcggagcaaggcggggactcgcacccgcatccgagagcgcggaggtcgcgcagc
ccgggagaagggagcctccggcggtgcttcctagagtcacagtgcgctgtctcctttggctgag
30 gagagtgtcctggccccgagtcctatcgaggaaaatg (Seq ID No: 197)

Homo sapiens myelin basic protein (MBP):

ccgcctcttttcccagatgccccggggagggaggacaacaccttcaaagacaggccctctgagtc
cgacgagctccagaccatccaagaagacagtgccagccacctccgagagcctggatgtgatg
(Seq ID No: 198)

35 Homo sapiens melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor) (MCLR):

cattcttcccaggacctcagcgcagccctggcccaggaaggcaggagacagaggccaggacggtcc
agaggtgtcgaaatgtcctggggacctgagcagcagccaccagggaagaggcagggaggagctga
ggaccaggcttggttgtagaatccctgagcccaggcggtagatgccaggaggtgtctggactggc
40 tgggccatgcctgggctgacctgtccagccaggagaggggtgtgagggcagatctgggggtgccca
gatggaaggaggcaggcatgggggacaccaaggccccctggcagcaccatgaactaagcaggaca
cctggaggggaagaactgtggggacctggaggcctccaacgactccttctgcttcctggacagga
ctatg (Seq ID No: 199)

Homo sapiens malic enzyme 1, NADP(+)-dependent, cytosolic (ME1):

45 gggcctttcccagtgcgggccgcccggccacagctgcagtcagcaccgtcaccacagcagcatccg

ccgcctgcaccgcgcgtgcggcccgccccggcctgacccccgcgcggaacccggcgccagccatg
(Seq ID No: 200)

Homo sapiens myocyte enhancer factor 2C (MEF2C):

5 agctctctgctcgtctcgtcgcagtcacagacacttgagcacacgcgtacaccagacatcttcg
ggctgctattggattgactttgaaggttctgtgtgggtcgccgtggctgcatgtttgaatcaggtg
gagaagcacttcaacgctggacgaagtaaagattattgttatttttttttctctctctctct
ctcttaagaaaggaaaatatcccaaggactaatctgatcgggtcttccttcattcaggaacgaatgc
aggaatttgggaactgagctgtgcaagtgtgaagaaggagatttgtttggaggaaacaggaaaga
gaaagaaaagggaagaaaaatacataatttcagggacgagagagagaagaaaaacggggactatg
10 (Seq ID No: 201)

Homo sapiens mannosyl (alpha-1,3)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase (MGAT1):

15 agcccttcttggggaagtcagctaccagcagcctgtagtctcctcggtacccaccctcaccgcctg
gggtcccatggtgagacagctgggtgggcatcaggcttctgcagagggccaggccggaggagctg
ggcgaggagagtggtggtcctggttgcaccggcctcgtggaatccaggcctcagacctgac
gctggcgaaactggctctgtgcgtggagccccctggtcttctgcgtctgtcctcctccggccaga
ctttactcctggctcagcgacaggtatttgctatggaagagctgtccctccctccctcgggtgggc
ctgggtccacctccacctcctcttcagggtccgcaccttcccttttaaaacaccagccggggcg
20 agaccgttctaggcttttccatgggtgcttccgcaaagcttgtgaccgagtccttccgcctagg
gctggtgggcctccctgctggtaggctctctcttcgcttttcttactcagaactgaagctctcatt
cccacccaccaaggaaaaacaaaagggaagaagccacagctggccccggcttgctttggcacagg
tgtttccccccggccccccgctcgggcaccttggttctgttctgtccctgccccacgcgacctgg
ggctcccacccgggtcctcagcctccctgggttgggggtggggggaactggctcccagcccttggc
25 ctagggtttggtgaacgccttcttggaactgcggggccacttcaggcgcggtccaggctgggcag
ctgcgtgagggccgagggcaggggtgggtcgggcgtccacctcagggttgccgacaggagcc
ggaaagccgactcccgaagtgggtcctgggaaaacttgggtcctgggtgactgagaagcggcg
gggaaaggagggcgggccaggagggggcctggcgagcgcggccggggggcggggcgcgcggg
gctgtcgggtcacgccccctcagtcgccccgccccgccccgctgcggggaagggccaggttgccc
30 gccccggcgtccggccccggcgcgccgcagaaagggtggcgagtcgaaaggcgaggcgccgcgg
cagcgttgggacgcgcctgggcaccgggctcgtccctgcgccccggagcaggccaagttcgggg
ccaggacgtcgggaggacctggtgcatggctgcctcctaataccatagtcagaggagcatccct
aggactgcgggcaaggagccgggcaagcccagggcagccttgaaccgtcccttggtgcctcc
ccggtgggggacaggatg (Seq ID No: 202)

35 Homo sapiens mitogen-activated protein kinase kinase kinase 11 (MAP3K11):

ctgcctcccgcggggccaaagtacaaaggaggaggaagaaggagcggggtcggagccgtc
ggggccaaaggagacggggccaggaacaggcagtcctcgcccaactgcggagcctccctccacccc
ctgcgcaaaaagacccaaccggagttgaggcgtgcccctgaaggccccaccttacacttggcggg
40 ggccggagccaggctcccaggactgtccagaaccgagggaaagctcgggtcctccaagctagcca
tggtgaggcgccggaggccccggggccccaccccccggtgaccacactgccttgggtgcctc
ctccagaagcccagatgcggggggcggggagacaacactcctggctcccagagaggcgtgggtc
tggggctgaggggcaggggccggatgccaggttcggggactagggccttggcagccagcggggt
ggggaccacgggcacccagagaaggtcctccacacatcccagcgccgggtcccgccatg
45 (Seq ID No: 203)

Homo sapiens membrane protein, palmitoylated 1, 55kDa (MPP1):

ccgccttctccgcagccccgcaggccccgggcccctgtcattcccagcgtgcctgtcttgcgttc
cagtgttccagcttctgcgagatg (Seq ID No: 204)

Homo sapiens v-myc myelocytomatosis viral oncogene homolog (avian) (MYC):

5 ggccctttataatgcgaggggtctggacggctgaggacccccgagctgtgctgctcgcgccgcccac
cgccggggccccggcgcgtccctggctccccctcctgcctcgagaagggcagggttctcagaggcttg
gcgggaaaaaagaacggaggggagggatcgcgctgagtataaaagccgggttttcggggctttatctaa
ctcgctgtagtaattccagcgagaggcagagggagcgagcgggcgccggctagggtggaagagcc
ggcgagcagagctgcgctgcgggcgctcctgggaagggagatccggagcgaatagggggcttcgcc
tctggcccagccctcccgctgatccccagccagcggtccgcaacccttgccgcattccacgaaact
ttgcccatagcagcgggcgggcacttttgactggaacttacaacacccgagcaaggacgcgactct
10 ccgacgcggggaggctattctgccatttggggacacttccccgcccgtgccaggaccgcttct
ctgaaaggctctccttgacgctgcttagacgctg (Seq ID No: 205)

Homo sapiens nuclear cap binding protein subunit 1, 80kDa (NCBP1):
tggcctctcggttccgcggcgcaccggagggcagcatg (Seq ID No: 206)

Homo sapiens necdin homolog (mouse) (NDN):

15 cttcctctccaggaatccgcggaggagcgaggtcgaagagctcctggacgcagaggccctgcc
cttgccagacggcgagacatg (Seq ID No: 207)

Homo sapiens NADH dehydrogenase

(ubiquinone) 1 beta subcomplex, 5, 16kDa (NDUFB5):

ccttcttctcctgcccgtagtagccatg (Seq ID No: 208)

20 Homo sapiens NADH dehydrogenase (ubiquinone) Fe-S protein 4, 18kDa
(NADH-coenzyme Q reductase) (NDUFS4):

ccgtcctttcatcctggcggttgctgcagcaagatg (Seq ID No: 209)

Homo sapiens nuclear factor of kappa light polypeptide gene enhanc
er in B-cells 2 (p49/p100) (NFKB2):

25 tgccctttccccggccaagcccaactccggatctcgtctctccaccggatctcaccgcccacaccg
gacaggcggttgaggaggcgggcggtctaaaattctgggaagcagaacctggccggagccactaga
cagagccgggcctagcccagagacatg (Seq ID No: 210)

Homo sapiens non-metastatic cells 2, protein (NM23B) expressed in
(NME2):

30 gccctctctccgcccgggtccccgggtgtggtggctgcaccagctctctgctctcccagcgcagc
gccgcccggcccccctccagcttcccggacctg (Seq ID No: 211)

Homo sapiens nucleophosmin (nucleo-
lar phosphoprotein B23, numatrin) (NPM1):

35 gcgtcctttccctgggtgtgattccgtcctgcgcggttggtctctggagcagcgttcttttatctcc
gtccgccttctctctacctaagtgcgtgccgccaccgatg (Seq ID No: 212)

Homo sapiens 5'-nucleotidase, ecto (CD73) (NT5E):

cattcctttttagaaaaaacccgtgcctcgaatgaggcgagactcagagaggaccaggcgcgggg
cggacccctccaattccttccctcgcgcccccgaaagagcggcgcaccagcagccgaactgccggcg
cccaggctccctgggtccggccgggatgcggccggtaccgctccccgcccgggaacaacctctccac
40 tcttctgacgggagctgggtgccagccgacagccgcgcaggccgctccaggctgaagaaggaccgccc
tcgggtgacgtcggaacttgccgctggccgccaagccggcctccaggctgaagaaggaccgccc
cggccttgaccggggccccgccccctccagccggggcaccgagccccggccctagctgctcgccct
actcgccggcactcgccgggtcgcccgctttcgcaaccagttcacgcgccacagctatg
(Seq ID No: 213)

Homo sapiens phosphatidylethanolamine binding protein 1 (PEBP1):
gcgctcttcccgagccagtgctgctgagctctccgcgtcgctctgtcgcccgcgctggcctaccgc
ggcactcccggtgcacgctctgcttggcctcgccatg (Seq ID No: 214)

Homo sapiens poly(A) binding protein, cytoplasmic 1 (PABPC1):
5 gcttcccccttctccccggcggttagtgctgagagtgcgagtggtgctccgggctcggaacacac
atttattattaaaaaatccaaaaaaatctaaaaaatcttttaaaaaacccccaaaaaatttaca
aaaaatccgcgtctcccccgccggagacttttatttttttcttctcttttataaaataaccgg
tgaagcagccgagaccgacccgccccgcccgcggccccgcagcagctccaagaaggaaccaagagac
cgaggccttcccgctgcccggaccgacaccgcccctcgctccccgcggcagccggcagccag
10 cggcagtggtatcgaccccggttctgcgggcgttgagtagttttcaattccggttgatttttgtccct
ctgcgcttgctccccgctccccctcccccggtcgcggccccagccccggcactcgctctcctcct
ctcacggaaaggtcgcggtctgtggcctgcgggcagccgtgccgagatg
(Seq ID No: 215)

Homo sapiens proprotein convertase subtilisin/kexin type 2
15 (PCSK2):
cgctctttctctccggtacacacagctccccacattcgacccctgcccgcgcgcccgggcccgcctg
actgcacggcttccccctccagccagatgctggagaacacacactgattcgctgctttccaagacc
tgttcagtctctttctctatacaaagatttttttaaaaactatatataagaattctttatttgcac
cctccctccgagtcctcctgctccgcccagcctgcgcgcctcctagcaccacttttcaactcccaaaga
20 aggatg (Seq ID No: 216)

Homo sapiens phosphogluconate dehydrogenase (PGD):
gggtctttccctcactcgctcctccgcgcgtcgccgctcttcggttctgctctgtccgcccgcctg
(Seq ID No: 217)

Homo sapiens phosphoglucomutase 1 (PGM1):
25 cgctccccctttccctcccgccggacctgccaggaggtgggctggcgcgaggaggggccctgtcc
cctgtccctttaaggaggaggggccaaacgcccggcctagagtgcggcgtagccccacccgcccgtgc
cctcaccccagagcagctgcagcctcagccggccgcccctccgcccagccaagtccgcccgtctgac
ccccggcagcaagtcgccaccatg (Seq ID No: 218)

Homo sapiens solute carrier family 25 (mitochon-
30 drial carrier; phosphate carrier), member 3 (SLC25A3):
cggcctctgtgagccgcaacctttccaagggagtggttggtgatcgccatcttagggagtgagtg
tggccgggcttctcctgtggcggttggtgggagcggagcccagagctcctgtggggccgctgctt
tggcgggtgggcccagccgggagcagcctctttcgaaggccgctgacctctcaagggcggtggag
acgggaagggaaaaggccccgggttggggtccagggcgccggtaacgttaaccggcgcccttgccgt
35 cctctaaccgctcgctccctcctccctagaaagatg (Seq ID No: 219)

Homo sapiens pim-1 oncogene (PIM1):
cctcccccttactcctggctgcggggcgagccgggcgtctgctgcagcggccgcggtggctgagga
ggcccagaggagtcggtggcagcggcgggcgggaccggcagcagcagcagcagcagcagcagc
agcaaccactagcctcctgccccgcggcgtgcgcacgagccccacgagccgctcaccccgccgt
40 tctcagcgtgcccgaacccgctggcgcgccctcccgccgagtcccggcagcgcctcagttgt
cctccgactcgccctcggccttccgcgccagccgcagccacagccgcaacgccacccgcagccaca
gccacagccacagccccaggcatagccttcggcacagccccgggtccgggtcctgcggcagctcct
ctgggcacccgtccctgcgcccagatcctggaggttgggatg (Seq ID No: 220)

Homo sapiens pyruvate kinase, muscle (PKM2):
45 ggatctcttcgtctttgcagcgtagcccagtcggtcagcgcggaggtgagcgggtgcaggaggct
acgccatcagtcccccaccaagggccagtcgcccggctagtgcggaatccccggcgcgccggccggcc
ccgggcacgcaggcaggcgggcgccagcatccagggcgtctgggatgcagtgaggctcagagagagg

agaacggctcctcacgcctggggcctgctcttcagaagtccccagcgccgttccttccagatcagg
acctcagcagccatg (Seq ID No: 221)

Homo sapiens pleiomorphic adenoma gene-like 1 (PLAGL1):

5 cggcctcctcggcgagccatcctcttggctgccgcggcgcaaagcccacggcatctgccattt
gtcattcagcccgctcgggtaccgccccgagccttgatttagacacggctggggcgctgctctggcctc
actctccgggcggtgctggacggacggacggacggggcagccgtgctcacagctcagcagcgcg
ggccttggcgcgcggggcgcttccccgggtcgccgtcatggccgcggaggtggcacgcccagcg
cctcgctgagctccgggggtcgtcgccccgcagggttgcgtgcacgtctaattgtggctgctgcc
10 tcgtgtcacatctgaaactcatctgtacctcacttagaaagtgggttctgattagacaagacttttc
gttgacgtcgacagaaacctaattgggaccattgaagaattccaaacagggtatttgcataaggaatca
gaggagttaatcttgtctcttctcacagggttgaaatcttcagacaaacttctgggaggactcggtc
cctgcctcgagcagatgttccctgtcactcagtaggcataatg (Seq ID No: 222)

Homo sapiens phospholipase D2 (PLD2):

15 tgctctcttggctccggaacccccgcgggcgctggctccgtctgccagggatg
(Seq ID No: 223)

Homo sapiens proteolipid protein 2 (colonic epithelium-enriched)
(PLP2):

cccccttcccgccagacggcgggcaagacagctgggtgtacagcgtcctcgaaaccacgagcaag
tgagcagatcctccgaggcaccagggtactccagcccatgccatg (Seq ID No: 224)

20 Homo sapiens pinin, desmosome associated protein (PNN):

cagtcctttcgcgcctcggcgggcgccatagcccggtcggcctgtaaagcagctctcaagcctgc
cgcagggagaagatg (Seq ID No: 225)

Homo sapiens phosphoribosyl pyrophosphate amidotransferase (PPAT):
ggtccttccacgtgctttcggcgggcagatg (Seq ID No: 226)

25 Homo sapiens protein phosphatase 1, catalytic subunit, gamma isozy
me (PPP1CC):

tgttcttctcgtgggttccagtggggagagaaggaggaagtagggagcgggggtggcaggggggggac
ccgcccgggtgctgccaccgcccaccaccgcctctgctcgtggcggtgggaaaggaggtgtgag
tcccgggcgcgagccggcgggcgccgctgcgggaggggtcggcggtgggaaggcgatg
30 (Seq ID No: 227)

Homo sapiens protein phosphatase 1, regulatory subunit 8 (PPP1R8):
cggctcttccagtttcccgcgctgcttagggcgcgccaaatgggagggggagacgcaagatg
(Seq ID No: 228)

Homo sapiens protein phosphatase 6, catalytic subunit (PPP6C):

35 cggcctccgcccgtgccgcccgcgtgctacagccgcccgcgctgttgccgcggcttggttatt
cttaaaatg (Seq ID No: 229)

Homo sapiens protein kinase C substrate 80K-H (PRKCSH):

ctttctttctgcagcaggaaccgcggctgctggacaagaggggtgcgggtggatactgacctttgct
ccggcctcgtcgtgaagacacagcgcattctcccgcgtgtaggttccctccacagaaccggtttcg
40 ggcctcagagcgtctggtgagatg (Seq ID No: 230)

Homo sapiens mitogen-activated protein kinase 6 (MAPK6):

cgcccccttctcctcgccctctctcgggggtcgggggttacatggcgggcactgcggcaaagcgaga
gcctcggagacgccgctgccgcccagcacagccggagacctgagccgacactgggggagtcgcgga
gccccgcactctctcgatgagtcggagaagtcccgttgatcagagtaagatggacggtagctttg

attgtgattgtggtgagctggagccacctgatcactaacaaaagacatcttctgttaaccaacagc
cgccagggtcttctgttgaaataaatatatagcaacaaaggaaaaaagaagcaaacggaaatag
tgcttaccagcaccttagaatgatgctgctcaggaccagtccaacactgaatgtatctgcactgtg
aggagaatgttcatagaagcctgttgtgtgcatattttattcacatttttgttaaattgttaaattcgt
5 ttagcacggtaattctgagtgcacagtatgtcattttcattccgtttgagtttcttgttttcgttaaa
tgtctgcagagttgctgccccctttcttgaactatgagtactgcaatctttttaattctcaatatga
atagagctttttgagcttttaaatctaagggggaactcgacaggcctgtttggcatatgcaatgaaca
tcaagaaaccatcttgcgtgtggaagcataattatttttcttctccctttttgaaagatctttcctt
ttgatgccagttttcttccctgtttacacaagttcaatttgaaaggaaaaggcaatagtaagggtt
10 tcaaaatg (Seq ID No: 231)

Homo sapiens phosphoribosyl pyrophosphate synthetase 2 (PRPS2):
cctcccccttccctacatctagccgcgcgctttcccgctcccgcagcagcagcctcccgcgtcgt
gtcgtgttgctccgcacctcctccgcgcgcgcgcgcgcctcggagttccgcgccccaccatg
(Seq ID No: 232)

15 Homo sapiens phosphoribosyl pyrophosphate synthetase-associated pr
oteins 1 (PRPSAP1):
ttgcctctggctctgaggcggcgccggcgccggcgctgcgaaggctcggccgctgtagtcagtgggtg
ggggtgcgaaggcgcagcagcctcggagctctcccgcgttgccgcagagtttctcagcgccttcccc
acccaaaccggggtctcgcagtcggaagcactcagagtgagccccgcgcggggccggtcgtaacc
20 gcgcgcggggccggacgatg (Seq ID No: 233)

Homo sapiens proteasome (prosome, macropain) subunit, beta type, 5
(PSMB5): agttctttctgcccacactagacatg (Seq ID No: 234)

Homo sapiens proteasome (pro-
some, macropain) 26S subunit, non-ATPase, 13 (PSMD13):
25 tgttcttctgtgccgggggtcttctctgctgtcatg (Seq ID No: 235)

Homo sapiens protein tyrosine phosphatase, receptor type, N
(PTPRN):
cagccccctctggcaggctcccgcacagcgtcgtcgcggctccggccccgggagcgcagcgcgccggagct
cggaagatg (Seq ID No: 236)

30 Homo sapiens RAB3A, member RAS oncogene family (RAB3A):
ctccctttgcaggacgtcacggaggactgcaggggctgagccgctgctgccgcgcgcgcgcgcga
gccccacatcaacgcaccggggtcctgtcaccgccaccgcaaaaaagtcaccgcgcgctagggtcg
ccgttgcatcggtgcagggaagatg (Seq ID No: 237)

35 Homo sapiens RNA binding motif, single stranded interacting protei
n 2 (RBMS2): ctctctctctctctctctctcgtcgttccctaacattaaagagaaaatg
(Seq ID No: 238)

Homo sapiens reticulocalbin 1, EF-hand calcium binding domain
(RCN1):
40 gcgccccctctgctccggctcggggcgggcactggcggagggactggccagtcccctcctccgcgcc
ggcccaaccctgtcgtgcccgcgcgtccgagtcctccattcccagctgcccgtgttgctcgtc
gctcagcgtctccctctcggccgcctctcctcgggacgatg (Seq ID No: 239)

Homo sapiens radixin (RDX):
ccgccttttccgcggaggcgcgcgagcggccatattgcggagctgtctgcggtgggcgggcgccct
ctcgtctccgcggcccagcgtcgcaccaccgcttctccctccctgtcgcagccgcgcgcgcgcgc

cagcgccccagccacacgcccggcgggcagaagccgcccgcctctccggaaagtgataacagaattca
ttgaagtggagaatttttaagaaggtacaaaaagagaaagaaatg (Seq ID No: 240)

Homo sapiens replication factor C (activator 1) 1, 145kDa (RFC1):
5 tgcctttcttgacttcgcgggagaagttgttggcggaatggatcctgagcctcgataacagatt
cctcaaccggcccaccgcccagccagccagcgcttcctcctggggctgcatg
(Seq ID No: 241)

Homo sapiens ring finger protein 4 (RNF4):
gcatctttctcgaggagctctcctgggcggtgaagaaggagcttcttctccggagtgcgcccggcg
gtggcgctgcgacctaactagctccaggttaggcccagctttgcggaagcagcggaacttgaa
10 aatactggaaatctgtccggatccaaattatgttgaagccagatgagtaaccagagggcatgaaa
ggttgagaacatttgacttccctgcaaacccttggtatagatcacttctttctgtaggaaaggaa
aggcaccaaagagcacaatg (Seq ID No: 242)

Homo sapiens ribophorin I (RPN1): tgctcttcccgggtcatg
(Seq ID No: 243)

15 Homo sapiens ribosomal protein S27a (RPS27A):
cgttcttctcttttcgatccgcatctgcggtggagccgccaccaaagatg (Seq ID No: 244)

Homo sapiens secreted and transmembrane 1 (SECTM1):
cttccttttagcgtgaaccgcggtgcggtgcctcccgtgaaaataataaattcacccgtcacgcttg
ttgtgaacgcgggtggttcccgaaacttgaggcttcccgtaaaccagctccttccctcatctggg
20 aggtgggtcccgcgcggttccgcgcctcctcctggccctccctctcgtgtctttcattttcct
ggggctccggggcgcggaagctgcatcccagaggagcgcgctccaggagcggaaccgggagtggt
tcaagagccagtgaacaaggaccaggggcccgaagtccaccagccatg (Seq ID No: 245)

Homo sapiens small glutamine-rich tetratricopeptide repeat
(TPR)-containing, alpha (SGTA):
25 ctttcttttgcgaggcgctcgcgccctggggccggggccggggcgccaccgcggtgcgcaagcgcaa
ccgtcggtgggtcggggatcggtcgctgagaggtatcacctcttctgggctcaagatg
(Seq ID No: 246)

Homo sapiens SH3 domain binding glutamic acid-rich protein like
(SH3BGL):
30 agttctccttccaccttccccacccttctctgccaaccgctgtttcagcccctagctggattcca
gccattgctgcagctgctccacagcccttttcaggacccaaacaaccgcagccgctgttccagga
tg (Seq ID No: 247)

Homo sapiens solute carrier family 1 (gluta-
mate/neutral amino acid transporter), member 4 (SLC1A4):
35 cgccctcctacttccccgtctgcgtccgcgttcgcggtcccgtttgcatcatccccgtctgcgtc
cggttgcgggtcccgtttgcatcatctccagccggcggtgctccaggagggtggcgcgatc
ctctccgcccgcggtccaaccgcactctgcgctctcctcgcctttctcgcacctgctcctgcg
ccaggcccggagacccccggggcggttcccagaacctgcggagcacaactggccgaccgacccat
tcattgggaaccccgctcttttgccagagcccacgtcccctgccaccttagctcggagcggcggtg
40 agcgccatg (Seq ID No: 248)

Homo sapiens small nuclear RNA activating complex, polypeptide 2,
45kDa (SNAPC2): ctgccttttctgagcggcatg (Seq ID No: 249)

Homo sapiens sorting nexin 1 (SNX1): ctatctctcga-
taaagttgttgttgcggcttccgcccgggtggaagaagatg (Seq ID No: 250)

Homo sapiens signal recognition particle 54kDa (SRP54):

ctatctctcatctttccgctcttagctgggagtgctccgcctagtcacttttcttaaggtggctcg
tcgaggcctgacttcttccccgaaatcacgtccctagacagcctcctattttaccactaactttac
tcctgcagttattcagcggtaggaaactgaaacaaaaaccagtgtgaagcaagtaaacaatctaaac
5 tgtttcaggagccgcgtagaaggaacgcggcgggtgtgccccggaagcggaagtagattctcctata
gaaaggctggactacgcggagtggtgacgttttctcattgggcggaagggttcgctggcactccgtt
ggtcttccagctggtgggagttgacgacgtggtgctgggcgttgggaccctactttatctagttcg
ggaagttgggtgtggggtcatacctgtctgtctgctcccagctttcttgggtttcttccgacggc
gtggggcctcgctaaggaattcccggcccctcagggccacggcttttagcgggtgtcttttgcgagtt
10 cttcgtaagtacatcttaaagctgtcaagatg (Seq ID No: 251)

Homo sapiens signal sequence receptor, beta (translo-
con-associated protein beta) (SSR2):

cggctcttccgatgctgacgctctcttctgtctttgtggctccggaaggcggttgggatgcca
cgatg (Seq ID No: 252)

15 Homo sapiens signal transducer and activator of transcription 6, i
nterleukin-4 induced (STAT6):

ttttcttttttggtggtggtggtggaagggggaggtgctagcagggccagccttgaactcgctgga
cagagctacagacctatggggcctggaagtgccgctgagaaagggagaagacagcagaggggttg
ccgaggcaacctccaagtcccagatcatg (Seq ID No: 253)

20 Homo sapiens suppressor of Ty 4 homolog 1 (S. cerevisiae)
(SUPT4H1): tgttcttcccatcggcgaagatg (Seq ID No: 254)

Homo sapiens transcription factor 7 (T-cell specific, HMG-box)
(TCF7):

ggtccttccccctaaaacttggcactgccgatactcccagcccgttcccttcccaagtcaggaacttg
25 caggggaccccttggcaattcttttctctcaagagcagacagccttcagtcccagccgctgccag
ggctggtgtgtctgaccagctgtggtttttccaggcctgaaggccccggagtgcaccagcggcat
g (Seq ID No: 255)

Homo sapiens TEA domain family member 4 (TEAD4):

cagtctcctccccgaggtgccggtggccccgccgccaactccctccggctccctccctcccgccgcg
30 gcgcgcattctattccagccctcattccgcgcattccagcgtcctcctcgcacactcgaggccagg
gggcccggaggccgcagctccggcgccgcgcgcgtcccgccaggtgagaggcgcccgccgcgcgc
accgcggggcgccctcacgggcccgcgcgcgcgcacgcgcgcgcagccgactcgcgcgcgcgtgct
cggctgctctttttctttccgcccgcgcgttcccgcccttgacctctgcgctccgacgcgctccgt
cccgacctctggcttccctccgcgctccggcgctgctcgctgccccctctccgcttccctcctgtc
35 cgccccgcgctccctcctcgcctccgggtgactcactcctccaggaataggatccccgtgtttt
cccgtcagtcctcattctgggaaaactcctccctccgcgcgctccgctccgctccgctgggcgcacc
ggggccggtcggcgcggggtgggcttgccccgcggccccgccttactgcgcgcgcgcgtcgggccc
cggccggagcccggctctgcgcgctgacgccctgtcgtccccgcagaacgatcgccgcggccggaa
gagttggcgctcggggcggactccttggaaactggcttagcgcacccatcccaccttcccgcaccc
40 gggaccggtcggaacgagctgattgccgctacatcaagctccggacaggaagaccgcaccagg
aagcaggtctccagccacatccaggtgctggctcgtcgaaagctcgcgagatccaggccaagcta
aaggaccaggcagctaaggacaaggccctgcagagcatg (Seq ID No: 256)

Homo sapiens G protein-coupled receptor 137B (GPR137B):

ttttctttcctccagtctcggggctgcaggctgagcgcgatgcgcggagacccccgcgggggcggc
45 ggcggccgtgagccccgatg (Seq ID No: 257)

Homo sapiens tumor protein, translationally-controlled 1 (TPT1):

cggccttttccgcccgcctccccctccccccgagcgcgcgctccggctgcaccgcgctcgcctccgag

tttcaggctcgtgctaagctagcgccgctcgtcgtctcccttcagtcgccatcatg
(Seq ID No: 258)

Homo sapiens ubiquitin A-52 residue ribosomal protein fusion product 1 (UBA52): ctatcttctttttcttcagcgaggcgccgagctgacgcaaacatg
5 (Seq ID No: 259)

Homo sapiens ubiquinol-cytochrome c reductase core protein II (UQCRC2):
cggcctccgccaccatcttgctttcctttaatccggcagtgaccgtgtgtcagaacaatcttgaat
catg (Seq ID No: 260)

10 Homo sapiens ubiquitin specific peptidase 1 (USP1):
ctgcctttcgtgtctctgcagcgtggagactggaaccggcaatttcaaaggacgccacgttcaatc
gcagcgtggcgcgggcgaggctaaaacacgggggtcctgagactgaggaaaacgcgccaagtgc
ccctcgggtggcgagtgctaaagaccctagcgggttcaggcggttcggcgagcggggcccgtgcttgt
tgcgctcctggctctccggggcgggcgagatgggcgcgctcccggtatgtagtgggtgttgggt
15 gcaagacgggagcgagcggcggtcgggggttccgctcttgggagcggtatggctactccccgcggg
gagggcgagccgaccagattttcctggggcggggacccggcggtcggggcgaggactcacctg
tcgacccacactcattcgggttgacttgccggcgctaccgcgcggacttcgctttgggccatg
accagatataattggtgattacaactttcctctataaattaactcttgacactccttgggatttga
agaaaaaatg (Seq ID No: 261)

20 Homo sapiens voltage-dependent anion channel 2 (VDAC2):
gtgtctccttcacttcgacctcagctgctggagctgcagcccgaccgcgagcgtgccaaagcggct
tcagcagctagcggagcgggtggcgggcgccccctcaggacaccaccagattccctcttcccgcg
gcctcgccatg (Seq ID No: 262)

Homo sapiens vimentin (VIM):
25 gcctcttctccgggagccagtcgcgcgccaccgcgcgcccaggccatcgccaccctccgcagcca
tg (Seq ID No: 263)

Homo sapiens very low density lipoprotein receptor (VLDLR):
ccccctccccgctgctcaccctcgtctccggcgccgcccgggtgcgggtgctccgctaccggctcct
ctccgttctgtgctctcttctgctctcggtctccccacccccctctcccttccctcctctcccttgc
30 ctccccctctctgcagcgctgcattattttctgcccgcaggctcggcttgactgctgctgcagc
ccggggaggtggctgggtgggtggggaggagactgtgcaagttgtaggggaggggtgccctcttc
ttccccgctcccttccccgcgaactccttccccctccttctccccctttccccctccccgccccac
cttcttctccttcttcggaaggactggtaacttgcgtgcggagcgaacggcgggcgggcgggcggc
ggcggcaccatccaggcgggcaccatg (Seq ID No: 264)

35 Homo sapiens wingless-type MMTV integration site family, member 10
B (WNT10B):
agtcctttgctcgccggttgctagctctctcgatcaactccctcccttccctcccttctctccc
ggcgggcgcgggcggtggtggaagcgggtgaagaggagtgcccgggccctggaagaatgcggctct
gacaaggggacagaaccagcgagctctccccacggtttaagcagcactagtgaagcccagggaac
40 ccaaccgtgcctgtctcggaacccgcacccaaaccactggaggtcctgatcgtctgccaccgga
gcctccgggcttcgacatg (Seq ID No: 265)

Homo sapiens CCHC-type zinc finger, nucleic acid binding protein (CNBP):
cagcctctaccttgcgagccgtcttccccaggcctgcgtccgagctctccgccgctgcggggcccgt
45 ccgacgcggaagatctgactgcagccatg (Seq ID No: 266)

Homo sapiens zinc finger protein 43 (ZNF43):

gggcctttgtctctggtgcagttggagctctgctctcgtcttcgttcttctgtgtcctctgctg
ctagaggtccagcctctgtggctctgtgacctgcggtattgggggatccacagctaagacgccag
gacccccggaagcctagaaatg (Seq ID No: 267)

5 Homo sapiens zinc finger protein 74 (ZNF74):

cagtccttttgtgggagtcgggtctgtccacttgccggtccctcagaccgtcggcggtctctgtcc
gcttcgggacctgtccgctggtcgtccgctccgatggctcctggccgcggaaccttaggcctgg
ccctggtctccgagcgcggttcgccgggaggagcgtgtggcggggtgtgccggggcgtagtg
gcccagcatggggctgagcctggtgtggggagtgggtatctgcggagccggcctgaacccccacctc
10 agccgggcgcggggaggggctccgtgctgtgatcgtgcagctgtgagcgcgtggccgccccgcg
gggctccgctgcaggccccctcagccccaggagcagtactcgtcttccagggcctgccctggatcct
ggaggctacacagctgccactcctcctggggagggtgccgtggaggccatg
(Seq ID No: 268)

Homo sapiens zinc finger protein 85 (ZNF85):

15 gggcctttgtctctcgtgcagcctgagctctaggtcttgttttccctgctttgtgttttctgctc
gtggacgcccagcctctgtggcctgtggcctgcaggtattgggagatccacagctaagacgccg
gacccccggaagcctagaaatg (Seq ID No: 269)

Homo sapiens zinc finger protein 91 (ZNF91):

20 gggcctttgtctctcgtgcgcgaggatttccaggctcgcacttcaactgctctgtgtcctctgct
ccaggaggcccagcctgtgtggcctgtgacctgcaggtattggagagccacagctaagatg
(Seq ID No: 270)

Homo sapiens zinc finger protein 141 (ZNF141):

25 gggcttttgcgtctggtactaccagaccgcggttaggggttcatctctctgcgttctcagttg
tgggaggccttggtgattcggccacagcctcagcctccgtcgtctgtgacctgcgggtattggat
gattggtagctaagactcccgaatacttcagaagtggggaaatg (Seq ID No: 271)

Homo sapiens zinc finger protein 205 (ZNF205):

tggtcttttctagctctgaaatagaaaatg (Seq ID No: 272)

Homo sapiens transmembrane protein 187 (TMEM187):

30 ctcccttttgcggagatttgaatttccccagcgaggcgagtgaggcgaaatacccgatggtgata
gctggccttttgcgcgaataactgaaaaaggcagaacggttccctccgctggcgccagccaatcagca
ggactcctgccttcttcggggcaaggctgcagcatctgcctcggaatcacgaaatcacggggct
tctttctgctggctcagccgggaggccagagtgttctgcagaggctgcgtattgaaggctgctct
ctgaagctccctgccccagggtcacgcccgggttccagatg (Seq ID No: 273)

Homo sapiens histone cluster 2, H2be (HIST2H2BE):

35 acttcttttcttggttaagccgcgtttgtactgtgtcttaccatg (Seq ID No: 274)

Homo sapiens solute carrier family 25 (mitochon-
drial carrier; oxoglutarate carrier), member 11 (SLC25A11):

40 ccgcctttgcgctgcgcgcctgcgccgcgcggcttccagcgggtgtcggacctgagagctggag
gggctgcgcgcgcctcgtctgttgcgcgcgcgggtgtcaccttgggcgcgagcggggccgcgcg
cgcacgggacccggagccgagggccattgagtggcgatg (Seq ID No: 275)

Homo sapiens tyrosylprotein sulfotransferase 2 (TPST2):

45 cctcccccttccccgggtggggcggtggagagccgggagtcgctgggtgcgtggggctgcctcgc
cgctctcgcacgggctctgccagcagacagccttggcacacaggcacaagggctggagcccaga
gatgagagtgcceaagggagatgtgagcctggcggtgcccgttaacctgtcgtgaagccccag
aagcggggcctcaggccaggcctaccctgcctccggcccagcatg (Seq ID No: 276)

Homo sapiens sorbin and SH3 domain containing 2 (SORBS2):

aagcctcttttatacatctcttcaggggaagagagaagcaatgggcatgtagtatacaatgatcac
agccacgcaggcctgcaagctgccttttggacaggctgttgactgccgttccaattagctgattgg
agaatgtggaatgcagagtataatgctgcatactgctatcaggcagcagcaaaggtttttgtct
5 tgggaaggcaagctttccctgcaatattatctcagcagctccctagctgcttaccctgaaaacgag
ggatccaaacggagggtgttgactctgctaacgctggcctgtgcgtggctgtggcatatgagcg
gcaggtctgaaaaagcaggtgtgtgctgggacgggactggactggaacgcaggcggacgctctcg
ggtttacctgcttccctgttaacagattgtgggctcccagggcataatgtctgcagctgaggccgag
gcggagaaggggcttccctgagcgtcccagtagcactgacagagacacttgattggacttaattcta
10 aacctctggagttcaagaccttttaaaaagggttaaataaacaatctctacatgtaaaaggccact
gactcctacttctctgtatagagcaactgttgaaactcagctgcctgtaggaaaactgaagacttt
aataacaaactctccaaggtgaaaatg (Seq ID No: 277)

Homo sapiens G protein-coupled receptor 65 (GPR65):

gtttctcttcttgacttgatgcaggcacagatttatcaagctcctcagtcaacaaacacatcacccg
15 gaagaaatatggaaggaaaggaatttttaaaaggaaataccaatctctgtgcaacaaagccttgta
tattcatgtttgcaccaatctactgtgagatttatgaagaaaaacaaattgcggacaactctctat
gtacacttacaaatgcctcagttgatgcttgtgggctgtttgtcagcgttctgtgataatgaacac
atggacttctgtttattaaattcagttgacccttttagccaattgccaggagcctggatttttact
tccaactgctgatactgtgtgtaaaaattgatctacatccacccttttaaagcattgatgaattaat
20 tagaacttttagacaacaaagaaaaattgaaaaagaattctcagtaaaagcgaattcgatgttcaaa
acaaactacaaagagacaagacttctctgtttacttttctaagaactaatataattgctaccttaaa
aaggaaaaaatg (Seq ID No: 278)

Homo sapiens nipsnap homolog 1 (C. elegans) (NIPSNAP1):

gggccttctctgcaacctttgcggtccaacatg (Seq ID No: 279)

25 Homo sapiens inhibitor of kappa light polypeptide gene enhancer in
B-cells, kinase complex-associated protein (IKBKAP):

gcttctttgcagcgttccagcgttttccctggaggggcgctccatccttgaggcctagtgcctg
cggagagagagcgggagccgcggacagagacgcgtgcgcaattcggagccgactctgggtgcggac
tgtgggagctgactctgggtagccggctgcgcgtggctggggaggcgaggccggacgcacctctgt
30 ttgggggtcctcagagattaatgattcatcaagggatagttgtacttgtctcgtgggaatcacttc
atcatg (Seq ID No: 280)

Homo sapiens COP9 constitutive photomorphogenic homolog subunit 3
(Arabidopsis) (COPS3): ctgccttcgccgctcgggcccgggggaaaacatg
(Seq ID No: 281)

35 Homo sapiens pirin (iron-binding nuclear protein) (PIR):

ccgcctcctctaggccgcccggccgcaagcgtgagtcacggtgaggctactggaccacactctc
ttaacctgccctccctgactcgtcccgcggtcttctcgctcaccgccgctaagggtccag
gtgccgctaccgcagcgtgagtacctggggctcctgcaggggtccactagccctccatcctctaca
gctcagcatcagaacactctcttttttagactccgatatg (Seq ID No: 282)

40 Homo sapiens THO complex 5 (THOC5):

ccttccttacttccggttctctatgggtgcgcgggcaagctttgctccgcctccggcagtggttac
tcccgggtgccaggttcttggagctgtgaggaggaacaacatg (Seq ID No: 283)

Homo sapiens RuvB-like 1 (E. coli) (RUVBL1):

gggcctttgcaaaattgccctagtaacggccgcatggtaactcaggcgccgggcgactgtccttag
45 ctgctgggttttccacgctggttttagctcccggcgtctgcaaaatg (Seq ID No: 284)

Homo sapiens Kruppel-like factor 7 (ubiquitous) (KLF7):

tttccttttttagttgactgaaacaaaacaaaagggccactggatgtctgccttcttggggg
gtgagccagacagactgacaaacaaacagcccaactgtgttcgggggagggtttcgctcccgtt
ttgcccggcagcagcagcatg (Seq ID No: 285)

5 Homo sapiens USO1 vesicle docking protein homolog (yeast) (USO1):
gctccctttttgccttcaaccttcgagccgccacgtaatgccacgtccccgcgcagcgcacatcttg
gccgctgctggcggtgtttccgggcttagagggctggagtggccgccgagttggaggcggtggtg
gcagcagtaggagtggttagagtgcgggattgggggccaggccctgcggagggcgggggaagttgt
cttctttttttccggagggcggttaaacctggtggctgaacggcaagatg

10 (Seq ID No: 286)

Homo sapiens unc-5 homolog C (C. elegans) (UNC5C):

cccccttttgccctgcctttggagaaagtggagtgtggcgcttggttgtcgttatttcttcgga
ctgcttcgcggtgcacggattcagcttctgccagtggggctttcagctgtttgcgcgtctctctg
tccccctccccctcccccggcacacctctgtctacgatg (Seq ID No: 287)

15 Homo sapiens RNA terminal phosphate cyclase domain 1 (RTCD1):
gcttcttccgctttctcgtcaggctcctgcgccccaggcatgaaccaaggtttctgaactactggg
cgggagccaacgtctcttcttctcccgctctggcgaggctttgtcgtcgcgggctgggccccag
ggtgtcccccatg (Seq ID No: 288)

20 Homo sapiens eukaryotic translation initiation factor 3, subunit A
(EIF3A):

ggctccttccctttccgtctcttgccggctgggcgcgggcgactgctggcgaggcgcggtgggacctt
acgctggttcccccttcgtctcctctcccgccccgggccactagagagttcgtgacgccgggtgag
ctgagcctgccgccaagatg (Seq ID No: 289)

25 Homo sapiens eukaryotic translation initiation factor 3, subunit D
(EIF3D):

gtttcctcttttctggtttctcaagagtgtgctgctgctaacgcggtccccggcacgcaccatctgt
tgccatccccggccggccgaggccattgcagattttggaagatg (Seq ID No: 290)

Homo sapiens eukaryotic translation initiation factor 3, subunit F
(EIF3F): ccgcctccttcttctcgcacaagatg (Seq ID No: 291)

30 Homo sapiens eukaryotic translation initiation factor 3, subunit G
(EIF3G): cgctctctggccgggcttgggctgctggagaataactttttgcatg
(Seq ID No: 292)

Homo sapiens eukaryotic translation initiation factor 3, subunit H
(EIF3H): gtttctcttcttctctgtctgcttggaagatg (Seq ID No: 293)

35 Homo sapiens eukaryotic translation initiation factor 3, subunit I
(EIF3I): aaaccttttccggtcttactcacgttgccgcttctcgcgtcacagccgggatg
(Seq ID No: 294)

Homo sapiens eukaryotic translation initiation factor 3, subunit J
(EIF3J): ctccctctcacacagctcacaccggctcgagatg (Seq ID No: 295)

40 Homo sapiens poly(A) binding protein, cytoplasmic 4 (induc-
ible form) (PABPC4):

ccgcctctctccgccccgggtcgctgccgcctccgcgctttcgggcttcgcagcctgaggaaaaa
aagagaaaaagataaaaaaatctgaaaacgcttcaaaatctgaaaaaaaaaaggaaaagaaaa

- aacgaatcctcggagaacccgcggggaagtcactttcgtacgcttccggcctgccccgcgcccgc
gcccgcagcgcttggcgctccgtcgggtctccgtccgtcgggtccgggggtgagccgcccgcggccgc
ccgtgccctcccccgctcgggccccgagccccgcgccccgcgctgccccggcgcaccacgtgtc
cgtgctgcccttcgcccgcggccggggtcgcggagtcggcgcccacaaagatttggtttccctc
5 tgccccggcggttgtaatcttaaaccgcccggagcccgagggcctatatattatagagaaacgcgtgtc
cccgaggccgcggtgggcagcgtccgggtcgctcttaaaggatttttacccttcggaaggggattc
cccgtttaatttttttctactttgattttttgaaatttgagcttcgcaccaggaccgcggagaa
gtgcaaagtcgcggggaggggccgtattgtgcggagagccttttgtctgcgggtgctgcggccgtggg
agccggcccccgctcccggtttccgtcccgctctccaagcccgcgactccagctcgtcctcgccgc
10 gccggtgccacctgtgagccgcggcggggccccgggctccgaaggcgcccctttgtcctgcggcgg
gcccgataagaagtcctcctggcggggctcgggggtggtggggggcggggagatg
(Seq ID No: 296)
- Homo sapiens receptor-interacting serine-threonine kinase 2
(RIPK2):
15 agctctttcgcggcgctacggcggttgccaccagtctctagaaaagaagtcagctctggttcggaga
agcagcggctggcggtggggccatccggggaatgggcgcccctcgtgacctagtgttgcggggcaaaaa
gggtcttgccggcctcgctcgtgcaggggctatctgggcgctgagcgcggtgggagccttg
gagccgcgcgagcagggggcacaccgggaaccggcctgagcgcccgggaccatg
(Seq ID No: 297)
- 20 Homo sapiens neuropilin 1 (NRP1):
ctttcttttctccaagacgggctgaggattgtacagctctaggcggagttggggctcttcggatcg
cttagattctcctctttgctgcatttccccccacgtcctcgttctcccgcgctcgcctgcggaccc
ggagaagggagaatg (Seq ID No: 298)
- Homo sapiens guanine monphosphate synthetase (GMPS):
25 tggctttctctcccgcggcgctggggccccgcgctccgctgctgttgctccattcggcgcttttctg
gcggctggctcctctccgctgcgggctgctcctcgaccaggcctccttctcaacctcagccgcgg
cgccgacccttcgggcaccctcccgcggcgctcgtactgtcgccgtcaccgcgcgggctccggcc
ctggccccgatg (Seq ID No: 299)
- 30 Homo sapiens far upstream element (FUSE) binding protein 1
(FUBP1):
ttttcttttctttcttagctgtagctgagaggaagtctctgaacaggcggcagcggctcttatagt
gcaaccatg (Seq ID No: 300)
- Homo sapiens eukaryotic translation initiation factor 2B, subunit
5 epsilon, 82kDa (EIF2B5):
35 gatcctttttgtcccctactgcgtgcgggtggcagcttccttgcggaagtggtagccgtgagagaag
aagatg (Seq ID No: 301)
- Homo sapiens eukaryotic translation initiation factor 2, subunit 2
beta, 38kDa (EIF2S2):
40 gtttcctttcgcgtgatgcaagagcctagtgcgggtgggagaggtatcggcaggggcagcgtgc
cgccggggcctggggctgacctctgacttcccgtccgtgccgagcccactcgagccgcagccat
g (Seq ID No: 302)
- Homo sapiens adaptor-related protein complex 1, sigma 2 subunit
(AP1S2):
45 cctcccctctccgcctaagcctgccctatgccagccgggtgtcctccccacagcaccacggcttct
cttctcagcacggcgacaggggttccccttcgcgcgcggccgcggccgccaagctccgcgcg
cgccgcgggcccgcggccgcatg (Seq ID No: 303)

Homo sapiens suppression of tumorigenicity 13 (colon carcinoma) (Hsp70 interacting protein) (ST13):
cgcccccttctgcgcggtcacgccgagccagcgctgggcctggaaccgggcccgtagccccccag
tttcgcccaccacctccctaccatg (Seq ID No: 304)

- 5 Homo sapiens solute carrier family 7 (cationic amino acid transporter, y⁺ system), member 7 (SLC7A7):
ctcccttttcttaaagtcttggggtgagagagaagagaggctagggtggggcatggaggacacagag
agagagagtgtgtgtattccttccccgctactgtcctgtcctcagctaacttgctctgggacagc
tccccagggtacagatactgcactcagctgactgtcctttcttctgggcccctgggtcccagagc
10 agagctgacaaaggagattcctgagagagcaccttcttatcacagaaagtgtgagccaagagctc
ctagctgcccccttttgcagatgtgaagggccagtgaaccttggaccagatgggttgcttaatactc
ctttccccctccctcactccttcttttgcgggtgcctcacctcctccacccttcttgcttaaatc
cataggcatttgtctggccttcccttttactgtctgggtggaaggaggagcatcagaccacagatc
ctggaaggcacttctctccctgactgtgtcactgccgtgagaacctgcttatatccaggacc
15 aaggaggcaatgccaggaagctggtgaagggtttcctctcctccaccatg
(Seq ID No: 305)

Homo sapiens paired box 2 (PAX2):
ctcccttttctcctcaagtcctgaagttgagtttgagaggcgacacggcgggcgccgcgctgc
tccgctcctctgcctcccatg (Seq ID No: 306)

- 20 Homo sapiens 5-aminoimidazole-4-carboxamide ribonucleotide formyltransferase/IMP cyclohydrolase (ATIC):
agccctcctacctgcgcacgtggtgccgcccgtgtgtcctcccgtcgcctgaacctcagtgccgtg
cagccatg (Seq ID No: 307)

- Homo sapiens ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1):
25 ccttcttttgcggctcggccatttgtcccagtcagtcggaggctgcggctgcagaagtaccgcct
gcggagtaactgcaaagatg (Seq ID No: 308)

- Homo sapiens cyclin G1 (CCNG1):
cggcccccttcggctccgagctgacctgatcagggccgagttgtctcggcgccgctgccgaggcct
30 ccaccaggacagtccccctccccgggcctctctcctcttgcctacgagtcacctctcctcgtagg
cctctcggatctgatatcgtggggtgaggtgagcaggccccggggagggtggttaccgctgaggagc
tgagtcctctgtcaagatg (Seq ID No: 309)

- Homo sapiens cadherin 16, KSP-cadherin (CDH16):
agctctcttcttgccttggcagctggaccaaggagccagctcttgggcgctggagggcctgtcctga
35 ccatg (Seq ID No: 310)

- Homo sapiens cyclin-dependent kinase inhibitor 1B (p27, Kip1) (CDKN1B):
ttttcttcttctcgtcagcctcccttccaccgccatattgggccactaaaaaagggggctcgtcttt
tcgggggtgtttttctccccctccccgttgcctcagcgtctgcgactccgacgccggca
40 aggtttggagagcggtgggttcgcgggaccgcgggcttgacccgcccagactcggacgggctt
tgccaccctctccgcttgcttggtccccctctcctctccgccctcccgctcgcagctccatttgatc
agcggagactcggcgggccgggcccgggcttccccgcagccccctgcgcgctcctagagctcggggccg
tggtcgtcggggctctgtgtcttttggctccgagggcagtcgctgggcttccgagaggggttcggg
ctgcgtaggggcgctttgttttgggttttgggttttggagagtgagagagggcggtcgtgca
45 gaccgggagaaagatg (Seq ID No: 311)

Homo sapiens chimerin (chimaerin) 2 (CHN2):

tctcctcttcttcttcttgtgtgtgctgcgagcggagttggggcggaggggagaagggggaggtcgctc
tgtctgtccgtctcccgccgctctgcccgtctactcgaagtgcggcgggagagggcgggagccca
ggagaggggtgcgggagctggcggggcggtcggagctgccaggacgccctggtcccagccgcgcac
5 aggggagcgtggacggcagaggggctcggcgggagccgagatccgccgtcccggctgccctcgg
cctccctctgtctccacctacccccctgacacccatagaaaagcgtgcaaaggcgcggagcgggacg
gaaaccacaaataaataagcggcggcggcagcgcgtcatctggtggagcaggaagtgcaggcagagt
ccggaggctggtgctttctgcgcgtccccaggactttgccatgggctgggggcccgcggaggtcgcg
agcggccgggagggcagcggcggcggcgtccgcaccggggctgagcagcagcgcgcgagggg
10 cgcgcgagatg (Seq ID No: 312)

Homo sapiens citrate synthase (CS):

gggcctccttgaggaccccggtggtggcgcgcgcgggttcgtctactctttccttcagccgct
cctttcaaccttgtcaaccgctcggcgcggcctctggtgcagcggcggcgtcctgttcttgcg
cagctctctcccttcttacctccccaccagatcccgagatcgcccgcctatggctttacttactg
15 cggcgcggcgtccttggaaccaaggcaccagtggtgcaagtactagctgagcatttgggagatgc
ttgtcttacttggtggttcttctcctgctgctggggaaaaggaaatgcattcttcttcttgc
agccggcatgccagtgttctccacgaatttgaaagacatattggctgacctgatacctaagga
gcaggccagaattaagactttcaggcagcaacatggcaagacggtggtgggcaaatacactgtgga
catg (Seq ID No: 313)

20 Homo sapiens cathepsin S (CTSS):

atttcttttcaagtcaattgaactgaaatctccttgttgccttgaaatcttagaagagagcccact
aattcaaggactcttactgtgggagcaactgctggttctatcacaatg (Seq ID No: 314)

Homo sapiens deoxynucleotidyltransferase, terminal (DNTT):

cagtctccctcccttctggagacaccaccagatgggccagccagaggcagcagcagcctcttccca
25 tg (Seq ID No: 315)

Homo sapiens dual specificity phosphatase 3 (DUSP3):

cgctctccgctcgcttgcctcctgcccggcggtgcagggccccgcgcgcgcatg
(Seq ID No: 316)

30 Homo sapiens coagulation factor II (thrombin) receptor-like 2
(F2RL2):

catcctttccctgcggaggaccagggcaagtttctgcctgcacggcacaggagagcaaaacttcta
cagacagaccaaggcttccatttgctgctgacacatggaactgaggtgaaattgtgctccatgatt
ttacagatttcataacgtttaagagacgggactcaggtcatcaaaatg (Seq ID No: 317)

35 Homo sapiens Fc fragment of IgG, receptor, transporter, alpha
(FCGRT): cgctctctcagcatg (Seq ID No: 318)

Homo sapiens guanylate binding protein 2, interferon-inducible
(GBP2):

ttacctcttttcttgtctctcgtcaggtctctgacattgacagagcctggacgttggaggaagcc
ccaggacgttggaggggtaaagtaaaagtccacagttaccgtgagagaaaaagagggagaaagca
40 gtgcagccaaactcggaagaaaagagaggagaaaaggactcgactttcacattggaacaaccttc
tttcagtgctaaaggatctctgatctggtggaacaacacctggacatg (Seq ID No: 319)

Homo sapiens G protein pathway suppressor 1 (GPS1):

cgctctttctcccttcagcagccagccagctctgtgtcagggtcggggggtgcagaaagtcaggac
agaatg (Seq ID No: 320)

Homo sapiens general transcription factor IIF, polypeptide 2, 30kDa (GTF2F2):

gttcctcttttctcctcggttcccagtggttctggcaggttaaggaacgccggctcttcgcctctcagcg
cggcttgctcttctgttccggacgcccgctcctcagccctgcggctcctggggctcgtgctgcatcc
5 cgcacgcctccaccggctgcagacccatg (Seq ID No: 321)

Homo sapiens glycogenin 1 (GYG1):

cgctccctcccgggtgccggcttctctgagtcaccaacctgaggctgccccggccgcctgcgcaccc
ggcagcaccatg (Seq ID No: 322)

Homo sapiens heat shock 70kDa protein 9 (mortalin) (HSPA9):

10 agctctttgccgtcggagcgcttggttgctgcctcgtactcctccatttatccgccatg
(Seq ID No: 323)

Homo sapiens iron-responsive element binding protein 2 (IREB2):

cttccttcttttctccttgcagtcgcctgtcttctccccgtcttccctgccccggcctcccc
ttcttcccccgctggccccctccccggaggataatatggtctccggcgatg
15 (Seq ID No: 324)

Homo sapiens origin recognition complex, subunit 1 (ORC1):

ccaccttcttttcatcttagtgagacacacgctttggctcctggctttcggcccgtagttgtagaa
ggagccctgctggtgcaggttagaggtgccgcatcccccgagctctcgaagtggaggcggttagga
aacggagggttgctggctagccggaggaagctttggagccggaagccatg
20 (Seq ID No: 325)

Homo sapiens RAB1A, member RAS oncogene family (RAB1A):

cattcctttctttcgattaccctggtgcgagagtcagggcgggcggtgcggcagcaagggcggc
ggtggcgggcgggcagctgcagtgcacatg (Seq ID No: 326)

Homo sapiens cytohesin 2 (CYTH2):

25 gagtcttttcagcgctgaggactggcgctgaggaggcggtggctcccggggcggtttgagcggg
ctcaccagagcccggggccaacgcggatccaggcccgactggcgggaccgccccggattccccgc
ggccttcttagccgcatg (Seq ID No: 327)

Homo sapiens COP9 constitutive photomorphogenic homolog subunit 2 (Arabidopsis) (COPS2): atttctcctccccctcccgccaagatg

30 (Seq ID No: 328)

Homo sapiens solute carrier family 9 (so-

dium/hydrogen exchanger), member 3 regulator 1 (SLC9A3R1):

ggtcctctctcggctcctcgcggtcgcggcgccgacggttctgggacacctgcttgcttggcc
cgtccggcggtcagggttctctgctgcgtcctccggttcgctggacgggaagaagggtgggccc
35 tcccgctcccgctccccatcggaacccaagtcgcgctcgtgacctcgcagggcgagatg
(Seq ID No: 329)

Homo sapiens peptidase (mitochondrial processing) beta (PMPCB):

ctaccttcttctagcagaaatg (Seq ID No: 330)

Homo sapiens RAB3D, member RAS oncogene family (RAB3D):

40 cggcccttctcctcgccttctggcgagcccgcgctggatccgggtggctgcaggctgctggcttc
tgcggtgcggggtcggttcgcggccaggcccaagccgcagcgagttcacaggcggaacccctgc
aggcggcgccccctacgcgaggtcacccctgggaaggagcgcagcccacccggccccctccgcatcc
gagcaggacgcccgtctcctctcctgaggatttcagggtctcctgtcccaggaggcttgtgcca
gatg (Seq ID No: 331)

Homo sapiens ATP-binding cassette, sub-family B (MDR/TAP):
tcttctctcggttcctctttcctcgctcaagatg (Seq ID No: 332)

5 Homo sapiens N-acylsphingosine amidohydrolase (acid ceramidase) 1
(ASAHL): ggctcttctttgcctctgctggagtcgggggagtggcgttggtgctagagcgatg
(Seq ID No: 333)

Homo sapiens cytochrome c oxidase subunit VIc (COX6C):
ttttccttttagtcaggaaggacgttggtgttgaggttagcatcacgtatcaaggacagtaactacca
tg (Seq ID No: 334)

10 Homo sapiens COX15 homolog, cytochrome c oxidase assembly protein
(yeast) (COX15):
gcttctcttttcttggcggaggagggagaccacagagccctgggttggtggaagaggtggctgttc
cctgtcatcagtatg (Seq ID No: 335)

15 Homo sapiens c-src tyrosine kinase (CSK):
cccccttccccgcctttcttccctccgcgacccgggcccgtgcgctccctgcctctgcctg
gcggtccctcctccctctccttgacccatacctctttgtaccgcacccctggggacccctgcg
ccctccctccctccctgacgcgatggaccgtcccgaggccgctgatgccgcccgcggcgaggtg
gcccggaccgcagtgcaccaagagagctctaattggtaccaagtgcaggttggtttactgtgact
cggggacgccagagctcctgagaagatg (Seq ID No: 336)

20 Homo sapiens versican (VCAN):
gagcctttctggggaagaactccaggcgtgcggacgcaacagccgagaacattaggtgttgaggac
aggagctgggaccaagatcttcggccagccccgcacccctcccgcatcttcagcaccgtcccgcac
cctccgcacccctcccgggccaccacgcttctctatgtgaccgcctgggcaacgccgaaccagtc
cgcgacgcgtgcagtgaattttccccccaaactgcaataagccgccttccaaggccaagatg
(Seq ID No: 337)

25 Homo sapiens dystroglycan 1 (dystrophin-associated glycoprotein 1)
(DAG1):
gcgctcttaggcttggcggtggcgggcgggcagcttcgcgccgaatccccggggagcggcggtg
gcggcgtcctggggccaggaggagcgaacacctgcgcggtcctcccgccggcgtgggctctgtg
tgctccgggatggagcaggtgtgcagaggggtgagaaccagctctgggaccaagtcaactgtctcc
30 ttacttagcaagactatcgacttgagcaaacttggaacctgggatg (Seq ID No: 338)

Homo sapiens DEAD (Asp-Glu-Ala-Asp) box helicase 5 (DDX5):
ccccctcttttggttacagacgtgagggctctttggagacgtaaacatctccgagtggcgaggggtg
ggcggggctgggcttgggaaagggcggggtggcttgcttgaggtgtggaaagaccagaagaaggtg
aggtcaagagagtgacgaatgaggcattccaatggtgggtgggcccctgacctgagagagtggcgcg
35 gggaggggtgaaagcgcggcgatcctggaacgccagcgggcgttgcgccctatgcgcgaggggcgg
ggcgattaggtcatagagcggctcccagcgttccctgcggcgtaggaggcggtccagactataaaa
gcggctgcgggaaagcggccggcacctcattcatttctaccggtctctagtagtgacgcttcggct
ggtgtcatcggtgtccttccctccgctgccgcccccgcaaggcttcgccgtcatcgaggccatttcc
agcgacttgtcgacgcttttctatatacttcggttcccccgccaaccgcaaccattgacgccatg
40 (Seq ID No: 339)

Homo sapiens desmoplakin (DSP):
gctcctctgcgcccttgccgcctccgagccacagctttcctcccgtcctgccccggcccgctcg
ccgtctccgcgctcgcagcggcctcgggagggcccaggtagcgagcagcgacctcgcgagccttcc
gcactcccggccggttccccggcgtccgcctatccttgccccctccgctttctccgcgcgggc
45 cgctcgcttatgcctcggcgctgagccgctctcccgaattgcccgcgacatg
(Seq ID No: 340)

Homo sapiens glutamyl-prolyl-tRNA synthetase (EPRS):

cttccttttcgcggggtcctccgtagttctggcagcagccaggcgtactgacaggtggaccagcgga
ctggtggagatg (Seq ID No: 341)

Homo sapiens acyl-CoA synthetase long-chain family member 4

5 (ACSL4):

gctcctcctcgtcccagcgctagcgggcacgcggttcctttttgcgagctttccgagtgccaggcg
ccggccggctgcgaagacgcggtgggcccctccgattgaaatcacagaagatattcgtgttct
tcttaagagaaaaagaggacatttttagctttctcagttgaaggcgtactttattgtcggcttccaa
agattactaacttttatctgtatcactaagattgaactgccttggtgtactgtattcttactgc
10 tgcttctattattgccttcttcagcacaataaggctttcaaaagccaaagaataacaagaataag
caccatttttagaagcctttccactatg (Seq ID No: 342)

Homo sapiens fibroblast activation protein, alpha (FAP):

tggctccttttcaacgggttttcacagatccagtgacccacgctctgaagacagaattagctaacttt
caaaaacatctggaaaaatg (Seq ID No: 343)

15 Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acet
ylgalactosaminyltransferase 3 (GalNAc-T3) (GALNT3):

ctgcctctccaggcaacgcgggaggcccgaggcgggaaggcaggaggcggcgggaggaggagctct
actgagccgcaactgtggcgacagcaaccggagtcgcagccgcccacctgcacctggcgcttag
cccacgtccagcgctgcccggcgccgcttcccggccaccctgcctgcccacccggccaggtacta
20 ccattaaagataccttcttctcagcaaatctatgataaaaaatataagtaacagaagaagaataa
ctgttattttgtcaagtgacaagcttttaatgtcagaatg (Seq ID No: 344)

Homo sapiens glypican 3 (GPC3):

acgtctcttgctcctcagggccactgccaggcttgccgagtcctgggactgctctcgctccggctg
ccactctcccgcgctctccttagctccctgcgaagcaggatg (Seq ID No: 345)

25 Homo sapiens interleukin enhancer binding factor 2, 45kDa (ILF2):

acgcctcttcagttgtctgctactcagaggaagggcggttggtgcggcctccattgttcgtgttt
taaggcgccatg (Seq ID No: 346)

Homo sapiens nucleosome assembly protein 1-like 1 (NAP1L1):

gggtcttttttagcgccatctgctcgcggcgccgctcctgctcctcccgtgctgctgccgctgc
30 cgccctgagtcactgctgcgagctccggccgctggctccccatactagtcgccgatatttggga
gttcttacaacatg (Seq ID No: 347)

Homo sapiens asparaginyl-tRNA synthetase (NARS):

cgctctctgatgcaacgccggaatcgcggaaccgcccgggtgcacgttgagtcataagacggcgctc
ggtgttgagctctgtgtccttgagggtgaccagggccactgcaggcatg (Seq ID No: 348)

35 Homo sapiens NADH dehydrogenase

(ubiquinone) 1 alpha subcomplex, 10, 42kDa (NDUFA10):

cgtccccttgggctccttgatcctgagctgaccgggtagccatg (Seq ID No: 349)

Homo sapiens NADH dehydrogenase (ubiquinone) Fe-S protein 2, 49kDa
(NADH-coenzyme Q reductase) (NDUFS2):

40 ttctccttcccgcagctctgcagccggagtaagatg (Seq ID No: 350)

Homo sapiens NADH dehydrogenase (ubiquinone) Fe-S protein 5, 15kDa
(NADH-coenzyme Q reductase) (NDUFS5):

catccttttacggcaggcgctccgctcgctagctagtcgttctgaagcggcgccagagaagagtca
agggcacgagcatcgggtagccatg (Seq ID No: 351)

Homo sapiens phosphoenolpyruvate carboxykinase 2 (mitochondrial) (PCK2):

ccctcctttttaagcgctcccgccagcctctgctgtggctcgcttcgccgcgctccctccttccc
cgcttccatacctccccggctccgctcggttcttgccaccccgagccctgccaggtgccat

5 g (Seq ID No: 352)

Homo sapiens serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6):

ctcccttcgcgctccggacgggacggtagctcgagacccgggactccgcccgcctccccgcgag
tatttgaggtccggggcggtccggcgctctgcccgcggttctgctcgctcgctccccgcgtctgg

10 agtctgccatcatg (Seq ID No: 353)

Homo sapiens Rab geranylgeranyltransferase, alpha subunit (RABGGTA):

ttctctcctcagacttcaagggctaccactggacccttccccctgtcttgaaccctgagccggcacc
atg (Seq ID No: 354)

15 Homo sapiens Rab geranylgeranyltransferase, beta subunit (RABGGTB): ctctctcctttccctgtagacatg (Seq ID No: 355)

Homo sapiens small nuclear ribonucleoprotein polypeptide A (SNRPA):

agttctctccgcacgcgggctggagaagcgggtcctacgcacgctttgttgctcgcgctttgcctcc
gtccttgcccctactccgccttacctgacttccttttcggaggaagatccttgagcagccgacgt

20 tgggacaaaggatttgagaaacccagggttaaagtcacgtttttcctcctttaagacttacctca
acattcactccatg (Seq ID No: 356)

Homo sapiens sterol regulatory element binding transcription factor 2 (SREBF2):

cgccctttctgtgcgggcgccggggcgcaacgcaaacatggcgggcggtggcaccgcgtcggtgaggg
ggtgcccggggcggggttgctcggtgtcatggcggtggcgacggcaccgccccgcgtctccctga

25 gcgggacggcagggggggcttctgctgagccggggcgatg (Seq ID No: 357)

Homo sapiens translin (TSN):

ctgccctttggacgcgcgcctcggttccgaacgcagcggacggcgccctcaggcagcgcgggcgga
gcccgtcctccggcgcgccgcgagcctcgaggaccctagcgacggctcgtggcgtaagaccggggg
gacgcggcggttagcgggcgccgttgcgattgattgcgtggttgctgcggcgctccacttccttgg

30 ccgcccttgctacactggctgattgttgctgcagccggcgccatg (Seq ID No: 358)

Homo sapiens Fanconi anemia, complementation group G (FANCG):

ccaccctttctcgaggctgtggcctccgcgagagccgagcgggcccgcaccgcccggccgtgcgactg
ccccagtcagacacgacccccggcttctagcccgcctaagcctgtttgggggttgctgactcgtttcc
tccccgagtttcccgcggggaactaactcttcaagaggaccaaccgcagcccagagcttcgcagacc

35 cggccaaccagaggcgaggttgagagcccggcgggggcggggagagagcgtcccatctgtcctgg
aaagcctggggcggttgattgggaccccgagagaagcaggggagctcggcggggtgcagaagtgcc
caggccccctccccgctgggggtgggagcttgggcaggccagcttcacccttcctaagtcgcttct

40 ggtctccgggcccagcctcggccaccatg (Seq ID No: 359)

Homo sapiens DEAD (Asp-Glu-Ala-Asp) box polypeptide 39B (DDX39B):

ttccctccttcgtcgctgttgctgccgccatacgcgctctccctgttttagctcttctgttagaaat
agtatctttgttttcccttctgtgttcctcaatcccctactcttcaccccttgttttcacctatgtt
gcgagaacccatccagatcccccttcccttcttccctgcccggcccagttatg

45 (Seq ID No: 360)

Homo sapiens RAB11A, member RAS oncogene family (RAB11A):
ccgccctttcgtcctcgcggccgcgaatg (Seq ID No: 361)

Homo sapiens SPARC-like 1 (hevin) (SPARCL1):
5 agctctttcccttttgggttgcaagcactgcctgtaaagccctcgcagtgagaggccagcctgctag
ggaaatccaggaatctgcaacaaaaacgatgacagtctgaaatactctctggtgccaacctccaaa
ttctcgtctgtcacttcagacccccactagttgacagagcagcagaatttcaactccagtagactt
gaatatgcctctgggcaaagaagcagagctaacgaggaaaggatttaaagagtttttcttggggtg
tttgtcaaacttttattccctgtctgtgtgcagaggggattcaacttcaatttttctgcagtggct
ctgggtccagcccccttacttaaagatctggaaagcatg (Seq ID No: 362)

10 Homo sapiens cyclin B2 (CCNB2):
ctcccttttcagtccgcgtccctccctgggcccgggctggcactcttgccctccccgtccctcatg
(Seq ID No: 363)

Homo sapiens cytochrome c oxidase subunit VIIa polypeptide 2 like
(COX7A2L): ggtccttctctggggcggtcgcgttggcagcggatgcgggaagccggactctg-
15 ggcgtcatg (Seq ID No: 364)

Homo sapiens lysophosphatidic acid receptor 2 (LPAR2):
cgccctctcagcaaccgcacagggcgacccggacgtctaccgctcccgcgcagtcgccgggc
catgggcctcgagcccgcgccgaacccccgcgagcccgcttgtctgcggcgtgactggaggccca
gatg (Seq ID No: 365)

20 Homo sapiens adaptor-related protein complex 4, mu 1 subunit
(AP4M1):
cgttcttttgttccggggccgcagggcggggagggcccgactttcgccgtcttcttgtctactctc
cagaacggccatg (Seq ID No: 366)

Homo sapiens budding uninhibited by benzimidazoles 3 homolog
(yeast) (BUB3):
25 cttcctctccgcctccttcgcctagcctgcgagtgttctgaggggaagcaaggaggcgggcgggcc
gcagcgagtggcgagttagtgaaacgttgcttctgaggggagcccaagatg
(Seq ID No: 367)

Homo sapiens DEAD (Asp-Glu-Ala-Asp) box helicase 21 (DDX21):
30 ctacctcttctctccacgcggttgagaagaccggtcggcctgggcaacctgcgctgaagatg
(Seq ID No: 368)

Homo sapiens solute carrier family 33 (ace-
tyl-CoA transporter), member 1 (SLC33A1):
tgctctctgccgattgatagcagcgagagctggaggtgttggtcgggagaccagccgttcgatac
35 ccgccgcaggtaggagctgggtttccatcctggcaccacggcacacacctccagcctcgagcccggc
gctgctgcccgggggtctccttcaggctctttgacgcggttccagggggcacctatccaggcatcc
tctgggcctctagccagaggactggctcccggcttcagcactccgggctgcagtaagaagtgcctt
tatcgtcttgagccctgccaccatcccgtgaaccaccgaaacctgggtccagcgcgacagccttgg
acctgggactggacggatccaaaacgtcagcctcggccccccacagacggggctctgcacgtct
40 ctgatatg (Seq ID No: 369)

Homo sapiens G protein-coupled receptor 37 like 1 (GPR37L1):
tgctcttctctgggctggctgtctcctgctcatccagccatg (Seq ID No: 370)

Homo sapiens neuronal regeneration related protein homolog (rat)
(NREP):

ctgtctttctagcatgttgccctttttcaaccacatttgtgtttcaggtgtagagaggagagagag
tgaacagggagcggggcttttgtctgttggtctccctggactgaagagagggagaatagaagccca
agactaagattctcaaatg (Seq ID No: 371)

- 5 Homo sapiens vesicle-associated membrane protein 3 (cellubrevin)
(VAMP3): gcttctctgctgaccctctctcgtcgccgctgccgcccgcagctgccaaaatg
(Seq ID No: 372)

Homo sapiens synaptosomal-associated protein, 29kDa (SNAP29):
cctccttctgtttcccagaccgagagccgcgccggcaccatg (Seq ID No: 373)

- 10 Homo sapiens lon peptidase 1, mitochondrial (LONP1):
ccccctcttctccgctagggccagctccctgaagcggctgtttcgagccacgcgccccatcgggta
ccgaggcacgcgcccggcgctcacgtgctgtttcgcgccgagcggaatgacgcgagttgtgtgagcc
gccagtatggccgggctatg (Seq ID No: 374)

- 15 Homo sapiens kinesin family member 3B (KIF3B):
ctgtctctccccatccggggcagcgggggaatggctgagccaggggttcgccgcccccgccgcccgc
gccgcccgcgcccgcgcccgcgcccgcgctttcggctcgggcctcaggaccgtagcatcctgaga
cattttgaattgacacttctcaagatttgactggatcagagttcatcatg
(Seq ID No: 375)

- 20 Homo sapiens transmembrane 9 superfamily member 2 (TM9SF2):
cttcctttatctctggcgcccttgtagtcgtctccgagactccccaccctccttccctcttgacc
ccctaggtttgattgccctttccccgaacaactatcatg (Seq ID No: 376)

Homo sapiens cytosolic iron-sulfur protein assembly 1 (CIAO1):
gagcctctgtcggccgcggaagcctggagtggcggtacgcagacgcgcggtgagaccgctgt
ctgctcagcggactctgcccgcacacctccccctgcgtcgggcccagacatg
(Seq ID No: 377)

- 25 Homo sapiens GRB2-related adaptor protein 2 (GRAP2):
caccctctttcagagtgtgtacatggaagacagcacaagtgatccatactctgaaatgcagtaac
tctgatgcttgaatttgtctcccttcttgccagaaaggattctaataactcgggtgtcaaagccaag
acataaactcaacccttctcttccaaaagcttcacgttacagcatg (Seq ID No: 378)

- 30 Homo sapiens leupaxin (LPXN):
gtacctttctcggggtgtctgcgttaactgccagacttgcccttggtttggtcagatgacacctcct
ctgggactggctagccagcgttcatg (Seq ID No: 379)

- Homo sapiens SH3-domain binding protein 5 (BTK-associated)
(SH3BP5):
35 tttcctctgctccgcccgcggccggaggtatccgcacgcggcagctgcgtctccggggtgtcggccc
cggcggtccccgaccgtgcccggtgtggcgaggcggctccagcccagcctgtggcagccgcgac
ccccggggcgctccggagccactgcgcggcgcgctgcccggctgcctgcatg
(Seq ID No: 380)

Homo sapiens phosphatidylinositol glycan anchor biosynthesis, clas
s B (PIGB): ctttcttccgccttaggaaggtggcgccaggatg (Seq ID No: 381)

- 40 Homo sapiens lipopolysaccharide-induced TNF factor (LITAF):
cggcccttttctcggggcgcccagaggccagctcagacctcccggctcgacaggcggcgcgggcg
gcggtgagtgcggcgcggggacgccccggcgcggggaccagcgggagacagcggggggcccgggtggc
gccagcacctgctgggggccccgggcactgagcccttggtggggcctcctgggatgccagggggc

gcgggtcgggtcgcgggcatcgaggcgcgcgaggcggtgggggcccggccggggcggggtccgg
cctcccagcgctggtcccggccgcgtctccggttggttcagctcctgcgtcccagagtggccga
tcgcgcgtggcggggtcgtccggccccaccgaacgagcgcccttcgcggcccgcgcgtcccc
tccccggagaggacggccctgggcttttttagaaaaaggcgcgattctctctagtgactcaggttg
5 agatttccagaaatatccccgggggttcagaaacaaaacaaaacaaaaaaccccaacga
attcccaaatgctatttgccaaacatttgacttctaggggcgcggtaccgcgttttctctccctg
ccccgcgacttcgcgcaagatccgggaaggacaccgcggccctgggagaccctggggaggtga
aaatcagagagcgaagcgggcggtggcccttaggcctgaccctccccgcgggtaaggcgggcac
cccgcgagcgcgagggtcctcttactgctgatggcaccagctctgggccagacgcgcgtcaccg
10 tccaccgcccgtgctgggtaaaatg (Seq ID No: 382)

Homo sapiens etoposide induced 2.4 mRNA (EI24):
ccacccttcggctctgggccccgcctcgtggtgcgggctggttcttcgcgctcgcccgacttccc
agcgcccccggtgcggccccggcatgccagtgcgggcgagcgccccggccctggaagcgccccg
gcggagctggcctgcggtgggctaggggcaggcgcgagccgcggcgcgagctgtggatccttc
15 atgatgagagatttggggacacttctctctcctgtgtgtagttgatagtttgggtggtgaagagatg
(Seq ID No: 383)

Homo sapiens chromosome 14 open reading frame 2 (C14orf2):
tgacctttccgagttggctgcagatttgtggtgcgttctgagccgtctgtcctgcgccaagatg
(Seq ID No: 384)

20 Homo sapiens peroxiredoxin 6 (PRDX6):
attcctccgcgcgctgggacaggctgcttcttcgccagaaccaaccggttgcttgcgtgtcccagcg
gcgccccctcatcacgcgtcgccatg (Seq ID No: 385)

Homo sapiens solute carrier family 29 (nucleo-
side transporters), member 1 (SLC29A1):
25 ctctcttccgccccggcgccacacgggtcaggcccgcgcgggctgcgctctccagctgtggcta
tggccccagccccagatgaggagggagagaactagggcccgagccctgggaatttccgtcccc
caccaagtccgatgctcactccaaagtctcagcaggccctgaggggaggagctgtcagccaggg
aaaaccgagaacaccatcaccatg (Seq ID No: 386)

Homo sapiens heterogeneous nuclear ribonucleoprotein F (HNRNPF):
30 cgaccttctgcggggccggcggtccgaggctgctggagtgcggtgagcaggccgcgggaacgct
gccgtcaccttgtctcggggctcggcgctgcttccccgcaaaaacacgtttaccgcgcgccccggc
ctccaccttgcggaagggaacccaccacttgatttctgttgcaggttgagaacaaaaacat
gcacctggagtttccccggagccctctgcgtggttgagcttcggtggaatttcggggctcttggct
gccagccgcgcttgctggttagcaacagaaaccagtcctgctgcctccgtggacatttcattacc
35 atccagaagtgtctccactgaaggcatccgtggttgtttttaagccacaaaaaagccacacccaa
gatcacctgacaccaccctgacaagtgtccatg (Seq ID No: 387)

Homo sapiens islet cell autoantigen 1, 69kDa (ICA1):
ccgcccccttccctcgccttcggctgacgctgacgtcggtatgagtgatccggaggagcgtccgac
cgcgccggggaggctcctgggggccccgggctccgaggtataataataactatcctctcatgcttt
40 tttcctgcccccttctcccaaatcatcaacaatagaagaagaagaaaacatg
(Seq ID No: 388)

Homo sapiens PWP2 periodic tryptophan protein homolog (yeast)
(PWP2): gtgtctctgtgggccccgcccgggttgagctgcggcacacgtgcgacggccgtgatg
(Seq ID No: 389)

45 Homo sapiens glutaminyl-tRNA synthetase (QARS):
gtttcttttagtttccggtgtctctgcaatg (Seq ID No: 390)

Homo sapiens stearoyl-CoA desaturase (delta-9-desaturase) (SCD):
cggcctctgtctcctccccctcccgcccttacctccacgcgggaccgcccgcgccagtcaactcct
cgcacttttggccctgcttggcagcggataaaaagggggctgaggaaataccggacacggtcaccctg
tgccagctctagccttttaaattcccggtcggggacctccacgcaccgcggctagcgccgacaacc
5 agctagcgtgcaaggcgccgcggctcagcgcgtaccggcgggcttcgaaaccgcagtcctccggcg
accccgaaactccgctccggagcctcagccccctggaaagtgatcccgcatccgagagccaagatg
(Seq ID No: 391)

Homo sapiens fragile X mental retardation, autosomal homolog 1
(FXR1): cggcctttgcggttccaacatg (Seq ID No: 392)

10 Homo sapiens musculin (MSC):
tagccttttcaaaaggcgcagcttacccgcggtgcgcgcggattctggacttgggcgccaactcgta
gtccacgctccccggggtcagcagaggggcgctcacgctctcgccaccacctcgcttttctcacc
cgcgcttcccggtcgtgggttttagtcttccctggagcgctctctggcctccgctccgccagga
gcggaaggcggagacagcgagactggccagggggaggaaagaggacgcgtgtgggcaagggggac
15 aacgggatg (Seq ID No: 393)

Homo sapiens RNA binding motif protein 8A (RBM8A):
cgacctttcccctctgcgacagtttcccgaggtacctagtgtctgagcggcacagacgagatctcg
atcgaaggcgagatg (Seq ID No: 394)

20 Homo sapiens heparan sulfate (glucosamine) 3-O-sulfotransferase 1
(HS3ST1):
ggctctctgcgccttggcagccaggagtcgcccgcacgaccgcccgggtctcagtgggtgcctgcgc
cttctccccgcccgcctgccccgggccatccagaaacttgctctaccgcgcgggtgctcggcag
tgctgcccattggccagcccaggagcctatttagggcgccggacgggctggacagaggcgcggtc
agtaattgaaggcctgaaacgcccattgtgccactgactaggaggttccctgctgcggcacttcat
25 gaccagcggcgcgcggcccagtgaagccaccgtggtgtccagcatg (Seq ID No: 395)

Homo sapiens solute carrier family 12 (potas-
sium/chloride transporters), member 6 (SLC12A6):
ctgtctctttagtaggcagggatcacagtctgaaacgacagcaaggaagaggtaggcagggaacta
actggaaggaagtttaatacacagaaagagcaaagtattatctaactataacaatg
30 (Seq ID No: 396)

Homo sapiens apelin receptor (APLNR):
cttctccagggtctggagaacccagaggcagctcctcctgagtgtctgggaaggactctgggcac
ttcagcccttcttactctctgaggctcaagccagaaattcaggctgcttgagagtggtgacaga
gccacggagctggtgtccctgggaccctctgcccgtcttctctccactccccagcatg
35 (Seq ID No: 397)

Homo sapiens calpain 1, (mu/I) large subunit (CAPN1):
cgctcttctggttgggccctgccctgagctgccaccgggaagccagcctcagggaactgcagcgac
ccccaaacacccctccccagcatg (Seq ID No: 398)

Homo sapiens cyclin C (CCNC):
40 cttcttttgcgcgtcgcgcggcggagcggagctcgagccgagctgatttgatcgaggagcgcggtt
accggacgggctgggtctatggtcgctccgcgggcccgtccgcggctggtgcttttttatcaggg
caagctgtgttccatg (Seq ID No: 399)

Homo sapiens glutamate dehydrogenase 1 (GLUD1):
45 cttctccctagtcgcggggagctctgagaaagcgcgcctgtttcgcgaccatcacgcacctccct
ccgcttgtggccatg (Seq ID No: 400)

Homo sapiens guanine nucleotide binding protein-like 1 (GNL1):

cctccttcctcgccgccccggcgccctctcggtgccactggctctcacgtgccagtagccccacccc
gcatcatcctctcgctcgctcctggagggaagtgactatatctcccccgccgcttccatcgcc
gccgcgccggttaattctgtcgggcccgccgctgacgtcacctgctagccccgcctcctctaggg
5 cccgggccccctgcggcgggggctgccccggggggcagtcagttgaggcgggagctcggcgagg
ggcgggccaggtgactggtccggggccatg (Seq ID No: 401)

Homo sapiens lysophosphatidic acid receptor 4 (LPAR4):

aggcctttttgtgtcctgtttgctaaaggcatgcgggctacagcattcaagagagggagtcgttaa
caaagggaaagagataaatgtaaataagctcacatttacagaatgagcgggtttgcagtaaaaagct
10 gcggcagcccagagtctgctacttttaggtgggctaacctttccctgtaaaaaaaaaaaaaaaa
aaaaaaaaaaaaaatggataaaaatatgcacttccaaagggcagttgccatttacatgtttatta
gctaattatctacaggcatcagcacattctctcatctagcacactctttcttggggaggaaaatat
ttcctaccggtccatagtgtcagagtgggtgaaccctgcagccagcaggcctcctgaaaaaaagt
ccatg (Seq ID No: 402)

Homo sapiens G protein-coupled receptor kinase 5 (GRK5):

gctcctctttgcagagggggaaactcttgggctgagagcaggaataatgcggtaggcaaggcgggc
tgctggctcccccggtccggcagcagcgggcagcccgagcagcggcagcagcagcggcagcac
cccaggcgctgacagccccgcccggcggtccgttgctgaccgcccactgtcaatg
15 (Seq ID No: 403)

Homo sapiens glutamic-pyruvate transaminase
(alanine aminotransferase) (GPT):

agccctttctgtccctcccagtgaggccagctgcggtgaagaggggtgctctcttgccctggagttcc
ctctgctacggctgccccctcccagccctggcccactaagccagaccagctgtcgccattccac
ttctggctcctgccacctcctgagctgccttcccgcctgggtctgggtagagtcatg
20 (Seq ID No: 404)

Homo sapiens hydroxyacyl-CoA dehydrogenase (HADH):

gggtctcctcgctgtcgccgcccgtgcccacaccatg (Seq ID No: 405)

Homo sapiens high density lipoprotein binding protein (HDLBP):

tcttctcctttaccaagatggcggttgctccctgtttcgccacagttcctaccttatgagctcggt
30 tttcttatgcttataagagtgaacagcaaaagctggcaggctgacagaggcgccctcaggacgga
ccttctggctactgaccgttttgctgtggttttccggattgtgtgtaggtgtgagatcaaccatg
(Seq ID No: 406)

Homo sapiens histidine triad nucleotide binding protein 1 (HINT1):

gttcctcccttcttccgagcctctcctctggccgcccgcgaggagagaggccgagatg

35 (Seq ID No: 407)

Homo sapiens heat shock 70kDa protein 1A (HSPA1A):

ctacctttttcgagagtgactcccgttggtcccaaggcttcccagagcgaacctgtgcggctgcagg
caccggcgcgctcagtttccggcggtccggaaggaccgagctcttctcgcggtatccagtgttccgtt
tccagcccccaatctcagagcggagccgacagagagcaggggaaccggcatg
40 (Seq ID No: 408)

Homo sapiens nucleolin (NCL):

cagtcttttcgctcagttctcgagctctcgctggccttcgggtgtacgtgctccgggatcttcagca
cccggcgccgcatcgccgtcgcttggttcttcttggaactcatctgcgccacttgctccgcttcaca
ctccgcccgcacatg (Seq ID No: 409)

Homo sapiens nuclear factor, interleukin 3 regulated (NFIL3):

ccgccccctttctttctcctcgccggcccgagagcaggaacacgataacgaaggaggcccaacttca
ttcaataaggagcctgacggatttatcccagacggtagaacaaaaggaagaatattgatggatttt
aaaccagagtttttaagagcttgagaatacggggaaattaatttgttctctacacacatagata
5 gggtaagggttgtttctgatg (Seq ID No: 410)

Homo sapiens protein phosphatase 1, regulatory subunit 3C
(PPP1R3C):

cagtctctcccagcgaccgcccgggggcaaggcctggagctgtggttcgaatttgtgcaggcagc
gggtgctggcttttaggtccgcccgcctctctgcctaag (Seq ID No: 411)

10 Homo sapiens protein tyrosine phosphatase, non-receptor type 14
(PTPN14):

agttctttccaactttttctcggcggagtgagcgcagcgggcccagactcgggggcaggttgctgt
gcttctccgggctcagccgctgctctcctggctcaggtcctcggggagccctagacagacatcaa
gtggccactggcgctccttccccctcccagctgagccatcctccccggcctcctcgggcgggacagc
15 cccgtgcttaggtttttctccttttctccccgggtgcgcctctgctcggactctcgcgcgggac
gcgcggaacactccctcccctttcgccctcctgcggtccttcccttcgccccctcctccgccagtc
actggaatcaattccgtggggaatcgggtccgcccgcgcgaaggacagcctttccgcgcgggactc
cggggcgccacggggggccatgtaagcagctatcttccagagggccacactgggcatggacaccctt
ttccctgcctggaggagcacaggtgatagtgaattttccagtcacgaaactgctaaggccatctc
20 aggggcgtgtgcgccagcataggcgggcggcgtccgaggaccacatagccatg
(Seq ID No: 412)

Homo sapiens selenoprotein P, plasma, 1 (SEPP1):

ctttcttttaagttagataacaatcagctcaggggtttgctctgcttgcaaggctcactgcaagaatg
aacattgaactttggactatacctgaggggtgaggtaaacaacaggactataaatatcagagtgtg
25 ctgctgtggctttgtggagctgccagagtaaagcaaagagaaaggaagcaggcccggttgaagtgg
ttgtgacaaccccagcaatg (Seq ID No: 413)

Homo sapiens serine hydroxymethyltransferase 2 (mitochondrial)
(SHMT2):

agctcttctcgcgcacgttctccgaacgggtcttcttccgacagcttgctgccctagaccagagt
30 tgggtggctggacctcctgcgacttccgagttgcgatg (Seq ID No: 414)

Homo sapiens tyrosine kinase with immunoglobulin-like and EGF-like
domains 1 (TIE1):

tttctcttctccccagcaccgacccacactgaccaacacaggctgagcagtcaggccacagca
tctgacccccaggccagctcgtcctggctggcctgggtcggcctctggagtatg
35 (Seq ID No: 415)

Homo sapiens coiled-coil domain containing 6 (CCDC6):

cctcctttccccagcccgccgcccagcatg (Seq ID No: 416)

Homo sapiens nuclear receptor coactivator 4 (NCOA4):

ggacctttcgcactcgggtcaggggtaaagcagcctgtcgttgccgggcagctgggtgagtcgggt
40 acctggcctgtgaggagcagtgaggagaatg (Seq ID No: 417)

Homo sapiens chromatin assembly factor 1, subunit B (p60)
(CHAF1B):

gtgcctctgactgtccgggtccctccagcattttgcagctttctcctgtcttgaagaagtagaacg
gtgcccagaaaacgtttttcccttcgagactcaggaggatgaaagtcacttgtgaaatagcc
45 tggcacaacaaggagcccgtgtacagcctggacttccagcatg (Seq ID No: 418)

Homo sapiens 3'-phosphoadenosine 5'-phosphosulfate synthase 1 (PAPSS1):

agccccgccccgctcgctggcctgccctcctcttgctaccctcccggcgagagaacccccggctgc
tcagcgcgctccgcggtcatg (Seq ID No: 419)

5 Homo sapiens Fas apoptotic inhibitory molecule 3 (FAIM3):

tgccctcctcttgctaccctcccggcgagagaacccccggctgctcagcgcgctccgcggtcatg
(Seq ID No: 420)

Homo sapiens N-acetylated alpha-linked acidic dipeptidase 2 (NAALAD2):

10 cagcctcctgccagcgcgctctctgtttctctgcagccccgaagctcgcaatgtagcaggcgccc
caagctcggtcctcaagaagccatggcggaatccaggggctgtgtacctttggatgtgcttggc
tgctgcgctggcatctttcctgatgggatttatggtgggtaagt (Seq ID No: 421)

Homo sapiens abl-interactor 1 (ABI1): ctgtctctttaacgcgagaggaag-
cgatgcagaggggtggaaaatg (Seq ID No: 422)

15 Homo sapiens potassium voltage-gated channel, Isk-related family,
member 3 (KCNE3):

cttccttttctgccttctctcctgctttctagctctgggctttccagctccgaagtcaatactga
gatccagatgtgtccagagacatcctgaagaggctcgggggtggaggagccttagtgtgtccaca
aagggactcctgaaactgactgagagccagt (Seq ID No: 423)

20 Homo sapiens target of myb1 (chicken)-like 1 (TOM1L1):
ggccctctggcgctaccatg (Seq ID No: 424)

Homo sapiens ubiquitin-like modifier activating enzyme 2 (UBA2):
cgcccttccccaccgcttccggcgcggtcggttctccgcctccgctccgcccggctcgt
ggttgtcccgcctatg (Seq ID No: 425)

25 Homo sapiens scavenger receptor class B, member 2 (SCARB2):

ctccctccttgcatgttgatccctggcggtgcggccccggccccggtgagcggcgacagaat
g (Seq ID No: 426)

Homo sapiens insulin induced gene 1 (INSIG1):

30 actcctcctttccccgccccgctccgttcggagagccggcgggcgccctctcggccaggaa
gcgcctcttgagcgcgtgtgaccgatg (Seq ID No: 427)

Homo sapiens kinesin family member C3 (KIFC3):

35 aggcctcttctgaggctctaggtgccccagtagcagggccttctgcagcaaggccgggaactgctg
caccattggtgtgttttaccttaagggactccaggcagcttccttgctgggaagatattcatttgc
tgggggtggggctgggggtgcagaggtaggaagtgctgtggctagaaggcggcctggccagcgagta
ggtggtggagcgagtgcagagcgtgtgcgctgtaaacagtgtgagtgcatg
(Seq ID No: 428)

Homo sapiens LIM domain kinase 2 (LIMK2):

40 aggcctcttctgaggctctaggtgccccagtagcagggccttctgcagcaaggccgggaactgctg
caccattggtgtgttttaccttaagggactccaggcagcttccttgctgggaagatattcatttgc
tgggggtggggctgggggtgcagaggtaggaagtgctgtggctagaaggcggcctggccagcgagta
ggtggtggagcgagtgcagagcgtgtgcgctgtaaacagtgtgagtgcatgtgcgccagcgctgca
aggacacggtaagggatgtacatgtattgtctcgtgagtaagagcttggtgtgtgtgttgggatggg
aagacacgtactggtatgagagcccgcgtgagaagtgtatgtgtgagtactcgcgtggaagtttgc
cactcggggttgaggctgtgcaaaagtacgcctgggtcaccaggtgtggggctgtgtgggctgcct

cgtgtgtgccagcccgtgtgcaggcctgttttgtgagagccttcagggaaacgcatgagcacgtgtg
ccagtgcgagtgcgggacgcggggagggcgaggagaccgagtgggagggccccgcgaaggagtggga
gtgggagtgggagtgcggcgaggagacctgcggggcgcgccccgggctgacgcgtgcgcgccagtgcg
cggtgagtgcgggcgcgcgccgcgcccccgccggggtcggagccggttgccatgggaacgcgc
5 cgcgccccgagttaatcatttcctgtggaaagtgtgcgggagggcgcgagcgggctggccgagga
ggagggcgggcggtggagctgcctcctgcggcgggcgggcgggccgagccccgggcgctgcgg
cgacgcctggatcctgcctccgccaggccgggctgcctgggtgccccgaggaggctgctgagccccag
gccatg (Seq ID No: 429)

10 Homo sapiens lectin, mannose-binding, 1 (LMAN1): cctcctccgcgttcca-
gaatccaagatg (Seq ID No: 430)

Homo sapiens MRE11 meiotic recombination 11 homolog A
(S. cerevisiae) (MRE11A):

cgttctctcccgcggaattcaggtttacggccctgcgggttctcagaggcaagttcagaccgtgtt
gttttcttttcacggatcctgccttttcttccgaaaagaagacagccttgggtcgcgattgtggg
15 gcttcgaagagtccagcagtgggaatttctagaatttggaaatcagatgcattttctgacatttgag
tacagtaccaggggttcttgagagaagaacctgggtcccagaggagcttgactgaccataaaaatg
(Seq ID No: 431)

Homo sapiens nascent polypeptide-associated complex alpha subunit
(NACA):

20 cttccttctgcaacaggcggtgggtcacgctctcgctcgggtctttctgccgccatcttggttcgcg
ttccctgcacagtaagtactttctgtgcgctactgtctatccgcagccatccgcctttctttcgg
gctaagccgccccggggactgagagttaaggagagttggaggctttactgggccacagggttccta
ctcgccccctgggcctccggacaaaatgggggtctgcggttggtgtcctggcaaaagcagggtagaag
ggctgcggggcgggccagaatccgagcctgcagagatgggagcagttgcagtggttgagggcgga
25 gaggagtgcgtcttgttttggaactgcttcacaggatccagaaaaggaaatg
(Seq ID No: 432)

Homo sapiens claudin 11 (CLDN11):

cgcccttcgcgctgagctcgcagcctccggcgcccacctccacctccagtgtcccgcctcgggcc
gtcgccctccagcggctcgcgagcgtgggagacgtacctgggcaggcactgtccagcccaggccca
30 ggcacagccgtgagggggcgaggcacggggacatcctggcgccaccatg (Seq ID No: 433)

Homo sapiens retinoblastoma binding protein 4 (RBBP4):

ccgcccctcccgaacgctcgaccccaggattcccccggtcgcctgccccgcatg
(Seq ID No: 434)

Homo sapiens acyl-CoA synthetase medium-chain family member 3
(ACSM3):

35 ccctcttcttttagactgccacgaggaaaaagcagatgtgagaactcaagggttcagggctgctcttc
taagaaacaagtctgccataatctccatctgtgttggaatctgttaactaatgaactggctctgt
gcaaactctgagtgtaaagcttccaacaagactgatg (Seq ID No: 435)

Homo sapiens syndecan binding protein (syntenin) (SDCBP):

40 cgctctcttacactcgggcctcagaagtccgtgccagtgaccggaggcgggcgggcgagcggttc
cttgtgggctagaagaatcctgcaaaaatg (Seq ID No: 436)

Homo sapiens serum/glucocorticoid regulated kinase 1 (SGK1):

agtccttctcattccttgcccccgcccaaggctctcttcaccttcccccggggggtcctctcgttt
tctgtctccaaatgctggcttccgcctttcctcccccgcttatttacttaattaaggccctggg
45 gctgcacccacccggcagctccttcgggggtgtggccgaagagctccgagggcggggctgaccgag
ccatattcgggcggtggccgggtgattgggtgagggcggggcctgcgcagggggcggggctgca

ggtttggccccgcagggagcgcagctggcgccgctgggagctggtggcgcggcgcaggtcccggc
cgagtgtggcgagcagtggtggcggttcccatcgccatgcgcgggggtgggtgcccgaaggtt
gcatgatggaatttgaacattacttcaagaggttttgatatttgattagtttaattgggtttgtcc
tctgctgactgtttcttcggatgcatttttgggtgtgctcttgagggattaaatg

5 (Seq ID No: 437)

Homo sapiens Wolf-Hirschhorn syndrome candidate 2 (WHSC2):

cgtccttccggtctcggctttgccacaaagcttcccgaagacgcggccgctacccggagacgcgg
tcgccacccagaagcgtctctccgggaagccccgctcgtgggaccgcgccacctgcgcgcctctg
cggccgcgagcccgacgggcgcgccatgttggggctctagcgagggacgcgtaggtgtcttcata
agatg (Seq ID No: 438)

10

Homo sapiens nuclear receptor subfamily 1, group H, member 3
(NR1H3):

cagtccttttgcaagagctgctaagagcgcgtgggtaaggagaggaaggggagagacatggaacttg
gctggtctgcagggaaatgccactgttttggccgggagtagggggcgaggtaggcgggagaggggg
tgcccggtggtgggagagccagcctggtggagaagctgcctgtgggcgggggtgaggaggggagg
gctgtggtcaccaggcaggaaggaggggtggcctgacctcggcagtccttccctcagcctttc
cccaaattgctacttctctggggctccaggtctgcttgtgctcagctccagctcactggctggcc
accgagacttctggacaggaaactgcaccatcctcttctccagcaagggggtccagagactgcc
caccaggaagtctggtggcctggggatttgggtgggtctgctccttag (Seq ID No: 439)

15

20 Homo sapiens glypican 6 (GPC6):

cctcctttctccttccctcttgctccagtgactgtctccaggatttctctcttcttctatttcagga
ggactctcacaggctcccacagcctgtgttaagctgaggtttcccttagatctcgtatatcccaa
cacatacctccacgcacacacatccccaagaacctcgagctcacaccaacagacacacgcgcgcat
acacactcgctctcgcttgctccatctccctcccggggagccggcgcgctccacactttgcccgc
acactccggcgagccgagcccgagcgcgtccaggattctgcggtcggaactcggttgagctct
gaacccccatggtggttttttaaacacttcttttcttcttctcctcgtttgattgcaccgtttc
catctgggggtagaggagcaaggcagcagccttccagccagccctgttggttgccatcgctcc
atctggcttataaaaagtttgctgagcgcagtcagagggctgcgctgctcgtccctcggctggca
gaaggggggtgacgctgggcagcggcgaggagcgcgcgctgctctggcggggtttcggttgagg
ggcaaggtgaagagcgcaccggccgtggggttaccgagctggatttgatgttgaccatg
(Seq ID No: 440)

25

30

Homo sapiens peptidylprolyl isomerase F (PPIF):

cggccttctgggcgcgcgcgacgtcagtttgattctgtgttctccccgcccgtgtcccggccgac
ccgcgcccgcgatg (Seq ID No: 441)

35

Homo sapiens ARP1 actin-related protein 1 homolog A, centractin al
pha (yeast) (ACTR1A): agttccttccccagaaggagagattcctctgccatg
(Seq ID No: 442)

Homo sapiens tripartite motif containing 28 (TRIM28):

ggctctttctgcgagcgggcgcgcgggcgagcgggtgtgcttgcttggtggcgcggtgctgggt
ttcggcggcggtgaggaagaagcgcgggcggcgccctcgaggagcgagcaggcagcagttggccg
tgccgtagcagcgtcccgcgcgcgggcgggcagcggcccaggaggcgcgtggcggcgctcggcctcg
cggcggcggcggcgagcggcccagcagttggcggcgagcgcgtctgcgctgcgcggcgggccc
cgcgccccctctccccccctgggcgcccccggcggcggtgtgaatg (Seq ID No: 443)

40

Homo sapiens aminoadipate-semialdehyde synthase (AASS):

cggccttccatcccagtttcttctaggaattcgagacctcccctgcagcgactcggaagattcgag
gcggcgggggacaagtcggcgccccagagcggacgagtcaccaggtgtcaagatg
(Seq ID No: 444)

45

Homo sapiens cornichon homolog (Drosophila) (CNIH):
ccgccttttctccgctggcaacggcgccgctccccgctcctcctccccagccatg
(Seq ID No: 445)

5 Homo sapiens M-phase phosphoprotein 10
(U3 small nucleolar ribonucleoprotein) (MPHOSPH10):
ctcccttcccttgcatgctgcattgtgtcgggagttgctgacagccatg (Seq ID No: 446)

Homo sapiens ubiquitin specific peptidase like 1 (USPL1):
ccgccttccctagtgagacgcgagtgggggaggagcagtcgaggggaacgtgggttgaaacgttgc
aactaggggtggagatcaagctggaacaggagttccgatcgaccggtaccaagaaggggagtgccc
10 gcggcagggttcattgaaaaaatccttagtgatattgacatgtctcaagtacataaattagccaa
tgactcggaatg (Seq ID No: 447)

Homo sapiens solute carrier family 23 (nucleo-
base transporters), member 1 (SLC23A1):
tggcctttgtcaagtcacccctcttctcctcaggaactgctcaaacctgtgccccaaagatg
15 (Seq ID No: 448)

Homo sapiens splicing factor 3b, subunit 4, 49kDa (SF3B4):
ggatctctttcgccatg (Seq ID No: 449)

Homo sapiens DnaJ (Hsp40) homolog, subfamily A, member 2 (DNAJA2):
ctgtctccctcggcctgtgccgcccgcgacgccgcttggtgggcccgcactccgctctgtctgcttcg
20 ccaccttctccccgagcactgcccggccggccgcatg (Seq ID No: 450)

Homo sapiens calicin (CCIN):
catcctctcttccaccctctcttctccctggtcaaccgctctgcaaacaacctcaatctgatccc
acaggcctgagaaagtctgctctccagtacctgctgctgatctgtttcagccgacaagaggcacca
tg (Seq ID No: 451)

25 Homo sapiens mannosidase, beta A, lysosomal (MANBA):
ctgcctttcgatctctccacatctcgggtggcgcgggatctcaagatg (Seq ID No: 452)

Homo sapiens microtubule-associated protein 1B (MAP1B):
aatcctttctcctgccgcagtgagaggagcggccggagcgagacacttcgccgaggcacagcagc
cggcaggatg (Seq ID No: 453)

30 Homo sapiens malate dehydrogenase 1, NAD (soluble) (MDH1):
gagccttttctcgctaaccgcctcgcctctccgagtcagttccgcggtagaggtgacctgactc
tctgaggctcattttgcaattgttgaattgtccccgcagttttcaatcatg
(Seq ID No: 454)

Homo sapiens microfibrillar-associated protein 1 (MFAP1):
35 gtttctctatcagtcgcgcagctgtgttcgcggactcaggtggaaggaatttcttctcttcggttga
cgttgctggtgttactgtttggaattagtcagtttccgggaatcaccgtcgtgcatcaacatg
(Seq ID No: 455)

Homo sapiens chaperonin containing TCP1, subunit 3 (gamma) (CCT3):
40 ggttctctctctccagaaggttctgccggttccccagctctgggtacccggctctgcatcgcgctc
gcatg (Seq ID No: 456)

Homo sapiens tubulin, alpha 1a (TUBA1A):
caacctctcctcttcgtctccgcatcagctcggcagtcgcgaagcagcaaccatg
(Seq ID No: 457)

Homo sapiens CD164 molecule, sialomucin (CD164):
ctttctcccgaacgccagcgctgaggacacgatg (Seq ID No: 458)

Homo sapiens cysteine-rich secretory protein 3 (CRISP3):
ctctctctgcaccttccttctgtcaatagatg (Seq ID No: 459)

5 Homo sapiens SMYD family member 5 (SMYD5):
cggcctccatgtgcgacgtgttctccttctgctgggcggtggcgggccgcgcggggtctccgtgg
aagtccgtttcgtgagcagcgccaaggtgaggtcggggcggggtcctgccgggagcctctccccagt
ccggccatg (Seq ID No: 460)

10 Homo sapiens kelch repeat and BTB (POZ) domain containing 10
(KBTBD10):
ctgccttttttacagctagacctgtgtgctgcaaggagctaaggccttcagtgtccccttccttacc
caggtttctcacagaatg (Seq ID No: 461)

Homo sapiens aldo-keto reductase family 1, member A1 (alde-
hyde reductase) (AKR1A1):
15 ccgcccccttgccacgcccacgtggccagcgccacctgcctcattgtgccaggagttctccaaacc
cgcgctgcggagtgagtgaccaagttccggccagttcgacctcgaggatccagaggtggagacggt
actacctccagctctgttttccatcccccttcaggtccttctcgggagggcggaaggcggtcca
ccctgcgcgtgatcctttatgcccgccccctgccctccctcgggtggaacttccccctcaccgc
cagacttaagctgaggatcggttgatctctggcggggtgcagaactgagcccaggccacagtaccc
20 tattcacgctctgtgcttggtgccaaggtttcaagtatcctccgcctcagcctgccagggtgctg
agattacatgtatgagccactgcacctggaaaggagccagaaatgtgaagtgctagctgaaggatg
agcagcagctagccaggcaaagggggcaatg (Seq ID No: 462)

Homo sapiens TRK-fused gene (TFG):
25 tgttcttccccacctgccacgtacagagcccaagttctcgctaggcttgttggggtcagcgcgatt
ggccggggggcgcgcgagcctgagcgaggtgcgggcggtcgcggaagggaaccgagggggcggtg
accaccgctccccgcgacgccccagtcagtgccctcgcgcccgccattcagcggagacctgcg
gagaggcgggcgggcgggcctccgcaagccgtctttctctagagttgtatatatagaacatcctgg
agtccaccatg (Seq ID No: 463)

Homo sapiens 3'(2'), 5'-bisphosphate nucleotidase 1 (BPNT1):
30 catccttctcaaaagacttattgacagtgccaaagctcggtactggacacaacgagggacctgggt
ctacgataacgcgcttttgcctcctgaagtgtctttggtccaacggttgttccagagtgtaccat
g (Seq ID No: 464)

Homo sapiens guanine nucleotide binding protein (G protein):
35 ttttctctctctcttttcaactgcaaggcgggcggcaggagaggttggtggtgctagtttctctaagcca
tccagtgccatcctcgtcgctgcagcgacacacgctctcgccgcccgcctg
(Seq ID No: 465)

Homo sapiens major histocompatibility complex, class II, DM alpha
(HLA-DMA):
40 caccctctcggggagggagttggggaagctggggttggtgggttagctcctacctactgtgtg
gcaagaaggatg (Seq ID No: 466)

Homo sapiens transmembrane protein 50B (TMEM50B):
tctccttctcgcgcgcgccctgaagtcggcggtggcggtttgaggaagctgggatacagcatttaa
tgaaaaatttatgcttaagaagtaaaaatg (Seq ID No: 467)

Homo sapiens lactoperoxidase (LPO):

cagtcttttctgctaagcctcagcgtctcctccaagccacatcaaaatctttccttctgggccttt
cccagaagtgaattcttgctggaaggtataaaagaccagctcctccaagcagagcaactccctggc
tgccgtgaaaagacaaggcactgggcagtgatg (Seq ID No: 468)

- 5 Homo sapiens NEL-like 2 (chicken) (NELL2): ctgcctttacaacagagggga-
gacgatggactgagctgatccgcaccatg (Seq ID No: 469)

Homo sapiens nucleobindin 1 (NUCB1): cgccctctgcggtgaaggagagacca-
cactgccatg (Seq ID No: 470)

Homo sapiens paired box 9 (PAX9):

- 10 aagcctctttcatcggggcacagacttccttttacttcttcttttgcctctcgcctcctcctcc
tgggaagaagcggaggcgccggcggtcgccgggatagcaacaggccgggactgaggcggtgcg
gaaagtttctgctctgggagtgcggaactggggccgggttggtgtactgctcggagcaatg
(Seq ID No: 471)

Homo sapiens cyclin-dependent kinase 16 (CDK16):

- 15 cgccctttattcttgctcggcctcgccacagagagcaaatcagattggctgggcgacaacctcaaa
ggcggggctgcacacgttcactacgggaatgaggtagcgggtggagggggcagttggcggggata
ggcgcctctagctaaggtggttaaaggccaataactcttcaggctgcctctcctcgaaaagtcatt
tctcggaacctttaaaatgccttctcctcccaagcacctcaagggactagaactgagtgttcatt
tgtcttttttctccttgcaaaagtcctggttgccaccatggggatgtaccaagtgagaccgagta
20 gggggaacgagtggtgattgacgcgccaggttactggccactgctcacctaggcgctagcaactt
ctgccaagatcggaactgagtactaaacagcctccacagttctcctggtgccgtctccggcttgg
cgccgcctcctcctctgggctcgcatggccgcgtcccctcccgtgcggacgggtccttggtag
atg (Seq ID No: 472)

Homo sapiens serpin peptidase inhibitor, clade E

- 25 (nexin, plasminogen activator inhibitor type 1), member 2 (SER-
PINE2):
ctgcctctttccggctgtgacctcctcgccgcccggcgttggtgctgcgtcctccgactccccgcgc
cgccgagaccaggctcccgtcctggttgccgcccgcacccgctccgcccggcccccctggggatcc
agcgagcgcggtcgtccttggtggaaggaacctg (Seq ID No: 473)

Homo sapiens pancreatic lipase-related protein 1 (PNLIPRP1):

- 30 aactcctttccccctgctgtgacgtacaggtgaggtaaacagtactgaagtccaggcgctcggtgc
tactgctctggcaatgcccggtgagactgaattatgtttaaattttattgtagatg
(Seq ID No: 474)

Homo sapiens peripherin (PRPH):

- 35 ggctccttcccagccccggcctagctctgcgaacgggtgactgcccatccttggccgcaatg
(Seq ID No: 475)

Homo sapiens RAD21 homolog (S. pombe) (RAD21):

gacccttttccccctccccgggcccagcccgcccaactcccagcggagagcaagggttttcttct
gttttcatagccagccagaacaatg (Seq ID No: 476)

- 40 Homo sapiens signal sequence receptor, delta (SSR4):
tttcttttctccttaggcagagaagaggcgatg (Seq ID No: 477)

Homo sapiens tissue factor pathway inhibitor (lipopro-
tein-associated coagulation inhibitor) (TFPI):

ctccctctttgctctaacagacagcagcgactttaggctggataatagtcaaattcttacctcgct

ctttcactgctagtaagatcagattgcgtttctttcagttactcttcaatcgccagtttcttgatc
tgcttctaaaagaagaagtagagaagataaatcctgtcttcaatacctggaaggaaaaacaaaata
acctcaactccgttttgaaaaaacattccaagaactttcatcagagattttacttagatg
(Seq ID No: 478)

- 5 Homo sapiens ubiquinol-cytochrome c reductase binding protein
(UQCRB): gcttctctttctggtcaaaatg (Seq ID No: 479)

Homo sapiens mitogen-activated protein kinase kinase kinase 12
(MAP3K12):

- 10 ccgccttttgtgctgcgccgagagccccgagggccagtggtcaccatcataccaggggccag
aggcgatg (Seq ID No: 480)

Homo sapiens sushi-repeat containing protein, X-linked (SRPX):
tggctctttcggtctcctgcccggggaagcgcgctgcccagggcgagctaagcgccccg
ctcgccatg (Seq ID No: 481)

- 15 Homo sapiens aminopeptidase puromycin sensitive (NPEPPS):
ccccctctccctccctccttgcgggccctccctccctccctccgcccccttccccgtagg
cagcccgcccgccagtcgcccgcacgcctccttcccagcccctagcgctccgggtgggtctctc
ccccgccccccaggtcccccggtcgtctctcctccggcggtcgcccgcgctcggtggatg
(Seq ID No: 482)

Homo sapiens fibulin 5 (FBLN5):

- 20 tcgccttctgcccggcgctcgagccgagcgcgggcgagggaagggtctcctcccagcgccgagc
actgggcccctggcagacgcccccaagattggttgtaggagtgtagccagttggtgagcgctgtaatc
tgaaccagctgtgtccagactgaggccccatttgcatgtttaacatacttagaaaatgaagtgtt
catttttaacattcctcctccaattgggtttaatgctgaattactgaagagggtgaagcaaaaccag
gtgcttgcgctgagggtctgtagtggtgggaggaccccgcgctctccccgtgtcctctccacg
25 actcgctcgccccctctggaataaaacaccccgagccccgagggcccagaggaggccgacgtgcc
cgagctcctccgggggtcccgcgcgagctttcttctcgcttctcgcatctcctcctcgcgcgctct
tgacatg (Seq ID No: 483)

Homo sapiens lysophospholipase I (LYPLA1):

cgctcttccctccgcttgcgctgtgagctgaggcggtgtatg (Seq ID No: 484)

- 30 Homo sapiens high mobility group nucleosomal binding domain 4
(HMGN4):
tcgtcttctctgtcttagggctggtgctggccctgcccacgcctagggctccggcgcgctcacgggc
ctcagctgggattcccgcgcccctcgagcgccacgagactcgacatcttccaggaacagcggtg
aggaggacagaagacccaacaggactgctcaagccacctgcaacactgctgctaccatg
35 (Seq ID No: 485)

Homo sapiens eukaryotic translation initiation factor 3, subunit M
(EIF3M):

agttcccttttccggctcggcgtggtcttgcgagtgagtggtccgctgtgcccgggcctgcaccatg
(Seq ID No: 486)

- 40 Homo sapiens Sec23 homolog A (S. cerevisiae) (SEC23A):
cctcctcttgacgtggcagaggcgccagccatg (Seq ID No: 487)

Homo sapiens cartilage associated protein (CRTAP):

cgctccttttcccttctcctccctcccttttcccttctccttcgctcccttcccttcccttccg
cgggcgcatg (Seq ID No: 488)

Homo sapiens vesicle amine transport protein 1 homolog
(T. californica) (VAT1):

ccgccccctcccgtggtatcccgagccgcggctcttcccagcgcgttccgccttcccagctgtgc
actctccatccagctgtgcgctctcgtcgggagtcaccagccatg (Seq ID No: 489)

5 Homo sapiens importin 7 (IPO7):

gcttctcttttctttcgcgcgggttgccgctgcggagcgcggcgggtccatgtgcgcagtgagtg
cgctattcctggccagtagcaccgagccccgggttgaccgagtcgcgcgtgcgatg
(Seq ID No: 490)

Homo sapiens ATG7 autophagy related 7 homolog (S. cerevisiae)
(ATG7):

10 gctcctttgcgacgcgcgcgcttcccagtggaagcgcgggcaggaccgcgttgcgatccggg
gcgcgcgcctcagagagagctgtggttgccggaagttgagcggcggtaagtgcgcgcggcggcg
aggggtgtagtggggtcttgctgggcccgggtttggaggcctggagtcaggggcgagctcgccagg
agggcgagggtcacagcaagtctcaggatcctcctctgccagtttctgggtggtccttctcctcc
15 agggactcactgattccggctggcgccttctgctgtagccgcgtcccctcagactggttcagtc
ggggtcttctgacttggaagctcgtgctgatttcctaagtcagcccctcctgtcctcttggtaggc
agtgtcagaatcttcagtggttggaacacgggagatgggacatttggttcccagcctggctgtgt
ctggatttgctgtctctggcacgttccctccccatctaagctgcttttccatctgcaaaatgggaa
tgataatccgccatttggttaagtgaggaggttaataagtttactttctgagaaagaagattctc
20 gattccttggttacagggttagaaactaatg (Seq ID No: 491)

Homo sapiens dynactin 2 (p50) (DCTN2):

cgctccctttgcccgcgccttagcccggaaccagcctctcccctaccggaacaccggcc
ccggctccaccgaggcccggtccccagccgctctcgcgcgcgcatg (Seq ID No: 492)

Homo sapiens acidic (leu-

25 cine-rich) nuclear phosphoprotein 32 family, member B (ANP32B):
agcccccttttccctccatgggttctctccgctcccgtagtaacttggtccgggggctccgctc
gcctgcccgcacgcgcgcgcgcccagaccgcgcgcgcgctccgcccgtagcaaacccttcc
gacggccctcgtgcgcaagccgggacgcctctccccctccgccccgcgcggaaagttaagtt
tgaagaggggggaagaggggaacatg (Seq ID No: 493)

30 Homo sapiens protein C receptor, endothelial (PROCR):

acttctcttttccctagactgcagccagcggagcccgcagccggcccagccaggaaccaggtcc
ggagcctcaacttcaggatg (Seq ID No: 494)

Homo sapiens actin related protein 2/3 complex, subunit 1A, 41kDa
(ARPC1A):

35 cgctccctctgggcttccgctcctccgcccgcgcccagcgagcctgttcgcgtcgactgccagag
tccgcgaatcctccgctccgagcccgtccggactccccgatcccagcttctctcctttgaaaac
actaagaataatg (Seq ID No: 495)

Homo sapiens chaperonin containing TCP1, subunit 4 (delta) (CCT4):

40 agggcccccttctccgctccgcctcctcccagcgcggcgccgctttctggaaggttcgtgaaggc
agtgagggttaccgttattacactgcggccggccagaatccgggtccatccgtccttcccagacc
aaccagacacagcggagtttgccatg (Seq ID No: 496)

Homo sapiens Niemann-Pick disease, type C2 (NPC2):

gcttctttcccagcttggaacttcgttatccgcgatg (Seq ID No: 497)

Homo sapiens phosphoribosylaminoimidazole carboxylase, phosphoribo
45 sylaminoimidazole succinocarboxamide synthetase (PAICS):

5

10

15

35

40

Homo sapiens ecotropic viral integration site 2B (EVI2B):
 ttttcctttcttagccaaatcaccaaatgtccagttagaacaagaatttagcattctgcaaaaga
 agttaacagctgagataacgaggaaatattctgaaatg (Seq ID No: 503)

Homo sapiens guanine nucleotide binding protein

(G protein), alpha inhibiting activity polypeptide 3 (GNAI3):

ggttcttctgggcgctaaggagctgacggagagggccaccgcccagcaatagacgggtgcctcagc
ctgccgagccgcagtttccgtggtgtgagtgagtcggggcccggtgtcccctctcccgcgcgcgcca
tg (Seq ID No: 504)

Homo sapiens polymerase (DNA directed), eta (POLH):

cggcccttcgcagcggggcgcgctgtcagacctcagtctggcggctgcattgctggggcgcgccgctc
tcgtctgatccctgctggggacggttgcccgggcaggatcctttacgatcccttctcggtttctcc
gtcgtcacagggaataaatctcgtcgaactcactggaccgctcctagaaaggcgaaaagatatt
caggagcccttccattttccttccagtaggcaccgaaccagcattttcggcaaccgctgctggca
gttttgccagggtgtttgttaccttgaaaaatg (Seq ID No: 505)

Homo sapiens solute carrier family 2 (facili-
tated glucose transporter), member 1 (SLC2A1):

cgctctctggcaagaggcaagaggtagcaacagcgagcgtgccggctcgtagtcgcgggtccccga
gtgagcacgccaggagcaggagaccaaacgacgggggtcggagtcagagtcgcagtgaggagtc
cggaccggagcacgagcctgagcgggagagcgccgctcgacgcccgtcgccaccgcgtaccg
cgagccagagccaccagcgcagcgtgccatg (Seq ID No: 506)

Homo sapiens zinc finger protein 138 (ZNF138):

gggtctttgtctcgtgcagcgggtgctgcaggtctggccttcacttttctgcgtcctcttactcc
tagaggcccagcctctgtggcgtgtgatctggttattgggagattcacagctaagacgccaggat
ccccggaagcctagaaatg (Seq ID No: 507)

Homo sapiens ubiquitin specific peptidase 3 (USP3):

ctttctttgacgcaagggctcgagacgcagccgcccgtcggccgagcgccggctagaagcgacacc
agacggagcctccggagttcctccgccccacctcgccgggtcctggagccgcagtcctcccagct
gccctcctcgtggccatg (Seq ID No: 508)

Homo sapiens calcium channel, voltage-dependent, gamma subunit 3
(CACNG3):

ctgtcttttctccagttttgagcgggggtgtcgggagcaggcggagagctttcctgagaggctgtgg
aagcagtgaacactcttctcagcgggtcgccctccagcagtgtatttttgccatccgcccctcac
ccccagcacacgcgctcgcacacacacgcacgcacgcacacacacacacacactcacacaga
gacctctctgggttttctttgccttgagtctcccggggtgtgagaagccaggcgcgtctcaaaccg
agctggcagctccaggctccggagccatgccctgcacggaccctcgtctttaccacgctcctgagg
aatgaaaggaaccagggaccctcagaaggcagcagtgatgcggaccaacccccggagcctgcac
ccttccgagggccataggcgaccagggaactggagagagctccagaaaggaaatcccagctttcc
caaagtccctgtggatgctgacaaaaggagacctgaatttttgaagagcctgtactagggttacc
ggctgcagagtgattttcccctccggcactgactctcccctccaacccccagccgtccagagtac
catgaagaattatg (Seq ID No: 509)

Homo sapiens guanine nucleotide binding protein

(G protein), beta 5 (GNB5): ttccctctccgctgcgtccccgcgcgaagatg

(Seq ID No: 510)

Homo sapiens chaperonin containing TCP1, subunit 8 (theta) (CCT8):

cttctccgcggtcttccgagcgggtcgcgtaactgcttctgcaggctggccatg
(Seq ID No: 511)

Homo sapiens prostaglandin E synthase 3 (cytosolic) (PTGES3):

cgctctttccgcgcgggtgcattctggggcccgaggctcgagcccgcgctgccgcccgtcgccctgagg
gaagcgagaagaggccgcgaccggagagaaaaagcggagtcgccaccggagagaagtcgactccct

agcagcagccgcccagagagggcccgcccaccagttcgcccggtccccctgccccgttcacaatg
(Seq ID No: 512)

Homo sapiens zinc finger protein 266 (ZNF266):

5 ttttcttctctggtggcggtttgggcttaatacagctttggcgaggtcggatgacgggtgggagccag
cggtggaaggggtggcgaaagtaccggtttgccccaggccgcccaggggctccttagagagacct
tgctgctccgctcgcgctccgcccggggcgcggggtcctcctggcgccgaggttcaaaaagcc
actcgagttgtcactgcgacggccctggggccaggagccgtttcgggatctgtcaacaacgagttt
tcgtcgttcgaatcaggttgactggtccttcatcccccaatctcccgtacctggcgagtccagct
cgtcgcggaatgctaagaaaagagtgaatgcaagctgagacaaaaatatggtatgatttagcc
10 atactgaaggggaaggaaataagagctgggcaaagcattctgtgaattggctgactccacttctat
ggtgagagagaggagtgcatacaagattactcccagtagagatggtttcagcatggtggccagctct
ggtctcagactcctgacctcaagtgatccaccacctcggcctccaaaatgctgggattacaggt
ataagccactgtgcctggccaaagataccgttaaccctggataaagagaatggaggttacctctgt
ccgtgtagattcctaagctgtcctggagtgatccttgaggataaaggaaaggtgctttgaagcacat
15 tcagccatcagccctgtgggatggcagccactgatttgtcctatgggtctttacagggaccagctct
gccttcaagaaaagacagaagtagaaaggggtgggtggctgactgtctgacaaattggtatcaggtat
gcaggaagtatatccttctccaaaatatcatacttgcatcaccaggtagacacatttcttctaca
cagaattatcttcagagcttcttaaagcaaataaagcctgcttcaaggactgagtcctagtcgaa
ttcccggaaggagtgaggcctgtcatattgtgtttatctagcatctgctcaagagtggtgctgcagt
20 ggagggaaatcagatgacctccagctctggttggtgttacatacaatcatgtgtaagaagtgccatt
caagccgtgtcactggaggggactgacagtgagattcagtgacttttgatgatctggctgtggact
tcaccccagaagaatggactttactggacccaactcagagaaacctctacagagatgtgatg
(Seq ID No: 513)

25 Homo sapiens methylenetetrahydrofolate dehydrogenase
(NADP+ dependent) 2, methenyltetrahydrofolate cyclohydrolase
(MTHFD2): gcttccctcccggcgagtcaccggcgcggtctatg (Seq ID No: 514)

Homo sapiens chemokine (C-C motif) receptor 9 (CCR9):
cttcctttctcgtgttggttatcgggtagctgctgctcagaaccacaaagcctgcccctcatccc
30 aggcagagagcaaccagctctttccccagacactgagagctgggtggtgctgctgtcccagggag
agttgcatcgccctccacagagcaggcttgcatctgactgaccaccatg
(Seq ID No: 515)

Homo sapiens heat shock 105kDa/110kDa protein 1 (HSPH1):
cctcccccttttgggtcggttagttcagcgccggcgccggtgtgagccgcccagagtgaggcagg
35 caaccgaggtgaggagcgacctgaggaggtgagccccgctttctccaggggttcttatcagcc
agccgcccgtgtccccgggggagtaggaggtcctgacaggccgcccgtgtctgtgtccttctg
agtgtcagaggaacggccagaccccgccgggagcagaacgcccagggcagaaagcggcggc
aggagaagcaggcagggggccggaggacgcagaccgagacccgaggcgaggcgaccgcgagccg
gccatg (Seq ID No: 516)

40 Homo sapiens StAR-related lipid transfer
(START) domain containing 10 (STARD10):
tggtcctttcttttatgattcacaaggaatgacctcttcatcgccctctcctaattcagtcctcac
aacagtccttttacaaatgggacaacagggttagaggaagttaggcagatttccagcatcatagaga
gtaaaggaccagggaaggatcaggattcaaggactgcacccaggctctgcttccagcttgctgtgt
gactttgggttaattttgttcccttagggaaactgagctttctcatttgtaaatgaaacaggctgtt
45 gggaggatcaaatgagatccaggggtgaaaacagcttagtttactttcaggaatttaccacgcgg
tatataaaggcaaaatattattatagtcaggtgattgtagattgaggaaccatttctcattctg
caaattgcaaactgagggcccaaagagggacaggggcttgccccaggtctcagcaggctgtgagc
aagagctaaagcctaatactcctgcttgggctggagcccttcttgtacccaggggtcagtg

tctttgttggtacaggcttagattgactgactgtaccctgagaacctaggggagtcctgttccc
aattcttctcctacccccaccttggcctgatggaggaagacctgctgtgttgagatgagcaccag
agccaagaagctgaggaggatctggagaattctggaggaagaggagagtgttgctggagctgtaca
gacctgcttctcaggtcccaggaaggtggcgtcagcatctgcagccgcgtcgacgttgtcggagc
ctccgcggaggaccaggagagccggactaggaccagggccctgggcctccccacactccccatg
(Seq ID No: 517)

Homo sapiens UTP14, U3 small nucleolar ribonucleoprotein, homolog A (yeast) (UTP14A): ctttccttcggtccggttcttggtccatgtgagagaagctggctgctgaaatg (Seq ID No: 518)

10 Homo sapiens SUB1 homolog (S. cerevisiae) (SUB1):
ggttctctgtcagtcgcgagcgaacgaccaagagggtgttcgactgctagagccgagcgaagcgat
g (Seq ID No: 519)

Homo sapiens minichromosome maintenance complex component 5 (MCM5):

15 ccgcctcttgtttttccgcgaaactcggcggctgagcgtggaggttcttgtctcccctggtttgt
gaagtgcggaaaaccagaggcgcagtcattg (Seq ID No: 520)

Homo sapiens RNA binding motif (RNP1, RRM) protein 3 (RBM3):
tactctttatcaatcgctcttccggcgcagccccgtccctgtttttgtgctcctccgagctcgctg
ttcgctccgggttttttacgttttaatttccaggacttgaactgccatg (Seq ID No: 521)

20 Homo sapiens KDEL
(Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor
1 (KDELRL1):
ctccccctctcgtctctcctccctcttcccggtccagctccgccgagctccagcctttgctccc
cctcccaaagtccctccccggagcggagcgcacctagggtccctcttccgtccccccagcccagc
25 taccggttcagaccagcagcctcgggggcacccccccgccagcctgctcctccctcccgctcagccc
tqccagggttccccagccatg (Seq ID No: 522)

Homo sapiens StAR-related lipid transfer (START) domain containing 3 (STARD3):

30 agatcttcttccgctctgagggcgtactgagggccgaggccggactgcggttggggcggaagag
ccggggccgtggctgacatggagcagccctgctgctgaggccgcgcctccccgcctgaggtggg
ggccaccaggatg (Seq ID No: 523)

Homo sapiens heterogeneous nuclear ribonucleoprotein A0 (HNRNPA0):
 cggcctctttgtgtggtgccagatagggagcggaggtggcggcggcgcggtagcgggtggcctt
 gttgtcttcagttctctcggtcgcccttagcggcaccgctcccttccctcccccttctctc
 tcttccttccctccctcccttcccttttcccttccccgtcgggtgagcggcgggggtggctccag
 caacggctggggcccaagctgtgtagaggccttaaccaacgataacggcggcgacggcgaaacctcg
 gagctcgagggcgggggcaaggcccgggccttgagatg (Seq ID No: 524)

Homo sapiens chromobox homolog 1 (CBX1):

40 ggctcttttgttcggctgaggggaggggcggttgccggggcctgcggtacgccgcttcagtgaagg
acgccactgcggccaccggttgctgccttcctgggcgccactccccaggcgaccgcagcgac
gcgccagcagcgacacccgattcctctcggtcttgggcgctgctctgaggtgaggagcccgct
ggaggcgggagagctgggggagggggcgcgggcgggcgggcgggagccctgcgtgagggaaacg
cgctttcgaggcggaggttaggagcggggagcgcgcccggtccagcgtcctgcttctccgcttcc
cgcgctgagctcttcgcctgtcgctgaggcgtcggtgccagctgcgtgaaggatggagagggcggg
45 gcggaatcctgagccagagactgagtgcctgggggtgggcccagcacttgggggcccgtcttcgg
ggccgggtggtctggaacaatgttgcttggctggcggtgcgggatagggcggaaggggacagg

cttgaggccttgatagggcgtgaggaggcgcatagcaccgcacaacccgaggtttgtaactgtattc
ggaagacgccgggtccggctgggactgccagaggaacctggctttgcaggactacggaggagtaac
gtcgagtgaattggaagagggcccgaggccgcacaagcagcgtcaccctttacaccagaaagctgg
cgggcactatg (Seq ID No: 525)

- 5 Homo sapiens myeloid/lymphoid or mixed-lineage leukemia (tritho-
rax homolog, Drosophila); translocated to, 11 (MLLT11):
cgcccttcttaggaggggctgcattgcaggggagagtgaactgacagactcagtcactgaagagg
gaaaaggagtgagaagacaaagccgtcaaagcccaacagctttgtattttctccagcccggcgcag
accccgagctcccaggcactccctccatctttggaacacgccagtaattgattgataacaggaa
10 gctatg (Seq ID No: 526)

Homo sapiens interferon-induced protein 44-like (IFI44L):
ttttctttctttcctagagtctctgaagccacagatctcttaagaactttctgtctccaaaccgtg
gctgctcgataaatcagacagaacagttaatctcaatttaagcctgatctaaccctagaaacag
atatagaacaatg (Seq ID No: 527)

- 15 Homo sapiens cyclin I (CCNI):
acttcttctccttccctctcttccctccctccccagccttcccccgcgagcggacgcggcagc
gcctctgtctcgctttttcttatttttccccccttcccttcttttttttttttcttttcttt
tctccctccccctttcaccatttccctcggaggcgcttcccccgggcaggggcagagccggt
ctcacccccgcctctccccggccccgcgcgcctatggcgagagggagccccctcccaaccggg
20 ctcgagcggcgggcgccctcaggccgggggtcatcatggaactaattcgctgaccgaccagcggcc
gcagccgtgctccccgctcgagcgccagcgcccgcgcccccgatccgcttccccctttct
ccctcctcagttggccgagtcgtcccgcgcgaccgcctccgcgcgccctatgagaatgaggtggt
acgggcccccgatgaccccgctcaccactgtgaggcctacagctctgccccggaggaggaggag
gaggaagaggaggagaaggtagctacagcaagctgggtagcaggcagatccaaaggatatcatg
25 (Seq ID No: 528)

Homo sapiens methionyl aminopeptidase 2 (METAP2):
cattccctcgcgctctctcgggcaacatg (Seq ID No: 529)

- Homo sapiens leukocyte immunoglobulin-like receptor, subfamily B
(with TM and ITIM domains), member 4 (LILRB4):
30 gtctctttgtcctgccggcactgaggactcatccatctgcacagctggggcccctgggaggagacg
ccatg (Seq ID No: 530)

Homo sapiens destrin (actin depolymerizing factor) (DSTN):
gggtctctcggtcccgagccgtgaggaggacggtctgcatactcgctgccgcgggtccctccc
ccgctccctgcgaccgccgcggaagatg (Seq ID No: 531)

- 35 Homo sapiens eukaryotic translation initiation factor 2D (EIF2D):
gggcccttttcgcgccgggccccagcatggctgccccacggctgaggcctggcagctgctgcg
ccctcgctttcttgacattccctggcttctgtgctcttccccaggccaccccagcagacatg
(Seq ID No: 532)

- 40 Homo sapiens histamine N-methyltransferase (HNMT): ctgtcttttctca-
gaaaaccaaataatg (Seq ID No: 533)

- Homo sapiens ras-related C3 botulinum toxin substrate 1
(rho family, small GTP binding protein Rac1) (RAC1):
gtttctctgcagttttcctcagctttgggtgggtggccgctgccgggcatcggttccagtcgcg
agggcgaggcgcggtggacagcgccccggcaccagcgccccgccccgcaagccgcgcgcccc
45 tccgccccgagccccggcgttccctatctcagcgccctgcgcggccgccccgagcga

gcggccctgatgcaggccatcaagtgtgtggtggtgggagacggaaacaagaatctcagtgttaacc
cgagcaaaatcgcgctctcagcgcttgcttgatagagctgtaggtaaaacttgccactgatcag
ttacacaaccaatgcatttcctggagaatatatccctactgtctttgacaattattctgccaatgt
tatg (Seq ID No: 534)

- 5 Homo sapiens signal recognition particle 72kDa (SRP72):
tcgtctcctccaagatg (Seq ID No: 535)

Homo sapiens zinc finger protein 33B (ZNF33B):
ccgcctttccttttgtttgtctcacgttttgctgggagggcggtcccgggatttcaggggtctacc
ggctctcttatggcgaatgcaaccggaagagagagtggctgtatcttcagagttgtctccgtctt
10 tccaagaacagaacaaaatg (Seq ID No: 536)

Homo sapiens zinc finger protein 16 (ZNF16):
gcctcctttccaagcgcgacccgttgaggctctgtcatg (Seq ID No: 537)

Homo sapiens zinc finger protein 33A (ZNF33A):
ccgcctttccttttgtttttctcaggttttgctgggagggcggtcccgggatttcaaggggtctacg
15 cgctttttctatggcgaatgcaaccgacgagggagtggtgtatcttcagagttgtctccgtctt
tccaagaacagaacaaaatg (Seq ID No: 538)

Homo sapiens butyrophilin, subfamily 3, member A3 (BTN3A3):
ctttctttttcctttcttcggaatgagagactcaaccataatagaaagaatggagaactattaacc
accattcttcagtgggctgtgattttcagaggggaataactaagaaatggttttccatactggaacc
20 caaaggtaaagacactcaaggacagacatttttggcagagctgctcactccttgctcagctcagtt
ttctgtgcttggaacctctgggcccacacctggccatg (Seq ID No: 539)

Homo sapiens butyrophilin, subfamily 2, member A2 (BTN2A2):
ctctttgggatgctttgttgtctggtggtgactgtgcccatgggtgagttgtatcggaatcgctc
atgtgaggatcagaggggaaaagaaaacagaggcctctggtctctgcctgccctgggtgctcatg
25 (Seq ID No: 540)

Homo sapiens nudix (nucleo-
side diphosphate linked moiety X)-type motif 21 (NUDT21):
acgcctcctcttgcgctgtcctgttaatggcgggcagtagccgctgaggggattgcagataaccgc
ttccgcacggggaaagtctaccctgcctgccactttctgctcgccgctcagcgccggagctcgcca
30 gcatg (Seq ID No: 541)

Homo sapiens stathmin-like 2 (STMN2):
tgctctttctctagcacgggtcccactctgcagactcagtgcttattcagtcttctctctcgctct
ctccgctgctgtagccggacctttgccttcgccactgctcagcgtctgcacatccctacaatg
(Seq ID No: 542)

35 Homo sapiens katanin p60 (ATPase containing) subunit A 1 (KATNA1):
caccctcttccgcgctcccgcagcgacctcgctcccggggcgacgcccgcgtgcgccagagt
cgccgaggtcgctcccggcaccggaagtgacctggcggtttgtcttcaaattctcggcgagcag
gagccgcgccggcaggtggtgttgacgattgaactgggcagtagctggggccgtgagcggagagcaa
agtgggctggactgggtcaggccctccttctcgtgctgccgggatctccactccgccaatcccctgt
40 gcctggcggttggcggtttcccagaggagcttgggcccgcagcttacagttgaacatg
(Seq ID No: 543)

Homo sapiens butyrophilin, subfamily 3, member A2 (BTN3A2):
ctttctctttttcctttcttcgggatgagaggctaagccataatagaaagaatggagaattattga
ttgaccgtctttattctgtgggctctgattctccaatgggaataccaagggtggttttccatact

ggaacccaaaggtaaagacactcaaggacagacatttttggcagagcatagatg
(Seq ID No: 544)

Homo sapiens CLK4-associating serine/arginine rich protein
(CLASRP):

5 cggcctttcatttccgcttccggtgcgggccgcgcgagcgcagcgggtgggagggcggcgaccagc
cggttgaggccccaggccttggcctcaccacaatg (Seq ID No: 545)

Homo sapiens clathrin, light chain A (CLTA):

ctccctcctggcgcttgtcctcctctcccagtcggcaccacagcgggtggctgccgggctgggtgtc
gggtgggtcggttgggttttgtctcaccggttgggtgcccgttcagttgcccgccatg
10 (Seq ID No: 546)

Homo sapiens NADH dehydrogenase (ubiquinone) flavoprotein 1, 51kDa
(NDUFV1):

gcgtctctatcgcgccagttcctcagcctcagtgctatgaaggtgacagcgtgaggtgacccatct
ggcccgcgcgatg (Seq ID No: 547)

15 Homo sapiens signal sequence receptor, gamma (translo-
con-associated protein gamma) (SSR3):

gggcctttgccgccttggcgccggctctacgttccctgttctcgcctgcagctccgccatg
(Seq ID No: 548)

Homo sapiens valosin containing protein (VCP):

20 gcttcccttccgatgattcggctcttctcggctcagtcagcgaagcgtctgcgaccgtcgtttg
agtcgtcgtgccgctgccgctgccactgccactgccacctcgcgatcaggagccagcgttgttc
gcccagcgcctcgtgccggtgggaggaagcgagaggggaagccgcttgcgggttctgcgccgctgc
tcgcccaccgcctggaagagccgagccccggcccagtcggctcgttggccaccgctcgttagccgtta
cccgcgggcccgcacagccgcggccgggagaggcgcgccatg (Seq ID No: 549)

25 Homo sapiens zinc finger protein 195 (ZNF195):

gggcctttgtcccagacagagctccacttctgtccccgcggctctgtgtcccctgctagccgtagg
tcgtgtgacccgcaggcaccgggagatccagaagtgaacgccaggctctctggaggccaggagat
g (Seq ID No: 550)

Homo sapiens testis-specific kinase 2 (TESK2):

30 cagtctttcgcggcccgggagctcagcagagctaccagctgccctgttggcttcgctggtcggatc
gtcctcctggccccgcaaacaggcggggggagcgggccccgactgtggggccatggcagtagtctc
ctcgttcgcgcgcgcgctagcctagctgagtcgcgggcttctgcgctaggggtcccaccgcctc
cgcaggctaaggagccgctgccaccaacgagctgtgagggttactatgctccctctttgccgccgt
ctcctcctcttgcgcgcgaggacccctctggctgctcagtcctgcctcagtgtaaaccagaag
35 agaagtaaaattcaacaaaaatttatgtgtggagttccttcttaaaagaagaaaaaagtgattatt
tagactatg (Seq ID No: 551)

Homo sapiens family with sequence similarity 107, member A
(FAM107A):

40 agccctccttgctagtctgggacttcccgggtggagtgaggaacccagcaacacgctcctgacttcc
cttcccaaggactcgacctgagaaggacacagcagtcctctgaatttcagtcctcctctttgatgt
gaagaaaatgaaaagctgaacagttgtggaactgtggatagagttagacaataaggccgcatg
(Seq ID No: 552)

Homo sapiens serine/threonine kinase receptor associated protein
(STRAP):

45 ccctccctccctttccctccctcgtcgactgttgcttgctggctcgcagactccctgacccctccct

cacccctccctaacctcggtgccaccggattgcccttcttttctgttgcccagcccagccctagt
gtcagggcggggctggagcagcccgaggcactgcagcagaagagagaaaagacaacgacgaccc
tcagctcgccagtcggtcgctggcttcgccgccgcatg (Seq ID No: 553)

Homo sapiens mitochondrial ribosomal protein L3 (MRPL3):

5 ctttctttccgctcgagagagcatcgggccggcgaccgttccggcgccattgcgaaaacttcccca
cggctactgctccacgtggcggtggcggtgggactccctgaaagcagagcggcagggcgcccga
agtcgtgagtcgagtccttccgggctaatacatg (Seq ID No: 554)

Homo sapiens zinc fingers and homeoboxes 1 (ZHX1):

10 ctcccttccccctccgccccggacggccgctggggcgcgccctctcctcgacccccaccctga
gtccccacactccgcggggaccagagctgctgaggccctttgcgggcccgccgagcggttccgg
gtttagggttcacaggtcagagttgactccctgaaaagtgcagccggtttgaaatgcaagatggcg
gcggcgtggcgctgagaggcgcgggccctgcaggagaagacagactgctgctttggacctgtt
ggtaatgatggcctgagctaaacatctaactagaaggatacccttccatttcaaagaacagaatg
15 ctaaggaagctgtggcaagtgttgagttgtgcttcaaaaatttcagaaattcagcagttttta
tctgccaacaataagctctttacttgattgcacatgagaaagctgctaatagacttgttgagca
caaaaatggacttgaagaaccaaagccattgttttcaaatgaagaacactgaacagttttaagcc
tcgatgctttttaatacaccactgagcttttctcataacatcagaatg (Seq ID No: 555)

Homo sapiens calcium binding protein P22 (CHP):

20 ccttccttccctccctccttccctcctgtcgccgtctcttctggcgccgctgctcccggaggagct
cccggcacggcgatg (Seq ID No: 556)

Homo sapiens ecdysoneless homolog (Drosophila) (ECD):

ctttctctcaggatttccgctggcttcaggttccggtcaggcgtcgggacagagcctgatccaggc
ttcgggcgccggtggcagctctcgatcagctctcgagtcggagaggcggttaaggaaaggtgcc
cagcagagacgcgaaggagaggccctagaaccttttcaaagaagaatg (Seq ID No: 557)

25 Homo sapiens V-set and immunoglobulin domain containing 4 (VSIG4):
gagcctctttggttagcaggaggctggaagaaaggacagaagtagctctggctgtgatg
(Seq ID No: 558)

Homo sapiens prohibitin 2 (PHB2):

30 tgccctttctttcgccagccttacggggccgaaccctcgtgtgaaggggtgcagtacctaagccgga
gcggggtagaggcgggcgccgaccccccttctgacctccagtgcgcggcctcaagatcagacatg
(Seq ID No: 559)

Homo sapiens signal transducer and activator of transcription 1, 9
1kDa (STAT1):

35 ctgccttttctcctgcccgggtagtttgcctttcctgcgcagagctctgcggaggggctcggctgcac
cggggggatcgcgctggcagacccagaccgagcagaggcgacccagcgctcgaggagaggctg
caccgcccgcggcccgctagcccttccggatcctgcgcgcaaaaagtttcatttgctgtatgcc
atcctcgagagctgtctaggttaacgttcgcactctgtgtatataacctcgacagtcttggcacct
aacgtgctgtgcgtagctgctcctttgggtgaatccccaggcccttggtggggcacaaggtggcag
gatg (Seq ID No: 560)

40 Homo sapiens heat shock protein 90kDa alpha (cytoso-
lic), class B member 1 (HSP90AB1):

agctctctcgagtcactccggcgagtggtgggactgtctgggtatcggaagcaagcctacgttg
ctcactattacgtataatccttttcttttcaagatg (Seq ID No: 561)

Homo sapiens cancer susceptibility candidate 3 (CASC3):

45 cgttctccgtaagatg (Seq ID No: 562)

Homo sapiens nuclear cap binding protein subunit 2, 20kDa (NCBP2):
gcttctctgcactatg (Seq ID No: 563)

5 Homo sapiens non-POU domain containing, octamer-binding (NONO):
cgctcttttctcgggacgggagaggccgtgtagcgctcgccgttactccgaggagataccagtcggt
agaggagaagtcgaggttagaggggaactgggaggcactttgctgtctgcaatcgaagttgaggggtg
caaaaatg (Seq ID No: 564)

Homo sapiens lectin, galactoside-binding, soluble, 9 (LGALS9):
atttctttgttaagtcgttccctctacaaaggacttcctagtgggtgtgaaaggcagcgggtggcca
cagaggcggcgagagatg (Seq ID No: 565)

10 Homo sapiens chaperonin containing TCP1, subunit 5 (epsilon)
(CCT5): cggtctccgccggttggggggaagtaattccggttgttgaccatg
(Seq ID No: 566)

15 Homo sapiens haloacid dehalogenase-like hydrolase domain containin
g 1 (HDHD1): cttcctcctcgccccacccagacccagaaggcgccaccatg
(Seq ID No: 567)

Homo sapiens glutamate dehydrogenase 2 (GLUD2):
cttccttcctagtcgcggggagtcgagaaagcgacactgttcgcgaccgtcacgcacccctcct
ccgcctgccgcgatg (Seq ID No: 568)

20 Homo sapiens general transcription factor IIIC, polypeptide 3, 102
kDa (GTF3C3): ggttctctgtcccgggttctgggggtgcacagacagaccctgtaaactg
(Seq ID No: 569)

Homo sapiens general transcription factor IIIC, polypeptide 5, 63k
Da (GTF3C5):
25 gggtcctcgcgtggctagtaggagagactgggtgcttgccccgcccgggtggactaactcgcttaatt
ttaaataaaaagtcgaggacacggcggtcggttttcccgaagacatgggccctcccatgggccattt
gctccctggaggccctcgcgtcttgctgagcccggggagttaggatgacgcgagcgggtgagggagc
ccggaacgattccttcgcggaacaattgaggcgaggcctttgggagtactttgtgggacggaccct
ggcgggccctgccagacgcacagggatg (Seq ID No: 570)

30 Homo sapiens ancient ubiquitous protein 1 (AUP1):
ccgccttcccaagagcccctgcggccgggcgcgaaaatggcggcgggcgacggccggggcgctcc
tgaagcagcagttatg (Seq ID No: 571)

Homo sapiens coatomer protein complex, subunit gamma 2 (COPG2):
cggccttctcgcagcctcttccgctcgccggctgcggcgccctgggacggttgcggtgggtctgggc
gctgggaagtcgtccaagatg (Seq ID No: 572)

35 Homo sapiens apoptosis antagonizing transcription factor (AATF):
cggctctctggcggagtcggggaatcggatcaaggcgagaggatccggcaggggaaggagcttcgggg
ccgggggttgggcccacatttacgtgcgcgaagcggagtggaacgggagctggtgacgatg
(Seq ID No: 573)

40 Homo sapiens integrator complex subunit 6 (INTS6):
tctcctctttctccaccacctcgggccccggtgtccccggccagcactatg
(Seq ID No: 574)

Homo sapiens F-box and leucine-rich repeat protein 4 (FBXL4):
tcttccttccgggtcgcgctaggccgggcttgcggcggttggtgccgcatctagagagtcggggagc

cgccccgcacccaggccttctcgcgctgcctggcgctgggaagcccgcggcgcgcgcctctcc
cggaccctgcagggtaaaagaatgtcacatgtcagcatttgtacctgaagtcagcatgcaaagttc
agggtacctggatgaatgccaaacttttgcatttcccatgtgtatcctgtgaccattctatctggga
acatccttcaaagagttcatgcatcttactgaggacacctgaccttttgaagcttcataattcaca
tctagatg (Seq ID No: 575)

Homo sapiens guanine nucleotide binding protein
(G protein), gamma 3 (GNG3):

gctccttctagcatccttcatccttcagggtaccagccatccagacagtgccttgagctgcagaaact
gagaccagacctctggcctggccctccccaggggcctccttctgtatagtcactgcttctgcatca
gatactttcagctgcaactccctactgggtggggcaccatttcaggcagaagggttttggtaccct
ccactgaccctacaccagggtgctactgccgcttgtggcttcaggatg
(Seq ID No: 576)

Homo sapiens histidyl-tRNA synthetase 2, mitochondrial (putative)
(HARS2): aggccttttgttctgtcccggaaagccggcgctcctgccgcgcgatg

(Seq ID No: 577)

Homo sapiens interleukin enhancer binding factor 3, 90kDa (ILF3):

cctcctcctcctcttctcgccattgcagttggaccagcagcccgcgcgccacgcgctggcttttg
ggggcagaccccgcgggctgtggcaggagggcggcggcgcgctgcggtcgaagaaggggacgc
cgacaagagttgaagtattgataacaccaaggaactctatcacaatttgaaaagataagcaaaagt
ttgatttccagacactacagaagaagtaaaaatg (Seq ID No: 578)

Homo sapiens polymerase I and transcript release factor (PTRF):

gtttcctctgctctccgctctcgccgctagctctcctcccttccgctcctgcttctctccgggtc
tcccgctccagctccagccccaccggccggtcccgccacggctccgggtagccatg
(Seq ID No: 579)

Homo sapiens 5'-3' exoribonuclease 2 (XRN2):

tgccctctgccgctgctcccgtctcttttggttacgctcgtcagccggtcggccgcccgcctccagcc
gtgtgccgctatg (Seq ID No: 580)

Homo sapiens 2-hydroxyacyl-CoA lyase 1 (HACL1):

ccgcctcttccctcccggttggttaaggcagttggttgccctcctgtccgctcagaggtgcagtacca
gaggtggcgctgctgccgatttcgcggttgcttgctggatgattccgcttgtttgccggctgcgctg
agtgttagagcttttcgggtggaagatg (Seq ID No: 581)

Homo sapiens zinc finger protein 346 (ZNF346):

ggctctctaccggtgaggggttgcggggaagatg (Seq ID No: 582)

Homo sapiens microtubule-associated protein, RP/EB family, member
3 (MAPRE3):

cagtctctgtgcgttgaagccggagaccgcggcggcctcagcgaggaccctccgccccggagccgc
cggccggagccgcagcctctgccgcagcgccccgccacctgtcccctccccctccgcctccgccc
gagccgcctcgtgcactctggggatg (Seq ID No: 583)

Homo sapiens splicing factor 3b, subunit 3, 130kDa (SF3B3):

gtgcctttttccgcccgcgccaccagaatgtccctgtcttgaggtctaattggcggacgccagtat
gttgaggttggtgggtggttaagttttgaaggaggtagcatccgttgatatccacaccatcctt
ctcgctgcaggctttcttgactccgtactgttggtgtaaccaaggcctggaggtctgggtggctc
aggtttctgcagccatg (Seq ID No: 584)

Homo sapiens spondin 2, extracellular matrix protein (SPON2):
ctgcctctcgcgtggaggccaggccgtgcagcatcgaagacaggaggaactggagcctcattggccg
gcccggggcgccggcctcgggcttaaataggagctccgggctctggctgggacccgaccgctgccg
gccgcgctcccgcgtgctcctgccgggtgatg (Seq ID No: 585)

- 5 Homo sapiens solute carrier family 13 (so-
dium/sulfate symporters), member 4 (SLC13A4):
ttttcttttctgctttgcaggcccaggctcaaggcaaattataagtagggaaccaatttgaggga
agacatgtgaacagagttaaggtaccacgtcctgggagcgaccagcagccccacctgaagtccgca
tgcaactctgacaagctcaggtgcttgttttaaggaaaggggctactagagctttaccaacagcga
10 gccaggtgggagatgaaacaggtactccccaaaataggatcatccgagggaggaaaactgatggag
agcacaatgtgctctgagcgttttttaagtgttttaagctttttaaatgatttcttcaaggccgagca
gcagcagcaaagggtgtggcttaaggattaagggggtttctgctgacacctagaatgaagttaactc
tattactaatcaagccgagagaggagggccactatgccccggtttatcatcctttcccagttcctttt
tgctggtcacaaaacgatgctcatcaatcccacctaaagcaggaggccaggagcccagcctcttgt
15 agaaacagcgagggtataactgccctcccgttctgcccccaagacgaaggaggactctcggaagcc
aagaaagggttaagaagtctttctggatagagagcagtgcccaggcaggaagcctttcgccggcag
agcgggggtccaaggacgagctggagaggacagaggcgcatg (Seq ID No: 586)

- Homo sapiens PRP6 pre-mRNA processing factor 6 homolog
(S. cerevisiae) (PRPF6): attcctttccttcctagccttggtcgctcgccgccaccatg
20 (Seq ID No: 587)

Homo sapiens eukaryotic translation initiation factor 3, subunit K
(EIF3K):
ccacctcttctgttcccgtccttgaggacgccgtgcggggtcagtggttagcctccagccctgggt
gtggaaggcgacagaagtcacg (Seq ID No: 588)

- 25 Homo sapiens ataxin 10 (ATXN10):
ccccctcccccgcggcgcgtctcctcctcccgccctgaggcgagctctgggctcagcctagagctct
ccggcgcgcgagcgttcagggcagcgcgggctgcagcgcgcgcggttagggctgtgtaggg
cgaggcctccccctcctcctcgccatcctactcctcctcctcgtcatcctcccccttcgtcctc
ctcgcttctcctcctcctcgtcaggctcgacccagctgtgagcggcaagatg
30 (Seq ID No: 589)

- Homo sapiens secretogranin III (SCG3):
cttccttctcacttcctctgcaggagggagcgagagtaaagctacgccctggcgcgagctctccg
cgtcacaggaacttcagcacccacagggcgagcagcgctcccctctacctggagacttgactcccg
cgcgcccccaacctgcttatcccttgaccgtcgagtgctcagagatcctgcagccgcccagtcgccg
35 cccctctcccgccccacacccacctcctggctcttctgtttttactcctccttttcattcataa
caaaagctacagctccaggagcccagcgccgggctgtgacccaagccgagcgtggaagaatg
(Seq ID No: 590)

- Homo sapiens polymerase (DNA directed), mu (POLM):
cttccttccgtctcgtcggagtttccctctgcgttcgctccgcgctgctggaggctgtcgtccca
40 atg (Seq ID No: 591)

- Homo sapiens epsin 1 (EPN1):
cctccttctgtttgcttcccgtctcctcggcggtccccctccccgcccggctctccgcgccccttc
tgggcgggcgggcgggcgagccgtcggcggtgcggccctccttgcttcgtgctgctgcccgtggcc
cggcgcacgtcccgcgacaccgaggccgagcggggaggggctgaccgccatgacccccagagc
45 ccggcgtgagggggccgagatgcggtgacctgccagcacctgccgcagccttcgtccgggagtcgc
cccatctctccacgcacggggccctgtgcccttgctgctgcagccgggacccatg
(Seq ID No: 592)

- 40 Homo sapiens osteoglycin (OGN):
catcctctaagcttttaaatattgcttcgatggctcgaatttttatttccagggaaaaagagagtt
ttgtcccacagtcagcaggccactagtttattaacttccagtcaccttgatttttgctaaaatg
(Seq ID No: 602)

Homo sapiens NIN1/RPN12 binding protein 1 homolog (S. cerevisiae) (NOB1): gctcccctctcacgcagccaacatg (Seq ID No: 603)

Homo sapiens nudix (nucleo-
side diphosphate linked moiety X)-type motif 5 (NUDT5):
5 catccttttagcaccgcgagaggcgccggtgtttcgagccgtggcaccggcatcggctgacactgc
tgctccagctagttatttcgtcctcttccggttcttcacccctacaccttgagggtgaacttctca
cctgagggctgtaaagactcgtttgaaaatg (Seq ID No: 604)

Homo sapiens WD repeat domain 91 (WDR91): cgtccctcaccgcac-
caccctaagacgctagcgctgcgatg (Seq ID No: 605)

10 Homo sapiens nuclear transcription factor Y, gamma (NFYC):
gggctctgcattgcccgactccgtaggagcgcgggggcggtcctgctcttctcctggactcctgag
cagagttgtcgagatg (Seq ID No: 606)

Homo sapiens protein phosphatase 2, regulatory subunit A, alpha (PPP2R1A):
15 ccgcccttccttcttctcccagcattgccccccacgtttcagcacagcgctggccgcagctctga
caggaaagggacggagccaagatg (Seq ID No: 607)

Homo sapiens vesicle-associated membrane protein 2 (synapto-
brevin 2) (VAMP2):
20 ccatctttccgtcccgggcagccagcgccagtcggagccagcgcgagccgcccgcgccatcactgc
cgctgccaaagtctccaccgcgtgcccccgccatg (Seq ID No: 608)

Homo sapiens transmembrane protein 5 (TMEM5):
gattctctttccgcccgcgtccatggcggtggatgcctgactggaagcccagtgaggatg
(Seq ID No: 609)

25 Homo sapiens UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltrans-
ferase 3 (B3GNT3):
aactctttcttcggctcgcgagctgagaggagcaggttagaggggcagaggcgggactgtcgtctgg
ggagccgcccaggaggtcctcaggccgacccagaccctggctggccaggatg
(Seq ID No: 610)

30 Homo sapiens SEC11 homolog A (S. cerevisiae) (SEC11A):
gcgccctttccctgcccgggtgtcctgctcgccgtccccgccatg (Seq ID No: 611)

Homo sapiens RUN and SH3 domain containing 1 (RUSC1):
ctccctccccgcgcccgcgtcctctcccgccctacaggccctagcagggcaggcgagggtgagcgc
ggccatccccgcgtcccggagttccgggatcctggagtcgcgtagttcgtggtccttcgcccgggtgtccc
cgagagcccagcggctgtggatg (Seq ID No: 612)

35 Homo sapiens aryl hydrocarbon receptor interacting protein-like 1 (AIPL1): cctccctttctcctgcagccatg (Seq ID No: 613)

Homo sapiens tumor necrosis factor, alpha-induced protein 8 (TNFAIP8):
40 cctccttttctccccgcgggtctaaccgcgcttggttaaggtccgcgggaaccggtgagccaccg
agagagcagagaactcggcgccgccaacagcccagctcgcgcttcagcgtcccggcgccgtcgcg
ccactcctccgatg (Seq ID No: 614)

Homo sapiens staphylococcal nuclease and tudor domain containing 1 (SND1):

gcgtctcttttcgctccgtgtcccgtgctgctcctgtgagcgccccggcgagtcctgctccggtccacc
gtccgcagctggttagccagcctgccctcgctcgaactccctttcaccaacaccgacacccacatt
gacacctccagtcgggccagccgctccactcgttgccctttgcatctccacacatg
(Seq ID No: 615)

- 5 Homo sapiens DNA segment on chromosome 4
(unique) 234 expressed sequence (D4S234E):

cgccctcttttggtcgccccctccccaaccagcactaaggagcaccctgctctggtctccgccac
caccagcgctccttggaacctcccccaaaccttgaacgtcctcaggacccccaggtagcgcg
ggcgcgctgcgggcggggacctctctgcacctccccgcacccctgggggtcgctctgtccctacg
10 gtccccgcctccccctttctcctttctaagcgctcgcgcccaggcgccgccccggggtggcgcgagc
ccgcagccctcccgctccgggccccctccgcgctccgagacccccctgggggcgcgctcctctcccg
ctccccgtttccctcccccggtcagggcgggcgcggtgggtcccaggggaggctcccgcccagcccc
gcactcctttgtgcgggcgggcgggcgctgctcaagggtggaggcgcgggccacacgcgcgccacca
ccccgcgagcaccagcccccgaggagaggaaggagggcgggcgcgaggaggaggaggagggcg
15 cgtggagcccaatcgcttcgctcccccttcccggtccgcgcgcgggcgccgctccgccattgctgcg
agcaggagcaggagacgcggagctcggagcgctcagctgacctgccggagccgggctgggctgca
gcctcggagctcccggaacgatg (Seq ID No: 616)

Homo sapiens growth hormone inducible transmembrane protein
(GHITM):

20 acgtccttttcgatgttgctcatgcagtgcgcccggaggaactgtgctctttgaggccgacgctagg
ggcccggaaagggaactgcgaggcgaaggtagccggggaccgagcatttcagatctgctcggtaga
cctggtgcaccaccaccatg (Seq ID No: 617)

Homo sapiens stress-associated endoplasmic reticulum protein 1
(SERP1):

25 tttccttctcttttactccgcgctcacggcgggcgccaaagcgggcgggcgagggcgggcggagaac
gaccggcgggccagttctcttctcctcgccacctgccccgctcggtcagtcagtcggcgggccggc
gcccggcttggtgctcagacctcgcgcttgcgggcgcccaggccagcgccgtagctagcgtctggc
ctgagaacctcgggcgctccggcgggcggggcaccacgagccgagcctcgagcggtccagaggag
gcaggcgagtgagcgagtcggaggggtggccggggcaggtggtggcgccgccaagatg
30 (Seq ID No: 618)

Homo sapiens ADP-ribosylation factor interacting protein 1 (AR-
FIP1):

35 cggctctcctcacttccggcttcgctgctcttggttctggttctggaggctgggttgagaggtcgcc
ggtccgactgtcctcgggggttggtcagtgatgaatttgtagacagctgcagttgctccccgccccg
agcagccgaggagctaccatg (Seq ID No: 619)

Homo sapiens tumor necrosis factor receptor superfamily, member 21
(TNFRSF21):

40 ccgccccttcggcgccaccacgtgtgtccctgcgcccgggtggccaccgactcagtccttcgcccga
cagctctgggcagcgaggagggtggttggaagcttcgctatgggaagttgttcctt
tgctctctcgcgcccagtcctcctccttggttctcctcagccgctgtcggaggagagcaccggag
acgcgggctgcagtcgcggggcttctccccgctggcgggcgccgctgggcaggtgctgagc
gcccctagagcctccttgccgctcctcctctgcccggccgcagcagtgacatgggggtgttg
aggtagatgggctccgggccccgggaggcgggcggtggatgcggcgctgggcagaagcagcccgcat
tccagctgccccgcgccccggggcgccccctgcgagtcgccgggttcagccatg
45 (Seq ID No: 620)

Homo sapiens sushi-repeat containing protein, X-linked 2 (SRPX2):

ccccctcttctgcagcagacggactgagttcctctaataccctgtgttccttctcccccatctttct

aaaacccttctctgagagaggaataactatagcttcagggataatatagctttaaggaaacttttg
gcagatgtggacgtcgtaacatctgggcagtggttaacagaatcccggaggccgggacagaccagga
gccactcgttctaggaatgttaaagtagaagggttttttccaattgatgagaggagcagagaggaag
gagaaagaggaggagagagaaaaagggcacaaaataccataaaacagatcccatatttctgcttcc
5 cctcactttttagaagttaattgatggctgacttctgaaagtcactttcctttgacctggtacttca
ggccatatacatcttttcttgtctccataatcctccctttcaaggatg (Seq ID No: 621)

Homo sapiens HIV-1 Tat specific factor 1 (HTATSF1):
acctcccttttctctgctcagctccagcgtcatttcggcctcttagttcttctgaacctgctcctg
agctaggtaggaaacatg (Seq ID No: 622)

10 Homo sapiens trafficking protein particle complex 2 (TRAPPC2):
gggtctcttccgcggaaactgacattgcgtttccgttgctcggcctccactgcaggagccatata
tgaagaccatg (Seq ID No: 623)

Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acet
ylgalactosaminyltransferase 5 (GalNAc-T5) (GALNT5):

15 ccaccttttcttgggcttgttaggaaggtggacatgggctcccggagacaagacaagtgatatggtg
aactgttcgggtggctggaatcaactgctcctggagtgacctaaggccagtgtttatcagaacttag
ccagggccagccaagcaggcacagatgctctgctatgaaatgccacgcaggcagagactgacaagc
ggtaggaactgagctttccccttggactgctgcttccctgctgtgttcaggggaggggtcactttc
20 tggcaactctgctgctgctgctgctgctgctgctacttcagcttccctctccactcaaggttaagcag
gctaaggggagggcaggctgctagggaaagctttgtaccatg (Seq ID No: 624)

Homo sapiens transmembrane protein 97 (TMEM97): tggcccctcttctca-
catcagcgggtccaggcccaaccgacagactatg (Seq ID No: 625)

Homo sapiens EH-domain containing 2 (EHD2):

25 cgtcctccccgctccggggccccacccggctcagacggctccggacgggaccgagcacaggccgc
tccgcgggcgcttcggatcctcgcgggacccaccctctcccagcctgccagcccgtgcagccg
ccagcgcgccccgtcggcagctctccatctgcagctctctccgtgaaccccgtagcggtgtgcag
ccaccatg (Seq ID No: 626)

Homo sapiens tubulin tyrosine ligase-like family, member 4
(TTLL4):

30 cgccctcttcttccagactctcggtctgtccgctgggggcgcgcgcggtgtgtggcaggcggcagc
ggcgctggcgccgagtgcgcttgctacgcgtggcggtgcgtggttgctaggggcgctgaggctg
ccgggtagcccagcaggccgaggaggagtagcgtggagccggtgccgagccggggcgaagctgg
atcccctagatagactgtcttcaagctcactgatattttcctctgcttgatccattgtgctgttga
gagcctctagtaaatttttccagactgacagacttcaaggatgcagctgctactaccggagggtgtgt
35 ggcaccttacctcagcaaggccatgagaccgtgtggccatgatgtgggcccctcatg
(Seq ID No: 627)

Homo sapiens basic leucine zipper and W2 domains 1 (BZW1):

40 acctctccctcctcctggcggttagttccggctgcagaggagacaccgccgcagttgccggtacatc
ggggattttctggctctttcctcttcgccttaaattcgggtgtcttttatg
(Seq ID No: 628)

Homo sapiens centrosomal protein 57kDa (CEP57):

45 ttgcccttttctgtgtaagctgtgagcgtaggcgccctgagggggtgtgttgagggggtttccaag
cccagcaccagcacccttgcccttttccatcaggggttcagcctagggtccccgctggtggcgcc
tcccagctcttgagagaagagcacgagaacctagaccgccccgaagtgcggagaccccctgggcag
gctgaaagatg (Seq ID No: 629)

Homo sapiens family with sequence similarity 115, member A
(FAM115A):

ctgccctttgcctcctgggcggaagctgcttcctcctgggaacaaccgcctcccgctccttagca
ggttgctactgccccgaaccgcgctgcagggaacagcggggcaaacagtgagtggggttcagcgt
5 agactctggaccaggagaggcccggtgaccgaggcctgggccccggaaccaatagagccatg
(Seq ID No: 630)

Homo sapiens ATG13 autophagy related 13 homolog (S. cerevisiae)
(ATG13):

agccctctttcacccccccccccggccattaccgaagcgatgaaaacaaacactaacgatggcg
10 gcgccgggaagcgaccggctgctgggcttaaggcgggagtgaccgcttaaccagtgagggaagcac
tgaagagcgccagtcgacgtgggtgcgacaactcgcggagtcttaggagcaaaacgtctggggcct
gcgagccaggacccttctgaagccttaggtgtctatcggcgacgtgtacggctactgcagctccgg
agcgcggaaccctcagccaggaggcgcggctggctcgggtcccagggtcccggcctccgtaatgagagc
ccggaaccactctttgtgccgcagcttcgcagcatcttgactcaagtgattctcctgcctcagcc
15 tcctgagtagctgggactacagattcctataggcaatg (Seq ID No: 631)

Homo sapiens sorting nexin 17 (SNX17):

ccgccttccacatcgcatcgagggtcccaaatggcgagtgaggctgcggggactcgtgagc
agcggagggggagcgtgcagagccgctgcggccctcacagtcaggagcccgccgctgccgtgccgt
aggaacatg (Seq ID No: 632)

20 Homo sapiens phytanoyl-CoA 2-hydroxylase interacting protein (PHY-
HIP):

cgttcttttcccttctctgcctctctctcctccacgctgctttgatttcgctctttgcctctcttc
ttgcgctgctcagctgggaacatcgctctcaccaggggcagcagcgacgcgctgcacagccagacag
gagctggctgcggggcatggaagcagcctccttggcagccgggagaggagcaagcgcagccactg
25 cccgtgaccagcgctccggctgctgtccctgcccgggagctcatccacgcagaggtctctccct
gtcctccctgcgagcttttctctgcagagcccagtgaggccagtcctccacaggagacaaccctga
cgggagcatg (Seq ID No: 633)

Homo sapiens translocase of outer mitochondrial membrane 20 homolo
g (yeast) (TOMM20):

30 cggcctttctgtgttctcctggcccgccgctcgggtgtgagctgcgccgaccgctctgaggggttcg
tggcccaccgctccttcgcggtccctgccgccaccgtccacgctcagcgtttagagaagatg
(Seq ID No: 634)

Homo sapiens KIAA0141 (KIAA0141):

35 cggcctttctagcgcgtgtccaagggttggtctcgcgctttcggctgcgagctctctgtggtgct
ggcagcgacatg (Seq ID No: 635)

Homo sapiens janus kinase and microtubule interacting protein 2
(JAKMIP2):

ctccctcctttaaacagcttctccgggtctcagcatgggcttccagggcagcgattgaggagacct
taccaaggagcaccacacagtagatgctgagacatcgctactccaggataagaaacagtaacatggc
40 agcactgcttgaaagaaattaaaaaccaacagactccatttagaaaggaacaatg
(Seq ID No: 636)

Homo sapiens EPM2A (laforin) interacting protein 1 (EPM2AIP1):

cctcctctccccttgccgcctttctaacgttgccctgctcttggtggcctcccgcagaatg
(Seq ID No: 637)

45 Homo sapiens centrosomal protein 170kDa (CEP170):

cgttctttgccgttaccgctatgtgtggggcggtgtgtggaataacggtattgccagcggagctga

5 Homo sapiens suppressor of Ty 7 (S. cerevisiae)-like (SUPT7L):
aggcctctcgaggtccagacagccgcccagccgctctgcgacgcagcagtgaatagtgtggtacc
tccttgctcggttcaggtccagacctccccgtcttcgggctgccctgaacgtcaggcgacctcag
gacctgtgattggcgccctgcgccggcgaccgtgaccgaggaaacccctggagggacttgggcat
tccttgggctccgtgcctgttcttcgtgctcctttcgggcaaggatctcacattatcagtccttga
10 ccgacacagaatgcctggcatttgataaatgtttgttgaaacttgaagagacatatggacaatg
(Seq ID No: 639)

Homo sapiens non-SMC condensin I complex, subunit D2 (NCAPD2):
ttttccttttcatttcagcctgactgccggaatcagagccgcgggtgagatccccagccctgtgag
cctgtaggagtagaatg (Seq ID No: 640)

15 Homo sapiens ring finger protein 10 (RNF10):
 gggtctttgagatgctgtttggcgactcgtcgccattcccggagcaggtcggcctcggcccagggg
 cgagtatccgttgctgtgtcgagacactagtccccgacaccgagacagccagccctctcccctgc
 ctgcggcgaggagagcgtgtccggccggccggccggggctcgcgcaacctccctcgctcccc
 tcccccgagcctccgccccgcaggccccggccggactcccagagccccggcctcctcgtcctcg
 20 gtcgccgctgccgcccgggcttaacagccccgtccgccgttctcttcttagtttgagaagccaagg
 aaggaaacagggaataatgtcgccatgaaggccgagaaccgctgccgcccgcgacccccgcgggcc
 ctgaacgccatgagcctgggtccccgcgcgcccgtccgctccgactgccgtcgcgcgagggcc
 ccggttgatg (Seq ID No: 641)

25 Homo sapiens PAN2 poly(A) specific ribonuclease subunit homolog
(S. cerevisiae) (PAN2):
agcccttcttgattggaagaagcgctcggacccggctccttggcgccgtagtggttaggttgagc
cctaggcggtgggggagaactggggaaactggaatttcccgcggagctgacagcgcttgcgctcccc
ctactcgtttctaattccacgcgctccaaaatatccgccatggagaaatcttggccaggatgtccat
tctaggcccatcggtgctgtcttggctgaaggttgggtcaggcatctaaagggaactgtggtaaggga
30 gqgtgtgacacaggtgtaagctgccatcgctcatcatg (Seq ID No: 642)

Homo sapiens CD302 molecule (CD302):
gctcctctccggccgcgagccgctgccgccacccgcacccgccgtcatg
(Seq ID No: 643)

35 Homo sapiens NSA2 ribosome biogenesis homolog (S. cerevisiae)
(NSA2):
gactcttttctgtcccggcctgcggtggtgtgggcttgtgggtctttgagacccgaaaattgagagc
gttttcgcactccagcggctgctcctggcggctctgcggccgtcaccatg
(Seq ID No: 644)

Homo sapiens DIS3 mitotic control homolog (S. cerevisiae) (DIS3):
40 acgccttttgctggaagagcgctgctggggtaggattctgcgcggcgaggcaagatg
(Seq ID No: 645)

Homo sapiens caspase recruitment domain family, member 8 (CARD8):
cctcctctgcgagcggttattttcaaaagaagttgagaaccagagaaaccgacctaaggggattctcc
catttgcccgctcctaccctaagtcaccacctgctgcttttctggagcgcttaccagtgaaccaag
45 aggaacagaacacagagcagcctggcagtggtccaagcaacaagcctccgctcctccttctctgcacc

ctggggctcctgaaactcacatgggtaaaaaagatacagtaaagacataaataccacatttgacaa
atg (Seq ID No: 646)

Homo sapiens epsin 2 (EPN2):

5 ccgcctctcgagcgctgccggtggccgcagcggcgccccacgcccggaggagcagagtgtt
catttctgtgtcgggcacagtgttaagtgtctgggtgctcactggtgatgaggcagatgaaggttac
caaacttgtggacaggagcctcatatcagagacgtggacctcactgtagcctgggtcatggcttcca
gcttttcgaatctgaggctccaaaggaggaaatgaccattcagggatcttactccagcttgattac
ggagactgaaccttcatagggtgcgcacttaccaaggacaggaagggtttctctgtttgaagggtt
taaacttataacaagaaaataaaaatg (Seq ID No: 647)

10 Homo sapiens pyridoxal-dependent decarboxylase domain containing 1
(PDXDC1):

ccgcctctcaaccatcagggttcggcagcccgcgccgcgcccctggcagctcctcctcttctccgccc
cgccggcgcgggcgcgggggacgtcagcgctgccagcggtggaaggagctgccccgcggggagga
ggaagtagagcccgggaccgcccaggccaccaccgcccgcctcagccatg (Seq ID No: 648)

15 Homo sapiens nicotinamide nucleotide adenyltransferase 2
(NMNAT2):

ccttcctttctccctctgcagacacaacgagacacaaaaagagaggcaacccttagaccaccgcga
aggaccatctgcaccatg (Seq ID No: 649)

Homo sapiens mitochondrial ribosomal protein S27 (MRPS27):

20 tgttccttttggtacgctccaagatg (Seq ID No: 650)

Homo sapiens leucine-rich repeats and calponin homology
(CH) domain containing 1 (LRCH1):

25 tcccctccttcagcgccctttcggtggagcactgcggcactcagccccagctgccgttttcccctc
gcggggaacgctgtgacccccccgcaggagcgggcggggtgggggggcccgggagaagatg
(Seq ID No: 651)

Homo sapiens PAS domain containing serine/threonine kinase (PASK):
gctcctttccgtggtgtgtagccggttggtgacccctgcctgatccagttgtagagttggaa
gcttggcagttggcctcccttcttcccatg (Seq ID No: 652)

30 Homo sapiens megalencephalic leukoencephalopathy with subcortical
cysts 1 (MLC1):

cttcctttcctagttgggttctgacagctccgaggcagtggtttacacaaccaacacgaaacattt
ctacgatccaccgattcctccctcattgatattcaggaagcagctctccttcccctgccttcag
ctcaagtttgctgagcttttgtttcatttgtaatacttcttgctggaagtcctcaccagagac
cagtgtcccaacggcagagcagcggggagataaagaactggtgacacgtggctgtacattcagc
35 acagctgtggtgtccccaagtgccatg (Seq ID No: 653)

Homo sapiens RRS1 ribosome biogenesis regulator homolog
(S. cerevisiae) (RRS1):

40 ctttcttttccgattgggcatcccgcatctgcacgtggttatgctgccggagtttgggccgcca
ctgtaggaaaagtaacttcagctgcagcccaaagcgagtgagccgagccggagccatg
(Seq ID No: 654)

Homo sapiens formin binding protein 4 (FNBP4):

cgctctctgctcgcgcttgggctcgcatg (Seq ID No: 655)

Homo sapiens peptidylprolyl isomerase domain and WD repeat contain
ing 1 (PPWD1): gcgccttttctgacgatgcgaacaacatg (Seq ID No: 656)

Homo sapiens sorting and assembly machinery component 50 homolog (S. cerevisiae) (SAMM50):
ccgccttctgcccctcagcagcagacgctctgtcccgcccgggcagctctgagggcagcggtgga
gaggaaccatg (Seq ID No: 657)

- 5 Homo sapiens Yip1 domain family, member 3 (YIPF3):
gcttctcctttttgtgttccggccgatccacctctcctcgaccctggacgtctaccttccggagg
cccacatcttgcccactccgcgcgccccgtagcgcggtttcagcgacgggagccctcaaggga
atg (Seq ID No: 658)

- 10 Homo sapiens tectonin beta-propeller repeat containing 1 (TECPRI):
caccctcttggccgggtccccgggagggccgggtccgctcctcccgacgccgaggacctaccaccgc
gacttcgccccgccccggcgccggccaggacctgatgtcgcttttgaacagccccctgcacctggc
agccagcgagctactgtagtaggcattgccgactgtttgcataccggatgggagtgacagtgtaat
agaaaaacaagcaagaaaccttttaggtaggactcctaaggctcagaggaagttacctccagccgc
tgccatg (Seq ID No: 659)

- 15 Homo sapiens DDB1 and CUL4 associated factor 12 (DCAF12):
ccttccctttcccggtcaagtccttctctcttcttcttccgcctatctttttctgct
gccgctccgggtccgggccattttccggggccggcgccactaagggtgcgcggccccggggccagta
tatgaccgcgcgtctgtatctcttccgcttcccccgccccatgtggctgcggggccgcggcgccgc
tgccactatg (Seq ID No: 660)

- 20 Homo sapiens chromosome 3 open reading frame 17 (C3orf17):
ccgcctttcgtaagtccccccgcctcgcatg (Seq ID No: 661)

Homo sapiens LETM1 domain containing 1 (LETMD1):
caacctcttctctcccgcttctctcgctgtgaagatg (Seq ID No: 662)

- 25 Homo sapiens chordin-like 2 (CHRD2):
ctcccttctgctggaccttcccttctctccatctctccctcctttccccgcgttctctttccac
ctttctcttcttccaccttagacctcccttctctgcccctcctttcctgcccaccgctgcttccctgg
cccttctccgaccccgctctagcagcagacctcctggggtctgtgggttgatctgtggccccctgtg
cctccgtgtccttttctctcccttctcccgactccgctcccgaccagcgccctgacctgggg
aaaggatg (Seq ID No: 663)

- 30 Homo sapiens CCR4-NOT transcription complex, subunit 10 (CNOT10):
actcctctagccggaacctgggggccccggagccgggtaggcacagagttgtcctcgagggtccag
gacagcgccagcccgccggcgaggatcaggggccacgccacctgcagggaagaacccgagtcgaag
cggaagatg (Seq ID No: 664)

- 35 Homo sapiens THUMP domain containing 3 (THUMP3):
cttcctcttgagttgagggccggcgccgagccggacttcaggcggtctctgtggcgagcccatct
tgctccctctccaggccctttaccgctccctaggattccggggccctgtaggtgggagttgggag
acgacagtactgcttttaagagacagtgttagggatcttggaagcacagccaacatg
(Seq ID No: 665)

- 40 Homo sapiens nipsnap homolog 3A (C. elegans) (NIPSNAP3A):
gctcctttccactcgggaaaccttcagaggagtctcagaaaggacacggctggctgcttttctcag
cgccgaagccgcgcatg (Seq ID No: 666)

Homo sapiens CAP-GLY domain containing linker protein 3 (CLIP3):
gccctcctctccgccccccccctgtcggcgtctgggctcgtcccttctctctgtctccct
tgctcccccatcacgtcccctgacaccgacacccattgctcccacagtctccccagtctccact

ttgggtccccagcgctgtctgcccaggatttgcctgaaggctgcccccaactctgcacccgcccc
cgagggccaccgaggaccatg (Seq ID No: 667)

Homo sapiens ring finger protein 167 (RNF167):

5 cacccttcccgaagtttttctgtcacctgtgttaggctccgtcccctttccgcgttttatccccgt
accagaaaaggatacatttagtgcctcccaccagctccactaaacgggttgatatctcattctt
tgagttggtgttcttccccggcgcccccatgtagctgggaagtgggacctgggggtggttgacc
cctgggatacctaaaggaggggcagggagggcgagaaactccgcttctgctccttgctaccaggacg
cgcggcctcctcagcctctttcctcccgtgccatg (Seq ID No: 668)

Homo sapiens polymerase (RNA) II (DNA directed) polypeptide M
(POLR2M):

10 cgttcttccgggaaaatggcgactcccgtcgtgccccggagtcaccgccgtccgcggatccggcg
ctagtagcggggcctgccgaggaagccgagtgcccgccgcgcgcagcctcagcccgcgcagaat
g (Seq ID No: 669)

Homo sapiens dihydroxyacetone kinase 2 homolog (S. cerevisiae)
(DAK):

15 tcgcctctttccgccagcgcccgaggaccggatgagagcgcacgcttcgggggtctccgggaagt
cgcggcgccttcggatgtggcgatgcggccgtgagccggcgggggaggtgctgctgctgcctcca
ctgtactcagaccaggtagcacaggattgtccatcctccagcagctcagtgaacggtgtgaact
cagcctgtttcagagcctccacaccatg (Seq ID No: 670)

Homo sapiens RNA polymerase II associated protein 1 (RPAP1):

20 cgatctctgcggggcaagatggcgccgcccagacaggcctggagcacggatgaataagaggggaacc
cccacacggagacactgctggagagagtcgtactggggaggcagctggagcagcaagatg
(Seq ID No: 671)

Homo sapiens torsin A interacting protein 1 (TOR1AIP1):

25 cctcctcttttggtgcctccagccaggaggcgggagcgatccacagcagctgaccagctcaggcac
tgctctctcacagccctcaagacacaccatgggcccagaggcaggtttgctacacagcagcgacg
acgcaggcggcgcccccagcgactcgcaactgcctccctgaccacagcggccaccgcccacaccc
ccgagaagccatcgccaccaccggcaggagaacctaggggtccataaagccatcttcgcgatcgact
aaagctacgtcaacaactatg (Seq ID No: 672)

Homo sapiens SERPINE1 mRNA binding protein 1 (SERBP1):

30 cccctctctcggcccgccatcttgtgggaagagctgaagcaggcgctcttggtcggcgcgggcc
cgctgcaatccgtggaggaacgcgcgcgcgagccaccatcatg (Seq ID No: 673)

Homo sapiens N-acetyltransferase 9 (GCN5-related, putative)
(NAT9): caccctttctgcgggggacgatttcgtcggtaggtgctaccatg

35 (Seq ID No: 674)

Homo sapiens ribosomal L1 domain containing 1 (RSL1D1):

gcgcctcttcacgaggtggaaacaagatg (Seq ID No: 675)

Homo sapiens SH3 domain containing, Ysc84-like 1 (S. cerevisiae)
(SH3YL1): cttcctcttctgggcagcctcgggacggggcgccgcggccggggcgggcagcatg

40 (Seq ID No: 676)

Homo sapiens methylmalonic aciduria (cobala-
min deficiency) cb1D type, with homocystinuria (MMADHC):

acttcctttgcctgctcaccgccagcgtaggtgctaccaccgctgccgtcgccgcgcgcatctttga

tggcaggaagagtccggttctgggacagctggagacagtggtggtgactgaaataactttaccaa
ggaaagctattttgcgaactatcttctccagcggagatg (Seq ID No: 677)

Homo sapiens glioma tumor suppressor candidate region gene 2
(GLTSCR2): agttcttcctttgacaagatg (Seq ID No: 678)

5 Homo sapiens DDB1 and CUL4 associated factor 8 (DCAF8):
cagtcttctcagcacatcgctcgaaacggggccggaagcgtggcagcgcaggcgcaagcgcaga
gagcggaggcggtggtggtggcgccgctggccagttccttcagtgaatctacagacctatcttct
caggagctcagcctggccttacttcagtgataaaaggaggaaaggctggctacagcaaaccattc
caagatg (Seq ID No: 679)

10 Homo sapiens UBX domain protein 1 (UBXN1):
ctttcttctcgtcgggtgttcccggctgctatagagccgggtgagagagcgcagcgcggcggtg
tgtcgagggcggttgctcgcgctgacccttcccgccctccttctcgtcacacaccagggtccccg
cggaagccgcggtgtcggcgccatg (Seq ID No: 680)

Homo sapiens antizyme inhibitor 1 (AZIN1):
15 ccgccttctcacactttcaggctctgatcgcgccgcagtttttcttttcttctgccgtcgcc
ttctctgcctcttctcatccttctcgtctgctgctctgcagtgtgacgagtcgaatcctcttc
ccaccagcccgcgcccttcttcttttgctgctggttctatttctccttcggccgcggcgcca
ctgctgcacacagctggtgtcgggtgccgcgcttttaccaccaagtcggtcccgagcctatggccc
aggccgccttgggtatttctgctcaaggtaaccacatccctctttaaaaattccgcccgaagag
20 aagacgctttaccgactcttggggcggttatctcacggcgaactttctgaccaagtatacacta
cccagagggcctaggagaagtgtgtatagagagcagttcgacttcaacgctgagccaccttggga
acctagctgatgataggggggttccatctcccaacttgccatggagggtcttcacttcagaaatcc
aagactcataattcatccagcttgggtgtcaagtgggctgttgctgccagaattatcttgtgattatt
tgagagatgtatcagtttcttctgaagtacaatcaactgtagaagcctttagtagcagtttgttgca
25 tattctaaggaccagacataggcttgggtggccgctctcttgtctttcctgggttatgactttcgg
ctttgtggaatacggctgagatg (Seq ID No: 681)

Homo sapiens cell division cycle 40 homolog (S. cerevisiae)
(CDC40): gcctcttcttcttccgcctggcagggctcgcgcagaagatttgttgccgtcatg
(Seq ID No: 682)

30 Homo sapiens stathmin-like 3 (STMN3):
gcgctctccagcctccgcaggcccaaccgcccagcaccatg (Seq ID No: 683)

Homo sapiens nudix (nucleo-
side diphosphate linked moiety X)-type motif 13 (NUDT13):
35 tttctcttttgtgctgattcctgaggactaggaaggtgccccgaaaagaattcagagacctgaca
atg (Seq ID No: 684)

Homo sapiens calcium homeostasis modulator 2 (CALHM2):
ctctcttttctggagtttagattagtctgaagccgccaccagccccaggccccgtgcagaagaaaa
gcgaggagggaacggcgaggccgcccgtgccctgcaccgccctcctggaggccacttgagaggtcc
ggccccgaggaggccatggccacaagtgccacagctggccccagggttgccagcgtcgctacagcc
40 cagaccaaggcagaataatctccgatgagctggtggcaccgctgagccttgggtctcaccagggc
ttcctgttgctggcaggcggggtggagcggagctgctgggaggctgctggataggagaggggtcac
ggctgcggaagaggaggttcttcgggacaccgctggatggacacggcaaggaaacaccaggccaac
cacagctggggataaaaatagcacaaccacaccctgccgtccagcgcctccagcctgtgcccttc
ctagtaccaccagcaaccatcaatcccgtctcctcctgcctcctctcctgcaatccaccccgccac
45 gactatcgccatg (Seq ID No: 685)

Homo sapiens NMD3 homolog (S. cerevisiae) (NMD3):

tcttctctgtggcggagacagccaggttggcagctgacgggacagccgggtctatatttgttgcg
gttttcagcaaattccagggtggtctggaggcgcaaaacttaaggcatacagaacgatg
(Seq ID No: 686)

- 5 Homo sapiens ATPase, H⁺ transporting, lysosomal 50/57kDa, V1 subunit H (ATP6V1H):

gcgcctctgtcattctactgcgccgcccctggcttccttctacctgtgcgccctcaacgtctcct
tggtgcgggacccgcttcactttcggctcccggagtctccctccactgctcagacctctggacctg
acaggagacgcctacttggtctctgacgcggcgccccagccggctgtgtccccggcgccccggacc
10 acctccctgccggctttgggtgcgttggtggggtcccgaggattcgcgagatttggtgaaagacat
tcaagattacgaagtttagatg (Seq ID No: 687)

Homo sapiens DPH5 homolog (S. cerevisiae) (DPH5):

gggccttttctctgcacggagccggcgcttttgcagttgcttctgcggaaaggtggtagttaagaa
tttgtaaaggccagagaactacctacgattctctcagcggtctctcttctcctcaagtttgaaatg
15 (Seq ID No: 688)

Homo sapiens polymerase (RNA) I polypeptide D, 16kDa (POLR1D):

cctcctccctccttcctcctcgcgccttcctcctcggtcggtccttgcttctctgcttcgcctccgc
gcctcgcgctatgggacagagccccgatccgccagcaccacctgaggatccagaaaccgccccag
cgatg (Seq ID No: 689)

- 20 Homo sapiens HMP19 protein (HMP19):

ctgtcctttcagcaccacaagctcgggctgaggagggaggactcctggccgtcctcctcctcttca
aattggcttgaaatcttctctgacccccacgagtgacagcacagtctgggaagaaaggcgtaaggat
g (Seq ID No: 690)

Homo sapiens adiponectin receptor 1 (ADIPOR1):

25 gcgcccccttcggcgcggggagggcgctgaagatcggggcgctcggccgcaggccgcctccagcg
ccgcgggatgtagcgcggggacccgcgccccagcagagcccgctgcccggttgcttaccatc
agaggagatctctgccccctggggctgagagacccaacctttccccaagctgaagctgcagggt
attgaggtaccagccagatg (Seq ID No: 691)

Homo sapiens SH3-domain GRB2-like endophilin B1 (SH3GLB1):

30 ttttcccttgggaccgggtccacacggcggggtcgccgtccatctccggctcgcccgcggggccc
catcgctgacgttagcgggcgttctccgagccgactgacctatccttggcgctgcgcgcgcgct
tgttctcctccctcgccccgccttcctcctcccggttcacggaaacgacagctgcgggtgcggggc
tggcgccgcctccctccacctaccacgtctgcctcgccgctctagccctgcgccccagcccgcc
gcggcacctccgctcgccgcccgtaggtcgggcggtccgccccggtgcgcctaggatg
35 (Seq ID No: 692)

Homo sapiens anterior pharynx defective 1 homolog A (C. elegans) (APH1A):

gtccccctcttcggcttccgtagaggaagtggcgcgggaccttcatttgggggtttcggttccccccct
tccccctccccggggtctgggggtgacattgcaccgcgccccctcgtaggggtcgcggttgccacccca
40 cgcggaactccccagctggcgcgccccctccatttgctgtcctgggtcaggccccccaccccccttcc
cacctgaccagccatg (Seq ID No: 693)

Homo sapiens RNA binding motif protein, X-linked 2 (RBMX2):

ctgcctttcccgggcgctgattcctgagtgctgagcgcgaaaccgaggagatg
(Seq ID No: 694)

Homo sapiens family with sequence similarity 82, member B (FAM82B):

atctccttttagccccgccccgctccgtagctgcctgaagtagtgcagggtcagccccgaagttgca
ggtcattg (Seq ID No: 695)

- 5 Homo sapiens UTP11-like, U3 small nucleolar ribonucleoprotein, (yeast) (UTP11L): tgatcttttccaaggctgtacagacatg (Seq ID No: 696)

Homo sapiens chromosome 14 open reading frame 166 (C14orf166):
cgccctctcgccgctcgccggtgctgcgctcccgctccacctcgcttcttctctcccgccga
ggccgggggaccagagcgagaagcggggaccatg (Seq ID No: 697)

- 10 Homo sapiens transmembrane emp24 protein transport domain containi
ng 5 (TMED5):
gcttctcttttcggaggagtggttcgcccgcgcccgcggccgcccacctggagtttcttcagactccag
atttcctgtcaaccacgaggagtcagagaggaaacgcggagcggagacaacagtacctgacgcc
tctttcagccccgggatcgccccagcagggatg (Seq ID No: 698)

- 15 Homo sapiens coatomer protein complex, subunit zeta 1 (COPZ1):
gtttcttttgccggtccacgtcggcaccagctgcggggcaagat (Seq ID No: 699)

Homo sapiens mitochondrial ribosomal protein S16 (MRPS16):
ggttctttctgtgtttgttctctgccttgccaaggccgtagagctggtgctgctgggtagcggggc
tctccgaggagccgcacgcccgcggcaccatg (Seq ID No: 700)

- 20 Homo sapiens charged multivesicular body protein 3 (CHMP3):
ctacctccttttcgccccgccccgagcggtgcccgtgacctgctggcgccgggggaactga
aagccggaaggggcaagacgggttcagttcgtcatggggctggttggaagaccaggagaagccg
cccaaagaactgatatccaaagagaagaagaaaaagtgaacgatctgtgaaagatgctgccaaga
agggccagaaggatgtctgcatagtcttgccaaggagatg (Seq ID No: 701)

- 25 Homo sapiens RNA binding motif protein 7 (RBM7):
cgaccttttgccaggttagggagggggcgacgctgagatg (Seq ID No: 702)

Homo sapiens eukaryotic translation initiation factor 3, subunit L (EIF3L): cgctctttccggcggtgctcgcaagcgaggcagccatg (Seq ID No: 703)

- 30 Homo sapiens zinc finger protein 706 (ZNF706):
ccttcctttccctccggcgtcctctcccgccctctcgcgctgcactgtctctccgacgcaagact
gtccccggccccgatatg (Seq ID No: 704)

Homo sapiens androgen-induced 1 (AIG1):
cgccctccttgccgcccagccggtccaggcctctggcgaacatg (Seq ID No: 705)

- 35 Homo sapiens interleukin-1 receptor-associated kinase 4 (IRAK4):
cgcccttcgcgcgcttcttagttcggtggttcttctgtcgccggcttcagcagccccgcgccc
ggcaggaatagaagatg (Seq ID No: 706)

- 40 Homo sapiens transmembrane protein 66 (TMEM66):
cgttccttcgcccgcgcccagggtagcgggtgtagctgcgcagcgtcgcgcgctaccgcacccag
gttcggcccgtaggcgtctggcagccccgcgccatcttcatcgagcgccatg
(Seq ID No: 707)

Homo sapiens carboxypeptidase Q (CPQ):
ccgcctctcgccccgcggcctggccggcaagcagggtgcagtcacggggcgcgcgaggggccc

cagcccagtcaggggtgtggccgcccaccgtaaggctaggccgcgagcttagtcctgggagccg
cctccgtcgccgccgtcagagccgccctatcagattatcttaacaagaaaaccaactggaaaaaa
aatg (Seq ID No: 708)

5 Homo sapiens hydroxysteroid (17-beta) dehydrogenase 12 (HSD17B12):
cgctcttttctattcacgaaggtagtgaggcctagtggaaagccatg (Seq ID No: 709)

Homo sapiens protein phosphatase methylesterase 1 (PPME1):
cctcccctcgatg (Seq ID No: 710)

Homo sapiens HemK methyltransferase family member 1 (HEMK1):
 10 cccctttccggcaggctactgggctccgcccacacacctcccggtggttcctaaacgccagct
 cggagcaatccccttgggctggagccaaatccctgctgtgattttaaggaagaccggcagggtccgg
 gccccaaagggtcaacccacacacatccccgcacttttctgtatgcaggcctgcgagcgtagagg
 gagtggaattcacagcctccccacccatccgcagggggtctcctgggaggaaccaccagcgatagg
 aacactgaagctgggctacggcgtccgcccagccttttcttaaaggcgccgaccccggaagcggg
 15 gcgtccgagggagcgcgcgacgggccacgcacgtccgggcgtccagttcggggcagcttctccggc
 tgggtgggtgggtggggcagcctttcaggcagggtggcaaccaactatatctgaggaccagagccat
 tttggggcaccagagcttgtgacctctccatctccaccagctgggtccagggggccactctcagca
 ctcacctcagcagctgacatcataaagcagacttgggaaacctggaagcactctggagaacctttcc
 ctgagacatg (Seq ID No: 711)

20 Homo sapiens N(alpha)-acetyltransferase 38, NatC auxiliary subunit
(NAA38):
cgccctttcagttctgcttgctgctcggcaccgctgcgttaccgcgaaccgccgggccgaacagcat
g (Seq ID No: 712)

Homo sapiens cleavage and polyadenylation specific factor 3, 73kDa (CPSF3):

25 ggttcttcccttttttattttaccgggtggctgtgcttccaatttaggaagaccccggcgacctgttcc
tcacccccgccttcgccctcacacttttcgggatg (Seq ID No: 713)

Homo sapiens dynactin 4 (p62) (DCTN4): tcgcctcctccctccccaagatg
(Seq ID No: 714)

30 Homo sapiens hydroxysteroid (17-beta) dehydrogenase 11 (HSD17B11):
gttcctccttgctctcgccctactctttctgggtgtagatcgagctaccctctaaaagcagttta
gagtggtaaaaaaaaaaaaaaacacaccaaacgctcgagccacaaaagggatg
(Seq ID No: 715)

Homo sapiens YTH domain family, member 2 (YTHDF2):

35 tagtctttccaggtgttagtcgaaacctcgtaggtgcgacctggtcgccccaaacccctaggcct
taatcctggggcggtggggcgaggagccgtgagcacggcttccgctcctccaatccgccagagg
gcgcagcgccggcctctcccttccgggggttcttcgcgcgcggcccttccgcgtgggtgagtga
atgtgagagtcagcgctcgcgccgcgcgcgcgcgcgcctccgctgttcggcgctctgctttaggc
ggtggggggcgggcgcgcgcgtaaaagcatagagacgggcattgagctcttgggctagagcgctgc
cgagtcggagccggagcctgagccgcgcgctgtgtctccgctgcgtccgccgagggcccccgagtgt
40 cagggacaaaagcctccgcctgctccgcgagccggggctcatctgccgcgcgcgcgcgctgagga
gagttccgcccgctcgccgcccgtgaggaatctgagagccatg (Seq ID No: 716)

Homo sapiens tubulin, epsilon 1 (TUBE1):
agctctctagcagagcgccgttgctgggggaatgcagaagcgggcgcgggctagcaagctcccgga
gccgcgcgcaccaccatg (Seq ID No: 717)

Homo sapiens ubiquitin interaction motif containing 1 (UIMC1):
cctccttttcttctcagcgggtccgcggcccgctactctccgggagggcgcttcccagcgccaa
ggtaggcctctcccagcgcggggcgcccttctgatgccggggtgtgtctctcgcgacgcgggg
gtgggctccggacgcggggctggccttgccgaagtcgggggtgggtccctccggacgcggaagt
5 ggctcgggatgcggggctgggacctcccgaattccggggcggaattccggacgcggggaccggccat
tactggtgccgggttgggcttctccagatgccggggctgggtccttcccaaggttgagacaaaagg
atg (Seq ID No: 718)

Homo sapiens TNF receptor-associated protein 1 (TRAP1):
ccgcccccttcccatcggtgtacgggtcccgcgtggctgcgcgcggcgctctgggagtagcacatg
10 (Seq ID No: 719)

Homo sapiens cereblon (CRBN): cagcctcctttgcgggtaaacagacatg
(Seq ID No: 720)

Homo sapiens ribosomal L24 domain containing 1 (RSL24D1):
cttcctctcaagcttggcggttgggtggggttacacgcgggttcaacatg
15 (Seq ID No: 721)

Homo sapiens leucine carboxyl methyltransferase 1 (LCMT1):
taccctcttctgttgccttctccctgtggctcgcgcggtcccccgcggcccgctcgaccccgcttcc
atgtccctggcgacacagctcccaggaacctccacgcccattggccactaggcagagggaatcctc
tatcacctcctgctgttccacctcgagctgcgacgcagacgacgaggcgctgcgcggcacctgcga
20 agatg (Seq ID No: 722)

Homo sapiens RAB14, member RAS oncogene family (RAB14):
cccccttcttttgtggtccggccattgacgagggtgacaggaaacctgtgcaggagcgccgcca
tcttgaccagcccaggaagatactgaggagacagagcagtcaccgctgccactgctactgc
cgctactgctgcggcgcgctctgcacctctcggcctgccagtgtacctgcggcgccctcggtcgac
25 cgccccgcggccctctcccgctgcgtccgcactcctgttccctgggtcctgacgccccctcccgccc
ggaaagctgccagccaccagcaacccccagtgccaccatg (Seq ID No: 723)

Homo sapiens Enah/Vasp-like (EVL):
cttccttttctgttttgggttttaagtaggctataaaaaatcaagttgctgtcttccagagggtctgtg
gtcctctgatcaacataggctggtgggagtagcaggactcgctcctcagggttccctgtgctgcc
30 cttttcagccatg (Seq ID No: 724)

Homo sapiens LIM domain and actin binding 1 (LIMA1):
ctctcttccctctccctctccctctgcgggtggatgctttctccatgtggcaaggctgtaactg
ttcacagctgtctgaaacagcagtgaccaggagcagcttgagttttaactttcattttacaaag
aacaacatgtttgaatgtttcagcaggcaagttataactggcatctacttctgttcttctagaac
35 accgaaaatctctccagcacttttagaaaggggaccctgactgtgttaaagaagaagtgggagaac
ccagggtgggagcagagtctcacacagactctctacggaacagcagcactgagattaggcacaga
gcagaccatcctcctgctgaagtgaagccacgctgcttctggagccaaagctgaccaagaagaa
caaatccacccagatctagactcagggtcacctcctgaagccctcgttcagggtcgatatccccac
atcaaggacggtgaggatcttaagaccactcaacagaaagtaaaaaaatg
40 (Seq ID No: 725)

Homo sapiens ubiquitin-fold modifier conjugating enzyme 1 (UFC1):
gtttctcttgcgccctgggtccaagatg (Seq ID No: 726)

Homo sapiens coatomer protein complex, subunit beta 1 (COPB1):
cacccttccacgtcagccaaggactctggagccgcccgcgctgctgcggttcatagccgga
45 gtagacggagccgcagtagacggatccgcggctgcaccaaaccactgccctcggagcctggtagt

gggccacaagccccaggtcccagaggcggtgggtcgggcagagtcggaagaactggctttctag
ctggaagatgcggaaggggagcgactaggccgcttgctgctgggcctggcagaagggaccggattt
tctggcatccttaaatcttgtgtcaaggattggttataataaccagaaaccatg
(Seq ID No: 727)

- 5 Homo sapiens transmembrane protein 9 (TMEM9):
gggtcttttgcggctgcagcgggctttaggtgtccggctttgctggccagcaagcctgataagc
atg (Seq ID No: 728)

- Homo sapiens shisa homolog 5 (Xenopus laevis) (SHISA5):
10 ctttctttttctccaaaaggggaggaaattgaaactgagtggccacgatgggaagaggggaagcc
caggggtacaggaggcctctgggtgaaggcagaggctaactg (Seq ID No: 729)

- Homo sapiens transmembrane protein 69 (TMEM69):
gtgcctttccagtggaacctgggctgttgttgcggttgttttcttctctccgtgcaacgctggcaa
gtctcaaagtcgccacagaaacatgccctgattcagtgccctctgcttagctgtaacatgttaac
agaactacctggcatcttctgaacaagactttcaataggggccagtatg
15 (Seq ID No: 730)

Homo sapiens kelch repeat and BTB (POZ) domain containing 4
(KBTBD4): agatcttcttccgggcgacgtggagccggaagcggaggttccgggctccgggatg
(Seq ID No: 731)

- Homo sapiens pipecolic acid oxidase (PIPOX):
20 cgtccttttagccgggagcctgtctttgcttgcccttgagggctctgtggctgtggggctga
gtggcatcatg (Seq ID No: 732)

- Homo sapiens blocked early in transport 1 homolog
(S. cerevisiae)-like (BET1L):
25 agctctttccccgcgactgcgccacgtctgaggcggctgtggccgcgtcggtgtccgcgtcgagga
gccggggcagggcacgatg (Seq ID No: 733)

Homo sapiens zinc finger protein 581 (ZNF581):
ttctctctttcggccggcgcccgccagttcctggggcacacccagaggtccccttctcgccgccc
tgcaactgcgagggtagcccgggcgcttgagtcgcccggacctgagaggtgctgcactgggg
ctcagccagccctccggatg (Seq ID No: 734)

- 30 Homo sapiens armadillo repeat containing, X-linked 1 (ARMCX1):
cgtccttctaatacttagtcttctggttccggttgcaactcttctatagcccagagggcgagagg
gcctgtggcctgggggaaggaggacgaggttctgcctggatcccagcagtaggacgctgtgccatt
tgggaacaaaggaatagctgcttggaatccctgcagatcttggggccggaggccagtcacaacct
tggagcaggaagaaacgcaaagttgtcaagaaccaagtcgagctgcctcagagccggcccgagta
35 gctgcagactccgcccgcgacgtgtgcgcgcttctctggggcagagcgagcctgttttgtgctcgg
gttaagagatttgtcccagctataccatg (Seq ID No: 735)

- Homo sapiens spastic paraplegia 21 (auto-
somal recessive, Mast syndrome) (SPG21):
40 cggcctcccgacgcaccgcgagcctgctgtgcccgtgggtcccagtgctccgcccggcccc
gaccggggcccagccgctccacggcccgcgctcgtagtgagcgaagagcggcctcctgaaggag
gggaaggagcgtgggggcccggccacggcaggattaacctccatttcagctaactcatg
(Seq ID No: 736)

Homo sapiens staufen, RNA binding protein, homolog 1 (Drosophila)
(STAU1):

tctcccttttttcttcttcttccctcctcgccgccaccgcccaggaccgcccggccgggggacg
agctcggagcagcagccagagtttattaaccacttaacctctcagaactgaacaaagacaacattg
ttcctggaacgccctcttttttaaaaaagaaagcataaccctactgtagaactaaatgcactgtgc
atg (Seq ID No: 737)

- 5 Homo sapiens adducin 2 (beta) (ADD2):
cggccttttgtcagcgcgcagggccaggagagctctcatttctcccagcctcgtgcgggaaatgg
ctttaattctgacggcagggctgtgagggactagcgggaaccgagccttttgtcaaggaactgcg
gcgctcgggtggccagtcacccccgccgcgcggagccgctgcactgctgggggatctcccagcagct
ctgacgagcgcgggctgcagcatgggcagaaaacgctgccctgcagattagctgggtggattttt
10 aagcgcacccccacccccaaaaccataaaaataacaaaaccaaccgcagtgccgcagccggagatag
ctaagatgccgcgcagaggtttccacctggatgtttgaggttgtgtagatgtggccggcacccttg
agagtggagctaggggtgcagactgagcagtgaaacagaaggagccttgacagggctgggccagc
ctcccagattccaggagcgaattgcaaaccaccgggaaaatg (Seq ID No: 738)

- Homo sapiens WD repeat domain 1 (WDR1):
15 ccgccttccggctccagtcctccgggctcggcctcggcgaggtgtaattcgcagcgcgggcccggccc
cggaggctctcggcgagcgcggcgcggttaacaagtgggcgaggatg (Seq ID No: 739)

- Homo sapiens family with sequence similarity 20, member A
(FAM20A):
20 cgacctctacttccacctctggccccaagtacagcgcagctgcggcctcgggagcgcgccgcgggg
gtgcccgtgcaccggccgcgcctcctccctggcgcgggactcggccgcagctgcctcggaccccg
cacgatcgtgcacaacttttcccgaaccgagccccggactgaaccggctggcggcagccacagcgg
gtcgagctccaagttgcaggccctcttcgccaccgcgtgtacaacgtcccggaggagccgcctct
cctgggagccgaggactcgctcctggccagccaggaggcgtgcggtattaccggagggaaggtggc
ccgctggaacaggcctcagttcctgcttttgaaaggaagagggggagctgtgtagccctgaggcct
25 ccttgcaactctgttttccaagctttgcacatcttccgaatttcttcttcaaagtctaccctaag
aaatatcagacaattttccaagtggtgcttcatgaacttctgggaggtgcttcacagtttctgcaaa
tgattgattgaattttcactttgaaaaataatactttaaggcgacacaagatg
(Seq ID No: 740)

- Homo sapiens kelch domain containing 4 (KLHDC4):
30 ttttctttcctggtgtcccgtcgcggcttgggaccggcaagatg (Seq ID No: 741)

Homo sapiens calcium channel flower domain containing 1 (CACFD1):
tgctccctctcccacaaggcagcgcgcggctcggacgcggccggctaccgagccctttgtgaggg
ctgtgagctgcgcctgacggtggcaccatg (Seq ID No: 742)

- Homo sapiens zinc finger, CCHC domain containing 8 (ZCCHC8):
35 gaatcttttccacagcccaaatg (Seq ID No: 743)

- Homo sapiens kelch-like 24 (Drosophila) (KLHL24):
gtttcctttgttgtgagctgcggcagagactggtggctggaggagacgccggcgctggagagtgcg
ctgcgcgcgcccgccgctgagggaccgcggggttagccactgctggctgcttccagtgttcgccgag
aggtaccgggggtgacagctccgggaccggccgaaaggcgaggaaccggtgtggaaattaaaagaa
40 cacacataattttgactggggctttgatcaaccaaagctaaaaagccacataaagaagatccctaa
tagtcatttctcaacaattatatagtcaactgatgtaacaatg (Seq ID No: 744)

Homo sapiens FtsJ homolog 3 (E. coli) (FTSJ3): ctccccctttccaccatg
(Seq ID No: 745)

- Homo sapiens dymeclin (DYM):
45 gcttccctcttctctcgcgcctcctggcctccgcaccgacgcggcccgggctggagccgagccgg

ggccgagctgcaggccggaccggagccggatctgtacccgctgagacgtggaaacatggaggcctg
agccggtgtgcgccacctgggctgcggcgccgacagcgacttctcctgacccctctgccaccctcc
catccgtccgcgggtccgtggagctggagcagatccccagccggctgagacaggttgtcttttg
aatgcaggtttaaggacaaattatctgcttaagctagaagatg (Seq ID No: 746)

- 5 Homo sapiens zinc finger protein 280D (ZNF280D):
cctcctcttttctcctcctcctcagggtccagtcaggccgatccgctccgctcacggaaggaaaac
agaaataacttgctggcttgctgagtcacatgtacttaggtgacaatttacagaaagtcacctc
tgcagcttgatg (Seq ID No: 747)

- 10 Homo sapiens ankyrin repeat domain 10 (ANKRD10):
cgttcctttgtgctgcggcgccggttctcagtcctccccgacgcgtcctctaggccagcgagcc
ccgcgctctccggtgacggaccatg (Seq ID No: 748)

- Homo sapiens SWT1 RNA endoribonuclease homolog (S. cerevisiae)
(SWT1):
15 ctctcctttggcttggggctccggagttgccactgccgccggcgctggttaagcttttcaggatg
(Seq ID No: 749)

Homo sapiens leucine rich repeat containing 49 (LRRC49):
tgacctctttcgggtctctttgaatctccgctgtagcgtcacctggaaggcagatctaacagagaa
cctggactgtctcctatcatg (Seq ID No: 750)

- 20 Homo sapiens F-box and leucine-rich repeat protein 12 (FBXL12):
ccgccttctggacttggctcttagttcccgatcgcgcccaaatcacgcctcagccacctcccgaag
cctctcactgcctcagccacgctttccaggctggtttctggtccccatccgcggctggtccggccc
tgggaccgaatcacttcccagcgagaggaaggtcaaatttctcgaccggctacgggaaggtcgcg
ccgcgcctgtcagccgcctcggcgccccaggaccctcggggtctctttaaccggaagcggaag
tgctgtcggcgggatcatg (Seq ID No: 751)

- 25 Homo sapiens WD repeat domain 55 (WDR55): cagtccttctcagcatg
(Seq ID No: 752)

- Homo sapiens zinc finger protein 3 (ZNF3):
30 cgttctttgttctgtccccgggtgtgtgggtctgtgacagggtccaacagggcctgggtccgtgtccg
gtcccccaaatctgtcgtccctgccccaggcattggcatcaacaaaagtcagaattcccgggaac
ttgaacagaggctgctaaattcccagtaattgctcctttggccttctagggtactgacttcaaagaa
ggaaggaaagaatcaggcagtgcttctcattctcttttaaaacccgcttcccgtgagtcgtgcac
ccaggagaccagagagcaccttgcccttccatg (Seq ID No: 753)

- Homo sapiens tetratricopeptide repeat domain 27 (TTC27):
35 ggttcttctcctaggcggaagccagaccagagagcgtgctgtttttcccagggtgccccgcgctg
ctgttatggccgctccttgaggtagtatccgcacatggaattctaggggccgcaggtgtatttacg
gtaactgtcgccactagatttcagcgcctttggactctcctgttttcttcttttggttgactcc
cgtgtggccctcgtgggagcctgttttggctgcagcggtgtctggggtgatg
(Seq ID No: 754)

- 40 Homo sapiens THUMP domain containing 1 (THUMP1):
gtttctctttcctctcagtttgccacacacatg (Seq ID No: 755)

Homo sapiens ankyrin repeat and KH domain containing 1 (ANKHD1):
tgctcttctcgttcccagatcagcggcgccggtgaccgcgagtgggtcggcaccgtctccggctc
cgggtgcgaacaatg (Seq ID No: 756)

Homo sapiens syntabulin (syntaxin-interacting) (SYBU):
cctcctcctggacggcgccgagcgccgagcgagccggcgccgagcgccgagcgatg
(Seq ID No: 757)

5 Homo sapiens coiled-coil-helix-coiled-coil-helix domain containing
3 (CHCHD3):
gcgcccttctccttgcttctgggggtcgtggccttgctcccgctgtgccccgaaagaatccaggccc
ttccacgcgcgtgtgggtgccccggcccgaaagtgtcgtggttccccgctaggtctccgctgggg
caggaaccggaatcatg (Seq ID No: 758)

10 Homo sapiens HAUS augmin-like complex, subunit 4 (HAUS4):
cctccttcgtcgccgctctagtgcactttcggctccttcccccttccccgggcctttcagcttggtc
tttccgggcctcgcttccccagccctgcgccccggcccgaaacgagaggttccggagccccggcg
ggcggttctggtgttagacgtgctggccagcccgccccagccgaggttctcggcaccgcctt
gagagcttcagctgccccaggattagaatcccaagaaaatcaaag (Seq ID No: 759)

15 Homo sapiens solute carrier family 41, member 3 (SLC41A3):
ccgcctctttcccgccgcgcctgggaggggacccgggctgccaggcgccagctgtgccagatg
(Seq ID No: 760)

Homo sapiens phosphatidylinositol glycan anchor biosynthesis, clas
s V (PIGV):
20 cttcctttccagcctcccgccctcgtctgcttccggccctgtggcctgggtggggctctgcaggctc
cctcgggagtggtccttggggcgtggccctctgggaggcctgaggagctcaatcctggtagcaa
caccctgaattcctgggtggtgaaaggatg (Seq ID No: 761)

Homo sapiens poly (ADP-ribose) polymerase family, member 16
(PARP16):
25 agttcctttatccctgggcccacctccccgcccagcccggtccaggcctcggtctctctcttcg
gcgccgagccgcggccagaccccgccagaggacacttgctcgccacgttctcaccctgtcatctc
agccccctgcctagctccacccaggttgggaacccggccctgacggccattgtccgcggggcc
cagccccgcgctgaacgcacgctcgcccttggccctaaccagcgctctaccccggaacgcgca
gtgacctgggatg (Seq ID No: 762)

30 Homo sapiens thioredoxin-like 4B (TXNL4B):
gtttcttttctgcgcttgctgcttttctgttcggtttccttccccgctagcggggccacgagggttg
ctaggcaacagccctgggtgacttggtcttagggctcctgtccggcttggggctgatgaaaggagc
tgtccgcgcccgggtcttccgagaagtgggtgctgacagccacaaagtgaaggagtgaggcgg
cgtggacgagtaaggagtgacagtgaggattcacatttgggttatttcaagatg
(Seq ID No: 763)

35 Homo sapiens slingshot homolog 3 (Drosophila) (SSH3):
cgtccttctcgtgctgccccgagcaggttgcgggggtgagggagggggccgtgccccggtgc
cagcccaggtgctcgggcctggctccatg (Seq ID No: 764)

40 Homo sapiens zinc finger protein 692 (ZNF692):
ctccctctggggcgccggcctcagttccgggctacagcagccgagccgagaggcaccgtttcttc
ttaaagagaaacgctgcgcgcgcgaggtggggccctgtcttccagcagctccgggcctgctcgt
aggcccgaggcgagggcgagggcgaggtgggggtgagggcgctggggcgccacagcctctggt
gcacatg (Seq ID No: 765)

45 Homo sapiens tRNA-histidine guanylyltransferase 1-like
(S. cerevisiae) (THG1L): tggccctttcctttccgcgtgtagaatg
(Seq ID No: 766)

Homo sapiens solute carrier family 25, member 38 (SLC25A38):

tctcccttctacagagttcctccggcgcttctccacccgggatacacagaacctcatctccta
cgggtgctgaagcctgcagcagggcaggatgggcaggagagcagagccgcggagtctgcggcgcggg
tgaagagcggcgcgtaattcccgagcaagattgttccgcgcccgcagcccctggactagcaggat
5 ccgaaccccgggcggtgctgcttataaggcgcagacgtcagagagcccgcggttaaagcgctcg
cctggctagcgccacccctagccttcttcaaggcctccagggtgggcccagcgcccgctcgacg
gcaccctgggcccagaggactcggggcctcatctccaatg (Seq ID No: 767)

Homo sapiens WD repeat domain 13 (WDR13):

agttctttctgatagcaggcagccatcttgctggagcctgagaaagggaggagagacagaaggaa
10 ccggcgacagtggctctcagggccgctccggggggcctcaagaaccggaggcagccccggaggtgg
ccccgatcccggtatgctcttgatctgagaagggaaggcggaggggcgggggacaagatggg
tggagaatgtcaagcaaggaatgctaggcgggggagggcggttgctatggcgactggggagggcg
gtgtctgttctgaatcgctgtgtgtcaccggcgctgccaggaagggcagggtgggggtgatga
ccatggtaacacccggggggagttcgtgacatctccggcgcgaggggactcgatgtctatggcaa
15 tggtcgcctggtggaagggacggaactagatcccttcgctcgggacgctcacattccaggcccttg
tcctgcaggctgccggggcgacacgccagaggaggaggccggggaatg
(Seq ID No: 768)

Homo sapiens chromosome 1 open reading frame 123 (Clorf123):

ccgccttttacgacgcgccggaagcaacggcaagggcggcagccagcaccgggaggagagggcta
20 ccatg (Seq ID No: 769)

Homo sapiens chromosome 20 open reading frame 11 (C20orf11):

ctgcctccttctactcgggcgccccggcgccgacacctctcccagcccaggagaggctgcggag
ccgcagccgccagaccgcgcagcgggaggcaggttccgcacgaaataaatcagaatg
(Seq ID No: 770)

25 Homo sapiens zinc finger protein 446 (ZNF446):

ttcccttttggggacagatcccgaagttcgagcatccctcggataggccgggtgtcaggcctggt
ctctcaggcccgctccaggcccatcttgacgattccaagaccaccccttgagcaagaatg
(Seq ID No: 771)

Homo sapiens mitofusin 1 (MFN1):

30 ccgccctttgccactccccctgcctcctctccgcctttaacttctcggaagatgaggcagtttgg
catctgtggccgagttgctgttgccgggtgatagttggagcggagacttagcataatg
(Seq ID No: 772)

Homo sapiens phosphotyrosine interaction domain containing 1
(PID1):

35 agtcctctcgagctgcgccaggacagccggcgcgcgccgtgccacaagttgccggcagctgag
cgccgcgcctcctcctgctcgcagccccctacgccaccccgcgcggtggccagcgccaggacgc
acatcccgcggacaccgaccccgatgtaaagcgggaccccgccccctcgcccccgcgcgatcg
acagtctcgccagcgtctcctctgcaaaacccagggtggaagatgtggcagccggccacggagc
gcctgcaggagagatttgcagacacagaagcggcacagagaaggccattgtgaagatcaaggcaga
40 aaccggagttatggcatcataagccaaggaatg (Seq ID No: 773)

Homo sapiens pleckstrin homology domain interacting protein
(PHIP):

45 tttcctcctcctcctcctccgcctccgcgcggttgcttgaatggtggagccgaagctcggctcgt
gaacacacactgacagctatagggcaggcgggcgccacgtccccgcttcccctcggcggggggtg
tcccgctcggcgccctgaagtgaccataaacatg (Seq ID No: 774)

Homo sapiens LIM and senescent cell antigen-like domains 2
(LIMS2):

tggccttttttgggcgtctccctgctccgcggcccggtggcgggcgggcgctcggtggcggt
gcagcagcagaggagaccgcggcaaccccggaacccagggtcggcgtcgctgccaccatg

5 (Seq ID No: 775)

Homo sapiens SCY1-like 2 (S. cerevisiae) (SCYL2):

aggtcttttagtctttttccccctcccttactcttcgtccccgggtccctcccccccccttt
ccttctagctccgacgtttgcgccgcggggcgggcgaggatatggagtaaagccagagtcagtg
gccaggcacgaaggcagagcaggaacagccaggaggcggtttattagggggcggggggaaagagcc
10 ccagcacgcgccctcctggaagaaggaagaggttaagtgaccggccgcggccaccgaccgacctcc
tcaccggcggtctctcgcctgggctcccgagccggcgaggagggaatggaggactcgcgcccg
gttaggcctccaggggcgctcaggctgggtgggtgttgctggtgacgggctgcccggcgccggc
cgggcgatcggcggctcggcgcccgcgcaaagcggggtggacgagcagcgagctccggggagcgga
tccgagagggccgagtcctcgaaagaggccttgaggcgacgggagaccgggatcgaaagtcagctg
15 ccggagggagagccccccatgccggctcgagagctcggggttcggtgggtggagaacgtagtacct
tcggggacattggacactactctaggaccgggtaactataactaccaatattgcagccatg
(Seq ID No: 776)

Homo sapiens ring finger protein 31 (RNF31):

caccctctctcctagtagtacttctgttctcggctaaccctggcgctgggcccgggggctggagagtga
20 ccgtgggtctgagtgaactggggcggtgctggtggcgccgggtgggcctcaaagccgggcaccagacg
ggagggcggcgctcgggccgcgctgcccgcgcccgggtcctggcgggcggcgaggctggggctg
actcctgcctcaggatg (Seq ID No: 777)

Homo sapiens mediator complex subunit 9 (MED9): cgacctctggctaacc-
tcccccgagccatg (Seq ID No: 778)

25 Homo sapiens ATP5S-like (ATP5SL): cggccccttcgggttacgaaaccttagcaa-
gatg (Seq ID No: 779)

Homo sapiens GPN-loop GTPase 2 (GPN2):

tctccttttgcgcgacaggtctcagctgttccgcctgaggcgagtgacgctggccgccaacgagg
tatacgtactgggaccctcgccctcagctctcgtctcggcgcggtacctgccccgttttccctgt
30 gaggtagctgctccgggcccgcgggcccgaatg (Seq ID No: 780)

Homo sapiens transmembrane protein 48 (TMEM48):

cggctctctgtacgccctagactaggggcccgcctctccatg (Seq ID No: 781)

Homo sapiens ankyrin repeat and zinc finger domain containing 1
(ANKZF1):

35 ttgtcctcttcgctgctccgtagtgacggggattgttggttgagaaatccggcaatcgacctga
ggacttgagagccgctcagctcccgggacgtttggagctgctgctaaataatttctgctcagccat
g (Seq ID No: 782)

Homo sapiens notchless homolog 1 (Drosophila) (NLE1):

ggctctttctcctccacgtggggacgcaggatg (Seq ID No: 783)

40 Homo sapiens cell division cycle associated 8 (CDCA8):

cgctctctctcactggcacagcgaggttttgctcagcccttgctcgggaccgcagcctccgccga
gcgccatg (Seq ID No: 784)

Homo sapiens polymerase (RNA) III (DNA directed) polypeptide E
(80kD) (POLR3E):

cgctccccccacgtgtccgcccggagtttctccaccagcaacatggccgcccgcctgagaggagagc
cgggcccgcgcgtctctgcagcccgcgggtaactgggcccgttgccgcccgtccgcgctcggcccc
gcggagagatcgagctgaaggactgcgcggctggctctcctctagtatg (Seq ID No: 785)

Homo sapiens armadillo repeat containing 1 (ARMC1):

5 gagcctttgccgcacgccttcgctctttggctccctgagttagtcgcgttgcttgcatcgcc
gcggccggggctgcgaaccgaagggtcgctccgcgcgcctgggtctctacctcatccgtaggtg
tgccctgatggtgtggcaggctctggactcctaaagctctggagcgaatttaagattttattcat
gtgcatggcatagaagatg (Seq ID No: 786)

Homo sapiens transmembrane protein 33 (TMEM33):

10 ccgtctttctggaacaccgctttgatctcggcgggtgcgggacaggtacctcccggctgctgcggg
tgccctggatccagtcggctgcaccaggcgagcgagacccttcctgggtggaggctcagagttccg
gcagggtgcatccggcctgtgtgtggcgcgaggcaggaagccggtaccgggtcctggccccagc
gctgacgttttctctcccctttctctctcttcgcgggttgccgctcgcagacgctagtgtgagcc
cccatg (Seq ID No: 787)

Homo sapiens pyridoxamine 5'-phosphate oxidase (PNPO):

15 ccttccttccccggggtagaagtcagggtgagaaattggttcgaactcaaaggaaccagtgcc
gggccacagccgggtcacgtggccggcgccccccatg (Seq ID No: 788)

Homo sapiens golgi phosphoprotein 3-like (GOLPH3L):

20 attccttctctgcatcgaaggatcaggaagtttgctctctgcgtggctaagtttttcacctact
aggacgggggtggggtggggagaaacaggtgtccttctaaaatacagcacaagctacagcctgcgtc
cagccataaccaggagtaacatcagaaacaggtgagaatg (Seq ID No: 789)

Homo sapiens regulator of chromosome condensation (RCC1) and BTB
(POZ) domain containing protein 1 (RCBTB1):

25 cgctcctcctcttcgctgcccgtgggacccgcgcctcgcctcgcacttctgcgcccattggagcttc
ggagatccctgcggtcccgcgggacggcgcggcagcagctgacctcgcagacaggatcttgctctc
ttgccagactggaatacagtgggtgtgaacacgggtcactgcagcctcaacctcctggactcagag
atgtcggccttatttataggaattgcttgaagccagagtcatg (Seq ID No: 790)

Homo sapiens leprecan-like 1 (LEPREL1):

30 cgtccctttaagagcggctggccaggcacggcctccgcctctcagtacgcggagcgccggcggtca
cctggggctcgcggagcggccagatcgcggcggagtcggcgcgcttccccgaggggaaggtgggaga
ggggaccgcgagcgggtgccccgaagccctctcgagcgtaaccgtcccgcgcctctctgaggcg
gaggatg (Seq ID No: 791)

Homo sapiens hedgehog acyltransferase (HHAT):

35 ctgtctcttggtcaggcttgaggcctccgagcagcaacatcgctcccaattataccccgttgag
catcttcagatcttccactcttttcacaacgcaatcaaaatcttcgtaccttttgtagtagtga
tctctgtaagttgctttacaattcataaagtttattctatttgatcttcactctaatttacaaga
aaagcaggggaagtctattttctgtttttacagaggtgtacagggaggctcacaggggctaagttcaca
cagtaagccctcgaagctgccagggtgcaaagcccacctctttccaccgcaccgaactacctcc
tttcgcctacaaaacgtaggtggggaccactggtgttggaatgacggcccacctcgagtttcagggt
40 gacttccactctgcaattaacttgaggcagccccagacctgcaatgaacacacgggtgggggaga
gatatgcacgccagggtcagtgggaaaccaacagccgaggggtgagcggggctaggggccccgggccc
gccggcggggcaaacgcgggttcagaaacgcaggccgcgctctggcccggccccctgcagcagcacgg
cctgctcgccatcgcccggagagcgccgcgggttcccagatccgggcgcggaggggcgcgcgggcac
ggcggcaggggcgtgctcggaggacgcgcgctgcgctgctcctccaaagggcagctccgggggaaa
45 gagggtggcgtcccgggggaagcccgcagccgcgcgatgtcgtggtgactcggaagtgccgaaag
aggggtgttggaactcgcggcgcgctgaacgttgccgtcgccgcccggggacagcccggaga
aactctcagcgtaggcatcggaaccttcgtgccaaaggagccatg (Seq ID No: 792)

Homo sapiens chromosome 11 open reading frame 57 (C11orf57):

cctcctttttctcccaaaccacttcttccccctacccccgccacgcgaggctgcggcgcacggt
atgggtgtgtttgtgtgtatattgtgtggggagggcggtttggagggaagggtaccgggagctccgag
gccgctggggaacagggatcccgggtgacaaagatggggatatttcctctgtcttccacttgaaac
5 ctcaacccccgcttcaggctccctagatactttctggggcccaaccgaaggccgtagccatccaaa
gcgttcccagcctttctggggagtgaacttaccctccgggttcgtcctagaggagcgtgagcggg
gaatgccaggtcaaccgggctgtccgaattccgccccggctcagcctccggcctcagtcggggag
agagatctgcctgtcgggtctgggctgggggaaacgcggcagtggcctggggcacagggtgagggcag
agtaaccagtgagggaaggctgcgttttccagaaggactcgggtgaagctgcagagctgcctttgagc
10 cctgactccttggttctctgggtcggaggagatcttgtaatggagtgggttcttcgtctcactagca
agatgcctgatttcctcaggatcaagggtgaagaatg (Seq ID No: 793)

Homo sapiens high mobility group 20A (HMG20A):

agtccttcgccgcattggggcaaaataatcccttcatttttgtgaagggtaccgtggaaaatatttc
atttttcttctcaccggagcaattgtaaatgctatgcggtaagaggagttacctgtggaaagggtg
15 ttaagagattaggtaaagaaaaggaaaggacaccaaataaagtgcgcggaagaatttttgtcca
gctgtgagacgacgagtgcgtagtggaaggcgattgagaggggctgaggggaattgtcctctgtgg
aagggactttcttttggccctaggcccccttctgccccctgtcgtcagcagagtctctacaaggaag
ataacggactgtaaaattctataaagcaaaagctacacatcacttgacaccatacaccatcttggtt
acataatgaagagagatg (Seq ID No: 794)

20 Homo sapiens checkpoint with forkhead and ring finger domains, E3
ubiquitin protein ligase (CHFR):
atgtctcttgacagcggcgggcgagccgggttcggggttcggcgcgggggcggggatgtgaatcc
cgatg (Seq ID No: 795)

Homo sapiens nucleoporin 133kDa (NUP133):

25 ccatctcttcccttaggtgtttaagttccgcgcgcaggccaggctgcaacctgacggccagatccc
tcgctgtccttagtcgctgctccttgagatcatg (Seq ID No: 796)

Homo sapiens CNDP dipeptidase 2 (metallopeptidase M20 family)
(CNDP2): cttccttccaagaaccttcgagatctgcgggtctgggggtctggttgaaagatg
(Seq ID No: 797)

30 Homo sapiens oxoglutarate dehydrogenase-like (OGDHL):
gcaccccttccgcgcagccccctgacctgcagcctccggacctcgctgcagcgcggacccggccccg
cccggccgaatg (Seq ID No: 798)

Homo sapiens transmembrane protein 30A (TMEM30A):

35 ccgcctcttccgctctacagcggagggtggctgtggcggtggcgctggtggctgcggcgggcgggc
ggcagcggcgctcgagcgggttctgtcagggtcagccggcgggccccctgggtgggtccacctgcaa
atcgcgagcggcgccccagggtatcgatg (Seq ID No: 799)

Homo sapiens elongation protein 2 homolog (S. cerevisiae) (ELP2):
gcgtctcttgtttgtgcggctgaccagttggcgacatg (Seq ID No: 800)

Homo sapiens WD repeat domain 12 (WDR12):

40 cgttcttttctttgtattttccgcctctcgcctctctctaaaagccgcagtttagaggcgagatttag
gaaaaacctctgccgagtgagcctctgggtgggaatatgtatgagaaaaaaaaactggcaaggcgt
tagtcaagcaaagctgaaggcagaggaaatttgatatctggctggagtctagaggatttaatgcaa
ataagatactctgagggcagcgtggcaaaaaaagactacaattcccggtgggtcacagcgtttgaga
agcgatgctttctgagacttgtagtaactaggagctgtgtttgaactatccaggctcaggacagcc
45 tcttgaaaaaaaaatttttattaataaagcggatttgagtgggatctttttcctaatacgattacgg
gccacacgtatgggaagaattctaacaatgattaaaggacatgctacctttacgactatccttt

tctaatacgatgactcctaaatctaggagtaggtagtcgatgtttgtggtctgggcgtctgtagaag
ggcaacctcgtgctttctgcagaggagaccggaggggcagaaggcagagtccaggcttagactgcag
ttcctcgcttacctgtgcagtctaattttgagctgcctctttgtagtcttaaaaggcaggagcttc
gtgttggtgggtctgctaaccggtacgtttccgtgggcaagtcgtgtgtactcctcgccatg

5 (Seq ID No: 801)

Homo sapiens tetratricopeptide repeat domain 17 (TTC17):
cgacctcttcaagatggcgggcccggagactagcttccgcttccggtgtgagcggcccgccggg
ggggcaagatg (Seq ID No: 802)

Homo sapiens proline rich 11 (PRR11):
10 ttttctttatggcgtgggagaggccacagcccgactccatcgactcccccggtcttagactaaa
atcatg (Seq ID No: 803)

Homo sapiens TBC1 domain family, member 23 (TBC1D23):
ctccctctttcttcccctctggggaagctcagtgctggacttccgaagaccttttacgacattgag
tctcgagattggtctcagcgccggatccacttttcggcaaagtgacgtggacgtcaacagcaatg
15 (Seq ID No: 804)

Homo sapiens leucine rich repeat neuronal 3 (LRRN3):
gctcctctctggggagtggagggtgttcagttattaatgaccgctgagcaggcagcaccatgtcag
tgtgacaactgatcgggtgaacgatgcaccactaaccaccatggaaacaaggaaaaataaagccag
ctcacaggatctctcttccactggattgagagcctcagcctgccgactgagaaaaagagttccagga
20 aaaagaaggaatcccggctgcagcctcctgccttctttatattttaaaatagagagataagattg
cgtgcatgtgtgcatatctatagatatatattttgtacactttgttacacagacacacaaatgcacc
tatttataaccgggcaagaacacaaccatgtgattatctcaaccaaggaactgaggaatccagcacg
caaggacatcggagggtgggctagcactgaaactgcttttcaagcatcatgctgtattcctgcaaa
tactgaagaagcatgggattttaaataattttacttctaaataaatgaattactcaatctcctatgac
25 catctatacatactccaccttcaaaaagtacatcaatattatatcattaaggaaatagtaaccttc
tcttctccaatatgcatgacattttttggacaatgcaattgtggcactggcacttatttcagtgaag
aaaaactttgtggttctatggcattcatcatttgacaaatgcaagcatcttccttatcaatcagct
cctattgaacttactagcactgactgtggaatccttaagggccattacatttctgaagaagaaag
ctaagatg (Seq ID No: 805)

Homo sapiens MIS18 binding protein 1 (MIS18BP1):
30 ggccctctctccgcgcggagccgagccggaactgcggcagtcctctccctgccaggctcttcatcca
aggtttctgtggatcccttctgaagttctatctgaaaattgcgcttaagtgaattttctgtagaa
gaacttggttgctacttttcttgtcaagatg (Seq ID No: 806)

Homo sapiens LMBR1 domain containing 1 (LMBRD1):
35 ccgccccctttaaccttttaggggtgcgcgggtgcagtatatctcgcgctctctccccctttccccctcc
cctttccccaccccgggcgctcaggttggtctggaccggaagcgaagatg
(Seq ID No: 807)

Homo sapiens ST6 (alpha-
40 pha-N-acetyl-neuraminy-2,3-beta-galactosyl-1,3)-N-acetylgalactosa
minide alpha-2,6-sialyltransferase 1 (ST6GALNAC1): cttcctcta-
gaacccgacccaccaccatg (Seq ID No: 808)

Homo sapiens spermatogenesis associated 7 (SPATA7):
gctcctctttttccagtcctccactgcgggggtgggcccggccgcgggaaggaccgaaggggatac
agcgtgtccctgcggcggtgcaagaggactaagcatg (Seq ID No: 809)

Homo sapiens zinc finger, C4H2 domain containing (ZC4H2):
aggcctctccaagcccctaccgcacaggctcatagccccaagcccggaggaggtggctacattgtg
tctattgtatcccttggtgtgtttgtacatctctcgggacgtgaaattgacagtgaaaagta
tg (Seq ID No: 821)

Homo sapiens BAI1-associated protein 2-like 1 (BAIAP2L1):
cttcctctggcgcggtccggccgcttctcctctgctcctcgaagaaggccagggcggcgctgccgc
aagttttgacattttcgcagcggagacgcgcgcgggcactctcgggccgacggctgcggcggcggc
cgaccctccagagccccttagtcgcgccccggccctcccgctgcccgagtcggcgggccacgagg
cccagcgcgctcctcccgcgcttgctcgcgccggcgggccgcagccatg (Seq ID No: 822)

Homo sapiens solute carrier family 25, member 40 (SLC25A40):
cgtccttctcgcgcctcgctctggccctgcaggttggtttccgcctctacccgcctccattccg
ttgctctctcagtcctcagacccgggctctcggtccgcgcgttcaggtcttggcgcagcctcagaga
gttggcgcggctctgtgttgaccaaactagtggatgcagttagcgcggagcccgccccgcccgcg
tcaccaggggtattccgccttctaggtttgccaggactgccggccctgcagctgccttctgcccc
aggtttttggtactgatgttacaaacaataaaatattggagcatagagttgaagaacagactcaa
accaggtttttatttaattagttaaaaaatatg (Seq ID No: 823)

Homo sapiens protocadherin alpha subfamily C, 2 (PCDHAC2):
 tttcctttttccctccccctggagctgtagcggcagcagcagcaggaagccgagccgggttgagcga
 ctcgaggagcgagcggaggagctggaatatggggagtcagcgaggacgggtggggccaggagcccttg
 ggagggcctacggagggagcggccccaggcgctttctagagcgtgagcgggtgggggagcaggcgca
 ggggtggcacgagcggaggcggggcccgggcgctggggcacggctggggaagctgccgcctccggccc
 tgccccgctgcctccgcgcgcggccagtggtatg (Seq ID No: 824)

Homo sapiens chondroitin polymerizing factor 2 (CHPF2):
gttcctttttgggttagctttggcagttattgagttttacttcctcctcttttttagtggagacaga
ccataatcccagtgtagtgaaattgattgtttcattttattaccgttttggctgggggttagttcc
gacaccttcacagttgaagagcaggcagaaggagttgtgaagacaggacaatcttcttggggatgc
tggtcctggaagccagcgggcctcgctctgtctttggcctcattgaccccaggttctctggttaaa
actgaaagcctactactggcctggtgcccatcaatccattgatccttgaggctgtgccctggggc
accacctggcagggcctaccaccatg (Seq ID No: 825)

Homo sapiens thioredoxin-related transmembrane protein 3 (TMX3):
gcttctcttccgctccgggtcggctccgtttccctttccgggcgggcaggcggcggaacccagtgt
ctttatccctcttttgcacagtcagcttctgcagctctccgggctagcatg
(Seq ID No: 826)

Homo sapiens ras homolog family member F (in filopodia) (RHOF):
cgacctcttggtccgctagtgcccggcgcgcgcgcgcagtgctgcgggctccgggcaatg
(Seq ID No: 827)

Homo sapiens amyloid beta
(A4) precursor protein-binding, family B, member 1 interacting pro
tein (APBB1IP):
ctttctctcaggaactccactcccaactgacaggtgctatttccagccagtcctatgctgttgca

Homo sapiens roundabout, axon guidance receptor, homolog 4 (Drosophila) (ROBO4):
ccttccctcttctactgtgagctcagagcagcaggacaaagtgctcgggacaaggacatagggctga
gagtagccatg (Seq ID No: 829)

Homo sapiens translocase of outer mitochondrial membrane 7 homolog (yeast) (TOMM7):
acctcctttccctttcggattcccgacgctgtggttgctgtaaggggtcctccctgcgccacacgg
ccgtcgccatg (Seq ID No: 830)

Homo sapiens major histocompatibility complex, class II, DR alpha (HLA-DRA):
 ttttcttttattcttgtctgttctgcctcactcccgagctctactgactcccaacagagcgcccaa
 gaagaaaatg (Seq ID No: 831)

Homo sapiens protein arginine methyltransferase 8 (PRMT8):
cctcctctactatctcggatcaccaaaccttgccggctcttatg (Seq ID No: 832)

[illegible]

Homo sapiens BarH-like homeobox 1 (BARHL1):
agcccttttg gatcta atgcgcagaggaggttg gccagagctcccgggctcccccaaggctgaac
tccgtccaagggtcccgcaggctccctgccgccttccccatgccagccgcagctaggggcaggg
gcagcggcggtgtgggttggggtgggtggggagcttttggggaggacaggtcgcagcttggctat
g (Seq ID No: 834)

Homo sapiens intraflagellar transport 46 homolog (Chlamydomonas) (IFT46):
 ttatctttttgcctagcgactgacaacaggctggttgcttgcgctggaatcctaaagtggcctggc
 ttgagactggagtgagaccccagccctaggctgggggttctttccattatagaggagacggattca
 gaagggctacagaccaaggttggtgaaaaccagacatatgatgagcgtctagagattaacgactcc
 gaagagggttgcaagtattttatactccaaccccaagacaccaaggacttcctcgttctgcccattt
 cctaacaaggctatg (Seq ID No: 835)

Homo sapiens dual specificity phosphatase 22 (DUSP22):

Homo sapiens phosphoribosyl transferase domain containing 1

Homo sapiens arrestin, beta 1 (ARRB1):

Homo sapiens leucine zipper transcription factor-like 1 (LZTFL1):

45 taccctccttccccattttctgtggtccaactaccctcggcgatcccaggcttggcggggcacgc
ctggcctctccgttcctttaggctgccgcgcgtgcctgccgccatg (Seq ID No: 843)

Homo sapiens phospholipid scramblase 4 (PLSCR4):

agccctcccttcgcgcgcttactttgtttataacttgaaaaatcctctccgtctcccttcctgc
ctcctttcctttccctttcctctgccagtacaactagaccggcgctgtggcgctccccggtgccag
cattctgcggggcaggcggttaattggaattcttcaaaatg (Seq ID No: 844)

- 5 Homo sapiens ectonucleoside triphosphate diphosphohydrolase 7
(ENTPD7):

cctccttcggctgggcaaggggcccgcggggagcagctcgggactgaaccgagagggtgccgaagga
accggcgggcgcttgatcccgcgtgcagacgtaggagatgcctgggacaaggaggccaccttctca
gggcaaaagaaaaagaaggtgacaggcggtgagaccaccaaggaacccatg

- 10 (Seq ID No: 845)

Homo sapiens fascin homolog 3, actin-bundling protein, testicular
(Strongylocentrotus purpuratus) (FSCN3):

agttctctctgggaacatctggtgggtactacaggccctattccaggccctatggcctgtggaacc
tcaccacgggggggagggtgggcccagacggagacatcacctgtggtgtcagcccatg

- 15 (Seq ID No: 846)

Homo sapiens X-prolyl aminopeptidase (aminopeptidase P) 1, soluble
(XPNPEP1):

cctccttcgcgcggcccttcgcgggtgatcagctggtctgcgctcccctgacgtgggctggggc
acgtcaccgccgaatg (Seq ID No: 847)

- 20 Homo sapiens REX4, RNA exonuclease 4 homolog (S. cerevisiae)
(REXO4):

gggtctcttcggagtcttttcttgacgggggtccctgcggtgggtgtgtttcggcctggcctggg
caggcgcttgctgctgccaggcgccggggccggggaggccgggtctcgggtggccgcccggccag
gcgctggacggcagcaggatg (Seq ID No: 848)

- 25 Homo sapiens LYR motif containing 4 (LYRM4): ttttctttccaaaatg
(Seq ID No: 849)

Homo sapiens DEAD (Asp-Glu-Ala-Asp) box polypeptide 24 (DDX24):

ggttcttctactcgcgactgacggagctgcggtggcgctctccacacgaaccatg
(Seq ID No: 850)

- 30 Homo sapiens transmembrane protein 159 (TMEM159):

ccttcttctcttctgttctcctcctgcctctcttcgcttcgcctgcaaacgcggtgggggctgctc
ggcggtcaggagcaggttaccctccgtctgcatgccaccatcaaggtatgaggatggtagaagct
ctcgtcgaaccagatggatgaagaccactaacggcttttgtttcctctggtaacagcaagagacag
agcgacatgagagattggaccgcgggctgcactggagaatttactggtaggataattcatccctaa
35 agagattgaagtgaagcttcagaatg (Seq ID No: 851)

Homo sapiens NDRG family member 4 (NDRG4):

cggcctccgcccctgcagccgcgggcacgcggaggggctcctggctgccgcacctgcacccgcgc
gtcggcgggcgccgaagccccgctccccgcctgcgcgtctgtctcgtccgcacatctccgcggcctcct
gctccacgacgtgaccatg (Seq ID No: 852)

- 40 Homo sapiens pre-B-cell leukemia homeobox interacting protein 1
(PBXIP1):

ttttcttctcgggctgcaaacaaaggggaagcctgcaacaagttaagctgaagaccgaagcaagagc
tggttcaggtggcagccacagcagcctcaggacctcagcaactatg (Seq ID No: 853)

Homo sapiens twisted gastrulation homolog 1 (Drosophila) (TWSG1):
ctgtctctttaaggtgcccgaggctcgcgggcgctgcgctgaggggacggcgggaggcgcggcctg
gcctcgactcaaagccgcccagcgcgccccgggctcgccgacccggcggggatctaggggtgg
gcgacttcgcgggaccgtggcgcatgtttcctgggagttactgatcatcttctttgaagaacatg
5 (Seq ID No: 854)

Homo sapiens zinc finger protein 286A (ZNF286A):
gtcccccttgtgaggccccgggatgggaggtgcccggttccccaggacagcttcaagcggtaggg
acagacatctgaggaccagcctcagggatgctgtccccgggcttccaggctccagcgccgtagga
ctgaggcagactccacggtgagaaagagacccgatctaaccaggcctttcatcagagcccaggag
10 ggaaggcaggaagtgggaccacgaggccccggggggttctaactcgtctggccaggagatctgaa
ttgggggtgaagagcagaatctccagaacaaggaggaggtggtgatcatg (Seq ID No: 855)

Homo sapiens S100 calcium binding protein A14 (S100A14):
gctcctcctgtcttgtctcagcggtgccaacagatcatgagccatcagctcctctggggccagct
ataggacaacagaactctcaccaaaggaccagacacagtgggcaccatg (Seq ID No: 856)

15 Homo sapiens ANKHD1-EIF4EBP3 readthrough (ANKHD1-EIF4EBP3):
tgctcttctcgttcccagatcagcgggcggtgaccgcgagtgggtcggcaccgtctccggctc
cggtgccaacaatg (Seq ID No: 857)

Homo sapiens KIAA1143 (KIAA1143): ctgtctttaccagagctaccatg
(Seq ID No: 858)

20 Homo sapiens neuroligin 4, X-linked (NLGN4X):
ctctctttttcttgcagaaccgtctctctcccttctctgtctcttagcacagagctcttattcagc
cactagcttggcccttctctgcttcaattgtaatgcttgttctgcccgtccacagactattggcggc
agaaacaacgaatttctccaaactaggcggtgttggtggctcttgcattcctctggatgaggaaa
tctagttgggggttccagaagggaaggctcctgggctttcaatacatcctcctgaatcatacct
25 cgtttcgggttccctagaaaaatctggacgtgtaaaaagaactcttaacggccgatgcagctcttc
caaagctaaggctgccttggagttttcataagaaattgtccctggaggtgttggtgatcacagct
tccttggagcattgcagttgctggaatccagtttcaggattaagggagggctgcctccttgcaatg
ggctgccaaagaaaacggctgtgcttgttcttaacctcaggctctgtctgtgatcagctctgagagtc
tctcccaggtctactgctccctggaaagccctatctctctgcaggctcgctctgggctttgtctc
30 cttggagccacatcactgggacagctgtggatgtggatgcagatttgaaccatg
(Seq ID No: 859)

Homo sapiens mitochondrial antiviral signaling protein (MAVS):
ccgcctcctcgctgcgggaagggtcctgggccccgggcgggcggtcgccaggtctcagggccggggg
taccgagctctgtttctctcagtcacatccacccttcatggggccagagccctctctccagaatc
35 tgagcagcaatg (Seq ID No: 860)

Homo sapiens serine incorporator 1 (SERINC1):
ctgtctccatcttgtctgtatccgctgctcttgtgacgttgtggagatg (Seq ID No: 861)

Homo sapiens KIAA1324 (KIAA1324):
cctcccccttttttccgccttctgccagcagaagcagcagccgcagcacctgagccgctactgccg
40 ctactcaggacaacgctatg (Seq ID No: 862)

Homo sapiens synaptotagmin IV (SYT4):
ggacctccctctttgctcctcctgttccaggagctgggtgccctgggctctgcgctgttgttttc
agcgctccgaaagccggcgcttgagatccaggcaagtgaatccagccaggcagttttcccttcagc
acctcggacagaacacgcagtaaaaaatg (Seq ID No: 863)

Homo sapiens pyruvate dehydrogenase phosphatase catalytic subunit 2 (PDP2):

cttccttctgagctgggtcctgactagggaccgcctgggtgaggtgaggacctggtggccgcagt
tgtggcactgtgcgaggcgctgaactgaccggacggagcggcggtgtggcctcgccagctggt
5 ttaaaaatatccttttttctgaaggaacacatttgctggtatagtttcagaatg
(Seq ID No: 864)

Homo sapiens gephyrin (GPHN):

ctatcctttcctctcagtcctgccatctagctgccttgggtctcgcgctccgcagagcggtccgac
actctccggcctcggttctgcccctccgcgcgctctccccgtgcggccaccgcgcccccaagctt
10 gcctccttcttgccggaacttggggccgcgcgacctgactccttccccctccgcggaccgcgcact
cccggcgcggcctctccccacgcaggccaccgtgactctgtggcctccccctccttccccgctc
tcctcgcgcttctctggctccctagctgtcgcgctctcctcggcgagcgcgctcccgggccgcgcg
ctccgggctccggtttctcccggtcctgtcagtgcggtgactgcgctgggaacatg
(Seq ID No: 865)

15 Homo sapiens deltex homolog 2 (Drosophila) (DTX2):

ccttctcctgagagtgcggagccacagccagagccctgccaggccgagccggagctgcagcccgag
cgcggtggtgccctcagccccgtcctcttctcctcagcctcggtgccttggatttgtgtcgc
tgagtcagcaagcctttcagatttgcgcggtttttgttgtttgtggtttgtatcaagatgggaact
caaacaagtcattcctcctaaggagctggtgtcttcatccagaagggacagtttgtgccagctctc
20 cagagagaaaaggatctggtactgttctggagtggcctgtagcagacactgaaccaccagccagct
gcatttgttctcctggaagtcattgccaaactctgccagtcacactgggggtcccagagaagtcaag
atctgccggaggcgctgggcaatgaccccggaactccaggccagaggggtctgaagctgtttggga
aagcagcgggactccttgggaagatg (Seq ID No: 866)

Homo sapiens melanoma antigen family E, 1 (MAGEE1):

25 ctgcctttttcaccacctctaatttcagcttcagcagttgcttggaaactttggttctggcagcagc
agcaacatcattaccgctagcggcagttttgtgcccaggacacctacacacctcccgctcctctgc
cagatcgcgggcctgtcggtgtctgctcctacacgccaacgcgggtgggcaggaccatg
(Seq ID No: 867)

Homo sapiens G protein-coupled receptor 107 (GPR107):

30 cgccctttcaccgccgagctgggcgggagaggaagcggctggtgatgctggaacaaacatg
(Seq ID No: 868)

Homo sapiens PDZ and LIM domain 1 (PDLIM1):

cgctcttttctccgacagctgccgggggtgccctgcaagctgttccgcgcgctcctgcccgctctgtcc
ccgcgggtcgtcgcccgcacagccgcgccatg (Seq ID No: 869)

35 Homo sapiens thymosin beta 10 (TMSB10):

cgctcttttgttttcttctgctgcagcaacgcgagtgaggagcaccaggatctcgggctcggaacgagac
tgcacggattgttttaagaaaatg (Seq ID No: 870)

Homo sapiens phospholipid scramblase 1 (PLSCR1):

40 agacccttttcagacccttttccggctgacttctgagaaggttgccgagcagctgtgcccggcagct
ctagaggcgcagaagaggaagccatcgccctggccccggctctctggacctgtctcgctcgggagc
ggaaacagcggcagccagagaactgttttaaatcatg (Seq ID No: 871)

Homo sapiens eukaryotic translation elongation factor 1 beta 2 (EEF1B2):

45 gggctcctttttcctctcttccagcgtggggcgcccacaatttgcgcgctctctttctgctgctcccc
agctctcggatacagccgacacccatg (Seq ID No: 872)

Homo sapiens pyrophosphatase (inorganic) 1 (PPA1):

ggctctctccttgtcagtcggcgccgcgtgcgggctggtggctctgtggcagcggcgggcggcagga
ctccggcactatg (Seq ID No: 873)

Homo sapiens X-ray repair complementing defective repair in Chinese
5 e hamster cells 5 (double-strand-break rejoining) (XRCC5):

ggctctttccgctatctgccgcttgtccaccggaagcgagttgcgacacggcagggttcccgcccg
aagaagcgaccaaaagcgccctgaggaccggcaacatg (Seq ID No: 874)

Homo sapiens GATA zinc finger domain containing 1 (GATAD1):

10 gatccctttccagtcctgcttccagtgctcgggccagggaatcctggcctccgctgaggagc
cggcggaacccgcttcccgctccacggggcagcgccagcggcctggctcttccacggcagctcc
gtgccgacgctctcaccgctcttccctatcgccgggagtgggggccgaccagggggcgccgggct
accgtccgccattcccgctgtctctgcgcccgcgggggcccgcagccggccaccatg
(Seq ID No: 875)

Homo sapiens enolase-phosphatase 1 (ENOPH1):

15 ccgccttttccagttccaggtgtgcagaagtgtcctctcccccacgcgcggcgggctgcacttggtc
gctggctccgagatcgcgcggggcccgcggaagcccaagacggtaccgggggcccgcagccgcagcc
ggcgccgcctccgcctccccaacagcaggccgagtccttagcatccggtagggaaatg
(Seq ID No: 876)

Homo sapiens regulation of nuclear pre-mRNA domain containing 1B
20 (RPRD1B):

agctctttccgggggcccggggaactactctccttgctcgtctgtctccttcgaagtgcctctgc
gcgaggttcagagcggcgccgcctccaaagggacgggttttctagagctccgacgcctctcggtgc
ccctctgctccggcccttgccctttgacctcgtctctcgcgccaggggtgagaggtcggtggccatc
25 ttgtggcgggcggcgcgggcggtgttactgcggagaccatccctcccccttctcgcacccctgg
cagtctgtcagtcggtaaaaagtcccgagcctgtcaggtgagggcccgccctcgtgccgtcgtc
ttccgcgcgactggcgggcccaggccgctccctgcccgggctcactgccgccaccatg
(Seq ID No: 877)

Homo sapiens family with sequence similarity 60, member A
(FAM60A):

30 ctatcttttctagacaaggcagttgaggaggaggagcgcttgagggggactggcctggcgtgcact
ccgcacctcggggacattattgcgcggtggaacggctgcttttggaaggcacaacttcctgaatgga
ccatgactcccaccaaagatccctgtctctgattcaccaaacagcttcaaccctgaaaccaggagc
agaagttgacaacatctgagtggaacagctaattgacctaagacttcagaccagactattgccaga
agaaaagatg (Seq ID No: 878)

Homo sapiens MID1 interacting protein 1 (MID1IP1):

35 gggcctttttatctcggtgctgccgggggagggcgggaggagagacaccaggggtggccctgagcgc
cggcgacaccttttctggactataaattgagcacctgggatgggtagggggccaacgcagtcaccg
ccgtccgcagtcacagtcagccactgaccgcagcagcgcccttgcgtagcagccgcttgacgcga
gaacactgaattgccaacgagcaggagaggtctcaaggcgcaagaggaggccagggctcgaccaca
40 gagcaccctcagccatcgcgagtttccgggcgccaagccaggagaagccgcccacccgcagggc
cggctctgccagcgagacgagagttggcgaggcgaggagtgccgggaatcccgccacaccggcta
tagccagggccccagcgcgggccttgagagcgcggtgaaggcgggcatccccttgaccggccgac
catcccgtgcccctgcgtccctgcgtccaacgtccgcgcggccaccatg
(Seq ID No: 879)

Homo sapiens transmembrane protein 35 (TMEM35):

45 ctctccctttgtcattctagctgcctgctgcctccgcagcgtccccccagctctccctgtgctaac

tgctgcaccttggacagagcgggtgcgcaaatcagaaggattagttgggacctgccttggcgacc
ccatg (Seq ID No: 880)

Homo sapiens Fc fragment of IgG, low affinity IIa, receptor (CD32)
(FCGR2A): cttcctcttttctaagcttgtctctttaaaccactggacgttggcacagtgtg-
5 ggatg (Seq ID No: 881)

Homo sapiens tribbles homolog 2 (Drosophila) (TRIB2):

ctttctcttttggtttggcttctaacgcgttgggactgagtcgcgcgcgtgagctccccgaagact
gcacaaactaccgcgggtcctccgccccgtctgcgattcggaagccggcctgggggtcgcgtcgg
gagccctggcgctgcagctccgcaccttagcagcccggtactcatccagatccacgcgggggaca
10 cacacacagagtaactaaaagtgcggcgattctgcacatcgccgactgcttggggtaacaaaaag
accgagttgcctgccgaccgaggacccccgggagccgggtcggagcagacgaggtatccggcgg
cgccatttgggggttctaactctttctccacgcagccctcttctgtccctccctctcgtc
ccttttaaaatcagtggcaccgaggcgctgcagccgactcgccagcgactcatctctccagcgg
gttttttttggttgtcgtgtgcgatacctcacactcatg (Seq ID No: 882)

15 Homo sapiens family with sequence similarity 3, member A (FAM3A):
cgctctctccgggggaggagcgggtcggcgggcctgacaggaacctccctgaccgagcccacgtc
tccccagggccagagaaatctccggcccgcccgcatcgccagccccagggccggaggaacggcc
cgagcccaggagaaccacatcttcgtcccagccccggaggctcctgtgggcaagatcgtgagcaa
cgggttcttgaggccccctcctggccaggcagggtttccccgcgcgtttccgaggagccctgcctgg
20 ccgggcggtggacaaacaggtcgtagcaccgatcgcgccccccccagcaggggtcccgcacagg
cttgccttgacccccacccaaacctgtccttcgcgtttgcccccaaacagtgcacttgccggcgg
tcccaaccagcaggagaagtggacatg (Seq ID No: 883)

Homo sapiens exocyst complex component 4 (EXOC4):

ggctctccccgcgtccaagatg (Seq ID No: 884)

25 Homo sapiens ELOVL fatty acid elongase 5 (ELOVL5):

gcgcttctcttcccatcgcggggtcctagccaccggtgtctccttctacatccgcctctgcgc
cggctgccaccgcgctccctccgcgcgcgcttgctgctgctcaaagctgctgccgcccctt
gggctaaaaggttttcaaag (Seq ID No: 885)

Homo sapiens apolipoprotein B mRNA editing enzyme, catalytic polyp
30 eptide-like 3G (APOBEC3G):

ctttctctttccctttgcaattgccttgggtcctgcccgcacagagcggcctgtctttatcagaggt
ccctctgccagggggaggggccccagagaaaaccagaaagagggtgagagactgaggaagataaagc
gtcccagggcctcctacaccagcgctgagcaggaagcgggaggggcatgactacgaggccctgg
gaggtcacttttagggagggtgtcctaaaaccagaagcttgagcagaaagtgaaccctgggtgct
35 ccagacaaagatcttagtcgggactagccggccaaggatg (Seq ID No: 886)

Homo sapiens gamma-aminobutyric acid (GABA) B receptor, 1
(GABBR1):

gctcctcctcctccctccgctcggtcagtcagtcgcgcgaggagagtcgcgcgttggcggcgacgggtg
gcgagagccgcggggggcgtaggaagccaaccttccctgcttctccggggccctcgccccctcctc
40 ccacaaaatcagggatggaggcgctcccgccaccctcttagcagccctccccaggaaaagtgt
ccccctgagctcctaacgctccccaacagctaccctgccccccacgccatg
(Seq ID No: 887)

Homo sapiens cofilin 2 (muscle) (CFL2):

cctccttctcctccagtgccacagagccgaagcccagagctgccgcccagccacagccgagggga
45 ctatg (Seq ID No: 888)

Homo sapiens DEAH (Asp-Glu-Ala-His) box polypeptide 35 (DHX35):
tgacctttttaccccaacatg (Seq ID No: 889)

Homo sapiens resistance to inhibitors of cholinesterase 8 homolog
A (C. elegans) (RIC8A): ccgccttccccggcgcccatg (Seq ID No: 890)

- 5 Homo sapiens FK506 binding protein 10, 65 kDa (FKBP10):
agttctttgtagtgcctccctcagactctaacacactcagcctggccccctcctctattgcaacc
ccctcccccgctcctcccgccaggccagctcagttctccagccccattccacgtggaccagcc
agggcggggtagggaaagaggacaggaagagggggagccagttctgggaggcgggggaaggagg
ttggtggcgactccctcgctcgccctcactgccggcggtcccaactccaggcaccatg
10 (Seq ID No: 891)

Homo sapiens small ArfGAP 1 (SMAP1):
cctcctcccggttccagctgccgctgccgcttctgggctgagtcgccccgcggtcccgcgggcgcc
aggtgcgttccactctgcccggtccagccagcgctccgcccgcgtagctgccccaggctccccg
ccccgctgccgagatg (Seq ID No: 892)

- 15 Homo sapiens chromosome 14 open reading frame 93 (C14orf93):
cctcctttttgcacacacacgaatacaaaagagccatacagacttcggatgccggaaggctccttctg
aatcccttccctgttcccttaggttgcaactagtcgggggttccatgctggggggcagaaggatgct
ctctaccgtctgaaaccgttccatcaggaaggccttgatttgtgatgtgctaggagagcacaggatc
tgcaaatagaaggcacctgtctcccttctgcaggccgaggagaggccgcatggactgtgtgcttc
20 ttcattggcttgtttactcttctttcacagaccctacagcttggggcctgggctcctctgaccatcc
tcattgagaaaggaaagtgagtcagagaagttgatgcttccctacctgttgagcgggccagcagt
gtaagcgtgggttgttactgccccatccgcatg (Seq ID No: 893)

- Homo sapiens brevican (BCAN):
cgccctcttccgaatgtcctgcggccccagcctctcctcacgctcgcgcagttctccgccgagttct
25 cagctgcagctgcaggactgagccgtgcacccggaggagacccccggaggaggcgacaaacttcgc
agtgcgcgacccaacccagccttgggtagcctgcagcatg (Seq ID No: 894)

- Homo sapiens H2.0-like homeobox (HLX):
cggcctctcttctcagtgccggcgaggagaagcgaaagcggatcgctcctcggtgccgcccgttct
ccgggactcgcgcgccccctccccgcgcgccacccacccagtcgggtggactgcggcagccgcgc
30 ggctcaccgccgagcatg (Seq ID No: 895)

Homo sapiens v-rel reticuloendotheliosis viral oncogene homolog A
(avian) (RELA):
ccgcctctggcgaatggctcgtctgtagtgcacgccggggccagctgcgacccccggccccgccc
ccgggacccccggccatg (Seq ID No: 896)

- 35 Homo sapiens zinc finger protein 277 (ZNF277):
cctcccttttcttttctgccgggtaatg (Seq ID No: 897)

- Homo sapiens globoside alpha-1,3-N-acetylgalactosaminyltransferase
1 (GBGT1):
cttctcttttctgtctggccccgccccgctgcctgcctgctccaggctccacctgcgccgcc
40 gatgcccggtatcgcgggggccaggccagctgagtcctgttttccgcgcgggggtggcgccccct
ccaaccgtcctaacgcggggccggcagcaaggagtgttctgggacctcagagaccaggctcagag
cctgacatccctgcgaggggacagcctcatccgcccaggccagtggggtctctacaagtgccag
gctcaggtgcagccccagcaatg (Seq ID No: 898)

Homo sapiens FXYP domain containing ion transport regulator 6 (FXYP6): ggtcctcctgggagtcctcgaggaggaccggctgtgcagacgccatg (Seq ID No: 899)

gccacctggcgggccgctccggtgcgccctgcccgcgcttttactgacaggcgctgttccccacag
ccagcgccgcccgcacgtcccagctctcggccaacggagctgcgcgcggtgacctttccgagc
ccagcgcgatg (Seq ID No: 909)

Homo sapiens stimulated by retinoic acid gene 6 homolog (mouse)
(STRA6):

5 ctaccctttcatctctgcaactccttccctgggcctcccttctggtgtgtctgtgggtctgtc
taggtgggcttgaggaaaggggaaggaagggcgctctctttaggcagctcagactggacaagccttc
tttgaatggtcctttgaacacacgcctgctggtggttggtcagacagatgcgccagcgaggacc
ccggggccccaaggggacagctatctctgcaggaccagtgcgatg (Seq ID No: 910)

10 Homo sapiens 5-azacytidine induced 2 (AZI2):

cagccccctttccggctgagagctcatccacacttccaatcactttccggagtgttccccctccct
ccggcccgtgtggtcccgacggcgggcctgggtctcgcgcgctattgctgggtaacgggccttc
tctcgctcgggcccgccccctcctgcctcggtcgtccctcctccagaacgtcccgggctcctgc
cgagtcagaagaaatgggactccctccgcgacgtgcccggagcagctcccttcgctgtggaagcgg
15 cgggtgtcttcgaagaaaccggaagcccggtggtgaccttggcgacctgggttgttttcgggtcgtt
tccaaacactaaggaatcgaaactcggcgcccttggggcgccctacgtagcctggcttctggtt
gtcatg (Seq ID No: 911)

Homo sapiens polymerase (RNA) I polypeptide E, 53kDa (POLR1E):

20 acgccttttccggcccgcagcgcgccctgggctcccgcggtgtttaaagtgcgcttgtggctgctg
ctgtcttaactcctgtgcttggcgacagacaggcgagatg (Seq ID No: 912)

Homo sapiens mitochondrial ribosomal protein S25 (MRPS25):

agtcctttctcgtcgtgctcggtcgcggcccggtggggctggccccgccaccgttgccgcatg
(Seq ID No: 913)

Homo sapiens TRM2 tRNA methyltransferase 2 homolog A

25 (S. cerevisiae) (TRMT2A):
cggcctccgcccgcacgcgctggcggaactaagagtggctggcgaagcgagcgggcgcgggcccc
tggcgggcgggcggtacagcccaagcctgagaccggacctgagcatcgaggttcgagtcgccgc
ccgcctggggcgaagccgggggtggcgggcagctcgcggcgttgaccggctctgtgagcacctc
ccctctgagcacttcccttgtgacaggccacttcccttgtgacaggcccaggacgaggtggccagg
30 cggcccccatggcgctccctggtctaggcggaagaaccgcctgggcgatg (Seq ID No: 914)

Homo sapiens lipid phosphate phosphatase-related protein type 2
(LPPR2):

35 ccctccctccacctcgaggtctgcgcgcgcgccaggccccggccgaccgcgtctcggtcttcgcg
tctgccagcctggctggcagtcctgtctgtccatcccgcgcgccccgggagcttaggcggagcggg
ggctcaggcgggcgggcctcgacgcgagtgagtgctcggtggttgggtgctggaccagagtgccct
accctcgctgcttggcctcagtttccacatctgcacaatgggggtgaccatccctgcctgctg
gctgccaggagcggctgtgagtccttcaggcgtggatgcagcctgggggaagccatagggcgcttcc
acaggcctggccttcaccatg (Seq ID No: 915)

Homo sapiens chromosome 11 open reading frame 1 (C11orf1):

40 gaacctttttcacctcgtctgaaatg (Seq ID No: 916)

Homo sapiens microtubule associated monooxygenase, calponin and LIM
domain containing 1 (MICAL1):

cgccctcccaccgctcagacctggttgccagcccaacaggaagcgccctcccggcttcggagc
cgccgccactcatctctgccagctgctgccctccccaggaggcctccatg
45 (Seq ID No: 917)

Homo sapiens kinesin light chain 2 (KLC2):

gctcctttaaggcagcgaacgggccaagagaagcgtgtttcgccccctccgacgccaccgaggtag
cggttcacctttaaggcggcgcgggggtgctgggaaggccggcgggatggaggcggcgggaccg
gctcgcggtgctgggtccgggtgaagcgggaggcagccagagtcggagccgggcccagaccagg
5 cgcaggccccggcgcccgcctgcccgcaccctcgtcctcacagacgccacagccatg
(Seq ID No: 918)

Homo sapiens DNA cross-link repair 1B (DCLRE1B):

acttcctttttctgcccactctggtaacttattgctctgctgggctctttcccttagggctctctgg
ccctgttcttgccccagcatgacttttatcgggacgccgttggtggaagcctcacgcaggagccctg
10 ccccgctggagaagatcccaactggtgactccaaccctaccaccatg (Seq ID No: 919)

Homo sapiens armadillo repeat containing, X-linked 5 (ARMCX5):

gctcctcccactgccgttggtggtaacgcggacgtggaagaacctcgtctgctggaggaaaaggtag
atgttaaattgtaactacgcgcgaggttctgaggagccctgggaacaggaaggagaaaagaatacc
aaaagtgacaacagtttgccaatcgagtccttaattctgataaagcggttatctcgtcttgagtcc
15 caggtgcccagagtcaatccccatacacagccgcgccattgacctcagagtccttggtgctgactgtct
gttctgctgctgtatgacacagcacctcgaggcaaggaaataagaaaactgcctctgatccaagc
agagaaggctcgcctgtagatctgctgtagggcttgctcaccattggaagcaaggctcctacttcagt
ggcagatctggtggccttgagtggtgaagaccaccaccctccacagggtgggcccctgcacag
ccatccttccctaccttgagtgagcttcctctgcatgttttctatatcactggcagagcctgtagt
20 tggaaaggggacagagtgactactggactttgtgtgaaaacaccaaccgggacaaaacttcagtc
aggctgagacgggtgggggtatataacttgctccttacgttaaacttggaacatg
(Seq ID No: 920)

Homo sapiens chromosome 12 open reading frame 43 (C12orf43):

aatcctttgcggtggttcaagatg (Seq ID No: 921)

25 Homo sapiens vacuolar protein sorting 33 homolog A (S. cerevisiae)
(VPS33A):

ggtcctcccgtaggaaccggcgactcggttggcgttggtggggcaggggggtggtggagcaagatg
(Seq ID No: 922)

Homo sapiens arginine/serine-rich coiled-coil 2 (RSRC2):

30 gggcctcctcgcccttctgcccacccgggtctctcgcgcgagcgatttagtctgaggcgaagcttcg
gagcgggcggtactgttgaaagcgacaagtggaggcgccgctctagcgggccgggactctgaactat
ggcggctagtgatacagagcgagatggactagccccagaaaagacatcaccagatagagataagaa
aaaagagcagtcagaagtatctgtttctcctagagcttcaaaacatcattattcaagatcacgac
aaggtcaagagaaagaaaacgaaagtcaagataatgaaggaagaaaacacaggagccggagcagaag
35 caaagagcgtgcttatgcggaagagactgaactgaagacgctgcagactcagatagcaaaataat
aagcctacttcatgataagggaagaagacatgaatccaaagataaatcctctaagaaacataagtc
tgaggaacataatgacaaagaacattcttctgataaaggaagagagcgactaaattcatctgaaaa
tggtgaggacaggcacaacgcaaagaaagaaagtcacaaagaggcagaagtcactcaagatctag
gtctcgtgaaagacgccatcgtagtagaagcaggagcggaagaagtctcgatccaggagtaggga
40 gcggaagaaatcgagatccagaagcagagagaggaagaaatcgagatccagaagcagggaagaaa
acggcggatcaggctcgttcccgtcaagatcaagacacaggcataggactagaagcaggagtag
gacaaggagtaggagtcgagatagaagaagagaattgaaaagccgagaagatttagcagaagttt
aagccggactccaagtccacctcccttcagaggcagaaaacacagcaatg (Seq ID No: 923)

Homo sapiens integrator complex subunit 3 (INTS3):

45 ccgccttccccccccccgccccttccactatggccgcttctgtgtggtgtggggagacgctggctct
ccccgtcctcccatagcgcttattgcctcaccctcaccctcctaggggcccggatccaaaggcgctgc
actccccaagccttggggcatcagccaggaagggttctacctcctaattcaggggagagactcct

cttttccccccacggggaaaagaggcagaaacttaggggtttccctcctttcttaggggtcagacgc
tcttaggggtccacttcttcaggggcggaagcctctcctacccttcccataggggacagggccttta
ccccactgtacttcggagccaacgcctttccctcagcactgccacccagagtcaggaccagagg
actgtgccttcgcccccaacgcaggcgcggtcttttgagaggaggaggagtgaggagacaggg
5 gcccttgctctccctccccaacttgttcctcttgccccccagtccttggaatccagagatcccg
atatctaggactgtccatccatccactccctgaccttttcccggctcctgggtgcagccatg
(Seq ID No: 924)

Homo sapiens spermatogenesis associated, serine-rich 2 (SPATS2):
tctcctttcctcttctcagacccgggagcgctccgggacgcggagcccgagctggggcgacgaggc
10 gattgcgggggctgggctagctgctggctaccaatattctactttctgtctctatgaatgtgact
accctggttacctcatataatctccctggaaaaggagacatgaatgtctgcaatgatacttctga
caagaagttgatacaagaaaaggaaaggagattaacagctagtgcagaaatttcgaacagcagga
tttcgtattttttgcttccaactgcacacttccgttgccacttttaaatcagagatacctacact
caaaaccagacaaggcaaaaggatacttttcttgatatatttttgagatcgaagaaacgacaatg
15 (Seq ID No: 925)

Homo sapiens fibroblast growth factor receptor 1 (FGFR1):
ccgccccctttcacctcctggctccctccccgggagatccgcgcccccttggggtctcccctcccttccc
tccgtccgctctcctgcgccccctccctgcgctcgtcccgccgctcttcccgcgcccactttt
cctccaactcgcgctcgggagctggcgaggcgggcggtcctcaggtcagtttgaaaaggagga
20 tcgagctcactgtggagtatccatggagatgtggagccttgtcaccaacctctaactgcagaactg
ggatg (Seq ID No: 926)

Homo sapiens FUN14 domain containing 2 (FUNDC2):
ctccctcttccgctgccgcccgtgggaatg (Seq ID No: 927)

Homo sapiens ganglioside induced differentiation associated protei
25 n 1-like 1 (GDAP1L1): cctccttctttcctgcctctgattccgggctgtcatg
(Seq ID No: 928)

Homo sapiens chromosome 19 open reading frame 43 (C19orf43):
agtcctttgcgcggcacctggcgacaaaatg (Seq ID No: 929)

Homo sapiens MIS12, MIND kinetochore complex component, homolog
30 (S. pombe) (MIS12):
ccctctcttctccaccagccaacgtccgggaaaaaacgagtaagtacagggttccttctgccaatccc
cgccggccacagctaactttcccgcccgccctttctgtcataattgaggtgtccacaaccagcc
aatcaggaacgcgagagtatcccgcttttgcttctcgtcgccgaggcgcgatcagtcggaatttt
ggggagccaaccgcgcgtctgtccctggcaagccagcgggcggtttaaaggaggtggcggggaagcc
35 tgtgtgtgcttcaaactcgtcacctcatggctcgtccggtaagtgtcggggcagcattttctct
gaggaggagcggggacgggcgagactggcataagcgtcttcgcgaggagcaaggcggcctgtggg
tcggcctcaccgccctccgacctgaagatcccagcatgcagcgcgggcgggggcccgacggaa
gccgggagccggccggaagcagttcctgcgctctggcttctgggtcctgtcctgcgcgatcgcggg
gtcttagacagctcaactcgccgagatgacctgggcacctctgcgttgaaatcggaataactgatc
40 aagccgcatttattctgtctcaggaactctaagtctagcagagaagatgaggcggtagaagttca
tcaatggccttggtggaggacaagcaaattgaggacattggcaacggagtgatcaaaatgatagat
catgaggcctaaaatgaataaggaaagaagagaagtggcagaggctgagaacagaaagagaggggtg
gaggggctgtaaatcttgaagattaggggtataatatgagtatatgggtaagaattggaagaattgt
gtaggaggcagtagtcaaaaagtagaagcagtttggaagagtagttacaaatatcaagagccaggt
45 ggctaaaagggtggagctataggtcattgaagctcaagaaactgagtctctagggcatttggttaagt
catctgtctagacttcaaagttgtctaggatgataattcagaagactgatctgtgccaaagtcaca
ggtttttccagactgaaaacaacatagcaaaataagccaagatg (Seq ID No: 930)

Homo sapiens DEAD (Asp-Glu-Ala-Asp) box polypeptide 50 (DDX50):
cttcctttcacgctgctcgctgcccgtaggtggttgtggccactgtgcccggagggaggcggcggtg
gccagtaatg (Seq ID No: 931)

5 Homo sapiens chromosome 7 open reading frame 25 (C7orf25):
cggcctctgctgcacgcgcctgctgctcgcgctcgcggttctggcgctgccggaataatgctga
cagcatg (Seq ID No: 932)

Homo sapiens KxDL motif containing 1 (KXD1):
ccgccctttcctgctgctgacttaacgcacgcaagcggctccagggtacgtccccgccacgcgcgct
cgcaggatcggtgctggtgacgtttcgccggcgcgggcgccatcccgaagcgcgagcaaggccg
10 ccagatgtgcaggcagcggaggaggagaaagagatg (Seq ID No: 933)

Homo sapiens defective in sister chromatid cohesion 1 homolog
(S. cerevisiae) (DSCC1):
acttctttcttgcccgcgaagcccgagccaccggcgcgcgggactcctagaccggcgctgc
gatg (Seq ID No: 934)

15 Homo sapiens zinc finger protein 426 (ZNF426):
cgttccttttgtagcgcggctgtgagcgcctgagagtccttttgcccttcagagttaaggcctca
ctggcctgggaaaataattgctgccttttgcatccgcgcttggtccgtccccaggatcttcccgt
tcaggacctggcgatttctgagtggtccggaatccaataaccctgtttaagagggaatggagat
tgccactgtccatttagattaatgaggtgtcctgaagtgatggtgacatcaatgaaaggagggttc
20 tgacacgttctcacctcgcggtg (Seq ID No: 935)

Homo sapiens TATA box binding protein
(TBP)-associated factor, RNA polymerase I, D, 41kDa (TAF1D):
caacccttttcttcgcacggttgaggagggtcggtggttatcgagggttgaggggctgaggtcg
ggagggtggtgtgtacagagctctaggacaccaggccagtcgcgggttttgggccgaggcctgggt
25 tacaagcagcaagtgcgcggttggggccactgcgaggccggttttagaaaactgtttaaaacaaga
gcaattgatg (Seq ID No: 936)

Homo sapiens PHD finger protein 1 (PHF1):
ccgcctcctcctcctgcgcgtgcccgtgctttggctgctgctcatagccccagagccgcccggga
cggaggggctgggctggggacccccggcctcgccctgcacgccccccacgcccggacgtgccc
30 tctccgcgcgggggactcgcctaggtctcctacgtctgcccctgcccggctcccggcgggcccccagc
tgtcaccggccccccaggatgcaatg (Seq ID No: 937)

Homo sapiens family with sequence similarity 134, member A
(FAM134A):
cccccttcgcctgacgcgccccggcgggcgccgcgcagccctggctcctcgcgggctcgggcgg
35 cggctgcggcggggctatg (Seq ID No: 938)

Homo sapiens membrane bound O-acyltransferase domain containing 7
(MBOAT7):
ccgcctcctttccggagcccgtctgttccccttcgggtccaaagcttttggtcctccttgttccg
agcccgaaggcccgcccttcacgtactcggagctcggatcccagtggtgacctggactcgaatcc
40 cgttgccgactcgcgctctcggcttctgctccggggttcttccctgcccggccggggccctgacc
gtggcttcttccccggcctgatctgcgcagcccggcgggcgcccagaaggagcaggcgggcgggg
gcgcgctgggcgggggaggcggtggccggagctgcggcggaagcgggctgggactgctcgggccgc
tcctgcccggcgagcagctcagaccatg (Seq ID No: 939)

45 Homo sapiens major facilitator superfamily domain containing 11
(MFSD11):

acgcccccttttttgcctcagccgtcagccccgtctccgtctgaagagtgccttctgccctcatttgcc
tctccctgtgaccccgccccctcagactccgctgcgtcgtctctcgccccgtccagccgttcct
gactgctcttcgcccggagtccgcttcccaacccccctttcgccagagccccgagagctccgtcggctc
tgctcctggcggattgtcagtggcttcgccccgaggagagctgactgccctgggctgctgcctcc
5 ggcagagctgagccaaatg (Seq ID No: 940)

Homo sapiens thiamine triphosphatase (THTPA):

ctcccttccccctctgtgggtcccgcgaggagactctcgggctttgaggtgagacctgaagttccg
ctggccggtagtgtagcaggaaagggcaggtcctccgggtcgtgagccagtagcctcctgggggtg
gcaaggtgtagagagggggcggttgaaaggacaccgcctaccggcctgctttctaggggtctctt
10 tggattgaggacatcagcagcagtgaagggattttactggagacctgtcactgtcagagccttaa
aatatcaccgacggggccttaattgtcaccgaggtagagagaaaagggcagtagccctagagactat
tgcgacacagtgtgccccctcataagttttccaggagggttctgtactgagttgacgccccagg
agctgagcaccaggctttgcctccttggaactcagcaaacgtttgttcagccaattgcaggtagc
atg (Seq ID No: 941)

15 Homo sapiens acyl-CoA synthetase short-chain family member 3
(ACSS3):

tactcccttccctcaggccccaggaagttgcaagagtaccatttgtcgcacactcggggaccgcgg
gtggccggaggagatg (Seq ID No: 942)

Homo sapiens chromosome 6 open reading frame 211 (C6orf211):

20 gctcctccttcgcgcggtaccgcctctgtttctgcggcgattgaacagccgagctttgcggccgg
gatcgcggaagtgatg (Seq ID No: 943)

Homo sapiens transmembrane protein 204 (TMEM204):

atttcctctctgctgagagccagggaagggcagctctgcgcacacgggctccctgcagcagccac
tctgctttccaggaccggccaactgccttgaggcatccacacagggggccaggcagcagagga
25 gctgtgaacccgctccacaccggccaccctgccggagcctggcactcacagcaggccggtgctaa
ggagtgtggcgcgggctcgactcccactgctgcggcctcccgagtgactctgttttccactgctg
caggcgagaagaggcacgcgcggcacaggccggcctccgcttcccggaagacggcgactcctgg
ccctgggttcttctgctgctgccaccctctgctccctgggatgggccccgaggcgagcagcttcagc
acaggcctggccctgctccaggtgcaggaaggaggataaggccgggcccagaggcgccacacctgg
30 accatcccatgggcctccgcccgcgcgccccgaggatgagtggtgatgtcctctagccaccctta
gcagcgtcggctctccctggacgtgcggccgcggactgggacttggctttctccggataagcggcg
gcaccggcgctcagcgatg (Seq ID No: 944)

Homo sapiens DEAH (Asp-Glu-Ala-His) box polypeptide 40 (DHX40):

35 tcgtctttccctcccatctcctcagatcggtggacgtgctgcctccactcggggccaggtctat
g (Seq ID No: 945)

Homo sapiens importin 4 (IPO4): cctcccccttttcggcccagtagcggcggtcag-
ttgctgccatg (Seq ID No: 946)

Homo sapiens N-acetyltransferase 10 (GCN5-related) (NAT10):

40 ccttctctttcggagttgttccgtgctccacgtgcttcccccttctccactggctgggatcccccg
ggctcggggcgagtaataattttcaccatg (Seq ID No: 947)

Homo sapiens lin-28 homolog A (C. elegans) (LIN28A):

aaccctttgccttcggacttctccggggccagcagccgccccgaccagggggccggggccacgggt
cagccgacgaccatg (Seq ID No: 948)

45 Homo sapiens CAP-GLY domain containing linker protein family, memb
er 4 (CLIP4):

cggcctttcctccgcgcccccgctccccagccggccgctccgagaggacccggaggaggcaggtg
gctttctagaagatg (Seq ID No: 949)

Homo sapiens zinc finger, AN1-type domain 1 (ZFAND1):
ccgcccccttacggcgccggagagatg (Seq ID No: 950)

5 Homo sapiens GTPase, IMAP family member 6 (GIMAP6):
cctcccttttttctacttccgaggctgcaaagtgaacacgcagactcttctgactcaggaaggccgg
tgctcctaccacttccctgttcctccatctccagcggacactgctctttcaagggcaggtctccag
cccagctctctgaaaacattttgctgaaaatataagcaaacatcggccttgtcctccttgtgttca
10 taaactgtggaagcttttctctgcctcctccgtgagagtgcgtggccgggagaccagaaacgtggt
cctttctcttgctgtgagctggtgcagagatg (Seq ID No: 951)

Homo sapiens thioredoxin domain containing 15 (TXNDC15):
cttcctccggctggcagcagcactcgcgtagccgtgcgccgattgcctctcggcctgggcaatg
(Seq ID No: 952)

15 Homo sapiens asparagine-linked glycosylation 9, alpha-1,2-mannosyl
transferase homolog (S. cerevisiae) (ALG9):
aattcttttttccccaggcttgccatg (Seq ID No: 953)

Homo sapiens glutathione S-transferase, C-terminal domain containi
ng (GSTCD):
20 acttccctttttccgggtccgcccggattatgaatgacggccggcgcgagtattttccacataaggtg
gctgtcggtttttctcctggcgtctgtggaggcgagtggtctgcgggcagcagctcccagaggcagc
cttgaattccagctcggactgggcccgggaaggcgagggcgccaggtcgccgacacgctcacgca
ccctccctgctggccgcgctctgcgaccaggtgaccaatgaaagaagaaaatg
(Seq ID No: 954)

Homo sapiens CXADR-like membrane protein (CLMP):
25 actccttttttctttccaaacagggaaaagtgttccacgaagcggtagcgcctttccgcctcgcgtt
ttcctccctgaccctgggtcccggctcccgtccgggcccagctggtggggcgagcgccgggagccc
atctgccccaggggcacggggcgcggggcccggctcccggccggcacatggctgcagccacctcgc
gcgacccccagggcgccgcgcccagctcgcccaggtccgtcggaggcgccggcgcccgccggagc
caagcagcagctgagcggggaagcgcccgcgtccggggatcgggatg (Seq ID No: 955)

30 Homo sapiens nonhomologous end-joining factor 1 (NHEJ1):
cctcctcttgcggtgggggaaagcggcctcttactctaggcctttcggtttgcgcgagcgggagc
gaaagcgtgcgtgcccgaagagagtgggcgctctcgcggccgctgacgatg
(Seq ID No: 956)

Homo sapiens gametogenetin binding protein 2 (GNBP2):
35 cctccttcttccactccccgcggcgagcggctgactgccgtagaggaaacgacattcggagct
gcgctcccggccaggccggccctgacgcgggctcgtcagccagtaacaggagcagaggtgggag
ttagcgaggcgaccacgaaaacggtgaaggtcggaaccgacagcctcctccgagaagggcaggagc
tgggaggaggcggcagcggcgggcgagaaacagcagcggcgggcgggcgggcagctgggaggagg
tggtgacggtggcaacggcagcgtcggggacgatg (Seq ID No: 957)

40 Homo sapiens zinc finger protein 672 (ZNF672):
ctttctcttttagccccgcctgcttcccggctccagctggggccggagaggctgagtgggttggtac
gctgctcgctggcctcccagcttcccagcaaccggtgacactgccgcgccagactgaccactag
ccgacgcgggagaggggacaggagcgtgacctcccatcccgaggggcccggacgctcggggcgct
ccccgctccccccactcggaggccgcgcgcgcttagcccccttctcgtctccccgccccagttcc
45 cgcagtcggggaggcgggggtcggcagccggctgagtgggaaccgcgcggtgtctgaggaggcagt

- cggcgaccggtttccacttcaagcgtgacccttttgccctgtgggatgagctccagcatgggggtgag
gtacagaagagagacttgaagagcgtgccttgggactcaagcgccaaacctgtaccctagcgagtg
tcctactccgcatccgtaatggaaggaaatgcacatcttactccagaggcacaagaggaggacatc
ccatgcggctactcctgcccagcgtggtggggcagcagaagctccagagcccagacttgagggtc
5 acggtgcagggtgaacctggccacagctcaccctggaacagccacaatgtctgccccttagagaag
aacctgaaatcagaccagtttttgcggcctccccctttcctctctgttacagtgccctttccagg
ccttaagagaagtaaaacttagctgcagcgccaggaggtggacccagagtgtagtggtggcagcgtt
ccctgtgaacccgtcctcaccatg (Seq ID No: 958)
- Homo sapiens N(alpha)-acetyltransferase 60, NatF catalytic subunit
10 (NAA60):
ccgcctccgtcccggctgcggccccctgcgggttacataactcgttgcgggctccgcgcggtccac
ttcccggctcccttcgcctccaggatgcgctgagccctacaacacccccagcgggccgcgggtccc
ccacgaggtgtgaatg (Seq ID No: 959)
- Homo sapiens transcription elongation factor A (SII)-like 4
15 (TCEAL4):
tgccctctgtccccgcgggtggtctcgtctgctccggttccctgggctcctaattcttgggtccagc
ttcttccagggtcagtgtgcgggcttccacgctgccagcggaacactggaatggcggaaggggaac
gggtctgcgcgtctgttgttcccagcgtctgcgaagcctgaaaaggaggagcaacctgtccagaa
tccccgcaggacaggaaaaggaggggaaatctcgacatg (Seq ID No: 960)
- Homo sapiens progesterone and adiponectin receptor family member VI
20 (PAQR6):
tcccccttctgttccccactccccgcccaggcctggcccgcctgcctggccactcttccctccatcag
cctggctggcagcagccttggactccgcccgtggagccctgggctgttgaccaccagcttagga
gcaccaccaagctctgggtaagggaagctcaccttctggggctcttctgggaaaatagagggtcaac
25 gtggaggtaccaggccaccatgctcagttctcaagctgcccgaacttcttcaagtccaccaggtccc
ccgggtgttctgggaagatggcatcatgtctggctaccgcccaccagctcggttctggactg
tgtctcagctccttccagatgaccaacgagacgggtcaacatctggactcacttccctgcccacctg
gtgaggggaggtctgccccaggccgcggccttgagctcagaggggtaccaggcgggcagggac
cgtccaggcccacgggctgcagcggcagtcgcgggggtccgcggcggcctgagcacgcgcccgcg
30 caggtacttccctgtggcggtcctggcgctggcgggcgccccggccttccgtgaggagccgtacca
ctggccgctgctggtcttccctgctgcccgcctgcctctaccccttcgcgtcgtgctgcgcgcacac
cttcagctccatg (Seq ID No: 961)
- Homo sapiens DENN/MADD domain containing 2D (DENND2D):
catccttcttctgctcaaccactgggtgcacaggatggaaacttctattccctctctggaagacagcg
35 cgtggcttggcttcacagagttgtggctggagaccgaagcagcccccttctcaggcttactgtcac
cagttctgtctgtttaggggagaggggagtcgctctgtcctgaaggccagagatg
(Seq ID No: 962)
- Homo sapiens family with sequence similarity 188, member A
(FAM188A):
40 ccttcttcttctcctgcctcaccttccaattcgtttgcccgcgcggtcccgagctgctgtttccgg
agttgcccttccccatgttccggggcaggagtcgcaaaagcgaagatccgcccgcgggttccctca
tcatg (Seq ID No: 963)
- Homo sapiens neurensin 2 (NRSN2):
ccgccttctgctcggcgagacagcaggcagagagatgaggaaactgagaccagaaaaggtggaagc
45 acttgtctaaggtcacgcctccaggaagcagtggtgtccacgactccagtccaagtgggtcaggctcc
agagcccacagttcccaggggtccatg (Seq ID No: 964)

Homo sapiens tripartite motif containing 46 (TRIM46):

agccctcctcacacccccactgggctcctgcattaagcccggggttcgcagccgcagccgggatcg
ggcaccacaggggcgggcgggcacggtagggccatg (Seq ID No: 965)

Homo sapiens target of EGR1, member 1 (nuclear) (TOE1):

5 catcctctctgggaattttaccgatgcccagaacgcccttctttccccacacgaccctctcctagt
ctaactcctgggctgctttaagctcagctcaggcagcgtcaccttctctggaagcccaaacc
gccacccactacccgctacccgcggccacgctgatgaagacagcagaacacggaggccccgcgt
tccgcgcgcgagagcaggagagaaagattacctccgcgagctctagcgcgcccggctttccggcg
cactccagggggcgtggctcgggtccacccgggctgcgagccggcagcacaggccaataggcaatt
10 agcgcgcgccaggctgccttccccgcgcggaccgggacgtctgaacggaagttcgacccatcgg
cgacccgacggcgagacccccgcccatccccgactgcctgaaccgcgccaggagacggaccgcaag
tccagcgtacccacagacgactcaggcgggagacgagcgggtgtcatg (Seq ID No: 966)

Homo sapiens DBF4 homolog B (S. cerevisiae) (DBF4B):

15 cgttcttttaggggtggagccggcaggaaatttaaactgaagccgcggccgaaaacgccaagagat
tgatgctgtagctgccctgagataaccaggactgtggaatcggaagagctcatggagctcgcgaa
tgtaatacggaggcctctgaggaaggagtacggaggccgagaaggagccggcatttgatg
(Seq ID No: 967)

Homo sapiens myc target 1 (MYCT1): atttccttttatg (Seq ID No: 968)

Homo sapiens myosin XIX (MYO19):

20 ggttcctttcctcactgcacgctcttgccccctctcttttctctcctgccgctgttcttccgcgcg
cctgacctggcccgcccgctttccagctctggccgggcggggctgaagcacggcggtcggggc
gtgggaccgtgttcacaccctttccagaaattcttggttgtaaccgcgaaaccgactggagcagg
agctgggagaactggagaaaactgctctaattctacttgactccagctaggagctgatgtgtcatc
gtaataacatttgagagcgctttcacaggcgctggagtgtgtgtgagattcctccagaactg
25 agccctttgttggaaccataccccagcccatgggtcccatgactagggtggatagtactccttgatc
tcctgcaaccacagaaccctggctgaccactttgaaggaggatg (Seq ID No: 969)

Homo sapiens KIAA0226-like (KIAA0226L):

30 cctccccctttctgctgttaccgggagcgcggtggccacggaacgctgcccgagccgcgcgagggg
ggacccgacgcgcggcgtttaccagcgcagcgttccaccgctcgggtttggctggataaaataaa
aaatggggatattgacctcctgtcactactgcattgactttgatgggtttccaatcattactttctc
ctctgtgtcaatctgcctcttcgagaaattcatactcctgaatagctctccagacccccagctggc
catgtggtgagttcagggcccaaatcaagtagtaccagcaatcagggaaactcctatctgttttgaa
tggaattcacaccagccacaagcctggaaagatg (Seq ID No: 970)

Homo sapiens MUS81 endonuclease homolog (S. cerevisiae) (MUS81):

35 ctccctcttcccccgccccgcctgggcccaggtgttcgaatcccgactccagaactggcggcgctcc
cagtcgccgcgggcgtggagcgcggaggaccgcctcgggctcatg (Seq ID No: 971)

Homo sapiens zinc finger protein 430 (ZNF430):

40 gggcctttgtccctcgctgtggcctgagctccaggtctcgtcttcagcgtctgtgtcctctgctc
ctagagggtccaggtctgtggccctgtgaccgcaggtattgggagatctacagctaagacgccag
gaaccctggaagcctagaaatg (Seq ID No: 972)

Homo sapiens mutS homolog 5 (E. coli) (MSH5):

gctccttttgaggctcgtggcggtcgggtcagcggggcggttctccacctgtagcgtcaggtta
ctgaaaaggcgggaaaacgctgcgatggcggcagctgggggaggaggaagataagcgcgtgaggct
ggggctcctggcgctgggttggcagaggcagagacataagacgtgcacgactcggccacagggccc
45 tcagacccttcttccaaaggagcctccaagctcatg (Seq ID No: 973)

Homo sapiens proline rich 3 (PRR3): gcccttcctcac-
taccctccaaatcccgtgcagccattgccgcagacacgatg (Seq ID No: 974)

Homo sapiens sirtuin 2 (SIRT2):

cgccctttaccaacatggctgctgacgccacgccttctgggactcgtagtccggtcctcgcgcgct
5 ttcttacctaactggggcgctctgggtgtgtacgaaagcgctctgcggccgaatgtctgctga
gagttgtagttctgtgccctatcacggccactccatttctggtgccgtcacgggacagagcagtc
ggtgacaggacagagcagtcggtgacgggacacagtggttggtgacgggacagagcggtcggtgac
agcctcaagggcttcagcaccgcgcccatggcagagccagaccgactcagattcagactctgaggg
aggagccgctggtggagaagcagacatg (Seq ID No: 975)

10 Homo sapiens KIAA1715 (KIAA1715):

ttgtctctctgtcagtgggcggtgctgcctgctctggaggcaggctgggcggtggcgccgagact
ggcggggggtggacgcccgggccgggctgcgcccgttcttgacgtgtgaattcctttggacaatt
gatgatatttatcattgtgccagtttctacaaataaaagatg (Seq ID No: 976)

Homo sapiens proline-rich transmembrane protein 1 (PRRT1):

15 ctgccttcatctctccatctctgcgctgctgcggctgcgccatccagcaccagactccagcacc
ggccgaggacccccactccggctgcagggaccctgtccagcgagaccgcaggcatg
(Seq ID No: 977)

Homo sapiens t-complex 1 (TCPl):

20 ccgcccccttccccggagcctcacttccgtcacagtcctgtttctctccctggtgtccctgcctctt
tttcttccccgcccgtgccccgcggccggggccggggcagccgggaagcgggtgggggtggtgtgttac
ccagtagctcctgggacatcgctcgggtacgtccacgccgtcgcagccactgctgtggtcgccgg
tcggccgaggggcccgcgatactggttgcccgcgggtgtaagcagaattcgacgtgtatcgctgccgt
caagatg (Seq ID No: 978)

Homo sapiens Yip1 domain family, member 5 (YIPF5):

25 cgttctttggccctgtgacacgtagcaacggggctggttcagggtctgaaacagagtttgggggtt
gtttgggattagtgaaactactgcctttgccgccagcgcagcctcagagtttgattatttgcaatg
(Seq ID No: 979)

Homo sapiens glucose-fructose oxidoreductase domain containing 2
(GFOD2):

30 cctccctttccagagccccagttccttagaaaccaggcggcgcgttcccgggtggcggcgccttg
actccccggggcccgcgcatccccgccagccttcttaaggcggtgggtggcccccgagaccccgtc
ggacccatggtttccagtgacgcggagtgggcgatgccagcgtgccaggagccatgtctgacca
ggacgtttggaagatcatatccatgccagaggctcttgtaggagatgagttggttaaagagagagg
ctgggatg (Seq ID No: 980)

35 Homo sapiens apolipoprotein L, 2 (APOL2):

ttccctttcgaattccagggtatatctgggaggccggaggacgtgtctggttattacacagatgca
cagctggacgtgggatccacacagctcagaacagttggatcttgctcagtcctctgtcagaggaaga
tcccttggaacaagaggaccctgccttggtgtgagagtgaggaagaggaagctggaacgaggggta
aggaacaccttccagtcctggacagtactggagagctccaaggaaagcccctcggttaaccagccg
40 ctggcaccatg (Seq ID No: 981)

Homo sapiens microtubule-associated protein 4 (MAP4):

ccgcctccctgcgccccgcccctccggctagctcgctgggtcccggtcctcccgacgtctcctac
ctcctcacggctcttcccggcgctctcctggctcccttctgccccagctccgtctcggcggcgcg
ggcagttgcagtggtgcagaatg (Seq ID No: 982)

Homo sapiens exonuclease NEF-sp (LOC81691): cttccttctttgccaggca-gacgcccgtttagccgttgggaaccgttgagaatccgcatg (Seq ID No: 983)

Homo sapiens ST6 (alpha-
pha-N-acetyl-neuraminyl-2,3-beta-galactosyl-1,3)-N-acetylgalactosa
minide alpha-2,6-sialyltransferase 5 (ST6GALNAC5):
5 ctgtctctaatactctgcaacagccgcgcttcccggtcccgcggtcccgcgcgatctgccgcg
gccggctgctgggcaaaaatcagagccgcctccgccccattacccatcatggaaaccctccaggaa
aaagtggccccggacgcgcgagcctgaggattctgcacaaaagaggtgccaaaatg
(Seq ID No: 984)

10 Homo sapiens heterogeneous nuclear ribonucleoprotein A1 (HNRNPA1):
tgctcctttctgcccgtggacgccgcgaagaagcatcgtaaaagtctctcttcaccctgccgtca
tg (Seq ID No: 985)

Homo sapiens zinc finger protein 93 (ZNF93):
15 gggtcctttgtctctcgggtgcagccggagctccaggtctcctcttcactactctgtgtcctgtgct
cctacaggcccagcctctgtggccctgtgacctgcaggtattgggagatccacagctaagacacca
ggaccctggaagcctagaaatg (Seq ID No: 986)

Homo sapiens N-terminal EF-hand calcium binding protein 3 (NE-
CAB3):
20 cggcctctagccacaccgagtcggccgcggcggtccagggtcggcagcaaccgcagccgagcccag
cgggtggcggcgccatg (Seq ID No: 987)

Homo sapiens splicing factor 3b, subunit 5, 10kDa (SF3B5):
cattcttctgacgagcgcgacactggagcttccgcgcggtggcttcactctcctgtaaaacgcta
gagcggcgagttgttacctgcgtcctctgacctgagagcgaaggggaaagcggcgagatg
(Seq ID No: 988)

25 Homo sapiens INO80 complex subunit B (INO80B): gtcccctttcctcgcag-
gacctcatg (Seq ID No: 989)

Homo sapiens polyamine modulated factor 1 binding protein 1
(PMFBP1):
30 ctttcttctccttggcttatattagggataggggatgtggtttgttacaaaggatgagtatgttga
tagcttctcattccttgaactattctgcaggtttataacaaagctcagaaaataactaaagggttaa
ggagaattgagagctgccaaggaaatg (Seq ID No: 990)

Homo sapiens pseudouridylate synthase 3 (PUS3):
cttccttttctcgaaacgcgcgcggccggctgccggaaaacagggcagacctgtatggttcgttt
attcctgggggttgatcatatcatg (Seq ID No: 991)

35 Homo sapiens heterogeneous nuclear ribonucleoprotein D
(AU-rich element RNA binding protein 1, 37kDa) (HNRNPD):
tattcttttttagtgacagcgggagagagcgggagtgctgcgcgcgcgagagtgaggaggcgaagggg
gcaggccaggagagggcgcaggagcctttgcagccacgcgcgccttccctgtcttggtgtgcttc
gagaggtagagcgggcgcgcggcagcggcgggattactttgctgctagtttcgggttcgcggcagc
40 ggcggtgtagctctcggcggcagcggcggagacactagcactatg (Seq ID No: 992)

Homo sapiens GABA(A) receptor-associated protein like 1
(GABARAPL1):
atttctccatctggctctcctctacctccaggcagggtcacccgagatccccgccccgaaccccc
ctgcacactcgcccagcgcgtgttgccccggagcggacgtttctgcagctattctgagcacacct

tgacgtcggctgagggagcgggacaggggtcagcggcgaaggaggcaggccccgcgcggggatctcg
gaagccctgcggtgcatcatg (Seq ID No: 993)

Homo sapiens chromosome 22 open reading frame 13 (C22orf13):

5 ccttcctttccccagtggttgagcgcggtctcgcctccgcttcctcctcactccgcctgccggctgg
gaaactagggcaccagtacgatagttccggcaccggaaaagagggctgatgactgggccccggggc
cgccgcaacgacccttggggccggcaaagagccagagaggggtgctcacacttccaagcaccacaca
ccaaggacaggctggacggcaaggcggagacgcggggcttggggccctcagaccggggacagcagga
ggttggggccaagggccaggacttcccgtcacaatttcatattgttgatcccgccaccgccaggttaag
gggggccctgagtgaggctaggtatctggtacggataaagttaggtatagagtagagcggctgcc
10 gctcagggttatccctaaagacagttggaggagagttgcttggggccctcggggatgcactgggcgg
gatcagggttacacctaggactggcaaaagagcgggaccggcagaggcggggcttggcgaaggg
acgagcctctattcaggaaatgcacgagcttggggcggggctcaaagaaagggcggggcttccg
ggggccgcgtcctggtgagctgcgcgtctgcgcgaggattgggcgagaggggtggggccactcaacg
ctgaggcggcgcaatggccggagcagacttaaatcaagaggctggggacctctaagatcaaagtttg
15 gggcggggcctaaggagggggcggggcctccagattcgagacctggaagggtggggcggcgcttg
gggcggccctgccgcgcctccggttctccctccgcagcggcggtggcgagagaaggaaactcg
acacgcaccgaccgcccctcccgccccagccgaagcgggaagctgtagcccgtctggggccggggcca
tgggcgccccgcgcgcgcccgggtcatg (Seq ID No: 994)

Homo sapiens lon peptidase 2, peroxisomal (LONP2):

20 ggctctttttgacagccccagtgcgaaaggctgccagcatg (Seq ID No: 995)

Homo sapiens RNA binding motif protein 4B (RBM4B):

ggttctctctgacgtgggagccgcgctcgctgccgccaccggaggctcttgtcaggatg
(Seq ID No: 996)

Homo sapiens protocadherin alpha 3 (PCDHA3):

25 aggtcttttccacaaaagaataacagcgtgcattacgtattcagatactgctttgcttcctcct
ctctaaaatttaacaccgaggagttaagaaatgaagataaggaactcgaattatttttaaacttt
ggatcaatgtaaaggcaatctaataatttggaataacttgcaatg (Seq ID No: 997)

Homo sapiens RAB34, member RAS oncogene family (RAB34):

30 gcctctccttggggcccttctctcccccttccctccctgctggttcctggcatcgccagatgct
gcgagcagctctccgattccccatcaccaattcggtggcgtctccgagaccgcggactcccgtag
ggtccccgtggccccgagttgtagtcgggacaccccgccgcggtgatcgctgggtctccacgcg
ccgggtcgctgacgcggatccggcctcggcgccttctcagggcgccctgcaaggccgcaggcagg
atg (Seq ID No: 998)

Homo sapiens cell division cycle associated 7 (CDCA7):

35 gctcctcctgctgtgggaccgctgaccgcgcggctgctccgctctccccgctccaagcgccgatct
gggcaccgccaccagcatg (Seq ID No: 999)

Homo sapiens ArfGAP with GTPase domain, ankyrin repeat and PH domain 3 (AGAP3):

40 gggcttttttaggagagcactgctgcagccggcagtgagagcctggggcaggagacagggagaaaa
ctccggcagcagggtggtctctagggctgacctcgagcctggggacaggggagcctatgccgcac
tgaaggcgggacgctgtaagcaggagcagctgggcctgggcggactcctcgccaatcagcctcg
gtcagcagcaccctcaggcgcagggcactgtttgggcattgcctagagatccgacaccccgcccag
atcagcgcagggaggcgaaagcgacagccgggcgcgggaggagaccagggcagctgtcccctccgc
gaggttggccctcgaggcaatgcgggtgggggtggtgaggaggcggaaggccgaggctgagtgg
45 gaggggccccggcgccagggtggagcgcgcggctcgggggtggaggctgcagagccagcagcgga
gagggggcgggggcgccggggcgccggcgagggggcggggggaggggggctcggg
ctgcgtgtgccggagccggcgggggcgggcggtgctgcgcagtgacgcggggggagggcctgggccc

Homo sapiens cyclin B1 (CCNB1):

0 cattctctgcgaccggcagccgccaatgggaagggagtgagtgccacgaacaggccaataaggagg
gagcagtgcggggttttaaatctgaggctaggctggctcttctcggcgctgctgcggcggaacggctg
ttggtttctgctgggtgtaggtccttggctggctcgggcctccgggtgttctgcttctccccgctgag
ctgctgcctggtgaagaggaagccatg (Seq ID No: 1002)

Homo sapiens protocadherin gamma subfamily B, 7 (PCDHGB7):
cagcctctagcctgggattccctgcgcgagccaacaacagaaaaagaaaaccagctccacacagagg
ctccgggctgcgcgagaccttgcccagcacaccagattgccagctccgagaccgggactcctcctg
tcttggggccgaatgctcttttagcgcggtagagtgcactttctccaactggaaaagcggggaccca
gcgagaacccgagcggaacgatg (Seq ID No: 1004)

Homo sapiens zinc finger, CCHC domain containing 7 (ZCCHC7):
ccgtccctctacgcgttttggttcccggttggtgcttcctgttcgcagctgcggcacttcaaggtt
actgactttttatg (Seq ID No: 1006)

0 Homo sapiens forty-two-three domain containing 1 (FYTTD1):
cgctccctcggtagcgggcgggctgcgtagcgagtgaggaggcaggcctgcgactccggccttgt
ccgcgcccgcctctcggcgcgacgtctccagccatg (Seq ID No: 1008)

ctccctttctctcagcatcttcttggtagcctgcctgtaggtgaagaagcaccagcagcatccatg

gcctgtcttttggcttaacacttatctcctttggctttgacagcggacggaatagacctcagcagc
ggcgtggtgaggacttagctgggacctggaatcgtatcctcctgtgttttttcagactccttgga
attaaggaatgcaattctgccaccatg (Seq ID No: 1009)

5 Homo sapiens GTPase activating Rap/RanGAP domain-like 3 (GARNL3):
cagccctttttgcaaata (Seq ID No: 1010)

Homo sapiens DCN1, defective in cullin neddylation 1, domain conta
ining 5 (S. cerevisiae) (DCUN1D5):
gagcctcttgcttgctgtgactgggtggagctgccgcgctgtccgcggttatctcctcccgggtgagaa
cgaaccgcagtgctccaccggcgaggagccagccctgtcccggtcagagaaagacgacgaggatacc
10 tgggagcgggcgccggcgccgggctgggcccgcgcccgggtgcgggctggcgactctgctcctccgcttgc
tgctgtctctgggaactgggtgccagcgtgaggggcttcagcggacagggaccccttccccgg
ctccccctgccaccctgccggggaggcggaagatg (Seq ID No: 1011)

Homo sapiens alkB, alkylation repair homolog 7 (E. coli) (ALKBH7):
tgccctctctcatgaccccgctccgggattatg (Seq ID No: 1012)

15 Homo sapiens nitric oxide associated 1 (NOA1):
ccgccccctttggagctacttctcatg (Seq ID No: 1013)

Homo sapiens BTB (POZ) domain containing 10 (BTBD10):
tcgcctcttcgcattgtgagctctcgcggttaagaggctgaggagccggcctgcaacctgccggggc
ggctccgctacgcgcagccgcctcagtggttccctccacagccacctccggagggatctggctgag
20 gaggaagtggaggtgtcactggccccggcctttgccccaatcttgtgtgggactgaagggggact
acaggttcgagagttatgggtgctacatgtgtgctttcagagcagtagtgtaggaagcttggagt
gggatg (Seq ID No: 1014)

Homo sapiens zinc finger protein 397 (ZNF397):
cggctctttgtggcttgagctcgggggtgggtggctcatttcttgccgctcctgggcttcgcggaa
25 agaagagattactcacactccttcgcaagcacagaaccagttgtactgagctttttgctaagctgt
ttcagccaagaatg (Seq ID No: 1015)

Homo sapiens mitochondrial ribosomal protein L45 (MRPL45):
gctcccttcccggcggcctttgcggaacaagatg (Seq ID No: 1016)

Homo sapiens AKT1 substrate 1 (proline-rich) (AKT1S1):
30 cttccttctccatattgtatactggaattgaagccaaggaggtaccattttgctcgagggcatggc
ctaagccggtcagctaaggccatgttaatacggggctgtcccatctctctgcggggcgcgacagct
ggaagagccgaacggataagagaagaggaggtgagaggagctgtacaccacaagaggcactgaggg
actcaggataacgggatgaagccgtcagtgccccagaaacgaagcggccccggacgaatttctga
gtcaccgtcgcgagaaagcgggctgagccgccattttgaagcctggcaaaccgaagcaagaaatgc
35 tgccgtgttgatctttgccagccttcgtgccgaatgggagcaggttggaggaggaggagagccaat
atacactatgggctgattaagcccgggttggtgccatgttggttaacgagcaccgatttctctact
tttgtcgaagaagtttattgtgggtcagggacgtcaggtcgttgccttcgtttactgtggtcatg
attgagcatatgaggacggccattattgttgggggcaaattgaaatgctctaggcggggccatttt
tcttaggggcaagctgtcgtcacccttgtcaactggttcggatgaagccctgtggccgccatctt
40 gatctcgggcgccccgataagggaggcggagtgtgcggagaggaggcggggcaactgcgcggacg
tgacgcaaggcgccgcatgtcttttgaggcggtgacggcgccgggcccggccatgctggctacgg
gcacggcgcggtatg (Seq ID No: 1017)

Homo sapiens transmembrane protein 101 (TMEM101): ctgccctttcccaa-
gatg (Seq ID No: 1018)

Homo sapiens eukaryotic translation elongation factor 1 delta
(guanine nucleotide exchange protein) (EEF1D):

ggccctccctttcatcagtccttcccgcgtccgccgattcctcctccttggtcgccgcgtccttggc
tggcgtagagacagggtttcaacgtgttagccaggatggctcagtcctccagaccctgtgatccg
5 cccgcctcggcctcccaaagtgttgggattacaggtgtgagccaccgtgcctggccgaggctcctt
cttttatg (Seq ID No: 1019)

Homo sapiens ADP-ribosylation factor GTPase activating protein 2
(ARFGAP2):

cgccctccccgccgtggattggcccgcggcgaggaccgcagccgcggttgtgtctgggaaggaga
10 gaaaatg (Seq ID No: 1020)

Homo sapiens junctophilin 4 (JPH4):

atttctctcctccttgggggtctcagtgcatctccttctcctctctgctgcctcctcctcaccg
aagggttagcggacaccatccttttctgcttggggacccaccaccaccgcgaacactgccgctg
tctcttcttcaccgtatccttctctaccaccctccttctcttctcttctccttgcctttaa
15 tctgcttggccagcctccccgcgtgatgctgggatggagcaaacattgatttgtgctgggatggaa
tcggaattttgatttatttttcctctcccaaccataagaagaaaaataataaaaaacacccctc
ttgagagccccctcccccttgcacccagctccagctccttctcctatctccatccaaggcaga
tttttccccctacactattctcatcttccccacccttgcactacctcgccccccaccagcct
gctcctccagctggggagagaggggactctccggactccccacctttcctctctgggttggagca
20 gtctctccggaaggggagggggttggcttgcgggagggagggtggaggtatcctgccatg
gatgctgtgccggggaggcagcctgagccccagccacatgagacgccgaagaaccggggcagagg
ggtcctgacagcagccagggaacgggtgccctacgattctgccagccccctctcaggaccccc
aactgccatccacactcgacacttcgggggttagccactcaggatgaggggtccggcctgcctgc
cctcgttggggcccccccgcccggtctaaactgccccgccccgaggcctcgcccggtcc
25 aaggccccagcaggctctccagtcaccaggatgcgtgagccgcgggggggtgaggccgcgcaa
ctacatgcatg (Seq ID No: 1021)

Homo sapiens embryonal Fyn-associated substrate (EFS):

ttttctttctcctcctccaaccttggcggaggccacgactcaggcgccacagctgggggctagagg
ccgcggaccatggtgctggggcagccaccgctgaagtcagcaaaaccgagcctggcctgaggcaggc
30 tgcgcgggaggccaaagccatg (Seq ID No: 1022)

Homo sapiens GH3 domain containing (GHDC):

cgctccttctttctggccgatgtgtgctgagaccagagtcaccaggggtctccgtcacgtgcc
aggagtaggcagaagtgggctgtgacagatcaggaaacagagctcagtcagccactaaattgct
cagggccctacagctaacaagcggcagaggcaggatctgcactcaggagctgcttggagatg
35 (Seq ID No: 1023)

Homo sapiens acrosin binding protein (ACRBP):

ggctctctctgcggcttggcccgttagaggcggcttgtgtccacgggacgcgggcggatcttctcc
ggccatg (Seq ID No: 1024)

Homo sapiens jagunal homolog 1 (Drosophila) (JAGN1):

agttctcttcacggagccgcgcggtgcggggcgcaaatagggtcagtgggccgcttggcggtgt
cgttgcgggtaccaggtccgcgtgaggggttcgggggttctgggcaggcacaatg
40 (Seq ID No: 1025)

Homo sapiens ligand of numb-protein X 1, E3 ubiquitin protein liga
se (LNx1):

gttcctttcctgggcatcagcttgcctgctctcagcctaagctctctcgccaaccgtgggtggctcc
ttgcgttccatcctctcatctgagaatcagagagcataatcttcttacgggcccgtgatttat
taacgtggcttaatctgaaggttctcagtc aaattcttctgtgatctactgattgtgggggcatggc
45

aaggtttgcttaaaggagcttggctgggttgggcccttgtagctgacagaaggtggccaggagaa .
ggcagcacactgctcggagaatg (Seq ID No: 1026)

Homo sapiens cyclin-dependent kinase 2 interacting protein (CINP):
tctccttctacggatatctgtggaccttatg (Seq ID No: 1027)

- 5 Homo sapiens splA/ryanodine receptor domain and SOCS box containin
g 2 (SPSB2):
gcttctttccgcccggctccttcagaggcccgcgacctccagggctgggaagtcaaccgagctcc
cttcaggtcaatccaaactggagctcaactttcagaagagaaagacgccccagcaagcctctttc
ggggagtcctctagctcctcacctccatg (Seq ID No: 1028)

- 10 Homo sapiens Berardinelli-Seip congenital lipodystrophy 2 (seipin)
(BSCL2):
cctcctcctttcctccctctactctgacacagcacttagcacctgaatcttcgtttctctcccagg
gacctccattttccatatccaggaaaatgtgatgcccacaggtatcagcgtctggatcgccact
tcacgttttagccacaagtgactcagtggaagatccagagtcaacagaggctcgtcaggaagatg
15 (Seq ID No: 1029)

Homo sapiens tubulin, alpha 1c (TUBA1C):
caccctttcactacttctccccggactccttggtagtctgttagtgggagatccttggtgccgtc
ccttcgctccttcaccgcccagaccccttcaagttctagtcatg (Seq ID No: 1030)

- 20 Homo sapiens 1-acylglycerol-3-phosphate O-acyltransferase 9 (AG-
PAT9):
tttccttctctcttcccttcgagaggtgagtgccgggctcggcgctctgctcctggagctcccg
cgggactgctggggacagggactgctgtggcgctcggccctccactgcggacctctcctgagtgg
gtgcgccgagtcatg (Seq ID No: 1031)

- 25 Homo sapiens 1-acylglycerol-3-phosphate O-acyltransferase 1 (lyso-
phosphatidic acid acyltransferase, alpha) (AGPAT1):
gcccctttctttccttcgcttctcttttagagaatgtccggattgctattggactttggagcgta
tggctccaaatcaactcattggctaaaacttgacggaaaatggtggttaggtggccagaatg
(Seq ID No: 1032)

- 30 Homo sapiens abhydrolase domain containing 14B (ABHD14B):
cggcctcttcccagcgcttctcctccggccccaggtcaccgcccagcacgcgctgcttcccgtctg
cgcgagtccacgcagctccccagatcaagaagctgaggccccaggttacacactaaagtaaatggc
agaggcagaaataacacctatgtcctcctgaccccaaggcatgttcttaaagttctggaaacctcc
tggaggcttcccttgctgctcctctgggactgccaccctgggcagggtgttctgtggcccctcatca
tcgtggttttgaaccacaggcccttcaccagcacagcagcagcagcagcatg
35 (Seq ID No: 1033)

- Homo sapiens protein tyrosine phosphatase, non-receptor type 5
(striatum-enriched) (PTPN5):
catcctcccgcagcctgcccgcctgctcgccggcgcccggagcccgcctctggccgcttgcttttt
gctgagaaagcttctgcccctggaagatggcacccttccccatccagacaccttggaatg
40 (Seq ID No: 1034)

- Homo sapiens carbonyl reductase 4 (CBR4):
cttctcctttttcacggcgcttgcattactattgtgcggctgcaggaggtgtcgagcggcgttat
ttttttttgcgggtttgcctttttttttcttttttttttttttgaaccgcggttggttaaaagcct
gagggaacctggagaggggctcccactccctaccctctttcctccgagtttgtagtccgagatg
45 (Seq ID No: 1035)

Homo sapiens zinc finger CCCH-type containing 10 (ZC3H10):
 ggctctttgtcgaagctagaggaccggcaggcggcagcagcaactacggcggcggcggcagaaaccc
 agcagcgatgtggaggtggagacccacaggagccccggacttcacctgagctacctcagtggtcac
 caagagtggcaagataaagaaaaccctgagttgggcgggaccaggatg (Seq ID No: 1036)

- 5 Homo sapiens poly (ADP-ribose) polymerase family, member 10 (PARP10):
 ccgtctttcagttttcactttttgttttctgctcccagcagggttaggcttgctgaggggcaggcac
 aggagtcttggtgagctcatggcctgaggctgcctagcggccacggggaatg
 (Seq ID No: 1037)

- 10 Homo sapiens RNA pseudouridylate synthase domain containing 4 (RPUSD4): ccgcccttccttgtaagatg (Seq ID No: 1038)

Homo sapiens family with sequence similarity 73, member B (FAM73B):

- 15 ctgcccttccgcagcgatggcatcccggtgagtatcgccccggccgagcccccaaggcgggcgg
 gcagcggcggcaggccgggacttgagcggaggaccgagtaggcgcaggtgtccgggccaacagga
 ccaggaaggtgtcggggttggaatgagtggtaccgggcccgggacggtgcgagaggggtgccttg
 cttgggagcggaaacgagaaggtacttgggtcaggagagtgatgcccgggcctggaacgtggcgggg
 attggagcaggcgcgcaggtaccgatccgaggcggggagagcaccgggatggaaggagcaggcg
 tgccggccgtgagcggcgcagaggggtacctggctctgtggaggggcccctctggtatgtgtgtccc
 20 tgtccttctggggcgtggatggtgcctgggaccagctggcaaccagttgaagacgttctccttgg
 aagctcttggccctgaggactttgcctggggcattggccctgcatg (Seq ID No: 1039)

Homo sapiens protein phosphatase 1, regulatory subunit 15B (PPP1R15B):

- 25 gcgtctcttccggcgtctaggggggtgtcctgccggcgcgcgggcccctgcggccattttgggcttc
 gcttccaccgcaccagcggcctaccagtccttccggtatcgcgttgctcaggggcttttcaacc
 ctctgtcagtcggaaaaccatcgccgaggccgtggggggactcctatccatggtgttgaaagcgtcg
 agccgactagggaaacctccttccccgccaggatggaagtcgcatcagtcgccgcctattgcgcggg
 ctgttcttccctgtgttctgccgcccgtgccgcatctcgtgccctctgtggcttttctgctggct
 cgaagatcggcctggagcagcgcagccaccgctgggcaaggccgagactctgtaggcttctccga
 30 atcccgtcgacctccagccgctgagcgcgcggccctacctgagagactgtcaagaaaaaggagat
 g (Seq ID No: 1040)

Homo sapiens family with sequence similarity 104, member A (FAM104A): ccctctcttcgcggagcggcgcgcgtagcttccatccgcagctgccatg
 (Seq ID No: 1041)

- 35 Homo sapiens PRP38 pre-mRNA processing factor 38 (yeast) domain containing A (PRPF38A):
 agccctttacactacggtgtttccggccttcaagatgggtcgcctaagctgttttagtgaaacttcttc
 cacctttctccattcctctaggtgctttttctgaacctggatgtgaggcattaaaggatccgacgg
 aaatagaattgaaggcattctaaaatg (Seq ID No: 1042)

- 40 Homo sapiens synaptotagmin-like 1 (SYTL1):
 cctcctccgtgtggggcagctgctggctgggctgcctgttgagtcagccttcttccctcacggctc
 ttctcccggtccctgaaactcggctgccaggggagctggagccacctgcgaaggtgtcctcccata
 ctggaccctacaggaagctccgtgtgccagctggggcacagccccagctgatg
 (Seq ID No: 1043)

- 45 Homo sapiens ubiquitin associated and SH3 domain containing B (UB-ASH3B):

gctccttttctttttgatccattcaaaaattactcattgcaaattcccggactgctagggcgagga
gaggggaagggggcgaggagacaggggtactgcaggcgagagctgggggcagccgggggcccag
tggctgaggctgggtcccgagcgccgcttgccggcggttctggctcctgtggcctcaccaggaagc
gtcagagtcccgacactggggaagctcggagcgccgctccgctgcgcgcgctcctgctggctc
5 tgggtccccgagccccctccccctggcccagcccgaactccctccttcccgaaccatccggctcg
ggctccttccctggcgatggctggccgctgagccatg (Seq ID No: 1044)

Homo sapiens transmembrane protein 241 (TMEM241):
ccgtctctggggcggtgctgcccgtgcccgtgctgctgctgctgctggggggtcgggcgccaggggga
tttgggcaggcaccgtggatccccgagaagggggacgagttgacagatg (Seq ID No: 1045)

10 Homo sapiens ataxia, cerebellar, Cayman type (ATCAY):
gagcctctgccagccctgagctgggaagaagcagctacctcggaggcagggcgcgagggcgggcg
cgatgagagggggcgagcccgagcccccgctggggagcccaccgctaaccctgcacccccacca
ccccctgcacaaaagagctggcgggcgctggccacgtcgccctgggtgaccttctcgatgcagaa
tccgccccctgcgagcatcctcttctcctaggtctgaaggcccggggagcgtgagcgatgccag
15 ctgcacccggggcagggctcgcccttggtttgccagtaaggaggagagggctgtctcagctgcagaggg
gtcatccctgcttcaagccagtgcctcttcccagctcccatg (Seq ID No: 1046)

Homo sapiens ELL associated factor 1 (EAF1):
attcctctctacccccacgcagaggagagaacttgcttctggaccgggtgggtgcccggctcggc
tctccttgtcttccagagcgggtggcccggaagcacagtcctcccagacgccagcgccagaagctcg
20 gatcgcggtgcaccgggagagcgccgatctgggtgcgaggcaggtgcgggggcatg
(Seq ID No: 1047)

Homo sapiens tripartite motif containing 5 (TRIM5):
gttcctctaggaaaattcctttgtgcagatcaggcccggtgattgggtgagtgaatcctaaccacgt
cttccctggcctgtcttcaactcttctccccagaatcaccacttctgcactgggtgtctgaagggtga
25 ttgagtgattttgtggagggcagaagtaggaagtctttgggacaaaactgtatttaccttgggatc
tgtgaacaagaggaacctcagcagccaggacaggcaggagcagtggaatagctactatg
(Seq ID No: 1048)

Homo sapiens wingless-type MMTV integration site family, member 3A
(WNT3A): cgccctctcgcgcggcgatg (Seq ID No: 1049)

30 Homo sapiens chromosome 16 open reading frame 45 (C16orf45):
ctccctccctgcagcccgcaacgggaatggagtaaaggagaccgctcgacctggccacggggatc
agcgatg (Seq ID No: 1050)

Homo sapiens zinc finger protein 502 (ZNF502):
cattcttccggtttcagaagttaaggctgggtgtcctggccccagtcacactctgggagcgccctgcg
35 ccgctccgcggagagtcctggtgatctcacagtgaataatgtttgctgaccttgacattgacaaac
tgctgacagctcagatgatccatgattggaaggatgtgggtcatcaccaagatgtctttctttctcc
ggttcccagttttccagacctgaagtgttttccaatcaaagcgaagagacgatctgtggatg
(Seq ID No: 1051)

Homo sapiens armadillo repeat containing 6 (ARMC6):
40 ggctctcttgcgcaagcgcgctgtccgcttcttctggggcgagcgtctggaggcaaaacatttccc
tgctggggggcgggcaccacgtgagcgtcccgggaagggcgggcaaaagacgcctccgtcgcgcacga
gggtggcctcggttggtttaccttggttcgcggtcgtccttggttatcgtgagcgtccgcgagctctc
tgggaggccaagcctagggggcgccacagcgctgcgcgcgtacggcgggcgggaaggggctagaggc
ggctccctgggtgacaaccgcgcgccccacctttccccacgtggccgcgaagaccgggtcaggagc
45 atctatcgggtgcacgccaacatcaacacaggcgaagatg (Seq ID No: 1052)

Homo sapiens post-GPI attachment to proteins 3 (PGAP3):
gctcctcccccggcggcgagccagggagaaaggatg (Seq ID No: 1053)

5 Homo sapiens histone cluster 3, H2a (HIST3H2A):
tgccctcttgttttttagtctcgcttttcggttgccgttgcttttttccttgactcggaaatg
(Seq ID No: 1054)

10 Homo sapiens ethanolaminephosphotransferase 1
(CDP-ethanolamine-specific) (EPT1):
ggctctcctaccttctcgggcagcccagtccttgccatccttgcccagccggtgtggtgcttgtgtg
gtcacagccttgtagccgggagtcgctgccgagtgggcgctcagttttcgggtcgtcatg
(Seq ID No: 1055)

15 Homo sapiens F-box and leucine-rich repeat protein 5 (FBXL5):
ccgcctctgcccccggcggcgaggggtgtctatggagaggcggcggcgcggtgctgaggcggaggct
gaggcagtgggcagtgggcgccctttcctgaagaagtggacgtcttcaccgccccacactggcggatg
aagcagctgggtggggctctactgcgacaagctttctaaaaccaatttttccaacaacaacgatttc
20 cgtgctcttctgcagtcctttgtatgctactttcaaggagttcaaaatgcatgagcagattgaaaat
gaatacattattggtttgcttcaacaacgcagccagaccatttataatgtacattctgacaataaa
ctctccgagatgcttagcctctttgaaaagggactgaagaatgttaagcctactactgttgactgg
aagccttaccaataacataaaacaatcgaataacaattatttcatgtattatatgtaaaatatata
tactggattcttacagtaagaatgaatatgaacagttaaattatgcaaaacaactgaaagagagat
25 tggaggctttttacaagagatttttcttcctcacatg (Seq ID No: 1056)

Homo sapiens major histocompatibility complex, class II, DP alpha
1 (HLA-DPA1):
ctgcctccactcggcctcagttcctcatcactgttccctgtgctcacagtcacatcaattatagacccc
acaacatg (Seq ID No: 1057)

25 Homo sapiens secretory carrier membrane protein 1 (SCAMP1):
tcgtctctctctctgcgcctgggtcgggtgggtgacgccgagagccagagagatg
(Seq ID No: 1058)

30 Homo sapiens chromosome 15 open reading frame 57 (C15orf57):
ccgccccctccgatttctcctcgggctacaggcgacagagctgagccaagcgtttactgggcagctg
ttacggtaagtgaggaggggctgggggtgccagcgttttgatctccactctggcccgccccgg
aataccacatagaggccttgggacctgattcatcccgctccagacagccctagagacctgagcgact
gaggcctgggatctggacgccggaatttctcgtggttctggacgccctgcctgggctcagatt
ccaaatg (Seq ID No: 1059)

35 Homo sapiens WD repeat and FYVE domain containing 2 (WDFY2):
cctcctctttagtgggcgccggttgcatcccaggctcgtggcggttttggtgcctgaagcagggag
cgcgagtcgttcccgagagaggcgccaggctatgctcgccggtttccggcggtccgctccggcc
agccagagtctctgtctcaacctgtgtccgtgctccagcagtcctcagccccggcccccgggcgc
ggttggcgggcggcgccccaggcgcgccccctcctccgatg (Seq ID No: 1060)

40 Homo sapiens topoisomerase (DNA) I, mitochondrial (TOP1MT):
cgctctttcccgaggctggcagatg (Seq ID No: 1061)

Homo sapiens intraflagellar transport 122 homolog (Chlamydomonas)
(IFT122):
ctttcccttttcggacatgcgcgctcggagcaaggcgccctcgcaactcagcttaccgcgcatgtacg
ttgccaggggtaacgcaggtagccaaagtggcttgtggagtggcgaccggttagtgaggcggttgct

gagacagacgctgagggcggttaggaggagcccagaccgtaagggagccgtgatg
(Seq ID No: 1062)

Homo sapiens mitochondrial ribosomal protein L53 (MRPL53):
agttcttccggggcgagggtcaccatg (Seq ID No: 1063)

5 Homo sapiens T-cell activation RhoGTPase activating protein (TA-
GAP):
ccgccccttcgcttataatgcagagcatgtgaagggagaccggctcggtctctctctctccagtg
gactagaaggagcagagagttatgctgtttctccattctttacagctcaccggatgtaaaagaac
tctggctagagaccctccaaggacagaggcacagccacacgggagtgaatccacccctggacagt
10 cagccgcaatactgatgaagctgagaagcagccacaatgcttcaaaaacactaaacgccaataata
tggagacactaatcgaatgtcaatcagaggggtgatatcaaggaacatcccctgttggcatcatgtg
agagtgaagacagtatttgccagctcattggacattctcactattctatgccttaaaggcccttca
acggaagggatattcaggagagcagccaacgagaaagcccgtgaaggagctgaaggaggagctcaac
tctggggatgcggtggatctggagaggctccccgtgcacctcctcgctgtggtctttaaggacttc
15 ctcaagaagtatccccggaagctactttcaagcgacctctttgaggagtggatg
(Seq ID No: 1064)

Homo sapiens phosphoserine aminotransferase 1 (PSAT1):
ggctctccttggtgactcaccgccctggccgcgcaccatg (Seq ID No: 1065)

Homo sapiens CD97 molecule (CD97):
20 cccctccttcataaagtccctggcctcgggacagcctgcacagctgcctagcctgtggagacggga
cagccctgtccactcactctttccctgccgtcctgccggcagctccaaccatg
(Seq ID No: 1066)

Homo sapiens protein tyrosine phosphatase, non-receptor type 2
(PTPN2):
25 cgctctccccggatcgtgcggggcctgagcctctccgccggcgaggtctgctcgcgccagctcg
ctcccgagccatg (Seq ID No: 1067)

Homo sapiens chromosome 20 open reading frame 112 (C20orf112):
gcccctctccccgggacgcccggcgggcagcagcagcagcagcagctggagctgtggggctgtcac
cgccggccgccccgctcactcgcggtatcccgaccgccatctccgctcgcttcagcccaggatg
30 agacttctgtgagcagcgaggattttgatatg (Seq ID No: 1068)

Homo sapiens APEX nuclease (multifunctional DNA repair enzyme) 1
(APEX1):
cacccttctttgtgctcggggttaggaggagctaggctgccatcgggcccgtgcagatacgggggttg
ctcttttgtcataagaggggcttcgctggcagctctgaacggcaagcttgagtcaggacccttaat
35 taagatcctcaattggctggagggcagatctcgcgagtagggcaacgcggtaaaaatattgcttcg
gtgggtgacgcggtacagctgcccaagggcgttcgtaacgggaatg (Seq ID No: 1069)

Homo sapiens intermediate filament family orphan 1 (IFFO1):
tttcctcttgagccatcatgcacatctgactgcagccccagcagcccttcttctctgtctgact
gctcttcttctcgatttcttcttctgttctgccttctcggtttgcagccctgaccccgctgtgtgtc
40 tggcccttggtgactgtccgtgtttctgttctgtcattgttaactgtgacttttctctctgtctgc
cccccttctactggttcatgttctctccccattcccacctctctgcccggcctcccgtccccg
cccttctctctcatgcacccggcctcgctctctgtagtctctgcacttgtctccattaaggtccca
tccatg (Seq ID No: 1070)

Homo sapiens neuralized homolog 2 (Drosophila) (NEURL2):
45 cagtcttctcccgccccttctttggtccctacggacctggggggcggtggcggtcaatgccgggt

caaggtccgcgggacctgcagatcgtagcccgggcgacgcgatcagatgatcctgttggtggaacgg
ctaagttgtaggcgggatggctgagaaagcggcgctaggacccccgggcagaggctcggggaagg
agtcaggggggaaatgccttacaaggtgccttgcggtcaccatcattgccgcgcgccccaaatag
ccccggcgccagctggcctgcctatggccgagagatg (Seq ID No: 1071)

5 Homo sapiens drebrin 1 (DBN1):
ctccctctttccctccctcctcctccgtccgccggtccgtccgcgcgtctgtccgttcggcccggt
ccggcccgaagcatg (Seq ID No: 1072)

Homo sapiens WW domain containing adaptor with coiled-coil (WAC):
cagcctcccttatttagtcgcgatggcttccctcgcgccccaccgtcctcttcggaaggcggct
ccctccctgcgagcccgagccctgagatcagcctcgagcaggcgcccgagcgaactatccct
aaacgggaacggcgggtggccgactcgcgagtgaggaaaagaaggaaagggcagactggtcgcgaag
agaagatccaggcctcagaggaggagaaaggccggagccagccgaggtttgccgagggcggtgttc
cggacccgcgcggtgcggggaggaaggccgaggggtgggagaggaggggcccggcggaactgccga
ggtttcccgaaggcggcagcgtccgagttgcccgatgtagtgtggtggagcggcagcggcggcacc
agcggcggcgcggcggcgggaggaggaggaggagaagaaggaccaggcggcggcagcagcggcgg
cggcggggggaggaggggaggaggcggcggagcaggaggaggagaaggcgaggaggcagtcgctc
tcgcgggggctgagccggacgcgtcgtcttgccccccctcccccggttcgcggtgcccgcgtgtag
ttggcgccgctgccccggctgagagtgcgcgtggtgtgcacggaggggagatggccccgggagcgccg
gcgccagtaactgggagctgatgagagtcgccgagggcgcgccggggcccaggtgccggggctgcc
gccgccgcgcgcgcgcgcgcctgcgcgcccgcgcgcctttcgcgccgctctccccccctcccga
cacacactcacaggccqggcattgatg (Seq ID No: 1073)

Homo sapiens kelch-like 6 (Drosophila) (KLHL6):
cgctccttcagtctcgaatg (Seq ID No: 1074)

25 Homo sapiens GTPase, IMAP family member 1 (GIMAP1):
cagcctttctgcactcacagccgaagggaaagcagcaggttggggcttcttgtggccaacttcagag
cctgtcaccaggaaaggttaagcatg (Seq ID No: 1075)

Homo sapiens RAB24, member RAS oncogene family (RAB24):
cgccctctagccccctccgcgggagtcgcggcgctgcgggtaggagccgggttgcgggagacccc
aggttcggttgggattcccagccagaacggagcttaagccgggagcgagcgaatgacggagtag
30 cgagctgcacggcggcgctgctgcgctgttgaggacgctgtcccgcgcgctcccaggccgccccgag
gcttggggtcttcgaaggataatcggcgcccggggccgaacagcgggggcacacggggcgctgccg
aagtgcaggccacggccagagctcgagcccgacgcgctgtctggagtcgtaggacctgacgtgg
ctgaagcggccccgggagcatg (Seq ID No: 1076)

35 Homo sapiens adaptor-related protein complex 2, alpha 1 subunit
(AP2A1):
agccctccccgcggcggctcggctccttggcgtgcctggggtcctttccgcccgggtccccgctt
gccagccccgcgtgctctgtgccctgtccggccaggcctggagccgacaccacgccatcatg
(Seq ID No: 1077)

Homo sapiens copine IV (CPNE4):

40 ctcctcttttctcagtagcctcctctttactctccgagttaactgagagccgacctgacatctcc
aacattttcacctcttcccccccccacccgagaatggagtcagggtttccggagagaccga
actctgctctcagcacctttccagccgctgttgctaaactgacctcgaggacgagaggggaagg
aggtgcgacgccccttacatcagtagcataactaccacaccaaccacctccacttcaaagccggatt
45 ttgcatcctggggggcgggacagacctcgtcccgggctgaattctctctccactcttcgagattggc
acaccagaatg (Seq ID No: 1078)

Homo sapiens synaptosomal-associated protein, 25kDa (SNAP25):
ctgtcttttccttccctccctgctcggcggtccaccacagttgcaacctgcagaggcccgagaaac
acaaccctcccgagaagcccaggtccagagccaaaccgctcactgacccccagcccagggcgccca
gccactccccaccgctaccatg (Seq ID No: 1079)

- 5 Homo sapiens cAMP responsive element binding protein 3-like 4
(CREB3L4):
aggtctcttgactctttccgcctttgtttacaaccctgccatgatctccctcttgcaaaagcgagg
gctacagaacaggcattcaggagtcctgtgctccagtcacagccttttctgttcttcagctaggag
acaccaaaccctcaggaagatttactatagctaagagaaaaactgcagcagaaagggcgcggtacc
10 tacttcttaaatccgtttgtggaccctcagactcttagtccccctactcccagatacagcgccct
accgtggctcctggcaaggtggcatccactttttagtaagcatg (Seq ID No: 1080)

Homo sapiens leucine-rich pentatricopeptide repeat containing
(LRPPRC): ctgtccttctggcggagcgtgcttcccgctgcggggacgttcgagcaatg
(Seq ID No: 1081)

- 15 Homo sapiens zinc finger protein 418 (ZNF418):
cgttctctggttagcgaccatttttggttaatgttgggtgtgtttctgcggtttgtgaggtgagaggc
gctggagctatgggtccgaaccgcggtgtctgaaccagaaggtgaagagtccttcttctgtgcaca
gaggcagatcttaggccccgtaacggcgcccgccgctcccggcagtgtttccccgcgtactcggg
atggcgggcgccgcgtgaggtcctccggctcaggcatcatctggctgcaaagaagagaacacactg
20 tgtttgagggaggaggaaggaggatcagagtttaactcctgccataatg
(Seq ID No: 1082)

Homo sapiens tetratricopeptide repeat domain 14 (TTC14):
gtttcttccgcttctgtaccacccggctcaagtagcggacacggaacagggaactatcagcccgt
cggcctccggggccctgcattctctagccatg (Seq ID No: 1083)

- 25 Homo sapiens BMP binding endothelial regulator (BMPER):
agcccttttgcactgtgagctgcggcagctgagcagaggcggcgggcgaggacctgcagtcgccag
ggattccctccaggtgacgatg (Seq ID No: 1084)

- Homo sapiens zinc finger protein 384 (ZNF384):
cccccttttgcgttccggcgctccgccttctctccgcagagctcttctctgagcctgttgggggg
30 agggagggggcggtggaggaactggggttcgcgggagcacgagctgcagcaccacttccgggtgag
tgcaaggggagggcagcaaggagggggggccaccactacctcgcgcccccgccctgcgggtgtct
cgcgcgcttccgtgcgtgtgagtggtgtgtctcgtccagaagtgcgtgcccgcgcgctg
cgcttgcgctttttccctccctcgccttcttggctcctccaccctcctcggtccctccttt
cccagcaaacgccccctcccgcgccctggctcaggctctggcgcccgccgagccgtcgccgccc
35 gaaagtccagagccctggaaaggagaaggaataagacggcaggaggaagagagagagagggtaga
atg (Seq ID No: 1085)

- Homo sapiens RAD51-like 3 (S. cerevisiae) (RAD51L3):
ctctccttttctcctccggcagccagcgcgctgtgtcctctctaggaaggggtaggggagggggcg
ctggagaggacccccgcgaatgccacgtgacgtgcagtccttggggctgttccggcctgcgg
40 ggaacatg (Seq ID No: 1086)

Homo sapiens CD99 molecule-like 2 (CD99L2):
gctcctcctccgctcctcctcggcctcccttccggcgctctcgcgctaactgtgctcctccggg
gccctccgcctgctcccagccatg (Seq ID No: 1087)

- Homo sapiens glucosamine-6-phosphate deaminase 2 (GNPDA2):
45 gcgcctttatctgcatccgggtccgtgggattcgcgctccactgggtcagctggggctcgctctcggg

tggttggtggtgttgcttgttcccgctgttccagcgctcgaagaaccattgggtctgccggtttgaact
tggtctggaagctgtgcgtcaccgtaatg (Seq ID No: 1088)

Homo sapiens methionyl-tRNA synthetase 2, mitochondrial (MARS2):
ccgcctcctccgcttgcgccggtctgcacatg (Seq ID No: 1089)

- 5 Homo sapiens chromosome 12 open reading frame 57 (C12orf57):
tttcctttccgctcccagggcggttggaacggttgtaggacgtggctctttattcgtgagttttc
cattacctccgctgaacctagagcttcagacgcctatg (Seq ID No: 1090)

Homo sapiens tRNA-yW synthesizing protein 3 homolog
(S. cerevisiae) (TYW3):

- 10 ggaccttttcggccaccgctcgcttcaatatggctgccccagggagagacgaggctaccatgaag
gagccgagcgcagaccctgagtcgcgtcacccatg (Seq ID No: 1091)

Homo sapiens Sp1 transcription factor (SP1):

- ctccctcctccttccccccccctccctgtccgggtccgggttcgcttgccctcgtcagcgctccgcgtt
tttcccgcccccccccaaccccccgacaggaccccccttgagcttgccctcagctgccaccatg
15 (Seq ID No: 1092)

Homo sapiens histidine triad nucleotide binding protein 3 (HINT3):
cgccctctagtggcagccggttttgaggccggcctccggccttgaagttcctcacccgctctcctt
ccctctcccaaagcctggatcacgcgccagcgctcaggcgagggcgacgtctcgaggtaaaacgg
aggaggtgcgggacgcggagactgcgcgggcccggtagccctggagaggccgaggctctaggccgc
20 gagggcggggtgcaatg (Seq ID No: 1093)

Homo sapiens M-phase specific PLK1 interacting protein (MPLKIP):
agttctctgcggagggccggttgatacagttccgggtgggagaacgcggctgcgaggttttcggctt
tggctcctgatatg (Seq ID No: 1094)

Homo sapiens palmitoyl-protein thioesterase 2 (PPT2):

- 25 cacccttccccccgccaccgtgggttccagacttgggataagtaaacagcgggtggagcgaggcct
acggacccaggccaggtgggagtcctgcactcttcaaggggctgggctgctgctcacgggtattaa
agaactccgcgttggtcatggctgaggcgatgcattaggaagatcctggacctagagaacaagtcc
ccgaacgctgagttggaggcgggacttcgggtgcgcggttgccgggagcatg
(Seq ID No: 1095)

- 30 Homo sapiens BCL2-like 14 (apoptosis facilitator) (BCL2L14):
aagcctcttttcaggctgagtcctaaacctgaagaaagtttagagcctggggctctaaactacctg
agtctttccaaacgacaagccaagaagacctgttgaaagtttctcttaagtttcgtggagagaga
ctcaggtatagaaatataccttactgccacctgacctgaagcagaagaaatcacagacagcttccag
accaggcccaacatg (Seq ID No: 1096)

- 35 Homo sapiens galactose mutarotase (aldose 1-epimerase) (GALM):
acgccccttctcctgtaaaacttgggtcgccctctagcttagcgagcgctggagtttgaagagcgggc
agtggctgcacacgcctaaactttccctatg (Seq ID No: 1097)

Homo sapiens carboxymethylenebutenolidase homolog (Pseudomonas)
(CMBL):

- 40 cttccttcccttccccgactttgcagattttcttccccagggcctccctcctccacctctccgcc
ccctccgggcttggctctccagaggctacgactggagccactgggtcccgcaggatccccgcgtc
ctcggtgcgcgcgtccacgtccctctcgctccccgccggcgccacgcgcctcctctgggttcg
gcctccgcgcggtgcagcgcagtctcaggccgcgggacaagcccgacttaaatctctgcaatg
(Seq ID No: 1098)

Homo sapiens chromosome 7 open reading frame 31 (C7orf31):

cgtcctttctcccgcccccgccctgacctgccagctccaccgggcccgtaggtgaggacgacctcaaa
attcctcggccccgcgaaggccgagctgcggggaggggagggagggcgcggtcccgcagcgcgcc
caggctcatgtcccaggtatgtccagacccccgaggcaccgcttgagggcagtgacagcccgtga
5 ggctcggcctcgacccctggcacccttggtcccagctacgcccgtcctggccttcccccaagtcc
gagagagaggtgggatttctccccgacgcagttggaaaccgggaatcccctttagggctccgttcgt
gctgcactactgactccaccatctgcaaagggttcttgtccagaatccccgaaggctttaggaca
gcgcttattttgttgatgaagagtctctaatttttcggaaagaccacaggctaaaagtcaagttgt
gccttttttagccaagaagcatg (Seq ID No: 1099)

10 Homo sapiens secretory carrier membrane protein 5 (SCAMP5):

cggccttttcggcagccgaacggcgcgagcttcaggacaaagaggtgtgggcaggccactggggcc
agctggtaacatcatg (Seq ID No: 1100)

Homo sapiens mitogen-activated protein kinase 10 (MAPK10):

15 tgctccttttcggttgccatagcaacccattccccaaagccctctgtccgtctcctctggttaggttc
cacaatggtacaggcagcatcacgctgcacaatggtttccaggcagtgaaagaggggtgattcagca
agccactcttcttctattttctttaacctcccccttactttttatttttatgggggtgggtggtgc
ttgctatatgcttacctttttctttttcttttttcttttttataaaatttcttttttgcctcacc
ctcaattcctaggggcttgagtgtgatttaagattgggttttcttggaatcacctgtccatcggtta
attttaaacaatctccatatctccaaagaatctcttccatgttagtctggaatgtgggttaatgaaa
20 aacaagtagggaggatttctggggcaaacactgccggatcaggatcgtagttctcaggcacggaat
ggctagtgtgagaaacaccaacagcaggcccatctcagatcttcactatggcaacttatgcaagaa
actgttgaattagaccggtttcctatagatgagaaaccatacaagctgtggtatttatgagcctcc
atttcttatactactgcagtgaaccaacattggatgtgaaaattgccttttgtcagggtgtgtgttc
cttacaggtaaaacaagggttcgataaacaagtggatgtgtcatatattgccaacattacaaca
25 tg (Seq ID No: 1101)

Homo sapiens beta-site APP-cleaving enzyme 2 (BACE2):

cgtcctccccgcgcgcgcgggtcccgggtgcgcgcccattccctgcccgcagccccgcgcgcggccg
agtcgctgagccgcggctgccggacgggacgggacgggctaggctgggcgcgccccccgggccccg
ccgtgggcatg (Seq ID No: 1102)

30 Homo sapiens SWI/SNF related, matrix associated, actin dependent r
egulator of chromatin, subfamily d, member 1 (SMARCD1):

acgccttttccgctagtcgccccgctctatcccatagtctcgtgccctgagcctcccgtgccggc
cggccggccgggggaacaggcgggctcgggggcgctcgggggcggggggagttccggttccg
gttctttgtgcggctgcatcggcggtccgggaagatg (Seq ID No: 1103)

35 Homo sapiens family with sequence similarity 175, member A

(FAM175A): cgtcctcttgtgtagcctgaggcggcggttagcatg (Seq ID No: 1104)

Homo sapiens adenosine deaminase domain containing 1 (tes-
tis-specific) (ADAD1):

40 aggcctcttttgaaagatgcggccctgaccctgtgaacctcgcgagagcggcctgaagcgagagg
ttgaggctgggaggtgagaaaatg (Seq ID No: 1105)

Homo sapiens acyl-CoA synthetase short-chain family member 2
(ACSS2):

gcccctctacggaggccccgcctctagttcggcctgttttctcagtcgccggcaccgcgcgacccg
caaaggcggccgcgggttctaggaacttgacgtgatg (Seq ID No: 1106)

Homo sapiens multiple coagulation factor deficiency 2 (MCFD2):
cttccttactcaccggtgtccggaaggtgaacgctgcgctcgggctgcctcgctgttacctcc
gccgccgggcatg (Seq ID No: 1107)

5 Homo sapiens SPOC domain containing 1 (SPOCD1):
gctcctttttcagctagtgggtggaaccccaggaggggaaaactcaggaagcccagggcccggtgttg
tgcttttggcccaggtaggtggacagacatg (Seq ID No: 1108)

Homo sapiens LY6/PLAUR domain containing 1 (LYPD1):
agttccttcagtctcagccgccaactccggaggcgcggtgctcggcccgaggagcgagcgaggagg
agcagagacccgcagccgggagcccagcgcgggcgatgcaggctccgcgagcggcacctgcggct
10 cctctaagctacgaccgtcgtctccgcggcagcagcgcgggccccagcagcctcggcagccacagc
cgctgcagccggggcagcctccgctgctgctgcctcctctgatgcgcttgccctctcccgggccccg
ggactccgggagaatg (Seq ID No: 1109)

Homo sapiens cytochrome b5 domain containing 1 (CYB5D1):
cattctttcatactgcctcctcccttggttttctgtctcagagagatagtctgtcctaaatatccc
15 atgtagcccaggccactgaattaaaacggagcgtattcgttctctgccccaccccgcaactcctga
aagcggcgcaactcaattacttgatccttatatgccccacgcgggactcatactacgtttcccggtg
aacacgtgcagtccaaaccccgccccctgatatttatctcagtggacggtggccggaaaaggacaat
ggtttccatgtcagcggataaacgctctccctcggctcccgagcgcgacggaggtcgtagtagta
gtgagtacgtgctgaggagcaaaggagtaaccaagagatccagtgaccgacagagcaagagccatg
20 (Seq ID No: 1110)

Homo sapiens synaptoporin (SYNPR):
tctcctcctttgcttcataaaaagaggggacaagtggctggtgctgtggacagagaagctttat
tagtatgagacaacctctattttctttcaggagaggggaagtggattatcaattcttttgtaa
(Seq ID No: 1111)

25 Homo sapiens heterogeneous nuclear ribonucleoprotein U-like 1
(HNRPUL1):
ccccccctttcccccttcgcctcctgacaggaaaggtttaagggggacagagccctgggaggccgg
gccgggctcggggggccaccccgggggcccgggccatg (Seq ID No: 1112)

Homo sapiens schlafen family member 5 (SLFN5):
30 ggttctctgctctggacttgggaggctccgttgctgctcccgaggaggagacgcgctgccgaggag
aaccagcgggagaaacatttcaggataggaataggccaagtgctgagaagatg
(Seq ID No: 1113)

Homo sapiens MAS-related GPR, member F (MRGPRF):
ccatctcttccagcaggagagggtctactctgagctcctattttccaaggctccggggccgcgctc
35 ggcgctggcctgctgccccggcggggtccgcggccggaggcgggagtcacaggaagagccctccac
aaaaggaggcctcggcggatcaggacagctgcaggtgggtgtgcagactggtgagctgccagcagg
ggcccagacgcgccaggcctggagatg (Seq ID No: 1114)

Homo sapiens ubiquitin-like domain containing CTD phosphatase 1
(UBLCP1):
40 cggctctctcagcggccgggtttctgcgtccgctgccgcaggttccaccgcgctccaggtat
ttctgaaggaaagctgcttcctcatatgtttcaagaatg (Seq ID No: 1115)

Homo sapiens Rab interacting lysosomal protein-like 2 (RILPL2):
cctccttttccgttgctcccttcgcgccccaaaccacatcctggagcgcaactctccagcgtggctgg
cagcggggacgggtgcgcgggggcgagggcccaagagtcgcgtgcgcggcccccttgaccatcccc
45 cggggccacccccggggccgcgctgattgggcaggtagggactctgccagcggaaagtgttgggtg

5 acttctttttccgtgggagtaaggaagtgccttttgaatgaggtactgagggccaaggtgttggaagt
 tcctaattctttcctcggttaactgtgaaactctgcgtattgggaaggcctggcctcagtcacag
 gccaggagaggtactggacgccgcgcacgcactcgtctgccagcgaggcccaaagggggaagcctag
 cggagctcagtggtgcagctgctggcctctgggccgctacttgtaataccatg
 (Seq ID No: 1117)

Homo sapiens solute carrier family 2 (facilitated glucose transporter), member 12 (SLC2A12):
cactcttcttttagcatgctattatggggaaagtgaccactcctgggagcgggggtgggtcggggcgg
tttggtggcggggaagcggtgtaacttctacgtgaccatg (Seq ID No: 1119)

Homo sapiens tripartite motif containing 11 (TRIM11):
gctcctcttctgcccgcacatccgggacccctacgtcccgcgtcccccgagcgctcggagcctacgc
gccagcgctaccgaaaccagagtcctgcgccttgagtccccgcgccccggagcccgagcacc
gggagtcggcgagcctcgcgccccggagtgcccagacctgcgcgcgcgcacccggatacccgcgctc
cccgcgagctgccgaggccgcccgcgcgcgcgcgcggacagtaccgccttctctccctctgtccg
cgccatg (Seq ID No: 1121)

Homo sapiens zinc finger protein 626 (ZNF626):
cggcctttgtctctcgctgcagtcagagctccaggtctggttcttctcctaagggcccaggctgtg
tggtcccggtgtcctgcaggtattgggagatccacagctaagacaccgggacctcctggaagccaaa
aatg (Seq ID No: 1123)

Homo sapiens solute carrier family 25, member 43 (SLC25A43):
cggtcttccggcccggaatcggggtcgaatg (Seq ID No: 1124)

Homo sapiens crystallin, zeta (quinone reductase)-like 1 (CRYZL1):
ggctctctgacgaaggactggaaggtggcggtggtgaaggtgcaggccgttggggcggtcagagg
caggtgactatg (Seq ID No: 1125)

5 Homo sapiens mitogen-activated protein kinase kinase kinase 7
(MAP3K7):
ctgcctctacccccgccacggatcgccgggtagtaggactgcgcgggtccaggctgagggtcggtc
cggaggcggtggtggcgcggtctcaccggattgtccgggtggcaccgttcccgccccaccgggc
gccgcgagggatcatg (Seq ID No: 1126)

10 Homo sapiens septin 6 (SEPT6):
ctttctctttgtcggaggagctcctctgtttcctgtgcagtagctcccgttgcggcggcaccctg
gcagccctggcgagcgcaggagcatg (Seq ID No: 1127)

Homo sapiens myotrophin (MTPN):
ctgcctctcctcggccaggcggaacctctctgctgggcccgggtggccgcaaaagaactttctttct
ccgccccgaacggtcgccgcggccaactgcctcgccgctggcagcctaaccctccttctcttct
15 tctcctctccggttcgcgcggccctgcctccctctcgccggcggcacatccgcttgctgctgccac
cgctcctcatcttctgcccggccaaccggcctgccccgctgcagtcatg
(Seq ID No: 1128)

Homo sapiens annexin A11 (ANXA11):
ccctcccttgactgcctctggcacctggggcagccgcgcccgcggagttttccgcccggcgctga
20 cggctgctgcgcccgcggctccccagtgccccgagtgccccgcgggccccgcgagcgggagtgga
cccagcccctaggcagaaccaggcgccgcgcccgggacgcccgcggagagagccactcccgccca
cgtcccatttgcgcccctcgctcggagtgcccggtggccagggtatttggtgacctgctggtttaa
actattgtcttagttaattttgtgctgcttaacaaaatatcacagactgagtaatttataagcaa
tagtagcttatttggtcacagttctggaggctgagaagatcgtaggctgcatctggcaagggcc
25 ttcttgctgcttcataacatggcagaagacatcatgcgggtgtgtgtctggggaagagacttacag
aagtggagttgctgagtc aaagatctaaccatg (Seq ID No: 1129)

Homo sapiens RNA binding protein, fox-1 homolog (C. elegans) 1
(RBFox1):
ttttctttctttcctctcccggcggttgatgagtgcttggtccttgacagaagggatttggtccca
30 gctttgtagttcggaagaagttgggtctatagatttccccctaactctccattgatgtgttgagct
tcagaggggaataataactctacgtaaagcatg (Seq ID No: 1130)

Homo sapiens prefoldin subunit 5 (PFDN5):
cttcctcttcgtaagtgcggccttcccaacatg (Seq ID No: 1131)

Homo sapiens high mobility group AT-hook 1 (HMGA1):
35 cgctctttttaagctcccctgagccggtgctgcgctcctctaattgggactccgagccggggctat
ttctggcgctggcgcggtccaagaaggcatccgcatttgctaccagcggcgccgcggcgagcc
aggccggtcctcagcgcccagcaccgcccgtcccggcaaccggagcgcgccaccgcaggccggcg
ccgagctcgcgcatccagccatcactcttccacctgctccttagagaagggaagatg
(Seq ID No: 1132)

40 Homo sapiens zinc finger protein 323 (ZNF323):
cggccttttgcggttgatcggtcattgggtgctgcagccccgccacctgttccgtagcttgccggt
gccccgaaggtgtcttctcctaaggaagattaaatcagaaaattttaaatcacagttatcccttta
cttaaagccagagtaagccttccaaattaacccagggaatg (Seq ID No: 1133)

Homo sapiens tumor protein p53 inducible protein 3 (TP53I3):
45 ctttctcttctcttagcagcaccagcttgcccacccatgctcaagatgggcgggatgccagcctg

ttacataaatgtgccaaaagcctggccatgcctggaaaatggaccaatccgcccgcgaagaggttg
ggtctcgttccctagagagaaggaagtttctctccttgaagtgaagagctagaatcgcactttctg
tcaagctgagagaaaagactcttttccagaggctaaaaggacaagaaaatctgatttgcttgcttct
aactttgcgtttttaaagggggaaggaggaaaggaaagagggggagggtggttctgcttagccccac
5 cctccgggtaccccggtccagccgtccattccggtggaggcagaggcagtcctggggctctggg
gctcgggctttgtcacccgggacccgcaggagccagaaccactcggcgccgctggtgcatgggagg
ggagccggggccaggaacaatatg (Seq ID No: 1134)

Homo sapiens ceramide synthase 5 (CERS5):

10 ccgctccccgcgggttccggttggtgtggcggcagctgacgcttgtggcggcggtggcttcgggg
tggcgtaagatg (Seq ID No: 1135)

Homo sapiens TRAF3 interacting protein 2 (TRAF3IP2):

tgttcttctacttacctgggcccggagaaggtggagggagacgagaagccgccgagagccgactac
cctccgggcccagtcgtctgtctgctcgtggtggatctaagaaactagaatg
(Seq ID No: 1136)

15 Homo sapiens Smith-Magenis syndrome chromosome region, candidate 7
(SMCR7):

ggtccttcacgttccattcccaggctggtctgagctccggggccgtggtcccgtgcctcctccgg
tcgtcgtgcggaagctgcgacgcaggcagaccatg (Seq ID No: 1137)

Homo sapiens mitochondrial ribosomal protein L10 (MRPL10):

20 cattcttccggtggagatggctgcggccgtggcggggatgctgcgagggggtctcctgccccaggc
gggctagagtgcagtggcatg (Seq ID No: 1138)

Homo sapiens proteasome (pro-
some, macropain) subunit, alpha type, 1 (PSMA1):

25 acttctctgtagatcgctgagcgatactttcggcagcacctccttgattctcagttttgctggagg
ccgcaaccaggcccgcgcgccaccatg (Seq ID No: 1139)

Homo sapiens sorting nexin 5 (SNX5):

cggtctttctctagacgcgtcttgctgggagagtgtccgttgcttcccgtccgtgtcgcgccctg
cggttgccggcctcctcgtggagcggagcaaggccaggcggccctgctcgagtcccgcgtcgcca
tg (Seq ID No: 1140)

30 Homo sapiens zinc finger protein 276 (ZNF276):

gggccccctccgcgcgtactgcgggccccacgggtgttagtgggcgggggcggcagagtccgggtgg
gttgctcgacgagcgggacctcttcgccgtcttgagacggggctggcgagaagggccccctcacg
gagttgcatgggctctaaccgcggcagccaggccccctctctacgtgagaccccgccccctcc
cctttctgcagcccgcgccacctgcgcgcgcgtggcctccgcggcgccgtgcccggccccgcgc
35 ctccgtctcccacggagcaggccgggctctcgccatg (Seq ID No: 1141)

Homo sapiens zinc finger protein 561 (ZNF561):

ccatcttttccggcgctggctcctctccgtcagtgcggtttcgccctttatggtgggtggagtctgcc
caggctgtggaccgcaaataaccctgtacaaagaggaatggagattgcctctatccacctagattc
ataagctggcctgaggtgatcttggcatcaaggaagggatgcacatcatcacaccatcagcttcag
40 agaatg (Seq ID No: 1142)

Homo sapiens mucin 7, secreted (MUC7):

ctttctcttcttttgccttctagttaccatcctcaaaggattggctaaaagcaagcaactggattga
acaccctaagaagaagattcacactgcaccaggagacatcagaaagaatg
(Seq ID No: 1143)

Homo sapiens threonyl-tRNA synthetase (TARS):

gcgccctttcgattgcatcagctggtccagccgaggccaagtcgccccggtcgctagccacctcccacc
cgctcttggctcctctcctctagggcgctcgctttcggttctctcatcgcttcgtcgttcgccaa
tg (Seq ID No: 1144)

- 5 Homo sapiens ATPase, Na⁺/K⁺ transporting, alpha 3 polypeptide (ATP1A3):

cagcctctgtgcggtgggaccaacggacggacggacggacgcgcgcacctaccgagggcgcgggcg
tgcagaggctcccagcccaagcctgagcctgagcccgccccgaggtccccgccccgccccgctggc
tctctcgccgaggagccgccaagatg (Seq ID No: 1145)

- 10 Homo sapiens chromosome 11 open reading frame 46 (C11orf46):

cgtcctctcagtggttagcgcggggactggctgggaagcggtcggtcgagtgtggcctgtgtggact
cgcatcttggccgaagccgggaggagagctcaagctaagggtgatcagcccatgacctaaacc
tccagacaaaataaaacggaaaatttgctagaatcaagaatg (Seq ID No: 1146)

Homo sapiens chromosome 17 open reading frame 45 (C17orf45):

- 15 tgaccttttcattcccgttggttatggaggtaggctctctaggaatctgggagtagtagctgggggg
caagagcaaataaagagctcgagcttctgtggtctctggggagatg (Seq ID No: 1147)

Homo sapiens AHA1, activator of heat shock 90kDa protein ATPase homolog 2 (yeast) (AHS2):

- 20 gggccttctgagcagtttctgggagctgcgaacgcgcgccccggggctcggcggccggaaacgctg
gcttcggagccttaggcgcgcggtcttcttctgttttccgccagtcacgcgcgcatggccaag
tggggccaggggaacccccactggatcgtggaggagcgggaggacgggaccaacgtgaacaactgg
cgctggcgcggtggtggcggtcctcttccgggatctggggaggggccgggcccgggagccggggc
tgccctgggggtctgtgcggggccgccccgggggagggggcagggggccgccccccctcagctgctgg
acgcaggggtcggtccttcgctctcggtcgggagagtccttgagtacggagaccggctaggaggg
25 ttgcagctgcctctttttgaaagtgggttgggccccaaagagtgaactccgacagaccttccact
cccaccgtctgtggcctgagggccttcccttctcctcccgccccccctctggatgtttcggggag
ttagaaggagctggattgagagactgtgttagggcgggggtatggaacgtagtggaaaggcgag
aaatttggtatctcagttcgcgccaccgccgagggcgctcccgcgagccgggcccctctgtgagtga
gacaagctccccttctttacgcgcctcacctggcgcggtgggagaggtcggcagccctccgccc
30 agaacctccggaaggatgtcctctgcctcgcctctggccggggctgtggtccctccaggccgt
cgaggggatgctgaggccggtccccagaggagcatgacttggctggtccggaggagctctgagggc
atgggcaatcttggctcgtgcaacctcagcttccagagttcaagcgagtctcctgcttcagcctc
atgagttagctgggactacagatgcgtgccactacgtccgtctgatgtttgtatttttagtagagac
agggtttcaccatgttgggtcaggctgctctcgaactccagatctcgtgatccgccccgctgggcct
35 actaaagtgtggtgattacaggcgtgagctagatctgacttctagtgtcctagccttggcccgat
ggacatgtcatttctctcagctcgtttctgtcccctaaagtgagaatattgcttgggaagattaca
ttagacgatgtatatgcgaagacacttgatagctggtattgtcatgattctgattagttcactact
gctactttccctgtggccttaggctttgcctatttccagtgggcgagctagctagatcctcctcct
taaataagccagtggttttaagacagaataactacttgcatagtggaacaataatcttaaagaact
40 gagcaggatgaaaagaatttgatagaaagcaggtttgaggagcacattggaggttggcaggtttcg
aggctgcttgagaggacttgggccgatctgggctgggcttgagcgtgacctggcaccaggcagg
tggatcccagctggggcttccattcacgacttctgggtccctggcaggacagagcgggatgccacc
agcttgtccaaagggaagttccaggagctcctggtgggcatcggttggtgagaatgacgctggccgc
ggcgagatcaacgagttgaagcaggtggaaggggaggcttcgtgcagcagccgcaaaggaaagctg
45 attttcttctatgagtgaacatcaaactgggctggaaaggcatcggttaaagaatctggagtgaag
cacaagggttgattgaaatacccaatcttctgaggaaaatgaagtagatgacactgagaattta
caacgggaatg (Seq ID No: 1148)

Homo sapiens GrpE-like 2, mitochondrial (E. coli) (GRPEL2):
ctgcctctcagcccaaattggaacatg (Seq ID No: 1149)

Homo sapiens xyloside xylosyltransferase 1 (XXYLT1):
ccgccccctttcatggccgcccgcctggcgccggggctaagtggccgcccggcggtccgggtaccgagg
5 gctctcccgcgttgctggcaccgctggcgccgggtctcgtagcgcgcatg
(Seq ID No: 1150)

Homo sapiens chromosome 7 open reading frame 60 (C7orf60):
cctcctctggctgctgcctccgcagctccctcctcctacccacctcctccatctggggagcgctct
gcgggggcctgagggggcgccggcgccggcgccggctgcgatatg (Seq ID No: 1151)

10 Homo sapiens tetratricopeptide repeat domain 39B (TTC39B):
ccctcctttgcgctgggctgagccagagccgagagcaggggtcggctctgagttccctgcttggt
ttttgggtggcagcagccagaggaggaatg (Seq ID No: 1152)

Homo sapiens motile sperm domain containing 2 (MOSPD2):
15 cacccttctctgtctacctctgggcccggactgccgggtgatgagatactcggtcggcgacggtaga
acgggcgacggcgacaaccgcaatcacatccacgacgggtgatcatg (Seq ID No: 1153)

Homo sapiens major facilitator superfamily domain containing 6-like (MFSD6L):
ggccccctttcgggtccaacggcaggacctgggggctgtggccggggggcgccggttgacctggtgacc
gcggcgccgccccagaccggggggcgagtcctcactcgctccgagccccgggtcccccaagcctccct
20 cccgggtacctggggccgcccgcctcctgcgccagctccgccctccgctcggcccaggcctgacag
agccccgcagccatg (Seq ID No: 1154)

Homo sapiens consortin, connexin sorting protein (CNST):
cttcctctctagccgcccagtgctctatgctccgcgggtcgccggccgcccagcctccagccggccagc
cgcgaggggtgcgagagggaggcgggggcggaaggcgagaggtgtctcctccaccggagccaggg
25 gagaccgagcaagctccgtgacagcacgtcggccgccatgtcgccgagtggggctggaaacagac
ccggcgcccagcggtagccctccttgccgctccgattcccagacatggaaggtctttaatgtaact
ttaaatgggttcaccaaaggatgctctaag (Seq ID No: 1155)

Homo sapiens zinc finger protein 92 (ZNF92):
gggcctttgtctctcgtgcagccggcgctccacgtctagtcttccactgctctgcgtcctgtgctg
30 ataaaggctcgccgctgtgacctgttacctgcaagaacttgagggttcacagctaagacgccagg
acccccctggaagcctagaaatg (Seq ID No: 1156)

Homo sapiens DnaJ (Hsp40) homolog, subfamily C, member 18 (DNAJC18):
cccccttctctttcagcctcgggcacgggggaggctcggcggacctgctgattgggaaccgatatg
35 (Seq ID No: 1157)

Homo sapiens polymerase (RNA) I polypeptide D, 16kDa (POLR1D):
cctcctccctccttccgctcctccgcgccttccgctcgggtcgggtccttgcttccctgcttcgcctccgc
gcctcgcgctatgggacagagccccgatccgccagcaccacctgaggatccagaaaccgccccag
cgatg (Seq ID No: 1158)

40 Homo sapiens ring finger protein 182 (RNF182):
acctccctcccctcccaggcgccgcccgcagccggagcgggtcccgggccctgggcccgcgcccggcc
aggaagaaataacttgtgttggtgcatttccagggtgctaccagagctcaaggctgtcacctggt
cttgcccagaagagccgttcttagaggcaggacttgatgaaggcttctcctgctgatggaataggtt
tgctagagctggccttggaattagaaccttcatgtggcctttataaatatgcgtttgagacagag

ttatatgcagaagttgaaaatgcctggaagatttctggtttctttcactacttatcctgccttttt
gcatcgctgccagatttgatgatatgatattcagaggggcaccttaatcaaagccattcttcaac
aagaccacactggcataagattgcacacataattcaagatg (Seq ID No: 1159)

Homo sapiens transmembrane protein 18 (TMEM18):

5 cctcctctgtggattctggccaggccgggttcggcggttgctgtgagagcgggcttcccaacacca
tg (Seq ID No: 1160)

Homo sapiens Hermansky-Pudlak syndrome 4 (HPS4):

aggcctctctgccgcgcgcgcaggtacggggcagaagtcgcaggtacccagctgctgccacattt
ctggtccagagtcccgaaccccgagcactgggatgcctggctactccgagccaaggcactgatgtt
10 tgaactggaaacttcaaaacgtttaataagagtcttcaggatgggtttgaactagacaagctagaa
atttctttagaacaccagctctagcatgcatctccacttttggctttcctggagaggagcttgaa
gaggtggttctgcagacagccacagtgatacttaggaaaccagaggaatggatttgacttttctgc
taggattctctgttatagtttctccctgagttgtaagaggcatggaaatatacatgaaactgaaga
acctgcaaggaagggaagtggaactttccatgctgagtgaaaactaaccaagtggcagttgtgact
15 gaaaacactgaaacctaccacgtccagattcactggattgggggatagaggaacggtcacagctag
ggagaaagaagtataccggaaaagaaaacctaataagagaatgaggatgactgcacagtagat
g (Seq ID No: 1161)

Homo sapiens PTK7 protein tyrosine kinase 7 (PTK7):

agctccttttctgagcccgccgcgatg (Seq ID No: 1162)

20 Homo sapiens kelch repeat and BTB (POZ) domain containing 6
(KBTBD6):

agttctcctgggcgcctagcattgtcgcccacgctgcagtagcggcttctgcggtccaagccagc
gggtcctgtgaaggcgagcagacgcggagaaaggacgcgggagtgagagaggggtgagtcagccact
gtctaaacgataacgggagcggtctgcggggtaggggtgaattcagtaaattgggctcgtgctgc
25 tgtctcttcggagacgctgctatcttagcgctcagcgagggaaggttgaggaggagccagagccggg
tcctgcagcgtttctcgccatcagcgcccgtcgccatctccaccatg (Seq ID No: 1163)

Homo sapiens sperm antigen with calponin homology and coiled-coil
domains 1 (SPECC1):

30 ctttctttgactggagcggaccgcggacgcaaccgcctcgccagccggagccagcgcgagctcg
gcacgggtggacaccgggtccgaggccggcaagccggctggtgcccagctcgccaagcatg
(Seq ID No: 1164)

Homo sapiens ST6 (alpha-
pha-N-acetyl-neuraminyl-2,3-beta-galactosyl-1,3)-N-acetylgalactosa
minide alpha-2,6-sialyltransferase 3 (ST6GALNAC3):

35 ggtcccttatttgatctgcgggaatgtgggctggagaggtcctgccgtggtaccagcctccagc
ctgccccaggaactgcccctgaccaggcgcgcccgctgctcggtggcaggagggccggcgagcgc
ccatg (Seq ID No: 1165)

Homo sapiens transportin 1 (TNPO1):

40 gattctctttgttccgcagccatttcaggccccggacaggaggcagtgccgcttcggccgaaggcc
cgagcggccgaggcgtctgaggatg (Seq ID No: 1166)

Homo sapiens heat shock 70kDa protein 8 (HSPA8):

cttccttcgttattggagccaggcctacaccccagcaaccatg (Seq ID No: 1167)

Homo sapiens hyaluronoglucosaminidase 1 (HYAL1):

45 ggctccttcctccaggagtctctggtgcagctggggtggaatctggccaggccctgcttaggcccc
catcctgggggtcaggaaatttgaggataaggcccttcagccccaaggacatcctggctgccatac

ctgctcctgacttctcagggctggcagtcacgactgggagggcatggcgcccacgctgggccttca
actgggacaccaaggacatttaccggcagcgctcacgggcactggtagggcacagcacctgatt
ggccagctcctcaggtggagggcagtagccagggaccagttccagggagctgcacgggcctggatg
(Seq ID No: 1168)

- 5 Homo sapiens STE20-related kinase adaptor alpha (STRADA):
agtcctcccgtcgccccactgcgcatggcaggttgctactcccctccagcaaccggctctggcg
gcggcgcggcagtaaaactgaggaggcggagccaagacggctcggggctgcttgctaactccaggaa
caggtttaagtttttgaaactgaagtaggcctacacagtaggaactcatg
(Seq ID No: 1169)

- 10 Homo sapiens transmembrane protein 161B (TMEM161B):
ccctctctttcgctgtttgagagtctctcggctcaaggaccgggaggttaagaggtttgggactgcc
ccggcaactccaggggtgtctgtgtccacgacctatcctaggcgccatg (Seq ID No: 1170)

Homo sapiens Usher syndrome 1C (autosomal recessive, severe)
(USH1C):

- 15 ggctctttccagctcctggcagccgggacccgaaggaacgggtcgtgcaacgacgcagctggacc
tgcccagccatg (Seq ID No: 1171)

Homo sapiens interleukin 12 receptor, beta 1 (IL12RB1):
cagtcttttctccttgctcagcttcaatgtgttcggagtggggacgggggtggctgaacctcgag
gtggcagagaggctcccctggggctgtggggctctacgtggatccgatg

- 20 (Seq ID No: 1172)

Homo sapiens Meis homeobox 2 (MEIS2):
atcccttctctcttttctgttcgcccccttctccttgcctctttttccctttccacccccctcctc
tgttctccctcacctcctgcgccccctcccccttcccggttctgacagtacgatgagctgccccca
ttacggcgggatg (Seq ID No: 1173)

- 25 Homo sapiens G elongation factor, mitochondrial 2 (GFM2):
ttttcttttctgttagatacattgccttttgcttaggctggcgctcgagacttgaggccgttgcaga
ctttggcgcggtcgcgcctcctgcttcaagagcccagcggtgagagctggcctgcggcacgcggc
ctaattgccagacagtaacagtttgaggatcaagatg (Seq ID No: 1174)

Homo sapiens lamin A/C (LMNA):

- 30 gagcctttgccccggcgctcggtagctcagtggtcgcgggagcgccgcacctacaccagccaaccca
gatcccagaggtccgacagcgcccgccagatccccacgcctgccaggagcaagccgagagccagc
cggccggcgcaactccgactccgagcagttctgtctcttcgacccgagccccgcgcctttccggga
cccctgccccgcgggcagcgctgccaacctgccggccatg (Seq ID No: 1175)

Homo sapiens calcium/calmodulin-dependent protein kinase II delta
(CAMK2D):

- 35 cgctctttctctcgccgcgcccgtcttgaagccgcgcgggctcgtgagcagcgcgaggccgccaagg
tgctcgtcttcgcccagcgcgtgcgccccgcggagggaagccggcctcgggcgcgcacgctcgt
cggagccccggcgcgccccgcgctgagcctgctgacagcgccgctgggctcaggctgtccgctc
tgggctccgcggcctcggccccgctgcaactccacctccgccccctcggaactccctcccctctgctt
40 ctactcctcctgctccagtgcggatcgtttcgcaactgcttgccactcgtcccgtgcctggctgtt
tttccatttcccgccccctcttcttgagtactttacccccctgcatttggggacagggactggaaa
aggggcgggtggagcgtccagtggagaagaaggaagcgaggcccgagaggaggaggatcggcgg
actgtggggaggagacccacgccaccctttctggatctcccctcccgccccgcctcgcgac
actccctcgcgggcgagctactttcggaccaggaagtaagagcgccctgggtgacagcgccgcg
45 gggccagtcccggggttagccgcgcgtctgctcgcttctggtccgtcgcgctcccagccagggcac
agcccggaaccgaggatg (Seq ID No: 1176)

Homo sapiens calcium/calmodulin-dependent protein kinase II gamma (CAMK2G):

ccgtctcctcctcttgcctccctcgggccggcgggcggtgactgtgcaccgacgtcggcgcgggctgc
accgcccgcgtccgcccgcgccagcatg (Seq ID No: 1177)

5 Homo sapiens interleukin 15 (IL15):

ttttcttttcgccaggggttgggactccgggtggcaggcgccccgggggaatcccagctgactcgct
cactgccttcgaagtcggcgccccccgggagggaactgggtggccgcaccctcccggctgcgggtg
gctgtcgccccccaccctgcagccaggactcgatggagaatccattccaatatatggccatgtggc
tctttggagcaatgttccatcatgttccatgctgctgacgtcacatggagcacagaaatcaatgtt
10 agcagatagccagcccatacaagatcgttttcaactagtggccccactgtgtccggaattgatggg
ttcttgggtctcactgacttcaagaatgaagccgcggaccctcgcggtgagtgttacagctcttaag
gtggcgcatctggagtttgttcttctgatgttcggatgtgttcggagtcttcttcttctggtggg
ttcgtggtctcgctggctcaggagtgaagctacagaccttcgcggaggcattgtggatggatggct
gctggaaacccttgccatagccagctcttcttcaataacttaaggatttaccgtggctttgagtaa
15 tgagaatttcgaaaccacatttgagaagtatcttccatccagtgtctacttgtgtttacttctaata
gtcatttttctaactgaagctggcattcatgtcttcatcttgggatgcagctaataataccagttgg
ccaaagcacctaactatagttatataatctgactctcagttcagttttactctactaatgcctt
catg (Seq ID No: 1178)

Homo sapiens protein O-fucosyltransferase 1 (POFUT1):

20 gtccctccttccctccccgactgtgcgcgcggctgggtcgggttccccgggcccagacatg
(Seq ID No: 1179)

Homo sapiens calpain 3, (p94) (CAPN3):

cactctcttttctctctccctctggcatgcatgctgctggtaggagaccccccaagtcaacattgctt
cagaaatcccttagcactcatttctcaggagaacttatggcttcagaatcacagctcgggtttttta
25 gatggacataacctgtacgaccttctgatgggctttcaactttgaactggatgtggacacttttct
ctcagatgacagaattactccaacttcccccttgcagttgcttcttcttgaaggtagctgtat
cttattttctttaaaaagctttttcttccaaagccacttgccatg (Seq ID No: 1180)

Homo sapiens PTK2B protein tyrosine kinase 2 beta (PTK2B):

30 agcccttttactcagccacagcctccggagccgttgccacacctacctgcccggccgacttacctgt
acttgccgcccgtcccggctcacctggcggtgcccaggagtagtcgctggagtccgcgcctccctg
ggactgcaatgtgccgatcttagctgctgctgagaggatg (Seq ID No: 1181)

Homo sapiens ST6 beta-galactosamide alpha-2,6-sialyltransferase 1 (ST6GAL1):

35 cttccttcccttctccagtccttccactgtgcgtcttctgtcccccggttcttccccagcggacccc
tctttcgagactccctagtgggggtccccagctcccggcgatcctgcccttgccgagcgcgttttc
tgagtcacctgggggaggggagtcctgggcagggccgggctggggaagacgcctggggcactgcc
cggcgtaacaaagggagccgataccgaccggcggtgggcgcggagcgggcggccgccaccgagcgt
gctgagcaaccgcagcctccgcggccgagagtgcagcagcaaggggagagccagttgctgcagagc
cctgcaaccagcagtcacaggagaagtggatgtcatggagcccagctgaaatggactggcccc
40 cttgagcctgtcccaagccctggtgccaggtgtccatccccgtgctgagatgagttttgatcatcc
tgagaaaaatgggccttggcctgcagacccaataaaccttccctcccatggataatagtgttaatt
cctgaggacctgaagggcctgccgcccctgggggattagccagaagcagatgatcatgacgcagtc
ctgaggtttaatggggcacccacagccaacttccaacaagatgtggggcacaaaaactaccattcgc
ctgatg (Seq ID No: 1182)

45 Homo sapiens ubiquitin-conjugating enzyme E2Q family member 2 (UBE2Q2):
ctcccccttcgcgcgccggctcccccttcgcgcgccctcccgcggagatgagggg-
gaagatg (Seq ID No: 1183)

Homo sapiens membrane magnesium transporter 1 (MMGT1):
gcttcttttgcgtgggctgctgctccttcggcatcatg (Seq ID No: 1184)

Homo sapiens PAP associated domain containing 4 (PAPD4):

5 cggctcttcgggtgtctttgacaggggttttctacgcgcgtttttcggcgactttttgctcttcgc
ttdttgccaaccgcccccaaccttctatatccttgacgcccctaccttttcttggtgctcctccc
ctggcagccgtgaggggggtagatctcagccggagccggagctgggcctagctgtcccacgggcc
accactacctcctttgggttcgggagaaagctacgaccaagtacgcccagctcgggccttagaactt
ctgaacgggcagtgcggttaggccctgcttagcccttcccggaggacacctgacaaaagaggaag
atagtcttgggacccttgcatggtgtttcaaaggggtggtgaagaactaaggtagaagaatacatgt
10 tcacttccagtgacaagagcatg (Seq ID No: 1185)

Homo sapiens chromosome 3 open reading frame 23 (C3orf23):

ctcccttctggtgtactgggtgggaggtggaactagtcggacaaagccctcgctcggacccttgc
cagaactcaattaatggatgcctcgaagttgacgtacatatattcagaaatg
(Seq ID No: 1186)

15 Homo sapiens mucosa associated lymphoid tissue lymphoma translocat
ion gene 1 (MALT1):

cgcccttttgcgcggctggcgcgccagccggccaggctcccctcggcaaacctgtctaattgggg
cggggagcggagcttctcctctgagggccgtgccgcgctgccagatttggttcttcgcccctgcc
tccgcggtcggaggcgagcggaaggtgccccggggccgaggcccgtagcggggcggggcgggagcc
20 ccggcagtcgggggtcgccggcgagggccatg (Seq ID No: 1187)

Homo sapiens UDP glycosyltransferase 3 family, polypeptide A2
(UGT3A2):

ctaccttaccacagccagtgcttttggcgactgaggtgcacaggggtcccttagccggggcgag
ggcgcgagcccaggctgagatccgcggcttccgtagaagtgagcatg (Seq ID No: 1188)

25 Homo sapiens sodium channel, voltage-gated, type IV, beta subunit
(SCN4B):

cctcctctcgtctcttgcggcctaactttcccgagccccgaccggcgcgagagctccggggtag
ctttgtggccgaacgccgacctcggggcgagagcgcggtgtgccagtatcccatccccgcgacc
cccgcgcgtccggagagaacaggactatg (Seq ID No: 1189)

30 Homo sapiens JAZF zinc finger 1 (JAZF1):

tcccctctgcctcccggtggctcctcgtctctcttccatctctctcgtccccctctccctccgtccc
gtcctcgcgcgtccccctcaccccgctctctccccctccccagccccctctctcctcacccacc
cggcctccctccctccctcgcggcgccggcgctcgcagagccgacaccaggggggctctcgatgta
gcaccatg (Seq ID No: 1190)

35 Homo sapiens chromosome 15 open reading frame 55 (C15orf55):

ttcccttcttggatccctgtgcactactggagccaggttactctgggtcctggacctgactgcc
tcattctggaggcttccagacagccacagttagtgcccaaacctgagaggatg
(Seq ID No: 1191)

Homo sapiens ras homolog family member C (RHOC):

40 cgccctctcttctcgcagcctgggaacttcagccggctggagccccaccatg
(Seq ID No: 1192)

Homo sapiens CTP synthase II (CTPS2):

cattctcttttcttttcttctctcctgagcgctcctgcagttcctggggcgtagtaggggatcca
caagcgtttgtgaccagtgaagttctttacaaggggtgagatctgcacgggaggacccgagcgaggg
45 tctcggcttgccaggaagccgggggttccccgggaagcggtggagttcacccgcgcactcgaagtgc

tttgcaaaattatatctgggtgttggcaccagccactattctgccaatg
(Seq ID No: 1193)

Homo sapiens PRP4 pre-mRNA processing factor 4 homolog B (yeast)
(PRPF4B):

5 agctcttttcttcttcttccacttcccctaccctccaccgtccgggagccgccaccgccgcc
gaggagtcaggaagttcaagatg (Seq ID No: 1194)

Homo sapiens molybdenum cofactor synthesis 2 (MOCS2):

gcgcttttgcggccgtgattcgggtcccgtgtcctagggcggtggtgcccgtgtgccaggtaagg
gtggcggtgtgtcgtgcgggctgggtgcggagccctcctcgacgtgtctctcccgcccttccct
10 ccacataccagccttgggtcagtcggacctcccactagcccccaacctggccggcggtcttgggtt
cgggggcgccccgccccgccccgggcccccttctgtctccgggctttactgcgactgccccagc
agaagtgcgggtcctctccgagaactcttgtcagctcacggcagcaaggacggactcgttctgaagg
cgctccaccttttatgaccacctctttcccagattattcgttttgatgaagctaaaattttaatc
15 taaaaagaaatgcacctcatggagaattcttgtgaagaactgtgcttcatctgtggatttctacac
ccttgatcatttgcacacctgtaattatttgcgtaaagagttggttgcacggagtgacaggttgaag
tattgtatttgcacaaagtgtgaaataacaggagttcggttcagagaccatttctgtgcctcaag
aaataaaagcgttgcagctgtggaaggagatagaaactcgacatcctggattggctgatgtagaa
atcagataatatattgctgttcgtcaagaatatg (Seq ID No: 1195)

20 Homo sapiens cat eye syndrome chromosome region, candidate 1
(CECR1): tttcctttttccggaggggagatg (Seq ID No: 1196)

Homo sapiens solute carrier family 13 (so-
dium-dependent citrate transporter), member 5 (SLC13A5):
ctgccccctactcgtctcgcggccagtcctcctcccgcgcatg (Seq ID No: 1197)

Homo sapiens armadillo repeat containing, X-linked 3 (ARMCX3):
25 agtccttcttgtcctgggtcgttgttcccgtctgagtaccagctccccactgcctgagggcgggcc
ggcctgcggcgaggaggaaggaagaggagaaggaaattgtccgaatccctgcagtggttccaa
gcctctcccggtggccagtccttctgtaggttgcggcacaacgccaggcaaaagaagaggaagga
atttaatcctaactcgggtggaggtcgatttgagggctctgctgtagcaggtggctccgcttgaagcga
30 gggaggaagtttctccgatcagtagagattggaaagattgttgggagtggtcacaccactagggaa
aagaagaagggcgaaactgcttgtcttgaggaggtcaacccccagaatcagctcttgtggccttga
agtggctgaagacgatcacctccacaggcttgagcccagtcacacagccttctccccagcctg
agtactactctattccttgggtccctgctattgtcggggacgattgcatg
(Seq ID No: 1198)

Homo sapiens armadillo repeat containing, X-linked 2 (ARMCX2):
35 cgtcctcctctgggtaccaactctattgcgcagctcgtgcccgtgcgtttaaaccaggcgaggagg
aggaggagaaaattccccagattcgggcaggcccgacccccacattccgtcctgttttgagagga
ggagggaagagaaataaacgtggcagcgcataagaaggccagcaggagactgctttccagacacct
ccggcccacacagccgttcacccccgtcttttcagtcctggaaaaggaattcgggtctgtccttag
gatgaagctctaactgaactgaagtaaggagaaacagccttgaatctttggaggggtctgtcttccct
40 tttgggtctgtgcaactgcagctacagtggaataaagcaaaactgctcttgatcccaggccctgccc
taagcctcagcagaacttgtaagcctaaactgaagagcctcaccggacgagcaggcatcccttaa
ccttaagcaatccagttccacgcccctggatcagtgataaaccagctgcaccatg
(Seq ID No: 1199)

Homo sapiens UBA domain containing 2 (UBAC2):
45 cgccctctggggctccgagcccggcgggaccatgttcaccagcaccggctccagtggtctgtga
gtaccggcctccgccatcctggctgccccctacacgccaccctaggcacctctttgaggaggctgg
ggcagcggggaccctcgggtttgccggaggtggtggggccgaccctccagaccggcggtccgaaccc

tgctagttcccggtcttgggggtcagcggaaaccgccccatttcggcctggagggggaatgggg
acaaagccccgcccgcgccccgacccacctggtatcccagggtgctctgccaggagtctcttg
gggcccgtgcaagtgggcaggtgccctggtgttctcgtgggcccggccccaggccctttgcggagcg
tgtgccgcgtgaaggaaggggcccgtcccccttaccatgccccattcttttaggcttgggggaccg
5 aactaactcccccgccccacttgcaaagttcagcctccgctttagaagctgacctctcagtttc
acttgatg (Seq ID No: 1200)

Homo sapiens cancer susceptibility candidate 4 (CASC4):
cctcctccctcggccggccctggggccgtgtccgcccggcaactccagccgaggcctgggcttctg
cctgcaggtgtctgcggcgaggccccctagggtagagcccgatttgggcccatg
10 (Seq ID No: 1201)

Homo sapiens protein phosphatase, Mg²⁺/Mn²⁺ dependent, 1G (PPM1G):
cgctccctcacagctcccgtcccgttaccgcctcctggccggcctcgcgctttccaccggcacctt
gcgctcggctcgcgccgcccggcctgctcctgccgcgcgacccccggggcttcggctccggcacggg
tcgcgcccagctttcctgcacctgaggccgcccggccagccgcccagc
15 (Seq ID No: 1202)

Homo sapiens StAR-related lipid transfer
(START) domain containing 13 (STARD13):
ctttcttttttaaaaatcgctgggtctgttgagctgtcctgggctgggtgccttgctctttgactga
gactggagacagacggcaacagccacaggcagactgaggtggcaataggaaatctgccgagatg
20 (Seq ID No: 1203)

Homo sapiens tubulin, beta class I (TUBB):
gattctcccgctcccagccccggcgacgcgcgccccgcccagcctgctttccctccgcgcctc
ccctctcctttctccctctcagaaccttcctgccgtcgctttgcacctcgctgctccagcctctg
gggcgcattccaaccttcagcctgcgacctgcggagaaaaaaattacttattttcttgcccat
25 acataccttgaggcgagcaaaaaattaaattttaaccatg (Seq ID No: 1204)

Homo sapiens cytochrome P450, family 4, subfamily X, polypeptide 1
(CYP4X1):
tttccttcttcccgcgagtcagaagcttcgcgagggcccagagaggcggtgggggagcccta
cgccagctccggggcgggagaaagcccaccctctcccgcgccccaggaaaccgcccggcgttcggcgc
30 tgcgagagccatg (Seq ID No: 1205)

Homo sapiens doublecortin (DCX):
ttttctttctctcagcatctccaccaaccagcagaaaaccggtgagtggggcttttaagtgattt
tcaagaagaatgtaacagatgtcaaagggaagcacaaggcaaagcctgctctctctgtctctc
tgtctcctcttctcctttttgccttattctatccgattttttccctaagcttctacctgggattt
35 tcctttggaaaagtctctgaggttcacccaaaatag (Seq ID No: 1206)

Homo sapiens protein phosphatase 2, regulatory subunit B', gamma
(PPP2R5C):
ttgtcttttttttttaactaaaatggaggctggtttcttgctttaaggagccattgcctttcc
cgctgaagtctagatg (Seq ID No: 1207)

Homo sapiens solute carrier family 9, subfamily B
(cation proton antiporter 2), member 2 (SLC9B2):
ccacctttccgggggaagccacgcgcaccaggcatcgcacgcggctctgcacccgcgcccgggac
ctgaaaccggcgaggggcacacggggctgccgctgcgggccccggaccaaccatgcttactccg
gagcctgtaccggcgccgacgggtcggacctccctgcgcggtgtcgcccagcgggttcgtgcgaaa
45 ggccggggccgactacacgcgggtgccgcgcctgagaccgtttatctgcagtcaacgcagcctcccg
gctcagcctgggaagatgcgcgaatcgggaaccccagagcgcgggtggctagaccgggctccgcgcg

ctccccacagcccctttcctaatacgttcagacggagcctggctcgacttcgccggagactgccaga
tctcgttctcttccctgtgtcatcttcttaattataaataatg (Seq ID No: 1208)

Homo sapiens hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor) (HIF1A):

5 caccctcttcgtcgttcggccagtgtgtcgggctgggcccctgacaagccacctgaggagaggctc
ggagccggggcccgaccccgcgattgccgcccgttctctctagtctcacgaggggtttcccgcc
tcgcacccccacctctggacttgccctttccttctcttctccgctgtggagggagccagcgcttag
gccggagcgagcctgggggcccgcgcgtgaagacatcgcggggaccgattcaccatg
(Seq ID No: 1209)

10 Homo sapiens interleukin 21 receptor (IL21R):

cctcctcttctccccactctgcacatgcggctgggtggcagccagcggcctcagacagaccact
ggcgtctctctgtgtagtgaccgtaagctcggcgtctggccctctgctgctctccctgagtgtg
gctgacagccacgcagctgtgtctgtctgtcggcccgatccctgctgcccgcctggta
ccttcttggcgtctcttctctgtctgtctgtggtggacacctgctggaggccagctgcc
15 cgtcatcagagtgcaggtcttatgacagcctgattgggtgactcgggctgggtgtggattctcacc
ccaggcctctgctgcttctcagaccctcatctgtcacccccacgctgaaccagctgccacccc
cagaagcccatcagactgccccagcacacggaatggatttctgagaaagaagccgaaacagaaga
tgaggcaatgaggctgcgagaggtagagtgattttccctcgggtgactcaactgggacgtagcaggt
cgggcagtcaagccaggtgaccccatg (Seq ID No: 1210)

20 Homo sapiens DDB1 and CUL4 associated factor 4 (DCAF4):

tggtctttccgggtccttgacgcttcgctccaactcctgcagagctgagccggaggggaatccgg
aaggacacgctgaacaggtctgactcccgggcagcacagcccgcctcacgattccggccacggtga
tgacgagtctccgtcaacctcgtctggcacagctgggacctcctctgtgccagagctacctgggtt
ttactttgaccctgaaaagaaacgctacttcgcttgcctcctggacataacaactgcaacccct
25 gacgaaagagagcatccggcagagagatg (Seq ID No: 1211)

Homo sapiens oxidation resistance 1 (OXR1):

ccgcctcttgtgaggcgcgcgagccgcctcccctgggtcaggtctgatgggcccgtgggcgcgct
agtgggtggccgccaccgccgaaaccgtcgacctcctgggccccagttccgcgtccagccccgcggc
agcatg (Seq ID No: 1212)

30 Homo sapiens cut-like homeobox 1 (CUX1): cccctctctat-
cagccgctcactccgtctcaatatgtctcaagatg (Seq ID No: 1213)

Homo sapiens atlastin GTPase 1 (ATL1):

ctcccttttctccccactccttcccaccagcgccacagcaacatcctcagagtctgagcgaactg
cgcccagcgcgggcacggagcctcccaccgccagcaacctgcggccccggagaaggcagcgagcgc
35 agtgacagcgccctaccgccaccagctcctggaccaccatg (Seq ID No: 1214)

Homo sapiens chemokine-like factor superfamily 5 (CKLFSF5):

ctgccttctctccgggggcccctgtgggcaagcctcctgcttcaatttcaggtttctcgaagtgcct
tcttgcctcctgtctgtttcccatcctgccagatttctgtttctcttgcctgggcttttggcagtag
ggggtgtgttggtgggcccctacgaagatg (Seq ID No: 1215)

40 Homo sapiens transmembrane emp24 protein transport domain containi
ng 7 (TMED7):

aggccttttccgcttctcttttacctccccaggtccgcccgtctgcgcccctcacaggaagccgga
gggtcgtctgatcccgaatctccacaggcgtgaacctgctctgctgtgtatctttgcggggtgg
cctgcgctgaggcctgccgcgcggtgagtcgcgcagacctgacctgcgtctcgcagctcgggt
45 tgaggccgcgcgccttctcggtatg (Seq ID No: 1216)

Homo sapiens ubiquitin-conjugating enzyme E2D 3 (UBE2D3):

cttcctttaccttctctcccatggtctccttccggttctcgatgcttctctgagcctaagggtttcc
gccactcgttcacccctccccccagctcatgatcctcctcctcccccgccctcctggtccaatctc
cgatctgttttagtaagaaggtgctgttccgagaagaaggaaaagggttgacacgtattcactcgg
5 ccccggaagtggaagcaagccgtctggcttcggcctcacatcgggtcttggtcgggacggcggc
gttggcggactgatccggcggtgaagagaggccgggaagttaaactttagaccaccacctccgc
tcttcccgctcacctcgcctccctcgggcccgaagcacgggtacagaggctgttggtggctttgc
cacgccacccccacccaccccgatcggcggtgtcttaaggacctggattcatcaggggctcttcg
ggcctgtgagtgctgatctgctcgtttttgcaaaaggcgctgtgtctggcagagctggtgt
10 gagacgagacaatcctgccccgcccgggataatcaagagttttggccggacctttgagcataca
ccgagagagtgaggagccagacgacaagcacacactatg (Seq ID No: 1217)

Homo sapiens zinc finger protein 595 (ZNF595):

tttcctctggctcctgcgagggcttggtttagggttcagctctctgcgttctcgggtccgggagg
cctcgggtgattcagccacagcctctgcctccggttgctctgtgacctgagggtattggacaatttg
15 tagctaagactcccggataccctgaagtcgggaaatg (Seq ID No: 1218)

Homo sapiens acyl-CoA synthetase medium-chain family member 2B
(ACSM2B):

tgctctcttccaaggctgtaggagttctggagctgctggctggagaggagggtggacgaagctctc
tccagaaagacatcctgagaggacttggcagcctgcagatggcctattgtgggaccttgtgatcat
20 gcctgaacatg (Seq ID No: 1219)

Homo sapiens SRSF protein kinase 2 (SRPK2):

tttccttttatagcaccattgaatcccagtcctaacagaagtactgcgaatcttgtggcctcattc
tgaacaaaagggttagagaagaaaaatctcttgatataaggcttgaaagcaaggcgaggcaatct
tggttgatgaatattttctgatttttccagaaatcaagcagaagattgagctgctgatg
25 (Seq ID No: 1220)

Homo sapiens synaptophysin-like 1 (SYPL1):

tgcccttctctgccaccgggctgctctggtctcgtcggtccctcctccgccccgctcgtcctgact
ctctctccctcctttcctcagaggatg (Seq ID No: 1221)

Homo sapiens thioredoxin reductase 1 (TXNRD1):

aaccctttcacctcagttttcttactccggcatttgcagcagagcgaaagggtggtcgagtcctga
aggaggggcctgatgtcttcatcattctcaaatcttgaagctctgctcggtgaaaccagacaa
agccgcgagcccagggtgaggagcacgcgggggacggcctgccggcggggacgacagcattgcgcc
tgggtgcagcagtgctgctcggggaagggaagatattttaaggcgtgtctgagcagacggggag
gcttttccaaaccaggcagcttcgtggcgtgtgcggtttcgaccgggtcacacaaagcttcagca
35 tgtcatgtggcttatcaggagggcagacttcaaaagctactaaaaatg (Seq ID No: 1222)

Homo sapiens minichromosome maintenance complex component 7
(MCM7):

tgctcttccgcgcggcgccgcggagagagctgcggccccggggggcggtgcctgggatccggagct
tcgctcgggccccgggaaaggcggcagtggtggtggtatcgcggtgtctctgggtgtgatggccaatg
40 gctggactggctcccgcctggtggcggaggaatcccagctgtgaagcggctggaatccgggccc
gtgcttctttgttactaagagcggaagcgatggcgggagcgggggtgggtgcggtggcgggggtg
cgggtggcggaggtcccgggtgaaatcaggggctaaggggacccaaagaaggcgggggatcatagggg
tggaagaaagctgagaaccttgagaccggagtgtagggggccaacgggggaaggcgctagaattt
taaactaaagtagggaccggaattcccctggggagatgttgatggccctgtgactgccacgggc
45 tctttattcttcgctggttagaaacagacttgtgaaaaagagttatgccactttggggagacttc
gaaaagggttaagaagttcttacaagagttctaccaggatgatgaactcgggaagaagcagttcaag
tatgggaaccagttggttcggctggtcatcggaacaggtggctctgtatgtggacctggacgac

gtagccgaggatgacccccgagttggtggactcaatttgtgagaatgccaggcgctacgcgaagctc
tttgctgatgccgtacaagagctgctgcctcagtacaaggagaggggaagtggtaaataaagatgtc
ctggacgtttacattgagcatcggctaattgatggagcagcggagtcgggaccctgggatggtccga
agccccagaaccagtagcctgctgaactcatgcgcagattgtgagtggctctctgtcgggaaagat
5 gtagggattggttctccaggatcttgtttgtgactgttttctccccttagtgagctgtattttcaa
ggccctagcagcaacaagcctcgtgtgatccgggaagtgcgggctgactctgtggggaagttggta
actgtgcgtggaatcgtcactcgtgtctctgaagtcaaaccaagatg (Seq ID No: 1223)

Homo sapiens pre-B-cell colony enhancing factor 1 (PBEF1):

tttccccctctccccctcctccgccgaccgagcagtgacttaagcaacggagcgcggtgaagctca
10 tttttctccttctctcgcagccgcgccaggagctcgcggcgcgcggccccctgtcctccggcccag
atg (Seq ID No: 1224)

Homo sapiens cyclin B1 interacting protein 1, E3 ubiquitin protein
ligase (CCNB1IP1):

ctttctttccctctccgttttgggtgggctggttgaagatgaaatccactgaggagggaagtccagc
15 accctgtgtgccagtcacagaactggcccatctgtagacccccctgaaaatcatatgggcttgattt
ggatattctcaacagaaagggttaaaggctgatggtacctaagcctggtacttgaattttgatca
agataagctgccttaagttctcttcattacacaaatgatcctagataattgatagatcctgtgggt
caactggatttctagatagaagctggattcatgtgatgccagaggagtaaaatttcaagagactga
aaccagatctgagtttcgctgttccagctctggacctctttgggtgctgtaaactcctggatatactgt
20 agatgagtactgcgtttttcttttatggcctctcttcagcttctggagacctcactatcctattat
g (Seq ID No: 1225)

Homo sapiens STEAP family member 3, metalloredutase (STEAP3):

ccgccttcgccgcggaccttcagctgccgcggtcgcctccgagcggcgggccgcagagatgacattt
attcatttttatgcatcctgggttctactggtcgtcccacctcagttcctgtagcaaagagacttga
25 gtctgagccactaattatcacccgtgaggtttcctccccgagcaggaagcagcaggccagagctgc
gctctctcagtgactctccaaccaagcatcagtcaccactcccgggtccagcccctgtggccaaga
gctggcgtgcaggctgcgggaggcagctggctgtgcaagaccctggcaggggccctcgcctcctgag
aaaccgagagtcagaaccaaagccaggctgtcctggttgagactgagccagaaagggtgggtcac
ctcacggtgaggctgtcagtgacctgagagcctcagaccctcacgtcagccggatg
30 (Seq ID No: 1226)

Homo sapiens nicotinamide nucleotide transhydrogenase (NNT):

tgttcttccgggttgaggcgcagcgcgcggggcccaagcccgggtctgccagcgcgacgtcctc
tcgcggccctcagggcacagcccaaggctgtcagcctcccggcccagtgatttgccttcaaggaaa
ctggggagtcagaaaattgggaactcatatcaacatg (Seq ID No: 1227)

35 Homo sapiens SHC

(Src homology 2 domain containing) transforming protein 1 (SHC1):
gtccctctccctcccaggacttctgtgactcctgggccacagaggtccaaccaggctaagggcct
ggggataccccctgcctggcccccttgcccaaactggcagggggggccaggctgggcagcagcccct
ctttcacctcaactatg (Seq ID No: 1228)

40 Homo sapiens bromodomain containing 8 (BRD8): cggcccttcca-
gaccgtctctcctcagggttgagacttcggggccaagatg (Seq ID No: 1229)

Homo sapiens ring finger protein 13 (RNF13):

tcgcctcttttagtaggtcgggtgagtgtagtgtgcagggaagagacgcgtcagcgcagggccagg
cccggccgggggcagcccggcagccgaatcttgggtactctgtcccaacagccggagcagatcag
45 accgaccggccctgcccgtcgggtcccgcgcctccagaccctacggtctccgtttctaggggcac
atggttagcggcaggcgcacagccaatccactttgcccagcctgccccttctctgccaagagca
gcttcttcagccgcgtccagttccgcagacgcctgccccacctgctcttcccttccagggaaga

cggatcacgcggccaagaacgagactcgcaaactgggcatttctccgagccgggctagagcaagta
gcgagactccgcgtgagagtgggaaagagccttaacaggcaaccatgttgcccagtggggttttctg
tgcctttgggtgcggaccaatgaggcgcgtggggcgggacttccgcttcgcctaggtgttgctcgtc
cctgctagtactccgggctgtgggggtcggtgcggatattcagtcataaagcagggtagggaactt
5 ctcccgcagcgacgcggctggcaagactgtttgtgttgcgggggccggacttcaagagagaaagag
agagtgggcagacatcgaagccaaacagcagtatcccggaagcactcatgcaactttggtggcggc
cactcagttttctctgccagtgtctggtgattttacaacgagatg (Seq ID No: 1230)

Homo sapiens aldolase A, fructose-bisphosphate (ALDOA):

10 ccgcctcctgcgccccttccgaggctaaatcggtgcgttcctctcggaacgcgcgcagaaag
gggtcctggtgacgagtcctcggttctctccttgaatccactcgccagcccgcgcctctgcccgc
cgaccctgcacaccgccttctctgtgccaggaacttgctactaccagcaccatg
(Seq ID No: 1231)

Homo sapiens LY6/PLAUR domain containing 6 (LYPD6):

15 cgctccttccctgagctcccgggctccggcagcgcgtggcgggcgccgcattgcacactctggg
ggcgccgcagtggtcgtgggatggggcagcgggctgcagctggcgccggaatccgcgcgcagccc
gggtgcaagttctctcctgttgccctgagtgccactcccaggccctctgtatgagtgacacttca
gtctgccatg (Seq ID No: 1232)

Homo sapiens butyrophilin, subfamily 3, member A1 (BTN3A1):

20 cagtctctgctttcttttcccttcttccagaaggagatttaaccatagtagaaagaatggagaac
tattaactgcctttcttctgtgggctgtgattttcagaggggaatgctaagaggtgattttcaatg
ttgggactcaaaggtgaagacactgaaggacagaatttttggcagaggaaagatcttcttcggtca
ccatacttgagttagctctagggaggtggaggtttccatttgggaattctatagcttcttccaggtc
atagtgtctgccccccaccttccagtatctcctgatatgcagcatgaatg
(Seq ID No: 1233)

25 Homo sapiens lipoic acid synthetase (LIAS):

ctgtcctttcccgaggattagcgatccctcaaccctgcactgcgctagtcctaaagaggaaatg
(Seq ID No: 1234)

Homo sapiens C-type lectin domain family 7, member A (CLEC7A):

30 gattctcttttgtccacagacagtcattctcaggagcagaaagaaaagagctcccaaagctatatc
tattcaggggctctcaagaacaatg (Seq ID No: 1235)

Homo sapiens CD247 molecule (CD247):

actccttttctcctaaccgtcccggccaccgctgcctcagcctctgcctcccagcctctttctgag
ggaaaggacaagatg (Seq ID No: 1236)

Homo sapiens myeloid zinc finger 1 (MZF1):

35 aagccttttctccattttgcggtctaggaagtagcagaggccccttctgtaggaggttgccatgga
gacgcggtggggcaccgacggagttctaataacgcggcgtgattggtgcaggatcctgctaattctca
ggaaggcccgtagagaagtgaggaacacgtggtggggggcatgcgcgatctggtaggcggtgctgc
cgtctgttgtagctgagaggcttgccatgccgacgcacggattcagaggcgggagcatgggaaga
agcggccaggagtatgacctgatcattgcgaccaccgctaggggaaggaggagagggtgtagaaa
40 cggggacgagggtgggggaagggaaggagggcgtcagagctggtgcgcggagcatcctgggagacg
tagtcagcgggagggggaagtcgaagactgcgcgtgctcaggagcgcggagcggcccgtgagcg
cagaggggcagacactggcctcagatacctgacctggtaccctctatg (Seq ID No: 1237)

Homo sapiens E2F transcription factor 6 (E2F6):

45 cctcctctttttccgtctgcgtcgggagctcccgggcacgtgaggccgtgccgcgtttactggcgg
gcgggacggcctagccgggcggcctcggagggaagccgcggaccccttaggtgctgggccccttgg

aaatcggcgcgctggggggcggtgctcgagctgagcgcgagagggcgaggagctcgtggggtgcga
ggggagcaggacgcccgccgggcagcatg (Seq ID No: 1238)

Homo sapiens zinc finger protein 621 (ZNF621):

cgcccttccggctcggcctttagttagtgaccagctcctcggcggttctgcagagcgtgggtttcag
cgagttctacgtgccaggtccgcccgggtgccggcttctcgtgcccctggcggtcgtcagcccc
cactaccctgaacttgggtcccaatggcgggccgcccctccttcaccgcgaccgtgggcatctggg
5 cctcgccgaagccgtcaaggtgggtgctcgggttctagagcccggtgtccagccctttgccaccga
ggcctgatcctcttttctgccctaaagaacttgccctgacagcctctggctcccgctcttgaggat
cttgcttgtccaaaccagaagacagtgcaggaagccaggggacatccgcatg
(Seq ID No: 1247)

Homo sapiens family with sequence similarity 73, member A
10 (FAM73A): ccgccttctccatg (Seq ID No: 1248)

Homo sapiens RNA binding motif protein 43 (RBM43):

ccgcccttttcttctgtagcctccaaggagctggaacaaaaacgaaaccaaaccctgcctgctcg
ctcctctccccatcgccctgcgttccgctggttggtgggcttctgcgggcgctgagggcgctctcc
cctccgcatg (Seq ID No: 1249)

15 Homo sapiens spermatogenesis and centriole associated 1 (SPATC1):
caccctccttcagcccaggcaaggcctggggcctgggcagcctccaggtgcagtgcctcccgtg
ggcgcacccttgccactgccccagggcatg (Seq ID No: 1250)

Homo sapiens carbonic anhydrase XIII (CA13):

ctttctcttccttcacccccgagggacatg (Seq ID No: 1251)

20 Homo sapiens transglutaminase 2
(C polypeptide, protein-glutamine-gamma-glutamyltransferase)
(TGM2):

cgctctccgcctcggcagtgccagccgcccagtggtcgcacttgagggtctcgccgcccagtggaag
gagccaccgccccgcccagcatg (Seq ID No: 1252)

25 Homo sapiens NOP2/Sun domain family, member 4 (NSUN4):
atttcctttcccttttttctgctcgtgtcccgccgggtggcgctcaccacctccccggaacacgcga
gtctcctgtcgcggttcgggtcggaattaccccggtggagcacgccgatatg
(Seq ID No: 1253)

Homo sapiens mitochondrial ribosome recycling factor (MRRF):

30 gagtctttccttagtaacctgggcgatagctgtggatgtttccaaggattgtcttcagtcag
(Seq ID No: 1254)

Homo sapiens PHD finger protein 17 (PHF17):

cttcctccataacaagccaaacgccagaccgagagtgccctccgtgcgcgagtgcccgggtgtgtgcg
cgccggcgagagcagggggcccgccgggtccccgcccgcgcggcccgaactcatgcagctccgag
35 cgagcgagcggcgcccagcccagcgccctcgccgaaccctccgcagcagggtgcctgctgtttcc
cggggagatcatg (Seq ID No: 1255)

Homo sapiens prolylcarboxypeptidase (angiotensinase C) (PRCP):

cctccttttgcgcctcccacccgcactgcagctctccagcctgagccatg
(Seq ID No: 1256)

40 Homo sapiens proteolipid protein 1 (PLP1):

aagcccttttcatgtcaggagaagaggacaaagatactcagagagaaaaagtaaaagaccgaagaa
ggaggctggagagaccagatccttccagctgaacaaagtccagccacaaagcagactagccagccg
gctacaattggagtcagagtcctcaagacatg (Seq ID No: 1257)

Homo sapiens coiled-coil domain containing 80 (CCDC80):

cagccttctcactcctcactgagtcactctgaacgtgctaaaatgggaaggaggcggtgttttgc
tgatctgttaaattcttagtgaagtttcttgatttccagtggtgctgtgtttgagtttggttt
ggagcaaaactgaggtagtcctaacatttctgggactgaatccaggcaagagaaagaagaaaaa
5 agaagaaaaagaggaggaaaaaggtagggagaaataaaggaggagagaaagcacagtgaagaaaa
aaaaagtccttttgcacatcacattcctgtgttttccctcagcctggaaaacatattaatcccag
tgcttttacgcccggaaacaaagagactaagccagactatgggggaaaggagataagaaggatcc
tggaacttttaaagagggaagagtgagattcagaaatcgccaggactggactttaagggacgtcct
gtgtcagcacaagggactggcacacacagacacacgagaccgaggagaaactgcagacaaatggag
10 atacaaagacttagaaggacagctcctttcacctcatcctacttgtccagaaggtaaaaagacaca
gccagaaagaaaaggcatcggtcagctctcagatcaggacaggctgtggatctgtggcggtactc
tgaaagctggagctgcagcacacccctttgtattgctcacctcggtaaagagagagagggtgg
gaggaaaagtagttcatctaggaaactgtcctgggaaccaaacttctgatttctttgcaaccctc
tgcatccatctctatgagccaccattggattacacaatg (Seq ID No: 1258)

15 Homo sapiens chromosome 20 open reading frame 44 (C20orf44):

cgaccttttgcgcctgcgcccccttgccagtctttcgccggcaaaaggaggacgtagaaaagg
gacaccggaaactcactcttcacccggaaatggttattgaggaacatg (Seq ID No: 1259)

Homo sapiens tryptophanyl tRNA synthetase 2, mitochondrial
(WARS2): cgcccttctcaagatg (Seq ID No: 1260)

20 Homo sapiens myotubularin related protein 2 (MTMR2):

ctttccctgtgctgccctgccgcgcatggagaagagctcagctgcgagagtcttggctcccag
ccggcgccggctcgcccgccagcgtggactccttgtccagttaatgtgttaagagccattgacat
ttgaagatcatcagaagtgaagataaaacatctcaaaaattataattgcctccacttctcattcag
agaattcagtgcatataaaaatcagcttctgttgatcatcagattccatttcaacttctgccgaca
25 acttttctcctgatttgaggagggagctctcgctctatcccctaggctggagtgattggcgccatc
tcggctcatttgcaacctctgtctcccggttcaagcgattctcctgcctcagcttcccgaggagc
tgggattacaggctcctgagggagcttaacaagttagcagaaatg (Seq ID No: 1261)

Homo sapiens reticulon 3 (RTN3):

cgccctctagctgcgctcggtgagtcagtcagtcgtgcgagtcctgcctcggagcaggcggagt
30 aaagggacttgagcgagccagttgccggattattctatttcccctccctctctcccgcctcgtatc
tcttttcaccccttctcccaccctcgctcgctagccatg (Seq ID No: 1262)

Homo sapiens G protein-coupled receptor 56 (GPR56):

gtccctccctctccgcactagctgtctgccctgccctgccgtaggagatgggctgggagcctccca
cgctctccagctcactcggcaggcagcggggaccagggtggcagggttaagcctctgggggtggat
35 cctgaaagggtgggtccagccgcctggccctgcgtgggaccctccacctggcagcagacagggtctcg
ctctgtcacacaggctggagtgagtggtgtgatcttggctcatcgtaacctccacctcccgggtt
caagtgattctcatgcctcagcctcccagtagctgggattacagggtggtgacttccaagagtgac
tccgtcggaggaaaatg (Seq ID No: 1263)

40 Homo sapiens immunoglobulin superfamily containing leucine-rich re
peat (ISLR):

gctcctccctgccgcctcctctcagtggtatggttccaggcacccctgtctggggcagggagggcaca
ggcctgcacatcgaagggtgggtgggaccaggctgcccctcgccccagcatccaagtcctcccttg
ggcgccggtggccctgcagactctcagggttaaggtcctctgttgccttttgggtccaccttagaa
gaggctccgcttgactaagagtagcttgaaggaggcaccatg (Seq ID No: 1264)

45 Homo sapiens glycoprotein M6A (GPM6A):

atttcttttccccatttttaaagcaagcaagacttgtgaatcatagtgctctgctcctgggatt
cagaccaaatttccccccaaaattctcaggctatttgtttgaatacctgcttacagtgggtacacaa

tgggcagctttgagaagaaaaattgataatcttcacggaagagtaatttgaatgaaattacacttg
acagcctgtctccaagcaaaacaagaggaacgagggagcctgagctaagctctgaggacttgcccaa
gccactgctgttgagcttcccaggaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaac
accagttttccaacatctaattgagcttttgattaattccgtgtaccagattctactgaagaaag
gtagccatg (Seq ID No: 1265)

Homo sapiens splicing factor 1 (SF1):

ctccctctttgtgctctcgcgccgcgcgcgcgcgcgcgtgagaggacgggctccgcgcgctccg
gcagcgcattcgggtccccctcccccgaggaggttggaaggagaagccgcgcagaggaaaagca
ggtgccgggtgctgtccccgggggcgccatg (Seq ID No: 1266)

10 Homo sapiens cell cycle associated protein 1 (CAPRIN1):

ccgcccctcgcgaccagagggctgctggctggctaagtccctcccgctccgggtctcgccctcac
taggagcgggtctcggtgcagcgggacagggcgaagcggcctgcgcccacggagcgcgcgacactg
cccgaagggaaccgccacccttgccccctcagctgcccaactcgtgatttccagcggcctccgcgcg
cgcacgatg (Seq ID No: 1267)

15 Homo sapiens hypothetical protein FLJ90297 (LOC388152):

ctgccctcttgcggtgccccggccacccccgggcggttgtagccgggtgcgcgggggtggctggggct
acgtgcagagctgtcgcgaggccggaacagcagcgggtgaagccccctcggtcggccgagaccgccc
tgcccattgctcgctcgggttgccgcgcgtttagccgcagccgctgctgccgcgcggggggagag
gcagcctattgtctttctccgcggcgaaggtgaggagctgtctcggtcggccccggggggagccc
cgggagccgcacggagatggaggaggacatctggacagtgagcaggaggcgccctcggcccatg
(Seq ID No: 1268)

Homo sapiens kelch-like ECH-associated protein 1 (KEAP1):

cgccctctcccgccctccttttcggggcgtcccgaggcgcgtccccaaccgacaaccaagacccgc
aggccacgcagccctggagccgaggcccccgacggcggaggcgcgcgggtcccctacagccaa
ggtccctgagtgccagaggtggtggtgttgcttatcttctggaaccccatg
(Seq ID No: 1269)

Homo sapiens F-box protein 38 (FBXO38):

ctccctctcaaccacaataaacaggcgagggtcggcgtaggtactttgaactcaagtaaacaaaag
ggaagattttctcgttgatactggagactgcacaacaatg (Seq ID No: 1270)

30 Homo sapiens musculoskeletal, embryonic nuclear protein 1
{MUSTN1):

agatcttttccagcagctgctgcctgccagagaggcgcttcagagaccagcgcttacacaatac
ccaccatg (Seq ID No: 1271)

Homo sapiens QKI, KH domain containing, RNA binding (QKI):

35 cctcctctccggcgggcgggcgggcgggcgggcgggcgaggtagctgcggagcctggaatatg
(Seq ID No: 1272)

Homo sapiens protein phosphatase 1, catalytic subunit, beta isoform (PPP1CB):

40 ggccctctcttgtttattttatttttccgtgggtgcctccgagtgtagcgcgctctcgctacc
ggcggggagggggtgggggagggccgggaaaagggggagttggagccgggtcgaaacgcgc
gtgacttgtaggtgagagaacgcgagccgtcgccgcagcctccgccgagagaagcccttgttc
cgctgctgggaaggagagtctgtgccgacaagatg (Seq ID No: 1273)

Homo sapiens methyltransferase like 21B (METTL21B):

45 cagcctctaccccgctccggatccgggatctgagcgccggccgcggtgccaggcactcccttggc
gggccggatg (Seq ID No: 1274)

Homo sapiens adaptor-related protein complex 3, mu 1 subunit (AP3M1):

cgcccttctcggcttctccagcttcggtaggagaggatccggcgccgaatcactgactggcacagg
tggtgggatatgtgtcacttgggtcaccaggtggagtgagtgaggcgcaatcttagctctctaca
5 gcgtcgatcttctcctggtgctcaagcaattctcctgcttcctcctgagtacctaggactaca
gaaaatg (Seq ID No: 1275)

Homo sapiens muscleblind-like splicing regulator 1 (MBNL1):

cagtcttttactgcagctgaatgagttgtggcgcccaaatgctcccatgacaaggagctgacaa
gttccattttccgtcgcgggcatcttggaatcatgactcccaaatgccttgggcaacttggtcgac
10 agtggggccgctctgaaaaaaaaatgtgagaggttgggtactaagaagtgcctttcctgacgtctc
tgctgcttggaaaccgcttctagagcagctctctgcttttgccttgccttgcctgagctagactgtga
cgacagcacatccaccctccacctctagcccagacacccccatttctacttataatcaagagaaaa
gctctaagtatctggcattgccttaggctgcttttagtggttaaaagaaaagtttgctgaaaaagtaa
gatatcttctgccaggaaatcaaggaggaaaaaaaaaatcattttctcgattttgctctaaactgc
15 tgcattctgtctatgccaaactaatcaataccgattgcaccaccaaactccattgcaaattcagctg
tgaggagattccctttcagacaactttgctgaaagcagcttggaaattcggtgtcgaaaggtctgc
cacgttttcatgcttgcattttgggctccaaattggcactgggaaggggttactgagagcacaagg
ctgataccaggccctacttttaaacggttcatctacttacaatcctagtatttctctaaaaaccaa
acctctttgaattaacagtttcatgctgtgaatttctagtgggagatcttttccttgatattgacg
20 acacaattttccatgtacttttaaacgaggagtggggaaaagtattttgaggggacattttcatc
atcagttcagcttttttttttgggtgttgccttttttgggggggttgggtttgttgggtttcact
gaaacatttaactacctgtaaaatctaaacatg (Seq ID No: 1276)

Homo sapiens lipid phosphate phosphatase-related protein type 1 (LPPR1):

cagccttttgctcttttcctttcattaaacaaacaggagatcctgaaacctggacctgtgcaagct
25 gcagcgccaggaggaggcagcgagggaagcagagcgcgggatggcgccagcgggcatctgtgatc
ccgcgcacctccgccccacggcgcgcgacaaacacggacacacacatacacacactcgcgacaca
cactcgacacaaacacacactcgtaacgcccgcgcgctcgctcgccggttgcctctcccacgcaa
gcggaatgcagcagcgctggagagcgtgtctcgaccgcccgcctgaatgtacctcgctcccggga
30 gccggacggccagtagggcgactggaggacgtccgctgcgggagcctggacagtttttgacgg
tgagctcttgctatatgtgtgagaaatg (Seq ID No: 1277)

Homo sapiens muscleblind-like splicing regulator 2 (MBNL2):

ctgtctttgcttcatctgaaggtaaaaattttccagatacggcagacggctttcagagtacaat
aaacaggggaatgagaactatttacatggaagtttctttctcatgatgcggtggagaagcctcgcc
35 acttggttctgccagatgttcttgggttactgtaaatgggaaggacaggcagagctaaacaagg
ttatcatttaaaagtgcctgtgtgaagtcacttttgctggaaaactgcagcttgggagctttctt
gtattcacatcccactcttctgtcaagtacactttaccctgaccttatgagtggatgaagatacct
cagttgtctgactttgccaatgtcttaatttcagaatttaaaaaggggaaagaaaaacatcctgct
aaaatatgaacatctgagtgcttattttccaacatcgtaataagctgtgagcgtcagcattaaat
40 attctccaaggagtgccatgatattgaagtcactttattaataacagctgtatctgcaaaacagt
caagagactcggacggtgaaagccagagatgacactgagcatgcttttattgcggcctaccatctt
taagtgggacatattgattgatgagtgattgcctgtccatacactctctcatcatcctgttccttg
gattggacttcactaagcaatttatcactcaccttcagacttacatgtgggagttttcacaacagt
agttttggaatcattagaacttggaattgatttcatttaacagaaacaaacagcccaattact
45 ttatcacatg (Seq ID No: 1278)

Homo sapiens chromosome 3 open reading frame 25 (C3orf25):

gcgcctttcgcacgacttgagttacggtttatctgataccgcggtaccctacgcaagcaagccc
acatcgacacacattcacacacgcccttcagcacccccctcccagcaccacgaccatg
(Seq ID No: 1279)

Homo sapiens testis expressed 19 (TEX19):

cctcctcctttccctgggtgccacatgaacagagacaccaggatgctctcctgagaccacagcaa
ctgcagaagctgaagacatttccagaagttcaagcttccaccctctgcagggtcccccactgagctgg
gacccaggtcatccacccccaccccaaatccctggataggaaacccctttctcctcctgctccttgt
5 ccccttcacccctgccgccagcatcctactggcctcagcacctgtggccagaccgtccaagatcc
tctgaaggcccagctcttgctgtccaccccggcagtaggcaggcagcctggccatg
(Seq ID No: 1280)

Homo sapiens protein kinase C, beta (PRKCB):

gcctccctccccgcagctggggccagcggtgccaaagcgcagctggacgagcggcagcagctgggc
10 gagtgcagagccccggctccgcgcgcgcggccgcccagagccggcgcaggggaagcgcggccg
cggtgcagcagcggccgcgcctccgcgcctcccgccgcagcccgcgggtcccgcgggccccg
ggccggcacctctcgggctccggctccccgcgcgaagatg (Seq ID No: 1281)

Homo sapiens protein kinase N1 (PKN1):

ccctccctccgcgcggggacccctggcgggcggcaggaggacatg (Seq ID No: 1282)

15 Homo sapiens hemochromatosis type 2 (juvenile) (HFE2):

ccttctctgggtccctgacctcagtgagacagcagccggcctggggacctgggggagacacggagg
acccctggctggagctgaccacagagtagggaatcatggctggagaattggatagcagagtaat
gtttgacctctggaaacatcacttacagggtctccgggtcaaaattcactaggtaggaggggtcatca
gctgggaagaaccggcgccctgggaacctggctggataggtatg (Seq ID No: 1283)

20 Homo sapiens ribosomal protein L9 (RPL9):

cgttctttctttgctgcgtctactgcgagaatg (Seq ID No: 1284)

Homo sapiens ribosomal protein L3 (RPL3):

cggcctctaccggcgggatttgatggcgtgatg (Seq ID No: 1285)

Homo sapiens ribosomal protein L4 (RPL4):

25 acttccttttctgtggcagcagccgggctgagaggagcgtggctgtctcctctctccgccatg
(Seq ID No: 1286)

Homo sapiens ribosomal protein L5 (RPL5):

tggcccttttccccccccctagcgccgctgggcctgcaggtctctgtcgcagcagcggacgcgggtc
tctgttccgcaggatg (Seq ID No: 1287)

30 Homo sapiens ribosomal protein L6 (RPL6): aattctctttcccatcttgcaa-
gatg (Seq ID No: 1288)

Homo sapiens ribosomal protein L7 (RPL7): cttcctctttttccggctggaac-
catg (Seq ID No: 1289)

Homo sapiens ribosomal protein L7a (RPL7A):

35 ctttcttttctctctcctcccgccgccaagatg (Seq ID No: 1290)

Homo sapiens ribosomal protein L11 (RPL11): ctttctcttctgctctccat-
catg (Seq ID No: 1291)

Homo sapiens ribosomal protein L12 (RPL12):

40 cggcctctcggctttcggctcggaggaggccaaggtgcaacttccttcggctcgtcccgaatccggg
ttcatccgacaccagccgcctccaccatg (Seq ID No: 1292)

245

Homo sapiens ribosomal protein L13 (RPL13):

gcttcctttccgctcggctgttttctctgcgcaggagccgcagggccgtaggcagccatg
(Seq ID No: 1293)

Homo sapiens ribosomal protein L23 (RPL23):

5 acttccttttttctttttccggcggttcaagatg (Seq ID No: 1294)

Homo sapiens ribosomal protein L18 (RPL18):

cgttctctctttccggacctggccgagcaggaggcgccatcatg (Seq ID No: 1295)

Homo sapiens ribosomal protein L18a (RPL18A):

acttccttttgcggtggcggcgaacgcggagagcacgccatg (Seq ID No: 1296)

10 Homo sapiens ribosomal protein L19 (RPL19):

agctctttcctttcgctgctgcggccgcagccatg (Seq ID No: 1297)

Homo sapiens ribosomal protein L21 (RPL21):

gcctctttcctttcgggccgaaccgccatcttccagtaattcgccaaaatg

(Seq ID No: 1298) Homo sapiens ribosomal protein L22 (RPL22):

15 acctcccttttctaactccgctgccgcatg (Seq ID No: 1299)

Homo sapiens ribosomal protein L23a (RPL23A): agacccttttcacaagatg
(Seq ID No: 1300)

Homo sapiens ribosomal protein L17 (RPL17):

20 cgctcttcctctttccctaagcagcctgagggttgactggattggtgaggcccggtgtggctacttc
tgtggaagcagtgctgtagttactggaagataaaagggaaagcaagcccttggtgggggaaagtat
ggctgcgatgatggcatttcttaggacacctttggattaataatgaaaacaactactctctgagca
gctgttcgaatcatctgatatttatactgaatgagttactgtaagtacgtattgacagaattacac
tgtactttcctctaggtgatctgtgaaaatg (Seq ID No: 1301)

Homo sapiens ribosomal protein L24 (RPL24):

25 ttctctctttccttttcgccatcttttgtctttccgtggagctgtcgccatg
(Seq ID No: 1302)

Homo sapiens ribosomal protein L26 (RPL26):

agttctcttcccttttgcgccatcacccaagcgggagcggccaaaatg
(Seq ID No: 1303)

30 Homo sapiens ribosomal protein L27 (RPL27):

ctttcctttttgctggtagggccgggtggttgctgccgaaatg (Seq ID No: 1304)

Homo sapiens ribosomal protein L30 (RPL30):

aagtctttccttttctcgttccccggccatcttagcggtgctgttggttgggggcccgtcccgtcc
taaggcaggaagatg (Seq ID No: 1305)

35 Homo sapiens ribosomal protein L27a (RPL27A):

ccttcctttttcgtctgggctgccaacatg (Seq ID No: 1306)

Homo sapiens ribosomal protein L28 (RPL28):

cttcctctttccgtctcaggtcgccgctgcgaaggagccgccgcatg
(Seq ID No: 1307)

Homo sapiens ribosomal protein L29 (RPL29):

cagccccctttctcttccggttctaggcgcttcgggagccgcggttatggtgcagacatg

(Seq ID No: 1308)

Homo sapiens ribosomal protein L31 (RPL31):

5 cgctcttcctttccaacttggacgctgcagaatg (Seq ID No: 1309)

Homo sapiens ribosomal protein L32 (RPL32):

ccgtcccttctctcttccctcggcgctgcctacggaggtggcagccatctccttctcggcatcatg

(Seq ID No: 1310)

Homo sapiens ribosomal protein L35a (RPL35A):

10 cgtccttctcttaccgccatcttggctcctgtggaggcctgctgggaacgggacttctaaaaggaa
ctatg (Seq ID No: 1311)

Homo sapiens ribosomal protein L37 (RPL37):

ccttctcttccggtctttctggtctcggccgcagaagcgagatg (Seq ID No: 1312)

Homo sapiens ribosomal protein L37a (RPL37A):

15 gcgtctcttcctttctggtcgcgacctaggtcgcggcgacatg (Seq ID No: 1313)

Homo sapiens ribosomal protein L38 (RPL38):

cgttctttttcgtccttttccccgggttgctgcttgctgtgagtgctcttagggtgatacgtgggtg

agaaaggtcctggtccgcgccagagcccagcgcgcctcgtcgccatg (Seq ID No: 1314)

Homo sapiens ribosomal protein L39 (RPL39):

20 ccctcctcttcttcttccgccatcgtggtgtgttcttgactccgctgctcgccatg

(Seq ID No: 1315)

Homo sapiens ribosomal protein, large, P0 (RPLP0):

aggcccttctctcgccaggcgtcctcgtggaagtgcacatcgtctttaaacctgcgtggcaatccc

tgacgcaccgcgctgatg (Seq ID No: 1316)

25 Homo sapiens ribosomal protein, large, P1 (RPLP1):

cggtccttccgaggaagctaaggctgcgttgggtgaggccctcacttcattccggcgactagcacc

gcgtccggcagcgcgccagccctacactcgcccgcgccatg (Seq ID No: 1317)

Homo sapiens ribosomal protein, large, P2 (RPLP2):

30 ccttccttttctcctgtcgccaccgaggtcgcacgcgtgagacttctccgccgcctccgccgca

gacgccgcgcgcatg (Seq ID No: 1318)

Homo sapiens ribosomal protein S3 (RPS3):

acttccttttcttccagcggagcgcggcggaagatg (Seq ID No: 1319)

Homo sapiens ribosomal protein S3A (RPS3A):

ccgcccttttggctctctgaccagcaccatg (Seq ID No: 1320)

35 Homo sapiens ribosomal protein S4, X-linked (RPS4X):

ggctcctctttccttgcttaacgcagccatg (Seq ID No: 1321)

Homo sapiens ribosomal protein S4, Y-linked 1 (RPS4Y1):

gattctcttccgtcgcagagtttcgccatg (Seq ID No: 1322)

Homo sapiens ribosomal protein S5 (RPS5):

40 ttttcttccagttaaaagtgttgcccgcggcgcgcgcgcctcttctgtctgtaccagggcggcg

cgtggtctacgccgagtgacagagacgctcaggctgtgttctcaggatg
(Seq ID No: 1323)

Homo sapiens ribosomal protein S6 (RPS6):
ggccctcttttccgtggcgccctcgaggcggttcagctgcttcaagatg (Seq ID No: 1324)

5 Homo sapiens ribosomal protein S7 (RPS7):
gggtctcttcctaagccggcgctcggcaagttctcccaggagaaagccatg
(Seq ID No: 1325)

Homo sapiens ribosomal protein S8 (RPS8):
gtttctctttccagccagcgccgagcgatg (Seq ID No: 1326)

10 Homo sapiens ribosomal protein S9 (RPS9):
gcgctcttttctcagtgaccgggtggttgcttaggcgcagacggggaagcggagccaacatg
(Seq ID No: 1327)

Homo sapiens ribosomal protein S10 (RPS10):
gctccttctttccagccccggtaccggaccctgcagccgcagagatg (Seq ID No: 1328)

15 Homo sapiens ribosomal protein S11 (RPS11):
ctgccccctttcttttttttcaggcgccgggaagatg (Seq ID No: 1329)

Homo sapiens ribosomal protein S12 (RPS12):
aggcctctttccctgcccgcgcgagtcgcgcggaggcggaggcttgggtgcgttcaagattcaac
ttcacccgtaaccacccgcatg (Seq ID No: 1330)

20 Homo sapiens ribosomal protein S13 (RPS13):
cgctctcctttcggtgcctgatcgccgccatcatg (Seq ID No: 1331)

Homo sapiens ribosomal protein S15 (RPS15):
cgatctcttctgaggatccggcaagatg (Seq ID No: 1332)

Homo sapiens ribosomal protein S15a (RPS15A):
25 cgtcctctttccgccatctttccgcgcgggtgagtagcactctctgagagctccaatttcatccgt
ctgccatcggcgccatcctgcaatctaagccacaatg (Seq ID No: 1333)

Homo sapiens ribosomal protein S16 (RPS16):
ctttccttttccggttgccgcgcgcggtgaggttgcttagtccacgctcggagccatg
(Seq ID No: 1334)

30 Homo sapiens ribosomal protein S19 (RPS19):
cgttccctttcccctggctggcagcgcgaggccgcacgatg (Seq ID No: 1335)

Homo sapiens ribosomal protein S20 (RPS20):
ccacccttttctttttgaggaagacgcggtcgtaagggctgaggatttttggtccgcacgctcctg
ctcctgactcacgcgtgttcgctctcgccgaggaacaagtcggtcaggaagcccgcgcgaacagc
35 catg (Seq ID No: 1336)

Homo sapiens ribosomal protein S21 (RPS21):
gcttcctttctctctcgcgcgcggtgtggtggcagcaggcgcagcccagcctcgaaatg
(Seq ID No: 1337)

Homo sapiens ribosomal protein S23 (RPS23):
40 gcttctctctttcgctcaggcccggtggcgccgacaggatg (Seq ID No: 1338)

Homo sapiens ribosomal protein S25 (RPS25):
gcttcctttttgtccgacatcttgacgaggctgcggtgtctgctgctattctccgagcttcgcaat
g (Seq ID No: 1339)

5 Homo sapiens ribosomal protein S26 (RPS26):
ccgtctcctctctccggtccgtgcctccaagatg (Seq ID No: 1340)

Homo sapiens ribosomal protein S27 (RPS27):
cgctcctttccggcggtgacgacctacgcacacgagaacatg (Seq ID No: 1341)

Homo sapiens ribosomal protein S28 (RPS28):
actcctctccgccagaccgccgcgcgcccatcatg (Seq ID No: 1342)

10 Homo sapiens ribosomal protein S29 (RPS29):
gcttcttccttttacctcgttgactgctgagagcaagatg (Seq ID No: 1343)

Homo sapiens ribosomal protein L15 (RPL15):
agctctttcctttccgtctggcggcagccatcaggttaagccaagatg (Seq ID No: 1344)

15 Homo sapiens ribosomal protein S2 (RPS2):
cgttcttcttttccgacaaaacaccaaagatg (Seq ID No: 1345)

Homo sapiens ribosomal protein L14 (RPL14):
gggtcttcttctctcgcctaacgcgcgaacatg (Seq ID No: 1346)

Homo sapiens ribosomal protein S14 (RPS14):
ctctctttccggtgtggagtctggagacgacgtgcagaaatg (Seq ID No: 1347)

20 Homo sapiens ribosomal protein L10 (RPL10):
gcgcctctttcccttcggtgtgccactgaagatcctggtgtcgccatg (Seq ID No: 1348)

Homo sapiens ribosomal protein L10a (RPL10A):
tagtctcttttccggttagcgcgcggtgagaagccatg (Seq ID No: 1349)

25 Homo sapiens ribosomal protein L35 (RPL35):
tcctctttccctcggagcgggcggcggttgcggttggtgcagcaatg
(Seq ID No: 1350)

Homo sapiens ribosomal protein L13a (RPL13A):
cctcctcctttccaagcggctgccgaagatg (Seq ID No: 1351)

30 Homo sapiens ribosomal protein L36 (RPL36):
cagcccttccgccacggccgtctctggagagcagcagccatg (Seq ID No: 1352)

Homo sapiens ribosomal protein L36a (RPL36A):
gtttctttctttccgcgccgatagcgctcacgcaagcatg (Seq ID No: 1353)

35 Homo sapiens ribosomal protein L41 (RPL41):
tcgcc tttctctcggccttagcgccatttttttgaaacctctgcgccatg
(Seq ID No: 1354)

Homo sapiens ribosomal protein S18 (RPS18):
cgctctcttctccacaggaggcctacacgccgcgcttggtgctgcagccatg
(Seq ID No: 1355)

Homo sapiens ribosomal protein S24 (RPS24):
ggttctcttttctccttggctgtctgaagatagatcgccatcatg (Seq ID No: 1356)

Homo sapiens ribosomal protein L8 (RPL8):
tttctcttttcggccgcgctggtgaacaggtaggtcatccttgcggccttgcggcgcatg
5 (Seq ID No: 1357)

Homo sapiens ribosomal protein L34 (RPL34):
cttcctcttccggggacggtgtctgcaggtatg (Seq ID No: 1358)

Homo sapiens ribosomal protein S17 (RPS17):
gtttcctcttttaccaggaccgccaacatg (Seq ID No: 1359)

10 Homo sapiens ribosomal protein SA (RPSA):
ctgtcttttccgtgctacctgcagaggggtccatacggcggttgttctggattcccgtcgtaactta
aagggaattttcacatg (Seq ID No: 1360)

Homo sapiens eukaryotic translation initiation factor 3, subunit C (EIF3C):
15 cttctctctcggcgtttccgctgtcagggccctgcgggtgtgactcgcgggctcagctggtccggcc
gtagcacctccgcgcgctcgccatg (Seq ID No: 1361)

Homo sapiens poly(A) binding protein, cytoplasmic 1 (PABPC1):
cgctctctcctctcacggaaaggtcgcggcctgtggccctgcgggcagccgtgccgagatg
(Seq ID No: 1362)

20 Homo sapiens tubulin, beta 1 class VI (TUBB1):
cactcccttccaaaagcatgacaggcagaaagcagagaagggccaggactggctgagggcggggag
ctgggcctctggggtggacacacccttggtcacattgtgagggtagcttggttggccagtcccacc
actgcagtgaccacagttgtgttgggtcacaccagtgaaccgaagctctggattctgagagtctg
aggattccgtgaagatctcagacttgggctcagagcaaggatg (Seq ID No: 1363)

25 PpLuc(GC) - A64N64
GGGAGAAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGCGCCCTTCTA
CCCGCTGGAGGACGGGACCGCCGCGGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCT
GGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTGACATCACCTACGCGGA
GTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAA
30 CCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGTGGGCGC
CCTCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCT
GAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAA
GATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAA
GACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGG
35 CTTC AACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGAT
CATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGC
CTGCGTGCGCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACAC
CGCCATCCTGAGCGTGGTGGCCTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTA
CCTCATCTGCGGCTTCCGGGTGGTCTGTATGTACCGGTTCGAGGAGGAGCTGTTCTCTGCG
40 GAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCTGTTACGCTTCTT
CGCCAAGAGCACCCCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGG
GGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGG
CATCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGG
GGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGA
45 CCTGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGCGAGCTGTGCGTGCGGGGGCC
GATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGA

CGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGT
CGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGA
GAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGA
CGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAGACCATGACGGA
5 GAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGG
CGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGAT
CCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGAT
CTAA
AAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGC
10 CACCAGAATT (SEQ ID No: 1364)

PpLuc(GC) - albumin7 - A64N64

GGGAGAAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTA
15 CCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCT
GGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTGACATCACCTACGCGGA
GTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAA
CCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGTGTTGGGCGC
CCTCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCT
20 GAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAA
GATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAA
GACCGACTACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGG
CTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGAT
CATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGC
25 CTGCGTGCCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACAC
CGCCATCCTGAGCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTA
CCTCATCTGCGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTCTGCG
GAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTT
CGCCAAGAGCACCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGG
30 GGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGG
CATCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGGATCCTGATCACCCCGAGGG
GGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGA
CCTGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCC
GATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGA
35 CGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGT
CGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGA
GAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGA
CGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAGACCATGACGGA
GAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGG
40 CGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGAT
CCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCAT
CACATTTAAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAG
CTTATTCATCTCTTTTCTTTTCTGTTGGTGTAAAGCCAACACCCTGTCTAAAAAACATA
AATTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAA
45 CCTAGATCTAA
AAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTT
TCAGAGCCACCAGAATT (SEQ ID No: 1365)

50 RPL32RPL32 - PpLuc(GC) - A64N64

GGGGCGCTGCCTACGGAGGTGGCAGCCATCTCCTTCTCGGCATCAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCGCTGGAGGACGGGACCGCC

GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCGTC
5 GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTC
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
10 GGCTTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCG
TTCCACCACGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCCTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTACGTTCTTCGCCAAGAGCACCCCTGATCGAC
15 AAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
20 AACAAACCCGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
25 GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAA
AATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID No: 1366)

30

RPL32 - PpLuc(GC) - albumin7 - A64N64
GGGGCGCTGCCTACGGAGGTGGCAGCCATCTCCTTCTCGGCATCAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
35 GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCGTC
GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
40 CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTC
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
GGCTTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
45 CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCG
TTCCACCACGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCCTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTACGTTCTTCGCCAAGAGCACCCCTGATCGAC
AAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
50 GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTG

GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
AACAAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
5 AACATCTTCGACGCCGGCGTGCCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCC
10 TACCATGAGAATAAGAGAAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTT
TCGTTGGTGTAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCT
CTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAA
AAAAAATGCATCCCC
CCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID
15 No: 1367)

5'UTR of human ribosomal protein Large 32 (RPL32) lacking the 5'
terminal oligopyrimidine tract
GGCGCTGCCTACGGAGGTGGCAGCCATCTCCTTCTCGGCATC (SEQ ID No. 1368)
20 Human albumin 3'UTR
CATCACATTT AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA
AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAC
ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA AAAATGGAAA
GAATCT (SEQ ID No: 1369)

25 3'UTR of Homo sapiens hemoglobin, alpha 1 (HBA1)
gctggagcctcggtggccatgcttcttggcccttgggcctccccccagcccctcctccccttcctg
caccgctacccccgtggtctttgaataaagtctgagtgggcggc (SEQ ID No: 1370)

30 3'UTR of Homo sapiens hemoglobin, alpha 2 (HBA2)
gctggagcctcggtagccgttctcctgcccgtgggcctcccaacgggcctcctcccctccttg
caccggcccttctggtctttgaataaagtctgagtgggcag (SEQ ID No: 1371)

3'UTR of Homo sapiens hemoglobin, beta (HBB)
35 Gctcgctttcttctgtgtccaatttctattaaaggttcccttgttccctaagtccaactactaaact
gggggatattatgaaggccttgagcatctggattctgcctaataaaaaacattttattttcattgc
(SEQ ID No: 1372)

3'UTR of Homo sapiens tyrosine hydroxylase (TH)
40 gtgcacggcgtccctgagggcccttcccaacctcccctggtcctgcactgtcccggagctcagggc
ctggtgaggggctgggtcccgggtgcccccatgcctcctgctgccaggctcccactgcccctg
cacctgcttctcagcgcaacagctgtgtgtgcccgtggtgaggttgtgctgctgtggtgaggtcc
tgtcctggctcccagggctcctgggggctgctgcactgcctccgccttccctgacactgtctgct
gcccacatcaccgtcacaataaaagaaactgtggtctcta (SEQ ID No: 1373)

45 3'UTR of Homo sapiens arachidonate 15-lipoxygenase (ALOX15)
gcgtcgccaccctttgggtatttcagcccccatcaccacagccacaagctgacccttctgtggtta
tagccctgcccctcccaagtcccaccctcttcccatgtcccaccctccctagaggggcaccttttca
tggtctctgcaccagtgaaacacattttactctagaggcatcacctgggaccttactcctctttcc

ttccttcctcctttcctatcttccttcctctctctcttcctctttcttcattcagatctatatggc
aaatagccacaattatataaatcattttcaagactagaataggggatataatacatattactccac
acctttttatgaatcaaataatgatttttttgttgttgttaagacagagtctcactttgacacccagg
ctggagtgacagtgggtgccatcaccacgggtcactgcagcctcagcgtcctggggtcaaataatgatcct
5 cccacctcagcctcctgagtagctgggactacagggtcatgccatcatgccagctaataatttttt
tattttcgtggagacggggcctcactatgttgccctagggtggaaataggatttttgaacccaaattg
agtttaacaataataaaaaagttgttttacgctaagatggaaaagaactaggactgaactatttta
aataaaatattggc (SEQ ID No: 1374)

10 3'UTR of Homo sapiens collagen, type I, alpha 1 (COL1A1)
actccctccatcccaacctggctccctccacccaaccaactttccccccaacccggaaacagaca
agcaacccaaactgaacccccctcaaaagccaaaaaatgggagacaatttcacatggacttttgaaa
atatttttttcccttgcaattcatctctcaaaacttagtttttatctttgaccaaccgaacatgacca
aaaaccaaagtgcattcaaccttaccaaaaaaaaaaaaaaaaaaaaaagaataaataaacttttt
15 aaaaaaggaagcttgggtccacttgcttgaagacccatgcgggggtaagtccctttctgcccgttgg
gcttatgaaacccaatgctgccctttctgctcctttctccacaccccccttggggcctccctcc
actccttcccaaactctgtctcccagaagacacaggaaacaatgtattgtctgccagcaatcaaa
ggcaatgctcaaacacccaagtggccccccacccctcagcccgtcctgcccggccagcaccacccagg
cctgggggacctgggggttctcagactgccaaagaagccttgccatctggcgtcccatgggtcct
20 gcaacatctcccttctgtttttgaggggtcatgcgggggagccaccagccctcactgggttcg
gaggagagtcaggaagggccacgacaaaagcagaacatcgatttggggaacgcgtgtcaatccct
tgtgccgcaggggtggggcgggagagactgttctgttccttgtgtaactgtgttgctgaaagactac
ctcgttcttctgtcttgatgtgtcaccggggcaactgcctgggggcggggatgggggcaggggtggaag
cggctccccattttataccaaaggtgctacatctatgtgatgggtgggggtggggaggggaatcactg
25 gtgctatagaaattgagatgccccccagggccagcaaatgttcctttttgttcaaagtctattttt
attccttgataatttttctttttttttttttttttttttgtggatggggacttgtgaatttttctaaag
gtgctatttaacatgggaggagagcgtgtgcgggtccagcccagcccgtgctcactttccaccct
ctctccacctgcctctggttctcaggcctctgctctccgacctctctcctctgaaacctcctcc
acagctgcagcccatecctcccggtccctcctagctctgtcctgcgtcctctgtccccgggtttcag
30 agacaacttcccaaagcacaagcagtttttccccctaggggtgggaggaagcaaaagactctgta
cctattttgtatgtgtataataatttgagatgtttttaattattttgattgctggaataaagcatg
tggaatgacccaacataa (SEQ ID No: 1375)

albumin7 3'UTR

35 CATCACATTTAAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTT
ATTCATCTCTTTTCTTTTTCGTTGGTGTAAGCCAACACCCTGTCTAAAAACATAAATTTCTTT
AATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCT (SEQ ID No:
1376)

40 Human albumin 3'UTR + poly(A) sequence

CATCACATTT AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA
AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA AAAATGGAAA
GAATCTAGAT CTAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA
45 AAAAAAAAAA AAAAAA (SEQ ID No: 1377)

254

Human albumin 3'UTR fragment 1

AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC
ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT
TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATT (SEQ ID No: 1378)

5

Human albumin 3'UTR fragment 2

CATCACATTT AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA
AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG (SEQ ID No:
1379)

10

Human albumin 3'UTR fragment 3

AAAAGCATCT CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC
ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC (SEQ ID No:
1380)

15

Human albumin 3'UTR fragment 4

CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT
CTTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT (SEQ ID No:
1381)

20

Human albumin 3'UTR fragment 5

TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT
GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT (SEQ ID No:
1382)

25

Human albumin 3'UTR fragment 6

AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC
CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT (SEQ ID No:
1383)

30

Human albumin 3'UTR fragment 7

TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG
TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT (SEQ ID No:
1384)

35

Human albumin 3'UTR fragment 8

AAGCTTATTC ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA (SEQ ID No:
1385)

40

Human albumin 3'UTR fragment 9

ATCTGTTTTT CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT
TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA AAAATGGAAA (SEQ ID No:
1386)

45

Human albumin 3'UTR fragment 10

CAGCCTACCA TGAGAATAAG AGAAAGAAAA TGAAGATCAA AAGCTTATTC ATCTGTTTTT
CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT
TGCCTCTTTT CTCTGTGCTT CAATTAATAA A (SEQ ID No: 1387)

5

Human albumin 3'UTR fragment 11

TGAAGATCAA AAGCTTATTC ATCTGTTTTT CTTTTCGTT GGTGTAAAGC CAACACCCTG
TCTAAAAAAC ATAAATTTCT TTAATCATTT TGCCTCTTTT CTCTGTGCTT CAATTAATAA
A (SEQ ID No: 1388)

10

Human albumin 3'UTR fragment 12

CTTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC ATAAATTTCT TTAATCATTT
TGCCTCTTTT CTCTGTGCTT CAATTAATAA A (SEQ ID No: 1389)

15

Human albumin 3'UTR fragment 13

AAGCTTATTC ATCTGTTTTT CTTTTCGTT GGTGTAAAGC CAACACCCTG TCTAAAAAAC
(SEQ ID No: 1390)

Albumin7 3'UTR - poly(A) sequence - poly(C) sequence - HL

20 CATCACATTTAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTT
ATTCATCTCTTTTTCTTTTTCGTTGGTGTAAAGCCAACACCCTGTCTAAAAACATAAATTTCTTT
AATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAA
AAATGCATCCC
CCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID
25 No: 1391)

Albumin7 3'UTR - poly(A) sequence - poly(C) sequence

30 CATCACATTTAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTT
ATTCATCTCTTTTTCTTTTTCGTTGGTGTAAAGCCAACACCCTGTCTAAAAACATAAATTTCTTT
AATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAA
AAATGCATCCC
CCCCCCCCCCCCCCCCCCCCCCCCCCCC (SEQ ID No: 1392)

35 Center, α -complex-binding portion of the 3'UTR of an α -globin gene
GCCCGATGGGCCTCCCAACGGGCCCTCCTCCCCTCCTTGACCG (SEQ ID NO: 1393)

Histone stem-loop

CAAAGGCTCTTTTCAGAGCCACCA (SEQ ID NO: 1394)

40

ATP synthase lipid-binding protein, mitochondrial (atp5g2)

tagttt ctcctctcga acgccaggtg gagcaaccgg ccggataccg ccacagccct
ggcaggcggc gctgtgatg (SEQ ID NO: 1395)

45

RPL35 - PpLuc(GC) - albumin7 - A64N64

GGGGAGCGGGCGGCGGCGTGTGCGGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCGGCGAGCA
GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA

GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGTGGGCGCCCTCTTCATCGGCGTGGCCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
GGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
5 CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
10 CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
15 CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
20 GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
25 CGGCAAGATCGCCGTGTAAGACTAGTGATCACATTTAAAAGCATCTCAGCCTACCATGA
GAATAAGAGAAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTTTCTTTTGGT
GTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
TGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAAAAAA
AAATGCATCCCCCCCCCCCC
30 CCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1396)

RPL21 - PpLuc(GC) - albumin7 - A64N64
GGGGCCGGAACCGCCATCTTCCAGTAATTCGCCAAAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
35 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGTGGGCGCCCTCTTCATCGGCGTGGCCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
40 GGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
45 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
50 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA

CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
5 CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGA
10 GAATAAGAGAAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTTTCTGTTGGT
GTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
TGTGCTTCAATTAATAAAAAATGGAAGAACCTAGATCTAAAAAAAAAAAAAAAAAAAAA
AAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1397)
15

ATP5A1 - PpLuc(GC) - albumin7 - A64N64
GGGCGGCTCGGCCATTTTGTCCAGTCAGTCCGGAGGCTGCGGCTGCAGAAGTACCGCCT
GCGGAGTAACTGCAAAGAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCG
GCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAG
20 CGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTGCACATC
ACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGC
CTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCG
GTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAG
CGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAG
25 GGCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATC
ATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCGTGACCAGCCAC
CTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACC
ATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCG
CACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATC
30 ATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCACCACGGCTTCGGCATGTTACG
ACCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAG
CTGTTCTTCCGAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCCTG
TTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGACCTGTGAACCTGCACGAG
ATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTTC
35 CACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGGATCCTGATC
ACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCC
AAGGTGGTGGACCTGGACACCGGCAAGACCTGGGCGTGAACCAGCGGGGCGAGCTGTGC
GTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTC
ATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCAC
40 TTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCG
GCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGG
CTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGCTGGAGCACGGCAAG
ACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAG
CTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGAC
45 GCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAA
GACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATG
AAGATCAATAGCTTATTCATCTCTTTTTCTTTTTCTGTTGGTGTAAAGCCAACACCCTGTC
TAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAA
AATGGAAGAAGACCTAGATCTAA
50 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCC
AAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1398)

HSD17B4 - PpLuc(GC) - albumin7 - A64N64

GGGTCCCGCAGTCGGCGTCCAGCGGCTCTGCTTGTTTCGTGTGTGTGTCGTTGCAGGCCTT
ATTCAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGCGCCCTTCTACC
5 CGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGG
TGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGT
ACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACC
ACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCC
TCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGA
10 ACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGA
TCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATATGGACAGCAAGA
CCGACTACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCT
TCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCA
TGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCT
15 GCGTGCGCTTCTCGCACGCCCCGGGACCCCCATCTTCGGCAACCAGATCATCCCGGACACCG
CCATCCTGAGCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACC
TCATCTGCGGCTTCCGGGTGGTCCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTCTGCGGA
GCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTTCG
CCAAGAGCACCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGG
20 GCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCA
TCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGGATCCTGATCACCCCGAGGGGG
ACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACC
TGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGA
TGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACG
25 GCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCG
ACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGA
GCATCCTGCTCCAGCACCCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACG
ACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGGAGCACGGCAAGACCATGACGGAGA
AGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCG
30 TGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCC
GCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCA
CATTTAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAAGAAAATGAAGATCAATAGCT
TATTCATCTCTTTTCTTTTCTGTTGGTGTAAAGCCAACACCCTGTCTAAAAAACATAAA
TTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACC
35 TAGATCTAAAAAAGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTC
AGAGCCACCAGAATT (SEQ ID NO: 1399)

AIG1 - PpLuc(GC) - albumin7 - A64N64

GGGCCGCCCAGCCGGTCCAGGCCTCTGGCGAACAAAGCTTGAGGATGGAGGACGCCAAGAA
40 CATCAAGAAGGGCCCGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCT
CCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCCA
CATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGC
CATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCT
45 GCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGCGAA
CGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGT
GTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCAT
CCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACAC
GTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTT
50 CGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAA

GGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCCCCAT
CTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCACGG
CTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCCTGATGTA
CCGGTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCT
5 GCTCGTGCCGACCCTGTTGAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGACCT
GTCGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGC
CGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCAC
GAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGT
CCCGTTCTTCGAGGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAACCA
10 GCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGA
GGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTG
GGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGG
CTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCCAACATCTTCGA
CGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGT
15 GCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGT
GACCACCGCCAAGAAGCTGCCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCT
GACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGG
CAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAGCATCTCAGCCTACCATGAGAA
TAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTTTCGTTGGTGTA
20 AAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTCTGT
GCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAA
AAATGCATCCCCCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1400)

25 COX6C - PpLuc(GC) - albumin7 - A64N64
GGAGTCAGGAAGGACGTTGGTGTTGAGGTTAGCATACGTATCAAGGACAGTAACTACCAA
GCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCCGGCGCCCTTCTACCCGCTGG
AGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGG
GCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCG
30 AGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGA
TCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCA
TCGGCGTGGCCGTGCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCA
TGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGA
ACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACT
35 ACCAGGGCTTCCAGTCGATGTACAGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACG
AGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACA
GCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGC
GCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCC
TGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTTACGACCCTGGGCTACCTCATCT
40 GCGGCTTCCGGGTGGTCTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTTGCAGGAGCCTGC
AGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGA
GCACCCTGATCGACAAGTACGACCTGTGGAACCTGCACGAGATCGCCAGCGGGGGCGCCC
CGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCC
AGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACA
45 AGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACA
CCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCA
TGAGCGGCTACGTGAACAACCCGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGC
TGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGC
TGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCC
50 TGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACCGCG
GCGAGCTGCCGGCCGCGGTGGTGGTGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGA

TCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCCGGGCGGGCGTGGTGT
TCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGA
TCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTA
AAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCA
5 TCTCTTTTTCTTTTTCTGTTGGTGTAAAGCCAACACCCTGTCTAAAAACATAAATTTCTT
TAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATC
TAAA
AAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCC
ACCAGAATT (SEQ ID NO: 1401)

ASA1 - PpLuc(GC) - albumin7 - A64N64

GGGCCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCTAGAGCGAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
15 ACCGACGCCCACATCGAGGTGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGTGTTGGCGCCCTCTTCATCGGCGTGGCCGTC
GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
20 CTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTC
CCGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
GGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCCTTCTCGCACGCC
CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGGCG
25 TTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCCTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTTCGCCAAGAGCACCCTGATCGAC
AAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
30 ACCGAGACCACGAGCGGATCCTGATCACCCCCGAGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
35 AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACCGCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTGAGGACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
40 AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAGCATCTCAGCC
TACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTT
TCGTTGGTGTAAGCCAACACCCTGTCTAAAAACATAAATTTCTTTAATCATTTTGCCT
CTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAAAAAAAAA
AAATGCATCCCC
45 CCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID
NO: 1402)

mRPL21 - PpLuc(GC) - albumin7 - A64N64

GGGGCCCGCCGAGCCATCTTCCAGTAACTCGCCAAAAAGCTTGAGGATGGAGGACGCCAA
50 GAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCGGCGAGCA

GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGC
5 GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
GGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
10 GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCC
CATCTTCGGCAACCAGATCATCCCGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGA
15 CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
20 GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGTGAGGACGCGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
25 GGTGACCACCGCCAAGAAGCTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGA
GAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTTCATCTCTTTTTCTTTTTCTGTTGGT
GTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
30 TGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1403)

mRPL35A - PpLuc(GC) - albumin7 - A64N64

35 GGGCCATCTTGGCGCCTGTGGAGGCCTGCTGGGAACAGGACTTCTAACAGCAAGTAAGCT
TGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGG
ACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCA
CGATCGCCTTCACCGACGCCACATCGAGGTGACATCACCTACGCGGAGTACTTCGAGA
TGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCG
40 TGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGTGGGCGCCCTCTTCATCG
GCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGG
GGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACG
TGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACC
AGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGT
45 ACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCA
GCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCT
TCTCGCACGCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGA
GCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCG
GCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAGCTGTTTCTGCGGAGCCTGCAGG
50 ACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCA
CCCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGC

TGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGTTCCACCTCCCGGGCATCCGCCAGG
GCTACGGCCTGACCGAGACCACGAGCGGATCCTGATCACCCCGAGGGGACGACAAGC
CGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCG
GCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGA
5 GCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGC
ACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGA
AGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGC
TCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCG
AGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCG
10 TCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCG
TGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCC
TGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAA
GCATCTCAGCCTACCATGAGAATAAGAGAAAAGAAAATGAAGATCAATAGCTTATTCATCT
CTTTTTCTTTTTCTGTTGGTGTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAA
15 TCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAA
AAA
AATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACC
AGAATT (SEQ ID NO: 1404)

20 RPL35 - PpLuc (GC) - A64N64
GGGGAGCGGGCGGCGGCGTGTGGCGGCTTGTGCAGCAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGGCGGGCACGATCGCCTTACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
25 GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
GGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
30 CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCCTGCC
GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCCCTGAT
35 GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
40 GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
45 CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGTGAGGACGCGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAA
50 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCC
CCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1405)

RPL21 - PpLuc (GC) - A64N64

5 GGGGCCGGAACCGCCATCTTCCAGTAATTCGCCAAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGC
10 CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGCTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
15 GGTGTTCTGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTCTGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
20 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
25 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
30 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGCCGGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
35 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCC
CCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1406)

35 ATP5A1 - PpLuc (GC) - A64N64

GGGCGGCTCGGCCATTTTGTCCAGTCAGTCCGGAGGCTGCGGCTGCAGAAGTACCGCCT
GCGGAGTAAGTCAAAGAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCG
GCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAG
CGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATC
40 ACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGC
CTGAACACCAACCACCGGATCGTGCTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCG
GTGCTGGGCGCCCTCTTCATCGGCGTGCCGTCGCCCCGGCGAACGACATCTACAACGAG
CGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAG
GGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATC
45 ATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCAC
CTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACC
ATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCG
CACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATC
ATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTACG
50 ACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAG

20 GGGTCCCGCAGTCCGGCGTCCAGCGGCTCTGCTTGTTCGTGTGTGTGTCGTTGCAGGCCCTT
ATTCAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCCGGCGCCCTTCTACG
CGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGG
TGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGT
ACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACC
25 ACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGCTGCTGGGCGCC
TCTTCATCGGCGTGGCCGTCGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGA
ACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGA
TCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATATGGACAGCAAGA
CCGACTACCAGGGCTTCCAGTCGATGTACAGTTTCGTGACCAGCCACCTCCCGCCGGGCT
30 TCAACGAGTACGACTTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCA
TGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCT
GCGTGCGCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCG
CCATCCTGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACC
TCATCTGCGGCTTCCGGGTGGTCCTGATGTACCGGTTTCGAGGAGGAGCTGTTTCTGCGGA
35 GCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCG
CCAAGAGCACCTGATCGACAAGTACGACCTGTTCGAACCTGCACGAGATCGCCAGCGGGG
GCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCA
TCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGG
ACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACC
40 TGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGA
TGATCATGAGCGGCTACGTGAACAACCCGAGGCCACCAACGCCCTCATCGACAAGGACG
GCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCG
ACCGGCTGAAGTCGTGATCAAGTACAAGGGCTACCAGGTGGCGCCGCGGAGCTGGAGA
GCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACG
45 ACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGAGACACGGCAAGACCATGACGGAGA
AGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCG
TGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCC
GCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCT
AAA
50 AAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCA
CCAGAATT (SEQ ID NO: 1408)

AIG1 - PpLuc (GC) - A64N64

GGGCCGCCCAGCCGGTCCAGGCCTCTGGCGAACAAGCTTGAGGATGGAGGACGCCAAGAA
CATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCT
5 CCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCCA
CATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGC
CATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCT
GCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCCGTCGCCCCGGCGAA
CGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGT
10 GTTCGTGAGCAAGAAGGGCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCAT
CCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACAC
GTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTT
CGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAA
GGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCAT
15 CTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCACGG
CTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCCTGATGTA
CCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCT
GCTCGTGCCGACCCTGTTACGTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGACCT
GTCGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGC
20 CGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCAC
GAGCGGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGT
CCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAACCA
GCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGA
GGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTG
25 GGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGG
CTACCAGGTGGCGCCGGCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGA
CGCCGGCGTGCGCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGT
GCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGT
GACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCT
30 GACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGG
CAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCC
CCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1409)

35 COX6C - PpLuc (GC) - A64N64

GGAGTCAGGAAGGACGTTGGTGTTGAGGTTAGCATACGTATCAAGGACAGTAACTACCAA
GCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGG
AGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGG
GCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCG
40 AGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGA
TCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCA
TCGGCGTGCCGTCGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCA
TGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGA
ACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACT
45 ACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACG
AGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACA
GCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGC
GCTTCTCGCACGCCCCGGGACCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCC
TGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTTACGACCCTGGGCTACCTCATCT
50 GCGGCTTCCGGGTGGTCCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGC

AGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGA
GCACCCTGATCGACAAGTACGACCTGTGGAACCTGCACGAGATCGCCAGCGGGGGCGCCC
CGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCACCTCCCGGGCATCCGCC
AGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACA
5 AGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACA
CCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCA
TGAGCGGCTACGTGAACAACCCGAGGGCCACCAACGCCCTCATCGACAAGGACGGCTGGC
TGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGC
TGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCC
10 TGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCG
GCGAGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAGACCATGACGGAGAAGGAGA
TCGTGCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGT
TCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGA
TCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAA
15 AATG
CATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAA
TT (SEQ ID NO: 1410)

ASAH1 - PpLuc (GC) - A64N64

20 GGGCCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCTAGAGCGAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
25 GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTC
GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTC
30 CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
GGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCG
TTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCCTGATGTACCGTTTCGAGGAGGAGCTGTTCTCGCGAGCCTGCAGGACTACAAGATC
35 CAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTTCGCCAAGAGCACCTGATCGAC
AAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTG
40 GCGGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
45 GCGGTGGTGGTGTGTTGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAA
AAATGCATCCCCCCCCC
50 CCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT (SEQ ID NO: 1411)

5'UTR of human ribosomal protein Large 35 (RPL35) lacking the 5' terminal oligopyrimidine tract

GGAGCGGGCGGCGGCGTTGGCGGCTTGTGCAGCA (SEQ ID NO: 1412) 5' UTR of human ribosomal protein Large 21 (RPL21) lacking the 5' terminal oli-

gopyrimidine tract

GGCCGGAACCGCCATCTTCCAGTAATTCGCCAAA (SEO ID NO: 1413)

5'UTR of human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) lacking the 5' terminal oligopyrimidine tract

CGCGCTCGGCCATTTTGTCCCAGTCAGTCCGGAGGCTGCGGCTGCAGAAGTACCGCCTGCGGAGTA
ACTGCAAAG (SEQ ID NO: 1414)

5'UTR of human hydroxysteroid (17-beta) dehydrogenase 4 (HSD17B4) lacking the 5' terminal oligopyrimidine tract

GTCCCGCAGTCGGCGTCTGAGCGGCTCTGCTTGTTCGTGTGTGTCGTTGCAGGCCTTATTC
(SEQ ID NO: 1415)

5'UTR of human androgen-induced 1 (AIG1) lacking the 5' terminal oligopyrimidine tract

GCCGCCCAGCCGGTCCAGGCCTCTGGCGAAC (SEO ID NO: 1416)

5'UTR of human cytochrome c oxidase subunit VIc (COX6C) lacking the 5' terminal oligopyrimidine tract

AGTCAGGAAGGACGTTGGTGTGAGGTTAGCATACGTATCAAGGACAGTAACTACC (SEQ ID NO: 1417)

5'UTR of human N-acylsphingosine amidohydrolase (acid ceramidase) 1 (ASAH1) lacking the 5' terminal oligopyrimidine tract

GCCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCTAGAGCG (SEQ ID NO: 1418)

5'UTR of mouse ribosomal protein Large 21 (mRPL21) lacking the 5' terminal oligopyrimidine tract

GGCCGCCGCAGCCATCTTCCAGTAACTCGCCAAA (SEO ID NO: 1419)

5'UTR of mouse ribosomal protein large 35A (mRPL35A) lacking the 5' terminal oligopyrimidine tract

GCCATCTTTGGCGCCTGTGGAGGCCTGCTGGGAACAGGACTTCTAACAGCAAGT (SEQ ID NO: 1420)

Mouse ribosomal protein Large 21 (mRPL21)

TCCTCCTTTTCGGCCGCCGAGCCATCTTCCAGTAACTCGCCAAAATGCCATCTTCCAGTAACTCGC
CAAATG (SEQ ID NO: 1421)

mouse ribosomal protein large 35A (mRPL35A)

CTTCCTCTTTCCGCCATCTTGGCGCCTGTGGAGGCCTGCTGGGAACAGGACTTCTAACAGCAAGTA
TG (SEQ ID NO: 1422)

The claims defining the invention are as follows:

1. An artificial nucleic acid molecule comprising:
 - a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene;
 - b. at least one open reading frame (ORF); and
 - c. at least one 3'UTR element comprising or consisting of a nucleic acid sequence derived from a 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.
2. The artificial nucleic acid molecule according to claim 1, wherein the at least one 3'UTR element comprises or consist of a nucleic acid sequence which is derived from the 3'UTR of a vertebrate albumin gene or from a variant thereof, the 3'UTR of a mammalian albumin gene or from a variant thereof, or the 3'UTR of a human albumin gene or from a variant thereof.
3. The artificial nucleic acid molecule according to claims 1 or 2, wherein the at least one 3'UTR element exhibits a length of at least about 50 nucleotides, at least about 75 nucleotides, at least about 100 nucleotides, of at least about 125 nucleotides, or at least about 150 nucleotides.
4. The artificial nucleic acid molecule according to any one of claims 1-3, wherein the 5'UTR element and the open reading frame are heterologous and wherein the 5'UTR lacks the 5'TOP motif or the 5'UTR does not contain a functional 5'TOP motif.
5. The artificial nucleic acid molecule according to claim 1-4, wherein the 5'UTR element is suitable for increasing protein production from the artificial nucleic acid molecule.
6. The artificial nucleic acid molecule according to any one of claims 1-5, wherein the 5'UTR element does not comprise a TOP-motif.
7. The artificial nucleic acid molecule according to any one of claims 1-6, wherein the nucleic acid sequence which is derived from the 5'UTR of a TOP gene is derived from the 5'UTR of a eukaryotic TOP gene or from a variant thereof, the 5'UTR of a plant or animal TOP gene or from a variant thereof, the 5'UTR of a chordate TOP

gene or from a variant thereof, the 5'UTR of a vertebrate TOP gene or from a variant thereof, or the 5'UTR of a mammalian TOP gene or from a variant thereof, or the 5'UTR of a human TOP gene, or a variant thereof.

- 5 8. The artificial nucleic acid molecule according to any of claims 1-7, wherein the at least one 3'UTR element and the at least one 5'UTR element act at least additively or synergistically to increase protein production from said artificial nucleic acid molecule.
9. The artificial nucleic acid molecule according to any one of claims 1-8 further comprising a poly(A) sequence and/or a polyadenylation signal.
- 10 10. The artificial nucleic acid molecule according to claim 9, wherein the polyadenylation signal comprises the consensus sequence NN(U/T)ANA, with N = A or U.
11. The artificial nucleic acid molecule according to any one of claims 1-10, wherein the 5'UTR element
- 15 is derived from a 5'UTR of a TOP gene encoding a ribosomal Large protein (RPL) or from a variant of a 5'UTR of a TOP gene encoding a ribosomal Large protein (RPL), wherein the 5'UTR lacks the 5'TOP motif or the 5'UTR does not contain a functional 5'TOP motif;
- 20 comprises or consists of a nucleic acid having an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to a nucleic acid sequence extending from nucleotide position 5 to the nucleotide position immediately 5' to the start codon of a nucleic acid sequence according to any of SEQ ID NOs: 1-1363, 1395, 1421 or 1422, or to a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence, which
- 25 has an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to a nucleic acid sequence extending from nucleotide position 5 to the nucleotide position immediately 5' to the start codon of a nucleic acid sequence according to any of SEQ ID NOs: 1-1363, 1395, 1421 or 1422, or to a corresponding
- 30 RNA sequence; or
- comprises or consists of a nucleic acid sequence having an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to the 5'UTR of a nucleic acid sequence according to any of SEQ ID NOs. 67, 259, 1284-1318, 1344,

1346, 1348-1354, 1357, 1358, 1421 or 1422 or to a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to the 5'UTR of a nucleic acid sequence according to SEQ ID No. SEQ ID NOs: 67, 259, 1284-1318, 1344, 1346, 1348-1354, 1357, and 1358, 1421 or 1422 or to a corresponding RNA sequence.

- 5
12. The artificial nucleic acid molecule according to claim 11, wherein the corresponding RNA sequence lacks the 5'TOP motif.
- 10
13. The artificial nucleic acid molecule according to any one of claims 1-12, wherein the 5'UTR element comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), an ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, an hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), an androgen-induced 1 gene (AIG1), cytochrome c oxidase subunit VIc gene (COX6C), or a N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof; derived from the 5' UTR of a vertebrate ribosomal protein Large 32 gene (RPL32), a vertebrate ribosomal protein Large 35 gene (RPL35), a vertebrate ribosomal protein Large 21 gene (RPL21), a vertebrate ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a vertebrate hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a vertebrate androgen-induced 1 gene (AIG1), a vertebrate cytochrome c oxidase subunit VIc gene (COX6C), or a vertebrate N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof; derived from the 5'UTR of a mammalian ribosomal protein Large 32 gene (RPL32), a ribosomal protein Large 35 gene (RPL35), a ribosomal protein Large 21 gene (RPL21), a mammalian ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a mammalian hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a mammalian androgen-induced 1 gene (AIG1), a mammalian cytochrome c oxidase subunit VIc gene (COX6C), or a mammalian N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof; or derived from the 5'UTR of a human ribosomal protein Large 32 gene (RPL32), a human ribosomal protein Large 35 gene (RPL35), a
- 15
- 20
- 25
- 30
- 35

human ribosomal protein Large 21 gene (RPL21), a human ATP synthase, H⁺ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle (ATP5A1) gene, a human hydroxysteroid (17-beta) dehydrogenase 4 gene (HSD17B4), a human androgen-induced 1 gene (AIG1), a human cytochrome c oxidase subunit VIc gene (COX6C), or a human N-acylsphingosine amidohydrolase (acid ceramidase) 1 gene (ASAH1) or from a variant thereof.

14. The artificial nucleic acid molecule according to claim 13, wherein the 5'UTR element does not comprise the 5'TOP motif of said gene.

15. The artificial nucleic acid molecule according to any one of claims 1-14, wherein the 5'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to the nucleic acid sequence according to SEQ ID NO. 1368 or SEQ ID NOs 1412-1420, or to a corresponding RNA sequence, or wherein the at least one 5'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, at least about 50%, at least about 60%, least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to the nucleic acid sequence according to SEQ ID No. 1368 or SEQ ID NOs 1412-1420, or to a corresponding RNA sequence.

16. The artificial nucleic acid molecule according to any one of claims 1-15, wherein the at least one 3'UTR element comprises or consists of a nucleic acid sequence which has an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to a nucleic acid sequence selected from SEQ ID NOs. 1369-1393 or to a corresponding RNA sequence, or wherein the at least one 3'UTR element comprises or consists of a fragment of a nucleic acid sequence which has an identity of at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99% to a nucleic acid sequence selected from SEQ ID NOs. 1369-1393 or to a corresponding RNA sequence.

17. The artificial nucleic acid molecule according to claim 13 to 15, wherein the fragment consists of a continuous stretch of nucleotides corresponding to a continuous stretch of nucleotides in the full-length sequence, which represents at least 20%, at least 30%,

at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% of the full-length sequence the fragment is derived from.

- 5 18. The artificial nucleic acid molecule according to any one of claims 1-17, wherein the open reading frame does not code for a reporter protein, such as a GFP protein, a luciferase protein, a globin protein, human growth hormone, albumin or human albumin.
- 10 19. The artificial nucleic acid molecule according to any one of claims 1-18, further comprising at least one of a 5'-cap structure, a poly(C) sequence and an IRES-motif.
20. The artificial nucleic acid molecule according to claim 19, wherein the open reading frame is at least partially G/C modified.
21. The artificial nucleic acid molecule according to claim 20, wherein the G/C content of the open reading frame is increased compared to the wild type open reading frame.
- 15 22. The artificial nucleic acid molecule according to any one of claims 1-21, wherein the open reading frame comprises a codon-optimized region or the open reading frame is codon-optimized.
23. The artificial nucleic acid molecule according to any one of claims 1-22, wherein the artificial nucleic acid molecule is an RNA.
24. A vector comprising:
- 20 a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene;
- b. at least one open reading frame (ORF) and/or at least one cloning site; and
- 25 c. at least one 3'UTR element, which comprises or consists of a nucleic acid sequence derived from a 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.
25. The vector according to claim 24, wherein the 5'UTR element and the open reading frame are heterologous.
26. The vector according to claim 24 or 25, which is an RNA or DNA vector.

27. The vector according to any one of claims 24-26, which is a plasmid vector or a viral vector.
28. The vector according to any one of claims 24-27, which comprises or codes for an artificial nucleic acid molecule according to any one of claims 1-23.
- 5 29. The vector according to any one of claims 32-36, which is a circular molecule.
30. The vector according to claim 29, wherein the ORF, the poly(A) sequence or the 3'UTR element of the coding strand is followed in 5'→3' direction by a restriction site for linearization of the circular vector molecule.
- 10 31. A cell comprising the artificial nucleic acid molecule according to any one of claims 1-23 or the vector according to any one of claims 24-30.
- 32 A cell according to claim 31, wherein the cell is a mammalian cell.
33. A cell according to claim 32, wherein the mammalian cell is an isolated human cell.
34. A pharmaceutical composition comprising the artificial nucleic acid molecule according to any one of claims 1-23, the vector according to any one of claims 24-30, or the cell according to any one of claims 31-33.
- 15 35. The pharmaceutical composition according to claim 34, further comprising one or more pharmaceutically acceptable diluents and/or excipients and/or one or more adjuvants.
- 20 36. Use of the artificial nucleic acid molecule according to any one of claims 1-23, the vector according to any one of claims 24-30, the cell according to any one of claims 31-33, or the pharmaceutical composition according to claim 34 or 35 for the manufacture of a medicament.
- 25 37. Use of the artificial nucleic acid molecule according to any one of claims 1-31, the vector according to any one of claims 24-30, the cell according to any one of claims 31-33, or the pharmaceutical composition according to claim 34 or 35 as a vaccine or in gene therapy.
38. A method for treating or preventing a disorder comprising administering the artificial nucleic acid molecule according to any one of claims 1-23, the vector according to any one of claims 24-30, the cell according to any one of claims 31-33,

or the pharmaceutical composition according to claim 34 or 35 to a subject in need thereof.

- 5
39. A method of treating or preventing a disorder comprising transfection of a cell with the artificial nucleic acid molecule according to any one of claims 1-23 or the vector according to any one of claims 24-30.
40. The method according to claim 39, wherein transfection of a cell is performed *in vitro* or *ex vivo* and the transfected cell is administered to a subject in need thereof.
41. The method according to claim 40, wherein the subject is a human patient.
- 10
42. The method according to any one of claims 38-41 wherein the method is a vaccination method or a gene therapy method.
43. A method for increasing protein production from an artificial nucleic acid molecule, comprising the step of providing the artificial nucleic acid molecule with:
- 15
- a. at least one 5'-untranslated region element (5'UTR element) which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene, and
- b. at least one 3'UTR element which comprises or consists of a nucleic acid sequence derived from the 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.
- 20
44. The method according to claim 43, wherein the 5'UTR element does not comprise a TOP-motif.
45. The method according to claim 43 or 44, wherein the 5'UTR lacks the 5'TOP motif or the 5'UTR does not contain a functional 5'TOP motif.
- 25
46. Use of a 5'UTR element which comprises or consists of a nucleic acid sequence which is derived from the 5'UTR of a TOP gene or which is derived from a variant of the 5'UTR of a TOP gene for increasing protein production from a nucleic acid molecule, wherein the 5'UTR element is used in combination with a 3'UTR element, which comprises or consists of a nucleic acid sequence derived from the 3'UTR of an albumin gene or from a variant of a 3'UTR of an albumin gene.

- 5
- 47. Use according to claim 46, wherein the 5'UTR element does not comprise a TOP-motif.
 - 48. The use according to claim 45 or 46, wherein the 5'UTR lacks the 5'TOP motif or the 5'UTR does not contain a functional 5'TOP motif.
 - 49. A kit comprising an artificial nucleic acid molecule according to any one of claims 1-23, the vector according to any one of claims 24-30, or the cell according to any one of claims 31-33, and/or the pharmaceutical composition according to claim 34 or 35.

Nucleotide sequence of PpLuc(GC) – A64N64

GGGAGAAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTA
CCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCT
5 GGTGCCGGGCACGATCGCCTTCACCGACGCCCACATCGAGGTCGACATCACCTACGCGGA
GTACTTCGAGATGAGCGTGC GCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAA
CCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGC
CCTCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCT
GAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAA
10 GATCCTGAACGTGCAGAAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAA
GACCGACTACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGG
CTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGAT
CATGAACAGCAGCGGCAGCACC GGCCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGC
CTGCGTGCGCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACAC
15 CGCCATCCTGAGCGTGGTGGCCTTCCACCACGGCTTCGGCATGTTACGACCCCTGGGCTA
CCTCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTTGCG
GAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCGTTCAGCTTCTT
CGCCAAGAGCACCCGTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGG
GGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCACCTCCCGGG
20 CATCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGG
GGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGA
CCTGGACACCGGCAAGACCCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCC
GATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGA
CGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGT
25 CGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGA
GAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGA
CGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGCTGGAGCACGGCAAGACCATGACGGA
GAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGG
CGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGAT
30 CCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGAT
CTAA
AAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGC
CACCAGAATT

35

Fig. 1

Nucleotide sequence of PpLuc(GC) – albumin7 – A64N64

5 GGGAGAAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTA
CCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCT
GGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTGACATCACCTACGCGGA
GTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAA
CCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGC
CCTCTTCATCGGCGTGCCGTGCGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCT
10 GAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAA
GATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAA
GACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGG
CTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGAT
CATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGC
15 CTGCGTGCGCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACAC
CGCCATCCTGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTACGACCTGGGCTA
CCTCATCTGCGGCTTCCGGGTGGTCCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTTGCG
GAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCTGTTACGCTTCTT
CGCCAAGAGCACCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGG
20 GGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCACCTCCCGGG
CATCCGCCAGGGCTACGGCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGG
GGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGA
CCTGGACACCGGCAAGACCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCC
GATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGA
25 CGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGT
CGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGA
GAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGA
CGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGCTGGAGCACGGCAAGACCATGACGGA
GAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGG
30 CGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGAT
CCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCAT
CACATTTAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAG
CTTATTCATCTCTTTTTCTTTTTCGTTGGTGTAAGCCAACACCCTGTCTAAAAAACATA
AATTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAGAA
35 CCTAGATCTAAAAA
AAA
AAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTT
TCAGAGCCACCAGAATT

Fig. 2

Nucleotide sequence of RPL32 – PpLuc(GC) – A64N64

GGGGCGCTGCCTACGGAGGTGGCAGCCATCTCCTTCTCGGCATCAAGCTTGAGGATGGAG
5 GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCCGTC
10 GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCTC
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
15 GGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCG
TTCCACCACGGCTTCGGCATGTTACGACCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCTTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTTACGCTTCTTCGCCAAGAGCACCTTGATCGAC
20 AAGTACGACCTGTGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
25 AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
GCGGTGGTGGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
30 GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAA
AAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCCCCAAAGGCTCTTTTCAGAGCCACCAGAATT

Nucleotide sequence of RPL32 – PpLuc(GC) – albumin7 – A64N64

GGGGCGCTGCCTACGGAGGTGGCAGCCATCTCCTTCTCGGCATCAAGCTTGAGGATGGAG
5 GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
GGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCCCGTC
10 GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCTC
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
15 GGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCG
TTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCCTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTTACGCTTCTTCGCCAAGAGCACCTGATCGAC
20 AAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
25 AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
30 GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
CCGAAGGGCCTGACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGACATCACATTTAAAGCATCTCAGCC
TACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTCTTTT
TCGTTGGTGTAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCT
35 CTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAA
AAATGCATCCCC
CCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 4

5/30


PpLuc from mRNA with combined UTRs (HDF)

Fig. 5

Nucleotide sequence of RPL35 – PpLuc(GC) – albumin7 – A64N64

GGGGAGCGGGCGGCGGCGTTGGCGGCTTGTGCAGCAAAGCTTGAGGATGGAGGACGCCAA
 GAACATCAAGAAGGGCCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
 5 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTACCGACGC
 CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
 GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
 CCTGCAGTTCTTCATGCCGGTGTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGC
 GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
 10 GGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
 CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
 CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
 CTTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
 GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
 15 CATCTTCGGCAACCAGATCATCCCGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
 CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTTGAT
 GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
 GCTGCTCGTGCCGACCCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGACAAGTACGA
 CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
 20 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
 CACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
 GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCCTGGGCGTGAA
 CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
 GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
 25 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
 GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
 CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
 GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
 GGTGACCACCGCCAAGAAGCTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
 30 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
 CGGCAAGATCGCCGTGTAAGACTAGTGATCACATTTAAAAGCATCTCAGCCTACCATGA
 GAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTTTCTTTTGGT
 GTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
 TGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAAAAAA
 35 AAATGCATCCCCCCCCCCCC
 CCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 6

Nucleotide sequence of RPL21 – PpLuc(GC) – albumin7 – A64N64

GGGGCCGGAACCGCCATCTTCCAGTAATTTCGCCAAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
5 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCGCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
10 GGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
15 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
20 GGCCGTGGCCAAGCGGTTCCACCTCCCGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
25 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
30 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGA
GAATAAGAGAAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTCTTTTTCGTTGGT
GTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
TGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAAAAAA
35 AAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 7

Nucleotide sequence of atp5a1 – PpLuc(GC) – albumin7 – A64N64

GGGCGGCTCGGCCATTTTGTCCCAGTCAGTCCGGAGGCTGCGGCTGCAGAAGTACCGCCT
GCGGAGTAACTGCAAAGAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCG
5 GCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAG
CGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATC
ACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGC
CTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCG
GTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAG
10 CGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAG
GGCCTGCAGAAATCCTGAACGTGCAGAAAGCTGCCCATCATCCAGAAGATCATCATC
ATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCAC
CTCCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCCGGAGAGCTTCGACCGGGACAAGACC
ATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCG
15 CACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATC
ATCCCGGACACCGCCATCCTGAGCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACAG
ACCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAG
CTGTTCTTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCTG
TTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGACCTGTGAACCTGCACGAG
20 ATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTC
CACCTCCCGGGCATCCGCCAGGGCTACGGCTGACCGAGACCACGAGCGCGATCCTGATC
ACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCC
AAGGTGGTGGACCTGGACACCGGCAAGACCTGGGCGTGAACCAGCGGGGCGAGCTGTGC
GTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTC
25 ATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCAC
TTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCG
GCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGG
CTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAG
ACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAG
30 CTGCCGGGGCGGCGTGGTGTTCGTGGACGAGTCCCGAAGGGCCTGACCGGGAAGCTCGAC
GCCCCGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAA
GACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATG
AAGATCAATAGCTTATTCATCTCTTTTCTTTTCTTTTCTGTTGGTGTAAAGCCAACACCCTGTC
TAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAA
35 AATGGAAAGAACCTAGATCTAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCC
AAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 8

Nucleotide sequence of HSD17B4 – PpLuc(GC) – albumin7 – A64N64

GGGTCCCGCAGTCGGCGTCCAGCGGCTCTGCTTGTTCGTGTGTGTGTCGTTGCAGGCCCTT
 ATTCAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCCCTTCTACC
 5 CGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGG
 TGCCGGGCACGATCGCCTTACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGT
 ACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACC
 ACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCC
 TCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGA
 10 ACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGA
 TCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATATGGACAGCAAGA
 CCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGGCT
 TCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCA
 TGAACAGCAGCGGCAGCACCGGCCGTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCT
 15 GCGTGCGCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCG
 CCATCCTGAGCGTGGTGGCGTTCACCACGGCTTCGGCATGTTACGACCCTGGGCTACC
 TCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTTGCGGA
 GCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTTCG
 CCAAGAGCACCTTGATCGACAAGTACGACCTGTCAACCTGCACGAGATCGCCAGCGGGG
 20 GCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCA
 TCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGG
 ACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACC
 TGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGA
 TGATCATGAGCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACG
 25 GCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCG
 ACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGA
 GCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACG
 ACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGGAGCACGGCAAGACCATGACGGAGA
 AGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCG
 30 TGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCC
 GCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCA
 CATTTAAAAGCATCTCAGCCTACCATGAGAATAAGAGAAAAGAAAATGAAGATCAATAGCT
 TATTCATCTCTTTTCTTTTTCGTTGGTGTAAAGCCAACACCCTGTCTAAAAAACATAAA
 TTTCTTTAATCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACC
 35 TAGATCTAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTC
 AGAGCCACCAGAAAT

Fig. 9

Nucleotide sequence of AIG1 – PpLuc(GC) – albumin7 – A64N64

GGGCCGCCCAGCCGGTCCAGGCCTCTGGCGAACAAGCTTGAGGATGGAGGACGCCAAGAA
CATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCT
5 CCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCCA
CATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGC
CATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCT
GCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTCGCCCCGGCGAA
CGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGT
10 GTTCGTGAGCAAGAAGGGCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCAT
CCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACAC
GTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTT
CGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAA
GGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCAT
15 CTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGGCGTTCCACCACGG
CTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTA
CCGGTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCT
GCTCGTGCCGACCCTGTTACGTTCTTCGCCAAGAGCACCTGATCGACAAGTACGACCT
GTCGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGC
20 CGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCAC
GAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGT
CCC GTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTGGGCGTGAACCA
GCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGA
GGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTG
25 GGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGG
CTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGA
CGCCGGCGTGCGCCGGGCTGCCGGACGACGACGCGCGGCGAGCTGCCGGCCGCGGTGGTGGT
GCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGT
GACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCT
30 GACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGG
CAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCCTACCATGAGAA
TAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTCTTTTCTGTTGGTGTA
AAGCCAACACCCTGTCTAAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTCTGT
GCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAAAAAAAAAA
35 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 10

11/30

Nucleotide sequence of COX6C – PpLuc(GC) – albumin7 – A64N64

GGAGTCAGGAAGGACGTTGGTGTGAGGTTAGCATACGTATCAAGGACAGTAACTACCAA
GCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGG
5 AGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGG
GCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCG
AGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGA
TCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCA
TCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCA
10 TGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGA
ACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACT
ACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACG
AGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACA
GCAGCGGCAGCACCGGCCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGC
15 GCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCC
TGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTTACGACCCTGGGCTACCTCATCT
GCGGCTTCCGGGTGGTCCCTGATGTACCGGTTTCGAGGAGGAGCTGTTTCTGCGGAGCCTGC
AGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTACGCTTCTTCGCCAAGA
GCACCCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCC
20 CGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCC
AGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACA
AGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACA
CCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCA
TGAGCGGCTACGTGAACAACCCGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGC
25 TGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGC
TGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCC
TGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCG
GCGAGCTGCCGGCCGCGGTGGTGGTGTGAGCACGGCAAGACCATGACGGAGAAGGAGA
TCGTGCACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCCGGGGCGGCGTGGTGT
30 TCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGA
TCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGATCACATTTA
AAAGCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCA
TCTCTTTTTCTTTTTCTTTTGGTGTAAAGCCAACACCCTGTCTAAAAAACATAAATTTCTT
TAATCATTTTGCCTCTTTTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCCTAGATC
35 TAAAAAATAA
AAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCC
ACCAGAATT

Fig. 11

12/30

Nucleotide sequence of ASAH1 – PpLuc(GC) – albumin7 – A64N64

GGGCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCTAGAGCGAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCC
5 GCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTC
GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
10 CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCATCATCCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTCTGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCT
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
GGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
15 CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGGCG
TTCCACCACGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCTGATGTACCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTTACGCTTCTTCGCCAAGAGCACCCCTGATCGAC
AAGTACGACCTGTGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
20 GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
25 ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCGCGGCTGGCCGGGCTGCCGGACGACGACGCGCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
30 CCGAAGGGCCTGACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAAGCATCTCAGCC
TACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTTCTTTT
TCGTTGGTGTAAAGCCAACACCCTGTCTAAAAACATAAATTTCTTTAATCATTTTGCCT
CTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAA
35 AAATGCATCCCC
CCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 12

13/30

Nucleotide sequence of mRPL21 – PpLuc(GC) – albumin7 – A64N64

GGGGCCGCCGCGAGCCATCTTCCAGTAACTCGCCAAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
5 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCGCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
10 GGTGTTCTGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTCTGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCC
15 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
20 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
25 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
30 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTGATCACATTTAAAAGCATCTCAGCCTACCATGA
GAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCTCTTTTCTTTTTCGTTGGT
GTAAAGCCAACACCCTGTCTAAAAACATAAATTTCTTTAATCATTTTGCCTCTTTTCTC
TGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAAAAAATAAAAAAAAAAAAAA
35 AAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

Fig. 13

Nucleotide sequence of mRPL35A – PpLuc(GC) – albumin7 – A64N64

GGGCCATCTTGGCGCCTGTGGAGGCCTGCTGGGAACAGGACTTCTAACAGCAAGTAAGCT
TGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGCGCCCTTCTACCCGCTGGAGG
5 ACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCA
CGATCGCCTTCACCGACGCCACATCGAGGTGCACATCACCTACGCGGAGTACTTCGAGA
TGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCG
TGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCG
GCGTGGCCGTCGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGG
10 GGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACG
TGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACC
AGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGGCTTCAACGAGT
ACGACTTTCGTCCCGGAGAGCTTCGACCCGGGACAAGACCATCGCCCTGATCATGAACAGCA
GCGGCAGCACCGGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCT
15 TCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGA
GCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCG
GCTTCCGGGTGGTCTGTATGTACCGGTTTCGAGGAGGAGCTGTTCTCGCGGAGCCTGCAGG
ACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCA
CCCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGC
20 TGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGTTCCACCTCCCGGGCATCCGCCAGG
GCTACGGCTGACCGAGACCACGAGCGGATCCTGATCACCCCGAGGGGGACGACAAGC
CGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCG
GCAAGACCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGA
GCGGCTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGC
25 ACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGA
AGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGC
TCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGGCTGCCGGACGACGACGCCGGCG
AGCTGCCGGCCGCGGTGGTGGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCG
TCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCG
30 TGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCC
TGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTGCATCACATTTAAAA
GCATCTCAGCCTACCATGAGAATAAGAGAAAGAAAATGAAGATCAATAGCTTATTCATCT
CTTTTTCTTTTCGTTGGTGTAAGCCAACACCCTGTCTAAAAACATAAATTTCTTTAA
TCATTTTGCCTCTTTTCTCTGTGCTTCAATTAATAAAAAATGGAAAGAACCTAGATCTAA
35 AA
AATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACC
AGAATT

Nucleotide sequence of RPL35 – PpLuc(GC) – A64N64

GGGGAGCGGGCGGGCGGCGTTGGCGGCTTGTGCAGCAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
5 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGCGCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
10 GGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCCGGGACCC
15 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTCTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
20 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
25 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
30 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCC
CCCCCCCCCCCCAAAGGCTCTTTTCAGAGCCACCAGAATT

35

Fig. 15

Nucleotide sequence of RPL21 – PpLuc(GC) – A64N64

GGGGCCGGAACCGCCATCTTCCAGTAATTCGCCAAAAAGCTTGAGGATGGAGGACGCCAA
GAACATCAAGAAGGGCCCGGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCA
5 GCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTACCGACGC
CCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGA
GGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAG
CCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTCGCCCCGGC
GAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGT
10 GGTGTTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCAT
CATCCAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTA
CACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAG
CTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCC
GAAGGGGGTGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCC
15 CATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCA
CGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTTGAT
GTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGC
GCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGA
CCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGA
20 GGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGAC
CACGAGCGCGATCCTGATCACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGT
GGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAA
CCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCC
GGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTA
25 CTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAA
GGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTT
CGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGT
GGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCA
GGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGG
30 CCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGG
CGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCC
CCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

35

Fig. 16

Nucleotide sequence of atp5a1 – PpLuc(GC) – A64N64

GGGCGGCTCGGCCATTTTGTCCCAGTCAGTCCGGAGGCTGCGGCTGCAGAAGTACCGCCT
GCGGAGTAACTGCAAAGAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCG
5 GCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAG
CGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCACATCGAGGTCGACATC
ACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGC
CTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCG
GTGCTGGGCGCCCTCTTCATCGGCGTGCCGTCGCCCCGGCGAACGACATCTACAACGAG
10 CGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAG
GGCCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATC
ATGGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACACGTTCTGTGACCAGCCAC
CTCCCCCGGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACC
ATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCG
15 CACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATC
ATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTACG
ACCCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAG
CTGTTCTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTG
TTCAGCTTCTTCGCCAAGAGCACCTGATCGACAAGTACGACCTGTGAACTGCACGAG
20 ATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTC
CACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATC
ACCCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCC
AAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGC
GTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGAGGCCACCAACGCCCTC
25 ATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCAC
TTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCG
GCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGCCGGG
CTGCCGGACGACGACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGGTGGAGCACGGCAAG
ACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAG
30 CTGCGGGGCGCGTGTTCTGTTGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGAC
GCCCCGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAA
GACTAGTAGATCTAAA
AAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCT
CTTTTCAGAGCCACCAGAATT
35

Fig. 17

Nucleotide sequence of HSD17B4 – PpLuc(GC) – A64N64

GGGTCCCGCAGTCGGCGTCCAGCGGCTCTGCTTGTTCGTGTGTGTGTGTCGTTGCAGGCCTT
ATTCAAGCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGCGCCCTTCTACC
5 CGCTGGAGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGG
TGCCGGGCACGATCGCCTTCACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGT
ACTTCGAGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACC
ACCGGATCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGTGGGCGCCC
TCTTCATCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGA
10 ACAGCATGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGA
TCCTGAACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGA
CCGACTACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCT
TCAACGAGTACGACTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCA
TGAACAGCAGCGGCAGCACCGGCCTGCCGAAGGGGTGGCCCTGCCGCACCGGACCGCCT
15 GCGTGCGCTTCTCGCACGCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCG
CCATCCTGAGCGTGGTGGCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACC
TCATCTGCGGCTTCCGGGTGGTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTCGCGGA
GCCTGCAGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCCTGTTACGCTTCTTCG
CCAAGAGCACCTGATCGACAAGTACGACCTGTCAACCTGCACGAGATCGCCAGCGGGG
20 GCGCCCCGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCA
TCCGCCAGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCCGAGGGGG
ACGACAAGCCGGGCGCCGTGGGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACC
TGGACACCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGA
TGATCATGAGCGGTACGTGAACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACG
25 GCTGGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCG
ACCGGCTGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGA
GCATCCTGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACG
ACGCCGGCGAGCTGCCGGCCGCGGTGGTGGTGTGAGCACGGCAAGACCATGACGGAGA
AGGAGATCGTCGACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCG
30 TGGTGTTCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCC
GCGAGATCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCT
AAA
AAAATGCATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCA
CCAGAATT
35

Fig. 18

Nucleotide sequence of AIG1 – PpLuc(GC) – A64N64

GGGCCGCCCAGCCGGTCCAGGCCTCTGGCGAACAAGCTTGAGGATGGAGGACGCCAAGAA
CATCAAGAAGGGCCCGCGCCCTTCTACCCGCTGGAGGACGGGACCGCCGGCGAGCAGCT
5 CCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTCACCGACGCCCA
CATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGCCTGGCCGAGGC
CATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCGGAGAACAGCCT
GCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTCGCCCCGGCGAA
CGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAGCCGACCGTGGT
10 GTTCGTGAGCAAGAAGGGCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCAT
CCAGAAGATCATCATATGACAGCAAGACCGACTACCAGGGCTTCCAGTCGATGTACAC
GTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTCCCGGAGAGCTT
CGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACCGGCCTGCCGAA
GGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCCCGGGACCCCAT
15 CTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGCCGTTCCACCACGG
CTTCGGCATGTTACGACCTGGGCTACCTCATCTGCGGCTTCCGGGTGGTCTGATGTA
CCGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATCCAGAGCGCGCT
GCTCGTGCCGACCCTGTTACGTTCTTCGCCAAGAGCACCTGATCGACAAGTACGACCT
GTCGAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAGGTGGGCGAGGC
20 CGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTGACCGAGACCAC
GAGCGGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTGGGCAAGGTGGT
CCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCTGGGCGTGAACCA
GCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTGAACAACCCGGA
GGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGACATCGCCTACTG
25 GGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATCAAGTACAAGGG
CTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCCAACATCTTCGA
CGCCGGCGTGCGCGGGCTGCCGGACGACGACGCGGCGAGCTGCCGGCCGCGGTGGTGGT
GCTGGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTGGCCAGCCAGGT
GACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTCCCGAAGGGCCT
30 GACCGGGAAGCTCGACGCCCCGGAAGATCCGCGAGATCCTGATCAAGGCCAAGAAGGGCGG
CAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATGCATCCCCCCCCCCCCCCCCCCCC
CCCCCCCCCAAAGGCTCTTTTCAGAGCCACCAGAATT

35

Fig. 19

20/30

Nucleotide sequence of COX6C – PpLuc(GC) – A64N64

GGAGTCAGGAAGGACGTTGGTGTGAGGTTAGCATACGTATCAAGGACAGTAACTACCAA
GCTTGAGGATGGAGGACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGG
5 AGGACGGGACCGCCGGCGAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGG
GCACGATCGCCTTCACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCG
AGATGAGCGTGCGCCTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGA
TCGTGGTGTGCTCGGAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCA
TCGGCGTGGCCGTGCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCA
10 TGGGGATCAGCCAGCCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGA
ACGTGCAGAAGAAGCTGCCCATCATCCAGAAGATCATCATCATGGACAGCAAGACCGACT
ACCAGGGCTTCCAGTCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACG
AGTACGACTTTCGTCCCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACA
GCAGCGGCAGCACCGGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGC
15 GCTTCTCGCACGCCCCGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCC
TGAGCGTGGTGCCGTTCCACCACGGCTTCGGCATGTTACGACCCTGGGCTACCTCATCT
GCGGCTTCCGGGTGGTCTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTTGCAGGAGCCTGC
AGGACTACAAGATCCAGAGCGCGCTGCTCGTGCCGACCTGTTTCAGCTTCTTCGCCAAGA
GCACCCTGATCGACAAGTACGACCTGTGCAACCTGCACGAGATCGCCAGCGGGGGCGCCC
20 CGCTGAGCAAGGAGGTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCC
AGGGCTACGGCCTGACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACA
AGCCGGGCGCCGTGGGCAAGGTGGTCCCGTTCTTCGAGGCCAAGGTGGTGGACCTGGACA
CCGGCAAGACCCTGGGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCA
TGAGCGGCTACGTGAACAACCCGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGC
25 TGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGC
TGAAGTCGCTGATCAAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCC
TGCTCCAGCACCCCAACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCG
GCGAGCTGCCGGCCGCGGTGGTGGTGCTGGAGCACGGCAAGACCATGACGGAGAAGGAGA
TCGTGCACTACGTGGCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGT
30 TCGTGGACGAGGTCCCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGA
TCCTGATCAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAA
AAATG
CATCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAA
TT
35


Fig. 20

Nucleotide sequence of ASAH1 – PpLuc(GC) – A64N64

GGGCCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCTAGAGCGAAGCTTGAGGATGGAG
GACGCCAAGAACATCAAGAAGGGCCCGGCGCCCTTCTACCCGCTGGAGGACGGGACCGCC
5 GGCAGCAGCTCCACAAGGCCATGAAGCGGTACGCCCTGGTGCCGGGCACGATCGCCTTC
ACCGACGCCCACATCGAGGTCGACATCACCTACGCGGAGTACTTCGAGATGAGCGTGCGC
CTGGCCGAGGCCATGAAGCGGTACGGCCTGAACACCAACCACCGGATCGTGGTGTGCTCG
GAGAACAGCCTGCAGTTCTTCATGCCGGTGCTGGGCGCCCTCTTCATCGGCGTGGCCGTC
10 GCCCCGGCGAACGACATCTACAACGAGCGGGAGCTGCTGAACAGCATGGGGATCAGCCAG
CCGACCGTGGTGTTCGTGAGCAAGAAGGGCCTGCAGAAGATCCTGAACGTGCAGAAGAAG
CTGCCCCATCATCCAGAAGATCATCATATGGACAGCAAGACCGACTACCAGGGCTTCCAG
TCGATGTACACGTTTCGTGACCAGCCACCTCCCGCCGGGCTTCAACGAGTACGACTTCGTC
CCGGAGAGCTTCGACCGGGACAAGACCATCGCCCTGATCATGAACAGCAGCGGCAGCACC
GGCCTGCCGAAGGGGGTGGCCCTGCCGCACCGGACCGCCTGCGTGCGCTTCTCGCACGCC
15 CGGGACCCCATCTTCGGCAACCAGATCATCCCGGACACCGCCATCCTGAGCGTGGTGGCG
TTCCACCACGGCTTCGGCATGTTACGACCCCTGGGCTACCTCATCTGCGGCTTCCGGGTG
GTCTGATGTACCGGTTTCGAGGAGGAGCTGTTCTGCGGAGCCTGCAGGACTACAAGATC
CAGAGCGCGCTGCTCGTGCCGACCCTGTTTCAGCTTCTTCGCCAAGAGCACCCCTGATCGAC
AAGTACGACCTGTCAACCTGCACGAGATCGCCAGCGGGGGCGCCCCGCTGAGCAAGGAG
20 GTGGGCGAGGCCGTGGCCAAGCGGTTCCACCTCCCGGGCATCCGCCAGGGCTACGGCCTG
ACCGAGACCACGAGCGCGATCCTGATCACCCCGAGGGGGACGACAAGCCGGGCGCCGTG
GGCAAGGTGGTCCCCTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCCCTG
GGCGTGAACCAGCGGGGCGAGCTGTGCGTGCGGGGGCCGATGATCATGAGCGGCTACGTG
AACAACCCGGAGGCCACCAACGCCCTCATCGACAAGGACGGCTGGCTGCACAGCGGCGAC
25 ATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTCGACCGGCTGAAGTCGCTGATC
AAGTACAAGGGCTACCAGGTGGCGCCGGCCGAGCTGGAGAGCATCCTGCTCCAGCACCCC
AACATCTTCGACGCCGGCGTGGCCGGGCTGCCGGACGACGACGCCGGCGAGCTGCCGGCC
GCGGTGGTGGTGTGAGCACGGCAAGACCATGACGGAGAAGGAGATCGTCGACTACGTG
GCCAGCCAGGTGACCACCGCCAAGAAGCTGCGGGGCGGCGTGGTGTTCGTGGACGAGGTC
30 CCGAAGGGCCTGACCGGGAAGCTCGACGCCCGGAAGATCCGCGAGATCCTGATCAAGGCC
AAGAAGGGCGGCAAGATCGCCGTGTAAGACTAGTAGATCTAAAAAAAAAAAAAAAAAAAA
AAATGCATCCCCCCCCCCCC
CCCCCCCCCCCCCCCCCAAGGCTCTTTTCAGAGCCACCAGAATT

35

Fig. 21






Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30