



(86) Date de dépôt PCT/PCT Filing Date: 2015/08/12
(87) Date publication PCT/PCT Publication Date: 2016/02/18
(85) Entrée phase nationale/National Entry: 2017/02/13
(86) N° demande PCT/PCT Application No.: US 2015/044883
(87) N° publication PCT/PCT Publication No.: 2016/025615
(30) Priorité/Priority: 2014/08/12 (US62/036,438)

(51) Cl.Int./Int.Cl. *C12N 9/64* (2006.01),
C07K 14/745 (2006.01), *C12N 15/12* (2006.01),
C12N 15/57 (2006.01), *C12N 15/79* (2006.01),
C12N 5/10 (2006.01), *C12N 9/74* (2006.01),
C12P 21/06 (2006.01)

(71) Demandeurs/Applicants:
BAXALTA INCORPORATED, US;
BAXALTA GMBH, CH

(72) Inventeurs/Inventors:
BOHM, ERNST, AT;
HORLING, FRANZISKA, AT;
KOEHN, JADRANKA, AT;
DOCKAL, MICHAEL, AT

(74) Agent: SMART & BIGGAR

(54) Titre : PRODUCTION DE FACTEUR X FONCTIONNEL ENTIEREMENT TRAITE DANS UN SYSTEME
D'EXPRESSION DE MAMMIFERE SECRETANT DE LA FURINE
(54) Title: PRODUCTION OF FULLY PROCESSED AND FUNCTIONAL FACTOR X IN A FURIN-SECRETING
MAMMALIAN EXPRESSION SYSTEM

(57) **Abrégé/Abstract:**

Disclosed herein are methods for production of fully-processed mature Factor X in an expression system producing a controlled amount of furin between 50 U/mL and 300 U/mL of culture supernatant. Also disclosed are transformed cells, expression systems, and expression vectors for the expression of furin and Factor X.



PRODUCTION OF FULLY PROCESSED AND FUNCTIONAL FACTOR X IN A FURIN-SECRETING MAMMALIAN EXPRESSION SYSTEM

FIELD

[0001] Disclosed herein are expression systems, transformed cells, and methods related thereto, for expression of fully-processed and active recombinant Factor X in the presence of furin.

BACKGROUND

[0002] Human coagulation Factor X (FX), activated FX (FXa), and variants thereof, are used as therapeutic agents in blood coagulation disorders including, but not limited to, hemophilia and von Willebrand disease. FX, a vitamin K-dependent serine protease, is synthesized as a single chain precursor protein in the endoplasmic reticulum, with subsequent intracellular proteolytic furin cleavage in the Golgi apparatus before secretion by the producing cell into the blood stream, or into the culture medium in case of recombinant expression. Three furin cleavage sites in FX are responsible for proper FX proteolytic processing. The mature form of FX is a disulfide-linked two-chain molecule consisting of a heavy and light chain, formed after cleavage of the precursor protein. Further modifications of the molecule include γ -carboxylation of the light chain and N- and O-linked glycosylation of the activation peptide which is attached to the heavy chain.

[0003] Besides FX, further Vitamin K-dependent coagulation factors bearing the consensus recognition site Arg-X-Lys/Arg-Arg (SEQ ID NO:4) are substrates of the ubiquitously expressed endoprotease furin, also known as paired basic amino acid residue-cleaving enzyme (PACE). Adequate proteolytic processing of recombinant proteins of the coagulation cascade are impaired in cell culture expression systems due to intracellular processing limitations at high yield expression. Similar to von Willebrand Factor and coagulation Factor IX (FIX) which exhibit insufficient proteolytic processing at high expression rates in recombinant mammalian cells, FX secretion in low producing CHO cell clones is characterized by fully processed FX, whereas high producing clones comprise unprocessed single chain FX and multiple unprocessed forms of FX light chain, in addition to the correctly processed FX heavy and light chain species. Types and degrees of unprocessed FX light chain varied among individual cell clones and under different cell culture conditions such as cell density. Additional *in vivo* furin co-expression or post-cell culture *in vitro* furin incubation is needed to support the endogenous furin proteolytic machinery, facilitating intact protein cleavage.

[0004] Furin co-expression is indispensable for the expression of fully processed FX at high yield. However, to date no threshold level of furin has been reported that would ensure a high percentage of intact processed FX in cell culture systems. High levels of furin are toxic, therefore levels of furin expression by FX-producing mammalian expression systems must be balanced between levels that are toxic, yet potentially process 100% of the FX precursor protein, and those that are too low, resulting in healthy cell cultures in which suboptimal processed FX is produced.

SUMMARY

[0005] Disclosed herein is a system for expressing furin and a human Factor X (FX) in the same cell line and thereby providing a critical furin concentration in the culture supernatant for the generation of fully processed and fully active FX, while maintaining the viability of the culture.

[0006] Thus, disclosed herein is a transformed cell comprising a nucleotide sequence encoding a human furin, such that the transformed cell expresses and secretes functional furin into a culture supernatant, wherein the functional furin is secreted at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for between about 36 and about 78 hours. In one embodiment, the transformed cells further comprise a nucleotide sequence encoding a protein cleavable by furin and exhibiting an Arg-(Lys/Arg)-Arg motif. In another embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on different expression vectors. In another embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on the same expression vector.

[0007] Also provided is a eukaryotic protein expression system comprising a cell line suitable for expression of mammalian proteins; a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and a second expression vector adapted for expression of a protein by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a protein cleavable by furin and exhibiting an Arg-(Lys/Arg)-Arg motif, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for between about 36 and about 78 hours.

[0008] Also provided is a eukaryotic protein expression system comprising a cell line suitable for expression of mammalian proteins; a first expression vector adapted for expression of human furin and a protein cleavable by furin, and exhibiting an Arg-(Lys/Arg)-Arg motif, by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide and a nucleotide sequence encoding the protein cleavable by furin, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for between about 36 and about 78 hours. In one embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on different expression vectors. In another embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on the same expression vector.

[0009] Also provided is a transformed cell comprising a first nucleotide sequence encoding a human furin and a second nucleotide sequence encoding a human FX, such that the transformed cell expresses and secretes functional furin and FX into a culture supernatant, wherein the furin is secreted at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for between about 36 and about 78 hours and at least 85% of the FX is fully processed. In one embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the FX are on different expression vectors. In another embodiment, the nucleotide sequence encoding human furin and the nucleotide sequence encoding the FX are on the same expression vector.

[0010] Also provided is a eukaryotic protein expression system comprising a cell line suitable for expression of mammalian proteins; a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and a second expression vector adapted for expression of FX by the cell line, wherein the second expression vector includes a nucleotide sequence encoding FX, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for between about 36 and about 78 hours.

[0011] Further provided is a eukaryotic protein expression system comprising a cell line suitable for expression of mammalian proteins; a first expression vector adapted for expression of human furin and FX, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide and a nucleotide sequence encoding FX, wherein the cell

line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for between about 36 and about 78 hours.

[0012] Also provided is a method of preparing a recombinant protein comprising transfecting a cell line suitable for expression of mammalian proteins with a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and transfecting the cell line with a second expression vector adapted for expression of a protein by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a protein exhibiting an Arg-(Lys/Arg)-Arg motif; wherein the cell line transfected with the first and the second expression vectors expresses and secretes functional human furin at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for between about 40 and about 80 hours or about 36 and about 78 hours. In one embodiment, the cell line is transfected with the first expression vector and the second expression vector substantially simultaneously. In another embodiment, the cell line is transfected with the first expression vector and cells secreting stable levels of furin are obtained prior to transfecting the cell line with the second expression vector. In yet another embodiment, the cell line is transfected with the second expression vector and cells secreting stable levels of the protein are obtained prior to transfecting the cell line with the first expression vector.

[0013] In one embodiment, the protein is von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z. In another embodiment, the protein is Factor X.

[0014] Also provided is a method of preparing a recombinant protein comprising transfecting a cell line suitable for expression of mammalian proteins with a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and transfecting the cell line with a second expression vector adapted for expression of FX by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a FX polypeptide; wherein the cell line transfected with the first and the second expression vectors expresses and secretes functional human furin at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for between about 36 and about 78 hours. In one embodiment, the cell line is transfected with the first expression vector and the second expression vector substantially simultaneously. In another embodiment, the cell line is transfected with the first expression vector and cells secreting stable levels of furin are obtained prior to transfecting the cell line with the second expression vector. In yet another embodiment, the cell line is transfected with the

second expression vector and cells secreting stable levels of the protein are obtained prior to transfecting the cell line with the first expression vector.

[0015] In another embodiment, the cells are capable of secreting functional furin into the culture supernatant at a concentration of at least about 50 to about 60 U/mL after culture for between about 36 and about 78 hours and wherein at least 90% of the FX is fully processed. In another embodiment, the cells are capable of secreting functional furin into the culture supernatant at a concentration of at least about 90 to about 100 U/mL after culture for between about 36 and about 78 hours and wherein at least 95% of the FX is fully processed.

[0016] Also provided is a recombinant FX produced by a transformed cell disclosed herein.

[0017] Further provided is a recombinant FX produced by an expression system disclosed herein.

[0018] Also provided is a recombinant FX produced by a method disclosed herein.

[0019] Also provided is an expression system for recombinant FX adapted to secrete furin into a culture supernatant at a concentration of between about 50 U/mL and about 300 U/mL after culture for between about 36 and about 78 hours.

[0020] Also provided is a method of producing mature, fully-processed FX comprising an expression system secreting furin into a culture supernatant at a concentration between about 50 U/mL and about 300 U/mL after culture for between about 36 and about 78 hours.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] FIG. 1A depicts the RCL.012-74.pD3H-Furin expression vector and FIG. 1B depicts the nucleotide sequence of the vector (SEQ ID NO:1). The human furin sequence is underlined and the start and stop codons are double underlined.

[0022] FIG. 2 depicts the nucleotide sequence of human furin (SEQ ID NO:2). Start and stop codons are double underlined.

[0023] FIG. 3 depicts the amino acid sequence of human furin (SEQ ID NO:3).

[0024] FIG. 4 depicts the degree of fully processed Factor X (FX) in cultures. Densitometric quantification was conducted of a FX Western blot under reducing conditions and stained with a polyclonal anti-FX antibody. The clones (clone ID 42 to 52) exhibit up to 4 species of FX with varying pixel intensities including the unprocessed FX single chain (box 1, 5, 9, etc.), the FX heavy chain (box 2, 6, 10, etc.), the unprocessed propeptide-containing FX light chain

(box 3, 7, 11, etc.) and the processed FX light chain (box 4, 8, 12, etc.). The pixel intensity of boxes 45-48 was determined for background subtraction.

[0025] FIG. 5 depicts the secreted furin concentration and the percentage of fully processed FX/total FX in culture. A dose-response relationship exists between the secreted furin concentration in the cell culture supernatant (determined with a furin activity assay) and the % fully processed FX/total FX (determined by densitometric quantification of respective bands in Western blots).

[0026] FIG. 6 depicts an analysis of furin dose and fully processed FX. Data (circles) with cell specific (dark lines) and population average (light lines) predicted fully processed FX/total FX (%) as a function of furin concentration with the $E_{\max}$ model.

[0027] FIGs. 7A-D depict an $E_{\max}$ model validity test. FIG. 7A depicts residual of response versus predicted response where data points are scattered symmetrically around zero indicating no systematic trend. FIG. 7B depicts a normal Q-Q plot for the residuals indicating that the assumption of normal distributed errors hold as data points are scattered around the line of identity. FIG. 7C depicts residual of response versus cell line where data points are scattered symmetrically around zero indicating no systematic trend. FIG. 7D depicts observed and predicted values plotted against each other indicating a good fit of the data as the data points are scattered symmetrically around the line of identity.

[0028] FIG. 8 depicts a null model for testing the hypothesis that the degree of FX processing is independent of furin concentration. Data (circles) and the fitted null model with intercepts only assuming that processed FX is independent from the furin concentration (cell specific and population averages fits as dark and light lines, respectively).

[0029] FIG. 9 depicts a dose-response curve and calculated furin minimum concentrations to yield 90% and 95% processed FX. The population average predicted of fully processed FX/total FX (%) as a function of furin concentration (black line) along with furin concentrations that give 90% and 95% fully processed FX/total FX obtained by numerical optimization of the fitted model.

DETAILED DESCRIPTION

[0030] Provided herein are transformed cells, eukaryotic expression systems, methods for producing recombinant proteins, and recombinant proteins made by the methods, all directed to expression of furin and Factor X (FX) in the same cell line and thereby providing a critical furin

concentration in the culture supernatant for the generation of fully processed and mature FX, while maintaining the viability of the culture.

[0031] Common to the transformed cells, eukaryotic expression systems and methods for production of recombinant proteins is the ability of the host cell to produce consistent levels of recombinant furin. Furin is necessary for cleavage of certain mammalian proteins, including FX, from a precursor protein form to a mature, fully processed form. Low concentrations of furin in the culture supernatant of the expression system result in accumulation of propeptide-containing and other non- or partially-processed forms of the protein. Concentrations of furin that are too high result in impaired growth of the host cells and ultimately cell death.

[0032] As used herein, the term “furin” includes full-length furin as well as any furin fragment capable of cleavage of the consensus recognition site Arg–X–Lys/Arg–Arg. Active truncated forms of furin are known in the art and are suitable for use in the instant disclosure. Non-limiting examples of suitable furin fragments can be found in U.S. 6,210,926 and Preininger et al. (Cytotechnology 30:1-15, 1999), both of which are incorporated by reference herein for all they disclose regarding truncated forms of recombinant furin.

[0033] Also within the scope of the present disclosure are variants of the furin and Factor X proteins disclosed herein. For example, conservative amino acid changes may be made, which although they alter the primary sequence of the protein or peptide, do not normally alter its function. Conservative amino acid substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; and lysine, arginine; phenylalanine, tyrosine.

[0034] Also included are fusion proteins, or other modifications, or FX which have increased half-life after administration to a subject. Examples of such would be fusions with an immunoglobulin Fc domain, an albumin domain, an extended (XTEN) recombinant polypeptide (see US 8,673,860 which is incorporated by reference herein for all it discloses regarding XTEN polypeptides), poly Glu or poly Asp sequences, transferrin, or a PAS (Pro Ala Ser)-containing polypeptides attached to the FX sequence.

[0035] Modifications (which do not normally alter primary sequence) include *in vivo*, or *in vitro* chemical derivatization of polypeptides, e.g., acetylation, or carboxylation. Also included are modifications of glycosylation, e.g., those made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing or in further processing steps; e.g. by exposing the polypeptide to enzymes which affect glycosylation, e.g., mammalian glycosylating or

deglycosylating enzymes. Also embraced are sequences which have phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine. Proteins can also be modified chemically after purification with water soluble biocompatible polymers, e.g., polyethylene glycol, polysialic acid, or hydroxyethyl starch.

[0036] Also included are polypeptides which have been modified using ordinary molecular biological techniques so as to improve their resistance to proteolytic degradation or to optimize solubility properties. Analogs of such polypeptides include those containing residues other than naturally occurring L-amino acids, e.g., D-amino acids or non-naturally occurring synthetic amino acids. The peptides of the invention are not limited to products of any of the specific exemplary processes listed herein.

[0037] The disclosure herein is generally directed to systems, transformed cells, expression vectors, and methods for producing at least one recombinant mammalian protein which is post-translationally processed by furin (a recombinant furin-requiring mammalian protein). The mammalian protein is one or more of von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z. In another embodiment, the protein is FX.

[0038] The concentration of furin in the culture supernatant is targeted within an optimum range for production of mature, fully-processed proteins while maintaining viability of the culture after a defined period of culture. Thus, a useful concentration of furin in a culture supernatant for the production of a mature, fully-processed mammalian protein is between about 50 U/mL and about 400 U/mL, between about 50 U/mL and about 350 U/mL, between about 50 U/mL and about 300 U/mL, between about 50 U/mL and about 250 U/mL, between about 50 U/mL and about 200 U/mL, between about 50 U/mL and about 175 U/mL, between about 50 U/mL and about 150 U/mL, between about 50 U/mL and about 125 U/mL, or between about 50 U/mL and about 100 U/mL. In one embodiment, the concentration of furin in the culture supernatant is not less than 50 U/mL.

[0039] In other embodiment, the useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein after a defined period of culture is between about 50 U/mL and about 60 U/mL, between about 55 U/mL and about 65 U/mL, between about 60 U/mL and about 70 U/mL, between about 65 U/mL and about 75 U/mL, between about 70 U/mL and about 80 U/mL, between about 75 U/mL and about 85 U/mL, between about 80 U/mL and about 90 U/mL, between about 85 U/mL and about 95 U/mL, between about 90 U/mL and about 95 U/mL, between about 95 U/mL and about 105 U/mL, between about 100 U/mL and about 110 U/mL, between about 115 U/mL and about 125 U/mL,

between about 120 U/mL and about 130 U/mL, between about 125 U/mL and about 135 U/mL, between about 130 U/mL and about 140 U/mL, between about 135 U/mL and about 145 U/mL, between about 140 U/mL and about 150 U/mL, between about 145 U/mL and about 155 U/mL, between about 150 U/mL and about 160 U/mL, between about 155 U/mL and about 165 U/mL, between about 160 U/mL and about 170 U/mL, between about 165 U/mL and about 175 U/mL, between about 170 U/mL and about 180 U/mL, between about 175 U/mL and about 185 U/mL, between about 180 U/mL and about 190 U/mL, between about 185 U/mL and about 195 U/mL, or between about 190 U/mL and about 200 U/mL. In another embodiment, the useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein after a defined period of culture is between about 50 U/mL and about 60 U/mL, or about 57 U/mL. In another embodiment, the useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein is between about 90 U/mL and about 100 U/mL, or about 96 U/mL.

[0040] In other embodiments, the useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein after a defined period of culture is less than about 400 U/mL, less than about 375 U/mL, less than about 350 U/mL, less than about 325 U/mL, less than about 300 U/mL, less than about 275 U/mL, less than about 250 U/mL, less than about 225 U/mL, less than about 200 U/mL, less than about 175 U/mL, less than about 150 U/mL, less than about 125 U/mL, or less than about 100 U/mL.

[0041] In other embodiments, the useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein after a defined period of culture is more than about 50 U/mL, more than about 60 U/mL, more than about 70 U/mL, more than about 80 U/mL, more than about 90 U/mL, more than about 100 U/mL, more than about 110 U/mL, more than about 120 U/mL, more than about 130 U/mL, more than about 140 U/mL, more than about 150 U/mL, more than about 160 U/mL, more than about 170 U/mL, more than about 180 U/mL, more than about 190 U/mL, or more than about 200 U/mL.

[0042] For the purposes of the present disclosure, the levels of furin in culture supernatants disclosed herein are generated within a period of time from about 12 hours to about 96 hours after the initiation of the culture (after culture for about 12 hours to about 96 hours) and reflect the levels of furin which accumulate in the culture supernatant during that period. In other embodiments, the desired levels of furin in culture supernatants are reached within about 18 hours to about 90 hours, about 24 hours to about 84 hours, about 30 hours to about 78 hours, about 36 to about 72 hours, about 40 hours to about 80 hours, about 42 hours to about 68

hours, or about 48 hours to about 72 hours after the initiation of the culture, or after culture for the indicated period of time.

[0043] Alternatively, the levels of furin in culture supernatants disclosed herein are expressed as a concentration of furin secreted by a quantity of cells per volume of culture supernatant per day. In a non-limiting example, the concentration of furin is expressed as U/10⁶ cells/day. In other embodiment, a useful concentration of furin in the culture supernatant for the production of a mature, fully-processed mammalian protein is between about 20 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 25 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 30 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 35 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 40 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 45 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 50 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 55 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 60 U/10⁶ cells/day and about 75 U/10⁶ cells/day, between about 20 U/10⁶ cells/day and about 70 U/10⁶ cells/day, between about 20 U/10⁶ cells/day and about 65 U/10⁶ cells/day, between about 20 U/10⁶ cells/day and about 60 U/10⁶ cells/day, between about 20 U/10⁶ cells/day and about 55 U/10⁶ cells/day, between about 25 U/10⁶ cells/day and about 55 U/10⁶ cells/day, between about 25 U/10⁶ cells/day and about 50 U/10⁶ cells/day, between about 25 U/10⁶ cells/day and about 45 U/10⁶ cells/day, or between about 25 U/10⁶ cells/day and about 40 U/10⁶ cells/day.

[0044] The concentration of furin in the culture supernatant is sufficient to process at least about 75% of the mammalian precursor protein to a mature, functional protein. The protein is any protein translated as a precursor protein and processed into a mature form, at least in part, by the actions of furin. In one embodiment, the protein is FX. In other embodiments, the furin concentration is sufficient to process at least about 80% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 82% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 84% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 86% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 88% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 90% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 92% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 93% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 94% of the mammalian FX precursor

protein to a mature, functional FX protein, to process at least about 95% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 96% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 97% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 98% of the mammalian FX precursor protein to a mature, functional FX protein, to process at least about 99% of the mammalian FX precursor protein to a mature, functional FX protein, or to process 100% of the mammalian FX precursor protein to a mature, functional FX protein.

[0045] As used herein, the term “precursor protein” refers to a precursor protein that is inactive and is turned into an active form by cleavage and, optionally, other post-translational modifications in the cell after synthesis.

[0046] Thus, provided herein are transformed cells adapted for secretion of both furin and a mammalian protein, such as FX. The transformed cells can be any eukaryotic cell suitable for secretion of mammalian proteins, regardless of whether the cells produce endogenous furin. Suitable cell lines for generation of the transformed cells include, but are not limited to, Chinese hamster ovary (CHO) cells, human embryonic kidney cells, primate kidney cells (e.g., COS cells, HEK293), fibroblasts (e.g., murine fibroblasts), and mouse myeloma cells (e.g., NSO-GS). Suitable cell lines are capable of high level expression of mammalian proteins and are capable of post-translational modifications, e.g., glycosylation, formation of disulfide bonds, phosphorylation, and γ -carboxylation. Methods for selecting and culturing host cells and for inducing the host cells to express a polypeptide are generally known to the person skilled in the art.

[0047] Also disclosed herein are expression systems comprising cells suitable for production of mammalian proteins and at least one expression vector adapted for expression of at least one mammalian protein. Eukaryotic expression vectors are generally available for expression in mammalian cells. In order to enable furin and a mammalian protein, such as FX, to be expressed according to the methods disclosed herein, nucleotide sequences encoding the proteins are introduced into a eukaryotic cell by means of transfection, transformation or infection with an expression vector, whereby the polypeptides are expressed. The expression of the furin and/or mammalian proteins can be either transient or stable. The furin and mammalian nucleotide sequences are present as a plasmid, or as a part of a viral or non-viral expression vector. Particularly suitable viral vectors include, but are not limited to, baculoviruses, vaccinia viruses, adenoviruses, cytomegaloviruses, adeno-associated viruses, replication-competent

lentiviruses (RCL), and herpes viruses. Non-limiting examples of viral eukaryotic expression vectors include Rc/CMV, Rc/RSV, RCL, and SV40 vectors. Exemplary non-viral eukaryotic expression vectors include, but are not limited to, virosomes, liposomes, cationic lipids, plasmids, and polylysine-conjugated DNA. Exemplary plasmid expression vectors include, but are not limited to, pSLX, pcDNA, and others known to persons of ordinary skill in the art.

[0048] In another embodiment, disclosed herein is an expression vector comprising a furin-encoding nucleotide sequence, a mammalian protein-encoding nucleotide sequence, such as an FX-encoding nucleotide sequence, or a combination thereof. In one embodiment, both the furin and the protein sequences are expressed from a single expression vector. In another embodiment, the furin sequence and the protein sequences are expressed from different expression vectors. In one embodiment, if the furin and protein nucleotide sequences are expressed from the same expression vector, they are optionally separated by an internal ribosome entry site (IRES). The genes can be expressed from one or more promoters. Furthermore, nucleotide sequences encoding for each protein can be oriented in opposite directions on the plasmid or oriented in the same direction. The expression vectors further comprise selectable elements and other regulatory sequences for effective production of mammalian proteins as is understood by persons of ordinary skill in the art.

[0049] Also disclosed herein are expression vectors that allow expression of furin and other mammalian proteins by use of recombinase-mediated cassette exchange.

[0050] If furin and the mammalian protein are expressed from different expression vectors, then the expression vectors will have different selection markers so that cells transformed with the vector can be selected. Such selected cells may then be isolated and grown into monoclonal cultures

[0051] Promoters which permit constitutive, regulatable, tissue-specific, cell type-specific, cell cycle-specific, or metabolism-specific expression in eukaryotic cells are suitable, for example, for expression in mammalian cells. Regulatable elements are promoters, activator sequences, enhancers, silencers and/or repressor sequences. Examples of regulatable elements which permit constitutive expression in eukaryotes are promoters which are recognized by RNA polymerase III or viral promoters, cytomegalovirus (CMV) enhancer, CMV promoter, SV40 promoter or long terminal repeat (LTR) promoters, e.g. derived from MMTV (mouse mammary tumor virus) and other viral promoter and activator sequences which are derived from, for example, hepatitis B virus (HBV), hepatitis C virus (HCV), herpes simplex virus (HSV), human papilloma virus (HPV), Epstein-Barr virus (EBV), heat shock promoters, or human

immunodeficiency virus (HIV). Examples of regulatable elements which permit inducible expression in eukaryotes are the tetracycline operator in combination with an appropriate repressor. The expression of furin and mammalian protein nucleotide sequences can also take place under the control of tissue-specific, or protein-specific, promoters. Non-limiting examples of protein-specific promoters are FX gene promoters or furin gene promoters.

[0052] In certain embodiments, the cells are transformed with another protein in addition to furin and the mammalian protein. In one embodiment, the additional protein is vitamin K epoxide reductase (VKOR). In certain embodiments, the additional protein is expressed from the same expression vector as one, or both, of furin and the mammalian protein, or the additional protein is expressed from a different expression vector.

[0053] Also disclosed herein are expression systems comprising host cells and one or more expression vectors adapted to express furin and at least one additional mammalian protein, e.g. FX.

[0054] Also disclosed herein are methods of producing fully processed recombinant furin-requiring mammalian proteins, such as FX. In one embodiment, a stable, recombinant furin-producing cell line is produced and subsequently transfected with an expression vector containing the nucleotide sequence for at least one furin-requiring mammalian protein. Stable, recombinant furin-producing cell lines can be established and stored for transfection with an expression vector containing the nucleotide sequence for at least one furin-requiring mammalian protein as needed. Alternatively, expression vectors for furin and for the furin-requiring mammalian protein, such as FX, can be transfected into the host cells within about 30 minutes, about 60 minutes, about 2 hours, about 6 hours, about 12 hours, or about 24 hours of each other. In another embodiment, two or more expression vectors are transfected into the host cells substantially simultaneously. For the purposes of the present disclosure, substantially simultaneously refers to any time period that is less than or equal to 1 hour.

[0055] Transformed cells are selected according to the selection markers present in the expression vector(s) to produce stable pools of transformed cells and then the pools are optionally cloned to yield stable clones. The stable clones produce between about 50 U/mL and about 300 U/mL, between about 50 U/mL and about 400 U/mL, between about 50 U/mL and about 350 U/mL, between about 50 U/mL and about 300 U/mL, between about 50 U/mL and about 250 U/mL, between about 50 U/mL and about 200 U/mL, between about 50 U/mL and about 175 U/mL, between about 50 U/mL and about 150 U/mL, between about 50 U/mL and about 125 U/mL, or between about 50 U/mL and about 100 U/mL of furin in the culture

supernatant after about 36 to about 78 hours, about 36 to about 72 hours, about 40 hours to about 78 hours, about 42 hours to about 68 hours, or about 48 hours to about 72 hours after the initiation of the culture, or after culture for the indicated period of time. Furthermore, the stable clones yield more than 80% fully-processed and active recombinant mammalian protein, such as FX, of all the recombinant protein, such as FX, produced by the transformed cells.

[0056] Also disclosed herein is an expression system for recombinant furin and recombinant FX secreting furin into the culture supernatant at an accumulated concentration of between about 50 U/mL and about 300 U/mL after about 36 to about 78 hours of culture.

[0057] Also disclosed herein is a method of producing mature, fully-processed FX comprising use of an expression system secreting furin into the culture supernatant at an accumulated concentration between about 50 U/mL and about 300 U/mL after about 36 to about 78 hours of culture.

[0058] Also encompassed herein are recombinant mammalian proteins produced by the claimed methods and any full-processed mammalian recombinant FX.

EXAMPLES

Example 1. Production of fully processed and fully active recombinant Factor X by defined levels of furin

[0059] For FX expression, the mammalian expression plasmid pSLX containing either human codon-optimized FX or both, human codon-optimized FX and human codon-optimized vitamin K epoxide reductase (FX/VKOR), separated by an internal ribosome entry site (IRES), was used. Constructs for Chinese hamster ovary (CHO)-S and CHO-DG44 expression systems included geneticin selection and dihydrofolate reductase (dhfr) selection, respectively. For furin expression, the mammalian expression plasmid pcDNA3.1 containing human full length furin in combination with hygromycin as selection marker was used (FIG. 1A).

[0060] Initially, CHO-derived cell lines (CHO-S and CHO DG44) were transfected with the FX or FX/VKOR constructs to generate stable pools and subsequently the pools were subjected to subcloning to generate stable clones. In a second round of transfection and subcloning, a selected number of FX- or FX/VKOR-expressing clones were each super-transfected with furin resulting in stable pools and stable clones expressing FX/furin or FX/VKOR/furin.

[0061] Stable recombinant FX-producing CHO-S and CHO-DG44 cell lines were grown in animal component-free media, in shaker flasks for about 42 to about 72 hours and with starting

cell numbers of 0.3×10^6 or 0.5×10^6 cells/mL. CHO-S cells were maintained in PowerCHO®-CD media (Lonza BioWhittaker) supplemented with 4 mM glutamine, 500 µg/mL geneticin, 500 µg/mL hygromycin and 5 µg/mL vitamin K1. CHO-DG44 cells were maintained in OptiCHO™-CD media (Life Technologies) supplemented with 6 mM glutamine, 500 nM methotrexate (MTX) and 5 µg/mL vitamin K1.

[0062] The harvested cell culture supernatant was analyzed by Western blotting under reducing conditions to determine the quality of recombinant human FX using a polyclonal goat anti-human FX or polyclonal sheep anti-human FX (Affinity Biologicals). Densitometric analysis of the Western blots enabled the quantification of the different species of correctly processed FX, termed heavy chain FX (HC) and light chain FX (LC), and inadequately cleaved FX species, termed single chain FX (SC) and propeptide-containing light chain FX (PP-LC).

[0063] For FX quantification, the cell culture supernatant was analyzed with ELISA to determine the FX concentration and with the FXa chromogenic assay using Russell's viper venom (RVV) as activator to determine the concentration of active FX. These assays were calibrated using plasma-derived FX (Hyphen Biomed). The specific activity is given in %, by dividing the concentration of active FX by the concentration of total FX multiplied by 100. For furin quantification, active furin was determined in a furin fluorogenic assay calibrated against a furin reference material (New England Biolabs).

[0064] For statistical analysis, fully processed FX/total FX (%) was modeled as a function of furin concentration using the $E_{\max}$ model on CHO-DG44 transfection pools (A), CHO-S transfection pools (B), and CHO-S single cell-derived clones (C). This model is used for statistical evaluation of dose-response studies. The $E_{\max}$ model uses four parameters (E_0 , $E_{\max}$, ED_{50} , and n) to model FX as a function of furin as follows:

$$y = E_0 + (x^n \cdot E_{\max}) / (ED_{50}^n + x^n)$$

where y refers to the fully-processed FX/total FX and x refers to the furin concentration. The parameter E_0 refers to the basal effect corresponding to the response when the furin concentration is zero, $E_{\max}$ to the maximum effect attributable to the furin concentration, ED_{50} to the furin concentration which produces half of $E_{\max}$, and the parameter n represents the slope (Hill factor) determining the steepness of the curve.

[0065] To account that fully-processed FX/total FX approaches 100% if furin approaches an infinite large concentration, the $E_{\max}$ model was modified to a function with three parameters to be estimated as follows;

$$y = E_0 + (x^n \cdot (100 - E_0) / (ED_{50}^n = x^n))$$

[0066] This model was fitted to the data taking the variability among the three different cell lines into account using a non-linear mixed effects model by allowing the parameters E_0 and n to vary between the different cell lines by also modeling these two parameters as random effects.

[0067] Model diagnostics was done to validate the applied model. A comparison of the fitted $E_{\max}$ model with the null model using the likelihood ratio test was performed to determine statistical evidence for the $E_{\max}$ model estimating the percentage of fully processed FX of total FX depending on the furin concentration.

[0068] Results

[0069] The CHO-based heterologous expression system for human FX, comprising CHO-DG44 transfection pools (A), CHO-S transfection pools (B), and CHO-S single cell-derived clones (C), was used as a basis to study the effect of furin expression on human FX processing following different transfection strategies. Transfection pools, as well as clones, additionally expressed VKOR which had no impact on the study.

[0070] After an incubation period of two to three culture days, the cell culture supernatant was subjected to a series of analyses, including Western blot analysis under reducing conditions, furin activity assay, ELISA and RVV assay (Table 1). On average, FX-producing CHO pools and clones revealed a FX specific activity of over 50%, partly reaching 100% (Table 1). Western blot analyses showed that the recombinant FX was inadequately processed to different degrees, as shown by two incompletely processed forms of FX (i.e. the propeptide-containing FX light chain and the FX single chain) besides the fully processed, propeptide-free FX light chain and FX heavy chain (FIG. 4). By means of densitometric analysis of these four species of FX, the percentage of fully processed FX, i.e. FX light chain plus FX heavy chain in relation to total FX, ranged between 30% and almost 100% in cell culture supernatants (Table 1, FIG. 4). Furthermore, no pre-activation was observed, which would be visible as heavy chain band shortened by the size of the missing activation peptide. Also evaluated was whether the concentration of secreted furin had an influence on the degree of processed FX by plotting these two parameters (FIG. 5). As shown in FIG. 5, only partial processing of FX is feasible with low concentrations of secreted furin (< 20 U/mL), whereas higher levels of secreted furin correlate with better processed FX.

Table 1: Data summary of FX and furin co-expressing cell lines: shown are FX productivities, furin productivities and titers, and percentages of fully processed FX/total FX measured for each cell line.

Clone / Pool ID	Expressed Recombinant Proteins	Cell line (pool or clone)	Furin conc. (U/mL)	Final cell density [10 ⁶ cells/mL]	Furin Specific Productivity [U/10 ⁶ cells/day]	FX specific activity [%]	Fully processed FX/total FX (%)
1	FX	CHO-DG44 pool	8.03	2.015	2.22	5.64	30.34
2	FX/Furin	CHO-DG44 pool	8.05	1.155	3.45	7.58	45.69
3	FX/VKOR	CHO-DG44 pool	7.88	1.436	2.86	6.44	31.05
4	FX/VKOR/Furin	CHO-DG44 pool	7.88	1.029	3.68	11.16	40.08
5	FX	CHO-DG44 pool	2.74	1.518	0.95	9.04	41.42
6	FX/Furin	CHO-DG44 pool	11.59	0.952	5.71	14.12	71.78
7	FX/VKOR	CHO-DG44 pool	2.38	1.152	1.02	15.14	36.92
8	FX/ VKOR /Furin	CHO-DG44 pool	2.82	0.795	1.58	29.37	57.17
9	FX	CHO-DG44 pool	5.36	2.438	1.26	9.23	34.46
10	FX/Furin	CHO-DG44 pool	9.94	1.503	3.48	12.23	59.30
11	FX/VKOR	CHO-DG44 pool	4.88	2.162	1.27	14.57	33.12
12	FX/ VKOR /Furin	CHO-DG44 pool	4.78	1.372	1.80	20.48	46.98
13	FX	CHO-DG44 pool	8.07	2.005	2.24	47.68	40.93
14	FX/Furin	CHO-DG44 pool	8.83	1.851	2.62	64.80	60.54
15	FX/VKOR	CHO-DG44 pool	8.42	1.843	2.50	45.48	33.98
16	FX/ VKOR /Furin	CHO-DG44 pool	6.72	2.220	1.71	58.04	47.46
17	FX	CHO-DG44 pool	2.59	1.246	1.05	78.72	46.40
18	FX/Furin	CHO-DG44 pool	55.91	1.396	20.75	98.49	73.47
19	FX/VKOR	CHO-DG44 pool	2.59	1.207	1.07	70.43	47.61

Clone / Pool ID	Expressed Recombinant Proteins	Cell line (pool or clone)	Furin conc. (U/mL)	Final cell density [10 ⁶ cells/mL]	Furin Specific Productivity [U/10 ⁶ cells/day]	FX specific activity [%]	Fully processed FX/total FX (%)
20	FX/ VKOR /Furin	CHO-DG44 pool	3.48	1.431	1.27	93.42	60.04
21	FX	CHO-DG44 pool	5.96	2.035	1.63	71.19	37.77
22	FX/Furin	CHO-DG44 pool	27.57	2.226	7.00	104.40	71.85
23	FX/VKOR	CHO-DG44 pool	4.40	1.827	1.32	69.50	42.84
24	FX/ VKOR /Furin	CHO-DG44 pool	5.20	2.283	1.29	75.60	49.19
25	FX/VKOR	CHO-S pool	0.00	2.224	0.00	90.59	63.96
26	FX/VKOR/Furin	CHO-S pool	51.34	1.828	25.20	99.57	88.97
27	FX/VKOR	CHO-S pool	0.00	2.851	0.00	49.99	57.75
28	FX/VKOR/Furin	CHO-S pool	62.84	2.248	26.13	48.73	87.78
29	FX/VKOR	CHO-S pool	0.00	3.584	0.00	81.39	68.02
30	FX/VKOR/Furin	CHO-S pool	66.98	2.362	26.75	66.57	89.64
31	FX/VKOR	CHO-S pool	0.00	2.870	0.00	69.04	67.02
32	FX/VKOR/Furin	CHO-S pool	55.08	2.295	22.52	52.91	89.99
33	FX/VKOR	CHO-S pool	0.00	2.645	0.00	59.62	71.13
34	FX/Furin	CHO-S pool	116.89	2.083	50.52	51.40	86.56
35	FX	CHO-S pool	0.00	2.230	0.00	67.49	62.65
36	FX/Furin	CHO-S pool	46.62	1.846	22.18	19.29	82.99
37	FX	CHO-S pool	0.00	2.049	0.00	11.47	51.69
38	FX/Furin	CHO-S pool	48.41	1.457	27.61	<25.33	77.46
39	FX	CHO-S pool	0.00	2.265	0.00	95.04	45.87
40	FX/Furin	CHO-S pool	60.72	1.566	32.80	27.69	84.27

Clone / Pool ID	Expressed Recombinant Proteins	Cell line (pool or clone)	Furin conc. (U/mL)	Final cell density [10 ⁶ cells/mL]	Furin Specific Productivity [U/10 ⁶ cells/day]	FX specific activity [%]	Fully processed FX/total FX (%)
41	FX	CHO-S pool	0.00	2.018	0.00	35.63	49.28
42	FX/VKOR/Furin	CHO-S clone	113.45	3.704	26.99	90.30	95.91
43	FX/VKOR/Furin	CHO-S clone	63.41	3.133	17.45	92.26	96.23
44	FX/VKOR/Furin	CHO-S clone	92.86	2.893	27.37	98.55	98.12
45	FX/VKOR/Furin	CHO-S clone	173.37	3.430	44.11	79.46	98.42
46	FX/VKOR/Furin	CHO-S clone	23.54	2.816	7.10	86.31	91.69
47	FX/VKOR/Furin	CHO-S clone	105.42	2.918	30.84	78.63	95.14
48	FX/VKOR/Furin	CHO-S clone	33.77	3.542	8.35	80.02	92.21
49	FX/VKOR/Furin	CHO-S clone	135.62	2.777	41.39	94.46	95.18
50	FX/VKOR/Furin	CHO-S clone	82.03	2.193	30.46	48.80	93.75
51	FX/VKOR/Furin	CHO-S clone	8.06	2.280	2.90	67.11	52.38
52	FX/VKOR/Furin	CHO-S clone	81.02	2.889	23.91	85.37	95.61

[0071] To understand the influence of furin on FX processing and to provide statistical support of the data, fully processed FX of total FX (%) was modeled as a function of furin concentration using the $E_{\max}$ model on cell lines A, B and C (FIG. 6). Four model diagnostics plots, indicating a good fit of the model to the data, are provided in FIGs. 7A-D. A comparison of the fitted $E_{\max}$ model with the null model using the likelihood ratio test resulted in a p-value <0.0001 , providing statistical evidence for a higher percentage of fully processed FX of total FX depending on a higher furin concentration (FIG. 8), once again proving validity of the data. Based on the statistical analysis, the estimated furin concentrations to be produced by the production cell line as detected in the cell culture medium together with FX resulting in equal or higher 90% and equal or higher 95% fully processed FX of total FX were at least 57 U/mL and at least 96 U/mL, respectively (FIG. 9).

[0072] In summary, the data provides a defined minimal level of secreted furin (at least 57 U/mL and at least 96 U/mL) in the cell culture supernatant that is required for sufficient FX processing (equal or higher 90% and equal or higher 95%).

[0073] In biotechnological processes that express high levels of recombinant protein, furin overexpression may be used to obtain fully processed zymogens. With our invention, we provide for the first time a defined minimum of secreted furin warranting for high FX processing (≥ 57 U/mL to achieve at least 90% fully processed FX and ≥ 96 U/mL furin for at least 95% fully processed FX). This finding is particularly beneficial to fermentation processes expressing recombinant FX, FXa and variants from human and animal species, where the furin level may be used as an indicator for adequate processing of the FX precursor protein, and as target for cell line and process development.

[074] Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." As used herein the terms "about" and "approximately" means within 10 to 15%, preferably within 5 to 10%. Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain

errors necessarily resulting from the standard deviation found in their respective testing measurements.

[075] The terms “a,” “an,” “the” and similar referents used in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. Recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the invention.

[076] Groupings of alternative elements or embodiments of the invention disclosed herein are not to be construed as limitations. Each group member may be referred to and claimed individually or in any combination with other members of the group or other elements found herein. It is anticipated that one or more members of a group may be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

[077] Certain embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Of course, variations on these described embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventor expects skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

[078] Specific embodiments disclosed herein may be further limited in the claims using consisting of or consisting essentially of language. When used in the claims, whether as filed or added per amendment, the transition term “consisting of” excludes any element, step, or ingredient not specified in the claims. The transition term “consisting essentially of”

limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s). Embodiments of the invention so claimed are inherently or expressly described and enabled herein.

[079] Furthermore, numerous references have been made to patents and printed publications throughout this specification. Each of the above-cited references and printed publications are individually incorporated herein by reference in their entirety.

[080] In closing, it is to be understood that the embodiments of the invention disclosed herein are illustrative of the principles of the present invention. Other modifications that may be employed are within the scope of the invention. Thus, by way of example, but not of limitation, alternative configurations of the present invention may be utilized in accordance with the teachings herein. Accordingly, the present invention is not limited to that precisely as shown and described.

What is claimed is:

1. A transformed cell comprising:
a nucleotide sequence encoding a human furin, such that the transformed cell expresses and secretes functional furin into a culture supernatant, wherein the furin is secreted at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for about 36 to about 78 hours.
2. The transformed cell of claim 1, further comprising a nucleotide sequence encoding a protein cleavable by furin and exhibiting an Arg-(Lys/Arg)-Arg motif.
3. The transformed cell of claim 2, wherein the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on different expression vectors.
4. The transformed cell of claim 2, wherein the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on the same expression vector.
5. The transformed cell of claim 2, wherein the protein is von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z.
6. The transformed cell of claim 5, wherein the protein is Factor X.
7. The transformed cell of claim 6, wherein the functional furin is secreted at a concentration of at least about 50 to about 60 U/mL in the culture supernatant.
8. The transformed cell of claim 7, wherein at least 90% of the Factor X produced by the transformed cell is fully processed.
9. The transformed cell of claim 6, wherein the functional furin is secreted at a concentration of at least about 90 to about 100 U/mL in the culture supernatant.
10. The transformed cell of claim 9, wherein at least 95% of the Factor X produced by the transformed cell is fully processed.
11. A transformed cell comprising:
a first nucleotide sequence encoding a human furin and a second nucleotide sequence encoding a human Factor X, such that the transformed cell expresses and

secretes functional furin and Factor X into a culture supernatant, wherein the furin is secreted at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for about 36 to about 78 hours and at least 85% of the Factor X is fully processed.

12. The transformed cell of claim 11, wherein the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on different expression vectors.

13. The transformed cell of claim 11, wherein the nucleotide sequence encoding human furin and the nucleotide sequence encoding the protein are on the same expression vector.

14. The transformed cell of claim 11, wherein the functional furin is secreted at a concentration of at least about 50 to about 60 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

15. The transformed cell of claim 14, wherein at least 90% of the Factor X produced by the transformed cell is fully processed.

16. The transformed cell of claim 11, wherein the functional furin is secreted at a concentration of at least about 90 to about 100 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

17. The transformed cell of claim 16, wherein at least 95% of the Factor X produced by the transformed cell is fully processed.

18. The transformed cell of either of claims 1 or 11, wherein the furin concentration in the culture supernatant reaches about 50 to about 300 U/mL after culture for about 42 to about 72 hours.

19. A eukaryotic protein expression system comprising:
a cell line suitable for expression of mammalian proteins;
a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and
a second expression vector adapted for expression of a protein by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a protein cleavable by furin and exhibiting an Arg-(Lys/Arg)-Arg motif,

wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for about 36 to about 78 hours.

20. The eukaryotic protein expression system of claim 19, wherein the protein is von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z.

21. The eukaryotic protein expression system of claim 20, wherein the protein is Factor X.

22. The eukaryotic protein expression system of claim 21, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of at least about 50 to about 60 U/mL after culture for about 36 to about 78 hours.

23. The eukaryotic protein expression system of claim 22, wherein at least 90% of the Factor X produced by the expression system is fully processed.

24. The eukaryotic protein expression system of claim 21, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of at least about 90 to about 100 U/mL after culture for about 36 to about 78 hours.

25. The eukaryotic protein expression system of claim 24, wherein at least 95% of the Factor X produced by the expression system is fully processed.

26. A eukaryotic protein expression system comprising:
a cell line suitable for expression of mammalian proteins;
a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and
a second expression vector adapted for expression of Factor X by the cell line, wherein the second expression vector includes a nucleotide sequence encoding Factor X,
wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for about 36 to about 78 hours.

27. The eukaryotic protein expression system of claim 26, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of at least about 50 to about 60 U/mL after culture for about 36 to about 78 hours.
28. The eukaryotic protein expression system of claim 27 wherein at least 90% of the Factor X produced by the expression system is fully processed.
29. The eukaryotic protein expression system of claim 26, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of at least about 90 to about 100 U/mL after culture for about 36 to about 78 hours.
30. The eukaryotic protein expression system of claim 29, wherein at least 95% of the Factor X produced by the expression system is fully processed.
31. A eukaryotic protein expression system comprising:
a cell line suitable for expression of mammalian proteins;
a first expression vector adapted for expression of human furin and a protein cleavable by furin, and exhibiting an Arg-(Lys/Arg)-Arg motif, by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide and a nucleotide sequence encoding the protein cleavable by furin, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for about 36 to about 78 hours.
32. The eukaryotic protein expression system of claim 31, wherein the protein is von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z.
33. The eukaryotic protein expression system of claim 32, wherein the protein is Factor X.
34. The eukaryotic protein expression system of claim 33, wherein the cell line is capable of secreting functional furin into a culture supernatant at a concentration of at least about 50 to about 60 U/mL after culture for about 36 to about 78 hours.
35. The eukaryotic protein expression system of claim 34, wherein at least 90% of the Factor X produced by the expression system is fully processed.

36. The eukaryotic protein expression system of claim 33, wherein the cell line is capable of secreting functional furin into a culture supernatant at a concentration of at least about 90 to about 100 U/mL after culture for about 36 to about 78 hours.

37. The eukaryotic protein expression system of claim 36, wherein at least 95% of the Factor X produced by the expression system is fully processed.

38. A eukaryotic protein expression system comprising:
a cell line suitable for expression of mammalian proteins;
a first expression vector adapted for expression of human furin and Factor X, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide and a nucleotide sequence encoding Factor X, wherein the cell line is capable of secreting functional furin into the culture supernatant at a concentration of about 50 U/mL to about 300 U/mL after culture for about 36 to about 78 hours.

39. The eukaryotic protein expression system of claim 38, wherein the cell line is capable of secreting functional furin into a culture supernatant at a concentration of at least about 50 to about 60 U/mL after culture for about 36 to about 78 hours.

40. The eukaryotic protein expression system of claim 39, wherein at least 90% of the Factor X produced by the expression system is fully processed.

41. The eukaryotic protein expression system of claim 38, wherein the cell line is capable of secreting functional furin into a culture supernatant at a concentration of at least about 90 to about 100 U/mL after culture for about 36 to about 78 hours.

42. The eukaryotic protein expression system of claim 41, wherein at least 95% of the Factor X produced by the expression system is fully processed.

43. The eukaryotic protein expression system of any of claims 19, 26, 31 or 38, wherein the furin concentration in the culture supernatant reaches about 50 to about 300 U/mL after culture for about 42 to about 72 hours.

44. A method of preparing a recombinant protein comprising:
transfecting a cell line suitable for expression of mammalian proteins with a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and

transfecting the cells line with a second expression vector adapted for expression of a protein by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a protein exhibiting an Arg-(Lys/Arg)-Arg motif;

wherein the cell line transfected with the first and the second expression vectors expresses and secretes a functional human furin at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

45. The method of claim 44, wherein the protein is von Willebrand Factor, Factor II, Factor IX, Factor X, Protein C, Protein S, or Protein Z.

46. The method of claim 45, wherein the protein is Factor X.

47. The method of claim 45, wherein the functional furin is secreted by the cell line at a concentration of at least about 50 to about 60 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

48. The method of claim 47, wherein at least 90% of the Factor X produced by the cell line is fully processed.

49. The method of claim 45, wherein the functional furin is secreted by the cell line at a concentration of at least about 90 to about 100 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

50. The method of claim 49, wherein at least 95% of the Factor X produced by the cell line is fully processed.

51. The method of claim 44, wherein the cell line is transfected with the first expression vector and the second expression vector substantially simultaneously.

52. The method of claim 44, wherein the cell line is transfected with the first expression vector and cells secreting stable levels of furin are obtained prior to transfecting the cell line with the second expression vector.

53. The method of claim 44, wherein the cell line is transfected with the second expression vector and cells secreting stable levels of the protein are obtained prior to transfecting the cell line with the first expression vector.

54. A method of preparing a recombinant protein comprising:

transfecting a cell line suitable for expression of mammalian proteins with a first expression vector adapted for expression of human furin by the cell line, wherein the first expression vector includes a nucleotide sequence encoding a human furin polypeptide; and

transfecting the cells line with a second expression vector adapted for expression of Factor X by the cell line, wherein the second expression vector includes a nucleotide sequence encoding a Factor X polypeptide;

wherein the cell line transfected with the first and the second expression vectors expresses and secretes a functional human furin at a concentration of about 50 U/mL to about 300 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

55. The method of claim 54, wherein the functional furin is secreted by the cell line at a concentration of at least about 50 to about 60 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

56. The method of claim 55, wherein at least 90% of the Factor X produced by the cell line is fully processed.

57. The method of claim 54, wherein the functional furin is secreted by the cell line at a concentration of at least about 90 to about 100 U/mL in the culture supernatant after culture for about 36 to about 78 hours.

58. The method of claim 57, wherein at least 95% of the Factor X produced by the cell line is fully processed.

59. The method of claim 55, wherein the cell line is transfected with the first expression vector and the second expression vector substantially simultaneously.

60. The method of claim 55, wherein the cell line is transfected with the first expression vector and cells secreting stable levels of furin are obtained prior to transfecting the cell line with the second expression vector.

61. The method of claim 55, wherein the cell line is transfected with the second expression vector and cells secreting stable levels of the protein are obtained prior to transfecting the cell line with the first expression vector.

62. The method of either of claims 44 or 54, wherein the furin concentration in the culture supernatant reaches about 50 to about 300 U/mL after culture for about 42 to about 72 hours.
60. A fully-processed recombinant Factor X produced by the transformed cell of any one of claims 1-18.
61. A fully-processed recombinant Factor X produced by the expression system of any one of claims 19-43.
62. A fully-processed recombinant Factor X produced by the method of any one of claims 44-62.
63. An expression system for recombinant FX adapted to secrete furin into a culture supernatant at a concentration of between about 50 U/mL and about 300 U/mL after culture for about 36 to about 78 hours.
64. A method of producing mature, fully-processed FX comprising an expression system secreting furin into a culture supernatant at a concentration between about 50 U/mL and about 300 U/mL after culture for about 36 to about 78 hours.
65. A method of treatment of a coagulation disorder comprising:
administering to a subject in need thereof a fully-processed recombinant Factor X of any of claims 60-62, wherein as a result of the administration, the coagulation disorder is treated.
66. The method according to claim 65, wherein the coagulation disorder is hemophilia or von Willebrand disease.




FIG. 1A

GACGGATCGGGAGATCTCCCGATCCCCCTATGGTCGACTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCAGTATCTGCTCCCT
GCTTGCTGTTGGAGGTCGCTGAGTAGTGC CGAGCAAAATTTAAGCTACAACAAGGCAAGGCTTGACCGACAATTCATGAAGAATCT
GCTTAGGTTAGGCGTTTGGCGTGTGCGATGTACGGCCAGATATACGCGTTGACATTGATTTATGACTAGTTATTAATAGTAAT
CAATTACGGGTCAATTAGTTTCATAGCCCATATATGAGTTCCGCGTTACATAAATTACGGTAAATGGCCCGCTGCTGACCGCCCAAC
GACCCCGCCCATTGACGTC AATAATGACGTATGTTCCCATAGTAACGCCAATAGGACTTTCATTGACGTC AATGGTGACTATTT
ACGGTAAACTGCCACTTGGCAGTACATCAAGTGTAATGCAAGTACGCCCTATTGACGTCAATGACGGTAAATGGCCCGCT
GGCATTATGCCCAGTACATGACCTTATGGGACTTTTCCTACTTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCCG
TTTTGGCAGTACATCAATGGCGGTGATAGCGGTTTGACTCACGGGGATTTCCAAAGTCTCCAACTTGACGTC AATGGAGTTTGT
TTGGCACCAAAATCAACGGGACTTTC AAAATGTCGTAACAACCTCCGCCATTGACGCAAAATGGCGGTAGCGGTACGGTGGAGG
TCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAAACCACTGCTTACTGCTTATCGAAATTAATACGACTCATATAGGAGACCCAA
GCTGGCTAGCGTTTAAACTTAAAGCTTGGTACCGAGCTCGGATCCACTAGTCCAGTGTGGTGAATTTCTGCAGATATCCAGCACAGTGGC
GGCCGCTAGGAGCTGAGGCCCTGGTTGCTATGGGTAGCAGCAACAGGAACCTTGGTCTCTAGCAGCTGATGCTCAGGGCCAGAA
GGTCTTCACCAACACGTGGGCTGTGCGCATCCCTGGAGGCCAGCGGTGGCAACAGTGTGGACGGAAAGCATGGGTTCTCAACCTGG
GCCAGATCTTTCGGGACTATTACCACTTCTGGCATCGAGGAGTGACGAAGCGTCCCTGTGCGCTCACCGCCCGGCA CAGCCGGCTG
CAGAGGGAGCCTCAAGTACAGTGGCTGGAACAGCAGGTGGCAAGCGGACTAAACGGGACGTGTAC CAGGAGCCACAGACCCCAA
GTTTCTCAGCAGTGGTACCTGTCTGGTGTCACTCAGCGGACCTG AATGTGAAGCGGCTGGCGCAGGCTACACAGGGCACGGCA
TTGTGGTCTCCATTCTGGACGATGGCATCGAGAAAGAACCCCGACTTGGCAGGCAATTAATGATCCTGGGGCCAGTTTTGATGTCAAT
GACCAGGACCCCTGACCCCGAGCCTCGGTACACACAGATGAATGACAACAGGCACCGGTGTGCGGGGAAGTGGCTGCGGTGGC
CAACAAACGGTGTCTGTGGTGTAGGTGTGGCTACAACGCCCGCATTTGGAGGGGTGCGCATGCTGGAATGGCGAGGTGACAGATGCAGTGG
AGGCACGCTCGCTGGGCTGAACCCCAACCATCCACATCTACAGTGCCAGCTGGGGCCCCGAGGATGACGGCAAGACAGTGGATGGG
CCAGCCCGCTCGCGAGGAGGCCCTTCTTCGGTGGGTAGCCAGGGCCGAGGGGCTGGGCTCCATCTTTGTCTGGCCCTCGGGGAA
CGGGGCCGGGAACATGACAGCTGCAACTGCGACGGCTACACCAACAGTATCTACACGCTGTCCATCAGCAGCGCCACG CAGTTTGGCA
ACGTGCCGTGGTACAGCGAGGCCCTGCTCGTCCACACTGGCCACGACCTACAGCAGTGGCAAC CAGAAATGAGAAGCAGATCGTGACGACT
GACTTGCGGCAGAACTGCACGGAGTCTCACACGGGCACCTCAGCCTCTGCCCTTAGCAGCCGGCATCATTTGCTCTCACCCTGGAGGC
CAATAAGAACCTCACATGGCGGACATGCAACACCTGGTGTACAGACCTCGAAGCCAGCCACCTCAATGCCAACGACTGGGCCACCA
ATGGTGTGGCCGGAAGTGAGCCACTCATATGGCTACGGGCTTTTGGACCGCAGCGCCATGTTGGCCCTGCCCCAGAAATTGGACCCACA
GTGGCCCCCAGCGGAAGTGCAATCATCGACATCCTCACCGAGCCCAAAGACATCGGGAAACGGCTCGAGGTCCGGAAGACCGTGACCGC
GTGCCCTGGCGAGCCCAACACATCACTCGGCTGGAGCACGCTCAGCGGCTCACCTGTCTATAATCGCCGTGGCGACCTGGCCA
TCCACCTGGTCAGCCCATGGGCACCCGCTCCACCTGCTGGCAGCCAGGCCACATGACTACTCCGCAGATGGGTTTAATGACTGGGCC
TTCATGACAACTCATTCCTGGGATGAGGATCCCTCTGGCGAGTGGTCCTAGAGATTGAAAACACCAGCGAAGCCAACTATGGGAC
GCTGACCAAGTTCACCTCGTACTATGGCACCGCCCTGAGGGCTGCCGTACCTCCAGAAAGCAGTGGCTGCAAGACCTCACGT

FIG. 1B

3/14

CCAGTCAGGCCTGTGTGTGCGAGGAAGCCTTCTCCCTGCCACAGAAAGAGCTGTGTCCAGCACCTGCCCTCCAGGCTTCCGCCCCCAA
GTCCCTCGATACGCACTATAGCACCGAGAAATGACGTGGAGACCATCCGGCCAGCGTCTGCGCCCTGCCACGCTCATGTGCCACATG
CCAGGGCCCGCCCTGACAGACTGCCCTCAGCTGCCCCAGCCACGCCCTCCTTGGACCCCTGTGGAGCAGACTTGTCTCCCGGCAAAAGCCAGA
GCAGCCGAGAGTCCCGCCACAGCAGCAGCCACCTCGGCTGCCCCCGAGGTGAGGCGGGCAACGGCTGCGGGCAGGGCTGCTGCC
TCACACCTGCTGAGGTGGTGGCCGGCTCAGCTGGCCCTTCATCGTGTGCTCTTCCCTGCTCCTGTCAGCTGCGCTC
TGGCTTTAGTTTTCCGGGGGTGAAGGTGTACACCAATGGACCGTGCCCTCACTCTCTACAAGGGCTGCCCTGAAAGCCTGGCAGGAGG
AGTGCCCGCTCTGACTCAGAAAGAGGACGAGGCGGGCGAGAGGACCGCTTTATCAAAAGACCAGAGCGCCCTCTGTATCTAGAGGGCCC
GTTTAAACCCGCTGATCAGCCTCGACTGTGCCCTTCTAGTTGCCAGCCATCTGTGTGTTTGGCCCTCCCGGTGCCCTTCTGTGACCTGGA
AGGTGCCACTCCCACTGTCTCTTCCCTAATAAATGAGGAAATGCAATCGCATTTGTCTGAGTAGGTGTCATTTCTATTTCTGCGGGGTGGG
TGGGCGAGGACAGCAAGGGGAGGATTTGGGAACAAATAGCAGGCATGCTGGGATGCCGTGGCTCTATGCTTCTGAGGCCGGAAGA
ACCAGCTGGGCTCTAGGGGTATCCCAAGCGGCTGTAGCGGCGCATTAAGCGGCGGTGTGGTTACGCCGAGCGTGACCGC
TACACTTGGCAGCGCCCTAGCGCCCGCTCTTTCGCTTCTTCCCTTCTCGCCACGTTCCCGGCTTTTCCCGTCAAGCTCTAA
ATCGGGGCATCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAACTTGATTAGGGTGATGGTTCAAGTAGTGGG
CCATCGCCCTGATAGACGGTTTTTTCGCCCTTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACCTGGAAACAACACT
CAACCTATCTCGGTCTATTCTTTTGATTTTATAAGGATTTTGGGATTTTCGGCTATTGGTTAAAAAATGAGCTGATTTAAACAAAAAT
TTAAACGCGAATTAAATCTGTGGAAATGTGTGTCAGTTAGGGTGTGAAAGTCCCGAGGCTCCCGAGGAGGCAAGATATGCAAAAGCATG
CATCTCAATTAGTCAGCAACCAGGTGTGAAAGTCCCGAGGCTCCCGAGCAGAGAGTATGCAAAAGCATGCACTCTCAATTAGTCAGC
AACCATAGTCCCGCCCTAACTCCGCCCATCCCGCCCTAACTCCGCCCATTTCCGCCCATTTCCGCCCATGGCTGACTAATTTT
TTATTTATGCAGAGCCGAGGCGCCCTCTGCTCTGAGCTATTCCAGAAAGTAGTGAGGAGGCTTTTGTGGAGGCTTAGGCTTTTGCAAA
AAGCTCCCGGAGCTTGTATATCCATTTTCGGATCTGATCAGCAGTGAATAAAGCTGAACCTCACCGGACGCTCTGTCGAGAAGTT
TCTGATCGAAAGCTTCGACAGCGCTCTCCGACCTGATGCAAGTCTCGTGCTTTCAGCTTTCGATGTAGGAGGCG
GTGGATATGTCCTGCGGGTAAATAGCTGCGCCGATGGTTCTACAAAGATCGTTATCGGCACCTTTGCATCGGCGCGCTCCCG
ATTCCGGGAAGTGCTTGACATTGGGGAATTGAGCGAGGCCCTGACCTATTGCTATCTCCCGCGTGCCAGGGTGTCAAGTTCAGACCT
GCCTGAAACCGAACTGCCCGCTGTCTGCGAGCCGGTCCGGAGGCCATGGATCGGTGCGGCCGATCTTAGCCAGACGAGCGGGT
TCGGCCCATTCGGACCGCAAGGAATCGGTCAATACACTACATGCGGTGATTTTCATATGCGCGATTGCTGATCCCCATGTGTATCACTGG
CAAACTGTGATGGACGACACCGTCAGTGCGTCCGTGCGCAGGCTCTCGATGAGCTGATGCTTTGGGCGGAGGACTGCCCGAAGTCCG

FIG. 1B (cont.)

4/14

FIG. 1B (cont.)

5/14

ATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAAGTAAGTTGGCCGCAGTGTTATCACTCATGGTTATGGC
AGCACTGCATAATTCTCTTACTGTCAATGCCATCCGTAAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAAATAGT
GTATGCGGCGACCGAGTTGCTCTTTGCCCGCGCTCAATACGGGATAAATACCGGCCACATAGCAGAACTTTTAAAGTGCTCATCATTTGGA
AAACGTTCTTCGGGCGGAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCAACCCAACTGATCTTC
AGCATCTTTTACTTTTCAACAGCGGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGGCGACACGGAAAT
GTTGAATACTCATACTCTCTCTCTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCATGAGCGGATACATAATTGGAATGTATT
TAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAGTGCCACCTGACGTC

FIG. 1B (cont.)


ATGGAGCTGAGGCCCTGGTTGCTATGGGTGGTAGCAGCAACAGGAACCTTGGTCCTGTCTAGCAGCTGATGCTCAGGGCCAGAAAGGTCTT
CACCAACACGTGGGCTGTGCGCATCCCTGGAGGCCCAGCGGTGCCAACAGTGTGGCAAGCAAGCATGGGTTCTCTCAACCTGGGCCAGA
TCTTCGGGGAATAATTACCACTTCTGGCATCGAGGAGTGACGAAGCGGTCCCTGTGCTCCCTCACCGCCCGGCACAGCCGGCTGCAGAGG
GAGCCTCAAGTACAGTGGCTGGAAACAGCAGGTGGCAAGCGACGGAATAACGGGACGTGTACAGGAGCCACAGACCCCAAAGTTTCC
TCAGCAGTGGTACCTGTCTGTGTCTCACTCAGCGGACCTGAATGTGAAGCGGCTGTGGCGCAGGGCTACACAGGCACGGCATTTGTGG
TCTCCATTTCTGGACGATGGCATCGAGAAAGAACCCGGACTTGGCAGGCAATTATGATCCTGGGCCAGTTTGTATGTC AATGACCAAG
GACCCGTGACCCCGAGCCTCGGTACACACAGATGAATGACAACAGGCAAGGATGTCGGGGAAGTGGCTGCGTGCCAAACAA
CGGTGTCGTGTGTAGGTGTGGCTTACAACGCCCATTTGGAGGGTGGCATGCTGGATGGCGAGGTGACAGATGCAGTGGAGGCAC
GCTCGCTGGGCTGAACCCCAACACATCCACATCTACAGTGCCAGCTGGGGCCCGAGGATGACGGCAAGACAGTGGATGGGCCAGCC
CGCCTCGCCGAGGAGCCCTTCTTCCTGGGCTTAGCCAGGGCCGAGGGGCTGGGCTCCATCTTTGTCTGGGCCCTCGGGGAAACGGGG
CCGGGAACATGACAGCTGCAACTGCCGACGGCTACACCAACAGTATCTACACGCTGTCCATCAGCAGCCACGCAGTTTGGCAACGTGC
CGTGTACACGCGAGGCCCTGCTCGTCCACACTGGCCACGACCTACAGCAGTGGCAACCCAGAAATGAGAAGCAGATCGTGACGACTGACTG
CGGCAGAAAGTGCACGGAGTCTCACACGGGCACCTCAGCCCTCTGCCCTTAGCAGCCGGCATATTGCTCTCACCCCTGGAGGCCAATAA
GAACCTCACATGGCGGACATGCAACACCTGGTGGTACAGACCTCGAAGCCAGCCCACTCAATGCCAACGACTGGGCCACCAATGGTG
TGGGCCGGAAGTGAGCCACTCATATGGCTACGGGCTTTTGGACGCAAGGCCATGGTGCCCTGGCCAGAAATTGGACCAACAGTGGCC
CCCCAGCGGAAGTGCAATCATCGACATCCTCACCGAGCCCAAGACATCGGGAACGGCTCGAGGTGGGAAGACCGTGACCGCGTGCCCT
GGCGAGCCCAACACATCCTCGGCTGGAGCACGCTCAGGCGCGGCTCACCTGTCTCTATAATCGCCGTGGCGACCTGGCCATCCACC
TGGTGAGCCCCATGGGCACCCGCTCCACCTGCTGGCAGCCAGGCCACATGACTACTCCGCAGATGGGTTTAATGACTGGGCTTCATG
ACAACTCATTCCTGGGATGAGGATCCCTCTGGCGAGTGGTCTTAGAATAACACAGCGAAGCCAAACAATACTATGGGACGCTGAC
CAAGTTCACCTCGTACTCTATGGCACCGCCCTGAGGGGCTGCCGTACCTCAGAAAGCAGTGGCTGCAAGACCTTCACGTCCAGTC
AGGCCCTGTGTGTGCGAGGAAGCTTCTCCCTGCACCAAGACCTGTGTCCAGCACTGCCCTCCAGGCTTCGCCCCCAAGTCCTC
GATACGCACTATAGCACCGAGAAATGACGTGGAGACCATCCGGGCCAGCGTCTGCGCCCTGCCACGCTCATGTGCCACATGCCAGGG
GCCGGCCCTGACAGACTGCCCTCAGCTGCCCCAGCCACGCTCCTTGGACCTGTGGAGCAGACTTGCTCCCGGCAAGCCAGAGCAGCC
GAGAGTCCCAGCCACAGCAGCCACCTCGGCTGCCCTGGAGGTGGAGGCGGGCAACGGCTGCGGGCAGGGCTGCTGCCCTCACAC
CTGCCCTGAGGTGGCCGGCCCTCAGCTGGCCCTTCATCGTGTGGTCTTTCGTCACTGTCTTCTGGTCTCTGACGTGCGCTCTGGCTT
TAGTTTTCGGGGTGAAGGTGTACACCATGGACCGTGGCTCATCTCCTACAAGGGGCTGCCCCCTGAAGCCTGGCAGGAGGAGTGCC
CGTCTGACTCAGAAAGAGGACGAGGGCCGGGCGAGAGGACCGCCTTTATCAAGACCAAGCGCCCTCTGA

FIG. 2

MELRPWLLWVVAATGTLVLLAADAQQQKVFTNTWAVRIPGCPAVANSVARKHGFLNLGQIFGDYVHFVWHRGVTKRSLSPHPRHRLQR
EPQVQWLEQQVAKRRRTKRDVYQEPTDPKFPQQWYLSGVTQRDLNVKAAWAQGYTGHGIVVSIILDDGIEKNHPDDLAGNYDPGASFVNDQ
DPPQPRYTMNDNRHGTRCAGEVAAVANNGVCGVGAYNARIIGVVRMLDGEVTDAVEARSLGLNPNHHIYASWGPEDDDGKTVDGPA
RLAEAEFFRGVSQGRGGLGSI FVWASGNGGREHDS CNCDGYTNSIYTLSISSATQFCGNVPWYSEACSSTLATYSSGNQNEKQIVTTDL
RQKCTESHGTSASAPLAAGIIALTLEANKNLTWRDMQHLVVQTSKPAHLNANDWATNGVGRKVSHSYGYGLLDAGAMVALAQNWTTVA
PQRKCIIDILTEPKDIGKRLEVRKTVTACLGEPNHITRLEHAQARLTLSYNRRGDLAIHLVSPMGTRSTLLAARPHDYSADGFNDWAFM
TTHSWDEDPGSEWVLEIENTSEANNYGTLTKFTLVLYGTAPEGLPVPPESSGCKTLTSSQACVVCEEGLSLHQSVCVQHCPPGFAPQVL
DTHYSTENDVETIRASVCAPCHASCATCQGPALTDCLSCPSHASLDPVEQTCSRQSQSSRESPPQQQPPRPPPEVEAGQRLRAGLLPSH
LPEVVAGLSCAFI VLVFVTVFLVLQLRSGFSFRGVKVVYTMDRGLISYKGLPPEANQHECPSDSEDEGRGERTAFIKDQSAI

FIG. 3

FIG. 4







FIG. 5

10/14

FIG. 6



11/14


Residual vs. Predicted

FIG. 7A


Normal Q-Q plot for residuals

FIG. 7B

12/14




FIG. 7C





FIG. 7D

FIG. 8






FIG. 9