



(51) International Patent Classification: Not classified  
(21) International Application Number: PCT/IL2011/000426

(22) International Filing Date: 1 June 2011 (01.06.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 61/350,073 1 June 2010 (01.06.2010) US

(71) Applicant (for all designated States except US): **TODOS MEDICAL LTD.** [IL/IL]; 12 Kineret Street, POB 293, 70100 Airport City (IL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KAPELUSHNIK, Joseph** [IL/IL]; 385 Moshav Neve Ilan, 90850 Moshav Neve Ilan (IL). **MORDECHAI, Shaul** [IL/IL]; 21 Erez Street, 84965 Omer (IL). **NATHAN, Ilana** [IL/IL]; 5 Sirpad Street, 84965 Omer (IL). **ZELIG, Udi** [IL/IL]; 85455 Kibbutz Nir Yitzhak (IL). **ZIGDON, Rami** [IL/IL]; 6 Asher Street, 43553 Ra'anana (IL).

(74) Agents: **SANFORD T. COLB & CO.** et al.; P.O. Box 2273, 76122 Rehovot (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: DIAGNOSIS OF CANCER

FIG. 1A



(57) Abstract: A method is provided comprising, obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition. Other embodiments are also provided.

WO 2011/151825 A2

## DIAGNOSIS OF CANCER

### **CROSS-REFERENCES TO RELATED APPLICATIONS**

The present application claims the priority of US Provisional Application 61/350,073 to Kapelushnik et al., filed June 1, 2010, which is incorporated herein by reference.

### **5 FIELD OF EMBODIMENTS OF THE INVENTION**

Embodiments of the present invention relate generally to diagnosis and monitoring of cancer, and particularly to methods for diagnosis and monitoring of malignant solid tumors.

### **BACKGROUND**

Infrared spectroscopy is a technique based on the absorption or reflection of infrared  
10 radiation by chemical substances; each chemical substance having unique absorption spectra. Fourier Transform Infrared (FTIR) spectroscopy is used to identify biochemical compounds and examine the biochemical composition of a biological sample. Typically, FTIR spectra are composed of absorption bands each corresponding to specific functional groups related to cellular components such as lipids, proteins, carbohydrates and nucleic acids. Processes such as  
15 carcinogenesis may trigger global changes in cancer cell biochemistry resulting in differences in the absorption spectra when analyzed by FTIR spectroscopy techniques. Therefore, FTIR spectroscopy is commonly used to distinguish between normal and abnormal tissue by analyzing the changes in absorption bands of macromolecules such as lipids, proteins, carbohydrates and nucleic acids. Additionally, FTIR spectroscopy may be utilized for evaluation of cell death  
20 mode, cell cycle progression and the degree of maturation of hematopoietic cells.

### **SUMMARY OF EMBODIMENTS OF THE INVENTION**

In some applications of the present invention, methods and systems are provided for the diagnosis and monitoring of multiple types of malignant neoplasms, particularly malignant solid tumors.

25 Additionally or alternatively, some applications of the present invention comprise diagnosis and monitoring of a pre-malignant condition.

Typically "Total Biochemical Infrared Analysis" (TBIA) of blood-derived mononuclear cells is used to diagnose a solid tumor. For example, some applications of the present invention comprise analysis by infrared (IR) spectroscopy, e.g., FTIR spectroscopy and microspectroscopy, of global biochemical properties of blood-derived mononuclear cells for the  
5 detection of solid tumors. As provided by some applications of the present invention, FTIR Optical Diagnosis Technology (FODT) analysis of biochemical changes in a Peripheral Blood Mononuclear Cells (PBMC) sample of a patient can indicate the presence of a solid tumor and/or a pre-malignant condition.

For some applications, biochemical analysis of PBMC obtained from cancer patients and  
10 from control individuals who do not suffer from a malignant solid tumor, e.g., healthy controls, is conducted using FTIR microspectroscopy techniques. In accordance with some applications of the present invention, PBMC from a plurality of cancer patients each suffering from a solid tumor (e.g., in the breast, pancreas, lung, head and neck, prostate, ovary, and gastrointestinal tract) is analyzed by FTIR microspectroscopy techniques. Subsequently, the FTIR spectra  
15 (absorption and/or reflection) of the PBMC samples of the cancer patients are compared to the FTIR spectra of PBMC samples obtained from the controls.

The inventors have identified that the PBMC samples obtained from cancer patients suffering from a malignant solid tumor produce FTIR spectra that differ from those of the control individuals who do not suffer from a malignant solid tumor, allowing distinguishing between the  
20 cancer patients and controls. Thus, some applications of the present invention can be used to diagnose cancer patients suffering from various types of malignancies, particularly solid tumors. Importantly, the distinction by FTIR spectroscopy between controls and patients suffering from solid tumors is typically performed based on analysis of PBMC and not of the actual tumor cells.

For some applications, a data processor analyzes the IR spectrum, e.g., the FTIR  
25 spectrum, of the PBMC sample of a subject. Information from the data processor is typically fed into an output unit that generates a result indicative of the presence of a solid tumor and/or a pre-malignant condition, based on the infrared (IR) spectrum. Additionally, the data processor is typically configured to calculate a second derivative of the infrared (IR) spectrum of the PBMC

sample and, based on the second derivative of the infrared (IR) spectrum, to generate an output indicative of the presence of a solid tumor.

Additionally, the inventors have identified that PBMC obtained from each cancer patient suffering from a solid tumor produced an FTIR spectrum having a unique spectral pattern which is characteristic of the type of malignancy, e.g., breast, lung, prostate or gastrointestinal, and distinct from spectra of other malignancy types.

For some applications, analysis by IR spectroscopy, e.g., FTIR spectroscopy, of the biochemistry of PBMC or any other blood-derived cells is used for the screening of large populations, aiding in the early detection of solid tumors. FTIR spectroscopy (and microspectroscopy) is typically a simple, reagent-free and rapid method suitable for use as a screening test for large populations. Early detection of cancer generally enables early intervention and treatment, contributing to a reduced mortality rate.

There is therefore provided in accordance with some applications of the present invention a method including:

obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition.

For some applications, generating the output includes generating the output indicative of the presence of the solid tumor.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4$  cm<sup>-1</sup>,  $798 \pm 4$  cm<sup>-1</sup>,  $809 \pm 4$  cm<sup>-1</sup>,  $814 \pm 4$  cm<sup>-1</sup>,  $875 \pm 4$  cm<sup>-1</sup>,  $997 \pm 4$  cm<sup>-1</sup>,  $1001 \pm 4$  cm<sup>-1</sup>,  $1015 \pm 4$  cm<sup>-1</sup>,  $1103 \pm 4$  cm<sup>-1</sup>,  $1118 \pm 4$  cm<sup>-1</sup>,  $1162 \pm 4$  cm<sup>-1</sup>,  $1221 \pm 4$  cm<sup>-1</sup>,  $1270 \pm 4$  cm<sup>-1</sup>,  $1283 \pm 4$  cm<sup>-1</sup>,  $1295 \pm 4$

cm-1,  $1315 \pm 4$  cm-1,  $1341 \pm 4$  cm-1,  $1367 \pm 4$  cm-1,  $1392 \pm 4$  cm-1,  $1429 \pm 4$  cm-1,  $1440 \pm 4$  cm-1,  $1445 \pm 4$  cm-1 and  $1455 \pm 4$  cm-1.

For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

5 For some applications, analyzing includes assessing the characteristic at at least three wavenumbers selected from the group.

For some applications, assessing the characteristic includes analyzing a band of the IR spectrum surrounding at least one wavenumber selected from the group.

10 For some applications, analyzing the sample includes obtaining a second derivative of the infrared (IR) spectrum of the sample.

For some applications, the infrared (IR) spectrum includes an absorption spectrum, and obtaining the infrared (IR) spectrum includes obtaining the absorption spectrum.

For some applications, the infrared (IR) spectrum includes a reflection spectrum, and obtaining the infrared (IR) spectrum includes obtaining the reflection spectrum.

15 For some applications, generating the output includes indicating via the output whether the solid tumor is a first type of solid tumor or a second type of solid tumor.

For some applications, the solid tumor includes a solid tumor in tissue selected from the group consisting of: head and neck, esophagus, and pancreas, and generating the output includes generating an output indicative of the presence of a solid tumor in tissue selected from the group.

20 For some applications, the solid tumor includes a solid tumor in tissue selected from the group consisting of: breast, gastrointestinal tract, prostate, and lung, and generating the output includes generating an output indicative of the presence of a solid tumor in tissue selected from the group.

25 For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $752 \pm 4$  cm-1,  $1030 \pm 4$  cm-1,  $1046 \pm 4$  cm-1,  $1128 \pm 4$  cm-1, and  $1237 \pm 4$  cm-1, and generating includes generating an output indicative of the presence of a tumor in the breast tissue.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $797 \pm 4 \text{ cm}^{-1}$ ,  $830 \pm 4 \text{ cm}^{-1}$ ,  $893 \pm 4 \text{ cm}^{-1}$ ,  $899 \pm 4 \text{ cm}^{-1}$ ,  $1128 \pm 4 \text{ cm}^{-1}$ ,  $1298 \pm 4 \text{ cm}^{-1}$ ,  $1354 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $1725 \pm 4 \text{ cm}^{-1}$ ,  $1738, \pm 4 \text{ cm}^{-1}$ , and  $3013 \pm 4 \text{ cm}^{-1}$ , and generating includes generating an output  
5 indicative of the presence of a tumor in the gastrointestinal tract tissue.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $780 \pm 4 \text{ cm}^{-1}$ ,  $797 \pm 4 \text{ cm}^{-1}$ ,  $851 \pm 4 \text{ cm}^{-1}$ ,  $874 \pm 4 \text{ cm}^{-1}$ ,  $881 \pm 4 \text{ cm}^{-1}$ ,  $913 \pm 4 \text{ cm}^{-1}$ ,  $923 \pm 4 \text{ cm}^{-1}$ ,  $958 \pm 4 \text{ cm}^{-1}$ ,  $968, \pm 4 \text{ cm}^{-1}$ ,  $1044 \pm 4 \text{ cm}^{-1}$ ,  $1085 \pm 4 \text{ cm}^{-1}$ ,  $1191 \pm 4 \text{ cm}^{-1}$ ,  $1241 \pm 4 \text{ cm}^{-1}$ ,  $1344 \pm 4 \text{ cm}^{-1}$ ,  
10  $1373 \pm 4 \text{ cm}^{-1}$ ,  $1417 \pm 4 \text{ cm}^{-1}$ ,  $1458 \pm 4 \text{ cm}^{-1}$ ,  $1469 \pm 4 \text{ cm}^{-1}$ ,  $1692 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $1728 \pm 4 \text{ cm}^{-1}$ ,  $2852 \pm 4 \text{ cm}^{-1}$ , and  $2984 \pm 4 \text{ cm}^{-1}$ , and generating includes generating an output indicative of the presence of a tumor in the lung tissue.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $828 \pm 4 \text{ cm}^{-1}$ ,  $932 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1059 \pm 4 \text{ cm}^{-1}$ ,  $1299 \pm 4 \text{ cm}^{-1}$ ,  $1302 \pm 4 \text{ cm}^{-1}$ ,  $1403 \pm 4 \text{ cm}^{-1}$ ,  $1454 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $2979, \pm 4 \text{ cm}^{-1}$ , and  $3013 \pm 4 \text{ cm}^{-1}$ , and generating includes generating an output  
15 indicative of the presence of a tumor in the prostate tissue.

There is further provided in accordance with some applications of the present invention a method including:

20 obtaining an infrared (IR) spectrum of a sample of white blood cells by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition.

For some applications, generating the output includes generating the output indicative of  
25 the presence of the solid tumor.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

There is additionally provided in accordance with some applications of the present invention a method for monitoring the effect of an anti-cancer treatment on a subject undergoing anti-cancer treatment for a solid tumor, for use with a first Peripheral Blood Mononuclear Cells (PBMC) sample separated from blood of the subject that was obtained prior to initiation of the treatment and a second PBMC sample separated from blood of the subject that was obtained after initiation of the treatment, the method including:

obtaining IR spectra of the first and second PBMC samples by analyzing the first and second PBMC samples by IR spectroscopy; and

based on the IR spectra, generating an output indicative of the effect of the treatment.

For some applications, analyzing the first and second PBMC samples by IR spectroscopy includes analyzing the samples by Fourier Transformed Infrared spectroscopy, and obtaining the IR spectra includes obtaining Fourier Transformed Infrared (FTIR) spectra.

For some applications, analyzing the first and second PBMC samples by infrared (IR) spectroscopy includes analyzing the first and second PBMC samples by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

For some applications, the method includes obtaining an IR spectrum of a third PBMC sample separated from blood of the subject that was obtained following termination of the treatment, by analyzing the third PBMC sample by IR spectroscopy.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $798 \pm 4 \text{ cm}^{-1}$ ,  $809 \pm 4 \text{ cm}^{-1}$ ,  $814 \pm 4 \text{ cm}^{-1}$ ,  $875 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1001 \pm 4 \text{ cm}^{-1}$ ,  $1015 \pm 4 \text{ cm}^{-1}$ ,  $1103 \pm 4 \text{ cm}^{-1}$ ,  $1118 \pm 4 \text{ cm}^{-1}$ ,  $1162 \pm 4 \text{ cm}^{-1}$ ,  $1221 \pm 4 \text{ cm}^{-1}$ ,  $1270 \pm 4 \text{ cm}^{-1}$ ,  $1283 \pm 4 \text{ cm}^{-1}$ ,  $1295 \pm 4 \text{ cm}^{-1}$ ,  $1315 \pm 4 \text{ cm}^{-1}$ ,  $1341 \pm 4 \text{ cm}^{-1}$ ,  $1367 \pm 4 \text{ cm}^{-1}$ ,  $1392 \pm 4 \text{ cm}^{-1}$ ,  $1429 \pm 4 \text{ cm}^{-1}$ ,  $1440 \pm 4 \text{ cm}^{-1}$ ,  $1445 \pm 4 \text{ cm}^{-1}$ , and  $1455 \pm 4 \text{ cm}^{-1}$ .

For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

For some applications, analyzing includes assessing the characteristic at at least three wavenumbers selected from the group.

There is yet additionally provided in accordance with some applications of the present invention, a method including:

- 5       obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample; and  
      based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition.

10       For some applications, generating the output includes generating the output indicative of the presence of the solid tumor.

There is still additionally provided in accordance with some applications of the present invention, a system for diagnosing a solid tumor, including:

- a data processor, configured to analyze an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample of a subject; and  
15       an output unit, configured to generate an output indicative of the presence of a solid tumor, based on the infrared (IR) spectrum.

For some applications, the data processor is configured to calculate a second derivative of the infrared (IR) spectrum of the PBMC sample and, based on the second derivative of the infrared (IR) spectrum, to generate an output indicative of the presence of a solid tumor.

For some applications, the IR spectrum includes a Fourier Transformed Infrared (FTIR) spectrum, and the data processor is configured to calculate a second derivative of the FTIR spectrum.

- 20       For some applications, the data processor is configured to analyze the infrared (IR) spectrum by assessing a characteristic of the PBMC sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $798 \pm 4 \text{ cm}^{-1}$ ,  $809 \pm 4 \text{ cm}^{-1}$ ,  $814 \pm 4 \text{ cm}^{-1}$ ,  $875 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1001 \pm 4 \text{ cm}^{-1}$ ,  $1015 \pm 4 \text{ cm}^{-1}$ ,  $1103 \pm 4 \text{ cm}^{-1}$ ,  $1118 \pm 4 \text{ cm}^{-1}$ ,  $1162 \pm 4 \text{ cm}^{-1}$ ,  $1221 \pm 4 \text{ cm}^{-1}$ ,  $1270 \pm 4 \text{ cm}^{-1}$ ,  $1283 \pm 4 \text{ cm}^{-1}$ ,  $1295 \pm 4 \text{ cm}^{-1}$ ,  $1315 \pm 4 \text{ cm}^{-1}$ ,  $1341 \pm 4 \text{ cm}^{-1}$ ,  $1367 \pm 4 \text{ cm}^{-1}$ ,  $1392 \pm 4 \text{ cm}^{-1}$ ,  $1429 \pm 4 \text{ cm}^{-1}$ ,  $1440 \pm 4 \text{ cm}^{-1}$ ,  $1445 \pm 4 \text{ cm}^{-1}$ , and  $1455 \pm 4 \text{ cm}^{-1}$ .

For some applications, the data processor is configured to analyze the infrared (IR) spectrum by assessing the characteristic at at least two wavenumbers selected from the group.

For some applications, the data processor is configured to analyze the infrared (IR) spectrum by assessing the characteristic at at least three wavenumbers selected from the group.

5           There is still further provided in accordance with some applications of the present invention, a computer program product for administering processing of a body of data, the product including a computer-readable medium having program instructions embodied therein, which instructions, when read by a computer, cause the computer to:

          obtain an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) by  
10   analyzing the PBMC by infrared spectroscopy; and  
          based on the infrared spectrum, generate an output indicative of the presence of a solid tumor.

          There is yet provided in accordance with some applications of the present invention, a method including:

15           obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
          based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in a breast tissue of a subject.

          For some applications, analyzing includes assessing a characteristic of the sample at at  
20   least one wavenumber selected from the group consisting of:  $752 \pm 4$  cm<sup>-1</sup>,  $1030 \pm 4$  cm<sup>-1</sup>,  $1046 \pm 4$  cm<sup>-1</sup>,  $1128 \pm 4$  cm<sup>-1</sup>, and  $1237 \pm 4$  cm<sup>-1</sup>.

          For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

          For some applications, analyzing includes assessing the characteristic at at least three  
25   wavenumbers selected from the group.

          For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

There is still provided in accordance with some applications of the present invention a method including:

- 5           obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
            based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in tissue of a gastrointestinal tract of a subject.

10           For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $797 \pm 4 \text{ cm}^{-1}$ ,  $830 \pm 4 \text{ cm}^{-1}$ ,  $893 \pm 4 \text{ cm}^{-1}$ ,  $899 \pm 4 \text{ cm}^{-1}$ ,  $1128 \pm 4 \text{ cm}^{-1}$ ,  $1298 \pm 4 \text{ cm}^{-1}$ ,  $1354 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $1725 \pm 4 \text{ cm}^{-1}$ ,  $1738, \pm 4 \text{ cm}^{-1}$ , and  $3013 \pm 4 \text{ cm}^{-1}$ .

            For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

- 15           For some applications, analyzing includes assessing the characteristic at at least three wavenumbers selected from the group.

            For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

- 20           For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

            There is additionally provided in accordance with some applications of the present invention, a method including:

- obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC)  
25           sample by analyzing the sample by infrared spectroscopy; and  
            based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in lung tissue of a subject.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $780 \pm 4 \text{ cm}^{-1}$ ,  $797 \pm 4 \text{ cm}^{-1}$ ,  $851 \pm 4 \text{ cm}^{-1}$ ,  $874 \pm 4 \text{ cm}^{-1}$ ,  $881 \pm 4 \text{ cm}^{-1}$ ,  $913 \pm 4 \text{ cm}^{-1}$ ,  $923 \pm 4 \text{ cm}^{-1}$ ,  $958 \pm 4 \text{ cm}^{-1}$ ,  $968, \pm 4 \text{ cm}^{-1}$ ,  $1044 \pm 4 \text{ cm}^{-1}$ ,  $1085 \pm 4 \text{ cm}^{-1}$ ,  $1191 \pm 4 \text{ cm}^{-1}$ ,  $1241 \pm 4 \text{ cm}^{-1}$ ,  $1344 \pm 4 \text{ cm}^{-1}$ ,  
5  $1373 \pm 4 \text{ cm}^{-1}$ ,  $1417 \pm 4 \text{ cm}^{-1}$ ,  $1458 \pm 4 \text{ cm}^{-1}$ ,  $1469 \pm 4 \text{ cm}^{-1}$ ,  $1692 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $1728 \pm 4 \text{ cm}^{-1}$ ,  $2852 \pm 4 \text{ cm}^{-1}$ , and  $2984 \pm 4 \text{ cm}^{-1}$ .

For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

For some applications, analyzing includes assessing the characteristic at at least three  
10 wavenumbers selected from the group.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes  
15 analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

There is further additionally provided in accordance with some applications of the present invention, a method including:

obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and

20 based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in a prostate tissue of a subject.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $828 \pm 4 \text{ cm}^{-1}$ ,  $932 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1059 \pm 4 \text{ cm}^{-1}$ ,  $1299 \pm 4 \text{ cm}^{-1}$ ,  $1302 \pm 4 \text{ cm}^{-1}$ ,  $1403 \pm 4 \text{ cm}^{-1}$ ,  $1454 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $2979, \pm 4 \text{ cm}^{-1}$ , and  $3013 \pm 4 \text{ cm}^{-1}$ .  
25

For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

For some applications, analyzing includes assessing the characteristic at at least three wavenumbers selected from the group.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the  
5 infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

There is further provided in accordance with some applications of the present invention, a method including:

10 obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample from a cancer patient by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of a stage of the cancer.

For some applications, analyzing includes assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $865 \pm 4 \text{ cm}^{-1}$ ,  $897 \pm 4 \text{ cm}^{-1}$ ,  $924$   
15  $\pm 4 \text{ cm}^{-1}$ ,  $1030 \pm 4 \text{ cm}^{-1}$ ,  $1047 \pm 4 \text{ cm}^{-1}$ ,  $1191 \pm 4 \text{ cm}^{-1}$ , and  $1238 \pm 4 \text{ cm}^{-1}$ .

For some applications, analyzing includes assessing the characteristic at at least two wavenumbers selected from the group.

For some applications, analyzing includes assessing the characteristic at at least three wavenumbers selected from the group.

20 For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and obtaining the infrared (IR) spectrum includes obtaining a Fourier Transformed Infrared (FTIR) spectrum.

For some applications, analyzing the sample by infrared (IR) spectroscopy includes analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

25 The present invention will be more fully understood from the following detailed description of embodiments thereof, taken together with the drawings, in which:

### BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A-C are graphs representing FTIR absorption spectra and the second derivative of absorption spectra and analysis thereof, based on PBMC samples from several cancer patients and controls, derived in accordance with some applications of the present invention;

5 Figs. 2A-C are graphs representing FTIR absorption spectra and the second derivative of absorption spectra of PBMC from a pancreatic cancer patient compared to PBMC from healthy controls, derived in accordance with some applications of the present invention;

Figs. 3A-C are graphs showing the second derivative of spectra of PBMC from several cancer patients and healthy controls, derived in accordance with some applications of the present invention, and a table summarizing the main biochemical changes induced in PBMC of the  
10 cancer patients, as observed by FTIR microspectroscopy, as derived in accordance with some applications of the present invention; and

Fig. 4 shows second derivative spectra of PBMC from a breast cancer patient, a gastrointestinal cancer patient with a history of breast cancer, and healthy controls, as derived in  
15 accordance with some applications of the present invention; and

Figs. 5A-G are graphs representing the second derivative spectra of PBMC and analysis thereof, based on PBMC samples from cancer patients suffering from various types of solid tumors, derived in accordance with some applications of the present invention; and

Figs. 6A-C are graphs representing the second derivative spectra and analysis thereof, of  
20 PBMC samples from cancer patients suffering from different stages of cancer, derived in accordance with some applications of the present invention.

### DETAILED DESCRIPTION OF THE EMBODIMENTS

Some applications of the present invention comprise diagnosis of a solid tumor by techniques of IR spectroscopy, e.g., FTIR microspectroscopy (MSP) techniques. For some  
25 applications, FTIR Optical Diagnosis Technology (FODT) is used to diagnose a solid tumor based on biochemical properties of Peripheral Blood Mononuclear Cells (PBMC) of a subject. Some applications of the present invention comprise obtaining a blood sample from a subject and analyzing mononuclear cells from the sample by FTIR-MSP techniques for the detection of a

solid tumor. Typically, a PBMC sample of a patient suffering from a solid tumor is identified as exhibiting FTIR spectra that are different from FTIR spectra produced by PBMC from a subject who does not suffer from a solid tumor (for some applications, the control group may include subjects suffering from a pathology that is not a solid tumor). Accordingly, some applications  
5 of the present invention provide a useful method for the detection of cancer, specifically solid tumors. FTIR spectra of PBMC obtained from a cancer patient with a solid tumor generally reflect biochemical changes which occur in the PBMC of the patient in response to the tumor.

For some applications, methods of the present invention are used to determine a stage of the cancer.

10 For some applications, methods of the present invention can be used to provide monitoring and follow up of cancer patients during and after treatment, e.g., chemotherapy treatment. Typically, changes in FTIR spectra of PBMC of solid-tumor patients who are undergoing treatment can indicate biochemical changes in the cells in response to the treatment. This biochemical information can provide insight into the effect of treatment on the patient  
15 and/or the tumor.

In some applications of the present invention, analysis of PBMC by FTIR-MSP is used to detect a type of solid tumor. Typically, each type of malignant solid tumor produces distinct FTIR spectra of the PBMC, which are unique to the type of solid tumor. This can be due to each type of solid tumor inducing specific biochemical changes in PBMC.

## 20 **METHODS USED IN SOME EMBODIMENTS OF THE PRESENT INVENTION**

A series of protocols are described hereinbelow which may be used separately or in combination, as appropriate, in accordance with applications of the present invention. It is to be appreciated that numerical values are provided by way of illustration and not limitation. Typically, but not necessarily, each value shown is an example selected from a range of values  
25 that is within 20% of the value shown. Similarly, although certain steps are described with a high level of specificity, a person of ordinary skill in the art will appreciate that other steps may be performed, *mutatis mutandis*.

In accordance with some applications of the present invention, the following methods were applied:

**Obtaining patient and control populations**

All studies were approved by the local Ethics Committee of the Soroka University Medical Center and conducted in accordance with the Declaration of Helsinki. Qualified personnel obtained informed consent from each individual participating in this study.

- 5 The patient population included cancer patients (n=63) diagnosed with primary solid tumors as set forth in the following Table I:

**Table I:**

		Control	Disease site				
			Breast	GI	Lung	Pro-state	Other
Gender	Male	24	1	9	7	5	2
	Female	22	24	9	1	0	5
Disease stage	I	0	9	1	0	2	2
	II	0	12	3	0	0	0
	III	0	3	4	1	0	3
	IV	0	1	10	7	3	2
Total		46	25	18	8	5	7

10

The patient population categorized under "other" included six patients each diagnosed with a different type of primary tumor. Among the "other" group are patients suffering from primary tumors such as pancreas (n=2), head and neck (n=1) and esophagus (n=1).

- 15 The diagnosis of cancer was determined by clinical, surgical, histological, and pathologic diagnosis. The pathologic stage of the tumor was determined according to tumor-node-metastasis (TNM) classification, as described in "TNM Classification of Malignant Tumours", by Sobin LH. et al., 7th Edition, New York: John Wiley, 2009.

- 20 A control group (n=46) included healthy volunteers who underwent detailed clinical questioning to exclude a possible pathology, at the Soroka University Medical Center and Ben-Gurion University.

### **Collection of blood samples**

1-2 ml of peripheral blood was collected in 5 ml EDTA blood collection tubes (BD Vacutainer® Tubes, BD Vacutainer, Toronto) from patients and healthy controls using standardized phlebotomy procedures. Samples were processed within two hours of collection.

### 5 **Extraction of peripheral blood mononuclear cells (PBMC)**

Platelet-depleted residual leukocytes obtained from cancer patients and healthy controls were applied to Histopaque 1077 gradients (Sigma Chemical Co., St. Louis, Missouri, USA) following manufacturer's protocol to obtain PBMC.

10 The cells were aspirated from the interface, washed twice with isotonic saline (0.9% NaCl solution) at 250 g, and resuspended in 5 µl fresh isotonic saline. 1.5 µl of washed cells were deposited on zinc selenide (ZnSe) slides to form approximately a monolayer of cells. It is noted that any other suitable slide may be used, e.g., reflection measurements may be carried out using a gold slide. The slides were then air dried for 1 h under laminar flow to remove water. The dried cells were then measured by FTIR microscopy.

### 15 **FTIR-Microspectroscopy**

Fourier Transform Infrared Microspectroscopy (FTIR-MSP) and Data Acquisition Measurements on cell cultures were performed using the FTIR microscope IR scope 2 with a liquid-nitrogen-cooled mercury-cadmium-telluride (MCT) detector, coupled to the FTIR spectrometer Bruker Equinox model 55/S, using OPUS software (Bruker Optik GmbH, 20 Ettlingen, Germany). For some of the experiments, Fourier Transform Infrared Microspectroscopy (FTIR-MSP) and Data Acquisition Measurements were performed using the FTIR microscope Nicolet Centaurus with a liquid-nitrogen-cooled mercury-cadmium-telluride (MCT) detector, coupled to the FTIR spectrometer Nicolet iS10, OMNIC software (Nicolet, Madison, WI) using OPUS software (Bruker Optik GmbH, Ettlingen, Germany). Essentially the 25 same results were obtained with each of these microscopes.

To achieve high signal-to-noise ratio (SNR), 128 coadded scans were collected in each measurement in the wavenumber region 700 to 4000 cm<sup>-1</sup>. The measurement site was circular with a diameter of 100 µm and a spectral resolution of 4 cm<sup>-1</sup> (0.482 cm<sup>-1</sup> data spacing). To

reduce cell amount variation and facilitate proper comparison between different samples, the following procedures were adopted.

1. Each sample was measured at least five times at different spots.
2. Analog to Digital Converter (ADC) rates were empirically chosen between 2000 to 3000 counts / sec (providing measurement areas with similar cellular density).
3. The obtained spectra were baseline corrected using the rubber band method, with 64 consecutive points, and normalized using vector normalization in OPUS software as described in an article entitled "Early spectral changes of cellular malignant transformation using Fourier transformation infrared microspectroscopy," by Bogomolny et al., 2007. *J Biomed Opt.*12:024003.

In order to obtain precise absorption values at a given wavenumber with minimal background interference, the second derivative spectra were used to determine concentrations of bio-molecules of interest. This method is susceptible to changes in full width at half maximum (FWHM) of the IR bands. However, in the case of biological samples, all cells from the same type are composed from similar basic components which give relatively broad bands. Thus, it is possible to generally neglect the changes in bands FWHM as described in an article entitled "Selenium alters the lipid content and protein profile of rat heart: An FTIR microspectroscopy study," by Toyran et al., *Arch.Biochem.Biophys.* 2007 458:184-193.

#### **Statistical analysis:**

Statistical analysis was performed using STATISTICA software (STATISTICA, StatSoft, Inc., Tulsa, OK) and the student T-test. P-values < 0.05 were considered significant.

### **EXPERIMENTAL DATA**

The experiments described hereinbelow were performed by the inventors in accordance with applications of the present invention and using the techniques described hereinabove.

The experiments presented hereinbelow with reference to Example 1 and Example 2 demonstrate that in accordance with some applications of the present invention, analysis of PBMC samples by FTIR-MSP techniques can be used for diagnosis of a solid tumor based on the FTIR-MSP spectral pattern at selected wavenumbers.

### Example 1

In a set of experiments, PBMC samples from 46 healthy controls were analyzed by FTIR-MSP, and a typical FTIR-MSP spectral pattern was established for control PBMC. Additionally, PBMC samples from 63 cancer patients suffering from multiple types of solid tumors were subjected to FTIR-MSP analysis and compared to the control FTIR-MSP spectral pattern. The PBMC samples were obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP, in accordance with the protocols described hereinabove with reference to FTIR-MSP.

Reference is made to Figs. 1A-C, which are graphs representing FTIR absorption spectra and the second derivative of the absorption spectra and analysis thereof, for PBMC samples from 63 cancer patients suffering from solid tumors and 46 healthy controls, derived in accordance with some applications of the present invention.

FTIR-MSP analysis of peripheral blood mononuclear cells (PBMC) typically generated spectra in the region of 4000-700  $\text{cm}^{-1}$ . The spectra are composed of several absorption bands, each corresponding to specific functional groups of specific macromolecules. Fig. 1A shows an average of the FTIR-MSP spectra of PBMC samples of healthy controls and cancer patients in the regions of 4000-700  $\text{cm}^{-1}$ , after baseline correction and vector normalization. The spectra are composed of several absorption bands, each corresponding to specific functional groups of specific macromolecules such as lipids, proteins, and carbohydrates /nucleic acids. The main absorption bands are marked. For example, the region 3000-2830  $\text{cm}^{-1}$  contains symmetric and antisymmetric stretching of  $\text{CH}_3$  and  $\text{CH}_2$  groups, which correspond mainly to proteins and lipids respectively. The region 1700-1500  $\text{cm}^{-1}$  corresponds to amide I and amide II, which contain information regarding the secondary structures of proteins. The region 1300-1000  $\text{cm}^{-1}$  includes the symmetric and antisymmetric vibrations of  $\text{PO}_2^-$  groups. 1000-700  $\text{cm}^{-1}$  is a 'finger print' region which contains several different vibrations, corresponding to carbohydrates, lipids, nucleic acids and other bio-molecules, as described in an article by Mantsch M and Chapman D. entitled "Infrared spectroscopy of bio molecules," John Wiley New York 1996. The FTIR spectrum was typically analyzed by tracking changes in absorption (intensity and/or shift) of these macromolecules.

Table II represents some of the main IR absorption bands for PBMC cells, and their corresponding molecular functional groups:

**Table II:**

Wavenumber (cm <sup>-1</sup> ± 4)	Assignment
2958	$\nu_{as}$ CH <sub>3</sub> , mostly proteins, lipids
2922	$\nu_{as}$ CH <sub>2</sub> , mostly lipids, proteins
2873	$\nu_s$ CH <sub>3</sub> , mostly proteins, lipids
2854	$\nu_s$ CH <sub>2</sub> , mostly lipids, proteins
~1,656	Amide I $\nu$ C=O (80%), $\nu$ C-N (10%), $\delta$ N-H
~1,546	Amide II $\delta$ N-H (60%), $\nu$ C-N (40%)
1400	$\nu$ COO <sup>-</sup> , $\delta$ s CH <sub>3</sub> lipids, proteins
1313	Amide III band components of proteins
1240	$\nu_{as}$ PO <sub>2</sub> <sup>-</sup> , phosphodiester groups of nucleic acids
1170	C-O bands from glycomaterials and proteins
1155	$\nu$ C-O of proteins and carbohydrates
1085	$\nu_s$ PO <sub>2</sub> <sup>-</sup> of nucleic acids, phospholipids, proteins
1053	$\nu$ C-O & $\delta$ C-O of carbohydrates
996	C-C & C-O of ribose of RNA
967	C-C & C-O of deoxyribose skeletal motions of DNA
780	sugar-phosphate Z conformation of DNA
740	$\nu$ N=H of Thymine

Reference is made to Fig. 1B. In order to achieve effective comparison between the PBMC samples of the cancer patients and the controls, the second derivative of the baseline-corrected, vector-normalized FTIR-MSP spectra was used. Results are presented in Fig. 1B. As shown, the second derivative spectra of PBMC samples from the cancer patients differed significantly from the second derivative spectra of PBMC samples from the healthy controls in the spectral region of 1340-1260  $\text{cm}^{-1}$ . The mean  $\pm$  SEM is represented by the hashed region (for control) and the dotted region (for cancer) as shown in the exploded view in Fig. 1B.

Reference is made to Fig. 1C, which is a graph representing values of the second derivative of absorption spectra at wavenumbers A1-A23 of PBMC samples from cancer patients compared to PBMC samples from healthy controls, derived in accordance with some applications of the present invention. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC samples from the cancer patients differed significantly from the second derivative analysis of FTIR-MSP spectra from PBMC of healthy controls.

Table III lists the wavenumbers shown in Fig. 1C. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between healthy controls and cancer patients.

**Table III:**

<i>Control vs. Cancer</i>							
	Wavenumber (cm-1 ± 4)		Wavenumber (cm-1 ± 4)		Wavenumber (cm-1 ± 4)		Wavenumber (cm-1 ± 4)
<b>A1</b>	765	<b>A7</b>	1001	<b>A13</b>	1270	<b>A19</b>	1392
<b>A2</b>	798	<b>A8</b>	1015	<b>A14</b>	1283	<b>A20</b>	1429
<b>A3</b>	809	<b>A9</b>	1103	<b>A15</b>	1295	<b>A21</b>	1440
<b>A4</b>	814	<b>A10</b>	1118	<b>A16</b>	1315	<b>A22</b>	1445
<b>A5</b>	875	<b>A11</b>	1162	<b>A17</b>	1341	<b>A23</b>	1455
<b>A6</b>	997	<b>A12</b>	1221	<b>A18</b>	1367		

**Example 2**

In this set of experiments, PBMC from a single pancreatic cancer patient was subjected to FTIR-MSP analysis and compared to a control FTIR-MSP spectral pattern based on PBMC from 27 healthy controls. Results are presented in Figs. 2A-C. The PBMC was obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP in accordance with the protocols described hereinabove with reference to FTIR-Microspectroscopy.

Fig. 2A shows representative FTIR-MSP spectra of PBMC of a healthy control compared to FTIR-MSP spectra of PBMC of a pancreatic cancer patient after baseline correction and Min-Max normalization to amide II. Each spectrum represents the average of five measurements at

different sites for each sample. The spectra are composed of several absorption bands, each corresponding to specific functional groups of specific macromolecules such as lipids, proteins, carbohydrates and nucleic acids. The main absorption bands are marked. The FTIR spectrum was analyzed by tracking changes in absorption (intensity and/or shift) of these macromolecules.

5 As shown in Fig. 2A, the FTIR-MSP spectra derived from analysis of PBMC from the pancreatic cancer patient exhibited a different spectral pattern when compared to the FTIR-MSP spectra of PBMC of healthy controls.

Reference is made to Figs. 2B-C. In order to increase accuracy and achieve effective comparison between the PBMC sample of the pancreatic cancer patient and healthy controls, the second derivative of the baseline-corrected, vector-normalized FTIR spectra was used. Results are presented in Figs. 2B-C. The main absorption bands are marked. As shown, the second derivative spectral pattern of PBMC from the pancreatic cancer patient differed significantly from an average FTIR-MSP spectral pattern of PBMC of the healthy controls. The mean  $\pm$  SEM for the controls is represented by the hashed region as shown in the exploded view in Figs. 2B-C.

15 Reference is now made to Example 3 - Example 5. The experiments presented hereinbelow with reference to Example 3 - Example 5 demonstrate that in accordance with some applications of the present invention, analysis of PBMC samples by FTIR-MSP techniques is used to detect a type of solid tumor. Typically, each type of malignant solid tumor produces distinct FTIR spectra of the PBMC which are unique to the type of solid tumor.

### 20 Example 3

In this set of preliminary experiments, PBMC from cancer patients suffering from solid tumors, and PBMC from healthy controls was analyzed by FTIR-MSP. The population of cancer patients for this set of experiments comprised a total of 5 patients suffering from the following solid tumors: Breast (n=1), lung (n=1), pancreas (n=1), head and neck (n=1) and esophagus (n=1). The PBMC was obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP in accordance with the protocols described hereinabove with reference to FTIR-Microspectroscopy.

The results show that the FTIR-MSP spectral pattern of all the cancer patients differs from those of the healthy controls. The results additionally show that each type of malignancy produces a distinct spectral absorption pattern of the PBMC, which is unique to each type of solid tumor.

5 Reference is made to Figs. 3A-B. In order to increase accuracy and achieve effective comparison between PBMC samples of cancer patients and healthy controls, the second derivative of the baseline-corrected, vector-normalized FTIR spectra was used. Results are presented in Fig. 3A. As shown in Fig. 3A, the second derivative spectral pattern of PBMC from each one of the cancer patients differed from PBMC of the healthy controls. For example,  
10 the FTIR-MSP spectrum of PBMC of the lung cancer patient is distinct from control by exhibiting decreased absorbance of  $\text{CH}_2$ , which corresponds to cellular lipids; a decrease in  $\nu_{\text{as}} \text{PO}_2^-$ ; a shift to a higher wavenumber at  $967 \text{ cm}^{-1}$ , which corresponds to deoxyribose skeletal motions of DNA; and an increase in RNA absorption.

15 Additionally, each spectrum has its own unique spectral pattern, which is distinct from control, and which is characteristic of each type of malignancy.

Reference is made to Fig. 3B, which is a graph showing an analysis of the second derivative data shown in Fig. 3A. Fig. 3B represents the change in value of each type of cancer relative to the control, as derived by analysis by FTIR-MSP of PBMC from each patient. As shown, each cancer patient exhibited a spectrum that differed from the control.

20 Fig. 3C is a table summarizing the main biochemical changes induced in PBMC of cancer patients suffering from different types of solid tumors, as observed by FTIR-MSP (shown in Figs. 3A-B) and analyzed in accordance with some applications of the present invention. Peak intensities which indicate absorption were calculated for each spectrum to reveal the main biochemical changes characteristic of each type of tumor.

#### 25 Example 4

In this set of experiments, PBMC from a breast cancer patient and a gastrointestinal cancer patient were analyzed by FTIR-MSP, and compared to analysis of PBMC from healthy controls. It is to be noted that the breast cancer patient suffers from a primary breast tumor, and the gastrointestinal cancer patient suffers from a primary gastrointestinal tumor. The

gastrointestinal cancer patient has a history of breast cancer, and a pathological evaluation of the gastrointestinal tumor showed a breast cancer phenotype rather than a gastrointestinal phenotype.

First, peripheral blood was extracted from the two cancer patients and 27 healthy controls. The PBMC was obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP in accordance with the protocols described hereinabove with reference to FTIR-Microspectroscopy.

The results show that the FTIR-MSP spectral pattern of PBMC from the cancer patients differs significantly from those of the healthy controls. The results additionally show that PBMC of the breast cancer patient and the gastrointestinal cancer patient (with a history of breast cancer) exhibited a similar spectral pattern that was distinct from the control. This can be explained by the gastrointestinal tumor being shown by pathological evaluation to have a phenotype characteristic of a breast tumor. In some cases of breast cancer, malignant cells break away from the primary breast tumor and spread to other parts of the body. These cells may remain inactive for years before they begin to grow again. It is possible that a tumor, although located remotely from the original tumor site, triggers biochemical changes in the PBMC that are similar to those triggered by the original tumor. It is to be noted that the spectral absorbance pattern of PBMC of both the breast cancer patient and the gastrointestinal cancer patient (who has a history of breast cancer) differ from the spectral pattern produced by PBMC of patients suffering from other types of solid tumors.

Fig. 4 shows second derivative spectra of PBMC from the breast cancer patient and from the gastrointestinal cancer patient (the gastrointestinal cancer patient having a history of breast cancer and the gastrointestinal tumor exhibiting a phenotype characteristic of a breast tumor) compared to PBMC of healthy controls, as derived in accordance with some applications of the present invention. As shown, the FTIR-MSP second derivative spectra exhibit significant differences between PBMC from the cancer patients and PBMC from healthy controls. These spectral differences typically represent molecular changes in the PBMC of the cancer patients and allow distinguishing between the healthy control and the cancer patients. In addition, PBMC from both the primary breast cancer patient and the gastrointestinal cancer patient exhibited a similar FTIR-MSP spectral pattern. Accordingly, some applications of the present invention can

be used to diagnose the type and/or the origin of solid tumor of a patient based on the unique FTIR-MSP spectra produced by analysis of the patient's PBMC. This is consistent with Example 3, which showed that PBMC from patients with different types of solid tumors each produced a distinct FTIR spectral pattern with its own set of characteristic bands. Additionally, the molecular changes which trigger the changes in the FTIR-MSP spectra of breast cancer patients, including patients with breast cancer history, can serve as biomarkers to diagnose breast cancer. Additionally, some applications of the present invention can be used to select effective treatment based on the origin of a tumor. It is to be noted that the type of solid tumor diagnosed in accordance with applications of the present invention is not limited to breast tumors, but may include any other type of solid tumors.

#### Example 5

In this set of experiments, PBMC samples from cancer patients suffering from various types of solid tumors were analyzed by FTIR-MSP. The FTIR-MSP spectral pattern of each type of solid tumor was compared to the FTIR-MSP spectral pattern of the other solid tumors, allowing distinguishing between different types of solid tumors. The population of cancer patients for this set of experiments comprised a total of 63 patients suffering from solid tumors, in accordance with Table I. The PBMC was obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP, in accordance with the protocols described hereinabove with reference to FTIR-Microspectroscopy.

The results show that each type of solid tumor produces a distinct spectral absorption pattern of the PBMC, which is unique to each type of solid tumor, allowing distinguishing between different types of solid tumors.

Reference is made to Figs. 5A-C. In order to increase accuracy and achieve effective comparison between PBMC samples of the various types of cancer patients, the second derivative of the baseline-corrected, vector-normalized average FTIR absorption spectra was used. Results are presented in Figs. 5A-C, showing the second derivative of several regions of the spectra (the main absorption bands are marked). As shown in Figs. 5A-C, the second

derivative spectra of each type of solid tumor produced a distinct spectral absorption pattern of the PBMC, which is unique to each type of solid tumor.

Reference is made to Figs. 5D-G, which are a series of graphs representing values of the second derivative of absorption spectra of each type of solid tumor compared to the second derivative of absorption spectra of the other solid tumors, derived in accordance with some applications of the present invention.

Fig. 5D is a graph representing values of the second derivative of absorption spectra of PBMC samples from the breast cancer patients (n=25) compared to PBMC samples from the cancer patients suffering from other types of solid tumors that do not include breast cancer (n=38), at wavenumbers B1 - B5. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC from the breast cancer patients differed significantly from the second derivative analysis of FTIR-MSP spectral patterns from PBMC of other cancer patients who do not have breast cancer.

Table IV lists the wavenumbers shown in Fig. 5D. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between breast cancer patients and cancer patients who do not have breast cancer.

**Table IV:**

<i>Breast vs. Non Breast</i>	
	Wavenumber (cm-1 ± 4)
<b>B1</b>	752
<b>B2</b>	1030
<b>B3</b>	1046
<b>B4</b>	1128
<b>B5</b>	1237

Fig. 5E is a graph representing values of the second derivative of absorption spectra of PBMC samples from the gastrointestinal cancer patients (n=18) compared to PBMC samples

from the cancer patients suffering from other types of solid tumors that do not include gastrointestinal tumors (n=45), at wavenumbers C1 – C11. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC from the gastrointestinal cancer patients differed significantly from the second derivative analysis of FTIR-MSP spectral pattern from PBMC of other cancer patients who do not have gastrointestinal cancer.

Table V lists the wavenumbers shown in Fig. 5E. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between gastrointestinal cancer patients and cancer patients who do not have gastrointestinal cancer.

**Table V:**

<i>GI vs. Non GI</i>	
	Wavenumber (cm-1 ± 4)
<b>C1</b>	797
<b>C2</b>	830
<b>C3</b>	893
<b>C4</b>	899
<b>C5</b>	1128
<b>C6</b>	1298
<b>C7</b>	1354
<b>C8</b>	1714
<b>C9</b>	1725
<b>C10</b>	1738
<b>C11</b>	3013

Fig. 5F is a graph representing values of the second derivative of absorption spectra of PBMC samples from the lung cancer patients (n=8) compared to PBMC samples from the cancer patients suffering from other types of solid tumors that do not include lung tumors (n=55) at

wavenumbers D1 – D24. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC from the lung cancer patients differed significantly from the second derivative analysis of FTIR-MSP spectral patterns from PBMC of other cancer patients who do not have lung cancer.

- 5 Table VI lists the wavenumbers shown in Fig. 5F. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between lung cancer patients and cancer patients who do not have lung cancer.

**Table VI:**

<i>Lung vs. Non Lung</i>	
	Wavenumber ( $\text{cm}^{-1} \pm 4$ )
<b>D1</b>	765
<b>D2</b>	780
<b>D3</b>	797
<b>D4</b>	851
<b>D5</b>	874
<b>D6</b>	881
<b>D7</b>	913
<b>D8</b>	923
<b>D9</b>	958
<b>D10</b>	968
<b>D11</b>	1044
<b>D12</b>	1085
<b>D13</b>	1191
<b>D14</b>	1241
<b>D15</b>	1344

<b>D16</b>	1373
<b>D17</b>	1417
<b>D18</b>	1458
<b>D19</b>	1469
<b>D20</b>	1692
<b>D21</b>	1714
<b>D22</b>	1728
<b>D23</b>	2852
<b>D24</b>	2984

5

10

Fig. 5G is a graph representing values of the second derivative of absorption spectra of PBMC samples from the prostate cancer patients (n=5) compared to PBMC samples from the cancer patients suffering from other types of solid tumors that do not include prostate cancer (n=58) at wavenumbers E1 – E11. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC from the prostate cancer patients differed significantly from the second derivative analysis of FTIR-MSP spectral pattern from PBMC of other cancer patients who do not have prostate cancer.

Table VII lists the wavenumbers shown in Fig. 5G. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between prostate cancer patients and cancer patients who do not have prostate cancer.

20

**Table VII:**

<i>Prostate vs. Non Prostate</i>	
	Wavenumber (cm-1 ± 4)
<b>E1</b>	828
<b>E2</b>	932
<b>E3</b>	997
<b>E4</b>	1059
<b>E5</b>	1299
<b>E6</b>	1302
<b>E7</b>	1403
<b>E8</b>	1454
<b>E9</b>	1714
<b>E10</b>	2979
<b>E11</b>	3013

Reference is now made to Example 6. The experiments presented hereinbelow with reference to Example 6 demonstrate that in accordance with some applications of the present invention, analysis of PBMC samples by FTIR-MSP techniques is used for staging cancer. Typically, each stage of cancer produces distinct FTIR spectra of the PBMC.

**Example 6**

In this set of experiments, PBMC samples from cancer patients suffering from different stages of cancer due to solid tumors were analyzed by FTIR-MSP and a second derivative of the average of the spectra was obtained for each of the stages of cancer, allowing distinguishing between different stages of cancer. The population of cancer patients for this set of experiments comprised a total of 63 patients suffering from the different stages (stages one and two (n=29), and stages three and four (n=34)) of cancer, as described in Table I.

The PBMC was obtained by preliminary processing of the peripheral blood in accordance with the protocols described hereinabove with reference to extraction of peripheral blood mononuclear cells (PBMC). The PBMC samples were then analyzed by FTIR-MSP in accordance with the protocols described hereinabove with reference to FTIR-Microspectroscopy.

5 The results show that early stages of cancer produce spectral absorption pattern of the PBMC, that are different than those produced by PBMC samples taken from patients with advanced stages of cancer, allowing distinguishing among different stages of cancer, in particular, between early and advanced stages.

10 Reference is made to Figs. 6A-C, which are graphs showing the second derivative spectra of PBMC and analysis thereof, based on PBMC samples from cancer patients suffering from different stages of cancer, derived in accordance with some applications of the present invention.

15 Fig. 6A is a graph representing the second derivative of baseline-corrected, vector-normalized average FTIR absorption spectra of PBMC samples obtained from cancer patients in different stages of the disease. As shown, PBMC of each stage of cancer produced a distinct spectral absorption pattern of the PBMC.

20 Fig. 6B is a graph representing the second derivative of baseline-corrected, vector-normalized average FTIR absorption spectra of PBMC samples obtained from breast cancer patients in different stages of the disease. As shown, PBMC of each stage of breast cancer produced a distinct spectral absorption pattern of the PBMC, and particularly, the early stages (stages one and two) were distinct from the more advanced stages (stages three and four). It is to be noted that breast cancer is shown by way of illustration and not limitation, and that the scope of the present includes staging of any type of solid tumor by techniques described herein.

25 Fig. 6C is a graph representing values of the second derivative of absorption spectra of PBMC samples from stage one and two (n=29) cancer patients, compared to stage three and four (n=34) cancer patients at wavenumbers F1 – F7. Statistical analysis was performed and P-values are provided. As shown, the second derivative of PBMC from the cancer patients with early stages of cancer (stages one and two) differed significantly from the second derivative analysis of FTIR-MSP spectral pattern of cancer patients with advanced stages of cancer (stage three and four).

Table VIII lists the wavenumbers shown in Fig. 6A. Typically, PBMC samples were analyzed by FTIR-MSP techniques using these wavenumbers to distinguish between early stages (stage one and two) and more advanced stages (three and four) of cancer.

5 **Table VIII:**

<i>Initial Stages vs. Advanced Stages</i>	
	Wavenumber (cm-1 $\pm$ 4)
F1	865
F2	897
F3	924
F4	1030
F5	1047
F6	1191
F7	1238

10

15

Reference is made to Figs. 1-6 and Examples 1-6. It is to be noted that techniques described herein with reference to use of peripheral blood mononuclear cells (PBMC) may be applied to any type of white blood cell (WBC) or a combination of types of white blood cells. For example, analysis by FTIR microscopy techniques may be performed on any type of white blood cell, including but not limited to a total population of white blood cells (e.g., as obtained by red blood cell lysis).

20

Reference is still made to Figs. 1-6 and Examples 1-6.

The data obtained by analysis of the PBMC samples may be further analyzed by any suitable method known in the art, e.g., Artificial Neural Network and/or Cluster Analysis, and/or Principal Component Analysis, and/or Linear Discriminant Analysis (LDA) e.g., Fisher's Linear

25

Discriminant Analysis (FLDA), Quadratic Discriminant Analysis, and/or Non Linear Discriminant Analysis.

For example, data obtained in accordance with applications of the present invention may be analyzed by an artificial neural network (ANN). Several biomarkers shown in Tables II-VII  
5 which are statistically significant ( $p < 0.05$ ) may be served as an input vector for the ANN analysis.

It is further noted that the scope of the present invention includes the use of only one wavenumber (representing one biomarker) for detection and/or monitoring of a solid tumor, as well as the use of two, three, four, or more wavenumbers.

10 Additionally, the scope of the present invention includes using any IR spectral feature or any feature derived from analysis of an IR spectral feature (e.g., any type of peak analysis), to indicate the presence of a solid tumor.

It is also noted that the scope of the present invention is not limited to any particular form or analysis of IR spectroscopy. For example, IR spectroscopy may include Attenuated Total  
15 Reflectance (ATR) spectroscopy techniques.

Although applications of the present invention are described hereinabove with respect to spectroscopy, microspectroscopy, and particularly FTIR spectroscopy, the scope of the present invention includes the use of analysis techniques with data obtained by other means as well (for example, using a monochromator or an LED, at specific single wavenumbers, and/or FTIR  
20 imaging).

It will additionally be understood by one skilled in the art that aspects of the present invention described hereinabove can be embodied in a computer running software, and that the software can be supplied and stored in tangible media, e.g., hard disks, floppy disks, a USB flash drive, or compact disks, or in intangible media, e.g., in an electronic memory, or on a network  
25 such as the Internet.

It will be appreciated by persons skilled in the art that the present invention is not limited to what has been particularly shown and described hereinabove. Rather, the scope of the present invention includes both combinations and subcombinations of the various features described

hereinabove, as well as variations and modifications thereof that are not in the prior art, which would occur to persons skilled in the art upon reading the foregoing description.

## CLAIMS

1. A method comprising:  
obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
5 based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition.
2. The method according to claim 1, wherein generating the output comprises generating the output indicative of the presence of the solid tumor.
3. The method according to claim 1, wherein analyzing the sample by infrared (IR)  
10 spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.
4. The method according to claim 3, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared  
15 microspectroscopy (FTIR-MSP).
5. The method according to claim 1, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4$  cm<sup>-1</sup>,  $798 \pm 4$  cm<sup>-1</sup>,  $809 \pm 4$  cm<sup>-1</sup>,  $814 \pm 4$  cm<sup>-1</sup>,  $875 \pm 4$  cm<sup>-1</sup>,  $997 \pm 4$  cm<sup>-1</sup>,  $1001 \pm 4$  cm<sup>-1</sup>,  $1015 \pm 4$  cm<sup>-1</sup>,  $1103 \pm 4$  cm<sup>-1</sup>,  $1118 \pm 4$  cm<sup>-1</sup>,  $1162 \pm 4$  cm<sup>-1</sup>,  $1221 \pm 4$  cm<sup>-1</sup>,  $1270 \pm 4$  cm<sup>-1</sup>,  $1283 \pm 4$  cm<sup>-1</sup>,  $1295 \pm 4$  cm<sup>-1</sup>,  $1315 \pm 4$  cm<sup>-1</sup>,  $1341 \pm 4$  cm<sup>-1</sup>,  $1367 \pm 4$  cm<sup>-1</sup>,  $1392 \pm 4$  cm<sup>-1</sup>,  $1429 \pm 4$  cm<sup>-1</sup>,  $1440 \pm 4$  cm<sup>-1</sup>,  $1445 \pm 4$  cm<sup>-1</sup> and  $1455 \pm 4$  cm<sup>-1</sup>.  
20
6. The method according to claim 5, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.
7. The method according to claim 5, wherein analyzing comprises assessing the  
25 characteristic at at least three wavenumbers selected from the group.
8. The method according to claim 5, wherein assessing the characteristic comprises analyzing a band of the IR spectrum surrounding at least one wavenumber selected from the group.

9. The method according to claim 1, wherein analyzing the sample comprises obtaining a second derivative of the infrared (IR) spectrum of the sample.
10. The method according to claim 1, wherein the infrared (IR) spectrum includes an absorption spectrum, and wherein obtaining the infrared (IR) spectrum comprises obtaining the absorption spectrum.
11. The method according to claim 1, wherein the infrared (IR) spectrum includes a reflection spectrum, and wherein obtaining the infrared (IR) spectrum comprises obtaining the reflection spectrum.
12. The method according to claim 1, wherein generating the output comprises indicating via the output whether the solid tumor is a first type of solid tumor or a second type of solid tumor.
13. The method according to claim 1, wherein the solid tumor includes a solid tumor in tissue selected from the group consisting of: head and neck, esophagus, and pancreas, and wherein generating the output comprises generating an output indicative of the presence of a solid tumor in tissue selected from the group.
14. The method according to claim 1, wherein the solid tumor includes a solid tumor in tissue selected from the group consisting of: breast, gastrointestinal tract, prostate, and lung, and wherein generating the output comprises generating an output indicative of the presence of a solid tumor in tissue selected from the group.
15. The method according to claim 14, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $752 \pm 4 \text{ cm}^{-1}$ ,  $1030 \pm 4 \text{ cm}^{-1}$ ,  $1046 \pm 4 \text{ cm}^{-1}$ ,  $1128 \pm 4 \text{ cm}^{-1}$ , and  $1237 \pm 4 \text{ cm}^{-1}$ , and wherein generating comprises generating an output indicative of the presence of a tumor in the breast tissue.
16. The method according to claim 14, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $797 \pm 4 \text{ cm}^{-1}$ ,  $830 \pm 4 \text{ cm}^{-1}$ ,  $893 \pm 4 \text{ cm}^{-1}$ ,  $899 \pm 4 \text{ cm}^{-1}$ ,  $1128 \pm 4 \text{ cm}^{-1}$ ,  $1298 \pm 4 \text{ cm}^{-1}$ ,  $1354 \pm 4 \text{ cm}^{-1}$ ,  $1714 \pm 4 \text{ cm}^{-1}$ ,  $1725 \pm 4 \text{ cm}^{-1}$ ,  $1738 \pm 4 \text{ cm}^{-1}$ , and  $3013 \pm 4 \text{ cm}^{-1}$ , and wherein generating comprises generating an output indicative of the presence of a tumor in the gastrointestinal tract tissue.

17. The method according to claim 14, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of: 765  $\pm$  4 cm<sup>-1</sup>, 780  $\pm$  4 cm<sup>-1</sup>, 797  $\pm$  4 cm<sup>-1</sup>, 851  $\pm$  4 cm<sup>-1</sup>, 874  $\pm$  4 cm<sup>-1</sup>, 881  $\pm$  4 cm<sup>-1</sup>, 913  $\pm$  4 cm<sup>-1</sup>, 923  $\pm$  4 cm<sup>-1</sup>, 958  $\pm$  4 cm<sup>-1</sup>, 968,  $\pm$  4 cm<sup>-1</sup>, 1044  $\pm$  4 cm<sup>-1</sup>, 1085  $\pm$  4 cm<sup>-1</sup>, 1191  $\pm$  4 cm<sup>-1</sup>, 1241  $\pm$  4 cm<sup>-1</sup>, 1344  $\pm$  4 cm<sup>-1</sup>, 1373  $\pm$  4 cm<sup>-1</sup>, 1417  $\pm$  4 cm<sup>-1</sup>, 1458  $\pm$  4 cm<sup>-1</sup>, 1469  $\pm$  4 cm<sup>-1</sup>, 1692  $\pm$  4 cm<sup>-1</sup>, 1714  $\pm$  4 cm<sup>-1</sup>, 1728  $\pm$  4 cm<sup>-1</sup>, 2852  $\pm$  4 cm<sup>-1</sup>, and 2984  $\pm$  4 cm, and wherein generating comprises generating an output indicative of the presence of a tumor in the lung tissue.
18. The method according to claim 14, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of: 828  $\pm$  4 cm<sup>-1</sup>, 932  $\pm$  4 cm<sup>-1</sup>, 997  $\pm$  4 cm<sup>-1</sup>, 1059  $\pm$  4 cm<sup>-1</sup>, 1299  $\pm$  4 cm<sup>-1</sup>, 1302  $\pm$  4 cm<sup>-1</sup>, 1403  $\pm$  4 cm<sup>-1</sup>, 1454  $\pm$  4 cm<sup>-1</sup>, 1714  $\pm$  4 cm<sup>-1</sup>, 2979,  $\pm$  4 cm<sup>-1</sup>, and 3013  $\pm$  4 cm<sup>-1</sup>, and wherein generating comprises generating an output indicative of the presence of a tumor in the prostate tissue.
19. A method comprising:  
obtaining an infrared (IR) spectrum of a sample of white blood cells by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid tumor or a pre-malignant condition.
20. The method according to claim 19, wherein generating the output comprises generating the output indicative of the presence of the solid tumor.
21. The method according to claim 19, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.
22. The method according to claim 21, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

23. A method for monitoring the effect of an anti-cancer treatment on a subject undergoing anti-cancer treatment for a solid tumor, for use with a first Peripheral Blood Mononuclear Cells (PBMC) sample separated from blood of the subject that was obtained prior to initiation of the treatment and a second PBMC sample separated from blood of the subject that was obtained  
5 after initiation of the treatment, the method comprising:  
    obtaining IR spectra of the first and second PBMC samples by analyzing the first and second PBMC samples by IR spectroscopy; and  
    based on the IR spectra, generating an output indicative of the effect of the treatment.
24. The method according to claim 23, wherein analyzing the first and second PBMC  
10 samples by IR spectroscopy comprises analyzing the samples by Fourier Transformed Infrared spectroscopy, and wherein obtaining the IR spectra comprises obtaining Fourier Transformed Infrared (FTIR) spectra.
25. The method according to claim 24, wherein analyzing the first and second PBMC  
15 samples by infrared (IR) spectroscopy comprises analyzing the first and second PBMC samples by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).
26. The method according to claim 23, further comprising obtaining an IR spectrum of a third PBMC sample separated from blood of the subject that was obtained following termination of the treatment, by analyzing the third PBMC sample by IR spectroscopy.
27. The method according to claim 3, wherein analyzing comprises assessing a characteristic  
20 of the sample at at least one wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $798 \pm 4 \text{ cm}^{-1}$ ,  $809 \pm 4 \text{ cm}^{-1}$ ,  $814 \pm 4 \text{ cm}^{-1}$ ,  $875 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1001 \pm 4 \text{ cm}^{-1}$ ,  $1015 \pm 4 \text{ cm}^{-1}$ ,  $1103 \pm 4 \text{ cm}^{-1}$ ,  $1118 \pm 4 \text{ cm}^{-1}$ ,  $1162 \pm 4 \text{ cm}^{-1}$ ,  $1221 \pm 4 \text{ cm}^{-1}$ ,  $1270 \pm 4 \text{ cm}^{-1}$ ,  $1283 \pm 4 \text{ cm}^{-1}$ ,  $1295 \pm 4 \text{ cm}^{-1}$ ,  $1315 \pm 4 \text{ cm}^{-1}$ ,  $1341 \pm 4 \text{ cm}^{-1}$ ,  $1367 \pm 4 \text{ cm}^{-1}$ ,  $1392 \pm 4 \text{ cm}^{-1}$ ,  $1429 \pm 4 \text{ cm}^{-1}$ ,  $1440 \pm 4 \text{ cm}^{-1}$ ,  $1445 \pm 4 \text{ cm}^{-1}$ , and  $1455 \pm 4 \text{ cm}^{-1}$ .
28. The method according to claim 27, wherein analyzing comprises assessing the  
25 characteristic at at least two wavenumbers selected from the group.
29. The method according to claim 27, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.

30. A method comprising:  
obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid  
5 tumor or a pre-malignant condition.
31. The method according to claim 30, wherein generating the output comprises generating the output indicative of the presence of the solid tumor.
32. A system for diagnosing a solid tumor, comprising:  
a data processor, configured to analyze an infrared (IR) spectrum of a Peripheral Blood  
10 Mononuclear Cells (PBMC) sample of a subject; and  
an output unit, configured to generate an output indicative of the presence of a solid tumor, based on the infrared (IR) spectrum.
33. The system according to claim 32, wherein the data processor is configured to calculate a second derivative of the infrared (IR) spectrum of the PBMC sample and, based on the second  
15 derivative of the infrared (IR) spectrum, to generate an output indicative of the presence of a solid tumor.
34. The system according to claim 33, wherein the IR spectrum includes a Fourier Transformed Infrared (FTIR) spectrum, and wherein the data processor is configured to calculate a second derivative of the FTIR spectrum.
35. The system according to claim 32, wherein the data processor is configured to analyze the infrared (IR) spectrum by assessing a characteristic of the PBMC sample at at least one  
20 wavenumber selected from the group consisting of:  $765 \pm 4 \text{ cm}^{-1}$ ,  $798 \pm 4 \text{ cm}^{-1}$ ,  $809 \pm 4 \text{ cm}^{-1}$ ,  $814 \pm 4 \text{ cm}^{-1}$ ,  $875 \pm 4 \text{ cm}^{-1}$ ,  $997 \pm 4 \text{ cm}^{-1}$ ,  $1001 \pm 4 \text{ cm}^{-1}$ ,  $1015 \pm 4 \text{ cm}^{-1}$ ,  $1103 \pm 4 \text{ cm}^{-1}$ ,  $1118 \pm 4 \text{ cm}^{-1}$ ,  $1162 \pm 4 \text{ cm}^{-1}$ ,  $1221 \pm 4 \text{ cm}^{-1}$ ,  $1270 \pm 4 \text{ cm}^{-1}$ ,  $1283 \pm 4 \text{ cm}^{-1}$ ,  $1295 \pm 4 \text{ cm}^{-1}$ ,  $1315 \pm 4 \text{ cm}^{-1}$ ,  $1341 \pm 4 \text{ cm}^{-1}$ ,  $1367 \pm 4 \text{ cm}^{-1}$ ,  $1392 \pm 4 \text{ cm}^{-1}$ ,  $1429 \pm 4 \text{ cm}^{-1}$ ,  $1440 \pm 4 \text{ cm}^{-1}$ ,  $1445 \pm 4 \text{ cm}^{-1}$ , and  $1455 \pm 4 \text{ cm}^{-1}$ .
36. The system according to claim 35, wherein the data processor is configured to analyze  
25 the infrared (IR) spectrum by assessing the characteristic at at least two wavenumbers selected from the group.

37. The system according to claim 35, wherein the data processor is configured to analyze the infrared (IR) spectrum by assessing the characteristic at at least three wavenumbers selected from the group.
38. A computer program product for administering processing of a body of data, the product comprising a computer-readable medium having program instructions embodied therein, which instructions, when read by a computer, cause the computer to:
- obtain an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) by analyzing the PBMC by infrared spectroscopy; and
- based on the infrared spectrum, generate an output indicative of the presence of a solid tumor.
39. A method comprising:
- obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and
- based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in a breast tissue of a subject.
40. The method according to claim 39, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $752 \pm 4 \text{ cm}^{-1}$ ,  $1030 \pm 4 \text{ cm}^{-1}$ ,  $1046 \pm 4 \text{ cm}^{-1}$ ,  $1128 \pm 4 \text{ cm}^{-1}$ , and  $1237 \pm 4 \text{ cm}^{-1}$ .
41. The method according to claim 40, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.
42. The method according to claim 40, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.
43. The method according to claim 39, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.

44. The method according to claim 43, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).
45. A method comprising:  
5 obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in tissue of a gastrointestinal tract of a subject.
46. The method according to claim 45, wherein analyzing comprises assessing a  
10 characteristic of the sample at at least one wavenumber selected from the group consisting of: 797  $\pm$  4 cm<sup>-1</sup>, 830  $\pm$  4 cm<sup>-1</sup>, 893  $\pm$  4 cm<sup>-1</sup>, 899  $\pm$  4 cm<sup>-1</sup>, 1128  $\pm$  4 cm<sup>-1</sup>, 1298  $\pm$  4 cm<sup>-1</sup>, 1354  $\pm$  4 cm<sup>-1</sup>, 1714  $\pm$  4 cm<sup>-1</sup>, 1725  $\pm$  4 cm<sup>-1</sup>, 1738,  $\pm$  4 cm<sup>-1</sup>, and 3013  $\pm$  4 cm<sup>-1</sup>.
47. The method according to claim 46, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.
- 15 48. The method according to claim 46, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.
49. The method according to claim 45, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier  
20 Transformed Infrared (FTIR) spectrum.
50. The method according to claim 49, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).
51. A method comprising:  
25 obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in lung tissue of a subject.

52. The method according to claim 51, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of: 765 ± 4 cm<sup>-1</sup>, 780 ± 4 cm<sup>-1</sup>, 797 ± 4 cm<sup>-1</sup>, 851 ± 4 cm<sup>-1</sup>, 874 ± 4 cm<sup>-1</sup>, 881 ± 4 cm<sup>-1</sup>, 913 ± 4 cm<sup>-1</sup>, 923 ± 4 cm<sup>-1</sup>, 958 ± 4 cm<sup>-1</sup>, 968, ± 4 cm<sup>-1</sup>, 1044 ± 4 cm<sup>-1</sup>, 1085 ± 4 cm<sup>-1</sup>, 1191 ± 4 cm<sup>-1</sup>, 1241 ± 4 cm<sup>-1</sup>, 1344 ± 4 cm<sup>-1</sup>, 1373 ± 4 cm<sup>-1</sup>, 1417 ± 4 cm<sup>-1</sup>, 1458 ± 4 cm<sup>-1</sup>, 1469 ± 4 cm<sup>-1</sup>, 1692 ± 4 cm<sup>-1</sup>, 1714 ± 4 cm<sup>-1</sup>, 1728 ± 4 cm<sup>-1</sup>, 2852 ± 4 cm<sup>-1</sup>, and 2984 ± 4 cm.

53. The method according to claim 52, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.

54. The method according to claim 52, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.

55. The method according to claim 51, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.

56. The method according to claim 55, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

57. A method comprising:  
obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of the presence of a solid tumor in a prostate tissue of a subject.

58. The method according to claim 57, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of: 828 ± 4 cm<sup>-1</sup>, 932 ± 4 cm<sup>-1</sup>, 997 ± 4 cm<sup>-1</sup>, 1059 ± 4 cm<sup>-1</sup>, 1299 ± 4 cm<sup>-1</sup>, 1302 ± 4 cm<sup>-1</sup>, 1403 ± 4 cm<sup>-1</sup>, 1454 ± 4 cm<sup>-1</sup>, 1714 ± 4 cm<sup>-1</sup>, 2979, ± 4 cm<sup>-1</sup>, and 3013 ± 4 cm<sup>-1</sup>.

59. The method according to claim 58, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.

60. The method according to claim 58, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.
61. The method according to claim 57, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.
62. The method according to claim 61, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).
63. A method comprising:  
obtaining an infrared (IR) spectrum of a Peripheral Blood Mononuclear Cells (PBMC) sample from a cancer patient by analyzing the sample by infrared spectroscopy; and  
based on the infrared spectrum, generating an output indicative of a stage of the cancer.
64. The method according to claim 63, wherein analyzing comprises assessing a characteristic of the sample at at least one wavenumber selected from the group consisting of:  $865 \pm 4 \text{ cm}^{-1}$ ,  $897 \pm 4 \text{ cm}^{-1}$ ,  $924 \pm 4 \text{ cm}^{-1}$ ,  $1030 \pm 4 \text{ cm}^{-1}$ ,  $1047 \pm 4 \text{ cm}^{-1}$ ,  $1191 \pm 4 \text{ cm}^{-1}$ , and  $1238 \pm 4 \text{ cm}^{-1}$ .
65. The method according to claim 64, wherein analyzing comprises assessing the characteristic at at least two wavenumbers selected from the group.
66. The method according to claim 64, wherein analyzing comprises assessing the characteristic at at least three wavenumbers selected from the group.
67. The method according to claim 63, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared (FTIR) spectroscopy, and wherein obtaining the infrared (IR) spectrum comprises obtaining a Fourier Transformed Infrared (FTIR) spectrum.
68. The method according to claim 67, wherein analyzing the sample by infrared (IR) spectroscopy comprises analyzing the sample by Fourier Transformed Infrared microspectroscopy (FTIR-MSP).

FIG. 1A






FIG. 1C








FIG. 2C







FIG. 3C

Wavenumber ( $\text{cm}^{-1}$ )	Breast Cancer	Lung Cancer	Head & Neck Cancer	Pancreas Cancer	Esophagus Cancer
2871	↓	↓			
2850	↓	↓		↓	
1390	↓	↓	↓		
1369		↓			
1354				↑	
1346		↓		↑	
1317		↑		↓	↑
1311		↑			
1286					↑
1271	↑	↑		↑	↑
1240	↓	↓	↓		
1223	↓	↓			
1173	↓			↑	
1155	↓	↓	↑		↑
1126		↑		↑	
1101	↓	↓	↓		↓
1086	↓		↑	↓	↓
1049	↓			↓	
1047	↓			↓	↓
1024			↑	↓	↑
995	↓				
970	↓	↑	↑	↑	↓
930				↑	
925					↑
918			↓	↑	
893		↑			
876			↑		
852				↑	↑
839		↑		↓	↓
822	↑			↑	
810		↑		↑	
795	↓				
779	↓		↓		↓
740	↓	↓	↓		

FIG. 4




FIG. 5A





FIG. 5B



9/13

FIG. 5C




FIG. 5D




FIG. 5E





FIG. 5F



11/13

FIG. 5G




FIG. 6A




FIG. 6B




FIG. 6C

