



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2727156 A1 2009/12/10

(21) **2 727 156**

(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2009/06/02
(87) Date publication PCT/PCT Publication Date: 2009/12/10
(85) Entrée phase nationale/National Entry: 2010/12/06
(86) N° demande PCT/PCT Application No.: NL 2009/050301
(87) N° publication PCT/PCT Publication No.: 2009/148306
(30) Priorité/Priority: 2008/06/06 (EP08157758.7)

(51) Cl.Int./Int.Cl. *G21G 1/06* (2006.01)

(71) **Demandeur/Applicant:**
TECHNISCHE UNIVERSITEIT DELFT, NL

(72) **Inventeurs/Inventors:**
WOLTERBEEK, HUBERT THEODOOR, NL;
BODE, PETER, NL

(74) **Agent:** BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : PROCEDE POUR LA PRODUCTION DE ⁹⁹MO SANS SUPPORT AJOUTÉ
(54) Title: A PROCESS FOR THE PRODUCTION OF NO-CARRIER ADDED ⁹⁹MO

(57) Abrégé/Abstract:

The present invention relates to a process for the production of no-carrier added ⁹⁹Mo by neutron activation of ⁹⁸Mo thereby reaching specific radioactivity which allow the use of such produced ⁹⁹Mo as an option for the ⁹⁹Mo produced by the fission of ²³⁵U. This has been achieved by taking advantage of the recoil of the ⁹⁹Mo nuclei upon the capture of neutrons by the ⁹⁸Mo containing compound. These recoiled nuclei are no longer chemically bound to the ⁹⁸Mo containing compound allowing further specific separation. Preferred ⁹⁸Mo containing compounds are molybdenum(0)hexacarbonyl [(Mo (CO) ₆] and molybdenum (VI) di oxodioxinate [C₄H₃ (O) -NC₅H₃]₂-MoO₂.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 December 2009 (10.12.2009)(10) International Publication Number
WO 2009/148306 A1(51) International Patent Classification:
G21G 1/06 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/NL2009/050301(22) International Filing Date:
2 June 2009 (02.06.2009)(25) Filing Language:
English(26) Publication Language:
English(30) Priority Data:
08157758.7 6 June 2008 (06.06.2008) EP(71) Applicant (for all designated States except US): **TECHNISCHE UNIVERSITEIT DELFT** [NL/NL]; Stevinweg 1, NL-2628 CN Delft (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WOLTERBEEK, Hubert Theodoor** [NL/NL]; Stompwijkseweg 57B, NL-2266 GD Leidschendam-Voorburg (NL). **BODE, Peter** [NL/NL]; Zwartsluisje 37, NL-3284 LK Zuid-Beijerland-Korendijk (NL).(74) Agent: **Kupecz, Árpád**; Weteringschans 96, NL-1017 XS Amsterdam (NL).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2009/148306 A1

(54) Title: A PROCESS FOR THE PRODUCTION OF NO-CARRIER ADDED ^{99}MO (57) Abstract: The present invention relates to a process for the production of no-carrier added ^{99}Mo by neutron activation of ^{98}Mo thereby reaching specific radioactivity which allow the use of such produced ^{99}Mo as an option for the ^{99}Mo produced by the fission of ^{235}U . This has been achieved by taking advantage of the recoil of the ^{99}Mo nuclei upon the capture of neutrons by the ^{98}Mo containing compound. These recoiled nuclei are no longer chemically bound to the ^{98}Mo containing compound allowing further specific separation. Preferred ^{98}Mo containing compounds are molybdenum(0)hexacarbonyl [(Mo (CO) ₆] and molybdenum (VI) di oxodioxinate [C₄H₃ (O) -NC₅H₃] ₂-MoO₂.

A process for the production of no-carrier added ^{99}Mo

The present invention relates to a process for the production of no-carrier added ^{99}Mo .

According to the current practice, ^{99}Mo with high specific radioactivity is produced by fission of fissile actinide targets (^{233}U , ^{235}U , ^{239}Pu etc), mostly using ^{235}U , wherein ^{99}Mo is one of the fission products of high yield (ca. 6%). However, next to this ^{99}Mo a range of other further fission products are produced as well. The consequence of this production route is that the production requires handling of nuclear fuel, wherein ^{99}Mo has to be isolated and purified from the other fission products. Furthermore, the prior art process involves a final storage of the co-produced additional fission products. This total implicates that only few production sites of ^{99}Mo exist with the required production licenses. In turn, this makes that the world-production of ^{99}Mo - $^{99\text{m}}\text{Tc}$ generators (used in medical radio-imaging) is based on only a very few sites, wherein any problem in one of the current sites immediately endangers the continuity of the necessary supply.

Now the present invention aims to provide a process for the production of ^{99}Mo of high specific radioactivity, wherein the above-mentioned disadvantages are removed.

The present invention enables the production of no-carrier added ^{99}Mo by neutron activation of ^{98}Mo , thereby achieving specific radioactivity which allows the use of such produced ^{99}Mo as a favorable option (alternative) for the ^{99}Mo production by means of the fission of ^{235}U . This high specific radioactivity is obtained according to the invention by taking advantage of the recoil of the ^{99}Mo nuclei upon the capture of neutrons by the ^{98}Mo containing nuclei. The mentioned recoiled nuclei are no longer chemically bound to the target matrix and thus allow for specific separation.

Accordingly the present invention relates to a process for the production of no-carrier added ^{99}Mo of high specific radioactivity, characterized in that an ^{98}Mo containing chemical compound is bombarded with neutrons and the

resulting ^{99}Mo radioactivity which is incorporated in said compound is separated.

It has been surprisingly found that by bombarding ^{98}Mo containing chemical compound with neutrons, ^{99}Mo with high specific radioactivity may be obtained without the disadvantages of the prior art fission of ^{235}U . Obviously, next to ^{99}Mo no additional fission products are formed.

There are two options for the process for the present invention.

According to the first option, said ^{99}Mo radioactivity, incorporated in said compound, is a) transferred into a liquid in which only the produced ^{99}Mo dissolves, or b) transferred into a liquid in which said compound has a high solubility which liquid is mixed with a second liquid wherein said compound does not dissolve and the "loose" ^{99}Mo nuclei are transferred into said second liquid phase.

Thus, after bombarding the ^{98}Mo containing chemical compound with neutrons the produced ^{99}Mo radioactivity incorporated in said compound is transferred into a liquid in which only the produced ^{99}Mo dissolves or into a first liquid having a high solubility for said compound having ^{99}Mo radioactivity.

Said first liquid is mixed with a second liquid, wherein the "loose" ^{99}Mo nuclei are transferred by extraction into a second liquid phase, wherein the compound does not dissolve.

Preferred ^{98}Mo containing compounds are molybdenum(0)hexacarbonyl [$(\text{Mo}(\text{CO})_6)$] and molybdenum(VI)dioxo-dioxinate [$\text{C}_4\text{H}_3(\text{O})-\text{NC}_5\text{H}_3\right]_2\text{-MoO}_2$.

Next to these preferred molybdenum compounds the following molybdenum compounds may be used.

Cycloheptatrienemolybdenum(0)tricarbonyl [$(\text{C}_7\text{H}_8)\text{Mo}(\text{CO})_3$],

d. purple cryst. powder (Across Organics);

Molybdenum(0)hexacarbonyl [$(\text{Mo}(\text{CO})_6)$], white, crystalline powder (Across Organics);

Methylcyclopentadienylmolybdenum(I)tricarbonyl, dimer [$(\text{CH}_3)_2-(\text{C}_5\text{H}_5)\right]_2\text{-Mo}_2(\text{CO})_6$ d. purple, crystalline powder (Across Organics);

Propylcyclopentadienylmolybdenum(I)tricarbonyl, dimer

$[\text{CH}_3\text{CH}_2\text{CH}_2 - (\text{C}_5\text{H}_5)]_2\text{-Mo}_2(\text{CO})_6$ d. brown, crystall. Powder (Across Organics);

Cyclopentadienylmolybdenum(II) tricarbonyl dimer $[(\text{C}_5\text{H}_5) -$

5 $\text{Mo}(\text{CO})_3]_2$, d. purple cryst. powder (Across Organics);

Pentamethylcyclopentadienyl-molybdenum(V) dicarbonyl dimer

$[(\text{CH}_3)_5 - (\text{C}_5\text{H}_5) - \text{Mo}(\text{CO})_2]_2$ olive-green crystalline powder;

Molybdenum(VI) dioxo-Bis(acetylacetonato) $[(\text{CH}_3\text{COCH}=\text{C}(\text{O}-)$
 $\text{CH}_3)]_2\text{-MoO}_2$, white, cristalline powder (Sigma Aldrich, USA).

10 **Molybdenum(VI) dioxo-dioxinate** $[(\text{C}_4\text{H}_3(\text{O}) - \text{NC}_5\text{H}_3)]_2\text{-MoO}_2$, orange-yellow cristalline powder, was synthesized according to the method as described in Vogel et.al. [xxx].

Molybdenum(IV) disulfide $[\text{MoS}_2]$, d. grey powder, 325 Mesh (Across Organics);

15 **Molybdenum disilicide** $[\text{MoSi}_2]$, d. grey powder, 325 Mesh (Alfa Aesar GmbH, Karlsruhe, Germany) ;

Molybdenum nanoparticles (~ 100 nm), d. grey powder, (Johnson & Matthey, USA)

Potassium molybdenum(VI)-hexacyanoferrate $[\text{KMo}[\text{Fe}^{\text{III}}(\text{CN})_6]]$, d.

20 brown, crystalline powder was synthesized according to the method as described by Sebesta et. al. [yy]

Preferred first liquid is an organic solvent dichloromethane (CH_2Cl_2), whereas the second preferred liquid is an aqueous phase of different pH (2-12) prepared in 50 mM ammonium acetate buffer.

Other suitable first liquids are chloroform (CH_3Cl), benzene (C_6H_6), toluene ($\text{CH}_3-\text{C}_6\text{H}_5$).

Other suitable second liquids are aqueous solutions of acidic solution HCl (0.05 M), alkaline solution NaOH (0.05 M), chelating solutions Na_2EDTA (0.05 M), $\text{Na}_3\text{citrate}$ (0.05 M), oxidizing solution H_2O_2 (0.02 M) in HCl (0.05 M), reducing solution (NaHSO_3 (0.05 M), saline solution NaCl (0.9% w/w), neutral buffer solution NH_4Ac (0.05 M ; pH 7.3)).

According to a second optional variant of claim 1, a 35 ^{98}Mo containing compound is transferred into an irradiation container containing 1) a liquid in which only the produced ^{99}Mo dissolves, or 2) a liquid in which the compound dissolves, as well as the liquid (non-mixable with the first liquid) in which the ^{99}Mo dissolves and the compound does not

dissolve, the container is, under continuous shaking, irradiated with neutrons in an external neutron beam, resulting in transfer of the recoiled ^{99}Mo on-line from one to another liquid phase.

5 Also by using of this variant, the disadvantages of the prior art fission process are removed.

It is noted that the present process is new and not obvious over the current techniques, because the current techniques did not significantly increase the molybdenum 10 specific radioactivity due to non-suited Mo-compounds and/or non-suited extraction protocols. The prior art technique predominantly by fission of nuclear fuel (^{235}U) was until now used worldwide for large-scaled production of no-carrier added ^{99}Mo with the disadvantages as mentioned herein before.

15 It is noted that the recoil-production of ^{99}Mo leads to ^{99}Mo with the required high specific radioactivity without the otherwise obligatory processing of nuclear fuel accompanied by the disadvantages as mentioned before.

Furthermore, it is noted that currently there are no 20 production options other than by the fission-produced ^{99}Mo which lead to comparable specific radioactivity. Because for the fission-produced ^{99}Mo , production facilities should process nuclear fuel and only a small number of facilities worldwide have the required licenses as mentioned before. The 25 proposed production by recoil- ^{99}Mo from neutron irradiation of enriched ^{98}Mo targets implies that many more facilities worldwide could start up the production of high radioactivity ^{99}Mo .

Although the principle of the recoil (Szilard-Chalmers) reactions is known, it is surprising that by using 30 the right chemical compounds and experimental conditions, such as the availability of a neutron beam of adequate density ^{99}Mo of high specific radioactivity may be obtained by the invention. Therefore, the present process is not only new, but also inventive over the current ^{99}Mo producing technique.

Further, there are no disadvantages of the present process apart from the necessary entrance to a neutron source coupled to a radiochemical infrastructure.

Further to the process options according to claims 2

and 6, it is remarked that according to claim 2 the bombardment of the ^{98}Mo chemical compound with neutrons occurs in the reactor, whereas according to option disclosing claim 6 the bombardment occurs outside the reactor in a neutron beam.

5 It is noted that the present process is not limited to the production of ^{99}Mo but it may be used for other products which at the moment are mainly produced through the ^{235}U fission process.

10 The process of the invention is also suitable for the production of $^{90}\text{Sr} \rightarrow ^{90}\text{Y}$; $^{103}\text{Ru} \rightarrow ^{103m}\text{Ru}$; $^{132}\text{Te} \rightarrow ^{132}\text{I}$; $^{137}\text{Cs} \rightarrow ^{137m}\text{Ba}$ and $^{140}\text{Ba} \rightarrow ^{140}\text{La}$.

The invention will be further explained by means of the following examples.

15 EXPERIMENTAL approach 1

Szilard Chalmers reaction:

Example 1. Irradiation of the molybdenum complexes

20 - 1500 mg of Mo(0)hexacarbonyl and Mo(VI)dioxo-
20 dioxinate were sealed in a polyethylene capsule, and irradiated via the pneumatic facility in the Hoger Onderwijs Reactor of Delft University of Technology, having a neutron fluence rate of $5.0 \times 10^{12} \text{ cm}^{-2} \cdot \text{s}^{-1}$ for a suitable length of time (15 minutes to 5 hours). Some of the irradiations were also
25 carried out in the in-core radiation facility, which has a considerable higher fluence rate ($2.4 \times 10^{13} \text{ cm}^{-2} \cdot \text{s}^{-1}$), but a different neutron fluence rate profile (ratio of thermal to fast neutron fluence rates) compared to pneumatic facility. In the case of short irradiations (15 - 30 minutes), the
30 radiochemical separation of ^{99}Mo was carried out 1 h after the end of irradiation, while in the case of longer irradiations, the separation was carried out 2 hours after the end of irradiation so as to allow the decay of shorter ^{101}Mo and ^{101}Tc with shorter half lives.

35

Example 2. Liquid-Liquid extraction of organomolybdenum targets

After irradiation the target was dissolved in 50ml of purified organic liquid (dichloromethane (CH_2Cl_2), chloro-

form (CH_3Cl), benzene (C_6H_6), toluene ($\text{CH}_3\text{-C}_6\text{H}_5$)). 2.0 ml aliquots from the stock solution were contacted with equal volumes of aqueous phase of different pH (2 - 12), prepared in 50mM ammonium acetate buffer. The pH of the buffer solutions was maintained by adding dilute acetic acid or ammonia solutions. Further, the following aqueous solutions were used: acidic solution HCl (0.05 M), alkaline solution NaOH (0.05 M), chelating solutions Na_2EDTA (0.05 M), $\text{Na}_3\text{citrate}$ (0.05 M), oxidizing solution H_2O_2 (0.02 M) in HCl (0.05 M), reducing solution (NaHSO_3 (0.05 M), saline solution NaCl (0.9% w/w), neutral buffer solution NH_4Ac (0.05 M ; pH 7.3). Experiments were also carried out with MilliQ water as aqueous phase. Kinetic studies on the solvent extraction of molybdenum from the organic solution into ammonium acetate. Experiments were also carried out with MilliQ water as aqueous phase. Kinetic studies on the solvent extraction of ^{99}Mo from dichloromethane into ammonium acetate buffer solution were carried out to optimize the time of equilibration for subsequent studies. In this experiment the samples were removed from the roller-bed at different time intervals ranging from 5 minutes to one hour. It was observed that the extraction yield of ^{99}Mo reached a constant value after 15 minutes, while that of total molybdenum increased up to 30 minutes of shaking time. Thus the highest enrichment factor was obtained for a shaking time of 15 minutes. In view of this the subsequent extractions were carried out with a shaking time of 15 minutes (Tomar et.al. ,2008). After shaking the solutions for 15 minutes, the samples were centrifuged at 3000rpm (Jouan) for 5 minutes to obtain clear separation of phases. Subsequently 1.0 mL aliquots from the aqueous layer were taken for measurement of the ^{99}Mo radioactivity by gamma counting as well as determination of total molybdenum concentration. In the case of the dichloromethane stock solution, 0.2 mL aliquots ($n = 3$) were first treated with *aqua regia* (3x 1.0 mL concentrated HCl, plus 1x 1.0 mL concentrated HNO_3) which after gamma counting were diluted up to 10mL for determination of total Mo content (ICP-OES).

Example 3. Analysis

The ^{99}Mo radioactivities of the organic phase, the aqueous phases and the dichloromethane-Mo stock solution were measured as follows:

5 The gamma-ray spectrometric measurement was carried out using a shielded well type NaI(Tl) counter coupled to a 2048 multichannel pulse height analyzer (Wallac). The peak at 140 keV due to $^{99\text{m}}\text{Tc}$ was used as an indication for the radioactivity of ^{99}Mo . Counting of the samples was carried out 24

10 hours after the radiochemical separation so as to obtain equilibrium between $^{99\text{m}}\text{Tc}$ and ^{99}Mo . The net peak area of 140 keV was obtained by linear subtraction of Compton background. The counting time was adjusted so as to obtain at least 10000 counts under the 140 keV peak.

15 The total molybdenum concentration in the aqueous samples as well as the aqua regia destructed dichloromethane stock solutions were measured using Inductively Coupled Plasma Optical Emission Spectrometer (Perkin Elmer ICP-OES 4300DV). The emission lines at 202.031 nm, 203.845 nm and

20 204.597 nm were used for the measurement of molybdenum concentration. The instrument was calibrated for Molybdenum using a ICP-OES standard solution (Merck, Ultrapure 1.000 g Mo.L $^{-1}$), which was suitably diluted to obtain standard solutions in the range of 0.05 to 2.5 $\mu\text{g.mL}^{-1}$ Mo.

25 The specific radioactivity of ^{99}Mo (expressed in cpm/mg total Mo) in the aqueous phase and the stock solution was obtained from the ratio of the gamma activity and total Mo concentration. The enrichment factor was calculated as the ratio of specific activity of ^{99}Mo in the separated aqueous

30 phase to that in the organic phase.

EXPERIMENTAL approach 2

Example 4

35 This experimental approach is based on the same chemical principles as the first approach. However, the liquid-liquid extraction is now performed simultaneously with the neutron bombardment. After completion of the irradiation/liquid-liquid extraction, the entire solution is proc-

essed in the same way as described in the above.

In this approach, benzene or toluene are the preferred phases for dissolution of the Mo compound since irradiation of dichloromethane or chloroform results in production of a very high and unpractical ^{38}Cl radioactivity besides intense high energy prompt gamma-radiation during the irradiation.

The advantage of the neutron beam irradiation is that the compound is exposed to a considerable smaller associated gamma-ray dose than during the irradiation 'in' the reactor. The gamma-radiation (resulting from the fission processes in the reactor) has, to some extent, a reverse effect to the recoil process (described as 'annealing'). Another advantage is that also compounds may be considered risky for reactor irradiation because of possible chemical decomposition and formation of gaseous compounds which is unwanted for safety considerations. Such effects are almost negligible during beam irradiation and impose risks of a considerable smaller extent.

A disadvantage of the neutron beam irradiation is the lower neutron intensity and therefore the lower ^{99}Mo yield.

Examples 1, 2 and 3 relate to option according to claim 2 and example 4 relates to option according to claim 6.

It should be noted that the invention is not limited to the above-mentioned disclosure, examples or the claims.

CLAIMS

1. A process for the production of no-carrier added ⁹⁹Mo of high specific radioactivity, **characterized** in that an ⁹⁸Mo containing chemical compound is bombarded with neutrons and the resulting ⁹⁹Mo radioactivity which is incorporated in 5 said compound is separated.

2. The process of claim 1, **characterized** in that said ⁹⁹Mo radioactivity, incorporated in said compound, is transferred a) into a liquid in which only the produced ⁹⁹Mo dissolves, or b) transferred into a first liquid, in which 10 said compound has a high solubility which liquid is mixed with a second liquid wherein said compound does not dissolve and the "loose" ⁹⁹Mo nuclei are transferred into said second liquid phase and removed.

3. The process of claim 1 or 2, **characterized** in 15 that said ⁹⁸Mo containing chemical compound is molybdenum(0)hexacarbonyl[(Mo(CO)₆] or molybdenum(VI)dioxo-dioxinate[C₄H₃(O)-NC₅H₃]₂-MoO₂.

4. The process of claims 1-3, **characterized** in that 20 said first liquid is dichloromethane.

5. The process of claims 1-3, **characterized** in that the second liquid is an aqueous phase of different pH (2-12) prepared in 50 mM ammonium acetate buffer.

6. The process of claims 1 and 3-5, **characterized** in 25 that a non-dissolvable ⁹⁸Mo containing compound is transferred into an irradiation container 1) containing the liquid in which only the produced ⁹⁹Mo dissolves, or 2) containing both the liquid in which the compound does dissolve, as well as the liquid in which only the ⁹⁹Mo dissolves, the container is, under continuous shaking, irradiated with neutrons in an 30 external neutron beam, resulting in transfer of the recoiled ⁹⁹Mo on-line from one to another liquid phase.

7. The process of claim 6, **characterized** in that the ⁹⁸Mo containing chemical compound is as defined in claim 3.