



(43) International Publication Date
20 September 2012 (20.09.2012)

(10) International Publication Number
WO 2012/125724 A1

- (51) **International Patent Classification:**
C07C 57/00 (2006.01)
- (21) **International Application Number:**
PCT/US2012/029077
- (22) **International Filing Date:**
14 March 2012 (14.03.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/452,357 14 March 2011 (14.03.2011) US
- (71) **Applicant (for all designated States except US):** **BETH ISRAEL DEACONESS MEDICAL CENTER** [US/US]; 330 Brookline Avenue, Boston, MA 02215 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **LU, Kun, Ping** [US/US]; 68 Maplewood Avenue, Newton, MA 02459 (US). **WEI, Shuo** [—/US]; 134 Thornton Road, Chestnut Hill, MA 02467 (US). **ZHOU, Xiao, Zhen** [US/US]; 68 Maplewood Avenue, Newton, MA 02459 (US).
- (74) **Agents:** **BIEKER-BRADY, Kristina** et al.; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** METHODS AND COMPOSITIONS FOR THE TREATMENT OF PROLIFERATIVE DISORDERS

(57) **Abstract:** The invention features methods of treating a proliferative disorder characterized by elevated Pin1 marker levels and/or reduced Pin1 Ser71 phosphorylation in a subject by administering a retinoic acid compound. Additionally, the invention features methods of treating proliferative disorders (e.g., proliferative disorders characterized by elevated Pin1 marker levels) by administering a retinoic acid compound in combination with another anti-proliferative compound. Finally, the invention also features methods including high-throughput screens for discovering and validating Pin1 inhibitors.



WO 2012/125724 A1

METHODS AND COMPOSITIONS FOR THE TREATMENT OF PROLIFERATIVE DISORDERS

Statement as to Federally Sponsored Research

5 This invention was made with government support under grant numbers NIH GM058556, NIH AG017870, and NIH CA122434. The government has certain rights in the invention.

Field of the Invention

10 In general, this invention relates to the treatment of proliferative disorders (e.g., proliferative disorders characterized by elevated Pin1 marker levels) with retinoic acid compounds.

Background of the Invention

15 The increased number of cancer cases reported in the United States, and, indeed, around the world, is a major concern. Currently there are only a handful of detection and treatment methods available for some specific types of cancer, and these provide no absolute guarantee of success. In order to be most effective, these treatments require not only an early detection of the malignancy, but a reliable assessment of the severity of the malignancy.

20 It is apparent that the complex process of tumor development and growth must involve multiple gene products. It is therefore important to define the role of specific genes involved in tumor development and growth and identify those genes and gene products that can serve as targets for the diagnosis, prevention, and treatment of cancers.

25 In the realm of cancer therapy, it often happens that a therapeutic agent that is initially effective for a given patient becomes, over time, ineffective or less effective for that patient. The very same therapeutic agent may continue to be effective over a long period of time for a different patient. Further, a therapeutic agent that is effective, at least initially, for some patients can be completely ineffective or even harmful for other patients. Accordingly, it would be useful to identify genes and/or gene products that represent prognostic genes with respect to a given therapeutic agent or class of therapeutic agents. It then may be possible to determine which patients will benefit from particular therapeutic regimen and, importantly, determine when, if ever, the therapeutic regime begins to lose its effectiveness for a given patient. The ability to make such predictions would make it possible to discontinue a therapeutic regime

30

that has lost its effectiveness well before its loss of effectiveness becomes apparent by conventional measures.

Recent advances in the understanding of molecular mechanisms of oncogenesis have led to exciting new drugs that target specific molecular pathways. These drugs have transformed cancer treatments, especially for those caused by some specific oncogenic events, such as Herceptin for breast cancer, caused by HER2/Neu, and Gleevec for chronic myelogenous leukemia caused by Bcr-Abl. However, it has been increasingly evident that, in many individual tumors, there are a large number of mutated genes that disrupt multiple interactive and/or redundant pathways. Thus, intervening in a single pathway may not be effective. Furthermore, cancer resistance to molecularly targeted drugs can develop through secondary target mutation or compensatory activation of alternative pathways, so-called “oncogenic switching.” Thus, a major challenge remains how to simultaneously inhibit multiple oncogenic pathways either using a combination of multiple drugs, with each acting on a specific pathway, or using a single drug that concurrently blocks multiple pathways. The results disclosed herein suggest that Pin1 inhibitors might have a major impact on treating cancers, especially aggressive and/or drug-resistant cancers.

We and others have shown that Pin1 is prevalently overexpressed in human cancers and that high Pin1 marker levels correlate with poor clinical outcome in many cancers. In contrast, the Pin1 polymorphism that reduces Pin1 expression is associated with reduced cancer risk in humans. Significantly, Pin1 activates at least 19 oncogenes/growth enhancers, including β -catenin, cyclin D1, NF- κ B, c-Jun, c-fos, AKT, A1B1, HER2/Neu, Mcl-1, Notch, Raf-1, Stat3, c-Myb, Hbx, Tax, and v-rel, and also inactivates at least 12 tumor suppressors/growth inhibitors, including PML, SMRT, FOXOs, RAR α , and Smad (Figs 1A and 1B). Whereas Pin1 overexpression causes cell transformation and tumorigenesis, Pin1 knockdown inhibits cancer cell growth in cell cultures and mice. Pin1-null mice are highly resistant to tumorigenesis induced either by oncogenes such as activated Ras or HER2/Neu, or tumor suppressors such as p53. Thus, Pin1 inhibitors might have the desired property to suppress numerous oncogenic pathways simultaneously for treating cancers, especially those aggressive and/or drug-resistant cancers.

Summary of the Invention

In one aspect, the invention provides a method of treating a proliferative disease in a subject by administering a retinoic acid compound (e.g., a deuterated compound) to the subject in an amount sufficient to treat the subject, wherein the subject is determined to have elevated levels of a Pin1 marker (e.g., Ser71 phosphorylation) prior to the administration.

In another aspect, the invention features a method of treating a proliferative disease in a subject by determining Pin1 marker levels (e.g., reduced Ser71 phosphorylation) in a sample (e.g., tumor samples, blood, urine, biopsies, lymph, saliva, phlegm, and pus) from the subject and administering a retinoic acid compound to the subject if sample is determined to have elevated Pin1 marker levels.

5 In any of the foregoing aspects, the method can also include the administration of a second anti-proliferative compound (e.g., at a low dosage) or anti-cancer compound (e.g., an anti-angiogenic compound). The second compound can be administered separately, or in a single formulation with, the retinoic acid compound. The second anti-proliferative compound can be, e.g., MK-2206, ON 013105, RTA 402, BI 2536, Sorafenib, ISIS-STAT3Rx, a microtubule inhibitor, a topoisomerase inhibitor, a
10 platin, an alkylating agent, an anti-metabolite, paclitaxel, gemcitabine, doxorubicin, vinblastine, etoposide, 5-fluorouracil, carboplatin, altretamine, aminoglutethimide, amsacrine, anastrozole, azacitidine, bleomycin, busulfan, carmustine, chlorambucil, 2-chlorodeoxyadenosine, cisplatin, colchicine, cyclophosphamide, cytarabine, cytoxan, dacarbazine, dactinomycin, daunorubicin, docetaxel, estramustine phosphate, floxuridine, fludarabine, gentuzumab, hexamethylmelamine, hydroxyurea,
15 ifosfamide, imatinib, interferon, irinotecan, lomustine, mechlorethamine, melphalen, 6-mercaptopurine, methotrexate, mitomycin, mitotane, mitoxantrone, pentostatin, procarbazine, rituximab, streptozocin, tamoxifen, temozolomide, teniposide, 6-thioguanine, topotecan, trastuzumab, vincristine, vindesine, and/or vinorelbine. Additionally, or alternatively, any of the foregoing methods can include determining Pin1 marker levels in the sample after the administration of a retinoic acid compound.

20 In any of the foregoing aspects, the retinoic acid compound may selected from 13-cis-retinoic acid, all-trans-retinoic acid, retinol, retinyl acetate, retinal, AC-55649, acitretin or any of the compounds listed in Fig. 2A, Fig. 2B, Fig. 10A, Fig. 10D, and Table 1.

The elevated Pin1 marker level of any of the foregoing methods can be due to, e.g., an inherited trait or a somatic mutation.

25 The proliferative disorder of any of the foregoing methods can be, e.g., leukemias, polycythemia vera, lymphomas, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors. Specifically, the proliferative disorder can be e.g., acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic
30 leukemia, chronic lymphocytic leukemia), Hodgkin's disease, non-Hodgkin's disease, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma,

Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

The invention also features a method for identifying a Pin1 ligand including: (i) incubating Pin1 protein with a fluorescently labeled probe (e.g., HF488-FP or FITC-FP labeled peptide probes), forming a Pin1-probe complex; (ii) adding a test compound to the incubation; and (iii) determining whether any substantial portion of the probe is displaced from the Pin1-probe complex by the test compound, such displacement indicating that the test compound is a Pin1 ligand. The invention also features a method for identifying a Pin1 modulating compound including: (i) incubating human-derived cancer cells (e.g., breast cancer cells) that have Neu/Erb2 gene amplification; (ii) applying a test compound to the cell; and (iii) determining whether Neu/Erb2 overexpression is reduced, wherein reduction of Neu/Erb2 overexpression indicates that the test compound is a Pin1 modulating compound.

By the term "proliferative disorder" is meant a disorder characterized by inappropriate accumulation of a cell population in a tissue (e.g., by abnormal cell growth). This inappropriate accumulation may be the result of a genetic or epigenetic variation that occurs in one or more cells of the cell population. This genetic or epigenetic variation causes the cells of the cell population to grow faster, die slower, or differentiate slower than the surrounding, normal tissue. The cell population includes cells of hematopoietic, epithelial, endothelial, or solid tissue origin.

As used herein, the term "abnormal cell growth" is intended to include cell growth which is undesirable or inappropriate. Abnormal cell growth also includes proliferation which is undesirable or inappropriate (e.g., unregulated cell proliferation or undesirably rapid cell proliferation). Abnormal cell growth can be benign and result in benign masses of tissue or cells, or benign tumors. Many art-recognized conditions are associated with such benign masses or benign tumors including diabetic retinopathy, retrolental fibrioplasia, neovascular glaucoma, psoriasis, angiofibromas, rheumatoid arthritis, hemangiomas, and Kaposi's sarcoma. Abnormal cell growth can also be malignant and result in malignancies, malignant masses of tissue or cells, or malignant tumors. Many art-recognized conditions

and disorders are associated with malignancies, malignant masses, and malignant tumors including cancer and carcinoma.

As used herein, the term “tumor” is intended to encompass both in vitro and in vivo tumors that form in any organ of the body. Tumors may be associated with benign abnormal cell growth (e.g., benign tumors) or malignant cell growth (e.g., malignant tumors). The tumors which are described herein are preferably sensitive to the Pin1 inhibitors of the present invention. Examples of the types of tumors intended to be encompassed by the present invention include those tumors associated with breast cancer, skin cancer, bone cancer, prostate cancer, liver cancer, lung cancer, brain cancer, cancer of the larynx, gallbladder, pancreas, rectum, parathyroid, thyroid, adrenal, neural tissue, head and neck, colon, stomach, bronchi, kidneys.

As used herein, the term “Pin1 marker” refers to a marker which is capable of being indicative of Pin1 activity levels in a sample of the invention. Pin1 markers include nucleic acid molecules (e.g., mRNA, DNA) which corresponds to some or all of a Pin1 gene, peptide sequences (e.g., amino acid sequences) which correspond to some or all of a Pin1 protein, nucleic acid sequences which are homologous to Pin1 gene sequences, peptide sequences which are homologous to Pin1 peptide sequences, antibodies to Pin1 protein, substrates of Pin1 protein, binding partners of Pin1 protein, and activity of Pin1.

By “elevated levels of a Pin1 marker” is meant a level of Pin1 marker that is altered thereby indicating elevated Pin1 activity. “Elevated levels of a Pin1 marker” include levels at least 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 500%, 1000%, or greater than, or 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% less than the marker levels measured in a normal, disease free subject or tissue.

By the term “retinoic acid compound” is meant a compound that is either (a) the diterpene retinoic acid, or a derivative thereof, or (b) a compound having the structure $R^1-Ar^1-L^1Ar^2-L^2-C(=O)R^3$ (Formula (I)). Exemplary retinoic acid compounds described herein (including derivatives thereof) include the compounds identified in Figs. 2A, 2B, 10A, 10D, and Table 1. The term “diterpene retinoic acid” encompasses any stereoisomer of retinoic acid (e.g., the retinoic acid may be in the all-trans configuration (ATRA) or one or more of the double bonds may be in the *cis* configuration (e.g., 13-*cis* retinoic acid (13cRA)). Derivatives of the diterpene retinoic acid include reduced forms such as retinal, retinol, and retinyl acetate. In Formula (I), each of Ar^1 and Ar^2 is, independently, optionally substituted aryl or an optionally substituted heteroaryl; R^1 is H, an optionally substituted alkyl group, an optionally substituted alkenyl group, or an optionally substituted alkynyl group; each of L^1 and L^2 is selected,

independently from a covalent bond, an optionally substituted C₁₋₁₀ alkylene, an optionally substituted C₂₋₁₀ alkenylene (e.g., -CH=CH-, -COCH=CH-, -CH=CHCO-, a dienyl group, or a trienyl group), optionally substituted C₂₋₁₀ alkynylene (e.g., -C≡C-), or -(CHR⁴)_nCONR⁵-, -NR⁵CO-, where n is 0 or 1, R⁴ is H or OH, and R⁵ is H or optionally substituted alkyl; and R³ is H, OR⁴ or N(R⁴)₂, where each R⁴ is selected, independently, from H, optionally substituted alkyl, or optionally substituted heteroalkyl.

Any of the chemical groups, functional groups, or substituents described herein may be deuterated if the chemical group, functional group, or substituent has -H. As used herein, when a particular position in a compound of this invention is designated as being “deuterated” or “having deuterium,” it is understood that the position contains deuterium or includes deuterium (the element deuterium is represented by the letter “D” in chemical structures and formulas and indicated with a lower case “d” in chemical names, according to the Boughton system). When any of the position is deuterated, it is understood that the abundance of deuterium at that position is substantially greater than the natural abundance of deuterium, which is 0.015%. In certain embodiments, a composition of the invention has a minimum isotopic enrichment factor of at least 5 (0.075% deuterium incorporation), e.g., at least 10 (0.15% deuterium incorporation). In other embodiments, a composition has an isotopic enrichment factor of at least 50 (0.75% deuterium incorporation), at least 500 (7.5% deuterium incorporation), at least 2000 (30% deuterium incorporation), at least 3000 (45% deuterium incorporation), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium incorporation), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), or at least 6600 (99% deuterium incorporation).

As used herein, the term “isotopic enrichment factor” refers to the ratio of the isotopic abundance of a composition to the natural abundance of the specified isotope. For example, deuterium has a natural abundance of 0.015%. A compound with, for example, 45% deuterium incorporation at a specified position, has an isotopic enrichment factor of 3000 at that site relative to the natural abundance of deuterium.

Table 1. List of compounds structurally similar to retinoic acid

CID	IUPAC	Other names
444795	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Retinoic acid; tretinoin; Vitamin A acid

25145416	(2Z,4E,6Z,8Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
23275881	(2Z,4Z,6E,8Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
12358678	(2E,4E,6E,8Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL44478; CHEBI:168407; AC-540
10881132	(2Z,4Z,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10638113	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
9861147	(2E,4Z,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
9796370	(2E,4Z,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	1tyr; (11Z)-retinoic acid; 11-cis-Retinoic acid
6603983	(2E,4Z,6E,8Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Tocris-0695; Lopac-R-2625; Lopac-R-3255
6419708	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	9,13-di-cis-RA; 9,13-Di-cis-retinoic acid; 9-cis,13-cis-Retinoic acid
5282379	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Isotretinoin; 13-cis-Retinoic acid; Accutan
449171	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Alitretinoin; Panretin; 9-CIS-RETINOIC ACID
5538	3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Spectrum_001676; SpecPlus_000696; AC1L1KKH

54305566	2,4-dideuterio-7-methyl-3-(trideuteriomethyl)-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54305565	9-[3,3-dideuterio-6,6-dimethyl-2-(trideuteriomethyl)cyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10566385	(2E,4E,6Z,8E)-7-methyl-3-(trideuteriomethyl)-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10518761	(2E,4E,6Z,8E)-7-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)-3-(tritritiomethyl)nona-2,4,6,8-tetraenoic acid	
10470200	(2E,4Z,6Z,8E)-4,5-dideuterio-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10425032	(2E,4E,6Z,8E)-4,5-dideuterio-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10357701	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-4,5-ditritiocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10267048	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	
10086398	(2Z,4Z,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	
10086397	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-3,4-ditritiocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10063649	(2E,4E,6Z,8E)-9-[2,6-dimethyl-6-(trideuteriomethyl)cyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10040620	(2E,4E,6Z,8E)-9-(4,5-dideuterio-2,6,6-trimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10017935	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	
10017822	(2E,4E,6Z,8E)-9-(3,4-dideuterio-2,6,6-trimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
9995220	(2E,4Z,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	
9972327	(2Z,4Z,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-	

	4,5-ditritionona-2,4,6,8-tetraenoic acid	
9972326	(2E,4Z,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	
9839397	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-5-trititionona-2,4,6,8-tetraenoic acid	
6913160	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-5-trititionona-2,4,6,8-tetraenoic acid	Retinoic-11-t acid; AC1OC7MJ; all-trans-(11-3H)-Retinoic acid
6913136	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-4,5-ditritionona-2,4,6,8-tetraenoic acid	AC1OC7KP; Retinoic-11,12-t2 acid; 11,12-3H-Retinoic acid
6913131	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)-5,6-ditritionona-2,4,6,8-tetraenoic acid	AC1OC7KA; Retinoic-10,11-t2 acid; all-trans-(10,11-3H2)-Retinoic acid
6439661	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
134262	3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	SHGAZHPCJJPHSC-SPLUINJESA-N; FDEFF7D13961B76 6CC9FE8A74062324 3
56684147	(2E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,6,8-trienoic acid	
54219808	3,6,7-trimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53936974	3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,6,8-trienoic acid	
53740187	3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trienoic acid	

44725022	(Z)-3-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]hept-2-enoic acid	AC1Q2V68; (2Z)-3-[(E)-2-(2,6,6-trimethylcyclohex-1-en-1-yl)ethenyl]hept-2-enoic acid
21590819	(2Z,4E,8E)-3-methyl-7-methylidene-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,8-trienoic acid	CHEMBL182393
11738545	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)deca-2,4,6,8-tetraenoic acid	
10518336	(2E,4E,8E)-3-methyl-7-methylidene-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,8-trienoic acid	CHEMBL426963
10380944	(2E,4E,6Z,8E)-3-ethyl-7-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10335106	(2E,4E,6E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trienoic acid	CHEMBL487208
10286439	(2E,4E,6Z,8E)-7-ethyl-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10149682	(2E,4E,6Z,8E)-3,6,7-trimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10041353	(2E,4E,6E,8E)-3-ethyl-7-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
6439749	(2E,4E,6E,8E)-9-(2-ethyl-6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	SRI 2712-24; 2,4,6,8-Nonatetracenoic acid,
5496917	(2E,4Z,6E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trienoic acid	AC1NUZ8L
5326825	(2Z,4Z,6E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trienoic acid	AC1NS159
4136524	3-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]hept-2-enoic acid	AC1N4YDA
135317	9-(2-ethyl-6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
54525370	13-(2,6,6-trimethylcyclohexen-1-yl)trideca-2,4,6,8,10,12-	

	hexaenoic acid	
54472611	4,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54398880	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclopenten-1-yl]penta-2,4-dienoic acid	
54044750	11-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	
53876852	3,7-dimethyl-9-(2,4,6,6-tetramethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53790569	9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53743104	5,9-dimethyl-11-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	
44579060	(2E,4E,6Z,8E)-9-(2-butyl-6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	CHEMBL518436
44393163	(2Z,4E,8E)-7-methylidene-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,8-trienoic acid	
25141345	(2E,4E,6E,8E)-9-(2-butyl-6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
19609253	(2E,4E)-3-methyl-5-[2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclopenten-1-yl]penta-2,4-dienoic acid	
14731990	(2E,4E,6E,8E)-7-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
11141121	(2E,4E,6E,8E)-4,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10712359	(2E,4E,6Z)-3-methyl-7-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]undeca-2,4,6-trienoic acid	
10474100	(2E,4E,6E,8E,10E,12E)-3,7,11-trimethyl-13-(2,6,6-trimethylcyclohexen-1-yl)trideca-2,4,6,8,10,12-hexanoic acid	
10426543	(E,4E)-3-methyl-4-[3-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohex-2-en-1-ylidene]but-2-enoic acid	
10358907	(Z,4E)-3-methyl-4-[(4E)-3-methyl-4-[(2,6,6-trimethylcyclohexen-1-	

	yl)methylidene]cyclohexa-2,5-dien-1-yl)methylidene)	
10314319	(2E,4E,6E,8E,10E)-5,9-dimethyl-11-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	CHEMBL225948
10286753	(2E,4E,6Z,8E)-7-tert-butyl-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10266931	(2E,4E,6Z)-3-methyl-7-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]deca-2,4,6-trienoic acid	CHEMBL507779
10125803	(2E,4E,6Z)-3-methyl-7-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]deca-2,4,6-trienoic acid	
10087786	(Z,4E)-3-methyl-4-[3-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohex-2-en-1-ylidene]but-2-enoic acid	
10015486	(2E,4E,6E)-5-methyl-7-(2,6,6-trimethylcyclohexen-1-yl)hepta-2,4,6-trienoic acid	
9929074	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
9860303	(2E,4E,6E,8E)-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
5355027	(2E,4E)-3-methyl-5-(2,6,6-trimethylcyclohexen-1-yl)penta-2,4-dienoic acid	C15 acid; AC1NS6O9; NSC23978
167095	3-methyl-5-(2,6,6-trimethylcyclohexen-1-yl)penta-2,4-dienoic acid	AC1L4ZB4
56606832	3,7-dimethyl-9-(9,9,11-trimethylspiro[2.5]oct-10-en-10-yl)nona-2,4,6,8-tetraenoic acid	
54548815	3,7,11,11-tetramethyldodeca-2,4-dienoic acid	
54515105	7-methyl-3-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]nona-2,5-dienoic acid	YLWKTERFWUXE BW-UHFFFAOYSA-N; 005B26AC36D10A0 C9DB5EF006864943 F
54358950	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-	

	yl)ethenyl]cyclohepten-1-yl]penta-2,4-dienoic acid	
54353726	3,7,11,11-tetramethyltrideca-2,4-dienoic acid	
54193713	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cycloocten-1-yl]penta-2,4-dienoic acid	
53946778	2,3,7-trimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53944823	9-(6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	JAIGDKSXLVOFM H-UHFFFAOYSA-N; F42136BEED6C5A3 745B9BA23356D783 0
53921377	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexen-1-yl]penta-2,4-dienoic acid	
44579100	(2E,4E,6Z,8E)-9-[6,6-dimethyl-2-(2-methylpropyl)cyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	CHEMBL476773
44579056	(2E,4E,6E,8E)-9-[6,6-dimethyl-2-(2-methylpropyl)cyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	CHEMBL476348
44314230	(2Z,5E)-7-methyl-3-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]nona-2,5-dienoic acid	CHEMBL75548; CHEBI:220121
25011742	(2E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,8-dienoic acid	
22646220	(2E,4E,6E,8E)-2,3-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
20830941	(2E,4E,6E,8E)-2,3-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
19609240	(2E,4E)-3-methyl-5-[(1Z)-2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cycloocten-1-yl]penta-2,4-dienoic acid	
18977383	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,5,6,6-tetramethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
15125883	(2Z,4E,6E,8E)-2,3,7-trimethyl-9-(2,6,6-trimethylcyclohexen-1-	

	yl)nona-2,4,6,8-tetraenoic acid	
15125882	(2E,4E,6E,8E)-2,3,7-trimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL153895; 14-methyl-all-trans-retinoic acid; LMPR01090034
11266097	(2Z,4E,8E)-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,8-trien-6-ynoic acid	
11000660	(2E,4E,6Z,8E)-9-(6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10733921	(2E,4E,6Z)-7-(8,8-dimethyl-4,5,6,7-tetrahydro-3H-naphthalen-2-yl)-3-methylocta-2,4,6-trienoic acid	
10636975	(2E,4E,6E,8E)-9-(6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10591236	(2E,4E,6Z)-7-(4a,8-dimethyl-4,5,6,7-tetrahydro-3H-naphthalen-2-yl)-3-methylocta-2,4,6-trienoic acid	
10404132	(Z,4E)-3-methyl-4-[(4E)-3-methyl-4-[(2,6,6-trimethylcyclohexen-1-yl)methylidene]cyclohex-2-en-1-ylidene]but-2-enoic acid	
10314318	(E,4E)-3-methyl-4-[(4E)-3-methyl-4-[(2,6,6-trimethylcyclohexen-1-yl)methylidene]cyclohex-2-en-1-ylidene]but-2-enoic acid	
10215224	(2E,4E,6Z,8E)-3-methyl-7-propan-2-yl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10193246	(2E,4E)-3-methyl-6-[1-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclopropyl]hexa-2,4-dienoic acid	
9841547	(2E,4E)-3-methyl-5-[2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohepten-1-yl]penta-2,4-dienoic acid	
9830767	(2Z,4E,6Z,8E)-9-(6,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
9819335	(2E,4E)-3-methyl-5-[2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexen-1-yl]penta-2,4-dienoic acid	Ro 25-6603; 173792-73-9
56667667	(2E,4E,6Z,8E)-3,7-dimethyl-9-(6-methyl-3-prop-1-en-2-ylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL455993; CHEMBL455994

54758572	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	9-cis-Retinoate; CPD-13549
54426679	2,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54325149	6-chloro-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53702687	6-iodo-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
29986894	(2E,4Z,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	ZINC22066351
29927144	(2E,4E,6E,8Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	ZINC21992287
24916820	(2E,4E,6E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trienoate	2g78
24771817	3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	CHEBI:15036
21917290	(2E,4E,6E,8E)-9-(5-tert-butyl-2,6,6-trimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
19609245	(2E,4E,6E,8E)-6-chloro-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
19609224	(2E,4E,6E,8E)-6-iodo-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10924150	(2E,4E,6Z,8E)-9-(2,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10613228	(2E,4E,6E,8E)-9-(2,6-dimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
10469989	(2E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,6,8-trien-4-ynoic acid	
10334998	(2E,4E)-3-methyl-5-[2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclopropyl]penta-2,4-dienoic acid	
9904356	(2Z,4E,6Z)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-	

	2,4,6-trien-8-ynoic acid	
7364357	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	AC1OKKW8; ZINC12661824;
7048538	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	13-cis-retinoate; ZINC03792789
6440565	2E,4E,6E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6-trien-8-ynoic acid	7,8-Dehydroretinoic acid; 7,8-Didehydroretinoic acid
6419707	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	Retinoate; all-trans-Retinoate; Tretinoine
5771658	(Z)-3-(2,6,6-trimethylcyclohexen-1-yl)prop-2-enoic acid	NSC-202789; AC1NY9IQ; NCGC00014560
5383969	(E)-3-(2,6,6-trimethylcyclohexen-1-yl)prop-2-enoic acid	NSC202789; NSC-20278
5353358	(2Z,4E)-3-methyl-6-(2,7,7-trimethyl-3-methylidene-1,4,5,6-tetrahydroinden-2-yl)hexa-2,4-dienoic acid	AC1NS43Q
5289278	(2E,4E)-3-methyl-6-[(2R)-2,7,7-trimethyl-3-methylidene-1,4,5,6-tetrahydroinden-2-yl]hexa-2,4-dienoic acid	NSC202789; 3-(2,6,6-trimethyl-1-cyclohexen-1-yl)acrylic acid; AC1L77HZ
305742	3-(2,6,6-trimethylcyclohexen-1-yl)prop-2-enoic acid	NSC202789; 3-(2,6,6-trimethyl-1-cyclohexen-1-yl)acrylic acid; AC1L77HZ
1851	3-methyl-6-(2,7,7-trimethyl-3-methylidene-1,4,5,6-tetrahydroinden-2-yl)hexa-2,4-dienoic acid	AC1L1CDO
54399542	6-bromo-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	

54233476	3,7-dimethyl-5-oxo-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,6,8-trienoic acid	
54033110	2,5,9-trimethyl-11-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	
53936708	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethynyl]cyclopenten-1-yl]penta-2,4-dienoic acid	
44314320	(2Z,4E)-3-methyl-5-[2-[(E)-2-(3,3,6,6-tetramethylcyclohexen-1-yl)ethenyl]cyclopropyl]penta-2,4-dienoic acid	CHEMBL73973; CHEBI:220303
44314319	(2E,4E)-3-methyl-5-[2-[(E)-2-(3,3,6,6-tetramethylcyclohexen-1-yl)ethenyl]cyclopropyl]penta-2,4-dienoic acid	CHEMBL74331; CHEBI:220301
22373193	(2E,4E)-3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethynyl]cyclopenten-1-yl]penta-2,4-dienoic acid	
21145248	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
20151571	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
19609231	(2E,4E,6E,8E)-6-bromo-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
16727824	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	All-trans-Retinoic acid & 9-cis-Retinoic Acid
11015604	(2E,4E,6E,8E,10E,12E,14E,16E)-2,6,11,15-tetramethyl-17-(2,6,6-trimethylcyclohexen-1-yl)-3-tritioheptadeca-2,4,6,8-trimethylcyclohexen-1-yl)-3-tritioheptadeca-2,4,6,8,10,12,14,16-octaenoic acid	
10406618	(2E,4Z,6E,8E,10E,12E)-2,7,11-trimethyl-13-(2,6,6-trimethylcyclohexen-1-yl)trideca-2,4,6,8,10,12-hexanoic acid	
9976193	(2E,4E,6E,8E,10E,12E)-2,7,11-trimethyl-13-(2,6,6-trimethylcyclohexen-1-yl)trideca-2,4,6,8,10,12-hexanoic acid	
9843074	(2E,4E,6E)-3-methyl-7-(4,4,7,7-tetramethyl-2-pentyl-1,3,5,6-tetrahydroinden-2-yl)hepta-2,4,6-trienoic acid	
6439881	(2Z,4E,6Z,8E)-9-(3,3-difluoro-2,6,6-trimethylcyclohexen-1-yl)-	DFRA; 4,4-

	3,7-dimethylnona-2,4,6,8-tetraenoic acid	Difluororetinoic acid; AC1O5SM
6436320	(2E,4E,6Z,8E,10E,12E,14E,16E)-2,6,11,15-tetramethyl-17-(2,6,6-trimethylcyclohexen-1-yl)heptadeca-2,4,6,8,10,12,14,16-octaenoic acid	AC1O5LFK; beta-apo-8'-Carotenoic acid; 8'-Apo-beta,psi-carotenoic acid
5387557	(2Z)-2-[5-(2,6,6-trimethylcyclohexen-1-yl)-3-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexanoic acid	NSC624510; AC1NTSHG; AC1Q5T6Y
5366642	(2E,4E,6E,8E)-9-(3,3-difluoro-2,6,6-trimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	4,4-Difluororetinoic acid; AC1NSNWF; 4,4-Difluororetinoic acid (all-trans)
361473	2-[5-(2,6,6-trimethylcyclohexen-1-yl)-3-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohex-2-en-1-yl)heptadeca-2,4,6,8,10,12,14,16-octaenoic acid	AC1L7IQC; NCI60_007432; 2-[5-(2,6,6-trimethylcyclohexen-1-yl)-3-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohex-2-en-1-ylidene]acetic acid
146218	9-(3,3-difluoro-2,6,6-trimethylcyclohexen-1-yl)-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
56660872	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2-methyl-5-prop-1-en-2-ylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL457645; CHEMBL513434
54587023	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3S,6R)-3-methyl-6-prop-1-en-2-ylcyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773351
54586043	(2E,4E,6Z)-3-methyl-7-[(3R,6S)-3-methyl-6-propan-2-ylcyclohexen-1-yl]octa-2,4,6-trienoic acid	CHEMBL1773361
54310202	7-ethyl-3,11-dimethyltrideca-2,4-dienoic acid	

54177995	8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	OZUIXDDSOLQKN K-UHFFFAOYSA-N; 982DADEA9DC557 9A132BDF2AD7FA 647A
54012267	3,8,12-trimethyltrideca-2,4-dienoic acid	
53787191	3,8,13-trimethyltetradeca-2,4-dienoic acid	
53743194	4-methyl-6-(2,6,6-trimethylcyclohexen-1-yl)hex-2-enoic acid	
53710521	3,7,13-trimethyltetradeca-2,4-dienoic acid	
53707670	3,7-dimethyl-8-(3-methyl-2-propan-2-ylcyclohex-2-en-1-ylidene)octa-2,4,6-trienoic acid	BYHSFJNWVLBCI M-UHFFFAOYSA-N; 14B10A34153F37A6 6327788679FAC42F
53666154	3,7,11-trimethyltrideca-2,4-dienoic acid	
53438161	3,7,11-trimethyltetradeca-2,4-dienoic acid	
53427754	7,7-dimethylcosa-2,4-dienoic acid	
52952998	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3R,6S)-3-methyl-6-prop-1-en-2-ylcyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773352
44631433	(2Z,4E)-3-methyl-5-(2,2,4-trimethylcyclohex-3-en-1-yl)penta-2,4-dienoic acid	FZFFLFPGBIXCKI- STRRHFTISA-
44291210	(2Z,4Z,6Z,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	CHEMBL43954
44290946	(2E,4Z,6Z,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	CHEMBL43833; CHEBI:167938
24845989	sodium (2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	LS-143475
23670222	potassium (2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	

23665641	sodium (2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	Sodium retinoate; Retinoic acid, sodium salt; Vitamin A acid sodium sal
23265304	(2E,4E)-3-methyl-5-(2,2,4-trimethylcyclohex-3-en-1-yl)penta-2,4-dienoic acid	
21437585	(2E,4E)-3,8,12-trimethyltrideca-2,4-dienoic acid	
21437539	(2E,4E)-3,8,13-trimethyltetradeca-2,4-dienoic acid	
21437504	(2E,4E)-3,7,13-trimethyltetradeca-2,4-dienoic acid	
21158960	(2E,4E)-7,7-dimethylcosa-2,4-dienoic acid	
20270951	(6E,8E)-2,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,3,6,8-tetraenoic acid	
19609232	(2E,4E)-3-methyl-5-[2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexen-1-yl]penta-2,4-dienoic acid	
11130378	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohex-2-en-1-yl)nona-2,4,6,8-tetraenoic acid	
11066537	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohex-2-en-1-yl)nona-2,4,6,8-tetraenoic acid	
10470917	(2Z,4E,6Z,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	
10402558	(2Z,4E,6E,8E)-3,7-dimethyl-8-(3-methyl-2-propan-2-ylcyclohex-2-en-1-ylidene)octa-2,4,6-trienoic acid	
10357464	(2E,4E,6Z,8E)-3,7-dimethyl-8-(3-methyl-2-propan-2-ylcyclohex-2-en-1-ylidene)octa-2,4,6-trienoic acid	
10086191	(2E,4E,6E,8E)-3,7-dimethyl-8-(3-methyl-2-propan-2-ylcyclohex-2-en-1-ylidene)octa-2,4,6-trienoic acid	CHEMBL333032; CHEBI:299410
10086189	(2Z,4E,6Z,8E)-3,7-dimethyl-8-(3-methyl-2-propan-2-ylcyclohex-2-en-1-ylidene)octa-2,4,6-trienoic acid	
9972952	(2Z,4E,6E,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	CHEMBL44582; CHEBI:168408
9972949	(2E,4E,6Z,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-	

	3,7-dimethylocta-2,4,6-trienoic acid	
9883342	(2E,4E,6E,8E)-8-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3,7-dimethylocta-2,4,6-trienoic acid	CHEMBL46398; CHEBI:168441
5372326	(E)-3-methyl-5-(2,6,6-trimethylcyclohexen-1-yl)pent-2-enoic acid	AC1NSY3I; 2-Pentenoic acid, 3-methyl-5-(2,6,6-trimethyl-1-cyclohexenyl); (E)-3-methyl-5-(2,6,6-trimethylcyclohexen-1-yl)pent-2-enoic acid
445560	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohex-2-en-1-yl)nona-2,4,6,8-tetraenoic acid	AC1L9I79
56667221	(2E,4E,6Z,8E)-3,7-dimethyl-9-(3-methyl-6-propan-2-ylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL508378
54585066	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(1S,4R,5R)-4,6,6-trimethyl-3-bicyclo[3.1.1]hept-2-enyl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773358
54585064	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3R)-3-methyl-6-propan-2-ylidenecyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773355
54582176	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3S)-3-methyl-6-propan-2-ylidenecyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773354
54581148	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(1R,2R,5S)-2-methyl-5-propan-2-yl-3-bicyclo[3.1.0]hex-3-enyl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773360
54542310	3,4,4-trimethyltetradec-2-enoic acid	
54521054	3,4,4-trimethyloctadec-2-enoic acid	
54518673	3,7-dimethyl-9-(2,6,6-trimethyl-5-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54348687	3,7,10,11-tetramethyldodeca-2,4-dienoic acid	
54325421	3,4,4-trimethylheptadec-2-enoic acid	

54316493	3,4,4-trimethylpentadec-2-enoic acid	
54305044	2-ethyl-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54265680	3,7,11,15-tetramethylhexadeca-2,4-dienoic acid	
54194359	3,7-dimethyl-9-(2,6,6-trimethyl-4-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
54170467	3,7,11,15-tetramethylhexadeca-2,4,6,14-tetraenoic acid	
54167172	3,4,4-trimethylhexadec-2-enoic acid	
54105865	3,7,7,11,11-pentamethyldodec-2-enoic acid	
54064253	2-ethyl-5,9-dimethyl-3-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	
53961371	3,7,11-trimethyldodeca-2,4,11-trienoic acid	
53936602	9-[5-(2-cyclohexylethyl)-2,6,6-trimethylcyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
53825233	3,7,11,15,19-pentamethylcosa-2,4,6,10,18-pentaenoic acid	
53801569	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethynyl]cyclohepten-1-yl]penta-2,4-dienoic acid	
53725805	3,7-dimethyldodeca-2,4-dienoic acid	
53700416	3,7,11,15-tetramethylhexadeca-2,4,6-trienoic acid	
52953080	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3S,6R)-3-methyl-6-propan-2-ylcyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773353
52952997	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(1R,4S,5S)-4,6,6-trimethyl-3-bicyclo[3.1.1]hept-2-enyl]nona-2,4,6,8-tetraenoic acid	CHEMBL1773357
52921782	(2E,5R,10E,12E)-3,5,15-trimethyl-7-methylidenehexadeca-2,10,12-trienoic acid	LMFA01020367; 16:3(2E,10E,12E)(3Me,5Me[R],7My,15Me)
46178652	(2E,4E)-5-[(1R)-2,2-dimethyl-6-methylidenecyclohexyl]-3-methylpenta-2,4-dienoic acid	
44579059	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,2,6-trimethylcyclohexyl)nona-2,4,6,8-tetraenoic acid	CHEMBL451158

25147656	(2E,4E,6Z,8E)-3,7-dimethyl-9-[(3R,6S)-3-methyl-6-propan-2-ylcyclohexen-1-yl]nona-2,4,6,8-tetraenoic acid	CHEMBL508378
22168242	(2E,4E,6E,10E)-3,7,11,15,19-pentamethylcosa-2,4,6,10,18-pentaenoic acid	
22168239	(2E,4E,6E)-3,7,11,15-tetramethylhexadeca-2,4,6-trienoic acid	
22168234	(2E,4E,6E)-3,7,11,15-tetramethylhexadeca-2,4,6,14-tetraenoic acid	
21764469	(2E,4E)-3-methyl-5-[(1R)-2,6,6-trimethylcyclohex-2-en-1-yl]penta-2,4-dienoic acid	
21650797	acetyl (2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraeneperoxoate	
21525820	(2E,4E)-7,11,11-trimethyldodeca-2,4-dienoic acid	
21525806	(2E,4E)-3,7-dimethyldodeca-2,4-dienoic acid	
21291068	(E)-3,4,4-trimethylhexadec-2-enoic acid	
21291063	(E)-3,4,4-trimethyltetradec-2-enoic acid	
21291060	(E)-3,4,4-trimethylpentadec-2-enoic acid	
21291047	(E)-3,4,4-trimethylheptadec-2-enoic acid	
21291045	(E)-3,4,4-trimethyloctadec-2-enoic acid	
20830940	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,5,6,6-tetramethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	
20306860	(2E,4E)-3,7,11-trimethyldodeca-2,4,11-trienoic acid	
20027300	azanium (2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	
19609235	(2E,4E)-2-iodo-3-methyl-5-(2,6,6-trimethylcyclohexen-1-yl)penta-2,4-dienoic acid	
19606927	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-4-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
18977382	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,5,6,6-tetramethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoate	
16061319	(2Z,4E,6Z,8E)-7-(hydroxymethyl)-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	19-Hydroxy-13-cis-retinoic acid;

		LMPR01090029
16061318	(2E,4E,6Z,8E)-7-(hydroxymethyl)-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	19-Hydroxy-all-trans-retinoic acid; LMPR01090028
15125888	(2E,4E,6E,8E)-2-ethyl-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL154239
11747707	(2E,4E,6Z,8E)-3,7-dimethyl-9-(6-methylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
11602784	(2E,4E)-3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethynyl]cyclohepten-1-yl]penta-2,4-dienoic acid	
10516342	(2E,4E,6E,8E)-3,7-dimethyl-9-(6-methylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
10354668	(Z,4E)-4-(3-ethyl-2-propan-2-ylcyclohex-2-en-1-ylidene)-3-methylbut-2-enoic acid	
10053647	(2Z,4Z,6E,8E,10E,12E,14E,16E,18E,20E,22E,24E)-2,6,10,14,19,23-hexamethyl-25-(2,6,6-trimethylcyclohexen-1-yl)pentacos-2,4,6,8,10,12,14,16,18,20,22,24-dodecaenoic acid	
9995780	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-5-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Oxo-13-cis-retinoate; 4-keto-13-cis-retinoate
9949957	(2E,4E,6E,8E)-3,7-dimethyl-8-[3-(2-methylpropyl)-2-propan-2-ylcyclohex-2-en-1-ylidene]octa-2,4,6-trienoic acid	
9948768	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-5-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
9829386	(2E,4Z,6E,8E,10E,12E,14E,16E,18E,20E,22E,24E)-2,6,10,14,19,23-hexamethyl-25-(2,6,6-trimethylcyclohexen-1-yl)pentacos-2,4,6,8,10,12,14,16,18,20,22,24-dodecaenoic acid	
6477090	(2Z,4Z,6Z,8E,10Z,12Z,14E,16Z,18Z,20E,22Z,24E)-2,6,10,14,19,23-hexamethyl-25-(2,6,6-trimethylcyclohexen-1-	AC1O53P5; 3',4'-Didehydro-,psi.-

	yl)pentacos-2,4,6,8,10, 12,14,16,18,20,22,24-dodecaenoic acid	caroten-16'-oic acid
6439734	(2Z,4E,6Z,8E)-3,7-dimethyl-9-(2,2,6-trimethylcyclohexyl)nona- 2,4,6,8-tetraenoic acid	7,8-Dihydroretinoic acid
6437018	(2Z,4E)-3,7,11-trimethyldodeca-2,4-dienoic acid	AC1O5MUO; EINECS 258-354-9
6437016	(2E,4E)-3,7,11-trimethyldodeca-2,4-dienoic acid	AC1O5MUI; ChEMBL37590
5476505	(2E,4E)-3-methyl-5-(2,6,6-trimethylcyclohex-2-en-1-yl)penta-2,4- dienoic acid	AC1O5MUI; ChEMBL37590
5460164	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,2,6-trimethylcyclohexyl)nona- 2,4,6,8-tetraenoic acid	Retinyl ester; all- trans-Retinyl ester
5281248	(2E,4E,6E,8E,10E,12E,14E,16E,18E,20E,22E,24E)- 2,6,10,14,19,23-hexamethyl-25-(2,6,6-trimethylcyclohexanoic acid	NSC635690; Torularhodin; AC1NQY9
637039	2E,4E,6E,8E,10E,12E,14E,16E,18E,20E)-2,6,10,15,19- pentamethyl-21-(2,6,6-trimethylcyclohexen-1-yl)hexanoic acid	Neurosporaxanthin; all-trans- Neurosporaxanthin
428485	3-methyl-5-(2,6,6-trimethylcyclohex-2-en-1-yl)penta-2,4-dienoic acid	AC1L8LML; 3- methyl-5-(2,6,6- trimethylcyclohex-2- en-1-yl)penta-2,4- dienoic acid
103723	3,7,11-trimethyldodeca-2,4-dienoic acid	
94165	2,6,10,14,19,23-hexamethyl-25-(2,6,6-trimethylcyclohexen-1- yl)pentacos-2,4,6,8,10,12,14,16,18,20,22, 24-dodecanenoic acid	AC1L3RN8; NCI60_011910
56661049	(2E,4E,6Z,8E)-3,7-dimethyl-9-(4,4,6,6-tetramethyl-2- bicyclo[3.1.1]hept-2-enyl)nona-2,4,6,8-tetraenoic acid	CHEMBI.455992
54581147	(2E,4E,6Z,8E)-9-[(1S,5R)-6,6-dimethyl-4-bicyclo[3.1.1]hept-3- enyl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	CHEMBL1773359
54478024	3,4,4-trimethylnon-2-enoic acid	

54476971	3,4,4-trimethylundec-2-enoic acid	
54287870	3-formyl-7-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	RVKZSGIKOAAYYJ -UHFFFAOYSA-N; 293564D2B64FAC5 F524A1B691CBF7C 6B
54116397	3,7-dimethyl-2-propan-2-yl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	NKQIYDSGIYJXSA -UHFFFAOYSA-N; 5597749F477D668D 55E163C44DA1F3E B
54073647	3,4,4-trimethyldec-2-enoic acid	
53995964	3-methyl-5-[2-[2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexyl]penta-2,4-dienoic acid	
53919798	3,4,4-trimethyldodec-2-enoic acid	
53889922	3,7-dimethyl-9-(2,4,4,6,6-pentamethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53887460	4-(hydroxymethyl)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
53854796	3-methyl-6-(3,3,7,7-tetramethyl-3a,4,5,6-tetrahydroinden-2-ylidene)hexa-2,4-dienoic acid	
53754609	2-ethyl-5,9-dimethyl-11-(2,6,6-trimethylcyclohexen-1-yl)undeca-2,4,6,8,10-pentaenoic acid	
50925583	(2E,4E,6E,8E)-9-[(1R,2R,4aS,8aR)-1,6-dimethyl-2-propyl-4a,5,8,8a-tetrahydro-2H-naphthalen-1-yl]-8-methylnona-2,4,6,8-tetraenoic acid	
45039634	(2E,4E,6E,8E)-9-[6,6-dimethyl-3-oxo-2-(trideuteriomethyl)cyclohexen-1-yl]-3,7-dimethylnona-2,4,6,8-tetraenoic acid	
21291081	(E)-3,4,4-trimethyldec-2-enoic acid	
21291044	(E)-3,4,4-trimethyldodec-2-enoic acid	
21291042	(E)-3,4,4-trimethylnon-2-enoic acid	

21291032	(E)-3,4,4-trimethylundec-2-enoic acid	
19384872	(E)-4-[(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenyl]oxy-4-oxobut-2-enoic acid	
16061321	(2Z,4E,6Z,8E)-7-formyl-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	19-Oxo-9-cis-retinoic acid; LMPR01090031
16061320	(2E,4E,6Z,8E)-7-formyl-3-methyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	19-Oxo-all-trans-retinoic acid; LMPR01090030
15125894	(2E,4E,6E,8E)-3,7-dimethyl-2-propan-2-yl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL153894
10043037	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,4,4,6,6-pentamethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	CHEMBL103068
9972939	(2E,4E,6Z,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
9906064	(2E,4E)-3-methyl-5-[(1R)-2-[(E)-2-(2,6,6-trimethylcyclohexen-1-yl)ethenyl]cyclohexyl]penta-2,4-dienoic acid	
9902057	(2Z,4E,6Z,8E)-4-(hydroxymethyl)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	
6437087	(2Z,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	Oxoretinoic acid; 4-Oxo-isotretinoin
6437063	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	4-Oxoretinoic acid; 4-Ketoretinoic acid
447276	(2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexa-1,3-dien-1-yl)nona-2,4,6,8-tetraenoic acid	Vitamin A2 acid; 3,4-Didehydroretinoic acid
104857	3,7-dimethyl-9-(2,6,6-trimethyl-3-oxocyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	

The term “aryl,” as used herein, represents a mono- or bicyclic C₆-C₁₄ group with $[4n + 2]$ π electrons in conjugation and where n is 1, 2, or 3. Aryl groups also include ring systems where the ring system having $[4n + 2]$ π electrons is fused to a non-aromatic cycloalkyl or a non-aromatic heterocyclyl.

Phenyl is an aryl group where n is 1. Aryl groups may be unsubstituted or substituted with, e.g., 1, 2, 3, or 4 substituent groups as defined herein. Still other exemplary aryl groups include, but are not limited to, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, fluorenyl, indanyl, and indenyl.

The term “cycloalkyl,” as used herein, represents a monovalent saturated or unsaturated non-aromatic cyclic hydrocarbon group from three to ten carbons, unless otherwise specified, and is exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]heptyl, and the like. In some embodiments, the cycloalkyl is a polycyclic (e.g., adamantyl). Cycloalkyl groups may be unsubstituted or substituted with, e.g., 1, 2, 3, or 4 substituent groups as defined herein.

The term “heteroaryl,” as used herein, represents an aromatic (i.e., containing $4n+2$ pi electrons within the ring system) 5- or 6-membered ring containing one, two, three, or four heteroatoms independently selected from the group consisting of nitrogen, oxygen, and sulfur, as well as bicyclic, tricyclic, and tetracyclic groups in which any of the aromatic ring is fused to one, two, or three heterocyclic or carbocyclic rings (e.g., an aryl ring). Exemplary heteroaryls include, but are not limited to, furan, thiophene, pyrrole, thiadiazole (e.g., 1,2,3-thiadiazole or 1,2,4-thiadiazole), oxadiazole (e.g., 1,2,3-oxadiazole or 1,2,5-oxadiazole), oxazole, isoxazole, isothiazole, pyrazole, thiazole, triazole (e.g., 1,2,4-triazole or 1,2,3-triazole), pyridine, pyrimidine, pyrazine, triazine (e.g., 1,2,3-triazine, 1,2,4-triazine, or 1,3,5-triazine), 1,2,4,5-tetrazine, indolyl, quinoliny, isoquinoliny, benzimidazolyl, benzothiazolyl, and benzoxazolyl. Heteroaryls may be unsubstituted or substituted with, e.g., 1, 2, 3, or 4 substituent groups as defined herein.

The term “heterocyclyl,” as used herein represents a non-aromatic 5-, 6- or 7-membered ring, unless otherwise specified, containing one, two, three, or four heteroatoms independently selected from the group consisting of nitrogen, oxygen, and sulfur. Heterocyclyl groups may be unsubstituted or substituted with, e.g., 1, 2, 3, or 4 substituent groups as defined herein.

Where a group is substituted, the group may be substituted with 1, 2, 3, 4, 5, or 6 substituent groups. Optional substituent groups include, but are not limited to: C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, halogen (-F, -Cl, -Br, or -I), azido ($-N_3$), nitro ($-NO_2$), cyano ($-CN$), acyloxy ($-OC(=O)R'$), acyl ($-C(=O)R'$), alkoxy ($-OR'$), amido ($-NR'C(=O)R''$ or $-C(=O)NRR'$), amino ($-NRR'$), carboxylic acid ($-CO_2H$), carboxylic ester ($-CO_2R'$), carbamoyl ($-OC(=O)NR'R''$ or $-NRC(=O)OR'$), hydroxy ($-OH$), oxo ($=O$), isocyano ($-NC$), sulfonate ($-S(=O)_2OR$), sulfonamide ($-S(=O)_2NRR'$ or $-NRS(=O)_2R'$), or sulfonyl ($-S(=O)_2R$), where each R or R' is selected, independently, from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, or heteroaryl. In some embodiments, the substituent groups themselves may be further substituted with, for example, 1, 2, 3, 4, 5, or 6 substituents as defined

herein. For example, a C₁₋₆ alkyl, aryl, or heteroaryl group may be further substituted with 1, 2, 3, 4, 5, or 6 substituents as described herein.

The retinoic acid compounds of the invention inhibit Pin1 activity (e.g., as determined by the fluorescence polarization-based displacement assay or PPIase assay as describe herein). This inhibition can be, e.g., greater than 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, or greater.

The language “anti-proliferative compound” is intended to include chemical reagents which inhibit the growth of proliferating cells or tissues wherein the growth of such cells or tissues is undesirable. Chemotherapeutic agents are well known in the art (as well as described herein), and are typically used to treat neoplastic diseases, tumors, and cancers.

“Treatment,” as used herein, is defined as the application or administration of a therapeutic agent (e.g., a retinoic acid compound) to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has a disease, a symptom of disease or a predisposition toward a disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease, the symptoms of disease or the predisposition toward disease, or to slow the progression of the disease.

As used herein, the terms “sample” and “biological sample” include samples obtained from a mammal or a subject containing Pin1 which can be used within the methods described herein, e.g., tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Typical samples from a subject include tissue samples, tumor samples, blood, urine, biopsies, lymph, saliva, phlegm, pus, and the like.

By a “low dosage” or “low concentration” is meant at least 5% less (e.g., at least 10%, 20%, 50%, 80%, 90%, or even 95%) than the lowest standard recommended dosage or lowest standard recommended concentration of a particular compound formulated for a given route of administration for treatment of any human disease or condition. For example, a low dosage of an anti-proliferative compound formulated for oral administration will differ from a low dosage of an anti-proliferative compound formulated for intravenous administration.

Brief Description of the Drawings

Figure 1A is a schematic showing relationship between Pin1 and the activity of the indicated genes.

Figure 1B is a schematic showing the role of Pin1 in the indicated signal transduction pathways.

Figure 2A is a schematic of the chemical structure of 13-cis-retinoic acid and all-trans retinoic acid.

Figure 2B is a schematic showing additional retinoic acid compounds.

Figure 3 is a pair of graphs showing the ability of 13-cis-retinoic acid and all-trans retinoic acid at different concentrations to compete with a fluorescence-labeled Pin1 peptide inhibitor probe BIKL-488 for binding Pin1 after an incubation of the indicated amount of time.

Figure 4 is a pair of graphs showing inhibition of Pin1 by cis-retinoic acid and trans-retinoic acid at the indicated concentrations.

Figure 5 is a photograph indicating both cis and trans retinoic acids can bind to Pin1 and compete away DAPK, which binds to the Pin1 catalytic domain.

Figures 6A is a graph showing specific residues in the Pin1 active site that are important for interacting with the detecting probe BIKL-488.

Figure 6B is a graph showing specific residues in the Pin1 active site that are important for interacting with retinoic acid.

Figure 7A is a pair of graphs showing cell viability of the indicated cell lines as a function of concentration of cis-retinoic acid or trans-retinoic acid.

Figure 7B is a series Western blots showing expression of the indicated protein in cells treated with of either cis-retinoic acid or trans-retinoic acid in the indicated cell lines.

Figure 8A is a photograph indicating the amount of Pin1 or GAPDH mRNA produced in the presence of the indicated compound as measured in an RT-PCR experiment.

Figure 8B is a Western blot showing Pin1 and tubulin protein levels in cells treated with the indicated compounds

Figure 8C is a Western blot showing Pin1 and tubulin protein levels in cells treated with cycloheximide and the indicated compound.

Figure 9A is a pair of graphs showing cell viability as a function of the concentration of all trans-retinoic acid in the indicated cell types.

Figure 9B is a Western blot showing Pin1 protein levels in the indicated cell types.

Figure 9C is Western blot showing Pin1 protein levels in the indicated cell types treated with either 13cRA or ATRA.

5 Figure 10A is a series of schematics showing the indicated retinoic acid compounds and β -carotene.

Figure 10B is a graph showing the concentration of free particles of Pin1 as a function of concentration of the indicated compounds.

Figure 10C is a table summarizing the results of Fig. 10B.

Figure 10D is a series of schematics showing compounds that modulate the retinoic acid receptor.

10 Figure 11A is a schematic of the chemical structure of all-trans retinoic acid and acitretin.

Figure 11B is a graph showing the concentration of free particles of Pin1 as a function of concentration of the indicated compounds.

Figure 11C is a Western blot showing Pin1 and tubulin protein levels in cells treated with the different concentrations of the indicated compound.

15 Figure 11D is a graph showing cell viability as a function of the concentration of the indicated compounds.

Figure 12 is a graph showing cell number as a function of the concentration of the indicated compound or combination of compounds. For data points corresponding to the combination of compounds, the concentration value corresponds to the concentration of each individual compound in the
20 combination.

Figure 13A is a Western blot showing Pin, Her2, and actin protein levels in human breast cancer cells treated with the siRNA to inhibit Pin1.

Figure 13B is a graph showing cell number as a function of inhibiting Pin1 in cells by siRNA.

Detailed Description of the Invention

In general, the invention features methods of treating a proliferative disorder characterized by elevated Pin1 marker levels in a subject by administering a retinoic acid compound. Additionally, the invention features methods of treating proliferative disorders (e.g., proliferative disorders characterized by elevated Pin1 marker levels) by administering a retinoic acid compound in combination with one or more additional anti-proliferative compounds or other anti-cancer therapies.

Inhibitors of Pin1 (e.g., retinoic acid compounds) are useful for treating proliferative disorders (e.g., disorders characterized by increased Pin1 activity). Furthermore, because Pin1 acts in several different oncogenic pathways, Pin1 inhibition would be expected to behave synergistically with many anti-proliferative compounds

I. Pin1

Phosphorylation on serine/threonine-proline motifs restrains cis/trans prolyl isomerization, and also creates a binding site for the essential protein Pin1. Pin1 binds and regulates the activity of a defined subset of phosphoproteins, as well as participating in the timing of mitotic progression. Both structural and functional analyses have indicated that Pin1 contains a phosphoserine/threonine-binding module that binds phosphoproteins, and a catalytic activity that specifically isomerizes the phosphorylated phosphoserine/threonine-proline. Both of these Pin1 activities are essential for Pin1 to carry out its function in vivo.

Pin1 is overexpressed in human cancer samples and the levels of Pin1 are correlated with the aggressiveness of tumors. We have found that a potent anticancer reagent with an unknown mechanism potently and reversibly inhibits Pin1 isomerase activity. Moreover, inhibition of Pin1 by various approaches, including the Pin1 inhibitor, Pin1 antisense polynucleotides, or genetic depletion, kills human and yeast dividing cells by inducing premature mitotic entry and apoptosis. Thus, upon phosphorylation, Pin1 latches onto phosphoproteins and twists the peptide bond next to the proline, which regulates the function of phosphoproteins and participates in controlling the timing of mitotic progression.

Pin1 is highly conserved and contains a protein-interacting module, called WW domain, and a catalytically active peptidyl-prolyl isomerase (PPIase). Pin1 is structurally and functionally distinct from members of two other well-characterized families of PPIases, the cyclophilins and the FKBP. PPIases are ubiquitous enzymes that catalyze the typically slow prolyl isomerization of proteins, allowing

relaxation of local energetically unfavorable conformational states. Phosphorylation on Ser/Thr residues immediately preceding Pro not only alters the prolyl isomerization rate, but also creates a binding site for the WW domain of Pin1. The WW domain acts a novel phosphoserine-binding module targeting Pin1 to a highly conserved subset of phosphoproteins. Furthermore, Pin1 displays a unique phosphorylation-dependent PPIase that specifically isomerizes phosphorylated Ser/Thr-Pro bonds and regulates the function of phosphoproteins.

Taken together, these results indicate that the Pin1 subfamily of enzymes is a diagnostic and therapeutic target for diseases characterized by uncontrolled cell proliferation, primarily malignancies.

II. Measurement of PIN1 marker levels

The present invention pertains to the treatment of proliferative diseases identified as coinciding with elevated Pin1 marker levels with retinoic acid compounds. In some aspects, the invention features the determination of Pin1 marker levels in a subject; where retinoic acid is administered in subjects where Pin1 marker levels are determined to be elevated. In other aspects, the invention can also feature the measurement of Pin1 marker levels subsequent to the administration of retinoic acid compounds in order to evaluate the progress of therapy in treating the proliferative disorder.

Accordingly, one aspect of the present invention relates to diagnostic assays for measuring levels of Pin1 marker, as well as Pin1 activity, in the context of a biological sample (e.g., tumor samples, blood, urine, biopsies, lymph, saliva, phlegm, and pus) to thereby determine whether an individual is a candidate for treatment with a retinoic acid compound. The invention features both treatment of subjects exhibiting symptoms of a proliferative disorder and individuals at risk for developing a proliferative disorder.

Diagnostic Assays

An exemplary method for detecting the presence or absence of Pin1 protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting Pin1 protein or a nucleic acid (e.g., mRNA, genomic DNA) that encodes Pin1 protein such that the presence of Pin1 protein or nucleic acid is detected in the biological sample. A preferred agent for detecting Pin1 mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to Pin1 mRNA or DNA. The nucleic acid probe can be, for example, a Pin1 nucleic acid or a corresponding nucleic acid such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length which is capable of specifically hybridizing under stringent

conditions to Pin1 mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

A preferred agent for detecting Pin1 marker is an antibody capable of binding to Pin1 protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled," with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin.

With respect to antibody-based detection techniques, one of skill in the art can raise anti-Pin1 antibodies against an appropriate immunogen, such as isolated and/or recombinant Pin1 or a portion or fragment thereof (including synthetic molecules, such as synthetic peptides) using no more than routine experimentation. Synthetic peptides can be designed and used to immunize animals, such as rabbits and mice, for antibody production. The nucleic and amino acid sequence of Pin1 is known (Hunter et al., WO 97/17986 (1997); Hunter et al., U.S. Pat. Nos. 5,952,467 and 5,972,697, the teachings of all of which are hereby incorporated by reference in their entirety) and can be used to design nucleic acid constructs for producing proteins for immunization or in nucleic acid detection methods or for the synthesis of peptides for immunization.

Conditions for incubating an antibody with a test sample can vary depending upon the tissue or cellular type. Incubation conditions can depend on the format employed in the assay, the detection methods employed, and the type and nature of the antibody used in the assay. One skilled in the art will recognize that any one of the commonly available immunological assay formats (such as radioimmunoassays, enzyme-linked immunosorbent assays, diffusion based Ouchterlony, or rocket immunofluorescent assays) can readily be adapted to employ the antibodies of the present invention. Examples of such assays can be found in Chard, "An Introduction to Radioimmunoassay and Related Techniques," Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock et al., "Techniques in Immunocytochemistry," Academic Press, Orlando, Fla. Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, "Practice and Theory of enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology," is Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The detection method of the invention can be used to detect Pin1 mRNA, protein, or genomic DNA in a biological sample in vitro as well as in vivo. For example, in vitro techniques for detection of Pin1 mRNA include Northern hybridizations and in situ hybridizations. In vitro techniques for detection of Pin1 protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, immunofluorescence, or quantitative sequencing reactions. In vitro techniques for detection of Pin1 genomic DNA include Southern hybridizations. The detection of genomic mutations in Pin1 (or other genes that effect Pin1 marker levels) can be used to identify inherited or somatic mutations. Furthermore, in vivo techniques for detection of Pin1 protein include introducing into a subject a labeled anti-Pin1 antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In another embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting Pin1 marker such that the presence of Pin1 marker is detected in the biological sample, and comparing the presence of Pin1 marker in the control sample with the presence of Pin1 marker in the test sample.

The immunological assay test samples of the present invention may include cells, protein or membrane extracts of cells, blood or biological fluids such as ascites fluid or brain fluid (e.g., cerebrospinal fluid). The test sample used in the above-described method is based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is capable with the system utilized. The invention also encompasses kits for detecting the presence of Pin1 in a biological sample. For example, the kit can comprise a labeled compound or agent capable of detecting Pin1 protein or mRNA in a biological sample; means for determining the amount of Pin1 in the sample; and means for comparing the amount of Pin1 in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect Pin1 protein or nucleic acid.

Pin1 marker levels can also be measured in an assay designed to evaluate a panel of target genes, e.g., a microarray or multiplex sequencing reaction. In the embodiments of the invention described herein, well known biomolecular methods such as northern blot analysis, RNase protection assays,

southern blot analysis, western blot analysis, in situ hybridization, immunocytochemical procedures of tissue sections or cellular spreads, and nucleic acid amplification reactions (e.g., polymerase chain reactions) may be used interchangeably. One of skill in the art would be capable of performing these well-established protocols for the methods of the invention. (See, for example, Ausubel, et al., "Current
 5 Protocols in Molecular Biology," John Wiley & Sons, NY, N.Y. (1999)).

Diagnostic assays can be carried out in, e.g., subjects diagnosed or at risk of a proliferative disorder. Such disorders include, without limitation, leukemias (e.g., acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic
 10 myelocytic leukemia, chronic lymphocytic leukemia), polycythemia vera, lymphoma (Hodgkin's disease, non-Hodgkin's disease), Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors such as sarcomas and carcinomas (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma,
 15 rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer,
 20 testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma).

25 *Prognostic Assays*

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant Pin1 expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with Pin1 marker
 30 (e.g., a proliferative disorder). Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant Pin1 expression or activity in which a test sample is obtained from a subject and Pin1 protein or nucleic acid (e.g., mRNA, genomic DNA) is detected, wherein the presence of

Pin1 protein or nucleic acid is diagnostic for a subject having or at risk of developing a Pin1-associated disorder and are, therefore, susceptible to treatment with a retinoic acid compound.

Furthermore, the present invention provides methods for determining whether a subject can be effectively treated with a retinoic acid compound for a disorder associated with aberrant Pin1 expression or activity in which a test sample is obtained and Pin1 protein or nucleic acid expression or activity is detected (e.g., wherein the abundance of Pin1 protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder Pin1-associated disorder).

In one embodiment, the present invention provides methods for determining Pin1 post-translational modifications. For example, phosphorylation of Pin1 on Ser71 in the catalytic active site by the tumor suppressor DAPK1 completely inhibits Pin1 catalytic activity and cell function to promote oncogenesis. More importantly, phosphorylation of Pin1 on Ser71 in the catalytic active site also prevents retinoic acid compounds from binding to Pin1 active site and induce Pin1 degradation and to inhibit Pin1 function. Therefore, by detecting reduced Ser71 phosphorylation using phospho-specific Pin1 antibodies that we have generated can be a method to select patients for RA treatments and to explain some patients may not respond to RA. Because aberrantly proliferating cells exhibit reduced Ser71 phosphorylation, these cells are more sensitive to RA treatments compared to normal cells.

The methods of the invention can also be used to detect genetic alterations in a Pin1 gene, thereby determining if a subject with the altered gene is at risk for a disorder associated with the Pin1 gene and, consequently, a candidate for retinoic acid therapy. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a Pin1-protein, or the mis-expression of the Pin1 gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a Pin1 gene; 2) an addition of one or more nucleotides to a Pin1 gene; 3) a substitution of one or more nucleotides of a Pin1 gene, 4) a chromosomal rearrangement of a Pin1 gene; 5) an alteration in the level of a messenger RNA transcript of a Pin1 gene, 6) aberrant modification of a Pin1 gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a Pin1 gene, 8) a non-wild type level of a Pin1-protein, 9) allelic loss of a Pin1 gene, and 10) inappropriate post-translational modification of a Pin1-protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting alterations in a Pin1 gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject, e.g., a cardiac tissue sample.

In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g., U.S. Pat. Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran et al. (1988) Science 241:1077-1080; and Nakazawa et al. (1994) Proc. Natl. Acad. Sci. USA 91:360-364), the latter of which can be particularly useful for detecting point mutations in the Pin1-gene (see Abravaya et al. (1995) Nucleic Acids Res 23:675-682). This method can include the steps of collecting a sample from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a Pin1 gene under conditions such that hybridization and amplification of the Pin1-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli, J. C. et al., (1990) Proc. Natl. Acad. Sci. USA 87:1874-1878), transcriptional amplification system (Kwoh, D. Y. et al, (1989) Proc. Natl. Acad. Sci. USA 86:1173-1177), Q-Beta Replicase (Lizardi, P. M. et al. (1988) Bio-Technology 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

In an alternative embodiment, mutations in a Pin1 gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Pat. No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in Pin1 can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M. T. et al. (1996) Human Mutation 7: 244-255; Kozal, M. J. et al. (1996) Nature Medicine 2: 753-759). For example, genetic mutations in Pin1 can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M. T. et al. supra.

Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the Pin1 gene and detect mutations by comparing the sequence of the sample Pin1 with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxam and Gilbert ((1977) *Proc. Natl. Acad. Sci. USA* 74:560) or Sanger ((1977) *Proc. Natl. Acad. Sci. USA* 74:5463). It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen et al. (1996) *Adv. Chromatogr.* 36:127-162; and Griffin et al. (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the Pin1 gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers et al. (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes formed by hybridizing (labeled) RNA or DNA containing the wild-type Pin1 sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton et al. (1988) *Proc. Nat Acad Sci USA* 85:4397; Saleeba et al. (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes)

in defined systems for detecting and mapping point mutations in Pin1 cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu et al. (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a Pin1 sequence, e.g., a wild-type Pin1
5 sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Pat. No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in Pin1 genes. For example, single strand conformation polymorphism (SSCP) may be used to detect
10 differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita et al. (1989) *Proc Natl. Acad. Sci. USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; and Hayashi (1992) *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control Pin1 nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of
15 even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen et al. (1991) *Trends Genet.* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE)
20 (Myers et al. (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature
25 gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and
30 then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki et al. (1986) *Nature* 324:163); Saiki et al. (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different

mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs et al. (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner et al. (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini et al. (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, e.g., in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a Pin1 gene.

Furthermore, any cell type or tissue in which Pin1 is expressed may be utilized in the prognostic assays described herein.

As with the diagnostic assay described above, prognostic assays of Pin1 activity can be included as part of a panel of target genes.

Additional methods of detecting Pin1 activity and diagnosing Pin1 related disorders are disclosed in U.S. Patent Application Publication Nos.: 2009/0258352, 2008/0214470, 2006/0074222, 2005/0239095, US2002/0025521, U.S. Patent No. 6,495,376, and PCT Application Publication No. WO02/065091, each of which is hereby incorporated by reference in its entirety.

The present invention also features methods and compositions to diagnose, treat and monitor the progression of a disorder described herein (e.g., a cellular proliferation disorder) by detection and measurement of, for example, Pin1 substrates (or any fragments or derivatives thereof) containing a phosphorylated Ser/Thr-Pro motif in a cis or trans conformation, as described in U.S. Patent Provisional Application No. 61/255,431, which is hereby incorporated by reference in its entirety. The methods can include measurement of absolute levels of the Pin1 substrate (examples of which are listed in Tables 2, 3

and 4) in a *cis* or *trans* conformation as compared to a normal reference., using conformation specific antibodies. For example, a serum level or level in a biopsy of a Pin1 substrate in the *cis* or *trans* conformation that is less than 5 ng/ml, 4 ng/ml, 3 ng/ml, 2 ng/ml, or less than 1 ng/ml serum or a biopsy is considered to be predictive of a good outcome in a patient diagnosed with a disorder (e.g., a disorder associated with a deregulation of Pin1 activity). A serum level of the substrate in the *cis* or *trans* conformation that is greater than 5 ng/ml, 10 ng/ml, 20 ng/ml, 30 ng/ml, 40 ng/ml, or 50 ng/ml is considered diagnostic of a poor outcome in a subject already diagnosed with a disorder, e.g., associated with a deregulation of Pin1 activity.

For diagnoses based on relative levels of substrate in a particular conformation (e.g., a Pin1 substrate in the *cis* or *trans* conformation), a subject with a disorder (e.g., a disorder associated with a deregulation of PPase activity) will show an alteration (e.g., an increase of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more) in the amount of the substrate in, for example, the *cis* conformation. A normal reference sample can be, for example, a prior sample taken from the same subject prior to the development of the disorder or of symptoms suggestive of the disorder, a sample from a subject not having the disorder, a sample from a subject not having symptoms of the disorder, or a sample of a purified reference polypeptide in a given conformation at a known normal concentration (i.e., not indicative of the disorder).

Standard methods may be used to measure levels of the substrate in any bodily fluid, including, but not limited to, urine, blood, serum, plasma, saliva, amniotic fluid, or cerebrospinal fluid. Such methods include immunoassay, ELISA, Western blotting, and quantitative enzyme immunoassay techniques.

For diagnostic purposes, the conformation-specific antibodies may be labeled. Labeling of the antibody is intended to encompass direct labeling of the antibody by coupling (e.g., physically linking) a detectable substance to the antibody, as well as indirect labeling the antibody by reacting the antibody with another reagent that is directly labeled. For example, the antibody can be labeled with a radioactive or fluorescent marker whose presence and location in a subject can be detected by standard imaging techniques.

The diagnostic methods described herein can be used individually or in combination with any other diagnostic method described herein for a more accurate diagnosis of the presence or severity of a disorder (e.g., a cellular proliferation disorder or a neurological disorder). Examples of additional methods for diagnosing such disorders include, e.g., examining a subject's health history, immunohistochemical staining of tissues, computed tomography (CT) scans, or culture growths.

Monitoring the effects of retinoic acid treatment, and disease progression

In one embodiment, the present invention features a method for monitoring the effectiveness of treatment of a subject with a retinoic acid compound comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the compound; (ii) detecting the level of expression or activity of a Pin1 protein, Pin1 phosphorylation on Ser71, mRNA, or genomic DNA in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the Pin1 protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the Pin1 protein, mRNA, or genomic DNA in the pre-administration sample with the Pin1 protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the retinoic acid compound to the subject accordingly. According to such an embodiment, Pin1 expression, phosphorylation or activity may be used as an indicator of the effectiveness of the retinoic acid compound, even in the absence of an observable phenotypic response.

In another embodiment, the diagnostic methods described herein can also be used to measure the levels of, for example, polypeptides (e.g., Pin1 substrates listed in Tables 2, 3 and 4) with pSer/Thr-Pro motifs in the *cis* or *trans* conformation using conformation specific antibodies. The methods can include repeated measurements, using conformation specific antibodies, for diagnosing the disorder and monitoring the treatment or management of the disorder. In order to monitor the progression of the disorder in a subject, subject samples can be obtained at several time points and conformation specific antibodies can be used to monitor the levels of *cis* and *trans* isomers of Pin1 substrates (listed in Tables 2, 3 and 4). For example, the diagnostic methods can be used to monitor subjects during chemotherapy (e.g., therapy with a retinoic acid compound or other agent described herein). In this example, serum samples from a subject can be obtained before treatment with a chemotherapeutic agent, again during treatment with a chemotherapeutic agent, and again after treatment with a chemotherapeutic agent. In this example, the level of Pin1 substrate with a pSer/Thr-Pro motif in the *cis* conformation in a subject is closely monitored using the conformation-specific antibodies of the invention and, if the level of Pin1 substrate with a pSer/Thr-Pro motif in the *cis* conformation begins to increase during therapy, the therapeutic regimen for treatment of the disorder can be modified as determined by the clinician (e.g., the dosage of the therapy may be changed or a different therapeutic may be administered). The monitoring methods of the invention may also be used, for example, in assessing the efficacy of a particular drug or therapy in a subject, determining dosages, or in assessing progression, status, or stage of the infection.

Table 2. General Pin1 Substrates

Substrate (GenBank Accession Number)	Targeting Site(s)	Functional Consequence of PPIase Activity of Pin1 Upon Substrate
<u>G2/M and Mitotic Regulation</u>		
NIMA (P11837)	-	Regulation of mitotic function
RAB4 (NP_004569)	-	-
CDC25 (AAA58417)	pThr48/67-Pro	Dephosphorylation and regulation of activity
WEE1 (NP_003381)	pT186-P	Inhibition of WEE1 activity
PLK1 (P53350)	-	-
MYT1 (NP_004194)	-	-
CDC27 (AAH11656)	-	-
CENP-F (P49454)	-	-
Incenp (NP_064623)	-	-
RPB1 (CAA65619)	pSer5-Pro	Regulation of CTD dephosphorylation
NHERF-1 (AAA80218)	pSer279/301-P	Dephosphorylation
KRMP1 (NP_057279)	pT-1604-P	Regulation of mitotic function
CK2 (NP_808227)	Multiple pSer/Thr-Pro sites	Inhibition of kinase activity
TopoII α (NP_001058)	-	Inhibition or induction of phosphorylation
DAB2 (NP_001334)	-	Dephosphorylation
p54nrb (CAA72157)	Multiple pSer/Thr-Pro sites	-

Sil (CAC14001)	Multiple pSer/Thr-Pro sites	Regulation of function
EMI1 (NP_036309)	pS10-P	Stabilization
<u>G1/S Regulation</u>		
Cyclin D1 (NP_444284)	pT286-P	Stabilization and nuclear localization
Ki67	pT234-P	-
c-Myc (CAA46984)	pT58-P	Dephosphorylation and destabilization
Cyclin E (P24864)	pS384-P	Destabilization
<u>Growth and Oncogenic Signaling</u>		
c-Jun (AAH06175)	pS63/73-P	Transactivation
B-catenin (P35222)	pS246-P	Stabilization, protein interaction, and transactivation
Cf-2 (NP_034298)	-	Destabilization
NF- κ B (AAH33210)	pT254-P	Stabilization, protein interaction, and transactivation
RAF1 (AAA60247)	Multiple pSer/Thr-Pro sites	Dephosphorylation and prolonging activation
c-Fos (CAA24756)	Multiple pSer/Thr-Pro sites	Transactivation
RAR α (NP_001019980)	pS77-P	Stabilization and transactivation
AIB1/SRC-3	-	Transactivation and destabilization
HBx (NP_110380)	pS41-P	Stabilization and potentiation
STAT3 (NP_998827)	pS727-P	Transactivation

<u>DNA Damage, Oxidative Stress Response, and Apoptosis</u>		
p53 (BAC16799)	Multiple pSer/Thr-Pro sites	Stabilization and transactivation
Bcl-2 (NP_000648)	pS70/87-P	-
p73 (CAA72221)	Multiple pSer/Thr-Pro sites	Stabilization and transactivation
BimEL (AAC39593)	pS65-P	Stabilization
p66 ^{Shc} (AAH14158)	-	Mitochondrial import
CHE1 (P06276)	-	Destabilization
<u>Neuronal Survival and Degeneration</u>		
Tau (NP_058519)	pT231-P pT212-P	Dephosphorylation and protein interaction
APP (P05067)	pT668-P	Promotes non-amyloidogenic APP processing and reduces A β production
APP fragment	pT668-P	Increases A β production from C99 APP fragment
Synphilin-1 (AAD30362)	pS211/215-P	Protein interaction
Gephyrin (CAC81240)	pS188/194/200-P	Protein interaction
MCL1 (CAI15504)	pT163-P	Stabilization
<u>Immune Response and Asthma</u>		
NFAT (NP_666017)	-	
AUF1 (NP_112738)	-	Protein interaction
IRF3 (AAH71721)	pS339-P	Destabilization
BTK (CAI42359)	pS21/115-P	Destabilization

<u>Others</u>		
SIN2-RPD3	-	Reduces histone deacetylases
hSpt5 (NP_001124297)	-	

Table 3A Pin1 targets where Pin1 prevents protein from degradation

	PROTEIN NAME	PIN1 FUNCTION	PIN1 BINDING MOTIF	KINASE/ PHOSPHATASE	PATHWAYS/ DISEASES/ MECHANISM
1	p53	1. Stabilization; Increase p53 promoter binding activity 2. Pin1 in formation of HMWC and stabilizes p53	Ser33, Ser315, Thr181; Pro82	Chk2	Genotoxic stress, DNA damage Trophoblast invasiveness
2	Cyclin D1	Stabilization; localization and transcription	Thr286	Block Socs-1 ub-mediated proteolysis	Cell proliferation/cancers

3	Tau	1. Dephosphorylated at Thr231 2. Pin1 knockdown or KO increased WT tau protein stability in vitro	Thr231		Tauopathy
4	β-Catenin	Stability; localization and transactivation	Ser246	Could stabilize β -catenin by inhibiting GSK-3 β dependent degradation	Cell proliferation/cancers
5	c-Jun	Transcriptional activity	Ser63/Ser73	JNK	Breast cancer; AML
6	p65/NF-kB	Nuclear translocation; stability	Thr254		Cytokines; Hepatocyte NF-kappaB activation
7	p73	Stabilizing, transcriptional activity	Ser412, Thr442 and Thr482	c-Abl and p300	Genotoxic stress

8	Synphilin-1	Facilitates Lewy Body formation; stabilizes alpha-synuclein	Ser211 and Ser215	casein kinase II	Parkinson disease
9	c-Fos	Transcriptional activation	C-terminal	ERK	
10	Sil	No impact on Sil spindle checkpoint			Unknown
11	p54nrb		Thr412, Thr430 and Thr452	Cdk1	
12	Bruton tyrosine kinase (Btk)	Mediates Btk degradation	Ser21 an Ser115		Tyrosine kinase
13	AUF1	Regulates GM-CSF mRNA; AUF1, AU rich element-binding protein	Ser83		

14	BIM_{EL}	Stabilize BIM _{EL} and induce apoptosis	Ser65	JIP3, MKK7 and JNK	Neuronal apoptosis
15	Mcl-1 (Myeloid cell leukemia sequence-1)	1. Pin1 inhibits Mcl-1 ubiquitination 2. Stabilizes Mcl-1	Thr163; Thr92	JNK3 induces Mcl-1 degradation by counting the protective binding of Pin1 Erk phosphorylates Thr92 and Thr163	Oligodendrocyte apoptosis; Mcl-1 mediated chemoresistance; breast cancer
16	HBx; Hepatitis B virus encoded protein X	Pin1 overexpression increased the protein stability of HBx	Ser41	Pin1 binds HBx and enhance hepatocarcinogenesis in HBV-infected hepatocytes	Hepatocarcinogenesis

17	Origin recognition complex, subunit 1 (ORC1)	Prevents degradation of ORC1 by inactivating mitotic APC complex		APC; Topo II	Mitosis; chromosome segregation and for reprogramming replicons
18	Bcl-2	1. Induce changes in the bioactivity of Bcl-2; 2. Prevents dephosphorylation of Bcl-2		Possibly mediated by cdc2	
19	Erb2	Stabilize ErbB2	Ubiquitinated erbB2	ErbB2 pathway; ubiquitin mediated degradation	Her2-positive breast cancer
20	PPARγ	Prevents the polyubiquitination of PPAR γ through ubiquitin- proteasome pathway	Ser84	Ras mediated kinase	Macrophages mediated atherosclerosis

21	Cep55	Increased Cep55 stability	Ser425, Ser428	Cdk1, Plk1	Mitosis and cytokinesis
22	Spt23	Ess1 stabilizes Spt23	Ser654		Unsaturated fatty-acid synthesis
23	p27	Pin1 protects p27 from degradation through polyubiquitination mechanism.	Thr187	Cdk2	Cell cycle; Cancer
24	Akt	Regulates Akt protein stability	Thr92, Thr450		Oncogenesis
25	HTLV Tax protein	Increased Tax protein expression; inhibits Tax protein degradation	Ser160		Pathogenesis of Human T-cell leukemia virus type 1 (HTLV-1) related diseases
26	Nanog	Leads to Nanog stability	Ser52, Ser65		Stem cells pluripotency; cell renewal

27	Viral Integrase	Stabilized phospho-HIV-1 integrase	Ser57	JNK	HIV-1 cDNA integration and infection
----	--------------------	---------------------------------------	-------	-----	---

Table 3B Pin1 targets where Pin1 enhances degradation of phosphorylated proteins

	PROTEIN NAME	PIN1 FUNCTION	PIN1 BINDING MOTIF	KINASE/PHOSPHATASE	PATHWAYS/DISEASES/MECHANISM
1	c-Myc	Enhances c-Myc degradation	Thr58		Tumorigenesis
2	Cyclin E	Depletion of Pin1 upregulates cellular level of cyclin E	Ser384	Cyclin E-Cdk2 complex	Cell cycle, genomic instability and tumorigenesis
3	SRC-3/AIB1	Enhances SRC-3 degradation	N/A	Steroid receptor	
4	RAR α	Induces its degradation	Ser77		Retinoic acid receptor;
5	IRF3	Promote its degradation via the ubiquitin-proteasome pathway.	Ser339		Host antiviral responses during virus infection

6	Che-1	Conformational changes induced by Pin1 are requested for Che-1/HDM2 interaction	Thr144		p53 transcription; DNA damage apoptotic pathway
7	Pim-1 protein kinase	Binding of Pin1 leads to a decrease in the protein level of Pim-1	N/A	PP2A	Elevated in lymphomas leukemias and prostate cancer through c-Myc pathway
8	Promyelocytic leukemia protein (PML)	Binds to phosphorylated C terminus of PML and enhances PML degradation	C-terminal of PML (Ser403, Ser505, Ser518 and Ser527)		Breast cancer; hydrogen peroxide-induced death; cell proliferation
9	FOXO	A novel negative FOXO regulator, interconnecting FOXO phosphorylation and monoubiquitination in response to cellular stress			Oxidative stress

		to regulate p27				
10	Silencing mediator for retinoic acid and thyroid hormone receptor (SMRT)	Pin1 destabilizes SMRT	Ser1241, Thr1445, Ser1469	Her2/Neu/ErbB2 receptor; Cdk2	Human cancer	
11	TRF-1	Pin1 inhibition resulted in decrease TRF1 degradation;	Thr149	Cdk	Cancer; ageing	
12	G protein-coupled receptor kinase 2 (GRK2)	Promotes GRK2 degradation	Ser670	CDK2-cyclinA	Cell cycle progression; p53 response and the induction of apoptosis	
13	SF-1	Pin1 promotes SF-1	Ser203	CDK7	Gonadotropin beta-	

		ubiquitination and degradation			subunit gene transcription
14	Sulfotransferase 4A1 (SULT4A1)	Pin1 destabilizes SULT4A1	Thr8, Thr11	ERK1 and PP2A	Metabolism of endogenous and exogenous compounds
15	Smad2/Smad3	Reduced Smad2/3 protein levels	Thr179, Ser204, Ser208, Ser213 (Smad3 linker domain)	Smurf2 with Smads and enhanced Smad ubiquitination	TGF- β signaling
16	MEF2C	Pin1 decreases MEF2C stability	Ser98/Ser110		Muscle terminal differentiation

Table 3C Pin1 targets where Pin1 regulates target phosphorylation/dephosphorylation/other modifications

	PROTEIN NAME	PIN1 FUNCTION	PIN1 BINDING MOTIF	KINASE/ PHOSPHATASE	PATHWAYS/ DISEASES/ MECHANISM
1	Tau	Enhance dephosphorylation at Thr231	Thr231		Neuronal differentiation; stress induced; Alzheimer disease
2	Raf-1	Dephosphorylation by PP2A	N.A.	Ras/MAP Kinase	
3	Type-1 protein phosphatase Inhibitor-2 (I-2)	Cycling phosphorylation in mitotic	Thr72	CDK1-cyclin B	Entry and exit in mitotic

4	TGF- β 1 mRNA	Decay, accumulation and translation of TGF- β 1 mRNA in Eosinophils			Chronic asthma
5	K-homology splicing regulator protein (KSRP)	parathyroid hormone (PTH) mRNA stability	Ser181		

Table 4. Pin1-targets for promoting tumorigenesis

Substrates	Function	Binding Sites	Effects	Activation/ Inactivation
AIB1/SRC3	Transactivator	-	Activity	+
Akt	Protein kinase	pThr ^{92/450} -Pro	Stability	+
Bax	Apoptosis	pThr ¹⁶⁷ -Pro	Activity	-
Bcl-2	Antiapoptotic protein	-	Stability	-
Btk	Tyrosine kinase	pSer ^{21/115} -Pro	Stability	-
β-catenin	Transcription factor	pSer ²⁴⁶ -Pro	Localization, stability	+
C/EBP	Transcription factor	-	Activity	-
Cyclin D1	Transcription factor	pThr ²⁸⁶ -Pro	Localization, stability	+
Daxx	Apoptosis	pSer ¹⁷⁸ -Pro	Stability	-
FAK	Tyrosine kinase	pSer ⁹¹⁰ -Pro	Activity	+
c-Fos	Transcription factor	-	Activity	+
FOXO4	Transcription factor	-	Localization, activity	-
GRK2	G protein receptor	pSer ⁶⁷⁰ -Pro	Stability	-
Hbx	Transactivator	pSer ⁴¹ -Pro	Activity, stability	+
c-Jun	Transcription factor	pSer ^{63/73} -Pro	Activity, stability	+
Mcl-1	Apoptosis	pThr ^{92/163} -Pro	Stability	+
c-Myb	Transactivator	pSer ⁵²⁸ -Pro	Activity	+

Neu	Growth factor receptor	-	Stability	+
NF-κB	Transcription factor	pThr ²⁵⁴ -Pro	Localization, stability	+
Notch1	Growth factor	-	Activity	+
p70S6K	Ribosomal S6 kinases	-	Activity	+
p53	Transcription factor	-	Activity, stability	-*
Plk1	Mitotic kinase	-	Binding activity	+
PML	Transcription factor	pSer ^{403/505/518/527} - Pro	Stability	-
Raf-1	Protein kinase	-	Activity	+
RARα	Transcriptional regulator	pSer ⁷⁷ -Pro	Stability	-
V-Rel	Transcription factor	pThr ²⁵⁴ -Pro	Localization, stability	+
Smad	Transactivator	-	Stability	-
SMRT	Transcriptional co-repressor	pSer ^{1241/1469} , Thr ¹⁴⁴⁵ -Pro	Stability	-
Stat3	Transcription factor	pSer ⁷²⁷ -Pro	Activity	+
Tax	Viral oncoprotein	pSer ¹⁶⁰ -Pro	Activity, stability	+
Pin2/TRF 1	Telomere regulation	pThr ¹⁴⁹ -Pro	Stability	-

III. Methods of Treatment

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a proliferative disorder (e.g., a disorder associated with increased Pin1 expression or activity) with a retinoic compound.

5 Certain embodiments of the invention feature formulation of a retinoic acid compound for, e.g., controlled or extended release. Many strategies can be pursued to obtain controlled and/or extended release in which the rate of release outweighs the rate of metabolism of the therapeutic compound. For example, controlled release can be obtained by the appropriate selection of formulation parameters and ingredients (e.g., appropriate controlled release compositions and coatings). Examples include single or
10 multiple unit tablet or capsule compositions, oil solutions, suspensions, emulsions, microcapsules, microspheres, nanoparticles, patches, and liposomes. The release mechanism can be controlled such that the retinoic acid compound is released at period intervals, the release could be simultaneous, or a delayed release of one of the agents of the combination can be affected, when the early release of one particular agent is preferred over the other.

15 Certain embodiments of the invention feature a deuterated retinoic acid compound that is made by replacing some or all hydrogen with deuterium using state of the art techniques (e.g., as described herein and at www.concertpharma.com).

Prophylactic Methods

20 In one aspect, the invention provides a method for preventing a proliferative disorder in a subject by administering to the subject a retinoic acid compound. Subjects at risk for a disease which is caused or contributed to by aberrant Pin1 expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a retinoic acid compound can occur prior to the manifestation of symptoms characteristic of the Pin1 aberrancy, such
25 that a disease or disorder is prevented or, alternatively, delayed in its progression.

Combination Therapies

Anti-proliferative and other anti-cancer compounds (e.g., anti-angiogenic compounds) are useful for treating proliferative disorders in combination with the retinoic acid compounds of the invention.

30 Anti-proliferative agents that can be used in combination with a retinoic acid compound include, without limitation, microtubule inhibitors, topoisomerase inhibitors, platins, alkylating agents, and anti-metabolites. Particular anti-proliferative agents that are useful in the methods and compositions of the invention include, without limitation, paclitaxel, gemcitabine, doxorubicin, vinblastine, etoposide, 5-

fluorouracil, carboplatin, altretamine, aminoglutethimide, amsacrine, anastrozole, azacitidine, bleomycin, busulfan, carmustine, chlorambucil, 2-chlorodeoxyadenosine, cisplatin, colchicine, cyclophosphamide, cytarabine, cytoxan, dacarbazine, dactinomycin, daunorubicin, docetaxel, estramustine phosphate, floxuridine, fludarabine, gentuzumab, hexamethylmelamine, hydroxyurea, ifosfamide, imatinib, interferon, irinotecan, lomustine, mechlorethamine, melphalen, 6-mercaptopurine, methotrexate, mitomycin, mitotane, mitoxantrone, pentostatin, procarbazine, rituximab, streptozocin, tamoxifen, temozolomide, teniposide, 6-thioguanine, topotecan, trastuzumab, vincristine, vindesine, and vinorelbine. The ability of a compound to inhibit the growth of a neoplasm can be assessed using known animal models.

Compounds which are known to interact with other proteins implicated in Pin1 signaling pathways can also be useful in combination with a retinoic acid compound (see, e.g., the targets and compounds in Table 5).

Table 5.

Target	Target Class	Representative Antagonist
AKT	Kinase	MK-2206
Cyclin D1	Cyclin	ON 013105
HER2/Neu (ErbB-2)	Kinase	Herceptin
NF- κ B	Transcription Factor	RTA 402
Plk	Kinase	BI 2536
Raf-1	Kinase	Sorafenib
Stat3	Transcription Factor	ISIS-STAT3Rx
α 5 β 1-SIS-STAT	Adhesion	Nucleic Acid-Based Rx in Enzon Program at Santaris

Such compounds can act synergistically with a retinoic acid compound. Additionally, co-administration with a retinoic acid compound may result in the efficacy of the anti-proliferative compound at lower (and thus safer) doses (e.g., at least 5% less (e.g., at least 10%, 20%, 50%, 80%, 90%, or even 95%) less than when the anti-proliferative compound is administered alone.

Therapy according to the invention may be performed alone or in conjunction with another therapy and may be provided at home, the doctor's office, a clinic, a hospital's outpatient department, or a hospital. Treatment optionally begins at a hospital so that the doctor can observe the therapy's effects closely and make any adjustments that are needed, or it may begin on an outpatient basis. The duration of

the therapy depends on the type of disease or disorder being treated, the age and condition of the patient, the stage and type of the patient's disease, and how the patient responds to the treatment. Additionally, a person having a greater risk of developing an proliferative disease may receive treatment to inhibit or delay the onset of symptoms.

5 Routes of administration for the various embodiments include, but are not limited to, topical, transdermal, nasal, and systemic administration (such as, intravenous, intramuscular, subcutaneous, inhalation, rectal, buccal, vaginal, intraperitoneal, intraarticular, ophthalmic, otic, or oral administration). As used herein, "systemic administration" refers to all nondermal routes of administration, and specifically excludes topical and transdermal routes of administration.

10 In combination therapy (e.g., a retinoic acid compound with a second anti-proliferative agent), the dosage and frequency of administration of each component of the combination can be controlled independently. For example, one compound may be administered three times per day, while the second compound may be administered once per day. Alternatively, one compound may be administered earlier and the second compound may be administered later. Combination therapy may be given in on-and-off
15 cycles that include rest periods so that the patient's body has a chance to recover from any as yet unforeseen side effects. The compounds may also be formulated together such that one administration delivers both compounds.

Each compound of the combination may be formulated in a variety of ways that are known in the art. For example, the first and second agents may be formulated together or separately. Desirably, the
20 first and second agents are formulated together for the simultaneous or near simultaneous administration of the agents. Such co-formulated compositions can include the two drugs together in the same pill, ointment, cream, foam, capsule, liquid, etc. It is to be understood that, when referring to the formulation of combinations of the invention, the formulation technology employed is also useful for the formulation of the individual agents of the combination, as well as other combinations of the invention. By using
25 different formulation strategies for different agents, the pharmacokinetic profiles for each agent can be suitably matched.

The individually or separately formulated agents can be packaged together as a kit. Non-limiting examples include kits that contain, e.g., two pills, a pill and a powder, a suppository and a liquid in a vial, two topical creams, ointments, foams etc. The kit can include optional components that aid in the
30 administration of the unit dose to patients, such as vials for reconstituting powder forms, syringes for injection, customized IV delivery systems, inhalers, etc. Additionally, the unit dose kit can contain instructions for preparation and administration of the compositions. The kit may be manufactured as a single use unit dose for one patient, multiple uses for a particular patient (at a constant dose or in which

the individual compounds may vary in potency as therapy progresses); or the kit may contain multiple doses suitable for administration to multiple patients ("bulk packaging"). The kit components may be assembled in cartons, blister packs, bottles, tubes, and the like.

IV. Experimental Results

We have identified a series of nanomolar inhibitors that binds only to the catalytically active PPIase domain of Pin1. Upon electroporation or microinjection, these Pin1 peptide inhibitors block HeLa division, and this block is rescued by co-injected Pin1, indicating that the compounds are highly specific and potent. Using the above Pin1 peptide inhibitors, we have developed a high-throughput screen for identifying Pin1 inhibitors using a single-step fluorescence polarization-based displacement assay (FP-HTS). The FP-HTS detects molecules that compete for the substrate binding to the catalytic active site, measures ligand binding under equilibrium conditions, and does not suffer from product inhibition. The HF488 fluorescent probes for the FP assay contain only four residue core structure of Bth-L-phos.Thr-Pip-Nal (pTide), with a K_d of 258 nM for Pin1, was synthesized by Anaspec. We performed FP-HTS in a 384-well plate format with full length Pin1 and produced robust FP, resulting in a 6-7 fold increase in polarization degree values, using a Synergy II plate reader. This novel FP-HTS showed robust and reproducible performance. The assay can tolerate up to 10% DMSO. The Z' is around 0.70 and is consistent for day-to-day performance. The coefficient of variation is in the range of 4-5%. More importantly, we have shown that both pTide, the unlabeled Pin1 peptide as the positive control in this project, and juglone, the Pin1 small molecule inhibitor, displaced HF488 probe from Pin1. Although it is difficult to determine the K_d for the covalent and irreversible inhibitor juglone, the K_d for pTide was ~250 nM, similar to that derived from PPIase-based assays.

We used the FP-HTS with a 5 nM probe and 200 nM Pin1 to conduct a pilot screen on a selected set of chemical libraries. We obtained the resulting potential positive hits and grouped them into 3 classes according to the Z-score, which is folds of standard deviation below the mean. The top strong chemical is the clinically used drug 13-cis-retinoic acid (cis-RA) (Fig. 2A). cis-RA is particularly attractive for the following reasons. 1) cis-RA is not listed in promiscuous inhibitor databases 2) all-trans retinoic acid (trans-RA), the pairing compound of cis-RA, is currently used as oral prescription for patients with acute promyelocytic leukemia (AML); 3) both cis-RA and trans-RAs are used as a medication in the phase II/III clinical trial for breast cancer, and more because of its ability to kill rapidly dividing cells; and 4) although it has been reported that RAs target on retinoic acid receptor (RAR), the exact mechanism of the anticancer action is unknown. As a result, we examined whether the anti-cancer effect of RAs depends on RAR in the breast cancer cells. RAR α knockdown can only partially rescue-cis-RA-mediated cell

death, indicating that RAs may have unidentified “off-target effect.” To confirm that RAs indeed target Pin1, we examined cis-RA and trans-RA in the FP assay, and, surprisingly, found that trans-RA displayed even more prominent Pin1 inhibition than cis-RA dose and that cis-RA would eventually catch up with trans-RA in the long-term incubation with Pin1, likely due to resonance-mediated cis-trans conversion (Fig. 3). In the PPIase assay, Pin1 activity was blocked by either cis- or trans-RA in a dose-dependent manner (Fig. 4). These data confirm that the interaction between RAs and Pin1 is specific and not due to aggregation. Furthermore, both trans and cis RA blocked the association between Pin1 and DAPK1 in a dose-dependent manner, with trans being more potent (Fig. 5). These results indicate that RAs binds to Pin1 C-terminal catalytic domain because DAPK1 is known to bind this domain (Lee et al., 2011 Mol Cell in press). To determine which amino acid residue in the Pin1 catalytic domain that are important for retinoic acid binding, point mutations of Pin1 including K63A, S67E, R68/69A, H59A or S71E completely or significantly abolished trans-RA binding to Pin1 (Fig. 6A and B). Together, these data indicate that RAs inhibit Pin1 by occupying its catalytic PPIase pocket in the C-terminus and that phosphorylation of Ser67 or Ser71 inhibits RA binding to Pin1.

To further discern the causal relationship between the anti-proliferative effect of RAs on Pin1, cell viability of three breast cancer cell lines was tested with different dosages of cis- or trans-RA, of which SKBR3 and T47D exhibited preferential sensitivity to RAs with an IC50 in the nano-molar range, while the normal cell line MCF10A remained unaffected (Fig. 7A). This discrepancy between cell lines was correlated with the RAs’ ability to suppress Pin1 expression. Pin1 level was decreased by treatment of RAs in drugs-responsive SKBR3 and T47D, but not in drugs-irresponsive normal cells, MCF10A (Fig. 7B), in which Pin1 target protein, cyclin D1, was served as biomarkers of in vivo Pin1 activity. Moreover, RAs did not alter Pin1 mRNA level, but did reduce Pin1 protein stability, suggesting that RAs interact with Pin1, result in Pin1 degradation, and subsequently leads to anti-proliferation of breast cancer cells (Figs. 8A-C). To further confirm this premise, wild-type mouse embryonic fibroblast (WT MEF) and Pin1 knockout MEF (Pin1 KO MEF) were used to test trans-RA-mediated cell viability (Figs. 9A-C). As expected, Pin1 KO MEF was more resistant to trans-RA than WT MEF due to lack of drug target. In addition, Pin1 KO MEF stably expressing WT Pin1, but not W34/K63A Pin1 mutant, enabled cells to re-sensitize trans-RA. These results indicate that RAs-mediated cell death is at least in part dependent on Pin1.

As trans-RA is selected as the current lead compound, it is important to identify the essential moiety of RAs on Pin1 interaction for further lead optimization. Many commercially available retinoids were tested for Pin1 inhibition. Only those with end carboxylic group maintained Pin1 inhibition in a cell culture model, which are cis-RA, trans-RA, AC-55649, and acitretin (Figs. 10A, 10B, 11A, and 11B).

Other RA compounds that are metabolized to contain an end carboxylic group are also likely to be useful in treating proliferative disorders (Fig. 10D). In addition, compounds that modulate the retinoic acid receptor (Figs. 10D and 11A and compounds listed in Table 1) are also likely to be useful in treating proliferative disorders.

5 In addition, we have developed Cell-based assays to screen and validate Pin1 inhibitor hits. We have shown 1) that Pin1 is highly expressed in HER2-positive human breast cancer tissues; 2) that Pin1 inhibition almost completely suppresses HER2 overexpression on cell surface in human HER2+ breast cancer cell lines such as AU565 and SKBR3 cells; 3) that Pin1 inhibition greatly increases the sensitivity of HER2+ breast cancer cells to the mTOR inhibitor, but not to the HER2 inhibitor, suggesting that Pin1
10 might act on Her2 to regulate cell growth; 4) that Pin1 acts on Neu and multiple substrates in Neu-mediated oncogenic pathway; and 5) that Pin1 knockout in mice inhibits breast cancer development induced by activated Her2. Therefore, Pin1 is essential for maintaining HER2 overexpression and growth of human HER2+ breast cancer cells. Given that HER2 expression on cell surface and cell growth are readily assayed on 384-well format, we can test the ability of the hits to repress HER2 overexpression and
15 cell growth of HER2_ AU565 and SKBR3 cells, which will be treated with Pin1 prodrugs or hits, and then immunostained with Alexa 488-anti-HER2 monoclonal antibody (BioLegend), followed by automated microscopy.

We have further shown that combinations of therapeutic compounds including retinoic acid compounds are useful for treating cancer, e.g., cancer characterized by elevated Pin1 activity. Fig. 12
20 shows results obtained from the treatment of breast cancer cells overexpressing Pin1 with ATRA or Doxorubicin or their combination, followed by counting cancer cell numbers. The results show that ATRA and Doxorubicin combination dramatically increases anticancer potency and reduce the dose of each drug to inhibit cancer cell growth. Therefore, ATRA can drastically reduce dose and toxicity of Doxorubicin and other chemotherapeutic drugs.

25 We have also shown that Pin1 inhibition, using siRNA, dramatically reduces Neu/Erb2 overexpression and cell proliferation of human breast cancer cells that have Neu/Erb2 gene amplification (Figs. 13A and 13B). This provides us with a method to identify a Pin1 modulatory compound by applying a test compound to human-derived cancer cells that have Neu/Erb2 gene amplification, and determining the effect of the test compound on Neu/Erb2 overexpression and cell proliferation.

Other Embodiments

All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure that come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth, and follows in the scope of the claims.

Other embodiments are within the claims.

What is claimed is:

1. A method of treating a proliferative disease in a subject, said method comprising administering a retinoic acid compound to said subject in an amount sufficient to treat said subject, wherein said subject is determined to have elevated levels of a Pin1 marker prior to said administration.
2. A method of treating a proliferative disease in a subject, said method comprising determining Pin1 marker levels in a sample from said subject and administering a retinoic acid compound to said subject if sample is determined to have elevated Pin1 marker levels.
3. The method of claims 1 or 2, further comprising the administration of a second anti-proliferative compound.
4. The method of claim 1, wherein said Pin1 marker is reduced Ser71 phosphorylation of Pin1.
5. The method of any one of claims 1-4, further comprising determining Pin1 marker levels in said sample after said administration of a retinoic acid compound.
6. The method of any one of claims 1-5, wherein said retinoic acid compound is selected from the group consisting of 13-cis-retinoic acid and all-trans-retinoic acid.
7. The method of any one of claims 1-5, wherein said retinoic acid compound is selected from the group consisting of retinol, retinyl acetate, retinal, and AC-55640.
8. The method of any one of claims 1-5, wherein said retinoic acid compound is selected from compounds structurally similar to retinoic acid, preferably those listed in Figs. 2A, 2B, 10A, 10D, and Table 1.
9. The method of any one of claims 1-8, wherein said retinoic acid compound is deuterated.
10. The method of any one of claims 1-9, wherein said sample is selected from the group consisting of tumor samples, blood, urine, biopsies, lymph, saliva, phlegm, and pus.
11. The method of any one of claims 1-10, wherein said elevated Pin1 marker level is due to an inherited trait or a somatic mutation.
12. The method of any of claims 3 or 5-11 wherein said second anti-proliferative compound is selected from the group consisting of MK-2206, ON 013105, RTA 402, BI 2536, Sorafenib, and ISIS-STAT3Rx.

13. The method of any of claims 3 or 5-11, wherein said second anti-proliferative compound is selected from the group consisting of microtubule inhibitors, topoisomerase inhibitors, platins, alkylating agents, and anti-metabolites.

14. The method of any of claims 3 or 5-11, wherein said second anti-proliferative compound is selected from the group consisting of paclitaxel, gemcitabine, doxorubicin, vinblastine, etoposide, 5-fluorouracil, carboplatin, altretamine, aminoglutethimide, amsacrine, anastrozole, azacitidine, bleomycin, busulfan, carmustine, chlorambucil, 2-chlorodeoxyadenosine, cisplatin, colchicine, cyclophosphamide, cytarabine, cytoxan, dacarbazine, dactinomycin, daunorubicin, docetaxel, estramustine phosphate, floxuridine, fludarabine, gentuzumab, hexamethylmelamine, hydroxyurea, ifosfamide, imatinib, interferon, irinotecan, lomustine, mechlorethamine, melphalen, 6-mercaptopurine, methotrexate, mitomycin, mitotane, mitoxantrone, pentostatin, procarbazine, rituximab, streptozocin, tamoxifen, temozolomide, teniposide, 6-thioguanine, topotecan, trastuzumab, vincristine, vindesine, and vinorelbine.

15. The method of any of claims 3 or 5-14, wherein said anti-proliferative compound is administered at a low dosage or at different time.

16. The method of any of claims 3 or 5-14, wherein said anti-proliferative compound is formulated in such as a liposomal formulation or a controlled release formulation.

17. The method of any of claims 3 or 5-14, wherein said retinoic acid compound and said anti-proliferative compound are formulated together.

18. The method of any of the preceding claims, wherein said proliferative disorder is selected from the group consisting of leukemias, polycythemia vera, lymphomas, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors.

19. The method of claim 18, wherein said proliferative disorder is selected from the group consisting of acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, Hodgkin's disease, non-Hodgkin's disease, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland

carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

20. A method for identifying a Pin1 ligand comprising of:

- (i) incubating Pin1 protein with a fluorescently labeled probe, forming a Pin1-probe complex;
- (ii) adding a test compound to the incubation; and
- (iii) determining whether any substantial portion of the probe is displaced from the Pin1-probe complex by the test compound, such displacement indicating that the test compound is a Pin1 ligand.

21. A method for identifying a Pin1 modulating compound comprising of:

- (i) incubating human-derived cancer cells that have Neu/Erb2 gene amplification;
- (ii) applying a test compound to the cell; and
- (iii) determining whether Neu/Erb2 overexpression is reduced, wherein reduction of Neu/Erb2 overexpression indicates that the test compound is a Pin1 modulating compound.

Fig. 1B






Fig. 2A




13-cis-retinoic acid (13cRA)



All-trans-retinoic acid (ATRA)

Fig. 2B






Fig. 3





Fig. 4

Fig. 5







Fig. 6


A



B

Fig. 7

A



B




Fig. 8






Fig. 9 A



B C



A

13cRA

ATRA

Retinol

Retinyl acetate

Retinal

AC-55649

β -carotene

Chemical structures of various retinoids and carotenoids are shown, including 13cRA, ATRA, Retinol, Retinyl acetate, Retinal, AC-55649, and β -carotene. The structures are arranged vertically, with their names labeled to the left. The structures are drawn in a skeletal format, showing the carbon backbone and functional groups. The structures are arranged in a column, with the names of the compounds written to the left of each structure. The structures are drawn in a skeletal format, showing the carbon backbone and functional groups. The structures are arranged in a column, with the names of the compounds written to the left of each structure.




Fig. 10 D





Fig. 10 D continued









Fig. 11





Fig. 12

Fig. 13



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/29077

A. CLASSIFICATION OF SUBJECT MATTER IPC(8) - C07C 57/00 (2012.01) USPC - 554/221 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC(8): C07C 57/00 (2012.01) USPC- 554/221 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched USPC- 514/559, 560; 554/224; 435/7.23 (text search) Find search terms below Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) PubWEST (US Pat, PgPub, EPO, JPO), GoogleScholar (PL, NPL), FreePatentsOnline (US Pat, PgPub, EPO, JPO, WIPO, NPL) Search terms: retinoic acid Pin 1 fluorescently labeled probes erb2 erbb2 her2 neu Peptidyl-Prolyl-Isomerase rotamase retinoic erb fluorescent label		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	GIANNI et al. Inhibition of the Peptidyl-Prolyl-Isomerase Pin1 Enhances the Responses of Acute Myeloid Leukemia Cells to Retinoic Acid via Stabilization of RARA and PML-RARA. Cancer Res. 1 February 2009 (01.02.2009) , Vol 69 (3), pp 1016-1026. Abstract, pg 1016, col 2, para 1; pg 1025, col 1, para 3; pg 1023, col 2, para 3 to pg 1024, col 1, para 1; pg 1025, col 2, para 3	1-3 ---- 4
Y	US 2004/0176912 A1 (SOWADSKI et al.) 09 September 2004 (09.09.2004) para [0107], [0145]-[0149], [0153]	4, 20-21
Y	WO 2005/027727 A2 (LU et al.) 31 March 2005 (31.03.2005) pg 31, ln 24 - pg 32, ln 10; pg 34, ln 1 - pg 35, ln 7.	20
Y	LAM et al. 'Prolyl isomerase Pin1 is highly expressed in Her2-positive breast cancer and regulates erbB2 protein stability', Molecular Cancer 2008, Vol 7:9, pages 1-12. abstract; pg 3, col 2, para 2 - pg 4, col 2, para 1	21
A	US 2006/0018899 A1 (KAO et al.) 26 January 2006 (26.01.2006) para [0232]- 0234]	1-4, 20-21
☐ Further documents are listed in the continuation of Box C. ☐		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 07 June 2012 (07.06.2012)		Date of mailing of the international search report 18 JUL 2012
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201		Authorized officer: Lee W. Young PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/29077

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5-19
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.