

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 February 2012 (23.02.2012)

PCT

(10) International Publication Number
WO 2012/022814 A1

(51) International Patent Classification:
C07K 16/32 (2006.01) *A61P 35/00* (2006.01)
A61K 39/395 (2006.01)

(21) International Application Number:
PCT/EP2011/064407

(22) International Filing Date:
22 August 2011 (22.08.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/375,408 20 August 2010 (20.08.2010) US

(71) Applicant (for all designated States except US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ELIS, Winfried** [DE/DE]; MORPHOSYS AG, Lena-Christ-Strasse 48, 82152 Planegg (DE). **ETTENBERG, Seth** [US/US]; Novartis Institutes for BioMedical Research, Inc., 250 Massachusetts Avenue, Cambridge, Massachusetts 02139 (US). **GARNER, Andrew Paul** [GB/US]; Novartis Institutes for BioMedical Research, Inc., 250 Massachusetts Avenue, Cambridge, Massachusetts 02139 (US). **HAUBST, Nicole** [DE/DE]; MORPHOSYS AG, Lena-Christ-Strasse 48, 82152 Planegg (DE). **KUNZ, Christian Carsten Silvester** [DE/DE]; MORPHOSYS AG, Lena-Christ-Strasse 48, 82152 Planegg (DE). **REISINGER SPRAGUE, Elizabeth Anne** [US/US]; Novartis Institutes for BioMedical Research, Inc., 250 Massachusetts Avenue, Cambridge, Massachusetts 02139 (US).

(74) Agent: **VON SPRECHER, Georg**; NOVARTIS PHARMA AG, Patent Department, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTIBODIES FOR EPIDERMAL GROWTH FACTOR RECEPTOR 3 (HER3)

(57) Abstract: The present invention relates to antibodies or fragments thereof that target a conformational epitope of a HER receptor. In particular, the invention relates to antibodies or fragments thereof that target a conformational epitope of HER3 receptor and compositions and methods of use thereof.



WO 2012/022814 A1

ANTIBODIES FOR EPIDERMAL GROWTH FACTOR RECEPTOR 3 (HER3)

Cross-Reference to Related Applications

This application claims priority to US provisional application number 61/375,408, filed August 20, 2010, the contents of which are included in their entirety.

5 Field of the Invention

This invention relates generally to antibodies or fragments thereof which interact with HER family of receptors, e.g., HER3 receptor. In particular, it relates to antibodies or fragments thereof that recognize a conformational epitope of HER receptor (e.g., HER3) comprising residues from both domains 2 and 4 resulting in inhibition of both ligand-dependent and
10 ligand-independent signal transduction. The invention also relates to antibodies and fragments thereof that bind to HER receptors (e.g., HER3 receptor) concurrently with a ligand (e.g., neuregulin), whilst preventing ligand-induced activation of signal transduction.

Background of the Invention

15 The human epidermal growth factor receptor 3 (ErbB3, also known as HER3) is a receptor protein tyrosine kinase and belongs to the epidermal growth factor receptor (EGFR) subfamily of receptor protein tyrosine kinases, which also includes EGFR (HER1, ErbB1), HER2 (ErbB2, Neu), and HER4 (ErbB4) (Plowman *et al.*, (1990) Proc. Natl. Acad. Sci. U.S.A. 87:4905-4909; Kraus *et al.*, (1989) Proc. Natl. Acad. Sci. U.S.A. 86:9193-9197; and Kraus
20 *et al.*, (1993) Proc. Natl. Acad. Sci. U.S.A. 90:2900-2904). Like the prototypical epidermal growth factor receptor, the transmembrane receptor HER3 consists of an extracellular ligand-binding domain (ECD), a dimerization domain within the ECD, a transmembrane domain, an intracellular protein tyrosine kinase-like domain (TKD) and a C-terminal phosphorylation domain. Unlike the other HER family members, the kinase domain of HER3 displays very
25 low intrinsic kinase activity.

The ligands neuregulin 1 (NRG) or neuregulin 2 bind to the extracellular domain of HER3 and activate receptor-mediated signaling pathway by promoting dimerization with other dimerization partners such as HER2. Heterodimerization results in activation and
30 transphosphorylation of HER3's intracellular domain and is a means not only for signal diversification but also signal amplification. In addition, HER3 heterodimerization can also occur in the absence of activating ligands and this is commonly termed ligand-independent

HER3 activation. For example, when HER2 is expressed at high levels as a result of gene amplification (e.g. in breast, lung, ovarian or gastric cancer) spontaneous HER2/HER3 dimers can be formed. In this situation the HER2/HER3 is considered the most active ErbB signaling dimer and is therefore highly transforming.

5

Increased HER3 has been found in several types of cancer such as breast, lung, gastrointestinal and pancreatic cancers. Interestingly, a correlation between the expression of HER2/HER3 and the progression from a non-invasive to an invasive stage has been shown (Alimandi *et al.*, (1995) *Oncogene* 10:1813-1821; DeFazio *et al.*, (2000) *Cancer* 87:487-498; 10 Naidu *et al.*, (1988) *Br. J. Cancer* 78:1385-1390). Accordingly, agents that interfere with HER3 mediated signaling are needed.

Summary of the Invention

The invention is based on the discovery of antigen binding proteins (e.g., antibodies or fragments thereof) that bind to a conformational epitope of HER3 receptor comprising amino 15 acid residues within domain 2 and domain 4 of HER3. This binding of the antibodies or fragments thereof with domain 2 and domain 4 stabilizes the HER3 receptor in an inactive or closed conformation such that HER3 activation is inhibited. Surprisingly, binding of the antibodies or fragments thereof with this conformational epitope blocks both ligand- 20 dependent (e.g. neuregulin) and ligand-independent HER3 signaling pathways. Furthermore, antibody mediated inhibition of ligand induced signaling occurs without blocking ligand binding (i.e. both ligand and antibody can bind HER3) presumably because HER3 cannot undergo the conformational rearrangements required for activation.

Accordingly, in one aspect, the invention pertains to an isolated antibody or fragment thereof that binds to an inactive state of a HER receptor, wherein the antibody or fragment thereof 25 blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof stabilizes the HER receptor in an inactive state.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, and 30 wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof binds to the inactive state of the HER receptor. In one embodiment, the antibody or fragment

thereof binds to the active state of the HER receptor and drives it into the inactive state. In another embodiment, the antibody or fragment thereof stabilizes the HER receptor in the inactive state. The HER receptor is selected from the group consisting of HER1, HER2, HER3 and HER4. The antibody is selected from the group consisting of a monoclonal
5 antibody, a polyclonal antibody, a chimeric antibody, a humanized antibody, and a synthetic antibody.

An another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, wherein
10 binding of the antibody stabilizes the HER receptor in an inactive state, and wherein a HER ligand can concurrently bind to a ligand binding site on the HER receptor. In one embodiment, HER ligand binding to the ligand binding site fails to induce a conformational change in the HER receptor to an active state. In another embodiment, HER ligand binding to the ligand binding site fails to activate signal transduction.

15 In one embodiment, the HER ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, neuregulin 3, neuregulin 4, betacellulin, heparin-binding epidermal growth factor, epiregulin, epidermal growth factor, amphiregulin, and transforming growth factor alpha.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that
20 recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, wherein binding of the antibody stabilizes the HER receptor in an inactive state such that the HER receptor fails to dimerize with a co-receptor to form a receptor-receptor complex. The failure to form a receptor-receptor complex prevents activation of both ligand-dependent and ligand-
25 independent signal transduction.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, wherein
30 binding of the antibody to the HER receptor allows dimerization with a co-receptor to form an inactive receptor-receptor complex. The formation of the inactive receptor-receptor complex prevents activation of ligand-independent signal transduction.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that binds to an inactive conformation of a HER3 receptor, wherein the antibody blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state.

5 In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof binds
10 to the inactive state of the HER3 receptor. In another embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in the inactive state. The antibody is selected from the group consisting of a monoclonal antibody, a polyclonal antibody, a chimeric antibody, a humanized antibody, and a synthetic antibody.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that
15 recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein binding of the antibody stabilizes the HER3 receptor in an inactive state, and wherein a HER3 ligand can concurrently bind to a ligand binding site on the HER3 receptor. In one
20 embodiment, HER3 ligand binding to the ligand binding site fails to induce a conformational change in the HER3 receptor to an active state. In another embodiment, HER3 ligand binding to the ligand binding site fails to activate signal transduction. In one embodiment, the HER3 ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, betacellulin, heparin-binding epidermal growth factor, and epiregulin.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that
25 recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof binds
30 to the inactive state of the HER3 receptor. In another embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that binds a conformational epitope of HER3 receptor, wherein the conformational epitope comprises

amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein domain 2 comprises a dimerization loop, and wherein the antibody or fragment blocks both ligand-dependent and ligand-independent signal transduction. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state. In one embodiment, the conformational epitope comprises amino acid residues 265-277, 315 (of domain 2), 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) or a subset thereof. In one embodiment, the VH of the antibody or fragment thereof binds to at least one of the following HER3 residues: Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597. In one embodiment, the VL of the antibody or fragment thereof binds to at least one of the following HER3 residues: Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of first HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the first HER receptor, wherein binding of the antibody or fragment thereof to the first HER receptor in the absence of a HER receptor ligand reduces ligand-independent formation of a first HER receptor-second HER receptor protein complex in a cell which expresses first HER receptor and second HER receptor. In one embodiment, the antibody or fragment thereof stabilizes the first HER receptor in an inactive state such that the first HER receptor fails to dimerize with the second HER receptor to form a first HER receptor-second HER receptor protein complex. In one embodiment, the failure to form a first HER receptor-second HER receptor protein complex prevents activation of signal transduction. In one embodiment, the first HER is selected from the group consisting of HER1, HER2, HER3, and HER4. In one embodiment, the second HER is selected from the group consisting of HER1, HER2, HER3, and HER4.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of HER3, wherein binding of the antibody or fragment thereof to the HER3 receptor in the absence of a HER3 ligand reduces ligand-independent formation of a HER2-HER3 protein complex in a cell which expresses HER2 and HER3. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state such that the HER3 receptor fails to dimerize with the HER2 receptor to form a HER2-HER3 protein complex. In one embodiment, the failure to

form a HER2-HER3 protein complex prevents activation of signal transduction. In one embodiment, the antibody or fragment thereof stabilizes HER3 receptor in inactive state such that the HER3 receptor can still dimerize with HER2 but forms an inactive HER2-HER3 protein complex. In one embodiment, the formation of an inactive HER2-HER3 protein complex prevents activation of signal transduction.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of first HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the first HER receptor, wherein binding of the antibody or fragment thereof to the first HER receptor in the presence of a HER ligand reduces ligand-dependent formation of a first HER receptor-second HER receptor protein complex in a cell which expresses first HER receptor and second HER receptor. In one embodiment, the antibody or fragment thereof stabilizes the first HER receptor in an inactive state such that the HER receptor fails to dimerize with the second HER receptor in the presence of a first HER ligand to form a first HER receptor- second HER receptor protein complex. In one embodiment, the failure to form a first HER receptor-second HER receptor protein complex prevents activation of signal transduction. In one embodiment, the HER ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, neuregulin 3, neuregulin 4, betacellulin, heparin-binding epidermal growth factor, epiregulin, epidermal growth factor, amphiregulin, and transforming growth factor alpha. In one embodiment, the first HER is selected from the group consisting of HER1, HER2, HER3, and HER4. In one embodiment, the second HER is selected from the group consisting of HER1, HER2, HER3, and HER4.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of HER3, wherein binding of the antibody or fragment thereof to the HER3 receptor in the presence of a HER3 ligand reduces ligand-dependent formation of a HER2-HER3 protein complex in a cell which expresses HER2 and HER3. The ligand is selected from the group consisting of neuregulin 1 (NRG), and neuregulin 2. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state such that the HER3 receptor fails to dimerize with the HER2 receptor in the presence of a HER3 ligand to form a HER2-HER3 protein complex. In one embodiment, the failure to form a HER2-HER3 protein complex prevents activation of signal transduction.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of HER3, and wherein the antibody or fragment thereof inhibits phosphorylation of HER3 as assessed by HER3 ligand-independent phosphorylation assay. In one embodiment, the HER3 ligand-independent phosphorylation assay uses HER2 amplified cells, wherein the HER2 amplified cells are SK-Br-3 cells.

In another aspect, the invention pertains to an isolated antibody or fragment thereof that recognizes a conformational epitope of HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of HER3, and wherein the antibody or fragment thereof inhibits phosphorylation of HER3 as assessed by HER3 ligand-dependent phosphorylation assay. In one embodiment, the HER3 ligand-dependent phosphorylation assay uses MCF7 cells stimulated with neuregulin (NRG).

In another aspect, the invention pertains isolated antibody or fragment thereof to a HER3 receptor, having a dissociation (K_D) of at least $1 \times 10^7 M^{-1}$, $10^8 M^{-1}$, $10^9 M^{-1}$, $10^{10} M^{-1}$, $10^{11} M^{-1}$, $10^{12} M^{-1}$, $10^{13} M^{-1}$. In one embodiment, the antibody or fragment thereof inhibits phosphorylation of HER3 as measured by in vitro binding to human HER3 in a phosphorylation assay selected from the group consisting of phospho-HER3 and phospho-Akt.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor, that cross competes with an antibody described in Table 1; an antibody or fragment thereof that interacts with (e.g., by binding, steric hindrance, stabilizing/destabilizing, spatial distribution) the same epitope as an antibody described in Table 1. In one embodiment, the antibody or fragment thereof is a monoclonal antibody. In another embodiment, the antibody or fragment thereof is a human or humanized antibody. In another embodiment, the antibody or fragment thereof is a chimeric antibody. In one embodiment, the antibody or fragment thereof comprises a human heavy chain constant region and a human light chain constant region. In one embodiment, the antibody or fragment thereof is a single chain antibody. In another embodiment, the antibody or fragment thereof is a Fab fragment. In yet another embodiment, the antibody or fragment thereof is a scFv. In one embodiment, the antibody or fragment thereof binds to both human HER3 and cynomolgus HER3. In one embodiment, the antibody or fragment thereof is an IgG isotype. In another embodiment, the antibody or

fragment thereof comprises a framework in which amino acids have been substituted into the antibody framework from the respective human VH or VL germline sequences.

In one aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 comprising 1, 2, 3, 4, 5, or 6 CDRs calculated by Kabat or Chothia of any of the antibodies in
5 Table 1.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor comprising a heavy chain CDR3 selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 10, SEQ ID NO: 22, SEQ ID NO: 28, SEQ ID NO: 40, SEQ ID NO: 46, SEQ ID NO: 58, SEQ ID NO: 64, SEQ ID NO: 76, SEQ ID NO: 82, SEQ ID NO: 94, SEQ ID NO:
10 100, SEQ ID NO: 112, SEQ ID NO: 118, SEQ ID NO: 130, SEQ ID NO: 136, SEQ ID NO: 148, SEQ ID NO: 166, SEQ ID NO: 184, SEQ ID NO: 202, SEQ ID NO: 220, SEQ ID NO: 238, SEQ ID NO: 256, SEQ ID NO: 274, SEQ ID NO: 292, SEQ ID NO: 310, SEQ ID NO: 328, SEQ ID NO: 346, and SEQ ID NO: 364.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3
15 receptor which antibody comprises a VH comprising SEQ ID NO: 15 and a VL comprising SEQ ID NO: 14, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 33 and a VL comprising SEQ ID NO: 32, or an amino acid sequence with 97-99% identity thereof.

20 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 51 and a VL comprising SEQ ID NO: 50, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3
25 receptor which antibody comprises a VH comprising SEQ ID NO: 69 and a VL comprising SEQ ID NO: 68, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 87 and a VL comprising SEQ ID NO: 86, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3
30 receptor which antibody comprises a VH comprising SEQ ID NO: 105 and a VL comprising SEQ ID NO: 104, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 123 and a VL comprising SEQ ID NO: 122, or an amino acid sequence with 97-99% identity thereof.

5 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 141 and a VL comprising SEQ ID NO: 140, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 159 and a VL comprising SEQ ID NO: 158, or an amino acid sequence with 97-99% identity thereof.

10 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 177 and a VL comprising SEQ ID NO: 176, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 195 and a VL comprising
15 SEQ ID NO: 194, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 213 and a VL comprising SEQ ID NO: 212, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3
20 receptor which antibody comprises a VH comprising SEQ ID NO: 231 and a VL comprising SEQ ID NO: 230, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 249 and a VL comprising SEQ ID NO: 248, or an amino acid sequence with 97-99% identity thereof.

25 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 267 and a VL comprising SEQ ID NO: 266, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3
30 receptor which antibody comprises a VH comprising SEQ ID NO: 285 and a VL comprising SEQ ID NO: 284, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 303 and a VL comprising SEQ ID NO: 302, or an amino acid sequence with 97-99% identity thereof.

5 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 321 and a VL comprising SEQ ID NO: 320, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 339 and a VL comprising SEQ ID NO: 338, or an amino acid sequence with 97-99% identity thereof.

10 In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 357 and a VL comprising SEQ ID NO: 356, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 375 and a VL comprising
15 SEQ ID NO: 374, or an amino acid sequence with 97-99% identity thereof.

In another aspect, the invention pertains to an isolated antibody or fragment thereof comprising a variable heavy chain sequence having SEQ ID NO: 493.

In another aspect, the invention pertains to an isolated antibody or fragment thereof comprising a variable light chain sequence having SEQ ID NO: 494.

20 In another aspect, the invention pertains to an isolated antibody or fragment thereof comprising a variable heavy chain sequence having SEQ ID NO: 493 and a variable light chain sequence having SEQ ID NO: 494.

In another aspect, the invention pertains to an isolated antibody or fragment thereof to HER3 receptor with a variant heavy chain variable region comprising CDR1, CDR2, and CDR3,
25 wherein variant has at least one to four amino acid changes in one of CDR1, CDR2, or CDR3.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 2; CDR2 of SEQ ID NO: 3; CDR3 of SEQ ID NO: 4; a light chain variable region CDR1 of SEQ ID NO: 5; CDR2 of SEQ ID NO: 6; and CDR3 of SEQ ID NO: 7.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 20; CDR2 of SEQ ID NO: 21; CDR3 of SEQ ID NO: 22; a light chain variable region CDR1 of SEQ ID NO: 23; CDR2 of SEQ ID NO: 24; and CDR3 of SEQ ID NO: 25.

- 5 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 38; CDR2 of SEQ ID NO: 39; CDR3 of SEQ ID NO: 40; a light chain variable region CDR1 of SEQ ID NO: 41; CDR2 of SEQ ID NO: 42; and CDR3 of SEQ ID NO: 43.

- 10 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 56; CDR2 of SEQ ID NO: 57; CDR3 of SEQ ID NO: 58; a light chain variable region CDR1 of SEQ ID NO: 59; CDR2 of SEQ ID NO: 60; and CDR3 of SEQ ID NO: 61.

- 15 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 74; CDR2 of SEQ ID NO: 75; CDR3 of SEQ ID NO: 76; a light chain variable region CDR1 of SEQ ID NO: 77; CDR2 of SEQ ID NO: 78; and CDR3 of SEQ ID NO: 79.

- 20 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 92; CDR2 of SEQ ID NO: 93; CDR3 of SEQ ID NO: 94; a light chain variable region CDR1 of SEQ ID NO: 95; CDR2 of SEQ ID NO: 96; and CDR3 of SEQ ID NO: 97.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 110; CDR2 of SEQ ID NO: 111; CDR3 of SEQ ID NO: 112; a light chain variable region CDR1 of SEQ ID NO: 113; CDR2 of SEQ ID NO: 114; and CDR3 of SEQ ID NO: 115.

- 25 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 128; CDR2 of SEQ ID NO: 129; CDR3 of SEQ ID NO: 130; a light chain variable region CDR1 of SEQ ID NO: 131; CDR2 of SEQ ID NO: 132; and CDR3 of SEQ ID NO: 133.

- 30 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 146; CDR2 of SEQ ID NO:

147; CDR3 of SEQ ID NO: 148; a light chain variable region CDR1 of SEQ ID NO: 149; CDR2 of SEQ ID NO: 150; and CDR3 of SEQ ID NO: 151.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 164; CDR2 of SEQ ID NO: 165; CDR3 of SEQ ID NO: 166; a light chain variable region CDR1 of SEQ ID NO: 167; CDR2 of SEQ ID NO: 168; and CDR3 of SEQ ID NO: 169.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 182; CDR2 of SEQ ID NO: 183; CDR3 of SEQ ID NO: 184; a light chain variable region CDR1 of SEQ ID NO: 185; CDR2 of SEQ ID NO: 186; and CDR3 of SEQ ID NO: 187.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 200; CDR2 of SEQ ID NO: 201; CDR3 of SEQ ID NO: 202; a light chain variable region CDR1 of SEQ ID NO: 203; CDR2 of SEQ ID NO: 204; and CDR3 of SEQ ID NO: 205.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 218; CDR2 of SEQ ID NO: 219; CDR3 of SEQ ID NO: 220; a light chain variable region CDR1 of SEQ ID NO: 221; CDR2 of SEQ ID NO: 222; and CDR3 of SEQ ID NO: 223.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 236; CDR2 of SEQ ID NO: 237; CDR3 of SEQ ID NO: 238; a light chain variable region CDR1 of SEQ ID NO: 239; CDR2 of SEQ ID NO: 240; and CDR3 of SEQ ID NO: 241.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 254; CDR2 of SEQ ID NO: 255; CDR3 of SEQ ID NO: 256; a light chain variable region CDR1 of SEQ ID NO: 257; CDR2 of SEQ ID NO: 258; and CDR3 of SEQ ID NO: 259.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 272; CDR2 of SEQ ID NO: 273; CDR3 of SEQ ID NO: 274; a light chain variable region CDR1 of SEQ ID NO: 275; CDR2 of SEQ ID NO: 276; and CDR3 of SEQ ID NO: 277.

In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 290; CDR2 of SEQ ID NO: 291; CDR3 of SEQ ID NO: 292; a light chain variable region CDR1 of SEQ ID NO: 293; CDR2 of SEQ ID NO: 294; and CDR3 of SEQ ID NO: 295.

- 5 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 308; CDR2 of SEQ ID NO: 309; CDR3 of SEQ ID NO: 310; a light chain variable region CDR1 of SEQ ID NO: 311; CDR2 of SEQ ID NO: 312; and CDR3 of SEQ ID NO: 313.

- 10 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 326; CDR2 of SEQ ID NO: 327; CDR3 of SEQ ID NO: 328; a light chain variable region CDR1 of SEQ ID NO: 329; CDR2 of SEQ ID NO: 330; and CDR3 of SEQ ID NO: 331.

- 15 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 344; CDR2 of SEQ ID NO: 345; CDR3 of SEQ ID NO: 346; a light chain variable region CDR1 of SEQ ID NO: 347; CDR2 of SEQ ID NO: 348; and CDR3 of SEQ ID NO: 349.

- 20 In another aspect, the invention pertains to an isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 362; CDR2 of SEQ ID NO: 363; CDR3 of SEQ ID NO: 364; a light chain variable region CDR1 of SEQ ID NO: 365; CDR2 of SEQ ID NO: 366; and CDR3 of SEQ ID NO: 367.

In one embodiment, the fragment of an antibody binds to HER3 is selected from the group consisting of Fab, F(ab₂)', F(ab)₂', scFv, VHH, VH, VL, dAbs.

- 25 In another aspect, the invention pertains to a pharmaceutical composition comprising an antibody or fragment and a pharmaceutically acceptable carrier. In one embodiment, the pharmaceutical composition further comprising an additional therapeutic agent, such as an antibody, a small molecule, an mTOR inhibitor or a PI3Kinase inhibitor. In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a HER1 inhibitor including, but is not limited to, Matuzumab (EMD72000),
Erbix®/Cetuximab, Vectibix®/Panitumumab, mAb 806, Nimotuzumab, Iressa®/Gefitinib,
30 CI-1033 (PD183805), Lapatinib (GW-572016), Tykerb®/Lapatinib Ditosylate, Tarceva®/
Erlotinib HCL (OSI-774), PKI-166, and Tovok®.

In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a HER2 inhibitor including, but is not limited to, Pertuzumab, Trastuzumab, MM-111, neratinib, lapatinib or lapatinib ditosylate /Tykerb®.

5 In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a HER3 inhibitor including, but is not limited to, MM-121, MM-111, IB4C3, 2DID12 (U3 Pharma AG), AMG888 (Amgen), AV-203(Aveo), MEHD7945A (Genentech); small molecules that inhibit HER3.

In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a HER4 inhibitor.

10 In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a PI3 kinase inhibitor including, but is not limited to, GDC 0941 BEZ235, BMK120 and BYL719.

In one embodiment, the pharmaceutical composition comprises the antibody or fragment thereof of the invention and a mTOR inhibitor including, but is not limited to,
15 Temsirolimus/Torisel®, ridaforolimus / Deforolimus, AP23573, MK8669, everolimus /Affinitor®. In another aspect, the invention pertains to a method of treating a cancer comprising selecting a subject having an HER3 expressing cancer, administering to the subject an effective amount of a composition comprising an antibody or fragment thereof selected from any one of the previous claims. In one embodiment, the subject is a human.

20 In another aspect, the invention pertains to a method of treating a cancer comprising selecting a subject having an HER3 expressing cancer, administering to the subject an effective amount of a composition comprising an antibody or fragment thereof selected from any one of the previous claims, wherein the cancer is selected from the group consisting of breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer,
25 pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors, schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, melanoma. In one embodiment, the cancer is breast cancer.

30 In another aspect, the invention pertains to a method of treating a cancer comprising selecting a subject having an HER3 expressing cancer, administering to said subject an effective

amount of a composition comprising an a combination of antibodies or fragments thereof disclosed in Table 1 that binds to HER3.

In another aspect, the invention pertains to a method of treating a cancer comprising selecting a subject having an HER3 expressing cancer, administering to said subject an effective
5 amount of a composition comprising an antibody or fragment thereof that binds to HER3 and inhibits HER3 ligand-dependent signal transduction and ligand-independent signal transduction.

In another aspect, the invention pertains to use of an antibody or fragment thereof of any one of the previous claims in the manufacture of a medicament for the treatment of a cancer
10 mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway selected from the group consisting of breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors, schwannoma, head and neck cancer, bladder
15 cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and melanoma.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 15 and VL of SEQ ID NO: 14 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

20 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 33 and VL of SEQ ID NO: 32 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 51 and VL of SEQ ID NO: 50 for use in treating a cancer mediated by a HER3 ligand-dependent signal
25 transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 69 and VL of SEQ ID NO: 68 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 87 and VL
30 of SEQ ID NO: 86 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 105 and VL of SEQ ID NO: 104 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

5 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 123 and VL of SEQ ID NO: 122 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 141 and VL of SEQ ID NO: 140 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

10 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 151 and VL of SEQ ID NO: 158 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 177 and VL of SEQ ID NO: 176 for use in treating a cancer mediated by a HER3 ligand-dependent
15 signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 195 and VL of SEQ ID NO: 194 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 213 and
20 VL of SEQ ID NO: 212 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 231 and VL of SEQ ID NO: 230 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

25 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 249 and VL of SEQ ID NO: 248 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 267 and
30 VL of SEQ ID NO: 266 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 285 and VL of SEQ ID NO: 284 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

5 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 303 and VL of SEQ ID NO: 302 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 321 and VL of SEQ ID NO: 320 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

10 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 339 and VL of SEQ ID NO: 338 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 357 and VL of SEQ ID NO: 356 for use in treating a cancer mediated by a HER3 ligand-dependent
15 signal transduction or ligand-independent signal transduction pathway.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 375 and VL of SEQ ID NO: 374 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway. In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 15 and VL of SEQ ID NO: 14
20 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 33 and VL of SEQ ID NO: 32 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 51 and VL of SEQ ID NO: 50 for use as a medicament.

25 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 69 and VL of SEQ ID NO: 68 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 87 and VL of SEQ ID NO: 86 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 105 and
30 VL of SEQ ID NO: 104 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 123 and VL of SEQ ID NO: 122 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 141 and VL of SEQ ID NO: 140 for use as a medicament.

5 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 159 and VL of SEQ ID NO: 158 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 177 and VL of SEQ ID NO: 176 for use as a medicament.

10 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 195 and VL of SEQ ID NO: 194 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 213 and VL of SEQ ID NO: 212 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 231 and VL of SEQ ID NO: 230 for use as a medicament.

15 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 249 and VL of SEQ ID NO: 248 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 267 and VL of SEQ ID NO: 266 for use as a medicament.

20 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 285 and VL of SEQ ID NO: 284 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 303 and VL of SEQ ID NO: 302 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 321 and VL of SEQ ID NO: 320 for use as a medicament.

25 In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 339 and VL of SEQ ID NO: 338 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 357 and VL of SEQ ID NO: 356 for use as a medicament.

In another aspect, the invention pertains to an antibody having VH of SEQ ID NO: 375 and VL of SEQ ID NO: 374 for use as a medicament.

Brief Description of Figures

Figure 1: Representative MOR10701 SET curves obtained with human, mouse, rat and cyno
5 HER3

Figure 2: SK-Br-3 cell binding determination by FACS titration

Figure 3: HER3 domain binding ELISA

Figure 4: Hydrogen deuterium exchange epitope mapping. A) HER3 ECD peptides recovered
10 following HDX-MS analysis are indicated by dashed lines. Potential N-linked glycosylation sites are highlighted. B) The relative degree of deuteration observed in peptides identified by MS. C) Protected residues mapped onto the published HER3 crystal structure.

Figure 5: A) Surface representation of the HER3/MOR09823 and HER3/ MOR09825 x-ray
15 crystal structures. HER3 (in lighter gray) is in the closed conformation, and MOR09823 or MOR09825 (in darkest gray) bind to both domains 2 and 4. B). Surface view of HER3 from the HER3/MOR09823 structure shown in a similar orientation as (A). MOR09823 was omitted for clarity. C) HER3/MOR09823 structure illustrated as a ribbon structure, viewed at a 90° rotation from panels (A), (B) and (D). D) A ribbon representation of the inactive HER3 conformation recognized by MOR09823 Fab with a close up view of the domain 2/domain 4 interface, highlighting the HER3 residues that are within 5Å of the Fab. E) Mutant HER3/
20 MOR10703 binding determination by ELISA titration.

Figure 6: Inhibition of ligand induced (A) or ligand-independent (B) HER3 phosphorylation.

Figure 7: Inhibition of HER3 dependent downstream signaling pathways in *HER2* amplified cell lines.

Figure 8: The impact of HER3 inhibition upon cell growth in A) BT-474 and B) neuregulin
25 stimulated MCF7 cells.

Figure 9: The effect of MOR09823 and MOR09825 upon neuregulin binding to MCF7 cells.

Figure 10: Impact of MOR09823 binding upon HER3/ neuregulin complex formation as assessed by Biacore™. No antibody (black bars), MOR09823 (white bars), 105.5 (grey) & control IgG (striped bars).

Figure 11: MOR09823 mediated inhibition of (A) ligand independent (BT-474) and (B) ligand dependent (BxPC3) HER3 signaling *in vivo*.

Figure 12: The impact of MOR10701 and MOR10703 upon BT-474 tumor growth.

Figure 13: The impact of MOR10701 and MOR10703 upon BxPC3 tumor growth.

- 5 Figure 14: MOR10703 *in vitro* drug combination isobolograms (A) MOR09823/ trastuzumab, (B) MOR09823/ lapatinib, (C) MOR10703/ BEZ235, (D) MOR10703/ BKM120, (E) MOR10703/ BYL719, (F) MOR10703/ RAD001, (G) MOR10703/ cetuximab and (H) MOR10703/ erlotinib.

Figure 15: MOR10701 or MOR10703 *in vivo* combinations with (A) trastuzumab and (B) erlotinib in BT-474 and L3.3.

10

Detailed Description of the Invention

Definitions

In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

- 15 The phrase “signal transduction” or “signaling activity” as used herein refers to a biochemical causal relationship generally initiated by a protein-protein interaction such as binding of a growth factor to a receptor, resulting in transmission of a signal from one portion of a cell to another portion of a cell. For HER3, the transmission involves specific phosphorylation of one or more tyrosine, serine, or threonine residues on one or more proteins in the series of
- 20 reactions causing signal transduction. Penultimate processes typically include nuclear events, resulting in a change in gene expression.

- A “HER receptor” is a receptor protein tyrosine kinase which belongs to the HER receptor family and includes EGFR, HER2, HER3 and HER4 receptors and other members of this family to be identified in the future. The HER receptor will generally comprise an
- 25 extracellular domain, which may bind an HER ligand; a lipophilic transmembrane domain; a conserved intracellular tyrosine kinase domain; and a carboxyl-terminal signaling domain harboring several tyrosine residues which can be phosphorylated. Preferably the HER receptor is native sequence human HER receptor.

The terms “HER1,” “ErbB1,” “epidermal growth factor receptor” and “EGFR” are used interchangeably herein and refer to EGFR as disclosed, for example, in Carpenter *et al.* Ann. Rev. Biochem. 56:881-914 (1987), including naturally occurring mutant forms thereof (e.g. a deletion mutant EGFR as in Humphrey *et al.*, (1990) PNAS (USA) 87:4207-4211). erbB1
 5 refers to the gene encoding the EGFR protein product.

The terms “HER2” and “ErbB2” and are used interchangeably herein and refer to human HER2 protein described, for example, in Semba *et al.*, (1985) PNAS (USA) 82:6497-6501 and Yamamoto *et al.*(1986) Nature 319:230-234 (Genebank accession number X03363). The
 10 term “erbB2” refers to the gene encoding human ErbB2 and “neu” refers to the-gene encoding rat p185^{neu}.

The terms “HER4” and “ErbB4” herein refer to the receptor polypeptide as disclosed, for example, in EP Pat Appln No 599,274; Plowman *et al.*, (1993) Proc. Natl. Acad. Sci. USA, 90:1746-1750; and Plowman *et al.*, (1993) Nature, 366:473-475, including isoforms thereof,
 15 e.g., as disclosed in WO99/19488, published Apr. 22, 1999.

The term “HER3” or “HER3 receptor” also known as “ErbB3” as used herein refers to mammalian HER3 protein and “her3” or "erbB3" refers to mammalian her3 gene. The preferred HER3 protein is human HER3 protein present in the cell membrane of a cell. The human her3 gene is described in U.S. Pat. No. 5,480,968 and Plowman *et al.*, (1990) Proc.
 20 Natl. Acad. Sci. USA, 87:4905-4909.

Human HER3 as defined in Accession No. NP_001973 (human), and represented below as SEQ ID NO: 1. All nomenclature is for full length, immature HER3 (amino acids 1-1342). The immature HER3 is cleaved between positions 19 and 20, resulting in the mature HER3 protein (20-1342 amino acids).

25 mrandalqvl gllfslargs evgnsqavcp gtlnglsvtg daenqyqtly klyercevvm
 gnleivltgh nadlsflqwi revtgyvlva mnefstlplp nlrvvrgtqv ydgkfaifvm
 lnyntnssha lrqlrltqlt eilsggvyie kndklchmdt idwrdivrdr daeivvkdng
 rscppchevc kgrcwpggse dcqtltktic apqcnghcfg pnpnqcchde caggcsgppd
 tdcfacrhfn dsqacvprcp qplvynkltf qlepnphtky qyggvcvasc phnfvvdqts

cvracppdkm evdknglkmc epcggglepka cegtgsgrf qtvdsnidg fvnctkilgn
 ldflitgng dpwhkipald peklnvfrtv reitgylmq swpphmhfs vfnlittigg
 rslrnrgfsl limklnvts lgrslkeis agriyisanr qlcyhhslnw tkvlrgptee
 rldikhnrpr rdcvaegkvc dplcssggcw gpgpgqlsc rnysrggvcv thenflngep
 5 refaheaecf schpecqpmc gtatengsgs dtcaqcahfr dgphcvsscp hgvlgakgpi
 ykypdvq nec rpchenctqg ckqpelqdcl gqtlvligkt hltmaltvia glvvimmlg
 gtflywrgrr iqnkramrry lergesiepl dpsekankvl arifketelr klkvlgsqvf
 gtvhkgvwip egesikipvc ikviedksgr qsfqavtdhm laigsldhah ivrlglcpg
 sslqlvtqyl plgsldhvr qhrgalgpql llwngvqiak gmyyleehgm vhrnlaarnv
 10 llkspsqvqv adfgvadllp pddkqllyse aktpikwmal esihfgkyth qsdvwsygv
 vwelmtfgae pyaglraev pdllekgerl aqqictidv ymvvmkcwmi denirptfke
 laneftrmar dpprylvikr esgpgiapgp ephgltnkkl eevelepeld lddleaeed
 nlattlgsa lslpvgtlr prgsqslsp ssgympmnqg nlgesqesa vsgssercpr
 pvslhpmprg clasesegh vtgseaelqe kvsmcrsr rrsprprgds ayhsqrhsl
 15 tpvtplspg leedvngyv mpdthlkgtp ssregtlssv glssvlgteee ededeeyeym
 nrrrhspph pprpsleel gyeymdvgsd lsaslgstqs cplhpvpimp tagttdedy
 eymnrqrdgg gpggdyamg acpaseqgye emrafqgpg qaphvhyarl ktlrsleatd
 safdnpywh srlfpkanaq rt (SEQ ID NO: 1)

The term "HER ligand" as used herein refers to polypeptides which bind and activate HER
 20 receptors such as HER1, HER2, HER3 and HER4. Examples of HER ligands include, but are
 not limited to neuregulin 1 (NRG), neuregulin 2, neuregulin 3, neuregulin 4, betacellulin,
 heparin-binding epidermal growth factor, epiregulin, epidermal growth factor, amphiregulin,

and transforming growth factor alpha. The term includes biologically active fragments and/or variants of a naturally occurring polypeptide.

The term “HER3 ligand” as used herein refers to polypeptides which bind and activate HER3. Examples of HER3 ligands include, but are not limited to neuregulin 1 (NRG) and neuregulin
5 2, betacellulin, heparin-binding epidermal growth factor, and epiregulin. The term includes biologically active fragments and/or variants of a naturally occurring polypeptide.

The “HER-HER protein complex” is a noncovalently associated oligomer containing a HER
co- receptors in any combination (e.g., HER1-HER2, HER1-HER3, HER1-HER4, HER2-
HER3, HER3- HER4, and the like). This complex can form when a cell expressing both of
10 these receptors is exposed to a HER ligand e.g., NRG, or when a HER receptor is active or overexpressed.

The “HER2-HER3 protein complex” is a noncovalently associated oligomer containing HER2
receptor and the HER3 receptor. This complex can form when a cell expressing both of these
receptors is exposed to a HER3 ligand e.g., NRG or when HER2 is active/overexpressed

15 The phrase “HER3 activity” or “HER3 activation” as used herein refers to an increase in oligomerization (e.g. an increase in HER3 containing complexes), HER3 phosphorylation, conformational rearrangements (for example those induced by ligands), and HER3 mediated downstream signaling.

The term “stabilization” or “stabilized” used in the context of HER3 refers to an antibody or
20 fragment thereof that directly maintains (locks, tethers, holds, preferentially binds, favors) the inactive state or conformation of HER3 without blocking ligand binding to HER3, such that ligand binding is no longer able to activate HER3. Assays described in the Examples can be used to measure ligand binding to a stabilized HER3 receptor, e.g., Biacore assay.

The term “ligand-dependent signaling” as used herein refers to the activation of HER (e.g.,
25 HER3) via ligand. HER3 activation is evidenced by increased oligomerization (e.g. heterodimerization) and/ or HER3 phosphorylation such that downstream signaling pathways (e.g. PI3K) are activated. The antibody or fragment thereof can statistically significantly reduce the amount of phosphorylated HER3 in a stimulated cell exposed to the antigen binding protein (e.g., an antibody) relative to an untreated (control) cell, as measured using
30 the assays described in the Examples. The cell which expresses HER3 can be a naturally

occurring cell line (e.g. MCF7) or can be recombinantly produced by introducing nucleic acids encoding HER3 protein into a host cell. Cell stimulation can occur either via the exogenous addition of an activating HER3 ligand or by the endogenous expression of an activating ligand.

- 5 The antibody or fragment thereof which “reduces neregulin-induced HER3 activation in a cell” is one which statistically significantly reduces HER3 tyrosine phosphorylation relative to an untreated (control) cell, as measured using the assays described in the Examples. This can be determined based on HER3 phosphotyrosine levels following exposure of HER3 to NRG and the antibody of interest. The cell which expresses HER3 protein can be a naturally
10 occurring cell or cell line (e.g. MCF7) or can be recombinantly produced.

The term “ligand-independent signaling” as used herein refers to cellular HER3 activity (e.g phosphorylation) in the absence of a requirement for ligand binding. For example, ligand-independent HER3 activation can be a result of HER2 overexpression or activating mutations in HER3 heterodimer partners such as EGFR and HER2. The antibody or fragment thereof
15 can statistically significantly reduce the amount of phosphorylated HER3 in a cell exposed to the antigen binding protein (e.g., an antibody) relative to an untreated (control) cell. The cell which expresses HER3 can be a naturally occurring cell line (e.g. SK-Br-3) or can be recombinantly produced by introducing nucleic acids encoding HER3 protein into a host cell.

The term “blocks” as used herein refers to stopping or preventing an interaction or a process,
20 e.g., stopping ligand-dependent or ligand-independent signaling.

The term “recognize” as used herein refers to an antibody or fragment thereof that finds and interacts (e.g., binds) with its conformational epitope.

The phrase “concurrently binds” as used herein refers to a HER ligand that can bind to a ligand binding site on the HER receptor along with the HER antibody. This means that both
25 the antibody and antibody can bind to the HER receptor together. For the sake of illustration only, the HER3 ligand NRG, can bind to the HER3 receptor along with the HER3 antibody. Assay to measure concurrent binding of the ligand and antibody are described in the Examples section (e.g., Biacore).

The term “fails” as used herein refers to an antibody or fragment thereof that does not do a
30 particular event. For example, an antibody or fragment thereof that “fails to activate signal

transduction” is one that does not trigger signal transduction; an antibody or fragment thereof that “fails to induce a conformational change” is one that does not cause a structural alteration in the HER receptor; an antibody or fragment thereof that stabilizes the HER receptor in an inactive state such that the HER receptor “fails to dimerize” is one that does not form protein-
5 protein complexes.

The term “antibody” as used herein refers to whole antibodies that interact with (e.g., by binding, steric hindrance, stabilizing/destabilizing, spatial distribution) an HER3 epitope and inhibit signal transduction. A naturally occurring “antibody” is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each
10 heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into
15 regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant
20 regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system. The term “antibody” includes for example, monoclonal antibodies, human antibodies, humanized antibodies, camelised antibodies, chimeric antibodies, single-chain Fvs (scFv), disulfide-linked Fvs (sdFv), Fab
25 fragments, F (ab') fragments, and anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The antibodies can be of any isotype (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass.

Both the light and heavy chains are divided into regions of structural and functional
30 homology. The terms “constant” and “variable” are used functionally. In this regard, it will be appreciated that the variable domains of both the light (VL) and heavy (VH) chain portions determine antigen recognition and specificity. Conversely, the constant domains of the light chain (CL) and the heavy chain (CH1, CH2 or CH3) confer important biological properties

such as secretion, transplacental mobility, Fc receptor binding, complement binding, and the like. By convention the numbering of the constant region domains increases as they become more distal from the antigen binding site or amino-terminus of the antibody. The N-terminus is a variable region and at the C-terminus is a constant region; the CH3 and CL domains
5 actually comprise the carboxy-terminus of the heavy and light chain, respectively.

The phrase “antibody fragment”, as used herein, refers to one or more portions of an antibody that retain the ability to specifically interact with (e.g., by binding, steric hindrance, stabilizing/destabilizing, spatial distribution) an HER3 epitope and inhibit signal transduction. Examples of binding fragments include, but are not limited to, a Fab fragment, a monovalent
10 fragment consisting of the VL, VH, CL and CH1 domains; a F(ab)₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consisting of the VH and CH1 domains; a Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a dAb fragment (Ward *et al.*, (1989) Nature 341:544-546), which consists of a VH domain; and an isolated complementarity determining region
15 (CDR).

Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird *et al.*, (1988)
20 Science 242:423-426; and Huston *et al.*, (1988) Proc. Natl. Acad. Sci. 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term “antibody fragment”. These antibody fragments are obtained using conventional techniques known to those of skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies.

25 Antibody fragments can also be incorporated into single domain antibodies, maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, e.g., Hollinger and Hudson, (2005) Nature Biotechnology 23:1126-1136). Antibody fragments can be grafted into scaffolds based on polypeptides such as Fibronectin type III (Fn3) (see U.S. Pat. No. 6,703,199, which describes fibronectin polypeptide monobodies).

30 Antibody fragments can be incorporated into single chain molecules comprising a pair of tandem Fv segments (VH-CH1-VH-CH1) which, together with complementary light chain

polypeptides, form a pair of antigen binding regions (Zapata *et al.*, (1995) Protein Eng. 8:1057-1062; and U.S. Pat. No. 5,641,870).

The term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin or otherwise interacting with a molecule. Epitopic determinants generally
5 consist of chemically active surface groupings of molecules such as amino acids or carbohydrate or sugar side chains and can have specific three-dimensional structural characteristics, as well as specific charge characteristics. An epitope may be “linear” or “conformational.”

The term “linear epitope” refers to an epitope with all of the points of interaction between the
10 protein and the interacting molecule (such as an antibody) occur linearly along the primary amino acid sequence of the protein (continuous). Once a desired epitope on an antigen is determined, it is possible to generate antibodies to that epitope, e.g., using the techniques described in the present invention. Alternatively, during the discovery process, the generation and characterization of antibodies may elucidate information about desirable epitopes. From
15 this information, it is then possible to competitively screen antibodies for binding to the same epitope. An approach to achieve this is to conduct cross-competition studies to find antibodies that competitively bind with one another, e.g., the antibodies compete for binding to the antigen. A high throughput process for “binning” antibodies based upon their cross-competition is described in International Patent Application No. WO 2003/48731. As will be
20 appreciated by one of skill in the art, practically anything to which an antibody can specifically bind could be an epitope. An epitope can comprises those residues to which the antibody binds.

The term “conformational epitope” refers to an epitope in which discontinuous amino acids that come together in three dimensional conformation. In a conformational epitope, the points
25 of interaction occur across amino acid residues on the protein that are separated from one another. In one embodiment, the epitope is that described in Examples of this specification. In one embodiment, the conformational epitope is defined by (i) HER3 amino acid residues 265-277 and 315 (of domain 2) and (ii) HER3 amino acid residues 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) of SEQ ID NO: 1, or a subset thereof. As will be appreciated
30 by one of skill in the art, the space that is occupied by a residue or side chain that creates the shape of a molecule helps to determine what an epitope is.

Generally, antibodies specific for a particular target antigen will preferentially recognize an epitope on the target antigen in a complex mixture of proteins and/or macromolecules.

Regions of a given polypeptide that include an epitope can be identified using any number of epitope mapping techniques, well known in the art. See, e.g., Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66 (Glenn E. Morris, Ed., 1996) Humana Press, Totowa, New Jersey. For example, linear epitopes may be determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the peptides with antibodies while the peptides are still attached to the supports. Such techniques are known in the art and described in, e.g., U.S. Patent No. 4,708,871 ; Geysen *et al.*, (1984) Proc. Natl. Acad. Sci. USA 8:3998-4002; Geysen *et al.*, (1985) Proc. Natl. Acad. Sci. USA 82:78-182; Geysen *et al.*, (1986) Mol. Immunol. 23:709-715. Similarly, conformational epitopes are readily identified by determining spatial conformation of amino acids such as by, e.g., hydrogen/deuterium exchange, x-ray crystallography and two-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols, supra. Antigenic regions of proteins can also be identified using standard antigenicity and hydrophathy plots, such as those calculated using, e.g., the Omega version 1.0 software program available from the Oxford Molecular Group. This computer program employs the Hopp/Woods method, Hopp *et al.*, (1981) Proc. Natl. Acad. Sci USA 78:3824-3828; for determining antigenicity profiles, and the Kyte-Doolittle technique, Kyte *et al.*, (1982) J.MoI. Biol. 157:105-132; for hydrophathy plots.

The term “paratope” as used herein refers to the general structure of a binding region that determines binding to an epitope. This structure influences whether or not and in what manner the binding region might bind to an epitope. Paratope can refer to an antigenic site of an antibody that is responsible for an antibody or fragment thereof, to bind to an antigenic determinant. Paratope also refers to the idiotope of the antibody, and the complementary determining region (CDR) region that binds to the epitope. In one embodiment, the paratope is the region of the antibody that binds to the conformational epitope comprising (i) HER3 amino acid residues 265-277 and 315 (of domain 2), and (ii) HER3 amino acid residues 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) of SEQ ID NO: 1, or a subset thereof. In one embodiment, the paratope is the region of the antibody that comprises the CDR sequences. In one embodiment, the paratope comprises the sequences listed in Table 1. In one embodiment, the paratope comprises at least one amino acid residue that binds with HER3 residues: Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276,

His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597. In one embodiment, the paratope comprises at least one amino acid residue that binds with HER3 residues: Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615. As will be appreciated by one of skill in the art, the paratope of any
5 antibody, or variant thereof, can be determined in the manner set forth by the present application.

The phrases “monoclonal antibody” or “monoclonal antibody composition” as used herein refers to polypeptides, including antibodies, antibody fragments, bispecific antibodies, etc. that have substantially identical to amino acid sequence or are derived from the same genetic
10 source. This term also includes preparations of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope.

The phrase “human antibody”, as used herein, includes antibodies having variable regions in which both the framework and CDR regions are derived from sequences of human origin.
15 Furthermore, if the antibody contains a constant region, the constant region also is derived from such human sequences, e.g., human germline sequences, or mutated versions of human germline sequences or antibody containing consensus framework sequences derived from human framework sequences analysis, for example, as described in Knappik *et al.*, (2000) J Mol Biol 296:57-86). The structures and locations of immunoglobulin variable domains, e.g.,
20 CDRs, may be defined using well known numbering schemes, e.g., the Kabat numbering scheme, the Chothia numbering scheme, or a combination of Kabat and Chothia (see, e.g., Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services (1991), eds. Kabat *et al.*; Lazikani *et al.*, (1997) J. Mol. Bio. 273:927-948); Kabat *et al.*, (1991) Sequences of Proteins of Immunological Interest, 5th edit., NIH Publication no.
25 91-3242 U.S. Department of Health and Human Services; Chothia *et al.*, (1987) J. Mol. Biol. 196:901-917; Chothia *et al.*, (1989) Nature 342:877-883; and Al-Lazikani *et al.*, (1997) J. Mol. Biol. 273:927-948.

The human antibodies of the invention may include amino acid residues not encoded by human sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro
30 or by somatic mutation in vivo, or a conservative substitution to promote stability or manufacturing).

The phrase “human monoclonal antibody” as used herein refers to antibodies displaying a single binding specificity which have variable regions in which both the framework and CDR regions are derived from human sequences. In one embodiment, the human monoclonal antibodies are produced by a hybridoma which includes a B cell obtained from a transgenic nonhuman animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

The phrase “recombinant human antibody”, as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom, antibodies isolated from a host cell transformed to express the human antibody, e.g., from a transfectoma, antibodies isolated from a recombinant, combinatorial human antibody library, and antibodies prepared, expressed, created or isolated by any other means that involve splicing of all or a portion of a human immunoglobulin gene, sequences to other DNA sequences. Such recombinant human antibodies have variable regions in which the framework and CDR regions are derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies can be subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the V_H and V_L regions of the recombinant antibodies are sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

Specific binding between two entities means a binding with an equilibrium constant (K_A) (k_{on}/k_{off}) of at least $10^2 M^{-1}$, at least $5 \times 10^2 M^{-1}$, at least $10^3 M^{-1}$, at least $5 \times 10^3 M^{-1}$, at least $10^4 M^{-1}$, at least $5 \times 10^4 M^{-1}$, at least $10^5 M^{-1}$, at least $5 \times 10^5 M^{-1}$, at least $10^6 M^{-1}$, at least $5 \times 10^6 M^{-1}$, at least $10^7 M^{-1}$, at least $5 \times 10^7 M^{-1}$, at least $10^8 M^{-1}$, at least $5 \times 10^8 M^{-1}$, at least $10^9 M^{-1}$, at least $5 \times 10^9 M^{-1}$, at least $10^{10} M^{-1}$, at least $5 \times 10^{10} M^{-1}$, at least $10^{11} M^{-1}$, at least $5 \times 10^{11} M^{-1}$, at least $10^{12} M^{-1}$, at least $5 \times 10^{12} M^{-1}$, at least $10^{13} M^{-1}$, at least $5 \times 10^{13} M^{-1}$, at least $10^{14} M^{-1}$, at least $5 \times 10^{14} M^{-1}$, at least $10^{15} M^{-1}$, or at least $5 \times 10^{15} M^{-1}$.

The phrase “specifically (or selectively) binds” to an antibody (e.g., a HER3 binding antibody) refers to a binding reaction that is determinative of the presence of a cognate antigen (e.g., a human HER3) in a heterogeneous population of proteins and other biologics. In addition to the equilibrium constant (K_A) noted above, an HER3 binding antibody of the

invention typically also has a dissociation rate constant (K_D) (k_{off}/k_{on}) of less than $5 \times 10^{-2}M$, less than $10^{-2}M$, less than $5 \times 10^{-3}M$, less than $10^{-3}M$, less than $5 \times 10^{-4}M$, less than $10^{-4}M$, less than $5 \times 10^{-5}M$, less than $10^{-5}M$, less than $5 \times 10^{-6}M$, less than $10^{-6}M$, less than $5 \times 10^{-7}M$, less than $10^{-7}M$, less than $5 \times 10^{-8}M$, less than $10^{-8}M$, less than $5 \times 10^{-9}M$, less than $10^{-9}M$, less than $5 \times 10^{-10}M$, less than $10^{-10}M$, less than $5 \times 10^{-11}M$, less than $10^{-11}M$, less than $5 \times 10^{-12}M$, less than $10^{-12}M$, less than $5 \times 10^{-13}M$, less than $10^{-13}M$, less than $5 \times 10^{-14}M$, less than $10^{-14}M$, less than $5 \times 10^{-15}M$, or less than $10^{-15}M$ or lower, and binds to HER3 with an affinity that is at least two-fold greater than its affinity for binding to a non-specific antigen (e.g., HSA).

In one embodiment, the antibody or fragment thereof has dissociation constant (K_d) of less than 3000 pM, less than 2500 pM, less than 2000 pM, less than 1500 pM, less than 1000 pM, less than 750 pM, less than 500 pM, less than 250 pM, less than 200 pM, less than 150 pM, less than 100 pM, less than 75 pM, less than 10 pM, less than 1 pM as assessed using a method described herein or known to one of skill in the art (e.g., a BIAcore assay, ELISA, FACS, SET) (Biacore International AB, Uppsala, Sweden). The term " K_{assoc} " or " K_a ", as used herein, refers to the association rate of a particular antibody-antigen interaction, whereas the term " K_{dis} " or " K_d ", as used herein, refers to the dissociation rate of a particular antibody-antigen interaction. The term " K_D ", as used herein, refers to the dissociation constant, which is obtained from the ratio of K_d to K_a (i.e. K_d/K_a) and is expressed as a molar concentration (M). K_D values for antibodies can be determined using methods well established in the art. A method for determining the K_D of an antibody is by using surface plasmon resonance, or using a biosensor system such as a Biacore[®] system.

The term "affinity" as used herein refers to the strength of interaction between antibody and antigen at single antigenic sites. Within each antigenic site, the variable region of the antibody "arm" interacts through weak non-covalent forces with antigen at numerous sites; the more interactions, the stronger the affinity.

The term "avidity" as used herein refers to an informative measure of the overall stability or strength of the antibody-antigen complex. It is controlled by three major factors: antibody epitope affinity; the valence of both the antigen and antibody; and the structural arrangement of the interacting parts. Ultimately these factors define the specificity of the antibody, that is, the likelihood that the particular antibody is binding to a precise antigen epitope.

The term "valency" as used herein refers to the number of potential target binding sites in a polypeptide. Each target binding site specifically binds one target molecule or specific site

(i.e., epitope) on a target molecule. When a polypeptide comprises more than one target binding site, each target binding site may specifically bind the same or different molecules (e.g., may bind to different molecules, e.g., different antigens, or different epitopes on the same molecule).

5 The phrase “antagonist antibody” as used herein refers to an antibody that binds with HER3 and neutralizes the biological activity of HER3 signaling, e.g., reduces, decreases and/or inhibits HER3 induced signaling activity, e.g., in a phospho-HER3 or phospho-Akt assay. Examples of assays are described in more details in the examples below. Accordingly, an antibody that “inhibits” one or more of these HER3 functional properties (e.g., biochemical,
10 immunochemical, cellular, physiological or other biological activities, or the like) as determined according to methodologies known to the art and described herein, will be understood to relate to a statistically significant decrease in the particular activity relative to that seen in the absence of the antibody (e.g., or when a control antibody of irrelevant specificity is present). An antibody that inhibits HER3 activity effects such a statistically
15 significant decrease by at least 10% of the measured parameter, by at least 50%, 80% or 90%, and in certain embodiments an antibody of the invention may inhibit greater than 95%, 98% or 99% of HER3 functional activity as evidenced by a reduction in the level of cellular HER3 phosphorylation.

The phrase “isolated antibody” refers to an antibody that is substantially free of other
20 antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds HER3 is substantially free of antibodies that specifically bind antigens other than HER3). An isolated antibody that specifically binds HER3 may, however, have cross-reactivity to other antigens. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

25 The phrase “conservatively modified variant” applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large
30 number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the

corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are “silent variations,” which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid that encodes a polypeptide is implicit in each described sequence.

For polypeptide sequences, “conservatively modified variants” include individual substitutions, deletions or additions to a polypeptide sequence which result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention. The following eight groups contain amino acids that are conservative substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (see, *e.g.*, Creighton, *Proteins* (1984)). In some embodiments, the term “conservative sequence modifications” are used to refer to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody containing the amino acid sequence.

The terms “cross-compete” and “cross-competing” are used interchangeably herein to mean the ability of an antibody or other binding agent to interfere with the binding of other antibodies or binding agents to HER3 in a standard competitive binding assay.

The ability or extent to which an antibody or other binding agent is able to interfere with the binding of another antibody or binding molecule to HER3, and therefore whether it can be said to cross-compete according to the invention, can be determined using standard competition binding assays. One suitable assay involves the use of the Biacore technology (*e.g.* by using the BIAcore 3000 instrument (Biacore, Uppsala, Sweden)), which can measure the extent of interactions using surface plasmon resonance technology. Another assay for measuring cross-competing uses an ELISA-based approach.

The term “optimized” as used herein refers to a nucleotide sequence has been altered to encode an amino acid sequence using codons that are preferred in the production cell or

organism, generally a eukaryotic cell, for example, a cell of *Pichia*, a cell of *Trichoderma*, a Chinese Hamster Ovary cell (CHO) or a human cell. The optimized nucleotide sequence is engineered to retain completely or as much as possible the amino acid sequence originally encoded by the starting nucleotide sequence, which is also known as the “parental” sequence.

5 Standard assays to evaluate the binding ability of the antibodies toward HER3 of various species are known in the art, including for example, ELISAs, western blots and RIAs. Suitable assays are described in detail in the Examples. The binding kinetics (e.g., binding affinity) of the antibodies also can be assessed by standard assays known in the art, such as by Biacore analysis, or FACS relative affinity (Scatchard). Assays to evaluate the effects of the
10 antibodies on functional properties of HER3 (e.g., receptor binding assays, modulating the Her pathway) are described in further detail in the Examples.

The phrases “percent identical” or “percent identity,” in the context of two or more nucleic acids or polypeptide sequences, refers to two or more sequences or subsequences that are the same. Two sequences are "substantially identical" if two sequences have a specified
15 percentage of amino acid residues or nucleotides that are the same (*i.e.*, 60% identity, optionally 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identity over a specified region, or, when not specified, over the entire sequence), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection.
20 Optionally, the identity exists over a region that is at least about 50 nucleotides (or 10 amino acids) in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides (or 20, 50, 200 or more amino acids) in length.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference
25 sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

30 A “comparison window”, as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be

compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith and Waterman, (1970) *Adv. Appl. Math.* 2:482c, by
5 the homology alignment algorithm of Needleman and Wunsch, (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman, (1988) *Proc. Nat'l. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see,
10 *e.g.*, Brent *et al.*, (2003) *Current Protocols in Molecular Biology*).

Two examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, (1977) *Nuc. Acids Res.* 25:3389-3402; and Altschul *et al.*, (1990) *J. Mol. Biol.* 215:403-410, respectively. Software for performing BLAST analyses is publicly available
15 through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word
20 hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to
25 calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN
30 program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) or 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, (1989) *Proc. Natl. Acad. Sci. USA*

89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin and Altschul, (1993) Proc. Natl. Acad. Sci. USA 90:5873-5787).

5 One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less
10 than about 0.01, and most preferably less than about 0.001.

The percent identity between two amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller, (1988) Comput. Appl. Biosci. 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two
15 amino acid sequences can be determined using the Needleman and Wunsch (1970) J. Mol. Biol. 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package (available at www.gcg.com), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

20 Other than percentage of sequence identity noted above, another indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides
25 differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequence.

30 The phrase "nucleic acid" is used herein interchangeably with the term "polynucleotide" and refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide

analogues or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogues include, without limitation, phosphorothioates, phosphoramidates, methyl
5 phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, as
10 detailed below, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, (1991) *Nucleic Acid Res.* 19:5081; Ohtsuka *et al.*, (1985) *J. Biol. Chem.* 260:2605-2608; and Rossolini *et al.*, (1994) *Mol. Cell. Probes* 8:91-98).

The phrase “operably linked” refers to a functional relationship between two or more
15 polynucleotide (*e.g.*, DNA) segments. Typically, it refers to the functional relationship of a transcriptional regulatory sequence to a transcribed sequence. For example, a promoter or enhancer sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a
20 transcribed sequence are physically contiguous to the transcribed sequence, *i.e.*, they are *cis*-acting. However, some transcriptional regulatory sequences, such as enhancers, need not be physically contiguous or located in close proximity to the coding sequences whose transcription they enhance.

The terms “polypeptide” and “protein” are used interchangeably herein to refer to a polymer
25 of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. Unless otherwise indicated, a particular polypeptide sequence also implicitly encompasses conservatively modified variants thereof.

30 The term “subject” includes human and non-human animals. Non-human animals include all vertebrates, *e.g.*, mammals and non-mammals, such as non-human primates, sheep, dog, cow,

chickens, amphibians, and reptiles. Except when noted, the terms “patient” or “subject” are used herein interchangeably.

The term “anti-cancer agent” means any agent that can be used to treat a cell proliferative disorder such as cancer, including cytotoxic agents, chemotherapeutic agents, radiotherapy
5 and radiotherapeutic agents, targeted anti-cancer agents, and immunotherapeutic agents.

“Tumor” refers to neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

The term “anti-tumor activity” means a reduction in the rate of tumor cell proliferation, viability, or metastatic activity. A possible way of showing anti-tumor activity is show a
10 decline in growth rate of abnormal cells that arises during therapy or tumor size stability or reduction. Such activity can be assessed using accepted in vitro or in vivo tumor models, including but not limited to xenograft models, allograft models, MMTV models, and other known models known in the art to investigate anti-tumor activity.

The term “malignancy” refers to a non-benign tumor or a cancer. As used herein, the term
15 “cancer” includes a malignancy characterized by deregulated or uncontrolled cell growth. Exemplary cancers include: carcinomas, sarcomas, leukemias, and lymphomas. The term “cancer” includes primary malignant tumors (e.g., those whose cells have not migrated to sites in the subject's body other than the site of the original tumor) and secondary malignant tumors (e.g., those arising from metastasis, the migration of tumor cells to secondary sites that are
20 different from the site of the original tumor).

Various aspects of the invention are described in further detail in the following sections and subsections.

Structure and Mechanism of Activation of the HER Receptors

25 All four HER receptors have an extracellular ligand-binding domain, a single trans-membrane domain and a cytoplasmic tyrosine kinase-containing domain. The intracellular tyrosine kinase domain of HER receptors is highly conserved, although the kinase domain of HER3 contains substitutions of critical amino acids and therefore lacks kinase activity (Guy *et al.*, (1994): PNAS 91, 8132-8136). Ligand-induced dimerisation of the HER receptors induces
30 activation of the kinase, receptor transphosphorylation on tyrosine residues in the C-terminal tail, followed by recruitment and activation of intracellular signalling effectors (Yarden and

Sliwkowski, (2001) *Nature Rev* 2, 127-137; Jorissen *et al.*, (2003) *Exp Cell Res* 284, 31-53.

The crystal structures of the extracellular domains of HERs have provided some insight into the process of ligand-induced receptor activation (Schlessinger, (2002) *Cell* 110, 669-672).

5 The extracellular domain of each HER receptor consists of four subdomains: Subdomain I and III cooperate in forming the ligand-binding site, whereas subdomain II (and perhaps also subdomain IV) participates in receptor dimerisation via direct receptor-receptor interactions. In the structures of ligand-bound HER1, a β hairpin (termed the dimerisation loop) in subdomain II interacts with the dimerisation loop of the partner receptor, mediating receptor
10 dimerisation (Garrett *et al.*, (2002) *Cell* 110, 763-773; Ogiso *et al.*, (2002) *Cell* 110, 775-787). In contrast, in the structures of the inactive HER1, HER3 and HER4, the dimerisation loop is engaged in intramolecular interactions with subdomain IV, which prevents receptor dimerisation in the absence of ligand (Cho and Leahy, (2002) *Science* 297, 1330-1333; Ferguson *et al.*, (2003) *Mol Cell* 12, 541-552; Bouyan *et al.*, (2005) *PNAS* 102, 15024-15029).
15 The structure of HER2 is unique among the HERs. In the absence of a ligand, HER2 has a conformation that resembles the ligand-activated state of HER1 with a protruding dimerisation loop, available to interact with other HER receptors (Cho *et al.*, (2003) *Nature* 421, 756-760; Garrett *et al.*, (2003) *Mol Cell* 11, 495-505). This may explain the enhanced heterodimerisation capacity of HER2.

20 Although the HER receptor crystal structures provide a model for HER receptor homo- and heterodimerisation, the background for the prevalence of some HER homo- and heterodimers over others (Franklin *et al.*, (2004) *Cancer Cell* 5, 317-328) as well as the conformational role of each of the domain in receptor dimerisation and autoinhibition (Burgess *et al.*, (2003) *Mol Cell* 12, 541-552; Mattoon *et al.*, (2004) *PNAS* 101, 923-928) remains somewhat unclear. As
25 described below, the HER3 X-ray crystal structure provides more insights.

HER3 Structure and Conformational Epitopes

A conformational epitope to which antigen binding proteins, e.g., anti-HER3 antibodies bind is provided herein. For the first time, the three dimensional structure of a truncated form
30 (residues 20-640) of the extracellular domain of HER3 complexed with an antibody have been shown. The HER3-MOR09823 Fab complex and the HER3-MOR09825 have been determined at 3.2Å and 3.4 Å resolution, respectively, and shown in Figure 5A. The disclosure herein also shows for the first time an antibody or fragment thereof that binds to an

inactive state of HER3 and stabilizes the receptor in the inactive state. The antibodies of the invention also permit concurrent binding of a HER3 ligand, such as neuregulin with the HER3 receptor.

Although not bound to provide a theory, one possible model for the mechanism of action is that HER3 typically exists in an inactive (closed, tethered) or active (open) state. Ligand binding induces a conformational change such that HER3 exists in the active (open) state which is capable of binding heterodimer partners resulting in activation in downstream signaling. Antibodies such as MOR09823 bind the inactive (tethered) state of HER3 but do not block the ligand binding site. Antibodies such as MOR09823 inhibit HER3 by preventing the ligand induced structural rearrangements required for HER3 to transition to the active conformation, thereby preventing signal transduction. In one embodiment, the antibodies of the invention or fragments thereof bind the inactive (tethered) state of HER3 but do not block the ligand binding site. In another embodiment, the antibodies or fragments thereof inhibit HER3 by preventing the ligand-induced structural rearrangements required for HER3 to transition to the active conformation, thereby preventing signal transduction. In another embodiment, the antibody or fragment thereof stabilizes (directly maintains, locks, tethers, holds, preferentially binds, or favors) HER3 receptor in the inactive state or conformation. In one embodiment, the inactive HER3 receptor may be susceptible to preferential internalization or degradation such that it leads to loss of cell surface HER3 receptors. The biological data presented in the Examples section supports these embodiments.

The crystals of HER3 may be prepared by expressing a nucleotide sequence encoding HER3 or a variant thereof in a suitable host cell, and then crystallising the purified protein(s) in the presence of the relevant HER3 targeted Fab. Preferably the HER3 polypeptide contains the extracellular domain (amino acids 20 to 640 of the human polypeptide or a truncated version thereof, preferably comprising amino acids 20-640) but lacks the transmembrane and intracellular domains.

HER3 polypeptides may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), histidine (HIS), hexahistidine (6HIS), GAL4 (DNA binding and/or transcriptional activation domains) and beta-galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal

of fusion protein sequences.

After expression, the proteins may be purified and/or concentrated, for example by immobilised metal affinity chromatography, ion-exchange chromatography, and/or gel
5 filtration.

The protein(s) may be crystallised using techniques described herein. Commonly, in a crystallisation process, a drop containing the protein solution is mixed with the crystallisation buffer and allowed to equilibrate in a sealed container. Equilibration may be achieved by
10 known techniques such as the “hanging drop” or the “sitting drop” method. In these methods, the drop is hung above or sitting beside a much larger reservoir of crystallization buffer and equilibration is reached through vapor diffusion. Alternatively, equilibration may occur by other methods, for example under oil, through a semi-permeable membrane, or by free-interface diffusion (See e.g., Chayen *et al.*, (2008) *Nature Methods* 5, 147 - 153).

15
Once the crystals have been obtained, the structure may be solved by known X-ray diffraction techniques. Many techniques use chemically modified crystals, such as those modified by heavy atom derivatization to approximate phases. In practice, a crystal is soaked in a solution containing heavy metal atom salts, or organometallic compounds, e.g., lead chloride, gold
20 thiomalate, thimerosal or uranyl acetate, which can diffuse through the crystal and bind to the surface of the protein. The location(s) of the bound heavy metal atom(s) can then be determined by X-ray diffraction analysis of the soaked crystal. The patterns obtained on diffraction of a monochromatic beam of X-rays by the atoms (scattering centres) of the crystal can be solved by mathematical equations to give mathematical coordinates. The diffraction
25 data are used to calculate an electron density map of the repeating unit of the crystal. Another method of obtaining phase information is using a technique known as molecular replacement. In this method, rotational and translational algorithms are applied to a search model derived from a related structure, resulting in an approximate orientation for the protein of interest (See Rossmann, (1990) *Acta Crystals A* 46, 73-82). The electron density maps are used to establish
30 the positions of the individual atoms within the unit cell of the crystal (Blundel *et al.*, (1976) *Protein Crystallography*, Academic Press).

The present disclosure describes for the first time, the three-dimensional structure of HER3 and a Fab of an anti-HER3 antibody. The approximate domain boundaries of extracellular

domain of HER3 are as follows; domain 1: amino acids 20-207; domain 2: amino acids 208-328; domain 3: amino acids 329-498; and domain 4: amino acids 499-642. The three-dimensional structure of HER3 and the antibody also allows the identification of target binding sites for potential HER3 modulators. Preferred target binding sites are those involved in the activation of HER3. In one embodiment, the target binding site is located within domain 2 and domain 4 of HER3. Thus an antibody or fragment thereof which binds to either domain 2 or domain 4, and preferably to both domains can modulate HER3 activation by either preventing the domains from dissociation from each other or by modifying the relative positions of the domains. Thus binding an antibody or fragment thereof to amino acid residues within domain 2 or domain 4 may cause the protein to adopt a conformation that prevents activation. The disclosure herein also shows for the first time an antibody or fragment thereof that can concurrently bind with a HER3 ligand, such as neuregulin.

In some embodiments, the antibody or fragment thereof recognize a specific conformational state of HER3 such that the antibody or fragment thereof prevents HER3 from interacting with a co-receptor (including, but not limited to, HER1, HER2 and HER4). In some embodiments, the antibody or fragment thereof prevents HER3 from interacting with a co-receptor by stabilizing the HER3 receptor in an inactive or closed state. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor by binding to amino acid residues within domain 2 and domain 4 of HER3. In this inactive state, the dimerization loop located within domain 2 is not exposed and therefore unavailable for dimerization with other co-receptors (including, but not limited to, HER1, HER2 and HER4). In some embodiments, the antibody or fragment thereof binds to human HER3 protein having a conformational epitope comprising (i) HER3 amino acid residues 265-277 and 315 (of domain 2) and (ii) HER3 amino acid residues 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) of SEQ ID NO: 1, or a subset thereof. In some embodiments, the antibody or fragment thereof binds to amino acids within or overlapping amino acid residues 265-277 and 315 (of domain 2) and (ii) HER3 amino acid residues 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) of SEQ ID NO: 1. In some embodiments, the antibody or fragment thereof binds to amino acids within (and/or amino acid sequences consisting of) amino acids 265-277 and 315 (of domain 2) and (ii) HER3 amino acid residues 571, 582-584, 596-597, 600-602, 609-615 (of domain 4) of SEQ ID NO: 1, or a subset thereof. In some embodiments, the antibody or fragment thereof binds to the conformational epitope such that it restricts the mobility of domain 2 and domain 4, stabilizing it in an inactive or closed conformation. The failure to form the active conformation results in failure to activate signal transduction. In some embodiments, the

antibody or fragment thereof binds to the conformational epitope such that it occludes the dimerization loop within domain 2, thereby rendering it unavailable for receptor-receptor interaction. The failure to form homo- or heterodimers results in failure to activate signal transduction.

5 In another aspect, the antibody or fragment thereof binds a conformational epitope of HER receptor, such as a HER3 receptor. In one embodiment, the antibody or fragment thereof stabilizes the HER3 receptor in the inactive state. In another embodiment, the antibody or fragment thereof binds to the active state of the HER3 receptor and drives it into the inactive state as the inactive state. Thus, the antibody or fragment thereof can bind to either the active
10 or inactive state of HER3, but favors the formation of the inactive state and drives the active state of HER3 into the inactive state, resulting in a failure to activate signal transduction.

In another aspect, the antibody or fragment thereof binds a conformational epitope of HER receptor, such as a HER3 receptor where binding of the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state such that the HER3 receptor fails to dimerize
15 with a co-receptor to form a receptor-receptor complex. The failure to form a receptor-receptor complex prevents activation of both ligand-dependent and ligand-independent signal transduction.

In another aspect, the antibody or fragment thereof binds a conformational epitope of HER receptor such as a HER3 receptor, where binding of the antibody or fragment thereof to the
20 HER3 receptor allows dimerization with a co-receptor to form an inactive receptor-receptor complex. The formation of the inactive receptor-receptor complex prevents activation of ligand-independent signal transduction. For example, in ligand-independent signal transduction, HER3 may exist in an inactive state, however the overexpression of HER2 causes HER2-HER3 complex formation, however these resulting complexes are inactive and
25 prevent activation of ligand-independent signal transduction.

The depicted structure also allows one to identify specific core HER3 amino acid residues for the interaction interface of an antibody or fragment thereof (e.g., MOR09823) with HER3.

This was defined as residues that are within 5Å of the MOR09823 protein VH chain. The core
30 residues are as follows: Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597.

The structures can also be used to identify boundary HER3 amino acid residues for the interaction interface with an antibody or fragment thereof (e.g., MOR09823). These residues can be HER3 residues that were 5-8Å from the MOR09823 protein VH chain. The boundary residues are as follows: Pro262, Val264, Tyr265, Phe270, Leu272, Thr278, Lys314, Gly316, Glu321, Asn566, Ser568, Gly569, Ser570, Thr572, Arg580, Asp581, Gly582, Gly595, Gly598, Ile600.

The depicted structure also allows one to identify specific core HER3 amino acid residues for the interaction interface of an antibody or fragment thereof (e.g., MOR09823) with HER3. This was defined as residues that are within 5Å of the MOR09823 protein VL chain. The core residues are as follows: Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615.

The structures were also used to identify boundary HER3 amino acid residues for the interaction interface with an antibody or fragment thereof (e.g., MOR09823). These residues were HER3 residues that were 5-8Å from the MOR09823 protein VL chain. The boundary residues are as follows: Asn266, Thr269, Asp571, Arg580, Asp581, His584, Pro590, Ala596, Pro599, Tyr601, Tyr603, Asp605, Gln607, Cys610, Asn616, Cys617, Cys621, Gly623, Pro624.

As can be seen in Tables 11 and 12 (MOR09823) and Tables 13 and 14 (MOR09825), respectively, the heavy chain is mainly involved in the antigen binding protein's binding to amino acid residues within domain 2 of the epitope with fewer interactions with amino acid residues of domain 4, while the light chain is mainly involved with binding to amino acid residues within domain 4 of the epitope with fewer interactions with amino acid residues within domain 2.

As such, one of skill in the art, given the present teachings, can predict which residues and areas of the antigen binding proteins can be varied without unduly interfering with the antigen binding protein's ability to bind to HER3.

Core interaction interface amino acids were determined as being all amino acid residues with at least one atom less than or equal to 5Å from the HER3 partner protein. 5Å was chosen as the core region cutoff distance to allow for atoms within a van der Waals radius plus a possible water-mediated hydrogen bond. Boundary interaction interface amino acids were

determined as all amino acid residues with at least one atom less than or equal to 8Å from the HER3 partner protein but not included in the core interaction list.

In some embodiments, any antigen binding protein that binds to, covers, or prevents MOR09823 from interacting with any of the above residues can be employed to bind to or neutralize HER3. In some embodiments, the antibodies or fragments thereof binds to or interacts with at least one of the following HER3 residues (SEQ ID NO: 1): Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597. In some embodiments, the antibodies and fragments thereof binds to or interacts with at least one of the following HER3 residues (SEQ ID NO: 1): Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615. In some embodiments, the antibodies or fragments thereof binds to or interacts with at least one of the following HER3 residues (SEQ ID NO: 1): Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597, Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615. In some embodiments, the antibodies or fragments thereof binds to or interacts with a combination of the following HER3 residues (SEQ ID NO: 1): Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597, Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615. In some embodiments, the antibodies or fragments thereof binds to or interacts with all of the following HER3 residues (SEQ ID NO: 1): Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597, Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615. In some embodiments, the antibody or fragment thereof is within 5 angstroms of one or more of the above residues. In some embodiments, the antibody or fragment thereof is 5 to 8 angstroms from one or more of the above residues. In some embodiments, the antibody or fragment thereof interacts, blocks, or is within 8 angstroms of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, or 50 of the above residues.

The availability of 3D structures for the HER3 and the complex of HER3:MOR09823, for example, provides the framework to explore other HER3 antibodies in more detail. The 3D

structure of HER3 allows the epitopes for monoclonal antibodies to be mapped and their mode of action inferred, since some inhibit, some stimulate and others have no effect on cell growth. The conformational epitope for MOR09823 has been located to the domains 2 and 4 of HER3. The availability of the 3D structures of this receptor will facilitate the determination of the precise mechanism of action of these inhibitory agents and the design of new approaches to interfering with HER3 receptor function. In one embodiment, the antibodies of the invention bind to the same conformational epitope as MOR09823.

In some embodiments, the conformational epitope bound by any of the antibodies listed in Table 1 is especially useful. In certain embodiments, a HER3 conformational epitope can be utilized to isolate antibodies or fragments thereof that bind to HER3. In certain embodiments, a HER3 conformational epitope can be utilized to generate antibodies or fragments thereof which bind to HER3. In certain embodiments, a HER3 conformational epitope can be utilized as an immunogen to generate antibodies or fragments thereof that bind to the HER3 conformational epitope. In certain embodiments, a HER3 conformational epitope can be administered to an animal, and antibodies that bind to HER3 can subsequently be obtained from the animal.

In some embodiments, the domain(s)/region(s) containing residues that are in contact with or are buried by an antibody can be identified by mutating specific residues in HER3 (e.g., a wild-type antigen) and determining whether antibody or fragment thereof can bind the mutated or variant HER3 protein or measure changes of affinity from wild-type. By making a number of individual mutations, residues that play a direct role in binding or that are in sufficiently close proximity to the antibody such that a mutation can affect binding between the antibody and antigen can be identified. From a knowledge of these amino acids, the domain(s) or region(s) of the antigen (HER3) that contain residues in contact with the antibody or covered by the antibody can be elucidated. Mutagenesis using known techniques such as alanine-scanning can help define functionally relevant epitopes. Mutagenesis utilizing an arginine/glutamic acid scanning protocol can also be employed (see, e.g., Nanevicz *et al.*, (1995), *J. Biol. Chem.* 270(37):21619-21625 and Zupnick *et al.*, (2006), *J. Biol. Chem.* 281(29):20464-20473). In general, arginine and glutamic acids are substituted (typically individually) for an amino acid in the wild-type polypeptide because these amino acids are charged and bulky and thus have the potential to disrupt binding between an antigen binding

protein and an antigen in the region of the antigen where the mutation is introduced. Arginines that exist in the wild-type antigen are replaced with glutamic acid. A variety of such individual mutants can be obtained and the collected binding results analyzed to determine what residues affect binding. A series of mutant HER3 antigens can be created, with each mutant antigen having a single mutation. Binding of each mutant HER3 antigen with various HER3 antibodies or fragments thereof can be measured and compared to the ability of the selected an antibody or fragments thereof to bind wild-type HER3 (SEQ ID NO: 1).

An alteration (for example a reduction or increase) in binding between an antibody or fragment thereof and a mutant or variant HER3 as used herein means that there is a change in binding affinity (e.g., as measured by known methods such as Biacore testing or the bead based assay described below in the examples), EC_{50} , and/or a change (for example a reduction) in the total binding capacity of the antigen binding protein (for example, as evidenced by a decrease in B_{max} in a plot of antigen binding protein concentration versus antigen concentration). A significant alteration in binding indicates that the mutated residue is involved in binding to the antibody or fragment thereof.

In some embodiments, a significant reduction in binding means that the binding affinity, EC_{50} , and/or capacity between an antibody or fragments thereof and a mutant HER3 antigen is reduced by greater than 10%, greater than 20%, greater than 40%, greater than 50%, greater than 55%, greater than 60%, greater than 65%, greater than 70%, greater than 75%, greater than 80%, greater than 85%, greater than 90% or greater than 95% relative to binding between the an antibody or fragment thereof and a wild type HER3 (e.g., SEQ ID NO: 1).

In some embodiments, binding of an antibody or fragments thereof is significantly reduced or increased for a mutant HER3 protein having one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) mutations as compared to a wild-type HER3 protein (e.g., SEQ ID NO: 1).

Although the variant forms are referenced with respect to the wild-type sequence shown in SEQ ID NO: 1, it will be appreciated that in an allelic or splice variants of HER3 the amino acids could differ. Antibodies or fragments thereof showing significantly altered binding (e.g., lower or higher binding) for such allelic forms of HER3 are also contemplated.

In addition to the general structural aspects of antibodies, the more specific interaction between the paratope and the epitope may be examined through structural approaches. In one embodiment, the structure of the CDRs contribute to a paratope, through which an antibody is able to bind to an epitope. The shape of such a paratope may be determined in a number of ways. Traditional structural examination approaches can be used, such as NMR or x-ray crystallography. These approaches can examine the shape of the paratope alone, or while it is bound to the epitope. Alternatively, molecular models may be generated in silico. A structure can be generated through homology modeling, aided with a commercial package, such as InsightII modeling package from Accelrys (San Diego, Calif.). Briefly, one can use the sequence of the antibody to be examined to search against a database of proteins of known structures, such as the Protein Data Bank. After one identifies homologous proteins with known structures, these homologous proteins are used as modeling templates. Each of the possible templates can be aligned, thus producing structure based sequence alignments among the templates. The sequence of the antibody with the unknown structure can then be aligned with these templates to generate a molecular model for the antibody with the unknown structure. As will be appreciated by one of skill in the art, there are many alternative methods for generating such structures in silico, any of which may be used. For instance, a process similar to the one described in Hardman et al., issued U.S. Pat. No. 5,958,708 employing QUANTA (Polygen Corp., Waltham, Mass.) and CHARM (Brooks *et al.*, (1983), J. Comp. Chem. 4:187) may be used (hereby incorporated in its entirety by reference).

Not only is the shape of the paratope important in determining whether and how well a possible paratope will bind to an epitope, but the interaction itself, between the epitope and the paratope is a source of great information in the design of variant antibodies. As appreciated by one of skill in the art, there are a variety of ways in which this interaction can be studied. One way is to use the structural model generated, perhaps as described above, and then to use a program such as InsightII (Accelrys, San Diego, Calif.), which has a docking module, which, among other things, is capable of performing a Monte Carlo search on the conformational and orientational spaces between the paratope and its epitope. The result is that one is able to estimate where and how the epitope interacts with the paratope. In one embodiment, only a fragment, or variant, of the epitope is used to assist in determining the relevant interactions. In one embodiment, the entire epitope is used in the modeling of the interaction between the paratope and the epitope.

Through the use of these modelled structures, one is able to predict which residues are the most important in the interaction between the epitope and the paratope. Thus, in one embodiment, one is able to readily select which residues to change in order to alter the binding characteristics of the antibody. For instance, it may be apparent from the docking models that the side chains of certain residues in the paratope may sterically hinder the binding of the epitope, thus altering these residues to residues with smaller side chains may be beneficial. One can determine this in many ways. For example, one may simply look at the two models and estimate interactions based on functional groups and proximity. Alternatively, one may perform repeated pairings of epitope and paratope, as described above, in order to obtain more favorable energy interactions. One can also determine these interactions for a variety of variants of the antibody to determine alternative ways in which the antibody may bind to the epitope. One can also combine the various models to determine how one should alter the structure of the antibodies in order to obtain an antibody with the particular characteristics that are desired.

The models determined above can be tested through various techniques. For example, the interaction energy can be determined with the programs discussed above in order to determine which of the variants to further examine. Also, coulombic and van der Waals interactions are used to determine the interaction energies of the epitope and the variant paratopes. Also site directed mutagenesis is used to see if predicted changes in antibody structure actually result in the desired changes in binding characteristics. Alternatively, changes may be made to the epitope to verify that the models are correct or to determine general binding themes that may be occurring between the paratope and the epitope.

As will be appreciated by one of skill in the art, while these models will provide the guidance necessary to make the antibodies and variants thereof of the present embodiments, it may still be desirable to perform routine testing of the in silico models, perhaps through in vitro studies. In addition, as will be apparent to one of skill in the art, any modification may also have additional side effects on the activity of the antibody. For instance, while any alteration predicted to result in greater binding, may induce greater binding, it may also cause other structural changes which might reduce or alter the activity of the antibody. The determination of whether or not this is the case is routine in the art and can be achieved in many ways. For example, the activity can be tested through an ELISA test. Alternatively, the samples can be

tested through the use of a surface plasmon resonance device.

HER3 Antibodies

The present invention provides antibodies that recognize a conformational epitope of HER3.

5 The invention is based on the surprising finding that a class of antibodies against HER3, block both ligand-dependent and ligand-independent HER3 signal transduction pathways. A class of antibodies that bind to the particular conformation epitope of HER3 is disclosed in Table 1. In one embodiment, the antibodies inhibit both ligand-dependent and ligand-independent HER3 signalling. In another embodiment, the antibodies bind to HER3 and do not block
10 HER ligand binding to the ligand binding site (i.e. both ligand and antibody can bind HER3 concurrently).

The present invention provides antibodies that specifically bind a HER3 protein (*e.g.*, human and/or cynomologus HER3), said antibodies comprising a VH domain having an amino acid sequence of SEQ ID NO: 15, 33, 51, 69, 87, 105, 123, 141, 159, 177, 195, 213, 231, 249, 267,
15 285, 303, 321, 339, 357, and 375. The present invention provides antibodies that specifically bind a HER3 protein (*e.g.*, human and/or cynomologus HER3), said antibodies comprising a VL domain having an amino acid sequence of SEQ ID NO: 14, 32, 50, 68, 86, 104, 122, 140, 158, 176, 194, 212, 230, 248, 266, 284, 302, 320, 338, 356, and 374. The present invention also provides antibodies that specifically bind to a HER3 protein (*e.g.*, human and/or
20 cynomologus HER3), said antibodies comprising a VH CDR having an amino acid sequence of any one of the VH CDRs listed in Table 1, *infra*. In particular, the invention provides antibodies that specifically bind to a HER3 protein (*e.g.*, human and/or cynomologus HER3), said antibodies comprising (or alternatively, consisting of) one, two, three, four, five or more VH CDRs having an amino acid sequence of any of the VH CDRs listed in Table 1, *infra*.

25 Other antibodies of the invention include amino acids that have been mutated, yet have at least 60, 70, 80, 90, 95, or 98 percent identity in the CDR regions with the CDR regions depicted in the sequences described in Table 1. In some embodiments, it includes mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated in the CDR regions when compared with the CDR regions depicted in the sequence described
30 Table 1, while still maintaining their specificity for the original antibody's epitope

Other antibodies of the invention include amino acids that have been mutated, yet have at least 60, 70, 80, 90, 95, or 98 percent identity in the framework regions with the framework

regions depicted in the sequences described in Table 1. In some embodiments, it includes mutant amino acid sequences wherein no more than 1, 2, 3, 4, 5, 6, or 7 amino acids have been mutated in the framework regions when compared with the framework regions depicted in the sequence described Table 1, while still maintaining their specificity for the original antibody's epitope. The present invention also provides nucleic acid sequences that encode VH, VL, the full length heavy chain, and the full length light chain of the antibodies that specifically bind to a HER3 protein (e.g., human and/or cynomolgus HER3).

The HER3 antibodies of the invention bind to the conformational epitope of HER3 comprising amino acid residues from domain 2 and domain 4 of HER3.

10

Table 1: Examples of HER3 Antibodies of the Present Invention

SEQ ID NUMBER	Ab region	
MOR09823		
SEQ ID NO: 2 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 3 (Kabat)	HCDR2	VTGAVGRITYYPDSVKG
SEQ ID NO: 4 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 5 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 6 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 7 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 8 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 9 (Chothia)	HCDR2	GAVGR
SEQ ID NO: 10 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 11 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 12 (Chothia)	LCDR2	GAS
SEQ ID NO: (Chothia) 13	LCDR3	YSSFPT
SEQ ID NO: 14	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSGTDFTLTISLQPEDFAVYYCQQYSSFPPTFGQ GTKVEIK
SEQ ID NO: 15	VH	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV TGAVGRITYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGD EGFDIWGQGTLLVTVSS
SEQ ID NO: 16	DNA VL	GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGGAGCGTGGGTGATCGTG TGACCATTACCTGCAGAGCGAGCCAGGGTATTTCTAATTTGGCTGGCTTGGTACCA GCAGAAACCAGGTAAAGCACCGAAACTATTAATTTATGGTGCTTCTTCTTTGCAA AGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTTATTTGCCAGCAGTATTC TTCTTTTCTACTACTCTTTGGCCAGGGTACGAAAGTTGAAATTTAAA
SEQ ID NO: 17	DNA VH	CAGGTGCAATTTGGTGGAAAGCGGCGGCGCCTGGTGCAACCGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCCTGGGAAGGGTCTCGAGTGGGTGAGCGTTACTGGTGCTGTTGGT CGTACTTATATCTCTGATCTGTAAAGGTCGTTTTACCATTTACGTTGATAATF CGAAAAACACCCTGTATCTGCAAAATGAACAGCCTGCGTGCGGAAGATACGGCCGT GTATTATTTGCGCGGCTTGGGGTGTAGGGTTTTGATATTTGGGGCCAAAGGCACC CTGGTGACGGTTAGCTCA
SEQ ID NO: 18	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSGTDFTLTISLQPEDFAVYYCQQYSSFPPTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSVTLTSLKADYEKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 19	Heavy IgG1	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV TGAVGRITYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGD EGFDIWGQGTLLVTVSSASTKGPSVFLPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICN VNHKPSNTKVDKRVKPKCDKHTHTCPPCPAPELGGPSVFLFPPKPKDTL MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR

		VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
MOR09824		
SEQ ID NO: 20 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 21 (Kabat)	HCDR2	VISAWGHVKYYADSVKG
SEQ ID NO: 22 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 23 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 24 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 25 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 26 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 27 (Chothia)	HCDR2	SAWGHV
SEQ ID NO: 28 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 29 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 30 (Chothia)	LCDR2	GAS
SEQ ID NO: 31 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 32	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYQCQQYSSFPTTFGQ GTKVEIK
SEQ ID NO: 33	VH	QVQLVESGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV ISAWGHVKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWG DEGFDIWGQGLTVVSS
SEQ ID NO: 34	DNA VL	GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGGAGCGTGGGTGATCGTG TGACCATTACCTGCAGAGCGAGCCAGGGTATTTCTAATTGGCTGGCTTGGTACCA GCAGAAACCAGGTAAAGACCCGAACTATTAATTTATGGTGTCTTCTTTGCAA AGCGGGTCCCCTCCCCTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCTGA CCATTAGCAGCCTGCACCTGAAGACTTTCGGTTTATTTATGCCAGCAGTATTC TTCTTTTCTACTACCTTTGGCCAGGGTACGAAAGTTGAAATTTAA
SEQ ID NO: 35	DNA VH	CAGGTGCAATTGGTGGAAAGCGCGCGCCCTGGTGCAACCGGGCGGCAGCCTGC GTCTGAGCTGCGCGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCTGGGAAGGGTCTCGAGTGGGTGAGCGTTATTTCTGCTTGGGGT CATGTTAAGTATTATGCTGATTCTGTTAAGGGTCGTTTTACCATTTACGTGATA ATTCGAAAAACCCCTGTATCTGCAAAATGAACAGCCTGCGTGCGGAAGATACGGC CGTGTATTATTGCGCGCGTTGGGGTGTAGGGTTTTGTATTTGGGGCCAAGGC ACCTGGTGACGGTTAGCTCA
SEQ ID NO: 36	Light Kappa	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYQCQQYSSFPTTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFFYPREAKVQWKV DNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 37	Heavy IgG1	QVQLVESGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV ISAWGHVKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWG DEGFDIWGQGLTVVSSASTKGPSVFPPLAPSSKSTSGGTAALGLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
MOR09825		
SEQ ID NO: 38 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 39 (Kabat)	HCDR2	AINSQGKSTYYADSVKG
SEQ ID NO: 40 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 41 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 42 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 43 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 44 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 45 (Chothia)	HCDR2	NSQGKS
SEQ ID NO: 46 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 47 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 48 (Chothia)	LCDR2	GAS
SEQ ID NO: 49 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 50	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYQCQQYSSFPTTFGQ

		GTKVEIK
SEQ ID NO: 51	VH	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSA INSQGKSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGLVTVSS
SEQ ID NO: 52	DNA VL	GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCAGCGTGGGTGATCGTG TGACCATTACCTGCAGAGCGAGCCAGGGTATTTCTAATTGGCTGGCTTGGTACCA GCAGAAACCAGGTAAAGCACCAGAACTATTAATTTATGGTGCTTCTCTTTGCAA AGCGGGTCCCCTCCCCTTTAGCGGCTCTGGATCCGGCACTGATTTACCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTATTGCCAGCAGTATTC TTCTTTTCTACTACTCTTTGGCCAGGGTACGAAAGTTGAAATTTAAA
SEQ ID NO: 53	DNA VH	CAGGTGCAATTGGTGGAAAGCGGCGGCGCCTGGTGCACCAGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCCTGGGAAGGGTCTCGAGTGGGTGAGCGCTATTAATTTCTCAGGGT AAGTCTACTTATTATGCTGATTCTGTTAAGGGTCTTTTTACCATTTACGTTGATA ATTGAAAAACACCCTGTATCTGCAAATGAACAGCCTGCGTGCAGGAAGATACGGC CGTGTATTATTGCGCGCCTTGGGGTATGAGGGTTTTGATATTTGGGGCCAAGGC ACCCTGGTGACGGTTAGCTCA
SEQ ID NO: 54	Light Kappa	DIQMTQSPSSLSASVGRVITITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPFTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 55	Heavy IgG1	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSA INSQGKSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGLVTVSSASTKGPSVFPLAPSSKSTSGGTAAALGLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYF LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDLSD DGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYHTQKLSLSLSPGK
MOR09974		
SEQ ID NO: 56 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 57 (Kabat)	HCDR2	VINPSGNFTNYADSVKG
SEQ ID NO: 58 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 59 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 60 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 61 (Kabat)	LCDR3	QQYSSFPFT
SEQ ID NO: 62 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 63 (Chothia)	HCDR2	NPSGNF
SEQ ID NO: 64 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 65 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 66 (Chothia)	LCDR2	GAS
SEQ ID NO: 67 (Chothia)	LCDR3	YSSFPFT
SEQ ID NO: 68	VL	DIQMTQSPSSLSASVGRVITITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPFTFGQ GTKVEIK
SEQ ID NO: 69	VH	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV INPSGNFTNYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGLVTVSS
SEQ ID NO: 70	DNA VL	GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCAGCGTGGGTGATCGTG TGACCATTACCTGCAGAGCGAGCCAGGGTATTTCTAATTGGCTGGCTTGGTACCA GCAGAAACCAGGTAAAGCACCAGAACTATTAATTTATGGTGCTTCTCTTTGCAA AGCGGGTCCCCTCCCCTTTAGCGGCTCTGGATCCGGCACTGATTTACCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTATTGCCAGCAGTATTC TTCTTTTCTACTACTCTTTGGCCAGGGTACGAAAGTTGAAATTTAAA
SEQ ID NO: 71	DNA VH	CAGGTGCAATTGGTGGAAAGCGGCGGCGCCTGGTGCACCAGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCCTGGGAAGGGTCTCGAGTGGGTGAGCGTTATTAATCTTCTGGT AATTTTACTAATTATGCTGATTCTGTTAAGGGTCTTTTTACCATTTACGTTGATA ATTGAAAAACACCCTGTATCTGCAAATGAACAGCCTGCGTGCAGGAAGATACGGC CGTGTATTATTGCGCGCCTTGGGGTATGAGGGTTTTGATATTTGGGGCCAAGGC ACCCTGGTGACGGTTAGCTCA
SEQ ID NO: 72	Light Kappa	DIQMTQSPSSLSASVGRVITITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPFTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYKHKVYACEVTHQG LSSPVTKSFNRGEC

SEQ ID NO: 73	Heavy IgG1	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV INPSGNFTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWG DEGFDIWGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTVSWNSGALTSVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYIC NVNHKPSNTKVKDRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYF LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDS DGSFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGK
MOR10452		
SEQ ID NO: 74 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 75 (Kabat)	HCDR2	NTSPIGYTYAGSVKG
SEQ ID NO: 76 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 77 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 78 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 79 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 80 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 81 (Chothia)	HCDR2	SPIGY
SEQ ID NO: 82 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 83 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 84 (Chothia)	LCDR2	GAS
SEQ ID NO: 85 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 86	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPTTFGQ GTKVEIK
SEQ ID NO: 87	VH	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSN TSP IGYTYAGSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGD EGFDIWGQGT LVTVSS
SEQ ID NO: 88	DNA VL	GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCGAGCGTGGGTGATCGTG TGACCATTACCTGCAGAGCGAGCCAGGGTATTTCTAATTGGCTGGCTGGTACCA GCAGAAACCAGGTAAAGCACCGAACTATTAATTTATGGTGCTTCTCTTTGCAA AGCGGGTCCCCTCCCCTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATATATGCCAGCAGTATTC TCTTTTCTACTACCTTTGGCCAGGGTACGAAAGTTGAAATTTAAA
SEQ ID NO: 89	DNA VH	CAGGTGCAATTTGGTGGAAAGCGCGCGCGCCTGGTGCAACCGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCCTGGGAAGGGTCTCGAGTGGGTGAGCAATACTTCTCCTATTGGT TATACTTATATGCTGGTCTGTTAAGGGTCTGTTTACCATTTACGCTGATAATT CGAAAAACACCCTGTATCTGCAAATGAACAGCCTGCGTGGGAAAGATACGGCCGT GTATTATTGCGCGCCTTGGGGTGTGAGGGTTTTGATATTTGGGGCCAAGGCACC CTGGTGACGGTTAGCTCA
SEQ ID NO: 90	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPTTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQG LSSPVTKSFNRGEA
SEQ ID NO: 91	Heavy Chain (only VH and CH1 domains)	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSN TSP IGYTYAGSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGD EGFDIWGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF EPVTVSWNSGALTSVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICN VNHKPSNTKVKDKVEPKS
MOR10701		
SEQ ID NO: 92 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 93 (Kabat)	HCDR2	VTGAVGRSTYYPDSVKG
SEQ ID NO: 94 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 95 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 96 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 97 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 98 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 99 (Chothia)	HCDR2	GAVGRS
SEQ ID NO: 100 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 101 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 102 (Chothia)	LCDR2	GAS
SEQ ID NO: 103 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 104	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTFLTISSLQPEDFAVYYCQQYSSFPTTFGQ

		GTKVEIK
SEQ ID NO: 105	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV TGAVGRSTYYPDSVKGRFTISRDN SKNTLYLQMNLSRAEDTAVYYCARWG DEGFDIWGQGLVTVSS
SEQ ID NO: 106	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGCGACAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGCAAGGCCCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTCACCCTGA CCATCAGCAGCCTGCAGCCCAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 107	DNA VH	GAGGTGCAATTGCTGGAAGCGCGGAGGCGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCGCCGCCAGCGGCTTCACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGACAGGCGCCGTGGGC AGAAGCACCTACTACCCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGGGGCCGAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCCTGGTCACCGTCAGCTCA
SEQ ID NO: 108	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTLTISLQPEDFATYYCQQYSSFPFTTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 109	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV TGAVGRSTYYPDSVKGRFTISRDN SKNTLYLQMNLSRAEDTAVYYCARWG DEGFDIWGQGLVTVSSASTKGPSVFPLAPSSKSTSGGTAAALGLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC NVNHKPSNTKVDKRVPEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYV LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDL DGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYHTQKSLSLSPGK
MOR10702		
SEQ ID NO: 110 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 111 (Kabat)	HCDR2	VI SAWGHVKYYADSVKG
SEQ ID NO: 112 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 113 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 114 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 115 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 116 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 117 (Chothia)	HCDR2	SAWGHV
SEQ ID NO: 118 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 119 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 120 (Chothia)	LCDR2	GAS
SEQ ID NO: 121 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 122	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTLTISLQPEDFATYYCQQYSSFPFTTFGQ GTKVEIK
SEQ ID NO: 123	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV ISAWGHVKYYADSVKGRFTISRDN SKNTLYLQMNLSRAEDTAVYYCARWG DEGFDIWGQGLVTVSS
SEQ ID NO: 124	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGCGACAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGCAAGGCCCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTCACCCTGA CCATCAGCAGCCTGCAGCCCAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 125	DNA VH	GAGGTGCAATTGCTGGAAGCGCGGAGGCGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCGCCGCCAGCGGCTTCACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGATCAGCGCCTGGGGC CACGTGAAGTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTTCGGGGCCGAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCCTGGTCACCGTCAGCTCA
SEQ ID NO: 126	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTLTISLQPEDFATYYCQQYSSFPFTTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 127	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSV

		ISAWGHVKYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGT LVTVSSASTKGP SVFPLAPSSKSTSGGTAALGLVKDYF PEPVTVSWNSGALTSVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEL LGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
MOR10703		
SEQ ID NO: 128 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 129 (Kabat)	HCDR2	AIN SQGKSTYYADSVKG
SEQ ID NO: 130 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 131 (Kabat)	LCDR1	RASQGISNWL A
SEQ ID NO: 132 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 133 (Kabat)	LCDR3	QQYSSFPPT
SEQ ID NO: 134 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 135 (Chothia)	HCDR2	NSQGKS
SEQ ID NO: 136 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 137 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 138 (Chothia)	LCDR2	GAS
SEQ ID NO: 139 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 140	VL	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTLTITSSLQPEDFATYYCQQYSSFPPTFGQ GTKVEIK
SEQ ID NO: 141	VH	EVQLLESGGGLVQPGGSLRLS CAASGFTFSSYAMSWVRQAPGKLEWVSA INSQGKSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGT LVTVSS
SEQ ID NO: 142	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGCGCAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGCAAGGCCCAAGCTGCTGATCTACGGCCAGCTCCCTGCAG AGCGCGTGCACAGCAGATTCAGCGCCAGCGGCTCCGGCACCGACTTCACCTGA CCATCAGCAGCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 143	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGCCTGGTGCAGCCTGGCGGCAGCCTGA GACTGCTTGCAGCCAGCGGCTTCACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCAACAGCCAGGGC AAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCCTGGTCACCGTCAGCTCA
SEQ ID NO: 144	Light Kappa	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYG ASSLQSGVPSRFRSGSGSDFTLTITSSLQPEDFATYYCQQYSSFPPTFGQ GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQG LSSPVTKSFNRGEC
SEQ ID NO: 145	Heavy IgG1	EVQLLESGGGLVQPGGSLRLS CAASGFTFSSYAMSWVRQAPGKLEWVSA INSQGKSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWG DEGFDIWGQGT LVTVSSASTKGP SVFPLAPSSKSTSGGTAALGLVKDYF PEPVTVSWNSGALTSVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEL LGGPSVFLFPPKPKDT LMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
MOR10703 N52S		
SEQ ID NO: 146 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 147 (Kabat)	HCDR2	AI <u>SS</u> QGKSTYYADSVKG
SEQ ID NO: 148 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 149 (Kabat)	LCDR1	RASQGISNWL A
SEQ ID NO: 150 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 151 (Kabat)	LCDR3	QQYSSFPPT
SEQ ID NO: 152 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 153 (Chothia)	HCDR2	<u>SS</u> QGKS
SEQ ID NO: 154 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 155 (Chothia)	LCDR1	SQGISNW

SEQ ID NO: 156 (Chothia)	LCDR2	GAS
SEQ ID NO: 157 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 158	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIK
SEQ ID NO: 159	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI <u>S</u> SQG KSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 160	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCCAAGCAGATTCAGCGGCAGCGGCTCCGGCACCGACTTACCCTGA CCATCAGCAGCCTGCAGCCCCAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 161	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGCTTTCGCCCGCCAGCGGCTTACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCAGCAGCCAGGGC AAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCACCGTCAGCTCA
SEQ ID NO: 162	Light Kappa	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPREAKVQWKVDNALQSGNSQESVTE QDSKDSITYLSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 163	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI <u>S</u> SQG KSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKSC DKHTCTPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKS LSLSPGK
MOR10703 N52G		
SEQ ID NO: 164 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 165 (Kabat)	HCDR2	AI <u>G</u> SQGGKSTYYADSVKG
SEQ ID NO: 166 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 167 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 168 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 169 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 170 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 171 (Chothia)	HCDR2	<u>G</u> SQGGKS
SEQ ID NO: 172 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 173 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 174 (Chothia)	LCDR2	GAS
SEQ ID NO: 175 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 176	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIK
SEQ ID NO: 177	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI <u>G</u> SQG KSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 178	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA

		GCAGAAGCCCGGCAAGGCCCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTCACCCTGA CCATCAGCAGCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAGGTGGAATCAAG
SEQ ID NO: 179	DNA VH	GAGGTGCAATTGCTGGAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCGCCGACGCGGCTTACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCGGCAGCCAGGGC AAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACCCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCCACCGTCAGCTCA
SEQ ID NO: 180	Light Kappa	DIQMTQSPSSLSASVGDRTVITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGQGTKVEIKRTV AAPSVFI FPPSDEQLKSGTASVCLLNFPYFPAKLVQWKVDNALQSGNSQESVTE QDSKDYSLSSLTLSKADYKHKVYACEVTHQGLSSPVTKS FNRGEC
SEQ ID NO: 181	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI GS QG KSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALT GVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSC DKHTCTPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SAKAGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKS LSLSPGK
MOR10703 N52S_S52aN		
SEQ ID NO: 182 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 183 (Kabat)	HCDR2	AI SN QGGKSTYYADSVK
SEQ ID NO: 184 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 185 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 186 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 187 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 188 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 189 (Chothia)	HCDR2	SN QGKS
SEQ ID NO: 190 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 191 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 192 (Chothia)	LCDR2	GAS
SEQ ID NO: 193 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 194	VL	DIQMTQSPSSLSASVGDRTVITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGQGTKVEIK
SEQ ID NO: 195	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI SN QG KSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 196	DNA VL	GATATCCAGATGACCCAGAGCCCGCAGCAGCTGAGCGCCAGCGTGGGCGACAGAG TGACCATCACCTGTCCGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGCAAGGCCCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTCACCCTGA CCATCAGCAGCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAGGTGGAATCAAG
SEQ ID NO: 197	DNA VH	GAGGTGCAATTGCTGGAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCGCCGCGCAGCGGCTTACCTTCAGCAGTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCAGCAACCAGGGC AAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACCCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC

		ACCCTGGTCACCGTCAGCTCA
SEQ ID NO: 198	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPREAKVQWKVDNALQSGNSQESVTE QDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 199	Heavy IgG1	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI SN QG KSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSC DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP AIEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYTQKS LSLSPGK
MOR10703 A50V_N52S		
SEQ ID NO: 200 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 201 (Kabat)	HCDR2	VI SSQ GKSTYYADSVK
SEQ ID NO: 202 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 203 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 204 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 205 (Kabat)	LCDR3	QQYSSFPFT
SEQ ID NO: 206 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 207 (Chothia)	HCDR2	S SSQ GKS
SEQ ID NO: 208 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 209 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 210 (Chothia)	LCDR2	GAS
SEQ ID NO: 211 (Chothia)	LCDR3	YSSFPFT
SEQ ID NO: 212	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIK
SEQ ID NO: 213	VH	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI VI SSQ KSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 214	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGCGACAGAG TGACCATCACCTGTTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGCAAGGCCCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCAGCAGATTCAGCGGCAGCGGCTCCGGCACCGACTTCACCCTGA CCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAAGGTGGAAATCAAG
SEQ ID NO: 215	DNA VH	GAGGTGCAATTGCTGGAAGCGCGGAGGCGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTTCGGCCGCGCAGCGGCTTCACCTTCAGCAGCTACGCCATGAGCTGGGT CGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTATCAGCAGCCAGGGC AAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCACCGTCAGCTCA
SEQ ID NO: 216	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPREAKVQWKVDNALQSGNSQESVTE QDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 217	Heavy IgG1	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAI VI SSQ KSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT

		GVHTFPAVLQSSGLYLSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKS LSLSPGK
MOR10703 A50V_N52G		
SEQ ID NO: 218 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 219 (Kabat)	HCDR2	<u>V</u> I <u>G</u> SQ GKSTYYADSVK G
SEQ ID NO: 220 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 221 (Kabat)	LCDR1	RASQGISNWL A
SEQ ID NO: 222 (Kabat)	LCDR2	GASSLQ S
SEQ ID NO: 223 (Kabat)	LCDR3	QQYSSFP TT
SEQ ID NO: 224 (Chothia)	HCDR1	GTFSSY
SEQ ID NO: 225 (Chothia)	HCDR2	<u>G</u> SQ GK S
SEQ ID NO: 226 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 227 (Chothia)	LCDR1	SQGISN W
SEQ ID NO: 228 (Chothia)	LCDR2	GAS
SEQ ID NO: 229 (Chothia)	LCDR3	YSSFP T
SEQ ID NO: 230	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFLTITISLQPEDFATYYCQQYSSFP TTFGQGTKVEIK
SEQ ID NO: 231	VH	EVQLLES GGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVS <u>V</u> I <u>G</u> SQ G KSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQ G TLVTVSS
SEQ ID NO: 232	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGCAGAG TGACCATCACCTGTCCGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCACAAGCAGATTCAGCGGCAGCGGCTCCGGCACCGACTCACCTGA CCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 233	DNA VH	GAGGTGCAATGTGTTGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGC CGCGCCAGCGGCTTACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTTCATCGGCAGCCAGGGC AAGAGCCTACTACCGGACAGCGTGAAGGGCCGGTTACCATCAGCCGGGACA ACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCACCGTCAGCTCA
SEQ ID NO: 234	Light Kappa	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFLTITISLQPEDFATYYCQQYSSFP TTFGQGTKVEIKRTV AAPSVFI FPPSDEQLKSGTASVCLLN NFYPREAKVQWKVDNALQSGNSQESVTE QDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 235	Heavy IgG1	EVQLLES GGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVS <u>V</u> I <u>G</u> SQ G KSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQ G TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTS GVHTFPAVLQSSGLYLSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKS LSLSPGK

MOR10703 S52aA		
SEQ ID NO: 236 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 237 (Kabat)	HCDR2	AINAQQGKSTYYADSVKG
SEQ ID NO: 238 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 239 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 240 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 241 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 242 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 243 (Chothia)	HCDR2	NAQQKS
SEQ ID NO: 244 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 245 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 246 (Chothia)	LCDR2	GAS
SEQ ID NO: 247 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 248	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSGSGTDFLTISSLQPEDFATYYCQQYSSFPSTFGQGTKVEIK
SEQ ID NO: 249	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAINAQQKSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQGLTVVSS
SEQ ID NO: 250	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGCGCACAGAGTGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCTGGTATCAGCAGAAGCCCGCAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAGAGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTCACCCCTGACCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAGCAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 251	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGAGACTGTCCTGCGCGCCAGCGGCTCACCTTCAGCAGCTACGCCATGAGCTGGGTCCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCAACGCCAGGGCAAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCAGGACACCGCCGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGCACCTGGTCACCGTCAGCTCA
SEQ ID NO: 252	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSGSGTDFLTISSLQPEDFATYYCQQYSSFPSTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSITLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 253	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAINAQQKSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQGLTVTVSSASTKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKLSLSPPGK
MOR10703 S52aT		
SEQ ID NO: 254 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 255 (Kabat)	HCDR2	AINTQQGKSTYYADSVKG
SEQ ID NO: 256 (Kabat)	HCDR3	WGDEGFDI

SEQ ID NO: 257 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 258 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 259 (Kabat)	LCDR3	QYSSFPTT
SEQ ID NO: 260 (Chothia)	HCDR1	GTFSSY
SEQ ID NO: 261 (Chothia)	HCDR2	N <u>T</u> QGKS
SEQ ID NO: 262 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 263 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 264 (Chothia)	LCDR2	GAS
SEQ ID NO: 265 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 266	VL	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSGSGTDFLTITISLQPEDFATYYCQYSSFPTTFGQGTKVEIK
SEQ ID NO: 267	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAIN <u>T</u> QGKSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQGLVTVSS
SEQ ID NO: 268	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGAGAGTGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCAGCAGAAGCCCGCAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAGAGCGGCGTGCCAAGCAGATTCAGCGGCAGCGGCTCCGGCACCGACTTACCCTGACCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAGCAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 269	DNA VH	GAGGTGCAATTGCTGGAAGCGGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGAGACTGTCTTGC CGCGCCAGCGGCTTACCTTCAGCAGCTACGCCATGAGCTGGGTCCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGCCATCAACACCCAGGGCAAGAGCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACACAGCAAGAACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCCGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGCACCTGGTCAACCGTCAGCTCA
SEQ ID NO: 270	Light Kappa	DIQMTQSPSSLSASVGRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSGSGTDFLTITISLQPEDFATYYCQYSSFPTTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPYPRKAVQWVDNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 271	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSAIN <u>T</u> QGKSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQGLVTVSSASTKGPVFPFLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSVHTFPFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKSCDKTHTCPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPLVDSGDSFFLYSKLTVDKSRWQQGNV FSCVMHEALHNHYTQKSLSLSPGK
MOR10701 R555		
SEQ ID NO: 272 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 273 (Kabat)	HCDR2	VTGAVG <u>S</u> STYYPDSVKG
SEQ ID NO: 274 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 275 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 276 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 277 (Kabat)	LCDR3	QYSSFPTT
SEQ ID NO: 278 (Chothia)	HCDR1	GTFSSY

SEQ ID NO: 279 (Chothia)	HCDR2	GAVG <u>SS</u>
SEQ ID NO: 280 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 281 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 282 (Chothia)	LCDR2	GAS
SEQ ID NO: 283 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 284	VL	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSSFPPTTFGQGTKVEIK
SEQ ID NO: 285	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG <u>S</u> STYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 286	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGGAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGCGTGCACAAGCAGATTAGCGGCAGCGGCTCCGGCACCGACTCACCTGA CCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 287	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCCTCGCCAGCGGCTTACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGACAGGCGCCGTGGGC AGCAGCACCTACTACCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACCCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCCACCGTCAGCTCA
SEQ ID NO: 288	Light Kappa	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSSFPPTTFGQGTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPYPRKAVQWVVDNALQSGNSQESVTE QDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 289	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG <u>S</u> STYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEPKSC DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKF NHWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPPLVDSGDSFFLYSKLTVDKSRWQQGNVFSQSVMHEALHNHYTQKS LSLSPGK
MOR10701 R55G		
SEQ ID NO: 290 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 291 (Kabat)	HCDR2	VTGAVG <u>SS</u> STYYPDSVKG
SEQ ID NO: 292 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 293 (Kabat)	LCDR1	RASQGISNWLA
SEQ ID NO: 294 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 295 (Kabat)	LCDR3	QQYSSFPPT
SEQ ID NO: 296 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 297 (Chothia)	HCDR2	GAVG <u>SS</u>
SEQ ID NO: 298 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 299 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 300 (Chothia)	LCDR2	GAS

SEQ ID NO: 301 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 302	VL	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSSFPPTTFGGQTKVEIK
SEQ ID NO: 303	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG <u>G</u> STYYPDSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS
SEQ ID NO: 304	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGGCAGAGAG TGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCA GCAGAAGCCCGCAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAG AGCGGCGTGCACAAGCAGATTCAGCGGCAGCGGCTCCGGCACCGACTCACCCCTGA CCATCAGCAGCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG CAGCTTCCCCACCCTTCGGCCAGGGCACCAGGTGGAAATCAAG
SEQ ID NO: 305	DNA VH	GAGGTGCAATTGCTGGAAAGCGCGGAGGCGCTGGTGCAGCCTGGCGGCAGCCTGA GACTGTCTTGCGCCCGCAGCGGCTTCACCTTCAGCAGCTACGCCATGAGCTGGGT CCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGACAGGCGCCGTGGGC GGAAGCACCTACTACCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACA ACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCAGGACACCGC CGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGC ACCTGGTCCACCGTCAGCTCA
SEQ ID NO: 306	Light Kappa	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSSFPPTTFGGQTKVEIKRTV AAPSVFI FPPSDEQLKSGTASVVCLLNFPREAKVQWKVDNALQSGNSQESVTE QDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 307	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG <u>G</u> STYYPDSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGLVLDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSC DKHTHTCPPELLEGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NHWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKS LSLSPGK
MOR10701 R55K		
SEQ ID NO: 308 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 309 (Kabat)	HCDR2	VTGAVG <u>K</u> STYYPDSVKG
SEQ ID NO: 310 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 311 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 312 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 313 (Kabat)	LCDR3	QQYSSFPPT
SEQ ID NO: 314 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 315 (Chothia)	HCDR2	GAVG <u>K</u> S
SEQ ID NO: 316 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 317 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 318 (Chothia)	LCDR2	GAS
SEQ ID NO: 319 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 320	VL	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQ SGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSSFPPTTFGGQTKVEIK
SEQ ID NO: 321	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG <u>K</u> STYYPDSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARWGDEGFDIWGQG TLVTVSS

SEQ ID NO: 322	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGCGACAGAGTGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCAGCAGAAGCCCGGCAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAGAGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTACCCTGACCATCAGCAGCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAGCAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 323	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGAGACTGTCTTGCGCCGCCAGCGGCTTACCTTACAGCAGCTACGCCATGAGCTGGGTCCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGACAGGCGCCGTGGGCAAAAGCACCTACTACCCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACAACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCCGCGTGTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGCACCTGGTCACCGTCAGCTCA
SEQ ID NO: 324	Light Kappa	DIQMTQSPSSLSASVGDVITITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQSGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSLTITLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 325	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVGKSTYYPDSVKGRFTISRDN SKNTLYLQMNLSRAEDTAVYYCARWGDEGFDIWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKSKDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHEALHNHYTQKSLSLSPGK
MOR10701 deletion S56		
SEQ ID NO: 326 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 327 (Kabat)	HCDR2	VTGAVGRTYYPDSVKG
SEQ ID NO: 328 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 329 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 330 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 331 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 332 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 333 (Chothia)	HCDR2	GAVGRT
SEQ ID NO: 334 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 335 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 336 (Chothia)	LCDR2	GAS
SEQ ID NO: 337 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 338	VL	DIQMTQSPSSLSASVGDVITITCRASQGISNWLAWYQQKPKAPKLLIYGASSLQSGVPSRFRSGSGSDFTLTITSSLPEDFATYYCQQYSSFPFTTFGGQTKVEIK
SEQ ID NO: 339	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSVTGAVGRTYYPDSVKGRFTISRDN SKNTLYLQMNLSRAEDTAVYYCARWGDEGFDIWGQGLVTVSS
SEQ ID NO: 340	DNA VL	GATATCCAGATGACCCAGAGCCCCAGCAGCCTGAGCGCCAGCGTGGGCGACAGAGTGACCATCACCTGTCGGGCCAGCCAGGGCATCAGCAACTGGCTGGCCTGGTATCAGCAGAAGCCCGGCAAGGCCCAAGCTGCTGATCTACGGCGCCAGCTCCCTGCAGAGCGGCGTGCCAAGCAGATTACAGCGGCAGCGGCTCCGGCACCGACTTACCCTGACCATCAGCAGCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAGCAGCTTCCCCACCACCTTCGGCCAGGGCACCAAGGTGGAATCAAG
SEQ ID NO: 341	DNA VH	GAGGTGCAATTGCTGGAAAGCGGCGGAGGCTGGTGCAGCCTGGCGGCAGCCTGAGACTGTCTTGCGCCGCCAGCGGCTTACCTTACAGCAGCTACGCCATGAGCTGGGTCCGCCAGGCCCTGGCAAGGGACTGGAATGGGTGTCCGTGACAGGCGCCGTGGGCAAAAGCACCTACTACCCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACAACAACAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCCGT

		GTACTACTGTGCCAGATGGGGCGACGAGGGCTTCGACATCTGGGGCCAGGGCACC CTGGTCACCGTCAGCTCA
SEQ ID NO: 342	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKGKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFAVYQCQQYSSFPFTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPYPREAKVQWKVDNALQSGNSQESVTE QDSKDYSLSSLTITLSKADYKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 343	Heavy IgG1	EVQLLESGGGLVQPGGSLRLSCLASGFTFSSYAMSWVRQAPGKLEWVSVTGAVG RTYYPDSVKGRFTISRDNKNTLYLQMNLSRAEDTAVYFCARWGDEGFDIWGQGT LTVVSSASTKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVTPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSCD KTHHTCPPCPAPELGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAP IEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTFPVLDSDGSGFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSL SLSPGK
MOR12609		
SEQ ID NO: 344 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 345 (Kabat)	HCDR2	VINGLGYTTFYADSVKGG
SEQ ID NO: 346 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 347 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 348 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 349 (Kabat)	LCDR3	QQYSSFPFTT
SEQ ID NO: 350 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 351 (Chothia)	HCDR2	NGLGYT
SEQ ID NO: 352 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 353 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 354 (Chothia)	LCDR2	GAS
SEQ ID NO: 355 (Chothia)	LCDR3	YSSFPFT
SEQ ID NO: 356	VL	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKGKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFAVYQCQQYSSFPFTTFGGQTKVEIK
SEQ ID NO: 357	VH	QVQLVESGGGLVQPGGSLRLSCLASGFTFSSYAMSWVRQAPGKLEWVSVINGLG YTFYADSVKGRFTISRDNKNTLYLQMNLSRAEDTAVYFCARWGDEGFDIWGQGT TLVTVSS
SEQ ID NO: 358	DNA VL	GATATCCAGATGACCCAGAGCCCGCTAGCCTGAGCGCAGCGTGGGTGATCGTG TGACCATACCTGCAGAGCGAGCCAGGGTATTTCTAATTGGCTGGCTTGGTACCA GCAGAAACCAGGTAAAGCACCGAACTATTAATTTATGGTGTCTTCTTTGCAA AGCGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTTATGCCAGCAGTATTC TTCTTTTCTACTACCTTTGGCCAGGGTACGAAAGTTGAAATTTAA
SEQ ID NO: 359	DNA VH	CAGGTGCAATTTGGTGGAAAGCGCGCGCGCTGGTGCAACCGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCTGGGAAGGGTCTCGAGTGGGTGAGCGTTATTTAATGGTCTTGGT TATACTACTTTTTATGCTGATTCTGTTAAGGGTCGTTTTACCATTTACGATGATA ATTCGAAAAACACCCTGTATCTGCAAAATGAACAGCCTGCGTGCGGAAGATACGGC CGTGATTTATGCGCGCGTTGGGGTGTAGGGTTTTGATATTTGGGGCCAAGGC ACCTGGTGACGGTTAGCTCA
SEQ ID NO: 360	Light Kappa	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPKGKAPKLLIYGASSLQ SGVPSRFRSGSGSGTDFTLTITSSLPEDFAVYQCQQYSSFPFTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVVCLLNFPYPREAKVQWKVDNALQSGNSQESVTE QDSKDYSLSSLTITLSKADYKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 361	Heavy IgG1	QVQLVESGGGLVQPGGSLRLSCLASGFTFSSYAMSWVRQAPGKLEWVSVINGLG YTFYADSVKGRFTISRDNKNTLYLQMNLSRAEDTAVYFCARWGDEGFDIWGQGT TLVTVSSASTKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG

		GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC DKHTTCPPELGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYTQKS LSLSPGK
MOR12610		
SEQ ID NO: 362 (Kabat)	HCDR1	SYAMS
SEQ ID NO: 363 (Kabat)	HCDR2	GTGPGYGGTYYPDSVKG
SEQ ID NO: 364 (Kabat)	HCDR3	WGDEGFDI
SEQ ID NO: 365 (Kabat)	LCDR1	RASQGISNWL
SEQ ID NO: 366 (Kabat)	LCDR2	GASSLQS
SEQ ID NO: 367 (Kabat)	LCDR3	QQYSSFPTT
SEQ ID NO: 368 (Chothia)	HCDR1	GFTFSSY
SEQ ID NO: 369 (Chothia)	HCDR2	GPYGG
SEQ ID NO: 370 (Chothia)	HCDR3	WGDEGFDI
SEQ ID NO: 371 (Chothia)	LCDR1	SQGISNW
SEQ ID NO: 372 (Chothia)	LCDR2	GAS
SEQ ID NO: 373 (Chothia)	LCDR3	YSSFPT
SEQ ID NO: 374	VL	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTISLQPEDFAVYQCQQYSSFPTTFGGQTKVEIK
SEQ ID NO: 375	VH	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSGTGPY GTYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYVCARWDEGFDIWGQGT LTVVSS
SEQ ID NO: 376	DNA VL	GATATCCAGATGACCCAGAGCCCCTAGCCTGAGCGCAGCGTGGGTGATCGTG TGACCATACCTGCAGAGCGAGCCAGGGTATTTCTAATTTGGCTGGCTGGTACCA GCAGAAACCAGGTAAAGCACCGAACTATTAATTTATGGTGTCTCTCTTTGCAA AGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCTGA CCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTATTGCCAGCAGTATTC TTCTTTTCTACTACCTTTGGCCAGGGTACGAAAGTTGAAATTTAAA
SEQ ID NO: 377	DNA VH	CAGGTGCAATTTGGTGGAAAGCGCGCGGCGCTGGTGCACCGGGCGGCAGCCTGC GTCTGAGCTGCGCGGCCCTCCGGATTTACCTTTAGCAGCTATGCGATGAGCTGGGT GCGCCAAGCCCCCTGGGAAGGGTCTCGAGTGGGTGAGCGGTACTGGTCTTATGGT GGTACTTATATCTGATCTGTTAAGGGTCTTTTACCATTTACAGTGATAATTT CGAAAAACACCCTGTATCTGCAATGAACAGCCTGCGTGCAGGAAAGATAACGGCCGT GTATTATGCGCGCGTTGGGGTGTGAGGGTTTTGATATTTGGGGCCAAGGCACC CTGGTGACGGTTAGCTCA
SEQ ID NO: 378	Light Kappa	DIQMTQSPSSLSASVGRVITTCRASQGISNWLAWYQQKPKAPKLLIYGASSLQ SGVPSRFRSGSGSDFTLTISLQPEDFAVYQCQQYSSFPTTFGGQTKVEIKRTV AAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTE QDSKDYSLSSLTLSKADYKHKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 379	Heavy IgG1	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSGTGPY GTYYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYVCARWDEGFDIWGQGT LTVVSSASTKGPSVFLPAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCD KHTTCPPELGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKF WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAP IEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYTQKSL LSLSPGK

Other antibodies of the invention include those where the amino acids or nucleic acids encoding the amino acids have been mutated, yet have at least 60, 70, 80, 90, 95, 96, 97, 98, and 99 percent identity to the sequences described in Table 1. In some embodiments, it include mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated in the variable regions when compared with the variable regions depicted in the sequence described in Table 1, while retaining substantially the same therapeutic activity.

Since each of these antibodies or fragments thereof can bind to HER3, the VH, VL, full length light chain, and full length heavy chain sequences (amino acid sequences and the nucleotide sequences encoding the amino acid sequences) can be "mixed and matched" to create other HER3-binding antibodies of the invention. Such "mixed and matched" HER3-binding antibodies can be tested using the binding assays known in the art (*e.g.*, ELISAs, and other assays described in the Example section). When these chains are mixed and matched, a VH sequence from a particular VH/VL pairing should be replaced with a structurally similar VH sequence. Likewise a full length heavy chain sequence from a particular full length heavy chain / full length light chain pairing should be replaced with a structurally similar full length heavy chain sequence. Likewise, a VL sequence from a particular VH/VL pairing should be replaced with a structurally similar VL sequence. Likewise a full length light chain sequence from a particular full length heavy chain / full length light chain pairing should be replaced with a structurally similar full length light chain sequence. Accordingly, in one aspect, the invention provides an isolated monoclonal antibody or fragment thereof having: a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 15, 33, 51, 69, 87, 105, 123, 141, 159, 177, 195, 213, 231, 249, 267, 285, 303, 321, 339, 357, and 375; and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 14, 32, 50, 68, 86, 104, 122, 140, 158, 176, 194, 212, 230, 248, 266, 284, 302, 320, 338, 356, and 374; wherein the antibody specifically binds to HER3 (*e.g.*, human and/or cynomologus).

In another aspect, the present invention provides HER3-binding antibodies that comprise the heavy chain and light chain CDR1s, CDR2s and CDR3s as described in Table 1, or combinations thereof. The amino acid sequences of the VH CDR1s of the antibodies are shown in SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296, 308, 314, 326, 332, 344, 350, 362, and 368. The amino acid sequences of the VH CDR2s of the antibodies and are shown in SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111,

117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369. The amino acid sequences of the VH CDR3s of the antibodies are shown in SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 76, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370. The amino acid sequences of the VL CDR1s of the antibodies are shown in SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137, 149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371. The amino acid sequences of the VL CDR2s of the antibodies are shown in SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372. The amino acid sequences of the VL CDR3s of the antibodies are shown in SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 133, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373. The CDR regions are delineated using the Kabat system (Kabat *et al.*, (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Chothia *et al.*, (1987) J. Mol. Biol. 196:901-917; Chothia *et al.*, (1989) Nature 342: 877-883; and Al-Lazikani *et al.*, (1997) J. Mol. Biol. 273, 927-948).

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 2; a CDR2 of SEQ ID NO: 3; a CDR3 of SEQ ID NO: 4; a light chain variable region CDR1 of SEQ ID NO: 5; a CDR2 of SEQ ID NO: 6; and a CDR3 of SEQ ID NO: 7.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 20; a CDR2 of SEQ ID NO: 21; a CDR3 of SEQ ID NO: 22; a light chain variable region CDR1 of SEQ ID NO: 23; a CDR2 of SEQ ID NO: 24; and a CDR3 of SEQ ID NO: 25.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 38; a CDR2 of SEQ ID NO: 39; a CDR3 of SEQ ID NO: 40; a light chain variable region CDR1 of SEQ ID NO: 41; a CDR2 of SEQ ID NO: 42; and a CDR3 of SEQ ID NO: 43.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 56; a CDR2 of SEQ ID NO: 57; a CDR3 of SEQ ID NO: 58; a light chain variable region CDR1 of SEQ ID NO: 59; a CDR2 of SEQ ID NO: 60; and a CDR3 of SEQ ID NO: 61.

- 5 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 74; a CDR2 of SEQ ID NO: 75; a CDR3 of SEQ ID NO: 76; a light chain variable region CDR1 of SEQ ID NO: 77; a CDR2 of SEQ ID NO: 78; and a CDR3 of SEQ ID NO: 79.

- 10 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 92; a CDR2 of SEQ ID NO: 93; a CDR3 of SEQ ID NO: 94; a light chain variable region CDR1 of SEQ ID NO: 95; a CDR2 of SEQ ID NO: 96; and a CDR3 of SEQ ID NO: 97.

- 15 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 110; a CDR2 of SEQ ID NO: 111; a CDR3 of SEQ ID NO: 112; a light chain variable region CDR1 of SEQ ID NO: 113; a CDR2 of SEQ ID NO: 114; and a CDR3 of SEQ ID NO: 115.

- 20 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 128; a CDR2 of SEQ ID NO: 129; a CDR3 of SEQ ID NO: 130; a light chain variable region CDR1 of SEQ ID NO: 131; a CDR2 of SEQ ID NO: 132; and a CDR3 of SEQ ID NO: 133.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 146; a CDR2 of SEQ ID NO: 147; a CDR3 of SEQ ID NO: 148; a light chain variable region CDR1 of SEQ ID NO: 149; a CDR2 of SEQ ID NO: 150; and a CDR3 of SEQ ID NO: 151.

- 25 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 164; a CDR2 of SEQ ID NO: 165; a CDR3 of SEQ ID NO: 166; a light chain variable region CDR1 of SEQ ID NO: 167; a CDR2 of SEQ ID NO: 168; and a CDR3 of SEQ ID NO: 169.

- 30 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 182; a CDR2 of SEQ ID NO: 183; a CDR3 of SEQ ID NO:

184; a light chain variable region CDR1 of SEQ ID NO: 185; a CDR2 of SEQ ID NO: 186; and a CDR3 of SEQ ID NO: 187.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 200; a CDR2 of SEQ ID NO: 201; a CDR3 of SEQ ID NO: 202; a light chain variable region CDR1 of SEQ ID NO: 203; a CDR2 of SEQ ID NO: 204; and a CDR3 of SEQ ID NO: 205.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 218; a CDR2 of SEQ ID NO: 219; a CDR3 of SEQ ID NO: 220; a light chain variable region CDR1 of SEQ ID NO: 221; a CDR2 of SEQ ID NO: 222; and a CDR3 of SEQ ID NO: 223.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 236; a CDR2 of SEQ ID NO: 237; a CDR3 of SEQ ID NO: 238; a light chain variable region CDR1 of SEQ ID NO: 239; a CDR2 of SEQ ID NO: 240; and a CDR3 of SEQ ID NO: 241.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 254; a CDR2 of SEQ ID NO: 255; a CDR3 of SEQ ID NO: 256; a light chain variable region CDR1 of SEQ ID NO: 257; a CDR2 of SEQ ID NO: 258; and a CDR3 of SEQ ID NO: 259.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 272; a CDR2 of SEQ ID NO: 273; a CDR3 of SEQ ID NO: 274; a light chain variable region CDR1 of SEQ ID NO: 275; a CDR2 of SEQ ID NO: 276; and a CDR3 of SEQ ID NO: 277.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 290; a CDR2 of SEQ ID NO: 291; a CDR3 of SEQ ID NO: 292; a light chain variable region CDR1 of SEQ ID NO: 293; a CDR2 of SEQ ID NO: 294; and a CDR3 of SEQ ID NO: 295.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 308; a CDR2 of SEQ ID NO: 309; a CDR3 of SEQ ID NO: 310; a light chain variable region CDR1 of SEQ ID NO: 311; a CDR2 of SEQ ID NO: 312; and a CDR3 of SEQ ID NO: 313.

In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 326; a CDR2 of SEQ ID NO: 327; a CDR3 of SEQ ID NO: 328; a light chain variable region CDR1 of SEQ ID NO: 329; a CDR2 of SEQ ID NO: 330; and a CDR3 of SEQ ID NO: 331.

- 5 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 344; a CDR2 of SEQ ID NO: 345; a CDR3 of SEQ ID NO: 346; a light chain variable region CDR1 of SEQ ID NO: 347; a CDR2 of SEQ ID NO: 348; and a CDR3 of SEQ ID NO: 349.

10 In a specific embodiment, an antibody that binds to HER3 comprises a heavy chain variable region CDR1 of SEQ ID NO: 362; a CDR2 of SEQ ID NO: 363; a CDR3 of SEQ ID NO: 364; a light chain variable region CDR1 of SEQ ID NO: 365; a CDR2 of SEQ ID NO: 366; and a CDR3 of SEQ ID NO: 367.

In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO. 15 and VL of SEQ ID NO: 14. In a specific embodiment, an antibody that binds to HER3
15 comprises a VH of SEQ ID NO: 33 and VL of SEQ ID NO: 32. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 51 and VL of SEQ ID NO: 50. In a specific embodiment, an antibody that binds to HER3 comprises a SEQ ID NO: 69 and VL of SEQ ID NO: 68. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 87 and VL of SEQ ID NO: 86. In a specific embodiment, an antibody
20 that binds to HER3 comprises a VH of SEQ ID NO: 105 and VL of SEQ ID NO: 104. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 123 and VL of SEQ ID NO: 122. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 141 and VL of SEQ ID NO: 140. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 159 and VL of SEQ ID NO:
25 158. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 177 and VL of SEQ ID NO: 176. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 195 and VL of SEQ ID NO: 194. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 213 and VL of SEQ ID NO: 212. In a specific embodiment, an antibody that binds to HER3 comprises a VH
30 of SEQ ID NO: 231 and VL of SEQ ID NO: 230. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 249 and VL of SEQ ID NO: 248. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 267 and VL of SEQ ID NO: 266. In a specific embodiment, an antibody that binds to HER3 comprises a VH

of SEQ ID NO: 285 and VL of SEQ ID NO: 284. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 303 and VL of SEQ ID NO: 302. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 321 and VL of SEQ ID NO: 320. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 339 and VL of SEQ ID NO: 338. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 357 and VL of SEQ ID NO: 356. In a specific embodiment, an antibody that binds to HER3 comprises a VH of SEQ ID NO: 375 and VL of SEQ ID NO: 374. In one embodiment, the HER3 antibodies are antagonist antibodies. In certain embodiments, an antibody that binds to HER3 is an antibody that is described in Table 1.

As used herein, a human antibody comprises heavy or light chain variable regions or full length heavy or light chains that are "the product of" or "derived from" a particular germline sequence if the variable regions or full length chains of the antibody are obtained from a system that uses human germline immunoglobulin genes. Such systems include immunizing a transgenic mouse carrying human immunoglobulin genes with the antigen of interest or screening a human immunoglobulin gene library displayed on phage with the antigen of interest. A human antibody that is "the product of" or "derived from" a human germline immunoglobulin sequence can be identified as such by comparing the amino acid sequence of the human antibody to the amino acid sequences of human germline immunoglobulins and selecting the human germline immunoglobulin sequence that is closest in sequence (*i.e.*, greatest % identity) to the sequence of the human antibody. A human antibody that is "the product of" or "derived from" a particular human germline immunoglobulin sequence may contain amino acid differences as compared to the germline sequence, due to, for example, naturally occurring somatic mutations or intentional introduction of site-directed mutations. However, in the VH or VL framework regions, a selected human antibody typically is at least 90% identical in amino acids sequence to an amino acid sequence encoded by a human germline immunoglobulin gene and contains amino acid residues that identify the human antibody as being human when compared to the germline immunoglobulin amino acid sequences of other species (*e.g.*, murine germline sequences). In certain cases, a human antibody may be at least 60%, 70%, 80%, 90%, or at least 95%, or even at least 96%, 97%, 98%, or 99% identical in amino acid sequence to the amino acid sequence encoded by the germline immunoglobulin gene. Typically, a recombinant human antibody will display no more than 10 amino acid differences from the amino acid sequence encoded by the human germline immunoglobulin gene in the VH or VL framework regions. In certain cases, the

human antibody may display no more than 5, or even no more than 4, 3, 2, or 1 amino acid difference from the amino acid sequence encoded by the germline immunoglobulin gene. Different germlined versions using the VH and VL germline sequences for a representative number of HER3 antibodies is shown in Table 2, using Kabat. The CDR positions are highlighted in boldface. The notation used in the Tables with germlined sequences is as follows: MOR10701-VH_3-07 means MOR10701 CDR loops in framework regions of VH germline sequence 3-07 (nomenclature is according to Vbase), MOR10703-VK_L1 means CDR from MOR10703 in germline framework regions from VK_L1, where VK is the kappa light chain.

10 Table 2: Different germlined versions of a selected number of representative antibodies

SEQ ID NUMBER	Sequence Name	Amino Acid Sequence
	MOR10701 VH domain	
SEQ ID NO: 380	MOR10701-VH_3-07	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVAVTGAVGRST YYPD SVKGRFTISRDN AKNSLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 381	MOR10701-VH_3-09	EVQLVESGGGLVQPGRSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISRDN AKNSLYLQMN SLRAEDTALYYCAK WGDEGF DI
SEQ ID NO: 382	MOR10701-VH_3-11	QVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WIRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISRDN AKNSLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 383	MOR10701-VH_3-13	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQATGKGLEWVSVTGAVGRST YYPDS VKGRFTISR ENAKNSLYLQMN SLRAGDTAVYYCARWGDEGF DI
SEQ ID NO: 384	MOR10701-VH_3-15	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVVGVTGAVGRST YYPDS VKGRFTISR DDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGF DI
SEQ ID NO: 385	MOR10701-VH_3-20	EVQLVESGGGVVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISRDN AKNSLYLQMN SLRAEDTALYHCARWGDEGF DI
SEQ ID NO: 386	MOR10701-VH_3-21	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISRDN AKNSLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 387	MOR10701-VH_3-23	EVQLLESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCAK WGDEGF DI
SEQ ID NO: 388	MOR10701-VH_3-30	QVQLVESGGGVVQPGRSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVAVTGAVGRST YYPD SVKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCAK WGDEGF DI
SEQ ID NO: 389	MOR10701-VH_3-30.3	QVQLVESGGGVVQPGRSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVAVTGAVGRST YYPD SVKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 390	MOR10701-VH_3-30.5	QVQLVESGGGVVQPGRSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVAVTGAVGRST YYPD SVKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCAK WGDEGF DI
SEQ ID NO: 391	MOR10701-VH_3-33	QVQLVESGGGVVQPGRSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVAVTGAVGRST YYPD SVKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 392	MOR10701-VH_3-43	EVQLVESGGVVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNSLYLQMN SLRTEDTALYYCAK WGDEGF DI
SEQ ID NO: 393	MOR10701-VH_3-48	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISRDN AKNSLYLQMN SLRDEDTAVYYCARWGDEGF DI
SEQ ID NO: 394	MOR10701-VH_3-49	EVQLVESGGGLVQPGRSLRLSCTASGFTFSS YAMS WFRQAPGKGLEWVVGVTGAVGRST YYPDS VKGRFTISR DGSKSIAYLQMN SLKTEDTAVYYCTR WGDEGF DI
SEQ ID NO: 395	MOR10701-VH_3-53	EVQLVETGGGLIQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 396	MOR10701-VH_3-64	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNTLYLQMN SLRAEDMAVYYCARWGDEGF DI
SEQ ID NO: 397	MOR10701-VH_3-66	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNTLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 398	MOR10701-VH_3-72	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVVGVTGAVGRST YYPD SVKGRFTISR DDSKNSLYLQMN SLKTEDTAVYYCARWGDEGF DI
SEQ ID NO: 399	MOR10701-VH_3-73	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQASGKGLEWVVGVTGAVGRST YYPDS VKGRFTISR DDSKNTAYLQMN SLKTEDTAVYYCTR WGDEGF DI
SEQ ID NO: 400	MOR10701-VH_3-74	EVQLVESGGGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLVWVSVTGAVGRST YYPDS VKGRFTISR DNKNTLYLQMN SLRAEDTAVYYCARWGDEGF DI
SEQ ID NO: 401	MOR10701-VH_3-d	EVQLVESRGLVQPGGSLRLSCAASGFTFSS YAMS WVRQAPGKGLEWVSVTGAVGRST YYPDS VKGRFTISR DNSKNTLHLQMN SLRAEDTAVYYCK WGDEGF DI
	MOR10703 VH domain	

SEQ ID NO: 402	MOR10703-VH_3-07	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 403	MOR10703-VH_3-09	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAEDTALYYCAKWGDEGFDI
SEQ ID NO: 404	MOR10703-VH_3-11	QVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 405	MOR10703-VH_3-13	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQATGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAGDTAVYYCARWGDEGFDI
SEQ ID NO: 406	MOR10703-VH_3-15	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVGAAINSQQGKSTYYADS VKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTWGDEGFDI
SEQ ID NO: 407	MOR10703-VH_3-20	EVQLVESGGGVVVRPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAEDTALYHRCARWGDEGFDI
SEQ ID NO: 408	MOR10703-VH_3-21	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 409	MOR10703-VH_3-23	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKWGDEGFDI
SEQ ID NO: 410	MOR10703-VH_3-30	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYAD SVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKWGDEGFDI
SEQ ID NO: 411	MOR10703-VH_3-30.3	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYAD SVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 412	MOR10703-VH_3-30.5	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYAD SVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKWGDEGFDI
SEQ ID NO: 413	MOR10703-VH_3-33	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYAD SVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 414	MOR10703-VH_3-43	EVQLVESGGVVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNKNSLYLQMNSLRTEDEALYYCAKWGDEGFDI
SEQ ID NO: 415	MOR10703-VH_3-48	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNAKNSLYLQMNSLRDEDTAVYYCARWGDEGFDI
SEQ ID NO: 416	MOR10703-VH_3-49	EVQLVESGGGLVQPGGSLRLSCTASGFTFSSYAMSWVRQAPGKGLEWVGAAINSQQGKSTYYADS VKGRFTISRDSKSIAYLQMNSLKTEDTAVYYCTRWGDEGFDI
SEQ ID NO: 417	MOR10703-VH_3-53	EVQLVETGGGLIQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 418	MOR10703-VH_3-64	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEYVVAAINSQQGKSTYYADSV KGRFTISRDNKNTLYLQMGSLRAEDMAVYYCARWGDEGFDI
SEQ ID NO: 419	MOR10703-VH_3-66	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 420	MOR10703-VH_3-72	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVGAAINSQQGKSTYYADS VKGRFTISRDDSKNSLYLQMNSLKTEDTAVYYCARWGDEGFDI
SEQ ID NO: 421	MOR10703-VH_3-73	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQASGKGLEWVGAAINSQQGKSTYYADS VKGRFTISRDDSKNTAYLQMNSLKTEDTAVYYCTRWGDEGFDI
SEQ ID NO: 422	MOR10703-VH_3-74	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLVWVAAINSQQGKSTYYADS VKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARWGDEGFDI
SEQ ID NO: 423	MOR10703-VH_3-d	EVQLVESRGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAAINSQQGKSTYYADS VKGRFTISRDNKNTLHLQMNSLRAEDTAVYYCKKWGDEGFDI
	MOR10701 VK domain	
SEQ ID NO: 424	MOR10701-VKI_O12 (same as MOR10701 wt)	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 425	MOR10701-VKI_O2	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 426	MOR10701-VKI_O18	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GGSDFTFTISLQPEDATYYCQQYSSFPPT
SEQ ID NO: 427	MOR10701-VKI_O8	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GGSDFTFTISLQPEDATYYCQQYSSFPPT
SEQ ID NO: 428	MOR10701-VKI_A20	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKVPKLLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDVATYYCQQYSSFPPT
SEQ ID NO: 429	MOR10701-VKI_A30	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKRLIYGASSLQSGVPSRFSGS GGSEFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 430	MOR10701-VKI_L14	NIQMTQSPSAMSASVGDRTITCRASQGISNWLAWFQQKPGKVPKHLIYGASSLQSGVPSRFSGS GSGSEFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 431	MOR10701-VKI_L1	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWFQQKPGKAPKSLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 432	MOR10701-VKI_L15	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPEKAPKSLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 433	MOR10701-VKI_L4	AIQLTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSG SGDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 434	MOR10701-VKI_L18	AIQLTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSG SGDFTLTISLQPEDFATYYCQQYSSFPPT
SEQ ID NO: 435	MOR10701-VKI_L5	DIQMTQSPSSVSASVGDRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GGSDFTLTISLQPEDFATYYCQQYSSFPPT

SEQ ID NO: 436	MOR10701-VKI_L19	DIQMTQSPSSVSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 437	MOR10701-VKI_L8	DIQLTQSPSFLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTEFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 438	MOR10701-VKI_L23	AIRMTQSPFSLASVGDVRTITCRASQGISNWLAWYQQKPAKAPKLLIYGASSLQSGVPSRFSGS GSGTDYTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 439	MOR10701-VKI_L9	AIRMTQSPSSFASTGDRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISCLQSEDFATYYCQQYSSFPTT
SEQ ID NO: 440	MOR10701-VKI_L24	VIWMTQSPSLLSASTGDRVTISCRASQGISNWLAWYQQKPGKAPPELLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQSEDFATYYCQQYSSFPTT
SEQ ID NO: 441	MOR10701-VKI_L11	AIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 442	MOR10701-VKI_L12	DIQMTQSPSTLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTEFTLTISLQPDDEFATYYCQQYSSFPTT
	MOR10701 VK domain	
SEQ ID NO: 443	MOR10703-VKI_O12 (same as MOR10703 wt)	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 444	MOR10703-VKI_O2	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 445	MOR10703-VKI_O18	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTTISLQPEDIATYYCQQYSSFPTT
SEQ ID NO: 446	MOR10703-VKI_O8	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTTISLQPEDIATYYCQQYSSFPTT
SEQ ID NO: 447	MOR10703-VKI_A20	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKVPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDVATYYCQQYSSFPTT
SEQ ID NO: 448	MOR10703-VKI_A30	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKRLIYGASSLQSGVPSRFSGS GSGTEFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 449	MOR10703-VKI_L14	NIQMTQSPSAMSASVGDVRTITCRASQGISNWLAWFQQKPGKVPKHLIYGASSLQSGVPSRFSGS GSGSDFEFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 450	MOR10703-VKI_L1	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWFQQKPGKAPKSLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 451	MOR10703-VKI_L15	DIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPEKAPKSLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 452	MOR10703-VKI_L4	AIQLTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS SGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 453	MOR10703-VKI_L18	AIQLTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS SGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 454	MOR10703-VKI_L5	DIQMTQSPSSVSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 455	MOR10703-VKI_L19	DIQMTQSPSSVSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 456	MOR10703-VKI_L8	DIQLTQSPSFLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTEFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 457	MOR10703-VKI_L23	AIRMTQSPFSLASVGDVRTITCRASQGISNWLAWYQQKPAKAPKLLIYGASSLQSGVPSRFSGS GSGTDYTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 458	MOR10703-VKI_L9	AIRMTQSPSSFASTGDRVTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISCLQSEDFATYYCQQYSSFPTT
SEQ ID NO: 459	MOR10703-VKI_L24	VIWMTQSPSLLSASTGDRVTISCRASQGISNWLAWYQQKPGKAPPELLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQSEDFATYYCQQYSSFPTT
SEQ ID NO: 460	MOR10703-VKI_L11	AIQMTQSPSSLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQYSSFPTT
SEQ ID NO: 461	MOR10703-VKI_L12	DIQMTQSPSTLSASVGDVRTITCRASQGISNWLAWYQQKPGKAPKLLIYGASSLQSGVPSRFSGS GSGTEFTLTISLQPDDEFATYYCQQYSSFPTT

Table 3: JH segments

SEQ ID NO: 462	JH1	WGQGTLVTVSS
SEQ ID NO: 463	JH2	WGRGTLTVSS
SEQ ID NO: 464	JH3	WGQGTMTVSS
SEQ ID NO: 465	JH4	WGQGTLVTVSS
SEQ ID NO: 466	JH5	WGQGTLVTVSS
SEQ ID NO: 467	JH6	WGQGTTVTVSS

Table 4: JK segments

SEQ ID NO: 468	JK1	FGQGTKVEIK
SEQ ID NO: 469	JK2	FGQGTKLEIK
SEQ ID NO: 470	JK3	FGPGTKVDIK
SEQ ID NO: 471	JK4	FGGGTKVEIK
SEQ ID NO: 472	JK5	FGQGTRLEIK

Any combination of the VH-germlined sequences with a JH segments can be used.

Representative examples of combinations are shown in Table 5.

5 Table 5: Representative examples of combinations of the VH-germlined sequences with a JH segments.

SEQ ID NO: 473	MOR10701-VH_3-15_JH1	EVQLVESGGGLV K PGGSLRLS CAASGFTFSSYAMS WVRQAPGKGLEWVGV TGAVGRST YYPDSVKGR FTISRDDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGFDI WGQGT LVTVSS
SEQ ID NO: 474	MOR10701-VH_3-15_JH3	EVQLVESGGGLV K PGGSLRLS CAASGFTFSSYAMS WVRQAPGKGLEWVGV TGAVGRST YYPDSVKGR FTISRDDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGFDI WGQGT MVTVSS
SEQ ID NO: 475	MOR10703-VH_3-15_JH1	EVQLVESGGGLV K PGGSLRLS CAASGFTFSSYAMS WVRQAPGKGLEWV GAINS QGKSTY YADSVKGR FTISRDDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGFDI WGQGT LVTVSS
SEQ ID NO: 476	MOR10703-VH_3-15_JH3	EVQLVESGGGLV K PGGSLRLS CAASGFTFSSYAMS WVRQAPGKGLEWV GAINS QGKSTY YADSVKGR FTISRDDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGFDI WGQGT MVTVSS

10 Any combination of the VL-germlined sequences with a JK segments can be used.

Representative examples of combinations are shown in Table 6.

15 Table 6: Representative examples of combinations of the VK-germlined sequences with a JK segments

SEQ ID NO: 477	MOR10701-VKI_O2_JK1	DIQMTQSPSSLSASVGD RVTITCRASQGISN WLAWYQQKPGKAP KLLIYGASSLQ SGVPSR FSGSGSGTDFLT ISSLPEDFATYYCCQQYSSFP TTFGQGTKVEIK
SEQ ID NO: 478	MOR10701-VKI_O2_JK4	DIQMTQSPSSLSASVGD RVTITCRASQGISN WLAWYQQKPGKAP KLLIYGASSLQ SGVPSR FSGSGSGTDFLT ISSLPEDFATYYCCQQYSSFP TTFGGGTKVEIK
SEQ ID NO: 479	MOR10703-VKI_A20_JK4	DIQMTQSPSSLSASVGD RVTITCRASQGISN WLAWYQQKPGK V P KLLIYGASSLQ SGVPSR FSGSGSGTDFLT ISSLPEDVATYYCCQQYSSFP TTFGGGTKVEIK
SEQ ID NO: 480	MOR10703-VKI_A20_JK1	DIQMTQSPSSLSASVGD RVTITCRASQGISN WLAWYQQKPGK V P KLLIYGASSLQ SGVPSR FSGSGSGTDFLT ISSLPEDVATYYCCQQYSSFP TTFGQGTKVEIK

20 Once VH has been combined with JH and VK with JK, then any combination of VH or JH with VK or JK, can be used. In one embodiment, any of the VH germlined regions can be combined with any of the VK (VL) germlined regions for each antibody. A representative number of examples of combinations is shown in Table 7.

25 Table 7: Representative examples of combinations of germlined sequences

	Combination 1	
SEQ ID NO: 481	MOR10701-VH 3-15_JH3	EVQLVESGGGLV K PGGSLRLS CAASGFTFSSYAMS WVRQAPGKGLEWVGV TGAVGRSTY YPDSVKGR FTISRDDSKNTLYLQMN SLKTEDTAVYYCTT WGDEGFDI WGQGT MVTVSS

SEQ ID NO: 482	MOR10701-VKI_A30_JK4	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKAPKRLIYGASSLQSGVPSR FSGSGSGTEFTLTISSLQPEDFATYYCQQYSSFPITTFGGGTKVEIK
	Combination 2	
SEQ ID NO: 483	MOR10701-VH_3-30_JH1	QVQLVESGGGVVQPGRSRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAVTGAVGRST YYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAK WGDEGFDI WGQGLTVTVSS
SEQ ID NO: 484	MOR10701-VKI_L1_JK2	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWFQQKPGKAPKSLIYGASSLQSGVPSR FSGSGSGTDFLTISLQPEDFATYYCQQYSSFPITTFGGGQTKLEIK
	Combination 3	
SEQ ID NO: 485	MOR10701-VH_3-30_JH2	QVQLVESGGGVVQPGRSRLSCAASGFTFSSYAMSWVRQAPGKGLEWVAVTGAVGRST YYPDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAK WGDEGFDI WGRGLTVTVSS
SEQ ID NO: 486	MOR10701-VKI_L1_JK2	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWFQQKPGKAPKSLIYGASSLQSGVPSR FSGSGSGTDFLTISLQPEDFATYYCQQYSSFPITTFGGGQTKLEIK
	Combination 4	
SEQ ID NO: 487	MOR10703-VH_3-20_JH5	EVQLVESGGGVVVRPGGSLRSLCAASGFTFSSYAMSWVRQAPGKGLEWVA AINSQGKSTY YADSVKGRFTISRDNKNTLYLQMNSLRAEDTALYHCAR WGDEGFDI WGQGLTVTVSS
SEQ ID NO: 488	MOR10703-VKI_L15_JK3	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPEKAPKSLIYGASSLQSGVPSR FSGSGSGTDFLTISLQPEDFATYYCQQYSSFPITTFGGPGTKVDIK
	Combination 5	
SEQ ID NO: 489	MOR10703-VH_3-33_JH2	QVQLVESGGGVVQPGRSRLSCAASGFTFSSYAMSWVRQAPGKGLEWVA AINSQGKSTY YADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAR WGDEGFDI WGRGLTVTVSS
SEQ ID NO: 490	MOR10703-VKI_A20_JK1	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKVPKLLIYGASSLQSGVPSR FSGSGSGTDFLTISLQPEDVATYYCQQYSSFPITTFGGGQTKVEIK
	Combination 6	
SEQ ID NO: 491	MOR10703-VH_3-33_JH3	QVQLVESGGGVVQPGRSRLSCAASGFTFSSYAMSWVRQAPGKGLEWVA AINSQGKSTY YADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAR WGDEGFDI WGQGMVTVSS
SEQ ID NO: 492	MOR10703-VKI_A20_JK2	DIQMTQSPSSLSASVGDRTITCRASQGISNWLAWYQQKPGKVPKLLIYGASSLQSGVPSR FSGSGSGTDFLTISLQPEDVATYYCQQYSSFPITTFGGGQTKLEIK

In one embodiment, the invention pertains to a heavy chain variable region comprising a sequence of Xaa₁-HCDR1-Xaa₂-HCDR2-Xaa₃-HCDR3-Xaa₄ where the heavy chain HCDR1, HCDR2, HCDR3 are any heavy chain CDRs selected from Tables 1 and 2. For illustrative purposes only, the sequence can be:

Xaa₁ - SYAMS - Xaa₂ - AINSQGKSTYYADSVKG - Xaa₃ - WGDEGFDI - Xaa₄ (SEQ ID NO: 493), where,

Xaa₁ is framework region of any 30 amino acids;

Xaa₂ is framework region of any 14 amino acids;

10 Xaa₃ is framework region of any 32 amino acids;

Xaa₄ is framework region of any 11 amino acids;

In one embodiment, the invention pertains to a light chain variable region comprising a sequence of Xaa₁-LCDR1-Xaa₂-LCDR2-Xaa₃-LCDR3-Xaa₄, where the light chain LCDR1,

LCDR2, LCDR3 are any light chain CDRs selected from Tables 1 and 2. For illustrative purposes only, the sequence can be:

Xaa₁ - RASQGISNWL A - Xaa₂ - GASSLQS - Xaa₃ - QQYSSFPTT - Xaa₄ (SEQ ID NO: 494), where,

- 5 Xaa₁ is a framework region of any 23 amino acids;
- Xaa₂ is a framework region of any 15 amino acids;
- Xaa₃ is a framework region of any 32 amino acids; and
- Xaa₄ is a framework region of any 10 amino acids.

The antibodies disclosed herein can be derivatives of single chain antibodies, diabodies,
 10 domain antibodies, nanobodies, and unibodies. A "single-chain antibody" (scFv) consists of a single polypeptide chain comprising a VL domain linked to a VH domain, wherein VL domain and VH domain are paired to form a monovalent molecule. Single chain antibody can be prepared according to method known in the art (see, for example, Bird *et al.*, (1988) Science 242:423-426 and Huston *et al.*, (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883). A
 15 "disbud" consists of two chains, each chain comprising a heavy chain variable region connected to a light chain variable region on the same polypeptide chain connected by a short peptide linker, wherein the two regions on the same chain do not pair with each other but with complementary domains on the other chain to form a bispecific molecule. Methods of preparing diabodies are known in the art (See, e.g., Holliger *et al.*, (1993) Proc. Natl. Acad. Sci. USA 90:6444-6448, and Poljak *et al.*, (1994) Structure 2:1121-1123). Domain antibodies
 20 (dAbs) are small functional binding units of antibodies, corresponding to the variable regions of either the heavy or light chains of antibodies. Domain antibodies are well expressed in bacterial, yeast, and mammalian cell systems. Further details of domain antibodies and methods of production thereof are known in the art (see, for example, U.S. Pat. Nos.
 25 6,291,158; 6,582,915; 6,593,081; 6,172,197; 6,696,245; European Patents 0368684 & 0616640; WO05/035572, WO04/101790, WO04/081026, WO04/058821, WO04/003019 and WO03/002609. Nanobodies are derived from the heavy chains of an antibody. A nanobody typically comprises a single variable domain and two constant domains (CH2 and CH3) and retains antigen-binding capacity of the original antibody. Nanobodies can be prepared by
 30 methods known in the art (See e.g., U.S. Pat. No. 6,765,087, U.S. Pat. No. 6,838,254, WO 06/079372). Unibodies consist of one light chain and one heavy chain of a IgG4 antibody.

Unibodies may be made by the removal of the hinge region of IgG4 antibodies. Further details of unibodies and methods of preparing them may be found in WO2007/059782.

Homologous antibodies

In yet another embodiment, the present invention provides an antibody or fragment thereof comprising amino acid sequences that are homologous to the sequences described in Table 1, and said antibody binds to a HER3 protein (*e.g.*, human and/or cynomologus HER3), and retains the desired functional properties of those antibodies described in Table 1.

For example, the invention provides an isolated monoclonal antibody (or a functional fragment thereof) comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises an amino acid sequence that is at least 80%, at least 90%, or at least 95% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 15, 33, 51, 69, 87, 105, 123, 141, 159, 177, 195, 213, 231, 249, 267, 285, 303, 321, 339, 357, and 375; the light chain variable region comprises an amino acid sequence that is at least 80%, at least 90%, or at least 95% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 14, 32, 50, 68, 86, 104, 122, 140, 158, 176, 194, 212, 230, 248, 266, 284, 302, 320, 338, 356, and 374; the antibody binds to HER3 (*e.g.*, human and/or cynomologus HER3) and neutralizes the signaling activity of HER3, which can be measured in a phosphorylation assay or other measure of HER signaling (*e.g.*, phospo-HER3 assays, phospo-Akt assays, cell proliferation, and ligand blocking assays as described in the Examples). Also included within the scope of the invention are variable heavy and light chain parental nucleotide sequences; and full length heavy and light chain sequences optimized for expression in a mammalian cell. Other antibodies of the invention include amino acids or nucleic acids that have been mutated, yet have at least 60, 70, 80, 90, 95, or 98% percent identity to the sequences described above. In some embodiments, it include mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated by amino acid deletion, insertion or substitution in the variable regions when compared with the variable regions depicted in the sequence described above.

In other embodiments, the VH and/or VL amino acid sequences may be 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth in Table 1. In other embodiments, the VH and/or VL amino acid sequences may be identical except an amino acid substitution in no more than 1,2,3,4 or 5 amino acid position. An antibody having

VH and VL regions having high (*i. e.*, 80% or greater) identity to the VH and VL regions of the antibodies described in Table 1 can be obtained by mutagenesis (*e.g.*, site-directed or PCR-mediated mutagenesis), followed by testing of the encoded altered antibody for retained function using the functional assays described herein.

- 5 In other embodiments, the variable regions of heavy chain and/or light chain nucleotide sequences may be 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth above.

As used herein, “percent identity” between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity equals number of identical
10 positions/total number of positions x 100), taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

- 15 Additionally or alternatively, the protein sequences of the present invention can further be used as a “query sequence” to perform a search against public databases to, for example, identify related sequences. For example, such searches can be performed using the BLAST program (version 2.0) of Altschul *et al.*, (1990) *J.Mol. Biol.* 215:403-10.

Antibodies with Conservative Modifications

- 20 In certain embodiments, an antibody of the invention has a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein one or more of these CDR sequences have specified amino acid sequences based on the antibodies described herein or conservative
25 modifications thereof, and wherein the antibodies retain the desired functional properties of the HER3-binding antibodies of the invention.

Accordingly, the invention provides an isolated HER3 monoclonal antibody, or a fragment thereof, consisting of a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein: the heavy chain variable region CDR1 amino acid sequences are selected from the
30 group consisting of SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296,

308, 314, 326, 332, 344, 350, 362, and 368, and conservative modifications thereof; the heavy chain variable region CDR2 amino acid sequences are selected from the group consisting of SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111, 117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369 and conservative modifications thereof; the heavy chain variable region CDR3 amino acid sequences are selected from the group consisting of SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 76, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370 and conservative modifications thereof; the light chain variable regions CDR1 amino acid sequences are selected from the group consisting of SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137, 149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371 and conservative modifications thereof; the light chain variable regions CDR2 amino acid sequences are selected from the group consisting of SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372, and conservative modifications thereof; the light chain variable regions of CDR3 amino acid sequences are selected from the group consisting of SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 133, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373, and conservative modifications thereof; the antibody or fragment thereof specifically binds to HER3, and neutralizes HER3 activity by inhibiting a HER signaling pathway, which can be measured in a phosphorylation assay or other measure of HER signaling (e.g., phospo-HER3 assays, phospo-Akt assays, cell proliferation, and ligand blocking assays as described in the Examples).

Antibodies That Bind to the Same Epitope

The present invention provides antibodies that interacts with (e.g., by binding, steric hindrance, stabilizing/destabilizing, spatial distribution) the same epitope as do the HER3-binding antibodies described in Table 1 and Fig. 7. Additional antibodies can therefore be identified based on their ability to cross-compete (e.g., to competitively inhibit the binding of,

in a statistically significant manner) with other antibodies of the invention in HER3 binding assays. The ability of a test antibody to inhibit the binding of antibodies of the present invention to a HER3 protein (*e.g.*, human and/or cynomologus HER3) demonstrates that the test antibody can compete with that antibody for binding to HER3; such an antibody may, according to non-limiting theory, bind to the same or a related (*e.g.*, a structurally similar or spatially proximal) epitope on the HER3 protein as the antibody with which it competes. In a certain embodiment, the antibody that binds to the same epitope on HER3 as the antibodies of the present invention is a human monoclonal antibody. Such human monoclonal antibodies can be prepared and isolated as described herein.

- 10 In one embodiment, the antibody or fragments thereof binds to both domain 2 and domain 4 of HER3 to hold the HER3 in an inactive conformation which prevents exposure of an dimerization loop present within domain 2. This prevents heterodimerization with other family members, such as HER1, HER2, and HER4. The antibodies or fragments thereof inhibit both ligand dependent and ligand-independent HER3 signal transduction.
- 15 In another embodiment, the antibody or fragment thereof binds to both domain 2 and domain 4 of HER3 and without blocking the concurrent binding of a HER3 ligand such as neuregulin. While not required to provide a theory, it is feasible that the antibody or fragment thereof binding to both domain 2 and domain 4 of HER3, holds HER3 in an inactive conformation without blocking the ligand binding site on HER3. Thus a HER3 ligand (*e.g.*, neuregulin) is able to bind to HER3 at the same time as the antibody or fragment thereof.
- 20

The antibodies of the invention or fragments thereof inhibit both ligand dependent and independent activation of HER3 without preventing ligand binding. This is considered advantageous for the following reasons:

- (i) The therapeutic antibody would have clinical utility in a broad spectrum of tumors than an antibody which targeted a single mechanism of HER3 activation (*i.e.* ligand dependent or ligand independent) since distinct tumor types are driven by each mechanism.
- (ii) The therapeutic antibody would be efficacious in tumor types where both mechanisms of HER3 activation are simultaneously involved. An antibody targeting a single mechanism of HER3 activation (*i.e.* ligand dependent or ligand independent) would display little or no efficacy in these tumor types
- 25
- 30

(iii) The efficacy of an antibody which inhibits ligand dependent activation of HER3 without preventing ligand binding would be less likely to be adversely affected by increasing concentrations of ligand. This would translate to either increased efficacy in a tumor type driven by very high concentrations of HER3 ligand or a reduced drug resistance liability where resistance is mediated by up-regulation of HER3 ligands.

(iv) An antibody which inhibits HER3 activation by stabilizing the inactive form would be less prone to drug resistance driven by alternative mechanisms of HER3 activation.

Consequently, the antibodies of the invention may be used to treat conditions where existing therapeutic antibodies are clinically ineffective.

10 **Engineered and Modified Antibodies**

An antibody of the invention further can be prepared using an antibody having one or more of the VH and/or VL sequences shown herein as starting material to engineer a modified antibody, which modified antibody may have altered properties from the starting antibody.

An antibody can be engineered by modifying one or more residues within one or both variable regions (*i. e.*, VH and/or VL), for example within one or more CDR regions and/or within one or more framework regions. Additionally or alternatively, an antibody can be engineered by modifying residues within the constant region(s), for example to alter the effector function(s) of the antibody.

One type of variable region engineering that can be performed is CDR grafting. Antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see, *e.g.*, Riechmann *et al.*, (1998) *Nature* 332:323-327; Jones *et al.*, (1986) *Nature* 321:522-525; Queen *et al.*, (1989) *Proc. Natl. Acad., U.S.A.* 86:10029-10033; U.S. Patent No. 5,225,539 to Winter, and U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen *et al.*)

Accordingly, another embodiment of the invention pertains to an isolated HER3 binding monoclonal antibody, or fragment thereof, comprising a heavy chain variable region comprising CDR1 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296, 308, 314, 326, 332, 344, 350, 362, and 368; CDR2 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111, 117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369; CDR3 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 76, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370, respectively; and a light chain variable region having CDR1 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137, 149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371; CDR2 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372; and CDR3 sequences consisting of an amino acid sequence selected from the group consisting of SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 135, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373, respectively. Thus, such antibodies contain the VH and VL CDR sequences of monoclonal antibodies, yet may contain different framework sequences from these antibodies. Such framework sequences can be obtained from public DNA databases or published references that include germline antibody gene sequences. For example, germline DNA sequences for human heavy and light chain variable region genes can be found in the "Vase" human germline sequence database (available on the Internet at www.mrc-cpe.cam.ac.uk/vbase), as well as in Kabat *et al.*, (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Chothia *et al.*, (1987) J. Mol. Biol. 196:901-917; Chothia *et al.*, (1989) Nature 342:877-883; and Al-Lazikani *et al.*, (1997) J. Mol. Biol. 273:927-948; Tomlinson *et al.*, (1992) J. Mol. Biol. 227:776-798; and Cox *et al.*, (1994) Eur. J Immunol. 24:827-836; the contents of each of which are expressly incorporated herein by reference.

An example of framework sequences for use in the antibodies of the invention are those that are structurally similar to the framework sequences used by selected antibodies of the invention, *e.g.*, consensus sequences and/or framework sequences used by monoclonal antibodies of the invention. The VH CDR1, 2 and 3 sequences, and the VL CDR1, 2 and 3 sequences, can be grafted onto framework regions that have the identical sequence as that found in the germline immunoglobulin gene from which the framework sequence derive, or the CDR sequences can be grafted onto framework regions that contain one or more mutations as compared to the germline sequences. For example, it has been found that in certain instances it is beneficial to mutate residues within the framework regions to maintain or enhance the antigen binding ability of the antibody (see *e.g.*, U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen *et al*).

Another type of variable region modification is to mutate amino acid residues within the VH and/or VL CDR1, CDR2 and/or CDR3 regions to thereby improve one or more binding properties (*e.g.*, affinity) of the antibody of interest, known as "affinity maturation." Site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the mutation(s) and the effect on antibody binding, or other functional property of interest, can be evaluated in *in vitro* or *in vivo* assays as described herein and provided in the Examples. Conservative modifications (as discussed above) can be introduced. The mutations may be amino acid substitutions, additions or deletions. Moreover, typically no more than one, two, three, four or five residues within a CDR region are altered.

Accordingly, in another embodiment, the invention provides isolated HER3 binding monoclonal antibodies, or fragment thereof, consisting of a heavy chain variable region having: a VH CDR1 region consisting of an amino acid sequence selected from the group having SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296, 308, 314, 326, 332, 344, 350, 362, and 368 or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296, 308, 314, 326, 332, 344, 350, 362, and 368; a VH CDR2 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111, 117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369 or an amino acid sequence having one, two, three, four or five amino acid

substitutions, deletions or additions as compared to SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111, 117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369; a VH CDR3 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 76, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 76, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370; a VL CDR1 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137, 149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137, 149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371; a VL CDR2 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372; and a VL CDR3 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 135, 139, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 135, 139, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373.

Grafting Antibody Fragments Into Alternative Frameworks or Scaffolds

A wide variety of antibody/ immunoglobulin frameworks or scaffolds can be employed so long as the resulting polypeptide includes at least one binding region which specifically binds to HER3. Such frameworks or scaffolds include the 5 main idiotypes of human immunoglobulins, or fragments thereof, and include immunoglobulins of other animal species, preferably having humanized aspects. Novel frameworks, scaffolds and fragments continue to be discovered and developed by those skilled in the art.

In one aspect, the invention pertains to generating non-immunoglobulin based antibodies using non- immunoglobulin scaffolds onto which CDRs of the invention can be grafted. Known or future non-immunoglobulin frameworks and scaffolds may be employed, as long as they comprise a binding region specific for the target HER3 protein (*e.g.*, human and/or cynomologus HER3). Known non-immunoglobulin frameworks or scaffolds include, but are not limited to, fibronectin (Compound Therapeutics, Inc., Waltham, MA), ankyrin (Molecular Partners AG, Zurich, Switzerland), domain antibodies (Domantis, Ltd., Cambridge, MA, and Ablynx nv, Zwijnaarde, Belgium), lipocalin (Pieris Proteolab AG, Freising, Germany), small modular immuno-pharmaceuticals (Trubion Pharmaceuticals Inc., Seattle, WA), maxybodies (Avidia, Inc., Mountain View, CA), Protein A (Affibody AG, Sweden), and affilin (gamma-crystallin or ubiquitin) (Scil Proteins GmbH, Halle, Germany).

The fibronectin scaffolds are based on fibronectin type III domain (*e.g.*, the tenth module of the fibronectin type III (¹⁰ Fn3 domain)). The fibronectin type III domain has 7 or 8 beta strands which are distributed between two beta sheets, which themselves pack against each other to form the core of the protein, and further containing loops (analogous to CDRs) which connect the beta strands to each other and are solvent exposed. There are at least three such loops at each edge of the beta sheet sandwich, where the edge is the boundary of the protein perpendicular to the direction of the beta strands (see US 6,818,418). These fibronectin-based scaffolds are not an immunoglobulin, although the overall fold is closely related to that of the smallest functional antibody fragment, the variable region of the heavy chain, which comprises the entire antigen recognition unit in camel and llama IgG. Because of this structure, the non-immunoglobulin antibody mimics antigen binding properties that are similar in nature and affinity to those of antibodies. These scaffolds can be used in a loop randomization and shuffling strategy *in vitro* that is similar to the process of affinity maturation of antibodies *in vivo*. These fibronectin-based molecules can be used as scaffolds where the loop regions of the molecule can be replaced with CDRs of the invention using standard cloning techniques.

The ankyrin technology is based on using proteins with ankyrin derived repeat modules as scaffolds for bearing variable regions which can be used for binding to different targets. The ankyrin repeat module is a 33 amino acid polypeptide consisting of two anti-parallel α -helices and a β -turn. Binding of the variable regions is mostly optimized by using ribosome display.

5 Avimers are derived from natural A-domain containing protein such as HER3. These domains are used by nature for protein-protein interactions and in human over 250 proteins are structurally based on A-domains. Avimers consist of a number of different "A-domain" monomers (2-10) linked via amino acid linkers. Avimers can be created that can bind to the target antigen using the methodology described in, for example, U.S. Patent Application
10 Publication Nos. 20040175756; 20050053973; 20050048512; and 20060008844.

Affibody affinity ligands are small, simple proteins composed of a three-helix bundle based on the scaffold of one of the IgG-binding domains of Protein A. Protein A is a surface protein from the bacterium *Staphylococcus aureus*. This scaffold domain consists of 58 amino acids, 13 of which are randomized to generate affibody libraries with a large number of ligand
15 variants (See *e.g.*, US 5,831,012). Affibody molecules mimic antibodies, they have a molecular weight of 6 kDa, compared to the molecular weight of antibodies, which is 150 kDa. In spite of its small size, the binding site of affibody molecules is similar to that of an antibody.

Anticalins are products developed by the company Pieris ProteoLab AG. They are derived
20 from lipocalins, a widespread group of small and robust proteins that are usually involved in the physiological transport or storage of chemically sensitive or insoluble compounds. Several natural lipocalins occur in human tissues or body liquids. The protein architecture is reminiscent of immunoglobulins, with hypervariable loops on top of a rigid framework. However, in contrast with antibodies or their recombinant fragments, lipocalins are composed
25 of a single polypeptide chain with 160 to 180 amino acid residues, being just marginally bigger than a single immunoglobulin domain. The set of four loops, which makes up the binding pocket, shows pronounced structural plasticity and tolerates a variety of side chains. The binding site can thus be reshaped in a proprietary process in order to recognize prescribed target molecules of different shape with high affinity and specificity. One protein of lipocalin
30 family, the bilin-binding protein (BBP) of *Pieris Brassicae* has been used to develop anticalins by mutagenizing the set of four loops. One example of a patent application describing anticalins is in PCT Publication No. WO 199916873.

Affilin molecules are small non-immunoglobulin proteins which are designed for specific affinities towards proteins and small molecules. New affilin molecules can be very quickly selected from two libraries, each of which is based on a different human derived scaffold protein. Affilin molecules do not show any structural homology to immunoglobulin proteins.

5 Currently, two affilin scaffolds are employed, one of which is gamma crystalline, a human structural eye lens protein and the other is "ubiquitin" superfamily proteins. Both human scaffolds are very small, show high temperature stability and are almost resistant to pH changes and denaturing agents. This high stability is mainly due to the expanded beta sheet structure of the proteins. Examples of gamma crystalline derived proteins are described in
10 WO200104144 and examples of "ubiquitin-like" proteins are described in WO2004106368.

Protein epitope mimetics (PEM) are medium-sized, cyclic, peptide-like molecules (MW 1-2kDa) mimicking beta-hairpin secondary structures of proteins, the major secondary structure involved in protein-protein interactions.

In some embodiments, the Fabs are converted to silent IgG1 format by changing the Fc
15 region. For example, antibodies in Table 1 can be converted to IgG format.

Human or humanized antibodies

The present invention provides fully human antibodies that specifically bind to a HER3 protein (*e.g.*, human and/or cynomologus/ mouse/rat HER3). Compared to the chimeric or humanized antibodies, the human HER3 -binding antibodies of the invention have further
20 reduced antigenicity when administered to human subjects.

The human HER3-binding antibodies can be generated using methods that are known in the art. For example, the humanizing technology used to converting non-human antibodies into engineered human antibodies. U.S. Patent Publication No. 20050008625 describes an *in vivo* method for replacing a nonhuman antibody variable region with a human variable region in an
25 antibody while maintaining the same or providing better binding characteristics relative to that of the nonhuman antibody. The method relies on epitope guided replacement of variable regions of a non-human reference antibody with a fully human antibody. The resulting human antibody is generally unrelated structurally to the reference nonhuman antibody, but binds to the same epitope on the same antigen as the reference antibody. Briefly, the serial
30 epitope-guided complementarity replacement approach is enabled by setting up a competition in cells between a "competitor" and a library of diverse hybrids of the reference antibody ("test antibodies") for binding to limiting amounts of antigen in the presence of a reporter

system which responds to the binding of test antibody to antigen. The competitor can be the reference antibody or derivative thereof such as a single-chain Fv fragment. The competitor can also be a natural or artificial ligand of the antigen which binds to the same epitope as the reference antibody. The only requirements of the competitor are that it binds to the same
5 epitope as the reference antibody, and that it competes with the reference antibody for antigen binding. The test antibodies have one antigen-binding V-region in common from the nonhuman reference antibody, and the other V-region selected at random from a diverse source such as a repertoire library of human antibodies. The common V-region from the reference antibody serves as a guide, positioning the test antibodies on the same epitope on
10 the antigen, and in the same orientation, so that selection is biased toward the highest antigen-binding fidelity to the reference antibody.

Many types of reporter system can be used to detect desired interactions between test antibodies and antigen. For example, complementing reporter fragments may be linked to antigen and test antibody, respectively, so that reporter activation by fragment
15 complementation only occurs when the test antibody binds to the antigen. When the test antibody- and antigen-reporter fragment fusions are co-expressed with a competitor, reporter activation becomes dependent on the ability of the test antibody to compete with the competitor, which is proportional to the affinity of the test antibody for the antigen. Other reporter systems that can be used include the reactivator of an auto-inhibited reporter
20 reactivation system (RAIR) as disclosed in U.S. Patent Application Ser. No. 10/208,730 (Publication No. 20030198971), or competitive activation system disclosed in U.S. Patent Application Ser. No. 10/076,845 (Publication No. 20030157579).

With the serial epitope-guided complementarity replacement system, selection is made to identify cells expresses a single test antibody along with the competitor, antigen, and reporter
25 components. In these cells, each test antibody competes one-on-one with the competitor for binding to a limiting amount of antigen. Activity of the reporter is proportional to the amount of antigen bound to the test antibody, which in turn is proportional to the affinity of the test antibody for the antigen and the stability of the test antibody. Test antibodies are initially selected on the basis of their activity relative to that of the reference antibody when expressed
30 as the test antibody. The result of the first round of selection is a set of "hybrid" antibodies, each of which is comprised of the same non-human V-region from the reference antibody and a human V-region from the library, and each of which binds to the same epitope on the antigen as the reference antibody. One of more of the hybrid antibodies selected in the first

round will have an affinity for the antigen comparable to or higher than that of the reference antibody.

In the second V-region replacement step, the human V-regions selected in the first step are used as guide for the selection of human replacements for the remaining non-human reference antibody V-region with a diverse library of cognate human V-regions. The hybrid antibodies selected in the first round may also be used as competitors for the second round of selection. The result of the second round of selection is a set of fully human antibodies which differ structurally from the reference antibody, but which compete with the reference antibody for binding to the same antigen. Some of the selected human antibodies bind to the same epitope on the same antigen as the reference antibody. Among these selected human antibodies, one or more binds to the same epitope with an affinity which is comparable to or higher than that of the reference antibody.

Using one of the mouse or chimeric HER3-binding antibodies described above as the reference antibody, this method can be readily employed to generate human antibodies that bind to human HER3 with the same binding specificity and the same or better binding affinity. In addition, such human HER3-binding antibodies can also be commercially obtained from companies which customarily produce human antibodies, *e.g.*, KaloBios, Inc. (Mountain View, CA).

Camelid antibodies

Antibody proteins obtained from members of the camel and dromedary (*Camelus bactrianus* and *Camelus dromaderius*) family including new world members such as llama species (*Lama pacos*, *Lama glama* and *Lama vicugna*) have been characterized with respect to size, structural complexity and antigenicity for human subjects. Certain IgG antibodies from this family of mammals as found in nature lack light chains, and are thus structurally distinct from the typical four chain quaternary structure having two heavy and two light chains, for antibodies from other animals. See PCT/EP93/02214 (WO 94/04678 published 3 March 1994).

A region of the camelid antibody which is the small single variable domain identified as VHH can be obtained by genetic engineering to yield a small protein having high affinity for a target, resulting in a low molecular weight antibody-derived protein known as a "camelid nanobody". See U.S. patent number 5,759,808 issued June 2, 1998; see also Stijlemans *et al.*, (2004) *J Biol Chem* 279:1256-1261; Dumoulin *et al.*, (2003) *Nature* 424:783-788;

Pleschberger *et al.*, (2003) Bioconjugate Chem 14:440-448; Cortez-Retamozo *et al.*, (2002) Int J Cancer 89:456-62; and Lauwereys *et al.*, (1998) EMBO J 17:3512-3520. Engineered libraries of camelid antibodies and antibody fragments are commercially available, for example, from Ablynx, Ghent, Belgium . (e.g., US20060115470; Domantis (US20070065440, 5 US20090148434). As with other antibodies of non-human origin, an amino acid sequence of a camelid antibody can be altered recombinantly to obtain a sequence that more closely resembles a human sequence, *i.e.*, the nanobody can be “humanized”. Thus the natural low antigenicity of camelid antibodies to humans can be further reduced.

The camelid nanobody has a molecular weight approximately one-tenth that of a human IgG 10 molecule, and the protein has a physical diameter of only a few nanometers. One consequence of the small size is the ability of camelid nanobodies to bind to antigenic sites that are functionally invisible to larger antibody proteins, *i.e.*, camelid nanobodies are useful as reagents detect antigens that are otherwise cryptic using classical immunological techniques, and as possible therapeutic agents. Thus yet another consequence of small size is 15 that a camelid nanobody can inhibit as a result of binding to a specific site in a groove or narrow cleft of a target protein, and hence can serve in a capacity that more closely resembles the function of a classical low molecular weight drug than that of a classical antibody.

The low molecular weight and compact size further result in camelid nanobodies being extremely thermostable, stable to extreme pH and to proteolytic digestion, and poorly 20 antigenic. Another consequence is that camelid nanobodies readily move from the circulatory system into tissues, and even cross the blood-brain barrier and can treat disorders that affect nervous tissue. Nanobodies can further facilitated drug transport across the blood brain barrier. See U.S. patent application 20040161738 published August 19, 2004. These features combined with the low antigenicity to humans indicate great therapeutic potential. Further, 25 these molecules can be fully expressed in prokaryotic cells such as *E. coli* and are expressed as fusion proteins with bacteriophage and are functional.

Accordingly, a feature of the present invention is a camelid antibody or nanobody having high affinity for HER3. In certain embodiments herein, the camelid antibody or nanobody is naturally produced in the camelid animal, *i.e.*, is produced by the camelid following 30 immunization with HER3 or a peptide fragment thereof, using techniques described herein for other antibodies. Alternatively, the HER3-binding camelid nanobody is engineered, *i.e.*, produced by selection for example from a library of phage displaying appropriately mutagenized camelid nanobody proteins using panning procedures with HER3 as a target as

described in the examples herein. Engineered nanobodies can further be customized by genetic engineering to have a half life in a recipient subject of from 45 minutes to two weeks. In a specific embodiment, the camelid antibody or nanobody is obtained by grafting the CDRs sequences of the heavy or light chain of the human antibodies of the invention into nanobody or single domain antibody framework sequences, as described for example in PCT/EP93/02214. In one embodiment, the camelid antibody or nanobody binds to at least one of the following HER3 residues : Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597. In one embodiment, the camelid antibody or nanobody binds to at least one of the following HER3 residues : Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615.

Bispecific Molecules and Multivalent Antibodies

In another aspect, the present invention features biparatopic, bispecific or multispecific molecules comprising an HER3-binding antibody, or a fragment thereof, of the invention. An antibody of the invention, or fragments thereof, can be derivatized or linked to another functional molecule, *e.g.*, another peptide or protein (*e.g.*, another antibody or ligand for a receptor) to generate a bispecific molecule that binds to at least two different binding sites or target molecules. The antibody of the invention may in fact be derivatized or linked to more than one other functional molecule to generate biparatopic or multi-specific molecules that bind to more than two different binding sites and/or target molecules; such biparatopic or multi-specific molecules. To create a bispecific molecule of the invention, an antibody of the invention can be functionally linked (*e.g.*, by chemical coupling, genetic fusion, non-covalent association or otherwise) to one or more other binding molecules, such as another antibody, antibody fragment, peptide or binding mimetic, such that a bispecific molecule results.

Further clinical benefits may be provided by the binding of two or more antigens within one antibody (Coloma *et al.*, (1997); Merchant *et al.*, (1998); Alt *et al.*, (1999); Zuo *et al.*, (2000); Lu *et al.*, (2004); Lu *et al.*, (2005); Marvin *et al.*, (2005); Marvin *et al.*, (2006); Shen *et al.*, (2007); Wu *et al.*, (2007); Dimasi *et al.*, (2009); Michaelson *et al.*, (2009)). (Morrison *et al.*, (1997) Nature Biotech. 15:159-163; Alt *et al.* (1999) FEBS Letters 454:90-94; Zuo *et al.*, (2000) Protein Engineering 13:361-367; Lu *et al.*, (2004) JBC 279:2856-2865; Lu *et al.*, (2005) JBC 280:19665-19672; Marvin *et al.*, (2005) Acta Pharmacologica Sinica 26:649-658; Marvin *et al.*, (2006) Curr Opin Drug Disc Develop 9:184-193; Shen *et al.*, (2007) J

Immun Methods 218:65-74; Wu *et al.*, (2007) Nat Biotechnol. 11:1290-1297; Dimasi *et al.*, (2009) J Mol Biol. 393:672-692; and Michaelson *et al.*, (2009) mAbs 1:128-141.

The bispecific molecules of the present invention can be prepared by conjugating the constituent binding specificities, using methods known in the art. For example, each binding specificity of the bispecific molecule can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-S-acetyl-thioacetate (SATA), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC) (see *e.g.*, Karpovsky *et al.*, (1984) J. Exp. Med. 160:1686; Liu *et al.*, (1985) Proc. Natl. Acad. Sci. USA 82:8648). Other methods include those described in Paulus (1985) Behring Ins. Mitt. No. 78:118-132; Brennan *et al.*, (1985) Science 229:81-83), and Glennie *et al.*, (1987) J. Immunol. 139: 2367-2375). Conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, IL).

When the binding specificities are antibodies, they can be conjugated by sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In a particularly embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, for example one, prior to conjugation.

Alternatively, both binding specificities can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific molecule is a mAb x mAb, mAb x Fab, Fab x F(ab')₂ or ligand x Fab fusion protein. A bispecific molecule of the invention can be a single chain molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific molecules may comprise at least two single chain molecules. Methods for preparing bispecific molecules are described for example in U.S. Patent Number 5,260,203; U.S. Patent Number 5,455,030; U.S. Patent Number 4,881,175; U.S. Patent Number 5,132,405; U.S. Patent Number 5,091,513; U.S. Patent Number 5,476,786; U.S. Patent Number 5,013,653; U.S. Patent Number 5,258,498; and U.S. Patent Number 5,482,858.

Binding of the bispecific molecules to their specific targets can be confirmed by, for example, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (REA), FACS analysis,

bioassay (*e.g.*, growth inhibition), or Western Blot assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (*e.g.*, an antibody) specific for the complex of interest.

In another aspect, the present invention provides multivalent compounds comprising at least two identical or different fragments of the antibodies of the invention binding to HER3. The antibody fragments can be linked together via protein fusion or covalent or non covalent linkage. Tetravalent compounds can be obtained for example by cross-linking antibodies of the antibodies of the invention with an antibody that binds to the constant regions of the antibodies of the invention, for example the Fc or hinge region. Trimerizing domain are described for example in Borean patent EP 1012280B1. Pentamerizing modules are described for example in PCT/EP97/05897.

In one embodiment, a biparatopic/bispecific binds to amino acid residues within domain 2 and domain 4 of HER3.

In another embodiment, the invention pertains to dual function antibodies in which a single monoclonal antibody has been modified such that the antigen binding site binds to more than one antigen, such as a dual function antibody which binds both HER3 and another antigen (*e.g.*, HER1, HER2, and HER4). In another embodiment, the invention pertains to a dual function antibody that targets antigens having the same conformation, for example an antigen that has the same conformation of HER3 in the “closed” or “inactive” state. Examples of antigens with the same conformation of HER3 in the “closed” or “inactive” state include, but are not limited to, HER1 and HER4. Thus, a dual function antibody may bind to both HER3 and HER1; HER3 and HER4, or HER1 and HER4. The dual binding specificity of the dual function antibody may further translate into dual activity, or inhibition of activity. (See *e.g.*, Jenny Bostrom *et al.*, (2009) *Science*: 323; 1610-1614).

25 **Antibodies with Extended Half Life**

The present invention provides for antibodies that specifically bind to HER3 protein which have an extended half-life *in vivo*.

Many factors may affect a protein’s half life *in vivo*. For examples, kidney filtration, metabolism in the liver, degradation by proteolytic enzymes (proteases), and immunogenic responses (*e.g.*, protein neutralization by antibodies and uptake by macrophages and dendritic cells). A variety of strategies can be used to extend the half life of the antibodies of the

present invention. For example, by chemical linkage to polyethyleneglycol (PEG), reCODE PEG, antibody scaffold, polysialic acid (PSA), hydroxyethyl starch (HES), albumin-binding ligands, and carbohydrate shields; by genetic fusion to proteins binding to serum proteins, such as albumin, IgG, FcRn, and transferring; by coupling (genetically or chemically) to other
5 binding moieties that bind to serum proteins, such as nanobodies, Fabs, DARPins, avimers, affibodies, and anticalins; by genetic fusion to rPEG, albumin, domain of albumin, albumin-binding proteins, and Fc; or by incorporation into nanocarriers, slow release formulations, or medical devices.

To prolong the serum circulation of antibodies *in vivo*, inert polymer molecules such as high
10 molecular weight PEG can be attached to the antibodies or a fragment thereof with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C-terminus of the antibodies or via epsilon-amino groups present on lysine residues. To pegylate an antibody, the antibody, or fragment thereof, typically is reacted with polyethylene glycol (PEG), such as a reactive ester or aldehyde derivative of PEG, under conditions in
15 which one or more PEG groups become attached to the antibody or antibody fragment. The pegylation can be carried out by an acylation reaction or an alkylation reaction with a reactive PEG molecule (or an analogous reactive water-soluble polymer). As used herein, the term "polyethylene glycol" is intended to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol or
20 polyethylene glycol-maleimide. In certain embodiments, the antibody to be pegylated is an aglycosylated antibody. Linear or branched polymer derivatization that results in minimal loss of biological activity will be used. The degree of conjugation can be closely monitored by SDS-PAGE and mass spectrometry to ensure proper conjugation of PEG molecules to the antibodies. Unreacted PEG can be separated from antibody-PEG conjugates by size-
25 exclusion or by ion-exchange chromatography. PEG-derivatized antibodies can be tested for binding activity as well as for *in vivo* efficacy using methods well-known to those of skill in the art, for example, by immunoassays described herein. Methods for pegylating proteins are known in the art and can be applied to the antibodies of the invention. See for example, EP 0 154 316 by Nishimura *et al.* and EP 0 401 384 by Ishikawa *et al.*

30 Other modified pegylation technologies include reconstituting chemically orthogonal directed engineering technology (ReCODE PEG), which incorporates chemically specified side chains into biosynthetic proteins via a reconstituted system that includes tRNA synthetase and tRNA. This technology enables incorporation of more than 30 new amino acids into biosynthetic

proteins in *E.coli*, yeast, and mammalian cells. The tRNA incorporates a nonnative amino acid any place an amber codon is positioned, converting the amber from a stop codon to one that signals incorporation of the chemically specified amino acid.

Recombinant pegylation technology (rPEG) can also be used for serum half-life extension.

5 This technology involves genetically fusing a 300-600 amino acid unstructured protein tail to an existing pharmaceutical protein. Because the apparent molecular weight of such an unstructured protein chain is about 15-fold larger than its actual molecular weight, the serum half-life of the protein is greatly increased. In contrast to traditional PEGylation, which requires chemical conjugation and repurification, the manufacturing process is greatly
10 simplified and the product is homogeneous.

Polysialylation is another technology, which uses the natural polymer polysialic acid (PSA) to prolong the active life and improve the stability of therapeutic peptides and proteins. PSA is a polymer of sialic acid (a sugar). When used for protein and therapeutic peptide drug delivery, polysialic acid provides a protective microenvironment on conjugation. This increases the
15 active life of the therapeutic protein in the circulation and prevents it from being recognized by the immune system. The PSA polymer is naturally found in the human body. It was adopted by certain bacteria which evolved over millions of years to coat their walls with it. These naturally polysialylated bacteria were then able, by virtue of molecular mimicry, to foil the body's defense system. PSA, nature's ultimate stealth technology, can be easily produced
20 from such bacteria in large quantities and with predetermined physical characteristics. Bacterial PSA is completely non-immunogenic, even when coupled to proteins, as it is chemically identical to PSA in the human body.

Another technology include the use of hydroxyethyl starch ("HES") derivatives linked to antibodies. HES is a modified natural polymer derived from waxy maize starch and can be
25 metabolized by the body's enzymes. HES solutions are usually administered to substitute deficient blood volume and to improve the rheological properties of the blood. Hesylation of an antibody enables the prolongation of the circulation half-life by increasing the stability of the molecule, as well as by reducing renal clearance, resulting in an increased biological activity. By varying different parameters, such as the molecular weight of HES, a wide range
30 of HES antibody conjugates can be customized.

Antibodies having an increased half-life *in vivo* can also be generated introducing one or more amino acid modifications (*i.e.*, substitutions, insertions or deletions) into an IgG constant

domain, or FcRn binding fragment thereof (preferably a Fc or hinge Fc domain fragment). See, *e.g.*, International Publication No. WO 98/23289; International Publication No. WO 97/34631; and U.S. Patent No. 6,277,375.

Further, antibodies can be conjugated to albumin in order to make the antibody or antibody
5 fragment more stable *in vivo* or have a longer half life *in vivo*. The techniques are well-known in the art, see, *e.g.*, International Publication Nos. WO 93/15199, WO 93/15200, and WO 01/77137; and European Patent No. EP 413,622.

The HER3 antibody or a fragment thereof may also be fused to one or more human serum albumin (HSA) polypeptides, or a portion thereof. HSA, a protein of 585 amino acids in its
10 mature form, is responsible for a significant proportion of the osmotic pressure of serum and also functions as a carrier of endogenous and exogenous ligands. The role of albumin as a carrier molecule and its inert nature are desirable properties for use as a carrier and transporter of polypeptides *in vivo*. The use of albumin as a component of an albumin fusion protein as a carrier for various proteins has been suggested in WO 93/15199, WO 93/15200, and EP 413
15 622. The use of N-terminal fragments of HSA for fusions to polypeptides has also been proposed (EP 399 666). Accordingly, by genetically or chemically fusing or conjugating the antibodies or fragments thereof to albumin, can stabilize or extend the shelf-life, and/or to retain the molecule's activity for extended periods of time in solution, *in vitro* and/or *in vivo*.

Fusion of albumin to another protein may be achieved by genetic manipulation, such that the
20 DNA coding for HSA, or a fragment thereof, is joined to the DNA coding for the protein. A suitable host is then transformed or transfected with the fused nucleotide sequences, so arranged on a suitable plasmid as to express a fusion polypeptide. The expression may be effected *in vitro* from, for example, prokaryotic or eukaryotic cells, or *in vivo e.g.* from a transgenic organism. Additional methods pertaining to HSA fusions can be found, for
25 example, in WO 2001077137 and WO 200306007, incorporated herein by reference. In a specific embodiment, the expression of the fusion protein is performed in mammalian cell lines, for example, CHO cell lines. Altered differential binding of an antibody to a receptor at low or high pHs is also contemplated to be within the scope of the invention. For example, the affinity of an antibody may be modified such that it remains bound to its receptor at a low
30 pH, *e.g.*, the low pH within a lysosome, by modifying the antibody to include additional amino acids such as a histine in a CDR of the antibody (See *e.g.*, Tomoyuki Igawa *et al.* (2010) *Nature Biotechnology*; 28, 1203-1207).

Antibody Conjugates

The present invention provides antibodies or fragments thereof that specifically bind to a HER3 protein recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, preferably to a polypeptide of at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids) to generate fusion proteins. In particular, the invention provides fusion proteins comprising an antibody fragment described herein (*e.g.*, a Fab fragment, Fd fragment, Fv fragment, F(ab)₂ fragment, a VH domain, a VH CDR, a VL domain or a VL CDR) and a heterologous protein, polypeptide, or peptide. Methods for fusing or conjugating proteins, polypeptides, or peptides to an antibody or an antibody fragment are known in the art. See, *e.g.*, U.S. Patent Nos. 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, and 5,112,946; European Patent Nos. EP 307,434 and EP 367,166; International Publication Nos. WO 96/04388 and WO 91/06570; Ashkenazi *et al.*, (1991) Proc. Natl. Acad. Sci. USA 88:10535-10539; Zheng *et al.*, (1995) J. Immunol. 154:5590-5600; and Vil *et al.*, (1992) Proc. Natl. Acad. Sci. USA 89:11337- 11341.

Additional fusion proteins may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as “DNA shuffling”). DNA shuffling may be employed to alter the activities of antibodies of the invention or fragments thereof (*e.g.*, antibodies or fragments thereof with higher affinities and lower dissociation rates). See, generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458; Patten *et al.*, (1997) Curr. Opin. Biotechnol. 8:724-33; Harayama, (1998) Trends Biotechnol. 16(2):76-82; Hansson *et al.*, (1999) J. Mol. Biol. 287:265-76; and Lorenzo and Blasco, (1998) Biotechniques 24(2):308- 313 (each of these patents and publications are hereby incorporated by reference in its entirety). Antibodies or fragments thereof, or the encoded antibodies or fragments thereof, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. A polynucleotide encoding an antibody or fragment thereof that specifically binds to a HER3 protein may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. Moreover, the antibodies or fragments thereof can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially

available. As described in Gentz *et al.*, (1989) Proc. Natl. Acad. Sci. USA 86:821-824, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin (“HA”) tag, which corresponds to an epitope derived from the influenza hemagglutinin protein
5 (Wilson *et al.*, (1984) Cell 37:767), and the “flag” tag.

In other embodiments, antibodies of the present invention or fragments thereof conjugated to a diagnostic or detectable agent. Such antibodies can be useful for monitoring or prognosing the onset, development, progression and/or severity of a disease or disorder as part of a clinical testing procedure, such as determining the efficacy of a particular therapy. Such
10 diagnosis and detection can be accomplished by coupling the antibody to detectable substances including, but not limited to, various enzymes, such as, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as, but not limited to, streptavidin/biotin and avidin/biotin; fluorescent materials, such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine,
15 dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent materials, such as, but not limited to, luminol; bioluminescent materials, such as but not limited to, luciferase, luciferin, and aequorin; radioactive materials, such as, but not limited to, iodine (¹³¹I, ¹²⁵I, ¹²³I, and ¹²¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹⁵In, ¹¹³In, ¹¹²In, and ¹¹¹In), technetium (⁹⁹Tc), thallium (²⁰¹Tl), gallium (⁶⁸Ga, ⁶⁷Ga), palladium (¹⁰³Pd),
20 molybdenum (⁹⁹Mo), xenon (¹³³Xe), fluorine (¹⁸F), ¹⁵³Sm, ¹⁷⁷Lu, ¹⁵⁹Gd, ¹⁴⁹Pm, ¹⁴⁰La, ¹⁷⁵Yb, ¹⁶⁶Ho, ⁹⁰Y, ⁴⁷Sc, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, ⁹⁷Ru, ⁶⁸Ge, ⁵⁷Co, ⁶⁵Zn, ⁸⁵Sr, ³²P, ¹⁵³Gd, ¹⁶⁹Yb, ⁵¹Cr, ⁵⁴Mn, ⁷⁵Se, ¹¹³Sn, and ¹¹⁷Tm; and positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions.

The present invention further encompasses uses of antibodies or fragments thereof conjugated
25 to a therapeutic moiety. An antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, *e.g.*, a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, *e.g.*, alpha-emitters. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells.

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety or drug
30 moiety that modifies a given biological response. Therapeutic moieties or drug moieties are not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein, peptide, or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas

exotoxin, cholera toxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, an anti-angiogenic agent; or, a biological response modifier such as, for example, a lymphokine. In one embodiment, the anti-HER3 antibody, or
5 a fragment thereof, conjugated to a therapeutic moiety, such as a cytotoxin, a drug (e.g., an immunosuppressant) or a radiotoxin. Such conjugates are referred to herein as “immunoconjugates”. Immunoconjugates that include one or more cytotoxins are referred to as “immunotoxins.” A cytotoxin or cytotoxic agent includes any agent that is detrimental to (e.g., kills) cells. Examples include taxon, cytochalasin B, gramicidin D, ethidium bromide,
10 emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, t. colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1 - dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents also include, for example, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-
15 fluorouracil decarbazine), ablating agents (e.g., mechlorethamine, thioepa chloraxnbucil, meiphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin, anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and
20 anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). (See e.g., Seattle Genetics US20090304721).

Other examples of therapeutic cytotoxins that can be conjugated to an antibody of the invention include duocarmycins, calicheamicins, maytansines and auristatins, and derivatives thereof. An example of a calicheamicin antibody conjugate is commercially available
25 (Mylotarg™; Wyeth-Ayerst).

Cytotoxins can be conjugated to antibodies of the invention using linker technology available in the art. Examples of linker types that have been used to conjugate a cytotoxin to an antibody include, but are not limited to, hydrazones, thioethers, esters, disulfides and peptide-containing linkers. A linker can be chosen that is, for example, susceptible to cleavage by low
30 pH within the lysosomal compartment or susceptible to cleavage by proteases, such as proteases preferentially expressed in tumor tissue such as cathepsins (e.g., cathepsins B, C, D).

For further discussion of types of cytotoxins, linkers and methods for conjugating therapeutic agents to antibodies, see also Saito *et al.*, (2003) *Adv. Drug Deliv. Rev.* 55:199-215; Trail *et al.*, (2003) *Cancer Immunol. Immunother.* 52:328-337; Payne, (2003) *Cancer Cell* 3:207-212; Allen, (2002) *Nat. Rev. Cancer* 2:750-763; Pastan and Kreitman, (2002) *Curr. Opin. Investig. Drugs* 3:1089-1091; Senter and Springer, (2001) *Adv. Drug Deliv. Rev.* 53:247-264.

Antibodies of the present invention also can be conjugated to a radioactive isotope to generate cytotoxic radiopharmaceuticals, also referred to as radioimmunoconjugates. Examples of radioactive isotopes that can be conjugated to antibodies for use diagnostically or therapeutically include, but are not limited to, iodine¹³¹, indium¹¹¹, yttrium⁹⁰, and lutetium¹⁷⁷.

Method for preparing radioimmunconjugates are established in the art. Examples of radioimmunoconjugates are commercially available, including ZevalinTM (DEC Pharmaceuticals) and BexxarTM (Corixa Pharmaceuticals), and similar methods can be used to prepare radioimmunoconjugates using the antibodies of the invention. In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo *et al.*, (1998) *Clin Cancer Res.* 4(10):2483-90; Peterson *et al.*, (1999) *Bioconjug. Chem.* 10(4):553-7; and Zimmerman *et al.*, (1999) *Nucl. Med. Biol.* 26(8):943-50, each incorporated by reference in their entireties.

Techniques for conjugating therapeutic moieties to antibodies are well known, see, *e.g.*, Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies 84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16 (Academic Press 1985), and Thorpe *et al.*, (1982) *Immunol. Rev.* 62:119-58.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not

limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Antibody Combinations

5 An another aspect, the invention pertains to HER3 antibodies, or fragments thereof of the invention used with other therapeutic agents such as another antibodies, small molecule inhibitors, mTOR inhibitors or PI3Kinase inhibitors. Examples include, but are not limited to, the following:

10 *HER1 inhibitors:* The HER3 antibodies or fragments thereof can be used with HER1 inhibitors which include, but are not limited to, Matuzumab (EMD72000), Erbitux®/Cetuximab (Imclone), Vectibix® /Panitumumab (Amgen), mAb 806, and Nimotuzumab (TheraCIM), Iressa® /Gefitinib (Astrazeneca); CI-1033 (PD183805) (Pfizer), Lapatinib (GW-572016) (GlaxoSmithKline), Tykerb® /Lapatinib Ditosylate (SmithKlineBeecham), Tarceva® / Erlotinib HCL (OSI-774) (OSI Pharma) , and PKI-166
15 (Novartis), and N-[4-[(3-Chloro-4-fluorophenyl)amino]-7-[[3"S"]-tetrahydro-3-furanyl]oxy]-6-quinazoliny]-4(dimethylamino)-2-butenamide, sold under the tradename Tovok® by Boehringer Ingelheim).

20 *HER2 inhibitors:* The HER3 antibodies or fragments thereof can be used with HER2 inhibitors which include, but are not limited to, Pertuzumab (sold under the trademark Omnitarg®, by Genentech), Trastuzumab (sold under the trademark Herceptin® by Genentech/Roche), MM-111, neratinib (also known as HKI-272, (2E)-N-[4-[[3-chloro-4-[(pyridin-2-yl)methoxy]phenyl]amino]-3-cyano-7-ethoxyquinolin-6-yl]-4-(dimethylamino)but-2-enamide, and described PCT Publication No. WO 05/028443), lapatinib or lapatinib ditosylate (sold under the trademark Tykerb® by GlaxoSmithKline.

25 *HER3 inhibitors:* The HER3 antibodies or fragments thereof can be used with HER3 inhibitors which include, but are not limited to, MM-121, MM-111, IB4C3, 2DID12 (U3 Pharma AG), AMG888 (Amgen), AV-203(Aveo), MEHD7945A (Genentech), and small molecules that inhibit HER3.

30 *HER4 inhibitors:* The HER3 antibodies or fragments thereof can be used with HER4 inhibitors.

PI3K inhibitors: The HER3 antibodies or fragments thereof can be used with PI3 kinase inhibitors which include, but are not limited to, 4-[2-(1H-Indazol-4-yl)-6-[[4-(methylsulfonyl)piperazin-1-yl]methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine (also known as GDC 0941 and described in PCT Publication Nos. WO 09/036082 and WO 09/055730), 2-Methyl-2-[4-[3-methyl-2-oxo-8-(quinolin-3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile (also known as BEZ 235 or NVP-BEZ 235, and described in PCT Publication No. WO 06/122806), BMK120 and BYL719.

mTOR inhibitors: The HER3 antibodies or fragments thereof can be used with mTOR inhibitors which include, but are not limited to, Temsirolimus (sold under the tradename Torisel® by Pfizer), ridaforolimus (formally known as deferolimus, (1R,2R,4S)-4-[(2R)-2-[(1R,9S,12S,15R,16E,18R,19R,21R, 23S,24E,26E,28Z,30S,32S,35R)-1,18-dihydroxy-19,30-dimethoxy-15,17,21,23, 29,35-hexamethyl-2,3,10,14,20-pentaoxo-11,36-dioxa-4-azatricyclo[30.3.1.04,9] hexatriaconta-16,24,26,28-tetraen-12-yl]propyl]-2-methoxycyclohexyl dimethylphosphinate, also known as Deforolimus, AP23573 and MK8669 (Ariad Pharm.), and described in PCT Publication No. WO 03/064383), everolimus (RAD001) (sold under the tradename Afinitor® by Novartis), One or more therapeutic agents may be administered either simultaneously or before or after administration of a HER3 antibody or fragment thereof of the present invention.

Methods of Producing Antibodies of the Invention

(i) Nucleic Acids Encoding the Antibodies

The invention provides substantially purified nucleic acid molecules which encode polypeptides comprising segments or domains of the HER3-binding antibody chains described above. Some of the nucleic acids of the invention comprise the nucleotide sequence encoding the HER3 antibody heavy chain variable region, and/or the nucleotide sequence encoding the light chain variable region. In a specific embodiment, the nucleic acid molecules are those identified in Table 1. Some other nucleic acid molecules of the invention comprise nucleotide sequences that are substantially identical (*e.g.*, at least 65, 80%, 95%, or 99%) to the nucleotide sequences of those identified in Table 1. When expressed from appropriate expression vectors, polypeptides encoded by these polynucleotides are capable of exhibiting HER3 antigen binding capacity.

Also provided in the invention are polynucleotides which encode at least one CDR region and usually all three CDR regions from the heavy or light chain of the HER3-binding antibody set

forth above. Some other polynucleotides encode all or substantially all of the variable region sequence of the heavy chain and/or the light chain of the HER3-binding antibody set forth above. Because of the degeneracy of the code, a variety of nucleic acid sequences will encode each of the immunoglobulin amino acid sequences.

- 5 The nucleic acid molecules of the invention can encode both a variable region and a constant region of the antibody. Some of nucleic acid sequences of the invention comprise nucleotides encoding a mature heavy chain variable region sequence that is substantially identical (*e.g.*, at least 80%, 90%, or 99%) to the mature heavy chain variable region sequence of a HER3 antibody set forth in Table 1. Some other nucleic acid sequences comprising nucleotide
10 encoding a mature light chain variable region sequence that is substantially identical (*e.g.*, at least 80%, 90%, or 99%) to the mature light chain variable region sequence of a HER3 antibody set forth in Table 1.

The polynucleotide sequences can be produced by *de novo* solid-phase DNA synthesis or by PCR mutagenesis of an existing sequence (*e.g.*, sequences as described in the Examples
15 below) encoding an HER3-binding antibody or its binding fragment. Direct chemical synthesis of nucleic acids can be accomplished by methods known in the art, such as the phosphotriester method of Narang *et al.*, (1979) *Meth. Enzymol.* 68:90; the phosphodiester method of Brown *et al.*, (1979) *Meth. Enzymol.* 68:109; the diethylphosphoramidite method of Beaucage *et al.*, (1981) *Tetra. Lett.*, 22:1859; and the solid support method of U.S. Patent
20 No. 4,458,066. Introducing mutations to a polynucleotide sequence by PCR can be performed as described in, *e.g.*, *PCR Technology: Principles and Applications for DNA Amplification*, H.A. Erlich (Ed.), Freeman Press, NY, NY, 1992; *PCR Protocols: A Guide to Methods and Applications*, Innis *et al.* (Ed.), Academic Press, San Diego, CA, 1990; Mattila *et al.*, (1991) *Nucleic Acids Res.* 19:967; and Eckert *et al.*, (1991) *PCR Methods and*
25 *Applications* 1:17.

Also provided in the invention are expression vectors and host cells for producing the HER3-binding antibodies described above. Various expression vectors can be employed to express the polynucleotides encoding the HER3-binding antibody chains or binding fragments. Both
30 viral-based and nonviral expression vectors can be used to produce the antibodies in a mammalian host cell. Nonviral vectors and systems include plasmids, episomal vectors, typically with an expression cassette for expressing a protein or RNA, and human artificial chromosomes (see, *e.g.*, Harrington *et al.*, (1997) *Nat Genet* 15:345). For example, nonviral vectors useful for expression of the HER3-binding polynucleotides and polypeptides in

mammalian (*e.g.*, human) cells include pThioHis A, B & C, pcDNA3.1/His, pEBVHis A, B & C, (Invitrogen, San Diego, CA), MPSV vectors, and numerous other vectors known in the art for expressing other proteins. Useful viral vectors include vectors based on retroviruses, adenoviruses, adenoassociated viruses, herpes viruses, vectors based on SV40, papilloma virus, HBP Epstein Barr virus, vaccinia virus vectors and Semliki Forest virus (SFV). See, Brent *et al.*, (1995) *supra*; Smith, *Annu. Rev. Microbiol.* 49:807; and Rosenfeld *et al.*, (1992) *Cell* 68:143.

The choice of expression vector depends on the intended host cells in which the vector is to be expressed. Typically, the expression vectors contain a promoter and other regulatory sequences (*e.g.*, enhancers) that are operably linked to the polynucleotides encoding an HER3-binding antibody chain or fragment. In some embodiments, an inducible promoter is employed to prevent expression of inserted sequences except under inducing conditions. Inducible promoters include, *e.g.*, arabinose, lacZ, metallothionein promoter or a heat shock promoter. Cultures of transformed organisms can be expanded under noninducing conditions without biasing the population for coding sequences whose expression products are better tolerated by the host cells. In addition to promoters, other regulatory elements may also be required or desired for efficient expression of an HER3-binding antibody chain or fragment. These elements typically include an ATG initiation codon and adjacent ribosome binding site or other sequences. In addition, the efficiency of expression may be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, *e.g.*, Scharf *et al.*, (1994) *Results Probl. Cell Differ.* 20:125; and Bittner *et al.*, (1987) *Meth. Enzymol.*, 153:516). For example, the SV40 enhancer or CMV enhancer may be used to increase expression in mammalian host cells.

The expression vectors may also provide a secretion signal sequence position to form a fusion protein with polypeptides encoded by inserted HER3-binding antibody sequences. More often, the inserted HER3-binding antibody sequences are linked to a signal sequences before inclusion in the vector. Vectors to be used to receive sequences encoding HER3-binding antibody light and heavy chain variable domains sometimes also encode constant regions or parts thereof. Such vectors allow expression of the variable regions as fusion proteins with the constant regions thereby leading to production of intact antibodies or fragments thereof. Typically, such constant regions are human.

The host cells for harboring and expressing the HER3-binding antibody chains can be either prokaryotic or eukaryotic. *E. coli* is one prokaryotic host useful for cloning and expressing

the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which typically contain expression control sequences compatible with the host cell
5 (*e.g.*, an origin of replication). In addition, any number of a variety of well-known promoters will be present, such as the lactose promoter system, a tryptophan (*trp*) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters typically control expression, optionally with an operator sequence, and have ribosome binding site sequences and the like, for initiating and completing transcription and translation. Other
10 microbes, such as yeast, can also be employed to express HER3-binding polypeptides of the invention. Insect cells in combination with baculovirus vectors can also be used.

In some preferred embodiments, mammalian host cells are used to express and produce the HER3-binding polypeptides of the present invention. For example, they can be either a hybridoma cell line expressing endogenous immunoglobulin genes (*e.g.*, the 1D6.C9
15 myeloma hybridoma clone as described in the Examples) or a mammalian cell line harboring an exogenous expression vector (*e.g.*, the SP2/0 myeloma cells exemplified below). These include any normal mortal or normal or abnormal immortal animal or human cell. For example, a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed including the CHO cell lines, various Cos cell lines, HeLa cells,
20 myeloma cell lines, transformed B-cells and hybridomas. The use of mammalian tissue cell culture to express polypeptides is discussed generally in, *e.g.*, Winnacker, FROM GENES TO CLONES, VCH Publishers, N.Y., N.Y., 1987. Expression vectors for mammalian host cells can include expression control sequences, such as an origin of replication, a promoter, and an enhancer (see, *e.g.*, Queen *et al.*, (1986) *Immunol. Rev.* 89:49-68), and necessary processing
25 information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. These expression vectors usually contain promoters derived from mammalian genes or from mammalian viruses. Suitable promoters may be constitutive, cell type-specific, stage-specific, and/or modulatable or regulatable. Useful promoters include, but are not limited to, the metallothionein promoter, the constitutive adenovirus major late promoter, the dexamethasone-inducible MMTV promoter, the SV40
30 promoter, the MRP polIII promoter, the constitutive MPSV promoter, the tetracycline-inducible CMV promoter (such as the human immediate-early CMV promoter), the constitutive CMV promoter, and promoter-enhancer combinations known in the art.

Methods for introducing expression vectors containing the polynucleotide sequences of interest vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts. (See generally Sambrook, *et al.*, supra). Other methods include, *e.g.*, electroporation, calcium phosphate treatment, liposome-mediated transformation, injection and microinjection, ballistic methods, virosomes, immunoliposomes, polycation:nucleic acid conjugates, naked DNA, artificial virions, fusion to the herpes virus structural protein VP22 (Elliot and O'Hare, (1997) Cell 88:223), agent-enhanced uptake of DNA, and *ex vivo* transduction. For long-term, high-yield production of recombinant proteins, stable expression will often be desired. For example, cell lines which stably express HER3-binding antibody chains or binding fragments can be prepared using expression vectors of the invention which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth of cells which successfully express the introduced sequences in selective media. Resistant, stably transfected cells can be proliferated using tissue culture techniques appropriate to the cell type.

(ii) Generation of monoclonal antibodies of the invention

Monoclonal antibodies (mAbs) can be produced by a variety of techniques, including conventional monoclonal antibody methodology *e.g.*, the standard somatic cell hybridization technique of Kohler and Milstein, (1975) Nature 256:495. Many techniques for producing monoclonal antibody can be employed *e.g.*, viral or oncogenic transformation of B lymphocytes.

An animal system for preparing hybridomas is the murine system. Hybridoma production in the mouse is a well established procedure. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art. Fusion partners (*e.g.*, murine myeloma cells) and fusion procedures are also known.

Chimeric or humanized antibodies of the present invention can be prepared based on the sequence of a murine monoclonal antibody prepared as described above. DNA encoding the heavy and light chain immunoglobulins can be obtained from the murine hybridoma of interest and engineered to contain non-murine (*e.g.*, human) immunoglobulin sequences

using standard molecular biology techniques. For example, to create a chimeric antibody, the murine variable regions can be linked to human constant regions using methods known in the art (see *e.g.*, U.S. Patent No. 4,816,567 to Cabilly *et al.*). To create a humanized antibody, the murine CDR regions can be inserted into a human framework using methods known in the art.

5 See *e.g.*, U.S. Patent No. 5225539 to Winter, and U.S. Patent Nos. 5530101; 5585089; 5693762 and 6180370 to Queen *et al.*

In a certain embodiment, the antibodies of the invention are human monoclonal antibodies. Such human monoclonal antibodies directed against HER3 can be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse
10 system. These transgenic and transchromosomal mice include mice referred to herein as HuMAb mice and KM mice, respectively, and are collectively referred to herein as "human Ig mice."

The HuMAb mouse[®] (Medarex, Inc.) contains human immunoglobulin gene miniloci that encode un-rearranged human heavy (μ and γ) and κ light chain immunoglobulin sequences,
15 together with targeted mutations that inactivate the endogenous μ and κ chain loci (see *e.g.*, Lonberg *et al.*, (1994) *Nature* 368(6474): 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or κ , and in response to immunization, the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgG κ monoclonal (Lonberg *et al.*, (1994) *supra*; reviewed in Lonberg, (1994)
20 *Handbook of Experimental Pharmacology* 113:49-101; Lonberg and Huszar, (1995) *Intern. Rev. Immunol.* 13:65-93, and Harding and Lonberg, (1995) *Ann. N. Y. Acad. Sci.* 764:536-546). The preparation and use of HuMAb mice, and the genomic modifications carried by such mice, is further described in Taylor *et al.*, (1992) *Nucleic Acids Research* 20:6287-6295; Chen *et al.*, (1993) *International Immunology* 5:647-656; Tuailleon *et al.*, (1993) *Proc. Natl.*
25 *Acad. Sci. USA* 94:3720-3724; Choi *et al.*, (1993) *Nature Genetics* 4:117-123; Chen *et al.*, (1993) *EMBO J.* 12:821-830; Tuailleon *et al.*, (1994) *J. Immunol.* 152:2912-2920; Taylor *et al.*, (1994) *International Immunology* 579-591; and Fishwild *et al.*, (1996) *Nature Biotechnology* 14:845-851, the contents of all of which are hereby specifically incorporated by reference in their entirety. See further, U.S. Patent Nos. 5,545,806; 5,569,825; 5,625,126;
30 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay; U.S. Patent No. 5,545,807 to Surani *et al.*; PCT Publication Nos. WO 92103918, WO 93/12227, WO 94/25585, WO 97113852, WO 98/24884 and WO 99/45962, all to Lonberg and Kay; and PCT Publication No. WO 01/14424 to Korman *et al.*

In another embodiment, human antibodies of the invention can be raised using a mouse that carries human immunoglobulin sequences on transgenes and transchromosomes such as a mouse that carries a human heavy chain transgene and a human light chain transchromosome. Such mice, referred to herein as "KM mice", are described in detail in PCT Publication WO
5 02/43478 to Ishida *et al.*

Still further, alternative transgenic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise HER3-binding antibodies of the invention. For example, an alternative transgenic system referred to as the Xenomouse (Abgenix, Inc.) can be used. Such mice are described in, *e.g.*, U.S. Patent Nos. 5,939,598; 6,075,181; 6,114,598;
10 6, 150,584 and 6,162,963 to Kucherlapati *et al.*

Moreover, alternative transchromosomal animal systems expressing human immunoglobulin genes are available in the art and can be used to raise HER3-binding antibodies of the invention. For example, mice carrying both a human heavy chain transchromosome and a human light chain transchromosome, referred to as "TC mice" can be used; such mice are
15 described in Tomizuka *et al.*, (2000) Proc. Natl. Acad. Sci. USA 97:722-727. Furthermore, cows carrying human heavy and light chain transchromosomes have been described in the art (Kuroiwa *et al.*, (2002) Nature Biotechnology 20:889-894) and can be used to raise HER3-binding antibodies of the invention.

Human monoclonal antibodies of the invention can also be prepared using phage display
20 methods for screening libraries of human immunoglobulin genes. Such phage display methods for isolating human antibodies are established in the art or described in the examples below. See for example: U.S. Patent Nos. 5,223,409; 5,403,484; and 5,571,698 to Ladner *et al.*; U.S. Patent Nos. 5,427,908 and 5,580,717 to Dower *et al.*; U.S. Patent Nos. 5,969,108 and 6,172,197 to McCafferty *et al.*; and U.S. Patent Nos. 5,885,793; 6,521,404; 6,544,731;
25 6,555,313; 6,582,915 and 6,593,081 to Griffiths *et al.*

Human monoclonal antibodies of the invention can also be prepared using SCID mice into which human immune cells have been reconstituted such that a human antibody response can be generated upon immunization. Such mice are described in, for example, U.S. Patent Nos. 5,476,996 and 5,698,767 to Wilson *et al.*

30 (iii) Framework or Fc engineering

Engineered antibodies of the invention include those in which modifications have been made to framework residues within VH and/or VL, *e.g.* to improve the properties of the antibody. Typically such framework modifications are made to decrease the immunogenicity of the antibody. For example, one approach is to "backmutate" one or more framework residues to the corresponding germline sequence. More specifically, an antibody that has undergone somatic mutation may contain framework residues that differ from the germline sequence from which the antibody is derived. Such residues can be identified by comparing the antibody framework sequences to the germline sequences from which the antibody is derived. To return the framework region sequences to their germline configuration, the somatic mutations can be "backmutated" to the germline sequence by, for example, site-directed mutagenesis. Such "backmutated" antibodies are also intended to be encompassed by the invention.

Another type of framework modification involves mutating one or more residues within the framework region, or even within one or more CDR regions, to remove T cell -epitopes to thereby reduce the potential immunogenicity of the antibody. This approach is also referred to as "deimmunization" and is described in further detail in U.S. Patent Publication No. 20030153043 by Carr *et al.*

In addition or alternative to modifications made within the framework or CDR regions, antibodies of the invention may be engineered to include modifications within the Fc region, typically to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding, and/or antigen-dependent cellular cytotoxicity. Furthermore, an antibody of the invention may be chemically modified (*e.g.*, one or more chemical moieties can be attached to the antibody) or be modified to alter its glycosylation, again to alter one or more functional properties of the antibody. Each of these embodiments is described in further detail below. The numbering of residues in the Fc region is that of the EU index of Kabat.

In one embodiment, the hinge region of CH1 is modified such that the number of cysteine residues in the hinge region is altered, *e.g.*, increased or decreased. This approach is described further in U.S. Patent No. 5,677,425 by Bodmer *et al.* The number of cysteine residues in the hinge region of CH1 is altered to, for example, facilitate assembly of the light and heavy chains or to increase or decrease the stability of the antibody.

In another embodiment, the Fc hinge region of an antibody is mutated to decrease the biological half-life of the antibody. More specifically, one or more amino acid mutations are introduced into the CH2-CH3 domain interface region of the Fc-hinge fragment such that the antibody has impaired Staphylococcal protein A (SpA) binding relative to native Fc-hinge domain SpA binding. This approach is described in further detail in U.S. Patent No. 6,165,745 by Ward *et al.*

In yet other embodiments, the Fc region is altered by replacing at least one amino acid residue with a different amino acid residue to alter the effector functions of the antibody. For example, one or more amino acids can be replaced with a different amino acid residue such that the antibody has an altered affinity for an effector ligand but retains the antigen-binding ability of the parent antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Patent Nos. 5,624,821 and 5,648,260, both by Winter *et al.*

In another embodiment, one or more amino acids selected from amino acid residues can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Patent Nos. 6,194,551 by Idusogie *et al.*

In another embodiment, one or more amino acid residues are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in PCT Publication WO 94/29351 by Bodmer *et al.*

In yet another embodiment, the Fc region is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or to increase the affinity of the antibody for an Fc γ receptor by modifying one or more amino acids. This approach is described further in PCT Publication WO 00/42072 by Presta. Moreover, the binding sites on human IgG1 for Fc γ R1, Fc γ R2, Fc γ R3 and FcRn have been mapped and variants with improved binding have been described (see Shields *et al.*, (2001) J. Biol. Chem. 276:6591-6604).

In still another embodiment, the glycosylation of an antibody is modified. For example, an aglycosylated antibody can be made (*i.e.*, the antibody lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the antibody for 'antigen'. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. For example, one or more amino acid

substitutions can be made that result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the antibody for antigen. Such an approach is described in further detail in U.S. Patent Nos. 5,714,350 and 6,350,861 by Co *et al.*

5 Additionally or alternatively, an antibody can be made that has an altered type of glycosylation, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNac structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the
10 antibody in a host cell with altered glycosylation machinery. Cells with altered glycosylation machinery have been described in the art and can be used as host cells in which to express recombinant antibodies of the invention to thereby produce an antibody with altered glycosylation. For example, EP 1,176,195 by Hang *et al.* describes a cell line with a functionally disrupted FUT8 gene, which encodes a fucosyl transferase, such that antibodies
15 expressed in such a cell line exhibit hypofucosylation. PCT Publication WO 03/035835 by Presta describes a variant CHO cell line, Lecl3 cells, with reduced ability to attach fucose to Asn(297)-linked carbohydrates, also resulting in hypofucosylation of antibodies expressed in that host cell (see also Shields *et al.*, (2002) J. Biol. Chem. 277:26733-26740). PCT
Publication WO 99/54342 by Umana *et al.* describes cell lines engineered to express
20 glycoprotein-modifying glycosyl transferases (*e.g.*, beta(1,4)-N acetylglucosaminyltransferase III (GnTIII)) such that antibodies expressed in the engineered cell lines exhibit increased bisecting GlcNac structures which results in increased ADCC activity of the antibodies (see also Umana *et al.*, (1999) Nat. Biotech. 17:176-180).

In another embodiment, the antibody is modified to increase its biological half-life. Various
25 approaches are possible. For example, one or more of the following mutations can be introduced: T252L, T254S, T256F, as described in U.S. Patent No. 6,277,375 to Ward. Alternatively, to increase the biological half life, the antibody can be altered within the CH1 or CL region to contain a salvage receptor binding epitope taken from two loops of a CH2 domain of an Fc region of an IgG, as described in U.S. Patent Nos. 5,869,046 and 6,121,022
30 by Presta *et al.*

(iv) Methods of Engineering Altered Antibodies

As discussed above, the HER3-binding antibodies having VH and VL sequences or full length heavy and light chain sequences shown herein can be used to create new HER3-binding antibodies by modifying full length heavy chain and/or light chain sequences, VH and/or VL sequences, or the constant region(s) attached thereto. Thus, in another aspect of the invention, the structural features of a HER3-binding antibody of the invention are used to create structurally related HER3-binding antibodies that retain at least one functional property of the antibodies of the invention, such as binding to human HER3 and also inhibiting one or more functional properties of HER3. For example, one or more CDR regions of the antibodies of the present invention, or mutations thereof, can be combined recombinantly with known framework regions and/or other CDRs to create additional, recombinantly-engineered, HER3-binding antibodies of the invention, as discussed above. Other types of modifications include those described in the previous section. The starting material for the engineering method is one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. To create the engineered antibody, it is not necessary to actually prepare (*i.e.*, express as a protein) an antibody having one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. Rather, the information contained in the sequence(s) is used as the starting material to create a "second generation" sequence(s) derived from the original sequence(s) and then the "second generation" sequence(s) is prepared and expressed as a protein.

Accordingly, in another embodiment, the invention provides a method for preparing a HER3-binding antibody consisting of: a heavy chain variable region antibody sequence having a CDR1 sequence selected from the group consisting of SEQ ID NOs: 2, 8, 20, 26, 38, 44, 56, 62, 74, 80, 92, 98, 110, 116, 128, 134, 146, 152, 164, 170, 182, 188, 200, 206, 218, 224, 236, 242, 254, 260, 272, 278, 290, 296, 308, 314, 326, 332, 344, 350, 362, and 368; a CDR2 sequence selected from the group consisting of SEQ ID NOs: 3, 9, 21, 27, 39, 45, 57, 63, 75, 81, 93, 99, 111, 117, 129, 135, 147, 153, 165, 171, 183, 189, 201, 207, 219, 225, 237, 243, 255, 261, 273, 279, 291, 297, 309, 315, 327, 333, 345, 351, 363, and 369; and/or a CDR3 sequence selected from the group consisting of SEQ ID NOs: 4, 10, 22, 28, 40, 46, 58, 64, 75, 82, 94, 100, 112, 118, 130, 136, 148, 154, 166, 172, 184, 190, 202, 208, 220, 226, 238, 244, 256, 262, 274, 280, 292, 298, 310, 316, 328, 334, 346, 352, 364, and 370; and a light chain variable region antibody sequence having a CDR1 sequence selected from the group consisting of SEQ ID NOs: 5, 11, 23, 29, 41, 47, 59, 65, 77, 83, 95, 101, 113, 119, 131, 137,

149, 155, 167, 173, 185, 191, 203, 209, 221, 227, 239, 245, 257, 263, 275, 281, 293, 299, 311, 317, 329, 335, 347, 353, 365, and 371; a CDR2 sequence selected from the group consisting of SEQ ID NOs: 6, 12, 24, 30, 42, 48, 60, 66, 78, 84, 96, 102, 114, 120, 132, 138, 150, 156, 168, 174, 186, 192, 204, 210, 222, 228, 240, 246, 258, 264, 276, 282, 294, 300, 312, 318, 330, 336, 348, 354, 366, and 372; and/or a CDR3 sequence selected from the group consisting of
5 SEQ ID NOs: 7, 13, 25, 31, 43, 49, 61, 67, 79, 85, 97, 103, 115, 121, 133, 139, 151, 157, 169, 175, 187, 193, 205, 211, 223, 229, 241, 247, 259, 265, 277, 283, 295, 301, 313, 319, 331, 337, 349, 355, 367, and 373; altering at least one amino acid residue within the heavy chain variable region antibody sequence and/or the light chain variable region antibody sequence to
10 create at least one altered antibody sequence; and expressing the altered antibody sequence as a protein. The altered antibody sequence can also be prepared by screening antibody libraries having fixed CDR3 sequences or minimal essential binding determinants as described in US20050255552 and diversity on CDR1 and CDR2 sequences. The screening can be performed according to any screening technology appropriate for screening antibodies from
15 antibody libraries, such as phage display technology.

Standard molecular biology techniques can be used to prepare and express the altered antibody sequence. The antibody encoded by the altered antibody sequence(s) is one that retains one, some or all of the functional properties of the HER3-binding antibodies described herein, which functional properties include, but are not limited to, specifically binding to
20 human and/or cynomologus HER3; the antibody binds to HER3 and neutralizes HER3 biological activity by inhibiting the HER signaling activity in a phospho-HER assay.

The functional properties of the altered antibodies can be assessed using standard assays available in the art and/or described herein, such as those set forth in the Examples (*e.g.*, ELISAs).

25 In certain embodiments of the methods of engineering antibodies of the invention, mutations can be introduced randomly or selectively along all or part of an HER3-binding antibody coding sequence and the resulting modified HER3-binding antibodies can be screened for binding activity and/or other functional properties as described herein. Mutational methods have been described in the art. For example, PCT Publication WO 02/092780 by Short
30 describes methods for creating and screening antibody mutations using saturation mutagenesis, synthetic ligation assembly, or a combination thereof. Alternatively, PCT Publication WO 03/074679 by Lazar *et al.* describes methods of using computational screening methods to optimize physiochemical properties of antibodies.

Characterization of the Antibodies of the Invention

The antibodies of the invention can be characterized by various functional assays. For example, they can be characterized by their ability to neutralize biological activity by inhibiting HER signaling in a phospho-HER assay as described herein, their affinity to a
5 HER3 protein (*e.g.*, human and/or cynomologus HER3), the epitope binning, their resistance to proteolysis, and their ability to block HER3 downstream signaling. Various methods can be used to measure HER3 -mediated signaling. For example, the HER signaling pathway can be monitored by (i) measurement of phospho-HER3; (ii) measurement of phosphorylation of HER3 or other downstream signaling proteins (*e.g.* Akt), (iii) ligand blocking assays as
10 described herein, (iv) heterodimer formation, (v) HER3 dependent gene expression signature, (vi) receptor internalization, and (vii) HER3 driven cell phenotypes (*e.g.* proliferation).

The ability of an antibody to bind to HER3 can be detected by labelling the antibody of interest directly, or the antibody may be unlabelled and binding detected indirectly using various sandwich assay formats known in the art.

15 In some embodiments, the HER3-binding antibodies of the invention block or compete with binding of a reference HER3-binding antibody to a HER3 polypeptide or protein. These can be fully human HER3-binding antibodies described above. They can also be other mouse, chimeric or humanized HER3-binding antibodies which bind to the same epitope as the reference antibody. The capacity to block or compete with the reference antibody binding
20 indicates that a HER3-binding antibody under test binds to the same or similar epitope as that defined by the reference antibody, or to an epitope which is sufficiently proximal to the epitope bound by the reference HER3-binding antibody. Such antibodies are especially likely to share the advantageous properties identified for the reference antibody. The capacity to block or compete with the reference antibody may be determined by, *e.g.*, a competition
25 binding assay. With a competition binding assay, the antibody under test is examined for ability to inhibit specific binding of the reference antibody to a common antigen, such as a HER3 polypeptide or protein. A test antibody competes with the reference antibody for specific binding to the antigen if an excess of the test antibody substantially inhibits binding of the reference antibody. Substantial inhibition means that the test antibody reduces specific
30 binding of the reference antibody usually by at least 10%, 25%, 50%, 75%, or 90%.

There are a number of known competition binding assays that can be used to assess competition of a HER3-binding antibody with the reference HER3-binding antibody for

binding to a HER3 protein. These include, *e.g.*, solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahli *et al.*, (1983) *Methods in Enzymology* 9:242-253); solid phase direct biotin-avidin EIA (see Kirkland *et al.*, (1986) *J. Immunol.* 137:3614-3619);
5 solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow & Lane, *supra*); solid phase direct label RIA using I-125 label (see Morel *et al.*, (1988) *Molec. Immunol.* 25:7-15); solid phase direct biotin-avidin EIA (Cheung *et al.*, (1990) *Virology* 176:546-552); and direct labeled RIA (Moldenhauer *et al.*, (1990) *Scand. J. Immunol.* 32:77-82). Typically, such an assay involves the use of purified antigen bound to a solid surface or
10 cells bearing either of these, an unlabelled test HER3-binding antibody and a labelled reference antibody. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test antibody. Usually the test antibody is present in excess. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and
15 antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur.

To determine if the selected HER3-binding monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (*e.g.*, reagents from Pierce, Rockford, IL). Competition studies using unlabeled monoclonal antibodies and
20 biotinylated monoclonal antibodies can be performed using a HER3 polypeptide coated-ELISA plates. Biotinylated MAb binding can be detected with a strep-avidin-alkaline phosphatase probe. To determine the isotype of a purified HER3-binding antibody, isotype ELISAs can be performed. For example, wells of microtiter plates can be coated with 1 $\mu\text{g/ml}$ of anti-human IgG overnight at 4°C. After blocking with 1% BSA, the plates are
25 reacted with 1 $\mu\text{g/ml}$ or less of the monoclonal HER3-binding antibody or purified isotype controls, at ambient temperature for one to two hours. The wells can then be reacted with either human IgG1 or human IgM-specific alkaline phosphatase-conjugated probes. Plates are then developed and analyzed so that the isotype of the purified antibody can be determined.

To demonstrate binding of monoclonal HER3-binding antibodies to live cells expressing a
30 HER3 polypeptide, flow cytometry can be used. Briefly, cell lines expressing HER3 (grown under standard growth conditions) can be mixed with various concentrations of a HER3-binding antibody in PBS containing 0.1% BSA and 10% fetal calf serum, and incubated at 4°C for 1 hour. After washing, the cells are reacted with Fluorescein-labeled anti-human IgG

antibody under the same conditions as the primary antibody staining. The samples can be analyzed by FACScan instrument using light and side scatter properties to gate on single cells. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells can be stained exactly as described above and examined by
5 fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

HER3-binding antibodies of the invention can be further tested for reactivity with a HER3 polypeptide or antigenic fragment by Western blotting. Briefly, purified HER3 polypeptides or fusion proteins, or cell extracts from cells expressing HER3 can be prepared and subjected
10 to sodium dodecyl sulfate polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens are transferred to nitrocellulose membranes, blocked with 10% fetal calf serum, and probed with the monoclonal antibodies to be tested. Human IgG binding can be detected using anti-human IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, MO).

15 A number of readouts can be used to assess the efficacy, and specificity, of HER3 antibodies in cell-based assays of ligand-induced heterodimer formation. Activity can be assessed by one or more of the following:

(i) Inhibition of ligand-induced heterodimerisation of HER2 with other EGF family members in a target cell line, for example MCF-7 breast cancer cells. Immunoprecipitation of HER2
20 complexes from cell lysates can be performed with a receptor-specific antibody, and the absence/presence of other EGF receptors and their biologically relevant ligands within the complex can be analysed following electrophoresis/Western transfer by probing with antibodies to other EGF receptors.

25 (ii) Inhibition of the activation of signaling pathways by ligand-activated heterodimers. Association with HER3 appears key for other members of the EGF family of receptors to elicit maximal cellular response following ligand binding. In the case of the kinase-defective HER3, HER2 provides a functional tyrosine kinase domain to enable signaling to occur following binding of growth factor ligands. Thus, cells co-expressing HER2 and HER3 can be
30 treated with ligand, for example heregulin, in the absence and presence of inhibitor and the effect on HER3 tyrosine phosphorylation monitored by a number of ways including immunoprecipitation of HER3 from treated cell lysates and subsequent Western blotting using anti-phosphotyrosine antibodies (see Agus *op. cit.* for details). Alternatively, a high-

throughput assay can be developed by trapping HER3 from solubilized lysates onto the wells of a 96-well plate coated with an anti-HER3 receptor antibody, and the level of tyrosine phosphorylation measured using, for example, europium-labelled anti-phosphotyrosine antibodies, as embodied by Waddleton *et al.*, (2002) *Anal. Biochem.* 309:150-157.

5

In a broader extension of this approach, effector molecules known to be activated downstream of activated receptor heterodimers, such as mitogen-activated protein kinases (MAPK) and Akt, may be analysed directly, by immunoprecipitation from treated lysates and blotting with antibodies that detect the activated forms of these proteins, or by analysing the ability of these proteins to modify/activate specific substrates.

10

(iii) Inhibition of ligand-induced cellular proliferation. A variety of cell lines are known to co-express combinations of ErbB receptors, for example many breast and prostate cancer cell lines. Assays may be performed in 24/48/96-well formats with the readout based around DNA synthesis (tritiated thymidine incorporation), increase in cell number (crystal violet staining) etc.

15

A number of readouts can be used to assess the efficacy, and specificity, of HER3 antibodies in cell-based assays of ligand-independent homo- and heterodimer formation. For example, HER2 overexpression triggers ligand-independent activation of the kinase domain as a result of spontaneous dimer formation. Over expressed HER2 generates either homo- or heterodimers with other HER molecules such as HER1, HER3 and HER4.

20

Ability of antibodies or fragments thereof to block *in vivo* growth of tumour xenografts of human tumour cell lines whose tumorigenic phenotype is known to be at least partly dependent on ligand activation of HER3 heterodimer cell signaling e.g. BxPC3 pancreatic cancer cells etc. This can be assessed in immunocompromised mice either alone or in combination with an appropriate cytotoxic agent for the cell line in question. Examples of functional assays are also described in the Example section below.

25

30 **Prophylactic and Therapeutic Uses**

The present invention provides methods of treating a disease or disorder associated with the HER3 signaling pathway by administering to a subject in need thereof an effective amount of

the antibodies of the invention. In a specific embodiment, the present invention provides a method of treating or preventing cancers (e.g., breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors, schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer and melanoma) by administering to a subject in need thereof an effective amount of the antibodies of the invention. In some embodiments, the present invention provides methods of treating or preventing cancers associated with a HER signaling pathway by administering to a subject in need thereof an effective amount of the antibodies of the invention.

In a specific embodiment, the present invention provides methods of treating cancers associated with a HER signaling pathway that include, but are not limited to breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and melanoma.

HER3 antibodies can also be used to treat or prevent other disorders associated with aberrant or defective HER signaling, including but are not limited to respiratory diseases, osteoporosis, osteoarthritis, polycystic kidney disease, diabetes, schizophrenia, vascular disease, cardiac disease, non-oncogenic proliferative diseases, fibrosis, and neurodegenerative diseases such as Alzheimer's disease.

Suitable agents for combination treatment with HER3-binding antibodies include standard of care agents known in the art that are able to modulate the ErbB signaling pathway. Suitable examples of standard of care agents for HER2 include, but are not limited to Herceptin and Tykerb. Suitable examples of standard of care agents for EGFR include, but are not limited to Iressa, Tarceva, Erbitux and Vectibix. Other agents that may be suitable for combination treatment with HER3-binding antibodies include, but are not limited to those that modulate receptor tyrosine kinases, G-protein coupled receptors, growth/ survival signal transduction pathways, nuclear hormone receptors, apoptotic pathways, cell cycle and angiogenesis.

Diagnostic Uses

In one aspect, the invention encompasses diagnostic assays for determining HER3 protein and/or nucleic acid expression as well as HER3 protein function, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) or from individual afflicted with cancer, or is at risk of developing cancer.

Diagnostic assays, such as competitive assays rely on the ability of a labelled analogue (the "tracer") to compete with the test sample analyte for a limited number of binding sites on a common binding partner. The binding partner generally is insolubilized before or after the competition and then the tracer and analyte bound to the binding partner are separated from the unbound tracer and analyte. This separation is accomplished by decanting (where the binding partner was preinsolubilized) or by centrifuging (where the binding partner was precipitated after the competitive reaction). The amount of test sample analyte is inversely proportional to the amount of bound tracer as measured by the amount of marker substance. Dose-response curves with known amounts of analyte are prepared and compared with the test results in order to quantitatively determine the amount of analyte present in the test sample. These assays are called ELISA systems when enzymes are used as the detectable markers. In an assay of this form, competitive binding between antibodies and HER3-binding antibodies results in the bound HER3 protein, preferably the HER3 epitopes of the invention, being a measure of antibodies in the serum sample, most particularly, neutralizing antibodies in the serum sample.

A significant advantage of the assay is that measurement is made of neutralizing antibodies directly (*i.e.*, those which interfere with binding of HER3 protein, specifically, epitopes). Such an assay, particularly in the form of an ELISA test has considerable applications in the clinical environment and in routine blood screening.

Another aspect of the invention provides methods for determining HER3 nucleic acid expression or HER3 protein activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (*e.g.*, drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (*e.g.*, the genotype of the individual examined to determine the ability of the individual to respond to a particular agent.)

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs) on the expression or activity of HER3 protein in clinical trials.

Pharmaceutical Compositions

To prepare pharmaceutical or sterile compositions including a HER3-binding antibodies (intact or binding fragments), the HER3-binding antibodies (intact or binding fragments) is mixed with a pharmaceutically acceptable carrier or excipient. The compositions can additionally contain one or more other therapeutic agents that are suitable for treating or preventing cancer (breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and melanoma).

Formulations of therapeutic and diagnostic agents can be prepared by mixing with physiologically acceptable carriers, excipients, or stabilizers in the form of, *e.g.*, lyophilized powders, slurries, aqueous solutions, lotions, or suspensions (see, *e.g.*, Hardman *et al.*, (2001) Goodman and Gilman's The Pharmacological Basis of Therapeutics, McGraw-Hill, New York, N.Y.; Gennaro (2000) Remington: The Science and Practice of Pharmacy, Lippincott, Williams, and Wilkins, New York, N.Y.; Avis, *et al.* (eds.) (1993) Pharmaceutical Dosage Forms: Parenteral Medications, Marcel Dekker, NY; Lieberman, *et al.* (eds.) (1990) Pharmaceutical Dosage Forms: Tablets, Marcel Dekker, NY; Lieberman, *et al.* (eds.) (1990) Pharmaceutical Dosage Forms: Disperse Systems, Marcel Dekker, NY; Weiner and Kotkoskie (2000) Excipient Toxicity and Safety, Marcel Dekker, Inc., New York, N.Y.).

Selecting an administration regimen for a therapeutic depends on several factors, including the serum or tissue turnover rate of the entity, the level of symptoms, the immunogenicity of the entity, and the accessibility of the target cells in the biological matrix. In certain embodiments, an administration regimen maximizes the amount of therapeutic delivered to the patient consistent with an acceptable level of side effects. Accordingly, the amount of biologic delivered depends in part on the particular entity and the severity of the condition being treated. Guidance in selecting appropriate doses of antibodies, cytokines, and small molecules are available (see, *e.g.*, Wawrzynczak (1996) Antibody Therapy, Bios Scientific Pub. Ltd, Oxfordshire, UK; Kresina (ed.) (1991) Monoclonal Antibodies, Cytokines and

Arthritis, Marcel Dekker, New York, N.Y.; Bach (ed.) (1993) *Monoclonal Antibodies and Peptide Therapy in Autoimmune Diseases*, Marcel Dekker, New York, N.Y.; Baert *et al.*, (2003) *New Engl. J. Med.* 348:601-608; Milgrom *et al.*, (1999) *New Engl. J. Med.* 341:1966-1973; Slamon *et al.*, (2001) *New Engl. J. Med.* 344:783-792; Beniaminovitz *et al.*, (2000) *New Engl. J. Med.* 342:613-619; Ghosh *et al.*, (2003) *New Engl. J. Med.* 348:24-32; Lipsky *et al.*, (2000) *New Engl. J. Med.* 343:1594-1602).

Determination of the appropriate dose is made by the clinician, e.g., using parameters or factors known or suspected in the art to affect treatment or predicted to affect treatment.

10 Generally, the dose begins with an amount somewhat less than the optimum dose and it is increased by small increments thereafter until the desired or optimum effect is achieved relative to any negative side effects. Important diagnostic measures include those of symptoms of, e.g., the inflammation or level of inflammatory cytokines produced.

15 Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular

20 compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors known in

25 the medical arts.

Compositions comprising antibodies or fragments thereof of the invention can be provided by continuous infusion, or by doses at intervals of, e.g., one day, one week, or 1-7 times per week. Doses may be provided intravenously, subcutaneously, topically, orally, nasally,

30 rectally, intramuscular, intracerebrally, or by inhalation. A specific dose protocol is one involving the maximal dose or dose frequency that avoids significant undesirable side effects. A total weekly dose may be at least 0.05 $\mu\text{g}/\text{kg}$ body weight, at least 0.2 $\mu\text{g}/\text{kg}$, at least 0.5 $\mu\text{g}/\text{kg}$, at least 1 $\mu\text{g}/\text{kg}$, at least 10 $\mu\text{g}/\text{kg}$, at least 100 $\mu\text{g}/\text{kg}$, at least 0.2 mg/kg, at least 1.0 mg/kg, at least 2.0 mg/kg, at least 10 mg/kg, at least 25 mg/kg, or at least 50 mg/kg (see, e.g.,

Yang *et al.*, (2003) *New Engl. J. Med.* 349:427-434; Herold *et al.*, (2002) *New Engl. J. Med.* 346:1692-1698; Liu *et al.*, (1999) *J. Neurol. Neurosurg. Psych.* 67:451-456; Portielji *et al.*, (2003) *Cancer Immunol. Immunother.* 52:133-144). The desired dose of antibodies or fragments thereof is about the same as for an antibody or polypeptide, on a moles/kg body weight basis. The desired plasma concentration of the antibodies or fragments thereof is about, on a moles/kg body weight basis. The dose may be at least 15 μg , at least 20 μg , at least 25 μg , at least 30 μg , at least 35 μg , at least 40 μg , at least 45 μg , at least 50 μg , at least 55 μg , at least 60 μg , at least 65 μg , at least 70 μg , at least 75 μg , at least 80 μg , at least 85 μg , at least 90 μg , at least 95 μg , or at least 100 μg . The doses administered to a subject may number at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12, or more.

For antibodies or fragments thereof of the invention, the dosage administered to a patient may be 0.0001 mg/kg to 100 mg/kg of the patient's body weight. The dosage may be between 0.0001 mg/kg and 20 mg/kg, 0.0001 mg/kg and 10 mg/kg, 0.0001 mg/kg and 5 mg/kg, 0.0001 and 2 mg/kg, 0.0001 and 1 mg/kg, 0.0001 mg/kg and 0.75 mg/kg, 0.0001 mg/kg and 0.5 mg/kg, 0.0001 mg/kg to 0.25 mg/kg, 0.0001 to 0.15 mg/kg, 0.0001 to 0.10 mg/kg, 0.001 to 0.5 mg/kg, 0.01 to 0.25 mg/kg or 0.01 to 0.10 mg/kg of the patient's body weight.

The dosage of the antibodies or fragments thereof of the invention may be calculated using the patient's weight in kilograms (kg) multiplied by the dose to be administered in mg/kg. The dosage of the antibodies or fragments thereof of the invention may be 150 $\mu\text{g}/\text{kg}$ or less, 125 $\mu\text{g}/\text{kg}$ or less, 100 $\mu\text{g}/\text{kg}$ or less, 95 $\mu\text{g}/\text{kg}$ or less, 90 $\mu\text{g}/\text{kg}$ or less, 85 $\mu\text{g}/\text{kg}$ or less, 80 $\mu\text{g}/\text{kg}$ or less, 75 $\mu\text{g}/\text{kg}$ or less, 70 $\mu\text{g}/\text{kg}$ or less, 65 $\mu\text{g}/\text{kg}$ or less, 60 $\mu\text{g}/\text{kg}$ or less, 55 $\mu\text{g}/\text{kg}$ or less, 50 $\mu\text{g}/\text{kg}$ or less, 45 $\mu\text{g}/\text{kg}$ or less, 40 $\mu\text{g}/\text{kg}$ or less, 35 $\mu\text{g}/\text{kg}$ or less, 30 $\mu\text{g}/\text{kg}$ or less, 25 $\mu\text{g}/\text{kg}$ or less, 20 $\mu\text{g}/\text{kg}$ or less, 15 $\mu\text{g}/\text{kg}$ or less, 10 $\mu\text{g}/\text{kg}$ or less, 5 $\mu\text{g}/\text{kg}$ or less, 2.5 $\mu\text{g}/\text{kg}$ or less, 2 $\mu\text{g}/\text{kg}$ or less, 1.5 $\mu\text{g}/\text{kg}$ or less, 1 $\mu\text{g}/\text{kg}$ or less, 0.5 $\mu\text{g}/\text{kg}$ or less, or 0.5 $\mu\text{g}/\text{kg}$ or less of a patient's body weight.

Unit dose of the antibodies or fragments thereof of the invention may be 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.1 mg to 12 mg, 0.1 mg to 10 mg, 0.1 mg to 8 mg, 0.1 mg to 7 mg, 0.1 mg to 5 mg, 0.1 to 2.5 mg, 0.25 mg to 20 mg, 0.25 to 15 mg, 0.25 to 12 mg, 0.25 to 10 mg, 0.25 to 8 mg, 0.25 mg to 7 m g, 0.25 mg to 5 mg, 0.5 mg to 2.5 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1 mg to 12 mg, 1 mg to 10 mg, 1 mg to 8 mg, 1 mg to 7 mg, 1 mg to 5 mg, or 1 mg to 2.5 mg.

The dosage of the antibodies or fragments thereof of the invention may achieve a serum titer of at least 0.1 µg/ml, at least 0.5 µg/ml, at least 1 µg/ml, at least 2 µg/ml, at least 5 µg/ml, at least 6 µg/ml, at least 10 µg/ml, at least 15 µg/ml, at least 20 µg/ml, at least 25 µg/ml, at least 50 µg/ml, at least 100 µg/ml, at least 125 µg/ml, at least 150 µg/ml, at least 175 µg/ml, at least 200 µg/ml, at least 225 µg/ml, at least 250 µg/ml, at least 275 µg/ml, at least 300 µg/ml, at least 325 µg/ml, at least 350 µg/ml, at least 375 µg/ml, or at least 400 µg/ml in a subject.

Alternatively, the dosage of the antibodies or fragments thereof of the invention may achieve a serum titer of at least 0.1 µg/ml, at least 0.5 µg/ml, at least 1 µg/ml, at least, 2 µg/ml, at least 5 µg/ml, at least 6 µg/ml, at least 10 µg/ml, at least 15 µg/ml, at least 20 .mu.g/ml, at least 25 µg/ml, at least 50 µg/ml, at least 100 µg/ml, at least 125 µg/ml, at least 150 µg/ml, at least 175 µg/ml, at least 200 µg/ml, at least 225 µg/ml, at least 250 µg/ml, at least 275 µg/ml, at least 300 µg/ml, at least 325 µg/ml, at least 350 µg/ml, at least 375 µg/ml, or at least 400 µg/ml in the subject.

Doses of antibodies or fragments thereof of the invention may be repeated and the administrations may be separated by at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or at least 6 months.

An effective amount for a particular patient may vary depending on factors such as the condition being treated, the overall health of the patient, the method route and dose of administration and the severity of side affects (see, e.g., Maynard *et al.*, (1996) A Handbook of SOPs for Good Clinical Practice, Interpharm Press, Boca Raton, Fla.; Dent (2001) Good Laboratory and Good Clinical Practice, Urch Publ., London, UK).

The route of administration may be by, e.g., topical or cutaneous application, injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial, intracerebrospinal, intralesional, or by sustained release systems or an implant (see, e.g., Sidman *et al.*, (1983) Biopolymers 22:547-556; Langer *et al.*, (1981) J. Biomed. Mater. Res. 15:167-277; Langer (1982) Chem. Tech. 12:98-105; Epstein et al., (1985) Proc. Natl. Acad. Sci. USA 82:3688-3692; Hwang *et al.*, (1980) Proc. Natl. Acad. Sci. USA 77:4030-4034; U.S. Pat. Nos. 6,350,466 and 6,316,024). Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the injection. In addition, pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. See, e.g., U.S. Pat. Nos.

6,019,968, 5,985,320, 5,985,309, 5,934,272, 5,874,064, 5,855,913, 5,290,540, and 4,880,078; and PCT Publication Nos. WO 92/19244, WO 97/32572, WO 97/44013, WO 98/31346, and WO 99/66903, each of which is incorporated herein by reference their entirety.

5 A composition of the present invention may also be administered via one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Selected routes of administration for antibodies or fragments thereof of the invention include intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other parenteral routes of administration, for example by injection or
10 infusion. Parenteral administration may represent modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion. Alternatively, a
15 composition of the invention can be administered via a non-parenteral route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically. In one embodiment, the antibodies or fragments thereof of the invention is administered by infusion. In another embodiment, the multispecific epitope
20 binding protein of the invention is administered subcutaneously.

If the antibodies or fragments thereof of the invention are administered in a controlled release or sustained release system, a pump may be used to achieve controlled or sustained release (see Langer, *supra*; Sefton, (1987) *CRC Crit. Ref Biomed. Eng.* 14:20; Buchwald *et al.*,
25 (1980), *Surgery* 88:507; Saudek *et al.*, (1989) *N. Engl. J. Med.* 321:574). Polymeric materials can be used to achieve controlled or sustained release of the therapies of the invention (see e.g., *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, (1983) *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; see also Levy *et al.*, (1985) *Science* 228:190; During *et al.*, (1989) *Ann. Neurol.* 25:351; Howard *et al.*, (1989) *J. Neurosurg.* 71:105); U.S. Pat. No. 5,679,377; U.S. Pat. No. 5,916,597; U.S. Pat. No. 5,912,015; U.S. Pat. No. 5,989,463; U.S. Pat. No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO
30 99/20253. Examples of polymers used in sustained release formulations include, but are not

limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In one embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. A controlled or sustained release system can be placed in proximity of the prophylactic or therapeutic target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138 (1984)).

Controlled release systems are discussed in the review by Langer, (1990), *Science* 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more antibodies or fragments thereof of the invention. See, e.g., U.S. Pat. No. 4,526,938, PCT publication WO 91/05548, PCT publication WO 96/20698, Ning *et al.*, (1996), *Radiotherapy & Oncology* 39:179-189, Song *et al.*, (1995) *PDA Journal of Pharmaceutical Science & Technology* 50:372-397, Cleek *et al.*, (1997) *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854, and Lam *et al.*, (1997) *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760, each of which is incorporated herein by reference in their entirety.

If the antibodies or fragments thereof of the invention are administered topically, they can be formulated in the form of an ointment, cream, transdermal patch, lotion, gel, shampoo, spray, aerosol, solution, emulsion, or other form well-known to one of skill in the art. See, e.g., Remington's *Pharmaceutical Sciences and Introduction to Pharmaceutical Dosage Forms*, 19th ed., Mack Pub. Co., Easton, Pa. (1995). For non-sprayable topical dosage forms, viscous to semi-solid or solid forms comprising a carrier or one or more excipients compatible with topical application and having a dynamic viscosity, in some instances, greater than water are typically employed. Suitable formulations include, without limitation, solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, and the like, which are, if desired, sterilized or mixed with auxiliary agents (e.g., preservatives, stabilizers, wetting agents, buffers, or salts) for influencing various properties, such as, for example, osmotic pressure. Other suitable topical dosage forms include sprayable aerosol preparations wherein the active ingredient, in some instances, in combination with a solid or liquid inert carrier, is packaged in a mixture with a pressurized volatile (e.g., a gaseous propellant, such as freon) or in a

squeeze bottle. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well-known in the art.

5 If the compositions comprising antibodies or fragments thereof are administered intranasally, it can be formulated in an aerosol form, spray, mist or in the form of drops. In particular, prophylactic or therapeutic agents for use according to the present invention can be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant (e.g., dichlorodifluoromethane,
10 trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges (composed of, e.g., gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

15 Methods for co-administration or treatment with a second therapeutic agent, e.g., a cytokine, steroid, chemotherapeutic agent, antibiotic, or radiation, are known in the art (see, e.g., Hardman *et al.*, (eds.) (2001) Goodman and Gilman's The Pharmacological Basis of Therapeutics, 10th ed., McGraw-Hill, New York, N.Y.; Poole and Peterson (eds.) (2001)
20 Pharmacotherapeutics for Advanced Practice: A Practical Approach, Lippincott, Williams & Wilkins, Phila., Pa.; Chabner and Longo (eds.) (2001) Cancer Chemotherapy and Biotherapy, Lippincott, Williams & Wilkins, Phila., Pa.). An effective amount of therapeutic may decrease the symptoms by at least 10%; by at least 20%; at least about 30%; at least 40%, or at least 50%.

25 Additional therapies (e.g., prophylactic or therapeutic agents), which can be administered in combination with the antibodies or fragments thereof of the invention may be administered less than 5 minutes apart, less than 30 minutes apart, 1 hour apart, at about 1 hour apart, at about 1 to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to
30 about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, at about 12 hours to 18 hours apart, 18 hours to 24 hours apart, 24 hours to 36 hours apart, 36 hours to 48 hours apart, 48

hours to 52 hours apart, 52 hours to 60 hours apart, 60 hours to 72 hours apart, 72 hours to 84 hours apart, 84 hours to 96 hours apart, or 96 hours to 120 hours apart from the antibodies or fragments thereof of the invention. The two or more therapies may be administered within one same patient visit.

5

The antibodies or fragments thereof of the invention and the other therapies may be cyclically administered. Cycling therapy involves the administration of a first therapy (e.g., a first prophylactic or therapeutic agent) for a period of time, followed by the administration of a second therapy (e.g., a second prophylactic or therapeutic agent) for a period of time, optionally, followed by the administration of a third therapy (e.g., prophylactic or therapeutic agent) for a period of time and so forth, and repeating this sequential administration, i.e., the cycle in order to reduce the development of resistance to one of the therapies, to avoid or reduce the side effects of one of the therapies, and/or to improve the efficacy of the therapies.

10

15 In certain embodiments, the antibodies or fragments thereof of the invention can be formulated to ensure proper distribution in vivo. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds of the invention cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g., U.S. Pat. Nos. 4,522,811; 5,374,548; and 20 5,399,331. The liposomes may comprise one or more moieties which are selectively transported into specific cells or organs, thus enhance targeted drug delivery (see, e.g., Ranade, (1989) *J. Clin. Pharmacol.* 29:685). Exemplary targeting moieties include folate or biotin (see, e.g., U.S. Pat. No. 5,416,016 to Low et al); mannosides (Umezawa *et al.*, (1988) *Biochem. Biophys. Res. Commun.* 153:1038); antibodies (Bloeman *et al.*, (1995) *FEBS Lett.* 25 357:140; Owais *et al.*, (1995) *Antimicrob. Agents Chemother.* 39:180); surfactant protein A receptor (Briscoe *et al.*, (1995) *Am. J. Physiol.* 1233:134); p 120 (Schreier *et al.*, (1994) *J. Biol. Chem.* 269:9090); see also K. Keinanen; M. L. Laukkanen (1994) *FEBS Lett.* 346:123; J. J. Killion; I. J. Fidler (1994) *Immunomethods* 4:273.

25

30 The invention provides protocols for the administration of pharmaceutical composition comprising antibodies or fragments thereof of the invention alone or in combination with other therapies to a subject in need thereof. The therapies (e.g., prophylactic or therapeutic agents) of the combination therapies of the present invention can be administered concomitantly or sequentially to a subject. The therapy (e.g., prophylactic or therapeutic agents) of the

30

combination therapies of the present invention can also be cyclically administered. Cycling therapy involves the administration of a first therapy (e.g., a first prophylactic or therapeutic agent) for a period of time, followed by the administration of a second therapy (e.g., a second prophylactic or therapeutic agent) for a period of time and repeating this sequential
5 administration, i.e., the cycle, in order to reduce the development of resistance to one of the therapies (e.g., agents) to avoid or reduce the side effects of one of the therapies (e.g., agents), and/or to improve, the efficacy of the therapies.

The therapies (e.g., prophylactic or therapeutic agents) of the combination therapies of the
10 invention can be administered to a subject concurrently. The term "concurrently" is not limited to the administration of therapies (e.g., prophylactic or therapeutic agents) at exactly the same time, but rather it is meant that a pharmaceutical composition comprising antibodies or fragments thereof of the invention are administered to a subject in a sequence and within a
15 time interval such that the antibodies of the invention can act together with the other therapy(ies) to provide an increased benefit than if they were administered otherwise. For example, each therapy may be administered to a subject at the same time or sequentially in any order at different points in time; however, if not administered at the same time, they should be administered sufficiently close in time so as to provide the desired therapeutic or prophylactic effect. Each therapy can be administered to a subject separately, in any
20 appropriate form and by any suitable route. In various embodiments, the therapies (e.g., prophylactic or therapeutic agents) are administered to a subject less than 15 minutes, less than 30 minutes, less than 1 hour apart, at about 1 hour apart, at about 1 hour to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about
25 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, 24 hours apart, 48 hours apart, 72 hours apart, or 1 week apart. In other embodiments, two or more therapies (e.g., prophylactic or therapeutic agents) are administered to a within the same patient visit.

30

The prophylactic or therapeutic agents of the combination therapies can be administered to a subject in the same pharmaceutical composition. Alternatively, the prophylactic or therapeutic agents of the combination therapies can be administered concurrently to a subject in separate

pharmaceutical compositions. The prophylactic or therapeutic agents may be administered to a subject by the same or different routes of administration.

The invention having been fully described, it is further illustrated by the following examples and claims, which are illustrative and are not meant to be further limiting.

5

Examples

Example 1: Methods, Materials and Screening for Antibodies

(i) Cell Lines

BXPC-3, SK-Br-3, BT-474, MDA-MB-453, FaDu and MCF-7 cell lines were purchased from
10 ATCC and routinely maintained in growth media supplemented with 10% fetal bovine serum (FBS).

(ii) Generation of Recombinant Human, Cyno, Mouse and Rat HER3 Vectors

Murine HER3 extracellular domain was PCR amplified from mouse brain cDNA (Clontech) and sequence verified by comparison with Refseq NM_010153. Rat HER3 ECD was reverse
15 transcribed from Rat-2 cell mRNA and sequence verified by comparison with NM_017218. Cynomolgus HER3 cDNA template was generated using RNA from various cyno tissues (Zyagen Laboratories), and the RT-PCR product cloned into pCR[®]-TOPO-XL (Invitrogen) prior to sequencing of both strands. Human HER3 was derived from a human fetal brain cDNA library (Source) and sequence verified by comparison with NM_001982.

20 To generate tagged recombinant proteins, human, mouse, rat and cyno HER3 was PCR amplified using Pwo Taq polymerase (Roche Diagnostics). Amplified PCR products were gel purified and cloned into a pDonR201 (Invitrogen) gateway entry vector that had previously been modified to include an in-frame N-terminal CD33 leader sequence and a C-terminal TAG, e.g., FLAG TAG. The TAG allows purification of monomeric proteins via an anti-
25 TAG monoclonal antibody. The target genes were flanked with AttB1 and AttB2 allowing recombination into Gateway adapted proprietary destination vectors (e.g., pcDNA3.1) using the Gateway[®] cloning technology (Invitrogen). Recombination reactions were performed using a Gateway LR reaction with proprietary destination vectors containing a CMV promoter to create the TAG expression vectors, although any commercially available vector can be
30 used.

Further recombinant HER3 proteins were generated that fused the HER3 ECD upstream of a C-terminal Factor X cleavage site and the human IgG hinge and Fc domain to create an Fc-tagged protein. To achieve this, the various HER3 ECD's were PCR amplified and cloned into a vector (e.g., pcDNA3.1) modified to contain an in-frame C-terminal fusion of Factor X site-
 5 Hinge-hFc. The generated open reading frame was flanked with AttB1 and AttB2 sites for further cloning with the Gateway[®] recombinant cloning technology (Invitrogen). An LR Gateway reaction was used to transfer HER3-Fc into a destination expression construct containing a CMV promoter. HER3 point mutation expression constructs were generated using standard site directed mutagenesis protocols and the resultant vectors sequence verified.

10 Table 8. Generation of HER3 expression vectors. HER3 amino acid numbering is based on NP_001973 (human), NP_034283 (mouse) and NP_058914 (rat).

Name	Description
Hu HER3	CD33-[Human HER3, residues 20-640]-TAG
Mu HER3	CD33-[Murine HER3, residues 20-643]-TAG
Rat HER3	CD33-[Rat HER3, residues 20-643]-TAG
Cyno HER3	CD33-[Cyno HER3, residues 20-643]-TAG
HER3 D1-2	CD33-[Human HER3, residues 20-329]-TAG
HER3 D2	CD33-[Human HER3, residues 185-329]- TAG
HER3 D3-4	CD33-[Human HER3, residues 330-643]- TAG
HER3 D4	CD33-[Human HER3, residues 496-643]- TAG
Hu HER3-Fc	[Human HER3, residues 1-643]-Fc
Mu HER3-Fc	[Murine HER3, residues 1-643]-Fc
Cyno HER3-Fc	[Cyno HER3, residues 1-643]-Fc
Rat HER3-Fc	[Rat HER3, residues 1-643]-Fc
HER3 D2-Fc	[Human HER3 residues 207-329]-Fc
HER3 K267A	CD33-[Human HER3, residues 20-640, K267A]-TAG
HER3 L268A	CD33-[Human HER3, residues 20-640, L268A]-TAG
HER3 K267A/ L268A	CD33-[Human HER3, residues 20-640, K267A/ L268A]-TAG

(iii) Expression of Recombinant HER3 Proteins

The desired HER3 recombinant proteins were expressed in HEK293 derived cell lines previously adapted to suspension culture and grown in a Novartis proprietary serum-free medium. Small scale expression verification was undertaken in transient 6-well-plate
5 transfection assays on the basis of lipofection. Large-scale protein production via transient transfection and was performed at the 10- 20 L scale in the Wave™ bioreactor system (Wave Biotech). DNA Polyethylenimine (Polysciences) was used as a plasmid carrier at a ratio of 1:3 (w:w). The cell culture supernatants were harvested 7-10 days post transfection and concentrated by cross-flow filtration and diafiltration prior to purification.

10 *(iv) Tagged Protein Purification*

Recombinant tagged HER3 proteins (e.g., TAG-HER3) were purified by collecting the cell culture supernatant and concentrating 10-fold by cross-flow filtration with a 10 kDa cut off filter (Fresenius). An anti-TAG column was prepared by coupling an anti-TAG monoclonal antibody to CNBr activated Sepharose 4B at a final ratio of 10 mg antibody per mL of resin.

15 Concentrated supernatant was applied to a 35ml anti-Tag column at a flow rate of 1- 2 mL/minute. After base-line washing with PBS, bound material was eluted with 100 mM glycine (pH 2.7), neutralized and sterile filtered. Protein concentrations were determined by measuring the absorbance at 280 nm and converting using a theoretical factor of 0.66 AU/mg. The purified protein was finally characterized by SDS-PAGE, N-terminal sequencing and
20 LC-MS.

(v) Fc Tag Purification

Concentrated cell culture supernatant was applied to a 50 ml Protein A Sepharose Fast Flow column at a flow rate of 1 ml/min. After baseline washing with PBS, the column was washed with 10 column volumes of 10 mM NaH₂PO₄/ 30% (v/v) Isopropanol, pH 7.3 followed by 5
25 column volumes of PBS. Finally, bound material was eluted with 50 mM Citrate/140 mM NaCl (pH 2.7), neutralized and sterile filtered.

(vi) Generation of Over-Expressing Cell Lines

To generate a cell line that expresses high levels of HER3 on the cell surface, a mammalian expression vector was constructed containing an insert coding for a CD33 leader sequence
30 upstream of amino acid residues 20-667 of human HER3 fused in-frame to amino acid residues 669-1210 of human EGFR. When expressed in mammalian cells the resultant

chimeric protein contains an N-terminal HER3 extracellular and transmembrane domain and a C-terminal EGFR cytoplasmic domain. The HER3/1 vector was transfected into CHO-S cells (Invitrogen) and stable pools generated following antibiotic selection. The resultant cell line (CHO HER3/1) expressed high levels of HER3 extracellular domain on its cell surface.

5 (vii) *HuCAL GOLD[®] Pannings*

For the selection of antibodies recognizing human HER3 multiple panning strategies were employed. Therapeutic antibodies against human HER3 protein were generated by selection of clones having high binding affinities, using as the source of antibody variant proteins a commercially available phage display library, the MorphoSys HuCAL GOLD[®] library. The phagemid library is based on the HuCAL[®] concept (Knappik *et al.*, (2000) J Mol Biol 296:57-86) and employs the CysDisplay[®] technology for displaying the Fab on the phage surface (WO01/05950 to Lohning).

For the isolation of anti-HER3 antibodies, standard as well as RapMAT panning strategies were performed using solid phase, solution, whole cell and differential whole cell panning approaches.

15 (viii) *Solid Phase Panning*

To identify anti-HER3 antibodies a variety of solid phase panning strategies were performed using differing recombinant HER3 proteins. To perform each round of solid phase panning, Maxisorp plates (Nunc) were coated with HER3 protein. Tagged proteins were either captured using plates previously coated with anti-Fc (goat or mouse anti-human IgG, Jackson Immuno Research), anti-Tag antibody or via passive adsorption. The coated plates were washed with PBS and blocked. Coated plates were washed twice with PBS prior to the addition of HuCAL GOLD[®] phage-antibodies for 2 hours at room temperature on a shaker. Bound phages were eluted were added to *E. coli* TG-1 and incubated for phage infection. Subsequently infected bacteria were isolated and plated on agar plates. Colonies were scraped off the plates and phages were rescued and amplified. Each HER3 panning strategy comprised of individual rounds of panning and contained unique antigens, antigen concentrations and washing stringency.

25 (ix) *Solution Phase Panning*

Each round of solution phase panning was performed using various biotinylated recombinant HER3 proteins in the presence or absence of neuregulin 1- β 1 (R&D Systems). Proteins were

biotinylated using the EZ-link sulfo-NHS-LC biotinylation kit (Pierce) according to the manufacturers instructions. 800µl of Streptavidin linked magnetic beads (Dynabeads, Dynal) were washed once with PBS and blocked overnight with Chemiblocker (Chemicon). HuCAL GOLD[®] phage-antibodies and the appropriate biotinylated HER3 were incubated in a reaction tube. Streptavidin magnetic beads were added for 20 minutes and were collected with a magnetic particle separator (Dynal). Bound phages were eluted from the Dynabeads by adding DTT containing buffer to each tube and added to *E. coli* TG-1. Phage infection was performed in an identical manner to that described in solid phase panning. Each HER3 panning strategy comprised of individual rounds of panning and contained unique antigens, antigen concentrations and washing stringency.

(x) Cell based panning

For cell pannings, HuCAL GOLD[®] phage-antibodies were incubated with approximately 10⁷ cells on a rotator for 2 hours at room temperature, followed by centrifugation. The cell pellet was isolated phages were eluted from the cells The supernatant was collected and added to *E. coli* TG-1 culture continued by the process described above . Two cell based strategies were employed to identify anti-HER3 antibodies:

- a) Whole cell panning: In this strategy a variety of intact cell lines were used as the antigens.
- b) Differential whole cell panning: In this strategy the antigens sequentially consisted of cells and recombinant HER3 proteins (see 1981.09 as an example). The cell based pannings were performed as described above whilst solid phase panning protocols were employed when recombinant proteins were utilized as antigens. The washes were conducted using PBS (2-3X) and PBST (2-3X).

(xi) RapMAT[™] library generation and pannings

In order to increase antibody binding affinity whilst maintaining library diversity the second round output of both solution and solid phase pannings were entered into the RapMAT[™] process whilst the third round output of the whole cell and differential whole cell panning strategies were entered (Prassler *et al.*, (2009) Immunotherapy; 1: 571-583. RapMAT[™] libraries were generated by sub-cloning Fab-encoding inserts of phages selected via panning into the display vector pMORPH[®]25_bla_LHC and were further digested to either generate H-CDR2 RapMAT[™] libraries and L-CDR3 RapMAT[™] libraries by using specific restriction

enzymes. The inserts were replaced with TRIM maturation cassettes (Virnekas *et al.*, (1994) Nucleic Acids Research 22:5600- 5607) for H-CDR2 or L-CDR3 according to pool composition. Library sizes were estimated to range between 8×10^6 - 1×10^8 clones. RapMAT antibody-phage were produced and subjected to two further rounds of solution, solid phase or cell based panning using the experimental methods described previously.

Example 2: Transient expression of anti-HER3 IgG's

Suspension adapted HEK293-6E cells were cultivated in a BioWave20 to a density of approximately 2×10^6 viable cells/ mL. The cells were transiently transfected with the relevant sterile DNA: PEI-MIX and further cultivated. Seven days after transfection, cells were removed by crossflow filtration using Fresenius filters (0.2 μ m). The cell free material was concentrated with crossflow filtration using a 10kDa cut off filter (Fresenius) and the concentrate was sterile filtered through a stericup filter (0.22 μ m). The sterile supernatant was stored at 4°C.

Example 3: Purification of anti-HER3 IgG

The purification of IgG was performed on a ÄKTA 100 explorer Air chromatography system at 6°C in a cooling cabinet, using a XK16/20 column with 25 mL of self-packed MabSelect SuRe resin (all GE Healthcare). All flow rates were 3.5 mL/min, except for loading, at a pressure limit of 5 bar. The column was equilibrated with 3 column volumes of PBS prior to loading the filtered fermentation supernatant at 2.0 mL/min. The column was washed with 8 column volumes of PBS. IgG was eluted with a pH gradient, starting at 50 mM citrate/70 mM NaCl (pH 4.5), going linearly down in 12 Column volumes to 50 mM citrate/70 mM NaCl (pH 2.5), followed by a 2 column volume constant step of the same pH 2.5 buffer. The IgG containing fractions were pooled and immediately neutralized and sterile filtered (Millipore Steriflip, 0.22 μ m). OD₂₈₀ was measured and the protein concentration calculated based on the sequence data. The pools were separately tested for aggregation (SEC-MALS) and purity (SDS-PAGE and MS).

Example 4: Expression and Purification of HuCAL®-Fab Antibodies in E. coli

Expression of Fab fragments encoded by pMORPH®X9_Fab_MH in TG-1 cells was carried out in shaker flask cultures using 500 mL of 2x YT medium supplemented with 34 μ g/mL chloramphenicol. Cultures were shaken at 30 °C until the OD_{600nm} reached 0.5. Expression was induced by addition of 0.75 mM IPTG (isopropyl- β -D-thiogalactopyranoside) for 20

hours at 30 °C. Cells were disrupted using lysozyme. His₆-tagged Fab fragments were isolated via IMAC (Bio-Rad). Buffer exchange to 1x Dulbecco's PBS (pH 7.2) was performed using PD10 columns. Samples were sterile filtered (0.2 μm). Protein concentrations were determined by UV-spectrophotometry. The purity of the samples was analyzed in denaturing, reducing 15% SDS-PAGE. The homogeneity of Fab preparations was determined in native state by size exclusion chromatography (HP-SEC) with calibration standards

Example 5: HER3 Antibody Affinity (K_D) Measurements by Solution Equilibrium Titration (SET)

Affinity determination in solution was essentially performed as previously described (Friguet *et al.*, (1985) J Immunol Methods 77:305-19). In order to improve the sensitivity and accuracy of the SET method, it was transferred from classical ELISA to ECL based technology (Hanel *et al.*, (2005) Anal biochem 339:182-84).

Unlabeled HER3-Tag (human, rat, mouse or cyno) described previously was used for affinity determination by SET.

The data was evaluated with XLfit software (ID Business Solutions) applying customized fitting models. For K_D determination of each IgG the following model was used (modified according to Piehler, *et al* (Piehler *et al.*, (1997) J Immunol Methods 201:189-206).

$$y = \frac{2B_{\max}}{[IgG]} \left(\frac{[IgG]}{2} - \frac{\left(\frac{x + [IgG] + K_D}{2} - \sqrt{\frac{(x + [IgG] + K_D)^2}{4} - x[IgG]} \right)^2}{2[IgG]} \right)$$

[IgG]: applied total IgG concentration

x: applied total soluble antigen concentration (binding sites)

B_{max}: maximal signal of IgG without antigen

K_D: affinity

Example 6: Antibody Cell Binding Determination by FACS

The binding of antibodies to endogenous human antigen expressed on human cancer cells was accessed by FACS. In order to determine antibody EC₅₀ values SK-Br-3 cells were harvested

with accutase and diluted to 1×10^6 cells/mL in FACS buffer (PBS/ 3% FBS/ 0.2% NaN_3). 1×10^5 cells/ well were added to each well of a 96-well plate (Nunc) and centrifuged at 210 g for 5 minutes at 4 °C before removing the supernatant. Serial dilutions of test antibodies (diluted in 1:4 dilution steps with FACS buffer) were added to the pelleted cells and incubated for 1 hour on ice. The cells were washed and pelleted three times with 100 μL FACS buffer. PE conjugated goat anti-human IgG (Jackson ImmunoResearch) diluted 1/200 with FACS buffer were added to the cells and incubated on ice for 1 hour. Additional washing steps were performed three times with 100 μL FACS buffer followed by centrifugation steps at 210 g for 5 minutes at 4°C. Finally, cells were resuspended in 200 μL FACS buffer and fluorescence values were measured with a FACSArray (BD Biosciences). The amount of cell surface bound anti-HER3 antibody was assessed by measuring the mean channel fluorescence.

Example 7: HER3 Domain and Mutant Binding

96-well Maxisorp plates (Nunc) were coated overnight at 4 °C with 200 ng of the appropriate recombinant human protein (HER3-Tag, D1-2- Tag, D2- Tag, D3-4- Tag, D4- Tag, HER3 K267A- Tag, HER3 L268A-Tag, HER3 K267A/ L268A and a tagged irrelevant control). All wells were then washed three times with PBS/ 0.1% Tween-20, blocked for one hour with PBS/ 1% BSA/ 0.1% Tween-20 and washed three times with PBS/ 0.1% Tween-20. Anti-HER3 antibodies were added to the relevant wells up to a final concentration of 10 $\mu\text{g}/\text{mL}$ were added to the appropriate wells and incubated at room temperature for two hours. Plates were washed three times with PBS/ 0.1% Tween-20 prior to the addition of the appropriate peroxidase linked detection antibody diluted 1/10000 in PBS/ 1% BSA/ 0.1% Tween-20. The detection antibodies used were goat anti-mouse (Pierce, 31432), rabbit anti-goat (Pierce, 31402) and goat anti-human (Pierce, 31412). Plates were incubated at room temperature for one hour before washing three times with PBS/ 0.1% Tween-20. 100 μl TMB (3,3', 5,5' tetramethyl benzidine) substrate solution (BioFfx) was added to all wells for 6 minutes before stopping the reaction with 50 μl 2.5% H_2SO_4 . The extent of HER3 antibody binding to each recombinant protein was determined by measuring the OD_{450} using a SpectraMax plate reader (Molecular Devices). Where appropriate, dose response curves were analyzed using Graphpad Prism.

30

Example 8: HER3 Epitope Mapping Using Hydrogen/ Deuterium Exchange Mass Spectrometry

Materials

D₂O buffer was made by dissolving 25 mM TBS (pH 7.5)/ 500 mM NaCl in heavy water (Sigma). The reduction solution was 50mM formate buffer (pH 4) 500mM TCEP and the quenching solution 0.5% (v/v) trifluoroacetic acid (TFA) in water. Buffer A was 0.25% formic acid/ 10% methanol/ 10% ethylene glycol in water, and buffer B was 0.25% formic acid in acetonitrile. All chemicals were purchased from Sigma, and HPLC grade solvents were from Fisher Scientific.

10 *Liquid Handling and Chromatography*

Automated hydrogen-deuterium exchange mass spectrometry (HDX MS) experiments were designed based upon methods and equipment described by Wales *et al.*, (2006) Anal. Chem. 78:1005-1014). In short, all liquid handling operations used a Pal HTS liquid-handler (LEAP Technologies) housed in a refrigerated enclosure maintained at 2°C. A 6-port injection valve and a wash station were mounted on the liquid-handler rail and facilitated sample injection into the chromatographic system and syringe washing. The chromatographic system, consisted of an additional 10-port valve, a 2.1 mm x 30 mm Poroszyme pepsin column (Applied Biosystems), a reverse-phase 0.5 mm x 2 mm Cap Trap cartridge (Michrom Bioresources), and a self-packed electrospray emitter as analytical column (100 µm x ~60 mm, Kinetex 2.6 µm C18, Phenomenex). The 10-port valve head, the trap cartridge and the analytical column were housed in a separate enclosure constructed from aluminum and maintained at -5°C by peltier stacks. Valves and columns were configured in such a way as to allow in-line protein digestion, peptide desalting, and reversed-phase chromatography prior to introduction of the sample into the electrospray ionization (ESI) source of the mass spectrometer (LTQ-Orbitrap, Thermo Scientific).

The fluid streams required for operation were provided by two separate HPLC pumps. The first HPLC (Surveyor MS pump, Thermo Scientific) delivered buffer A at a constant flow rate of 125 µL/min and was used to transfer sample through the immobilized pepsin cartridge onto the reversed-phase trap cartridge mounted across the 10-port valve. After the loading and desalting period, the 10-port valve was switched to elute the sample with the help of a gradient pump (AQUITY UPLC, Waters) from the reversed-phase trap cartridge, through the analytical column and into the ion source of the mass spectrometer. The immobilized enzyme

cartridge was isolated to waste during gradient elution. The gradient pump delivered linear gradient segments of 0 to 40% mobile phase B over 35 minutes at 5 $\mu\text{L}/\text{min}$ and 40 to 95% mobile phase B at 5 $\mu\text{L}/\text{min}$ over 10 minutes. The gradient flow from the pump was split at the 10-port valve using a passive splitter so that the actual flow through the trap cartridge and analytical column for gradient elution was $\sim 1 \mu\text{L}/\text{min}$. The entire chromatographic run was 70 minutes long including washing and equilibration steps.

Mass Spectrometry

For the purpose of identification of proteolytic fragments resulting from online digestion several data-dependent MS/MS experiments were performed. For these acquisitions, tandem MS spectra were acquired with the LTQ analyzer of the LTQ-Orbitrap hybrid mass spectrometer. Precursor mass selection was based on MS scans acquired by the Orbitrap analyzer. Single stage MS acquisitions performed for the purpose of deuteration level determination were acquired at a resolution of 60,000 by the Orbitrap (over m/z 400-2000) analyzer.

Preparation of Protein and Protein:Fab Complexes

HER3 protein was prepared by diluting 50 μg HER3-Tag with 25 mM TBS (pH 7.5)/ 500 mM NaCl to yield a final volume of 50 μL . Protein:Fab complexes were prepared by mixing 50 μg HER3-Tag in a 1:1 molar ratio with the Fab's studied. Protein:Fab mixtures were then diluted to a final volume of 50 μL with 25 mM TBS (pH 7.5)/ 500 mM NaCl.

Protein:Fab complexes were prepared and allowed to incubate for at least 2 hours at 4°C. Four 96-well plates containing sample, diluent, quench, and reduction solutions were loaded into the liquid-handler before the start of each experimental. For on-exchange experiments 50 μL of HER3 or HER3:Fab complex was diluted with 150 μL D_2O buffer. The mixture was reduced by adding 200 μL reduction buffer for 1 minute before quenching with 600 μL of quench buffer. The total volume after all liquid handling steps was $\sim 1\text{mL}$. Once mixed, the quenched solution was injected into the chromatographic system where it was automatically digested, separated and analyzed by LCMS. The average change in deuteration between sample and control was calculated as the difference between the deuterium uptake levels of the sample and control.

Data Processing

The Orbitrap RAW files were converted into mzXML files using an in-house program (RawXtract). Subsequently, tandem MS acquisitions were searched using SEQUEST (Yates Lab, Scripps Research Institute, La Jolla, CA) and search results were automatically filtered using DTASelect 2.0 (Yates Lab, Scripps Research Institute, La Jolla, CA). Using the peptide sequence identifications, an in-house written program was used to automatically extract single-ion chromatograms for each identified sequence and generate average spectra across the chromatographic peak. Average spectra were smoothed and centroided. The level of deuterium uptake was taken as the difference in mass between a deuterated sample and non-deuterated reference. Processed data was manually validated and adjusted to correct inaccuracies and errors from automated processing steps. Deuterium uptake levels were assigned to each residue of the protein sequence by delocalizing the deuterium content across each peptide (i.e., dividing the observed deuteration level by the number of amino acids in that peptide). If a residue was covered by more than one peptide, the normalized deuterium uptakes of all peptides covering that residue were averaged.

Example 9: X-ray crystallographic structure determination of the human HER3/MOR09823 Fab and human HER3/MOR09825 Fab complexes

The present example presents the crystal structure of full length HER3 bound to the Fab fragment of MOR09823 and the Fab fragment of MOR09825, determined at 3.2Å and 3.4Å resolution, respectively. Tagged human HER3 was further purified on a HiLoad 26/60 Superdex 200 PrepGrade column (GE Healthcare) equilibrated in PBS (pH 7.3). *E. coli* expressed MOR09823 and MOR09825 Fabs were isolated by lysing cells with lysozyme and His₆-tagged Fab fragments were captured on a HisTrap_HP (GE Healthcare) column. MOR09823 Fab-fragments were further purified by gel filtration chromatography using a Superdex 75 16/60 column (GE Healthcare) equilibrated in 25 mM Tris (pH 7.5), 150 mM NaCl.

HER3 Fab complexes were prepared by mixing excess Fab with tagged HER3 in a molar ratios of 1.3-1.8:1 (concentration estimated by absorbance at 280 nm using calculated extinction coefficients of 0.9 and 1.4 (mg/ml)⁻¹cm⁻¹ for HER3 and Fab, respectively) and purifying the complexes on a Superdex 200 10/300 column (GE Healthcare) equilibrated in 25 mM Tris (pH 7.5), 150 mM NaCl. Peak fractions were analyzed by SDS-PAGE and LCMS. For each complex, fractions containing both HER3 and Fab in an approximate

equimolar ratio were pooled and concentrated. HER3/MOR09823 crystals were grown at 293K by sitting drop vapor diffusion from drops containing 150 nl HER3/ MOR09823 complex and 150 nl of reservoir solution (100 mM sodium citrate pH 5.6, 20% PEG 4000 and 20% isopropanol). Crystals were transferred to reservoir solution containing additional 8% glycerol and flash cooled in liquid nitrogen. HER3/MOR09825 crystals were grown at 293K by sitting drop vapor diffusion from drops containing 150 nl HER3/ MOR09825 complex and 150 nl of reservoir solution (100 mM bis-tris pH 6.5, 16% PEG 10,000). Crystals were transferred to 100 mM bis-tris pH 6.5, 18% PEG 10,000 and 22% glycerol and flash cooled in liquid nitrogen.

10 Data were collected at beamline 17-ID at the Advanced Photon Source (Argonne National Laboratory). HER3/MOR09823 Fab complex data were processed and scaled at 3.2Å using HKL2000 (HKL Research Inc) in space group I222 with cell dimensions a=124.16, b=139.44, c=180.25 Å, with good statistics. The HER3/ MOR09823 Fab structure was solved by molecular replacement using Phaser (McCoy *et al.*, (2007) J. Appl. Cryst. 40:658-674) with

15 fragments of a Fab and the published HER3 ECD structure 1mb6 as search models. The final model, which contains 1 molecule of the HER3/MOR09823 Fab complex per asymmetric unit, was built in COOT (Emsley & Cowtan (2004) Acta Cryst. 60:2126-2132) and refined to R and R_{free} values of 19.0 and 24.5%, respectively, with an rmsd of 0.010 Å and 1.37° for bond lengths and bond angles, respectively, using BUSTER (Global Phasing, LTD). Residues

20 of HER3 that contain atoms within 5Å of any atom in MOR09823 Fab as identified in PyMOL (Schrödinger, LLC) are listed in Tables 11 and 12. HER3/MOR09825 Fab complex data were processed and scaled at 3.4Å using autoPROC (Global Phasing, LTD) in space group I222 with cell dimensions a=124.23, b=140.94, c=180.25 Å, with good statistics. The HER3/ MOR09825 Fab structure was solved by molecular replacement using Phaser (McCoy

25 *et al.*, (2007) J. Appl. Cryst. 40:658-674) with the HER3/MOR09823 Fab structure as a search model. The final model, which contains 1 molecule of the HER3/MOR09825 Fab complex per asymmetric unit, was built in COOT (Emsley & Cowtan (2004) Acta Cryst. 60:2126-2132) and refined to R and R_{free} values of 18.8 and 24.9%, respectively, with an rmsd of 0.009 Å and 1.21° for bond lengths and bond angles, respectively, using BUSTER (Global Phasing,

30 LTD). Residues of HER3 that contain atoms within 5Å of any atom in MOR09825 Fab as identified in PyMOL (Schrödinger, LLC) are listed in Tables 13 and 14.

Example 10: Phospho-HER3 *in vitro* cell assays.

MCF-7 cells were routinely maintained in DMEM/F12, 15mM HEPES, L-glutamine, 10% FCS and SK-Br-3 in McCoy's 5a, 10% FCS, 1.5mM L-glutamine. Sub-confluent MCF7 or SK-Br-3 cells grown in complete media were harvested with accutase (PAA Laboratories) and resuspended in the appropriate growth media at a final concentration of 5×10^5 cells/ mL. 100 μ L of cell suspension was then added to each well of a 96-well flat bottomed plate (Nunc) to give a final density of 5×10^4 cells/ well. MCF7 cells were allowed to attach for approximately 3 hours before the media was exchanged for starvation media containing 0.5% FBS. All plates were then incubated overnight at 37°C prior to treatment with the appropriate concentration of HER3 antibodies (diluted in the appropriate media) for 80 minutes at 37°C. MCF7 cells were treated with 50 ng/mL neuregulin 1- β 1 EGF domain (R&D Systems) for the final 20 minutes to stimulate HER3 phosphorylation. All media was gently aspirated and the cells washed with ice-cold PBS containing 1mM CaCl₂ and 0.5 mM MgCl₂ (Gibco). The cells were lysed by adding 50 μ L ice-cold lysis buffer (20 mM Tris (pH8.0)/ 137 mM NaCl/ 10% Glycerol/ 2mM EDTA/ 1% NP-40/ 1 mM sodium orthovanadate/, Aprotinin (10 μ g/mL)/ Leupeptin (10 μ g/mL)) and incubated on ice with shaking for 30 minutes. Lysates were then collected and spun at 1800 g for 15 minutes at 4°C to remove cell debris. 20 μ L of lysate was added to a pre-prepared capture plate.

HER3 capture plates were generated using a carbon plate (Mesoscale Discovery) coated overnight at 4°C with 20 μ L of 4 μ g/mL MAB3481 capture antibody (R&D Systems) diluted in PBS and subsequently blocked with 3% bovine serum albumin in 1x Tris buffer (Mesoscale Discovery)/ 0.1% Tween-20. HER3 was captured from the lysate by incubating the plate at room temperature for one hour with shaking before the lysate was aspirated and the wells washed with 1x Tris buffer (Mesoscale Discovery)/ 0.1% Tween-20. Phosphorylated HER3 was detected using 0.75 μ g/mL biotinylated anti-phosphotyrosine antibody (R&D Systems) prepared in 1% BSA/ 1x Tris/ 0.1% Tween-20 by incubating with shaking at room temperature for 1 hour. The wells were washed four times with 1x Tris/ 0.1% Tween-20 and biotinylated proteins were detected by incubating with S-Tag labelled Streptavidin (Mesoscale Discovery) diluted in 1% BSA/ 1x Tris/ 0.1% Tween-20 for one hour at room temperature. Each well was aspirated and washed four times with 1x Tris/ 0.1% Tween-20 before adding 20 μ L of Read buffer T with surfactant (Mesoscale Discovery) and the signal quantified using a Mesoscale Sector Imager. Antibodies MOR06391 or MOR03207 were included in signalling experiments as isotype controls.

Example 11: Phospho-Akt (S473) *in vitro* cell assays.

Sub-confluent SK-Br-3 and BT-474 cells grown in complete media were harvested with accutase (PAA Laboratories) and resuspended in the appropriate growth media at a final concentration of 5×10^5 cells/ mL. 100 μ L of cell suspension was then added to each well of a
5 96-well flat bottomed plate (Nunc) to yield a final density of 5×10^4 cells/ well. All plates were then incubated overnight at 37°C prior to treatment with the appropriate concentration of HER3 antibodies (diluted in the appropriate media) for 80 minutes at 37°C. All media was gently aspirated and the cells washed with ice-cold PBS containing 1mM CaCl₂ and 0.5 mM MgCl₂ (Gibco). The cells were lysed by adding 50 μ L ice-cold lysis buffer (20 mM Tris
10 (pH8.0)/ 137 mM NaCl/ 10% Glycerol/ 2mM EDTA/ 1% NP-40/ 1 mM sodium orthovanadate/ Aprotinin (10 μ g/mL)/ Leupeptin (10 μ g/mL)) and incubated on ice with shaking for 30 minutes. Lysates were then collected and spun at 1800 g for 15 minutes at 4°C to remove cell debris. 20 μ L of lysate was added to a multi-spot 384-well Phospho-Akt carbon plate (Mesoscale Discovery) that had previously been blocked with 3% BSA/ 1x Tris/ 0.1%
15 Tween-20. The plate was incubated at room temperature for two hours with shaking before the lysate was aspirated and the wells washed four times with 1x Tris buffer (Mesoscale Discovery)/ 0.1% Tween-20. Phosphorylated Akt was detected using 20 μ L of SULFO-TAG anti-phospho-Akt (S473) antibody (Mesoscale Discovery) diluted 50-fold in 1% BSA/ 1x Tris/ 0.1% Tween-20 by incubating with shaking at room temperature for 2 hours. The wells
20 were washed four times with 1x Tris/ 0.1% Tween-20 before adding 20 μ L of Read buffer T with surfactant (Mesoscale Discovery) and the signal quantified using a Mesoscale Sector Imager. Antibodies MOR06391 or MOR03207 were included in signalling experiments as isotype controls.

Example 12: Cell-line proliferation assays.

25 SK-Br-3 cells were routinely cultured in McCoy's 5A medium modified, supplemented with 10% fetal bovine serum and BT-474 cells were cultured in DMEM supplemented with 10% FBS. Sub-confluent cells were trypsinized, washed with PBS, diluted to 5×10^4 cells/ mL with growth media and plated in 96-well clear bottom black plates (Costar 3904) at a density of 5000 cells/ well. The cells were incubated overnight at 37°C before adding the appropriate
30 concentration of HER3 antibody (typical final concentrations of 10 or 1 μ g/mL). The plates were returned to the incubator for 6 days before assessing cell viability using CellTiter-Glo (Promega). 100 μ L of CellTiter-Glo solution was added to each well and incubated at room temperature with gentle shaking for 10 minutes. The amount of luminescence was determined

using a SpectraMax plate reader (Molecular Devices). The extent of growth inhibition obtained with each antibody was calculated by comparing the luminescence values obtained with each HER3 antibody to a standard isotype control antibody (MOR06391).

For proliferation assays MCF-7 cells were routinely cultured in DMEM/ F12 (1:1) containing 4 mM L-Glutamine/ 15mM HEPES/ 10% FBS. Sub-confluent cells were trypsinized, washed with PBS and diluted to 1×10^5 cells/ mL with DMEM/ F12 (1:1) containing 4 mM L-Glutamine/ 15mM HEPES/ 10 μ g/mL Human Transferrin/ 0.2% BSA. Cells were plated in 96-well clear bottom black plates (Costar) at a density of 5000 cells/ well. The appropriate concentration of HER3 antibody (typical final concentrations of 10 or 1 μ g/mL) was then added. 10 ng/mL of NRG1- β 1 EGF domain (R&D Systems) was also added to the appropriate wells to stimulate cell growth. The plates were returned to the incubator for 6 days before assessing cell viability using CellTiter-Glo (Promega). The extent of growth inhibition obtained with each antibody was calculated by subtracting the background (no neuregulin) luminescence values and comparing the resulting values obtained with each anti-HER3 antibody to a standard isotype control antibody (MOR06391).

Example 13: Ligand blocking cell assays

MCF-7 cells cultured in MEM supplemented with 10% FBS and 1 μ g/mL insulin (Sigma) were rinsed and collected in a small volume of FACSmax cell dissociation buffer (Genlantis) prior to the addition of 5 mL of FACS buffer (PBS/ 1% FBS/ 0.1% sodium azide). The cell density was counted and adjusted to a final concentration of 1×10^6 cells/ mL. 100 μ L of cell suspension was added to each well of a 96-well plate and the cells pelleted via centrifugation (220g, 3 minutes, 4°C). Cell pellets were resuspended in 100 μ L of the appropriate test antibodies diluted in FACS buffer (typical final antibody concentrations ranged from 100 to 0.1 nM) and the plate incubated on ice for 45 minutes. The ligand blocking antibody MAB3481 (R&D Systems) was included as a positive control. Cells were washed twice with staining buffer prior to adding 10 nM NRG1- β 1 EGF domain (R&D Systems) diluted in FACS buffer and incubating on ice for 45 minutes. Cells were washed twice with staining buffer and bound neuregulin detected by incubating the cells with 10 nM anti-human NRG1- β 1 EGF domain antibody (R&D Systems) on ice for 45 minutes. Cells were washed twice with staining buffer and incubated on ice for 45 minutes with PE-linked anti-goat antibody (Jackson ImmunoResearch) diluted 1/500 with FACS buffer. Cells were then pelleted via centrifugation and the pellet resuspended in 200 μ L FACS buffer. To quantify each sample

10,000 live cells were counted on a LSR II Flow Cytometer (BD Biosciences) and the amount of cell surface bound neuregulin was assessed by measuring the mean channel fluorescence.

Example 14: Ligand blocking biochemical assay

The present method includes utility of a Surface plasmon resonance (SPR)-based biosensor (Biacore™ GE Healthcare, Uppsala, Sweden) to examine the ability of HER3/ antibody
5 complexes to bind neuregulin.

Biacore™ utilizes the phenomenon of surface plasmon resonance (SPR) to detect and measure binding interactions. In a typical Biacore experiment, one of the interacting molecules (neuregulin) is immobilized on a matrix while the interacting partner (HER3) is
10 flowed over the surface. A binding interaction results in an increase in mass on the sensor surface and a corresponding direct change in the refractive index of the medium in the vicinity of the sensor surface. Changes in refractive index or signal are recorded in resonance units (R.U.) Signal changes due to association and dissociation of complexes are monitored in a non-invasive manner, continuously and in real-time, the results of which are reported in the
15 form of a sensorgram.

Biacore™ T100 (GE Healthcare, Uppsala, Sweden) was used to conduct all experiments reported herein. Sensor surface preparation and interaction analyses were performed at 25°C. Buffer and Biacore reagents were purchased from GE Healthcare. Running buffer containing 10mM HEPES, pH7.4/ 150mM NaCl, 0.05% P20, 0.5% BSA was utilized throughout the
20 assay.

NRG-1β1 extracellular domain (R&D Systems) was incubated on ice for 45 minutes with EZ-link Sulfo-NHS-LC-LC-Biotin (Pierce) at a molar ratio of 5:1. The reaction was quenched via the addition excess ethanolamine and uncoupled biotin removed from the biotinylated-NRG using desalt spin columns (Zeba). Biotinylated-NRG was captured onto a sensor chip CAP
25 pre-immobilized with approximately 3000 R.U. of ssDNA-streptavidin (Biotin CAPture kit) to yield neuregulin surface densities in the range 400 – 600 R.U. A reference flowcell was generated by omitting biotinylated-NRG from the injection steps such that only ssDNA-streptavidin was present on the flowcell surface.

HER3/ antibody complexes were generated by incubating 10nM human HER3-Fc with
30 increasing concentrations (0 – 50nM) of the appropriate test antibody for 15 minutes at room temperature prior to incubating in the Biacore™ at 10°C. Interaction analyses were performed

by injecting HER3/ antibody complexes over reference and neuregulin surfaces in series for 180 seconds at a flow-rate of 60 μ L/min. Complex dissociation was monitored for 180 seconds at a flow rate of 60 μ L/min. Surface regeneration was performed at the end of each analysis cycle using a 120 second injection of 8M guanidine: 1M NaOH (3:1) followed by a 120
5 second injection of 30% acetonitrile/ 0.25M NaOH at a flow rate of 30 μ L/min.

Example 15: *In vivo* PD studies

BxPC3 and BT-474 cells were cultured and implanted in female athymic nu/nu Balb/C mice (Harlan Laboratories) as described in Examples 16 and 17.

Once tumors had reached an appropriate size, animals were examined for tumor quality.
10 Animals with ulcerated tumors or animal with fluid-filled tumors were excluded from the study. The remaining animals were dosed intravenously with antibody via lateral tail vein injection. At the given time points, animals were euthanized via CO₂ asphyxiation and whole blood was collected via cardiac puncture and placed into a 1.5 mL Eppendorf collection tube. Tumor tissue was immediately dissected, placed into a screw-top polypropylene sample tube
15 and snap frozen in liquid nitrogen. Tissue was stored at -80°C until lysates were prepared.

Example 16: *In vivo* BT-474 efficacy studies

BT-474 cells were cultured in DMEM containing 10% heat-inactivated fetal bovine serum without antibiotics until the time of implantation.

One day before cell inoculation, female athymic nu/nu Balb/C mice (Harlan Laboratories)
20 were implanted subcutaneously with a sustained release 17 β -estradiol pellet (Innovative Research of America) to maintain serum estrogen levels. One day after 17 β -estradiol pellet implantation, 5 x10⁶ cells were injected orthotopically into the 4th mammary fatpad in a suspension containing 50% phenol red-free matrigel (BD Biosciences) in Hank's balanced salt solution. The total injection volume containing cells in suspension was 200 μ L. 20 days
25 following cell implantation animals with a tumor volume of approximately 200 mm³ were enrolled in the efficacy study. In general, a total of 10 animals per group were enrolled in efficacy studies.

For single-agent studies, animals were dosed intravenously via lateral tail vein injection with either MOR10701 or MOR10703. An initial loading dose of 40 mg/kg was given for the first
30 dose. After the initial dose, animals were on a 20 mg/kg, every other day schedule for the

duration of the study. For combination studies, animals were dosed with either MOR10701 or MOR10703 (20mg/kg, iv, q2d) and a sub-optimal dose of trastuzumab (1mg/kg, iv, 2qw).

For the duration of the studies, tumor volume was measured by calipering twice per week. Percent treatment/ control (T/C) values were calculated using the following formula:

5
$$\% T/C = 100 \times \Delta T/\Delta C \text{ if } \Delta T > 0$$

where:

T = mean tumor volume of the drug-treated group on the final day of the study;

ΔT = mean tumor volume of the drug-treated group on the final day of the study – mean tumor volume of the drug-treated group on initial day of dosing;

10 C = mean tumor volume of the control group on the final day of the study; and

ΔC = mean tumor volume of the control group on the final day of the study – mean tumor volume of the control group on initial day of dosing.

Body weight was measured twice per week and dose was body weight adjusted. The % change in body weight was calculated as $(BW_{\text{current}} - BW_{\text{initial}})/(BW_{\text{initial}}) \times 100$. Data is
15 presented as percent body weight change from the day of treatment initiation.

All data were expressed as mean \pm standard error of the mean (SEM). Delta tumor volume and body weight were used for statistical analysis. Between groups comparisons were carried out using a one-way ANOVA followed by a post hoc Tukey. For all statistical evaluations the level of significance was set at $p < 0.05$. Significance compared to the vehicle control group is
20 reported.

Example 17: *In vivo* BxPC3 efficacy studies

BxPC3 cells were cultured in RPMI-1640 medium containing 10% heat-inactivated fetal bovine serum without antibiotics until the time of implantation.

Female athymic nu/nu Balb/C mice (Harlan Laboratories) were implanted subcutaneously
25 with 10×10^6 cells in a mixture of 50% phosphate buffered saline with 50% matrigel. The total injection volume containing cells in suspension was 200 μL . Once tumors had reached approximately 200mm^3 in size, animals were enrolled in the efficacy study. In general, a total

of 10 animals per group were enrolled in studies. Animals were excluded from enrollment if they exhibited unusual tumor growth characteristics prior to enrollment.

Animals were dosed intravenously via lateral tail vein injection. An initial loading dose of 40 mg/kg was given for the first dose. After the initial dose, animals were on a 20 mg/kg, every other day schedule for the duration of the study (25 days under treatment). Tumor volume and T/C values were calculated as previously detailed.

Example 18: Phospho-Akt (S473) *in vivo* PD assays.

Approximately 50 mm³ frozen tumor (e.g. BT-474 or BXPC-3) tissue was thawed on ice and 100- 300 µL of T-PER buffer (Pierce) containing phosphatase (Roche) and protease inhibitors (Roche) was added to each sample. The volume of lysis buffer added was dependent upon the size of the tumor sample. The tissue was broken down using a 1.5 mL pestle (Fisher Scientific) and the resultant suspensions were incubated on ice for 15 minutes before being frozen overnight at -80 °C. Samples were thawed and spun for 15 minutes at 13000 g, 4 °C prior to quantifying the supernatant protein concentration by BCA assay (Thermo Scientific). Tissue supernatants were diluted with lysis buffer (Mesoscale Discovery) and 25 µg added to a multi-spot 96-well Phospho-Akt carbon plate (Mesoscale Discovery) that had previously been blocked with Blocking Solution-A (Mesoscale Discovery). The plate was incubated at room temperature for one hour with shaking before the lysate was aspirated and the wells washed four times with Tris Wash buffer (Mesoscale Discovery). Phosphorylated Akt was detected using 25 µL of SULFO-TAG anti-phospho-Akt (S473) antibody (Mesoscale Discovery) diluted in antibody dilution buffer by incubating with shaking at room temperature for one hour. The wells were washed four times with Tris Wash buffer before adding 150 µL of Read buffer T (with surfactant) (Mesoscale Discovery) and the signal quantified using a Mesoscale Sector Imager.

Example 19: Phospho HER3 (Y1197) *in vivo* PD assays

Approximately 50 mm³ frozen tumor (e.g. BXPC-3) tissue was thawed on ice and 100- 300 µL of T-PER buffer (Pierce) containing phosphatase (Roche) and protease inhibitors (Roche) was added to each sample. The tissue was broken down using a 1.5 mL pestle (Fisher Scientific) and the resultant suspensions were incubated on ice for 15 minutes before being frozen overnight at -80 °C. Samples were thawed and spun for 15 minutes at 13000 g, 4 °C prior to quantifying the supernatant protein concentration by BCA assay (Thermo Scientific). Tissue supernatants were diluted with lysis buffer and 150 µg added to a multi-spot 96-well

carbon plate (Mesoscale Discovery) that had previously been coated overnight with 4 µg/mL MAB3481 (R&D Systems) and blocked with 3% milk. The plate was incubated at room temperature for two hours with shaking before the lysate was aspirated and the wells washed four times with Tris Wash buffer (Mesoscale Discovery). Phosphorylated HER3 was bound using anti-HER3 pY1197 diluted 1/8000 with blocking buffer. Following incubation at room temperature for one hour the wells were washed with Tris Wash buffer and the anti-pY1197 antibody detected using S-Tag labelled anti-rabbit antibody (Mesoscale Discovery) diluted 1/1000 in blocking buffer by incubating with shaking at room temperature for one hour. The wells were washed four times with Tris Wash buffer before adding 150 µl of 1/4 diluted Read buffer T (with surfactant) (Mesoscale Discovery) and the signal quantified using a Mesoscale Sector Imager.

Example 20: *In vitro* drug combination studies

To assess the ability of HER3-targeted antibodies to combine with targeted therapies MOR09825 or MOR10703 were combined with trastuzumab, lapatinib, BEZ235, BKM120, BYL719, RAD001, erlotinib and cetuximab in cell viability assays. Approximately 1000-1500 SK-Br-3 (McCoy's), MDA-MB-453 (RPMI), FaDu (EMEM) or L3.3 (RPMI) cells were seeded into 384-well plates in the appropriate culture media supplemented with 2% FBS and allowed to adhere overnight at 37°C. The appropriate drug combinations (typical final drug concentrations for lapatinib, BKM120, and BYL719 ranged from 3µM to 13 nM; for RAD001 ranged from 27nM to 0.0041nM; for erlotinib ranged from 1 µM to 0.0025nM; for MOR1073 ranged from 100nm to 0.01nm; for cetuximab ranged from 100nM to 0.0015nM; and for trastuzumab ranged from 300nM to 0.046nM)) were subsequently added to the wells such that each plate contained a dose response curve of each drug in a two-dimensional matrix. The plates were returned to the incubator for 3- 6 days before assessing cell viability using CellTiter-Glo (Promega). CellTiter-Glo solution was added to each well and incubated at room temperature with gentle shaking for 10 minutes. The amount of luminescence was determined using a SpectraMax plate reader (Molecular Devices). The extent of growth inhibition obtained with each combination was calculated and combination activity highlighted using the Loewe additivity model.

Example 21: In vivo drug combination studies in L3.3 Cells.

Pancreatic L3.3 cells were cultured in DMEM medium containing 10% heat-inactivated fetal bovine serum until the time of implantation. Female Foxn1 nude mice (Harlan Laboratories) were implanted subcutaneously with 3×10^6 cells in FBS free DMEM. The total injection
5 volume containing cells in suspension was 100 μ L. 12 days following cell implantation, animals were enrolled in the efficacy study with a mean tumor volume of approximately 100mm^3 for all groups. In general, a total of 8 animals per group were enrolled in studies. Animals were excluded from enrollment if they exhibited unusual tumor growth characteristics prior to enrollment.

10 Animals were dosed intravenously with MOR10703 via lateral tail vein injection on a 20 mg/kg, every other day schedule for the duration of the study (14 days under treatment). Erlotinib was dosed at 50mg/kg (PO) on a daily schedule either as a single-agent or in combination with MOR10703. Tumor volume and T/C values were calculated as previously detailed.

15 Results and Discussion

Collectively, these results show that a class of antibodies bind to amino acid residues within domain 2 and domain 4 of a conformational epitope of HER3 and stabilizes HER3 in an inactive or closed conformation. Binding of these antibodies inhibits both ligand-dependent and ligand-independent signaling. These antibodies are also able to bind concurrently with a
20 HER3 ligand.

(i) Affinity Determination

Antibody affinity was determined by solution equilibrium titration (SET) as described above. The results are summarized in Table 9 and example titration curves for MOR10701 are contained in Figure 1. The data indicate that a number of antibodies were identified that
25 tightly bound human, cyno, rat and murine HER3.

Table 9: K_D values of anti-HER3 IgGs as determined by solution equilibrium titration (SET).
Hu (human), Cy (cynomolgus), Mu (murine) and ra (rat)

MOR#	SET K_D (pM)			
	hu HER3-Tag	cy HER3- Tag	mu HER3- Tag	ra HER3- Tag
09823	9	4	2	11
09824	3	3	2	7
09825	25	56	24	96
09974	350	200	120	n.d.
10701	4	4	6	10
10702	3	3	5	6
10703	26	23	20	40
12609	10	n.d	n.d	n.d
12610	37	n.d	n.d	n.d
10703 N52S	57	n.d	n.d	n.d
10703 N52G	60	n.d	n.d	n.d
10703_A50V_N52S	16	n.d	n.d	n.d
10703_A50V_N52G	22	n.d	n.d	n.d
10701 R55G	18	n.d	n.d	n.d
10701 R55K	11	n.d	n.d	n.d

(ii) *SK-Br-3 Cell EC₅₀ Determination*

- 5 The ability of the identified antibodies to bind HER3 expressing cells was determined by calculating EC_{50} values for their binding to the *HER2* amplified cell line SK-Br-3 (see Figure 2 and Table 10).

Table 10: FACS EC_{50} values of anti-HER3 IgG on SK-Br-3 cells. n.d. (not determined)

MOR#	SK-Br-3 FACS EC_{50} (pM)
09823	630
09824	324
09825	839
09974	n.d.
10701	n.d.
10702	n.d.
10703	2454

(iii) HER3 Domain Binding

A subset of anti-HER3 antibodies were characterized for their ability to bind the various extracellular domains of human HER3 in an ELISA assay. To achieve this, the extracellular domain of HER3 was divided into its four constitutive domains and various combinations of these domains were cloned, expressed and purified as independent proteins as described above. Using this strategy the following domains were successfully generated as soluble proteins: domains 1 and 2 (D1-2), domain 2 (D2), domains 3 and 4 (D3-4) and domain 4 (D4). A number of internally generated mouse anti-human HER3 antibodies (8D7, 1F5 and 8P2) were also tested as positive controls to demonstrate the integrity of each isolated domain.

As shown in Figure 3 MOR09823 and MOR09825 were both observed to successfully bind the HER3 extracellular domain, but little binding to the isolated domains was observed in this assay with these antibodies. There are several possible explanations for this binding pattern:

- a) MOR09823 and MOR09825 may bind a linear epitope that spans a domain boundary thus part of the binding epitope would be lost when the domains were expressed as isolated proteins.
- b) MOR09823 and MOR09825 may bind a non-linear epitope that bridges multiple domains. Consequently, separation of HER3 into its component units may destroy the binding site.
- c) The shape/conformation of HER3 may be a component of the binding of MOR09823 and MOR09825 to HER3 such that only the full-length extracellular domain of HER3 is capable of adopting this shape/ conformation whilst the isolated domains cannot fully assume this conformation.

(vi) HER3 Epitope Mapping Using Hydrogen/ Deuterium Exchange Mass Spectrometry

The HER3 epitope was further explored by HDX-MS analysis of HER3 ECD in the presence and absence of Fab versions of MOR09823, MOR09824, MOR09825 and MOR09974. Figure 4A shows that in the absence of bound Fab, approximately 69% of the HER3 ECD sequence was covered by at least one peptide. Gaps in coverage may be due to glycosylation of residues within these regions or insufficient reduction of disulphide bonds in cysteine rich regions, which is particularly apparent in domain 2. Interestingly, although each Fab yielded individual protection patterns, one region of strong protection was consistently observed with MOR09823, MOR09824, MOR09825 and MOR09974 (see Figure 4B) indicating that these highly related family of antibodies bind HER3 in an identical manner. The strongest

protection was observed for domain 2 residues 269-286 (TFQLEPNPHTKYQYGGVC) (SEQ ID NO: 146) indicating that residues in this vicinity may be important for mAb binding. Mapping of the Fab protected residues onto the published HER3 crystal structure (Cho & Leahy, (2002) Science 297:1330-1333) highlights that residues 269-286 are within and proximal to a functionally important β -hairpin loop within domain 2 (see Figure 4C).

(vii) *HER3/MOR09823 crystal structure*

The 3.2Å resolution x-ray crystal structure of MOR09823 Fab fragment bound to the HER3 extracellular domain was solved to further define the HER3 epitope that is recognized by this family of related antibodies (see Figure 5A). In addition, the 3.4Å structure of MOR09825

Fab fragment bound to human HER3 was resolved. In both the MOR09823/HER3 and MOR09825/HER3 crystal structures, HER3 is in the tethered (inactive) conformation (see Figure 5A, B, C and D). This conformation is characterized by a significant interaction interface between domains 2 and 4 mediated by a β -hairpin dimerization loop in domain 2.

The observed conformation of HER3 is similar to that previously described by Cho *et al.* (Cho & Leahy, (2002), Science 297:1330-1333) who published the crystal structure of the HER3 extra-cellular domain in the absence of neuregulin. Since neuregulin can activate HER3, the tethered conformation of HER3 is presumed to be inactive. Similar tethered conformations have also been observed when the related EGFR family members HER4 (Bouyain *et al.*, (2005) Proc. Natl. Acad. Sci. USA, 102:15024-15029) and HER1 (Ferguson *et al.*, (2003)

Molec. Cell 11:507-517) have been crystallized.

The spatial relationships between domains 1 to 4 of HER3 in the inactive (tethered) state are significantly different from that of the extended (active) state. This finding is based upon the crystal structures of the related EGFR family members HER2 and ligand-bound HER1 (Cho *et al.*, (2003) Nature 421:756-760; Ogiso *et al.*, (2002) Cell 110:775-787; Garrett *et al.*, (2002) Cell 110:763-773) both of which are in an extended (active) state. In the extended state, the domain 2 β -hairpin dimerization loop is released from its inhibitory interaction with 4 and is thus free to interact with its dimerization partner proteins. Thus, the domain 2 β -hairpin dimerization loop is functionally important both in maintaining the tethered (inactive) state and in mediating dimerization of EGF receptors in the extended state, leading to activation of the intracellular kinase domain. The MOR09823/HER3 and MOR09825/HER3 crystal structures (see Figure 5) therefore suggest that both MOR09823 and MOR09825 function by stabilizing the inactive conformation of HER3.

The crystal structure also revealed that the HER3 epitope recognized by both MOR09823 and MOR09825 is a non-linear epitope that includes residues from both domains 2 and 4 (see Figure 5C and D, Tables 11, 12, 13 and 14). The HER3 epitope recognized by this family of highly related antibodies can therefore be defined as:

5 Domain 2: residues 265-277, 315

Domain 4 residues: 571, 582-584, 596-597, 600-602, 609-615

Binding of both domains 2 and 4 by MOR09823 or MOR09825 would consequently stabilize the tethered conformation of HER3 thus antagonizing its ability to signal.

The MOR09823/ MOR09825 binding mode observed in the crystal structure is consistent with our other epitope mapping studies. Specifically, the ELISA domain binding experiments demonstrate that the affinity of MOR09823 and MOR09825 are significantly greater for the intact HER3 extracellular protein than for any isolated domains (e.g. D1, D1-D2, D3, or D3-D4 fragments) (see Figure 3). There is also agreement with the HER3 HDX-MS data (see Figure 4B), which identifies domain 2 β -hairpin as part of the antibody recognition epitope. Finally, both crystal structures indicate that the ligand-binding surface of HER3, which has been mapped by analogy to HER1 to domains 1 and 3 (Ogiso *et al.*, (2002) Cell, 110:775-787; Garrett *et al.*, (2002) Cell, 110:763-773) is not occluded by either MOR09823 or MOR09825 binding (see Figure 5B). This is consistent with our findings that neither MOR09823 nor MOR09825 block neuregulin binding to MCF7 cells (see Figure 9) and that HER3/MOR09823 complexes can bind to immobilized neuregulin in biacore studies (see Figure 10).

Table 11: Interactions between MOR09823 Fab heavy chain and human HER3. Fab VH residues are numbered based upon their linear amino acid sequence (SEQ ID NO: 15). HER3 residues are numbered based upon NP_001973. HER3 residues shown have at least one atom within 5Å of an atom in the MOR09823 Fab.

MOR09823 Fab			Human HER3		
Residue	Number	Chain	Residue	Number	Domain
Ser	30	VH	Pro	276	2
Ser	31	VH	Pro	274	2
			Asn	275	2
			Pro	276	2
Tyr	32	VH	Pro	276	2

			His	277	2
Ala	33	VH	Asn	266	2
			Leu	268	2
Ser	35	VH	Leu	268	2
Val	50	VH	Leu	268	2
			Thr	269	2
Gly	52	VH	Glu	273	2
			Thr	269	2
Ala	53	VH	Glu	273	2
			Pro	274	2
Val	54	VH	Glu	273	2
Tyr	58	VH	Pro	583	4
			Asp	571	4
			His	584	4
			Thr	269	2
			Gln	271	2
Asn	73	VH	Asn	315	2
Ser	74	VH	Asn	315	2
Trp	98	VH	Leu	268	2
			Lys	267	2
			Asn	266	2
Asp	100	VH	Ala	596	4
			Lys	597	4
			Pro	276	2
			His	277	2
Glu	101	VH	Lys	267	2
			Lys	597	4
Phe	103	VH	Leu	268	2

Table 12: Interactions between MOR09823 Fab light chain and human HER3. Fab VL residues are numbered based upon their linear amino acid sequence (SEQ ID NO: 14). HER3 residues are numbered based upon NP_001973. HER3 residues shown have at least one atom within 5Å of an atom in the MOR09823 Fab.

5

MOR09823 Fab			Human HER3		
Residue	Number	Chain	Residue	Number	Domain
Gln	27	VL	Arg	611	4
			Glu	609	4
Gly	28	VL	Arg	611	4
			Pro	612	4
Ile	29	VL	Pro	612	4
Ser	30	VL	Pro	612	4
			Cys	613	4
			His	614	4
			Glu	615	4
Asn	31	VL	Glu	615	4
			Cys	613	4
Trp	32	VL	Lys	267	2
			Tyr	265	2
			Pro	612	4
			Cys	613	4
			Ile	600	4
			Lys	602	4
Tyr	49	VL	Lys	597	4
Gly	66	VL	Glu	615	4
Ser	67	VL	His	614	4
			Glu	615	4
Gln	89	VL	Leu	268	2
Tyr	91	VL	Lys	267	2
			Leu	268	2
			Phe	270	2
Ser	92	VL	Phe	270	2
			Lys	602	4
			Pro	612	4
Ser	93	VL	Phe	270	2
			Glu	609	4
Phe	94	VL	Phe	270	2
			Leu	268	2
			Gly	582	4
			Pro	583	4

Thr	96	VL	Leu	268	2
-----	----	----	-----	-----	---

Table 13: Interactions between MOR09825 Fab heavy chain and human HER3. Fab VH residues are numbered based upon their linear amino acid sequence (SEQ ID NO: 51). HER3 residues are numbered based upon NP_001973. HER3 residues shown have at least one atom within 5 Å of an atom in the MOR09825 Fab.

MOR09825 Fab			Human HER3		
Residue	Number	Chain	Residue	Number	Domain
Ser	30	VH	Asn	315	2
Ser	31	VH	Pro	274	2
			Pro	276	2
Tyr	32	VH	Pro	276	2
			His	277	2
Ala	33	VH	Asn	266	2
			Thr	269	2
Ser	35	VH	Leu	268	2
Trp	47	VH	Leu	268	2
Ala	50	VH	Leu	268	2
Asn	52	VH	Glu	273	2
			Gln	271	2
			Thr	269	2
Ser	53	VH	Glu	273	2
			Pro	274	2
Gln	54	VH	Glu	273	2
			Pro	274	2
Ser	57	VH	Gln	271	2
Tyr	59	VH	Pro	583	4
			Asp	571	4
			His	584	4
			Thr	269	2
			Gln	271	2
Asn	74	VH	Asn	315	2
Trp	99	VH	Leu	268	2
			Lys	267	2
			Asn	266	2

Asp	101	VH	Ala	596	4
			Lys	597	4
			Pro	276	2
			His	277	2
Glu	102	VH	Lys	267	2
			Lys	597	4
Phe	104	VH	Leu	268	2

Table 14: Interactions between MOR09825 Fab light chain and human HER3. Fab VL residues are numbered based upon their linear amino acid sequence (SEQ ID NO: 50). HER3 residues are numbered based upon NP_001973. HER3 residues shown have at least one atom within 5Å of an atom in the MOR09825 Fab.

MOR09825 Fab			Human HER3		
Residue	Number	Chain	Residue	Number	Domain
Gln	27	VL	Arg	611	4
Gly	28	VL	Arg	611	4
			Pro	612	4
Ile	29	VL	Pro	612	4
Ser	30	VL	Pro	612	4
			Cys	613	4
			His	614	4
			Glu	615	4
Asn	31	VL	Glu	615	4
			His	614	4
			Cys	613	4
Trp	32	VL	Lys	267	2
			Tyr	265	2
			Pro	612	4
			Cys	613	4
			Ile	600	4
			Lys	602	4
Tyr	49	VL	Lys	597	4
Gly	66	VL	Glu	615	4
Ser	67	VL	His	614	4
			Glu	615	4

Gln	89	VL	Leu	268	2
Tyr	91	VL	Lys	267	2
			Leu	268	2
			Phe	270	2
Ser	92	VL	Phe	270	2
			Lys	602	4
			Pro	612	4
			Arg	611	4
Ser	93	VL	Phe	270	2
			Glu	609	4
Phe	94	VL	Phe	270	2
			Gly	582	4
			Pro	583	4
Thr	96	VL	Leu	268	2

Visual inspection of the MOR09823/ MOR09825 crystal structures highlighted that HER3 residues Lys267 and Leu268 formed multiple interactions with various antibody CDR's suggesting that they may be important for antibody binding. Consequently, Lys267 and/ or Leu268 were mutated to alanine, expressed and the resultant recombinant proteins purified in order to assess their impact upon antibody binding. ELISA binding assays indicated that mutation of either Lys267 or Leu268 abolished MOR10703 binding to HER3 (Figure 5F) suggesting that both residues are an integral part of the HER3 epitope and thus supporting the proposed interactions between MOR09823/ MOR09825 and HER3.

10 (viii) Inhibition of Cell Signaling

To ascertain the effect of anti-HER3 antibodies upon ligand dependent HER3 activity MCF7 cells were incubated with IgG prior to stimulation with neuregulin. Example inhibition curves are illustrated in Figure 6A and summarized in Table 15. The effect of anti-HER3 antibodies upon HER2- mediated HER3 activation was also studied using the *HER2* amplified cell line SK-Br-3 (Figure 6B and Table 15).

Table 15: pHER3 IC₅₀ and extent of inhibition values of anti-HER3 IgG in MCF7, and SK-Br-3 cells.

MOR#	MCF7 pHER3		SK-Br-3 pHER3	
	IC ₅₀ (pM)	% inhibition	IC ₅₀ (pM)	% inhibition
09823	181	89	56	59
09824	103	91	110	64
09825	399	80	169	66
09974	3066	69	1928	67
10701	n.d.	n.d.	370	74
10702	n.d.	n.d.	n.d.	n.d.
10703	333	80	167	69
12609	5	86	241	71
12610	126	84	192	75

To determine whether inhibition of HER3 activity impacted downstream cell signaling Akt, phosphorylation was also measured in *HER2* amplified cells following treatment with anti-HER3 antibodies (see Figure 7 and Table 16).

Table 16: pAkt (S⁴⁷³) IC₅₀ and extent of inhibition values of anti-HER3 IgG in SK-Br-3 BT-474 and MCF7 cells.

MOR#	SK-Br-3 pAkt		BT-474 pAkt	MCF7 pAkt	
	IC ₅₀ (pM)	% inhibition	% inhibition	IC ₅₀ (pM)	% inhibition
09823	55	92	57	n.d.	n.d.
09824	62	93	46	n.d.	n.d.
09825	156	91	69	294	79
09974	814	85	n.d.	n.d.	n.d.
10701	n.d.	n.d.	59	n.d.	n.d.
10702	n.d.	n.d.	55	n.d.	n.d.
10703	70	89	62	449	79

In summary MOR09823, MOR09824, MOR09825, MOR09974, MOR10701, MOR10702 MOR10703, MOR12609 and MOR12610 are each capable of inhibiting cellular HER3 activity in both a ligand-dependent and ligand-independent manner.

(ix) Inhibition of Proliferation

- 5 Since MOR09823, MOR09824, MOR09825, MOR09974, MOR10701, MOR10702 and MOR10703 all inhibited HER3 activity and downstream signaling they were tested for their ability to block ligand dependent and independent *in vitro* cell growth (Example data is shown in Figure 8 and summarized in Table 17). The anti-HER3 antibodies tested were all effective inhibitors of cell proliferation.
- 10 Table 17: Inhibition of proliferation following treatment with 10 µg/ml anti-HER3 IgG in SK-Br-3, BT-474 and MCF7 cells.

MOR#	% Inhibition		
	SK-Br-3	BT-474	MCF7
09823	39	39.8	82
09824	33	36.8	82
09825	41	37.2	63
09974	35	n.d.	20
10701	n.d.	43.6	n.d.
10702	n.d.	43.8	n.d.
10703	35	41.6	81

(x) Ligand blocking assessment

- The ability of the described anti-HER3 antibodies to block ligand binding was assessed by
 15 examining the binding of neuregulin to MCF7 cells previously treated with either MOR09823 or MOR09825. The presence of either MOR09823 or MOR09825 had no significant effect upon the ability of neuregulin to bind MCF7 cells whilst the positive control used in the experiment (Mab3481) was capable of profoundly interfering with neuregulin binding (see Figure 9). These results are consistent with the crystal structure since MOR09823 interacts
 20 with domains 2 and 4 whilst the major contact points for HER3's interaction with neuregulin are hypothesized to be primarily clustered within domains 1 and 3. Given that neuregulin is capable of binding the inactive conformation of HER3 (Kani *et al.*, (2005) *Biochemistry* 44:

15842-15857) it is probable that MOR09823 and MOR09825 function by preventing the HER3 domain rearrangements necessary for signaling or by interfering with receptor dimerization.

(xi) Ligand blocking assessment (biochemical)

5 To explore whether MOR09823 and neuregulin can bind HER3 concurrently a biochemical assay was established using Biacore™ technology. Interaction analyses were performed by capturing biotinylated neuregulin on the surface of a Biacore™ sensor chip CAP (GE Healthcare) utilizing a Biotin CAPture kit (GE Healthcare). HER3 complexes were generated by incubating human HER3-Fc with increasing concentrations of either MOR09823, 105.5
10 (Thermo Scientific) or human IgG. Preformed HER3/ antibody complexes were injected over reference and active surfaces and the interaction of HER3 with neuregulin observed.

Control IgG had no effect upon HER3/ neuregulin complex formation whilst 105.5 was observed to significantly inhibit the ability of HER3 to bind neuregulin confirming its description as a ligand blocking antibody (Figure 10). In contrast HER3/ MOR09823
15 complexes were capable of binding neuregulin demonstrating that MOR09823 does not prevent ligand binding. Interestingly, a dose-dependent increase in RU values was uniquely observed when MOR09823/ HER3 complexes were injected. This data indicates that a trimeric complex containing neuregulin, HER3 and MOR09823 is generated on the chip surface. The ability of this trimeric complex to form is predicted by the HER3/ MOR09823
20 crystal structure since MOR09823 binding does not occlude the ligand binding site of HER3 suggesting that binding of neuregulin and MOR09823 are not mutually exclusive.

In another embodiment, the antibody or fragment thereof binds to both domain 2 and domain 4 of HER3 and without blocking the concurrent binding of a HER3 ligand such as neuregulin. While not required to provide a theory, it is feasible that the antibody or fragment thereof
25 binding to both domain 2 and domain 4 of HER3, holds HER3 in an inactive conformation without blocking the ligand binding site on HER3. Thus a HER3 ligand (e.g., neuregulin) is able to bind to HER3 at the same time as the antibody.

The antibodies of the invention or fragments thereof inhibit both ligand dependent and independent activation of HER3 without preventing ligand binding. This is considered
30 advantageous for the following reasons:

(i) The therapeutic antibody would have clinical utility in a broad spectrum of tumors than an antibody which targeted a single mechanism of HER3 activation (i.e. ligand dependent or ligand independent) since distinct tumor types are driven by each mechanism.

(ii) The therapeutic antibody would be efficacious in tumor types where both mechanisms of HER3 activation are simultaneously involved. An antibody targeting a single mechanism of HER3 activation (i.e. ligand dependent or ligand independent) would display little or no efficacy in these tumor types

(iii) The efficacy of an antibody which inhibits ligand dependent activation of HER3 without preventing ligand binding would be less likely to be adversely affected by increasing concentrations of ligand. This would translate to either increased efficacy in a tumor type driven by very high concentrations of HER3 ligand or a reduced drug resistance liability where resistance is mediated by up-regulation of HER3 ligands.

(iv) An antibody which inhibits HER3 activation by stabilizing the inactive form would be less prone to drug resistance driven by alternative mechanisms of HER3 activation.

Consequently, the antibodies of the invention may be used to treat conditions where existing therapeutic antibodies are clinically ineffective.

(xii) In vivo inhibition of HER3 activity and effect upon tumor growth

To determine the *in vivo* activity of the described anti-HER3 antibodies, MOR09823 was tested in both BxPC-3 and BT-474 tumor models. MOR09823 was demonstrated to inhibit HER3 activity as evidenced by a significant reduction in tumor pHER3 levels (Figure 11). Signaling downstream of HER3 was similarly inhibited as demonstrated by reduced pAkt levels in both BxPC-3 and BT-474 (Figure 11). In a HER2 driven BT-474 efficacy study, repeated MOR10701 treatment yielded a 74% inhibition of tumor growth (see Figure 12A) whilst MOR10703 yielded 83% inhibition. In the BxPC3 tumor growth model, both MOR10701 and MOR10703 very effectively inhibited ligand driven tumor growth (see Figure 13).

(xiii) In vitro drug combinations and impact upon cell growth.

Since tumor cell growth is frequently driven by multiple signaling pathways we assessed whether combinations of MOR09823 or MOR10703 with various targeted agents would be of benefit in blocking cell proliferation. The targeted agents chosen primarily inhibited HER2 (trastuzumab, lapatinib) EGFR (cetuximab, erlotinib), PI3K/mTOR (BEZ235), PI3K

(BKM120), PIK3CA (BYL719) and mTOR (RAD001) since these targets are commonly activated in human tumors. Isobologram analysis (see Figure 14) indicated that MOR09823 and MOR10703 displayed synergistic drug combinations with trastuzumab, lapatinib, erlotinib, cetuximab, BEZ235, BKM120, BYL719 and RAD001. This data suggests that inhibition of HER3 signaling is of particular benefit to inhibitors that target receptor tyrosine kinases or the PI3K signaling pathway.

(xiv) In vivo MOR10703 drug combinations

Since HER3 inhibition combined with receptor tyrosine kinase targeted agents *in vitro* we assessed the impact of either MOR10701 or MOR10703 in combination with trastuzumab and erlotinib *in vivo*. In BT-474 xenografts (see Figure 15A), combination of either MOR10701 or MOR10703 (20mg/kg) with a sub-optimal dose of trastuzumab (1mg/kg) was sufficient to induce tumor regressions (%T/C= -50 and -37 respectively). In L3.3 pancreatic xenografts, combination of MOR10703 (20mg/kg) with daily erlotinib (50mg/kg) resulted in tumor stasis (%T/C= 3, see Figure 15B). In both models, the combination of two drugs was significantly more efficacious than either drug alone thus supporting our earlier *in vitro* finding of the benefit of combining HER3-targeted antibodies with ErbB-targeted agents.

In summary, the unique ability of this family of antibodies to stabilize the inactive conformation of HER3 results in significant *in vivo* efficacy in models where HER3 is activated in either a ligand dependent or independent manner. Furthermore, HER3 inhibition by this family of antibodies appears beneficial in combination with a wide variety of targeted therapies.

Incorporation By reference

All references cited herein, including patents, patent applications, papers, text books, and the like, and the references cited therein, to the extent that they are not already, are hereby incorporated herein by reference in their entirety.

Equivalents

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The foregoing description and examples detail certain preferred embodiments of the invention and describe the best mode contemplated by the inventors. It will be appreciated, however, that no matter how detailed the foregoing may appear in text, the invention may be practiced in many ways and the invention should be construed in accordance with the appended claims and any equivalents thereof.

Claims:

1. An isolated antibody or fragment thereof that binds to an inactive state of a HER receptor, wherein the antibody blocks both ligand-dependent and ligand-independent signal transduction.
5
2. The isolated antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof stabilizes the HER receptor in an inactive state.
3. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, and wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction.
10
4. The isolated antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof binds to the inactive state of the HER receptor.
5. The isolated antibody or fragment thereof of claim 4, wherein the antibody or fragment thereof stabilizes the HER receptor in the inactive state.
15
6. The isolated antibody or fragment thereof of claim 2, wherein the HER receptor is selected from the group consisting of HER1, HER2, HER3 and HER4.
7. The isolated antibody or fragment thereof of claim 3, wherein the antibody is selected from the group consisting of a monoclonal antibody, a polyclonal antibody, a chimeric antibody, a humanized antibody, and a synthetic antibody.
20
8. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, wherein binding of the antibody stabilizes the HER receptor in an inactive state, and wherein a HER ligand can concurrently bind to a ligand binding site on the HER receptor.
25
9. The isolated antibody or fragment thereof of claim 8, wherein HER ligand binding to the ligand binding site fails to induce a conformational change in the HER receptor to an active state.

10. The isolated antibody or fragment thereof of claim 8, wherein HER ligand binding to the ligand binding site fails to activate signal transduction.
11. The isolated antibody or fragment thereof of claim 8, wherein the HER ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, neuregulin 3, neuregulin 4, betacellulin, heparin-binding epidermal growth factor, epiregulin, epidermal growth factor, amphiregulin, and transforming growth factor alpha.
12. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER receptor, wherein binding of the antibody stabilizes the HER receptor in an inactive state such that the HER receptor fails to dimerize with a co-receptor to form a receptor-receptor complex.
13. The isolated antibody or fragment thereof of claim 12, wherein the failure to form a receptor-receptor complex prevents activation of signal transduction.
14. The isolated antibody or fragment thereof of claim 12, wherein signal transduction is ligand-independent signal transduction.
15. The isolated antibody or fragment thereof of claim 12, wherein signal transduction is ligand-dependent signal transduction.
16. An isolated antibody or fragment thereof that binds to an inactive conformation of a HER3 receptor, wherein the antibody blocks both ligand-dependent and ligand-independent signal transduction.
17. The isolated antibody or fragment thereof of claim 16, wherein the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state.
18. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction.
19. The isolated antibody or fragment thereof of claim 18, wherein the antibody or fragment thereof binds to the inactive state of the HER3 receptor.
20. The isolated antibody or fragment thereof of claim 18, wherein the antibody or fragment thereof stabilizes the HER3 receptor in the inactive state.

21. The isolated antibody or fragment thereof of claim 18, wherein the antibody is selected from the group consisting of a monoclonal antibody, a polyclonal antibody, a chimeric antibody, a humanized antibody, and a synthetic antibody.
22. An isolated antibody or fragment thereof that recognizes a conformational epitope of a
5 HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein binding of the antibody stabilizes the HER3 receptor in an inactive state, and wherein a HER3 ligand can concurrently bind to a ligand binding site on the HER3 receptor.
23. The isolated antibody or fragment thereof of claim 22, wherein HER3 ligand binding to
10 the ligand binding site fails to induce a conformational change in the HER3 receptor to an active state.
24. The isolated antibody or fragment thereof of claim 22, wherein the HER3 ligand binding to the ligand binding site fails to activate signal transduction.
25. The isolated antibody or fragment thereof of claim 22, wherein the HER3 ligand is
15 selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, betacellulin, heparin-binding epidermal growth factor, and epiregulin.
26. An isolated antibody or fragment thereof that recognizes a conformational epitope of a
20 HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof blocks both ligand-dependent and ligand-independent signal transduction.
27. The isolated antibody or fragment thereof of claim 26, wherein the antibody or fragment thereof binds to the inactive state of the HER3 receptor.
28. The isolated antibody or fragment thereof of claim 27, wherein the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state.
- 25 29. An isolated antibody or fragment thereof that binds a conformational epitope of HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein the antibody or fragment thereof binds to the inactive HER3 receptor, and wherein the antibody or fragment blocks both ligand-dependent and ligand-independent signal transduction.

30, The isolated antibody or fragment thereof of claim 28, wherein the antibody or fragment thereof stabilizes the HER3 receptor in an inactive state.

31. The isolated antibody of claim 29, wherein the conformational epitope comprises amino acid residues 265-277, 315 (of domain 2), 571, 582-584, 596-597, 600-602, 609-615 (of domain 4), or a subset thereof.

32. The isolated antibody of claim 29, wherein the VH of the antibody or fragment thereof binds to at least one of the following HER3 residues: Asn266, Lys267, Leu268, Thr269, Gln271, Glu273, Pro274, Asn275, Pro276, His277, Asn315, Asp571, Pro583, His584, Ala596, Lys597.

33. The isolated antibody of claim 29, wherein the VL of the antibody or fragment thereof binds to at least one of the following HER3 residues: Tyr265, Lys267, Leu268, Phe270, Gly582, Pro583, Lys597, Ile600, Lys602, Glu609, Arg611, Pro612, Cys613, His614, Glu615.

34. An isolated antibody or fragment thereof that recognizes a conformational epitope of first HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the first HER receptor, wherein binding of the antibody or fragment thereof to the first HER receptor in the absence of a HER receptor ligand reduces ligand-independent formation of a first HER receptor-second HER receptor protein complex in a cell which expresses first HER receptor and second HER receptor.

35. The isolated antibody or fragment thereof of claim 34, wherein antibody or fragment thereof stabilizes first HER receptor in an inactive state such that the first HER receptor fails to dimerize with the second HER receptor to form a first HER receptor-second HER receptor protein complex.

36. The isolated antibody or fragment thereof of claim 35, wherein the failure to form a first HER receptor-second HER receptor protein complex prevents activation of signal transduction.

37. The isolated antibody or fragment thereof of claim 35, wherein the first HER is selected from the group consisting of HER1, HER2, HER3, and HER4.

38. The isolated antibody or fragment thereof of claim 35, wherein the second HER is selected from the group consisting of HER1, HER2, HER3, and HER4.

39. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein binding of the antibody or fragment thereof to the HER3 receptor in the absence of a HER3 ligand reduces ligand-independent formation of a HER2-HER3 protein complex in a cell which expresses HER2 and HER3.
40. The isolated antibody or fragment thereof of claim 39, wherein antibody or fragment thereof stabilizes the HER3 receptor in an inactive state such that the HER3 receptor fails to dimerize with the HER2 receptor to form a HER2-HER3 protein complex.
41. The isolated antibody or fragment thereof of claim 40, wherein the failure to form a HER2-HER3 protein complex prevents activation of signal transduction.
42. An isolated antibody or fragment thereof that recognizes a conformational epitope of first HER receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the first HER receptor, wherein binding of the antibody or fragment thereof to the first HER receptor in the presence of a HER ligand reduces ligand-dependent formation of a first HER receptor-second HER receptor protein complex in a cell which expresses first HER receptor and second HER receptor.
43. The isolated antibody or fragment thereof of claim 42, wherein antibody or fragment thereof stabilizes the first HER receptor in an inactive state such that the HER receptor fails to dimerize with the second HER receptor in the presence of a first HER ligand to form a first HER receptor- second HER receptor protein complex.
44. The isolated antibody or fragment thereof of claim 43, wherein the failure to form a first HER receptor- second HER receptor protein complex prevents activation of signal transduction.
45. The isolated antibody or fragment thereof of claim 42, wherein the HER ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, neuregulin 3, neuregulin 4, betacellulin, heparin-binding epidermal growth factor, epiregulin, epidermal growth factor, amphiregulin, and transforming growth factor alpha.
46. The isolated antibody or fragment thereof of claim 42, wherein the first HER is selected from the group consisting of HER1, HER2, HER3, and HER4.

47. The isolated antibody or fragment thereof of claim 42, wherein the second HER is selected from the group consisting of HER1, HER2, HER3, and HER4.
48. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, wherein binding of the antibody or fragment thereof to the HER3 receptor in the presence of a HER3 ligand reduces ligand-dependent formation of a HER2-HER3 protein complex in a cell which expresses HER2 and HER3.
49. The isolated antibody or fragment thereof of claim 48, wherein antibody or fragment thereof stabilizes the HER3 receptor in an inactive state such that the HER3 receptor fails to dimerize with the HER2 receptor in the presence of a HER3 ligand to form a HER2-HER3 protein complex.
50. The isolated antibody or fragment thereof of claim 49, wherein the failure to form a HER2-HER3 protein complex prevents activation of signal transduction.
51. The isolated antibody or fragment thereof of claim 48, wherein ligand is selected from the group consisting of neuregulin 1 (NRG), neuregulin 2, betacellulin, heparin-binding epidermal growth factor, and epiregulin.
52. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof inhibits phosphorylation of HER3 as assessed by HER3 ligand-independent phosphorylation assay.
53. The isolated antibody or fragment thereof of claim 52, wherein the HER3 ligand-independent phosphorylation assay uses HER2 amplified cells, wherein the HER2 amplified cells are SK-Br-3 cells.
54. An isolated antibody or fragment thereof that recognizes a conformational epitope of a HER3 receptor, wherein the conformational epitope comprises amino acid residues within domain 2 and domain 4 of the HER3 receptor, and wherein the antibody or fragment thereof inhibits phosphorylation of HER3 as assessed by HER3 ligand-dependent phosphorylation assay.

55. The isolated antibody or fragment thereof of claim 54, wherein the HER3 ligand-dependent phosphorylation assay uses stimulated MCF7 cells in the presence of neuregulin (NRG).
56. An isolated antibody or fragment thereof to a HER3 receptor, having a dissociation (K_D) of at least $1 \times 10^7 M^{-1}$, $10^8 M^{-1}$, $10^9 M^{-1}$, $10^{10} M^{-1}$, $10^{11} M^{-1}$, $10^{12} M^{-1}$, $10^{13} M^{-1}$.
57. The isolated antibody or fragment thereof of claim 56, wherein the antibody or fragment thereof inhibits phosphorylation of HER3 as measured by an in vitro phosphorylation assay selected from the group consisting of phospho-HER3 and phospho-Akt.
58. An isolated antibody or fragment thereof to HER3 receptor, that cross-competes with an antibody described in Table 1.
59. An isolated antibody or fragment thereof, that binds to the same conformational epitope as an antibody described in Table 1.
60. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody is selected from the group consisting of a monoclonal antibody, chimeric antibody, a single chain antibody, an Fab, and an scFv.
61. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody is a human.
62. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody comprises a human heavy chain constant region and a human light chain constant region.
63. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody or fragment binds to both human HER3 and cynomolgus HER3.
64. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody or fragment is an IgG isotype.
65. The isolated antibody or fragment thereof of any of the above claims, wherein the antibody comprises a framework in which amino acids have been substituted into the antibody framework from the respective human VH or VL germline sequences.
66. An isolated antibody or fragment thereof to HER3 receptor comprising 1, 2, 3, 4, 5, or 6 CDRs calculated by Kabat or Chothia of any of the antibodies in Table 1.

67. An isolated antibody or fragment thereof to HER3 receptor comprising a heavy chain CDR3 selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 10, SEQ ID NO: 22, SEQ ID NO: 28, SEQ ID NO: 40, SEQ ID NO: 46, SEQ ID NO: 58, SEQ ID NO: 64, SEQ ID NO: 76, SEQ ID NO: 82, SEQ ID NO: 94, SEQ ID NO: 100, SEQ ID NO: 112, SEQ ID NO: 118, SEQ ID NO: 130, SEQ ID NO: 136, SEQ ID NO: 148, SEQ ID NO: 166, SEQ ID NO: 184, SEQ ID NO: 202, SEQ ID NO: 220, SEQ ID NO: 238, SEQ ID NO: 256, SEQ ID NO: 274, SEQ ID NO: 292, SEQ ID NO: 310, SEQ ID NO: 328, SEQ ID NO: 346, and SEQ ID NO: 364.
68. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 15 and a VL comprising SEQ ID NO: 14, or an amino acid sequence with 97-99% identity thereof.
69. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 33 and a VL comprising SEQ ID NO: 32, or an amino acid sequence with 97-99% identity thereof.
70. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 51 and a VL comprising SEQ ID NO: 50, or an amino acid sequence with 97-99% identity thereof.
71. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 69 and a VL comprising SEQ ID NO: 68, or an amino acid sequence with 97-99% identity thereof.
72. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 87 and a VL comprising SEQ ID NO: 86, or an amino acid sequence with 97-99% identity thereof.
73. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 105 and a VL comprising SEQ ID NO: 104, or an amino acid sequence with 97-99% identity thereof.
74. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 123 and a VL comprising SEQ ID NO: 122, or an amino acid sequence with 97-99% identity thereof.

75. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 141 and a VL comprising SEQ ID NO: 140, or an amino acid sequence with 97-99% identity thereof.
76. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a
5 VH comprising SEQ ID NO: 159 and a VL comprising SEQ ID NO: 158, or an amino acid sequence with 97-99% identity thereof.
77. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 177 and a VL comprising SEQ ID NO: 176, or an amino acid sequence with 97-99% identity thereof.
- 10 78. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 195 and a VL comprising SEQ ID NO: 194, or an amino acid sequence with 97-99% identity thereof.
79. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a
15 VH comprising SEQ ID NO: 213 and a VL comprising SEQ ID NO: 212, or an amino acid sequence with 97-99% identity thereof.
80. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 231 and a VL comprising SEQ ID NO: 230, or an amino acid sequence with 97-99% identity thereof.
81. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a
20 VH comprising SEQ ID NO: 249 and a VL comprising SEQ ID NO: 248, or an amino acid sequence with 97-99% identity thereof.
82. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 267 and a VL comprising SEQ ID NO: 266, or an amino acid sequence with 97-99% identity thereof.
- 25 83. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 285 and a VL comprising SEQ ID NO: 284, or an amino acid sequence with 97-99% identity thereof.
84. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a
30 VH comprising SEQ ID NO: 303 and a VL comprising SEQ ID NO: 302, or an amino acid sequence with 97-99% identity thereof.

85. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 321 and a VL comprising SEQ ID NO: 320, or an amino acid sequence with 97-99% identity thereof.
86. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a
5 VH comprising SEQ ID NO: 339 and a VL comprising SEQ ID NO: 338, or an amino acid sequence with 97-99% identity thereof.
87. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 357 and a VL comprising SEQ ID NO: 356, or an amino acid sequence with 97-99% identity thereof.
- 10 88. An isolated antibody or fragment thereof to HER3 receptor which antibody comprises a VH comprising SEQ ID NO: 375 and a VL comprising SEQ ID NO: 374, or an amino acid sequence with 97-99% identity thereof.
89. An isolated antibody or fragment thereof comprising a variable heavy chain sequence having SEQ ID NO: 493.
- 15 90. An isolated antibody or fragment thereof comprising a variable light chain sequence having SEQ ID NO: 494.
91. An isolated antibody or fragment thereof comprising a variable heavy chain sequence having SEQ ID NO: 493 and a variable light chain sequence having SEQ ID NO: 494.
92. An isolated antibody or fragment thereof to HER3 receptor with a variant heavy chain
20 variable region comprising CDR1, CDR2, and CDR3, wherein the variant has at least one to four amino acid changes in one of CDR1, CDR2, or CDR3.
93. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 2; CDR2 of SEQ ID NO: 3; CDR3 of SEQ ID NO: 4; a light chain variable region CDR1 of SEQ ID NO: 5; CDR2 of SEQ ID NO: 6; and CDR3 of SEQ ID NO: 7.
- 25 94. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 20; CDR2 of SEQ ID NO: 21; CDR3 of SEQ ID NO: 22; a light chain variable region CDR1 of SEQ ID NO: 23; CDR2 of SEQ ID NO: 24; and CDR3 of SEQ ID NO: 25.
95. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1
30 of SEQ ID NO: 38; CDR2 of SEQ ID NO: 39; CDR3 of SEQ ID NO: 40; a light chain

variable region CDR1 of SEQ ID NO: 41; CDR2 of SEQ ID NO: 42; and CDR3 of SEQ ID NO: 43.

96. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 56; CDR2 of SEQ ID NO: 57; CDR3 of SEQ ID NO: 58; a light chain
5 variable region CDR1 of SEQ ID NO: 59; CDR2 of SEQ ID NO: 60; and CDR3 of SEQ ID NO: 61.

97. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 74; CDR2 of SEQ ID NO: 75; CDR3 of SEQ ID NO: 76; a light chain
10 variable region CDR1 of SEQ ID NO: 77; CDR2 of SEQ ID NO: 78; and CDR3 of SEQ ID NO: 79.

98. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 92; CDR2 of SEQ ID NO: 93; CDR3 of SEQ ID NO: 94; a light chain
variable region CDR1 of SEQ ID NO: 95; CDR2 of SEQ ID NO: 96; and CDR3 of SEQ ID NO: 97.

15 99. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 110; CDR2 of SEQ ID NO: 111; CDR3 of SEQ ID NO: 112; a light chain
variable region CDR1 of SEQ ID NO: 113; CDR2 of SEQ ID NO: 114; and CDR3 of SEQ ID NO: 115.

100. An isolated antibody or fragment thereof, comprising a heavy chain variable region
20 CDR1 of SEQ ID NO: 128; CDR2 of SEQ ID NO: 129; CDR3 of SEQ ID NO: 130; a light
chain variable region CDR1 of SEQ ID NO: 131; CDR2 of SEQ ID NO: 132; and CDR3 of
SEQ ID NO: 133.

101. An isolated antibody or fragment thereof, comprising a heavy chain variable region
25 CDR1 of SEQ ID NO: 146; CDR2 of SEQ ID NO: 147; CDR3 of SEQ ID NO: 148; a light
chain variable region CDR1 of SEQ ID NO: 149; CDR2 of SEQ ID NO: 150; and CDR3 of
SEQ ID NO: 151.

102. An isolated antibody or fragment thereof, comprising a heavy chain variable region
30 CDR1 of SEQ ID NO: 164; CDR2 of SEQ ID NO: 165; CDR3 of SEQ ID NO: 166; a light
chain variable region CDR1 of SEQ ID NO: 167; CDR2 of SEQ ID NO: 168; and CDR3 of
SEQ ID NO: 169.

103. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 182; CDR2 of SEQ ID NO: 183; CDR3 of SEQ ID NO: 184; a light chain variable region CDR1 of SEQ ID NO: 185; CDR2 of SEQ ID NO: 186; and CDR3 of SEQ ID NO: 187.
- 5 104. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 200; CDR2 of SEQ ID NO: 201; CDR3 of SEQ ID NO: 202; a light chain variable region CDR1 of SEQ ID NO: 203; CDR2 of SEQ ID NO: 204; and CDR3 of SEQ ID NO: 205.
105. An isolated antibody or fragment thereof, comprising a heavy chain variable region
10 CDR1 of SEQ ID NO: 218; CDR2 of SEQ ID NO: 219; CDR3 of SEQ ID NO: 220; a light chain variable region CDR1 of SEQ ID NO: 221; CDR2 of SEQ ID NO: 222; and CDR3 of SEQ ID NO: 223.
106. An isolated antibody or fragment thereof, comprising a heavy chain variable region
15 CDR1 of SEQ ID NO: 236; CDR2 of SEQ ID NO: 237; CDR3 of SEQ ID NO: 238; a light chain variable region CDR1 of SEQ ID NO: 239; CDR2 of SEQ ID NO: 240; and CDR3 of SEQ ID NO: 241.
107. An isolated antibody or fragment thereof, comprising a heavy chain variable region
20 CDR1 of SEQ ID NO: 254; CDR2 of SEQ ID NO: 255; CDR3 of SEQ ID NO: 256; a light chain variable region CDR1 of SEQ ID NO: 257; CDR2 of SEQ ID NO: 258; and CDR3 of SEQ ID NO: 259.
108. An isolated antibody or fragment thereof, comprising a heavy chain variable region
CDR1 of SEQ ID NO: 272; CDR2 of SEQ ID NO: 273; CDR3 of SEQ ID NO: 274; a light chain variable region CDR1 of SEQ ID NO: 275; CDR2 of SEQ ID NO: 276; and CDR3 of SEQ ID NO: 277.
- 25 109. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 290; CDR2 of SEQ ID NO: 291; CDR3 of SEQ ID NO: 292; a light chain variable region CDR1 of SEQ ID NO: 293; CDR2 of SEQ ID NO: 294; and CDR3 of SEQ ID NO: 295.
110. An isolated antibody or fragment thereof, comprising a heavy chain variable region
30 CDR1 of SEQ ID NO: 308; CDR2 of SEQ ID NO: 309; CDR3 of SEQ ID NO: 310; a light

chain variable region CDR1 of SEQ ID NO: 311; CDR2 of SEQ ID NO: 312; and CDR3 of SEQ ID NO: 313.

111. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 326; CDR2 of SEQ ID NO: 327; CDR3 of SEQ ID NO: 328; a light chain variable region CDR1 of SEQ ID NO: 329; CDR2 of SEQ ID NO: 330; and CDR3 of SEQ ID NO: 331.

112. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 344; CDR2 of SEQ ID NO: 345; CDR3 of SEQ ID NO: 346; a light chain variable region CDR1 of SEQ ID NO: 347; CDR2 of SEQ ID NO: 348; and CDR3 of SEQ ID NO: 349.

113. An isolated antibody or fragment thereof, comprising a heavy chain variable region CDR1 of SEQ ID NO: 362; CDR2 of SEQ ID NO: 363; CDR3 of SEQ ID NO: 364; a light chain variable region CDR1 of SEQ ID NO: 365; CDR2 of SEQ ID NO: 366; and CDR3 of SEQ ID NO: 367.

114. A fragment of an antibody of anyone of the previous claim that binds to HER3 selected from the group consisting of; Fab, F(ab₂)', F(ab)₂', scFv, VHH, VH, VL, dAbs.

115. A pharmaceutical composition comprising an antibody or fragment thereof selected from any one of the previous claims and a pharmaceutically acceptable carrier.

116. The pharmaceutical composition of claim 115, further comprising an additional therapeutic agent.

117. The pharmaceutical composition of claim 116, wherein the additional therapeutic agent is selected from the group consisting of an HER1 inhibitor, a HER2 inhibitor, a HER3 inhibitor, a HER4 inhibitor, an mTOR inhibitor and a PI3 Kinase inhibitor.

118. The pharmaceutical composition of claim 117, wherein the additional therapeutic agent is a HER1 inhibitor selected from the group consisting of Matuzumab (EMD72000), Erbitux®/Cetuximab, Vectibix® /Panitumumab, mAb 806, Nimotuzumab, Iressa® /Gefitinib, CI-1033 (PD183805), Lapatinib (GW-572016), Tykerb® /Lapatinib Ditosylate, Tarceva® / Erlotinib HCL (OSI-774), PKI-166, and Tovok®; a HER2 inhibitor selected from the group consisting of Pertuzumab, Trastuzumab, MM-111, neratinib, lapatinib or lapatinib ditosylate /Tykerb®; a HER3 inhibitor selected from the group consisting of, MM-121, MM-111,

IB4C3, 2DID12 (U3 Pharma AG), AMG888 (Amgen), AV-203(Aveo), MEHD7945A (Genentech) and small molecules that inhibit HER3; and a HER4 inhibitor.

119. The pharmaceutical composition of claim 117, wherein the additional therapeutic agent
5 is an mTOR inhibitor selected from the group consisting of Temsirolimus/Torisel®,
ridaforolimus / Deforolimus, AP23573, MK8669, everolimus /Affinitor® .

120. The pharmaceutical composition of claim 117, wherein the additional therapeutic agent
10 is a PI3 Kinase inhibitor selected from the group consisting of GDC 0941, BEZ235, BMK120
and BYL719.

121. A method of treating a cancer comprising selecting a subject having an HER3
expressing cancer, administering to the subject an effective amount of a composition
comprising an antibody or fragment thereof selected from any one of the previous claims.

15 122. The method of claim 121, wherein the subject is a human.

123. A method of treating a cancer mediated by a HER signalling pathway comprising
selecting a subject having an HER3 expressing cancer, administering to the subject an
effective amount of a composition comprising an antibody or fragment thereof selected from
any one of the previous claims, wherein the cancer is selected from the group consisting of
20 breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer,
gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid
leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors ,
schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear
cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and
25 melanoma.

124. The method of claim 123, wherein said cancer is breast cancer.

125. A method of treating a cancer comprising selecting a subject having an HER3
expressing cancer, administering to said subject an effective amount of a composition
comprising an a combination of antibodies or fragments thereof disclosed in Table 1 that
30 binds to HER3.

126. A method of treating a cancer comprising selecting a subject having an HER3
expressing cancer, administering to said subject an effective amount of a composition

comprising an antibody or fragment thereof that binds to HER3 and inhibits HER3 ligand-dependent signal transduction and ligand-independent signal transduction.

127. An antibody or fragment thereof of any one of the previous claims for use as a medicament.

5 128. An antibody or fragment thereof of any one of the previous claims for use in the treatment of a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway selected from the group consisting of breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia,
10 osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors, schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and melanoma.

129. An antibody having VH of SEQ ID NO: 15 and VL of SEQ ID NO: 14 for use in treating
15 a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

130. An antibody having VH of SEQ ID NO: 33 and VL of SEQ ID NO: 32 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

131. An antibody having VH of SEQ ID NO: 51 and VL of SEQ ID NO: 50 for use in treating
20 a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

132. An antibody having VH of SEQ ID NO: 69 and VL of SEQ ID NO: 68 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

25 133. An antibody having VH of SEQ ID NO: 87 and VL of SEQ ID NO: 86 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

134. An antibody having VH of SEQ ID NO: 105 and VL of SEQ ID NO: 104 for use in
30 treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

135. An antibody having VH of SEQ ID NO: 123 and VL of SEQ ID NO: 122 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
136. An antibody having VH of SEQ ID NO: 141 and VL of SEQ ID NO: 140 for use in
5 treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
137. An antibody having VH of SEQ ID NO: 151 and VL of SEQ ID NO: 158 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
- 10 138. An antibody having VH of SEQ ID NO: 177 and VL of SEQ ID NO: 176 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
139. An antibody having VH of SEQ ID NO: 195 and VL of SEQ ID NO: 194 for use in
15 treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
140. An antibody having VH of SEQ ID NO: 213 and VL of SEQ ID NO: 212 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
141. An antibody having VH of SEQ ID NO: 231 and VL of SEQ ID NO: 230 for use in
20 treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
142. An antibody having VH of SEQ ID NO: 249 and VL of SEQ ID NO: 248 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
- 25 143. An antibody having VH of SEQ ID NO: 267 and VL of SEQ ID NO: 266 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.
144. An antibody having VH of SEQ ID NO: 285 and VL of SEQ ID NO: 284 for use in
30 treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

145. An antibody having VH of SEQ ID NO: 303 and VL of SEQ ID NO: 302 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

5 146. An antibody having VH of SEQ ID NO: 321 and VL of SEQ ID NO: 320 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

147. An antibody having VH of SEQ ID NO: 339 and VL of SEQ ID NO: 338 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

10

148. An antibody having VH of SEQ ID NO: 357 and VL of SEQ ID NO: 356 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

149. An antibody having VH of SEQ ID NO: 375 and VL of SEQ ID NO: 374 for use in treating a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway.

15

150. An antibody having VH of SEQ ID NO: 15 and VL of SEQ ID NO: 14 for use as a medicament.

151. An antibody having VH of SEQ ID NO: 33 and VL of SEQ ID NO: 32 for use as a medicament.

20

152. An antibody having VH of SEQ ID NO: 51 and VL of SEQ ID NO: 50 for use as a medicament.

153. An antibody having VH of SEQ ID NO: 69 and VL of SEQ ID NO: 68 for use as a medicament.

25 154. An antibody having VH of SEQ ID NO: 87 and VL of SEQ ID NO: 86 for use as a medicament.

155. An antibody having VH of SEQ ID NO: 105 and VL of SEQ ID NO: 104 for use as a medicament.

30 156. An antibody having VH of SEQ ID NO: 123 and VL of SEQ ID NO: 122 for use as a medicament.

157. An antibody having VH of SEQ ID NO: 141 and VL of SEQ ID NO: 140 for use as a medicament.
158. An antibody having VH of SEQ ID NO: 159 and VL of SEQ ID NO: 158 for use as a medicament.
- 5 159. An antibody having VH of SEQ ID NO: 177 and VL of SEQ ID NO: 176 for use as a medicament.
160. An antibody having VH of SEQ ID NO: 195 and VL of SEQ ID NO: 194 for use as a medicament.
161. An antibody having VH of SEQ ID NO: 213 and VL of SEQ ID NO: 212 for use as a
10 medicament.
162. An antibody having VH of SEQ ID NO: 231 and VL of SEQ ID NO: 230 for use as a medicament.
163. An antibody having VH of SEQ ID NO: 249 and VL of SEQ ID NO: 248 for use as a medicament.
- 15 164. An antibody having VH of SEQ ID NO: 267 and VL of SEQ ID NO: 266 for use as a medicament.
165. An antibody having VH of SEQ ID NO: 285 and VL of SEQ ID NO: 284 for use as a medicament.
166. An antibody having VH of SEQ ID NO: 303 and VL of SEQ ID NO: 302 for use as a
20 medicament.
167. An antibody having VH of SEQ ID NO: 321 and VL of SEQ ID NO: 320 for use as a medicament.
168. An antibody having VH of SEQ ID NO: 339 and VL of SEQ ID NO: 338 for use as a medicament.
- 25 169. An antibody having VH of SEQ ID NO: 357 and VL of SEQ ID NO: 356 for use as a medicament.
170. An antibody having VH of SEQ ID NO: 375 and VL of SEQ ID NO: 374 for use as a medicament.

171. Use of an antibody or fragment thereof of any one of the previous claims for the manufacture of a medicament for the treatment of a cancer mediated by a HER3 ligand-dependent signal transduction or ligand-independent signal transduction pathway selected from the group consisting of breast cancer, colorectal cancer, lung cancer, multiple myeloma, ovarian cancer, liver cancer, gastric cancer, pancreatic cancer, prostate cancer, acute myeloid leukemia, chronic myeloid leukemia, osteosarcoma, squamous cell carcinoma, peripheral nerve sheath tumors, schwannoma, head and neck cancer, bladder cancer, esophageal cancer, glioblastoma, clear cell sarcoma of soft tissue, malignant mesothelioma, neurofibromatosis, renal cancer, and melanoma.

10

15




Fig. 1




Fig. 2




Fig. 3

ΔD [HER3: Fab] vs HER3




Fig. 4B




Fig. 4C





Fig. 5A






Fig. 5B






Fig. 5E

Fig. 6A

Inhibition of HER3-phosphorylation in stimulated MCF7 cells




Fig. 6B

Inhibition of HER3-phosphorylation in SKBR3 cells




Fig. 7A




Fig. 7B




Fig. 8A





Fig. 8B






Fig. 9




Fig. 10

Fig. 11A

MOR0923 treated BT474 xenografts




Fig. 11B

MOR09823 treated BxPC3 xenografts




Fig. 12A





Fig. 12B






Fig. 13





Fig. 14

G



H





Fig. 14 (cont.)

Fig. 15A

BT474 Tumor Growth Curve



L3.3 Tumor Growth Curve




Fig. 15B

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/064407

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/32 A61K39/395 A61P35/00
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
A61K C07K
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/077028 A2 (U3 PHARMA AG [DE]; AMGEN INC [US]; ROTHE MIKE [DE]; TREDER MARTIN [DE]) 12 July 2007 (2007-07-12) the whole document, especially page 14, line 22 - page 15, line 14; examples 8, 11, 13-25; table 4	1-171
X	WO 2006/087637 A2 (HELLENIC PASTEUR INST [GR]; MAMALAKI AVGI [GR]; BELIMEZI MARIA [GR]; P) 24 August 2006 (2006-08-24) the whole document, especially page 3, lines 16-28; page 50, lines 11-21; page 51, line 24 - page 52, line 5; page 54, lines 6-20	1,2,6,115
	----- -/--	

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>
--	--

Date of the actual completion of the international search 27 October 2011	Date of mailing of the international search report 09/11/2011
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Bayer, Annette

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/064407

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2008/019290 A2 (ASTRAZENECA AB [SE]; CARTILAGE SUE [GB]; DONG JIANYING [US]) 14 February 2008 (2008-02-14) the whole document, especially examples 4, 6, 9-13; paragraphs [0227]-[0230] -----	1,2,6, 115-118
A	FRANKLIN M C ET AL: "Insights into ErbB signaling from the structure of the ErbB2-pertuzumab complex", CANCER CELL, CELL PRESS, US, vol. 5, no. 4, 1 April 2004 (2004-04-01), pages 317-328, XP002372929, ISSN: 1535-6108, DOI: 10.1016/S1535-6108(04)00083-2 the whole document -----	1-171
A	CHO HYUN-SOO ET AL: "Structure of the extracellular region of HER3 reveals an interdomain tether", SCIENCE (WASHINGTON D C), vol. 297, no. 5585, 23 August 2002 (2002-08-23), pages 1330-1333, XP002662196, ISSN: 0036-8075 the whole document -----	1-171
A	BURGESS A W ET AL: "An open-and-shut case? Recent insights into the activation of EGF/ErbB receptors", MOLECULAR CELL, CELL PRESS, CAMBRIDGE, MA, US, vol. 12, no. 3, 1 September 2003 (2003-09-01), pages 541-552, XP002396190, ISSN: 1097-2765, DOI: 10.1016/S1097-2765(03)00350-2 the whole document, especially page 549, last full paragraph - page 550, second full paragraph -----	1-171
A	KANI KIAN ET AL: "The extracellular domains of ErbB3 retain high ligand binding affinity at endosome pH and in the locked conformation", BIOCHEMISTRY, vol. 44, no. 48, December 2005 (2005-12), pages 15842-15857, XP002662197, ISSN: 0006-2960 the whole document -----	1-171
	-/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/064407

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MCKAY BROWN ET AL: "Tolerance to single, but not multiple, amino acid replacements in antibody V-H CDR2: A means of minimizing B cell wastage from somatic hypermutation?", JOURNAL OF IMMUNOLOGY, AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US, vol. 156, no. 9, 1 January 1996 (1996-01-01), pages 3285-3291, XP002649029, ISSN: 0022-1767 the whole document -----</p>	66-90,92

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/064407

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2007077028	A2	12-07-2007	AR 056857 A1	24-10-2007
			AU 2006332065 A1	12-07-2007
			CA 2633222 A1	12-07-2007
			CN 101365723 A	11-02-2009
			CN 102174105 A	07-09-2011
			EA 200801621 A1	30-12-2008
			EP 1984402 A2	29-10-2008
			JP 2009521913 A	11-06-2009
			KR 20080099235 A	12-11-2008
			TW 200804420 A	16-01-2008
			US 2008124345 A1	29-05-2008
			US 2010183631 A1	22-07-2010
			WO 2007077028 A2	12-07-2007
			ZA 200805474 A	27-05-2009
WO 2006087637	A2	24-08-2006	AU 2006215307 A1	24-08-2006
			CA 2598535 A1	24-08-2006
			EP 1891115 A2	27-02-2008
			JP 2008529549 A	07-08-2008
			KR 20070115996 A	06-12-2007
			US 2010047230 A1	25-02-2010
			WO 2006087637 A2	24-08-2006
WO 2008019290	A2	14-02-2008	AU 2007281774 A1	14-02-2008
			CA 2659631 A1	14-02-2008
			CN 101522717 A	02-09-2009
			EP 2054444 A2	06-05-2009
			JP 2009545325 A	24-12-2009
			KR 20090058512 A	09-06-2009
			RU 2009107494 A	10-09-2010
			US 2010158926 A1	24-06-2010
			WO 2008019290 A2	14-02-2008
			ZA 200900753 A	29-09-2010