PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4:		(11) International Publication Number: WO 88/05650
A61K 6/00	A1	(43) International Publication Date: 11 August 1988 (11.08.88)
(21) International Application Number: PCT/HU (22) International Filing Date: 29 January 1988 (•	H-1368 Budapest (HU).
(31) Priority Application Number: (32) Priority Date: 3 February 1987 ((33) Priority Country:	360/ (03.02.8	pean patent), DK, FR (European patent), GB (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European
(71) Applicant (for all designated States except US) AL FINOMVEGYSZERGYÁR [HU/HU]; T 53, H-1441 Budapest (HU).	: REA Celepes	Fublished With international search report.
(72) Inventors; and (75) Inventors/Applicants (for US only): CSÁNY [HU/HU]; Gorkij fasor 40, H-1068 Budape CSÁNYI, Gábor [HU/HU]; Reviczky u. 5. Budapest (HU). BALOGH, Tibor [HU/HU] Károly krt. 12/c, H-1138 Budapest (HU). NA zló [HU/HU]; Pasaréti út 67, H-1026 Budape	est (HU , H-10 ; Rób GY, La). 8 t

(54) Title: NEW DENTAL COMPOSITION AND A PROCESS FOR THE PREPARATION THEREOF

(57) Abstract

Dental composition comprising an anti-osteoporotic agent, an antiphlogistic agent, optionally a natural or synthetic chemotherapeutic agent or local anaesthetic, furthermore one or more known auxiliary agent(s). The invention also relates to a process for the preparation of the above composition.

FOR THE PURPOSES OF INFORMATION ONLY

 $Codes \ used \ to \ identify \ States \ party \ to \ the \ PCT \ on \ the \ front \ pages \ of \ pamphlets \ publishing \ international \ applications \ under \ the \ PCT.$

ΑT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	ΙT	Italy	NO	Norway
BJ	Вепіп	JP ·	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden.
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark	MC	Monaco	US	United States of Americ
FI	Finland	MG	Madagascar	CS	Ourse praces of Americ

NEW DENTAL COMPOSITION AND A PROCESS FOR THE PREPARATION THEREOF

The invention relates to a new dental composition and to a process for the preparation thereof.

As known, caries is the most widespread endemic disease. It is quite obvious to one skilled in the art that a great deal of teeth could be saved by a preservative treatment performed in due time. In practice, however, patients turn to the dentist only when they suffer from an untolerable pain, i.e. when at least one complication exists, too.

10

15

20

25

30

Pain is caused first by pulp hyperaemia, i.e. by the reversible inflammation of the vessels and intravascular nerves of the dental radix. The volume of the narrow pulp canal, closed between hard walls, cannot increase, thus the arteries exert a hard pressure on the veins, which results in an early appearance of venal hyperaemia. At the stage of pulp hyperaemia, however, caries does not reach the pulp, since it is separated from it by a thin healthy layer of dentin.

When caries proceeds, an infectious inflammation appears in the pulp (pulpitis). Of the various forms of pulpitis only the main types are mentioned now. When an acute inflammation exists and the pulp is only partially damaged, dentists endeavour to save the living tooth. When inflammation gets, however, cronic and the pulp is completely damaged, the goal of dental treatment is to render the further treatment painless, to exclude additional complications and to ensure a symptom-free recovery.

In its final stage, pulpitis leads to gangrene. Tooth necrosis does not terminate, however, in the pulp. The defensive mechanism of the organism can act against infections only at territories with good blood supply. Periodontium, which surrounds the tooth, suits this requirement, consequently the next stage is the appearance of periodontitis. At the appearance of this disorder, causing pains even upon touch, pulp itself is no longer sensitive, the subjective symptoms are, however, very painful. The task of dentists is to ensure the quick alleviation of this complaint in the acute stage of inflammation. In chronic stages, however, additional complications must be taken into account. Inflammation may spread over the periodontal bony substance

10

15

20

25

30

where it may lead to pathologic osteoporoses (granuloma, cysts).

Several attempts have been made to defend the exacerbation of the above diseases. According to Adler and Záray

/Konzerváló fogászat (Preservative Dentistry) p. 172 (Kedicina, Budapest, 1972) (in Hungarian) pulp canal must be sterilized completely. It is very important that the surrounding tissues be free of inflammation. Therefore pulp canal has been treated with various sterilizing agents, antibiotics and antiphlogistics, utilized most frequently as a liquid composition. A disadvantage of this method is that prolonged activity cannot be ensured.

Utilization of powders or removable pastes in such treatments has also been attempted /Fogorvosi Szemle (Dentists' Review) 64, 404 (1971) (in Hungarian)7, but prolonged effect could not be achieved.

Hungarian patent No. 167,308 suggests the use of pointers comprising a sterilizing (i.e. oxidizing) or antiphlogistic agent and/or an antibiotic in admixture with an inert plastic carrier. Penicillin, chloramphenicol and oxytetracyclin are mentioned as antibiotics, whereas hydrocortisone is mentioned as antiphlogistic agent. The major disadvantage of this method is that it can be applied only for the treatment of excavated pulp canal, furthermore it cannot act against osteoporosis coupled with other complications.

The invention aims at providing a new composition applicable for the treatment of caries and for the healing of acute, subacute or chronic inflammatory lesions of pulp, dental radix or the bony substance surrounding them.

10

15

20

25

30

Our extensive experimental work has led to the unexpected recognition that a particular combination, i.e. a combination of ipriflavone (7-isopropoxy-isoflavone) and an antiphlogistic agent can be utilized with good results for the complex treatment of teeth. Depending on the conditions of use, this combination can be supplemented with chemotherapeutic agents, local anaesthetics and known auxiliary agents.

The above combination can be applied successfully in all stages of pulpitis and periodontitis.

Based on the above, the invention relates to a new dental composition comprising an anti-osteoporotic agent, preferably ipriflavone, an antiphlogistic agent, optionally a natural or synthetic chemotherapeutic agent or a local anaesthetic, furthermore one or more known auxiliary agent(s).

The invention also relates to a process for the preparation of the above dental composition. According to the invention 0.1 to 30 % by weight of ipriflavone, 0.5 to 15 % by weight of an antiphlogistic agent, preferably triamcinolone, and optionally one or more natural or synthetic chemotherapeutic agent(s) or local anaesthetic(s) are admixed with auxiliary agents commonly applied in dentistry. As synthetic chemotherapeutic agents e.g. sulphonamides or trimethoprim \(\frac{7}{2}, 4\)-diamino-5-(3',4',5'-trimethoxybenzyl)-pyrimidine? whereas as natural chemotherapeutic agent e.g. doxycyclin base (an antibiotic) or a salt thereof can be applied. As local anaesthetic \(\frac{7}{2}\)required e.g. to prepare a coating for protecting dental stump? lidocaine can be used. Of the sulphonamides sulphamethoxazole is particularly preferred. The known auxiliary agents for dental purposes are

10

15

e.g. zinc oxide, calcium oxide, bismuth oxide, titanium dioxide, eugenol, epoxy resin and dental lacquers.

The utilization of more than one active agent in a dental filler has led to particularly advantageous and unexpected results.

The antimicrobial agent with a broad spectrum of activity is utilized to exterminate the pathogens which cause inflammation. The antimicrobial agent alone does not act on the other components of the inflammatory process.

Triamcinolone, utilized as antiphlogistic agent, exerts a very strong effect on the inflammatory process: it suppresses the cedema and stops the other inflammatory components, resulting in a quick alleviation of pain and in the cessation of inflammation. It is known, however, that this substance may also enhance the activity of osteoklasts causing osteoporosis in the bony substance e.g. when a triamcinolone-containing filling agent is introduced into the pulp canal after the removal of pulp.

Ipriflavone, one of the members of the combination according to the invention, supplements the effects of the two former components in a particularly advantageous manner: it is able to suppress osteoporosis even alone, furthermore it antagonizes the osteoklast activity-increasing effects of steroidal antiphlogistics.

By combining these three types of active agents a series of therapeutic dental fillers, suitable for preserving the teeth in any stage of pulpitis and periodontitis, can be obtained.

The most important advantage provided for by the invention is that the combination can be utilized in several

25

30

ŝ

20

10

15

types of dental diseases. The new combinations can be utilized for the following purposes:

1/ In the treatment of pulp hyperaemia and acute partial pulpitis a dental cement comprising 0.1 to 10 %, preferably 0.7 to 2 % of the combination according to the invention should be placed onto the pulp-side wall of the cavity formed by the dentist, wherefrom the active agents enter the pulp and result in a very quick relief of pain and in the suppression of inflammation. Doxycyclin, mentioned as an example of the chemotherapeutic agents, accumulates in the tooth, resulting in a prolonged effect. Upon the effect of ipriflavone the formation of secondary dentin is also to be expected.

2/ In chronic total pulpitis the treatment is performed in two steps. First the treatment described in point 1/ above is performed in order to attain a painless stage, thereafter the pulp canal is excavated to prevent the spreading of inflammation, and in the second treatment step the method described in point 3/ below is performed.

3/ In the treatment of periodontitis the therapeutic paste (admixed with appropriate auxiliary agents) is applied into the previously cleaned and dried pulp canal. The antiphlogistic and antimicrobial effects appear quickly, and a very quick relief of pain is to be expected.

4/ When the complications of periodontitis are to be avoided or the already existing complications should be suppressed, the reversal of osteoporoses already formed (ostitis diffusa, granuloma, cysts) is to be expected upon a prolonged application of the anti-osteoporotic therapeutic paste. The final root filling is prepared after the termina-

25

30

20

10

15

20

25

30

Ţ,

tion of the complete bony defects, utilizing an antibiotic-free composition comprising anti-osteoporotic agents, X-ray contrast materials and carriers.

The invention is elucidated in detail by the aid of the following non-limiting Examples.

Example 1

Preparation of a therapeutic cement

A mixture of 2 g of doxycyclin hyclate, 0.8 g of triamcinolone, 2.0 g of ipriflavone, 80 g of zinc oxide and 15.2 g of calcium oxide (Component "A") is admixed with eugenol (Component "B") in a ratio to obtain a composition with a consistency required in dentistry.

Example 2

Preparation of a therapeutic cement

One proceeds as described in Example 1 with the difference that doxycyclin hyclate is replaced by 2.2 g of trimethoprim.

Example 3

Preparation of a therapeutic paste for sterilizing pulp canal

0.2 g of ipriflavone is admixed with 0.2 g of doxy-cyclin base, 0.08 g of triamcinolone and 10 g of zinc oxide, and the mixture is completed with eugenol in an amount required to obtain a paste.

Example 4

Preparation of a therapeutic paste for sterilizing pulp canal

One proceeds as described in Example 3 with the difference that sulphamethoxazole is substituted for doxy-cyclin base.

Example 5

Preparation of a root filler

0.5 g of ipriflavone are admixed with 0.1 g of triamcinolone, 4.0 g of bismuth oxide, 2.0 g of titanium dioxide and 3.4 g of zinc oxide, and epoxy resin is added to the mix- * ture in an amount to obtain a paste.

Example 6

Preparation of a coating composition for protecting dental stump

A mixture of 0.3 g of ipriflavone, 0.1 g of triamcinolone and 0.2 g of lidocaine base is dissolved in 5 ml of acetone. This solution can be utilized for the preparation of a single coating or multiple coatings.

10

15.

20

25

Ē

What we claim is:

- l. A dental composition which comprises an anti-osteoporotic agent, an antiphlogistic agent, optionally a natural or synthetic chemotherapeutic agent or local anaesthetic, furthermore one or more known auxiliary agent(s).
- 2. A composition as claimed in claim 1, which comprises ipriflavone (7-isopropoxy-isoflavone) as anti-osteoporotic agent, triamcinolone as antiphlogistic agent, an antibiotic, preferably doxycyclin or a salt thereof, a sulphonamide or trimethoprim as a chemotherapeutic agent and lidocaine as local anaesthetic.
- 3. A composition as claimed in claim 1 or 2, which comprises 0.1 to 30 % by weight of ipriflavone, 0.5 to 15 % by weight of triamcinolone, optionally 0.1 to 10 % by weight of doxycyclin or a salt thereof, optionally 1 to 20 % by weight of lidocaine, and 35 to 99.4 % by weight of one or more known auxiliary agent(s).
- 4. A process for the preparation of a dental composition, characterized in that 0.1 to 30 % by weight of ipriflavone, 0.5 to 15 % by weight of triamcinolone, optionally 0.1 to 10 % by weight of doxycyclin or a salt thereof and optionally 1 to 20 % by weight of lidocaine are admixed with 35 to 99.4 % by weight of one or more known auxiliary agent(s).
- 5. A method for protecting pulp of carious teeth, characterized in that a composition as claimed in claim 1 is utilized.
- 6. A method for preserving the vitality of pulp in early stages of pulpitis, characterized in that a composition as claimed in claim 1 is utilized.

- 7. A method for sterilizing pulp canal, characterized 'in that a composition as claimed in claim l is utilized.
- 8. A method for preparing a final root filling with the provision of an X-ray contrast agent, characterized in that a composition as claimed in claim 1 is utilized.

INTERNATIONAL SEARCH REPORT

International Application No PCT/HU 88/00003

. CLASSIS	CATION OF SUBJECT MATTER (if several classification)	ation symbols apply, indicate all) *	
According to	nternational Patent Classification (IPC) or to both Nation	al Classification and IPC	
TPC ⁴ :	A 61 K 6/00		
II. FIELDS S			
	Minimum Documental		
Classification S	ystem Cla	assification Symbols	
	4		
Int.Cl.	A 61 K 6/00,6/02		
	Documentation Searched other that to the Extent that such Documents ar	n Minimum Documentation re included in the Fields Searched ⁹	
III. DOCUME	NTS CONSIDERED TO BE RELEVANT	ories of the relevant nessages 12	Relevant to Claim No. 13
ategory •	Citation of Document, 11 with Indication, where appro-	•	
	E, A1, 3 539 202 (LION COF 22.05.86).	RP.) 22 May 1986	(1)
A I	E, A1, 2 629 694 (TAKAHASH O January 1977 (20.01.77).	HI et al.)	(1)
A 1	E, A, 2 321 215 (MOCHIDA S 5 November 1973 (15.11.73)	SEIYAKU)	(1)
A I	E, B, 1 202 937 (L.D.CAUL) 4 October 1965 (14.10.65).	(COMPANY)	(1)
A I	E, B, 1 195 436 (HAAS et a 24.06.65).	al.) 24 June 1965	(1)
A. I	DE, C, 667 808 (LINGNER-WER 938 (21.11.38).	RKE) 21 November	(1)
A C	B, A, 1 465 406 (SVEN-GUNN) 3 February 1977 (23.02.77)	NAR HESSELGREN)).	(1)
A U	JS, A, 4 121 940 (MICHEL et .978 (24.10.78).	t al.) 24 October	(1)
"A" docum consider earlier filing docum which citatio "O" docum other "P" docum later t	sent which may throw doubts on priority claim(s) or is cited to establish the publication date of another in or other special reason (as specified) sent referring to an oral disclosure, use, exhibition or means sent published prior to the international filing date but han the priority date claimed	"T" later document published after or priority date and not in concided to understand the principle invention. "X" document of particular relevations of the considered novel (involve an inventive step document of particular relevation of the considered to involve document is combined with or ments, such combination being in the art. "A" document member of the same	ple or theory underlying the ince; the claimed invention or cannet be considered to ince; the claimed invention e an inventive step when the or more other such docup obvious to a person skilled a patent family
14 A	April 1988 (14.04.88)	25 April 1988	(25.04.88)
international	Searching Authority	Signature of Authorized Officer	
ATTST	RIAN PATENT OFFICE	Milling	

III. DOCU	MENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET		
ategory *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No	
·			
A	US, A, 3 913 229 (DRISKELL et al.) 21 October 1975 (21.10.75).	(1)	
	□ □ □		
1		-	
ļ			
-		• .	
-			
		· ·	
-			
		-	
	·		

Anhang zum internationalen Recherchenbericht über die internationale Patentanmeldung Nr.

In diesem Anhang sind die Mitglieder der Patentfamilien der im obengenannten internationalen Recherchenbericht angeführten Patentdokumente angegeben. Diese Angaben dienen nur zur Unterrichtung und erfolgen ohne Gewähr.

Annex to the International Search Report on International Patent Application No.PCT/HU 88/0003

This Annex lists the patent family members relating to the patent documents cited in the above-mentioned International search report. The Austrian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Annexe au rapport de recherche internationale relatif à la demande de brevet international n°.

La présente annexe indique les membres de la famille de brevets relatifs aux documents de brevets cités dans le rapport de recherche internationale visé ci-dessus. Les renseignements fournis sont donnés à titre indicatif et n'engagent pas la responsabilité de l'Office autrichien des brevets.

Im Recherchenbericht angeführtes Patent- dokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veroffentlichung Publication date Date de publication	
DE-A1-3 539 202	22/05/1986	GB-A1- 2 167 299 JP-A2-61-112 028 US-A - 4 725 428	29/05/1986 30/05/1986 16/02/1988	
DE-A1-2 629 694 -B2- -C3-	20/01/1977 27/09/1979 04/06/1980	DK-A - 2 911/76 DK-B - 145 448 DK-C - 145 448 JP-A2-52-004 687 JP-B4-60-050 763 US-A - 4 080 212	31/12/1976 22/11/1982 18/04/1983 13/01/1977 11/11/1985 21/03/1978	
DE-A -2 321 215	15/11/1973	CA-A1- 960 107 CH-A - 548 749 DE-A1- 2 259 490 BR-A0- 7 303 064 CA-A1- 1 032 692 FR-A1- 2 182 186 FR-B1- 2 182 186 GB-A - 1 426 901 JP-A2-49-005 143 JP-B4-51-037 092 US-A - 3 829 973	31/12/1974 15/05/1974 18/10/1973 29/08/1974 06/06/1978 07/12/1973 28/05/1976 03/03/1976 17/01/1974 13/10/1976 20/08/1974	
DE-B -1 202 937	14/10/1965	None		
DE-B -1 195 436	24/06/1965	None ·		
DE-C - 667 808	21/11/1938	None		

GB3A-1 465 406	23/02/1977	NO-C - 141 SE-B - 379	726 726 970 970 492 604 495 145 145 479	08/08/1974 23/10/1978 02/04/1979 30/08/1974 28/07/1978 23/10/1974 02/05/1983 07/08/1974 15/10/1979 23/01/1980 13/10/1975 22/01/1976 22/06/1976
US-A-4 121 940	24/10/1978	CA-A1- 1 104 JP-A2-53-133		30/06/1981 21/11/1978
US-A-3 913 229	21/10/1975	BE-A1- 825 CA-A1- 1 042	801 701 933 749 749 781 489	26/08/1976 16/06/1975 21/11/1978 15/09/1978 28/08/1975 19/09/1975 04/11/1977 08/09/1977 30/09/1975 14/03/1983