



US 20140056879A1

(19) **United States**

(12) **Patent Application Publication**
LAZAR

(10) **Pub. No.: US 2014/0056879 A1**

(43) **Pub. Date: Feb. 27, 2014**

(54) **FC VARIANTS THAT EXTEND ANTIBODY
HALF-LIFE**

(60) Provisional application No. 61/392,115, filed on Oct.
12, 2010.

(71) Applicant: **XENCOR, INC.**, MONROVIA, CA
(US)

Publication Classification

(72) Inventor: **GREGORY A. LAZAR**, ARCADIA,
CA (US)

(51) **Int. Cl.**
C07K 16/22 (2006.01)

(73) Assignee: **XENCOR, INC.**, MONROVIA, CA
(US)

(52) **U.S. Cl.**
CPC **C07K 16/22** (2013.01)
USPC ... **424/133.1**; 530/387.3; 536/23.53; 435/328

(21) Appl. No.: **13/973,934**

(22) Filed: **Aug. 22, 2013**

(57) **ABSTRACT**

Related U.S. Application Data

(63) Continuation of application No. 13/272,143, filed on
Oct. 12, 2011, now Pat. No. 8,546,543.

The invention relates generally to compositions and methods
for altering the serum half-life in vivo of an antibody.

Figure 1

CH1																									
EU Index	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138				
IgG1	A	S	T	K	G	P	S	V	F	P	L	A	P	S	S	K	S	T	S	G	G				
IgG2	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S				
IgG3	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	G	G				
IgG4	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S				
EU Index	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159				
IgG1	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N				
IgG2	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N				
IgG3	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N				
IgG4	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N				
EU Index	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180				
IgG1	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y				
IgG2	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y				
IgG3	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y				
IgG4	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y				
EU Index	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201				
IgG1	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	Q	T	Y	I	C	N				
IgG2	S	L	S	S	V	V	T	V	P	S	S	N	F	G	T	Q	T	Y	T	C	N				
IgG3	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	Q	T	Y	T	C	N				
IgG4	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	K	T	Y	T	C	N				
EU Index	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220						
IgG1	V	N	H	K	P	S	N	T	K	V	D	K	K/R	V	E	P	K	S	C						
IgG2	V	D	H	K	P	S	N	T	K	V	D	K	T	V	E	R	K	C	C						
IgG3	V	N	H	K	P	S	N	T	K	V	D	K	R	V	E	L	K	T	P						
IgG4	V	D	H	K	P	S	N	T	K	V	D	K	R	V	E	S	K	Y	G						
Hinge											Fc														
EU Index	221				222	223	224	225	226	227	228														
IgG1	D				K	T	H	T	C	P	P														
IgG2					V		E		C	P	P														
IgG3	L	G	D	T	T	H	T	C	P	R	C	P	E	P	K	S	C	D	T	P	P				
IgG4							P	P	C	P	S														
EU Index																									
IgG1																									
IgG2																									
IgG3																									
IgG4																									
EU Index													229	230	231	232	233	234	235	236					
IgG1													C	P	A	P	E	L	L	G					
IgG2													C	P	A	P	P	V	A						
IgG3													C	P	A	P	E	L	L	G					
IgG4													C	P	A	P	E	F	L	G					

Figure 1 (continued)

CH2																												
EU Index	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257							
IgG1	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P							
IgG2	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P							
IgG3	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P							
IgG4	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P							
EU Index	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278							
IgG1	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y							
IgG2	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	Q	F	N	W	Y							
IgG3	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	Q	F	K	W	Y							
IgG4	E	V	T	C	V	V	V	D	V	S	Q	E	D	P	E	V	Q	F	N	W	Y							
EU Index	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299							
IgG1	V	D	G	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	Y	N	S	T							
IgG2	V	D	G	VM	E	V	H	N	A	K	T	K	P	R	E	E	Q	F	N	S	T							
IgG3	V	D	G	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	Y	N	S	T							
IgG4	V	D	G	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	F	N	S	T							
EU Index	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320							
IgG1	Y	R	V	V	S	V	L	T	V	L	H	Q	D	W	L	N	G	K	E	Y	K							
IgG2	F	R	V	V	S	V	L	T	V	V	H	Q	D	W	L	N	G	K	E	Y	K							
IgG3	F	R	V	V	S	V	L	T	V	L	H	Q	D	W	L	N	G	K	E	Y	K							
IgG4	Y	R	V	V	S	V	L	T	V	L	H	Q	D	W	L	N	G	K	E	Y	K							
EU Index	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340								
IgG1	C	K	V	S	N	K	A	L	P	A	P	I	E	K	T	I	S	K	A	K								
IgG2	C	K	V	S	N	K	G	L	P	A	P	I	E	K	T	I	S	K	T	K								
IgG3	C	K	V	S	N	K	A	L	P	A	P	I	E	K	T	I	S	K	T	K								
IgG4	C	K	V	S	N	K	G	L	P	S	S	I	E	K	T	I	S	K	A	K								
CH3																												
EU Index	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361							
IgG1	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	DE	E	LM	T	K	N							
IgG2	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	E	E	M	T	K	N							
IgG3	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	E	E	M	T	K	N							
IgG4	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	Q	E	E	M	T	K	N							
EU Index	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382							
IgG1	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E							
IgG2	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E							
IgG3	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E							
IgG4	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E							
EU Index	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403							
IgG1	S	N	G	Q	P	E	N	N	Y	K	T	T	P	P	V	L	D	S	D	G	S							
IgG2	S	N	G	Q	P	E	N	N	Y	K	T	T	P	P	M	L	D	S	D	G	S							
IgG3	S	S	G	Q	P	E	N	N	Y	N	T	T	P	P	M	L	D	S	D	G	S							
IgG4	S	N	G	Q	P	E	N	N	Y	K	T	T	P	P	V	L	D	S	D	G	S							
EU Index	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424							
IgG1	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	G	N	V	F	S							
IgG2	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	G	N	V	F	S							
IgG3	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	G	N	I	F	S							
IgG4	F	F	L	Y	S	R	L	T	V	D	K	S	R	W	Q	E	G	N	V	F	S							
EU Index	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445							
IgG1	C	S	V	M	H	E	AG	L	H	N	H	Y	T	Q	K	S	L	S	L	S	P							
IgG2	C	S	V	M	H	E	A	L	H	N	H	Y	T	Q	K	S	L	S	L	S	P							
IgG3	C	S	V	M	H	E	A	L	H	N	R	F	T	Q	K	S	L	S	L	S	P							
IgG4	C	S	V	M	H	E	A	L	H	N	H	Y	T	Q	K	S	L	S	L	S	L							
EU Index	446	447																										
IgG1	G	K																										
IgG2	G	K																										
IgG3	G	K																										
IgG4	G	K																										

Figure 2

Kappa constant light chain (C κ) (SEQ ID NO:1)

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD
SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

IgG1 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:2)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPS
VFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGN
VFSCSVMHEALHNHYTQKSLSLSPGK

IgG2 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:3)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSNFGTQTYTCNVNHHKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLF
PPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRV
SVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPGK

IgG3 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:4)

ASTKGPSVFPLAPCSRSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYTCNVNHHKPSNTKVDKRVELKTPLGDTTHTCPRCPEPKSCD
TPPPCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCPAPELLGGPSVFLFPPKPKDTL
MISRTPEVTCVVDVSHEDPEVQFKWYVDGVEVHNAKTKPREEQYNSTFRVSVLTVLHQ
DWLNGKEYKCKVSNKALPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGF
YPSDIAVEWESSGQPENNYNTTPPMLDSDGSFFLYSKLTVDKSRWQQGNIFSCSVMHEAL
HNRFTQKSLSLSPGK

IgG4 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:5)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYTCNVNHHKPSNTKVDKRVESKYGPPCPSCPAPEFLGGPSVFL
FPPKPKDTLMISRTPEVTCVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRV
VSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF
SCSVMHEALHNHYTQKSLSLGLGK

IgG1/2 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:6)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSV
FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFR
VSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQ
VSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNV
FSCSVMHEALHNHYTQKSLSLSPGK

Figure 3

IgG2 434S (SEQ ID NO:7)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSNFGTQTYTCNVDHKPSNTKVDKTVKCCVECPAPPVAGPSVFLF
PPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVV
SVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTPMLDSDGSFFLYSKLTVDKSRWQQGNVFS
CSVMHEALHSHYTQKSLSLSPGK

IgG1/2 434S (SEQ ID NO:8)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSV
FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFR
VVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQ
VSLTCLVKGFYPSDIAVEWESNGQPENNYKTPMLDSDGSFFLYSKLTVDKSRWQQGNV
FSCVMHEALHSHYTQKSLSLSPGK

Figure 4

Anti-VEGF VH (SEQ ID NO:9)

EVQLVESGGGLVQPGGSLRLSCAASGYFTFTNYGMNWVRQAPGKGLEWVGWINTYTGEPT
YAADFRRFTFSLDTSKSTAYLQMNSLR AEDTAVYYCAKYPHYGSSHWYFDVWGQGT
TVSS

Anti-VEGF VL (SEQ ID NO:10)

DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVPSRF
SGSGSGTDFLTISLQPEDFATYYCQQYSTVPWTFGQGTKVEIK

Figure 5




Figure 6




Figure 7




Figure 8a




Figure 8b




Figure 9

Fc variant	IgG	n	Half-Life			Cmax		AUC		Clearance	
			(day)			(ug/mL)		(day*ug/mL)		(mL/day/kg)	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
Study 1											
WT (Native IgG1)	IgG1	6	3	0.5	1.0	28.5	7.3	69.8	8.4	29.1	4.2
P257L	IgG1	6	1.9	0.4	0.6	18.1	8.4	38	11.3	59.5	28.8
P257N	IgG1	6	0.9	0.1	0.3	30.1	4.7	38.2	6.1	53.4	8.8
V308F	IgG1	6	4.9	0.3	1.6	30.1	9.4	129.2	11	15.6	1.3
T250Q/M428L	IgG1	6	8.4	2.8	2.8	22.3	3.8	186	52.5	12	5.4
M252Y/S254T/ T256E	IgG1	4	10.9	1.2	3.6	34.6	10.1	330.8	57.3	6.2	1.1
N434A	IgG1	6	5.5	2.1	1.8	24.5	6.2	143.1	26.9	14.3	2.2
Study 2											
WT (Native IgG1)	IgG1	7	3.9	0.52	1.0	36	5.6	92	13	22	2.9
V308F	IgG1	7	6.8	0.99	1.7	28	8.7	152	34.6	14	3.4
M252T/S254T/ T256E	IgG1	7	11.3	1.2	2.9	25	7	241	64.2	8.9	2.6
Study 3											
WT (Native IgG1)	IgG1	6	2.8	0.3	1.0	27	6	69	10	29.4	4.6
V308F	IgG1	6	5.9	0.4	2.1	33	3	173	33	11.8	1.9
M252T/S254T/ T256E	IgG1	6	10.4	1.5	3.7	34	6	317	67	6.6	1.5
N434S	IgG1	6	7.7	1.5	2.8	33	11	228	75	10	4.6
V259I/V308F	IgG1	6	9.2	1.5	3.3	36	5	262	47	7.9	1.4
V308F/M428L	IgG1	6	11.2	1.1	4.0	38	6	307	50	6.7	1.1
M252Y/V308F	IgG1	6	5.7	0.4	2.0	40	6	140	24	14.7	2.6
M428L/N434S	IgG1	6	12	2.9	4.3	37	9	400	112	5.5	2
V308F/N434S	IgG1	6	8.6	0.3	3.1	28	3	195	25	10.4	1.3
V259I/V308F/ M428L	IgG1	6	13.3	2.7	4.8	37	6	383	68	5.3	0.8

Figure 9 (continued)

Fc variant	IgG	n	Half-Life			Cmax		AUC		Clearance	
			(day)			(ug/mL)		(day*ug/mL)		(mL/day/kg)	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
Study 4											
WT (Native IgG1)	IgG1	6	2.9	0.4	1.0	32.5	3.1	72.7	5.8	27.6	2.3
M252Y/S254T/T256E	IgG1	6	11.3	1.8	3.9	40.6	5.8	376.5	60.7	5.4	0.8
V308F/N434M	IgG1	6	4.9	0.8	1.7	38.6	5.4	153.6	21.3	13.2	1.9
V259I/V308F	IgG1	6	7.5	0.8	2.6	37.9	3.8	234.5	23.1	8.6	0.9
V308F	IgG1	6	6.1	0.4	2.1	37.8	1.9	172.2	16.7	11.7	1.2
V308F/Y319I	IgG1	6	6	0.5	2.1	38.1	6.5	206.5	71	10.5	3.1
M428L/N434S	IgG1	6	11.8	0.6	4.1	42	4.6	392.4	52.2	5.2	0.7
V308F/M428L/N434S	IgG1	6	9	0.4	3.1	31.8	7.9	204.6	34.2	10	1.8
V259I/V308F/N434S	IgG1	6	8.5	1.8	2.9	33.4	7.3	254.9	29	7.9	0.9
V259I/V308F/M428L	IgG1	6	10.9	0.6	3.8	35	7.5	295.1	54.5	7	1.3
Study 5											
WT (Native IgG2)	IgG2	6	5.9	0.9	1.0	43.8	5	212.3	33.8	9.6	1.5
N434S	IgG2	6	12.2	1.3	2.1	39.1	6.4	381.6	91.6	5.5	1.2
V259I/V308F	IgG2	6	10.2	0.6	1.7	42.3	1.3	342	21.2	5.9	0.4
M428L/N434S	IgG2	6	16.5	1.1	2.8	48	5.8	603.3	72.9	3.4	0.4

Figure 10a

	Fold cyno FcRn			Fold human FcRn		
	n	Mean	SD	n	Mean	SD
WT IgG1		1.00			1.00	
M252Y/S254T/T256E	2	11.95	2.23	10	11.37	2.90
V259I/V308F	2	8.60	1.86	12	9.79	2.72
M428L/N434S	3	12.18	1.23	14	17.28	6.70
V259I/V308F/M428L	4	12.43	0.57	13	19.94	10.46

Figure 10b




Figure 11




Figure 12

Fc variant	IgG	n	Half-Life (day)			Cmax (ug/mL)		AUC Day*ug/mL		Clearance mL/day/kg	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
WT (Native IgG1)	IgG1	3	9.7		1.0	101.2		822.9		4.9	
M252Y/S254T/T256E	IgG1	3	24.2	1.6	2.5	111.2	13.9	1919	210.1	2.1	0.2
V259I/V308F	IgG1	3	16.2	6.4	1.7	102.8	3.2	1352.6	367	3.1	0.9
M428L/N434S	IgG1	3	31.1	7.9	3.2	126	17	2660.9	791	1.6	0.6
V259I/V308F/M428L	IgG1	3	25.1	5.9	2.6	124.6	14.3	2301.6	922.5	1.9	0.8

Figure 13

K248Y	R255W	K288N	T307K	D312K	Q438W
D249G	R255P	K288H	T307H	D312Y	Q438H
I253H	V284D	K288Q	T307L	K338R	Q438N
I253N	V284E	K288Y	T307F	K338H	Q438S
I253L	H285D	K288F	T307W	A378S	Q438G
I253T	H285E	K288W	V309W	A378V	Q438Y
I253V	N286Q	K288L	V309R	A378G	Q438F
I253Q	N286F	K288I	V309K	S426W	Q438I
I253M	N286D	T307M	V309D	S426H	Q438D
I253Y	N286E	T307E	V309E	S426L	Y436I
I253F	N286G	T307R	V309F	S426V	Y436V
I253W	N286L	T307D	V309H	S426Y	Y436W
R255H	N286P	T307G	V309I	S426M	
R255I	N286Y	T307I	V309P	S426I	
R255Q	N286W	T307N	V309Q	S426F	
R255E	N286I	T307P	V309L	S426T	
R255F	N286V	T307Q	V309Y	M428I	
R255L	N286R	T307S	V309M	M428V	
R255S	N286K	T307V	D312E	H433G	
R255G	N286H	T307Y	D312R	Y436F	

Figure 14

IgG	Variant	k_{off} (1/s)	IgG	Variant	k_{off} (1/s)
IgG1	WT	0.365	IgG1/2	N434S/H285E	0.126
IgG1/2	N434S	0.117	IgG1/2	N434S/N286Q	0.126
IgG1/2	M428L/N434S	0.0416	IgG1/2	N434S/N286F	0.12
IgG1/2	N434S/K248Y	0.176	IgG1/2	N434S/N286D	0.0991
IgG1/2	N434S/D249G	0.142	IgG1/2	N434S/N286E	ND
IgG1/2	N434S/I253H	ND	IgG1/2	N434S/N286G	0.124
IgG1/2	N434S/I253N	ND	IgG1/2	N434S/N286L	0.094
IgG1/2	N434S/I253L	ND	IgG1/2	N434S/N286P	0.122
IgG1/2	N434S/I253T	ND	IgG1/2	N434S/N286Y	0.103
IgG1/2	N434S/I253V	0.98	IgG1/2	N434S/N286W	0.1
IgG1/2	N434S/I253Q	ND	IgG1/2	N434S/N286I	0.148
IgG1/2	N434S/I253M	0.818	IgG1/2	N434S/N286V	0.143
IgG1/2	N434S/I253Y	ND	IgG1/2	N434S/N286R	0.0988
IgG1/2	N434S/I253F	ND	IgG1/2	N434S/N286K	0.272
IgG1/2	N434S/I253W	ND	IgG1/2	N434S/N286H	0.149
IgG1/2	N434S/R255H	0.189	IgG1/2	N434S/K288N	0.147
IgG1/2	N434S/R255I	0.0943	IgG1/2	N434S/K288H	0.121
IgG1/2	N434S/R255Q	0.186	IgG1/2	N434S/K288Q	0.162
IgG1/2	N434S/R255E	0.213	IgG1/2	N434S/K288Y	0.162
IgG1/2	N434S/R255F	0.136	IgG1/2	N434S/K288F	0.158
IgG1/2	N434S/R255L	0.107	IgG1/2	N434S/K288W	0.154
IgG1/2	N434S/R255S	0.169	IgG1/2	N434S/K288L	0.164
IgG1/2	N434S/R255G	0.189	IgG1/2	N434S/K288I	0.17
IgG1/2	N434S/R255W	0.107	IgG1/2	N434S/T307M	0.0653
IgG1/2	N434S/R255P	0.0965	IgG1/2	N434S/T307E	0.0702
IgG1/2	N434S/V284D	0.116	IgG1/2	N434S/T307R	0.193
IgG1/2	N434S/V284E	0.0981	IgG1/2	N434S/T307D	0.0749
IgG1/2	N434S/H285D	0.117	IgG1/2	N434S/T307G	0.0818

Figure 14 (continued)

IgG	Variant	k_{off} (1/s)	IgG	Variant	k_{off} (1/s)
IgG1/2	N434S/T307I	0.111	IgG1/2	N434S/D312Y	0.163
IgG1/2	N434S/T307N	0.125	IgG1/2	N434S/K338R	0.141
IgG1/2	N434S/T307P	0.0702	IgG1/2	N434S/K338H	0.142
IgG1/2	N434S/T307Q	0.0488	IgG1/2	N434S/A378S	0.0902
IgG1/2	N434S/T307S	ND	IgG1/2	N434S/A378V	0.0593
IgG1/2	N434S/T307V	0.092	IgG1/2	N434S/A378G	0.179
IgG1/2	N434S/T307Y	0.0847	IgG1/2	N434S/S426W	0.275
IgG1/2	N434S/T307K	0.211	IgG1/2	N434S/S426H	0.134
IgG1/2	N434S/T307H	0.107	IgG1/2	N434S/S426L	0.173
IgG1/2	N434S/T307L	0.137	IgG1/2	N434S/S426V	0.0896
IgG1/2	N434S/T307F	0.087	IgG1/2	N434S/S426Y	0.22
IgG1/2	N434S/T307W	0.0627	IgG1/2	N434S/S426M	0.149
IgG1/2	N434S/V309W	0.0832	IgG1/2	N434S/S426I	0.0867
IgG1/2	N434S/V309R	0.329	IgG1/2	N434S/S426F	0.285
IgG1/2	N434S/V309K	0.617	IgG1/2	N434S/S426T	0.0784
IgG1/2	N434S/V309D	0.359	IgG1/2	N434S/M428I	0.117
IgG1/2	N434S/V309E	0.138	IgG1/2	N434S/M428V	0.153
IgG1/2	N434S/V309F	0.168	IgG1/2	N434S/H433G	0.121
IgG1/2	N434S/V309H	0.364	IgG1/2	N434S/Y436F	0.118
IgG1/2	N434S/V309I	0.13	IgG1/2	N434S/Q438W	0.119
IgG1/2	N434S/V309P	0.16	IgG1/2	N434S/Q438H	0.164
IgG1/2	N434S/V309Q	0.227	IgG1/2	N434S/Q438N	0.113
IgG1/2	N434S/V309L	0.137	IgG1/2	N434S/Q438S	0.106
IgG1/2	N434S/V309Y	0.129	IgG1/2	N434S/Q438G	0.146
IgG1/2	N434S/V309M	0.11	IgG1/2	N434S/Q438Y	0.139
IgG1/2	N434S/D312E	0.11	IgG1/2	N434S/Q438F	0.122
IgG1/2	N434S/D312R	0.231	IgG1/2	N434S/Q438I	0.104
IgG1/2	N434S/D312K	0.221	IgG1/2	N434S/Q438D	0.104

Figure 15




Figure 16

IgG	Variant	k_{on} (1/Ms)	k_{off} (1/s)	K_A (1/M)	K_D (M)	Fold K_D IgG1 WT	Fold K_D IgG1/2 N434S	Fold k_{off} IgG1 WT	Fold k_{off} IgG1/2 N434S
IgG1	WT	7.40E+04	0.258	2.87E+05	3.49E-06	1.00	0.10	1.00	0.28
IgG1	N434S	2.12E+05	0.124	1.71E+06	5.83E-07	5.99	0.61	2.08	0.59
IgG1	M428L/N434S	2.06E+05	0.0407	5.06E+06	1.98E-07	17.63	1.78	6.34	1.81
IgG1/2	WT	3.80E+04	0.291	1.31E+05	7.66E-06	0.46	0.05	0.89	0.25
IgG1/2	N434S	2.08E+05	0.0735	2.83E+06	3.53E-07	9.89	1.00	3.51	1.00
IgG1/2	M428L/N434S	2.00E+05	0.0338	5.92E+06	1.69E-07	20.65	2.09	7.63	2.17
IgG1/2	Q311I/N434S	2.49E+05	0.0365	6.82E+06	1.47E-07	23.74	2.40	7.07	2.01
IgG1/2	Q311L/N434S	2.40E+05	0.0554	4.34E+06	2.31E-07	15.11	1.53	4.66	1.33
IgG1/2	Q311M/N434S	2.26E+05	0.0487	4.64E+06	2.15E-07	16.23	1.64	5.30	1.51
IgG1/2	Q311V/N434S	2.12E+05	0.0367	5.77E+06	1.73E-07	20.17	2.04	7.03	2.00
IgG1/2	N434S/Y436H	1.14E+05	0.214	5.32E+05	1.88E-06	1.86	0.19	1.21	0.34
IgG1/2	N434S/Y436I	2.12E+05	0.0413	5.14E+06	1.94E-07	17.99	1.82	6.25	1.78
IgG1/2	N434S/Y436K	1.54E+05	0.213	7.23E+05	1.38E-06	2.53	0.26	1.21	0.35
IgG1/2	N434S/Y436L	1.60E+05	0.097	1.65E+06	6.05E-07	5.77	0.58	2.66	0.76
IgG1/2	N434S/Y436V	1.96E+05	0.055	3.56E+06	2.81E-07	12.42	1.26	4.69	1.34
IgG1/2	N434S/Y436W	2.20E+05	0.0754	2.91E+06	3.44E-07	10.15	1.03	3.42	0.97
IgG1	Y436V	1.99E+05	0.136	1.46E+06	6.83E-07	5.11	0.52	1.90	0.54
IgG1	Y436W	2.72E+05	0.192	1.42E+06	7.06E-07	4.94	0.50	1.34	0.38
IgG1/2	N434S/R255I	1.86E+05	0.0726	2.57E+06	3.90E-07	8.95	0.91	3.55	1.01
IgG1/2	N434S/V284E	2.04E+05	0.0741	2.75E+06	3.63E-07	9.61	0.97	3.48	0.99
IgG1/2	N434S/N286D	1.82E+05	0.077	2.36E+06	4.23E-07	8.25	0.83	3.35	0.95
IgG1/2	N434S/N286L	1.98E+05	0.0738	2.69E+06	3.72E-07	9.38	0.95	3.50	1.00
IgG1/2	N434S/N286R	1.99E+05	0.0795	2.50E+06	4.00E-07	8.73	0.88	3.25	0.92
IgG1/2	N434S/T307Q	2.42E+05	0.0397	6.10E+06	1.64E-07	21.28	2.15	6.50	1.85
IgG1/2	N434S/V309W	2.62E+05	0.0852	3.07E+06	3.26E-07	10.71	1.08	3.03	0.86
IgG1/2	N434S/A378V	1.77E+05	0.0528	3.35E+06	2.99E-07	11.67	1.18	4.89	1.39
IgG1/2	N434S/S426V	2.63E+05	0.074	3.56E+06	2.81E-07	12.42	1.26	3.49	0.99
IgG1/2	N434S/S426I	1.81E+05	0.0809	2.23E+06	4.48E-07	7.79	0.79	3.19	0.91
IgG1/2	N434S/S426T	2.41E+05	0.0713	3.38E+06	2.96E-07	11.79	1.19	3.62	1.03
IgG1/2	Y436I	1.76E+05	0.123	1.43E+06	6.99E-07	4.99	0.51	2.10	0.60
IgG1/2	Y436V	ND	ND	ND	ND				
IgG1/2	Y436W	5.04E+05	0.16	3.15E+06	3.17E-07	11.01	1.11	1.61	0.46

Figure 17




Figure 18




Figure 19

A378T	A378M	S426G	S426Q
A378I	A378F	S426A	S426K
A378L	A378Y	S426P	S426R
A378E	A378R		

Figure 20

T307Q/N434S/Q311I	N434S/Y436V/A378V
T307Q/N434S/Q311V	N434S/Y436V/A378T
T307Q/N434S/A378V	N434S/Y436V/A378I
T307Q/N434S/S426V	N434S/Y436V/A378L
T307Q/N434S/Y436I	N434S/Y436V/S426G
T307Q/N434S/Y436V	N434S/Y436V/S426A
	N434S/Y436V/S426Q
Q311I/N434S/A378V	N434S/Y436V/S426P
Q311I/N434S/A378T	N434S/Y436V/S426L
Q311I/N434S/A378I	
Q311I/N434S/A378L	M428L/N434S/T307Q
Q311I/N434S/S426V	M428L/N434S/Q311I
Q311I/N434S/Y436I	M428L/N434S/Q311V
Q311I/N434S/Y436V	M428L/N434S/A378V
Q311I/S426V	M428L/N434S/S426V
Q311I/Y436I	M428L/N434S/Y436I
Q311I/Y436V	M428L/N434S/Y436V
Q311I/S426V/Y436I	
Q311I/S426V/Y436V	V308P/N434S
	A378T/N434S
Q311V/N434S/A378V	A378I/N434S
Q311V/N434S/S426V	A378L/N434S
Q311V/N434S/Y436I	A378E/N434S
Q311V/N434S/Y436V	A378M/N434S
	A378F/N434S
S426V/N434S/A378V	A378Y/N434S
S426V/N434S/A378T	A378R/N434S
S426V/N434S/A378I	S426G/N434S
S426V/N434S/A378L	S426A/N434S
S426V/N434S/Y436I	S426P/N434S
S426V/N434S/Y436V	S426Q/N434S
	S426K/N434S
N434S/Y436I/A378V	S426R/N434S
N434S/Y436I/A378T	
N434S/Y436I/A378I	T307Q/Q311I
N434S/Y436I/A378L	T307Q/Q311V
	Q311I
	Q311V

Figure 21

Antibody	KD (M)	Fold IgG1/2
IgG1/2_WT	1.18E-06	1.0
N434S	4.89E-07	2.4
M428L/N434S	2.15E-07	5.5
Q311I/N434S	1.92E-07	6.1
N434S/Y436I	3.42E-07	3.5
T307Q/N434S/Q311I	7.49E-08	15.8
Q311I/N434S/A378I	1.72E-07	6.9
Q311I/N434S/S426V	1.80E-07	6.6
Q311I/N434S/Y436I	1.06E-07	11.1
Q311I/N434S/Y436V	1.65E-07	7.2
Q311I/Y436I	5.23E-07	2.3
Q311I/Y436V	5.68E-07	2.1
Q311I/S426V/Y436I	4.74E-07	2.5
Q311V/N434S/S426V	1.94E-07	6.1
Q311V/N434S/Y436I	1.23E-07	9.6
Q311V/N434S/Y436V	2.06E-07	5.7
S426V/N434S/Y436I	3.10E-07	3.8
N434S/Y436I/A378I	2.24E-07	5.3
M428L/N434S/T307Q	8.90E-08	13.3
M428L/N434S/Q311I	7.49E-08	15.8
M428L/N434S/Y436I	9.16E-08	12.9
V308P/N434S	1.46E-07	8.1

Figure 22




Figure 23





Figure 24

Group	Variant	n	Individual mice t _{1/2} (days)					Average t _{1/2} (days)	St. Dev. (days)
			n1	n2	n3	n4	n5		
7349	IgG1/2_WT	4	2.9	2.5	3.2	2.8		2.9	0.3
7350	IgG1/2_N434S	4	6.3	7.7	7.3	6.5		7.0	0.7
7368	IgG1/2_Q311I/N434S	5	9.4	8.7	8.9	11.6	7.9	9.3	1.4
7377	IgG1/2_Q311V/N434S	5	9.4	8.7	8.9	8.2	8.4	8.7	0.4
7437	IgG1/2_N434S/Y436I	4	9.2	11.2	14.3	10.0		11.2	2.2
7447	IgG1/2_N434S/Y436V	4	9.4	9.5	8.5	5.7		8.3	1.8
7626	IgG1/2_M428L/N434S	4	16.1	14.8	13.7	9.3		13.5	3.0

Figure 25



FC VARIANTS THAT EXTEND ANTIBODY HALF-LIFE

[0001] This application claims the benefit under 35 U.S.C. 119 to U.S. Provisional Application Ser. No. 61/392,115, filed Oct. 12, 2010, and is also a continuation-in-part of U.S. Ser. No. 12/341,769, filed Dec. 22, 2008, which are both incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] The present application relates to optimized IgG immunoglobulin variants that extend antibody half-life in vivo, and their application, particularly for therapeutic purposes.

BACKGROUND OF THE INVENTION

[0003] Antibodies are immunological proteins that bind a specific antigen. In most mammals, including humans and mice, antibodies are constructed from paired heavy and light polypeptide chains. Each chain is made up of individual immunoglobulin (Ig) domains, and thus the generic term immunoglobulin is used for such proteins. Each chain is made up of two distinct regions, referred to as the variable and constant regions. The light and heavy chain variable regions show significant sequence diversity between antibodies, and are responsible for binding the target antigen. The constant regions show less sequence diversity, and are responsible for binding a number of natural proteins to elicit important biochemical events. In humans there are five different classes of antibodies including IgA (which includes subclasses IgA1 and IgA2), IgD, IgE, IgG (which includes subclasses IgG1, IgG2, IgG3, and IgG4), and IgM. The distinguishing feature between these antibody classes is their constant regions, although subtler differences may exist in the V region. IgG antibodies are tetrameric proteins composed of two heavy chains and two light chains. The IgG heavy chain is composed of four immunoglobulin domains linked from N- to C-terminus in the order VH—CH1-CH2-CH3, referring to the heavy chain variable domain, heavy chain constant domain 1, heavy chain constant domain 2, and heavy chain constant domain 3 respectively (also referred to as VH—C γ 1-C γ 2-C γ 3, referring to the heavy chain variable domain, constant gamma 1 domain, constant gamma 2 domain, and constant gamma 3 domain respectively). The IgG light chain is composed of two immunoglobulin domains linked from N- to C-terminus in the order VL-CL, referring to the light chain variable domain and the light chain constant domain respectively.

[0004] In IgG, a site on Fc between the C γ 2 and C γ 3 domains mediates interaction with the neonatal receptor FcRn. Binding to FcRn recycles endocytosed antibody from the endosome back to the bloodstream (Raghavan et al., 1996, *Annu Rev Cell Dev Biol* 12:181-220; Ghetie et al., 2000, *Annu Rev Immunol* 18:739-766, both entirely incorporated by reference). This process, coupled with preclusion of kidney filtration due to the large size of the full-length molecule, results in favorable antibody serum half-lives ranging from one to three weeks. Binding of Fc to FcRn also plays a key role in antibody transport. The binding site on Fc for FcRn is also the site at which the bacterial proteins A and G bind. The tight binding by these proteins is typically exploited as a means to purify antibodies by employing protein A or protein G affinity chromatography during protein purification. Thus the fidelity of this region on Fc is important for both the clinical properties of antibodies and their purification. Avail-

able structures of the rat Fc/FcRn complex (Burmeister et al., 1994, *Nature*, 372:379-383; Martin et al., 2001, *Mol Cell* 7:867-877, both entirely incorporated by reference), and of the complexes of Fc with proteins A and G (Deisenhofer, 1981, *Biochemistry* 20:2361-2370; Sauer-Eriksson et al., 1995, *Structure* 3:265-278; Tashiro et al., 1995, *Curr Opin Struct Biol* 5:471-481, all entirely incorporated by reference), provide insight into the interaction of Fc with these proteins. The FcRn receptor is also responsible for the transfer of IgG to the neonatal gut and to the lumen of the intestinal epithelia in adults (Ghetie and Ward, *Annu. Rev. Immunol.*, 2000, 18:739-766; Yoshida et al., *Immunity*, 2004, 20(6):769-783, both entirely incorporated by reference).

[0005] Antibodies have serum half-lives in vivo ranging from one to three weeks. This favorable property is due to the preclusion of kidney filtration due to the large size of the full-length molecule, and interaction of the antibody Fc region with the neonatal Fc receptor FcRn. Binding to FcRn recycles endocytosed antibody from the endosome back to the bloodstream (Raghavan et al., 1996, *Annu Rev Cell Dev Biol* 12:181-220; Ghetie et al., 2000, *Annu Rev Immunol* 18:739-766, both entirely incorporated by reference).

[0006] Other properties of the antibody may determine its clearance rate (e.g. stability and half-life) in vivo. In addition to antibody binding to the FcRn receptor, other factors that contribute to clearance and half-life are serum aggregation, enzymatic degradation in the serum, inherent immunogenicity of the antibody leading to clearing by the immune system, antigen-mediated uptake, FcR (non-FcRn) mediated uptake and non-serum distribution (e.g. in different tissue compartments).

[0007] There is a need for antibody modifications that improve the pharmacokinetic properties of antibodies. The present application meets these and other needs and provides novel engineered variants in the constant regions to improve serum half-life.

BRIEF SUMMARY OF THE INVENTION

Problem to be Solved

[0008] Accordingly, one problem to be solved is to increase serum half life of antibodies by altering the constant domains, thus allowing the same constant regions to be used with different antigen binding sequences, e.g. the variable regions including the CDRs, and minimizing the possibility of immunogenic alterations. Thus providing antibodies with constant region variants with extended half-life provides a modular approach to improving the pharmacokinetic properties of antibodies, as described herein. In addition, due to the methodologies outlined herein, the possibility of immunogenicity resulting from the FcRn variants is significantly reduced by importing variants from different IgG isotypes such that serum half-life is increased without introducing significant immunogenicity.

SUMMARY

[0009] In one aspect, the present invention provides an antibody comprising a variant Fc region as compared to a parent Fc region, wherein the variant Fc region comprises a first mutation which is a serine at position 434 and a second mutation selected from the group: an isoleucine at position 311, a valine at position 311, an isoleucine at position 436, and a valine at position 436. In a further aspect, the antibody

has increased serum half-life as compared to an antibody comprising the parent Fc region, wherein numbering is according to the EU index. In a still further aspect, the antibody has increased binding affinity to a human FcRn receptor as compared to an antibody comprising the parent Fc region, wherein numbering is according to the EU index.

[0010] In one embodiment, the present invention provides nucleic acids encoding any of the variant Fc regions described herein.

[0011] In a further embodiment, the present invention provides host cells comprising a nucleic acid encoding any of the variant Fc regions described herein.

[0012] In one aspect, the present invention provides methods of administering an antibody to a subject, where the antibody comprises a variant Fc region as compared to a parent Fc region, wherein the variant Fc region comprises a first mutation which is a serine at position 434 and a second mutation selected from the group: an isoleucine at position 311, a valine at position 311, an isoleucine at position 436, and a valine at position 436, where the antibody has increased serum half-life as compared to an antibody comprising the parent Fc region, and wherein numbering is according to the EU index.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] FIG. 1. Sequence alignments of human IgG constant heavy chains. Gray indicates differences from IgG1, and boxed residues indicate common allotypic variations in the human population.

[0014] FIG. 2. (SEQ ID NO: 1-6) Amino acid sequences of constant regions used in the invention.

[0015] FIG. 3. (SEQ ID NO: 7-8) Amino acid sequences of exemplary variant constant regions.

[0016] FIG. 4. (SEQ ID NO: 9-10) Amino acid sequences of VH and VL variable regions used in the invention.

[0017] FIG. 5. Relative VEGF binding by WT and select variant IgG1 anti-VEGF antibodies. The plot shows the Biacore response units (RU) at the end of the association phase for binding of antibody analyte to immobilized VEGF antigen. Anti-Her2 IgG1 antibody was used as a negative control.

[0018] FIG. 6. Biacore sensorgrams of WT and variant IgG1 antibodies to immobilized human FcRn at low (6.0) and high (7.4) pH.

[0019] FIG. 7. FcRn binding affinities of WT and select variant IgG1 antibodies to human FcRn at pH 6.0 as determined by Biacore. The graph shows a plot of the pseudo-affinity constant (K_a^*), on a log scale.

[0020] FIG. 8a-8b. In vivo pharmacokinetics of WT and variant antibodies in mFcRn^{-/-} hFcRn⁺ mice. The graphs plot the serum concentration of antibody versus time after a single intravenous dose. FIG. 8a shows data from one of the 4 studies carried out with IgG1 antibodies (Study 3), and FIG. 8b shows data from a study carried out with IgG2 antibodies (Study 5).

[0021] FIG. 9. Fitted PK parameters from all in vivo PK studies carried out in mFcRn^{-/-} hFcRn⁺ mice with variant and WT antibodies, n represents the number of mice per group, with Mean and standard deviation (SD) data provided for PK parameters. Half-Life represents the beta phase that characterizes elimination of antibody from serum. C_{max} is the maximal observed serum concentration, AUC is the area under the concentration time curve, and clearance is the clearance of antibody from serum. Fold half-life is calculated as

the half-life of variant antibody over that of the WT IgG1 or IgG2 parent within each study.

[0022] FIG. 10a-10b. Relative binding of variant IgG1 anti-VEGF antibodies to cynomolgus monkey and human FcRn as determined by Biacore. FIG. 10a shows the data in tabular form. FIG. 10b shows a plot of the data.

[0023] FIG. 11. In vivo pharmacokinetics of WT and variant antibodies in cynomolgus monkeys. The graphs plot the serum concentration of antibody versus time after a single intravenous dose.

[0024] FIG. 12. Fitted PK parameters from the in vivo PK study in cynomolgus monkeys with variant and WT antibodies. Parameters are as described in FIG. 9.

[0025] FIG. 13. Designed Fc variants. Variants are screened using the described methods in order to obtain modifications that extend the in vivo half-life of antibodies.

[0026] FIG. 14. Screen of engineered Fc variants for binding to human FcRn. The table shows the off-rate (k_{off}) for binding of each variant to human FcRn at pH 6.0 by Biacore.

[0027] FIG. 15. Graph of k_{off} for screened Fc variants (data plotted are from FIG. 14).

[0028] FIG. 16. Affinities of select variants for human FcRn at pH 6.0 based on a FcRn concentration series Biacore experiment. The data show the on and off kinetic rate constants (k_{on} and k_{off} respectively), and the association and dissociation equilibrium constants (K_A and K_D respectively), as well as the fold improvement in K_D and k_{off} relative to WT IgG1 and IgG1/2N434S.

[0029] FIG. 17. Plot off affinities of select variants for human FcRn at pH 6.0 as determined by Biacore. Values are plotted from FIG. 16.

[0030] FIG. 18. Plot off affinities of select variants for human FcRn at pH 6.0 as determined by Biacore. Values are plotted from FIG. 16.

[0031] FIG. 19. Designed single variants

[0032] FIG. 20. Designed combination variants

[0033] FIG. 21. Affinities of select variants for human FcRn at pH 6.0 based on a FcRn concentration series Biacore experiment. The data show the dissociation equilibrium constant K_D , as well as the fold improvement in K_D relative to the IgG1/2 parent.

[0034] FIG. 22. Plot off affinities of select variants for human FcRn at pH 6.0 as determined by Biacore. Values are plotted from FIG. 21.

[0035] FIG. 23. In vivo pharmacokinetics of IgG1/2 variant antibodies in mFcRn^{-/-} hFcRn⁺ mice. The graphs plot the serum concentration of antibody versus time after a single intravenous dose.

[0036] FIG. 24. Fitted half-lives ($t_{1/2}$) from the PK study in hFcRn⁺ mice, n=number of mice, columns n1-n5 show the half-lives of the individual mice in each group, and the average half-life and standard deviation are shown in the last two columns.

[0037] FIG. 25. Scatter plot of half-life results from PK study in hFcRn⁺ mice. Data are plotted from FIG. 24. Each dot represents the $t_{1/2}$ of an individual mouse within each variant group, and the line indicates the average.

DETAILED DESCRIPTION OF THE INVENTION

I. Overview

[0038] The present invention discloses the generation of novel variants of Fc domains, including those found in antibodies, Fc fusions, and immuno-adhesions, which have an

increased binding to the FcRn receptor. As noted herein, binding to FcRn results in longer serum retention in vivo. These variants of Fc domains are also referred to herein as “Fc variants” or “FcRn variants”.

[0039] The substitutions in the Fc domains (“Fc substitutions”) are chosen such that the resultant proteins (“Fc proteins”) show improved serum half-life in vivo as compared to the wild type protein. In order to increase the retention of the Fc proteins in vivo, the increase in binding affinity must be at around pH 6 while maintaining lower affinity at around pH 7.4. Although still under examination, Fc regions are believed to have longer half-lives in vivo, because binding to FcRn at pH 6 in an endosome sequesters the Fc (Ghetie and Ward, 1997 *Immunol Today*. 18(12): 592-598, entirely incorporated by reference). The endosomal compartment then recycles the Fc to the cell surface. Once the compartment opens to the extracellular space, the higher pH, ~7.4, induces the release of Fc back into the blood. In mice, Dall’Acqua et al. showed that Fc mutants with increased FcRn binding at pH 6 and pH 7.4 actually had reduced serum concentrations and the same half life as wild-type Fc (Dall’Acqua et al. 2002, *J. Immunol.* 169:5171-5180, entirely incorporated by reference). The increased affinity of Fc for FcRn at pH 7.4 is thought to forbid the release of the Fc back into the blood. Therefore, the Fc mutations that will increase Fc’s half-life in vivo will ideally increase FcRn binding at the lower pH while still allowing release of Fc at higher pH. The amino acid histidine changes its charge state in the pH range of 6.0 to 7.4. Therefore, it is not surprising to find His residues at important positions in the Fc/FcRn complex.

[0040] An additional aspect of the invention is the increase in FcRn binding over wild type specifically at lower pH, about pH 6.0, to facilitate Fc/FcRn binding in the endosome. Also disclosed are Fc variants with altered FcRn binding and altered binding to another class of Fc receptors, the FcγR’s (sometimes written FcγR’s) as differential binding to FcγRs, particularly increased binding to FcγRIIb and decreased binding to FcγRIIb, has been shown to result in increased efficacy.

[0041] In addition, many embodiments of the invention rely on the “importation” of substitutions at particular positions from one IgG isotype into another, thus reducing or eliminating the possibility of unwanted immunogenicity being introduced into the variants. That is, IgG1 is a common isotype for therapeutic antibodies for a variety of reasons, including high effector function. IgG2 residues at particular positions can be introduced into the IgG1 backbone to result in a protein that exhibits longer serum half-life.

[0042] In other embodiments, non-isotypic amino acid changes are made, to improve binding to FcRn and/or to increase in vivo serum half-life, and/or to allow accommodations in structure for stability, etc. as is more further described below.

[0043] As will be appreciated by those in the art and described below, a number of factors contribute to the in vivo clearance, and thus the half-life, of antibodies in serum. One factor involves the antigen to which the antibody binds; that is, antibodies with identical constant regions but different variable regions (e.g. Fv domains), may have different half-lives due to differential ligand binding effects. However, the present invention demonstrates that while the absolute half life of two different antibodies may differ due to these antigen specificity effects, the FcRn variants described herein, can transfer to different ligands to give the same trends of increas-

ing half-life. That is, in general, the relative “order” of the FcRn binding/half life increases will track to antibodies with the same variants of antibodies with different Fvs as is discussed herein.

II. Description of the Invention

A. Antibodies

[0044] The present invention relates to the generation of Fc variants of antibodies, generally therapeutic antibodies. As is discussed below, the term “antibody” is used generally. Antibodies that find use in the present invention can take on a number of formats as described herein, including traditional antibodies as well as antibody derivatives, fragments and mimetics, described below. In general, the term “antibody” includes any polypeptide that includes at least one constant domain, including, but not limited to, CH1, CH2, CH3 and CL.

[0045] Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one “light” (typically having a molecular weight of about 25 kDa) and one “heavy” chain (typically having a molecular weight of about 50-70 kDa). Human light chains are classified as kappa and lambda light chains. The present invention is directed to the IgG class, which has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. Thus, “isotype” as used herein is meant any of the subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. It should be understood that therapeutic antibodies can also comprise hybrids of isotypes and/or subclasses. For example, as shown herein, the present invention covers Fc variant engineering of IgG1/G2 hybrids.

[0046] The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition, generally referred to in the art and herein as the “Fv domain” or “Fv region”. In the variable region, three loops are gathered for each of the V domains of the heavy chain and light chain to form an antigen-binding site. Each of the loops is referred to as a complementarity-determining region (hereinafter referred to as a “CDR”), in which the variation in the amino acid sequence is most significant. “Variable” refers to the fact that certain segments of the variable region differ extensively in sequence among antibodies. Variability within the variable region is not evenly distributed. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called “hypervariable regions” that are each 9-15 amino acids long or longer.

[0047] Each VH and VL is composed of three hypervariable regions (“complementary determining regions,” “CDRs”) and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4.

[0048] The hypervariable region generally encompasses amino acid residues from about amino acid residues 24-34 (LCDR1; “L” denotes light chain), 50-56 (LCDR2) and 89-97 (LCDR3) in the light chain variable region and around about 31-35B (HCDR1; “H” denotes heavy chain), 50-65 (HCDR2), and 95-102 (HCDR3) in the heavy chain variable region; Kabat et al., SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, 5th Ed. Public Health Ser-

vice, National Institutes of Health, Bethesda, Md. (1991) and/or those residues forming a hypervariable loop (e.g. residues 26-32 (LCDR1), 50-52 (LCDR2) and 91-96 (LCDR3) in the light chain variable region and 26-32 (HCDR1), 53-55 (HCDR2) and 96-101 (HCDR3) in the heavy chain variable region; Chothia and Lesk (1987) J. Mol. Biol. 196:901-917. Specific CDRs of the invention are described below.

[0049] Throughout the present specification, the Kabat numbering system is generally used when referring to a residue in the variable domain (approximately, residues 1-107 of the light chain variable region and residues 1-113 of the heavy chain variable region) (e.g. Kabat et al., supra (199.1)).

[0050] The CDRs contribute to the formation of the antigen-binding, or more specifically, epitope binding site of antibodies. "Epitope" refers to a determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. Epitopes are groupings of molecules such as amino acids or sugar side chains and usually have specific structural characteristics, as well as specific charge characteristics. A single antigen may have more than one epitope.

[0051] The epitope may comprise amino acid residues directly involved in the binding (also called immunodominant component of the epitope) and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen binding peptide; in other words, the amino acid residue is within the footprint of the specifically antigen binding peptide.

[0052] Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. Conformational and nonconformational epitopes may be distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

[0053] An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Antibodies that recognize the same epitope can be verified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen, for example "binning."

[0054] As will be appreciated by those in the art, a wide variant of antigen binding domains, e.g. Fv regions, may find use in the present invention. Virtually any antigen may be targeted by the IgG variants, including but not limited to proteins, subunits, domains, motifs, and/or epitopes belonging to the following list of target antigens, which includes both soluble factors such as cytokines and membrane-bound factors, including transmembrane receptors: 17-1A, 4-1BB, 4Dc, 6-keto-PGF1a, 8-iso-PGF2a, 8-oxo-dG, A1 Adenosine Receptor, A33, ACE, ACE-2, Activin, Activin A, Activin AB, Activin B, Activin C, Activin RIA, Activin RIA ALK-2, Activin RIB ALK-4, Activin RIIA, Activin RIIIB, ADAM, ADAM10, ADAM12, ADAM15, ADAM17/TACE, ADAMS, ADAM9, ADAMTS, ADAMTS4, ADAMTS5, Addressins, aFGF, ALCAM, ALK, ALK-1, ALK-7, alpha-1-antitrypsin, alpha-V/beta-1 antagonist, ANG, Ang, APAF-1, APE, APJ, APP, APRIL, AR, ARC, ART, Artemin, anti-Id, ASPARTIC, Atrial natriuretic factor, av/b3 integrin, Ax1, b2M, B7-1, B7-2, B7-H, B-lymphocyte Stimulator (BlyS), BACE, BACE-1, Bad, BAFF, BAFF-R, Bag-1, BAK, Bax, BCA-1, BCAM, Bcl, BCMA, BDNF, b-ECGF, bFGF, BID,

Bik, BIM, BLC, BL-CAM, BLK, BMP, BMP-2 BMP-2a, BMP-3 Osteogenin, BMP-4 BMP-2b, BMP-5, BMP-6 Vgr-1, BMP-7 (OP-1), BMP-8 (BMP-8a, OP-2), BMPR, BMPR-1A (ALK-3), BMPR-1B (ALK-6), BRK-2, RPK-1, BMPR-II (BRK-3), BMPs, b-NGF, BOK, Bombesin, Bone-derived neurotrophic factor, BPDE, BPDE-DNA, BTC, complement factor 3 (C3), C3a, C4, C5, C5a, C10, CA125, CAD-8, Calcitonin, cAMP, carcinoembryonic antigen (CEA), carcinoma-associated antigen, Cathepsin A, Cathepsin B, Cathepsin C/DPPI, Cathepsin D, Cathepsin E, Cathepsin H, Cathepsin L, Cathepsin O, Cathepsin S, Cathepsin V, Cathepsin X/Z/P, CBL, CCI, CCK2, CCL, CCL1, CCL11, CCL12, CCL13, CCL14, CCL15, CCL16, CCL17, CCL18, CCL19, CCL2, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCL26, CCL27, CCL28, CCL3, CCL4, CCL5, CCL6, CCL7, CCL8, CCL9/10, CCR, CCR1, CCR10, CCR10, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CD1, CD2, CD3, CD3E, CD4, CD5, CD6, CD7, CD8, CD10, CD11a, CD11b, CD11c, CD13 CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD27L, CD28, CD29, CD30, CD30L, CD32, CD33 (p67 proteins), CD34, CD38, CD40, CD40L, CD44, CD45, CD46, CD49a, CD52, CD54, CD55, CD56, CD61, CD64, CD66e, CD74, CD80 (B7-1), CD89, CD95, CD123, CD137, CD138, CD140a, CD146, CD147, CD148, CD152, CD164, CEACAM5, CFTR, cGMP, CINC, *Clostridium botulinum* toxin, *Clostridium perfringens* toxin, CKb8-1, CLC, CMV, CMV UL, CNTF, CNTN-1, COX, C-Ret, CRG-2, CT-1, CTACK, CTGF, CTLA-4, CX3CL1, CX3CR1, CXCL, CXCL1, CXCL2, CXCL3, CXCL4, CXCL5, CXCL6, CXCL7, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, CXCL14, CXCL15, CXCL16, CXCR, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, cytokeratin tumor-associated antigen, DAN, DCC, DcR3, DC-SIGN, Decay accelerating factor, des(1-3)-IGF-I (brain IGF-I), Dhh, digoxin, DNAM-1, Dnase, Dpp, DPPIV/CD26, Dtk, ECAD, EDA, EDA-A1, EDA-A2, EDAR, EGF, EGFR (ErbB-1), EMA, EMMPRIN, ENA, endothelin receptor, Enkephalinase, eNOS, Eot, eotaxin1, EpCAM, Ephrin B2/EphB4, EPO, ERCC, E-selectin, ET-1, Factor IIa, Factor VII, Factor VIIIc, Factor IX, fibroblast activation protein (FAP), Fas, FcR1, FEN-1, Ferritin, FGF, FGF-19, FGF-2, FGF3, FGF-8, FGFR, FGFR-3, Fibrin, FL, FLIP, Flt-3, Flt-4, Follicle stimulating hormone, Fractalkine, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, FZD10, G250, Gas 6, GCP-2, GCSF, GD2, GD3, GDF, GDF-1, GDF-3 (Vgr-2), GDF-5 (BMP-14, CDMP-1), GDF-6 (BMP-13, CDMP-2), GDF-7 (BMP-12, CDMP-3), GDF-8 (Myostatin), GDF-9, GDF-15 (MIC-1), GDNF, GDNF, GFAP, GFRa-1, GFR-alpha1, GFR-alpha2, GFR-alpha3, GTR, Glucagon, Glut 4, glycoprotein IIb/IIIa (GP IIb/IIIa), GM-CSF, gp130, gp72, GRO, Growth hormone releasing factor, Hapten (NP-cap or NIP-cap), HB-EGF, HCC, HCMV gB envelope glycoprotein, HCMV) gH envelope glycoprotein, HCMV UL, Hemopoietic growth factor (HGF), Hep B gp120, heparanase, Her2, Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), herpes simplex virus (HSV) gB glycoprotein, HSV gD glycoprotein, HGFA, High molecular weight melanoma-associated antigen (HMW-MAA), HIV gp120, HIV IIIB gp120 V3 loop, HLA, HLA-DR, HMI.24, HMFG PEM, HRG, Hrk, human cardiac myosin, human cytomegalovirus (HCMV), human growth hormone (HGH), HVEM, 1-309, IAP, ICAM, ICAM-1, ICAM-3, ICE, ICOS, IFNg, Ig, IgA receptor, IgE, IGF, IGF binding proteins, IGF-1R, IGFBP, IGF-I, IGF-II, IL, IL-1,

IL-1R, IL-2, IL-2R, IL-4, IL-4R, IL-5, IL-5R, IL-6, IL-6R, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-18R, IL-23, interferon (INF)-alpha, INF-beta, INF-gamma, Inhibin, iNOS, Insulin A-chain, Insulin B-chain, Insulin-like growth factor 1, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha4/beta1, integrin alpha4/beta7, integrin alpha5 (alphaV), integrin alpha5/beta1, integrin alpha5/beta3, integrin alpha6, integrin beta1, integrin beta2, interferon gamma, IP-10, I-TAC, JE, Kallikrein 2, Kallikrein 5, Kallikrein 6, Kallikrein 11, Kallikrein 12, Kallikrein 14, Kallikrein 15, Kallikrein L1, Kallikrein L2, Kallikrein L3, Kallikrein L4, KC, KDR, Keratinocyte Growth Factor (KGF), laminin 5, LAMP, LAP, LAP (TGF-1), Latent TGF-1, Latent TGF-1 bp1, LBP, LDGF, LECT2, Lefty, Lewis-Y antigen, Lewis-Y related antigen, LFA-1, LFA-3, Lfo, LIF, LIGHT, lipoproteins, LIX, LKN, Lptn, L-Selectin, LT-a, LT-b, LTB4, LTBP-1, Lung surfactant, Luteinizing hormone, Lymphotoxin Beta Receptor, Mac-1, MAdCAM, MAG, MAP2, MARC, MCAM, MCAM, MCK-2, MCP, M-CSF, MDC, Mer, MET-ALLOPROTEASES, MGDF receptor, MGMT, MHC (HLA-DR), MIF, MIG, MIP, MIP-1-alpha, MK, MMAC1, MMP, MMP-1, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-2, MMP-24, MMP-3, MMP-7, MMP-8, MMP-9, MPIF, Mpo, MSK, MSP, mucin (Muc1), MUC18, Muellierian-inhibiting substance, Mug, MuSK, NAIP, NAP, NCAD, N-Cadherin, NCA 90, NCAM, NCAM, Neprilysin, Neurotrophin-3, -4, or -6, Neurturin, Neuronal growth factor (NGF), NGFR, NGF-beta, nNOS, NO, NOS, Npn, NRG-3, NT, NTN, OB, OGG1, OPG, OPN, OSM, OX40L, OX40R, p150, p95, PADPr, Parathyroid hormone, PARC, PARP, PBR, PBSE, PCAD, P-Cadherin, PCNA, PDGF, PDGF, PDK-1, PECAM, PEM, PF4, PGE, PGE, PGI2, PGJ2, PIN, PLA2, placental alkaline phosphatase (FLAP), PIGF, PLP, PP14, Proinsulin, Prorelaxin, Protein C, PS, PSA, PSCA, prostate specific membrane antigen (PSMA), PTEN, PTHrp, Ptk, PTN, R51, RANK, RANKL, RANTES, RANTES, Relaxin A-chain, Relaxin B-chain, renin, respiratory syncytial virus (RSV) F, RSV Fgp, Ret, Rheumatoid factors, RLIP76, RPA2, RSK, S100, SCF/KL, SDF-1, SERINE, Serum albumin, sFRP-3, Shh, SIGIRR, SK-1, SLAM, SLPI, SMAC, SMDF, SMOH, SOD, SPARC, Stat, STEAP, STEAP-II, TACE, TACI, TAG-72 (tumor-associated glycoprotein-72), TARC, TCA-3, T-cell receptors (e.g., T-cell receptor alpha/beta), TdT, TECK, TEM1, TEM5, TEM7, TEM8, TERT, testicular PLAP-like alkaline phosphatase, Tfr, TGF, TGF-alpha, TGF-beta, TGF-beta Pan Specific, TGF-beta R1 (ALK-5), TGF-beta RII, TGF-beta RIIB, TGF-beta RIII, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta4, TGF-beta5, Thrombin, Thymus Ck-1, Thyroid stimulating hormone, Tie, TIMP, TIQ, Tissue Factor, TMEFF2, Tmpo, TMPRSS2, TNF, TNF-alpha, TNF-alpha beta, TNF-beta2, TNF α , TNF-RI, TNF-RII, TNFRSF10A (TRAIL R1 Apo-2, DR4), TNFRSF10B (TRAIL R2 DR5, KILLER, TRICK-2A, TRICK-B), TNFRSF10C (TRAIL R3 DR1, LIT, TRID), TNFRSF10D (TRAIL R4 DcR2, TRUND), TNFRSF11A (RANK ODF R, TRANCE R), TNFRSF11B (OPG OCIF, TR1), TNFRSF12 (TWEAK R FN14), TNFRSF13B (TACI), TNFRSF13C (BAFF R), TNFRSF14 (HVEM ATAR, HveA, LIGHT R, TR2), TNFRSF16 (NGFR p75NTR), TNFRSF17 (BCMA), TNFRSF18 (GITR AITR), TNFRSF19 (TROY TAJ, TRADE), TNFRSF19L (RELT), TNFRSF1A (TNF RI CD120a, p55-60), TNFRSF1B (TNF RII CD120b, p75-80), TNFRSF26 (TNFRH3), TNFRSF3 (LTbR TNF RUT, TNFC R), TNFRSF4 (OX40 ACT35, TXGP1 R), TNFRSF5 (CD40

p50), TNFRSF6 (Fas Apo-1, APT1, CD95), TNFRSF6B (DcR3 M68, TR6), TNFRSF7 (CD27), TNFRSF8 (CD30), TNFRSF9 (4-1BB CD137, ILA), TNFRSF21 (DR6), TNFRSF22 (DcTRAIL R2 TNFRH2), TNFRST23 (Dc-TRAIL R1 TNFRH1), TNFRSF25 (DR3 Apo-3, LARD, TR-3, TRAMP, WSL-1), TNFSF10 (TRAIL Apo-2 Ligand, TL2), TNFSF11 (TRANCE/RANK Ligand ODF, OPG Ligand), TNFSF12 (TWEAK Apo-3 Ligand, DR3 Ligand), TNFSF13 (APRIL TALL2), TNFSF13B (BAFF BLYS, TALL 1, THANK, TNFSF20), TNFSF14 (LIGHT HVEM Ligand, LTg), TNFSF15 (TL1A/VEGI), TNFSF18 (GITR Ligand AITR Ligand, TL6), TNFSF1A (TNF-a Conectin, DIF, TNFSF2), TNFSF1B (TNF-b LTa, TNFSF1), TNFSF3 (LTb TNFC, p33), TNFSF4 (OX40 Ligand gp34, TXGP1), TNFSF5 (CD40 Ligand CD154, gp39, HIGM1, IMD3, TRAP), TNFSF6 (Fas Ligand Apo-1 Ligand, APT1 Ligand), TNFSF7 (CD27 Ligand CD70), TNFSF8 (CD30 Ligand CD153), TNFSF9 (4-1BB Ligand CD137 Ligand), TP-1, t-PA, Tpo, TRAIL, TRAIL R, TRAIL-R1, TRAIL-R2, TRANCE, transferring receptor, TRF, Trk, TROP-2, TSG, TSLP, tumor-associated antigen CA 125, tumor-associated antigen expressing Lewis Y related carbohydrate, TWEAK, TXB2, Ung, uPAR, uPAR-1, Urokinase, VCAM, VCAM-1, VECAD, VE-Cadherin, VE-cadherin-2, VEGFR-1 (flt-1), VEGF, VEGFR, VEGFR-3 (flt-4), VEGIVIM, Viral antigens, VLA, VLA-1, VLA-4, VNR integrin, von Willebrands factor, WIF-1, WNT1, WNT2, WNT2B/13, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, WNT16, XCL1, XCL2, XCR1, XCR1, XEDAR, XIAP, XPD, and receptors for hormones and growth factors.

[0055] In some embodiments, the engineering of Fc domains described herein is done to therapeutic antibodies. A number of antibodies that are approved for use, in clinical trials, or in development may benefit from the Fc variants of the present invention. These antibodies are herein referred to as “clinical products and candidates”. Thus in a preferred embodiment, the engineered constant region(s) of the present invention may find use in a range of clinical products and candidates. For example, a number of antibodies that target CD20 may benefit from the Fc engineering of the present invention. For example the Fc variants of the present invention may find use in an antibody that is substantially similar to rituximab (Rituxan®, IDEC/Genentech/Roche) (see for example U.S. Pat. No. 5,736,137), a chimeric anti-CD20 antibody approved to treat Non-Hodgkin’s lymphoma; HuMax-CD20, an anti-CD20 currently being developed by Genmab, an anti-CD20 antibody described in U.S. Pat. No. 5,500,362, AME-133 (Applied Molecular Evolution), hA20 (Immunomedics, Inc.), HumaLYM (Intracel), and PRO70769 (PCT/US2003/040426, entitled “Immunoglobulin Variants and Uses Thereof”). A number of antibodies that target members of the family of epidermal growth factor receptors, including EGFR (ErbB-1), Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), may benefit from Fc engineered constant region(s) of the invention. For example the Fc engineered constant region(s) of the invention may find use in an antibody that is substantially similar to trastuzumab (Herceptin®, Genentech) (see for example U.S. Pat. No. 5,677,171), a humanized anti-Her2/neu antibody approved to treat breast cancer; pertuzumab (rhuMab-2C4, Omnitarg™), currently being developed by Genentech; an anti-Her2 antibody described in U.S. Pat. No. 4,753,894; cetuximab (Erbix®,

Imclone) (U.S. Pat. No. 4,943,533; PCT WO 96/40210), a chimeric anti-EGFR antibody in clinical trials for a variety of cancers; ABX-EGF (U.S. Pat. No. 6,235,883), currently being developed by Abgenix-Immimex-Amgen; HuMax-EGFr (U.S. Ser. No. 10/172,317), currently being developed by Genmab; 425, EMD55900, EMD62000, and EMD72000 (Merck KGaA) (U.S. Pat. No. 5,558,864; Murthy et al. 1987, Arch Biochem Biophys. 252(2):549-60; Rodeck et al., 1987, J Cell Biochem. 35(4):315-20; Kettleborough et al., 1991, Protein Eng. 4(7):773-83); ICR62 (Institute of Cancer Research) (PCT WO 95/20045; Modjtahedi et al., 1993, J. Cell Biophys. 1993, 22(1-3): 129-46; Modjtahedi et al., 1993, Br J Cancer. 1993, 67(2):247-53; Modjtahedi et al, 1996, Br J Cancer, 73(2):228-35; Modjtahedi et al, 2003, Int J Cancer, 105(2):273-80); TheraCIM hR3 (YM Biosciences, Canada and Centro de Immunologia Molecular, Cuba (U.S. Pat. No. 5,891,996; U.S. Pat. No. 6,506,883; Mateo et al, 1997, Immunotechnology, 3(1):71-81); mAb-806 (Ludwig Institute for Cancer Research, Memorial Sloan-Kettering) (Jungbluth et al. 2003, Proc Natl Acad Sci USA. 100(2):639-44); KSB-102 (KS Biomedix); MR1-1 (IVAX, National Cancer Institute) (PCT WO 0162931A2); and SC100 (Scancell) (PCT WO 01/88138). In another preferred embodiment, the Fc engineered constant region(s) of the present invention may find use in alemtuzumab (Campath®, Millenium), a humanized monoclonal antibody currently approved for treatment of B-cell chronic lymphocytic leukemia. The Fc engineered constant region(s) of the present invention may find use in a variety of antibodies that are substantially similar to other clinical products and candidates, including but not limited to muromonab-CD3 (Orthoclone OKT3®), an anti-CD3 antibody developed by Ortho Biotech/Johnson & Johnson, ibritumomab tiuxetan (Zevalin®), an anti-CD20 antibody developed by IDEC/Schering AG, gemtuzumab ozogamicin (Mylotarg®), an anti-CD33 (p67 protein) antibody developed by Celltech/Wyeth, alefacept (Amevive®), an anti-LFA-3 Fc fusion developed by Biogen), abciximab (ReoPro®), developed by Centocor/Lilly, basiliximab (Simulect®), developed by Novartis, palivizumab (Synagis®), developed by MedImmune, infliximab (Remicade®), an anti-TNF α antibody developed by Centocor, adalimumab (Humira®), an anti-TNF α antibody developed by Abbott, Humicade™, an anti-TNF α antibody developed by Celltech, etanercept (Enbrel®), an anti-TNF α Fc fusion developed by Immunex/Amgen, ABX-CBL, an anti-CD147 antibody being developed by Abgenix, ABX-IL8, an anti-IL8 antibody being developed by Abgenix, ABX-MA1, an anti-MUC18 antibody being developed by Abgenix, Pemtumomab (R1549, 90Y-muHMF1), an anti-MUC1 in development by Antisoma, Therex (R1550), an anti-MUC1 antibody being developed by Antisoma, AngioMab (A81405), being developed by Antisoma, HuBC-1, being developed by Antisoma, Thioplatin (AS1407) being developed by Antisoma, Antegren® (natalizumab), an anti- α -4- β -7 (VLA-4) and α -4- β -7 antibody being developed by Biogen, VLA-1 mAb, an anti-VLA-1 integrin antibody being developed by Biogen, LTBR mAb, an anti-lymphotoxin beta receptor (LTBR) antibody being developed by Biogen, CAT-152, an anti-TGF- β 2 antibody being developed by Cambridge Antibody Technology, J695, an anti-IL-12 antibody being developed by Cambridge Antibody Technology and Abbott, CAT-192, an anti-TGF β 1 antibody being developed by Cambridge Antibody Technology and Genzyme, CAT-213, an anti-Eotaxin1 antibody being developed by Cam-

bridge Antibody Technology, LymphoStat-B™ an anti-Blys antibody being developed by Cambridge Antibody Technology and Human Genome Sciences Inc., TRAIL-R1mAb, an anti-TRAIL-R1 antibody being developed by Cambridge Antibody Technology and Human Genome Sciences, Inc., Avastin™ (bevacizumab, rhuMAb-VEGF), an anti-VEGF antibody being developed by Genentech, an anti-HER receptor family antibody being developed by Genentech, Anti-Tissue Factor (ATF), an anti-Tissue Factor antibody being developed by Genentech, Xolair™ (Omalizumab), an anti-IgE antibody being developed by Genentech, Raptiva™ (Efalizumab), an anti-CD11a antibody being developed by Genentech and Xoma, MLN-02 Antibody (formerly LDP-02), being developed by Genentech and Millenium Pharmaceuticals, HuMax CD4, an anti-CD4 antibody being developed by Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Genmab and Amgen, HuMax-Inflam, being developed by Genmab and Medarex, HuMax-Cancer, an anti-Heparanase I antibody being developed by Genmab and Medarex and Oxford GeoSciences, HuMax-Lymphoma, being developed by Genmab and Amgen, HuMax-TAC, being developed by Genmab, IDEC-131, and anti-CD40L antibody being developed by IDEC Pharmaceuticals, IDEC-151 (Clenoliximab), an anti-CD4 antibody being developed by IDEC Pharmaceuticals, IDEC-114, an anti-CD80 antibody being developed by IDEC Pharmaceuticals, IDEC-152, an anti-CD23 being developed by IDEC Pharmaceuticals, anti-macrophage migration factor (MIF) antibodies being developed by IDEC Pharmaceuticals, BEC2, an anti-idiotypic antibody being developed by Imclone, IMC-1C11, an anti-KDR antibody being developed by Imclone, DC101, an anti-flk-1 antibody being developed by Imclone, anti-VE cadherin antibodies being developed by Imclone, CEA-Cide™ (labetuzumab), an anti-carcinoembryonic antigen (CEA) antibody being developed by Immunomedics, LymphoCide™ (Epratuzumab), an anti-CD22 antibody being developed by Immunomedics, AFP-Cide, being developed by Immunomedics, MyelomaCide, being developed by Immunomedics, LkoCide, being developed by Immunomedics, ProstaCide, being developed by Immunomedics, MDX-010, an anti-CTLA4 antibody being developed by Medarex, MDX-060, an anti-CD30 antibody being developed by Medarex, MDX-070 being developed by Medarex, MDX-018 being developed by Medarex, Osidem™ (IDM-1), and anti-Her2 antibody being developed by Medarex and Immuno-Designed Molecules, HuMax™-CD4, an anti-CD4 antibody being developed by Medarex and Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Medarex and Genmab, CNTO 148, an anti-TNF α antibody being developed by Medarex and Centocor/J&J, CNTO 1275, an anti-cytokine antibody being developed by Centocor/J&J, MOR101 and MOR102, anti-intercellular adhesion molecule-1 (ICAM-1) (CD54) antibodies being developed by MorphoSys, MOR201, an anti-fibroblast growth factor receptor 3 (FGFR-3) antibody being developed by MorphoSys, Nuvion® (visilizumab), an anti-CD3 antibody being developed by Protein Design Labs, HuZAF™, an anti-gamma interferon antibody being developed by Protein Design Labs, Anti- α 5 β 1 Integrin, being developed by Protein Design Labs, anti-IL-12, being developed by Protein Design Labs, ING-1, an anti-Ep-CAM antibody being developed by Xoma, and MLN01, an anti-Beta2 integrin antibody being developed by Xoma, an p1-ADC antibody being developed by

Seattle Genetics, all of the above-cited references in this paragraph are expressly incorporated herein by reference.

[0056] The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Kabat et al. collected numerous primary sequences of the variable regions of heavy chains and light chains. Based on the degree of conservation of the sequences, they classified individual primary sequences into the CDR and the framework and made a list thereof (see SEQUENCES OF IMMUNOLOGICAL INTEREST, 5th edition, NIH publication, No. 91-3242, E. A. Kabat et al., entirely incorporated by reference).

[0057] In the IgG subclass of immunoglobulins, there are several immunoglobulin domains in the heavy chain. By “immunoglobulin (Ig) domain” herein is meant a region of an immunoglobulin having a distinct tertiary structure. Of interest in the present invention are the heavy chain domains, including, the constant heavy (CH) domains and the hinge domains. In the context of IgG antibodies, the IgG isotypes each have three CH regions. Accordingly, “CH” domains in the context of IgG are as follows: “CH1” refers to positions 118-220 according to the EU index as in Kabat. “CH2” refers to positions 237-340 according to the EU index as in Kabat, and “CH3” refers to positions 341-447 according to the EU index as in Kabat. As shown herein and described below, the variants of the present invention can include substitutions in one or more of the CH regions, as well as the hinge region, discussed below.

[0058] It should be noted that the sequences depicted herein start at the CH1 region, position 118; the variable regions are not included except as noted. For example, the first amino acid of SEQ ID NO: 2, while designated as position “1” in the sequence listing, corresponds to position 118 of the CH1 region, according to EU numbering.

[0059] Another type of Ig domain of the heavy chain is the hinge region. By “hinge” or “hinge region” or “antibody hinge region” or “immunoglobulin hinge region” herein is meant the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 236 (G236 in IgG1), wherein the numbering is according to the EU index as in Kabat. In some embodiments, for example in the context of an Fc region, the lower hinge is included, with the “lower hinge” generally referring to positions 226 or 230. As noted herein, Fc variant variants can be made in the hinge region as well.

[0060] The light chain generally comprises two domains, the variable light domain (containing the light chain CDRs and together with the variable heavy domains forming the Fv region), and a constant light chain region (often referred to as CL or Ck).

[0061] Another region of interest for additional substitutions, outlined below, is the Fc region. By “Fc” or “Fc region” or “Fc domain” as used herein is meant the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, the Fc domain comprises immunoglobulin domains Cγ2 and Cγ3 (Cγ2 and

Cγ3) and the lower hinge region between Cγ1 (Cγ1) and Cγ2 (Cγ2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. In some embodiments, as is more fully described below, amino acid modifications are made to the Fc region, for example to alter binding to one or more FcγR receptors or to the FcRn receptor.

[0062] In some embodiments, the antibodies are full length. By “full length antibody” herein is meant the structure that constitutes the natural biological form of an antibody, including variable and constant regions, including one or more modifications as outlined herein.

[0063] Alternatively, the antibodies can be a variety of structures, including, but not limited to, antibody fragments, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as “antibody mimetics”), chimeric antibodies, humanized antibodies, antibody fusions (sometimes referred to as “antibody conjugates”), and fragments of each, respectively.

[0064] In one embodiment, the antibody is an antibody fragment, as long as it contains at least one constant domain which can be engineered. Specific antibody fragments include, but are not limited to, (i) the Fab fragment consisting of VL, VH, CL and CH1 domains, (ii) the Fd fragment consisting of the VH and CH1 domains, (iii) F(ab')₂ fragments, a bivalent fragment comprising two linked Fab fragments (vii) single chain Fv molecules (scFv), wherein a VH domain and a VL domain are linked by a peptide linker which allows the two domains to associate to form an antigen binding site (Bird et al., 1988, Science 242:423-426, Huston et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:5879-5883, entirely incorporated by reference), (iv) “diabodies” or “triabodies”, multivalent or multispecific fragments constructed by gene fusion (Tomlinson et al., 2000, Methods Enzymol. 326:461-479; WO94/13804; Holliger et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90:6444-6448, all entirely incorporated by reference).

[0065] Other antibody fragments that can be used include fragments that contain one or more of the CH1, CH2, CH3, hinge and CL domains of the invention that have been engineered. For example, Fc fusions are fusions of the Fc region (CH2 and CH3, optionally with the hinge region) fused to another protein. A number of Fc fusions are known the art and can be improved by the addition of the variants of the invention. In the present case, antibody fusions can be made comprising CH1; CH1, CH2 and CH3; CH2; CH3; CH2 and CH3; CH1 and CH3, any or all of which can be made optionally with the hinge region, utilizing any combination of Fc variants described herein.

B. Chimeric and Humanized Antibodies

[0066] In some embodiments, the antibody can be a mixture from different species, e.g. a chimeric antibody and/or a humanized antibody. In general, both “chimeric antibodies” and “humanized antibodies” refer to antibodies that combine regions from more than one species. For example, “chimeric antibodies” traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human. “Humanized antibodies” generally refer to non-human antibodies that have had the variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire

antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018, Jones, 1986, *Nature* 321:522-525, Verhoeyen et al., 1988, *Science* 239:1534-1536, all entirely incorporated by reference. "Backmutation" of selected acceptor framework residues to the corresponding donor residues is often required to regain affinity that is lost in the initial grafted construct (U.S. Pat. No. 5,530,101; U.S. Pat. No. 5,585,089; U.S. Pat. No. 5,693,761; U.S. Pat. No. 5,693,762; U.S. Pat. No. 6,180,370; U.S. Pat. No. 5,859,205; U.S. Pat. No. 5,821,337; U.S. Pat. No. 6,054,297; U.S. Pat. No. 6,407,213, all entirely incorporated by reference). The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin, and thus will typically comprise a human Fc region. Humanized antibodies can also be generated using mice with a genetically engineered immune system, Roque et al., 2004, *Biotechnol. Prog.* 20:639-654, entirely incorporated by reference. A variety of techniques and methods for humanizing and reshaping non-human antibodies are well known in the art (See Tsurushita & Vasquez, 2004, *Humanization of Monoclonal Antibodies*, *Molecular Biology of B Cells*, 533-545, Elsevier Science (USA), and references cited therein, all entirely incorporated by reference). Humanization methods include but are not limited to methods described in Jones et al., 1986, *Nature* 321:522-525; Riechmann et al., 1988, *Nature* 332:323-329; Verhoeyen et al., 1988, *Science*, 239:1534-1536; Queen et al., 1989, *Proc Natl Acad Sci USA* 86:10029-33; He et al., 1998, *J. Immunol.* 160: 1029-1035; Carter et al., 1992, *Proc Natl Acad Sci USA* 89:4285-9; Presta et al., 1997, *Cancer Res.* 57(20):4593-9; Gorman et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4181-4185; O'Connor et al., 1998, *Protein Eng* 11:321-8, all entirely incorporated by reference. Humanization or other methods of reducing the immunogenicity of nonhuman antibody variable regions may include resurfacing methods, as described for example in Roguska et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:969-973, entirely incorporated by reference. In one embodiment, the parent antibody has been affinity matured, as is known in the art. Structure-based methods may be employed for humanization and affinity maturation, for example as described in U.S. Ser. No. 11/004,590. Selection based methods may be employed to humanize and/or affinity mature antibody variable regions, including but not limited to methods described in Wu et al., 1999, *J. Mol. Biol.* 294:151-162; Baca et al., 1997, *J. Biol. Chem.* 272(16): 10678-10684; Rosok et al., 1996, *J. Biol. Chem.* 271(37): 22611-22618; Rader et al., 1998, *Proc. Natl. Acad. Sci. USA* 95: 8910-8915; Krauss et al., 2003, *Protein Engineering* 16(10):753-759, all entirely incorporated by reference. Other humanization methods may involve the grafting of only parts of the CDRs, including but not limited to methods described in U.S. Ser. No. 09/810,510; Tan et al., 2002, *J. Immunol.* 169:1119-1125; De Pascalis et al., 2002, *J. Immunol.* 169:3076-3084, all entirely incorporated by reference.

[0067] In one embodiment, the antibodies of the invention can be multispecific antibodies, and notably bispecific antibodies, also sometimes referred to as "diabodies". These are antibodies that bind to two (or more) different antigens, or

different epitopes on the same antigen. Diabodies can be manufactured in a variety of ways known in the art (Holliger and Winter, 1993, *Current Opinion Biotechnol.* 4:446-449, entirely incorporated by reference), e.g., prepared chemically or from hybrid hybridomas. In some cases, multispecific (for example bispecific) antibodies are not preferred.

[0068] In one embodiment, the antibody is a minibody. Minibodies are minimized antibody-like proteins comprising a scFv joined to a CH3 domain. Hu et al., 1996, *Cancer Res.* 56:3055-3061, entirely incorporated by reference. In the present instance, the CH3 domain can be engineered to improve binding to FcRn and/or increase in vivo serum half-life. In some cases, the scFv can be joined to the Fc region, and may include some or the entire hinge region.

[0069] The antibodies of the present invention are generally isolated or recombinant. "Isolated," when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a cell or cell culture from which it was expressed. Ordinarily, an isolated polypeptide will be prepared by at least one purification step. An "isolated antibody," refers to an antibody which is substantially free of other antibodies having different antigenic specificities.

[0070] "Specific binding" or "specifically binds to" or is "specific for" a particular antigen or an epitope means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target.

[0071] Specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KD for an antigen or epitope of at least about 10^{-4} M, at least about 10^{-5} M, at least about 10^{-6} M, at least about 10^{-7} M, at least about 10^{-8} M, at least about 10^{-9} M, alternatively at least about 10^{-10} M, at least about 10^{-11} M, at least about 10^{-12} M, or greater, where KD refers to a dissociation rate of a particular antibody-antigen interaction. Typically, an antibody that specifically binds an antigen will have a KD that, is 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for a control molecule relative to the antigen or epitope.

[0072] Also, specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KA or Ka for an antigen or epitope of at least 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for the epitope relative to a control, where KA or Ka refers to an association rate of a particular antibody-antigen interaction.

C. Fc Variants

[0073] The present invention relates to the generation of Fc variants of antibodies. As discussed above, by "Fc" or "Fc region" or "Fc domain" as used herein is meant the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, the Fc domain comprises immunoglobulin domains C γ 2 and C γ 3 (C γ 2 and C γ 3) and the lower hinge region between C γ 1 (C γ 1) and C γ 2 (C γ 2). Although the boundaries of the Fc region may

vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. In some embodiments, as is more fully described below, amino acid modifications are made to the Fc region, for example to alter binding to one or more FcγR receptors or to the FcRn receptor.

[0074] The present invention relates to the generation of Fc variants of antibodies to form “Fc variant antibodies”. Fc variants are made by introducing amino acid mutations into the parent molecule. “Mutations” in this context are usually amino acid substitutions, although as shown herein, deletions and insertions of amino acids can also be done and thus are defined as mutations.

[0075] The Fc variant antibodies of the invention show increased binding to FcRn and/or increased in vivo serum half-life. By “FcRn” or “neonatal Fc Receptor” as used herein is meant a protein that binds the IgG antibody Fc region and is encoded at least in part by an FcRn gene. The FcRn may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. As is known in the art, the functional FcRn protein comprises two polypeptides, often referred to as the heavy chain and light chain. The light chain is beta-2-microglobulin and the heavy chain is encoded by the FcRn gene. Unless otherwise noted herein, FcRn or an FcRn protein refers to the complex of FcRn heavy chain with beta-2-microglobulin. In some cases, the FcRn variants bind to the human FcRn receptor, or it may be desirable to design variants that bind to rodent or primate receptors in addition, to facilitate clinical trials.

[0076] A variety of such substitutions are known and described in U.S. Ser. Nos. 12/341,769; 10/672,280; 10/822,231; 11/124,620; 11/174,287; 11/396,495; 11/538,406; 11/538,411; and 12/020,443, each of which is incorporated herein by reference in its entirety for all purposes and in particular for all outlined substitutions.

[0077] In some embodiments, an Fc variant antibody can be engineered to include any of the following substitutions, alone or in any combination: 436I, 436V, 311I, 311V, 428L, 434S, 428L/434S, 259I, 308F, 259I/308F, 259I/308F/428L, 307Q/434S, 434A, 434H, 250Q/428L, M252Y/S254T/T256E, 307Q/434A, 307Q//380A/434A, and 308P/434A, 311I/434S, 311V/434S, 434S/436I, 434S/436V. Numbering is EU as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0078] In further embodiments, an Fc variant antibody can be engineered to include any of the following substitutions, alone or in any combination: 248Y, 249G, 253H, 253N, 253L, 253T, 253V, 253Q, 253M, 253Y, 253F, 253W, 255H, 255I, 255Q, 255E, 255F, 255L, 255S, 255G, 255W, 255P, 284D, 284E, 285D, 285E, 286Q, 286F, 286D, 286E, 286G, 286L, 286P, 286V, 286W, 286I, 286V, 286R, 286K, 286H, 288N, 288H, 288Q, 288Y, 288F, 288W, 288L, 288I, 307M, 307E, 307R, 307D, 307G, 307I, 307N, 307P, 307Q, 307S, 307V, 307Y, 307K, 307H, 307L, 307F, 307W, 309W, 309R, 309K, 309D, 309E, 309F, 309H, 309I, 309P, 309Q, 309L, 309Y, 309M, 312E, 312R, 312K, 312Y, 338R, 338H, 378S, 378V, 378G, 426W, 426H, 426L, 426V, 426Y, 426M, 426I, 426F, 426T, 428I, 428V, 433G, 436F, 438W, 438I, 438N, 438S, 438G, 438Y, 438F, 438I, 438D, 436I, 436V, and 436W,

where the numbering is according to the EU Index as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0079] In some embodiments, the Fc variant of the invention includes at least two modifications, wherein one of said modifications is the substitution N434S, and the other of said modifications is a substitution selected from one or more of the following: 248Y, 249G, 253H, 253N, 253L, 253T, 253V, 253Q, 253M, 253Y, 253F, 253W, 255H, 255I, 255Q, 255E, 255F, 255L, 255S, 255G, 255W, 255P, 284D, 284E, 285D, 285E, 286Q, 286F, 286D, 286E, 286G, 286L, 286P, 286Y, 286W, 286I, 286V, 286R, 286K, 286H, 288N, 288H, 288Q, 288Y, 288F, 288W, 288L, 288I, 307M, 307E, 307R, 307D, 307G, 307I, 307N, 307P, 307Q, 307S, 307V, 307Y, 307K, 307H, 307L, 307F, 307W, 309W, 309R, 309K, 309D, 309E, 309F, 309H, 309I, 309P, 309Q, 309L, 309Y, 309M, 312E, 312R, 312K, 312Y, 338R, 338H, 378S, 378V, 378G, 426W, 426H, 426L, 426V, 426Y, 426M, 426I, 426F, 426T, 428I, 428V, 433G, 436F, 438W, 438H, 438N, 438S, 438G, 438Y, 438F, 438I, 438D, 436I, 436V, and 436W, where the numbering is according to the EU Index as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0080] In some embodiments, an Fc variant antibody can be engineered to include any of the following substitutions, alone or in any combination: 378T, 378E, 378I, 378L, 378M, 426G, 426A, 426Q, 426P, 426K, and 426R, where the numbering is according to the EU Index as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0081] In some embodiments, an Fc variant antibody can be engineered to include any of the substitutions in any of the following positions alone or in any combination: 307, 311, 378, 426, 428, 434, and 436. In further embodiments, the substitutions are as follows, again alone or in any combination: 307Q, 308P, 311I, 311V, 378E, 378F, 378I, 378L, 378M, 378R, 378T, 378V, 378Y, 426A, 426G, 426K, 426L, 426P, 426Q, 426R, 426V, 428L, 434S, 436I, and 436V. In still further embodiments, the Fc variant antibody is engineered to include any of the of the following substitutions, alone or in any combination of the following or with any other of the substitutions discussed herein: T307Q/N434S/Q311I, T307Q/N434S/Q311V, T307Q/N434S/A378V, T307Q/N434S/S426V, T307Q/N434S/Y436I, T307Q/N434S/Y436V, Q311I/N434S/A378V, Q311I/N434S/A378T, Q311I/N434S/A378I, Q311I/N434S/A378L, Q311I/N434S/S426V, Q311I/N434S/Y436L, Q311I/N434S/Y436V, Q311V/S426V, Q311I/Y436I, Q311I/Y436V, Q311I/S426V/Y436I, Q311V/S426V/Y436V, Q311V/N434S/A378V, Q311V/N434S/S426V, Q311V/N434S/Y436I, Q311V/N434S/Y436V, S426V/N434S/A378V, S426V/N434S/A378T, S426V/N434S/A378I, S426V/N434S/A378L, S426V/N434S/Y436I, S426V/N434S/Y436V, N434S/Y436I/A378V, N434S/Y436I/A378T, N434S/Y436I/A378I, N434S/Y436I/A378L, N434S/Y436V/A378V, N434S/

Y436V/A378T, N434S/Y436V/A378I, N434S/Y436V/A378L, N434S/Y436V/S426G, N434S/Y436V/S426A, N434S/Y436V/S426Q, N434S/Y436V/S426P, N434S/Y436V/S426L, M428L/N434S/T307Q, M428L/N434S/Q311I, M428L/N434S/Q311V, M428L/N434S/A378V, M428L/N434S/S426V, M428L/N434S/Y436I, M428L/N434S/Y436V, V308P/N434S, A378I/N434S, A378I/N434S, A378L/N434S, A378E/N434S, A378M/N434S, A378F/N434S, A378Y/N434S, A378R/N434S, S426G/N434S, S426A/N434S, S426P/N434S, S426Q/N434S, S426K/N434S, S426R/N434S, T307Q/Q311I, and T307Q/Q311V, where the numbering is according to the EU Index as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0082] In some embodiments, an Fc variant antibody can be engineered to include at least two of the following substitutions, alone or in any combination: 248Y, 249G, 253H, 253N, 253L, 253T, 253V, 253Q, 253M, 253Y, 253F, 253W, 255H, 255I, 255Q, 255E, 255F, 255L, 255S, 255G, 255W, 255P, 284D, 284E, 285D, 285E, 286Q, 286F, 286D, 286E, 286G, 286L, 286P, 286Y, 286W, 286I, 286V, 286R, 286K, 286H, 288N, 288H, 288Q, 288Y, 288F, 288W, 288L, 288I, 307M, 307E, 307R, 307D, 307G, 307I, 307N, 307P, 307Q, 307S, 307V, 307Y, 307K, 307H, 307L, 307F, 307W, 309W, 309R, 309K, 309D, 309E, 309F, 309H, 309I, 309P, 309Q, 309L, 309Y, 309M, 312E, 312R, 312K, 312Y, 338R, 338H, 378S, 378V, 378G, 426W, 426H, 426L, 426V, 426Y, 426M, 426I, 426F, 426T, 428I, 428V, 433G, 436F, 438W, 438H, 438N, 438S, 438G, 438Y, 438F, 438I, 438D, 436I, 436V, and 436W, where the numbering is according to the EU Index as in Kabat, and it is understood that the substitution is non-native to the starting molecule. As has been shown previously, these FcRn substitutions work in IgG1, IgG2 and IgG1/G2 hybrid backbones, and are specifically included for IgG3 and IgG4 backbones and derivatives of any IgG isoform as well.

[0083] In other embodiments, no Fc variants are made in the variable region(s) of the antibodies. This is to be distinguished from affinity maturation substitutions in the variable region(s) that are made to increase binding affinity of the antibody to its antigen. That is, an Fc variant in the variable region(s) is generally significantly “silent” with respect to binding affinity.

III. Other Amino Acid Substitutions

[0084] As will be appreciated by those in the art, the Fc variant antibodies of the invention can contain additional amino acid substitutions in addition to the substitutions discussed above.

[0085] In some embodiments, amino acid substitutions are imported from one isotype into the Fc variant antibody.

[0086] In the hinge region (positions 233-236), changes can be made to increase effector function. That is, IgG2 has lowered effector function, and as a result, amino acid substitutions at these positions from PVA(deletion) can be changed to ELLG, and an additional G327A variant generated as well.

[0087] In the CH3 region, a mutation at position 384 can be made, for example substituting a non-native serine.

[0088] Additional mutations that can be made include adding either N-terminal or C-terminal (depending on the structure of the antibody or fusion protein) “tails” or sequences of one or more Fc amino acids; for example, glutamic acids and

aspartic acids can be added to the CH3 O-terminus; generally, from 1 to 5 amino acids are added.

Properties of the Fc Variant Antibodies of the Invention

[0089] The Fc variant antibodies of the invention display increased binding to FcRn and/or increased in vivo serum half-life.

[0090] In addition, some variants herein are generated to increase stability. As noted herein, a number of properties of antibodies affect the clearance rate (e.g. stability for half-life) in vivo. In addition to antibody binding to the FcRn receptor, other factors that contribute to clearance and half-life are serum aggregation, enzymatic degradation in the serum, inherent immunogenicity of the antibody leading to clearing by the immune system, antigen-mediated uptake, FcR (non-FcRn) mediated uptake and non-serum distribution (e.g. in different tissue compartments).

IV. Optional and Additional Fc Engineering

Fc Engineering

[0091] In addition to substitutions made to increase binding affinity to FcRn and/or increase serum half life, other substitutions can be made in the Fc region, in general for altering binding to FcγR receptors.

[0092] By “Fc gamma receptor”, “FcγR” or “FcγgammaR” as used herein is meant any member of the family of proteins that bind the IgG antibody Fc region and is encoded by an FcγR gene. In humans this family includes but is not limited to FcγRI (CD64), including isoforms FcγRIa, FcγRIb, and FcγRIc; FcγRII (CD32), including isoforms FcγRIIa (including allotypes H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and FcγRIIc; and FcγRIII (CD16), including isoforms FcγRIIIa (including allotypes V158 and F158) and FcγRIIIb (including allotypes FcγRIIIb-NA1 and FcγRIIIb-NA2) (Jefferis et al., 2002, *Immunol Lett* 82:57-65, entirely incorporated by reference), as well as any undiscovered human FcγRs or FcγR isoforms or allotypes. An FcγR may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. Mouse FcγRs include but are not limited to FcγRI (CD64), FcγRII (CD32), FcγRIII-1 (CD16), and FcγRIII-2 (CD16-2), as well as any undiscovered mouse FcγRs or FcγR isoforms or allotypes.

[0093] There are a number of useful Fc substitutions that can be made to alter binding to one or more of the FcγR receptors. Substitutions that result in increased binding as well as decreased binding can be useful. For example, it is known that increased binding to FcγRIIIa generally results in increased ADCC (antibody dependent cell-mediated cytotoxicity; the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause lysis of the target cell. Similarly, decreased binding to FcγRIIb (an inhibitory receptor) can be beneficial as well in some circumstances. Amino acid substitutions that find use in the present invention include those listed in U.S. Ser. No. 11/124,620 (particularly FIG. 41), U.S. Ser. Nos. 11/174,287, 11/396,495, 11/538,406, all of which are expressly incorporated herein by reference in their entirety and specifically for the variants disclosed therein. Particular variants that find use include, but are not limited to, 236A, 239D, 239E, 332E, 332D, 239D/332E, 267D, 267E, 328F, 267E/328F, 236A/332E, 239D/332E/330Y, 239D, 332E/330L and 299T.

V. Other Antibody Modifications

[0094] In addition to substitutions made to increase binding affinity to FcRn and/or increase in vivo serum half life, additional antibody modifications can be made, as described in further detail below.

Affinity Maturation

[0095] In some embodiments, one or more amino acid modifications are made in one or more of the CDRs of the antibody. In general, only 1 or 2 or 3-amino acids are substituted in any single CDR, and generally no more than from 4, 5, 6, 7, 8 or 9 or 10 changes are made within a set of CDRs. However, it should be appreciated that any combination of no substitutions, 1, 2 or 3 substitutions in any CDR can be independently and optionally combined with any other substitution.

[0096] In some cases, amino acid modifications in the CDRs are referred to as "affinity maturation". An "affinity matured" antibody is one having one or more alteration(s) in one or more CDRs which results in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In some cases, although rare, it may be desirable to decrease the affinity of an antibody to its antigen, but this is generally not preferred.

[0097] Affinity maturation can be done to increase the binding affinity of the antibody for the antigen by at least about 10% to 50-100-150% or more, or from 1 to 5 fold as compared to the "parent" antibody. Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by known procedures. See, for example, Marks et al., 1992, *Biotechnology* 10:779-783 that describes affinity maturation by variable heavy chain (VH) and variable light chain (VL) domain shuffling. Random mutagenesis of CDR and/or framework residues is described in: Barbas, et al. 1994, *Proc. Nat. Acad. Sci. USA* 91:3809-3813; Shier et al., 1995, *Gene* 169:147-155; Yelton et al., 1995, *J. Immunol.* 155:1994-2004; Jackson et al., 1995, *J. Immunol.* 154(7):3310-9; and Hawkins et al., 1992, *J. Mol. Biol.* 226:889-896, for example.

[0098] Alternatively, amino acid modifications can be made in one or more of the CDRs of the antibodies of the invention that are "silent", e.g. that do not significantly alter the affinity of the antibody for the antigen. These can be made for a number of reasons, including optimizing expression (as can be done for the nucleic acids encoding the antibodies of the invention).

[0099] Thus, included within the definition of the CDRs and antibodies of the invention are variant CDRs and antibodies; that is, the antibodies of the invention can include amino acid modifications in one or more of the CDRs of Ab79 and Ab19. In addition, as outlined below, amino acid modifications can also independently and optionally be made in any region outside the CDRs, including framework and constant regions.

ADC Modifications

[0100] In some embodiments, the Fc variant antibodies of the invention are conjugated with drugs to form antibody-drug conjugates (ADCs). In general, ADCs are used in oncology applications, where the use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents allows for the targeted delivery of the drug moiety to tumors, which

can allow higher efficacy, lower toxicity, etc. An overview of this technology is provided in Ducry et al., *Bioconjugate Chem.*, 21:5-13 (2010), Carter et al., *Cancer J.* 14(3):154 (2008) and Senter, *Current Opin. Chem. Biol.* 13:235-244 (2009), all of which are hereby incorporated by reference in their entirety.

[0101] Thus the invention provides Fc variant antibodies conjugated to drugs. Generally, conjugation is done by covalent attachment to the antibody, as further described below, and generally relies on a linker, often a peptide linkage (which, as described below, may be designed to be sensitive to cleavage by proteases at the target site or not). In addition, as described above, linkage of the linker-drug unit (LU-D) can be done by attachment to cysteines within the antibody. As will be appreciated by those in the art, the number of drug moieties per antibody can change, depending on the conditions of the reaction, and can vary from 1:1 to 10:1 drug:antibody. As will be appreciated by those in the art, the actual number is an average.

[0102] Thus the invention provides Fc variant antibodies conjugated to drugs. As described below, the drug of the ADC can be any number of agents, including but not limited to cytotoxic agents such as chemotherapeutic agents, growth inhibitory agents, toxins (for example, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (that is, a radioconjugate) are provided. In other embodiments, the invention further provides methods of using the ADCs.

[0103] Drugs for use in the present invention include cytotoxic drugs, particularly those which are used for cancer therapy. Such drugs include, in general, DNA damaging agents, anti-metabolites, natural products and their analogs. Exemplary classes of cytotoxic agents include the enzyme inhibitors such as dihydrofolate reductase inhibitors, and thymidylate synthase inhibitors, DNA intercalators, DNA cleavers, topoisomerase inhibitors, the anthracycline family of drugs, the vinca drugs, the mitomycins, the bleomycins, the cytotoxic nucleosides, the pteridine family of drugs, diynes, the podophyllotoxins, dolastatins, maytansinoids, differentiation inducers, and taxols.

[0104] Members of these classes include, for example, methotrexate, methopterin, dichloromethotrexate, 5-fluorouracil, 6-mercaptopurine, cytosine arabinoside, melphalan, leurosine, leurosideine, actinomycin, daunorubicin, doxorubicin, mitomycin C, mitomycin A, caminomycin, aminopterin, tallysomycin, podophyllotoxin and podophyllotoxin derivatives such as etoposide or etoposide phosphate, vinblastine, vincristine, vindesine, taxanes including taxol, taxotere retinoic acid, butyric acid, N8-acetyl spermidine, camptothecin, calicheamicin, esperamicin, ene-dynes, duocarmycin A, duocarmycin SA, calicheamicin, camptothecin, maytansinoids (including DM1), monomethylauristatin E (MMAE), monomethylauristatin F (MMAF), and maytansinoids (DM4) and their analogues.

[0105] Toxins may be used as antibody-toxin conjugates and include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) *J. Nat. Cancer Inst.* 92(19): 1573-1581; Mandler et al (2000) *Bioorganic & Med. Chem. Letters* 10:1025-1028; Mandler et al (2002) *Bioconjugate Chem.* 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:8618-8623), and calicheamicin (Lode et al (1998) *Cancer Res.* 58:2928; Hinman et al (1993) *Cancer Res.* 53:3336-3342). Toxins may exert

their cytotoxic and cytostatic effects by mechanisms including tubulin binding, DNA binding, or topoisomerase inhibition.

[0106] Conjugates of an Fc variant antibody and one or more small molecule toxins,

[0107] such as a maytansinoids, dolastatins, auristatins, a trichothecene, calicheamicin, and CC1065, and the derivatives of these toxins that have toxin activity, are contemplated.

[0108] Maytansinoids

[0109] Maytansine compounds suitable for use as maytansinoid drug moieties are well known in the art, and can be isolated from natural sources according to known methods, produced using genetic engineering techniques (see Yu et al (2002) PNAS 99:7968-7973), or maytansinol and maytansinol analogues prepared synthetically according to known methods. As described below, drugs may be modified by the incorporation of a functionally active group such as a thiol or amine group for conjugation to the antibody.

[0110] Exemplary maytansinoid drug moieties include those having a modified aromatic ring, such as: C-19-dechloro (U.S. Pat. No. 4,256,746) (prepared by lithium aluminum hydride reduction of ansamycin P2); C-20-hydroxy (or C-20-demethyl)+/-C-19-dechloro (U.S. Pat. Nos. 4,361,650 and 4,307,016) (prepared by demethylation using *Streptomyces* or *Actinomyces* or dechlorination using LAH); and C-20-demethoxy, C-20-acyloxy (—OCOR), +/-dechloro (U.S. Pat. No. 4,294,757) (prepared by acylation using acyl chlorides) and those having modifications at other positions

[0111] Exemplary maytansinoid drug moieties also include those having modifications such as: C-9-SH (U.S. Pat. No. 4,424,219) (prepared by the reaction of maytansinol with H₂S or P₂S₅); C-14-alkoxymethyl(demethoxy/CH₂OR) (U.S. Pat. No. 4,331,598); C-14-hydroxymethyl or acyloxymethyl (CH₂OH or CH₂OAc) (U.S. Pat. No. 4,450,254) prepared from *Nocardia*; C-15-hydroxy/acyloxy (U.S. Pat. No. 4,364,866) (prepared by the conversion of maytansinol by *Streptomyces*); C-15-methoxy (U.S. Pat. Nos. 4,313,946 and 4,315,929) (isolated from *Trewia nudiflora*); C-18-N-demethyl (U.S. Pat. Nos. 4,362,663 and 4,322,348) (prepared by the demethylation of maytansinol by *Streptomyces*); and 4,5-deoxy (U.S. Pat. No. 4,371,533) (prepared by the titanium trichloride/LAH reduction of maytansinol).

[0112] Of particular use are DM1 (disclosed in U.S. Pat. No. 5,208,020, incorporated by reference) and DM4 (disclosed in U.S. Pat. No. 7,276,497, incorporated by reference). See also a number of additional maytansinoid derivatives and methods in 5,416,064, WO/01/24763, 7,303,749, 7,601,354, U.S. Ser. No. 12/631,508, WO02/098883, 6,441,163, 7,368,565, WO02/16368 and WO04/1033272, all of which are expressly incorporated by reference in their entirety.

[0113] ADC's containing maytansinoids, methods of making same, and their therapeutic use are disclosed, for example, in U.S. Pat. Nos. 5,208,020; 5,416,064; 6,441,163 and European Patent EP 0 425 235 B1, the disclosures of which are hereby expressly incorporated by reference, Liu et al., Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described ADCs comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an in vivo tumor growth assay.

[0114] Chari et al., Cancer Research 52:127-131 (1992) describe ADCs in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an

antigen on human colon cancer cell lines, or to another murine monoclonal antibody TA. 1 that binds the HER-2/neu oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested in vitro on the human breast cancer cell line SK-BR-3, which expresses 3×10⁵ HER-2 surface antigens per cell. The drug conjugate achieved a degree of cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

[0115] Auristatins and Dolastatins

[0116] In some embodiments, the ADC comprises an Fc variant antibody conjugated to dolastatins or dolastatin peptidic analogs and derivatives, the auristatins (U.S. Pat. Nos. 5,635,483; 5,780,588). Dolastatins and auristatins have been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke et al (2001) Antimicrob. Agents and Chemother. 45(12):3580-3584) and have anticancer (U.S. Pat. No. 5,663,149) and antifungal activity (Pettit et al (1998) Antimicrob. Agents Chemother. 42:2961-2965). The dolastatin or auristatin drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl) terminus of the peptidic drug moiety (WO 02/088172).

[0117] Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, disclosed in "Senter et al, Proceedings of the American Association for Cancer Research, Volume 45, Abstract Number 623, presented Mar. 28, 2004 and described in United States Patent Publication No. 2005/0238648, the disclosure of which is expressly incorporated by reference in its entirety.

[0118] An exemplary auristatin embodiment is MMAE (shown in FIG. 10 wherein the wavy line indicates the covalent attachment to a linker (L) of an antibody drug conjugate; see U.S. Pat. No. 6,884,869 expressly incorporated by reference in its entirety).

[0119] Another exemplary auristatin embodiment is MMAF, shown in FIG. 10 wherein the wavy line indicates the covalent attachment to a linker (L) of an antibody drug conjugate (US 2005/0238649, 5,767,237 and 6,124,431, expressly incorporated by reference in their entirety).

[0120] Additional exemplary embodiments comprising MMAE or MMAF and various linker components (described further herein) have the following structures and abbreviations (wherein Ab means antibody and p is 1 to about 8):

[0121] Typically, peptide-based drug moieties can be prepared by forming a peptide bond between two or more amino acids and/or peptide fragments. Such peptide bonds can be prepared, for example, according to the liquid phase synthesis method (see E. Schroder and K. Lubke, "The Peptides", volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry. The auristatin/dolastatin drug moieties may be prepared according to the methods of: U.S. Pat. No. 5,635,483; U.S. Pat. No. 5,780,588; Pettit et al (1989) J. Am. Chem. Soc. 111:5463-5465; Pettit et al (1998) Anti-Cancer Drug Design 13:243-277; Pettit, G. R., et al Synthesis, 1996, 719-725; Pettit et al (1996) J. Chem. Soc. Perkin Trans. 1 5:859-863; and Doronina (2003) Nat Biotechnol 21(7):778-784.

[0122] Calicheamicin

[0123] In other embodiments, the ADC comprises an antibody of the invention conjugated to one or more calicheamicin molecules. For example, Mylotarg is the first commercial

ADC drug and utilizes calicheamicin γ 1 as the payload (see U.S. Pat. No. 4,970,198, incorporated by reference in its entirety). Additional calicheamicin derivatives are described in U.S. Pat. Nos. 5,264,586, 5,384,412, 5,550,246, 5,739,116, 5,773,001, 5,767,285 and 5,877,296, all expressly incorporated by reference. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Pat. Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, 5,877,296 (all to American Cyanamid Company). Structural analogues of calicheamicin which may be used include, but are not limited to, γ 1I, α 2I, α 2L, N-acetyl- γ 1I, PSAG and 011 (Hinman et al., Cancer Research 53:3336-3342 (1993), Lode et al., Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents to American Cyanamid). Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

[0124] Duocarmycins

[0125] CC-1065 (see U.S. Pat. No. 4,169,888, incorporated by reference) and duocarmycins are members of a family of antitumor antibiotics utilized in ADCs. These antibiotics appear to work through sequence-selectively alkylating DNA at the N3 of adenine in the minor groove, which initiates a cascade of events that result in apoptosis.

[0126] Important members of the duocarmycins include duocarmycin A (U.S. Pat. No. 4,923,990, incorporated by reference) and duocarmycin SA (U.S. Pat. No. 5,101,038, incorporated by reference), and a large number of analogues as described in U.S. Pat. Nos. 7,517,903, 7,691,962, 5,101,038; 5,641,780; 5,187,186; 5,070,092; 5,070,092; 5,641,780; 5,101,038; 5,084,468, 5,475,092, 5,585,499, 5,846,545, WO2007/089149, WO2009/017394A1, 5,703,080, 6,989,452, 7,087,600, 7,129,261, 7,498,302, and 7,507,420, all of which are expressly incorporated by reference.

VI. Other Cytotoxic Agents

[0127] Other antitumor agents that can be conjugated to the antibodies of the invention include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Pat. Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Pat. No. 5,877,296).

[0128] Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), *momordica charantia* inhibitor, curcin, cretin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published Oct. 28, 1993.

[0129] The present invention further contemplates an ADC formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

[0130] For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of radio-con-

jugated antibodies. Examples include At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32, Pb212 and radioactive isotopes of Lit.

[0131] The radio- or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as Tc99m or I123, Re186, Re188 and In111 can be attached via a cysteine residue in the peptide, Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker et al (1978) Biochem. Biophys. Res. Commun. 80: 49-57 can be used to incorporate Iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

[0132] For compositions comprising a plurality of antibodies, the drug loading is represented by p, the average number of drug molecules per Antibody. Drug loading may range from 1 to 20 drugs (D) per Antibody. The average number of drugs per antibody in preparation of conjugation reactions may be characterized by conventional means such as mass spectroscopy, ELISA assay, and HPLC. The quantitative distribution of Antibody-Drug-Conjugates in terms of p may also be determined.

[0133] In some instances, separation, purification, and characterization of homogeneous Antibody-Drug-conjugates where p is a certain value from Antibody-Drug-Conjugates with other drug loadings may be achieved by means such as reverse phase HPLC or electrophoresis. In exemplary embodiments, p is 2, 3, 4, 5, 6, 7, or 8 or a fraction thereof.

[0134] The generation of Antibody-drug conjugate compounds can be accomplished by any technique known to the skilled artisan. Briefly, the Antibody-drug conjugate compounds can include an Fc variant antibody as the Antibody unit, a drug, and optionally a linker that joins the drug and the binding agent.

[0135] A number of different reactions are available for covalent attachment of drugs and/or linkers to binding agents. This can be accomplished by reaction of the amino acid residues of the binding agent, for example, antibody molecule, including the amine groups of lysine, the free carboxylic acid groups of glutamic and aspartic acid, the sulfhydryl groups of cysteine and the various moieties of the aromatic amino acids. A commonly used non-specific methods of covalent attachment is the carbodiimide reaction to link a carboxy (or amino) group of a compound to amino (or carboxy) groups of the antibody. Additionally, bifunctional agents such as dialdehydes or imidoesters have been used to link the amino group of a compound to amino groups of an antibody molecule.

[0136] Also available for attachment of drugs to binding agents is the Schiff base reaction. This method involves the periodate oxidation of a drug that contains glycol or hydroxy groups, thus forming an aldehyde which is then reacted with the binding agent. Attachment occurs via formation of a Schiff base with amino groups of the binding agent. Isothiocyanates can also be used as coupling agents for covalently attaching drugs to binding agents. Other techniques are known to the skilled artisan and within the scope of the present invention.

[0137] In some embodiments, an intermediate, which is the precursor of the linker, is reacted with the drug under appropriate conditions. In other embodiments, reactive groups are used on the drug and/or the intermediate. The product of the

reaction between the drug and the intermediate, or the derivatized drug, is subsequently reacted with an Fc variant antibody of the invention under appropriate conditions.

[0138] It will be understood that chemical modifications may also be made to the desired compound in order to make reactions of that compound more convenient for purposes of preparing conjugates of the invention. For example a functional group e.g. amine, hydroxyl, or sulfhydryl, may be appended to the drug at a position which has minimal or an acceptable effect on the activity or other properties of the drug

VII. Linker Units

[0139] Typically, the antibody-drug conjugate compounds comprise a Linker unit between the drug unit and the antibody unit. In some embodiments, the linker is cleavable under intracellular or extracellular conditions, such that cleavage of the linker releases the drug unit from the antibody in the appropriate environment. For example, solid tumors that secrete certain proteases may serve as the target of the cleavable linker; in other embodiments, it is the intracellular proteases that are utilized. In yet other embodiments, the linker unit is not cleavable and the drug is released, for example, by antibody degradation in lysosomes.

[0140] In some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (for example, within a lysosome or endosome or caveola). The linker can be, for example, a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. In some embodiments, the peptidyl linker is at least two amino acids long or at least three amino acids long or more.

[0141] Cleaving agents can include, without limitation, cathepsins B and D and plasmin, all of which are known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells (see, e.g., Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123). Peptidyl linkers that are cleavable by enzymes that are present in CD38-expressing cells. For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a Phe-Leu or a Gly-Phe-Leu-Gly linker (SEQ ID NO: 11). Other examples of such linkers are described, e.g., in U.S. Pat. No. 6,214,345, incorporated herein by reference in its entirety and for all purposes.

[0142] In some embodiments, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker or a Phe-Lys linker (see, e.g., U.S. Pat. No. 6,214,345, which describes the synthesis of doxorubicin with the val-cit linker).

[0143] In other embodiments, the cleavable linker is pH-sensitive, that is, sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acid-labile linker that is hydrolyzable in the lysosome (for example, a hydrazone, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like) may be used. (See, e.g., U.S. Pat. Nos. 5,122,368; 5,824,805; 5,622,929; Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123; Neville et al., 1989, *Biol. Chem.* 264:14653-14661.) Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome. In certain embodiments, the hydrolyzable linker is a thioether linker (such as, e.g., a thioether attached to the therapeutic agent via an acylhydrazone bond (see, e.g., U.S. Pat. No. 5,622,929).

[0144] In yet other embodiments, the linker is cleavable under reducing conditions (for example, a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-5-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), SPDB (N-succinimidyl-3-(2-pyridyldithio)butyrate) and SMPT (N-succinimidyl-oxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio) toluene)-, SPDB and SMPT. (See, e.g., Thorpe et al., 1987, *Cancer Res.* 47:5924-5931; Wawrzynczak et al., In *Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer* (C. W. Vogel ed., Oxford U. Press, 1987. See also U.S. Pat. No. 4,880,935.)

[0145] In other embodiments, the linker is a malonate linker (Johnson et al., 1995, *Anticancer Res.* 15:1387-93), a maleimidobenzoyl linker (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10): 1299-1304), or a 3'-N-amide analog (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10):1305-12).

[0146] In yet other embodiments, the linker unit is not cleavable and the drug is released by antibody degradation. (See U.S. Publication No. 2005/0238649 incorporated by reference herein in its entirety and for all purposes).

[0147] In many embodiments, the linker is self-immolative. As used herein, the term "self-immolative Spacer" refers to a bifunctional chemical moiety that is capable of covalently linking together two spaced chemical moieties into a stable tripartite molecule. It will spontaneously separate from the second chemical moiety if its bond to the first moiety is cleaved. See for example, WO 2007/059404A2, WO06110476A2, WO05112919A2, WO2010/062171, WO09/017,394, WO07/089,149, WO 07/018,431, WO04/043493 and WO02/083180, which are directed to drug-cleavable substrate conjugates where the drug and cleavable substrate are optionally linked through a self-immolative linker and which are all expressly incorporated by reference.

[0148] Often the linker is not substantially sensitive to the extracellular environment. As used herein, "not substantially sensitive to the extracellular environment," in the context of a linker, means that no more than about 20%, 15%, 10%, 5%, 3%, or no more than about 1% of the linkers, in a sample of antibody-drug conjugate compound, are cleaved when the antibody-drug conjugate compound presents in an extracellular environment (for example, in plasma).

[0149] Whether a linker is not substantially sensitive to the extracellular environment can be determined, for example, by incubating with plasma the antibody-drug conjugate compound for a predetermined time period (for example, 2, 4, 8, 16, or 24 hours) and then quantitating the amount of free drug present in the plasma.

[0150] In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (that is, in the milieu of the linker-therapeutic agent moiety of the antibody-drug conjugate compound as described herein). In yet other embodiments, the linker promotes cellular internalization when conjugated to both the auristatin compound and the Fc variant antibodies of the invention.

[0151] A variety of exemplary linkers that can be used with the present compositions and methods are described in WO 2004-010957, U.S. Publication No. 2006/0074008, U.S. Publication No. 20050238649, and U.S. Publication No. 2006/0024317 (each of which is incorporated by reference herein in its entirety and for all purposes).

VIII. Drug Loading

[0152] Drug loading is represented by *p* and is the average number of Drug moieties per antibody in a molecule. Drug loading ("p") may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more moieties (*D*) per antibody, although frequently the average number is a fraction or a decimal. Generally, drug loading of from 1 to 4 is frequently useful, and from 1 to 2 is also useful. ADCs of the invention include collections of antibodies conjugated with a range of drug moieties, from 1 to 20. The average number of drug moieties per antibody in preparations of ADC from conjugation reactions may be characterized by conventional means such as mass spectroscopy and, ELISA assay.

[0153] The quantitative distribution of ADC in terms of *p* may also be determined. In some instances, separation, purification, and characterization of homogeneous ADC where *p* is a certain value from ADC with other drug loadings may be achieved by means such as electrophoresis.

[0154] For some antibody-drug conjugates, *p* may be limited by the number of attachment sites on the antibody. For example, where the attachment is a cysteine thiol, as in the exemplary embodiments above, an antibody may have only one or several cysteine thiol groups, or may have only one or several sufficiently reactive thiol groups through which a linker may be attached. In certain embodiments, higher drug loading, e.g. $p > 5$, may cause aggregation, insolubility, toxicity, or loss of cellular permeability of certain antibody-drug conjugates. In certain embodiments, the drug loading for an ADC of the invention ranges from 1 to about 8; from about 2 to about 6; from about 3 to about 5; from about 3 to about 4; from about 3.1 to about 3.9; from about 3.2 to about 3.8; from about 3.2 to about 3.7; from about 3.2 to about 3.6; from about 3.3 to about 3.8; or from about 3.3 to about 3.7. Indeed, it has been shown that for certain ADCs, the optimal ratio of drug moieties per antibody may be less than 8, and may be about 2 to about 5. See US 2005-0238649 A1 (herein incorporated by reference in its entirety).

[0155] In certain embodiments, fewer than the theoretical maximum of drug moieties are conjugated to an antibody during a conjugation reaction. An antibody may contain, for example, lysine residues that do not react with the drug-linker intermediate or linker reagent, as discussed below. Generally, antibodies do not contain many free and reactive cysteine thiol groups which may be linked to a drug moiety; indeed most cysteine thiol residues in antibodies exist as disulfide bridges. In certain embodiments, an antibody may be reduced with a reducing agent such as dithiothreitol (DTT) or tricarboxylethylphosphine (TCEP), under partial or total reducing conditions, to generate reactive cysteine thiol groups. In certain embodiments, an antibody is subjected to denaturing conditions to reveal reactive nucleophilic groups such as lysine or cysteine.

[0156] The loading (drug/antibody ratio) of an ADC may be controlled in different ways, e.g., by: (i) limiting the molar excess of drug-linker intermediate or linker reagent relative to antibody, (ii) limiting the conjugation reaction time or temperature, (iii) partial or limiting reductive conditions for cysteine thiol modification, (iv) engineering by recombinant techniques the amino acid sequence of the antibody such that the number and position of cysteine residues is modified for control of the number and/or position of linker-drug attachments (such as thioMab or thioFab prepared as disclosed herein and in WO2006/034488 (herein incorporated by reference in its entirety)).

[0157] It is to be understood that where more than one nucleophilic group reacts with a drug-linker intermediate or linker reagent followed by drug moiety reagent, then the resulting product is a mixture of ADC compounds with a distribution of one or more drug moieties attached to an antibody. The average number of drugs per antibody may be calculated from the mixture by a dual ELISA antibody assay, which is specific for antibody and specific for the drug. Individual ADC molecules may be identified in the mixture by mass spectroscopy and separated by HPLC, e.g. hydrophobic interaction chromatography.

[0158] In some embodiments, a homogeneous ADC with a single loading value may be isolated from the conjugation mixture by electrophoresis or chromatography.

Methods of Determining Cytotoxic Effect of ADCs

[0159] Methods of determining whether a Drug or Antibody-Drug conjugate exerts a cytostatic and/or cytotoxic effect on a cell are known. Generally, the cytotoxic or cytostatic activity of an Antibody Drug conjugate can be measured by: exposing mammalian cells expressing a target protein of the Antibody Drug conjugate in a cell culture medium; culturing the cells for a period from about 6 hours to about 5 days; and measuring cell viability. Cell-based in vitro assays can be used to measure viability (proliferation), cytotoxicity, and induction of apoptosis (caspase activation) of the Antibody Drug conjugate.

[0160] For determining whether an Antibody Drug conjugate exerts a cytostatic effect, a thymidine incorporation assay may be used. For example, cancer cells expressing a target antigen at a density of 5,000 cells/well of a 96-well plated can be cultured for a 72-hour period and exposed to 0.5 μ Ci of 3 H-thymidine during the final 8 hours of the 72-hour period. The incorporation of 3 H-thymidine into cells of the culture is measured in the presence and absence of the Antibody Drug conjugate.

[0161] For determining cytotoxicity, necrosis or apoptosis (programmed cell death) can be measured. Necrosis is typically accompanied by increased permeability of the plasma membrane; swelling of the cell, and rupture of the plasma membrane. Apoptosis is typically characterized by membrane blebbing, condensation of cytoplasm, and the activation of endogenous endonucleases. Determination of any of these effects on cancer cells indicates that an Antibody Drug conjugate is useful in the treatment of cancers.

[0162] Cell viability can be measured by determining in a cell the uptake of a dye such as neutral red, trypan blue, or ALAMAR™ blue (see, e.g., Page et al., 1993, Intl. J. Oncology 3:473-476). In such an assay, the cells are incubated in media containing the dye, the cells are washed, and the remaining dye, reflecting cellular uptake of the dye, is measured spectrophotometrically. The protein-binding dye sulforhodamine B (SRB) can also be used to measure cytotoxicity (Skehan et al., 1990, J. Natl Cancer Inst. 82:1107-12).

[0163] Alternatively, a tetrazolium salt, such as MTT, is used in a quantitative colorimetric assay for mammalian cell survival and proliferation by detecting living, but not dead, cells (see, e.g., Mosmann, 1983, J. Immunol. Methods 65:55-63).

[0164] Apoptosis can be quantitated by measuring, for example, DNA fragmentation. Commercial photometric methods for the quantitative in vitro determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucle-

otides in fragmented DNA) and ELISA-based assays, are described in Biochemica, 1999, no. 2, pp. 34-37 (Roche Molecular Biochemicals).

[0165] Apoptosis can also be determined by measuring morphological changes in a cell. For example, as with necrosis, loss of plasma membrane integrity can be determined by measuring uptake of certain dyes (e.g., a fluorescent dye such as, for example, acridine orange or ethidium bromide). A method for measuring apoptotic cell number has been described by Duke and Cohen, *Current Protocols in Immunology* (Coligan et al. eds., 1992, pp. 3.17.1-3.17.16). Cells also can be labeled with a DNA dye (e.g., acridine orange, ethidium bromide, or propidium iodide) and the cells observed for chromatin condensation and margination along the inner nuclear membrane. Other morphological changes that can be measured to determine apoptosis include, e.g., cytoplasmic condensation, increased membrane blebbing, and cellular shrinkage.

[0166] The presence of apoptotic cells can be measured in both the attached and "floating" compartments of the cultures. For example, both compartments can be collected by removing the supernatant, trypsinizing the attached cells, combining the preparations following a centrifugation wash step (e.g., 10 minutes at 2000 rpm), and detecting apoptosis (e.g., by measuring DNA fragmentation). (See, e.g., Piazza et al., 1995, *Cancer Research* 55:3110-16).

[0167] In vivo, the effect of a therapeutic composition of the Fc variant antibody of the invention can be evaluated in a suitable animal model. For example, xenogenic cancer models can be used, wherein cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice (Klein et al., 1997, *Nature Medicine* 3: 402-408). Efficacy can be measured using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like.

[0168] The therapeutic compositions used in the practice of the foregoing methods can be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is generally non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, Remington's *Pharmaceutical Sciences* 16th Edition, A. Osal., Ed., 1980).

[0169] Glycosylation

[0170] Another type of covalent modification is alterations in glycosylation. In another embodiment, the antibodies disclosed herein can be modified to include one or more engineered glycoforms. By "engineered glycoform" as used herein is meant a carbohydrate composition that is covalently attached to the antibody, wherein said carbohydrate composition differs chemically from that of a parent antibody. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing effector function. A preferred form of engineered glycoform is afucosylation, which has been shown to be correlated to an increase in ADCC function, presumably through tighter binding to the FcγRIIIa receptor. In this context, "afucosylation" means that the majority of the antibody produced in the host cells is substantially devoid of fucose, e.g. 90-95-98% of the generated antibodies do not have appreciable fucose as a compo-

nent of the carbohydrate moiety of the antibody (generally attached at N297 in the Fc region). Defined functionally, afucosylated antibodies generally exhibit at least a 50% or higher affinity to the FcγRIIIa receptor.

[0171] Engineered glycoforms may be generated by a variety of methods known in the art (Umaña et al., 1999, *Nat Biotechnol* 17:176-180; Davies et al., 2001, *Biotechnol Bioeng* 74:288-294; Shields et al., 2002, *J Biol Chem* 277: 26733-26740; Shinkawa et al., 2003, *J Biol Chem* 278:3466-3473; U.S. Pat. No. 6,602,684; U.S. Ser. No. 10/277,370; U.S. Ser. No. 10/113,929; PCT WO 00/61739A1; PCT WO 01/29246A1; PCT WO 02/31140A1; PCT WO 02/30954A1, all entirely incorporated by reference; (Potelligent® technology [Biowa, Inc., Princeton, N.J.]; GlycoMAb® glycosylation engineering technology [Glycart Biotechnology AG, Zürich, Switzerland]). Many of these techniques are based on controlling the level of fucosylated and/or bisecting oligosaccharides that are covalently attached to the Fc region, for example by expressing an IgG in various organisms or cell lines, engineered or otherwise (for example Lee-13 CHO cells or rat hybridoma YB2/O cells, by regulating enzymes involved in the glycosylation pathway (for example FUT8 [α1,6-fucosyltransferase] and/or β1-4-N-acetylglucosaminyltransferase III [GnTIII]), or by modifying carbohydrate(s) after the IgG has been expressed. For example, the "sugar engineered antibody" or "SEA technology" of Seattle Genetics functions by adding modified saccharides that inhibit fucosylation during production; see for example 20090317869, hereby incorporated by reference in its entirety. Engineered glycoform typically refers to the different carbohydrate or oligosaccharide; thus an antibody can include an engineered glycoform.

[0172] Alternatively, engineered glycoform may refer to the IgG variant that comprises the different carbohydrate or oligosaccharide. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

[0173] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0174] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antibody amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating

the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0175] Another means of increasing the number of carbohydrate moieties on the antibody is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxy-proline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330 and in Aplin and Wriston, 1981, CRC Crit. Rev. Biochem., pp. 259-306, both entirely incorporated by reference.

[0176] Removal of carbohydrate moieties present on the starting antibody (e.g. post-translationally) may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, Arch. Biochem. Biophys. 259:52 and by Edge et al., 1981, Anal. Biochem. 118:131, both entirely incorporated by reference. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, Meth. Enzymol. 138:350, entirely incorporated by reference. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, J. Biol. Chem. 257:3105, entirely incorporated by reference. Tunicamycin blocks the formation of protein-N-glycoside linkages.

[0177] Another type of covalent modification of the antibody comprises linking the antibody to various nonproteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in, for example, 2005-2006 PEG Catalog from Nektar Therapeutics (available at the Nektar website) U.S. Pat. No. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337, all entirely incorporated by reference. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody to facilitate the addition of polymers such as PEG. See for example, U.S. Publication No. 2005/0114037A1, entirely incorporated by reference.

IX. Nucleic Acids and Host Cells

[0178] Included within the invention are the nucleic acids encoding the Fc variant antibodies of the invention. In the case where both a heavy and light chain constant domains are included in the Fc variant antibody, generally these are made using nucleic acids encoding each, that are combined into standard host cells (e.g. CHO cells, etc.) to produce the tetrameric structure of the antibody. If only one Fc variant engineered constant domain is being made, only a single nucleic acid will be used.

X. Antibody Compositions for In Vivo Administration

[0179] The use of the Fc variant antibodies of the invention in therapy will depend on the antigen binding component; e.g. in the case of full length standard therapeutic antibodies, on the antigen to which the antibody's Fv binds. That is, as will be appreciated by those in the art, the treatment of specific diseases can be done with the additional benefit of increased half life of the molecule. This can result in a variety of benefits, including, but not limited to, less frequent dosing (which can lead to better patient compliance), lower dosing, and lower production costs.

[0180] Formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

[0181] The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to provide antibodies with other specificities. Alternatively, or in addition, the composition may comprise a cytotoxic agent, cytokine, growth inhibitory agent and/or small molecule antagonist. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0182] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

[0183] The formulations to be used for in vivo administration should be sterile, or nearly so. This is readily accomplished by filtration through sterile filtration membranes.

[0184] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include

semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ -ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

[0185] When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

XI. Administrative Modalities

[0186] The antibodies and chemotherapeutic agents of the invention are administered to a subject, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous or subcutaneous administration of the antibody is preferred,

XII. Treatment Modalities

[0187] In the methods of the invention, therapy is used to provide a positive therapeutic response with respect to a disease or condition. By “positive therapeutic response” is intended an improvement in the disease or condition, and/or an improvement in the symptoms associated with the disease or condition. For example, a positive therapeutic response would refer to one or more of the following improvements in the disease: (1) a reduction in the number of neoplastic cells; (2) an increase in neoplastic cell death; (3) inhibition of neoplastic cell survival; (5) inhibition (i.e., slowing to some extent, preferably halting) of tumor growth; (6) an increased patient survival rate; and (7) some relief from one or more symptoms associated with the disease or condition.

[0188] Positive therapeutic responses in any given disease or condition can be determined by standardized response criteria specific to that disease or condition. Tumor response can be assessed for changes in tumor morphology (i.e., overall tumor burden, tumor size, and the like) using screening techniques such as magnetic resonance imaging (MRI) scan, x-radiographic imaging, computed tomographic (CT) scan, bone scan imaging, endoscopy, and tumor biopsy sampling including bone marrow aspiration (BMA) and counting of tumor cells in the circulation.

[0189] In addition to these positive therapeutic responses, the subject undergoing therapy may experience the beneficial effect of an improvement in the symptoms associated with the disease.

[0190] Thus for B cell tumors, the subject may experience a decrease in the so-called B symptoms, i.e., night sweats, fever, weight loss, and/or urticaria. For pre-malignant conditions, therapy with an Fc variant therapeutic agent may block and/or prolong the time before development of a related malignant condition, for example, development of multiple myeloma in subjects suffering from monoclonal gammopathy of undetermined significance (MGUS).

[0191] An improvement in the disease may be characterized as a complete response. By “complete response” is intended an absence of clinically detectable disease with normalization of any previously abnormal radiographic studies, bone marrow, and cerebrospinal fluid (CSF) or abnormal monoclonal protein in the case of myeloma.

[0192] Such a response may persist for at least 4 to 8 weeks, or sometimes 6 to 8 weeks, following treatment according to the methods of the invention. Alternatively, an improvement in the disease may be categorized as being a partial response. By “partial response” is intended at least about a 50% decrease in all measurable tumor burden (i.e., the number of malignant cells present in the subject, or the measured bulk of tumor masses or the quantity of abnormal monoclonal protein) in the absence of new lesions, which may persist for 4 to 8 weeks, or 6 to 8 weeks.

[0193] Treatment according to the present invention includes a “therapeutically effective amount” of the medicaments used. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result.

[0194] A therapeutically effective amount may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the medicaments to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

[0195] A “therapeutically effective amount” for tumor therapy may also be measured by its ability to stabilize the progression of disease. The ability of a compound to inhibit cancer may be evaluated in an animal model system predictive of efficacy in human tumors.

[0196] Alternatively, this property of a composition may be evaluated by examining the ability of the compound to inhibit cell growth or to induce apoptosis by in vitro assays known to the skilled practitioner. A therapeutically effective amount of a therapeutic compound may decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

[0197] Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary

dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0198] The specification for the dosage unit forms of the present invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[0199] The efficient dosages and the dosage regimens for the Fc variant antibodies used in the present invention depend on the disease or condition to be treated and may be determined by the persons skilled in the art.

[0200] An exemplary, non-limiting range for a therapeutically effective amount of an Fc variant antibody used in the present invention is about 0, 1-100 mg/kg, such as about 0.1-50 mg/kg, for example about 0.1-20 mg/kg, such as about 0.1-10 mg/kg, for instance about 0.5, about such as 0.3, about 1, or about 3 mg/kg. In another embodiment, the antibody is administered in a dose of 1 mg/kg or more, such as a dose of from 1 to 20 mg/kg, e.g. a dose of from 5 to 20 mg/kg, e.g. a dose of 8 mg/kg.

[0201] A medical professional having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, a physician or a veterinarian could start doses of the medication employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

[0202] In one embodiment, the Fc variant antibody is administered by infusion in a weekly dosage of from 10 to 500 mg/kg such as of from 200 to 400 mg/kg. Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours.

[0203] In one embodiment, the Fc variant antibody is administered by slow continuous infusion over a long period, such as more than 24 hours, if required to reduce side effects including toxicity.

[0204] In one embodiment the Fc variant antibody is administered in a weekly dosage of from 250 mg to 2000 mg, such as for example 300 mg, 500 mg, 700 mg, 1000 mg, 1500 mg or 2000 mg, for up to 8 times, such as from 4 to 6 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours. Such regimen may be repeated one or more times as necessary, for example, after 6 months or 12 months. The dosage may be determined or adjusted by measuring the amount of compound of the present invention in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the antigen binding region of the Fc variant antibody.

[0205] In a further embodiment, the Fc variant antibody is administered once weekly for 2 to 12 weeks, such as for 3 to 10 weeks, such as for 4 to 8 weeks.

[0206] In one embodiment, the Fc variant antibody is administered by maintenance therapy, such as, e.g., once a week for a period of 6 months or more.

[0207] In one embodiment, the Fc variant antibody is administered by a regimen including one infusion of an Fc variant antibody followed by an infusion of an Fc variant

antibody conjugated to a radioisotope. The regimen may be repeated, e.g., 7 to 9 days later.

[0208] As non-limiting examples, treatment according to the present invention may be provided as a daily dosage of an antibody in an amount of about 0.1-100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 after initiation of treatment, or any combination thereof, using single or divided doses of every 24, 12, 8, 6, 4, or 2 hours, or any combination thereof.

[0209] In some embodiments the Fc variant antibody molecule thereof is used in combination with one or more additional therapeutic agents, e.g. a chemotherapeutic agent. Non-limiting examples of DNA damaging chemotherapeutic agents include topoisomerase I inhibitors (e.g., irinotecan, topotecan, camptothecin and analogs or metabolites thereof, and doxorubicin); topoisomerase II inhibitors (e.g., etoposide, teniposide, and daunorubicin); alkylating agents (e.g., melphalan, chlorambucil, busulfan, thiopeta, ifosfamide, carmustine, lomustine, semustine, streptozocin, decarbazine, methotrexate, mitomycin C, and cyclophosphamide); DNA intercalators (e.g., cisplatin, oxaliplatin, and carboplatin); DNA intercalators and free radical generators such as bleomycin; and nucleoside mimetics (e.g., 5-fluorouracil, capecitabine, gemcitabine, fludarabine, cytarabine, mercaptopurine, thioguanine, pentostatin, and hydroxyurea).

[0210] Chemotherapeutic agents that disrupt cell replication include: paclitaxel, docetaxel, and related analogs; vincristine, vinblastin, and related analogs; thalidomide, lenalidomide, and related analogs (e.g., CC-5013 and CC-4047); protein tyrosine kinase inhibitors (e.g., imatinib mesylate and gefitinib); proteasome inhibitors (e.g., bortezomib); NF- κ B inhibitors, including inhibitors of I κ B kinase; antibodies which bind to proteins overexpressed in cancers and thereby downregulate cell replication (e.g., trastuzumab, rituximab, cetuximab, and bevacizumab); and other inhibitors of proteins or enzymes known to be upregulated, over-expressed or activated in cancers, the inhibition of which downregulates cell replication.

[0211] In some embodiments, the antibodies of the invention can be used prior to, concurrent with, or after treatment with Velcade® (bortezomib).

EXAMPLES

[0212] Examples are provided below to illustrate the present invention. These examples are not meant to constrain the present invention to any particular application or theory of operation. For all constant region positions discussed in the present invention, numbering is according to the EU index as in Kabat (Kabat et al., 1991, *Sequences of Proteins of Immunological Interest*, 5th Ed., United States Public Health Service, National Institutes of Health, Bethesda, entirely incorporated by reference). Those skilled in the art of antibodies will appreciate that this convention consists of nonsequential numbering in specific regions of an immunoglobulin sequence, enabling a normalized reference to conserved positions in immunoglobulin families. Accordingly, the positions of any given immunoglobulin as defined by the EU index will not necessarily correspond to its sequential sequence.

Example 1

DNA Construction, Expression, and Purification of Fc Variants

[0213] Amino acid modifications were engineered in the Fc region of IgG antibodies to improve their affinity for the neonatal Fc receptor FcRn. Variants were screened in the context of a number of different human IgG constant chains (FIG. 2), including IgG1, IgG2, and a hybrid IgG sequences that contains the CH1 and upper hinge of IgG1 and the Fc region of IgG2. It will be appreciated by those skilled in the art that because of the different interactions of the IgG1 and IgG2 Fc region with FcγRs and complement, these different parent Fc regions will have different FcγR- and complement-mediated effector function properties. Exemplary sequences of Fc variants in the context of these parent IgG constant chains are shown in FIG. 3.

[0214] Fc variants were engineered in the context of an antibody targeting vascular endothelial factor (VEGF). The heavy and light chain variable regions (VH and VL) are those of a humanized version of the antibody A4.6.1, also referred to as bevacizumab (Avastin®), which is approved for the treatment of a variety of cancers. The amino acid sequences of the VH and VL regions of this antibody are shown in FIG. 4.

[0215] Genes encoding the heavy and light chains of the anti-VEGF antibodies were constructed in the mammalian expression vector pTT5. Human IgG1 and IgG2 constant chain genes were obtained from IMAGE clones and subcloned into the pTT5 vector. The IgG1/2 gene was constructed using PCR mutagenesis. VH and VL genes encoding the anti-VEGF antibodies were synthesized commercially (Blue Heron Biotechnologies, Bothell Wash.), and subcloned into the vectors encoding the appropriate CL, IgG1, IgG2, and IgG1/2 constant chains. Amino acid modifications were constructed using site-directed mutagenesis using the QuikChange® site-directed mutagenesis methods (Stratagene, La Jolla Calif.). All DNA was sequenced to confirm the fidelity of the sequences.

[0216] Plasmids containing heavy chain gene (VH—Cγ1-Cγ2-Cγ3) were co-transfected with plasmid containing light chain gene (VL—Cκ) into 293E cells using lipofectamine (Invitrogen, Carlsbad Calif.) and grown in FreeStyle 293 media (Invitrogen, Carlsbad Calif.). After 5 days of growth, the antibodies were purified from the culture supernatant by protein A affinity using the MabSelect resin (GE Healthcare). Antibody concentrations were determined by bicinchoninic acid (BCA) assay (Pierce).

Example 2

Fc Variant Antibodies Maintain Binding to Antigen

[0217] The fidelity of the expressed variant antibodies was confirmed by demonstrating that they maintained specificity for antigen. VEGF binding was monitored using surface plasmon resonance (SPR, Biacore), performed using a Biacore 3000 instrument. Recombinant VEGF (VEGF-165, Pepro-Tech, Rocky Hill, N.J.) was adhered to a CM5 chip surface by coupling with N-hydroxysuccinimide/N-ethyl-N'-(-3-dimethylamino-propyl) carbodiimide (NHS/EDC) using standard methods. WT and variant antibodies were injected as analytes, and response, measured in resonance units (RU), was acquired. The dissociation phase was too slow to measure true equilibrium constants, and so relative binding was deter-

mined by measuring RU's at the end of the association phase, which should be proportional to the protein concentration (which is held constant in the experiment) and the association rate constant. The data (FIG. 5) show that the variant anti-VEGF antibodies maintain binding to antigen, in contrast to the negative control anti-Her2 antibody which does not bind VEGF.

Example 3

Measurement of Binding to Human FcRn

[0218] Binding of variant antibodies to human FcRn was measured at pH 6.0, the pH at which it is naturally bound in endosomes. Vectors encoding beta 2 microglobulin and His-tagged alpha chain genes of FcRn were constructed, co-transfected into 293T cells, and purified using nickel chromatography. Antibody affinity for human FcRn (hFcRn) at pH 6.0 was measured on a Biacore 3000 instrument by coupling human FcRn to a CM5 chip surface using standard NHS/EDC chemistry. WT and variant antibodies were used in the mobile phase at 25-100 nM concentration and response was measured in resonance units. Association and dissociation phases at pH 6.0 were acquired, followed by an injection of pH 7.4 buffer to measure release of antibody from receptor at the higher pH. A cycle with antibody and buffer only provided baseline response, which was subtracted from each sample sensorgram.

[0219] FIG. 6 shows Biacore sensorgrams for binding of native IgG1 and select Fc variant antibodies to human FcRn at the two relevant pH's. The data show that wild-type and variant antibodies bind readily to FcRn chip at pH 6.0 and dissociate slowly at that pH, as they would in the endosome, yet release rapidly at pH 7.4, as they would upon recycling of endosome to the membrane and exposure to the higher pH of serum.

[0220] The FcRn association/dissociation curves did not fit to a simple Langmuir model, possibly due to the antibody and receptor multi-valency or chip heterogeneity. Pseudo-K_a values (referred to as K_a*) were determined by fitting to a conformational change model with the change in refractive index (RI) fixed at 0 RU. These values for select variant antibodies are plotted in FIG. 7. The relative affinity of each variant as compared to its parent IgG was calculated according to the equation Fold=(WT K_a*/Variant K_a*).

Example 4

Pharmacokinetic Experiments in Human FcRn Knock-in Mice

[0221] To test the ability of select variants to improve half-life in vivo, pharmacokinetic experiments were performed in B6 mice that are homozygous knock-outs for murine FcRn and heterozygous knock-ins of human FcRn (mFcRn^{-/-}, hFcRn^{+/+}) (Petkova et al., 2006, Int. Immunol 18(12): 1759-69, entirely incorporated by reference), herein referred to as hFcRn or hFcRn⁺ mice. A single, intravenous tail vein injection of anti-VEGF antibody (2 mg/kg) was given to groups of 4-7 female mice randomized by body weight (20-30 g range). Blood (~50 ul) was drawn from the orbital plexus at each time point, processed to serum, and stored at -80° C. until analysis. Study durations were 28 or 49 days.

[0222] Antibody concentrations were determined using two ELISA assays. In the first two studies (referred to as Study 1 and Study 2), goat anti-human Fc antibody (Jackson

Immune research) was adhered to the plate, wells were washed with PBST (phosphate buffered saline with 0.05% Tween) and blocked with 3% BSA in PBST. Serum or calibration standards were added, followed by PBST washing, addition of europium labeled anti-human IgG (Perkin Elmer), and further PBST washing. The time resolved fluorescence signal was collected. For Studies 3-5, serum concentration was detected using a similar ELISA, but recombinant VEGF (VEGF-165, PeproTech, Rocky Hill, N.J.) was used as capture reagent and detection was carried out with biotinylated anti-human kappa antibody and europium-labeled streptavidin. PK parameters were determined for individual mice with a non-compartmental model using WinNonLin (Pharsight Inc, Mountain View Calif.). Nominal times and dose were used with uniform weighing of points. The time points used (lambda Z ranges) were from 4 days to the end of the study, although all time points were used for the faster clearing mutants, P257N and P257L.

[0223] Five antibody PK studies in mFcRn^{-/-} hFcRn⁺ mice were carried out. FIG. 8a-8b shows serum concentration data for WT and variant IgG1 (Study 3) and IgG2 (Study 5) antibodies respectively. Fitted PK parameters from all in vivo PK studies carried out in mFcRn^{-/-} hFcRn⁺ mice are provided in FIG. 9, PK data include half-life, which represents the beta phase that characterizes elimination of antibody from serum, C_{max}, which represents the maximal observed serum concentration, AUC, which represents the area under the concentration time curve, and clearance, which represents the clearance of antibody from serum. Also provided for each variant is the calculated fold improvement or reduction in half-life relative to the IgG1 or IgG2 parent antibody [Fold half-life=half-life(variant)/half-life (WT)].

Example 5

Pharmacokinetic Experiment in Nonhuman Primates

[0224] The PK properties of biologics in non-human primates are well-established to be predictive of their properties in humans. A PK study was carried out in cynomolgus monkeys (*macaca fascicularis*) in order to evaluate the capacity of the variant anti-VEGF antibodies to improve serum half-life in non-human primates.

[0225] In preparation for a PK study in cynomolgus monkeys, binding of the variant antibodies to cynomolgus (cyno) FcRn (cFcRn) at pH 6.0 was measured. cFcRn was constructed, expressed, and purified as described above for human FcRn. Binding of variant anti-VEGF antibodies to cFcRn was measured using Biacore as described above. The data is provided in FIG. 10a-10b. The results show that the variants improve affinity for cyno FcRn similarly as they do for human FcRn. Dissociation at the higher pH (7.4) was also very rapid (data not shown), similar to as was observed for binding to human FcRn. These results are not surprising given the high sequence homology of the human and cyno receptors (FcRn alpha chain 96%, beta-2-microglobulin 91%).

[0226] The PK of the variants were studied in vivo in non-human primates. Male cynomolgus monkeys (*macaca fascicularis*, also called crab-eating Macaque) weighing 2.3-5.1 kg were randomized by weight and divided into 5 groups with 3 monkeys per group. The monkeys were given a single, 1 hour peripheral vein infusion of 4 mg/kg antibody. Blood samples (1 ml) were drawn from a separate vein from 5

minutes to 90 days after completion of the infusion, processed to serum and stored at -70 C. Animals were not harmed during these studies.

[0227] Antibody concentrations were determined using the VEGF capture method as described above. PK parameters were determined by fitting the concentrations versus time to a non-compartmental model as was done in the mouse PK studies. However, time points from day 10 to day 90 were used for PK parameter determinations. The PK results are plotted in FIG. 11, and the fitted parameters are provided in FIG. 12. The results show that the variants enhanced the in vivo half-life of antibody up to 3.2-fold. In the best case (the 428L/434S variant) half-life was extended from 9.7 days to 31.1 days. The PK results obtained in cynomolgus monkeys are consistent with those obtained in mFcRn^{-/-} hFcRn⁺ mice, validating the hFcRn mouse model as a system for assessing the in vivo PK properties of the variants, and supporting the conclusions from those studies.

Example 6

Engineering Additional Fc Variants that Extend In Vivo Half-Life

[0228] New variants were engineered to further screen for modifications that extend in vivo half-life. Designed substitutions are shown in FIG. 13, Variants were constructed in the context of an antibody with the bevacizumab variant region and IgG1/2 N434S constant region; however, variants may be constructed in any IgG isotype. Variants were constructed, expressed, and purified as described above.

[0229] Variants were screened for binding to human FcRn at pH 6.0 by Biacore. Anti-VEGF antibodies were captured to a VEGF coupled chip, a single concentration of FcRn analyte was flowed over the chip (association phase), and then buffer was washed over to dissociate FcRn analyte (dissociation phase). The dissociation off-rate (k_{off}) was determined by fitting the dissociation phase data to a 1:1 Langmuir dissociation model. The results are shown numerically in FIG. 14, and plotted in FIG. 15. As can be seen, a number of the designed variants improve FcRn binding (reduce the k_{off}) relative to the IgG1/2 N434S background.

[0230] The FcRn binding of select variants was tested using the same Biacore format but with an analyte (FcRn) concentration series in order to obtain accurate equilibrium dissociation constants (K_D s). Sensorgrams at multiple analyte concentrations were fit globally to a 1:1 Langmuir binding model. Resulting kinetic and equilibrium binding constants are presented in FIG. 16, along with the Fold K_D and Fold k_{off} relative to IgG1 WT and IgG1/2 N434S respectively. Data are plotted in FIGS. 17 and 18. A number of variants, comprising several different substitutions improve binding to human FcRn. Beneficial substitutions include but are not limited to T307Q, Q311I, Q311V, A378V, S426V, S426T, Y436I, and Y436V. The successful engineering at positions 378 and 426 is surprising giving that they are more distal to the FcRn binding interface on Fc.

[0231] Additional substitutions were designed to explore whether other substitutions at these positions may provide improved binding to FcRn. These variants are listed in FIG. 19. Based on the collective data, a new library of variant combinations was designed to further screen for Fc variants that have the potential to extend antibody half-life in vivo. These variants are listed in FIG. 20.

[0232] Variants were constructed, expressed, and purified as described above. Variants were screened for binding to human FcRn at pH 6.0 by Biacore. Anti-VEGF antibodies were captured to a VEGF coupled chip, a single concentration of FcRn analyte was flowed over the chip (association phase), and then buffer was washed over to dissociate FcRn analyte (dissociation phase). The dissociation off-rate (k_{off}) was determined by fitting the dissociation phase data to a 1:1 Langmuir dissociation model. The results are shown numerically in FIG. 21, and plotted in FIG. 22. As can be seen, a number of the designed variants improve FcRn binding (reduce the k_{off}).

[0233] In vivo pharmacokinetic properties of the variants were tested in hFcRn⁺ mice as described above. A single, intravenous tail vein injection of anti-VEGF antibody (2

mg/kg) was given. Antibody concentrations were determined using the VEGF-capture ELISA assay described above. FIG. 23 shows mean serum concentration data for Fc variant and IgG1/2 parent anti-VEGF antibodies. Fitted half-lives of individual mice and means are provided in FIG. 24. FIG. 25 shows a scatter plot of the fitted half-lives from the individual mice in the study. The results indicate that the engineered Fc variants extend half-life relative to the parent IgG1/2 antibody, as well as the IgG1/2 N434S variant.

[0234] Whereas particular embodiments of the invention have been described above for purposes of illustration, it will be appreciated by those skilled in the art that numerous variations of the details may be made without departing from the invention as described in the appended claims. All references cited herein are incorporated in their entirety.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
1          5          10          15
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
20          25          30
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
35          40          45
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
50          55          60
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
65          70          75          80
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
85          90          95
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
100          105

```

<210> SEQ ID NO 2

<211> LENGTH: 330

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1          5          10          15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20          25          30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35          40          45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50          55          60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65          70          75          80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

```

-continued

85					90					95				
Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Cys
			100					105					110	
Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
			115				120					125		
Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Cys
			130				135					140		
Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn
							150							160
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
				165					170					175
Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
				180				185					190	Leu
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser
			195				200					205		Asn
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
			210				215					220		Gly
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu
							230							240
Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe
				245					250					255
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
				260				265					270	Asn
Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe
				275				280				285		Phe
Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly
				290				295				300		Asn
Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr
							310					315		320
Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys					
				325					330					

<210> SEQ ID NO 3

<211> LENGTH: 326

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg
1				5					10				15		
Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
			20					25					30		
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
			35					40					45		
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
			50				55					60			
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe	Gly	Thr	Gln	Thr
			65				70					75			80
Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
				85				90					95		
Thr	Val	Glu	Arg	Lys	Cys	Cys	Val	Glu	Cys	Pro	Pro	Cys	Pro	Ala	Pro
			100					105					110		
Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp

-continued

115					120					125					
Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
130						135					140				
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly
145					150					155					160
Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn
				165					170					175	
Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val	His	Gln	Asp	Trp
			180					185					190		
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro
	195					200					205				
Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg	Glu
210					215						220				
Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn
225					230					235					240
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile
			245						250					255	
Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr
			260					265					270		
Thr	Pro	Pro	Met	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys
	275					280						285			
Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
	290					295					300				
Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu
305					310					315					320
Ser	Leu	Ser	Pro	Gly	Lys										
				325											

<210> SEQ ID NO 4

<211> LENGTH: 377

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg
1				5					10					15	
Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
			20					25					30		
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
		35					40					45			
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
	50					55					60				
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr
65					70					75				80	
Tyr	Thr	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
			85					90						95	
Arg	Val	Glu	Leu	Lys	Thr	Pro	Leu	Gly	Asp	Thr	Thr	His	Thr	Cys	Pro
			100					105					110		
Arg	Cys	Pro	Glu	Pro	Lys	Ser	Cys	Asp	Thr	Pro	Pro	Pro	Cys	Pro	Arg
		115					120					125			
Cys	Pro	Glu	Pro	Lys	Ser	Cys	Asp	Thr	Pro	Pro	Pro	Cys	Pro	Arg	Cys
	130					135					140				
Pro	Glu	Pro	Lys	Ser	Cys	Asp	Thr	Pro	Pro	Pro	Cys	Pro	Arg	Cys	Pro

-continued

145	150	155	160
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys	165	170	175
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val	180	185	190
Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Lys Trp Tyr	195	200	205
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu	210	215	220
Gln Tyr Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Leu His	225	230	235
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys	245	250	255
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln	260	265	270
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met	275	280	285
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro	290	295	300
Ser Asp Ile Ala Val Glu Trp Glu Ser Ser Gly Gln Pro Glu Asn Asn	305	310	315
Tyr Asn Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu	325	330	335
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Ile	340	345	350
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln	355	360	365
Lys Ser Leu Ser Leu Ser Pro Gly Lys	370	375	

<210> SEQ ID NO 5
 <211> LENGTH: 327
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 5

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg	1	5	10	15
Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr	20	25	30	
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser	35	40	45	
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser	50	55	60	
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr	65	70	75	80
Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys	85	90	95	
Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro	100	105	110	
Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys	115	120	125	
Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val				

-continued

130	135	140
Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp 145 150 155 160		
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe 165 170 175		
Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp 180 185 190		
Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu 195 200 205		
Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg 210 215 220		
Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys 225 230 235 240		
Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp 245 250 255		
Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys 260 265 270		
Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser 275 280 285		
Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser 290 295 300		
Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser 305 310 315 320		
Leu Ser Leu Ser Leu Gly Lys 325		

<210> SEQ ID NO 6

<211> LENGTH: 329

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: IgG1/2 constant heavy chain (CH1-hinge-CH2-CH3)

<400> SEQUENCE: 6

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys 1 5 10 15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr 20 25 30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser 35 40 45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser 50 55 60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr 65 70 75 80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys 85 90 95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys 100 105 110
Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys 115 120 125
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val 130 135 140
Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr 145 150 155 160

```
<210> SEQ ID NO 7
<211> LENGTH: 326
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IqG2 434S
```

Ala 1	Ser	Thr	Lys 5	Gly	Pro	Ser	Val	Phe 10	Pro	Leu	Ala	Pro	Cys 15	Ser	Arg
Ser	Thr	Ser	Glu 20	Ser	Thr	Ala	Ala	Leu 25	Gly	Cys	Leu	Val	Lys 30	Asp	Tyr
Phe	Pro	Glu 35	Pro	Val	Thr	Val	Ser 40	Trp	Asn	Ser	Gly	Ala 45	Leu	Thr	Ser
Gly 50	Val	His	Thr	Phe	Pro 55	Ala	Val	Leu	Gln	Ser	Ser 60	Gly	Leu	Tyr	Ser
Leu 65	Ser	Ser	Val	Val	Thr 70	Val	Pro	Ser	Ser	Asn 75	Phe	Gly	Thr	Gln	Thr 80
Tyr	Thr	Cys	Asn 85	Val	Asp	His	Lys	Pro	Ser 90	Asn	Thr	Lys	Val	Asp 95	Lys
Thr	Val	Glu	Arg 100	Lys	Cys	Cys	Val	Glu 105	Cys	Pro	Pro	Cys	Pro 110	Ala	Pro
Pro	Val	Ala 115	Gly	Pro	Ser	Val	Phe 120	Leu	Phe	Pro	Pro	Lys 125	Pro	Lys	Asp
Thr 130	Leu	Met	Ile	Ser	Arg 135	Thr	Pro	Glu	Val	Thr	Cys 140	Val	Val	Val	Asp
Val 145	Ser	His	Glu	Asp 150	Pro	Glu	Val	Gln	Phe	Asn 155	Trp	Tyr	Val	Asp	Gly 160
Val	Glu	Val	His 165	Asn	Ala	Lys	Thr	Lys 170	Pro	Arg	Glu	Glu	Gln	Phe	Asn 175

-continued

```

Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
      180                      185                      190

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
      195                      200                      205

Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
      210                      215                      220

Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
      225                      230                      235                      240

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
      245                      250                      255

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
      260                      265                      270

Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
      275                      280                      285

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
      290                      295                      300

Ser Val Met His Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu
      305                      310                      315                      320

Ser Leu Ser Pro Gly Lys
      325

```

```

<210> SEQ ID NO 8
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IgG1/2 434S

```

```

<400> SEQUENCE: 8

```

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20      25      30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35      40      45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50      55      60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
      65      70      75      80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      100     105     110

Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
      115     120     125

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
      130     135     140

Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
      145     150     155     160

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
      165     170     175

Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His
      180     185     190

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys

```

-continued

195	200	205
Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln 210 215 220		
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met 225 230 235 240		
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro 245 250 255		
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn 260 265 270		
Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu 275 280 285		
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val 290 295 300		
Phe Ser Cys Ser Val Met His Glu Ala Leu His Ser His Tyr Thr Gln 305 310 315 320		
Lys Ser Leu Ser Leu Ser Pro Gly Lys 325		

<210> SEQ ID NO 9
 <211> LENGTH: 123
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Anti-VEGF VH

<400> SEQUENCE: 9

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly 1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr 20 25 30
Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45
Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe 50 55 60
Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr 65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95
Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val 100 105 110
Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser 115 120

<210> SEQ ID NO 10
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Anti-VEGF VL

<400> SEQUENCE: 10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly 1 5 10 15
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr 20 25 30
Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

-continued

35	40	45
Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly		
50	55	60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
65	70	75
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp		
85	90	95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys		
100	105	

<210> SEQ ID NO 11
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Peptidyl Linker

<400> SEQUENCE: 11

Gly Phe Leu Gly
1

1. An antibody comprising a variant Fc region as compared to a parent Fc region, wherein said variant Fc region comprises a first mutation which is a serine at position 434 and a second mutation selected from a group consisting of: an isoleucine at position 311, a valine at position 311, an isoleucine at position 436, and a valine at position 436, wherein said antibody has increased serum half-life as compared to an antibody comprising said parent Fc region, and wherein numbering is according to the ELI index.

2. An antibody comprising a variant Fc region as compared to a parent Fc region, wherein said variant Fc region comprises a first mutation which is a serine at position 434 and a second mutation selected from a group consisting of: an isoleucine at position 311, a valine at position 311, an isoleucine at position 436, and a valine at position 436, wherein said antibody has increased binding affinity to a human FcRn receptor as compared to an antibody comprising said parent Fc region, and wherein numbering is according to the EU index.

3. A nucleic acid encoding the variant Fc region of claim 1 or 2.

4. A host cell comprising the nucleic acid of claim 21.

5. A method comprising administering an antibody to a subject, said method comprising administering an antibody comprising a variant Fc region as compared to a parent Fc region, wherein said variant Fc region comprises a first mutation which is a serine at position 434 and a second mutation selected from a group consisting of: an isoleucine at position 311, a valine at position 311, an isoleucine at position 436, and a valine at position 436, wherein said antibody has increased serum half-life as compared to an antibody comprising said parent Fc region, and wherein numbering is according to the EU index.

* * * * *